PROCEEDINGS

OF THE

NEW ENGLAND ZOÖLOGICAL CLUB

A NEW RAIL FROM SOUTHERN CALIFORNIA.

BY OUTRAM BANGS.

The large rail that inhabits the marshes of southern California differs very much from *Rallus obsoletus* of the marshes of San Francisco Bay, and quite as much from *R. beldingi* of Lower California. It is not a common bird in collections; still there are several skins in the National Museum, several in Mr. William Brewster's collection, and one in my own. All of these I have examined, and I find very little variation among them. Its characters are so well marked and so constant that I regard it as a distinct species, and I propose to call it

Rallus levipes1 sp. nov.

Type, from Newport Landing, Los-Angeles Co., California, Q adult, no. 678, coll. of E. A. and O. Bangs, collected Feb. 23, 1886, by F. Stephens.

Characters.—Much smaller than either R. obsoletus or R. beldingi; bill much more slender; tarsus and foot smaller than in either. In color it differs from R. obsoletus in being much darker above — more olive and less grayish, brown; in having breast and sides of neck deep cinnamon-rufous instead of grayish cinnamon, this color extending well up on sides of neck and meeting the color of upper parts sharply; ground color of flanks, etc., darker—less grayish; a gray patch behind eye; superciliary streak white instead of rusty. From R. beldingi it differs, in color, in having the back feathers much less decidedly streaked; breast, etc., less pinkish or salmon-colored; flanks, etc., browner—

^{· 1} Levipes - light-footed.

without dusky bars bordering the white ones, the white bars wider; superciliary streak white instead of rusty.

Color.—Above, olive brown, broadly striped with blackish brown; breast, etc., deep cinnamon-rufous, extending over sides of neck and meeting color of upper parts sharply; flanks, etc., olive, broadly banded with white; centre of belly and throat whitish; superciliary streak white; a gray patch on side of head behind eye.

Measurements.— Type, Q adult; wing, 152.; tail, 58.2; tarsus, 49.; culmen, 58. mm. Topotype, Q adult, no. 47,848, coll. of Wm. Brewster: wing, 141.; tail, 56.6; tarsus, 45.; culmen, 54.2 mm.



Bangs, Outram. 1899. "A new rail from southern California." *Proceedings of the New England*

Zoo

200

logical Club 1, 45-46.

View This Item Online: https://www.biodiversitylibrary.org/item/45427

Permalink: https://www.biodiversitylibrary.org/partpdf/144972

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.