

A WHOOPING CRANE FROM THE PLEISTOCENE OF NORTH FLORIDA

STORRS L. OLSON¹

Department of Pathobiology
The Johns Hopkins University
School of Hygiene and Public Health
Baltimore, Maryland 21205

In a small collection of bird fossils given to me for study by Dr. Richard Ohmes of Chaires, Florida, there is a distal two-thirds of a right tibiotarsus of a Whooping Crane (*Grus americana*). It was collected in 1969 by Dr. Ohmes from river-bottom deposits in a branch of the Aucilla River on the west side of Ward Island in the NE $\frac{1}{4}$ sec. 13, T. 4 S., R. 3 E., Jefferson County, Florida, and is now number 170878 in the paleontological collections of the U.S. National Museum. Because of its manner of deposition, it is not possible to age the bone precisely, but it was associated with numerous remains of Pleistocene to Recent vertebrates including such forms as *Megalonyx*, *Mammut*, *Tapirus*, *Equus*, *Camelops*, and *Geochelone*, which are either extinct or no longer present in Florida, and it is safe to assume its contemporaneity with these animals. Analysis of the other avian elements is not yet complete but so far indicates a mixture of fresh and salt water avifaunas of extant species.

Recent Florida records of Whooping Cranes have generally been subject to doubt (Sprunt 1954), but Hallman (1965) has published photographs of a mounted specimen said to have been obtained in 1927 or 1928, 12 miles N of St. Augustine, St. Johns County. Other Florida fossil records are from Ich-tucknee River, Columbia County; Seminole Field,

Pinellas County; and Melbourne, Brevard County (Wetmore 1931). In addition, Brodkorb (pers. comm.) informs me that his record (Neill et al. 1956) of a Sandhill Crane (*Grus canadensis*), based on a proximal portion of a left carpometacarpus from Lemon Bluff on the St. Johns River, Volusia County, actually pertains to *G. americana*. The Aucilla River specimen represents the westernmost locality for the state and is the first indication of the species in the panhandle. The Whooping Crane is also known from Pleistocene fossils or prehistoric remains from California, North Dakota, Illinois, Idaho, Michigan, Kentucky, Virginia (Brodkorb 1967), and Arizona (Cra-craft 1968).

I would like to thank D. Bruce Means and Pierce Brodkorb for their assistance with this note, and Tall Timbers Research Station for providing facilities and support for myself and Means to carry out research at the Aucilla site.

LITERATURE CITED

- BRODKORB, P. 1967. Catalogue of fossil birds. Part 3 (Ralliformes, Ichthyornithiformes, Charadriiformes). Bull. Fla. State Mus. 11:99-220.
- CRA-CRAFT, J. 1968. The Whooping Crane from the Lower Pleistocene of Arizona. Wilson Bull. 80: 490.
- HALLMAN, R. C. 1965. Record of Whooping Crane (*Grus americana*) killed in St. Johns County, Florida. Fla. Natur. 38:23.
- NEILL, W. T., H. J. CUT, AND P. BRODKORB. 1956. Animal remains from four preceramic sites in Florida. Amer. Antiquity 21:383-395.
- SPRUNT, A. 1954. Florida bird life. Coward-Mc-Cann, New York.
- WETMORE, A. 1931. The avifauna of the Pleistocene in Florida. Smithsonian Misc. Coll. 85:1-41.

¹Present address: Department of Vertebrate Zoology, Smithsonian Institution, Washington, D.C. 20560.