



## Spatial and temporal abilities of proxies used to detect pre-Columbian Indigenous human activity in Amazonian ecosystems

C.N.H. McMichael<sup>a,\*</sup>, C. Levis<sup>b,c,d</sup>, W.D. Gosling<sup>a</sup>, A.B. Junqueira<sup>e</sup>, D.R. Piperno<sup>f</sup>, E.G. Neves<sup>g</sup>, F. Mayle<sup>h</sup>, M. Peña-Claros<sup>b</sup>, F. Bongers<sup>b</sup>

<sup>a</sup> Department of Ecosystem and Landscape Dynamics, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, the Netherlands

<sup>b</sup> Forest Ecology and Forest Management Group, Wageningen University & Research, Wageningen, the Netherlands

<sup>c</sup> Post-graduate Program in Ecology, Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil

<sup>d</sup> Post-graduate Program in Ecology, Universidade Federal de Santa Catarina, Florianópolis, Brazil

<sup>e</sup> Institut de Ciència i Tecnologia Ambientals, Universitat Autònoma de Barcelona, Barcelona, Spain

<sup>f</sup> Department of Anthropology, Smithsonian National Museum of Natural History, Washington, D.C., and Smithsonian Tropical Research Institute, Balboa, Panama

<sup>g</sup> Laboratory of Tropical Archaeology, Museum of Archaeology and Ethnology, University of São Paulo, São Paulo, Brazil

<sup>h</sup> Department of Geography and Environmental Science, University of Reading, Reading, United Kingdom

### ARTICLE INFO

Handling Editor: Dr Mira Matthews

#### Keywords:

Archaeology  
Paleoecology  
Ethnography  
Human impacts  
Plant genetics  
Biological collections  
Past human activity  
Remote sensing

### ABSTRACT

Humans have been modifying ecosystems since before the Holocene began ca. 12,000 years ago, even in Neotropical regions. The Amazon was once thought to be 'pristine' and only lightly impacted by Indigenous people before European colonization in the Americas (e.g., pre-Columbian); however, multiple lines of evidence have shown that Indigenous human activities over the past millennia have left ecological legacies on modern ecosystems. We review the various lines of evidence used to reconstruct pre-Columbian Indigenous human activity in Amazonia, and assess the spatial and temporal resolution and limits of each one of them. We suggest that a multi-proxy approach is always preferred, and that lines of evidence that cover overlapping yet discrete spatial and temporal scales can provide a robust and comprehensive assessment of the nuances of pre-Columbian Indigenous human activities in Amazonia, and how they affect modern ecosystems.

### 1. Introduction

Humans are now considered the major driving force in many abiotic and biotic processes on Earth, and it has been suggested that this has created a distinct geological era called the Anthropocene (e.g., [Crutzen, 2002](#); [Di Marco and Santini, 2015](#); [Gallardo et al., 2015](#); [Halpern et al., 2008](#); [Lewis and Maslin, 2015](#); [Pachauri et al., 2015](#); [Vitousek et al., 1997](#)). Humans contributed to the extinction of Pleistocene megafauna across the Neotropics ([Barnosky and Lindsey, 2010](#); [Rozas-Davila et al., 2016, 2021](#)), and have been modifying landscapes in Neotropical ecosystems for at least the last 12,000 years ([Ellis et al., 2021](#); [Roosevelt, 2013](#)). Forms of human activities, including plant cultivation and domestication, have also occurred in Neotropical regions for at least 10,000 years ([Lombardo et al., 2020](#); [Piperno, 2011](#); [Roberts et al., 2017](#)). Many of these activities, particularly those that have occurred over the last 2000 years, likely left persistent effects, or ecological legacies, on

Neotropical ecosystems that are still visible today ([Furquim et al., 2023](#); [Levis et al., 2017](#); [McMichael, 2021](#); [McMichael et al., 2023](#); [Ross, 2011](#)).

Reconstructing human-environment interactions through time is particularly important in the ca. 6 million km<sup>2</sup> of Amazonian rainforests, which hold a large proportion of Earth's biodiversity ([Olson et al., 2001](#)). The suggestion that pre-Columbian Indigenous human influence played a large role in shaping the biodiversity observed in today's ecosystems has generated an important debate (e.g. [Balée, 2010](#); [Clement et al., 2015](#); [Levis et al., 2017](#); [Piperno et al., 2021](#); [Piperno et al., 2019](#); [Roosevelt, 2013](#)). Most scholars from various disciplines agree that the pre-Columbian Indigenous Peoples of Amazonia influenced its ecosystems to some degree, but the intensity, cultural variability, spatial extensiveness, spatial variability, temporal duration and continuity remain debated ([Barlow et al., 2012](#); [Bush et al., 2015](#); [Clement et al., 2015](#); [Heckenberger et al., 2003, 2008](#); [Levis et al., 2012](#),

\* Corresponding author. Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, 904 Science Park, 1098 XH, Amsterdam, the Netherlands.  
E-mail address: [c.n.h.mcmichael@uva.nl](mailto:c.n.h.mcmichael@uva.nl) (C.N.H. McMichael).


2017; McMichael et al., 2012a; Piperno et al., 2015). Much of the controversy results from the overall paucity of data in the region, differences in the interpretation of the same datasets, and on the ecological heterogeneity of this vast area. Even the largest ecological datasets within Amazonia reflect mostly relatively easily accessible areas (McMichael et al., 2017). Addressing aspects of this debate has important implications for how we conserve and manage modern ecosystems, as well as for Indigenous sovereignty.

Ecologists, paleoecologists, archaeologists, anthropologists, ethnographers, Indigenous peoples, and local communities can all provide valuable information about the activities of pre-Columbian Indigenous people and their influence on Amazonian vegetation and landscapes. These lines of information, however, come from a variety of sources, are measured at different spatial and temporal scales, and the capabilities of reconstructing pre-Columbian human activities vary between them. Integrating different data and knowledge sources, however, is vital for understanding multifaceted and complex problems, and to achieve a more reliable and robust assessment of the influence of pre-Columbian activities on Amazonian ecosystems (Mayle and Iriarte, 2014; McClellan et al., 2015; Tengö et al., 2014).

A large part of data interpretation is recognizing the advantages and limitations of the various proxies, methods, and settings used to reconstruct pre-Columbian Indigenous human activities and their effect on tropical systems. Here we review proxies (i.e., lines of evidence) and settings (i.e., depositional environments or archival material) commonly used to assess pre-Columbian human activities and influences on Amazonian ecosystems, with a specific focus on the spatial and temporal resolution (the potential sampling frequency of the archival material) and spatial and temporal limits (the total amount of space or time potentially captured within the record). We also discuss the detectability and variability of proxies used to assess pre-Columbian Indigenous human activities and their influences on Amazonian ecosystems (Fig. 1). Our goal is to provide a spatio-temporal framework for the various proxies used to assess long-term Indigenous human activity in Amazonian ecosystems so that future work can recognize the advantages and limitations of all of the proxies and integrate them more comprehensively.

## 2. Soils as local-scale archives of pre-Columbian Indigenous human activities and vegetation change

Soils contain a wealth of valuable information and proxies that can be used to reconstruct pre-Columbian human activities and the influence of Indigenous Peoples on the vegetation. Archaeological sites, artefacts, macrofossils, and microfossils that directly document pre-Columbian human activities are all found on, or in, tropical terrestrial soils (Figs. 1 and 2). Information from archaeological surveys is beginning to



**Fig. 1.** The array of proxies that can be used to detect pre-Columbian Indigenous human activity in Amazonia, and the source, or archival material from which they are derived. The proxies are shown on a gradient from those which are used specifically to assess human activities (far right) to those which are used to assess both human and non-human processes (far left). ADE = Amazonian Dark Earths.



**Fig. 2.** Examples of directly documented evidence of human activity found in Amazonian soils. The top panel shows an archaeological excavation, and the middle panel shows ceramic artifacts that were uncovered during the excavation (Photos: Bernardo Oliveira/Instituto Mamirauá). The bottom panel shows a Google Earth image of earthworks that were uncovered in southwestern Amazonia following deforestation of the landscape.

be compiled into online databases, including the AmazonArch (Amazonian Archaeological Sites Network), which contains the geographical location and basic archaeological information for over 10,000 sites (Clement et al., 2015; Riris and Arroyo-Kalin, 2019; WinklerPrins and Aldrich, 2010; <https://sites.google.com/view/amazonarch>). The variability and diversity of archaeological evidence includes sites containing artifacts such as lithics, ceramics, rock paintings, or petroglyphs, earthworks, and anthropogenic soils (called Amazonian Dark Earths, or ADE) (e.g., Neves et al., 2021) (examples of ceramics and

earthworks shown in Fig. 2). At some of these sites, pre-Columbian people have modified soil texture, nutrient content, and stable isotopes, leaving persistent legacies of soil properties for thousands of years (Glaser and Birk, 2012; Glaser and Woods, 2004; Lehman et al., 2003).

Phytoliths are microscopic bodies of silica that are produced in the cells of many plant species and often preserved in high quantities in soil archives where pollen and macrofossils have largely decayed (Piperno, 2006) (example shown in Fig. 3). Phytolith analysis is commonly performed on samples obtained at archaeological sites, but also in soils that are not associated with archaeological settings such as lake sediments and terrestrial soils (Fig. 1). Phytoliths often reflect localized plant decay, particularly in terrestrial soils, as they do not have intrinsic dispersal mechanisms that would carry them considerable distances from the depositional environment as can be the case with pollen. Studies of phytoliths in surface soils from tropical forests in Panama showed that phytolith movement, via fire, or surface water transport, can be as little as 25–30 m from their source area (Piperno, 1988) or up to 100–120 m from their source plant (Piperno, pers. Comm.; Piperno and McMichael, manuscript in preparation). Further, lakes with in-flowing streams may contain phytoliths from considerable distances, and in fluvial forest soils from Brazil, phytolith movement from areas a substantial distance upstream is indicated (Watling et al., 2016). Seasonally flooded savannas might be expected to be similar.

Phytoliths directly document different types of vegetation, such as savanna, open forests with bamboo, evergreen, semi-evergreen, and deciduous forest, and early successional growth typical of human disturbance. Phytoliths also directly detect some major and now-minor crops and other economic plants, (e.g., maize [*Zea mays* L.], *Cucurbita* spp. [squashes and gourds], manioc [*Manihot esculenta* Cranz], arrowroot [*Maranta arundinacea* L.], and various palm species) (Carson et al., 2014; Dickau et al., 2013; Lombardo et al., 2020; McMichael et al., 2012a, 2012b; Piperno, 2006; Watling et al., 2016; Whitney et al., 2013, 2014). All palms, grasses, and sedges, and many arboreal basal angiosperms and eudicotyledons, produce high phytolith numbers diagnostic to at least the family, and not uncommonly, genus levels (Huisman et al., 2018; Morcote-Ríos et al., 2015, 2016; Piperno, 2006; Piperno and McMichael, 2023; Piperno et al., 2019; Witteveen et al., 2022). The high phytolith production of palms and grasses also means that their absence in the record actually represents the absence of these taxa in the vegetation (Piperno et al., 2015). Not all plants produce phytoliths, however, and some taxa will remain undetectable in reconstructions (Piperno, 2006). This is the case with some major economically important trees, e.g., *Bertholletia excelsa* (Brazil nut), *Annona* spp. (soursop, cherimoya), guava (*Psidium guajava* L.).

Starch grains (example shown in Fig. 3) can identify certain cultivars, such as maize (*Zea mays* L.), squashes (*Cucurbita* spp.), manioc (*Manihot esculenta* Cranz), and other tuber crops (e.g., Pearsall et al., 2004; Piperno, 2006; Piperno, 2011). Some crops that do not produce phytoliths, such as peanuts (*Arachis hypogaea* L.) and chili peppers (*Capsicum* spp.), have diagnostic starch grains (e.g., Dickau et al., 2007; Ezell et al., 2006; Piperno, 2006). Beans, including *Phaseolus* species, and some palm trees that were used by pre-Columbian Indigenous Peoples also produce identifiable starch grains (Watling et al., 2018). The starch grains are usually isolated from ceramics or stone tools found at archaeological sites (Fig. 1) (Iriarte et al., 2004; Pearsall et al., 2004; Watling et al., 2018; Young et al., 2023). Macrobotanical remains, often carbonized, are typically recovered from site sediments and along with the starch grains and phytoliths, provide empirical evidence for the diets and lifestyles of pre-contact Indigenous Peoples (e.g., Furquim et al., 2021; Watling et al., 2018).

Macroscopic charcoal fragments (e.g., >500 µm) found in soils, including at archaeological sites, represent localized past fire events (e.g., Rhodes, 1998; Whitlock and Larsen, 2002) (example shown in Fig. 3). In the aseasonal forests with a limited dry season (for instance, those found in northwestern Amazonia), forest fire almost always starts with human intervention (Fig. 1) (Bush et al., 2008; Malhi et al., 2008). The




Fig. 3. Microfossils found in soils in Amazonia that can document human activity and environmental changes. The top panel shows phytoliths from the tree *Licania micrantha* Miq. (Chrysobalanaceae), which are diagnostic to at least the genus level (Piperno and McMichael, 2023) (Photo: Dolores Piperno). The middle panel shows starch grains of *Phaseolus vulgaris* L. (beans) (Photo: Dolores Piperno). The bottom panel shows charcoal isolated from Amazonian soils (Photo: Crystal McMichael).


presence of charcoal in these aseasonal forests indicates human activity (Bush et al., 2008), but escaped wildfires in these areas can also occur during extreme droughts (Flores et al., 2017). In drier areas, i.e., the seasonal forests that border savannas in eastern Amazonia, fires are less dependent on human ignition, but are still more frequent when humans are present (Alencar et al., 2004; Maezumi et al., 2015, 2018b; Nepstad et al., 2004; Power et al., 2016; Ramos-Neto and Pivello, 2000). Paleocological and archaeological data show that fire frequency across Amazonia was more frequent in the late Holocene, when climate was wetter than the early-to mid-Holocene (Arroyo-Kalin and Riris, 2021; McMichael and Bush, 2019; Nascimento et al., 2022), highlighting that the primary source of ignition was human activity.

Charcoal retains diagnostic morphological features of the plant from which it is derived, and can be used to identify types of plants, e.g., woody versus non-woody taxa (Bodin et al., 2020; Di Pasquale et al., 2008; Orvis et al., 2005). Charcoal morphology can sometimes provide taxonomic identification to the family-level and sometimes genus or species level in tropical ecosystems, and has been used to characterize land use and successional forests at archaeological sites (Bachelet and Scheel-Ybert, 2017; Bodin et al., 2019; Cartwright, 2015; Fernandes Caromano et al., 2013; Goulart et al., 2017; Iriarte et al., 2020; Scheel-Ybert et al., 2014). The chemical properties (i.e., FTIR spectroscopy) of charcoal fragments can also be used to infer burn temperature of the fire events, and can also distinguish plant types (e.g. woody vs grassy material) that were burned (Gosling et al., 2019; Maezumi et al., 2021).

With soil surveys to reconstruct past fire events, replicate soil cores are typically collected from a given site (100 m–200 m radius) (Hammond et al., 2006; McMichael et al., 2012a, 2012c) to account for the uneven deposition of charcoal that occurs on localized scales after burning of vegetation (e.g. Lynch et al., 2004). Thus, the repeated absence of charcoal from soil cores located in close proximity can be confidently interpreted that the sampled area truly lacked fire rather than the sampling was unable to detect the fire (McMichael et al., 2012a, 2015). Replicate soil cores can also indicate whether large tracts of vegetation were burned or repeatedly burned, especially if some of the particles are  $^{14}\text{C}$  AMS dated (Feldpausch et al., 2022; Heijink et al., 2022; McMichael et al., 2012a; Sanford and Horn, 2000; Whitlock and Larsen, 2002). Replicate cores are often also analysed to look at how vegetation change has occurred over relatively small geographic scales or along environmental gradients (e.g., Heijink et al., 2022; McMichael et al., 2012a; McMichael et al., 2012b; Watling et al., 2017).

Stable carbon and nitrogen isotopes are commonly used to infer vegetation dynamics (Fig. 1) (de Freitas et al., 2001; Pessenda et al., 1998). Stable carbon isotopes from soils have also been used to infer landscape transformations by pre-Columbian Indigenous Peoples. These are particularly useful for documenting changes or shifts between C3 and C4 grass assemblages and vegetation changes on pre-Columbian raised fields (Iriarte et al., 2010; McKey et al., 2010; Watling et al., 2017). The analysis of stable carbon and nitrogen isotopes from bone collagen has also been used to reconstruct dietary changes in pre-Columbian Indigenous Peoples (e.g., Colonese et al., 2020; Roosevelt, 1989).

Soils have a very high spatial resolution as multiple samples can be collected within meters of each other (i.e., high potential sampling frequency) and a very high spatial limit because they are found almost everywhere in terrestrial systems (Fig. 4). The temporal limit of soils is also very high; soils capture evidence from modern times to thousands of years ago (Fig. 4). The uppermost 1 m of Amazonian soils typically represent the last several thousand years (Piperno, 2016; Piperno et al., 2021). Soils, however, have a low temporal resolution due to processes such as soil formation, erosion, and bioturbation, and establishing age-depth relationships is not always possible (e.g., Mayle and Iriarte, 2014; Sanford and Horn, 2000) (Fig. 4a). Radiocarbon ( $^{14}\text{C}$  AMS) dates from archaeological sites or soil microfossils usually have a 2-sigma precision of ca. 100 years (Neves et al., 2004; Piperno, 2016; Schaan et al., 2012; Taylor and Bar-Yosef, 2016). When multiple dates are



**Fig. 4.** The spatial and temporal characteristics of data sources used to assess past human activities and vegetation change. (a) Each type of archival material color-coded and plotted in relation to its spatial resolution (the highest possible sampling frequency of archival material across space) and temporal resolution (the highest possible sampling frequency of archival material through time). (b) The types of archival materials are plotted in relation to their spatial and temporal limits (the total amount of space or time captured within an archive). Dotted lines indicate the potential range of resolution/limit achieved from each data source. Crosses are centered on the most common resolution/limit achieved by sampling efforts on the data sources in Amazonia.

obtained from the same core, however, general trends in fire or vegetation of older to younger within soil cores can often be established (e.g., Hill et al., 2023; McMichael et al., 2012a; McMichael et al., 2012c; Piperno et al., 2021).

### 3. Lake sediments as local-to regional-scale archives of pre-Columbian Indigenous human activity and vegetation change

Like soils, lake sediments also contain microfossils that can be used to document pre-Columbian human activities and the resulting legacies on ecosystems. Lakes sufficiently old for palaeoecological studies are rare across much of Amazonia, limiting the spatial resolution of palaeovegetation reconstructions (Bush and Silman, 2007), although the temporal resolution can sometimes be high (Fig. 4). Unlike soils, lake sediments typically have continuous deposition and thus robust stratigraphic integrity, and age-depth relationships can be derived that place temporal frameworks on human activities and environmental change. Most lake sediment records from Amazonia contain samples analysed at

centennial scale temporal resolution (e.g., every century to several hundred years) (Nascimento et al., 2022). In rare deep lakes with anoxic conditions, however, the sediments may retain sub-decadal stratigraphy, allowing an almost continuous insight into the local dynamics of human activity and forest recovery (Fig. 4) (Åkesson et al., 2021; Bush et al., 2016, 2021a). The temporal limit of lake sediment records varies depending on lake type, local depositional environment and preservation conditions (Fig. 4). Few sites in Amazonia extend back to the Pleistocene (Mayle et al., 2000; Whitney et al., 2011), as most non-riverine lakes were formed (and sedimentation began) in the mid-to late-Holocene periods (ca. 8000-4000 years ago) (e.g. Bush and McMichael, 2016; Bush et al., 2007; Carson et al., 2014; Nascimento et al., 2022; Urrego et al., 2013).

Charcoal abundances and their changes within a sedimentary sequence are typically used to infer changes in the amount of biomass burning in the surrounding landscape (e.g. Marlon et al., 2013; Marlon et al., 2016); equated to fire severity (Keeley, 2009). Recent work has also estimated burn temperatures from charcoal found within lake sediments, and shown how vegetation composition and traits change depending on the intensity of the fire (Nascimento et al., 2023). Charcoal particles are deposited into lake sediments from airborne or terrestrial sources, and assessing size classes of charcoal particles is commonly used to distinguish local from regional input (Clark and Royall, 1996; Sanford and Horn, 2000). The relationships between charcoal abundance and biomass burned, and the source area of charcoal particles for Amazonian lakes remain poorly documented and need further exploration.

Organic macrofossils and microfossils (e.g., pollen and spores) that decay in soils typically preserve in lake sediments. Fungal spores can be associated with fire and thus human activity (Fig. 1) (Brugger et al., 2016; Loughlin et al., 2018). Tree and shrub pollen can generally be identified to a more specific taxonomic level than phytoliths, although the inverse is true for herbaceous taxa (especially grasses and sedges) and palm taxa (Piperno, 2006). Over 1000 pollen morphotypes from Amazonia have been identified and catalogued (Bush and Weng, 2007), with up to over 100 pollen types being identified within a single pollen sample (Åkesson et al., 2021). Pollen from domesticated maize (*Zea mays*) (see example in Fig. 5) can be reliably identified because wild *Zea* does not occur in South America, but major crop plants such as manioc (*Manihot esculenta*) and sweet potato (*Ipomoea batatas*) cannot be differentiated from their wild varieties using pollen because the latter are native to South America, and taxonomic distinctions cannot reliably be made (Mayle and Iriarte, 2014; Whitney et al., 2012).

The spatial representation of pollen and phytolith data depends on site-specific characteristics of the lake. Phytoliths from lake sediment cores have varying source areas that depend on lake size and the presence of in-flowing streams; the assemblages are often mixtures of these source areas (Carson et al., 2014; Mayle and Iriarte, 2014; Piperno, 2006; Plumpton et al., 2020; Whitney et al., 2013, 2014). Pollen assemblages also reflect a combination of local and regional inputs, and this varies, not only depending upon lake size and the presence or absence of inflowing streams, but also on the relative proportion of wind-pollinated taxa in the parent vegetation (Bush et al., 2021b; Jacobson and Bradshaw, 1981). In the forest-savanna ecotone regions of Bolivia, where the dominant forest taxa (e.g. Moraceae) have wind-dispersed pollen, large lakes may have a pollen source area of up to 40 km (Whitney et al., 2019). However, in many areas of Amazonia, closed-canopy forests are dominated by insect-pollinated taxa, where very small lakes register much smaller pollen source areas of potentially only 1–2 km<sup>2</sup> (Blaus et al., 2023). Differentiating between local- and regional-scale human land-use and deforestation is possible, however, with pollen analyses from tight clusters of small lakes (Bush et al., 2007) or pairs of small and large lakes (Carson et al., 2014). It is also important, where possible, to pair lake sediments from small lakes with nearby archaeological sites to provide matching spatial resolution and a continuous temporal framework of past land use (Carson et al., 2014;




Fig. 5. Microfossils found in Amazonian lakes that can indicate human activity or environmental change. The top panel shows a pollen grain of *Zea mays* (corn), and the bottom panel shows a diatom, *Discostella steligera* (Cleve & Grun.) Houk & Klee, which can indicate water turbidity or lake level. (Photos: Majoi de Novaes Nascimento).

Mayle and Iriarte, 2014; Whitney et al., 2014).

Diatoms are siliceous microalgae that are found in water bodies that provide information about environmental or hydrological conditions (e.g., Battarbee, 1986; Benito et al., 2018) (Fig. 1, for example see Fig. 5). Diatoms are commonly assessed in lake sediment reconstructions, and can indicate changes in hydrology (e.g., lake level recorded by changing proportions of deep versus shallow water taxa) or water quality that are related to climate dynamics (Castro et al., 2013; Nascimento et al., 2021) (Fig. 1). Diatoms, however, can also provide information about pre-Columbian human activity in Amazonia (Fig. 1). They have been used to document changes in wetland management (Duncan et al., 2021), and nutrient status and productivity (Bush et al., 2016). Diatom assemblages can also be used to parse apart climatic and human-induced vegetation changes in lake sediment records (e.g., Bush et al., 2000) (Fig. 1).

Lake sediments also contain stable isotopes and chemical elements that can shed light on pre-Columbian human activity (Fig. 1) (Hodell et al., 1995, 2005). For example, Ca<sup>++</sup> and K<sup>+</sup> concentrations were used to provide information on lake level changes related to climatic fluctuations alongside human activity (Bush et al., 2000; Sahoo et al., 2019). More recently, micro-X-ray fluorescence (XRF) has become a standard tool in paleolimnology, including multivariate analysis of XRF data (Parsons et al., 2018) or ratios of cation concentrations, such as Ca/Ti (proxy for drought), Fe/Mn (proxy for lake depth) or Rb/Sr (proxy for grain size) (Davies et al., 2015). XRF data have been used to reconstruct human-induced soil runoff and erosion (Åkesson et al., 2019), and to place human activities in a context of environmental

change (Aniceto et al., 2014; Maezumi et al., 2018b; Rodríguez-Zorro et al., 2015). Sediment color, which reflects abundances of organic material and clays, can also provide information on environmental rhythms or human-induced change (Bush et al., 2000, 2017; Rodbell et al., 1999).

#### 4. Ethnographic, ethnohistorical and ethnoecological data provide insights into pre-Columbian Indigenous human activity and vegetation change

Ethnographic, ethnohistorical and ethnoecological studies can be used to assess, document, interpret and obtain insights from Indigenous and local knowledge systems (for example see Fig. 6). Indigenous and local knowledge systems provide valuable information that can be used to interpret pre-Columbian human activities (Fig. 1) (Cassino et al., 2019), such as: (i) resource use and management, including past and modern distributions of plant resources (Cassino et al., 2019; Levis et al., 2018, 2020); (ii) how lifestyles and resource management systems influence, and are influenced by landscapes (Balée, 2006); (iii) the technological and labour constraints on resource use (Junqueira et al., 2016); and (iv) the population densities that can be sustained in different ecosystems and by different production systems (Heckenberger et al., 2008).

Ethnography refers to the in-depth description of everyday life and



Fig. 6. Examples of ethnographic studies and working with Indigenous and local people in Amazonia. Top panel shows researcher Carolina Levis conducting an ethnoecological study with a local resident of the Tapajós National Forest of Brazil (Photo taken in 2014 by Bernardo Flores). Bottom panel shows researchers Paul Colinvaux (right), Paulo de Oliveira (second from right) and Melanie Reidinger (left) exchanging knowledge of microfossils and Amazonian plants with members of the Siona ethnic group (Photo taken in 1988 by Mark Bush).

practice of a given culture or society [Oxford (2016); see also Hammersley and Atkinson (2007) for a broader definition], and ethnohistory combines ethnography with the scrutiny of historical records and other sources of information (Axtell, 1979). Ethnoecology is the study of people's interactions with their environment, often with special attention to current knowledge and practices concerning resource use and management, including the subdisciplines of ethnobotany and ethnozoology (Martin, 2004). The subdiscipline of ethnoarchaeology involves ethnographic studies performed by archaeologists with an explicit focus on material culture (Politis, 2014). Much of the culture and resource management practices of the current inhabitants of Amazonia have been inherited from pre-Columbian populations, but transformed to different extents because of the heterogeneous and discontinuous history of human occupation in the region (Cleary, 2001; Denevan, 2001). For example, European arrival caused a massive die-off of Indigenous populations (Denevan, 1992), and many of the remaining groups were fragmented, displaced, or enslaved during European colonization (Dobyns, 1966). Thus, the projection of ethnographic data to past lifestyles and production systems requires caution (McClenachan et al., 2015). Ethnographic data cannot always be tied to human activity at specific times in the past, but local Indigenous knowledge is crucial to locating ancestral forests, anthropogenic soils and other signs of human activity, especially in remote regions (Franco-Moraes et al., 2019; Kopenawa and Albert, 2023).

Indigenous Peoples are the descendants of native ethnic groups that retain historical and cultural connections with pre-Columbian Indigenous societies, though major social disruptions and upheavals occurred with European colonizations (Cook, 1998; Livi-Bacci, 2016). Most ethnographic and ethnoecological studies were done by non-Indigenous scientists. Recent collaborative studies with contemporary Indigenous people have posited that Amazonian ecosystems have been transformed by an extensive and long-term network of social relations and interconnections between humans (e.g., Heckenberger et al., 2008; Ribeiro et al., 2023). Future research should promote participatory and collaborative approaches, as Indigenous people and their knowledge systems can inspire new ideas to enhance understanding of human activities and vegetation changes (for example see Fig. 6).

Archaeological evidence combined with observation of Indigenous production systems suggests, for example, that pre-Columbian production systems in some regions seem to have been much more based on agroforestry and on the management of forest and aquatic resources compared with colonial systems (Maezumi et al., 2018a; Moraes, 2015; Neves, 2013; Shepard et al., 2020). Ethnographic work on current soil and waste management systems has also been essential to our understanding of the processes that led to the formation of Amazonian Dark Earth (ADE) (Hecht, 2003; Schmidt et al., 2014; Winklerprins, 2009).

The variance and scarcity of ethnographic data limits extrapolations of locally derived information to other regions within Amazonia. For example, while major crops like maize and manioc were grown in most regions of the basin, some minor crops, or useful native species, that are culturally or economically important in a certain region may not be so in others. Instead of a direct projection of the present into the past, ethnographical research provides *insights* to further understand ancient Indigenous livelihoods, resource management strategies and their potential impacts in past and current landscapes (McClenachan et al., 2015). Ethnographic studies also provide valuable information on Indigenous resource use and societal practice since European colonization, and facilitate disentangling the effects of pre- and post-Colonial human activities in current landscapes (Forline, 2008). Ethnographic data can be paired with linguistics, as it is known that groups within the same language families are more likely to share similar resource management systems, e.g., the Arawak (Eriksen and Danielsen, 2014). Historic distributions of Indigenous languages (e.g., Eriksen, 2011), may thus facilitate ethnographic projections across space since European colonization.

Historical documents originating during the early colonization of

Amazonia may also provide insight into pre-Columbian Indigenous land use systems. Francisco Orellana led the first expedition down the Amazon River in 1540 CE, which was recorded by Gaspar de Carvajal (Medina, 1934). Early colonists moved in and established Jesuit missions later in the 1600s CE in several regions (Reeve, 1994), and the Amazonian Rubber Boom occurred from ca. 1850–1920 CE (Hecht and Cockburn, 2010; Weinstein, 1983). All these events had corresponding historical documents that recorded aspects of Indigenous Peoples and their interactions with the colonists.

The spatial resolution of ethnographic records and historical data has the potential to be very high (Fig. 4). There is also a high potential temporal resolution of both historical and ethnographic records, though available data through time are relatively sparse. The temporal limit of historical documents encompasses only the last several hundred years since European arrival, and the temporal limit of ethnographic records is bound by the memories of local residents, although it can extend much further back in time through knowledge transmission across generations (Fig. 4b). Because of the fragmentation and upheaval of Indigenous populations after European colonization, it remains contentious as to how far back in time ethnography can be extrapolated back. In some cases there is a strong cultural and historical continuity between ancient and contemporary Indigenous Peoples, but in other cases contemporary Indigenous Peoples have inhabited their current lands for only several decades (e.g., the Kichwa People of Andean descent that now inhabit areas of lowland Ecuador).

##### 5. Modern datasets provide insight to pre-Columbian Indigenous human activities and vegetation change: biological collections and remote sensing data

Biological collection records include plant and animal surveys (inventories), biological or ecological monitoring networks (e.g., Anderson-Teixeira et al., 2015; Malhi et al., 2002; ter Steege et al., 2013), and herbarium or museum specimens (e.g., Feeley and Silman, 2011) (for example see Fig. 7). These biological collection records are used to assess the presence, absence, and abundances of plants and animals in modern ecosystems (or those during the historic period). Although biological records provide valuable information about past (pre- and post-colonial) activities, they do not directly measure pre-Columbian Indigenous influences on ecosystems (Fig. 1). Biological records need linkages with archaeological, paleoecological, genetic, or ethnographic data on the degree and form of past human activities to infer cause and effect relationships (e.g., Heijink et al., 2022; Heijink et al., 2020; Levis et al.,



Fig. 7. Researcher Carolina Levis measures and identifies a piquiá tree (*Carycar villosum*) with a local resident of the Tapajós National Forest of Brazil (Photo: Bernardo Flores).

2017; Piperno et al., 2021).

Past and current human activities are increasingly recognized as factors influencing species' distributions (Boivin et al., 2016; Di Marco and Santini, 2015; Gallardo et al., 2015; Guisan and Thuiller, 2005; Halpern et al., 2008). Humans have modified the distribution range and abundance of several plant species, expanding the distribution of useful and domesticated plants more often than plants without a documented use to humans (Balée, 1989; Levis et al., 2017; Coelho et al., 2021). For instance, manioc (*Manihot esculenta*) was cultivated by Indigenous people throughout the Holocene (Piperno, 2011). Currently, domesticated manioc populations are cultivated throughout the tropics, while the direct ancestor of domesticated manioc (*M. flabellifolia*) is limited to South America (Olsen and Schaal, 1999). Differences between the natural distribution and the human-modified distribution of cultivated species, and especially those with domesticated populations can indicate past human activities. Several palms and trees that are used for food are abundant in plant assemblages of archaeological sites (Balée, 1989; Junqueira et al., 2010). Modern plant inventories that are closer to archaeological sites also tend to have a higher abundance and diversity of useful and domesticated plants (Levis et al., 2012, 2017; Thomas et al., 2015).

Plant genetic material (DNA) is typically derived from plants but can also be found in soils and sediments. Genetic material from plant remains can be linked with past events in human history, such as plant domestication and migrations (Fig. 1) (e.g. Clement, 1988a; Clement, 1988b; Gutaker and Burbano, 2017; Moreira et al., 2017; Roullier et al., 2013). Genetic studies can also provide information about species with populations that were domesticated by humans, because selection and cultivation of desirable phenotypes results in changes in morphology, physiology, and genotype of descendent populations (Emshwiller, 2006; Harlan, 1992; Olsen and Schaal, 2001). The whole set of selected phenotypic changes in a species is termed its domestication syndrome, which can be studied with morphological, chemical, archaeobotanical, and molecular genetic methods (e.g. Emshwiller, 2006; Meyer et al., 2012; Smith, 2006).

Species with populations with some degree of domestication show patterns of morphological variation and genetic diversity and structure across geographical space that result from domestication events, dispersal and subsequent diversification (Meyer and Purugganan, 2013). Economically important domesticates are more likely to show dramatic morphological changes, such as a 2000% increase in fruit size from wild source populations of peach palm (*Bactris gasipaes*) compared with some domesticated populations (Clement, 1988b). Dispersal events are often accompanied by other natural and human selection pressures, resulting in diversification of uses, variation in morphology, chemical composition and physiology (Meyer and Purugganan, 2013), and adaptation to domesticated landscapes (Clement, 1999).

Until very recently, plant geneticists worked exclusively with living plants or samples collected over the last 200–300 years (e.g., Roullier et al., 2013). Over the last decade, new molecular genetic methods have allowed the extraction and analysis of DNA from archaeobotanical remains (Wales et al., 2014). Ancient DNA (aDNA) is increasing the ability to document the genetic history of plants, can differentiate crops from their wild ancestors, or estimate genetic change and migration of domesticated plants or cultivars over time (Freitas et al., 2003; McLachlan and Clark, 2005; Piperno, 2011). It is now even possible to extract and date aDNA recovered from pollen grains found in lake sediments and historical samples (Gutaker and Burbano, 2017; Parducci et al., 2017). aDNA is denatured very rapidly in tropical lake sediments (due to the hot, wet conditions), though extraction has been successful in the African tropics (Bremond et al., 2017). Geographic representation of aDNA samples is patchy, although many crop plants have been databased.

Biological collection records can be collected from anywhere on Earth, so their potential spatial limit is endless (Fig. 4). To date, however, sampling covers only a small portion of the 6 million km<sup>2</sup> of

Amazonia (Carvalho et al., 2023; ter Steege et al., 2013). Ground based surveys of plant and animal distributions and abundances can also have high spatial resolution, with hierarchies of transects or 1-ha plots commonly clustered within a region (e.g., ter Steege et al., 2013). Biological inventories can also be measured at yearly frequencies, and have high temporal resolution, but most repeated censuses only span a few decades (Malhi et al., 2002; Phillips et al., 1994) (Fig. 4).

Ground-based biological collections and satellite imagery are often paired in modern ecological and global change studies. Satellite imagery has also been used to infer soil and vegetation legacies of pre-Columbian land use (Iriarte, 2016; Palace et al., 2017; Thayn et al., 2011). Satellite data from Landsat, Sentinel, and MODIS, typically capture landscape features at 30-m to 1-km spatial resolution (Fig. 4). Landsat has been used to detect legacies of pre-Columbian land use (Heckenberger et al., 2003; Söderström et al., 2016), and MODIS has been used to detect or predict Amazonian Dark Earth (ADE) (Palace et al., 2017; Thayn et al., 2011). This detection is possible because pre-Columbian Indigenous Peoples permanently changed the soil characteristics, which affects the types of vegetation that can grow on those soils (Junqueira et al., 2011). The differences in modern biomass between ADE and forested non-ADE sites within the same region is also detectable using MODIS satellite imagery (Palace et al., 2017).

Remotely sensed data has low temporal limits (Fig. 4). Satellite images are only available for the last few decades, although aerial images may extend further back in time. However, they have higher spatial limits than biological collections, and usually have Amazonian-wide spatial coverage (Fig. 4).

## 6. Dating data sources and proxies used in assessing pre-Columbian Indigenous human activity and vegetation change

Different scientific disciplines assessing pre-Columbian Indigenous human activities place differing emphases on dating or age control of their data sources or proxies. In soils, archaeologists and paleoecologists typically use  $^{14}\text{C}$  AMS dating (radiocarbon dating) to obtain ages on specific material(s) of interest. In archaeological surveys, multiple ages are usually derived from specific horizons of interest where artifacts are recovered (e.g., Roosevelt et al., 1996). Sometimes, however, ages of specific horizons are inferred from a known type of pottery or artifact that has been recovered and dated from another location (e.g., McEwan, 2001). Paleoecologists will typically obtain dates from individual charcoal fragments or conglomerations of phytoliths recovered from soil cores or profiles (e.g., Heijink et al., 2022; McMichael et al., 2012a; Piperno et al., 2021; Watling et al., 2017). In both archaeological and paleoecological surveys in soils, repeated dating across sites can help determine the synchronicity of events across space. Because of soil bioturbation, age-depth models are not applied to soil profiles. General stratigraphic trends, however, are often intact (Piperno et al., 2021; Watling et al., 2017).

Lake sediments typically retain stratigraphic integrity, and age-depth models (e.g., Blaauw and Christen, 2011) are often used to reconstruct a temporally continuous sedimentary sequence. The age-depth models typically require fewer  $^{14}\text{C}$  dates than may be obtained with soil surveys, but the more dates obtained on a sequence, the more confidence can be placed in the model. With lake sediments, specific layers of change or markers of human activity can be directly dated to provide additional confidence for the timing of those events. Additional temporal control can also be placed on the younger sections of lake sediment cores using  $^{210}\text{Pb}$  dating, which can be used on sediments less than 150 years old (e.g., Sanchez-Cabeza and Ruiz-Fernández, 2012). Age-depth models can incorporate mixtures of  $^{210}\text{Pb}$  and  $^{14}\text{C}$  dates (Aquino-López et al., 2018), which can be particularly useful for increasing confidence in the ages of sediments around the period of European Contact.

The other data sources included in this review are not based on laboratory dating of materials or sediments. Historical records have specific ages corresponding to the dates they were produced. It is not

possible to place specific dates or ages on Indigenous knowledge, or its interpretation through the studies of ethnography, ethnohistory, or ethnoecology. It is possible, however, to correlate some historical events or personal events, such as the arrival of missionaries in an Indigenous community or abandonment of a village with changes in resource use and management. This information is crucial to understanding how pre-Columbian people lived, even though specific ages are difficult to obtain.

Modern datasets, including living plants and museum specimens, may have specific dates recorded. Inferring an age when pre-Columbian Indigenous people shaped plant abundances or plant growth patterns (for instance) is possible with the direct dating of living materials using a combination of dendrochronology, radiocarbon analysis, stable isotope analysis and DNA analysis (Caetano-Andrade et al., 2020). Recent advances in genetic methods have ages inferred also from molecular clocks and DNA-based dating method for ancient genomes (Kistler et al., 2020).

## 7. Discussion: moving forward

Understanding the interactions of pre-Columbian people and their environments in Amazonia is important for sustainability science, conservation biology and cultural anthropology (Levis et al., 2017; Mayle and Iriarte, 2014; McMichael et al., 2017; Roberts et al., 2017; Szabó and Hédli, 2011; Watling et al., 2017; WinklerPrins and Levis, 2020). Here we have provided a review of the most commonly used sources and proxies for reconstructing pre-Columbian human activity in Amazonia (Fig. 1) and have described the associated advantages and limitations of each by assessing their potential spatial and temporal resolution and limits (Fig. 4). We acknowledge that a comprehensive assessment of all proxies of Indigenous human activity is beyond the scope of this manuscript. For example, proxies such as lipid biomarkers have been used to assess the components of anthropogenic soils (Glaser, 2007). Fecal biomarkers are a newly emerging proxy that is being used to detect past human activity (Argiriadis et al., 2018; Zocattelli et al., 2017). These proxies have not yet been used in Amazonian systems, though show great promise in other geographic regions. The preservation of biomarkers and sterols in the humid tropics, however, is likely poorer than in other areas.

Assessments of the long-term Indigenous history in Amazonia would be stronger if 'absence data' from all lines of evidence were reported or archived in data repositories. Ecological datasets, including plant inventory records, include the presence, *absence*, and abundance of species within a given study area (e.g. Hubbell, 1979; ter Steege et al., 2013), allowing for more robust statistical analyses than presence-only analysis often applied to archaeological datasets (e.g., McMichael et al., 2014a; McMichael et al., 2017). To fully understand the impact of pre-Columbian people in Amazonian landscapes, the reporting of 'absence data' (i.e., when there is no evidence of past human activity) is crucial. For example, when identifying earthworks using remotely sensed data, the total area surveyed and total number of earthworks found should be reported so that site densities can be calculated and compared across regions. The entire area sampled and information on the *absence* of ADEs using field-based surveys is also rarely reported, and the varying densities of ADEs across the landscape cannot yet be calculated. The same approach should apply for archaeological surveys, in which generally a wide area is initially surveyed before determining excavation locations. We suggest that efforts to compile and build repositories of archaeological information should develop guidelines and protocols for reporting and documenting absence data.

The 'absence' of evidence of human activities from paleoecological proxies, including charcoal, pollen, and phytoliths is reported. However, because these lines of evidence come from a limited amount of sample material, it is possible that they are present but not detected. Thus, the absence of evidence is not necessarily evidence of absence. This issue is partially overcome with repeated sampling in the case of soil cores (i.e., multiple cores collected and analysed per locality) (e.g., Heijink et al.,



2022; McMichael et al., 2015; McMichael et al., 2012a; Piperno et al., 2021), and with continuous sampling and the analyses of multiple cores in a region in the case of lake sediments (e.g., Bush et al., 2007; Carson et al., 2014).

Several predictive models have been developed for various types of archaeological features in Amazonia and for the overall likely distribution of pre-Columbian Indigenous Peoples in the region, which provide targets for future archaeological surveys in the ca. 6 million km<sup>2</sup> of Amazonian forests (McMichael et al., 2014a, 2014b, 2017; Souza et al., 2018; Walker et al., 2023). Due to the lack of 'absence data', the only approaches available for these predictions are models that require presence-only data (i.e., that do not require absence data). Absence data of all types would help to validate and refine these models on both regional and continental-wide scales, and open doors to an array of additional modelling approaches that could be employed (McMichael et al., 2017). All types of models and other macro-paleoecological and macro-archaeological syntheses would also further benefit by including more precision in the geographic coordinates of localities being studied (i.e., to ca. ± 200 m spatial resolution), and more precision in the time bins of analysis.

Perhaps the best way to strengthen assessments of past human activity is via an interdisciplinary approach, pairing multiple proxies and multiple types of data sources (Carson et al., 2014; Mayle and Iriarte, 2014; Watling et al., 2017; Whitney et al., 2014). Without this pairing, it can be difficult to determine whether the observed pattern is related to human activity or other forcing mechanisms. Ideally, though not necessarily, those pairings should be spatially overlapped. For example, Levis et al. (2012) paired plant distribution information with charcoal recovered from soils beneath them, and found a higher percentage of useful species in plots that contained higher amounts of charcoal in the vicinity of archaeological sites. It is also well documented how phytolith and pollen data complement one another in paleoecological and archaeological reconstructions (Åkesson et al., 2021; Mayle and Iriarte, 2014; Piperno, 2006). Phytoliths tend to be more sensitive to detecting cultivation, forest openings, and some basal angiosperm and eudicotyledon tree taxa, particularly in wet closed canopy forests (as opposed to the savanna ecotone regions), whereas pollen can detect changes in tree taxon abundances that remain undetectable in phytolith analyses (Åkesson et al., 2021; Piperno and McMichael, 2023). Phytoliths identify basal angiosperm and eudicotyledon taxa that pollen does not, and the two are highly complementary when they can be studied together (Piperno and McMichael, 2023). Phytolith analysis has also been paired with starch grain and stable carbon isotope analysis in archaeological settings for a more comprehensive view of the diet and lifestyle of pre-Columbian Indigenous people in Amazonia (Iriarte et al., 2010; McKey et al., 2010). The pairing of proxies, or using multi-proxy approaches, can also aid in detectability of past human influences whereas single proxies may lack detectability of specific lines of evidence.

To move forward, the integration of data needs to occur across proxies, data sources, and consider both spatial and temporal scales (Fig. 4). Either within or between archives, a multi-proxy approach is more sensitive to detecting past human activity than a single-proxy approach, providing more confidence in conclusions on the presence or absence of pre-Columbian Indigenous human activity and the impact that they caused on the landscape (Fig. 1) (Clement et al., 2015; Iriarte, 2016; Mayle and Iriarte, 2014; Piperno, 2006). Recent research has already begun to integrate: (i) lake sediment data with soil survey data from areas within the watershed (McMichael et al., 2012b), (ii) terrestrial soil archives with archaeological sites containing earthworks (Watling et al., 2017), (iii) lake sediment records, even with lakes of different sizes reflecting different source areas, with archaeological sites (Carson et al., 2014; Maezumi et al., 2018b; Whitney et al., 2013), (iv) geospatial patterns of plant distributions with archaeological sites across Amazonia (Levis et al., 2017). As trees in Amazonia can also be directly dated (Brienen and Zuidema, 2006; Chambers et al., 1998; Schöngart et al., 2015), pairing the age of the modern forest with archaeological,

paleoecological and historical data could also prove invaluable. In a recent example, Caetano Andrade et al. (2019) integrated dendrochronology and historical data to evaluate Indigenous and traditional management of a Brazil nut (*Bertholletia excelsa*) stand near an archaeological site south of Manaus.

If the advantages and limitations of data used to infer pre-Columbian Indigenous human activity are recognized and acknowledged, particularly regarding spatial and temporal scale (Fig. 4), then disagreement among existing and future datasets may be minimized. We also highlight the potential and importance of bridging ethnography and ethnoecology with historical records, archaeological data, and paleoecological data. Importantly, but often not considered, the voices and knowledge of Indigenous Peoples should also be integrated into scientific research designs (Trisos et al., 2021). Together, these recommendations can advance the understanding of the complexity and variation of pre-Columbian Indigenous human influences in tropical ecosystems, such as Amazonia.

#### Author contributions

All authors conceived the ideas.

CNHM led the writing, with input from all coauthors.

#### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Data availability

No data were used for the research described in the article.

#### Acknowledgements

CNHM is funded by a European Research Council Starting Grant (ERC-StG 853394) and the Netherlands Organisation for Scientific Research (NWO) award ALWOP.322. CL thanks the Brazilian National Research Council (CNPq) for a doctoral research scholarships (proc. No. 141652/2014–4 and 201573/2014–8) and CNPq and Capes for a post-doctoral fellowships (proc. No. 159440/2018–1, 88887.474568/2020 and 400369/2021–4). ABJ received funding from *Terra Preta Program* funded by the Interdisciplinary Research and Education Fund (INREF) of Wageningen University, from the Netherlands Organisation for INTERNATIONAL COOPERATION in Higher Education (NUFFIC), and from the European Research Council under an ERC Consolidator Grant (FP7-771056-LICCI). DRP thanks the Smithsonian Institution. We would all like to thank Bernardo Oliveira, Bernardo Flores, Majoi de Novaes Nascimento, and Mark Bush for photo contributions.

#### References

- Åkesson, C.M., Matthews-Bird, F., Bitting, M., Fennell, C.-J., Church, W.B., Peterson, L.C., Valencia, B.G., Bush, M.B., 2019. 2,100 years of human adaptation to climate change in the High Andes. *Nat. Ecol. & Evol.* 1–9.
- Åkesson, C.M., McMichael, C.N., Raczka, M.F., Huisman, S.N., Palmeira, M., Vogel, J., Neill, D., Veizaj, J., Bush, M.B., 2021. Long-term ecological legacies in western Amazonia. *J. Ecol.* 109, 432–446.
- Alencar, A.A.C., Solorzano, L.A., Nepstad, D.C., 2004. Modeling forest understory fires in an eastern Amazonian landscape. *Ecol. Appl.* 14, 139–149.
- Anderson-Teixeira, K.J., Davies, S.J., Bennett, A.C., Gonzalez-Akre, E.B., Muller-Landau, H.C., Joseph Wright, S., Abu Salim, K., Almeyda Zambrano, A.M., Alonso, A., Baltzer, J.L., 2015. CTPS-Forest GEO: a worldwide network monitoring forests in an era of global change. *Global Change Biol.* 21, 528–549.
- Aniceto, LakeK., Moreira-Turcq, P., Cordeiro, R.C., Quintana, I., Fraizy, P., 2014. Hydrological Changes in West Amazonia over the Past 6 Ka Inferred from Geochemical Proxies in the Sediment Record of a Floodplain. *TurcqProcedia Earth and Planetary Science* 10, 287–291DOI. <https://doi.org/10.1016/j.proeps.2014.08.065>.

- Aquino-López, M.A., Blaauw, M., Christen, J.A., Sanderson, N.K., 2018. Bayesian analysis of 210 Pb dating. *J. Agric. Biol. Environ. Stat.* 23, 317–333.
- Argiriadis, E., Battistell, D., McWethy, D.B., Vecchiato, M., Kirchgörg, T., Kehrwald, N. M., Whitlock, C., Wilmshurst, J.M., Barbante, C., 2018. Lake sediment fecal and biomass burning biomarkers provide direct evidence for prehistoric human-lit fires in New Zealand. *Sci. Rep.* 8, 12113.
- Arroyo-Kalin, M., Riris, P., 2021. Did pre-Columbian populations of the Amazonian biome reach carrying capacity during the Late Holocene? *Phil. Transac. Royal Soc. B* 376, 20190715.
- Axtell, J., 1979. Ethnohistory: an historian's viewpoint. *Ethnohistory* 26, 1–13.
- Bachelet, C., Scheel-Ybert, R., 2017. Landscape and firewood selection in the santa elina rock shelter (mato grosso, Brazil) during the holocene. *Quat. Int.* 431, 52–60.
- Balée, W., 1989. The culture of Amazonian forests. In: Posey, D.A., Balée, W. (Eds.), *Resource Management in Amazonia: Indigenous and Folk Strategies*. New York Botanical Garden, New York, pp. 1–21.
- Balée, W., 2006. The research Program of historical ecology. *Annu. Rev. Anthropol.* 35, 75–98.
- Balée, W., 2010. Contingent diversity on anthropic landscapes. *Diversity* 2, 163–181.
- Barlow, J., Gardner, T.A., Lees, A.C., Parry, L., Peres, C.A., 2012. How pristine are tropical forests? An ecological perspective on the pre-Columbian human footprint in Amazonia and implications for contemporary conservation. *Biol. Conserv.* 151, 45–49.
- Barnosky, A.D., Lindsey, E.L., 2010. Timing of Quaternary megafaunal extinction in South America in relation to human arrival and climate change. *Quat. Int.* 217, 10–29.
- Battarbee, R.W., 1986. *Handbook of Holocene Palaeoecology and Palaeohydrology*. John Wiley and Sons, New York.
- Benito, X., Fritz, S.C., Steinitz-Kannan, M., Tapia, P.M., Kelly, M.A., Lowell, T.V., 2018. Geo-climatic factors drive diatom community distribution in tropical South American freshwaters. *J. Ecol.* 106, 1660–1672.
- Blaauw, M., Christen, J.A., 2011. Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Anal.* 6, 457–474.
- Blaus, A., McMichael, C., Raczka, M., Herrick, C., Palace, M., Witteveen, N., Nascimento, M., Bush, M., 2023. Amazonian pollen assemblages reflect biogeographic gradients and forest cover. *J. Biogeogr.* Volume 50 Number 11, pages 1926–1938.
- Bodin, S.C., Molino, J.-F., Odonne, G., Bremond, L., 2020. Unraveling pre-Columbian occupation patterns in the tropical forests of French Guiana using an anthracological approach. *Veg. Hist. Archaeobotany* 29, 567–580.
- Bodin, S.C., Scheel-Ybert, R., Beauchêne, J., Molino, J.-F., Bremond, L., 2019. CharKey: an electronic identification key for wood charcoals of French Guiana. *IAWA J.* 40, 75–82.
- Boivin, N.L., Zeder, M.A., Fuller, D.Q., Crowther, A., Larson, G., Erlandson, J.M., Denham, T., Petraglia, M.D., 2016. Ecological consequences of human niche construction: examining long-term anthropogenic shaping of global species distributions. *Proc. Natl. Acad. Sci. USA* 113, 6388–6396.
- Bremond, L., Favier, C., Ficotola, G.F., Tossou, M.G., Akouégninou, A., Gielly, L., Giguet-Covex, C., Oslisly, R., Salzmann, U., 2017. Five thousand years of tropical lake sediment DNA records from Benin. *Quat. Sci. Rev.* 170, 203–211.
- Brienen, R.J.W., Zuidema, P.A., 2006. The use of tree rings in tropical forest management: projecting timber yields of four Bolivian tree species. *For. Ecol. Manag.* 226, 256–267.
- Brugger, S.O., Gobet, E., van Leeuwen, J.F., Ledru, M.-P., Colombaroli, D., van der Knaap, W., Lombardo, U., Escobar-Torrez, K., Finsinger, W., Rodrigues, L., 2016. Long-term man–environment interactions in the Bolivian Amazon: 8000 years of vegetation dynamics. *Quat. Sci. Rev.* 132, 114–128.
- Bush, M., Correa-Metrio, A., McMichael, C., Sully, S., Shadik, C., Valencia, B., Guilderson, T., Steinitz-Kannan, M., Overpeck, J., 2016. A 6900-year history of landscape modification by humans in lowland Amazonia. *Quat. Sci. Rev.* 141, 52–64.
- Bush, M., Nascimento, M., Åkesson, C., Cárdenes-Sandí, G., Maezumí, S., Behling, H., Correa-Metrio, A., Church, W., Huisman, S., Kelly, T., Mayle, F., McMichael, C.N.H., 2021a. Widespread reforestation before European influence on Amazonia. *Science* 372, 484–487.
- Bush, M.B., Correa-Metrio, A., van Woesik, R., Collins, A., Hanselman, J., Martinez, P., McMichael, C.N., 2021b. Modern pollen assemblages of the Neotropics. *J. Biogeogr.* 48, 231–241.
- Bush, M.B., Correa-Metrio, A., Woesik, R., Shadik, C.R., McMichael, C.N., 2017. Human disturbance amplifies amazonian el niño–southern oscillation signal. *Global Change Biol.* 23, 3181–3192.
- Bush, M.B., McMichael, C.H., Piperno, D.R., Silman, M.R., Barlow, J.B., Peres, C.A., Power, M.J., Palace, M.W., 2015. Anthropogenic influence on Amazonian forests in prehistory: an ecological perspective. *J. Biogeogr.* 42, 2277–2288.
- Bush, M.B., McMichael, C.N., 2016. Holocene variability of an Amazonian hyperdominant. *J. Ecol.* 104, 1370–1378.
- Bush, M.B., Miller, M.C., De Oliveira, P.E., Colinvaux, P.A., 2000. Two histories of environmental change and human disturbance in eastern lowland Amazonia. *Holocene* 10, 543–554.
- Bush, M.B., Silman, M.R., 2007. Amazonian exploitation revisited: ecological asymmetry and the policy pendulum. *Front. Ecol. Environ.* 5, 457–465.
- Bush, M.B., Silman, M.R., de Toledo, M.B., Listopad, C.R.S., Gosling, W.D., Williams, C., de Oliveira, P.E., Krisel, C., 2007. Holocene fire and occupation in Amazonia: records from two lake districts. *Phil. Transac. Royal Soc. London. Ser. B* 362, 209–218.
- Bush, M.B., Silman, M.R., McMichael, C., Saatchi, S., 2008. Fire, climate change and biodiversity in Amazonia: a Late-Holocene perspective. *Phil. Trans. Biol. Sci.* 363, 1795–1802.
- Bush, M.B., Weng, C., 2007. Introducing a new (freeware) tool for palynology. *J. Biogeogr.* 34, 377–380.
- Caetano Andrade, V.L., Flores, B.M., Levis, C., Clement, C.R., Roberts, P., Schöngart, J., 2019. Growth rings of Brazil nut trees (*Bertholletia excelsa*) as a living record of historical human disturbance in Central Amazonia. *PLoS One* 14, e0214128.
- Caetano-Andrade, V.L., Clement, C.R., Weigel, D., Trumbore, S., Boivin, N., Schöngart, J., Roberts, P., 2020. Tropical trees as time capsules of anthropogenic activity. *Trends Plant Sci.* 25, 369–380.
- Carson, J.F., Whitney, B.S., Mayle, F.E., Iriarte, J., Prümers, H., Soto, J.D., Watling, J., 2014. Environmental impact of geometric earthwork construction in pre-Columbian Amazonia. *Proc. Natl. Acad. Sci. USA* 111, 10497–10502.
- Cartwright, C.R., 2015. The principles, procedures and pitfalls in identifying archaeological and historical wood samples. *Ann. Bot.* 116, 1–13.
- Carvalho, R.L., Resende, A.F., Barlow, J., França, F.M., Moura, M.R., Maciel, R., Alves-Martins, F., Shutt, J., Nunes, C.A., Elias, F., 2023. Pervasive gaps in Amazonian ecological research. *Curr. Biol.* 33, 3495–3504.
- Cassino, M.F., Alves, R.P., Levis, C., Watling, J., Junqueira, A.B., Shock, M.P., Ferreira, M.J., Caetano Andrade, V.L., Furquim, L.P., Coelho, S.D., 2019. Ethnobotany and Ethnoecology Applied to Historical Ecology, Methods and Techniques in Ethnobiology and Ethnoecology. Springer, pp. 187–208.
- Castro, D.F., Oliveira, P.E., Rossetti, D.F., Pessenda, L.C., 2013. Late Quaternary landscape evolution of northeastern Amazonia from pollen and diatom records. *An Acad. Bras Ciências* 85, 35–55.
- Chambers, J.Q., Higuchi, N., Schimel, J.P., 1998. Ancient trees in Amazonia. *Nature* 391, 135–136.
- Clark, J.S., Royall, P.D., 1996. Local and regional sediment charcoal evidence for fire regimes in presettlement north-eastern America. *J. Ecol.* 84, 365–382.
- Cleary, D., 2001. Towards an environmental history of the Amazon: from prehistory to the nineteenth century. *Lat. Am. Res. Rev.* 65–96.
- Clement, C.R., 1988a. Domestication of the peibaybe palm (*Bactris gasipaes*): past and present. *Adv. Econ. Bot.* 6, 155–174.
- Clement, C.R., 1988b. Domestication of the peibaybe palm (*Bactris gasipaes*): past and present. In: Balick, M.J. (Ed.), *The Palm - Tree of Life. Biology, Utilization and Conservation*. The New York Botanical Garden, New York, pp. 155–174.
- Clement, C.R., 1999. 1492 and the loss of Amazonian crop genetic resources. I. The relation between domestication and human population decline. *Econ. Bot.* 53, 188–202.
- Clement, C.R., Denevan, W.M., Heckenberger, M.J., Junqueira, A.B., Neves, E.G., Teixeira, W.G., Woods, W.L., 2015. The domestication of Amazonia before European conquest. *Proc. Royal Soc. B* 282, 20150813.
- Coelho, S.D., Levis, C., Baccaro, F.B., Figueiredo, F.O., Pinassi Antunes, A., Ter Steege, H., et al., 2021. Eighty-four per cent of all Amazonian arboreal plant individuals are useful to humans. *PLoS One* 16 (Issue 10), e0257875.
- Colone, A.C., Winter, R., Brandi, R., Fossile, T., Fernandes, R., Soncin, S., McGrath, K., Von Tersch, M., Bandeira, A.M., 2020. Stable isotope evidence for dietary diversification in the pre-Columbian Amazon. *Sci. Rep.* 10, 16560.
- Cook, N.D., 1998. *Born to die: disease and New World conquest*. Cambridge University Press, pp. 1492–1650.
- Crutzen, P.J., 2002. The “anthropocene”. In: *Journal de Physique IV (Proceedings)*. EDP sciences, pp. 1–5.
- Davies, S.J., Lamb, H.F., Roberts, S.J., 2015. Micro-XRF core scanning in palaeolimnology: recent developments. *Micro-XRF Stud. Sediment Cores: Appl. Non-destruct. Tool Environ. Sci.* 189–226.
- de Freitas, H.A., Pessenda, L.C.R., Aravena, R., Gouveia, S.E.M., Ribeiro, A.S., Boulet, R., 2001. Late quaternary vegetation dynamics in the southern Amazon basin inferred from carbon isotopes in soil organic matter. *Quat. Res.* 55, 39–46.
- Denevan Ann, W.M., 1992. The Pristine Myth - the Landscape of the America in 1492. *Assoc. Am. Geograph.* Vol. 82 (Issue 3), 369–385.
- Denevan, W.M., 2001. *Cultivated Landscapes of Native Amazonia and the Andes*. Oxford University Press, Oxford.
- Di Marco, M., Santini, L., 2015. Human pressures predict species' geographic range size better than biological traits. *Global Change Biol.* 21, 2169–2178.
- Di Pasquale, G., Marziano, M., Impagliazzo, S., Lubritto, C., De Natale, A., Bader, M.Y., 2008. The Holocene treeline in the northern Andes (Ecuador): first evidence from soil charcoal. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 259, 17–34.
- Dickau, R., Ranere, A.J., Cooke, R.G., 2007. Starch grain evidence for the preceramic dispersals of maize and root crops into tropical dry and humid forests of Panama. *Proc. Natl. Acad. Sci. USA* 104, 3651–3656.
- Dickau, R., Whitney, B.S., Iriarte, J., Mayle, F.E., Soto, J.D., Metcalfe, P., Street-Perrott, F.A., Loader, N.J., Ficken, K.J., Killeen, T.J., 2013. Differentiation of neotropical ecosystems by modern soil phytolith assemblages and its implications for palaeoenvironmental and archaeological reconstructions. *Rev. Palaeobot. Palynol.* 193, 15–37.
- Dobyns, H.F., 1966. Estimating aboriginal American population I: an appraisal of techniques with a new hemispheric estimate. *Curr. Anthropol.* 7, 395–416.
- Duncan, N.A., Loughlin, N.J., Walker, J.H., Hocking, E.P., Whitney, B.S., 2021. Pre-Columbian fire management and control of climate-driven floodwaters over 3,500 years in southwestern Amazonia. *Proc. Natl. Acad. Sci. USA* 118, e202206118.
- Ellis, E.C., Gauthier, N., Goldewijk, K.K., Bird, R.B., Boivin, N., Díaz, S., Fuller, D.Q., Gill, J.L., Kaplan, J.O., Kingston, N., 2021. People have shaped most of terrestrial nature for at least 12,000 years. *Proc. Natl. Acad. Sci. USA* 118.
- Emshwiller, E., 2006. Genetic data and plant domestication. In: Zeder, M.A., Bradley, D. G., Emshwiller, E., Smith, B.D. (Eds.), *Documenting Domestication: New Genetic and Archaeological Paradigms*. University of California Press, Berkeley, California, USA, pp. 99–122.
- Eriksen, L., 2011. *Nature and Culture in Prehistoric Amazonia: Using GIS to Reconstruct Ancient Ethnogenetic Processes from Archaeology, Linguistics, Geography, and Ethnohistory*. Lund University.

- Eriksen, L., Danielsen, S., 2014. The Arawakan matrix. In: O'Connor, L., Muysken, P. (Eds.), *The Native Languages of South America: Origins, Developments, Typology*. Cambridge University Press, Cambridge, UK, pp. 152–176.
- Ezell, K.C., Pearsall, D.M., Zeidler, J.A., 2006. Root and tuber phytoliths and starch grains document manioc (*Manihot esculenta*) arrowroot (*Maranta arundinacea*) and llerén (*Calathea* sp.) at the real alto site Ecuador. *Econ. Bot.* 60, 103–120.
- Feeley, K.J., Silman, M.R., 2011. Keep collecting: accurate species distribution modelling requires more collections than previously thought. *Divers. Distrib.* 17, 1132–1140.
- Feldpausch, T.R., Carvalho, L., Macario, K.D., Ascough, P.L., Flores, C.F., Coronado, E.N. H., Kalamandeen, M., Phillips, O.L., Staff, R.A., 2022. Forest fire history in Amazonia inferred from intensive soil charcoal sampling and radiocarbon dating. *Front. Forests Glob. Change* 5, 815438.
- Fernandes Caromano, C., Matthews Cascon, L., Góes Neves, E., Scheel-Ybert, R., 2013. Revealing fires and rich diets: macro-and micro-archaeobotanical analysis at the Hatahara Site, Central Amazonia. *Tipití: Journal of the Society for the Anthropology of Lowland South America* 11, 40–51.
- Flores, B.M., Holmgren, M., Xu, C., Van Nes, E.H., Jakovac, C.C., Mesquita, R.C., Scheffer, M., 2017. Floodplains as an Achilles' heel of Amazonian forest resilience. *Proc. Natl. Acad. Sci. USA* 114, 4442–4446.
- Forline, L.C., 2008. Putting history back into historical ecology: some perspectives on the recent human ecology of the Amazon Basin. *J. Ecol. Anthropol.* 12, 69–74.
- Franco-Moraes, J., Baniwa, A.F.M.B., Costa, F.R.C., Lima, H.P., Clement, C.R., Shepard, G.H., 2019. Historical landscape domestication in ancestral forests with nutrient-poor soils in northwestern Amazonia. *For. Ecol. Manag.* 446, 317–330.
- Freitas, F.O., Bendel, G., Allaby, R.G., Brown, T.A., 2003. DNA from primitive maize landraces and archaeological remains: implications for the domestication of maize and its expansion into South America. *J. Archaeol. Sci.* 30, 901–908.
- Furquim, L.P., Neves, E.G., Shock, M.P., Watling, J., 2023. The constructed biodiversity, forest management and use of fire in ancient Amazon: an archaeological testimony on the last 14,000 Years of indigenous history. In: *Global Ecology in Historical Perspective: Monsoon Asia and beyond*. Springer, pp. 259–281.
- Furquim, L.P., Watling, J., Hilbert, L.M., Shock, M.P., Prestes-Carneiro, G., Calo, C.M., Py-Daniel, A.R., Brandão, K., Pugliese, F., Zimpel, C.A., 2021. Facing change through diversity: resilience and diversification of plant management strategies during the mid to late holocene transition at the Monte Castelo Shellmound, SW Amazonia. *Quaternary* 4, 8.
- Gallardo, B., Zieritz, A., Aldridge, D.C., 2015. The importance of the human footprint in shaping the global distribution of terrestrial, freshwater and marine invaders. *PLoS One* 10, e0125801.
- Glaser, B., 2007. Prehistorically modified soils of central Amazonia: a model for sustainable agriculture in the twenty-first century. *Phil. Transac. Royal Soc.: B* 362, 187–196.
- Glaser, B., Birk, J.J., 2012. State of the scientific knowledge on properties and genesis of Anthropogenic Dark Earths in Central Amazonia (terra preta de Índio). *Geochem. Cosmochim. Acta* 82, 39–51.
- Glaser, B., Woods, W.I., 2004. *Amazonian Dark Earths: Explorations in Space and Time*. Springer-Verlag, Berlin.
- Gosling, W.D., Cornelissen, H.L., McMichael, C.N.H., 2019. Reconstructing past fire temperatures from ancient charcoal material. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 520, 128–137.
- Goulart, A.C., Macario, K.D., Scheel-Ybert, R., Alves, E.Q., Bachelet, C., Pereira, B.B., Levis, C., Junior, B.H.M., Marimon, B.S., Quesada, C.A., 2017. Charcoal chronology of the Amazon forest: a record of biodiversity preserved by ancient fires. *Quat. Geochronol.* 41, 180–186.
- Guisan, A., Thuiller, W., 2005. Predicting species distribution: offering more than simple habitat models. *Ecol. Lett.* 8, 993–1009.
- Gutaker, R.M., Burbano, H.A., 2017. Reinforcing plant evolutionary genomics using ancient DNA. *Curr. Opin. Plant Biol.* 36, 38–45.
- Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'agrosa, C., Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., 2008. A global map of human impact on marine ecosystems. *Science* 319, 948–952.
- Hammersley, M., Atkinson, P., 2007. *Ethnography: Principles in Practice*. Routledge, Oxford, UK.
- Hammond, D.S., ter Steege, H., van der Borg, K., 2006. Upland soil charcoal in the wet tropical forests of Central Guyana. *Biotropica* 39, 153–160.
- Harlan, J.R., 1992. *Crops & Man*, 2 ed. American Society of Agronomy & Crop Science Society of America, Madison, WI.
- Hecht, S.B., 2003. Indigenous soil management and the creation of amazonian Dark earths: implications of kayapó practice. In: Lehmann, J., Kern, D.C., Glaser, B., Wodos, W.I. (Eds.), *Amazonian Dark Earths: Origin Properties Management*. Springer Netherlands, Dordrecht, pp. 355–372.
- Hecht, S.B., Cockburn, A., 2010. *The Fate of the Forest: Developers, Destroyers, and Defenders of the Amazon*. University of Chicago Press.
- Heckenberger, M., Russell, J., Fausto, C., Toney, J., Schmidt, M., Pereira, E., Franchetto, B., Kuikuro, A., 2008. Pre-Columbian urbanism, anthropogenic landscapes, and the future of the Amazon. *Science* 321, 1214–1217.
- Heckenberger, M.J., Kuikuro, A., Kuikuro, U.T., Russell, J.C., Schmidt, M., Fausto, C., Franchetto, B., 2003. Amazonia 1492: pristine forest or cultural parkland? *Science* 301, 1710–1714.
- Heijink, B.M., Mattijs, Q.A., Valencia, R., Philip, A.L., Piperno, D.R., McMichael, C.N., 2022. Long-term fire and vegetation change in northwestern Amazonia. *Biotropica* 55, 197–209.
- Heijink, B.M., McMichael, C.N., Piperno, D.R., Duivenvoorden, J.F., Cárdenas, D., Duque, A., 2020. Holocene increases in palm abundances in north-western Amazonia. *J. Biogeogr.* 47, 698–711.
- Hill, J., Black, S., Soto, D., Chavez, E., Vos, V., Mayle, F., 2023. Differing local-scale responses of Bolivian Amazon forest ecotones to middle Holocene drought based upon multiproxy soil data. *J. Quat. Sci.* 38 (6), 970–990.
- Hodell, D.A., Brenner, M., Curtis, J.H., 2005. Terminal Classic drought in the northern Maya lowlands inferred from multiple sediment cores in Lake Chichancanab (Mexico). *Quat. Sci. Rev.* 24, 1413–1427.
- Hodell, D.A., Curtis, J.H., Brenner, M., 1995. Possible role of climate in the collapse of ancient Maya civilization. *Nature* 357, 391–394.
- Hubbell, S.P., 1979. Tree dispersion, abundance, and diversity in a tropical dry forest. *Science* 203, 1299–1309.
- Huisman, S.N., Raczka, M.F., McMichael, C.N.H., 2018. Palm phytoliths of mid-elevation andean forests. *Front. Ecol. Evolution* 6, 193.
- Iriarte, J., 2016. Investigating amazonian Dark earths as agro-ecosystems and their impact on the regional landscapes of the lower Amazon. In: Stenborg, P. (Ed.), *Beyond Waters: Archaeology and Environmental History of the Amazonian Inland*. University of Gothenburg, Gothenburg, pp. 71–86.
- Iriarte, J., Elliott, S., Maezumi, S.Y., Alves, D., Gonda, R., Robinson, M., de Souza, J.G., Watling, J., Handley, J., 2020. The origins of Amazonian landscapes: plant cultivation, domestication and the spread of food production in tropical South America. *Quat. Sci. Rev.* 248, 106582.
- Iriarte, J., Glaser, B., Watling, J., Wainwright, A., Birk, J.J., Renard, D., Rostain, S., McKey, D., 2010. Late Holocene Neotropical agricultural landscapes: phytolith and stable carbon isotope analysis of raised fields from French Guianan coastal savannahs. *J. Archaeol. Sci.* 37, 2984–2994.
- Iriarte, J., Holst, I., Marozzi, O., Listopad, C., Alonso, E., Rinderknecht, A., Montaña, J., 2004. Evidence for cultivar adoption and emerging complexity during the mid-Holocene in the La Plata basin. *Nature* 432, 614–617.
- Jacobson, G.L., Bradshaw, R.H.W., 1981. The selection of sites for paleovegetational studies. *Quat. Res.* 16, 80–96.
- Junqueira, A., Almekinders, C., Stomph, T.-J., Clement, C., Struik, P., 2016. The role of Amazonian anthropogenic soils in shifting cultivation: learning from farmers' rationales. *Ecol. Soc.* 21.
- Junqueira, A.B., Shepard, G.H., Clement, C.R., 2010. Secondary forests on anthropogenic soils in Brazilian Amazonia conserve agrobiodiversity. *Biodivers. Conserv.* 19, 1933–1961.
- Junqueira, A.B., Shepard, G.H., Clement, C.R., 2011. Secondary forests on anthropogenic soils of the middle madeira river: valuation, local knowledge, and landscape domestication in Brazilian Amazonia. *Econ. Bot.* 1–15.
- Keeley, J.E., 2009. Fire intensity, fire severity and burn severity: a brief review and suggested usage. *Int. J. Wildland Fire* 18, 116–126.
- Kistler, L., Bieker, V.C., Martin, M.D., Pedersen, M.W., Ramos Madrigal, J., Wales, N., 2020. Ancient plant genomics in archaeology, herbaria, and the environment. *Annu. Rev. Plant Biol.* 71, 605–629.
- Kopenawa, D., Albert, B., 2023. *The Falling Sky, the Falling Sky*. Harvard University Press.
- Lehman, J., Kern, D.C., Glaser, B., Woods, W.I., 2003. *Amazonian Dark Earths: Origin, Properties, Management*. Kluwer Academic Publisher, Dordrecht, The Netherlands.
- Levis, C., Costa, F.R.C., Bongers, F., Peña-Claros, M., Clement, C.R., Junqueira, A.B., Neves, E.G., Tamañá, E.K., Figueiredo, F.O.G., Salomão, R.P., Castilho, C.V., Magnusson, W.E., Phillips, O.L., Guevara, J.E., Sabatier, D., Molino, J.-F., López, D. C., Mendoza, A.M., Pitman, N.C.A., Duque, A., Vargas, P.N., Zartman, C.E., Vasquez, R., Andrade, A., Camargo, J.L., Feldpausch, T.R., Laurance, S.G.W., Laurance, W.F., Killeen, T.J., Nascimento, H.E.M., Montano, J.C., Mostacedo, B., Amaral, L.L., Guimarães Vieira, I.C., Brienen, R., Castellanos, H., Terborgh, J., Carim, M.d.J.V., Guimarães, J.R.d.S., Coelho, L.d.S., Matos, F.D.d.A., Wittmann, F., Mogollón, H.F., Damasco, G., Dávila, N., García-Villacorta, R., Coronado, E.N.H., Emilio, T., Filho, D.d.A.L., Schiatti, J., Souza, P., Targhetta, N., Comiskey, J.A., Marimon, B.S., Marimon, B.-H., Neill, D., Alonso, A., Arroyo, L., Carvalho, F.A., de Souza, F.C., Dallmeier, F., Pansonato, M.P., Duivenvoorden, J.F., Fine, P.V.A., Stevenson, P.R., Araujo-Murakami, A., Aymard, C.G.A., Baraloto, C., do Amaral, D. D., Engel, J., Henkel, T.W., Maas, P., Petronelli, P., Revilla, J.D.C., Stropp, J., Daly, D., Gribel, R., Paredes, M.R., Silveira, M., Thomas-Caesar, R., Baker, T.R., da Silva, N.F., Ferreira, L.V., Peres, C.A., Silman, M.R., Cerón, C., Valverde, F.C., Di Fiore, A., Jimenez, E.M., Mora, M.C.P., Toledo, M., Barbosa, E.M., Bonates, L.C.d.M., Arboleda, N.C., Farias, E.d.S., Fuentes, A., Guillaumet, J.-L., Jørgensen, P.M., Malhi, Y., de Andrade Miranda, I.P., Phillips, J.F., Prieto, A., Rudas, A., Ruschel, A. R., Silva, N., von Hildebrand, P., Vos, V.A., Zent, E.L., Zent, S., Cintra, B.B.L., Nascimento, M.T., Oliveira, A.A., Ramirez-Angulo, H., Ramos, J.F., Rivas, G., Schöngart, J., Sierra, R., Tirado, M., van der Heijden, G., Torre, E.V., Wang, O., Young, K.R., Baider, C., Cano, A., Farfan-Rios, W., Ferreira, C., Hoffman, B., Mendoza, C., Mesones, I., Torres-Lezama, A., Medina, M.N.U., van Andel, T.R., Villarroel, D., Zagt, R., Alexiades, M.N., Balslev, H., Garcia-Cabrera, K., Gonzales, T., Hernandez, L., Huamantupa-Chuquimaco, I., Manzatto, A.G., Milliken, W., Cuenca, W.P., Pansini, S., Pauletto, D., Arevalo, F.R., Reis, N.F.C., Sampaio, A.F., Giraldo, L.E.U., Sandoval, E.H.V., Gamarra, L.V., Vela, C.I.A., ter Steege, H., 2017. Persistent effects of pre-Columbian plant domestication on Amazonian forest composition. *Science* 355, 925–931.
- Levis, C., de Souza, P.F., Schiatti, J., Emilio, T., da Veiga Pinto, J.L.P., Clement, C.R., Costa, F.R., 2012. Historical human footprint on modern tree species composition in the purus-madeira interfluvium, central Amazonia. *PLoS One* 7, e48559.
- Levis, C., Flores, B.M., Moreira, P.A., Luiz, B.G., Alves, R.P., Franco-Moraes, J., Lins, J., Konings, E., Peña-Claros, M., Bongers, F., 2018. How people domesticated Amazonian forests. *Front. Ecol. Evolution* 5, 171.
- Levis, C., Peña-Claros, M., Clement, C.R., Costa, F.R.C., Alves, R.P., Ferreira, M.J., Figueiredo, C.G., Bongers, F., 2020. Pre-Columbian soil fertilization and current

- management maintain food resource availability in old-growth Amazonian forests. *Plant Soil* 450, 29–48.
- Lewis, S.L., Maslin, M.A., 2015. Defining the anthropocene. *Nature* 519, 171–180.
- Livi-Bacci, M., 2016. The depopulation of upper Amazonia in colonial times. *Rev. Indias* 76, 419–448.
- Lombardo, U., Iriarte, J., Hilbert, L., Ruiz-Pérez, J., Capriles, J.M., Veit, H., 2020. Early Holocene crop cultivation and landscape modification in Amazonia. *Nature* 190–193.
- Loughlin, N.J., Gosling, W.D., Mothes, P., Montoya, E., 2018. Ecological consequences of post-Columbian indigenous depopulation in the Andean–Amazonian corridor. *Nat. Ecol. & Evolution* 2, 1233–1236.
- Lynch, J.A., Clark, J.S., Stocks, B.J., 2004. Charcoal production, dispersal, and deposition from the Fort Providence experimental fire: interpreting fire regimes from charcoal records in boreal forests. *Can. J. For. Res.* 34, 1642–1656.
- Maezumi, S., Power, M., Mayle, F., McLaughlan, K., Iriarte, J., 2015. The Effects of Past Climate Variability on Fire and Vegetation in the Cerrado Savanna Ecosystem of the Huanchaca Mesetta. Noel Kempff Mercado National Park, NE Bolivia.
- Maezumi, S.Y., Alves, D., Robinson, M., de Souza, J.G., Levis, C., Barnett, R.L., de Oliveira, E.A., Urrego, D., Schaan, D., Iriarte, J., 2018a. The legacy of 4,500 years of polyculture agroforestry in the eastern Amazon. *Nat. Plants* 4, 540–547.
- Maezumi, S.Y., Gosling, W.D., Kirschner, J., Chevalier, M., Cornelissen, H.L., Heinecke, T., McMichael, C.N., 2021. A Modern Analogue Matching Approach to Characterize Fire Temperatures and Plant Species from Charcoal. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 110580.
- Maezumi, S.Y., Whitney, B.S., Mayle, F.E., de Souza, J.G., Iriarte, J., 2018b. Reassessing climate and pre-Columbian drivers of paleofire activity in the Bolivian Amazon. *Quat. Int.* 488, 81–94.
- Malhi, Y., Phillips, O.L., Lloyd, J., Baker, T., Wright, J., Almeida, S., Arroyo, L., Frederiksen, T., Grace, J., Higuchi, N., Killeen, T., Laurance, W.F., Leano, C., Lewis, S., Meir, P., Menteagudo, A., Neill, D., Nunez Vargas, P., Panfil, S.N., Patino, S., Pitman, N., Quesada, C.A., Rudas-Li, A., Solomao, R., Saleska, S., Silva, N., Silveira, M., Sombroek, W.G., Valencia, R., Vasquez Martinez, R., Vieira, I.C.G., Vinceti, B., 2002. An international network to monitor the structure, composition and dynamics of Amazonian forests (RAINFOR). *J. Veg. Sci.* 13, 439–450.
- Malhi, Y., Roberts, J.T., Betts, R.A., Killeen, T.J., Li, W., Nobre, C.A., 2008. Climate change, deforestation, and the fate of the Amazon. *Science* 319, 169–172.
- Marlon, J.R., Bartlein, P.J., Daniiau, A.-L., Harrison, S.P., Maezumi, S.Y., Power, M.J., Tinner, W., Vanniére, B., 2013. Global biomass burning: a synthesis and review of Holocene paleofire records and their controls. *Quat. Sci. Rev.* 65, 5–25.
- Marlon, J.R., Kelly, R., Daniiau, A.-L., Vanniére, B., Power, M.J., Bartlein, P., Higuera, P., Blarquez, O., Brewer, S., Brücher, T., 2016. Reconstructions of biomass burning from sediment charcoal records to improve data-model comparisons. *Biogeosciences* 13, 3225–3244.
- Martin, G.J., 2004. *Ethnobotany: a Methods Manual*. Routledge, London.
- Mayle, F.E., Burbridge, R., Killeen, T.J., 2000. Millennial-scale dynamics of southern Amazonian rain forests. *Science* 290, 2291–2294.
- Mayle, F.E., Iriarte, J., 2014. Integrated palaeoecology and archaeology—A powerful approach for understanding pre-Columbian Amazonia. *J. Archaeol. Sci.* 51, 54–64.
- McClenachan, L., Cooper, A.B., McKenzie, M.G., Drew, J.A., 2015. The importance of surprising results and best practices in Historical Ecology. *Bioscience* 65, 932–939.
- McEwan, C., 2001. *Unknown Amazon: Culture in Nature in Ancient Brazil*. British Museum, London.
- McKey, D., Rostain, S., Iriarte, J., Glaser, B., Birk, J., Holst, I., Renard, D., 2010. Pre-Columbian agricultural landscapes, ecosystem engineers, and self-organized patchiness in Amazonia. *Proc. Natl. Acad. Sci. USA* 107, 7823–7828.
- McLachlan, J.S., Clark, J.S., 2005. Molecular indicators of tree migration capacity under rapid climate change. *Ecology* 86, 2088–2098.
- McMichael, C., Palace, M., Bush, M., Braswell, B., Hagen, S., Neves, E., Silman, M., Tamañana, E., Czarnecki, C., 2014a. Predicting pre-Columbian anthropogenic soils in Amazonia. *Proc. Biol. Sci.* 281, 20132475.
- McMichael, C., Piperno, D., Neves, E., Bush, M., Almeida, F., Mongeló, G., Eyjolfsson, M.B., 2015. Phytolith assemblages along a gradient of ancient human disturbance in western Amazonia. *Front. Ecol. Evolution* 3, 141.
- McMichael, C., Piperno, D.R., Bush, M.B., Silman, M.R., Zimmerman, A.R., Raczka, M.F., Lobato, L.C., 2012a. Sparse pre-Columbian human habitation in western Amazonia. *Science* 336, 1429–1431.
- McMichael, C.H., Bush, M.B., Piperno, D.R., Silman, M.R., Zimmerman, A.R., Anderson, C., 2012b. Spatial and temporal scales of pre-Columbian disturbance associated with western Amazonian lakes. *Holocene* 22, 131–141.
- McMichael, C.H., Correa Metrio, A., Bush, M.B., 2012c. Pre-Columbian fire regimes in lowland tropical rainforests of southeastern Peru. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 342–343, 73–83.
- McMichael, C.H., Palace, M.W., Golightly, M., 2014b. Bamboo-dominated forests and pre-Columbian earthwork formations in south-western Amazonia. *J. Biogeogr.* 41, 1733–1745.
- McMichael, C.N., 2021. Ecological legacies of past human activities in Amazonian forests. *New Phytol.* 229, 2492–2496.
- McMichael, C.N., Bush, M.B., 2019. Spatiotemporal patterns of pre-Columbian people in Amazonia. *Quat. Res.* 92, 53–69.
- McMichael, C.N., Vink, V., Heijink, B.M., Witteveen, N.H., Piperno, D.R., Gosling, W.D., Bush, M.B., 2023. Ecological legacies of past fire and human activity in a Panamanian forest. *Plants, People, Planet* 5, 281–291.
- McMichael, C.N.H., Matthews-Bird, F., Farfan-Rios, W., Feeley, K.J., 2017. Ancient human disturbances may be skewing our understanding of Amazonian forests. *Proc. Natl. Acad. Sci. USA* 114, 522–527.
- Medina, J., 1934. *The discovery of the Amazon according to the account of Friar Caspar de Carvajal and other documents*. American Geographical Society, New York.
- Meyer, R.S., DuVal, A.E., Jensen, H.R., 2012. Patterns and processes in crop domestication: an historical review and quantitative analysis of 203 global food crops. *New Phytol.* 196, 29–48.
- Meyer, R.S., Purugganan, M.D., 2013. Evolution of crop species: genetics of domestication and diversification. *Nat. Rev. Genet.* 14, 840–852.
- Moraes, C.d.P., 2015. O determinismo agrícola na arqueologia amazônica. *Estud. Avançados* 29, 25–43.
- Morcote-Ríos, G., Bernal, R., Raz, L., 2016. Phytoliths as a tool for archaeobotanical, palaeobotanical and palaeoecological studies in Amazonian palms. *Bot. J. Linn. Soc.* 182, 348–360.
- Morcote-Ríos, G., Giraldo-Cañas, D., Raz, L., 2015. Catálogo ilustrado de fitolitos contemporáneos con énfasis arqueológico y paleoecológico I. Gramíneas amazónicas de Colombia. Universidad Nacional de Colombia, Bogotá, Colombia.
- Moreira, P.A., Mariac, C., Zekraoui, L., Couderc, M., Rodrigues, D.P., Clement, C.R., Vigouroux, Y., 2017. Human management and hybridization shape treegourd fruits in the Brazilian Amazon Basin. *Evol. Appl.* 10 (6), 577–589.
- Nascimento, M.N., Heijink, B.M., Bush, M.B., Gosling, W.D., McMichael, C.N.H., 2022. Early to mid-Holocene human activity exerted gradual influences on Amazonian forest vegetation. *Phil. Trans. Biol. Sci.* 377, 20200498.
- Nascimento, M.d.N., McMichael, C.N.H., Kleijwegt, Z., Åkesson, C., Gredal, C., Maezumi, S.Y., Bush, M.B., 2023. Fire in the clouds: How changing land use shaped an Andean biodiversity hotspot. *Quat. Sci. Rev.* 317, 108278.
- Nascimento, M.N., Peters-Schulze, G., Martins, G.S., Cordeiro, R.C., Turcq, B., Moreira, L. S., Bush, M.B., 2021. Limnological response to climatic changes in western Amazonia over the last millennium. *Front. Biogeogr.* 13.
- Nepstad, D., Lefebvre, P., Silva, U.L.D., Tomasella, J., Schlesinger, P., Solorzano, L., Moutinho, P., Ray, D., Benito, J.G., 2004. Amazon drought and its implications for forest flammability and tree growth: a basin-wide analysis. *Global Change Biol.* 10, 704–717.
- Neves, E.G., 2013. Was agriculture a key productive activity in pre-colonial Amazonia? The stable productive basis for social equality in the central Amazon. In: Brondizio, E.S., Moran, E.F. (Eds.), *Human-Environment Interactions*. Springer, Dordrecht, pp. 371–388.
- Neves, E.G., Furquim, L.P., Levis, C., Rocha, B.C., Watling, J.G., Almeida, F.O., Betancourt, C.J., Junqueira, A.B., Moraes, C.P., Morcote-Ríos, G., 2021. Peoples of the Amazon before European colonization. *Sci. Panel Amazon Amazon Assess. Rep.* Chapter 8, 1–40.
- Neves, E.G., Petersen, J.B., Bartone, R.N., Heckenberger, M.J., 2004. The timing of *terra preta* formation in the central Amazon: archaeological data from three sites. In: Glaser, B., Woods, W.I. (Eds.), *Amazonian Dark Earths: Explorations in Space and Time*. Springer, Berlin, pp. 125–133.
- Olsen, K.M., Schaal, B.A., 1999. Evidence on the origin of cassava: phylogeography of *Manihot esculenta*. *Proc. Natl. Acad. Sci. USA* 96, 5586–5591.
- Olsen, K.M., Schaal, B.A., 2001. Microsatellite variation in cassava (*Manihot esculenta*, Euphorbiaceae) and its wild relatives: further evidence for a southern Amazonian origin of domestication. *Am. J. Bot.* 88, 131–142.
- Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V.N., Underwood, E.C., D'Amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C., 2001. Terrestrial ecoregions of the world: a new map of life on earth. *Bioscience* 51, 933–938.
- Orvis, K.H., Lane, C.S., Horn, S.P., 2005. Laboratory production of vouchered reference charcoal from small wood samples and non-woody plant tissues. *Palynology* 29, 1–11.
- Oxford, 2016. *Oxford Dictionaries*. Oxford University Press.
- Pachauri, R.K., Meyer, L., Plattner, G.-K., Stocker, T., 2015. IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC.
- Palace, M.W., McMichael, C.N.H., Braswell, B.H., Hagen, S.C., Bush, M.B., Neves, E., Tamañana, E., Herrick, C., Frolking, S., 2017. Ancient Amazonian populations left lasting impacts on forest structure. *Ecosphere* 8, e02035 n/a.
- Parducci, L., Bennett, K.D., Ficetola, G.F., Alsos, I.G., Suyama, Y., Wood, J.R., Pedersen, M.W., 2017. Ancient plant DNA in lake sediments. *New Phytol.* 214, 924–942.
- Parsons, L., LeRoy, S., Overpeck, J., Bush, M., Cárdenes-Sandí, G., Saleska, S., 2018. The threat of multi-year drought in western Amazonia. *Water Resour. Res.* 54, 5890–5904.
- Pearsall, D.M., Chandler-Ezell, K., Zeidler, J.A., 2004. Maize in ancient Ecuador: results of residue analysis of stone tools from the Real Alto site. *J. Archaeol. Sci.* 31, 423–442.
- Pessenda, L.C.R., Gomes, B.M., Aravena, R., Ribeiro, A.S., Boulet, R., Gouveia, S.E.M., 1998. The carbon isotope record in soils along a forest-cerrado ecosystem transect: implications for vegetation changes in the Rondonia state, southwestern Brazilian Amazon region. *Holocene* 8, 599–603.
- Phillips, O., Hall, P., Gentry, A.H., Sawyer, S., Vasquez, R., 1994. Dynamics and species richness of tropical rain forests. *Proc. Natl. Acad. Sci. USA* 91, 2805–2809.
1988. *Phytolith Analysis: An Archaeological and Geological Perspective*, 1988. Piperno Publisher: Academic Press.
- Piperno, D.R., 2006. *Phytoliths: A Comprehensive Guide for Archaeologists and Paleoeologists*. Alta Mira Press, Lanham, MD.
- Piperno, D.R., 2011. The origins of plant cultivation and domestication in the New World tropics. *Curr. Anthropol.* 52, S453–S470.
- Piperno, D.R., 2016. Phytolith radiocarbon dating in archaeological and paleoecological research: a case study of phytoliths from modern Neotropical plants and a review of the previous dating evidence. *J. Archaeol. Sci.* 68, 54–61.

- Piperno, D.R., McMichael, C., 2023. Phytoliths in modern plants from Amazonia and the Neotropics at large: II. Enhancement of eudicotyledon reference collections. *Quat. Int.* 655, 1–17.
- Piperno, D.R., McMichael, C., Bush, M.B., 2015. Amazonia and the Anthropocene: what was the spatial extent and intensity of human landscape modification in the Amazon Basin at the end of prehistory? *Holocene* 25, 1588–1597.
- Piperno, D.R., McMichael, C.H., Pitman, N.C., Andino, J.E.G., Paredes, M.R., Heijink, B. M., Torres-Montenegro, L.A., 2021. A 5,000-year vegetation and fire history for tierra firme forests in the Medio Putumayo-Algodón watersheds, northeastern Peru. *Proc. Natl. Acad. Sci. USA* 118 (40), e2022213118.
- Piperno, D.R., McMichael, C.N., Bush, M.B., 2019. Finding forest management in prehistoric Amazonia. *Anthropocene* 26, 100211.
- Plumpton, H.J., Mayle, F.M., Whitney, B.S., 2020. Paleoecological potential of phytoliths from lake sediment records from the tropical lowlands of Bolivia. *Rev. Palaeobot. Palynol.* 275, 104113.
- Politis, G.G., 2014. Ethnoarchaeology: Approaches to Fieldwork, *Field Archaeology from Around the World: Ideas and Approaches*. Springer, pp. 83–88.
- Power, M., Whitney, B., Mayle, F., Neves, D., de Boer, E., Maclean, K., 2016. Fire, climate and vegetation linkages in the Bolivian Chiquitano seasonally dry tropical forest. *Phil. Trans. R. Soc. B* 371, 20150165.
- Ramos-Neto, M.B., Pivello, V.R., 2000. Lightning fires in a Brazilian savanna National Park: rethinking management strategies. *Environ. Manag.* 26, 675–684.
- Reeve, M.-E., 1994. Regional interaction in the Western Amazon: the early colonial encounter and the Jesuit years: 1538-1767. *Ethnohistory* 41, 106–138.
- Rhodes, A.N., 1998. A method for the preparation and quantification of microscopic charcoal from terrestrial and lacustrine sediment cores. *Holocene* 8, 113–117.
- Ribeiro, F.A.N., Moraes, C.d.P., Valle, R.B.M., 2023. Through zo'ë paths and capoeiras: networks of relations in the guianas and the "sparse population" question. In: *Historical Ecology and Landscape Archaeology in Lowland South America*. Springer, pp. 51–85.
- Riris, P., Arroyo-Kalin, M., 2019. Widespread population decline in South America correlates with mid-Holocene climate change. *Sci. Rep.* 9, 1–10.
- Roberts, P., Hunt, C., Arroyo-Kalin, M., Evans, D., Boivin, N., 2017. The deep human prehistory of global tropical forests and its relevance for modern conservation. *Nat. Plants* 3, 17093.
- Rodbell, D.T., Seltzer, G.O., Anderson, D.M., Abbott, M.B., Enfield, D.B., Newman, J.H., 1999. An ~15,000-year record of El Niño-driven alluviation in southwestern Ecuador. *Science* 283, 516–520.
- Rodríguez-Zorro, P.A., Enters, D., Hermanowski, B., da Costa, M.L., Behling, H., 2015. Vegetation changes and human impact inferred from an oxbow lake in southwestern Amazonia, Brazil since the 19th century. *J. South Am. Earth Sci.* 62, 186–194.
- Roosevelt, A., 1989. Resource management in Amazonia before the conquest: beyond ethnographic projection. *Adv. Econ. Bot.* 30–62.
- Roosevelt, A.C., 2013. The Amazon and the anthropocene: 13,000 Years of human influence in a tropical rainforest. *Anthropocene* 4, 69–87.
- Roosevelt, A.C., Lima da Costa, M., Lopes Machado, C., Michab, M., Mercier, N., Valladas, H., Feathers, J., Barnett, W., Imazio da Silveira, M., Henderson, A., Sliva, J., Chernoff, B., Reese, D.S., Holman, J.A., Toth, N., Schick, K., 1996. Paleolithic cave dwellers in the Amazon: the peopling of the Americas. *Science* 272, 373–384.
- Ross, N.J., 2011. Modern tree species composition reflects ancient Maya "forest gardens" in northwest Belize. *Ecol. Appl.* 21, 75–84.
- Roullier, C., Benoit, L., McKey, D.B., Lebot, V., 2013. Historical collections reveal patterns of diffusion of sweet potato in Oceania obscured by modern plant movements and recombination. *Proc. Natl. Acad. Sci. U.S.A.* 110, 2205–2210.
- Rozas-Davila, A., Correa-Metrio, A., McMichael, C.N., Bush, M.B., 2021. When the grass wasn't greener: megafaunal ecology and paleodroughts. *Quat. Sci. Rev.* 266, 107073.
- Rozas-Davila, A., Valencia, B.G., Bush, M.B., 2016. The functional extinction of Andean megafauna. *Ecology* 97, 2533–2539.
- Sahoo, P.K., Guimarães, J.T.F., Souza-Filho, P.W.M., Powell, M.A., da Silva, M.S., Moraes, A.M., Alves, R., Leite, A.S., Júnior, W.N., Rodrigues, T.M., 2019. Statistical analysis of lake sediment geochemical data for understanding surface geological factors and processes: an example from Amazonian upland lakes, Brazil. *Catena* 175, 47–62.
- Sanchez-Cabeza, J., Ruiz-Fernández, A., 2012. 210Pb sediment radiochronology: an integrated formulation and classification of dating models. *Geochem. Cosmochim. Acta* 82, 183–200.
- Sanford, R.L., Horn, S.P., 2000. Holocene rain-forest wilderness: a Neotropical perspective on humans as an exotic, invasive species. In: Cole, D., McCool, S.F. (Eds.), *Wilderness Science in a Time of Change*. US Dept. Of Agriculture, Forest Service. Proc. RMRS-P-000. Rocky Mountain Research Station, Ogden, UT.
- Schaan, D., Pärssinen, M., Saunalauma, S., Ranzi, A., Bueno, M., Barbosa, A., 2012. New radiometric dates for pre-Columbian (2000 - 700 BP) earthworks in western Amazonia, Brazil. *J. Field Archaeol.* 37, 132–142.
- Scheel-Ybert, R., Beauclair, M., Buarque, A., 2014. The forest people: landscape and firewood use in the Araruama region, southeastern Brazil, during the late Holocene. *Veg. Hist. Archaeobotany* 23, 97–111.
- Schmidt, M.J., Rapp Py-Daniel, A., de Paula Moraes, C., Valle, R.B.M., Caromano, C.F., Teixeira, W.G., Barbosa, C.A., Fonseca, J.A., Magalhães, M.P., Silva do Carmo Santos, D., da Silva e Silva, R., Guapindaia, V.L., Moraes, B., Lima, H.P., Neves, E.G., Heckenberger, M.J., 2014. Dark earths and the human built landscape in Amazonia: a widespread pattern of anthroposol formation. *J. Archaeol. Sci.* 42, 152–165.
- Schöngart, J., Gribel, R., Ferreira da Fonseca-Junior, S., Haugaasen, T., 2015. Age and growth patterns of Brazil nut trees (*Bertholletia excelsa* bonpl.) in Amazonia, Brazil. *Biotropica* 47, 550–558.
- Shepard Jr., G.H., Clement, C.R., Lima, H.P., dos Santos, G.M., de Paula Moraes, C., Neves, E.G., 2020. Ancient and Traditional Agriculture in South America: Tropical Lowlands. *Oxford Research Encyclopedia of Environmental Science*.
- Smith, B.D., 2006. Documenting domesticated plants in the archaeological record. In: Zeder, M.A., Bradley, D.G., Emshwiller, E., Smith, B.D. (Eds.), *Documenting Domestication. New Genetic and Archaeological Paradigms*. University of California Press, Berkeley, pp. 15–24.
- Söderström, M., Eriksson, J., Isendahl, C., Schaan, D.P., Stenborg, P., Rebellato, L., Piikki, K., 2016. Sensor mapping of amazonian Dark earths in deforested croplands. *Geoderma* 281, 58–68.
- Souza, J.G., Schaan, D.P., Robinson, M., Barbosa, A.D., Aragão, L.E., Marimon Jr., B.H., Marimon, B.S., Silva, I.B., Khan, S.S., Nakahara, F.R., 2018. Pre-Columbian earth-builders settled along the entire southern rim of the Amazon. *Nat. Commun.* 9, 1125.
- Szabó, P., Hédl, R., 2011. Advancing the integration of history and ecology for conservation. *Conserv. Biol.* 25, 680–687.
- Taylor, R.E., Bar-Yosef, O., 2016. *Radiocarbon Dating: an Archaeological Perspective*. Routledge.
- Tengö, M., Brondizio, E.S., Elmqvist, T., Malmer, P., Spierenburg, M., 2014. Connecting diverse knowledge systems for enhanced ecosystem governance: the multiple evidence base approach. *Ambio* 43, 579–591.
- ter Steege, H., Pitman, N.C.A., Sabatier, D., Baraloto, C., Salomão, R.P., Guevara, J.E., Phillips, O.L., Castilho, C.V., Magnusson, W.E., Molino, J.-F., Monteagudo, A., Núñez Vargas, P., Montero, J.C., Feldpausch, T.R., Coronado, E.N.H., Killeen, T.J., Mostacedo, B., Vasquez, R., Assis, R.L., Terborgh, J., Wittmann, F., Andrade, A., Laurance, W.F., Laurance, S.G.W., Marimon, B.S., Marimon, B.-H., Guimarães Vieira, I.C., Amaral, I.L., Brien, R., Castellanos, H., Cárdenas López, D., Duivenvoorden, J.F., Mogollón, H.F., Matos, F.D.d.A., Dávila, N., García-Villacorta, R., Stevenson Diaz, P.R., Costa, F., Emilio, T., Levis, C., Schiatti, J., Souza, P., Alonso, A., Dallmeier, F., Montoya, A.J.D., Fernandez Piedade, M.T., Araujo-Murakami, A., Arroyo, L., Gribel, R., Fine, P.V.A., Peres, C.A., Toledo, M., Aymard, C.G.A., Baker, T.R., Cerón, C., Engel, J., Henkel, T.W., Maas, P., Petronelli, P., Stropp, J., Zartman, C.E., Daly, D., Neill, D., Silveira, M., Paredes, M. R., Chave, J., Lima Filho, D.d.A., Jørgensen, P.M., Fuentes, A., Schöngart, J., Cornejo Valverde, F., Di Fiore, A., Jimenez, E.M., Peñuela Mora, M.C., Phillips, J.F., Rivas, G., van Andel, T.R., von Hildebrand, P., Hoffman, B., Zent, E.L., Malhi, Y., Prieto, A., Rudas, A., Ruschell, A.R., Silva, N., Vos, V., Zent, S., Oliveira, A.A., Schutz, A.C., Gonzales, T., Trindade Nascimento, M., Ramirez-Angulo, H., Sierra, R., Tirado, M., Umaña Medina, M.N., van der Heijden, G., Vela, C.I.A., Vilanova Torre, E., Vriesendorp, C., Wang, O., Young, K.R., Baidar, C., Balslev, H., Ferreira, C., Mesones, I., Torres-Lezama, A., Urrego Giraldo, L.E., Zagt, R.W., Alexiades, M.N., Hernandez, L., Huamantupa-Chuquimaco, I., Milliken, W., Palacios Cuenca, W., Pausetto, D., Valderrama Sandoval, E., Valenzuela Gamarra, L., Dexter, K.G., Feeley, K., Lopez-Gonzalez, G., Silman, B.R., 2013. Hyperdominance in the amazonian tree flora. *Science* 342, 1243092.
- Thayn, J.B., Price, K.P., Woods, W.I., 2011. Locating Amazonian Dark Earths (ADE) using vegetation vigour as a surrogate for soil type. *Int. J. Rem. Sens.* 32, 6713–6729.
- Thomas, E., Alcázar Caicedo, C., McMichael, C.H., Corvera, R., Loo, J., 2015. Uncovering spatial patterns in the natural and human history of Brazil nut (*Bertholletia excelsa*) across the Amazon Basin. *J. Biogeogr.* 42, 1367–1382.
- Trisos, C.H., Auerbach, J., Katti, M., 2021. Decoloniality and anti-oppressive practices for a more ethical ecology. *Nat. Ecol. & Evol.* 5, 1205–1212.
- Urrego, D.H., Bush, M.B., Silman, M.R., Niccum, B.A., De La Rosa, P., McMichael, C.H., Hagen, S., Palace, M., 2013. Holocene fires, forest stability and human occupation in south-western Amazonia. *J. Biogeogr.* 40, 521–533.
- Vitousek, P.M., Mooney, H.A., Lubchenco, J., Melillo, J.M., 1997. Human domination of earth's ecosystems. *Science* 277, 494–498.
- Wales, N., Andersen, K., Cappellini, E., Avila-Arcos, M.C., Gilbert, M.T.P., 2014. Optimization of DNA recovery and amplification from non-carbonized archaeological remains. *PLoS One* 9, e86827.
- Walker, R.S., Ferguson, J.R., Olmeda, A., Hamilton, M.J., Elghammer, J., Buchanan, B., 2023. Predicting the geographic distribution of ancient Amazonian archaeological sites with machine learning. *PeerJ* 11, e15137.
- Watling, J., Iriarte, J., Mayle, F.E., Schaan, D., Pessenda, L.C.R., Loader, N.J., Street-Perrott, F.A., Dickau, R.E., Damasceno, A., Ranzi, A., 2017. Impact of pre-Columbian "geoglyph" builders on Amazonian forests. *Proc. Natl. Acad. Sci. USA* 114, 1868–1873.
- Watling, J., Iriarte, J., Whitney, B., Consuelo, E., Mayle, F., Castro, W., Schaan, D., Feldpausch, T.R., 2016. Differentiation of neotropical ecosystems by modern soil phytolith assemblages and its implications for palaeoenvironmental and archaeological reconstructions II: southwestern Amazonian forests. *Rev. Palaeobot. Palynol.* 226, 30–43.
- Watling, J., Shock, M.P., Mongeló, G.Z., Almeida, F.O., Kater, T., De Oliveira, P.E., Neves, E.G., 2018. Direct archaeological evidence for Southwestern Amazonia as an early plant domestication and food production centre. *PLoS One* 13, e0199868.
- Weinstein, B., 1983. *The Amazon Rubber Boom, 1850-1920*. Stanford University Press.
- Whitlock, C., Larsen, C., 2002. Charcoal as a fire proxy. *Track. Environ. Change Using Lake Sediments* 75–97.
- Whitney, B.S., Dickau, R., Mayle, F.E., Soto, J.D., Iriarte, J., 2013. Pre-Columbian landscape impact and agriculture in the Monumental Mound region of the Llanos de Moxos, lowland Bolivia. *Quat. Res.* 80, 207–217.
- Whitney, B.S., Dickau, R., Mayle, F.E., Walker, J.H., Soto, J.D., Iriarte, J., 2014. Pre-Columbian raised-field agriculture and land use in the Bolivian Amazon. *Holocene*, 0959683613517401.
- Whitney, B.S., Mayle, F.E., Punyasena, S.W., Fitzpatrick, K.A., Burn, M.J., Guillen, R., Chavez, E., Mann, D., Pennington, R.T., Metcalfe, S.E., 2011. A 45 kyr palaeoclimate

- record from the lowland interior of tropical South America. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 307, 177–192.
- Whitney, B.S., Rushton, E.A., Carson, J.F., Iriarte, J., Mayle, F.E., 2012. An improved methodology for the recovery of *Zea mays* and other large crop pollen, with implications for environmental archaeology in the Neotropics. *Holocene* 22, 1087–1096.
- Whitney, B.S., Smallman, T.L., Mitchard, E.T., Carson, J.F., Mayle, F.E., Bunting, M.J., 2019. Constraining pollen-based estimates of forest cover in the Amazon: a simulation approach. *Holocene* 29, 262–270.
- WinklerPrins, A., 2009. Sweep and char and the creation of amazonian Dark earths in homegardens. In: Woods, W.I., Teixeira, W.G., Lehmann, J., Steiner, C., WinklerPrins, A., Rebellato, L. (Eds.), *Amazonian Dark Earths: Wim Sombroek's Vision*. Springer Netherlands, Dordrecht, pp. 205–211.
- WinklerPrins, A.M.A., Levis, C., 2020. Reframing pre-European Amazonia through an anthropocene lens. *Ann. Assoc. Am. Geogr.* 111, 858–868.
- WinklerPrins, A.M.G.A., Aldrich, S.P., 2010. Locating amazonian Dark earths: creating an interactive GIS of known locations. *J. Lat. Am. Geogr.* 9, 33–50.
- Witteveen, N., Hobus, C., Philip, A., Piperno, D., McMichael, C., 2022. The variability of Amazonian palm phytoliths. *Rev. Palaeobot. Palynol.* 300, 104613.
- Young, D.N., Duncan, N.A., Walker, J.H., 2023. Starch grain analysis of ceramic residue from forest islands associated with raised fields in west central Mojos, Bolivia. *J. Archaeol. Sci.: Report* 47, 103761.
- Zocatelli, R., Lavrieux, M., Guillemot, T., Chassiot, L., Le Milbeau, C., Jacob, J., 2017. Fecal biomarker imprints as indicators of past human land uses: source distinction and preservation potential in archaeological and natural archives. *J. Archaeol. Sci.* 81, 79–89.