

DESCRIPTION OF A NEW WARBLER FROM YUCATAN.

R. (C)

By ROBERT RIDGWAY.

Granatellus sallæi boucardi, subsp. nov.

Granatellus sallæi, SALV. & GODM., Biol. Centr.-Am. Aves. I., 1881, 161 (part; spec's ex Yucatan).—BOUCARD, P. Z. S., 1883, 441 (Yucatan).

SP. CHAR.—Similar to *G. sallæi*, SCL., but with chin and throat light plumbeous-gray instead of deep plumbeous, the upper parts decidedly paler plumbeous, the red of the breast paler, and the wing shorter.

Adult ♂ (type, No. 81760, U. S. Nat. Mus., Yucatan, 1879, A. Boucard): Above bluish plumbeous, the sides of the crown bordered by a broad but rather indistinct black line, beneath which, and extending from the upper posterior portion of the eye over the auriculars to the occiput, is a broad stripe of white; auriculars and cheeks lighter plumbeous than the upper parts, and the lores, malar region, chin, and throat still paler, or light ash-gray. Entire breast, middle of abdomen (longitudinally), anal region, and crissum fine pale rose-red; lining of wing, sides, and flanks, pure white. Tail black, the feathers edged, especially towards the base, with bluish plumbeous; tips of two outer rectrices narrowly white, and shaft of outer rectrix entirely white. Wing, 2.30; tail, 2.40; culmen, .55; depth of bill at base, .20; tarsus, .75; middle toe, .45.

An adult ♂ of *G. sallæi*, SCL., from Protrero, near Cordoba, Mexico (No. 41599, U. S. Nat. Mus.), measures as follows: Wing, 2.50; tail, 2.40; culmen, .55; depth of bill at base, .20; tarsus, .75; middle toe, .45. There is, therefore, no difference in dimensions except in the length of the wing, but the two forms may be easily distinguished by the marked difference in their coloration.

DESCRIPTION OF TWO NEW BIRDS FROM COSTA RICA.

By ROBERT RIDGWAY.

1. *Cyanocorax cucullatus*, sp. nov.

SP. CHAR.—Similar to *C. ornatus* (LESS.), but smaller (the wing about half an inch shorter), the blue hood without white border (except anteriorly), and the tibiæ blue instead of black.

Adult (type, No. 101845, U. S. Nat. Mus., Navarro, Costa Rica, October 30, 1882; Juan Cooper, collector; received from José C. Zeledon): Campanuliform patch covering upper part of nape, occiput, and hinder part of crown rich sky-blue, gradually passing anteriorly into silvery white; rest of head and neck, together with upper part of jugulum and back deep black, gradually changing, on breast, back, and scapulars to Berlin blue, this in turn changing, on remaining portions, above and below, to deep Paris blue, the tail with a greenish cast in certain lights.