

THE VARIATION EXHIBITED BY THAMNOPHIS ORDINOIDES (BAIRD AND GIRARD), A GARTER SNAKE INHABITING THE SAN FRANCISCO PENINSULA.

By JOSEPH C. THOMPSON,
Surgeon, United States Navy.

INTRODUCTION.

The more one reviews the literature of the garter snakes of North America, the more one becomes impressed with the necessity of ascertaining the complete range of variation that may be exhibited by each of the well-established species in the genus. To determine this it will be requisite to collect a fair series of specimens from a restricted locality and to record its slightest variations. Similar studies will have to be made of sets from widely separated regions. In order to facilitate the comparison of the data an endeavor should be made to conform to a uniform method of tabulating the figures and presenting the facts.

In the past *T. ordinoides* has been divided into as many as eight species and subspecies; this alone implies that the form is subject to considerable variation. What this variation really amounts to can only be appreciated by one who has seen large series from all parts of its range, for it is difficult to believe that the dwarfed and sombre-hued examples from the north, with the minimum scale count of 17-15, are one and the same species as the large brilliantly colored specimens from the south or the veritable giant from the Santa Clara Valley, whose body alone measures 1,040 mm. and has a maximum scale formula of 21-23-21-19-17.

Dr. Ruthven in his Memoir on the Garter Snakes of North America¹ recognizes *Thamnophis elegans* (Baird and Girard) 1853, and *Thamnophis ordinoides* (Baird and Girard) 1852, as distinct species. In speaking of *T. elegans*² he writes: ". . . its western limit can not be drawn exactly owing to the fact that it intergrades with another form . . ." referring to *T. ordinoides*. In speaking of

¹ 1908, Bull. 61, U. S. National Museum.

² *Idem*, p. 143.

*T. ordinoides*¹ he states: "I believe that it is impossible to fix the exact eastern and southern boundary of *ordinoides* for the reason that it intergrades with *elegans* throughout the entire length of its range." From this two inferences may be drawn: These species are either valid and the specimens are amenable to unquestionable separation, or our recognition of them as two distinct forms must give way to the fact that there is complete intergradation, hence they are not distinct and require to be united.

The initial step in the direction of attempting to ascertain the final status of these two species was taken when 50 specimens of *T. ordinoides* from the Sausalito Peninsula were subjected to a critical examination.² The present contribution offers the data obtained from a similar number of specimens captured in Golden Gate Park on the San Francisco peninsula along with a comparison of the two series. As a further illustration of the need of these and similar studies there is appended the record of 80 specimens from twelve different localities.

METHODS.

In order that the data be correctly coordinated it is necessary to ascertain the highest scale row count that obtains in the genus. In *Thamnophis* the maximum is 23 rows. The paired rows are designated by permanent numbers from I to XI and the median by M, the count being made from without inward.

When the number of scale rows is decreased the sequence of suppression is constant and is as follows:

- 23 rows, V row suppressed, leaving,
- 21 rows, VI row suppressed, leaving,
- 19 rows, IV row suppressed, leaving,
- 17 rows, VII row suppressed, leaving,
- 15 rows, which are continued to the vent.

In dealing with a specimen in which the maximum count is 21 rows it is necessary to regard the V row as suppressed constructively. In enumerating the rows one must count I, II, III, IV (V suppressed), VI, VII, VIII, IX, X, XI, and the median. In such a specimen when the 21 rows are reduced posteriorly to 19 rows it will be found to be due to the termination of the fifth row in actual counting, but this row in terms of the generic count is the VI row and must be so recorded. When the 19 rows are decreased to 17 it is due to the ending of the fourth row in actual counting, which is also the IV row in the generic sense. When the 17 rows are reduced to 15 it is due to the loss of the fourth row in actual counting, which in this case is the VII row in terms of the generic count. Constant attention to these details is imperative.

¹ 1908, Bull. 61, U. S. National Museum, p. 149.

² 1914, Thompson, Proc. U. S. Nat. Mus., vol. 47, pp. 351-360.

VARIATION IN NUMBER OF DORSAL SCALE ROWS.

This species presents the widest range of variation in the number of scale rows that is to be found in the group. There are at present known in the genus eleven distinct scale formulae, ten of which have been recorded in *T. ordinoides*. These ten and the frequency of occurrence of the five combinations found in the San Francisco series are as follows:

	Per cent.		Per cent.
21-23-21-19-17	0	19-17-15-13.....	0
21-19-17.....	12	17-19-17-15.....	0
19-21-19-17.....	56	17-15.....	0
19-21-19-17-15.....	8	15-17-15.....	0
19-17.....	16	Asymmetrical.....	4
19-17-15.....	4		

The influence of sex upon the variations in this character is clearly demonstrated. Among the specimens that vary from what may be assumed to be the normal count of 19-21-19-17, those having the lower counts are prone to be males, and those with the higher counts are largely females. This increase in the number of scale rows in the female is associated with the need for the increase in the diameter of the abdominal cavity when carrying the young.

	Males.	Females.
	<i>Per cent.</i>	<i>Per cent.</i>
21-19-17	17	83
19-21-19-17....	39	61
19-21-19-17-15	50	50
19-17.....	86	14
19-17-15	67	33

The data showing the gastrostege levels on the right and left side of the body at which the added rows begin and the suppressed rows end may be learned from the following tables:

SCALE FORMULA 21-19-17.

This is next to the highest count recorded in the species. In this series it occurs in 12 per cent of the specimens of which 16 per cent are males.

U.S.N.M. No.	Sex.	Ventrals.	Scale rows.		
			21-VI.	19-IV.	17 cont.
52176.....	Male	158	68 76	92 92
53575.....	Female..	151	71, IV 74	98, VI 99
52190.....	...do....	153	70 70	92 95
52159.....	...do....	155	71 76	89 88
53545.....	...do....	155	65, IV 71	82, VI 87
53577.....	...do....	157	71 73	99 101

SCALE FORMULA 19—21—19—17.

This formula may be regarded as the normal for this immediate region. It is found in 56 per cent of the specimens; of these, 39 per cent are males.

MALE.

U.S.N.M. No.	Ventrals.	Scale rows.			
		19+VI.	21—VI.	19—IV.	17 cont.
52184.....	154	26 28	51 42	77 76
53583.....	157	32 30	51 58	78 78
52163.....	158	47 41	53 43	79 80
53573.....	158	27 25	60 65	83 86
52179.....	159	24 27	68 74	89 89
52180.....	159	24 37	60 60	86 86
52166.....	160	26 24	70 70	86 87
52174.....	160	27 24	62 63	82 83
53581.....	161	34 26	44 32	87 82
52181.....	162	32 32	69 62	86 86
52162.....	162	30 31	71 75	92 93

FEMALE.

The influence of sex on this character is evident. On the average the VI row tends to be longer in that it begins further forward and extends more to the rear. The IV row also ends further back. This lengthening of the scale rows is associated with the stouter body in the female.

U.S.N.M. No.	Ventrals.	Scale rows.			
		19+VI.	21—VI.	19—IV.	17 cont.
52170.....	147	32 32	50 50	76 76
53541.....	149	23 22	60 61	87 86
52161.....	150	32 44	55 53	82 82
52175.....	150	24 23	61 62	80 82
52171.....	151	31 36	40 55	79 79
52160.....	152	13 13	68 71	90 93
52168.....	153	26 26	58 58	83 84
52165.....	153	30 41	51 51	81 79
53576.....	153	22 23	68 67	88 90
53542.....	153	26 28	54 53	82 86
53585.....	154	24 24	65 68	88 92
52167.....	155	29 27	64 65	89 87
52164.....	155	22 20	62 64	85 92
52185.....	156	21 13	68 71	91 95
52158.....	157	27 23	63 71	89 88
52178.....	160	27 26	67 59	88 89
52186.....	166	12 15	75 75	91 92

It will be noted that No. 52170 is bilaterally symmetrical. It is comparatively rare for the scale rows to arise and to be suppressed at exactly the same gastrostege levels.

SCALE FORMULA 19—21—19—17—15.

This is not a common variation. It occurs in 8 per cent of the series, and of these 50 per cent are males. In specimens with this formula there are five zones on the body, each with a different count. In two examples the VII rows reappeared just anterior to the vent, thereby increasing the count to 17 in this zone.

U.S.N.M. No.	Sex.	Ventrals.	Scale rows.								
			19+VI.		21—VI.		19—IV.	17—VII.	15 cont.		
52182....	Male.....	156	30	30	54	51	77	77	121	124
52173....	do.....	156	30	28	51	62	84	86	122	129
53580....	Female..	158	28	35	49	50	83	81	130	141
52172....	do.....	151	31	43	57	52	81	79	135	135

SCALE FORMULA 19—17.

This formula is found next in frequency to the normal. It occurs in 14 per cent of the series, and of these 86 per cent are males. Two specimens in this table exhibited bilateral symmetry, the rows terminating on each side at exactly the same level.

U.S.N.M. No.	Sex.	Ventrals.	Scale rows.		
			19—IV.	17 cont.	
52169.....	Male.....	152	75	75
53578.....	do.....	156	81	81
53579.....	do.....	157	80	80
53584.....	do.....	159	78	80
53543.....	do.....	159	78	79
53544.....	do.....	159	81	82
52187.....	Female....	152	83	83

SCALE FORMULA 19—17—15.

This is the least frequent count in the series. It occurs in but 6 per cent, of which 66 per cent are males.

In both these specimens the VII rows reappear a short distance anterior to the vent.

U.S.N.M. No.	Sex.	Ventrals.	Scale rows.				
			19—IV.	17—VII.	15 cont.		
52177.....	Male.....	153	77	78	135	138
53582.....	do.....	156	74	76	112	118
52188.....	Female....	156	78	79	135	137

BILATERALLY ASYMMETRICAL.

U.S.N.M. No.	Sex.	Ventrals.	Scale rows.			
			19+VI.	21-VI.	19-IV.	17 cont.
52183.....	Male.....	158	<i>rt.</i> 33	<i>lt.</i> 51	81 82
52189.....	Female....	159	{28 41	{34 43	} 87 87

It will be noted that in both these cases the asymmetry consists in an absence of the VI row on the left side of the body. It is this row that is normally added to the 19 row zone anteriorly and raises the count to 21 rows. In the male the VI row exists on the right side between the level of the thirty-third and the fifty-first gastrostege. In the female this row is also present only on the right side; it differs in being interrupted, one series extending from the twenty-eighth to the thirty-fourth, and the other from the forty-first to the forty-third gastrostege.

There is one important question bearing on these dermal characters that requires to be settled, and that is whether the number and extent of the scale rows are definitely fixed from birth and remain the same throughout the life of the individual, or if their number and extent may be increased as a sequel to abundant food, corpulency, pregnancy, and old age. The fact seems to be, that when a series of these garter snakes from the same locality is examined, a suspiciously large percentage of the low-scale row counts are found in the young and half-grown examples.


OTHER VARIATIONS.

All of the dermal characters show more or less variation. The scale rows in the zones of transition from the neck to the body and between the body and the tail are too irregular and complicated to permit the making of satisfactory records. There is some variation in the size and shape of the rostral, frontal, and parietal shields that depends upon the age of the specimen. The numerical variations in other characters are very important to ascertain, for they confirm beyond dispute, as Doctor Ruthven has demonstrated, the genetic relationships among the species.

VARIATION IN GASTROSTEGES.

Attention is drawn to the fact that the minimum and maximum counts have been found in the female specimens. In the males the range of variation is but 55 per cent of the total and centers nearly around the mean for the series.

Diagram showing the variation in the number of gastrosteges in specimens from San Francisco.




Females.

Males.

Range of variation in 50 specimens.....	147-166
Males, 23 specimens.....	152-162
Females, 27 specimens.....	147-166

Diagram showing the variation in the number of gastrosteges in Sausalito specimens.



Female.

Male.

Range of variations in 59 specimens.....	145-164
Males, 34 specimens.....	150-164
Female, 25 specimens.....	145-161

An occasional specimen is seen in which some of the gastrosteges are incomplete in that they fail to reach across the body. These incomplete shields usually extend nearly to the median line. Where there are several on one side the asymmetry is often compensated for by there being nearly an equal number on the opposite half of the body.

One specimen, a female, from San Francisco, Cat. No. 53574, U.S.N.M. (Orig. No. T-136), that is not included in the series, has nine incomplete gastrosteges on the right and none on the left side of the body. It is further abnormal in having a low gastrostege count of 149 (right), a scale formula of 21-19-17-15, and eight infralabial shields on the left side.

VARIATION IN ANAL.


The normal condition is for this shield to be entire. There are no specimens in this series in which it is divided.

VARIATION IN UROSTEGES.

These plates are normally paired. An occasional specimen may be found in which from one to two of the urosteges remain entire. When this occurs, these undivided shields are regularly situated near the base of the tail. All showing this variation are males.

Paired throughout, 94 per cent; 1 to 2 entire, 6 per cent.

Diagram showing the variation in the number of urosteges in specimens from San Francisco.




Range of variation in 42 specimens:

Males, 18 specimens	74—85
Females, 24 specimens	66—80

Further along when the table comparing these specimens with the series from Sausalito is reviewed; it will be noted that in both localities the range in the number of urosteges is from 66 to 85. These bare figures fail to bring out one salient feature—the higher average that exists in the Sausalito set. This difference may be seen at a glance if the preceding and following diagrams be compared.

Diagram showing the variation in the number of urosteges in specimens from Sausalito.



Range of variation in 32 specimens:

Males, 19 specimens	76—85
Females, 13 specimens	66—85

Over one-half of these from Sausalito have the tail docked, while less than one-fifth of the snakes from Golden Gate Park are similarly crippled. In grazing land this is frequently due to their being trodden upon by cattle. Where field rodents are plentiful the tail is often bitten off and devoured by these animals.

VARIATION IN PREOCULAR.

The normal condition is a single preocular. Where two exist it is due to the fragmentation of the lower one-third of the normal shield. Where three are found it is due to a middle plate which has become separated from the anterior superior shield. In the majority of specimens the lower portion of the preocular is of a lighter tint and frequently is indented at the margins. In the specimens showing variation in this character 25 per cent are males.

	Per cent.
1 normal.....	92
1-2 asymmetrical.....	4
2 bilateral.....	2
2-3 asymmetrical.....	2

VARIATION IN POSTOCULARS.

In this series the variation in the postocular shields is limited, much more so than in the Sausalito specimens. When the number is reduced to two it is due to the fusing of the normal middle and inferior shields; when increased to four it is due to the dividing of the inferior shield. In the specimens showing variation in this character 50 per cent are males.

	Per cent.
3 normal.....	96
3-2 asymmetrical.....	2
3-4 asymmetrical.....	2

VARIATION IN ANTERIOR TEMPORAL.

This shield is subject to but slight variation. When there are two it is due to a separation of the antero-external angle of the parietal. This part of the parietal is at times dented or partially incised. The single specimen showing an aberration in this character was a male.

	Per cent.
1 normal.....	98
1-2 asymmetrical.....	2

VARIATION IN POSTERIOR TEMPORALS.

These shields exhibit considerable irregularity as to shape and size. Their number may be increased to three; very rarely they become fused into one. The abnormal specimens were all females.

	Per cent.
2 normal.....	92
2-3 asymmetrical.....	4
3 bilateral.....	4

VARIATION IN SUPRALABIALS.

The normal is 8 supralabials with the fourth and fifth bordering the orbit. When the number is reduced to 7 it is due to the fusing of the normal second and third shields (66 per cent), or the sixth and seventh shields (33 per cent). None of the specimens in this set showed the increase to 9 supralabials which is known to occur at times in this species. In the specimens showing variation in this character 20 per cent are males.

	Per cent.
8 normal.....	90
8-7 asymmetrical.....	10

VARIATION IN INFRALABIALS.

The normal count is 10 infralabials. When the number is decreased to 9 it is due to the fusing of the normal third and fourth shields (82 per cent), or the normal seventh and eighth (18 per cent). When the number is further reduced to 8 it is due to the fusing of these same pairs—namely, the third and fourth and the seventh and

eighth. None in this set showed the increase to 11 infralabials which is known to occur in specimens from this part of the State. Among the specimens showing variation in this character 27 per cent are males.

	Per cent.
10 normal.....	78
10-9 asymmetrical.....	14
9 symmetrical.....	2
9-8 asymmetrical.....	4
8 symmetrical.....	2

One specimen, Cat No. 53574, U.S.N.M. (Orig. No. T-136), a female, shows marked asymmetry, there being 10 shields on one and 8 shields on the other side. This is the specimen in which many of the gastrosteges are incomplete.

VARIATION IN GENEIALS.

The anterior and the posterior pairs may be equal in length, one pair may be longer or shorter than the other, and it often occurs that there is much discrepancy between the length of the right and left posterior shields. In spite of these facts, this purely book character—the relative length of the anterior in terms of the posterior pair of geneials—has been made use of in the attempt to separate this complex Pacific Coast garter snake into endless species.

VARIATION IN GULARS.

The gular shields that lie between the posterior geneials and the first gastrostegae are irregularly paired anteriorly and azygos posteriorly. The normal count is two paired and two unpaired shields. The variation ranges from four paired and one unpaired to one unpaired and four azygos shields.

SUMMARY OF VARIATIONS.

The following may be assumed to be the normal conditions:

	Per cent.
Scale rows, 19-21-19-17.....	56
Preocular, 1.....	90
Postoculars, 3.....	96
Anterior temporal, 1.....	98
Posterior temporals, 2.....	92
Supralabials, 8.....	90
Infralabials, 10.....	78
Anal, entire.....	100
Urosteges, paired.....	94

The following table shows the percentage of normal individuals and the percentage of those that are abnormal in one or more characters:

	Per cent.
Normal in all characters.....	38
Abnormal in one character.....	30
Abnormal in two characters.....	22
Abnormal in three characters.....	8
Abnormal in four characters.....	2

Tabulated summary.

U.S.N.M. No.	Sex.	Scales.	Oculars.		Temporals.		Labials.		Gast.	An.	Urost.
			Pre.	Post.	Ant.	Post.	Supra.	Infra.			
52176	Male	21-19-17							158		81
52190	Female.	21-19-17				3			153		74
52159	do.	21-19-17	2					9-10	155		?
53575	do.	21-19-17				3			151		73
53577	do.	21-19-17							157		68
53545	do.	21-19-17		3-4					155		75
52184	Male	19-21-19-17							154		78
52163	do.	19-21-19-17							158		?
52179	do.	19-21-19-17							159		76 (III)
52180	do.	19-21-19-17							159		78
52166	do.	19-21-19-17							160		79
52174	do.	19-21-19-17							160		78
52181	do.	19-21-19-17						10-9	162		? (II-IV)
52162	do.	19-21-19-17							162		?
53573	do.	19-21-19-17							158		?
53583	do.	19-21-19-17						9	157		?
53581	do.	19-21-19-17							161		77
52170	Female.	19-21-19-17				3-2			147		82
52161	do.	19-21-19-17							150		66
52175	do.	19-21-19-17							150		?
52171	do.	19-21-19-17							151		70
52160	do.	19-21-19-17	2-3		2-1				152		67
52168	do.	19-21-19-17							153		74
52165	do.	19-21-19-17						10-9	153		70
52167	do.	19-21-19-17	2-1						155		?
52164	do.	19-21-19-17							155		68
52185	do.	19-21-19-17							156		67
52158	do.	19-21-19-17							156		72
52178	do.	19-21-19-17							157		?
52178	do.	19-21-19-17							160		71
52186	do.	19-21-19-17							166		74
53576	do.	19-21-19-17							153		71
53585	do.	19-21-19-17				3-2		9-10	154		68
53541	do.	19-21-19-17							149		68
53542	do.	19-21-19-17		4-3					153		68
52182	Male	19-21-19-17-15					7-8		156		85
52173	do.	19-21-19-17-15							156		82
52172	Female.	19-21-19-17-15							152		72
53580	do.	19-21-19-17-15							151		70
52169	Male	19-17					8-7	10-9	152		74
53584	do.	19-17						9-8	159		70
53579	do.	19-17	1-2				7-8	8-9	157		79
53543	do.	19-17							159		80
53544	do.	19-17		3-2				9-10	159		74
53578	do.	19-17							156		79 (III)
52187	Female.	19-17							152		80
52177	Male	19-17-15						9-10	153		83
53582	do.	19-17-15					8-7	8	156		85
52188	Female.	19-17-15							156		79
52183	Male	Asymmetrical							158		80
52189	Female.	Asymmetrical					8-7		159		79

For the sake of clearness the normal records are indicated by dashes.

This table brings to light an interesting condition: Specimens that have the normal scale row count of 19-21-19-17 are prone to be normal in the remaining characters in that only 33 per cent of them present any variations; on the other hand, in those having an abnormal scale row count 66 per cent, possess other variations from the normal. This is an analogous condition to that shown in the study of degenerates among the human species. Individuals are much more liable to have several stigmata than but a single earmark of faulty heredity.

INFLUENCE OF SEX ON VARIATION.

In zoology it is a fairly well established point that in a given species the range of variation is regularly greater in a series of males than in a series of female specimens. In the anthropoid apes and in the

human species this is particularly noticeable, the female remaining nearer the infantile and generalized type than the male.

When the influence of sex is examined for *T. ordinoides*, it is found that the opposite condition prevails. The following list shows that a greater number of variations occur among the female specimens:

	Male.	Female.
Abnormal as to—	<i>Per cent.</i>	<i>Per cent.</i>
Scale rows.....	45	55
Preocular.....	25	75
Post oculars.....	33	67
Anterior temporal.....	100	0
Posterior temporal.....	0	100
Supralabials.....	20	80
Infralabials.....	27	73

VARIATION IN COLOR.

In the young the ground color is almost invariably a dark olive; the dorsal and lateral rows of spots are large, regular, and sharply defined; the median stripe is pale yellow; the underside is greenish grey; no specimens are seen with even a trace of red.

In the adults the ground color varies in different examples, there being many hues of dull brick red, olive brown, and dark olive; the dorsal and lateral spots are clearly marked, in some the dorsal series is partly fused; the median stripe is sharply defined, varying from pale yellow to dark orange, and when of the latter color it is at times dotted with salmon; the lateral stripe is yellowish or greenish grey, in some this line is also dotted with salmon; the underside is usually a uniform bluish or greenish grey, and in a few specimens there are a few irregular reddish spots.

The most striking chromatic character in this set is the absence of any examples having the striped pattern, those in which the ground color is a solid dark brown, without spots or red, and with the median and lateral stripes bright and sharply defined. In having but the one color pattern this set from Golden Gate Park is distinctive. Series from over a score of different localities have been studied and each set regularly contained from two to three separate designs.

FOOD.

The food was found to consist almost entirely of slugs, of the family *Arionidæ*. Two of the largest snakes had eaten small rodents, and several had remains of salamanders, *Batrachoseps*, and *Autodax* in the stomach.

The garden slugs in and about San Francisco are abundant and are most destructive to small and tender cultivated plants. As the Bay Region has not proved favorable for the establishing of toad colonies,

the garter snakes assume a rôle of particular economic interest. They should become a special object of care and protection on the part of florists and gardeners.

COMPARISON BETWEEN SAN FRANCISCO AND SAUSALITO SERIES.

There is presented for comparison in the following parallel columns a summary of the variations in the series from the two localities on either side of the Golden Gate:

	San Francisco.	Sausalito.
	<i>Per cent.</i>	<i>Per cent.</i>
Scale rows:		
21—23—21—19—17		
21—19—17	12	6
19—21—19—17	56	66
19—21—19—17—15	8	
19—17	14	10
19—17—15	6	12
19—17—15—13		
17—19—17—15		
17—15		
15—17—15		
Asymmetrical	4	4
Preocular:		
1 normal	92	86
1-2 asymmetrical	4	10
2 bilateral	2	4
2-3 asymmetrical	2	
Postoculars:		
3 normal	94	80
3-2 asymmetrical	2	10
3-4 asymmetrical	4	6
2 bilateral		2
4 bilateral		2
Anterior temporal:		
1 normal	98	94
1-2 asymmetrical	2	4
2 bilateral		2
Posterior temporals:		
2 normal	92	82
2-3 symmetrical	4	12
3 bilateral	4	6
Supralabials:		
8 normal	90	94
8-7 asymmetrical	10	4
8-9 asymmetrical		2
Infralabials:		
10 normal	78	74
10-11 asymmetrical		6
10-9 asymmetrical	14	10
10-8 asymmetrical		2
9 bilateral	2	4
9-8 asymmetrical	4	4
8 bilateral	2	
Anal:		
Entire normal	100	90
Divided		10
Urostegees:		
Paired normal	94	72
1 to 4 entire	6	28
Gastrostegees (plates)	147-166	145-164
Urostegees (plates)	66-85	66-85

The following table accentuates the difference in the two series, the San Francisco group containing fewer specimens that vary from what may be assumed to be the normal for the region.

	San Francisco.	Sausalito.
	<i>Per cent.</i>	<i>Per cent.</i>
Normal in all characters.....	38	14
Abnormal in one character.....	30	44
Abnormal in two characters.....	22	20
Abnormal in three characters.....	8	14
Abnormal in four characters.....	2	6
Abnormal in five characters.....	2

The color characters of the two sets differ to a marked degree. All the specimens captured or seen in Golden Gate Park have the pronounced spotted pattern. Of those from the Sausalito peninsula about two-thirds of the examples are similarly marked, though in darker tones. The remaining one-third have the striped pattern; in these the ground is a dark brown, the dorsal and lateral spots are absent, the median and lateral stripes are distinct, and are without any red on the underside.

GEOGRAPHIC VARIATION.

In the Memoir of Doctor Ruthven it was pointed out that the specimens of this species became smaller, the head shields reduced in number, and the scale rows less numerous as the range extended to the north. The various sets presented in the following tables confirm these findings. The one additional fact brought out is that these dwarfed members of the species are prone to be found in localities having lower temperatures, rather than merely a higher latitude.

MERCED COUNTY, CALIFORNIA.

Scale formulas 21—23—21—19—17
21—19—17

Museum. Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.					Gulars.		Pro.	Pto.	Supral.	Infral.				
						21+V.	23-V.	21-VI.	19-IV.	17 cont.	Prs.	Azy.								
13635.....	Los Baños.....	♂+♀	152	ε	71	25	25	56	80	83	129	125	1	4	1	3	ε	10	
13636.....	do.....	♂+♀	158	ε	77	29	28	53	39	81	82	116	122	1	3	1	2	ε	9
13637.....	do.....	154	(?)	27	28	63	60	81	85	112	115	2	3	3	ε	10	
13638.....	do.....	156	ε	69	78	81	98	98	11-10	
13640.....	do.....	153	ε	72	76	80	93	97	11	
36071.....	do.....	154	ε	(?)	47	40	49	48	83	83	104	105	1	11-9	
Univ. Cal. 5428.	Gadwell.....	154	ε	(?)	28	30	60	62	81	82	114	111	1	4	1	3	ε	11

The color of the specimens from Los Baños is of one pattern: The ground a uniform light-brownish olive; faint trace of small dorsal spots; these tend to encroach upon the median stripe; the median stripe indicated by a paler tint than the ground color; below no spots.

MONTEREY COUNTY, CALIFORNIA.

Scale formulas 19-21-19-17
19-17
19-17-15

Museum Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.					15 cont.	Frs. Gulars. Azy.	Pro.	Pto.	Supral.	Infral.			
						19+VI.	21-VI.	19-IV.	17-VII.										
13756.....	Carmel.....	♂	157	♂	(?)	36	39	54	55	79	81	1	3	1	3	8	10
13760.....	do.....	♂	152	♂	(?)	39	42	56	55	85	85	1	3	1	3	8	10
13761.....	do.....	♂	149	♂	(?)	81	81	147	142	1	3	8	10
13764.....	Monterey.....	♂	147	♂	76	78	80	1	3	1	3	8	10
13765.....	do.....	♂	155	♂	(?)	84	85	131	143	1	3	8	10
13757.....	Carmel.....	♂	142	♂	65	62	56	97	99	1	3	8	10
13758.....	do.....	♂	155	♂	68	33	33	71	74	89	91	1	3	1	3	8	10
13759.....	do.....	♂	146	♂	69	17	20	66	62	85	87	1	3	1	3	8	10

The color of this set is as follows: The ground varies from a very dark red-brown to a dark brown; the dorsal and lateral rows of spots absent or when visible ranging from a mere shade darker than the ground color to sharply defined marks; the median stripe uniform canary yellow or mixed yellow and salmon; the lateral stripe yellowish green or nearly solid salmon.

SAN MATEO COUNTY, CALIFORNIA.

Scale formulas 19-17
19-17-15
17-15

Museum U.S.N.M.	Locality.	Sex.	G.	A.	U.	Scale rows.			Frs. Gulars. Azy.	Pro.	Pto.	Supral.	Infral.
						19-IV.	17-VII.	15 cont.					
50307.....	Half Moon Bay.....	♂	157	♂	83	82	82	1	3	3	9-10
53538.....	San Andreas.....	♂	151	♂	81	81	81	1	3	3	10
53540.....	do.....	♂	144	♂	75	74	75	133	126	1	3	3	10
53539.....	do.....	♂	141	♂	78	79	2	3	7	8-9

One of the most interesting garter snakes on record is Cat. No. 53539, U.S.N.M. (Orig. No. K. 103), collected by H. C. Kellers, United States Navy. It is the only specimen known from south of Eureka with the minimum formula of 17-15. Associated with this low scale count is a reduction in the supralabial shields to 7, a condition also found as far as this species is concerned only in the north.

ALAMEDA COUNTY, CALIFORNIA.

Scale formulas 21-19-17
19-21-19-17
19-17

Museum No. U.S.N.M.	Locality.	Sex.	G.	A.	U.	Scale rows.				Gulars.		Pro.	Pto.	Supral.	Intral.							
						19+VI.	21-VI.	19-IV.	17 cont.	Pts.	Azy.											
52194.....	Alameda.....	♂	162	ε	(?)	28	33	63	67	90	93	1	3	3	2	3	3	8	8	10	
52195.....	do.....	♂	161	ε	ε	36	28	29	63	75	88	89	1	3	3	1	3	3	3	8	10
52196.....	do.....	♂	162	ε	ε	84	81	83	105	104	1	4	1	1	3	3	3	8	10	
Cal. Ac. Sci. 13223.	San Leandra.....	♂	162	ε	(?)	28	27	61	66	89	85	2	2	1	1	2	3	8	8	9	
Univ. Cal. 5417	Berkeley.....	♂	158	ε	ε	80	23	26	67	66	90	91	2	2	2	2	3	3	3	8	10
5418.....	do.....	♂	163	ε	ε	82	25	27	69	69	88	89	2	2	2	1	3	3	3	8	10
5419.....	do.....	♂	158	ε	ε	78	81	83	3	1	1	1	3	3	3	8	10	

The prevailing arrangement of color in this set is as follows: Ground dark olive brown; dorsal and lateral spots usually present in the young and absent in the adult; median stripe pale yellow in the young, orange or vermillion in the adult; lateral stripe light olive gray, with or without salmon; below uniform greenish gray, or with salmon spots.

SONOMA COUNTY, CALIFORNIA.

Scale formulas 19-17
19-17-15

Museum Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.			Gulars.		Pro.	Pto.	Supral.	Intral.			
						19-IV.	17-VII.	15 cont.	Pts.	Azy.							
28021.....	Seaggs.....	♂	158	ε	90	65	86	125	123	1	3	1	3	3	10
28025.....	Springs.....	♂	160	ε	86	79	83	138	133	1	2	1	3	3	10
28029.....	do.....	♂	162	ε	87	88	91	1	3	1	3	3	10
28019.....	do.....	♂	150	ε	71	80	82	2	2	1	2	3	10
28024.....	do.....	♂	150	ε	(?)	77	79	1	3	1	2	3	9 V + VI

The color pattern of this set is as follows: The ground is a dark olive brown; the dorsal and lateral rows of spots are usually absent or but faintly defined; the median stripe bright yellow; the lateral stripe yellowish gray; below uniform olive gray.

MENDOCINO COUNTY, CALIFORNIA.

Scale formulas 19-21-19-17
19-17-15

Museum Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.					Gulars.		Pro.	Pto.	Supral.	Infral.
						19+VI.	21-VI.	19-IV.	17-VII.	15 cont.	Prs.	Azy.				
28302.....	Comptche	♂	152	ε	75	81 84	- III 128 133	1 3	1	3	∞	10	
28303.....do.....	♂	154	ε	85	83 83	141 141	1 3	1	3	∞	10	
28304.....do.....	♂	153	ε	76	83 83	103 118	1 3	1	3	∞	9 III+IV	
28306.....do.....	♂	157	ε	(?)	23	26	51 61	84 85	1 3	1	3	∞	10	
28308.....do.....	♂	148	ε	(?)	77 79	130 137	1 4	1	3	∞	9 III+IV	
28307.....do.....	♂	149	ε	(?)	13	26	68 63	89 90	2 2	1	3	∞	10	

The color design of this set is as follows: The ground varies from olive brown to olive gray; dorsal and lateral spots when present small and indistinct; the median stripe distinct and yellowish or merely indicated as a lighter tint than the ground color; lateral stripe usually poorly defined.

DEL NORTE COUNTY, CALIFORNIA.

Scale formulas 19-21-19-17
19-17

Museum Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.				Gular.		Pro.	Pto.	Supral.	Infral.
						19+VI.	21-VI.	19-IV.	17 cont.	Prs.	Azy.				
29076	Regua.....	♂	156	ε	79 (II-III)	29 31	58 54	78 82	1 3	1	3	∞	10	
29081do.....	♂	155	ε	80	77 79	1	3	∞	9	
29073do.....	♂	157	ε	(?)	28 26	60 60	88 89	1	3	∞	8 II+III	
29080do.....	♂	147	ε	(?)	74 75	1	3	∞	VIII+IX	
29082do.....	♂	153½	ε	65	84 86	1	3	∞	9	

The colors are dull; ground dark brown; spots absent or indicated by darker tones; median stripe dull orange, lateral yellowish grey. This set begins to show the marked reduction in size that occurs to the north; the largest, No. 29081, measures 640 mm.; tail, 161; head, from tip of snout to behind angle of jaw, 22.5; snout, 7.2; eye, 3.5.

CURRY COUNTY, OREGON.

Scale formulas 17-19-17-15
17-15

Museum Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.				Gulas.				Supral.	Infral.		
						17+IV.	19-IV.	17-VII.	15 cont.	Pfs.	Azy.	Pto.	Pto.				
29272	Harbor.....	♂	139	€	70	27 29	40 29	27 33	41 48	85 81	1	3	1	3	7	8 III+IV
29274do.....	♂	146	€	77	77	82	1	3	1	3	7	9
29275do.....	♂	148	€	77	76	79	1	3	1	3	7	9
29268do.....	♂	153	€	77	74	75	2	3	1	3	7	9
29273do.....	♂	140	€	80	80	1	1	1	3	8 (II)	9
29277do.....	♂	140	€	77	77	1	1	1	3	8 (II)	9
29278do.....	♂	1	€	84	83	1	1	1	3	7	8 (III+IV)
29269do.....	♂	2	€	66	73	75	1	1	1	3	7	8 (III+IV)
29276do.....	♂	2	€	85	87	1	3	1	3	7	8 rt, VII+VIII 8 lt, VIII+IX
29271do.....	♂	5	€	59	45	56	97	101	1	3	1	3	7	9
29279do.....	♂	5	€	64	79	81	1	3	1	3	7	8 III+IV
29270do.....	♂	7	€	70	39	39	74	76	2	1	1	3	7	8 III+IV

The color pattern of this set is quite simple; ground light olive brown, or brown; spots usually absent, or indicated as darker shades of brown; stripes indistinct or sharply marked, median brick red, ateral faintly paler than ground color; below bluish grey, with or without red.

It will be noted that the predominant type of scale formula is the one with 17 rows anteriorly and 15 posteriorly, this occurring in 75 per cent of the specimens. Those in the 17-19-17-15 class are either asymmetrical or have very few scales in the added IV row. All are of small size, the average being about 400 mm. in length.

In looking over the tables which will follow of series from localities to the north, it will be observed that the specimens are larger and the increased scale formula of 17-19-17-15 again prevails as it does to the immediate south. Incidentally, southwestern Oregon is one of the coldest portions of the Pacific coast region, and the localities mentioned to the north are actually much warmer. It becomes evident that the reduced size and lower scale counts in *T. ordinoides* are correlated with colder climate rather than mere extension of the range to more northern latitudes.

TILLAMOOK COUNTY, OREGON

Scale formulas 17-19-17-15
17-15

Museum Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.					Gulars			Supral.		Infral.						
						17+IV.	19-IV.	17-VII.	15 cont.	Prs.	Azy.	Pro.										
															Pto.							
29688	Tillamook	♂	145	€	60	27	31	57	57	87	85	2	2	1	3*	7	8	II	10 VIII III	9 VIII	
29693do.....	♂	146	€	(?)	90	87	1	2	1	3	7	8	II	8	
29690do.....	♂	149	€	71	29	27	52	50	93	94	1	2	2	3	7	8	II	8	
29692do.....	♂	142	€	59	26	37	56	60	101	100	1	1	2	3	7	8	II	9 III	8	
29694do.....	♂	146	€	56	26	26	65	63	120	120	1	2	1	3	6V+VI	7	8	II	8
29697do.....	♂	150	€	(?)	90	89	2	2	1	3	7	8	II	8	
29695do.....	♂	152	€	(?)	38	43	93	95	1	2	1	3	7	8	II	8	

This set has the normal scheme for the northern limits; ground brown; spots absent or small; median stripe orange or yellowish green. The largest (No. 29694) measures 496 mm.

CHEHALIS COUNTY, WASHINGTON.

Scale formulas 17-19-17-15
17-15

Museum Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.					Gulars			Supral.		Infral.					
						17+IV.	19-IV.	17-VII.	15 cont.	Prs.	Azy.	Pro.									
																	Pto.				
29940	Melbourne.	♂	139	€	(?)	30	27	53	53	82	78	1	2	1	3	6	III+IV	7	9 VII	8
29938do.....	♂	143	€	63	30	22	58	57	90	87	2	2	3	7	7	7	9 VII	8
29930do.....	♂	145	€	67	29	36	51	54	98	96	1	1	3	7	7	7	9 VII	8
29932do.....	♂	147	€	63	30	30	55	55	89	88	1	1	3	7	7	7	9 VII	8
29937do.....	♂	147	€	61	99	97	1	1	3	7	7	7	9 VII	8
24101	Montesano.	♂	150	€	(?)	86	90	0	4	1	3	7	7	7	9 VII	8
29931	Melbourne.	♂	141	€	(?)	23	24	63	60	85	92	2	1	3	7	7	7	9 VII	8
29935do.....	♂	141	€	(?)	27	26	66	63	90	90	1	1	3	7	7	7	9 VII	8
29936do.....	♂	143	€	(?)	26	26	66	55	96	88	1	1	3	7	7	7	9 VII	8
29933do.....	♂	144	€	60	41	31	41	31	91	83	1	3	7	6	III+IV	7	9 VII	8
29939do.....	♀	144	€	63	20	13	75	77	122	125	1	3	7	9	II	8	9	VII
24102	Montesano.	♂	145	€	59	25	23	70	76	121	122	1	2	1	3	7	7	7	7 VII+VIII	8
29934	Melbourne.	♂	146	€	57	23	23	56	52	84	83	1	3	7	7	7	7	7 VII+VIII	8
24103	Montesano.	♀	148	€	59	26	28	69	70	115	110	1	3	7	7	8,	7	III+IV	8

The color pattern of these northern specimens is generally dull, not pronounced; ground olive grey; dorsal and lateral spots small and faint; median stripe barely a shade paler than the ground color, in others sharply defined.

PACIFIC COUNTY, WASHINGTON.

Scale formulas 17-19-17-15
17-15

Museum No. Cal. Ac. Sci.	Locality.	Sex.	G.	A.	U.	Scale rows.				Gulars				Infral.						
						17+IV.	19-IV.	17-VI.	15 cont.	Prs.	Axy.	Pro.	Pto.		Supral.					
29925	Trap Creek	♂	144	ε	(?)	22	22	58	54	107	103	1	3	2	3	7	9 VII	8	
29924do	♂	146	ε	72	35	31	54	56	113	116	1	3	3	3	7	8	8	
29923do	♂	146	ε	60	108	106	1	3	1	3	3	8	8	
29926do	♂	146	ε	(?)	+V	25	26	66	69	94	100	3	0	1-2	3	7	9 VII	8

This set has markedly somber hues; the ground dark brown; dorsal and lateral spots indicated by darker shades; median stripe narrow, pale yellow; lateral stripe distinct; below dark grey. The measurements of the largest specimen, No. 29923, are as follows: Total length, 516 mm.; tail, 115; head, 17; snout, 4.8; eye, 2.6 mm.

SUMMARY OF SCALE FORMULAE IN THAMNOPHIS ORDINOIDES.

A tabulated summary of the occurrence of the different scale formulae in these various sets will show at a glance the prevailing types for each locality; to facilitate comparison they are given as percentages.

Scale formulas.	California.									Oregon.		Washington.		
	Merced.	Monterey.	San Mateo.	Alameda.	San Francisco.	Sausalito.	Marin.	Sonoma.	Mendocino.	Del Norte.	Curry.	Tillamook.	Chehalis.	Pacific.
21-23-21-19-17.....	72
21-19-17.....	28	14	12	6	33
19-21-19-17.....	50	71	56	66	33	40
19-21-19-17-15.....	8
19-17.....	12	50	14	16	10	63	60
19-17-15.....	37	25	4	12	66	37	66
19-17-15-13.....	2
17-19-17-15.....	16	57	82	21
17-15.....	25	76	28	18	79
15-17-15.....	1
Asymmetrical.....	4	4	8	14

This table brings out several instructive facts. The decrease in the number of rows in specimens from more northern regions is not a perfectly regular one, each locality having a decidedly different average. In the same locality there is never a single type. Where two or more formulas exist they are never equally divided among the specimens, one type regularly predominating to a marked degree.

NOTE ON SCALE FORMULA IN THAMNOPHIS PARIETALIS (SAY) AND THAMNOPHIS COCCINUS (HALLOWELL).

Another species of garter snake found in the same region with *T. ordinoides* is *Thamnophis parietalis* (Say) 1823; this form is noteworthy in possessing but one type of squamation, there being 19 rows anteriorly and 17 posteriorly. As this formula is a fairly prevalent one in *T. ordinoides* it will prove of interest to compare a small set of each species from the same locality. Specimens have been chosen from Skaggs Springs, Sonoma County, California, owing to this place being approximately in the middle of the area occupied by *T. ordinoides*.

SKAGGS SPRINGS, SONOMA COUNTY, CALIFORNIA.

T. ORDINOIDES.

Museum Cal. Ac. Sci.	Sex.	G.	Scale rows.	
			19-IV.	17 cent.
27940.....	Male.....	157	80	79
28029.....	..do.....	162	88	91
28024.....	Female..	150	77	79
28019.....	..do.....	150	80	82
27938.....	..do.....	160	84	85

T. PARIETALIS.

28026.....	Male.....	157	100	90
28023.....	Female..	151	89	88
28028.....	..do.....	154	80	82
28027.....	..do.....	157	84	85
28022.....	..do.....	157	90	87

This table shows that there is a tendency in *T. parietalis* for the 19-row zone to be continued a little further down the body. This, however, is merely a slight difference in the averages of two small sets.

The only remaining species found along the Northwest Coast is *T. coccinus* (Hallowell) 1852, this also presents the one scale count of 19-17.

TRASK, TILLAMOOK COUNTY, OREGON.

T. COCCINUS.

Museum Cal. Ac. Sci.	Sex.	G.	Scale rows.	
			19-IV.	17 cent.
29740.....	Male.....	167	87	91
29738.....	..do.....	162	88	89
29741.....	..do.....	160	83	84
29737.....	Female..	159	90	89
29734.....	..do.....	154	92	94

This table shows that the IV row does not extend quite so far down the body as in *parietalis*.

SUMMARY OF SCALAE FORMULAS IN GENUS THAMNOPHIS.

Doctor Ruthven recognizes 19 species of garter snakes in North America. Among these there are known 11 distinct combinations of scale row counts. The occurrence of the various combinations among the different species may be seen from the following table:

Species.	21-23-21-19-17.	21-19-17.	19-21-19-17.	19-21-19-17-15.	19-17.	19-17-15.	19-17-15-13.	17-19-17.	17-19-17-15	17-15.	15-17-15.
<i>Megalops</i>	*	*	*		*						
<i>Marciana</i>		*	*								
<i>Rudix</i>		*	*		*						
<i>Butleri</i>					*			*			
<i>Proximus</i>					*			*			
<i>Saurita</i>					*						
<i>Sackeni</i>					*						
<i>Angustirostris</i>	*	*	*								
<i>Elegans</i>		*	*			*					
<i>Ordinoides</i>	*	*	*	*	*	*	*		*	*	*
<i>Hammondi</i>		*	*								
<i>Melanogaster</i>			*								
<i>Scalaris</i>			*								
<i>Phenax</i>			*		*				*		
<i>Eques</i>					*				*		
<i>Sumichrasti</i>						*		*		*	
<i>Parietalis</i>					*						
<i>Coccinus</i>					*						
<i>Sirtalis</i>					*						

Inspection of this table shows that the formula of 19-17 is the most frequent, as it occurs in 12 species; also that there are eight species with only one scale row count, which in these cases is regularly one of the following three: 21-19-17, 19-21-19-17, or 19-17; and, finally, that in *Thamnophis ordinoides*, there are to be found twice as many combinations as are to be seen in any of the other species.

If *ordinoides* is a valid species it offers an example at variance with the prevailing condition in which it is the species with the widest ranges that vary the most, for *ordinoides* has one of the smallest areas of distribution in the genus. If the converse is true, which is most probable, and it ultimately proves to be inseparable from *elegans*, it will then conform to the rule in having a wide distribution and a large variability.

The handling of large series of garter snakes will inevitably result in bringing about a readjustment and broadening of one's concept of what may be implied by Wallace's phrase: "The variability of a species in the state of nature."