

PROCEEDINGS OF THE UNITED STATES NATIONAL MUSEUM



SMITHSONIAN INSTITUTION  
U. S. NATIONAL MUSEUM

---

---

Vol. 105

Washington : 1956

No. 3359

---

---

TYPE SPECIES OF THE GENERA AND SUBGENERA OF  
PARASITIC WASPS COMPRISING THE SUPERFAMILY  
PROCTOTRUPOIDEA (ORDER HYMENOPTERA)

---

By C. F. W. MUESEBECK<sup>1</sup> and LUELLE M. WALKLEY<sup>1</sup>

---

Introduction

The International Commission on Zoological Nomenclature emphasized the importance of genotype fixation by the adoption of an amendment to Article 25 of the Rules, effective Jan. 1, 1931, requiring that the description of any new genus be accompanied by "a definite, unambiguous designation of the type species." The steady and rapid expansion of our knowledge of insects has resulted in increasingly critical taxonomic study with constant adjustment of concepts at all levels of classification. Since the genus is one of the most significant of the various categories involved, it must be nicely defined in order that classifications may be developed on solid bases and the nomenclature will be as stable as possible. Essentially the definition of the genus must be based

---

<sup>1</sup> U. S. Department of Agriculture, Entomology Research Branch, Insect Identification and Parasite Introduction Section.

on its type species, a requirement which has often been ignored in the past, with the result that erroneous generic concepts have gradually become widely accepted in certain cases. Sometimes it is difficult and time consuming to ascertain the type species of a given genus. Much searching of the literature may be necessary, and the required literature is not always at hand. For these reasons the availability of genotype catalogs, prepared after exhaustive searching and thorough study, should have a stabilizing influence and should aid in the gradual betterment of our classifications.

Students of Hymenoptera are not in agreement concerning the limits of the Proctotrupoidea, and we recognize the difficulties in justifying the inclusion in this superfamily of such isolated groups as the Evaniidae, the Gasteruptiidae, and the Aulacidae. Nevertheless, for the purposes of this catalog we have considered it most practical to include them under the superfamily name Proctotrupoidea as was done in "Hymenoptera of America North of Mexico, Synoptic Catalog" (U. S. Dept. Agr., Agr. Monogr. 2, 1951). The present paper contains 735 generic and subgeneric names published through 1953 (a few later), including 72 that are errors or emendations of previously published names, and including also the following new names proposed to replace pre-occupied names: *Aulacomastus*, *Dorachia*, *Eccinetus*, *Nesopria*, and *Philoplanes*. Omitted are the following errors, most of them obviously typographical, that have appeared in the Zoological Record; in each case the year cited is the year covered by the particular volume of the Zoological Record and not the year of its publication.

ERROR	CORRECT NAME
<i>Acanthoscelis</i> , 1893, Ins. p. 215	<i>Acanthoscelio</i> Ashmead
<i>Aclysta</i> , 1913, Ins. p. 311	<i>Aclista</i> Foerster
<i>Alacinus</i> , 1908, Ins. p. 300	<i>Aulacinus</i> Westwood
<i>Anommatrum</i> , 1913, Ins. p. 312	<i>Anommatium</i> Foerster
<i>Aritomellus</i> , 1939, Ins. p. 303	<i>Atritomellus</i> Kieffer
<i>Ashmeadopsia</i> , 1914, Ins. p. 200	<i>Ashmeadopria</i> Kieffer
<i>Carina</i> , 1905, Ins. p. 222	<i>Carinia</i> Kieffer
<i>Catoteleia</i> , 1925, Ins. p. 336	<i>Calotelea</i> Westwood
<i>Christoteleia</i> , 1914, Ins. p. 200	<i>Chrestoteleia</i> Kieffer
<i>Cineta</i> , 1913, p. 313	<i>Cinetus</i> Jurine
<i>Coelogaster</i> , 1893, Ins. p. 216	<i>Coelopelta</i> Ashmead
<i>Erimicrops</i> , 1913, Ins. p. 320	<i>Trimicrops</i> Kieffer
<i>Heteroteleia</i> , 1920, Ins. p. 182	<i>Holoteleia</i> Kieffer
<i>Hoplotileia</i> , 1914, Ins. p. 201	<i>Hoploteleia</i> Ashmead
<i>Lagnyodes</i> , 1906, Ins. p. 308	<i>Lagnyodes</i> Foerster

<i>Loxotropa</i> , 1869, p. 321	<i>Loxotropa</i> Foerster
<i>Limnodyles</i> , 1902, Ins. p. 200	<i>Limnodytes</i> Marchal
<i>Loxotraps</i> , 1906, Ins. p. 308	<i>Loxotropa</i> Foerster
<i>Lygocerus</i> , 1878, Ins. p. 154	<i>Lygocerus</i> Foerster
<i>Mista</i> , 1921, Ins. p. 163	<i>Miota</i> Foerster
<i>Onyteleia</i> , 1910, Ins. p. 304	<i>Oxyteleia</i> Kieffer
<i>Parabalus</i> , 1910, Ins. p. 305	<i>Parabacus</i> Kieffer
<i>Paramegischia</i> , 1882, Ins. p. 139	<i>Pammegischia</i> Provancher
<i>Paramesus</i> , 1907, Ins. p. 288	<i>Paramesius</i> Westwood
<i>Paratrimorus</i> , 1910, Ins. p. 305	<i>Paratrimorus</i> Kieffer
<i>Phaenomorphus</i> , 1919, Ins. p. 167	<i>Phaenoserphus</i> Kieffer
<i>Procinctus</i> , 1914, Ins. p. 203	<i>Procinetus</i> Kieffer
<i>Prophanus</i> , 1923, Ins. p. 205	<i>Prophanurus</i> Kieffer
<i>Protimorus</i> , 1908, Ins. p. 304	<i>Protrimorus</i> Kieffer
<i>Psilonima</i> , 1913, Ins. p. 318	<i>Psilomma</i> Foerster
<i>Telemonus</i> , 1888, Ins. p. 201	<i>Telenomus</i> Haliday
<i>Teleus</i> , 1908, Ins. p. 305	<i>Teleas</i> Latreille
<i>Xestonatus</i> , 1926, Ins. p. 338	<i>Xestonotus</i> Foerster

We have seen the original publications for all the references cited. However, we have not succeeded in ascertaining the precise dates of publication of two papers by Kieffer that contain supposedly new generic names published elsewhere by him at about the same time. In *The Transactions of the Linnean Society of London*, series 2, Zoology, volume 15, pages 43–80, Kieffer published as new 11 generic names in the Proctotrupeoidea, some of which were apparently validated earlier in his generic classification of the Diapriidae, published in Wytzman's "Genera Insectorum" (fasc. 124). The latter work carries the date 1911, and in later publications Kieffer used 1911 in references to it. Neave's "Nomenclator Zoologicus," however, gives 1912 as the year of publication, and that is probably correct. It seems that the volume appeared very early in 1912, early enough to antedate the other paper mentioned above, which was published in March 1912. Accordingly we have treated the "Genera Insectorum" publication as the earlier.

There is also uncertainty as to the exact date of publication of a paper by Kieffer in *Annales de la Société Scientifique de Bruxelles* (vol. 33, 1909) which contained descriptions of five new species of *Paraclista*, a genus Kieffer described as new in André's "Species des Hyménoptères d'Europe & d'Algérie" (vol. 10, p. 476). The publication date given for that section of André's volume 10 is Sept. 15, 1909; and since the Smithsonian Institution's copy of the other paper is stamped as having been received Oct. 4, 1909, the two may have appeared at virtually the same time. In the absence of precise information concerning the date

of publication of the paper in *Annales de la Société Scientifique de Bruxelles*, and since Kieffer clearly intended it should be so, we are considering the paper in André's volume 10 to have priority.

We have not attempted to show all published generic synonymy but have given only that which is generally accepted and with which we agree. The type citations are shown in the original combinations even where the author designating a type used a different generic name.

Literature citations to periodicals are abbreviated but citations to separate works are given in full with one exception: André, *Species des Hyménoptères d'Europe & d'Algérie*. Kieffer's work published in it is cited so frequently that the citation is abbreviated as Kieffer *in* André, *Spec. Hym. Eur. Alg.*

### Superfamily PROTOTRUPOIDEA

#### **Abothropria** Kieffer

1913. Kieffer, *in* Alluaud and Jeannel, *Voyage . . . en Afrique Orientale*. *Insectes hyménoptères*, vol. 1, p. 23.

TYPE: *Abothropria nigra* Kieffer, 1913, by monotypy and original designation.

#### **Acanopsilus** Kieffer

1908. Kieffer, *in* André, *Spec. Hym. Eur. Alg.*, vol. 10, pp. 364, 426.

TYPE: *Acanopsilus clavatus* Kieffer, 1908, by monotypy.

#### **Acanosema** Kieffer

1908. Kieffer, *in* André, *Spec. Hym. Eur. Alg.*, vol. 10, pp. 360, 367, 407. Four species.

TYPE: *Acanosema rufum* Kieffer, 1908, by original designation.

#### **Acanthinevania** Bradley

1908. Bradley, *Trans. Amer. Ent. Soc.*, vol. 34, pp. 163, 172. Thirty-two species and one variety.

TYPE: *Evania princeps* Westwood, 1841, by original designation.

#### **Acanthobetyla** Dodd

1926. Dodd, *Trans. Roy. Soc. South Australia*, vol. 50, p. 299.

TYPE: *Acanthobetyla mirabilis* Dodd, 1926, by monotypy and original designation.

#### **Acantholapitha** Cameron

1912 (July). Cameron, *Soc. Ent.*, vol. 27, no. 15, p. 70. No species.

1912 (Aug.). Cameron, Soc. Ent., vol. 27, no. 16, p. 74. One species.

TYPE: *Acantholapitha nigricollis* Cameron, 1912. First included species.

#### **Acanthomiota** Jansson

1942. Jansson, Ent. Tidskr., vol. 63, p. 210.

TYPE: *Acanthomiota oxylaboides* Jansson, 1942, by monotypy.

#### **Acanthopria** Ashmead

1896. Ashmead, Proc. Zool. Soc. London, 1895, p. 804.

TYPE: *Acanthopria crassicornis* Ashmead, 1896, by monotypy and original designation.

#### **Acanthopsilus** Kieffer

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 380, 394.

TYPE: *Oxylabis marshalli* Kieffer, 1907, by monotypy.

Proposed as a subgenus of *Oxylabis* Foerster.

#### **Acanthoscelio** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 210, 211, 236. Two species.

TYPE: *Acanthoscelio americanus* Ashmead, 1893, by original designation.

#### **Acanthoserphus** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, p. 384.

TYPE: *Acanthoserphus albicoxa* Dodd, 1915, by monotypy and original designation.

#### **Acanthoteleia** Kieffer

1910. Kieffer, Ann. Soc. Ent. France, vol. 78, 1909, p. 311.

TYPE: *Acanthoteleia nigriclavus* Kieffer, 1910, by monotypy.

#### **Acerota** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, p. 107. No species.

1887. Ashmead, Can. Ent., vol. 19, p. 128. Two species.

TYPE: *Acerota caryae* Ashmead, 1887. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 707. *Acerota evanescens* Kieffer, 1914, designated by Kieffer, 1914, in André, Spec. Hym. Eur. Alg., vol. 11, p. 369, was included later.

**Acidopria** Kieffer

1913. Kieffer, *Insecta*, vol. 3, p. 442. Three species.

TYPE: *Acidopria variicornis* Kieffer, 1913. Designated by Kieffer, 1916, *Das Tierreich*, Lief. 44, p. 165.

**Acidopsilus** Kieffer

1909. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 33, p. 340.

TYPE: *Acidopsilus longicornis* Kieffer, 1909, by monotypy.

**Aclista** Foerster

1856. Foerster, *Hymenopterologische Studien*, Heft 2, pp. 128, 131, 133, 134, 135. No species.

1873. Marshall, *A catalogue of British Hymenoptera*; *Oxyura*, p. 9. Three species, one of them doubtfully included.

TYPE: *Acoretus scutellaris* Thomson, 1858. Designated by Muesebeck and Walkley, 1951, in Muesebeck, et al., *U. S. Dept. Agr., Agr. Monogr. No. 2*, p. 685.

**Aclistoides** Dodd

1916. Dodd, *Trans. Roy. Soc. South Australia*, vol. 40, p. 11.

TYPE: *Aclistoides retractus* Dodd, 1916, by monotypy and original designation.

**Acoloides** Howard

1890. Howard, *Insect life*, vol. 2, p. 269.

TYPE: *Acoloides saitidis* Howard, 1890, by monotypy.

**Acolomorpha** Dodd

1914. Dodd, *Can. Ent.*, vol. 46, p. 60.

TYPE: *Acolomorpha minuta* Dodd, 1914, by monotypy and original designation.

**Acolus** Foerster (= *Holacolus* Kieffer; *Holalcus* Risbec, error)

1856. Foerster, *Hymenopterologische Studien*, Heft 2, pp. 100, 102. No species.

1859. Thomson, *Öfv. Vet.-Akad. Förh.*, vol. 15, 1858, p. 422. Two species.

TYPE: *Acolus opacus* Thomson, 1859. Designated by Ashmead, 1903, *Journ. New York Ent. Soc.*, vol. 11, p. 90. *Acolus xanthogaster* Ashmead, 1893, designated by Brues, 1908, in Wytsman, *Genera insectorum*, fasc. 80, p. 16, and *Acolus piceiventris* Kieffer, 1908, designated by Kieffer, 1910, in Wytsman, *Genera insectorum*, fasc. 80B, p. 102, were included later. *A. piceiventris* was described by Kieffer from a specimen in the Foerster collection and is the species upon which the genus was based. However, in a recent decision (*Bull. Zool. Nom.*, vol. 4, 1950, pp. 160, 346)

the International Commission on Zoological Nomenclature agreed that only the first species referred to a genus described without originally included species shall be eligible for type designation.

Isogenotypic with *Holacolus* Kieffer.

#### **Acoretus** Haliday

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. p. 169.

TYPE: *Belyta rufopetiolata* Nees, 1834, by monotypy.

Proposed as a subgenus of *Belyta* Jurine. Considered synonym of *Achista* Foerster.

#### **Acropiesta** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 129, 135. No species.

1883. Cameron, Trans. Ent. Soc. London, 1883, p. 195. One species doubtfully included and therefore not available as type.

1893. Ashmead, U. S. Nat. Mus. Bull. 45, p. 353. Two species.

TYPE: *Acropiesta flavicauda* Ashmead, 1893. Designated by Muesebeck and Walkley, 1951, in Muesebeck, et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 689. *Acropiesta collaris* Foerster, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 353, was a nomen nudum.

#### **Acutibaesus** Meunier [fossil]

1916. Meunier, Zeitschr. Deutschen Geol. Gesellsch., vol. 68, pp. 392, 395.

TYPE: *Acutibaesus bellicosus* Meunier, 1916, by monotypy.

#### **Adelioneiva** Fischer

1940. Fischer, Rev. de Ent., vol. 11, p. 397.

TYPE: *Adelioneiva concolor* Fischer, 1940, by monotypy and original designation.

#### **Adeliopria** Ashmead

1902. Ashmead, Biol. Bull., vol. 3, p. 15.

TYPE: *Adeliopria longii* Ashmead, 1902, by monotypy and original designation.

#### **Aegyptoscelio** Priesner

1951. Priesner, Bull. Inst. Fouad I du Desert, vol. 1, p. 133.

TYPE: *Aegyptoscelio frequens* Priesner, 1951, by monotypy and original designation.

#### **Afrevania** Benoit

1953. Benoit, Rev. Zool. Bot. Africaines, vol. 48, p. 259.

TYPE: *Afrevania leroysi* Benoit, 1953, by monotypy and original designation.

**Agonophorus** Dahlbom

1858. Dahlbom, Öfv. Vet.-Akad. Förh., vol. 14, 1857, p. 289.  
No species.

No species have been referred to this genus up to the present time.

**Aholcus** Kieffer

1913. Kieffer, in Alluaud and Jeannel, Voyage . . . en Afrique Orientale. Insectes hyménoptères, vol. 1, p. 4.

TYPE: *Aholcus monticola* Kieffer, 1913, by monotypy and original designation.

**Aleria** Marshall

1874. Marshall, Ent. Monthly Mag., vol. 10, p. 208.

TYPE: *Aleria flavibarbis* Marshall, 1874, by monotypy.

Considered synonym of *Scelio* Latreille.

**Allogryon** Kieffer

1910. Kieffer, in Wytzman, Genera insectorum, fasc. 80B, p. 95. Thirty-two species.

TYPE: *Prosacantha caraborum* Riley, in Ashmead, 1893. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 697.

Proposed as a subgenus of *Hoplogryon* Ashmead. Considered synonym of *Trimorus* Foerster.

**Allomicrops** Kieffer

1914. Kieffer, Das Tierreich, Lief. 42, pp. xv, 69, 138.

TYPE: *Ceraphron abnormis* Perkins, 1910, by monotypy.

**Allophanurus** Kieffer

1912. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 8, 11. Two species.

TYPE: *Telenomus vibius* Walker, 1838. Designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 23.

**Alloteleia** Kieffer

1917. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 15, p. 59.

TYPE: *Alloteleia appendiculata* Kieffer, 1917, by monotypy and original designation.

**Allotropa** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 106, 109.

TYPE: *Inostemma mecrida* Walker, 1835, by monotypy.

**Amblyaspis** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 107, 112. Three species included by bibliographic reference.



TYPE: *Platygaster tritici* Curtis, 1831. Designated by Kieffer, 1926, *Das Tierreich*, Lief. 48, p. 611. *Amblyaspis aliena* Foerster MS., designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 266, and by Fouts, 1924, Proc. U. S. Nat. Mus., vol. 63, art. 15a, p. 3, was a nomen nudum taken from a label on a specimen in the Foerster collection.

#### **Amblyconus** Kieffer

1913. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 221. No species.

1914. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 325. One species.

TYPE: *Amblyconus quadridens* Kieffer, 1914. First included species.

#### **Amblyscelio** Kieffer

1913. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 225, 258.

TYPE: *Amblyscelio striaticeps* Kieffer, 1913, by monotypy.

#### **Amitus** Haldeman (= *Elaptus* Forbes)

1850. Haldeman, Amer. Journ. Sci., ser. 2, vol. 9, p. 109.

TYPE: *Amitus aleurodinis* Haldeman 1850 (= *Amitus corni* Haldeman, error), by monotypy.

Isogenotypic with *Elaptus* Forbes through synonymy.

#### **Anaclista** Kieffer

1916. Kieffer, *Das Tierreich*, Lief. 44, pp. xxv, 352, 462. Two species.

TYPE: *Anaclista holotoma* Kieffer, 1909, by original designation.

#### **Anacoryphe** Debauche

1947. Debauche, Bull. Ann. Soc. Ent. Belgique, vol. 83, p. 274.

TYPE: *Anacoryphe orchymonti* Debauche, 1947, by monotypy and original designation.

#### **(*Anaulacus* Semenov, not MacLeay, 1825, replaced by *Semenowia* Kieffer, 1903, also preoccupied) = *Semenovius* Bradley**

1903. Semenov, Rev. Russe Ent., vol. 3, p. 173. Two species, one of them doubtfully included.

TYPE: *Aulacus sibiricola* Semenov, 1894. Designated by Bradley, 1908, Trans. Amer. Ent. Soc., vol. 34, p. 120.

#### **Anectadius** Kieffer

1905. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 29, p. 187. Two species.

TYPE: *Anectadius striolatus* Kieffer, 1905. Designated by Kieffer, 1926, *Das Tierreich*, Lief. 48, p. 608.

**Anectata** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 129, 136. No species.

1887. Ashmead, Can. Ent., vol. 19, p. 198. One species.

TYPE: *Anectata hirtifrons* Ashmead, 1887. First included species.

Considered synonym of *Aclista* Foerster.

**Aneurhynchus** Westwood (= *Mythras* Haliday, in Curtis, nomen nudum; *Mithras* Agassiz, emendation; *Mythras* Brullé; *Aneurhynchus* Maneval and *Aneurhynchus* Brullé, emendations)

1832. Westwood, Journ. London Edinburgh Phil. Mag., ser. 3, vol. 1, p. 129.

TYPE: *Aneurhynchus galesiformis* Westwood, 1832, by monotypy.

*Mythras* Haliday, in Curtis, nomen nudum, was validated by Brullé, in Lepeletier, 1846, Histoire naturelle des insectes, Hyménoptères, vol. 4, p. 612, through citation as a synonym of *Aneurhynchus*.

**Aneurobaeus** Kieffer

1912. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 87, 91.

TYPE: *Baeus apterus* Bugnion and Popoff, 1910, by monotypy and original designation.

**Aneuron** Brues

1910. Brues, Bull. Wisconsin Nat. Hist. Soc., vol. 8, p. 49.

TYPE: *Aneuron anormis* Brues, 1910, by monotypy and original designation.

Considered synonym of *Platygaster* Latreille.

**Aneuropria** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2 (42), p. 35. No species.

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 898. One species.

TYPE: *Aneuropria clavata* Kieffer, 1911. First included species.

**Aneuroscelio** Kieffer

1913. Kieffer, in Alluaud and Jeannel, Voyage . . . en Afrique Orientale. Insectes hyménoptères, vol. 1, p. 14.

TYPE: *Aneuroscelio rufipes* Kieffer, 1913, by monotypy and original designation.

(*Aneurhynchus* Berland, emendation) = *Aneurhynchus* Westwood

1940. Maneval, in Berland, Faune de la France, vol. 7, Hyménoptères, p. 103.

(*Aneurynchus* Brullé, emendation) = *Aneurhynchus* Westwood

1846. Brullé, in Lepeletier, Histoire naturelle des insectes. Hyménoptères, vol. 4, p. 612.

(*Anisoptera* Herrich-Schaeffer, not Berthold, 1827) = *Laches* Gistel

1840. Herrich-Schaeffer, Nomenclator entomologicus, Heft 2, pp. 57, 69.

TYPE: *Anisoptera egregia* Herrich-Schaeffer, 1840, by monotypy.

Considered synonym of *Psilus* Panzer.

**Anommatium** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 130, 131, 140. No species.

1904. Mayr, Verh. Zool.-Bot. Gesellsch. Wien, vol. 54, p. 592. One species.

TYPE: *Anommatium ashmeadi* Mayr, 1904. First included species.

**Anopedias** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 108, 114. No species.

1859. Thomson, Öfv. Vet.-Akad. Förh., vol. 16, p. 79. Three species.

TYPE: *Anopedias obscurus* Thomson, 1859. Designated by Ashmead, 1903, Journ. New York Ent. Soc., vol. 11, p. 97. *Anopedias lacustris* Kieffer, 1926, designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 701, was included later. *Anopedias tritomus* Thomson, 1859, was designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 716, through oversight of Ashmead's 1903 citation.

Considered synonym of *Leptacis* Foerster.

**Anoxylabis** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 343.

TYPE: *Anoxylabis laticeps* Kieffer, 1909, by monotypy.

**Antarctopria** Brues

1920. Brues, in Tillyard, Australasian Antarctic Expedition . . . Scientific reports, ser. c, vol. 5, pt. 8, p. 17; Brues, *ibid.*, app. A, p. 27.

TYPE: *Antarctopria latigaster* Brues, 1920, by monotypy and original designation.

**Anteris** Foerster (= *Auteris* Ashmead, error; *Trichacolus* Kieffer) 1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 101, 103. No species.

1859. Thomson, Öfv. Vet.-Akad. Förh., vol. 15, 1858, p. 421. Two species.

TYPE: *Anteris bilineatus* Thomson, 1859, by present designation. *Anteris rufitarsis* Foerster MS., designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 224, was a nomen nudum. *Anteris virginiensis* Ashmead, 1893, designated by Brues, 1908, in Wytsman, Genera insectorum, fasc. 80, p. 36, and *Anteris simulans* Kieffer, 1908, designated by Kieffer, 1910, in Wytsman, Genera insectorum, fasc. 80B, p. 80, were included later. *A. simulans* was described by Kieffer from the specimen in the Foerster collection upon which the genus was based, but under the recent modification of Opinion 46 by the International Commission on Zoological Nomenclature (Bull. Zool. Nomencl., vol. 4, 1950, pp. 160, 346) only the first included species shall be eligible for designation as type of a genus described without contained species.

Isogenotypic with *Trichacolus* Kieffer.

**Anteromorpha** Dodd

1913. Dodd, Trans. Roy. Soc. South Australia, vol. 37, pp. 131, 145. Two species.

TYPE: *Anteromorpha australica* Dodd, 1913, by original designation.

**Antropria** Kieffer

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 694, 743.

TYPE: *Antropria pedestris* Kieffer, 1910, by monotypy.

**Antrosclio** Kieffer

1913. Kieffer, in Alluaud and Jeannel, Voyage . . . en Afrique Orientale. Insectes hyménoptères, vol. 1, p. 15.

TYPE: *Antrosclio lucifugax* Kieffer, 1913, by monotypy and original designation.

**Aparamesius** Kieffer

1913. Kieffer, Insecta, vol. 3, p. 436. Four species.

TYPE: *Aparamesius carinatus* Kieffer, 1913, by original designation.

**Apegus** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 101, 104.

TYPE: *Apegus leptocerus* Foerster, 1856, by monotypy.

**Apegusoneura** Cameron

1912. Cameron, Soc. Ent., vol. 27, No. 15, p. 69. Three species.

TYPE: *Apegusoneura striolatus* Cameron, 1912. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 703.

Considered synonym of *Hoploteleia* Ashmead.

**Aphanogmus** Thomson

1858. Thomson, Öf. Vet.-Akad. Förh., vol. 15, pp. 287, 305. Five species.

TYPE: *Aphanogmus fumipennis* Thomson, 1858. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 132.

**Aphanomerella** Dodd

1913. Dodd, Mem. Queensland Mus., vol. 2, p. 339.

TYPE: *Aphanomerella ovi* Dodd, 1913, by monotypy and original designation.

**Aphanomerus** Perkins

1905. Perkins, Haw. Sugar Planters Assoc. Exp. Sta., Div. Ent., Bull. 1, pp. 200, 201. Four species.

TYPE: *Aphanomerus bicolor* Perkins, 1905, by original designation.

**(*Aphanurus* Kieffer, not Looss, 1907) = *Microphanurus* Kieffer**

1912. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 10, 69. Nineteen species, one of them doubtfully included.

TYPE: *Teleas semistriatus* Nees, 1834, by original designation.

**Apteroscelio** Kieffer

1913. Kieffer, in Alluaud and Jeannel, Voyage . . . en Afrique Orientale. Insectes hyménoptères, vol. 1, p. 17.

TYPE: *Apteroscelio montanus* Kieffer, 1913, by monotypy and original designation.

**Aradophagus** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 138, 166.

TYPE: *Aradophagus fasciatus* Ashmead, 1893, by monotypy.

**Aratala** Dodd

1927. Dodd, Mem. Queensland Mus., vol. 9, p. 74.

TYPE: *Aratala globiceps* Dodd, 1927, by monotypy and original designation.

**Archaebelyta** Meunier [fossil]

1923. Meunier, Misc. Ent., vol. 26, No. 11-12, p. 84.

TYPE: *Archaebelyta superba* Meunier, by monotypy.

**Archaeoscelio** Brues [fossil]

1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 74, p. 88.  
Two species.

TYPE: *Archaeoscelio rugosus* Brues, 1940, by original designation.

**(Artibolus** Haliday) = **Synacra** Foerster

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. p. 173. Two species.

TYPE: *Diapria brachialis* Nees, 1834, by present designation.  
Proposed as a subgenus of *Platymischus* Westwood. Isogenotypic with *Synacra* Foerster.

**Ashmeadopria** Kieffer

1912. Kieffer, in Wytzman, Genera insectorum, fasc. 124, 1911, pp. 8, 10, 59. Fifty-five species.

TYPE: *Diapria verticillata* Latreille, 1805. Designated by Mani, 1941, Catalogue of Indian insects, Pt. 26, Serphoidea, p. 44.

Considered synonym of *Trichopria* Ashmead.

**Asolcus** Nakagawa

1900. Nakawaga, Spec. Rep. Imp. Agric. Exp. Sta., Japan, 6, p. 17.

TYPE: *Asolcus nigripedius* Nakagawa, by monotypy.

**Asolenopsia** Kieffer

1921. Kieffer, An. Soc. Cient. Argentina, vol. 91, p. [36].

TYPE: *Asolenopsia rufa* Kieffer, 1921, by monotypy and original designation.

**Atelopsilus** Kieffer

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 360.

TYPE: *Pantolyta brunnea* Ashmead, 1893, by monotypy and original designation.

Proposed as a subgenus of *Rhynchopsilus* Kieffer.

**Atomopria** Kieffer

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 695. No species.

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 875. Two species.

TYPE: *Atomopria rufithorax* Kieffer, 1911. Designated by Kieffer, 1911, in André, Spec. Hym. Eur. Alg., vol. 10, p. 875.

**Atrichopria** Kieffer

1910. Kieffer, Ent. Rundschau, vol. 27, p. 55. Two species.

TYPE: *Atrichopria rufa* Kieffer, 1910. Designated by Kieffer, 1912, in Wytsman, Genera insectorum, fasc. 124, 1911, p. 54.

**Atritomellus** Kieffer (= *Atritomus* Foerster, 1878, not Reitter, 1877).

1914. Kieffer, Das Tierreich, Lief. 42, pp. xv, 139, 141.

TYPE: *Atritomus coccophagus* Foerster, 1878, by substitution of *Atritomellus* for *Atritomus* Foerster.

**(Atritomus** Foerster, not Reitter, 1877) = **Atritomellus** Kieffer

1878. Foerster, Verh. Naturh. Ver. Preuss. Rheinl., vol. 35, p. 56.

TYPE: *Atritomus coccophagus* Foerster, 1878, by monotypy and original designation.

**Aulacinus** Westwood

1868. Westwood, Trans. Ent. Soc. London, 1868, p. 331.

TYPE: *Aulacus (Aulacinus) moerens* Westwood, 1868, by monotypy.

Proposed as a subgenus of *Aulacus* Jurine.

**Aulacites** Cockerell [fossil]

1916. Cockerell, Proc. U. S. Nat. Mus., vol. 51, p. 102. Two species.

TYPE: *Aulacites secundus* Cockerell, 1916, by original designation.

**Aulacofoenus** Kieffer

1911. Kieffer, Ann. Soc. Ent. France, vol. 80, p. 177.

TYPE: *Hyptiogaster szepligetii* Kieffer, 1904, by monotypy and original designation.

**Aulacomastus**, new name (= *Aulacostethus* Philippi, 1873, not Waterhouse, 1869, not Uhler, 1871; *Aulacosthethus* Schletterer, error; *Aulacostathus* Dalla Torre, error; *Aulacosthetus* Kieffer, error)

TYPE: *Aulacostethus rubriventer* Philippi, 1873, by substitution of *Aulacomastus* for *Aulacostethus* Philippi.

**Aulacopria** Kieffer

1904. Kieffer, Bull. Soc. Hist. Nat. Metz, vol. 23, p. 54.

TYPE: *Aulacopria formicarum* Kieffer, 1904, by monotypy.

**(Aulacostathus** Dalla Torre, error for **Aulacostethus** Philippi) = **Aulacomastus**, new name

1902. Dalla Torre, Catalogus hymenopterorum, vol. 3, pt. 2, p. 1062.

- (*Aulacostethus* Philippi, not Waterhouse, 1869, not Uhler, 1871)  
 = *Aulacomastus*, new name  
 1873. Philippi, Stett. Ent. Zeit., vol. 34, p. 302.  
 TYPE: *Aulacostethus rubriventer* Philippi, 1873, by monotypy.
- (*Aulacostethus* Schletterer, error for *Aulacostethus* Philippi)  
 = *Aulacomastus*, new name  
 1890. Ann. Naturh. Hofmus. Wien, vol. 4, p. 523.
- (*Aulacosthetus* Kieffer, error for *Aulacostethus* Philippi) = *Aulacomastus*, new name  
 1912. Kieffer, Das Tierreich, Lief. 30, p. 370.
- Aulacus** Jurine  
 1807. Jurine, Nouvelle méthode de classer les hyménoptères et les diptères, p. 89.  
 TYPE: *Aulacus striatus* Jurine, 1807, by monotypy.
- Aulatopria** Brèthes  
 1927. Brèthes, Rev. Indus. Agr. Tucumán, vol. 17, pp. 163, 164.  
 TYPE: *Aulatopria tucumana* Brèthes, 1927, by monotypy.  
 Also published later in 1927, Bull. Ent. Res., vol. 18, p. 206, as English translation.
- Austronia** Riek  
 1955. Riek, Australian Journ. Zool., vol. 3, p. 263. Three species.  
 TYPE: *Austronia nitida* Riek, 1955, by original designation.
- Austroscelio** Dodd  
 1914. Dodd, Proc. Roy. Soc. Queensland, vol. 26, p. 93. Three species.  
 TYPE: *Sparasion nigricoxa* Dodd, 1914, by original designation.
- Austroserphus** Dodd  
 1933. Dodd, Proc. Linn. Soc. New South Wales, vol. 58, p. 275.  
 TYPE: *Austroserphus albofasciatus* Dodd, 1933, by monotypy and original designation.
- (*Auteris* Ashmead, error) = *Anteris* Foerster  
 1903. Ashmead, Journ. New York Ent. Soc., vol. 11, p. 92.
- Auxopaedeutes** Brues  
 1903. Brues, Trans. Amer. Ent. Soc., vol. 29, p. 126.  
 TYPE: *Auxopaedeutes sodalis* Brues, 1903, by monotypy.
- (*Axestonotus* Kieffer, new name for *Xestonotus* Foerster, 1856, not Leconte, 1853) = *Xestonotidea* Gahan  
 1926. Kieffer, Das Tierreich, Lief. 48, pp. xxxi, 561, 625.  
 TYPE: *Xestonotus andriciphilus* Ashmead, 1887, by substitution of *Axestonotus* for *Xestonotus* Foerster.



**Bactropria** Kieffer

1910. Kieffer, Ann. Soc. Ent. France, vol. 78, 1909, p. 339.

TYPE: *Bactropria brasiliensis* Kieffer, 1910, by monotypy.

**(Baeceura** Ashmead, error) = **Baeoneura** Foerster

1887. Ashmead, Ent. Amer., vol. 3, p. 99.

**Baeomorpha** Brues [fossil]

1937. Brues, in Carpenter, Univ. Toronto Stud., Geol. ser., vol. 40, p. 41.

TYPE: *Baeomorpha dubitata* Brues, 1937, by monotypy and original designation.

**Baeoneura** Foerster (= *Baeceura* Ashmead, error)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 100, 102. No species.

1887. Ashmead, Ent. Amer., vol. 3, p. 99. Two species.

TYPE: *Baeoneura floridana* Ashmead, 1887, by present designation.

Considered synonym of *Macroteleia* Westwood.

**Baeoneurella** Dodd

1914. Dodd, Trans. Roy. Soc. South Australia, vol. 38, p. 124. Four species.

TYPE: *Baeoneura giraulti* Dodd, 1913, by original designation.

Considered synonym of *Eumicrosoma* Gahan.

**Baeus** Haliday (= *Hyperbaeus* Foerster)

1833. Haliday, Ent. Mag., vol. 1, p. 270.

TYPE: *Baeus seminulum* Haliday, 1833, by monotypy.

**Bakeria** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2 (42), p. 34. No species.

1906. Kieffer, Berliner Ent. Zeitschr., vol. 50, 1905, p. 282. One species.

TYPE: *Bakeria complanata* Kieffer, 1906. First included species.

**(Balyta** Ashmead, error) = **Belyta** Jurine

1893. Ashmead, U. S. Nat. Mus. Bull. 45, p. 366.

**Baryconus** Foerster (= *Hoploteleia* Ashmead)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 101, 104. No species.

1887. Ashmead, Ent. Amer., vol. 3, p. 118. One species.

TYPE: *Baryconus floridanus* Ashmead, 1887. First included species. *Baryconus oecanthi* Riley, in Ashmead, 1893, designated by Brues, 1908, in Wytzman, Genera insectorum, fasc. 80, p. 31, and *Baryconus rufipes* Kieffer, 1903, designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 480, were included later.

Isogenotypic with *Hoploteleia* Ashmead.

**Baryteleia** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxix, 273, 544.  
Three species.

TYPE: *Macroteleia nigriceps* Kieffer, 1905, by original designation.

**Basalys** Westwood

1833. Westwood, London Edinburgh Phil. Mag., ser. 3, vol. 3, p. 343.

TYPE: *Basalys fumipennis* Westwood, 1833, by monotypy.

**Bebelus** Gistel (= *Oxyurus* Lamarck, 1817, not Rafinesque, 1810)

1848. Gistel, Naturgeschichte des Tierreichs, p. x.

TYPE: *Sparasion frontalis* Latreille, 1805, by substitution of *Bebelus* for *Oxyurus* Lamarck.

Considered synonym of *Sparasion* Latreille.

(*Belita* Haliday, emendation) = **Belyta** Jurine

1857. Nat. Hist. Rev., vol. 4, Proc. p. 169.

**Belyta** Jurine (= *Belita* Haliday, emendation; *Balyta* Ashmead, error)

1807. Jurine, Nouvelle méthode de classer les hyménoptères et les diptères, p. 311.

TYPE: *Belyta bicolor* Jurine, 1807, by monotypy. *Belyta iridipennis* Lepeletier and Serville, 1825, was designated by Blanchard, 1849, in Cuvier, Règne animal [ed. 3], Insectes, text vol. 2, p. 156; atlas, pl. 115, fig. 6 (Blanchard erroneously cited Curtis as author of *iridipennis* in plate legend and Walker on plate itself).

(*Betula* Ashmead, error) = **Betyla** Cameron

1902. Ashmead, Journ. New York Ent. Soc., vol. 10, p. 245.

**Betyla** Cameron (= *Betula* Ashmead, error)

1889. Cameron, Mem. Proc. Manchester Lit. Phil. Soc., ser. 4, vol. 2, p. 12.

TYPE: *Betyla fulva* Cameron, 1889, by monotypy.

**Bothriopria** Kieffer

1905. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 29, pp. 132, 134.

TYPE: *Bothriopria saussurei* Kieffer, 1905, by monotypy.

**Bracalba** Dodd

1930. Dodd, Proc. Roy. Soc. Queensland, vol. 42, p. 78.  
Three species.

TYPE: *Bracalba laminata* Dodd, 1930, by original designation.

**Brachevania** Turner

1927. Turner, Ann. Mag. Nat. Hist., ser. 9, vol. 20, p. 553.

TYPE: *Brachevania kristenseni* Turner, 1927, by monotypy and original designation.

**Brachinostemma** Kieffer

1916. Kieffer, Zentralbl. Bakt. Parasitenk. Infektionsk. Abt. 2, vol. 46, p. 551.

TYPE: *Brachinostemma mediterranea* Kieffer, 1916, by monotypy and original designation.

**Brachygaster** Leach (= *Coranila* Westwood; *Semaeodogaster* Bradley)

1815. Leach, in Brewster, Edinburgh Encyclopaediae, vol. 9, p. 142.

TYPE: *Evania minuta* Olivier, 1791, by monotypy.

Isogenotypic with *Semaeodogaster* Bradley.

**Brachyscelio** Brues [fossil]

1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 74, p. 76.  
Two species.

TYPE: *Brachyscelio cephalotes* Brues, 1940, by original designation.

**(*Brachyscelio* Risbec, not Brues, 1940) = *Pachyscelio* Risbec**

1950. Risbec, Contribution à l'étude des Proctotrupidae. Trav. Lab. Ent. Sect. Soudan. Rech. Agron., p. 614.

TYPE: *Brachyscelio jeanneli* Risbec, 1950, by monotypy.

**Brachyserphus** Hellén

1941. Hellén, Notulae Ent., vol. 21, p. 42. Two species.

TYPE: *Codrux parvulus* Nees, 1834, by original designation.

**Bruchiola** Kieffer

1921. Kieffer, An. Soc. Cient. Argentina, vol. 91, p. 40.

TYPE: *Bruchiola formicaria* Kieffer, 1921, by monotypy.

**Bruchopria** Kieffer

1921. Kieffer, An. Soc. Cient. Argentina, vol. 91, p. 38. Two species.

TYPE: *Bruchopria pentatoma* Kieffer, 1921, by original designation.

**Bruesopria** Wing

1951. Wing, Trans. Roy. Ent. Soc. London, vol. 102, p. 195.  
Two species.

TYPE: *Bruesopria seeversi* Wing, 1951, by original designation.

**Brunnicophilus** Nixon

1931. Nixon, Ent. Rec. Journ. Variation, vol. 43, p. 83.

TYPE: *Brunnicophilus donisthorpei* Nixon, 1931, by monotypy and original designation.

(*Cacellus* Ashmead, new name for *Cacus* Riley, 1893, not Gistel, 1848, not Selys, 1854, not Costa, 1857) = *Oethecoctonus* Ashmead

1913. Ashmead, Journ. New York Ent. Soc., vol. 11, p. 92.

TYPE: *Cacus oecanthi* Riley, in Ashmead 1893, by substitution of *Cacellus* for *Cacus* Riley.

Unnecessarily proposed for *Cacus* Riley. *Oethecoctonus* Ashmead, 1900, had already been published as a replacement name.

(*Cacus* Riley, not Gistel, 1848, not Selys, 1854, not Costa, 1857) = *Oethecoctonus* Ashmead

1893. Riley, in Ashmead, U. S. Nat. Mus. Bull. 45, pp. 209, 210, 211, 223.

TYPE: *Cacus oecanthi* Riley in Ashmead, 1893, by monotypy and designation of Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 223.

**Caenoteleia** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxix, 266, 550.

TYPE: *Caloteleia elegans* Perkins, 1926, by monotypy.

(*Calliceras* Nees (= *Hadroceras* Foerster, new name) = *Ceraphron* Jurine

1834. Nees, Hymenopterorum ichneumonibus affinium, monographiae, vol. 2, p. 278. Ten species.

TYPE: *Calliceras sulcata* Nees, 1834. Designated by Mani, 1941, Catalogue of Indian insects, part 26, Serphoidea, p. 37. *Ceraphron carpenteri* Curtis, 1829, designated by Westwood, 1839, An introduction to the modern classification of insects, vol. 2, generic synopsis p. 77, was not among the originally included species.

Isogenotypic with *Ceraphron* Jurine through synonymy.

**Calliscelio** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 209, 218.

TYPE: *Calliscelio laticinctus* Ashmead, 1893, by monotypy and original designation.

(*Callitelea* Agassiz, emendation) = *Calotelea* Westwood

1946. Agassiz, Nomenclator zoologicus. Index universalis, pp. 60, 61.

**Calogalesus** Kieffer

1912. Kieffer, in Wytzman, Genera insectorum, fasc. 124, 1911, pp. 6, 43.

TYPE: *Calogalesus parvulus* Kieffer, 1912, by monotypy.

Also described as new by Kieffer, 1912, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 73.

#### **Caloptenobia** Riley

1878. Riley, Ann. Rep. U. S. Ent. Comm., Dep. Int., No. 1, p. 306.

TYPE: *Caloptenobia ovivora* Riley, 1878, by monotypy.

Considered synonym of *Scelio* Latreille.

#### **Calotelea** Westwood (= *Callitelea* Agassiz, emendation; *Caloteleia* Ashmead, emendation) [fossil]

1837. Westwood, in Hope, Trans. Ent. Soc. London, vol. 2, p. 55.

TYPE: *Calotelea aurantia* Hope, 1837, by monotypy.

#### **(*Caloteleia* Ashmead, emendation) = *Calotelea* Westwood**

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 209, 210, 212.

#### **Camptopsilus** Kieffer

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 359, 366. No species.

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 339. One species.

TYPE: *Camptopsilus nigriceps* Kieffer, 1909. First included species.

#### **Camptoteleia** Kieffer

1913. Kieffer, Insecta, vol. 3, p. 387. Two species.

TYPE: *Camptoteleia carinata* Kieffer, 1913. Designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 379.

#### **Carabiphagus** Morley

1929. Morley, Trans. Suffolk Nat. Soc., vol. 1, p. 40.

TYPE: *Proctotrupes laevifrons* Foerster, 1861, by monotypy. Considered synonym of *Phaenoserphus* Kieffer.

#### **Cardiopria** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 398, 429.

TYPE: *Cardiopria ventralis* Dodd, 1915, by monotypy and original designation.

#### **Cardiopsilus** Kieffer

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 360, 405.

TYPE: *Cardiopsilus productus* Kieffer, 1908, by monotypy.

#### **Carinafoenus** Crosskey

1953. Crosskey, Trans. Roy. Ent. Soc. London, vol. 104, pp. 355, 358. Five species.

TYPE: *Foenus rufus* Westwood, 1841, by original designation.

**Carinia** Kieffer

1905. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 29, p. 140.

TYPE: *Carinia nitida* Kieffer, 1905, by monotypy.

(*Catillus* Foerster, not Brongniart, 1822) = *Picstopleura* Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 107, 111, 144.

TYPE: *Platygaster catillus* Walker, 1834 (= *Catillus walkeri* Foerster, 1856), by monotypy.

**Ceranogmus** Risbec

1953. Risbec, Bull. Inst. Français d'Afrique Noire, vol. 15, p. 560.

TYPE: *Ceranogmus testaceus* Risbec, 1953, by monotypy.

**Ceraphron** Jurine (= *Calliceras* Nees (= *Hadrocera* Foerster, new name); *Cerataphron* Schulz, emendation)

1807. Jurine, Nouvelle méthode de classer les hyménoptères et les diptères, p. 303. Two species.

TYPE: *Ceraphron sulcatus* Jurine, 1807. Designated by the International Commission on Zoological Nomenclature, 1946, under suspension of the Rules, Opinion 174, p. [497]. All prior type designations were set aside by this opinion.

Isogenotypic with *Calliceras* Nees through synonymy.

**Ceraphron** Panzer

1805. Panzer, Faunae insectorum Germaniae initia . . . , Heft 97, No. 16.

This name has been suppressed by the International Commission on Zoological Nomenclature under suspension of the Rules (Opinion 174, 1946).

**Cerapsilus** Haliday

1829. Haliday, in Curtis, A guide to an arrangement of British insects, column 109.

TYPE: *Psilus sericeicornis* Spinola, 1808, by monotypy. Eight names were listed but seven of them were nomina nuda.

(*Cerastopsilus* Kieffer, error) = *Ceratopsilus* Kieffer

1913. Kieffer, Insecta, vol. 3, pp. 461, 490.

**Ceratacis** Thomson

1859. Thomson, Öfv. Vet.-Akad. Förh., vol. 16, pp. 69, 78.

TYPE: *Ceratacis flavipes* Thomson, 1859, by monotypy.

Considered synonym of *Leptacis* Foerster.

(*Cerataphron* Schulz, emendation) = *Ceraphron* Jurine

1906. Schulz, Spolia hymenopterologica, p. 152.

**Ceratobaeoides** Dodd

1913. Dodd, Mem. Queensland Mus., vol. 2, p. 337. Two species.

TYPE: *Ceratobaeoides hackeri* Dodd, 1913, by original designation.

**Ceratobaeus** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 167, 175. Two species.

TYPE: *Ceratobaeus cornutus* Ashmead, 1893, by original description.

**Ceratopria** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 407, 428. Six species.

TYPE: *Ceratopria longiceps* Ashmead, 1893, by original designation. *Ceratopria longicornis* Ashmead, designated by Ashmead, 1903, Journ. New York Ent. Soc., vol. 11, p. 31, was an error for *Ceratopria longiceps* Ashmead.

Considered synonym of *Trichopria* Ashmead.

**Ceratopsilus** Kieffer (= *Cerastopsilus* Kieffer)

1913. Kieffer, Insecta, vol. 3, pp. 462 (fig. 12), 493.

TYPE: *Ceratopsilus rufipes* Kieffer, 1913, by monotypy and original designation.

**Ceratoteleia** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, p. 121. Thirteen species.

TYPE: *Caloteleia grenadensis* Ashmead, 1895. Designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 501.

**Chaetanteris** Priesner

1951. Priesner, Bull. Inst. Fouad I du Desert, vol. 1, p. 136.

TYPE: *Chaetanteris serraticeps* Priesner, 1951, by monotypy and original designation.

**Chalcidoptarella** Enderlein

1909. Enderlein, Stett. Ent. Zeit., vol. 70, p. 262. Two species.

TYPE: *Evaniellus chalcidipennis* Enderlein, 1905, by original designation.

**(Cheirocerus** Ashmead, lapsus) = **Lygocerus** Foerster

1881. Ashmead, Trans. Amer. Ent. Soc., vol. 9, Mon. Proc. p. xxxiv.

**(Chlidonia** Herrich-Schaeffer, not Lamouroux, 1824, not Huebner, 1825)

1838. Herrich-Schaeffer, in Panzer, Faunae insectorum Germaniae initia . . . , Heft 157, No. 24. Eight species.

TYPE: *Chlidonia moniliata* Herrich-Schaeffer, 1838. Designated by Viereck, 1914, U. S. Nat. Mus. Bull. 83, p. 31.

Viereck followed Dalla Torre, 1898, Catalogus hymenopterorum, vol. 4, p. 228, in synonymizing *Chlidonia* with *Hormius* Nees, 1818 (Braconidae), on the incorrect supposition that *Chlidonia moniliata* Herrich-Schaeffer was identical with *Bracon moniliatus* Nees, the type of *Hormius*.

Although *Chlidonia* Herrich-Schaeffer is preoccupied it is not renamed since it is considered a synonym of *Spilomicrus* Westwood.

**Chrestoteleia** Kieffer (= *Crestoteleia* Kieffer, error)

1913. Kieffer, Insecta, vol. 3, p. 388.

TYPE: *Chrestoteleia bakeri* Kieffer, 1913, by monotypy and original designation.

**Chromoteleia** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 209, 211, 219.

TYPE: *Chromoteleia semicyanea* Ashmead, 1893, by monotypy and original designation.

**Cinetus** Jurine (= *Cinntus* Curtis, error)

1807. Jurine, Nouvelle méthode de classer les hyménoptères et les diptères, p. 310. No species.

1825. Lepeletier and Serville, Encyclopédie Méthodique, Zool., vol. 10, p. 210. One species.

TYPE: *Cinetus iridipennis* Lepeletier and Serville, 1825. First included species. Also designated by Kieffer, 1910, Genera insectorum, fasc. 107, p. 36. *Cinetus gracilipes* Curtis, 1831, designated by Curtis, 1831, British entomology, vol. 8, folio 380, was included later. *Belyta bicolor* Jurine, 1807, was erroneously designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 682.

(*Cinntus* Curtis, error) = *Cinetus* Jurine

1829. Curtis, A guide to an arrangement of British insects, column 109.

**Clinopria** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2 (42), pp. 30, 37.

TYPE: *Clinopria fusca* Kieffer, 1905, by monotypy.

**Codrus** Panzer

1801. Panzer, Faunae insectorum Germainae initia . . . ,



Heft 85. No. 9. Two species.

TYPE: *Codrus niger* Panzer, 1801. Designated by Morice and Durrant, 1915, *Trans. Ent. Soc. London*, 1915, p. 435. *Codrus apterygynus* Haliday, 1839, designated by Ashmead, 1893, *U. S. Nat. Mus. Bull.* 45, p. 344, was not originally included.

**Coelopelta** Ashmead

1893. Ashmead, *U. S. Nat. Mus. Bull.* 45, pp. 263, 264, 289.

TYPE: *Coelopelta mirabilis* Ashmead, 1893, by monotypy and original designation.

**Coelopria** Kieffer

1910. Kieffer, *in* André, *Spec. Hym. Eur. Alg.*, vol. 10, pp. 716, 747.

TYPE: *Coelopria maura* Kieffer, 1910, by monotypy.

**Coenopria** Kieffer

1905. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 29, p. 132.

TYPE: *Coenopria fuscipennis* Kieffer, 1905, by monotypy.

**Cologlyptus** Crawford

1910. Crawford, *Proc. U. S. Nat. Mus.*, vol. 38, p. 123.

TYPE: *Cologlytus kiefferi* Crawford, 1910, by monotypy and original designation.

**Conostigmoides** Dodd

1914. Dodd, *Trans. Roy. Soc. South Australia*, vol. 38, pp. 88, 94.

TYPE: *Eumegaspilus erythrothorax* Ashmead, 1893, by monotypy and original designation.

**Conostigmus** Dahlbom

1858. Dahlbom, *Öfv. Vet.-Akad. Förh.*, vol. 14, 1857, p. 291. No species.

1907. Kieffer, *in* André, *Spec. Hym. Eur. Alg.*, vol. 10, p. 80. Many species.

TYPE: *Megaspilus alutaceus* Thomson, 1858. Designated by Muesebeck and Walkley, 1951, *in* Muesebeck et al., *U. S. Dept. Agr., Agr. Monogr. No. 2*, p. 670.

Proposed as a subgenus of *Megaspilus* Westwood.

**Copelus** Provancher

1881. Provancher, *Nat. Canadien*, vol. 12, p. 206.

TYPE: *Copelus paradoxus* Provancher, 1881, by monotypy. Considered synonym of *Helorus* Latreille.

**Coptera** Say

1836. Say, *Boston Journ. Nat. Hist.*, vol. 1, p. 281.

TYPE: *Coptera polita* Say, 1836, by monotypy. Considered synonym of *Psilus* Panzer.

**(Coranila Westwood) = Brachygaster Leach**

1843. Westwood, Trans. Ent. Soc. London, vol. 3, p. 240.

TYPE: *Evania minuta* Olivier, 1791. *Coranila* Leach MS. was validated and its type fixed by Westwood through citation as a synonym of *Brachygaster* Leach.

**Corynopria Haliday**

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. p. 170. Three species.

TYPE: *Diapria petiolaris* Nees, 1834. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 676.

Proposed as a subgenus of *Diapria* Latreille. Considered synonym of *Monelata* Foerster.

**Cracinopria Fouts**

1924. Fouts, Proc. Ent. Soc. Washington, vol. 26, p. 162. Four species.

TYPE: *Trichopria marylandica* Fouts, 1920, by original designation.

**Crassifoenus Crosskey**

1953. Crosskey, Trans. Roy. Ent. Soc. London, vol. 104, p. 355. Two species.

TYPE: *Gasteruption macronyx* Schletterer, 1889, by original designation.

**Cremastobaeus Ashmead**

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 210, 211, 228.

TYPE: *Cremastobaeus bicolor* Ashmead, 1893, by monotypy and original designation.

**Cremastoscielio Dodd**

1913. Dodd, Trans. Roy. Soc. South Australia, vol. 37, pp. 131, 156. Two species.

TYPE: *Cremastoscielio flavipes* Dodd, 1913, by original designation.

**(Crestoteleia Kieffer, error) = Chrestoteleia Kieffer**

1916. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 14, p. 181.

**Cryptoserphus Kieffer**

1907. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 288. No species.

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 313. Ten species.

TYPE: *Serphus (Cryptoserphus) longicalcar* Kieffer, 1908. Designated by Kieffer, 1908, in André, Spec. Hym. Eur. Alg., vol. 10, p. 314.

Proposed as a subgenus of *Serphus* Schrank.

**Cyathopria** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 383.

TYPE: *Cyathopria rufiventris* Kieffer, 1909, by monotypy.

**Cyphacolus** Priesner

1951. Priesner, Bull. Inst. Fouad I du Desert, vol. 1, p. 123.

TYPE: *Cyphacolus veniprivus* Priesner, 1951, by monotypy and original designation.

**(Dasynopeas** Apstein, error) = **Disynopeas** Kieffer

1926. Apstein, in Kieffer, Das Tierreich, Lief. 48, p. 879.

**Dendrocerus** Ratzeburg

1852. Ratzeburg, Die Ichneumonien der Forstinsecten . . . , vol. 3, p. 180.

TYPE: *Dendrocerus lichtensteini* Ratzeburg, 1852, by monotypy.

**(Deraiodontus** Bradley, error) = **Deraiodontus** Bradley

1901. Bradley, Trans. Amer. Ent. Soc., vol. 27, p. 321.

**Deraiodontus** Bradley (= *Deraiodontus* Bradley, error)

1901 (May). Bradley, Ent. Student, vol. 2, p. 29. No species.

1901 (Nov.). Bradley, Trans. Amer. Ent. Soc., vol. 27, pp. 320, 321. Three species.

TYPE: *Aulacus montanus* Cresson, 1879. Designated by Bradley, Trans. Amer. Ent. Soc., vol. 27, p. 321.

Considered synonym of *Pristaulacus* Kieffer.

**Diapria** Latreille (= *Tropidopria* Ashmead; *Tropidobria* Brèthes, error)

1796. Latreille, Précis caractères généraux des insectes . . . , p. 110. No species.

1802. Latreille, Histoire naturelle . . . crustacés et des insectes, vol. 3, p. 309. Two species.

TYPE: *Ichneumon conicus* Fabricius, 1775. Designated by Latreille, 1810, Considérations générales . . . des insectes, p. 436. *Psilus elegans* Jurine, 1807, designated by Westwood, 1839, An introduction to the modern classification of insects, vol. 2, generic synopsis p. 75, and *Diapria verticillata* Latreille, 1805, designated by Ashmead, 1903, Journ. New York Ent. Soc., vol. 11, p. 31, were included later.

Isogenotypic with *Tropidopria* Ashmead.

**Dibaryconus** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxvii, 270, 486. Four species.

TYPE: *Baryconus sordidus* Dodd, 1914, by original designation.

**Dichacantha** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, pp. 118, 147.  
TYPE: *Trimorus luteus* Cameron, 1888, by monotypy and original designation.

Considered synonym of *Scelio* Latreille.

**Dichogmus** Thomson (= *Dichognus* Ashmead, error)

1858. Thomson, Öfv. Vet.-Akad. Förh., vol. 15, pp. 287, 301.  
TYPE: *Dichogmus dimidiatus* Thomson, 1858, by monotypy.

**(Dichognus** Ashmead, error) = **Dichogmus** Thomson

1903. Ashmead, Journ. New York Ent. Soc., vol. 11, p. 34.

**Dichoteleas** Kieffer

1907. Kieffer, Berliner Ent. Zeitschr., vol. 51, p. 297.  
TYPE: *Dichoteleas rugosus* Kieffer, 1907, by monotypy.

**Dicroscelio** Kieffer

1913. Kieffer, in Alluaud and Jeannel, Voyage . . . en Afrique Orientale. Insectes hyménoptères, vol. 1, p. 16.

TYPE: *Dicroscelio flavipes* Kieffer, 1913, by monotypy and original designation.

**Dicroteleia** Kieffer

1908. Kieffer, Leyden Mus. Notes, vol. 30, p. 92.  
TYPE: *Dicroteleia rugosa* Kieffer, 1908, by monotypy.

**Digalesus** Kieffer

1914. Kieffer, Philippine Journ. Sci., vol. 9, p. 303.  
TYPE: *Digalesus flavipes* Kieffer, 1914, by monotypy and original designation.

**Dilapitha** Kieffer

1914. Kieffer, Philippine Journ. Sci., vol. 9, p. 293. Two species.  
TYPE: *Dilapitha albipes* Kieffer, 1914, by original designation.

**Dilobopria** Kieffer

1914. Kieffer, Insecta, vol. 4, p. 190.  
TYPE: *Dilobopria rufipes* Kieffer, 1914, by monotypy and original designation.

**Diphora** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 130, 140.  
TYPE: *Diphora westwoodii* Foerster, 1856, by monotypy.

**Diphoropria** Kieffer

1905. Kieffer, Bull. Soc. Hist. Nat. Metz, vol. 24, p. 103.  
TYPE: *Diphoropria rufipes* Kieffer, 1905, by monotypy.

**Diplatygaster** Kieffer

1926. Kieffer, *Das Tierreich*, Lief. 48, pp. xxxvi, 563, 843.

TYPE: *Platygaster formicarius* Kieffer, 1904, by monotypy.

**Disaulacinus** Kieffer

1910. Kieffer, *Bull. Soc. Ent. France*, 1910, p. 350. No species.

1911. Kieffer, *Ann. Soc. Ent. France*, vol. 80, pp. 214, 224. One species.

TYPE: *Disaulacinus flavimanus* Kieffer, 1911. First included species.

**Discelio** Kieffer

1908. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 32, pp. 116, 124. Two species.

TYPE: *Scelio thoracicus* Ashmead, 1900. Designated by Kieffer, 1926, *Das Tierreich*, Lief. 48, p. 343.

Considered synonym of *Scelio* Latreille.

**Disognus** Foerster

1856. Foerster, *Hymenopterologische Studien*, Heft 2, p. 99.

TYPE: *Proctotrupes areolator* Haliday, 1839, by monotypy.

**Disphaeron** Dahlbom

1837. Dahlbom, *Isis*, Heft 3, column 175.

TYPE: *Aulacus arcticus* Dahlbom, 1837 (= *Disphaeron arcticus* Dahlbom, 1837), by monotypy.

Dahlbom described *Aulacus arcticus* from the Zetterstedt collection and stated that it had been labeled *Disphaeron arcticus* by Zetterstedt. Thus he validated the name *Disphaeron*.

Considered synonym of *Aulacus* Jurine.

**Dissacolus** Kieffer

1926. Kieffer, *Das Tierreich*, Lief. 48, pp. xvii, 133, 155.

TYPE: *Acolus bidentatus* Dodd, 1914, by monotypy.

**Dissolcoides** Dodd

1913. Dodd, *Arch. Naturg.*, vol. 79, A6, p. 179.

TYPE: *Dissolcoides exsertus* Dodd, 1913, by monotypy and original designation.

**Dissolcus** Ashmead

1893. Ashmead, *U. S. Nat. Mus. Bull.* 45, pp. 138, 164.

TYPE: *Dissolcus nigricornis* Ashmead 1893, by monotypy and original designation.

**Dissoxylabis** Kieffer

1909. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 33, p. 342.

TYPE: *Dissoxylabis hirtipes* Kieffer, 1909, by monotypy.

**Disynopeas** Kieffer (= *Dasynopeas* Apstein, error)

1916. Kieffer, Zentralbl. Bakt. Parasitenk. Infektionskr.,  
Abt. 2, vol. 46, p. 591.

TYPE: *Disynopeas lasiopterae* Kieffer, 1916, by monotypy  
and original designation.

**Doddiella** Kieffer

1913. Kieffer, Bol. Lab. Zool. Portici, vol. 7, p. 109.

TYPE: *Doddiella nigriceps* Kieffer, 1913, by monotypy and  
original designation.

**Dolichofoenus** Kieffer

1910. Kieffer, Ann. Soc. Ent. France, vol. 79, p. 77. Two  
species.

TYPE: *Foenus raphidioides* Westwood, 1851, by original  
designation.

**Dolichopria** Kieffer

1910. Kieffer, in Wissenschaftliche Ergebnisse der deut-  
schen Zentral-Afrika-Expedition 1907-08 . . . , vol. 3  
(Zool.), Lief. 2, p. 99.

TYPE: *Dolichopria gracilis* Kieffer, 1910, by monotypy.

**Dolichotrypes** Crawford and Bradley

1911. Crawford and Bradley, Proc. Ent. Soc. Washington,  
vol. 13, p. 124.

TYPE: *Dolichotrypes hopkinsi* Crawford and Bradley, 1911,  
by monotypy and original designation.

Considered synonym of *Leptacis* Foerster.

**Doliopria** Kieffer

1910. Kieffer, Ent. Rundschau, vol. 27, No. 8, p. 48. No  
species.

1910. Kieffer, Ent. Rundschau, vol. 27, No. 9, p. 54. One  
species.

TYPE: *Doliopria flavipes* Kieffer, 1910. First included species.

**Dorachia**, new name (= *Rhacodia* Herrich-Schaeffer, 1838, not  
Huebner [1825])

TYPE: *Diapria striolata* Nees, 1834, by substitution of  
*Dorachia* for *Rhacodia* Herrich-Schaeffer.

**Duarina** Dodd

1926. Dodd, Proc. Linn. Soc. New South Wales, vol. 51, p.  
377.

TYPE: *Duarina venustella* Dodd, 1926, by monotypy and  
original designation.

**Duta** Nixon

1933. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 12, p. 306.

TYPE: *Holoteleia tenuicornis* Dodd, 1920, by monotypy and  
original designation.

**Dyscritobaeus** Perkins

1910. Perkins, Fauna Hawaiiensis, vol. 2, p. 621.

TYPE: *Dyscritobaeus comitans* Perkins, 1910, by monotypy.

**Eccinetus**, new name (= *Procinetus* Kieffer, 1910, not Foerster, 1868).

TYPE: *Cinetus radiatus* Kieffer, 1905, by substitution of *Eccinetus* for *Procinetus* Kieffer.

**Echinoteleas** Risbec

1954. Risbec, Bull. Inst. Français d'Afrique Noire, vol. 16, p. [1035].

TYPE: *Echinoteleas auratus* Risbec, by monotypy and original designation.

**Ecitonetes** Brues

1902. Brues, Amer. Nat., vol. 36, p. 370.

TYPE: *Ecitonetes subapterus* Brues, 1902, by monotypy.

**Ectadius** Foerster (= *Polymecus* Foerster)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 108, 113, 114, 144.

TYPE: *Platygaster craterus* Walker, 1835, by monotypy.  
Considered synonym of *Leptacis* Foerster.

**Eflatounina** Priesner

1951. Priesner, Bull. Inst. Fouad I du Desert, vol. 1, p. 126.

TYPE: *Eflatounina gryontoides* Priesner, 1951, by monotypy and original designation.

**(Elaptus** Forbes) = *Amitus* Haldeman

1885. Forbes, Rep. Illinois State Ent., No. 14, 1884, p. 110.

TYPE: *Elaptus aleurodis* Forbes, 1885, by monotypy.

Isogenotypic with *Amitus* Haldeman through synonymy.

Evidently error for mymarid genus *Alaptus* Westwood to which Forbes wrongly considered his species to belong. Moreover, *Elaptus* Forbes is preoccupied by *Elaptus* Pascoe, 1867, in Coleoptera.

**Electrofoenus** Cockerell [fossil]

1917. Cockerell, Amer. Journ. Sci., vol. 44, p. 364.

TYPE: *Electrofoenus gracilipes* Cockerell, 1917, by monotypy and original designation.

**Electroteleia** Brues [fossil]

1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 74, p. 80.

TYPE: *Electroteleia stigmatica* Brues, 1940, by monotypy and original designation.

**Elgonia** Risbec

1950. Risbec, Contribution à l'étude des Proctotrupidae. Trav. Lab. Ent. Sect. Soudan. Rech. Agron., p. 549.

TYPE: *Elgonia maxillosa* Risbec, 1950, by monotypy.

Considered synonym of *Apegus* Foerster.

**Elysoceraphron** Szelényi

1936. Szelényi, Ann. Mus. Nat. Hungarici, vol. 30, Zool., p. 64.

TYPE: *Elysoceraphron hungaricus* Szelényi, 1936, by monotypy and original designation.

**Embidobia** Ashmead

1895. Ashmead, Journ. Trinidad Field Nat. Club, vol. 2, p. 264.

TYPE: *Embidobia urichi* Ashmead, 1895, by monotypy.

**Encyrtoscelio** Dodd

1914. Dodd, Proc. Roy. Soc. Queensland, vol. 26, p. 119.

TYPE: *Encyrtoscelio mirissimus* Dodd, 1914, by monotypy and original designation.

**Enneascelio** Kieffer

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 293.

TYPE: *Enneascelio exaratus* Kieffer, 1910, by monotypy and original designation.

Again described as new by Kieffer, 1912, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 54.

**Entomacis** Foerster (= *Glyphidopria* Haliday)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 121, 123. No species.

1873. Marshall, A catalogue of British Hymenoptera; Oxyura, p. 10. Two species.

TYPE: *Diapria* (*Glyphidopria*) *platyptera* Haliday, 1857. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 673. *Hemilexis* (*Entomacis*) *cordata* Kieffer, 1911, designated by Kieffer, 1916, Das Tierreich, Lief. 44, p. 40, was included later.

Isogenotypic with *Glyphidopria* Haliday.

**Entomia** Herrich-Schaeffer (= *Entomius* Haliday, emendation)

1840. Herrich-Schaeffer, Nomenclator entomologicus, Heft 2, p. 127; pl. 6, fig. 21.

TYPE: *Entomia campanulata* Herrich-Schaeffer, 1840, by monotypy.

Considered synonym of *Ismarus* Haliday.

**(Entomius** Haliday, emendation) = **Entomia** Herrich-Schaeffer

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. p. 169.

Designation of *Ismarus halidayi* Foerster, 1850, as type by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 681, was unnecessary since *Entomia* Herrich-Schaeffer is a valid name and not a nomen nudum as previously supposed.



**Entomopria** Kieffer

1912. Kieffer, in Wytsman, *Genera insectorum*, fasc. 124, 1911, pp. 4, 18.

TYPE: *Corynopria solida* Thomson, 1858, by monotypy.

Again described as new by Kieffer, 1912, *Trans. Linn. Soc. London*, ser. 2, Zool., vol. 15, p. 68.

**Eoxestonotus** Debauche

1947. Debauche, *Bull. Ann. Soc. Ent. Belgique*, vol. 83, p. 267. Two species.

TYPE: *Eoxestonotus pini* Debauche, 1947, by original designation.

Considered synonym of *Xestonotidea* Gahan.

**(Epimeces** Westwood, not Billberg, 1820) = **Parepimeces** Kieffer

1833. Westwood, *Mag. Nat. Hist.*, vol. 6, p. 421. Two species.

TYPE: *Epimeces ensifer* Westwood, 1833, by elimination. In establishing *Sactogaster*, Foerster, 1856, *Hymenopterologische Studien*, Heft 2, p. 113, transferred to it *Epimeces ventralis* Westwood, 1833, leaving only *ensifer* Westwood in *Epimeces*.

**Epinomus** Ghesquière

1948. Ghesquière, *Rev. Zool. Bot. Africaines*, vol. 40, p. 324.

TYPE: *Epinomus anoplocnemidis* Ghesquière, 1948, by monotypy and original designation.

**(Episceuastes** Gistel) = **Pelecinius** Latreille

1848. Gistel, *Naturgeschichte des Tierreichs*, p. x.

TYPE: *Ichneumon polycerator* Fabricius, 1776, by substitution of *Episceuastes* for *Pelecinius* Latreille.

Unnecessarily proposed for *Pelecinius* Latreille which Gistel thought to be preoccupied by *Pelecinius* Moench in botany.

**Eremioscelio** Priesner

1951. Priesner, *Bull. Inst. Fouad I du Desert*, vol. 1, p. 129.

TYPE: *Eremioscelio cydnoides* Priesner, 1951, by monotypy and original designation.

**(Eriodorus** Agassiz, error) = **Erodorus** Walckenaer

1846. Agassiz, *Nomenclator zoologicus. Index universalis*, p. 143.

**Eriopria** Kieffer

1910. Kieffer, in André, *Spec. Hym. Eur. Alg.*, vol. 10, pp. 693, 744. Three species, one of them doubtfully included.

TYPE: *Eriopria nigra* Kieffer, 1910. Designated by Kieffer, 1912, in Wytsman, *Genera insectorum*, fasc. 124, 1911, p. 31.

**Eritrissomerus** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 263, 264, 298.

TYPE: *Eritrissomerus cecidomyiae* Ashmead, 1893, by monotypy and original designation.

**Erodorus** Walckenaer (= *Eriodoros* Agassiz, error)

1802. Walckenaer, Faune Parisienne, Insectes, vol. 2, p. 47.

TYPE: *Erodorus bimaculatus* Walckenaer, 1802, by monotypy.

Considered synonym of *Proctotrupes* Latreille.

**Eufoenus** Szépligeti

1903 (August). Szépligeti, Ann. Mus. Nat. Hungarici, vol. 1, p. 365. Four species.

TYPE: *Gasteruption antennale* Schletterer, 1889. Designated by Bradley, 1909, Deut. Ent. Zeitschr., 1909, p. 38.

Considered synonym of *Hyptiogaster* Kieffer, which was published Mar. 6, 1903.

**Euhoplopria** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 399, 414. Three species.

TYPE: *Euhoplopria carinatifrons* Dodd, 1915, by original designation.

**Eulagynodes** Girault

1917. Girault, New Javanese Hymenoptera, p. 9. (Private publication.)

TYPE: *Eulagynodes bicolor* Girault, 1917, by monotypy and original designation.

**(Eumegaspilus** Schulz, emendation) = **Eumegaspilus** Ashmead

1906. Schulz, Spolia hymenopterologica, p. 152.

**Eumegaspilus** Ashmead (= *Eumegaspilus* Schulz, emendation)

1888. Ashmead, Can. Ent., vol. 20, pp. 43, 49. Two species.

TYPE: *Eumegaspilus canadensis* Ashmead, 1888. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 670. *Eumegaspilus erythrothorax* Ashmead, 1893, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 119, was not originally included (see *Conostigmoides* Dodd).

Considered synonym of *Conostigmus* Dahlbom.

**Eumicrosoma** Gahan

1913. Gahan, Proc. U. S. Nat. Mus., vol. 46, p. 442.

TYPE: *Eumicrosoma beneficum* Gahan, 1913, by monotypy and original designation.

**Euplacopria** Ferrière

1929. Ferrière, Zool. Anz., vol. 82, p. 157.

TYPE: *Euplacopria mutilata* Ferrière, 1929, by monotypy.

**Eurostemma** Szelényi

1938. Szelényi, Folia Ent. Hungarica, vol. 3, pp. 99, 102.

TYPE: *Inostemma europus* Walker, 1838, by monotypy and original designation.

**Euxestonotus** Fouts

1925. Fouts, Proc. Ent. Soc. Washington, vol. 27, p. 98.  
Four species.

TYPE: *Platygaster error* Fitch, 1861, by original designation.  
Considered synonym of *Xestonotidea* Gahan.

**Evania** Fabricius (= *Evenia* Schletterer, error)

1775. Fabricius, Systema entomologiae . . . , p. 345. Two species.

TYPE: (*Sphex appendigaster* Fabricius) = *Ichneumon appendigaster* Linnaeus. Designated by Latreille, 1810, Considérations générales . . . des insectes, p. 436.

**Evaniella** Bradley

1905. Bradley, Can. Ent., vol. 37, p. 64. Three species.

TYPE: (*Evania unicolor* Ashmead, 1901, not Say, 1859) = *Evaniella semaeoda* Bradley, 1908, by original designation.

**Evaniellus** Enderlein

1905. Enderlein, Zool. Anz., vol. 28, pp. 700, 701, 714. Four species.

TYPE: *Evaniellus peruanus* Enderlein, 1905, by original designation.

**Evaniscus** Szépligeti

1903. Szépligeti, Ann. Mus. Nat. Hungarici, vol. 1, pp. 375, 378.

TYPE: *Evaniscus tibialis* Szépligeti, 1903, by monotypy.

**(*Evenia* Schletterer, error) = *Evania* Fabricius**

1889. Schletterer, Ann. Naturh. Hofmus., vol. 4, p. 118.

**Exallonyx** Kieffer

1904. Kieffer, Bull. Soc. Hist. Nat. Metz, vol. 23, p. 34.  
Eleven species.

TYPE: *Exallonyx formicarius* Kieffer, 1904, by original designation.

Considered synonym of *Codrux* Panzer.

**Fahringeria** Kieffer

1921. Kieffer, Brotéria, Rev. Luso-Braziliera, Zool. ser., vol. 19, p. 68.

TYPE: *Fahringeria synergorum* Kieffer, 1921, by monotypy.

**(*Foenus* Couper, error) = *Foenus* Fabricius**

1870. Couper, Can. Ent., vol. 2, p. 110.

**Fidiobia** Ashmead (= *Rosneta* Brues)

1894. Ashmead, Journ. Cincinnati Soc. Nat. Hist., vol. 17, p. 170.

TYPE: *Fidiobia flavipes* Ashmead, 1894, by monotypy.

Isogenotypic with *Rosneta* Brues through synonymy.

(*Foenus* Fabricius; *Feonus* Couper, error; *Phoenus* Schletterer, emendation) = **Gasteruption** Latreille

1798. Fabricius, Supplementum entomologiae systematicae, pp. 210, 240. Two species.

TYPE: *Ichneumon jaculator* Linnaeus, 1758. Designated by Latreille, 1810, Considérations générales . . . des insectes, p. 436. *Ichneumon assectator* Linnaeus, 1758, was designated by Curtis, 1832, British entomology, folio 423.

Isogenotypic with *Gasteruption* Latreille.

**Fusicornia** Risbec

1950. Risbec, Contribution à l'étude des Proctotrupidae. Trav. Lab. Ent. Sect. Soudan. Rech. Agron., p. 606.

TYPE: *Fusicornia bambeyi* Risbec, 1950, by monotypy.

**Galesimorpha** Brues [fossil]

1910. Brues, Bull. Comp. Zool., vol. 54, p. 12.

TYPE: *Galesimorpha wheeleri* Brues, 1910, by monotypy and original designation.

(*Galesus* Haliday) = **Psilus** Panzer

1829. Haliday, in Curtis, A guide to an arrangement of British insects, column 108.

TYPE: *Psilus cornutus* Panzer, 1801, by monotypy. Three names were listed but two of them were nomina nuda. *Galesus fuscipennis* Curtis, 1831, designated by Kieffer, 1912, in Wytzman, Genera insectorum, fasc. 124, 1911, p. 40, was one of the nomina nuda.

Isogenotypic with *Psilus* Panzer.

(*Gasteruption* Dominique, emendation) = **Gasteruption** Latreille

1893. Dominique, Bull. Soc. Sci. Nat. Ouest France, vol. 3, p. 198.

**Gasteruption** Latreille (= *Gastryptium* Agassiz, *Gasteryption* Schletterer, *Gasteruption* Dominique, *Hyptiagaster* Dominique, *Gasteruption* Schulz, *Gastrhyption* Schulz, emendations; *Foenus* Fabricius)

1796. Latreille, Précis caractères génériques des insectes, p. 113. No species.

1802. Latreille, Histoire naturelle . . . crustacés et des insectes, vol. 3, p. 329. Two species by bibliographic reference through synonymy of *Gasteruption* with *Foenus*.

TYPE: *Ichneumon jaculator* Linnaeus, 1758. Designated by Westwood, 1839, An introduction to the modern classification of insects, vol. 2, generic synopsis p. 56, in his treatment of *Foenus* Fabricius. *Ichneumon assecutor* Linnaeus, 1758, was designated by Mani, 1939, Catalogue of Indian insects, pt. 24, Evaniidae, p. 9.

Isogenotypic with *Foenus* Fabricius.

(*Gasteruption* Schulz, emendation) = *Gasteruption* Latreille 1906. Schulz, Spolia hymenopterologica, p. 133.

(*Gasteryption* Schletterer, emendation) = *Gasteruption* Latreille 1890. Schletterer, Ann. Naturh. Hofmus., vol. 4, p. 375.

(*Gastrhyption* Schulz, emendation) = *Gasteruption* Latreille 1911. Schulz, Zool. Ann., vol. 4, p. 55.

#### **Gastrotrypes** Brues

1922. Brues, Proc. Amer. Acad. Arts Sci., vol. 57, p. 270. Two species.

TYPE: *Gastrotrypes spatulatus* Brues, 1922, by original designation.

#### **Gastroxylabis** Ogloblin

1954. Ogloblin, Rev. Chilena Ent., vol. 3, 1953, p. 69. Two species.

TYPE: *Gastroxylabis rubrosignata* Ogloblin, 1954, by original designation.

(*Gastryptium* Agassiz, emendation) = *Gasteruption* Latreille

1846. Agassiz, Nomenclator zoologicus. Index universalis, pp. 159, 160.

#### **Geodiapria** Kieffer

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 707. No species.

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 896. One species.

TYPE: *Geodiapria longiceps* Kieffer, 1911. First included species.

#### **Gita** Nixon

1933. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 12, p. 309.

TYPE: *Gita infortunata* Nixon, 1933, by monotypy and original designation.

(*Glyphidopria* Haliday) = *Entomacis* Foerster

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. p. 172. Two species.

TYPE: *Diapria* (*Glyphidopria*) *platyptera* Haliday, 1857. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 673.

Proposed as a subgenus of *Diapria* Latreille. Isogenotypic with *Entomacis* Foerster.

**Glyptonota** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 122, 126. No species.

1893. Ashmead, U. S. Nat. Mus. Bull. 45, p. 401. One species.

TYPE: *Glyptonota nigriclavata* Ashmead, 1893. First included species.

**Glyptoteleia** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxvii, 272, 487.

TYPE: *Baryconus bisulcatus* Kieffer, 1910, by monotypy.

**Govinda** Nixon

1933. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 12, p. 465. Five species.

TYPE: *Govinda mila* Nixon, 1933, by original designation.

**Gryon** Haliday

1833. Haliday, Ent. Mag., vol. 1, p. 271.

TYPE: *Gryon misellus* Haliday, 1833, by monotypy.

**Gryonella** Dodd

1914. Dodd, Trans. Roy. Soc. South Australia, vol. 38, pp. 75, 84. Two species.

TYPE: *Gryonella crawfordi* Dodd, 1914, by original designation.

**Gryonoides** Dodd

1920. Dodd, Trans. Ent. Soc. London, 1919, p. 360. Three species.

TYPE: *Gryonoides pulchellus* Dodd, 1920, by original designation.

**(Habropelta Schulz, emendation) = Habropelte Thomson**

1906. Schulz, Spolia hymenopterologica, p. 152.

**Habropelte Thomson (= Habropelta Schulz, emendation)**

1858. Thomson, Öfv. Vet.-Akad. Förh., vol. 15, pp. 287, 288. Three species.

TYPE: *Ceraphron tibialis* Boheman, 1832. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 104. *Ceraphron scutellaris* Boheman (cited as "Dalb." by Ashmead), 1832, of which *Ceraphron tibialis* Boheman is considered a synonym, was designated by Ashmead, 1903, Journ. New York Ent. Soc., vol. 11, p. 33.

Considered synonym of *Megaspilus* Westwood.

**Habroteleia** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2 (42), p. 14.

TYPE: *Habroteleia flavipes* Kieffer, 1905, by monotypy.

(*Hadroceras* Foerster, new name for *Calliceras* Nees) = *Ceraphron* Jurine

1840. Foerster, Beiträge zur Monographie der Pteromalinen Nees, Heft 1, p. xliv.

TYPE: *Calliceras sulcatus* Nees, 1834, by substitution of *Hadroceras* for *Calliceras* Nees.

Unnecessarily proposed for *Calliceras* Nees, 1834, which Foerster thought to be preoccupied by *Callicera* Meigen, 1822.

**Hadronotellus** Kieffer

1917. Kieffer, Ent. Medd., ser. 2, vol. 11, p. 341.

TYPE: *Hadronotellus pedester* Kieffer, 1917, by monotypy and original designation.

Considered synonym of *Hadronotus* Foerster.

**Hadronotoides** Dodd

1913. Dodd, Arch. Naturg., vol. 79, A6, p. 171.

TYPE: *Hadronotoides pentatomus* Dodd, 1913, by monotypy and original designation.

**Hadronotus** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 101, 105. Two names, both nomina nuda.

1861. Foerster, Programm der Realschule zu Aachen . . . , p. xli. One species.

TYPE: *Hadronotus exsculptus* Foerster, 1861. First included species. *Hadronotus laticeps* Foerster, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 229, was a nomen nudum. *Hadronotus laticeps* Kieffer, 1908, designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 453, and by Mani, 1941, Catalogue of Indian insects, Pt. 26, Serphoidea, p. 26, was included later.

**Hadrophanurus** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xvi, 15, 130.

TYPE: *Telenomus* (?) *pennsylvanicus* Ashmead, 1893, by monotypy and original designation.

**Helorus** Latreille

1802. Latreille, Histoire naturelle . . . crustacés et des insectes, vol. 3, p. 309.

TYPE: *Helorus ater* Latreille, 1802, by monotypy. *Sphex anomalipes* Panzer, 1798, of which *Helorus ater* is

considered a synonym, was designated by Latreille, 1810, *Considérations générales . . . des insectes*, p. 437.

#### **Hemifoenus** Kieffer

1911. Kieffer, *Ann. Soc. Ent. France*, vol. 80, pp. 177, 182.

TYPE: *Hemifoenus brevithorax* Kieffer, 1911, by monotypy.

#### **Hemigalesus** Kieffer

1913. Kieffer, *Insecta*, vol. 3, p. 433. Four species.

TYPE: *Hemigalesus niger* Kieffer, 1913, by original designation.

#### **Hemilexis** Foerster

1856. Foerster, *Hymenopterologische Studien*, Heft 2, pp. 122, 123, 127. No species.

1887. Ashmead, *Can. Ent.*, vol. 19, p. 196. One species.

TYPE: *Hemilexis (mellipetiola!) = mellipetiolata* Ashmead, 1887. First included species. *Diapria (Glyphidopria) platyptera* Haliday, 1857, designated by Ashmead, 1903, *Journ. New York Ent. Soc.*, vol. 11, p. 29, and *Hemilexis californica* Ashmead, 1893, designated by Kieffer, 1916, *Das Tierreich*, Lief. 44, p. 46, were included later.

#### **Hemilexodes** Ashmead

1893. Ashmead, *U. S. Nat. Mus. Bull.* 45, pp. 386, 399.

TYPE: *Hemilexodes floridana* Ashmead, 1893, by monotypy and original designation.

#### **Hemilexomyia** Dodd

1920. Dodd, *Proc. Linn. Soc. New South Wales*, vol. 45, p. 443.

TYPE: *Hemilexomyia abrupta* Dodd, 1920, by monotypy and original designation.

#### **Hemimorus** Cameron

1912. Cameron, *Soc. Ent.*, vol. 27, No. 17, p. 77.

TYPE: *Hemimorus clavicornis* Cameron, 1912, by monotypy. Considered synonym of *Trimorus* Foerster.

#### **Hemisius** Westwood

1833. Westwood, *London Edinburgh Phil. Mag.*, ser. 3, vol. 2, p. 445.

TYPE: *Hemisius minutus* Westwood, 1833, by monotypy.

#### **Heptascelio** Kieffer

1916. Kieffer, *Brotéria*, *Rev. Luso-Brazileira*, *Zool. ser.*, vol. 14, p. 58.

TYPE: *Heptascelio lugens* Kieffer, 1916, by monotypy and original designation.



**Heterogryon** Kieffer

1926. Kieffer, *Das Tierreich*, Lief. 48, pp. xxvi, 271, 446, 448. Fourteen species.

TYPE: *Plastogryon sagax* Kieffer, 1908, by present designation.

Proposed as a subgenus of *Plastogryon* Kieffer.

**Heteropria** Kieffer

1905. Kieffer, *Ann. Mus. Civ. Stor. Nat. Genova*, ser. 3, vol. 2 (42), pp. 23, 36.

TYPE: *Heteropria compressipes* Kieffer, 1905, by monotypy.

**Hexapria** Kieffer

1905. Kieffer, *Ann. Mus. Civ. Stor. Nat. Genova*, ser. 3, vol. 2 (42), pp. 27, 34.

TYPE: *Hexapria fuscoclavata* Kieffer, 1905, by monotypy.

**(Holacolus** Kieffer; *Holalcus* Risbec, error) = *Acolus* Foerster

1912. Kieffer, in André, *Spec. Hym. Eur. Alg.*, vol. 11, pp. 89, 94, 106. Two species.

TYPE: *Acolus opacus* Thomson, 1859, by present designation. Isogenotypic with *Acolus* Foerster.

**(Holalcus** Risbec, error for *Holacolus* Kieffer) = *Acolus* Foerster

1950. Contribution à l'étude des Proctotrupidae. *Trav. Lab. Ent. Sect. Soudan. Rech. Agron.*, p. 577.

**Holoteleia** Kieffer

1908. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 32, pp. 120, 169. Four species.

TYPE: *Baryconus (Holoteleia) bicolor* Kieffer, 1908, by original designation. This name, preoccupied by *Baryconus bicolor* Harrington, 1889, has been replaced by *Holoteleia nigriceps* Kieffer, 1926.

Proposed as a subgenus of *Baryconus* Foerster.

**Homophanurus** Kieffer

1912. Kieffer, in André, *Spec. Hym. Eur. Alg.*, vol. 11, pp. 9, 36.

TYPE: *Homophanurus hofmanni* Kieffer, 1912, by monotypy.

**Hoplogryon** Ashmead

1893. Ashmead, *U. S. Nat. Mus. Bull.* 45, pp. 181, 200. Nine species.

TYPE: *Prosacantha minutissima* Ashmead, 1887, by original designation.

Considered synonym of *Trimorus* Foerster.

**Hoplopria** Ashmead

1893. Ashmead, *U. S. Nat. Mus. Bull.* 45, pp. 385, 386, 388.

TYPE: *Hoplopria pulchripennis* Ashmead, 1893, by monotypy and original designation.

**Hoplopriella** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 399, 416. Thirteen species.

TYPE: *Hoplopriella bicoloricornis* Dodd, 1915, by original designation.

**(Hoploteleia** Ashmead) = **Baryconus** Foerster

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 210, 211, 227.

TYPE: *Baryconus floridanus* Ashmead, 1887, by monotypy and original designation.

Isogenotypic with *Baryconus* Foerster.

**(Hympocampsis** Szelényi, error) = **Hypocampsis** Foerster

1938. Szelényi, Folia Ent. Hungarica, vol. 3, p. 100.

**(Hyperbaeus** Foerster) = **Baeus** Haliday

1856. Foerster, Hymenopterologische Studien, Heft 2, p. 144.

TYPE: *Baeus seminulum* Haliday, 1833, by substitution of *Hyperbaeus* for *Baeus* Haliday.

Unnecessarily proposed for *Baeus* Haliday which Foerster thought to be preoccupied by *Baea* Commerson in botany.

**Hypocampsis** Foerster (= *Hympocampsis* Szelényi, error)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 108, 115. No species.

1859. Thomson, Öfv. Vet.-Akad. Förh., vol. 16, pp. 70, 82. Three species.

TYPE: *Hypocampsis hyalinata* Thomson, 1859. Designated by Ashmead, 1903, Journ. New York Ent. Soc., vol. 11, p. 98. *Hypocampsis compressicornis* Thomson, 1859, was designated by Debauche, 1947, Bull. Ann. Soc. Ent. Belgique, vol. 83, p. 283. Szelényi, 1938, Folia Ent. Hungarica, vol. 3, p. 100, maintained that the species upon which Foerster founded *Hypocampsis* is *Platygaster contorticornis* Ratzeburg, 1844, and that is apparently correct, but under a recent decision of the International Commission on Zoological Nomenclature (Bull. Zool. Nom., vol. 4, 1950, pp. 160, 346) only the first species to be included shall be eligible for designation as type of a genus originally published without included species.

**Hyptia** Illiger (= *Hyptiam* Schuckard, error)

1807. Illiger, in Rossi, Fauna Etrusca sistens Insecta, vol. 2, p. 82; Illiger, Magazin für Insektenkunde, vol. 6, p. 192.

TYPE: *Evania petiolata* Fabricius, 1793, by monotypy.

In Fauna Etrusca . . . Illiger wrote “. . . quod *Hyptiam* voco,” which accounts for Shuckard's adoption of that spelling.

(*Hyptiagaster* Dominique, emendation) = **Gasteruption** Latreille 1893. Dominique, Bull. Soc. Sci. Nat. Ouest France, vol. 3, p. 198.

(*Hyptiam* Shuckard, error) = **Hyptia** Illiger

1841. Shuckard, Entomologist, vol. 1, pp. 117, 120.

Illiger, 1807, in Rossi, Fauna Etrusca, II, p. 82, wrote “. . . quod *Hyptiam* voco,” which accounts for Shuckard's adoption of that spelling.

**Hyptiagaster** Kieffer

1903 (March 6). Kieffer, Bull. Soc. Ent. France, 1903, pp. 93, 94. Eleven species.

TYPE: *Gasteruption crassiceps* Schletterer, 1889, by original designation. *Gasteruption antennale* Schletterer, 1889, was designated by Kieffer later in 1903, in André, Spec. Hym. Eur. Alg., vol. 7 bis, p. 381, and *Foenus australis* Westwood, 1843, was designated by Crosskey, 1953, Trans. Ent. Soc. London, vol. 104, p. 363.

See *Eufoenus Szépligetii*, which was published August 1903 and is considered a synonym of *Hyptiagaster* Kieffer.

(*Hyptiagaster* Mani, error) = **Hyptiogastrites** Cockerell [fossil]

1939. Mani, Catalogue of Indian insects, Pt. 24, Evaniidae, p. 2.

**Hyptiogastrites** Cockerell (= *Hyptiagaster* Mani, error) [fossil]

1917. Cockerell, Ann. Ent. Soc. Amer., vol. 10, p. 19.

TYPE: *Hyptiogastrites electrinus* Cockerell, by monotypy.

**Idiotypa** Foerster (= *Mionopria* Haliday)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 122, 125. No species.

1873. Marshall, A catalogue of British Hymenoptera; *Oxyura*, p. 11. Two species.

TYPE: *Diapria (Mionopria) maritima* Haliday, 1857. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 403.

Isogenotypic with *Mionopria* Haliday.

**Idris** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 102, 105.

TYPE: *Idris flavicornis* Foerster, 1856, by monotypy. *Idris laeviceps* Ashmead, 1893, designated by Brues, 1908, in Wytzman, Genera insectorum, fasc. 80, p. 41, was included later.

**Immsia** Cameron

1913. Cameron, Indian For. Rec., 1912, vol. 4, p. 104.

TYPE: *Immsia carinifrons* Cameron, 1913, by monotypy.

Considered synonym of *Microphanurus* Kieffer.

**Inocerota** Szelényi

1939. Szelényi, Ann. Mus. Nat. Hungarici, vol. 32, Zool., p. 121. Two species.

TYPE: *Inocerota discessus* Szelényi, 1939, by original designation.

**Inostemma** Haliday

1833. Haliday, Ent. Mag., vol. 1, p. 270.

TYPE: *Psilus boscii* Jurine, 1807, by monotypy.

**Interaulacus** Bradley

1908. Bradley, Trans. Amer. Ent. Soc., vol. 34, pp. 120, 123.

TYPE: *Interaulacus kiefferi* Bradley, 1908, by monotypy and original designation.

**Iphitrachelus** Walker

1835. Walker, Ent. Mag., vol. 3, p. 273.

TYPE: *Iphitrachelus lar* Walker, 1835, by monotypy.

**Ismarus** Haliday

1835. Haliday, Ent. Mag., vol. 2, p. 467.

TYPE: *Cinetus dorsiger* Curtis, 1831, by monotypy.

**Isoeybus** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 108, 114. Five species, one of them included by bibliographic reference.

TYPE: *Platygaster grandis* Nees, 1834. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 327. *Platygaster cotta* Walker, 1835, was designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 709, through oversight of Ashmead's citation.

**Isolia** Foerster

1878. Foerster, Verh. Naturh. Ver. Preuss. Rheinl., vol. 35, p. 46. No species.

No species have been referred to this genus up to the present time. *Isolia*, accordingly, remains a genus dubium.

**Isorhombus** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 107, 112, 113. No species.

1887. Ashmead, Can. Ent., vol. 19, p. 129. One species.

TYPE: *Isorhombus hyalinipennis* Ashmead, 1887. First included species.

Considered synonym of *Platygaster* Latreille.

**(*Isostasis* Provancher, error) = *Isostasius* Foerster**

1887. Provancher, Additions et corrections au volume II de la fauna entomologique du Canada . . . Hyménoptères, p. 183.

***Isostasius* Foerster (= *Isostasis* Provancher, error)**

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 106, 109.

TYPE: *Platygaster punctiger* Nees, 1834, by monotypy.

***Kiefferopria* Brèthes**

1927. Brèthes, Ent. Mitt., vol. 16, p. 333.

TYPE: *Kiefferopria horni* Brèthes, 1927, by monotypy.

***Labidopria* Wasmann**

1925. Wasmann, Abh. Theor. Biol., Heft 19, p. 94.

TYPE: *Labidopria longicornis* Wasmann, 1925, by monotypy.

***Labolips* Haliday**

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. p. 173.

TYPE: *Labolips innupta* Haliday, 1857, by monotypy.

***Laches* Gistel (= *Anisoptera* Herrich-Schaeffer, 1840, not Berthold, 1827)**

1848. Gistel, Naturgeschichte des Thierreichs, p. viii.

TYPE: *Anisoptera egregia* Herrich-Schaeffer, 1840, by substitution of *Laches* for *Anisoptera* Herrich-Schaeffer.

Considered synonym of *Psilus* Panzer.

***Lagynodes* Foerster**

1840. Foerster, Beiträge zur Monographie der Pteromalinen Nees, Heft 1, p. xlv.

TYPE: *Lagynodes rufus* Foerster, 1840, by monotypy. *Ceraphron pallidus* Boheman, 1832, was designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 120.

***Lamproteleia* Kieffer**

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 293. Two species.

TYPE: *Lamproteleia fasciatipennis* Kieffer, 1910, by original designation.

Described again as new by Kieffer, 1912, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 63.

***Lapitha* Ashmead**

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 209, 211, 222.

TYPE: *Lapitha spinosa* Ashmead, 1893, by monotypy and original designation.

***Lapithoides* Nixon**

1933. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 12, p. 549.

TYPE: *Lapithoides semiramis* Nixon, 1933, by monotypy and original designation.

**Leaiopria** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 398, 426.

TYPE: *Leaiopria termitarii* Dodd, 1915, by monotypy and original designation.

**Ledouxopria** Risbec

1953. Risbec, Bull. Inst. Français d'Afrique Noire, vol. 15, p. 551.

TYPE: *Ledouxopria africana* Risbec, 1953, by monotypy.

**Lepidopria** Kieffer (= *Lepidoria* Brues, error)

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 695. No species.

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 869. One species.

TYPE: *Lepidopria pedestris* Kieffer, 1911. First included species.

(*Lepidoria* Brues, not Mulsant and Verreaux, 1866, error) =

**Lepidopria** Kieffer

1916. Brues, Psyche, vol. 23, p. 126.

**Lepidoscelio** Kieffer

1905. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 29, p. 129.

TYPE: *Lepidoscelio fuscipennis* Kieffer, 1905, by monotypy.

**Leptacis** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 107, 112. Four species.

TYPE: *Ichneumon tipulae* Kirby, 1798. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 270.

**Leptorhaptus** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 129, 137. No species.

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 346, 349. Two species.

TYPE: *Leptorhaptus conicus* Ashmead, 1893. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 685. *Leptorhaptus abbreviatus* Foerster, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 350, was a nomen nudum.

**Leptoteleia** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, pp. 120, 163.

TYPE: *Baryconus oecanthi* Riley, in Ashmead, 1893, by monotypy through bibliographic reference.

(*Limnodytes* Marchal, not Duméril and Bibron, 1841) = *Tiphodytes* Bradley

1900. Marchal, Bull. Soc. Ent. France, 1900, p. 328. Nomen nudum.

1900. Marchal, Ann. Soc. Ent. France, vol. 69, pp. 172, 174.

TYPE: *Limnodytes gerriphagus* Marchal, 1900, by monotypy and original designation.

(*Linkia* Kieffer, error, not Nardo, 1834) = *Linkiola* Kieffer

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 702.

*Linkiola* Kieffer (= *Linkia* Kieffer, error, 1910, not Nardo, 1834)

1910. Kieffer, Ent. Rundschau, vol. 27, p. 39.

TYPE: *Linkiola crassipes* Kieffer, 1910, by monotypy.

In 1912, in Wytzman, Genera insectorum, fasc. 124, 1911, p. 26, Kieffer shows *Linkia* to have been an error for *Linkiola* Kieffer. It was inadvertently published before *Linkiola* but is preoccupied.

*Liophanurus* Kieffer

1912. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 9, 61. Thirty-seven species, three of them doubtfully included.

TYPE: *Telenomus pilosomatis* Ashmead, 1893, by original designation.

Considered synonym of *Telenomus* Haliday.

*Lipoglyptus* Crawford

1910. Crawford, Proc. U. S. Nat. Mus., vol. 38, p. 123.

TYPE: *Lipoglyptus primus* Crawford, 1910, by monotypy and original designation.

*Lithobelyta* Cockerell [fossil]

1921. Cockerell, Ann. Mag. Nat. Hist., ser. 9, vol. 7, p. 22.

TYPE: *Lithobelyta reducta* Cockerell, 1921, by monotypy.

(*Loboscelidea* Rye, emendation) = *Loboscelidia* Westwood

1876. Rye, in Zool. Rec., vol. 11, 1874, p. 552.

*Loboscelidia* Westwood (= *Loboscelidoidea* Rye, emendation; *Loboscelidea* Rye, emendation)

1874. Westwood, Thesaurus entomologicus oxoniensis . . . , p. 171.

TYPE: *Loboscelidia rufescens* Westwood, 1874, by monotypy.

(*Loboscelidoidea* Rye, emendation) = *Loboscelidia* Westwood

1876. Rye, in Zool. Rec., vol. 11, 1874, pp. 365, 552.

*Lophopria* Kieffer

1910. Kieffer, in Wissenschaftliche Ergebnisse der deutschen Zentral-Afrika-Expedition 1907-08 . . . , vol. 3 (Zool.), Lief. 2, p. 97.

TYPE: *Lophopria crassiclava* Kieffer, 1910, by monotypy.

(*Loxoptera* Mani, error) = *Loxotropa* Foerster

1941. Mani, Catalogue of Indian insects, Pt. 26, Serphoidea, pp. 39, 40, 41, 47.

*Loxotropa* Foerster (= *Loxoptera* Mani, error)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 122, 123, 126.

TYPE: *Psilus antennatus* Jurine, 1807, by monotypy. *Loxotropa acolutha* Foerster, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 412, was a nomen nudum. *Diapria dispar* Nees, 1834, was designated by Mani, 1941, Catalogue of Indian insects, Pt. 26, Serphoidea, p. 47.

*Lygocerus* Foerster (= *Cheirocerus* Ashmead, error)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 97, 99. No species.

1868. Marshall, Ent. Monthly Mag., vol. 5, p. 158. Five species.

TYPE: *Ceraphron ramicornis* Boheman, 1832. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 107.

*Lyteba* Thomson

1858. Thomson, Öfv. Vet.-Akad. Förh., vol. 15, pp. 155, 180.

TYPE: *Belyta bisulca* Nees, 1834, by monotypy.

Considered synonym of *Oxylabis* Foerster.

*Macrogyron* Nixon

1936. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 17, No. 97, p. 116. No species.

1936. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 17, No. 98, p. 185. Three species.

TYPE: *Macrogyron pluto* Nixon, 1936. Designated by Nixon, 1936, Ann. Mag. Nat. Hist., ser. 10, vol. 17, No. 98, p. 185.

*Macrohynniss* Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 129, 136. No species.

1904. Mayr, Verh. Zool.-Bot. Gesellsch. Wien, vol. 54, p. 593. One species.

TYPE: *Macrohynniss lepidus* Mayr, 1904. First included species.

*Macrostigma* Rondani

1877. Rondani, Boll. Soc. Ent. Italiana, vol. 9, p. 184.

TYPE: *Macrostigma aphidum* Rondani, 1877, by monotypy.



**Macroteleia** Westwood (= *Macrotelia* Agassiz)

1835. Westwood, Proc. Zool. Soc. London, vol. 3, p. 70.

TYPE: *Macroteleia cleonymoides* Westwood, 1835, by monotypy.**(Macrotelia** Agassiz, emendation) = **Macroteleia** Westwood

1846. Agassiz, Nomenclator zoologicus. Index universalis, p. 221.

**Mallateleia** Dodd

1913. Dodd, Trans. Roy. Soc. South Australia, vol. 37, pp. 131, 151.

TYPE: *Mallateleia giraulti* Dodd, 1913, by monotypy and original designation.**Mallateleicoides** Dodd

1913. Dodd, Trans. Roy. Soc. South Australia, vol. 37, pp. 131, 152.

TYPE: *Mallateleicoides splendida* Dodd, 1913, by monotypy and original designation.**Malvina** Cameron

1889. Cameron, Mem. Proc. Manchester Lit. Phil. Soc., ser. 4, vol. 2, p. 13.

TYPE: *Malvina punctata* Cameron, 1889, by monotypy.**Mandraka** Risbec

1953. Risbec, Mem. Inst. Scient. Madagascar, vol. 3, p. 343.

TYPE: *Mandraka pauliani* Risbec, by monotypy.**Mantara** Dodd

1920. Dodd, Trans. Ent. Soc. London, 1919, p. 379.

TYPE: *Mantara bifurcata* Dodd, 1920, by monotypy and original designation.**Mantibaria** Kirby

1900. Kirby, in Andrews, A monograph of Christmas Island . . . , p. 82.

TYPE: *Mantibaria anomala* Kirby, 1900, by monotypy.**Marshalliella** Kieffer

1913. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 222, pl. 8, fig. 6.

TYPE: *Marshalliella oxygaster* Marshall, 1913, in Kieffer, by monotypy.**Martinica** Risbec

1950. Risbec, Contribution à l'étude des Proctotrupidae. Trav. Lab. Ent. Sect. Soudan. Rech. Agron., p. 533.

TYPE: *Martinica antillensis* Risbec, 1950, by monotypy.**(Megacolus** Priesner, not Cameron, 1903) = **Philoplanes**, new name

1951. Priesner, Bull. Inst. Fouad I du Desert, vol. 1, p. 121.  
 TYPE: *Megacolus desertorum* Priesner, 1951, by monotypy and original designation.

(*Megalospilus* Schulz, emendation) = *Megaspilus* Westwood

1906. Schulz, Spolia hymenopterologica, p. 152.

**Megaplastopria** Ashmead

1903. Ashmead, Journ. New York Ent. Soc., vol. 11, p. 31.

TYPE: *Megaplastopria brasiliensis* Ashmead, 1903, by monotypy and original designation.

**Megaspilidea** Ashmead

1888. Ashmead, Can. Ent., vol. 20, pp. 48, 49.

TYPE: *Megaspilidea minuta* Ashmead, 1888, by monotypy. Considered synonym of *Ceraphron* Jurine.

**Megaspilodes** Ashmead

- 1888 (March). Ashmead, Can. Ent., vol. 20, p. 48. Two names, one of them a nomen nudum.

TYPE: *Ceraphron armatus* Say, 1836, by monotypy. Also designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 671, who at that time thought both originally included names to be valid. Ashmead intended to establish *Megaspilodes* as a monotypic genus with *fuscipennis* Ashmead as type but unfortunately the paper in which he first described the genus was not published until June, 1888 (Kans. State Agr. College Bull. No. 3, App. p. ii).

Considered synonym of *Megaspilus* Westwood.

**Megaspilus** Westwood (= *Megalospilus* Schulz, emendation)

1829. Westwood, in Stephens, The nomenclature of British insects . . . , p. 37. Sixteen names, thirteen of them nomina nuda.

TYPE: *Ceraphron dux* Curtis, 1829. Designated by Westwood, 1839, An introduction to the modern classification of insects, vol. 2, generic synopsis p. 77. *Ceraphron abdominalis* Boheman, 1832, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 112, was not originally included.

**Merriwa** Dodd

1920. Dodd, Trans. Ent. Soc. London, 1919, p. 332.

TYPE: *Merriwa quadridentata* Dodd, 1920, by monotypy and original designation.

**Mesaulacinus** Martynov [fossil]

1925. Martynov, Bull. Acad. des Sci. U. R. S. S., ser. 6, vol. 19, p. 757.

TYPE: *Mesaulacinus oviformis* Martynov, 1925, by monotypy.

**Mesohelorus** Martynov [fossil]

1925. Martynov, Bull. Acad. des Sci. U. R. S. S., ser. 6, vol. 19, p. 758.

TYPE: *Mesohelorus muchini* Martynov, 1925, by monotypy.

**Mesoteleia** Kieffer

1917. Kieffer, Brotéria, Rev. Luso-Brazilera, Zool. ser., vol. 15, p. 51.

TYPE: *Mesoteleia pallida* Kieffer, 1917, by monotypy and original designation.

**Metaclisis** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 106, 109.

TYPE: *Metaclisis areolatus* Foerster, 1856, by monotypy.

**Metanopedias** Brues

1910. Brues, Bull. Amer. Mus. Nat. Hist., vol. 28, p. 79.

TYPE: *Metanopedias sicarius* Brues, 1910, by monotypy.

**Meuselia** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 391. Two species.

TYPE: *Meuselia fuscicornis* Kieffer, 1909, by original designation.

**Micraulacinus** Kieffer

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 350. No species.

1911. Kieffer, Ann. Soc. Ent. France, vol. 80, pp. 214, 219. One species.

TYPE: *Micraulacinus elegans* Kieffer, 1911. First included species.

**Micrevania** Benoit

1952. Benoit, Mém. Inst. Sci. Madagascar, ser. E., vol. 1, p. 273. Two species.

TYPE: *Micrevania minuscula* Benoit, by original designation.

**Microceraphron** Szelényi

1935. Szelényi, Allattani Közlemények, vol. 32, pp. 140, 142.

TYPE: *Microceraphron subterraneus* Szelényi, 1935, by monotypy.

**Microgalesus** Kieffer

1912. Kieffer, in Wytsman, Genera insectorum, fasc. 124, 1911, pp. 6, 43.

TYPE: *Microgalesus quadridens* Kieffer, 1912, by monotypy.  
Again described as new by Kieffer, 1912, Trans. Linn. Soc.  
London, ser. 2, Zool., vol. 15, p. 74.

**Microphanurus** Kieffer (= *Aphanurus* Kieffer, 1912, not Looss, 1907)

1926. Kieffer, Das Tierreich, Lief. 48, pp. xv, 16, 91.

TYPE: *Teleas semistriatus* Nees, 1834, by substitution of  
*Microphanurus* for *Aphanurus* Kieffer.

(**Microps** Haliday, not Dahl, 1823, not Wagler, 1828, not Wagler, 1830)

1833. Haliday, Ent. Mag., vol. 1, p. 272.

TYPE: *Microps rubi* Haliday, 1833, by monotypy.

Although preoccupied, *Microps* Haliday is not renamed since  
it is considered a synonym of *Lagynodes* Foerster.

**Microteleia** Kieffer

1910. Kieffer, in Voeltzkow, Reise in Ostafrika 1903-05,  
vol. 2, Heft 5, p. 531.

TYPE: *Microteleia nitida* Kieffer, 1910, by monotypy.

(**Microtelenomus** Kieffer, error) = **Mirotelenomus** Dodd

1926. Kieffer, Das Tierreich, Lief. 48, pp. 15, 19.

**Mimopria** Holmgren

1908. Holmgren, Zool. Anz., vol. 33, p. 346.

TYPE: *Mimopria ecitophila* Holmgren, 1908, by monotypy.

(**Mionopria** Haliday) = **Idiotypa** Foerster

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. pp. 170, 172.

TYPE: *Diapria (Mionopria) maritima* Haliday, 1857, by mon-  
otypy.

Proposed as a subgenus of *Diapria* Latreille. Isogenotypic  
with *Idiotypa* Foerster.

**Miota** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp.  
131, 141. No species.

1890. Ashmead, Colorado Biol. Assoc. Bull. No. 1, p. 12.  
One species.

TYPE: *Miota glabra* Ashmead, 1890. First included species.  
*Miota compressa* Kieffer, 1910, designated by Kieffer,  
1910, in Wytzman, Genera insectorum, fasc. 107, p.  
40, was included later.

Considered synonym of *Cinetus* Jurine.

**Miotella** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 379.

TYPE: *Miotella tenuicornis* Kieffer, 1909, by monotypy.

**Miramiblyaspis** Dodd

1914. Dodd, Arch. Naturg., vol. 79, A8, p. 91 (published February 1914).

TYPE: *Miramiblyaspis mirabilis* Dodd, 1914, by monotypy. Again described as new by Dodd, December 1914, Ent. News, vol. 25, p. 455, with the single species, *Miramiblyaspis mirabilis* Dodd, designated as type. This was evidently the original description but it happened to be published later.

**Mirobaeoides** Dodd

1914. Dodd, Trans. Roy. Soc. South Australia, vol. 38, pp. 59, 74.

TYPE: *Mirobaeoides tasmanicus* Dodd, 1914, by monotypy and original designation.

**Mirobaeus** Dodd

1914. Dodd, Trans. Roy. Soc. South Australia, vol. 38, pp. 59, 73.

TYPE: *Mirobaeus bicolor* Dodd, 1914, by monotypy and original designation.

**Mirotelenomus** Dodd (= *Microtelenomus* Kieffer, error)

1913. Dodd, Trans. Roy. Soc. South Australia, vol. 37, pp. 158, 173.

TYPE: *Mirotelenomus abnormis* Dodd, 1913, by monotypy.

**Misocyclops** Kieffer

1914. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 353, 362, Pl. xiii, figs. 4, 5.

TYPE: *Platygaster ornatus* Kieffer, 1914, by monotypy.

**(Mithras** Agassiz, emendation of *Mythras* Haliday, nomen nudum) = **Aneurhynchus** Westwood

1846. Agassiz, Nomenclator zoologicus. Index universalis, pp. 235, 244.

*Mythras* Haliday was validated by Brullé, in Lepeletier, 1846, Histoire naturelle des insectes, Hyménoptères, vol. 4, p. 612, through citation as a synonym of *Aneurhynchus* Westwood.

**Monelata** Foerster (= *Monoleta* Maneval, error)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 123, 127. No species.

1873. Marshall, A catalogue of British Hymenoptera; Oxyura, p. 13. Three species.

TYPE: *Diapria parvula* Nees, 1834. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 441.

**Moninostemma** Kieffer

1914. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 357.

TYPE: *Inostemma festus* Walker, 1839, by monotypy and original designation.

**Monocrita** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 106, 109.

TYPE: *Inostemma atinas* Walker, 1835, by monotypy.

**(Monoleta** Maneval, error) = **Monelata** Foerster

1940. Maneval, in Berland, Faune de la France, vol. 7, Hyménoptères, pp. 102, 103, 106.

**Monomachus** Klug

1841. Klug, in Germar, Zeitschr. Ent., vol. 3, p. 378.

TYPE: *Pelecinus fuscator* Perty, 1833, by monotypy. (*Monomachus gladiator* (Klug) Westwood) = *Monomachus gladiator* Brullé, 1846, designated by Ashmead, 1902, Journ. New York. Ent. Soc., vol. 10, p. 243, was included later.

**Monoteleia** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxix, 272, 545. Three species.

TYPE: *Macroteleia grenadensis* Ashmead, 1900, by original designation.

**Monoxylabis** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 345.

TYPE: *Monoxylabis flavimanus* Kieffer, 1909, by monotypy.

**Myrmecopria** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 407, 446.

TYPE: *Loxotropa mellea* Ashmead, 1887, by monotypy and original designation.

**(Mythras** Brullé, validation of *Mythras* Haliday, nomen nudum)

= **Aneurhynchus** Westwood

1846. Brullé, in Lepeletier, Histoire naturelle des insectes. Hyménoptères, vol. 4, p. 612.

TYPE: *Aneurhynchus galesiformis* Westwood, 1832. *Mythras* Haliday MS. was validated and its type fixed by Brullé through citation as synonym of *Aneurhynchus* Westwood.

**(Mythras** Haliday, nomen nudum; *Mithras* Agassiz, emendation)

= **Aneurhynchus** Westwood

1829. Haliday, in Curtis, A guide to an arrangement of British insects, column 108.

Validated by Brullé, 1846, in Lepeletier, *Histoire naturelle des insectes. Hyménoptères*, vol. 4, p. 612, through citation as a synonym of *Aneurhynchus* Westwood.

**Nanopria** Kieffer

1913. Kieffer, in Alluaud and Jeannel, *Voyage . . . en Afrique Orientale. Insectes hyménoptères*, vol. 1, p. 26.

TYPE: *Nanopria fuscipes* Kieffer, 1913, by monotypy and original designation.

**Nardo** Nixon

1938. Nixon, *Ann. Mag. Nat. Hist.*, ser. 11, vol. 1, p. 278. Two species.

TYPE: *Nardo cumaeus* Nixon, 1938, by original designation.

**Nasdia** Nixon

1942. Nixon, *Ann. Mag. Nat. Hist.*, ser. 11, vol. 9, p. 462.

TYPE: *Nasdia prosper* Nixon, 1942, by monotypy and original designation.

**Neanteris** Szelényi, nomen nudum

1937. Szelényi, in Moczar, *A Koszegi Muz. Kozlemenyei, Pubs. Musei Ginsiensis*, ser. 1, no. 5, p. 73.

Cited with *impiger* Szelényi, new species, also a nomen nudum.

**Neaulacus** Bradley

1908. Bradley, *Trans. Amer. Ent. Soc.*, vol. 34, pp. 119, 121, 126. Five species, one of them doubtfully included.

TYPE: *Aulacus occidentalis* Cresson, 1879, by original designation.

Proposed as a subgenus of *Pristaulacus* Kieffer.

**Neivapria** Borgmeier

1939. Borgmeier, *Rev. de Ent.*, vol. 10, p. 543.

TYPE: *Neivapria penicillata* Borgmeier, 1939, by monotypy and original designation.

**Neobetyla** Dodd

1915. Dodd, *Trans. Roy. Soc. South Australia*, vol. 39, pp. 389, 396. Two species.

TYPE: *Neobetyla pulchricornis* Dodd, 1915, by original designation.

**Neoceraphron** Ashmead (= *Neocerataphron* Schulz, emendation)

1893. Ashmead, *U. S. Nat. Mus. Bull.* 45, pp. 123, 136.

TYPE: *Ceraphron macroneurus* Ashmead, 1887, by monotypy.

(*Neocerataphron* Schulz, emendation) = **Neoceraphron** Ashmead

1906. Schulz, *Spolia hymenopterologica*, p. 152.

**Neodiapria** Kieffer

1916. Kieffer, *Das Tierreich*, Lief. 44, pp. xxi, 346.

TYPE: *Tropidopria erynniae* Kieffer, 1910, by monotypy.

**Neolygocerus** Ishii

1951. Ishii, *Oyo-Dobutsugaku-Zasshi* (Japanese Soc. Applied Zool.), vol. 16, p. [93].

TYPE: *Neolygocerus koyamae* Ishii, 1951, by monotypy.

**Neopria** Dodd

1915. Dodd, *Trans. Roy. Soc. South Australia*, vol. 39, pp. 398, 429. Three species.

TYPE: *Neopria trifoveata* Dodd, 1915, by original designation.

**Neoscelio** Dodd

1913. Dodd, *Trans. Roy. Soc. South Australia*, vol. 37, pp. 131, 138.

TYPE: *Neoscelio gloriosus* Dodd, 1913, by monotypy and original designation.

**Neoteleia** Dodd

1913. Dodd, *Arch. Naturg.*, vol. 79, A6, p. 169.

TYPE: *Neoteleia punctata* Dodd, 1913, by monotypy and original designation.

**Neotelenomus** Dodd

1913. Dodd, *Trans. Roy. Soc. South Australia*, vol. 37, pp. 158, 171. Five species.

TYPE: *Neotelenomus anthereae* Dodd, 1913, by original designation.

**Nesopria**, new name (= *Tropidopsis* Ashmead, 1893, not Brauer and Bergenstamm, 1890)

TYPE: *Tropidopsis clavata* Ashmead, 1893, by substitution of *Nesopria* for *Tropidopsis* Ashmead.

**Neuraulacinus** Kieffer

1910. Kieffer, *Bull. Soc. Ent. France*, 1910, p. 350. No species.

1911. Kieffer, *Ann. Soc. Ent. France*, vol. 80, pp. 214, 216. Three species.

TYPE: *Neuraulacinus braconiformis* Kieffer, 1911. Designated by Kieffer, 1911, *Ann. Soc. Ent. France*, vol. 80, p. 217. *Neuraulacinus vespiformis* Kieffer, 1911, was designated by Kieffer, 1912, *Das Tierreich*, Lief. 30, pl. 358.

**Neurocacus** Kieffer

1913. Kieffer, *Insecta*, vol. 3, p. 428.

TYPE: *Neurocacus philippinensis* Kieffer, 1913, by monotypy and original designation.



**Neurogalesus** Kieffer

1907. Kieffer, Berliner Ent. Zeitschr., vol. 51, 1906, p. 298.

TYPE: *Neurogalesus carinatus* Kieffer, 1907, by monotypy.

**Neuropria** Kieffer

1904. Kieffer, Bull. Soc. Hist. Nat. Metz, vol. 23, p. 53.

TYPE: *Neuropria sociabilis* Kieffer, 1904, by monotypy.

**Neuroscelio** Dodd

1913. Dodd, Arch. Naturg., vol. 79, A6, p. 170.

TYPE: *Neuroscelio nervalis* Dodd, 1913, by monotypy and original designation.

**Neuroteleia** Kieffer

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 293. Two species.

TYPE: *Neuroteleia rufa* Kieffer, 1910, by original designation.

Again described as new by Kieffer, 1912, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 61.

**Nirupama** Nixon

1935. Nixon, Trans. Roy. Ent. Soc. London, vol. 83, p. 93. Two species.

TYPE: *Nirupama morpheus* Nixon, 1935, by original designation.

**Nothoserphus** Brues

1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 73, p. 263.

TYPE: *Nothoserphus mirabilis* Brues, 1940, by monotypy and original designation.

**Notilena** Brèthes

1913. Brèthes, An. Mus. Nac. Hist. Nat. Buenos Aires, vol. 24, p. 84.

TYPE: *Notilena gallardi* Brèthes, 1913, by monotypy and original designation.

**Notoxoides** Ashmead

1903. Ashmead, Journ. New York Ent. Soc., vol. 11, p. 30.

TYPE: *Notoxoides brasiliensis* Ashmead, 1903, by monotypy and original designation.

**Notoxopria** Kieffer

1910. Kieffer, Ent. Rundschau, vol. 27, p. 39.

TYPE: *Notoxopria rufa* Kieffer, 1910, by monotypy.

**Nyleta** Dodd

1926. Dodd, Trans. Roy. Soc. South Australia, vol. 1, p. 300.

TYPE: *Nyleta striaticeps* Dodd, 1926, by monotypy and original designation.

**Odontacolus** Kieffer

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 294.

TYPE: *Odontacolus longiceps* Kieffer, 1910, by monotypy and original designation.

Again described as new by Kieffer, 1912, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 54.

**Odontaulacus** Kieffer

1903. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 7 bis, p. 382. Two species.

TYPE: *Aulacus rufitarsis* Cresson, 1864. Designated by Bradley, 1908, Trans. Amer. Ent. Soc., vol. 34, p. 120. *Aulacus editus* Cresson, 1880, was designated by Kieffer, 1912, Das Tierreich, Lief. 30, p. 365.

**Odontofoenus** Kieffer

1910. Kieffer, Ann. Soc. Ent. France, vol. 79, p. 77.

TYPE: *Gasteruption humerale* Schletterer, 1889, by monotypy and original designation.

**Odontopria** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2(42), pp. 24, 34, 39. Two species.

TYPE: *Odontopria temporalis* Kieffer, 1905. Designated by Kieffer, 1912, in Wytsman, Genera insectorum, fasc. 124, 1911, p. 29.

**Odontopsilus** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 341.

TYPE: *Odontopsilus tenuicornis* Kieffer, 1909, by monotypy.

**Odontoscelio** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2(42), p. 13.

TYPE: *Odontoscelio striatifrons* Kieffer, 1905, by monotypy.

**Oethecoctonus** Ashmead (= *Cacus* Riley, 1893, not Gistel, 1848, not Selys, 1854, not Costa, 1857; *Cacellus* Ashmead)

1900. Ashmead, Can. Ent., vol. 32, p. 368.

TYPE: *Cacus oecanthi* Riley, 1893, by substitution of *Oethecoctonus* for *Cacus* Riley.

**Oleisoprister** Bradley

1901. Bradley, Trans. Amer. Ent. Soc., vol. 27, pp. 320, 324. Five species.

TYPE: *Aulacus firmus* Cresson, 1879, by original designation.

**Opazon** Haliday

1857. Haliday, Nat. Hist. Rev., vol. 4, Proc. p. 170.

TYPE: *Belyta* (*Opazon*) *parvula* Haliday, 1857, by monotypy. *Belyta* (*Opazon*?) *ciliata* Thomson, 1858, was desig-

nated by Kieffer, 1916, *Das Tierreich*, Lief. 44, p. 378, since he considered *B. (O.) parvula* Haliday a nomen nudum.

Proposed as a subgenus of *Belyta* Jurine.

**Opisthacantha** Ashmead (= *Raia* Ashmead)

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 209, 211, 221.

TYPE: *Opisthacantha mellipes* Ashmead, 1893, by monotypy and original designation.

**Oreiscelio** Kieffer (= *Oreiscelio* Kieffer, emendation)

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 293.

TYPE: *Oreiscelio sechellensis* Kieffer, 1910, by monotypy and original designation.

(*Oreiscelio* Kieffer, emendation) = **Oreiscelio** Kieffer

1912. Kieffer, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 58. Emendation of *Oreiscelio* Kieffer, and described as new. The type species was also emended to read *Oreiscelio seychellensis* Kieffer, 1910.

(*Orthopria* Kieffer) = **Planopria** Kieffer

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 983, 984. Twenty-two species, 16 of them included by bibliographic reference.

TYPE: *Diapria californica* Ashmead, 1893. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 678.

Proposed as a subgenus of *Trichopria* Ashmead. Isogenotypic with *Planopria* Kieffer, 1908.

**Oxylabis** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 128, 130, 133, 139. Two species.

TYPE: *Cinetus picipes* Nees, 1834. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 681. *Belyta bisulca* Nees, 1834, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 357, was not originally included.

**Oxyphanurus** Kieffer

1926. Kieffer, *Das Tierreich*, Lief. 48, pp. [xiii], 19.

TYPE: *Telenomus charmus* Walker, 1839, by monotypy.

**Oxypria** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, p. 37.

TYPE: *Oxypria thoracica* Kieffer, 1908, by monotypy.

**Oxyscelio** Kieffer

1907. Kieffer, Zeitschr. Hym. Dipt., vol. 7, p. 310.

TYPE: *Oxyscelio foveatus* Kieffer, 1907, by monotypy.

**Oxyteleia** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, p. 118. Four species.

TYPE: *Caloteleia bidentata* Kieffer, 1905. Designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 516.

**(Oxyurus** Lamarck, not Rafinesque, 1810, not Lamarck, 1816) = **Bebelus** Gistel

1817. Lamarck, Histoire naturelle des animaux sans vertèbres, vol. 4, p. 128. Seven species.

TYPE: *Sparasion frontalis* Latreille, 1805. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 701.

Considered synonym of *Sparasion* Latreille.

**Pachyscelidris** Szelényi

1941. Szelényi, Zool. Anz., vol. 134, pp. 158, 163.

TYPE: *Pachyscelidris aptera* Szelényi, 1941, by monotypy and original designation.

**Pachyscelio** Risbec (= *Brachyscelio* Risbec, 1950, not Brues, 1940)

1954. Risbec, Bull. Inst. Français d'Afrique Noire, vol. 16, p. 552.

TYPE: *Brachyscelio jeanneli* Risbec, 1950, by substitution of *Pachyscelio* for *Brachyscelio* Risbec.

**Palaeoteleia** Cockerell [fossil]

1915. Cockerell, Proc. Acad. Nat. Sci. Philadelphia, vol. 66, 1914, p. 637.

TYPE: *Palaeoteleia oxyura* Cockerell, 1915, by monotypy.

**Palpoteleia** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxix, 272, 547. Five species.

TYPE: *Psiloteleia atra* Kieffer, 1926, by original designation.

**Pammegischia** Provancher

1882. Provancher, Nat. Canadien, vol. 13, p. 302.

TYPE: *Pammegischia burquei* Provancher, 1882, by monotypy.

Considered synonym of *Aulacus* Jurine.

**Pantoclis** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 129, 137. No species.

1861. Foerster, Programm der Realschule zu Aachen, p. xliii. One species.

TYPE: *Pantoclis barycera* Foerster, 1861. First included species. *Belyta brevis* Nees, 1834, designated by Ash-

mead, 1893, U. S. Nat. Mus. Bull. 45, p. 366, was included later.

Considered synonym of *Aclista* Foerster.

(*Pantolysa* Foerster, error) = *Pantolyta* Foerster

1861. Foerster, Programm der Realschule zu Aachen, p. xliii.

*Pantolyta* Foerster (= *Pantolysa* Foerster, error)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 128, 130, 135, 136. No species.

1861. Foerster, Programm der Realschule zu Aachen, p. xliii. One species.

TYPE: *Pantolyta atrata* Foerster, 1861. First included species. *Belyta heterocera* Haliday, 1857, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 383, and *Pantolyta pallida* Kieffer, 1908, by Kieffer, 1908, in André, Spec. Hym. Eur. Alg., vol. 10, p. 428, were included later.

*Pantolytoidea* Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 389, 390. Four species.

TYPE: *Pantolytoidea splendida* Dodd, 1915, by original designation.

*Pantolytomyia* Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 389, 395.

TYPE: *Pantolytomyia ferruginea* Dodd, 1915, by monotypy and original designation.

*Pantopiasta* Maneval

1939. Maneval, Bull. Soc. Ent. France, vol. 44, pp. 169, 170.

TYPE: *Cinetus flaviventris* Thomson, 1858, by monotypy and original designation.

*Parabaeus* Kieffer

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 294.

TYPE: *Parabaeus ruficornis* Kieffer, 1910, by monotypy and original designation.

Also described as new by Kieffer, 1912, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 53.

*Parabetyla* Brues

1922. Brues, Psyche, vol. 29, pp. 222, 224.

TYPE: *Parabetyla spinosa* Brues, 1922, by monotypy and original designation.

*Paraclista* Kieffer

1909 (September 15). Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 476. Four species.

- TYPE: *Belyta brachyptera* Thomson, 1858. Designated by Kieffer, 1910, in Wytzman, *Genera insectorum*, fasc. 107, p. 23.
- Kieffer described five new species in this genus in *Ann. Soc. Sci. Bruxelles*, vol. 33, p. 390, 1909, received by the Smithsonian Institution, Washington, D. C., Oct. 4, 1909, but precise date of publication is unknown.
- Proposed as a subgenus of *Belyta* Jurine.
- Paracodrus** Kieffer
1907. Kieffer, in André, *Spec. Hym. Eur. Alg.*, vol. 10, pp. 272, 273. Three species.
- TYPE: *Paracodrus bethyliformis* Kieffer, 1907, by original designation.
- (*Paracyclops* Maneval, not Claus, 1893) = *Urocyclops* Maneval
1936. Maneval, *Bull. Ann. Soc. Ent. Belgique*, vol. 76, p. 56.
- TYPE: *Paracyclops bettyae* Maneval, 1936, by monotypy and original designation.
- Parafoenus** Kieffer
1910. Kieffer, *Bull. Soc. Ent. France*, 1910, p. 350. No species.
1911. Kieffer, *Ann. Soc. Ent. France*, vol. 80, pp. 213, 215. One species.
- TYPE: *Parafoenus formosus* Kieffer, 1911. First included species.
- Paragryon** Kieffer
1908. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 32, pp. 189, 199. Six species, five by bibliographic reference.
- TYPE: *Paragryon pedestris* Kieffer, 1908. Designated by Kieffer, 1926, *Das Tierreich*, Lief. 48, p. 234.
- Paramesius** Westwood
1832. Westwood, *London Edinburgh Phil. Mag.*, ser. 3, vol. 1, p. 129.
- TYPE: *Paramesius rufipes* Westwood, 1832, by monotypy.
- Paranteris** Kieffer
1910. Kieffer, *Bull. Soc. Ent. France*, 1910, p. 292. Six species.
- TYPE: *Paranteris nigriclava* Kieffer, 1910, by original designation.
- Again described as new by Kieffer, 1912, *Trans. Linn. Soc. London*, ser. 2, *Zool.*, vol. 15, p. 65.
- Parapegus** Kieffer
1908. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 32, pp. 148 (footnote), 149. Two species.

TYPE: *Apegus (Parapegus) punctatus* Kieffer, 1908. Designated by Kieffer, 1910, in Wytzman, *Genera insectorum*, fasc. 80B, p. 86.

Proposed as a subgenus of *Apegus* Foerster.

#### **Parascelio** Dodd

1920. Dodd, *Trans. Ent. Soc. London*, 1919, p. 341.

TYPE: *Parascelio undulatus* Dodd, 1920, by monotypy and original designation.

#### **Paraspilomicrus** Johnston and Tiegs

1921. Johnston and Tiegs, *Proc. Roy. Soc. Queensland*, vol. 33, pp. 103, 104, 123.

TYPE: *Paraspilomicrus froggatti* Johnston and Tiegs, 1921, by monotypy and original designation.

#### **Paratelenomus** Dodd

1914. Dodd, *Proc. Roy. Soc. Queensland*, vol. 26, p. 121.

TYPE: *Telenomus bicolor* Dodd, 1914, by monotypy and original designation.

#### **Parateloipsis** Whittaker

1930. *Proc. Ent. Soc. Washington*, vol. 32, p. 73.

TYPE: *Parateloipsis canadensis* Whittaker, 1930, by monotypy and original designation.

#### **Paratrimorus** Kieffer

1908. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 32, pp. 117, 146.

TYPE: *Paratrimorus perplexus* Kieffer, 1908, by monotypy.

#### **Parepimeces** Kieffer (= *Epimeces* Westwood, 1833, not Billberg, 1820)

1926. Kieffer, *Das Tierreich*, Lief. 48, pp. xxxiv, 563, 760.

TYPE: *Epimeces ensifer* Westwood, 1833, by substitution of *Parepimeces* for *Epimeces* Westwood.

#### **Parevania** Kieffer

1907. Kieffer, *Berliner Ent. Zeitschr.*, vol. 51, 1906, p. 270.

TYPE: *Parevania semirufa* Kieffer, 1907, by monotypy.

#### **(Paridris** Brèthes, not Kieffer, 1908)

1917. Brèthes, *An. Zool. Aplicada*, vol. 4, p. 27.

TYPE: *Paridris chilensis* Brèthes, 1917, by monotypy.

No replacement name has been published and none appears needed since the genus is considered a synonym of *Telenomus* Haliday.

#### **Paridris** Kieffer

1908. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 32, p. 122. Three species.

TYPE: *Idris laeviceps* Ashmead, 1893. Designated by Kieffer, 1926, *Das Tierreich*, Lief. 48, p. 421.

**Parinostemma** Kieffer

1914. Kieffer, *in* André, Spec. Hym. Eur. Alg., vol. 11, p. 355.

TYPE: *Inostemma quinda* Walker, 1842, by monotypy and original designation.

**Paroxylabis** Kieffer

1907. Kieffer, Brotéria, Rev. Sci. Nat., Zool. ser., vol. 6, p. 16.

TYPE: *Paroxylabis semirufa* Kieffer, 1907, by monotypy. Again described as new by Kieffer, 1908, *in* André, Spec. Hym. Eur. Alg., vol. 10, pp. 359, 367.

**Passalida** Brèthes

1914. Brèthes, Nunquam otiosus, Buenos Aires, p. 2.

TYPE: *Passalida spinifera* Brèthes, 1914, by monotypy and original designation.

Considered synonym of *Amitus* Haldeman.

**Pegoteleia** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxviii, 272, 510. Fourteen species.

TYPE: *Baryconus calopterus* Kieffer, 1910, by original designation.

**Pelecinopteron** Brues [fossil]

1933. Brues, Bernsteinforschungen, Heft 3, 1932, pp. 17, 19.

TYPE: *Pelecinopteron tubuliforme* Brues, 1933, by monotypy and original designation.

**Pelecinus** Latreille (= *Episceuastes* Gistel)

1800. Latreille, *in* Duméril, Bull. Sci. Soc. Philom. Paris, vol. 2, No. 44, p. 155. Two species, one of them included by bibliographic reference.

TYPE: *Ichneumon polycerator* Fabricius, 1776. Designated by Latreille, 1810, Considérations générales . . . des insectes, p. 436. Latreille considered *Ichneumon polyturator* Drury, 1773, and *I. polycerator* Fabricius as the same species but he used the Fabrician name.

**(*Pentacantha* Ashmead, not Stål, 1871) = Propentacantha** Kieffer

1888. Ashmead, Can. Ent., vol. 20, p. 51.

TYPE: *Pentacantha canadensis* Ashmead, 1888, by monotypy. Considered synonym of *Trisacantha* Ashmead.

**Pentapria** Kieffer

1905 (October). Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2 (42), p. 34. No species.



1905 (November). Kieffer, Bull. Soc. Hist. Nat. Metz, ser. 2, vol. 12, p. 105. One species.

TYPE: *Pentapria conjugens* Kieffer, 1905. First included species. *Pentapria punctaticeps* Kieffer, 1906, designated by Kieffer, 1912, in Wytsman, Genera insectorum, fasc. 124, 1911, p. 34, was included later.

**Petalosema** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxiv, 267, 358. Five species.

TYPE: *Chromoteleia rufithorax* Kieffer, 1907, by present designation.

**Pezopria** Kieffer

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 697, 712. No species.

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 885. One species.

TYPE: *Pezopria fuscicornis* Kieffer, 1911. First included species.

**Phaedroteleia** Kieffer

1916. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 14, p. 182. Two species.

TYPE: *Phaedroteleia armata* Kieffer, 1916. Designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 418.

**Phaenopria** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 407, 436. Eight species.

TYPE: *Phaenopria minutissima* Ashmead, 1893, by original designation.

**Phaenoserphus** Kieffer

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 289, 298. Thirteen species and three varieties, as well as two species included doubtfully.

TYPE: *Proctotrupes curtipennis* Haliday, 1839. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 664.

Proposed as a subgenus of *Serphus* Schrank.

**Phanuomorpha** Dodd

1926. Dodd, in Kieffer, Das Tierreich, Lief. 48, p. 556. Nomen nudum.

Still a nomen nudum.

**Phanuromyia** Dodd (= *Pharunomyia* Dodd, error)

1914. Dodd, Proc. Roy. Soc. Queensland, vol. 26, p. 121.

TYPE: *Pharunomyia* (!) *rufobasalis* Dodd, 1914, by monotypy and original designation.**Phanuropsis** Girault

1916. Girault, Entomologist, vol. 49, p. 198.

TYPE: *Phanuropsis semiflaviventris* Girault, 1916, by monotypy and original designation.**Phanurus** Thomson

1860. Thomson, Öfv. Vet.-Akad. Förh., vol. 17, pp. 169, 172. Ten species.

TYPE: *Phanurus angustatus* Thomson, 1860. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 139.Considered synonym of *Telenomus* Haliday.**(Pharunomyia** Dodd, error) = **Phanuromyia** Dodd

1914. Dodd, Proc. Roy. Soc. Queensland, vol. 26, p. 121.

**Philolestes** Kieffer

1922. Kieffer, An. Soc. Cient. Argentina, vol. 94, p. 205.

TYPE: *Philolestes rufus* Kieffer, 1922, by monotypy and original designation.**Philolestoides** Ferrière

1929. Ferrière, Zool. Anz., vol. 82, p. 160.

TYPE: *Philolestoides wasmanni* Ferrière, 1929, by monotypy.**Philoplanes**, new name (= *Megacolus* Priesner, 1951, not Cameron, 1903)TYPE: *Megacolus desertorum* Priesner, 1951, by substitution of *Philoplanes* for *Megacolus* Priesner.**Phoenoteleia** Kieffer

1916. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 14, p. 62.

TYPE: *Phoenoteleia rufa* Kieffer, 1916, by monotypy and original designation.**(Phoenus** Schletterer, emendation of *Foenus* Fabricius) = **Gasteruption** Latreille

1890. Schletterer, Ann. Naturh. Hofmus. Wien, vol. 4, p. 375.

**Piestopleura** Foerster (= *Catillus* Foerster 1856, not Brongniart, 1822)

1856. Foerster, Hymenopterologische Studien, Heft 2, p. 144.

TYPE: *Platygaster catillus* Walker, 1835, by substitution of *Piestopleura* Foerster for *Catillus* Foerster.**Plagioscelio** Kieffer

1916. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 14, p. 185. Two species.

TYPE: *Plagioscelio rufescens* Kieffer, 1916, by original designation.

**Planopria** Kieffer (= *Orthopria* Kieffer)

1908. Kieffer, Bull. Soc. Hist. Nat. Metz, vol. 25, p. 19. Sixteen species included by bibliographic reference.

TYPE: *Diapria californica* Ashmead, 1893. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 678.

Isogenotypic with *Orthopria* Kieffer. Considered synonym of *Trichopria* Ashmead.

(**Planopria** Kieffer, 1910, not Kieffer, 1908) = **Planopriella** Kieffer

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 695. No species.

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 895. One species.

TYPE: *Planopria pedestris* Kieffer, 1911. First included species.

**Planopriella** Kieffer (= *Planopria* Kieffer, 1910, not Kieffer, 1908)

1912. Kieffer, in Wytsman, Genera insectorum, fasc. 124, 1911, pp. 4, 48.

TYPE: *Planopria pedestris* Kieffer, 1911, by substitution of *Planopriella* for *Planopria* Kieffer, 1910.

**Plastobelyta** Kieffer

1906. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 30, p. 116.

TYPE: *Plastobelyta gallicola* Kieffer, 1906, by monotypy. Doubtfully referred to the Proctotrupoidea.

**Plastogryon** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, pp. 119, 141. Four species and one variety.

TYPE: *Plastogryon foersteri* Kieffer, 1908. Designated by Brues, 1908, in Wytsman, Genera insectorum, fasc. 80, p. 51.

**Plastomicrops** Kieffer

1906. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 30, p. 145. One species and one variety.

TYPE: *Plastomicrops acuticornis* Kieffer, 1906, by present designation.

**Platycephron** Kieffer

1906. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 30, p. 141. Two species.

TYPE: *Platycephron muscidarum* Kieffer, 1906, by present designation.

**Platygaster** Latreille

1809. Latreille, Genera crustaceorum et insectorum . . . , vol. 4, p. 31.

TYPE: *Scelio ruficornis* Latreille, 1805, by monotypy. "*P. rufipes* Latr.," cited as type by Ashmead, 1903, Journ. New York Ent. Soc., vol. 11, p. 98, was undoubtedly a lapsus for *ruficornis*.

**Platygastoides** Dodd

1913. Dodd, Can. Ent., vol. 45, p. 346.

TYPE: *Platygastoides mirabilis* Dodd, 1913, by monotypy and original designation.

**Platyllostropa** Szelényi

1938. Szelényi, Ann. Mus. Nat. Hungarici, vol. 31, Zool., p. 126.

TYPE: *Platyllostropa gallicola* Szelényi, 1938, by monotypy and original designation.

**Platymischoides** Ashmead

1901. Ashmead, Fauna Hawaiiensis, vol. 1, p. 296.

TYPE: *Platymischoides molokaiensis* Ashmead, 1901, by monotypy.

**Platymischus** Westwood

1832. Westwood, London Edinburgh Phil. Mag., ser. 3, vol. 1, p. 128.

TYPE: *Platymischus dilatatus* Westwood, 1832, by monotypy.

**Platyscelio** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2 (42), p. 11.

TYPE: *Platyscelio pulchricornis* Kieffer, 1905, by monotypy.

**Platystasius** Nixon

1937. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 19, p. 372. Two species.

TYPE: *Platystasius strangaliophagus* Nixon, 1937, by original designation.

**Platyteleia** Dodd

1913. Trans. Roy. Soc. South Australia, vol. 37, pp. 131, 153.

TYPE: *Platyteleia latipennis* Dodd, 1913, by monotypy and original designation.

**Platytelenomus** Dodd

1914. Dodd, Ent. News, vol. 25, p. 126.

TYPE: *Platytelenomus planus* Dodd, 1914, by monotypy and original designation.

**Plesiobaecus** Kieffer

1913. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 229, 282.

TYPE: *Plesiobaecus hospes* Kieffer, 1913, by monotypy.

**Pleuropria** Kieffer

1905. Ann. Soc. Sci. Bruxelles, vol. 29, pp. 132, 136.

TYPE: *Pleuropria maculipennis* Kieffer, 1905, by monotypy.

**Plutofoenus** Kieffer

1911. Kieffer, Ann. Soc. Ent. France, vol. 80, p. 177.

TYPE: *Gasteruption paraguayense* Schrottky, 1906, by monotypy and original designation.

**Plutopria** Kieffer

1910. Kieffer, Ent. Rundschau, vol. 27, p. 48.

TYPE: *Plutopria luctuosa* Kieffer, 1910, by monotypy.

**Polydiapria** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 398, 427.

TYPE: *Polydiapria atriceps* Dodd, 1915, by monotypy and original designation.

**Polygnotus** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 103, 115.

TYPE: *Platygaster striolatus* Nees, 1834, by monotypy and original designation.

Considered synonym of *Platygaster* Latreille.

**(*Polymecus* Foerster) = *Ectadius* Foerster**

1856. Foerster, Hymenopterologische Studien, Heft 2, p. 144.

TYPE: *Platygaster craterus* Walker, 1835, by substitution of *Polymecus* Foerster for *Ectadius* Foerster.

Unnecessarily proposed for *Ectadius* Foerster which Foerster considered preoccupied by *Ectadium* E. Meyer in botany.

**Polypeza** Foerster (= *Propantolyta* Kieffer)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 123, 127. No species.

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 385, 387. One species.

TYPE: *Polypeza pergandei* Ashmead, 1893. First included species. *Polypeza foersteri* Kieffer, 1910, designated by Kieffer, 1910, in André, Spec. Hym. Eur. Alg., vol. 10, p. 718, was included later.

Isogenotypic with *Propantolyta* Kieffer.

**Polypria** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 399, 410.

TYPE: *Polypria nigriventris* Dodd, 1915, by monotypy and original designation.

**Pristaulacus** Kieffer

1900. Kieffer, Ann. Soc. Ent. France, vol. 68, 1899, p. 813.  
Four species.

TYPE: *Pristaulacus chlapowskii* Kieffer, 1900. Designated by Kieffer, 1903, in André, Spec. Hym. Eur. Alg., vol. 7 bis, p. 455.

**Pristomicrops** Kieffer

1906. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 30, p. 146.

TYPE: *Pristomicrops clavatus* Kieffer, 1906, by monotypy.

**Probaryconus** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, pp. 118, 165, 168.

TYPE: *Baryconus (Probaryconus) spinosus* Kieffer, 1908, by monotypy.

Proposed as a subgenus of *Baryconus* Foerster.

**Probelyta** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 364.

TYPE: *Probelyta alticola* Kieffer, 1909, by monotypy.

**Probetyla** Brues

1922. Brues, Psyche, vol. 29, p. 222.

TYPE: *Probetyla subaptera* Brues, 1922, by monotypy and original designation.

**Procacus** Kieffer

1910. Kieffer, Ann. Soc. Ent. France, vol. 78, 1909, p. 319.

TYPE: *Procacus striatigena* Kieffer, 1910, by monotypy.

(*Procinetus* Kieffer, not Foerster, 1868) = *Eccinetus*, new name

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 638 (footnote).

TYPE: *Cinetus radiatus* Kieffer, 1905, by monotypy and original designation.

(*Proctotripes* Provancher, error) = *Proctotrupes* Latreille.

1883. Provancher, Petite faune entomologique du Canada . . . , p. 558.

(*Proctotropis* Gistel, emendation) = *Proctotrupes* Latreille.

1848. Gistel, Naturgeschichte des Thierreichs, p. 143.

**Proctotrupes** Latreille (= *Serphus* Schrank, suppressed; *Proctotrupes* Rafinesque, emendation; *Proctotrypes* Agassiz, emendation; *Proctotropis* Gistel, emendation; *Proctotripes*, error)

1796. Latreille, Précis caractères génériques des insectes . . . , p. 103. No species.

1802. Latreille, Histoire naturelle . . . crustacés et des insectes, vol. 3, p. 308. One species.

TYPE: *Proctotrupes brevipennis* Latreille, 1802. First included species. Designated by the International Commission on Zoological Nomenclature under suspension of the Rules, Opinion 178, 1946, p. [547]. *Codrus pallipes* Jurine, 1807, designated by Latreille, 1810, *Considérations générales . . . des insectes*, p. 437, *Codrus niger* Panzer, 1801, designated by Westwood, 1839, *An introduction to the modern classification of insects*, vol. 2, generic synopsis p. 76, *Ichneumon gravidator* Linnaeus, 1758, designated by Curtis, 1839, *British entomology*, vol. 16, folio 744, and *Proctotrupes areolator* Haliday, 1839, designated by Blanchard, 1849, *in* Cuvier, *Règne animal* [ed. 3], *Insectes*, text vol. 2, p. 156; atlas, pl. 115, fig. 4, were included later.

(*Proctotrypes* Agassiz, emendation) = *Proctotrupes* Latreille.

1846. Agassiz, *Nomenclator zoologicus. Index universalis*, p. 309.

(*Proctrupes* Rafinesque, emendation) = *Proctotrupes* Latreille

1815. Rafinesque, *Analyse*, p. 125.

*Proctotrypites* Meunier [fossil]

1919. Meunier, *Jahrb. Preuss. Geol. Landesanst. Bergakad.*, Berlin, vol. 39, pt. 1, Heft 1, p. 145.

TYPE: *Proctotrypites rottensis* Meunier, 1919, by monotypy.

*Prodendrocerus* Kieffer

1907. Kieffer, *in* André, *Spec. Hym. Eur. Alg.*, vol. 10, p. 11. No species.

1909. Kieffer, *in* Wytsman, *Genera insectorum*, fasc. 94, p. 6. One species.

TYPE: *Dendrocerus ratzeburgi* Ashmead, 1904. First included species.

*Prolapitha* Kieffer

1908. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 32, p. 117.

TYPE: *Prolapitha nigriceps* Kieffer, 1908, by monotypy and original designation.

*Proleptacis* Kieffer

1926. Kieffer, *Das Tierreich*, Lief. 48, pp. xxi, 562, 632. Four species.

TYPE: *Leptacis foersteri* Kieffer, 1914, by original designation.

*Promeuselia* Kieffer

1910. Kieffer, *in* Wytsman, *Genera insectorum*, fasc. 107, pp. 4, 15, 16.

- TYPE: *Rhynchopsilus clausus* Kieffer, 1908, by monotypy.  
Proposed as a subgenus of *Rhynchopsilus* Kieffer.
- (Propantolyta Kieffer) = Polypeza Foerster**  
1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 697, 709.
- TYPE: *Polypeza pergandei* Ashmead, 1893, by monotypy and original designation.  
Isogenotypic with *Polypeza* Foerster.
- Propentacantha Kieffer (= Pentacantha Ashmead, 1888, not Stål, 1871)**  
1926. Kieffer, Das Tierreich, Lief. 48, pp. xxi, 173, 241.  
TYPE: *Pentacantha canadensis* Ashmead, 1888, by substitution of *Propentacantha* for *Pentacantha* Ashmead.  
Considered synonym of *Trisacantha* Ashmead.
- Propentapria Dodd**  
1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 399, 425.  
TYPE: *Propentapria multifoveata* Dodd, 1915, by monotypy and original designation.
- Prophanurus Kieffer**  
1912. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 9, 37. Twenty species, one of them doubtfully included.  
TYPE: *Teleas phalaenarum* Nees, 1834, by original designation.  
Considered synonym of *Telenomus* Haliday.
- Proplatygaster Kieffer**  
1904. Kieffer, Rev. Chilena, vol. 8, p. 144.  
TYPE: *Proplatygaster rufipes* Kieffer, 1904, by monotypy.
- Proplatyscelio Brues [fossil]**  
1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 74, p. 85.  
TYPE: *Proplatyscelio depressus* Brues, 1940, by monotypy and original designation.
- Propsilomma Kieffer**  
1916. Kieffer, Das Tierreich, Lief. 44, pp. xxiv, 351, 422.  
TYPE: *Psilomma columbianum* Ashmead, 1893, by monotypy.
- Prosacantha Nees**  
1834. Nees, Hymenopterorum ichneumonibus affinium, monographiae, vol. 2, p. 294. Three species.  
TYPE: *Prosacantha longicornis* Nees, 1834. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 185.



**Prosactogaster** Kieffer

1914. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 352, 362, pl. 13, figs. 6 and 7, pl. 17, figs. 5-7. Two species.

TYPE: *Platygaster lineatus* Kieffer, 1906, by present designation. *Prosactogaster floricola* Kieffer, 1916, designated by Kieffer, 1916, Zentralbl. Bakt. Parasitenk. Infektionsk., Abt. 2, vol. 46, p. 570, and *Prosactogaster umbraculi* Kieffer, 1916, designated by him, 1926, Das Tierreich, Lief. 48, p. 762, were included later.

**Prosamblyaspis** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xxx, 561, 610.

TYPE: *Amblyaspis flavosignatus* Kieffer, 1912, by monotypy.

**Prosanteris** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, pp. 121, 136. Nine species.

TYPE: *Anteris nigriceps* Ashmead, 1893. Designated by Kieffer, 1910, in Wytsman, Genera insectorum, fasc. 80B, p. 87.

**Prosapegus** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, pp. 121, 147.

TYPE: *Anteris elongata* Ashmead, 1887, by monotypy and original designation.

**Prosevania** Kieffer

1911. Kieffer, Ann. Soc. Ent. France, vol. 80, p. 157. Three species.

TYPE: *Evania (Prosevania) afra* Kieffer, 1911. Designated by Viereck, 1914, U. S. Nat. Mus. Bull. 83, p. 123. *Evania (Prosevania) subtangens* Kieffer, 1911, was designated by Hedicke, 1939, Hymenopterorum catalogus, pars 9, p. 28, who probably considered Viereck's "monobasic" designation invalid.

Proposed as a subgenus of *Evania* Fabricius.

**Prosinostemma** Kieffer

1914. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 356, 378.

TYPE: *Prosinostemma nigerrima* Kieffer, 1914, by monotypy.

**Prosoxylabis** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 344.

TYPE: *Prosoxylabis maculipennis* Kieffer, 1909, by monotypy.

**Prosparasion** Kieffer

1913. Kieffer, Boll. Lab. Zool. Portici, vol. 7, p. 190.

TYPE: *Prosparasion coeruleum* Kieffer, 1913, by monotypy and original designation.

**Prospilomicrus** Kieffer

1910. Kieffer, in Voeltzkow, Reise in Ostafrika, 1903-05, vol. 2, Heft 5, p. 532. Two species.

TYPE: *Prospilomicrus fuscicornis* Kieffer, 1910, by original designation.

Also published by Kieffer Mar. 1, 1910, without included species, in André, Hym. Eur. Alg., vol. 10, pp. 702, 710.

**Prosynacra** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2, pp. 35, 37. No species.

1910 (January 28). Kieffer, Boll. Lab. Zool. Portici, vol. 4, p. 108. One species and one nomen nudum.

TYPE: *Prosynacra nigriceps* Kieffer, 1910. First included species. Kieffer actually based this genus on *Prosynacra giraudi* Kieffer, which he designated as type in 1912, in Wytsman, Genera insectorum, fasc. 124, p. 13, 1911; but unfortunately the description of this species was not published until Mar. 1, 1910.

**Prosynopeas** Kieffer

1916. Kieffer, Zentralbl. Bakt. Parasitenk. Infektionskr., Abt. 2, vol. 46, p. 554. Two species.

TYPE: *Synopeas (Prosynopeas) dactylidis* Kieffer, 1916. Designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 694.

Proposed as a subgenus of *Synopeas* Foerster.

**Protelenomus** Kieffer

1908. Kieffer, Bull. Soc. Hist. Nat. Metz, vol. 25, p. 6.

TYPE: *Protelenomus flavicornis* Kieffer, 1908, by monotypy.

**Proteroscelio** Brues [fossil]

1937. Brues, in Carpenter, Univ. Toronto Studies, Geol. ser., No. 40, p. 39.

TYPE: *Proteroscelio antennalis* Brues, 1937, by monotypy and original designation.

**Protofoenus** Cockerell [fossil]

1917. Cockerell, Ann. Ent. Soc. Amer., vol. 10, p. 19.

TYPE: *Protofoenus swinhoei* Cockerell, by monotypy.

**Protrimorus** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, p. 146.

TYPE: *Trimorus americanus* Ashmead, 1893, by monotypy.

**Prozelotypa** Kieffer

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 367.

TYPE: *Prozelotypa longicornis* Kieffer, 1909, by monotypy.

**Pseudanteris** Fouts

1927. Fouts, Proc. Ent. Soc. Washington, vol. 29, p. 177.

TYPE: *Pseudanteris insignis* Fouts, 1927, by monotypy and original designation.

**Pseudaphanomerus** Szélenyi

1941. Szélenyi, Zool. Anz., vol. 134, p. 160.

TYPE: *Pseudaphanomerus hyalinatus* Szélenyi, 1941, by monotypy and original designation.

**(Pseudevania** Kieffer, error) = **Zeuxevania** Kieffer

1903. Kieffer, Zeitschr. Hym. Dipt., vol. 3, p. 111.

**Pseudobaeus** Perkins

1910. Perkins, Fauna Hawaiiensis, vol. 2, p. 620.

TYPE: *Pseudobaeus peregrinus* Perkins, 1910, by monotypy.

**Pseudoceraphron** Dodd

1924. Dodd, Trans. Roy. Soc. South Australia, vol. 48, p. 175.

TYPE: *Pseudoceraphron pulex* Dodd, 1924, by monotypy and original designation.

**Pseudofoenus** Kieffer

1902. Kieffer, in Wytzman, Genera insectorum, fasc. 2, p. 6. Three species, two of them doubtful.

TYPE: *Gasteruption pedunculatum* Schletterer, 1889. Designated by Kieffer, 1903, in André, Spec. Hym. Eur. Alg., vol. 7 bis, p. 381.

**Pseudotelenomus** Costa Lima

1928 (Sept.). Costa Lima, Compt. Rend. Soc. Biol., vol. 99, p. 880.

TYPE: *Pseudotelenomus pachycoris* Costa Lima, 1928, by monotypy and original designation.

Again described as new by Costa Lima, December 1928, Bol. Mus. Nac. Rio de Janeiro, vol. 4, p. 52.

**Psilacolus** Kieffer

1908. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 32, pp. 179, 180. Two species included by bibliographic reference.

TYPE: *Acolus xanthogaster* Ashmead, 1893. Designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 152.

Considered synonym of *Acoloides* Howard.

**Psilanteris** Kieffer

1916. Kieffer, Brotéria, Rev. Luso-Brazilcira, Zool. ser., vol. 14, p. 177. Five species.

TYPE: *Anteris bicolor* Kieffer, 1908, by original designation.

**Psilaulacus** Kieffer

1910. Kieffer, Bull. Soc. Ent. France, 1910, p. 350. No species.

1911. Kieffer, Ann. Soc. Ent. France, vol. 80, pp. 215, 230. One species.

TYPE: *Psilaulacus annulatus* Kieffer, 1911. First included species.

**(*Psilloma* Cameron, error) = *Psilomma* Foerster**

1881. Cameron, Trans. Ent. Soc. London, 1881, p. 557.

**Psilobaeus** Kieffer

1926. Kieffer, Das Tierreich, Lief. 48, pp. xvii, 132, 151.

TYPE: *Baeus curvatus* Kieffer, 1910, by monotypy.

**Psilomma** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 128, 130, 132. No species.

1861. Foerster, Programm der Realschule zu Aachen, p. xliii. One species.

TYPE: *Psilomma fusciscapis* Foerster, 1861. First included species. *Psilomma tenuicornis* Kieffer, 1908, designated by Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, p. 419, was included later. *P. tenuicornis* was described from the specimen in the Foerster collection upon which the genus was based, but under a recent decision of the International Commission on Zoological Nomenclature (Bull. Zool. Nom., vol. 4, 1950, pp. 160, 346) only the first species to be included shall be eligible for designation as type of a genus originally published without included species.

**Psilommella** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 389, 392.

TYPE: *Psilommella petiolata* Dodd, 1915, by monotypy and original designation.

**Psilopria** Kieffer

1905. Kieffer, Ann. Mus. Civ. Stor. Nat. Genova, ser. 3, vol. 2 (42), pp. 29, 35, 37.

TYPE: *Psilopria rufitarsis* Kieffer, 1905, by monotypy.

**Psiloteleia** Kieffer

1910. Kieffer, in Voeltzkow, Reise in Ostafrika, 1903-05, vol. 2, Heft 5, p. 530.

TYPE: *Psiloteleia striatigena* Kieffer, 1910, by monotypy.

**Psilus** Panzer (= *Galesus* Haliday)

1801. Panzer, Faunae insectorum Germaniae initia . . . , Heft 83, No. 11.

TYPE: *Psilus cornutus* Panzer, 1801, by monotypy.

Isogenotypic with *Galesus* Haliday.

#### **Pycnaulacus** Cushman

1930. Cushman, Proc. U. S. Nat. Mus., vol. 76, art. 25, 1929, p. 17.

TYPE: *Pycnaulacus brevicaudus* Cushman, 1930, by monotypy and original designation.

Considered synonym of *Aulacus* Jurine.

#### **(*Raia* Ashmead, not Cuvier, 1798, not Delaroché, 1809) = *Opisthacantha* Ashmead**

1893. Ashmead, U. S. Nat. Mus. Bull. 45, p. 221.

TYPE: *Opisthacantha mellipes* Ashmead, 1893. *Raia* Ashmead was validated and its type fixed by Ashmead through citation as a synonym of *Opisthacantha* Ashmead. Although it is preoccupied a new name is not required.

#### **(*Rhacodia* Herrich-Schaeffer, not Huebner [1825]) = *Dorachia*, new name**

1838. Herrich-Schaeffer, in Panzer, Faunae insectorum Germaniae initia . . . , Heft 157, No. 23. Five species.

TYPE: *Diapria striolata* Nees, 1834, by present designation.

#### **Rhacoteleia** Cameron

1905. Cameron, Spolia zeylanica, vol. 3, No. 10, p. 72.

TYPE: *Rhacoteleia pilosa* Cameron, 1905, by monotypy.

Considered synonym of *Hoploteleia* Ashmead.

#### **Rhopalopria** Kieffer

1912. Kieffer, in Wytsman, Genera insectorum, fasc. 124, 1911, pp. 8, 61.

TYPE: *Rhopalopria vulgaris* Kieffer, 1912, by monotypy.

Also described as new by Kieffer, 1912, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 71.

#### **Rhynidinofoenus** Bradley

1909. Bradley, Deutsche Ent. Zeitschr., 1909, pp. 38, 39. Two species.

TYPE: *Rhynidinofoenus kaweahensis* Bradley, 1909, by original designation.

#### **Rhynchopria** Kieffer

1910. Kieffer, in Wissenschaftliche Ergebnisse der deutschen Zentral-Afrika-Expedition . . . , vol. 3 (Zool.) Lief. 2, p. 99.

TYPE: *Rhynchopria tritoma* Kieffer, 1910, by monotypy.

#### **Rhynchopsilus** Kieffer

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 360, 400. Two species.

TYPE: *Rhynchopsilus apertus* Kieffer, 1908. Designated by Kieffer, 1910, in Wytsman, Genera insectorum, fasc. 107, p. 15.

**Rielia** Kieffer

1910. Kieffer, Boll. Lab. Zool. Portici, vol. 4, p. 107.

TYPE: *Rielia manticida* Kieffer, 1910, by monotypy.

Considered synonym of *Mantibaria* Kirby.

**Rieliomorpha** Dodd

1913. Dodd, Trans. Roy. Soc. South Australia, vol. 37, pp. 131, 155.

TYPE: *Rieliomorpha mantis* Dodd, 1915, by monotypy and original designation.

Considered synonym of *Mantibaria* Kirby.

**Roena** Cameron

1905. Cameron, Rec. Albany Mus., vol. 1, p. 225.

TYPE: *Roena cariniscutis* Cameron, 1905, by monotypy.

**Romilius** Walker

1842. Walker, Ann. Mag. Nat. Hist., vol. 10, p. 274. Two species.

TYPE: *Romilius zotale* Walker, 1842. Designated by Kieffer, 1910, in Wytsman, Genera insectorum, fasc. 80B, p. 70.

**Ropronia** Provancher (= *Roptronia* Ashmead, error)

1886. Provancher, Additions et corrections au volume II de la faune entomologique du Canada . . . Hyménoptères, pp. 152, 154.

TYPE: *Ropronia pediculata* Provancher, 1886, by monotypy.

(**Roptronia** Ashmead, error) = **Ropronia** Provancher

1898. Ashmead, Proc. Ent. Soc. Washington, vol. 4, p. 132.

(**Rosneta** Brues) = **Fidiobia** Ashmead

1908. Brues, Bull. Wisconsin Nat. Hist. Soc., vol. 6, p. 157.

TYPE: *Rosneta tritici* Brues, 1908, by monotypy and original designation.

Isogenotypic with *Fidiobia* Ashmead, 1895, through synonymy.

**Sacespalus** Kieffer

1917. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 15, p. 61.

TYPE: *Sacespalus rugosiceps* Kieffer, 1917, by monotypy and original designation.

**Sactogaster** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 108, 113. Six species.

TYPE: *Epimeces ventralis* Westwood, 1833. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 284. *Sactogaster pisi* Foerster, 1856, was designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 659.

#### **Scapopria** Kieffer

1913. Kieffer, Insecta, vol. 3, p. 441.

TYPE: *Scapopria atriceps* Kieffer, 1913, by monotypy and original designation.

#### **Sceliacantha** Dodd

1913. Dodd, Arch. Naturg., vol. 79, A6, p. 173.

TYPE: *Sceliacantha quadrispinosa* Dodd, 1913, by monotypy and original designation.

#### **Sceliacanthella** Dodd

1913. Dodd, Arch. Naturg., vol. 79, A6, p. 174.

TYPE: *Sceliacanthella parvipennis* Dodd, 1913, by monotypy and original designation.

#### **Scelio** Latreille (= *Scelionus* Rafinesque, emendation; *Serlion* Say, error; *Scelion* Blanchard, emendation)

1805. Latreille, Histoire naturelle . . . crustacés et des insectes, vol. 13, p. 226. Three species.

TYPE: *Scelio rugosulus* Latreille, 1805. Designated by Latreille, 1810, Considérations générales . . . des insectes, p. 436.

#### **Scelioliria** Brèthes

1916. Brèthes, An. Mus. Nac. Buenos Aires, vol. 27, p. 413.

TYPE: *Scelioliria mariae* Brèthes, 1916, by monotypy and original designation.

#### **Sceliomorpha** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 210, 211, 239. Two species, one of them doubtfully included.

TYPE: *Sceliomorpha longicornis* Ashmead, 1893, by original designation.

#### **(Scelion** Blanchard, emendation) = **Scelio** Latreille

1849. Blanchard, in Cuvier, Règne animal [ed. 3], insectes, text vol. 2, p. 157; atlas, pl. 116, fig. 4.

#### **(Scelionus** Rafinesque, emendation) = **Scelio** Latreille

1815. Rafinesque, Analyse, p. 125.

#### **Sceliotrachelus** Brues

1908. Brues, in Wytsman, Genera insectorum, fasc. 80, p. 13.

TYPE: *Sceliotrachelus braunsi* Brues, 1908, by monotypy and original designation.

**Schizogalesus** Kieffer

1911. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 832, 833. Fifteen species and one variety.

TYPE: *Galesus* (*Schizogalesus*) *punctatus* Kieffer, 1911. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 675.

Proposed as a subgenus of *Galesus* Curtis. Considered synonym of *Psilus* Panzer.

**Schizopria** Kieffer

1912. Kieffer, in Wytsman, Genera insectorum, fasc. 124, 1911, pp. 5, 17. Two names, both nomina nuda.

1912. Kieffer, Trans. Linn. Soc. London, ser. 2, Zool., vol. 15, p. 68. Two species.

TYPE: *Schizopria fallax* Kieffer, 1912. Designated by Kieffer, 1916, Das Tierreich, Lief. 44, p. 38.

**Scorpioteleia** Ashmead

1897. Ashmead, Can. Ent., vol. 29, p. 53.

TYPE: *Scorpioteleia mirabilis* Ashmead, 1897, by monotypy.

**Scyphodon** Brues

1925. Brues, Treubia, vol. 6, p. 93.

TYPE: *Scyphodon anomalum* Brues, 1925, by monotypy and original designation.

Doubtfully referred to the Proctotrupoidea.

(*Semaecodogaster* Bradley, new name for *Brachygaster* Stephens, 1829, not Meigen, 1826) = *Brachygaster* Leach

1905. Bradley, Can. Ent., vol. 37, p. 63. Two species.

TYPE: *Evania minuta* Olivier, 1791. Present designation. Bradley erroneously considered *Brachygaster* Leach, 1815, to be a nomen nudum, and to have been first validated by Stephens, 1829.

Isogenotypic with *Brachygaster* Leach.

(*Semaemya* Enderlein, error) = *Semaemyia* Bradley

1909. Enderlein, Stett. Ent. Zeit., vol. 70, p. 259.

*Semaemyia* Bradley (= *Semaemya* Enderlein, error)

1908. Bradley, Trans. Amer. Ent. Soc., vol. 34, pp. 163, 180. Fifteen species.

TYPE: *Semaemyia kiefferi* Bradley, 1908, by original designation.

Considered synonym of *Brachygaster* Leach.

**Sembilanocera** Brues [fossil]

1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 74, p. 70.

TYPE: *Sembilanocera clavata* Brues, 1940, by monotypy and original designation.



**Semenovius** Bradley (= *Semenovia* Kieffer, 1903, not Weise, 1889)

1908. Bradley, Trans. Amer. Ent. Soc., vol. 34, pp. 120, 123.

TYPE: *Aulacus sibiricola* Semenov, 1894, by substitution of *Semenovius* for *Semenovia* Kieffer.

(*Semenovia* Kieffer, 1903, not Weise, 1889, new name for *Anaulacus* Semenov, 1903, not MacLeay, 1825) = **Semenovius** Bradley

1903. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 7 bis, pp. 382, 720.

TYPE: *Aulacus sibiricola* Semenov, 1894, by substitution of *Semenovia* for *Anaulacus* Semenov.

(*Serlion* Say, error) = **Scelio** Latreille

1828. Say, Contr. Maclur. Lyc. Philadelphia, vol. 1, p. 80.

**Serphites** Brues [fossil]

1937. Brues, in Carpenter, Toronto Univ. Studies, Geol. ser., No. 40, p. 33.

TYPE: *Serphites paradoxus* Brues, 1937, by monotypy and original designation.

(*Serphus* Schrank) = **Proctotrupes** Latreille

1780. Schrank, Schrift. Berlin. Gesellsch. Naturf. Freunde, vol. 1, p. 307.

Suppressed in favor of *Proctotrupes* Latreille by the International Commission on Zoological Nomenclature, 1946, under suspension of the Rules (Opinion 178). Established originally as a monobasic genus with *Serphus brachypterus* Schrank, 1780, as type.

**Shreemana** Nixon

1933. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 12, p. 300.

TYPE: *Shreemana sera* Nixon, 1933, by monotypy and original designation.

(*Sinaris* Gregor, error) = **Synarsis** Foerster

1939. Gregor, Sborn. Ent. Odd. Nár. Mus. Praze, vol. 17, p. 132.

**Solenopsia** Wasmann

1899. Wasmann, Zoologica, Stuttgart, Heft 26, pp. 53, 126.

TYPE: *Solenopsia imitatrix* Wasmann, 1899, by monotypy.

**Solenopsiella** Dodd

1915. Dodd, Trans. Roy. Soc. South Australia, vol. 39, pp. 398, 428.

TYPE: *Solenopsiella distincta* Dodd, 1915, by monotypy and original designation.

(*Sparaison* Brues, error) = *Sparasion* Latreille

1908. Brues, in Wytsman, *Genera insectorum*, fasc. 80, pp. 41, 56.

*Sparasion* Latreille (= *Sparaison* Brues, error)

1802. Latreille, *Histoire naturelle . . . crustacés et des insectes*, vol. 3, p. 316.

TYPE: *Sparasion cephalotes* Latreille, 1802, by monotypy. *Ceraphron cornutus* Jurine, 1807, designated by Latreille, 1810, *Considérations générales . . . des insectes*, p. 436, and *Sparasion frontalis* Latreille, 1805, designated by Curtis, 1830, *British entomology*, vol. 7, folio 317, were included later.

*Spilomicrus* Westwood

1832. Westwood, *London Edinburgh Phil. Mag.*, ser. 3, vol. 1, p. 129.

TYPE: *Spilomicrus stigmatalis* Westwood, 1832, by monotypy.

*Stictoteleia* Kieffer

1926. Kieffer, *Das Tierreich*, Lief. 48, pp. xxix, 272, 546. Two species.

TYPE: *Macroteleia virginicensis* Ashmead, 1893, by original designation.

*Streptopria* Maneval

1939. Maneval, *Bull. Soc. Ent. France*, vol. 44, p. 168.

TYPE: *Streptopria rozieri* Maneval, 1939, by monotypy and original designation.

*Styлаclista* Dodd

1915. Dodd, *Trans. Roy. Soc. South Australia*, vol. 39, p. 389.

TYPE: *Styлаclista notabilis* Dodd, 1915, by monotypy and original designation.

(*Stylidodon* Ashmead, error) = *Stylidolon* Ashmead

1902. Ashmead, *Journ. New York Ent. Soc.*, vol. 10, p. 245.

*Stylidolon* Ashmead (= *Stylidodon* Ashmead, error)

1897. Ashmead, *Can. Ent.*, vol. 29, p. 53.

TYPE: *Stylidolon politum* Ashmead, 1897, by monotypy.

*Stylopria* Kieffer

1914. Kieffer, *Philippine Journ. Sci.*, vol. 9, p. 307.

TYPE: *Stylopria rufa* Kieffer, 1914, by monotypy and original designation.

*Styloteleia* Kieffer

1916. Kieffer, *Brotéria, Rev. Luso-Brazileira, Zool. ser.*, vol. 14, p. 184.

TYPE: *Styloteleia rufescens* Kieffer, 1916, by monotypy and original designation.

### Symphytoria Kieffer

1910. Kieffer, in André Spec. Hym. Eur. Alg., vol. 10, pp. 694, 739. Five species.

TYPE: *Symphytoria fulva* Kieffer, 1910. Designated by Kieffer, 1912, in Wytsman, Genera insectorum, fasc. 124, 1911, p. 32.

### Synacra Foerster (= *Artibolus* Haliday)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 128, 130, 134. No species.

1873. Marshall, A catalogue of British Hymenoptera; Oxyura, p. 10. One species but two names, one of them validated in synonymy.

TYPE: *Diapria brachialis* Nees, 1834. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 400.

Isogenotypic with *Artibolus* Haliday.

### Synarsis Foerster (= *Sinaris* Gregor, error)

1878. Foerster, Verh. Naturh. Ver. Preuss. Rheinl., vol. 35, p. 57.

TYPE: *Synarsis pulla* Foerster, 1878, by monotypy and original designation.

### Synopeas Foerster (= *Synopias* Ashmead, error)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 108, 114. No species.

1859. Thomson, Öfv. Vet.-Akad. Förh., vol. 16, p. 71. Thirteen species.

TYPE: *Synopeas inermis* Thomson, 1859. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., U. S. Dept. Agr., Agr. Monogr. No. 2, p. 716. *Synopeas prospectus* Foerster, 1861, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 285, *Synopeas melampus* Foerster, 1861, designated by Ashmead, 1903, Journ. New York Ent. Soc., vol. 11, p. 97, and *Platygaster sosis* Walker, 1835, designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 667, were included later.

Considered synonym of *Leptacis* Foerster.

### (*Synopias* Ashmead, error) = *Synopeas* Foerster

1896. Ashmead, Proc. Zool. Soc. London, 1895, p. 802.

### Syntelesia Fouts

1927. Fouts, Proc. Ent. Soc. Washington, vol. 29, p. 178.

TYPE: *Syntelesia coracina* Fouts, 1927, by monotypy and original designation.

**Szepligetella** Bradley

1908. Bradley, Trans. Amer. Ent. Soc., vol. 34, pp. 163, 172.

TYPE: *Evania sericea* Cameron, 1883, by monotypy and original designation.

**Tanyzonus** Marshall

1892. Marshall, Ent. Monthly Mag., vol. 28, p. 276.

TYPE: *Tanyzonus bolitophilae* Marshall, 1892, by monotypy.

**Teleas** Latreille (= *Telias* Leach, emendation)

1809. Latreille, Genera crustaceorum et insectorum . . . , vol. 4, p. 32. Four species.

TYPE: *Scelio clavicornis* Latreille, 1805. Designated by Latreille, 1810, Considérations générales . . . des insectes, p. 436.

**Telenomoides** Dodd

1913. Dodd, Trans. Roy. Soc. South Australia, vol. 37, pp. 158, 168. Seven species.

TYPE: *Telenomoides flavipes* Dodd, 1913, by original designation.

Considered synonym of *Hadronotus* Foerster.

**Telenomus** Haliday

1833. Haliday, Ent. Mag., vol. 1, p. 271. Four species.

TYPE: *Telenomus brachialis* Haliday, 1833. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 142.

**(Telias** Leach, emendation) = **Teleas** Latreille

1815. Leach, in Brewster, Edinburgh encyclopaediae, vol. 9, p. 145.

**(Telopsilus** Dalla Torre, emendation) = **Telospilus** Dahlbom

1898. Dalla Torre, Catalogus hymenopterorum, vol. 5, p. 522.

**Telospilus** Dahlbom (= *Telopsilus* Dalla Torre, emendation)

1858. Dahlbom, Öfv. Vet.-Akad. Förh., vol. 14, 1857, p. 290. No species.

No species have been referred to this genus up to the present time.

**Tetrabaesus** Kieffer

1912. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 87.

TYPE: *Aphanomerus americanus* Brues, 1908, by monotypy and original designation.

**Tetraconus** Szépligeti

1903. Szépligeti, Ann. Mus. Nat. Hungarici, vol. 1, pp. 388, 389.

TYPE: *Tetraconus mocsáryi* Szépligeti, 1903, by monotypy.

**Tetramopria** Wasmann

1899. Wasmann, *Zoologica*, Stuttgart, Heft 26, pp. 56, 127.  
Two species.

TYPE: *Tetramopria aurocincta* Wasmann, 1899. Designated by Ashmead, 1903, *Journ. New York Ent. Soc.*, vol. 11, p. 31.

**Tetrapsilus** Kieffer

1908. Kieffer, in André, *Spec. Hym. Eur. Alg.*, vol. 10, pp. 367, 397.

TYPE: *Tetrapsilus filicornis* Kieffer, 1908, by monotypy. Considered synonym of *Zygotia* Foerster.

**Tetraulacinus** Kieffer

1910. Kieffer, *Bull. Soc. Ent. France*, 1910, p. 350. No species.

1911. Kieffer, *Ann. Soc. Ent. France*, vol. 80, pp. 214, 224. One species.

TYPE: *Pristaulacus rufobalteatus* Cameron, 1907. First included species.

**Thaumatevania** Ceballos

1935. Ceballos, *Eos*, vol. 10, 1934, p. 233.

TYPE: *Thaumatevania ensifera* Ceballos, 1935, by monotypy and original designation.

**Thelepte** Nixon

1931. Nixon, *Eos*, vol. 7, p. 379.

TYPE: *Thelepte serapis* Nixon, 1931, by monotypy and original designation.

**Therinopsilus** Kieffer

1909. Kieffer, *Ann. Soc. Sci. Bruxelles*, vol. 33, p. 365. Two species.

TYPE: *Therinopsilus pubescens* Kieffer, 1909. Designated by Kieffer, 1910, in Wytzman, *Genera insectorum*, fasc. 107, p. 14.

**(Thliboneura** Thomson) = *Trichosteresis* Foerster

1858. Thomson, *Öfv. Vet.-Akad. Förh.*, vol. 15, pp. 287, 288. Four species.

TYPE: *Ceraphron glaber* Boheman, 1832. Designated by Muesebeck and Walkley, 1951, in Muesebeck et al., *U. S. Dept. Agr., Agr. Monogr. No. 2*, p. 671.

Isogenotypic with *Trichosteresis* Foerster.

**Thomsonina** Hellén

1941. Hellén, *Notulae Ent.*, vol. 21, p. 40.

TYPE: *Proctotrupes boops* Thomson, 1858, by monotypy and original designation.

**Thoron Haliday**

1833. Haliday, Ent. Mag., vol. 1, p. 271.

TYPE: *Thoron metallicus* Haliday, 1833, by monotypy.

**Tiphodytes Bradley (= *Limnodytes* Marchal, 1900, not Duméril and Bibron, 1841)**

1902. Bradley, Can. Ent., vol. 34, p. 179.

TYPE: *Limnodytes gerriphagus* Marchal, 1901, by substitution of *Tiphodytes* for *Limnodytes* Marchal.

**Tomoteleia Kieffer**

1917. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 15, p. 50.

TYPE: *Tomoteleia trifasciata* Kieffer, 1917, by monotypy and original designation.

**Trachelopteron Brues [fossil]**

1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 74, p. 86.

TYPE: *Trachelopteron angulipenne* Brues, 1940, by monotypy and original designation.

**Trichacis Foerster (= *Trichasis* Thomson, emendation)**

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 108, 115. Three species.

TYPE: *Platygaster pisis* Walker, 1835. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 294. *Platygaster remulus* Walker, 1835, was designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 711.

**Trichacoides Dodd**

1914. Dodd, Ent. News, vol. 25, p. 416.

TYPE: *Trichacoides scutellaris* Dodd, 1914, by monotypy and original designation.

**(*Trichacolus* Kieffer) = *Anteris* Foerster**

1912. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, pp. 89, 107. Two species.

TYPE: *Anteris bilineatus* Thomson, 1859, by present designation.

Isogenotypic with *Anteris* Foerster.

**Trichanteris Kieffer**

1910. Kieffer, in Wytzman, General insectorum, fasc. 80B, p. 88.

TYPE: *Prosanteris foveatifrons* Kieffer, 1908, by monotypy. Proposed as a subgenus of *Prosanteris* Kieffer.

**(*Trichasis* Thomson, emendation) = *Trichacis* Foerster**

1859. Thomson, Öfv. Vet.-Akad. Förh., vol. 16, pp. 70, 78.

**Trichasius** Provancher

1887. Provancher, Additions et corrections au volume II de de la faune entomologique du Canada . . . Hyménoptères, p. 209.

TYPE: *Trichasius clavatus* Provancher, 1887, by monotypy.

**Trichofoenus** Kieffer

1910. Kieffer, Ann. Soc. Ent. France, vol. 79, p. 77. Four species.

TYPE: *Gasteruption merceti* Kieffer, 1904, by original designation.

**Tricholeptacis** Kieffer

1914. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 357.

TYPE: *Amblyaspis verticillatus* Ashmead, 1894, by monotypy and original designation.

**Trichopria** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 407, 431. Nine species.

TYPE: *Trichopria pentaplasta* Ashmead, 1893, by original designation.

**Trichosteresis** Foerster (= *Thliboneura* Thomson)

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 97, 99. Two species.

TYPE: *Ceraphron glaber* Boheman, 1832. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 107. *Trichosteresis foersteri* Kieffer, 1907, designated by Kieffer, 1909, in Wytzman, Genera insectorum, fasc. 94, p. 14, was not originally included.

Isogenotypic with *Thliboneura* Thomson.

**Trichoteleia** Kieffer

1910. Kieffer, in Voeltzkow, Reise in Ostafrika, 1903-05, vol. 2, Heft 5, p. 530.

TYPE: *Trichoteleia albidipes* Kieffer, 1910, by monotypy.

**Triclavus** Brèthes

1916. Brèthes, An. Mus. Nac. Buenos Aires, vol. 27, p. 411.

TYPE: *Triclavus bonariensis* Brèthes, 1916, by monotypy.

**Trigonfoenus** Kieffer

1911. Kieffer, Ann. Soc. Ent. France, vol. 80, p. 177.

TYPE: *Gasteruption trianguliferum* Kieffer, 1911, by monotypy and original designation.

**Trimicrops** Kieffer

1906. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 30, p. 142. One species and two varieties.

TYPE: *Trimicrops claviger* Kieffer, 1906, by present designation.

**Trimorus** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 101, 104. Two species.

TYPE: *Gryon nanno* Walker, 1836. Designated by Ashmead, 1903, Journ. New York Ent. Soc., vol. 9, p. 87. *Gryon nanno* Walker and *Gryon phlias* Walker, 1836, were both cited as type by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 138.

**Triognus** Marshall

1874. Marshall, Ent. Ann., 1874, p. 134.

TYPE: *Triognus furcifer* Marshall, 1874, by monotypy. Considered synonym of *Lagynodes* Foerster.

**Triplatygaster** Kieffer

1913. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 11, p. 178.

TYPE: *Platygaster contorticornis* Ratzeburg, 1844, by monotypy.

Proposed as a subgenus of *Platygaster* Latreille.

**Trisacantha** Ashmead (= *Trissacantha* Ashmead, emendation)

1887. Ashmead, Ent. Amer., vol. 3, p. 117.

TYPE: *Trisacantha americana* Ashmead, 1887, by monotypy.

**Trisinostemma** Kieffer

1914. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 11, p. 357.

TYPE: *Inostemma braesia* Walker, 1839, by monotypy and original designation.

(*Trissacantha* Ashmead, emendation) = **Trisacantha** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 180, 182.

**Trissevania** Kieffer

1913. Kieffer, in Alluaud and Jeannel, Voyage . . . en Afrique Orientale, Insectes hyménoptères, vol. 1, p. 33.

TYPE: *Trissevania anemotis* Kieffer, 1913, by monotypy and original designation.

**Trissolcus** Ashmead

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 138, 161. Six species.

TYPE: *Trissolcus brochymenae* Ashmead, 1893, by original designation. *Telenomus (euschristus)* = *euschisti* Ashmead, 1888, was designated by Kieffer, 1926, Das Tierreich, Lief. 48, p. 126.



**Trissoscelio** Kieffer

1917. Kieffer, Brotéria, Rev. Luso-Brazileira, Zool. ser., vol. 15, p. 52. Three species.

TYPE: *Trissoscelio nigriceps* Kieffer, 1917, by original designation.

**Triteleia** Kieffer

1906. Kieffer, Berliner Ent. Zeitschr., vol. 50, 1905, p. 264.

TYPE: *Triteleia punctaticeps* Kieffer, 1906, by monotypy.

**Tritopria** Kieffer

1910. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 717, 748.

TYPE: *Tritopria lusitanica* Kieffer, 1910, by monotypy.

**Tropaulacus** Bradley

1908. Bradley, Trans. Amer. Ent. Soc., vol. 34, pp. 120, 125.

TYPE: *Tropaulacus torridus* Bradley, 1908, by monotypy and original designation.

(*Tropidobria* Brèthes, error for *Tropidopria* Ashmead) = **Diapria** Latreille

1927. Brèthes, Ent. Mitt., vol. 16, p. 334.

(*Tropidopria* Ashmead; *Tropidobria* Brèthes, error) = **Diapria** Latreille

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 406, 407, 416. Five species.

TYPE: *Ichneumon conicus* Fabricius, 1775, by original designation.

Isogenotypic with *Diapria* Latreille.

**Tropidopsilus** Kieffer

1908. Kieffer, in André, Spec. Hym. Eur. Alg., vol. 10, pp. 358, 366. No species.

1909. Kieffer, Ann. Soc. Sci. Bruxelles, vol. 33, p. 338. One species.

TYPE: *Tropidopsilus laticeps* Kieffer, 1909. First included species.

(*Tropidopsis* Ashmead, not Brauer and Bergenstamm, 1889) = **Nesopria**, new name

1893. Ashmead, U. S. Nat. Mus. Bull. 45, pp. 386, 402.

TYPE: *Tropidopsis clavata* Ashmead, 1893, by monotypy and original designation.

**Urocyclops** Maneval (= *Paracyclops* Maneval, 1936, not Claus, 1893)

1936. Maneval, Bull. Ann. Soc. Ent. Belgique, vol. 76, p. 142.

TYPE: *Paracyclops bettyae* Maneval, by substitution of *Urocyclops* for *Paracyclops* Maneval.

**Uroscelio** Kieffer

1914. Kieffer, Philippine Journ. Sci., vol. 9, p. 291.

TYPE: *Uroscelio luteipes* Kieffer, 1914, by monotypy and original designation.

**Uroteleia** Brues [fossil]

1940. Brues, Proc. Amer. Acad. Arts Sci., vol. 74, p. 87.

TYPE: *Uroteleia synthetica* Brues, 1940, by monotypy and original designation.

**Vanhornia** Crawford

1909. Crawford, Proc. Ent. Soc. Washington, vol. 11, p. 63.

TYPE: *Vanhornia eucnemidarum* Crawford, 1909, by monotypy and original designation.

**Vardhana** Nixon

1933. Nixon, Ann. Mag. Nat. Hist., ser. 10, vol. 12, p. 320.

TYPE: *Vardhana selene* Nixon, 1933, by monotypy and original designation.

**Vectevania** Cockerell [fossil]

1922. Cockerell, Can. Ent., vol. 54, p. 33.

TYPE: *Vectevania vetula* Cockerell, 1922, by monotypy.

**Viennopria** Jansson

1953. Jansson, Ent. Tidskr., vol. 74, p. 105.

TYPE: *Viennopria priesneri* Jansson, 1953, by monotypy.

**Xanthopria** Brues

1915. Brues, Psyche, vol. 22, p. 9. Two species.

TYPE: *Xanthopria opaca* Brues, 1915, by original designation.

**Xenomerus** Walker

1836. Walker, Ent. Mag., vol. 3, pp. 342, 355.

TYPE: *Xenomerus ergenna* Walker, 1836, by monotypy.

**Xenopria** Fouts

1939. Fouts, Proc. Ent. Soc. Washington, vol. 41, p. 260.  
Three species.

TYPE: *Xenopria columbiana* Fouts, 1939, by original designation.

**Xenoteleia** Kieffer

1913. Kieffer, Insecta, vol. 3, p. 390.

TYPE: *Xenoteleia flavipennis* Kieffer, 1913, by monotypy and original designation.

**Xenotoma** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 129, 137. No species.

1861. Foerster, Programm der Realschule zu Aachen, p. xliv. One species.

TYPE: *Xenotoma nigricoxis* Foerster, 1861. First included species. *Belyta bicolor* Jurine, 1807, as identified by Nees, 1834, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 362, was included later.

Considered synonym of *Aclista* Foerster.

**Xestonotidea** Gahan (= *Xestonotus* Foerster, 1856, not Leconte, 1853; *Axestonotus* Kieffer)

1919. Gahan, Proc. U. S. Nat. Mus., vol. 56, p. 524.

TYPE: *Xestonotus andriciphilus* Ashmead, 1887, by substitution of *Xestonotidea* for *Xestonotus* Foerster. *Xestonotidea foersteri* Gahan, 1919, was wrongly designated by Gahan, 1919, Proc. U. S. Nat. Mus., vol. 56, p. 524.

(*Xestonotus* Foerster, not Leconte, 1853) = *Xestonotidea* Gahan

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 107, 112. No species.

1887. Ashmead, Can. Ent., vol. 19, p. 128. One species.

TYPE: *Xestonotus andriciphilus* Ashmead, 1887. First included species. *Xestonotus refulgens* Foerster, designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 265, was a nomen nudum.

**Xyalopria** Kieffer

1907. Kieffer, Berliner Ent. Zeitschr., vol. 51, 1906, p. 300.

TYPE: *Xyalopria nigriceps* Kieffer, 1907, by monotypy.

**Zacranium** Ashmead

1901. Ashmead, Fauna Hawaiiensis, vol. 1, p. 295.

TYPE: *Zacranium oahuense* Ashmead, 1901, by monotypy.

**Zacrita** Foerster

1878. Foerster, Verh. Naturh. Ver. Preuss. Rheinl., vol. 35, p. 46.

TYPE: *Zacrita longicornis* Foerster, 1878, by monotypy and original designation.

**Zelotypa** Foerster

1856. Foerster Hymenopterologische Studien, Heft 2, pp. 130, 141. No species.

1890. Ashmead, Colo. Biol. Assoc. Bull. 1, p. 12. One species.

TYPE: *Zelotypa coloradensis* Ashmead, 1890. First included species.

Considered synonym of *Aclista* Foerster.

**Zeuxevania** Kieffer (= *Pseudevania* Kieffer, error)

1902. Kieffer, in Wytsman, Genera insectorum, fasc. 2, pp. 2, 4.

TYPE: *Evania dinarica* Schletterer, 1886, by monotypy.

**Zygota** Foerster

1856. Foerster, Hymenopterologische Studien, Heft 2, pp. 128, 131, 133, 135. No species.

1873. Marshall, A catalogue of British Hymenoptera; Oxyura, p. 9. Two species.

TYPE: *Belyta abdominalis* Nees, 1834. Designated by Ashmead, 1893, U. S. Nat. Mus. Bull. 45, p. 372.

## Species index

(In this index the generic or subgeneric name under which a species may be found in the catalog is the last name in each line. Where that is not the genus or subgenus in which the species was originally described, the original generic name (or generic and subgeneric names when a subgeneric name was originally used) is given in brackets. Specific synonymy is in italics.)

- abbreviatus* Foerster, *nomen nudum*,  
*Leptorhaptus*, 364  
*abdominalis* Boheman, [*Ceraphron*]  
*Megaspilus*, 368  
*abdominalis* Nees, [*Belyta*] *Zygota*,  
 410  
*abnormis* Dodd, *Mirotelenomus*, 371  
*abnormis* Perkins, [*Ceraphron*] *Allo-*  
*microps*, 326  
*abrupta* Dodd, *Hemilexomyia*, 358  
*acoluta* Foerster, *nomen nudum*,  
*Loxotropa*, 366  
*acuticornis* Kieffer, *Plastomicrops*, 385  
*afra* Kieffer, [*Evania* (*Prosevania*)]  
*Prosevania*, 391  
*africana* Risbec, *Ledouxopria*, 364  
*albicoxa* Dodd, *Acanthoserphus*, 323  
*albidipes* Kieffer, *Trichoteleia*, 405  
*albipes* Kieffer, *Dilapitha*, 346  
*albofasciatus* Dodd, *Austroserphus*,  
 334  
*aleurodinis* Haldeman, *Amitus*, 327  
*aleurodis* Forbes, *Elaptus*, 349  
*aliena* Foerster, *nomen nudum*, *Am-*  
*blyaspis*, 327  
*alticola* Kieffer, *Probelyta*, 388  
*alutaceus* Thomson, [*Megaspilus*] *Con-*  
*ostigmus*, 343  
*americana* Ashmead, *Trisacantha*, 406  
*americanus* Ashmead, *Acanthoscelio*,  
 323  
*americanus* Ashmead, [*Trimorus*] *Pro-*  
*trimorus*, 392  
*americanus* Brues, [*Aphanomerus*]  
*Tetrabaesus*, 402  
*andriciphilus* Ashmead, *Xestonotus*,  
 409  
*andriciphilus* Ashmead, [*Xestonotus*]  
*Axestonotus*, 334  
*andriciphilus* Ashmead, [*Xestonotus*]  
*Xestonotidea*, 409  
*anemotis* Kieffer, *Trissevania*, 406  
*angulipenne* Brues, *Trachelopteron*,  
 404  
*angustatus* Thomson, *Phanurus*, 384  
*annulatus* Kieffer, *Psilaulacus*, 394  
*anomala* Kirby, *Mantibaria*, 367  
*anomalipes* Panzer, [*Sphex*] *Helorus*,  
 357  
*anomalum* Brues, *Scyphodon*, 398  
*anoplocnemidis* Ghesquière, *Epinomus*,  
 351  
*anormis*, Brues, *Aneuron*, 328  
*antennale* Schletterer, [*Gasteruption*]  
*Eufoenus*, 352  
*antennale* Schletterer, [*Gasteruption*]  
*Hyptiogster*, 361  
*antennalis* Brues, *Proteroscelio*, 392  
*antennatus* Jurine, [*Psilus*] *Loxotro-*  
*pa*, 366  
*anthereae* Dodd, *Neotelenomus*, 374  
*antillensis* Risbec, *Martinica*, 367  
*apertus* Kieffer, *Rhynchopsilus*, 396  
*aphidum* Rondani, *Macrostigma*, 366  
*appendiculata* Kieffer, *Alloteleia*, 326  
*appendigaster* Fabricius, [*Sphex*] *Eva-*  
*nia*, 353  
*appendigaster* Linnaeus, [*Ichneumon*]  
*Evania*, 353  
*aptera* Szélényi, *Pachyscelidris*, 378  
*apterus* Bugnion and Popoff, [*Baeus*]  
*Aneurobaeus*, 328  
*apterygynus* Haliday, *Codrus*, 343  
*arcticus* Dahlbom, [*Aulacus*] *Disphae-*  
*ron*, 347  
*arcticus* Dahlbom, *Disphaeron*, 347  
*areolator* Haliday, *Proctotrupes*, 389  
*areolator* Haliday, [*Proctotrupes*] *Di-*  
*sognus*, 347  
*areolatus* Foerster, *Metaclisis*, 369  
*armata* Kieffer, *Phaedroteleia*, 383  
*armatus* Say, [*Ceraphron*] *Megaspi-*  
*lodes*, 368  
*ashmeadi* Mayr, *Anommatium*, 329  
*assectator* Linnaeus, [*Ichneumon*]  
*Foenus*, 354  
*assectator* Linnaeus, [*Ichneumon*]  
*Gasteruption*, 355  
*ater* Latreille, *Helorus*, 357  
*atinas* Walker, [*Inostemma*] *Monoc-*  
*rita*, 372  
*atra* Kieffer, [*Psiloteleia*] *Palpoteleia*,  
 378  
*atrata* Foerster, *Pantolyta*, 379  
*atriceps* Dodd, *Polydiapria*, 387  
*atriceps* Kieffer, *Scapopria*, 397  
*aurantia* Hope, *Caloteleia*, 339  
*auratus* Risbec, *Echinoteleas*, 349  
*aurocincta* Wasmann, *Tetramopria*,  
 403  
*australica* Dodd, *Anteromorpha*, 330  
*australis* Westwood, [*Foenus*] *Hyptio-*  
*gaster*, 361

- bakeri Kieffer, Chrestoteleia, 342  
 bambeyi Risbec, Fusicornia, 354  
 barycera Foerster, Pantoclis, 378  
 bellicosus Meunier, Acutibaesus, 325  
 beneficum Gahan, Eumicrosoma, 352  
 bethyloformis Kieffer, Paracodrus, 380  
 bettyae Maneval, Paracyclops, 380  
 bettyae Maneval, [Paracyclops] Uro-  
 cyclops, 407  
 bicolor Ashmead, Cremastobaesus, 344  
 bicolor Dodd, Mirobaesus, 371  
 bicolor Dodd, [Telenomus] Paratele-  
 nomus, 381  
 bicolor Girault, Eulagynodes, 352  
 bicolor Harrington, [Baryconus] Hol-  
 oteleia, 359  
 bicolor Jurine, Belyta, 336  
 bicolor Jurine, [Belyta] Cinetus, 342  
 bicolor Kieffer, [Anteris] Psilanteris,  
 393  
*bicolor* Kieffer, [Baryconus (Holotele-  
 leia)] Holoteleia, 359  
 bicolor (Jurine) Nees, [Belyta] Xen-  
 otoma, 409  
 bicolor Perkins, Aphanomerus, 331  
 bicoloricornis Dodd, Hoplopriella, 360  
 bidentata Kieffer, [Caloteleia] Oxyte-  
 leia, 378  
 bidentatus Dodd, [Acolus] Dissacolus,  
 347  
 bifurcata Dodd, Mantara, 367  
 bilineatus Thomson, Anteris, 330  
 bilineatus Thomson, [Anteris] Tricha-  
 colus, 404  
 bimaculatus Walckenaer, Erodorus,  
 352  
 bisulca Nees, [Belyta] Lyteba, 366  
 bisulca Nees, [Belyta] Oxylabis, 377  
 bisulcatus Kieffer, [Baryconus] Glyp-  
 toteleia, 356  
 bolitophilae Marshall, Tanyzonus, 402  
 bonariensis Brèthes, Triclavus, 405  
 boops Thomson, [Proctotrupes] Thom-  
 sonina, 403  
 boscii Jurine, [Psilus] Inostemma, 362  
 brachialis Haliday, Telenomus, 402  
 brachialis Nees, [Diapria] Artibolus,  
 332  
 brachialis Nees, [Diapria] Synacra,  
 401  
 brachyptera Thomson, [Belyta] Para-  
 clista, 380  
 brachypterus Schrank, Serphus, 399  
 braconiformis Kieffer, Neuraulacinus,  
 374  
 braesia Walker, [Inostemma] Trisi-  
 nostemma, 406  
 brasiliensis Ashmead, Megaplastopria,  
 368  
 brasiliensis Ashmead, Notoxoides, 375  
 brasiliensis Kieffer, Bactropria, 335  
 braunsi Brues, Sceliotrachelus, 397  
 brevicaudus Cushman, Pycnaulacus,  
 395  
 brevipennis Latreille, Proctotrupes,  
 389  
 brevis Nees, [Belyta] Pantoclis, 378  
 brevithorax Kieffer, Hemifoenus, 358  
 brochymenae Ashmead, Trissolcus,  
 406  
 brunnea Ashmead, [Pantolyta] Ate-  
 lopsilus, 332  
 burquei Provancher, Pammegischia,  
 378  
 californica Ashmead, [Diapria] Or-  
 thopria, 377  
 californica Ashmead, [Diapria] Plan-  
 opria Kieffer, 385  
 californica Ashmead, Hemilexis, 358  
 calopterus Kieffer, [Baryconus] Pego-  
 teleia, 382  
 campanulata Herrich-Schaeffer, En-  
 tomia, 350  
 canadensis Ashmead, Eumegaspilus,  
 352  
 canadensis Ashmead, Pentacantha,  
 382  
 canadensis Ashmead, [Pentacantha]  
 Propentacantha, 390  
 canadensis Whittaker, Paratelopsilus,  
 381  
 caraborum Riley, [Prosacantha] Allo-  
 gryon, 326  
 carinata Kieffer, Camptoteleia, 339  
 carinatifrons Dodd, Euhoplopria, 352  
 carinatus Kieffer, Aparamesius, 330  
 carinatus Kieffer, Neurogalesus, 375  
 carinifrons Cameron, Immsia, 362  
 cariniscutis Cameron, Roena, 396  
 carpenteri Curtis, [Ceraphron] Calli-  
 ceras, 338  
 caryae Ashmead, Acerota, 323  
 catillus Walker, [Platygaster] Catil-  
 lus, 340  
 catillus Walker, [Platygaster] Pies-  
 topleura, 384  
 cecidomyiiae Ashmead, Eritrissomerus,  
 352  
 cephalotes Brues, Brachyscelio, 337  
 cephalotes Latreille, Sparasion, 400  
 chalcidipennis Enderlein, [Evaniellus]  
 Chalcopterella, 341  
 charmus Walker, [Telenomus] Oxy-  
 phanurus, 377  
 chilensis Brèthes, Paridris Brèthes,  
 381  
 chlapowskii Kieffer, Pristaulacus, 388  
 ciliata Thomson, [Belyta (Opazon)]  
 Opazon, 376  
 clausus Kieffer, [Rhynchopsilus] Pro-  
 meuselia, 390  
 clavata Ashmead, Tropicopsis, 407  
 clavata Ashmead, [Tropicopsis] Neso-  
 pria, 374  
 clavata Brues, Sembilanocera, 398  
 clavata Kieffer, Aneurhopria, 328  
 clavatus Kieffer, Acanopsilus, 322  
 clavatus Kieffer, Pristomicrops, 388  
 clavatus Provancher, Trichasius, 405  
 clavicornis Cameron, Hemimorus, 358  
 clavicornis Latreille, [Scelio] Teleas,  
 402

- claviger* Kieffer, *Trimicrops*, 406  
*cleonymoides* Westwood, *Macroteleia*, 367  
*coccophagus* Foerster, *Atritomus*, 333  
*coccophagus* Foerster, [*Atritomus*] *Atritomellus*, 333  
*coeruleum* Kieffer, *Prosparasion*, 392  
*collaris* Foerster, *nomen nudum*, *Acropiasta*, 325  
*coloradensis* Ashmead, *Zelotypa*, 409  
*columbiana* Fouts, *Xenopria*, 408  
*columbianum* Ashmead, [*Psilomma*] *Propsilomma*, 390  
*comitans* Perkins, *Dyscritobaenus*, 349  
*complanata* Kieffer, *Bakeria*, 335  
*compressa* Kieffer, *Miota*, 370  
*compressicornis* Thomson, *Hypocampsis*, 360  
*compressipes* Kieffer, *Heteropria*, 359  
*concolor* Fischer, *Adelioneiva*, 325  
*conicus* Ashmead, *Leptorhaptus*, 364  
*conicus* Fabricius, [*Ichneumon*] *Diapria*, 345  
*conicus* Fabricius, [*Ichneumon*] *Tropidopria*, 407  
*conjungens* Kieffer, *Pentapria*, 383  
*contorticornis* Ratzeburg, [*Platygaster*] *Hypocampsis*, 360  
*contorticornis* Ratzeburg, [*Platygaster*] *Triplatygaster*, 406  
*coracina* Fouts, *Synteleia*, 401  
*cordata* Kieffer, [*Hemilexis* (*Entomacis*)] *Entomacis*, 350  
*corni* Haldeman, *Amitus*, 327  
*cornutus* Ashmead, *Ceratobaenus*, 341  
*cornutus* Jurine, [*Ceraphron*] *Sparasion*, 400  
*cornutus* Panzer, *Psilus*, 395  
*cornutus* Panzer, [*Psilus*] *Galesus*, 354  
*cotta* Walker, [*Platygaster*] *Isocybus*, 362  
*crassiceps* Schletterer, [*Gasteruption*] *Hyptiogaster*, 361  
*crassiclava* Kieffer, *Lophopria*, 366  
*crassicornis* Ashmead, *Acanthopria*, 323  
*crassipes* Kieffer, *Linkiola*, 365  
*craterus* Walker, [*Platygaster*] *Ectadius*, 349  
*craterus* Walker, [*Platygaster*] *Polymecus*, 387  
*crawfordi* Dodd, *Gryonella*, 356  
*cumaeus* Nixon, *Nardo*, 373  
*curtipennis* Haliday, [*Proctotrupes*] *Phaenoserphus*, 383  
*curvatus* Kieffer, [*Baeus*] *Psilobaenus*, 394  
*cydnoides* Priesner, *Eremioscelio*, 351  
*dactylidis* Kieffer, [*Synopeas* (*Prosynopeas*)] *Prosynopeas*, 392  
*depressus* Brues, *Proplatyscelio*, 390  
*desertorum* Priesner, *Megacolus*, 368  
*desertorum* Priesner, [*Megacolus*] *Philoplanes*, 384  
*dilatatus* Westwood, *Platymischus*, 386  
*dimidiatus* Thomson, *Dichogmus*, 346  
*dinariae* Schletterer, [*Evania*] *Zeuxevania*, 409  
*discessus* Szelényi, *Inocerota*, 362  
*dispar* Nees, [*Diapria*] *Loxotropa*, 366  
*distincta* Dodd, *Solenopsiella*, 399  
*donisthorpei* Nixon, *Brunnicophilus*, 338  
*dorsiger* Curtis, [*Cinetus*] *Ismarus*, 362  
*dubitata* Brues, *Baemorpha*, 335  
*dux* Curtis, [*Ceraphron*] *Megaspilus*, 368  
*ecitophila* Holmgren, *Mimopria*, 370  
*editus* Cresson, [*Aulacus*] *Odontaulacus*, 376  
*egregia* Herrich-Schaeffer, *Anisoptera*, 329  
*egregia* Herrich-Schaeffer, [*Anisoptera*] *Laches*, 363  
*electrinus* Cockerell, *Hyptiogasterites*, 361  
*elegans* Jurine, [*Psilus*] *Diapria*, 345  
*elegans* Kieffer, *Micraulacinus*, 369  
*elegans* Perkins, [*Caloteleia*] *Caenoteleia*, 338  
*elongata* Ashmead, [*Anteris*] *Prosaepagus*, 391  
*ensifer* Westwood, *Epimeces*, 351  
*ensifer* Westwood, [*Epimeces*] *Parepimeces*, 381  
*ensifera* Ceballos, *Thaumatovania*, 403  
*ergenna* Walker, *Xenomerus*, 408  
*error* Fitch, [*Platygaster*] *Euxestonotus*, 353  
*erynniae* Kieffer, [*Tropidopria*] *Neodiapria*, 374  
*erythrothorax* Ashmead, *Eumegaspilus*, 352  
*erythrothorax* Ashmead, [*Eumegaspilus*] *Conostigmoides*, 343  
*eucnemidarum* Crawford, *Vanhornia*, 408  
*europus* Walker, [*Inostemma*] *Eurostemma*, 353  
*euschisti* Ashmead, [*Telenomus*] *Trissolcus*, 406  
*euschristus* Ashmead, [*Telenomus*] *Trissolcus*, 406  
*evanescens* Kieffer, *Acerota*, 323  
*exaratus* Kieffer, *Enneascelio*, 350  
*exsculptus* Foerster, *Hadronotus*, 357  
*exsertus* Dodd, *Dissolcoides*, 347  
*fallax* Kieffer, *Schizopria*, 398  
*fasciatipennis* Kieffer, *Lamproteleia*, 363  
*fasciatus* Ashmead, *Aradophagus*, 331  
*ferruginea* Dodd, *Pantolytomyia*, 379  
*festus* Walker, [*Inostemma*] *Moninostemma*, 372  
*filicornis* Kieffer, *Tetrapsilus*, 403  
*firmus* Cresson, [*Aulacus*] *Oleisopriester*, 376  
*flavibarbis* Marshall, *Aleria*, 326

- flavicauda Ashmead, Acropiesta, 325  
 flavicornis Foerster, Idris, 361  
 flavicornis Kieffer, Protelenomus, 392  
 flavimanus Kieffer, Disaulacinus, 347  
 flavimanus Kieffer, Monoxylabis, 372  
 flavipennis Kieffer, Xenoteleia, 408  
 flavipes Ashmead, Fidiobia, 354  
 flavipes Dodd, Cremastoscilio, 344  
 flavipes Dodd, Telenomoides, 402  
 flavipes Kieffer, Dicroscelio, 346  
 flavipes Kieffer, Digalesus, 346  
 flavipes Kieffer, Doliopria, 348  
 flavipes Kieffer, Habroteleia, 357  
 flavipes Thomson, Ceratacis, 340  
 flaviventris Thomson, [Cinetus] Pantopiasta, 379  
 flavosignatus Kieffer, [Amblyaspis] Prosamblyaspis, 391  
 floricola Kieffer, Prosectogaster, 391  
 floridana Ashmead, Baeoneura, 335  
 floridana Ashmead, Hemilexodes, 358  
 floridanus Ashmead, Baryconus, 335  
 floridanus Ashmead, [Baryconus] Hoploteleia, 360  
 foersteri Gahan, Xestonotidea, 409  
 foersteri Kieffer, [Leptacis] Proleptacis, 389  
 foersteri Kieffer, Plastogryon, 385  
 foersteri Kieffer, Polypeza, 387  
 foersteri Kieffer, Trichosteresis, 405  
 formicaria Kieffer, Bruchiola, 337  
 formicarius Kieffer, [Platygaster] Diplatygaster, 347  
 formicarius Kieffer, Exallonyx, 353  
 formicarium Kieffer, Aulacopria, 333  
 formosus Kieffer, Parafoenus, 380  
 foveatifrons Kieffer, [Prosanteris] Trichanteris, 404  
 foveatus Kieffer, Oxyscelio, 377  
 frequens Priesner, Aegyptoscelio, 325  
 froggatti Johnston and Tiegs, Paraspilomicrus, 381  
 frontalis Latreille, Sparasion, 400  
 frontalis Latreille, [Sparasion] Bebelus, 336  
 frontalis Latreille, [Sparasion] Oxyurus, 378  
 fulva Cameron, Betyla, 336  
 fulva Kieffer, Symphytopria, 401  
 fumipennis Thomson, Aphanogmus, 331  
 fumipennis Westwood, Basalys, 336  
 fureifer Marshall, Triogmus, 406  
 fusca Kieffer, Clinopria, 342  
 fuscator Perty, [Pelecinius] Monomachus, 372  
 fuscicornis Kieffer, Meuselia, 369  
 fuscicornis Kieffer, Pezopria, 383  
 fuscicornis Kieffer, Prospilomicrus, 392  
 fuscipennis Ashmead, Megaspilodes, 368  
 fuscipennis Curtis, *nomen nudum*, Galesus, 354  
 fuscipennis Kieffer, Coenopria, 343  
 fuscipennis Kieffer, Lepidoscelio, 364  
 fuscipes Kieffer, Nanopria, 373  
 fusciscapis Foerster, Psilomma, 394  
 fuscoclavata Kieffer, Hexapria, 359  
 galesiformis Westwood, Aneurhynchus, 328  
 galesiformis Westwood, [Aneurhynchus] Mythras, 372  
 gallardi Brèthes, Notilena, 375  
 gallicola Kieffer, Plastobelyta, 385  
 gallicola Szélényi, Platyllostropa, 386  
 gerriphagus Marchal, Limnodytes, 365  
 gerriphagus Marchal, [Limnodytes] Tiphodytes, 404  
 giraudi Kieffer, Prosynacra, 392  
 giraulti Dodd, [Baeoncura] Baeoncur-ella, 335  
 giraulti Dodd, Mallateleia, 367  
 glaber Boheman, [Ceraphron] Thiboneura, 403  
 glaber Boheman, [Ceraphron] Trichosteresis, 405  
 glabra Ashmead, Miota, 370  
 gladiator Brullé, Monomachus, 372  
 gladiator (Klug) Westwood, Monomachus, 372  
 globiceps Dodd, Aratala, 331  
 gloriosus Dodd, Neoscelio, 374  
 gracilipes Cockerell, Electrofoenus, 349  
 gracilipes Curtis, Cinetus, 342  
 gracilis Kieffer, Dolichopria, 348  
 grandis Nees, [Platygaster] Isocybus, 362  
 gravidator Linnaeus, [Ichneumon] Proctotrupes, 389  
 grenadensis Ashmead, [Caloteleia] Ceratoteleia, 341  
 grenadensis Ashmead, [Macroteleia] Monoteleia, 372  
 gryontoides Priesner, Efflatounina, 349  
 hackeri Dodd, Ceratobaeoides, 341  
 halidayi Foerster, [Ismarus] Entomius, 350  
 heterocera Haliday, [Belyta] Pantolyta, 379  
 hirtifrons Ashmead, Anectata, 328  
 hirtipes Kieffer, Dissoxylabis, 347  
 hofmanni Kieffer, Homophanurus, 359  
 holotoma Kieffer, [Aclista] Anaclista, 327  
 hopkinsi Crawford and Bradley, Dolichotrypes, 348  
 horni Brèthes, Kiefferopria, 363  
 hospes Kieffer, Plesiobaeus, 386  
 humerale Schletterer, [Gastruption] Odontofoenus, 376  
 hungaricus Szélényi, Elysoceraphron, 350  
 hyalinata Thomson, Hypocampsis, 360  
 hyalinatus Szélényi, Pseudaphanomerus, 393  
 hyalinipennis Ashmead, Isorhombus, 362  
 imitatrix Wasmann, Solenopsia, 399  
 impiger Szélényi, *nomen nudum*, Neanteris, 373



- inermis Thomson, Synopeas, 401  
 infortunata Nixon, Gita, 355  
 innupta Haliday, Labolips, 363  
 insignis Fouts, Pseudanteris, 393  
 iridipennis Lepeletier and Serville,  
 Cinetus, 336, 342  
 jaculator Linnaeus, [Ichneumon] Foe-  
 nus, 354  
 jaculator Linnaeus, [Ichneumon] Gas-  
 teruption, 355  
 jeanneli Risbec, Brachyscelio, 337  
 jeanneli Risbec, [Brachyscelio] Pach-  
 yscelio, 378  
 kaweahensis Bradley, Rhydinofoenus,  
 395  
 kiefferi Bradley, Interaulacus, 362  
 kiefferi Bradley, Semaemyia, 398  
 kiefferi Crawford, Cologlyptus, 343  
 koyamae Ishii, Neolygocerus, 374  
 kristenseni Turner, Brachevania, 337  
 lacustris Kieffer, Anopedias, 329  
 laeviceps Ashmead, Idris, 361  
 laeviceps Ashmead, [Idris] Paridris  
 Kieffer, 381  
 laevifrons Foerster, [Proctotrupes]  
 Carabiphagus, 339  
 laminata Dodd, Bracalba, 336  
 lar Walker, Iphitrachelus, 362  
 lasiopterae Kieffer, Disynopeas, 348  
 laticeps Foerster, *nomen nudum*, Had-  
 ronotus, 357  
 laticeps Kieffer, Anoxylabis, 329  
 laticeps Kieffer, Hadronotus, 357  
 laticeps Kieffer, Tropidopsilus, 407  
 laticinctus Ashmead, Calliscelio, 338  
 latigaster Brues, Antaretropria, 330  
 latipennis Dodd, Platyteleia, 386  
 lepidus Mayr, Macrohynnis, 366  
 leptocerus Foerster, Apegus, 331  
 leroyi Benoit, Afrevania, 325  
 lichtensteinii Ratzeburg, Dendrocerus,  
 345  
 lineatus Kieffer, [Platygaster] Pro-  
 sactogaster, 391  
 longicalcar Kieffer, [Serphus (Crypto-  
 serphus)] Cryptoserphus, 344  
 longiceps Ashmead, Ceratopria, 341  
 longiceps Kieffer, Geodiapria, 355  
 longiceps Kieffer, Odontacolus, 376  
 longicornis Ashmead, Ceratopria, 341  
 longicornis Ashmead, Sceliomorpha,  
 397  
 longicornis Foerster, Zacrita, 409  
 longicornis Kieffer, Acidopsilus, 324  
 longicornis Kieffer, Prozelotypa, 393  
 longicornis Nees, Prosacantha, 390  
 longicornis Wasmann, Labidopria, 363  
 longii Ashmead, Adeliopria, 325  
 lucifugax Kieffer, Antroscelio, 330  
 luctuosa Kieffer, Plutopria, 387  
 lugens Kieffer, Heptascelio, 358  
 lusitanica Kieffer, Tritopria, 407  
 luteipes Kieffer, Uroscelio, 408  
 luteus Cameron, [Trimorus] Dicha-  
 cantha, 346  
 macroneurus Ashmead, [Ceraphron]  
 Neoceraphron, 373  
 macronyx Schletterer, [Gasteruption]  
 Crassifoenus, 344  
 maculipennis Kieffer, Pleuropria, 387  
 maculipennis Kieffer, Prosoxylabis,  
 391  
 manticida Kieffer, Rielia, 396  
 mantis Dodd, Rieliomorpha, 396  
 mariae Brèthes, Scelioliria, 397  
 maritima Haliday, [Diapria (Miono-  
 pria)] Idiotypa, 361  
 maritima Haliday, [Diapria (Miono-  
 pria)] Mionopria, 370  
 marshalli Kieffer, [Oxylabis] Acan-  
 thopsilus, 323  
 marylandica Fouts, [Trichopria] Cra-  
 cinopria, 344  
 maura Kieffer, Coelopria, 343  
 maxillosa Risbec, Elgonia, 349  
 meecrida Walker, [Inostemma] Allo-  
 tropa, 326  
 mediterranea Kieffer, Brachinostem-  
 ma, 337  
 melampus Foerster, Synopeas, 401  
 mellea Ashmead, [Loxotropa] Myr-  
 mecopria, 372  
 mellipes Ashmead, Opisthacantha, 377  
 mellipes Ashmead, [Opisthacantha]  
 Raia, 395  
 mellipetiola Ashmead, Hemilexis, 358  
 mellipetiolata Ashmead, Hemilexis, 358  
 merceti Kieffer, [Gasteruption] Trich-  
 ofoenus, 405  
 metallicus Haliday, Thoron, 404  
 mila Nixon, Govinda, 356  
 minuscula Benoit, Micrevania, 369  
 minuta Ashmead, Megaspilidea, 368  
 minuta Dodd, Acolomorpha, 324  
 minuta Olivier, [Evania] Brachy-  
 gaster, 337  
 minuta Olivier, [Evania] Coranila,  
 344  
 minuta Olivier, [Evania] Semaecodo-  
 gaster, 398  
 minutissima Ashmead, Phaenopria,  
 383  
 minutissima Ashmead, [Prosacantha]  
 Hoplogryon, 359  
 minutus Westwood, Hemisius, 358  
 mirabilis Ashmead, Coelopelta, 343  
 mirabilis Ashmead, Scorpioteleia, 398  
 mirabilis Brues, Nothoserphus, 375  
 mirabilis Dodd, Acanthobetyla, 322  
 mirabilis Dodd, Miramblyaspis, 371  
 mirabilis Dodd, Platygastoides, 386  
 mirissimus Dodd, Encrytoscelio, 350  
 misellus Haliday, Gryon, 356  
 mocsáryi Szépligeti, Tetraconus, 402  
 moerens Westwood, [Aulacus (Aula-  
 cinus)] Aulacinus, 333  
 molokaiensis Ashmead, Platymischoi-  
 des, 386  
 moniliata Herrich-Schaeffer, Chlido-  
 nia, 342

- moniliatus Nees, [Bracon] Chlidonia, 342  
 montanus Cresson, [Aulacus] Deraiodontus, 345  
 montanus Kieffer, Apteroscelio, 331  
 monticola Kieffer, Aholcus, 326  
 morpheus Nixon, Nirupama, 375  
 muchini Martynov, Mesohelorus, 369  
 multifoveata Dodd, Propentapria, 390  
 muscidarum Kieffer, Platyceraphron, 385  
 mutilata Ferrière, Euplacopria, 352  
 nanno Walker, [Gryon] Trimorus, 406  
 nervalis Dodd, Neuroscelio, 375  
 niger Kieffer, Hemigalesus, 358  
 niger Panzer, Codrus, 343  
 niger Panzer, [Codrus] Proctotrupes, 389  
 nigerrima Kieffer, Prosinostemma, 391  
 nigra Kieffer, Abothropria, 322  
 nigra Kieffer, Eriopria, 351  
 nigriceps Ashmead, [Anteris] Prosanteris, 391  
 nigriceps Kieffer, Camptopsilus, 339  
 nigriceps Kieffer, Doddiella, 348  
 nigriceps Kieffer, Holoteleia, 359  
 nigriceps Kieffer, [Macroteleia] Baryteleia, 336  
 nigriceps Kieffer, Prolapitha, 389  
 nigriceps Kieffer, Prosynacra, 392  
 nigriceps Kieffer, Trissoscelio, 407  
 nigriceps Kieffer, Xyalopria, 409  
 nigriclava Kieffer, Paranteris, 380  
 nigriclavata Ashmead, Glyptonota, 356  
 nigriclavus Kieffer, Acanthoteleia, 323  
 nigricollis Cameron, Acantholapitha, 323  
 nigricornis Ashmead, Dissolcus, 347  
 nigricoxa Dodd, [Sparasion] Austroscelio, 334  
 nigricoxis Foerster, Xenotoma, 409  
 nigripedius Nakagawa, Asolcus, 332  
 nigriventris Dodd, Polypria, 387  
 nitida Kieffer, Carinia, 340  
 nitida Kieffer, Microteleia, 370  
 nitida Riek, Austronia, 334  
 notabilis Dodd, Stylaclista, 400  
 oahuense Ashmead, Zacranium, 409  
 obscurus Thomson, Anopediias, 329  
 occidentalis Cresson, [Aulacus] Neaulacus, 373  
 oecanthi Riley, Baryconus, 335  
 oecanthi Riley, [Baryconus] Leptoteleia, 364  
 oecanthi Riley, Cacus, 338  
 oecanthi Riley, [Cacus] Cacellus, 338  
 oecanthi Riley, [Cacus] Oethecoctonus, 376  
 opaca Brues, Xanthopria, 408  
 opacus Thomson, Acolus, 324  
 opacus Thomson, [Acolus] Holacolus, 359  
 orchymonti Debauche, Anacoryphe, 327  
 ornatus Kieffer, [Platygaster] Miso-cyclops, 371  
 ovi Dodd, Aphanomerella, 331  
 oviformis Martynov, Mesaulacinus, 369  
 ovivora Riley, Caloptenobia, 339  
 oxygaster Marshall, Marshalliella, 367  
 oxylaboides Jansson, Acanthomiota, 323  
 oxyura Cockerell, Palaeoteleia, 378  
 pachycoris Costa Lima, Pseudotelenomus, 393  
 pallida Kieffer, Mesoteleia, 369  
 pallida Kieffer, Pantolyta, 379  
 pallidus Boheman, [Ceraphron] Lagynodes, 363  
 pallipes Jurine, [Codrus] Proctotrupes, 389  
 paradoxus Brues, Serphites, 399  
 paradoxus Provancher, Copelus, 343  
 paraguayense Schrottky, [Gasterup-tion] Plutofoenus, 387  
 parvipennis Dodd, Sceliacanthella, 397  
 parvula Haliday, [Belyta (Opazon)] Opazon, 376, 377  
 parvula Nees, [Diapria] Monelata, 371  
 parvulus Kieffer, Calogalesus, 339  
 parvulus Nees, [Codrus] Brachyserphus, 337  
 pauliani Risbec, Mandraka, 367  
 pedester Kieffer, Hadronotellus, 357  
 pedestris Kieffer, Antropria, 330  
 pedestris Kieffer, Lepidopria, 364  
 pedestris Kieffer, Paragryon, 380  
 pedestris Kieffer, Planopria Kieffer, 1910, 385  
 pedestris Kieffer, [Planopria] Planopriella, 385  
 pediculata Provancher, Ropronia, 396  
 pedunculatum Schletterer, [Gasterup-tion] Pseudofoenus, 393  
 penicillata Borgmeier, Neivapria, 373  
 pennsylvanicus Ashmead, [?Telenomus] Hadrophanurus, 357  
 pentaplasta Ashmead, Trichopria, 405  
 pentatoma Kieffer, Bruchopria, 337  
 pentatomus Dodd, Hadronotoides, 357  
 peregrinus Perkins, Pseudobaeus, 393  
 pergandei Ashmead, Polypeza, 387  
 pergandei Ashmead, [Polypeza] Propantolyta, 390  
 perplexus Kieffer, Paratrimorus, 381  
 peruanus Enderlein, Evaniellus, 353  
 petiolaris Nees, [Diapria] Corynopria, 344  
 petiolata Dodd, Psilommella, 394  
 petiolata Fabricius, [Evania] Hyptia, 361  
 phalaenarum Nees, [Teleas] Prophanurus, 390  
 philippinensis Kieffer, Neurocacus, 374  
 phlias Walker [Gryon] Trimorus, 406  
 piceiventris Kieffer, Acolus, 324

- picipes Nees, [Cinetus] Oxylabis, 377  
 pilosa Cameron, Rhacoteleia, 395  
 pini Debauche, Eoxestonotus, 351  
 pisi Foerster, Sactogaster, 397  
 pisis Walker, [Platygaster] Trichacis, 404  
 planus Dodd, Platytenomus, 386  
 platyptera Haliday [Diapria (Glyphidopria)] Entomacis, 350  
 platyptera Haliday, [Diapria (Glyphidopria)] Glyphidopria, 355  
 platyptera Haliday, [Diapria (Glyphidopria)] Hemilexis, 358  
 pluto Nixon, Macrogyron, 366  
 polita Say, Coptera, 343  
 politum Ashmead, Styldolon, 400  
 polycerator Fabricius, [Ichneumon] Epiceuastes, 351  
 polycerator Fabricius, [Ichneumon] Pelecinus, 382  
 polyturator Drury, [Ichneumon] Pelecinus, 382  
 priesneri Jansson, Viennopria, 408  
 primus Crawford, Lipoglyptus, 365  
 princeps Westwood, [Evania] Acanthinevania, 322  
 productus Kieffer, Cardiopsilus, 339  
 prospectus Foerster, Synopeas, 401  
 prosper Nixon, Nasdia, 373  
 pubescens Kieffer, Therinopsilus, 403  
 pulchellus Dodd, Gryonoides, 356  
 pulchricornis Dodd, Neobetyla, 373  
 pulchricornis Kieffer, Platyscelio, 386  
 pulchripennis Ashmead, Hoplopria, 359  
 pulex Dodd, Pseudoceraphron, 393  
 pulla Foerster, Synarsis, 401  
 punctata Cameron, Malvina, 367  
 punctata Dodd, Neoteleia, 374  
 punctaticeps Kieffer, Pentapria, 383  
 punctaticeps Kieffer, Triteleia, 407  
 punctatus Kieffer, [Apegus (Parapegus)] Parapegus, 381  
 punctatus Kieffer, [Galesus (Schizogalesus)] Schizogalesus, 398  
 punctiger Nees, [Platygaster] Isostasius, 363  
 quadridens Kieffer, Amblyconus, 327  
 quadridens Kieffer, Microgalesus, 370  
 quadridentata Dodd, Merriwa, 368  
 quadrispinosa Dodd, Scelicantha, 397  
 quinda Walker, [Inostemma] Parinostemma, 382  
 radiatus Kieffer, [Cinetus] Eccinetus, 349  
 radiatus Kieffer, [Cinetus] Procinetus, 388  
 ramicornis Boheman, [Ceraphron] Lygocerus, 366  
 raphidioides Westwood [Foenus] Dolichofoenus, 348  
 ratzeburgi Ashmead, [Dendrocerus] Prodendrocerus, 389  
 reducta Cockerell, Lithobelyta, 365  
 refulgens Foerster, *nomen nudum*, Xestonotus, 409  
 remulus Walker, [Platygaster] Trichacis, 404  
 retractus Dodd, Aclistoides, 324  
 rottensis Meunier, Proctotrypites, 389  
 rozieri Maneval, Streptopria, 400  
 rubi Haliday, Microps, 370  
 rubriventer Philippi, Aulacostethus, 334  
 rubriventer Philippi, [Aulacostethus] Aulacomastus, 333  
 rubrosignata Ogloblin, Gastroxylabis, 355  
 rufa Kieffer, Asolenopsia, 332  
 rufa Kieffer, Atrichopria, 333  
 rufa Kieffer, Neuroteleia, 375  
 rufa Kieffer, Notoxopria, 375  
 rufa Kieffer, Phoenoteleia, 384  
 rufa Kieffer, Stylopria, 400  
 rufescens Kieffer, Plagioscelio, 385  
 rufescens Kieffer, Styloleia, 401  
 rufescens Westwood, Loboscelidia, 365  
 ruficornis Kieffer, Parabaeus, 379  
 ruficornis Latreille, [Scelio] Platygaster, 386  
 rufipes Ashmead, Platygaster, 386  
 rufipes Kieffer, Aneurosclio, 328  
 rufipes Kieffer, Baryconus, 335  
 rufipes Kieffer, Ceratopsilus, 341  
 rufipes Kieffer, Dilobopria, 346  
 rufipes Kieffer, Diphropria, 346  
 rufipes Kieffer, Proplatygaster, 390  
 rufipes Westwood, Parnamesius, 380  
 rufitarsis Cresson, [Aulacus] Odon-taulacus, 376  
 rufitarsis Foerster, *nomen nudum*, Anteris, 330  
 rufitarsis Kieffer, Psilopria, 394  
 rufithorax Kieffer, Atomopria, 332  
 rufithorax Kieffer, [Chromoteleia] Petalosema, 383  
 rufiventris Kieffer, Cyathopria, 345  
 rufobalteatus Cameron, [Pristaulacus] Tetraulacinus, 403  
 rufobasalis Dodd, Phanuromyia, 384  
 rufopetiolata Nees, [Belyta] Acoretus, 325  
 rufum Kieffer, Acanosema, 322  
 rufus Foerster, Lagynodes, 363  
 rufus Kieffer, Philolestes, 384  
 rufus Westwood, [Foenus] Carinafoenus, 339  
 rugosa Kieffer, Dicroteleia, 346  
 rugosiceps Kieffer, Saccespalus, 396  
 rugosulus Latreille, Scelio, 397  
 rugosus Brues, Archaeosclio, 332  
 rugosus Kieffer, Dichoteleas, 346  
 sagax Kieffer, [Plastogryon] Heterogryon, 359  
 saitidis Howard, Acoloides, 324  
 saussurei Kieffer, Bothriopria, 336  
 scutellaris Boheman, [Ceraphron] Habropelte, 356  
 scutellaris Dodd, Trichacoides, 404  
 scutellaris Thomson, [Acoretus] Ac-  
 lista, 324  
 sechellensis Kieffer, Oreiscelio, 377

- secundus* Cockerell, *Aulacites*, 333  
*seeveri* Wing, *Bruesopria*, 337  
*selene* Nixon, *Vardhana*, 408  
*semaeoda* Bradley, *Evaniella*, 353  
*semicyanea* Ashmead, *Chromoteleia*, 342  
*semiflaviventris* Girault, *Phanuropsis*, 384  
*seminulum* Haliday, *Baeus*, 335  
*seminulum* Haliday, [*Baeus*] *Hyperbaeus*, 360  
*semiramis* Nixon, *Lapithoides*, 363  
*semirufa* Kieffer, *Parevania*, 381  
*semirufa* Kieffer, *Paroxylabis*, 382  
*semistriatus* Nees, [*Teleas*] *Aphanurus*, 331  
*semistriatus* Nees, [*Teleas*] *Microphanurus*, 370  
*sera* Nixon, *Shreemana*, 399  
*serapis* Nixon, *Thelepte*, 403  
*sericea* Cameron, [*Evania*] *Szepligetella*, 402  
*sericeicornis* Spinola [*Psilus*] *Cerapsilon*, 340  
*serraticeps* Priesner, *Chaetanteris*, 341  
*seychellensis* Kieffer, *Oriscelio*, 377  
*sibiricola* Semenov, [*Aulacus*] *Anaulacus*, 327  
*sibiricola* Semenov, [*Aulacus*] *Semenovius*, 399  
*sibiricola* Semenov, [*Aulacus*] *Semenowia*, 399  
*sicarius* Brues, *Metanopedias*, 369  
*simulus* Kieffer, *Anteris*, 330  
*sociabilis* Kieffer, *Neuropria*, 375  
*sodalis* Brues, *Auxopaeutes*, 334  
*solida* Thomson, [*Corynopria*] *Entomopria*, 351  
*sordidus* Dodd, [*Baryconus*] *Dibaryconus*, 345  
*sisis* Walker, [*Platygaster*] *Synopeas*, 401  
*spatulatus* Brues, *Gastrotrypes*, 355  
*spilosomatis* Ashmead, [*Telenomus*] *Liophanurus*, 365  
*spinifera* Brèthes, *Passalida*, 382  
*spinosa* Ashmead, *Lapitha*, 363  
*spinosa* Brues, *Parabetyla*, 379  
*spinosus* Kieffer, [*Baryconus* (*Probaryconus*)] *Probaryconus*, 388  
*splendida* Dodd, *Mallateleioides*, 367  
*splendida* Dodd, *Pantolytoidea*, 379  
*stigmatica* Brues, *Electroteleia*, 349  
*stigmaticalis* Westwood, *Spilomicrus*, 400  
*strangaliophagus* Nixon, *Platystasius*, 386  
*striaticeps* Dodd, *Nyleta*, 375  
*striaticeps* Kieffer, *Amblyscelio*, 327  
*striatifrons* Kieffer, *Odontoscelio*, 376  
*striatigena* Kieffer, *Procacus*, 388  
*striatigena* Kieffer, *Psiloteleia*, 394  
*striatus* Jurine, *Aulacus*, 334  
*striolata* Nees, [*Diapria*] *Dorachia*, 348  
*striolata* Nees, [*Diapria*] *Rhacodia*, 395  
*striolatus* Cameron, *Apegusoneura*, 331  
*striolatus* Kieffer, *Anectadius*, 327  
*striolatus* Nees, [*Platygaster*] *Polygnotus*, 387  
*subaptera* Brues, *Probetyla*, 388  
*subapterus* Brues, *Ecitonetes*, 349  
*subtangens* Kieffer, [*Evania* (*Prosevania*)] *Prosevania*, 391  
*subterraneus* Szelényi, *Microceraphron*, 369  
*sulcata* Nees, *Calliceras*, 338  
*sulcata* Nees, [*Calliceras*] *Hadroceras*, 357  
*sulcatus* Jurine, *Ceraphron*, 340  
*superba* Meunier, *Archabelyta*, 332  
*swinehoi* Cockerell, *Protofoenus*, 392  
*synergorum* Kieffer, *Fahringeria*, 353  
*synthetica* Brues, *Uroteleia*, 408  
*szepligetii* Kieffer, [*Hyptiogaster*] *Aulacofoenus*, 333  
*tasmanicus* Dodd, *Mirobaeoides*, 371  
*temporalis* Kieffer, *Odontopria*, 376  
*tenuicornis* Dodd, [*Holoteleia*] *Duta*, 348  
*tenuicornis* Kieffer, *Miotella*, 370  
*tenuicornis* Kieffer, *Odontopsilus*, 376  
*tenuicornis* Kieffer, *Psilomma*, 394  
*termitarii* Dodd, *Leaiopria*, 364  
*testaceus* Risbec, *Ceranogmus*, 340  
*thoracica* Kieffer, *Oxypria*, 377  
*thoracicus* Ashmead, [*Scelio*] *Discelio*, 347  
*tibialis* Boheman, [*Ceraphron*] *Habropelte*, 356  
*tibialis* Szépligeti, *Evaniscus*, 353  
*tipulae* Kirby, [*Ichneumon*] *Leptacis*, 364  
*torridus* Bradley, *Tropaulacus*, 407  
*trianguliferum* Kieffer, [*Gasterurpation*] *Trigonofoenus*, 405  
*trifasciata* Kieffer, *Tomoteleia*, 404  
*trifoveata* Dodd, *Neopria*, 374  
*tritici* Brues, *Rosneta*, 396  
*tritici* Curtis, [*Platygaster*] *Amblyaspis*, 327  
*tritoma* Kieffer, *Rhynchopria*, 395  
*tritomus* Thomson, *Anopedias*, 329  
*tubuliforme* Brues, *Pelecinopterum*, 382  
*tucumana* Brèthes, *Aulatopria*, 334  
*umbraculi* Kieffer, *Prosactogaster*, 391  
*undulatus* Dodd, *Parascelio*, 381  
*unicolor* Ashmead, [*Evania*] *Evaniella*, 353  
*urichi* Ashmead, *Embidobia*, 350  
*variicornis* Kieffer, *Acidopria*, 324  
*veniprivus* Priesner, *Cyphacolus*, 345  
*ventralis* Dodd, *Cardiopria*, 339  
*ventralis* Westwood, *Epimeces*, 351  
*ventralis* Westwood, [*Epimeces*] *Sactogaster*, 397  
*venustella* Dodd, *Duarina*, 348

verticillata Latreille, Diapria, 345	virginiensis Ashmead, [Macroteleia]
verticillata Latreille, [Diapria] Ash-	Stictoteleia, 400
meadopria, 332	vulgaris Kieffer, Rhopalopria, 395
verticillatus Ashmead, [Amblyaspis]	walkeri Foerster, Catillus, 340
Tricholeptacis, 405	wasmanni Ferrière, Philolestoides, 384
vespiformis Kieffer, Neuraulacinus,	westwoodii Foerster, Diphora, 346
374	wheeleri Brues, Galesimorpha, 354
vetula Cockerell, Vectevania, 408	xanthogaster Ashmead, Acolus, 324
vibius Walker, [Telenomus] Allopha-	xanthogaster Ashmead, [Acolus] Psi-
nurus, 326	lacolus, 393
virginiensis, Ashmead, Anteris, 330	zotale Walker, Romilius, 396