A MATHEMATICAL AND STATISTICAL ANALYSIS OF THE -3/2 POWER RULE OF SELF-THINNING IN EVEN-AGED PLANT POPULATIONS

A Dissertation
Presented for the
Doctor of Philosophy
Degree

The University of Tennessee, Knoxville

Donald Ernest Weller
June 1985

Dedicated to my grandparents

Della Florence (Miller) Weller

and

Everett Earl Weller

ACKNOWLEDGMENTS

The successful completion of a doctoral program requires the assistance of many individuals. In my case, particular thanks are due to Bob Gardner, Hank Shugart, Jr., Don DeAngelis, and Bob O'Neill for their guidance in this and other projects undertaken during my graduate study at Oak Ridge National Laboratory.

Committee members Tom Hallam, Dewey Bunting, and Cliff Amundsen also provided invaluable assistance in designing my graduate program and preparing this dissertation.

I thank the Graduate Program in Ecology for providing the basic matrix for my graduate education. My fellow graduate students, who are to numerous to be listed here, provided an intellectually stimulating atmosphere that was a essential component of the graduate school experience. Dewey Bunting and Cornell Gilmore helped me to avoid numerous bureaucratic pitfalls and facilitated my graduate program in many other ways. Ray Millemann provided similar assistance in coordinating the interactions between the University of Tennessee and Oak Ridge National Laboratory. Carole Hom, Yetta Jager, and Tom Smith helped me to meet important deadlines after my departure from Tennessee.

The staff and management of Oak Ridge National Laboratory, particularly the Environmental Sciences Division, maintained an outstanding working environment that included the routine provision of facilities and services unavailable to most graduate students. The Word Processing Center of the Environmental Sciences Division prepared the final copy of the dissertation.

I thank my family for encouraging my education, particularly my parents Lowell and Eloise Weller and my grandparents Everett and Della Weller. My wife Debbie deserves special thanks for her love, friendship, encouragement, and continued patience during the difficult stages of my graduate career.

I thank the scientists whose work provided the background for this study. They developed the ideas and data vital to a review of this kind, and many sent additional information to extend and clarify published reports. Since this group is too numerous to be listed here, the "Literature Cited" section should be considered an extended list of acknowledgements. Without the studies cited there, my project would have had neither motivation nor basis.

This research was supported by the National Science
Foundation's Ecosystems Studies Program under Interagency Agreement
BSR-8315185 with the U.S. Department of Energy, under contract
No. DE-AC05-840R21400 with Martin Marietta Energy Systems, Inc.

ABSTRACT

The self-thinning rule for even-aged plant populations (also called the -3/2 power law or Yoda's law) is reviewed. This widely accepted but poorly understood generalization predicts that, through time, growth and mortality in a crowded population trace a straight thinning line of slope -3/2 in a log-log plot of average plant weight versus plant density. The evidence for this rule is examined, then reanalyzed to objectively evaluate the strength of support for the rule. Mathematical models are constructed to produce testable predictions about causal factors.

Major problems in the evidence for the thinning rule include inattention to contradictory data, lack of hypothesis testing, inappropriate curve-fitting techniques, and the use of an invalid data transformation. When these problems are corrected, many data sets thought to corroborate the rule do not demonstrate any size-density relationship. Also, the variations among thinning slopes and intercepts are much greater than currently accepted, many slopes disagree quantitatively with the thinning rule, and thinning slope and intercept differ among plant groups.

The models predict that thinning line slope is determined by the allometry between area occupied and plant weight, while the intercept is also related to the density of biomass per unit of space occupied and the partitioning of resources among competing individuals. Statistical tests confirm that thinning slope is correlated with several measures of plant allometry and that variations in thinning slope among plant groups reflect allometric differences.

The ultimate thinning line, which describes the overall size-density relationship among populations of many species, is a trivial geometric consequence of packing objects onto a surface. This cause differs from the factors positioning the self-thinning lines of individual populations, so the existence of an overall relationship is not relevant to the thinning rule.

The evidence does not support acceptance of the self-thinning rule as a quantitative biological law. The slopes and intercepts of size-density relationships are variable, and the slopes can be explained by simple geometric arguments.

vii

TABLE OF CONTENTS

CHA	PTER	PAGE
INT	RODUCTION	1
1.	LITERATURE REVIEW Statement of the Self-thinning Rule Evidence for the Self-thinning Rule Effects of Environmental Factors on Self-thinning Lines Explanations for the Self-thinning Rule Interpretation of the Self-thinning Constant K	11
2.	SIMPLE MODELS OF SELF-THINNING Introduction Model Formulation Basic Model for the Spatial Constraint Enhanced Model with Additional Constraints Model Analysis and Results Basic Model Enhanced Model Discussion Basic Model Enhanced Model	37 38 38
3.	SIMULATION MODEL OF SELF-THINNING Introduction	47 53 57
4.	SOME PROBLEMS IN TESTING THE SELF-THINNING RULE Introduction	77 83 87

viii

CHAF	PTER																											PAGE
5.	NEW TE Intro Metho Resu Disc	oduc ods lts	tio •	n •	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	102 102 103 109 120
6.	SELF-TI Intro Expe	oduc	tic	n		•		•						•		•			•	•		•						128 128
		ant ods lts	A11 •	on •	et	ry • •	•	•	•	•	•	•	•	•	•	•	•	•	:	•	•	•	•	•	•		•	128 133 134 151
7.		oduc ls o uris prov	tic f t tic ed	n the Mo	od ode	ve el l	ra	11 •		i iz	:e-	·de	ens	it		Re	•la	et	ior	nsk •	nip	•	•					158 158 159 159 161 164
8.	CONCLU	SION	•	•	•		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	179
LITE	ERATURE	CIT	ED	•	•	•	•		•	•	•	•		•				•	•			•			•		•	184
APPE	ENDIXES APPEND APPEND	IX A		ΕX	PE	RΙ	ΜE	ΝT	ΑL	Α	ND	F	ΙE	LD) [(AC	A											198 199 218
VITA	٠																											240

LIST OF TABLES

TABLE		Р	AGE
3.1.	Competition Algorithms in the Simulation Model		51
3.2.	Simulation Model Parameters and Their Reference Values	•	56
3.3.	Self-thinning Lines for the Simulation Experiments	•	59
3.4.	Thinning Line Statistics for the Simulation Experiments	•	60
4.1.	Three Examples of the Sensitivity of the Fitted Self- thinning Line to the Points Chosen for Analysis	•	81
5.1.	Statistical Distributions of Thinning Line Slope and Intercept	•	110
5.2.	Comparisons of Thinning Line Slope and Intercept Among Plant Groups	•	113
5.3.	Spearman Correlations of Shade Tolerance with Thinning Line Slope and Intercept in the Forestry Yield Data .	•	115
5.4.	Thinning Line Slopes and Intercepts of Species for which Several Thinning Lines Were Fit from Experimental or Field Data	•	116
5.5.	Statistics for Thinning Line Slope and Intercept of Species for which Several Thinning Lines Were Fit from Forestry Yield Data	•	121
6.1.	Statistical Distributions of Allometric Powers for Experimental and Field Data	•	136
6.2.	Statistical Distributions of Allometric Powers for Forestry Yield Table Data	•	137
6.3.	Correlations and Regressions Relating Transformed Thinning Slope to Allometric Powers	•	140
6.4.	Tests for Differences in Allometric Powers Among Groups in the Experimental and Field Data	•	144

TABLE		PAGE
6.5.	Tests for Differences in Allometric Powers Among Groups in the Forestry Yield Table Data	147
6.6.	Spearman Correlations of Shade Tolerance with Allometric Powers in the Forestry Yield Table Data	149
7.1.	Parameter Ranges for Monte Carlo Analysis of the Improved Model	165
7.2.	Ranges of Variation of Variables Derived in the Monte Carlo Analysis of the Improved Model	166
7.3.	Statistical Distributions of κ and log κ	173
A.1.	Sources of Experimental and Field Data	200
A.2.	Self-thinning Lines Fit to Experimental and Field Data	203
A.3.	Ranges of Log B and Log N Used to Fit Self-thinning Lines to Experimental and Field Data	207
A.4.	Fitted Allometric Relationships for Experimental and Field Data	211
A.5.	Self-thinning Lines from Experimental and Field Data Cited by Previous Authors as Support for the Self-thinning Rule	216
B.1.	Sources of Forestry Yield Table Data	219
В.2.	Self-thinning Lines Fit to Forestry Yield Table Data	221
в.3.	Fitted Allometric Relationships for Forestry Yield Table Data	230
B.4.	Self-thinning Lines from Forestry Yield Table Data Cited by Previous Authors in Support of the Self-thinning Rule	239

LIST OF FIGURES

FIGUR	E	PAGE
1.1.	Four possible growth stages for an idealized even-aged plant population	9
2.1.	Power functions of the crowding index	27
2.2.	Vertical distance between the zero isocline of biomass growth and the self-thinning line	35
2.3.	Basic model fitted to data from a Pinus strobus plantation	36
2.4.	Analysis of a model with three constraints on population growth	39
2.5.	Effect of a reduction in illumination on population trajectories in a two-constraint model	45
3.1.	Typical dynamic behavior of the simulation model	58
3.2.	Variations in the self-thinning trajectory due to stochastic factors in the simulation model	62
3.3.	Effect of the allometric power, p, on the self-thinning line	63
3.4.	Effect of the density of biomass in occupied space, d, on the self-thinning line	64
3.5.	Effect of the competition algorithm on the self-thinning line	66
3.6.	Simulations for three parameters that did not affect the self-thinning line	68
3.7.	Dynamics of the weight distributions in four simulations with different competition algorithms	73

FIGUR	E	PAGE
4.1.	Two forestry yield tables showing variation in thinning line parameters with site index	78
4.2.	Three examples of the sensitivity of self-thinning line parameters to the points chosen for analysis	80
4.3.	Example of a data set with two regions of linear behavior	82
4.4.	Thinning lines fit to one data set by four methods	85
4.5.	Example of a spurious reported self-thinning line	91
4.6.	Example of potential bias in data editing in the log $\overline{w}\text{-log N}$ plane	92
4.7.	Example of deceptive straightening of curves in the log $\bar{w}\text{-log N}$ plane	94
4.8.	Potential misinterpretation of experimental results due to deceptive effects of the log $\overline{w}\text{-log N}$ plot	95
5.1.	Histograms for the slopes and intercepts of fitted thinning lines	112
6.1.	Histograms of allometric powers of the experimental and field data	138
6.2.	Histograms of allometric powers of the forestry yield table data	139
6.3.	Observed relationships between transformed thinning slope and allometric powers in the experimental and field data	141
6.4.	Observed relationships between transformed thinning slope and allometric powers in the forestry yield table data .	143
7.1.	Monte Carlo analysis of the improved model for the overall size-density relationship	167

xiii

FIGURE		PAGE
7.2.	Previous analysis of the overall size-density relationship among thinning lines	170
7.3.	Gorham's analysis of the overall size-density relationship among stands of different species	
7.4.	Histograms of $\log \kappa$	174

xiv

LIST OF ABBREVIATIONS AND MATHEMATICAL SYMBOLS

ABBREVIATIONS

SYMBOL	MEANING
ANOVA	analysis of variance
BSLA	basal area of tree bole at breast height
CI	confidence interval
CV	coefficient of variation
DBH	diameter of tree bole at breast height
EFD	experimental and field data
%EV	percentage of the summed variance of two
FYD	variables explained by bivariate PCA forestry yield table data
GMR	geometric mean regression
ln	natural logarithm
log, log ₁₀	common logarithm
PCA	principal component analysis
RGR	relative growth rate
RMR	relative mortality rate
RHS	right hand side of an equation or inequality
SD	standard deviation
SE	standard error

SYMBOL	MEANING
VOI	volume of influence; space filled
ZOI	zone of influence; area covered
-	(bar) indicates average value, e.g. $\bar{\text{w}}$, $\bar{\text{a}}$
α	sign of proportionality
∞	infinite; without limit
^	indicates a statistical estimate of a parameter, e.g. $\hat{\beta}$, $\hat{\alpha}$

xvi

MATHEMATICAL SYMBOLS

SYMBOL	CHAPTER	MEANING
a	1-3,6-7	growing surface area occupied by a plant
aloss	3	area of a plant ZOI lost to a competitor
А	3	total area covered by a population
Aplot	3	size (area) of a simulated plot
b	1,3	constant in a power function relating maintenance cost to weight; cost = b wq
В	all	total population biomass; stand biomass
B _{max}	2	maximum biomass that a population can attain; carrying capacity for biomass
С	2,3,7	constant in a power function relating ZOI radius to weight
сl	2,3,7	constant defined by $c_1 = \pi c^2$
c ₂	2	constant correcting for systematic differences between the $\overline{a}\text{-}\overline{w}$ and a-w power relationships
c ₃	2	constant related to the allowable overlap between plant ZOIs
C(N,B)	2	function for crowding at given biomass and density
Cov(X,Y)4	covariance of variables X and Y
d	3	density of biomass per unit volume occupied
D _{ij}	3	distance between the $i^{\mbox{th}}$ and $j^{\mbox{th}}$ plants in a population
f	2	a constant in the crowding function $C(N,B)$

xvii

SYMBOL	CHAPTER	MEANING
F	5,6	the variance ratio computed in ANOVA
90	2	maximum relative growth rate of biomass
91	1,3	maximum assimilation rate per unit of ground area covered
G(N,B)	2	function for the growth rate of population biomass
G _r (N,B)	2	multiplier function reducing RGR_B with increasing crowding
Н	5,6	test statistic from Kruskal-Wallis analysis
h	2-7	plant height
K	all	constant in the self-thinning rule equation
m	2	mortality constant
M(N,B)	2	function for population mortality rate
M _r (N,B)	2	multiplier function producing higher relative mortality with increasing crowding
N	all	plant density in individuals per unit area
n	1-7	size of a sample for statistical analysis
n _t	3	number of plants remaining in a simulation at time t
n_0	3	initial number of plants in a simulation
p	1-7	power relating ZOI radius to plant weight; R = c w^p
pest	6	power relating ZOI radius plant weight, estimated by mathematical transformation of a fitted thinning slope; $p_{est} = 0.5 / (1-\beta)$

xviii

SYMBOL	CHAPTER	MEANING
Р	1-7	statistical significance level; probability of a Type I statistical error
q	3	power relating maintenance cost to weight; cost = b wq
r	3 - 7	correlation coefficient
r ²	3-7	coefficient of determination
r_S	3,6	Spearman rank correlation coefficient
R	1-7	ZOI radius
RGR	1-3	relative growth rate
rgr _B	1-3	relative growth rate of total biomass
$RGR_{\overline{W}}$	1-3	relative growth rate of average weight
RGR _{min}	3	minimum individual RGR allowing survival
RMR	1-3	relative mortality rate
$s_{\bar{w}_0}$	3	standard deviation of weights at time O
s _{xx} , s _{yy} , s _{xy}	4	sums of squares and cross products for X and Y
t	2,3	time
Var(X)	4	variance of variable X
V	1-3,6-7	volume of an individual plant VOI
W	all	weight of an individual plant
W	all	average plant weight for a population

SYMBOL	CHAPTER	MEANING
w _O	3	initial plant weight at time O
₩max	2	maximum individual weight
Х, Ү	4	general symbols for two variables involved in a bivariate relationship
α	1-6	common log of K (α = log K); intercept at log N = 0 of the self-thinning line
α3.35	3	<pre>intercept of the self-thinning line at log N = 3.35</pre>
β	all	power in the B-N thinning equation B = K N^{β} ; slope of the log B-log N self-thinning line
βt	2	<pre>instantaneous slope of the self-thinning trajectory at time t</pre>
β_{yx}	4	slope of a linear function giving Y in terms of X
Υ	all	power in the $\overline{w}-N$ thinning equation $\overline{w}=K$ NY; slope of the log $\overline{w}-\log N$ self-thinning line
Δ log K	2	difference between the log N = 0 intercepts of the straight lines in the log B-log N plane representing the asymptotic population trajectory and the zero isocline of biomass increase
Δt	3	simulation time step
(∆t) _m	3	simulation time step over which RGR's are averaged to identify plants with untenably low growth rates
ε	3	expansion factor to correct for edge effects in the simulation model
η	3	sample angle in edge correction equation
ζ	7	parameter of the lognormal distribution; if w is lognormal, ζ is the mean of ln w

SYMBOL	CHAPTER	MEANING
θ1 - θ5	2	parameters controlling the suddenness of onset of model constraints
λ	4	ratio of the residual variances in Y and X around a true linear relationship
κ	7	quantity defined for a stand by κ = B N $^{0.5};$ an index of stand shape and biomass density per unit of volume occupied
ρ	7	ratio of the weights of the largest and smallest individuals in a population
σ	7	parameter of the lognormal distribution; if w is lognormal, σ is the standard deviation of ln w
τ	3,7	ratio of plant height to ZOI radius
Φ0, Φ1	6	constant and power, respectively, in an allometric power relationship between two plant dimensions
ΦBw	6	allometric power relating BSLA to weight; B \propto w ΦB w
$^\Phi$ dw	6	allometric power relating density to weight; d \propto w $^{\Phi dw}$ (density of biomass per unit of volume occupied)
$^\PhiDw$	6	allometric power relating DBH to weight; DBH \varpropto w ΦD W
$^\Phi$ h B	6	allometric power relating height to BSLA; h \propto BSLA $^\Phi$ hB
[⊕] hD	6	allometric power relating height to DBH; h \propto DBH $^\Phi$ hD
$^\Phi$ hw	6	allometric power relating height to weight; h $\propto w^{\Phi h w}$

SYMBOL	CHAPTER	MEANING
Ψ	2	slope of a straight line in the log B-log N plane representing points where d log B / d log N = Ψ
ω	2	ratio of mortality and growth constants; $\omega = m / g_0$

INTRODUCTION

The title of a recent review article "Ecology's law in search of a theory" (Hutchings 1983) concisely summarizes the status of a nypothesis variously referred to as the self-thinning rule, the -3/2 power law, or Yoda's law. This rule states that as growth and mortality proceed in a crowded even-aged plant population, average weight (\bar{w}) and the number of plants per unit area (N) are related by a simple power equation $\bar{w} = K N^{\gamma}$, where $\gamma = -3/2$ and K is a constant. Widespread acceptance of this rule by plant ecologists is based on many observations of this power relationship in plant populations ranging from mosses to trees, but the reasons for the rule's apparent generality are not well understood (Hutchings and Budd 1981a, Westoby 1981, White 1981, Hutchings 1983).

The theoretical importance of the self-thinning rule is evidenced by the published statements of plant ecologists. White (1981) called it one of the best documented generalizations of plant demography, and further states that its empirical generality is now beyond question. Westoby (1981) considers it the most general principle of plant demography and suggests that it be elevated beyond the status of an empirical generalization to take a "central place in the concepts of population dynamics." Hutchings and Budd (1981a) emphasized the uniqueness of its precise mathematical formulation to a science where most general statements can be stated in only vaguer, qualitative terms. To many ecologists, the rule is

sufficiently well verified to be considered a scientific law (Yoda et al. 1963, Dirzo and Harper 1980, Lonsdale and Watkinson 1982, Malmberg and Smith 1982, Hutchings 1983). Harper (quoted in Hutchings 1983) called it "the only generalization worthy of the name of a law in plant ecology" and McIntosh (1980) agreed that, if substantiated, a self-thinning law could well be the first basic law demonstrated for ecology. As such, it would help to fill the need for verified laws in a discipline that has been hindered by a lack of laws and other regularities from which to develop the body of comprehensive theory that is the hallmark of a mature science (Johnson 1977, McIntosh 1980).

There have been many proposed extensions and applications of the self-thinning rule. Although it was originally observed in monocultures, evidence now suggests that aggregate measurements of even-aged two-species mixtures (White and Harper 1970, Bazzaz and Harper 1976, Malmberg and Smith 1982) and many-species mixtures (White 1980, Westoby and Howell 1981) also conform. Some authors have even attempted to apply the thinning rule to animal populations (Furnas 1981, Wethey 1983). The rule can be a used to compare the site qualities or histories of plant populations growing at different sites, since relative positions along the curve $\overline{w} = K N^{-3/2}$ give a ranking of the fertilities experienced by equal-aged populations (Yoda et al. 1963), or of population ages if fertility is constant among sites (Barkham 1978). The existence of the -3/2 power relationship among a set of population measurements has been interpreted as evidence that natural self-thinning is

occurring, canopies are closed, growth and mortality are ongoing, and competition is the cause of mortality (Barkham 1978), while the absence of the -3/2 relationship has been cited as evidence that density independent factors are the important causes of mortality (Schlesinger 1978). The rule can also be a useful management tool in forestry (see Yoda et al. 1963, Drew and Flewelling 1977, and Japanese language papers cited therein), or in other applications requiring predictions of the limits of biomass production for a given species at any density (Hutchings 1983).

In view of the popularity, theoretical importance, and applicability of the self-thinning rule, the troublesome lack of a verified explanatory theory continues to motivate further research. The original objective of the present study was to examine the causes of the self-thinning rule by analyzing a detailed simulation model of a generalized plant population and extracting testable predictions about the effects of different biological parameters on the constants γ and K of the power rule equation. The central question was how the processes of growth and mortality operate to eliminate large among-species variations in size, shape, physiological capability, strategy, and other important factors so that all species obey the same power rule. The model developed for this purpose did not predict such constancy, rather it suggested that the power γ should vary from the idealized value of -3/2, with its exact value determined by the relationship between ground area covered and plant weight in a given stand. This hypothesis is neither new (see Miyanishi et. al 1979) nor well supported in previous studies (Westoby 1976, Mohler et al. 1978, White 1981).

Among the ideas considered after reaching this conclusion was the thought that the previous analyses were somehow wrong. When the evidence behind the self-thinning rule was carefully examined, some troubling problems were discovered. They motivated a new analysis of the data supporting the thinning rule and some new tests of the simple hypothesis suggested by the simulation model. Together, the mathematical models and data analyses evolved into six lines of investigation presented in Chapters 2 through 7 of this report.

Chapter 2. Simple, spatially-averaged models of growth and mortality are developed to formalize in a dynamic model the hypothesis that the power of the self-thinning equation is determined by a relationship between ground area covered and plant weight, and to examine the interaction between this effect and constraints imposed by a carrying capacity and a maximum individual size.

Chapter 3. A detailed, spatially explicit computer simulation model is analyzed to consider the effects of additional population parameters on the power rule relationship, and to verify that the conclusions of the simpler models are not artifacts of spatial averaging.

Chapter 4. Some difficulties in the selection, analysis, and interpretation of self-thinning data are discussed along with the implications for the self-thinning rule. Remedies for some problems are suggested.

Chapter 5. Biomass and density data for 488 self-thinning relationships are statistically analyzed to evaluate support for the self-thinning rule and to estimate the variability of the power equation parameters K and γ . Variations in these parameters among plant groups are considered.

Chapter 6. The biomass and density data from Chapter 5 are combined with plant shape measurements from the same stands and used to test some predictions of the hypothesis that the power of the self-thinning equation is determined by a relationship between ground area covered and plant weight.

Chapter 7. Some hypotheses are presented to explain the overall relationship between size and density. (This relationship emerges when measurements of many crowded plant stands ranging from small herbs to trees are combined to estimate a single power equation relating stand size and stand density across the entire plant kingdom--Gorham 1979, White 1980). The relevance of this overall relationship to the self-thinning rule for individual plant populations is considered.

These efforts are united by an overall goal of developing and testing a unified theoretical framework for understanding observed size-density relationships and evaluating their scientific importance.

CHAPTER 1

LITERATURE REVIEW

Statement of the Self-thinning Rule

The self-thinning rule describes a relationship between size and density in even-aged plant populations that are crowded but actively growing. In the absence of competition from other populations and of significant density-independent stresses (drought, fire, etc.), mortality or "thinning" is caused by the stresses of competition within the population, hence the term "self-thinning." Yoda et al. (1963) observed a general relationship among successive measurements of average weight and density taken after the start of self-thinning. When the logarithm of average weight is plotted vertically and the logarithm of plant density horizontally, the points form a straight line with a slope near -3/2 represented by the equation

$$\log \tilde{W} = \gamma \log N + \log K, \tag{1.1}$$

where \bar{w} is average weight (in g), N is plant density (in individuals/m²), γ = -3/2, and K is a constant. Antilogarithmic transformation of this relationship gives the power equation

$$\bar{w} = K N^{\gamma}$$
 (1.2)

Since $\bar{w} = B/N$, (where B is total stand biomass or yield in g/m^2), these equations relating average weight to density are

mathematically equivalent to the following relationships between stand biomass and density:

$$\log B = \beta \log N + \log K \tag{1.3}$$

and

$$B = K N^{\beta}, \qquad (1.4)$$

with β = -1/2. The parameters γ and β are related by β = γ + 1.

Yoda et al. (1963) named this relationship the "-3/2 power law of self-thinning" because of the mathematical form of equation 1.2 and the value of the power γ . Any of the above relationships can be referred to as the self-thinning equation, while the linear equations 1.1 and 1.3 give the self-thinning line, or simply the thinning line. γ and β are called self-thinning powers or exponents in reference to equations 1.2 and 1.4, but the names self-thinning slope or thinning slope are also common because these parameters are the slopes of the linear equations 1.1 and 1.3. The logarithm of the constant K is often called the self-thinning intercept because it gives the intersection of the thinning line with the vertical line log N = 0.

The self-thinning rule does not necessarily apply throughout the history of a plant population. When populations are established at combinations of average weight and density well below the self-thinning line, the initial competitive stresses can be absorbed through plastic changes in shape, growth can proceed with little

mortality, and a nearly vertical line is traced in a log-log plot of size versus density. As competition becomes more severe, growth slows and some plants die after exhausting their capacities to adjust to their neighbors' intrusions. The population's path in log size-log density plot bends toward the self-thinning line, then begins to move along it as growth and mortality progress (Hutchings and Budd 1981a, White 1981). The curves traced by populations of differing initial densities converge on the same self-thinning line (Yoda et al. 1963, White 1980, 1981). Movement along the self-thinning line can not continue indefinitely because the population will eventually approach the environmental carrying capacity, which places an upper limit on total stand biomass (Hutchings and Budd 1981a, Peet and Christensen 1980, Lonsdale and Watkinson 1983b, Watkinson 1984). Limits on individual growth, such as a genetically or mechanically defined maximum size or the end of the growing season, can also deflect the population from the thinning line (Harper and White 1971, Peet and Christensen 1980).

The history of an even-aged population can, then, be divided into up to four stages: (1) a period of initial establishment, rapid growth, and low mortality; (2) a period of adherence to the self-thinning rule; (3) a period when constant biomass is maintained at the carrying capacity; and (4) a period of population degeneration, when growth does not replace the biomass lost through mortality. Figure 1.1 shows the idealized representions of these stages as paths in the log \bar{w} -log N and log B-log N planes.






Figure 1.1. Four possible growth stages for an idealized even-aged plant population. (1) initial rapid growth and low mortality, (2) self-thinning rule, (3) constant biomass at carrying capacity, and (4) terminal degeneration or senescence. In the $\log \bar{w} - \log N$ plane, (a) the respective slopes of the last three stages are -3/2, -1, and 0, while the corresponding slopes in the $\log B - \log N$ plane (b) are -1/2, 0, and +1. In both plots, the line representing the first stage is nearly vertical.

The self-thinning rule describes a certain balance between the rates of growth and mortality, which are linked so that the ratio of the relative growth rate to the relative mortality rate is held constant (Harper and White 1971, Westoby and Brown 1980) at the value of the self-thinning slope, that is,

$$\frac{RGR_{\bar{W}}}{RMR} = \frac{d\bar{w} / (\bar{w} dt)}{dN / (N dt)} = \frac{d \log \bar{w}}{d \log N} = \gamma, \qquad (1.5)$$

where ${\rm RGR}_{\widetilde{W}}$ is the relative growth rate of average weight and RMR is the relative mortality rate (Hozumi 1977). The equivalent relation for the relative growth rate of biomass, ${\rm RGR}_{\rm B}$, is

$$\frac{RGR_B}{RMR} = \frac{dB / (B dt)}{dN / (N dt)} = \frac{d \log B}{d \log N} = \beta.$$
 (1.6)

Mortality during self-thinning is concentrated among smaller individuals that have been suppressed, possibly because of smaller seed size, later germination, lower growth rate, or close neighbors (Ross and Harper 1972, Kays and Harper 1974, Ford 1975, Hutchings and Barkham 1976, Bazzaz and Harper 1976, Harper 1977, Mohler et al. 1978, Rabinowitz 1979, Hutchings and Budd 1981a, Lonsdale and Watkinson 1983a). These deaths permit a net biomass production among the remaining plants so that total population biomass increases (Ford 1975, Hutchings and Barkham 1976, Westoby 1981).

The self-thinning line can also be interpreted as a constraint separating possible combinations of biomass and density from impossible ones. Biomass and density combinations below or on the

line are possible, while combinations above the line do not occur (Yoda et al. 1963; White 1980, 1981; Westoby and Howell 1981). This means that thinning lines need not be measured by following individual populations through time, rather data for a given population type from several plots of different ages can be used, provided that the plots do not differ in important biotic or environmental factors that would alter the position of the thinning line. This method has been used since the earliest studies (Yoda et al. 1963), and is particularly important for studying long-lived trees (White 1981).

Evidence for the Self-thinning Rule

The self-thinning rule is an empirical statement generalized from repeated observations of linear relationship between log \overline{w} -log N with an estimated slope near -3/2. The supporting examples include populations ranging in size from small herbs to large trees collected from experimental or natural conditions or from forestry yield tables. Yoda et al. (1963) presented ten data sets which exhibit such a relationship. White and Harper (1970) added five additional examples, plus some evidence from forestry thinning tables. White (1980) presented 36 more examples and mentioned unpublished data for 80 additional cases. White's paper has been widely cited by other authors, along with the study of Gorham (1979), as firmly establishing the generality of the -3/2 rule (Hutchings 1979, Dirzo and Harper 1980, Furnas 1981, Westoby

and Howell 1982, Lonsdale and Watkinson 1982, 1983a, Watkinson et al. 1983). Tables A.5 and B.5 give references to studies presenting additional corroborative examples. Most of the weight measurements in these studies are of aboveground plant parts only, but some also include roots. This inclusion raises the value of K but does not affect the slope of the thinning line (Watkinson 1980, Westoby and Howell 1981).

The limited range of variation of thinning line slopes and intercepts are also considered strong evidence for the thinning rule. White (1980) reported that thinning slopes vary between -1.3 and -1.8 when $\log \bar{w}$ is fitted to $\log N$. This range is considered remarkably invariant (Watkinson et al. 1980, Furnas 1981, Lonsdale and Watkinson 1982, Hutchings 1983); however, several exceptions have been reported (Ernst 1979, O'Neill and DeAngelis 1981, Sprugel 1984). The constant K is usually limited to 3.5 \leq log K \leq 4.4 and values outside this range are biologically significant (White 1980, 1981). Grasses can give higher log K values between 4.5 (Kays and Harper 1974) and 6.67 (Lonsdale and Watkinson 1983a), possibly because their erect, linear leaves allow deeper light penetration through the canopy and greater plant biomass per unit volume (Lonsdale and Watkinson 1983a, Watkinson 1984). Some values of log K greater than 4.4 have also been reported for herbaceous dicots (Westoby 1976, Westoby and Howell 1981, Lonsdale and Watkinson 1983a).

Further support for the self-thinning rule has come from two studies that examined the relationship between size and density among stands of different species (Gorham 1979, White 1980). This relationship is referred to here as the overall size-density relationship to distinguish it from the self-thinning lines of particular populations. Gorham (1979) reported that measurements of 65 stands of 29 species (including mosses, reeds, herbs, and trees) form a straight line of slope -1.5 in the $\log \bar{w}$ - $\log N$ plane, while White (1980) showed that the individual self-thinning lines of 27 different species are closely grouped around a common linear trend of slope -1.5 (see Chapter 7). These results are considered evidence that self-thinning rule applies over a wide range of plant types and growth forms (Gorham 1979, White 1980, 1981, Westoby 1981, Malmberg and Smith 1982, Hutchings 1983). Even some types of plant populations that do not trace straight trajectories in the $\log \bar{w}$ - $\log N$ plane, such as the shoots of clonal perennial species, still seem to be constrained below the ultimate thinning line described by Gorham's study (Hutchings 1979).

<u>Effects of Environmental Factors on Self-thinning Lines</u>

Ecologists have examined the effects of the availability of essential resources, such as light and mineral nutrients, on self-thinning. Plants grown at low levels of illumination thin faster (Harper 1977) and reach maximum biomass levels sooner (Hutchings and Budd 1981b) than populations grown with higher illumination. Decreased illumination also lowers the intercept of

the self-thinning line (White 1981, Hutchings and Budd 1981a, 1981b, Westoby and Howell 1981), possibly due to a decrease in the density of plant matter per unit of occupied space (Lonsdale and Watkinson 1982, 1983a) or to more rapid mortality among shorter plants (Hutchings 1983). The thinning slope is not affected by mild reductions in illumination, but changes from the typical value of -3/2 to -1 under severely lowered light treatments have been observed (White and Harper 1970, Kays and Harper 1974, Harper 1977, Furnas 1981). However, other experiments with equally severe light reductions report no change in the thinning slope (Westoby and Howell 1981, Hutchings and Budd 1981b). Possible reasons for these ambiguous results are considered by Westoby and Howell (1982) and Lonsdale and Watkinson (1982). Since the level of illumination affects the position of the thinning line, light has been implicated as the limiting factor whose availability controls the rate of self-thinning (Kays and Harper 1974, Harper 1977, Westoby and Howell 1982, Lonsdale and Watkinson 1982).

High levels of mineral nutrients increase growth and mortality rates and the rate at which populations approach and follow the self-thinning line. The slope and position of that line are insensitive to difference in fertility (Yoda et al. 1963, White and Harper 1970, Harper 1977, White 1981, Westoby 1981, Hutchings and Budd 1981a); however, there is some evidence against this established view. White (1981) mentioned unpublished data which suggests that fertilizer treatments in forest stands may systematically alter log K, while Hara (1984) showed that forestry

yield tables can indicate different thinning lines for stands grown on sites of different quality. Furnas (1981) saw an effect of soil fertility on thinning line position in his own experiments and in those of Yoda et al., which have interpreted as lacking a fertility effect (Yoda et al. 1963, White and Harper 1970).

Explanations for the Self-thinning Rule

Yoda et al. (1963) derived a simple, geometric explanation of the self-thinning from two assumptions: (1) plants of a given species are always geometrically similar regardless of habitat, size, or age; and (2) mortality occurs only when the total coverage of a plant stand exceeds the available area then acts to maintain 100% cover. The first assumption allows the ground area, a, covered by a plant to be be expressed mathematically as a power function of plant weight, a $\propto w^{2/3}$, while the second assumption implies that the average area covered is inversely proportional to density, that is, $\bar{a} \propto 1/N$. Combining these two equations and adding a constant of proportionality, K, gives the thinning rule equation $\bar{w} = K N^{-3/2}$. Starting from the Clark and Evans (1954) equation for nearest neighbor distance, White and Harper (1970) developed an alternative derivation that renders the second assumption unnecessary, but still implicitly requires the first.

The assumption that plant shape is invariant is not tenable, so these derivations of the thinning rule are unsatisfactory (White 1981, Furnas 1981). Miyanishi et al. (1979) attempted to reconcile these simple geometric models with the fact of varying plant shapes

in their generalized self-thinning law, which states that the power of the thinning equation depends on the proportionality between plant weight and ground area covered. Their hypothesis can be stated mathematically by setting the area covered proportional to w^{2p} , where p can deviate from 1/3 to represent changes in shape with increasing size (allometric growth). With this modification, the Yoda logic gives $\overline{w} = K N^{-1/(2p)}$ and the thinning slope is $\gamma = -1/(2p)$, which equals -3/2 only if shape is truly invariant (isometric growth, p = 1/3). Westoby (1976) used similar logic to predict that the thinning slope should be -1 in the special case of plants that grow radially, but not in height.

The implication of these modified geometric arguments that the self-thinning slope should be dependent on plant allometry has not been supported by experimental tests. Westoby's (1976) experiment with plants that grow only radially gave a thinning slope near -3/2, not the expected value of -1; however, White (1981) discredited this result, claiming that that the species used really does grow in height. Mohler et al. (1978) use allometric data for two tree species to predict thinning slopes of -2.17 and -1.85, which disagree with the idealized value of -1.5 and with the respective measured slopes of -1.21 and -1.46. In the most extensive review of the allometric theory to date, White (1981) applied an allometric model to available data for trees and predicted thinning slopes between -2.05 and -0.78. His discussion of this result implies that the allometric model is faulty in predicting of thinning slopes outside the range -1.8 to -1.3 that he considers acceptably close to

-1.5. White concluded that any theory predicting thinning slopes significantly different from -1.5 is not useful or realistic.

Several non-allometric explanations of explanations of the self-thinning rule have also been proposed. Westoby (1977) hypothesized that self-thinning is related to leaf area rather than to plant weight, but this theory has been discredited (White 1977, Gorham 1979, Hutchings and Budd 1981b). Mohler et al. (1978) suggested that the value of -3/2 is maintained by a mutual adjustment of plant allometry and stand structure during self-thinning, while Furnas (1981) proposed that the -3/2 value is determined by the fact that limiting resources for plant growth are distributed in a three-dimensional volume. Jones (1982) hypothesized that the -3/2 thinning relationship derives from growth and mortality acting to maintain a constant total plot metabolic rate.

Recently, mathematical models have been used to develop even more theories. Pickard (1983) presented three different models deriving the -3/2 value from a mixture of allometric theory and physiological considerations, such as the fraction of photysynthate allocated to biomass increase, the amount of structural and vascular overhead incurred by spatial extension, and the rise in proportionate maintenance costs with increasing weight.

Charles-Edwards (1984) combined his basic hypothesis that each plant requires a minimum flux of assimilate to grow and persist with additional assumptions about the mathematical representation of growth and mortality to produce an explanation of self-thinning.

Perry (1984) derived the self-thinning curve from a physiological model in which the relationship between leaf area and weight, the decrease in photosynthetic efficiency with crowding, maximum plant size, and age are all important factors that must obey certain mutual constraints if the model is to give thinning slopes and intercepts within the ranges that have been actually observed.

In light of the apparent failure of the allometric theory and the paucity of experimental tests of other theories, no satisfactory explanation of self-thinning rule has yet emerged (White 1980, 1981, Westoby 1981, Hutchings and Budd 1981a, Hutchings 1983). It is a "crude statement of constraint whose underlying rationale remains elusive" (Harper as quoted in Hutchings 1983).

Interpretation of the Self-thinning Constant K

Plant ecologists are also interested in interpreting the constant K and its observed range of variation. K has been presented as a species constant invariant to changes in all environmental conditions except the level of illumination (Hickman 1979, Hozumi 1980, White 1981, Hutchings 1983). Many authors regard K as a parameter related to plant architecture (Harper 1977, Gorham 1979, Hutchings and Budd 1981a, Lonsdale and Watkinson 1983a), but some have proposed that K is insensitive to plant morphology (Westoby 1976, Furnas 1981). White (1981) suggested that K is a rough approximation of the density of biomass in the volume of space occupied by plants and can be considered as a weight to volume conversion, but Lonsdale and Watkinson (1983a) provided evidence

against this hypothesis. Lonsdale and Watkinson (1983a) concluded that plant geometry, particularly leaf shape and disposition, do influence thinning intercepts. Harper (1977) speculated the pyramidally shaped trees have higher intercepts than round crowned trees. Westoby and Howell (1981) and Lonsdale and Watkinson (1983a) have hypothesized that shade tolerant plants should have higher thinning intercepts than intolerant plants. Understanding of K is in a similar status as understanding of the -3/2 power: no general theory explaining the variations in K has yet been developed (Hutchings and Budd 1981a).

CHAPTER 2

SIMPLE MODELS OF SELF-THINNING

Introduction

Most proposed explanations of the self-thinning rule have been based on simple geometric models of the way plants occupy the growing surface (Yoda et al. 1963, White and Harper 1970, Westoby 1976, Miyanishi et al. 1979, Mohler et al. 1978, White 1981), but these models suffer from two major limitations. Since they are not related to time dynamics, they can not provide an interpretation of the thinning line as a time trajectory (Hozumi 1977, White 1981). Also, they can not explain deviations from the self-thinning rule, such as the alteration of thinning slopes in deep shade (Westoby 1977).

Two dynamic models are developed here to remedy these deficiencies. A basic model, in which growth is constrained only by the limited availability of growing space, gives an asymptotic linear trajectory in the log B-log N plane. The slope and intercept of the model self-thinning line are related to the parameters and assumptions of the model to develop biological interpretations for the slope and intercept. The second model incorporates two additional growth constraints: a upper limit on the size of individuals (Harper and White 1971, Peet and Christensen 1980) and a maximum total population biomass or carrying capacity (Peet and Christensen 1980, Hutchings and Budd 1981a, Lonsdale and Watkinson

1983b, Watkinson 1984). The general behavior of this model is related to observed phases of population growth (White 1980), and the effects of heavy shading on trajectories in the log B-log N plane are considered.

Model Formulation

Basic Model for the Spatial Constraint

The first step in formulating a dynamic model of self-thinning is selecting a set of state variables to represent the plant population at any time t. Animal populations have been successfully modeled with a single state variable, such as the total number of animals, because animals are relatively uniform in size and simple counts provide rough estimates of total biomass, growth rates, and productivity (Harper 1977). However, similar-aged plants can vary up to 50,000-fold in size and reproductive output, so measurements of both numbers and biomass are essential for understanding plant populations (White and Harper 1970, Harper 1977, White 1980, Westoby 1981). Average weight has been the measure of population biomass in most self-thinning analyses, but this popular choice entails some serious statistical difficulties that can be avoided if total biomass is used (see Chapter 4). Accordingly, the models developed here employ two state variables, plant density and stand biomass, as a minimum reasonable representation of a plant population. Density, N(t), and biomass, B(t), are measured in individuals per unit area and weight per unit area, respectively.

In an even-aged population with no recruitment, plant density and stand biomass change only through growth and mortality. If the population is sufficiently large, these processes can be modeled by two differential equations,

$$\frac{dN}{dt} = -M(N,B) \tag{2.1}$$

and

$$\frac{dB}{dt} = G(N,B) , \qquad (2.2)$$

where M and G are functions for the rates of mortality and growth when the population state is (N,B). The simplest choices for these functions would apply to a young population of widely spaced plants. In this case plants would not interfere with one another, growth would be approximately exponential, and mortality would be zero (ignoring density-independent causes of mortality). Equations 2.1 and 2.2 would become

$$\frac{dN}{dt} = 0 ag{2.3}$$

and

$$\frac{dB}{dt} = g_0 B, \qquad (2.4)$$

where \mathbf{g}_0 is the exponential growth constant in units of time⁻¹.

The deleterious effects of crowding can be represented in this model by assuming that competition reduces the growth rate and increases the mortality rate. The reduction in growth rate can be modeled by multiplying the exponential growth rate by a function $G_r(N,B) \leq 1 \text{ that decreases as either B or N increases, that is,}$

$$\frac{dB}{dt} = g_0 B G_r(N,B) . \qquad (2.5)$$

To specify a similar function for the increase in mortality with crowding, assume that populations undergoing the same level of crowding stress have the same level of per capita mortality, then define $M(N,B) = m \ M_r(N,B)$ with m constant. The relative mortality rate is, then,

$$\frac{dN}{N dt} = -m M_{r}(N,B) , \qquad (2.6)$$

which is constant for any given level of crowding stress M_r . M_r is near zero in a widely spaced population and increases as either B or N increases.

Further model development requires some mathematical representation of crowding. A reasonable assumption is that crowding depends on the amount of space actually occupied by the population relative to the total available space. Assume that the

area, a, occupied by an individual is related to its weight, w, by a power function

$$a = c_1 w^{2p}$$
, (2.7)

with $0 \le p \le 0.5$. Plants that grow only upwards are represented by p=0, while p=0.5 gives pure radial growth. The special case of isometric growth where shape does not vary with size is given by p=1/3. The constant c_1 is inversely related to the density of biomass per unit of occupied space. This constant also depends on initial plant shape, with initially shorter but fatter plants having higher values than taller, thinner ones. Now assume a similar relationship between the average area occupied and the average plant weight

$$\bar{a} = c_2 c_1 \bar{w}^{2p}$$
, (2.8)

with constant c_2 correcting for any systematic differences between the \bar{a} - \bar{w} relationship and the a-w relationship for individuals. The total area, A, occupied by N individuals is proportional to the product of N and the average area occupied,

$$A = c_3 N \bar{a} = c_3 c_2 c_1 N \bar{w}^{2p} . \qquad (2.9)$$

The constant $c_3 > 0$ is related to the allowable overlap between neighboring plants. If overlap is extensive, c_3 is less than one and the total area occupied is less that the product of the number

of plants and the average area occupied. If plants touch but do not overlap, c_3 is near one and A approximately equals N \bar{a} . A shade-intolerant population would have a higher value of c_3 than a shade-tolerant one. Since \bar{w} = B/N, the total area covered can also be expressed as

$$A = c_3 c_2 c_1 N^{1-2p} B^{2p} . (2.10)$$

Now divide this expression for the total area occupied by the available area, which is simply one because density is already scaled to individuals per unit area. This gives a general crowding index, C(N,B):

$$C(N,B) = f N^{1-2p} B^{2p}$$
, (2.11)

with $f = c_3 c_2 c_1$.

The constant, f, in this crowding function subsumes several factors, including a weight to volume conversion (the density of biomass in occupied space) and information on initial plant shape from constant c_1 of equation 2.7, and a correction from c_2 in equation 2.8 for any systematic differences between the \bar{a} - \bar{w} relationship and the a-w relationship for individuals. Also, f includes information from constant c_3 on how much overlap between the zones of influence of plants is permissible.

The crowding index, C(N,B) can now be used to further specify the functions G_r and M_r by defining

$$M_{r}(N,B) = C(N,B) \tag{2.12}$$

and

$$G_{r}(N,B) = 1 - C(N,B)$$
 (2.13)

The differential equation model now becomes

$$\frac{dN}{dt} = -m N C(N,B) = -m f N^{1-2p} B^{2p}$$
 (2.14)

and

$$\frac{dB}{dt} = g_0 B [1 - C(N,B)] = g_0 B [1 - f N^{1-2p} B^{2p}]. \qquad (2.15)$$

Two additional parameters can be used to adjust how rapidly the growth and mortality rates respond to changes in crowding. The linear equations 2.12 and 2.13 are special cases of more general power functions of C,

$$M_{r}(N,B) = C(N,B)^{\theta}$$
 (2.16)

and

$$G_r(N,B) = 1 - C(N,B)^{\theta_3},$$
 (2.17)

where both θ_1 and θ_3 are greater than zero (Figure 2.1). The use of such power functions in plant growth models is discussed in Barnes (1977). With these modifications, equations 2.14 and 2.15 become

$$\frac{dN}{dt} = -m N (f N^{1-2p} B^{2p})^{\theta} 1$$
 (2.18)




Figure 2.1. Power functions of the crowding index. (a) plots the function $M_r = C^{\theta\,l}$ against C for the indicated values of θ_l while (b) plots $G_r = 1 - C^{\theta\,3}$. The powers θ_l and θ_3 adjust how suddenly each function changes near the critical value C = 1.

and

$$\frac{dB}{dt} = g_0 B [1 - (f N^{1-2p} B^{2p})^{\theta}] . \qquad (2.19)$$

Parameters θ_1 and θ_3 allow additional flexibility in representing the plant population. Biologically, they could be related to adaptability to crowding (plasticity) or to initial planting arrangement. Regardless of the exact values of θ_1 and θ_3 , some growth reduction and mortality increase will occur even at low levels of crowding. This is reasonable for natural populations because the random distribution of seedlings places some plants unusually close to their neighbors and competition begins well before the total plot surface is used.

Enhanced Model with Additional Constraints

The basic model can be modified to incorporate other constraints on plant growth. Here, limitations on individual plant weight and total population biomass are added to the basic spatial constraint. It is assumed that approach toward any of the three constraints reduces the population's growth rate, but only the spatial constraint and the carrying capacity affect the mortality rate. Also, the deleterious effects of three constraints are assumed to be additive. The augmented model is

$$\frac{dN}{dt} = -m N \left[(f N^{1-2p} B^{2p})^{\theta} \right] + (\frac{B}{B_{max}})^{\theta} 2$$
 (2.20)

and

$$\frac{dB}{dt} = g_0 B \left[1 - (f N^{1-2p} B^{2p})^{\theta 3} - (\frac{B}{B_{max}})^{\theta 4} - (\frac{B}{N w_{max}})^{\theta 5}\right], \quad (2.21)$$

where B_{max} is the carrying capacity, w_{max} is the maximum individual weight, and θ_1 through θ_5 control how abruptly the rates respond to changes in the level of each constrained quantity.

Model Analysis and Results

Basic Model

Although the basic model of equations 2.18 and 2.19 is derived from very simple assumptions, it can not be solved to give explicit equations for N(t) and B(t). However, it is amenable to isocline analysis, a technique discussed in many introductory ecology texts. This method is applied here to the simple case of $\theta_1 = \theta_3 = 1$ represented by equations 2.14 and 2.15. First, note that the growth rate of population biomass is zero when the right hand side (RHS) of equation 2.15 is zero, that is,

$$g_0 B [1 - f B^{2p} N^{1-2p}] = 0.$$
 (2.22)

On log transformation and algebraic manipulation, this yields

$$\log B = \left(-\frac{1}{2p} + 1\right) \log N - \frac{1}{2p} \log f$$
, (2.23)

the equation of a straight line of slope $\beta = -1/(2p) + 1$ when

ordinate log B is plotted against abscissa log N. This slope must be negative or zero because $0 \le p \le 0.5$, so that $-1/(2p) \le -1$.

Restrictions on the path of the model population in the log B-log N plane (the self-thinning curve) can be deduced from this zero isocline, which divides the log B-log N plane into two regions. Below the isocline, population biomass is increasing and trajectories move upward, while above the isocline, biomass is decreasing and trajectories move downward. Since dN/dt is strictly negative, the force of mortality is always decreasing density and moving the population leftward in the plane. Because the isocline slants up toward the left of the plane, it must intercept the downward and leftward path of any population starting above the isocline. Such a population steadily approaches the isocline, then crosses it to enter the lower half of log B-log N plane. A population below the zero isocline must move upward (dB/dt > 0), but can not grow through the isocline because dB/dt is zero along that line. Three possibilities remain: the population trajectory could approach the zero isocline asymptotically, remain a constant distance from the isocline, or move leftward faster than upward, thus moving away from the isocline.

The potential ambiguity in the behavior of the model population below the zero isocline can be eliminated as follows. The instantaneous direction, β_t , of the population's path at any point in the log B-log N plane is given by

$$\frac{d \log B}{d \log N} = \frac{dB / (B dt)}{dN / (N dt)} = \beta_t, \qquad (2.24)$$

a relationship that can be used to identify points of the log B-log N plane where the slope of the population trajectory is less than or equal to any given value Ψ , that is, where $\beta_t \leq \Psi$. Combining this inequality with equations 2.18 and 2.19 in the ratio of equation 2.24 yields

$$\frac{g_0 \left[1 - f N^{1-2p} B^{2p}\right]}{-m f N^{1-2p} B^{2p}} \leq \Psi. \tag{2.25}$$

Algebraic manipulation and log transformation of this expression gives a relationship for log B in terms of log N:

$$\log B < (-\frac{1}{2p} + 1) \log N - \frac{1}{2p} \log f + \frac{1}{2p} \log (\frac{g_0}{g_0 - m\Psi})$$
 (2.26)

The first two terms of the right hand side (RHS) of this equation simply give equation 2.23 for the zero isocline. The third term is a constant added to the intercept of the zero isocline since g_0 , m, and Ψ are constants. The equality in 2.26, which is the locus of points where trajectories take a given slope Ψ , defines a straight line parallel to the zero isocline. In fact, the zero isocline equation 2.23 is the special case of the more general equation 2.26 with $\Psi=0$.

This general relationship can be used to find regions of the plane below the zero isocline where a population's instantaneous trajectory is steeper (more negative) than the slope of the zero isocline, so that the population is moving closer to the isocline.

Substituting Ψ = β = -1/(2p) + 1 (the slope of the zero isocline) in equation 2.26 gives

$$\log B \leq \left(-\frac{1}{2p} + 1\right) \log N - \frac{1}{2p} \log f + \frac{1}{2p} \log \left(\frac{g_0}{g_0 - m[-1/(2p) + 1]}\right) \cdot (2.27)$$

Since $g_0 > 0$ and m > 0 while $0 \le p \le 0.5$, the third term on the RHS is negative and equality defines a straight line parallel to but below the zero isocline. This lower line is asymptotically approached by all populations. Below this asymptotic trajectory, a population's path is steeper than the asymptotic trajectory, while the path of a population above the asymptotic trajectory is less steep than the asymptotic path. In both cases, the population's trajectory must continuously move closer to the asymptotic trajectory. Thus, the asymptotic trajectory has both attributes of the self-thinning line: populations approach it from any starting point in the log B-log N plane, and it is a boundary between allowable and unallowable biomass-density combinations. This second conclusion follows because populations can never grow through the asymptotic trajectory from below, and even if the model was started above the thinning line, mortality and negative growth would drive the population trajectory toward the thinning line and out of the region of the plane representing untenable biomass-density states.

Equation 2.27 also gives the slope and intercept of the self-thinning line in terms of the model parameters. The slope is -1/(2p) + 1, a function of the single parameter p which relates area

occupied to plant weight. Only if p = 1/3 is the model thinning slope equal to the value of β = -1/2 predicted by the self-thinning rule. The thinning line intercept is given by the last two terms of equation 2.27

$$\log K = -\frac{1}{2p} \log f + \frac{1}{2p} \log(\frac{g_0}{g_0 - m[-1/(2p)+1]}).$$
 (2.28)

The value of log K depends on all the model parameters, but the first term of the RHS depends only on p and f. If the second term is small relative to the first, then log K is mainly determined by these two parameters. To determine when this condition is satisfied, define a new quantity, $\Delta \log K$, as the second term of equation 2.28

$$\Delta \log K = \frac{1}{2p} \log(\frac{g_0}{g_0 - m[-1/(2p)+1]}),$$
 (2.29)

which is the contribution to the intercept of the self-thinning line of the second term of equation 2.28 and is also the vertical distance between the asymptotic population trajectory and the isocline of zero biomass increase. Since \mathbf{g}_0 and \mathbf{m} are both constants, \mathbf{m} can be expressed as the product of \mathbf{g}_0 and a constant \mathbf{w} , that is, $\mathbf{m} = \mathbf{w} \ \mathbf{g}_0$. With this substitution, equation 2.29 gives $\Delta \log K$ in terms of \mathbf{p} and \mathbf{w}

$$\Delta \log K = -\frac{1}{2p} \log(1 + \omega \left[\frac{1}{2p} - 1\right])$$
 (2.30)

This expression is negative or zero, so the thinning line is always below the zero isocline or equal to it. Figure 2.2 plots Δ log K against log ω for several choices of p and shows that most combinations of ω and p give Δ log K \leq 1. Only when p is small (p < 1/3) or ω is large (ω > 10) does Δ log K exceed one, and Δ log K is much less than one for most reasonable parameter values. Thus, log K values of four or more are primarily determined by f and p as specified by the first term of equation 2.28.

Several features of this analysis are illustrated in Figure 2.3a, in which equations 2.14 and 2.15 are fitted to a Pinus strobus plantation remeasured nine times between 12 and 51 years after planting (lot 2B, Spurr et al. 1957). By repeated trials, the parameter values $g_0 = 0.5$, m = 0.0475, p = 0.29, and f = 0.006were found to give a visually good fit to the log B-log N data when the model was started at the first data point and solved numerically using the LSODE differential equation solver (Hindmarsh 1980). The resulting solution demonstrates that the model can represent actual population data quite well. Isopleths of equation 2.26 for four values of Ψ are also shown, including the zero isocline of biomass growth (log B = $-0.724 \log N + 3.83$) given by $\Psi = 0$, and the asymptotic self-thinning line (log B = -0.724 log N + 3.78) given by $\omega = \beta = -1/(2p) + 1 = -0.724$. The self-thinning line is -0.50 log units below the zero isocline, that is, $\triangle \log K = -0.05$. Figure 2.3b shows how model solutions for different initial states converge on the asymptotic self-thinning line.




Figure 2.2. Vertical distance between the zero isocline of biomass growth and the self-thinning line. This distance, $\Delta \log K$, is plotted against the ratio, ω , of the mortality constant to the growth constant (ω = m/g_0) for the indicated values of the parameter p. $\Delta \log K$ is the contribution of g_0 and m to the self-thinning intercept and is less than one when m \leq g_0.




Figure 2.3. Basic model fitted to data from a <u>Pinus strobus</u> plantation. Model parameters are g_0 = 0.5, m = 0.0475, f = 0.006, and p = 0.29. In (a) data points (Spurr et al. 1957) are circled, the solid line is the solution of the model starting from the first point, the dashed line is the dB/dt = 0 isocline log B = -0.724 log N + 3.83, and the dotted lines are loci where trajectories have the indicated instantaneous slopes. The line marked "-0.724" is the model self-thinning line log B = -0.724 log N + 3.78. In (b), the dotted lines are model solutions starting from different initial conditions and converging on the asymptotic thinning line (solid).

Enhanced Model

Analysis of the model with added constraints on total biomass and individual weight is more difficult, because the equation for the isocline of zero biomass growth,

$$[f N^{1-2p} B^{2p}]^{\theta_3} + [\frac{B}{B_{max}}]^{\theta_4} + [\frac{B}{N w_{max}}]^{\theta_5} = 1, \quad (2.31)$$

is too complex to solve for log B in terms of log N. However, the effect of each constraint can be considered independently by removing two of the three terms on the RHS. This analysis gives three straight lines in the log B-log N plane. The equation associated with the first term on the RHS is simply equation 2.23 for the zero isocline of the basic model, while the remaining two terms give

$$\log B = \log B_{\text{max}} \tag{2.32}$$

and

$$\log B = \log N + \log w_{\text{max}}. \tag{2.33}$$

The three terms are additive, so the actual zero isocline lies beneath the lowest of the three straight lines in the log B-log N plane, and the lowest line dominates the position and slope of the true isocline. Where two of the lines cross, the actual zero isocline undergoes a gradual transition from one slope to another. The abruptness of the transition is controlled by the parameters θ_1 , θ_2 , and θ_3 with higher values giving more abrupt transitions.

These behaviors of the enhanced model are illustrated in Figure 2.4, which was constructed with the parameters estimated for the Pinus strobus plantation, combined with hypothetical values of $10^4~\rm g/m^2$ and $10^8~\rm g$, respectively, for the new parameters $B_{\rm max}$ and $W_{\rm max}$. Parameters θ_1 through θ_5 were all set to two. The equation of the true zero isocline 2.31 was found numerically using the ZEROIN computer subroutine (Forsythe et al. 1977) to solve

$$[f N^{1-2p} B^{2p}]^{\theta_1} + [\frac{B}{B_{max}}]^{\theta_2} + [\frac{B}{N w_{max}}]^{\theta_3} - 1 = 0$$
 (2.35)

for B at different values of N. The exact model solution for one set of initial conditions was calculated numerically using LSODE.

Discussion

Basic Model

Analysis of the basic model has shown that the self-thinning rule can be derived in a dynamic model as a consequence of the limited availability of growing area. The power relationship between biomass and density (the straight line relationship between log B and log N) follows directly from an assumed power relationship between the area occupied by a plant and plant weight. This assumption is reasonable because power functions between plant measurements have been repeatedly demonstrated in many disciplines




Figure 2.4. Analysis of a model with three constraints on population growth. Parameters g_0 , m, f, and p are as in Figure 2.2, while $B_{max}=10^5$, $w_{max}=10^8$, and θ_1 through θ_5 all equal 2. Dotted lines are dB/dt=0 isoclines for each constraint considered independently. The maximum individual weight, the carrying capacity, and the spatial constraint are labelled " w_{max} ", " B_{max} ", and "f,p", respectively. The dashed line is the true isocline for all three constraints and the solid line is a model solution for 450 years. Regions I, II, and III are, respectively, portions of the plane where model behavior is dominated by the spatial constraint, the carrying capacity, and the maximum individual weight.

including forestry (Reinke 1933, Curtis 1971) and plant ecology (Whittaker and Woodwell 1968), and such relationships provide the basis for widely used methods of nondestructive sampling of plant populations (see references in Hutchings 1975). When traced to this rather commonplace origin, the power equation form of the self-thinning rule is unremarkable.

The slope of the model thinning line is determined only by the power, p, of the area-weight relationship and equals the classic thinning rule slope of β = -1/2 only if p = 1/3. Plant growth is typically not isometric (Mohler et al. 1978, Furnas 1981, White 1981) so p is not generally 1/3, and the model predicts that thinning slopes should vary systematically with plant allometry. This hypothesis has not been supported in previous reviews and experimental tests (Westoby 1976, Mohler et al. 1978, White 1981), so the causal mechanisms formalized in the model developed here have been discredited as possible explanations for the self-thinning rule. However, some new analyses presented in Chapters 5 and 6 show that measured self-thinning slopes do vary significantly from the idealized value of the thinning rule, and that these deviations can be correlated with differences in plant allometry, thus verifying a major prediction of the model developed here.

Even for this very simple model, the exact value of the self-thinning constant, K, depends on all the model parameters and can not be interpreted as a function of a single measurement, such as biomass density in space (as discussed in White 1981 and Lonsdale and Watkinson 1983a). Biomass density in space is a component of

the parameter f, but its relationship to log K is confounded by other components of f, such as initial plant shape and the degree of tolerable overlap between plants, and by the dependence of log K on the allometric power p. Since thinning slopes do vary with plant allometry (Chapter 6), direct comparisons of thinning intercepts of experimental thinning lines are not clearly interpretable because the comparison of the constants among power relationships is not meaningful unless the relationships have the same power (White and Gould 1965). Experimental interpretation of K will, then, require a careful statistical analysis relating K to several factors, including plant allometry, initial plant shape, the density of biomass per unit of occupied space, and some measure of allowable overlap between plants, such as shade tolerance.

The position of the model self-thinning line is also affected by the rate constants \mathbf{g}_0 and \mathbf{m} , but these effects are relatively small over a large range of reasonable parameter values. The primary effect in the log B-log N plane of \mathbf{g}_0 and \mathbf{m} is to determine the rate and direction of approach toward the thinning line. By concentrating on the thinning line, these rate dynamics are deemphasized in favor of a focus on the limitations imposed by available growing space and plant geometry.

The model also predicts that the parameters of the self-thinning line are not species constants invariant to changes in all environmental factors except illumination (as proposed by Yoda et al. 1963, Hickman 1979, Hozumi 1980, White 1981, and Hutchings 1983). Any factor that could affect the density of biomass in

occupied space or the degree of allowable overlap between plants would also affect the thinning constant K, while environmental factors affecting plant allometry would also change both K and the self-thinning slope. The remarkable abilities of plants to vary their sizes, shapes, canopy densities, etc. in response to environmental and competitive factors are well-documented (see references in Harper 1977), and allometric relationships for a species also vary significantly among sites (Hutchings 1975). Peet and Christensen (1980) reported that the thinning characteristics of a pine stand could be permanently altered if the initial density is very high and speculated that some aspects of tree geometry were fixed by the initial growing conditions. Such permanent effects of initial density would be another cause for variation in thinning line parameters among populations of a species. The prediction that thinning line parameters are not species constants is tested in Chapters 4 though 6.

Enhanced Model

The model with additional constraints on total biomass and individual plant weight can explain most general features of population dynamics in the log B-log N plane. The four phases of population growth that were discussed in Chapter 1 and diagrammed in Figure 1.1, page 9 are reproduced by the model in Figure 2.4, with the sharp corners of idealized behavior replaced by gradual transitions from one growth phase to another. The four phases are explained by the model as the sequential operation of three

constraints on population growth: the spatial constraint, the carrying capacity, and the maximum individual size. The transition from adherence to the self-thinning rule to constant biomass at the carrying capacity, shown by movement from region I to region II of Figure 2.4, has been discussed by White and Harper (1970) and Hutchings and Budd (1981a). Experimental evidence of this transition has been reported by Schlesinger and Gill (1978), Lonsdale and Watkinson (1983b), Watkinson (1984), and Peet and Christensen (1980). The last of these papers also reported that thinning trajectories eventually become even less steep than the constant biomass line, as shown in the transition from region II to region III in Figure 2.4.

The modified model can also explain the ambiguous results of experiments comparing the self-thinning lines of deeply shaded populations to those of populations grown under better illumination. Some investigators have concluded that reduced illumination lowers thinning lines and shifts the slope in the log B-log N plane from -1/2 to 0 (White and Harper 1970, Kays and Harper 1974, Lonsdale and Watkinson 1982), while others have observed no change in slope when illumination is decreased (Westoby and Howell 1981, Hutchings and Budd 1981b).

The enhanced model (equations 2.20 and 2.21) is used here to investigate the effects of decreased illumination on the log B-log N trajectory, but the constraint on individual weight is eliminated to simplify the analysis. It is unlikely that plants would reach their greatest potential size during short experiments under reduced

illumination, so this constraint would be irrelevant. A reduction in illumination could shift the positions of the two remaining constraining lines in the log B-log N plane. As a first approximation, a certain percentage reduction in illumination might be expected to reduce the carrying capacity by the same percentage. It is more difficult to estimate the effect of a light reduction on the spatial constraint line. Lower illumination can stimulate plants to emphasize height growth rather than radial growth (Harper 1977), so that the allometric power p relating area covered to weight might decrease, resulting in a steeper thinning line. However, illumination changes may also alter the density of biomass in occupied space and change the thinning intercept (Lonsdale and Watkinson 1982). Since species differ widely in their shade tolerances, meristem placements, and other important physiological and morphological factors, the actual effect on the spatial constraint line should vary among species. Regardless of these particulars, the carrying capacity will become the dominant constraint over a wider range of plant densities under reduced illumination as long as the spatial constraint line does not drop too drastically.

The consequences for the observed log B-log N trajectory of a drop in total supportable biomass with reduced illumination are shown in Figure 2.5. In this particular example, the carrying capacity drops by one log unit when illumination is reduced by 90%, but the spatial constraint line is not affected. With the light reduction, the lowered carrying capacity dominates model behavior




Figure 2.5. Effect of a reduction in illumination on population trajectories in a two-constraint model. A hypothetical 90% drop in illumination changes the carrying capacity from 10^5 to 10^4 (dotted lines labelled "B_{max}") but does not affect the spatial constraint line (dotted line labelled "f,p"). Other model parameters are as in Figure 2.4. The dashed line is the true zero isocline for both constraints. Model solutions (solid lines) starting from the same initial conditions are shown for the two values of B_{max}.

over a wider range of densities when compared to model behavior at the higher light level. This is clear when the complete trajectories are examined over four orders of magnitude of density; however, the response revealed to an experimenter by measurements taken over more limited densities would depend on the range observed. An investigator sampling over 0.5 < log N < 1.5 from populations behaving as shown in Figure 2.5 would see virtually no difference between the high light and reduced light trajectories, but measurements over $-1.0 < \log N < -0.5$ would reveal a major drop in both the intercept and the slope of a straight line through the data. Thus, the ambiguous and seemingly contradictory results from experiments on reduced illumination are explainable by this model. Confusion has resulted because the various studies have used different plants observed over different densities and so focused on different regions of the overall behavior revealed in Figure 2.5. A similar theory and some supporting evidence were presented by Lonsdale and Watkinson (1982).

CHAPTER 3

SIMULATION MODEL OF SELF-THINNING

Introduction

The models of Chapter 2 yield predictions about the effects of plant geometry and restricted resources on self-thinning, but some of the underlying assumptions merit further consideration since they may be over-simplifications. These highly aggregated and spatially averaged models do not explicitly consider the sizes and locations of individuals or the interactions between them, but plants compete with immediate neighbors not average population conditions (Schaffer and Leigh 1976, Harper 1977). A simulation model which represents these factors in a more realistic, detailed way is analyzed here to check if its responses to variations in plant allometry, the density of biomass in occupied space, and the degree of tolerable overlap between plants are similar to the responses of the simpler models. As in Chapter 2, the principal objective is to discover which model parameters and processes determine the linearity, slope, and intercept of the self-thinning line, then use these results to provide explanations for observed self-thinning relationships.

Model Formulation

The simulation model is based on simple equations for the growth and geometry of an individual plant. Each plant in the

population is represented by separate copies of these equations, and the entire group is united into a single system by some rules for partitioning resources in areas where neighboring plants interact. It is assumed that the growth of a plant is related to the ground area it covers, since this is the region over which light, water, and mineral resources are obtained (Aikman and Watkinson 1980). The basic equation for the growth in weight, w, of a single plant is

$$\frac{dw}{dt} = g_1 a - b w^q , \qquad (3.1)$$

where g_1 represents the maximum assimilation rate (in $g/m^2/unit$ time) sustainable per unit of ground area, and a is the area of ground covered by the plant, also called the zone of influence (ZOI--Bella 1971). The second term on the right hand side (RHS) of equation 3.1 represents the cost of maintaining existing plant biomass, which is assumed to be a power function of weight specified by the constants b and q.

To determine the ground area covered by an individual, assume that plants, on the average, occupy cylindrical volumes of space called the volume of influence (VOI). Plant weight can then be related to the dimensions of this cylinder by

$$w = vd = \pi R^2 hd$$
, (3.2)

where v, R, and h are the volume, radius, and height of the VOI and d is the average density of biomass in that volume. Changes in

plant geometry with plant size can be modeled by assuming that the radius of the VOI is a power function of weight

$$R = c w^p, (3.3)$$

where c and p are constants. As in Chapter 2, values of p are between 0.0 and 0.5, and c depends on initial plant shape and the density of biomass in occupied space. Assuming that the height to radius ratio, τ , is known for plants of some weight $\overline{\mathbf{w}}_0$, equations 3.2 and 3.3 can be combined and solved for c to relate this constant to parameters with more direct biological interpretations

$$c = \left[\frac{\bar{w}_0}{\pi \tau d}\right]^{1/3} \frac{1}{\bar{w}_0^p} = \frac{\bar{w}_0^{(1/3 - p)}}{(\pi \tau d)^{1/3}}.$$
 (3.4)

Thus, c is related to the density of biomass in occupied space, d, and to values of τ and \overline{w}_0 that specify plant shape at a particular weight. Equation 3.3 can be used to develop a formula relating area covered to plant weight

$$a = \pi c^2 w^2 p = c_1 w^2 p$$
 (3.5)

Since $c_1 = \pi c^2$, c_1 can also be related to \bar{w}_0 , p, τ , and d through equation 3.4. Combining equations 3.1 and 3.5 gives a general equation for the growth rate of an isolated individual

$$\frac{dw}{dt} = g_1 c_1 w^{2p} - b w^q . ag{3.6}$$

A plant can not fully exploit the resources within its ZOI if some resources are preempted by neighbors attempting to occupy the same area, so the model partitions the resources associated with a particular section of ground among the plants that are competing for it. This is accomplished by a pairwise comparison of all plants. Since the spatial location of each plant and the radius of a ZOI centered at that point are known, simple trigonometry can be used to determine if two plants overlap. If so, the area of ground surface shared is calculated and a fraction of the shared area is subtracted from the ZOI of each member of the pair to represent the loss of resources to its competitor. Equation 3.6 becomes

$$\frac{dw}{dt}i = g_1 \left[c_1 w_i^{2p} - \sum_{\substack{j=1\\j\neq i}}^{n} a_{loss,ij}\right] - b w^q, \quad (3.7)$$

where $a_{loss,ij}$ is the area lost by plant i to plant j and n_t is the number of plants remaining at time t. The model includes six different algorithms for dividing areas of overlap between two plants. Table 3.1 gives names and formulas for all six. The fractions in Table 3.1 are multiplied by the actual area of overlap to obtain the a_{loss} term of equation 3.6 for each pairwise comparison. Since the area controlled by a plant cannot be negative, a_i (the bracketed term in equation 3.7) is set to zero if the total area lost by a plant exceeds the area of its ZOI.

The direct application of equation 3.7 is precluded by edge effects, which can bias the results if the simulated plot is

Table 3.1. Competition Algorithms in the Simulation Model.

Algorithm Number ^a	Method of Partitioning Contested Areas	Fraction Allo	ocated to Plant j
1	Divided in proportion to weight	w _i + w _j	w _j w _i + w _j
2	Larger plant gets allb	1	0
3	Neither plant gets any	0	0
4	Divided in proportion to height	h _i h _i + h _j	hj hi + hj
5	Divided equally	<u>1</u> 2	<u>1</u>
6	Half divided equally; half in proportion to weight	$\frac{1}{2} + \frac{1}{2} \frac{w_i}{w_i + w_j}$	$\frac{1}{2} + \frac{1}{2} \frac{w_j}{w_i + w_j}$

 $^{\,^{\}mbox{\scriptsize bFr}}_{\mbox{\scriptsize ractions}}$ are given for the case where plant i is larger than plant j.

intended to represent a sample of a larger population. These effects arise because some of the competitors of plants near the plot edge lie off the plot and are not observed. The linear expansion algorithm of Martin et al. (1977) is used to remove this effect. The calculated area lost by each plant i to its competitor j ($a_{loss,ij}$) is multiplied by an expansion factor

$$\varepsilon_{ij} = 2\pi / \eta_{ij}$$
, (3.8)

before being summed into equation 3.7. The quantity η_{ij} , called the sample angle, is calculated for each pair of plants by centering on plant i a circle of radius D_{ij} (the distance between plants i and j). The angle subtending the portion of this circle that lies on the simulated plot is η_{ij} . This angle is 2 π when the entire circle is on the plot and less than 2 π when part of the circle falls outside the plot boundary, so $\varepsilon_{ij} \geq 1$. Multiplication of each observed competitive loss by an expansion factor gives a statistically unbiased estimate of the additional loss to unobserved competitors outside the simulated plot, but this estimate is valid only if the radius of the largest ZOI is less than the plot radius (Martin et al. 1977); therefore, all simulations were halted before this condition was violated.

The persistence of an individual plant requires a favorable balance between the rates of assimilation and respiration. This requirement can be modeled by killing plants whose average relative growth rates over a selected time interval fall below a threshold value, RGR_{min} (Aikman and Watkinson 1980). For the simulations

presented here, the interval was 0.1 time units and the threshold value was 0, that is, all plants not satisfying the inequality

$$RGR_{i} = \frac{1}{0.1} \quad In \quad \frac{w_{i,t}}{w_{i,t-0.1}} \geq 0$$
 (3.9)

were eliminated from the simulation ($w_{i,t}$ is the weight of plant i at time t).

The initial conditions for a simulation were specified by selecting an initial number of plants and a location and weight for each plant. Initial weights for plants were chosen randomly from a normal distribution with mean \bar{w}_0 and standard deviation $s_{\bar{w}_0}$, while plant locations were assigned from a uniform random distribution on a circular plot of specified area. The model was typically started with 200 plants, but 600 was the initial population size when the dynamics of the size distribution were of particular interest and larger sample sizes were needed to obtain accurate estimates of the moments of that distribution.

Model Analysis

The model was analyzed in three steps: (1) verification that the dynamics resemble real self-thinning behavior, (2) estimation of the variations in the self-thinning trajectory strictly due to stochastic model elements, and (3) evaluation of simulation experiments in which a parameter was varied among simulations to determine the effect of that parameter on self-thinning. A reference set of parameters was defined to generate the model

solutions used in steps one and two. The values chosen (Table 3.1) are arbitrary, but within biologically realistic limits. Except for special cases detailed in the results, the model solutions used here were linear over the interval $3.0 \le \log N \le 3.7$, so regression of log B against log N was applied to data within this interval to estimate the slope, $\hat{\beta}$, and intercept, $\hat{\alpha} = \log K$, of a self-thinning line for each simulation. The intercept of each regression line at log N = 3.35, $\hat{\alpha}_{3.35}$, was also calculated to provide a measure of thinning line position near the center of range of data. Coefficients of determination were so high (all $r^2 > 0.97$) that the regression estimates of $\hat{\beta}$ and $\hat{\alpha}$ were identical to those from principal component analysis. PCA is preferable to regression when the two methods give different results (Chapter 4). The time required for the density of individuals to fall from the initial value of 10,000 to 1000 was recorded as a measure of the average rate of self-thinning.

Because the model is complex, a controlled procedure was used to identify model parameters that affect the self-thinning trajectory. In each of six simulation experiments, a parameter was varied across 5-15 simulations and the resulting group of self-thinning trajectories was compared to a control group of ten trajectories generated by the reference parameter set. This control group was analyzed to estimate the ranges of variation in $\hat{\beta}$, $\hat{\alpha}$, and $\hat{\alpha}_{3.35}$ strictly due to stochastic model factors. Values of $\hat{\beta}$, $\hat{\alpha}$, or $\hat{\alpha}_{3.35}$ well outside these ranges in an experimental

simulation indicated that the parameter being studied did alter the self-thinning line beyond the normal limits of stochastic variation and therefore has a role in positioning the self-thinning line. seven parameters that have entries in the "Range of Variation" column of Table 3.2 were investigated in these simulation experiments, but one experiment considered parameters b and q together. For each experimental group, log B-log N plots were used to visually compare the variation among thinning trajectories to the variation among the control trajectories. The mean, range, and coefficient of variation (CV) of $\hat{\beta}$, $\hat{\alpha}$, and $\hat{\alpha}_{3,35}$ were also tabulated for each simulation experiment and compared to the same statistics for the control group. For five of the experiments, Spearman correlation coefficients, r_s , (Sokal and Rohlf 1981) between the experimental parameter and $\hat{\beta}$, $\hat{\alpha}$, and $\hat{\alpha}_{3.35}$ were calculated and tested for statistical significance to determine if these thinning line descriptors varied systematically with the experimental parameter. Although this application of statistical tests to simulation results may initially seem contrived, it is appropriate for a model with stochastic factors. Chance variations among a small number of stochastic simulations can produce meaningless correlations in the same manner as in real data. high correlations were found, further analysis was done to estimate the precise relationship between the slope or intercept and the model parameter.

Table 3.2. Simulation Model Parameters and Their Reference Values.

Symbol	Meaning	Units	Reference Value	Range of Variation	
р	Allometric power relating ZOI radius to weight		0.333	0.27- 0.49	
d	Density of biomass in occupied space	g/m ³	6200	631 - 39810	
	Competition algorithm		1	1-6	
91	Maximum assimilation rate per unit of ground area covered	g/m ² /time	25	10-50	
b	Constant relation metabolic rate to weight	(g/m ²)(1/q) 0.00147	1.47x10 ⁻¹⁰ - 147*	
q	Power relating metabolic rate to weight		2	0.5-3.5*	
Aplot	Plot area	_m 2	0.02	0.000632- 0.0632**	
τ	Height to radius ratio for VOI's of plants of weight $\bar{\mathtt{w}}_0$		1		
RGR _{min}	Minimum survivable relative growth rate		0		
Δt	Time step	time	0.1		
(∆t) _m	Time interval over which RGR's are averaged	time	0.1		
n_0	Initial number of plants		200		
\bar{w}_0	Initial average weight	g	0.0001		
s _{₩0}	Standard deviation of initial weight	g	5x10 ⁻⁵		

 $[\]ensuremath{^{\star}}\ensuremath{\text{b}}$ and q were varied together in a single set of simulations (see text).

 $^{^{\}star\star}A_{\text{plot}}$ was varied to give initial plant densities from 3160 to 316,000 plants/m².

Results

The simulation model mimics several behaviors of real plant populations (Figure 3.1). Initially uncrowded populations trace a nearly vertical path in the log B-log N plane, but eventually bend toward and move along a negatively sloped self-thinning line. Model populations show other behaviors of real plant monocultures, including nearly logistic increase in biomass (Hutchings and Budd 1981a), approximately exponential mortality during self-thinning (Yoda et al. 1963, Harper 1977), and the development of skewed size distributions from initially symmetric distributions (White and Harper 1970, Hutchings and Budd 1981a).

Information on individual simulations is presented in Table 3.3, including the experimental parameter value (if any) and the three thinning line descriptors $\hat{\beta}$, $\hat{\alpha}$, and $\hat{\alpha}_{3.35}$. All the thinning line regressions were based on at least eight points, but most used between 20 and 70. Coefficients of determination, r^2 , were uniformly high ($r^2 \geq 0.97$), confirming that trajectories were well described by a straight line over the range $3.0 \leq \log N \leq 3.7$. The thinning line descriptors are further summarized in Table 3.4, where means, ranges, and coefficients of variation are given for each of six experimental or control groups, along with Spearman correlations with the experimental parameter (if any). Table 3.3 also gives the time required for the density to fall from 10,000 to 1000. Systematic changes in these times within the simulation experiments show that all parameters affected the




Figure 3.1. Typical dynamic behavior of the simulation model. (a) is a log B-log N self-thinning plot, (b) shows the logistic increase in total population biomass, and (c) shows approximately exponential mortality commencing with the onset of thinning around time 20. The reference parameters (Table 3.2) were used.

Table 3.3. Self-thinning Lines for the Simulation Experiments.

			Estimated Thinning Line					
Parameter Par Varied		Time to og N = 3*	Slope I	ntercept ດີ	Intercept at $\log N = 3.35$ $\alpha_{3.35}$			
None (Figure 3.2)		70 55 54 67 59 58 59 59 58 60	-0.69 -0.55 -0.55 -0.65 -0.62 -0.59 -0.72 -0.64 -0.54	4.94 4.48 4.46 4.83 4.73 4.62 5.05 4.79 4.45 4.90	2.64 2.63 2.63 2.67 2.66 2.65 2.62 2.65 2.64 2.66			
p (Figure 3.3)	0.27 0.29 0.31 0.33 0.335 0.37 0.39 0.41 0.43 0.45 0.47	426 203 106 65 8 57 41 27 19 14 11 8 7 6	-0.98 -0.91 -0.69 -0.70 -0.61 -0.57 -0.42 -0.35 -0.28 -0.24 -0.22 -0.09 -0.11	6.69 6.23 5.22 5.02 4.68 4.42 3.71 3.36 2.97 2.70 2.54 2.00 1.97	3.41 3.17 2.91 2.66 2.64 2.50 2.30 2.18 2.02 1.89 1.80 1.68 1.58			
d (Figure 3.4)	631 1000 1585 2512 3981 6310 10000 15850 25120 39810	6 9 15 22 36 60 90 158 251 378	-0.45 -0.55 -0.56 -0.60 -0.61 -0.63 -0.63 -0.55	3.17 3.66 3.90 4.25 4.47 4.70 4.93 5.16 5.11 5.44	1.65 1.83 2.03 2.26 2.45 2.65 2.83 3.06 3.27 3.45			
Competition algorithm (Figure 3.5)	1 2 3 4 5 6	59 36 65 84 >100 44	-0.68 -0.70 -0.64 -0.57 -0.62 -0.74	4.93 4.76 4.60 4.65 4.89 4.99	2.64 2.43 2.47 2.73 2.82 2.52			
91 (Figure 3.6a)	10 15 20 25 30 35 40 45	>100 >100 71 55 52 37 38 36 30	-0.54 -0.71 -0.64 -0.59 -0.64 -0.63 -0.59 -0.71	4.43 5.03 4.79 4.60 4.80 4.76 4.62 5.06 4.79	2.63 2.65 2.64 2.62 2.65 2.63 2.65 2.68 2.65			
q (Figure 3.6b)	0.5 1.0 1.5 2.0 2.5 3.0 3.5	58 64 59 57 61 73	-0.62 -0.64 -0.64 -0.58 -0.67 -0.68	4.71 4.80 4.78 4.59 4.88 4.92	2.64 2.64 2.65 2.64 2.62 2.65			

 $^{^{\}star}\text{The time}$ required for plant density to drop from its initial value of log N = 4.0 to log N = 3.0.

Table 3.4. Thinning Line Statistics for the Simulation Experiments.

Figure	Parameter Varied	Slope ĝ			Intercept â				Intercept at log N = 3.35 $\hat{\alpha}_{3.35}$				
		Mean	Range	CV	rĝ	Mean	Range	CV	rĝ	Mean I	Range	CV	rs
3.2	none	-0.62	0.18	10.2		4.72	0.59	4.5		2.64	0.05	0.6	
3.3	p	-0.48	0.89	61.2	0.99* (<0.0001)	3.96	4.72	39.2	-1.00* (<0.0001)	2.37	1.83	24.4	-1.00* (<0.0001)
3.4	d	-0.58	0.17	9.2	-0.49 (0.15)	4.48	2.28	16.3	0.99* (<0.0001)	2.55	1.80	24.0	-1.00* (<0.0001)
3.5	Competition algorithm	-0.66	0.16	8.9		4.80	0.38	3.3		2.60	0.39	5.9	
3.6a	91	-0.63	0.17	8.9	-0.20 (0.61)	4.76	0.63	4.2	0.25 (0.52)	2.64	0.06	0.6	0.40 (0.28)
3.6b	q	-0.64	0.10	5.7	-0.60 (0.21)	4.78	0.34	2.5	0.60 (0.21)	2.64	0.03	0.4	0.26 (0.62)

^aSpearman correlation of the thinning line descriptor with the model parameter. The significance level of the correlation is given in parentheses.

^{*}Significant at the 95% confidence level ($P \le 0.05$).

rate of self-thinning, except q, which had no effect, and initial density, which was not amenable to this analysis. The log B-log N plot of the control group simulations (Figure 3.2) shows that there are variations in the self-thinning trajectory that can only be attributed to persistent effects of the stochastic initial conditions. However, the slopes of the self-thinning lines are quite similar and the lines are all positioned within a narrow vertical band.

Only the allometric power, p, altered the self-thinning slope. This effect is demonstrated visually by comparing the experimental thinning diagram (Figure 3.3a) to Figure 3.2 and quantitatively by the high Spearman correlation between p and $\hat{\beta}$ ($r_s = 0.99$, P < 0.0001). The relationship between $\hat{\beta}$ and 1/p is linear (Figure 3.3b), and the regression equation $\hat{\beta} = -0.555(1/p) + 1.05$ ($r^2 = 0.98$, P < 0.0001, 95% CI for slope = [-0.604, -0.506], 95% CI for intercept = [0.91, 1.18]) was very close to the ideal linear relationship $\hat{\beta} = -0.50$ (1/p) + 1.00 developed in Chapter 2. The parameter p also altered the self-thinning intercept, $\hat{\alpha}$, ($r_s = -1.00$, P < 0.0001), which was also linearly related to (1/p) (Figure 3.3c), again as predicted in Chapter 2. The regression equation was $\hat{\alpha} = 2.97(1/p) - 4.18$ ($r^2 = 0.99$, P < 0.0001, 95% CI for slope = [2.80, 3.13], 95% CI for intercept = [-4.65, -3.72]).

The parameter d, the density of biomass in occupied space, affected the position of the self-thinning line (Figure 3.4a), as measured by the thinning intercept $\hat{\alpha}$ (r_s =0.99, P < 0.0001).




Figure 3.2. Variations in the self-thinning trajectory due to stochastic factors in the simulation model. The ten runs all used the reference parameter set (Table 3.2) and differed only in the initial plant locations and weights.






Figure 3.3. Effect of the allometric power, p, on the self-thinning line. (a) shows 13 simulations with p values between 0.27 (highest curve) and 0.49 (lowest curve). In (b), the fitted thinning slopes, $\hat{\beta}$, (Table 3.4) are plotted against (1/p). The solid regression line is $\hat{\beta} = -0.555(1/p) + 1.05$ ($r^2 = 0.98$, P < 0.0001, 95% CI for slope = [-0.604, -0.506], 95% CI for intercept = [0.91, 1.18]). The dotted line is $\beta = -0.5$ (1/p) + 1 (Chapter 2). (c) plots the thinning intercept, $\hat{\alpha}$, against (1/p) and the regression line $\hat{\alpha} = 2.97(1/p) - 4.18$ ($r^2 = 0.99$, P < 0.0001, 95% CI for slope = [2.80, 3.13], 95% CI for intercept = [-4.65, -3.72]).




Figure 3.4. Effect of the density of biomass in occupied space, d, on the self-thinning line. (a) shows trajectories for ten values of d between 631 (lowest curve) and 39810 (highest curve), while (b) plots the fitted thinning intercept, $\hat{\alpha}$, (Table 3.4) against log d. The regression line is α = 1.19 log d + 0.09 (r^2 = 0.97, P < 0.0001).

The relationship between $\hat{\alpha}$ and $\log d$ is apparently linear (Figure 3.4b), as predicted in Chapter 2. The regression equation is $\hat{\alpha}=1.186\log d+0.0891$ ($r^2=0.97,\ P<0.0001$). Thinning slope was not affected by d, as shown by the low correlation between d and $\hat{\beta}$ ($r_s=-0.49,\ P=0.15$) and by the fact the range and CV of $\hat{\beta}$ in the experimental group (0.17 and 9.2%, respectively) were not greater than the corresponding values for the control group (0.18 and 10.2%).

The thinning lines are more widely spread among the six simulations using the different competition algorithms (Figure 3.5) than among control group simulations, indicating that thinning line position is affected by this algorithm. Spearman correlations could not be calculated for this simulation experiment because the competition algorithm is a nominal variable that can not be ranked, but further support does come from a comparison of the ranges of variation of $\hat{\alpha}_{3,35}.$ This range is eight times larger among the experimental simulations than in the control group, and the CV is almost ten times larger. A similar comparison of $\hat{\alpha}$ values seems to contradict this conclusion; however, when the interpretations $\hat{lpha}_{3/35}$ and \hat{lpha} differ, $\hat{lpha}_{3/35}$ is the more reliable measure of thinning line position because it lies in the middle of the range of the data while $\hat{\alpha}$ is an extrapolation of the regression lines well outside the data (Sokal and Rohlf 1981). A lack of effect of the competition algorithm on thinning slope is indicated by the similar means and ranges for $\hat{\beta}$ in the control and experimental groups.




Figure 3.5. Effect of the competition algorithm on the self-thinning line. Curves for six simulations are are marked with competition algorithm identification numbers (Table 3.1).

The assimilation rate, g_1 , had no effect on the self-thinning trajectory, as indicated by similarity of Figure 3.6a to Figure 3.2, the near equality of the ranges and CVs of the thinning line descriptors to the same statistics for the control group, and the low Spearman correlations of g_1 with the thinning line descriptors.

Metabolic cost parameters b and q were likewise unimportant in positioning the self-thinning line (Figure 3.6b). In this simulation experiment, the power q relating maintenance cost to weight was varied from 0.5 to 3.5. Simultaneously, the parameter b was adjusted so that the initial maintenance cost (second term of equation 3.1) for a plant of weight \bar{w}_0 (10⁻⁴ g) was constant across the seven simulations. However, as growth increased average weight above \bar{w}_0 , differences in maintenance cost due to differences in q became important. Interpretation of this simulation experiment is complicated by the two trajectories that are decreasing in both biomass and density by the end of the simulation. The explanation is that maintenance costs affect the maximum possible size for a model plant, which can be calculated by setting equation 3.6 equal to zero and solving for w_{max} to obtain

$$w_{\text{max}} = [g_1 c_1 / b_1]^{1/(q-2p)}$$
 (3.10)

The differences in q across the simulations led to differences in w_{max} and this limitation became important in two simulations; however, all the simulations followed a common path through the log B-log N plane until the two populations became limited by




Figure 3.6. Simulations for three parameters that did not affect the self-thinning line. In (a) parameter g_1 was varied from 10 to 50 $g/m^2/t$ ime unit across ten simulations. (b) was generated by using six values of parameter q between 0.5 and 3.0. Two curves are marked with the associated values of q (Table 3.4). (c) shows thinning trajectories for five simulations started at initial densities of 3160, 10000, 31600, 100000, and 316000 plants/ m^2 .

 w_{max} and diverged sharply from the common path. Before fitting the thinning line in the q = 3.0 simulation, data points with log N values below 3.4 were eliminated. No thinning line was fitted to the q = 3.5 simulation since the maximum weight limitation took effect before the linear portion of the thinning trajectory was established. The visually evident lack of effect of q on the thinning line estimates is further supported by the failure of $\hat{\beta}$, $\hat{\alpha}$, and $\hat{\alpha}_{3.35}$ to exceed the limits established in the control group and by the low Spearman correlations between q and the thinning line descriptors.

Variations in initial density also had no effect on the position of the self-thinning line (Figure 3.6c). The five different initial densities were created by placing 200 plants on different sized plots. This was the only way to vary initial density over a large range because model limitations precluded direct manipulations of the initial number of plants. The maximum initial number was 600 and simulations were stopped when less than 20 plants remained because the thinning trajectory made undesirably sharp jumps at low population sizes. Although the resulting trajectories start from different points in the plane, all converge on the same thinning line, at least within the limits of the stochastic variation seen in Figure 3.2. A more quantitative analysis was not possible because there was no common range of linear behavior over which to fit self-thinning lines, so this simulation experiment is not included in Tables 3.3 and 3.4.

Discussion

This analysis has identified only one parameter, the power p relating area occupied to individual weight, that affected the slope of the model self-thinning line. Three parameters affected thinning line position: p; the density of biomass in occupied space, d; and the competition algorithm. These results agree with the simpler model of Chapter 2 and together suggest that the regularities of self-thinning can be explained by the shape-dependent occupation of space. The linearity and slope of the self-thinning line are determined by the power relationship relating space occupied to plant size, and the thinning intercept is related to at least two additional factors. The implications of these results have already been discussed in Chapter 2. The possibility that the conclusions of Chapter 2 were biased due to an over-simplified, spatially homogenous model can now be discarded because the detailed representation of individual size, location, and competitive interactions in the simulation model led to the same results. Although spatially averaged models are not useful for many applications in plant ecology (Schaffer and Leigh 1976), they are appropriate tools for investigating self-thinning because of the unusual degree of spatial uniformity that is present in even-aged monospecific stands.

Model parameters representing resource availability and utilization efficiency (maximum assimilation rate \mathbf{g}_1) and

metabolic costs (q and b) affected only the rate of self-thinning, not the slope or position of the thinning line. Initial density also did not affect the thinning line. However, the model did not allow the parameters of individual shape and density, p and d, to vary in response to these important environmental factors. Real plants can respond to environmental limitations by varying their shapes (Hutchings 1975, Harper 1977) and canopy densities (Lonsdale and Watkinson 1983a), and density stress can cause permanent alterations in plant geometry (Peet and Christensen 1980). The exact responses would vary among species. Environmental factors could, then, induce changes in the self-thinning line, not because the factors are key determinants of the thinning line, rather because they affect the thinning line indirectly by altering the growth parameters of the plants.

The important effects of the competition algorithm on the rate of self-thinning and the position of the thinning line are significant new results of this analysis. The competition algorithm is related to the allowable overlap between plants (Chapter 2) and to the degree of asymmetry in competitive interactions, so measures of these factors should be related to the self-thinning line. This prediction is tested in Chapters 5 and 6, where the relationships of shade tolerance with thinning slope and intercept are considered.

The persistent effect of the stochastic initial conditions seen in Figure 3.2 is also interesting. Since all parameters were held constant, these differences could only be attributed to random variations in the initial weight distributions and initial plant

locations. This result suggests that even if all other sources of variation could be eliminated from thinning experiments, observed self-thinning lines could still differ in position because of these stochastic factors.

Some results of this analysis are relevant to other theories about the causes of the self-thinning rule. The unimportance of metabolic parameters b and q in positioning the self-thinning line arques against the hypothesis that the self-thinning rule derives from a 2/3 power relationship between metabolic costs and plant weight (Jones 1982), and against an important role in fixing the self-thinning slope for the fraction of assimilate devoted to maintenance (Pickard 1983). The simulations of Figure 3.5 indirectly address another theory that the -3/2 exponent of the $\overline{w}-N$ relationship is maintained during thinning by mutual adjustment of plant allometry and stand structure (the distribution of individual sizes--Mohler et al. 1978). Figure 3.7 shows how the frequency distributions of individual plant weight change with time for the four simulations generated by competition algorithms 1, 2, 3, and 5. These time plots of the first four moments of the weight distribution (average, coefficient of variation, skewness, and kurtosis) show that the four competition algorithms lead to widely different dynamics of stand structure. If the mutual adjustment hypothesis is correct, thinning slopes should also be different since stand structure varies without any possibility of compensatory adjustments in plant allometry (parameter p was 1/3 for all four




Figure 3.7. Dynamics of the weight distributions in four simulations with different competition algorithms. Algorithm numbers (Table 3.1) were 1 (solid line), 2 (dashed line), 3 (dotted line), and 4 (chain-dotted line). The mean, coefficient of variation, skewness, and kurtosis of each of the weight distributions are shown in (a) through (d). Methods of calculation are given in Sokal and Rohlf (1981).

simulations). However, if plant allometry sets the thinning slope, the simulations should give the same thinning slope despite the differences in competitive regime and size distribution dynamics. Since thinning slope did not vary among the four simulations, the allometric hypothesis is supported and the mutual adjustment hypothesis is not.

CHAPTER 4

SOME PROBLEMS IN TESTING THE SELF-THINNING RULE

Introduction

Many self-thinning lines with slopes near $\gamma=-3/2$ have now been reported, and their sheer number is considered strong evidence for the self-thinning rule, or even a self-thinning law (see references in Hutchings 1983). However, an analysis of size-density data is not simply a matter of regressing log \overline{w} against log N and so demonstrating the self-thinning rule. Some important analytical difficulties must be fully discussed before the large body of evidence is embraced as convincing proof for the rule. The principal problem areas are (1) the data selected to test the hypothesis, (2) the points used to estimate a thinning line, (3) the curve fitting methods used, (4) the choice between the log \overline{w} -log N or log B-log N formulations of the rule, and (5) the conclusions drawn from the results. Recommendations for resolving some difficulties are presented here, and the implications for acceptance of the self-thinning rule are discussed.

<u>Selecting Test Data</u>

Many data sets have been reported to exhibit a linear relationship between $\log \bar{w}$ and $\log N$ with a slope near $\gamma = -3/2$, but it is usually easy to find evidence for a hypothesis, regardless of whether or not it is generally true. Therefore, the mere

existence of such evidence does not verify the hypothesis. Rigorous verification instead comes from <u>failure</u> of the opposite endeavor, to find evidence that contradicts or falsifies the hypothesis (Popper 1963). The emphasis on compiling corroborative evidence for the thinning rule has diverted attention from data that do not conform. Violations of the rule have been discussed only for shoot populations of clonal perennials (Hutchings 1979) and for thinning under very low illumination (Westoby and Howell 1982, Lonsdale and Watkinson 1982), but there are other violations besides these special cases, such as thinning slopes of $\hat{\gamma} = -2.59$ and -4.5 for tropical trees <u>Shorea robusta</u> and <u>Tectona grandis</u> (O'Neill and DeAngelis 1981), $\hat{\gamma} = -1.2$ the temperate tree <u>Abies balsamea</u> (Sprugel 1984), and $\hat{\gamma} = -3.2$ for seedling populations of the woodland herb Allium ursinum (Ernst 1979).

Information contradicting the self-thinning rule has been missed even in the very sources from which supporting evidence has been drawn. For example, data for one stand of <u>Pinus strobus</u> (Spurr et al. 1957) have been repeatedly cited in self-thinning studies (Hozumi 1977, 1980, Hara 1984), and a self-thinning slope of $\hat{\gamma}$ = -1.7 has been fit (White 1980). However, the report of Spurr et al. also presented data for a second stand which gives a thinning slope of $\hat{\gamma}$ = -2.11 (Table A.2). This second, unreported stand contradicts the self-thinning rule in two ways: the thinning slope is quite different from the predicted value, and both the slope and intercept change from stand to stand. A second example refers to a yield table for Pinus ponderosa (Meyer 1938) reported to give a

thinning line of log \bar{w} = -1.33 log N + 4.06 (White 1980). However, the log B-log N thinning plot for the complete yield table (Figure 4.1a) shows that 13 thinning lines could be fit since information is given for 13 values of site index, a general measure of site including soil composition, fertility, slope, aspect, and climate (Bruce and Schumacher 1950). The existence of the twelve unreported thinning lines contradicts two tenets of the self-thinning rule: the thinning line is not independent of site quality and the thinning intercepts are not species constants. The data from some yield tables even give different thinning slopes for different site indexes, as shown in Figure 4.1b for Sequoia sempervirens (Lindquist and Palley 1963).

Editing Data Sets

A second major problem in testing the self-thinning rule arises because the thinning line is an asymptotic constraint approached only as stands become sufficiently crowded. To estimate the slope and position of a thinning line using linear statistics, data points from populations that are not limited by the hypothesized linear constraint must be eliminated. These would include points from young populations that have not yet reached the thinning line, older stands understocked because of poor establishment or density-independent mortality, and senescent stands. Failure to eliminate such points will bias the thinning line estimates (Mohler et al. 1978), but when the data are confounded by biological variability and measurement errors, recognition and elimination of




Figure 4.1. Two forestry yield tables showing variation in thinning line parameters with site index. Dotted lines connect data points and solid lines are PCA thinning lines. (a) shows data for Pinus ponderosa (Meyer 1938). Thinning slopes, $\hat{\beta}$, for ten site indexes were near -0.31 (between -0.307 and -0.326), but intercepts ranged from 3.75 to 4.18. (b) shows data for Sequoia sempervirens (Lindquist and Palley 1963). Thinning slopes for six site indexes ranged from -4.15 to -1.81 while intercepts ranged from -1.93 to 2.67. Tables B.1 and B.2 give additional information.

spurious points is difficult. Since there is no <u>a priori</u> estimate of the thinning line position, decisions to eliminate data points must be made <u>a posteriori</u> (Westoby and Howell 1982). This is true even if detailed field notes (Mohler et al. 1978) or mortality curves (Hutchings and Budd 1981a) are available to aid the process. With such <u>a posteriori</u> manipulations, no thinning analysis can be done in a strictly objective way. Figure 4.2 presents three data sets that illustrate these problems. Each plot shows how the slope, intercept, r^2 , significance level, and confidence interval all change with the points used to fit the thinning line (Table 4.1). The results are very sensitive to certain points, yet there is no objective way to decide whether or not to include those points.

The sensitivity of thinning line parameters to the choice of data points also has important statistical implications. The uncertainties about including or excluding some points should be counted in forming confidence intervals and performing tests of significance. However, existing statistical methods do not take such uncertainties into account, so estimated r^2 values are too high and confidence intervals are too narrow.

Some data sets show more that one region of linear behavior and so present the analyst with still another subjective decision: Which linear region is relevant to the self-thinning rule? Figure 4.3 presents a yield table for <u>Populus deltoides</u> (Williamson 1913) that illustrates this problem. White (1980) fit the thinning line $\bar{w} = -1.8 \log N + 3.08$ through the data for ages 7 to 15, while




Figure 4.2. Three examples of the sensitivity of self-thinning line parameters to the points chosen for analysis. In each plot, several thinning lines (capital letters) are fitted to different combinations of points (numbers) as indicated in Table 4.1. Data in (a), (b), and (c) are for Populus tremuloides (Pollard 1971, 1972), Triticum sp. (Puckridge and Donald 1967, White and Harper 1970), and Tagetes patula (Ford 1975).

Table 4.1. Three Examples of the Sensitivity of the Fitted Selfthinning Line to the Points Chosen for Analysis.

							PCA Thinning L	_ine ^e	
		Points					Slope	Intercept	
Figure ^a	Line ^b	Included ^C	r ²	Pd	Ē	3	95% CI	â	
4.5a	 А В	1-2 1-3	1.00	0.16	-0.1 -0.2			3.73 3.66	
	С	1-3	0.94*				T 0 40 0 077	3.65	
							[-0.48, -0.07]		
	D	2-4	0.99*	0.010	-0.5	3	[-0.64,-0.42]	3.72	
4.5b	Α	1-10	0.63*	0.0061	-0.2	20	[-0.33,-0.08]	3.32	
	В	1-3,5-10	0.86*	0.0003	-0.2	24	[-0.32, -0.15]	3.43	
	С	5-10	0.91*	0.0030	-0.3	31	[-0.46, -0.18]	3.70	
	C D E F	6-10	0.98*	0.0008	-0.3	36	[-0.45, -0.28]	3.84	
	Ε			0.0027			[-0.25, -0.08]	3.26	
	F	1-3,5-8		0.027			[-0.24,-0.02]	3.17	
	•	. 0,0 0	0.00	0.027	•	. •	[0.2., 0.0.]	••••	
4.5c	Α	1-10	0.34	0.075	-0.4	41		4.36	
	В	1-8	0.06	0.54	+0.2			2.22	
	Č	3-10		0.056	-0.			5.38	
	Ď	4,6,9,10	0.84	0.079	-0.3			4.03	
	U	7,0,2,10	0.04	0.073	-0.	,,		1.00	

aSpecies names and references are given in the legend of Figure 4.2.

CThe numbers of data points in Figure 4.2 used to fit the thinning line.

dThe statistical significance of the log B-log N correlation.

 $^{\mbox{\scriptsize e}}\mbox{\sc The method of fitting the thinning line by principal component analysis is discussed in Chapter 5.$

bThese letters label the fitted lines in Figure 4.2.

^{*}Significant at the 95% confidence level ($P \le 0.05$).




Figure 4.3. Example of a data set with two regions of linear behavior. White (1980) fit the dotted line $\log \bar{w} = -1.80 \log N + 3.08$ to yield table data (solid line) for juvenile stands (7-15 years old) of Populus deltoides (Williamson 1913). Older stands follow the dashed line $\log B = -0.20 \log N + 3.85$ fit to 25-50 year old stands (Table B.2)

the present study has fit a line of snallower slope, $\log B = -0.20$ $\log N + 3.85$, to the data for ages 25 through 50. The line fit by White may be more typical of the steep ascent of juvenile populations through the $\log B - \log N$ plane, while the second line may represent the self-thinning behavior of more mature stands.

Fitting the Self-thinning Line

Most self-thinning lines have been estimated by linear regression of $\log \bar{w}$ against $\log N$, but regression is inappropriate for thinning data because $\log N$ is not a good independent variable, that is, it is neither measured without error nor controlled by the experimenter. Although principal component analysis (PCA--Mohler et al. 1978) and geometric mean regression (GMR--Gorham 1979) have been proposed as more appropriate fitting methods, the true self-thinning line is actually not estimable. In general, the slope, $\hat{\beta}_{yx}$, of a linear bivariate relationship for Y in terms of X is given by the linear structural relationship (Madansky 1959, Moran 1971, Jolicoeur 1975),

$$\hat{\beta}_{yx} = \frac{s_{yy} - \lambda s_{xx} + [(s_{yy} - \lambda s_{xx})^2 + 4 \lambda s_{xy}^2]}{2 s_{xy}}, (4.1)$$

where $s_{\chi\chi}$, $s_{\chi\chi}$, and $s_{\chi\chi}$ are sums of squares and cross products corrected for the mean and λ is the ratio of the error variance in Y to the error variance in X. These error variances are the

residuals around the true linear relationship between Y and X, not the total sample variances. Particular values of λ give the standard methods as special cases of this general equation: $\lambda = \infty$ (no error in X) gives $\hat{\beta}_{yx} = s_{xy}/s_{xx}$, the regression of Y against X; λ = 0 (no error in Y) gives $\hat{\beta}_{yx} = s_{yy}/s_{xy}$, as obtained from regressing X against Y; λ = 1 (equal marginal variances of X and Y) gives the PCA solution; and $\lambda = s_{yy}/s_{xx}$ (marginal variances in the same ratio as the total sample variances) gives the GMR solution, $\hat{\beta}_{vx} = sign(s_{xv}) s_{vv}/s_{xx}$. If the relationship between X and Y is strongly linear (high r^2), then all solutions are similar, but as the association becomes less strict (lower r^2), the discrepancies among the solutions increase and results become more sensitive to λ . Figure 4.4 shows these four particular solutions for a typical thinning data set (Mohler et al. 1978). Although this data set showed a very significant log B-log N relationship ($r^2 = 0.40$, P < 0.0001), the four solutions ranged from $\hat{\beta}$ = -0.34 to -0.87, and the regression of log B against log N gave -0.34 while PCA gave -0.41.

Two sources of variation are reflected by λ : measurement errors, which can be estimated by replication, and natural biological variability, which can not be estimated (Ricker 1973, 1975). Since λ can not be known, the true solution of equation 4.1 is unestimable (Ricker 1975, Sprent and Dolby 1980) and the slope of the self-thinning line must be based on some assumed value of λ . For most self-thinning data, there is no basis to




Figure 4.4. Thinning lines fit to one data set by four methods. The data are for <u>Prunus pensylvanica</u> (Mohler et al. 1978). Lines were fit to the 34 square points by regression of log B on log N, regression of log N on log B, geometric mean regression, and principal components analysis (lines A, D, C, and B, respectively). Equations for the four lines were, respectively, log B = -0.34 log N + 3.88, log B = -0.87 log N + 4.19, log B = -0.55 log N + 4.00, and log B = -0.41 log N + 3.92, while confidence intervals for the slopes were [-0.50, -0.19], [-1.58, -0.60], [-0.72, -0.42], and [-0.61, -0.23]. For all four lines, $r^2 = 0.40$ and P < 0.0001.

assume the marginal variation of log N is less than that of log B (or vice versa), so the most reasonable assumption is that the two marginal variances are approximately equal, giving λ = 1 and leading to the PCA solution of equation 4.1. Regression analysis implicitly assumes that one of the variables is error free and λ takes one of the extreme possible values λ = 0 and λ = ∞ (Moran 1971). Since this extreme assumption is clearly untrue for self-thinning data, many reported thinning lines estimated by regression analysis are potentially in error.

The use of PCA has created yet another statistical problem: many studies now report the percentage of variance explained (%EV) by the first principal component (PC) rather than the correlation coefficient, r, or r^2 . Unlike r^2 , which ranges from 0 to 1, %EV ranges from 0.5 to 1 because the first PC always explains at least 50% of the total variation, even if the two variables are completely uncorrelated. Therefore, %EV values are always higher than r² values. This has been misinterpreted by some authors as an indication that PCA is more reliable then regression. Actually, the exact method of fitting a straight line (Y-X regression, X-Y regression, PCA, or GMR) is irrelevant to the calculation and interpretation of correlations or coefficients of determination since these measure the strength of linear association rather than the position of any particular line in the plane. The reported measure of association should always be r or r^2 , regardless of the fitting method used (Sprent and Dolby 1980).

Choosing the Best Mathematical Representation

The recommended use of PCA focuses attention on another question: Should the thinning rule be tested by relating $\log \bar{w}$ to $\log N$ or by relating $\log B$ to $\log N$? The two choices are mathematically equivalent (Chapter 1), and when the regression of $\log B$ against $\log N$ is compared to the regression of $\log \bar{w}$ on $\log N$ for a set of data, the slopes differ by exactly one $(\beta = \gamma + 1)$ and the confidence intervals have identical widths. However, neither of these conditions holds when $\log \bar{w} - \log N$ PCA is compared to $\log B - \log N$ PCA, so one CI may include the predicted slope of the self-thinning rule while the other CI does not. Although mathematically equivalent, the $\log \bar{w} - \log N$ and $\log B - \log N$ formulations of self-thinning rule are <u>not</u> statistically equivalent when appropriate curve fitting methods are used, so an explicit decision is required: Which formulation is more appropriate for testing the self-thinning rule?

The log B-log N formulation is the correct choice because the log \overline{w} -log N alternative suffers from two major limitations. Changes in average weight can be misleading because average weight increases when small individuals die, even if the survivors do not actually gain weight (Westoby and Brown 1980). Average size increases through two processes--growth of living plants and elimination of small plants--so that the average size of the stand increases more rapidly than the sizes of individuals composing it (Bruce and

Schumacher 1950). Attempts to relate $\log \bar{w}$ to $\log N$ are actually correlating some combination of growth and mortality with mortality, so the results are difficult to interpret. However, total stand biomass only increases through growth, so a correlation of $\log B$ with $\log N$ directly addresses the growth-mortality relationship and focuses attention on the extent to which mortality permits a more than compensatory increase in the size of the survivors.

The second shortcoming of the $\log \bar{w}$ - $\log N$ analysis is more serious and damaging to the case for the self-thinning rule: the analysis is statistically invalid, gives biased results, and leads to unjustified conclusions. To understand why, consider the methods used to measure plant biomass. Stand biomass is often measured directly by harvesting all the plants in a stand and weighing them as a single group. Even when each individual is weighed (or individual weights estimated from a relationship between weight and some plant dimension) stand biomass is still estimated directly as the sum of the individual weights. Average weight is then derived from the original stand measurements by dividing the biomass by the density. In general, there are "serious drawbacks" in analyzing such derived ratios (Sokal and Rohlf 1981), but these problems are particularly acute when the ratio is correlated with one of the variables from which it was derived. Such an analysis gives correlations that have been variously called "spurious" (Pearson 1897, as cited in Snedecor and Cochran 1956), "artificial" (Riggs 1963), and "forced" (Gold 1977).

A high correlation between $\log \overline{w}$ and $\log N$ is both unsurprising and meaningless because log N is used to calculate log \overline{w} . This is most easily seen when the original measurements of log B and log N are unrelated so that the sample correlation coefficient is low and not statistically significant. After deriving average weight from $\log \bar{w} = \log B - \log N$, $\log \bar{w}$ is a function of $\log N$ even though log B was not. The variance of log \overline{w} is Var(log B) + Var(log N) -2 Cov(log B, log N) (Snedecor and Cochran 1956), which reduces to Var(log B) + Var(log N) because the covariance of unrelated variables is zero. Thus, the variance of $\log w$ is higher than the variance of log B, and all of the additional variation is directly attributable to log N. Since log N explains more of the variance in $\log \bar{w}$ than in $\log B$, the correlation between $\log \bar{w}$ and $\log N$ is higher and more significant than the correlation between log B and log N. Although mathematically real, this higher correlation does not represent an increase in the information content of the data, but is a "wonderful tool for misleading the unwary" (Gold 1977).

The deceptive effects of this data transformation are also present in simple plots of the data. This is important because self-thinning data must be edited to remove extraneous points before fitting a thinning line. Since plots of the data are essential tools for recognizing such points, a data transformation that creates artificial linear trends in the plot will obviously disrupt the editing procedure. In the most extreme case, the distorted log \bar{w} -log N plot may suggest a linear relationship when none was present in the original log B-log N data. More subtle errors arise

when a real log B-log N correlation exists, but points that are not associated with the constraint of the thinning line are mistakenly included in fitting the thinning line.

A few examples will illustrate how the deceptive effects of the data transformation pervade the existing evidence for the self-thinning rule. In the first example, the log \overline{w} -log N plot (Figure 4.5a) for a study of <u>Trifolium pratense</u> (Black 1960) shows a linear trend and high correlation ($r^2 = 0.76$) between log \overline{w} and log N among nine data points purported to form a thinning line that agrees with the self-thinning rule (White and Harper 1970). However, the untransformed log B-log N plot (Figure 4.5b) shows the true situation: there is no significant negative correlation between log B and log N for the nine points, and time trajectories of stands cut steeply across the proposed thinning line rather than approaching it asymptotically.

Figure 4.6 illustrates a less extreme case where a linear trend is present, but the log \overline{w} -log N plot gives a distorted impression of which points lie along the constraining line. The log B-log N plot of the data (<u>Chenopodium album</u>, Yoda et al. 1963) shows an apparent linear constraint which could be estimated by fitting a line through the 13 square data points in Figure 4.6b. The remaining 14 points are distant from the constraint and should be removed before curve fitting, but this is hidden in the distorted log \overline{w} -log N plot (Figure 4.6a), where the artificial linearization causes the extraneous data points to fall in line with the others. Fitting a thinning line through all 27 data points by PCA of log \overline{w} against




Figure 4.5. Example of a spurious reported self-thinning line. (a) White and Harper (1970) fit the solid regression line $\log \bar{w} = -1.33 \log N + 3.86$ (n = 9, $r^2 = 0.76$, P = 0.003) to data for Trifolium pratense (Black 1960). In (b) the nine points are in the box and other points omitted by White and Harper are also shown. The original measurements of log B and log N for the nine points are uncorrelated ($r^2 = 0.13$, P = .33). Time trajectories for different initial densities (dotted lines) cut through the proposed thinning line and do not approach it.




Figure 4.6. Example of potential bias in data editing in the log \bar{w} -log N plane. Data are for <u>Chenopodium album</u> (Yoda et al. 1963). In the log \bar{w} -log N plot (a) all 27 data points follow a common linear trend, represented by the dotted line. The equation of this line, estimated from PCA of log B and log N, is log B = -1.33 log N + 3.94 (r^2 = 0.90, P < 0.0001, 95% CI for $\hat{\beta}$ = [-1.53, -1.16]). When the artificial enhancement of linearity is removed on examination of the log B-log N plot of (b), an apparent linear constraint is still evident; however, many data points fall relatively far from the constraining line (triangles). A new (solid) PCA line through the 13 points closer to the border of the constrained region has the equation log B = -0.41 log N + 5.15 (r^2 = 0.93, P < 0.0001, 95% CI for $\hat{\beta}$ = [-0.48, -0.33]), and the inadequacy of the dotted line is revealed.

log N gives log \bar{w} = -1.33 log N + 3.94 (r^2 = 0.90, P < 0.0001, 95% CI for slope = [-1.53, -1.16]), while the 13 points actually near the constraint in the log B-log N plot give log B = -0.41 log N + 5.15 (r^2 = 0.93, P < 0.0001, 95% CI for slope = [-0.48, -0.33]). The distorted log \bar{w} -log N plot changes the selection of relevant points, the estimated thinning line, and its comparison to the self-thinning rule.

The transformation can artificially straighten data that are actually curved in the log B-log N plane. The straight lines in the log \overline{w} -log N plane of Figure 4.7a seem to be a reasonable fits to the data (<u>Fagopyrum esculentum</u>--Furnas 1981), but the log B-log N plot (Figure 4.7b) reveals the true curvature of the data and the inadequacy of the straight line model. This curve straightening deception is important in editing data sets with juvenile or senescent stands, which often curve gradually toward or away from the thinning line.

A final example shows how the log \overline{w} -log N plot can lead to questionable conclusions about the effects of an experimental treatment on self-thinning. To evaluate the effects of a fertilizer treatment on self-thinning behavior, five plots of <u>Erigeron</u> <u>canadensis</u> received different fertilizer applications in a ratio of 5:4:3:2:1 before seeds were planted (Yoda et al. 1963). The conclusion that the thinning trajectory was insensitive to soil fertility seems justified in the log \overline{w} -log N plot (Figure 4.8a), where the five treatments seem to approach the same thinning line despite the large differences in fertility. However, the




Figure 4.7. Example of deceptive straightening of curves in the log \bar{w} -log N plane. Data are from an experiment with <u>Fagopyrum esculentum</u> where the circle population received five times as much fertilizer as the square population (Furnas 1981). Both trajectories appear reasonably linear in the log \bar{w} -log N plot (a), and the two lines log \bar{w} = -1.50 log N + 4.837 and log \bar{w} = -1.50 log N + 4.622 reported by Furnas seem to fit the data well. The log B-log N plot (b) reveals the the inadequacy of the straight line model for these data.




Figure 4.8. Potential misinterpretation of experimental results due to deceptive effects of the log \bar{w} -log N plot. Numbers marking data points (<u>Erigeron canadensis</u>--Yoda et al. 1963) indicate the relative amount of fertilizer applied before planting. Yoda et al. concluded that the overall trend of the data show a slope of -1.5 (chain-dot line) in the log \bar{w} -log N plot (a). White (1980) reported the dotted regression line log \bar{w} = -1.66 log N + 4.31 (r^2 = 0.93, P < 0.0001, 95% CI for slope = [-1.86, -1.47]). In (b) the present study fit the PCA line (solid line) log B = -1.04 log N + 5.70 (r^2 = 0.99, P < 0.0001, 95% CI for slope = [1.12, -0.96]).

log B-log N (Figure 4.8b) plot suggests a very different interpretation. The populations do respond to fertility differences and initially follow very different trajectories, but the differences gradually disappear and are gone around the fourth harvest, about 4.5 months after planting. It seems reasonable that over this long interval, leaching and plant uptake removed fertilizer from the soil, all five treatments approached a background fertility level, and only then did the treatments converge on a common trajectory.

The log B-log N plot also suggests a thinning line that is very different from the expected value of β = -1/2. This plot reveals a fact that is obscured in the log \overline{w} -log N plot: all five plots lost biomass between harvests two and three, which is not surprising because these harvests were made during the winter. Because winter conditions apparently interfered with growth, early harvests should not be included in estimating a thinning line. PCA analysis of harvests four through six gives log B = -1.04 log N + 5.70 (r^2 = 0.99, P < 0.0001, 95% CI for slope = [-1.12, -0.96]). This slope is not close to the hypothesized value of β = -1/2 and is statistically different from -1/2 (P < 0.0001).

Testing Agreement with the Self-thinning Rule

Evaluation of the self-thinning rule has been hindered by the lack of an objective definition of how close to the predicted value a thinning slope must be to agree quantitatively with the rule.

White (1980) reported that many thinning slopes fall between

 $\hat{\gamma}$ = -1.8 and $\hat{\gamma}$ = -1.3 and presented all these values as examples of the same quantitative rule. However, this arbitrary range has no objective basis and its limits represent very different predictions about population growth. A population following a thinning trajectory of slope $\gamma = -1.3$ over a 100-fold decrease in density will increase its biomass about fourfold, while a population following a trajectory of slope $\gamma = -1.8$ will increase its biomass about 40-fold over the same 100-fold density decrease. Although the two populations show qualitatively similar behavior in following a linear trajectory in the log size-log density plane, they show a tenfold difference in biomass response to the same degree of density decrease and thus would not seem to obey the same quantitative rule. Over a 1000-fold density decrease, one population shows an eightfold biomass increase, while the other thinning population shows a 250-fold biomass increase and the discrepancy between the two thinning regimes increases exponentially as larger amounts of density decrease are considered.

A statistical test of the hypothesis that an observed thinning slope is equal to the predicted value provides a more objective test of agreement with the self-thinning rule. The confidence interval for the slope of a fitted thinning line can be compared to the predicted value. If the CI includes the predicted value, then the data can not be said to contradict the rule and the slope is close to the idealized value within the limits of resolution of the data. If the CI does not contain the predicted slope, then the data are in quantitative disagreement with the self-thinning rule.

This statistical test of agreement with the self-thinning rule should be accompanied by tests of relevant alternative hypotheses. Only data sets that can discriminate among alternatives can provide convincing support for the self-thinning rule. Two alternative hypotheses are particularly relevant: the null hypotheses that no competitive effect was observed, and the hypothesis that a carrying capacity keeps stand biomass below a fixed level. If the null hypothesis is true, there will be no significant relationship between log B and log N or a positive relationship. If a biomass is fixed at a constant level, there will again be no significant relationship between log B and log N. Only data sets that have a statistically significant negative correlation between log B and log N can clearly reject these alternatives. Data sets that do not reject the null hypothesis do not support the self-thinning rule or any other hypothesis about the course of plant competition. Such data do not even demonstrate the presence of competition.

Unfortunately, the interpretation of hypothesis tests is confounded by the uncertainties of data editing. The analysis can not account for these uncertainties, so the confidence interval of the fitted thinning slope is too narrow and the probability of making a Type I statistical error (rejecting a true null hypothesis) is increased by some unknown amount. A test with the 95% confidence interval actually gives less than 95% confidence.

Discussion

The problems with the evidence for the self-thinning rule can be grouped into three categories of solvability: (1) omissions that can be rectified by analyzing and interpreting a greater breadth of information, (2) problems that can be fixed by using the best available statistical methods, and (3) unsolvable problems. The inattention to potentially contradictory information can be remedied by analyzing new data and by integrating existing contradictory examples into the body of data used to test the rule and delimit ranges of variation for thinning line parameters.

There are three areas where improvement of analytical methods is possible: the analysis should relate log B to log N because the alternative of relating log \overline{w} to log N is statistically invalid; PCA should be used rather than regression, which relies on unrealistic assumptions about the error structure of the data; and statistical tests of hypothesis should be used to interpret the results. The importance of making these improvements can be partially evaluated by quantifying the prevalence of less desirable methods. Table A.5 lists the methods used to estimate 76 reported self-thinning lines for experimental and field data (EFD) and Table B.4 gives similar information for 16 thinning lines for forestry yield table data (FYD). All of the FYD and 69 (91%) of EFD thinning lines were estimated by relating log \overline{w} to log N. For this large majority of studies, the points selected for analysis may not be the best choices, the correlations are inflated, spurious relationships may

have been created and interpreted, and the interpretation of the effects of experimental treatments and environmental factors may be biased. Few studies have included explicit statistical tests of the hypotheses that the estimated thinning slope is zero or equal to the value predicted by the thinning rule. Even if they had, the results of the tests would be meaningless because of the spurious component of the relationship between log \bar{w} and log N present in all studies that used the log \bar{w} -log N analysis.

The majority of studies have not used the best curve fitting methods. All of the FYD thinning lines were fit by regression, while 37 (49%) of the EFD thinning lines were regressions and only 15 (20%) used the more appropriate technique of PCA. Nineteen (25%) were graphical analyses that used no statistics, four (5%) assumed axiomatically that the thinning slope was $\gamma = -3/2$, and one (1%) one did not specify the statistical method.

The third category of unsolvable problems includes the need to edit data before fitting the self-thinning line. Editing of data before addressing some hypothesis should be suspect because it reduces the objectivity of the analysis and affects the confidence levels of statistical tests, but no alternative procedure for thinning analysis has been developed. However, these deleterious effects of editing are seriously compounded when the original data are transformed to average weights before editing, which can lead to false linear trends and disguise the need to remove certain points from consideration. These effects can be avoided by analyzing data in log B-log N form.

The prevalence of some correctable problems in analysis and interpretation of self-thinning data suggests that a major reanalysis of the evidence is required. Such an analysis is presented in Chapter 5.

CHAPTER 5

NEW TESTS OF THE SELF-THINNING RULE

Introduction

This chapter presents some new analyses of the support for the self-thinning rule and the extent of variation among self-thinning lines. This is the first effort to critically and comprehensively review the full spectrum of available evidence, and it reveals patterns that are not apparent when the evidence is judged on a piece-by-piece basis. No analysis can be perfect because the data must be edited before fitting the thinning line, resulting in some loss of objectivity and distortion of significance tests (Chapter 4). However, the analysis can be done as rigorously and consistently as possible by: (1) eliminating a priori biases, (2) using log B-log N data to avoid compounding the problems of editing with the artificial linearity and spurious correlation of $\log \overline{w}$ - $\log N$ plots, (3) using the best method for estimating the self-thinning line (principal component analysis of log B and log N), and (4) using statistical tests to objectively interpret the results.

This analysis considers 488 self-thinning lines, 137 from experimental and field data and 351 from forestry yield tables. This is the largest sample of thinning lines considered in any single study and includes many data sets not previously analyzed in a self-thinning context. The new data are important in guarding

against any a priori bias toward the self-thinning rule. Five different analyses address the constancy of observed thinning lines and their agreement with the self-thinning rule: (1) Statistical hypothesis tests are used to test for significant relationships between log B and log N and for quantitative agreement with the self-thinning rule. (2) The frequency distributions of thinning slope and intercepts are prepared to provide a complete description of the observed variations. (3) Plant groups are compared to determine if thinning line parameters vary among groups or if a single thinning rule applies to all plants. (4) The constancy of thinning line parameters with respect to a functional measure of plant performance is tested by checking for significant correlations of thinning slope and intercept with shade tolerance. (5) Finally, when several thinning lines were estimated for a particular species, the results are compared to determine if thinning slope and intercept are species constants.

Methods

Data were collected from two classes of studies:

(1) experimental and field studies (EFD) and forestry yield tables (FYD). Thirty-nine sources of EFD and 51 sources of FYD were examined (Tables A.1 and B.1). Many of these sources reported more than one study or yield table, so 95 studies were represented in the EFD and the FYD contained 77 yield tables. Some of these contained information for several thinning lines, so that a total of 488 thinning lines were fit, 137 for EFD and 351 for FYD. To ensure

comparability among studies, stand biomass values were converted to common units of g/m^2 while plant densities were converted to individuals/ m^2 . For forestry data, stand yields in volumes of wood were converted to weights per unit of forest area (g/m^2) by multiplying by the wood density of the species (Table B.1).

Fifty-six different species were represented in monospecific studies in the EFD, along with seven different two-species combinations, three multi-species mixtures, and two studies of monogeneric populations of unspecified species. The FYD considered 32 different single-species forests, two types of two-species forests, four multi-species forest types, and seven monogeneric forests of unspecified species. Some of the single-species forest types were examined in both the EFD and FYD. When these duplications are accounted for, a total of 78 different monospecific population types were considered. The multi-species stands were included because the thinning rule has been proposed to also apply for two species stands (White and Harper 1970, Bazazz and Harper 1976, Malmberg and Smith 1980) and multi-species stands (White 1980, Westoby and Howell 1981).

Some of the EFD studies presented data only in graphical form. In an initial attempt to acquire these data, requests for listings were sent to the authors of 12 studies and two data sets were successfully completed. The remaining graphical data sets were reconstructed from graphs by measuring the horizontal and vertical positions of each data point and axis tic mark to the nearest 0.05 mm with a micrometer. Regressions relating labeled axis values

to the micrometer measurements were used to recover the actual data values. Coefficients of determination for these regressions were uniformly high $(r^2 > 0.98)$, indicating that the graphs were of good quality and that this method was adequate for recovering the data. For individual numbers published in both written and graphical form, the reconstructed values were within 5% of the true values. Slopes, intercepts, and r^2 values estimated from reconstructed data were within 2% of the published values when reanalyzed with the points and fitting methods used by the authors, again suggesting good recovery of the original information. Reconstructed data has also been used in other studies of self-thinning (White and Harper 1970, White 1980, Lonsdale and Watkinson 1983a). Forty-two of the 95 EFD studies required some reconstruction from graphical data. These are indicated by the inclusion of figure numbers in the reference column of Table A.l. No reconstruction was necessary for the FYD.

Many sources gave data for stands grown under different conditions of light, fertilization, site quality, initial density, and other factors that may affect the thinning line. Data from the different conditions could be used to to estimate separate thinning lines or pooled to estimate a single line. To simplify the present analysis and preserve comparability with the original reports, the EFD were reanalyzed with the same grouping used by the original authors, except where those authors later decided that the separate thinning lines were not different. Table A.l gives the conditions that subdivided some studies into several thinning trajectories.

Thinning lines were fitted separately for different site indexes in the FYD, except where the source clearly indicated that the same relationships were used to develop the tables for different site indexes.

Log B-log N plots were examined to select points for fitting thinning lines. In many cases, the selection of points used by the original authors was not changed. However, log B-log N plots of some data sets revealed that points included in the original $\log \overline{w}$ - $\log N$ analysis were not closely associated with the thinning line and these points were removed in this analysis. Point selections were never altered when the sources gave justification for particular choices or validated their choices with field notes or mortality curves (Mohler et al. 1978, Hutchings and Budd 1981b). A complete count of the number of data sets for which the point selections were changed here is not possible because many studies did not report how many data points or which ones were used. Of 63 previously cited thinning trajectories (Table A.5), 33 indicated which points were used. Twenty-four of these were analyzed here with no changes and the point selections of the remaining 9 were altered. These studies can be identified by comparing the sample size columns of Tables A.2 and A.5.

Each edited data set was analyzed for the strength and position of linear trend. The strength was measured by calculating the Pearson correlation coefficient (Sprent and Dolby 1980), and a straight line was fit by principal component analysis (PCA)

(Jolicoeur 1973, 1975). Thinning intercepts, $\hat{\alpha}$, were calculated from $\hat{\beta}$ and the mean values of log B and log N using

$$\hat{\alpha} = \overline{\log B} - \hat{\beta} \overline{\log N} , \qquad (5.1)$$

were $\overline{\log B}$ is the average value of $\log B$ and $\overline{\log N}$ is the average of $\log N$.

The statistical significance of each correlation in the EFD was examined by testing the null hypothesis that log B and log N were uncorrelated, and 95% confidence intervals for the PCA slopes (Jolicoeur and Heusner 1971) were used to test agreement with the value of β = -1/2 predicted by the self-thinning rule. With low correlation or few data, the PCA confidence limits can be imaginary, but this is correctly interpreted as the absence of a significant relationship (Jolicoeur 1973). The 95% confidence limits for the intercept were calculated by applying equation 5.1 to the confidence limits of $\hat{\beta}$.

None of these statistical tests were applied to the FYD, because yield tables are the predictions of curve fitting procedures. The biological variability and measurement errors inherent in the original forestry data are absent in a yield table, so statistical inferences drawn from a single thinning line would be invalid and the high r^2 values (Table B.2) provide only a crude index of low variability around the fitted line. However, a large sample of FYD thinning lines can be examined to look at the statistical distributions and ranges of variation of thinning line parameters, but in all the following analyses, the EFD and FYD are

considered separately because of the different levels of statistical testing that were possible in the two groups.

Univariate statistics were computed to describe the frequency distributions of $\hat{\beta}$ and $\hat{\alpha}$. In the EFD, 55 data sets gave biomass measurements based on aboveground parts only while 20 also included roots. ANOVA and nonparametric Kruskal-Wallis tests (Sokal and Rohlf 1981) were used to compare $\hat{\beta}$ and $\hat{\alpha}$ between the two groups and determine if their statistical distributions should be examined separately.

The thinning lines were then divided into broad plant groups. Six categories were used for the EFD: herbaceous monocots, herbaceous dicots, temperate angiosperm trees, temperate gymnosperm trees, Australian trees (genus Eucalyptus), and tropical angiosperm trees (Table A.1). The FYD were divided into three categories: temperate gymnosperms, temperate angiosperms, and Eucalypts (Table B.1). ANOVA and Kruskal-Wallis tests were done to determine if $\hat{\beta}$ and $\hat{\alpha}$ differed significantly among the groups. Spearman rank correlations were also calculated for the FYD to relate $\hat{\beta}$ and $\hat{\alpha}$ to shade tolerance, a common forestry measure summarizing the ability of a species to regenerate in the presence of competition (Baker 1949, Harlow et al. 1978). These correlations were tested to determine if thinning line parameters varied significantly with shade tolerance. Finally, tables were prepared for all species examined in more than one study. For the EFD, the 95% confidence intervals of the thinning line parameters were compared to test for significant differences among the estimates.

Results

Detailed information for each thinning line, including the number of points, experimental conditions, correlation, thinning slope and intercept, confidence intervals, etc. are given in Tables A.2 and A.3 for the EFD and Table B.2 for the FYD.

Sixty-three of the thinning lines were cited in previous studies as demonstrations of the self-thinning rule (Table A.5), but did not provide strong support for the rule when tested statistically. Nineteen (30%) did not show any significant correlation between log B and log N at the 95% confidence level. The remaining 44 (70%) were significant, but the slopes of 20 (32%) were statistically different from the thinning rule prediction β = -1/2, with 7 (11%) greater than and 13 (21%) less than -1/2. Only 24 (38%) of the slopes were both statistically significant and not different from the predicted value. Thirty-one additional thinning lines from data not previously analyzed in a self-thinning study also showed a significant log B-log N correlation. When combined with the 44 from the previously cited group, this gave at total of 75 significant thinning lines. Of these 75, 34 (45%) were not statistically different from -1/2. The remaining 41 (55%) were different, with 14 (19%) greater than and 27 (36%) less than -1/2.

Univariate statistics for the slopes and intercepts of these 75 thinning lines are presented for two groups: data sets with biomass measurements of aboveground parts only and data sets with biomass measurements that included roots (Table 5.1). ANOVA

Table 5.1. Statistical Distributions of Thinning Line Slope and Intercept.

	Shoot Bi	Forestry Table Bole Bi	Data omass					
Statistic	Ŗ	â	β	â	ß	â	β̂	â
n	55	55	20	20	75	75	351	351
Mean	-0.863	4.08	-0.804	4.48	-0.847	4.18	-0.876	3.45
Std. dev.	0.701	1.12	0.862	1.62	0.742	1.27	1.024	1.04
CV(%)	81	27	107	36	88	30	117	30
Std. err.	0.095	0.15	0.193	0.36	0.086	0.15	0.055	0.06
Skewness	-2.57	0.33	-2.59	1.89	-2.51	1.25	-4.36	-4.30
Kurtosis	7.14	2.03	7.21	6.86	6.59	5.64	21.04	23.45
Range	3.662	6.27	3.56	8.53	3.66	8.65	8.014	8.60
Percentiles								
0 (Min.)	-3.808	1.28	-3.760	1.40	-3.808	1.28	-8.132	-4.18
1	*	*	*	*	*	*	-6.641	-2.53
5	-2.838	1.64	-3.687	1.48	-2.838	1.65	-2.411	-1.90
10	-1.589	2.94	-2.207	3.15	-1.589	3.07	-1.268	2.74
25	-0.986	3.65	-0.951	3.89	-0.986	3.70	-0.848	3.33
50 (Median)	-0.649	3.84	-0.468	4.13	-0.622	3.97	-0.618	3.68
75	-0.503	4.63	-0.332	4.91	-0.465	4.63	-0.480	3.95
90	-0.316	5.37	-0.226	6.31	-0.299	5.58	- 0.370	4.12
95	-0.249	6.53	-0.205	9.75	-0.220	6.53	-0.318	4.27
99	*	*	*	*	*	*	-0.222	4.39
100 (Max.)	-0.146	7.54	-0.204	9.93	-0.146	9.93	-0.119	4.42

aThe three categories under this heading are based on the method of biomass measurement. "Shoot Biomass" includes thinning trajectories from biomass measurements of aboveground parts only. "Total Biomass" trajectories are based on biomass measurements that include aboveground and belowground parts. "All Data Sets" gives statistics for both of these groups combined.

^{*}Indicates that the 1st (or 99^{th}) percentile value is identical to the minimum (or maximum) due to small sample size.

detected no significant differences between the two group means for both $\hat{\beta}$ (F_{1.73} = 0.09, P = 0.76) and for $\hat{\alpha}$ (F_{1,73} = 1.44, P = 0.23). Lack of significance was also obtained in the Kruskal-Wallis tests for $\hat{\beta}$ (H₂ = 3.02, P = 0.08) and $\hat{\alpha}$ (H₂ = 1.90, P = 0.16), so the two groups were pooled and their composite statistical distribution is also given. Table 5.1 also includes analogous statistics for the 351 thinning lines fitted to data sets in the FYD. Figure 5.1 gives histograms for $\hat{\beta}$ and $\hat{\alpha}$. The distributions were also compared to the accepted ranges of variation $-0.8 \le \hat{\beta} \le -0.3$ and $3.5 \le \hat{\alpha} \le 4.4$ (White 1980). For the EFD, 24 (32%) of the values of $\hat{\beta}$ were outside this range, with 22 (29%) less than -0.8 and 2 (3%) greater than -0.3. Thirty-four (46%) $\hat{\alpha}$ values were more extreme, with 14 (19%) less than 3.5 and 20 (27%) greater than 4.5. Among the FYD, 105 (30%) $\hat{\beta}$ values were more extreme, with 102 (29%) less than -0.8 and 3 (1%) greater than -0.3. Of the FYD $\hat{\alpha}$ values, 124 (36%) were more extreme with 122 (35%) less than 3.5 and 2 (1%) greater than 4.4.

Comparisons among broad plant groups showed that thinning parameters are not constant across the entire plant kingdom. ANOVA and Kruskal-Wallis tests detected significant differences in both $\hat{\beta}$ and $\hat{\alpha}$ among the six groups in the EFD and among the three groups of the FYD (Table 5.2). Because the Eucalypt group of the FYD was so extreme, two-way comparisons between the angiosperm and gymnosperm groups were also done. The differences in $\hat{\beta}$ between these two more similar groups were also statistically significant. For $\hat{\alpha}$, the ANOVA detected no significant differences between the




Figure 5.1. Histograms for the slopes and intercepts of fitted thinning lines. (a) and (b) show the distributions of slope and intercept, respectively, for log B-log N thinning lines in the experimental and field data. (c) and (d) show the same distributions for thinning lines in the forestry yield table data.

Table 5.2. Comparisons of Thinning Line Slope and Intercept Among Plant Groups.

Group	n	S1 Mean	ope β Median	Interc Mean	ept & Median
Experiment	: <u>al a</u>	nd Field	Data		
Herbaceous monocots Herbaceous dicots Temperate angiosperm trees Temperate gymnosperm trees <u>Eucalyptus</u> trees Tropical angiosperm trees All of the above	8 25 15 19 4 4 75		-0.65 -0.53 -0.65 -1.03	4.45 5.17 3.78 3.79 2.87 2.20 4.18	4.24 5.09 3.72 3.88 3.07 2.21 3.97
Tests for Significant		erences	Among Si	x Groups	
ANOVA Â Kruskal-Wallis Â		F(5,69) F(5,69) H(5) = 1 H(5) = 4	= 10.9 7.9	P < 0. P < 0. P = 0. P < 0.	.0001* .0031*
Forestry	<u>Yiel</u>	d <u>Table</u>	<u>Data</u>		
Temperate angiosperm trees Temperate gymnosperm trees Eucalyptus trees All temperate trees All of the above	58 281 12 339 351	-0.80 -3.90	-0.61 -4.39 -0.61	3.54 1.09	3.56 3.72 1.79 3.68 3.68
Tests for Differences Be	tween	Gymnosp	erms and	Angiospe	rms
ANOVA É Kruskal-Wallis	<u>.</u>			P = 0 P = 0 P = 0 P = 0	.8 .052
Tests for Differ	ence	s Among	Three Gro	oups	
ANOVA É Kruskal-Wallis	\$ & \$.	F(2,348 H(2) =) = 38.9 14.9		.0001* .0006*
Ĝ	i	H(2) =	11.9	P = 0	.0027*

^{*}Significant at the 95% confidence level (P \leq 0.05)

two groups while the Kruskal-Wallis analysis did. This indicates that the mean values of $\hat{\alpha}$ are similar, but one of the groups has a disproportionate of number extreme values of $\hat{\alpha}$ (in this case, the gymnosperm group has more high values), causing the rank based Kruskal-Wallis test to detect a difference.

Thinning slope and intercept were both correlated with shade tolerance in the FYD. The correlation analysis was done separately for the angiosperms and gymnosperms because of the observed differences in $\hat{\beta}$ and $\hat{\alpha}$ between these two groups. The Eucalypt group was not analyzed because shade tolerances were not available. For the 46 angiosperm thinning trajectories analyzed, $\hat{\beta}$ was significantly correlated with shade tolerance but thinning intercept was not (Table 5.3). Both $\hat{\beta}$ and $\hat{\alpha}$ were significantly correlated with tolerance in the gymnosperms; however, the sign of the correlation of $\hat{\beta}$ with shade tolerance was opposite to that observed for the angiosperms, further justifying the separate analyses of the two groups.

Thinning line parameters were not constant for species considered in more than one study. Both $\hat{\beta}$ and $\hat{\alpha}$ show considerable variation within species in the EFD (Table 5.4). In some cases, parameter estimates for a given species are quite different, but the confidence intervals for the estimates are so large that the differences are not statistically significant. In other cases, the differences are statistically significant, as indicated by non-overlap of the 95% confidence intervals. Eight (20%) of the 40 possible pairwise within-species comparisons of $\hat{\beta}$

Table 5.3. Spearman Correlations of Shade Tolerance with Thinning Line Slope and Intercept in the Forestry Yield Data.

T	Means	for Sha		rman lation											
Thinning Parameter	1	2	3	4	5	r_S	pb								
	Temperate Angiosperms														
Slope β̂		-0.547 (18)				-0.52* (46)	0.0002								
Intercept \hat{lpha}	3.632 (10)	3.437 (18)				-0.19 (46)	0.22								
		Tempe	rate Gy	mnosper	ms										
Slope β̂		-0.748 (78)				0.35 * (267)	<0.0001								
Intercept $\hat{\alpha}$	3.123 (32)	3.438 (78)				0.57 * (267)	<0.0001								

 $^{{\}tt aSample}$ sizes are given in parentheses.

 $^{{}^{\}rm b}{\rm Statistical}$ significance level.

^{*}Significant at the 95% confidence level (P \leq 0.05).

Table 5.4. Thinning Line Slopes and Intercepts of Species for which Several Thinning Lines Were Fit from Experimental or Field Data.

				Thinning Slope	Line Para	Intercept					
Id. Code ^b	Condition ^C	r ²	β̂	95% CI	Diff.d	â	95% CI	Diff.d			
· · · · · · · · · · · · · · · · · · ·				Abies sachalinens	i <u>s</u>						
A 19A B 24T C 119A		0.841 0.839 0.971	-0.649 -0.465 -2.786	[-0.776,-0.535]* [-0.965,-0.104] [-5.645,-1.772]*	C	4.16 4.39 1.71	[4.15, 4.17 [4.07, 4.83 [-0.49, 2.49]	j c			
				Beta vulgaris							
A 33T B 34T C 43A D 43A E 43A F 43A G 43A	18% L.I. 25% L.I. 37% L.I. 55% L.I. 100% L.I.	0.592 0.645 0.957 0.916 0.940 0.934 0.698	-1.335 -2.304 -0.662 -0.692 -0.668 -0.649 -0.648	[-3.355,-0.648]* [-5.478,-1.348]* [-0.839,-0.509]* [-0.973,-0.470] [-0.886,-0.486] [-0.838,-0.487] [-1.415,-0.197]	CDEF	6.38 9.93 4.79 5.12 5.09 5.22 5.30	[4.54, 11.80 [7.08, 19.38 [4.17, 5.50 [4.22, 6.25 [4.39, 5.94 [4.59, 5.95 [3.55, 8.27	CDEF B B B B			

=

Table 5.4. (continued)

					Thinning Slope	Line Para	ameters ^d	Interc	ept	
	d. de ^b	Condition ^C	r ²	ĝ	95% CI	Diff.d	â	95%	CI	Diff.d
					<u>Er</u>	igeron ca	nadensi	<u>s</u>		
A B	15T 21T		0.930 0.987	-0.621 -1.038	[-0.688,-0.558]* [-1.121,-0.962]*	B A	4.36 5.70	[4.19, [5.43,	4.55] 6.00]	B A
					<u>Eu</u>	calyptus	regnans	<u>'-</u>		
A B	98A 98A	S.I. 28.9 S.I. 33.5	0.971 0.964		[-5.012,-1.559]* [-2.132,-0.549]*		1.39 3.44	[-1.63, [2.31,		
						Lolium pe	erenne			
A B C D E	38A 91A 91T 92A 92A	100% L.I. 100% L.I. 100% L.I. 23% L.I. 44% L.I.	0.549 0.908 0.854 0.776 0.786		[-0.674,-0.034] [-0.543,-0.319] [-0.330,-0.163]* [-1.509,-0.011] [-1.273,-0.027]		3.79 4.80 4.20 4.33 4.28	[2.88, [4.37, [3.87, [2.28, [2.48,	4.54]	

_

Table 5.4. (continued)

. .				Thinning Slope	MICCEL 3		ntercept		
Id. Code ^b	Condition ^C	r ²	β̂	95% CI	Diff.d	â	95% CI [Diff.d
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,					Picea a	bies_	to the terminal to the termina	· · · · · · · · · · · · · · · · · · ·	,
A 137A B 137T				[-0.462,-0.383]* [-0.476,-0.392]*		3.90 3.97	[3.88, [3.95,	3.92] 3.99]	
					<u>Pinus</u> st	robus			
A 8A B 8A C 93A	Lot 2B Lot 2C	0.986 0.987 0.955	-0.724 -1.116 -0.954	[-0.830,-0.628]* [-1.278,-0.976]* [-1.189,-0.764]*	В	3.78 3.34 3.44	[3.74, [3.22, [3.25,	3.43]	
					<u>Pinus</u> t	aeda			
A 82A B 102A		0.468 0.939		[-0.499,-0.130]* [-0.837,-0.526]*			[4.11, [3.23,		

Table 5.4. (continued)

					Thinning Line Parameters ^a									
I Co	d. de ^b	Condition ^C	r ²	β	Slope 95% CI	Diff.d	â	Inter		Diff.d				
	<u> </u>		,		<u>Tri</u>	folium sub	terrane	um		* , <u>, , , , , , , , , , , , , , , , , ,</u>				
A B	10A 35A	Full light	0.836 0.716	-0.473 -0.622	[-0.660,-0.310] [-0.928,-0.382]		4.60 5.17	[3.97, [4.33,	5.33] 6.23					

and a_{0} species for which more than one statistically significant thinning line (P \leq 0.05) were fit are included (see Table A.2 for actual significance levels).

 b Table A.1 associates the numeric part of each Id. code with a particular species and study. The letter indicates the type of biomass measurements made: A = aboveground parts only, T = aboveground and belowground parts both included.

 ${}^{\text{C}}\text{See}$ Table A.1 and the references given there for more information on condition.

dThe letters in this column identify table entries (see letters at left of table) for the same species that are significantly different from the current estimate of $\hat{\beta}$ or $\hat{\alpha}$. Statistical disagreement (at P \leq 0.05) between estimates is indicated by non-overlap of confidence intervals.

*Indicates slopes that are significantly different (P \leq 0.05) from the value β = -1/2 predicted by the self-thinning rule.

were significant, as were 9 (23%) of the 40 possible comparisons of $\hat{\alpha}$. Since the confidence level is 95%, 2 of 40 comparisons should be different by chance alone, but both analysis gave a least 4 times this number. For the FYD, individual thinning line estimates (Table B.2) were not retabulated because so many thinning trajectories (264) were involved and because 95% confidence intervals could not be calculated. Instead, the sample sizes, means, standard deviations, coefficients of variation, minima, maxima, and ranges are given for $\hat{\beta}$ and $\hat{\alpha}$ for each species (Table 5.5). Although significance tests were not possible, the large observed ranges again indicate that $\hat{\beta}$ and $\hat{\alpha}$ are not species constants.

Discussion

The predictions of the self-thinning rule have been tested here through five different analyses: (1) statistical tests of the hypothesis that the slope of the self-thinning line is β = -1/2, (2) examination of the frequency distributions and ranges of variation of thinning line slope and intercept, $\hat{\alpha}$, (3) tests for variation in $\hat{\beta}$ and $\hat{\alpha}$ among plant groups, (4) tests for correlations of $\hat{\beta}$ and $\hat{\alpha}$ with shade tolerance, and (5) examination of the variation in $\hat{\alpha}$ and $\hat{\beta}$ for particular species. Although many data sets do show the predicted region of linear association between log B and log N, all five analyses indicate that the slope of this relationship does not take the same value β = -1/2 across the entire plant kingdom. Therefore, the evidence

Table 5.5. Statistics for Thinning Line Slope and Intercept of Species for which Several Thinning Lines Were Fit from Forestry Yield Data.

	Number		Thinning Slope β							Thinning Intercept $\hat{\alpha}$						
Species ^a	Yield Tables	Thin. Lines	Mean	SDc	CAq	Min.	Max.	Range	Mean	SDc	CAq	Min.	Max.	Range		
	***************************************		Sp	ecies	Consid	ered in	More T	han <u>One</u>	Yield T	able						
Alnus rubra	3	8	-0.41	0.19	46	-0.65	-0.12	0.53	3.46	0.42	12	2.69	4.08	1.39		
Picea glauca	2	9	-0.75	0.30	39	-1.28	-0. 52	0.76	3.59	0.31	9	3.01	3.83	0.82		
Pinus banksiana	3	10	-0.79	0.39	50	-1.73	-0.24	1.49	3.06	0.27	9	2.51	3.51	1.00		
Pinus echinata	4	20	-0.72	0.23	32	-0.97	-0.40	0.56	3.53	0.35	10	3.14	4.05	0.91		
Pinus elliotti	2	1]	-0.56	0.16	28	-0.83	-0.38	0.45	3.62	0.34	9	3.22	3.99	0.77		
Pinus palustris	2	14	-1.04	0.22	21	-1.28	-0.80	0.48	3.01	0.49	16	2.20	3.52	1.32		
Pinus ponderosa	3	27	-0.72	0.66	92	-2.44	-0.31	2.13	3.36	0.90	27	1.08	4.18	3.10		
Pinus resinosa	2	8	-1.07	0.48	45	-1.97	-0.63	1.34	3.46	0.47	14	2.73	4.21	1.48		
Pinus strobus	3	12	-0.74	0.18	25	-1.07	-0.5 3	0.54	3.68	0.15	4	3.43	3.88	0.45		
Pinus taeda	3	17	-0.69	0.20	28	-0.90	-0.25	0.65	3.48	0.34	10	3.05	4.10	1.05		
Pseudotsuga menziesii	3	20	-0.60	0.15	25	-0.71	-0.26	0.45	3.80	0.22	6	3.59	4.26	0.67		
Sequoia sempervirens	2	10	-2.68	1.45	54	-4. 75	-1.22	3.53	1.20	2.24	186	-1.93	3.44	5.37		
Tsuga neterophylla	4	37	-0.44	0.08	18	-0.61	-0.36	0.25	4.21	0.14	3	3.93	4.42	0.48		

Table 5.5. (continued)

	Number			Thinning Slope β						Thinning Intercept \hat{lpha}					
_	Yield Tables	Thin. Lines	Mean	SDc	CAq	Min.	Max.	Range	Mean	SDc	cvd	Min.	Max.	Range	
		· · · · · · · · · · · · · · · · · ·	Sp	ecies	Consid	ered in	More T	han <u>One</u>	Yield Ta	able	* - 16 - 17 - 17 - 17	dr 81111			
Abies balsamea	1	4	-0.59	0.00	1	-0.60	-0.59	0.01	3.80	0.02	1	3.77	3.81	0.05	
Abies concolor	1	7	-0.57	0.01	2	-0.59	-0.56	0.03	3.84	0.13	4	3.67	3.98	0.31	
Castanea dentata	1	3	-0.65	0.01	2	-0.66	-0.64	0.02	3.62	0.06	2	3.56	3.67	0.11	
Chamaecyparis thyoides	1	6	-0.53	0.01	2	- 0.54	-0.51	0.03	3.80	0.14	4	3.53	3.89	0.36	
Eucalyptus globus	1 .	4	-5. 80	1.86	32	-8.13	-3.80	4.33	1.06	0.98	93	-0.21	2.06	2.27	
Eucalyptus microtheca	1	3	-6.84	0.39	6	-7. 22	-6.44	0.78	-3.67	0.51	14	-4.18	- 3.17	1.01	
Eucalyptus sieberi	1	3	-0.62	0.07	12	-0.69	-0.55	0.15	4.12	0.03	1	4.09	4.14	0.06	
<u>Liriodendron</u> tulipifer		3	-1.15	0.12	10	- 1.24	-1. 02	0.23	2.69	0.18	7	2.53	2.88	0.35	
<u>Liquidambar</u> styraciflu	<u>a</u>]	6	-0.39	0.03	8	-0.42	- 0.33	0.09	3.87	0.11	3	3.74	4.03	0.30	
<u>Picea mariana</u>]	3	-0.83	0.04	5	-0.85	-0. 78	0.07	3.72	0.01	0	3.72	3.72	0.01	
<u>Picea</u> <u>rubrens</u>	1	5	-0.54	0.00	1	- 0.55	-0.54	0.01	3.86	0.10	3	3 .7 2	3.97	0.25	
<u>Pinus</u> <u>monticola</u>	1	4	-0.79	0.02	2	-0. 80	-0.77	0.04	3.88	0.11	3	3.75	4.01	0.25	
Populus tremuloides	1	4	-0.33	0.07	23	-0.43	-0.27	0.16	3.62	0.12	3	3.46	3.74	0.28	
Thuja occidentalis	1	6	- 0.35	0.04	11	-0.41	-0.30	0.11	3.58	0.08	2	3.46	3.67	0.21	

^aOnly results from monsopecific yield tables are included (Tables B.1 and B.2).

bGives the number of yield tables for the species and the total number of thinning lines fit.

CStandard deviation.

dCoefficient of variation expressed as a percentage.

does not support the hypothesis that all plants obey the same quantitative self-thinning rule. These analyses also confirm predictions about the variations in thinning parameters from previous studies and from the models presented here.

The statistical tests of the 63 thinning lines reported to demonstrate the self-thinning rule show that this body of evidence does not strongly support the rule. Many of the reported high correlations between log W and log N proved spurious when the log B-log N data were reanalyzed, and 30% of the thinning lines did not even show a significant relationship between biomass and density. There could be two basic explanations for this result: the data are too variable or too few to detect the existence of a true relationship between log B and log N, or there really is no relationship, as when stand biomass is maintained at a carrying capacity. Either way, the data do not support the self-thinning rule. Seventy percent of the thinning lines did show the predicted significant linear relationship between log B and log N, but 32% of the thinning slopes were significantly different from $\beta = -1/2$ and disagreed quantitatively with the thinning rule. Deviations of the thinning slope from the predicted value are particularly important because β is the exponent of a power relationship, $B = K N^{\beta}$ so small differences in β represent large differences in the predictions of the equation (Chapter 4). Only 38% of the thinning slopes were both significantly different from 0 and not significantly different from $\beta = -1/2$. Because of the inherent ambiguities of statistical testing, these represent two

possibilities: a true relationship of slope β = -1/2 is present, or some different true slope is present but the data are so variable as to obscure this difference. In fact, many data sets were too variable to be useful in resolving alternative hypothesis about the slope of the thinning line. Strictly speaking, the most that can be claimed is that these 38% of the thinning lines do represent statistically significant relationships between log B and log N and the slopes are close to β = -1/2 within the resolution of the data.

In short, 30% of the data sets did not demonstrate <u>any</u> relationship between biomass and density while another 32% disagreed quantitatively with the -1/2 slope predicted by the thinning rule. This gives 62% of the data sets that were either useless for testing the rule or in quantitative disagreement with it, and only 38% that potentially support for the rule.

These tests must be interpreted with an important caveat: the true confidence level of each test is less than the nominal 95% because the necessary step of editing the data to fit the thinning line increases the probability of a Type I statistical error (rejection of a true null hypothesis) by some unknown amount. Some additional percentage of the thinning lines actually showed no correlation between log B and log N, while some slopes that tested as significantly different from β = -1/2 are actually different at some lower confidence level. At present, all we can do is acknowledge and decry this limitation and argue that the analysis is still the most objective that is currently possible. Since the

other four analyses also show that thinning slopes vary widely from $\beta = -1/2$, confidence in the test results seems justified.

The frequency distributions of thinning slope and intercept show greater variations in these parameters than suggested by the often cited ranges $-0.8 \le \hat{\beta} \le -0.3$ and $3.5 \le \hat{\alpha} \le 4.4$ proposed by White (1980). Both distributions have single modes near their accepted values of $\beta = -1/2$ and $\alpha = 4$, but more extreme values are also present. The tendencies for $\hat{\beta}$ to be near -1/2 and for $\hat{\alpha}$ to be near 4 reflect the fact that values near the mode of a distribution are more frequently observed than extreme values. The commonness of modal values does not indicate that a limited range is rigidly imposed by a biological law. Seventy percent of the 426 $\hat{\beta}$ values in the combined EFD and FYD were within White's range, as were 42% of the $\hat{\alpha}$ values, so these ranges do include a large percentage of the observations, but not all. The greater variation observed here has a simple explanation, White's ranges were based on 36 data sets collected precisely because their thinning slopes were close to $\beta = -1/2$, but attempts were made here to avoid prejudicing the analysis with this criterion.

The observed departures of the thinning slope from β = -1/2 for trees in both the EFD and FYD support Sprugel's (1984) speculation that a thinning slope of γ = -3/2 (β = -1/2) is the exception rather than the rule for woody plants. Since herbaceous species also showed such departures, Sprugel's prediction applies for all plants. White (1981) has suggested that thinning slopes steeper than γ = -2 (β = -1) are prima facie evidence of

significant departure from the classic thinning rule. Seventeen (23%) of the EFD and 45 (13%) of the FYD slopes meet this criterion.

Additional evidence of variation in thinning line parameters was found in the existence of statistically significant differences in thinning line parameters among plant groups, and in the observation of significant correlations between thinning line parameters and shade tolerance (as predicted by Westoby and Howell 1981, Lonsdale and Watkinson 1983a). These results argue against a single, quantitative thinning rule for all plants (as claimed by White 1981, Hutchings 1983). Thinning line slope and intercept were not even constant among thinning lines for a particular species, so these parameters are not species constants (as suggested by Mohler et al. 1978, Hozumi 1980, White 1981, Hutchings 1983). This result complements evidence that thinning slope and intercept can vary within a single yield table when data from different site indexes are compared (Chapter 4). This has also been reported by Hara (1984), and Furnas (1981) has shown experimentally that the position of a thinning line can respond to changes in nutrient availability. Unfortunately, the biological interpretation of α is confounded because thinning slopes are variable. The constants of power equations have direct biological interpretations only if the powers of the relationships are identical (White and Gould 1965).

The existence of variation in thinning slope and intercept supports the the models of Chapters 2 and 3, which predicted that the slope of the thinning line depends on how the shape of the space occupied by a plant changes with growth, while the intercept is

additionally affected by the density of biomass in occupied space and the way interacting plants partition areas of overlap. If these models are correct, then thinning slope should vary among species because different species have different shapes and densities of biomass per unit of space (Mohler et al. 1978, Furnas 1981, Lonsdale and Watkinson 1983a). Thinning parameters should also vary within a species because plants change shape and canopy density in response to growing conditions (Harper 1977). Between-species and within-species variations are both present in the data analyzed here, just as predicted. The observed correlations of thinning line parameters with shade tolerance also confirm the importance of differences in allowable overlap between plants in positioning the thinning line.

CHAPTER 6

SELF-THINNING AND PLANT ALLOMETRY

Introduction

The documentation of significant variation among the slopes of self-thinning lines (Chapter 5) verifies one major prediction of the mathematical model (Chapter 2), but the true test of the model comes in determining if the variations in self-thinning slope can be related to differences in plant allometry. Here three questions are addressed that provide such tests: (1) Are thinning slope and intercept correlated with available measures of plant allometry? (2) Do differences in thinning slope among plant groups (Chapter 5) reflect corresponding differences in plant allometry? (3) Do the correlations of thinning slope with shade tolerance (Chapter 5) reflect corresponding correlations between shade tolerance and plant allometries? The results and conclusions are related to those of previous studies to explain why allometric models have been discounted (Westoby 1976, Mohler et al. 1978, White 1981).

Expected Relationships between Thinning Slope and Plant Allometry

The model predicts that the slope of the self-thinning line in the log B-log N plane is $\beta = -1/[2p] + 1$ (or equivalently $\gamma = -1/[2p]$ in the log \bar{w} -log N plane), where p is the allometric power relating the radius of the zone of influence (ZOI) to weight according to the power equation R $\propto w^p$ (Chapter 2). This

Prediction is referred to here as the allometric hypothesis. Hypothetically, it could be tested by measuring both $\hat{\beta}$ and \hat{p} for several populations and testing statistically to determine if $\hat{\beta}$ and \hat{p} follow the expected relationship. However, the actual measurement of ZOIs is problematic. It is difficult to define precisely where the ZOI of an individual ends, especially in highly competitive situations where the zones may be tightly packed and overlapping to varying degrees. Even when the ZOI is defined as the canopy area (White 1981), the actual measurement is still difficult and less accurate than other common measurements, such as height or bole diameter at breast height (DBH). Consequently, few data are available for relating the ZOI to weight and no published data were found where both \hat{p} and $\hat{\beta}$ could be estimated simultaneously.

Plant ecologists and foresters routinely measure some parameters of plant shape, such as height, DBH, or bole basal area (BSLA), that can be used to fit allometric equations. Thinning slope can be related to these other measures of plant allometry. For example, the allometric equation $\bar{h} \propto \bar{w}^{\Phi h W}$ could be fit to average height and average weight measurements of the same stands used to estimate a self-thinning slope. Values of $\hat{\Phi}_{hw}$ for several species could then be correlated with $\hat{\beta}$ to test if the two parameters are significantly related.

Geometric models can be further analyzed to suggest a functional form for the β - Φ_{hw} relationship. Assume that plant height, the density of biomass in occupied space, and the ZOI radius all vary with plant weight according to allometric power functions

h \propto w^{Φ hw}, d \propto w^{Φ dw}, and R \propto w^P. If the volume of space occupied by a plant (VOI) is approximately cylindrical, then the volume is $v = \pi R^2$ h and plant weight is $w = v d = \pi R^2$ h d. Since $w \propto R^2$ h d and R, h, and d are allometrically related to weight, the two sets of equations can be combined to give $w \propto w^{2p} w^{\Phi hw} w^{\Phi dw}$ and the allometric powers are constrained by

$$2p + \Phi_{hw} + \Phi_{dw} = 1$$
 (6.1)

The predicted relationship $\beta = -1/(2p) + 1$ can be solved for p to give

$$P_{\text{est}} = \frac{1}{2(1-\beta)}$$
, (6.2)

where the symbol p_{est} emphasizes that p is not directly measured but derived by mathematical transformation of measured $\hat{\beta}$ values. Combining equations 6.1 and 6.2 gives a relationship between p_{est} , Φ_{hw} , and Φ_{dw} that should obtain if the allometric hypothesis is true

$$P_{\text{est}} = \frac{1}{2} \left[1 - (\phi_{\text{hw}} + \phi_{\text{dw}}) \right] . \tag{6.3}$$

This relationship is useful because of its linear form, simple geometric derivation, and clear representation of the compromises inherent in plant growth: allocation of more resources to height growth (higher Φ_{hw}) or to packing more biomass in space already occupied (higher Φ_{dw}) leaves fewer resources for expanding the

ZOI radially (lower p). Less radial expansion in turn means less conflict with neighbors, less self-thinning, and a steeper (more negatively sloped) thinning line.

To focus on the compromises between height growth and radial growth, assume that d is fairly constant ($\Phi_{\rm dw}$ = 0). Equation 6.3 simplifies to

$$P_{\text{est}} = -\frac{1}{2} \Phi_{\text{hw}} + \frac{1}{2} ,$$
 (6.4)

which defines a triangular region in the p_{est} - Φ_{hw} plane that is bounded by the two axes p_{est} = 0 and Φ_{hw} = 0 and the by the line p_{est} = -0.5 Φ_{hw} + 0.5. Empirical measurements of p_{est} and $\hat{\Phi}_{hw}$ would lie around this line, except that Φ_{dw} is not generally zero because the density of biomass in occupied space does vary with plant size (Lonsdale and Watkinson 1983a). Since measurements of d through time are not generally available for estimating $\hat{\Phi}_{dw}$, equation 6.3 can not be used and variation in Φ_{dw} will contribute to the errors in the simpler model of equation 6.4. Ignoring Φ_{dw} and fitting a linear relationship between p and $\hat{\Phi}_{hw}$ will produce a line in the p_{est} - Φ_{hw} plane below equation 6.4; however, a negative correlation between p_{est} and $\hat{\Phi}_{hw}$ will still support the allometric hypothesis, particularly if the functional form of the relationship approximates equation 6.4.

Other measurements besides height are commonly available for trees and can be used to fit allometric relationships to weight $(\overline{DBH} \propto \overline{w}^{\Phi Dw} \text{ or } \overline{BSLA} \propto \overline{w}^{\Phi Bw}) \text{ that can be related to } \beta. \text{ However,}$

DBH and BSLA are measures of the tree bole, not the VOI, so the expected relationship of $\Phi_{\mbox{\rm DW}}$ or $\Phi_{\mbox{\rm RW}}$ can not be simply derived from the geometry of the VOI. Also, mechanical considerations require that bole diameter increase to support any increase in tree weight, whether the growth is upward, radial, or simply an increase in the density of biomass in the VOI. Hence the DBH-weight or BSLA-weight allometries should not be be sensitive measures of change in the VOI shape and should not necessarily correlate strongly with p_{est} . DBH and height data can be combined to fit another allometric relation $\overline{h} \propto \overline{DBH}^{\Phi hD}$. Although interpretation of Φ_{hD} is again confounded because DBH is not a measure of the VOI, h is a VOI dimension so some expectations for the $p_{\rm est}$ - $\Phi_{\rm hD}$ relationship can be deduced geometrically. If the VOI expands only radially (p = 0.5 and Φ_{hw} = 0 in equation 6.4), then Φ_{hD} will be zero since height is constant. If the VOI expands both radially and upward (p < 0.5 and $\Phi_{\rm hw}$ > 0), then DBH will increase to support the additional weight and $\Phi_{\mbox{\scriptsize hD}}$ will be positive. Thus, a line representing the relationship in a p- $\Phi_{h\bar{l}l}$ plot would pass through the point $(\Phi_{h\Omega}, p) = (0,0.5)$ and would be negatively sloped. observed negative correlation between p_{est} and $\hat{\Phi}_{hD}$ would, then, support the allometric hypothesis. Furthermore, BSLA is simply related to DBH by BSLA \propto DBH², so the above logic also applies for the relationship between $p_{\rm est}$ and $\Phi_{\rm hR}$. Although the functional form of the relationship of $p_{\rm est}$ to $\Phi_{\rm hD}$ or $\Phi_{\mbox{\scriptsize hB}}$ is not geometrically obvious, it is reasonable to use a linear model unless this assumption proves inappropriate.

Methods

Information to test the allometric hypothesis was available in the sources of experimental and field data (EFD) and forestry yield table data (FYD) analyzed in Chapter 5 since average stand height, DBH, or BSLA were often reported along with stand biomass and density. All height and DBH measurements were converted to common units of m, while basal areas were converted to m² and divided by the density of individuals to give average basal area per tree. Many studies did not measure one or more of these dimensions, so fewer data were available for estimating allometric relationships. Of the 75 data sets in the EFD that showed significant relationships between log B and log N (Chapter 5), 31 had accompanying measurements of stand height, 8 reported DBH, and 29 reported BSLA. Of the 351 EFD thinning lines, 325 had height measurements, 334 had DBH, and 318 had BSLA.

Dimensional measurements from the same stands used to estimate $\hat{\beta}$ were log transformed and analyzed with principal component analysis (PCA, Chapter 5) to fit allometric equations relating log \bar{h} , log $\bar{D}BH$, and log $\bar{B}SLA$ to log \bar{w} ; log \bar{h} to log $\bar{D}BH$; and log \bar{h} to log $\bar{B}SLA$. For each proposed allometric relationship in the EFD, the null hypothesis that the two variables were uncorrelated was tested and only relationships significant at the 95% confidence level were retained for further analysis. Similar statistical tests for the FYD were not possible (see Chapter 5) so the EFD and FYD were again analyzed separately.

Values of $\hat{\beta}$ (Chapter 5) were transformed to p_{est} values and correlated with the five allometric powers using both Spearman and Pearson correlation calculations. The correlation coefficients were tested and when a significant relationship was found, linear regression was used to estimate an equation for the relationship. Principal component analysis was not used because the objective was to predict p_{est} from an independent allometric parameter rather than to estimate a functional relationship between two variables. With this objective, regression is the appropriate technique despite errors in the independent variable (Ricker 1973, Sokal and Rohlf 1982).

Univariate descriptive statistics and histograms were calculated for each allometric parameter, and ANOVA and Kruskal-Wallis tests of differences among plant groups were performed. Spearman correlations of shade tolerance with each allometric power for the temperate angiosperm and temperate gymnosperm groups of the FYD were also calculated. Additional information on the groups and methods is given in Chapter 5.

Results

Table A.3 presents allometric powers fitted to data sets in the EFD. The PCA slope and intercept of each allometric equation are given, along with the correlation and statistical significance of the relationship. For the five allometric powers, $\hat{\Phi}_{hw}$, $\hat{\Phi}_{Dw}$, $\hat{\Phi}_{Bw}$, $\hat{\Phi}_{hD}$, and $\hat{\Phi}_{hB}$, there were 23, 8, 23, 6, and 19, respectively, statistically significant relationships. Table B.3

gives the results of allometric regressions for the forestry yield data. Values of $p_{\rm est}$ calculated from fitted thinning slopes (Chapter 5) are included in Tables A.2 and B.2. Tables 6.1 and 6.2 give univariate statistics for $p_{\rm est}$ and the five allometric powers in the EFD and FYD, while Figures 6.1 and 6.2 present the corresponding histograms.

Two of the correlations between the transformed thinning slope and the allometric powers of the EFD were statistically significant (P < 0.05--Table 6.3, Figure 6.3). Values of $\hat{\Phi}_{hw}$ were negatively correlated with p_{est} (r = -0.55, P < 0.0026) and 30% of the observed variation in p_{est} was explained by the regression $p_{est} = -0.710 \; \hat{\Phi}_{hw} + 0.501$. This equation does not differ significantly in either slope or intercept from the predicted equation $p = -0.5 \; \Phi_{hw} + 0.5$. The failure of $\hat{\Phi}_{Dw}$ and $\hat{\Phi}_{hD}$ to correlate significantly with p_{est} is inconclusive because of the small sample sizes (n \leq 8). The allometric power $\hat{\Phi}_{Bw}$ also showed no significant correlation with p_{est} , despite a larger sample size, but BSLA was not not expected to be a good indicator of thinning slope. The allometry $\hat{\Phi}_{hB}$ showed a statistically significant correlation with p_{est} (r = -0.44, P = 0.032) and the regression explained 19% of the variation in p_{est} .

All five correlations between p_{est} and the allometric powers in the FYD were significant (P < 0.0001, Table 6.3), but the high confidence levels were partly due to the large sample sizes (n > 309). The coefficients of determination give a more realistic assessment of the ability of these allometric parameters

Table 6.1. Statistical Distributions of Allometric Powers for Experimental and Field Data.

	Allometric Parameter										
Statistic	p _{est}	$\hat{^\Phi}$ hw	$\hat{\Phi}_{DW}$	${}^{\hat{\Phi}}_{Bw}$	$^{\hat{\Phi}}$ h $^{ extsf{D}}$	Φ̂hB					
n	75	28	7	28	6	24					
Mean	0.298	0.317	0.341	0.795	1.070	0.401					
Std. dev.	0.075	0.068	0.038	0.108	0.134	0.118					
CV (%)	25	21	11	14	13	29					
Std. err.	0.009	0.013	0.015	0.020	0.055	0.024					
Skewness	-0.75	-0.31	-0.53	-0.15	-0.57	0.07					
Kurtosis	0.444	-0.52	0.66	0.23	-0.44	-1.13					
Range	0.332	0.262	0.118	0.503	0.355	0.386					
Percentiles											
0 (Min.)	0.104	0.184	0.274	0.542	0.860	0.229					
1	*	*	*	*	*	*					
5	0.130	0.189	*	0.589	*	0.229					
10	0.194	0.202	*	0.651	*	0.241					
25	0.252	0.269	0.321	0.704	0.959	0.273					
50 (Median)	0.308	0.321	0.347	0.827	1.078	0.399					
75	0.341	0.377	0.374	0.866	1.202	0.490					
90	0.385	0.398	*	0.893	*	0.573					
95	0.410	0.425	*	1.003	*	0.608					
99	*	*	*	*	*	*					
100 (Max.)	0.436	0.446	0.392	1.045	1.215	0.615					

^{*}Indicates percentiles that are identical to the minimum (or maximum) values due to small sample size.

Table 6.2. Statistical Distributions of Allometric Powers for Forestry Yield Table Data.

	Allometric Parameter									
Statistic	p _{est}	$\hat{\Phi}$ hw \cdot	ΦDw	Φ̂Βw	$^{\hat{\Phi}}$ h $^{ m D}$	∳hB				
n Mean Std. dev. C.V. (%) Std. err. Skewness Kurtosis Range Percentiles O (min.) 1 5 10 25 50 (Median) 75 90 95 99 100 (Max.)	351 0.298 0.067 22 0.004 -1.46 2.91 0.392 0.054 0.065 0.147 0.220 0.271 0.309 0.338 0.365 0.379 0.409 0.447	325 0.274 0.059 22 0.003 0.352 0.666 0.399 0.118 0.137 0.183 0.209 0.234 0.271 0.310 0.357 0.357 0.371 0.426 0.517	334 0.368 0.057 16 0.003 0.626 11.4 0.644 0.165 0.202 0.259 0.303 0.340 0.375 0.402 0.424 0.432 0.484 0.810	318 0.747 0.108 14 0.006 1.19 18.8 1.341 0.330 0.338 0.584 0.655 0.697 0.750 0.810 0.855 0.868 0.940 1.670	323 0.770 0.230 30 0.013 1.12 1.99 1.435 0.327 0.353 0.473 0.545 0.595 0.733 0.873 1.072 1.225 1.570 1.762	309 0.387 0.130 34 0.007 1.77 5.74 0.927 0.131 0.163 0.232 0.274 0.294 0.362 0.438 0.535 0.617 0.958 1.058				




Figure 6.1. Histograms of allometric powers of the experimental and field data. (a) through (f) show, respectively, the statistical distributions of the transformed thinning slope, p_{est} , and the allometric powers $\widehat{\Phi}_{hw}$, $\widehat{\Phi}_{Dw}$, $\widehat{\Phi}_{Bw}$, $\widehat{\Phi}_{hD}$, and $\widehat{\Phi}_{hB}$.




Figure 6.2. Histograms of allometric powers of the forestry yield table data. (a) through (f) show, respectively, the statistical distributions of the transformed thinning slope, p_{est} , and the allometric powers $\hat{\Phi}_{hw}$, $\hat{\Phi}_{Dw}$, $\hat{\Phi}_{Bw}$, $\hat{\Phi}_{hD}$, and $\hat{\Phi}_{hB}$.

Table 6.3. Correlations and Regressions Relating Transformed Thinning Slope to Allometric Powers.

		C		0		Linear Regression Equation					
Allo-		Spearm Correla		Pearson Correlation			Slop	e	Intercept		
netric Power	n	rs	Р	r	Р	r²	Â	95% CI	â	95% CI	
				Exp	erimental	and Fie	ld Data				
φ̂ hw	28	-0.51	0.0056*	-0.55	0.0026*	0.30	-0.71	[-1.14,-0.26]	0.50	[0.36, 0.64]	
φύν Φβω Φhù	7	-0.57	0.18	-0.63	0.13	0.40	-0.84	[-2.02, 0.34]	0.53	[0.13, 0.94	
ADW Pew	28	0.20	0.30	0.14	0.49	0.02	0.11	[-0.21, 0.44]	0.19	[-0.08, 0.45	
∧on ⊅ hù	6	-0.14	0.79	0.15	0.78	0.02	0.06	[-0.45, 0.56]	0.18	[-0.36, 0.72	
^A hB	24	-0.46	0.024*	-0.44	0.032*	0.19	-0.35	[-0.67,-0.03]	0.41	[0.28, 0.55	
				For	restry Yie	<u>ld Tabl</u>	e <u>Data</u>				
A Priw	325	-0.44	<0.0001*	-0.46	<0.0001*	0.21	-0.54	[-0.65,-0.43]	0.45	[0.41, 0.48	
Anw ADW ABW ADD ADD ADD ADD ADD ADD ADD AD	334	0.47	<0.0001*	0.48	<0.0001*	0.23	0.58	[0.47, 0.69]	0.09	[0.04, 0.13	
ΛDW D _{ELW}	318	0.55	<0.0001*	0.59	<0.0001*	0.35	0.36	[0.30, 0.41]	0.03	[-0.01, 0.07	
^º™ ⊅ n∩	323	-0.57	<0.0001*	-0.67	<0.0001*	0.45	-0.20	[-0.23,-0.18]	0.45	[0.43, 0.47	
^''D ₽ _{nR}	309	-0.56	<0.0001*	-0.71	<0.0001*	0.50	-0.36	[-0.40,-0.32]	0.44	[0.42, 0.45	

^{*}Significant at the 95% confidence level (P \leq 0.05).






Figure 6.3. Observed relationships between transformed thinning slope and allometric powers in the experimental and field data. (a) shows p_{est} , plotted against the $\hat{\Phi}_{hw}$, the average height-average weight allometric power. The solid line is the regression $p_{est} = -0.71 \hat{\Phi}_{hw} + 0.50$, which is not significantly different (Table 6.3) from the predicted relationship $p = -0.5^{\circ}\hat{\phi}_{hw} + 0.50^{\circ}$ (dotted line). (b) shows p_{est} plotted against $\hat{\Phi}_{hB}$, the average height-average BSLA allometry. The solid regression line is p_{est} = -0.35 $\hat{\Phi}_{hB}$ + 0.41.

to predict p_{est} : 21% to 50% of the variance in p_{est} was explained. Even $\hat{\Phi}_{DW}$ and $\hat{\Phi}_{BW}$, which were not expected to relate well with p_{est} , were actually significantly correlated to p_{est} and explained 23% and 35%, respectively, of its variance. The relationship of p_{est} to $\hat{\Phi}_{hw}$ explained 21% of the variance in p_{est} , and the regression line $p_{est} = -0.54$ $\hat{\Phi}_{hw} + 0.445$ was close to the expected equation p = -0.5 $\Phi_{hw} + 0.5$ and did not differ significantly in slope from this line. The regression intercept of -0.445 was significantly less than expected, but the difference was small: the regression intercept was 11% below the expected 0.5. The position of the observed relationship below the expected one in the p_{est} - Φ_{hw} plane is partly due to the invalid but necessary assumption that $\Phi_{dw} = 0$. Figure 6.4 shows the relationships of p_{est} with $\hat{\Phi}_{hw}$, $\hat{\Phi}_{hD}$, and $\hat{\Phi}_{hB}$.

Small group sizes hindered the comparisons among the six groups of the EFD (herbaceous monocots, herbaceous dicots, temperate angiosperm trees, temperate gymnosperm trees, Eucalypts, and tropical angiosperm trees--see Table 6.4). Height was the most commonly measured plant dimension, yet only 28 of the 75 data sets that showed a significant log B-log N relationship also had height data and showed a significant log \bar{h} -log \bar{w} relationship. Group sizes were as low as one or two observations, so the absence of differences in $\hat{\Phi}_{hW}$ among the six groups is inconclusive. A significant difference among the four tree groups was observed for $\hat{\Phi}_{hB}$, and the mean values of $\hat{\beta}$ and $\hat{\Phi}_{hB}$ for three groups were ranked in accordance with the allometric hypothesis: more




Figure 6.4. Observed relationships between transformed thinning slope and allometric powers in the forestry yield table data. (a) shows the transformed thinning slope, p_{est} , plotted against $\hat{\Phi}_{hw}$, the average height-average weight allometry. The solid line is the regression $p_{est} = -0.54~\hat{\Phi}_{hw} + 0.45$ while the dotted line is the predicted relationship $p = -0.5~\hat{\Phi}_{hw} + 0.50$. (b) shows p_{est} plotted against $\hat{\Phi}_{hD}$, the average height-average DBH allometry. The regression line is $p_{est} = -0.20~\hat{\Phi}_{hD} + 0.45$. (c) shows p_{est} plotted against $\hat{\Phi}_{hB}$, the average height-average BSLA allometry, and the regression line is $p_{est} = -0.36~\hat{\Phi}_{hB} + 0.44$. Table 6.3 gives further statistical information.

Table 6.4. Tests for Differences in Allometric Powers Among Groups in the Experimental and Field Data.

			Groi	up Means ^a		Tests for Differences Among Groups					
0.7.2	Не	rbs	Trees				A11 G	roups ^b	Temperate Trees Angio. vs. Gymno.		
Allo- metric Power	Mono- cots.	Di- cots.	Temperate Angio.	Temperate Gymno.	Eucal.	Tropical Angio.	ANOVAC	Kruskal- Wallis ^d	ANOVAC	Kruskal- Wallis ^d	
Ŗ	-0.44 (8)	-0.74 (25)	-0.65 (15)	-0.87 (19)	-1.26 (4)	-2.56 (4)	7.68(5,69) P<0.0001*	17.9(5) P=0.0031*	0.90(1,32) P=0.35	0.37(1) P=0.54	
â	4.45 (8)	5.17 (25)	3.78 (15)	3.79 (19)	2.87 (4)	2.20 (4)	10.9(5,69) P<0.0001*	41.1(5) P<0.0001*	0.00(1,32) P=0.97	1.43(1) P=0.23	
p _{est}	0.36 (8)	0.30 (25)	0.32 (15)	0.30 (19)	0.24 (4)	0.16 (4)	5.77(5,69) P=0.0002*	17.9(5) P=0.0031*	0.57(1,32) P=0.46	0.37(1) P=0.54	
$\hat{\Phi}_{\sf hw}$	0.30	0.33	0.27 (9)	0.35 (10)	0.36 (2)	0.30 (4)	1.88(5,22) P=0.14	7.13(5) P=0.21	7.51(1,17) P=0.014*	5.61(1) P=0.018*	
ĜD w			0.27 (1)	0.33 (3)	0.37 (3)		10.62(2,4) P=0.025*	4.00(2) P=0.14	13.7(1,2) P=0.066	1.80(1) P=0.18	
Φ̂Βw			0.82 (8)	0.79 (12)	0.76 (4)	0.81 (4)	0.25(3,24) P=0.86	1.44(3) P=0.70	0.32(1,18) P=0.58	0.38(1) P=0.54	

Table 6.4. (continued)

		Group Means ^a						Tests for Differences Among Groups					
	Не	Herbs Trees				All Groups ^b		Temperate Trees Angio. vs. Gymno.					
Allo- metric Power	Mono- cots.	Di- cots.	Temperate Angio.	Temperate Gymno.	Eucal.	Tropical Angio.	ANOVAC	Kruskal- Wallis ^d	ANOVAC	Kruskal- Wallis ^d			
φ̂hD				1.14 (3)	1.01		1.56(1,4) P=0.28	1.19(1) P=0.28					
Ŷ ŶhB			0.28 (7)	0.46 (10)	0.51 (3)	0.38 (4)	7.35(3,20) P=0.0017*	12.0(3) P=0.0076*	20.4(1,15) P=0.0004*	9.17(1) P=0.0025*			

aGroup sizes are given in parentheses.

 $^{\text{CF}}$ ratios for the ANOVAs are given, along with the degrees of freedom (in parentheses). P values are the statistical significance levels.

dThe H statistic from the Kruskal-Wallis test is given, along with the degrees of freedom (in parentheses) used to estimate the statistical significance level, P, by the χ^2 approximation.

bTests for differences among all groups for which data were available.

^{*}Indicates significance at the 95% confidence level (P \leq 0.05).

A more conclusive analysis is possible for two tree groups, temperate angiosperms and temperate gymnosperms, that had larger group sizes (n \geq 7) for $\hat{\beta}, \; \hat{\Phi}_{hw}, \; \text{and} \; \hat{\Phi}_{hB}.$ The difference in $\hat{\beta}$ between the two groups was not statistically significant; however, the gymnosperms seemed to have steeper self-thinning slopes (mean $\hat{\beta}$ = -0.87) than did the angiosperms (mean $\hat{\beta}$ = -0.61). If the allometric hypothesis is correct, this trend should be accompanied by a greater emphasis in height growth relative to radial growth in the gymnosperms than in the angiosperms. Such trends were present and statistically significant. For the gymnosperms, $\hat{\Phi}_{hw}$ (mean $\hat{\Phi}_{hw}$ = 0.35) was significantly greater than for the angiosperms (mean $\hat{\Phi}_{hw}$ = 0.27). In addition, $\hat{\Phi}_{hB}$ for the gymnosperms (mean $\hat{\Phi}_{hB}$ = 0.46) significantly exceeded the angiosperm value (mean $\hat{\Phi}_{hB}$ = 0.28).

The larger number of thinning trajectories and the more complete availability of dimensional data in the FYD permitted a more conclusive analysis of among-group differences (Table 6.5). The three groups of trees were temperate angiosperms, temperate gymnosperms, and Eucalypts, with mean $\hat{\beta}$ values of -0.60, -0.80,

Table 6.5. Tests for Differences in Allometric Powers Among Groups in the Forestry Yield Table Data.

					Differences e Groups	Among Groups Angiosperms vs. Gymnosperms		
Allo- metric	Gr	oup Mean	s ^a		Kruskal-		Kruskal-	
Power	Angio.	Gymno.	Eucal.	ANOVAP	Wallis ^C	ANOVAD	Wallis ^C	
Ŗ	-0.60	-0.80	-3.90	79.9(348)	14.9	4.85(337)	3.77	
	(58)	(281)	(12)	P<0.0001	P=0.0006	P=0.028	P=0.052	
â	3.50	3.54	1.09	38.9(348)	11.9	0.13(337)	8.30	
	(58)	(281)	(12)	P<0.0001	P=0.0027	P=0.8	P=0.004	
pest	0.32	0.30	0.17	29.1(348)	14.9	7.28(337)	3.77	
	(58)	(281)	(12)	P<0.0001	P=0.0006	P=0.0073	P=0.052	
[^] hw	0.22	0.28	0.30	23.8(322)	36.3	49.9(314)	36.9	
	(47)	(260)	(9)	P<0.0001	P<0.0001	P<0.0001	P<0.0001	
$\hat{\Phi}_{Dw}$	0.40	0.36	0.33	11.4(331)	20.3	18.3(307)	18.1	
	(53)	(269)	(12)	P<0.0001	P<0.0001	P<0.0001	P<0.0001	
[∳] B w	0.85	0.73	0.73	25.3(315)	44.6	50.2(320)	44.8	
	(44)	(265)	(9)	P<0.0001	P<0.0001	P<0.0001	P<0.0001	
[∳] hD	0.56	0.80	0.98	28.1(320)	51.4	49.9(312)	49.9	
	(45)	(269)	(9)	P<0.0001	P<0.0001	P<0.0001	P<0.0001	
[∳] hB	0.27	0.40	0.53	42.9(306)	21.8	36.2(301)	41.4	
	(38)	(265)	(6)	P<0.0001	P<0.0001	P<0.0001	P<0.0001	

aGroup sizes are given in parentheses.

bF ratios for ANOVA are given, along with the denominator degrees of freedom (in parentheses). The three and two-group tests have, respectively, two and one degree(s) of freedom in the numerator of the F ratio. P values are the statistical significance levels.

 $^{\text{C}}\text{The H}$ statistic from the Kruskal-Wallis test is given. Two and one degrees(s) of freedom , respectively, were used in the three-group and two-group tests to estimate statistical significance levels, P, by the χ^2 approximation.

and -3.90, respectively. These means were significantly different, both when all three groups were considered and when only the temperate angiosperms and temperate gymnosperms are compared (Chapter 5). The allometric hypothesis predicts that the allometric powers should be ranked in the same order: angiosperms with the lowest rate of height growth relative to radial growth, gymnosperms intermediate, and Eucalypts highest. This prediction is verified for all three of the allometric measures, $\hat{\Phi}_{\text{hw}}$, $\hat{\Phi}_{\text{hD}}$, and $\hat{\Phi}_{\text{hB}}$. The groups show the expected rankings and the differences among groups are statistically significant.

The observed correlations of shade tolerance with β in the FYD (Chapter 5) were accompanied by corresponding correlations of tolerance with the allometric powers (Table 6.6). For the gymnosperms, a significant positive correlation of β with tolerance was accompanied by negative correlations of shade tolerance with $\hat{\Phi}_{hw}$, $\hat{\Phi}_{hD}$, and $\hat{\Phi}_{hB}$, so shallower thinning slopes were associated with less emphasis on height growth relative to radial growth as predicted by the allometric hypothesis. The angiosperms showed very different behavior: thinning slope was negatively correlated with tolerance. This result is still consistent with the allometric hypothesis because the negative correlation between shade tolerance and β is accompanied by positive correlations of tolerance with ${\stackrel{\wedge}{\Phi}}_{\sf hw}$, ${\stackrel{\wedge}{\Phi}}_{\sf h\Omega}$, and $\hat{\tilde{\Psi}}_{\mathsf{hR}}.$ However, the positive correlations of tolerance with $\hat{\Phi}_{\mathsf{hD}}$ and $\hat{\Phi}_{\mathsf{hB}}$ are not statistically significant, while the correlation between tolerance and the $\overset{\wedge}{\Phi}_{f hw}$ allometry is of

Table 6.6. Spearman Correlations of Shade Tolerance with Allometric Powers in the Forestry Yield Table Data.

	Mea	ins for Sh	oups ^a	Spearman Correlation			
Parameter	1	2	3	4	5	r_{S}	рb
		Temper	ate Angio	osperms	0.35.46	1.12 . 19.14 . 1 . 1 . 1 . 1 . 1 . 1 . 1 . 1 . 1	
â	-0.391 (10)	-0.547 (18)				-0.52* (46)	0.0002
â	3.632 (10)		3.517 (18)			-0.19 (46)	0.22
Pest	0.364 (10)	0.335 (18)	0.299 (18)			-0.52* (46)	0.0002
$\overset{\wedge}{\Phi}$ hw	0.191 (7)	0.220 (18)	0.231 (16)			0.29 (41)	0.070
${}^{\wedge}_{Dw}$	0.350 (10)		0.404 (18)			0.19 (44)	0.21
Φ Bw	0.835 (9)	0.906 (12)	0.811 (16)			-0.18 (37)	0.28
$\overset{\wedge}{\Phi}$ h $ exttt{D}$	0.502 (7)	0.551 (16)	0.577 (16)			0.20 (39)	0.22
^Λ ^Φ hB	0.241 (6)	0.249 (12)				0.25 (34)	0.14
Wood Density	430 . (10)	522 . (18)				0.92 * (46)	<0.0001

Table 6.6. (continued)

	Mea	ans for Sh	Spearman Correlation				
Parameter	1	2	3	4	5	r_{S}	рb
		Ten	perate G	ymnosperm	IS.		
β	-0.916 (32)	-0.748 (78)	-0.642 (47)		-0.459 (41)	0.35* (267)	<0.0001
â	3.123 (32)	3.438 (78)	3.732 (47)	3.280 (69)	4.172 (41)	0.57 * (267)	<0.0001
pest	0.266 (32)	0.300 (78)	0.308 (47)	0.279 (69)	0.344 (41)	0.35 [*] (267)	<0.0001
^Ф hw	0.299 (32)	0.285 (78)	0.274 (47)	0.301 (69)	0.247 (41)	-0.24* (267)	<0.0001
$\hat{\Phi}_{Dw}$	0.348 (32)	0.345 (78)	0.378 (47)	0.354 (69)	0.411 (41)	0.46 [*] (267)	<0.0001
$^{\wedge}_{ar{\Phi}}$ Bw	0.721 (32)	0.702 (78)	0.758 (46)	0.699 (68)	0.821 (41)	0.40 * (265)	<0.0001
Φ̂hD	0.865 (32)	0.834 (78)	0.745 (47)	0.885 (69)	0.603 (41)	-0.36* (267)	<0.0001
$\hat{\Phi}_{nB}$	0.420 (32)	0.410 (78)	0.370 (46)	0.457 (68)	0.301 (41)	-0.31* (265)	<0.0001
Wood Density	564 . (32)	553 . (78)	500. (47)	415. 5 (69)		-0.42* (267)	<0.0001

aSample sizes are given in parentheses.

bStatistical significance level.

^{*}Significant at the 95% confidence level (P < 0.05).

borderline significance (P = 0.07). These low confidence levels probably reflect the small sample sizes available for the angiosperms. The 46 angiosperm thinning trajectories considered in this analysis were drawn from only 15 different yield tables. The tables for mixed hardwood forests that contributed many estimates of $\hat{\beta}$ could not be used because average shade tolerances could not be assigned to these diverse communities.

Discussion

These results strongly support the allometric hypothesis. Logical deductions from the basic hypothesis β = -1/(2p) + 1 predicted that the height-weight, height-DBH, and height-BSLA allometric powers would be negatively correlated with the transformed thinning slope. This prediction was repeatedly verified by significant observed correlations of p_{est} with all three allometric powers of the FYD and with $\hat{\Phi}_{hw}$ and $\hat{\Phi}_{hB}$ in the EFD. Only in the single case of $\hat{\Phi}_{hD}$ in the EFD were the results inconclusive because of the small sample size (n = 6). An equation was also predicted for the expected relationship between p_{est} and Φ_{hw} (p_{est} = -0.5 Φ_{hw} + 0.5) and the observed relationships in the EFD and FYD were both close to this prediction.

The allometric hypothesis is further supported by the association of differences in thinning slope among groups with corresponding differences in plant allometry, as expected if allometric factors determine β . This correspondence was also observed when the correlation of shade tolerance with $\hat{\beta}$ was

compared to the correlations of tolerance with the allometric powers. The results of these two analyses were not as statistically significant as the results of direct correlation of $p_{\rm est}$ with the allometric parameters because these two analyses are less powerful and group sizes were small. Nevertheless, several significant trends were documented that did verify the predictions of the allometric hypothesis.

The results also address some questions posed in previous studies. Lonsdale and Watkinson (1983a) questioned whether plants growing at densities high enough to thin actually have different shapes. The variation documented in the univariate distributions of the allometric powers, the significant differences in these powers among groups, and the significant correlations of these powers with shade tolerance indicate that there are important variations in the shapes of plants undergoing self-thinning. Miyanishi et al. (1979) and Perry (1984) posed another important question: Are observed variations in the thinning slope due to experimental errors or biological reality? In fact, variations in self-thinning slope are systematically related to variations in plant allometry, so the deviations from the idealized value β = -1/2 are real, not simply errors in observing a biological constant.

There are several reasons why this study has found strong evidence for the allometric hypothesis while previous studies (Westoby 1976, Mohler et al. 1978, White 1981) did not. A major problem was the accepted belief that all thinning slopes are near the value predicted by the self-thinning rule, so that the goal was

to show how the diverse allometries of real plants are resolved to give the same thinning slope. Allometric models that failed to do this were judged to be in contradiction of empirical evidence (White 1981). New information on the true variability among thinning slopes (Chapters 4 and 5) leads naturally to a new emphasis: Can the observed variation be attributed to variation in plant allometry? The present results indicate that it can. Previous studies have also been hindered by small sample sizes. One of the major studies (Mohler et al. 1978) considered only two species for which both allometric data and self-thinning lines were available. The present results show so much variation around a relationship between thinning slope and an allometric parameter (Figures 6.3 and 6.4) that a sample of two points would be useless in detecting the trend. More data were available here because theoretical models were carefully analyzed to predict the relationship between thinning slope and allometries involving frequently measured plant dimensions, such as the height, DBH, and BSLA. Previous studies (Mohler et al. 1978, White 1981) have tried to predict thinning slope directly from some allometry involving crown area, which is more difficult to measure and less frequently reported than height or DBH.

Another factor in the present success is the method of fitting allometric powers. Typically two variables, such as height and weight, are measured for a sample of individuals and the allometric equation is fit to these data, so the allometry describes a relationship among individual plants. In contrast, the thinning

line is a relationship based on aggregate measurements of density and biomass for whole populations rather than for individuals. Here allometric powers were fit to aggregate stand variables, such as average height and average weight, so the results are more commensurate with thinning slope than allometric powers derived at the individual plant level. These whole-stand allometric powers also directly address the time-dynamics of shape change with stand growth, while allometries measured from a sample of individual plants focus more on the static size structure within a single sample. A dynamic emphasis is more relevant because self-thinning is a dynamic, whole-stand process. The whole-stand approach also permits allometric powers to be fit to the same stands of plants used to estimate $\hat{\beta}$. Other studies (Mohler et al. 1978, White 1981) have compared thinning slopes to allometric data from a small subsample of individuals or from completely different sources. This introduces confounding variations because allometric relationships vary significantly with site, time, and other factors (Hutchings 1975). Thinning slopes are also not species constants (Chapters 4 and 5).

Previous studies of self-thinning and allometry also had other problems. Westoby (1976) tested the allometric derivation of the self-thinning rule by measuring the thinning slope of prostrate cultivars of <u>Trifolium subterraneum</u>. He predicted that the plants would grow only radially and give log w-log N thinning slopes near -1, but the observed thinning slope was near the more typical value $\gamma = -3/2$. This was interpreted as a falsification of the

allometric theory. However, the experiment and conclusion have been questioned by White (1981), who does not accept the assumption that height remains constant for this species. The high variability around the relationships observed here demonstrates the futility of testing the allometric hypothesis with a single observation. The most detailed previous review of allometry and self-thinning (White 1981) also has some inconsistencies. This analysis predicted the range of variation in thinning slopes from two tree allometries, the weight-DBH allometry and the crown area-DBH allometry. While the effects of variation in the crown area-DBH allometry on thinning slope were carefully considered, the relationship between weight and DBH was fixed at w \propto DBH^{2.5}, despite evidence in the same report that the allometric power actually ranges from 1.8 to 3.3. In view of this omission, the derived ranges for thinning slopes are suspect, as are the conclusions drawn from these ranges.

There are, then, many reasons why this study found significant relationships between plant allometry and self-thinning slope while previous studies did not. However, the successes of this study were less than hoped. Several statistically significant relationships between thinning slope and allometric powers were found, but the most predictive explained only 50% of the variation among thinning slopes. There are at least five major sources of variation that reduced the explanatory power of the regressions: errors in estimating the thinning slope, errors introduced by the invalid assumption that the density of biomass in occupied space does not vary with size, differences in measurement methods among studies,

differences in the criteria for counting individuals, and errors from using independent variables that are only indirectly related to p, the allometry hypothesized to actually determine β .

The potentially serious errors from a posteriori manipulations required in estimating β have already been discussed (Chapter 4), as have the errors from incorrectly assuming that Φ_{hd} = 0. Differences in measurement methods also reduced explanatory power. For example, the regression model predicting p_{est} from $\hat{\Phi}_{hw}$ for the FYD is derived from studies that variously defined stand height as the height of the tallest tree, the average of of dominant trees only, the average of dominant and codominant trees, or the average of all trees. Other variations come from different criteria for deciding which individuals should be counted. In forest surveys, individuals below a certain DBH are commonly ignored, and this threshold varies with species, stand size, and the purpose of the study. Studies also differ in including certain plant parts in biomass measurements. Ecologists tend to include small twigs, roots, and leaves, while foresters tend to ignore these parts and report only the economically important volume of harvestable wood. These wood volumes vary further with the method of harvesting and the uses being considered (lumber, pulp, poles, etc.).

The indirect method of testing the allometric hypothesis also reduces the explanatory power of statistical models. The allometric powers Φ_{hw} , Φ_{hD} , and Φ_{hB} are only indirectly related to the power p that is hypothesized to determine the thinning slope, and these indirect relationships are confounded by structural and

mechanical constraints on plant form. When the uncertainties due to measurement errors, differences in defining measurements, and the use of indirect allometric measures are all combined, the relatively low coefficients of determination observed in relating thinning slope to allometric powers are no longer surprising. Rather, the ability to simply detect the statistically significant correlations between thinning slope and allometric measures is a major positive result.

The statistical models demonstrate the importance of allometric factors in determining the thinning slope, but the significant proportion of unexplained variance admits the possibility that other factors may also affect β , such as the physiological parameters considered in some recent models (Pickard 1983, Perry 1984).

Some of the present results are also of interest beyond the context of the self-thinning rule. For example, one analysis of the FYD indicated opposite syndromes of adaptation to shading in the gymnosperm and angiosperm trees of the northern temperate forests. Among the gymnosperms, more tolerant trees exhibit shallower thinning slopes, greater height growth relative to radial growth, and wood that is less dense. On the other hand, the angiosperms show steeper thinning slopes with increasing shade tolerance (suggesting less height growth relative to radial growth) and greater wood density. Further analysis of this dichotomy in the adaptive responses to shading between the two groups of trees may yield insight into the origin and evolution of the two taxa and may contribute to general theories of tree strategies and forest dynamics.

CHAPTER 7

THE OVERALL SIZE-DENSITY RELATIONSHIP

Introduction

This chapter considers the overall relationship between size and density among stands of different species and evaluates its relevance to the self-thinning rule. This relationship has been observed by plotting $\log \overline{w}$ against $\log N$ for stands of species ranging from herbs to trees (Gorham 1979), and by plotting self-thinning lines for many species on a single $\log \overline{w}$ - $\log N$ plot (White 1980). In both cases, the data are grouped around the line $\log \overline{w} = -3/2 \log N + 4$. This trend is referred to here as the overall relationship to distinguish it from the self-thinning lines of individual populations.

The existence of the overall relationship has been interpreted as strong evidence for the self-thinning rule. It has been claimed to show that: (1) The model for intraspecific thinning of Yoda et al. (1963) also applies to interspecific weight-density relationships (Gorham 1979). (2) The same self-thinning law applies to all plants (Hutchings and Budd 1981a, Hutchings 1983). (3) The slope and position of the thinning line are insensitive to plant geometry (Furnas 1981, Hutchings and Budd 1981a). (4) The empirical generality of the self-thinning rule is beyond question (White 1981). The causes of the overall relationship are examined here to determine if the existence of the relationship actually supports such interpretations.

Models of the Overall Size-density Relationship

Heuristic Model

The overall relationship between size and density can be deduced from basic principles using a heuristic model for fully-stocked stands of different plant species. Assume that all the plants in a given stand are the same size and shape and that each plant occupies a cylindrical volume of space called its volume of influence (VOI). Plant weight and the volume and dimensions of this cylinder are related by

$$w = v d = (\pi R^2 h) d$$
, (7.1)

where w is plant weight (in g), v is the size of the VOI (in m^3), d is the density of plant material in the VOI (in g/m^3), and R and h are the radius and height, respectively, (in m) of the VOI. The ground area occupied by each plant is π R² and if all ground area is covered the density of plants per m^2 is simply

$$N = 1 / (\pi R^2) . (7.2)$$

The shape of the plants in a stand can be represented by the height to VOI radius ratio, τ = h/R. Since h = τ R, equation 7.1 can be rewritten as

$$w = (\pi R^3 \tau) d$$
, (7.3)

which can be combined with equation 7.2 to give

$$w = [\tau d / \sqrt{\pi}] N^{-3/2}, \qquad (7.4)$$

for a particular stand.

If plants in all stands had exactly the same values of τ and d, then the overall relationship among all stands would reduce to $\overline{w} = K \ N^{-3/2}$ where $K = \tau \ d \ / \sqrt{\pi}$. Actually, τ and d will vary among stands. Gorham's (1979) data can be used to estimate a range for τ . Assume that VOIs are cylindrical and that the base area, \overline{a} , of a VOI is 1/N, then substitute and solve $\overline{a} = \pi \ \overline{R}^2$ to obtain $\overline{R} = (\pi \ N)^{-1/2}$. The estimated \overline{R} values can then be divided into the reported average heights to yield τ , which ranges from 2 to 12 for Gorham's 65 stands. Measurements of biomass per unit of canopy volume can provide estimates of d, the density of biomass in occupied space. For trees, the measurements range from 600 to 40000 g/m³, with most falling between 4000 and 13000 g/m³ (White 1980). Lonsdale and Watkinson (1983a) reported values between 1500 and 5200 g/m³ for three herbaceous species. Combining the ranges for τ and d with the log-transformed version of equation 7.4,

$$\log \bar{w} = -3/2 \log N + \log(\tau d / \sqrt{\pi})$$
 , (7.5)

yields equations for the minimum and maximum average weights possible at any plant density. The minimum values of 2 and 600 for τ and d, respectively, give $\log \varpi = -3/2 \log N + 2.83$, while the respective maximum values of 12 and 40000 give the parallel line $\log \overline{w} = -3/2 \log N + 5.43$. These two lines define a region of the $\log \overline{w} - \log N$ plane enclosing all fully-stocked stands of identically

sized plants. Gorham's (1979) line, $\log \bar{w} = -1.49 \log N + 3.99$, is parallel to the boundary lines in the approximate center of the enclosed region. The principal axis line through a random sample of points distributed uniformly between the model boundary lines would have a slope of -3/2 and would be near Gorham's line.

Improved Model

The assumption that all plants in each stand are the same size and shape is clearly invalid, so this section analyzes a more complex model which represents each stand as a population with a lognormal distribution of individual weights. Within each stand, base area is represented by a simple power function of weight. Lognormal weight distributions have been repeatedly observed in plant stands (Koyama and Kira 1956, Obeid et al. 1967, White and Harper 1970, Hutchings and Budd 1981a). Power functions relating plant dimensions to weight have been validated and applied by both ecologists (Whittaker and Woodwell 1968, Hutchings 1975) and foresters (Bruce and Schumacher 1950). The weight distribution of a stand is specified by the lognormal distribution parameters ζ and σ , and the mean weight is

$$\bar{w} = e^{(\zeta + 0.5\sigma^2)}$$
 (7.6)

(Johnson and Kotz 1970). The area occupied by an individual is obtained by assuming that all plants fill a cylindrical VOI to a density of d $\rm g/m^3$, and that the height to radius ratio of a plant

of average weight is τ . Solving $\bar{w} = \pi \; \bar{R}^3 \; \tau \; d$ for \bar{R} then gives the base radius of a plant of average weight

$$\bar{R}_{\bar{W}} = \left[\frac{\bar{w}}{\pi \tau d}\right]^{(1/3)}, \qquad (7.7)$$

and the base area of the plant of average weight is simply $\pi\ \bar{R}^2.$ Across an entire stand, R is related to w by

$$R = c w^p , \qquad (7.8)$$

where c and p are constants, so that base area a is

$$a = \pi c^2 w^{2p}$$
 (7.9)

Equations 7.7 and 7.9 can now be combined to fix the value of c at c = \sqrt{a} / ($\sqrt{\pi}$ \overline{w}^p) and yield a general formula for the base area of any plant

$$a = \pi \left[\frac{1}{\bar{w}^p} \left(\frac{\bar{w}}{\pi + d} \right)^{1/3} \right]^2 w^{2p}$$
, (7.10)

which simplifies to a = c_1 w^{2p}, where c_1 = π c². Since In w is normally distributed with mean ζ and standard deviation σ (Johnson and Kotz 1970) and c_1 and p are constants, In a = $\ln (c_1 w^{2p}) = \ln c_1 + 2 p \ln w$ is normally distributed with mean $\ln c_1 + 2 p \zeta$ and standard deviation $2 p \sigma$. The expected value of a is

$$\bar{a} = e^{[\ln c_1 + 2p\zeta + 0.5 (2p\sigma)^2]},$$
 (7.11)

which simplifies to

$$\bar{a} = c_1 e^{2p (\zeta + p\sigma^2)}$$
 (7.12)

Assuming $N = 1 / \bar{a}$, equation 7.12 gives

$$N = 1 / [c_1 e^{2p (\zeta + p\sigma^2)}], \qquad (7.13)$$

the plant density for a stand at 100% cover specified by parameters ζ , σ , d, and p.

The overall log w-log N relationship for this model was estimated by using Monte Carlo techniques to select parameters for 500 plant stands, then calculating the resulting densities (equation 7.13) and fitting a relationship between $\log \overline{w}$ and $\log N$. Parameters ζ and σ were actually derived from two parameters with more direct biological interpretations: the average weight \bar{w} and ρ , which measures the range of weights within a population. Individual weights in a crowded plant population commonly span about two orders of magnitude (White 1980). This information was applied to the distribution of ln w by assuming that the ninety-ninth percentile weight is approximately ρ times the first percentile value, where ρ ranged from 10 to 1000 to allow for deviations from the reported value of 100. Since In w is normally distributed, the ninety-ninth percentile value is 4.652 standard deviations above the first percentile value and $e^{4.652\sigma} = \rho$. With this information, ζ and σ can be calculated sequentially from $\sigma = \rho/4.652$ and $\zeta = \ln \bar{w} - 0.5 \sigma^2$.

Values of the input parameters log τ , p, log d, log \overline{w} , and $\log \rho$ were chosen from uniform random distributions on the intervals given in Table 7.1. Statistics for the 500 derived values of ζ , σ , \bar{R} , \bar{h} , \bar{a} , N, and B are given in Table 7.2, along with values of a constant κ calculated from κ = B \sqrt{N} . Since \bar{w} was chosen a priori and random parameter variations were imposed in deriving N, \bar{w} is a truly independent variable and the correct method for estimating an overall linear relationship between log w and log N is regression of log N against log \bar{w} . The regression line was $\log N = -0.654 \log \bar{w} + 2.799 (r^2 = 0.96, P < 0.0001, 95\% CI for$ slope = [-0.67, -0.64]), which can be rearranged to give $\log \bar{w} = -1.53 \log N + 4.28 (95\% CI for slope = [-1.56, -1.49]), or$ equivalently, log B = -0.53 log N + 4.28 (95% CI for slope =[-0.56, -0.49]). Figure 7.1 shows the 500 hypothetical plant stands and this fitted relationship after transformation to log B-log N form to facilitate comparison with real plant data.

Discussion

The models show that the linear overall relationship between log size and log density results from the simple geometry of packing three-dimensional objects onto a two-dimensional surface. The slope of -3/2 in the log \bar{w} -log N plane follows directly from assuming a cylindrical shape for the VOI. The volume of any cylinder is a cubic function of the base radius while the base area is a function of the same radius squared, so that the ratio of these two powers is 3/2. This result is robust to changes in the basic cylindrical

Table 7.1. Parameter Ranges for Monte Carlo Analysis of the Improved Model.

Parameter	Parameter Minimum	limits ^a Maximum	Antilog Minimum	limits ^b Maximum
log ₩	-2.00	7.00	0.01	107
log τ	0.30	1.08	2	12
р	0.05	0.45		
log d	2.78	4.60	600	40000
log x	1.00	3.00	10	1000

aparameter values for 500 plant stands were chosen from uniform random distributions on the indicated intervals.

bThese columns give the antilogs of the limits of parameters for which uniform random distributions of the log values were used.

Table 7.2. Ranges of Variation of Variables Derived in the Monte Carlo Analysis of the Improved Model.

			Percentiles					
Variable ^a	Mean	St. Dev.	Min. O	5	Median 50	95	Max. 100	
ζ	4.94	6.07	-5.48	-4.25	4.80	14.5	15.9	
σ	1.00	0.28	0.496	0.551	1.01	1.43	1.48	
R	0.657	1.21	0.00238	0.00557	0.126	3.08	8.99	
log R̃	-0.889	0.881	-2.62	-2.25	-0.90	0.49	0.95	
h	3.57	6.68	0.0105	0.0278	0.598	18.1	42.0	
log h	-0.185	0.891	-1.98	-1.56	-0.22	1.26	1.62	
ā	6.38	24.0	0.0000188	0.000112	0.0598	33.6	258.7	
log ā	-1.24	1.76	-4.73	-3.94	-1.22	1.53	2.41	
N	1680	5632	0.00387	0.0298	16.7	8893	53270	
log N	1.24	1.76	-2.41	-1.53	1.22	3 . 95	4.73	
B	31110	72750	34.1	117	3927	142500	727600	
log B	3.62	0 . 98	1.53	2.07	3.59	5.15	5.86	
κ	35350	44340	1042	2150	18590	127100	294400	
log κ	4.24	0.55	3.02	3.33	4.27	5.10	5.47	

 $^{^{\}rm a}{\rm S}$ tatistics are calculated for 500 plant stands with input parameters chosen randomly from the ranges in Table 7.1.




Figure 7.1. Monte Carlo analysis of the improved model for the overall size-density relationship. The data points represent 500 hypothetical plant stands characterized by parameter values chosen from uniform random distributions (Table 7.1). The solid line is the overall relationship log B = -0.53 log N + 4.28 (r^2 = 0.96, P < 0.0001, 95% CI for slope = [-0.56,-0.49]) fit by regressing log N against log \bar{w} and transforming the resulting equation into an expression for log B as a function of log N. The dotted lines with intercepts of 3.33 and 5.10 are parallel to this line and enclose 90% of the data points. The dashed line is the relationship log B = -0.49 + 3.99 fitted to data from 65 stands of 29 plant species by Gorham (1979).

model, since other geometric solids that might represent real VOIs (cones, pyramids, spheroids, rectangular solids, hexagonal solids, and any general prismatoids) show the same power relationships of volume and base area to a linear dimension of the base. The variation around the overall trend is related to the ranges of possible values for plant shape and the density of biomass in occupied space. The more complex model yields similar predictions, thus demonstrating that the heuristic results are not artifacts of assuming that all plants in a given stand are the same size and shape. Both models can be used to understand why the overall relationship obtains and both show that the origin of this relationship is different from the allometric factors that determine the slopes of single species thinning lines (Chapter 6).

The multi-species plots of Gorham (1979) and White (1980) address a fundamentally different question from an analysis of the self-thinning lines of individual populations. The former address the overall relationship determined by simple geometry while the latter addresses the time dynamics of space occupation by a single growing population. Because different phenomena are being examined, the existence of the overall relationship does not provide evidence for or against the self-thinning rule. It certainly does not show that self-thinning lines are insensitive to plant geometry. Instead, geometric differences are reflected by variations around the overall relationship.

This interpretation would still admit the hypothesis that the self-thinning trajectories of individual stands all approximate the

line $\log \bar{w} = -3/2 \log N + 4$ and so coincide with the overall relationship despite the differences in causality. This might seem justified on examining plots like that of White (1980) (Figure 7.2a), where thinning lines from many species seem to lie along a single line. However, in addressing real data, the log w-log N diagram gives a distorted view in which the appearance of linearity and high correlation is artificially enhanced while variations among the data are hidden (Chapter 4). When the thinning lines are examined in an unbiased log B-log N diagram (Figure 7.2b) an overall relationship with a slope of -1/2 is present, but individual self-thinning lines vary widely in both slope and position around that trend. The same type of distortion is evident in Gorham's analysis. The log W-log N plot (Figure 7.3a) shows the fitted line $\log w = -1.49 \log N + 3.99$ and dotted lines enclosing 75% of the 65 stands examined (as in Gorham 1979). This plot appears to admit little potential variation from the overall trend and seems to justify the conclusion self-thinning lines must fall in a narrow band paralleling the overall relationship. However, the log B-log N plot of the same data (Figure 7.3b) with the dotted lines enclosing 90% of the observed stands shows that individual self-thinning lines can vary widely in slope and position from the overall trend yet still fall within the 90% limits. Thus, the overall relationship is consistent with different thinning slopes and intercepts for particular populations. In fact, the hypothesis that all self-thinning lines closely approximate log B = -1/2 log N + 4 has already beendisproven (Chapter 5).




Figure 7.2. Previous analysis of the overall size-density relationship among thinning lines. (a) shows 65 previously cited self-thinning lines drawn by applying a reported thinning slope and intercepts (Table A.5) over the range of log N values covered by the thinning line (Table A.3). A similar plot of 31 thinning lines was constructed by White (1980). The dotted line is the overall relationship log \bar{w} = -1.49 log N + 3.99 (Gorham 1979). (b) shows the same 65 self-thinning lines and Gorham's line replotted in the log B-log N plane.




Figure 7.3. Gorham's analysis of the overall size-density relationship among stands of different species. (a) shows data for 65 fully-stocked stands of species as analyzed by Gorham (1979). The solid line is $\log \bar{w} = -1.49 \log + 3.99$ and the dotted lines enclose 75% of the observations. In (b) the same data and fitted line are shown in a log B-log N plot, along with dotted lines enclosing 90% of the plant stands.

Further verification of the models for the overall relationship can be obtained by comparing the predicted limits of variation around the relationship to the observed limits. This can be done by calculating a value κ for a sample of real plant stands using

$$\kappa = B \sqrt{N} , \qquad (7.14)$$

which can be interpreted as the intercept at log N = 0 of the line $\log B = -1/2 \log N + \log \kappa$ passing through the point ($\log N$, $\log B$). Log κ is in some ways similar to log K, the intercept of a self-thinning line. Both κ and K are constants of power equations and both are related to plant shape and the density of biomass in occupied space. However, the two constants are associated with fundamentally different phenomena. K relates to the time-dynamic self-thinning line determined by the particular allometry of a given stand while κ relates to the overall relationship between size and density among all plant stands. Different symbols are used here to preserve this important distinction. Values of κ were calculated here for the 1033 individual stands used to construct 75 self-thinning lines from experimental and field data (EFD) and the 3330 stands used to estimate 351 self-thinning lines for forestry yield table data (FYD). Chapter 5 details the data sources and self-thinning analyses.

Table 7.3 presents summary statistics for calculated values of κ and $\log \kappa$ for both the EFD and the FYD while Figure 7.4 gives histograms for the distributions of $\log \kappa$. The range of $\log \kappa$

Table 7.3. Statistical Distributions of κ and $\log \kappa$.

		Experimental and Field Data ^a				Forestry Yield Table data			
	Shoot biomass		Total t	Total biomass		All data sets		Bole Biomass	
Statistic	K	log ĸ	К	log ĸ	κ	log ĸ	κ	log κ	
n	700		333		1033		3330		
Mean	21830	4.01	32600	4.14	25300	4.05	7746	3.82	
Std. dev.	43350	0.47	70440	0.46	53800	0.47	4196	0.27	
CV (%)	199	12	216	11	213	12	54	7	
Std. err.	1638	0.02	3860	0.03	1674	0.01	72.7	0.005	
Skewness	5.00	0.78	3.92	1.37	4.73	0.93	0.87	-0.77	
Kurtosis	30.1	1.13	15.8	1.79	25.1	1.40	1.60	0.80	
Range	385300	3.49	479900	2.58	481000	3.58	30110	2.00	
Percentiles									
0 (Min.)	125	2.10	1257	3.10	125	2.10	308	2.49	
1 ` ′	1010	3.00	3049	3.48	1671	3.22	1045	3.02	
5	2815	3.45	4160	3.62	3074	3.49	2203	3.34	
10	3461	3.54	4593	3.66	3992	3.60	2798	3.45	
25	5115	3.71	7013	3.85	5437	3.74	4312	3.63	
50 (Median)	7690	3.89	11030	4.04	8508	3.93	7325	3.86	
75	17010	4.23	18290	4.26	17310	4.24	10460	4.02	
90	52560	4.72	55890	4.75	52830	4.72	13460	4.13	
95	82470	4.92	194500	5.29	91810	4.96	14820	4.17	
99	280100	5.45	394200	5.60	305700	5.49	18320	4.26	
100 (Max.)	385500	5.59	481200	5.68	481200	5.68	30420	4.48	

aThe three categories under this heading are based on the method of biomass measurement. "Shoot Biomass" includes thinning trajectories from biomass measurements of aboveground parts only. "Total Biomass" trajectories are based on biomass measurements that include aboveground and belowground parts. "All Data Sets" gives statistics for both of these groups combined.




Figure 7.4. Histograms of log κ . (a) shows the distribution of log κ values for the 1033 data points of the experimental and field data, while (b) gives the distribution of log κ for the 3330 data points of the forestry yield table data.

predicted by the heuristic model is $2.83 \le \log \kappa \le 5.43$ when the parameter ranges $600 \le d \le 40000$ and $2 \le \tau \le 12$ were used. Considering the extreme simplicity of the model and the crudity of the estimated parameter ranges, the agreement between the predicted range and the range observed for the EFD (2.10 < $\log \kappa$ < 5.68) is remarkable. Agreement is likewise good with the predicted range 3.01 < $\log \kappa$ < 5.47 of the more complex model. Values of $\log \kappa$ for the FYD are consistently lower than the model predictions because many tree parts were not included in the wood volumes from which stand biomasses were estimated. The close agreement of predicted and observed κ ranges, despite the crude model parameter estimates, supports the adequacy of the model as a representation of the overall relationship. The fact that simple models produce the overall relationship with very minimal information suggests that this relationship is, indeed, a symptom of a very simple geometric phenomenon and not an area of profound biological interest. Equation 7.4 can be transformed to log B = -1/2 log N + $\log(\tau d / \sqrt{\pi})$ and combined with equation 7.12 to give $\log \kappa =$ $\log(\tau d / \sqrt{\pi})$ and $\kappa = \tau d / \sqrt{\pi}$. According to this equation, κ can be interpreted as an composite measure of shape and density factors. The median values of $\log \kappa$ near 4 observed in both the EFD and FYD corresponds to a value of 5600 g/m^3 for the product τd .

The self-thinning rule has also been judged important in defining an ultimate thinning line which even constrains populations, such as clonal perennial herbs, that do not obey the self-thinning rule (Hutchings 1979). Theories have been proposed to

explain the different patterns of biomass-density dynamics of clonal herbs (Hutchings and Barkham 1976, Hutchings 1979, Pitelka 1984), but all assume that plant populations are eventually constrained to lie below a line of slope -3/2 in the log \bar{w} -log N plane. Hutchings (1979) proposed the line log \bar{w} = -3/2 log N + 4.3 while White (as cited in Pitelka 1984) has suggested that few plant populations exceed the size-density combinations defined by log \bar{w} = -3/2 log N + 5. White (1981) stated that the constraint imposed by this ultimate thinning line demonstrates the importance of the self-thinning rule, even for species whose stand dynamics do not necessarily trace a line of slope -3/2 in the log \bar{w} -log N plane.

The ultimate thinning line is explained by the simple models, which predict that all stands must lie below the line defined by extreme values of certain model parameters. In the heuristic model, stand biomass can be increased by raising plant height so that taller, more massive columns of biomass rest on the same piece of ground. Alternatively, more biomass can be packed into each cubic meter of biological volume by increasing d. Structural requirements and genetic limitations must ultimately constrain how tall and thin plants can become and still remain upright in the face of gravity, wind, and rain. Energetic constraints must limit the density of biomass per unit of ground area. No photoautotroph can pack more grams of living biomass onto a given surface than can be maintained by the conversion of the radiant energy falling on that surface. Also, natural selection would not favor plants that waste energy maintaining biomass beyond the amount that would give adequate

structural support, maximal capture of resources, and maximum reproduction.

The adequacy of the simple models for explaining the ultimate thinning line is supported by the good agreement between observed maximal values of $\log \kappa$ and the maximal values predicted from crude quesses about the maximum values of τ and d. The ninety-ninth percentile for $\log \kappa$ in the EFD is approximately 5.0, supporting the assertion that most stands fall below the line $\log w = -3/2 \log N + 5.0$ (White, as cited in Pitelka 1984). However, the slope of this line is a consequence of simple geometry while its intercept is determined by energetic and structural constraints. A thorough exploration and quantitative evaluation of the general constraints on structure and biomass accumulation across the entire plant kingdom would be of great biological interest (and very difficult), but the existence of the ultimate thinning line of slope -3/2 is unsurprising. The term "ultimate thinning line" is also a misnomer. The constraint is not related to self-thinning dynamics and should hold regardless of how growth and mortality proceed in a given stand. In fact, a stand can approach this line by decreasing in average weight or actually increasing the density of individuals per unit area (Pitelka 1984), rather than by a mortality process as implied by the term "thinning."

The overall relationship between size and density among stands of plants with different thinning behaviors also has significance for efforts to construct single species thinning lines. Since it is often difficult to obtain successive measurements of a given stand

through time, thinning lines have been commonly estimated from single measurements of stands of different ages. If the stands are grown under equivalent conditions and actually follow the same thinning trajectory, this method will give good results. However, if the stands are following different thinning lines, possibly due to genetic variations in plant geometry or differences in illumination or site fertility, then the thinning analysis will tend to reveal the overall relationship rather that the particular thinning trajectory of any of the stands. For example, Mohler et al. (1978) presented thinning data for Prunus pensylvanica and reported a thinning slope of -1.46 from fitting log \bar{w} against log N by principal component analysis. However, their data were collected from four sites in New Hampshire that were up to 100 km apart (Marks 1974). Elevations at these sites ranged from 340 to 570 m on north, south, and southeast facing slopes. At some sites, Prunus pensylvanica shared dominance with an important codominant Populus tremuloides. It seems likely that the large differences in slope, aspect, and other factors among sites many kilometers apart might mean that the stands would in fact follow different thinning trajectories. The analysis of Mohler et al. (1978) which gave a slope near -3/2 may then have revealed the overall relationship among stands following different thinning trajectories rather that the particular time trajectory of one of the stands.

CHAPTER 8

CONCLUSION

This study has examined the self-thinning rule from both theoretical and empirical perspectives and made major advances in explaining observed size-density relationships. These include:

(1) developing and analyzing explanatory models to produce testable hypothesis about underlying causes, (2) considering some important but largely unrecognized difficulties in testing the self-thinning rule and suggesting remedies for some problems, (3) completing a major analysis of self-thinning data to quantify the extent of variation in thinning line slope and intercept, (4) relating this variation to variations in plant allometry to verify that allometric factors are important in positioning the thinning line, and (5) developing models to explain the overall size-density relationship and evaluate its relevance to the self-thinning rule. The results of this study provide a basic explanatory theory for the rule and clarify its scientific importance.

Several hypotheses about the cause of the self-thinning rule were derived from two mathematical models: a spatially averaged two-equation model and a simulation model that details individual sizes, locations, and competitive interactions. These hypotheses included: (1) Self-thinning lines are linear because plant dimensions and plant size are related by power functions. (2) The slope of the thinning line is determined by the allometry between the area occupied by an individual and its weight (the allometric

hypothesis). (3) The intercept of the thinning line is complexly related to plant allometry, the density of biomass in occupied space, and the partitioning of contested areas among competing individuals. Although the allometric hypothesis is not new (Westoby 1976, Miyanishi et al. 1979), the present models fill a major gap by deriving the hypothesis from time dynamic models rather than from ad hoc geometric arguments. The dynamic models can explain aspects of the thinning rule that simple geometric arguments can not, such as the dual nature of the thinning line as a time trajectory and a constraint separating feasible biomass-density combinations from impossible ones (Hozumi 1977, White 1981) and the change in thinning slope under low illumination (Westoby 1977). The models developed here emphasize a simple biological interpretation rather than an exact mathematical solution and relate the slope and intercept of the thinning line to meaningful ecological parameters rather than to arbitrary constants; therefore, the models are heuristically useful and yield hypotheses that can be tested with measurable biological data.

Chapter 4 considered the difficulties of testing the self-thinning rule and charted an optimal course for this task.

Many potentially serious problems, such as the need for a posteriori data editing, do not presently have a good solution. Such problems have been previously discussed (Mohler et al. 1978, Hutchings and Budd 1981b), but their gravity has not been fully appreciated.

These limitations have been explored here so that their implications for the acceptance of the rule can be evaluated. Other aspects of

the analysis do have an optimal solution and much of the data supporting the self-thinning rule has not been analyzed in the best possible way. The problems include inattention to contradictory data, an invalid data transformation, inappropriate fitting techniques, and lack of hypotheses testing. In view of these problems, the results and interpretations of many studies are questionable.

Chapter 5 presented the most exhaustive analysis of the self-thinning rule to date. Biomass and density data from previous self-thinning studies were reanalyzed to ensure that the selections of data, curve-fitting methods, and statistical interpretations were the most appropriate, and additional data not previously analyzed in a self-thinning context were included to broaden the body of test data. The data from previous studies did not provide strong support for the self-thinning rule. Many data sets that were claimed to corroborate the rule did not show any significant relationship between size and density, or gave a thinning slope different from the thinning rule prediction. The present analysis is also the most complete description of the variation in thinning slope and intercept, which are more variable than currently accepted. greater variability was evidenced by by variations beyond currently accepted limits, by statistically significant differences in thinning slope and intercept among plant groups, and by significant correlations of slope and intercept with shade tolerance for forest trees.

The most important result of this study was the observation that variations in thinning slope are correlated with variations in plant allometry (Chapter 6). Such correlations were found in directly relating thinning slopes to allometric powers and in comparing among-group differences in thinning slope to among-group differences in allometric powers. This result supports the allometric hypothesis and verifies that the self-thinning rule is at least partly explainable by simple allometric arguments, despite previous failures to validate geometric models (Westoby 1976, Mohler et al. 1978, White 1981).

Finally, the slope of the overall size-density relationship among stands of different species was shown to be a trivial consequence of the geometry of packing objects onto a surface and of the limitation of the plant shape and the density of biomass in occupied space to biologically reasonable values. As such, the overall relationship represents a phenomenon different from the individual self-thinning line and does not provide evidence for or against the self-thinning rule.

The models suggest some additional analyses of self-thinning lines and tests of the hypotheses considered here. Some controlled experiments measuring thinning slope and several plant dimensions for species of differing allometric properties would be appropriate for verifying the evidence for the allometric hypothesis. It would also be instructive to gather data on plant allometry, density of biomass in occupied space, and the extent of overlap between

neighbors for several experimental populations to determine if these factors can explain the differences among thinning intercepts.

The results of this study have important implications for the scientific importance of the self-thinning rule. This rule does not qualify as an ecological law. The accepted constancy of the proposed law $\bar{w} = K N^{-3/2}$ (equivalently B = K N^{-1/2}) is based on a body of data, analysis, and interpretation that is flawed in many ways. The slopes and intercepts of thinning lines are actually quite variable and can be explained by simple geometric models. The variability in the thinning slope is particularly important because β is actually an exponent in the power equation $\beta = K N^{\beta}$ so small differences in $\boldsymbol{\beta}$ represent major differences in the predicted course of growth and mortality. Since the slopes of thinning lines do not take the same constant value, they do not provide evidence of the operation of a single quantitative law. Sprugel (1984) has also recently suggested that the log w-log N thinning slope of -3/2 for trees is actually the exception rather than the rule. The recommendation of Pickard (1983) seems appropriate: It would now be profitable to proceed beyond the self-thinning rule to the analysis of deeper questions.

LITERATURE CITED

LITERATURE CITED

- Aikman, D. P., and A. R. Watkinson. 1980. A model for growth and self-thinning in even-aged monocultures of plants. Annals of Botany 45:419-427.
- Ashe, W. W. 1915. Loblolly or North Carolina pine. Bulletin 24, North Carolina Geological and Economic Survey, Raleigh, North Carolina, USA.
- Baker, F. S. 1925. Aspen in the central Rocky Mountain region. Department Bulletin 1291, United States Department of Agriculture, Washington D.C., USA.
- Baker, F. S. 1949. A revised tolerance table. Journal of Forestry 47:179-181.
- Bakuzis, E. V., and H. L. Hansen. 1965. Balsam fir Abies balsamea (Linnaeus) Miller: a monographic review. University of Minnesota Press, Minneapolis, Minnesota, USA.
- Barkham, J. P. 1978. Pedunculate oak woodland in a severe environment: Black Tor Copse, Dartmoor. Journal of Ecology 66:707-740.
- Barnes, G. H. 1962. Yield of even-aged stands of western hemlock. Technical Bulletin 1273, United States Department of Agriculture, Washington, D.C., USA.
- Bazzaz, F. A., and J. L. Harper. 1976. Relationship between plant weight and numbers in mixed populations of <u>Sinapsis alba</u> (L.) and <u>Lepidium sativum</u> (L.). Journal of Applied Ecology 13:211-216.
- Beck, D. E. 1978. Growth and yield of white pine. Pages 72-90 in H. L. Williston and W. E. Balmer, editors. Proceedings: Symposium for the management of pines of the interior South. Technical Publication SA-TP2, United States Department of Agriculture, Washington, D.C., USA.
- Behre, C. E. 1928. Preliminary normal yield tables for second-growth western yellow pine in northeastern Idaho and adjacent areas. Journal of Agricultural Research 37:379-397.
- Bella, I. E. 1968. Jack pine yield tables for southeastern Manitoba. Publication 1207, Forestry Branch, Canadian Department of Fisheries and Forestry, Ottawa, Canada.

- Bella, I. E. 1971. A new competition model for individual trees. Forest Science 17:364-372.
- Black, J. N. 1960. An assessment of the role of planting density in competition between red clover (<u>Trifolium pratense</u> L.) and lucerne (<u>Medicago sativa</u> L.) in the early vegetative stage. Oikos 11:26-42.
- Boisen, A. T., and J. A. Newlin. 1910. The commercial hickories. Forest Service Bulletin 80, United States Department of Agriculture, Washington D.C., USA.
- Boudoux, M. 1978. Empirical yield tables for jack pine. Forestry Chronicle 54:216-219.
- Bruce, D. 1923. Preliminary yield tables for second-growth redwood. Bulletin 361, University of California Agriculture Experiment Station, Berkely, California, USA.
- Bruce, D., and F. X. Schumacher. 1950. Forest mensuration. McGraw-Hill Book Co., New York, New York, USA.
- Cannell, M. G. R. 1982. World forest biomass and primary production data. Academic Press, London, England.
- Cary, N. L. 1922. Sitka spruce: Its uses, growth, and management. Bulletin 1060, United States Department of Agriculture, Washington D.C., USA.
- Charles-Edwards, D. A. 1984. On the ordered development of plants 2. Self-thinning in plant communities. Annals of Botany 53:709-714.
- Christensen, N. L., and R. K. Peet. 1982. Secondary forest succession on the North Carolina Piedmont. Pages 230-345 in D. C. West, H. H. Shugart, and D. B. Botkin, editors. Forest succession: Concepts and application. Springer Verlag, New York, New York, USA.
- Clark, P. J., and F. C. Evans. 1954. Distance to nearest neighbor as a measure of spatial relationships in populations. Ecology 35:445-453.
- Curtis, R. O. 1971. A tree-area power function and related stand density measures for Douglas-fir. Forest Science 17:146-159.
- Dirzo, R., and J. L. Harper. 1980. Experimental studies on slug-plant interactions II. The effect of grazing by slugs on high density monoculturies of <u>Capsella bursa-pastoris</u> and <u>Poa annua</u>. Journal of Ecology 68:999-1011.

- Drew, T. J., and J. W. Flewelling. 1977. Some recent theories of yield-density relationships and their application to Monterrey pine plantations. Forest Science 23:517-534.
- Ernst, W. H.. 1979. Population biology of Allium ursinum in Northern Germany. Journal of Ecology 67:347-362.
- Eyre, F. H. 1944. Management of jack pine stands in the Lake states. Technical Bulletin 863, United States Department of Agriculture, Washington D.C., USA.
- Eyre, F. H., and P. Zehngraff. 1948. Red pine management in Minnesota. Circular 778, United States Department of Agriculture, Washington, D.C., USA.
- Foiles, M. W. 1956. Effects of thinning on a 55-year old western white pine stand. Journal of Forestry 54:130-132.
- Forbes, R. D. 1961. Forestry handbook. The Ronald Press Co., New York, New York, USA.
- Ford, E. D. 1975. Competition and stand structure in some even-aged plant monocultures. Journal of Ecology 63:311-333.
- Forsythe, G. E., M. A. Malcolm, and C. B. Mohler. 1977. Computer methods for mathematical computations. Prentice Hall, Inc., Englewood Cliffs, New Jersey, USA.
- Fox, G. D., and G. W. Kruse. 1939. A yield table for well-stocked stands of black spruce in northwestern Minnesota. Journal of Forestry 37:565-567.
- Frothingham, E. H. 1912. Second-growth hardwood forests in Connecticut. Forest Service Bulletin 96, United States Department of Agriculture Washington, D.C., USA.
- Frothingham, E. H. 1931. Timber growing and logging practice in the Southern Appalachian region. Technical Bulletin 250, United States Department of Agriculture, Washington D.C., USA.
- Furnas, R. E. 1981. A resource theory of self-thinning in plant populations. Ph.D. dissertation. Cornell University. Ithaca, New York, USA.
- Gevorkiantz, S. R., and W. A. Duerr. 1937. A yield table for northern hardwoods in the Lake states. Journal of Forestry 35:340-343.

- Gevorkiantz, S. R., and W. A. Duerr. 1939. Volume and yield of northern white cedar in the lake states: A progress report. Lake States Forest Experiment Station, Saint Paul, Minnesota, USA.
- Gold, H. J. 1977. Mathematical modeling of biological systems—an introductory guidebook. John Wiley and Sons, New York, New York, USA.
- Gorham, E. 1979. Shoot height, weight and standing crop in relation to density in monospecific plant stands. Nature 279:148-150.
- Haig, I. T. 1932. Second-growth yield, stand, and volume tables for the eastern white pine type. Technical Bulletin 323, United States Department of Agriculture, Washington D.C, USA.
- Hara, T. 1984. Modelling the time course of self-thinning in crowded plant populations. Annals of Botany 53:181-188.
- Harlow, W. M., E. S. Harrar, and F. M. White. 1978. Textbook of dendrology. McGraw-Hill Book Co., New York, New York, USA.
- Harper, J. L. 1977. Population biology of plants. Academic Press, New York, New York, USA.
- Harper, J. L., and J. White. 1971. The dynamics of plant populations. Pages 41-63 in den Boer, P. J., and G. R. Gradwell editors. Proceedings of the advanced study institute on dynamics of numbers in populations. Center for Agricultural Publishing. Oesterbeek, The Netherlands.
- Hatcher, R. J. 1963. A study of black spruce forests in northern Quebec. Publication 1018, Canadian Department of Forestry, Ottawa, Canada.
- Hickman, J. C. 1979. The basic biology of plant numbers.

 Pages 232-263 in O. T. Solbrig, S. Jain, G. B. Johnson, and
 P. H. Raven, editors. Topics in plant population biology,
 Columbia University Press, New York, New York, USA.
- Hillis, W. E., and A. G. Brown. 1978. Eucalypts for wood production. Commonwealth Scientific and Research Organization, Adelaide, Australia.
- Hindmarsh, A. C. 1980. LSODE and LSODI, two new initial value ordinary differential equation solvers. ACM (Association for Computing Machinery)-Signum Newsletter 15:10-11.

- Hiroi, T., and M. Monsi. 1966. Dry matter economy of <u>Helianthus</u> annuus communities growing at varying densities and light intensities. Journal of the Faculty of Science of Tokyo University 9:241-285.
- Hozumi, K. 1977. Ecological and mathematical considerations on self-thinning in even-aged pure stands I. Mean plant weight-density trajectory during course of self-thinning. Botanical Magazine, Tokyo 90:165-179.
- Hozumi, K. 1980. Ecological and mathematical considerations on self-thinning in even-aged pure stands II. Growth analysis of self-thinning. Botanical Magazine, Tokyo 93:146-166.
- Hozumi, K., and K. Shinozaki. 1970. Studies on the frequency distribution of the weight of individual trees in a forest stand II. Exponential distribution. Japanese Journal of Ecology 20:1-9.
- Hutchings, M. J. 1975. Some statistical problems associated with determinations of population parameters for herbaceous plants in the field. New Phytologist 74:349-363.
- Hutchings, M. J. 1979. Weight-density relationships in ramet populations of clonal perennial herbs, with special reference to the -3/2 power law. Journal of Ecology 67:21-33.
- Hutchings, M. J. 1983. Ecology's law in search of a theory. New Scientist 98:765-767.
- Hutchings, M. J., and J. P. Barkham. 1976. An investigation of shoot interactions in <u>Mercurialis perennis</u>, a rhizomatous perennial herb. Journal of Ecology 64:723-743.
- Hutchings, M. J., and C. S. Budd. 1981a. Plant competition and its course through time. Bioscience 31:640-645.
- Hutchings, M. J., and C. S. Budd, 1981b. Plant self-thinning and leaf area dynamics in experimental and natural monocultures. Oikos 36:319-325.
- Jacobs, M. R. 1979. Eucalypts for planting. FAO Forestry Series 11. Food and Agriculture Organization of the United Nations, Forestry Department, Rome, Italy.
- Jeffers, J. N. R. 1956. The yield of Hazel Coppice. Pages 12-30 in Utilization of Hazel Coppice. Bulletin 27, Forestry Commission, London, England.
- Johnson, P. L., editor. 1977. A ecosystem paradigm for ecology. Oak Ridge Associated Universities, Oak Ridge, Tennessee, USA.

- Johnson, N. L., and S. Kotz. 1970. Continuous univariate distributions-1. Houghton Mifflin Co., Boston, Massachusetts, USA.
- Jolicoeur, P. 1973. Imaginary confidence limits of the slope of the major axis of a bivariate normal distribution: A sampling experiment. Journal of the American Statistical Association 68:866-871.
- Jolicoeur, P. 1975. Linear regressions in fishery research: Some comments. Journal of the Fisheries Research Board of Canada 32:1491-1494.
- Jolicoeur, P., and A. A. Heusner. 1971. The allomery equation in the analysis of the standard oxygen consumption and body weight of the white rat. Biometrics 27:841-855.
- Jones, R. M. 1982. -3/2 power law. Bioscience 32:6.
- Kays, S., and J. L. Harper. 1974. The regulation of plant and tiller density in a grass sward. Journal of Ecology 62:97-105.
- Khil'mi, G. F. 1957. Theoretical forest biogeophysics. Israel Program for Scientific Translation, Ltd., Jerusalem, Israel.
- Kittredge, J., and S. R. Gevorkiantz. 1929. Forest possibilities of aspen lands in the Lake states. Technical Bulletin 60, University of Minnesota Agricultural Experimental Station, Saint Paul, Minnesota, USA.
- Koristan, C. F. 1931. Southern white cedar. Technical Bulletin 251, United States Department of Agriculture, Washington, D.C., USA.
- Koyama, H., and T. Kira. 1956. Frequency distribution of individual plant weight as affected by the interaction between plants (Intraspecific competition among higher plants VII). Journal of the Institute of Polytechnics, Osaka City University, Series D 7:73-94.
- Lindquist, J. L., and M. N. Palley. 1963. Empirical yield tables for young-growth redwood. Bulletin 796, California Agricultural Experiment Station, University of California, Berkeley, California, USA.
- Lonsdale, W. M., and A. R. Watkinson. 1982. Light and self-thinning. New Phytologist 90:431-445.
- Lonsdale, W. M., and A. R. Watkinson. 1983a. Plant geometry and self-thinning. Journal of Ecology 71:285-297.

- Lonsdale, W. M., and A. R. Watkinson. 1983b. Tiller dynamics and self-thinning in grassland habitats. Oecologia 60:390-395.
- Madansky, A. 1959. The fitting of straight lines when both variables are subject to error. Journal of the American Statistical Association 54:173-205.
- Malmberg, C., and H. Smith. 1982. Relationship between plant weight and density in mixed population of Medicago sativa and Trifolium pratense. Oikos 38:365-368.
- Marks, P. L. 1974. The role of pin cherry (<u>Prunus pensylvanica</u> L.) in the maintenance of stability in northern hardwood ecosystems. Ecological Monographs 44:73-88.
- Martin, G. L., A. R. Ek, and R. A. Monserud. 1977. Control of plot edge bias in forest stand growth simulation models. Canadian Journal of Forest Research 7:100-105.
- Marty, R. 1965. The mensurational characteristics of eastern white pine. Forest Service Research Note NE-40, Northeastern Forest Experiment Station, Upper Darby, Pennsylvania, USA.
- Mattoon, W. R. 1915. Shortleaf pine: Its economic importance and forest management. Bulletin 308, United States Department of Agriculture, Washington, D.C., USA.
- McArdle, R. E. 1930. The yield of douglas fir in the Pacific Northwest. Bulletin 201, United States Department of Agriculture, Washington, D.C., USA.
- McArdle, R. E., W. H. Meyer, and D. Bruce. 1949. The yield of douglas fir in the Pacific Northwest. Technical Bulletin 201, 1949 Revision, United States Department of Agriculture, Washington, D.C., USA.
- McCarthy, E. F. 1933. Yellow poplar characteristics, growth, and management. Technical Bulletin 356, United States Department of Agriculture Washington, D.C., USA.
- McIntosh, R. P. 1980. The background and some current problems of theoretical ecology. Synthese 43:195-255.
- Meyer, W. H. 1929. Yields of second growth spruce and fir in the Northeast. Technical Bulletin 142, United States Department of Agriculture, Washington, D.C., USA.
- Meyer, W. H. 1937. Yield of even-aged stands of sitka spruce and western hemlock. Technical Bulletin 544, United States Department of Agriculture, Washington, D.C., USA.

- Meyer, W. H. 1938. Yield of even-aged stands of ponderosa pine. Technical Bulletin 630, United States Department of Agriculture, Washington, D.C., USA.
- Miyanishi, K., A. R. Hoy, and P. B. Cavers. 1979. A generalized law of self-thinning in plant populations. Journal of Theoretical Biology 78:439-442.
- Mohler, C. L., P. L. Marks, and D. G. Sprugel. 1978. Stand structure and allometry of trees during self-thinning of pure stands. Journal of Ecology 66:599-614.
- Moran, P. A. P. 1971. Estimating structural and functional relationships. Journal of Multivariate Analysis 1:232-255.
- Obeid, M., D. Machin, and J. L. Harper. 1967. Influence of density on plant to plant variation in fiber flax, <u>Linum usitatissimum</u> L. Crop Science 7:471-473.
- O'Neill, R. V., and D. L. DeAngelis. 1981. Comparative productivity and biomass relations of forest ecosystems. Pages 411-449 in D. E. Reichle, editor. Dynamic properties of forest ecosystems. Cambridge University Press, Cambridge, England.
- Oshima, Y., M. Kimura, H. Iwaki, and S. Kuroiwa. 1958. Ecological and physiological studies on the vegetation of Mt. Shimagare I. Preliminary survey of the vegetation of Mt. Shimagare. Botanical Magazine, Tokyo 71:289-301.
- Patton, R. T. 1922. Red oak and White oak: A Study of growth and yield. Harvard Forest Bulletin 4, Harvard University, Cambridge, Massachusetts, USA.
- Peattie, D. C. 1950. A natural history of trees of eastern and central North America. Houghton Mifflin Co., Boston, Massachusetts, USA.
- Peattie, D. C. 1953. A natural history of western trees. Houghton Mifflin Co., Boston, Massachusetts, USA.
- Peet, R. K., and N. L. Christensen. 1980. Succession: A population process. Vegetatio 43:131-140.
- Perry, D. A. 1984. A model of physiological and allometric factors in the self-thinning curve. Journal of Theoretical Biology 106:383-401.
- Pickard, W. F. 1983. Three interpretations of the self-thinning rule. Annals of Botany 51:749-757.

- Pitelka, L. F. 1984. Application of the -3/2 power law to clonal herbs. American Naturalist 123:442-449.
- Pollard, D. F. W. 1971. Mortality and annual changes in distribution of above-ground biomass in an aspen sucker stand. Canadian Journal of Forest Research 1:262-266.
- Pollard, D. F. W. 1972. Above-ground dry matter production in three stands of trembling aspen. Canadian Journal of Forest Research 2:27-33.
- Popper, K. R. 1963. Conjectures and refutations: The growth of scientific knowledge. Harper and Row, New York, New York, USA.
- Puckridge, D. W., and C. M. Donald. 1967. Competition among wheat plants sown at a wide range of densities. Australian Journal of Agricultural Research 18:193-211.
- Rabinowitz, D. 1979. Bimodal distributions of seedling weight in relation to density of <u>Festuca paradoxa</u> Desv. Nature 277:297-298.
- Reineke, L. H. 1933. Perfecting a stand-density index for even-aged forests. Journal of Agricultural Research 46:627-638.
- Ricker, W. E. 1973. Linear regressions in fishery research. Journal of the Fisheries Research Board of Canada 30:409-434.
- Ricker, W. E. 1975. A note concerning Professor Jolicoeur's comments. Journal of the Fisheries Research Board of Canada 32:1494-1498.
- Riggs, D. S. 1963. The mathematical approach to physiological problems: A critical primer. M.I.T. Press, Cambridge, Massachusetts, USA.
- Ross, M. A., and J. L. Harper. 1972. Occupation of biological space during seedling establishment. Journal of Ecology 60:77-88.
- Schaffer, W. M., and E. G. Leigh. 1976. The prospective role of mathematical theory in plant ecology. Systematic Botany 1:209-232.
- Schlesinger, W. H. 1978. Community structure, dynamics and nutrient cycling in the Okefenokee cypress swamp forest. Ecological Monographs 48:43-65.
- Schlesinger, W. H., and D. S. Gill. 1978. Demographic studies of the chaparral shrub, <u>Ceanothus megacarpus</u>, in the Santa Ynez Mountains, California. <u>Ecology</u> 59:1256-1263.

- Schnurr, G. L. 1937. Yield, stand, and volume tables for even-aged upland oak forests. Technical Bulletin 560, United States Department of Agriculture, Washington, D.C., USA.
- Schumacher, F. X. 1926. Yield, stand, and volume tables for white fir in the California pine region. Bulletin 407, University of California Agriculture Experiment Station, Berkeley, California, USA.
- Schumacher, F. X. 1930. Yield, stand, and volume tables for Douglas fir in California. Bulletin 491, University of California Agriculture Experiment Station Berkeley, California.
- Schumacher, F. X., and T. S. Coile. 1960. Growth and yield of natural stands of the southern pines. T. S. Coile, Inc., Durham, North Carolina, USA.
- Show, S. B. 1925. Yield capacities of the pure yellow pine type on the east slope of the Sierra Nevada Mountains in California. Journal of Agricultural Research 31:1121-1135.
- Smith, J. H. G. 1968. Growth and yield in red alder in British Columbia. Pages 273-286 in J. M. Trappe, J. F. Franklin, R. F. Tarrout, and G. M. Hansen, editors. Biology of alder: Proceedings of a Symposium held at Northwest Scientific Association, Fortieth Annual Meeting, Pullman, Washington, April 14-15, 1967. United States Forest Service Pacific Northwest Forest Experiment Station, Portland, Oregon, USA.
- Snedecor, G. W., and W. G. Cochran. 1956. Statistical methods. The Iowa State University Press, Ames, Iowa, USA.
- Sokal, R. R., and F. J. Rohlf. 1981. Biometry. W. H. Freeman and Co., San Francisco, California, USA.
- Sprent, P., and G. R. Dolby. 1980. The geometric mean functional relationship. Biometrics 36:547-550.
- Sprugel, D. G. 1984. Density, biomass, productivity, and nutrient-cycling changes during stand development in wave-regenerated balsam fir forests. Ecological Monographs 54:165-186.
- Spurr, S. H., L. J. Young, B. V. Barnes, and E. L. Hughes. 1957. Nine successive thinnings in a Michigan white pine plantation. Journal of Forestry 55:7-13.
- Steill, W. M. 1976. White spruce: Artificial regeneration in Canada. Canadian Forestry Service, Department of the Environment, Ottawa, Canada.

- Steill, W. M., and A. B. Berry. 1973. Yield of unthinned red pine plantations at the Petawawa forest experiment station. Publication 1320, Canadian Forestry Service, Ottawa, Canada.
- Sterrett, W. D. 1915. The ashes: Their characteristics and management. Bulletin 229, United States Department of Agriculture, Washington, D.C., USA.
- Tadaki, Y., and T. Shidei. 1959. Studies on the competition of forest trees II. The thinning experiment on a small model stand of sugi (Cryptomeria japonica) seedlings. Journal of the Japanese Forestry Society 41:341-349.
- Taylor, R. F. 1934. Yield of second-growth western hemlock-sitka spruce stands in Southwestern Alaska. Technical Bulletin 412, United States Department of Agriculture, Washington, D.C., USA.
- Tepper, H. B., and G. T. Bamford. 1960. Thinning sweetgum stands in New Jersey. Forest Research Note NE-95, United States Department of Agriculture Northeastern Forest Experiment Station, Upper Darby, Pennsylvania, USA.
- Tseplyaev, V.P. 1961. The forests of the USSR. Israel Program for Scientific Translation, Ltd., Jerusalem, Israel.
- United States Department of Agriculture. 1929. Volume, yield, and stand tables for second-growth southern pines. Miscellaneous Publication 50. United States Department of Agriculture, Washington, D.C., USA.
- Vermont Agricultural Experiment Station. 1914. The management of second growth hardwoods in Vermont. Forest Service Publication 13, Vermont Agricultural Experiment Station, Burlington, Vermont, USA.
- Wahlenberg, W. G. 1946. Longleaf pine: Its use, ecology, regeneration, growth, and management. Charles Lathrop Park Forestry Foundation, Washington D.C., USA.
- Wahlenberg, W. G. 1955. Six thinnings in a 56-year-old pure white pine plantation at Biltmore. Journal of Forestry 53:331-339.
- Watkinson, A. R. 1980. Density-dependence in single-species populations of plants. Journal of Theoretical Biology 80:344-357.
- Watkinson, A. R. 1984. Yield-density relationships: the influence of resource availability on growth and self-thinning in populations of Vulpia fasciculata. Annals of Botany 53:469-482.

- Watkinson, A. R., W. M. Lonsdale, and L. G. Firbank. 1983. A neighborhood approach to self-thinning. Oecologia 56:381-384.
- Westoby, M. 1976. Self-thinning in <u>Trifolium subterraneum</u> not affected by cultivar shape. Australian Journal of Ecology 1:245-247.
- Westoby, M. 1977. Self-thinning driven by leaf area not by weight. Nature 265:330-331.
- Westoby, M. 1981. The place of the self-thinning rule in population dynamics. American Naturalist 118:581-587.
- Westoby, M. and L. Brown. 1980. The effect of clipping on self-thinning in <u>Trifolium pratense</u>. Australian Journal of Ecology 5:407-409.
- Westoby, M., and J. Howell. 1981. Self-thinning: The effect of shading on glasshouse populations of silver beets (Beta vulgaris). Journal of Ecology 69:359-365.
- Westoby, M., and J. Howell. 1982. Self-thinning in <u>Trifolium</u> subterraneum populations transferred between full daylight and shade. Journal of Ecology 70:615-621.
- Wethey, D. S. 1983. Intrapopulation variation in growth of sessile organisms: Natural populations of the intertidal barnacle Balanus balanoides. Oikos 40:14-23.
- White, J. 1977. Generalization of self-thinning of plant populations Comment on Westoby(1977). Nature 268:337.
- White, J. 1980. Demographic factors in populations of plants, Pages 21-48 in O. T. Solbrig, editor. Demography and Evolution in Plant Populations. University of California Press, Berkeley, California, USA.
- White, J. 1981. The allometric interpretation of the self-thinning rule. Journal of Theoretical Biology 89:475-500.
- White, J., and J. L. Harper. 1970. Correlated changes in plant size and number in plant populations. Journal of Ecology 58:467-485.
- White, J., and S. J. Gould. 1965. Interpretation of the coefficient in allometric equations. American Naturalist 99:5-18.

- Whittaker, R. H. and G. M. Woodwell. 1968. Dimension and production relations of trees and shrubs in the Brookhaven Forest, New York. Journal of Ecology 56:1-25.
- Williamson, A. W. 1913. Cottonwood in the Mississippi Valley. Bulletin 24, United States Department of Agriculture, Washington D.C., USA.
- Winters, R. K., and J. G. Osborne. 1935. Growth and yield of second-growth red gum in fully stocked stands on alluvial lands in the South. Occasional Paper 54, Southern Forest Experiment Station, New Orleans, Louisiana, USA.
- Yoda, K., T. Kira, H. Ogawa, and K. Hozumi. 1963. Self-thinning in overcrowded pure stands under cultivated and natural conditions. (Intraspecific competition among higher plants XI). Journal of the Institute of Polytechnics, Osaka City University, Series D 14:107-129.

APPENDIXES

APPENDIX A EXPERIMENTAL AND FIELD DATA

Table A.1. Sources of Experimental and Field Data

Id. Code ^a	Species	Group ^b	Study Type ^C		Conditions ^e	Reference ^f
1	Prunus pensylvanica	TTmA	N	А		Marks 1974, Tabs. 2,3
5	Populus tremuloides	TTmA	N	Α		Pollard 1971, Tab. 1
						Pollard 1972, Tab. 2
7	<u>Triticum</u> 9	HM	F	TD		Puckridge and Donald 1967, Fig. 1, Tab. 2;
	0	++ 0		-	0.7	White and Harper 1970, Fig. 5
8	Pinus strobus	TTmG	Р	Ţ	Plot	Spurr et al. 1957, Tabs. 1,2
10	Trifolium subterraneum	HD TT0	E	Ţ		Westoby and Howell 1982, Tab. 1
14	Liquidambar styraciflua	TTmA	N	T		Tepper and Bamford 1960, Tab. 1
15	Erigeron canadensis	HD	N	N		Yoda et al. 1963, Fig. 14
16	Plantago asiatica	HD	N	N		Yoda et al. 1963, Fig. 13
17 18	Amaranthus retroflexus	HD	N	N		Yoda et al. 1963, Fig. 16
19	Ambrosia artimestifolia	HD TTC	N	N		Yoda et al. 1963, Fig. 15
20	Abies sachalinensis Chenopodium album	TTmG HD	N N	N N		Yoda et al. 1963, Fig. 22 Yoda et al. 1963, Fig. 17
21	Erigeron canadensis	HD	F	T		Yoda et al. 1963, Fig. 17
22	Prunus pensylvanica	TTmA	N N	Å		Mohler et al. 1978
23	Abies balsamea	TTmG	N	Â		Mohler et al. 1978
24	Abies sachalinesis	TTmG	N	N		Hozumi and Shinozaki 1970, Tab. 2A
26	Betula ⁹	TTmA	N	N		Yoda et al. 1963, Fig. 23
27	Trifolium praetense	HD.	Ë	Ť		Westoby and Brown 1980, Fig. 2
28	Trifolium praetense	HD	Ē	Ť	Illumination	Hutchings and Budd 1981b, Fig. 4
29	Fagopyrum esculentum	HD	F	ΤĎ	1714/11/14/17/01	Yoda et al. 1963, Fig. 21
30	Trifolium praetense	HD	F	TD		Black 1960, Fig. 1, Tabs. 1-3
31	Medicago sativa	HD	F	ŤĎ		Black 1960, Fig. 1, Tabs. 1-3
32	Fagopyrum esculentum	HD	Ε	TD	Fertility	Furnas 1981, Fig. 4.1
33	Beta vulgaris	HD	Ε	TD	.,,	Furnas 1981, Fig. 4.2
34	Beta vulgaris	HD	E	T		Furnas 1981, Fig. 4.5
35	Trifolium subterraneum	HD	Ε	T		Westoby 1976, Fig. 1
36	Ceanothus megacarpus	S	N	Α		Schlesinger and Gill 1978, Fig. 3, Tab. 1
38	Lolium perenne	HM	Ε	TD	Illumination	Kays and Harper 1974, Fig. 4
39	Betula ⁹	TTmA	N	N	Site	Hozumi and Shinozaki 1970, Tab. 2B,2C
40	Tagetes patula	HD	Ε	ΤĐ		Ford 1975, Fig. 12
41	Quercus robur	TTmA	N	Α		Barkham 1978, Tab. 5
43	Beta vulgaris	HD	Ε	T	Illumination	Westoby and Howell 1981, Fig. 1
44	Helianthus annuus	HD	F	TD	Experiment; Illumination	Hiroi and Monsi 1966, Figs. 1,5,6, Tab. 2
45	Picea mariana	TTmG	N	Α		Hatcher 1963, Tab. 11
48	Corylus avellana	TTmA	С	А		Jeffers 1956, Tabs. 16-18
49	Fagopyrum esculentum	HD	Ε	TD		Furnas 1981, Fig. 4.6
50	Brassica juncea	HD	F	TD	Illumination	Furnas 1981, Fig. 4.10
51	Sinapsis alba and	HD	Ε	T		Bazzaz and Harper 1976, Fig. 1
E 2	Leoidium sativum	TT0	,-	-		Todaki and Shidoi 1050 Taba 3 2
52	Cryptomeria japonica	TTmG	£ E	T T		Tadaki and Shidei 1959, Tabs. 1-3
53	Brassica napus and	HD	Ľ	1		White and Harper 1970, Fig. 2
	Raphanus sativus					

Table A.1. (continued)

Ref							
Pinus monticola		Species	Group ^b			Conditions ^e	Reference ^f
Trifolium praetense	54		TTmG	N	Т		Foiles 1956, Tabs. 1,2
Malmberg and Smith 1982, Fig. 4	5.5		HD	F	т		Malmherg and Smith 1982, Fig. 4
Medicago sativa and							
Beta vulgaris and		Medicago sativa and					
Taxodium distichum	58	Beta vulgaris and	HD	F	T		Furnas 1981, Figs. 4.8,4.9
Abies verticity Abies Ab	80		TTmG	N	N		Schlesinger 1978, Fig. 3C
Pinus taeda		Abies veitchii and					
Pinus densiflora	82		TTmG	N	AD		
Festuca pratensis	84	Pinus densiflora	TTmG	N	N		
Chicorium endivium	86	Festuca pratensis	HM	Ε	T		Lonsdale and Watkinson 1983a, Figs. 1,20
Capsella bursa-pastoris		Agrostemma githago	HD		T		Lonsdale and Watkinson 1983a, Figs. 1,20
Poa annua	88	Chicorium endivium	HD	Ε	T		Lonsdale and Watkinson 1983a, Figs. 1,2C
Column Devence HM E TO Illumination Lonsdale and Watkinson 1982, Fig. 1	89	Capsella bursa-pastoris	HD		T	Grazing level	Dirzo and Harper 1980, Fig. 3
Colium perenne HM E T Illumination Lonsdale and Watkinson 1982, Fig. 5			HM		T	Grazing level	Dirzo and Harper 1980, Fig. 3
Pinus strobus		Lolium perenne	HM		TD	Illumination	Lonsdale and Watkinson 1982, Fig. 1
Wahlenberg 1955, Fig. 2						Illumination	
95 Eucalyptus obliqua TE P T Hillis and Brown 1978, Tab. 10.17 96 Eucalyptus pilularis TE N T Hillis and Brown 1978, Tab. 10.21 97 Eucalyptus regnans TE N T Hillis and Brown 1978, Tab. 10.24 98 Eucalyptus regnans TE P T Site Index Hillis and Brown 1978, Tab. 10.26 99 Eucalyptus deglupta TE P T Site Index Hillis and Brown 1978, Tab. 10.26 99 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.26 99 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.24 101 Eucalyptus deglupta TE P T Cannell 1982, Page 11 102 Pinus taeda TTMA N A Cannell 1982, Page 31 103 Fagus sylvatica TTMA N A Cannell 1982, Page 33 104 Fagus sylvatica TTMA N A Cannell 1982, Page 33	93	Pinus strobus	TTmG	Р	T		
Fagus sylvatica TimA N A Cannell 1982, Page 33	95	Eucalyptus obliqua	ΤE	Р	Т		
97 Eucalyptus regnans TE N T Hillis and Brown 1978, Tab. 10.24 98 Eucalyptus regnans TE P T Site Index Hillis and Brown 1978, Tab. 10.26 99 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.26 101 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.26 101 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.26 101 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.26 102 Pinus daglous deglupta TE P T Cannell 1982, Page 11 102 Pinus taeda TImA N A Cannell 1982, Page 32 103 Fagus sylvatica TImA N A Cannell 1982, Page 31 104 Fagus sylvatica TImA N A Cannell 1982, Page 33 105 Fagus sylvatica TImA N A Cannell 1982, Page 33 106 Ac	96	Eucalyptus pilularis	TE	N			
98 Eucalyptus regnans TE P T Site Index Hillis and Brown 1978, Tab. 10.26 99 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.34 101 Eucalyptus deglupta TE P T Hillis and Brown 1978, Tab. 10.34 101 Eucalyptus deglupta TE P A Cannell 1982, Page 11 102 Pinus taeda TTm6 N A Cannell 1982, Page 320 103 Fagus sylvatica TTmA N A Cannell 1982, Page 33 104 Fagus sylvatica TTmA N A Cannell 1982, Page 58 105 Fagus sylvatica TTmA N A Cannell 1982, Page 58 106 Acer spicatum TTmA N A Cannell 1982, Page 72 106 Acer spicatum TTmA N A Cannell 1982, Page 33 109 Mixed hardwoods TTmA N A Cannell 1982, Page 34 E. Canada) E. Canada TTmA	97		TE	N	T		
101 Eucalyptus grandis TE	98	Eucalyptus regnans	TE	Р	T	Site Index	
Pinus taeda	99	Eucalyptus deglupta	TE	Р	T		Hillis and Brown 1978, Tab. 10.34
103 Fagus Sylvatica TImA N A Cannell 1982, Page 31 104 Fagus Sylvatica TImA N A Cannell 1982, Page 58 105 Fagus Sylvatica TImA N A Cannell 1982, Page 58 106 Acer Spicatum TImA N A Cannell 1982, Page 72 107 Acer Spicatum TImA N A Cannell 1982, Page 33 109 Mixed hardwoods TImA N A Cannell 1982, Page 34 E. Canada Cannell 1982, Page 34 E. Canada Cannell 1982, Page 34 110 Pinus banksiana TImG N A Cannell 1982, Page 44 111 Shorea robusta TITA P A Cannell 1982, Page 79 112 Cyclobalanopsis TImA C A Cannell 1982, Page 108 113 Tectona grandis TITA P A Cannell 1982, Page 108 114 Abies veitchii TImG N A Cannell 1982, Page 129 115 Camellia japonica TImA N D Cannell 1982, Page 19 116 Eucalyptus obliqua TE N A Cannell 1982, Page 14 117 Abies firma and TImG N A Cannell 1982, Page 124 118 Cannell 1982, Page 124 119 Abies firma and TImG N A Cannell 1982, Page 124 110 Cannell 1982, Page 124 111 Abies firma and TImG N A Cannell 1982, Page 124 112 Cannell 1982, Page 124 113 Cannell 1982, Page 124 114 Abies firma and TImG N A Cannell 1982, Page 124 115 Cannell 1982, Page 124 116 Cannell 1982, Page 124 117 Cannell 1982, Page 124 118 Cannell 1982, Page 124 119 Cannell 1982, Page 124 110 Cannell 1982, Page 124 111 Cannell 1982, Page 124 112 Cannell 1982, Page 124 113 Cannell 1982, Page 124 114 Cannell 1982, Page 124 115 Cannell 1982, Page 124 116 Cannell 1982, Page 124 117 Cannell 1982, Page 124 118 Cannell 1982, Page 124 119 Cannell 1982, Page 124 110 Cannell 1982, Page 124 111 Cannell 1982, Page 124 112 Cannell 1982, Page 124 113 Cannell 1982, Page 124 114 Cannell 1982, Page 124 115 Cannell 1982, Page 124 116 Cannell 1982, Page 124 1		Eucalyptus grandis	TE	Р	Α		Cannell 1982, Page 11
Tima			TTmG	N	Α		
Tima		Fagus sylvatica	TTmA	N	Α		
106 Acer spicatum		Fagus sylvatica	TTmA	N	Α		
Mixed hardwoods TImA N A Cannell 1982, Page 34		Fagus sylvatica	TTmA	N	Α		Cannell 1982, Page 72
E. Canada) 110				N			
111 Shorea robusta TTrA P A Cannell 1982, Page 79	109		TTmA	N	А		Cannell 1982, Page 34
112 Cyclobalanopsis myrsinaefolia TTmA C A Cannell 1982, Page 108 113 Tectona grandis TTrA P A Cannell 1982, Page 83 114 Abies veitchii TTmG N A Cannell 1982, Page 129 115 Camellia japonica TTmA N D Cannell 1982, Page 99 116 Eucalyptus obliqua TE N A Cannell 1982, Page 14 117 Abies firma and TTmG N A Cannell 1982, Page 124	110	Pinus banksiana	TTmG	N	А		Cannell 1982, Page 44
myrsinaefolia myrsinaefolia 113 Tectona grandis TTrA P A Cannell 1982, Page 83 114 Abies veitchii TTmG N A Cannell 1982, Page 129 115 Camellia japonica TTmA N D Cannell 1982, Page 99 116 Eucalyptus obliqua TE N A Cannell 1982, Page 14 117 Abies firma and TTmG N A Cannell 1982, Page 124	111	Shorea robusta	TTrA	Р	Α		Cannell 1982, Page 79
113 Tectona grandis	112		TTmA	С	А		Cannell 1982, Page 108
114 Abies veitchii TTmG N A Cannell 1982, Page 129 115 Camellia japonica TTmA N D Cannell 1982, Page 99 116 Eucalyptus obliqua TE N A Cannell 1982, Page 14 117 Abies firma and TTmG N A Cannell 1982, Page 124	113		TTrA	Р	А		Cannell 1982, Page 83
115 Camellia japonica TTmA N D Cannell 1982, Page 99 116 Eucalyptus obliqua TE N A Cannell 1982, Page 14 117 Abies firma and TTmG N A Cannell 1982, Page 124							
116 <u>Eucalyptus obliqua</u> TE N A Cannell 1982, Page 14 117 <u>Abies firma</u> and TTmG N A Cannell 1982, Page 124							
117 Abies firma and TTmG N A Cannell 1982, Page 124					-		
isaga siebuluti		Tsuga sieboldii					

Table A.l. (continued)

Id. Code ^a	Species	Group ^b	Study Type ^C		Conditions ^e	Reference ^f
118	Eucalyptus tereticornis	TE	Р	А	•	Cannell 1982, Page 78
119	Abies sachalinensis	TTmG	Р	Α		Cannell 1982, Page 126
120	Cryptomeria japonica	TTmG	Р	Α		Cannell 1982, Page 151
121	Betula ⁹	TTmA	N	Α		Cannell 1982, Page 238
122	Cryptomeria japonica	TTmG	Р	Α		Cannell 1982, Page 146
123	Castanea sativa	ΤŢmA	С	Α		Cannell 1982, Page 239
124	Abies9	ΤΤmG	N	Α		Cannell 1982, Page 133
25	Pinus sylvestris	TTmG	Р	Α		Cannell 1982, Page 243
26	Quercus pubescens	TTmA	N	Α		Cannell 1982, Page 68
28	Pinus pumila	TTmG	N	Α		Cannell 1982, Page 176
130	Pinus nigra	TTmG	Р	Α		Cannell 1982, Page 243
131	Alnus rubra	TTmA	N	Α		Cannell 1982, Page 252
132	Cryptomeria japonica	TTmG	Р	Α		Cannell 1982, Page 146
33	Populus deltoides	T TmA	N	Α		Cannell 1982, Page 266
135	Pinus banksiana and mixed hardwoods	TTmG	N	А		Cannell 1982, Page 300
36	Picea abies	TTmG	Р	Α		Cannell 1982, Page 73
37	Picea abies	TTmG	N	Α		Cannell 1982, Page 361
1 38	Alnus incana and Salix9	TTmA	N	А		Cannell 1982, Page 251

^aAn arbitrary number assigned to facilitate cross-referencing among data tables.

bone of seven categories. HM = herbaceous monocots; HD = herbaceous dicots; S = shrubs; TTmA = trees, temperate angiosperms; TTmG = trees, temperate gymnosperms; TTrA = trees, tropical angiosperms; and TE = trees of genus Eucalyptus.

CLevel of experimental control. N = natural field populations, P = tree plantations, C = coppiced trees, E = outdoor experiments, and E = greenhouse or light chamber experiments.

dIndicates how observations of stands at different densities and biomasses were generated. T = repeated measurements of a single population (time series), A = observations of populations of different ages (age series), D = observations of populations started at different initial densities (density series), N = not specified.

eNatural conditions or experimental treatments that may have affected thinning line slope or position, so that separate thinning lines were fitted to subsets of the data points from a single source. "Plot" and "site" both refer to different physical locations at which plant populations were observed. "Illumination" means that different populations received to different intensities of light. "Grazing level" and "fertility" likewise indicate that different populations were subjected to different levels of the indicated stress or resource. "Experiment" refers to different experiments that were considered separately in the reference. "Site index" is a measure of site quality used by foresters, here the height in meters of the largest trees when the stand is 20 years old. See the references given for further details.

 ${}^{
m f}$ Tabular data were taken from the indicated table numbers. Figure numbers refer to graphs from which data were reconstructed.

9Generic name only indicates that the specific name was not given in the reference, or that more than one species from the genus were present.

Table A.2. Self-thinning Lines Fit to Experimental and Field Data

		No. of					PCA Self-ti	ninning	Line ^e	
		Points					Slope		Intercept	
Id. Code ^a	Condition ^b	n _T	n	r ²	р ^d	ĝ	95% CI ^f	â	95% CI	p _{est} q
1A 1T 5A 7A 8A 10A 14A 15T 16T 17T 18T 20T 21T 22T 22T 24T 24T 26A 27A	Lot 28 Lot 2C Full light	4 4 7 33 9 8 10 6 47 38 21 28 27 28 27 30 53 29 9	3 3 4 5 7 7 10 4 31 38 17 43 26 13 13 34 27 27	0.983 0.980 0.941 0.985 0.986 0.937 0.836 0.971 0.930 0.926 0.898 0.841 0.929 0.987 0.395 0.654 0.839	0.0829 0.0911 0.0300* 0.0001* 0.0001* 0.0001* 0.0015* 0.0001* 0.0001* 0.0001* 0.0001* 0.0001* 0.0001* 0.0001* 0.0001* 0.0001* 0.0001*	-0.429 -0.424 -0.264 -0.362 -0.724 -1.116 -0.473 -0.667 -0.472 -0.588 -0.539 -0.405 -1.038 -0.410 -0.204 -0.465 -1.033	[-0.484, -0.065]* [-0.448, -0.280]* [-0.488, -0.628]* [-1.278, -0.976]* [-0.660, -0.310] [-1.117, -0.348] [-0.518, -0.428] [-0.518, -0.428] [-0.703, -0.483] [-0.776, -0.535]* [-0.482, -0.332]* [-0.482, -0.332]* [-1.121, -0.962]* [-0.273, -0.138]* [-0.611, -0.234] [-0.273, -0.138]* [-0.695, -0.104] [-1.406, -0.760]*	3.77 3.84 3.65 3.85 3.78 4.60 3.71 4.36 3.98 4.06 5.15 5.15 3.80 4.16 5.15 3.82 3.83	[3.61, 3.67]	0.350 0.351 0.396 0.367 0.290 0.236 0.339 0.300 0.303 0.305 0.305 0.325 0.325 0.341 0.345 0.341 0.345
28A 28A 28A 29T 30A 31A	Low L.I. Med. Ł.I. High L.I.	5 10 14 14 25 22 22	2 7 8 7 9	1.000 0.394 0.137 0.861 0.954 0.137 0.180	0.1310 0.3666 0.0026* <0.0001* 0.3266 0.1938	-0.313 -0.851 -1.192 -0.627 -0.690 -0.608 -0.434	[-0.978, -0.360] [-0.836, -0.561]*	4.33 4.62 6.53 4.78 5.08 4.89 3.95	[3.95, 5.86] [4.68, 5.55]	0.381 0.270 0.228 0.307 0.296 0.311 0.349
32T 32T 33T 34T 35A 36A	20% N.C. 100% N.C.	7 7 10 10 22	5 10 10 13	0.003 0.024 0.592 0.645 0.716 0.226	0.9289 0.8027 0.0092* 0.0052* 0.0003* 0.1653	-0.024 -0.029 -1.335 -2.304 -0.622 -0.188	[-0.430, 0.362] [-3.355, -0.648]* [-5.478, -1.348]* [-0.928, -0.382] [-0.514, 0.105]	2.86 3.21 6.38 9.93 5.17 3.64	[1.88, 4.59] [4.54, 11.80] [7.08, 19.38] [4.33, 6.23] [3.49, 3.80]	0.488 0.486 0.214 0.151 0.308 0.421
38A 38A 39A 39A 39A 40A 41A 43A	30% L.I. 70% L.I. 100% E.I. Kitami Shirogane	20 20 20 6 7 14 5	7 8 8 5 7 10 5 7	0.054 0.422 0.549 0.752 0.043 0.343 0.930 0.957	0.6144 0.0814 0.0354* 0.0569 0.6554 0.0750 0.0081*	0.055 -0.452 -0.324 -0.429 0.051 -0.414 -0.618	[-0.227, 0.347] [-1.429, 0.112] [-0.674, -0.034] [-1.118, 0.030] [-0.247, 0.359] [-1.146, 0.069] [-1.005, -0.331]	1.98 4.16 3.79 4.85 4.69 4.36 3.84	[1.03, 2.89] [2.36, 7.28] [2.88, 4.89] [4.52, 5.07] [4.59, 4.80] [2.74, 6.82] [3.43, 4.15]	0.529 0.344 0.378 0.350 0.527 0.354 0.309
43A 43A	25% L.I. 37% L.I.	10 10	7 7 7	0.957 0.916 0.940	0.0001* 0.0007* 0.0003*	-0.662 -0.692 -0.668	[-0.839, -0.509]* [-0.973, -0.470] [-0.886, -0.486]	4.79 5.12 5.09	[4.17, 5.50] [4.22, 6.25] [4.39, 5.94]	0.301 0.296 0.300

Table A.2. (continued)

							PCA Self-th	inning Li	ine ^e	
		No. Poin					Slope	Ir	ntercept	
Id. Code ^a	Condition ^b	nŢ	n	r ²	Рq	ĝ	95% CI ^f	â	95% CI	p q est
43A 43A 44A 44A	55% L.I. 100% L.I. Ex. 2-60% L.I. Ex. 2-100% L.I.	11 9 27 27	8 7 4 4	0.934 0.698 0.838 0.780	<0.0001* 0.0192* 0.0844 0.1168	-0.649 -0.648 -0.868 -0.926	[-0.838, -0.487] [-1.415, -0.197]	5.22 [5.30 [4.69 5.14		7] 0.303 0.268 0.260
44T 44T 44T 44T 44T	Ex. 1-23% L.I. Ex. 1-60% L.I. Ex. 1-100% L.I. Ex. 2-60% L.I. Ex. 2-100% L.I.	29 32 32 27 27	10 8 6 4 4	0.157 0.834 0.364 0.842 0.764	0.2576 0.0015* 0.2048 0.0826 0.1259	-0.106 -0.313 -0.147 -0.867 -0.968	[-0.318, 0.098] [-0.460, -0.176]* [-0.453, 0.133]	2.57 [3.60 [3.42 [4.71 5.25	2.14, 3.0 3.28, 3.9 2.69, 4.2	5] 0.381
45A 48A 49T 50T 50T 51A 52A	15% L.I. 100% L.I.	32 7 12 12 22 30	25 7 10 12 22 24 3	0.391 0.589 0.089 0.391 0.021 0.131 0.829	0.0008* 0.0439* 0.4031 0.0296 0.5217 0.0819	-0.475 -0.721 -0.075 0.284 0.038 -0.304 -1.594	[-0.775, -0.232] [-2.545, -0.052] [-0.283, 0.126] 0.037, 0.568] [-0.085, 0.162] [-0.741, 0.048]	3.77 [3.72 [3.03 [0.97 [2.27 [3.64 [6.52	3.66, 3.6 3.53, 4.2 2.37, 3.7 0.19, 1.6 1.91, 2.6 2.45, 5.1	5] 0.291 2] 0.465 5] 0.698 2] 0.520
53T 54A 55A 56A 57A 58T		36 4 17 17 21 32	29 4 11 11 13 23	0.547 0.978 0.333 0.766 0.559	<pre>0.2711 <0.0001* 0.0113* 0.0633 0.0004* 0.0033* 0.4304</pre>	-0.391 -1.454 -1.879 -0.780 -1.017 -0.088	[-0.541, -0.255] [-2.504, -0.924]* [-0.596, 18.075] [-1.186, -0.488] [-2.011, -0.519]* [-0.334, 0.147]	4.31 [3.35 [3.87, 4.8 2.63, 3.7 -66.11, 4.8 4.65, 7.3 4.70, 10.3 1.88, 3.1	0] 0.35° 1] 0.204 1] 0.174 2] 0.281 2] 0.248
80A 81A 82A 84A 86A 87A		18 7 42 286 18 15	5 4 17 243 15 12	0.876 0.886 0.468 0.248 0.582 0.934	0.0192* 0.0588 0.0025* <0.0001* 0.0009* <0.0001*	-0.725 -0.615 -0.305 -0.529 -0.985 -0.651	[-1.506, -0.277] [-2.769, 0.123] [-0.499, -0.130]* [-0.653, -0.417] [-1.721, -0.562]* [-0.781, -0.535]*	4.23 [4.06 [4.21 [3.73 [7.54]	3.70, 4.5 3.87, 4.6 4.11, 4.3 3.61, 3.8 5.67, 10.8 5.05, 5.9	4 0.290 4 0.310 0 0.383 4 0.327 0 0.252
88A 89A 39A 90A 90A	Grazed Ungrazed Grazed Ungrazed	18 5 5 5 5	15 3 5 2 2	0.864 0.995 0.948 1.000 1.000	<0.0001* 0.0447* 0.0051*	-0.993 -0.612 -0.618 -0.870 -0.479	[-1.265, -0.779]* [-1.592, -0.089 [-0.932, -0.372	6.52 { 4.14 [4.28 [5.67 4.36	5.75, 7.5 2.09, 7.9 3.36, 5.4	9] 0.310 5] 0.309 0.267 0.338
91A 91A 91T 91T 92A 92A 92A 92A 93A	17% L.I. 100% L.I. 17% L.I. 100% L.I. 17% L.I. 23% L.I. 44% L.I.	25 25 20 20 5 5 7 8	13 10 13 10 5 5 5	0.464 0.908 0.293 0.854 0.760 0.776 0.786 0.712 0.955	0.0104* <0.0001* 0.0559 0.00539 0.0483* 0.0450* 0.0725 <0.0001*	-0.146 -0.427 -0.097 -0.245 -0.189 -0.544 -0.503 -0.474 -0.954	[-0.252, -0.042]* [-0.543, -0.319] [-0.199, 0.003] [-0.330, -0.163]* [-0.398, 0.006] [-1.509, -0.011] [-1.273, -0.027] [-1.589, 0.124] [-1.189, -0.764]*	2.81 [4.80 [2.66 [4.20 [2.69 [4.33 [4.28] 4.38 [3.44 [2.43, 3.2 4.36, 5.2 2.30, 3.0 1.96, 4.5 1.96, 8.0 2.28, 8.0 2.48, 7.2 2.08, 8.6 3.25, 3.6	7] 0.350 4] 0.456 4] 0.402 8] 0.421 6] 0.324 0] 0.333 7] 0.339

707

							PCA Self-th	ninning L	.ine ^e		
		No. c Point	of S ^C				Slope]	Intercept		
Id. Code ^a	Condition ^b	n _T	n	r ²	Pq	ĝ	95% CI ^f	â	95% C	ī	p _{est} q
95A 96A 97A 98A 98A 99A 101A 103T 105A 105A 105A 1105A 1105A 1105A 1105A 1107 111A 111A 111A 111A 111A 111A 111	S.I. 22.8 S.I. 28.9 S.I. 33.5	3 28 4 4 4 4 10 8 10 3 3 3 6 6 6 9 9 9 4 4 7 7 5 5 5 5 7 7 7 9	2 4 2 2 4 4 4 7 8 9 3 3 3 6 3 3 8 4 8 3 0 10 3 5 5 5 9 9 4 3 3 3 3 3 4 7 7 8 3 5 4	1.000 0.892 1.000 1.000 0.971 0.964 0.664 0.949 0.945 0.847 0.770 0.188 0.780 0.887 0.890 0.887 0.890 0.887 0.890 0.953 0.890 0.953 0.890 0.953 0.961 0.975 0.975 0.975 0.975 0.624 0.9772 0.9775	0.0554 0.0146* 0.0183* 0.0255* 0.6340 0.0001* 0.2038 0.1450 0.0011* 0.2560 0.0042* 0.3108 0.1777 0.3108 0.0002* 0.00270* 0.0167* 0.0189* 0.0001* 0.0552 0.0074* 0.1384 0.3819 0.4029 0.0144* 0.2665 0.1699 0.0292* 0.4200 0.0499* 0.0125*	-0.757 -1.262 -0.841 -3.005 -2.478 -1.066 -0.986 -29.542 -0.670 -0.958 -0.782 -0.467 -0.218 -0.204 -0.534 -0.650 -1.301 -1.326 -1.335 -0.299 -0.299 -0.587 -0.513 -0.089 -1.421 -1.389 -2.786 -0.109 -0.133 -0.099 -0.133 -0.099 -0.133 -0.099 -0.509 -0.509 -0.509 -0.509 -0.509 -0.509 -0.509 -0.509 -0.956	[-5.012, -1.559]* [-2.132, -0.549]* [-3.077, -0.309] [-4.882, 7.398] [-0.837, -0.526]* [-0.636, -0.318] [-0.889, -0.261] [-0.125, 0.539] [-5.766, -2.755]* [-2.495, -0.746]* [-3.430, -0.639]* [-3.430, -0.639]* [-0.396, -0.206]* [-0.397, -0.206]* [-0.397, -0.206]* [-0.397, -0.206]* [-0.391, -0.439] [-0.361, 0.171] [-5.645, -1.772]* [-0.352, 0.122] [-0.364, 0.084] [-1.149, -0.087] [-20.004, -0.005]	4.04 3.02 3.79 1.55 1.39 3.44 2.70 -27.20 3.42 3.77 4.14 3.46 3.48 4.04 4.28 1.28 1.28 1.28 3.02 3.10 4.07 4.16 4.04 4.38 3.02 3.10 4.07 4.16 4.04 4.38 3.97 4.17 4.07 4.16 4.07 4.17 4.07 3.58 3.70 3.58 3.70 4.71 4.71 4.71 4.71 4.71 4.71 4.71 4.71	[-1.63, [-0.84, [-1.16, [-3.23, [-0.18, [-0.19, [-0.19, [-0.19, [-0.19, [-0.49, [-0.49, [-0.49, [-0.49, [-0.49, [-0.49, [-0.56, [-3.41, [-0.56, [-0.56, [-3.41, [-0.56	2.48] 3.99] 3.84] 11.80] 3.59] 3.91] 4.40] 4.40] 4.64] 4.64] 4.64] 4.10]	0.285 0.221 0.272 0.125 0.144 0.252 0.016 0.299 0.255 0.281 0.341 0.415 0.332 0.615 1.106 0.104 0.385 0.315 0.330 0.459 0.207 0.214 0.385 0.315 0.330 0.459 0.207 0.209 0.132 0.441 0.331 0.331 0.331 0.331
121A 121T 122A		9 4	8 3 5	0.575 0.624 0.772	0.0292* 0.4200 0.0499*	-0.509 -0.509 -0.956	[-1.149, -0.087]	3.58 3.70 3.84	[3.09,	3.90] 4.01]	

Table A.2. (continued)

			_			PCA Self-thinning Line ^e						
		No. o Poin					Slope]	ntercept			
Id. Code ^a	Condition ^b	^{n}T	n	r ²	Pd	ĝ	95% CI ^f	â	95%	CI	p _{est}	
131A		6	6	0,956	0.0007*	-0.498	[-0.657, -0.356]	3.70	ſ 3.59,	3.81]	0.334	
131T		4	4	0.953	0.0236*	-0.224	[-0.384, -0.075]*	4.11	[3.93,		0.408	
132A		8	8	0.206	0.2580	-0.294	-	3.86	_	_	0.386	
133A		8	8	0.566	0.0312*	-0.188	[-0.362, -0.024]*	3.73	[3.63,	3.82]	0.421	
135A		5	5	0.854	0.0248*	~3.047	[-11.936, -1.628]*	4.63	3.54	11.44	0.124	
136A		3	3	0.714	0.3593	-0.284		4.00	,	-	0.389	
136T		3	3	0.763	0.3240	-0.331		4.05			0.376	
137A		17	12	0.983	<0.0001*	-0.422	T-0.462, -0.383]*	3.90	「 3.88.	3.92	0.352	
137T		17	12	0.982	<0.0001*	-0.433	Γ-0.4760.392 [†] *	3.97	Ĩ 3.95.	3.99	0.349	
138A		3	3	0.798	0.2969	-0.550		3.42	. ,	_	0.323	
138T		3	3	0.783	0.3082	-0.544		3.61			0.324	

 a Table A.l associates the numeric part of each Id. code with a particular species and study. The letter indicates the type of biomass measurements made: A = aboveground parts only, T = aboveground and belowground parts both included.

DSee Table A.1 and the references given there for further information on condition.

 $^{\text{C}}$ n_{\text{T}} is the total number of log 8-log N points reported for each code and condition. In the the number of points remaining after removing points not relevant to the thinning line. This is the number of points used to fit the PCA relationship between log B and log N.

 $^{
m d}$ The statistical significance of the correlation between log B and log N. * indicates significance at the 95% confidence level (P < 0.05).

^eThinning lines were fitted to log B-log N data using principal component analysis (PCA). Forumlas for this analysis and for computing 95% confidence limits are given in Jolicoeur and Heusner (1971).

f* indicates thinning slopes that are statistically different at the 95% confidence level from the thinning rule prediction of β = -1/2.

9Transformed values of the log B-log N thinning slope, $\hat{\beta}$, calculated from p_{est} = 0.5 / (1 - $\hat{\beta}$) (Chapter 6).

Table A.3. Ranges of Log B and Log N Used to Fit Self-thinning Lines to Experimental and Field Data.

т.(Log N ^C			Log B ^C		
Id. Code ^a	Condition ^b	n	Min.	Max.	Mean	Min.	Max.	Mean	
1A		3 3	0.08	1.20	0.61	3.27	3.76	3.51	
1T 5A		3 4	0.08	1.20	0.61 -0.14	3.34 3.50	3.83 3.97	3.58 3.68	
7A		5	2.71	3.67	3.07	2.51	2.86	2.74	
88	Lot 2B	5 7	-0.70	-0.16		3.89	4.29	4.12	
8A	Lot 2C	7		-0.43		3.83	4.33	4.10	
10A 14A	Full light	10 4	3.05 -0.91	4.74 -0.61	3.87 -0.77	2.36 4.12	3.17 4.31	2.77 4.23	
15T		31	1.38	3.16	2.78	2.33	3.41	2.64	
16T		38	2.47	4.05	3.29	2.01	2.88	2.43	
17T		17	1.61	3.25	2.49	2.17	3.18	2.60	
18T 19A		43	1.24	3.17	2.43	2.09	3.11	2.49	
20T		26 13	-1.06 2.98	1.38 5.12	0.07 3.87	3.51 3.12	5.08 3.98	4.11 3.58	
21T		13	3.03	3.96	3.55	1.65	2.55	2.01	
22 T		34	-0.10	1.65	0.59	3.28	4.33	3.68	
23T		23	-0.81	1.12	0.08	3.49	4.06	3.87	
24T 26A		5 27	0.50 -0.75	1.39 0.16	0.88 -0.24	3.79 3.66	4.18 4.62	3.98 4.07	
27A			2.83	3.12	2.98	3.35	3.44	3.40	
28A	Low L.I.	2 7	3.21	3.44	2.98 3.31	1.70	1.90	1.81	
28A	Med. L.I.	8 7	3.08	3.22	3.16	2.69	2.83	2.76	
28A	High L.I.	/	2.96	3.22	3.09	2.74	2.92	2.84	
29T 30A		9 9	2.80 3.37	3.75 3.90	3.16 3.66	2.44 2.49	3.10 2.78	2.90 2.66	
31A		11	3.00	3.74	3.46	2.28	2.62	2.46	
32T	20% N.C.	5	3.46	4.01	3.66	2.64	2.84	2.77	
32T	100% N.C.	5	3.09	3.90	3.42	3.03	3.17	3.12	
33T 34T		10 10	2.54 2.77	2.87 3.27	2.68 2.98	2.56 2.64	2.96 3.58	2.80 3.07	
35A		13	3.14	3.80	3.46	2.78	3.21	3.07	
36A		10	0.00	0.88	0.49	3.41	3.77	3.54	
38A	30% L.I.	7	2.80	3.91	3.24	1.99	2.26	2.16	
38A 38A	70% L.I.	8	2.67		3.19 3.13	2.43	3.07 3.07	2.72 2.78	
39A	100% L.I. Kitami	8 5	2.50 -0.86	3.82 -0.06		2.53 4.82	5.24	5.06	
39A	Shirogane	7	-0.84		-0.34	4.56	4.79	4.67	
40A	•	10	3.00	3.79	3.37	2.71	3.16	2.97	
41A	10% + T	5		-0.87		4.38	4.61	4.50	
43A 43A	18% L.I. 25% L.I.	7 7	3.61 3.68	4.41 4.42	4.03 4.03	1.82 2.02	2.44 2.65	2.12	
43A 43A	37% L.I.	7	3.47		3.88	2.16	2.80	2.50	
	· · · · · · · · · · · · · · · · · · ·	•		•					

Table A.3. (continued)

ل ـ ۲				Log N ^C			Log B ^C	·····
Id. Codea	Condition ^b	n	Min.	Max.	Mean	Min.	Max.	Mean
43A	55% L.I.	8	3.46	4.49	3.88	2.30	2.99	2.70
43A	100% L.I.	7	3.68	4.27	3.88	2.62	3.04	2.78
44A	Ex. 2-60% L.I.	4	2.05	2.45	2.17	2.59	2.95	2.80
44A 44T	Ex. 2-100% L.I. Ex. 1-23% L.I.	4	2.09 1.76	2.45	2.23 2.11	2.91 2.07	3.29 2.45	3.08 2.35
44T	Ex. 1-23% L.I. Ex. 1-60% L.I.	10 8	1.78	2.92 3.19	2.35	2.62	3.07	2.87
44T	Ex. 1-100% L.I.	6	2.35	3.06	2.59	2.96	3.16	3.04
44T	Ex. 2-60% L.I.	4	2.05	2.45	2.17	2.61	2.97	2.82
44T	Ex. 2-100% L.I.	4	2.09	2.45	2.23	2.93	3.32	3.10
45A		25	-0.78	0.06	-0.38	3.62	4.24	3.95
48A		7	-0.05	0.52	0.29	3.27	3.76	3.51
49T	350/ 1 7	10	2.95	3.62	3.29	2.70	2.85	2.79
50T	15% L.I.	12	2.17	3.34	2.76	1.46	2.05	1.76
50T 51A	100% L.I.	22 24	2.03 2.86	4.00 3.76	2.91 3.41	2.08 2.35	2.56 2.93	2.38 2.61
52A		3	1.83	1.93	1.88	3.45	3.58	3.53
53T		29	2.92	3.64	3.28	2.84	3.23	3.03
54A		4	-0.81	-0.58	-0.69	4.17	4.52	4.35
55A		11	3.66	3.92	3.80	2.40	2.82	2.55
56A		11	3.64	3.95	3.82	2.68	2.96	2.78
57A		13	3.51	3.92	3.77	2.55	2.94	2.74
58T		23	2.14	3.41	2.60	1.98	2.63	2.27
A08		5	-0.78	-0.51	-0.68	4.62	4.83	4.73
81A 82A		4 17	-0.32 -0.85	0.88 -0.05	0.27 -0.53	3.58 4.23	4.26 4.51	3.90 4.37
84A		243	-1.57		-0.98	3.74	4.78	4.25
86A		15	4.09	4.83	4.43	2.85	3.54	3.18
87A		12	3.25	4.20	3.77	2.75	3.45	3.04
A88		15	3.21	3.94	3.60	2.57	3.26	2.94
89A	Grazed	3	3.77	4.10	3.93	1.63	1.83	1.74
89A	Ungrazed	5	3.55	4.00	3.73	1.81	2.09	1.97
90A	Grazed	2	3.77	3.94	3.85	2.24	2.39	2.31
90A 91A	Ungrazed	2	3.68	3.94	3.81	2.48 2.13	2.60 2.50	2.54
91A	17% L.I. 100% L.I.	13 10	2.91 3.23	4.53 4.85	3.66 4.06	2.13	3.45	2.27 3.07
91T	17% L.I.	13	2.91	4.53	3.66	2.15	2.50	2.31
91T	100% L.I.	10	3.23	4.85	4.06	3.04	3.46	3.20
92A	17% L.I.	5	3.18	4.56	3.76	1.86	2.10	1.99
92A	23% L.I.	5	3.52	4.58		1.92	2.57	2.23
92A	44% L.I.	5	3.27	4.55		1.92	2.69	2.38
92A	100% L.I.	5	3.37	4.57		2.13	2.78	2.56
93A		8		-0.57		3.96	4.56	4.22
95A		2	-0.16	0.16	0.00	3.92	4.16	4.04

Table A.3. (continued)

Tal				Log N		Log BC			
Id. Code ^a	Condition ^b	n	Min.	Max.	Mean	Min.	Max.	Mean	
96A		4		-1.04		4.32	4.49	4.42	
97A	C T 22 0	2		-1.04		4.66 4.50	4.78	4.72 4.55	
98A 98A	S.I. 22.8 S.I. 28.9	2 4			-1.00 -1.19	3.82	4.61 4.65	4.33	
98A	S.I. 33.5	4			-1.07	4.32	4.74	4.58	
99A		7			-1.69	4.22	4.54	4.36	
101A		8			-1.06	3.35	4.60	3.99	
102A		9			-1.17	3.98	4.40	4.21	
103A		3			-0.74	4.30	4.63	4.48	
103T 104A		3 6 3 3			-0.74 -1.28	4.41 4.11	4.67 4.51	4.54 4.32	
104A 105A		3			-0.99	4.11	4.44	4.28	
105T		3			-0.99	4.26	4.49	4.34	
106A		8		0.52		3.14	3.62	3.40	
109A		4	-0.31	0.23	-0.11	3.35	3.60	3.47	
110A		8			-1.01	3.64	4.02	3.85	
110T		3			-0.57	3.85	4.04	3.97	
111A		10			-0.74	3.26	4.75	4.10	
111T 112A		10 3	0.08		-0.74 0.22	3.35 3.92	4.82 4.27	4.19 4.10	
113A		5			-1.05	4.07	4.80	4.4	
113T		5			-1.05	4.16	4.88	4.50	
114A		9	-0.92			3.26	4.44	3.98	
114T		9	-0.92			3.35	4.52	4.0	
115A		4			-0.28	4.08	4.28	4.2	
116A 117A		3 3			-1.16	4.48 4.55	4.58	4.58 4.58	
117A 118A		3			-1.14 -0.91	3.89	4.60 4.27	4.17	
118T		3 3			-0.91	3.94	4.31	4.17	
119A		4			-0.77	3.29	4.34	3.8	
120A		7	-0.72	0.61		3.91	4.17	4.0	
120T		7			-0.35	4.01	4.26	4.1	
121A		8			-0.77	3.81	4.24	3.9	
121T		3			0.74	3.91	4.34	4.0 4.0	
122A 123A		5 4	-0.44 0.15		-0.17 3 0.47	3.77 2.20	4.18 3.55	2.8	
123A 124A		4	-0.13			4.08	4.24	4.1	
125A		3			-0.68	3.79	4.09	3.9	
126A		5			-0.40	3.78	4.20	3.9	
128A		4	0.90			3.88	4.17	4.0	
130A		3			-0.56	3.46	4.07	3.8	
131A 131T		6 4			-0.72 -1.12	3.49 4.26	4.32 4.42	4.0 4.3	

Table A.3. (continued)

T				Log N ^C			Log BC		
Id. Code ^a	Condition ^b	n	Min.	Max.	Mean	Min.	Max.	Mean	
132A		8	-0.92	0.00	-0.35	3,76	4.18	3.96	
133A		8	-1.02	0.24	-0.56	3.62	3.93	3.83	
135A		5	0.60	0.99	0.77	1.60	2.90	2.29	
136A		3	-1.52	-0.83	-1.19	4.21	4.42	4.34	
136T		3	-1. 52	-0.83	-1.19	4.29	4.52	4.44	
137A		12	-1.07	0.14	-0.50	3.85	4.34	4.11	
137T		12	-1.07	0.14	-0.50	3.93	4.42	4.19	
138A		3	-0.34	0.70	0.07	3.03	3.65	3.38	
138T		3	-0.34	0.70	0.07	3.22	3.84	3.57	

a Table A.l associates the numeric part of each Id. code with a particular species and study. The letter indicates the type of biomass measurements made: A = aboveground parts only, T = aboveground and belowground parts both included.

bSee Table A.1 and the references given there for more information on condition.

CThe mean, minimum, and maximum are given for log B and log N over the n data points used to fit the thinning line for each code and condition.

Table A.4. Fitted Allometric Relationships for Experimental and Field Data.

		A ⁻	llometr	ric Relation	nship Fit	by PCA ^C
Id. Code ^a	Condition ^b	n	r ²	pď	$\hat{\Phi}_1$	$\hat{\Phi}_0$
height-	-weight allometry:	log	_ h =	φ̂ _l log w̄ +	$\hat{\Phi}_0$ ($\hat{\Phi}_1$ =	Ĝ _{hw})e
1A 1T 8A 44A 44A 44T 48A 86A 86A 98A 98A 98A 103A 105T 106A 110T 111A 111T 112A 113T 114A 115A 116A 120T 121A 121T 122A	Lot 2B Lot 2C Ex. 2-60% L.I. Ex. 2-100% L.I. Ex. 2-100% L.I. S.I. 28.9 S.I. 33.5	337744447451258337933633883003559943477835	0.997 0.996 0.997 0.998 0.977 0.919 0.975 0.919 0.973 0.899 0.951 1.000 0.993 0.994 0.980 0.993 0.991 0.959 0.959 0.959 0.959 0.959 0.959 0.977 0.938 0.939	<pre><0.0001* <0.0001* <0.0001* 0.0064* 0.0540 <0.0001* 0.0892 0.0533 <0.0001* 0.1292 <0.0001* <0.0002* <0.0002* <0.0002*</pre>	0.282 0.283 0.384 0.399 0.368 0.471 0.368 0.463 0.253 0.348 0.303 0.356 0.296 0.339 0.366 0.337 0.344 0.328 0.359 0.195 0.167 0.169 0.391 0.353 0.358 0.279 0.239 0.238 0.321 0.163 0.188 0.446 0.297 0.291 0.263 0.243 0.387	-0.091 -0.113 -0.610 -0.725 -0.202 -0.394 -0.209 -0.397 -0.086 -0.587 0.016 -0.167 -0.212 -0.418 -0.747 -0.576 -0.441 -0.544 -0.453 -0.640 0.191 0.424 0.404 -0.712 -0.429 -0.62 -0.619 -0.676 -0.295 -0.125 -0.144 -0.512 -0.541 0.253 0.366 -1.131 -0.361 -0.361 -0.361 -0.725

Table A.4. (continued)

		Α.	llometr	ic Relation	nship Fit	by PCA ^C
Id. Code ^a	Condition ^b	n	r ²	pd	ф 1	$\hat{\Phi}_0$
height-	weight allometry:	log	ĥ =	φ̂ _l log w̄ +	$\hat{\Phi}_0$ ($\hat{\Phi}_1$ =	ф̂ _{hw})е
123A 124A 125A 126A 128A 130A 131A 131T 132A 136A 136T 137A 137T 138A 138T		4 4 3 5 4 3 6 4 8 8 3 12 12 3 3	0.974 0.984 0.966 0.774 0.992 0.979 0.982 0.938 0.962 0.991 0.985 1.000	0.0131* 0.0082* 0.1189 0.0492* 0.0038* 0.0930 0.0002* 0.0093* <0.0001* 0.0621 0.0613 <0.0001* 0.0043* 0.0043* 0.0006*	0.398 0.371 0.436 0.293 0.382 0.365 0.288 0.184 0.322 0.203 0.203 0.282 0.272 0.304 0.302 0.228	-0.471 -0.718 -1.013 -0.554 -1.130 -0.682 -0.154 0.390 -0.473 0.337 -0.179 -0.154 -0.278 -0.291 -0.054 -0.100
DBH-we	ight allometry: 1	og DB	H = Φ1	$\log \bar{w} + \hat{\Phi}_0$	$(\hat{\Phi}_1 = \hat{\Phi}_{DV})$	√)e
8A 8A 14A 54A 81A 93A 95A 96A 97A 98A 98A 99A	Lot 2B Lot 2C S.I. 22.8 S.I. 28.9 S.I. 33.5	7 7 4 2 4 8 2 4 2 2 4 7	0.999 0.997 0.995 1.000 1.000 0.994 1.000 1.000 1.000 0.996 0.993	<0.0001* 0.0037* 0.0002* 0.0020* <0.0001*	0.321 0.328 0.274 0.278 0.395 0.347 0.169 0.310 0.317 0.422 0.374 0.347	
BSLA-we	eight allometry:	log \bar{B}	SLA =	φֿן log w̄ +	$\hat{\Phi}_0$ ($\hat{\Phi}_1 = \hat{\Phi}_0$	∱ _{Bw})e
8A 8A 14A 45A 52A 54A 93A	Lot 2B Lot 2C	7 7 4 25 3 4 8	0.998 0.999 0.999 0.963 0.997 0.997	<0.0001* 0.0006* <0.0001* 0.0344* 0.0014*	0.651 0.648 0.542 0.826 0.774 0.664 0.698	-5.030

Table A.4. (continued)

		/	Allometr	ic Relatio	onship Fit by	, PCA ^C
Id. Code ^a	Condition ^b	n	r ²	pd	^ 1	$\hat{\Phi}_0$
BSLA-we	eight allometry:	log i	BSLA = 🏟	log ₩ +	$\hat{\Phi}_0$ ($\hat{\Phi}_1 = \hat{\Phi}_{BV}$	v)e
95A 96A 97A 98A 98A 98A 101A 102A 103T 104A 105A 105T 109A 110T 111A 111T 112A 113A 113T 114A 115A 116A 117A 118A 118T 119A 121A 121A 121A 121A 121A 121A 121A	S.I. 22.8 S.I. 28.9 S.I. 33.5	10g 242244789336334831035577433333483534564	1.000 0.997 1.000 1.000 1.000 0.996 0.999 0.949 0.994 0.991 1.000 1.000 1.000 0.994 1.000 0.992 0.988 0.997 0.997 0.997 0.995 1.000 0.995 0.995 1.000 0.995 0.995 0.995 0.995 0.995 0.995 0.998	0.0016* 0.0002* 0.0001* 0.0001* 0.0001* 0.0950 0.0590 0.0590 0.0021* 0.0021* 0.0031* 0.0031* 0.0038* 0.0001* 0.0038* 0.0001* 0.0026* 0.0001* 0.0026* 0.0067* 0.0204* 0.1379 0.1283 0.0017* 0.0204* 0.1358 0.0017* 0.0001* 0.0564 0.0172* 0.0002* 0.0001* 0.0564 0.0172* 0.0002* 0.0001* 0.0053*	Φ0 (Φ1 = ΦBW 0.765 0.719 0.633 0.839 0.746 0.692 0.766 0.650 0.697 0.666 0.728 0.844 0.806 0.815 1.231 0.850 0.819 0.720 0.730 1.045 0.887 0.886 0.849 0.891 1.030 1.051 0.951 0.866 0.878 0.849 0.891 1.030 1.051 0.951 0.866 0.878 0.849 0.878 0.849 0.878	-5.467 -5.174 -4.820 -5.971 -5.466 -5.204 -5.630 -5.040 -5.091 -5.106 -5.478 -6.071 -5.813 -5.907 -7.368 -5.763 -5.856 -5.763 -5.856 -6.143 -6.212 -5.881 -6.212 -5.883 -6.190 -6.357 -6.357 -6.357 -6.333 -6.746 -6.357 -6.357 -6.333 -6.746 -6.357 -6.357 -6.333 -6.746 -6.357 -6.357 -6.333 -6.360 -5.789 -5.886
132A 133A 136A 136T		8 8 3 3	0.993 0.989 1.000	<0.0001* <0.0001* 0.0012* 0.0004*	0.871 0.874 0.822 0.794	-5.940 -5.961 -5.762 -5.688

Table A.4. (continued)

		A	llometr	ic Relatio	onship Fit by	y PCA ^C
Id. Codea	Condition ^b	n	r ²	pd	Ŷ 1	$\hat{\Phi}_0$
BSLA-we	eight allometry:	log B	SLA = 6	ົງ log พี +	$\hat{\Phi}_0$ ($\hat{\Phi}_1 = \hat{\Phi}_B$	w)e
137A 137T 138A 138T		12 12 3 3	0.998 0.998 1.000 1.000	<0.0001* <0.0001* 0.0084* 0.0121*	0.872 0.865 0.766 0.769	-6.062 -6.099 -5.295 -5.450
height-	DBH allometry:	log ñ	= φ 1	log DBH +	$\hat{\Phi}_0$ ($\hat{\Phi}_1 = \hat{\Phi}_h$	D)e
8A 8A 81A 93A 98A 98A 99A	Lot 2B Lot 2C S.I. 28.9 S.I. 33.5	7 7 4 8 3 3 7	0.993 0.991 0.997 0.976 0.999 0.995 0.999	<0.0001* <0.0001* 0.0016* <0.0001* 0.0201* 0.0455* <0.0001*	1.198 1.215 0.883 0.993 1.041 1.114 0.860	2.225 2.205 2.205 2.060 2.107 2.219 2.039
height-	BSLA allometry:	log ñ	= Φ η	log BSLA +	$\hat{\Phi}_0$ ($\hat{\Phi}_1 = \hat{\Phi}_h$	_B) e
8A 93A 98A 98A 99A 102A 103A 103T 104A 105A 105T 110A 111T 111A 111T 112A 113A 113T 114A 114T 115A 116A 117A 119A	Lot 2B Lot 2C S.I. 28.9 S.I. 33.5	7 7 8 3 3 7 9 3 3 6 3 3 8 3 10 0 3 5 5 7 7 4 3 3 4	0.991 0.993 0.978 0.999 0.994 0.990 1.000 0.987 0.961 0.956 0.408 0.905 0.905 0.901 0.901 0.941 0.967 0.999	<pre><0.0001* 0.1271 0.1271 <0.0001* 0.5586 <0.0001* 0.1483 0.0137* 0.0137* 0.0003* 0.0167* 0.2070</pre>	0.588 0.615 0.491 0.523 0.557 0.440 0.496 0.494 0.230 0.207 0.394 0.355 0.488 0.488 0.264 0.264 0.264 0.399 0.399 0.192 0.223 0.000 0.470	2.225 2.205 2.060 2.107 2.219 2.039 1.974 2.065 1.592 1.627 1.874 1.802 1.982 1.982 1.530 1.521 1.740 1.740 1.417 1.672 1.415 1.854

Table A.4. (continued)

			llometr	ic Relatio	nship Fit by	PCA ^C
Id. Code ^a	Condition ^b	n	r ²	Рd	Ŷı	$\hat{\Phi}_0$
height-	-BSLA allometry:	log h	i = Φη 1	og BSLA +	$\hat{\Phi}_0$ ($\hat{\Phi}_1 = \hat{\Phi}_{hB}$)	е
121A		8	0.941	<0.0001*	0.299	1.707
121T			0.954	0.1380	0.271	1.639
122A		3 5 3 4 5	0.960	0.0034*	0.462	1.964
123A		3	0.997	0.0347*	0.260	1.505
124A		4	0.990	0.0052*	0.239	1.365
126A		5	0.803	0.0396*	0.371	1.521
131A		6	0.993	<0.0001*	0.351	1.866
131T		4	0.998	0.0012*	0.252	1.735
132A		8	0.922	0.0002*	0.368	1.718
133A		8	0.933	<0.0001*	0.229	1.714
136A		8 3 3	0.991	0.0609	0.343	1.795
136T		3	0.991	0.0609	0.343	1.795
137A		12	0.986	<0.0001*	0.349	1.837
137T		12	0.986	<0.0001*	0.349	1.837
138A		3	1.000	0.0127*	0.297	1.521
138T	•	3	1.000	0.0127*	0.297	1.521

aTable A.l associates the numeric part of each Id. code with a particular species and study. The letter indicates the type of biomass measurements made: A = aboveground parts only, T = aboveground and belowground parts both included.

bSee Table A.1 and the references given there for more information on condition.

CThe general allometric relation is $\log Y = \widehat{\Phi}_1 \log X + \widehat{\Phi}_0$, where Y is $\log \widehat{h}$, $\log \widehat{DBH}$, or $\log \widehat{BSLA}$ and X is $\log \widehat{w}$, $\log \widehat{DBH}$, or $\log \widehat{BSLA}$. (DBH and BSLA here refer to the diameter and breast height and basal area at breast height, respectively, of the boles of individual trees). Y and X are paired as indicated in the sub-headings within the table. The number of data points and coefficient of determination (r²) are given for each allometric relationship. The slope and intercept ($\widehat{\Phi}_1$ and $\widehat{\Phi}_0$, respectively) are fit using principal component analysis (Jolicoeur and Heusner 1971).

 $^{d}\mathsf{The}$ statistical significance of the correlation between the two measurements log Y and log X, where Y and X are as explained in footnote c.

eThe mathematical symbols $\widehat{\Phi}_{hw}$, $\widehat{\Phi}_{Dw}$, $\widehat{\Phi}_{Bw}$, $\widehat{\Phi}_{hD}$, and $\widehat{\Phi}_{hB}$ are used in Chapter 6 to represent the five allometric powers in this table.

Table A.S. Self-thinning Lines from Experimental and Field Data Cited by Previous Authors as Support for the Self-thinning Rule.

					of nts ^e		T	hinning Line Eq Slope		rcept	
ld. Codeª	Conditionb	Dep. Var. ^C	Fitting Method ^d	nT	n	r ² or %EV ^f	βg	95% CI or SEh	â	SEh	Reference ⁱ
1T		W	Req				-0.43		3.85		White 1980
5A		W	Reg				-0.33		3.64		White 1980
7 A		W	Reg		10		-0.39		3.83		White 1980
8A	Lot 2B	W	Reg				-0.70		3.78		White 1980
1 OA	Full light	В	Reg	10	6		-0.29	0.21	3.96	0.75	Westoby and Howell 1982
14A		W	Reg	47			-0.66		3.65		White 1980
15T +		W	Gra	47			0.61		3.92		Yoda et al. 1963 White 1980
161		W	Reg Gra	38			-0.51		3.92		Yoda et al. 1963
+		W	Req	20			-0.48		3.89		White 1980
17T		ŵ	Gra	21			-0.40		3.07		Yoda et al. 1963
+		W	Req	_			-0.48		3.85		White 1980
18T		W	Gra	51							Yoda et al. 1963
+		W	Reg				-0.48		3.66		White 1980
19A		W	Gra	28							Yoda et al. 1963
+		W	Reg	0.7			-0.54		4.34		White 1980
20T		W	Gra	27			0.00		4.00		Yoda et al. 1963 White 1980
217		W	Reg Gra	30			-0.38		4.00		Yoda et al. 1963
+		W	Req	30	25		-0.66		4.31		White and Harper 1970;
		"	weg	30	LJ		.0.00		****		White 1980
22T		W	PCA	53	34	0.98*	-0.46				Mohler et al. 1978
23T		W	PCA	29	23	0.98	-0.22				Mohler et al. 1978
26A		W	Gra	46			-0.50				Yoda et al. 1963
+		W	Req				-0.63		3.94		White 1980
27T		В	Gra	20							Westoby 1980
28A	Low L.I.	W	PCA	10	7	0.95*	-0.74				Hutchings and Budd 1981b
28A	Medium L.I.	W W	PCA	14	8	0.95*	-0.91				Hutchings and Budd 19815
28A 29T	High L.I.	W	PCA Gra	14 25	7	>0.99*	-0.62				Hutchings and Budd 1981b Yoda et al. 1963
+		W	Req	65			-0.48		4.41		White 1980
30A		ŵ	Reg		9	0.76	-0.33		3.86		White and Harper 1970;
31A		W	Reg		11	0.89	-0.42		3.93		White 1980 White and Harper 1970;
			· ·			0.03	-0.42				White 1980
32T	20% N.C.	W	Ax		7				4.62	0.04	Furnas 1981
32T 33T	100% N.C.	W	Ax		7				4.84	0.05	Furnas 1981 Furnas 1981
331 &	Restricted Control	W W	Ax Ax		5 5				4.07	0.04	Furnas 1981
35A	CONCIOI	W	Req	22	13		-0.56	[-0.79,-0.33]	4.95	0.04	Westoby 1976
38A	70% L.I.	ŵ	Gra	20	1.5		-0.50	[-0.75,-0.55]	4.75		Kays and Harper 1974
38A	100% L.I.	W	Gra	20							Kays and Harper 1974
40A		W		8	4		-0.46				Ford 1975
41A		W	Gra	5	5						Barkham 1978
43A	18% L.I.	В	Reg	10	7	0.95	-0.67	0.18			Westoby and Howell 1981
43A	25% L.I.	В	Reg	10	7	0.92	-0.66	0.07			Westoby and Howell 1981
43A	37% L.I.	В	Reg	10	7	0.94	-0.65	0.08			Westoby and Howell 1981
43A	55% L.I.	В	Reg	11	8	0.93	-0.64	0.09			Westoby and Howell 1981 Westoby and Howell 1981
43A 44T	100% L.I. Ex. 1-60% L.I.	B W	Reg Reg	9	7 8	0.67 0.96	-0.56 -0.30	0.07	3.53		White and Harper 1970
441 44T	Ex. 1-100% L.I.		Reg		9	0.96	-0.30		3.84		White and Harper 1970
44T	Ex. 2-60% L.I.	. w	Reg		6	0.98	-0.33		4.34		White and Harper 1970
+	-x 00% -11.	W	PCA		U	>0.99*	-0.79	[-1.19,-0.65]	4.51		Lonsdale and Watkinson 19
	5 0 100% t T		Req		5	0.97	-0.84	,	4.96		White and Harper 1970
44T	Ex. 2-100% L.I.		IVE4								
44T + 45A	Ex. 2-100% L.1	. w w	PCA			>0.99*	-0.89	[-1.29,-0.45]	5.06		Lonsdale and Watkinson 19 White 1980

Table A.5. (continued)

				No.	of nts ^e			hinning Line Eq Slope		cept	
Id. Codea	Condition ^b	Dep. Var. ^C	Fitting Method ^d	n _T	n	r² or %EV ^f	ĝg	95% CI or SE ^h	â	SEh	Reference ⁱ
48A		W	Reg				-0.30		3.61		White 1980
51A	Low fertility	W	Reg	15			-0.44	0.35			Bazzaz and Harper 1976
&	High fertility		Reg	15			-0.64	0.19			Bazzaz and Harper 1976
52A		W	Gra	9							Tadaki and Shidei 1959
53T		W	Reg	36		0.87	-0.41		4.19		White and Harper 1970; White 1980
54A		W	Gra	4							Tadaki and Shidei 1959
55A		W	Rea	17							Malmberg and Smith 1982
56A		W	Req	34							Malmberg and Smith 1982
57A		W	Req	21		0.87	-0.75		5.58		Malmberg and Smith 1982
82A		W	Gra								Peet and Christensen 1980; Christensen and Peet 1982
84A		W	Gra								Yoda et al. 1963
86A		W	PCA	18	15	>0.99*	-0.78	[-1.11,-0.52]	6.67		Lonsdale and Watkinson 1983a
87A		W	PCA	15	12	>0.99*	-0.68	[-0.82,-0.56]	5.64		Lonsdale and Watkinson 1983a
88A		W	PCA	18	15	>0.99*	-0.79	[-1.08,-0.55]	5.82		Lonsdale and Watkinson 1983a
89A	Grazed	W	Reg	5	3		-0.60				Dirzo and Harper 1980
89A	Ungrazed	W	Reg	5	5	0.96	-0.80				Dirzo and Harper 1980
90A	Grazed	W	Gra	5							Dirzo and Harper 1980
90A	Ungrazed	W	Gra	5							Dirzo and Harper 1980
91A	100% L.I.	W	PCA	25		>0.99*	-0.43	[-0.51,-0.35]	4.84		Lonsdale and Watkinson 1982
917	100% L.I.	W	PCA	25		>0.99*	-0.26	[-0.33,-0.19]	4.29		Lonsdale and Watkinson 1982
92A	23% L.I.	W	PCA	5	5	>0.99*	-0.52	[-0.74,-0.30]	4.26		Lonsdale and Watkinson 1982
92A	44% L.I.	W	PCA	5 5	5	>0.99*	-0.49	[-0.71,-0.27]	4.26		Lonsdale and Watkinson 1982
92A	100% L.I.	W	PCA	5	5	>0.99*	-0.46	[-0.73,-0.19]	4.37		Lonsdale and Watkinson 1982

Table A.l associates the numeric part of each Id. code with a particular species and study. The letter indicates the type of biomass measurements made: A = aboveground parts only, T = aboveground and belowground parts both included. "+" indicates that the line is a duplicate analysis of the previous code and condition by different author. "&" indicates a treatment or condition analyzed separately in the reference above, but combined with the preceding condition in the reanalysis of this report (Table A.2).

 ${}^{\mathrm{b}}\mathsf{See}$ Table A.1 and the references given there for further information on condition.

 $^{\text{C"W"}}$ indicates that the data were analyzed with log $\overline{\text{W}}$ -log N plots or by fitting log $\overline{\text{W}}$ against log N, while "B" indicates that log B-log N plots or fitting was used.

dMethod of estimating the thinning line. "Gra" indicates graphical analysis only with no statistics, "Reg" indicates regression analysis, and "PCA" indicates principal component analysis. "Ax" indicates a method used by Furnas (1981), who assumed a priori that thinning slopes were -1.5 in the log \tilde{w} -log N plane, then used statistics to estimate the intercept under this axiom. Blanks indicate that the method of fitting was not specified.

 $^{
m en}_{
m T}$ is the total number of size-density points reported for each code and condition. In the the number of points remaining after removing points not relevant to the thinning line. This is the number of points used to fit the relationship between size and density. Blanks indicate that sample sizes were not given in the reference.

^fThis column contains coefficients of determination if the reference analyzed the data by correlation or regression. If the analysis was done by PCA, the number here is the percentage of the combined variance of the two variables explained by the first principal component (%EV).

9Most thinning slopes were originally given for data in $\log \overline{w}$ -log N form. Such slopes were converted here to log B-log N form by adding 1 (Chapter 1). This facilitates comparison with the analyses in Table A.2.

hSingle numbers in the column are standard errors. Paired numbers in brackets are 95% confidence limits.

[†]These are references for the self-thinning analysis only. References to the actual data are in Table A.1

^{*}Percent variance explained (see footnote f).

APPENDIX B FORESTRY YIELD TABLE DATA

Table B.1. Sources of Forestry Yield Table Data.

Id. ode ^a	Species	Group ^b	Shade Tolerance ^C	Wood Density (kg/m ³)d	Conditions ^e	Reference ^f
101	Pinus ponderosa	G	2	472	S.I.(ft) 100	Meyer 1938, Tabs. 3-6
02	Alnus rubra	Ā	2	480	S.I.(ft) 50	Smith 1968, Tab. 3
03	Abies concolor	G	4	360	S.I.(ft) 50	Schumacher 1926, Tab. 1
04	Castanea dentata	Α	3	737	S.I.(ft) 50	Frothingham 1912, Tabs. 17-19
05	Castanea dentata and Quercus ⁹	Α	3	737	S.I.(ft) 50	Frothingham 1912, Tabs. 20-22
06	Quercus	Α	3	737	S.I.(ft) 50	Frothingham 1912, Tabs. 23-25
07	Caryag	Α	2	657	S.I.(ft) 50	Boisen and Newlin 1910, Tab. 14
80	Pinus monticola	G	3	432	S.I.(ft) 50	Haig 1932, Tab. 1
)9	Populus tremuloides	Α	1	400	S.Q. (Ranks)	Baker 1925, Tabs. 14-17
10	Populus deltoides	A	1	384	S.I.(ft) 50	Williamson 1913, Tab. 3
11	Chamaecyparis thyoides	G	4	368	S.I.(ft) 50	Korstian 1931, Tabs. 22,25
12	<u>Tsuga heterophylla</u>	G	4	472	S.I.(ft) 100	Meyer 1937, Tabs. 1-5
13	Picea rubrens	G	4	448	S.I.(ft) 50	Meyer 1929, Tabs. 2-6
14	Pinus ⁹	G		513	S.Q. (Ranks)	Khil'mi 1957, Tabs. 13,14
5	<u>Picea</u> ^g	G		416	S.Q. (Ranks)	Khil'mi 1957, Tabs. 18,45
6	Quercusg	A		737	S.Q. (Ranks)	Khil'mi 1957, Tabs. 23,50
7	Pinus strobus	G	3	400	S.Q. (Ranks)	Marty 1965, Tab. 1
8	Pinus strobus	G	3	400	S.Q. (Ranks)	Marty 1965, Tab. 2
0	Pinus strobus	G	3	400	S.Q. (Ranks)	Marty 1965, Tab. 4
21	Pseudotsuga menziesii	G	3	513	S.I.(ft) 100	McArdle 1930, Tab. 12
2	Pinus resinosa	G	2	529	Spacing (Ranks)	Stiell and Berry 1973, Tabs. 4-8
23	Pinus ⁹	G		513	S.Q. (Ranks)	Tseplyaev 1961, Tab. 47
24	Cedrus9	G		448	S.Q. (Ranks)	Tseplyaev 1961, Tab. 81
25 27	Populus9	A	2	400	S.Q. (Ranks)	Tseplyaev 1961, Tab. 134
28	Pinus ponderosa	G	2 4	472	S.I.(ft) 100	Behre 1928, Tabs. 2-6
9	Thuja occidentalis Pinus taeda	G G		304 609	S.I.(ft) 160	Gevorkiantz and Duerr 1939, Tabs. 18-
30	Pinus elliottii	G	2	609	S.I.(ft) 50	Schumacher and Coile 1960, Tab. 1
31	Pinus palustris	G	3 1	609	S.I.(ft) 50	Schumacher and Coile 1960, Tab. 2
32	Pinus echinata	G	2	609	S.I.(ft) 50 S.I.(ft) 50	Schumacher and Coile 1960, Tab. 3 Schumacher and Coile 1960, Tab. 4
3	Pinus serotina	G	2	609	S.I.(ft) 50	Schumacher and Coile 1960, Tab. 5
34	Pinus taeda	G	2	609	S.I.(ft) 50	Ashe 1915, Tabs. 19,35,42
5	Pinus palustris	Ğ	i	609	S.I.(ft) 50	Wahlenberg 1946, Tabs. 4b,6c,7a,7,b
36	Quercus9 (Upland oaks)	Ä	3	737	S.I.(ft) 50	Schnurr 1937, Tab. 2
37	Picea glauca	Ğ	4	400	Spacing (Ranks)	Stiell 1976, Tabs. 10-13
38	Abies balsamea	Ğ	5	416	S.I.(ft) 65	Meyer 1929, Tabs. 35-49
39	Picea glauca	Ğ	4	400	S.I.(ft) 65	Meyer 1929, Tabs. 29-33
1)	Picea9 and Abies9	Ğ	4	416	S.I.(ft) 50	Bakuzis and Hansen 1965,
2		-	2	480	, .	Tabs. 89,91,93,95,97
13	Alnus rubra Alnus rubra	A A	2	480	S.I.(ft) 50 Crown width	Smith 1968, Tab. 5 Smith 1968, Tab. 6
4	Southern mixed hardwoods		L	641	S.I.(ft) 50	Frothingham 1931, Tab. 8
15	Northern mixed hardwood:			641	S.Q. (Ranks)	Gevorkiantz and Ouerr 1937, Tab. 3
16	Populus9 (Aspen)	> ^	1	464	S.I.(ft) 50	Kittredge and Gevorkiantz 1929, Tab.
17	Eucalyptus globus	É	•	889	S.Q. (Ranks)	Jacobs 1979, Tab. A3.5
18	Eucalyptus microtheca	Ē		801	S.Q. (Ranks)	Jacobs 1979, Tab. A3.13
19	Liriodendron tulipifera		2	448	S.I.(ft) 50	McCarthy 1933, Tab. 17
50	Fraxinus9	Ä	3	641	S.Q. (Ranks)	Sterrett 1915, Tab. 15
51	Pseudotsuga menziesii	Ĝ	3	513	none	McCardle et al. 1949, Tab. 25

Table B.1. (continued)

Id∶ Codeª	Species	Group ^b	Shade Tolerance ^C	Wood Density (kg/m ³) ^d	Conditions ^e	Reference ^f
252	Sequoia sempervirens	G	4	416	S.I.(ft) 100	Lindquist and Palley 1963, Tabs. 1,2,4-6
253	Pinus echinata	G	2	609	S.1.(ft) 50	Mattoon 1915, Tab. 14
254	Pinus echinata	G	2	609	S.I.(ft) 50	Mattoon 1915, Tab. 16
255	Picea sitchensis	G	4	424	S.I.(ft) 50	Cary 1922, Tabs. 6-9
256	Pinus banksiana	G	1	464	S.I.(ft) 50	Eyre 1944, Tab. 8
257	Pinus banksiana	G	1	464	S.I.(ft) 50	Bella 1968, Tabs. 2-6
258	Pinus resinosa	G	2 5	529	S.I.(ft) 50	Eyre and Zehngraff 1948, Tab. 10
259	Tsuga heterophylla	G	5	521	S.I.(ft) 100	Barnes 1962, Tabs. 3,5,7,9,12
260	Tsuga heterophylla	G	5	521	S.I.(ft) 100	Barnes 1962, Tabs. 4,6,8,10,13
261	Tsuga heterophylla	G	5	521	S.I.(ft) 100	Barnes 1962, Tabs. 4,6,8,10,14
262	Picea sitchensis and	G	4	472	S.I.(ft) 100	Taylor 1934, Tabs. 4-8
	Tsuga heterophylla					
263	Picea mariana	G	4	529	S.I.(ft) 50	Fox and Kruse 1939, Tab. 2
264	<u>Pinus</u> <u>banksiana</u>	G	1	464	S.I.(ft) 50	Boudoux 1978, Tabs. 3,4
265	Fraxinus americana	A	3	657	S.I.(ft) 50	Patton 1922, page 37
266	Pseudotsuga menziesii	G	3	512	S.I.(ft) 50	Schumacher 1930, Tabs. 2-5,7
267	Northern mixed hardwoods			641	S.I.(ft) 50	Forbes 1961, Tab. 14
268	Northern mixed hardwoods			641	S.I.(ft) 50	Vermont Ag. Expt. Sta. 1914, Tab. 8
269	<u>Quercus</u> ^g (Red oaks)	А	3	737	S.I.(ft) 50	Patton 1922, page 37
270	Pinus taeda	G	2	609	S.I.(ft) 50	USDA 1929, Tabs. 33-38
271	Pinus palustris	G	1	609	S.I.(ft) 50	USDA 1929, Tabs. 68-70
272	Pinus ponderosa	G	2	472	S.I.(ft) 100	Show 1925, Tab. 2
273	Pinus echinata	G	2	609	S.I.(ft) 50	USDA 1929, Tabs. 98-102
274	Pinus elliottii	G	3	609	S.I.(ft) 50	USDA 1929, Tabs. 130-134
276	Sequoia sempervirens	G	4	416	S.I.(ft) 50	Bruce 1923, Tabs. 1-3
277	<u>Liquidamber</u> styraciflua	Α	2	593	S.I.(ft) 50	Winters and Osborne 1935, Tabs. 4-8
278	Tsuga heterophylla	G	5	520	S.I.(ft) 100	Barnes 1962, Tab. 27
279	Eucalyptus delegatensis	E		617	S.I.(m) 50	Hillis and Brown 1978, Tab. 10.1
280	Eucalyptus regnans	Ε		689	S.I.(m) 50	Hillis and Brown 1978, Tab. 10.22
281	Eucalyptus sieberi	Ε		840	S.I.(m) 50	Hillis and Brown 1978, Tab. 10.27

aAn arbitrary number assigned to facilitate cross-referencing among tables.

Done of three categories. A = temperate angiosperms, E = trees of genus Eucalyptus, G = temperate gymnosperms.

CFrom references above or Baker (1949).

 $^{
m d}$ From references above or Peattie (1950, 1953). Used to convert stand volumes (m $^{
m 3}$ of wood per m $^{
m 2}$ of ground area) to stand biomasses (grams of wood per m $^{
m 2}$).

eDifferences in growing site or cultural practice that may affect thinning line. S.I. = height of trees (in either feet or meters, as indicated) when the stand is at the specified age (in years). This measure, called 'site index', is a widely used indicator of the quality of a site for tree growth. S.Q. = Site quality measured by ranks rather than by site index. Lower numbers are of lower quality. Spacing = ranked initial densities in tree plantations, with lower numbers indicating higher initial densities. Crown width = a ranked measure of canopy openness. Lower numbers indicate a more open canopy. See references for further details.

fData are taken from the indicated tables in the reference.

9Generic name only indicates that the specific name was not given or that more than one species from the genus were present.

Table B.2. Self-thinning Lines Fit to Forestry Yield Table Data.

T.,		No. Poi	of nts ^c		PCA T	hinnin	ıg Line ^e		Log N ^f			Log B ^f	-
Id. Code ^a	Cond. ^b	ηŢ	n	r ^d	β̂	â	p _{est}	Min.	Max.	Mean	Min.	Max.	Mean
Code ^a 201 201 201 201 201 201 201 201 201 20	Cond. b 40 50 60 70 80 90 100 110 120 130 140 150 160 80 96 112 30 40 50 60 70 80 90 62 71 80 90 62 71 80 58 68 80 52 64 75	nT 18 19 19 19 19 19 19 19 19 9 9 9 11 11	13 14 14 14 14 14 14 14 4 4 9 9 9 8 8 8 8 8 8 8 9 9 9	rd -0.998 -0.997 -1.000 -0.999 -0.999 -0.997 -0.993 -0.998 -0.998 -0.998 -0.998 -0.998 -0.999 -0.998 -0.999 -0.998 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999 -0.999	6 -0.308 -0.310 -0.324 -0.326 -0.318 -0.318 -0.318 -0.318 -0.479 -0.524 -0.653 -0.479 -0.563 -0.575 -0.585 -0.564 -0.558 -0.570 -0.580 -0.630 -0.630 -0.691 -0.698	â 3.75 3.79 3.83 3.95 4.08 4.12 4.18 3.90 4.01 3.30 3.59 3.67 3.89 3.95 3.64 3.56 3.64 3.56 3.64 3.56 3.67 3.31 3.50	Pest 0.382 0.382 0.378 0.377 0.378 0.379 0.380 0.383 0.379 0.334 0.338 0.328 0.303 0.340 0.3317 0.315 0.320 0.321 0.321 0.321 0.316 0.305 0.305 0.305 0.305 0.305 0.296 0.307 0.277 0.286 0.294	Min. -1.27 -1.38 -1.49 -1.57 -1.64 -1.71 -1.76 -1.80 -1.84 -1.87 -1.46 -1.49 -1.52 -1.38 -1.46 -1.54 -1.52 -1.38 -1.16 -1.23 -1.31 -1.23 -1.31 -1.23 -1.31 -1.23 -1.31 -1.23 -1.31 -1.23	Max. -0.49 -0.52 -0.69 -0.81 -0.91 -0.99 -1.06 -1.12 -1.17 -1.22 -1.26 -1.30 -1.33 -0.76 -0.86 -0.94 -0.59 -0.73 -0.83 -0.91 -0.99 -1.07 -1.15 -0.91 -0.92 -1.01 -0.72 -0.88 -0.95 -0.72 -0.88	Mean -0.97 -1.07 -1.18 -1.27 -1.35 -1.47 -1.52 -1.56 -1.59 -1.37 -1.40 -1.43 -1.15 -1.24 -1.31 -0.73 -0.87 -0.96 -1.05 -1.13 -1.21 -1.29 -1.04 -1.09 -1.17 -0.90 -1.17 -0.92 -1.04	Min. 3.90 3.93 4.01 4.09 4.17 4.25 4.33 4.41 4.48 4.55 4.61 4.71 3.79 3.98 4.07 4.00 4.10 4.25 4.40 4.51 4.58 4.63 4.13 4.22 4.32 4.32 4.32 4.32 4.32 4.32 4.3	Max. 4.14 4.21 4.27 4.34 4.41 4.456 4.63 4.70 4.77 4.71 4.19 4.27 4.39 4.54 4.64 4.39 4.54 4.64 4.39 4.54 4.64 4.39 4.54 4.64 4.39 4.54 4.64 4.39 4.54 4.64 4.39 4.54 4.39 4.54 4.39 4.54 4.39 4.42 4.39 4.42 4.39 4.42 4.39 4.43 4.39 4.39 4.39	Mean 4.05 4.11 4.17 4.25 4.32 4.40 4.62 4.69 4.66 4.71 4.76 4.05 4.17 4.25 4.19 4.33 4.48 4.59 4.33 4.48 4.59 4.33 4.48 4.59 4.33 4.48 4.59 4.33 4.49 4.33 4.49 4.33
207 208 208 208 208 209 209 209 210 211 211	49 40 50 60 70 1 2 3 4 136 20 30	11 8 8 8 7 9 11 13 44 17	11 6 6 6 6 4 6 7 8 28 14	-0.992 -0.997 -0.997 -0.998 -0.996 -0.937 -0.947 -0.985 -0.998 -0.998 -0.998	-0.787 -0.795 -0.803 -0.796 -0.766 -0.318 -0.282 -0.270 -0.433 -0.199 -0.537 -0.529	2.94 4.01 3.94 3.83 3.75 3.46 3.63 3.74 3.66 3.85 3.53 3.76	0.280 0.279 0.277 0.278 0.283 0.379 0.390 0.394 0.417 0.325 0.327	-1.79 -0.66 -0.84 -1.07 -1.27 -1.46 -1.57 -1.33 -2.10 -0.42 -0.53	-0.76 -0.13 -0.31 -0.53 -0.73 -0.97 -0.89 -0.92 -0.95 -1.49 0.26 0.16	-1.36 -0.49 -0.66 -0.89 -1.10 -1.20 -1.21 -1.16 -1.83 -0.17 -0.27	3.57 4.11 4.18 4.25 4.31 3.76 3.85 3.96 4.05 4.15 3.38 3.67	4.42 4.55 4.63 4.69 4.75 3.92 4.05 4.14 4.23 4.27 3.74 4.02	4.01 4.39 4.47 4.54 4.59 3.85 3.97 4.07 4.16 4.22 3.62 3.90

Table B.2. (continued)

Id.		No. Poi	of nts ^c		PCA T	hinnin	g Line ^e		Log N ^f			Log B ^f	
Code ^a	Cond.b	nΤ	n	rd	β	â	^p est	Min.	Max.	Mean	Min.	Max.	Mean
211 211 211 211	40 50 60 70	17 17 17 17	14 14 14 14	-0.999 -0.999 -0.998 -0.998	-0.510 -0.535 -0.534 -0.535	3.86 3.89 3.88 3.84	0.331 0.326 0.326 0.326	-0.65 -0.81 -1.02 -1.25	0.05 -0.14 -0.34 -0.56	-0.40 -0.56 -0.76 -0.99	3.83 3.96 4.05 4.13	4.19 4.32 4.42 4.49	4.07 4.19 4.29 4.37
212 212 212 212 212	60 80 100 120 140	18 19 19 19	14 15 15 15	-0.994 -0.993 -0.993 -0.994 -0.992	-0.509 -0.551 -0.542 -0.554 -0.572	3.95 3.97 4.03 4.05 4.03	0.331 0.322 0.324 0.322 0.318	-1.06 -1.19 -1.34 -1.50 -1.65	-0.50 -0.55 -0.68 -0.84 -1.03	-0.83 -0.93 -1.08 -1.24 -1.41	4.18 4.24 4.37 4.49 4.59	4.48 4.61 4.74 4.86 4.96	4.37 4.49 4.61 4.73 4.84
212 212 212 213	160 180 200 30	19 19 19 10	15 15 15 6	-0.991 -0.992 -0.993 -0.998	-0.556 -0.586 -0.565 -0.545	4.06 3.99 3.99 3.72	0.321 0.315 0.319 0.324	-1.80 -1.92 -2.06 -0.38	-1.17 -1.31 -1.43 -0.11	-1.55 -1.68 -1.81 -0.29	4.67 4.72 4.77 3.78	5.04 5.10 5.14 3.93	4.92 4.97 5.01 3.88
213 213 213 213 214	40 50 60 70 1	10 10 10 10 11	6 6 6 6	-0.996 -0.998 -0.997 -0.997 -0.998	-0.536 -0.542 -0.538 -0.540 -0.636	3.83 3.88 3.93 3.97 3.79	0.325 0.324 0.325 0.325 0.306	-0.65 -0.81 -0.91 -0.98 -0.96	-0.39 -0.54 -0.64 -0.71 -0.64	-0.57 -0.73 -0.82 -0.90 -0.82	4.04 4.18 4.28 4.36 4.19	4.18 4.32 4.42 4.50 4.39	4.13 4.27 4.37 4.45 4.31
214 214 214 214 214	2 3 4 5 6	13 13 13 13	9 9 9 9	-0.997 -0.997 -0.998 -0.999 -0.999	-0.638 -0.622 -0.625 -0.606 -0.608	3.79 3.83 3.85 3.91 3.95	0.305 0.308 0.308 0.311 0.311	-1.20 -1.33 -1.40 -1.45 -1.50	-0.73 -0.87 -0.97 -1.03 -1.09	-1.01 -1.15 -1.23 -1.29 -1.34	4.25 4.36 4.45 4.53 4.60	4.55 4.65 4.73 4.79 4.86	4.44 4.54 4.62 4.69 4.77
215 215 215 215 215	1 2 3 4	9 10 11 11	9 10 11 11	-0.999 -0.998 -0.995 -0.996	-0.994 -0.964 -0.971 -0.939	3.65 3.66 3.65 3.68	0.251 0.255 0.254 0.258	-0.75 -0.91 -1.04 -1.15	0.11 0.23 0.45 0.27	-0.46 -0.51 -0.55 -0.68	3.53 3.40 3.12 3.36	4.37 4.50 4.61 4.72	4.10 4.15 4.18 4.32
215 215 216 216 216	5 6 1 2 3]]]]]]]]	 	-0.997 -0.998 -1.000 -0.999 -0.998	-0.905 -0.904 -0.729 -0.739 -0.747	3.74 3.76 3.65 3.66 3.66	0.262 0.263 0.289 0.288 0.286	-1.22 -1.28 -1.48 -1.59 -1.68	0.07 -0.08 0.04 -0.17 -0.32	-0.79 -0.90 -0.96 -1.10 -1.21	3.62 3.79 3.61 3.76 3.87	4.81 4.90 4.72 4.81 4.89	4.45 4.58 4.35 4.47 4.56
217 217 217 218 218	54 64 75 50 60	10 10 10 10 10	6 6 6 10 10	-0.997 -0.997 -0.996 -0.997 -0.999	-0.562 -0.544 -0.528 -0.776 -0.803	3.77 3.82 3.87 3.58 3.59	0.320 0.324 0.327 0.281 0.277	-1.27 -1.35 -1.42 -1.09 -1.19	-0.72 -0.88 -1.00 -0.16 -0.27	-1.03 -1.14 -1.23 -0.79 -0.90	4.16 4.29 4.39 3.69 3.80	4.47 4.55 4.61 4.40 4.53	4.34 4.44 4.53 4.19 4.32
218 220 220 220 220 220	70 40 50 60 70	10 9 9 9	10 6 6 6 6	-0.998 -0.995 -0.992 -0.991 -0.987	-0.788 -1.065 -0.985 -0.879 -0.746	3.65 3.43 3.50 3.59 3.69	0.280 0.242 0.252 0.266 0.286	-1.26 -0.60 -0.72 -0.85 -0.99	-0.35 -0.31 -0.40 -0.49 -0.56	-0.97 -0.47 -0.57 -0.68 -0.78	3.91 3.74 3.88 4.00 4.09	4.62 4.07 4.20 4.32 4.41	4.41 3.93 4.07 4.19 4.28
220 220	80 90	9 9	6 6	-0.984 -0.979	-0.623 -0.528	3.79 3.88	0.308 0.327	-1.15 -1.30	-0.64 -0.71	-0.90 -1.00	4.16 4.22	4.48 4.54	4.35 4.41

Table B.2. (continued)

Id.		No. <u>Poi</u>	of nts		PCA T	hinnin	g Line ^e		Log N ^f			Log B ^f	
Code ^a	Cond.b	nŢ	n	r ^d	β̂	â	p _{est}	Min.	Max.	Mean	Min.	Max.	Mean
221	80	15	13	-0.996	-0.629	3.64	0.307	-1.21	-0.42	-0.94	3.88	4.38	4.23
221	90	15	13	-0.995	-0.655	3.66	0.302	-1.25	-0.50	-1.00	3.96	4.46	4.31
221	100	15	13	-0.996	-0.667	3.68	0.300	-1.31	-0.57	-1.05	4.03	4.53	4.38
221	110	15	13	-0.995	-0.684	3.69	0.297	-1.36	-0.64	-1.11	4.10	4.60	4.45
221	120	15	13	-0.994	-0.683	3.72	0.297	-1.43	-0.71	-1.17	4.17	4.67	4.52
221	130	15	13	-0.993	-0.695	3.72	0.295	-1.48	-0.77	-1.23	4.23	4.73	4.58
221	140	15	13	-0.992	-0.693	3.73	0.295	-1.54	-0.84	-1.29	4.27	4.77	4.62
221 221	150	15	13	-0.992	-0.697	3.73	0.295	-1.60	-0.90	-1.35	4.31	4.81	4.66
	160	15 15	13	-0.990	-0.696	3.72	0.295	-1.65	-0.96	-1.40	4.34	4.84	4.69
221 221	170 180	15	13 13	-0.990 -0.992	-0.694	3.71	0.295	-1.72	-1.02	-1.46	4.37	4.87	4.72
221	190	15	13		-0.696	3.68	0.295	-1.78	-1.08	-1.53	4.39 4.41	4.89 4.91	4.74 4.76
221	200	15	13	-0.991 -0.992	-0.700 -0.705	3.64 3.60	0.294 0.293	-1.85 -1.93	-1.16 -1.23	-1.60 -1.67	4.41	4.91	4.78
221	210	15	13	-0.992	-0.691	3.59	0.293	-2.01	-1.23	-1.75	4.45	4.95	4.80
222	1	35	8	-0.997	-0.631	4.21	0.307	-0.66	-0.44	-0.52	4.48	4.62	4.54
222	2	35	8	-0.997	-0.880	3.98	0.266	-0.72	-0.54	-0.61	4.45	4.61	4.51
222	3	35	8	-0.996	-1.242	3.62	0.223	-0.78	-0.64	-0.69	4.42	4.59	4.49
222	4	35	5	-0.999	-1.535	3.27	0.197	-0.85	-0.77	-0.80	4.44	4.57	4.50
222	5	35	5	-0.995	-1.975	2.73	0.168	-0.92	-0.86	-0.88	4.42	4.55	4.47
223	ĭ	15	13	-0.999	-0.338	4.02	0.374	-1.59	-0.84	-1.35	4.30	4.55	4.48
224	3	17	14	-0.995	-0.504	3.91	0.333	-1.36	-0.61	-1.14	4.23	4.60	4.49
225	ĺ	6	6	-0.998	-0.687	3.40	0.296	-1.32	-0.56	-1.02	3.78	4.31	4.09
227	40	16	13	-0.994	-0.511	2.99	0.331	-1.41	-0.79	-1.17	3.41	3.73	3.59
227	50	16	13	-0.995	-0.508	3.26	0.331	-1.46	-0.83	-1.22	3.70	4.02	3.88
227	60	16	13	-0.995	-0.508	3.41	0.332	-1.51	-0.88	-1.27	3.88	4.20	4.06
227	70	16	13	-0.995	-0.509	3.50	0.331	-1.57	-0.94	-1.33	4.00	4.32	4.18
227	80	16	13	-0.994	-0.509	3.55	0.331	-1.63	-1.01	-1.39	4.07	4.39	4.26
227	90	16	13	-0.994	-0.509	3.57	0.331	-1.70	-1.07	-1.46	4.13	4.45	4.31
227	100	16	13	-0.993	-0.508	3.60	0.332	-1.77	-1.14	-1.53	4.20	4.52	4.38
227	110	16	13	-0.994	-0.509	3.64	0.331	-1.84	-1.21	-1.60	4.27	4.59	4.45
227	120	16	13	-0.977	-0.509	3.67	0.331	-1.91	-1.28	-1.68	4.35	4.67	4.53
228	25	14	11	-0.998	-0.407	3.46	0.355	-0.55	0.06	-0.34	3.42	3.68	3.59
228	35	14	11	-0.998	-0.372	3.52	0.364	-0.75	-0.11	-0.53	3.56	3.80	3.72
228	45	15	12	-0.997	-0.368	3.56	0.366	-0.93	-0.10	-0.65	3.58	3.89	3.80
228 228	55 65	15 15	12 12	-0.998	-0.340	3.60	0.373	-1.07	-0.24	-0.79	3.67	3.96	3.87
228	75	15	12	-0.998 -0.998	-0.319 -0.301	3.64	0.379 0.384	-1.21 -1.34	-0.37	-0.93	3.75 3.81	4.02	3.94 3.98
229	60	7	7	-0.998	-0.301	3.67 3.05	0.384	-1.34	-0.49 -0.80	-1.05 -1.09	3.76	4.06 4.14	4.02
229	70	7	7	-0.999	-0.898	3.05	0.263	-1.23	-0.80	-1.19	3.86	4.24	4.12
229	80	7	7	-0.997	-0.895	3.03	0.264	-1.41	-0.99	-1.19	3.94	4.32	4.21
229	90	7	7	-0.997	-0.887	3.11	0.265	-1.48	-1.05	-1.34	4.03	4.41	4.30
229	100	7	7	-0.995	-0.888	3.13	0.265	-1.55	-1.12	-1.40	4.11	4.49	4.38
229	110	7	7	-0.992	-0.884	3.17	0.265	-1.60	-1.17	-1.45	4.18	4.56	4.45
229	120	7	7	-0.988	-0.880	3.21	0.266	-1.64	-1.21	-1.49	4.26	4.63	4.53

Table B.2. (continued)

1.4		No. Poi	of nts ^c		PCA T	hinnin	g Line ^e		Log N ^f			Log B ^f	
ld. Code ^a	Cond.b	nΤ	n	r^{d}	ß	â	pest	Min.	Max.	Mean	Min.	Max.	Mean
230	50	7	7	-0.999	-0.565	3.36	0.320	-1.09	-0.49	-0.90	3.64	3.98	3.87
230	60	7	7	-0.998	-0.602	3.36	0.312	-1.20	-0.67	-1.03	3.76	4.09	3.98
230	70	7	7	-0.999	-0.640	3.35	0.305	-1.30	-0.81	-1.14	3.88	4.19	4.08
230	80	7	7	-0.998	-0.680	3.34	0.298	-1.37	-0.93	-1.23	3.98	4.29	4.18
230	90	7	7	-0.998	-0.751	3.29	0.286	-1.43	-1.04	-1.31	4.08	4.38	4.28
230	100	7	7	-0.997	-0.833	3.22	0.273	-1.48	-1.13	-1.37	4.17	4.47	4.37
231	50	7	5	-1.000	-1.278	2.20	0.219	-1.36	-1.19	-1.29	3.72	3.93	3.85
231	60	7	5	-1.000	-1.283	2.34	0.219	-1. 37	-1.19	-1.30	3.87	4.09	4.00
231	70	7	5	-1.000	-1.274	2.48	0.220	-1.37	-1.20	-1.30	4.01	4.23	4.14
231	80	7	5	-1.000	-1.274	2.59	0.220	-1.38	-1.21	-1.31	4.12	4.34	4.25
231	90	7	5	-1.000	-1.278	2.68	0.220	-1.38	-1.21	-1.31	4.22	4.44	4.35
231	100	7	5	-1.000	-1.281	2.76	0.219	-1.39	-1.22	-1.32	4.32	4.54	4.45
232	40	7	7	-1.000	-0.937	3.18	0.258	-0.88	-0.37	-0.68	3.52	4.00	3.82
232	50	7	7	-1.000	-0.937	3.18	0.258	-1.05	-0.54	-0.85	3.68	4.16	3.98
232	60	7	7	-1.000	-0.927	3.19	0.260	-1.19	-0.68	-0.99	3.81	4.29	4.11
232	70	7	7	-1.000	-0.939	3.18	0.258	-1.30	-0.80	-1.11	3.92	4.40	4.22
232	80	7	7 7	-1.000	-0.944	3.17	0.257	-1.41	-0.90	-1.21	4.02	4.50	4.31
232	90	7		-1.000	-0.968	3.14	0.254	-1.50	-1.00	-1.30	4.09	4.58	4.39
232 233	100 50	7 7	7 7	-1.000	-0.961	3.15	0.255	-1.57	-1.07	-1.38	4.17	4.66	4.47
233	60	7	7	-1.000	-0.569	3.32	0.319	-1.02	-0.45	-0.84	3.58	3.90	3.79
233	70	7	7	-0.999 -0.999	-0.615	3.33	0.310	-1.10	-0.55	-0.93	3.67	4.01 4.11	3.90 4.00
233	80	7	7	-0.999	-0.664 -0.723	3.33 3.32	0.301 0.290	-1.17 -1.22	-0.63 -0.70	-1.00 -1.05	3.76 3.83	4.21	4.08
233	90	7	7	-0.998	-0.802	3.29	0.278	-1.25	-0.75	-1.10	3.90	4.30	4.17
233	100	7	7	-0.996	-0.898	3.23	0.263	-1.27	-0.73	-1.13	3.97	4.39	4.25
234	70	6	5	-0.989	-0.441	3.68	0.203	-1.43	-1.05	-1.28	4.14	4.31	4.25
234	84	9	7	-0.963	-0.246	4.10	0.401	-1.54	-1.07	-1.36	4.37	4.48	4.43
234	99	9	7	-0.962	-0.449	3.91	0.345	-1.57	-1.17	-1.41	4.44	4.60	4.54
235	40		18	-0.995	-0.888	2.97	0.265	-1.14	-0.28	-0.85	3.17	3.94	3.72
235	50		18	-0.999	-0.804	3.26	0.277	-1.17	-0.31	-0.89	3.50	4.19	3.97
235	60		18	-0.999	-0.813	3.41	0.276	-1.21	-0.35	-0.93	3.69	4.37	4.16
235	70		18	-0.999	-0.829	3.49	0.273	-1.25	-0.40	-0.97	3.81	4.51	4.29
235	80	18	18	-0.999	-0.854	3.52	0.270	-1.31	-0.45	-1.02	3.87	4.61	4.39
235	90	18	18	-0.998	-0.882	3.51	0.266	-1.36	-0.51	-1.08	3.93	4.69	4.46
235	100	18	18	-0.998	-0.897	3.50	0.264	-1.42	-0.57	-1.14	3.97	4.75	4.52
235	110	18	18	-0.997	-0.881	3.50	0.266	-1.49	-0.62	-1.20	4.02	4.78	4.56
236	40	10	9	-0.996	-0.772	3.29	0.282	-0.99	-0.09	-0.70	3.40	4.09	3.84
236	50	10	9	-0.996	-0.781	3.32	0.281	-1.10	-0.21	-0.81	3.52	4.21	3.96
236	60	10	9	-0.996	-0.780	3.34	0.281	-1.21	-0.32	-0.93	3.62	4.31	4.06
236	70	10	9	-0.996	-0.779	3.34	0.281	-1.32	-0.43	-1.04	3.70	4.39	4.14
236	80		10	-0.997	-0.833	3.26	0.273	-1.44	-0.22	-1.06	3.40	4.46	4.13
237	1		12	-0.988	-0.620	3.83	0.309	-0.81	-0.40	-0.55	4.07	4.31	4.17
237	2	28	8	-0.996	-0.702	3.70	0.294	-0.86	-0.61	-0.70	4.12	4.30	4.19
237	3	28	8	-0.994	-0.922	3.45	0.260	-0.91	-0.70	- 0.77	4.09	4.28	4.17

Table B.2. (continued)

* .		No. Poi	of nts ^c		PCA T	hinnin	g Line ^e		Log N ^f			Log B ^f	
Id. Code ^a	Cond.b	nŢ	n	r ^d	β̂	â	p _{est}	Min.	Max.	Mean	Min.	Max.	Mean
237	4	28	8	-0.995	-1.158	3.18	0.232	-0.95	-0.77	-0.84	4.07	4.27	4.15
237	5	28	5	-0.999	-1.280	3.01	0.219	-0.97	-0.86	-0.91	4.11	4.26	4.18
238	40	8	5	-0.997	-0.598	3.77	0.313	-0.54	-0.27	-0.44	3.93	4.10	4.03
238	50	8	5	-0.998	-0.591	3.80	0.314	-0.71	-0.43	-0.61	4.06	4.22	4.16
238	60	8	5	-0.998	-0.588	3.81	0.315	-0.86	-0.58	-0.76	4.16	4.32	4.26
238	70	8	5	-0.998	-0.589	3.81	0.315	-1.01	-0.73	-0.90	4.24	4.40	4.34
239	40	8	5	-0.999	-0.524	3.72	0.328	-0.71	-0.35	-0.58	3.90	4.10	4.02
239	50	8	5	-0.999	-0.524	3.78	0.328	-0.82	-0.45	-0.69	4.02	4.22	4.14
239	60	8	5	-0.998	-0.521	3.82	0.329	-0.94	-0.57	-0.80	4.11	4.31	4.23
239	70	8	5	-0.998	-0.524	3.82	0.328	-1.07	-0.71	-0.94	4.19	4.39	4.31
241	20	6	5	-0.990	-4.100	2.03	0.098	-0.25	-0.20	-0.23	2.84	3.08	2.99
241	30	6	5	-0.997	-4.571	1.86	0.090	-0.40	-0.35	-0.38	3.47	3.69	3.61
241 241	4 0 50	6 6	5	-0.984	-3.830	1.92	0.104	-0.51	-0.45	-0.49	3.66	3.89 3.96	3.80 3.87
241	60	6	5 5	-0.991	-3.935 -4.467	1.50	0.101	-0.62	-0.57	-0.60	3.73	4.00	3.91
242	56	6	3	-0.984 -0.963	-0.393	0.68 3.23	0.091 0.359	-0.74 -1.33	-0.69 -1.29	-0.72 -1.31	3.77 3.74	3.75	3.74
242	73	7	4	-0.964	-0.393	3.78	0.339	-1.33	-1.29	-1.28	3.74	3.98	3.97
242	73 90	8	4	-0.951	-0.119	4.08	0.433	-1.34	-1.22	-1.30	4.22	4.24	4.23
243	ĵ	12	7	-0.998	-0.551	2.69	0.322	-2.33	-1.89	-2.13	3.73	3.96	3.86
243	2	7	5	-0.969	-0.480	3.38	0.338	-1.69	-1.20	-1.46	3.96	4.17	4.08
244	70	6	3	-1.000	-1.352	2.58	0.213	-1.31	-1.25	-1.28	4.27	4.35	4.31
244	86	6	4	-1.000	-0.898	3.24	0.263	-1.38	-1.25	-1.32	4.36	4.48	4.42
244	102	6	5	-0.996	-0.753	3.49	0.285	-1.44	-1.23	-1.34	4.41	4.57	4.51
245	1	12	5	-0.995	-0.568	3.28	0.319	-1.63	-1.45	-1.54	4.10	4.20	4.15
245	2	11	5	-0.991	-0.437	3.54	0.348	-1.73	-1.53	-1.63	4.21	4.29	4.25
245	3	9	4	-0.998	-0.446	3.55	0.346	-1.79	-1.63	-1.72	4.28	4.35	4.32
246	30	4	4	-0.997	-0.874	3.08	0.267	-0.56	0.01	-0.29	3.07	3.56	3.34
246	50	6	4	-0.984	-0.470	3.51	0.340	-1.01	-0.49	-0.75	3.72	3.96	3.86
246	60	7	4	-0.998	-0.352	3.74	0.370	-1.24	-0.80	-1.04	4.02	4.17	4.11
246	7 0	7	4	-0.992	-0.348	3.81	0.371	-1.35	-0.91	-1.15	4.12	4.27	4.21
246	80	7	4	-0.997	-0.361	3.83	0.367	-1.44	-1.01	-1.24	4.19	4.34	4.28
247]	7	4	-0.990	-3.804	2.06	0.104	-0.57	-0.55	-0.56	4.13	4.22	4.18
247	2	7	4	-0.988	-4.973	1.53	0.084	-0.58	-0.56	-0.57	4.31	4.43	4.38
247	3	7	4	-0.994	-6.286	0.85	0.069	-0.60	-0.58	-0.59	4.47	4.61	4.55
247	4	7	4	-0.991	-8.132		0.055	-0.62	-0.59	-0.60	4.60	4.79	4.70
248	1	9	5	-0.996	-6.444		0.067	-1.07	-0.95	-1.02	2.98	3.69	3.40
248	2 3	10 11	6 7	-0.999	-6.855		0.064	-1.12	-0.97	-1.06	3.02	4.01	3.59 3.72
248 2 49	100	9	3	-0.992		-4.18	0.061 0.223	-1.16	-0.98	-1.09	2.98 4.11	4.22 4.22	4.17
249	110	9	3 3	-0.995 -0.998	-1.243 -1.188	2.53 2.64	0.223	-1.36 -1.39	-1.27 -1.30	-1.31 -1.34	4.11	4.22	4.17
249	120	9	3	-0.998	-1.188		0.229	-1.39	-1.30	-1.34	4.18	4.29	4.29
250	2	13	4	-0.999	-0.659	2.88 3.69	0.301	-1.44	-1.09	-1.13	4.23	4.47	4.44
250	3	13	8	-0.993	-0.549	3.82	0.323	-1.39	-1.10	-1.26	4.42	4.59	4.51
251	ĭ	27	-	-0.999	-0.645	3.74	0.304	-1.77	-0.42	-1.30	3.99	4.87	4.58
231	1	21	_ /	~0. 555	-0.045	3.74	0.304	-1.//	-0.42	-1.50	3.39	4.07	4.30

Table 8.2. (continued)

Code	7.1		No. of Points		PCA 1	hinnin	g Line ^e		Log N ^f			Log B	
252 140 9 5 -0.987 -4.533 -1.78 0.089 -1.42 -1.35 -1.35 3.82 4.84 4. 252 180 9 9 -0.999 -3.084 0.56 0.122 -1.43 -1.135 3.82 4.84 4. 252 200 9 9 -0.999 -2.392 1.66 0.147 -1.41 -1.07 -1.29 4.24 5.03 4. 252 220 9 9 -0.999 -2.032 1.25 0.164 -1.39 -1.03 -1.27 4.36 5.11 4. 253 47 14 10 -0.999 -0.888 3.35 0.269 -1.18 -0.99 -1.24 4.46 5.17 4. 2.53 4.7 14 10 -0.999 -0.888 3.35 0.269 -1.42 -0.39 -1.00 3.85 4.32 4.66 4. 2.53 66 15.14 -0.368 2.67	Id. Codeª	Cond.b	nŢ n	rd	β̂	â	p _{est}	Min.	Max.	Mean	Min.	Max.	Mean
252 160 9 9 -0.996 -4.748 -1.93 0.087 -1.43 -1.21 -1.35 3.82 4.84 4.97 4.94 4.25 252 180 9 9 -0.999 -2.392 1.66 0.147 -1.41 -1.07 -1.29 4.24 5.03 4.94 4.94 4.25 220 9 9 -0.999 -2.043 2.25 0.164 -1.39 -1.03 -1.27 4.36 5.11 4.25 6.61 1.18 -0.59 -0.999 -1.814 2.67 0.178 -1.38 -0.99 -1.24 4.66 5.17 4.25 3.40 0.099 -1.42 -0.43 3.85 4.35 4.25 4.66 5.11 4.0.998 -0.272 3.49 0.290 -1.42 -0.39 -1.00 3.77 4.48 4.2 2.54 1.00 3.17 4.42 4.2 4.2 4.39 4.19 4.2 4.43 4.2 4.56 4.2 4.2 4.2 </td <td></td> <td>4.55</td>													4.55
252 180 9 9 -0.999 -3.084 0.56 0.122 -1.43 -1.32 4.07 4.94 4. 252 220 9 9 -0.999 -2.092 1.66 0.147 -1.41 -1.07 -1.22 4.24 5.03 4. 252 220 9 9 -0.999 -2.083 2.25 0.164 -1.39 -1.03 -1.27 4.36 5.11 4. 253 47 14 10 -0.999 -0.858 3.35 0.269 -1.24 4.46 5.17 4. 253 66 15 14 -0.996 -0.618 3.60 0.309 -1.65 -0.36 -1.15 3.82 4.59 4. 254 60 12 12 -1.000 -0.947 3.24 0.257 -1.25 -0.57 -1.00 3.77 4.42 4. 254 61 13 9 -0.992 -0.691 3.64													4.60
252 200 9 9 -0.999 -2.392 1.66 0.147 -1.41 -1.07 -1.29 4.24 5.03 4. 252 220 9 9 -0.999 -2.043 2.25 0.164 -1.39 -1.03 -1.27 4.36 5.11 4. 253 47 14 10 -0.999 -0.888 3.35 0.269 -1.18 -0.999 -1.03 -1.27 4.46 5.17 4. 253 57 15 13 -0.998 -0.272 3.49 0.290 -1.42 -0.39 -1.00 3.77 4.48 4. 254 50 12 12 -1.000 -0.947 3.24 0.257 -1.25 -0.56 -1.15 3.82 4.59 4. 254 50 12 12 -1.000 -0.947 3.24 0.257 -1.25 -0.56 -1.33 -1.00 -1.19 4.22 4.56 4. <td< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>4.48</td></td<>													4.48
252 220 9 9 -0.999 -1.814 2.67 0.178 -1.38 -0.99 -1.27 4.36 5.17 4. 253 47 14 10 -0.999 -0.858 3.35 0.269 -1.18 -0.99 -1.024 4.46 5.17 4. 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.55 4.254 6.0 15 14 -0.996 -0.618 3.60 0.309 -1.65 -0.36 -1.15 3.82 4.59 4.254 4.254 61 13 9 -0.999 -0.691 3.64 0.296 -1.34 -1.00 -1.19 4.22 4.56 4.255 4.7 7 -0.999 -0.691 3.64 0.296 -1.34 -1.00 -1.19 4.32 4.66 4.255 4.7 4 -0.992 -0.680 3.51 0.266 -1.34 -1.00 -1.14													4.75
252 240 9 9 -0.999 -1.814 2.67 0.178 -1.38 -0.99 -1.24 4.46 5.17 4 253 47 14 10 -0.998 -0.2727 3.49 0.290 -1.42 -0.39 -1.00 3.77 4.48 4. 253 66 15 14 -0.996 -0.618 3.60 0.309 -1.65 -0.36 -1.15 3.82 4.59 4. 254 61 13 9 -0.999 -0.691 3.64 0.296 -1.34 -1.00 -1.19 4.32 4.56 4. 254 70 13 11 -0.992 -0.809 3.51 0.276 -1.44 -1.03 -1.28 4.32 4.66 4. 255 87 14 7 -0.993 -0.547 4.39 0.323 -1.76 -1.26 -1.56 5.06 5.33 5.63 3. 8.53 3.58 3.68 3. 256													4.84
253 47 14 10 -0.998 -0.727 3.49 0.290 -1.12 -0.39 -1.00 3.77 4.48 4. 253 66 15 14 -0.996 -0.618 3.60 0.309 -1.55 -0.36 -1.15 3.82 4.59 4. 254 50 12 12 -1.000 -0.947 3.24 0.257 -1.25 -0.57 -1.00 3.77 4.42 4. 254 61 13 9 -0.999 -0.609 3.51 0.276 -1.34 -1.00 -1.54 5.66 4. 255 87 14 7 -0.983 -0.547 4.39 0.323 -1.76 -1.26 -1.54 5.06 5.33 5. 256 40 7 4 -0.997 -0.688 2.91 0.268 -0.90 -0.51 -0.77 3.53 3.68 3. 256 63 7 4 -0.997	252												4.92
253 57 15 13 -0.996 -0.727 3.49 0.290 -1.42 -0.39 -1.00 3.77 4.48 4.254 50 15 14 -0.996 -0.618 3.60 0.309 -1.65 -0.36 -1.15 3.82 4.59 4.254 61 13 9 -0.999 -0.691 3.64 0.296 -1.34 -1.00 -1.19 4.32 4.56 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.6 4.25 4.56 4.4 4.52 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.22 4.56 4.	253												4.14
253 66 15 14 -0.996 -0.618 3.60 0.309 -1.65 -0.36 -1.15 3.82 4.59 4.254 4.50 1.21 -1.000 -0.947 3.24 0.257 -1.25 -0.57 -1.00 3.77 4.42 4.56 4.25 4.30 -1.09 -0.999 -0.691 3.64 0.296 -1.34 -1.00 -1.19 4.32 4.56 4.56 4.25 4.70 13 11 -0.992 -0.809 3.51 0.276 -1.44 -1.03 -1.28 4.32 4.66 4.6 4.255 8.7 4 -0.995 -0.577 3.03 0.282 -1.08 -0.67 -0.89 3.53 3.68 3.53 3.68 3.4 2.29 -0.688 2.91 0.268 -0.90 -0.578 -1.00 -0.78 -1.01 3.63 3.58 3.68 3.2 257 45 13 13 -1.000 -0.683 3.14 0.297 -1.28 -0.10 -0.95 3.13													4.22
254 50 12 12 -1.000 -0.947 3.24 0.257 -1.25 -0.57 -1.00 3.77 4.42 4.56 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.66 4.58 4.32 4.66 4.5 3.68 3.53 3.83 3.68 3.53 3.83 3.53 3.83 3.53 3.83 3.53 3.85 3.57 4.93		66	15 14	-0.996			0.309	-1.65	-0.36		3.82	4.59	4.31
254 70 13 11 -0.992 -0.809 3.51 0.276 -1.44 -1.03 -1.28 4.32 4.66 4. 255 87 14 7 -0.983 -0.547 4.39 0.323 -1.76 -1.26 -1.54 5.06 5.33 5.6 3. 256 40 7 4 -0.995 -0.773 3.03 0.282 -1.08 -0.67 -0.89 3.53 3.85 3. 256 66 7 4 -0.995 -0.673 3.03 0.282 -1.08 -0.67 -0.89 3.53 3.85 3. 257 45 13 13 -1.000 -0.683 3.04 0.297 -1.20 -0.78 -1.01 3.66 3.94 3. 257 49 13 11 -0.999 -0.655 3.00 0.302 -1.28 -0.10 3.59 3.99 3. 257 53 13 11 -0.999 -1.01					-0.947	3.24			-0.57	-1.00	3.77	4.42	4.18
255 87 14 7 -0.983 -0.547 4.39 0.323 -1.76 -1.26 -1.54 5.06 5.33 5.256 40 7 4 -0.997 -0.888 2.91 0.268 -0.90 -0.51 -0.72 3.35 3.68 3.256 53 7 4 -0.992 -0.683 3.14 0.297 -1.20 -0.78 -1.01 3.66 3.94 3.257 45 13 13 -1.000 -0.541 3.07 0.324 -1.28 -0.10 -0.95 3.13 3.76 3.257 49 13 11 -0.999 -0.655 3.00 0.302 -1.20 -0.55 -0.97 3.37 3.79 3.25 54 13 11 -0.999 -0.655 3.00 0.302 -1.20 -0.55 -0.97 3.37 3.79 3.25 57 54 13 11 -0.999 -0.011 2.98 0.249 -1.03 -0.55 -0.85 3.55 4.03													4.46
256 40 7 4 -0.997 -0.868 2.91 0.268 -0.90 -0.51 -0.72 3.35 3.68 3. 256 66 7 4 -0.995 -0.683 3.14 0.297 -1.20 -0.78 -1.01 3.66 3.94 3. 257 45 13 13 -1.000 -0.541 3.07 0.324 -1.28 -0.10 -0.95 3.13 3.76 3. 257 49 13 11 -0.999 -0.655 3.00 0.302 -1.20 -0.55 -0.97 3.37 3.79 3. 257 53 13 11 -1.000 -0.874 3.04 0.267 -1.08 -0.62 -0.91 3.59 3.99 3. 257 57 13 10 -1.000 -1.731 2.51 0.183 -0.91 -0.74 -0.84 3.79 4.09 3. 258 52 10 6 -0.991 -0.733 3.26 0.289<							0.276						4.55
256 53 7 4 -0.995 -0.773 3.03 0.282 -1.08 -0.67 -0.89 3.53 3.85 3. 256 66 7 4 -0.992 -0.683 3.14 0.297 -1.20 -0.78 -1.01 3.66 3.94 3. 257 45 13 13 -1.000 -0.541 3.07 0.324 -1.28 -0.10 -0.95 3.13 3.76 3. 257 53 13 11 -0.0999 -0.655 3.00 0.302 -1.08 -0.62 -0.91 3.59 3.99 3. 257 54 13 11 -0.999 -1.011 2.98 0.249 -1.08 -0.62 -0.91 3.59 3.99 3.29 257 54 13 11 -0.000 -1.731 2.51 0.183 -0.91 -0.74 -0.84 3.79 4.03 258 52 10 6 -0.998 </td <td></td> <td>5.23</td>													5.23
256 66 7 4 -0.992 -0.683 3.14 0.297 -1.20 -0.78 -1.01 3.66 3.94 3 257 45 13 13 -1.000 -0.541 3.07 0.324 -1.28 -0.10 -0.95 3.13 3.76 3 257 49 13 11 -0.999 -0.655 3.00 0.302 -1.20 -0.55 -0.97 3.37 3.79 3. 257 53 13 11 -0.999 -1.011 2.98 0.249 -1.03 -0.55 -0.91 3.59 3.99 3. 257 54 13 11 -0.999 -1.011 2.98 0.249 -1.03 -0.55 -0.85 3.55 4.03 3. 258 40 9 6 -0.733 3.26 0.289 -1.19 -0.87 -1.09 3.89 4.12 258 52 10 6 -0.997 -0.711													3.54
257 45 13 13 -1.000 -0.541 3.07 0.324 -1.28 -0.10 -0.95 3.13 3.76 3.257 49 13 11 -0.999 -0.655 3.00 0.302 -1.20 -0.55 -0.97 3.37 3.79 3.257 53 13 11 -1.000 -0.874 3.04 0.267 -1.08 -0.62 -0.91 3.59 3.99 3.257 57 13 10 -1.000 -1.731 2.51 0.183 -0.91 -0.74 -0.84 3.79 4.09 3.258 4.09 6 -0.991 -0.733 3.26 0.289 -1.19 -0.874 -1.09 3.89 4.12 4.258 60 10 6 -0.998 -0.824 3.19 0.274 -1.35 -1.04 -1.29 4.05 4.31 4.2 4.25 4.61 4.05 0.341 -1.59 -0.80 -1.27 4.40 4.78 4.2 259 100 17 14 -0.994 -0.466 4.05 <td< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>3.83</td></td<>													3.83
257 49 13 11 -0.999 -0.655 3.00 0.302 -1.20 -0.55 -0.97 3.37 3.79 3. 257 53 13 11 -1.000 -0.874 3.04 0.267 -1.08 -0.62 -0.91 3.59 3.99 3. 257 54 13 11 -0.999 -1.011 2.98 0.249 -1.03 -0.55 -0.85 3.55 4.03 3. 258 40 9 6 -0.991 -0.733 3.26 0.289 -1.19 -0.87 -1.09 3.89 4.12 4. 258 52 10 6 -0.998 -0.824 3.19 0.274 -1.35 -1.04 -1.24 4.05 4.31 4. 259 100 17 14 -0.999 -0.466 4.05 0.341 -1.59 -0.80 -1.27 4.40 4.78 4. 259 110 17 13 -0.999 -0.425 4.14 0.351 -1.64 -0.98 -1.36													3.58
257 53 13 11 -1.000 -0.874 3.04 0.267 -1.08 -0.62 -0.91 3.59 3.99 3. 257 54 13 11 -0.999 -1.011 2.98 0.249 -1.03 -0.55 -0.85 3.55 4.03 3. 258 40 9 6 -0.991 -0.733 3.26 0.289 -1.19 -0.87 -1.09 3.89 4.09 3. 258 52 10 6 -0.998 -0.824 3.19 0.274 -1.35 -1.04 -1.24 4.05 4.31 4. 258 60 10 6 -0.997 -0.711 3.40 0.292 -1.48 -1.17 -1.37 4.23 4.45 4. 259 100 17 13 -0.998 -0.425 4.14 0.351 -1.69 -1.03 -1.44 4.64 4.83 4. 259 120 17 13 -0.999 <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>3.64</td></t<>													3.64
257 57 13 10 -1.000 -1.731 2.51 0.183 -0.91 -0.74 -0.84 3.79 4.09 3.258 40 9 6 -0.991 -0.733 3.26 0.289 -1.19 -0.87 -1.09 3.89 4.12 4.25 4.05 4.31 4.24 4.25 4.05 4.31 4.25 4.45 4.25 4.45 4.25 4.45 4.25 4.45 4.25 4.45 4.25 4.45 4.25 4.45 4.25 4.45 4.25 4.14 0.351 -1.64 -0.80 -1.27 4.40 4.78 4.25 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4.25 1.00 17 13 -0.998 -0.425 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4.25 1.00 17 13 -0.999 -0.396 4.23 0.355 -1.69 -1.03 -1.40 4.54 4.83 4.25 0.360													3.84
258 40 9 6 -0.991 -0.733 3.26 0.289 -1.19 -0.87 -1.09 3.89 4.12 4.258 52 10 6 -0.998 -0.824 3.19 0.274 -1.35 -1.04 -1.24 4.05 4.31 4.25 60 10 6 -0.997 -0.711 3.40 0.292 -1.48 -1.17 -1.37 4.23 4.45 4. 259 100 17 14 -0.994 -0.466 4.05 0.341 -1.59 -0.80 -1.27 4.40 4.78 4. 259 110 17 13 -0.998 -0.425 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4.28 259 120 17 13 -0.999 -0.396 4.23 0.355 -1.69 -1.03 -1.40 4.64 4.90 4.25 9.360 -1.75 -1.11 -1.44 4.64 4.90 4.25 9.360 -1.75 -1.11	257	54	13 11	-0.999	-1.011	2.98	0.249	-1.03	-0.55	-0.85	3.55	4.03	3.84
258 52 10 6 -0.998 -0.824 3.19 0.274 -1.55 -1.04 -1.24 4.05 4.31 4.25 258 60 10 6 -0.997 -0.711 3.40 0.292 -1.48 -1.17 -1.37 4.23 4.45 4. 259 100 17 14 -0.994 -0.466 4.05 0.341 -1.59 -0.80 -1.27 4.40 4.78 4. 259 110 17 13 -0.998 -0.425 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4. 259 120 17 13 -0.999 -0.396 4.23 0.358 -1.72 -1.08 -1.44 4.64 4.90 4. 259 130 17 13 -0.999 -0.378 4.29 0.363 -1.75 -1.11 -1.47 4.68 4.93 4. 259 160 17 1				-1.000	-1.731		0.183	-0.91	-0.74	-0.84	3.79	4.09	3.97
258 60 10 6 -0.997 -0.711 3.40 0.292 -1.48 -1.17 -1.37 4.23 4.45 4. 259 100 17 14 -0.994 -0.466 4.05 0.341 -1.59 -0.80 -1.27 4.40 4.78 4. 259 110 17 13 -0.998 -0.425 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4. 259 120 17 13 -0.996 -0.407 4.19 0.355 -1.69 -1.03 -1.40 4.59 4.86 4. 259 130 17 13 -0.999 -0.396 4.23 0.358 -1.72 -1.08 -1.44 4.64 4.90 4. 259 140 17 13 -0.999 -0.378 4.29 0.363 -1.75 -1.11 -1.47 4.68 4.93 4. 259 150 17 13 -0.999 -0.378 4.31 0.363						3.26							4.06
259 100 17 14 -0.994 -0.466 4.05 0.341 -1.59 -0.80 -1.27 4.40 4.78 4.25 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4.25 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4.25 4.19 0.355 -1.69 -1.03 -1.40 4.59 4.86 4.25 0.360 -1.75 -1.10 -1.40 4.59 4.86 4.29 0.355 -1.72 -1.08 -1.40 4.59 4.86 4.29 0.360 -1.75 -1.11 -1.47 4.68 4.93 4.29 0.360 -1.75 -1.11 -1.47 4.68 4.93 4.29 0.363 -1.79 -1.14 -1.51 4.71 4.97 4.25 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.25 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.25 0.363 -1.81 -1.16 -1.53													4.22
259 110 17 13 -0.998 -0.425 4.14 0.351 -1.64 -0.98 -1.36 4.54 4.83 4.259 120 17 13 -0.996 -0.407 4.19 0.355 -1.69 -1.03 -1.40 4.59 4.86 4.259 130 17 13 -0.997 -0.396 4.23 0.358 -1.72 -1.08 -1.44 4.64 4.90 4.259 140 17 13 -0.999 -0.396 4.25 0.363 -1.75 -1.11 -1.47 4.68 4.93 4.25 0.363 -1.75 -1.11 -1.47 4.68 4.93 4.25 0.363 -1.79 -1.14 -1.51 4.71 4.97 4.25 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.25 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.25 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.25 0.363 -1.81 -1.18 -1.18 -1.18 -1.18<													4.37
259 120 17 13 -0.996 -0.407 4.19 0.355 -1.69 -1.03 -1.40 4.59 4.86 4. 259 130 17 13 -0.997 -0.396 4.23 0.358 -1.72 -1.08 -1.44 4.64 4.90 4. 259 140 17 13 -0.999 -0.390 4.25 0.360 -1.75 -1.11 -1.47 4.68 4.93 4. 259 150 17 13 -0.999 -0.378 4.29 0.363 -1.79 -1.14 -1.51 4.71 4.97 4. 259 160 17 13 -0.999 -0.379 4.31 0.363 -1.81 -1.16 -1.53 4.71 4.97 4. 259 170 17 13 -0.998 -0.383 4.32 0.362 -1.83 -1.18 -1.55 4.77 5.02 4. 259 180 17 13 -0.999 -0.377 4.37 0.363 -1.84													4.65 4.71
259 130 17 13 -0.997 -0.396 4.23 0.358 -1.72 -1.08 -1.44 4.64 4.90 4.259 259 140 17 13 -0.999 -0.390 4.25 0.360 -1.75 -1.11 -1.47 4.68 4.93 4.259 259 150 17 13 -0.999 -0.378 4.29 0.363 -1.79 -1.14 -1.51 4.71 4.97 4.27 259 160 17 13 -0.999 -0.379 4.31 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.27 259 170 17 13 -0.999 -0.378 4.32 0.362 -1.83 -1.18 -1.55 4.77 5.02 4.27 259 180 17 13 -0.999 -0.377 4.37 0.363 -1.84 -1.20 -1.57 4.80 5.05 4.25 259 190 17 13 -0.999 -0.377 4.37 0.363													4.76
259 140 17 13 -0.999 -0.390 4.25 0.360 -1.75 -1.11 -1.47 4.68 4.93 4.25 259 150 17 13 -0.999 -0.378 4.29 0.363 -1.79 -1.14 -1.51 4.71 4.97 4.25 259 160 17 13 -0.999 -0.379 4.31 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.25 259 170 17 13 -0.999 -0.378 4.32 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.25 259 180 17 13 -0.999 -0.376 4.35 0.363 -1.81 -1.18 -1.55 4.77 5.02 4.25 259 190 17 13 -0.999 -0.377 4.37 0.363 -1.86 -1.22 -1.59 4.82 5.07 4.25 259 200 17 13 -0.999 -0.375 4.40 0.364													4.80
259 150 17 13 -0.999 -0.378 4.29 0.363 -1.79 -1.14 -1.51 4.71 4.97 4.259 1.60 17 13 -0.999 -0.379 4.31 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.20 4.31 0.363 -1.81 -1.16 -1.53 4.74 4.99 4.20 4.25 1.81 -1.16 -1.53 4.74 4.99 4.20 4.25 1.81 -1.81 -1.55 4.77 5.02 4.25 4.25 1.22 -1.83 -1.18 -1.55 4.77 5.02 4.25 4.25 1.22 -1.83 -1.18 -1.55 4.77 5.02 4.25 4.25 1.28 -1.20 -1.57 4.80 5.05 4.25 4.25 1.28 -1.20 -1.57 4.80 5.05 4.25 4.25 1.20 -1.57 4.80 5.05 4.25 4.25 1.20 -1.57 4.80 5.05 4.25 4.25 1.20 </td <td></td> <td></td> <td></td> <td>-0.999</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>4.83</td>				-0.999									4.83
259 170 17 13 -0.998 -0.383 4.32 0.362 -1.83 -1.18 -1.55 4.77 5.02 4. 259 180 17 13 -0.999 -0.376 4.35 0.363 -1.84 -1.20 -1.57 4.80 5.05 4. 259 190 17 13 -0.999 -0.377 4.37 0.363 -1.86 -1.22 -1.59 4.82 5.07 4. 259 210 17 13 -1.000 -0.375 4.40 0.364 -1.89 -1.25 -1.62 4.87 5.11 5. 260 60 14 7 -0.999 -0.563 4.00 0.320 -0.85 -0.59 -0.72 4.33 4.48 4. 260 70 14 9 -0.999 -0.553 4.03 0.320 -0.96 -0.59 -0.72 4.33 4.48 4. 260 80 15 9 -0.997 -0.555 4.04<				-0.999	-0.378			-1.79	-1.14	-1.51	4.71		4.86
259 180 17 13 -0.999 -0.376 4.35 0.363 -1.84 -1.20 -1.57 4.80 5.05 4. 259 190 17 13 -0.999 -0.377 4.37 0.363 -1.86 -1.22 -1.59 4.82 5.07 4. 259 200 17 13 -0.999 -0.377 4.39 0.363 -1.87 -1.24 -1.60 4.85 5.10 4. 259 210 17 13 -1.000 -0.375 4.40 0.364 -1.89 -1.25 -1.62 4.87 5.11 5. 260 60 14 7 -0.999 -0.563 4.00 0.320 -0.85 -0.59 -0.72 4.33 4.48 4. 260 70 14 9 -0.999 -0.563 4.04 0.320 -0.96 -0.59 -0.79 4.33 4.48 4. 260 80 15 9 -0.997 -0.555 4.04 0.322 -1.08													4.89
259 190 17 13 -0.999 -0.377 4.37 0.363 -1.86 -1.22 -1.59 4.82 5.07 4. 259 200 17 13 -0.999 -0.377 4.39 0.363 -1.87 -1.24 -1.60 4.85 5.10 4. 259 210 17 13 -1.000 -0.375 4.40 0.364 -1.89 -1.25 -1.62 4.87 5.11 5. 260 60 14 7 -0.999 -0.563 4.00 0.320 -0.85 -0.59 -0.72 4.33 4.48 4. 260 70 14 9 -0.999 -0.563 4.03 0.320 -0.85 -0.59 -0.72 4.33 4.48 4. 260 80 15 9 -0.997 -0.555 4.04 0.322 -1.08 -0.73 -0.91 4.44 4.64 4. 260 90 15 9 -0.998 -0.513 4.11 0.331 -1.18													4.92
259 200 17 13 -0.999 -0.377 4.39 0.363 -1.87 -1.24 -1.60 4.85 5.10 4. 259 210 17 13 -1.000 -0.375 4.40 0.364 -1.89 -1.25 -1.62 4.87 5.11 5. 260 60 14 7 -0.999 -0.563 4.00 0.320 -0.85 -0.59 -0.72 4.33 4.48 4. 260 70 14 9 -0.999 -0.563 4.03 0.320 -0.96 -0.59 -0.79 4.34 4.57 260 80 15 9 -0.997 -0.555 4.04 0.322 -1.08 -0.73 -0.91 4.44 4.64 4. 260 90 15 9 -0.998 -0.513 4.11 0.331 -1.18 -0.79 -0.99 4.51 4.71 4. 260 100 15 9 -0.998 -0.480 4.15 0.338 -1.26 -0.88 -1.08 4.57 4.75 4. 260 110 15 9 -0.995 -0.455 4.20 0.344 -1.34 -0.95 -1.15 4.63 4.80 4.					-0.376								4.94
259 210 17 13 -1.000 -0.375 4.40 0.364 -1.89 -1.25 -1.62 4.87 5.11 5. 260 60 14 7 -0.999 -0.563 4.00 0.320 -0.85 -0.59 -0.72 4.33 4.48 4. 260 70 14 9 -0.999 -0.563 4.03 0.320 -0.96 -0.59 -0.79 4.36 4.57 4. 260 80 15 9 -0.997 -0.555 4.04 0.322 -1.08 -0.73 -0.91 4.44 4.64 4. 260 90 15 9 -0.998 -0.513 4.11 0.331 -1.18 -0.79 -0.99 4.51 4.71 4. 260 100 15 9 -0.998 -0.480 4.15 0.338 -1.26 -0.88 -1.08 4.57 4.75 4. 260 110 15 9 -0.995 -0.455 4.20 0.344 -1.34 -0.95 -1.15 4.63 4.80 4.					-0.3//								4.97 4.99
260 60 14 7 -0.999 -0.563 4.00 0.320 -0.85 -0.59 -0.72 4.33 4.48 4. 260 70 14 9 -0.999 -0.563 4.03 0.320 -0.96 -0.59 -0.79 4.36 4.57 4. 260 80 15 9 -0.997 -0.555 4.04 0.322 -1.08 -0.73 -0.91 4.44 4.64 4. 260 90 15 9 -0.998 -0.513 4.11 0.331 -1.18 -0.79 -0.99 4.51 4.71 4. 260 100 15 9 -0.998 -0.480 4.15 0.338 -1.26 -0.88 -1.08 4.57 4.75 4. 260 110 15 9 -0.995 -0.455 4.20 0.344 -1.34 -0.95 -1.15 4.63 4.80 4.													5.01
260 70 14 9 -0.999 -0.563 4.03 0.320 -0.96 -0.59 -0.79 4.36 4.57 4. 260 80 15 9 -0.997 -0.555 4.04 0.322 -1.08 -0.73 -0.91 4.44 4.64 4. 260 90 15 9 -0.998 -0.513 4.11 0.331 -1.18 -0.79 -0.99 4.51 4.71 4. 260 100 15 9 -0.998 -0.480 4.15 0.338 -1.26 -0.88 -1.08 4.57 4.75 4. 260 110 15 9 -0.995 -0.455 4.20 0.344 -1.34 -0.95 -1.15 4.63 4.80 4.													4.41
260 80 15 9 -0.997 -0.555 4.04 0.322 -1.08 -0.73 -0.91 4.44 4.64 4. 260 90 15 9 -0.998 -0.513 4.11 0.331 -1.18 -0.79 -0.99 4.51 4.71 4. 260 100 15 9 -0.998 -0.480 4.15 0.338 -1.26 -0.88 -1.08 4.57 4.75 4. 260 110 15 9 -0.995 -0.455 4.20 0.344 -1.34 -0.95 -1.15 4.63 4.80 4.													4.47
260 90 15 9 -0.998 -0.513 4.11 0.331 -1.18 -0.79 -0.99 4.51 4.71 4. 260 100 15 9 -0.998 -0.480 4.15 0.338 -1.26 -0.88 -1.08 4.57 4.75 4. 260 110 15 9 -0.995 -0.455 4.20 0.344 -1.34 -0.95 -1.15 4.63 4.80 4.													4.55
260 110 15 9 -0.995 -0.455 4.20 0.344 -1.34 -0.95 -1.15 4.63 4.80 4.		90			-0.513	4.11	0.331	-1.18			4.51	4.71	4.62
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1											4.57		4.67
260 120 15 9 -0.99/ -0.411 4.26 0.354 -1.41 -1.03 -1.23 4.68 4.84 4.													4.73
	260	120	15 9	-0.997	-0.411	4.26	0.354	-1.41	-1.03	-1.23	4.68	4.84	4.77

Table B.2. (continued)

* .1		No. Poin	of ts		PCA T	hinnin	g Line ^e		Log N ^f			Log B ^f	
Id. Code ^a	Cond.b	nΤ	n	rd	β̂	â	p _{est}	Min.	Max.	Mean	Min.	Max.	Mean
260	130	15	9	-0.998	-0.402	4.29	0.357	-1.46	-1.07	-1.27	4.72	4.88	4.80
260	140	15	9	-0.999	-0.382	4.33	0.362	-1.51	-1.13	-1.33	4.76	4.90	4.84
260	150	15	9	-0.999	-0.377	4.35	0.363	-1.55	-1.15	-1.36	4.78	4.93	4.86
260	160	15	9	-0.999	-0.370	4.37	0.365	-1.57	-1.18	-1.39	4.81	4.96	4.89
260	170	15	9	-0.994	-0.383	4.37	0.362	-1.59	-1.25	-1.42	4.83	4.98	4.91
260	180	15	9	-1.000	-0.358	4.42	0.368	-1.62	-1.23	-1.43	4.85	5.00	4.93
261	60	14	8	-0.999	-0.612	3.93	0.310	-0.85	-0.53	-0.70	4.26	4.46	4.36
261	70	14	8	-0.999	-0.597	3.97	0.313	-0.96	-0.65	-0.82	4.35	4.55	4.45
261	80	15	9	-0.997	-0.608	3.97	0.311	-1.08	-0.73	-0.91	4.40	4.62	4.52
261	90	15	9	-0.999	-0.543	4.05	0.324	-1.18	-0.79	-0.99	4.47	4.68	4.59
261	100		9	-0.997	-0.528	4.08	0.327	-1.26	-0.88	-1.08	4.53	4.74	4.65
261 261	110 120	15 15	9 9	-0.997	-0.499	4.12	0.333	-1.34	-0.95	-1.15	4.59	4.78	4.70 4.74
261	130	15		-0.997	-0.476	4.16	0.339	-1.41	-1.03	-1.23	4.64	4.83	4.74
261	140	15	9 9	-0.997 -0.995	-0.440 -0.423	4.22	0.347 0.351	-1.46 -1.51	-1.07 -1.13	-1.27 -1.33	4.68 4.72	4.86 4.89	4.82
261	150	15	9	-0.995	-0.423	4.25 4.28	0.351	-1.55	-1.13	-1.35	4.75	4.92	4.84
261	160	15	9	-0.999	-0.410	4.30	0.355	-1.57	-1.13	-1.39	4.78	4.94	4.87
262	70		13	-0.998	-0.584	4.02	0.316	-0.85	0.34	-0.43	3.80	4.49	4.27
262	80	13 1	-	-0.998	-0.580	4.01	0.316	-0.98	0.23	-0.56	3.86	4.56	4.34
262	90	13 1		-0.997	-0.595	3.99	0.314	-1.09	0.07	-0.68	3.92	4.62	4.39
262	100	13 1		-0.997	-0.595	3.99	0.313	-1.21	-0.04	-0.79	3.98	4.68	4.46
262	110	13 1		-0.998	-0.598	3.98	0.313	-1.31	-0.14	-0.89	4.04	4.74	4.52
262	120	13 1	3	-0.998	-0.598	3.98	0.313	-1.38	-0.23	-0.98	4.10	4.80	4.57
262	130	13 1	13	-0.998	-0.600	3.99	0.313	-1.46	-0.32	-1.07	4.15	4.85	4.63
262	140		3	-0.998	-0.599	3.99	0.313	-1.55	-0.39	-1.15	4.20	4.90	4.67
262	150		13	-0.9 9 8	-0.587	4.00	0.315	-1.63	-0.47	-1.23	4.25	4.94	4.72
263	26		4	-0.968	-0.848	3.72	0.271	-0.40	-0.31	-0.36	3.98	4.05	4.02
263	33	11	7	-0.998	-0.850	3.72	0.270	-0.60	-0.32	-0.48	3.98	4.22	4.12
263	39	11	9	-0.996	-0.776	3.72	0.281	-0.77	-0.32	-0.59	3.96	4.30	4.18
264	40	9	4	-0.936	-0.243	3.51	0.402	-0.94	-0.56	-0.72	3.63	3.73	3.68
264	50	11	7	-0.938	-0.551	3.41	0.322	-0.94	-0.60	-0.76	3.71	3.90	3.83
265	77	6	4	-0.979	-0.250	4.06	0.400	-1.38	-0.90	-1.15	4.28	4.39	4.35
266	60		8	-0.995	-0.256	4.16	0.398	-1.31	-0.96	-1.16	4.40	4.49	4.45
266	80		1]	-0.989	-0.336	4.14	0.374	-1.55	-0.93	-1.29	4.44	4.65	4.58
266 266	100 120	14 1 14 1	1 12	-0.993	-0.334	4.22	0.375	-1.73	-1.13	-1.48	4.58	4.79	4.71 4.78
266	140		8	-0.993 -0.964	-0.331 -0.566	4.26	0.376 0.319	-1.87	-1.16	-1.58 -1.37	4.62 4.44	4.87 4.84	4.78
267	42		8	-0.984	-0.719	3.93 3.62	0.319	-1.68 -0.96	-1.01 -0.63	-1.37 -0.79	4.44	4.84	4.71
268	54		7	-0.996	-0.719	3.71	0.362	-0.96	-0.65	-0.79	3.95	4.10	4.04
269	75		6	-0.994	-0.654	3.43	0.302	-1.57	-0.42	-1.09	3.68	4.42	4.15
270	60	14 1	-	-0.994	-0.597	3.68	0.302	-1.17	-0.42	-0.85	3.79	4.34	4.19
270	70		4	-0.994	-0.626	3.64	0.308	-1.30	-0.34	-0.98	3.83	4.43	4.26
270	80		4	-0.994	-0.627	3.65	0.307	-1.40	-0.45	-1.08	3.91	4.50	4.33
270	90	14 1		-0.991	-0.625	3.67	0.308	-1.48	-0.52	-1.16	3.97	4.56	4.40

Table B.2. (continued)

Id.		No. of Points		PCA T	hinnin	g Line ^e		Log N ^f		~~~	Log B ^f	
Codea	Cond.b	n _T n	rd	β̂	â	^p est	Min.	Max.	Mean	Min.	Max.	Mean
270 270	100 110	14 14 14 14	-0.990	-0.641	3.68	0.305	-1.55	-0.59	-1.22	4.02	4.63	4.46
270	120	14 14	-0.991 -0.991	-0.656 -0.646	3.69 3.73	0.302 0.304	-1.59 -1.63	-0.64 -0.68	-1.26 -1.31	4.08 4.13	4.73 4.74	4.52 4.58
271	40	18 18	-0.995	-0.888	2.97	0.265	-1.14	-0.28	-0.85	3.17	3.94	3.72
271	50	18 18	-0.999	-0.804	3.26	0.277	-1.17	-0.31	-0.89	3.50	4.19	3.97
271	60	18 18	-0.999	-0.813	3.41	0.276	-1.21	-0.35	-0.93	3.69	4.37	4.16
271	70	18 18	-0.999	-0.829	3.49	0.273	-1.25	-0.40	-0.97	3.81	4.51	4.29
271 271	80	18 18	-0.999	-0.854	3.52	0.270	-1.31	-0.45	-1.02	3.87	4.61	4.39
271	90 100	18 18 18 18	-0.998 -0.998	-0.882	3.51 3.50	0.266 0.264	-1.36	-0.51	-1.08	3.93	4.69	4.46
271	110	18 18	-0.996	-0.897 -0.881	3.50	0.264	-1.42 -1.49	-0.57 -0.62	-1.14 -1.20	3.97 4.02	4.75 4.78	4.52 4.56
272	50	8 6	-0.998	-1.656	2.12	0.188	-1.25	-1.08	-1.18	3.90	4.18	4.07
272	60	9 7	-0.956	-1.855	1.89	0.175	-1.36	-1.14	-1.25	3.94	4.35	4.21
272	71	9 7	-0.984	- 2.373	1.18	0.148	-1.41	-1.25	-1.34	4.11	4.50	4.37
272	81	10 8	-0.997	-2.439	1.08	0.145	-1.44	-1.26	-1.38	4.15	4.61	4.45
272 273	103 40	10 10 18 15	-0.999	-1.848	1.93	0.176	-1.49	-1.06	-1.37	3.88	4.69	4.46
273	40 50	18 15	-0.995 -0.988	-0.404 -0.437	3.85 3.90	0.356 0.348	-0.97	-0.06	-0.67	3.87	4.22	4.12 4.26
273	60	18 15	-0.990	-0.437	3.94	0.346	-1.11 -1.24	-0.34 -0.47	-0.82 -0.95	4.02 4.13	4.36 4.47	4.20
273	70	18 15	-0.993	-0.454	3.98	0.344	-1.34	-0.58	-1.06	4.22	4.57	4.46
273	80	18 15	-0.993	-0.414	4.05	0.354	-1.45	-0.70	-1.18	4.36	4.64	4.54
273	90	18 15	-0.994	-0.481	3.96	0.338	-1.59	-0.84	-1.31	4.34	4.70	4.59
273	100	18 15	-0.988	-0.471	3.95	0.340	-1.76	-1.00	-1.49	4.39	4.76	4.65
274 274	60 70	10 8 10 8	-0.999 -0.997	-0.378	3.96	0.363	-0.94	-0.42	-0.74	4.12	4.32	4.24
274	80	10 8	-0.997 -0.998	-0.399 -0.403	3.98 3.99	0.357 0.356	-1.08 -1.21	-0.56 -0.69	-0.89 -1.01	4.20 4.27	4.41 4.48	4.33 4.40
274	90	10 9	-0.998	-0.422	3.97	0.352	-1.32	-0.69	-1.08	4.26	4.54	4.43
274	100	10 9	-0.998	-0.440	3.95	0.347	-1.43	-0.81	-1.19	4.30	4.58	4.47
276	84	99	-0.998	-1.222	3.35	0.225	-1.06	-0.60	-0.88	4.09	4.62	4.43
276	95	9 9	-0.997	-1.220	3.42	0.225	-1.07	-0.62	-0.89	4.17	4.70	4.51
276 277	112	9 9	-0.998	-1.224	3.44	0.225	-1.10	-0.65	-0.93	4.24	4.77	4.58
277	70 80	5 4 9 7	-1.000 -0.996	-0.333	3.74	0.375	-1.05	-0.36	-0.75	3.86	4.09	3.98
277	90	10 9	-0.995	-0.362 -0.396	3.79 3.83	0.367 0.358	-1.40 -1.50	-0.52 -0.27	-1.09 -1.12	3.99 3.94	4.31 4.45	4.19 4.27
277	100	10 9	-0.993	-0.414	3.88	0.354	-1.54	-0.40	-1.20	4.06	4.54	4.38
277	110	10 9	-0.990	-0.418	3.95	0.353	-1.55	-0.46	-1.24	4.16	4.63	4.46
277	120	8 7	-0.994	-0.395	4.03	0.358	-1.51	-0.50	-1.18	4.24	4.65	4.50
278	1	33 19	-1.000	-0.373	4.31	0.364	-1.92	-1.29	-1.60	4.79	5.02	4.90
279 280	40 51	11 10 8 6	-0.993	-0.640	3.72	0.305	-1.53	-0.52	-1.14	4.02	4.67	4.45
281	27	8 6 5 5	-0.988 -0.992	-0.552 -0.691	3.83 4.09	0.322 0.296	-1.71 -0.89	-1.41 -0.17	-1.57 -0.57	4.60 4.18	4.76	4.70
281	30	5 5	-0.990	-0.621	4.11	0.298	-0.89	-0.17	-0.57	4.18	4.67 4.70	4.48 4.53
281	33	5 5	-0.979	-0.547	4.14	0.323	-1.11	-0.41	-0.79	4.34	4.71	4.58

Table B.2. (continued)

* 1		No. Poi	of nts		PCA T	hinnin	ıg Line ^e	A	Log N ^f			Log B	
Id. Code ^a	Cond.b	nŢ	n	r ^d	â	â	p _{est}	Min.	Max.	Mean	Min.	Max.	Mean
281 281 281	27 30 33	5 5 5	5	-0.992 -0.990 -0.979	-0.691 -0.621 -0.547	4.11	0.296 0.308 0.323	-1.00	-0.17 -0.29 -0.41	-0.68	4.27	4.67 4.70 4.71	4.53

 $aTable\ B.1$ associates each Id. code with a particular yield table.

 $^{^{\}mathrm{b}}\mathsf{See}$ Table B.1 and the references given there for more information on condition.

 $^{^{\}text{C}}\text{n}_{\text{T}}$ is the total number of log B-log N points given for each code and condition. n is the number of points remaining after removing points not relevant to the thinning line. This is the number of points used to fit the PCA relationship between log B and log N.

dThese correlation coefficients have little statistical meaning because the natural variability present in real forest measurements has been removed from yield table predictions. These values are included only as a crude index of variation around the fitted line.

 $e\hat{\beta}$ and $\hat{\alpha}$ are, respectively, the slope and intercept of the fitted PCA thinning line. Formulas for principle components analysis are given in Jolicoeur and Heusner (1971). p_{est} is a transformation of the thinning slope calculated by $p_{est} = 0.5$ / (1 - $\hat{\beta}$). See discussion in Chapter 6.

 $^{^{\}mbox{\scriptsize f}}$ The mean, minimum, and maximum are given for log B and for log N over the n data points used to fit each thinning line.

Table B.3. Fitted Allometric Relationships for Forestry Yield Table Data.

				Al	lomet	ric Rel	ationsh	ips F	it by P	rincipa	1 Com	ponent	Analysi	s ^C		
		_1	og ñ -	log ₩	_10	g DBH -	log w	log	BSLA -	log w	_10	g h - 1	og DBH	log	n - 10	g BSLA
Id. Code ^a	Cond.b	n	r	Ŷ _{h₩}	n	r	$\hat{\Phi}_{DW}$	n	r	$\hat{\Phi}_{BW}$	n	r	Ŷ _{hD}	n	r	ŶhΒ
201	40	13	0.988	0.159	13	1.000	0.381	13	1.000	0.764	13	0.985	0.417	13	0.986	0.207
201	50	14	0.994	0.188	14	0.998	0.381	14	1.000	0.763	14	0.993	0.492	14	0.993	0.246
201	60	14	0.997	0.215	14	1.000	0.381	14	1.000	0.755	14	0.998	0.564	14	0.997	0.284
201	70	14	1.000	0.234	14	1.000	0.377	14	1.000	0.754	14	1,000	0.620	14	0.999	0.310
201	80	14	1.000	0.250	14	1.000	0.379	14	1.000	0.757	14	1.000	0.660	14	1.000	0.331
201 201	90 100	14 14	1.000	0.257	14	1.000	0.379	14	1.000	0.759	14	1.000	0.679	14	1.000	0.339
201	110	14	0.999	0.271 0.278	14	1.000	0.381	14	1.000	0.761 0.765	14	0.999	0.711	14 14	0.999	0.363
201	120	14	1.000	0.278	14 14	1.000	0.384	14 14	1.000	0.759	14 14	0.999	0.724	14	0.999	0.363
201	130	14	1.000	0.281	14	0.999	0.380	14	0.999	0.758	14	0.999	0.736	14	0.999	0.369
201	140	4	0.999	0.273	4	1.000	0.333	4	1.000	0.668	4	1.000	0.821	4	1.000	0.409
201	150	4	0.999	0.275	4	1.000	0.339	4	1.000	0.676	4	1.000	0.810	4	1.000	0.406
201	160	4	1.000	0.271	4	1.000	0.326	4	1.000	0.656	4	1.000	0.832	4	1.000	0.414
202	80	9	0.999	0.231	9	1.000	0.347	-	1.000	0.050	9	0.998	0.665	-1	1,000	••••
202	96	9	0.999	0.229	9	1.000	0.392				g	0.999	0.584			
202	112	9	1.000	0.211	9	1.000	0.393				9	0.999	0.537			
203	30	8	0.995	0.215	8	1.000	0.322	8	1.000	0.662	8	0.994	0.670	8	0.994	0.325
203	40	8	0.998	0.208	8	1.000	0.332	8	1.000	0.653	8	0.997	0.626	8	0.997	0.318
203	50	8	0.993	0.219	8	1.000	0.322	8	1.000	0.656	8	0.991	0.681	8	0.991	0.333
203	60	8	0.997	0.213	8	1.000	0.330	8	1.000	0.664	8	0.995	0.644	8	0.996	0.320
203	70	8	0.996	0.209	8	1.000	0.332	8	1.000	0.669	8	0.995	0.630	8	0.995	0.313
203	80	8	0.998	0.216	8	1.000	0.327	8	1.000	0.660	8	0.998	0.661	8	0.999	0.328
203	90	8	0.999	0.222	8	1.000	0.329	8	1.000	0.660	8	0.999	0.675	8	0.999	0.336
204 204	62	7	0.997	0.205	7	0.999	0.427	7	1.000	0.849	7	0.995	0.480	7	0.996	0.241
204	71 80	8 8	0.995	0.189 0.174	8	1.000	0.408 0.404	8	1.000	0.818 0.818	8	0.993 0.997	0.462 0.431	8	0.994 0.997	0.230
205	58	8	0.998	0.174	8 8	1.000	0.444	8 8	1.000	0.884	8 8	0.997	0.431	8 8	0.997	0.282
205	68	8	0.996	0.234	8	1.000	0.455	8	1.000	0.915	8	0.996	0.515	8	0.996	0.256
205	80	8	0.997	0.192	8	1.000	0.449	8	1.000	0.898	8	0.997	0.429	8	0.997	0.214
206	52	9	0.999	0.264	9	0.999	0.369	9	1.000	0.730	9	0.998	0.715	9	0.998	0.362
206	64	9	0.994	0.233	9	1.000	0.408	9	1,000	0.824	9	0.995	0.572	9	0.994	0.283
206	75	9	0.998	0.217	9	1.000	0.428	9	1.000	0.857	9	0.998	0.508	9	0.998	0.254
207	49	11	0.995	0.258	11	0.993	0.333				11	0.982	0.770			
208	40	6	0.999	0.301	6	1.000	0.375	6	1.000	0.748	6	0.999	0.804	6	0.999	0.402
208	50	6	0.999	0.302	6	1.000	0.372	6	1.000	0.745	6	0.999	0.812	6	0.999	0.405
208	60	6	1.000	0.302	6	1.000	0.376	6	1.000	0.745	6	1.000	0.802	6	0.999	0.405
208	70	6	0.999	0.294	6	1.000	0.372	6	1.000	0.754	6	0.999	0.790	6	0.999	0.390
209	1				4	0.997	0.166	4	0.999	0.837						
209	2	_	0.05-		6	0.988	0.202	6	0.999	0.862	_		0 470	_	0.000	0 103
209	3	7	0.965	0.118	7	0.992	0.251	7	0.999	0.889	7	0.988	0.479	7	0.953	0.131
209	4	00	0.000	0.166	8	0.998	0.433	8	0.999	0.844	0.0	0.000	0 400			
210	136	28	0.998	0.166	28	0.998	0.392	3.4	1 000	0.700	28	0.999	0.422	1.4	1 000	0.070
211	20	14	1.000	0.217	14	0.999	0.406	14	1.000	0.780	14	1.000	0.535	14	1.000	0.278

Table B.3. (continued)

Id. ode ^a	Cond.b			log w	10	g DBH -	log w	loa	BSLA -	loa w	10	g h - 1	og DBH	Pof	h - 10	g BSLA
		n	r	$\widehat{\Phi}_{hw}$	n	r	φ _{Dw}	n	r	φ̂ _{Bw}	n	r	φ̂ _{h0}	n	r	φ̂hB
11	30	14	0.999	0.220	14	0.999	0.399	14	1.000	0.790	14	0.999	0.553	14	1.000	0.279
11 11	40	14	1.000	0.217	14	1.000	0.402	14	1.000	0.800	14	1.000	0.538	14	1.000	0.277
11	50 60	14 14	1.000	0.219	14 14	1.000	0.397	14 14	1.000	0.792 0.792	14 14	1.000	0.552 0.545	14 14	1.000	0.276
11	70	14	1.000	0.217	14	1.000	0.400	14	1.000	0.793	14	1.000	0.544	14	1.000	0.27
12	60	14	0.999	0.278	14	1.000	0.397	14	1.000	0.790	14	0.999	0.701	14	0.999	0.35
12	80	15	1.000	0.282	15	0.999	0.387	15	1.000	0.777	15	0.999	0.728	15	0.999	0.36
12	100	15	1.000	0.279	15	1.000	0.391	15	1.000	0.781	15	0.999	0.714	15	0.999	0.35
12 12	120	15	1.000	0.282	15	1.000	0.390	15	1.000	0.778	15	0.999	0.722	15	0.999	0.36
12	140 160	15 15	1.000	0.286 0.283	15 15	1.000	0.386 0.389	15 15	1.000	0.773 0.778	15 15	0.999	0.742 0.729	15 15	0.999	0.36
12	180	15	1.000	0.292	15	1.000	0.384	15	1.000	0.768	15	0.999	0.760	15	0.999	0.38
12	200	15	1.000	0.284	15	1.000	0.387	15	1.000	0.774	15	0.999	0.735	15	0.999	0.36
13	30	6	0.997	0.260	6	0.999	0.404	6	1.000	0.795	6	0.993	0.644	6	0.997	0.32
13	40	6	0.994	0.291	6	1.000	0.396	6	1.000	0.799	6	0.994	0.738	6	0.994	0.36
13 13	50 60	6 6	0.996	0.288 0.278	6 6	0.999	0.394 0.400	6 6	1.000	0.796 0.800	6 6	0.996	0.731 0.697	6 6	0.996	0.36
13	70	6	1.000	0.271	6	1.000	0.396	6	1.000	0.797	6	1.000	0.685	6	1.000	0.34
14	1	Ü	1.000	0.271	·	1.000	0.550	V	1.000	0.757		1.000	0.000	v		0.0.
14	2															
14	3															
14 14	4															
14	5 6															
15	ì															
15	2															
15	3															
15	4															
15 15	5 6															
16	1															
16	2															
16	3															
17	54	6	0.997	0.242	6	1.000	0.372	6	1.000	0.745	6	0.995	0.651	6	0.995	0.32
17	64	6	0.998	0.266	6	1.000	0.391	6	1.000	0.781	6	0.996	0.682	6	0.996	0.34
17 18	75 50	6 10	0.999	0.280	6 10	1.000	0.408	6 10	0.999	0.814 0.673	6 10	0.998	0.687 1.072	6 10	0.997	0.53
18	60	10	1.000	0.350	10	1.000	0.330	10	1.000	0.662	10	1.000	1.060	10	1.000	0.5
8	70	10	1.000	0.346	10	1.000	0.339	10	1.000	0.669	10	0.999	1.022	10	0.999	0.5
20	40	6	0.999	0.353	6	0.999	0.292	6	0.999	0.579	6	0.999	1.212	6	1.000	0.6
20	50	6	0.998	0.368	6	0.999	0.286	6	1.000	0.575	6	0.998	1.288	6	0.998	0.6
20 20	60 70	6 6	0.997	0.366 0.348	6 6	1.000	0.297	6 6	1.000	0.584 0.611	6 6	0.997	1.237	6 6	0.996 0.992	0.6

Table B.3. (continued)

				A1	lomet	ric Rel	ationsh	ips F	it by P	rincipa	Com	ponent	Analysi	s ^C		
		_1	og ñ -	log ŵ	10	g DBH -	log w	log	BSLA -	log w	10	g h - 1	og DBH	log	ñ - 1o	g BSLA
Id. Code ^a	Cond.b	n	r	φ̂hw	n	r	φ _{Dw}	n	r	δ̂Βw	n	r	φ̂hD	n	r	δ̂hB
220	80	6	0.993	0.321	6	0.999	0.321	6	0.999	0.642	6	0.988	1.008	6	0.987	0.499
220	90	6	0.988	0.292	6	0.999	0.340	6	0.999	0.674	6	0.979	0.866	6	0.980	0.432
221	80	13	0.998	0.216	13	1.000	0.415	13	0.999	0.825	13	0.996	0.519	13	0.996	0.261
221	90	13	0.999	0.223	13	0.999	0.408	13	0.999	0.821	13	0.998	0.546	13	0.998	0.272
221	100	13	0.999	0.227	13	0.999	0.410	13	0.999	0.821	13	0.997	0.552	13	0.997	0.276
221	110	13	0.999	0.226	13	0.999	0.406	13	0.999	0.814	13	0.997	0.556	13	0.997	0.277
221	120	13	0.999	0.231	13	0.999	0.410	13	0.999	0.817	13	0.997	0.563	13	0.997	0.282
221	130	13	0.999	0.231	13	0.999	0.407	13	0.999	0.813	13	0.996	0.566	13	0.996 0.996	0.283
221 221	140 150	13 13	0.998	0.232	13	0.999	0.410	13	0.999	0.815	13	0.996 0.996	0.565 0.569	13 13	0.996	0.284
221	160	13	0.998	0.233	13 13	1.000	0.409	13 13	0.999	0.813	13 13	0.996	0.509	13	0.996	0.286
221	170	13	0.999	0.234	13	0.999	0.408	13	0.999	0.814	13	0.995	0.572	13	0.995	0.285
221	180	13	0.998	0.233	13	1.000	0.406	13	0.999	0.813	13	0.995	0.576	13	0.996	0.285
221	190	13	0.999	0.235	13	0.999	0.404	13	0.999	0.814	13	0.996	0.578	13	0.996	0.288
221	200	13	0.998	0.235	13	0.999	0.406	13	0.999	0.812	13	0.996	0.577	13	0.996	0.288
221	210	13	0.998	0.233	13	0.999	0.407	13	0.999	0.816	13	0.996	0.572	13	0.995	0.285
222	1	8	0.999	0.324	.8	0.999	0.415	8	1.000	0.843	. 8	0.999	0.780	8	1.000	0.384
222	ż	8	1.000	0.349	8	0.999	0.401	8	1.000	0.817	8	0.999	0.872	8	1.000	0.428
222	3	8	1.000	0.383	8	1.000	0.395	8	1.000	0.801	8	0.999	0.969	8	1.000	0.478
222	4	5	1.000	0.404	5	0.999	0.392	5	1.000	0.780	5	1.000	1.030	5	1.000	0.517
222	5	5	1.000	0.425	5	0.999	0.384	5	1.000	0.749	5	0.999	1.106	5	1.000	0.567
223	1	13	0.998	0.223	13	1.000	0.423				13	0.998	0.527			
224	3	14	0.997	0.193	14	0.993	0.311				14	0.994	0.618			
225	1	6	0.994	0.248	6	1.000	0.404				6	0.993	0.616			
227	40	13	0.999	0.306	13	0.999	0.337	13	0.999	0.662	13	0.998	0.909	13	0.997	0.462
227	50	13	1.000	0.312	13	1.000	0.340	13	0.999	0.663	13	0.999	0.917	13	0.998	0.470
227	60	13	0.999	0.309	13	1.000	0.339	13	0.999	0.663	13	0.999	0.911	13	0.998	0.465
227	70	13	0.999	0.308	13	1.000	0.337	13	0.999	0.663	13	0.999	0.914	13	0.998	0.465
227	80	13	0.999	0.310	13	1.000	0.336	13	0.999	0.663	13	0.999	0.922	13	0.998	0.468
227	90	13	0.999	0.310	13	1.000	0.341	13	0.999	0.662	13	0.999	0.908	13	0.998	0.467
227	100	13	0.999	0.309	13	1.000	0.338	13	0.999	0.663	13	0.999	0.916	13	0.998	0.466
227	110	13	0.999	0.310	13	1.000	0.338	13	0.999	0.663	13	0.999	0.916	13	0.998	0.467
227 228	120	13	0.994	0.312	13	0.995	0.341	13	0.997	0.662	13	0.999	0.915	13	0.984	0.468
228	25 35	11	0.997	0.347	11	1.000	0.399]]	1.000	0.808	11	0.996	0.869	11	0.997	0.429
228	35 45	12	0.999	0.325	11 12	1.000	0.410	11 12	1.000	0.806 0.809	11 12	0.998	0.791	12	1.000	0.403
228	45 55													12	0.999	0.372
228	55 65	12 12	1.000	0.306 0.289	12 12	1.000	0.412	12 12	1.000	0.823	12 12	0.999	0.743	12	0.999	0.372
228	75	12	0.999	0.289	12	0.999	0.421	12	1.000	0.851	12	0.999	0.680	12	0.999	0.343
229	60	7	1.000	0.303	7	1.000	0.428	7	1,000	0.710	7	1.000	0.863	7	1.000	0.427
229	70	7	1.000	0.303	7	1.000	0.353	7	1.000	0.710	7	1.000	0.849	7	1.000	0.424
229	80	7	1.000	0.300	7	1.000	0.333	7	1.000	0.707	7	1.000	0.861	7	1.000	0.424
229	90	7	1.000	0.306	7	1.000	0.356	7	1.000	0.710	7	1.000	0.861	7	1.000	0.431
		,	1.000	0.500	,	1.000	0.550	,	1.000	3.7.10	′	1.000	0.001	,	1.000	0.701

Table B.3. (continued)

				All	lomet	ric Rel	ationsh:	ips F	it by P	rincipa	l Com	ponent	Analysi	s ^c		
		1	og ĥ -		10	g DBH -	log w	log	BSLA -	log w	10	g ñ - 1	og DBH	log	<u> </u>	g BSLA
Id. Code ^a	Cond.b	n	r	$\widehat{\Psi}$ hw	n	r	$\hat{\Phi}_{Dw}$	n	r	$\boldsymbol{\hat{\Phi}}_{Bw}$	n	r	$\hat{\Phi}_{hD}$	n	r	φ̂hΒ
229	100	7	1.000	0.301	7	1.000	0.358	7	1.000	0.707	7	1.000	0.841	7	1.000	0.426
229	110	7	1.000	0.304	7	1.000	0.353	7	1.000	0.708	7	1.000	0.860	7	1.000	0.429
229	120	7	1.000	0.306	7	1.000	0.361	7	1.000	0.710	7	0.999	0.848	7	1.000	0.431
230	50	7	1.000	0.297	7	1.000	0.360	7	1.000	0.727	7	1.000	0.823	7	1.000	0.408
230	60	7	1.000	0.285	7	1.000	0.366	7	1.000	0.733	7	1.000	0.778	7	1.000	0.389
230	70	7	1.000	0.286	7	1.000	0.368	7	1.000	0.745	7	1.000	0.778	7	1.000	0.384
230	80 90	7	1.000	0.272	7	1.000	0.376	7	1.000	0.763	7	1.000	0.724	7	1.000	0.357
230 230	100	7 7	1.000	0.265	7	1.000	0.384	7	1.000	0.773	7	1.000	0.691	7 7	1.000	0.343
231	50	5	0.999	0.260 0.236	7 5	1.000	0.394	7 5	1.000	0.784 0.728	7 5	1.000	0.659 0.639	5	1.000	0.324
231	60	5	0.999	0.253	5	1.000	0.358	5	1.000	0.728	5	1.000	0.706	5	0.999	0.351
231	70	5	0.999	0.249	5	1.000	0.366	5	1.000	0.728	5	0.999	0.678	5	0.999	0.342
231	80	5	0.999	0.247	5	1.000	0.361	5	1.000	0.729	5	0.999	0.685	5	0.999	0.340
231	90	5	1.000	0.253	5	1.000	0.360	5	1.000	0.726	5	0.999	0.702	5	1.000	0.348
231	100	5	1.000	0.249	5	1.000	0.355	5	1.000	0.724	5	1.000	0.702	5	1.000	0.345
232	40	7	1.000	0.293	7	1.000	0.337	7	1.000	0.661	7	1.000	0.870	7	1.000	0.443
232	50	7	1.000	0.296	7	1.000	0.337	7	1.000	0.666	7	1.000	0.877	7	1.000	0.445
232	60	7	1.000	0.290	7	1.000	0.330	7	1.000	0.666	7	1.000	0.879	7	1.000	0.436
232	70	7	1.000	0.295	7	1.000	0.331	7	1.000	0.664	7	1.000	0.893	7	1.000	0.445
232	80	7	1.000	0.297	7	1.000	0.328	7	1.000	0.661	7	1.000	0.906	7	1.000	0.449
232	90	7	1.000	0.290	7	1.000	0.329	7	1.000	0.655	7	1.000	0.883	7	1.000	0.443
232	100	7	1.000	0.292	7	1.000	0.327	7	1.000	0.656	7	1.000	0.892	7	1.000	0.444
233	50	7	0.999	0.247	7	1.000	0.346	7	1.000	0.701	7	0.999	0.714	7	1.000	0.352
233	60	7	1.000	0.249	7	1.000	0.349	7	1.000	0.697	7	1.000	0.712	7	1.000	0.356
233	70	7	1.000	0.255	7	1.000	0.351	7	1.000	0.703	7	1.000	0.727	7	1.000	0.363
233	80	7	1.000	0.251	7	1.000	0.350	7	1.000	0.703	7	1.000	0.718	7	1.000	0.357
233	90	7	1.000	0.247	7	1.000	0.353	7	1.000	0.699	7	1.000	0.698	7	1.000	0.353
233	100	7	1.000	0.252	7	1.000	0.353	7	1.000	0.702	7	1.000	0.715	7	1.000	0.360
234 234	70 84	5 7	0.999	0.292	5	0.996	0.315	5	0.999	0.768	5	0.995	0.925	5	0.999	0.380
234	99	7	0.989	0.236 0.192	7 7	1.000	0.377	5	0.999	0.916	7 7	0.985	0.629 0.475	5 5	0.998	0.200
235	40	18	0.989	0.192	18	0.999		5 18	0.996	0.765 0.735	18	0.984	0.475	18	0.998	0.416
235	50	18	0.997	0.305	18	0.998	0.366 0.382	18	0.999	0.735	18	0.997	0.835	18	0.998	0.410
235	60	18	0.991	0.321	18	0.998	0.382	18	0.998	0.756	18	0.998	0.843	18	0.997	0.437
235	70	18	0.994	0.321	18	0.998	0.362	18	0.999	0.751	18	0.998	0.847	18	0.998	0.427
235	80	18	0.994	0.310	18	0.999	0.365	18	0.999	0.739	18	0.997	0.853	18	0.997	0.421
235	90	18	0.994	0.309	18	0.998	0.358	18	0.999	0.725	18	0.997	0.865	18	0.997	0.426
235	100	18	0.995	0.308	18	0.999	0.353	18	0.999	0.721	18	0.997	0.873	18	0.998	0.427
235	110	18	0.994	0.306	18	0.999	0.354	18	0.999	0.723	18	0.997	0.868	18	0.996	0.425
236	40	9	0.994	0.319	9	1.000	0.380	9	1.000	0.749	9	0.993	0.844	9	0.993	0.426
236	50	9	0.995	0.292	9	1.000	0.371	9	1.000	0.745	9	0.993	0.788	9	0.993	0.391
236	60	9	0.997	0.272	9	1.000	0.375	9	1.000	0.747	9	0.997	0.725	9	0.996	0.364
236	70	9	0.997	0.266	9	1.000	0.379	9	1.000	0.747	9	0.997	0.702	9	0.996	0.355

Table B.3. (continued)

237 1 12 0.999 0.290 12 1.000 0.373 12 1.000 0.752 12 0.999 0.778 12 1.000 0.386 237 2 8 1.000 0.3300 8 0.999 0.349 8 1.000 0.7766 R 0.999 0.859 8 1.000 0.475 237 3 8 1.000 0.332 8 0.999 0.672 8 1.000 0.968 8 1.000 0.532 237 4 8 1.000 0.348 5 0.999 0.295 5 0.999 0.579 5 0.999 1.179 5 0.999 0.602 238 40 5 0.998 0.258 5 1.000 0.780 5 0.999 0.462 5 0.999 0.363 238 60 5 1.000 0.266 5 1.000 0.393 5 1.000 0.782					A1	lomet	ric Rel	ationsh:	ips F	it by P	rincipa	l Com	ponent	Analysi	s ^C		
Code ^a Cond, ^b n r \$\hat{\theta_{hw}} n \$\text{O} 0.437 2 1 0 0.998 0.757 1 0.998 0.701 10 0.998 0.352 237 3 8 1.000 0.322 8 1.000 0.340 8 0.999 0.438 8 0.999 0.438 8 0.999 0.438 8 0.999 1.046 8 1.000 0.478 237 4 8 1.000 0.340 8 0.999 0.259 5 0.999 1.016 8 1.000 0.638 8 0.999 1.029 0.532 233	* 1		_1	og ħ -	log ₩	10	g DBH -	log w	log	BSLA -	log w	10	g ĥ - 1	og DBH	log	ñ − 1о	g BSLA
237 1 12 0.999 0.290 12 1.000 0.373 12 1.000 0.752 12 0.999 0.439 8 1.000 0.300 0.999 0.439 8 1.000 0.909 0.839 8 1.000 0.425 8 1.000 0.762 8 1.000 0.475 8 1.000 0.340 8 0.999 0.638 8 0.999 1.010 0.475 8 1.000 0.475 8 1.000 0.999 1.019 5 0.999 0.630 8 1.000 0.965 8 1.000 0.425 5 0.999 0.630 8 0.999 1.104 8 1.000 0.425 5 0.999 0.630 5 0.999 0.257 5 0.999 0.258 5 1.000 0.333 5 1.000 0.780 5 0.999 0.668 5 0.999 0.333 5 1.000 0.736 5 1.099 0.		Cond.b	n	r	$\hat{\phi}_{hw}$	n	r	∳ _{Dw}	n	r	$\boldsymbol{\hat{\Phi}}_{Bw}$	n	r	ŶhD	n	r	$\hat{\Phi}_{hB}$
237 2 8 1.000 0.300 8 0.999 0.349 8 1.000 0.736 8 1.000 0.322 8 1.000 0.323 8 1.000 0.323 8 1.000 0.340 8 0.099 0.255 8 0.999 0.632 8 1.000 0.340 8 0.0999 0.625 8 1.000 0.472 237 5 5 1.000 0.348 5 0.999 0.638 8 0.999 1.066 8 1.000 0.582 238 40 5 0.998 0.257 5 0.999 0.300 0.782 5 0.999 0.658 5 0.999 0.362 238 60 5 0.999 0.258 5 1.000 0.782 5 0.999 0.658 5 0.999 0.629 5 0.999 0.650 5 0.999 0.322 239 60 5 0.999	236	80	10	0.998	0.267	10	1.000	0.381	10	1,000	0.757	10	0.998	0.701	10	0.998	0.352
237 3 8 1.000 0.322 8 1.000 0.333 8 0.999 0.632 8 1.000 0.475 237 5 5 1.000 0.348 5 0.999 0.255 5 0.999 0.638 8 0.999 1.179 5 0.999 0.632 238 40 5 0.998 0.258 5 1.000 0.779 5 0.999 0.668 5 0.999 0.666 5 0.999 0.626 5 1.000 0.780 5 0.999 0.668 5 0.999 0.666 5 1.000 0.395 5 1.000 0.782 5 0.999 0.666 5 0.999 0.266 5 1.000 0.382 5 1.000 0.788 5 0.999 0.669 5 0.999 0.299 239 50 5 0.999 0.221 5 1.000 0.372 5 0.999 0.232 </td <td>237</td> <td>1</td> <td>12</td> <td>0.999</td> <td>0.290</td> <td>12</td> <td>1.000</td> <td></td> <td>12</td> <td>1.000</td> <td>0.752</td> <td>12</td> <td>0.999</td> <td>0.778</td> <td>12</td> <td>1.000</td> <td>0.386</td>	237	1	12	0.999	0.290	12	1.000		12	1.000	0.752	12	0.999	0.778	12	1.000	0.386
237 3 8 1.000 0.322 8 1.000 0.333 8 0.999 0.632 8 1.000 0.475 237 5 5 1.000 0.348 5 0.999 0.255 5 0.999 0.638 8 0.999 1.179 5 0.999 0.632 238 40 5 0.998 0.258 5 1.000 0.779 5 0.999 0.668 5 0.999 0.666 5 0.999 0.626 5 1.000 0.780 5 0.999 0.668 5 0.999 0.666 5 1.000 0.395 5 1.000 0.782 5 0.999 0.666 5 0.999 0.266 5 1.000 0.382 5 1.000 0.788 5 0.999 0.669 5 0.999 0.299 239 50 5 0.999 0.221 5 1.000 0.372 5 0.999 0.232 </td <td>237</td> <td>2</td> <td>8</td> <td>1.000</td> <td></td> <td></td> <td>0.999</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>1.000</td> <td>0.425</td>	237	2	8	1.000			0.999									1.000	0.425
237 4 8 1,000 0.340 8 0,999 0.225 8 0,999 0.638 8 0,999 1,046 8 1,000 0.532 238 40 5 0.998 0.257 5 0,999 0.259 5 0,999 0.579 5 0,999 0.629 5 0,999 0.629 5 0,999 0.620 5 0,999 0.628 5 0,999 0.268 5 1,000 0.780 5 0,999 0.668 5 0,999 0.342 238 60 5 0,999 0.228 5 1,000 0.362 5 1,000 0.782 5 1,000 0.666 5 0,999 0.324 239 60 5 0,999 0,221 5 1,000 0.368 5 1,000 0.737 5 0,999 0,600 5 0,999 0,229 239 70 5 0,998 0,231	237					8											0.479
237 5 \$ 1,000 0.348 \$ 0.999 0.295 \$ 0.999 0.570 \$ 0.999 0.1779 \$ 0.999 0.1779 \$ 0.999 0.602 238 40 5 0.998 0.257 \$ 0.999 0.339 \$ 1.000 0.779 \$ 0.999 0.658 \$ 0.999 0.331 238 60 \$ 1.000 0.266 \$ 1.000 0.393 \$ 1.000 0.782 \$ 1.000 0.672 \$ 0.999 0.331 238 70 \$ 0.997 0.258 \$ 1.000 0.395 \$ 1.000 0.782 \$ 1.000 0.666 \$ 0.999 0.343 239 50 \$ 0.999 0.221 \$ 1.000 0.362 \$ 1.000 0.736 \$ 5 0.990 0.600 \$ 0.999 0.292 239 50 \$ 0.999 0.221 \$ 1.000 0.342 \$ 0.999 0.600 \$ 0.999 0.327 239 60 \$ 0.998 0.231 \$ 1.000 0.372 \$ 1.000 0.733 \$																	0.532
238 40 5 0.998 0.257 5 0.998 0.329 5 1.000 0.779 5 0.999 0.642 5 0.998 0.328 238 50 5 0.998 0.258 5 1.000 0.395 5 1.000 0.780 5 1.000 0.668 5 0.997 0.234 238 70 5 0.997 0.258 5 1.000 0.393 5 1.000 0.781 5 0.996 0.656 5 0.999 0.234 239 40 5 0.999 0.221 5 1.000 0.362 5 1.000 0.738 5 0.999 0.600 5 0.999 0.295 239 50 5 0.999 0.221 5 1.000 0.378 5 0.999 0.600 5 0.999 0.232 241 20 5 0.986 0.305 5 0.984 0.221																	
238 50 5 0.998 0.258 5 1.000 0.393 5 1.000 0.780 5 0.999 0.688 5 0.997 0.331 238 60 5 1.000 0.268 5 1.000 0.393 5 1.000 0.782 5 1.000 0.672 5 0.999 0.343 238 70 5 0.999 0.220 5 1.000 0.362 5 1.000 0.736 5 0.999 0.600 5 0.999 0.201 5 0.000 0.736 5 0.999 0.600 5 0.999 0.600 5 0.999 0.305 5 0.999 0.374 5 1.000 0.738 5 0.999 0.600 5 0.999 0.331 5 0.999 0.334 5 1.000 0.738 5 0.999 0.318 0.909 0.318 3 0.999 0.333 5 0.999 0.328																	
238 60 5 1,000 0,266 5 1,000 0,782 5 1,000 0,782 5 1,000 0,782 5 1,000 0,639 5 0,990 0,666 5 0,999 0,333 239 40 5 0,999 0,220 5 1,000 0,368 5 1,000 0,736 5 0,999 0,609 5 0,999 0,221 239 50 5 0,999 0,221 5 1,000 0,368 5 1,000 0,738 5 0,999 0,600 5 0,999 0,211 239 70 5 0,996 0,221 5 0,909 0,203 5 0,994 0,221 5 0,999 0,648 5 0,999 0,212 241 20 5 0,986 0,331 5 0,994 0,221 5 0,994 0,333 5 0,999 0,318 5 0,999 0,318																	
238 70 5 0.997 0.258 5 1.000 0.393 5 1.000 0.781 5 0.996 0.666 5 0.991 0.292 239 40 5 0.993 0.220 5 1.000 0.368 5 1.000 0.736 5 0.999 0.600 5 0.991 0.292 239 50 5 0.998 0.235 5 0.999 0.374 5 1.000 0.738 5 0.999 0.662 5 0.999 0.327 241 20 5 0.986 0.221 5 1.000 0.737 5 0.995 0.662 5 0.994 0.327 241 20 5 0.986 0.621 5 0.992 0.330 5 0.994 0.210 5 0.993 1.582 5 0.998 1.002 241 40 5 0.997 0.327 5 0.997 0.210																	
239 40 5 0.993 0.220 5 1.000 0.362 5 1.000 0.736 5 0.990 0.609 5 0.991 0.292 239 50 5 0.999 0.221 5 1.000 0.368 5 1.000 0.738 5 0.999 0.600 5 0.999 0.299 239 70 5 0.996 0.221 5 0.990 0.372 5 1.000 0.737 5 0.995 0.648 5 0.994 0.322 241 20 5 0.996 0.241 5 1.000 0.322 5 1.000 0.737 5 0.995 0.648 5 0.994 0.322 241 20 5 0.998 0.205 5 0.994 0.210 5 0.994 0.344 5 0.993 1.573 5 0.996 0.961 241 50 0.993 0.331 5																	
239 50 5 0.999 0.221 5 1.000 0.368 5 1.000 0.739 5 0.999 0.299 0.299 239 60 5 0.998 0.235 5 0.999 0.374 5 1.000 0.737 5 0.995 0.648 5 0.994 0.322 241 20 5 0.986 0.305 5 0.984 0.221 5 0.992 0.330 5 0.996 0.648 5 0.994 0.222 241 30 5 0.999 0.333 5 0.994 0.210 5 0.998 0.332 5 0.996 1.582 5 0.990 0.922 241 40 5 0.993 0.333 5 0.997 0.210 5 0.999 0.350 5 0.999 1.560 5 0.999 0.932 241 50 5 0.999 0.210 5 0.999						_									-		
239 60 5 0.998 0.235 5 0.999 0.374 5 1.000 0.738 5 0.996 0.29 5 0.997 0.318 239 70 5 0.996 0.241 5 1.000 0.372 5 1.000 0.738 5 0.995 0.648 5 0.994 0.305 5 0.992 0.030 5 0.995 0.648 5 0.994 0.210 5 0.998 0.332 5 0.999 0.333 5 0.992 0.210 5 0.994 0.344 5 0.993 1.582 5 0.998 0.932 241 50 5 0.999 0.331 5 0.992 0.210 5 0.994 0.344 5 0.993 1.560 5 0.998 0.932 241 60 5 0.998 0.358 5 0.999 0.203 5 0.991 0.337 5 0.994 1.762<																	
239 70 5 0.996 0.241 5 1.000 0.372 5 1.000 0.737 5 0.995 0.648 5 0.994 0.327 241 20 5 0.986 0.305 5 0.984 0.221 5 0.992 0.333 5 0.998 0.322 241 40 5 0.993 0.331 5 0.992 0.210 5 0.994 0.344 5 0.993 1.573 5 0.996 0.991 241 50 5 0.997 0.327 5 0.997 0.210 5 0.994 0.344 5 0.993 1.573 5 0.996 0.991 241 50 5 0.997 0.325 5 0.999 0.358 5 0.994 0.203 5 0.999 0.358 5 0.994 0.203 5 0.991 0.337 5 0.989 0.363 3 1.056															-		
241 20 5 0.986 0.305 5 0.984 0.221 5 0.992 0.330 5 0.999 0.333 5 0.994 0.210 5 0.998 0.332 5 0.998 1.002 2 0.993 1.582 5 0.998 1.002 2 0.994 0.340 5 0.9993 1.582 5 0.998 1.002 0.904 0.203 5 0.994 0.343 5 0.997 0.210 5 0.997 0.350 5 0.999 1.560 5 0.989 0.934 241 60 5 0.998 0.358 5 0.994 0.203 5 0.991 0.337 5 0.999 1.560 5 0.989 0.932 242 56 3 1.000 0.285 3 0.998 0.545 4 0.996 1.093 4 0.987 0.553 4 0.998 0.272 243 2 5																	
241 30 5 0.999 0.333 5 0.994 0.210 5 0.998 0.332 5 0.993 1.582 5 0.998 1.002 241 40 5 0.993 0.331 5 0.997 0.210 5 0.994 0.344 5 0.993 1.573 5 0.976 0.961 241 50 5 0.998 0.358 5 0.997 0.210 5 0.991 0.357 5 0.998 0.989 0.980 0.910 0.337 5 0.994 1.762 5 0.989 0.986 1.670 3 0.981 0.345 3 0.987 0.162 242 73 4 0.997 0.258 4 0.999 0.473 4 1.000 0.931 4 0.987 0.593 4 0.999 0.272 243 1 7 0.998 0.328 3 0.999 0.473 4 1.000 </td <td></td>																	
241 40 5 0.993 0.331 5 0.992 0.210 5 0.994 0.344 5 0.993 1.573 5 0.976 0.961 241 50 5 0.997 0.327 5 0.997 0.210 5 0.997 0.350 5 0.999 1.560 5 0.980 0.980 0.810 3 0.986 1.670 3 0.981 0.345 3 0.987 0.156 5 0.994 1.762 5 0.983 1.056 5 0.998 0.388 0.986 0.616 3 0.094 1.093 0.986 1.670 3 0.981 0.345 3 0.987 0.158 0.987 0.158 0.988 0.388 0.996 1.093 4 0.997 0.556 0.999 0.277 0.273 0.999 0.277 0.294 0.0999 0.388 3 0.999 0.763 3 1.000 0.845 3 1.000 0.432																	
241 50 5 0.997 0.327 5 0.997 0.210 5 0.997 0.350 5 0.999 1.560 5 0.989 0.934 241 60 5 0.998 0.358 5 0.994 0.203 5 0.991 0.337 5 0.994 1.762 5 0.983 1.058 242 56 3 1.000 0.285 3 0.980 0.810 3 0.986 1.670 3 0.981 0.345 3 0.987 0.168 242 73 4 0.997 0.258 4 0.995 0.545 4 0.997 0.533 4 0.995 0.546 4 0.997 0.546 4 0.997 0.546 4 0.997 0.568 0.999 0.473 4 1.000 0.931 4 0.997 0.568 0.999 0.277 243 2 5 0.998 0.318 3 0.999 </td <td></td>																	
241 60 5 0.998 0.358 5 0.994 0.203 5 0.991 0.337 5 0.994 1.762 5 0.983 1.058 242 56 3 1.000 0.285 3 0.980 0.810 3 0.986 1.670 3 0.981 0.345 3 0.985 0.292 242 73 4 0.999 0.258 4 0.995 0.545 4 0.996 1.693 4 0.987 0.598 0.162 243 1 7 0.998 0.225 2 0.999 0.328 3 0.999 0.473 4 1.000 0.931 4 0.999 0.546 4 0.999 0.227 243 2 5 0.988 0.179 0.388 3 0.999 0.763 3 1.000 0.845 3 1.000 0.432 244 102 5 0.999 0.388 5																	
242 56 3 1.000 0.285 3 0.980 0.810 3 0.986 1.670 3 0.981 0.345 3 0.987 0.165 242 73 4 0.997 0.317 4 0.995 0.545 4 0.996 1.093 4 0.997 0.553 4 0.999 0.277 243 1 7 0.998 0.179 2 2 5 0.988 0.179 2 244 70 3 0.999 0.328 3 0.998 0.388 3 0.999 0.763 3 1.000 0.845 3 1.000 0.431 244 86 4 0.999 0.329 4 1.000 0.411 4 1.000 0.837 4 0.999 0.881 4 1.000 0.432 244 102 5 0.999 0.417 5 0.999 0.844 5 1.000 0.804																	
242 73 4 0.967 0.317 4 0.995 0.545 4 0.996 1.093 4 0.987 0.593 4 0.985 0.294 242 90 4 0.999 0.258 4 0.999 0.473 4 1.000 0.931 4 0.999 0.546 4 0.999 0.277 243 1 7 0.998 0.328 3 0.998 0.388 3 0.999 0.763 3 1.000 0.845 3 1.000 0.430 244 70 3 0.999 0.329 4 1.000 0.411 4 1.000 0.837 4 0.999 0.841 5 1.000 0.432 244 102 5 0.999 0.315 5 0.999 0.388 5 1.000 0.804 5 1.000 0.754 5 1.000 0.752 4 0.999 0.812 4 0.995 0.812																	
242 90 4 0.999 0.258 4 0.999 0.473 4 1.000 0.931 4 0.999 0.546 4 0.999 0.277 243 1 7 0.998 0.225 5 0.988 0.179 5 0.999 0.763 3 1.000 0.845 3 1.000 0.430 244 70 3 0.999 0.329 4 1.000 0.411 4 1.000 0.837 4 0.999 0.801 4 1.000 0.432 244 102 5 0.999 0.315 5 0.999 0.417 5 0.999 0.884 5 1.000 0.754 5 1.000 0.432 245 1 5 0.999 0.438 5 1.000 0.804 5 1.000 0.804 5 1.000 0.804 6 1.000 0.804 9.995 0.812 4 0.995 0.408 4						3			3			3					
243 1 7 0.998 0.225 243 2 5 0.988 0.179 244 70 3 0.999 0.328 3 0.998 0.388 3 0.999 0.763 3 1.000 0.845 3 1.000 0.430 244 86 4 0.999 0.315 5 0.999 0.417 5 0.999 0.844 5 1.000 0.754 5 1.000 0.373 245 1 5 0.999 0.417 5 0.999 0.844 5 1.000 0.754 5 1.000 0.373 245 2 5 1.000 0.454 5 1.000 0.804 2 1 1.000 0.408 245 3 4 0.997 0.426 4 1.000 0.408 2 4 0.995 0.408 246 30 4 0.997 0.276 4 1.000			4	0.967	0.317	4	0.995	0.545	4	0.996	1.093	4	0.987	0.593	4	0.985	0.294
243 2 5 0.988 0.179 244 70 3 0.999 0.328 3 0.998 0.388 3 0.999 0.763 3 1.000 0.845 3 1.000 0.430 244 86 4 0.999 0.329 4 1.000 0.411 4 1.000 0.837 4 0.999 0.801 4 1.000 0.373 245 1 5 0.999 0.388 5 1.000 0.804 5 1.000 0.754 5 1.000 0.373 245 2 5 1.000 0.454 5 1.000 0.804 2 2 5 1.000 0.409 0.804 2 2 5 1.000 0.409 4 1.000 0.725 4 0.995 0.812 4 0.995 0.408 246 50 4 0.999 0.176 4 1.000 0.409 4 1.000 </td <td>242</td> <td>90</td> <td>4</td> <td>0.999</td> <td>0.258</td> <td>4</td> <td>0.999</td> <td>0.473</td> <td>4</td> <td>1.000</td> <td>0.931</td> <td>4</td> <td>0.999</td> <td>0.546</td> <td>4</td> <td>0.999</td> <td>0.277</td>	242	90	4	0.999	0.258	4	0.999	0.473	4	1.000	0.931	4	0.999	0.546	4	0.999	0.277
244 70 3 0.999 0.328 3 0.998 0.388 3 0.999 0.763 3 1.000 0.845 3 1.000 0.430 244 86 4 0.999 0.329 4 1.000 0.411 4 1.000 0.837 4 0.999 0.801 4 1.000 0.393 245 1 5 0.999 0.388 5 1.000 0.804 5 1.000 0.754 5 1.000 0.454 5 1.000 0.804 5 1.000 0.754 5 1.000 0.804 5 1.000 0.804 5 1.000 0.804 5 1.000 0.804 8 5 1.000 0.804 8 5 1.000 0.804 8 5 1.000 0.804 8 8 1.000 0.408 4 1.000 0.408 4 1.000 0.408 4 1.0099 0.532 4 0.995	243	1	7	0.998	0.225												
244 86 4 0.999 0.329 4 1.000 0.411 4 1.000 0.837 4 0.999 0.801 4 1.000 0.393 244 102 5 0.999 0.315 5 0.999 0.417 5 0.999 0.484 5 1.000 0.754 5 1.000 0.373 245 2 5 1.000 0.454 5 1.000 0.893 245 3 4 0.995 0.490 4 1.000 0.942 246 30 4 0.995 0.217 4 1.000 0.408 4 0.995 0.408 246 50 4 0.995 0.217 4 1.000 0.409 4 0.995 0.512 4 0.995 0.408 246 60 4 0.995 0.217 4 1.000 0.407 4 0.998 0.173 4 1.000 0.841 <t< td=""><td>243</td><td>2</td><td>5</td><td>0.988</td><td>0.179</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<>	243	2	5	0.988	0.179												
244 86 4 0.999 0.329 4 1.000 0.411 4 1.000 0.837 4 0.999 0.801 4 1.000 0.393 244 102 5 0.999 0.315 5 0.999 0.417 5 0.999 0.844 5 1.000 0.754 5 1.000 0.373 245 2 5 1.000 0.454 5 1.000 0.893 245 3 4 0.995 0.490 4 1.000 0.942 246 30 4 0.995 0.296 4 0.999 0.365 4 1.000 0.494 4 0.995 0.812 4 0.995 0.408 246 50 4 0.995 0.217 4 1.000 0.409 4 0.993 0.532 4 0.995 0.408 246 60 4 0.998 0.176 4 1.000 0.409 4 1.000	244	70	3	0.999	0.328	3	0.998	0.388	3	0.999	0.763	3	1.000	0.845	3	1.000	0.430
244 102 5 0.999 0.315 5 0.999 0.417 5 0.999 0.844 5 1.000 0.754 5 1.000 0.373 245 1 5 0.999 0.388 5 1.000 0.804 2 1 1.000 0.9754 5 1.000 0.373 245 2 5 1.000 0.454 5 1.000 0.893 2 2 4 0.995 0.490 4 1.000 0.942 2 4 0.995 0.409 4 1.000 0.725 4 0.995 0.812 4 0.995 0.261 246 50 4 0.995 0.217 4 1.000 0.407 4 1.000 0.855 4 0.995 0.532 4 0.992 0.261 246 60 4 0.999 0.176 4 1.000 0.427 4 1.000 0.855 4 0.998 <t></t>	244	86	4												4	1.000	0.393
245 1 5 0.999 0.388 5 1.000 0.804 2 893 4 0.995 0.490 4 1.000 0.893 4 0.995 0.490 4 1.000 0.725 4 0.995 0.408 2 4 0.995 0.408 4 0.995 0.217 4 1.000 0.409 4 1.000 0.804 4 0.993 0.532 4 0.992 0.266 246 50 4 0.999 0.176 4 1.000 0.409 4 1.000 0.804 4 0.993 0.532 4 0.992 0.266 246 60 4 0.999 0.176 4 1.000 0.427 4 1.000 0.855 4 0.998 0.412 4 0.999 0.206 246 80 4 0.998 0.173 4 1.000 0.811 4 0.998 0.402 4 0.998 0.222																	
245 2 5 1.000 0.454 5 1.000 0.893 245 3 4 0.995 0.490 4 1.000 0.942 246 30 4 0.995 0.217 4 1.000 0.409 4 1.000 0.804 4 0.995 0.406 246 50 4 0.995 0.217 4 1.000 0.409 4 1.000 0.855 4 0.999 0.412 4 0.992 0.263 246 60 4 0.999 0.176 4 1.000 0.427 4 1.000 0.855 4 0.999 0.412 4 0.999 0.203 246 70 4 0.998 0.173 4 1.000 0.431 4 1.000 0.855 4 0.998 0.402 4 0.999 0.203 247 1 4 1.000 0.330 4 0.995 0.333			•	0.355	0.0.5							J	1.000	0.,01	J		0.0.0
245 3 4 0.995 0.490 4 1.000 0.942 246 30 4 0.997 0.296 4 0.999 0.365 4 1.000 0.725 4 0.995 0.812 4 0.995 0.408 246 50 4 0.999 0.176 4 1.000 0.409 4 1.000 0.855 4 0.999 0.412 4 0.999 0.269 246 70 4 0.998 0.173 4 1.000 0.431 4 1.000 0.855 4 0.998 0.402 4 0.998 0.203 246 80 4 0.998 0.173 4 1.000 0.419 4 1.000 0.855 4 0.998 0.402 4 0.998 0.203 247 1 4 1.000 0.491 4 1.000 0.689 4 0.998 0.455 4 0.999 0.263						_											
246 30 4 0.997 0.296 4 0.999 0.365 4 1.000 0.725 4 0.995 0.812 4 0.995 0.402 246 50 4 0.995 0.217 4 1.000 0.409 4 1.000 0.804 4 0.993 0.532 4 0.999 0.266 246 60 4 0.999 0.176 4 1.000 0.427 4 1.000 0.855 4 0.999 0.412 4 0.999 0.206 246 80 4 0.998 0.173 4 1.000 0.411 4 1.000 0.855 4 0.998 0.402 4 0.999 0.202 246 80 4 0.998 0.191 4 1.000 0.841 4 0.998 0.455 4 0.998 0.455 4 0.999 0.222 247 1 4 1.000 0.333																	
246 50 4 0.995 0.217 4 1.000 0.409 4 1.000 0.804 4 0.993 0.532 4 0.992 0.263 246 60 4 0.999 0.176 4 1.000 0.427 4 1.000 0.855 4 0.999 0.412 4 0.998 0.203 246 70 4 0.998 0.173 4 1.000 0.431 4 1.000 0.855 4 0.998 0.402 4 0.998 0.203 246 80 4 0.998 0.171 4 1.000 0.431 4 1.000 0.841 4 0.998 0.455 4 0.998 0.455 4 0.999 0.252 247 1 4 1.000 0.309 4 0.999 0.333 4 1.000 0.689 4 0.994 1.167 4 1.000 0.564 247 2			4	0 997	0 296							1	0 995	0.812	1	n aas	0.408
246 60 4 0.999 0.176 4 1.000 0.427 4 1.000 0.855 4 0.999 0.412 4 0.999 0.202 246 70 4 0.998 0.173 4 1.000 0.431 4 1.000 0.855 4 0.998 0.402 4 0.998 0.202 246 80 4 0.998 0.191 4 1.000 0.412 4 0.998 0.402 4 0.998 0.202 247 1 4 1.000 0.498 4 1.000 0.681 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.998 0.455 4 0.999 0.228<																	
246 70 4 0.998 0.173 4 1.000 0.431 4 1.000 0.855 4 0.998 0.402 4 0.998 0.203 246 80 4 0.998 0.191 4 1.000 0.419 4 1.000 0.841 4 0.998 0.455 4 0.997 0.221 247 1 4 1.000 0.390 4 0.999 0.316 4 1.000 0.689 4 0.994 1.167 4 1.000 0.535 4 1.000 1.290 4 0.999 0.642 247 2 4 1.000 0.426 4 0.999 0.298 4 1.000 0.607 4 0.999 1.430 4 1.000 0.702 247 4 4 1.000 0.438 4 1.000 0.584 4 1.000 1.483 4 1.000 0.750 248 1																	
246 80 4 0.998 0.191 4 1.000 0.419 4 1.000 0.841 4 0.998 0.455 4 0.997 0.22 247 1 4 1.000 0.399 4 0.995 0.333 4 1.000 0.689 4 0.994 1.167 4 1.000 0.564 247 2 4 1.000 0.408 4 0.999 0.316 4 1.000 0.635 4 1.000 1.290 4 0.999 0.647 247 3 4 1.000 0.607 4 0.999 0.298 4 1.000 0.667 4 0.999 1.430 4 1.000 0.750 247 4 4 1.000 0.438 4 1.000 0.295 4 1.000 0.584 4 1.000 1.483 4 1.000 0.750 248 1 5 0.999 0.197 5 0.995 0.248 5 0.996 0.792 248 3 <																	
247 1 4 1.000 0.390 4 0.995 0.333 4 1.000 0.689 4 0.994 1.167 4 1.000 0.566 247 2 4 1.000 0.408 4 0.999 0.316 4 1.000 0.635 4 1.000 1.290 4 0.999 0.641 247 3 4 1.000 0.426 4 0.999 0.298 4 1.000 0.607 4 0.999 1.430 4 1.000 0.756 247 4 4 1.000 0.438 4 1.000 0.295 4 1.000 0.584 4 1.000 1.483 4 1.000 0.756 248 1 5 0.999 0.197 5 0.995 0.248 5 0.998 0.792 248 2 6 0.996 0.191 6 0.999 0.241 7 0.998 0.788																	
247 2 4 1.000 0.408 4 0.999 0.316 4 1.000 0.635 4 1.000 1.290 4 0.999 0.643 247 3 4 1.000 0.426 4 0.999 0.298 4 1.000 0.607 4 0.999 1.430 4 1.000 0.702 247 4 4 1.000 0.438 4 1.000 0.295 4 1.000 0.584 4 1.000 1.483 4 1.000 0.750 248 1 5 0.999 0.197 5 0.995 0.248 5 0.998 0.793 248 2 6 0.996 0.191 6 0.999 0.241 6 0.996 0.792 248 3 7 0.996 0.195 7 0.998 0.248 7 0.998 0.788																	
247 3 4 1.000 0.426 4 0.999 0.298 4 1.000 0.607 4 0.999 1.430 4 1.000 0.702 247 4 4 1.000 0.438 4 1.000 0.295 4 1.000 0.584 4 1.000 1.483 4 1.000 0.750 248 1 5 0.999 0.197 5 0.995 0.248 5 0.996 0.793 248 2 6 0.996 0.191 6 0.999 0.241 6 0.996 0.792 248 3 7 0.996 0.195 7 0.998 0.248 7 0.998 0.788																	
247 4 1.000 0.438 4 1.000 0.295 4 1.000 0.584 4 1.000 1.483 4 1.000 0.750 248 1 5 0.999 0.197 5 0.995 0.248 5 0.998 0.793 248 2 6 0.996 0.191 6 0.999 0.248 6 0.996 0.792 248 3 7 0.996 0.195 7 0.998 0.248 7 0.998 0.788		2															
248 1 5 0.999 0.197 5 0.995 0.248 5 0.998 0.793 248 2 6 0.996 0.191 6 0.999 0.241 6 0.996 0.792 248 3 7 0.996 0.195 7 0.998 0.248 7 0.998 0.788																	
248 2 6 0.996 0.191 6 0.999 0.241 6 0.996 0.792 248 3 7 0.996 0.195 7 0.998 0.248 7 0.998 0.788									4	1.000	0.584				4	1.000	0.750
248 3 7 0.996 0.195 7 0.998 0.248 7 0.998 0.788																	
												-					
249 100 3 0.995 0.151 3 1.000 0.399 3 1.000 0.875 3 0.996 0.380 3 0.994 0.173																	
	249	100	3	0.995	0.151	3	1.000	0.399	3	1.000	0.875	3	0.996	0.380	3	0.994	0.173

Table B.3. (continued)

				Al	lomet	ric Rel	ationsh:	ips F	it by P	rincipa	1 Com	ponent	Analysi	sc		
Id.		1	og ĥ -	log w	10	g DBH -	log w	log	BSLA -	log ₩	10	g h - 1	og DBH	<u>log</u>	h - 10	g BSLA
Code ^a	Cond.b	n	r	$\hat{\Phi}_{\text{hw}}$	n	r	$\boldsymbol{\hat{\Phi}}_{DW}$	n	r	$\boldsymbol{\hat{\Phi}}_{BW}$	n	r	$\hat{\phi}_{hD}$	n	r	ŶhB
249	110	3	1.000	0.158	3	0.999	0.409	3	1.000	0.874	3	0.999	0.386	3	1.000	0.181
249	120	3	0.998	0.155	3	1.000	0.365	3	1.000	0.881	3	0.997	0.425	3	0.998	0.176
250 250	2 3				4	0.999	0.442									
250	1	27	1.000	0.315	8	0.999	0.332				27	0.999	0.000			
252	120	3	0.999	0.342	27 3	1.000	0.394	3	1.000	0.598	27 3	0.999	0.800 1.115	3	0.998	0.571
252	140	5	1.000	0.342	5 5	1.000	0.299	5 5	0.999	0.601	5 5	0.999	1.228	5	0.999	0.611
252	160	9	0.999	0.379	9	0.999	0.312	9	0.991	0.704	9	1.000	1.214	9	0.994	0.535
252	180	9	0.999	0.384	9	0.998	0.333	9	0.998	0.665	9	0.999	1.152	9	0.999	0.578
252	200	9	0.999	0.376	9	0.997	0.341	9	0.997	0.683	9	0.999	1,101	9	0.999	0.550
252	220	9	0.999	0.368	9	0.997	0.346	9	0.996	0.687	9	0.999	1.064	9	0.998	0.535
252	240	9	0.999	0.357	9	0.997	0.348	9	0.996	0.698	9	0.998	1.024	9	0.998	0.510
253	47	10	0.999	0.173	10	1.000	0.251	10	1,000	0.582	10	1.000	0.690	10	0.999	0.297
253	57	13	0.999	0.192	13	1.000	0.259	13	0.999	0.668	13	0.999	0.740	13	0.999	0.287
253	66	14	0.994	0.192	14	0.998	0.259	14	0.999	0.721	14	0.999	0.742	14	0.996	0.266
254	50	12	1.000	0.212	12	0.997	0.238	12	1.000	0.651	12	0.998	0.888	12	0.999	0.325
254	61	9	0.999	0.247	9	1.000	0.287	9	0.998	0.688	9	0.999	0.862	9	0.997	0.359
254	70	1]	0.999	0.275	11	0.998	0.307	11	0.999	0.684	11	0.999	0.896	11	1.000	0.403
255	87	7	0.996	0.283	7	1.000	0.385		3 000	0.544	7	0.996	0.738		1 000	0 556
256 256	40 53	4	1.000	0.308	4	1.000	0.270	4	1.000	0.544	4	1.000	1.139	4	1.000	0.566
256	66	4	0.999	0.322	4 4	1.000	0.292	4	1.000	0.575 0.617	4	1.000	1.101	4	1.000	0.559
257	45	13	0.986	0.301	13	1,000	0.365	13	1.000	0.726	13	0.989	0.833	13	0.990	0.416
257	49	11	0.997	0.317	11	1.000	0.385	11	1.000	0.762	11	0.997	0.823	11	0.997	0.416
257	53	11	0.998	0.361	11	0.999	0.344	11	1.000	0.689	11	0.998	1.049	11	0.999	0.524
257	54	11	1.000	0.304	ii	1.000	0.357	11	1.000	0.707	11	0.999	0.853	ii	0.999	0.431
257	57	10	0.999	0.517	10	0.999	0.351	10	1.000	0.689	10	0.999	1.474	10	0.999	0.751
258	40	6	0.999	0.306	6	0.999	0.360	6	0.999	0.725	6	0.999	0.850	6	0.999	0.422
258	52	6	0.999	0.301	6	1.000	0.333	6	1.000	0.664	6	0.999	0.904	6	0.999	0.453
258	60	6	0.959	0.368	6	1.000	0.336	6	1.000	0.669	6	0.960	1.135	6	0.960	0.555
259	100	14	0.985	0.261	14	1.000	0.402	14	1.000	0.810	14	0.981	0.652	14	0.980	0.321
259	110	13	0.987	0.243	13	0.999	0.412	13	1.000	0.830	13	0.982	0.591	13	0.985	0.293
259	120	13	0.987	0.245	13	0.999	0.414	13	1.000	0.839	13	0.984	0.594	13	0.984	0.292
259	130	13	0.985	0.246	13	1.000	0.415	13	1.000	0.845	13	0.982	0.596	13	0.982	0.291
259	140	13	0.983	0.248	13	1.000	0.423	13	1.000	0.850	13	0.979	0.589	13	0.980	0.291
259 259	150 160	13	0.986	0.246 0.249	13	1.000	0.426	13	1.000	0.857	13	0.983	0.580	13	0.984	0.287
259	170	13 13	0.984	0.249	13 13	1.000	0.428 0.427	13	1.000	0.858	13 13	0.981	0.584 0.585	13	0.982	0.290
259	180	13	0.983	0.249	13	1.000	0.427	13 13	1.000	0.856 0.861	13	0.983	0.585	13	0.982	0.290
259	190	13	0.984	0.253	13	1.000	0.429	13	1.000	0.863	13	0.981	0.592	13	0.982	0.293
259	200	13	0.984	0.255	13	1.000	0.434	13	1.000	0.865	13	0.982	0.591	13	0.982	0.294
259	210	13	0.984	0.252	13	1.000	0.433	13	1.000	0.868	13	0.982	0.586	13	0.982	0.290
260	60	7	0.991	0.231	7	1.000	0.386	7	1.000	0.766	7	0.990	0.601	7	0.990	0.302

Table B.3. (continued)

				Al	lomet	ric Rel	ationsh:	ips F	it by P	rincipal	l Com	ponent	Analysi	s ^C		
* 1		_1	og h -	log ₩	10	g DBH -	log ₩	log	BSLA -	log w	10	g ĥ - 1	og DBH	<u>log</u>	ñ - 1o	g BSLA
Id. Code ^a	Cond.b	n	r	^ф hw	n	r	Φ̂Dw	n	r	φ̂Βw	n	r	φ̂hD	n	r	Φ̂hB
260	70	9	0.990	0.249	9	0.999	0.372	ō	1.000	0.773	9	0.987	0.671	9	0.990	0.322
260	80	9	0.986	0.247	9	0.993	0.413	9	1.000	0.776	9	0.978	0.595	9	0.984	0.318
260	90	9	0.992	0.238	9	0.999	0.395	9	1.000	0.790	9	0.988	0.603	Q.	0.990	0.301
260	100	9	0.989	0.243	9	1.000	0.403	9	1.000	0.810	9	0.986	0.606	9	0.987	0.300
260	110	9	0.989	0.245	9	0.999	0.412	9	1.000	0.820	9	0.984	0.595	9	0.985	0.298
260	120	9	0.986	0.249	9	1.000	0.430	9	1.000	0.844	9	0.987	0.583	9	0.983	0.295
260	130	9	0.987	0.244	9	1.000	0.425	9	1.000	0.847	9	0.988	0.578	9	0.985	0.288
260	140	9	0.985	0.251	9	0.999	0.441	9	1.000	0.858	9	0.988	0.574	9	0.983	0.293
260	150	9	0.985	0.244	9	1.000	0.423	9	1.000	0.857	9	0.985	0.579	9	0.984	0.284
260	160	9	0.985	0.245	9	1.000	0.425	9	1.000	0.860	9	0.986	0.580	9	0.985	0.285
260	170	9	0.981	0.253	9	0.998	0.440	9	1.000	0.857	9	0.987	0.580	9	0.979	0.295
260	180	9	0.988	0.244	9	1.000	0.427	9	1.000	0.864	9	0.989	0.574	9	0.988	0.282
261	60	8	0.978	0.227	8	1.000	0.366	8	1.000	0.745	8	0.975	0.625	8	0.980	0.306
261	70	8	0.985	0.226	8	0.999	0.362	8	1.000	0.755	8	0.979	0.625	8	0.986	0.299
261	80 90	9	0.989	0.246	9	0.993	0.400	9	1.000	0.750	9	0.980	0.611	9	0.986 0.988	0.327
261 261		9	0.989	0.233	9	0.999	0.387	9	1.000	0.774	9	0.985	0.604	9		0.301
261	100 110	9	0.990	0.233	9	1.000	0.390	9	1.000	0.784 0.796	9	0.986	0.599 0.619	9	0.987	0.296
261	120	9	0.992	0.254	9	1.000	0.410	9	1.000	0.796	9	0.991	0.619	9	0.988	0.311
261	130	9	0.991	0.234	9	1.000	0.412	9	1.000	0.807	9	0.991	0.520	9	0.988	0.299
261	140	9	0.988	0.247	9	1.000	0.414	9	1.000	0.834	9	0.991	0.599	9	0.984	0.303
261	150	9	0.988	0.233	9	1.000	0.429	9	1.000	0.834	9	0.987	0.594	9	0.986	0.295
261	160	9	0.987	0.243	9	1.000	0.413	9	1.000	0.836	9	0.988	0.592	9	0.988	0.291
262	70	13	0.996	0.361	13	0.998	0.413	13	1,000	0.740	13	0.990	0.929	13	0.996	0.487
262	80	13	0.996	0.363	13	0.997	0.382	13	1.000	0.740	13	0.985	0.949	13	0.996	0.489
262	90	13	0.996	0.368	13	0.998	0.391	13	1.000	0.737	13	0.988	0.949	13	0.996	0.499
262	100	13	0.996	0.365	13	0.997	0.383	13	1.000	0.737	13	0.988	0.953	13	0.996	0.495
262	110	13	0.996	0.370	13	0.997	0.386	13	1.000	0.736	13	0.986	0.960	13	0.995	0.503
262	120	13	0.996	0.368	13	0.996	0.381	13	1.000	0.735	13	0.985	0.964	13	0.996	0.501
262	130	13	0.996	0.368	13	0.997	0.385	13	1.000	0.735	13	0.986	0.955	13	0.996	0.500
262	140	13	0.996	0.372	13	0.997	0.388	13	1.000	0.736	13	0.986	0.958	13	0.996	0.506
262	150	13	0.991	0.359	13	0.996	0.380	13	0.997	0.722	13	0.982	0.949	13	0.991	0.498
263	26	4	1.000	0.250	4	0.994	0.275	4	0.996	0.596	4	0.990	0.905	4	0.994	0.418
263	33	7	0.998	0.281	ż	0.999	0.261	7	0.999	0.578	ż	0.997	1.079	7	0.997	0.487
263	39	9	0.999	0.317	9	1.000	0.269	9	0.999	0.600	9	1.000	1.179	9	0.999	0.527
264	40	4	0.998	0.128	4	0.993	0.172	4	0.999	0.838	4	0.998	0.738	4	0.995	0.152
264	50	7	0.994	0.182	7	1.000	0.232	7	1.000	0.740	7	0.995	0.786	7	0.991	0.245
265	77	4	0.965	0.137	4	1.000	0.419	4	1.000	0.840	4	0.958	0.327	4	0.959	0.163
266	60	8	0.996	0.209	8	1.000	0.438	8	1.000	0.873	8	0.997	0.478	8	0.996	0.240
266	80	11	0.997	0.238	11	1.000	0.431	11	1.000	0.862	11	0.995	0.553	11	0.996	0.276
266	100	11	0.993	0.232	- 11	1.000	0.415	11	1.000	0.833	11	0.993	0.559	11	0.993	0.278
266	120	12	0.984	0.245	12	1.000	0.410	12	1.000	0.821	12	0.984	0.601	12	0.984	0.298
266	140	8	0.995	0.342	8	1.000	0.385	8	0.999	0.770	8	0.992	0.891	8	0.992	0.443

Table B.3. (continued)

		Allometric Relationships Fit by Principal Component Analysis ^C														
Id. Code ^a	Cond.b	log h̄ − log w̄			log DBH − log w			log BSLA - log w			log h - log DBH			log h - log BSLA		
		n	r	$\hat{\Phi}_{hw}$	n	r	ŶDw	n	r	$\boldsymbol{\hat{\Phi}}_{BW}$	n	r	$\hat{\Phi}_{hD}$	n	r	Φ̂hB
267	42	8	0.987	0.138	8	0.999	0.418	8	1,000	0.830	8	0.982	0.329	8	0.983	0.165
268 269	54 75	7	0.993	0.147	7	0.999	0.346	_	0.000	0.701	7	0.991	0.426	_	0.000	0.224
270	60	6 14	0.982	0.187 0.327	6 14	0.998	0.395 0.377	6 14	0.998	0.791	6 14	0.969	0.472 0.873	6 14	0.969	0.234
270	70	14	0.991	0.327	14	1.000	0.377	14	1.000	0.742	14	0.989	0.873	14	0.993	0.439
270	80	14	0.994	0.320	14	0.999	0.365	14	1.000	0.723	14	0.993	0.881	14	0.993	0.439
270	90	14	0.991	0.321	14	1.000	0.362	14	1.000	0.726	14	0.991	0.893	14	0.990	0.442
270	100	14	0.993	0.315	14	1.000	0.364	14	1.000	0.716	14	0.993	0.870	14	0.990	0.439
270	110	14	0.990	0.313	14	0.999	0.359	14	1.000	0.709	14	0.991	0.877	14	0.989	0.441
270	120	14	0.991	0.311	14	1.000	0.357	14	1.000	0.714	14	0.991	0.877	14	0.989	0.436
271	40	18	0.997	0.305	18	0.998	0.366	18	0.999	0.735	18	0.997	0.835	18	0.998	0.416
271	50	18	0.991	0.331	18	0.999	0.382	18	0.998	0.761	18	0.996	0.872	18	0.997	0.437
271	60	18	0.994	0.321	18	0.998	0.382	18	0.999	0.756	18	0.998	0.843	18	0.998	0.426
271	70	18	0.994	0.320	18	0.998	0.379	18	0.999	0.751	18	0.998	0.847	18	0.998	0.427
271	80	18	0.994	0.310	18	0.999	0.365	18	0.999	0.739	18	0.997	0.853	18	0.997	0.421
271	90	18	0.994	0.309	18	0.998	0.358	18	0.999	0.725	18	0.997	0.865	18	0.997	0.426
271	100	18	0.994	0.311	18	0.999	0.353	18	0.999	0.723	18	0.997	0.884	18	0.997	0.431
271	110	18	0.994	0.306	18	0.999	0.354	18	0.999	0.723	18	0.997	0.868	18	0.996	0.425
272	50	6	0.989	0.267	6	0.999	0.312	6	0.999	0.623	6	0.981	0.859	6	0.981	0.426
272	60	7	0.997	0.308	7	0.994	0.287	7	0.994	0.573	7	0.983	1.071	7	0.983	0.532
272	71	7	0.998	0.319	7	0.997	0.269	7	0.997	0.537	7	0.992	1.186	7	0.992	0.591
272	81	8	0.998	0.384	8	0.998	0.303	8	0.998	0.609	8	0.998	1.267	8	0.998	0.630
272	103	10	0.989	0.294	10	0.997	0.306	10	0.997	0.613	10	0.997	0.967	10	0.997	0.482
273	40	15	0.998	0.267	15	0.996	0.321	15	1.000	0.716	15	0.996	0.829	15	0.996	0.372
273	50	15	0.999	0.272	15	0.999	0.350	15	0.999	0.701	15	0.997	0.774	15	0.997	0.387
273	60	15	0.999	0.278	15	0.999	0.347	15	0.999	0.695	15	0.997	0.800	15	0.996	0.398
273 273	70 80	15	0.998	0.276	15	0.999	0.341	15	0.999	0.693	15	0.995	0.809	15	0.995	0.397
273	90	15 15	0.998	0.280 0.272	15	0.999	0.356	15	0.999	0.713	15	0.998	0.786	15	0.999	0.400
273	100	15	0.999	0.272	15 15	0.998	0.346 0.338	15 15	1.000	0.680 0.685	15 15	0.997	0.785 0.781	15 15	0.998	0.400
274	60	8	0.999	0.269	8	1.000	0.368	8	1.000	0.763	8	0.999	0.731	8	1.000	0.352
274	70	8	0.999	0.261	8	1.000	0.359	8	1.000	0.750	8	0.999	0.728	8	0.999	0.348
274	80	8	0.999	0.260	8	1.000	0.358	8	1.000	0.749	8	0.999	0.727	8	1.000	0.348
274	90	9	0.998	0.270	9	0.999	0.369	9	1.000	0.750	9	1.000	0.733	9	0.998	0.360
274	100	9	0.998	0.273	g	0.998	0.370	9	1.000	0.741	g 9	1.000	0.738	ģ	0.998	0.368
276	84	9	0.996	0.411	9	0.998	0.329	g	0.999	0.660	9	0.991	1.255	9	0.991	0.622
276	95	9	0.997	0.415	9	0.998	0.330	ģ	0.999	0.662	9	0.993	1.260	ģ	0.993	0.625
276	112	9	0.997	0.415	9	0.998	0.331	9	0.998	0.662	9	0.992	1.256	9	0.992	0.625
277	70	4	0.999	0.233	4	1.000	0.402	4	1.000	0.777	4	0.999	0.581	4	0.999	0.301
277	80	7	1.000	0.223	7	0.999	0.366	7	1.000	0.761	7	0.999	0.609	7	1.000	0.293
277	90	9	0.999	0.224	9	0.999	0.359	9	1.000	0.756	9	1.000	0.625	9	0.999	0.297
277	100	9	0.998	0.220	9	0.997	0.358	9	0.999	0.749	9	1.000	0.613	9	0.995	0.293
277	110	9	0.996	0.213	9	0.997	0.362	9	0.999	0.748	9	1.000	0.591	9	0.992	0.284

Table B.3. (continued)

		log ĥ − log ₩			log DBH - log w		log BSLA - log w			log h - log DBH			log h - log BSLA			
Id. Code ^a	Cond.b	n	r	$\hat{\Phi}_{\text{hw}}$	n	r	$\boldsymbol{\hat{\Phi}}_{DW}$	n	r	∳ _{Bw}	n	r	$\hat{\Phi}_{hD}$	n	r	φ̂hΒ
277 278	120	7 19	0.997	0.195	7 19	0.999	0.349	7 19	0.999	0.760 0.870	7 19	1.000	0.560 0.547	7 19	0.994	0.256
279 280	40 51	10	0.996	0.202	10	1.000	0.392	10	1.000	0.788	10	0.994	0.515	10	0.995	0.256
281 281	27 30	Ü	0.330	0.200	5	1.000	0.393	5	1.000	0.786	Ü	0.333	0.552	Ů	0.333	0.270
281	33				5	1.000	0.401	5	1.000	0.807						

 $^{{}^{\}rm a}{}{}^{\rm Table}$ B.1 associates each Id. code with a particular yield table.

 $^{{}^{\}mathrm{b}}\mathsf{See}$ Table B.1 and the references given there for further information on condition.

CThe general formula for the allometric relation is $\log Y = \hat{\phi}_1 \log X + \hat{\phi}_0$, where Y is $\log \tilde{h}$, $\log \overline{OBH}$, or $\log \overline{BSLA}$ and X is $\log \tilde{w}$, $\log \overline{OBH}$, or $\log \overline{BSLA}$. \overline{OBH} and \overline{BSLA} are, respectively, the diameter at breast height and the basal area at breast height of the boles of individual trees. Y and X are paired as indicated in the table heading. The number of data points, correlation (r), and slope for each relationship are given. The correlation values have little statistical meaning because most of the natural variability in the original forest data has been removed from the yield table predictions. They are included here only as a crude index of variation around the fitted line. Values of the intercept, $\hat{\phi}_0$, were calculated, but were not used in the analyses and are not given here. Formulas for principal component analysis are given in Jolicoeur and Heusner (1971).

Table B.4 Self-thinning Lines from Forestry Yield Table
Data Cited by Previous Authors in Support of the
Self-thinning Rule.

-		No. Poi	of nts ^c	Thinn Lin	inq _d e	
Id. Code ^a	Condition ^b	nΤ	n	βe	â	Reference ^f
201				-0.33	4.06	White 1980
202,242,243				-0.54	3.46	White 1980
203				-0.57	3.98	White 1980
204				-0.63	3.54	White 1980 White 1980
206 207	49			-0.65 -0.73	3.53 3.06	White 1980 White 1980
208	43			-0.73	4.01	White 1980
209				-0.48	3.58	White 1980
210	136			-0.80	3.08	White 1980
211				-0.49	3.88	White 1980
212				-0.51	4.08	White 1980
213				-0.57	3.85	White 1980
216				-0.71	3.86	White 1980
217	54,64,75	30	26	-0.87	4 00	White 1981
221	maust a.f. 1	r	3	-0.54	4.00	White 1980
222	part of 1	5	3	-0.49		White 1981

aTable B.l associates each Id. code with a particular yield table. The presence of several codes in this column indicates that the reference did not specify which one of several yield tables from a single source was examined, making it impossible to determine which entry of Table B.l compares directly to the reported thinning line.

DSpecifies one of the conditions (site index, planting density, etc.) defined in Table B.l. Blanks indicate that the thinning reference did not state which condition or pooled combination of data from several conditions was analyzed. In these cases, it was not possible to determine which, if any, of the several thinning lines in Table B.2 from the same yield table is directly comparable with the one given above.

 $^{\text{C}}\text{n}_{T}$ is the total number of size-density data points given. n is the number of points remaining after removing points not relevant to the thinning line, so n is the number of points used to fit the thinning line. Blanks indicate that this information was not given in the thinning reference.

 ^{d}F it by regression of log \overline{w} against log N, or of log \overline{v} against log N for codes 217 and 222 (where \overline{v} is mean stem volume per tree).

 $^{\rm e}$ All slopes were reported for log $\overline{\rm w}{\rm -log}$ N thinning lines. The values above have been converted to log B-log N slopes by adding 1 (Chapter 1). This facilitates comparison with the analyses in Table B.2.

fThese are references for the self-thinning analysis only. Original data sources are given in Table B.1.

Donald Ernest Weller was born in Lansing, Michigan on September, 23, 1952. In 1957 he accompanied his family to their new home in Evansville, Indiana. There he benefited from 13 years of instruction in the public school system and graduated from Benjamin Bosse High School in June, 1970.

Beginning that fall, his college education advanced uneventfully in the quiet atmosphere of Wabash College, a "small liberal arts college for men," in Crawfordsville, Indiana. There, the unfortunate absence of feminine and urban distractions allowed Don to concentrate on his studies of biology, chemistry, and other subjects.

After a brief and unsatisfying foray into graduate level training in biochemistry at the California Institute of Technology in Pasadena, Don returned to Indiana and was employed at the Indiana University Medical School at Bloomington until July 1978. During this period, he developed interests in quantitative and theoretical ecology.

These interests motivated him to enroll in the Graduate Program in Ecology at The University of Tennessee, Knoxville where he was awarded a research assistantship from the Systems Ecology Group of the Environmental Sciences Division at Oak Ridge National Laboratory. While in Tennessee, Don met and married Deborah Mindel Goldsmith and earned a Ph.D. degree in ecology. In November, 1983 he accepted employment as a quantitative ecologist at the Smithsonian Environmental Research Center in Edgewater, Maryland.

Don is a member of Phi Beta Kappa, Phi Kappa Phi, the Ecological Society of America, the American Institute of Biological Sciences, and other professional and honorary organizations.