

# Elevated CO<sub>2</sub> increases the long-term decomposition rate of *Quercus myrtifolia* leaf litter

MYRA C. HALL\*, PETER STILING†, DANIEL C. MOON‡, BERT G. DRAKE§ and MARK D. HUNTER\*

\*Institute of Ecology, University of Georgia, Athens, GA 30602-2202, USA, †Department of Biology, University of South Florida, Tampa, FL 33620-5150, USA, ‡Department of Biology, University of North Florida, 4567 St. Johns Bluff Road South, Jacksonville, FL 32224, USA, §Smithsonian Environmental Research Center, PO Box 28, Edgewater, MD 21037, USA

## Abstract

Decomposition of *Quercus myrtifolia* leaf litter in a Florida scrub oak community was followed for 3 years in two separate experiments. In the first experiment, we examined the effects CO<sub>2</sub> and herbivore damage on litter quality and subsequent decomposition. Undamaged, chewed and mined litter generated under ambient and elevated (ambient + 350 ppm V) CO<sub>2</sub> was allowed to decompose under ambient conditions for 3 years. Initial litter chemistry indicated that CO<sub>2</sub> levels had minor effects on litter quality. Litter damaged by leaf miners had higher initial concentrations of condensed tannins and nitrogen (N) and lower concentrations of hemicellulose and C:N ratios compared with undamaged and chewed litter. Despite variation in litter quality associated with CO<sub>2</sub>, herbivory, and their interaction, there was no subsequent effect on rates of decomposition under ambient atmospheric conditions. In the second experiment, we examined the effects of source (ambient and elevated) of litter and decomposition site (ambient and elevated) on litter decomposition and N dynamics. Litter was not separated by damage type. The litter from both elevated and ambient CO<sub>2</sub> was then decomposed in both elevated and ambient CO<sub>2</sub> chambers. Initial litter chemistry indicated that concentrations of carbon (C), hemicellulose, and lignin were higher in litter from elevated than ambient CO<sub>2</sub> chambers. Despite differences in C and fiber concentrations, litter from ambient and elevated CO<sub>2</sub> decomposed at comparable rates. However, the atmosphere in which the decomposition took place resulted in significant differences in rates of decomposition. Litter decomposing under elevated CO<sub>2</sub> decomposed more rapidly than litter under ambient CO<sub>2</sub>, and exhibited higher rates of mineral N accumulation. The results suggest that the atmospheric conditions during the decomposition process have a greater impact on rates of decomposition and N cycling than do the atmospheric conditions under which the foliage was produced.

*Keywords:* ammonium, decomposition, elevated CO<sub>2</sub>, herbivory, Kennedy space center, litter quality, mass loss, nitrate, *Quercus myrtifolia*

Received 15 August 2005; revised version received 20 September 2005; accepted 21 September 2005

## Introduction

Carbon dioxide (CO<sub>2</sub>) concentrations in the atmosphere are currently increasing because of increased fossil fuel use and deforestation. These increases in CO<sub>2</sub> along with associated climate changes are expected to con-

tinue and to have significant impacts on terrestrial ecosystems (Houghton *et al.*, 2001). There is abundant evidence that increased levels of CO<sub>2</sub> change the chemical composition of plant foliage (Curtis *et al.*, 1989; Ceulemans & Mousseau, 1994; Koch & Mooney, 1996; Hall *et al.*, 2005b). The most commonly reported changes in CO<sub>2</sub>-grown plant foliage, increased C:N ratios and decreased nitrogen (N) concentrations, might be predicted to change the rate of decomposition and alter nutrient cycling because these variables are associated with decomposition in natural systems (Floate,

Correspondence: Mark D. Hunter, Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109-1048, USA. tel. +734 615 4917, e-mail: mdhunter@umich.edu

1970; Berg & Ekbohm, 1983; Seastedt, 1988; Taylor *et al.*, 1989). However, it remains to be seen whether changes in plant foliage induced under elevated CO<sub>2</sub> are retained in senesced leaves. Curtis *et al.* (1989) found that increased C:N ratios in the green foliage of the sedge, *Scirpus olneyi*, under enriched CO<sub>2</sub> were not maintained in the senesced foliage. Finzi *et al.* (2001) found no effect of elevated CO<sub>2</sub> on total nonstructural carbohydrates or N in green leaves or in leaf litter of five tree species. In our scrub oak ecosystem, previous work has shown that elevated CO<sub>2</sub> reduces foliar N concentrations and increases C:N ratios by an average of 6% and 7%, respectively, in green foliage across three oak species (Hall *et al.*, 2005b). However, the senesced litter of these oaks does not exhibit lower N concentrations or higher C:N ratios. Rather, the strongest CO<sub>2</sub> effects are seen in the secondary metabolites, particularly condensed tannins (Hall *et al.*, 2005a).

In other studies, chemical differences in senesced foliage between ambient and elevated CO<sub>2</sub> have been observed. Changes in N (Coûteaux *et al.*, 1991; Kemp *et al.*, 1994; Cotrufo *et al.*, 1994, 1998; Robinson *et al.*, 1997), C:N ratios (Cotrufo *et al.*, 1994, 1998; Ball & Drake, 1997; Robinson *et al.*, 1997; Lutze *et al.*, 2000; Parsons *et al.*, 2004) lignin:N ratios (Cotrufo *et al.*, 1994, 1998, 1999; Parsons *et al.*, 2004), condensed tannins (Cotrufo *et al.*, 1999; Parsons *et al.*, 2004), lignin (Cotrufo *et al.*, 1994, 1999; Henry *et al.*, 2005), and non-structural carbohydrates (Lutze *et al.*, 2000) have been found in litter from enriched CO<sub>2</sub> plants compared with the controls.

In addition to changes induced by elevated CO<sub>2</sub>, other factors can influence litter chemistry, one of which is the effects of insect herbivores. Insect herbivory can change the chemical composition of plant foliage, which may lead to changes in litter quality (Grace, 1986; Findlay *et al.*, 1996; Chapman *et al.*, 2003). Grace (1986) showed that gypsy moth defoliation resulted in increased N concentrations in Pennsylvania oak forest litter. Chapman *et al.* (2003) found that scale and moth herbivory increased litter N concentrations in pinyon pines and subsequently, C:N and lignin:N ratios decreased. Also scale herbivory increased annual needle litterfall. This herbivore-induced increase in litter quality translated into higher decomposition rates. Although there are a number of studies that have examined the effects of elevated CO<sub>2</sub> on insect herbivores (Lincoln & Couvet, 1986; Fajer *et al.*, 1991; Roth & Lindroth, 1995; Stiling *et al.*, 1999, 2002, 2003), we are not aware of any previous studies that have examined in detail the interactions among CO<sub>2</sub>, insect herbivory, litter chemistry and decomposition.

In this study, we conducted two experiments to explore the links among elevated CO<sub>2</sub>, herbivore da-

mage, litter chemistry, decomposition, and N dynamics. Specifically, we investigated whether direct effects of CO<sub>2</sub> on foliage quality and litter dynamics matter more or less than indirect effects of CO<sub>2</sub> on litter dynamics mediated by other changes in the environment (Dukes & Hungate, 2002). In the first experiment, we tested the effects of CO<sub>2</sub> growth conditions and herbivore damage on the decomposition of *Quercus myrtifolia* leaf litter. In the second experiment we tested the effects of CO<sub>2</sub> growth conditions (source) and CO<sub>2</sub> decomposition conditions (site) on the decomposition of *Q. myrtifolia* litter without regard to herbivore damage.

## Methods

### Study site

Our study site lies within a 2 ha native scrub-oak community located at Kennedy Space Center, Florida. This woody ecosystem is controlled by a natural fire return cycle of 8–12 years and the mature canopy is 3–5 m high. The last burn was in 1996 before site set up. The ecosystem is subtropical with a mean annual temperature of 22.4 °C and an annual average precipitation of 131 cm. Sixteen 3.6 m diameter plots, each enclosed with a clear polyester film open-top chamber 3.4 m in height, were utilized to control CO<sub>2</sub> levels. Chambers were overlaid on an octagonal framework of PVC pipe with a removable access door and frustrum to reduce dilution of air within the chamber by outside wind. All regrowth was cut to ground level in May 1996 and, since that time, the vegetation in eight of the chambers has been exposed to almost twice ambient CO<sub>2</sub> (ambient + 350 ppm V), whereas the other eight chambers have been exposed to ambient levels of CO<sub>2</sub>. The CO<sub>2</sub> is supplied 24 h a day. Monitoring and control of CO<sub>2</sub> injection into each chamber is carried out by infrared gas analysis in conjunction with manually adjusted needle valves. In ambient CO<sub>2</sub> chambers, the airflow is identical to that of the elevated CO<sub>2</sub> chambers but is not supplemented with CO<sub>2</sub>. The study species, *Q. myrtifolia*, accounts for 76% by biomass of the species composition within this community.

*Experimental Design.* Rates of decomposition focused on litter from the most common tree in the system, *Quercus myrtifolia*. All vegetation within chambers had been exposed to either ambient or elevated CO<sub>2</sub> for 6 years before this study. Litter was collected and pooled (by chamber) from litter trays in all chambers during April–May of 2002, when litter was most plentiful. The litter was sorted into four categories: mined, chewed, undamaged and miscellaneous damage (more than one type of damage or damage other than chewers or

miners). This resulted in eight litter types (elevated CO<sub>2</sub> and ambient CO<sub>2</sub> × 4 damage types). Litterbags consisted of 5 cm × 4 cm × 1.5 mm mesh and contained 0.1000–0.5000 g of uncut air-dried litter from each of the eight treatments depending on availability. Although litter bags exclude larger invertebrates from gaining access to litter (Hunter *et al.*, 2003), microbial populations, protozoa, microarthropods, and nematodes have access. In the present study, our primary aim is to compare among treatment groups and we make the assumption that any changes in decomposition caused by litter bags are equivalent across treatments.

*Experiment I.* The undamaged, chewed and mined litter (UCM) was used to examine the effects of 'source' (ambient or elevated CO<sub>2</sub>) and herbivore damage on litter decomposition. Litter bags were placed in six 3.6 m diameter open (unchambered) PVC rings adjacent to the CO<sub>2</sub> chambers, with three replicates of each treatment per ring ( $n = 18$  per ring). Litter bags were covered by a one leaf thick layer of litter, available from the forest floor in each ring. One set (replicate) was collected from each ring annually for 3 years (four dates including time zero, May 2002). Time zero samples were analyzed for litter chemistry, and sample sizes were 16 undamaged (seven ambient, nine elevated), 20 chewed, 20 mined (ten from each CO<sub>2</sub> treatment) with sample sizes based on litter availability from pooled samples.

*Experiment II.* Each of the 16 chambers received six bags (one for each collection date) of the miscellaneous damaged (MD) litter from each of the ambient and elevated CO<sub>2</sub> treatments ( $n = 12$  bags per chamber). PVC rings (12 cm diameter) were used to contain litter bags and prevent excessive movement of bags within the chambers under conditions of heavy rain. Three PVC rings were fixed into the bare soil of each chamber and the tops of the rings were approximately 0.25–0.50 cm above the soil surface. Two litter bags from ambient CO<sub>2</sub> and two litter bags from elevated CO<sub>2</sub> were placed in each PVC ring within chambers. Each ring was covered by a one leaf thick layer of litter, available from the forest floor in each chamber. One bag of each CO<sub>2</sub> treatment was removed from each chamber every 6 months for 3 years from time 6 months (November, 2002) to time 3 years (May 2005) (seven dates including time zero, May 2002). The total number of bags to be sampled was 36; however, some bags were not recoverable or had 100% mass loss and N analyses (below) could not be performed on all samples. Thus, sample sizes varied among collection dates. Actual sample sizes are noted on tables and figures. Time zero sample sizes were 10 ambient and 10 elevated from pooled samples. Our design allowed us to

consider the effects of both 'source' (ambient and elevated CO<sub>2</sub>) and 'site' (ambient and elevated) on the decomposition process. Owing to limited litter availability, herbivore damage effects were not included in this experiment.

On each collection date for both experiments, litter samples were weighed to determine mass loss. In addition during Experiment II, two N pools in the litter were followed over 3 years of decomposition: the mineral pool (NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>) and dissolved organic nitrogen (DON) pool (See methods).

*Initial litter chemistry.* Subsamples of litter used in both experiments were retained for measures of initial litter chemistry. The litter was air dried then ground to a fine powder and stored at -80 °C before to analysis. Cellulose, hemicellulose, and lignin concentrations were determined by sequential neutral detergent/acid detergent digestion on an Ankom fiber analyzer (Abrahamson *et al.*, 2003). Proanthocyanidins, an estimate of condensed tannin concentration, were assayed using *N*-butanol:HCl methods described in Rossiter *et al.* (1988). Total phenolics were estimated using the Folin–Denis assay (Swain, 1980), and gallotannins (hydrolysable tannins) were estimated using a potassium iodate technique developed by Bate-Smith (1977) and modified by Schultz & Baldwin (1982). The standards for tannin analyses were generated from bulk litter samples from our field site by multiple sequential washes of acetone extraction. Standards were not purified further. The percent dry weight N and carbon (C) were estimated using a Carlo-Erba NA1500 model C/N analyzer (Milan, Italy). (Carlo Erba, Milan, Italy)

*Decomposition.* Litter was air-dried before weighing and weights were used to determine mass loss. The air-dried litter was ground to a fine powder and stored at -80 °C before analysis for *N*-dynamics (Experiment II only).

Ground litter (0.05 g) was hydrated in 20 mL of distilled water and agitated overnight in a shaker at room temperature. The 'leachate' was then filtered through a 0.4 μL filter. The filtrate was analyzed for NO<sub>3</sub><sup>-</sup>-N and NH<sub>4</sub><sup>+</sup>-N using the automated cadmium reduction and phenate assays, respectively, on an Alpkem segmented-flow autoanalyzer (Alpkem RFA 300, Alpkem Corporation, Clackamas, OR, USA). Total dissolved nitrogen (TDN) of the filtrate was estimated following persulfate oxidation (Koroleff, 1983; Qualls, 1989) and DON was calculated as the difference between TDN and the sum of NO<sub>3</sub><sup>-</sup>-N and NH<sub>4</sub><sup>+</sup>-N.

The residue (anything that did not pass through the filter) was oven-dried at 60 °C for 24 h then used to estimate percent dry weight nitrogen (DWN) on a

Carlo-Erba NA1500 model C/N analyzer. Total N was calculated as DWN plus TDN.

**Data Analysis.** All data were tested for normality and log-transformed where appropriate. The initial litter chemistry was analyzed using ANOVA models generated by the GLM procedure of SAS 8.2 (SAS Institute, 1998). The UCM litter was analyzed using the repeated measures GLM (SAS Institute, 1998). The MD litter N was analyzed using the MIXED procedure of SAS 8.2 (SAS Institute, 1998). The Student–Neuman–Keuls (SNK) post hoc test ( $\alpha = 0.05$ ) was used to distinguish among treatment means.

## Results

### Experiment I – UCM litter

**Litter chemistry.** The initial chemical composition of undamaged, chewed, and mined *Q. myrtifolia* litter derived from elevated and ambient CO<sub>2</sub> chambers is shown in Table 1. Litter from elevated CO<sub>2</sub> chambers had lower concentrations of cellulose ( $F_{1,48} = 7.07$ ,  $P < 0.01$ ), and hemicellulose ( $F_{1,50} = 17.0$ ,  $P < 0.01$ ) than litter from the ambient chambers. The litter from the two CO<sub>2</sub> treatments had similar N concentrations as

well as similar C:N ratios. Similarly, lignin and phenolic concentrations, often used as indicators of litter quality, were unaffected by CO<sub>2</sub> levels (Table 1).

Herbivore damage was associated with variation in initial litter chemistry (Table 2). Condensed tannin concentrations were higher in mined litter than in undamaged or chewed litter ( $F_{2,50} = 6.30$ ,  $P < 0.01$ ) and N concentrations were also higher in mined litter than in undamaged litter ( $F = 6.82_{2,50}$ ,  $P < 0.01$ ). The C:N ratio was higher in undamaged litter than in mined litter ( $F_{2,50} = 10.4$ ,  $P < .01$ ) and hemicellulose was higher in undamaged litter than in either chewed or mined leaf litter ( $F_{2,50} = 6.25$ ,  $P < 0.01$ ; Table 2).

Only condensed tannin concentrations showed an interaction between CO<sub>2</sub> and herbivore damage (Fig. 1). Mined litter was consistent in the concentrations of condensed tannins between CO<sub>2</sub> treatments. Undamaged litter from elevated CO<sub>2</sub> chambers had lower concentrations of condensed tannins than litter from elevated CO<sub>2</sub> chambers whereas chewed litter was the opposite with higher concentrations of condensed tannins in chewed litter from elevated CO<sub>2</sub> chambers ( $F_{2,50} = 5.46$ ,  $P < 0.01$ ; Fig. 1).

**Litter Decomposition.** Despite initial differences in litter chemistry associated with CO<sub>2</sub> treatment (Table 1) and

**Table 1** Initial chemical composition of undamaged, chewed, and mined *Quercus myrtifolia* litter from ambient and elevated CO<sub>2</sub> across damage categories


Sample	Condensed Tannins (%)	Hydrolyzable Tannins (%)	Total Phenolics (%)	Nitrogen (%)	Carbon (%)	C:N	Cellulose (%)	Hemicellulose (%)	Lignin (%)
Elevated	26.0 (1.25)	27.3 (0.95)	28.0 (1.77)	0.58 (0.01)	50.6 (0.98)	88.3 (2.04)	19.9 (0.32)	13.2 (0.27)	8.91 (0.34)
Ambient	23.3 (1.67)	25.9 (0.97)	27.1 (1.07)	0.58 (0.01)	50.2 (0.11)	87.9 (1.97)	21.0 (0.16)	14.5 (0.22)	9.71 (0.19)
	NS	NS	NS	NS	NS	NS	$P = 0.003$	$P < 0.001$	NS

Values are means,  $n = 29$  for elevated and  $n = 27$  for ambient, with SE in parentheses.

**Table 2** Initial chemical composition of undamaged, chewed, and mined *Quercus myrtifolia* litter by damage category across CO<sub>2</sub> treatments

Sample	Condensed Tannins (%)	Hydrolyzable Tannins (%)	Total Phenolics (%)	Nitrogen (%)	Carbon (%)	C:N	Cellulose (%)	Hemicellulose (%)	Lignin (%)
Undamaged	22.4 (1.29)	26.1 (1.51)	27.5 (1.77)	0.54 (0.018)	51.3 (1.76)	95.5 (2.09)	20.8 (0.32)	14.7 (0.31)	8.60 (0.40)
Chewed	22.3 (1.74)	27.7 (1.24)	28.2 (1.37)	0.58 (0.02)	50.5 (0.14)	88.8 (2.19)	20.7 (0.23)	13.6 (0.35)	9.51 (0.31)
Mined	28.9 (1.79)	26.0 (0.81)	27.1 (2.22)	0.62 (0.01)	49.6 (0.24)	81.6 (2.0)	19.9 (0.43)	13.4 (0.28)	9.64 (0.35)
	$P = 0.004$	NS	NS	$P = 0.002$	NS	$P < 0.001$	NS	$P = 0.004$	NS

Values are means,  $n = 16$  undamaged,  $n = 20$  chewed,  $n = 20$  mined, with SE in parentheses.



**Fig. 1** Effects of herbivore damage and CO<sub>2</sub> treatment on concentrations of condensed tannins in *Quercus myrtifolia* leaf litter. Data are the means of nine samples (undamaged ambient) seven samples (undamaged elevated) or ten samples (chewed or mined from ambient or elevated CO<sub>2</sub>). Bars represent standard errors, and different letters denote significantly different means.


herbivore activity (Table 2, Fig. 1), neither CO<sub>2</sub> treatment ( $F_{2,56} = 0.18$ ,  $P = 0.83$ ; Fig. 2a) nor herbivore damage ( $F_{2,56} = 0.38$ ,  $P = 0.82$ ; Fig. 2b) influenced the rates of litter decomposition.

#### Experiment II – MD Litter

**Initial Chemistry.** The initial chemical composition of miscellaneous damage litter is summarized in Table 3. Litter from elevated CO<sub>2</sub> chambers had higher concentrations of C, hemicellulose and lignin than did the litter from ambient chambers (C:  $F_{1,17} = 15.9$ ,  $P < 0.01$ ; hemicellulose:  $F_{1,18} = 11.8$ ,  $P < 0.01$ ; lignin:  $F_{1,18} = 23.9$ ,  $P < 0.01$ ).

**Litter Decomposition.** The rates of decomposition were comparable for litter originating from elevated and ambient CO<sub>2</sub> chambers ( $F_{1,123} = 1.38$ ,  $P = 0.24$ ; Fig. 3a), however the CO<sub>2</sub> level under which litter was decomposing did have an effect on mass loss. Litter decomposition was accelerated under elevated CO<sub>2</sub> ( $F_{1,123} = 11.7$ ,  $P < 0.01$ ; Fig. 3b). This effect of decomposition site was most pronounced from 18 to 36 months; before 18 months, decomposition rates under elevated and ambient CO<sub>2</sub> were comparable.

Ammonium accumulated to higher levels ( $F_{1,121} = 12.7$ ,  $P < 0.01$ ) in litter decomposing under elevated CO<sub>2</sub> than in litter decomposing under ambient CO<sub>2</sub> (Fig. 4a). Likewise, nitrate concentrations accumulated more rapidly in litter decomposing under elevated CO<sub>2</sub> ( $F_{1,121} = 2.67$ ,  $P = 0.03$ ; Fig. 4b). There were no effects of source chamber on the accumulation of mineral N in litter. Total litter N concentrations tended to increase



**Fig. 2** Decomposition rates for (a) UCM *Quercus myrtifolia* litter derived from ambient and elevated CO<sub>2</sub> and decomposing under ambient conditions for 3 years and (b) by herbivore damage category decomposing under ambient conditions for 3 years. Bars represent standard errors. Data are the means of 18 samples for ambient and elevated CO<sub>2</sub> treatments and 12 for undamaged, chewed and mined samples.

over time ( $F_{5,118} = 41.4$ ,  $P < 0.01$ ; Table 4). In general DON and mineral pools also increased during decomposition (Table 4) but these increases in concentration largely reflect the relative loss of C. The percent of initial litter C remaining on each sample date ( $F_{5,118} = 22.58$ ,  $P < 0.01$ ; Fig. 5a) declined faster in litter decomposing under elevated CO<sub>2</sub>. The percent of initial litter N remaining on each sample date ( $F_{5,118} = 0.79$ ,  $P = 0.56$ ; Fig. 5b) did not differ between litter decomposing under ambient and elevated CO<sub>2</sub> and did not vary systematically over time.


#### Discussion

Contrary to expectations, the source of leaf litter (ambient or elevated), its degree of damage, and its initial chemistry did not predict decomposition rates. Rather, litter decomposing under elevated CO<sub>2</sub> decomposed more rapidly than litter decomposing under ambient CO<sub>2</sub> (Fig. 3b). Coupled with the increase in decomposi-

**Table 3** Initial chemical composition of miscellaneous damaged *Quercus myrtifolia* litter


Sample	Condensed tannins (%)	Hydrolyzable tannins (%)	Total phenolics (%)	Nitrogen (%)	Carbon (%)	C:N	Cellulose (%)	Hemicellulose (%)	Lignin (%)
Elevated	24.1 (2.59)	28.9 (1.80)	31.9 (2.47)	0.62 (0.02)	50.8 (0.09)	82.6 (3.27)	20.4 (0.37)	14.5 (0.13)	10.5 (0.44)
Ambient	23.6 (2.32)	25.8 (1.21)	29.9 (2.14)	0.63 (0.04)	49.5 (0.37)	80.7 (4.92)	20.7 (0.41)	13.4 (0.30)	8.07 (0.26)
	NS	NS	NS	NS	$P < 0.001$	NS	NS	$P = 0.003$	$P < 0.001$

Values are means,  $n = 10$ , with SE in parentheses.



**Fig. 3** Decomposition rates for MD *Quercus myrtifolia* litter for (a) litter from ambient and elevated CO<sub>2</sub> treatments (source) and (b) litter decomposing in ambient and elevated CO<sub>2</sub> treatments (site). Sample sizes vary by time and treatment and are as follows in order of time: Source–ambient = 16, 16, 15, 8, 5, 12; Source–elevated = 16, 16, 15, 12, 4, 12; Site–ambient = 16, 16, 16, 14, 7, 8; Site–elevated = 16, 16, 14, 6, 2, 16. Bars represent standard errors.

tion rate under elevated CO<sub>2</sub> was an increase in rate of mineral N accumulation in leaf litter (Fig. 4) and an increase in the rate of C loss (Fig. 5a). Litter C may, therefore, be driving the accumulation of mineral N. Given that overall levels of N varied little over the experiment (Fig. 5b), it is tempting to conclude that rates of N mineralization were higher under elevated than ambient CO<sub>2</sub>. However, we cannot rule out the



**Fig. 4** Ammonium concentrations (a) and nitrate concentrations (b) in litter decomposing under ambient and elevated CO<sub>2</sub> (site) for all collection dates of MD *Quercus myrtifolia* litter. Bars represent standard errors. Sample sizes varies by time and treatment and are as follows in order of time: ambient = 16, 16, 16, 13, 7, 7; elevated = 16, 16, 14, 5, 2, 16.

possibility that organic N export was balanced by mineral N import.

In this study (Experiment I), initial litter quality of UCM litter exhibited only minor differences between ambient and elevated CO<sub>2</sub> and none of the primary regulators (N, C:N ratio, lignin) of decomposition were affected by CO<sub>2</sub> concentrations. Rather, litter cellulose and hemicellulose concentrations were higher in litter

**Table 4** Total nitrogen and nitrogen pools in miscellaneous damaged litter of *Quercus myrtifolia* on each collection date from source and site of decomposition

Source Site	Ambient				Elevated			
	Ambient		Elevated		Ambient		Elevated	
Total N								
6	0.41	(0.05) 8	0.40	(0.03) 8	0.38	(0.07) 8	0.41	(0.05) 8
12	0.46	(0.03) 8	0.46	(0.03) 8	0.37	(0.03) 8	0.44	(0.06) 8
18	0.75	(0.05) 8	0.80	(0.04) 7	0.71	(0.03) 8	0.85	(0.05) 7
24	0.72	(0.03) 6	0.93	(0.16) 2	0.56	(0.04) 7	0.80	(0.11) 3
30	1.04	(0.08) 4	1.31	(.) 1	0.73	(0.04) 3	0.64	(.) 1
36	0.76	(0.12) 4	0.83	(0.10) 8	0.78	(0.08) 3	0.96	(0.07) 8
DON								
6	1.08	(0.13)	1.15	(0.19)	0.84	(0.07)	0.88	(0.07)
12	1.09	(0.12)	1.45	(0.28)	0.83	(0.04)	1.44	(0.28)
18	2.04	(0.35)	2.22	(0.47)	1.23	(0.13)	2.34	(0.42)
24	2.17	(0.44)	1.39	(0.40)	1.99	(0.41)	1.56	(0.24)
30	4.52	(0.26)	3.91	(.)	4.69	(0.44)	3.95	(.)
36	5.74	(0.26)	6.10	(0.43)	5.21	(0.15)	6.30	(0.57)
NO <sub>3</sub> <sup>-</sup> N								
6	0.024	(0.010)	0.004	(0.003)	0.002	(0.002)	0.0002	(0.0003)
12	0.003	(0.002)	0.020	(0.009)	0.014	(0.007)	0.004	(0.001)
18	0.016	(0.003)	0.014	(0.004)	0.016	(0.005)	0.011	(0.003)
24	0.012	(0.001)	0.026	(0.019)	0.014	(0.002)	0.016	(0.002)
30	0.021	(0.005)	0.051	(.)	0.015	(0.005)	0.028	(.)
36	0.020	(0.002)	0.022	(0.001)	0.013	(0.0004)	0.017	(0.001)
NH <sub>4</sub> <sup>+</sup> -N								
6	0.005	(0.005)	0.016	(0.014)	0.015	(0.008)	0.009	(0.007)
12	0.070	(0.021)	0.090	(0.038)	0.061	(0.034)	0.085	(0.033)
18	0.157	(0.036)	0.142	(0.043)	0.093	(0.032)	0.177	(0.031)
24	0.177	(0.079)	0.276	(0.276)	0.160	(0.076)	0.266	(0.096)
30	0.092	(0.016)	0.272	(.)	0.067	(0.034)	0.435	(.)
36	0.594	(0.009)	0.211	(0.125)	0.062	(0.015)	0.187	(0.053)


Total nitrogen is given as mean percent. DON, NO<sub>3</sub><sup>-</sup>N, and NH<sub>4</sub><sup>+</sup>-N are given in mg kg. Standard errors are in parentheses unless sample size = 1. Sample size is given following SE for Total nitrogen and are consistent for all subsequent measures of nitrogen. Source is the CO<sub>2</sub> level (ambient or elevated) in which litter originated (grew) and site is the CO<sub>2</sub> level (ambient or elevated) in which litter decomposed.

from ambient CO<sub>2</sub>. In line with other studies, which have documented little or no change in leaf litter quality (Finzi *et al.*, 2001), there was no change in decomposition rates. Conversely, MD litter (Experiment II) had higher concentrations of C, hemicellulose and lignin in litter originating from elevated CO<sub>2</sub> chambers compared with that from ambient CO<sub>2</sub>. Although lignin concentrations are often a good predictor of decomposition rates (Meetenmeyer, 1978), the origin of litter and, thus, litter quality did not affect decomposition rates in Experiment II. Overall, decomposition rates were unaffected by the source (ambient or elevated) of the litter regardless of whether or not litter quality was affected.

Although a number of studies have examined decomposition of litter from various CO<sub>2</sub> concentrations, few have examined the interactions between herbivory and

CO<sub>2</sub> and their effects on nutrient cycling. Prior work on the three oak species in the scrub oak system found that herbivore-damaged litter was lower in soluble phenolic concentrations and higher in lignin concentrations and lignin : N ratios (Hall *et al.*, 2005a). Whether these effects represented choice by herbivores or induction was not determined in the current or previous studies in the scrub oak system because herbivore densities could not be manipulated experimentally in the multiple-project chambers. Whether the result of induction or not, variation in litter quality associated with herbivore activity did not affect decomposition rates.

The higher rates of decomposition and N cycling in elevated CO<sub>2</sub> chambers that we observed in this study could have been the result of differences in microclimate or biota. Elevated CO<sub>2</sub> chambers at this site have



**Fig. 5** Dynamics of C (a) and N (b) concentration during decomposition of *Quercus myrtifolia* leaf litter decomposing under ambient and elevated CO<sub>2</sub> (site) for all collection dates of MD *Q. myrtifolia* litter. Bars represent standard errors. Sample sizes varies by time and treatment and are as follows in order of time: ambient = 16, 16, 16, 13, 7, 7; elevated = 16, 16, 14, 5, 2, 16.

greater productivity (Dijkstra *et al.*, 2002) and litter cover on the chamber floors than do control chambers (Stiling *et al.*, 2003), which may increase moisture levels above those in the ambient CO<sub>2</sub> chambers. As a radiative gas, CO<sub>2</sub> addition to chambers may have resulted in an increase in temperatures in elevated CO<sub>2</sub> chambers. If changes in the microclimate occurred, then activity rates of the biota may have been altered leading to increased decomposition and N accumulation. Unfortunately, no systematic comparisons of microclimate between ambient and elevated chambers have been made. Most reports of temperature differences related to open top chambers focus on interior and exterior temperature measurements (Drake *et al.*, 1989; Whitehead *et al.*, 1995). However, a review of open top chamber design and function by Leadley & Drake (1993) found no differences in long-term air temperature between elevated and ambient CO<sub>2</sub> chambers.

Changes in detritivore populations and activity levels other than those associated with microclimatic changes could also explain the acceleration of decomposition

and N accumulation seen under elevated CO<sub>2</sub>. Previous work in the scrub oak community at our site has demonstrated that fungal biomass in the soil is higher under elevated CO<sub>2</sub> than under ambient CO<sub>2</sub> (Klamer *et al.*, 2002). In a different system, Moscatelli *et al.* (2005) have reported that microbial biomass increases by 16% in soils under elevated CO<sub>2</sub>, with an apparent shift in enzyme activity towards fungal decomposition. How might such effects influence nutrient dynamics? There are two primary hypotheses concerning N availability under elevated CO<sub>2</sub>. The first suggests that increased C availability under elevated CO<sub>2</sub> could boost microbial biomass and lead to increased immobilization of N (Diaz *et al.*, 1993). Alternatively, Zak *et al.* (1993) suggested that increased C input would result in increased microbial activity leading to increased N availability. This latter suggestion is consistent with our results and some previous work (Cotrufo *et al.*, 1999; Cotrufo *et al.*, 2005). For example, under elevated CO<sub>2</sub>, decomposing *Populus* leaves accumulate N in a manner similar to that in our experiments (Cotrufo *et al.*, 2005). In the same study, the authors report that *Populus* litter decays more rapidly under elevated CO<sub>2</sub> than under ambient CO<sub>2</sub>. However, they also found that *Populus* litter grown under elevated CO<sub>2</sub> decomposed more slowly than that grown under ambient CO<sub>2</sub>, a 'source effect' that we did not find. It may be that there exists species-specific variation in decomposition depending on the source of litter (Cotrufo *et al.*, 2005). Couiteaux *et al.* (1991) reported changes in litter quality including lower N concentrations and higher C:N ratios in chestnut litter grown under elevated CO<sub>2</sub>. Litter from elevated CO<sub>2</sub> decomposed more slowly than did litter from ambient CO<sub>2</sub>. However, when soil fauna complexity was increased, litter from elevated CO<sub>2</sub> plots displayed higher decomposition rates than did ambient CO<sub>2</sub> litter.

Our major finding is that the site of decomposition (ambient or elevated CO<sub>2</sub>) has a greater impact on litter decomposition and N cycling than does variation in initial litter quality caused by elevated CO<sub>2</sub> or herbivore activity. As such, our results add to a growing body of work that suggests that increased concentrations of CO<sub>2</sub> may generate long-term shifts in decomposition processes and rates of nutrient cycling. Such shifts have the potential to change both ecosystem structure and function.

### Acknowledgments

This research was supported by the Office of Science (BER), U.S. Department of Energy, through the South East Regional Center of the National Institute for Global Environmental Change under Cooperative Agreement No. DE-FC03-90ER61010. We thank Chris Frost and Caralyn Zehnder for comments on a previous draft of this manuscript.

## References

- Abrahamson WG, Hunter MD, Melika G *et al.* (2003) Cynipid gall-wasp communities correlate with oak chemistry. *Journal of Chemical Ecology*, **29**, 209–223.
- Ball AS, Drake BG (1997) Short-term decomposition of litter produced by plants grown in ambient and elevated atmospheric CO<sub>2</sub> concentrations. *Global Change Biology*, **3**, 29–35.
- Bate-Smith EC (1977) Astringent tannins of *Acer* species. *Phytochemistry*, **16**, 1421–1426.
- Berg B, Ekbohm G (1983) Nitrogen immobilization in decomposing needle litter at variable carbon: nitrogen ratios. *Ecology*, **64**, 63–67.
- Ceulemans R, Mousseau M (1994) Tansley review no 71: effects of elevated atmospheric CO<sub>2</sub> on woody plants. *New Phytologist*, **127**, 425–446.
- Chapman SK, Hart SC, Cobb NS *et al.* (2003) Insect herbivory increases litter quality and decomposition: an extension of the acceleration hypothesis. *Ecology*, **84**, 2867–2876.
- Cotrufo MF, Briones MJJ, Ineson P (1998) Elevated CO<sub>2</sub> affects field decomposition rate and palatability of tree leaf litter: importance of changes in substrate quality. *Soil Biology and Biochemistry*, **30**, 1564–1571.
- Cotrufo MF, De Angelis P, Polle A (2005) Leaf litter production and decomposition in a poplar short-rotation coppice exposed to free air CO<sub>2</sub> enrichment (POPFACE). *Global Change Biology*, **11**, 971–982.
- Cotrufo MF, Ineson P, Rowland AP (1994) Decomposition of tree leaf litters grown under elevated CO<sub>2</sub>: effect of litter quality. *Plant and Soil*, **163**, 121–130.
- Cotrufo MF, Raschi A, Lanini M *et al.* (1999) Decomposition and nutrient dynamics of *Quercus pubescens* leaf litter in a naturally enriched CO<sub>2</sub> Mediterranean ecosystem. *Functional Ecology*, **13**, 343–351.
- Coûteaux M-M, Mousseau M, Célérier M-L *et al.* (1991) Increased atmospheric CO<sub>2</sub> and litter quality: decomposition of sweet chestnut leaf litter with animal food webs of different complexities. *Oikos*, **61**, 54–64.
- Curtis PS, Drake BG, Whigham DF (1989) Nitrogen and carbon dynamics in C<sub>3</sub> and C<sub>4</sub> estuarine marsh plants grown under elevated CO<sub>2</sub> in situ. *Oecologia*, **78**, 297–301.
- Diaz S, Grime JP, Harris J *et al.* (1993) Evidence of a feedback mechanism limiting plant response to elevated carbon dioxide. *Nature*, **364**, 616–617.
- Dijkstra P, Hymus G, Colavito D *et al.* (2002) Elevated atmospheric CO<sub>2</sub> stimulates aboveground biomass in a fire-regenerated scrub-oak ecosystem. *Global Change Biology*, **8**, 90–103.
- Drake BG, Leadley PW, Arp WJ *et al.* (1989) An open top chamber for field studies of elevated atmospheric CO<sub>2</sub> concentration on saltmarsh vegetation. *Functional Ecology*, **3**, 363–371.
- Dukes JS, Hungate BA (2002) Elevated carbon dioxide and litter decomposition in California annual grasslands: which mechanisms matter? *Ecosystems*, **5**, 171–183.
- Fajer ED, Bowers MD, Bazzaz FA (1991) Enriched CO<sub>2</sub> atmospheres and the growth of the buckeye butterfly, *Junonia coenia*. *Ecology*, **72**, 751–754.
- Findlay S, Carreiro M, Krischik V *et al.* (1996) Effects of damage to living plants on leaf litter quality. *Ecological Applications*, **6**, 269–275.
- Finzi AD, Allen AS, DeLucia EH *et al.* (2001) Forest litter production, chemistry, and decomposition following two years of free-air CO<sub>2</sub> enrichment. *Ecology*, **82**, 470–484.
- Floate MJS (1970) Decomposition of organic materials from hill soils and pastures. II. Comparative studies on the mineralization of carbon, nitrogen and phosphorus from plant materials and sheep faeces. *Soil Biology and Biochemistry*, **2**, 173–185.
- Grace JR (1986) The influence of gypsy moth on the composition and nutrient content of litter fall in a Pennsylvania oak forest. *Forest Science*, **32**, 855–870.
- Hall MC, Stiling P, Hungate BA *et al.* (2005a) Effects of elevated CO<sub>2</sub> and herbivore damage on litter quality in a scrub oak ecosystem. *Journal of Chemical Ecology*, **31**, 2343–2356.
- Hall MC, Stiling P, Moon DC *et al.* (2005b) Effects of elevated CO<sub>2</sub> on foliar quality and herbivore damage in a scrub oak ecosystem. *Journal of Chemical Ecology*, **31**, 267–286.
- Henry HAL, Cleland EE, Field CB *et al.* (2005) Interactive effects of elevated CO<sub>2</sub>, N deposition and climate change on plant litter quality in a California annual grassland. *Oecologia*, **142**, 465–473.
- Houghton JT, Ding Y, Griggs DJ *et al.* (2001) *Climate Change 2001, the scientific basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change (IPCC)*. Cambridge University Press, UK.
- Hunter MD, Adl S, Pringle CM *et al.* (2003) Relative effects of macroinvertebrates and habitat on the chemistry of litter during decomposition. *Pedobiologia*, **47**, 101–116.
- Kemp PR, Waldecker DG, Owensby CE *et al.* (1994) Effects of elevated CO<sub>2</sub> and nitrogen fertilization pretreatments on decomposition of tallgrass prairie leaf litter. *Plant and Soil*, **165**, 115–127.
- Klamer M, Roberts MS, Levine LH *et al.* (2002) Influence of elevated CO<sub>2</sub> on the fungal community in a coastal scrub oak forest soil investigated with terminal-restriction fragment length polymorphism analysis. *Applied and Environmental Microbiology*, **68**, 4370–4376.
- Koch GW, Mooney HA (1996) *Carbon Dioxide and Terrestrial Ecosystems*. Academic Press, San Diego.
- Koroleff F (1983) Simultaneous oxidation of nitrogen and phosphorus compounds by persulfate. In: *Methods of Seawater Analysis*, 2nd edn., eds Grasshoff K, Eberhardt M, Kremling K, pp. 168–169. Verlag Chemie, Weinheimer FRG.
- Leadley PW, Drake BG (1993) Open top chambers for exposing plant canopies to elevated CO<sub>2</sub> concentration and for measuring net gas exchange. *Vegetation*, **104/105**, 3–15.
- Lincoln DE, Couvet DSN (1986) Response of an insect herbivore to host plants grown in carbon dioxide enriched atmospheres. *Oecologia*, **69**, 556–560.
- Lutze JL, Gifford RM, Adams HN (2000) Litter quality and decomposition in *Danthonia richardsonii* swards in response to CO<sub>2</sub> and nitrogen supply over four years of growth. *Global Change Biology*, **6**, 13–24.
- Meentemeyer V (1978) Macroclimate and lignin control of litter decomposition rates. *Ecology*, **59**, 465–472.
- Moscatelli MC, Lagomarsino A, De Angelis P *et al.* (2005) Seasonality of soil biological properties in a poplar plantation growing under elevated atmospheric CO<sub>2</sub>. *Applied Soil Ecology*, **30**, 162–173.

- Parsons WFJ, Lindroth RL, Bockheim JG (2004) Decomposition of *Betula papyrifera* leaf litter under the independent and interactive effects of elevated CO<sub>2</sub> and O<sub>3</sub>. *Global Change Biology*, **10**, 1666–1677.
- Qualls RG (1989) Determination of total nitrogen and phosphorus in water using persulfate oxidation: a modification for small sample volumes using the method of Koroleff (1983). In: *The biogeochemical properties of dissolved organic matter in a hardwood forest ecosystem: their influence on the retention of nitrogen, phosphorus, and carbon*. Appendix A, pp. 131–138, Ph.D. Dissertation, University of Georgia Institute of Ecology, Athens, Georgia, USA. University Microfilms, Inc., no. DEX9003448.
- Robinson CH, Michelsen A, Lee JA *et al.* (1997) Elevated atmospheric CO<sub>2</sub> affects decomposition of *Festuca vivipara* (L.) Sm. litter and roots in experiments simulating environmental change in two contrasting arctic ecosystems. *Global Change Biology*, **3**, 37–49.
- Rossiter MC, Schultz JC, Baldwin IT (1988) Relationships among defoliation, red oak phenolics, and gypsy moth growth and reproduction. *Ecology*, **69**, 267–277.
- Roth SK, Lindroth RL (1995) Elevated atmospheric CO<sub>2</sub>: effects on phytochemistry, insect performance and insect-parasitoid interactions. *Global Change Biology*, **1**, 173–182.
- SAS Institute (1998) *SAS 8.2 for Windows*. SAS Institute, Cary, NY.
- Schultz JC, Baldwin IT (1982) Oak leaf quality declines in response to defoliation by gypsy moth larvae. *Science*, **217**, 149–151.
- Seastedt TR (1988) Mass, nitrogen, and phosphorus dynamics in foliage and root detritus of tallgrass prairie. *Ecology*, **69**, 59–65.
- Stiling P, Cattell M, Moon DC *et al.* (2002) Elevated atmospheric CO<sub>2</sub> lowers herbivore abundance, but increases leaf abscission rates. *Global Change Biology*, **8**, 658–667.
- Stiling P, Moon DC, Hunter MD *et al.* (2003) Elevated CO<sub>2</sub> lowers relative and absolute herbivore density across all species of a scrub-oak forest. *Oecologia*, **134**, 82–87.
- Stiling P, Rossi AM, Hungate BA *et al.* (1999) Decreased leaf-miner abundance in elevated CO<sub>2</sub>: reduced leaf quality and increased parasitoid attack. *Ecological Applications*, **9**, 240–244.
- Swain T (1980) The importance of flavonoids and related compounds in fern taxonomy and ecology: An overview of the symposium. *Bulletin of the Torrey Botanical Club*, **107**, 113–115.
- Taylor BR, Parkinson D, Parsons WFJ (1989) Nitrogen and lignin content as predictors of litter decay rates: a microcosm test. *Ecology*, **70**, 97–104.
- Whitehead D, Hogan KP, Rogers GND *et al.* (1995) Performance of large open-top chambers for long-term field investigations of tree response to elevated carbon dioxide concentration. *Journal of Biogeography*, **22**, 307–313.
- Zak DR, Pregitzer KS, Curtis PS *et al.* (1993) Elevated atmospheric CO<sub>2</sub> and feedback between carbon and nitrogen cycles. *Plant and Soil*, **151**, 105–117.