



Mammalian Faunal Zones of
the Bridger Middle Eocene

C. LEWIS GAZIN



SMITHSONIAN CONTRIBUTIONS TO PALEOBIOLOGY • NUMBER 26

SERIAL PUBLICATIONS OF THE SMITHSONIAN INSTITUTION

The emphasis upon publications as a means of diffusing knowledge was expressed by the first Secretary of the Smithsonian Institution. In his formal plan for the Institution, Joseph Henry articulated a program that included the following statement: "It is proposed to publish a series of reports, giving an account of the new discoveries in science, and of the changes made from year to year in all branches of knowledge." This keynote of basic research has been adhered to over the years in the issuance of thousands of titles in serial publications under the Smithsonian imprint, commencing with *Smithsonian Contributions to Knowledge* in 1848 and continuing with the following active series:

Smithsonian Annals of Flight
Smithsonian Contributions to Anthropology
Smithsonian Contributions to Astrophysics
Smithsonian Contributions to Botany
Smithsonian Contributions to the Earth Sciences
Smithsonian Contributions to Paleobiology
Smithsonian Contributions to Zoology
Smithsonian Studies in History and Technology

In these series, the Institution publishes original articles and monographs dealing with the research and collections of its several museums and offices and of professional colleagues at other institutions of learning. These papers report newly acquired facts, synoptic interpretations of data, or original theory in specialized fields. These publications are distributed by mailing lists to libraries, laboratories, and other interested institutions and specialists throughout the world. Individual copies may be obtained from the Smithsonian Institution Press as long as stocks are available.

S. DILLON RIPLEY
Secretary
Smithsonian Institution

Mammalian Faunal Zones of the Bridger Middle Eocene

C. Lewis Gazin

ISSUED

JAN 20 1976



SMITHSONIAN INSTITUTION PRESS

City of Washington

1976

ABSTRACT

Gazin, C. Lewis. Mammalian Faunal Zones of the Bridger Middle Eocene. *Smithsonian Contributions to Paleobiology*, number 26, 25 pages, 1976.—The zoning arrangement of the Bridger Middle Eocene as defined by W. D. Matthew in his 1909 monograph on the Carnivora and Insectivora of the Bridger Basin included a series of stratigraphic units lettered from A to E. The type section is in the western part of the basin but correlation of the sequence in the eastern part of the basin erred in that a very large area shown by Matthew as C, or upper Bridger, is actually B, or lower Bridger. As a consequence many of the mammalian remains collected in the eastern part of the basin were attributed to the wrong horizon. This was discovered in my faunal studies and verified by Wilmot Bradley's mapping of the Sage Creek White Layer, which is the base of Bridger C or upper Bridger.

A faunal list of the Mammalia recognized in the Bridger is given with type localities and their horizons, so far as known, and the number of specimens in the National Museum of Natural History collections from each of the two divisions, lower and upper. Following this a discussion of species is given in which the evidence for any species being restricted to one or the other of the stratigraphic divisions is cited, or such information demonstrating its occurrence in both levels, if this is not indicated by the National Museum of Natural History collections (under the catalog numbers of the old United States National Museum).

The Annotated Bibliography includes references to all papers in which recognized new mammalian families, genera, and species included in the Bridger faunas are described. Also included are papers in which stratigraphic and additional or detailed information on Bridger mammals is provided, with notations as to extent of coverage, and possible errors of detail or interpretation in certain cases.

OFFICIAL PUBLICATION DATE is handstamped in a limited number of initial copies and is recorded in the Institution's annual report, *Smithsonian Year*. SI PRESS NUMBER 6094. SERIES COVER DESIGN: The trilobite *Phacops rana* Green.

Library of Congress Cataloging in Publication Data

Gazin, Charles Lewis, 1904—

Mammalian faunal zones of the Bridger middle Eocene.

(Smithsonian contributions to paleobiology ; no. 26)

Bibliography: p.

Supt. of Docs. no.: SI1.30:26

1. Mammals, Fossil. 2. Paleontology—Eocene. 3. Paleontology—The West. I, Title. II. Series:

Smithsonian Institution. Smithsonian contributions to Paleobiology ; no. 26.

QE701.S56 no. 26 [QE881] 560'.8s [569'.09787'9] 75-619267

Contents

	<i>Page</i>
Introduction	1
Bridger Mammalia	2
Species Discussion	7
Annotated Bibliography	13

Mammalian Faunal Zones of the Bridger Middle Eocene

C. Lewis Gazin

Introduction

The outstanding monograph by W. D. Matthew, published in 1909, on the Carnivora and Insectivora of the Bridger Basin included a zoning arrangement based on field work done by Matthew and Walter Granger in 1902 and 1904. They defined a sequence of stratigraphic units lettered from A to E with the type section essentially in the western part of the basin. The units were defined, except for A, by prominent and widespread ash layers. Unfortunately, in correlating the sequence on the eastern side of the basin across the higher central tableland, they erred in assigning to Bridger C a large area of badlands that actually represent Bridger B. This became evident to me in comparisons of faunal materials, particularly of *Hyopsodus*, from various localities around the basin, and was so reported (Gazin, 1959:136).

In consultation with Wilmot Bradley of the U.S. Geological Survey, who had just finished mapping the Eocene beds in a large portion of southwestern Wyoming and adjacent parts of Colorado and Utah, he showed me his field map, which included precise location of the Sage Creek White Layer of Matthew, that defines the base of Bridger C. It shows clearly the wide distribution of lower Bridger in the eastern part of the basin, which the map published by Matthew shows as Bridger C. Bradley's completed map was published as a part of a

Geological Survey Professional Paper (496-A) in 1964.

The Sage Creek White Layer forms the cap rock of the mesalike uplands of the central part of the basin, extending north from Cedar Mountain to near the Lincoln Highway. It should be further noted that there is essentially no collecting ground for upper Bridger mammals above the Sage Creek White Layer to the north of the northerly slopes of Sage Creek Mountain, Cedar Mountain, and Twin Buttes, at least as far north and east as the Green River, although much farther north the upper beds have been preserved near the Continental Fault in the Tabernacle Butte area. The Sage Creek White Layer extends between Cedar Mountain and Twin Buttes at a level very little below the saddle between these two mountains so that Bridger B forms the more widespread collecting ground around the lower slopes to the north and east of Twin Buttes. The B horizon also extends around the south side of Twin Buttes and westward toward the town of Burnt Fork, near which the Sage Creek White Layer crosses Henry's Fork. In this area Bridger B is relatively thin as it tapers rapidly to the south between the Sage Creek White Layer and the underlying Green River beds.

The Sage Creek White Layer is the most important of the white layers named by Matthew, as it defines the boundary between upper and lower Bridger. Moreover, it is essentially different than the other white layers named by Matthew, such as the Lone Tree White Layer, in that besides containing appreciable volcanic ash it has a more lacustrine appearance, particularly toward the east,

C. Lewis Gazin, Department of Paleobiology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560.

resembling in some ways the Tipton Tongue of the Green River beds. This may well account for the significant faunal change that occurs at this horizon, where little or no change is noticeable with respect to the other ash layers.

Bridger A as a distinct unit is not clearly defined as separate from Bridger B. Bridger D, defined below by the base of the Lone Tree White Layer, shows no significant faunal differences from Bridger C, and Bridger E is essentially barren of fossil mammals. As a consequence there is little of significance in the zoning of the Bridger for purposes of mammalian paleontology other than the distinction of upper and lower. This was early recognized, and in 1934 H. E. Wood in his paper on the Hyrachyidae designated the lower Bridger (A and B) as the Black's Fork member and the upper Bridger (C and D) as the Twin Buttes member. The name designating the lower Bridger is entirely appropriate. The selection of the name Twin Buttes for the upper member, however, was somewhat unfortunate because of the extent of lower Bridger beds around the lower slopes of Twin Buttes.

Because of this error in stratigraphic correlation much of the Bridger material collected by Matthew and others may be incorrectly zoned, unless the localities are in the type section in the western part of the basin and clearly designated. As a consequence I have restricted my following tabulation of the number of specimens of the various Bridger species to the materials in the National Museum of Natural History (NMNH). These include C. W. Gilmore's 1930 collection, which was made only in the type section, and my parties' collections made during 11 different field seasons between 1940 and

1969. Since most of the types were collected by parties from other museums in earlier years, their horizon is indicated by locality name, if given, as well as horizon, if this is evident. The number of specimens of each species in the National Museum of Natural History is given in the horizon columns under the acronym of the old United States National Museum (USNM).

In the classification arrangement in general I have followed the authoritative work of George Simpson (1945) but with later modifications as seem appropriate, and more detailed arrangements in the lower categories by various workers as noted below. The Marsupialia follows Simpson's study (1928). For the classification of the Insectivora I have relied on various papers by Simpson, Robinson, and McKenna in revisions of Matthew's study (1909B). The Primates and Tillodontia are arranged according to my own (1958, 1953) investigations. The Taeniodonta and Pholidota follow Simpson (1945, 1931), but the latter as modified by Emry (1970). The Rodentia are essentially as arranged by Robert Wilson (1938) and Albert Wood (1962). The Carnivora follows Matthew closely, with certain modifications I (1968) have introduced. The Condylarthra are as arranged by me (1968). The Dinocerata follows Wheeler's (1961) revision. The classification of the Perissodactyla follows the studies of several authorities, including Osborn (1929), Peterson (1919), H. E. Wood (1934), and Kitts (1957), but with modifications, or reductions in species, such as the rhinos, resulting from my own study. The Artiodactyla arrangement is based largely on my own studies, and those of Sinclair (1914).

Bridger Mammalia

(type localities in parentheses)

	Type horizon	USNM referred	
		L	U
Marsupialia			
Didelphidae Gray, 1821			
<i>Peratherium</i> Aymard, 1850			
<i>P. marsupium</i> (Troxell), 1923d	?	6	2
<i>P. innominatum</i> (Simpson), 1928b (Millersville)	L	4	3
<i>P. knighti</i> McGrew, 1959 (Tabernacle Butte)	U	-	-
Insectivora			
Palaeoryctidae(?) Simpson, 1931b			
<i>Didelphodus</i> Cope, 1882b			

	Type horizon	USNM referred	
		L	U
<i>D. altidens</i> (Marsh), 1872 (near Henry's Fork)	U	-	-
Leptictidae Gill, 1872			
<i>Hypictops</i> Gazin, 1949			
<i>H. syntaphus</i> Gazin, 1949 (north of Lone Tree)	U	-	2
<i>Palaeictops</i> Matthew, 1899			
<i>P. bridgeri</i> (Simpson), 1959b (Tabernacle Butte)	U	-	-
Erinaceidae Bonaparte, 1838			
<i>Scenopagus</i> McKenna & Simpson, 1959			
<i>S. priscus</i> (Marsh), 1872 (near Henry's Fork)	U	5	3
<i>S. edenensis</i> (McGrew), 1959 (Eden Valley)	-	4	-
<i>Centetodon</i> Marsh, 1872			
<i>C. pulcher</i> Marsh, 1872 (near Henry's Fork)	U	-	-
<i>C.(?) bacchanalis</i> (McGrew), 1959 (Tabernacle Butte)	U	-	-
Adapisoricidae Schlosser, 1887			
<i>Entomolestes</i> Matthew, 1909b			
<i>E. grangeri</i> Matthew, 1909b (Grizzly Buttes)	L	-	-
<i>Talpavus</i> Marsh 1872			
<i>T. nitidus</i> Marsh, 1872 (near Henry's Fork)	U	-	-
Nyctitheriidae Simpson, 1928a			
<i>Nyctitherium</i> Marsh, 1872			
<i>N. velox</i> Marsh, 1872 (near Henry's Fork)	U	5	3
<i>N. serotinum</i> (Marsh), 1872 (Grizzly Buttes)	L	-	-
<i>N. dasypelix</i> (Matthew), 1909b (Grizzly Buttes)	L	1	-
<i>Pontifactor</i> West, 1974			
<i>P. bestiola</i> West, 1974 (Henry's Fork Divide)	U	-	-
Pantolestidae Cope, 1884			
<i>Pantolestes</i> Cope, 1872b			
<i>P. longicaudus</i> Cope, 1872 (Black's Fork)	L	5	11
<i>P. elegans</i> (Marsh), 1872 (near Henry's Fork)	U	3	13
<i>P. natans</i> Matthew, 1909b	U?	-	-
<i>P. phocipes</i> Matthew, 1909b	U	-	-
<i>P. intermedius</i> Matthew, 1909b	L	-	-
Microsopsidae Osborn, 1892			
<i>Microsops</i> Leidy, 1872a			
<i>M. elegans</i> (Marsh), 1871a (Grizzly Buttes)	L	96	-
<i>M. annectens</i> (Marsh), 1872 (near Henry's Fork)	U	-	9
<i>M. schlosseri</i> Wortman, 1903 (Henry's Fork)	U	-	1
Apatemyidae Matthew, 1909b			
<i>Apatemys</i> Marsh, 1972			
<i>A. bellus</i> Marsh 1872 (near Henry's Fork)	U	2	1
<i>A. bellulus</i> Marsh 1872 (near Henry's Fork)	U	1	-
<i>A. rodens</i> Troxell 1923c	?	1	1
Primates			
Notharctidae Trouessart, 1879			
<i>Notharctus</i> Leidy, 1870c			
<i>N. tenebrosus</i> Leidy, 1870c (Black's Fork)	L	136	-
<i>N. robustior</i> Leidy, 1872b (Henry's Fork)	U	-	64
<i>Smilodectes</i> Wortman, 1903			
<i>S. gracilis</i> (Marsh), 1871a (Grizzly Buttes)	L	110	-
Omomyidae Gazin, 1958			
<i>Omomys</i> Leidy, 1869b			
<i>O. carteri</i> Leidy, 1869b (Fort Bridger)	L	78	8
<i>Hemiacodon</i> Marsh, 1872			
<i>H. gracilis</i> Marsh, 1872 (near Henry's Fork)	U	-	90
<i>Washakius</i> Leidy, 1873			
<i>W. insignis</i> Leidy, 1873	?	21	1

	Type horizon	USNM referred	
		L	U
Anaptomorphidae Cope, 1883			
<i>Anaptomorphus</i> Cope, 1872e			
<i>A. aemulus</i> Cope, 1872e (Ham's Fork)	L	-	-
<i>A. wortmani</i> Gazin, 1958	?	-	-
<i>Uintanius</i> Matthew, 1915			
<i>U. ameghini</i> (Wortman), 1904	?	2	-
<i>Uintasorex</i> Matthew, 1909b			
<i>U. parvulus</i> Matthew, 1909b (Henry's Fork)	U	-	-
<i>Trogolemur</i> Matthew, 1909b			
<i>T. myodes</i> Matthew, 1909b (south of Granger)	L	-	-
Tillodontia			
Esthonychidae Cope, 1883			
<i>Trogosus</i> Leidy, 1871b			
<i>T. castoridens</i> Leidy, 1871b (Grizzly Buttes)	L	-	-
<i>T. hyracoides</i> (Marsh), 1873b (Granger Station)	L	2	-
<i>T. latidens</i> (Marsh), 1874b	?	-	-
<i>Tillodon</i> Gazin, 1953			
<i>T. fodiens</i> (Marsh), 1875b (Millersville)	L	2	-
Taeniodontia			
Stylinodontidae Marsh, 1875a			
<i>Stylinodon</i> Marsh, 1874b			
<i>S. mirus</i> Marsh, 1874b	?	-	1
Pholidota			
Metacheiromyidae Wortman, 1903			
<i>Metacheiromys</i> Wortman, 1903			
<i>M. marshi</i> Wortman, 1903 (Grizzly Buttes?)	L?	1	-
<i>M. tatusia</i> Osborn, 1904 (Grizzly Buttes)	L	-	-
<i>M. dasyptus</i> Osborn, 1904 (Grizzly Buttes)	L	-	-
<i>M. osborni</i> Simpson, 1931a (Lone Tree)	U	-	-
Epoicotheriidae Simpson, 1927			
<i>Tetrapassalus</i> Simpson, 1959a			
<i>T. mckennai</i> Simpson, 1959a (Tabernacle Butte)	U	-	-
<i>T. proius</i> West, 1973c (Sage Creek)	L	-	-
Rodentia			
Paramyidae Miller and Gidley, 1918			
<i>Paramys</i> Leidy, 1871d			
<i>P. delicatus</i> Leidy, 1871d (near Fort Bridger)	L	114	16
<i>P. delicatior</i> Leidy, 1871d (near Fort Bridger)	L	40	18
<i>Thisbemys</i> A. E. Wood, 1959b			
<i>T. corrugatus</i> A. E. Wood, 1959b (north of Cedar Mt.)	L	7	68
<i>T. plicatus</i> A. E. Wood, 1962 (Grizzly Buttes)	L	51	-
<i>Leptotomus</i> Matthew, 1910			
<i>L. parvus</i> A. E. Wood, 1959b (Grizzly Buttes)	L	39	-
<i>L. bridgerensis</i> A. E. Wood, 1962 (Henry's Fork)	U	-	7
<i>Microparamys</i> A. E. Wood, 1959b			
<i>M. minutus</i> (R. W. Wilson), 1937 (Millersville?)	L?	-	-
<i>M. wilsoni</i> A. E. Wood, 1962	U?	-	-
<i>M. wyomingensis</i> A. E. Wood, 1959b (Tabernacle Butte)	U	-	-
<i>Reithroparamys</i> Matthew, 1920			
<i>R. delicatissimus</i> (Leidy), 1871d (Grizzly Buttes)	L	17	-
<i>R. matthewi</i> A. E. Wood, 1962 (Henry's Fork)	U	-	-
<i>Pseudotomus</i> Cope, 1872b			
<i>P. hians</i> Cope, 1872b (Cottonwood Creek)	L	-	-
<i>P. robustus</i> (Marsh), 1872 (Grizzly Buttes)	L	-	2
<i>Ischyrotomus</i> Matthew, 1910			
<i>I. horribilis</i> A. E. Wood, 1962 (near Grizzly Buttes)	L	8	-

	Type horizon	USNM referred	
		L	U
<i>I. oweni</i> A. E. Wood, 1962 (north of Twin Buttes)	L	2	-
<i>I. superbus</i> (Osborn, Scott, and Speir), 1878 (Cottonwood Creek)	L	-	-
Cylindrodontidae Miller and Gidley, 1918			
<i>Mysops</i> Leidy, 1871d			
<i>M. minimus</i> Leidy, 1871d (Grizzly Buttes)	L	21	-
<i>M. parvus</i> (Marsh), 1872 (Grizzly Buttes)	L	44	4
<i>M. fraternus</i> Leidy, 1873	?	6	1
Sciuravidae Miller and Gidley, 1918			
<i>Sciuravus</i> Marsh, 1871b			
<i>S. nitidus</i> Marsh, 1871b (Grizzly Buttes)	L	175	3
<i>S. bridgeri</i> R. W. Wilson, 1938 (Millersville)	L	7	-
<i>S.(?) rarus</i> R. W. Wilson, 1938	?	-	-
<i>Tillomys</i> Marsh, 1872			
<i>T. senex</i> Marsh, 1872 (near Henry's Fork)	U	-	-
<i>T.(?) parvidens</i> (Marsh), 1872 (Grizzly Buttes?)	L	3	-
<i>Taxymys</i> Marsh, 1872			
<i>T. lucaris</i> Marsh, 1872 (near Henry's Fork)	U	1	2
<i>T.(?) progressus</i> R. W. Wilson, 1938	?	-	-
<i>Pauromys</i> Troxell, 1923a			
<i>P. perditus</i> Troxell, 1923a (Dry Creek)	L?	-	-
<i>P. schaubi</i> A. E. Wood, 1959a (Twin Buttes (high))	U	-	-
Carnivora, Creodonta			
Mesonychidae Cope, 1875			
<i>Mesonyx</i> Cope, 1872a			
<i>M. obtusidens</i> Cope, 1872a (Cottonwood Creek)	L	1	1
<i>Synoplotherium</i> Cope, 1872d			
<i>S. vorax</i> (Marsh), 1876c (Henry's Fork)	U	-	-
<i>Harpagolestes</i> Wortman, 1901			
<i>H. macrocephalus</i> Wortman, 1901 (Smith's Fork)	L	1	-
Oxyaenidae Cope, 1877			
<i>Patriofelis</i> Leidy, 1870a			
<i>P. ulta</i> Leidy, 1870a (Fort Bridger)	L	3	-
<i>P. ferox</i> (Marsh), 1872 (near Henry's Fork)	U	-	3
Hyaenodontidae Leidy, 1869a			
<i>Sinopa</i> Leidy, 1871b			
<i>S. rapax</i> Leidy, 1871b (Grizzly Buttes)	L	21	6
<i>S. pungens</i> (Cope), 1872b (Cottonwood Creek)	L	5	2
<i>S. major</i> Wortman, 1902 (Church Buttes?)	L?	-	3
<i>S. minor</i> Wortman, 1902 (Grizzly Buttes)	L	16	-
<i>S. grangeri</i> Matthew, 1906 (Cottonwood Creek)	L	4	-
<i>Tritemnodon</i> Matthew, 1906			
<i>T. agilis</i> (Marsh), 1872 (Grizzly Buttes)	L	13	-
Limnocyoniidae Gazin, 1946			
<i>Limnocyon</i> (Marsh), 1872			
<i>L. verus</i> Marsh, 1872 (Grizzly Buttes)	L	17	-
<i>Thinocyon</i> Marsh, 1872			
<i>T. velox</i> Marsh, 1872 (Grizzly Buttes)	L	25	-
<i>T. medius</i> (Wortman), 1902 (near Henry's Fork)	U	-	6
<i>T. mustelinus</i> Matthew, 1909b	L	15	-
<i>Machaeroides</i> Matthew, 1909b			
<i>M. eothen</i> Matthew, 1909b	L	2	-
Carnivora, Fissipeda			
Miacidae Cope, 1880			
<i>Viverravus</i> Marsh, 1872			
<i>V. gracilis</i> Marsh, 1872 (Grizzly Buttes)	L	11	5
<i>V. minutus</i> Wortman, 1901 (Dry Creek)	L	3	-

	Type horizon	USNM referred	
		L	U
<i>V. sicarius</i> Matthew, 1909b	L	2	-
<i>Miacis</i> Cope, 1872c			
<i>M. parvivorus</i> Cope, 1872c (Black's Fork)	L	18	-
<i>M. sylvestris</i> (Marsh), 1872 (near Henry's Fork)	U	-	1
<i>M. harger</i> (Wortman), 1901 (Henry's Fork)	U	-	-
<i>Oödetes</i> Wortman, 1901			
<i>O. herpestoides</i> Wortman, 1901	L?	2	-
<i>O. proximus</i> Matthew, 1909b (Grizzly Buttes)	L	-	-
<i>Uintacyon</i> Leidy, 1872d			
<i>U. edax</i> Leidy, 1872d	?	2	-
<i>U. vorax</i> Leidy, 1872d (Henry's Fork?)	U?	2	-
<i>U. jugulans</i> Matthew, 1909b (Henry's Fork)	U	-	3
<i>U. major</i> Matthew, 1909b	U?	-	-
<i>Vulpavus</i> Marsh, 1871b			
<i>V. palustris</i> Marsh, 1871b (near Fort Bridger)	L	2	-
<i>V. profectus</i> Matthew, 1909b (Grizzly Buttes)	L	5	-
<i>V. ovatus</i> Matthew, 1909b (Little Dry Creek)	L	-	-
<i>Palaearctonyx</i> Matthew, 1909b			
<i>P. meadi</i> Matthew, 1909b (Base of Twin Buttes)	L	-	-
Condylarthra			
Hyopsodontidae Lydekker, 1889			
<i>Hyopsodus</i> Leidy, 1870b			
<i>H. paulus</i> Leidy, 1870b (near Fort Bridger)	L	868	-
<i>H. minusculus</i> Leidy, 1873 (Little Dry Creek)	L	80	-
<i>H. marshi</i> Osborn, 1902 (Twin Buttes?)	U?	-	4
<i>H. despiciens</i> Matthew, 1909b (Henry's Fork)	U	-	162
<i>H. lepidus</i> Matthew, 1909b (Henry's Fork)	U	-	115
Dinocerata			
Uintatheriidae Flower, 1876			
<i>Uintatherium</i> Leidy, 1872c			
<i>U. anceps</i> (Marsh), 1871a (Sage Creek Divide)	U	-	7
<i>Bathyopsis</i> Cope, 1881			
<i>B. middleswarti</i> Wheeler, 1961 (near Sandy Creek)	L	-	-
Perissodactyla			
Equidae Gray, 1821			
<i>Orohippus</i> Marsh, 1872			
<i>O. pumilus</i> (Marsh), 1871a (near Marsh's Fork)	L	56	-
<i>O. agilis</i> Marsh, 1873b (Henry's Fork)	U	-	-
<i>O. major</i> Marsh, 1874a (Millersville)	L	9	-
<i>O. sylvaticus</i> (Leidy), 1870d (Henry's Fork)	U	-	9
<i>O. progressus</i> Granger, 1908 (Twin Buttes)	?	11	8
Brontotheriidae Marsh, 1873b			
<i>Palaeosyops</i> Leidy, 1870c			
<i>P. paludosus</i> Leidy, 1870c (Church Buttes)	L	23	-
<i>P. major</i> Leidy, 1871c (Grizzly Buttes)	L	7	-
<i>P. robustus</i> (Marsh), 1872 (near Henry's Fork)	U	-	17
<i>Limnohyops</i> Marsh, 1890			
<i>L. laticeps</i> (Marsh), 1872 (Marsh's Fork)	L?	-	-
<i>L. priscus</i> Osborn, 1908 (Grizzly Buttes)	L	1	-
<i>Telmatherium</i> Marsh, 1872			
<i>T. validum</i> Marsh, 1872 (near Henry's Fork)	U	-	2
<i>T. cultridens</i> (Osborn, Scott and Speir), 1878 (Henry's Fork Divide)	U	-	7
Mantoceras Hatcher, 1895			
<i>M. manteoceras</i> (Osborn), 1899 (Washakie)		-	7
<i>Mesatirhinus</i> Osborn, 1908a			
<i>M. megarhinus</i> (Earle), 1891a (Washakie)		-	5
<i>M. petersoni</i> Osborn, 1908a (Cat-tail Spring)	U	-	2

	Type horizon	USNM referred	
		L	U
Isectolophidae Peterson, 1919			
<i>Parisectolophus</i> Peterson, 1919			
<i>P. latidens</i> (Osborn, Scott, and Speir), 1878 (Henry's Fork)	U	8	3
Helaletidae Osborn, 1892			
<i>Helaletes</i> Marsh, 1872			
<i>H. nanus</i> (Marsh), 1871a (Grizzly Buttes)	L	89	4
<i>H. intermedius</i> (Osborn, Scott, and Speir), 1878 (Henry's Fork)	U	-	-
<i>Dilophodon</i> Scott, 1883			
<i>D. minusculus</i> Scott, 1883	?	-	2
Hyrachyidae H. E. Wood, 1927			
<i>Hyrachyus</i> Leidy, 1871a			
<i>H. modestus</i> (Leidy), 1870b (Smith's Fork)	L	55	2
<i>H. affinis</i> (Marsh), 1871a (near Marsh's Fork)	L	48	21
<i>H. eximius</i> Leidy, 1871c (Henry's Fork)	U	1	7
<i>H. princeps</i> Marsh, 1872 (near Henry's Fork)	U	-	3
Artiodactyla			
Dichobunidae Gill, 1872			
<i>Microsus</i> Leidy, 1870c			
<i>M. cuspidatus</i> Leidy, 1870c (Black's Fork)	L	12	3
<i>Homacodon</i> Marsh, 1872			
<i>H. vagans</i> Marsh, 1872 (near Henry's Fork)	U	-	4
<i>Antiacodon</i> Marsh, 1872			
<i>A. venustus</i> Marsh, 1872 (near Henry's Fork)	U	-	2
<i>A. pygmaeus</i> (Cope), 1872a (Cottonwood Creek)	L	20	-
<i>A. furcatus</i> (Cope), 1873a (Bluffs of upper-Green River)	?	-	-
<i>Helohyus</i> Marsh, 1872			
<i>H. lentus</i> (Marsh), 1871 (Henry's Fork)	U	-	1
<i>H. plicodon</i> Marsh, 1872 (Grizzly Buttes)	L	6	4
<i>H. validus</i> (Marsh), 1872 (near Henry's Fork)	U	-	-
<i>H. milleri</i> Sinclair, 1914 (Henry's Fork)	U	-	1
<i>Lophiohyus</i> Sinclair, 1914			
<i>L. alticeps</i> Sinclair, 1914 (Twin Buttes)	?	-	-
<i>Neodiapodexis</i> Atkins, 1970			
<i>N. emryi</i> Atkins, 1970 (Tabernacle Butte)	U	-	-

Species Discussion

The following discussion by species pertains to those forms that appear to be indicative of one or the other horizons, or where the evidence indicated in the foregoing table is inconclusive. Those species represented in the National Museum of Natural History collection by specimens from both horizons as shown above are not included. Otherwise, the sequential arrangement corresponds to the foregoing tabulation. (NMNH = National Museum of Natural History; AMNH = American Museum of Natural History.)

Peratherium knighti McGrew: The recorded occurrence of this species is so far only the upper Bridger at Tabernacle Butte.

Didelphodus altidens (Marsh): This species, rep-

resented by only a few specimens, seems from the evidence furnished by McKenna (see McKenna et al., 1962) to be upper Bridgerian.

Hypictops syntaphus Gazin: The type and one other known specimen are from the upper Bridger north of Lone Tree.

Palaeictops bridgeri (Simpson): The type from the upper Bridger is the only known specimen.

Scenopagus edenensis (McGrew): Although referred specimens, including the type of *Scenopagus mcgrewi*, are from the upper Bridger beds at Tabernacle Butte, the type of *S. edenensis* is from nearby Green River beds of lower Bridger equivalence.

Centetodon pulcher Marsh: The type from near Henry's Fork and referred material from Tabernacle Butte are upper Bridgerian, but McKenna

et al. (1962) referred to the species *Hypacodon praecursor*, the type of which is from Fort Bridger and may well be lower Bridger. Other Yale specimens of *C. pulcher* were not cited as to horizon.

Centetodon (?) *bacchanalis* (McGrew): The type and referred material of this species are recorded only from the upper Bridger at Tabernacle Butte.

Entomolestes grangeri Matthew: In Matthew's (1909b) description the type and several referred jaws were stated to be from the lower Bridger but in his faunal list he has the referred material as coming from C? and D. Robinson (1968) states that the type is the only specimen known to him.

Talpavus nitidus Marsh: Marsh cites several specimens but the locality, near Henry's Fork, is given for the type only. Robinson (1968) gives the locality for a referred specimen in the Yale collections as Dry Creek. The upper reaches of Dry Creek are in the upper Bridger but a large area drained by this creek on the east side of the Bridger Basin is lower Bridger, the area indicated by Matthew (1909b) as C.

Nyctitherium serotinum (Marsh): This species may well be restricted to the lower Bridger, but one of the specimens in the Yale collections, according to Robinson (1968), has no locality data.

Nyctitherium dasypelix (Matthew): Both the type and the one known referred specimen are from the lower Bridger at Grizzly Buttes.

Pontifactor bestiola West: The type is from the upper Bridger but the 26 referred specimens in the AMNH collections are stated by West (1974) to be from the lower Bridger. I suspect that the family represented is not the Nyctitheriidae but possibly the Mixodectidae.

Pantolestes natans Matthew: This species was regarded by Matthew as upper Bridgerian, but no localities are given. If either the type or referred material cited as C came from the eastern part of the basin, both horizons may be represented.

Pantolestes phocipes Matthew: The type is listed as Bridger D so is surely upper Bridger, but referred material is listed as C and D. If the C material is from the eastern side of the basin it may well be lower Bridger.

Pantolestes intermedius Matthew: Localities not given but listed only as B.

Microsycopops elegans (Marsh): This species is well represented and appears to be restricted to the lower Bridger.

Microsycopops annectens (Marsh): A good number of specimens of this species have been found, all listed as C or D. Wortman (1903) indicated that in the Marsh collection all are from Henry's Fork. Matthew (1909b) lists referred material in the AMNH as from C or D. While some of these might have come from beds on the east side of the basin mislabeled C, it seems improbable as there is no record of an occurrence in the lower beds on the west side, considering the number of specimens found elsewhere.

Microsycopops schlosseri Wortman: Wortman (1903) stated that the type and referred specimens in the Marsh collection are all from Henry's Fork. Matthew's (1909b) listing indicates C and D for referred materials in the AMNH. While probably restricted to the upper beds, Matthew's use of C leaves a doubt.

Notharctus tenebrosus Leidy: Robinson's (1957) work as well as mine (1958) has shown that *Notharctus tenebrosus* is limited to the lower Bridger.

Notharctus robustior Leidy: This species is characteristic of the upper Bridger and equivalent beds as shown by Robinson (1957) and by me (1958).

Smilodectes gracilis (Marsh): This species is actually limited to the lower Bridger and Green River equivalent. Robinson's (1957) including it in the upper Bridger is based on its frequent occurrence around Twin Buttes and elsewhere on the east side of the basin in beds of lower Bridger age, not C as indicated in Matthew's (1909b) map. The chart shown in my (1958) report is also in error for the same reason.

Hemiacodon gracilis Marsh: So far as known this species is not found in the lower Bridger.

Washakius insignis Leidy: This species is well represented in the lower Bridger, but is scarce in the upper beds. Simpson (in McGrew et al., 1959) has recorded it from the upper beds at Tabernacle Butte.

Anaptomorphus aemulus Cope: A comparatively rare species, but so far recorded only from the lower Bridger.

Anaptomorphus wortmani Gazin: One specimen is known to be lower Bridger, but the type and two others are from unknown localities in the Bridger.

Uintanius ameghini (Wortman): Remains of this species are rare but have been found in both upper and lower beds.

Uintasorex parvulus Matthew: In the Bridger

Basin this small species, though uncommon, appears restricted to the upper beds.

Trogolemur myodes Matthew: This very rare species may be limited to the lower Bridger, but tentative recognition of the genus in the upper Eocene at Badwater Creek makes this seem improbable.

Trogosus castoridens Leidy: No record later than lower Bridger.

Trogosus hyracoides (Marsh): No record later than lower Bridger.

Trogosus latidens (Marsh): Locality and horizon for the type not known, but well represented in the lower Bridger of the northern part of the basin, also recorded in the upper part of the Cathedral Bluffs tongue.

Tillodon fodiens (Marsh): Known only in the lower Bridger.

Stylinodon mirus Marsh: The locality for the type is not known. One specimen in the NMNH is from the upper Bridger; however, undetermined stylinodont material is known from the lower Bridger.

Metacheiromys marshi Wortman: This species may well be lower Bridger, although two specimens appear uncertain. Simpson (1931a) cites the type as "probably Bridger B at Grizzly Buttes." A referred specimen cited as "probably Bridger C, Dry Creek," may well be from the lower Bridger of Dry Creek in the area shown by Matthew (1909b) as C.

Metacheiromys tatusia Osborn: Possibly a lower Bridger form, but one of the three described specimens (see Simpson 1931a) is from an unknown locality.

Metacheiromys dasypus Osborn: Possibly a lower Bridger form, but the only referred specimen (Yale collection) is from an unknown locality, regarded by Simpson (1931a) as probably B.

Metacheiromys osborni Simpson: The only described specimen is from the upper Bridger.

Tetrapassalus mckennai Simpson: The type from the upper Bridger at Tabernacle Butte is the only known specimen.

Tetrapassalus proius West: The type from the lower Bridger along Sage Creek is the only known specimen.

Thisbemys plicatus A. E. Wood: All the known material is lower Bridger.

Leptotomus parvus A. E. Wood: Known only from the lower Bridger and beds of equivalent age.

Leptotomus bridgerensis A. E. Wood: Restricted to the upper Bridger and beds of equivalent age in the Washakie Basin.

Microparamys minutus R. W. Wilson: Type is believed to be from the lower Bridger but referred specimens are known from both levels.

Microparamys wilsoni A. E. Wood: Type and referred specimen are stated by Wood (1962) to be from the upper Bridger but no localities are given.

Microparamys wyomingensis A. E. Wood: Described from material found in the upper Bridger at Tabernacle Butte, but considered by West (1969a) to be represented in the lower Bridger as well.

Reithroparamys delicatissimus (Leidy): Known in the upper Bridger as well as the lower according to A. E. Wood (1962).

Reithroparamys matthewi A. E. Wood: Although the type is upper Bridger, referred material is from unknown locality.

Pseudotomus hians Cope: Although Matthew (1909b) lists referred material from upper and lower Bridger, A. E. Wood (1962) recognizes only the type which is from Cottonwood Creek and hence probably lower Bridger.

Ischyrotomus horribilis A. E. Wood: Type and referred specimens are from the lower Bridger, as represented in NMNH.

Ischyrotomus oweni A. E. Wood: Type and referred specimen in NMNH are cited by Wood (1962) as upper Bridger north of Twin Buttes. The locality is actually in Bridger B. The Princeton specimen cited is from the upper Bridger.

Ischyrotomus superbus (Osborn, Scott, and Speir): Although Matthew (1909b) lists this species as occurring in both upper and lower Bridger, A. E. Wood (1962) recognizes only the type in the hypodigm, which is from the lower beds along Cottonwood Creek.

Mysops minimus Leidy: This species may represent the lower Bridger only; however, Matthew (1909b) as with some other forms has it listed as represented throughout the Bridger. R. W. Wilson (1938) cites one specimen at Yale from an unknown locality.

Sciuravus bridgeri R. W. Wilson: All known specimens are apparently from the lower Bridger, as noted by R. W. Wilson (1938).

Sciuravus (?) *rarus* R. W. Wilson: Apparently

known only from the type specimen, which is from an unknown Bridger locality.

Tillomys senex Marsh: Found at Henry's Fork and Tabernacle Butte, but apparently not known from the lower Bridger.

Tillomys (?) *parvidens* (Marsh): According to R. W. Wilson (1938), material of this species is found in both upper and lower Bridger.

Taxymys (?) *progressus* R. W. Wilson: The type, evidently the only known specimen, is from an unknown Bridger locality.

Pauromys perditus Troxell: The type of this species, coming from Dry Creek, could be either upper or lower but I suspect it is lower because of the vastly greater area of lower Bridger involved in the Dry Creek drainage on the east side of the basin.

Pauromys schaubi A. E. Wood: The type, the only known specimen, is upper Bridgerian, as the red stratum at Twin Buttes is above the Sage Creek White Layer.

Synoplotherium vorax (Marsh): Apparently represented by only the type specimen, which is from the upper Bridger.

Harpagolestes macrocephalus Wortman: The type and referred specimen in the NMNH are from the lower Bridger. No others are recorded.

Patriofelis ulta Leidy: The type and referred specimens are all from the lower Bridger. One specimen stated by me (1957) to be from low in Bridger C north of Twin Buttes is actually from Bridger B.

Patriofelis ferox (Marsh): The type is from Henry's Fork and Matthew (1909b) cites all referred Bridger material in the AMNH as from C or D although no localities are given. It should be noted, however, that Wortman (1902), Matthew (1909b), and Thorpe (1923) list *Limnofelis latidens* Marsh as a synonym, the type of which comes from Grizzly Buttes.

Sinopa major Wortman: The type of this species is stated by Wortman to be from Church Buttes. This is doubted by Matthew (1909b) as all AMNH referred material is from Bridger D. The NMNH material is also upper Bridger.

Sinopa minor Wortman: The Yale type and referred material in both the AMNH and NMNH are from the lower Bridger. All but one specimen in the Yale collections are from the lower Bridger. The exception is stated by Wortman (1902) to be doubtfully from Henry's Fork.

Sinopa grangeri Matthew: The type and referred materials in the NMNH are all from the lower Bridger. Apparently no other specimens have been recorded.

Tritemnodon agilis (Marsh): The type and co-type at Yale were found at Grizzly Buttes and Church Buttes. Referred material in the AMNH (Matthew, 1909b) and the NMNH are all from the lower Bridger.

Limnocyon verus Marsh: Although the Yale type, the NMNH referred material, and the type of the synonym *Limnocyon riparius* are from the lower Bridger, Matthew (1909b) cites upper as well as lower Bridger specimens in the AMNH collections; however, he suspects that the upper Bridger material may represent a distinct species.

Thinocyon velox Marsh: Evidently all material of this species in the Yale collections (see Wortman, 1902, and Thorpe, 1923) and the AMNH as well as the NMNH is from the lower Bridger.

Thinocyon medius (Wortman): This may be an upper Bridger species as Matthew (1909b) cites it as represented in C and D but gives no localities. Thorpe (1923) and Denison (1938) also regard it as upper Bridger, but Wortman (1902), while citing the type as from Henry's Fork, says that other specimens were found in the lower part of the deposits. It is possible that Wortman's referred specimens are of a different species.

Thinocyon mustelinus Matthew: Matthew (1909b:300), while erroneously listing this species as *Thinocyon "minusus,"* cites the type as lower Bridger but gives no locality. Denison (1938) also cites the species as lower Bridger, as are the NMNH specimens.

Machaeroides eothen Matthew: This species is relatively rare but the type and paratype are from the lower Bridger, as are two NMNH specimens; however, Matthew (1909b) lists a referred tooth from the upper Bridger but no locality is given. Denison (1938) regards the latter as an advanced variant.

Viverravus minutus Wortman: This is evidently lower Bridger, although Matthew (1909a) lists the species from both upper and lower Bridger, possibly because the type was from Dry Creek. Later (1909b) he lists the type and all AMNH referred specimens from Bridger B. Wortman (1901) lists Yale referred specimens from Grizzly Buttes.

Viverravus sicarius Matthew: Matthew (1909b)

cites all specimens, including the type, as from lower Bridger. Thorpe (1923) notes that specimens of this species in the Yale collections are lower Bridger in age. Those in the NMNH are also lower Bridger.

Miacis parvivorus Cope: The type and all referred specimens in the AMNH, according to Matthew (1909b), are from the lower Bridger. This is true of the NMNH materials also. Thorpe (1923) recognized 8 specimens in the Yale collections but no horizon is mentioned.

Miacis sylvestris (Marsh): Although the Yale type is from the upper Bridger, Matthew (1909b) noted that one of two referred specimens in the AMNH collections was from high in Bridger B.

Miacis hargerii (Wortman): Most specimens were noted by Thorpe (1923) to be from the upper Bridger, including the type, but some he noted were from Grizzly Buttes, Millersville, and Church Buttes.

Oödictes herpestoides Wortman: No locality or horizon is given for the type by Wortman (1901), nor is the horizon cited by Matthew (1909b), although earlier Matthew (1909a) lists the species horizon as Bridger B. Thorpe (1923) notes referred material from Millersville, which is Bridger B.

Oödictes proximus Matthew: Evidently known only from the type, which is from the lower Bridger at Grizzly Buttes.

Uintacyon edax Leidy: The locality and horizon for the type is not known but referred materials in both the Yale (Thorpe, 1923) and the NMNH collections are from the lower Bridger.

Uintacyon vorax Leidy: Although the type was recorded by Leidy as from Henry's Fork, Matthew (1909b) questions this and lists AMNH specimens as from the lower Bridger, as are the NMNH specimens. Wortman (1901) also records a Yale specimen from the lower beds, but Thorpe (1923) cites several Yale specimens without locality data.

Uintacyon jugulans Matthew: Matthew (1909b) cites type and referred materials as from the upper Bridger, as are the NMNH specimens. It should be noted, however, that a Yale specimen from the lower Bridger, thought by Wortman (1901) to represent *U. vorax*, is regarded by Thorpe (1923) as being *U. jugulans*.

Uintacyon major Matthew: Matthew (1909b) states that all AMNH specimens came from the upper Bridger, but no localities are given. Thorpe

(1923) refers a lower Bridger specimen to this species.

Vulpavus palustris Marsh: This would appear to be a lower Bridger species as the Yale type as well as referred material in the AMNH (Matthew, 1909b) and the NMNH are from the lower horizon.

Vulpavus prosectus Matthew: AMNH type and referred material are lower Bridger, as are the NMNH specimens.

Vulpavus ovatus Matthew: Matthew (1909b) lists referred material as well as the type from the lower Bridger. It is also listed from the lower Bridger by McGrew (1970).

Palaeoarctonyx meadi Matthew: The type from the base of Twin Buttes is Bridger B, not C as Matthew (1909b) has listed it. Matthew states that referred material is from Henry's Fork.

Hyopsodus paulus Leidy: I (1968) find statistically that this species is characteristic of the lower Bridger.

Hyopsodus minusculus Leidy: This species is limited to the lower Bridger and is most characteristic of the localities from Millersville to the north (Gazin, 1968).

Hyopsodus marshi Osborn: This very large form appears from my (1968) statistical study to be typical of relatively late upper Bridgerian. The type is stated by Matthew (1909b) as either Twin Buttes or Henry's Fork. He reports that referred specimens are from C₂ of Twin Buttes, which would be Bridger B. I question Matthew's referring these to *H. marshi* as they may well represent large variants of *H. paulus*.

Hyopsodus despiciens Matthew: Characteristic of the upper Bridger according to my (1968) statistical study, and so regarded by Matthew (1909b).

Hyopsodus lepidus Matthew: The relatively small form appears characteristic of the upper Bridger. This is according to my (1968) study as well as Matthew's (1909b).

Uintatherium anceps (Marsh): This species as summarized by Wheeler (1961) is characteristic of the upper Bridger and beds of equivalent age in the Washakie Basin.

Bathyopsis middleswarti Wheeler: This species is characteristic only of the lower Bridger and appears concentrated in the more northerly areas.

Orohippus pumilus (Marsh): Kitts' (1957) review of this species shows it to be limited to the lower part of the Bridger.

Orohippus agilis Marsh: While of rather limited representation, this species appears to have been found only in the upper Bridger (Kitts, 1957). Matthew (1909b), however, erroneously lists all species of *Orohippus* as having been found in B, C, and D.

Orohippus major Marsh: Kitts cites the hypodigm of Yale and AMNH specimens as lower Bridger. This is true also of the NMNH specimens.

Orohippus sylvaticus (Leidy): The statement by Kitts (1957) covering the Yale materials as well as the NMNH collection shows this species to be limited to upper Bridger. Matthew (1909a) lists this species as only upper Bridger, and later (1909b), evidently an error, lists referred material in the AMNH as from all levels.

Palaeosyops paludosus Leidy: This titanotheres is evidently lower Bridger only, as indicated by Osborn (1929) as well as by the NMNH material.

Palaeosyops major Leidy: This species also appears to be limited to the lower Bridger according to Osborn (1929), as well as by the NMNH materials.

Palaeosyops robustus (Marsh): Limited to the upper Bridger as indicated by Osborn (1929). I have included here also the material of *P. leidyi*, which I cannot distinguish from this species. It is likewise upper Bridger.

Limnohyops laticeps (Marsh): The horizon for the type specimen is not known, but Osborn (1929) cites referred material as from the upper Bridger only.

Limnohyops priscus Osborn: Osborn (1929) cites type and a Princeton referred specimen as lower Bridger. A single specimen in the NMNH is also lower Bridger.

Telmatherium validum Marsh: Osborn recognized only the Yale type as belonging to this species. This specimen as well as the NMNH material is upper Bridger in age.

Telmatherium cultridens (Osborn, Scott, and Speir): This species may well be upper Bridger, but it should be noted that certain specimens cited by Osborn (1929) as Bridger C, or no level given, and one from Twin Buttes are actually uncertain as to horizon. It is not recognized, however, in definitely lower Bridger collections.

Manteoceras manteoceras (Osborn): This species is fairly well represented and appears to be limited to the upper Bridger and Washakie, as indicated by

Osborn (1929). It has not been reported from Bridger B.

Mesatirhinus megarhinus (Earle): A Washakie species, this form in the Bridger appears to be characteristic of the C₅ level according to Osborn (1929), although two specimens are from unknown levels. The NMNH specimens are all upper Bridger.

Mesatirhinus petersoni Osborn: All specimens are cited by Osborn (1929) as upper Bridger or Washakie. One specimen listed only as Bridger C₂ may be in question; however, there is no record of the species in the lower Bridger.

Helaletes nanus (Marsh): Recognized in both upper and lower levels of the Bridger. I have included here *Helaletes boops*, which appears to be a variant recognized only on upper premolars.

Helaletes intermedius (Osborn, Scott, and Speir): If Bridgerian, this relatively rare form would be from the upper level as listed by Matthew (1909a).

Dilophodon minusculus Scott: Evidently upper Bridger as well as Washakie in age, although the type locality is unknown. It has not been recorded from the lower Bridger.

Hyrachyus eximius Leidy: This species would appear to be limited to the upper Bridger, as indicated by H. E. Wood (1934), although I have referred one specimen to it that I found at the top of Bridger B, just below the Sage Creek White Layer to the north of Cedar Mountain.

Hyrachyus princeps Marsh: Although Troxell (1922b) cites the type as coming from Fort Bridger, Marsh states that he collected it himself near Henry's Fork. An upper Bridger species, as all the referred material, as noted by H. E. Wood (1934), is upper Bridger in age.

Homacodon vagans Marsh: The type and a plesio-type (after Sinclair, 1914) are both from the upper Bridger, as well as specimens in the NMNH. Matthew (1909a, b), however, lists the form as being both upper and lower Bridger.

Antiacodon venustus Marsh: The type and NMNH referred specimens are from the upper Bridger. Matthew (1909a) lists it only as upper Bridger.

Antiacodon pygmaeus (Cope): Matthew (1909b) lists this species as lower Bridger for both type and referred material in the AMNH, as are referred specimens in the NMNH.

Antiacodon furcatus (Cope): The horizon for the

type of this species is not certainly known but it may be low in Bridger B. Matthew (1909a) lists the species as lower Bridger, but later (1909b) leaves the type locality blank and lists no referred material in the AMNH.

Helohyus lentus (Marsh): In addition to the type a NMNH specimen is from the upper Bridger, and Matthew (1909b) lists a referred specimen in the AMNH as upper Bridger.

Helohyus validus (Marsh): Although the type is from the upper Bridger near Henry's Fork and McGrew (1959) records specimens from the upper Bridger at Tabernacle Butte, Sinclair (1914) refers

to a specimen in the AMNH from lower Bridger on Cottonwood Creek.

Helohyus milleri Sinclair: The recorded material, which apparently includes only the type in the AMNH and a referred specimen in the NMNH is from the upper Bridger.

Lophiohyus alticeps Sinclair: Apparently the only known specimen is the type from Twin Buttes, and as a consequence the level represented may be upper or lower Bridger.

Neodiadocodexis emryi Atkins: The only known specimen, the type, is from the upper Bridger at Tabernacle Butte.

Annotated Bibliography

The following annotated list of references to the Bridger Mammalia includes primarily those papers in which new taxonomic entities are defined or significant contributions are made to the definitions and relationships of the forms represented in the fauna. Also included are those papers which deal with the faunal zoning of the Bridger. Many papers

and texts that only repeat or summarize earlier reports and illustrations, or in which descriptions of other faunas include only comparisons or references to Bridger forms, for the most part are not included. Also omitted are references to descriptions of forms now regarded as synonyms unless important information is included.

Abel, Othenio

1922. *Lebensbilder aus der Tierwelt der Vorzeit*. Pages i-vi + 1-643, figures 1-507, 1 plate. Gustav Fischer: Jena. [While essentially a textbook treatment, Abel's book has a 48-page chapter devoted to the Bridger Basin and vicinity. About half of the 45 figures in this chapter pertain to the Bridger formation and its fauna, including skeletal remains and restorations.]

Allen, Joel A.

1877. Monographs of North America Rodentia: Sciuridae. Pages 631-940 in number 11 in F. V. Hayden, editor, *Report of the United States Geological Survey of the Territories*. [Included on pages 933-939 is a brief statement about each of the known Bridger rodents, which includes characteristics of genera and composition of the type of each species, its size, and where found.]

Atkins, Edward G.

1970. See West, Robert M., and Edward G. Atkins, 1970.
Aymard, Auguste
1850. Mammifères fossiles des calcaires du Puy. *Annales de la Société d'agriculture, sciences, arts et commerce du Puy*, 14:80-86. [The genus *Peratherium* is named.]

Bonaparte, Carlo L.

1838. Synopsis vertebratorum systematis. *Nuovi annali delle scienze naturali* (Bologna), 2:105-133. [The family Erinaceidae is named.]

Bradley, Wilmot H.

1964. Geology of the Green River Formation and Associated Eocene Rocks in Southwestern Wyoming and Adjacent Parts of Colorado and Utah. *Geological Survey Professional Paper*, 496-A:i-iv, A1-A86, figures 1-20, plates 1-3, tables 1-6. [Particularly important in showing the location by plane table survey of the Sage Creek White Layer, as well as the distribution of the Bridger, Green River, and Wasatch formations in the Bridger Basin.]

Butler, Percy M.

1952. The Milk Molars of Perissodactyla, with Remarks on Molar Occlusion. *Proceedings of the Zoological Society of London*, 121(4):777-817, figures 1-16. [Details of milk molars include discussion and figures of these teeth in Bridger species of *Palaeosyops* and *Hyrachyus*.]
1973. Molar Wear Facets of Early Tertiary North American Primates. *Symposium of the Fourth International Congress of Primatology*, 3:1-27, figures 1-12. [Discussion and figures of wear facets include Bridger species of *Notharctus*, *Omomyx*, and *Washakius*.]

Cope, Edward D.

- 1872a. Descriptions of Some New Vertebrata from the Bridger Group of the Eocene. *Proceedings of the American Philosophical Society*, 12:460-465. [Also, *Paleontological Bulletin*, number 1. Original description of the genus and species *Mesonyx obtusi-*

- dens, and the species *Antiacodon pygmaeus* as "*Lophiotherium pygmaeum*."] 1872b. Second Account of New Vertebrata from the Bridger Eocene. *Proceedings of the American Philosophical Society*, 12:466-468. [Also, *Palaeontological Bulletin*, number 2. Among the forms named as new are *Sinopa* ("*Stypholophus*") *pungens*, new species; *Pantolestes longicaudus*, new genus and species; and *Pseudotomus hians*, new genus and species (as an edentate).] 1872c. Third Account of New Vertebrata from the Bridger Eocene of the Wyoming Territory. *Proceedings of the American Philosophical Society*, 12:469-472. [Also, *Palaeontological Bulletin*, number 3. Included is the original description of the new genus and species *Miacis parvivorus*.] 1872d. Notices of New Vertebrata from the Upper Waters of Bitter Creek, Wyoming Territory. *Proceedings of the American Philosophical Society*, 12:483-486. [Also, *Palaeontological Bulletin*, number 6. The genus *Synoplotherium* is named.] 1872e. On a New Vertebrate Genus from the Northern Part of the Tertiary Basin of Green River. *Proceedings of the American Philosophical Society*, 12:554. [Also, *Palaeontological Bulletin*, number 8. Original description of the new genus and species *Anaptomorphus aemulus*.] 1873a. On Some Eocene Mammals Obtained by Hayden's Geological Survey of 1872. *Palaeontological Bulletin*, 12:1-6. [Among the various Bridger forms described as new, most of which were later regarded as synonyms, is included the new species *Antiacodon furcatus*.] 1873b. On the Extinct Vertebrata of the Eocene of Wyoming, Observed by the Expedition of 1872, with Notes on the Geology. Pages 545-649, plates 1-6, in F. V. Hayden, editor, *Sixth Annual Report of the United States Geological Survey of the Territories for 1872*. [Included is detailed description of many of the various Bridger Mammalia recognized in the collections.] 1873c. On the Short-footed Ungulata of the Eocene of Wyoming. *Proceedings of the American Philosophical Society*, 13:38-74, plates 1-4. [Outlines characters of Bridger species cited for *Uintatherium*. Compares various skeletal features of *Uintatherium* with other genera of Dinocerata, and Proboscidea as well.] 1873d. On the Flat-clawed Carnivora of the Eocene of Wyoming. *Proceedings of the American Philosophical Society*, 13:198-209. [Detailed description and measurements of *Mesonyx obtusidens*, also of *Synoplotherium* but based on a Washakie specimen.] 1873e. On the Osteology of the Extinct Tapiroid *Hyrachyus*. *Proceedings of the American Philosophical Society*, 13:212-224. [Detailed description with measurements of a skeleton of *Hyrachyus eximius*.] 1875. On the Supposed Carnivora of the Eocene of the Rocky Mountains. *Proceedings of the Academy of Natural Sciences, Philadelphia*, 1875:444-448. [Also, *Palaeontological Bulletin*, number 20. Names Creodonta and places it along with Tillodontia in the Insectivora. Also, names Mesonychidae, which he considers as representing the true Carnivora.] 1877. Report upon the Extinct Vertebrata Obtained in New Mexico by Parties of the Expedition of 1874. Part 2 in volume 4 in G. M. Wheeler, editor, *Report upon United States Geographical Surveys West of the One Hundredth Meridian*. iv + 370 pages, plates 22-83. [Oxyaenidae is named (page 89) and many Bridger mammals are discussed in comparisons.] 1880. On the Genera of Creodonta. *Proceedings of the American Philosophical Society*, 19:76-82. [Names the family Miacidae.] 1881. On the Vertebrata of the Wind River Eocene Beds of Wyoming. Pages 183-202 in part 8 in volume 6 in F. V. Hayden, editor, *Bulletin of United States Geological and Geographical Survey of the Territories*. [The genus *Bathyopsis* is named on a Wind River species.] 1882a. On *Uintatherium*, *Bathmodon* and *Trisodon*. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1882:294-300. [Discussion includes description and measurements of lower jaw from Henry's Fork that he refers to Leidy's *Uintatherium robustum*.] 1882b. Notes on Eocene Mammalia. *American Naturalist*, 16:522. [The genus *Didelphodus* is named with the Wasatchian species *Deltatherium absarokae* as the type.] 1883. On the Mutual Relations of Bunotherian Mammalia. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 35:77-83. [The families Anaptomorphidae and Esthonychidae are named.] 1884. The Vertebrata of the Tertiary Formations of the West, Book I. In volume 3 in F. V. Hayden, editor, *Report of the United States Geological Survey of the Territories*, pages i-xxxiv+1-1009, figures 1-38, plates 1-75a. [Includes a review of all the known Bridger forms in detail. Also, the family Pantolestidae is named.] 1884-1885. The Amblypoda. *American Naturalist*, 18:1110-1121, 1192-1202; 19:40-55, figures 1-35, plate 1. [Includes discussion and figures of uintatheres from the Bridger (in volume 19).] 1885. The Lemuroidea and Insectivora of the Eocene Period of North America. *American Naturalist*, 19:457-471, figures 1-18. [Discussion and figures include Bridger forms such as *Notharctus tenebrosus*, *Anaptomorphus aemulus*, and *Hyoopsodus paullus*.] Dawson, Mary R. 1961. The Skull of *Sciuravus nitidus*, a Middle Eocene Rodent. *Postilla*, 53:1-13, plates 1-5. [Description of a nearly complete skull and associated jaw from the Bridger.]

1962. A Sciuravid Rodent from the Middle Eocene of Wyoming. *American Museum Novitates*, 2075:1-5, 1 figure. [Bridger sciuravid upper teeth compared with known Bridger forms.]
- Dechaseaux, Colette
1958. Encephales de Dinocérates, pages 179-180, 1 figure; Encephales de Périssodactyles fossiles, pages 476-492, 18 figures; Encephale de Tillodontes, pages 506-507, 2 figures. In volume 2 in tome 6 in Pivctean, *Traité de paléontologie*. 962 pages, illustrated. Paris: Masson. [Brain casts of Bridger species of *Uintatherium*, *Orohippus*, *Palaeosyops*, and *Trogosus*, figured and discussed.]
- Denison, Robert H.
1938. The Broad Skulled Pseudocreoedi. *Annals of the New York Academy of Sciences*, 37:163-257, figures 1-32. [Several Bridger creodonts included in figures and discussion.]
- Dreverman, F.
1913. *Sinopa rapax* Leidy. *Bericht über die Senckenbergische naturforschende Gesellschaft in Frankfurt-am-Main*, 44:199-202, 4 figures. [Skull of *Sinopa grangeri*, the skeleton of *Sinopa rapax*, and the feet of *Hyaenodon* etc., figured.]
- Earle, Charles
- 1891a. On a New Species of *Palaeosyops* *Palaeosyops megarhinus*. *American Naturalist*, 25:45-47, 1 figure. [A Washakie specimen is made the type of a species now recognized as *Mesatirhinus megarhinus* and known also in the Bridger.]
- 1891b. *Palaeosyops* and Allied Genera. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1891: 105-117, figures 1-2. [Discusses known species of *Palaeosyops*, *Limnohyops*, and *Telmatherium* ("Telmatherium") as recognized on materials in various museum collections.]
1892. A Memoir upon the Genus *Palaeosyops* Leidy and Its Allies. *Journal of the Academy of Natural Sciences of Philadelphia*, series 2, 9 (part 3, article 6): 267-388, plates 10-14. [This includes the genera *Palaeosyops*, *Limnohyops*, and "Telmatherium" (= *Telmatherium*), principally Bridger material.]
1895. On the Supposed Case of Parallelism in the Genus *Palaeosyops*. *American Naturalist*, 29:622-626, plate 27. [Discusses parallelism in *Palaeosyops* in two lines through the Bridger and Uinta to *Titanotherium*.]
- Edinger, Tilly
1948. Evolution of the Horse Brain. *Geological Society of America Memoir* 25:i-x, 1-177, figures 1-24, plates 1-4, tables 1-11. [Describes in detail and figures endocranial casts of two Bridger specimens.]
1950. Frontal Sinus Evolution (particularly in Equidae). *Bulletin of the Museum of Comparative Zoology*, 103 (8):411-496, figures 1-7 (in text), figures 8-26 in plates 1-9. [Discussion and figures include Bridger *Uintatherium* and *Palaeosyops*.]
1956. Objets et résultats de la paléoneurologie. *Annales de Paléontologie*, 42:97-116, figures 1-18. Paris: Masson et Cie. [Bridger endocranial casts of *Thinoconyon*, *Palaeosyops*, and *Mesatirhinus* figured and involved in discussion.]
- Emmons, S. F.
1877. Green River Basin. Chapter 2 in volume 2, Descriptive Geology, in Clarence King, editor, *United States Geological Exploration of the Fortieth Parallel*. Pages 191-310, plates 1-9, and frontispiece. [Includes a lithologic description of Bridger as well as other formations in the basin and a list of vertebrate fossils (pages 249-250) described by Marsh and Leidy.]
- Emry, Robert J.
1970. A North American Oligocene Pangolin and Other Additions to the Pholidota. *Bulletin of the American Museum of Natural History*, 142 (6):457-510, figures 1-32. [Includes a discussion of the relationships of *Metacheiromys*.]
- Flower, Wm. Henry
1876. Hunterian Lectures on the Relations of Extinct to Existing Mammalia. *Nature*, 13:307-308, 327-328, 350-352, 387-388, 409-410, 449-450, 487-488, 513-514; 14:11. [Names *Uintatheriidae* (page 387).]
- Gazin, C. Lewis
1934. On the Priority of Specific Names for the Upper Bridger *Notharctus*. *Journal of Mammalogy*, 15 (1): 71. [Gave evidence for priority of *Notharctus robustior* Leidy, 1872, with respect to *Notharctus crassus* (Marsh), 1872.]
1941. Trailing Extinct Animals in Central Utah and the Bridger Basin of Wyoming. Pages 5-8 in *Explorations and Field-Work of the Smithsonian Institution in 1940*. Figures 6-10. [Includes description of certain collecting localities and successes in the Bridger, also photographs of some of the localities.]
1946. *Machaeroides eothen* Matthew, the Sabre-tooth Creodont of the Bridger Eocene. *Proceedings of the United National Museum*, 96 (3202):335-347, plates 45-46. [Skull, jaws, and certain limb bones described and relationships discussed. *Limnocyonidae* proposed as a family name.]
1949. A Leptictid Insectivore from the Middle Eocene Bridger Formation of Wyoming. *Journal of the Washington Academy of Sciences*, 39 (7):220-223, 1 figure. [*Hypictops syntaphus*, new genus and species described.]
1953. The Tillodontia: An Early Tertiary Order of Mammals. *Smithsonian Miscellaneous Collections*, 121 (10):i-vi, 1-110, figures 1-38, plates 1-16. [Includes a revision of the Bridger tillodonts and the new genus *Tillodon* is named.]

1955. A Review of the Upper Eocene Artiodactyla of North America. *Smithsonian Miscellaneous Collections*, 128 (8):i-iii, 1-96, plates 1-8, 2 charts. [Evolutionary chart, keys to genera, and discussions of relationships include Bridger forms. The Helohyinae proposed as a subfamily for Marsh's Helohyidae.]
1957. A Skull of the Bridger Middle Eocene Creodont, *Patriofelis ulta* Leidy. *Smithsonian Miscellaneous Collections*, 134 (8):1-20, plates 1-4. [In addition to the description of the skull and jaws found by the University of Utah, there is included a review of the various species of *Patriofelis*.]
1958. A Review of the Middle and Upper Eocene Primates of North America. *Smithsonian Miscellaneous Collections*, 136 (1):i-iii, 1-112, 1 chart, plates 1-14. [Includes a review of all the known Bridger forms, and among the new forms is the Bridger species *Anaptomorphus wortmani*. An error in the chart shows *Smilodectes* in the upper Bridger. Types and exceptional specimens figured.]
1959. Paleontological Exploration and Dating of the Early Tertiary Deposits in Basins Adjacent to the Uinta Mountains. Pages 131-138 in *Intermountain Association of Petroleum Geologists, 10th Annual Field Conference Guidebook*. 1 figure. [Includes a summary of the history of fossil mammal collecting in the Bridger and of zoning as initiated by Matthew. The error in zoning on the east side of the basin is corrected.]
1965. An Endocranial Cast of the Bridger Middle Eocene Primate *Smilodectes gracilis*. *Smithsonian Miscellaneous Collections*, 149 (4):1-14, plates 1-2. [Includes comparisons with *Notharctus* and European *Adapis*.]
1968. A Study of the Eocene Condylarthran Mammal *Hyopsodus*. *Smithsonian Miscellaneous Collections*, 153 (4):i-iv, 1-90, figures 1-10, plates 1-13. [In addition to a taxonomic review, a morphological study is made of the skeleton based on the Bridger species *Hyopsodus paulus*.]
- Gidley, James W.
1923. Paleocene Primates of the Fort Union, with Discussion of Relationships of Eocene Primates. *Proceedings of the United States National Museum*, 63 (1):1-38, plates 1-5. [Includes a restudy of *Notharctus* based on material described by Gregory.]
- Gill, Theodore
1872. Arrangement of the Families of Mammals and Synoptical Tables of Characters of the Subdivisions of Mammals. *Smithsonian Miscellaneous Collections*, 11 (1):i-vi, 1-98. [Names Dichobunidae and Leptictidae.]
- Gilmore, Charles W.
1931. Fossil Hunting in the Bridger Basin of Wyoming. Pages 13-20 in *Explorations and Field-Work of the Smithsonian Institution in 1930*. Figures 11-16. [Discussion of collecting localities and outstanding materials recovered. Includes photographs of localities.]
1943. Notes on Two Newly Mounted Fossil Vertebrate Skeletons in the United States National Museum. *American Journal of Science*, 241:764-766, 1 plate. [One of the skeletons is a nearly complete and articulated Bridger *Uintatherium*. Discussion includes vertebral formula. Mounted skeleton is illustrated.]
- Granger, Walter
1908. A Revision of the American Eocene Horses. *Bulletin of the American Museum of Natural History*, 24 (15): 221-264, figures 1-5, plates 15-18. [In addition to descriptions and illustrations of types of Bridger *Orohippus* species, *Orohippus progressus* is described as new.]
- Granger, Walter, and Wm. K. Gregory
1917. A Revision of the Eocene Primates of the Genus *Notharctus*. *Bulletin of the American Museum of Natural History*, 37 (34):841-859, tables 1-2, plates 103-107. [Reviews characters of all species, including Lost Cabin as well as Bridger. Most of the lower Bridger names have since been synonymized and others corrected.]
- Gray, John E.
1821. On the Natural Arrangement of Vertebrate Animals. *London Medical Repository*, 15 (1):296-310. [Named Didelphidae and Equidae.]
- Gregory, Wm. K.
1913. Relationship of the Tupaiidae and Eocene Lemurs, Especially *Notharctus*. *Bulletin of the Geological Society of America*, 24:247-252. [Discussion of *Entomolestes* as a tupaïid and *Notharctus* as a lemur.]
1915. On the Relationships of the Eocene Lemur *Notharctus* to the Adapidae and to Other Primates. *Bulletin of the Geological Society of America*, 26:419-425. [Gives resumé of earlier interpretations by others, and the anatomical evidence for his own conclusions.]
1920. On the Structure and Relations of *Notharctus*, an American Eocene Primate. *Memoirs of the American Museum of Natural History*, new series, 3 (2):49-243, figures 1-84, plates 23-59. [A detailed monograph on *Notharctus* based essentially on Bridger material, well illustrated and with extensive comparisons with other genera.]
1922. *The Origin and Evolution of the Human Dentition*. Pages i-xviii, 1-548, 353 figures, 15 plates. Baltimore: Williams and Wilkins Co. [Various Bridger primate dentitions discussed and figured, especially *Notharctus*.]
1929. The Musculature Anatomy and Restoration of the Titanotheres. In Osborn, 1929, *Titanotheres of Ancient Wyoming, Dakota, and Nebraska*. *United States Geological Survey Monograph*, 55:703-725, figures 640-662, plates 43-44. [Muscular anatomy

of Bridger titanotheres included in discussion and figures.]

Hatcher, J. B.

1895. On a New Species of *Diplacodon*, with a Discussion of the Relations of That Genus to *Telmatotherium*. *American Naturalist*, 29:1084-1090, figures 1-2, plates 38-39. [The genus *Manteoceras* is named in postscript, based on an upper Eocene species *T. validens*.]

Hayden, F. V.

1872. *Preliminary Report of the United States Geological Survey of Wyoming and Portions of Contiguous Territories* (2nd Annual Report). Parts I and II, pages 1-188, figures 1-20. [Includes a popular account of exploration in the Bridger area and description of rocks and formations encountered.]

Hill, Wm. C. Osman

1953. *Primates: Comparative Anatomy and Taxonomy, I: Strepsirhini*. Pages i-xxiii, 1-798, figures 1-199, plates 1-34. Edinburgh: University Press. [Includes summary descriptions of Bridger species of *Apatemys*, *Aphanolemur*, *Notharctus*, and *Uintasorex*.]
1955. *Primates: Comparative Anatomy and Taxonomy, II: Haplorhini: Tarsioidea*. Pages i-xx, 1-347, figures 1-49, plates 1-14. Edinburgh: University Press. [Includes summary descriptions of Bridger species of *Anaptomorphus*, *Hemicacodon*, *Washakius*, and *Uintanius*.]

Hubrecht, A. A. W.

1896. Die Keimblase von *Tarsius* ein Hilfsmittel zur schärferen definition gewisser saugethierordnungen. Pages 149-178 in Carl Gegenbaur, *Festschrift zum siebenzigsten geburtstage*. Figures a-l. [Extensive comparison of *Tarsius* and *Anaptomorphus homunculus*. Skull and teeth of latter figured (after Cope).]

King, Clarence

1878. Systematic Geology. Volume 1 in *United States Geological Exploration of the Fortieth Parallel*. Pages i-xi+1-803, 28 plates, 13 tables. [This includes about 12 pages (394-405) on the geology and topography of the Bridger Eocene (also Washakie) and a list of fossil vertebrates known as of 1878.]

Kitts, David B.

1897. A Revision of the Genus *Orohippus* (Perissodactyla, Equidae). *American Museum Novitates*, 1864:1-40, figures 1-9, tables 1-10. [Descriptions include numerical data and illustrations of teeth.]

Koenig, Karl J.

1960. Bridger Formation in the Bridger Basin, Wyoming. Pages 163-168 in *Wyoming Geological Association, 15th Annual Field Conference Guidebook*. Figures

1-3. [Discussion of the stratigraphy and lithology of the Bridger beds.]

Lane, Henry H.

1932. Variation in *Palaeosyops*. *The University of Kansas Science Bulletin*, 20 (18):319-326, 1 table. [Includes discussion of Osborn's stages of development in *Palaeosyops*.]

Lavocat, René

1958. Condylarthra, pages 1-27, figures 1-29; Dinocerata, pages 164-179, figures 1-18; Tillodontia, pages 499-505, figures 1-9; Taeniodonta, pages 508-521, figures 1-13. In volume 2 in tome 6 in Piveteau, *Traité de paléontologie*. [Discussion and figures of Bridger *Hyopsodus*, *Uintatherium*, *Trogosus* and *Stylinodon*.]

Leidy, Joseph

- 1869a. The Extinct Mammalian Fauna of Dakota and Nebraska, Including an Account of Some Allied Forms from Other Localities, Together with a Synopsis of the Mammalian Remains of North America. *Journal of the Academy of Natural Sciences of Philadelphia*, series 2, 7:1-472, plates 1-30. [Names Hyaenodontidae.]
- 1869b. Notice of Some Extinct Vertebrates from Wyoming and Dakota. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1869:63-67. [Includes original description of *Omomys carteri* (new genus and species).]
- 1870a. [Remarks on *Elasmosaurus platyurus* and Other Fossil Vertebrates.] *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1870:9-11. [Includes original description of *Patriofelis ulta* (new genus and species).]
- 1870b. [Remarks on a Collection of Fossils from the Western Territories.] *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1870:109-110. [Includes original descriptions of *Lophiodon* (= *Hyrachyus*) *modestus* (new species) and *Hyopsodus paulus* (new genus and species).]
- 1870c. [Descriptions of *Palaeosyops paludosus*, *Microsus cuspidatus*, and *Notharctus tenebrosus*.] *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1870:113-114. [All these are described as new genera and species.]
- 1870d. [Remarks on a Collection of Fossils from Table Mountain, California (also from the Bridger Basin, Wyoming).] *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1870:125-127. [The new species *Lophiotherium sylvaticum* is described (later referred to *Orohippus*).]
- 1871a. Report on the Vertebrate Fossils of the Tertiary Formations of the West. Pages 340-370 in part IV in F. V. Hayden, editor, *Preliminary Report of the United States Geological Survey of Wyoming and Portions of Contiguous Territories* (2nd Annual Report). [The genus *Hyrachyus* is named on the

- new species *Hyrachyus agrestis*. Also named is the new species *Hyrachyus agrarius*. Both species are synonyms of *Hyrachyus modestus*.]
- 1871b. Remains of Extinct Mammals from Wyoming. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1871:113-116. [*Sinopa rapax* and *Trogosus castoridens* named as new genera and species.]
- 1871c. Remarks on Fossil Vertebrates from Wyoming. *American Journal of Sciences*, 2:372-373. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1871:228-229. [The species *Hyrachyus eximius* and *Palaeosyops major* described as new.]
- 1871d. Notice of Some Extinct Rodents. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1871:230-232. [The genus *Paramys* and the three species *P. delicatus*, *P. delicatior*, and *P. delicatissimus* are described as new. Also described is the new genus and species *Mysops minimus*.]
- 1872a. Remarks on Fossils from Wyoming. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1872:19-21. [*Microsyops gracilis* is described as a new genus and species. Another form of *Microsyops* is described as *Palaeacodon verus* (= *M. elegans*).]
- 1872b. On the Fossil Vertebrates of the Early Tertiary of Wyoming. Pages 353-372 in F. V. Hayden, editor, *United States Geological Survey of Montana and Portions of Adjacent Territories* (5th Annual Report). [Bridger fauna is discussed by species and included is the original description of the species *Notharctus robustior*.]
- 1872c. On Some New Species of Fossil Mammalia from Wyoming. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1872:167-169. Also *American Journal of Science and Arts* (3)4:239-240. [Describes trip to Bridger Basin with Drs. Carter and Corson. Discusses material of *Palaeosyops* and *Hyrachyus* found. The genus *Uintatherium* is described as new on the basis of the new species *U. robustum*. Also describes *Uintamastix atrox* as new, which together with *U. robustum* are now regarded as synonyms of *Uintatherium anceps* (Marsh).]
- 1872d. Remarks on Fossils from Wyoming. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1872:277. [The new genus and species *Uintacyon edax* and a second new species *Uintacyon vorax* are named.]
1873. Contributions to the Extinct Vertebrate Fauna of the Western Territories. In volume I in F. V. Hayden, editor, *Report of the United States Geological Survey of the Territories*. 358 pages, plates 1-37. [Includes extensive description of the known Bridger Mammalia. Included are original descriptions of *Washakius insignis*, new genus and species; *Hyopsodus minusculus*, new species; and *Mysops fraternus*, new species.]
- Lydekker, Richard
1889. [No title.] Pages 889-1474 in volume 2 in Henry A. Nicholson and Richard Lydekker, *A Manual of Paleontology for Use of Students, with a General Introduction on the Principles of Paleontology*. 3rd Edition. Edinburgh and London: Wm. Blackwood and Sons. [Names Hyoposodontidae from Trouesart's Hyoposodinae of 1879.]
1904. The Ancestry of the Carnivora. *Knowledge*, 1 (new series):61-64, figures 1-5. [Discussion includes characteristics of Miacidae ("Viverravidae"), including *Vulpavus*, *Oödetes*, *Viverravus* (Bridger forms), and *Sinopa* in relation to modern Carnivora. *Vulpavus* and *Sinopa* figured.]
1910. Vertebrate Paleontology in 1909. *Science Progress*, 4:649-676. [Review of Matthew's 1909 memoir on Bridger. Includes discussion of some of Matthew's conclusions and refigures *Vulpavus projectus* and *Limnocyon verus*.]
- McGrew, Paul O.
1950. Tertiary Vertebrate Fossils of the Green River Basin. Pages 68-74 in *Wyoming Geological Association: 5th Annual Field Conference, Southwest Wyoming, Guidebook*. 2 figures, 1 map, 1 chart. [Includes brief discussion of history of investigation, characteristic mammal genera, and zoning. Also illustrations of upper teeth of five genera.]
- McGrew, Paul O., Jack E. Berman, Max K. Hecht, John M. Hummel, George G. Simpson, and Albert E. Wood
1959. The Geology and Paleontology of the Elk Mountain and Tabernacle Butte Area, Wyoming. *Bulletin of the American Museum of Natural History*, 117(3):117-176, figures 1-27, tables 1-15, plates 50-57. [Several localities for upper Bridger forms and two localities of lower Bridger materials in the Green River. Described as new species by McGrew are *Diacodon bacchanalis* and *Diacodon edenensis*, both since referred to *Scenopagus*. Also described as new by McGrew is *Peratherium knighti*. Wood described rodents found in the upper Bridger at Tabernacle Butte. Described as new are *Leptotomus parvus*, new species; *Microparamys*, new genus based on *Paramys minutus*; *Microparamys wyomingensis*, new species; and *Thisbemys corrugatus*, new genus and species.]
- McGrew, Paul O., and Raymond Sullivan
1970. The Stratigraphy and Paleontology of Bridger A. *University of Wyoming Contributions to Geology*, 9(2):66-85, figures 1-16. [A defense of the existence of Bridger A as distinct from Bridger B.]
- McKenna, Malcolm C.
1960. The Geolabidinae, a New Subfamily of Early Cenozoic Erinacoid Insectivores. *University of California Publications in Geological Sciences*, 37(2):131-164, figures 1-6. [In addition to describing the subfamily Geolabidinae under the Erinaceidae, he discussed the Bridger nyctitheriid *Myolestes dasypelix*, which he placed in his new subfamily. He also described from the Bridger a new genus and species *Hypacodon praecursor*, which Robinson (1962) synonymized with *Centetodon pulcher*.]

1963. Primitive Paleocene and Eocene Apatemyidac (Mammalia, Insectivora) and the Primate-Insectivore Boundary. *American Museum Novitates*, 2160:1-39, figures 1-5. [Discusses apatemyid position in Insectivora, also recognizes and figures the upper dentition in a Bridger specimen of *Apatemys*.]
- McKenna, Malcolm C., Peter Robinson, and Dwight W. Taylor
1962. Notes on Eocene Mammalia and Mollusca from Tabernacle Butte, Wyoming. *American Museum Novitates*, 2102:1-33, figures 1-9. [Includes discussion of relationships of various upper Bridger Insectivora from Tabernacle Butte and the presence of a possible chiropteran.]
- McKenna, Malcolm C., and George G. Simpson
1959. A New Insectivore from the Middle Eocene of Tabernacle Butte, Wyoming. *American Museum Novitates*, 1952:1-12, 1 figure, 1 table. [*Scenopagus mcgrewi* is described as a new genus and species. The species is later synonymized by Robinson with McGrew's species *Diacodon edenensis*, which is placed in the genus *Scenopagus*.]
- Marsh, Othniel C.
- 1871a. Notice of Some New Fossil Mammals from the Tertiary Formation. *American Journal of Science and Arts*, 2:35-44. [Described *Uintatherium* (as *Titanotherium?*) *anceps*, new species; *Hyrachyus* (as *Lophiodon*) *affinis*, new species; *Helaletes* (as *Lophiodon*) *nanus*, new species; *Orohippus* (as *Lophiodon*) *pumilus*, new species; *Helohyus* (as *Elothierium*) *lentus*, new species; *Smilodectes* (as *Hyopsodus*) *gracilis*, new species; *Microsyoops* (as *Limnotherium*) *elegans*, new species, all from the Bridger. Also included were other new names, since regarded as synonyms.]
- 1871b. Notice of Some New Fossil Mammals and Birds from the Tertiary Formations of the West. *American Journal of Science and Arts*, 2:120-127. [Described *Sciuravus nitidus*, new genus and species; and *Vulpavus palustris*, new genus and species, both from the Bridger. Other Bridger names proposed as new have since been regarded as synonyms of earlier names.]
1872. Preliminary Description of New Tertiary Mammals, Parts I-IV. *American Journal of Science and Arts*, 4:122-128, 202-224, and 504. [Described *Limnohyops* (as *Palaeosyoops*) *laticeps*, new species; *Telmatherium validum* (as *T. validus*), new genus and species; *Palaeosyoops* (as *Limnohyus*, new genus) *robustus*, new species; *Hyrachyus princeps*, new species; *Homacodon vagans*, new genus and species; *Limnocyon verus*, new genus and species; *Viverravus gracilis*, new genus and species; *Miacis* (as *Harpalodon*, new genus) *sylvestris*, new species; *Nyctitherium velox*, new genus and species; *Scenopagus* (as *Nyctitherium*) *priscus*, new species; *Talpavus nitidus*, new genus and species; *Patriofelis* (as *Limnofelis*, new genus) *ferox*, new species; *Tritemnodon* (as *Limnocyon*) *agilis*, new species; *Thinocyon velox*, new genus and species; *Orohippus*, new genus for *Lophiodon pumilus*; *Helohyus plicodon*, new genus and species; *Helohyus* (as *Thinotherium*, new genus) *validus* (as *valudum*), new species; *Pantolestes* (as *Anisacodon*, new genus) *elegans*, new species; *Centetodon pulcher*, new genus and species; *Antiacodon venustus*, new genus and species; *Microsyoops* (as *Bathrodon*) *annectens*, new species; *Hemiacodon gracilis*, new genus and species; *Didelphodus* (as *Centetodon*) *altidens*, new species; *Nyctitherium* (as *Nyctilestes*, new genus) *serotinum* (as *serotinus*) new species; *Helaletes boops*, new genus and species; *Pseudotomus* (as *Paramys*) *robustus*, new species; *Tillomys senex*, new genus and species; *Taxymys lucaris*, new genus and species; *Tillomys?* (as *Sciuravus*) *parvidens*, new species; *Mysops* (as *Tillomys*) *parvus*, new species; *Apatemys bellus*, new genus and species; *Apatemys bellulus*, new species. Several other new names proposed have since been regarded as synonyms of earlier names.]
- 1873a. On the Gigantic Fossil Mammals of the Order Dinocerata. *American Journal of Science and Arts*, 5:117-122, plates 1-2. [Includes discussion of characteristics of *Uintatherium* (as *Tinoceras* and *Dinoceras*) and figures of Bridger skull labeled "*Dinoceras mirabilis*".]
- 1873b. Notice of New Tertiary Mammals. *American Journal of Science and Arts*, 5:407-410, 485-488. [Described *Orohippus agilis*, new species; *Trogosus* (as *Tillotherium*, new genus) *hyrachoides*, new species; and named Brontotheriidae (as Brontotheridae).]
- 1874a. Notice of New Equine Mammals from the Tertiary Formation. *American Journal of Science and Arts*, 7:247-258, 5 figures. [Includes discussion of *Orohippus* and description of *Orohippus major*, new species.]
- 1874b. Notice of New Tertiary Mammals, III. *American Journal of Science and Arts*, 7:531-534. [Includes description of *Stylinodon mirus*, new genus and species, also *Trogosus* (as *Tillotherium*) *latidens*, new species.]
- 1875a. New Order of Eocene Mammals. *American Journal of Science and Arts*, 9:221. [Named the order Tillodontia and the family Stylinodontidae.]
- 1875b. Notice of New Tertiary Mammals, IV. *American Journal of Science and Arts*, 9:239-250. [Described as new *Tillodon* (as *Tillotherium*) *fodiens*, new species, from the Bridger, as well as new forms from other formations.]
- 1876a. Principal Characters of the Dinocerata. *American Journal of Science and Arts*, 11:163-168, 1 figure. [Description of skull, brain, and other skeletal portions of *Uintatherium* (as *Dinoceras mirabile*). Skull figure includes sketch of brain cavity.]
- 1876b. Principal Characters of the Tillodontia. *American Journal of Science and Arts*, 11:249-252, plates 8-9. [Description of skull, etc., of *Trogosus* (as *Tillotherium*) *fodiens*, more comprehensive than previously.]

- 1876c. Notice of New Tertiary Mammals, V. *American Journal of Science and Arts*, 12:401-404. [Includes description of *Synoplotherium* (as *Dromocyon vorax*, new species.)]
1881. Restoration of *Dinoceras mirabile*. *American Journal of Science*, 22:31-32, plate 2. [Discussion and drawing of articulated skeleton of *Uintatherium*. First printing of a much copied illustration.]
1885. The Gigantic Mammals of the Order Dinocerata. Pages 243-302 in *Fifth Annual Report of the United States Geological Survey*. Figures 36-137. [Detailed description of the skeleton of Bridger *Uintatherium*, under various synonyms of *U. anceps*, and comparisons with other large mammals.]
1886. Dinocerata: A Monograph of an Extinct Order of Gigantic Mammals. *Monographs of the United States Geological Survey*, 10:i-xviii, 1-243, plates 1-56. [A comprehensive study of the known materials of uintatheres, particularly Bridger, but including Wind River and Washakie forms as well. The skeletal details of Bridger material well illustrated.]
1890. Notice of New Tertiary Mammals. *American Journal of Science*, 39:523-525. [New genus *Limnohyops* is described on basis of species *Palaeosyops laticeps*.]
1894. Description of Tertiary Artiodactyles, *American Journal of Science*, 48:259-274. [Includes further description, including figures of tooth and foot material, of *Homacodon vagans* and *Helohyus plicodon*.]
1897. The Stylinodontia, a Suborder of Eocene Edentates. *American Journal of Science*, 3:137-146, figures 1-9. [Further description, including figures of teeth, cervical vertebrae, and limb material of *Stylinodon mirus*.]
- Matthew, Wm. D.
1899. A Provisional Classification of the Fresh-water Tertiary of the West. *Bulletin of the American Museum of Natural History*, 12:19-75. [Bridger species listed by locality. Also, the new genus *Palaeictops* is named with a Wind River species *Ictops bicuspis* as the type.]
1906. The Osteology of *Sinopa*, a Creodont Mammal from the Middle Eocene. *Proceedings of the United States National Museum*, 30:203-233, figures 1-20, 1 plate. [Detailed description with illustrations of the Bridger skeleton in the National Museum of Natural History. *Sinopa grangeri* is named as a new species and *Tritemnodon* is named as a new genus based on the species *Limnocyon agilis* Marsh.]
- 1909a. Faunal Lists of the Tertiary Mammalia of the West. *Bulletin of the United States Geological Survey*, 361:91-138. [Includes listing of Bridger mammals according to occurrence in upper and lower levels. Also includes geologic range of various genera.]
- 1909b. The Carnivora and Insectivora of the Bridger Basin, Middle Eocene. *Memoirs of the American Museum of Natural History*, 9(6):291-567, figures 1-118, plates 42-52. [Comprehensive study of the two orders and includes stratigraphic zoning of the Bridger from A to E with a listing of mammals encountered in each horizon. Mapping errs in the distribution of the Sage Creek White Layer. Among the forms described as new are: *Uintasorex parvulus*, new genus and species; *Trogolemur myodes*, new genus and species; *Nyctitherium* (as *Myolestes*, new genus) *dasyplex*, new species; *Entomolestes grangeri*, new genus and species; *Hyopsodus despiciens*, new species; *Hyopsodus lepidus*, new species; *Pantolestes natans*, new species; *Pantolestes phocipes*, new species; *Pantolestes intermedius*, new species; *Viverravus sicarius*, new species; *Uintacyon jugulans*, new species; *Uintacyon major*, new species; *Oödetes proximus*, new species; *Vulpavus profectus*, new species; *Vulpavus ovatus*, new species; *Palaearctonyx meadi*, new genus and species; *Thinocyon mustelinus* (*T. minimus* in listing), new species; *Machaeroides eothen*, new genus and species. Named new family Apatemyidae.]
1910. On the Osteology and Relationships of *Paramys*, and the Affinities of the Ischyromyidae. *Bulletin of the American Museum of Natural History*, 28(6):43-72, figures 1-19. [The genera *Ischyrotomus* and *Leptomus* are described as new subgenera based on upper Eocene species.]
1915. A Revision of the Lower Eocene Wasatch and Wind River Faunas, Part IV: Entelonychia, Primates, Insectivora (part). *Bulletin of the American Museum of Natural History*, 34(14):429-483, figures 1-52, plate 15. [Bridger forms involved in keys and discussions in this and other parts of the lower Eocene revision. *Uintanius* is named as a new genus with the new species *U. turriculorum* as the type, a synonym of *U. ameghini*.]
1920. A New Genus of Rodents from the Middle Eocene. *Journal of Mammalogy*, 1(4):168-169. [The new genus *Reithroparamys* is named for *Paramys delicatissimus*.]
- Merrill, George P.
1907. Catalogue of the Type and Figured Specimens of Fossils, Minerals, Rocks, and Ores in the Department of Geology, United States National Museum, Part II: Fossil Vertebrates; Fossil Plants; Minerals, Rocks, and Ores. *Bulletin of the United States National Museum*, 53(2):1-370. [Includes in the list the Bridger mammal types in the NMNH with original citations.]
- Miller, Gerrit S., Jr., and James W. Gidley
1918. Synopsis of the Supergeneric Groups of Rodents. *Journal of the Washington Academy of Sciences*, 8(13):431-448. [Names Paramyidae, Cylindrodontidae, and Scuravidae.]
- Osborn, Henry F.
1892. See Osborn and Wortman, 1892.
1898. The Extinct Rhinoceroses. *Memoirs of the American*

- Museum of Natural History*, 1(3):75-164, figures 1-49, plates 12A-20. [Discussion includes features of *Hyrachyus agrarius* (= *H. eximius*) as a hyracodont. Also, the skull and skeleton are figured.]
1899. See Matthew, W. D., 1899:47. [Names *Manteoceras* (as *Palaeosyops*) *manteoceras*, new species.]
1900. *Oxyaena* and *Patriofelis* Restudied as Terrestrial Creodonts. *Bulletin of the American Museum of Natural History*, 13(20):269-279, figures 1-8, plates 18-19. [The skeletal features of Bridger *Patriofelis* are reviewed and illustrated. He refutes Wortman's suggestion that certain *Patriofelis* characters indicate an aquatic habit.]
1902. American Eocene Primates, and the Supposed Rodent Family Mixodectidae. *Bulletin of the American Museum of Natural History*, 16(17):169-214, figures 1-40. [Discussion and illustrations include Bridger primates, hyopsodontids, and microsyopsids. The new Bridger species *Hyopsodus marshi* is described.]
1904. An Armadillo from the Middle Eocene (Bridger) of North America. *Bulletin of the American Museum of Natural History*, 20(12):163-165. [Names two new species, *Metacheiromys dasyptus* and *Metacheiromys tatusia*.]
- 1908a. New or Little Known Titanotheres from the Eocene and Oligocene. *Bulletin of the American Museum of Natural History*, 24(32):599-617, figures 1-21. [Among the forms named as new are *Limnohyops priscus*, new species, and *Mesatirhinus petersoni*, new genus and species.]
- 1908b. Coincident Evolution through Rectigradations and Fluctuations (third paper). *Science* (new series) 27: 749-752, figures 1-3. [Uses *Palaeosyops* and *Orohippus* as examples in discussing principles involved in dolichocephaly and brachycephaly. Drawings include skulls and teeth of *Palaeosyops* and teeth of *Orohippus*.]
1909. Cenozoic Mammal Horizons of Western North America. *Bulletin of the United States Geological Survey*, 361:1-90, figures 1-15, plates 1-3. [Includes a three-page section devoted to the Bridger zoning, A to E, and faunal characters.]
1929. The Titanotheres of Ancient Wyoming, Dakota, and Nebraska. *United States Geological Survey Monograph* 55(2 volumes):1-953, figures 1-797, plates 1-236. [Includes comprehensive discussion of Bridger titanotheres, including original descriptions. Well illustrated.]
- Osborn, Henry F., Wm. B. Scott, and Francis Speir, Jr.
1878. Palaeontological Report of the Princeton Scientific Expedition of 1877. *Contributions from the Museum of Geology and Archaeology of Princeton College*, 1:1-107, plates A, 1-10. [Description in detail of the materials of the various species obtained in the Bridger beds. Among the forms described as new are *Helaletes* (as *Hyrachyus*) *intermedius*, new species; *Ischyrotomus* (as *Paramys*) *superbus*, new species; *Telmatherium* (as *Leurocephalus*, new genus) *cultridens* new species; and *Parisectolophus* (as *Helaletes*) *latidens*, new species.]
- Osborn, Henry F., and Jacob L. Wortman
1892. Fossil Mammals of the Wahsatch and Wind River Beds: Collection of 1891. *Bulletin of the American Museum of Natural History*, 4(11):81-147, figures 1-18(+1), plate 4. [Discussions and comparisons involve Bridger species, also, the families Microsyopsidae and Helaletidae are named by Osborn.]
1895. Perissodactyls of the Lower Miocene White River Beds. *Bulletin of the American Museum of Natural History*, 7(12):343-375, figures 1-12, plates 8-11. [There is inserted description with illustrations of skull and teeth of a newly found Bridger specimen of *Hyrachyus agrarius* (= *H. modestus*).]
- Palmer, Theodore S.
1904. Index Generum Mammalium: A List of the Genera and Families of Mammals. *North American Fauna*, 23:1-984. [Includes all Bridger genera and family names up to 1904, also the type, author, reference, and derivation of each generic name is given, and the family arrangement.]
- Peterson, Olof A.
1919. Report upon the Material Discovered in the Upper Eocene of the Uinta Basin by Earl Douglass in the Years 1908-1909, and by O. A. Peterson in 1912. *Annals of the Carnegie Museum*, 12(2-4):40-168, figures 1-19, plates 34-47. [Discussions include Bridger forms. New forms named include the family Isectolophidae, and the genus *Parisectolophus* based on the Bridger species *Isectolophus* (originally *Helaletes*) *latidens*.]
- Piveteau, Jean
1961. Carnivora. Pages 641-820 in volume 1 in tome 6 in *Traité de paléontologie*. Figures 1-203. [Most of the better known Bridger Carnivora are figured and discussed, including endocranial cast of *Thinocyon velox*.]
- Radinsky, Leonard B.
1963. Origin and Early Evolution of North American Tapiroidea. *Yale University, Peabody Museum of Natural History Bulletin*, 17:1-106, figures 1-21, plates 1-4. [Bridger helaletids and isctolophids included in the study. Included in the lumping of certain genera and of certain species, the genus *Parisectolophus*, named by Peterson for *I. latidens*, is placed in synonymy with *Isectolophus*.]
1967. *Hyrachyus*, *Chasmotherium*, and the Early Evolution of the Helaletid Tapiroids. *American Museum Novitates*, 2313:1-23, figures 1-4, tables 1-5. [Principally concerned with Bridger *Hyrachyus eximius* and *Hyrachyus modestus*. Various species are placed in synonymy with one or the other of these two, including certain European forms. The Hyrachyidae is regarded as a subfamily of Helaletidae.]
- Robinson, Peter
1957. The Species of *Notharctus* from the Middle Eocene.

- Postilla* (Yale Peabody of Natural History), 28:1-27, figures 1-5, plates 1-2. [The Bridger species of *Notharctus* are reduced in number and arranged by horizon. *Smilodectes* (as *Notharctus*) *gracilis* is included in the upper Bridger because of its occurrence at Twin Buttes. The level involved is now known to be Bridger B.]
1963. Fused Cervical Vertebrae from the Bridger Formation (Eocene) of Wyoming. *University of Colorado Studies, Series in Geology*, 1:6-9, 1 figure. [Interpreted as representing an upper Bridger edentate.]
- 1968a. *Talpavus* and *Entomolestes* (Insectivora, Adapisoricidae). *American Museum Novitates*, 2339:1-7, 1 figure. [Describes characteristics of the Bridger species *Talpavus nitidus* and *Entomolestes grangeri*.]
- 1968b. Nyctitheriidae (Mammalia, Insectivora) from the Bridger Formation of Wyoming. *Contributions to Geology* (University of Wyoming), 7(2):129-138, plates 1-2. [Review of the genus *Nyctitherium* includes the species *N. velox* and *N. serotinum* and to these are added *N. dasypelix* the type of *Myolestes*. *Entomacodon angustidens* is placed in synonymy with *Nyctitherium serotinum* and *Entomacodon minutus* is regarded as a didelphid. *Nyctitherium curtidens* is considered a synonym of *Scenopagus priscus*.]
- Saban, Roger
1958. Palaeonodonta, pages 522-533, figures 1-4; Insectivora, pages 822-909, figures 1-37. In volume 2 in tome 6 in Piveteau, *Traité de paléontologie*. [Some genera of Bridger insectivores discussed and *Metacheiromys tatusia* figured and discussed.]
- Schaub, Samuel
1958. Simplicidentata. Pages 659-818 in volume 2 in tome 6 in Piveteau, *Traité de paléontologie*. Figures 1-283. [Various Bridger genera included in discussion. Bridger *Paramys* species, *Sciuravus nitidus*, *Taxymys lucaris*, and *Mysops parvus* figured.]
- Schlosser, Max
1887. Die Affen, Lemuren, Chiropteren, Insectivoren, Marsupialer, Creodonten, und Carnivoren des europäischen Tertiärs und deren Beziehungen zu ihren lebenden und fossilen aus europäischen Verwandten. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients* (Wien). 6:1-224, plates 1-5. [Names family Adapisoricidae.]
- Scott, Wm. B.
1883. On *Desmatotherium* and *Dilophodon*, Two New Eocene Lophiodonts. *Contributions from the E. M. Museum of Geology and Archaeology of Princeton College, Bulletin* 3(4):46-53, plate 8. [Includes description of *Dilophodon minusculus*, new genus and species, and figure of lower jaw.]
1888. On Some New and Little Known Creodonts. *Journal of the Academy of Natural Sciences of Philadelphia* (series 2), volume 9, part 2, article 4, pages 155-185, plates 5-7. [Extensive description and figures of skeleton of *Mesonyx obtusidens*.]
- Simpson, George G.
1927. A North American Oligocene Edentate. *Annals of the Carnegie Museum*, 17(2):283-298, plate 24. [Names the family Epoicotheriidae.]
- 1928a. A New Mammalian Fauna from the Fort Union of Southern Montana. *American Museum Novitates*, 297:1-15, figures 1-14. [Names the family Nyctitheriidae.]
- 1928b. American Eocene Didelphids. *American Museum Novitates*, 307:1-7, figures 1-5. [Describes the Bridger species *Peratherium innomatum*.]
- 1931a. *Metacheiromys* and the Edentata. *Bulletin of the American Museum of Natural History*, 59(6):295-381, figures 1-23. [Reviews the species of *Metacheiromys* and their association with other forms in the upper and lower Bridger. Describes in detail the skeleton of *Metacheiromys dasypus*. Names *Metacheiromys osborni*, new species.]
- 1931b. A New Classification of Mammals. *Bulletin of the American Museum of Natural History*, 59(5): 259-293. [Names family Palaeoryctidae.]
1940. Studies on the Earliest Primates. *Bulletin of the American Museum of Natural History*, 77(4):185-212, figures 1-8. [In addition to a discussion of the relationships of early primates, there is included a detailed description of a Bridger skeleton referred to *Hemiacodon*.]
1945. The Principles of Classification and a Classification of Mammals. *Bulletin of the American Museum of Natural History*, 85:i-xvi, 1-350. [With regard to the Bridger genera, except for a few amendments and additions, this remains the most logical arrangement.]
- 1959a. A New Middle Eocene Edentate from Wyoming. *American Museum Novitates*, 1950:1-8, figures 1-2. [Describes *Tetrapassalus mckennai*, new genus and species.]
- 1959b. Two New Records from the Bridger Middle Eocene Tabernacle Butte, Wyoming. *American Museum Novitates*, 1966:1-5, figures 1-2. [Describes *Palaeictops* (as *Diacodon*) *bridgeri*, new species, and records *Viverravus*, cf. *gracilis*.]
- Sinclair, Wm. J.
1906. Volcanic Ash in the Bridger Beds of Wyoming. *Bulletin of the American Museum of Natural History*, 22(15):273-280, plates 35-38. [Describes stratigraphic and lithologic features of the Bridger, although division of formation into lower, middle, and upper does not correspond to zoning of Matthew and Granger.]
1914. A Revision of the Bunodont Artiodactyla of the Middle and Lower Eocene of North America. *Bulletin of the American Museum of Natural History*, 33(21):267-295, figures 1-28. [The Bridger forms are reviewed and illustrated, and described as new are *Lophiohyus alticeps*, new genus and species, and *Helohyus milleri*, new species.]

Stehlin, Hans G., and Samuel Schaub

1951. Die Trigonodontie der Simplicidentaten Nager. *Schweizerische Palaeontologische Abhandlungen*, 67:1-385, figures 1-620. [Discussion and figures of dentitions of *Sciuravus nitidus*, *Mysops fraternus*, and *Taxymys lucaris* included.]
- Strauss, Wm. L., Jr.
1929. Studies of Primate Ilia. *American Journal of Anatomy*, 43:403-460, 8 figures. [Detailed study of *Notharctus ilium* is subject of 4-page appendix.]
- Szalay, Frederick S.
- 1969a. Uintasoricinae, a New Subfamily of Early Tertiary Mammals (?Primates). *American Museum Novitates*, 2363:1-36, figures 1-25. [The Bridger form *Uintasorex parvulus* is included in the subfamily, which he regards as part of the Microsypopsidae (as "Microsypopsidae").]
- 1969b. Mixodectidae, Microsypopsidae, and the Insectivore-Primate Transition. *Bulletin of the American Museum of Natural History*, 140(4):193-330, figures 1-28, tables 1-21, plates 17-57. [Various Bridger forms are involved in the study. It is noted that several of the National Museum of Natural History specimens figured are cited incorrectly as to formation or member and with names corresponding to the incorrect level. These are not misplaced legends as only two show the wrong catalog number. As a consequence his criteria for rearrangements and synonymy seem meaningless. The observed errors of fact and the revised determinations are as follows: Plate 34: figures 4-6, number 12138 is *Microsyps elegans* from the Bridger, not "*Microsyps*" *latidens* from the Knight. Plate 34: figures 7-9, no. 12140 is *Microsyps gracilis* from the Bridger, not "*Microsyps*" *latidens* from the Knight. Plate 36: figures 3 and 4, nos. 22110 and 22114 are from the Knight member of the Wasatch, not the New Fork, and which I regard as *Cynodontomys latidens* rather than *Microsyps*. Plate 36: figures 11 and 12 are of no. 22108, not 22107, which is from the Knight, not the New Fork, and which I regard as *Cynodontomys knightensis* rather than "*Microsyps*" *latidens*. Plate 37, figures 3-7, nos. 19322 and 22107 are from the Knight, not the New Fork, and which I regard as *Cynodontomys knightensis* rather than "*Microsyps*" *latidens*. Plate 39: figures 9 and 10, no. 17817 I regard as *Microsyps schlosseri*, not *Microsyps annectens*. Plate 40: figures 3-5 are of no. 22106, not 22107, and is from the Knight, not the New Fork member of the Wasatch, and which I regard as *Cynodontomys knightensis*, not "*Microsyps*" *latidens*. Plate 47: figures 1-5, no. 13440 I regard as representing *Microsyps gracilis* from an associated maxilla, not *Microsyps elegans*. I should also add that on page 248 he has listed "*Diacodexis*: Gazin, 1952, p. 71" as a synonym of *Microsyps*. This artiodactyl tooth cannot represent *Microsyps*.]
- Thorpe, Malcolm R.
1923. Notes on the Bridger (Eocene) Carnivora. *American Journal of Science* 5(3):23-39. [Discusses material and characteristics of various carnivora represented in the Marsh collection.]
- Tilney, Frederick
1931. Fossil Brains of Some Early Tertiary Mammals of North America. *Bulletin of the Neurological Institute of New York*, 1(3):430-505, figures 1-46. [Includes description and figures of brains of Bridger *Uintatherium*, *Palaeosyops*, and *Hyrachyus*.]
- Trouessart, Edouard-Louis
1879. Catalogue des mammifères vivants et fossiles. *Revue et magazine de zoologie pure et appliqué* (Paris), series 3, 7:219-285. [Names Notharctidae.]
- Troxell, Edward L.
- 1922a. *Helaletes* Redefined. *American Journal of Science*, 3(35):365-370, figures 1-3. [Describes skull of *Helaletes boops* and distinguishes species *Helaletes nanus* on upper premolars.]
- 1922b. The Genus *Hyrachyus* and Its Subgroups. *American Journal of Science*, 4(5):38-49, figures 1-5, plate 1. [Review of Bridger species of *Hyrachyus*.]
- 1923a. *Pauromys perditus*, a Small Rodent. *American Journal of Science*, 5(10):155-156, 1 figure. [*Pauromys perditus* is described as a new genus and species from the Bridger.]
- 1923b. The Eocene Rodents *Sciuravus* and *Tillomys*. *American Journal of Science*, 5(32):383-396, figures 1-23. [Reviews and figures dentitions of Bridger species of *Sciuravus* and *Tillomys*.]
- 1923c. The Apatemyidae. *American Journal of Science*, 5(40):503-506, figures 1-6. [Reviews Bridger species of *Apatemys* and adds *Apatemys rodens*, new species.]
- 1923d. A New Marsupial. *American Journal of Science*, 5(41):507-510, figures 1-4. [Describes the new marsupial *Peratherium* (as *Herpetotherium*) *marsupium*, new species.]
1926. *Smilodectes* or *Notharctus*. *American Journal of Science*, 11:423-428, figures 1-3. [Discusses confusion with regard to the relationships of *Smilodectes gracilis*, which he includes in *Notharctus*.]
- Van Valen, Leigh
1963. The Origin and Status of the Mammalian Order Tillodontia. *Journal of Mammalogy*, 44(3):364-373, plates 1-2. [Includes the order Tillodontia in the Condylarthra because he thinks (mistakenly) that *Esthonyx* looks like the creodont carnivore *Claenodon*.]
1966. Deltatheridia, a New Order of Mammals. *Bulletin of the American Museum of Natural History*, 132(1): 1-17, plates 1-8. [Among the Bridger forms mistakenly included in the order Deltatheridia are species belonging to the palaeoryctid insectivore, *Didelphodus*; the oxyaenid creodont, *Patriofelis*; the hyaenodontid creodonts *Sinopa* and *Tritemnodon*; and the limnocyonid creodonts *Limnocyon*, *Thinocyon*, and *Machaeroides*.]

Viret, Jean

1958. Perissodactyla. Pages 368–475 in volume 2 in tome 6 in Piveteau, *Traité de paléontologie*. Figures 1–112, plate 1. [Genera of horses, titanotheres, and tapiroids represented in the Bridger are discussed and figured.]
1961. Artiodactyla. Pages 887–1021, 1038–1084 in volume 1 in tome 6 in Piveteau, *Traité de paléontologie*. Figures 1–119, 136–166. [Includes discussion of some of the Bridger artiodactyles and *Homacodon vagans* is figured.]

West, Robert M.

- 1969a. *Paramys wyomingensis*, a Small Rodent from the Middle Eocene of Wyoming. *Journal of Paleontology*, 43(1):175–178, figures 1–2. [Regards Wood's *Microparamys wyominensis* as belonging in the genus *Paramys*.]
- 1969b. Biostratigraphy of Fluvial Sediments of the Upper Wasatch Formation in the Northern Green River Basin, Wyoming. *Contributions to Geology* (University of Wyoming), 8(number 2, part 2):184–196, 5 figures, 1 table. [Refers beds with a Bridgerian fauna, which are in the same depositional basin as the Bridger formation, to the Cathedral Bluffs tongue, which is essentially a late Wasatchian deposit in the Washakie and Great Divide Basins. See Bradley's map (1964) and the map in McGrew et al. (1959), also my report of 1959 (page 135).]
1972. Minimammals and Bridger Biostratigraphy. *Society of Vertebrate Paleontology Guidebook* (Field Conference on Tertiary Biostratigraphy of Southern and Western Wyoming), pages 40–50, figures 1–3. [Bridger fauna discussed by genera only. Regards local ecology as possibly more significant than vertical zoning.]
- 1973a. Geology and Mammalian Paleontology of the New Fork–Big Sandy Area, Sublette County, Wyoming. *Fieldiana Geology*, 29:i–viii, 1–193, figures 1–33, plates 1–16, tables 1–61, sections 1–14. [Includes discussion of Bridger species encountered at various localities. Reference to Robinson concerning occurrence of *Smilodectes* in the upper Bridger is an error which must result from Matthew's interpretation of zoning at Twin Buttes. Its occurrence there is lower not upper Bridger.]
- 1973b. Review of North American Eocene and Oligocene Apatemyidae (Mammalia: Insectivora). *Special Publications, The Museum, Texas Tech University*, 3:1–42, figures 1–20. [Lumps all Bridger and earlier apatemyid genera and species into one form, *Apatemys bellus*.]
- 1973c. An Early Middle Eocene Epoicotheriid (Mammalia) from Southwestern Wyoming. *Journal of Paleontology*, 47(5):929:931, 1 figure. [Describes *Tetrapassalus prius*, new species, on an indeterminate lower jaw fragment with one spikelike tooth preserved.]

1974. New North American Middle Eocene Nyctitheres (Mammalia, Insectivora). *Journal of Paleontology*, 48(5):983–987, 1 plate. [Names *Pontifactor bestiola*, new genus and species, from the upper Bridger.]

West, Robert M., and Edward G Atkins

1970. Additional Middle Eocene (Bridgerian) Mammals from Tabernacle Butte, Sublette County, Wyoming. *American Museum Novitates*, 2404:1–26, figures 1–9, tables 1–2. [Considerable discussion of *Phenacodus* material thought to occur in beds of Bridgerian age. A new artiodactyl from the upper Bridger, *Neodiapodexis emryi*, new genus and species, is described by Atkins.]

West, Robert M., and Mary R. Dawson

1973. Fossil Mammals from the Upper Part of the Cathedral Bluffs Tongue of the Wasatch Formation (Early Bridgerian), Northern Green River Basin, Wyoming. *Contributions to Geology* (University of Wyoming), 12(1):33–41, figures 1–4. [Reports about 40 species of Bridgerian mammals, mostly rodents from the "Green locality," purported to be from the upper part of the Cathedral Bluffs tongue, but not the same basin of deposition as the true Cathedral Bluffs tongue.]

Wheeler, Walter H.

1961. Revision of the Uintatheres. *Yale University, Peabody Museum of Natural History Bulletin*, 14:1–iv, 1–93, figures 1–9, plates 1–14. [Reduces the numerous names that have been applied to upper Bridger uintatheres to one, *Uintatherium anceps*, in the Bridger Basin. *Bathyopsis middleswarti*, new species, is named for the lower Bridger beds.]

Wilson, Robert W.

1937. Two New Eocene Rodents from the Green River Basin, Wyoming. *American Journal of Science*, 34: 447–456, figures 1–2. [Describes *Microparamys* (as *Paramys*) *minutus*, new species, from the Bridger, exact locality not known.]
1938. Review of Some Rodent Genera from the Bridger Eocene (Parts I–III). *American Journal of Science*, 35:123–137, 207–222, 297–304, figures 1–15. [Reviews the genera *Sciuravus*, *Mysops*, *Tillomys*, *Taxymys*, and *Pauromys*. Named as new species are *Sciuravus bridgeri*, *Sciuravus? rarus*, and *Taxymys? progressus*.]
1949. Early Tertiary Rodents of North America. *Carnegie Institution of Washington Publication*, 584: 67–164, figures 1–13. [Discussion of relationships involves known Bridger forms. Also, a chart shows distribution of species between upper and lower levels, but according to the original distribution of the zones according to work of Matthew and Granger.]

Wood, Albert E.

- 1959a. A New Sciuravid Rodent of the Genus *Pauromys* from the Eocene of Wyoming. *American Museum Novitates*, 1978:1–6, 1 figure. [*Pauromys schaubi*, new species, is described from the red stratum at Twin Buttes (upper Bridger).]

- 1959b. See McGrew, Paul O., Jack E. Berman, Max K. Hecht, John M. Hummel, George G. Simpson, and Albert E. Wood, 1959.
1962. Early Tertiary Rodents of the Family Paramyidae. *Transactions of the American Philosophical Society*, new series, 52(1):1-261, figures 1-91. [A comprehensive description of the known rodents of the Paramyidae. Among forms described as new are the Bridger species *Thisbemys plicatus*, *Leptotomus bridgerensis*, *Microparamys wilsoni*, *Reithroparamys matthewi*, *Ischyrotomus horribilis*, and *Ischyrotomus oweni*.]
- Wood, Horace E. II
1927. Some Early Tertiary Rhinoceroses and Hyracodonts. *Bulletins of American Paleontology*, 13(50):3-105, plates 1-7, tables 1-7. [Discusses *Hyrachyus*, names family Hyrachyidae.]
1934. Revision of the Hyrachyidae. *Bulletin of the American Museum of Natural History*, 67(5):181-295, tables 1-6, plates 20-24. [Detailed descriptions of members of the family Hyrachyidae. Recognizes 4 genera and 10 species of Bridger hyrachyids. Names Black's Fork member of the Bridger formation for Bridger A and B, and Twin Buttes member for Bridger C and D.]
- Wortman, Jacob L.
1894. Osteology of *Patriofelis*, a Middle Eocene Creodont. *Bulletin of the American Museum of Natural History*, 6(5):129-164, figures 1-5, 1 plate. [Detailed description of Bridger material of *Patriofelis*, with figures of skull and skeleton of *P. ferox*.]
- 1901-1902. Studies of Eocene Mammalia in the Marsh Collection, Peabody Museum, Part 1: Carnivora. *American Journal of Science*, 11:333-348, 437-450; 12:143-154, 193-206, 281-296, 377-382, 421-432; 13:39-46, 115-128, 197-206, 433-448; 14:17-23; figures 1-99, 11 plates. [Includes detailed description of Bridger Carnivora with illustrations. Described as new are *Harpagolestes macrocephalus*, new genus and species; *Sinopa minor*, new species; *Sinopa major*, new species; *Thinocyon* (as *Limnocyon*) *medius*, new species; *Viverravus minutus*, new species; *Miacis* (as *Valpavus*) *hargerii*, new species; and *Oödetes herpestoides*, new genus and species.]
- 1903-1904. Studies of Eocene Mammalia in the Marsh Collection, Peabody Museum, Part II: Primates. *American Journal of Science*, 15:163-176, 399-414, 419-436; 16:345-368; 17:23-33, 133-140, 203-214; figures 100-148, plates 11-12. [Includes detailed description of known Bridger primates with illustrations. Described as new are *Metacheiromys marshi*, new genus and species; *Microsyoops schlosseri*, new species; *Smilodectes*, new genus (for *Hyoopsodus gracilis*); and *Uintanius* (as *Omomys*) *ameghini*, new species.]
- Zittel, Karl A. von
1925. Mammalia (revised by Max Schlosser and Sir Arthur Smith Woodward). Volume III in *Text-Book of Paleontology*. Pages i-viii + 1-316, figures 1-374. London: Macmillan and Co. Limited. [Various Bridger genera summarized, some in detail. Some Bridger species figured.]

Publication in *Smithsonian Contributions to Paleobiology*

Manuscripts for serial publications are accepted by the Smithsonian Institution Press, subject to substantive review, only through departments of the various Smithsonian museums. Non-Smithsonian authors should address inquiries to the appropriate department. If submission is invited, the following format requirements of the Press will govern the preparation of copy.

Copy must be typewritten, double-spaced, on one side of standard white bond paper, with 1½" top and left margin, submitted in ribbon copy with a carbon or duplicate, and accompanied by the original artwork. Duplicate copies of all material, including illustrations, should be retained by the author. There may be several paragraphs to a page, but each page should begin with a new paragraph. Number consecutively all pages, including title page, abstract, text, literature cited, legends, and tables. The minimum length is 30 pages, including typescript and illustrations.

The *title* should be complete and clear for easy indexing by abstracting services. Taxonomic titles will carry a final line indicating the higher categories to which the taxon is referable: "(Ammonoidea: Goniatitidae)." Include an *abstract* as an introductory part of the text. Identify the *author* on the first page of text with an unnumbered footnote that includes his professional mailing address. A *table of contents* is optional. An *index*, if required, may be supplied by the author when he returns page proof.

Two *headings* are used: (1) text heads (boldface in print) for major sections and chapters and (2) paragraph sideheads (caps and small caps in print) for subdivisions. Further headings may be worked out with the editor.

In *taxonomic keys*, number only the first item of each couplet; if there is only one couplet, omit the number. For easy reference, number also the taxa and their corresponding headings throughout the text; do not incorporate page references in the key.

In *synonymy*, use the short form (taxon, author, date:page) with a full reference at the end of the paper under "Literature Cited." Begin each taxon at the left margin with subsequent lines indented about three spaces. Within an entry, use a period-dash (.—) to separate each reference. Enclose with square brackets any annotation in, or at the end of, the entry. For *references within the text*, use the author-date system: "(Jones, 1910)" and "Jones (1910)." If the reference is expanded, abbreviate the data: "Jones (1910:122, pl. 20: fig. 1)."

Simple *tabulations* in the text (e.g., columns of data) may carry headings or not, but they should not contain rules. Formal *tables* must be submitted as pages separate from the text, and each table, no matter how large, should be pasted up as a single sheet of copy.

Use the *metric system* instead of, or in addition to, the English system.

Illustrations (line drawings, maps, photographs, shaded drawings) can be intermixed throughout the printed text. They will be termed *Figures* and should be numbered consecutively; however, if a group of figures is treated as a single figure, the components should be indicated by lowercase italic letters on the illustration, in the legend, and in text references: "Figure 9b." If illustrations (usually tone photographs) are printed separately from the text as full pages on a different stock of paper, they will be termed *Plates*, and individual components should be lettered (Plate 9b) but may be numbered (Plate 9: figure 2). Never combine the numbering system of text illustrations with that of plate illustrations. Submit all legends on pages separate from the text and not attached to the artwork. An instruction booklet for the preparation of illustrations is available from the Press on request.

In the *bibliography* (usually called "Literature Cited"), spell out book, journal, and article titles, using initial caps with all words except minor terms such as "and, of, the." For capitalization of titles in foreign languages, follow the national practice of each language. Underscore (for italics) book and journal titles. Use the colon-parentheses system for volume, number, and page citations: "10(2):5-9." Spell out such words as "figures," "plates," "pages."

For *free copies* of his own paper, a Smithsonian author should indicate his requirements on "Form 36" (submitted to the Press with the manuscript). A non-Smithsonian author will receive 50 free copies; order forms for quantities above this amount with instructions for payment will be supplied when page proof is forwarded.

