

# Long prereproductive selection and divergence by depth in a Caribbean candelabrum coral

Carlos Prada<sup>1</sup> and Michael E. Hellberg

Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803

Edited by Mary Jane West-Eberhard, Smithsonian Tropical Research Institute, Ciudad Universitaria, Costa Rica, and approved December 19, 2012 (received for review May 25, 2012)

**Long-lived corals, the foundation of modern reefs, often follow ecological gradients, so that populations or sister species segregate by habitat. Adaptive divergence maintains sympatric congeners after secondary contact or may even generate species by natural selection in the face of gene flow. Such ecological divergence, initially between alternative phenotypes within populations, may be aided by immigrant inviability, especially when a long period separates larval dispersal and the onset of reproduction, during which selection can sort lineages to match different habitats. Here, we evaluate the strength of one ecological factor (depth) to isolate populations by comparing the genes and morphologies of pairs of depth-segregated populations of the candelabrum coral *Eunicea flexuosa* across the Caribbean. *Eunicea* is endemic to the Caribbean and all sister species co-occur. *Eunicea flexuosa* is widespread both geographically and across reef habitats. Our genetic analysis revealed two depth-segregated lineages. Field survivorship data, combined with estimates of selection coefficients based on transplant experiments, suggest that selection is strong enough to segregate these two lineages. Genetic exchange between the Shallow and Deep lineages occurred either immediately after divergence or the two have diverged with gene flow. Migration occurs asymmetrically from the Shallow to Deep lineage. Limited recruitment to reproductive age, even under weak annual selection advantage, is sufficient to generate habitat segregation because of the cumulative prolonged prereproductive selection. Ecological factors associated with depth can act as filters generating strong barriers to gene flow, altering morphologies, and contributing to the potential for speciation in the sea.**

delayed reproduction | ecological speciation | developmental plasticity | conditional phenotypes | octocoral

Natural selection for adaptation to different ecological conditions has become increasingly recognized as an important factor in the formation and maintenance of species (1, 2). Among the ecological factors that may cause adaptive divergence to result in reproductive isolation is immigrant inviability (3). Immigrant inviability results when populations in different environments are locally adapted, so that immigration to the “wrong” habitats results in increased mortality. The effects of this screening process may depend on the life history traits of the species being studied. In species with fast generation times, such as insects and annual plants, immigrant inviability can contribute to reproductive isolation (3). Much less is known about this process and speciation in general in long-lived organisms, which often provide the foundation of species-rich ecosystems.

On coral reefs, the most diverse marine ecosystems, such long-lived foundation species include corals, sponges, and sea fans. Like many other reef-dwelling organisms, the life cycle of these animals includes a planktonic larval stage with great dispersal potential (4), and sister taxa and even more species-rich clades often share the same geographic range (5, 6). The co-occurrence of closely related species with high dispersal capabilities creates a challenge for evolutionary biologists: How can new marine species emerge without obvious geographic isolation? Caribbean coral reefs in particular provide several examples of young clades that have diverged more recently than their immediate cousins in

the Eastern Pacific (6). Such species appear to have been generated by processes acting within the relatively small (~3 million km<sup>2</sup>) confines of the Caribbean.

One explanation for within-Caribbean speciation is that geological processes (e.g., sea level fluctuations) may have transiently divided populations long enough for new species to form. Recently diverged sister species may have then expanded their ranges when conditions allowed and come into secondary contact. For example, the rise of the Isthmus of Panama altered ocean circulation, causing novel conditions in the Caribbean basin that generated both widespread extinction and speciation in Caribbean mollusks and corals (7), species that later expanded their distributions to the entire basin. Geographic isolation is also evident between some populations of widespread taxa at large (1,000 km) (8) and small (<100 km) spatial scales (9, 10). Thus, new species may result from historic geographic isolation within the Caribbean. However, ecological segregation is also widespread on Caribbean coral reefs. Genetic differences have been detected between ecomorphs with overlapping ranges (11–15), and thus habitat diversity may aid marine speciation. For example, the most common coral on Caribbean reefs was regarded as one species, but fertilization, genetic, and morphological data suggest it is a complex of three species segregated in part by depth (5, 16). In fact, depth segregates cryptic species on reefs more often than any other factor (17).

Candelabrum corals of the genus *Eunicea* constitute one such group where sister species are segregated by depth (13). *Eunicea* constitutes the most diverse genus of anthozoans (anemones, corals, and their kin) on Caribbean reefs, with 15 species plus some undescribed forms (18). *Eunicea* species share a mutualistic relationship with algae of the genus *Symbiodinium* clade B (19), which restricts them to the photic zone. All *Eunicea* are gonochoric and reproduce sexually by spawning gametes (18, 20).

*Eunicea flexuosa* is an especially suitable species in which to test the effects of immigrant inviability associated with adaptation to depth in a wide geographical context because (i) its geographic range spans the Caribbean (18), (ii) it is common [ $>5$  colonies/m<sup>2</sup> (21)] and present in virtually all habitats and depths on coral reefs and rocky walls (18), (iii) depth-segregated colonies show extensive internal and external morphological differences (13, 22), (iv) transplant experiments show colonies are adapted to different depths (13), and (v) first reproduction does not occur until a size of 30 cm, at least 15 y after larval dispersal (20). Such delayed reproduction during sedentary asexual growth allows selection to act for a long time before genes can be mixed [as in the strawberry-coral model of Williams (23)] and thus may

Author contributions: C.P. and M.E.H. designed research; C.P. performed research; M.E.H. contributed new reagents/analytic tools; C.P. analyzed data; and C.P. and M.E.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. [KC310499–KC310687](https://doi.org/10.1073/pnas.1208931110) and [KC333998–KC335131](https://doi.org/10.1073/pnas.1208931110)).

<sup>1</sup>To whom correspondence should be addressed. E-mail: [cprada1@lsu.edu](mailto:cprada1@lsu.edu).

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1208931110/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1208931110/-DCSupplemental).


increase the intensity of immigrant inviability between depth-segregated populations.

Here, we use field observations to estimate the fraction of *E. flexuosa* that survive until the onset of reproduction, and then measured survivorship advantage of native over foreign colonies at each depth by reciprocal transplants to show that native colonies have an advantage over foreigners, an advantage magnified by the cumulative effect of long prereproductive selection. We then evaluate morphological and genetic differentiation at two depths across the Caribbean range of *E. flexuosa*. If immigrant inviability operates between depths and is enhanced by delayed reproduction, then genetic divergence will be higher between sympatric populations separated by depth than among those at the same depth separated by geography. We found two depth-segregated lineages, each showing little evidence of subdivision among geographic populations across the Caribbean. Genetic lineages correlate well with morphology but some mismatched individuals are found at low frequency in both lineages. Previously observed adaptive phenotypic differences between depths (13) may thus provide both material for divergent selection and a mechanism for reproductive isolation in *E. flexuosa*. More generally, selection acting on habitat-dependent alternative phenotypes may be especially likely to enhance phenotypic and genetic divergence in long-lived sessile clonal animals with delayed postdispersal reproduction.

## Results

**Survivorship to Reproductive Size is Low in Shallow Areas.** We estimated mean annual survivorship rates (averaged over 4 y) for colonies divided in three size categories (<10, 11–30, and >31 cm). Survivorship rates range from 69.1% for the smallest to 90.1% for the largest category (Table S1). Based on these values, the proportion of individuals (Fig. 1) that would survive to first reproduction (30 cm) is 1% and only 0.01% would reach full maturity (70 cm).

**Selection Is Strong Between Habitats.** We selected 40 adult colonies from shallow and 40 from deep habitats, divided them, and then transplanted the resulting clones (>25 cm) to both depths. We estimated annual survivorship rates based on those observed for the 2 y following transplantation and by pooling our data with previous estimates (13). Mean annual survivorship was highest (Table S2) for colonies transplanted to the same depth (87.1% and 94.0% for shallow and deep, respectively) and lowest for reciprocally transplanted colonies (78.5% and 63.0% for colonies transplanted to shallow and to deep, respectively). We estimated an annual increase in survivorship of native colonies over foreign ones of 9.9% in shallow areas and 33.0% in deep areas. Over the span of a generation (about 40 y), the survivorship advantage would be 98.3% for shallow areas and 99.9% for deep. Assuming a 1:1 ratio of settlers and Hardy–Weinberg proportions, we expect 96% of the mating events in shallow habitats to be between Shallow × Shallow, with only 4% between Shallow × Deep (hybrid crosses). In deep habitats, even fewer hybrid crosses are expected, one in a million.



**Fig. 1.** Age/size survivorship curve for *Eunicea flexuosa* (log scale). Age at first reproduction (20) is shown by the dotted line and full reproductive maturity (70) with the gray bar. Age calculated based on a 2-cm/y growth rate (13).

To compare our selection estimates with other studies, we translated them into difference in fitness between native and foreign colonies as in ref. 24. There are increases of 180% and 190% in fitness of native over foreigners for shallow and deep habitats, respectively. These estimates are based on a growth rate (1.8 cm/y) at the high end of those seen for *E. flexuosa* (13), the size (70 cm) at which colonies significantly contribute to reproduction (20), and adults showing the highest survivorship rates. Estimates based on faster growth rates, reproduction at younger ages, and survivorships from younger colonies result in greater asymmetry in the selective advantage between habitats.

**Two Genetic Lineages Are Segregated by Depth.** We sampled populations from four locations: Bahamas, Panama, Puerto Rico, and Curaçao (Table S3). At each location, we sampled adult colonies (>50 cm) at two depths, as earlier work in Puerto Rico suggested deep and shallow populations might be genetically differentiated (13).


All our analyses of genetic variation indicated unambiguously that there are two lineages within *E. flexuosa* that segregate between deep and shallow depths. Patterns of genetic subdivision revealed by STRUCTURE showed strong support for two clusters (Fig. 2*a*,  $\Delta L > 95\%$ ) that correlated closely with depth (Fig. 2*a*). The probability of membership to either cluster was high (>90%) for most colonies, although it dropped to 89–60% for 20 individuals (11%), perhaps as a result of hybridization between lineages. Similarly, analysis of molecular variance showed that depth explains most of the variation for both mitochondrial and nuclear markers (61% and 71%, respectively). Less variation was explained by geography (7% and 1% for mtDNA and nuclear markers, respectively) and within site variation (31% and 27% for mtDNA and nuclear markers, respectively). All fixation indices were significant (10,000 permutations) at the 0.01 level for both types of markers. Although haplotype networks for each gene did not sort completely (Fig. S1), they revealed two distinct lineages that largely correlate with depth (at least for *MSH* and *I3P*). For *EF1A* and *CT2*, haplotypes sampled from different depths mixed, but clinal variation was still evident.

To test the extent to which depth, geography, and their interaction have played roles in lineage splitting and to recover the evolutionary history of all populations, we used a combined gene trees/species tree (STEM) (25) and model-based inference approach. The Shallow and Deep lineages were well supported by STEM. The tree with the best corrected Akaike information criterion (AICc) score showed two lineages: one composed of all Shallow samples and one of all Deep (Fig. 2*c*). When we further split populations by depth and geography, the best tree still joins populations by depth, but was 218 times less likely than the all Shallow and all Deep tree. Conversely, if we grouped populations by geography (i.e., combine Panama Shallow and Panama Deep), the tree with the highest likelihood score was >1,000 times worse than the best tree.

Depth segregation was significant (Fisher's exact test,  $P < 0.0001$ ), but the degree of segregation varied among sites (Fig. 2*b*). In Puerto Rico and Bahamas, it was high, with >80% of colonies of each lineage segregated into either shallow or deep habitats. In Curaçao, the Shallow lineage dominated (100%) shallow environments, but was also common (50%) in deep areas. Thus, although the two lineages segregate over a depth gradient, the absolute depths defining the gradient vary among localities. In Puerto Rico, our sampling captured the ends, but in Curaçao and Bahamas the corresponding deep habitat may be at greater depths.

Panama was an extreme case in this regard: the Deep lineage dominated both deep and shallow environments, with the Shallow lineage present at only modest levels in shallow environments (even after adding an extra shallow collecting site). This Panamanian oddity was not unexpected: particulate matter around Bocas del Toro is high, which increases food for suspension feeders and attenuates light, allowing deep-water suspension feeders such





**Fig. 4.** Mean spindle length between Deep (black) and Shallow (light gray) lineages in different populations. The letters indicate significant differences for within lineage comparisons. The different letters indicate significant differences in an honestly significant difference (HSD) Tukey test after Bonferroni correction. D, Deep; S, Shallow.

the calcium carbonate particles that provide structure to the colony. Spindle length correlates well with other morphological characters (e.g., calice size, intercalice distance, branch thickness) and is the most divergent morphological character between deep and shallow colonies of *E. flexuosa* in Puerto Rico (13).

Patterns of morphological differentiation mirrored those seen genetically; lineage was a better predictor of spindle length than either geography or habitat (Table 1). All of the four best models had lineage as a variable, along with its interaction with either geography or habitat. When we excluded lineage or its interactions as variables, the remaining models were 100 times less likely than the best model with lineage included. Lineage fit the data better than depth, despite the two being highly correlated (Table 1). Although morphology and genetics match well, mismatched individuals occur at low frequency (2 of 97 and 7 of 92 for Shallow and Deep, respectively).

Once lineage is accounted for, some differences between locations remained (Fig. 4). For the Shallow lineage, colonies from Curaçao had larger spindles than colonies in Puerto Rico. For the Deep lineage, colonies in Panama had smaller spindles than for all other locations. This difference in spindle length in Panama made the morphological boundary between Shallow and Deep narrower and shifted shallower than in other locations. Thus, morphologies were diagnosable, correlated well with lineage, and segregated into shallow and deep habitats at local scales, but lineage by depth by locale interactions generated morphological overlap between taxa when populations were considered together.

## Discussion

Our study found that (i) selection is strong in both habitats for native over foreign colonies, (ii) colonies of the candelabrum coral

**Table 1. Model ranking for spindle length analysis using a general linear model**

Model	AIC	$\Delta_i$	ML	wi	Evidence ratio
GDL L*G	165.56		1.00	0.61	
GDL D*L G*L	167.31	1.75	0.42	0.25	2
GDL D*L G*D	170.47	4.91	0.09	0.05	12
GDL G*L G*D	171.08	5.52	0.06	0.04	16
Full model	171.08	5.52	0.06	0.04	16
GDL D*G	175.15	9.59	0.01	0.01	121
GL G*L	185.25	19.69	0	0	18,864
All other models	>188	>20	0	0	>10 <sup>5</sup>

G, geography; D, depth; L, lineage assignment from STRUCTURE clustering for  $K = 2$ ; ML, maximum likelihood; AIC, Akaike information criterion inferred from the generalized linear model fitting;  $\Delta_i$ , difference in AIC score with respect to the best model; model likelihoods, relative likelihood of the model given the data; wi, model probabilities; evidence ratio, fold difference in model probabilities against the best model.

*Eunicea flexuosa* occurring in shallow and deep habitat are segregated into two genetically and morphologically divergent lineages, (iii) genetic divergence is weak among geographic samples within each lineage, (iv) the Shallow and Deep lineages have exchanged genes since their initial divergence, and (v) migration between them has been higher from Shallow to Deep. Reciprocal transplantation suggests divergence between colonies found at different depths is adaptive and selection is strong enough to maintain the separation of the two lineages at the ends of the depth gradient. This divergent selection may by itself result in reproductive isolation via immigrant inviability (poor performance and viability of larvae settling at the wrong depth) and temporal isolation (spawning at different times) (29). We suggest divergence between Shallow and Deep lineages of *E. flexuosa* is driven by adaptation to different conditions at different depths (1), facilitated by developmental plasticity (30, 31) and prolonged opportunity for depth-specific selection before reproduction.

**Ecological Gradients, Adaptive Divergence, and Speciation.** Divergent selection along depth gradients is generated by associated factors such as the quantity and quality of light, the force of waves and currents, sediment load, the distribution of predators and mutualists, and the availability and composition of food. In response to this variation in environmental conditions, closely related species or populations found at different depths have been found to differ in their morphologies (5, 13), feeding strategies (32, 33), metabolic rates (33), symbiotic relationships (34), resistance to predators (35), and spawning times (16, 29). In *E. flexuosa*'s close relative *Plexaura homomalla* (*Eunicea* and *Plexaura* are paraphyletic), depth-segregated sister species increase heterotrophy and decrease photosynthesis as depth increases (32).

Such depth-related divergence seems to be correlated with the differentiation of closely related species, as in bryozoans (36). Depth gradients in light are hypothesized as a cause of speciation by sensory drive in cichlids (37) and factors associated with depth appear to have driven speciation in rockfish (38). Among marine sibling species reported by Knowlton (17), over one-half involved depth as a dividing factor, even though not all comparisons included depth. Thus, adaptive divergence by depth is consistent with a role for depth in maintaining and generating species differences in marine and aquatic settings.

These patterns associated with depth are analogous to those seen for other environmental gradients, such as altitude, salinity, or temperature. Such gradients generate adaptive divergence in vertebrates (39), invertebrates (40), and plants (41). Because such abiotic factors generate adaptive divergence across multiple species, they also generate gradients in biotic interactions (such as predator-prey interactions) (42), creating additional axes of complexity.

**High Gene Flow Within Species.** Divergence between depths at small scales (as little 200 m separate deep and shallow populations we sampled) in *E. flexuosa* contrasts with the high connectivity seen among populations within lineages across thousands of kilometers. Such high connectivity is not uncommon to marine species with planktonic larval dispersal (4), ranging to extremes in which gene flow spans the broadest expanses of the Pacific Ocean (43). Thus, adaptive divergence occurs here despite the potential for high levels of gene flow (39). Such adaptive divergence under high gene flow generates a pattern where neutral genetic divergence is primarily partitioned by habitats, with little geographic structure. Ecotypes are connected across vast geographical spans, thus producing a pattern of single ecological segregation as seen in *E. flexuosa* and shared by other broadcast spawners such as the star coral *Montastraea annularis* (5), some coral reef fishes (14), and the Hawaiian limpets *Cellana* (44).

In contrast, under low gene flow, genetic divergence first occurs geographically and then ecologically. The pattern has been detailed in the marine snail *Littorina saxatilis* (45) and is similar to those in other marine brooders where ecology has been proposed to drive speciation, such as the corals *Favia fragum* (15) and *Seriatopora hystrix* (46). Clonal brooders also

tend to reproduce at younger ages (smaller sizes) (47), increasing the role of geography in local adaptation. This difference between brooders and broadcasters has parallels in winged and wingless phytophagous insects. In pea aphids, segregation by host plant occurs at large geographical scales with little geographical differentiation (48). Conversely, in walking sticks, pairs of habitat-segregated ecotypes have occurred multiple times (3). Thus, the partition of genetic diversity during ecological speciation depends on the dispersal biology of the studied species.

**Divergence Between Habitat-Segregated Populations with Prolonged Prereproductive Selection.** Ecological filtering can help maintain species differences and contribute to reproductive isolation of populations occurring in different habitats (3). In *E. flexuosa*, adult colonies native to their depth survive far better than do foreign colonies, an advantage enhanced by the cumulative effect of many years of prereproductive selection. For a settling larva to reach adulthood, it must pass a series of ecological challenges until it reaches a safe size of at least 30 cm (>15 y) and begins reproductive activity. Significant genetic contributions to the population likely take far longer, however, not until attaining a colony size of 70 cm (>35 y), because such large colonies contribute 98% of all eggs in *E. flexuosa* (20).

The difference in fitness between native and foreign colonies that we found here is three times higher than any reported for similar transplant experiments (24). It also exceeds values used in mathematical models of ecological speciation in which adaptive divergence is sufficient to generate reproductive isolation in discrete habitats (49, 50) or along clines (51, 52). The long period between when a larva settles and when it becomes a colony that contributes genetically to the next generation thus decreases gene flow between depth-adapted lineages and enhances the opportunity for habitat-specific selection and divergent adaptation. Such immigrant inviability may be common in slowly maturing sessile species with broad dispersal [like some scrubs and trees (53, 54)]. In contrast, when adults are the primary dispersers, immigrant inviability would have to be exceptionally strong to have such an effect, as migrants can reproduce as soon as they settle, allowing locally adapted genotypes to mix with maladapted foreign ones.

Strong divergent selection between habitats could also stifle the early stages of divergence if dispersal is limited and a narrow range of phenotypes limit opportunities to colonize new habitats (31). In *E. flexuosa*, however, larval dispersal is high (Fig. 2) and colonies quite phenotypically plastic (13).

Temporal isolation may also emerge from adaptation to different depths and serve to further isolate depth-segregated populations. As for flowering plants segregated by altitude (41, 55), marine broadcasters segregated by depth often differ in their time of spawning, decreasing the probability of cross-fertilization between depth-segregated colonies (16, 29). Before spawning events, colonies in deep areas of *E. flexuosa* lack mature eggs compared with shallow ones (22). One-hour difference in the timing of spawning is sufficient for sperm to dilute to the point where crossbreeding is unlikely in other corals (16, 29). Although the exact traits that affect ecological fitness are unknown, population segregation by depth thus has the potential to increase reproductive isolation by pleiotropic interactions (31, 56) between fitness traits and those that control timing of reproduction, increasing the chance of genetic divergence in depth-segregated populations adapted to different ecological conditions.

**Speciation by Prolonged Selection on Adaptively Plastic Habitat-Specific Contrasting Forms: Divergence in the Sea.** Our data on *E. flexuosa* show how a developmentally plastic organism with ecologically variable phenotypes may diverge genetically to produce different species without geographic isolation, even in groups with widely dispersing propagules.

Habitat-specific alternative phenotypes, whether discrete or at the ends of a cline, represent different habitat-mediated sets of expressed genes (57). To the degree that they are independently expressed, they are independently exposed to habitat-specific

selection and can, given genetic variation in their form (57, 58), diverge genetically without reproductive isolation (30). Then, due to genetic accommodation or assimilation of the contrasting phenotypes (30, 58), they may diverge genetically such that migrants between habitats become effectively inviable, as suggested by our data and as hypothesized for speciation between alternative phenotypes (30, 59) and in clines (51, 52). A broader analysis of the evolution of plasticity in the genus *Eunicea* would provide a framework to further test this hypothesis.

As we discuss here, habitat-related divergence may apply especially in organisms with a long period of prereproductive selection, which affords an increased “opportunity for selection” (60), and in organisms that are resident for a prolonged period in a single place. In sessile clonal animals, independent selection on condition-sensitive (depth-sensitive) phenotypes is enhanced because immotile colonies cannot escape the selective factors of a given stable habitat. Selection is intense in juveniles and decreases exponentially with colony size, so that clonal organisms delay (for a couple of decades in *E. flexuosa*) sexual reproduction and allocate all resources to vegetative growth (23). Even once these younger colonies begin to reproduce, most gametes in a population will still come from the large, old (over one-half a century), well-adapted colonies that dominate sexual reproduction. These conditions characterize *E. flexuosa* and other sessile clonal organisms, such as anthozoans, sponges, long-lived trees (e.g., oaks and pines), and shrubs (e.g., *Ceanothus*) (23, 61), which invest heavily in vegetative growth before reproduction (23, 61). This process of habitat-related divergence without reproductive isolation may therefore be more common than usually appreciated. It can occur whether the contrasting phenotypes are initially genetically cued (51, 52) or environmentally induced (13, 30, 62), for it is the contrast between the different phenotypes and the underlying sets of expressed genes that, under selection, drives divergence—not the cues (genetic or environmental) that govern their expression.

Taken together, the combination of habitat-responsive plasticity, long prereproductive selection, sessile residence in distinctive habitats, and the potential for temporal separation of reproduction in different habitats, exemplified in *E. flexuosa*, may help solve the long-standing question of how speciation can occur in marine populations with widely dispersing larval forms.

## Materials and Methods

We estimated annual survivorship per size class in 1-m<sup>2</sup> quadrats. We monitored 161 quadrats spread across eight reefs in Puerto Rico over 4 y (*SI Materials and Methods*). We sampled survivorship advantage of native over foreign by reciprocally transplanting 40 adult (>25 cm) colonies from each depth to both shallow and deep areas (80 colonies per depth). Survivors were recorded annually for 2 y.

We sampled populations in Bahamas, Panama, Puerto Rico, and Curaçao for genetic analysis. At each location, we sampled 17–38 adult colonies (>50 cm) from each of two depths (Table S3). We developed three new nuclear sequence markers (Table S7) using sequences generated from a partial 454 run. Sequences from these three and a previously described mtDNA marker were generated using standard PCR procedures (*SI Materials and Methods*). We edited and assembled all sequences using standard procedures. We resolved haplotypes probabilistically (*SI Materials and Methods*) and corroborated inferred haplotypes by directly cloning 10% of our sequences. We recoded sequences as frequency data and used them to infer population subdivision using Bayesian clustering in STRUCTURE (63). We ran STRUCTURE without information of the origin of each individual, thereby reducing potential biases. We used the STRUCTURE cluster assignment for two subsequent analyses: STEM and IM.

We inferred gene genealogies and estimated likelihood scores for species trees using STEM 2.0 (25). We manipulated the settings file to estimate the likelihood score for the best tree for each of three arrangements: (i) populations divided by geography (four lineages), (ii) populations divided by depth (two lineages), and (iii) each population as an independent lineage (eight lineages). We then used the average likelihood scores across replicates to infer information theory statistics and rank all possible topologies (64).

We used IMA, the isolation with migration model (65), and model-based inference (64) to evaluate all possible scenarios of divergence. Our aim was to compare the support for strict isolation (zero migration) of the Shallow and

Deep lineages (as delineated by STRUCTURE) to that for a model of divergence with gene flow. We also evaluated the IMA model when individuals were partitioned by habitat and morphology. We then used model-based selection to calculate evidence ratios and rank all possible models (64).

We took images of each colony through a Diagnostic Instrument SPOT RT Slider CCD camera attached to a Leica MZ7 stereomicroscope. All morphological measurements were carried out at the Louisiana State University Socolofsky Microscopy Center. We fit the spindle length to a generalized linear model. We used a Gaussian distribution and geography, depth, and lineage as factors. We constructed all 17 possible models and used model based approaches to rank each model (64). All statistical tests were carried out in R.

- Schluter D (2009) Evidence for ecological speciation and its alternative. *Science* 323(5915):737–741.
- Darwin C (1859) *On the Origin of Species* (John Murray, London).
- Nosil P, Vines TH, Funk DJ (2005) Perspective: Reproductive isolation caused by natural selection against immigrants from divergent habitats. *Evolution* 59(4):705–719.
- Hellberg M (2009) Gene flow and isolation among populations of marine animals. *Annu Rev Ecol Syst* 40:291–310.
- Fukami H, et al. (2004) Geographic differences in species boundaries among members of the *Montastraea annularis* complex based on molecular and morphological markers. *Evolution* 58(2):324–337.
- Taylor MS, Hellberg ME (2005) Marine radiations at small geographic scales: Speciation in neotropical reef gobies (*Elacatinus*). *Evolution* 59(2):374–385.
- Jackson JBC, Jung P, Coates AG, Collins LS (1993) Diversity and extinction of tropical american mollusks and emergence of the isthmus of panama. *Science* 260(5114):1624–1626.
- Baums IB, Miller MW, Hellberg ME (2005) Regionally isolated populations of an imperiled Caribbean coral, *Acropora palmata*. *Mol Ecol* 14(5):1377–1390.
- DeBiasse MB, Richards VP, Shiyi MS (2010) Genetic assessment of connectivity in the common reef sponge, *Callyspongia vaginalis* (Demospongiae: Haplosclerida) reveals high population structure along the Florida reef tract. *Coral Reefs* 29(1):47–55.
- Taylor MS, Hellberg ME (2003) Genetic evidence for local retention of pelagic larvae in a Caribbean reef fish. *Science* 299(5603):107–109.
- Duran S, Rützler K (2006) Ecological speciation in a Caribbean marine sponge. *Mol Phylogenet Evol* 40(1):292–297.
- Brazeau DA, Harvell CD (1994) Genetic structure of local populations and divergence between growth forms in a clonal invertebrate, the Caribbean octocoral *Briareum asbestinum*. *Mar Biol* 119(1):53–60.
- Prada C, Schizas NV, Yoshioka PM (2008) Phenotypic plasticity or speciation? A case from a clonal marine organism. *BMC Evol Biol* 8:47.
- Rocha LA, Robertson DR, Roman J, Bowen BW (2005) Ecological speciation in tropical reef fishes. *Proc Biol Sci* 272(1563):573–579.
- Carlson DB, Budd AF (2002) Incipient speciation across a depth gradient in a scleractinian coral? *Evolution* 56(11):2227–2242.
- Knowlton N, Mate JL, Guzman HM, Rowan R, Jara J (1997) Direct evidence for reproductive isolation among the three species of the *Montastraea annularis* complex in Central America (Panama and Honduras). *Mar Biol* 127(4):705–711.
- Knowlton N (1993) Sibling species in the sea. *Annu Rev Ecol Syst* 24:189–216.
- Sanchez J (2009) Systematics of the candelabrum gorgonian corals (*Eunicea Lamouroux*; Plexauridae; Octocorallia; Cnidaria). *Zool J Linn Soc* 157(2):237–263.
- Santos SR, Taylor DJ, Coffroth MA (2001) Genetic comparisons of freshly isolated vs. cultured zooxanthellae: Implications for extrapolating to the intact symbiosis. *J Phycol* 37(5):900–912.
- Beiring EA, Lasker HR (2000) Egg production by colonies of a gorgonian coral. *Mar Ecol Prog Ser* 196:169–177.
- Yoshioka P (2009) Sediment transport and the distribution of shallow-water gorgonians. *Caribb J Sci* 45(2):254–259.
- Kim E, Lasker HR, Coffroth MA, Kim K (2004) Morphological and genetic variation across reef habitats in a broadcast-spawning octocoral. *Hydrobiologia* 530/531:423–432.
- Williams G (1975) *Sex and Evolution* (Princeton Univ Press, Princeton).
- Hereford J (2009) A quantitative survey of local adaptation and fitness trade-offs. *Am Nat* 173(5):579–588.
- Kubatko LS, Carstens BC, Knowles LL (2009) STEM: Species tree estimation using maximum likelihood for gene trees under coalescence. *Bioinformatics* 25(7):971–973.
- Opreko DM, Sanchez JA (2005) Caribbean shallow-water black corals (Cnidaria: Anthozoa: Antipatharia). *Caribb J Sci* 41(3):492–507.
- Guzmán HM, Guevara CA (1999) Coral reefs of Bocas del Toro, Panama: III. Distribution, structure, diversity and conservation status of reefs in Pastores, Cristobal, Popa and Cayo Agua islands. *Rev Biol Trop* 47(2):659–676.
- Lepais O, et al. (2009) Species relative abundance and direction of introgression in oaks. *Mol Ecol* 18(10):2228–2242.
- Levitan DR, et al. (2004) Mechanisms of reproductive isolation among sympatric broadcast-spawning corals of the *Montastraea annularis* species complex. *Evolution* 58(2):308–323.
- West-Eberhard MJ (2003) *Developmental Plasticity and Evolution* (Oxford Univ Press, New York).
- Fitzpatrick BM (2012) Underappreciated consequences of phenotypic plasticity for ecological speciation. *Int J Ecol* 2012:256017.
- Lasker HR, Gottfried MD, Coffroth MA (1983) Effects of depth on the feeding capabilities of two octocorals. *Mar Biol* 73(1):73–78.
- Lesser MP, et al. (2010) Photoacclimatization by the coral *Montastraea cavernosa* in the mesophotic zone: Light, food, and genetics. *Ecology* 91(4):990–1003.
- Kirk NL, Andras JP, Harvell CD, Santos SR, Coffroth MA (2009) Population structure of *Symbiodinium* sp. associated with the common sea fan, *Gorgonia ventalina*, in the Florida Keys across distance, depth, and time. *Mar Biol* 156(8):1609–1623.
- West JM, Harvell CD, Walls AM (1993) Morphological plasticity in a gorgonian coral (*Briareum asbestinum*) over a depth cline. *Mar Ecol Prog Ser* 94:61–69.
- Jackson J, McKinney F (1990) *Ecological Processes and Progressive Macroevolution of Marine Clonal Benthos. Causes of Evolution*, eds Ross R, Allmon W (Univ of Chicago Press, Chicago), pp 173–209.
- Seehausen O, et al. (2008) Speciation through sensory drive in cichlid fish. *Nature* 455(7213):620–626.
- Ingram T (2011) Speciation along a depth gradient in a marine adaptive radiation. *Proc Biol Sci* 278(1705):613–618.
- Whitehead A, Roach JL, Zhang S, Galvez F (2011) Genomic mechanisms of evolved physiological plasticity in killifish distributed along an environmental salinity gradient. *Proc Natl Acad Sci USA* 108(15):6193–6198.
- Sanford E, Kelly MW (2011) Local adaptation in marine invertebrates. *Annu Rev Mar Sci* 3:509–535.
- Hodges SA, Arnold ML (1994) Floral and ecological isolation between *Aquilegia formosa* and *Aquilegia pubescens*. *Proc Natl Acad Sci USA* 91(7):2493–2496.
- Freeman AS, Byers JE (2006) Divergent induced responses to an invasive predator in marine mussel populations. *Science* 313(5788):831–833.
- Lessios HA, Robertson DR (2006) Crossing the impassable: Genetic connections in 20 reef fishes across the eastern Pacific barrier. *Proc Biol Sci* 273(1598):2201–2208.
- Bird CE, Holland BS, Bowen BW, Toonen RJ (2011) Diversification of sympatric broadcast-spawning limpets (*Cellana* spp.) within the Hawaiian archipelago. *Mol Ecol* 20(10):2128–2141.
- Johannesson K (2001) Parallel speciation: A key to sympatric divergence. *Trends Ecol Evol* 16(3):148–153.
- van Oppen MJ, Bongaerts P, Underwood JN, Peplow LM, Cooper TF (2011) The role of deep reefs in shallow reef recovery: An assessment of vertical connectivity in a brooding coral from west and east Australia. *Mol Ecol* 20(8):1647–1660.
- Szmat AM (1986) Reproductive ecology of Caribbean reef corals. *Coral Reefs* 5:43–54.
- Ferrari J, West JA, Via S, Godfray HC (2012) Population genetic structure and secondary symbionts in host-associated populations of the pea aphid complex. *Evolution* 66(2):375–390.
- Thibert-Plante X, Hendry AP (2011) The consequences of phenotypic plasticity for ecological speciation. *J Evol Biol* 24(2):326–342.
- Giraud T, Gladieux P, Gavrilets S (2010) Linking the emergence of fungal plant diseases with ecological speciation. *Trends Ecol Evol* 25(7):387–395.
- Endler JA (1977) *Geographic Variation, Speciation, and Clines* (Princeton Univ Press, Princeton).
- Clarke BC (1966) The evolution of morpho-ratio clines. *Am Nat* 100(914):389–402.
- Savolainen V, et al. (2006) Sympatric speciation in palms on an oceanic island. *Nature* 441(7090):210–213.
- Alberto F, et al. (2010) Population differentiation of sessile oak at the altitudinal front of migration in the French Pyrenees. *Mol Ecol* 19(13):2626–2639.
- Hall MC, Willis JH (2006) Divergent selection on flowering time contributes to local adaptation in *Mimulus guttatus* populations. *Evolution* 60(12):2466–2477.
- Servedio MR, Van Doorn GS, Kopp M, Frame AM, Nosil P (2011) Magic traits in speciation: “Magic” but not rare? *Trends Ecol Evol* 26(8):389–397.
- Moczek AP, et al. (2011) The role of developmental plasticity in evolutionary innovation. *Proc Biol Sci* 278(1719):2705–2713.
- Waddington CH (1942) Canalization of development and the inheritance of acquired characters. *Nature* 150(3811):563–565.
- Pfennig DW, et al. (2010) Phenotypic plasticity’s impacts on diversification and speciation. *Trends Ecol Evol* 25(8):459–467.
- Crow J (1961) Mechanisms and trends in human evolution. *Daedalus* 90:416–431.
- Connell J (1973) Population ecology of reef-building corals. *Biology and Geology of Coral Reefs*, eds Jones A, Endean R (Academic, London), pp 205–245.
- Scoville AG, Pfrender ME (2010) Phenotypic plasticity facilitates recurrent rapid adaptation to introduced predators. *Proc Natl Acad Sci USA* 107(9):4260–4263.
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155(2):945–959.
- Anderson DR (2008) *Model Based Inference in the Life Sciences* (Springer, New York).
- Hey J, Nielsen R (2004) Multilocus methods for estimating population sizes, migration rates and divergence time, with applications to the divergence of *Drosophila pseudoobscura* and *D. persimilis*. *Genetics* 167(2):747–760.