



SMITHSONIAN INSTITUTION
UNITED STATES NATIONAL MUSEUM

Bulletin 100

VOLUME 4

CONTRIBUTIONS TO THE BIOLOGY OF THE
PHILIPPINE ARCHIPELAGO AND
ADJACENT REGIONS

FORAMINIFERA OF THE PHILIPPINE
AND ADJACENT SEAS

JOSEPH A. CUSHMAN

Of the Boston Society of Natural History



WASHINGTON
GOVERNMENT PRINTING OFFICE

1921

ADVERTISEMENT.

The scientific publications of the United States National Museum consist of two series, the *Proceedings* and the *Bulletins*.

The *Proceedings*, the first volume of which was issued in 1878, are intended primarily as a medium for the publication of original, and usually brief, papers based on the collections of the National Museum, presenting newly acquired facts in zoology, geology, and anthropology, including descriptions of new forms of animals, and revisions of limited groups. One or two volumes are issued annually and distributed to libraries and scientific organizations. A limited number of copies of each paper, in pamphlet form, is distributed to specialists and others interested in the different subjects, as soon as printed.

The dates of publication are recorded in the tables of contents of the volumes.

The *Bulletins*, the first of which was issued in 1875, consist of a series of separate publications comprising chiefly monographs of large zoological groups and other general systematic treatises (occasionally in several volumes), faunal works, reports of expeditions, and catalogues of type-specimens, special collections, etc. The majority of the volumes are octavos, but a quarto size has been adopted in a few instances in which large plates were regarded as indispensable.

Since 1902 a series of octavo volumes containing papers relating to the botanical collections of the Museum, and known as the *Contributions from the National Herbarium*, has been published as bulletins.

The present work forms No. 100, volume 4, of the *Bulletin* series.

WILLIAM DE C. RAVENEL,

*Administrative Assistant to the Secretary,
in charge of the United States National Museum.*

WASHINGTON, D. C., June, 1921.

TABLE OF CONTENTS.

	Page.
Introduction.....	21
Systematic part.....	35
Order Foraminifera.....	35
Family Astrorhizidae.....	35
Subfamily Astrorhizinae.....	35
Genus Astrorhiza.....	35
angulosa.....	35
granulosa.....	35
agglutinata.....	36
Genus Rhabdammina.....	36
abyssorum.....	36
var. radiata.....	37
irregularis.....	38
discreta.....	39
linearis.....	40
Genus Marsipella.....	40
gigantea.....	40
cylindrica.....	41
Genus Bathysiphon.....	41
filiformis.....	41
rufus.....	42
rufescens.....	43
flavidus.....	44
var. giganteus.....	44
papyraceus.....	45
Genus Rhizammina.....	45
indivisa.....	45
algaeformis.....	46
Subfamily Saccammininae.....	46
Genus Psammosphaera.....	46
fusca.....	46
parva.....	47
Genus Iridia.....	47
diaphana.....	47
Genus Saccammina.....	48
sphaerica.....	48
var. catenulata.....	48
Genus Proteonina.....	49
difflugiformis.....	49
fusiformis.....	49
testacea.....	50
Genus Technitella.....	50
legumen.....	50
Genus Tholosina.....	51
bulla.....	51

Systematic part—Continued.

Family Astrorhizidae—Continued.

	Page.
Subfamily Saccamininae—Continued.	
Genus <i>Thurammina</i>	52
<i>papyracea</i>	52
<i>papillata</i> , var. <i>castanea</i>	52
Subfamily Hyperammininae.....	53
Genus <i>Hyperammina</i>	53
<i>elongata</i>	53
<i>friabilis</i>	54
Genus <i>Saccorhiza</i>	54
<i>ramosa</i>	54
Genus <i>Tolypammina</i>	55
<i>vagans</i>	55
Genus <i>Dendrophrya</i>	56
<i>ramosa</i>	56
var. <i>robusta</i>	57
<i>attenuata</i>	57
Genus <i>Haliphysema</i>	58
<i>catenulata</i>	58
Genus <i>Jaculella</i>	59
<i>acuta</i>	59
<i>obtusa</i>	59
Genus <i>Sagenina</i>	60
<i>frondescens</i>	60
<i>divaricans</i>	60
Subfamily Ammodiscinae.....	61
Genus <i>Ammolagena</i>	61
<i>clavata</i>	61
Genus <i>Ammodiscus</i>	62
<i>incertus</i>	62
var. <i>discoideus</i>	63
<i>exsertus</i>	64
Family Lituolidae.....	64
Subfamily Aschemonellinae.....	64
Genus <i>Aschemonella</i>	64
<i>catenata</i>	64
Subfamily Reophacinae.....	65
Genus <i>Reophax</i>	65
<i>scorpiurus</i>	65
<i>pilulifer</i>	66
<i>distans</i>	66
<i>bacillaris</i>	67
<i>dentaliniformis</i>	68
<i>nodulosus</i>	68
var. <i>hormosinoides</i>	69
<i>guttifer</i>	69
<i>aduncus</i>	70
<i>cylindricus</i>	70
<i>pseudobacillaris</i>	71
<i>findens</i>	71
<i>spiculotestus</i>	72
<i>horridus</i>	73
<i>agglutinatus</i>	73
<i>bilocularis</i>	74

Systematic part—Continued.

Family Lituolidae—Continued.

Subfamily Aschemonellinae—Continued.

	Page.
Genus <i>Hormosina</i>	74
<i>globulifera</i>	74
<i>ovicula</i>	75
<i>normanii</i>	75
<i>ovaliformis</i>	76
<i>elongata</i>	76
<i>carpenteri</i>	77
Genus <i>Ilaplostiche</i>	78
<i>dubia</i>	78
Subfamily Trochammininae.....	79
Genus <i>Trochamminoides</i>	79
<i>proteus</i>	79
Genus <i>Haplophragmoides</i>	79
<i>canariensis</i>	79
<i>emaciatum</i>	80
<i>scitulum</i>	80
<i>subglobosum</i>	81
<i>ringens</i>	81
<i>grandiformis</i>	82
<i>sphaeriloculum</i>	83
Genus <i>Cribrostomoides</i>	83
<i>bradyi</i>	83
Genus <i>Cyclammmina</i>	84
<i>cancellata</i>	84
<i>compressa</i>	85
<i>pauciloculata</i>	86
<i>pusilla</i>	87
<i>bradyi</i>	88
Genus <i>Lituotuba</i>	88
<i>lituiformis</i>	88
Genus <i>Ammobaculites</i>	89
<i>agglutinans</i>	89
<i>tenuimargo</i>	90
<i>calcareum</i>	90
<i>cassis</i>	91
<i>cylindricus</i>	92
<i>reophaeiformis</i>	92
<i>pseudospirale</i>	94
Genus <i>Placopsilina</i>	95
<i>cenomana</i>	95
Genus <i>Trochammina</i>	95
<i>squamata</i>	95
<i>turbinata</i>	96
<i>globigeriniformis</i>	96
Genus <i>Ammochilostoma</i>	97
<i>rotundata</i>	97
Genus <i>Ammosphaeroidina</i>	98
<i>sphaeroidiniformis</i>	98
<i>grandis</i>	98
Genus <i>Sphaerammina</i>	99
<i>ovalis</i>	100

Systematic part—Continued.

Family Lituolidae—Continued.

Subfamily Trochammininae—Continued.

	Page.
Genus <i>Ammosphaerulina</i>	100
<i>adhaerens</i>	100
Genus <i>Nouria</i>	101
<i>harrisii</i>	101
<i>compressa</i>	102

Family Textulariidae..... 102

Subfamily Spiroplectinae..... 102

Genus <i>Spiroplectina</i>	102
<i>bulbosa</i>	102

Subfamily Textulariinae..... 103

Genus <i>Textularia</i>	103
<i>sagittula</i>	103
var. <i>atrata</i>	103
var. <i>fistulosa</i>	104
<i>gramen</i>	105
<i>hauerii</i>	105
<i>agglutinans</i>	106
var. <i>fistula</i>	107
<i>stricta</i>	107
<i>candeiana</i>	109
<i>porrecta</i>	109
<i>vertebralis</i>	110
<i>aperturalis</i>	111
<i>catenata</i>	112
<i>aspera</i>	112
<i>goësii</i>	113
<i>flintii</i>	113
<i>rugosa</i>	114
<i>siphonifera</i>	115
<i>semialata</i>	116
<i>foliacea</i>	117
<i>immensa</i>	118
<i>concava</i>	119
<i>jugosa</i>	119
<i>excavata</i>	120
<i>transversaria</i>	120
<i>crassisepta</i>	121
<i>pseudocarinata</i>	121
<i>abbreviata</i>	122
<i>conica</i>	123
<i>trochus</i>	124
<i>quadrilatera</i>	125
Genus <i>Bigenerina</i>	125
<i>nodosaria</i>	125
<i>arenacea</i>	126
<i>capreolus</i>	127
<i>pennatula</i>	127
Genus <i>Bolivina</i>	128
<i>dilatata</i>	128
<i>beyrichii</i>	128
var. <i>alata</i>	129

Systematic part—Continued.

Family Textulariidae—Continued.

Subfamily Textulariinae—Continued.

Genus *Bolivina*—Continued.

	Page.
robusta	129
schwageriana	130
nobilis	130
karreriana	131
var. <i>carinata</i>	132
hantkeniana	132
amygdalaeformis	133
semicostata	134
sculpturata	134
subangularis	135
limbata	135
pygmaea	136
punctata	136
compacta	137
aenariensis	137
pusilla	138
Genus <i>Pleurostomella</i>	138
<i>alternans</i>	138
<i>subnodosa</i>	139
Subfamily Verneuilininae	139
Genus <i>Verneuilina</i>	139
<i>polystropha</i>	139
<i>propinqua</i>	140
<i>bradyi</i>	141
<i>spinulosa</i>	141
<i>triquetra</i>	142
<i>affixa</i>	142
Genus <i>Valvulina</i>	142
<i>conica</i>	142
<i>fusca</i>	143
Genus <i>Chrysalidina</i>	144
<i>dimorpha</i>	144
Genus <i>Tritaxia</i>	145
<i>tricarinata</i>	145
<i>indiscreta</i>	145
Genus <i>Gaudryina</i>	146
<i>scabra</i>	146
<i>flintii</i>	146
<i>quadrangularis</i>	147
<i>triangularis</i>	148
<i>paupercula</i>	148
<i>bradyi</i>	149
<i>baccata</i>	150
<i>chilostoma</i>	150
<i>pseudofiliformis</i>	151
<i>apicularis</i>	151
<i>attenuata</i>	152
<i>robusta</i>	152
Genus <i>Tritaxilina</i>	153
<i>caperata</i>	153

Systematic part—Continued.

Family Textulariidae—Continued.

Subfamily Verneuiliniinae—Continued.

	Page.
Genus <i>Clavulina</i>	154
<i>communis</i>	154
<i>bradyi</i>	155
<i>angularis</i>	155
<i>parisiensis</i>	156
<i>difformis</i>	156
<i>rotundata</i>	157
<i>primaeva</i>	158
Subfamily <i>Bulimininae</i>	159
Genus <i>Bulimina</i>	159
<i>elegans</i> , var. <i>exilis</i>	159
<i>marginata</i>	159
<i>inflata</i>	160
<i>buchiana</i>	160
<i>aculeata</i>	161
<i>pupoides</i>	161
<i>pyrula</i>	162
var. <i>perversa</i>	163
var. <i>spinescens</i>	164
<i>ovata</i>	164
<i>affinis</i>	165
<i>elongata</i>	165
<i>subaffinis</i>	166
Genus <i>Buliminella</i>	167
<i>subteres</i>	167
<i>contraria</i>	167
<i>elegantissima</i>	168
Genus <i>Virgulina</i>	168
<i>subsquamosa</i>	168
<i>pauciloculata</i>	168
<i>schreibersiana</i>	169
<i>cornuta</i>	170
<i>subdepressa</i>	170
Subfamily <i>Cassidulininae</i>	171
Genus <i>Cassidulina</i>	171
<i>laevigata</i>	171
<i>subglobosa</i>	171
<i>crassa</i>	172
Genus <i>Ehrenbergina</i>	172
<i>serrata</i>	172
Family <i>Lagenidae</i>	173
Subfamily <i>Lageninae</i>	173
Genus <i>Lagena</i>	173
<i>globosa</i>	173
<i>laevis</i>	173
<i>laevigata</i>	174
<i>clavata</i>	174
<i>gracillima</i>	175
<i>elongata</i>	175
<i>apiculata</i>	176
<i>hispida</i>	176
var. <i>tubulata</i>	177

Systematic part—Continued.

Family Lagenidae—Continued.

Subfamily Lageninae—Continued.

Genus *Lagena*—Continued.

	Page.
hexagona, var. scalariformis.....	177
striata.....	177
var. haidingeri.....	178
var. strumosa.....	178
semistriata.....	179
sulcata.....	179
var. apiculata.....	180
acuticosta.....	180
gracilis.....	181
staphyllearia.....	181
alveolata, var. substriata.....	182
marginata.....	182
lagenoides.....	183
formosa.....	183
bicarinata.....	184
lamellata.....	184
Subfamily Nodosariinae.....	185
Genus <i>Nodosaria</i>	185
rotundata.....	185
laevigata.....	185
var. striatula.....	186
simplex.....	186
pyrula.....	187
var. semirugosa.....	187
var. longicostata.....	188
annulata.....	189
insecta.....	189
radicula.....	190
insolita.....	191
inflexa.....	191
soluta.....	192
communis.....	192
roemeri.....	193
filiformis.....	194
consobrina, var. emaciata.....	195
mucronata.....	195
japonica.....	196
antennula.....	197
subpolygona.....	198
catenulata.....	198
scalaris.....	199
var. separans.....	200
subscalaris.....	201
var. paucicostata.....	201
millettii.....	202
lepidula.....	203
var. hispidula.....	203
substriatula.....	204
pauciloculata.....	205
var. luzonensis.....	206

Systematic part—Continued.

Family Lagenidae—Continued.

Subfamily Nodosariinae—Continued.

Genus *Nodosaria*—Continued.

	Page.
<i>sublineata</i>	207
<i>perversa</i>	208
<i>subperversa</i>	208
<i>laevicostata</i>	209
<i>raphanus</i>	210
<i>obliqua</i>	210
<i>vertebralis</i>	211
<i>spirostriolata</i>	212
<i>subtenuata</i>	213
<i>hirsuta</i>	213
var. <i>aculeata</i>	214
Genus <i>Lingulina</i>	214
<i>costata</i>	214
<i>grandis</i>	215
Genus <i>Fronicularia</i>	216
<i>inaequalis</i>	216
<i>philippinensis</i>	216
<i>annularis</i>	217
var. <i>longistriata</i>	218
<i>plicata</i>	219
Genus <i>Triplasia</i>	219
<i>tricarinata</i>	219
Genus <i>Cristellaria</i>	220
<i>cultrata</i>	220
var. <i>decorata</i>	221
<i>reniformis</i>	221
<i>articulata</i>	222
<i>rotulata</i>	223
var. <i>umbonata</i>	223
<i>orbicularis</i>	224
var. <i>subcarinata</i>	225
var. <i>subumbonata</i>	226
var. <i>papillata</i>	226
<i>vortex</i>	227
<i>convergens</i>	228
<i>gibba</i>	228
<i>variabilis</i>	229
<i>tricarinella</i>	230
var. <i>spinipes</i>	230
<i>calcar</i>	231
<i>calcarata</i>	232
<i>echinata</i>	233
<i>papillosa</i>	234
<i>aculeata</i>	235
<i>submamelligera</i>	235
<i>expansa</i>	236
var. <i>planulata</i>	237
<i>denticulifera</i>	238
<i>gemma</i>	239

Systematic part—Continued.

Family Lagenidae—Continued.

Subfamily Nodosariinae—Continued.

Genus *Cristellaria*—Continued.

	Page.
<i>costata</i>	239
var. <i>multicosta</i>	240
var. <i>subdecorata</i>	241
var. <i>sublaevis</i>	242
<i>bradyi</i>	243
<i>cassis</i>	244
<i>cassinoides</i>	244
<i>tumido-costata</i>	245
var. <i>labyrinthica</i>	246
<i>dorso-costata</i>	247
<i>compressa</i>	247
<i>paucicostata</i>	248
<i>crepidula</i>	248
<i>schloenbachi</i>	249
<i>tenuis</i>	250
<i>helicina</i>	250
<i>helicinoides</i>	251
<i>italica</i>	252
var. <i>acuti-carinata</i>	253
<i>latifrons</i>	254
Genus <i>Marginulina</i>	254
<i>glabra</i>	254
<i>striatula</i>	255
<i>costata</i>	256
<i>philippinensis</i>	257
Genus <i>Vaginulina</i>	257
<i>legumen</i>	257
var. <i>elegans</i>	258
<i>spinigera</i>	259
<i>margaritifera</i>	259
<i>bradyi</i>	260
<i>linearis</i>	261
<i>acicula</i>	262
<i>protumida</i>	262
<i>patens</i>	263
Subfamily <i>Polymorphininae</i>	263
Genus <i>Polymorphina</i>	263
<i>regina</i>	263
<i>problema</i>	264
var. <i>indica</i>	265
<i>sororia</i>	265
<i>lactea</i>	265
var. <i>oblonga</i>	266
<i>compressa</i>	266
<i>gibba</i>	267
<i>elegantissima</i>	267
<i>oblonga</i>	268
Subfamily <i>Uvigerininae</i>	269
Genus <i>Uvigerina</i>	269
<i>pygmaea</i>	269
<i>tenuistriata</i>	269

Systematic part—Continued.

Family Lagenidae—Continued.

Subfamily Uvigerininae—Continued.

Genus *Uvigerina*—Continued.

Page.

<i>schwageri</i>	270
<i>angulosa</i>	271
<i>striata</i>	271
<i>selseyensis</i>	272
<i>brunnensis</i>	272
<i>aculeata</i>	273
<i>asperula</i>	274
<i>ampullacea</i>	274
<i>interrupta</i>	275
<i>canariensis</i>	275
<i>scrobiculata</i>	276
Genus <i>Siphogenerina</i>	276
<i>columellaris</i>	276
<i>bifrons</i>	277
var. <i>striatula</i>	278
<i>dimorpha</i>	279
<i>striata</i>	280
<i>raphanus</i>	280
var. <i>costulata</i>	281
Subfamily Ramulininae.....	282
Genus <i>Ramulina</i>	282
<i>globulifera</i>	282
Family <i>Chilostomellidae</i>	283
Genus <i>Chilostomella</i>	283
<i>ovoidea</i>	283
<i>grandis</i>	283
Genus <i>Allomorphina</i>	284
<i>trigona</i>	284
Family <i>Globigerinidae</i>	285
Genus <i>Globigerina</i>	285
<i>bulloides</i>	285
var. <i>triloba</i>	286
<i>cretacea</i>	287
<i>digitata</i>	288
<i>inflata</i>	289
<i>rubra</i>	289
<i>conglobata</i>	290
<i>sacculifera</i>	291
<i>aequilateralis</i>	293
var. <i>involuta</i>	294
<i>conglomerata</i>	295
Genus <i>Orbulina</i>	295
<i>universa</i>	295
Genus <i>Candeiana</i>	296
<i>nitida</i>	296
var. <i>triloba</i>	296
Genus <i>Sphaeroidina</i>	297
<i>bulloides</i>	297
<i>dehiscens</i>	297

Systematic part—Continued.

Family Globigerinidae—Continued.

	Page.
Genus <i>Pullenia</i>	299
<i>sphaeroides</i>	299
<i>quineloba</i>	299
<i>obliquiloculata</i>	299
Family Rotaliidae	301
Subfamily Spirillininae	301
Genus <i>Spirillina</i>	301
<i>limbata</i>	301
var. <i>denticulata</i>	302
var. <i>tuberculo-limbata</i>	302
Subfamily Rotalinae	303
Genus <i>Patellina</i>	303
<i>corrugata</i>	303
Genus <i>Discorbis</i>	303
<i>subobtusa</i>	304
<i>vilardeboana</i>	304
<i>orbicularis</i>	305
<i>bertheloti</i>	305
<i>allomorphinoides</i>	306
<i>rugosa</i>	306
<i>species</i>	307
Genus <i>Cymbalopora</i>	308
<i>poeyi</i>	308
var. <i>bradyi</i>	308
Genus <i>Tretomphalus</i>	309
<i>bulloides</i>	309
Genus <i>Planorbulina</i>	310
<i>larvata</i>	310
<i>acervalis</i>	311
Genus <i>Truncatulina</i>	312
<i>refulgens</i>	312
<i>lobatula</i>	313
<i>tenuimargo</i>	313
<i>variabilis</i>	314
<i>wuellerstorfi</i>	314
<i>haidingerii</i>	315
<i>akneriana</i>	316
<i>ungeriana</i>	317
var. <i>ornata</i>	317
<i>tenera</i>	318
<i>praecincta</i>	318
<i>margaritifera</i>	319
<i>culter</i>	320
<i>humulis</i>	321
<i>rostrata</i>	321
Genus <i>Siphonina</i>	322
<i>reticulata</i>	322
Genus <i>Anomalina</i>	322
<i>ariminensis</i>	322
<i>grosserugosa</i>	323
<i>ammonoides</i>	323

Systematic part—Continued.

Family Rotaliidae—Continued.

Subfamily Rotalinae—Continued.

Genus *Anomalina*—Continued.

	Page.
<i>polymorpha</i>	324
var. <i>cervicornis</i>	325
var. <i>siphonifera</i>	325
<i>coronata</i>	326
Genus <i>Pulvinulina</i>	326
<i>repanda</i>	326
<i>concentrica</i>	327
<i>punctulata</i>	328
<i>concamerata</i>	328
<i>auricula</i>	329
<i>oblonga</i>	330
<i>scabra</i>	330
<i>philippinensis</i>	331
<i>indica</i>	332
<i>berthelotiana</i> , var. <i>subornata</i>	333
<i>menardii</i>	333
var. <i>fimbriata</i>	335
<i>lateralis</i>	336
<i>tumida</i>	337
<i>canariensis</i>	338
<i>crassa</i>	338
<i>truncatulinoides</i>	339
<i>umbonata</i>	339
<i>exigua</i>	340
<i>pauperata</i>	340
<i>schreibersii</i>	341
<i>procera</i>	342
<i>elegans</i>	342
<i>partschiana</i>	344
Genus <i>Rotalia</i>	345
<i>beccarii</i>	345
<i>broeckhiana</i>	346
<i>orbicularis</i>	346
<i>schroeteriana</i>	347
<i>papillosa</i>	347
var. <i>compressiuscula</i>	348
<i>soldanii</i>	349
<i>calcar</i>	350
Genus <i>Calcarina</i>	350
<i>spengleri</i>	351
<i>defrancii</i>	353
<i>baculatus</i>	354
<i>hispida</i>	356
Genus <i>Siderolites</i>	357
<i>tetraedra</i>	358
Genus <i>Baculogypsina</i>	359
<i>sphaerulatus</i>	359
Genus <i>Gypsina</i>	360
<i>globulus</i>	360

Systematic part—Continued.

Family Rotaliidae—Continued.

Subfamily Rotalinae—Continued.

	Page.
Genus <i>Carpenteria</i>	360
<i>utricularis</i>	360
<i>proteiformis</i>	361
<i>monticularis</i>	362
<i>rhaphidodendron</i>	363
Genus <i>Polytrema</i>	363
<i>miniaceum</i>	363
Genus <i>Homotrema</i>	364
<i>rubrum</i>	364
Genus <i>Sporadotrema</i>	365
<i>cylindricum</i>	365
Family <i>Nummulitidae</i>	365
Genus <i>Nonionina</i>	365
<i>umbilicatula</i>	365
<i>pompilioides</i>	366
<i>boueana</i>	366
Genus <i>Polystomella</i>	367
<i>subnodosa</i>	367
<i>crispata</i>	368
<i>craticulata</i>	368
Genus <i>Amphistegina</i>	370
<i>lessonii</i>	370
<i>var. madagascariensis</i>	372
<i>var. radiata</i>	372
Genus <i>Operculina</i>	373
<i>gaimairdi</i>	375
<i>bartschi</i>	376
<i>var. plana</i>	377
<i>var. ornata</i>	378
<i>philippinensis</i>	378
<i>discoidalis</i>	379
<i>var. involuta</i>	380
<i>granulosa</i>	381
<i>elegans</i>	381
<i>ammonoides</i>	382
<i>venosa</i>	383
Genus <i>Heterostegina</i>	384
<i>depressa</i> , <i>var. tubercalata</i>	384
<i>suborbicularis</i>	385
Genus <i>Cycloclypeus</i>	386
<i>guembelianus</i>	386
Family <i>Miliolidae</i>	387
Subfamily <i>Cornuspirininae</i>	387
Genus <i>Cornuspira</i>	387
<i>foliacea</i>	387
<i>var. expansa</i>	388
<i>involvens</i>	389
<i>var. substriatula</i>	390
<i>lacunosa</i>	391
<i>carinata</i>	392
<i>crassisepta</i>	392

Systematic part—Continued.

Family Miliolidae—Continued.

Subfamily Cornuspirininae—Continued.

	Page.
Genus <i>Ophthalmidium</i>	393
<i>inconstans</i>	393
Genus <i>Spiroloculina</i>	394
<i>depressa</i>	394
<i>canaliculata</i>	395
<i>grateloupi</i>	396
var. <i>incisa</i>	397
<i>acutimargo</i>	398
<i>tenuimargo</i>	399
<i>tenuissima</i>	400
<i>millettii</i>	400
<i>planissima</i>	401
<i>tenuiseptata</i>	401
<i>circularis</i>	403
<i>orbis</i>	403
<i>robusta</i>	404
<i>validus</i>	404
<i>costifera</i>	405
var. <i>plena</i>	405
<i>scrobiculata</i>	406
<i>elegans</i>	406
<i>antillarum</i>	407
var. <i>angulata</i>	408
<i>convexiuscula</i>	409
<i>advena</i>	410
<i>affixa</i>	410
<i>arenaria</i>	411
Genus <i>Planispirina</i>	412
<i>contraria</i>	412
<i>striata</i>	413
<i>sphaera</i>	413
Genus <i>Vertebralina</i>	414
<i>striata</i>	414
var. <i>reticulosa</i>	414
Genus <i>Nodobacularia</i>	415
<i>tibia</i>	415
Subfamily <i>Quinqueloculininae</i>	415
Genus <i>Quinqueloculina</i>	415
<i>seminulum</i>	416
<i>vulgaris</i>	417
<i>lamarckiana</i>	418
<i>venusta</i>	420
<i>procera</i>	421
<i>apicula</i>	422
<i>limbata</i>	422
<i>crassa</i> , var. <i>subcuneata</i>	423
<i>recta</i>	423
<i>disparilis</i>	424
<i>boueana</i>	425
<i>bicornis</i> , var. <i>elegans</i>	425
<i>curta</i>	426

Systematic part—Continued.

Family Miliolidae—Continued.

Subfamily Quinqueloculinae—Continued.

Genus Quinqueloculina—Continued.

	Page.
bicarinata.....	428
ferussacii.....	429
rugosa.....	429
polygona.....	431
bradyana.....	432
contorta.....	432
sculpturata.....	433
reticulata.....	434
albatrossi.....	435
kerimbatica.....	437
var. reticulo-striata.....	437
var. philippinensis.....	438
parkeri.....	440
agglutinans.....	441
sclerotica.....	441
fusca.....	442
crassatina.....	443
alveoliniformis.....	443
Genus Massilina.....	444
secans, var. reticulata.....	444
var. tenuistriata.....	445
crenata.....	445
durrandii.....	446
arenaria.....	446
asperula.....	447
Genus Articulina.....	448
sagra.....	448
Genus Sigmoilina.....	448
sigmoidea.....	448
schlumbergeri.....	449
Genus Hauerina.....	450
bradyi.....	450
fragillissima.....	451
ornatissima.....	452
Genus Triloculina.....	452
trigonula.....	452
var. multistriata.....	453
advena.....	453
tricarinata.....	454
var. convexa.....	456
var. plicata.....	456
bertheliniana.....	457
terquemiana.....	458
insignis.....	458
oblonga.....	459
rotunda.....	460
valvularis.....	460
suborbicularis.....	461
cuneata.....	461
var. incisa.....	462

Systematic part—Continued.

Family Miliolidae—Continued.

Subfamily Quinqueloculininae—Continued.

Genus Triloculina—Continued.

	Page.
<i>circularis</i>	462
<i>linnaeana</i>	463
<i>rupertiana</i>	464
<i>tortuosa</i>	464
Genus <i>Flintina</i>	465
<i>bartschi</i>	465
<i>triquetra</i>	466
<i>bradyana</i>	467
Genus <i>Biloculina</i>	469
<i>depressa</i>	469
<i>murrhyna</i>	470
<i>serrata</i>	471
<i>sarsii</i>	471
<i>vespertilio</i>	472
<i>elongata</i>	473
<i>globulus</i>	474
<i>anomala</i>	474
<i>lucernula</i>	475
<i>pisum</i>	475
<i>denticulata</i>	476
var. <i>striolata</i>	477
<i>comata</i>	477
<i>millettii</i>	479
<i>irregularis</i>	479
Genus <i>Nevillina</i>	480
<i>coronata</i>	480
Genus <i>Peneroplis</i>	481
<i>pertusus</i>	481
<i>planatus</i>	481
<i>carinatus</i>	482
<i>arietinus</i>	482
<i>cylindraceus</i>	483
Genus <i>Orbiculina</i>	483
<i>adunca</i>	483
Genus <i>Orbitolites</i>	484
<i>marginalis</i>	484
<i>duplex</i>	485
<i>complanata</i>	486
var. <i>plicata</i>	487
Genus <i>Alveolina</i>	487
<i>boscii</i>	487
Explanation of plates.....	489
Index.....	591

LIST OF ILLUSTRATIONS.

	Page.
FIG. 1. <i>Bulimina pyrula</i> d'Orbigny. × 50. From <i>Albatross</i> station D5586. a-c, three views of same specimen.....	162
2. <i>Bulimina pyrula</i> d'Orbigny, var. <i>perversa</i> Cushman. From <i>Albatross</i> station D5654. a-c, three views of same specimen.....	163
3. <i>Bulimina pyrula</i> d'Orbigny, var. <i>spinescens</i> H. B. Brady. × 50. From <i>Albatross</i> station D5586.....	163
4. <i>Bulimina ovata</i> d'Orbigny. × 50. From <i>Albatross</i> station D5469...	164
5, 6. <i>Bulimina affinis</i> d'Orbigny. × 50. From <i>Albatross</i> station D5114. 5a, b, two views of same specimen. 6, another specimen of irreg- ular shape	165
7. <i>Bulimina subaffinis</i> , new species. Views from different sides. × 100. From station D5201.....	166
8. <i>Nodosaria pauciloculata</i> , var. <i>luzonensis</i> , new variety. Front view. × 100. From station D5220.....	206
9. <i>Cristellaria cultrata</i> , var. <i>decorata</i> Cushman. Side view. × 50. From station D5113.....	221
10. <i>Uvigerina scrobiculata</i> , new species. × 50. a, b, two views from different angles. From station D5333.....	276
11. <i>Globigerina aequilateralis</i> , var. <i>involuta</i> Cushman. × 50. a, side view; b, from an oblique angle; c, apertural view. From station D5255.....	293
12. <i>Truncatulina ungeriana</i> , var. <i>ornata</i> , new variety. × 100. a, dorsal view; b, ventral view. From station D5623.....	317
13. <i>Operculina bartschi</i> , new species. Side view. × 15. From station D5134.....	376
14. <i>Operculina bartschi</i> , var. <i>plana</i> , new variety. × 30. Side view. From station D5105.....	377
15. <i>Operculina philippinensis</i> , new species. Side view. × 15. From station D5134.....	379
16. <i>Operculina discoidalis</i> , var. <i>involuta</i> , new variety. Side view. × 20. From station D5172.....	380
17, 18. <i>Spiroloculina grateloupi</i> d'Orbigny. × 50. End views showing aperture and teeth. From <i>Albatross</i> station D5136.....	396
19, 20. <i>Quinqueloculina seminulum</i> (Linnaeus). × 20. From <i>Albatross</i> station D5152. 19, specimen with the tip of the tooth becoming bifid. 20, specimen with the tooth forked and much broadened toward the tip.....	416
21. <i>Quinqueloculina vulgaris</i> d'Orbigny. × 50. From <i>Albatross</i> station D5126. Opposite sides of same specimen.....	417
22, 23. <i>Quinqueloculina lamareckiana</i> d'Orbigny. × 50. From <i>Albatross</i> station D5220. 22, apertural view of young specimen with simple elongate tooth. 23, adult specimen which has developed a bifid tooth and a more rounded aperture.....	419
24. <i>Quinqueloculina rugosa</i> d'Orbigny. Aperture and undivided simple tooth of early stage. × 150. From <i>Albatross</i> station D5164...	430
25-33. <i>Quinqueloculina albatrossi</i> , new species. × 50. Various stages in the development of the apertural characters. 25, a simple tooth in a generally rounded aperture. 26-29, a bifid tooth, the aperture	

	Page.
gradually assuming an irregular form by the development of processes and reentrants. 30, one arm of the bifid tooth broadening, the other already joined to the side of the aperture. 31, 32, stages in the broadening of the tooth after the central opening is entirely surrounded. 33, a later stage where the sides have fused into the processes leaving a central aperture and six supplementary ones about it.....	435
FIG. 34. <i>Quinqueloculina kerimbatica</i> Heron-Allen and Earland, var. <i>philippinensis</i> , new variety, apertural view. $\times 50$. From <i>Albatross</i> station D5164.....	439
35, 36. <i>Triloculina tricarinata</i> d'Orbigny. 35, apertural view of broad, peculiarly shaped specimen. $\times 65$. From <i>Albatross</i> station D5160. 36, apertural view of a much larger specimen where the outer face has a double angle. $\times 35$. From <i>Albatross</i> station D5236..	454
37. <i>Triloculina tricarinata</i> d'Orbigny. Apertural view. $\times 50$. From <i>Albatross</i> station D5236.....	456
38-40. <i>Flintina bradyana</i> , new species. 38, specimen with simple bifid tooth. $\times 100$. From <i>Albatross</i> station D5141. 39, specimen in later stage with broadened tooth. $\times 50$. From <i>Albatross</i> station D5164. 40, larger specimen with reduced costae, the last-formed chamber smooth, the tooth sending out processes toward the wall at the sides of the aperture. $\times 50$	467
41-44. <i>Flintina bradyana</i> , new species. $\times 50$. From <i>Albatross</i> station D5133. 41a, side view; b, apertural view of young specimen with simple tooth. 42, aperture of older specimen with bifid tooth. 43, aperture of a still older specimen with the processes of the tooth beginning to extend in and the opening to have angles. 44, aperture of an adult where the arms have met to form a central ring, additional arms extended from either side of the aperture at the angles, and secondary arms being given off on the sides of the ring.....	468
45-47. <i>Biloculina depressa</i> d'Orbigny. 45, young specimen with elliptical aperture and elongate tooth. $\times 50$. From <i>Albatross</i> station D5236. 46, later stage showing the greater elongation of the aperture and tooth. $\times 100$. From <i>Albatross</i> station D5201. 47, adult specimen with very narrow curved aperture, largely occupied by the somewhat expanded tooth. $\times 40$. From <i>Albatross</i> station D5236.....	469
48. <i>Biloculina sarsii</i> Schlumberger. $\times 50$. From <i>Albatross</i> station D5220. a, front view; b, side view; c, apertural view of same specimen.....	471
49, 50. <i>Biloculina sarsii</i> Schlumberger. 49, aperture of early stage with a broad, rounded tooth and an elliptical opening. $\times 80$. From <i>Albatross</i> station D5178. 50, aperture of an adult specimen with very broad aperture, the tooth very narrow and elongate with the angles produced at each end. $\times 100$. From <i>Albatross</i> station D5133.....	472
51. <i>Biloculina comata</i> H. B. Brady. $\times 100$. Aperture of specimen from <i>Albatross</i> station D5236.....	477
52. <i>Biloculina comata</i> H. B. Brady. Outline of the chambers of a partially broken specimen, showing the early quinqueloculine stage with nearly smooth chambers. $\times 100$. From <i>Albatross</i> station D5236.....	478

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

By JOSEPH A. CUSHMAN.

Of the Boston Society of Natural History.

INTRODUCTION.

The collections represented by material included in the following report were made by the United States Bureau of Fisheries steamer *Albatross*. During several years the Philippine Expedition made collections covering the waters of the Archipelago and the surrounding waters of the China Sea and Pacific Ocean. In addition, collections were made in the region to the southward about Borneo, Celebes, and the larger East Indies. A part of this southern area is similar to that included in Millett's material, but his was almost entirely from comparatively shallow water, while our material has been largely from much deeper water. The *Challenger* did some dredging in the Philippine Archipelago, one station in 95 fathoms (174 meters) off Cebu, being thoroughly worked by Brady and referred to by others. With these exceptions there are few records for the recent material of the region. As a result, there are many interesting records and undescribed species.

A number of years have now been in part given to this report on the Philippine collections. As the collections include nearly 600 dredging stations, besides shallow-water material, the mechanical work, involving as it has the handling and sorting of many thousands of specimens, has been very time-consuming and laborious. The mounted material includes several thousand slides, including many thousands of actual specimens. These are in the collection of the United States National Museum.

The collections probably represent more completely the area covered than any similar collections for a region of equal size. The tropical character of much of the shallow-water material makes it very interesting from various points of view.

In the study of the collection it has been the purpose to give the distribution of such species in the region, and the tables give the data for these stations. In this way much detailed information is available for the causes affecting distribution of various genera and species. Something is given of the variation where this occurs, and in some cases changes in nomenclature are discussed where neces-

sary. The classification adopted is that already used in an earlier series of papers¹ and in the early parts of another paper now being issued.² In these papers will be found more detailed synonymy and the discussion of the classification and changes of name which have occurred in the earlier work. References to these papers and to the *Challenger* Report are given for nearly all the older species. The descriptions of many of the new species have already been published and references to the place of publication given, though for ease of reference the descriptions have been given in full for the species described as new from these collections.

Figures have been given for most of the species and all have been taken from material of this collection. Most of these are drawings, but a few are photographs. Many more figures could well have been used to illustrate the various species and varieties. It would have been of interest to show the developmental stages of numerous species, but this would have involved the drawing of many more figures and a considerable increase in the number of plates.

The work of completing the present report would have been impossible but for the very kindly help which has been unstintingly given. The Bureau of Fisheries under Commissioners George M. Bowers, Dr. Hugh M. Smith, and Deputy Commissioner H. F. Moore, has given unceasing aid in the furnishing of assistance in preparing the material and the work of getting the manuscript and figures in shape. The United States National Museum has taken care of the mounting of the slides of material and help in other ways. That the material is so rich is very largely due to careful work of Dr. Paul Bartsch during the time he had charge of the collecting work of the *Albatross* in the Philippines. Miss Jennie F. Cushman is responsible for the large series of species and specimens which are due to her painstaking work in patiently picking out the foraminifera from the mass of the bottom samples after they were carefully washed. A considerable portion of the figures were drawn by Mr. Edward Gilman, whose excellent work speaks for itself. Messrs. Charles B. Ames, S. N. F. Sanford, and Miss Margaret Birch did much of the card cataloging of records. Miss Jane M. Furber gave much time to the preparing of card records and the earlier part of the manuscript. Especially to Miss Shirley P. Beck are my thanks due for her very efficient and painstaking work in bringing together in systematic shape the various parts of the work, preparing the tables of distribution, and generally getting the manuscript into its final form. Without her careful work it would have been hardly possible to have gotten the whole into its final shape.

¹ Cushman, Monograph of the Foraminifera of the Pacific Ocean, Bull. 71, U. S. Nat. Mus., pts. 1-6, 1910-1917.

² Cushman, Foraminifera of the Atlantic Ocean, Bull. 104, U. S. Nat. Mus., pt. 1, 1918; pt. 2, 1920.

The lack of certain species is perhaps to be based on the lack of fineness of some of the material available, and if finer material had been available in some cases undoubtedly numerous records for smaller species would have been obtained. This is especially noticeable in the case of small things, such as *Lagena*, *Bolivina*, etc., which from the work of others should be abundant in the region.

Except for one or two stations in the Sulu Sea and in the region to the southward most of the material came from depths of less than 1,000 fathoms (1,829 meters), and much of it from shallow water. Some of the shallow water attached species of genera such as *Polytrema*, *Homotrema*, *Sporadotrema*, *Carpenteria*, etc., are undoubtedly abundant, but the larger mollusca and corals to which they would have been attached were not usually studied. Certain of these are probably widely distributed in shallow, warm waters, and where especially looked for were often found to be abundant.

The general relations of the material to that from other regions is interesting. The shallow water fauna, that inside of 30 fathoms (55 meters), is characteristically tropical, most of the genera being those of similar areas in the general Indo-Pacific region, but many of the species being distinct. In such locations the foraminifera make up a large proportion of the bottom samples. Such genera as *Calcarina*, *Baculogypsina*, *Siderolites*, *Operculina*, *Polystomella*, *Heterostegina*, *Cycloclypeus*, etc., as well as the Miliolidae, are abundant under such conditions.

There is an individuality to the fauna, however, which at once separates it from that of the Barrier Reef region of Australia or of the shallow water of the Indian Ocean, as will be noted under various genera.

In the somewhat deeper water, that from 100 to 300 fathoms (183 to 549 meters), there is developed a fauna which is very interesting and perhaps dominated largely by an exceptional development of the Lagenidae. *Nodosaria* and *Cristellaria* are very abundant and represented by numerous and well-developed species and varieties, many of which are new. A good development of *Fronidicularia* is an interesting find in the region. This fauna is interesting in comparison with the material obtained by the *Challenger* in 95 fathoms (173 meters), off Cebu. Some of the species which were described by Brady from this station have been almost unknown since until these *Albatross* dredgings were studied. These species have now been found to be abundant and characteristic in many parts of the Archipelago.

In various parts of the region but especially in the colder, deeper portions, there is an exceptional development of arenaceous foraminifera, especially *Astrorhizidae* and *Lituolidae*. These, even though under the Equator, have the characteristic genera and many of the

species of cold waters in high latitudes. This goes to further prove the wide distribution of these arenaceous forms in cold waters, temperature instead of depth having the greater control in their distribution.

However, many of the species of these arenaceous foraminifera were undescribed and their nearest allied species so far as present distribution is known are far away. These include species of *Hali-physema*, *Dendrophrya*, *Bathysiphon*, etc. They show what other observers have found, especially Heron-Allen and Earland in their work about the British Isles, that arenaceous species often occur in great abundance in very restricted localities. Certain species of the above genera at some of the stations made up a very large percentage of the foraminifera of the bottom material but were not found elsewhere.

One of the most interesting stations in the whole of the *Albatross* dredgings is D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao. The bottom material is of a peculiar brownish color and contains a very unusual assemblage of species, a large number of which have been described here as new. A large number of rare species have also occurred at this station which have not been found elsewhere in the region. A list of the species from this station is given, as well as from other stations of exceptional interest to show the general character of the foraminiferal fauna in some of the various parts of the area.

Globigerina-ooze is developed in some parts of the Sulu Sea and in the southern area. This is rather typical in most of its species but lacks certain ones. This probably may be due to currents and will be mentioned where the *Globigerinidae* are discussed.

One of the most interesting finds in the region is *Nevillina coronata*, a very rare species and of especial interest from its relationship to other *Miliolidae*. The great abundance of *Cycloclypeus carpenteri* in certain parts of the area was an unexpected find. It is the largest of the living calcareous foraminifera.

In general the foraminiferal fauna of the Philippine area is that of the Indo-Pacific, including the Hawaiian Islands on the east, the South Sea Archipelagoes, and Australia to the southward, and going westward to the East Coast of Africa, Madagascar, the Red Sea, and in some cases, the Mediterranean. This is only in general, however, and there are many characteristic species of the area which so far as is now known are limited to this region.

The systematic portion follows, giving the species with notes on their distribution in the area.

A LIST OF SPECIES FROM ALBATROSS STATION D5179, VICINITY OF
ROMBLON.

LATITUDE 12° 38' 15" NORTH; LONGITUDE 122° 06' 15" EAST. DEPTH, 37 FATHOMS
(68 METERS). BOTTOM TEMPERATURE, 75.7° F. (24.2° C.). CHARACTER OF
BOTTOM, HRD. S.

FAUNA OF SHALLOW WARM WATER WITHIN THE ARCHIPELAGO.

LITUOLIDAE:

Reophax scorpiurus.

TEXTULARIIDAE:

Textularia siphonifera.

rugosa.

conica.

gramen.

Verneuilina spinulosa.

LAGENIDAE:

Cristellaria vortex.

dorso-costata.

schloenbachi.

Polymorphina elegantissima.

GLOBIGERINIDAE:

Globigerina sacculifera.

aequilateralis.

var. involuta.

Pullenia obliquiloculata.

ROTALIIDAE:

Spirillina limbata.

var. denticulata.

var. tuberculo-limbata.

Truncatulina refulgens.

Sporadotrema cylindricum.

Homotrema rubrum.

Polytrema miniaceum.

Calcarina baculatus.

hispidus.

Carpenteria proteiformis.

Siderolites tetraedra.

Pulvinulina lateralis.

elegans.

menardii.

scabra.

concentrica.

NUMMULITIDAE:

Amphistegina lessonii.

var. madagascariensis.

var. radiata.

Operculina bartschi.

philippinensis.

Heterostegina suborbicularis.

Polystomella craticulata.

crispa.

Cyclocypeus guembelianus.

MILIOLIDAE:

- Spiroloculina convexiuscula*.
grateloupi.
canaliculata.
orbis.
antillarum.
- Quinqueloculina agglutinans*.
albatrossi.
alveoliniformis.
crassa, var. *subcuneata*.
parkeri.
rugosa.
- Massilina crenata*.
Triloculina circularis.
linneiana.
suborbicularis.
tortuosa.
tricarinata.
- Biloculina denticulata*.
 var. *striolata*.
- Alveolina boscii*.

A LIST OF SPECIES FROM ALBATROSS STATION D5201, SOGOD BAY,
SOUTHERN LEYTE ISLAND.

LATITUDE 10° 10' 00" NORTH; LONGITUDE 125° 04' 15" EAST. DEPTH, 554 FATHOMS
(1,012 METERS). BOTTOM TEMPERATURE, 52.8° F. (11.5° C.). CHARACTER OF
BOTTOM, GY. S, M.

FAUNA OF DEEPER, COOLER WATER WITHIN THE ARCHIPELAGO.

ASTRORHIZIDAE:

- Rhizammima algaeformis*.
Iridia diaphana.
Proteonina testacea.
Technitella legumen.

LITUOLIDAE:

- Reophax pilulifer*.
agglutinatus.
Ammobaculites calcareum.
cylindricus.
pseudospirale.

TEXTULARIIDAE:

- Textularia abbreviata*.
siphonifera.
quadrilatera.
porrecta.
- Gaudryina bradyi*.
Clavulina parisiensis.
Bolivina hantkeniana.
limbata.
punctata.
- Bulimina inflata*.
aculeata.
pyrula, var. *perversa*.
ovata.

TEXTULARIIDAE—Continued.

Bulimina pupoides.
pyrula.
subaffinis.

Virgulina cornuta.

LAGENIDAE:

Lagena globosa.

laevigata.
gracillima.

Cristellaria italica.

cassinoides.
echinata.
articulata.
submilligera.
expansa.
calcar.
orbicularis.
costata.
crepidula.
schloenbachi.
variabilis.
tricarinata.
calcarata.

Nodosaria pyrula.

var. *semirugosa*.
 var. *longicostata*.

filiformis.
soluta.
inflexa.
consobrina, var. *emaciata*.
mucronata.
scalaris.
 var. *separans*.
pauciloculata.
raphanus.
obliqua.
vertebralis.
hirsuta.

var. *aculeata*.

Fronicularia philippinensis.

annularis, var. *longistriata*.

Marginulina glabra.

striatula.

Siphogenerina bifrons.

var. *striatula*.

striata.

raphanus.

var. *costulata*.

Uvigerina pygmaea.

tenuistriata.

schwageri.

aculeata.

asperula.

Vaginulina patens.

GLOBIGERINIDAE:

- Globigerina inflata*.
 conglobata.
 sacculifera.
 aequilateralis, var. *involuta*.

- Orbulina universa*.
Sphaeroidina dehiscens.
Pullenia obliquiloculata.

CHILOSTOMELLIDAE:

- Chilostomella ovoidea*.
Allomorphina trigona.

ROTALIIDAE:

- Discorbis*, species?
Planorbulina larvata.
Anomalina polymorphina.
Truncatulina tenuimargo.
Rotalia soldanii.
 beccarii.

- Calcarina hispida*.
Pulvinulina crassa.
 menardii.
 auricula.

- Cymbalopora poeyi*.

NUMMULITIDAE:

- Amphistegina lessonii*.
Operculina bartschi.
 var. *plana*.
 philippinensis.
 discoidalis, var. *involuta*.
Heterostegina depressa, var. *tubercalata*.
Nonionina boueana.
Polystomella craticulata.
 crispa.

MILIOLIDAE:

- Cornuspira carinata*.
Spiroloculina grateloupi.
 canaliculata.
 tenuissima.
Nodobacularia tibia.
Quinqueloculina lamarckiana.
 vulgaris.
Sigmoilina sigmoidea.
 schlumbergeri.
Triloculina circularis.
 tricarinata.
Biloculina anomala.
 comata.
 depressa.
Orbitolites duplex.
 complanata.

A LIST OF SPECIES FROM ALBATROSS STATION D5236, PACIFIC OCEAN,
EAST COAST MINDANAO.

LATITUDE 8° 50' 45" NORTH; LONGITUDE 126° 26' 52" EAST. DEPTH, 494 FATHOMS
(903 METERS). BOTTOM TEMPERATURE 41.2° F. (5.1° C.). CHARACTER OF BOTTOM,
FNE. GY. S.

FAUNA OF COOL WATER IN MODERATE DEPTHS OUTSIDE THE ISLANDS.

ASTORRHIZIDAE:

- Rhabdammina abyssorum*.
 discreta.
 linearis.
Bathysiphon filiformis.
 rufus.
 rufescens.
Psammosphaera fusca.
 parva.
Protonina testacea.
Technitella legumen.
Thurammina papillata, var. *castanea*.
Saccorhiza ramosa.
Tolypammina vagans.
Ammolagena clavata.
Jaculella acuta.
 obtusa.
Ammodiscus incertus.

LITUOLIDAE:

- Reophax nodulosus*.
 pseudobacillaris.
 spiculotestus.
Hormosina elongata.
 ovaliformis.
Haplostiche dubia.
Haplophragmoides canariensis.
Cyclammina cancellata.
 pusilla.
 bradyi.
Lituotuba lituiformis.
Ammobaculites calcareum.
Placopsilina cenomana.
Trochammina squamata.
 globigeriniformis.
Ammosphaeroidina grandis.
Sphaerammina ovalis.
Trochamminoides proteus.

TEXTULARIIDAE:

- Textularia stricta*.
 flintii.
 quadrilatera.
 rugosa.
 excarata.
 foliacea.
 crassisepta.

TEXTULARIIDAE—Continued.

Gaudryina bradyi.
baccata.
chilostoma.

Clavulina communis.

Bigenerina arenacea.

capreolus.

pennatula.

Bolivina sculpturata.

Pleurostomella alternans.

subnodosa.

Verneuilina polystropha.

propinqua.

bradyi.

affixa.

Bulimina inflata.

pyrula, var. *perversa*.

Buliminella elegantissima.

contraria.

Cassidulina subglobosa.

LAGENIDAE:

Lagena staphyllearia.

alveolata, var. *substriata*.

Cristellaria italica.

cultrata.

Nodosaria laevigata.

var. *striatula*.

pyrula.

var. *semirugosa*.

filiformis.

soluta.

annulata.

insecta.

insolita.

inflexa.

communis.

consobrina, var. *emaciata*.

japonica.

antennula.

scalaris.

lepidula.

var. *hispidula*.

subperversa.

spirostriolata.

subtertenuata.

Siphogenerina columellaris.

bifrons, var. *striatula*.

striata.

Urigerina pygmaea.

tenuistriata.

aculeata.

asperula.

ampullacea.

Vaginulina legumen, var. *elegans*.

protumida.

GLOBIGERINIDAE:

- Globigerina inflata*.
conglobata.
sacculifera.
aequilateralis, var. *incoluta*.

- Orbulina universa*.
Sphaeroidina dehiscens.
Pullenia quinqueloba.

CHILOSTOMELLIDAE:

- Chilostomella grandis*.

ROTALIIDAE:

- Discorbis rugosa*.
subobtusa.
saucii.
Planorbulina larvata.
Anomalina grosserugosa.
ammonoides.
polymorpha, var. *cervicornis*.
 var. *siphonifera*.

- coronata*.
Truncatulina refulgens.
humulis.
ungeriana.
lobatula.
wuellerstorfi.
praecincta.

Rotalia soldanii.

- broeckhiana*.
calcar.

- Pulvinulina procera*.
schreibersii.
truncatulinoïdes.
pauperata.
tumida.
menardii.
punctulata.

Tetromphalus bulloides.

NUMMULITIDAE:

- Operculina bartschi*.
 var. *plana*.
 var. *ornata*.
Nonionina pompilioides.
umbilicatulata.

MILIOLIDAE:

- Cornuspira foliacea*.
involvens.
 var. *substriaatula*.
crassisepta.
Ophthalmidium inconstans.
Spiroloculina costifera.
 var. *plena*.
tenuiseptata.
Planispirina contraria.
Nodobacularia tibia.

MILIOLIDAE—Continued.

- Quinqueloculina apicula.*
vulgaris.
procera.
- Massilina arenaria.*
- Sigmoilina sigmoidea.*
schlumbergeri.
- Triloculina advena.*
insignis.
oblonga.
rotundata.
tricarinata.
 var. *plicata.*
trigonula, var. *multistriata.*
- Biloculina irregularis.*
lucernula.
comata.
depressa.
vespertilio.
- Flintina bartschi.*

A LIST OF SPECIES FROM ALBATROSS STATION D5650, GULF OF BONI.

LATITUDE 4° 53' 45" SOUTH; LONGITUDE 121° 29' 00" EAST. DEPTH, 540 FATHOMS (988 METERS). BOTTOM TEMPERATURE, 40.1° F. (3.5° C.). CHARACTER OF BOTTOM, GN. M.

FAUNA OF COLD WATER AT MODERATE DEPTHS IN THE SOUTHERN AREA.

ASTRORHIZIDAE:

- Astrorhiza granulosa.*
- Rhabdammina abyssorum*, var. *radiata.*
irregularis.
- Bathysiphon rufus.*
- Tholosina bulla.*
- Hyperammina elongata.*
- Tolypammina vagans.*
- Dendrophrya ramosa.*
- Ammolagena clavata.*
- Ammodiscus incertus.*

LITUOLIDAE:

- Reophae nodulosus.*
 var. *hormosinoides.*
bilocularis.
- Haplostiche dubia.*
- Cyclammina cancellata.*
compressa.
pauciloculata.
- Ammobaculites calcareum.*
cylindricus.
globigeriniformis.

TEXTULARIIDAE:

- Gaudryina flintii.*
- Clavulina communis.*
- Valvulina conica.*
- Cussidulina subglobosa.*

LAGENIDAE:

- Cristellaria italica*.
cassis.
latifrons.
rotulata.
submamilligera.
vortex.
helicina.
gibba.
Nodosaria soluta.
consobrina, var. *emaciata*.
japonica.
scalaris.
 var. *separans*.
sublineata.
raphanus.
Uvigerina aculeata.
asperula.
interrupta.

GLOBIGERINIDAE:

- Globigerina cretacea*.
digitata.
conglobata.
Sphaeroidina dehiscens.
Pullenia obliquiloculata.

ROTALIIDAE:

- Planorbulina larvata*.
Anomalina polymorphina.
Truncatulina praecincta.
Pulvinulina elegans.
canariensis.
pauperata.
tumida.
menardii.

MILIOLIDAE:

- Cornuspira carinata*.
Spiroloculina tenuiseptata.
robusta.
Massilina arenaria.
tricarinata.
valvularis.
Biloculina depressa.
vespertilio.

A LIST OF SPECIES FROM ALBATROSS STATION H4897, SULU SEA, OFF WESTERN MINDANAO.

LATITUDE 7° 46' 00" NORTH; LONGITUDE 122° 00' 00" EAST. DEPTH, 1,570 FATHOMS (2,871 meters). CHARACTER OF BOTTOM, GY. M., GLOB.

FAUNA OF DEEP WATER IN THE SULU SEA.

LITUOLIDAE:

- Reophax dentaliniformis*.
guttifer.
Haplophragmoides emaciatus.

LITUOLIDAE—Continued.

- Trochanmina squamata*.
globigeriniformis.

GLOBIGERINIDAE:

- Globigerina conglobata*.
sacculifera.
aequilateratus, var. *involuta*.
Sphaeroidina dehiscens.
Pullenia obliquiloculata.

MILIOLIDAE:

- Spiroloculina tenuiseptata*.
Sigmoilina schlumbergeri.

In many of the distribution tables no catalogue numbers appear, as mounts were not made in all cases. Material examined was recorded. In some of these cases, however, material was subsequently mounted and is available for study, even though the numbers do not appear in the tables.

A further study of these collections, especially in comparison with material from the Atlantic and other areas, will probably show that I have erred in referring many of the Philippine specimens to already described species. Notes on such cases will be made as the work progresses on the western Atlantic collections.

SYSTEMATIC TREATMENT.

Order FORAMINIFERA.

Family ASTRORHIZIDAE.

Subfamily ASTRORHIZINAE.

Genus ASTRORHIZA Sandahl, 1857.

ASTRORHIZA ANGULOSA H. B. Brady.

Astrorhiza angulosa H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 48, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 234, pl. 20, figs. 10-13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 20, fig. 1 (in text); Bull. 104; U. S. Nat. Mus., pt. 1, 1918, p. 10, pl. 3, fig. 2; pl. 4, figs. 1-3.

One specimen referable to this species was found in material dredged from *Albatross* station D5331, off western Luzon, 178 fathoms (325 meters); bottom temperature, 54.7° F. (12.6° C.).

Astrorhiza angulosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13310	U.S.N.M.	1	D5331 ...	° ' " ° ' "	178	° F. 54.7	s., sh., m...	Rare.

ASTRORHIZA GRANULOSA (H. B. Brady).

Plate 1, fig. 1.

Marsipella granulosa H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 36, pl. 3, figs. 8, 9.

Astrorhiza granulosa H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 48; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 234, pl. 20, figs. 14-23.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 21, fig. 5 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 12, pl. 5, fig. 4.

Specimens of this species occur in the material dredged by the *Albatross* south of the Equator. The wall in these specimens is very granular, the aperture and sometimes the tube partially closed by sponge spicules. The stations are as follows: D5613, off Celebes, 752 fathoms (1,375 meters), bottom temperature not given; D5650, Gulf of Boni, 540 fathoms (988 meters), bottom temperature 40.1° F. (4.5° C.); and D5668, Macassar Strait, 901 fathoms (1,648 meters), bottom temperature 38.2° F. (3.3° C.).

Astrorhiza granulosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13308	U.S.N.M.	D5550...	4 53 45 S.; 121 29 00 E....	540	° F. 40.1	gn. m.....	Few.
13309	U.S.N.M.	D5613...	0 42 00 S.; 121 44 00 E....	752	gy. m.....	Rare.
13305	U.S.N.M.	D5668...	2 28 15 S.; 118 49 00 E....	901	38.2	gy. m.....	Few.

ASTRORHIZA AGGLUTINATA, new species.

Description.—Test irregular in form, flattened, wall composed of agglutinated fragments of other foraminifera, at the type station mainly Globigerinidae; aperture near the border of the test.

Diameter up to 5 mm.

Type specimen.—(Cat. No. 13236, U.S.N.M.) from *Albatross* station D5630, south of Patiente Strait, in 569 fathoms (1,081 meters), where it is fairly common.

In its irregular form the species reminds one of *A. arenaria*.

Astrorhiza agglutinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13236	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E....	569	° F.	co. s., m....	Common.

Genus RHABDAMMINA W. B. Carpenter, 1869.

RHABDAMMINA ABYSSORUM W. B. Carpenter.

Plate 1, fig. 2.

Rhabdammina abyssorum M. Sars, Forh. Vid. Selsk. Christiana, 1868, p. 248 (*nomen nudum*).—W. B. CARPENTER, Ann. Mag. Nat. Hist., ser. 4, vol. 4, 1869, p. 288.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 266, pl. 21, figs. 1–13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 24, figs. 8–10, (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 15, pl. 6, fig. 1; pl. 7, fig. 1.

This species has occurred at more than 25 stations. The depths at these stations range from 124 to 940 fathoms (44 to 1,719 meters). There are two forms of this species met with in the dredgings—one with very coarse walls usually found in more shallow water and the other with fine material from deeper water. This division according to depth does not always hold true, however, as those with finely granular material are sometimes found in comparatively shallow water.

The distribution includes the following regions: East and south-east coasts of Mindanao; south and west coasts of Luzon; Palawan Passage; Jolo Sea, north of Tawi Tawi; between Negros and Siquijor;

north of Celebes; between Gillolo and Makyan Islands; Patiente Strait; Buton Strait; Gulf of Boni; and Flores Sea.

Rhabdammina abyssorum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperatures.	Character of bottom.	Abundance.
12740	U.S.N.M.	D5236...	80 50 45 N.; 126 26 52 E..	494	° F.	fne. gy. s...	Common.
13289	U.S.N.M.	8	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob. s. m.....	Common.
13969	U.S.N.M.	4	D5261...	12 30 55 N.; 121 54 24 E..	145	Few.
12743	U.S.N.M.	8	D5272...	14 00 00 N.; 120 22 30 E..	118	57.4	m. sh., Co...	Common.
12741								
13968	U.S.N.M.	3	D5277...	13 49 15 N.; 120 14 45 E..	80	58.6	fne. s.....	Few.
13970	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m. s.....	Rare.
13971	U.S.N.M.	2	D5331...	15 36 45 N.; 119 47 45 E..	178	54.7	s. sh. m.....	Rare.
12742	U.S.N.M.	D5348...	10 57 45 N.; 118 38 15 E..	375	56.1	Co., s.....	Frequent.
13972	U.S.N.M.	2	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.....	Rare.
13973	U.S.N.M.	2	D5425...	9 37 45 N.; 121 11 00 E..	495	49.4	gn. m., co. s.	Rare.
			D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Several.
13974	U.S.N.M.	6	D5429...	9 41 30 N.; 118 50 22 E..	766	gn. m.....	Several.
13975	U.S.N.M.	7	D5459...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.....	Several.
13976	U.S.N.M.	1	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.....	Rare.
13977	U.S.N.M.	6	D5449...	13 21 36 N.; 124 00 38 E..	300	Several.
			D5454...	13 12 00 N.; 123 50 30 E..	153
13978	U.S.N.M.	4	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.....	Few.
13979	U.S.N.M.	1	D5467...	15 35 27 N.; 123 37 18 E..	480	gy. m. (m. b.)	Rare.
12739	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m. (net).	Rare.
13980	U.S.N.M.	1	D5470...	13 37 30 N.; 123 41 09 E..	560	m.....	Rare.
13981	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gy. m. s.....	Few.
			D5570...	5 25 56 N.; 120 03 39 E..	277
13982	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Rare.
12738	U.S.N.M.	D5612...	0 38 00 N.; 121 45 40 E..	750	Common.
13983	U.S.N.M.	1	D5621...	0 15 00 N.; 124 35 00 E..	288	gy. & bk. s. (m. b.)	Rare.
			D5630...	0 56 30 S.; 128 05 00 E..	569
			D5640...	4 27 00 S.; 122 55 40 E..	24	s. brk. sh.
			D5651...	4 43 50 S.; 121 23 24 E..	700	38.7	gn. m.....
13984	U.S.N.M.	1	D5653...	3 32 40 S.; 120 31 30 E..	510	41.2	gy. m.....	Rare.
			D5668...	2 28 15 S.; 118 49 00 E..	901	38.2

RHABDAMMINA ABYSSORUM W. B. Carpenter, var. *RADIATA* Cushman.

Plate 1, fig. 3.

Rhabdammina abyssorum, var. *radiata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 652.

Description.—Test with a globular central chamber from which radiate numerous arms, with annular constrictions, largest diameter near the central chamber thence tapering toward the outer end; wall of fine texture, smooth, reddish-brown in color.

Type specimen.—Cat. No. 9101, U.S.N.M., from *Albatross* station D5654, Gulf of Boni, 805 fathoms (1,459 meters), bottom temperature 38.3° F. (3.3° C.).

Specimens were also frequent at D5650, 540 fathoms (988 meters), bottom temperature 40.1° F. (4.5° C.), and D5651, 700 fathoms (1,280 meters) bottom temperature 38.7° F. (3.7° C.), both in the Gulf of Boni, and D5640, Buton Strait, 24 fathoms (44 meters).

This certainly seems to be very different from the ordinary form of the species, the number of arms being as many as eight, the annular

constrictions and rapidly tapering arms also being distinctive characters. It was found only in this restricted region south of the Equator.

Rhabdammina abyssorum, var. *radiata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12744	U. S. N. M.	1	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Rare.
12736	U. S. N. M.	1	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.	Rare.
12735	U. S. N. M.	1	D5651...	4 43 50 S.; 121 23 24 E...	709	38.7	Rare.
11611	U. S. N. M.	1	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3	Rare.
12737	U. S. N. M.	1	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh ...	Rare.

RHABDAMMINA IRREGULARIS W. B. Carpenter.

Plate 1, fig. 5.

Rhabdammina irregularis W. B. CARPENTER, Proc. Roy. Soc. London, vol. 18, 1869, p. 60.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 268, pl. 21, fig. 9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., 1910, pt. 1, p. 26, figs. 11, 12 (in text); Bull. 104, U. S. Nat. Mus., 1918, pt. 1, p. 17, pl. 8, fig. 1.

In the region along the Equator and southward in the Gulf of Boni, Flores Sea, and Buton Strait this species is frequent at depths ranging from 24 to 1,560 fathoms (44 to 2,853 meters). It occurred at 20 stations in this area, the bottom temperatures, where given, ranging from 36.3° to 52.3° F. (2.3° C. to 11.2° C.). In one lot of material from D5385, Ragay Gulf, Luzon, 327 fathoms (598 meters), this species occurred with the bottom temperature recorded as 62.4° F. (16.8° C.). At some stations the great mass of the dredged material consisted of this species which was very similar to that from the Gulf of California. Some of the specimens show two sets of branches nearly an inch apart. Although the material was dredged in quantity it adds little to the knowledge of the structure or the initial portion of the test of this species. Evidently, as previously noted, there is a point of weakness just above the place of branching where the process of dredging seems sure to break the specimen. The following localities, mostly in the southern part of the region in deep water, had this species: Ragay Gulf, Luzon; between Gillolo and Makyan Islands; Buton Strait; Sibuko Bay, Borneo; between Gillolo and Kayoa Islands; Gulf of Boni; north of Celebes; Patiente Strait; and Flores Sea.

Rhabdammina irregularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13903	U.S.N.M.	1	D5385...	13 24 50 N.; 123 03 70 E...	327	62.4	gy. m.	Rare.
13904	U.S.N.M.	1	D5495...	9 06 30 N.; 125 00 20 E...	976	52.3	gy. m.	Rare.
12727	U.S.N.M.	1	D5591...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.	Rare.
12726	U.S.N.M.	1	D5609...	4 11 48 N.; 118 20 00 E...	260	gn. m.	Rare.
			D5612...	0 11 00 S.; 121 16 00 E...	1,092	36.3	Rare.
			D5612...	0 38 00 S.; 121 45 40 E...	750
13997	U.S.N.M.	1	D5621...	0 15 00 N.; 127 24 35 E...	298	gy. and bk. s. (m. b.).	Rare.
13998	U.S.N.M.	1	D5622 ..	0 19 20 N.; 127 28 30 E...	275	Rare.
			D5625...	0 07 00 N.; 127 28 00 E...	230	gy. m. fine. s.
			D5630...	0 56 30 S.; 128 05 00 E...	569	co. s. m.
12728	U.S.N.M.	1	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.	Rare.
			D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh.	Few.
12735								
12732								
12729	U.S.N.M.	10	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Common.
13995								
12733	U.S.N.M.	1	D5651...	4 43 50 S.; 121 23 24 E...	700	38.7	Rare.
13996	U.S.N.M.	5	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3	Few.
12734	U.S.N.M.	1	D5656...	3 17 40 S.; 120 36 45 E...	484	41.2	gy. m.	Rare.
13999	U.S.N.M.	1	D5658...	3 32 40 S.; 120 31 30 E...	510	41.2	gy. m.	Rare.
14000	U.S.N.M.	2	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.	Rare.
12731	U.S.N.M.	3	D5670...	1 19 00 S.; 118 43 00 E...	1,181	38.2	gy. m.	Few.

RHABDAMMINA DISCRETA H. B. Brady.

Rhabdopleura, species, G. M. DAWSON, Can. Nat., vol. 5, 1870, p. 177, fig. 7.

Rhabdopleura abyssorum G. M. DAWSON, Amer. Journ. Sci., vol. 1, 1871, p. 206, fig. 7.

Rhabdammina discreta H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 48; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 268, pl. 22, figs. 11–13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., 1910, pt. 1, p. 27, fig. 13 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 21, pl. 11, fig. 1.

At seven stations specimens referred to this species were found in rather small numbers. These stations are in the Sulu Sea; off the eastern coast of Luzon; and north of Celebes. In depth the stations are between 221 and 958 fathoms (404 and 1,752 meters), and the bottom temperatures from 41.2° to 50.1° F. (5.1° C. to 10° C.).

Rhabdammina discreta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12717	U.S.N.M.	1	D5127...	10 02 45 N.; 121 48 15 E...	958	50.1	gy. m., glob.	Rare.
			D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s.
			D5446...	12 43 51 N.; 124 59 18 E...	300	gn. m.
13992	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m. (net.).	Rare.
12716	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.	Rare.
12715	U.S.N.M.	2	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.	Few.
14885	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E...	221	gy. m., glob.	Rare.

RHABDAMMINA LINEARIS H. B. Brady.

Plate 1, fig. 4.

Rhabdammina linearis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 37, pl. 3, figs. 10, 11; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 269, pl. 22, figs. 1-6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., 1910, pt. 1, p. 28, figs. 14a-f (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 19, pl. 7, figs. 2-5.

Typical specimens of this species were obtained at nine stations with a rather different distribution from the preceding species, being more widely scattered: Off Manila Bay; Bouro Island; Jolo Sea; north of Celebes; Palawan Passage; and Patiente Strait. In depth these range from 297 to 1,105 fathoms (543 to 2,021 meters). The bottom temperatures where given range from 41.2° to 56.4° F. (5.1° C. to 13.5° C.).

Some of the specimens are very perfect and of large size. In a few cases the bottom material is recorded as coral sand, rather an unusual bottom condition for this species.

Rhabdammina linearis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8228	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Rare.
12725	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m. g.....	Rare.
12719	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Rare.
12720	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.....	Rare.
12721	U.S.N.M.	1	D5428...	9 13 00 N.; 118 51 15 E..	1,105	49.7	gy. m.....	Rare.
12722	U.S.N.M.	3	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Few.
12723	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E..	750	Rare.
12724	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E..	560	co. s., m.....	Rare.
			D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	

Genus MARSIPPELLA Norman, 1878.

MARSIPPELLA GIGANTEA Cushman.

Plate 1, fig. 6; plate 18, figs. 1, 2.

Marsipella gigantea CUSHMAN, Proc. U. S. Nat. Mus., vol. 42, 1912, p. 228, pl. 28, figs. 1, 2.

Description.—Test elongate, tubular, straight, or very gently and evenly curved, tapering somewhat from the narrow initial end, and again slightly tapering toward the apertural end; wall composed of elongate spicules firmly cemented by a reddish cement, spicules laid lengthwise of the test or slightly oblique, usually entire or nearly so; test of a reddish-brown color except near the apertural end, which is light gray.

Length up to 25 mm.; diameter in the widest portion, nearly 1 mm.

This species was described from *Albatross* station D5630, south of Patiente Strait, just south of the Equator, 569 fathoms (1,081 meters). One specimen was found at D5460, east coast Luzon,

565 fathoms (1,074 meters). The bottom temperature is not recorded for either of these stations.

This species is much larger than *M. cylindrica* H. B. Brady and has a tapering instead of a cylindrical form.

Marsipella gigantea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14005	U.S.N.M.	1	D5460 ..	° ' " ° ' "	565	° F.	gy. m.....	Rare.
13287	U.S.N.M.	3	D5630...	13 32 30 N.; 123 58 06 E..	569	co. s., m.....	Few.
8465	U.S.N.M.	2		0 56 30 S.; 128 05 00 E..				

MARSIPELLA CYLINDRICA H. B. Brady.

Marsipella cylindrica H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 714; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 265, pl. 24, figs. 20–22.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 30, figs. 15, 16 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 24, pl. 8, figs. 4–6; pl. 9, figs. 8, 9.

The only station at which this species was found is *Albatross* D5250, Gulf of Davao, 23 fathoms (42 meters).

But two specimens were found at this station, but those are typical in all respects.

Marsipella cylindrica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13285	U.S.N.M.	2	D5250...	° ' " ° ' "	23	° F.	co. s.....	Rare.
13286				7 05 05 N.; 125 39 45 E..				

Genus BATHYSIPHON G. O. Sars, 1871.

BATHYSIPHON FILIFORMIS G. O. Sars.

Plate 2, fig. 1.

Bathysiphon filiformis (M. Sars MS) G. O. Sars, Forh. Vid. Selsk. Christiania, 1871 (1872), p. 251.—NORMAN, Rep. Brit. Assoc., 1880, p. 389.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 248, pl. 26, figs. 15–20.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 31, text figs. 17–21 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 27, pl. 11, figs. 4, 5.

Description.—Test free, cylindrical, straight, or more often slightly curved; diameter uniform, tubular, open ends serving as apertures; wall composed of felted sponge spicules and fine sand in varying proportions, rather soft and even somewhat friable; color light gray or white, often with a superficial coating of a black deposit.

Diameter up to 2 mm.; length of Philippine specimens up to 15 mm.

Distribution.—This species occurred in considerable numbers from several stations as follows: *Albatross* station D5236, in 494 fathoms (903 meters) east coast of Mindanao; D5460, in 565 fathoms (1,074 meters), east coast of Luzon; D5468 in 569 fathoms (1,081 meters), east coast of Luzon; D5567, in 268 fathoms (490 meters), north of Tawi Tawi, bottom temperature 52° F. (11.1° C.); D5580, in 162 fathoms (297 meters), off Darvel Bay, Borneo, bottom temperature 55.8° F. (13.2° C.); D5585, in 476 fathoms (870 meters), Sibuko Bay, Borneo, bottom temperature 41.1° F. (5° C.); D5591, in 260 fathoms (476 meters), same locality; D5619, in 435 fathoms (796 meters), in Molucca Passage; D5622, in 275 fathoms (503 meters), between Gillolo and Makyan Islands; D5625, in 230 fathoms (421 meters), between Gillolo and Kayoa Islands, and D5668, in 901 fathoms (1,648 meters), in Flores Sea.

The specimens of the deeper, colder waters are larger and better developed than those in the shallower, warmer waters. Practically all the stations are in the deeper waters outside the protected waters of the Archipelago proper.

The texture of the wall is very different in appearance in this species from any of the others in the region.

Bathysiphon filiformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13404	U.S.N.M.	5	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s....	Common.
13415	U.S.N.M.	3	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.....	Few.
13958	U.S.N.M.	5	D5468...	13 35 39 N.; 123 40 28 E..	569	gn.m.(m.b.).	Rare.
13955	U.S.N.M.	1	D5567...	5 48 00 N.; 120 33 45 E..	268	52.0	fne. s.....	Rare.
13960	U.S.N.M.	2	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co....	Rare.
13961	U.S.N.M.	1	D5583...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Rare.
13962	U.S.N.M.	1	D5591...	4 11 48 N.; 118 38 20 E..	260	Rare.
13963	U.S.N.M.	3	D5619...	0 35 00 N.; 127 14 40 E..	435	fne. gy. s., m.	Rare.
13964	U.S.N.M.	2	D5622...	Makyan Id. (NE.); N. 66°	275	gy. m.....	Rare.
13965	U.S.N.M.	1	D5622...	W.				
13966	U.S.N.M.	4	D5625...	0 07 00 N.; 127 28 00 E..	230	gy. m., fne. s.	Few.
13967	U.S.N.M.	1	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	Rare.

BATHYSIPHON RUFUS de Folin.

Plate 2, fig. 2.

Bathysiphon rufum de FOLIN, Actes Soc. Linn. Bordeaux, vol. 40, 1886, p. 283, pl. 6, figs. 8a-c.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 267, pl. 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 32, fig. 22 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 29.

Description.—Test elongate, somewhat curved, decidedly tapering from the small initial end to the much larger last-developed portion; surface smooth and polished, wall comparatively thick, with numerous annular lines of growth and depressed areas, composed of sponge spicules and a very hard siliceous cement; color reddish-brown.

Length up to 10 mm.; diameter about 0.5 mm. in the widest part.

Distribution.—Specimens have occurred at several *Albatross* stations off the east coast of Luzon; Jolo Sea; Gulf of Tomini, Celebes; off Bouro Island; and Gulf of Boni. These stations range in depth from 310 to 1,181 fathoms (567 to 2,160 meters), the average being 744.3 fathoms (1,360 meters). The bottom temperatures where given average 41.2° F. (5.1° C.).

This species has a very different texture from the preceding, the wall being hard, the sponge spicules instead of being irregularly felted, are arranged longitudinally and cemented with a hard, siliceous cement with a smooth, polished surface.

Bathysiphon rufus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13413	U.S.N.M.	7	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Common.
13988	U.S.N.M.	3	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.....	Few.
13411	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Rare.
13985	U.S.N.M.	1	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.....	Rare.
13410	U.S.N.M.	1	D5468...	13 35 39 N.; 123 40 28 E..	569	gn. m.....	Rare.
13412	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	Few.
13414	U.S.N.M.	2	D5670...	1 19 00 N.; 118 43 00 E..	1,181	38.2	gy. m.....	Few.
13409	U.S.N.M.	1	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.....	Rare.
13407	U.S.N.M.	3	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s.....	Few.
13406	U.S.N.M.	1	D5127...	10 02 45 N.; 121 48 15 E..	958	50.1	gy. m., glob..	Rare.
13408	U.S.N.M.	3	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.....	Few.

BATHYSIPHON RUFESCENS Cushman.

Plate 2, figs. 3a, b.

Bathysiphon rufescens CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 651.

Description.—Test elongate, very slightly if at all curved, very slightly tapering, slender, wall marked by annular rings, surface rough, very light yellowish- or reddish-brown, dull.

Diameter up to 0.5 mm.; length up to 15 mm.

Distribution.—Type specimen (U.S.N.M. No. 9098) from *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean off eastern coast of Mindanao; bottom temperature 41.2° F. (5.1° C.). At this station the species was abundant. It also occurred at D5468, in 569 fathoms (1,081 meters), east coast of Luzon, and D5469, in 560 fathoms (1,024 meters), east coast of Luzon, D5639 in 1,560 fathoms (2,852 meters) in Molucca Sea. It was not found in the warmer, protected water of the Archipelago proper.

This species differs from *B. rufus*, which it most resembles in the rough, granular surface and the very slightly tapering, practically straight test.

Bathysiphons rufescens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11640	U.S.N.M.	7	D5236...	8 50 45 N.; 126 26 52 E....	494	41.2	fne. gy. s.....	Common.
13986	U.S.N.M.	4	D5468...	13 35 39 N.; 123 40 28 E....	569	gn. m.....	Few.
13987	U.S.N.M.	3	D5469...	13 37 30 N.; 123 41 09 E....	560	m.....	Few.
13405	U.S.N.M.	3	D5639...	3 54 50 S.; 123 27 20 E....	1,560	gy. m.....	Few.

BATHYSIPHON FLAVIDUS de Folin.

Bathysiphon flavidum de FOLIN, Actes Soc. Linn. Bordeaux, vol. 40, 1886, p. 279, pl. 6, figs. 5a-c.

Description.—Test elongate, subcylindrical, nearly straight, marked by annular lines and areas of growth, wall smooth, hard but not polished, light yellowish-brown except at the apertural end, which when complete is somewhat rounded and lighter colored.

Diameter 1–2 mm.; length up to 25 mm.

Distribution.—The species occurred usually in considerable numbers at the following *Albatross* stations: D5438, in 297 fathoms (543 meters), bottom temperature 46.2° F. (7.7° C.), and D5439, in 940 fathoms (1,719 meters), bottom temperature 36.7° F. (2.6° C.), both off the west coast of Luzon; D5447, in 310 fathoms (567 meters), bottom temperature 45.3° F. (7.2° C); D5460, in 565 fathoms (1,074 meters); D5468, in 569 fathoms (1,081 meters); D5469, in 500 fathoms (914 meters), and D5470, in 560 fathoms (1,024 meters), all off the east coast of Luzon; and D5622, in 275 fathoms (503 meters), between Gillolo and Makyan Islands.

This material seems to agree very well with de Folin's description and figure of this species. It is smooth but not polished, hard and not felted as in *B. filiformis*.

Bathysiphon flavidus—Material examined.

Cat. No.	Coil. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13949	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.	Rare.
13950	U.S.N.M.	4	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	s., Co., gn. m.	Few.
13951	U.S.N.M.	1	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.	Rare.
13952	U.S.N.M.	4	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m., Co., s	Few.
13953	U.S.N.M.	4	D5468...	13 35 39 N.; 123 40 28 E..	569	gn. m. (m. b.)	Few.
13954	U.S.N.M.	5	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m. (net)..	Few.
13955	U.S.N.M.	3	D5470...	13 37 30 N.; 123 41 09 E..	560	m.	Few.
			D5622...	Makyan Id. (N. E.), N. 66° W.	275	gy. m.	

BATHYSIPHON FLAVIDUS de Folin, var. GIGANTEUS Cushman.

Plate 2, figs. 4 a, b.

Bathysiphon flavidus de FOLIN, var. *giganteus* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 651.

Description.—Variety differing from the typical mainly in size, the test being much larger up to nearly 3 mm. in diameter and 40–50 mm. in length, the wall smooth and polished instead of dull, as in the smaller, typical form.

Distribution.—Type specimen (U.S.N.M. No. 9099) from *Albatross* station D5609, in 1,092 fathoms (3,131 meters), Gulf of Tomini, Celebes, bottom temperature 36.3° F. (2.3° C.). Specimens were also obtained from *Albatross* D5619, in 435 fathoms (796 meters), Molucca Passage; D5622, in 275 fathoms (503 meters), between Gillolo and

Makyan Islands; and D5668 in 901 fathoms (1,648 meters), Macassar Strait, bottom temperature 38.2° F. (3.3° C.).

Bathysiphon flavidus, var. *giganteus*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9099	U.S.N.M.	D5609...	00 11 00 S.; 121 16 00 E..	1,092	° F. 36.3	gn. m.....	
13956	U.S.N.M.	1	D5619...	00 35 00 S.; 127 14 40 E..	435	gy. m.....	Rare.
13957	U.S.N.M.	1	D5622...	Makyan Id. (N. E.), N. 60° W.	275	gy. m.....	Rare.
			D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	

BATHYSIPHON PAPYRACEUS Cushman.

Plate 2, figs. 5 a, b.

Bathysiphon papyraceus CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 651.

Description.—Test much elongate, irregularly curved, cylindrical, of even diameter; wall thin, friable, of sponge spicules with a small amount of cement, white in color, with the surface of light grayish-green in irregular bands; lines of growth very prominent.

Diameter up to 3 mm.; length up to 50 mm.

Distribution.—This species was extremely abundant at *Albatross* station D5247, in 135 fathoms (243 meters), in the Gulf of Davao. Type specimen, U.S.N.M. No. 9100.

It is very different from the other species of the genus in the very thin wall, which, however, is made up in the same way as the other friable species, and the irregular tube.

Bathysiphon papyraceus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9100	U.S.N.M.	10+	D5247...	7 02 N.; 125 38 45 E.....	135	° F.	m.....	Abundant.

Genus RHIZAMMINA H. B. Brady, 1879.

RHIZAMMINA INDIVISA H. B. Brady.

Plate 2, fig. 6.

Rhizammina indivisa H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 277, pl. 29, figs. 5-7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 34, fig. 24 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 32. pl. 12, figs. 7-10.

Tubular specimens with encrusting tests made up mostly of Globigerinidae and apparently belonging to this species were obtained from two stations, H4898, Sulu Sea, 221 fathoms (404 meters), and D5272, China Sea, off southern Luzon, 118 fathoms (216 meters). At the latter station the tubes were very common. At this station the bottom temperature was 57.4° F. (14.1° C.).

Rhizammia indivisa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12850	U.S.N.M.	3	D5272...	14 00 00 N.; 120 22 30 E..	118	° F. 57.4	m., sh., co. s. gy. m., glob.	Few.
			H4898...	7 43 45 N.; 122 03 45 E..	221		

RHIZAMMINA ALGAEFORMIS H. B. Brady.

Rhizammia algaeformis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 39, pl. 4, figs. 16, 17; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 274, pl. 28, figs. 1-11.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 272, pl. 15, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 33, fig. 23 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 31, pl. 11, figs. 2, 3.

There is a single specimen of this species from *Albatross* station D5201, Sogod Bay, southern Leyte, in 554 fathoms (1,012 meters). This is very similar to the specimens I have previously figured.³

Rhizammia algaeformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13239	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	° F. 52.8	gy. s., m.....	Rare.

Subfamily SACCAMMININAE.

Genus PSAMMOSPHAERA F. E. Schulze, 1875.

PSAMMOSPHAERA FUSCA F. E. Schulze.

Psammosphaera fusca F. E. SCHULZE, II Jahr. Comm. Wiss. Unt. deutsch. Meer in Kiel, 1875, p. 113, pl. 2, figs. 8a-f.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 249, pl. 18, figs. 1, 5-8 (not 2-4).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 35, figs. 25-28 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 35, pl. 13, figs. 1-6; pl. 14, figs. 1-3.

Both attached and free specimens have occurred in the dredgings, but the species has been noted at but three stations: D5236, 494 fathoms (903 meters), eastern coast of Mindanao; D5268, 170 fathoms (311 meters), Batangas Bay; and D5301, 208 fathoms (381 meters), China Sea. The bottom temperature at the latter station is given as 50.5° F. (10.2° C.), for station D5236, 41.2° F. (5.1° C.), and none is recorded for D5268.

Psammosphaera fusca—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13255	U.S.N.M.	D5301...	20 37 00 N.; 115 43 00 E..	208	° F. 50.5	gy. m., s.....	Rare.
13253	U.S.N.M.	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s.....	Rare.
13254	U.S.N.M.	D5268...	13 42 00 N.; 120 57 15 E..	170	s. p.....	Rare.

³ Bull. 104, pl. 11, fig. 2.

PSAMMOSPHAERA PARVA Flint.

Plate 2, fig. 7.

Psammospaera parva FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 268, pl. 9, fig. 1.—RHUMBLER, Arch. Protistk., vol. 3, 1903, p. 242, fig. 77 (in text).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 36, figs. 29, 30 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 35, pl. 12, figs. 4-6.

Psammospaera fusca (part) H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 250, pl. 18, figs. 2-4 (not 1, 5-8).

Typical specimens of this species occurred at D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (963 meters), bottom temperature 41.2° F. (5.1° C.); D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.); D5300, China Sea, vicinity of southern Luzon, 265 fathoms (485 meters), and D5601, Gulf of Tomini, Celebes, 765 fathoms (1,399 meters).

Psammospaera parva—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13256	U.S.N.M.	D5236...	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Rare.
13259	U.S.N.M.	D5601...	8 50 45 N.; 126 26 52 E..	765	s., glob., ptr.	Rare.
13258	U.S.N.M.	D5259...	1 13 10 N.; 125 17 05 E..	312	49.3	gy. m., glob.	Rare.
13257	U.S.N.M.	D5300...	11 57 30 N.; 121 42 15 E..	265	gy. m., s....	Rare.
				20 31 90 N.; 115 49 00 E..				

Genus IRIDIA Heron-Allen and Earland, 1914.

IRIDIA DIAPHANA Heron-Allen and Earland.

Iridia diaphana HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, pt. 12, 1914, p. 371, pl. 36.

This species, described by Heron-Allen and Earland from the Kerimba Archipelago, has not, so far as I have seen, occurred in its typical form in the Philippine material. There are two specimens, however, which have the thin membrane at the base, and are covered on the upper surface with sand grains and other foraminiferal tests. These may possibly belong to this species.

They are from D5201, Sogod Bay, southern Leyte, in 554 fathoms (1,012 meters), and D5282, China Sea, off southern Luzon, in 248 fathoms (454 meters).

Iridia diaphana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13322	U.S.N.M.	1	D5201...	° ' " ° ' "	554	° F. 52.8	gy. m., s....	Rare.
13323	U.S.N.M.	1	D5282...	10 10 00 N.; 125 04 15 E..	248	47.4	dk. gy. s....	Rare.
				13 53 00 N.; 120 26 45 E..				

Genus *SACCAMMINA* W. B. Carpenter, 1869.*SACCAMMINA SPHAERICA* G. O. Sars.

Saccammina sphaerica M. Sars, Forh. Vid. Selsk. Christiania, 1868 (1869), p. 248 (*nomen nudum*).—G. O. Sars, Forh. Vid. Selsk. Christiania, 1871, p. 250.—H. B. Brady, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 253, pl. 18, figs. 11–15, 17.—Cushman, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 39, figs. 33–36 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 44, pl. 16, figs. 4–5; pl. 19, figs. 2–5.

The typical form of this species has been seen but twice in this lot of material. It occurred at *Albatross* station D5260, off southeastern Mindoro, 234 fathoms (428 meters), bottom temperature 51.4° F. (10.7° C.), and D5447, off the eastern coast of Luzon, 310 fathoms (567 meters), bottom temperature 45.3° F. (7.2° C.).

Saccammina sphaerica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13221	U.S.N.M.		D5260...	° ' " ° ' "	234	° F. 51.4	gn. m., s.....	Rare.
13222	U.S.N.M.		D5447...	12 25 35 N.; 121 31 35 E... 13 28 00 N.; 123 46 18 E...	310	45.3	gn. m.....	Rare.

SACCAMMINA SPHAERICA G. O. Sars, var. *CATENULATA* Cushman.

Plate 2, fig. 8.

Saccammina sphaerica H. B. Brady (part), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 253, pl. 18, fig. 16.

Saccammina sphaerica, var. *catenulata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 652.

Description.—Wall of test like that of typical *S. sphaerica* in structure, but several chambers present, joined together in a linear series, the largest chamber adherent.

One specimen almost exactly like the figured specimen of Brady was found at *Albatross* station D5637, Bouro Island, 700 fathoms (1,280 meters). This seems to be different from the typical *S. sphaerica* and may be worthy of more than varietal rank. The several chambers attached by the largest make it a very different test. In the figure given by Brady a pebble forms the base for attachment; in this case the specimen is attached to another foraminifer.

Saccammina sphaerica, var. *catenulata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9450	U.S.N.M.		D5637...	° ' " ° ' "	700	° F.	gy. in.....	Rare.
				3 53 20 S.; 126 48 00 E....				

Genus *PROTEONINA* Williamson, 1858.*PROTEONINA DIFFLUGIFORMIS* (H. B. Brady).

Reophax difflugiformis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 51, pl. 4, figs. 3a, b; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 289, pl. 30, figs. 2-4 (not 1, 5).

Proteonina difflugiformis RHUMBLER, Arch. Protistk., vol. 3, 1903, p. 245, figs. 80a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 41, figs. 40, 41 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 47, pl. 21, figs. 1, 2.

The only stations at which this species occurred were *Albatross* D5126, Sulu Sea, vicinity southern Panay, 742 fathoms (1,357 meters), bottom temperature 49.5° F. (9.7° C.); D5300, 265 fathoms (485 meters), China sea, vicinity southern Luzon, no bottom temperature; and D5526, between Siquijor and Bohol Islands, 805 fathoms (1,459 meters), bottom temperature 52.3° F. (11.2° C.).

As this material is typical it is very probable that the species occurs more widely distributed, but is missing on account of the lack of very fine material from most of the stations.

Proteonina difflugiformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13437	U.S.N.M.	D5126...	10 34 45 N.; 121 47 30 E....	742	49.5	sft. gn. m ...	Rare.
13439	U.S.N.M.	D5526...	9 12 45 N.; 123 45 30 E....	805	52.3	gn. m., glob.	Rare.
13438	U.S.N.M.	D5300...	20 31 00 N.; 115 49 00 E....	265	gy. m., s.....	Rare.

PROTEONINA FUSIFORMIS Williamson.

Proteonina fusiformis WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 1, pl. 1, fig. 1.—RHUMBLER, Arch. Protistk., vol. 3, 1903, p. 248, fig. 84 (in text).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 41, fig. 39 (in text).

Reophax fusiformis H. B. BRADY, Denkschr. Kongl. Akad. Wiss. Wien, vol. 43, pt. 2, 1882, p. 99; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 290, pl. 30, figs. 7-11.

The only records for this species are two not entirely typical specimens from *Albatross* station D5551, Jolo Island. These have the general form of the species and are composed of sand grains and fine shell fragments.

These specimens are not unlike those figured by Heron-Allen and Earland,⁴ which they refer to *Thurammina papillata*.

Proteonina fusiformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
		2	D5551...	5 54 48 N.; 120 44 24 E....	193	53.3	fine s.....	Rare.

⁴ Journ. Roy. Micr. Soc., 1917, pl. 27, figs. 13-20.

PROTEONINA TESTACEA (Flint).

Reophax difflugiformis H. B. BRADY, var. *testacea* FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 273, pl. 16, fig. 1.

Proteonina testacea CUSHMAN, Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 49, pl. 20, figs. 1-4.

At the following six stations this species occurred, at some in very considerable numbers; D5201, Sogod Bay, southern Leyte, 554 fathoms (1,012 meters), bottom temperature, 52.8° F. (11.5° C.); D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature, 41.2° F. (5.1° C.); D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature, 49.3° F. (9.6° C.); D5281, China Sea, vicinity southern Luzon, 201 fathoms (368 meters), bottom temperature, 50.4° F. (10.2° C.); D5470, east coast of Luzon, San Bernardino Strait to San Miguel Bay, 560 fathoms (1,024 meters); and D5613, north of Celebes, 752 fathoms (1,375 meters).

These specimens, occurring in material containing a considerable number of Globigerinidae, are made almost entirely of such tests. The aperture is at the end of a definite and somewhat produced neck.

Proteonina testacea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15667	U.S.N.M.	10+	D5201...	10 10 00 N.; 125 04 15 E....	554	52.8	gy. m., s....	Common.
12812	U.S.N.M.	D5613...	0 42 00 S.; 121 44 00 E....	752	gy. m.	Common.
12809	U.S.N.M.	D5470...	13 37 30 N.; 123 41 09 E....	560	m.	Few.
			D5236...	8 50 45 N.; 126 26 52 E....	494	41.2	fine gy. s....	Few.
12811	U.S.N.M.	D5259...	11 57 30 N.; 121 42 15 E....	312	49.3	gy. m., glob.	Few.
12810	U.S.N.M.	D5281...	13 52 45 N.; 120 25 00 E....	201	50.4	dk. gy. s....	Few.

Genus TECHNITELLA Norman, 1878.

TECHNITELLA LEGUMEN Norman.

Technitella legumen NORMAN, Ann. Mag. Nat. Hist., ser. 5, vol. 1, 1878, p. 279, pl. 16, figs. 3, 4.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 246, pl. 25, figs. 8-12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 48, fig. 53 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 59, pl. 9, figs. 1, 2; pl. 10, fig. 1; pl. 16, figs. 7, 8; pl. 24, figs. 3-5; pl. 26, fig. 5.

Specimens of this species do not seem to be common in the region, but single specimens have occurred at six stations not very far apart. In all cases the specimens were typical. The stations are as follows: D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature, 52.8° F. (11.5° C.); D5236, eastern coast Mindanao, 494 fathoms (903 meters), bottom temperature, 41.2° F. (5.1° C.); D5242, Pujada Bay, 191 fathoms (350 meters), bottom temperature, 64.1° F. (17.8° C.); D5243, same general locality, 218 fathoms (399 meters), bottom temperature, 63.6° F. (17.5° C.); D5248, Gulf of Davao, 18 fathoms (33 meters); and D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature, 49.3° F. (9.6° C.). In all cases the bottom temperatures are high and not

what would be expected from the distribution of this species elsewhere.

These specimens show the typical structure of long acerate spicules arranged lengthwise of the test in the outer layer, and those of the inner layer arranged horizontally around the interior at right angles to those of the outside.

Technitella legumen—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12818	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E....	554	52.8	gy. s., m.....	Rare.
12819	U.S.N.M.	1	D5236	8 50 45 N.; 126 26 52 E....	494	41.2	fine gy. s.....	Rare.
		1	D5243	6 50 55 N.; 126 14 35 E....	218	63.6	gy. m.....	Rare.
		1	D5242	6 51 53 N.; 126 14 10 E....	191	64.1	soft gy. m.....	Rare.
11641	U.S.N.M.	1	D5248	7 07 25 N.; 125 40 24 E....	18	Co.....	Rare.
12820	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E....	312	49.3	gy. m., glob.	Rare.

Genus *THOLOSINA* Rumbler, 1895.

THOLOSINA BULLA (H. B. Brady).

Plate 3, figs. 1, 2; plate 4, fig. 4.

Placopsilina bulla H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 51; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 315, pl. 35, figs. 16, 17.

Tholosina bulla RHUMBLER, Nachr. kon. Ges. Wiss. Göttingen, 1895, p. 82.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 49, figs. 58a, b (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 63, pl. 25, fig. 6.

Specimens of this species attached to *Rhabdammina*, usually *R. irregularis*, and occasionally to *Hyperammina*, have been found at several stations. The specimens usually have a single, undivided chamber, but occasionally it is partially or wholly divided into two portions.

The typical form of the species somewhat broken and showing the interior is given in plate 4, figure 4. A very hispid variety is shown in plate 3, figure 1, and a form with a prominent aperture rising somewhat from the surface in plate 3, figure 2. This form of aperture seems always to be accompanied by a smooth surface.

The following localities are represented: Ragay Gulf, Luzon; Jolo Sea; between Negros and Siquijor, Sibuko Bay, Borneo; Gulf of Boni; and Flores Sea.

Tholosina bulla—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12826	U.S.N.M.	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.....	Rare.
12829	U.S.N.M.	D5650...	4 53 45 S.; 121 29 00 E....	540	40.1	gn. m.....	Rare.
12828	U.S.N.M.	D5585...	4 07 00 N.; 118 49 54 E....	476	41.1	gn. m.....	Rare.
12827	U.S.N.M.	D5424	9 37 05 N.; 121 12 37 E....	340	50.4	co. s.....	Rare.
12830	U.S.N.M.	D5658...	3 32 40 S.; 120 31 30 E....	510	41.2	gy. m.....	Rare.
12832	U.S.N.M.	D5259	11 57 30 N.; 121 42 15 E....	312	49.3	gy. m., glob.	Rare.
12833	U.S.N.M.	D5538...	9 08 15 N.; 123 23 20 E....	256	53.3	gn. m., s.....	Rare.
12834	U.S.N.M.	D5651...	4 43 50 S.; 121 23 24 E....	700	38.7	gn. m.....	Rare.

Genus **THURAMMINA** H. B. Brady, 1879.**THURAMMINA POPYRACEA** Cushman.

Plate 3, fig. 3.

Thurammina popyracea CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 637, pl. 79, fig. 4.

Description.—Test spherical, wall extremely thin and delicate, composed of fine sand grains, sponge spicules and a brownish cement; apertures very small and inconspicuous.

Diameter 1.5 mm.

Type specimen.—(Cat. No. 8511, U.S.N.M.) from *Albatross* station, D5613, north of Celebes, 752 fathoms (1,375 meters).

This is a large species with a paper-thin wall and very inconspicuous apertures.

Specimens were found at the following six stations: D5469, east coast of Luzon, 500 fathoms (914 meters); D5470, the neighboring station, 560 fathoms (1,024 meters); D5612, Gulf of Tomini, Celebes, 750 fathoms (1,370 meters); D5613, 752 fathoms (1,375 meters); D5637 south of Patiente Strait, 700 fathoms (1,280 meters); and D5668 Macassar Strait, 901 fathoms (1,648 meters). The bottom temperature was given in but one case—D5668, 38.2° F. (3.4 C.°)

Thurammina popyracea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12791	U.S.N.M.	1	D5469...	13 36 48 N.; 123 33 24 E...	500	° F.	gn. m.....	Rare.
12792	U.S.N.M.	2	D5470...	13 37 30 N.; 123 41 09 E...	560	m.....	Rare.
12793	U.S.N.M.	4	D5612...	0 38 00 S.; 121 45 40 E...	750	Few.
8511 12749	U.S.N.M.	3	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.....	Few.
12795	U.S.N.M.	2	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Few.
12796	U.S.N.M.	3	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.....	Few.

THURAMMINA PAPILLATA H. B. Brady, var. **CASTANEA** Heron-Allen and Earland.

Thurammina papillata H. B. BRADY, var. *castanea* HERON-ALLEN and EARLAND, Journ. Roy. Micr. Soc., 1917, p. 545, pl. 26, figs. 14-18; pl. 29, fig. 17.

There is a single specimen from D5236, Pacific Ocean, east coast of Mindano, in 494 fathoms (903 meters) which has a reddish-brown color and numerous blunt papillae, with a very fine opening at the end of each.

This is evidently the variety which Heron-Allen and Earland describe and figure in the above reference. No other specimens refer-

able either to this variety or to the species were found in the region, all others being referred to our species, *T. papyracea*.

Thurammia papillata, var. *castanea*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13238	U.S.N.M.	1	D5236...	° ' " ° ' "	494	° F. 41.2	fine gy. s. . . .	Rare.

Subfamily HYPERAMMININÆ.

Genus HYPERAMMINA H. B. Brady, 1878.

HYPERAMMINA ELONGATA H. B. Brady.

Plate 3, fig. 5.

Hyperammia elongata H. B. BRADY (part), Ann. Mag. Nat. Hist., ser. 5, vol. 1, 1878, p. 433, pl. 20, figs. 2a, b; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 257, pl. 23, figs. 4, 7 (not 9, 10).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 60, figs. 73, 74 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 74, pl. 29, fig. 4.

Tubular specimens referable to this species have been obtained at a number of stations well scattered throughout the region, depths varying from 248 to 1,105 fathoms (454 to 2,021 meters). As a rule the proloculum of specimens is missing, but was present in one complete specimen among others from D5470 in 560 fathoms (1,065 meters), off the east coast of Luzon. The smooth species, *H. laevigata*, was not found at all in any of the dredgings.

Hyperammia elongata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12797	U.S.N.M.	6	D5219...	° ' " ° ' "	530	° F. 50.8	gn. m.	Few.
			D5259...		312	49.3	gy. m.	
			D5282...		248	47.4	dk. gy. s.	
			D5428...		1,105	49.7	gy. m.	Few.
12798	U.S.N.M.	5	D5467...		480	gy. m. (m. b.).	
12799	U.S.N.M.	1	D5469...		500	gn. m. (net) ..	
12800	U.S.N.M.	3	D5470...		560	m.	
			D5523...		
			D5609...		1,092	36.3	gn. m.	Rare.
			D5630...		569	co. s., m.	
12801	U.S.N.M.	2	D5650...		540	40.1	gn. m.	

HYPERAMMINA FRIABILIS H. B. Brady.

Plate 3, fig. 4.

Hyperammina elongata H. B. BRADY (part), Ann. Mag. Nat. Hist., ser. 5, vol. 1, 1878, p. 433.

Hyperammina friabilis H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 258, pl. 23, figs. 1-3, 5, 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 62, fig. 76 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 75, pl. 29, figs. 1-3.

Very fine and complete specimens, showing the proloculum intact, were found at a number of stations in the region.

It is worth noting here that in the Atlantic *H. friabilis* was found in comparatively warm waters, while *H. subnodosa* was limited to very low temperatures. As all these stations in the Philippine area here noted have relatively high temperatures, the frequency of *H. friabilis* and complete lack of *H. subnodosa* is very significant.

The localities for this species in the region include China Sea, off Hongkong; Palawan Passage; Jolo Sea; eastern Palawan; between Leyte and Mindanao; between Negros and Siquijor; north of Celebes; off Bouro Island; and in Buton Strait.

Hyperammina friabilis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
15344	U. S. N. M.	6	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
12803	U. S. N. M.	1	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m. s.....	Rare.
12802	U. S. N. M.	4	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co. s.....	Few.
12804	U. S. N. M.	2	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Few.
12805	U. S. N. M.	3	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Few.
			D5428...	9 13 00 N.; 118 51 15 E...	1,105	49.7	gy. m.....	
12806	U. S. N. M.	3	D5495...	9 06 30 N.; 125 00 28 E...	976	52.3	gy. m.....	Few.
12807	U. S. N. M.	3	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s.....	Few.
12808	U. S. N. M.	10+	D5612...	0 38 00 S.; 121 45 40 E...	750	Common
			D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	
			D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk.sh...	

Genus SACCORHIZA Eimer and Fickert, 1899.

SACCORHIZA RAMOSA (H. B. Brady).

Plate 4, fig. 5.

Hyperammina ramosa H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 33, pl. 3, figs. 14, 15; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 261, pl. 23, figs. 15-19.

Saccorhiza ramosa EIMER and FICKERT, Zeitschr. Wiss. Zool., vol. 65, 1899, p. 670.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 65, fig. 81 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 81, pl. 30, figs. 3, 4.

Specimens of this species have been noted in material from 20 or more stations in the region. A considerable number of these stations were off the east coast of Luzon, in 310 to 569 fathoms (567 to 1,081 meters), bottom temperature, where recorded, 44.3° F. (6.8° C.).

In addition there were scattered stations covering the whole area rather well.

It will be noted that the bottom temperatures from some of these stations were relatively high and yet much of the material was from deep water. A number of specimens were obtained with the proloculum intact. There seem to be both microspheric and megalospheric forms with smaller but longer and more branching tubes, and the latter form of the specimens showed tubes twice branched. There is some considerable variation in regard to the comparative amount of included spicules.

Saccorhiza ramosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
			D5172...	6 03 15 N.; 120 35 30 E...	318	fine. s., sh.....	
			D5212...	12 04 15 N.; 124 04 36 E...	108	59.9	gy. s., m.....	
12776	U.S.N.M.	4	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s.....	Few
12777	U.S.N.M.	4		7 34 45 N.; 126 38 15 E...	380	43.0	gn. m.....	Few.
12768	U.S.N.M.	4	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.....	Few.
13940	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Few.
12770	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Rare.
12772	U.S.N.M.	3	D5439...	15 53 15 N.; 119 40 20 E...	940	36.7	gn. m.....	Few.
13941	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m. s.....	Few.
13942	U.S.N.M.	5	D5447...	13 28 00 N.; 123 46 18 E...	310	45.3	gn. m.....	Few.
12773	U.S.N.M.	3	D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m. (m. b.)	Rare.
13943	U.S.N.M.	3	D5468...	13 35 39 N.; 123 40 28 E...	569	gn. m. (m. b.)	Few.
27711	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m. (net.)	Rare.
			D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s.....	
13944	U.S.N.M.	1	D5535...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.....	Rare.
12774	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E...	750	Rare.
13945	U.S.N.M.	5	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.....	Few.
13946	U.S.N.M.	1	D5621...	0 15 00 N.; 127 24 35 E...	298	gy. & bk. s. (m. b.)	Few.
13947	U.S.N.M.	2	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.....	Few.
13775	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Few.
12762	U.S.N.M.	5	Nero 842	15 22 00 N.; 122 03 30 E...	1,330	br. m., s.....	Few.

Genus TOLYPAMMINA Rumbler, 1895.

TOLYPAMMINA VAGANS (H. B. Brady).

Plate 4, figs. 2, 3; plate 7, figs. 1, 2.

Hyperammima vagans H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 33, pl. 3, fig. 5; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 260, pl. 24, figs. 1-9.

Tolypammima vagans RHUMBLER, Nachr. kön. Ges. Wiss. Göttingen, 1895, p. 83; Zeitschr. allg. Phys., vol. 2, 1902, p. 281, fig. 97; Arch. Protistk., vol. 3, 1903, p. 277, figs. 125a, b (in text).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 67, fig. 85 (in text).

Serpulella vagans EDMER and FICKERT, Zeitschr. wiss. Zool., vol. 65, 1899, p. 674.

Girvanella vagans RHUMBLER, Foram. Plankton Exped., pt. 1, 1911, pl. 4, figs. 1, 2; pt. 2, 1913, p. 419.—CUSHMAN, Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 91, pl. 35, figs. 4, 5; pl. 36, fig. 1.

At 15 or more stations numerous typical specimens of this species have occurred. They have been found attached to *Rhabdammina* most frequently, but also have been noted upon *Hormosina glob-*

ulifera, *Trochammina turbinata*, *Haplophragmoides subglobosum*, *Ammodiscus incertus*, *Bathysiphon* and *Bigenerina*.

The localities include the east and west coasts of Luzon; east coast of Mindanao, Palawan Passage; Sibuko Bay, Borneo; north of Celebes; Molucca Sea; Gulf of Boni; and Flores Sea.

Tolypammina vagans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
5234 12779 12781	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fine. gy. s....	Common.
D5348...			7 07 25 N.; 125 40 24 E..	18	Co.....	
D5385...			13 24 50 N.; 123 03 70 E..	327	62.4	gy. m.....	
14002	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Rare.
12786	U.S.N.M.	1	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.....	Rare.
12785	U.S.N.M.	1	D5449...	13 21 36 N.; 121 00 30 E..	300	Rare.
14001	U.S.N.M.	1	D5468...	13 35 39 N.; 123 40 28 E..	569	gn. m. (m.b).	Rare.
12778	U.S.N.M.	2	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Rare.
12780	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E..	750	Rare.
			D5639...	3 54 08 S.; 123 27 20 E..	1,500	gy. m.....
12782	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Rare.
12783	U.S.N.M.	1	D5651...	4 43 50 S.; 121 23 24 E..	700	38.7	Rare.
			D5654...	3 42 00 S.; 120 45 50 E..	805	38.3
			D5658...	3 32 40 S.; 120 31 30 E..	310	41.2	gy. m.....
			D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....
12784	U.S.N.M.	2	D5670...	1 19 00 S.; 118 43 00 E..	1,181	38.2	gy. m.....	Rare.
14763	U.S.N.M.	1	Nero 838	15 25 00 N.; 122 12 40 E.	1,406	br., m., s....	Rare.

Genus *DENDROPHRYA* Str. Wright, 1861.

DENDROPHRYA RAMOSA Cushman.

Plate 18, figs. 7, 8.

Dendrophrya ramosa CUSHMAN, Proc. U. S. Nat. Mus., vol. 42, 1912, p. 227, pl. 28, figs. 7, 8.

Description.—Test large, laterally compressed, branching, wall with a chitinous base, over which there is a thick layer of mud or fine sand and scattered sponge spicules; apertures at the ends of the branching portions, elliptical.

Length 10 mm. or more.

Type specimen.—Cat. No. 8463, U.S.N.M., from *Albatross* station D5385, Ragay Gulf, Luzon, 327 fathoms (598 meters), with a bottom temperature of 62.4° F. (16.8° C.). Other stations at which this species occurred are D5438, west coast Luzon, 297 fathoms (543 meters), bottom temperature, 46.2° F. (7.8° C.); D5460, east coast Luzon, 565 fathoms (1,074 meters); D5470, 560 fathoms (1,065 meters); D5622, between Gillolo and Makyan Islands, Dutch East Indies, 275 fathoms (503 meters); D5630, south of Patiente Strait, 569 fathoms (1,081 meters); and D5650, Gulf of Boni, 540 fathoms (988 meters); bottom temperature, 40.1° F. (3.5° C.).

This species evidently belongs to the genus *Dendrophrya*, as described, but is much larger than either of the other two described species. The figured specimen is evidently only a fragment, as are

all the specimens seen, and the complete test must be considerably larger. There is some variation in the color and texture of the wall and the comparative amount of spicules, depending somewhat upon the bottom conditions. All the stations are in fairly deep water compared with the shore conditions under which the other two known species were found. Usually specimens were very numerous, making up the large mass of the dredged material at the type station.

Dendrophrya ramosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13105	U.S.N.M.	D5438	° ' " ° ' "	297	° F. 46.2	gn. m.	Few.
			D5460	15 54 42 N.; 119 44 42 E..	565	gy. m.	
13103	U.S.N.M.	D5630	13 32 30 N.; 123 58 06 E..	509	co. s., m.	Rare.
13102	U.S.N.M.	D5650	0 56 30 S.; 128 05 00 E..	540	40.1	gn. m.	Rare.
13104	U.S.N.M.	D5630	4 53 45 S.; 121 29 00 E..	569	co. s., m.	Rare.
13228 8463	U.S.N.M.	D5385	0 56 30 S.; 128 05 00 E..	327	62.4	gy. m.	Common.
			D5470	13 24 50 N.; 123 03 70 E..	500	m.	
			D5622	13 37 30 N.; 123 41 09 E..	275	gy. m.	
				Makyan Id. (NE.), N. 66° W.				

DENDROPHRYA RAMOSA Cushman, var. **ROBUSTA**, new variety.

Description.—This variety differs from the typical in having a coarser, broader test, with the branches less flattened and the main portion of the test broad and flattened.

Length 8 mm. (type specimen).

Type specimen.—(Cat. No. 13235, U.S.N.M.) from *Albatross* station D5637, south of Bouro Island, in 700 fathoms (1,280 meters).

This has a very different external appearance from the typical form of the species, mostly in its coarse and broad test.

Dendrophrya ramosa, var. *robusta*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13235 11625	U.S.N.M.	10+	D5637	° ' " ° ' "	700	° F.	gy. m.	Common.
				3 53 20 S.; 120 48 00 E....				

DENDROPHRYA ATTENUATA Cushman.

Plate 4, fig. 1.

Dendrophrya attenuata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 652.

Description.—Test elongate, compressed, consisting of a central, elongated body, with slightly projecting, apertural branches on either border, irregularly alternating; wall composed of sand grains with many spicules; color light gray.

Length up to 15 mm.

Distribution.—Type specimen (U.S.N.M. No. 9102) from *Albatross* station D5670 in 1,181 fathoms (2,160 meters), Macassar Strait, bottom temperature 38.2° F. (3.4° C.). It also occurred at D5622, in 275 fathoms (503 meters), between Gillolo and Makyan Islands and at D5619 in 435 fathoms (796 meters), Molucca Passage. At this last station it was very abundant, so much so that it might almost be called a *Dendrophrya attenuata* bottom, as this made up the great mass of the washed material at this station.

Dendrophrya attenuata.—*Material examined.*

Cat. No.	Coll. of—	No. of specimens.	Stations.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11621 9102	} U.S.N.M.	D5670...	1 19 00 S.; 118 43 00 E....	1,181	38.2	gy. m.....	Few.
			D5619...	0 35 00 N.; 127 14 40 E....	435	fine gy. s., m.	Abundant.
			D5622...	Makyan Id. (N.E.), N. 66° W.	275	gy. m.....	

Genus HALIPHYSEMA Bowerbank, 1862.

HALIPHYSEMA CATENULATA Cushman.

Plate 18, figs. 5, 6.

Haliphysema catenulata CUSHMAN, Proc. U. S. Nat. Mus., vol. 42, 1912, p. 228, pl. 28, figs. 5, 6.

Description.—Test elongate, tapering, with numerous constrictions as though segmented; wall arenaceous, with many included fragments of sponge spicules and with a crown of elongated spicules about the aperture; basal end with a sort of expanded portion like a proloculum, also made up largely of broken spicules; aperture terminal, circular, surrounded by the crown of erect spicules.

Length about 10 mm.

Type specimen.—Cat. No. 8464, U.S.N.M., from *Albatross* station D5630, south of Patiente Strait, Dutch East Indies, 569 fathoms (1,081 meters). The species was very common at this station, but was not met with elsewhere in the dredgings.

The spicules throughout the test except at the apertural tip are broken fragments, but about the aperture there are elongate spicules not entire, but so placed that they rise up above the aperture like a crown.

Haliphysema catenulata.—*Material examined.*

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11634 8464	} U.S.N.M.	10+	D5630...	0 56 30 S.; 128 05 00 E....	569	co. s., m....	Common.
				• • • • •		° F.		

Genus *JACULELLA* H. B. Brady, 1879.*JACULELLA ACUTA* H. B. Brady.

Plate 3, fig. 7.

Jaculella acuta H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 35, pl. 3, figs. 12, 13; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 225, pl. 22, figs. 14-18.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 70, figs. 90, 91 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 84, pl. 32, figs. 1-4.

Specimens of this species have been noted from five stations in the material examined: D5212, east of Masbate Island, 108 fathoms (198 meters), bottom temperature 59.9° F. (15.5° C.); D5236, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.); D5454, Jolo Sea, 153 fathoms (279 meters), and D5467, east coast of Luzon, 480 fathoms (878 meters); bottom temperatures in the last two cases not recorded.

The specimens were typical, and at station D5212 were fairly numerous.

Jaculella acuta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Stations.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13424	U.S.N.M.	D5454...	13 12 00 N.; 123 50 30 E...	153	Rare.
13423	U.S.N.M.	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s.	Rare.
13421	U.S.N.M.	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
13422	U.S.N.M.	D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m.	Rare.
13425	U.S.N.M.	D5212...	12 04 15 N.; 124 04 36 E...	108	59.9	gy. s., m.	Few.

JACULELLA OBTUSA H. B. Brady.

Plate 3, fig. 6.

Jaculella obtusa H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 714; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 256, pl. 22, figs. 19-22.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 71, fig. 92 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 85, pl. 32, fig. 5.

This species was seen in material from three stations: D5236, off the east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5291, China Sea, vicinity southern Luzon, 173 fathoms (316 meters), bottom temperature 51.5° F. (10.8° C.); and D5445, east coast of Luzon, 383 fathoms (699 meters), bottom temperature 44.3° F. (6.8° C.). The specimens from the last station were the most numerous and most nearly typical of any.

Jaculella obtusa—material examined.

Cat. No.	Coll. of—	No. of specimens.	Stations.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13420	U.S.N.M.	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s.	Rare.
13419								
13417								
13418	U.S.N.M.	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.	Few.
	U.S.N.M.	D5291...	13 29 40 N.; 121 09 45 E...	173	51.5	fne. bk. s.	Few.

Genus *SAGENINA* Chapman, 1900.*SAGENINA FRONDESCENS* (H. B. Brady).

Plate 9, fig. 1; plate 14, fig. 1.

Sagenella frondescens H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 41, pl. 5, fig. 1; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 278, pl. 28, figs. 14, 15.

Sagenina frondescens CHAPMAN, Journ. Linn. Soc., vol. 28, 1900, p. 4, pl. 1, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 71, fig. 93 (in text).

Occurring as single specimens, or, at most, rarely, this species has been noted from three stations: D5152, Sulu Archipelago, Tawi Tawi Group, 24–34 fathoms (44–62 meters); D5192, off north Cebu Island, 32 fathoms (58 meters); D5217, between Burias and Luzon, 105 fathoms (193 meters), bottom temperature 63.1° F. (17.2° C.).

As previous records indicate this seems to be a species frequenting shallow, warm water. It is easily overlooked and is probably much more common in the region than the records indicate.

Sagenina frondescens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Stations.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8230	U.S.N.M.	1	D5152...	° ' " ° ' "		° F.		
8231	U.S.N.M.	1	D5192...	5 22 55 N.; 120 15 45 E...	34	wh. s.	Rare.
12885				11 9 15 N.; 123 50 00 E...	32	gn. s.	Rare.
13337	U.S.N.M.	2	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s. ...	Rare.

SAGENINA DIVARICANS Cushman.

Plate 8.

Sagenina divaricans CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 437, fig. 1.

Description.—Test attached, tubular, very slender, uniform in size, branching at nearly regular intervals, with a wide angle, sometimes anastomosing; walls of coralline mud, somewhat roughened; apertures at the ends of the tubes; color, white.

Diameter of the tubes, 0.05 to 0.1 mm.

Type specimen.—(Cat. No. 8229, U.S.N.M.) from *Albatross* station 5145, vicinity of Jolo, Jolo Archipelago, 23 fathoms (42 meters); attached to hardened coral sand on the interior of a bivalve shell. It also occurred at D5276, China Sea, vicinity of southern Luzon, 18 fathoms (33 meters), attached to *Textularia*. A single specimen attached to *Textularia rugosa* was found in material from station D5134, 25 fathoms (46 meters), Sulu Archipelago, near Basilan Island. The two other stations in the area at which it has occurred are D5148, Sulu Archipelago, vicinity of Siasi, 17 fathoms (31 meters), and D5239, Pujada Bay and vicinity, 171 fathoms (312 meters).

Sagennina divaricans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13333	U.S.N.M.	3	D5134...	6 44 45 N.; 121 48 00 E..	25	fne. s.....	Few.
13334	U.S.N.M.	1	D5145...	6 4 30 N.; 120 59 30 E..	23	co. s., sh....	Rare.
8229	U.S.N.M.	1	D5148...	5 35 40 N.; 120 47 30 E..	17	co. s.....	Rare.
13335	U.S.N.M.	1	D5239...	6 49 8 N.; 126 15 12 E..	171	sft. gy. m....	Rare.
12884	U.S.N.M.	1	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s....	Rare.

Subfamily AMMODISCINAE.

Genus AMMOLAGENA Eimer and Fickert, 1899.

AMMOLAGENA CLAVATA (Parker and Jones).

Plate 6, figs. 1-4; plate 10, figs. 3, 4.

Trochammmina irregularis, var. *clavata* PARKER and JONES, Quart. Journ. Geol. Soc., vol. 16, 1860, p. 304.

Trochammmina irregularis (part) W. B. CARPENTER, PARKER, and JONES, Introd. Study Foram., 1862, p. 142, pl. 11, fig. 6.

Webbina clavata H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 711; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 349, pl. 41, figs. 12-16.

Ammolagena clavata EIMER and FICKERT, Zeitschr. Wiss. Zool., vol. 65, 1889, p. 673.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 68, figs. 86-89 (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 89. pl. 34, figs. 2-5; pl. 35, figs. 1-3.

Webbinella clavata RUMBLER, Arch. Protistk., vol. 3, 1903, pt. 229, fig. 55 (in text).

Both microspheric and megalospheric forms of this species have been obtained in this material, the microspheric form having a smaller but much longer tube than the megalospheric form; also the megalospheric form is much more common. The attachment is usually to smooth surfaces or ones that are comparatively so. I have noted its occurrence on the following genera: *Rhabdammina*, *Ammodiscus*, *Reophax*, *Cyclammmina*, *Haplophragmoides*, *Trochammmina*, *Bulimina*, *Cristellaria*, *Pulvinulina*, and *Biloculina*. It also occurs on bits of shell and various foreign bodies.

There is an interesting specimen from D5236, Pacific Ocean, east coast of Mindanao, attached to a species of *Quinqueloculina*, which apparently represents the first chamber only of *A. clavata*. This has the appearance of *Webbinella hemisphaerica*, but at the edge is a very small definite aperture. As *A. clavata* is common at this station this may represent the young stage of this species which has not yet developed the tubular chamber.

Ammolagena clavata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
8208	U. S. N. M.	1	D5236...	8 50 45 N.;	126 26 52 E..	494	41.2	fne. gy. s....	Common.
13431	U. S. N. M.	D5238...	7 34 45 N.;	126 38 15 E..	380	43	gn. m.....	Rare.
			D5282...	13 53 00 N.;	120 26 45 E..	248	47.4	dk. gy. s....	Rare.
14063	U. S. N. M.	1	D5447...	13 28 00 N.;	123 46 18 E..	310	45.3	gn. m.....	Rare.
13428	U. S. N. M.	1	D5459...	13 10 21 N.;	59 54 00 E..	201	Rare.
13426	U. S. N. M.	1	D5460...	13 32 30 N.;	123 58 06 E..	565	gy. m.....	Rare.
13429	U. S. N. M.	3	D5582...	4 19 54 N.;	58 38 00 E..	890	38.3	gy. m., fne. s.	Few.
12788	U. S. N. M.	1	D5591...	4 11 48 N.;	38 20 00 E..	260	Rare.
13427	U. S. N. M.	D5637...	3 33 20 S.;	48 00 00 E..	700	gy. m.....	Rare.
14004	U. S. N. M.	1	D5639...	3 54 50 S.;	123 27 20 E..	1,560	gy. m.....	Rare.
12789	U. S. N. M.	D5642...	4 31 40 S.;	122 49 42 E..	37	gy. m.....	Rare.
14886	U. S. N. M.	6	H4897...	7 46 00 N.;	122 00 00 E..	1,570	gy. m., glob..	Few.
13430									
13432	U. S. N. M.	10	D5236						Common.
13433									
13434									
12790	U. S. N. M.	3	D5650...	4 53 45 S.;	121 29 00 E..	540	40.1	gn. m.....	Few.
13435	U. S. N. M.	D5667...	5 48 00 N.;	120 33 45 E..	268	52.0	fne. s.....	Rare.
12787	U. S. N. M.	1							

¹ See No. 1 above.

Genus *AMMODISCUS* Reuss, 1861.*AMMODISCUS INCERTUS* (d'Orbigny).

Plate 5, figs. 1, 2.

Operculina incerta D'ORBIGNY, in De la Sagra. Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 49, pl. 6, figs. 16, 17.

Ammodiscus incertus H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 330, pl. 38, figs. 1-3.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 278, pl. 23, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 73, figs. 95, 96a, b (in text); Bull. 104, U. S. Nat. Mus., pt. 1, 1918, p. 95, pl. 39.

Ammodiscus tenuis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 51; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 332, pl. 38, figs. 4-6.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 279, pl. 23, fig. 1.

Specimens of this species were very numerous and were found at a large number of stations, especially in the southern portion of the area. Depths vary from 24-1,570 fathoms (44-2,871 meters) and bottom temperatures, were recorded, from 38.2° to 52.0° F. (3.4° C. to 11.1° C.). At several stations where a number of specimens were found a record was kept of the number of microspheric and megalospheric specimens, as follows:

	Micro-spheric.	Megalo-spheric.		Micro-spheric.	Megalo-spheric.
14.....	11	3	5.....	5	0
6.....	2	4	12.....	10	2
7.....	1	6	13.....	1	12
3.....	0	3	50.....	9	41
3.....	0	3	12.....	0	12
4.....	0	4	9.....	3	6
3.....	1	2	25.....	0	25
30.....	18	12			
3.....	0	3	199	61	138

If record had been kept of other cases where a few megalospheric specimens were found and no microspheric ones, the preponderance of megalospheric specimens would be much greater.

Ammodiscus incertus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13927	U.S.N.M.	6	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m....	Few.
12632	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Common.
12633		9						
12634	6							
12635	U.S.N.M.	10+	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Common.
13922	U.S.N.M.	3	D5268...	13 42 00 N.; 120 57 15 E...	170		s. p.	Few.
			D5272...	14 00 00 N.; 120 22 30 E...	118	57.4	m. sh., co. s..	
13923	U.S.N.M.	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s....	Rare.
		1	D5331...	15 36 45 N.; 119 47 45 E...	178	54.7	s., sh., m....	Rare.
12636	U.S.N.M.	7	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.	Few.
			D5385...	13 24 50 N.; 123 03 70 E...	327	62.4	gy. m.	
13919	U.S.N.M.	4	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.	Few.
13921	U.S.N.M.	1	D5428...	9 13 00 N.; 118 51 15 E...	1,105	49.7	gy. m.	Few.
13934	U.S.N.M.	2	D5429...	9 41 30 N.; 118 50 22 E...	766		gn. m.	Rare.
			D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.	
13924	U.S.N.M.	2	D5439...	15 58 15 N.; 119 40 20 E...	940	36.7	gn. m.	Rare.
			D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s....	
13920	U.S.N.M.	D5446...	12 43 51 N.; 124 53 18 E...	300		gn. m.	
		6	D5447...	13 28 00 N.; 123 46 18 E...	310	45.3	gn. m.	Few.
13950	U.S.N.M.	1	D5449...	13 21 36 N.; 124 00 30 E...	300		Rare.
12638	U.S.N.M.	3	D5460...	13 32 30 N.; 123 58 06 E...	565		gy. m.	Few.
13925	U.S.N.M.	1	D5467...	13 35 27 N.; 123 37 18 E...	480		gy. m. (m.b.)	Rare.
			D5567...	5 48 00 N.; 120 33 45 E...	268	52.0	fne. s.	
13933	U.S.N.M.	1	D5574...	5 30 45 N.; 120 07 57 E...	Rare.
12639	U.S.N.M.	6	D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.	Few.
13926	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gn. m.	Rare.
13929	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Rare.
12629	U.S.N.M.	10	D5630...	0 56 30 S.; 128 05 00 E...	569		co. s., m....	Common.
12637				D5637...	3 53 20 S.; 126 48 00 E...	700		gy. m.
13932	U.S.N.M.	3	D5639...	3 54 50 S.; 123 27 20 E...	1,500		gy. m.	Few.
13918	U.S.N.M.	1	D5640...	4 27 00 S.; 122 55 40 E...	24		s., brk. sh.	Rare.
13928	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Rare.
			D5656...	3 17 40 S.; 120 36 45 E...	484	41.2	gy. m.	
			D5658...	3 32 40 S.; 120 31 30 E...	510	41.2	gy. m.	Rare.
13948	U.S.N.M.	1	D5667...	2 56 00 S.; 118 47 30 E...	367	41.7	gy. s., m....	Rare.
13931	U.S.N.M.	3	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.	Few.
12630	U.S.N.M.	4	H4897...	7 46 00 N.; 122 00 00 E...	1,570		gy. m., glob.	Few.

AMMODISCUS INCERTUS d'Orbigny, var. DISCOIDEUS Cushman.

Plate 5, fig. 3.

Ammodiscus incertus, var. *discoideus* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 652.

Description.—Variety differing from the typical in having the sides rapidly increasing in diameter, the width of the coils in side view very much less than in the typical, and the border in apertural view very broadly rounded, the periphery oftentimes nearly straight in the central portion.

Diameter of microspheric specimens up to 2.5 mm.

Distribution.—Type-specimen (U.S.N.M. No. 9163) from *Albatross* station D5658 in 510 fathoms (933 meters), Gulf of Boni, bottom temperature 41.2° F. (5.1° C.). It also occurred at D5272, China Sea, vicinity southern Luzon, 118 fathoms (203 meters), bottom temperature, 57.4° F. (14.1° C.); D5582, in 890 fathoms (1,628 meters),

off Darvel Bay, Borneo, bottom temperature 38.3° F. (3.5° C.); D5585, Sibuko Bay, Borneo, 476 fathoms (871 meters), bottom temperature 41.1° F. (5° C.); D5591, in 260 fathoms (480 meters), Sibuko Bay, Borneo; D5651, in 700 fathoms (1,280 meters); and D5656, in 484 fathoms (885 meters), both in the Gulf of Boni.

The microspheric form of the variety is only about half the size of the microspheric form of the typical.

Ammodiscus incertus, var. *discoideus*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13935	U.S.N.M.	6	D5272...	° ' " ° ' "		° F.		
13936	U.S.N.M.	5	D5582...	14 00 00 N.; 120 22 30 E..	118	57.4	m., sh., co. s.	Few.
13178	U.S.N.M.	D5585...	4 19 54 N.; 118 58 33 E..	890	38.3	gy. m., fine. s.	Few.
13937	U.S.N.M.	2	D5591...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.	Rare.
13938	U.S.N.M.	2	D5591...	4 11 48 N.; 118 38 20 E..	260	Rare.
13933	U.S.N.M.	10	D5651...	4 43 50 S.; 121 23 24 E..	700	40.1	gy. m.	Common.
13939	U.S.N.M.	2	D5656...	3 17 40 S.; 120 36 45 E..	484	41.2	gy. m.	Few.
9103	U.S.N.M.	1	D5658...	3 32 40 S.; 120 31 30 E..	510	41.2	gy. m.	Few.

AMMODISCUS EXSERTUS Cushman.

Plate 5, fig. 4.

Ammodiscus exsertus CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 75, figs. 97a, b (in text).

The only station at which this species occurred is D5637, Bouro Island (south) and vicinity, 700 fathoms (1,280 meters).

Ammodiscus exsertus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13343	U.S.N.M.	1	D5637...	° ' " ° ' "		° F.		
				3 53 20 S.; 126 48 00 E..	700	gy. m.	Rare.

Family LITUOLIDAE.

Subfamily ASCHEMONELLINAE.

Genus ASCHEMONELLA H. B. Brady, 1879.

ASCHEMONELLA CATENATA (Norman).

Plate 6, fig. 5.

Astrorhiza catenata NORMAN, Proc. Roy. Soc. London, vol. 25, 1876, p. 213.

Aschemonella catenata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 42, pl. 14, figs. 12, 13; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 271, pl. 27, figs. 1-11; pl. 27A, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 81, figs. 111-113 (in text).

A single specimen here figured occurred at D5637, Bouro Island (south) and vicinity, 700 fathoms (1,280 meters). It was also

found at the neighboring station D5639, Molucca Sea, 1,560 fathoms (2,853 meters).

Aschemonella catenata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12601	} U.S.N.M.	1	D5637...	° ' " ° ' "	700	° F.	gy. m.	Rare.
12603			D5639...	3 53 20 S.; 126 48 00 E...	gy. m.	Rare.
12602	U.S.N.M.	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.	Rare.

Subfamily REOPHACINAE.

Genus REOPHAX Montfort, 1808.

REOPHAX SCORPIURUS Montfort.

Plate 6, fig. 6.

"*Orthoceras*"? SOLDANI, Testaceographica, vol. 1, 1795, p. 239, pl. 162, fig. k.
Reophax scorpiurus MONTFORT, Conch. Syst., vol. 1, 1808, p. 330, 83me genre.—
 H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 291, pl. 30, figs. 12-17.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 83, figs. 114-116 (in text).

Many specimens have been found in the material examined which may, in a general way, be referred to this species, but well defined material is not abundant. Specimens referred to this species have occurred in comparatively shallow water in various parts of the region from 24 to 510 fathoms (44 to 933 meters), with bottom temperatures, where recorded, 41.2° F. (5.1° C.), 44.3° F. (6.8° C.), 52.3° F. (10.1° C.), and 75.7° F. (24.2° C.). As a rule it seems to occur in warm and shallow water.

Reophax scorpiurus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12921	U.S.N.M.	D5096...	° ' " ° ' "	28	° F.	gy. m., s., sh.	Frequent.
D5100...			14 17 15 N.; 120 32 40 E...	35	gy. m., s., sh.	Few.	
D5133...			Id. off Panabutan Pt. N. 52° E., 1.50 miles.	38	gn. m., s.	Abundant.	
12927	U.S.N.M.	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.	Rare.
D5153...			5 18 10 N.; 120 02 55 E...	49	co. s., sh.	Few.	
D5178...			12 43 00 N.; 122 06 15 E...	78	fine s.	Abundant.	
12922	U.S.N.M.	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	brd. s.	Frequent.
14021	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Rare.
14022	U.S.N.M.	1	D5300...	° 31 00 N.; 115 49 00 E...	265	gy. m., s.	Rare.
12923	U.S.N.M.	D5318...	21 32 00 N.; 117 46 00 E...	340	s. br. Co.	Few.
12924	U.S.N.M.	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.	Rare.
D5590...			4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.	Few.	
D5640...			4 27 00 S.; 122 55 40 E...	24	s., brk. sh.	Frequent.	
12925	U.S.N.M.	D5658...	3 32 40 S.; 120 31 30 E...	510	41.2	gy. m.	Rare.

REOPHAX PILULIFER H. B. Brady.

Plate 12, fig. 1.

Reophax pilulifera H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 292, pl. 30, figs. 18-20.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 85, figs. 117, 118 (in text).

This species has occurred at 13 stations in the region, usually but few specimens occurring at any station.

The stations are all in less than 1,000 fathoms (1,829 meters), excepting one, D5637, in 1,560 fathoms (2,853 meters), while from the *Challenger* records Brady makes the statement that it "appears to be peculiar to deep water," most of the records from that material being of considerable depths.

The following localities are represented by the specimens: Sulu Sea, off southern Panay; China Sea, off Luzon; east coast of Luzon; between Siquijor and Bohol Islands; Sibuko Bay, Borneo; and between Gillolo and Makyan Islands.

After an examination of material from one of the original stations in the *Porcupine* dredgings it seems probable that many of the records for this species represent other species. The Philippine specimens are smaller than those of the *Porcupine* material and are not typical.

Reophax pilulifer—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11606	U.S.N.M.	D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m....	Rare.
15345	U.S.N.M.	3	D5127...	10 02 45 N.; 121 48 15 E..	958	50.1	gy. m., glob.	Rare.
			D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Few.
			D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.
12914	U.S.N.M.	D5301...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m., s....	Rare.
12905	U.S.N.M.	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.....	Rare.
			D5467...	13 35 27 N.; 123 37 18 E..	480	gy. m.(m.b.)	Frequent.
12916	U.S.N.M.	D5470...	13 37 30 N.; 123 41 09 E..	560	m.....	Few.
14029	U.S.N.M.	1	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn.m., glob..	Rare.
12920	U.S.N.M.	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Few.
12918	U.S.N.M.	D5622...	Makyan Id. (NE.) N. 66°W	275	gy. m.....	Rare.
12919	U.S.N.M.	D5639...	3 54 50 S.; 123 27 20 E..	1,560	gy. m.....	Rare.
12917	U.S.N.M.	D5523...	8 48 44 N.; 123 27 35 E..	Rare.

REOPHAX DISTANS H. B. Brady.

Plate 12, fig. 2.

Reophax distans H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 50; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 296, pl. 31, figs. 18-22.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 85, fig. 119 (in text).

A single specimen was found in material from D5621, between Gillolo and Makyan Islands, 298 fathoms (545 meters). At D5639, Molucca Sea, 1,560 fathoms (2,853 meters), it was abundant but most of the specimens were broken.

Owing to the weakness of the stolonlike connections between the chambers it is rare for more than two or three chambers to be found connected in dredged material. In the North Pacific material specimens were found which appeared to have the proloculum intact but the *Challenger* material showed none with the proloculum present. Single chambers with the stolonlike tubes are the most common condition.

Reophax distans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12949	U.S.N.M.	D5621...	0 15 00 N.; 127 24 35 E..	298 ° F.	gy. and bk. s.	Rare.
12948	U.S.N.M.	D5639...	3 54 50 S.; 123 27 30 E..	1,560	gy. m.....	Few.

REOPHAX BACILLARIS H. B. Brady.

Plate 12, fig. 3.

Reophax bacillaris H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 49; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 293, pl. 30, figs. 23, 24.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 86, fig. 120 (in text).

Specimens here referred to this species have been found at but four stations in the region, at one of which the specimens were in some numbers. The data for these stations show that its occurrence agrees with the rest of the Pacific records as to its being found at a considerable depth. The stations were: D5178, off Romblon, 73 fathoms (133 meters); D5460, east coast Luzon, 565 fathoms (1,074 meters); D5639, Molucca Sea, 1,560 fathoms (2,853 meters), bottom temperatures not given; and D5668, Macassar Strait, 901 fathoms (1,648 meters), bottom temperature 38.2° F. (3.4° C.).

An examination of *Valorous* material and other North Atlantic specimens has seemed to show that the typical *R. bacillaris* has a very definite distribution in the North Atlantic; also the size, form, and number of chambers, color, and in general the whole character of the species is very definite. The Philippine specimens are not typical, and it is probable that they do not represent this species. The discussion of the species is given in Bulletin 104, part 2, page 18.

Reophax bacillaris—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12947	U.S.N.M.	1	D5460...	13 32 30 N.; 123 58 06 E..	565 ° F.	gy. m.....	Rare.
8224	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	73	fine s.....	Rare.
12945	U.S.N.M.	1	D5639...	3 54 50 S.; 123 27 20 E..	1,560	gy. m.....	Rare.
12896	U.S.N.M.	8	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	Common.

REOPHAX DENTALINIFORMIS H. B. Brady.

Plate 12, fig. 4.

Reophax dentaliniformis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 49; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 293, pl. 30, figs. 21, 22.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 87, fig. 121 (in text).

Specimens from this region are not quite typical, having as a rule the chambers shorter and more rotund than in the type, but there is some variation. Most of the stations are in more than 250 fathoms (457 meters) and it is most frequent and best developed in material from those stations which have the greater depth and the lower temperatures.

Hitherto the records for this species have been almost exclusively from deep waters, but more than half of these noted here are less than 500 fathoms (914 meters).

The localities for this species include China Sea off Luzon; between Negros and Siquijor; Sibuko Bay, Borneo; Gulf of Tomini, Celebes; north of Celebes; and off Bouro Island.

Reophax dentaliniformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14028	U.S.N.M.	1	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.....	Rare.
12941	U.S.N.M.	3	D5612...	0 38 00 S.; 121 45 40 E..	750	Few.
12939	U.S.N.M.	1	D5589...	4 12 10 N.; 118 38 08 E..	260	45.7	fine. gy. s., gy. m.	Rare.
12937	U.S.N.M.	1	D5538...	9 08 15 N., 123 23 20 E..	256	53.3	gn. m., s.....	Rare.
12943	U.S.N.M.	1	D5670...	1 19 00 S.; 118 43 00 E..	1,181	38.2	gy. m.....	Rare.
12936	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Rare.
12944	U.S.N.M.	6	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	Frequent.
12938	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.....	Rare.
12942	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Rare.
12940	U.S.N.M.	3	D5612...	0 38 00 S.; 121 45 40 E..	750	Few.
14922	U.S.N.M.	1	H4897...	7 46 00 N.; 122 00 00 E..	1,570	gy. m., glob..	Rare.

REOPHAX NODULOSUS H. B. Brady.

Plate 12, fig. 5; plate 10, figs. 1, 2.

Reophax nodulosus H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 52, pl. 4, figs. 7, 8; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 294, pl. 31, figs. 1-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 87, fig. 122 (in text).

This species has occurred at several stations, usually in great numbers with sand, and showing the same range of variation as in the figures given by Brady in the *Challenger* Report. The specimens are large as a rule, and are in comparatively deep and cold water, as might be expected from the conditions of its occurrence elsewhere.

It is interesting to note, however, that at the shallowest station of this lot, D5589, in 260 fathoms (480 meters), it was very abundant and very variable, showing almost completely the various forms fig-

ured by Brady in the *Challenger* Report. At D5236 it occurred abundantly in 494 fathoms (903 meters).

Reophax nodulosus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12908	U.S.N.M.	D5460...	° ' " ° ' "	565	° F.		
12912	U.S.N.M.	D5670...	13 32 30 N.; 123 58 06 E..	gy. m.....	Rare.
12913	U.S.N.M.	D5670...	1 19 00 S.; 118 43 00 E..	1,181	38.2	gy. m.....	Few.
8225	} U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s....	Common.
12906								
12905								
12907	U.S.N.M.	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.....	Few.
12911	U.S.N.M.	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	Few.
12909	U.S.N.M.	D5589...	4 12 10 N.; 118 38 08 E..	260	45.7	fine gy. s....	Few.
12782	U.S.N.M.	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gy. m.....	Rare.

REOPHAX NODULOSUS H. B. Brady, var. HORMOSINOIDES, new variety.

Description.—Test large and coarse, chambers broad, slightly longer than the breadth, rounded, pyriform, the widest portion being about one-fourth of the way between the base and the aperture; proloculum rounded and broad, aperture large at the end of the broadly truncate chamber; color grayish-brown.

Length up to 10 mm. or more.

Type specimen.—(Cat. No. 13240, U.S.N.M.) from *Albatross* station D5650, Gulf of Boni, in 540 fathoms (988 meters). At this station it is common, but the typical form of this species does not appear.

It also occurred at D5639, Molucca Sea, in 1,560 fathoms (2,853 meters).

This variety is similar to the specimens figured by Brady.⁵

Reophax nodulosus, var. *hormosinoides*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12910	U.S.N.M.	7	D5639...	° ' " ° ' "	1,560	° F.		
11612	} U.S.N.M.	8	D5650...	3 54 50 S.; 123 27 20 E..	540	40.1	gy. m.....	Common.
13240								
				4 53 45 S.; 121 29 00 E..			gn. m.....	Common.

REOPHAX GUTTIFER H. B. Brady.

Plate 12, fig. 6.

Reophax guttifer H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 49; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 295, pl. 31, figs. 10–15.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 88, fig. 123 (in text).

Apparently from the material examined the species is rare in the region. Specimens have occurred at H4897, Sulu Sea, off western

⁵ *Challenger* Report, pl. 31, figs. 1 and 5.

Mindanao, 1,570 fathoms (2,871 meters), no bottom temperature given; and D5637, vicinity Bouro Island, 700 fathoms (1,280 meters), no bottom temperature given.

Reophax guttifer—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11609	U.S.N.M.	H4897...	° ' " ° ' "		° F.		
12951	U.S.N.M.	D5637...	7 43 45 N.; 122 03 45 E.. 3 53 20 S.; 126 48 00 E..	1,570 700	gy. m., glob. gy. m.....	Rare. Common.

REOPHAX ADUNCUS H. B. Brady.

Plate 19, fig. 3.

Reophax adunca H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 715; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 296, pl. 31, figs. 23-26.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 89, fig. 125 (in text).

In many ways this is an uncertain species to determine. Specimens from four stations may be referred to it: D5190, Tanon Strait, east coast of Negros Island, 295 fathoms (540 meters), bottom temperature 63° F. (17.2° C.); D5478, between Samar and Leyte, 57 fathoms (104 meters); D5481, the neighboring station, 61 fathoms (111 meters); and D5538, between Negros and Siquijor, 256 fathoms (468 meters), bottom temperature 53.3° F. (11.8° C.).

Reophax aduncus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12952	U.S.N.M.	D5190...	° ' " ° ' "		° F.		
12899	U.S.N.M.	D5538...	10 08 15 N.; 123 16 45 E.. 9 08 15 N.; 123 23 20 E..	295 256	63 53.3	gn. m..... gn. m., s....	Rare. Rare.
14030	U.S.N.M.	2	D5478...	10 46 24 N.; 125 16 30 E..	57	sh.....	Rare.
15346	U.S.N.M.	9	D5481...	10 27 30 N.; 125 17 10 E..	61	s., sh., g.....	Common.

REOPHAX CYLINDRICUS H. B. Brady.

Plate 13, fig. 1.

Reophax cylindrica H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 299, pl. 32, figs. 7-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 91, figs. 129-131 (in text).

At one station, D5276, China Sea, off southern Luzon, 18 fathoms (33 meters), specimens were found in considerable numbers which are apparently this species. They are slightly more constricted than shown in Brady's figures, but in all other respects are the same. The occurrence of this species, especially in such shallow water, at a low

latitude, is remarkable, as the other records are for depths of more than 1,500 fathoms (2,743 meters).

Reophax cylindricus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12950 14031	} U.S.N.M.	8	D5276...	° ' " ° ' "	18	° F.	sh., p., s....	Common.
13 49 15 N.; 120 14 45 E..							

REOPHAX PSEUDOBACILLARIS Cushman.

Reophax pseudobacillaris CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 438, figs. 2, 3.

Description.—Test elongate, arcuate, composed of a linear series of chambers; chambers shorter than broad in surface view, except the one last-formed, which appears nearly spherical, numerous, 16 to 20 or more; wall arenaceous, fairly coarse for the size of the test, slightly rough on the exterior; aperture small, in the middle of the terminal face of the chamber; color, dark reddish-brown.

Length up to 2 mm.

Specimens of this species have occurred at two stations; common at D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), 41.2° F. (5.1° C.), and rarely at D5613, Gulf of Tomini, off Celebes, 752 fathoms (1,375 meters), bottom temperature not given.

This species, while it has somewhat the appearance of *R. bacillaris* H. B. Brady, is much smaller, more arcuate, and has a dark reddish-brown color.

Reophax pseudobacillaris—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8226 12897 12898 12100	} U.S.N.M.	4	D5236...	° ' " ° ' "	494	° F.	fine gy. s....	Common.
8 50 45 N.; 126 26 52 E....				41.2				
	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E....	752	gy. m.....	Rare.

REOPHAX FINDENS (Parker).

Plate 13, fig. 4.

Lituola findens PARKER, (in Dawson's paper) Can. Nat., vol. 5, 1870, p. 177, p. 180, fig. 1.—SIDDALL, Proc. Chester Soc. Nat. Sci., pt. 2, 1878, p. 47.

Reophax findens SIDDALL, Cat. British Rec. Foram., 1879, p. 4; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 299, pl. 32, figs. 10, 11.

One specimen, which was typical in every way, was found in the Philippine material. Unfortunately the station number on the label

in the bottle had become so obliterated that it was impossible to read it. Therefore it can only be recorded as occurring in the region. The figure of this specimen is given.

Reophax findens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11608	1	° ' " ° ' "	° F.	Rare.

REOPHAX SPICULOTESTUS Cushman.

Plate 13, fig. 2; plate 11, fig. 1.

Reophax spiculotestus CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 438, fig. 4.

Description.—Test of medium size, composed of a linear series of elongate chambers, oval, in a straight or more often slightly curved line, contracted at the ends of each chamber, increasing rapidly in size toward the apertural end; perfect adult specimens with six or more chambers; wall thin, composed largely of siliceous sponge spicules, for the most part laid lengthwise of the test and firmly cemented; some sand grains present, usually neatly cemented; aperture fairly large; color, grayish.

Length up to 2 mm.

This species occurred at four stations in the region: D5210, off western Samar, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5410, between Cebu and Leyte Island, 385 fathoms (704 meters), no bottom temperature given; and D5654, Gulf of Boni, 805 fathoms (1,459 meters), bottom temperature 38.3° F. (3.4° C.).

This differs much in form and size from *R. spiculifer* H. B. Brady, as well as in the very different appearance of the test. Occurring as it does in company with an abundance of arenaceous species, it shows its selective power in the composition of its test.

Reophax spiculotestus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15348 12902 12901 8227	U.S.N.M.	1	D5201...	° ' " ° ' "	554	° F. 52.8	gy. s., m....	Rare.
12904 12903	U.S.N.M. U.S.N.M.	D5236... D5654... D5410 ..	8 50 45 N.; 126 26 52 E.... 3 42 00 S.; 120 45 50 E.... 10 28 45 N.; 124 05 30 E....	494 805 385	41.2 38.3	fine gy. s.... co. s..... gn. m.....	Common. Few. Few.

REOPHAX HORRIDUS Cushman.

Plate 13, fig. 3; plate 18, figs. 3, 4.

Reophax horrida CUSHMAN, Proc. U. S. Nat. Mus., vol. 42, 1912, p. 229, pl. 28, figs. 3, 4.

Description.—Test elongate, of several chambers, with a nearly straight axis, chambers gradually increasing in size; wall arenaceous, with an abundance of included sponge spicules so arranged as to give a bristly appearance to the exterior of the test; aperture terminal, with a well-developed neck.

Length about 3 mm.

This species occurred at D5582, vicinity of Darvel Bay, Borneo, 890 fathoms (1,628 meters), bottom temperature 38.3° F., (3.4° C.), and D5630, south of Patiente Strait, 569 fathoms (1,081 meters).

This species differs from others described in the character of the wall, the great number of included spicules bristling in all directions giving a spiny appearance to the wall.

Reophax horridus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8467	U.S.N.M.	1	D5582...	4 19 54 N.; 118 58 38 E....	890	38.3	gy. m., fine.s.	Rare.
11610	U.S.N.M.	3	D5630...	0 56 30 S.; 128 05 00 E....	569	co. s., m....	Few.

REOPHAX AGGLUTINATUS Cushman.

Plate 14, figs. 2a, b.

Reophax agglutinatus CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 637, pl. 79, fig. 6.

Description.—Test large and stout, composed of several chambers, nearly globular or slightly pyriform; wall composed almost entirely of agglutinated foraminiferal tests of *Globigerina* and *Pulvinulina*, held together by a light grayish cement; aperture with a slightly protuberant neck.

Length about 6 mm.

This species was described from *Albatross* station D5377, in 400 fathoms (732 meters), off Marinduque Island.

This is a large species and peculiar in its agglutinated test, made up almost entirely of other foraminiferal tests. In this respect it resembles *R. bilocularis* Flint, but is a typical uniserial *Reophax*.

It has also occurred at the following stations: D5201, Sogod Bay, southern Leyte, 554 fathoms (1,012 meters); D5259, off north-western Panay, 312 fathoms (571 meters); and D5582, Darvel Bay, Borneo, 890 fathoms (1,628 meters).

Reophas agglutinatus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15347	U.S.N.M.	7	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Common.
12946	U.S.N.M.	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
8512	U.S.N.M.	D5377...	13 26 00 N.; 122 19 00 E...	400	49.6	sft. gn. m....	Rare.
8467	U.S.N.M.	D5532...	4 19 54 N.; 118 58 38 E...	890	38.3	gy. m., fine.s.	Rare.

REOPHAS BILOCULARIS Flint.

Plate 12, fig. 7.

Reophas bilocularis FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 273, pl. 17, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 90, figs. 127 a, b (in text).

This species is very common in certain stations throughout the region. These stations range in depth from 208 to 976 fathoms (381 to 1,785 meters). A single specimen from D5342, Malampaya Sound, Palawan Island, in 14 to 25 fathoms (25 to 46 meters), is from the only shallow water station in which this species was found. The bottom temperature of these stations ranged from 41.1° to 53.9° F. (5° C. to 12.1° C.). The specimens from this region are more like the figures given by Flint than those which I have figured.⁶

Reophas bilocularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14023	U.S.N.M.	2	D5198...	9 40 50 N.; 123 39 45 E...	220	53.9	gn. m.	Few.
14024	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s....	Rare.
12934	U.S.N.M.	D5301...	20 37 00 N.; 115 53 00 E...	208	50.5	gy. m., s....	Few.
14025	U.S.N.M.	1	D5342...	10 56 55 N.; 119 17 24 E...	14-25	gy. m.	Rare.
14026	U.S.N.M.	4	D5424...	0 37 05 N.; 121 12 37 E...	340	50.4	co. s.	Few.
12933	U.S.N.M.	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Rare.
			D5469...	13 36 48 N.; 123 38 24 E...	500	gu. m. (net).	Rare.
12922	U.S.N.M.	D5495...	9 06 30 N.; 125 00 20 E...	976	52.3	gy. m.
12929	U.S.N.M.	D5523...	8 48 44 N.; 123 27 35 E...	Frequent.
12953	U.S.N.M.	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Few.
14027	U.S.N.M.	1	D5535...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.	Few.
12930	U.S.N.M.	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Few.
12931	} U.S.N.M.	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Few.
12932								

Genus HORMOSINA H. B. Brady, 1879.

HORMOSINA GLOBULIFERA H. B. Brady.

Plate 13, fig. 5.

Hormosina globulifera H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 60, pl. 4, figs. 4, 5; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 326, pl. 39, figs. 1-6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 93, figs. 136 a, b, 137 (in text).

Specimens of this species have occurred at a large number of stations in the region. These stations are either off the east coast of

⁶ Bull. 71, fig. 127.

Luzon or in the region south of the Equator, north of Celebes. These range in depth from 260 to 901 fathoms (475 to 1,648 meters), most of the stations, however, being over 500 fathoms (914 meters). The bottom temperatures of these stations range from 38.2° to 56.4° F. (3.4° C. to 13.5° C.), but are given in a very few of those stations at which the species occurs.

Hormosina globulifera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13907	U.S.N.M.	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Rare.
14017	U.S.N.M.	1	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.....	Rare.
14016	U.S.N.M.	1	D5449...	13 21 36 N.; 124 00 30 E..	300	Rare.
14020	U.S.N.M.	2	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.....	Rare.
14014	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E..	500	gy. m.(m.b.)	Rare.
			D5467...	13 35 27 N.; 123 37 18 E..	480	gy. m.(m.b.)	Rare.
14018	U.S.N.M.	1	D5468...	13 35 39 N.; 123 40 28 E..	569	gy. m.....	Rare.
			D5589...	4 12 10 N.; 118 38 08 E..	260	45.7	fine.gy.s.gy.m.	Rare.
13099	U.S.N.M.	D5612...	0 38 00 S.; 121 45 40 E..	750	Rare.
13096	U.S.N.M.	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Rare.
13098	U.S.N.M.	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m.....	Abundant.
13100	U.S.N.M.	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	Rare.
13101	U.S.N.M.						
14015	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E..	805	38.3	Rare.
14019	U.S.N.M.	8	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	Common.

HORMOSINA OVICULA H. B. Brady.

Plate 13, fig. 6.

Hormosina ovicula H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 61 pl. 4, fig. 6; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 327, pl. 39, fig. 79.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 95, figs. 138a, b (in text).

Single specimens which may be referable to this species occurred at two stations, D5348, Palawan Passage, 375 fathoms (686 meters), bottom temperature 56.4° F. (13.5° C.), and D5589, Sibuko Bay, Borneo, 260 fathoms (475 meters), bottom temperature 45.7° F. (7.6° C.). These are the only records for this species in the region.

Hormosina ovicula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14032	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Rare.
14033	U.S.N.M.	1	D5589...	14 12 10 N.; 118 38 08 E..	260	45.7	fine.gy.s.gy.m.	Rare.

HORMOSINA NORMANII H. B. Brady.

Plate 13, fig. 7.

Hormosina normanii H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 52; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 329, pl. 39, figs. 19–23.—CUSHMAN, Bull. 71, U. S. Nat. Mus., 1910, pt. 1, p. 95, fig. 139 (in text).

Specimens of this species which seem to be typical were found in some numbers at D5668, Macassar Strait, 901 fathoms (1,648 meters),

38.°2 F. (3.4° C.). This is the only station at which it occurred in the region.

Hormosina normanii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13093	U.S.N.M.	2	D5668...	° ' " ° ' " 2 28 15 S.; 118 49 00 E..	901	°F. 38.2	gy. m.....	Rare.

HORMOSINA OVALIFORMIS Cushman.

Plate 9, figs. 2, 3; plate 18, figs. 9, 10.

Hormosina ovaliformis CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 438, figs. 5, 6 (in text).

Hormosina monile CUSHMAN (not *H. monile* H. B. Brady), Proc. U. S. Nat. Mus., vol. 42, 1912, p. 229, pl. 28, figs. 9, 10.

Description.—Test composed of a straight or more often slightly arcuate series of chambers closely joined to one another, chambers evenly tapering at either end, or slightly oval; wall of fine sand and a reddish-brown cement, slightly roughened on the exterior, interior smooth; aperture small, rounded, at the end of the chamber, without a definite neck; color dark brick-red.

Length up to 3.5 mm.

Type specimen.—Cat. No. 8221, U.S.N.M., from *Albatross* station D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters), bottom temperature 41.2°F. (5.1°C.).

It has occurred elsewhere in the region at the following stations: D5539, between Negros and Siquijor, depth and bottom temperature not recorded, D5622, Makyan Island, 275 fathoms (503 meters), and D5656, Gulf of Boni, 484 fathoms (885 meters), bottom temperature 41.2° F. (5.1° C.).

Hormosina ovaliformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8221	U.S.N.M.	1	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E....	494	°F. 41.2	fne. gy. s....	Few.
8468	U.S.N.M.	1	D5539...	9 03 20 N.; 123 24 25 E....	Few.
14034	U.S.N.M.	1	D5622...	Makyan Id. (NE), N. 66° W	275	gy. m.....	Rare.
14035	U.S.N.M.	1	D5656...	3 17 40 S.; 120 36 45 E....	484	41.2	gy. m.....	Rare.

HORMOSINA ELONGATA Cushman.

Hormosina elongata CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 439, figs. 7 a, b.

Description.—Test elongate, nearly straight, composed of a series of elongate chambers; largest diameter near the base of each chamber,

thence gradually narrowing toward the apertural end; wall rather thick, composed of fine material with much cement, smoothly finished; aperture rather small, elliptical; color reddish-brown, lighter about the aperture.

Length 5 to 10 mm.

Several specimens of this species were found at D5236, off the east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5° 1 C.).

This species is much more elongate in regard to its entirety and in its individual chambers than other species of the genus. Its nearest related species seems to be *Hormosina carpenteri* H. B. Brady, but the curvature of the test and the shape of the chambers is very different.

Hormosina elongata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8263	U.S.N.M.	2	D5236...	8° 50' 45" N.; 126° 26' 52" E....	494	° F. 41.2	fine. gy. s....	Rare.

HORMOSINA CARPENTERI H. B. Brady.

Plate 15, fig. 5.

"*Moniliform Lituola*" CARPENTER, The Microscope, ed. 5, 1875, p. 531, fig. *f*.

Hormosina carpenteri H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 51.—

CARPENTER, The Microscope, ed. 6, 1881, p. 563, fig. *f*.—H. B. BRADY, Rep.

Voy. Challenger, Zoology, vol. 9, 1884, p. 327, pl. 39, figs. 14–18.

Description.—Test elongate, composed of several chambers arranged in an irregular line, pyriform, base rounded, tapering toward the apertural end, increasing slightly in size toward the apertural end of the test; wall composed of fine arenaceous material, often made hispid by included sponge spicules.

Length up to 12.5 mm. or more.

Distribution.—The only material I have had of this species was rather characteristic, from *Albatross* D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.).

Hormosina carpenteri—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15349	U.S.N.M.	1	D5259...	11° 57' 30" N.; 121° 42' 15" E....	312	° F. 49.3	gy. m., glob..	Rare.

Genus HAPLOSTICHE Reuss, 1861.

HAPLOSTICHE DUBIA (d'Orbigny).

Plate 13, figs. 8, 9.

"*Orthoceratia Zoophytica minuscula*" SOLDANI, Testaceographica, vol. 1, pt. 2, 1791, p. 93, pl. 98, fig. A.

Nodosaria dubia D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 252, No. 10.

Lituola dubia PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 4, vol. 8, 1871, p. 263, pl. 9, fig. 30.

Lituola soldanii JONES and PARKER, Quart. Journ. Geol. Soc., vol. 16, 1860, p. 307, no. 184.

Haplostiche soldanii H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 318, pl. 32, figs. 12-18.

Haplostiche dubia CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 96, figs. 140, 141 (in text).

This species is rather common in many parts of the region. Both microspheric and megalospheric forms of the species occurred usually at the same station. The microspheric form attains a much larger size, as is usual. The aperture in the larger specimens becomes decidedly complex, the teeth assuming a dendritic form. The smaller megalospheric specimens usually have a much simpler form of apertural teeth. In the development of a specimen there is a change from a simple tooth in the earlier chambers through gradual changes to the complex aperture of the adult.

In depth the stations range from 37 to 540 fathoms (67 to 987 meters), and bottom temperatures from 40.1° to 59.0° F. (3.5° to 15° C.), but most of the temperatures are above 50° F. (10° C.).

Haplostiche dubia—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature. °F.	Character of bottom.	Abundance.
14008	U.S.N.M.	5	D5110...	14 03 45 N.; 120 16 30 E...	135	59.0	dk. gy. m...	Few.
14007	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Few.
13160	U.S.N.M.	6	D5261...	12 30 55 N.; 121 34 24 E...	145	s., m.....	Common.
13159	U.S.N.M.	1						
14006	U.S.N.M.	2	D5272...	14 00 00 N.; 120 22 30 E...	118	57.4	m., sh., co. s.	Rare.
13161	U.S.N.M.	4	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.....	Few.
14012	U.S.N.M.	3	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Frequent.
13163	U.S.N.M.	1	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gn. m., co. s.	Few.
13165	U.S.N.M.	1						
14009	U.S.N.M.	10	D5523...	8 48 44 N.; 123 27 35 E...	Common.
14011	U.S.N.M.	2	D5565...	5 51 42 N.; 120 30 30 E...	243	52.3	s., ptr., sh....	Rare.
13162	U.S.N.M.	2	D5567...	5 48 00 N.; 120 33 45 E...	268	52.0	fine. s.....	Few.
13164	U.S.N.M.	1						
			D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.....	Rare.
14010	U.S.N.M.	1	D5642...	4 31 40 S.; 122 49 42 E...	37	gy. m.....	Rare.
14013	U.S.N.M.	3	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Rare.
			D5667...	2 56 00 S.; 118 47 30 E...	367	41.7	gy. s., m....	Abundant.

Subfamily TROCHAMMININAE.

Genus TROCHAMMINOIDES Cushman, 1910.

TROCHAMMINOIDES PROTEUS (Karrer).

Trochammina proteus KARRER, Sitz. kais. Akad. Wiss. Wien, vol. 52, (Abth. 1), 1865 (1866), p. 494, pl., fig. 8 (not 1-7).—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 341, pl. 40, figs. 1-3.

Trochamminoides proteus CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 98, figs. 142-144 (in text).

There is a single specimen which has less than the usual number of chambers in the last-formed whorl. Otherwise it seems typical in the character of its wall and general appearance.

Trochamminoides proteus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13154	U.S.N.M.	1	D5236...	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Rare.

Genus HAPLOPHRAGMOIDES Cushman, 1910.

HAPLOPHRAGMOIDES CANARIENSIS (d'Orbigny).

Nonionina canariensis d'ORBIGNY, in Barker, Webb, and Berthelot, Hist. Nat. Îles Canaries, vol. 2, pt. 2, 1839, "Foraminifères," p. 128, pl. 2, figs. 33, 34.

Placopsilina canariensis PARKER and JONES, Ann. Mag. Nat. Hist., ser. 2, vol. 19, 1857, p. 301, pl. 10, figs. 13, 14.

Lituola canariensis W. B. CARPENTER, PARKER, and JONES, Intr. Study Foram., 1862, pl. 6, figs. 39-41.

Lituola nautiloidea, var. *canariensis* PARKER and JONES (part), Trans. Roy. Soc. London, vol. 155, 1865, p. 406, pl. 15, figs. 45a, b; pl. 17, figs. 92-95.

Haplophragmium canariensis SIDDALL, Cat. Brit. Rec. Foram., 1879, p. 4.—H. B. BRADY, Denkschr. kais. Akad. Wiss. Wien, vol. 42, 1881, p. 99; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 310, pl. 35, figs. 1-5.

Haplophragmoides canariensis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 101, fig. 149 (in text).

This species occurred rarely in the region, being noted at but two stations: D5210, off western Samar, in 50 fathoms (91 meters), bottom temperature 76.3° F. (24.6° C.); and D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.). The specimens were not typical.

Haplophragmoides canariensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13081	U.S.N.M.	1	D5236...	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Rare.
13087	U.S.N.M.	1	D5210...	11 49 50 N.; 124 28 05 E...	50	76.3	fne. gy. s....	Rare.
13083	U.S.N.M.	1	D5210...	11 49 50 N.; 124 28 05 E...	50	76.3	fne. gy. s....	Rare.

HAPLOPHRAGMOIDES EMACIATUM (H. B. Brady).

Haplophragmium emaciatum H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 305, pl. 33, figs. 26-28.

Haplophragmium compressum MILLETT (not *Haplophragmium compressum* Goës), Journ. Roy. Micr. Soc., 1889, p. 359, pl. 5, fig. 8.

Haplophragmoides emaciatum CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 102, figs. 150-152 (in text).

This species occurred in the material from but four stations. These were all in deep water in the Sulu Sea, in bottom material of Globigerina-ooze off Celebes, and Bouro Island. The stations are: D5127, Sulu Sea, vicinity southern Panay, 958 fathoms (1,752 meters), bottom temperature 50.1° F. (10° C.); H4897, Sulu Sea, off western Mindanao, 1,570 fathoms (2,871 meters), no bottom temperature given; D5601, Gulf of Tomini, Celebes, 765 fathoms (1,399 meters), bottom temperature not recorded; and D5637, Bouro Island, 700 fathoms (1,280 meters).

Haplophragmoides emaciatum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13080	U.S.N.M.	1	D5127...	° ' " ° ' "	958	° F. 50.1	gy. m., glob..	Rare.
13029	U.S.N.M.	1	D5601...	10 02 45 N.; 121 48 15 E..	765	s., glob., ptr.	Rare.
14062	U.S.N.M.	1	D5637...	1 13 10 N.; 125 17 05 E..	700	gy. m.	Rare.
8216 4755	U.S.N.M.	H4897...	3 53 20 S.; 126 48 00 E..	1,570	gy. m., glob..	Few.
				7 46 00 N.; 122 00 00 E..				

HAPLOPHRAGMOIDES SCITULUM (H. B. Brady).

Haplophragmium scitulum H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 50; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 308, pl. 34, figs. 11-13.

Haplophragmoides scitulum CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 103, figs. 153-155 (in text).

Typical specimens of this species were found in material from two stations: D5175, Sulu Sea, southeast of Cagayanes Islands, depth and bottom temperature not given; and D5467, east coast Luzon, 480 fathoms (878 meters), bottom temperature not given.

Haplophragmoides scitulum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13086	U.S.N.M.	1	D5175...	° ' " ° ' "		° F.		
			D5467...	9 21 00 N.; 121 37 45 E..	480	gy. m. (m. b.)	Rare.
				13 35 27 N.; 123 37 18 E..				

HAPLOPHRAGMOIDES SUBGLOBOSUM (G. O. Sars).

Plate 15, figs. 1a, b.

Lituola subglobosa M. Sars, Forh. Vid. Selsk. Christiania, 1868 (1869), p. 250 (*nomen nudum*).—G. O. Sars, Forh. Vid. Selsk. Christiania, 1871, (1872), p. 253.

Haplophragmium subglobosum H. B. Brady, Denkschr. kais. Akad. Wiss. Wien, vol. 43, 1881, p. 100.

Haplophragmium latidorsatum H. B. Brady, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 307, pl. 34, figs. 7, 8, 10, 14 (?) (not fig. 9), and subsequent authors (not *Nonionina latidorsata* Bornemann, 1855).

Haplophragmoides subglobosum Cushman, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 105, figs. 162-164 (in text).

This is one of the commonest species of the genus in the region. Specimens are usually numerous at the stations where found and are typical, with slight variation in color and texture of the test according to the material of the bottom on which it occurred. Nearly all of the stations are in comparatively deep water. The depths vary from 80 to 1,560 fathoms (146 to 2,853 meters), nearly all of them being over 300 fathoms (549 meters). Bottom temperatures are recorded in but four of the many stations—44.0°, 44.3°, and 58.6° F. (6.4°, 6.8°, and 14.7° C.). These four stations were 310, 347, 383, and 80 fathoms (567, 635, 700, and 146 meters) respectively in depth. The localities include the following: East coast of Mindoro, China Sea off Luzon and off Formosa; east coast of Luzon; Sibuko Bay, Borneo; Patiente Strait; and Molucca Sea.

Haplophragmoides subglobosum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12765	U.S.N.M.	10+	D5121...	13 27 20 N.; 121 17 45 E...	108	° F.	dk. gn. m...	Common.
12766	U.S.N.M.	2	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine. s.	Rare.
12859	U.S.N.M.	3	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co.	Few.
			D5382...	13 15 20 N.; 122 45 30 E...	128	m.	Rare.
			D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.	Rare.
			D5446...	12 43 51 N.; 124 59 18 E...	300	gn. m.	Few.
12767	U.S.N.M.	10+	D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m. (m.b.)	Frequent.
			D5468...	13 35 39 N.; 123 40 28 E...	569	gn. m.	Rare.
			D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.	Common.
14063	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gy. m., s.	Rare.
12858	U.S.N.M.	1	D5591...	4 11 48 N.; 118 38 20 E...	260	Rare.
13077	U.S.N.M.	2	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.	Few.
13074	U.S.N.M.	6	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.	Frequent.

HAPLOPHRAGMOIDES RINGENS (H. B. Brady).

Plate 15, fig. 2.

Trochammina ringens H. B. Brady, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 57, pl. 5, figs. 12a, b; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 343, pl. 40, figs. 17, 18.

Ammochilostoma ringens EIMER and FICKERT, Zeitschr. wiss. Zool., vol. 65, 1889, p. 692.

Haplophragmoides ringens CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 107, fig. 166 (in text).

At one station, D5613, off Celebes, 752 fathoms (1,375 meters), bottom temperature unrecorded, numerous typical and very fine specimens of this species occurred, but it was not found in material elsewhere in the region.

Haplophragmoides ringens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13084	U.S.N.M.	8	D5613...	° ' " ° ' " 0 42 00 S.; 121 44 00 E..	752	° F.	gy. m.	Frequent.

HAPLOPHRAGMOIDES GRANDIFORMIS Cushman.

Plate 11, fig. 2.

Haplophragmoides grandiformis CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 440, fig. 11.

Description.—Test free, involute, planospiral; chambers low and broad, seven to nine in the last-formed coil, inflated; peripheral line of the test in side view considerably depressed at the sutures; wall composed of rather coarse sand grains with a considerable proportion of yellowish-brown cement, somewhat rough on the exterior; aperture an elongate, curved slit at the base of the apertural face of the chamber; color light brown.

Diameter about 3 mm.

Distribution.—Type specimen from *Albatross* station 5152, near the Tawi Tawi group, 34 fathoms (62 meters).

This is a large species somewhat resembling *H. canariensis*, but involute with broad low chambers.

It has occurred elsewhere in the region at D5277, China Sea, vicinity of southern Luzon, 80 fathoms (147 meters), and D5369, Molucca Sea, 1,560 fathoms (2,853 meters). Bottom temperatures were not recorded in any cases.

Haplophragmoides grandiformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8217	U.S.N.M.	1	D5152...	° ' " ° ' " 5 22 55 N.; 120 15 45 E..	34	° F.	wh. s.	Rare.
1406	U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E..	80	Rare.
14060	U.S.N.M.	1	D5369...	3 54 50 S.; 123 27 20 E..	1,560	gy. m.	Rare.

HAPLOPHRAGMOIDES SPHAERILOCULUM Cushman.

Plate 15, fig. 3.

Haplophragmoides sphaeriloculum CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 652.

Description.—Test biconvex, composed of about three coils, chambers comparatively few, five in each coil, subspherical, sutures deeply depressed; wall composed of fine sand grains with a reddish-brown cement, the sutures and umbilical depression more or less filled with a light-colored fine amorphous material; aperture a narrow slit at the base of the chamber.

Diameter about 2 mm.

Distribution.—Type specimen (U.S.N.M. No. 9104) from *Albatross* station D5637, in Pitt Passage, in 700 fathoms (1,280 meters).

This species is marked by the very convex, inflated, almost spherical chambers, with a fine grained, yellowish-brown wall.

Sidebottom has recorded this species from the east coast of Australia, in 465 fathoms (846 meters).⁷

Haplophragmoides sphaeriloculum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12764 9104 15350	U.S.N.M. U.S.N.M. U.S.N.M.	5 1	D5637...	° ' " ° ' " 3 53 20 S.; 126 48 00 E..	700	° F.	gy. m.	Few.

Genus CRIBROSTOMOIDES Cushman, 1910.

CRIBROSTOMOIDES BRADYI Cushman.

Plate 17, figs. 1a, b.

Haplophragmium latidorsatum H. B. BRADY (part) (not Bornemann), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 307, pl. 34, fig. 9.

Cribrostomoides bradyi CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 108, figs. 167, a, b (in text).

This species occurs at numerous stations and in considerable numbers, especially in the region south of the Equator, where so many arenaceous forms occur. Apparently both microspheric and megalospheric forms of the species occur here; the smaller meglospheric specimens do not show the cribriform aperture as well as the larger microspheric specimens. Also in the development of a single specimen, the aperture in the earlier chambers is simple and gradually assumes the cribriform structure of the adult. In these characters it agrees with what has already been noted in *Haplostiche dubia*. Almost all the stations at which the species occurred are in com-

⁷ Journ. Roy. Micr. Soc., 1918, p. 15, pl. 2, figs. 15, 16.

paratively deep water, between 700 and 900 fathoms (1,280 and 1,646 meters). At three stations specimens were found in 298, 312, and 476 fathoms (545, 571, and 871 meters). These are the only four stations less than 700 fathoms. The bottom temperatures were given in but four cases—38.2°, 38.3°, 41.1°, and 49.3° F. (3.4° C., 3.5° C., 5° C., and 9.6° C.). This last was from a station with a depth of but 312 fathoms (571 meters).

The localities represented are: Sibuko Bay, Borneo; north of Celebes; between Gillolo and Makyan Islands; off Bouro Island; Gulf of Boni; and Flores Sea.

Cribrostomoides bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.	
12658	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E ..	312	49.3	gy. m., glob.	Rare.	
			D5585...	4 07 00 N.; 118 49 54 E ..	476	41.1	gy. m.....	Few.	
12655	U.S.N.M.	2	D5612...	0 38 00 S.; 121 45 40 E ..	750	Few.	
12653	U.S.N.M.	4	D5613...	0 42 00 S.; 121 44 00 E ..	752	gy. m.....	Frequent.	
12657	U.S.N.M.	10+							
15351	U.S.N.M.	1	D5621...	0 15 00 S.; 127 24 35 E ..	298	gy. and bk. s. (m. b.).	Rare.	
12656	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E ..	700	gy. m.....	Few.	
			D5654...	3 42 00 S.; 120 45 50 E ..	805	38.3	Abundant.
			D5668...	2 28 15 S.; 118 49 00 E ..	901	38.2	gy. m.....	Common.

Genus CYCLAMMINA H. B. Brady, 1876.

CYCLAMMINA CANCELATA H. B. Brady.

Plate 16, figs. 1a, b.

"Nautiloid *Lituola*" W. B. CARPENTER, *The Microscope*, ed. 5, 1875, p. 536, figs. 274a-c (in text).

Cyclammina cancellata H. B. BRADY (MS.) in Norman, *Proc. Roy. Soc. London*, vol. 25, 1876, p. 214; *Rep. Voy. Challenger*, *Zoology*, vol. 9, 1884, p. 351, pl. 37, figs. 8-16.—CUSHMAN, *Bull.* 71, U. S. Nat. Mus., pt. 1, 1910, p. 110, figs. 168-170 (in text) (not fig. 171).

Description.—Test nautiloid, typically biconvex, but occasionally slightly more convex on one side than the other; chambers numerous, averaging about 13 chambers in the last-formed coil, ranging from 12 to 15; usually three full coils in the adult test; peripheral edge of test broadly rounded, approximately semicircular; sutures distinct but only slightly depressed, chambers little if at all inflated, curved anteriorly in a nearly even curve in side view; wall smooth and polished when perfect, when eroded, showing the perforate, cancellate appearance due to the exposure of the labyrinthic interior; aperture an elongate, curved slit at the base of the apertural face, supplemented by a clustered series of rounded pores occupying the central part of the apertural face, often becoming very numerous in large specimens; color reddish- or yellowish-brown, some of the individual chambers occasionally black.

Diameter up to 6.5 mm. in the microspheric form, averaging in the commoner megalospheric form 3 to 4 mm.

Distribution.—This species was found very abundantly at some stations: East coast of Mindanao; east coast of Luzon; China Sea; Jolo Sea; Molucca Passage; Patiente Strait; and Gulf of Boni. These stations range in depth from 208 to 940 fathoms (381 to 1,719 meters), the average being 560 fathoms (1,065 meters). Bottom temperatures range from 36.7° F. to 53° F. (2.6° C. to 11.7° C.), the average being 46.3° F. (7.8° C.)

This species may be characterized by the broadly rounded peripheral margin, the number of chambers averaging 13 to 15. The rotund tumid character of the test as a whole will distinguish it at once from any of the following. Both microspheric and megalospheric forms occur in this species and are similar except in size.

Cyclammina cancellata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8214	U.S.N.M.	3	D5236...	° ' " ° ' "	494	° F.	fne. gy. s....	Common.
12824	U.S.N.M.	10+		8 50 45 N.; 126 26 52 E..		41.2		
12825	U.S.N.M.	10+						
14036	U.S.N.M.	3	D5301...	20 37 00 N.; 115 53 00 E..	208	50.5	gy. m., s....	Few.
14040	U.S.N.M.	2	D5423...	9 38 30 N.; 121 11 00 E..	508	49.8	gy. m., co. s.	Few.
14041	U.S.N.M.	6	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.	Frequent.
14037	U.S.N.M.	5	D5425...	9 37 45 N.; 121 11 00 E..	495	49.4	gy. m., co. s.	Frequent.
14038	U.S.N.M.	1	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.	Few.
12869	U.S.N.M.	1	D5460...	13 32 30 N.; 123 58 06 E..	565	gn. m.	Few.
14042	U.S.N.M.	7	D5468...	13 35 39 N.; 123 40 28 E..	509	gn. m. (m. b.).	Frequent.
12868	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m. (net)..	Few.
14044	U.S.N.M.	6	D5470...	13 37 30 N.; 123 41 09 E..	560	m.	Frequent.
14039	U.S.N.M.	2	D5619...	0 35 00 N.; 127 14 40 E..	435	fne. gy. s., m.	Few.
12870	U.S.N.M.	4	D5529...	9 23 45 N.; 123 39 30 E..	441	53	gy. m., glob.	Few.
13440	U.S.N.M.	10	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m....	Frequent.
14043	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.	Few.

CYCLAMMINA COMPRESSA Cushman.

Plate 16, figs. 2a, b.

Cyclammina cancellata (part) CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 111, fig. 171 (not figs. 168-170) (in text).

Cyclammina compressa CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 653.

Description.—Test nautiloid, biconvex, compressed peripheral margin subacute, umbilicus often notably excavated; chambers numerous, 14 to 16 in the last-formed coil, usually 15; sutures subangular in the middle in side view, clearly depressed, surface smooth when perfect; aperture and color as in *C. cancellata*.

Diameter up to 3.5 mm.

Distribution.—Type specimen (U.S.N.M. No. 9105) from *Albatross* station D5470, in 560 fathoms (1,065 meters) off the east coast of Luzon. It was noted at numerous stations in the same locality, one

station off the west coast of Luzon; Darvel Bay and Sibuko Bay, Borneo; north of Celebes; Patiente Strait; Gulf of Boni, Flores and Macassar Strait. These stations range in depth from 208 to 1,092 fathoms (381 to 1,998 meters), the average depth being 601 fathoms (1,099 meters). Bottom temperatures range from 36.3° to 50.5° F. (2.3° C. to 10.2° C.) with the average 40.3° F. (4.6° C.).

This species seems to be in colder and deeper water than the preceding, from which it may be distinguished by its compressed form, larger number of chambers, depressed umbilical area, and subangular, somewhat depressed sutures.

Cyclammina compressa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14049	U.S.N.M.	10	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s....	Common.
14050	U.S.N.M.	2	D5439...	15 58 15 N.; 119 40 20 E...	940	36.7	gn. m.....	Few.
13250	U.S.N.M.	4	D5460...	13 32 30 N.; 123 58 06 E...	565	gy. m.....	Few.
13244	U.S.N.M.	1	D5468...	13 35 39 N.; 123 40 28 E...	569	gn. m.(m. b.)	Common.
12876	U.S.N.M.	10+	D5467...	13 35 27 N.; 123 37 18 E...	480	gn. m.(m. b.)	Common.
13246	U.S.N.M.	6	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.(m. b.)	Common.
13245	U.S.N.M.	10+	D5470...	13 37 30 N.; 123 41 09 E...	560	m.....	Frequent.
14051	U.S.N.M.	5	D5582...	4 19 54 N.; 118 58 38 E...	890	38.3	gy. m., fine. s.	Few.
14052	U.S.N.M.	7	D5591...	4 11 48 N.; 118 38 20 E...	260	Frequent.
13252	U.S.N.M.	2	D5609...	0 11 00 S.; 121 16 00 E...	1,092	36.3	gn. m.....	Few.
13249	U.S.N.M.	3	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.....	Few.
13248	U.S.N.M.	2	D5621...	0 15 00 N.; 127 24 35 E...	298	gy. and bk.s.	Few.
12875	U.S.N.M.	10	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.....	Fragment.
13243	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Rare.
14053	U.S.N.M.	2	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Few.
13251	U.S.N.M.	5	D5658...	3 17 40 S.; 120 36 45 E...	484	41.2	gy. m.....	Few.
13247	U.S.N.M.	4	D5668...	3 32 40 S.; 120 31 30 E...	510	41.2	gy. m.....	Few.
13247	U.S.N.M.	4	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.....	Rare.

CYCLAMMINA PAUCILOCLATA Cushman.

Plate 16, figs. 3 a, b.

Cyclammina pauciloculata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 653.

Description.—Test compressed, nautiloid, biconvex, peripheral margin subacute, umbilical region depressed, chambers 10 to 11 in number, sutures nearly straight to somewhat curved; surface smooth when perfect; supplementary apertural pores few in number.

Diameter up to 2 mm.

Distribution.—Type specimen (U.S.N.M. No. 9106) from *Albatross* station D5538, in 256 fathoms (468 meters), between Negros and Siquijor, bottom temperature 53.3° F. (11.8° C.). This species was abundant at this station, and was also found in the China Sea off southern Luzon; east coast of Luzon; off northern Mindanao; Gulf of Tomini, Celebes; and Gulf of Boni. These stations range in depth from 102 to 1,092 fathoms (187 to 1,998 meters), the average depth 530 fathoms (970 meters); bottom temperatures range from 36.3° to 59.6° F. (2.3° C. to 15.3° C.), the average being 44.0° F. (6.4° C.).

Both the bottom temperatures, which are much higher, and the depths, which are much less than the preceding species, show that it has a different habitat.

It may be most easily distinguished from *C. cancellata* by its compressed form, depressed umbilical region, and from both *C. cancellata* and *C. compressa* by its fewer chambers.

Cyclammina pauciloculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14045	U.S.N.M.	1	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fine.s., m., sh.	Rare.
13241	U.S.N.M.	1	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Rare.
14046	U.S.N.M.	1	D5447...	13 28 00 N.; 123 46 18 E...	310	45.3	gn. m.	Rare.
14047	U.S.N.M.	6	D5523...	8 48 44 N.; 123 27 35 E...	Common.
9106	U.S.N.M.	10	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s....	Common.
14048	U.S.N.M.	1	D5609...	0 11 00 S.; 121 16 00 E...	1,092	36.3	gn. m.	Rare.
13242	U.S.N.M.	10+	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.	Common.
12872	U.S.N.M.	4	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.	Few.
12871	U.S.N.M.	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Few.

CYCLAMMINA PUSILLA H. B. Brady.

Plate 16, figs. 4a, b.

Cyclammina pusilla H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 53; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 353, pl. 37, figs. 20-23.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 111, fig. 172 (in text).

This small species was found at five *Albatross* stations, as follows: D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.); D5247, Gulf of Davao, 135 fathoms (247 meters); D5590, Sibuko Bay, Borneo, 310 fathoms (567 meters), bottom temperature 44.3° F (6.8° C.); D5621, between Gillolo and Makyan Islands, 298 fathoms (545 meters); and D5659, Gulf of Boni, 702 fathoms (1,284 meters), bottom temperature 39.0° F. (4.1° C.).

It may be distinguished from all the preceding species by its much smaller size, its diameter being 1 mm. or less.

Cyclammina pusilla—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
14054	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E....	494	41.2	fine. gy. s....	Rare.
12823	U.S.N.M.	1	D5247...	7 02 00 N.; 125 38 45 E....	135	m.	Rare.
12874	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E....	310	44.3	gn. m., s....	Rare.
12873	U.S.N.M.	1	D5621...	0 15 00 N.; 127 24 35 E....	298	gy. and bk. s	Rare.
12822	U.S.N.M.	1	D5659...	5 33 20 S.; 120 47 10 E....	702	39.0	s., m.	Rare.

CYCLAMMINA BRADYI Cushman.

Trochammina trullissata H. B. BRADY (part), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 342, pl. 40, fig. 13 (not 14, 15).

Cyclammina bradyi CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 113, figs. 174a, b (in text).

The only record for this species in the whole region is a single specimen at *Albatross* station D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

Cyclammina bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12821	U.S.N.M.	1	D5236...	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Rare.

Genus LITUOTUBA Rhumbler, 1895.

LITUOTUBA LITUIFORMIS (H. B. Brady).

Trochammina lituiformis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 59, pl. 5, fig. 16; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 342, pl. 40, figs. 4-7.

Lituotuba lituiformis RHUMBLER, Nachr. kön. Ges. Wiss. Göttingen, 1895, p. 84.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 114, fig. 175 (in text).

There are two specimens from station D5236, Pacific Ocean, east coast of Mindanao, which are evidently this species. The division of the chambers is obscure. It consists of four coiled whorls and a short uncoiled portion. The earlier portion is not in one plane, but is more or less irregular, as is often the case in this species. The wall is composed of fine sand grains with numerous black specks and a light yellowish cement.

Lituotoba lituiformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13158 13331	U.S.N.M. U.S.N.M.	1 1	}D5236...	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Rare.

Genus AMMOBACULITES Cushman, 1910.

AMMOBACULITES AGGLUTINANS (d'Orbigny).

Plate 17, fig. 4.

Spirolina agglutinans D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 137, pl. 7, figs. 10-12.

Haplophragmium agglutinans H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 301, pl. 32, figs. 19-26.

Haplophragmium calcareum FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 275, pl. 19, fig. 1 (not *H. calcareum* d'Orbigny).

Ammobaculites agglutinans CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 115, fig. 176 (in text).

Throughout the region this species seems to occur at numerous stations, but never in great numbers, as do some of the other species of the genus. There is some variation in the color and material of which the test is made, according to bottom conditions. The stations are numerous, as noted, mostly in comparatively shallow and warm water. They range in depth from 14 to 750 fathoms (25 to 1,370 meters), most of the stations, however, being in less than 300 fathoms (549 meters), bottom temperatures, where recorded, ranging from 41.1° to 54.8° F. (5° C. to 12.6° C.), most of them being near 50° F. (10° C.) or more, the lower temperatures occurring at the deeper stations.

Most of the stations for this species are in the Archipelago proper, but a few of the records are from Sibuko Bay, Borneo, and north of Celebes.

Ammobaculites agglutinans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12457	U.S.N.M.	1	D5110...	° ' " ° ' "	135	39.0	dk. gy. m....	Few.
			D5214...	13 59 20 N.: 120 75 45 E..	218	51.4	gn. m.....	Rare.
			D5259...	12 25 18 N.: 123 37 15 E..	312	49.3	gy. m., glob.	Frequent.
			D5261...	11 57 30 N.: 121 42 15 E..	145		s. m.....	Frequent.
			D5282...	12 30 55 N.: 121 34 24 E..	248	47.4	dk. gy. s....	Few.
			D5342...	13 53 00 N.: 120 26 45 E..	14-25		gy. m.....	Several.
12458	U.S.N.M.	1	D5343...	10 56 55 N.: 119 17 24 E..			s., r.....	Rare.
			D5412...	10 51 35 N.: 119 23 24 E..	162	54.8	gn. m.....	Rare.
			D5438...	10 09 15 N.: 123 52 00 E..	267	46.2	gn. m.....	Rare.
12455	U.S.N.M.	1	D5469...	15 54 42 N.: 119 44 42 E..	500		gn.m.(net)...	Rare.
			D5470...	13 36 48 N.: 123 38 24 E..	560		m.....	Rare.
14071	U.S.N.M.	3	D5529...	13 37 40 N.: 123 41 09 E..	441	53.0	gy. m., glob.	Few.
12456	U.S.N.M.	1	D5537...	9 23 45 N.: 123 39 30 E..	254	53.5	gn. m.....	Few.
12459	U.S.N.M.	2	D5585...	4 07 00 N.: 118 49 54 E..	476	41.1	gy. m.....	Rare.
			D5586...	4 06 50 N.: 118 47 20 E..	347	44.0	gy. m.....	Rare.
12460	U.S.N.M.	1	D5589...	4 12 10 N.: 118 38 08 E..	260	45.7	fine gy.s., gy. m.	Rare.
			D5612...	0 38 00 S.: 121 45 40 E..	750			Rare.

AMMOBACULITES TENUIMARGO (H. B. Brady).

Haplophragmium tenuimargo H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 715; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 303, pl. 33, figs. 13-16.

Ammobaculites tenuimargo CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 117, figs. 180-183 (in text).

Four specimens of this species of rather smaller size than may be typical, but otherwise of the form and character of the specimens shown in the figures by Brady, occurred at D5420, between Cebu and Bohol Islands, 127 fathoms (233 meters); bottom temperature, 59° F. (15° C.).

Ammobaculites tenuimargo—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12480	U.S.N.M.	4	D5420...	9 49 35 N.; 123 45 00 E..	127	° F. 59.0	gn. m.....	Rare.

AMMOBACULITES CALCAREUM (H. B. Brady).

Plate 17, fig. 3.

Haplophragmium calcareum H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 302, pl. 33, figs. 5-12.

As might be expected from the records given in the *Challenger* Report, which were all but one from shallow water of the tropics, this species is common in the material from this region. It has occurred at a large number of stations and usually in considerable numbers. Specimens exceed in size those obtained by Brady, some of them measuring over 4 mm. in length. Apparently both microspheric and megalospheric forms of the species occur. The material of the test varies according to bottom conditions. It is always of fairly coarse material, but smoothly cemented, the proportion of colored cement being unusually large, as noted by Brady.

Most of the stations are in the Philippines proper, but a few are in the more southern region, in the Gulf of Boni, etc. In depth the stations range from 50 to 565 fathoms (91 to 1,074 meters), but three stations being less than 200 fathoms (366 meters), and few over 400 (732 meters). The range of depth compares rather well with that given from *Challenger* stations, at which this species occurred. Bottom temperatures range from 40.1° to 55.4° F. (3.5° C. to 15.2° C.), and at the shallowest station, 50 fathoms (91 meters), the bottom temperature was 76.3° F. (24.6° C.).

AMMOBACULITES CYLINDRICUS Cushman.

Plate 17, fig. 5.

Haplophragmium calcareum (part), H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, pl. 23, fig. 6.

Haplophragmium dyglutinans FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 275, pl. 19, fig. 2 (not *Spirolina agglutinans* d'Orbigny).

Ammobaculites cylindricus CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 441, figs. 15, 16.

Description.—Test free, elongate, cylindrical; early chambers planospiral, completely involute, with five to six chambers in each volution; later portion uncoiled, cylindrical, made up of a linear series of chambers; wall coarsely arenaceous, but the surface rather smoothly finished; aperture in the middle of the terminal face in the uncoiled portion simple; color, gray.

Length 2 mm.; diameter 0.5 to 0.75 mm.

This species has been found to occur at numerous stations throughout the region, ranging in depth from 18 to 554 fathoms (33 to 1,012 meters), most of the stations being between 200 and 400 fathoms (366 and 732 meters), bottom temperatures ranging from 40.1° to 53.3° F. (3.5° C. to 11.8° C.).

This species differs from the ordinary form of *A. calcareus* in its cylindrical shape, and from *A. agglutinans* in the greater proportion of coiled chambers, which are completely involute.

Ammobaculites cylindricus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.		Character of bottom.	Abundance.
						° F.			
12473	U.S.N.M.	3	D5120...	13 45 30 N.; 120 30 15 E...	393	43.7	gn. m., s....	Few.	
			D5156...	5 12 50 N.; 119 55 55 E...	18	fine s., sh....	Abundant.	
8205	U.S.N.M.	4	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Frequent.	
12472	U.S.N.M.	1	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m.....	Few.	
12477	U.S.N.M.	2							
12471	U.S.N.M.	5	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Frequent.	
14069	U.S.N.M.	1	D5260...	12 25 35 N.; 121 31 35 E...	234	51.4	gn. m., s....	Rare.	
			D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s....	Rare.	
12474	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.....	Rare.	
12475	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E...	360	Rare.	
			D5569...	5 23 15 N.; 120 15 30 E...	303	52.3	co. s.....	Few.	
			D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.....	Rare.	
12476	U.S.N.M.	5	D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fine gy. s., gy. m.	Few.	
			D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.....	Frequent.	
12478	U.S.N.M.	3	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Few.	
14070	U.S.N.M.	1							
12479	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E...	221	gy. m., glob.	Rare.	

AMMOBACULITES REOPHACIFORMIS Cushman.

Plate 11, fig. 3; plate 14, fig. 3.

Ammobaculites reophaciformis CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 440, figs. 12–14.

Description.—Test free, elongate-fusiform; early portion consisting of chambers arranged in a close-coiled planospiral series, much compressed, later portion straight, *Reophax*-like, consisting of a linear

series of chambers progressively increasing in size; wall coarsely arenaceous, rough, fairly thick, aperture circular, in the middle of the terminal face, often on a projecting, neck-like portion; color variable, usually white or gray.

Length up to 3.5 mm.; diameter 0.4 to 1 mm.

This species has proved to be one of the most common in the region. It is characteristic of the shallow water of the coral-reef region. It has occurred at a large number of stations ranging in depth from 14 to 318 fathoms (25 to 582 meters). Two stations at a greater depth were 258 and 318 fathoms (472 and 582 meters). In one case a single specimen occurred and in the other few. The great majority of the stations are in depths from 15 to 30 fathoms (27 to 55 meters). Bottom temperatures are not usually given at these shallow water stations, but where given they range from 75.7° to 76.3° F. (24.2° to 24.6° C.).

The early coiled portion separates the species from *Reophax*, but the later portion alone would be described as belonging to that genus. At first I took it to be the microspheric form of some species of *Reophax* but it is very abundant, and when perfect seems always to have the coiled early development. *Reophax scorpiurus*, which it in some ways resembles, is a common species in the same locality, but in the material examined was not seen to have a true close-coiled young in any case, although the young portion is often curved.

Ammobaculites reophaciformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14065	U.S.N.M.	3	D5096...	14 20 23 N.; 120 34 15 E...	28	° F.	gy. m., s., sh.	Abundant.
			D5097...	14 19 15 N.; 120 33 52 E...	30		gy. m., s., sh.	Abundant.
			D5100...	14 17 15 N.; 120 32 40 E...	35		gy. s.	Frequent.
12447	U.S.N.M.	3	D5105...	14 43 55 N.; 120 12 50 E...	25		Co.	Common.
			D5106...	14 23 55 N.; 120 32 33 E...	37		gy. m.	Common.
			D5134...	6 44 45 N.; 121 48 00 E...	25	(?)	fne. s.	Abundant.
			D5136...	6 04 20 N.; 120 59 20 E...	22		s., sh.	Rare.
			D5153...	5 18 10 N.; 120 02 55 E...	49		co. s., sh.	
8207	U.S.N.M.	4	D5156...	5 12 50 N.; 119 55 55 E...	18		fne. s., sh.	Abundant.
12453	U.S.N.M.	10+	D5161...	5 10 15 N.; 119 53 00 E...	16		fne. s., blk.sp	Rare.
12454	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318		inc. s., sh.	Few.
11637	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78		fne. s.	Frequent.
12452	U.S.N.M.	9	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.	Few.
15340	U.S.N.M.	1	D5181...	11 36 40 N.; 123 26 35 E...	26		m., fne. s.	Common.
			D5182...	11 30 40 N.; 123 23 20 E...	24		fne., s. m.	Rare.
			D5191...	10 29 45 N.; 123 31 15 E...	258	62.8	gn. m.	Rare.
12448	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32		gn. s.	Frequent.
12449	U.S.N.M.	5	D5206...	11 31 40 N.; 124 42 40 E...	32		gn. m.	Few.
14066	U.S.N.M.	2	D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fne. gy. s.	Frequent.
12451	U.S.N.M.	4	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.	Few.
			D5220...	13 38 00 N.; 121 58 00 E...	50		sft. gn. m.	Several.
12450	U.S.N.M.	1	D5256...	7 21 45 N.; 124 07 15 E...	158		m.	Rare.
			D5276...	13 49 15 N.; 120 14 45 E...	18		sh., p., s.	Frequent.
12444	U.S.N.M.	2	D5338...	11 33 45 N.; 119 24 45 E...	43		co. s., m.	Few.
			D5342...	10 56 55 N.; 119 17 24 E...	14-25		gy. m.	Few.
12445	U.S.N.M.	1	D5426...	9 12 00 N.; 118 23 00 E...	27		fne. gy. s.	Several.
12883	U.S.N.M.	5	D5478...	10 46 24 N.; 125 16 30 E...	57		sh.	Few.
15341	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co.	Rare.
14068	U.S.N.M.	1	D5642...	4 31 40 S.; 122 49 42 E...	37		gy. m.	Few.
12446	U.S.N.M.	5	D5642...	4 31 40 S.; 122 49 42 E...	162		gy. m.	Few.
14067	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E...	195	52.8	gn. m.	Rare.

AMMOBACULITES PSEUDOSPIRALE (Williamson).

Plate 19, figs. 1, 2.

Proteonina pseudospirale WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 2, pl. 1, figs. 2, 3.

Haplophragmium pseudospirale SIDDALL, Cat. British Recent Foram., 1879, p. 4.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 302, pl. 33, figs. 1-4.—HERON-ALLEN and EARLAND, Proc. Roy. Irish Acad., vol. 31, pt. 64, 1913, p. 45.

There are numerous specimens in the Philippine material which seem to be very close to this species, which is known from the North Atlantic. It has the general shape and character of the wall which characterizes British specimens, the only difference being in the more definite division of the chambers. The early part is coiled and the later part uncoiled, which should place it in the genus *Ammobaculites*. Many of the specimens seem somewhat like *A. calcareum*. It seems, however, distinct from that species at the stations at which it occurs. The material of the wall is either a dark material, probably some sort of sand or light calcareous material.

There are records for this species at 18 stations, ranging in depth from 88 to 554 fathoms (161 to 1,012 meters), the average 278 fathoms (508 meters), bottom temperatures range from 43° to 62.4° F. (6.1° to 16.8° C.), the average about 50° F. (10° C.). These stations include the following regions: Off Jolo; Verde Island Passage; Tanon Strait, east coast of Negros; off western Bohol; Sogod Bay, southern Leyte; east of Masbate; between Marinduque and Luzon; Pacific Ocean, east coast of Mindanao; off southeastern Mindoro; China Sea off southern Luzon; Ragay Gulf, Luzon; off northern Mindanao; between Negros and Siquijor; and to the south in Sibuko Bay, Borneo, and between Gillolo and Makyan Islands.

Ammobaculites pseudospirale—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12468	U.S.N.M.	3	D5120...	13 45 30 N.; 120 30 15 E...	393	43.7	gn. m., s.	Few.
12465	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318	fine s., sh.	Rare.
12469	U.S.N.M.	1	D5187...	9 16 45 N.; 123 21 15 E...	225	53.6	sft. gn. m.	Rare.
12462	U.S.N.M.	3	D5198...	9 40 50 N.; 123 39 45 E...	220	53.9	gn. m.	Few.
12466	U.S.N.M.	2 (10+)	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.	Frequent.
12467								
13312								
12461	U.S.N.M.	4	D5214...	12 25 18 N.; 123 37 15 E...	218	50.5	gn. m.	Few.
12463	U.S.N.M.	4	D5222...	13 38 30 N.; 121 42 45 E...	195	52.8	gn. m.	Few.
12464	U.S.N.M.	1	D5237...	8 09 06 N.; 126 31 45 E...	249	46.4	gn. m.	Rare.
12470	U.S.N.M.	5	D5260...	12 25 35 N.; 121 31 35 E...	312	49.3	gy. m., glob.	Few.
13313	U.S.N.M.	10+	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.	Common.
14756 8206	U.S.N.M.	10+	H4898...	7 43 45 N.; 122 03 45 E...	221	gy. m., glob.	Common.
13315								
13315	U.S.N.M.	10+	D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.	Common.
13314	U.S.N.M.	2	D5385...	13 24 50 N.; 123 03 70 E...	327	62.4	gy. m.	Few.
13316	U.S.N.M.	4	D5523...	8 48 44 N.; 123 27 35 E...	Few.
13317	U.S.N.M.	4	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Few.
13318	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.	Rare.
13319	U.S.N.M.	3	D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.	Few.
13320	U.S.N.M.	1	D5622...	Makyan Id. (NE.), N. 66° W.	275	gy. m.	Rare.

Genus *PLACOPSILINA* d'Orbigny, 1850.*PLACOPSILINA CENOMANA* d'Orbigny.

- Placopsilina cenomana* d'ORBIGNY, Prodr. Pal., vol. 2, 1850, p. 185, no. 758.—
H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 315, pl. 36, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 119, fig. 186 (in text).
Lituola cenomana JONES and PARKER, Quart. Journ. Geol. Soc., vol. 16, 1860, p. 302.
Lituola (Placopsilina) cenomana W. B. CARPENTER, PARKER, and JONES, Intr. Foram., 1862, p. 143, pl. 11, figs. 11-14.

This species has been found several times attached to various genera of foraminifera or to shells or pebbles. It is most frequent in shallow water, but the depths have ranged from 18 to 494 fathoms (33 to 903 meters). The only stations at which the bottom temperature is given are D5236, 494 fathoms (903 meters), 41.2° F. (5.1° C.), D5217, 105 fathoms (193 meters), 63.1° F. (17.2° C.), and D5538, 256 fathoms (468 meters), 53.3° F. (11.8° C.).

Placopsilina cenomana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12591	U.S.N.M.	3	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Few.
12586	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318	fne. s., sh...	Rare.
12587	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Common.
12590								
12593								
12588								
13262 8222	U.S.N.M.	8	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	ers. gy. s....	Common.
12589	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s. . .	Rare.
12592	U.S.N.M.	1	D5276...	13 49 50 N.; 120 14 45 E...	18	sh., p., s.....	Rare.
15669	U.S.N.M.	7	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s.....	Frequent.

Genus *TROCHAMMINA* Parker and Jones, 1860.*TROCHAMMINA SQUAMATA* Jones and Parker.

Plate 17, fig. 2; plate 22, fig. 6.

Trochammina squamata JONES and PARKER, Quart. Journ. Geol. Soc., vol. 16, 1860, p. 304.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 337, pl. 41, figs. 3a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 120, figs. 187a, b (in text).

Trochammina proteus KARRER (part), Sitz. kais. Akad. Wiss. Wien, vol. 52, 1865, p. 494, pl. 1, fig. 6 (not 1-5, 7, 8).

A few specimens of this species were found at the following stations: D5219, between Marinduque and Luzon, 530 fathoms (969 meters), bottom temperature 50.8° F. (10.4° C.); D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5670, Macassar Strait, 1,181 fathoms (2,160 meters), bottom temperature 38.2° F. (3.4° C.), and H4897, Sulu Sea off western Mindanao, 1,570 fathoms (2,871 meters), no bottom temperature recorded.

Trochammina squamata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12865	U.S.N.M.	(?)	D5219...	13 21 00 N.; 122 18 45 E...	530	50.8	gn. m.	
12866	U.S.N.M.	1	D5236...	8 45 50 N.; 126 26 52 E...	494	41.2	fne gy. s.	Rare.
12867	U.S.N.M.	1	D5670...	1 19 00 S.; 118 43 00 E...	1,181	38.2	gy. m.	Rare.
14752	U.S.N.M.	1	H4897...	7 46 00 N.; 122 00 00 E...	1,570	gy. m., glob.	Rare.

TROCHAMMINA TURBINATA (H. B. Brady).

Plate 15, fig. 4.

Haplophragmium turbinatum H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 50; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 312, pl. 35, figs. 9 a-c.

Trochammina turbinatum EIMER and FICKERT, Zeitschr. wiss. Zool., vol. 65, 1899, p. 695.

Trochammina turbinata CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 122, fig. 189 (in text).

The only records from the region are two specimens of this species, from *Albatross* station D5439, west coast of Luzon, in 940 fathoms (1,719 meters) and one specimen from D5637, Pitt Passage, in 700 fathoms (1,280 meters). These specimens are larger than the ordinary, and have the general form of the species, but the aperture, instead of being at the base of the apertural face, is small, semi-circular, and in the middle of the face.

Trochammina turbinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13321	U.S.N.M.	2	D5439...	15 58 15 N.; 119 40 20 E...	940	36.7	gn. m.	Rare.
14055	U.S.N.M.	1	D5637...	3 53.20 S.; 126 48 00 E...	700	gy. m.	Rare.

TROCHAMMINA GLOBIGERINIFORMIS (Parker and Jones).

Plate 11, figs. 4, 5.

Globigerina bulloides WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 56, pl. 5, figs. 116-118 (not *G. bulloides* d'Orbigny, 1826).

Lituola nautiloidea, var. *globigeriniformis* PARKER and JONES, Trans. Roy. Soc. London, vol. 155, 1865, p. 407, pl. 15, figs. 46, 47.

Lituola (*Haplophragmium*) *globigeriniformis* TERRIGI, Nuovi Lincei Atti, 1880, p. 175, pl. 1, fig. 3.

Haplophragmium globeriniforme SIDDALL, Cat. Brit. Rec. Foram., 1879, p. 4.—W. B. CARPENTER, The Microscope, ed. 6, 1881, p. 561, figs. 320 a, b.

Ammoglobigerina bulloides EIMER and FICKERT, Zeitschr. wiss. Zool., vol. 65, 1899, p. 704.

Trochammina globigeriniformis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 124, figs. 193, 194 (in text).

Fine large specimens of this species are found at a great number of stations in the region. Although not definitely proven from the two sizes that appear, it would seem that both microspheric and megalospheric specimens of this species may be present.

In depth the stations range from 260 to 1,570 fathoms (475 to 2,871 meters), bottom temperatures from 36.3° to 49.4° F. (2.3° to 9.7° C.).

From the known conditions under which this species is best developed it is not strange to find few records in the Philippines proper, but most of them from the deeper water to the south. It was not found at all among the islands, but off the Pacific coast of Mindanao and China Sea off Luzon. Southward it was found in Sibuko Bay, Borneo; north of Celebes; between Gillolo and Makyan Islands; Patiente Strait; and Flores Sea.

Trochammina globigeriniformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8235	} U.S.N.M.	4	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Frequent.
12851								
12857	U.S.N.M.	3	D5238 ..	7 34 45 N.; 126 38 15 E..	380	43.0	gn. m.	Few.
			D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.
14056	U.S.N.M.	1	D5425...	9 37 45 N.; 121 11 00 E..	495	49.4	gy. m., co. s.	Rare.
12855	U.S.N.M.	1	D5446...	12 43 51 N.; 124 59 18 E..	300	Rare.
12852	U.S.N.M.	2	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.	Rare.
			D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.	Few.
12853	U.S.N.M.	5	D5591...	4 11 48 N.; 118 38 20 E..	260	Few.
			D5609...	0 11 00 S.; 121 16 00 E..	1,092	36.3	gn. m.	Few.
12854	U.S.N.M.	10	D5513...	0 42 00 S.; 121 44 00 E..	752	gy. m.	Common.
12856	U.S.N.M.	2	D5621...	0 15 00 N.; 127 24 35 E..	298	gy. and bk.	Few.
							s. (m. b.).	
14957	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m.	Few.
14055	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.	Rare.
14059	U.S.N.M.	2	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.	Few.
			D5670...	1 19 00 S.; 118 43 00 E..	1,181	38.2	gn. m.	Few.
14753	U.S.N.M.	1	H4897...	7 46 00 N.; 122 00 00 E..	1,570	gy. m., glob.	Rare.

Genus **AMMOCHILOSTOMA** Eimer and Fickert, 1899.

AMMOCHILOSTOMA ROTUNDATA Cushman.

Ammochilostoma rotundata CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 637, pl. 79, fig. 1.

Description.—Test nearly spherical or somewhat elongate, of few chambers, the last-formed chamber nearly covering all the previously formed ones; chambers arranged in an irregular coil; wall fairly thick, of fine sand grains with a grayish-brown cement; aperture narrow and elongate, near the middle of the terminal face of the last-formed chamber, but remote from the border of the preceding chamber.

Diameter about 1 mm.

This species was found in considerable numbers at one station, D5613, off Celebes, 752 fathoms (1,375 meters); bottom temperature not given. It has also occurred in fewer numbers at D5612, Gulf of Tomini, Celebes, 750 fathoms (1,370 meters); D5654, Gulf of Boni, 805 fathoms (1,472 meters), bottom temperature 38.3° F. (3.5° C.); and D5668, Macassar Strait, 901 fathoms (1,648 meters), bottom temperature, 38.2° F. (3.4° C.).

Ammochilostoma rotundata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12594	U.S.N.M.	10+	D5612...	0 38 00 S.; 121 45 40 E....	750	° F.		Common.
12596								
8513								
11636	U.S.N.M.	10+	D5613...	0 42 00 S.; 121 44 00 E....	752		gy. m.....	Common.
12597								
12595	U.S.N.M.	4	D5654...	3 42 00 S.; 120 45 50 E....	805	38.3		Few.
12598								
12599	U.S.N.M.	2	D5668...	2 28 15 S.; 118 49 00 E....	901	38.2	gy. m.....	Few.

Genus AMMOSPHEROIDINA Cushman, 1910.

AMMOSPHEROIDINA SPHEROIDINIFORMIS (H. B. Brady).

Haplophragmium sphaeroidiniformis H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 313.

Ammospheroidina sphaeroidiniformis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 1, 1910, p. 128, fig. 202 (in text).

Specimens referable to this species were found at four stations: D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.); D5348, Palawan Passage, 375 fathoms (686 meters); D5467, east coast of Luzon, 480 fathoms (878 meters); and D5621, Mokuca Passage, 298 fathoms (545 meters).

Ammospheroidina sphaeroidiniformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13189	U.S.N.M.	6	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob..	Few.
13188								
	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	Co., s.....	Rare.
13187	U.S.N.M.	2	D5621...	0 15 00 N.; 127 24 35 E..	298		gy. m..... gy. and bk.s.	Rare.

AMMOSPHEROIDINA GRANDIS Cushman.

Plate 9, fig. 4; plate 16, fig. 5.

Ammospheroidina grandis CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 442, figs. 17-19 (in text).

Description.—Test large, globular, in adults usually made up of three visible chambers, one large one and two smaller ones on the opposite side; wall coarsely arenaceous, but with a fairly smooth exterior; aperture at the base of the largest chamber crescentic or semicircular, nearly opposite the suture marking the juncture of the walls of the two smaller chambers; color, grayish-brown.

Diameter up to 3 mm.

This species is very much more common than the preceding, being found at a number of stations in the area. These range in depth from 106 to 494 fathoms (194 to 903 meters), and bottom temperatures 41.2° to 50.4° F. (5.1° to 10.1° C.).

The question may be raised as to whether or not this may be the microspheric form of the preceding species, but from its occurrence in far greater numbers and at such a large number of stations it would seem not to justify this idea, being the opposite of the usual occurrence of the microspheric form of this species.

All the records for this species are in the Philippine Archipelago, none of the records being from the region to the southward.

Ammosphaeroidina grandis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8209	} U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s....	Abundant.
13179					312	49.3	gy. m., glob..	Rare.
13180	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E..	106	bk. s.	Rare.
15342	U.S.N.M.	1	D5382...	13 15 20 N.; 122 45 30 E..	128	m.	Rare.
13184	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.	Rare.
13182	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.	Rare.
13183	U.S.N.M.	1	D5446...	12 43 51 N.; 124 59 18 E..	300	Rare.
13186	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E..	Rare.
13185	U.S.N.M.	1	D5656...	3 17 40 S.; 120 36 45 E..	484	41.2	gy. m.	Rare.
14072	U.S.N.M.	2	D5591...	4 11 48 N.; 118 38 20 E..	260	gn. m., s....	Few.

Genus SPHAERAMMINA Cushman, 1910.

Sphaerammina CUSHMAN, Proc. U. S. Nat. Mus., vol. 33, 1910, p. 439.

Description.—Test composed of a series of chambers, the last one formed completely enveloping the preceding ones, but the axis straight; wall arenaceous.

Type of the genus.—*Sphaerammina ovalis* Cushman.

This genus strongly reminds one of *Ellipsoidina*, but there is an arenaceous wall, and the connections between the apertural ends of the chambers are indistinct or wanting. With its straight axis it belongs to the *Reophaeinae*, and holds a relation to *Reophae* and *Hormosina* similar to that which *Glandulina* holds to *Nodosaria*.

SPHAERAMMINA OVALIS Cushman.

Plate 11, figs. 6-8; plate 14, figs. 5a, b.

Sphaerammina ovalis CUSHMAN, Proc. U. S. Nat. Mus., vol. 38, 1910, p. 440, figs. 8-10.

Description.—Test oval or spherical, composed of a series of chambers, with a straight axis, each chamber as added extending back and enveloping the preceding; chambers oval or nearly spherical; wall of fine sand firmly cemented; aperture elliptical or rounded; color, grayish- or yellowish-brown.

Diameter 1 to 2 mm.

Specimens were common at the type station, D5236, Pacific Ocean, off eastern Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5621, between Gillolo and Makyan Island, 298 fathoms (545 meters); and also specimens which were typical in every way from D5637, near Bouro Island, 700 fathoms (1,280 meters), bottom temperatures not recorded in the last two cases.

Sphaerammina ovalis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.		
				° ' " ° ' "		° F.				
8233	U.S.N.M.	6	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s.	Few.		
14819	U.S.N.M.	3								
14809	U.S.N.M.	3				D5621...	0 15 00 N.; 127 24 35 E..	298	gy. and bk. s.	Few.
			D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.				

Genus AMMOSPHAERULINA Cushman, 1912.

AMMOSPHAERULINA ADHAERENS Cushman.

Plate 18, figs. 11, 12.

Ammosphaerulina adhaerens CUSHMAN, Proc. U. S. Nat. Mus., vol. 42, 1912, p. 229, pl. 28, figs. 11, 12.

Description.—Test spherical, adherent, wall arenaceous; composed of two or more chambers, each included by the one next-formed, eccentric; color light yellowish-brown.

Diameter about 0.75 mm.

Specimens of this species were found at *Albatross* station D5637, in the vicinity of Bouro Island, Dutch East Indies, 700 fathoms (1,280 meters), bottom temperature not recorded.

This is a peculiar form, somewhat like *Psammospaera*, but having chambers entirely included by the one last-formed. On account of its small size and attached character it may easily be overlooked. The figured specimen was attached to a specimen of *Saccorhiza ramosa*.

Ammosphaerulina adhaerens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8466	U.S.N.M.	1	D5637...	3° 53' 20" S.; 126° 48' 00" E..	700	° F.	gy. m.	Rare.

Genus *NOURIA* Heron-Allen and Earland, 1914.*NOURIA HARRISII* Heron-Allen and Esrland.

Nouria harrisii HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, pt. 12, 1914, p. 376, pl. 37, figs. 16–20.

The following is the original description of this species:

Test free, fusiform, cylindrical to subcylindrical in section, consisting of two to four chambers arranged in a polymorphine manner and constructed entirely of sponge spicules arranged in a single layer with their axes arranged more or less parallel to the long axis of the test. The ultimate chamber terminating in a somewhat produced neck, on which is situated the aperture of the test, which is a circular opening. The lines of demarcation between the separate chambers are marked by very slight sutural depressions, but are principally recognizable owing to a divergence of the angles at which the spicules of adjacent chambers are arranged (fig. 18). The septation of the internal cavity is very incomplete, but the different successive chambers are recognizable in section by these sutural depressions, which extend as an incomplete septum into the cavity of the test. It follows that each of these incomplete septa must originally have formed a portion of the external wall of the test which has been absorbed in the course of growth, the spicules, no doubt, being rebuilt into the new investing wall.

We first noticed this form more than 20 years ago in dredgings made by the late Captain Seabrook, and given to us by our old friend and correspondent, the late Mr. W. H. Harris, of Cardiff. The dredgings in which the specimens occurred were made off Cebu in the Philippine Islands (depth 120 fathoms, volcanic mud) and in the Java Sea (45 fathoms). Identical specimens have since been observed in a dredging from the Sahul Bank, in the Timor Sea (60 fathoms), and in a dredging off Old Providence Island in the Caribbean Sea (382 fathoms). The peculiar and highly characteristic construction of the test is the same in all these widely separated localities; and, as sponge spicules do not form any marked proportion of the bottom in any of the dredgings, the "selective" tendency must be regarded as exceptionally pronounced—almost as striking, in fact, as in our species *Technitella thompsoni*,⁸ which form a heterogeneous mass of available material, selects for the construction of its test nothing but the plates of an Ophiurid.

The skill, or "purpose," exhibited by this little organism in the building of its test reaches its most remarkable development in the construction of the aperture. The spicules designed to form the terminal portion of the shell are selected by the organism of such size and shape as to form a perfectly tapered neck with a circular aperture, round which the points of the spicules often form a regular fringe (figs. 17, 18, 20). It would appear that the organism utilizes only such spicules as are suitable for its purposes, owing to their tapering form, terminating in a sharp point, which allows the size of the aperture to be considerably reduced, as compared with the size which would have resulted from the utilization of broken or blunted spicules. The

⁸ Journ. Quekett Micr. Club, ser. 2, vol. 10, 1909, pp. 403–412, pls. 31–35.

spicules employed are invariably of a simple type, and this is very noticeable in the Cebu specimens, as hexactinellid spicules abound in that dredging.

Heron-Allen and Earland, in describing this species, had material from off Cebu; I have failed to find any specimens in the Philippine material I have had.

NOURIA COMPRESSA Heron-Allen and Earland.

Nouria compressa HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, pt. 12, 1914, p. 378, pl. 37, figs. 21-26.

The original description of this species is as follows:

Test free, highly compressed, consisting of 2 to 5 chambers rapidly increasing in size and arranged biserially, the walls of the test consisting of a single layer of sponge-spicules neatly cemented together into a smooth and finished surface. A grain of sand or glauconite is occasionally used to fill in a crevice between spicules (figs. 22 & 25). Marginal edges acute and usually smooth and unbroken, especially at the oral extremity of the shell. Less neat at the aboral end, where there is often a marginal fringe of projecting spicular points. Sutural lines nearly flush and obscure, but generally well marked, owing to the divergent angles at which the spicules are arranged in adjacent chambers. Aperture a neatly constructed terminal slit, sometimes with a slightly raised border or lip of cement (figs. 21, 23). Internal septa constructed of spicules and very often incomplete.

This species, described by Heron-Allen and Earland from the same station off Cebu, in 120 fathoms (220 meters), did not occur in any of the Philippine material I have had, but I have had very typical specimens from off the coast of Japan.

Family TEXTULARIIDAE.

Subfamily SPIROPLECTINAE.

Genus SPIROPLECTA Ehrenberg, 1844.

SPIROPLECTA BULBOSA Cushman.

Plate 20, fig. 1.

Spiroplecta bulbosa CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 5, figs. 1a, b (in text).

This species, originally described from the coast of Japan, has occurred at a single station, D5420, between Cebu and Bohol, 127 fathoms (233 meters); bottom temperature, 59.0° F. (15° C.).

The Philippine specimen seems to be typical.

Spiroplecta bulbosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14392	U.S.N.M.	1	D5420...	° ' " ° ' " 9 49 35 N.; 123 45 100 E..	127	° F. 59	Rare.

Subfamily TEXTULARIINAE.

Genus TEXTULARIA DeFrance, 1824.

TEXTULARIA SAGITTULA DeFrance.

Plate 20, figs. 2-4.

Polymorphum sagittula SOLDANI, Testaceographica, vol. 1, pt. 2, 1791, pp. 120-133, fig. T.

Textularia sagittula DEFANCE, Dict. Sci. Nat., vol. 32, 1824, p. 177, vol. 53, p. 344; Atlas, Conch., pl. 13, fig. 5.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 361, pl. 42, figs. 17, 18.

Typical material of this species was found at the following localities: Sulu Archipelago, Tawi Tawi Group; vicinity of Jolo; China Sea off southern Luzon; off northern Cebu; off western Samar; Verde Island Passage; between Negros and Siquijor; Mindoro Strait; east coast of Luzon; and vicinity of Darvel Bay, Borneo.

Some of the specimens measured as much as 4 mm. in length. Two interesting specimens are figured, plate 20, figs. 2-4, which had evidently been broken and started a new growth. In one case the axis of the new portion of the test is in the same line, but has been twisted about so that the chambers of the later part are at right angles to their former position. In the second the new portion of the test arises from the side of the earlier portion.

Such specimens show clearly that the test of the foraminifera is really an internal one, and that under certain conditions new growth may take place from any part.

Textularia sagittula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
12065	U.S.N.M.	6	D5096...	14 20 23 N.; 120 34 15 E...	28	gy. m., s., sh.	Few.
			D5110...	13 50 20 N.; 120 75 45 E...	135	59.0	dk. gy. m...	Common.
			D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Rare.
12064	U.S.N.M.	1	D5154...	5 14 50 N.; 119 58 45 E...	12	co. s.....	Rare.
12062	U.S.N.M.	3	D5192...	11 09 15 N.; 123 50 00 E...	71	gn. m.....	Few.
12061	U.S.N.M.	2	D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fine. gy. s...	Rare.
12063	U.S.N.M.	6	D5268...	13 42 00 N.; 120 57 15 E...	170	s. p.....	Frequent.
			D5333...	12 26 30 N.; 120 37 45 E...	310	73.8	s.....	Rare.
			D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Rare.
			D5538...	9 08 15 N.; 123 23 20 E...	257	53.3	gn. m., s.....	Rare.
			D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co...	Few.
12066	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.....	Rare.

TEXTULARIA SAGITTULA DeFrance, var. ATRATA Cushman.

Plate 20, fig. 5.

Textularia sagittula DEFANCE, var. *atrata* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 7, figs. 2-5 (in text).

This variety has occurred at the following localities in few numbers: Tara Island; Sulu Sea; off northern Cebu; off southern Luzon; China

Sea, off Hongkong and off Formosa; off northern Mindanao; Palawan Passage; and Gulf of Tomini, Celebes.

The peculiar dark condition of the material above the sutures was noted in all the specimens.

Specimens from *Albatross* station D5315 show excellent senescent characters. The last-formed chambers of the test show a lateral narrowing so that the width of this portion of the test is reduced to one-half that of its maximum development.

The type material of this variety was from *Albatross* station D4875, in the eastern channel of the Korean Straits, in 59 fathoms (107 meters).

Textularia sagittula, var. *atrata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12059	U.S.N.M.	2	D5133...	Id. off Panabutan Pt., N. 52 E.; 150 miles.	38	° F.	gn. m., s.....	Common.
12057	U.S.N.M.	1	D5142...	6 06 10 N.; 121 02 40 E...	21	co. s., sh.....	Rare.
			D5192...	11 09 15 N.; 123 50 00 E...	32	gn. m.....	Rare.
12055	} U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine. s.....	Rare.
12056								
12054	U.S.N.M.	3	D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh.....	Few.
12058	U.S.N.M.	5	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Common.
			D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co., s.....	Rare.
			D5523...	8 48 44 N.; 123 27 35 E...	Rare.
12100	U.S.N.M.	1	D5565...	5 51 42 N.; 120 30 30 E...	243	52.3	s., ptr., sh...	
12053	U.S.N.M.	3	D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.....	Few.
12051	U.S.N.M.	2	D5601...	1 13 10 N.; 125 17 05 E...	765	s., glob., ptr.	Rare.
12052	U.S.N.M.	2	Tara Island.....	

TEXTULARIA SAGITTULA DeFrance, var. FISTULOSA H. B. Brady.

Plate 20, fig. 6.

Textularia sagittula DEFRANCE, var. *fistulosa* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 362, pl. 42, figs. 19-22.

Brady mentions that this variety is principally found under tropical and subtropical conditions. It has occurred in the Philippine Expedition dredgings from two stations in the China Sea, off southern Luzon, D5105, 25 fathoms (46 meters), and D5300, in 265 fathoms (485 meters).

Textularia sagittula, var. *fistulosa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12060	U.S.N.M.	1	D5300...	0 ' ' ' 0 ' ' ' 20 31 00 N.; 115 49 00 E...	265	° F.	gy. m., s.....	Rare.
			D5105...	14 43 55 N.; 120 12 50 E...	25	

TEXTULARIA GRAMEN d'Orbigny.

Plate 20, fig. 7.

Textularia gramen D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 248, pl. 15, figs. 4, 6.—
H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 365, pl. 43, figs. 9,
10.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 8, figs. 6-8 (in text).

This is one of the most common species of the genus in the region and has been recorded from 28 stations, usually occurring in considerable numbers. As a rule it occurs in shallow water, but at two stations, D5636, Pitt Passage, it occurs in 1,262 fathoms (2,309 meters), and D5637, near Bouro Island, in 700 fathoms (1,280 meters) and 805 fathoms (1,472 meters). In each case a single specimen was found. The other stations, with a few exceptions, are mostly less than 50 fathoms (91 meters), the range being from 10 to 500 fathoms (18 to 914 meters). The range of temperature is from 38.3° to 75.7° F. (3.5° C. to 24.2° C.), the lower temperature corresponding with the deeper water.

Textularia gramen—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12123	U.S.N.M.	1	D5096...	14 20 23 N.; 120 34 15 E...	28	° F.	gy. m., s., sh.	Rare.
14177	U.S.N.M.	2	D5100...	14 17 15 N.; 120 32 40 E...	35		gy. s.	
12125	U.S.N.M.	3	D5110...	13 59 20 N.; 120 75 45 E...	135	59	dk. gy. m.	Few.
14187	U.S.N.M.	1	D5113...	13 51 30 N.; 120 50 30 E...	159		dk. gn. m.	Rare.
14181	U.S.N.M.	7	D5132...	Id. off Panabutan Pt. N. 15 W., 0.3 mile.	26		gn. m., s.	Few.
			D5142...	6 06 10 N.; 121 02 40 E...	21		co. s., sh.	Rare.
			D5143...	6 05 50 N.; 121 02 15 E...	19		co. s.	Rare.
14179	U.S.N.M.	4	D5159...	5 11 50 N.; 119 54 00 E...	10		co. s.	Few.
14186	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.	Rare.
14184	U.S.N.M.	2	D5267...	11 09 15 N.; 123 50 00 E...	32		gn. m.	
			D5213...	12 15 00 N.; 123 57 30 E...	80		s., m., sh.	Rare.
			D5218...	13 11 15 N.; 123 02 45 E...	20		crs. s.	
12121	U.S.N.M.	3	D5318...	21 32 00 N.; 117 46 00 E...	340		s., br. Co.	Few.
			D5267...	13 42 20 N.; 120 58 25 E...	170		p., s., sh.	
			D5333...	12 26 30 N.; 120 37 45 E...	310	73.6	s.	Rare.
14180	U.S.N.M.	4	D5381...	13 14 15 N.; 122 44 25 E...	88		co. s.	
			D5469...	13 36 48 N.; 123 38 24 E...	500		gn. m.	Rare.
12124	U.S.N.M.	3	D5478...	10 46 24 N.; 125 16 30 E...	57		sh.	Few.
			D5574...	5 30 45 N.; 120 07 57 E...	340			Rare.
12122	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.	Rare.
14188	U.S.N.M.	1	D5621...	0 15 00 N.; 127 24 35 E...	298		gy. and bk. s.	Rare.
14191	U.S.N.M.	1	D5636...	1 55 00 S.; 127 42 30 E...	1,262		gy. m., fine s.	Rare.
14182	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 00 E...	700		gy. m.	Rare.
14190	U.S.N.M.	3	D5640...	4 27 00 S.; 122 55 40 E...	24		s. brk. sh.	Few.
14178	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 56 E...	805	38.3		Rare.
14185	U.S.N.M.	1	D5658...	3 32 40 S.; 120 31 30 E...	510	41.2	gy. m.	Rare.
14183	U.S.N.M.	1		Binang Pool, Subin Bay...				

TEXTULARIA HAUERII d'Orbigny.

Plate 19, fig. 6.

Textularia hauerii D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 250, pl. 15, figs. 13-15.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 628, pl. 47, figs. 21-23.

The original description of this species is as follows:

A transition-form, which we figure, is generally distributed, and at the Stns. where it occurs is often one of the most abundant and typical. It seems to come

nearer to d'Orbigny's *T. hauerii* than to any other Textularian. Brady regarded *T. hauerii* as merely a modification of *T. gramen* (*ut supra*) characterized by less angular edges, but the Kerimba specimens are sufficiently marked and divergent from *T. gramen*, as represented in these dredgings, to render the separation of the specimens desirable. In the somewhat rough texture of the shell their affinities appear to lie rather in the direction of *T. candeiana* than of *T. gramen*.

The Kerimba specimens of *T. hauerii* have, as a rule, about eight pairs of chambers regularly increasing in breadth and thickness, and with the sutural lines somewhat depressed owing to the inflation of the chambers. Marginal edge lobulate and varying from rounded in the latter portion of the shell to acute or subcarinate in the initial portion. Surface-texture somewhat coarse, but neatly agglutinate. The species is most abundant at Stns. 1 and 12, reaching its best proportions at the latter. *T. hauerii* was recorded by d'Orbigny from the Tertiaries of Vienna. It is probably widely distributed, but has not been separated from the records of *T. gramen*.

In the above characterization of this species I heartily agree, and from two stations at least a similar form occurs. The specimens are from *Albatross* stations D5181, off eastern Panay in 26 fathoms (48 meters), and D5276, China Sea off southern Luzon in 18 fathoms (33 meters).

Textularia hauerii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14174	U.S.N.M.	2	D5181...	11 36 40 N.; 123 26 35 E...	26	° F.	m., fine s.....	Rare.
14173	U.S.N.M.	3	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Rare.

TEXTULARIA AGGLUTINANS d'Orbigny.

Plate 20, fig. 8.

Textularia agglutinans D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 136, pl. 1, figs. 17, 18, 32-34.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 363, pl. 43, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 9, figs. 10 a, b (in text).

Numerous records for this species, mostly from the southern portion of the area, have been obtained. These range in depth from 10 to 80 fathoms (18 to 146 meters), with a single record from D5574, north of Tawi Tawi Group, 340 fathoms (622 meters), and except for two other stations all the depths are for less than 40 fathoms (73 meters). Bottom temperatures are given in only two cases, these being 58.6° and 76.3° F. (14.7° C. and 24.6° C.).

The largest specimens are about 3 mm. in length.

Textularia agglutinans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14164	U.S.N.M.	3	D5097...	14 19 15 N.; 120 34 15 E...	28	° F.	gy. m., s.....	Few.
14170	U.S.N.M.	4	D5134...	6 44 45 N.; 121 48 00 E...	25	fne. s.....	Few.
14166	U.S.N.M.	6						
12093	U.S.N.M.	5	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Few.
14167	U.S.N.M.	2	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.....	Rare.
12098	U.S.N.M.	5	D5151...	5 24 40 N.; 120 27 15 E...	24	co. s., sh.....	Few.
			D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Few.
			D5154...	5 14 50 N.; 119 58 45 E...	12	co. s.....	Rare.
			D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.....	Rare.
14163	U.S.N.M.		3	D5161...	5 10 15 N.; 119 53 00 E...	16	fne. s.....
12097	U.S.N.M.	4	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. m.....	Few.
12096	U.S.N.M.	1	D5210...	11 49 55 N.; 124 28 05 E...	50	75.3	fne. gy. s.....	Rare.
14169	U.S.N.M.	2	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Few.
12099	U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fne. s.....	Rare.
14168	U.S.N.M.	10+	D5358...	6 06 40 N.; 118 18 15 E...	39	m.....	Frequent.
14172	U.S.N.M.		6	D5426...	9 12 00 N.; 118 28 00 E...	27	fne. gy. s.....
14165	U.S.N.M.	1	D5574...	5 30 45 N.; 120 07 57 E...	340	Rare.
			D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh.....	Few.
				Off Singapore.....	13	Rare.

TEXTULARIA AGGLUTINANS d'Orbigny, var. FISTULA Cushman.

Textularia agglutinans D'ORBIGNY, var. *fistula* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 10, fig. 11 (in text).

The only record for this variety is a single specimen from D5576, north of Tawi Tawi, 277 fathoms (506 meters). This is similar to the material which I have described from off the Hawaiian Islands and from off Japan.

Textularia agglutinans, var. *fistula*—Material examined.

Cat. No.	Collection of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
		1	D5576...	5 25 56 N.; 120 03 39 E...	277	° F. 53.3	s.....	Rare.

TEXTULARIA STRICTA Cushman.

Plate 21, fig. 1.

Textularia stricta CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 11, figs. 13a, b (in text).—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 20.

Spiroplecta sagittula, var. *fistulosa* CHAPMAN, Trans. New Zealand Inst., vol. 38, 1905 (1906), p. 87, pl. 3, fig. 4.

Numerous typical specimens were found at five stations: D5272, China Sea, vicinity southern Luzon, 118 fathoms (216 meters), bottom temperature 57.4° F. (14.1° C.); D5256, off southern Mindanao, 158 fathoms (289 meters), bottom temperature not given; D5282, China Sea, vicinity southern Luzon, 248 fathoms (454 meters), bottom temperature 47.4° F. (10.7° C.); D5318, China Sea, vicinity Formosa, 340 fathoms (556 meters), bottom temperature not given;

and D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

This is a large and striking species, often several millimeters in length and very straight, long, and narrow. It was originally described from *Albatross* station D4900 in the Eastern Sea off southwestern Japan, in 139 fathoms (254 meters). In this region many southern species range to the north, and it is not surprising to find this species somewhat more abundant in the Philippines.

The Philippine specimens are not as large as those from the type station off Japan. Sidebottom records this species from off the east coast of Australia in 465 fathoms (850 meters). It seems to be distinct from *T. porrecta* as developed in the Philippine region, and probably is a distinct Indo-Pacific species. The early chambers of the test are not always Spiroplectine, although they seem to be in the largest specimens, probably indicating that both the microspheric (Spiroplectine) and megalospheric forms occur. The following notes are from Sidebottom's paper:

Four occur, the largest having about forty chambers. They agree accurately with Cushman's description, excepting that the apertural end is not "somewhat acute." The same large tests occur at the *Challenger* St. 185, off Raine Island, but they are very roughly built.

Further investigation may prove that these are a variant of *Textularia agglutinans*, var. *porrecta* Brady.

Cushman's species appears to be identical with the forms recorded and figured by Chapman from Great Barrier Island, New Zealand, 110 fathoms, under the names *Spiroplecta sagittula* (Defrance) and *S. sagittula*, var. *fistulosa* Brady.⁹ It is a very large form, typical of the "Thomson Basin," and abundant in dredgings from this area. The early structure is in our experience usually, if not invariably, Spiroplectine, and Cushman's figure suggests a Spiroplectine form, though his description contains no reference to this feature. But the *Dart* specimens are of a megalospheric *Textularian* type, and this would seem to be proof of dimorphism. The species deserves to be carefully worked out.

I agree with Sidebottom that Chapman's material, especially the figured specimen, belongs to this species. It is the microspheric form as shown in the figure and by its being placed under *Spiroplecta*.

Textularia stricta—Material examined.

Cat. No.	Collection of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12067	U.S.N.M.	3	D5236...	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Few.
			D5256...	8 50 45 N.; 126 26 52 E..	158	m.....	
12068	} U.S.N.M.	10+	D5272...	7 21 45 N.; 124 07 15 E..	118	57.4	m., sh., co. s.	Common.
12069					14 00 00 N.; 120 22 30 E..			
12071	U.S.N.M.	1	D5282...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. gy. s....	Rare.
12070	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Rare.

⁹ Chapman, 1906, *Trans. Zealand Inst.*, vol. 33, 1905, p. 87, pl. 3, fig. 4.

TEXTULARIA CANDEIANA d'Orbigny.

Textularia candeiana D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 143, pl. 1, figs. 25-27.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 12, fig. 14-17 (in text).

Textularia sagittula, var. *candeiana* MILLETT, Journ. Roy. Micr. Soc., 1899, p. 562, pl. 7, fig. 12.

At two stations specimens of this species were rare. These were D5210, off western Samar, 50 fathoms (91 meters), bottom temperature 76.3° F. (24.6° C.), and D5630, south of Patiente Strait, 569 fathoms (1,081 meters), bottom temperature not recorded.

The previous North Pacific records for this species are off the Hawaiian Islands and Gaspar Straits. Millett records it in his Malay Archipelago paper in the region adjacent to that of our southern stations recorded here.

Textularia candeiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12048	U.S.N.M.	1	D5210...	° ' " ° ' "	50	° F. 76.3	fne. gy. s.	Rare.
12049	U.S.N.M.	1	D5630...	11 49 55 N.; 124 28 05 E.. 0 56 30 S.; 128 05 00 E..	569	co. s., m.	Rare.

TEXTULARIA PORRECTA H. B. Brady.

Plate 22, fig. 1.

Textularia agglutinans D'ORBIGNY, var. *porrecta* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 364, pl. 43, figs. 4a, b.

Textularia porrecta EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 269, pl. 6, figs. 17, 18.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 627.

Brady in the *Challenger* report refers to this variety as the "much elongated, subcylindrical, arenaceous *Textulariæ*," which are very common "in some localities, especially about the coral reefs of the tropics." In the shallow water dredgings in the Philippine region this form is extremely abundant among the reefs, ranking with *T. rugosa*, *T. vertebralis*, and other species in numbers. It is not always found in company with specimens which may be referred to *T. agglutinans*, and withal in the Philippine material seems worthy of specific rank.

Specimens occur at many stations under 50 fathoms (91 meters), but comparatively few show greater depths. The species is much more common in the southern half of the Archipelago. In two cases the temperature of the bottom is given for the shallow-water stations, 76.3° F. (24.6° C.) and 79.5° F. (26.3° C.). A few specimens were

found in material from the deeper stations, not in sufficient numbers to be classed as characteristic, but rather as an occasional specimen.

Heron-Allen and Earland record the species from the Kerimba Archipelago, off southeast Africa, apparently having the microspheric form with spiroplectine young.

Textularia porrecta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
14155	U.S.N.M.	4)	D5133...	Id. off Panabutan Pt., N. 52° E., 1.50 miles.	38	79.5	gn. m., s....	Few.
12102	U.S.N.M.	5)						
13137	U.S.N.M.	1	D5146...	5 46 40 N.; 120 48 50 E...	24	co. s., sh....	Rare.
12107	U.S.N.M.	1	D5192...	11 9 15 N.; 123 50 00 E...	32	gn. s.....	Rare.
12106	U.S.N.M.	2	D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fine. gy. s....	Rare.
14139	U.S.N.M.	3	D5217...	13 00 20 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Few.
12103	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Few.
14135	U.S.N.M.	3	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Few.
12105	U.S.N.M.	3	D5478...	10 46 24 N.; 125 16 30 E...	57	sh.....	Few.
13136	U.S.N.M.	6	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Few.
12101	U.S.N.M.	3	D5529...	9 23 45 N.; 123 39 30 E...	441	53	gy. m., glob.	Few.
14138	U.S.N.M.	2	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s....	Few.
12104	U.S.N.M.	4	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Few.
14140	U.S.N.M.	3	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.....	Few.
14141	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 45 E...	88	co., s.....	Rare.
14142	U.S.N.M.	2	D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.....	Rare.
14143	U.S.N.M.	1	D5426...	9 12 00 N.; 118 28 00 E...	27	fine. gy. s....	Rare.
14144	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E...	19	s., Co.....	Rare.
14145	U.S.N.M.	1	D5358...	6 06 40 N.; 118 18 50 E...	39	m.....	Rare.
14146	U.S.N.M.	10	D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s.....	Common.
14147	U.S.N.M.	2	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co....	Rare.
14148	U.S.N.M.	5	D5132...	Id. off Panabutan Pt., N. 15° W., 0.3 mile.	26	gn. m., s....	Few.
14149	U.S.N.M.	5	D5213...	12 15 00 N.; 124 04 36 E...	108	59.9	gy. s., m....	Few.
14150	U.S.N.M.	8	D5096...	14 20 23 N.; 120 34 15 E...	23	gy. m., s., sh.	Few.
14151	U.S.N.M.	1	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh., g....	Rare.
14152	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s....	Rare.
14153	U.S.N.M.	2	D5097...	14 19 15 N.; 120 33 52 E...	30	gy. m., s., sh.	Rare.
14154	U.S.N.M.	2	D5105...	14 43 55 N.; 120 12 50 E...	25	Rare.

TEXTULARIA VERTEBRALIS Cushman.

Plate 22, fig. 3; plate 24, fig. 1.

Textularia vertebralis CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 633, pl. 78, fig. 1.

Description.—Test elongate and tapering, chambers very numerous, in the adult portion rounded in cross-section, the distal portion of the chambers convex, the proximal portion concave behind the middle, chambers gradually increasing in size, wall composed of agglutinated sand usually fine and smoothly finished; aperture at the base of the inner margin of the chamber; color, gray.

Length up to 6 mm.

This is one of the most common species in the region, occurring in deeper water than the other common species, *T. rugosa*, *T. porrecta*, etc. The stations range from 80 to 742 fathoms (146 to 1,357 meters) in depth, the average 275 fathoms (503 meters), and the bottom temperatures where given from 43.3° to 59.6° F. (6.1° C. to 15.3° C.), most of them above 50° F. (10° C.).

This species is a very characteristic one, resembling a series of vertebrae, the chambers in adult specimens very numerous, as many as 60 having been counted in some specimens. It is also one of the largest species of the genus. At some of the stations it was very common.

Besides occurring in the Philippine Archipelago it occurred at several stations in the vicinity of Sibuko Bay, Borneo, and between Gillolo and Kayoa Islands.

Textularis vertebralis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
14101	U.S.N.M.	3	D5110...	13 21 30 N.	120 30 33 E.	220	° F.	gn. m.	Few.
14105	U.S.N.M.	1	D5126...	10 48 00 N.	121 47 30 E.	742	49.5	ft. gn. m.	Rare.
14112	U.S.N.M.	8	D5198...	9 40 50 N.	123 39 45 E.	220	53.9	gn. m.	Common.
12081 14111	U.S.N.M.	10+	D5259...	11 57 30 N.	121 42 15 E.	312	49.3	gy. m., glob.	Abundant.
14120	U.S.N.M.	1	D5260...	12 25 35 N.	121 31 35 E.	234	51.4	gn. m., s.	Few.
14108	U.S.N.M.	8	D5277...	13 56 55 N.	120 13 45 E.	80	53.6	fine s.	Frequent.
14113	U.S.N.M.	7	D5278...	14 00 10 N.	120 17 15 E.	102	59.6	fine s., m., sh.	Few.
14102	U.S.N.M.	8	D5281...	13 52 45 N.	120 25 00 E.	201	50.4	dk. gy. s.	Frequent.
14114	U.S.N.M.	1	D5291...	13 39 40 N.	121 00 45 E.	173	51.5	fine bk. s.	Rare.
12076 12080	U.S.N.M.	10+	D5301...	20 37 00 N.	115 43 00 E.	208	50.5	gy. m., s.	Frequent.
14100	U.S.N.M.	1	D5374...	13 46 45 N.	121 35 08 E.	190	gy. m.	Rare.
14167	U.S.N.M.	1	D5331...	15 36 45 N.	119 47 45 E.	178	54.7	s., sh., m.	Rare.
12079	U.S.N.M.	1	D5424...	9 37 05 N.	121 12 37 E.	340	50.4	co. s.	Frequent.
14115	U.S.N.M.	9	D5430...	9 49 40 N.	119 03 20 E.	464	50.4	glob. oz.	Rare.
14115	U.S.N.M.	2	D5454...	13 12 00 N.	123 50 30 E.	153	Rare.
12077 8501	U.S.N.M.	10+	D5512...	8 16 02 N.	123 58 26 E.	445	52.8	gy. m., fine s.	Frequent.
14106	U.S.N.M.	3	D5523...	8 48 44 N.	123 27 35 E.	Few.
14119	U.S.N.M.	5	D5529...	9 23 45 N.	123 39 30 E.	441	53	gy. m., glob.	Few.
12082	U.S.N.M.	10+	D5537...	9 11 00 N.	123 23 00 E.	254	53.5	gn. m.	Frequent.
14109	U.S.N.M.	3	D5538...	9 08 15 N.	123 23 20 E.	256	53.3	gn. m., s.
.....	D5548...	6 00 20 N.	120 45 35 E.	232	53.5	s., brk. sh.	Rare.
12078	U.S.N.M.	6	D5567...	5 48 00 N.	120 33 45 E.	268	52	fine s.	Few.
14121	U.S.N.M.	2	D5569...	5 33 15 N.	120 15 30 E.	303	52.3	co. s.	Rare.
14122	U.S.N.M.	1	D5570...	5 32 15 N.	120 12 57 E.	330	52.3	fine s., glob.	Rare.
.....	D5576...	5 25 56 N.	120 03 39 E.	277	53.3	s.	Few.
.....	D5586...	4 06 50 N.	118 47 20 E.	347	44	gy. m.	Rare.
14103	U.S.N.M.	4	D5589...	4 12 10 N.	118 38 08 E.	260	45.7	fine gy. s., gy. m.	Few.
14104	U.S.N.M.	1	4 10 50 N.	118 39 35 E.	310	44.3	gn. m., s.	Few.
14117	U.S.N.M.	2	D5592...	4 12 44 N.	118 27 44 E.	305	43.3	gn. m.	Few.
14118	U.S.N.M.	3	D5625...	0 07 00 N.	127 28 00 E.	230	gy. m., fine s.	Few.
.....	H4998...	7 43 45 N.	122 03 45 E.	221	gy. m., glob.	Few.

TEXTULARIA APERTURALIS Cushman.

Textularia solita (SCHWAGER), var. *inflata* GOËS, Bull. Mus. Comp. Zoöl., vol. 29, 1896, p. 42, pl. 5, figs. 1-3.

Textularia aperturalis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 20, figs. 34, 35 (in text).

The only record for the region is from D5259, 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.). Although the aperture was not double as in the type, nevertheless it was very oblique, more on one side than the other, and the whole shape of the test was similar to the figured specimens.

Textularia aperturalis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12141	U.S.N.M.	1	D5259...	° ' " ° ' " 11 57 30 N.; 121 42 15 E..	312	° F. 49.3	gy. m., glob.	Rare.

TEXTULARIA CATENATA Cushman.

Plate 23, fig. 5.

Textularia catenata CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 23, figs. 39, 40 (in text).

The species occurred at a single station, so far as noted, D5551, vicinity of Jolo Island, in 193 fathoms (353 meters), bottom temperature 53.3° F. (11.8° C.). The other North Pacific stations where this species has been recorded are very deep in comparison with this.

Textularia catenata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12142	U.S.N.M.	1	D5551...	° ' " ° ' " 5 54 48 N.; 120 44 24 E..	193	° F. 53.3	fine s.	Rare.

TEXTULARIA ASPERA H. B. Brady.

Plate 23, fig. 6.

Textularia aspera H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 715; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 367, pl. 44, figs. 9-13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 14, figs. 21-23 (in text).

From a single station, D5121, east coast of Mindoro, 108 fathoms (198 meters), a few specimens of this species were obtained. It was not noted elsewhere in the region.

Textularia aspera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12147	U.S.N.M.	2	D5121...	° ' " ° ' " 13 27 20 N.; 121 17 45 E..	108	° F.	dk. gn. m...	Rare.

TEXTULARIA GOËSI Cushman.

Plate 21, fig. 3.

Textularia sagittula DEFRANCE, var., GOËS, Kongl. Svensk. Vet. Akad. Handl., vol. 19, No. 4, 1882, pl. 5, figs. 150-158.

Textularia trochus H. B. BRADY (part), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 366, pl. 43, fig. 17 (not 15, 16, 18, 19); pl. 44, figs. 1-3 (not *T. trochus* d'Orbigny).

Textularia goësi CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 15, figs. 24a, b (in text).

At several stations material referable to this species was found representing the following localities: Sulu Archipelago, Tawi Tawi Group; China Sea, off southern Luzon; off southeastern Mindoro; north of Tawi Tawi; Mindoro Strait; vicinity of Jolo; and east coast of Luzon; San Bernardino Strait to San Miguel Bay.

Textularia goësi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12140	U.S.N.M.	4	D5110...	° ' " ° ' "	135	° F. 59.0	dk. gy. m...	Common.
			D5152...	13 59 20 N.; 120 75 45 E...	34	wh. s.....	Rare.
12135	U.S.N.M.	1	D5172...	5 22 55 N.; 120 15 45 E...	318	fne. s., sh.....	Rare.
			D5261...	6 03 15 N.; 120 35 30 E...	145	s., m.....	Rare.
12139	U.S.N.M.	1	D5268...	12 30 55 N.; 121 34 24 E...	170	s., p.....	Rare.
12138	U.S.N.M.	2	D5291...	13 42 00 N.; 120 57 15 E...	173	51.5	fne. bk. s.....	Rare.
			D5333...	13 29 40 N.; 121 00 45 E...	310	73.8	s.....	Rare.
12137	U.S.N.M.	1	D5469...	12 26 30 N.; 120 37 45 E...	500	gn. m.....	Rare.
12136	U.S.N.M.	1	D5569...	13 36 48 N.; 123 38 24 E...	303	52.3	co. s.....	Rare.

TEXTULARIA FLINTII Cushman.

Plate 22, fig. 4.

Textularia agglutinans FLINT (part), Rep. U. S. Nat. Mus., 1897 (1899), p. 284, pl. 29, fig. 4 (in part) (not *T. agglutinans* d'Orbigny).

Textularia rugosa BAGG, Proc. U. S. Nat. Mus., vol. 34, 1908, p. 131, (not *T. rugosa* (Reuss)).

Textularia flintii CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 21, figs. 36a, b (in text).

At three stations this species has been found: D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5301, China Sea, 208 fathoms (381 meters), bottom temperature 50.5° F. (10.2° C.); and D5586, vicinity Sibuko Bay, Borneo, 347 fathoms (569 meters), bottom temperature, 44.0° F. (6.4° C.). The material was typical and in considerable numbers.

I have already elsewhere recorded this species from off the Hawaiian Islands, off Guam, and off the Bonin Islands, so it might have been expected in the Philippine region.

Textularia flintii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12146	U.S.N.M.	3	D5236...	° / " ° / "		° F.		
12092	U.S.N.M.	1	D5301...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Few.
12091	U.S.N.M.	3	D5586...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s....	Rare.
				4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.	Rare.

TEXTULARIA RUGOSA (Reuss).

Plate 23, figs. 3, 4.

Plecanium rugosum REUSS, Sitz. Akad. Wiss. Wien, vol. 59, 1869, p. 453, pl. 1, figs. 3a, b.

Textularia rugosa H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 363, pl. 42, figs. 23, 24.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 625, pl. 47, figs. 7-9.

Throughout the region in shallow water this species is common. Except in a few cases the depths are less than 50 fathoms (91 meters) and the most of these less than 25 fathoms (33 meters).

This is the first record of the species from the North Pacific, but it has been found in shallow water about several of the South Sea Islands. It is a typical coral-reef species and at some shallow water stations very abundant.

The bottom temperatures of the shallow water stations were recorded but once, that being 75.7° F. (24.2° C.), 37 fathoms (68 meters).

There is a tendency in the largest specimens which may attain a length of 6 mm. to contract the test, showing definite traces of senescence. At the same time there is also a tendency to lessen the amount of the excavations at the base of the chambers. In some of the specimens these excavations are very deep and sharply sculptured, giving the whole test a very ornate appearance. There is some considerable variation in the size and depth of the excavations, and they may become filled with amorphous material in some cases.

Besides being widely scattered in shallow water in the Archipelago proper, it is found to the southward in both Sibuko and Darvel Bays, Borneo, but not in the deeper water of other stations in this region.

The specimen recorded by Flint from deep water off Bermuda does not seem to belong here. Heron-Allen and Earland record it from the Kerimba Archipelago off south eastern Africa, and note¹⁰ that they "have specimens from the Philippine Islands one-sixth inch in length."

¹⁰ Journ. Roy. Micr. Soc., 1911, p. 310.

Textularia rugosa.—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
							° F.		
14075	U.S.N.M.	6	D5109...	14 03 45 N.;	120 16 30 E..	10	Co.....	Few.
14074	U.S.N.M.	1	D5110...	13 59 20 N.;	120 75 45 E..	135	59.0	dk. gy. m...	Rare.
12089	U.S.N.M.	10+	D5134...	6 44 45 N.;	121 48 00 E..	25	fine. s.....	Abundant.
14086	U.S.N.M.	1	D5136...	6 04 20 N.;	120 59 20 E..	22	s., sh.....	Rare.
14073	U.S.N.M.	10+	D5137...	6 04 25 N.;	120 58 30 E..	20	s., sh.....	Abundant.
14076	U.S.N.M.	10+	D5141...	6 09 00 N.;	120 58 00 E..	29	co. s.....	Frequent.
12087	U.S.N.M.	10+	D5142...	6 06 10 N.;	121 02 40 E..	21	co. s., sh...	Rare.
12086	U.S.N.M.	10+	D5143...	6 05 50 N.;	121 02 15 E..	19	co. s.....	Frequent.
14077	U.S.N.M.	10+	D5144...	6 05 50 N.;	121 02 15 E..	19	co. s.....	Few.
14087	U.S.N.M.	5	D5147...	5 41 40 N.;	120 47 10 E..	21	co. s.....	Frequent.
14090	U.S.N.M.	10+	D5148...	5 35 40 N.;	120 47 30 E..	17	co. s.....	Few.
14085	U.S.N.M.	10+	D5151...	5 24 40 N.;	120 27 15 E..	24	co. s., sh...	Abundant.
12088									
12085	U.S.N.M.	9	D5152...	5 22 55 N.;	120 15 45 E..	34	wh. s.....	Common.
			D5159...	5 11 50 N.;	119 54 00 E..	10	co. s.....	Rare.
14091	U.S.N.M.	1	D5178...	12 43 00 N.;	122 06 15 E..	78	fine. s.....	Rare.
12090	U.S.N.M.	10+	D5179...	12 38 15 N.;	122 12 30 E..	37	75.7	hrd. s.....	Abundant.
14084	U.S.N.M.	1	D5192...	11 09 15 N.;	123 50 00 E..	32	gn. s.....	Few.
			D5217...	13 20 00 N.;	123 14 15 E..	105	63.1	crs. gy. s...	Few.
14089	U.S.N.M.	6	D5218...	13 11 15 N.;	123 02 45 E..	20	sh., p., s...	Few.
14088	U.S.N.M.	10+	D5276...	13 49 15 N.;	120 14 45 E..	18	sh., p., s...	Common.
14079	U.S.N.M.	2	D5278...	14 00 10 N.;	120 17 15 E..	102	59.6	fine. s., m., sh.	Few.
14078	U.S.N.M.	1	D5336...	11 37 45 N.;	119 46 00 E..	46	s., m.....	Rare.
12084	U.S.N.M.	2	D5478...	10 46 24 N.;	125 16 30 E..	57	sh.....	Rare.
			D5576...	5 25 56 N.;	120 03 39 E..	277	53.3	s.....	Few.
			D5580...	4 52 45 N.;	119 06 45 E..	162	55.8	br. s., Co...	Few.
12083	U.S.N.M.	2	D5589...	4 12 10 N.;	118 38 08 E..	260	45.7	fine. gy. s., gy. m.	Frequent.
14081	U.S.N.M.	1	D5640...	4 27 00 S.;	122 55 40 E..	24	s., brk. sh...	Rare.
				From exterior of pearl oyster.					Few.
14080	U.S.N.M.	1	D5236...	8 50 45 N.;	126 26 52 E..	494	41.2	fine. gy. s....	Rare.
14082	U.S.N.M.	4	D5149...	5 33 00 N.;	120 42 10 E..	10	Co., sh.....	Few.
14083	U.S.N.M.	1	D5570...	5 32 15 N.;	120 12 57 E..	330	52.3	fine. s., glob..	Rare.

TEXTULARIA SIPHONIFERA H. B. Brady.

Plate 21, figs. 4-7.

Textularia siphonifera H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 53; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 362, pl. 42, figs. 25-29.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 17, figs. 28, 29 (in text).
Spiroplecta siphonifera CHAPMAN, Proc. Roy. Soc. Victoria, vol. 22, 1909 (1910), p. 272, pl. 53, fig. 1.

This is one of the most abundant species of the genus in the shallow water of the region. The depth, except in six cases, is less than 90 fathoms (165 meters), most of them being less than 25 fathoms (46 meters). Single specimens have occurred in deeper water at D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature 53.8° F. (12.1° C.); D5569, north of Tawi Tawi Group, 303 fathoms (554 meters), bottom temperature 52.3° F. (10.1° C.); D5576, north of Tawi Tawi, 277 fathoms (506 meters), bottom temperature 53.3° F. (11.8° C.). The temperatures at the other stations are usually not given for the shallow water, but in the single case recorded it is 75.7° F. (24.2° C.).

As a rule there are six series of siphons, but specimens with eight series are not infrequent. Specimens were found in which one of the series of siphons was split into two and later on were united.

Both microspheric and megalospheric specimens were found at numerous stations. This seems to be a species which is peculiar to coral-reef regions as has been noted in its occurrence elsewhere.

In the remainder of the North Pacific it is known as far north as southern Japan and as far east as the Hawaiian Islands. Southward the records include the Admiralty and Friendly Islands, but it is not recorded from the Australian region except as a fossil, nor did I find it in the material I have had from Murray Island on the Great Barrier Reef of Australia. Chapman records it from Funafuti. Brady gives a record from the Gulf of Suez, but Heron-Allen and Earland do not record it from the Kerimba Archipelago off south-eastern Africa. Its distribution, therefore, while wide, is evidently restricted within this range.

Textularia siphonifera.—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14126	U.S.N.M.	2	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m...	Rare.
14131	U.S.N.M.	10+	D5134...	6 44 45 N.; 121 48 00 E..	25	fine s.....	Common.
12034	U.S.N.M.	8	D5137...	6 04 25 N.; 120 58 30 E..	20	s., sh.....	Abundant.
14124	U.S.N.M.	1	D5141...	6 09 00 N.; 120 58 00 E..	29	co. s.....	Rare.
12031	U.S.N.M.	10+	D5142...	6 06 10 N.; 121 02 40 E..	21	co. s., sh.....	Frequent.
12033	U.S.N.M.	9	D5151...	5 24 40 N.; 120 27 15 E..	24	co. s., sh.....	Common.
14123	U.S.N.M.	2	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Rare.
14128	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	53.8	gy. s., m.....	Rare.
12035	U.S.N.M.	10+	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Frequent.
12036	U.S.N.M.	1	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s.....	Few.
14127	U.S.N.M.	1	D5342...	10 56 55 N.; 119 17 24 E..	14-25	gy. m.....	Few.
			D5381...	13 14 15 N.; 122 44 25 E..	88	co. s.....	Rare.
12037	U.S.N.M.	1	D5386...	13 38 30 N.; 122 44 30 E..	287	62.4	Rare.
12038	U.S.N.M.	1	D5481...	10 27 30 N.; 125 17 10 E..	61	s., sh., g.....	Rare.
14130	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E..	Rare.
12032	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E..	303	52.3	co. s.....	Rare.
14129	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s.....	Rare.
12039	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Rare.
				East end of Midway Island, Jolo Jolo.	Rare.
14125	U.S.N.M.	10+	From interior of pearl oyster.	Common.

TEXTULARIA SEMIALATA Cushman.

Plate 24, figs. 2, 3 a, b.

Textularia semialata CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 634, pl. 80, figs. 6, 7.

Description.—Test much compressed, of numerous chambers, broader than high, the proximal outer angle of the adult chambers more or less projecting and extending backward, wall of fine sand, very smoothly finished; aperture at the inner margin of the chamber; color gray.

Length about 1 mm.

Type specimen.—Cat. No. 8504, U.S.N.M., from *Albatross* station D5214, east of Masbate Island, 218 fathoms (399 meters).

This species in some of its characters resembles *Bolivina beyrichi* Reuss. It is a true *Textularia*, however, and has many points in which it differs from that species. There is a considerable difference in the microspheric and megalospheric forms, both of which are here figured. The microspheric form is narrow at the beginning and continues this form for some time, finally broadening out somewhat. The megalospheric form starts almost at once to develop a broad test of fewer chambers, but each has the same characteristic shape of the chambers.

Textularia semialata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8504 12148 12149	U.S.N.M	3	D5214...	° ' "	218	° F. 51.4	gn. m.	Few.
' "				' "				

TEXTULARIA FOLIACEA Heron-Allen and Earland.

Plate 19, figs. 7 a, b.

Textularia foliacea HERON-ALLEN and EARLAND, Trans. Zool. Soc., London, vol. 20, 1915, p. 628, pl. 47, figs. 17-20.

The original description is as follows:

Test free, highly compressed, consisting of seven to nine pairs of chambers, regularly increasing in width so as to give a leaf-shaped outline to the shell. Sutural lines depressed, often strongly marked, but at times very obscure, and obliquely set, so that the tapering off of the ultimate pair of chambers gives the characteristic diamond or foliaceous outline. Median line of the shell depressed below the marginal edges, aperture small and regularly textularian. Test composed of sand-grains and other adventitious substances firmly and neatly cemented together, but with a rough external surface.

This is one of the most characteristic of the Kerimba Textulariidae, though not universally distributed. It occurs in the greatest abundance and in its best development at Stn. 1, and very fine examples were also obtained at Stn. 9. At Stns. 3 and 10 the specimens were less characteristic and showed a tendency to pass, by inflation of the chambers, into *T. hauerii*.

The affinities of our species are between *T. luculenta* Brady and *T. hauerii* or *gramen* d'Orbigny. It may be compared as regards its highly compressed and parallel-faced test with the *Textularia immensa* of Cushman¹¹ from which it differs only in the character of its aperture.

We have specimens from Timor Sea (Java, 50 fms.), where it is frequent, and from Vavau (S. Pacific, 16 fms.). We have also observed specimens in some of Brady's unsorted material from Fiji at Cambridge, and a single specimen among his specimens of *T. agglutinans* from "Coral Reef, Australia, 17 fms." It is probably therefore widely distributed in coral-reef areas.

Length, 1 to 1.5 mm.; breadth, 0.6; thickness, 0.3 mm.

¹¹ Proc. U. S. Nat. Mus., vol. 44, 1913, p. 633, pl. 79, fig. 2.

This species has occurred in very typical form in the Philippine material and was one of those laid aside in 1910 for later description. I agree with the authors that this is a very distinct species and the Philippine specimens are very close to the figured specimens of Heron-Allen and Earland.

Most of the stations at which it is recorded are in warm shallow water, where *Calcarina*, *Alveolina* and other shallow-water genera are abundant.

The species probably has a wide distribution in coral-reef regions in the Indo-Pacific.

Textularia foliacea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14196	U.S.N.M.	1	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.....	Rare.
14198	U.S.N.M.	3	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.....	Few.
14194	U.S.N.M.	4	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Few.
14193	U.S.N.M.	2	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Few.
14199	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	inc. gy. s.....	Few.
14200	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.....	Rare.
14192	U.S.N.M.	3	Tacloban Anchorage, P. I.	Few.

TEXTULARIA IMMENSA Cushman.

Plate 24, figs. 4a, b.

Textularia immensa CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 633, pl. 79, fig. 2.

Description.—Test large and broad, rhomboid, very much flattened, chambers low and long; wall thick, of rather coarse angular sand grains imbedded in an unusually large amount of light gray cement; aperture consisting of a series of small openings running from the inner margin of the apertural face to the highest point at the distal end of the test, about 20 in number; color gray.

Length up to 6 mm.; breadth up to 4 mm.; thickness 0.5 mm.

Type specimen.—Cat. No. 8502, U.S.N.M., from *Albatross* station, D5567, north of Tawi Tawi, 268 fathoms (495 meters), bottom temperature 52.0° F. (11.1° C.).

This is a very large species, and with its peculiar aperture is of more than usual interest. The whole test is rhomboid, very flattened, the two sides seeming to differ, one being slightly convex, the other slightly concave, the concave side having the sutures more prominent than the other.

It has occurred elsewhere in the region at station D5576, north of Tawi Tawi, 277 fathoms (506 meters), bottom temperature 53.3° F. (11.8° C.).

Textularia immensa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8502	U.S.N.M.	1	D5567...	5 48 00 N.; 120 33 45 E....	268	52	fne. s.....	Rare.
12145	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E....	277	53.3	s.....	Rare.

TEXTULARIA CONCAVA (Karrer).

Plecanium concavum KARRER, Sitz. kais. Akad. Wiss. Wien., vol. 58, 1868, p. 129, pl. 1, fig. 3.

Textularia concava H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 360, pl. 42, figs. 11, 12 (not pl. 43, fig. 11).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 22, fig. 38 (in text).

Material of this species occurred at but two stations—D5096, China Sea off southern Luzon, 28 fathoms (51 meters), and D5590, Sibuko Bay, Borneo, 310 fathoms (567 meters). It does not seem to be common in the North Pacific, the only other material I have had being from off the Hawaiian Islands.

Millett records it from the Malay region.

Textularia concava—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12109	U.S.N.M.	1	D5096...	14 20 23 N.; 120 34 15 E....	28	gy. m., s., sh.	Rare.
12108	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E....	310	44.3	gn. m., s.....	Rare.

TEXTULARIA JUGOSA H. B. Brady.

Textularia jugosa H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 358, pl. 42, figs. 7a, b.

A single specimen almost exactly like the figure given by Brady was found in material from D5551, vicinity of Jolo Island, 193 fathoms (353 meters), bottom temperature 53.3° F. (11.8° C.).

The *Challenger* material was from Torres Strait, 155 fathoms, and there are other records from the Indian Ocean and off Australia.

Textularia jugosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
		1	D5551...	5 54 48 N.; 120 44 24 E....	193	53.3	fne. s.....	Rare.

TEXTULARIA EXCAVATA Cushman.

Plate 24, figs. 5a, b.

Textularia excavata CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 634, pl. 79, fig. 5.

Description.—Test rhomboid, composed of comparatively few chambers, early portion with the sides angled, later chambers broadly rounded at sides, chambers obliquely set, each forming an angle of 90° with the opposite chamber, distal portion of chamber thickened and tending to roll back, giving an excavated appearance to the test, outer face of last-formed chamber in adult broadly rounded and convex; aperture at the base of the inner margin of the chamber; wall composed of rather fine sand with a grayish-brown color, due partly to the cement.

Length up to 2 mm. in adult specimens.

Type specimen.—Cat. No. 8503, U. S. N. M., from *Albatross* station D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters). At this station the species was common. Its shape and peculiar sculptured test will distinguish it from other species.

This species has also been found at D5318, China Sea, vicinity of Formosa, 340 fathoms (556 meters), and D5591, Sibuko Bay, Borneo, 260 fathoms (475 meters). Bottom temperatures were not recorded at either of these stations.

Textularia excavata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8503	} U.S.N.M.	10+	D5236...	° ' " ° ' "	494	° F.		
12127		1	D5318...	8 50 45 N.; 126 26 52 E...	41.2	fne. gy. s....	Common.	
12128		3	D5591 ..	21 32 00 N.; 117 46 00 E...	340	s., br. Co....	Rare.	
12126				4 11 48 N.; 118 38 20 E...	260		Few.	

TEXTULARIA TRANSVERSARIA H. B. Brady.

Plate 22, figs. 2 a-c.

Textularia transversaria H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 359, pl. 113, figs. 3-5.

A few specimens which compare closely with Brady's species occurred at D5572, north of Tawi Tawi, 334 fathoms (1,611 meters), bottom temperature 52.3° F. (11.2° C.). Brady's specimens were from this same faunal area, off Raine Island, Torres Strait, 155 fathoms (283 meters), and off Kandavu, Fiji Islands, 255 fathoms (466 meters).

Textularia transversaria—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12050	U.S.N.M.	2	D5572...	° ' " ° ' " 5 31 26 N.; 120 09 45 E..	334	° F. 52.3	s.....	Rare.

TEXTULARIA CRASSISEPTA Cushman.

Plate 23, fig. 1.

Textularia crassisepta CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 24, figs. 41 a, b (in text).

Specimens which were typical were found at three stations—D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5586, vicinity Sibuko Bay, Borneo, 347 fathoms (635 meters), bottom temperature 44° F. (6.6° C.); and D5121, east coast of Mindoro, 108 fathoms (198 meters), bottom temperature not given.

The type specimens were from off the Hawaiian Islands, *Albatross* H3007, in 323 fathoms (590 meters).

Textularia crassisepta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12144	U.S.N.M.	2	D5121...	° ' " ° ' " 13 27 20 N.; 121 17 45 E..	108	° F.	dk. gn. m....	Few.
14178	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s....	Few.
12143	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	44	gy. m.....	Rare.

TEXTULARIA PSEUDOCARINATA, new name.

Plate 22, fig. 5.

Textularia carinata H. B. BRADY, Rep. Voy. *Challenger*. Zoology, vol. 9, 1884, p. 360, pl. 42, figs. 15, 16 (not *T. carinata* d'Orbigny).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 17, figs. 26, 27a, b (in text).

From a study of d'Orbigny's descriptions and figures of this species with a series of recent Philippine specimens it seems that the recent tropical species is a different one from the fossil one in many ways. In this recent tropical species the carina is broad and has very prominent denticulations and a heavy layer of agglutinated material above the sutures connecting across the middle and united

with the material of the carina at either side. Brady records this species at *Challenger* station 209 in 95 fathoms (174 meters) off the Philippines. Our specimens are almost identical with the figured specimens in the *Challenger* Report. I have had material from the following stations: D5121, east coast of Mindoro, 108 fathoms (198 meters); D5272, China Sea, vicinity southern Luzon, 118 fathoms (203 meters), bottom temperature 57.4° F. (14.1° C.); D5261, off southeastern Mindoro, 145 fathoms (265 meters); D5543, northern Mindanao and vicinity, 162 fathoms (297 meters), bottom temperature 54.5° F. (12.5° C.); H4898, Sulu Sea off western Mindanao, 221 fathoms (404 meters).

Textularia pseudocarinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12073	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E..	108	dk. gn. m...	Rare.
12075	U.S.N.M.	1	D5261...	12 30 55 N.; 121 34 24 E..	145	s. m.....	Rare.
12074	U.S.N.M.	3	D5272...	14 00 00 N.; 120 22 30 E..	118	57.4	m...sh., co.s.	Rare.
12072	U.S.N.M.	4	D5543...	8 47 15 N.; 123 35 00 E..	162	54.5	s.....	Rare.
14219	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob.	Rare.

TEXTULARIA ABBREVIATA d'Orbigny.

Plate 21, figs. 2a, b.

Textularia abbreviata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 249, pl. 15, figs. 9-12 (7-12).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 14, figs. 20a, b (in text).

Specimens referred to this species are common in the dredgings throughout the region, occurring at a large number of stations. They are similar to the figure given in the North Pacific Monograph referred to above. It may be questioned whether the specimen I figured there really is the *T. abbreviata* of d'Orbigny, but throughout the Philippine region and beyond, the specimens agree well with my figured specimen. Most of the stations are in the Philippine Archipelago, but the species is also common in the vicinity of Sibuko Bay, Borneo. In depth the stations vary from 108 to 775 fathoms (198 to 1,417 meters), and the bottom temperature from 41.1° to 59.9° F. (5° to 15.5° C.).

Textularia abbreviata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
			D5115...	13 37 11 N.; 120 43 40 E...	340	
			D5172...	6 03 15 N.; 120 35 30 E...	318	fine. s., sh.	
			D5201...	19 19 99 N.; 125 94 15 E...	554	53.8	gy. s., m.	
13170	U.S.N.M.	1	D5203...	9 58 00 N.; 125 07 40 E...	775	52.9	gn. m.	Rare.
			D5212...	12 04 15 N.; 124 04 36 E...	108	59.9	gy. s., m.	
13169	U.S.N.M.	6	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m.	Few.
			D5219...	13 21 00 N.; 122 18 45 E...	530	50.8	gn. m.	
14209	U.S.N.M.	10+	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Common.
			D5260...	12 25 35 N.; 121 31 35 E...	234	51.4	gn. m.	
14211	U.S.N.M.	4	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co.	Few.
			D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.	
			D5388...	12 51 30 N.; 123 26 15 E...	226	51.4	sft. gn. m.	
			D5430...	9 49 40 N.; 119 03 20 E...	464	50	glob. oz.	
			D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.	
			D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.	
12119	U.S.N.M.	5	D5526...	9 12 45 N.; 123 45 30 E...	805	52.3	gn. m., glob.	Few.
14210	U.S.N.M.	10+	D5529...	9 23 45 N.; 123 39 30 E...	441	53.	gy. m., glob.	Common.
14208	U.S.N.M.	5	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Few.
			D5569...	5 32 15 N.; 120 12 57 E...	330	52.3	fine. s., glob.	
			D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.	
12117	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E...	347	44	gy. m.	Rare.
			D5589...	4 12 10 N.; 118 38 08 E...	260	Co.	
12118	U.S.N.M.	10+	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.	Frequent.
			D5591...	4 11 48 N.; 118 38 20 E...	260	

TEXTULARIA CONICA d'Orbigny.

Plate 25, figs. 2a-c.

Textularia conica D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 143, pl. 1, figs. 19, 20.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 365, pl. 43, figs. 13, 14; pl. 113, figs. 1a, b.

This species is a common one on tropical coral reefs. Most of the specimens are from stations less than 40 fathoms (73 meters) in depth, the bottom temperature in the only case given being 75.7° F. (24.2° C.). Except for the two stations in the vicinity of Darvel Bay, Borneo, the species has occurred in the Philippine Archipelago. At the following six stations the species has occurred in more than 40 fathoms (73 meters): D5217, between Burias and Luzon, 105 fathoms (193 meters), bottom temperature 63.1° F. (17.2° C.); D5311, China Sea, vicinity of Hongkong, 88 fathoms (161 meters); D5315, China Sea, vicinity of Formosa, 148 fathoms (271 meters), bottom temperature 54.4° F. (12.4° C.); D5481, between Samar and Leyte, vicinity of Surigao Strait, 61 fathoms (112 meters); D5579, vicinity of Darvel Bay, Borneo, 175 fathoms (320 meters), bottom temperature 55.3° F. (15.1° C.); D5580, vicinity of Darvel Bay, Borneo, 162 fathoms (297 meters), bottom temperature 55.8° F. (13.2° C.).

Textularia conica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12115	U.S.N.M.	7	D5106...	14 23 55 N.; 120 32 33 E...	37	gy. m.....	Common.
12110	U.S.N.M.	8	D5134...	6 44 45 N.; 121 48 00 E...	25	fine s.....	Common.
14162	U.S.N.M.	1	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.....	Few.
14157	U.S.N.M.	2	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.....	Rare.
12113	U.S.N.M.	2	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Frequent.
12111	U.S.N.M.	10	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Frequent.
14161	U.S.N.M.	3	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.....	Few.
12114	U.S.N.M.	5	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Frequent.
12112	U.S.N.M.	5	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Common.
12116	U.S.N.M.	9						
			D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh.	Few.
14156	U.S.N.M.	2	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Few.
14159	U.S.N.M.	1	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh., g.....	Few.
14160.	U.S.N.M.	1	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fine s., Co.....	Few.
14158.	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co.....	Few

TEXTULARIA TROCHUS d'Orbigny.

Plate 25, figs. 1a-c.

Textularia trochus D'ORBIGNY, Mém. Soc. géol. France, vol. 4, 1840, p. 45, pl. 4, figs. 25, 26.—VANDEN BROECK, Ann. Soc. Belg. Micr., vol. 2, 1876, p. 132, pl. 3, figs. 11, 12.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 366, pl. 43, figs. 15, 16, 18, 19 (not fig. 17).—HERON-ALLEN and EARLAND, Trans. Zool. Soc., London, vol. 20, 1915, p. 630, pl. 47, fig. 28.

The records for this species show that it is most common in warm shallow water, especially in the southern part of the Archipelago. The stations range from 18 to 277 fathoms (33 to 507 meters), only two of them, however, being greater than 37 fathoms (68 meters). These are D5212, 108 fathoms (198 meters), east of Masbate Island, and D5576, 277 fathoms (507 meters), north of Tawi Tawi. At both of these stations specimens were rare. In all cases the bottom temperatures were relatively high—59.9° F. (15.5° C.), 53.3° F. (11.8° C.), and 75.7° F. (24.2° C.).

Textularia trochus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12040	U.S.N.M.	7	D5106...	14 23 55 N.; 120 32 33 E...	37	gy. m.....	Frequent.
12042								
12041	U.S.N.M.	3	D5134...	6 44 45 N.; 121 48 00 E...	25	fine s.....	Few.
12043	U.S.N.M.	5	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Few.
12047	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Few.
12044	U.S.N.M.	4	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Few.
14134	U.S.N.M.	1	D5212...	12 04 15 N.; 124 04 36 E...	108	59.9	gy. s., m.....	Few.
12046	U.S.N.M.	4	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Few.
14132	U.S.N.M.	2	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Few.
12045	U.S.N.M.	3	D5576...	Jolo Jolo, P. I.	Few.
14133	U.S.N.M.	1						

TEXTULARIA QUADRILATERA Schwager.

Plate 23, figs. 2a, b.

Textularia quadrilatera SCHWAGER, *Novara-Exped.*, Geol. Theil, vol. 2, 1866, p. 253, pl. 7, fig. 10.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 358, pl. 42, figs. 8-12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 24, figs. 42-44 (in text).

Bolivina quadrilatera WRIGHT, Proc. Roy. Irish Acad., ser. 3, vol. 1, 1891, p. 475.

This species has occurred at a large number of stations, and its range of depth is very consistent with the records from other portions of the North Pacific. The stations vary in depth from 230 fathoms to 1,599 fathoms (420 to 2,775 meters), the majority of these being between 500 and 800 fathoms (914 and 1,463 meters).

Both microspheric and megalospheric forms occur in considerable numbers and are easily distinguished at a glance, the character of the initial portion of the test being very different—that of the microspheric form narrow, tapering to a point, while the megalospheric form is blunt and broad at the initial end. The spine at the initial end is not always present.

Textularia quadrilatera—Material examined.

Cat. No.	Coll. of—	No. of specim.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14096	U.S.N.M.	1	D5114...	13 36 11 N.; 129 45 26 E..	340	fine. s.....	Rare.
14098	U.S.N.M.	1	D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m.....	Rare.
12131	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	53.8	gy. s., m.....	Few.
			D5203...	9 58 00 N.; 125 07 40 E..	773	52.9	gn. m.....	Rare.
			D5235...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.....
12133	U.S.N.M.	7	D5284...	13 42 05 N.; 120 30 45 E..	422	42.3	gy. m., glob.	Frequent.
14093	U.S.N.M.	3	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m., s.....	Rare.
12129	U.S.N.M.	5	D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s.....	Few.
			D5446...	12 43 64 N.; 124 59 18 E..	300	gn. m.....	Few.
14220	U.S.N.M.	1	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.....	Rare.
14092	U.S.N.M.	3	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn. m., glob.	Few.
			D5612...	0 38 00 N.; 121 45 40 E..	750
12134	U.S.N.M.	6	D5613...	0 42 00 S.; 121 44 00 E..	752	Co.....	Few.
			D5625...	0 07 00 S.; 127 28 00 E..	230	gy. m., fine. s.	Rare.
			D5636...	1 55 00 S.; 127 42 30 E..	1,262	gy. m., fine. s.	Rare.
14099	U.S.N.M.	1	D5637...	3 53 20 S.; 120 48 00 E..	700	fine. s.....	Rare.
12132	U.S.N.M.	4	D5638...	3 47 15 S.; 126 23 40 E..	517	fine. gy. s.....	Few.
14094	U.S.N.M.	1	D5652...	4 35 00 S.; 121 23 06 E..	525	41.2	gn. m.....	Rare.
14097	U.S.N.M.	1	D5660...	5 36 30 S.; 120 49 00 E..	692	39.2	gy. m., s.....	Rare.
			D5666...	2 54 30 S.; 118 47 09 E..	272	47.5	gn. m.....	Rare.
14916	U.S.N.M.	1	Nero 855.	15 26 30 N.; 121 47 15 E..	1,599	gn. m.....	Rare.

Genus BIGENERINA d'Orbigny, 1826.

BIGENERINA NODOSARIA d'Orbigny.

Plate 26, fig. 2.

Bigenerina nodosaria D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 261, pl. 11 figs. 9-11; Modèles, 1826, No. 57.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 369, pl. 44, figs. 14-18.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 27, figs. 46-48 (in text).

Textularia agglutinans, var. *nodosaria* PARKER and JONES, Philos. Trans. Roy. Soc., vol. 155, 1865, p. 371, pl. 15, fig. 25; pl. 17, fig. 80.

This is the most common species of the genus in the region. The test is usually of coarse material, in most cases light colored, and at a few stations is covered by a nearly black coating. There are numerous records of its occurrence in depths varying from 13 to 508 fathoms (25 to 930 meters) with a number of stations less than 300 fathoms (549 meters). Where given, the bottom temperature ranges from 44.3° to 76.3° F. (6.8° to 24.6° C.).

All but one of the stations, that from Sibuko Bay, Borneo, are in the Archipelago, or in the China Sea; off southern Luzon, off western Bohol; between Marinduque and Luzon; off northwestern Panay; off Palawan; between Cebu and Bohol; Jolo Sea; between Negros and Siquijor and north of Tawi Tawi.

Bigenerina nodosaria—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14241	U.S.N.M.	5	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m....	Few.
14207	U.S.N.M.	7	D5198...	9 40 50 N.; 123 39 45 E..	220	53.9	gn. m.....	Few.
			D5210...	11 49 55 N.; 124 28 05 E..	50	76.3	fine. gy. s....	Common.
			D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m....	Frequent.
			D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Few.
			D5281...	13 52 45 N.; 120 25 00 E..	201	50.4	dk. gy. s....	Frequent.
12980	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.
			D5301...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m., s....	Frequent.
12981	U.S.N.M.	4	D5311...	21 33 00 N.; 116 15 00 E..	88	cr. s., sh....	Common.
14239	U.S.N.M.	7	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh....	Frequent.
14206	U.S.N.M.	10+	D5315...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Frequent.
14240	U.S.N.M.	5	D5342...	10 56 55 N.; 119 17 24 E..	14-25	gy. m.....	Frequent.
			D5420...	9 49 35 N.; 123 45 00 E..	127	59.0	Few.
			D5423...	9 38 30 N.; 121 11 00 E..	508	49.8	gy. m., co. s..	Frequent.
12979	U.S.N.M.	1	D5430...	9 49 40 N.; 119 03 20 E..	464	50.0	glob. oz....	Rare.
14238	U.S.N.M.	2	D5449...	13 21 36 N.; 124 00 30 E..	300	Rare.
			D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob.	Few.
14237	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gn. m., s....	Few.
12978	U.S.N.M.	2	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Rare.
12976	U.S.N.M.	6	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s....	Common.
12977			Off Singapore.....	13	Common.

BIGENERINA ARENACEA Bagg.

Bigenerina arenacea BAGG, Proc. U. S. Nat. Mus., vol. 34, 1908, p. 132, pl. 5, figs. 4-6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 29, figs. 50a, b (in text).

A single specimen very much like that figured¹² was found in the dredgings made at D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

Bagg's specimens were from off the reefs of the Hawaiian Islands.

Bigenerina arenacea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12985	U.S.N.M.	1	D5236...	° ' " ° ' "		° F.		
				8 45 50 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Rare.

¹² Bull. 71, pt. 2, fig. 50.

BIGENERINA CAPREOLUS (d'Orbigny).

Plate 26, figs. 1a, b.

Valvulina capreolus D'ORBIGNY, ANN. Sci. Nat., vol. 2, 1826, p. 264, pl. 11, figs. 5, 6; Modèles, 1826, No. 59.

Grammostomum capreolus PARKER and JONES, ANN. Mag. Nat. Hist., ser. 3, vol. 11, 1863, p. 93.

Bigenerina capreolus H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 372, pl. 45, figs. 1-4.

Several excellent specimens were obtained from material dredged at D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature, 41.2° F. (5.1° C.). In depth this agrees well with the *Challenger* records for this species. Apparently it is the first record for the Pacific Ocean.

Bigenerina capreolus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12983 12984	U.S.N.M.	5	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fne. gy. s....	Few.

BIGENERINA PENNATULA (Batsch).

Plate 25, figs. 3a, b.

"*Orthoceratia pupa*" SOLDANI, Testaceographica, vol. 1, pt. 2, 1791, p. 99, pl. 108, figs. D. E. F.

Nautilus (Orthoceras) pennatula BATSCH, Conch. des Seesandes, 1791, No. 13, pl. 4, figs. 13, a-d.

Bigenerina pennatula H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 373, pl. 45, figs. 5-8.

A single well-developed, typical specimen was obtained from material dredged at D5318; China Sea, vicinity Formosa, 340 fathoms (556 meters), bottom temperature not recorded. This is apparently the first record for this species in the Pacific. This species is also recorded at D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

Bigenerina pennatula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11638	U.S.N.M.	1 1	D5236... D5318..	° ' " ° ' " 8 50 45 N.; 126 26 52 E... 21 32 00 N.; 117 46 00 E....	494 340	° F. 41.2	fne. gy. s.... s., br. Co....	Rare. Rare.

Genus *BOLIVINA* d'Orbigny, 1839.*BOLIVINA DILATATA* Reuss.

Plate 26, fig. 6.

Bolivina dilatata REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1850, p. 381, pl. 48, fig. 15.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 418, pl. 52, figs. 20, 21.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 33, fig. 54 (in text).

A single specimen, evidently of this species, was obtained from D5580, vicinity Darvel Bay, Borneo, 162 fathoms (297 meters), bottom temperature 55.8° F. (13.8° C.). It occurred also at D5115, Balayan Bay and Verde Island Passage, 340 fathoms (556 meters); bottom temperature not recorded.

Bolivina dilatata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13015	U.S.N.M.	2	D5115...	° ' " ° ' "	°	° F.		Rare.
13014	U.S.N.M.	1	D5580...	13 37 11 N.; 120 43 40 E... 4 52 45 N.; 119 06 45 E...	340 162	55.8	br. s., Co....	Rare.

BOLIVINA BEYRICHI Reuss.

Bolivina beyrichi REUSS, Zeitschr. deutsch. geol. Ges., vol. 3, 1851, p. 83, pl. 6, fig. 51.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 422, pl. 53, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 34, fig. 56 (in text).

Specimens of this species were obtained from two stations: D5126, Sulu Sea, vicinity southern Panay, 742 fathoms (1,357 meters), bottom temperature 49.5° F. (19.8° C.); and D5300, China Sea, 265 fathoms (485 meters), bottom temperature not given. These specimens were of the typical narrow form.

Bolivina beyrichi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12996	U.S.N.M.	3	D5126...	° ' " ° ' "	°	° F.		Rare.
14362	U.S.N.M.	1	D5300...	10 34 45 N.; 121 47 30 E... 20 31 00 N.; 115 49 00 E....	742 265	49.5	sft. gn. m... gy. m., s....	Rare.

BOLIVINA BEYRICHI Reuss, var. ALATA (Seguenza).

Vulvulina alata SEGUENZA, Atti Accad. Gioenia Sci. Nat., ser. 2, vol. 18, 1862, p. 115, pl. 2, figs. 5, 5a.

Bolivina alata EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 296, pl. 8, fig. 27.

Bolivina beyrichi, var. *alata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 422, pl. 53, figs. 2-4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 35, figs. 57a, b (in text).

Brady in the *Challenger* report records this variety from off the Philippines in 95 fathoms (174 meters). I have noted it in material dredged at D5133, Sulu Sea, off western Mindanao, 38 fathoms (69 meters); D5259, off northwestern Panay, 312 fathoms (571 meters); bottom temperature 49.3° F. (9.6° C.); D5300, China Sea, vicinity of southern Luzon, 265 fathoms (484 meters); and D5438, west coast of Luzon, Manila Bay to Lingayen Gulf, 297 fathoms (543 meters), bottom temperature 46.2° F. (7.8° C.). The typical form of the species was also found at station D5300.

Bolivina beyrichi, var. *alata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12999	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E ..	312	49.3	gy. m., glob.	Rare.
12998	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E ..	265	gy. m., s....	Rare.
12997	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E ..	297	46.2	gn. m.....	Rare.
			D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	gn. m., s....	

BOLIVINA ROBUSTA H. B. Brady.

Bolivina robusta H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 57; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 421, pl. 53, figs. 7-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 36, figs. 59, 60 (in text).

Bolivina acaulis EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 295, pl. 8, figs. 28-30.

This species has been found at several stations. From these few records it would seem that it is a fairly common species in the region, as the most of the washed material did not contain small forms. The specimens found were of the form without the apical spine, with the basal reentrants few but distant.

The localities include stations between Panay and Negros; off northwestern Panay; China Sea, off Hongkong and off southern Luzon; off Jolo; and to the southward in Buton Strait and Macassar Strait.

Bolivina robusta—Material examined.

Cat. No.	Coll. of—	No. of species.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14222	U. S. N. M.	1	D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	gn. m.	Rare.
11663	U. S. N. M.	1	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Rare.
14223	U. S. N. M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s.	Few.
11665	U. S. N. M.	1	D5301...	20 37 00 N.; 115 43 00 E..	208	59.5	gy. m., s.	Rare.
11667	U. S. N. M.	1	D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s.	Rare.
14229	U. S. N. M.	1	D5514...	8 32 42 N.; 123 58 36 E..	697	52.3	gn. m., s.	Rare.
14226	U. S. N. M.	1	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn. m., glob.	Rare.
14225	U. S. N. M.	1	D5546...	6 06 48 N.; 121 20 32 E..	138	58.3	ine. co. s.	Rare.
			D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s.	Rare.
14224	U. S. N. M.	1	D5575...	5 28 30 N.; 120 02 27 E..	315	52.3	Co., s.	Few.
14227	U. S. N. M.	1	D5637...	3 53 20 S.; 126 45 00 E..	700	gy. m.	Rare.
14228	U. S. N. M.	3	D5638...	3 47 15 S.; 126 23 40 E..	517	ine. gy. s.	Few.
11666	U. S. N. M.	7	D5648...	5 35 00 S.; 122 20 00 E..	559	39.2	gn. m.	Common.
			D5665...	4 27 00 S.; 118 44 00 E..	1,008	m.	Rare.
11668	U. S. N. M.	3	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.	Few.

BOLIVINA SCHWAGERIANA H. B. Brady.

Bolivina schwageriana H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 58; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 425, pl. 53, figs. 24, 25.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 38, figs. 63a, b (in text).

This species occurs mostly in the southern portion of the region at depths varying from 162 to 1,262 fathoms (297 to 2,309 meters); bottom temperature, where recorded, 47.5° to 55.8° F. (10.8° to 13.2° C.).

Material is typical, with the exception of slight traces of costae at the initial end in some cases.

Only one of the stations is in the Archipelago, off the east coast of Luzon; the others, Darvel Bay, Borneo; Gulf of Tomini, Celebes; off Bouro Island; Pitt Passage; and Macassar Strait.

This is a fairly large species for this genus and is strikingly ornamented. It is apparently very closely restricted in the general Australian and East Indian region.

Bolivina schwageriana—Material examined.

Cat. No.	Coll. of—	No. of species.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11669	U. S. N. M.	1	D5443...	12 43 05 N.; 125 01 00 E..	241	51.3	co. s., sh.	Rare.
11670	U. S. N. M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.	Rare.
11671	U. S. N. M.	1	D5601...	1 13 10 N.; 125 17 05 E..	765	s. glob., ptr.	Rare.
11672	U. S. N. M.	1	D5636...	1 55 00 S.; 127 42 30 E..	1,262	gy. m., ine. s.	Rare.
11673	U. S. N. M.	1	D5638...	3 47 15 S.; 126 23 40 E..	517	ine. gy. s.	Rare.
11674	U. S. N. M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.	Rare.

BOLIVINA NOBILIS Hantken.

Bolivina nobilis HANTKEN, Magy. Kir. földt. int. évkönyve, vol. 4, 1875 (1876), p. 56, pl. 15, fig. 4; Mitth. Jahrb. Ung. geol. Anstalt, vol. 4, 1875 (1881), p. 65, pl. 15, fig. 4.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 424, pl. 53, figs. 14, 15.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 39, figs. 64a, b (in text).

The only records of this species are D5133, Sulu Sea off western Mindanao, 38 fathoms (70 meters), bottom temperature not given; D5268, Verde Island Passage and Batangas Bay, 170 fathoms (312 meters), bottom temperature not given; D5580, vicinity of Darvel Bay, Borneo, 162 fathoms (297 meters), bottom temperature 55.8° F. (13.2° C.); and D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F. (10.8° C.).

This seems to be largely confined to the region of the Indo-Pacific, being found eastward to the Hawaiian Islands, southward to Australia, and northward to the Philippines. Chapman has found it as a fossil in the Tertiary of Victoria. Egger records it from several localities, but his figures do not seem to be this species.

Bolivina nobilis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11651	U.S.N.M.	1	D5133...	° ' " ° ' " Island off Panabutan Print, N. 32° E., 1.50 miles.	38	° F.	gn. m., s.....	Rare.
11652	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Rare.
11653	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.....	Rare.
11654	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Rare.
11655	U.S.N.M.	1						

BOLIVINA KARRERIANA H. B. Brady.

Plate 26, fig. 4.

Bolivina karreriana H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 58; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 424, pl. 53, figs. 19–21.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 40, figs. 65a, b (in text).

Specimens occur rarely at several stations, one off the east coast of Luzon; the others to the southward of the Archipelago; off Darvel Bay, Borneo; between Gillolo and Makyan Islands; south of Bouru Island; and Macassar Strait. Depths range from 23 to 517 fathoms (42 to 946 meters).

Bolivina karreriana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12992	U.S.N.M.	1	D5250...	° ' " ° ' " 7 05 07 N.; 125 39 45 E..	23	° F.	co. s.....	Rare.
14236	U.S.N.M.	1	D5443...	12 34 05 N.; 125 01 00 E..	241	51.3	co. s., sh.....	Rare.
12991	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E..	383	43.3	gn. m., s.....	Rare.
			D5530...	9 26 45 N.; 123 38 30 E..	Rare.
12-93	U.S.N.M.	1	D5622...	Makyan Island (N. E.) N. 65° W.	275	gy. m.....	Rare.
12988	U.S.N.M.	1	D5638...	3 45 15 S.; 126 23 40 E..	517	fine. gy. s.....	Rare.
12989	U.S.N.M.	4	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Few.

BOLIVINA KARRERIANA H. B. Brady, var. CARINATA Millett.

Plate 27, fig. 1.

Bolivina karreriana H. B. BRADY, var. *carinata* MILLETT, Journ. Roy. Micr. Soc., 1900, p. 546, pl. 4, fig. 8.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 41, figs. 66a, b (in text).

Specimens of this variety and almost identical with the figure given by Millett occurred at two stations: D5255, Gulf of Davao, 160 fathoms (290 meters), bottom temperature not recorded; D5278, China Sea, vicinity southern Luzon, 102 fathoms (187 meters), bottom temperature 59.6° F. (15.3° C.) and D5443, east coast of Luzon, 241 fathoms (441 meters), bottom temperature 51.3° F. (10.6° C.). The typical form of the species was not found at any of these stations.

Bolivina karreriana, var. *carinata*—Material examined.

Cat. No.	Coll. of—	No. of spec- imens.	Station.	Locality.	Depth in fath- oms.	Bot- tom tem- per- ature.	Character of bottom.	Abundance.
12990	} U.S.N.M.	3	D5255...	7 03 00 N.; 125 39 00 E..	160	sft. m.....	Few.
12994		1	D5278...	14 00 10 N.; 120 17 15 E..	102	59.6	fne.s., m., sh.	Rare.
12995 11805		2	D5443...	12 43 05 N.; 125 01 06 E..	241	51.3	ors. s., sh....	Rare.

BOLIVINA HANTKENIANA H. B. Brady.

Plate 27, fig. 2.

Bolivina hantkeniana H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 58; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 424, pl. 53, figs. 16–18.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 42, figs. 68a, b (in text).

From the records this is the most common species of the genus in the region. Material is in considerable numbers and well distributed. From a study of the material, both microspheric and megalospheric forms of this species are present. In depth the range is from 26 to 1,262 fathoms (21 to 2,309 meters). The bottom temperatures, where given, range from 43.3° to 55.8° F. (6.1° to 13.2° C.).

In the southern portion of the area specimens were found at D5592, vicinity of Sibuko Bay, Borneo, 305 fathoms (558 meters); D5636, Pitt Passage, 1,262 fathoms (2,309 meters); and D5666, Macassar Strait, 272 fathoms (498 meters).

This seems to be definitely an Indo-Pacific species, now known as far east as the Hawaiian Islands, southward to Tahiti and Australia, and northward to southern Japan. Its western limits are not well known. Heron-Allen and Earland did not find it in the Kerimba Archipelago material off southeastern Africa.

Bolivina hantkeniana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14232	U.S.N.M.	1	D5132...	° ' " ° ' " Island off Panabutan Point, N. 15° W., 0.30 mile.	26	° F.	gn. m., s....	Rare.
13006	U.S.N.M.	2	D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	gn. m., s....	Few.
13010	U.S.N.M.	3						
13011	U.S.N.M.	1						
13004 13009	U.S.N.M.	10	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Abundant.
14231	U.S.N.M.	8	D5269...	13 35 45 N.; 120 58 30 E..	220	fne. s., p....	Abundant.
13005	U.S.N.M.	2	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., sh....	Rare.
13008	U.S.N.M.	4	D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s.....	Few.
			D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Rare.
14234	U.S.N.M.	1	D5437...	10 02 45 N.; 125 05 33 E..	732	52.3	gn. m.....	Rare.
13012	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co....	Rare.
14235	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Rare.
			D5636...	1 55 00 S.; 127 42 3 E..	1,262	gy. m., fac.s	Rare.
14233	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Rare.
			H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob.	Frequent.

BOLIVINA AMYGDALAEFORMIS H. B. Brady.

Plate 26, fig. 3.

Bolivina amygdalaeformis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 59; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 426, pl. 53, figs. 28, 29.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 42, figs. 69a, b (in text).

Specimens occurred rarely at five stations: D5132, Sulu Sea, off western Mindanao, 38 fathoms (70 meters), bottom temperature not given; D5178, vicinity of Romblon, 73 fathoms (133 meters); D5311, China Sea, vicinity Hongkong, 88 fathoms (161 meters); D5381, Raygay Gulf, Luzon, 88 fathoms (161 meters); bottom temperature not recorded in the latter cases; and D5572, north of Tawi Tawi, 334 fathoms (611 meters), bottom temperature 52.3° F. (11.2° C.). The *Challenger* obtained two specimens in 95 fathoms (173 meters), off the Philippine station 209.

This is another of the Indo-Pacific species, and from the published records rather more circumscribed in its distribution than many others of this same fauna.

Bolivina amygdalaeformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13018 13020	U.S.N.M.	6	D5133...	° ' " ° ' " Island off Panabutan Point, N. 52° E., 1.50 miles.	38	° F.	gn. m., s....	Few.
13013	U.S.N.M.	2	D5178...	12 43 00 N.; 122 06 15 E..	73	fne. s.....	Rare.
14394	U.S.N.M.	1						
13016	U.S.N.M.	1	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., sh....	Rare.
13017	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 45 E..	88	co. s.....	Rare.
13019	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s., sh.....	Rare.

BOLIVINA SEMICOSTATA Cushman.

Bolivina costata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 426 pl. 53, figs. 26, 27 (not *B. costata* d'Orbigny, 1839).

Bolivina semicostata CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 43, figs 70a, b (in text).

A single specimen occurred at D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F (10.8° C.); and another from D5443, east coast of Luzon, San Bernardino Strait to San Miguel Bay, bottom temperature 51.3° F. (10.6° C.). Some of the specimens obtained by Brady from the *Challenger* material were from off New Guinea.

Little is known of the distribution of this seemingly Indo-Pacific species.

Bolivina semicostata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13000	U.S.N.M.	1	D5443...	° ' " ° ' "		° F.		
11675	U.S.N.M.	1	D5666...	12 43 05 N.; 125 01 00 E.. 2 54 30 S.; 118 47 00 E..	241 272	51.3 47.5	co. s., sh..... gn. m.....	Rare. Rare.

BOLIVINA SCULPTURATA Cushman.

Plate 27, figs. 3 a, 6.

Bolivina sculpturata CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 635, pl. 80, fig. 2.

Description.—Test flattened, rhomboid, chambers nearly flat on the surface, the borders raised with a rounded rim, as are also the areas above the sutures, proximal end rounded, distal end tending to become uniserial; aperture on the terminal face of the last-formed chamber, on the inner face of the previously-formed ones; wall composed of finely granular material, yellowish-brown, raised borders of a whitish calcareous material.

Length about 1 mm.

Type specimen.—Cat. No. 8505, U.S.N.M., from *Albatross* station D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters).

The peculiar ornamentation of this species, with its terminal aperture as well as the materials of its construction, makes this species a distinctive one.

Bolivina sculpturata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8505	U.S.N.M.	1	D5236...	° ' " ° ' "		° F.		
				8 50 45 N.; 126 26 52 E....	494	41.2	fine. gy. s....	Rare.

BOLIVINA SUBANGULARIS H. B. Brady.

Bolivina subangularis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 59; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 427, pl. 53, figs. 32, 33.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 45, figs. 72-73 (in text).

Several typical specimens were obtained at D5121, east coast of Mindoro, 108 fathoms (198 meters), bottom temperature not given; D5255, Gulf of Davao, 100 fathoms (183 meters), bottom temperature not given; D5443, east coast of Luzon, 241 fathoms (441 meters), bottom temperature 51.3° F. (10.6° C.); and D5637, Bouro Island (south) and vicinity, 700 fathoms (1,280 meters), bottom temperature not given. Brady records this species from *Challenger* station 209, in 95 fathoms (173 meters), off the Philippines.

Except for the other *Challenger* station from which Brady described this species off Raine Island, Torres Strait, in 155 fathoms (283 meters), little else is known of the species. Millett records it from the Malay Archipelago.

Bolivina subangularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12987	U.S.N.M.	1	D5121...	° ' " ° ' "	108	° F.	dk. gn. m...	Rare.
12986	U.S.N.M.	3	D5255...	13 27 20 N.; 121 17 45 E...	100	sft. m.....	Rare.
13210	U.S.N.M.	1	D5443...	7 03 00 N.; 125 39 00 E...	241	51.3	gn. m.....	Rare.
13067	U.S.N.M.	1	D5637...	12 43 05 N.; 125 01 00 E...	700	gy. m.....	Rare.
				3 53 20 S.; 126 48 00 E...				

BOLIVINA LIMBATA H. B. Brady.

Plate 19, fig. 5.

Bolivina limbata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 57; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 419, pl. 52, figs. 26-28.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 47, figs. 78a, b, c (in text).

Several specimens were obtained at D5201, Sogod Bay, southern Leyte Island, in 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5133, Sulu Sea, off western Mindanao, 38 fathoms (70 meters), bottom temperature not given; D5139, vicinity of Jolo, 20 fathoms (37 meters), bottom temperature not given; D5145, vicinity of Jolo, 23 fathoms (42 meters), bottom temperature not given; and D5572, north of Tawi Tawi, 334 fathoms (611 meters), bottom temperature, 52.3° F. (11.2° C.).

Bolivina limbata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
11643	U.S.N.M.	1	D5133...	° ' " ° ' " Island off Panabutan Pt., N. 52° E., 1.50 miles.	38	° F.	gn. m., s....	Rare.
			D5139...	6 06 00 N.; 121 02 30 E...	20	co. s.....	Rare.
11645	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Rare.
			D5145...	6 05 50 N.; 121 02 15 E...	23	co. s., sh....	
11647	U.S.N.M.	3	D5162...	5 10 00 N.; 119 47 30 E...	230	52.9	grs. s., brk..	Rare.
14205	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s., sh..	Rare.
11646	U.S.N.M.	4	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
11644	U.S.N.M.	2	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s....	Rare.
11648	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.....	Rare.

BOLIVINA PYGMAEA H. B. Brady.

Bolivina pygmaea H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 57;
Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 421, pl. 53, figs. 5, 6.

A single typical specimen was obtained from D5514, northern Mindanao and vicinity, 697 fathoms (1,274 meters), bottom temperature 52.3° F. (10.1° C.); and another from D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature, 47.5° F. (8.5° C.). Apparently this is the first record of the species in the Pacific. The specimens were practically identical with Brady's figure of this species in the *Challenger* Report.

Bolivina pygmaea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14203	U.S.N.M.	1	D5514...	° ' " ° ' " 8 32 42 N.; 123 58 36 E...	697	° F. 52.3	gn. m., s....	Rare.
14390	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Rare.

BOLIVINA PUNCTATA d'Orbigny.

Plate 26, fig. 5.

Bolivina punctata D'ORBIGNY, Voyage Amér. Mérid., vol. 5, pt. 5, 1839, "Foraminifères", p. 63, pl. 8, figs. 10-12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 417, pl. 52, figs. 18, 19.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 32, figs. 53a, b (in text).

This species has been noted at but six stations: Apra Bay, Guam; D5096, China Sea, off southern Luzon, 28 fathoms (51 meters), bottom temperature not recorded; D5100, China Sea, off southern Luzon, 35 fathoms (64 meters), bottom temperature not given; D5192, off northern Cebu, 32 fathoms (59 meters), no bottom temperature; D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); and D5300,

China Sea, vicinity southern Luzon, 265 fathoms (485 meters), bottom temperature not given. The apparent rarity of this common species may be due to the lack of fine material already referred to.

Bolivina punctata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11656	U.S.N.M.	3	D5096...	° ' " ° ' "		° F.		
11657	U.S.N.M.	1	D5100...	14 20 23 N.; 120 34 15 E...	28	gy. m., s., sh.	Rare.
11658	U.S.N.M.	1	D5192...	14 17 15 N.; 120 32 40 E...	35	gy. s.	Rare.
11659	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Rare.
11660	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.	Rare.
11660	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s.	Rare.
11661	U.S.N.M.	1	Apra Bay, Guam.....	Rare.

BOLIVINA COMPACTA Sidebottom.

Plate 26, fig. 7.

Bolivina robusta H. B. BRADY, var. *compacta* SIDEBOTTOM, Mem. Proc. Manchester Lit. Philos. Soc., vol. 49, No. 5, 1905, p. 15, pl. 3, fig. 7.

Bolivina compacta CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 36, figs. 58a, b (in text).

At two stations this species was noted—D5546, vicinity of Jolo Island, 138 fathoms (252 meters), bottom temperature 58.3° F. (14.6° C.); and D5601, Gulf of Tomini, Celebes, 765 fathoms (1,399 meters). It was previously recorded from a *Nero* station near the Philippines.

Bolivina compacta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14230	U.S.N.M.	1	D5546...	° ' " ° ' "		° F.		
			D5601...	6 06 48 N.; 121 20 32 E....	138	58.3	fne. co. s....	Rare.
14913	U.S.N.M.	1	Nero 855	1 13 10 N.; 125 17 05 E....	765	s., glob., ptr.	Rare.

BOLIVINA AENARIENSIS (Costa).

Brizalina aenariensis COSTA, Atti Acad. Pont., vol. 7, 1856, p. 297, pl. 15, figs. 1A, B.

Bolivina aenariensis H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 711, table; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 423, pl. 53, figs. 10, 11.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 292, pl. 37, fig. 8.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 44, figs. 71a, b (in text).

Brady records this species from a single station off the Philippines, 95 fathoms (173 meters). It has also occurred at D5133, Sulu Sea, off western Mindanao, 38 fathoms (70 meters), bottom tempera-

ture not recorded, and D5318, China Sea, vicinity Formosa, 340 fathoms (622 meters), bottom temperature not recorded.

Bolivina aenariensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13002	U.S.N.M.	1	D5133...	Island off Panabutan Pt., N. 52° E., 1.50 miles.	38	° F.	gn. m., s....	Rare.
13001	U.S.N.M.	1	D5318...	21° 32' N.; 117° 46' E.....	340	s., br. Co....	Rare.

BOLIVINA PUSILLA Schwager.

Bolivina pusilla SCHWAGER, *Novara-Exped.*, Geol. Theil., pt. 2, 1866, p. 254, pl. 7, fig. 101.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 41, fig. 67 (in text).

This is one of the species described by Schwager from Kar Nicobar, so many of which are found in this general region. I have noted it from the North Pacific and from off the Hawaiian Islands and near Japan in fairly deep water. The only Philippine material is from *Albatross* station D5655, Gulf of Boni, 608 fathoms (1,112 meters).

Bolivina pusilla—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11662	U.S.N.M.	1	D5655...	° ' " ° ' " 3 34 10 S.; 120 50 30 E...	608	° F. 39.2	gy. m., fine s	Rare.

Genus PLEUROSOMELLA Reuss, 1860.

PLEUROSOMELLA ALTERNANS Schwager.

Pleurostomella alternans SCHWAGER, *Novara-Exped.*, Geol. Theil., vol. 2, 1866, p. 238, pl. 6, figs. 79, 80.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 412, pl. 51, figs. 22, 23.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 50, fig. 81 (in text).

The only station at which this species occurred in the region is *Albatross* station D5236, in 494 fathoms (903 meters), in the Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.). The material was typical. The only other record for the North Pacific is an *Albatross* station, D2806, off the Galapagos Islands. This Philippine record, therefore, greatly extends its known distribution in this ocean.

The *Challenger* stations were from the Ki Islands off New Guinea in 129 fathoms (236 meters) and a station south of the Low Archipelago in 2,075 fathoms (3,795 meters). Sidebottom records a single

immature specimen from the east coast of Australia in 465 fathoms (850 meters), and Chapman had it from 1,050 fathoms (1,920 meters) off Funafuti, and records it from the Arabian Sea. It seems to be a rather rare species.

Pleurostomella alternans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13330 13331	} U.S.N.M.	2	D5236...	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Rare.
8 50 45 N.; 126 26 52 E..								

PLEUROSTOMELLA SUBNODOSA (Reuss).

Nodosaria nodosa REUSS (part), Verst. Böhm. Kreid., pt. 1, 1845, p. 28, pl. 13, fig. 22.

Dentalina subnodosa REUSS (part), Haidinger's Nat. Abhandl., vol. 4, 1850, p. 24, pl. 1, fig. 9.

Pleurostomella subnodosa REUSS, Sitz. Akad. Wiss. Wien., vol. 40, 1860, p. 204, pl. 8, figs. 2a, b.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 412, pl. 52, figs. 12–13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 51, fig. 82 (in text).

This species was only found at *Albatross* station D5236 in 494 fathoms (903 meters), Pacific Ocean off east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.). The material was typical. It occurred at *Nero* station 842, in 1,330 fathoms (2,433 meters), off the coast of Luzon.¹³

Pleurostomella subnodosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13332	U.S.N.M.	1 1	D5236... <i>Nero</i> 842	° ' " ° ' "	494	° F. 41.2	fne. gy. s....	Rare.
8 50 45 N.; 126 26 52 E..								

Subfamily VERNEULININAE.

Genus VERNEULINA d'Orbigny, 1840.

VERNEULINA POLYSTROPIA (Reuss).

Plate 32, fig. 1.

Bulimina polystropha REUSS, Verst. Böhm. Kreid., pt. 2, 1845, p. 109, pl. 24, fig. 53.

Verneulina polystropha PARKER and JONES, Introd. Foram., 1862, p. 311.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 386, pl. 47, figs. 15–17.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 53, figs. 85a, b (in text).

¹³ Cushman, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 51.

This species was common at two adjacent stations—D5445, 383 fathoms (699 meters), and D5446, 300 fathoms (549 meters), both off the east coast of Luzon. It also occurred in fewer numbers at D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.), and D5589, Sibuko Bay, Borneo, and vicinity, 260 fathoms (475 meters), bottom temperature 45.7° F. (7.6° C.).

Verneuilina polystropha—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13263	U.S.N.M.	4	D5236...	8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fne. gy. s.	Few.
13266	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s.	Rare.
13265	U.S.N.M.	10+	D5446...	12 43 51 N.; 124 59 18 E..	300	gn. m.	Frequent.
13264	U.S.N.M.	1	D5589...	4 12 10 N.; 118 38 08 E..	260	45.7	fne. gy. s., gy. m.	Rare.

VERNEULINA PROPINQUA H. B. Brady.

Verneuilina propinqua H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 387, pl. 47, figs. 8–12 (not 13, 14).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 53, figs. 86a, b (in text).

Specimens of this species have occurred at six stations, all but one in the region south of the Equator. Where they occur this species seems to be abundant, and numerous specimens show definitely the attachment at the apertural end. One specimen shows the abnormal twinned condition.

The stations are D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5446, off eastern Luzon, 300 fathoms (549 meters); D5612, north of Celebes, 750 fathoms (1,370 meters); D5613, north of Celebes, 752 fathoms (1,375 meters), bottom temperature not recorded in either case; D5637, vicinity Bouro Island, 700 fathoms (1,280 meters), bottom temperature not recorded; D5654, Gulf of Boni, 805 fathoms (1,470 meters), bottom temperature 38.3° F. (3.5° C.).

Verneuilina propinqua—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13282	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fne. gy. s.	Common.
13283	U.S.N.M.	1	D5446...	12 43 51 N.; 124 59 18 E..	300	gn. m.	Rare.
13280	} U.S.N.M.	5	D5612...	0 38 00 S.; 121 45 40 E..	750	Few.
13281								
13284	U.S.N.M.	5	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.	Few.
13279	U.S.N.M.	5	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.	Few.
14245	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E..	805	38.3	Rare.

VERNEUILINA BRADYI Cushman.

Plate 27, fig. 4.

Verneuilina pygmaea H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 385, pl. 47, figs. 4-7 (not *Bulimina pygmaea* Egger).

Verneuilina bradyi CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 54, figs. 87a, b (in text).—PEARCEY, Trans. Roy. Soc. Edinburgh, vol. 49, 1914, p. 1013.

This species occurs in few numbers, but at numerous stations, the depth ranging from 23 to 525 fathoms (42 to 927 meters), bottom temperatures, where recorded, 41.1° to 64.1° F. (5° to 17.8° C.).

This is a very widely distributed species, but does not seem to occur in shallow water, as noted in the Pacific material about Japan. The same thing is true in the Philippine dredgings.

Verneuilina bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13277	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s....	Rare.
13275	U.S.N.M.	1	D5242...	6 51 53 N.; 126 14 10 E..	191	64.1	sft. gy. m....	Rare.
13274	U.S.N.M.	1	D5246...	6 29 15 N.; 126 18 45 E..	Rare.
			D5449...	7 06 06 N.; 125 40 08 E..	23	Co., s.....	Rare.
14247	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Rare.
			D5586...	4 06 40 N.; 118 47 20 E..	347	44.0	gy. m.....	Rare.
13276	U.S.N.M.	1	D5625...	0 07 00 N.; 127 28 00 E..	230	gy. m., fne. s	Rare.
14246	U.S.N.M.	1	D5652...	4 35 00 S.; 121 23 06 E..	525	41.2	gn. m.....	Rare.
			Philippine expedition without number.	Rare.

VERNEUILINA SPINULOSA Reuss.

Plate 27, fig. 5.

Verneuilina spinulosa REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1850, p. 374, pl. 47, fig. 12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 334, pl. 47, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 55, figs. 88a, b (in text).

Typical specimens occur at the following localities: Vicinity of Romblon Island, China Sea; vicinity southern Luzon; vicinity of Darvel Bay, Borneo; east coast of Luzon, San Bernardino Strait to San Miguel Bay; Jolo Island and vicinity; north of Tawi Tawi; Sibuko Bay, Borneo, and vicinity; Gulf of Tomini, Celebes.

Verneuilina spinulosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13270	U.S.N.M.	9	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Common.
13272	U.S.N.M.	1	D5284...	13 42 05 N.; 120 30 45 E..	422	42.3	gy. m., glob.	Rare.
14234	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.....	Rare.
14242	U.S.N.M.	1	D5546...	6 06 48 N.; 121 20 32 E..	138	58.3	fne. co. s....	Rare.
13269	U.S.N.M.	2	D5672...	5 31 26 N.; 120 09 45 E..	334	52.3	s.....	Rare.
13268								
13267	U.S.N.M.	1	D5589...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co....	Rare.
14221	U.S.N.M.	1						
14243	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Rare.
			D5601...	1 13 10 N.; 125 17 05 E..	755	s., glob. ptr.	Rare.
			D5637...	3 53 20 S.; 126 43 00 E..	700	gy. m.....	Rare.

VERNEULINA TRIQUETRA (Münster.)

Textularia triquetra MÜNSTER, (in Römer), Neues Jahrb. für Min., 1838, p. 384, pl. 3, fig. 19.

Textularia (Verneuilina) triquetra PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 11, 1863, p. 92.

Verneuilina triquetra H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 383, pl. 47, figs. 18-20.

The only record for this species in the region is D5586, vicinity Sibuko Bay, Borneo, 347 fathoms (635 meters), bottom temperature 44° F. (6.4° C.).

Brady records it from the Pacific in 210 and 255 fathoms (384 and 466 meters) off Kandavu, Fiji Islands.

Verneuilina triquetra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5586...	° ' " ° ' "	347	° F. 44	gy. m.....	
				4 06 50 N.; 118 47 20 E...				

VERNEULINA AFFIXA Cushman.

Plate 27, fig. 6.

Verneuilina propinqua H. B. BRADY (part), Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 387, pl. 47, figs. 13, 14 (not 8-12).—Goës (part), Bull. Mus. Comp. Zool., vol. 29, 1896, p. 38.

Verneuilina affixa CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 56, figs. 90, 91 (in text).

The only specimens of this species which were noted in the material examined were from D5650, Gulf of Boni, 540 fathoms (988 meters), bottom temperature, 40.1° F. (3.5° C.). They were of the form with the pointed apex and broad base.

Verneuilina affixa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13278	U.S.N.M.	2	D5650...	° ' " ° ' "	540	° F. 40.1	gn. m.....	Rare.
				4 53 45 S.; 121 29 00 E...				

Genus VALVULINA d'Orbigny, 1826.

VALVULINA CONICA Parker and Jones.

Plate 27, fig. 7.

Valvulina triangularis PARKER and JONES, Ann. Mag. Nat. Hist., ser. 2, vol. 19, 1857, p. 295, pl. 11, figs. 15, 16 (not *Valvulina triangularis* d'Orbigny).

Valvulina triangularis, var. *conica* PARKER and JONES, Philos. Trans. Roy. Soc., vol. 155, 1865, p. 406, pl. 15, fig. 27.

Valvulina conica M. Sars, Vid. Selsk. Forh., 1868, p. 249.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 392, pl. 49, figs. 15, 16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 58, figs. 93a, b, c (in text).

This has proved to be one of the commonest species in the whole region, and well distributed records for its occurrence have been obtained at about 50 stations. These range in depth from 24 to 1,560 fathoms (44 to 2,853 meters), but the great majority of these are between 250 and 500 fathoms (457 and 914 meters). Bottom temperatures range from 38.2° to 62.4° F. (3.4° to 16.8° C.), corresponding more or less with range of depth.

Valvulina conica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14272	U.S.N.M.	1	D5110	13 59 20 N., 120 75 45 E.	135	59.0	dk. gy. m.	Rare.
14271	U.S.N.M.	1	D5219	13 21 09 N., 122 18 45 E.	530	50.8	gn. m.	Rare.
			D5241	6 50 45 N., 126 14 38 E.	215		silt. gy. m.	Rare.
14265	U.S.N.M.	6	D5282	13 53 00 N., 120 26 45 E.	248	47.4	dk. gy. s.	Few.
14253	U.S.N.M.	1	D5301	20 37 00 N., 115 43 00 E.	208	50.5	gy. m., s.	Rare.
14357	U.S.N.M.	1	D5369	13 48 00 N., 121 43 00 E.	106		bk. s.	Rare.
14270	U.S.N.M.	3	D5370	13 44 15 N., 121 42 30 E.	159	54.3	silt. m.	Few.
			D5381	13 14 15 N., 122 44 45 E.	88		co. s.	Rare.
			D5385	13 24 50 N., 123 03 70 E.	327	62.4	gy. m.	Frequent.
14267	U.S. MN.	6	D5428	9 13 00 N., 118 51 15 E.	1,105	49.7	gy. m.	Few.
			D5429	9 41 30 N., 118 50 22 E.	766		gn. m.	Rare.
14255	U.S.N.M.	3	D5445	12 44 42 N., 124 59 50 E.	383	44.3	gn. m., s.	Few.
14256	U.S.N.M.	1	D5446	12 43 51 N., 124 59 18 E.	300		gn. m.	Rare.
14261	U.S.N.M.	8	D5447	13 28 00 N., 123 46 18 E.	310	45.3	gn. m.	Few.
14260	U.S.N.M.	5	D5449	13 21 36 N., 124 00 30 E.	300			Few.
14251	U.S.N.M.	2	D5450					
13295	U.S.N.M.	6	D5460	13 32 30 N., 123 58 06 E.	565		gy. m.	Few.
14266	U.S.N.M.	10+	D5467	13 35 27 N., 123 37 18 E.	480		gn. m. (m. b.)	Common.
14269	U.S.N.M.	1	D5468	13 35 39 N., 123 40 28 E.	569		gy. m.	Few.
13294	U.S.N.M.	10+	D5469	13 36 48 N., 123 38 24 E.	500		gn. m. (net).	Frequent.
13291								
14254	U.S.N.M.	5	D5470	13 37 30 N., 123 41 09 E.	560		m.	Few.
14262	U.S.N.M.	6	D5495	9 06 30 N., 125 00 20 E.	976	52.3	gy. m.	Few.
14273	U.S.N.M.	1	D5526	9 12 45 N., 123 45 30 E.	895	52.3	gn. m., glob.	Few.
14268	U.S.N.M.	2	D5538	9 08 15 N., 123 15 20 E.	256	53.3	gn. m., s.	Few.
14257	U.S.N.M.	4	D5582	4 19 54 N., 118 58 38 E.	890	38.3	gy. m., fine. s.	Few.
			D5585	4 07 00 N., 118 49 54 E.	476	41.1	gy. m.	Few.
14264	U.S.N.M.	10+	D5586	4 06 50 N., 118 47 20 E.	347	44.0	gn. m.	Frequent.
13293	U.S.N.M.	3	D5591	4 11 48 N., 118 38 20 E.	260			Few.
14250	U.S.N.M.	1						
14263	U.S.N.M.	10+	D5612	0 33 00 S., 121 45 40 E.	750			Several.
13242	U.S.N.M.	2	D5613	0 42 00 S., 121 44 00 E.	752		gy. m.	Few.
14252	U.S.N.M.	2						
			D5619	0 35 00 S., 127 14 40 E.	435		fine. gy. s., m.	Few.
			D5630	0 56 30 S., 128 05 00 E.	569		eo. s., m.	
13296	U.S.N.M.	10+	D5637	3 53 20 S., 126 48 00 E.	700		gy. m.	Frequent.
			D5639	3 54 50 S., 123 27 20 E.	1,560		gy. m.	Few.
14249	U.S.N.M.	3	D5640	4 27 00 S., 122 55 40 E.	24		s., bkr. sh.	Few.
13290	U.S.N.M.	2	D5650	4 53 45 S., 121 29 00 E.	540	40.1	gn. m.	Few.
14259	U.S.N.M.	10+	D5654	3 42 00 S., 125 45 50 E.	895	38.3		Frequent.
14258	U.S.N.M.	10+	D5658	3 32 40 S., 120 31 30 E.	510	41.2	gy. m.	Frequent.
14248	U.S.N.M.	1	D5648	2 28 15 S., 118 49 00 E.	901	38.2	gy. m.	Rare.

VALVULINA FUSCA (Williamson).

Plate 28, figs. 1a, b.

Rotalina fusca WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 55. pl. 5, figs. 114, 115.

Valvulina fusca M. Sars, Vid. Selsk. Forh., 1868, p. 249.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 392, pl. 49, figs. 13, 14.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 59, figs. 94, 95a, b, c (in text).

Valvulina triangularis, var. *austriaca* PARKER and JONES, Introd. Foram., 1862, p. 311.

Valvulina austriaca H. B. BRADY, Trans. Linn. Soc. London, vol. 24, 1864, p. 472.

This species is much less abundant than the preceding. It occurs at fewer stations and in fewer numbers. The range in depth is from 26 to 890 fathoms (48 to 1,628 meters); bottom temperature, from 38.3° to 53.3° F. (3.5° to 11.8° C.).

Valvulina fusca—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13301	U.S.N.M.	1	D5132...	Island off Panabutan Point, N. 15° W. 0.3 mile.	26	° F.	gn. m., s.	Rare.
13299 13300	} U.S.N.M.	2	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s.	Few.
			D5381...	13 14 15 N.; 122 44 25 E...	83	co. s.	Rare.
			D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.	Rare.
13303	U.S.N.M.	1	D5446...	8 48 44 N.; 123 27 35 E...	Few.
			D5469...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s.	Rare.
13302	U.S.N.M.	1	D5523...	4 19 54 N.; 118 58 38 E...	890	38.3	gy. m., fine s.	Rare.
13304	U.S.N.M.	3	D5538...	4 11 48 N.; 118 38 20 E...	260	Rare.
			D5582...	4 43 50 N.; 121 23 24 E...	700	38.7	Rare.
13298	U.S.N.M.	3	D5591...	12 43 51 N.; 124 59 18 E...	300	gn. m.	Rare.
13297	U.S.N.M.	1	D5651...	13 36 48 S.; 123 38 24 E...	500	gn. m. (net.)	Rare.

Genus *CHRYSALIDINA* d'Orbigny, 1846.

CHRYSALIDINA DIMORPHA H. B. Brady.

Chrysalidina dimorpha H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 54; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 388, pl. 46, figs. 20, 21.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 60, figs. 96, 97a, b (in text).

A few specimens occurred at one station, D5152, off Tawi Tawi Group, Sulu Archipelago, 34 fathoms (62 meters), bottom temperature not recorded.

This is an Indo-Pacific species known to the eastward from the Hawaiian Islands, southward off Australia, and westward to the east coast of Africa, Madagascar, Mauritius, and the Kerimba Archipelago. Its northward range includes this record from the Sulu Sea and Brady's from Hongkong Harbor.

Chrysalidina dimorpha—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5152...	5 22 55 N.; 120 15 45 E...	34	° F.	wh. s.	Few.

Genus TRITAXIA Reuss, 1860.

TRITAXIA TRICARINATA (Reuss).

Plate 28, figs. 2a, b.

Textularia tricarinata REUSS, Verst. Böhm. Kreid., pt. 1, 1845, p. 39, pl. 8, fig. 60.*Tritaxia tricarinata* REUSS, Sitz. Akad. Wiss. Wien, vol. 40, 1860, p. 228, pl. 12, figs. 1, 2.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 389, pl. 49, figs. 8, 9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 61, figs. 98a, 99 (in text).

This species occurred at a number of stations in the region from the following localities: Vicinity of Pujada Bay; Gulf of Davao; China Sea, vicinity of Hongkong; China Sea, vicinity of Formosa; Mindoro Strait; Jolo Sea; and east coast of Luzon.

Some of the specimens are exceptionally fine and large.

The only *Challenger* record for this species is from off Raine Island, Torres Strait, 155 fathoms (283 meters).

Tritaxia tricarinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13209	U.S.N.M.	10+	D5244...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m., s.....	Common.
13208	U.S.N.M.	7	D5247...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.....	Common.
13207	U.S.N.M.	2	D5301...	6 52 05 N.; 126 14 15 E..	171	gy. m.....	Few.
13202	U.S.N.M.	3	D5318...	7 02 00 N.; 125 38 45 E..	135	m.....	Few.
13206								
13204	U.S.N.M.	1	D5333...	12 26 30 N.; 120 37 45 E..	310	73.8	s.....	Rare.
13203	U.S.N.M.	3	D5425...	9 37 45 N.; 121 11 00 E..	495	49.4	gy. m., co. s.	Few.
13205	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.(net)..	Rare.

TRITAXIA INDISCRETA H. B. Brady.

Plate 28, fig. 3.

Clavulina indiscreta H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 55.*Tritaxia indiscreta* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 389, pl. 49, figs. 10, 11.

A specimen which closely agrees with Brady's figures of this species, both in character of the test and sutures, occurred at D5423, Jolo Sea, 508 fathoms (930 meters), bottom temperature, 49.8° F. (9.8° C.).

Brady's specimens were from *Challenger* station 174, off Kandavu, Fiji Islands, 210 fathoms (384 meters).

Tritaxia indiscreta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14204	U.S.N.M.	1	D5423...	9 38 30 N.; 121 11 E.....	508	49.8	gy. m., co. s.	Rare.

Genus GAUDRYINA d'Orbigny, 1839.

GAUDRYINA SCABRA H. B. Brady.

Plate 28, fig. 5.

Gaudryina pupoides H. B. BRADY (not *G. pupoides* d'Orbigny, 1840), Ann. Mag. Nat. Hist., ser. 4, vol. 6, 1870, p. 300, pl. 8, fig. 5.

Gaudryina scabra H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 381, pl. 46, fig. 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 62, figs. 100a, b, 101 (in text).

Specimens of this species have occurred at numerous stations well scattered throughout the region. The character of the test varies much, especially in the material of the included sponge spicules, which are often entirely absent. In depth the stations vary from 260 to 940 fathoms (476 to 1,719 meters). The bottom temperatures, where recorded, vary from 36.7° to 44.0° F. (2.6° to 6.4° C.).

In the Archipelago the records are from the east coast of Luzon. Southward it occurred off Darvel Bay and Sibuko Bay, Borneo; north of Celebes; between Gillolo and Makyan Islands; and Macassar Strait.

Gaudryina scabra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundanc.
11774	U.S.N.M.	2	D5439...	15 58 15 N.; 119 40 20 E...	940	36.7	gn. m.....	Few.
12701	U.S.N.M.	2	D5446...	12 43 51 N.; 124 59 18 E...	300	gn. m.....	Few.
14275	U.S.N.M.	6	D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m.....	Few.
11680	U.S.N.M.	4	D5468...	13 35 39 N.; 123 40 28 E...	569	gn. m.(m. b.)	Rare.
14276	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Rare.
11679	U.S.N.M.	2	D5470...	13 37 30 N.; 123 41 09 E...	560	gn. m.....	Rare.
14279	U.S.N.M.	2	D5582...	4 19 54 N.; 118 58 38 E...	890	38.3	gy. m., fine s.	Rare.
14280	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.....	Rare.
11678	U.S.N.M.	6	D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.....	Few.
14277	U.S.N.M.	3	D5591...	4 11 48 N.; 118 38 20 E...	260	Few.
12702	U.S.N.M.	3	D5612...	0 38 00 S.; 121 45 40 E...	750	Few.
14278	U.S.N.M.	4	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.....	Few.
14274	U.S.N.M.	1	D5621...	0 15 00 N.; 127 24 35 E...	298	gy. and bk. s.	Rare.
11677	U.S.N.M.	7	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.....	Frequent.

GAUDRYINA FLINTII Cushman.

Plate 29, fig. 1.

Gaudryina subrotundata FLINT (not *G. subrotundata* Schwager, 1866), Rep. U. S. Nat. Mus., 1897 (1899), p. 287, pl. 33, fig. 1.

Gaudryina rugosa GOËS (not *G. rugosa* d'Orbigny, 1840), Bull. Mus. Comp. Zoöl., vol. 29, 1896, p. 39.

Gaudryina flintii CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 63, figs. 102a, b, c (in text).

Specimens of this species have occurred at stations in the following localities: Jolo Sea; west coast of Luzon; vicinity of Darvel Bay, Borneo; south of Patiente Strait; and Gulf of Boni.

The Philippine specimens belong to this species and are not unlike those figured by Flint, and is evidently of wide distribution.

Gaudryina flintii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12707	U.S.N.M.	2	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Rare.
			D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.	Rare.
12689	U.S.N.M.	1	D5582...	4 19 54 N.; 118 58 38 E...	890	38.3	gy. m., fine. s.	Rare.
12708	U.S.N.M.	2	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.	Few.
12688	U.S.N.M.	5	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Common.
12687	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3	Rare.

GAUDRYINA QUADRANGULARIS Bagg.

Plate 29, fig. 2.

Gaudryina quadrangularis BAGG, Proc. U. S. Nat. Mus., vol. 34, 1908, p. 133, pl. 5, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 64, figs. 103a, b (in text).

Several specimens of this species occurred at D5113, China Sea, off southern Luzon, 159 fathoms (290 meters), bottom temperature not recorded; D5381, Ragay Gulf, Luzon, 88 fathoms (161 meters), bottom temperature not recorded; D5469, east coast of Luzon, San Bernardino Strait to San Miguel Bay, 500 fathoms (914 meters), bottom temperature not recorded; D5569, north of Tawi Tawi, 303 fathoms (554 meters), bottom temperature 52.3° F. (10.1° C.); D5575, north of Tawi Tawi, 315 fathoms (579 meters), bottom temperature 52.3° F. (10.1° C.); the neighboring station, D5576, 277 fathoms (506 meters), bottom temperature 53.3° F. (11.8° C.); D5585, Sibuko Bay, Borneo, and vicinity, 476 fathoms (871 meters), bottom temperature 41.1° F. (5° C.).

Material has previously been known only from the vicinity of the Hawaiian Islands. It is a large species and not easily overlooked; therefore its distribution probably is not wide unless it has not previously been distinguished from other species.

Gaudryina quadrangularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12674	U.S.N.M.	1	D5113...	13 48 22 N.; 120 50 30 E...	159	dk. gn. m.	Rare.
12676	U.S.N.M.	2	D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.	Rare.
12677	U.S.N.M.	1	D5469...	13 36 48 N.; 123 33 24 E...	500	gn. m. (net) ..	Rare.
12675	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.	Rare.
14290	U.S.N.M.	2	D5575...	5 28 30 N.; 120 02 27 E...	315	52.3	co., s.	Rare.
12679	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.	Rare.
12678	U.S.N.M.	5	D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.	Few.

GAUDRYINA TRIANGULARIS Cushman.

Gaudryina triangularis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 65, figs. 104a, b, c (in text).

The only station at which this species occurred was D5212, east of Masbate Island, 108 fathoms (198 meters); bottom temperature, 59.9° F. (15.5° C.).

The previous records for this species are off the Hawaiian Islands and off the Bonin Islands.

Gaudryina triangularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5212...	12 04 15 N.; 124 01 36 E..	108	° F. 59.9	gy. s., m....

GAUDRYINA PAUPERCULA Cushman.

Plate 29, figs. 4, 5.

Gaudryina paupercula CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 66, figs. 106a, b (in text).

At the following 11 stations specimens occurred which agree well with the type of the species: D5172, vicinity of Jolo, 318 fathoms (582 meters), bottom temperature not recorded; D5424, Jolo Sea, 340 fathoms (622 meters), bottom temperature 50.4° F. (10.1° C.); D5425, Jolo Sea, 495 fathoms (906 meters), bottom temperature 49.4° F. (9.7° C.); D5526, between Siquijor and Bohol Islands, 805 fathoms (1,472 meters), bottom temperature 52.3° F. (10.1° C.); D5569, north of Tawi Tawi, 303 fathoms (554 meters), bottom temperature, 52.3° F. (10.1° C.); D5570, north of Tawi Tawi, 330 fathoms (604 meters), bottom temperature 52.3° F. (10.1° C.); D5571, north of Tawi Tawi, 340 fathoms (622 meters), bottom temperature 52.3° F. (10.1° C.); D5575, north of Tawi Tawi, 315 fathoms (580 meters), bottom temperature 52.3° F. (10.1° C.); D5582, Darvel Bay, Borneo, 890 fathoms (1,628 meters), bottom temperature 38.3° F. (3.5° C.); D5584, Sibuko Bay, Borneo, 292 fathoms (535 meters), bottom temperature 44.3° F. (6.8° C.); and D5639, Molucca Sea, 1,560 fathoms (2,853 meters), bottom temperature not recorded.

The material from the Philippines serves to more clearly fix this as a very definite species. The short, broad form, arenaceous walls with smooth surface, well finished, are very apparent. The color is usually gray or brownish, but at D5172 very typically shaped specimens were found with tests of calcareous sand and white in color.

Gaudryina pauperula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12686	U.S.N.M.	4	D5172...	6 03 15 N.; 120 35 30 E...	318	fne. s., sh...	Few.
12684	U.S.N.M.	8	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Frequent.
12685	U.S.N.M.	2	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Rare.
12712	U.S.N.M.	1	D5526...	9 12 45 N.; 123 45 39 E...	805	52.3	gn. m., glob.	Rare.
12682	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.....	Rare.
12680	U.S.N.M.	3	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fne. s., glob.	Rare.
12681		2	D5571...	5 30 45 N.; 120 07 57 E...	340	52.3	s., sh.....	Rare.
12713	U.S.N.M.	1	D5575...	5 28 30 N.; 120 02 27 E...	315	52.3	co. s.....	Rare.
12683	U.S.N.M.	1	D5582...	4 19 54 N.; 118 53 38 E...	890	38.3	gy. m., fne. s.	Rare.
12714	U.S.N.M.	1	D5584...	4 17 40 N.; 118 57 42 E...	292	44.3	fne. s., gn. m.	Rare.
14289	U.S.N.M.	1	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.....	Rare.

GAUDRYINA BRADYI Cushman.

Plate 29, fig. 3.

Gaudryina pupoides H. B. BRADY (not *G. pupoides* d'Orbigny), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 378, pl. 46, figs. 1-4.

Gaudryina bradyi CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 67, figs. 107a-c (in text).—PEARCEY, Trans. Roy. Soc. Edinburgh, vol. 49, 1914, p. 1014.

There are numerous stations for this species in the area varying in depth from 300 to 805 fathoms (549 to 1,472 meters); bottom temperature, where recorded, ranging from 41.2° to 56.4° F. (5.1° to 13.5° C.).

These stations include the following localities: Sogod Bay, southern Leyte; between Marinduque and Luzon; east coast of Mindanao; off northwestern Panay; eastern and southern Luzon; off Formosa; Palawan Passage; Jolo Sea; between Siquijor and Bohol Islands; Molucca Passage; and off Bouro Island. Pearcey, after examining *Challenger* material, affirms the separation of this species.

Gaudryina bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12699	U.S.N.M.	6	D5201...	10 10 00 N.; 125 04 15 E...	554	53.8	gy. s., m....	Rare.
12700			D5219...	13 21 00 N.; 122 18 45 E...	530	50.8	gn. m.....	Rare.
14285			D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Few.
	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
			D5284...	13 42 05 N.; 120 30 45 E...	422	42.3	gy. m., glob.	Rare.
12698	U.S.N.M.	4	D5318...	21 32 00 N.; 117 46 00 E...	340	s. br. Co....	Frequent.
14282	U.S.N.M.	2						
14283	U.S.N.M.	5	D5348...	10 57 45 N.; 118 58 15 E...	375	56.4	co. s.....	Few.
14286	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E...	730	46.0	co. s.....	Rare.
12697	U.S.N.M.	1	D5423...	9 38 30 N.; 121 11 00 E...	508	49.8	gy. m., co. s.	Rare.
			D5446...	12 43 64 N.; 121 59 18 E...	300	gn. m.....	Few.
14281	U.S.N.M.	1	D5468...	13 35 39 N.; 123 40 28 E...	509	gn. m.....	Rare.
14284	U.S.N.M.	2	D5526...	9 12 45 N.; 123 45 30 E...	805	52.3	gn. m., glob.	Few.
14287	U.S.N.M.	1	D5618...	0 37 00 N.; 127 15 00 E...	417	gy. m.....	Rare.
12711	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E...	517	fne. gy. s....	Rare.

GAUDRYINA BACCATA Schwager.

Plate 29, fig. 6.

Gaudryina baccata SCHWAGER, *Novara-Exped.*, Geol. Theil, pt. 2, 1866, p. 200, pl. 4, figs. 12a, b.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 379, pl. 46, figs. 8–11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 68, figs. 108a, b (in text).

Excellent specimens agreeing well with the figures given by Brady occurred at three stations: D5211, east of Masbate Island, 155 fathoms (283 meters), bottom temperature 56.6° F. (13.7° C.); D5586, vicinity Sibuko Bay, 347 fathoms (635 meters), bottom temperature 44.0° F. (6.4° C.); and D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

Brady recorded this species from a *Challenger* station, 206, in 2,100 fathoms (3,841 meters), between China and the Philippines.

Gaudryina baccata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13210	U.S.N.M.	1	D5211...	11 51 35 N.; 124 14 00 E..	155	56.6	gn. m., s....	Rare.
13219	} U.S.N.M.	3	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne.gy.s.....	Rare.
13220								
12706	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy.m.....	Rare.

GAUDRYINA CHILOSTOMA (Reuss).

Textilaria chilostoma REUSS, Zeitschr. deutsch. geol. Ges., vol. 4, 1852, p. 18.

Gaudryina chilostoma REUSS, Denkschr. Akad. Wiss. Wien, vol. 25, 1866, p. 120, pl. 1, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 69, figs. 109a, b (in text).

Gaudryina pupoides, var. *chilostoma* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 379, pl. 46, figs. 5, 6.

Gaudryina pupoides, var. *chilostomella* EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 278, pl. 7, fig. 6 (?).

Fine typical specimens of this species occurred at three stations: D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5586, vicinity Sibuko Bay, Borneo, 347 fathoms (635 meters), bottom temperature 44.0° F. (6.4° C.), and D5592, Sibuko Bay, Borneo, 305 fathoms (558 meters), bottom temperature 43.3° F. (6.1° C.). Both this and the preceding species occurring at the same station, D5586, may raise the question as to whether or not these two are really distinct species.

Gaudryina chilostoma—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12704	U.S.N.M.	2	D5236...	° ' " ° ' "		° F.		
12705	U.S.N.M.	2	D5586...	8 50 45 N.; 126 26 52 E..	494	41.2	fine.gy.s.....	Rare.
12703	U.S.N.M.	2	D5592...	4 06 50 N.; 118 47 20 E..	347	44.0	gy.m.....	Rare.
				4 12 44 N.; 118 27 44 E..	305	43.3	gn.m.....	Rare.

GAUDRYINA PSEUDOFILIFORMIS Cushman.

Plate 29, fig. 8.

Gaudryina filiformis H. B. BRADY (not *G. filiformis* Berthelin), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 380, pl. 46, figs. 12a-c.

Gaudryina pseudofiliformis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 70, figs. 111a, b (in text).

Specimens of this species occurred at two stations: D5318, China Sea, vicinity Formosa, 340 fathoms (622 meters), bottom temperature not recorded; and D5586, vicinity Sibuko Bay, Borneo, 347 fathoms (635 meters), bottom temperature 44.0° F. (6.4° C.).

Some of the specimens show very well the peculiar form of aperture figured.¹⁴

Gaudryina pseudofiliformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12673	U.S.N.M.	2	D5318...	° ' " ° ' "		° F.		
12672	U.S.N.M.	1	D5586...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Rare.
				4 06 50 N.; 118 47 20 E..	347	44.0	gy.m.....	Rare.

GAUDRYINA APICULARIS Cushman.

Plate 29, fig. 7.

Gaudryina siphonella H. B. BRADY (not *G. siphonella* Reuss, 1851), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 382, pl. 46, figs. 17-19.—FLINT, Ann. Rep. U. S. Nat. Mus., 1897 (1899), p. 288, pl. 34, fig. 2.

Gaudryina apicularis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 69, figs. 110a, b (in text).

This species occurred at station D5268, Verde Island Passage and Batangas Bay, 170 fathoms (311 meters), bottom temperature not given. It is even more like the figures in the *Challenger* Report than those given in Bulletin 71.

Gaudryina apicularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14389	U.S.N.M.	1	D5268...	° ' " ° ' "		° F.		
				13 42 00 N.; 120 57 15 E..	170	s., p.....	Rare.

¹⁴Bull. 71, pt. 2, p. 70, fig. 111a.

GAUDRYINA ATTENUATA Cushman.

Plate 30, fig. 4.

Gaudryina attenuata CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 636, pl. 80, fig. 3.

Description.—Test elongate, composed of numerous chambers, early portion triserial, triangular in cross section with sharp angles, biserial chambers numerous, nearly as high as broad, in later development each chamber running nearly or quite to the opposite side of the test; wall rough, composed of rough sand grains and sponge spicules roughly cemented, aperture at first textularian, becoming more nearly circular in the last-formed chambers; color light gray.

Length about 3 mm.

Type specimen.—(Cat. No. 8508, U.S.N.M.) from *Albatross* station D5259, off northwestern Panay, 312 fathoms (571 meters).

This species differs from *G. filiformis* Berthelin in its sharply triangular basal portion and in its more loosely biserial form, as well as the form of the aperture. It differs from *G. pseudofiliformis* Cushman in the aperture, the very rough surface, and more loosely biserial character. The apertural end tends to become more attenuate, but not to assume a clavuline character.

The species has been found at two other stations in the region as follows: D5586, Sibuko Bay, Borneo, 347 fathoms (635 meters); and D5592, the neighboring station, 305 fathoms (558 meters).

Gaudryina attenuata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.								
8508	} U.S.N.M.	5	D5259...	11 57 30 N.; 121 42 15 E...	312	° F. 49.3	gy. m., glob..	Few.								
12896									} U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.....	Rare.
13261																

GAUDRYINA ROBUSTA Cushman.

Plate 30, figs. 1a-c.

Gaudryina robusta CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 636, pl. 78, fig. 2.

Description.—Test large, stout, early portion sharply triangular, triserial, later chambers biserial, rounded; wall made up of fine sand grayish in color, with numerous black specks; surface somewhat rugose, sutures slightly depressed, sides of early portion somewhat concave; aperture textularian, in a rather deep reëntrant of the inner border of the chamber.

Length 4 mm.; breadth 2 mm.

Type specimen.—(Cat. No. 8509, U.S.N.M.) from *Albatross* station D5612, Gulf of Tomini, Celebes, 750 fathoms (1,371 meters).

This is a large species, very robustly built, the early portion concavely triangular, the latter portion very much rounded. The specimen figured shows but a few of the biserial chambers.

The species has also occurred at D5637, south of Bouro Island, 700 fathoms (1,280 meters), and D5670, Macassar Strait, 1,181 fathoms (2,160 meters). Bottom temperature was recorded in the last case only—38.2° F. (3.4° C.).

Gaudryina robusta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
8509 12691 11683 12390 11682	U.S.N.M.	4	D5612...	0 38 00 S.; 121 45 40 E...	750 ° F.	Few.
		3	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.	Few.
	U.S.N.M.	1	D5670...	1 19 00 S.; 118 43 00 E...	1,181	38.2	gy. m.	Rare.

Genus TRITAXILINA Cushman, 1911.

TRITAXILINA CAPERATA (H. B. Brady).

Plate 28, figs. 4a, b.

Clavulina caperata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 54.

Tritaxia caperata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 390, pl. 49, figs. 1-7.

Tritaxilina caperata CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 71, figs. 112-113 (in text).

Excellent specimens showing the typical adult condition as given in Brady's figures, and broken ones showing the radiating labyrinthian portion of the interior, occurred at five stations: D5244, vicinity of Pujada Bay, 171 fathoms (313 meters), bottom temperature not recorded; D5261, off southeastern Mindoro, 145 fathoms (265 meters), bottom temperature not recorded; D5241, Pujada Bay and vicinity, 215 fathoms (393 meters), bottom temperature not recorded; D5277, China Sea, vicinity southern Luzon, 80 fathoms (146 meters), bottom temperature 58.6° F. (14.7° C.); and D5318, China Sea, vicinity Formosa, 340 fathoms (622 meters), bottom temperature not recorded. Brady records the species off the Philippine Islands in 95 fathoms (173 meters); and off Kandavu, Fiji Islands, 250 fathoms (457 meters).

Tritaxilina caperata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13195 13194 11613 13191 13196 13192 13193	U.S.N.M.	2	D5241...	6 50 45 N.; 126 14 38 E...	215 ° F.	s't. gy. m.	Rare.
		7	D5244...	6 52 05 N.; 126 14 15 E...	171	gy. m.	Frequent.
	U.S.N.M.	3	D5261...	12 30 55 N.; 121 34 24 E...	145	s., m.	Few.
	U.S.N.M.	3	D5277...	13 53 55 N.; 120 13 45 E...	80	58.6	fine s.	Few.
	U.S.N.M.	5	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co.	Few.

Genus CLAVULINA d'Orbigny, 1826.

CLAVULINA COMMUNIS d'Orbigny.

Plate 31, fig. 1.

Clavulina communis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 268; Foram. Foss. Vienne, 1846, p. 196, pl. 12, figs. 1, 2.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 394, pl. 48, figs. 1-13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 72, figs. 115-117 (in text).

Verneuilina communis JONES and PARKER, Quart. Journ. Geol. Soc., vol. 16, 1860, p. 303.

This species is common and rather widely distributed in the region. The specimens vary in numerous details. The color of the wall varies from a chalky white to a dark gray, is usually smooth, and occasionally somewhat roughened. Some of the specimens show excellently the change from a triserial through the biserial to the final uniserial arrangement of the chambers. Occasional specimens show well the external tubular aperture. Although not definitely worked out there seem to be indications of both microspheric and megalospheric forms of the species in the material, from the comparative size of the earlier chambers and the considerable difference in size of the adults.

The species occurs at a large number of stations in the region varying in depth from 50 to 1,560 fathoms (91 to 2,853 meters), most of the stations being in the neighborhood of 300 fathoms (549 meters), bottom temperatures ranging from 38.2° to 76.3° F. (3.4° to 24.6° C.), and at the single shallow water station, 50 fathoms (91 meters), the temperature was 76.3° F. (24.6° C.). Specimens, however, were common at this last station.

Clavulina communis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.				Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				°	"	°	"				
14306	U.S.N.M.	1	D5172...	6	03 15 N.	120	35 30 E.	318	°F.	fne. s., sh....	Rare.
			D5210...	11	49 55 N.	124	28 05 E.	50	76.3.	fne. gy. s....	Common.
12962	U.S.N.M.	1	D5219...	13	21 00 N.	122	18 45 E.	530	50.8.	gn. m.	Rare.
14298	U.S.N.M.	9	D5236...	8	50 45 N.	126	26 52 E.	494	41.2.	fne. gy. s....	Common.
12954	U.S.N.M.	10+	D5259...	11	57 30 N.	121	42 15 E.	312	49.3.	gn. m., glob..	Common.
12960											
14291	U.S.N.M.	4	D5260...	12	25 35 N.	121	31 35 E.	234	51.4.	gn. m., s....	Few.
			D5300...	20	31 00 N.	115	49 00 E.	265	gy. m., s....	Rare.
12959	U.S.N.M.	9	D5318...	21	32 00 N.	117	46 00 E.	340	s., br. Co....	Common.
14304	U.S.N.M.	3	D5349...	10	54 00 N.	118	26 20 E.	730	40.6.	co. s.	Few.
14305	U.S.N.M.	1	D5371...	13	49 40 N.	121	40 15 E.	83	gn. m.(m.b.)	Rare.
14293	U.S.N.M.	9	D5439...	15	58 15 N.	119	40 20 E.	940	gn. n.	Common.
14292	U.S.N.M.	1	D5468...	13	35 39 N.	123	40 28 E.	569	gn. m.(m.b.)	Rare.
			D5469...	13	36 48 N.	123	38 24 E.	500	gn. m.(net)	Rare.
12958	U.S.N.M.	1	D5521...	8	47 00 N.	123	22 30 E.	221	53.3.	fne. s.	Rare.
14294	U.S.N.M.	1	D5523...	8	48 44 N.	123	27 35 E.	Rare.
14295	U.S.N.M.	5	D5526...	9	12 45 N.	123	45 30 E.	805	52.3.	gn. m., glob.	Few.
14307	U.S.N.M.	1	D5529...	9	23 45 N.	123	39 30 E.	441	53.0.	gy. m., glob.	Rare.
14301	U.S.N.M.	4	D5537...	9	11 00 N.	123	23 00 E.	254	53.5.	gn. m.	Few.
14303	U.S.N.M.	1	D5543...	8	47 15 N.	123	35 00 E.	162	54.5.	s.	Rare.
14296	U.S.N.M.	1	D5572...	5	31 26 N.	120	09 45 E.	334	52.3.	s.	Rare.
12956	U.S.N.M.	6	D5586...	4	06 50 N.	118	47 20 E.	347	44.0.	gy. m.	Few.
14216	U.S.N.M.	1	D5590...	4	10 50 N.	118	39 35 E.	310	44.3.	gy. m., s....	Rare.
			D5623...	0	16 30 N.	127	30 00 E.	272	fne. s.	Rare.
14300	U.S.N.M.	1	D5625...	0	07 00 N.	127	28 00 E.	230	gy. m., fne. s.	Rare.
14299	U.S.N.M.	1	D5637...	3	53 20 S.	126	48 00 E.	700	gy. m.	Rare.
14302	U.S.N.M.	9	D5639...	3	54 50 S.	123	27 20 E.	1,560	gn. m.	Common
14297	U.S.N.M.	3	D5659...	4	53 45 S.	121	29 00 E.	540	40.1.	gn. m.	Few.
			D5660...	5	36 30 S.	120	40 00 E.	692	39.2.	gy. m., s....	Rare.
12963	U.S.N.M.	2	D5670...	1	19 09 S.	118	43 00 E.	1,181	38.2.	gy. m.	Rare.
			H4898...	7	43 45 N.	122	03 45 E.	gy. m., glob..	Few.

CLAVULINA BRADYI Cushman.

Plate 31, fig. 4.

Clavulina cylindrica H. B. BRADY (not *C. cylindrica* d'Orbigny, 1826), Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 396, pl. 48, figs. 32-38.

Clavulina bradyi CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 73, figs. 118a, b, 119 (in text).

At two stations specimens referable to this species occurred: D5300, China Sea, vicinity southern Luzon, 265 fathoms (484 meters); and D5642, Buton Strait, 37 fathoms (68 meters); bottom temperatures not recorded in either case.

Numerous specimens of this species occurred at a number of stations ranging in depth from 37 to 375 fathoms (68 to 686 meters), with bottom temperatures from 52.3° to 58.6° F. (10.1° to 14.7° C.). A large series was obtained at some stations showing excellent apertures with one, two, or more teeth, yet the structure of the early portion of the test shows that they are not *Haplostiche dubia*, which they in some ways resemble.

Clavulina bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12966	U.S.N.M.	7	D5277...	13 56 55 N.; 120 13 45 E...	80	°F.	fine.s.....	Few.
12381						58.6		
12965	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E...	265	gy.m.,s.....	Rare.	
12382	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.....	Rare.
14310	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E...	Rare.
12383	U.S.N.M.	2	D5543...	8 47 15 N.; 123 35 09 E...	162	54.5	s.....	Rare.
12964	U.S.N.M.	1	D5539...	5 33 15 N.; 120 15 30 E...	303	52.3	co.s.....	Rare.
12380	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fine.s.,glob..	Rare.
14309	U.S.N.M.	3	D5642...	4 31 40 S.; 122 49 42 E...	37	gy.m.....	Few.

CLAVULINA ANGULARIS d'Orbigny.

Plate 31, fig. 3.

Clavulina angularis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 268, pl. 12, fig. 7.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 396, pl. 48, figs. 22-24.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 74, figs. 120-122a, b (in text).

Valvulina angularis JONES and PARKER, Quart. Journ. Geol. Soc., vol. 16, 1860, p. 305.

Valvulina triangularis D'ORBIGNY, forma *clavulina* Goës, Kongl. Svensk. Vet. Akad. Handl., vol. 19, No. 4, 1882, p. 86, pl. 11, figs. 387-389.

Excellent specimens of this species occurred in the following regions: Vicinity of Jolo; China Sea off southern Luzon; China Sea, vicinity Hongkong; north of Tawi Tawi Group; vicinity Sibuko Bay, Borneo; Jolo Island and vicinity.

Clavulina angularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.		Character of bottom.	Abundance.
						° F.			
13146	U.S.N.M.	5	D5172...	6 03 15 N.; 120 35 30 E..	318	fne. s., sh...	Few.	
13145	U.S.N.M.	10+	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s.....	Frequent.	
13144	U.S.N.M.	1	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh.....	Rare.	
13147	U.S.N.M.	1	D5549...	6 01 15 N.; 120 44 20 E..	263	52.3	s., glob., for..	Rare.	
12975	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Rare.	
13148	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Rare.	
12974	U.S.N.M.	2	D5569...	5 33 15 N.; 120 15 30 E..	303	52.3	co. s.....	Rare.	

CLAVULINA PARISIENSIS d'Orbigny.

Plate 19, fig. 4.

Clavulina parisiensis d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 268; Modèles, 1826, No. 66.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 395, pl. 48, figs. 14–18.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 75, figs. 123–124 (in text).

Valvulina parisiensis PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 3, vol. 16, 1865, pp. 29, 35, pl. 1, fig. 26.

Specimens occur at several stations varying in depth from 80 to 554 fathoms (146 to 1,012 meters), with bottom temperatures from 44.3° to 59.° F. (6.8° C. to 15° C.). The specimens are of the slender form figured by Brady (pl. 48, figs. 17, 18), but specimens also occur with very rough exterior like those on plate 48, figures 14 to 16. There do not seem to be any connecting variations between these two forms as far as the material available shows.

Clavulina parisiensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.		Character of bottom.	Abundance.
						° F.			
14218	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	Rare.	
12968	U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E..	80	58.6	fne. s.....	Few.	
13140	U.S.N.M.	1	D5291...	13 29 40 N.; 121 00 45 E..	173	51.5	fne. bk. s.....	Rare.	
12967	U.S.N.M.	10+	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	Co., s.....	Frequent.	
			D5420...	9 49 35 N.; 123 45 00 E..	127	59.0	Few.	
			D5423...	9 38 30 N.; 121 11 00 E..	508	49.8	gy. m., co. s.	Few.	
12758	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E..	Rare.	
			D5551...	5 54 48 N.; 120 44 24 E..	193	53.3	fne. s.....	Rare.	
14308	U.S.N.M.	1	D5565...	5 51 42 N.; 120 30 30 E..	243	52.3	s., ptr., sh...	Rare.	
13150	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s.....	Rare.	

CLAVULINA DIFFORMIS (H. B. Brady).

Plate 31, figs. 2a, b.

Clavulina angularis d'ORBIGNY, var. *difformis* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 396, pl. 48, figs. 25–31.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 637, pl. 48, figs. 20–22.

The only record in the material dredged by the *Albatross* on this cruise which contains this species came from Apra Bay, Guam.

The specimens are quadrilateral, with an aperture like that figured by Brady.

Brady describes this as a variety of *angularis* from a single locality, Nares Harbor, Admiralty Island, in 17 fathoms (30 meters), in this same faunal region. The occurrence of this same form at Guam would indicate that this is something more than a casual variation, and it is here raised to specific rank.

Heron-Allen and Earland record this from a single station in the Kerimba Archipelago, off the southeastern coast of Africa, in water of 4 fathoms (7 meters) or less on inshore reefs.

This is a peculiar distribution at these widely isolated points, and it will be interesting to see how they are later filled in by future work.

Clavulina difformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
1315	U.S.N.M.	1	Apra Bay, Guam.....	Rare.

CLAVULINA ROTUNDATA Cushman.

Plate 30, figs. 8a, b.

Clavulina rotundata CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 635, pl. 79, fig. 3.

Description.—Test of comparatively few chambers, tapering apertural end broadly rounded, early portion triserial and conical, followed by two or three chambers biserially arranged, and the last-formed chambers uniserial; wall thick, composed of fine sand grains with a dark reddish-brown cement; smooth; aperture terminal, large and rounded, depressed, without a neck; color reddish-brown.

Length 2.5–3 mm.

Type specimen.—Cat. No. 8506, U.S.N.M, from *Albatross* station D5512, off northern Mindanao, 445 fathoms (814 meters).

It is not uncommon at this and other stations.

This species is much shorter than most other species of the genus, broadly rounded at the apertural end, the wall thick and resembling that found in *Hormosina* and some species of *Trochammina*, *Ammodiscus*, etc. The aperture, instead of having a neck, shows rather the lack of it, the actual opening into the last-formed chamber being in a slight depression.

It has also occurred at *Albatross* station D5388, between Burias and Luzon, 226 fathoms (414 meters), bottom temperature 51.4° F. (10.7° C.); and D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.).

Clavulina rotundata—Material examined.

Cat. No.	Coll. of—	No. of specimens	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12973	U.S.N.M.	4	D5259...	° ' " ° ' "	312	° F. 49.3	gy. m., glob.	Few
12971	U.S.N.M.	1	D5388...	12 51 30 N.; 123 26 15 E...	226	51.4	sft. gn. m....	Rare.
8506 12972	U.S.N.M.	5	D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy., m., fne.s.	Few.

CLAVULINA PRIMAeva Cushman.

Plate 30, figs. 2a, b, 3, 5-7.

Clavulina primaeva CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 635, pl. 80, figs. 4, 5.

Description.—Test elongate, cylindrical, slender, composed of numerous chambers; early portion triserial, median portion biserial, terminal portion uniserial; chambers high; wall thick, light gray in color, smooth, sutures slightly depressed; aperture small, with a slight neck, more marked in the young stages.

Length about 3 mm.

Type specimen.—(Cat. No. 8507, U.S.N.M.) from *Albatross* station D5585, vicinity of Sibuko Bay, Borneo, 476 fathoms (871 meters).

This species is peculiar in the retarded taking on of the typical uniserial character of chamber arrangement and in the length of time that the biserial condition is held. In this it is distinctly primitive, much more so than in any other known species. The early triserial condition, on the other hand, is not much longer held than in some other species of the genus. The figure of a young specimen shows the typical clavuline neck even while the biserial condition is still held.

The species also occurred at D5526, between Siquijor and Bohol Islands, in 805 fathoms (1,472 meters).

Clavulina primaeva—Material examined.

Cat. No.	Coll. of—	No. of specimens	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12970	U.S.N.M.	1	D5526...	° ' " ° ' "	805	° F. 52.3	gn. m., glob.	Rare.
8507	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gn. m.	Rare.

Subfamily BULIMININAE.

Genus BULIMINA d'Orbigny, 1826.

BULIMINA ELEGANS d'Orbigny, var. EXILIS H. B. Brady.

Bulimina elegans D'ORBIGNY, var. *exilis* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 399, pl. 50, figs. 5, 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 82, figs. 135a-c (in text).

A single typical specimen of this variety occurred at D5301, China Sea, vicinity Hongkong, 208 fathoms (381 meters), bottom temperature 50.5° F. (10.2° C.).

Bulimina elegans, var. *exilis*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12571	U.S.N.M.	1	D5301...	20° 37' N.; 115° 43' E.....	208	° F. 15.5	gy. m.....	Rare.

BULIMINA MARGINATA d'Orbigny.

Bulimina marginata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 269, No. 4, pl. 12, figs. 10-12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 405, pl. 51, figs. 3-5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 83, figs. 136a, b (in text).

Bulimina pupoides, var. *marginata* WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 62, pl. 5, figs. 126, 127.

Bulimina presli, var. *marginata* PARKER and JONES, Philos. Trans., Roy. Soc., vol. 155, 1865, p. 372, pl. 15, fig. 10; pl. 17, fig. 70.

Apparently this species is not common in the region, occurring at but four stations: D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.); D5622, between Gillolo and Makyan Islands, 275 fathoms (503 meters); D5655, Gulf of Boni, 608 fathoms (1,112 meters), bottom temperature 39.2° F. (4.3° C.); and D5618, Molucca Passage, 417 fathoms (763 meters), bottom temperature not recorded.

Bulimina marginata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12664	U.S.N.M.	1	D5259...	11° 57' 30" N.; 121° 42' 15" E..	312	° F. 49.3	gy. m., glob..	Rare.
			D5618...	0° 37' 00" N.; 127° 15' 00" E..	417	gy. m.....	
			D5622...	0° 19' 20" N.; 127° 28' 30" E..	275	gy. m.....	
			D5655...	3° 34' 10" S.; 120° 50' 30" E..	608	39.2	g. m., fine s..	

BULIMINA INFLATA Seguenza.

Plate 31, fig. 6.

Bulimina inflata SEGUENZA, Atti Accad. Gioenia Sci. Nat., ser. 2, vol. 18, 1862, p. 109, pl. 1, fig. 10.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 406, pl. 51, figs. 10–13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 84, figs. 137a, b (in text).

This species is fairly common in the region and well distributed. The stations range in depth from 195 to 805 fathoms (356 to 1,472 meters), and the bottom temperatures from 40.6° to 52.8° F. (4.7° to 11.5° C.). In the southern portion of the area it was found at D5590, vicinity of Sibuko Bay, Borneo, 310 fathoms (567 meters); D5652, Gulf of Boni, 525 fathoms (927 meters); and D5666, Macassar Strait, 272 fathoms (498 meters).

Bulimina inflata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14315	U.S.N.M.	1	D5115...	13 37 11 N.; 120 43 40 E..	340	Rare.
14314	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s.....	Rare.
12548	U.S.N.M.	3	D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	gn. m.....	Few.
12554	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	Few.
			D5214...	12 25 18 N.; 123 37 15 E..	218	51.4	gn. m.....	Common.
			D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.....	Frequent.
			D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.....	Frequent.
12551	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Few.
14313	U.S.N.M.	1	D5260...	12 25 35 N.; 121 31 35 E..	234	51.4	gn. m., s.....	Rare.
14316	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E..	730	40.6	Co., s.....	Rare.
12547	U.S.N.M.	3	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn. m., glob.	Rare.
12552								
12555	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s.....	Rare.
14311	U.S.N.M.	1	D5565...	5 51 42 N.; 120 30 30 E..	243	52.3	s., ptr., s.....	Rare.
14312	U.S.N.M.	2	D5652...	4 35 00 S.; 121 23 06 E..	525	41.2	gn. m.....	Few.
12553	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Few.
14914	U.S.N.M.	1	H4898...	7 43 45 N.; 123 03 45 E..	221	gy. m., glob.	Rare.
12549	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Rare.

BULIMINA BUCHIANA d'Orbigny.

Bulimina buchiana D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 186, pl. 11, figs. 15–18.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 407, pl. 51, figs. 18, 19.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 85, figs. 138a, b, (in text).

Bulimina prestli, var. *buchiana* PARKER and JONES, Philos. Trans., Roy. Soc. vol. 155, 1865, p. 374, pl. 17, fig. 71.

This species occurred at only one station—D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature, 47.5° F. (8.6° C.).

Bulimina buchiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12578	U.S.N.M.	3	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Rare.

BULIMINA ACULEATA d'Orbigny.

Plate 31, fig. 5.

Polymorpha pineiformis SOLDANI, Testaceographica, vol. 1, pt. 2, 1791, p. 118, pl. 127, fig. 1; pl. 130, fig. vv.

Bulimina aculeata d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 269.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 406, pl. 51, figs. 7-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 286, figs. 139a, b (in text).

Bulimina pupoides, var. *spinulosa* WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 62, pl. 5, fig. 128.

Bulimina presli, var. *aculeata* PARKER and JONES, Introd. Foraminifera, 1862, Appendix, p. 311.

This is the most common species of the genus in the region, the stations ranging in depth from 83 to 834 fathoms (151 to 1,525 meters), and the bottom temperature from 39.2° to 53.6° F. (4° to 12° C.) In the southern portion of the region it occurred at D5606, Gulf of Tomini, Celebes, 834 fathoms (1,525 meters); D5613, north of Celebes, 752 fathoms (1,375 meters); and D5655, Gulf of Boni, 608 fathoms (1,112 meters) bottom temperature 39.2° F. (4° C.).

Bulimina aculeata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12541	U.S.N.M.	2	D5114...	13 36 11 N.; 120 45 26 E.	340	fne. s.....	Rare.
14327	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E.	393	43.7	gn.m., s.....	Rare.
14326	U.S.N.M.	10+	D5112...	13 38 42 N.; 120 47 25 E.	177	52.4	dk.gn.m.....	Frequent.
			D5122...	13 21 30 N.; 120 30 33 E.	220	gn.m.....	Frequent.
14319	U.S.N.M.	2	D5126...	10 34 45 N.; 121 47 30 E.	742	sft.gn.m.....	Few.
			D5185...	10 05 45 N.; 122 18 30 E.	638	49.8	gn.m.....	Occasional.
12544	U.S.N.M.	4	D5187...	9 16 45 N.; 123 21 15 E.	225	53.6	sft.gn.m.....	Frequent.
12543	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E.	554	52.8	gy.s., m.....	Frequent.
14324	U.S.N.M.	9	D5216...	12 52 00 N.; 123 23 30 E.	215	51.9	gn.m.....	Frequent.
			D5214...	12 25 18 N.; 123 37 15 E.	218	51.4	gn.m.....	Frequent.
12538	U.S.N.M.	1	D5219...	13 21 00 N.; 122 18 45 E.	530	50.8	gn.m.....	Rare.
14321	U.S.N.M.	7	D5222...	13 38 30 N.; 121 42 45 E.	195	52.8	gn.m.....	Frequent.
14325			D5238...	7 34 45 N.; 126 38 15 E.	380	43.0	gn.m.....	Few.
12542	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E.	312	49.3	gy.m., glob..	Few.
14317	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E.	730	40.6	Co., s.....	Rare.
14318	U.S.N.M.	10+	D5260...	12 25 35 N.; 121 31 35 E.	234	51.4	gn.m., s.....	Frequent.
12540	U.S.N.M.	5	D5371...	13 49 40 N.; 121 40 15 E.	83	gn.m.(m.b.)	Frequent.
14322	U.S.N.M.	1	D5512...	8 16 02 N.; 123 58 26 E.	445	52.8	gy.m., fne.s.	Rare.
14323	U.S.N.M.	1	D5606...	0 16 28 N.; 121 33 30 E.	834	gn.m.....	Rare.
14328	U.S.N.M.	1	D5618...	0 37 00 N.; 127 15 00 E.	417	gy.m.....	Rare.
12545	U.S.N.M.	3	D5613...	0 42 00 S.; 121 44 00 E.	752	gn.m.....	Few.
14320	U.S.N.M.	1	D5655...	3 34 10 S.; 120 50 30 E.	608	39.2	gy.m., fne.s.	Rare.
			H4898...	7 43 45 N.; 122 03 45 E.	221	gy.m., glob..	Few.

BULIMINA PUPOIDES d'Orbigny.

Plate 31, fig. 8.

Bulimina pupoides d'ORBIGNY, Foraminifera, Vienne, 1846, p. 185, pl. 11, figs. 13, 14.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 400, pl. 50, figs. 15a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 80, figs. 132a - c (in text).

This is a widely distributed species according to published records. It has been noted at 13 stations in the area ranging from 232 to 805

fathoms (425 to 1,472 meters). Bottom temperatures were recorded from 10 of these stations, ranging from 40.1° to 52.8° F. (3.5° to 11.5° C.). The stations are off southeastern Mindoro; off Jolo; north of Tawi Tawi; Gulf of Tomini, Celebes; and Sibuko Bay, Borneo.

Other North Pacific stations are about the Hawaiian Islands and Galapagos Islands.

Bulimina pupoides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13396	U.S.N.M.	6	D5115...	13 37 11 N.; 120 43 40 E..	340	Few.
12584	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy.s., m.....	Rare.
12580	U.S.N.M.	3}	D5260...	12 25 35 N.; 121 31 35 E..	234	51.4	gn.m., s.....	Rare.
14352	U.S.N.M.	3}						
14329	U.S.N.M.	8	D5300...	20 31 00 N.; 115 49 00 E..	265	gy.m., s.....	Common.
12579	U.S.N.M.	1	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn.m., glob..	Rare.
14351	U.S.N.M.	1	D5548...	6 00 20 N.; 120 45 35 E..	232	53.5	s., brk., sh...	Rare.
12581	U.S.N.M.	3	D5589...	4 12 10 N.; 118 38 08 E..	260	45.7	fne.gys., gy.m	Rare.
14353	U.S.N.M.	10	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn.m., s.....	Common.
12583	U.S.N.M.	1	D5591...	4 11 48 N.; 118 38 20 E..	310	44.3	gn.m., s.....	Rare.
14354	U.S.N.M.	10	D5613...	0 42 00 S.; 121 44 00 E..	752	gy.m.....	Common.
12582	U.S.N.M.	3	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn.m.....	Few.
14355	U.S.N.M.	10	D5586...	4 06 50 N.; 118 47 20 E..	347	44	gy.m.....	Common.
14356	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s.....	Rare.

BULIMINA PYRULA d'Orbigny.

Bulimina pyrula d'ORBIGNY, Foram. Foss. Vienne, 1846, p. 184, pl. 11, figs. 9, 10.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 399, pl. 50, figs. 7-10.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 78, figs. 126a-c (in text).

There are 13 stations from which material was obtained. These range in depth from 208 to 805 fathoms (381 to 1,472 meters), the




FIG. 1.—BULIMINA PYRULA d'ORBIGNY. $\times 50$. FROM ALBATROSS STATION D5586. a-c. THREE VIEWS OF SAME SPECIMEN.

average 465 fathoms (850 meters). The 9 stations at which bottom temperature was recorded give a range from 38.3° F. to 53.8° F. (3.5° C. to 12° C.) with an average of 47° F. (8.3° C.).

The distribution covers the following areas: Sogod Bay, southern Leyte; off northwestern Panay; east coast of Luzon; north of Tawi Tawi; Sibuko Bay, Borneo; Gulf of Tomini, Celebes; off Bouro Island; and Gulf of Boni.

The only other records in the North Pacific are those from the coast of Japan.

Bulimina pyrula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
14339	U.S.N.M.	6	D5201...	10 10 00 N.; 125 04 15 E...	554	53.8	gy. s., m.....	Common.
14338	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob..	Rare.
12563	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.....	Rare.
14337	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E...	500	gy. m., (m.b)	Rare.
12561	U.S.N.M.	1	D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy. m., fine. s.	Rare.
14336	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fine. s., glob..	Rare.
14335	U.S.N.M.	9	D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.....	Common.
12585	U.S.N.M.	1	D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fine. gy. s.....	Rare.
14334	U.S.N.M.	9	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.....	Common.
14332	U.S.N.M.	9	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.....	Common.
14333	U.S.N.M.	5	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Few.
12562	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E...	517	fine. gy. s.....	Rare.
14340	U.S.N.M.	10	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3	Common.



FIG. 2.—BULIMINA PYRULA D'ORBIGNY, VAR. PERVERSA CUSHMAN. $\times 50$. FROM ALBATROSS STATION D5654. a-c. THREE VIEWS OF SAME SPECIMEN.

BULIMINA PYRULA d'Orbigny, var. *PERVERSA*, new variety.

Description.—Variety differing from the typical in the form which in the variety is broadest at the base, the initial end of the test hardly visible, successive chambers extending back to the initial end, whole test tapering to the broadly rounded apertural end; surface smooth and otherwise like the typical form of the species.

Type specimen.—(Cat. No. 14342, U. S. Nat. Mus.) from Albatross station D5591, in 260 fathoms (475 meters), Sibuko Bay, Borneo. Other localities are Sogod Bay, southern Leyte; between Marinduque and Luzon; east coast of Mindanao; between Negros and Siquijor; and the Gulf of Boni. These stations range in depth from 195 to 805 fathoms (356 to 1,472 meters), average 367 fathoms (672 meters). Bottom temperature of seven stations range from 38.3° F. to 53.5° F. (3.5° C. to 11.8° C.), with the average 47.3° F. (8.4° C.).

This variety in this material is easily distinguished from the typical.



FIG. 3.—BULIMINA PYRULA D'ORBIGNY, VAR. SPINESCENS H. B. BRADY. $\times 50$. FROM ALBATROSS STATION D5586.

Bulimina pyrula, var. *perversa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12558	U.S.N.M.	1	D5115...	13 37 11 N.; 120 43 40 E.	340	Rare.
12559	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E.	554	52.8	gy. s., m.....	Rare.
14344	U.S.N.M.	2						
14346	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E.	195	52.8	gn. m.....	Rare.
13395	U.S.N.M.	1	D5589...	4 12 10 N.; 118 38 08 E.	260	45.7	fine.gy.s.,gy.m.	Rare.
14341	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E.	254	53.5	gn. m.....	Rare.
12556	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E.	310	44.3	gu. m., s.....	Rare.
14342	U.S.N.M.	1	D5591...	4 11 48 N.; 118 38 20 E.	260	Rare.
12557	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 09 E.	700	gy. m.....	Rare.
14345	U.S.N.M.	7	D5654...	3 42 00 S.; 120 45 50 E.	805	38.3	Common.
14343	U.S.N.M.	3	D5236...	8 50 45 N.; 126 26 52 E.	494	41.2	fine.gy.s.....	Few.

BULIMINA PYRULA d'Orbigny, var. SPINESCENS H. B. Brady.

Bulimina pyrula d'ORBIGNY, var. *spinescens* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 400, pl. 50, figs. 11, 12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 78, figs. 128, 129 (in text).

The only material which can be referred to this variety is a single specimen from *Albatross* station, D5586, Sibuko Bay, Borneo, 347 fathoms (635 meters), bottom temperature, 44° F. (6.4° C.); and D5638, south of Bouro Island, in 517 fathoms (946 meters). They are of the form which I figured from off Japan, and are somewhat different in shape than that figured by Brady.

Bulimina pyrula, var. *spinescens*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14347	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E....	347	44.0	gy. m.....	Rare.
12560	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E....	517	fine.gy.s.....	Rare.

BULIMINA OVATA d'Orbigny.

Bulimina ovata d'ORBIGNY, Foram. Foss. Vienne, 1846, p. 185, pl. 9, figs. 13, 14.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 400, pl. 50, figs. 13a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 77, figs. 125a-c (in text).

Specimens are rare at three stations all in 500 or more fathoms (914 meters); D5201, Sogod Bay, southern Leyte, in 554 fathoms (1,012 meters); D5469, east coast of Luzon, in 500 fathoms (914 meters); and D5487, between Leyte and Mindanao, in 732 fathoms (1,339 meters). They are also found at two other stations in 200 or more fathoms (366 meters); D5216, between Burias and Luzon, 215 fathoms (393 meters); and D5591, Sibuko Bay, Borneo, 260 fathoms (476 meters).

This is a short species, in which even the last-formed chamber extends back to near the initial end of the test.



FIG. 4.—BULIMINA OVATA d'ORBIGNY. X50. FROM ALBATROSS STATION 15469.


Bulimina ovata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13348	U.S.N.M.	3	D5201...	10 10 00 N.; 125 04 15 E....	554	53.8	gy. s., m....	Few.
12566	U.S.N.M.	1	D5216...	12 52 00 N.; 123 23 30 E....	215	51.9	gn. m.....	Rare.
14350	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E....	500	gn. m.....	Rare.
14319	U.S.N.M.	1	D5487...	10 02 45 N.; 125 05 33 E....	732	52.3	gn. m.....	Rare.
12569	U.S.N.M.	1	D5591...	4 11 48 N.; 118 38 20 E....	260	Rare.

BULIMINA AFFINIS d'Orbigny.

Bulimina affinis D'ORBIGNY, in De la Sagra, Hist. Fis. Nat. Pol. Cuba, 1839, "Foraminifères," p. 109, pl. 2, figs. 25, 26.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 400, pl. 50, figs. 14a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 79, fig. 130 (in text).

There are only four records for this species from all the material I have examined; D5114, Balayan Bay and Verde Island Passage, 340 fathoms (622 meters); D5214, east of Masbate Island, 218 fathoms (399 meters); D5526, between Siquijor and Bohol Islands, 805 fathoms (1,472 meters); and D5590, Sibuko Bay, Borneo, 310 fathoms (567 meters), at all stations. From other North Pacific records it seems to occur in deep water.



FIGS. 5, 6.—BULIMINA AFFINIS D'ORBIGNY. $\times 50$. FROM ALBATROSS STATION D5114. 5a, b. TWO VIEWS OF SAME SPECIMEN. 6, ANOTHER SPECIMEN OF IRREGULAR SHAPE.

Bulimina affinis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14358	U.S.N.M.	1	D5114...	13 36 11 N.; 120 45 26 E....	340	fine. s.....	Rare.
14359	U.S.N.M.	1	D5214...	12 25 18 N.; 123 37 15 E....	218	51.4	gn. m.....	Rare.
12564	U.S.N.M.	1	D5526...	9 12 45 N.; 123 45 30 E....	805	52.3	gn. m., glob.	Rare.
12565	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E....	310	44.3	gn. m., s....	Rare.

BULIMINA ELONGATA d'Orbigny.

Bulimina elongata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 187, pl. 11, figs. 19, 20.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 401, pl. 51, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 79, figs. 131 a-d (in text).

The records for the region are two in the deeper water to the southward: D5613, Gulf of Tomini, Celebes, in 752 fathoms (1,375 meters), and D5654, Gulf of Boni, in 805 fathoms (1,472 meters)

and two in the north, D5301, China Sea, vicinity Hongkong, 208 fathoms (381 meters), and D5529, between Siquijor and Bohol Islands, 441 fathoms (812 meters). It was rather rare at all stations. The only other records for the North Pacific are off Japan.

Bulimina elongata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12575	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E....	208	50.5	gy. m., s....	Rare.
12574	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E....	441	53.0	gy. m., glob..	Rare.
14360	U.S.N.M.	3	D5613...	0 42 00 S.; 121 44 00 E....	752	gy. m.....	Rare.
14361	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E....	805	38.3	Rare.

BULIMINA SUBAFFINIS, new species.

Description.—Test fusiform, apical end sharply pointed, apertural end with a rounded point, bases of chambers gradually appearing somewhat above the apex, the last-formed one extending from the apertural end to about three-fourths of the way back on the test, surface smooth, aperture a comma-shaped slit with a tooth.




FIG. 7.—*BULIMINA SUBAFFINIS*, NEW SPECIES. VIEWS FROM DIFFERENT SIDES, $\times 100$. FROM STATION D5201.

Length about 1 mm. or slightly more.

Type specimen.—Cat. No. 14330, U.S.N.M., from *Albatross* station D5201, from Sogod Bay, southern Leyte in 554 fathoms (1,012 meters), bottom temperature 53.8° F.

(12.1° C.). It is also rare at D5639, Molucca Sea, in 1,560 fathoms (2,853 meters).

This sharply pointed species seems very distinct from the other species of the genus in the region.

Bulimina subaffinis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14330	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E....	554	53.8	gy. s., m....	Rare.
14331	U.S.N.M.	1	D5639...	3 54 50 S.; 123 27 20 E....	1,560	gy. m.....	Rare.

Genus *BULIMINELLA* Cushman, 1911.*BULIMINELLA SUBTERES* (H. B. Brady).

Bulimina presli, var. *elegantissima* PARKER and JONES, Philos. Trans., Roy. Soc., vol. 155, 1865, p. 374, pl. 15, figs. 12-17.

Bulimina elegantissima (var.) H. B. BRADY, Ann. Mag. Nat. Hist., ser. 5, vol. 1, 1878, p. 436, pl. 21, fig. 12.

Bulimina subteres H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 55; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 403, pl. 50, figs. 17, 18.

Buliminella subteres CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 89, figs. 142a, b (in text).

A single, generally typical specimen of this interesting species was obtained from D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.). This is the second record for the North Pacific for this species.

Buliminella subteres—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12660	U.S.N.M.	1	D5259...	° ' " N.; ° ' " E..	312	° F. 49.3	gy. m., glob.	Rare.

BULIMINELLA CONTRARIA (Reuss).

Plate 31, fig. 9.

Rotalina contraria REUSS, Zeitschr. deutsch. geol. Ges., vol. 3, 1851, p. 76, pl. 5, fig. 37.

Bulimina contraria H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 409, pl. 54, figs. 18a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 89, fig. 143 (in text).

Specimens of this species, very typical in every way, occurred at *Albatross* station D5236 in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.); D5349, in 730 fathoms (1,335 meters), Palawan Passage, bottom temperature 40.6° F. (4.7° C.); and D5618, in 417 fathoms (763 meters), Molucca Passage.

Buliminella contraria—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12663	U.S.N.M.	1	D5236...	° ' " N.; ° ' " E..	494	° F. 41.2	fne. gy. s....	Rare.
12662	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E..	730	40.6	co. s	Rare.
12661	U.S.N.M.	1	D5618...	0 37 00 N.; 127 15 00 E..	417	gy. m.	Rare.

BULIMINELLA ELEGANTISSIMA (d'Orbigny).

Bulimina elegantissima D'ORBIGNY, Foram. Amér. Mérid., 1839, p. 51, pl. 7, figs. 13, 14.—WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 64, pl. 5, figs. 134, 135.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 402, pl. 50, figs. 20-22.

This widely distributed species has occurred but once in the Philippine material, from the station at which so many rare species have been recorded—D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters).

Buliminella elegantissima—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12659	U.S.N.M.	1	D5236...	8° 45' 50" N.; 126° 26' 52" E...	494	° F. 41.2	fine gy. s....	Rare.

Genus VIRGULINA d'Orbigny, 1826.**VIRGULINA SUBSQUAMOSA Egger.**

Virgulina subsquamosa EGGER, Neues Jahrb., 1857, p. 295, pl. 12, figs. 19-21.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 415, pl. 52, figs. 9-11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 92, figs. 146a-c (in text).

At six stations specimens occurred which are referable to this species; D5115, Verde Island Passage, 340 fathoms (622 meters); D5300, China Sea, vicinity southern Luzon, 265 fathoms (484 meters); D5315, China Sea, vicinity Formosa, 148 fathoms (271 meters), bottom temperature 54.4° F. (12.4° C.); D5529, between Siquijor and Bohol Islands, 441 fathoms (812 meters), bottom temperature 53.0° F. (11.7° C.); D5592, Sibuko Bay, Borneo, and vicinity, 305 fathoms (558 meters), bottom temperature 43.3° F. (6.1° C.); and D5638, south of Bouro Island, 517 fathoms (946 meters).

Virgulina subsquamosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13363	U.S.N.M.	1	D5115...	13° 37' 11" N.; 120° 43' 40" E...	340	° F.	Rare.
13365	U.S.N.M.	1	D5300...	20° 31' 00" N.; 115° 49' 00" E...	265	gy. m., s....	Rare.
13364	U.S.N.M.	1	D5315...	21° 40' 00" N.; 116° 58' 00" E...	148	54.4	s., sh.....	Rare.
13360	U.S.N.M.	1	D5529...	9° 23' 45" N.; 123° 39' 30" E...	441	53	gy. m., glob.	Rare.
13359	U.S.N.M.	2	D5592...	4° 12' 44" N.; 118° 27' 44" E...	305	43.3	gn. m.....	Rare.
13361	U.S.N.M.	1	D5638...	3° 47' 15" S.; 126° 23' 40" E...	517	fine gy. s....	Rare.

VIRGULINA PAUCILOCLATA H. B. Brady.

Virgulina pauciloculata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 414, pl. 52, figs. 4, 5.

The only station at which specimens referable to this species have occurred is D5526, between Siquijor and Bohol Islands, 805 fathoms (1,472 meters), bottom temperature 52.3° F. (10.1° C.).

This is a very rare species, described by Brady in the *Challenger* report from off New Guinea, Torres Strait, and the Ki Islands in 3–129 fathoms (5–236 meters). Little more is known of its distribution. Egger records it from several localities, but his figures, unless possibly in one case, do not represent the species. The one Philippine specimen was typical.

Virgulina pauciloculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14393	U.S.N.M.	1	D5526...	° ' " ° ' " 9 12 45 N.; 123 45 30 E....	805	° F. 52.3	gn. m., glob.	Rare.

VIRGULINA SCHREIBERSIANA Czjzek.

Virgulina schreibersiana CZJZEK, Haidinger's Nat. Abhandl., vol. 2, 1848, p. 147, pl. 13, figs. 18–21.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 414, pl. 52, figs. 1–3.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 291, pl. 37, fig. 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 94, figs. 148a, b (in text).

I have obtained specimens of this species from six stations as follows: H4898, Sulu Sea, off western Mindanao, 221 fathoms (425 meters), bottom temperature not recorded; D5112, China Sea, off southern Luzon, 177 fathoms (324 meters), bottom temperature 52.4° F. (10.2° C.); D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.); D5523, northern Mindanao and vicinity, depth and temperature not recorded; D5529, between Siquijor and Bohol Islands, 441 fathoms (812 meters), bottom temperature 53° F. (6.5° C.); and D5120, Balayan Bay and Verde Island Passage, 393 fathoms (719 meters), bottom temperature 43.7° F. (6.5° C.).

Virgulina schreibersiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13370	U.S.N.M.	1	D5112...	° ' " ° ' " 13 45 15 N.; 120 46 30 E....	177	° F.		Rare.
13369	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E....	393	43.7	gn. m., s.	Rare.
13366	} U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E....	312	49.3	gy. m., glob.	Rare.
13367			D5523...	8 48 44 N.; 123 27 35 E....				Rare.
13368	U.S.N.M.	1	D5529...	9 23 45 N.; 133 39 30 E....	441	53	gy. m., glob.	
14921	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E....	221		gy. m., glob.	Rare.

VIRGULINA CORNUTA Cushman.

Plate 32, figs. 3-6.

Virgulina cornuta CUSHMAN, Proc. U. S. Nat. Mus., vol. 44, 1913, p. 637, pl. 80, fig. 1.

Description.—Test pointed at the initial end, broadly rounded at the apertural end, whole test curved, irregularly biserial, last-formed chambers very tumid, aperture a comma-shaped opening with a surrounding raised portion; wall thin and translucent, smooth.

Length about 0.8 mm.

Type specimen.—(Cat. No. 8510, U.S.N.M.) from *Albatross* station D5284, China Sea, near southern Luzon, 422 fathoms (775 meters). It has also occurred in typical form at D5112, China Sea, off southern Luzon, 177 fathoms (324 meters); D5114, in 340 fathoms (620 meters), off Verde Island Passage; D5201, Sogod Bay, southern Leyte, 554 fathoms (1,012 meters); and D5259, in 312 fathoms (571 meters), off northwestern Panay.

This species is peculiar in its horn-like shape, its very tumid last-formed chambers, and tapering curved early portion.

Virgulina cornuta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14201	U.S.N.M.	1	D5112...	13 48 22 N.; 120 47 25 E...	177	52.4	dk. gn. m....	Rare.
13372	U.S.N.M.	1	D5114...	13 36 11 N.; 120 45 26 E...	340	fine s.....	Rare.
14202	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Rare.
14202	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
8510	U.S.N.M.	1	D5284...	13 42 05 N.; 120 30 45 E...	422	42.3	gy. m., glob.	Rare.

VIRGULINA SUBDEPRESSA H. B. Brady.

Virgulina subdepressa H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 416, pl. 52, figs. 14-17.—EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 291, pl. 8, fig. 103.—Goës, Bull. Mus. Comp. Zool., vol. 29, 1896, p. 47.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 93, fig. 147 (in text).

This species is only known from comparatively deep water, and therefore it is not expected in any quantity in the Philippine material. The only record is from *Albatross* station D5257, southern Mindanao, in 28 fathoms (52 meters) proving an exception to the general rule.

Virgulina subdepressa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13371	U.S.N.M.	1	D5257...	7 12 12 N.; 124 12 15 E....	28	m.....	Rare.

Subfamily CASSIDULININAE.

Genus CASSIDULINA d'Orbigny, 1826.

CASSIDULINA LAEVIGATA d'Orbigny.

Plate 31, fig. 7.

Cassidulina laevigata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 282, pl. 15, figs. 4, 5; Modèles, 1826, No. 41.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 428, pl. 54, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 96, figs. 150a, b (in text),

The only stations at which this species was noted were D5156, Sulu Archipelago, Tawi Tawi Group, 18 fathoms (33 meters), D5172, vicinity of Jolo, 318 fathoms (582 meters), and D5301, China Sea, vicinity Hongkong, 208 fathoms (381 meters), bottom temperature 50.5° F. (10.2° C.).

Cassidulina laevigata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13394	U.S.N.M.	1	D5156...	° ' " ° ' "	18	° F.	fne. s., sh....	Rare.
			D5172...	5 12 50 N.; 119 55 55 E....	318	fne. s., sh....	Rare.
14391	U.S.N.M.	1	D5301...	6 03 15 N.; 120 35 30 E....	208	50.5	gy. m.	Rare.
				20 37 00 N.; 115 43 00 E....				

CASSIDULINA SUBGLOBOSA H. B. Brady.

Plate 32, fig. 2.

Cassidulina subglobosa H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 60; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 430, pl. 54, figs. 17a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 98, fig. 152 (in text).

At numerous stations well scattered over the region this species has occurred as follows: Balayan Bay and Verde Island Passage; Pacific Ocean; east coast of Mindanao; east coast of Luzon between San Bernardino Strait to San Miguel Bay; vicinity of Sibuko Bay, Borneo; between Gillolo and Makyan Islands; vicinity of Bourou Island; and in the Gulf of Boni.

There seem to be two well-defined forms of this species. The first of these is a form with few visible chambers, and similar to that figured by Brady in the *Challenger* Report (pl. 54, fig. 17); the other is a form with more numerous chambers and the apertural end broadly truncate and flattened, the aperture occupying the central portion. The latter form is found only in the region to the south and may represent a different species. A similar form is found off the coast of New Zealand.

Cassidulina subglobosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14217	U.S.N.M.	2	D5114...	13 36 11 N.; 120 45 26 E...	340	° F.	fne. s.....	Rare.
			D5162...	5 10 00 N.; 119 47 30 E...	230	52.9	crs. s.....	Rare.
13387	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Few.
13381	U.S.N.M.	1	D5247...	7 02 00 N.; 125 38 45 E...	135		m.....	Rare.
13389	U.S.N.M.	3	D5333...	12 26 30 N.; 120 37 45 E...	310	73.8	s.....	Few.
13384	U.S.N.M.	1	D5443...	12 43 05 N.; 125 01 00 E...	241	51.3	co. s., sh....	Rare.
			D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s....	Rare.
13386	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Rare.
13388	U.S.N.M.	1	D5621...	0 15 00 N.; 127 24 35 E...	298		gy. and bk. s.	Rare.
13382	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 00 E...	700		gy. m.....	Rare.
13385	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Rare.
13383	U.S.N.M.	1	D5652...	4 35 00 S.; 121 23 06 E...	525	41.2	gn. m.....	Rare.

CASSIDULINA CRASSA d'Orbigny.

Cassidulina crassa D'ORBIGNY, Foram. Amér. Mérid., 1839, p. 56, pl. 7, figs. 18-20;
Foram. Foss. Vienne, 1846, p. 213, pl. 21, figs. 42, 43.—CUSHMAN, Bull. 71,
U. S. Nat. Mus., pt. 2, 1911, p. 97, fig. 151 (in text).

There is a single specimen from D5300, China Sea off southern Luzon, which is characteristic.

Cassidulina crassa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13390	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E...	265	° F.	gy. m., s....	Rare.

Genus EHRENBERGINA Reuss, 1850.

EHRENBERGINA SERRATA Reuss.

Ehrenbergina serrata REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1850, p. 377, pl. 43, fig. 7.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 434, pl. 55, figs. 2-7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 2, 1911, p. 101, figs. 155a-d (in text).

This species occurred at seven stations, most of them in the southern portion of the area. They were D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.); D5267, off Batangas Bay, 170 fathoms (300 meters), bottom temperature not recorded; D5300, China Sea, vicinity southern Luzon, 265 fathoms (484 meters), bottom temperature not recorded; D5576, north of Tawi Tawi Group, 277 fathoms (506 meters), bottom temperature 53.3° F. (11.8° C.); D5660, Flores Sea, 692 fathoms (1,266 meters), bottom temperature 39.2° F. (4.3° C.); D5637, off Bouro Island, 700 fathoms (1,280 meters); D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F. (8.6° C.).

Ehrenbergina serrata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13398	U.S.N.M.	3	D5300...	° ' " ° ' "		° F.		
13397	U.S.N.M.	3	D5660...	20 31 00 N.; 115 49 00 E...	265	gy. m., s....	Few.
14363	U.S.N.M.	1	D5637...	5 36 30 S.; 120 49 00 E...	692	39.2	gy. m., s....	Few.
13400	U.S.N.M.	1	D5259...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Rare.
13401	U.S.N.M.	1	D5267...	11 57 39 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
13402	U.S.N.M.	1	D5576...	13 42 20 N.; 120 58 25 E...	170	p., s., sh....	Rare.
14364	U.S.N.M.	2	D5666...	5 25 56 N.; 120 03 39 E...	277	53.3	s.....	Rare.
				2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Rare.

Family LAGENIDAE.

Subfamily LAGENINAE.

Genus LAGENA Walker and Boys, 1784.

LAGENA GLOBOSA (Montagu).

"*Serpula (Lagena) laevis globosa*" WALKER and BOYS, Test. Min., 1784, p. 3, pl. 1, fig. 8.

Vermiculum globosum MONTAGU, Test. Brit., 1803, p. 523.

Lagena globosa BROWN, Illus. Rec. Conch. Great Britain and Ireland, ed. 1, 1827, pl. 1, fig. 37; ed. 2, 1884, p. 126, pl. 56, fig. 37.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 452, pl. 56, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 3, pl. 4, fig. 2.

This species occurred at several stations in the area: D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte Island, bottom temperature 52.8° F. (11.5° C.); D5268, in 170 fathoms (311 meters), off Batangas Bay; and D5666, in 272 fathoms (498 meters), off Celebes, bottom temperature 47.5° F. (8.6° C.).

Lagena globosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13058	U.S.N.M.	1	D5201...	° ' " ° ' "		° F.		
13057	U.S.N.M.	1	D5268...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
13056	U.S.N.M.	1	D5666...	13 42 00 N.; 120 57 15 E...	170	p., s., sh....	Rare.
				2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Rare.

LAGENA LAEVIS (Montagu).

"*Serpula (Lagena) laevis oralis*" WALKER and BOYS, Test. Min., 1784, p. 3, pl. 1, fig. 9.

Vermiculum laeve MONTAGU, Test. Brit., 1803, p. 524.

Lagena laevis WILLIAMSON, Ann. Mag. Nat. Hist., ser. 2, vol. 1, 1848, p. 12, pl. 1, figs. 1, 2.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 455.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 5, pl. 1, fig. 3; pl. 38, fig. 5.

Occurrence of this species was noted in material from two *Albarr* stations in the Philippines—D5259, in 312 fathoms (571 meters),

off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); and D5537, in 254 fathoms (464 meters) between Negros and Siquijor, bottom temperature 53.5° F. (11.8° C.).

Lagena laevis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13049	U.S.N.M.	1	D5259...	° ' " ° ' "	312	° F. 49.3	gy. m., glob.	Rare.
13050	U.S.N.M.	1	D5537...	11 57 30 N.; 121 42 15 E.. 9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.....	Rare.

LAGENA LAEVIGATA (Reuss).

Fissurina laevigata REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1849, p. 366, pl. 46, fig. 1.

Lagena laevigata TERRIGI, Atti Accad. Pont. Nuovi Lincei, vol. 33, 1880, p. 177, pl. 1, fig. 6.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 473, pl. 114, figs. 8a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 7, pl. 2, fig. 1.

The only records for the species in the Philippines are *Albatross* stations, D5172, in 318 fathoms (582 meters), vicinity of Jolo, and D5201, Sogod Bay, southern Leyte in 554 fathoms (1,012 meters).

The species was found to be common in the deeper waters of the North Pacific and seems to be widely distributed elsewhere, but it was very rare in the material examined.

Lagena laevigata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13052	U.S.N.M.	1	D5172... D5201 ..	° ' " ° ' "	318 554	° F. 52.8	fne. s., sh.... gy. s., m....	Rare. Rare.

LAGENA CLAVATA (d'Orbigny).

Oolina clavata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 24, pl. 1, figs. 2, 3.

Lagena clavata MACKIE, Recreative Science, vol. 1, 1859, p. 148, fig. 13.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 456.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 9, pl. 2, fig. 3.

This species was noted from two stations, *Albatross* D5114, in 340 fathoms (622 meters), Verde Island Passage and D5537, in 254 fathoms (464 meters), between Negros and Siquijor, bottom temperature 53.5° F. (11.8° C.).

Lagena clavata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13021	U.S.N.M.	1	D5114...	° ' " ° ' "	340	° F.	fine.s.....	Rare.
13022	U.S.N.M.	1	D5537...	13 36 11 N.; 120 45 26 E.. 9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.....	Rare.

LAGENA GRACILLIMA (Seguenza).

Amphorina gracilis COSTA, Atti Acad. Pont., vol. 7, 1856, p. 121, pl. 11, fig. 11.

Amphorina gracillima SEGUENZA, Foram. mon. Mioc. Messina, 1862, p. 51, pl. 1, fig. 37.

Lagena gracillima JONES, PARKER, and H. B. BRADY, Pal. Soc. Mon., vol. 19, 1866, p. 45, pl. 1, figs. 36, 37.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 456, pl. 56, figs. 21, 22, 24–26 [19?, 20?, 23?, 27?, 28?].—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 306, pl. 53, fig. 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913 p. 11, pl. 1, fig. 4.

The only Philippine stations for which records were obtained of this species are *Albatross* D5201, Sogod Bay, southern Leyte in 554 fathoms (1,012 meters), and D5638, in 517 fathoms (946 meters), entrance to Tifu Bay, Bouro Island.

Lagena gracillima—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13051	U.S.N.M.	1	D5201... D5638...	° ' " ° ' "	554 517	° F. 52.8	gy.s., m..... fine.gy.s.....	Rare.

LAGENA ELONGATA (Ehrenberg).

Miliola elongata EHRENBERG, Bericht preuss. Akad. Wiss. Berlin, 1844, p. 274, 1845, p. 371; Mikrogeologie, 1854, pl. 25, fig. 1.

Lagena elongata TATE and BLAKE, Yorkshire Lias, 1876, p. 454, pl. 18, figs. 9, 9a.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 457, pl. 56, fig. 29.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 12, pl. 1, fig. 5.

Very typical specimens of this species have occurred in material from the following *Albatross* stations in the Philippines: D5178, in 78 fathoms (143 meters), near Romblon; D5259, in 312 fathoms (571 meters), off northeastern Panay, bottom temperature 49.3° F. (9.6° C.); and D5660, in 692 fathoms (1,266 meters), in Flores Sea.

Lagena elongata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13061	U.S.N.M.	2	D5178...	° ' " ° ' "	78	° F.	fine.s.....	Rare.
13059	U.S.N.M.	1	D5259...	12 43 00 N.; 122 06 15 E.. 11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob..	Rare.
13060	U.S.N.M.	1	D5660...	5 36 30 S.; 120 49 00 E..	692	39.2	gy. m., s.....	Rare.

LAGENA APICULATA (Reuss).

Plate 52, fig. 6.

Oolina apiculata REUSS, in Haidinger's Nat. Abhandl., vol. 4, 1850, p. 22, pl. 1, fig. 1.

Lagena apiculata REUSS, Sitz. Akad. Wiss. Wien, vol. 46, 1862 (1863), p. 319, pl. 1, figs. 4-8, 10, 11.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 453, pl. 56, figs. 4, 15-18.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 13.

A single specimen of this species was obtained from *Albatross* station D5648 in 559 fathoms (1,021 meters), Buton Strait, bottom temperature 39.2° F. (4.0° C.).

From the North Pacific records this species seems to be found in deep and cold waters.

Lagena apiculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
		1	D5648...	° ' " ° ' " E... 5 35 00 S.; 122 20 00 E...	559	° F.	gn. m.	Rare.

LAGENA HISPIDA Reuss.

"*Sphaerulae hispidae*" SOLDANI, Testaceographia, vol. 2, 1798, p. 53, pl. 17 figs. V, X.

Lagena hispida REUSS, Zeitschr. deutsch. geol. Ges., vol. 10, 1858, p. 43; Sitz. Akad. Wiss. Wien, vol. 46, 1863, p. 335, pl. 6, figs. 77-79.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 459, pl. 57, figs. 1-4; pl. 59, figs. 2, 5.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 307, pl. 53, fig. 8.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 13, pl. 4, figs. 4, 5; pl. 5, fig. 1.

This species has occurred at the following Philippine stations, *Albatross* D5120, in 393 fathoms (719 meters), Verde Island Passage: bottom temperature 43.7° F. (6.5° C.); D5192, in 32 fathoms (59 meters), off northern Cebu Island; D5257, in 158 fathoms (289 meters), eastern Illana Bay, southern Mindanao; and D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.).

Lagena hispida—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' " E... 13 45 30 N.; 120 30 15 E..	393	° F. 43.7	gn. m., s.	Rare.
13045	U. S. N. M.	1	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.	Rare.
13046	U. S. N. M.	1	D5257...	7 22 12 N.; 124 12 15 E..	158	m.	Rare.
13047	U. S. N. M.	2	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob..	Rare.

LAGENA HISPIDA Reuss, var. TUBULATA Sidebottom.

Lagena hispida REUSS, var. *tubulata* SIDEBOTTOM, Journ. Quekett Micr. Club, ser. 2, vol. 11, 1912, p. 385, pl. 15, figs. 3-5.

Description.—Test with globular body and long, cylindrical neck, surface covered with long delicate spines with a symmetrical arrangement of larger tubular ones, variously placed.

Length up to 0.5 mm.

Distribution.—The only station of the present series from which this variety was obtained is *Albatross* D5666, in 272 fathoms (498 meters), off Celebes, bottom temperature 47.5° F. (8.6° C.).

The specimens described by Sidebottom were from the South-west Pacific. This is the first record from the North Pacific.

Lagena hispida, var. *tubulata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5666...	2 54 30 S.; 118 47 00 E..	272	° F. 47.5	gn. m.	Rare.

LAGENA HEXAGONA (Williamson), var. SCALARIFORMIS Williamson.

Entosolenia squamosa (MONTAGU), var. *scalariformis* WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 13, pl. 1, fig. 30.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 17, pl. 6, fig. 4.

The only occurrence in the region as far as noted was *Albatross* station D5445, in 383 fathoms (699 meters), off Atalaya Point, Batag Island, off east coast of Luzon, bottom temperature 44.3° F. (6.8° C.).

Lagena hexagona, var. *scalariformis*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13063	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E..	383	° F. 44.3	gn. m., s.	Rare.

LAGENA STRIATA (d'Orbigny).

Oolina striata D'ORBIGNY, Foram. Amér. Mérid., 1839, p. 21, pl. 5, fig. 12.

Lagena striata REUSS, Sitz. Akad. Wiss. Wien, vol. 46, pt. 1, 1862 (1863), p. 327, pl. 3, figs. 44, 45; pl. 4, figs. 46, 47.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 460, pl. 57, figs. 22, 24.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 19, pl. 7, figs. 4, 5.

The typical form of this species was noted from the following stations: *Albatross* H4898, in 221 fathoms (405 meters), Sulu Sea off western Mindanao; D5277, in 80 fathoms (146 meters), China Sea,

etc., off southern Luzon, bottom temperature 58.6° F. (14.7° C.); D5318, China Sea, vicinity Formosa, 340 fathoms (946 meters), and D5638, in 517 fathoms (622 meters), entrance to Tifu Bay, Bouro Island.

Lagena striata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13036	U.S.N.M.	1	D5277...	° ' " ° ' " 13 56 55 N.; 120 13 45 E..	80	° F. 58.6	sh., p., s., fne. s.	Rare.
13037	U.S.N.M.	1	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Rare.
13035	U.S.N.M.	1	D5638... H4898...	3 47 15 S.; 126 23 40 E.. 7 43 45 N.; 122 03 45 E..	517 221	fne. gy. s.... gy. m., glob.	Rare.

LAGENA STRIATA (d'Orbigny), var. **HADINGERI** (Czjzek).

Oolina haidingeri CZJZEK, Haidinger's Nat. Abhandl., vol. 2, 1847, p. 138, pl. 12, figs. 1, 2.

Lagena haidingeri REUSS, Sitz. Akad. Wiss. Wien, vol. 46, pt. 1, 1862 (1863), p. 326, pl. 3, fig. 41.

Lagena striata H. B. BRADY (part), Rep. Voy. Challenger, Zoology, vol. 9, 1884, pl. 57, figs. 29, 30.

Lagena striata, var. *haidingeri* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913; p. 19, pl. 7, fig. 6.

A single station is all the record for this variety in the Philippines, *Albatross* station D5178, in 78 fathoms (143 meters), off Romblon.

Lagena striata, var. *haidingeri*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13040	U.S.N.M.	4	D5178...	° ' " ° ' " 12 43 00 N.; 122 06 15 E..	78	° F.	fne. s.....	Few.

LAGENA STRIATA (d'Orbigny), var. **STRUMOSA** (Reuss).

Lagena strumosa REUSS, Zeitschr. geol. Ges., 1858, p. 434; Sitz. Akad. Wiss. Wien, vol. 46, pt. 1, (1862) 1863, p. 328, pl. 4, fig. 49.

Lagena striata (d'ORBIGNY), var. *strumosa* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 20, pl. 7, figs. 7-10.

I have had this variety from *Albatross* station D5178, in 78 fathoms (143 meters), off Romblon; and D5187, in 225 fathoms (412 meters), Tanon Strait, east coast of Negros Island, bottom temperature 53.6° F. (12° C.).

In the North Pacific material from about Japan this variety was found frequently, but it has been seen but twice in the Philippine material.

Lagena striata, var. *strumosa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13038	U.S.N.M.	3	D5178...	° / ' " ° / ' "	73	° F.	fne. s.	Few.
13039	U.S.N.M.	1	D5187...	12 43 00 N.; 122 06 15 E... 9 16 45 N.; 123 21 15 E..	225	53.6	sft. gn. m...	Rare.

LAGENA SEMISTRIATA Williamson.

Lagena striata, var. *semistriata* WILLIAMSON, Ann. Mag. Nat. Hist., ser. 2, vol. 1, 1848, p. 14, pl. 1, figs. 9, 10.

Lagena semistriata JONES, PARKER, and H. B. BRADY, Mon. Foram. Crag, 1866, p. 34, pl. 4, fig. 6.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 465, pl. 57, figs. 14, 16, 17.

Description.—Test pyriform, widest near the base and gradually tapering to the aperture, which is at the end of a fairly long neck; surface smooth except the basal portion, which is ornamented by a series of short longitudinal costae running back to the lower end of the test.

Length about 0.75 mm.

Distribution.—This species was found in material from a single Philippine station, *Albatross* D5551, in 193 fathoms (353 meters) off Jolo, bottom temperature 53.3° F. (11.8° C.).

This species is not recorded in the rest of the North Pacific.

Lagena semistriata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13032	U.S.N.M.	1	D5551...	° / ' " ° / ' "	193	° F.	fne. s.	Rare.
				5 54 48 N.; 120 44 24 E..		53.3		

LAGENA SULCATA (Walker and Jacob).

Serpula (Lagena) sulcata WALKER and JACOB, Adam's Essays, Kanmacher's ed., 1798, p. 634, pl. 14, fig. 5.

Lagena sulcata PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 351.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 462, pl. 57, figs. 23, 26, 33, 34.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 22, pl. 9, fig. 2.

This species seems to have occurred more often than most others of the genus in the Philippine material. It was recorded from *Albatross* stations D5178, in 78 fathoms (143 meters), off Romblon; D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); D5420, in 127 fathoms (233 meters), off Cruz Point, Bohol Island, bottom temperature 59° F.

(15° C.); D5443, in 241 fathoms (441 meters), off Atalaya Point, Batag Island, east coast of Luzon, bottom temperature 51.3° F. (10.7° C.); D5638, in 517 fathoms (946 meters) at the entrance to Tifu Bay, Bouro Island; and D5666, in 272 fathoms (498 meters), off Celebes, bottom temperature 47.5° F. (8.6° C.).

These stations are well scattered over the area of dredging operations, and probably the species is much more common than indicated here.

Lagena sulcata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12184	U.S.N.M.	2	D5178...	12 43 00 N.; 122 06 15 E..	78	fne. s.....	Rare.
13026	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Rare.
13027	U.S.N.M.	1	D5420...	9 49 35 N.; 123 45 00 E..	127	59.0	Rare.
13024	U.S.N.M.	1	D5443...	12 43 05 N.; 125 01 00 E..	241	51.3	co. s., sh.....	Rare.
12183	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E..	517	fne. gy. s.....	Rare.
13025	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Rare.

LAGENA SULCATA (Walker and Jacob), var. APICULATA Cushman.

Lagena sulcata, apiculate forms, H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, pl. 58, figs. 4, 17 (?).

Lagena sulcata, var. *apiculata* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 23, pl. 9, figs. 3, 4.

Specimens of this variety occurred at the following three *Albatross* stations: D5133, in 38 fathoms (70 meters), Sulu Sea off western Mindanao; D5178, in 78 fathoms (143 meters) off Romblon; and D5277, China Sea off southern Luzon, in 80 fathoms (146 meters), bottom temperature 58.6° F. (14.7° C.).

Lagena sulcata, var. *apiculata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13028	U.S.N.M.	1	D5133...	Island off Panabutan Point; N. 52° E. 1.50 miles.	38	fne. s.....	Rare.
12815	U.S.N.M.	3	D5178...	12 43 00 N.; 122 06 15 E..	78	fne. s.....	Rare.
13029	U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E..	80	58.6	fne. s., m., sh.	Rare.

LAGENA ACUTICOSTA Reuss.

Lagena acuticosta REUSS, Sitz. Akad. Wiss. Wien, vol. 44, pt. 1, 1861 (1862), p., 305, pl. 1, fig. 4; vol. 46, pt. 1, 1862 (1863), p. 331, pl. 5, fig. 63.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 464, pl. 57, figs. 31, 32; pl. 58, figs. 20 (?), 21.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 23, pl. 8, figs. 9, 10; pl. 23, fig. 2.

Records of the occurrence of this species in material from three *Albatross* stations in the region are as follows: D5120, in 393 fathoms (719 meters), Verde Island Passage, bottom temperature 43.7° F. (6.5° C.); D5133, in 38 fathoms (70 meters), Sulu Sea off western Mindanao; and D5178, in 78 fathoms (143 meters), off Romblon.

Lagena acuticosta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
13044	U.S.N.M.	1	D5120... D5133...	° ' " ° ' " 13 45 30 N.; 120 30 15 E.. Island off Panabutan Point, N. 52° E., 1.50 miles.	393 38	° F. 43.7	gn. m. fine. s.	Rare.
13023	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	fine. s.	Rare.

LAGENA GRACILIS Williamson.

Lagena gracilis WILLIAMSON, Ann. Mag. Nat. Hist., ser. 2, vol. 1, 1848, p. 13, pl. 1, figs. 3, 4.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 464, pl. 58, figs. 19, 22–24.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 24, pl. 8, figs. 5, 6.

Three *Albatross* stations gave records for this species as follows: H4914, in 464 fathoms (848 meters), China Sea off southern Luzon, bottom temperature 46.5° F. (8° C.); D5514, in 697 fathoms (1,274 meters), off northern Mindanao, bottom temperature 52.3° F. (11.2° C.); and D5666, off Celebes, bottom temperature 47.5° F. (8.6° C.).

No specimens of this species were noted in the shallow warmer water of the Archipelago.

Lagena gracilis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13054	U.S.N.M.	1	D5514...	° ' " ° ' " 8 32 42 N.; 123 58 36 E.	697	° F. 52.3	gn. m., s....	Rare.
13055	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E.	272	47.5	gu. m.	Rare.
14912	U.S.N.M.	1	H4914...	13 38 05 N.; 120 33 00 E.	464	46.5	gy. m., s....	Rare.

LAGENA STAPHYLLEARIA (Schwager).

Fissurina staphyllearia SCHWAGER, *Novara*-Exped., Geol. Theil., pt. 2, 1866, p. 209, pl. 5, fig. 24.

Lagena staphyllearia H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 474, pl. 59, figs. 8–11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 31, pl. 17, fig. 3.

The two *Albatross* stations from which this species was obtained are: D5185, in 638 fathoms (1,167 meters), between Panay and Negros, bottom temperature 49.8° F. (9.8° C.); and D5236, in 494

fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.). It also occurred, as has been already noted, at *Nero* station 862 in 1,550 fathoms (2,834 meters) off the east coast of Luzon. Apparently it does not occur, at least in any numbers, in the shallow waters of this region from which most of the material came.

Lagena staphyllearia—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13031	U.S.N.M.	1	D5185...	° ' " ° ' "		° F.		
13033	U.S.N.M.	1	D5236...	10 05 45 N.; 122 18 30 E... 8 50 45 N.; 126 26 52 E...	638 494	49.8 41.2	gn. m..... fine. gy. s....	Rare. Rare.

LAGENA ALVEOLATA H. B. Brady, var. *SUBSTRIATA* H. B. Brady.

Lagena auriculata H. B. BRADY, var. *substriata* H. B. BRADY, Quart. Journ. Micr. Soc., vol. 21, 1881, p. 61.

Lagena alveolata H. B. BRADY, var. *substriata* H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 488, pl. 60, fig. 34.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 34, pl. 18, fig. 5.

The only record for this variety is *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.).

Lagena alveolata, var. *substriata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13043	U.S.N.M.	1	D5236...	° ' " ° ' "		° F.		
				8 50 45 N.; 126 26 52 E....	494	41.2	fine. gy. s....	Rare.

LAGENA MARGINATA Walker and Boys.

Serpula (Lagena) marginata WALKER and BOYS, Test. Min., 1784, p. 2, pl. 1, fig. 7.

Vermiculum marginatum MONTAGU, Test. Brit., 1803, p. 524.

Entosolenia marginata WILLIAMSON (part) Ann. Mag. Nat. Hist., ser. 2, vol. 1, 1848, p. 17, pl. 2, figs. 15-17.

Lagena marginata BROWN, Illus. Conch. Great Britain, 1827, pl. 1, figs. 30, 31.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 476, pl. 59, figs. 21-23.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 307, pl. 54, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 37, pl. 22, figs. 1-7.

Although this is a very common and widely distributed species, I have but two records of its occurrence in this region, *Albatross* stations D5301, in 208 fathoms (381 meters), China Sea, vicinity of

Hongkong, bottom temperature 50.5° F. (10.8° C.), and D5349, in 730 fathoms (1,335 meters), Palawan Passage, bottom temperature 40.6° F. (4.7° C.).

Lagena marginata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13041	U.S.N.M.	1	D5301...	° ' " ° ' "		° F.		
				20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.....	Rare.
13042	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E...	730	40.6	Co., s.....	Rare.

LAGENA LAGENOIDES (Williamson).

Entosolenia marginata WALKER and BOYS, var. *lagenoides* WILLIAMSON, Rec. Foram. Great Britain, 1858, p. 11, pl. 1, figs. 25, 26.

Lagena lagenoides REUSS, Sitz. Akad. Wiss. Wien, vol. 46, 1862, p. 324, pl. 2, figs. 27, 28.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 479, pl. 60, figs. 6, 8, 9, 12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 39, pl. 16, fig. 2.

I have record of the occurrence of this species at but a single Philippine station, D5301, in 208 fathoms (381 meters), China Sea, vicinity of Hongkong, bottom temperature 50.5° F. (10.8° C.).

Lagena lagenoides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13053	U.S.N.M.	1	D5301...	° ' " ° ' "		° F.		
				20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.....	Rare.

LAGENA FORMOSA Schwager.

Lagena formosa SCHWAGER (part), *Novara*-Exped., Geol. Theil., vol. 2, 1886, p. 207, pl. 4, figs. 19a, 19d (not 19b, 19c).—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 41, pl. 11, fig. 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 41, pl. 11, fig. 6.

The only record for this species in the Philippine material which I have examined is *Albatross* station D5660, in 692 fathoms (1,266 meters) in Flores Sea, bottom temperature 39.2° F. (4° C.).

The scarcity of this species in this material bears out the generalization already made that it is a species of deep water. Brady noted this fact in the *Challenger* report, and it was very certainly true of the material of the North Pacific which I have had at my disposal. It will be noted that the bottom temperature at this station is much lower than in most cases in the Philippine Archipelago and is practically an ocean bottom temperature for this depth.

Lagena formosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13062	U.S.N.M.	1	D5660...	° ' " ° ' " 5 36 30 S., 120 49 00 E...	692	° F. 39.2	gy. m., s....	Rare.

LAGENA BICARINATA (Terquem).

Fissurina bicarinata TERQUEM, Mém. Soc. Géol. France, ser. 3, vol. 2, 1882, p. 31, pl. 1 (9), figs. 24a, b.

Lagena bicarinata BALKWILL and MILLET, Journ. Micr., vol. 3, 1884, p. 82, pl. 2, fig. 4.

But one station is recorded for this species in the Philippine material I have examined, *Albatross* station H4914, in 464 fathoms (848 meters), China Sea off southern Luzon, bottom temperature 46.5°F. (8° C).

Lagena bicarinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14919	U.S.N.M.	1	H4914...	° ' " ° ' " 13 38 05 N.; 120 33 00 E...	464	° F. 46.5	gy. m., s....	Rare.

LAGENA LAMELLATA Sidebottom.

Lagena lamellata SIDEBOTTOM, Journ. Quekett Micr. Club, ser 2, vol. 11, 1912, p. 396, pl. 16, figs. 24, 25.

Description.—Test pyriform, tapering to a slender neck with spiral markings, wall compound, outer coating thin and flaky, resting upon fine tubular spines which in turn come out from an inner thickened wall with a polygonal structure when denuded of spines and seen in surface view or from within.

Diameter about 0.3 mm.

Distribution.—The only Philippine material I have seen is from *Albatross* station D5187, in 225 fathoms (412 meters), Tanon strait east coast of Negros, bottom temperature 53.6° F. (12° C). This is the first record for the North Pacific. The specimens described by Sidebottom were from the South-west Pacific, depths of 505 and 484 fathoms (923 and 885 meters).

This is a complicated test and the specimens showed nothing which will add to the characters as shown by Sidebottom.

Lagena lamellata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5187...	° ' " ° ' " 9 16 45 N.; 123 21 15 E...	225	° F. 53.6	sft. gn. m....	Rare.

Subfamily NODOSARIINAE.

Genus NODOSARIA d'Orbigny, 1826.

NODOSARIA (GLANDULINA) ROTUNDATA (Reuss).

Glandulina rotundata REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1849, p. 366, pl. 46, fig. 2.

Nodosaria (Glandulina) rotundata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 491, pl. 61, figs. 17-19.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 308, pl. 54, fig. 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 47, pl. 28, fig. 6.

Rare specimens of this species were found in material from the following three stations, which are widely scattered in the region: D5192, off northern Cebu Island, 32 fathoms (59 meters); D5318, China Sea, vicinity of Formosa, 340 fathoms (622 meters); and D5580 vicinity of Darvel Bay, Borneo, 162 fathoms (297 meters), bottom temperature 55.8° F. (13.2° C.).

Nodosaria rotundata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12253	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E...	32	° F.	gn. s.....	Rare.
13354	U.S.N.M.	1	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co....	Rare.
12256	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br.s., Co.....	Rare.

NODOSARIA (GLANDULINA) LAEVIGATA d'Orbigny.

Plate 33, fig. 1.

"Cornu Hammonis erectum globosius" PLANCUS, Conch. Min., 1739, p. 16, pl. 13, fig. 1.

Nodosaria (Glandulina) laevigata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 252, pl. 10, figs. 1-3.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 490, pl. 61, figs. 20-22.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 308, pl. 55, fig. 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 47, pl. 24, figs. 1, 2.

Glandulina laevigata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 29, pl. 1, figs. 4, 5.

At 11 stations in the area specimens of this species were obtained, most of the specimens being without spines at the apical end. The stations ranged in depth from 32 to 752 fathoms (59 to 1,375 meters), the average depth being 315 fathoms (576 meters). Where bottom temperatures were given, they ranged from 41.2° F. to 56.4° F. (5.1° C. to 13.5° C.), the average temperature being 50.5° F. (10.8° C.).

Nodosaria laevigata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
15278	U.S.N.M.	1	D5178...	12 93 00 N.;	122 06 15 E...	78	° F.	fne. s.	Rare.
12197	U.S.N.M.	1	D5192...	11 09 15 N.;	123 50 00 E...	32	gn. s.	Rare.
12191	U.S.N.M.	1	D5216...	12 52 00 N.;	123 23 30 E...	215	51.9	gn. m.	Rare.
12198	U.S.N.M.	1	D5219...	13 21 00 N.;	122 18 45 E...	530	50.8	gn. m.	Rare.
12195	U.S.N.M.	1	D5220...	13 38 00 N.;	121 58 00 E...	50	sft. gn. m.	Rare.
12194 13065	U.S.N.M.	2	D5236...	8 50 45 N.;	126 26 52 E...	494	41.2	fne. gy. s.	Rare.
12196	U.S.N.M.	1	D5300...	20 31 00 N.;	115 49 00 E...	265	gy. m., s.	Rare.
12193	U.S.N.M.	1	D5318...	21 32 00 N.;	117 46 00 E...	340	s., br. Co.	Rare.
12192	U.S.N.M.	2	D5348...	10 57 45 N.;	118 38 15 E...	375	56.4	Co., s.	Rare.
12190	U.S.N.M.	1	D5571...	5 30 45 N.;	120 07 57 E...	340	52.3	s., sh.	Rare.
15279	U.S.N.M.	1	D5613...	0 42 00 S.;	121 44 00 E...	752	gy. m.	Rare.

NODOSARIA (GLANDULINA) LAEVIGATA d'Orbigny, var. STRIATULA Cushman.

Plate 33, fig. 2.

Nodosaria (Glandulina) laevigata D'ORBIGNY, var. *striatula* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 653.

Description.—Variety differing from the typical in having the surface ornamented with numerous very fine longitudinal costae.

Distribution.—Type specimen (No. 5107, U. S. N. M.) from *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.).

Brady figures a striate specimen of this species in the *Challenger* Report.

Nodosaria laevigata, var. *striatula*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
5107	U.S.N.M.	1	D5236...	8 50 45 N.;	126 26 52 E....	494	° F. 41.2	fne. gy. s.	Rare.

NODOSARIA SIMPLEX Silvestri.

Nodosaria simplex SILVESTRI, Atti Accad. Gioenia, Catania, ser. 3, vol. 7, 1872, p. 95, pl. 11, figs. 268–272.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 496, pl. 62, figs. 4, 5, and 6?.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 309, pl. 55, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 49, pl. 28, fig. 5.

The only record for this species in all the material examined from the region is off Singapore in 13 fathoms (23 meters), U.S.N.M. No. 15661. It was typical, however.

The *Challenger* specimens were from off the Ki Islands, 129 fathoms (236 meters), and off the west coast of New Zealand in 275 fathoms (503 meters). They had the peculiar initial chamber with a basal spine and the later chambers pyriform. A similar specimen was found off the Galapagos Islands in deep water. The Atlantic material recorded from about the British Isles does not seem from the published figures to be typical.

NODOSARIA PYRULA d'Orbigny.

Plate 33, figs. 3-5.

"*Orthoceras Monile*" SOLDANI, Testaceographia, vol. 2, 1798, p. 35, pl. 10, figs. b, c.

Nodosaria pyrula D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 253, No. 13.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 497, pl. 62, figs. 10-12.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 309, pl. 55, fig. 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 49, pl. 26, figs. 1-3.

Brady in the *Challenger* Report records this species from off the Philippines in 95 fathoms (174 meters). Both in its typical form and the two following varieties it has proved to be very common in the region, most of the stations, however, being in the Philippine Archipelago and occurring but once in the region to the southward. Some of the stations gave abundant and often very complete specimens, showing the very long, slender, pointed proloculum, very different from that usually found in this genus.

The stations range in depth from 218 to 559 fathoms (399 to 1,021 meters), the average 387 fathoms (708 meters). The bottom temperatures range from 41.2° to 53.8° F. (5.1° to 12.1° C.), the average being 48.6° F. (9.2° C.)

Nodosaria pyrula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12299	U.S.N.M.	3	D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s....	Rare.
12301	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	53.8	gy. s., m....	Few.
12302	U.S.N.M.	1	D5214...	12 25 18 N.; 123 37 15 E..	218	51.4	gn. m.....	Few.
12302	U.S.N.M.	1	D5219...	13 21 00 N.; 122 18 45 E..	530	50.8	gn. m.....	Rare.
12304	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Rare.
12303	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Few.
15127	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.
15128	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.....	Rare.
15129	U.S.N.M.	1	D5512...	8 16 02 N.; 123 58 26 E..	445	52.8	gy. m., fine. s.	Rare.
12300	U.S.N.M.	2	D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.....	Rare.
15130	U.S.N.M.	1	D5648...	5 35 00 S.; 122 20 00 E..	559	39.2	gn. m.....	Rare.

NODOSARIA PYRULA d'Orbigny, var. SEMIRUGOSA d'Orbigny.

Plate 33, figs. 6, 7.

Nodosaria semirugosa D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 34, pl. 1, figs. 20-23.—MILLETT, Journ. Roy. Micr. Soc., 1902, p. 515, pl. 11, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 50, pl. 26, figs. 4-8.

Nodosaria No. 35, VON SCHLICHT, Foram. Septarienthones von Pietzpuhl, 1870, p. 24, pl. 7, fig. 20.

Nodosaria stipitata REUSS, var. *costulata* REUSS, Sitz. Akad. Wiss. Wien, vol. 62, Abth. 1, 1870, p. 471.

Nodosaria costulata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 515, pl. 63, figs. 23-27.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 312, pl. 58, fig. 1.

In the *Challenger* Report Brady records this variety as *Nodosaria costulata* from off the Philippines in 95 fathoms (174 meters). It

can hardly be distinguished from typical *N. pyrula* in much of the material, all gradations existing between the two extremes and also toward the following variety.

At numerous stations throughout the Philippine Archipelago this variety has occurred usually with the typical form. The depths range from 83 to 554 fathoms (151 to 1,012 meters), average 302 fathoms (552 meters), the bottom temperatures range from 41.2° to 63.6° F (5.1° to 17.5° C.), average 52.2° F. (11.2° C.).

Nodosaria pyrula, var. *semirugosa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12308	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E.	108	dk. gn. m...	Rare.
12311	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E.	554	52.8	gy. s., m....	Rare.
12310	U.S.N.M.	10	D5219...	13 21 00 N.; 122 18 45 E.	530	50.8	gn. m.....	Common.
12314	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E.	494	41.2	fine. gy. s....	Rare.
12309	U.S.N.M.	1	D5243...	6 50 55 N.; 126 14 35 E.	218	63.6	gy. m.....	Rare.
12312	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E.	312	49.3	gy. m., glob..	Rare.
15134	U.S.N.M.	2	D5371...	13 49 40 N.; 121 40 15 E.	83	gn. m.....	Rare.
15153	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E.	441	53.0	gy. m., glob.	Rare.
12313	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E.	279	53.5	gn. m.....	Rare.
15135	U.S.N.M.	1	D5538...	9 03 15 N.; 123 23 20 E.	256	53.3	gn. m., s....	Few.
15136	U.S.N.M.	1	D5574...	5 30 45 N.; 120 07 57 E.	340	52.3	Rare.

NODOSARIA PYRULA d'Orbigny, var. LONGI-COSTATA Cushman.

Plate 33, figs. 8, 9.

Nodosaria pyrula d'ORBIGNY, var. *longi-costata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 5, 1917, p. 653.

Description.—Variety with the whole of the chamber body ornamented by longitudinal costae, even connecting across the connecting necks; aperture with the costae running to the edge of the opening.

Distribution.—Type specimen (U.S.N.M. No. 9108) from *Albatross* station D5388, in 226 fathoms (414 meters), between Burias and Luzon, bottom temperature 51.4° F. (10.7° C.). In addition to the type station the variety occurred at the following stations, often in considerable numbers: D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte, bottom temperature 52.8° F. (11.5° C.); D5214, in 218 fathoms (399 meters), east of Masbate Island, bottom temperature 51.4° F. (10.7° C.); D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); D5311, in 88 fathoms (161 meters), China Sea, off Hongkong; D5338, in 43 fathoms (78 meters), Palawan Passage; D5371, in 83 fathoms (151 meters), off Marinduque Island, and D5537, in 254 fathoms (464 meters), between Negros and Siquijor, bottom temperature 53.5° F. (11.8° C.).

This variety is found in company with var. *semirugosa* d'Orbigny, but seems distinct in form as well as in ornamentation.

Nodosaria pyrula, var. *longi-costata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15131 9108 12307	U.S.N.M.	3	D5201...	10 10 00 N.; 125 04 15 E...	554	° F. 52.8	gy. s., m....	Few.
12305	U.S.N.M.	1	D5214...	12 25 18 N.; 123 27 15 E...	218	51.4	gn. m.	Rare.
15132	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
12306	U.S.N.M.	1	D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh.	Rare.
			D5338...	11 33 45 N.; 119 24 45 E...	43	Co., s., m....	Rare.
			D5371...	13 49 40 N.; 121 40 15 E...	83	gn. m.	Rare.
15133	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Rare.

NODOSARIA ANNULATA (Terquem and Berthelin).

Plate 34, figs. 1, 2a, b.

Glandulina annulata TERQUEM AND BERTHELIN, Mém. Soc. géol. France, ser. 2, vol. 10, Mem. 3, 1875, p. 22, pl. 1, fig. 25.

Nodosaria radícula, var. *annulata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 496, pl. 62, figs. 1, 2.

Description.—Test subcylindrical, composed of several chambers inflated, with deeply constricted sutures, initial end rounded, wall smooth, last-formed chambers somewhat decreasing in size; aperture central, terminal, rounded with a slight lip.

Length about 1 mm. or somewhat more.

Distribution.—This species occurred at the following *Albatross* stations: D5236, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5278, in 102 fathoms (187 meters), China Sea, off southern Luzon, bottom temperature 59.6° F. (15.3° C.); D5592, in 305 fathoms (558 meters), Sibuko Bay, Borneo, bottom temperature 43.3° F. (6.1° C.); and D5626, in 265 fathoms (485 meters), between Gillolo and Kayoa Islands.

Brady had specimens from the stations off the Ki Islands in 129 fathoms (236 meters).

Our specimens resemble those of the *Challenger* Report, and this may be a distinctive species of the Indo-Pacific region.

Nodosaria annulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12286	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	° F. 41.2	fne. gy. s....	Rare.
12285	U.S.N.M.	1	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fne. s., sh., m.	Rare.
12284	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.	Rare.
12283	U.S.N.M.	1	D5626...	0 07 30 N.; 127 29 00 E...	265	gy. m., fnc. s.	Rare.

NODOSARIA INSECTA Schwager.

Plate 34, fig. 3.

Nodosaria insecta SCHWAGER, *Novara-Expéd.*, Geol. Theil., pt. 2, 1866, p. 224, pl. 5, figs. 53, 54.

Description.—Test elongate, straight or slightly curved, chambers numerous, inflated, sutures depressed, wall smooth, proloculum with

a long spinous projection placed asymmetrically, apertural end of the chamber with a slight neck.

Length up to 5 mm.

Distribution.—Specimens of this species were very common at Albatross station D5236, in 494 fathoms (903 meters) Pacific Ocean, off eastern coast of Mindanao, bottom temperature 41.2° F. (5.1° C.), and D5121 in 108 fathoms (198 meters), off the east coast of Mindoro.

Some of our specimens were almost identical with the specimens figured by Schwager. His specimens were from the Nicobar Islands—a fossil foraminiferal fauna, which, as has already been noted, is very closely related to that of the waters of the Philippines.

Nodosaria insecta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12274	U.S.N.M.	1	D5121...	° ' " ° ' "		° F.		
12275	U.S.N.M.	1	D5236...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m...	Few.
15280	U.S.N.M.	10+		8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Common.

NODOSARIA RADICULA (Linnaeus).

Plate 34, fig. 4.

“Cornu Hammonis erectum” PLANCUS, Conch. Min., 1739, p. 14, pl. 1, fig. 5.

Nautilus radicula LINNAEUS, Syst. Nat., ed. 12, 1767, pp. 1164, 285; (Gmelin's) ed. 13, 1788, vol. 1, pt. 6, p. 3373, No. 18.

Nodosaria radicula D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 252, No. 3; Modèles, 1826, No. 1.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 495, pl. 61, figs. 28–31.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 309, pl. 55, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 52.

Specimens of this species were obtained from the following four stations: H4898, Sulu Sea, off western Mindanao, 221 fathoms (405 meters); D5300, China Sea, vicinity southern Luzon, 265 fathoms (485 meters); D5586, Sibuko Bay, Borneo, and vicinity, 347 fathoms (635 meters), bottom temperature 44° F. (6.6° C.); D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature, 47.5° F. (8.6° C.).

Nodosaria radicula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12244	U.S.N.M.	2	D5300...	° ' " ° ' "		° F.		
15025	U.S.N.M.	1	D5586...	20 31 00 N.; 115 49 00 E...	265	gy, m., s....	Rare.
12245	U.S.N.M.	1	D5666...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.	Rare.
14926	U.S.N.M.	1	H4898...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.	Rare.
				7 43 45 N.; 122 03 45 E...	221	gy. m., glob..	Rare.

NODOSARIA INSOLITA Schwager.

Nodosaria insolita SCHWAGER, *Novara-Exped.*, Geol. Theil., pt. 2, 1866, p. 230, pl. 6, fig. 63.

Description.—Test elongate, tapering, slightly arcuate, composed of numerous inflated chambers, each much broader than long, sutures deeply constricted, wall smooth, apertural end with a slight neck.

Length up to 8 mm.

Distribution.—Typical specimens were obtained from *Albatross* stations D5176, in 260 fathoms (476 meters), Verde Island Passage, and an abundance of large specimens from D5236, in 494 fathoms (903 meters), east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.). Schwager's specimens were from the fossil beds of Kar Nicobar.

The aperture shown in Schwager's figure seems as though the specimen must have been broken. Our specimens agree very closely in all other characters.

Nodosaria insolita—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12186	U.S.N.M.	2	D5176...	13 35 15 N.; 120 53 20 E...	260	° F.	s.....	Rare.
12273	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s.....	Abundant.
15662	U.S.N.M.	1						

NODOSARIA INFLEXA Reuss.

Nodosaria inflexa REUSS, *Denkschr. Akad. Wiss. Wien*, vol. 25, 1866, p. 131, pl. 2, fig. 1; *Sitz. Akad. Wiss. Wien*, vol. 62, 1870, p. 472, No. 16.—VON SCHLICHT, *Foram. Septarienthones von Pietzpuhl*, 1870, pl. 38, fig. 3.—H. B. BRADY, *Rep. Voy. Challenger, Zoology*, vol. 9, 1884, p. 498, pl. 62, fig. 9.—CUSHMAN, *Bull. 71, U. S. Nat. Mus.*, pt. 3, 1913, p. 52, pl. 25, fig. 1.

Brady recorded this species in the *Challenger* Report from off the Philippines in 95 fathoms (174 meters). I have found material at four stations as follows: D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature, 52.8° F. (11.5° C.); D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), bottom temperature, 41.2° F. (5.1° C.); D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature, 49.3° F. (9.6° C.); D5487, between Leyte and Mindanao, 732 fathoms (1,339 meters), bottom temperature, 52.3° F. (11.2° C.).

Nodosaria inflexa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12298	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	° F.	gy. s., m.... fne. gy. s..... gy. m., glob. gn. m.....	Rare.
			D5236...	8 50 45 N.; 126 26 52 E...	494	52.8		
			D5259...	11 57 30 N.; 121 42 15 E...	312	41.2		
			D5487...	10 02 45 N.; 125 05 33 E...	732	49.3		

NODOSARIA SOLUTA (Reuss).

Plate 34, figs. 5, 6.

Dentalina soluta REUSS, Zeitschr. deutsch. geol. Ges., vol. 3, 1851, p. 60, pl. 3, figs. 4a, b.

Nodosaria soluta BORNEMANN, Zeitschr. deutsch. geol. Ges., vol. 7, 1855, p. 322, pl. 12, fig. 12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 503, pl. 62, figs. 13-16; pl. 64, fig. 28.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 310, pl. 56, fig. 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 53, pl. 26, figs. 9-11.

This species is one of the most common in the region, being found in the Philippine Archipelago and to the southward as well, off Borneo, in the Gulf of Boni, Flores Sea, Macassar Strait, etc. With but one exception, that of 78 fathoms (143 meters), the range is between 208 and 940 fathoms (381 and 1,719 meters) with the average 528 fathoms (966 meters). The bottom temperatures show a wide range from 36.7° to 62.4° F. (2.6° to 16.8° C.), the average being 47.3° F. (8.4° C.).

Nearly all the specimens show rather tumid, close-set chambers, the proximal half of each being much roughened, as shown in one of the figures of the *Challenger* Report.

Nodosaria soluta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.		Character of bottom.	Abundance.
						° F.			
15106	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	fne. s.....	Rare.	
			D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	gn. m.....	Few.	
12220	} U.S.N.M.	7	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	Few.	
15107			U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.
12223	U.S.N.M.	10	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s.....	Common.	
15109	U.S.N.M.	1	D5348...	10 57 45 N.; 118 33 15 E..	375	56.4	co. s.....	Few.	
12220	} U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m., s.....	Rare.	
					D5385...	13 24 50 N.; 123 03 70 E..	327	62.4	gy. m.....
15108	U.S.N.M.	1	D5277...	Rare.	
15110	U.S.N.M.	1	D5423...	9 38 30 N.; 121 11 00 E..	508	gy. m., co. s.	Rare.	
12231	U.S.N.M.	3	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.....	Rare.	
15111	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Few.	
15112	U.S.N.M.	2	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.....	Few.	
12222	U.S.N.M.	5	D5467...	13 35 27 N.; 123 37 18 E..	480	gy. m.....	Few.	
12229	U.S.N.M.	1	D5468...	13 35 39 N.; 123 40 38 E..	569	gn. m.....	Rare.	
12230	U.N.S.M.	9	D5470...	13 37 30 N.; 123 41 09 E..	560	m.....	Common.	
15113	U.S.N.M.	1	D5487...	10 02 45 N.; 125 05 33 E..	732	52.3	gn. m.....	Rare.	
12227	U.S.N.M.	3	D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob.	Few.	
15114	U.S.N.M.	5	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Few.	
15115	U.S.N.M.	2	D5583...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.....	Few.	
15116	U.S.N.M.	3	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m.	Few.	
15117	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	Few.	
15118	U.S.N.M.	8	D5650...	4 53 45 S.; 121 29 00 E..	700	40.1	gn. m.....	Few.	
15119	U.S.N.M.	1	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	Rare.	

NODOSARIA COMMUNIS d'Orbigny.

Plate 34, fig. 7.

Nodosaria (Dentalina) communis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 254, No. 35.

Dentalina communis D'ORBIGNY, Mém. Soc. Géol. France, vol. 4, 1840, p. 13, pl. 1, fig. 4.

Nodosaria communis REUSS, Verst. Böhm. Kreid., pt. 1, 1845, p. 28, pl. 12, fig. 21.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 504, pl. 62, figs. 19–22.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 310, pl. 56, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 54, pl. 28, figs. 1, 2.

Typical material which has been referred to this species has occurred at numerous stations throughout the region often in considerable numbers. With four exceptions the depths range from 162 to 494 fathoms (297 to 903 meters). The exceptions are four stations in shallower water in 34 to 88 fathoms (62 to 161 meters), and one station in the Molucca Sea in 1,560 fathoms (2,853 meters). Where given the bottom temperatures range from 41.1° to 55.8° F. (5° to 13.2° C.).

Nodosaria communis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15097	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E..	34	° F.	wh. s.....	Rare.
12260	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	gne. s.....	Rare.
			D5244...	6 52 05 N.; 126 14 15 E..	171	gy. m.....	Rare.
15281	U.S.N.M.	6	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.....	few.
12258	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m.....	Rare.
12261	U.S.N.M.	1	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., sh.....	Rare.
15101	U.S.N.M.	1	D5412...	10 09 15 N.; 123 52 00 E..	162	54.8	gn. m.....	Rare.
12259	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.....	Rare.
15102	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Few.
15103		3	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Few.
15164		4	D5625...	0 07 00 N.; 127 23 00 E..	230	gy. m.....	Few.
12262		1	D5639...	3 54 50 S.; 123 27 20 E..	1,560	gy. m.....	Rare.
15105	U.S.N.M.			Tara Island.....	
				Apra Bay, Guam.....	
15098	U.S.N.M.	2	D5198...	9 40 50 N.; 123 39 45 E..	220	53.9	gn. m.....	Rare.
15099	U.S.N.M.	2	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.....	Rare.
15100	U.S.N.M.	1	D5244...	6 52 05 N.; 126 14 15 E..	171	gy. m.....	Rare.

NODOSARIA ROEMERI (Neugeboren).

Dentalina roemeri NEUGEBOREN, Denkschr. Akad. Wiss. Wien, vol. 12, 1856, p. 82, pl. 2, figs. 13–17.

Nodosaria roemeri REUSS, Sitz. Akad. Wiss. Wien, vol. 62, 1870, p. 475.—VON SCHLICHT, Foram. Septerionthones von Pietzpuhl, 1870, pl. 10, figs. 21, 22, 24.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 505, pl. 63, fig. 1.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 310, pl. 56, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 55, pl. 24, figs. 4–6.

Although not common at any station, specimens which are fairly typical of this species have occurred at the following stations: D5112, China Sea, off southern Luzon, 177 fathoms (324 meters), bottom temperature 52.4° F. (11.3° C.); D5284, China Sea, vicinity southern Luzon, 422 fathoms (775 meters), bottom temperature 42.3° F. (5.7° C.); D5300, China Sea, vicinity southern Luzon, 265 fathoms (485 meters); D5301, China Sea, vicinity Hongkong, 208 fathoms (381 meters), bottom temperature 50.5° F. (10.8° C.); D5542, northern Mindanao and vicinity, 200 fathoms (366 meters),

bottom temperature 54.3° F. (12.3° C.); D5386, Ragay Gulf, Luzon, 287 fathoms (525 meters), bottom temperature 62.4° F. (16.8° C.); D5585, Sibuko Bay, Borneo, and vicinity, 476 fathoms (871 meters), bottom temperature 41.1° F. (5° C.); D5636, Pitt Passage, 1,262 fathoms (2,309 meters).

Nodosaria roemeri—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
12241	U.S.N.M.	1	D5112...	13 48 22 N.;	120 47 25 E..	177	52.4	dk. gn. m....	Rare.
12232	U.S.N.M.	2	D5284...	13 42 05 N.;	120 30 45 E..	422	42.3	gy. m., glob	Rare.
12239	U.S.N.M.	8	D5300...	20 31 00 N.;	115 49 00 E..	265	gy. m., s.....	Common.
12238	U.S.N.M.	1	D5301...	20 37 00 N.;	115 43 00 E..	208	50.5	gy. m., s.....	Rare.
12242	U.S.N.M.	1	D5386...	13 38 30 N.;	122 44 30 E..	287	62.4	s.....	Rare.
			D5542...	8 48 30 N.;	123 35 30 E..	200	54.3	fine s., brk. sh.	
12240	U.S.N.M.	1	D5585...	4 07 00 N.;	118 49 54 E..	476	41.1	gy. m.....	Rare.
			D5636...	1 55 00 S.;	127 42 30 E..	1,262	gy. m., fine s.	

NODOSARIA FILIFORMIS d'Orbigny.

Plate 34, fig. 9.

Nodosaria filiformis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 253, No. 14.—
H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 500, pl. 63, figs. 3-5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 55, pl. 27, figs. 1-4.

Specimens of *Nodosaria* which have elongated chambers as those shown in the figure are here referred to this species. At a few stations they are common, but as will be noted in the table are not usually found in any considerable numbers. The range in depth from 100-750 fathoms (183-1,370 meters) is the range of greatest abundance of *Nodosaria* and other Lagenidae.

Nodosaria filiformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
15059	U.S.N.M.	1	D5121...	13 27 20 N.;	121 17 45 E..	108	° F.	dk. gn. m....	Rare.
15060	U.S.N.M.	1	D5189...	9 56 30 N.;	123 15 00 E..	300	62.8	gn. m.....	Rare.
12263	} U.S.N.M.	5	D5201...	10 10 00 N.;	125 04 15 E..	554	52.8	fine gy. s.....	Frequent.
12267									
12268	U.S.N.M.	10	D5236...	8 50 45 N.;	126 26 52 E..	494	41.2	sft. m.....	Rare.
15061	U.S.N.M.	1	D5255...	7 03 00 N.;	125 39 00 E..	100	m., sh.....	Rare.
12269	U.S.N.M.	1	D5272...	14 00 00 N.;	120 22 30 E..	118	57.4	co. s., gn. m.	Rare.
15062	U.S.N.M.	1	D5410...	10 28 45 N.;	124 05 30 E..	385	s.....	Rare.
12265	U.S.N.M.	1	D5576...	5 25 56 N.;	120 03 39 E..	277	53.3	Rare.
12264	U.S.N.M.	2	D5612...	0 38 00 S.;	121 45 40 E..	750	Rare.
15063	U.S.N.M.	1	D5424...	9 37 05 N.;	121 12 37 E..	340	54	co. s.....	Rare.
15064	U.S.N.M.	1	D5446...	12 43 51 N.;	124 59 18 E..	300	gn. m.....	Rare.
15065	U.S.N.M.	1	D5567...	5 48 00 N.;	120 33 45 E..	268	52	fine s.....	Rare.
15066	U.S.N.M.	1	D5630...	0 56 30 S.;	128 05 00 E..	569	co. s., m.....	Rare.
15067	U.S.N.M.	9	D5637...	3 53 20 S.;	126 48 00 E..	700	gy. m.....	Common.

NODOSARIA CONSOBRINA d'Orbigny, var. EMACIATA Reuss.

Plate 34, fig. 8; plate 35, fig. 1.

Dentalina emaciata REUSS, Zeitschr. deutsch. geol. Ges., vol. 3, 1851, p. 63, pl. 3, fig. 9.

Nodosaria (D.) consobrina, var. *emaciata* REUSS, Denkschr. Akad. Wiss. Wien, vol. 25, 1865, p. 132, pl. 2, figs. 12, 13.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 502, pl. 62, figs. 25, 26.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 310, pl. 56, fig. 1.—CUSHMAN, Bull. 71 U. S. Nat. Mus., pt. 3, 1913, p. 56, pl. 27, fig. 9.

Throughout the region this species seems to occur often in very considerable numbers and represented by exceptionally fine large specimens. The depths range from 50 to 940 fathoms (91 to 1,719 meters), the average being 334 fathoms (611 meters). The bottom temperatures range from 36.7° to 55.8° F. (2.6° to 13.2° C.), the average being 49.6° F. (9.8° C.).

Nodosaria consobrina, var. *emaciata*—Material examined.

Cat- No.	Coll. of—	No. of speci- mens.	Station.	Locality.		Depth in fath- oms.	Bot- tom tem- pera- ture.	Character of bottom.	Abundance.
				° ' "	° ' "				
			D5120...	13 45 30 N.;	120 30 15 E..	393	43.7	gn. m., s....	Rare.
15026	U.S.N.M.	1	D5172...	6 03 15 N.;	120 35 30 E..	318	fine. s., sh....	Rare.
15027	U.S.N.M.	1	D5198...	9 40 50 N.;	123 39 45 E..	220	53.9	gn. m.	Few.
12279	U.S.N.M.	5	D5201...	10 10 00 N.;	125 04 15 E..	554	52.8	gy. s., m....	Few.
12276	U.S.N.M.	1	D5236...	8 50 45 N.;	126 26 52 E..	494	41.2	fine. gy. s....	Rare.
15028	U.S.N.M.	3	D5220...	13 38 09 N.;	121 58 00 E..	50	sft. gn. m....	Few.
			D5259...	11 57 30 N.;	121 42 15 E..	312	49.3	gy. m., glob.	Few.
12277	U.S.N.M.	1	D5301...	20 37 00 N.;	115 43 00 E..	208	50.5	gy. m., s....	Rare.
15029	U.S.N.M.	1	D5313...	21 30 00 N.;	116 43 00 E..	150	53.6	s.....	Rare.
15030	U.S.N.M.	2	D5374...	13 46 45 N.;	121 35 08 E..	190	gy. m.	Few.
15031	U.S.N.M.	1	D5381...	13 14 15 N.;	122 44 45 E..	88	co. s.	Rare.
15032	U.S.N.M.	1	D5424...	9 37 05 N.;	121 12 37 E..	340	50.4	co. s.	Fare.
15033	U.S.N.M.	1	D5438...	15 54 42 N.;	119 44 42 E..	297	46.2	gn. m.	Rare.
15034	U.S.N.M.	1	D5439...	15 58 15 N.;	119 40 20 E..	940	36.7	gm. m.	Rare.
15035	U.S.N.M.	4	D5529...	9 23 45 N.;	123 39 30 E..	441	53	gy. m., glob.	Frequent.
12281	U.S.N.M.	5	D5537...	9 11 00 N.;	123 23 00 E..	254	53.5	gn. m.	Common.
12278	U.S.N.M.	1	D5538...	9 08 15 N.;	123 23 20 E..	256	53.3	gn. m., s....	Rare.
15036	U.S.N.M.	2	D5570...	5 32 15 N.;	120 12 57 E..	330	52.3	fine. s., glob.	Few.
12280	U.S.N.M.		D5580...	4 52 45 N.;	119 06 45 E..	162	55.8	br. s., Co....	Rare.
			D5592...	4 12 44 N.;	118 27 44 E..	305	43.3	gn. m.	Rare.
15037	U.S.N.M.	3	D5630...	0 56 30 S.;	128 05 00 E..	569	co. s., m....	Few.
15038	U.S.N.M.	2	D5630...	4 53 45 S.;	121 29 00 E..	540	40.1	gn. m.	Rare.
15039	U.S.N.M.	1	H4898...	7 43 45 N.;	122 03 45 E..	221	gy. m., glob.	Rare.
				Near Cebu.....					Rare.

NODOSARIA MUCRONATA (Neugeboren).

"*Orthoceras intortum*" SOLDANI, Testaceographia, vol. 1, pt. 2, 1791, p. 98, pl. 105, fig. V.

Nodosaria (Dentalina) obliqua D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 254, No. 36; Modèles, 1826, No. 5 (not *N. obliqua* (Linnaeus)).

Dentalina mucronata NEUGEBOREN, Denkschr. Akad. Wiss. Wien, vol. 12, 1856, p. 83, pl. 3, figs. 8-11.

Nodosaria mucronata REUSS, Sitz. Akad. Wiss. Wien, vol. 62, 1870, p. 475, No. 30.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 506, pl. 62, figs. 27-31.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 311, pl. 57, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 56, pl. 24, fig. 3; pl. 25, fig. 2; pl. 27, figs. 5-7; pl. 35, fig. 6.

Specimens of this species occurred rarely at the following five stations: D5112, China Sea off southern Luzon, 177 fathoms (324 meters) bottom temperature 52.4° F. (11.3° C.); D5120, Balayan Bay and Verde Island Passage, 393 fathoms (719 meters), bottom temperature 43.7° F. (6.5° C.); D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5394, between Samar and Masbate, 153 fathoms (279 meters); D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F. (8.6° C.).

Nodosaria mucronata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12270	U.S.N.M.	1	D5112...	° ' " ° ' "	177	° F. 52.4	dk. gn. m...	Rare.
			D5120...	13 48 22 N.; 120 47 25 E..	393	43.7	gn. m., s....	Rare.
12271	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	Rare.
			D5394...	12 00 30 N.; 124 05 36 E..	153	gn. m.....	Rare.
12272	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Rare.

NODOSARIA JAPONICA Cushman.

Plate 35, fig. 2.

Nodosaria japonica CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913., p. 57, pl. 28, fig. 4.

This species was described from *Albatross* station D4900, off the southern portion of Japan, where numerous subtropical species seem to reach their northern limits. It is not surprising therefore to find as in other cases that this species occurs frequently in the Philippine region. It has occurred at numerous stations well scattered throughout the region ranging in depth mostly from 118 to 700 fathoms (203 to 1,280 meters), with two stations less than 100 fathoms (183 meters), with depths of 78 and 14 to 25 fathoms (143 and 25 to 46 meters), with the average 293 fathoms (536 meters). The bottom temperatures range from 50.4° to 57.4° F. (10.2° to 14.1° C.), with the average of 52.5° F. (11.3° C.). This is close to the bottom temperature of the original station, which was 52.9° F. (11.6° C.).

The Philippine specimens are not as large as those from southern Japan, but many of them are nearly 10 mm. in length and have the characteristic form. The localities are off Romblon; off western Bohol; between Marinduque and Luzon; China Sea, off southern Luzon; off Hongkong, and off Formosa; off Palawan, Jolo, and Tawi Tawi; and to the south near Patiente Strait; and in the Gulf of Boni.

Nodosaria japonica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15120	U.S.N.M.	3	D5178...	12 43 00 N.; 122 06 15 E ..	78	fne. s.....	Frequent.
12188	U.S.N.M.	1	D5198...	9 40 50 N.; 123 39 45 E ..	220	53.9	gn. m.....	Rare.
12159	U.S.N.M.	1	D5219...	13 21 00 N.; 122 18 45 E ..	530	50.8	gn. m.....	Rare.
12760	U.S.N.M.	2	D5236 ..	8 50 45 N.; 126 26 52 E ..	494	41.2	fne. gy. s.....	Rare.
15122	U.S.N.M.	1	D5272...	14 00 00 N.; 120 22 30 E ..	118	57.4	m., sh., co. s.	Rare.
12160	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E ..	170	s., p.....	Rare.
12161	U.S.N.M.	7	D5301...	20 37 00 N.; 115 43 00 E ..	208	50.5	gy. m., s.....	Frequent.
15123	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E ..	340	s., br. Co.....	Rare.
15124	U.S.N.M.	1	D5342 ..	10 56 55 N.; 119 17 24 E ..	14-25	gy. m.....	Rare.
12162	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E ..	340	50.4	co. s., m.....	Rare.
15125	U.S.N.M.	1	D5567...	13 35 27 N.; 123 27 18 E ..	480	gy. m.....	Rare.
			D5630...	0 56 30 S.; 128 05 00 E ..	569	co. s., m.....	Frequent.
12163	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 00 E ..	700	gy. m.....	Rare.
15126	U.S.N.M.	1	D5950...	4 53 45 S.; 121 29 00 E ..	540	40.1	gn. m.....	Rare.
15121	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E ..	312	49.3	gy. m., glob.	

NODOSARIA ANTENNULA Cushman.

Plate 35, fig. 3.

Nodosaria antennula CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 653.

Description.—Test elongate, slightly arcuate, composed of numerous short chambers, sutures projecting, of clear shell material, proloculum bulbous, apertural end of chamber tapering to a rounded point, wall smooth.

Length up to 8 mm.

Distribution.—Type specimens (No. 9109, U.S.N.M.) from *Albatross* station D5236 in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.). It also occurred at *Albatross* station D5121, in 108 fathoms (198 meters), east coast of Mindoro.

This species is a very peculiar one, the sutures being limbate, of clear shell material, standing out wider in side view than the chambers themselves, which are concaved between the sutures instead of inflated, as is usually the case in this genus.

Nodosaria antennula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12282	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E ..	108	dk. gn. m....	Rare.
9109	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E ..	494	41.2	fne. gy. s....	Common.
15282	U.S.N.M.	8						

NODOSARIA SUBPOLYGONA Cushman.

Plate 35, figs. 4a, 4b, 5.

Nodosaria subpolygona CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 654.

Description.—Test elongate, only very slightly tapering except in the microspheric form, polygonal in end view, chambers numerous, not at all inflated, sides parallel or very nearly so, sutures very slightly if at all depressed, apertural end with a short stout neck, aperture large, circular; surface ornamented by six raised ridges giving a hexagonal appearance in end view.

Length up to 5 mm.

Distribution.—Type specimens (U.S.N.M. No. 9110) from *Albatross* station D5318 in 340 fathoms (622 meters), China Sea, vicinity of Formosa.

Both microspheric and megalospheric forms of the species occur at this station, the former having numerous chambers and a tapering base, the latter with a very large proloculum and the succeeding chambers but slightly increasing in size and fewer than in the microspheric form.

The polygonal form with few ridges and the short neck with large aperture will distinguish this species.

Nodosaria subpolygona—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9110 15294	U.S.N.M.	2 1	D5318...	° ' " ° ' "	340	° F.	s., br. Co....	Rare.

NODOSARIA CATENULATA H. B. Brady.

Plate 35, fig. 7.

Nodosaria catenulata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 515, pl. 63, figs. 32-34.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 312, pl. 58, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 57, pl. 25, fig. 3.

Very typical specimens were found at the following Philippine stations: D5242, in 191 fathoms (350 meters), vicinity of Pujada Bay, bottom temperature 64.1° F. (17.8° C.); D5255, in 100 fathoms (183 meters), Gulf of Davao; D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); D5368, in 181 fathoms (331 meters), off Marinduque Island; and

D5572, in 334 fathoms (611 meters), north of Tawi Tawi, bottom temperature 52.3° F. (11.2° C.). The specimens were very abundant at one of these stations.

Brady described this species from the *Challenger* station in the Philippines, depth 95 fathoms (174 meters), and from off Raine Island, Torres Strait, 155 fathoms (283 meters). Sidebottom records it from the east coast of Australia in 465 fathoms (850 meters). Flint's specimen was from the Gulf of Mexico, where *N. vertebralis* is abundant and may be a form of that species with few costae, or the distribution of *N. catenulata* may include this region also.

Nodosaria catenulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12290	U.S.N.M.	1	D5242...	6 51 53 N.; 126 14 10 E....	191	64.1	sft. gy. m....	Rare.
12289	U.S.N.M.	1	D5255...	7 03 00 N.; 125 39 00 E....	100	sft. m.	Rare.
15283	U.S.N.M.	6	D5259...	11 57 30 N.; 121 42 15 E....	312	49.3	gy. m., glob..	Abundant.
12288	U.S.N.M.	1	D5368...	13 35 30 N.; 121 48 00 E....	181	gy. m.	Rare.
			D5572...	5 31 26 N.; 120 09 45 E....	334	52.3	s., sh.....	

NODOSARIA SCALARIS (Batsch).

Plate 35, fig. 6.

"*Orthoceria Ftosculi*" SOLDANI, Testaceographia, vol. 1, pt. 2, 1791, p. 91, pl. 95, figs. B-M.

Nautilus (Orthoceras) scalaris BATSCH, Conch. des Seesandes, 1791, No. 4, pl. 2, figs. 4a, b.

Nodosaria scalaris PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 340, pl. 16, figs. 2a, b, c.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 510, pl. 63, figs. 28-31.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 58, pl. 24, fig. 7.

Brady in the *Challenger* Report records this species in 95 fathoms, (174 meters), off the Philippine Islands. The typical form has been found to be abundant in this material, occurring off various coasts of Luzon; off Romblon; Sogod Bay, southern Leyte; east of Masbate; Verde Island Passage; between Leyte and Mindanao; north of Tawi Tawi; and farther south in Darvel Bay and Sibuko Bay, Borneo, and Buton Strait. The depths range from 24 to 890 fathoms (44 to 1,628 meters), with the average 274 fathoms (502 meters), and the bottom temperatures from 38.3° to 63.1° F. (3.5° to 17.2° C.), with the average 50.4° F. (10.2° C.).

Nodosaria scalaris—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15075	U.S.N.M.	1	D5100...	14 17 15 N.; 120 32 40 E....	35	gy. s.	Rare.
15076	U.S.N.M.	2	D5110...	13 59 20 N.; 120 75 45 E....	135	59.0	dk. gy. m.	Few.
12166	U.S.N.M.	2	D5112...	13 48 22 N.; 120 47 25 E....	177	52.4	dk. gn. m.	Few.
12207	U.S.N.M.	8	D5178...	12 43 00 N.; 122 06 15 E....	78	fine. s.	Frequent.
12205	U.S.N.M.	6	D5201...	10 10 00 N.; 125 04 15 E....	544	52.8	gy. s., m.	Frequent.
12209								
12206	U.S.N.M.	1	D5236...	12 15 00 N.; 123 57 30 E....	80	s. m., sh.	Frequent.
15079	U.S.N.M.	1	D5269...	8 50 45 N.; 126 26 52 E....	494	41.2	fine. gy. s.	Rare.
15080	U.S.N.M.	2	D5278...	13 39 50 N.; 120 59 30 E....	220	fine. s., p.	Rare.
15081	U.S.N.M.	2	D5278...	14 00 10 N.; 120 17 15 E....	102	59.6	fine. s., m., sh.	Few.
15081	U.S.N.M.	4	D5282...	13 53 00 N.; 120 26 45 E....	248	47.4	dk. gy. s.	Frequent.
15082	U.S.N.M.	2	D5300...	20 31 00 N.; 115 49 00 E....	265	gy. m., s.	Rare.
15083	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E....	150	53.6	s.	Few...
			D5318...	21 32 00 N.; 117 46 00 E....	340	s., br. Co.	Rare.
			D5487...	10 02 45 N.; 125 05 33 E....	732	52.3	gn. m.	Rare.
			D5572...	5 31 26 N.; 120 09 45 E....	334	52.3	Rare.
12167	U.S.N.M.	1	D5582...	4 19 54 N.; 118 58 38 E....	890	38.3	gy. m., fine. s.	Few.
15084	U.S.N.M.	4	D5592...	4 12 44 N.; 118 27 44 E....	305	43.3	gn. m.	Few.
15085	U.S.N.M.	1	D5640...	4 27 00 S.; 122 55 40 E....	24	s., brk. sh.	Few.
12208	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E....	540	40.1	gn. m.	Rare.
15077	U.S.N.M.	2	D5217...	13 20 00 N.; 123 14 15 E....	105	63.1	co. gy. s.	Rare.
15078	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E....	170	s., p.	Rare.
15086	U.S.N.M.	1	Tara Island, Philippine Islands.	Rare.

NODOSARIA SCALARIS (Batsch), var. SEPARANS H. B. Brady.

Nodosaria scalaris, var. *separans* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 511, pl. 64, figs. 16–19.

Description.—Variety with the last-formed chambers, one or more, remote, the neck of the previous chamber exposed between, the early chambers also often less inflated than in the typical.

Distribution.—This variety occurred at the following *Albatross* stations: D5201, in 544 fathoms (987 meters), Sogod Bay, southern Leyte, bottom temperature 52.8° F. (11.5° C.); D5254, Gulf of Davao, 28 fathoms (51 meters), no bottom temperature; D5282, in 248 fathoms (45½ meters), China Sea, off southern Luzon, bottom temperature 47.4° F. (8.5° C.); D5630, in 569 fathoms (1,040 meters), south of Patiente Strait, and D5650, Gulf of Davao, 540 fathoms (988 meters), bottom temperature 40.1° F. (4.5° C.).

The variety seems usually to be far less widely distributed than is the typical form.

Nodosaria scalaris, var. *separans*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15087	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	544	52.8	gy. s., m.	Rare.
12169	U.S.N.M.	2	D5254...	7 05 42 N.; 125 39 42 E..	28	s., Co.	Rare.
12168	U.S.N.M.	1	D5282...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. gy. s.	Rare.
12170	U.S.N.M.	2	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m.	Rare.
12172								
12171	U.S.N.M.	6	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.	Few.
12173								

NODOSARIA SUBSCALARIS Cushman.

Plate 36, figs. 2 a, b.

Nodosaria subscalaris CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 654.

Description.—Test elongate, composed of few chambers, mucronate at the apical end, earlier chambers close-set, the later ones often somewhat separated, inflated, sutures depressed, surface ornamented with numerous (up to 40) longitudinal costae, apertural end with a thick, tapering neck with 9 or 10 prominent, plate-like costae extending from the upper end of the ornamentation of the body of the chamber to the apertural end of the neck; aperture small, rounded.

Length up to 5 mm.

Distribution.—Type specimens (U.S.N.M. No. 9111) from *Albatross* station D5178, in 78 fathoms (143 meters), vicinity of Romblon, where the species was very abundant. It also occurred at D5277 in 80 fathoms (146 meters), China Sea vicinity of southern Luzon, bottom temperature 58.6° F. (14.7° C.).

This species differs from typical *N. scalaris* in the apertural characters especially, our species having a stout conical neck with very prominent longitudinal costae, while typical *scalaris* has a much narrower cylindrical neck with annular raised costae. This character is constant in the two species, as can be seen in specimens where the chambers are separated and the apertural characters of several chambers can be studied. The two species also do not occur mixed at the same station.

The apertural ornamentation and general characters suggest very strongly that of *Lagena striata* (d'Orbigny), var. *haidingeri* (Czjzek).¹⁵

Nodosaria subscalaris—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9111 12235 12234	} U.S.N.M.	10	D5178...	° ' " ° ' "	78	° F. 75.7	fine.s.....	Common.
		1	D5277...	13 56 55 N.; 120 13 45 E..	80	58.6	fine.s.....	Rare.

NODOSARIA SUBSCALARIS Cushman, var. PAUCI-COSTATA Cushman.

Plate 36, figs. 3, 4.

Nodosaria subscalaris CUSHMAN, var. *pauci-costata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 654.

Description.—Variety differing from the typical in the lesser number of costae (12–25) and in the apertural characters which in the variety consist of a stout nearly cylindrical neck, with about 15 costae, which are continuations of the costae of the body portion.

¹⁵ Cushman, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, pl. 7, fig. 6.

Distribution.—Type specimens (U.S.N.M. No. 9112) from *Albatross* station D5152, in 34 fathoms (62 meters), Tawi Tawi group, Sulu Archipelago. It has also occurred at D5217, in 105 fathoms (193 meters), between Burias and Luzon, bottom temperature 63.1° F. (17.2° C.); and D5133, in 38 fathoms (70 meters), Sulu Sea, off western Mindanao.

In the variety the reduction in number of the costae of the body portion of the chamber is accompanied by an increase in the number of the costae of the apertural neck and by the difference in shape of the neck as well.

Nodosaria subscalaris, var. *pauci-costata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12232	U.S.N.M.	2	D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	° F.	gn. m., s.....	Few.
15284	U.S.N.M.	4						
12295	U.S.N.M.	10	D5152...	5 22 55 N.; 120 15 45 E..	34	° F.	wh. s.....	Common.
15285	U.S.N.M.	6						
12233	U.S.N.M.	1	D5217...	13 00 20 N.; 123 14 15 E..	105	63.1	crs. gy. s.....	Rare.

NODOSARIA MILLETTII Cushman.

Plate 36, fig. 5.

Nodosaria millettii CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 654.

Nodosaria scalaris, var. *separans* MILLETT, Journ. Roy. Micr. Soc., 1902, p. 520, pl. 11, figs. 11, 12 (not var. *separans* H. B. BRADY).

Description.—Test elongate, nodose, consisting of few chambers, the early ones close-set, the latter ones remote, chambers pyriform or elongate-elliptical in side view, very finely costate, the basal portion sometimes slightly hispid, apertural necks very long and slender, aperture with a phialine lip.

Length up to 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9113) from *Albatross* station D5281, in 201 fathoms (368 meters), China Sea, off southern Luzon, bottom temperature 50.4° F. (10.2° C.).

This species found by Millett in this same general region, but farther south, seems to differ materially from *N. scalaris*, var. *separans*. The appearance of the test is at once different, more graceful, more transparent, and very much more delicate and finer in its surface markings. The form of the chamber is also much longer and more elliptical, the neck more slender, and the chambers more remote, while the end of the chamber, instead of having radiating slits as in var. *separans* and *scalaris*, has a definite, entire, phialine lip.

Nodosaria millettii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9113	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E..	201	° F. 50.4	dk. gy. s....	Rare.
12287	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.

NODOSARIA LEPIDULA Schwager.

Plate 36, fig. 6.

Nodosaria lepidula SCHWAGER, *Novara-Exped.*, Geol. Theil., pt. 2, 1866, p. 210, pl. 5, figs. 27, 28.

Description.—Test elongate, straight, tapering gradually to the acute initial end, chambers numerous, about as long as wide, chambers well separated by deep sutures, wall with an ornamentation consisting of a ring of short costae or spines about the widest portion of the chamber, aperture with a phialine lip, with a flange below the extreme top.

Length up to 2 mm.

Distribution.—Specimens were common at *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, off east coast of Mindanao, bottom temperature 41.2 ° F. (5.1° C.).

Some of the figures referred by Brady to *Sagrina* seem to belong to this species. Our specimens were nearly identical in all characters with those figured by Schwager from the Nicobar Islands.

Nodosaria lepidula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
15286	U.S.N.M.	6	D5236...	8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fine. gy. s....	Common.

NODOSARIA LEPIDULA Schwager, var. *HISPIDULA* Cushman.

Plate 36, fig. 7.

Nodosaria lepidula SCHWAGER, var. *hispidula* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 654.

Description.—Test differing from the typical in its larger size and the ornamentation, which in the variety consists of a series of interrupted fine costae and hispid roughenings of the surface over nearly the whole surface of the chamber.

Length up to 3 mm.

Distribution.—Type specimens (U.S.N.M. No. 9114) were found with the typical in considerable numbers at *Albatross* station D5236.

Nodosaria lepidula, var. *hispidula*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9114	U.S.N.M.	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fine-gy. s.....	Common.

NODOSARIA SUBSTRIATULA Cushman.

Plate 36, figs. 8, 9; plate 52, figs. 7-9.

Nodosaria subcanaliculata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 512, pl. 64, figs. 23, 24 (not *Dentalina subcanaliculata* Neugeboren).

Nodosaria substriatula CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 655.

Description.—Test usually consisting of four chambers, the proloculum with a globular body, an apical spine, and with definite, longitudinal costae; the second chamber much more elongate, flask-shaped with a long neck; when partially covered by the third chamber becoming subcylindrical, ornamented like the proloculum; third chamber still more elongate, with a long neck, the surface ornamentation consisting of short, broken, longitudinal striae; the fourth chamber similar but remote, a large part of the neck between the two chambers visible; final chamber with the apertural neck long and slender, smooth; the apertural end with four or more flange-like costae extending up and beyond the aperture and incurving somewhat over the aperture.

Length up to 2 mm.

Distribution.—Type specimen (U.S.N.M. No. 9115) from *Albatross* station D5123, in 283 fathoms (517 meters), east coast of Mindanao. Specimens also were obtained in material from D5112, in 177 fathoms (324 meters), China Sea, off southern Luzon, bottom temperature 52.4° F. (11.3° C.); D5120, in 393 fathoms (719 meters), off Balayan Bay, bottom temperature 43.7° F. (6.5° C.); D5122, east coast of Mindoro, 220 fathoms (404 meters), and D5123, same locality, 283 fathoms (517 meters); D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); D5260, off southeastern Mindoro, 234 fathoms (428 meters), bottom temperature 51.4° F. (10.7° C.); D5284, in 422 fathoms (775 meters), China Sea off southern Luzon, bottom temperature 42.3° F. (5.7° C.); D5300, in 265 fathoms (485 meters), same region, and D5420, in 127 fathoms (233 meters), between Cebu and Bohol, bottom temperature 59.0° F. (15° C.).

Brady had this species, apparently a single specimen, from 420 fathoms (728 meters) off Tahiti. He referred it provisionally to a

species described by Neugeboren from the Miocene of Transylvania. His wording is interesting:

Under the name *Dentalina subcanaliculata*, Neugeboren has figured two slender, curved, *Nodosaria* shells, which are characterized by their numerous segments, straight sutures, and a peculiar surface-ornament consisting of short, broken, longitudinal striae. The specimen represented in Pl. LXIV, figs. 23, 24 [*Challenger Report*], exhibits the same kind of external marking; and as the test is manifestly abnormal in point of form, it may be treated provisionally, in the absence of other indications of its affinity, as an aberrant modification of the species referred to.

With the single specimen at his disposal little else could be done by Brady, and certainly from a single specimen it might appear, as he says, that "the test is manifestly abnormal in point of form." However, I have had a considerable number of specimens from the above localities and the form is more than usually constant. The figure given by Brady is so well drawn that it might represent one of the specimens I have had. His specimen, however, did not have its aperture in complete preservation, as have a portion of those which I have been fortunate enough to obtain.

This is certainly a rather unique species in many ways, and is apparently rather widely distributed in the Indo-Pacific region, although it is a delicate one and easily broken.

Nodosaria substriatula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
12296	U.S.N.M.	1	D5112...	13 48 22 N.; 120 47 25 E..	177	52.4	dk. gn. m...	Rare.
			D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s....	
			D5122...	13 21 30 N.; 120 30 33 E..	220	gn. m.....	
11603 9115j	U.S.N.M.	2	D5123...	13 12 45 N.; 121 38 45 E..	283	gn. m.....	Rare.
			D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Rare.
12219	U.S.N.M.	1	D5290...	12 25 35 N.; 121 31 35 E..	234	51.4	gn. m., s....	Rare.
12216	U.S.N.M.	1	D5284...	13 42 05 N.; 120 30 45 E..	422	42.3	gy. m., glob.	Common.
			D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	
12217	U.S.N.M.	10	D5420...	9 49 35 N.; 123 45 00 E..	127	59.0	Rare.

NODOSARIA PAUCILOCULATA Cushman.

Plate 36, figs. 10-12.

Nodosaria pauciloculata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 655.

Description.—Test nodose, composed of few chambers, usually not more than five, two or three closely set, later ones remote, inflated, sutures even in the early chambers much depressed, wall ornamented by longitudinal costae, few and large, usually limited to the middle

portion of the chamber and becoming obsolete toward each end, neck long, swollen toward the top, then again rapidly contracted to the aperture; in the early chambers the neck with costae, in later ones with the costae limited to the last tapering portion near the aperture.

Length, up to 3.5 mm.

Distribution.—Type specimens (U.S.N.M. No. 9116) from *Albatross* station D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte Island, bottom temperature, 52.8° F. (11.5° C.).

Specimens were frequent at the type station, also at D5259, in 312 fathoms (571 meters), off north-western Panay, bottom temperature, 49.3° F. (9.6° C.); D5371, in 83 fathoms (151 meters), vicinity of Marinduque Island; and D5538, in 256 fathoms (468 meters), between Negros and Siquijor, bottom temperature, 53.3° F. (11.8° C.).



FIG. 8.—*NODOSARIA PAUCILOCLATA*, VAR. *LUZONENSIS*, NEW VARIETY. FRONT VIEW, $\times 100$. FROM STATION D5220.

This species very quickly loses its close-set character, but two and occasionally three chambers being so arranged, after which the chambers are at once remote. The costae, limited to the middle area of the chamber and the very characteristic shape of the apertural neck, are distinguishing characters. It is usually mucronate at the apical end. A young stage with three chambers is shown, as well as two adult five-chambered specimens, one with three, the other with two close-set early chambers.

Nodosaria pauciloculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12294	U.S.N.M.	2	D5112...	13 48 22 N.; 120 47 25 E..	177	52.4	dk. gn. m...	Rare.
12211	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s....	Rare.
9116	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Frequent.
15287	U.S.N.M.	1		13 21 00 N.; 122 18 45 E..	530	50.8	gn. m.....	Rare.
12213	U.S.N.M.	2	D5219...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.....	Rare.
12213	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.....	Rare.
12214	U.S.N.M.	4	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Frequent.
12215								
12216	U.S.N.M.	4	D5371...	13 49 40 N.; 121 40 15 E..	83	gn. m.....	Frequent.
12210	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gn. m., s....	Rare.
14925	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob.	Rare.

NODOSARIA PAUCILOCLATA Cushman, var. *LUZONENSIS*, new variety.

Description.—Test differing from the typical in having three closely set chambers followed by a single one, somewhat remotely placed; earlier chambers without the neck costae continuous.

Distribution.—Type specimen (U.S.N.M. No. 15288), from *Albatross* station D5220, between Marinduque and Luzon, 50 fathoms (91 meters).

This variety was common at this station, and seems to replace the typical, which was not found here.

Nodosaria pauciloculata, var. *luzonensis*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15288 15289	U.S.N.M.	8	D5220...	° ' " ° ' " 13 38 00 N.; 121 58 00 E..	50	° F.	sft. gn. m. . .	Common.

NODOSARIA SUBLINEATA H. B. BRADY.

Plate 37, fig. 1.

Nodosaria hispida, var. *sublineata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 508, pl. 63, figs. 19–22.

Description.—Test elongate, nodose, composed of few chambers, the early ones sometimes close-set, initial end mucronate, wall ornamented by longitudinal costae, more or less broken transversely, apertural neck cylindrical, with annular or spiral costae.

Length up to 6 mm.

Distribution.—Specimens referred to this species occurred at *Albatross* stations D5100, China Sea, off southern Luzon, 35 fathoms (66 meters), no bottom temperature; D5586, in 347 fathoms (635 meters), near Sibuko Bay, Borneo, bottom temperature 44° F (6.6° C.); and D5650, in 540 fathoms (988 meters), Gulf of Boni, bottom temperature 40.1° F. (4.5° C.). At the latter station the specimens were very numerous.

These specimens in some ways differ from those figured by Brady, but in the main seem to have the characters which he notes. Their relation to *N. hirsuta* does not seem to be close, some of the specimens suggesting *N. scalaris* much more closely. The specimen figured here shows the extreme form, with several close-set chambers and the costae but slightly broken. From this extreme the specimens range toward those figured by Brady. This possibly may prove not to be Brady's species, but a new one.

Nodosaria sublineata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12237	U.S.N.M.	1	D5100...	° ' " ° ' " 14 17 15 N.; 120 32 40 E..	35	° F.	gy. s.	Rare.
15290	U.S.N.M.	3	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.	Few.
12236	U.S.N.M.	10	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.	Common.

NODOSARIA PERVERSA Schwager.

Plate 37, fig. 2.

Nodosaria perversa SCHWAGER, *Novara-Exped.*, Geol. Theil., vol. 2, 1866, p. 212, pl. 5, fig. 29.—H. B. BRADY, *Rep. Voy. Challenger*, Zoology, vol. 9, 1884, p. 512, pl. 64, figs. 25-27.

Description.—Test elongate, subcylindrical, chambers comparatively few, inflated; sutures depressed, initial end broadly rounded; wall with numerous longitudinal costae, last-formed chambers usually smooth; apertural end with a short neck with a definite phialine lip.

Length 1 to 1.5 mm.

Distribution.—Specimens identical with Schwager's species occurred at *Albatross* station D5285, in 272 fathoms (498 meters), China Sea near southern Luzon, bottom temperature 46.5° F, (8° C.); and D5311, in 88 fathoms (161 meters), China Sea, vicinity of Hongkong.

Schwager's specimens were from the Nicobar Islands, the fossils of which, as has already been noted, are closely related to the present foraminiferal fauna of the Philippines. Brady's specimens were from off the Ki Islands, southwest of New Guinea, the fauna of which is also related closely to the fauna of the Philippines.

The specimens figured by Silvestri are closely allied if not identical with this species.¹⁶

Egger records this species from the west coast of Australia, Chapman from east of Tasmania in 1,122 fathoms, and Dakin as rare in the Gulf of Manaar, Ceylon.¹⁷

Nodosaria perversa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12291	U.S.N.M.	1	D5257...	° ' " ° ' "	28	° F.	m.	Rare.
12187	U.S.N.M.	1	D5285...	7 22 12 N.; 124 12 15 E..	272	46.5	sft. m.	Rare.
12292	} U.S.N.M.	2	} D5311...	13 39 36 N.; 120 32 55 E..	88	crs. s., sh....	Rare.
15293		1		21 33 00 N.; 116 15 00 E..				

NODOSARIA SUBPERVERSA Cushman.

Plate 37, fig. 3.

Nodosaria subperversa CUSHMAN, *Proc. U. S. Nat. Mus.*, vol. 51, 1917, p. 655.

Description.—Test elongate, subcylindrical, composed of several chambers, proloculum larger than the succeeding chambers; all chambers close-set, the breadth being much greater than the length,

¹⁶ Atti dell'Accad. Gioenia di Sci. Nat. di Catania, ser. 3, vol. 7, 1872, p. 65, pl. 4, figs. 148-158.

¹⁷ Egger, *Abh. Kön. Bay. Akad. München*, Cl. II, vol. 18, 1893, p. 152, pl. 11, fig. 42. Chapman, *Biol. Res. Endeavor*, vol. 3, pt. 1, 1915, p. 23. Dakin, *Ceylon Pearl-Oyster Fisheries*, *Suppl. Repts.*, pt. 5, 1906, p. 235.

sutures deeply constricted, wall finely marked by longitudinal striations, aperture without a neck.

Length up to 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9117) from *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.).

This species differs from *N. perversa* Schwager in its broader, shorter chambers and its very much finer ornamentation, as well as the lack of a definite apertural neck and the mucronate character of the proloculum.

Nodosaria subperversa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9117	U.S.N.M.	1	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E..	494	°F. 41.2	fne. gy. s....	Rare.

NODOSARIA LAEVICOSTATA Cushman.

Plate 37, fig. 4.

Nodosaria laevicostata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 656.

Description.—Test elongate, arcuate, composed of but few chambers, early ones more or less rounded, closely set, ornamented with longitudinal costae, later chambers pyriform, with the last-formed chamber often remote, wall smooth; apertural end tapering into an apertural neck.

Length up to 2 mm.

Distribution.—Type specimens (U.S.N.M. No. 9118) from *Albatross* station D5178, in 78 fathoms (143 meters), near Romblon.

This is a peculiarly formed species, the early, close-set chambers costate being nearly in a straight line, while the later smooth ones are considerably angled. None of the specimens seen had more than six chambers, the first three or four being close set and costate.

Nodosaria laevicostata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9118 15291	U.S.N.M.	2	D5178...	° ' " ° ' " 12 43 00 N.; 122 06 15 E..	78	°F.	fne. s.....	Rare.

NODOSARIA RAPHANUS (Linnaeus).

Plate 37, figs. 5-7.

"Cornu Hammonis erectum striatum" PLANCUS, Conch. Min., 1739, p. 15, pl. 1, fig. 6.

"Orthoceras minimum" etc., GAULTIERI, Index Test., 1742, pl. 19, fig. L.

Nautilus raphanus LINNAEUS, Syst. Nat., ed. 12, 1767, pp. 1164, 283; (Gmelin's) ed. 13, 1788, p. 3372, No. 16.

Orthocera raphanus LAMARCK, Anim. sans Vert., vol. 7, 1822, p. 593, No. 1; Tabl. Encycl. et Méth., pl. 465, figs. 2a, b, c.

Nodosaria raphanus PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 3, 1859, p. 477.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 512, pl. 64, figs. 6-10.—CUSHMAN, Bull. U. S. Nat. Mus., pt. 3, 1913, p. 59, pl. 26, figs. 12, 13.

Excellent specimens of this species were obtained at numerous stations well scattered throughout the region, but mostly in medium depths. The range of depths is from 80 to 554 fathoms (146 to 1,012 meters), with an average of 229 fathoms (419 meters), and the bottom temperatures from 53° F. to 58.6° F (11.7° C. to 14.7° C.), with an average of 55.6° F. (13.1° C.).

Some of the largest specimens measure as much as 25 mm., and often specimens show considerable constriction, usually toward the apertural end, but also occasionally in the middle of the test. This is one of the finest species in the region.

Nodosaria raphanus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		"F.		
12252	U.S.N.M.	2	D5198...	9 40 50 N.; 123 39 45 E..	220	54.3	gn. m.....	Rare.
12251	U.S.N.M.	1	D5201....	10 10 00 N.; 125 04 15 E..	554	53.8	gy. s., m....	Rare.
15088	U.S.N.M.	2	D5281....	12 30 55 N.; 121 34 24 E..	145	s. m.....	Few.
12248	U.S.N.M.	3	D5272....	14 00 00 N.; 120 22 30 E..	118	57.4	m., sh., co. s.	Common.
12249 12250	U.S.N.M.	2	D5277....	13 56 55 N.; 120 13 45 E..	80	58.6	fine. s.....	Rare.
15089	U.S.N.M.	1	D5313....	21 30 00 N.; 116 43 00 E..	150	53.6	s.....	Rare.
15090	U.S.N.M.	1	D5318....	21 32 00 N.; 117 46 00 E..	340	s. br. Co	Rare.
12247	U.S.N.M.	3	D5369....	13 48 00 N.; 121 43 00 E..	106	bk. s.....	Frequent.
15091	U.S.N.M.	1	D5370....	13 44 15 N.; 121 42 30 E..	159	54.3	sft. m.....	Rare.
15092	U.S.N.M.	6	D5374....	13 46 45 N.; 121 35 08 E..	190	gy. m.....	Common.
15093	U.S.N.M.	1	D5382....	13 15 20 N.; 122 45 30 E..	128	m.....	Rare.
12246	U.S.N.M.	1	D5454....	13 12 00 N.; 123 50 30 E..	153	Rare
15094	U.S.N.M.	1	D5523....	8 48 44 N.; 123 27 35 E..	Rare
15095	U.S.N.M.	1	D5529....	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m.....	Rare.
15096	U.S.N.M.	1	D5650....	4 54 45 S.; 121 29 00 E..	540	gn. m.....	Rare

NODOSARIA OBLIQUA (Linnaeus).

Plate 38, fig. 1.

"Orthoceras minimum" etc., GAULTIERI, Index Test., 1742, pl. 19, fig. N.
Nautilus obliquus LINNAEUS, Syst. Nat., ed. 12, 1767, p. 1163; (Gmelin's) ed. 13, 1788, p. 3372, No. 14.

Nodosaria (Dentalina) obliqua PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 3, 1859, p. 482.

Nodosaria obliqua H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 513, pl. 64, figs. 20-22.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 311, pl. 57, fig. 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 59, pl. 25, fig. 5,

Beautiful specimens of this species are well distributed over the area occurring at numerous stations: Off western Bohol; Sogod Bay, southern Leyte; between Marinduque and Luzon; Gulf of Davao; east coast of Luzon, off northern Mindanao; between Siquijor and Bohol; between Negros and Siquijor; off Sibuko Bay, Borneo; and Bouro Island. The depth of these stations ranges from 100 to 700 fathoms (183 to 1,280 meters), average depth about 400 fathoms (732 meters). The bottom temperatures where given range from 43.3° to 53.9° F. (6.1° to 12.1° C.), the average being 49.8° F. (9.8° C.).

While occurring at a large number of stations this is not so well represented in numbers as are other species of the genus.

Nodosaria obliqua—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12202	} U.S.N.M.	3	D5198...	9 40 50 N.; 123 39 45 E..	220	53.9	gn. m.....	Rare.
12204								
12200	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Few.
15068	U.S.N.M.	4	D5219...	13 21 00 N.; 122 18 45 E..	530	50.8	gn. m.....	Few.
15069	U.S.N.M.	1	D5255...	7 03 00 N.; 125 39 00 E..	100	sft. m.....	Rare.
12199	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E..	312	49.2	gy. m.....	Rare.
15070	U.S.N.M.	1	D5374...	30 46 45 N.; 121 35 08 E..	190	gy. m.....	Rare.
15071	U.S.N.M.	1	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.....	Rare.
12203	U.S.N.M.	2	D5465...	13 39 42 N.; 123 40 39 E..	500	gy. m.....	Rare.
12201	U.S.N.M.	1	D5470...	13 37 30 N.; 123 41 09 E..	560	m.....	Rare.
15072	U.S.N.M.	2	D5523...	8 48 44 N.; 123 27 35 E..	Rare.
15073	U.S.N.M.	2	D5529...	9 23 45 N.; 123 39 30 E..	441	51.0	gy. m.....	Rare.
				9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.....	Rare.
				4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Frequent.
15074	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	Rare.

NODOSARIA VERTEBRALIS (Batsch).

Plate 38, figs. 2, 3; plate 40, fig. 2.

Nautilus (Orthoceras) vertebralis BATSCH, Conch. des Seesandes, 1791, p. 3, No. 6, pl. 2, figs. 6a, b.

Nodosaria vertebralis H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 514, pl. 63, fig. 35; pl. 64, figs. 11-14.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 312, pl. 57, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 60, pl. 32, fig. 1.

This is one of the most abundant species in the area, occurring at some stations, as D5201, off southern Leyte, in great abundance. The specimens are well developed and show the distinct character of the sutures very clearly. The last-formed chamber is often much more constricted than the preceding ones, especially in adult specimens. The distribution includes the coast of Romblon; Tanon Strait; east coast of Negros; Sogod Bay, southern Leyte; off Burias; off Marinduque; Gulf of Davao; off southern Mindoro: Palawan Passage; between Samar and Leyte; Ragay Gulf, Luzon; and at various places off the coast of Mindanao.

The species seems to be confined to the Philippine Archipelago proper as far as the material examined shows, and to warm waters. The depths range from 26 to 554 fathoms (48 to 1,012 meters), with the average 129 fathoms (236 meters), and the bottom temperatures from 52.4° to 63.1° F. (11.3° to 17.2° C.), average 57.2° F. (14° C.). At the deepest station, 554 fathoms (1,012 meters), the bottom temperature was 52.8° F. (11.5° C.), a high temperature for this depth.

Nodosaria vertebralis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12153	U.S.N.M.	3	D5100...	° ' " ° ' "		° F.		
12150	U.S.N.M.	2	D5110...	14 17 15 N.; 120 32 40 E..	35	59.0	gy. s.....	Few.
12152	U.S.N.M.	1	D5112...	13 59 20 N.; 120 75 45 E..	135	52.4	dk. m.....	Few.
15045	U.S.N.M.	1	D5132...	13 48 22 N.; 120 47 25 E..	177		dk. gn. m....	Rare.
				Island off Panabutan	26		gn. m., s....	Rare.
15046	U.S.N.M.	6	D5133...	Point, N. 15 W. 0.3 mi. E.				
				Island off Panabutan	38		gn. m., s....	Frequent.
				Point, N. 52 E. 1.05 mi. E.				
15047	U.S.N.M.	10+	D5172...	6 03 15 N.; 120 35 30 E..	318		fine. s.....	Abundant.
12158	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78		fine. s.....	Rare.
15048	U.S.N.M.	3	D5189...	9 56 30 N.; 123 15 00 E..	300	62.8	gn. m.....	Few.
12151	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Rare.
15049	U.S.N.M.	2	D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s....	Few.
11601	} U.S.N.M.	10	D5220...	13 38 00 N.; 121 58 00 E..	50		sft. gn. m....	Abundant.
12157								
15050	U.S.N.M.	2	D5255...	7 03 00 N.; 125 39 00 E..	100		sft. m.....	Few.
15051	U.S.N.M.	7	D5261...	12 30 55 N.; 121 34 24 E..	145		s., m.....	Frequent.
15052	U.S.N.M.	5	D5272...	14 00 00 N.; 120 22 30 E..	118	57.4	m., sh., co. s.	Few.
15053	U.S.N.M.	1	D5311...	21 33 00 N.; 116 15 00 E..	88		crs. s., sh....	Rare.
12155	U.S.N.M.	1	} D5338...	11 33 45 N.; 119 24 45 E..	43		Co. s., m....	Frequent
15054	U.S.N.M.	3						
15055	U.S.N.M.	9						
12154	U.S.N.M.	5	D5382...	13 15 20 N.; 122 45 30 E..	128		co. s.....	Frequent.
15056	U.S.N.M.	3	D5481...	10 27 30 N.; 125 17 10 E..			m.....	Few.
12156	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	s., sh., g....	Few.
15057	U.S.N.M.	1	D5543...	8 47 15 N.; 123 35 00 E..	162	54.5	gn. m.....	Rare.
15058	U.S.N.M.	7	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	s.....	Rare.
							gn. m.....	Common.

NODOSARIA SPIROSTRIOLATA Cushman.

Plate 38, fig. 4.

Nodosaria spirostriolata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 656.

Description.—Test much elongate, very slightly tapering toward the initial end, which is broadly rounded, chambers numerous, short, elliptical in side view, sutures but slightly depressed, surface ornamented throughout with numerous (40–50) fine longitudinal costae, spirally arranged; apical end without a neck, broadly rounded.

Length 10 mm.

Distribution.—Type specimen (U.S.N.M. No. 9119) and others from *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.). It has also occurred at D5178, off Romblon, 78 fathoms (143 meters).

This is a large and striking species, with its very numerous fine spiral costae giving it a unique appearance.

Nodosaria spirostriolata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12297	U.S.N.M.	1	D5178...	° ' " ° ' "	78	° F.	fne. s.	Rare.
9119	} U.S.N.M.	7	D5236...	12 43 00 N.; 122 06 15 E..	494	41.2	fne. gy. s.	Common.
11602				8 50 45 N.; 126 26 52 E..				

NODOSARIA SUBTERTENUATA Schwager.

Plate 36, fig. 1.

Nodosaria subtertenuata SCHWAGER, *Novara-Exped.*, Geol. Theil., pt. 2, 1866, p. 235, pl. 6, fig. 74.—H. B. BRADY, *Rep. Voy. Challenger*, Zoology, vol. 9, 1884, p. 507, pl. 62, figs. 7, 8.

Description.—Test elongate, tapering, straight, composed of a few chambers, four to six, gradually increasing in size as added, elongate-elliptical, neck tapering and slender, wall distinctly roughened.

Length up to 2 mm.

Distribution.—One specimen, which was very typical, was obtained from *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.).

This species was described by Schwager from the upper Pliocene of Kar Nicobar. The *Challenger* specimens recorded by Brady were from the Ki Islands. Both the upper Pliocene of Kar Nicobar and the foraminiferal fauna of the Ki Islands seem to be closely related to the present foraminiferal fauna of the Philippines.

Nodosaria subtertenuata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15292	U.S.N.M.	1	D5236...	° ' " ° ' "	494	° F.	fne. gy. s.	Rare.
				8 50 45 N.; 126 26 52 E....		41.2		

NODOSARIA HIRSUTA d'Orbigny.

Plate 38, figs. 5 a, b, 6.

"*Orthoceratia quasi hispida*" SOLDANI, *Testaceographia*, vol. 2, 1798, p. 15, pl. 2, fig. P.

"*Orthoceratia hispida*" SOLDANI, *Testaceographia*, vol. 2, 1798, p. 36, pl. 11, figs. n-z, A, B.

Nodosaria hirsuta D'ORBIGNY, *Ann. Sci. Nat.*, vol. 7, 1826, p. 252, No. 7.—CUSHMAN, *Bull.* 71, U. S. Nat. Mus., pt. 3, 1913, p. 60, pl. 28, fig. 3.

Nodosaria hispida D'ORBIGNY, *Foram. Foss. Vienne*, 1846, p. 35, pl. 1 figs. 24, 25.—H. B. BRADY, *Rep. Voy. Challenger*, Zoology, vol. 9, 1884, p. 507, pl. 63, figs. 12-16.—FLINT, *Rep. U. S. Nat. Mus.*, 1897 (1899), p. 311, pl. 57, fig. 1.

The typical form of this species occurred at two stations—D5201, Sogod Bay, off southern Leyte, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.), and D5259, off northwestern Panay, 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.).

Nodosaria hirsuta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12179	U.S.N.M.	3	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s. m.	Rare.
12174	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob..	Rare.

NODOSARIA HIRSUTA d'Orbigny, var. *ACULEATA* d'Orbigny.

Plate 33, figs. 7, 8.

Nodosaria aculeata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 35, pl. 1, figs. 26, 27.

Description.—Variety with the test covered with coarse aculeate spines.

Distribution.—Specimens of this species occurred at several stations: Sogod Bay, southern Leyte; off northwestern Panay; China Sea, vicinity of southern Luzon; Palawan Passage, off northern Mindanao, and north of Tawi Tawi. The depth of these stations ranged from 201 to 554 fathoms (368 to 1,012 meters), average depth 356 fathoms (651 meters). Bottom temperatures ranged from 49.3° to 56.4° F. (9.6° to 13.5° C.), average temperature 52.2° F. (11.2° C.).

Nodosaria hirsuta var. *aculeata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12176	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s. m.	Few.
12179	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E...	302	49.3	gy. m.	Rare.
			D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s.	Frequent.
12180	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.	Rare.
12175	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E...	Rare.
12178	U.S.N.M.	1	D5571...	5 30 45 N.; 120 07 57 E...	340	52.3	s. sh.	Rare.

Genus *LINGULINA* d'Orbigny, 1826.

LINGULINA COSTATA d'Orbigny.

Plate 43, figs. 2a, b.

Lingulina costata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 62, pl. 3, figs. 1-5.—CARPENTER, PARKER, and JONES, *Introd. Foram.*, 1862, pl. 12, fig. 1.—KARRER, *Novara-Exped.*, *Geol. Theil.*, vol. 1, 1864, p. 75, pl. 16, fig. 3.—SCHWAGER, *Boll. R. Com. Geol. Ital.*, vol. 8, 1877, p. 25, pl. fig. 7.

Description.—Test composed of a few chambers in a uniserial arrangement, much overlapping, rapidly increasing in size as added,

whole test ovate, broadest toward the apertural end, slightly carinate laterally; surface ornamented by a series of longitudinal costae, those of each chamber independent of the adjacent ones; aperture long and narrow.

Diameter about 2 mm.

Distribution.—The figured specimen was obtained from *Albatross* station D5318, in 340 fathoms (622 meters), China Sea, vicinity of Formosa.

This is the first record for this species in the North Pacific.

Lingulina costata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15324	U.S.N.M.	1	D5318...	° ' " ° ' " 21 32 00 N.; 117 46 00 E...	340	° F.	s., br. Co.....	Rare.

LINGULINA GRANDIS Cushman.

Plate 43, figs. 1a, b.

Lingulina grandis CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 656.

Description.—Test large, compressed, early portion and in some specimens all but the final chamber distinctly keeled; chambers about eight in number, broad and short, somewhat inflated; sutures depressed, not noticed on the keel; wall smooth and shining, yellowish-brown or reddish-brown in color; aperture a very elongate slit extending half the width of the test.

Length up to 6 mm.; width 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9120) from *Albatross* station, D5268, in 170 fathoms (311 meters), Batangas Bay. Specimens were also obtained in material from D5178, in 78 fathoms (143 meters), vicinity of Romblon; D5217, in 105 fathoms (193 meters), between Burias and Luzon, bottom temperature 63.1° F. (17.2° C.); and D5589, in 260 fathoms (476 meters), Sibuko Bay, Borneo, bottom temperature 45.7° F. (7.6° C.).

This species in some ways resembles *L. carinata*, but it is several times as large and is differently shaped.

Lingulina grandis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
9120	U.S.N.M.	1	D5268...	° ' " ° ' " 13 42 00 N.; 120 57 15 E...	170	° F.	s., p.....	Rare.
11898	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.....	Rare.
11900	U.S.N.M.	3	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.....	Few.
11899	U.S.N.M.	1	D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fine. gy. s., gy. m.	Rare.

Genus FRONDICULARIA DeFrance, 1824.

There is a considerable development of this genus in the general region of the Philippines, probably better than in most recent gatherings, unless in the *Challenger* material dredged off the Ki Islands.

FRONDICULARIA INAEQUALIS Costa.

Plate 40, figs. 5, 6.

Frondicularia inaequalis COSTA, Mem. Accad. Sci. Napoli, vol. 2, 1855, p. 372, pl. 3, fig. 3.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 521, pl. 66, figs. 8-12.

Description.—Test compressed, irregularly elliptical, initial end usually narrow, bluntly pointed, proloculum subspherical, following chambers coiled in part or not at all, quickly giving place to typical V-shaped frondicularian chambers, highest in the center, thence gradually tapering to the pointed lower ends; test occasionally with a slight peripheral keel; sutures slightly depressed; aperture circular in the center of the periphery of the last-formed chamber; surface of test smooth and unornamented; wall translucent, thin.

Length 0.5 to 2 mm.

Distribution.—This delicate species has been previously recorded from the Atlantic and South Pacific, but not from the North Pacific. In the South Pacific, Brady in the *Challenger* Report recorded it from off New Zealand, 275 fathoms (503 meters); off Raine Island, Torres Strait, 155 fathoms (283 meters), and near the Ki Islands, 129 and 580 fathoms (236 and 1,061 meters). The two latter localities have already been mentioned as faunally closely related to the Philippines, so that it is natural to find *F. inaequalis* in this Philippine material. It has occurred at D5123, east coast of Mindoro, in 283 fathoms (517 meters), bottom temperature not given; and D5652, Gulf of Boni, 525 fathoms (927 meters), bottom temperature 41.2° F. (5.1° C.).

Frondicularis inaequalis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15334	U.S.N.M.	1	D5123...	° ' " ° ' "	283	° F.	gn. m.....	Rare.
15335	U.S.N.M.	1	D5652...	13 12 45 N.; 121 38 45 E... 4 35 00 S.; 121 23 06 E...	525	41.2	gn. m.....	Rare.

FRONDICULARIA PHILIPPINENSIS, new species.

Plate 39, figs. 1-3.

Description.—Test much compressed, comparatively large, roughly diamond-shaped, initial end bluntly pointed, the greatest breadth usually at about the lower third or fourth of the length, basal end

typically with a thin and narrow keel irregular at the edge or where well developed with several sharp toothlike processes; early chambers usually somewhat coiled, later ones typically bilateral and narrow not changing greatly in height, sutures usually nearly flush, of slightly clearer shell material; specimens occasionally flabelline throughout, last-formed chamber often with a square edge, with the peripheral angles slightly raised and the apertural end projecting in a slight knob, somewhat rounded; surface for the most part smooth and polished, the earlier chambers, however, typically ornamented with a series of irregular knobs or bosses with numerous irregular, short raised costae, running generally lengthwise of the test; aperture radiate, terminal.

Length up to 6 mm.

Distribution.—Type specimen (U.S.N.M. No. 15327) from D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.). It has also occurred in the following localities: D5259, off northwestern Panay; D5300, China Sea, vicinity southern Luzon; D5301, same locality, 208 fathoms (381 meters); D5425, Jolo Sea; D5429, eastern Palawan and vicinity; D5449, east coast of Luzon, San Bernardino Strait to San Miguel Bay.

Frondicularia philippinensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15327	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
15328	U.S.N.M.	9	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Common.
15329	U.S.N.M.	2	D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s....	Rare.
15330	U.S.N.M.	4	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s....	Few.
15331	U.S.N.M.	2	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Rare.
15332	U.S.N.M.	2	D5429...	9 41 30 N.; 118 50 22 E...	766	gn. m.	Rare.
15333	U.S.N.M.	1	D5449...	13 21 36 N.; 124 00 30 E...	300	Rare.

FRONDICULARIA ANNULARIS d'Orbigny.

Plate 39, fig. 5.

Frondicularia annularis d'ORBIGNY, *Foram. Foss. Vienne*, 1846, p. 59, pl. 2, figs. 44-47.

Description.—Test oval, much compressed, greatest width at the base, proloculum oval or elliptical, next few chambers not extending back as far as their predecessors, but soon equaling or exceeding the length backward; outer angle very broadly rounded, apertural end squarely truncated or sometimes with a projecting or rounded apertural knob; sutures but slightly depressed; early portions of test with numerous fine longitudinal raised costae parallel to the long axis of the test; remainder of the test smooth and polished.

Length equal to the diameter, about 3 mm.

Distribution.—The following localities are represented: East coast of Mindoro; China Sea off Formosa; Ragay Gulf, Luzon; and between Cebu and Leyte. It was not noted in the region to the southward. D'Orbigny's specimens were from the Miocene of Baden.

Fronicularia annularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12440	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m...	Rare.
12441	U.S.N.M.	1	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co...	Rare.
12437	U.S.N.M.	5	D5381...	13 14 15 N.; 122 44 25 E...	88	co. s.....	Common.
12439	} U.S.N.M.	8	D5382...	13 15 20 N.; 122 45 30 E...	128	m.....	Common.
15325								
12443	U.S.N.M.	1	D5408...	10 40 15 N.; 124 15 00 E...	159	55.4	gn. m.....	Rare.

FRONICULARIA ANNULARIS d'Orbigny, var. LONGISTRIATA, new variety.

Plate 39, fig. 6.

Description.—Variety differing from the typical in the form, which is much more elongate, narrower, the initial end pointed, and all or nearly all of the surface covered with the fine longitudinal costae.

Distribution.—Type specimen (U.S.N.M., No. 15326) from D5110, China Sea off southern Luzon, in 135 fathoms (247 meters), bottom temperature 59.0° F. (15° C.). It also occurred at D5201, Sogod Bay, southern Leyte Island, in 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.).

Chapman¹⁸ records a species of *Fronicularia* from off Great Barrier Island, Australia, as *F. reussi* Karrer, with the following notes:

Three examples of an ovate, striated *Fronicularia* were found off Great Barrier Island. They are almost exactly matched by Karrer's figured specimen from the Miocene of the Vienna basin. The narrowest of our specimens may also be compared with Karrer's *F. sculpta*, figured on the same plate as the above. These shells are obviously of the same type as the earlier-described *F. annularis* of d'Orbigny from the Miocene of Baden: this, however, is a generally broader form.

This appears to be the first occurrence of *F. reussi* in recent deposits.

Specimens, both of this form and the more typical form of *F. annularis*, were found in the Philippine material, but all are here referred to *F. annularis*, var. *longistriata*. It may be that Chapman's material is distinct from the Philippine material, but they appear to be very similar.

¹⁸ Trans. New Zealand Inst., vol. 38, 1905, p. 95, pl. 3, fig. 7.

Frondicularia annularis, var. *longistriata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15326	U.S.N.M.	1	D5110...	° ' " ° ' "	135	°F. 59.0	dk. gy. m....	Rare.
12442	U.S.N.M.	2	D5201...	13 59 20 N.; 129 75 45 E...	554	52.8	gy. s., m....	Rare.
15335	U.S.N.M.	1	F5523...	10 10 00 N.; 125 04 15 E...	Rare.
				8 48 44 N.; 123 27 35 E...				

FRONDICULARIA PLICATA, new species.

Plate 39, fig. 4; plate 40, fig. 4.

Description.—Test elongate, compressed, usually flabelline, border thickened, sutures irregular, periphery rounded, whole test irregularly triangular in shape; later chambers not extending back to the line of the previous chamber base, surface somewhat concave, smooth except for the initial portion, and the basal portion of the growing edge, which has a few large, irregular, rounded raised costae; aperture terminal, slightly projecting.

Length up to 4 mm. or more.

Distribution.—Type specimen (U.S.N.M. No. 15336) from D5301, China Sea, vicinity Hongkong, 208 fathoms (381 meters), bottom temperature 50.5° F. (10.2° C.).

The specimen figured (pl. 39, fig. 4) may be a smooth form of this species. It is from D5584, off Sibuko Bay, Borneo, and vicinity, 292 fathoms (535 meters), bottom temperature 44.3° F. (6.8° C.). It is however, much larger, being nearly 8 mm. in length.

Frondicularia plicata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15336	U.S.N.M.	2	D5301...	° ' " ° ' "	208	°F. 50.5	gy. m., s.	Rare.
15337	U.S.N.M.	1	D5584...	20 37 00 N.; 115 43 00 E...	292	44.3	fine. s., gy. m.	Rare.
				4 17 40 N.; 118 57 42 E...				

Genus TRIPLASIA Reuss, 1854.

TRIPLASIA TRICARINATA (d'Orbigny).

Vaginulina tricarinata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 258, No. 4; Modèles, No. 4.

Rhabdogonium tricarinatum H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 525, pl. 67, figs. 1-3.

Triplasia tricarinata CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 62, pl. 39, fig. 2.

Only two stations have been recorded in the Philippine material I have examined from which this species has been obtained. These

are *Albatross* stations D5301, in 208 fathoms (381 meters), China Sea, vicinity of Hongkong, bottom temperature 50.5° F. (10.2° C.); and D5638, in 517 fathoms (946 meters), entrance to Tifu Bay, Bouro Island.

Triplasia tricarinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11902	U.S.N.M.	1	D5301...	° ' " ° ' "	208	°F. 50.5	gy. m., s.....	Rare.
11901	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E...	517	fne. gy. s.....	Rare.

Genus *CRISTELLARIA* Lamarck, 1812.

CRISTELLARIA CULTRATA (Montfort).

"Cornu Hammonis" PLANCUS, Conch. Min., 1760, p. 120, pl. 1, fig. 12.

"Nautili (*Lenticulae marginatae*)" SOLDANI, Testaceographia, vol. 1, pt. 1, 1789, p. 54, pl. 33, fig. B.

Robulus cultratus MONTFORT, [?] Conch. Syst., vol. 1, 1808, p. 214, 54^e genre.

Robulina cultrata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 287, No. 1; Modèles, 1826, No. 82; Foram. Foss. Vienne, 1846, p. 96, pl. 4, figs. 14, 15.

Cristellaria cultrata PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 344, pl. 13, figs. 17, 18; pl. 16, fig. 5.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 550, pl. 70, figs. 4, 5, 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 64, pl. 29, fig. 4.

This species is well distributed in the area, including not only the Philippine Archipelago, but the China Sea, and the area to the south, including Patente Strait and Buton Strait.

The stations range in depth from 37 to 700 fathoms (68 to 1,280 meters), with the average depth 266 fathoms (486 meters), and the average of the bottom temperatures when given 49.1° F. (9.5° C.).

Cristellaria cultrata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		°F.		
15255	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318	fne. s., sh....	Rare.
11864	U.S.N.M.	2	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.....	Few.
11865	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s.....	Rare.
11859	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
15256	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	fne. bk. s.....	Rare.
15257	U.S.N.M.	1	D5291...	13 29 40 N.; 121 00 45 E...	173	51.5	dk. gy. s.....	Rare.
11860	U.S.N.M.	5	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.....	Few.
11862	U.S.N.M.	2	D5331...	15 36 45 N.; 119 47 45 E...	178	54.7	s., sh., m....	Few.
15258	U.S.N.M.	5	D5374...	13 46 45 N.; 121 35 08 E...	190	gy. m.....	Rare.
11861	U.S.N.M.	5	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Few.
11863	U.S.N.M.	3	D5454...	13 12 00 N.; 123 50 30 E...	153	Few.
15259	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Rare.
15261	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.....	Rare.
15663	U.S.N.M.	1	D5637...	3 53 20 S.; 126 43 00 E...	700	gy. m.....	Rare.
15260	U.S.N.M.	10+	D5642...	4 31 40 S.; 122 49 42 E...	37	gy. m.....	Common.

CRISTELLARIA CULTRATA (Montfort), var. DECORATA Cushman.

Cristellaria cultrata (MONTFORT), var. *decorata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 656.

Description.—Variety differing from the typical in the addition of a distinct type of ornamentation, consisting of a large umbilical knob above the umbilical region, and the sutures with costae which are gradually larger toward the periphery.

Distribution.—Type specimen (U.S.N.M. No. 9121) from *Albatross* station 5113 in 159 fathoms (290 meters), China Sea off southern Luzon. It also occurred at the following *Albatross* stations: D5121, in 108 fathoms (198 meters), east coast of Mindoro; D5222, in 195 fathoms (356 meters), between Marinduque and Luzon, bottom temperature 52.8° F. (11.5° C.); D5260, in 234 fathoms (428 meters), off southeastern Mindoro, bottom temperature 51.4° F. (10.7° C.); and D5590, in 310 fathoms (567 meters), Sibuko Bay, Borneo.



FIG. 9.—CRISTELLARIA CULTRATA, VAR. DECORATA CUSHMAN. SIDE VIEW, X 50. FROM STATION D5113.

Cristellaria cultrata, var. *decorata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13230 9121	} U.S.N.M.	4	D5113...	13 51 30 N.; 120 50 30 E..	159	dk. gn. m....	Rare.
11867		5	D5121...	13 27 20 N.; 121 17 45 E..	108	dk. gn. m....	Rare.
11868	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.	Rare.
11840	U.S.N.M.	1	D5260...	12 25 35 N.; 121 31 35 E..	234	51.4	gn. m., s.....	Rare.
11866	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s.....	Rare.

CRISTELLARIA RENIFORMIS d'Orbigny.

Cristellaria reniformis D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 88, pl. 3, figs. 39, 40.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 539, pl. 70, figs. 3a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 65, pl. 30, fig. 4; pl. 33, fig. 1.

Single specimens referable to this species were found at the following *Albatross* stations: D5439, in 940 fathoms (1,719 meters), west coast of Luzon, bottom temperature 36.7° F. (2.6° C.); D5613, in 752 fathoms (1,375 meters), Gulf of Tomini, Celebes; and D5668 in 901 fathoms (1,648 meters), Macassar Strait, bottom temperature 38.2° F. (3.4° C.).

These stations are all in comparatively deep water, with a cold bottom temperature, and compare rather well in conditions with most of those recorded in the *Challenger* report for this species in these two respects.

Cristellaria reniformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11773	U.S.N.M.	1	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.....	Rare.
11770	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Rare.
11771	U.S.N.M.	1	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	Rare.

CRISTELLARIA ARTICULATA (Reuss).

Plate 57, figs. 2-4.

Robulina articulata REUSS, Sitz. Akad. Wiss. Wien, vol. 48, 1863, p. 53, pl. 5, fig. 62.

Cristellaria articulata REUSS, Sitz. Akad. Wiss. Wien, vol. 62, 1870, p. 483.—
H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 547, pl. 69, figs. 10-12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 65, pl. 31, fig. 1.

Although few specimens have occurred at any one station, nevertheless numerous stations have yielded records for this species: China Sea, off Luzon; Sogod Bay, southern Leyte; off western Luzon; between Luzon and Marinduque; eastern Palawan; between Leyte and Mindanao; between Samar and Leyte; between Siquijor and Bohol; north of Tawi Tawi; Gulf of Tomini, Celebes, and Molucca Sea. With the exception of the last, all these stations are within the Archipelago. The depths range from 24 to 766 fathoms (44 to 1,400 meters), with the average depth 368 fathoms (673 meters), the average bottom temperature being 53.9° F. (12.1° C.).

A figure is given of a very attenuate specimen which was uncoiled almost from the beginning. It is a rather unusual form for this species.

Cristellaria articulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11824	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m....	Rare.
11822	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	sft. gn. m....	Rare.
15224	U.S.N.M.	1	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m....	Rare.
11823	U.S.N.M.	1	D5331...	15 36 45 N.; 119 47 45 E..	178	54.7	s. sh., m....	Few.
11821	U.S.N.M.	4	D5429...	9 37 30 N.; 118 48 30 E..	706	gn. m.....	Few.
11826	U.S.N.M.	2	D5493...	9 04 00 N.; 125 20 00 E..	478	52.1	gn. m.....	Rare.
11825	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob..	Rare.
11819	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E..	330	52.3	co. s.....	Rare.
11820	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Rare.
15225	U.S.N.M.	1	D5640...	4 27 00 S.; 122 55 40 E..	24	s., brk. sh.	Rare.
15172	U.S.N.M.	1	Tayabas Gulf, P. I.....

CRISTELLARIA ROTULATA (Lamarck).

"Cornu Hammonis seu Nautili" PLANCUS, Conch. Min., 1739, p. 13, pl. 1, fig. 3.
Lenticulites rotulata LAMARCK, Ann. Mus., vol. 5, 1804, p. 188, No. 3; vol. 8, 1806, pl. 62, fig. 11.

Cristellaria rotulata D'ORBIGNY, H. B. Brady, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 547, pl. 69, figs. 13a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 66, pl. 35, fig. 3.

Typical specimens of this species were obtained from numerous stations in the area; in the China Sea off Hongkong; between Burias and Luzon; off Marinduque; west and east coasts of Luzon; between Samar and Leyte; between Negros and Siquijor; north of Tawi Tawi; Sibuko Bay, Borneo; and Gulf of Boni.

These stations range in depth from 20 to 565 fathoms (37 to 1,033 meters), with an average depth of 357 fathoms (653 meters). The average bottom temperature where given was 50.3° F. (10.1° C.).

Cristellaria rotulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11777	U.S.N.M.	4	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m....	Few.
11778	U.S.N.M.	7	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Frequent.
11779	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E...	29	crs. s....	Rare.
16200	U.S.N.M.	10+	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s....	Common.
15173	U.S.N.M.	2	D5377...	13 26 00 N.; 122 19 00 E...	400	49.6	sft. gn. m....	Few.
11776	U.S.N.M.	5	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m....	Frequent.
15174	U.S.N.M.	1	D5460...	13 32 30 N.; 123 58 06 E...	565	gy. m....	Rare.
15175	U.S.N.M.	5	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m....	Frequent.
11780	U.S.N.M.	10	D5470...	13 37 30 N.; 123 41 09 E...	560	m.	Common.
11788			D5478...	10 46 24 N.; 125 16 30 E...	57	sh.	Rare.
			D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s....	Frequent.
15176	U.S.N.M.	2	D5569...	5 33 15 N.; 120 30 15 E...	303	52.3	co. s....	Few.
15177	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fine. s., glob.	Rare.
			D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fine. gy. s., gy. m.	Common.
			D5591...	4 11 48 N.; 118 38 20 E...	260	Common.
11787	U.S.N.M.	6	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m....	Common.

CRISTELLARIA ROTULATA (Lamarck), var. UMBONATA Cushman.

Plate 43, fig. 3.

Cristellaria rotulata (LAMARCK), var. *umbonata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 656.

Description.—Variety with the umbonal region greatly produced by a large mass of clear shell substance, standing out clearly above the rest of the test.

Distribution.—Type specimen (U.S.N.M. No. 9122) from *Albatross* station D5217, in 105 fathoms (193 meters), between Burias and Luzon, bottom temperature 63.1° F. (17.2° C.). Besides at the type station the variety was found at the following *Albatross* stations: D5184, in

565 fathoms (1,033 meters), between Panay and Negros, bottom temperature 49.8° F. (9.8° C.); D5185, in 638 fathoms (1,167 meters), same region and same bottom temperature; D5222, in 195 fathoms (356 meters), between Marinduque and Luzon, bottom temperature 52.8° F. (11.5° C.); D5425, in 495 fathoms (904 meters), Jolo Sea, bottom temperature 49.4° F. (9.7° C.); and D5470, east coast of Luzon, 560 fathoms (1,024 meters).

The variety is much less common than the typical form, but where it occurs seems to be distinct. At only one station was it associated with the typical form.

Cristellaria rotulata, var. *umbonata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11795	U.S.N.M.	1	D5184...	° ' " ° ' "	565	° F.		
11786	U.S.N.M.	1	D5185...	10 18 30 N.; 122 23 30 E...	638	49.8	gn. m.....	Rare.
13232 9122	U.S.N.M.	7	D5217...	10 05 45 N.; 122 18 30 E...	105	63.1	crs. gy. s....	Frequent.
11796	U.S.N.M.	3	D5222...	13 20 00 N.; 123 14 15 E...	195	52.8	gn. m.....	Few.
17775	U.S.N.M.	1	D5425...	13 38 30 N.; 121 42 45 E...	495	49.4	gy. m., co. s.	Rare.
11785	U.S.N.M.	2	D5470...	9 37 45 N.; 121 11 00 E...	560	m.....	Rare
				13 37 30 N.; 123 41 09 E...				

CRISTELLARIA ORBICULARIS (d'Orbigny).

"Nuclei conico rotundati" SOLDANI, Testaceographia, vol. 2, 1798, App., p. 138, pl. 1, fig. 12. p, P.

Robulina orbicularis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 288, pl. 15, figs. 8, 9.

Cristellaria orbicularis H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 549, pl. 69, fig. 17.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 67, pl. 36, figs. 4, 5.

This is one of the commonest species of *Cristellaria* in the region, being found in a large number of stations in the Philippine Archipelago proper; in the China Sea, off Formosa and off Hongkong; southward at several stations off Sibuko Bay, Borneo; in Molucca Passage; and between Gillolo and Kayoa Islands.

The depths range from 43 to 565 fathoms (79 to 1,033 meters), the average depth being 273 fathoms (499 meters). The average bottom temperature for the stations when recorded is 50.7° F. (10.4° C.).

The *Challenger* specimens were mostly from the South Pacific. Several distinct varieties occur in the region.

Cristellaria orbicularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
15226	U.S.N.M.	10+	D5113...	13 51 30 N.; 120 50 30 E...	159	dk. gn. m...	Common.
15227	U.S.N.M.	10+	D5114...	13 36 11 N.; 120 45 26 E...	340	fne. s.	Common.
15228	U.S.N.M.	6	D5198...	9 40 50 N.; 123 39 45 E...	220	53.9	gn. m.	Common.
11698	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m...	Few.
15229	U.S.N.M.	5	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m.	Few.
11699	U.S.N.M.	1	D5238...	7 34 45 N.; 126 38 15 E...	380	43.0	gn. m.	Rare.
15230	U.S.N.M.	10+	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Frequent.
15231	U.S.N.M.	6	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fne. s.	Frequent.
		5	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s...	Frequent.
15232	U.S.N.M.	6	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s...	Few.
15233	U.S.N.M.	7	D5291...	13 29 40 N.; 121 00 45 E...	173	51.5	fne. bk. s...	Frequent.
11700	U.S.N.M.	7	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s...	Frequent.
11702	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E...	150	53.6	s.	Rare.
			D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. g...	Few.
15234	U.S.N.M.	1	D5331...	15 36 45 N.; 119 47 45 E...	178	54.7	co. s., m...	Rare.
11701	U.S.N.M.	4	D5338...	11 33 45 N.; 119 24 45 E...	43	co. s., m...	Few.
15235	U.S.N.M.	6	D5369...	13 48 00 N.; 121 43 00 E...	106	bk. s.	Frequent.
15236	U.S.N.M.	2	D5371...	13 49 40 N.; 121 40 15 E...	83	gn. m.	Few.
			D5410...	10 28 45 N.; 124 05 30 E...	385	gn. m.
15237	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.	Rare.
15238	U.S.N.M.	1	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Rare.
11703	U.S.N.M.	5	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.	Few.
15239	U.S.N.M.	1	D5447...	13 25 00 N.; 123 46 18 E...	310	45.3	gn. m.	Rare.
11740	U.S.N.M.	10	D5449...	13 21 36 N.; 124 00 30 E...	300	Frequent.
15240	U.S.N.M.	2	D5453...	13 12 00 N.; 123 49 18 E...	146	Few.
11738	U.S.N.M.	1	D5460...	13 32 30 N.; 123 58 06 E...	575	gy. m.	Rare.
15242	U.S.N.M.	1	D5478...	10 46 24 N.; 125 16 30 E...	57	sh.	Rare.
			D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy. m., fne. s.	Frequent.
15243	U.S.N.M.	4	D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gn. m., glob.	Few.
11739	U.S.N.M.	10	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Common.
15244	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s...	Common.
			D5567...	5 48 00 N.; 120 33 45 E...	263	52.0	fne. s.	Common.
15245	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.	Rare.
15241	U.S.N.M.	6	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fne. s., glob.	Common.
15246	U.S.N.M.	1						
15247	U.S.N.M.	2	D5574...	5 30 45 N.; 120 07 57 E...	Few.
15248	U.S.N.M.	6	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.	Few.
			D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fne. gy. s., gy. m.	Common.
			D5591...	4 11 48 N.; 118 38 20 E...	260	Few.
			D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.	Frequent.
15249	U.S.N.M.	3	D5619...	0 35 00 S.; 127 14 40 E...	435	fne. gy. s., m.	Rare.
15250	U.S.N.M.	7	D5625...	0 07 00 N.; 127 28 00 E...	230	gy. m., fne. s.	Common.

CRISTELLARIA ORBICULARIS (d'Orbigny), var. SUBCARINATA Cushman.

Plate 44, fig. 1.

Cristellaria orbicularis (d'ORBIGNY), var. *subcarinata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 657.

Description.—Variety differing from the typical by having instead of a thin broad keel a very narrow one, which is thickened with a rounded periphery, otherwise like the typical.

Distribution.—Type specimen (U.S.N.M. No. 9123) from *Albatross* station D5654 in 805 fathoms (1,472 meters), Gulf of Boni, bottom temperature 38.3° F. (3.5° C.). Specimens were also obtained from material dredged at the following stations: D5465, in 500 fathoms (914 meters), and D5467 in 480 fathoms (878 meters), east coast of Luzon, and D5582, in 890 fathoms (1,628 meters), off Darvel Bay, Borneo, bottom temperature 38.3° F. (3.5° C.).

From the data given the variety seems to be developed in the deeper, colder waters of the region.

Cristellaria orbicularis, var. *subcarinata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
11737	U.S.N.M.	1	D5465...	° ' " ° ' "	500	° F.		Rare.
11704	U.S.N.M.	7	D5467...	13 39 42 N.; 123 40 39 E...	480		gy. m.....	Frequent.
11705	U.S.N.M.	1	D5582...	4 19 54 N.; 118 58 38 E...	890	38.3	gy. m., fne. s.	Rare.
15252								
13233	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3	Rare.
9123								
15253								

CRISTELLARIA ORBICULARIS (d'Orbigny), var. SUBUMBONATA Cushman.

Plate 44, fig. 3.

Cristellaria orbicularis (D'ORBIGNY), var. *subumbonata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 657.

Description.—Variety differing from the typical in its development of ornamental characters, having a cluster of closely set raised knobs over the umbilical region, giving a decidedly umbonate appearance in face view, remainder of the test smooth.

Distribution.—Type specimen (U.S.N.M. No. 9124) from *Albatross* station D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.). Specimens were also found in material from *Albatross* stations D5110, China Sea off southern Luzon, 135 fathoms (247 meters), bottom temperature 59.0° F. (15.0° C.); D5152, Sulu Archipelago, 34 fathoms (62 meters); and D5438, in 297 fathoms (543 meters), west coast of Luzon, bottom temperature 46.2° F. (7.8° C.)

Cristellaria orbicularis, var. *subumbonata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11707	U.S.N.M.	1	D5110...	° ' " ° ' "	135	° F.		Rare.
13229	U.S.N.M.	8	D5152...	5 22 55 N.; 120 15 45 E...	34	59.0	dk. gy. m....	Rare.
13234	U.S.N.M.	3	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
9124								
11706	U.S.N.M.	2	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Rare.

CRISTELLARIA ORBICULARIS (d'Orbigny), var. PAPILLATA Cushman.

Plate 44, fig. 2.

Cristellaria orbicularis (D'ORBIGNY), var. *papillata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 657.

Description.—Variety with the keel reduced in width and the earlier portion with an ornamentation consisting of fine papillae rather evenly scattered over the surface, in later growth gradually becoming peripheral and in the last-formed chambers gradually becoming obsolete.

Distribution.—Type specimen (U.S.N.M. No. 9125) from *Albatross* station D5152, in 34 fathoms (62 meters), Sulu Archipelago, Tawi Tawi group. Specimens were also obtained from *Albatross* station D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.).

This is a rather beautifully ornamented variety of this species and does not seem to be at all common in the region.

Cristellaria orbicularis, var. *papillata*—Material examined.

Cat. No.	Coll. of—	No. of specim. s.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9125 15251	} U.S.N.M.	5	D5152...	5 22 55 N.; 120 15 45 E...	34	°F.	wh. s.....	Frequent.
			D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	

CRISTELLARIA VORTEX (Fichtel and Moll).

“Nautili globuli” SOLDANI, Testaceographia, vol. 1, pt. 1, 1879, p. 66, pl. 59, fig. *tt*.

Nautilus vortex FICHEL and MOLL, Test. Micr., 1803, p. 33, pl. 2, figs. *d-i*.

Polystomella vortex BLAINVILLE, Man. de Malac., 1825, p. 389.

Robulina vortex D'ORBIGNY, Ann. Sci. Nat., vol. 7., 1826, p. 288, No. 4.

Cristellaria vortex PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 4, vol. 8, 1871, p. 240, pl. 10, fig. 82.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 548, pl. 69, figs. 14–16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 68, pl. 32, fig. 3.

Very fine specimens of this species have occurred in the *Albatross* material of the following localities: Off Romblon; between Burias and Luzon; Ragay Gulf and east coast of Luzon; between Samar and Leyte; north of Tawi Tawi; Buton Strait; and Gulf of Boni. These stations range in depth from 24 to 976 fathoms (44 to 1,785 meters) with the average 308 fathoms (564 meters); the bottom temperatures with an average of 57.8° F. (14.4° C).

Brady's *Challenger* material with one exception was from the Southern Pacific. The North Pacific material which I had, was from the Hawaiian Islands and Guam, and apparently this species is most at home in fairly warm waters of the Pacific.

Cristellaria vortex—Material examined.

Cat. No.	Coll. of—	No. of specim. s.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11763	} U.S.N.M.	8	D5178...	12 43 00 N.; 122 06 15 E...	78	° F.	inc. s.....	Frequent.
11764			D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Common.
11765	} U.S.N.M.	10+	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Few.
11766			D5382...	13 53 00 N.; 120 26 45 E...	128	m.....	Few.
11767	} U.S.N.M.	1	D5470...	13 37 30 N.; 123 41 09 E...	560	m.....	Rare.
15215			D5495...	9 06 30 N.; 125 00 20 E...	976	52.3	gy. m.....	Few.
	} U.S.N.M.	4	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	inc. s., glob.	Rare.
			D5640...	4 27 00 S.; 122 55 40 E...	24	s. brk., sh...	Few.
11768	} U.S.N.M.	3	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Rare.
15216								

CRISTELLARIA CONVERGENS Bornemann.

Plate 44, figs. 4a, b.

Cristellaria convergens BORNEMANN, Zeitschr. deutsch. Geol. Ges., vol. 7, 1885, p. 327, pl. 13, figs. 16, 17.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 546, pl. 69, figs. 6, 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 68, pl. 34, fig. 3.

This species has occurred at five stations within the area as follows: *Albatross* stations D5178, in 78 fathoms (143 meters), off Romblon; D5215, in 604 fathoms (1,105 meters), east of Masbate Island, bottom temperature 50.5° F. (10.2 C.); D5220, in 50 fathoms (91 meters), between Marinduque and Luzon; D5469, in 500 fathoms (914 meters), east coast of Mindanao; and D5523, off northern Mindanao. It was also found off Jolo Jolo.

At none of these stations were there but a few specimens of the species.

Cristellaria convergens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11876	U.S.N.M.	1	D5178...	° / " ° / "	78	° F.	fne. s.....	Rare.
11871	U.S.N.M.	1	D5215...	12 31 30 N.; 123 35 24 E..	604	50.5	gn. m.....	Rare.
11874	U.S.N.M.	8	D5220...	13 38 00 N.; 121 58 00 E..	50		sft. gn. m.....	Few.
11873	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500		gn. m.....	Rare.
11875	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E..			Rare.
15295	U.S.N.M.	3	Jolo Jolo, P. I.....			Few.

CRISTELLARIA GIBBA d'Orbigny.

Plate 45, fig. 1.

Cristellaria gibba D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères" p. 63, pl. 7, figs. 20, 21.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 546, pl. 69, figs. 8, 9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 69, pl. 35, fig. 1.

Only a very few records of this species in the area were noted. It occurred at *Albatross* stations D5121, east coast of Mindoro, 108 fathoms (198 meters); D5338, Mindoro Strait, 310 fathoms (567 meters), bottom temperature 73.8° F. (23.2° C.); D5333, in 43 fathoms (78 meters), and D5348, in 375 fathoms (686 meters), both in Palawan Passage, the latter station with a bottom temperature of 56.4° F. (13.5 C.); D5481, in 61 fathoms (112 meters), between Samar and Leyte, vicinity of Suriago Strait; and D5650 in 540 fathoms (988 meters), in the Gulf of Boni.

Cristellaria gibba—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11694	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E..	108	dk. gn. m...	Rare.
12013	U.S.N.M.	4	D5333...	11 33 45 N.; 119 24 45 E..	43	Co., s., m...	Rare.
11695	U.S.N.M.	1	D5338...	12 26 30 N.; 120 37 45 E..	310	73.8	Co., s., m...	Few.
12014	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	Co., s., m...	Few.
11697	U.S.N.M.	1	D5481...	10 27 30 N.; 125 17 10 E..	61	s., sh., g., g.	Rare.
11696	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Rare.

CRISTELLARIA VARIABILIS Reuss.

Cristellaria variabilis REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1849, p. 369, pl. 46, figs. 15, 16.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 541, pl. 68, figs. 11–16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 70, pl. 36, figs. 1–3.

Specimens of this species have been noted in material from the following *Albatross* stations in the area: D5201, Sogod Bay, southern Leyte, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5242, in 191 fathoms (350 meters), off Pujada Bay, bottom temperature 64.1° F. (17.8° C.); D5300, China Sea, off southern Luzon, 265 fathoms (485 meters); D5301, in 208 fathoms (381 meters), China Sea off Hongkong, bottom temperature 50.5° F. (10.2° C.); D5420, in 127 fathoms (233 meters), between Cebu and Bohol, bottom temperature 59° F. (15° C.); D5445, east coast of Luzon, 383 fathoms (699 meters), bottom temperature 44.3° F. (6.8° C.); D5601 in 765 fathoms (1,399 meters), also in the Gulf of Tomini, Celebes; and D5613 in 752 fathoms (1,375 meters) also in the Gulf of Tomini.

This is a rather distinctive species, easily distinguished by its few chambers in each volution and the peculiar fimbriate appearance of the area about the aperture, especially in well-developed specimens.

Cristellaria variabilis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11761	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Rare.
			D5242...	6 51 53 N.; 126 14 10 E..	191	64.1	sif. gy. m....	Rare.
11759	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.
11760	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m., s....	Rare.
11756	U.S.N.M.	1	D5420...	9 49 35 N.; 123 45 00 E..	127	59.0	Rare.
11766	U.S.N.M.	2	D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s....	Rare.
11758	U.S.N.M.	1	D5601...	1 13 10 N.; 125 17 05 E..	765	s., glob., ptr.	Rare.
11757	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Rare.

CRISTELLARIA TRICARINELLA Reuss.

Plate 50, figs. 3a, b.

Cristellaria tricarinella REUSS, Sitz. Akad. Wiss. Wien, vol. 46, 1862, p. 68, pl. 7, fig. 9; pl. 12, figs. 2-5.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 540, pl. 68, figs. 3, 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 72, pl. 34, figs. 1, 2.

Brady records this species from three *Challenger* stations in the western Pacific of which one is off the Philippines in 95 fathoms (174 meters). It has occurred at the following stations in the *Albatross* Philippine dredgings: D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte, bottom temperature, 52.8° F. (11.5° C.); D5211, in 155 fathoms (283 meters), east of Masbate Island, bottom temperature, 56.6° F. (13.7° C.); D5268, in 170 fathoms (311 meters), off Batangas Bay; D5318, in 340 fathoms (622 meters), China Sea off Formosa; D5592, in 305 fathoms (558 meters), off Sibuko Bay, Borneo, bottom temperature, 43.3° F. (6.1° C.).

This does not in all respects, especially those of the early development, agree very closely with the species of Reuss, and its limited area of distribution in the Indo-Pacific seems to indicate that it may be a distinct species.

The records outside the Philippine region include the *Challenger* records; off Raine Island, Torres Strait, 155 fathoms (283 meters), and off the west coast of New Zealand, 150 fathoms (270 meters) (Brady); off Great Barrier Island, New Zealand, 110 fathoms (200 meters); north of Enderby Island, New Zealand, 85 fathoms (153 meters); 40 miles south of Cape Wiles, South Australia (Chapman); from off Mauritius; off the west coast of Australia (Egger); and from off Ceylon (Dakin).

Cristellaria tricarinella—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
11750	U. S. N. M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.	Rare.
11753	U. S. N. M.	2	D5211...	11 51 35 N.; 124 14 00 E...	155	56.6	gn. m., s.	Rare.
11750	U. S. N. M.	1	D5268...	13 42 00 N.; 120 57 15 E...	170	p., s., sh.	Rare.
11752	U. S. N. M.	2	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co	Rare.
11749	U. S. N. M.	1	D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.	Rare.

CRISTELLARIA TRICARINELLA Reuss, var. SPINIPES Cushman.

Cristellaria tricarinella REUSS, var. *spinipes* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 72, pl. 33, fig. 2.

This variety was represented at three of the *Albatross* stations as follows: D5178, in 78 fathoms (143 meters), off Romblon; D5301, in 208 fathoms (381 meters), China Sea off Hongkong, bottom tempera-

ture 50.5° F. (10.2° C.); and D5590, in 310 fathoms (567 meters), Sibuko Bay, Borneo, bottom temperature 44.3° F. (6.8° C.).

The variety was described from *Albatross* station D4900 off southern Japan, a station with a bottom temperature of 52.9° F. (11.6° C.), where numerous species of the Philippine region seem to reach their northern limit.

Cristellaria tricarinella, var. *spinipes*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11747	U.S.N.M.	1	D5178...	° ' " ° ' "	78	° F.	fine.s.....	Rare.
11748	U.S.N.M.	1	D5301...	12 43 00 N.; 122 06 15 E...	208	50.5	gy.m.,s.....	Rare.
11746	U.S.N.M.	3	D5590...	20 37 00 N.; 115 43 00 E... 4 10 50 N.; 118 39 35 E...	310	44.3	gn.m.,s.....	

CRISTELLARIA CALCAR (Linnaeus).

Plate 45, figs. 2a, b.

"*Nautilus minimus non umbilicatus*" GAULTIERI, Index Test., 1742, pl. 19, fig. C.

"*Nautili (Lenticulæ radiatæ)*" SOLDANI, Testaceographia, vol. 1, pt. 1, 1789, p. 54, pl. 33, figs. aa, bb.

Nautilus calcar LINNAEUS, Syst. Nat., ed. 12, 1767, p. 1162, No. 272; (Gmelin's) ed. 13, 1788, p. 3370, No. 2.

Robulina calcar D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 99, pl. 4, figs. 18-20.

Cristellaria calcar PARKER and JONES, Ann. Mag. Nat. Hist., ser. 2, vol. 19, 1857, p. 289, pl. 10, figs. 10-12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 551, pl. 70, figs. 9-12 (not 13-15).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 72, pl. 32, fig. 4.

The great mass of material of *Cristellaria* which has been obtained from the Philippine material has made possible the demarcation of various species and the segregation of new forms with a much surer basis than would be the case with a few specimens. The abundance of specimens has resulted almost invariably in the clearness of lines of separation rather than in the obliteration of such lines. This has been the case here and the species *C. calcar* has been restricted to the test, comparatively small in size, with a few chambers in each whorl and with long acicular spines, one to each chamber, radially placed on the periphery, the peripheral carina being very narrow.

Such a test as described above occurred in more or less numbers in material from a number of stations in the area. Most of these stations are in the Philippine Archipelago proper, but a few are outside in the China Sea off Formosa, and off Sibuko Bay, Borneo. In the restricted Philippine area it has occurred off the east coast of Mindoro; Sogod Bay, southern Leyte; east of Masbate; between Marinduque and Luzon; between Siquijor and Bohol; and especially at various stations about Marinduque.

These stations range in depth from 50 to 554 fathoms (91 to 1,012 meters), the average depth being 239 fathoms (437 meters). The average temperature of the stations where this was recorded is 51.5° F. (10.8° C.).

Brady records this species off the Philippines in 95 fathoms (174 meters).

Cristellaria calcar—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15217	U.S.N.M.	7	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m....	Frequent.
11798	U.S.N.M.	7	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Frequent.
15218	U.S.N.M.	1	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m....	Rare.
11797	U.S.N.M.	5	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m....	Common.
11801	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s....	Rare.
			D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	
11802	U.S.N.M.	5	D5369...	13 48 00 N.; 121 43 00 F...	106	bk. s.....	Common.
15219	U.S.N.M.	1	D5370...	13 44 15 N.; 121 42 30 E...	159	54.3	sft. m.....	Rare.
11800	U.S.N.M.	1	D5371...	13 49 40 N.; 121 40 15 E...	83	gn. m.....	Rare.
15308	U.S.N.M.	7	D5382...	13 15 20 N.; 122 45 30 E...	128	m.....	Common.
11803	U.S.N.M.	2	D5465...	13 39 42 N.; 123 40 39 E...	500	gy. m.....	Few.
11220	U.S.N.M.	3	D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gy. m., glob.	Few.
11799	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Few.

CRISTELLARIA CALCARATA Cushman.

Plate 45, fig. 3.

Cristellaria calcarata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 657.

Description.—Test biconvex, but much compressed, closely coiled, about seven or eight chambers in the last-formed coil; sutures curved, and marked by raised ridges running to the umbonal region, where they unite in a central boss of clear shell material, which is typically excavated in the center, forming a ring of material; periphery with a narrow keel from which are rowel-like spines, the two sides often unequal in their angles; apertural face flattened or even concave, aperture radiate.

Diameter about 2 mm.

Distribution.—Type specimen (U.S.N.M. No. 9126) from *Albatross* station D5370, in 159 fathoms (291 meters), off Marinduque Island, bottom temperature 54.3° F. (12.3° C.). Besides this station it was found at various stations in the China Sea off southern Luzon; Sogod Bay, southern Leyte; off southeastern Mindoro; off northern Mindanao; and China Sea off Formosa.

The range of depths of these stations is from 28 to 554 fathoms (51 to 1,012 meters), the average depth 262 fathoms (480 meters). The average bottom temperature where given is 54.1° F. (12.2° C.).

This species is at once distinguished from *C. calcar* by its ornamentation, from *C. submamilligera* by its spines, and from *C. echinata*

by its type of spines and its lack of bosses or papillae. It is also more compressed than most of the species of *Cristellaria* with peripheral spines. It seems in the mass of material examined to be a very distinctive species.

Cristellaria calcarata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11809	U.S.N.M.	5	D5110...	13 59 20 N.; 120 75 45 E...	135	59.0	dk. gy. m...	Frequent.
11812				13 51 30 N.; 120 50 30 E...	159	dk. gy. m...	Frequent.
11810	U.S.N.M.	8	D5113...	13 51 30 N.; 120 50 30 E...	159	dk. gy. m...	Frequent.
11808	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gn. s., m....	Rare.
15296				7 22 12 N.; 124 12 15 E...	28	m.....	Rare.
15297	U.S.N.M.	1	D5257...	7 22 12 N.; 124 12 15 E...	28	m.....	Rare.
11806	U.S.N.M.	2	D5260...	12 25 35 N.; 121 31 35 E...	234	51.4	gn. m., s....	Few.
11807	U.S.N.M.	2	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s. sh.....	Few.
9126	U.S.N.M.	1	D5370...	13 44 15 N.; 121 42 30 E...	159	54.3	sft. m.....	Rare.
11811	U.S.N.M.	1	D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy. m., fine. s.	Rare.
11805	U.S.N.M.	1	Near Cebu.....	Rare.

CRISTELLARIA ECHINATA (d'Orbigny).

Plate 45, figs. 4a, b; plate 46, figs. 1a, b.

"Nautili Echinati" etc., SOLDANI, Testaceographia, vol. 1, pt. 1, 1789, p. 65, pl. 59, figs. qq, rr.

Nautilus calcar, var. ϵ , FICHEL and MOLL, Test. Micr., 1803, p. 74, pl. 12, figs. a, b, c.

Robulina echinata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 100, pl. 4, figs. 21, 22.

Cristellaria (Robulina) echinata, CARPENTER, PARKER, and JONES, Introd. Foram., 1862, pl. 12, fig. 3.

Cristellaria echinata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 554, pl. 71, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 73, pl. 34, fig. 5.

This beautiful species, allied with *C. calcar*, has been found to be a common one, both in the China Sea and in the more protected waters of the Archipelago. It has not been noted in the deeper waters of the southern area in which the *Albatross* dredged—Flores Sea, Gulf of Boni, etc. It apparently is most at home in warm waters of moderate depth. Brady records it from the Philippines in 95 fathoms (174 meters), one of the two stations from which the *Challenger* obtained it. Some of the localities at which the *Albatross* obtained the species are off southern and western Luzon; east and southeastern coasts of Mindoro; off western Bohol; Sogod Bay, southern Leyte; off northwestern Panay; off Marinduque; between Cebu and Bohol; Jolo Sea; eastern Palawan; between Samar and Leyte; north of Tawi Tawi, etc. At numerous stations it is recorded as common or frequent.

The stations except one range in depth from 108 to 766 fathoms (198 meters to 1,400 meters), average depth 320 fathoms (586 meters). The bottom temperatures average 52.4° F. (11.3° C.).

Cristellaria echinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
15171	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E...	10	Co.....	Rare.
15201	U.S.N.M.	3	D5110...	13 59 20 N.; 120 75 45 E...	135	59	dk. gy. m....	Few.
15202	U.S.N.M.	4	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m....	Frequent.
15203	U.S.N.M.	1	D5198...	9 40 50 N.; 123 39 45 E...	220	53.9	gn. m.....	Rare.
11998	U.S.N.M.	7	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Frequent.
11996	U.S.N.M.	7	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Common.
11937								
15204	U.S.N.M.	9	D5261...	12 30 55 N.; 121 34 54 E...	145	s., m.....	Frequent.
15295	U.S.N.M.	10+	D5272...	14 00 00 N.; 120 22 30 E...	118	57.4	m., sh., co. s.	Abundant.
11299	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s....	Rare.
15206	U.S.N.M.	3	D5331...	15 36 45 N.; 119 47 45 E...	178	54.7	s., sh., m....	Few.
11997	U.S.N.M.	8	D5374...	13 46 45 N.; 121 35 08 E...	190	gy. m.....	Rare.
12001								
15207	U.S.N.M.	3	D5412...	10 09 15 N.; 123 52 00 E...	162	54.8	gn. m.....	Few.
15208	U.S.N.M.	1	D5423...	9 38 30 N.; 121 11 00 E...	508	49.8	gy. m., co. s.	Rare.
12003	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Rare.
12002	U.S.N.M.	2	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Few.
15209	U.S.N.M.	1	D5429...	9 37 30 N.; 118 48 30 E...	766	gn. m.....	Few.
15210	U.S.N.M.	4	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Common.
12000	U.S.N.M.	7	D5493...	9 04 00 N.; 125 20 00 E...	478	52.1	gn. m.....	Rare.
15211	U.S.N.M.	1	D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy. m., fine s.	Frequent.
15212	U.S.N.M.	8	D5523...	8 48 44 N.; 123 27 35 E...	Rare.
15213	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gy. m., glob.	Frequent.
15214	U.S.N.M.	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.....	Rare.
15168	U.S.N.M.	1	Tayabas Gulf, P. I.	Rare.

CRISTELLARIA PAPILOSA (Fichtel and Moll).

Nautilus papillosus FICHTEL and MOLL, Test. Micr., 1803, p. 82, pl. 14, figs. a-c.

Cristellaria papillosa PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 5, 1860, p. 113, No. 17.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 553, pl. 70, fig. 16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 74, pl. 37, fig. 2.

Brady had this species from but two *Challenger* stations—one in the West Indies, the other in the Indo-Pacific.

Very fine specimens have occurred in material from the following four *Alabatross* stations in this area: D5301, in 208 fathoms (381 meters), China Sea, vicinity of Hongkong, bottom temperature 50.5° F. (10.2° C.); D5424, in 340 fathoms (622 meters), Jolo Sea, bottom temperature 50.4° F. (10.2° C.); D5589, in 260 fathoms (476 meters) off Sibuko Bay, bottom temperature 45.7° F. (7.6° C.); and D5592, in 305 fathoms (558 meters), off Sibuko Bay, Borneo, bottom temperature 43.3° F. (6.1° C.).

This is a beautifully ornamented species with the rows of beading gradually decreasing in size toward the periphery and in the later added portions.

Cristellaria papillosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11709	U.S.N.M.	1	D5301...	° ' " ° ' "	208	50.5	gy. m., s....	Rare.
11708	U.S.N.M.	1	D5424...	20 37 00 N.; 115 43 00 E..	340	50.4	co. s.....	Rare.
11711	U.S.N.M.	2	D5589...	9 37 05 N.; 121 12 37 E..	260	45.7	fine gy. s., gy. m.	Rare.
11710	U.S.N.M.	4	D5592...	4 12 10 N.; 118 38 08 E..	305	43.3	gn. m.....	Few.

CRISTELLARIA ACULEATA d'Orbigny.

Cristellaria aculeata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 292, No. 14.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 555, pl. 71, figs. 4, 5.

There are two specimens which are very close to the figures of this species as given by Brady. They occur as single specimens at two stations in water of medium depth. Brady's material dredged by the *Challenger* was from the West Indian region, but these specimens seem to be identical. They are not *C. gemmata*, which is much more common in the same region and which seems to be limited to the Indo-Pacific.

Cristellaria aculeata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12030	U.S.N.M.	1	D5282...	° ' " ° ' "	248	47.4	dk. gy. s.....	Rare.
14457 15664	U.S.N.M.	1	D5259...	13 53 00 N.; 120 26 45 E..	312	49.3	gy. m., glob.	Rare.

CRISTELLARIA SUBMAMILLIGERA Cushman.

Cristellaria mamilligera H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 553, pl. 70, figs. 17, 18 (not *C. mamilligera* Karrer).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 74, pl. 34, fig. 6a (not 6b which should read 5b).

Cristellaria submamilligera CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 657.

Description.—Test biconvex, close-coiled, periphery with a thin keel; sutures curved, limbate externally, ornamented by a raised ridge which ends near the umbilical region in a protuberant knob, often a distinct raised boss over the center of the umbilical region; keel entire and not denticulate in well-preserved specimens; wall between the raised ridges, smooth; aperture radiate.

Diameter up to 3 mm., usually less.

Distribution.—Type specimen (U.S.N.M. No. 9127) from *Albatross* station D5388, in 226 fathoms (414 meters), between Burias and Luzon, bottom temperature 51.4° F. (10.8° C.), where

the species was abundant. The species is a very common one in the Philippine Archipelago, being found also in the China Sea, but not recorded in the areas to the south. It seems to be characteristic of warm, not too deep water. Some of the general localities are as follows: Off southern Luzon; east coast of Mindoro; off western Bohol; Sogod Bay, southern Leyte; east of Masbate; between Marinduque and Luzon; off northwestern Panay; Ragay Gulf, Luzon; between Samar and Masbate; between Cebu and Bohol; Jolo Sea; between Negros and Siquijor; and north of Tawi Tawi.

The depth of these stations ranges from 30 to 554 fathoms (55 to 1,012 meters), the average being 215 fathoms (393 meters), and the average bottom temperature 52.4° F. (11.3° C.).

The species differs from *C. mamilligera* Karrer in the keel, which is entire and not regularly denticulate; the ornamentation, which consists of raised ridges with bosses and not of a series of knobs.

Brady had material from the Philippines in 95 fathoms (174 meters).

Cristellaria submamilligera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15178	U.S.N.M.	4	D5097...	14 19 15 N.; 120 34 15 E...	30	° F.	gy. m., s., sh.	Frequent.
15179	U.S.N.M.	10+	D5113...	13 51 30 N.; 120 50 30 E...	159	dk. gn. m.	Frequent.
15180	U.S.N.M.	4	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m.	Few.
15181	U.S.N.M.	2	D5198...	9 40 50 N.; 123 39 45 E...	220	53.9	gn. m.	Few.
11724 11725	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.	Rare.
15182	U.S.N.M.	1	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m.	Rare.
15183	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E...	195	52.8	gn. m.	Rare.
11790 11725 15184	U.S.N.M.	{ 5 9	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Frequent.
11585	U.S.N.M.	3	D5272...	14 00 00 N.; 120 22 30 E...	118	57.4	m., sh., co. s.	Few.
15186	U.S.N.M.	5	D5278...	14 00 10 N.; 120 17 15 E...	102	50.6	fine s., m., sh	Frequent.
11793	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s.	Rare.
15187	U.S.N.M.	7	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.	Frequent.
11789	U.S.N.M.	10	D5291...	13 29 40 N.; 121 00 45 E...	173	51.5	fine bk. s.	Frequent.
11791	U.S.N.M.	4	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.	Frequent.
15188	U.S.N.M.	3	D5369...	13 48 00 N.; 121 43 00 E...	106	bk. s.	Few.
11792	U.S.N.M.	2	D5371...	13 49 40 N.; 121 40 15 E...	83	gn. m.	Few.
15189	U.S.N.M.	6	D5374...	13 46 45 N.; 121 35 08 E...	190	gy. m.	Frequent.
15190	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 25 E...	88	co. s.	Rare.
15191	U.S.N.M.	9	D5388...	12 51 30 N.; 123 26 15 E...	226	51.4	sft. gn. m.	Abundant.
15192	U.S.N.M.	1	D5394...	12 00 30 N.; 124 05 30 E...	153	gn. m.	Rare.
15193	U.S.N.M.	1	D5412...	10 09 15 N.; 123 52 00 E...	162	54.8	gn. m.	Rare.
15194	U.S.N.M.	1	D5425...	9 37 45 N.; 121 11 00 E...	495	49.4	gy. m., co. s.	Few.
			D5512...	8 16 02 N.; 123 58 25 E...	445	52.8	gy. m., fine s.	Rare.
15195	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E...	Rare.
15196	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Rare.
15197	U.S.N.M.	2	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s.	Rare.
15198	U.S.N.M.	1	D5569...	5 13 15 N.; 120 15 30 E...	303	52.3	co. s.	Rare.
15199	U.S.N.M.	2	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Rare.

CRISTELLARIA EXPANSA Cushman.

Plate 46, figs. 2a, b.

Cristellaria expansa CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 658.

Description.—Test biconvex, with a very wide, prominent, very thin keel; chambers with a large alar projection on the umbilical

end extending beyond the umbilicus and onto the previous whorl; sutures with raised ribs ending in a knob near the umbilicus, remainder of surface smooth; aperture radiate, apertural face concave.

Diameter up to 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9128) from *Albatross* station D5467, in 480 fathoms (878 meters), east coast of Luzon. The species also occurred at numerous stations, mostly in the China Sea and in the Philippine Archipelago. The depth of these stations ranges from 88 to 554 fathoms (161 to 1,012 meters), the average being 260 fathoms (476 meters). The average bottom temperature is 52.6° F. (11.4° C.).

This species differs from *C. submamilligera* mainly in its great development of the alar projection on the inner border of the chamber.

Cristellaria expansa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12018	U.S.N.M.	4	D5113...	13 51 30 N.; 120 50 30 E...	159	dk. gn. m...	Few.
12022	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m...	Rare.
11692	U.S.N.M.	3	D5123...	13 12 45 N.; 121 38 45 E...	283	gn. m.	Few.
12024	U.S.N.M.	3	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m...	Rare.
12026	U.S.N.M.	9	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Common.
12021	U.S.N.M.	10+	D5272...	14 00 00 N.; 120 22 30 E...	118	57.4	m., sh., co. s.	Frequent.
12025	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s...	Rare.
12019	U.S.N.M.	10+	D5369...	13 48 00 N.; 121 43 00 E...	106	bk. s.	Frequent.
15221	U.S.N.M.	2	D5381...	13 14 15 N.; 122 44 25 E...	88	co. s.	Few.
12027	U.S.N.M.	4	D5382...	13 15 20 N.; 122 45 30 E...	128	gn. m.	Few.
12020	U.S.N.M.	1	D5410...	10 28 45 N.; 124 05 30 E...	385	m. m.	Rare.
9128	U.S.N.M.	1	D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m.	Rare.
				8 48 44 N.; 123 27 35 E...
12023	U.S.N.M.	2	D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gy. m., glob.	Rare.
11293	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Rare.
15222	U.S.N.M.	1	D5569...	5 33 15 N.; 120 12 57 E...	303	52.3	co. s.	Rare.
15223	U.S.N.M.	1	D5574...	5 30 45 N.; 120 07 57 E...	Rare.

CRISTELLARIA EXPANSA, var. PLANULATA Cushman.

Plate 46, figs. 3a, b.

Cristellaria expansa, var. *planulata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 658.

Description.—Variety differing from the typical mainly in the lack of ornamentation characteristic of the type, and if at all ornamented consisting of indistinct costae in the axis of growth; alar projections more distinct than in the type, due to the lack of ornamentation to hide the sutures, usually more compressed than the type form.

Distribution.—Type specimen (U.S.N.M. No. 9129) from *Albatross* station D5220, in 50 fathoms (91 meters), between Marinduque and Luzon.

Cristellaria expansa, var. *planulata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11617 9129 15301	U.S.N.M. U.S.N.M.	1 3	D5220... D5570...	° / " ° / " 13 38 00 N.; 121 58 00 E... 5 32 15 N.; 120 12 57 E...	50 330	° F. 52.3	sft. gn. m... fne. s., glob..	Rare. Few.

CRISTELLARIA DENTICULIFERA Cushman.

Cristellaria cultrata MONTFORT, dentate variety H. B. BRADY, Rep. Voy, *Challenger*, Zoology, vol. 9, 1884, p. 550, pl. 70, figs. 7, 8.

Cristellaria denticulifera CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 75, pl. 37, fig. 1.

Description.—Test biconvex, sutures curved; periphery marked by a keel which is denticulate, especially between the terminations of the sutures; the sutures somewhat limbate externally and with raised lines, terminating in a raised mass of clear shell material at the umbo; surface ornamented by a series of broken costae, running longitudinally and somewhat obliquely; especially well-developed on the earlier chambers, becoming obsolete in later growth.

Diameter up to 2 mm.

Distribution.—Typical specimens were obtained from *Albatross* stations: D5110, in 135 fathoms (247 meters), China Sea off southern Luzon, bottom temperature 59° F. (15° C.); D5132, in 26 fathoms (48 meters), Sulu Sea off western Mindanao; D5214, in 218 fathoms (399 meters), east of Masbate Island, bottom temperature 51.4° F. (10.8° C.); D 5313, in 150 fathoms (270 meters), China Sea off Hong-kong, bottom temperature 53.6° F. (12° C.); D5255, in 100 fathoms (183 meters), Gulf of Davao, D5565, in 243 fathoms (445 meters), between Jolo and Tawi Tawi, bottom temperature 52.3° F. (11.2° C.); and D5637, in 700 fathoms (1,280 meters), off Bouro Island.

Brady records *C. cultrata* in 95 fathoms off the Philippines, and it would be interesting to know whether this was the denticulate form he figures.

The species is very constant in its characters, and the figures given by Brady in the *Challenger* Report are good figures of it.

Cristellaria denticulifera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12029 12028	U.S.N.M. U.S.N.M.	1 1	D5110... D5132...	° / " ° / " 13 59 20 N.; 120 75 45 E... Island off Panabutan Point N.; 15° W., 30 mile.	135 26	° F. 59.0	dk. gy. m.... gn. m., s.....	Rare. Rare.
13326 13324 13325 15254 12017	U.S.N.M. U.S.N.M. U.S.N.M. U.S.N.M. U.S.N.M.	2 1 1 1 1	D5214... D5313... D5255... D5565... D5637...	12 25 18 N.; 123 37 15 E... 21 30 00 N.; 116 43 00 E... 7 03 00 N.; 125 39 00 E... 5 51 42 N.; 120 30 30 E... 3 53 20 S.; 126 48 00 E...	218 150 100 243 700	51.4 53.6 52.3	gn. m..... s..... sft. m..... s. ptr. sh... gy. m.....	Rare. Rare. Rare. Rare. Rare.

CRISTELLARIA GEMMATA H. B. Brady.

Cristellaria gemmata H. B. BRADY, Quart. Journ. Micr. Soc., vol. 21, 1881, p. 64; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 554, pl. 71, figs. 6, 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 75, pl. 34, fig. 7.

Brady describes this species from three Indo-Pacific stations of the *Challenger*, one of which was in the Philippines at a depth of 95 fathoms (174 meters).

Apparently this is a rather limited species, as it does not seem to occur or to be noted outside this region. I have had excellent material from several stations in the Philippines; China Sea off southern Luzon; between Marinduque and Luzon; off southeastern Mindoro; off Marinduque, it being especially well distributed. The 10 stations range in depth from 106 to 340 fathoms (194 to 622 meters), with the average 198 fathoms (363 meters). The average bottom temperature where given was 52.5° F. (11.3° C.).

Cristellaria gemmata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11685	U.S.N.M.	9	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m.	Frequent.
11686	U.S.N.M.	4	D5113...	13 51 30 N.; 120 50 30 E..	159	dk. gn. m.	Few.
15169	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.	Rare.
11691	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Rare.
11688	U.S.N.M.	7	D5261...	12 30 55 N.; 121 34 24 E..	145	s. m.	Frequent.
			D5282...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. gy. s.	
11690	U.S.N.M.	10+	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.	Frequent.
11689	U.S.N.M.	1	D5369...	13 48 00 N.; 121 43 00 E..	106	bk. s.	Few.
11687	U.S.N.M.	3	D5370...	13 44 15 N.; 121 42 30 E..	159	54.3	sft. m.	Few.
12016	U.S.N.M.	3	D5374...	13 46 45 N.; 121 35 08 E..	190	gy. m.	Few.

CRISTELLARIA COSTATA (Fichtel and Moll).

Plate 46, fig. 4; plate 47, fig. 1.

Nautilus costatus FICHEL and MOLL, Test. Micr., 1803, p. 47, pl. 4, figs. *g-i*.

Spincterulus costatus MONTFORT, Conch. Syst., 1808, p. 222, 56^e genr.

Lenticulina costata DEFRANCE, Dict. Sci. Nat., vol. 32, 1824, p. 182.

Robulina costata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 289, No. 13 [?].

Cristellaria costata PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 5, 1860, p. 113, No. 19.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 555, pl. 71, fig. 8 (not fig. 9).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 75, pl. 34, fig. 4.

Distribution.—This species with its attendant variations is one of the most striking constituents of the foraminiferal fauna of this area. With large suites of specimens it has been found possible to split the various forms into a typical form, with three varieties according to the trend which the variation develops. These varieties, while they seem almost like separate species, can be connected with one another and are really different modifications of the same type of ornamented test.

The typical form, consisting of a compressed test, close-coiled, with a few chambers, a rather acute peripheral border, and an ornamentation consisting of raised costal lines on the limbate sutures, with an independent set of costae between and running more nearly in the axis of growth. In the typical form these two sets of costae are more or less evenly balanced, as in the figure given by Brady (pl. 71, fig. 9) and as shown here.

The typical form of the species as here selected occurred at a number of stations in the area within the Archipelago proper; off Romblon; Sogod Bay, southern Leyte; at several stations off Marinduque Island; off northwestern Panay; and China Sea, off southern Luzon.

The stations range in depth from 50 to 554 fathoms (91 to 1,012 meters), with the average 215 fathoms (393 meters). The only two stations at which bottom temperatures were taken are 49.3° F. and 52.8° F. (9.6° C. and 11.5° C.).

Cristellaria costata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11833	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78	° F.	fine. s.....	Rare.
11834	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
11835 15302	U.S.N.M.	3	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m...	Frequent.
11838	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
15666	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s.....	Few.
11837 11836	U.S.N.M.	6	D5369...	13 48 00 N.; 121 43 00 E...	106	dk. s.....	Frequent.
11839	U.S.N.M.	2	D5374...	13 46 45 N.; 121 35 08 E...	190	gy. m.....	Few.

CRISTELLARIA COSTATA (Fichtel and Moll), var. MULTICOSTA Cushman.

Plate 47, figs. 2, 3.

Cristellaria costata (FICHTEL and MOLL), var. *multicosta* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 658.

Description.—Variety very much compressed, complanate, with the costae along the limbate sutures obsolete, the sutures slightly depressed; the ornamentation of each chamber consisting of a large number of fine raised costae spreading fan-like from the aperture as a center backward and covering the entire surface of the chamber; aperture radiate, protruded.

Diameter up to 4 mm.

Distribution.—Type specimen (U.S.N.M. No. 9130) from *Albatross* station D5538, in 256 fathoms (468 meters), between Negros and Siquijor, bottom temperature 53.3° F. (11.8° C.). Besides at the type station well-marked specimens of this variety have occurred at *Albatross* station D5178, in 78 fathoms (143 meters), off Romblon; D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom

temperature 49.3° F. (9.6° C.); D5278, in 102 fathoms (187 meters), China Sea off southern Luzon, bottom temperature 59.6° F. (15.3° C.); D5537, in 254 fathoms (465 meters), between Negros and Siquijor, bottom temperature 53.5° F. (11.8° C.); and D5579, in 175 fathoms (320 meters), off Darvel Bay, Borneo, bottom temperature 55.3° F. (12.9° C.).

This variety is one of the most beautifully sculptured of all the genus. It is an extreme modification of the costae of the chambers which are developed at the expense of those of the sutures.

Cristellaria costata, var. *multicosta*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15303	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78	fine s.	Rare.
11855	U.S.N.M.	4	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
11857	U.S.N.M.	1	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fine s., m., sh	Rare.
11858	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Rare.
13231	} U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s.	Rare.
9130								
11856	U.S.N.M.	2	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fine s., Co...	Rare.

CRISTELLARIA COSTATA (Fichtel and Moll), var. SUBDECORATA Cushman.

Plate 47, figs. 4, 5.

Cristellaria costata (FICHEL and MOLL), var. *subdecorata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 659.

Description.—Variety with the costae of the sutures the predominant factor in the ornamentation, the areas between with fine costae or striae, anastomosing; the costae of the sutures large and well-marked, sharply raised above the surface of the test, the umbilical region covered with a thick cluster of irregular bosses; periphery with traces of a keel between the sutures.

Diameter up to 4 mm.

Distribution.—Type specimen (U.S.N.M. No. 9131) from *Albatross* station D5454, in 153 fathoms (279 meters), east coast of Luzon. The variety was common at this station and occurred more or less frequently at the following *Albatross* stations: D5115, in 340 fathoms (622 meters), off Balayan Bay; D5121, in 108 fathoms (198 meters), east coast of Mindoro; D5191, Tanon Strait, in 258 fathoms (622 meters), bottom temperature 62.8° F. (17.1° C.); D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); D5261, off southeastern Mindoro, in 145 fathoms (265 meters); D5388, in 226 fathoms (414 meters), between Burias and Luzon, bottom temperature 51.4° F. (10.8° C.); D5291, in 173 fathoms (316 meters), China Sea off southern Luzon, bottom tem-

perature 51.5° F. (10.8° C.); D5412, in 162 fathoms (297 meters), between Cebu and Bohol, bottom temperature 54.8° F. (12.6° C.); and D5523, northern Mindanao.

This is the most beautifully ornamented variety of the species, and is full of detailed sculpture as shown in the figure. It has occurred in very large numbers at two of the stations.

Cristellaria costata, var. *subdecorata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
11842	U.S.N.M.	1	D5115...	13 37 11 N.; 120 43 40 E...	340	Rare.
11850	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m.	Rare.
11845	U.S.N.M.	2	D5191...	10 29 45 N.; 123 31 15 E...	258	62.8	gn. m.	Rare.
11847	U.S.N.M.	4	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
11841	U.S.N.M.	3	D5261...	12 30 55 N.; 121 34 24 E...	145	s. m.	Few.
11846	U.S.N.M.	1	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.	Rare.
11843	U.S.N.M.	1	D5388...	12 51 30 N.; 123 26 15 E...	226	51.4	sft. gn. m.	Rare.
15304	U.S.N.M.	10+	D5291...	13 29 40 N.; 121 00 45 E...	173	51.5	fne. bk. s.	Common.
11849	U.S.N.M.	1	D5412...	10 09 15 N.; 123 52 00 E...	162	54.8	gn. m.	Rare.
11844 9131	} U.S.N.M.	10+	D5454...	13 12 00 N.; 123 50 30 E...	153	Common.
11848								

CRISTELLARIA COSTATA (Fichtel and Moll), var. *SUBLAEVIS* Cushman.

Plate 49, fig. 1.

Cristellaria costata (FICHEL and MOLL), var. *sublaevis* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 659.

Description.—Variety differing from the typical in having the ornamentation of the test so developed that the limbate sutures and their resulting costae are developed at the complete expense of the other costae of the chambers or nearly so, the costae of the sutures being very high and ending on the inner border of each in a large raised knob; peripheral margin of the test keeled, somewhat interrupted at the sutures; wall of the chambers smooth or marked with obsolescent striae in some cases.

Diameter 3 to 4 mm.

Distribution.—Type specimen (U.S.N.M. No. 9132) from *Albatross* station D5374, in 190 fathoms (348 meters), off Marinduque Island. Specimens were also obtained from D5192, off northern Cebu Island, 32 fathoms (59 meters); D5212, in 108 fathoms (198 meters), east of Masbate Island, bottom temperature 59.9° F. (15.5° C.); D5523, off northern Mindanao; and D5586, Sibuko Bay, Borneo, 347 fathoms (635 meters), bottom temperature 44.0° F. (6.6° C.).

In this variety the costae of the chambers so highly developed in var. *multicosta* are almost entirely lacking and the development of the other set of ornamental costae is carried to a very high development. Without the typical form and variety *subdecorata* these two extremes would hardly be recognized as belonging to a single species.

Cristellaria costata, var. *sublaevis*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11851	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.	Rare.
11852	U.S.N.M.	1	D5212...	12 04 15 N.; 124 04 36 E..	108	59.9	gy. s., m.	Rare.
9132	} U.S.N.M.	1	D5374...	13 46 45 N.; 121 35 08 E..	190	gy. m.	Rare.
11853						
11854						
15305	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.	Rare.

CRISTELLARIA BRADYI Cushman.

Plate 48, figs. 1a, b.

Cristellaria costata H. B. BRADY (in part?), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 555, pl. 71, fig. 8.

Cristellaria bradyi CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 659.

Description.—Test ovate, thick, last few chambers more or less elongated; sutures hidden by the surface ornamentation, which consists of longitudinal rounded costae, more or less broken, especially toward the umbilical area, and becoming less distinct or entirely disappearing on the last-formed chambers; peripheral margin rounded; aperture radiate.

Length up to 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9133) from *Albatross* station D5313, in 150 fathoms (270 meters), China Sea off Hongkong, bottom temperature 53.6° F. (12.0° C.). Besides, it has occurred at various other stations in the China Sea, off Hongkong and off Formosa; off Jolo; Linapacan Strait; Palawan Passage; and between Jolo and Tawi Tawi.

These stations range in depth from 43 to 340 fathoms (78 to 622 meters), the average depth being 167 fathoms (305 meters). The bottom temperatures where given average 53.5° F. (11.8° C.).

This species differs from typical *C. costata* in that the longitudinal costae are the dominant ornamentation, running parallel with the axis of growth and are continuous over the whole test, while those of *C. costata* are limited to definite series, each set limited to a single chamber and independent of those of adjacent chambers.

Cristellaria bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11818	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E..	318	fne. s., sh.	Rare.
13227	} U.S.N.M.	5	D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s.	Frequent.
9133						
11815	U.S.N.M.	8	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh.	Frequent.
11814	U.S.N.M.	1	D5318...	21 32 00 N.; 117 46 00 E..	340	s. br., Co.	Rare.
11877	U.S.N.M.	1	D5336...	11 37 45 N.; 119 46 00 E..	46	s. m.	Rare.
11813	U.S.N.M.	10+	D5338...	11 33 45 N.; 119 24 45 E..	43	Co. s., m.	Common.
11816	U.S.N.M.	1	D5339...	11 22 00 N.; 119 12 00 E..	52	m.	Rare.
11817	U.S.N.M.	1	D5566...	5 52 12 N.; 120 31 00 E..	244	52.5	fne. s., sh.	Rare.

CRISTELLARIA CASSIS (Fichtel and Moll).

Plate 49, figs. 2a, b.

"Cornu Hammonis" PLANCUS, Conch. Min., 1760, p. 120, pl. 1, fig. 11.

"Litui crispatis et orbiculi" SOLDANI, Testaceographia, vol. 1, pt. 1, 1789, p. 63, pl. 56, figs. I, K, etc.

Nautilus cassis FICHEL and MOLL, Test Micr., 1803, p. 95, pl. 17.*Cristellaria cassis* LAMARCK, Tabl. Encycl. et Méthod., 1816, pl. 467, figs. 3a-d.—
H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 552, pl. 68, fig. 10.

Description.—Test much compressed, ovate, early chambers regularly coiled, later ones becoming elongate and showing a tendency toward uncoiling; umbilical region with prominent bosses and single bosses appear on the sutures near the peripheral edge; umbilical region broadest in front view, wall otherwise smooth; sutures little if at all depressed, of clear shell material; peripheral border with a definite, very thin keel; aperture radiate, at the peripheral margin of the chamber.

Length up to 4 mm. or more.

Distribution.—This species occurred at a number of *Albatross* stations in the region from the following localities: Jolo Sea; off eastern Palawan; several stations on the east coast of Luzon; Gulf of Tomini, Celebes; Patiente Strait; off Bouro Island and Gulf of Boni.

The depth of these stations ranges from 480 to 901 fathoms (887 to 1,648 meters), the average depth being 614 fathoms (1,123 meters). The bottom temperatures where given average 44.7° F. (7.0° C.).

At some of the stations off the east coast of Luzon this species was abundant, but at the others it was represented by but few specimens. It has not previously been recorded from the North Pacific.

Cristellaria cassis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11891	U.S.N.M.	1	D5425...	9 37 45 N.; 121 11 00 E..	495	49.4	gy. m. co., s.	Rare.
15162	U.S.N.M.	2	D5429...	9 37 30 N.; 118 48 30 E..	766	gn. m.	Few.
11893	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E..	500	gy. m.	Few.
11890	U.S.N.M.	10+	D5467...	13 35 37 N.; 123 37 18 E..	450	gy. m.	Abundant.
11894	U.S.N.M.	2	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.	Few.
11896	U.S.N.M.	10+	D5470...	13 37 30 N.; 123 41 09 E..	560	m.	Abundant.
15163	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E..	750	Rare.
11892	U.S.N.M.	4	D5630...	0 56 30 S.; 128 04 00 E..	569	co. s., m	Frequent.
11895								
15164	U.S.N.M.	1	D5637...	3 53 29 S.; 126 48 09 E..	700	gy. m.	Rare.
11897	U.S.N.M.	4	D5650...	4 53 45 S.; 121 29 09 E..	540	40.1	gn. m.	Few.
15165	U.S.N.M.	2	D5668...	2 23 15 S.; 118 49 00 E..	901	38.2	gy. m.	Rare.

CRISTELLARIA CASSINOIDES Cushman.

Plate 48, figs. 2 a, b.

Cristellaria cassinoides CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 659.

Description.—Test elongate, much compressed, arcuate, early chambers coiled, later ones uncoiling and forming an elongate growth

several times the length of the coiled portion, inner margin slightly keeled, peripheral margin bluntly rounded; sutures with a raised portion along the central portion of the peripheral half of the test, surface otherwise smooth and unornamented; aperture radiate.

Length up to 5 mm.

Distribution.—Type specimen (U.S.N.M. No. 9134) from *Albatross* station D5381, in 88 fathoms (161 meters), Raygay Gulf, Luzon. The species is fairly common in the area. Beside the type station it occurred at various stations: China Sea, off southern Luzon; east coast of Mindoro; off western Bohol; Sogod Bay, southern Leyte; off northwestern Panay; off Marinduque Island; Jolo Sea; north of Tawi Tawi; and Sibuko Bay, Borneo. At some of the stations specimens were abundant.

The stations range in depth from 88 to 554 fathoms (161 to 1,012 meters), the average being 236 fathoms (432 meters). The average bottom temperature is 51.9° F. (11.0° C.).

This is a species rather striking in its crozier form and peculiar but distinctive ornamentation.

Cristellaria cassinoides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11878	U.S.N.M.	10+	D5110...	13 59 20 N.; 120 75 45 E.	135	59.0	dk. gy. m.	Abundant.
11881	U.S.N.M.	1	D5221...	13 27 20 N.; 121 17 45 E.	108	59.0	dk. gn. m.	Rare.
11880	U.S.N.M.	1	D5198...	9 40 50 N.; 123 39 45 E.	220	53.9	gn. m.	Rare.
11882	U.S.N.M.	5	D5201...	10 10 00 N.; 125 04 15 E.	554	52.8	gy. s., m.	Frequent.
11885	U.S.N.M.	1	D5239...	11 57 30 N.; 121 42 15 E.	312	49.3	gy. m., glob.	Rare.
11884	U.S.N.M.	1	D5261...	12 30 55 N.; 121 34 24 E.	145		s., m.	Rare.
11883	U.S.N.M.	3	D5272...	14 00 00 N.; 120 22 30 E.	118	57.4	m., sh., co. s.	Few.
11886	U.S.N.M.	4	D5278...	14 10 09 N.; 120 17 15 E.	102	59.6	fine. s., m., sh.	Frequent.
15166	U.S.N.M.	1	D5282...	13 53 00 N.; 120 26 45 E.	248	47.4	dk. gy. s.	Rare.
11887	U.S.N.M.	2	D5374...	13 46 45 N.; 121 35 08 E.	190		gy. m.	Few.
			D5381...	13 14 15 N.; 122 44 25 E.	88		co. s.	Rare.
11889	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E.	340	50.4	co. s.	Rare.
15167	U.S.N.M.	1	D5569...	5 13 15 N.; 120 15 30 E.	303	52.3	co. s.	Rare.
11888	U.S.N.M.	2	D5576...	5 25 56 N.; 120 03 39 E.	277	53.3	s.	Few.
11879	U.S.N.M.	6	D5592...	4 12 44 N.; 118 27 44 E.	305	43.3	gn. m.	Frequent.

CRISTELLARIA TUMIDO-COSTATA Cushman.

Plate 49, figs. 4 a, b.

Cristellaria tumido-costata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 660.

Description.—Test very broadly biconvex, close-coiled; chambers comparatively few in the whorl; sutures marked by a series of very strong raised ribs uniting at the umbilicus; remaining surface between the sutures marked by a series of short, well-rounded, parallel costae, running obliquely to the axis of growth; peripheral portion of the test also marked by a series of costae parallel to the periphery, or nearly so; apertural face flattened or concave with a broad rim about the margin; aperture radiate at the peripheral angle of the chamber.

Diameter up to 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9136) from *Albatross* station D5570, in 330 fathoms (604 meters), north of Tawi Tawi, bottom temperature 52.3° F. (11.2° C.). It also occurred at D5548, in 232 fathoms (424 meters), off Jolo Island, bottom temperature 53.5° F. (11.8° C.), and D5579, in 175 fathoms (320 meters), off Darvel Bay, Borneo, bottom temperature 55.3° F. (12.9° C.).

This is one of the most beautifully sculptured species found in the Philippine region.

Cristellaria tumido-costata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	bottom.	Abundance.
11754	U.S.N.M.	1	D5548...	6 00 20 N.; 120 45 35 E...	232	53.5	s., brk. sh...	Rare.
9136	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fine s., glob...	Rare.
11755	U.S.N.M.	1	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fine s., Co...	Rare.

CRISTELLARIA TUMIDO-COSTATA, var. LABYRINTHICA Cushman.

Plate 50, figs. 1 a, b.

Cristellaria tumido-costata, var. *labyrinthica* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 660.

Description.—Variety differing from the typical in the shape, which is more tumid, the periphery more broadly rounded than in the typical, and the ornamentation, which consists of a peripheral series of costae, as in the typical; but the costae above the sutures obsolescent and the intermediary costae becoming the prominent sculpture, and becoming shorter, irregular, making a labyrinth-like network of raised ridges.

Distribution.—The variety was found only at *Albatross* station D5567 in 268 fathoms (491 meters), north of Tawi Tawi, bottom temperature 52.° F. (11.1° C.). Type (U.S.N.M. No. 9137).

Several specimens from this station show the relation between this variety and the typical form.

Cristellaria tumido-costata, var. *labyrinthica*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9137 15306	U.S.N.M. U.S.N.M.	1 2	D5567...	5 48 00 N.; 120 33 45 E...	268	52.0	fine s.....	Rare.

CRISTELLARIA DORSO-COSTATA Cushman.

Plate 48, figs. 3 a-c.

Cristellaria dorso-costata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 659.

Description.—Test composed of numerous chambers, the early ones close-coiled, the later ones becoming elongate and tending toward uncoiling; test thick, periphery broadly rounded; sutures little depressed; wall smooth except toward peripheral border, where there are several rounded costae running lengthwise of the test, becoming obsolete toward the later growth; aperture radiate.

Length up to 3.5 mm.

Distribution.—Type specimen (U.S.N.M. No. 9135) from *Albatross* station D5268, in 170 fathoms (311 meters), off Batangas Bay. Specimens also occurred in material from the following stations: D5134, in 25 fathoms (46 meters), Sulu Archipelago near Basilan Island; D5141, in 29 fathoms (53 meters); D5143, in 19 fathoms (35 meters); D5144, in 19 fathoms (35 meters), all off Jolo; D5147, 21 fathoms (39 meters), off Siasi; D5152, 34 fathoms (62 meters), off Tawi Tawi; D5179, off Romblon, 37 fathoms (68 meters); and D5218, 20 fathoms (37 meters) between Burias and Luzon.

This seems to be a rather constant species and its characters of ornamentation with its partially uncoiled form should make it easily distinguished. Apparently it is found in the shallower, warmer waters among the islands of the Archipelago from the data obtainable.

Cristellaria dorso-costata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12009	U.S.N.M.	2	D5134...	6 44 45 N.; 121 48 00 E...	25	fne. s.....	Rare.
12008	U.S.N.M.	4	D5141...	6 09 00 N.; 120 58 00 E...	29	co. s.....	Few.
12004	U.S.N.M.	10+	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Abundant.
12007	U.S.N.M.	3	D5144...	6 05 50 N.; 121 02 15 E...	19	co. s., sh.....	Few.
12010	U.S.N.M.	1	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.....	Rare.
12006	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Common.
12001	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	fne. s.....	Rare.
12005	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Rare.
9135	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Rare.

CRISTELLARIA COMPRESSA d'Orbigny.

Cristellaria compressa D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 86, pl. 3, figs. 32, 33.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 538, pl. 104, figs. 15, 16.

Description.—Test elongate, much compressed, early portion close-coiled, later chambers becoming uncoiled and arcuate; sutures very slightly if at all depressed, of clear shell material; periphery of the test with a thin keel; aperture on the peripheral margin of the chamber, radiate.

Length up to 6 mm.

Distribution.—The only station from which this species was obtained is *Albatross* station D5670 in 1,181 fathoms (2,160 meters), in Macassar Strait, with a bottom temperature of 38.2° F. (3.4° C.). It apparently has not previously been recorded from the Pacific.

The specimen and those figured by Brady seem almost like an extreme form of *C. cassis* (Fichtel and Moll), as they have very similar characters except that this is much more uncoiled than is *C. cassis* and lacks the bosses of the ornamentation.

Cristellaria compressa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11870 11867	} U.S.N.M.	2	D5670...	° ' " ° ' "	1,181	° F.	gy. m.	Rare.
				1 19 00 S.; 118 43 00 E....		38.2		

CRISTELLARIA PAUCICOSTATA Cushman.

Plate 49, figs. 3 a, b.

Cristellaria paucicostata CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 660.

Description.—Test much compressed, composed of few chambers; peripheral border with a prominent keel, the earlier portion of the test with two sharp raised costae toward the periphery and parallel to it, becoming obsolete on the later developed chambers; remainder of surface smooth; sutures slightly depressed; aperture radiate, somewhat protruded.

Length up to 2 mm.

Distribution.—Type specimen (U.S.N.M. No. 9138) from *Albatross* station D5586 in 347 fathoms (635 meters), off Sibuko Bay, Borneo, bottom temperature 44° F. (6.6° C.).

Cristellaria paucicostata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9138	U.S.N.M.	1	D5586...	° ' " ° ' "	347	° F.	gy. m.	Rare.
				4 06 50 N.; 118 47 20 E....		44.0		

CRISTELLARIA CREPIDULA (Fichtel and Moll).

Plate 50, figs. 4, 5.

Nautilus crepidula FICHEL and MOLL, Test. Micr., 1803, p. 107, pl. 19, figs. g-i.
Cristellaria crepidula D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 64, pl. 8, figs. 17, 18.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 542, pl. 67, figs. 17, 19, 20; pl. 68, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 70, pl. 29, figs. 5, 6; pl. 31, figs. 2-5.

The following localities have given records for this species: Off Pujada Bay; China Sea, off southern Luzon and off Hongkong; Palawan Passage; Sogod Bay, southern Leyte; Darvel Bay and Sibuko Bay, Borneo; and Macassar Strait. These stations range in depth from 88 to 730 fathoms (161 to 1,335 meters), the average depth being 237 fathoms (434 meters) and the average bottom temperature being 50.3° F. (10.1° C.).

Cristellaria crepidula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15307	U.S.N.M.	1	D5201...	° ' " ° ' "		° F.		
11829	U.S.N.M.	1	D5242...	10 10 00 N.; 125 04 15 E....	554	52.8	fine. gy. s., m.	Rare.
11830	U.S.N.M.	1	D5242...	6 51 53 N.; 126 14 10 E....	191	64.1	sit. gy. m	Rare.
11831	U.S.N.M.	4	D5300...	20 31 00 N.; 115 49 00 E....	265	gy. m., s....	Few.
11832	U.S.N.M.	1	D5311...	21 33 00 N.; 116 15 00 E....	88	crs. s., sh....	Rare.
11828	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E....	730	40.6	co. s	Rare.
11827	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E....	272	47.5	gn. m.....	Rare.

CRISTELLARIA SCHLOENBACHI Reuss.

Cristellaria schloenbachi REUSS, Sitz. Akad. Wiss. Wien, vol. 46, 1862, p. 65, pl. 6, figs. 14, 15.—H. P. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 539, pl. 67, fig. 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 77, pl. 36, fig. 6.

Specimens of this species have occurred at the following *Albatross* stations in the region: D5110, in 135 fathoms (247 meters), China Sea off southern Luzon, bottom temperature 59° F. (15° C.); D5112, in 177 fathoms (324 meters), same locality, bottom temperature 52.4° F. (11.3° C.); D5153, in 49 fathoms (90 meters), off Tawi Tawi; D5179, off Romblon, 37 fathoms (68 meters), bottom temperature 75.7° F. (24.2° C.); D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte, bottom temperature 52.8° F. (11.5° C.); D5277, in 80 fathoms (146 meters), China Sea off southern Luzon, bottom temperature 58.6° F. (14.7° C.); D5551, in 193 fathoms (353 meters) off Jolo, bottom temperature 53.3° F. (11.8° C.); and D5595 in 9 fathoms (16 meters), off Zamboanga, Mindanao.

Cristellaria schloenbachi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15298	U.S.N.M.	1	D5110...	° ' " ° ' "		° F.		
11722	U.S.N.M.	1	D5112...	13 59 20 N.; 120 75 45 E....	135	59.0	dk. gy. m...	Rare.
11720	U.S.N.M.	1	D5112...	13 48 22 N.; 120 47 25 E....	177	52.4	dk. gn. m...	Rare.
15299	U.S.N.M.	1	D5153...	5 18 10 N.; 120 02 55 E....	49	co. s., sh....	Rare.
11719	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E....	37	75.7	hrd. s....	Rare.
11723	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E....	554	52.8	gy. s., m....	Rare.
15300	U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E....	80	58.6	sh., p., s....	Rare.
11721	U.S.N.M.	1	D5551...	5 54 48 N.; 120 44 24 E....	193	53.3	fine. s.....	Rare.
			D5595...	6 54 00 N.; 122 04 30 E....	9

CRISTELLARIA TENUIS (Bornemann).

Plate 50, fig. 2.

Marginulina tenuis BORNEMANN, Zeitschr. deutsch. geol. Ges., vol. 7, 1855, p. 326, pl. 13, fig. 4.

Cristellaria tenuis REUSS, Sitz. Akad. Wiss. Wien, vol. 62, 1870, p. 479, No. 1.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 535, pl. 66, figs. 21–23.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 315, pl. 61, fig. 2.

Cristellaria perprocera SCHWAGER, *Novara-Exped.*, Geol. Theil., vol. 2, 1866, p. 241, pl. 6, fig. 84.

Description.—Test very long and slender, composed of a large number of chambers; early portion of the test compressed and coiled, the later portion becoming early uncoiled and gradually losing the compression until the final chambers are cylindrical and nodosarian; sutures of the early portion not incised, those of the later portion rather deeply constricted; wall smooth, aperture in the last-formed chamber, central, terminal, radiate.

Length up to 3.5 mm.

Distribution.—Specimens occurred at the following *Albatross* stations in the area: D5269, in 220 fathoms (403 meters), off Batangas Bay; D5301, in 208 fathoms (381 meters), China Sea, off Hongkong, bottom temperature 50.5° F. (10.2° C.); D5312, in 140 fathoms (250 meters), bottom temperature 57.5° F. (14.2° C.); also off Hongkong, D5622, in 275 fathoms (503 meters), between Gillolo and Makyan Islands; and D5648, in 559 fathoms (1,021 meters), Buton Strait.

The specimens I have had from this area are much like the specimens figured by Brady, perhaps more like them than like the other figures referred to above.

Cristellaria tenuis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11781	U.S.N.M.	1	D5269...	13 39 50 N.; 120 59 30 E...	220	° F.	fine s., p.....	Rare.
15309	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.....	Rare.
11783	U.S.N.M.	1	D5312...	21 30 00 N.; 116 32 00 E...	140	57.5	s., sml. sh.....	Rare.
11784	U.S.N.M.	1	D5622...	Makyan Island (NE.), N. 66° W.	275	gy. m.....	Rare.
11782	U.S.N.M.	1	D5648...	5 35 00 S.; 122 20 00 E...	559	39.2	gn. m.....	Rare.

CRISTELLARIA HELICINA Cushman.

Plate 51, figs. 4 a, b.

Cristellaria helicina CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 660.

Description.—Test unequally convex, close-coiled, chambers numerous, on one side coming to the umbilicus or nearly so, on the other coming only little farther than the periphery of the previous

whorl; in face view subcarinate; apertural face unequal; sutures ornamented by raised areas, obsolescent at the periphery, but gradually increasing in size toward the umbilical end, becoming in some cases a distinct boss toward the umbilicus, those of the earlier whorls fusing into a partial ring about the umbilicus; main surface of the chambers smooth and unornamented; aperture radiate and projecting on the peripheral margin of the chamber.

Diameter up to 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9139) from *Albatross* station D5272, in 118 fathoms (203 meters), China Sea, off southern Luzon, bottom temperature 57.4° F. (14.1° C.). Besides at the type station the species was obtained from the following *Albatross* stations: D5113, in 159 fathoms (291 meters), China Sea, off southern Luzon; D5282, in 248 fathoms (454 meters), also China Sea, off southern Luzon, bottom temperature 47.4° F. (8.5° C.); D5301, in 208 fathoms (381 meters), China Sea, off Hongkong, bottom temperature 50.5° F. (10.2° C.); D5424, in 340 fathoms (622 meters), Jolo Sea, bottom temperature 50.4° F. (10.2° C.); D5529, in 441 fathoms (812 meters), between Siquijor and Bohol, bottom temperature 53° F. (11.7° C.); D5570, in 330 fathoms (604 meters), north of Tawi Tawi, bottom temperature 52.3° F. (11.2° C.); and D5650, in 540 fathoms (988 meters), Gulf of Boni, bottom temperature 40.1° F. (4.5° C.).

This is a striking species, with its unequally biconvex form, the difference in the two sides in regard to the length of the chambers and the ornamentation.

Cristellaria helicina—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12015	U.S.N.M.	1	D5113...	13 51 30 N.; 120 50 30 E...	159	° F.	dk. gn. m....	Rare.
9139	} U.S.N.M.	2	D5272...	14 00 00 N.; 120 22 30 E...	118	57.4	m., sh.....	Rare.
11618								
11727	U.S.N.M.	1	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.....	Rare.
11728	U.S.N.M.	5	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.....	Few.
11729	U.S.N.M.	3	D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Rare.
11735	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gy. m., glob..	Rare.
11730	U.S.N.M.	2	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fine s., glob..	Rare.
11731	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Rare.

CRISTELLARIA HELICINOIDES Cushman.

Plate 51, figs. 1 a, b.

Cristellaria helicinoides CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 661.

Description.—Test unequally biconvex, especially in the later chambers; earlier chambers close-coiled, biconvex, numerous; later chambers becoming developed more strongly on one side of the axis

of growth and extending farther toward the umbilicus, giving the test a one-sided appearance; aperture in the last-formed chamber with a tubular neck; wall smooth and unornamented.

Diameter up to 2.5 mm.

Distribution.—Type specimen (U.S.N.M. No. 9140) from *Albatross* station D5301, in 208 fathoms (381 meters), China Sea, vicinity of Hongkong, bottom temperature 50.5° F. (10.2° C.). It also occurred at D5523, northern Mindanao, D5529, in 441 fathoms (812 meters), between Siquijor and Bohol Islands, bottom temperature 53° F. (11.8° C.), and D5537, in 254 fathoms (465 meters), between Negros and Siquijor, bottom temperature 53.5° F. (11.8° C.).

This species, while not distinguished by marks of ornamentation, is nevertheless striking by its change of axis in later growth and consequent helicoid appearance.

Cristellaria helicinoides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9140	U.S.N.M.	1	D5301...	° ' " ° ' "	208	° F. 50.5	gy. m., s....	Rare.
11734	U.S.N.M.	1	D5523...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m., s....	Rare.
11732	U.S.N.M.	1	D5529...	8 48 44 N.; 123 27 35 E..	441	53.0	gy. m., glob.	Rare.
11733	U.S.N.M.	1	D5537...	9 23 45 N.; 123 39 30 E..	254	53.5	gn. m.....	Rare.
				9 11 00 N.; 123 23 00 E..				

CRISTELLARIA ITALICA (Defrance).

Plate 51, fig. 2.

Saracenaria italica DEFANCE, Dict. Sci. Nat., vol. 32, 1824, p. 177; vol. 47, 1827, p. 344; Atlas, Conch., pl. 13, fig. 6.

Cristellaria (Saracenaria) italica D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 293, No. 26; Modèles, 1826, Nos. 19 and 85.

Cristellaria italica PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 3, vol. 16, 1865, pp. 21, 32, pl. 1, figs. 41, 42.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 544, pl. 68, figs. 17, 18, 20–23.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 78, pl. 33, fig. 3.

Specimens of this species occurred in material from more than 30 *Albatross* stations within the area: In the China Sea; Sulu Sea; off Romblon; Sogod Bay, southern Leyte; off Marinduque; east coast of Mindanao; off Panay; off southeastern Mindoro; Palawan Passage; both east and west coasts of Luzon; between Negros and Siquijor; off Sibuko Bay, Borneo; Gulf of Tomini, Celebes; Gulf of Boni; and Macassar Strait.

These stations range in depth from 38 to 752 fathoms (70 to 1,375 meters), with the average depth 288 fathoms (526 meters). The average bottom temperature was 50.2° F. (10.1° C.).

Some specimens were of extra size, measuring 6 to 8 mm. in length. At some stations they were very common, notably D5236, in 494 fathoms (903 meters), off eastern Mindanao, and D5220, in 50 fathoms (91 meters), between Marinduque and Luzon.

Cristelloria italica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15137	U.S.N.M.	2	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m....	Few.
15138	U.S.N.M.	2	D5133...	Islandoff Panabutan Point, N. 52; W. 1.5 mi.	38	gn. m., s....	Few.
15139	U.S.N.M.	4	D5178...	12 43 00 N.; 122 06 15 E..	78	fine. s.....	Few.
15140	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Rare.
15310	U.S.N.M.	2	D5219...	13 21 00 N.; 122 18 45 E..	530	50.8	gn. m.....	Rare.
11714	U.S.N.M.	10	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m....	Common.
15140	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.....	Few.
11712	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Common.
11716								
15141	U.S.N.M.	8	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Frequent.
15142	U.S.N.M.	1	D5261...	12 30 55 N.; 121 34 24 E..	145	s. m.....	Rare.
15143	U.S.N.M.	2	D5272...	14 00 00 N.; 120 22 30 E..	118	57.4	m., sh., co. s.	Few.
15144	U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E..	80	58.6	fine. s....	Rare.
15145	U.S.N.M.	2	D5278...	14 00 10 N.; 120 17 15 E..	102	59.6	fine. s., m., sh.	Few.
15146	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E..	201	50.4	dk. gy. s....	Rare.
15147	U.S.N.M.	6	D5301...	20 37 00 N.; 115 43 00 E..	208	50.5	gy. m., s....	Few.
15148	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s. br. Co.....	Rare.
15149	U.S.N.M.	8	D5318...	21 32 00 N.; 117 46 00 E..	340	co. s., m....	Common.
15150	U.S.N.M.	1	D5338...	11 33 45 N.; 119 24 45 E..	43	bk. s.....	Rare.
15151	U.S.N.M.	1	D5369...	13 48 00 N.; 121 43 00 E..	106	gy. m.....	Few.
15152	U.S.N.M.	2	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Few.
11713	U.S.N.M.	3	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.....	Rare.
15153	U.S.N.M.	6	D5465...	13 39 42 N.; 123 40 39 E..	500	gy. m.....	Few.
11717	U.S.N.M.	9	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.....	Frequent.
15154	U.S.N.M.	1	D5523...	8 43 44 N.; 123 27 35 E..	Rare.
15155	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.....	Rare.
15156	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E..	330	52.3	fine. s., glob.	Rare.
15158	U.S.N.M.	2	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Few.
15157	U.S.N.M.	3	D5589...	4 12 10 N.; 118 38 08 E..	260	45.7	fine. gy. s., gy. m.	Few.
15159	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Rare.
15160	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Few.
15161	U.S.N.M.	2	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Few.
11718	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Rare.

CRISTELLARIA ITALICA (Defrance), var. ACUTO-CARINATA Cushman.

Plate 51, figs. 3a, b.

Cristellaria italica (DEFRANCE), var. *acuto-carinata* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 661.

Description.—Test similar to typical *C. italica* in its general characters, but the angles greatly extended in thin carinae from apex to aperture, the lines of growth being apparent even on the carinae.

Length 3 mm.

Distribution.—Type specimen (U.S.N.M. No. 9141) from *Albatross* station D5523, off Mindanao, depth and bottom temperature not given.

This is rather extreme modification, with the thin carinae in side view nearly as wide as the body of the test.

Cristellaria italica, var. *acuto-carinata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9141	U.S.N.M.	1	D5523...	° ' " ° ' "	8 48 44 N.; 123 27 35 E..	° F.	Rare.

CRISTELLARIA LATIFRONS H. B. Brady.

Cristellaria latifrons H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 544, pl. 68, fig. 19; pl. 113, figs. 11 a, b.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 316, pl. 63, fig. 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 78, pl. 38, fig. 2.

Single specimens which may be referred to this species with some doubt have been obtained from the following stations: *Albatross* D5429, in 766 fathoms (1,401 meters), off eastern Palawan; D5445, east coast of Luzon, 383 fathoms (699 meters), bottom temperature 44.3° F. (6.8° C.); D5537, in 254 fathoms (465 meters), between Negros and Siquijor, bottom temperature 53.3° F. (11.8° C.); D5538, in 256 fathoms (468 meters), same locality and bottom temperature; D5551, in 193 fathoms (353 meters), off Jolo, bottom temperature 53.3° F. (11.8° C.); D5650, Gulf of Boni, in 540 fathoms (988 meters), bottom temperature 40.1° F. (4.5° C.).

The specimens from the first, third, and fourth stations had the chambers rather more closely set than the typical, and that from the last with the long apertural face, but rather more rounded than in the typical.

Cristellaria latifrons—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15170	U.S.N.M.	1	D5429...	° ' " ° ' "	9 41 30 N.; 118 50 22 E..	766	Rare.
15311	U.S.N.M.	1	D5445...	12 44 44 N.; 124 59 50 E..	353	44.3	gn. m., s.....	Rare.
11745	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E..	254	53.3	gn. m.....	Rare.
1517	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gn. m., s.....	Rare.
11744	U.S.N.M.	1	D5551...	5 54 48 N.; 120 44 24 E..	193	53.3	fine. s.....	Rare.
15665	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Rare.

Genus MARGINULINA d'Orbigny, 1826.

MARGINULINA GLABRA d'Orbigny.

Plate 41, fig. 1.

Marginulina glabra D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 259, No. 6; Modèles, 1826, No. 55.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 527, pl. 65, figs. 5, 6.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 313, pl. 60, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 79, pl. 23, fig. 3.

This species was found in material from several *Albatross* stations. These include the following localities: Sulu Sea, off western Mindanao; Sogod Bay, southern Leyte Island; between Marinduque and Luzon; Batangas Bay; China Sea, off southern Luzon; between Siquijor and Bohol Islands; between Negros and Siquijor; and Gulf of Boni.

The various stations range in depth from 170 to 805 fathoms (311 to 1,472 meters), with an average depth of 375 fathoms (686 meters). The bottom temperatures range from 41.2° F. to 53.3° F. (5.1° to 11.8° C.), the average being 50.9° F. (10.5° C.).

There seem to be two more or less distinctive forms in the material examined—one with short globular chambers in the final chambers, the other with more pyriform and elongate ones with a slight neck. Material was insufficient to definitely determine the constancy of these differences, although specimens easily fell into one or other of the two groupings.

Marginulina glabra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12320	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Rare.
12319	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Rare.
12318	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.
15323	U.S.N.M.	1	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn. m., glob.	Rare.
12316	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob.	Rare.
12315	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gn. m., s....	Rare.
12317	U.S.N.M.	1	D5657...	3 19 40 S.; 120 36 30 E..	492	41.2	gy. m.....	Rare.

MARGINULINA STRIATULA Cushman.

Plate 41, fig. 2.

Marginulina striatula CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 79, pl. 23, fig. 4.

Very typical material of this species was obtained from three *Albatross* stations: D5178, in 78 fathoms (143 meters), off Romblon; D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte Island, bottom temperature 52.8° F. (11.5° C.); and D5368, in 181 fathoms (331 meters), off Marinduque Island.

This species is easily distinguished from the preceding by form alone, as it has several chambers forming a moniliform series, with the axis straight and the coiled portion limited to the extreme base. The striations also help as well as the definite, well-developed neck with the four-radiate aperture.

Marginulina striatula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15322	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	fine s.	Rare.
12323	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Rare.
13190	U.S.N.M.	1	D5368...	13 35 30 N.; 121 48 00 E..	181	gy. m.	Rare.

MARGINULINA COSTATA (Batsch).

Plate 41, figs. 5-8.

Nautilus (*Orthoceras*) *costatus* BATSCH, Conch. des Seesandes, 1791, p. 2, pl. 1, figs. 1 a-g.

Marginulina costata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 528, pl. 65, figs. 10-13.—SHERBORN and CHAPMAN, Journ. Roy. Micr. Soc., 1839, p. 487, pl. 11, fig. 28.—TERRIGI, Mem. R. Com. Geol. Italia, vol. 4, 1891, p. 92, pl. 3, fig. 4.—FORNASINI, Mem. R. Accad. Sci. Istit. Bologna, ser. 5, vol. 3, 1893, p. 434, pl. 2, fig. 6; ser. 5, vol. 4, 1894, pp. 213, 214, 217, pl. 2, figs. 18-21.—JONES, Pal. Soc., 1896, p. 235, pl. 1, fig. 21.—SILVESTRI, Mem. Pont. Accad. Nuovi Lincei, vol. 12, 1896, p. 200, pl. 1, fig. 9; vol. 17, 1900, p. 273, pl. 6, fig. 22.—MILLETT (part), Journ. Roy. Micr. Soc., 1902, p. 526 (not pl. 11, fig. 20).

Marginulina raphanus D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 258, No. 1, pl. 10, figs. 7, 8; Modèles, 1826, No. 6.

Description.—Test elongate, subcylindrical, somewhat tapering, early chambers at least in the microspheric form in a partial coil, later ones uniserial and *Nodosaria*-like, chambers of early portion short and broad, later becoming comparatively longer; wall ornamented with 15 to 25 raised costae running lengthwise of the test.

Length up to 4 mm.

Distribution.—Typical material of this species was found in abundance in material from *Albatross* station D5178, in 78 fathoms (143 meters), off Romblon; and at D5277, 80 fathoms (146 meters), China Sea, off southern Luzon, bottom temperature 58.6° F. (14.7° C.).

In the large series obtained both microspheric and megalospheric specimens were present in considerable numbers, the latter predominating, as is usual where both forms are found together in quantity. In the microspheric form the early chambers are distinctly coiled, and in the uncoiling the oblique sutures can be traced for some considerable distance before sutures are developed at right angles to the axis of growth. In the megalospheric form various gradations are seen to the complete *Nodosaria* condition. Various stages are here shown in both the microspheric and megalospheric forms. This series is especially interesting in showing the relation between *Nodosaria* and *Marginulina*, as well as suggesting that a similar relation may be suspected between some of the oblique-sutured species of *Nodosaria* and certain similar uncoiled forms of *Cristellaria*.

Marginulina costata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12322	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E...	78	° F.	fine s.	Common.
12321 15321	U.S.N.M.	10+	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine s.	Common.

MARGINULINA PHILIPPINENSIS, new species.

Plate 53, figs. 2, 3a, b.

Description.—Test elongate, somewhat compressed, earliest chambers coiled, later ones uncoiled in a slightly curved series, all except the last two or three with a thin marginal carina, the last two or three chambers becoming inflated, nearly round in transverse section, well separated from one another, the last-formed one especially so; earliest portion with an irregular ornamentation consisting of numerous bosses or anastomosing costae; the remainder of the test with a few angled costae becoming often more numerous on the last-formed chamber; apertural slightly extended, aperture radiate.

Length 2 to 2.5 mm.

Distribution.—Type specimen (U.S.N.M. No. 14748), from *Albatross* station D5220, between Marinduque and Luzon, 50 fathoms (91 meters).

It has also occurred at station D5217, between Burias and Luzon, 105 fathoms (193 meters), bottom temperature, 63.1° F. (17.2° C.).

This is a common species, especially at D5220, where numerous specimens of a rather uniform character were obtained.

Marginulina philippinensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14747	U.S.N.M.	1	D5217...	13 00 20 N.; 123 14 15 E...	105	° F. 63.1	crs. gy. s. ...	Rare.
14748	U.S.N.M.	10	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m.	Common.

Genus VAGINULINA d'Orbigny, 1826.

VAGINULINA LEGUMEN (Linnaeus).

Plate 41, fig. 3.

Nautilus legumen LINNAEUS, Syst. Nat., ed. 10, 1758, p. 711, No. 248; ed. 12, 1767, p. 1164, No. 288.

Vaginulina legumen D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 257, No. 2.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 530, pl. 66, figs. 13–15.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 314, pl. 60, fig. 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 80, pl. 39, fig. 4.

The typical form with a rounded base or but slightly pointed and a considerably compressed contour occurred less commonly than the following variety. The typical form was found in material from the following stations: *Albatross* D5178, in 78 fathoms (143 meters), off Romblon; D5469, in 500 fathoms (914 meters), off Atulayan Island, off east coast of Luzon; D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); and D5580, in 162 fathoms (297 meters), near Darvel Bay, Borneo, bottom temperature 55.8° F. (13.2° C.).

Vaginulina legumen—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15314	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E...	78	° F.	fne. s.....	Common.
13351	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
13347	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	55.8	gn. m.....	Rare.
13352	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E...	162		br. s., Co....	Rare.

VAGINULINA LEGUMEN (Linnaeus), var. ELEGANS d'Orbigny.

Plate 41, fig. 4.

Vaginulina elegans D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 257, No. 1; Modèles, 1826, No. 54.—PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 3, vol. 16, 1865, p. 27, pl. 1, fig. 33.

Vaginulina legumen LINNAEUS, var. *elegans* FORNASINI, Boll. Soc. Geol. Ital., vol. 5, 1886, p. 25, pl. 1, figs. 1 (?), 2-8.

Description.—Test elongate, tapering, basal end with a stout spine; sutures very distinct, limbate, contour in end view less compressed than in the typical; last-formed chambers more distinct in outline than the earlier ones; wall smooth and transparent.

Distribution.—This variety was much more common, both in number of specimens and number of stations at which it occurred, than the typical. Although found at eight stations, none of these gave specimens of the typical form. The variety was found in material from the following stations: D5198, in 220 fathoms (403 meters), off western Bohol, bottom temperature 53.9° F. (12.1° C.); D5222, in 195 fathoms (356 meters), between Marinduque and Luzon; D5236, in 494 fathoms (903 meters), Pacific Ocean, off east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.); D5261, off southeastern Mindoro, 145 fathoms (265 meters); D5263, in 170 fathoms (311 meters), Batangas Bay; D5349, in 730 fathoms (1,335 meters), Palawan Passage, bottom temperature 40.6° F. (4.7° C.); D5467, in 480 fathoms (878 meters), off Atulayan Island, east coast of Luzon; and D5660, in 692 fathoms (1,266 meters), in Flores Sea.

It will be seen that this variety is well distributed over the area and for the most part is in the deeper waters.

Vaginulina legumen, var. *elegans*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13350	U.S.N.M.	1	D5198...	° ' " ° ' "		° F.		
13349	U.S.N.M.	1	D5222...	9 40 50 N.; 123 39 45 E...	220	53.9	gn. m.....	Rare.
15315	U.S.N.M.	10+	D5236...	13 38 30 N.; 121 48 15 E...	195		gn. m.....	Rare.
15316	U.S.N.M.	2	D5261...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s.....	Common.
12651	U.S.N.M.	1	D5268...	12 30 55 N.; 121 54 24 E...	145		s, m.....	Rare.
12650	U.S.N.M.	5	D5349...	13 42 00 N.; 120 57 15 E...	170		s, p.....	Rare.
13348	U.S.N.M.	1	D5467...	10 54 00 N.; 118 26 20 E...	730	40.6	Co., s.....	Few.
13346	U.S.N.M.	1	D5660...	13 35 27 N.; 123 37 18 E...	480		gy. m.....	Rare.
				5 36 30 S.; 120 49 00 E...	692	39.2	gy. m., s.....	Rare.

VAGINULINA SPINIGERA H. B. Brady.

Plate 42, fig. 1.

Marginulina species, WHITEAVES, Rep. British Association, Brighton Meeting, Trans., 1872, p. 144.

Vaginulina spinigera H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 63; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 531, pl. 67, figs. 13, 14.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 314, pl. 60, fig. 3.

Description.—Test compressed, tapering to the basal end which has two or more very long stout spines, chambers broad and short, early ones coiled, later becoming uniserial, but with the sutures angled; sutures distinct, limbate; aperture radiate.

Length without spines, up to 4 mm.

Distribution.—This species was obtained but once, *Albatross* station D5637, in 700 fathoms (1,280 meters), off Bouro Island. It has not been obtained from the North Pacific. Brady in his description gives it from the Ki Islands, among other localities. It was not found in the Philippine Archipelago.

Vaginulina spinigera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15317	U.S.N.M.	1	D5637...	° ' " ° ' "	700	° F.	gy. m.....	Rare.
				3 53 20 S.; 126 48 00 E...				

VAGINULINA MARGARITIFERA (Batsch).

Plate 42, fig. 2.

Nautilus (Orthoceras) margaritiferus BATSCH, Conch. des Seesandes, 1791, p. 3, pl. 4, figs. 12a-c.

Vaginulina margaritifera H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 532, pl. 66, fig. 16.

Description.—Test elongate, hardly tapering, initial end with a long slender spine, chambers numerous, limbate, sutures oblique, later

chambers more distinct and somewhat inflated, smooth; aperture radiate, on the dorsal side of the chamber.

Length up to 4 mm.

Distribution.—The only station from which I have had this species is *Albatross* D5612, in 750 fathoms (1,370 meters), just south of the Equator, in the Gulf of Tomini, Celebes.

The character of the last-formed chamber in the specimen figured here is rather unusual, but the earlier chambers are typical of this species. It has not been recorded from the North Pacific.

Vaginulina margaritifera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15318	U.S.N.M.	1	D5612...	° ' " ° ' " S.; 121 45 40 E..	750	° F.	Rare.

VAGINULINA BRADYI Cushman.

Plate 42, figs. 3, 4 a, b.

Vaginulina brukenthalii H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 532, pl. 66, figs. 18, 19 (not *V. brukenthalii* Neugeboren).

Vaginulina bradyi CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 661.

Description.—Test elongate, tapering toward the basal end, which is armed with one or more short spines, somewhat compressed, chambers numerous, distinct, early ones coiled, later ones in a linear series but with the sutures oblique throughout, limbate in the central portion, not keeled, in transverse view ovate; aperture at the dorsal side of the chamber, radiate.

Length up to 4 mm.

Distribution.—Type specimen (U.S.N.M. No. 9142) from *Albatross* station D5580, in 162 fathoms (297 meters), Darvel Bay, Borneo, bottom temperatures, 55.8° F. (13.2° C.). It also occurred at the following stations: D5313, in 150 fathoms (270 meters), China Sea, vicinity of Hongkong, bottom temperature 53.6° F. (12° C.); D5315, in 148 fathoms (271 meters), bottom temperature 54.4° F. (12.4° C.); and D5318, in 340 fathoms (622 meters), both in the China Sea, vicinity of Formosa.

This species which was obtained by the *Challenger* in Torres Strait and referred by Brady to the species described by Neugeboren from the Miocene of Transylvania seems to be distinct from that species although it resembles it in certain respects. Our recent species lacks the prominent keel of *B. brukenthalii*, and the greater compression of that species especially in the earlier portion. This new species is evidently a tropical one of the Indo-Pacific region.

Vaginulina bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13357	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	fne. s.....	Rare.
13358	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s.....	Rare.
13353	U.S.N.M.	1	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh.....	Rare.
13354	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Rare.
9142	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co....	Rare.

VAGINULINA LINEARIS (Montagu).

Plate 42, figs. 5-8.

Nautilus linearis MONTAGU, Test. Brit., Suppl., 1808, p. 87, pl. 30, fig. 9.*Dentalina legumen*, var. *linearis* WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 22, pl. 2, figs. 46-48.*Vaginulina linearis* PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 343, pl. 13, figs. 12, 13.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 532, pl. 67, figs. 10-12.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 314, pl. 61, fig. 1.

Description.—Test elongate, hardly tapering, early chambers coiled, later ones becoming uniserial and finally Nodosarian; sutures of early portion rather indistinct, close set, not depressed, of later portion becoming oblique, then directly transverse and toward the apertural end becoming depressed and the chambers inflated; test except the last-formed chambers ornamented with a series of raised costae usually subspiral; last-formed chamber smooth and unornamented; aperture radiate somewhat extended; becoming central and terminal.

Length up to 5 mm. or more.

Distribution.—This species was found at the following stations; *Albatross* D5133, in 38 fathoms (70 meters), Sulu Sea off western Mindanao; D5162, in 230 fathoms (420 meters), off Tawi Tawi group, bottom temperature 52.9° F. (11.6° C.); D5172, in 318 fathoms (582 meters), off Jolo; D5268 in 170 fathoms (311 meters), off eastern Mindoro; D5569, in 303 fathoms (554 meters), north of Tawi Tawi, bottom temperature 52.3° F. (11.2° C.); and D5585, in 476 fathoms (871 meters), off Sibuko Bay, Borneo, bottom temperature 41.1° F. (5° C.).

In certain respects the Philippine material differs from that of the Atlantic. The earlier portion is much more coiled and compressed and the latter portion in full grown specimens more Nodosarian. Recent Atlantic material which I have seems different and the Indo-Pacific material may be a distinct species. Both microspheric and megalospheric specimens occur in the material examined in about equal numbers.

Vaginulina linearis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15319	U.S.N.M.	1	D5133...	° ' " ° ' " Island off Panabutan Point, N. 52° E., 1.50 miles.	38	° F.	gn. m., s.	Rare.
12643	U.S.N.M.	1	D5162...	5 10 00 N.; 119 47 30 E..	230	52.9	crs. s., brk. sh	Rare.
15320	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E..	318	fine s., sh.	Rare.
12641	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.	Rare.
12640	U.S.N.M.	3	D5569...	5 33 15 N.; 120 15 30 E..	303	52.3	co. s.	Rare.
12642	U.S.N.M.	2	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.	Rare.

VAGINULINA ACICULA Cushman.

Plate 40, figs. 1a, b.

Vaginulina acicula CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 661.

Description.—Test elongate, slender, tapering from a point at the initial end to the largest chamber at the apertural end; initial end with one or more short spines; chambers very numerous, rather indistinct in the early portion with oblique sutures, later becoming transverse and slightly depressed; wall smooth and polished; aperture radiate, terminal.

Length up to 5 mm.

Distribution.—Type specimen (U.S.N.M. No. 9143) from *Albatross* station D5613, in 752 fathoms (1,375 meters) Gulf of Tomini, Celebes.

This is a very unusual species in its very slender form, very tapering apex, and much larger apertural end, the aperture becoming central and the last-formed chambers Nodosarian.

Vaginulina acicula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
9143	U.S.N.M.	1	D5613...	° ' " ° ' " 0 42 00 S.; 121 44 00 E....	752	° F.	gy. m.	Rare.

VAGINULINA PROTUMIDA (Schwager).

Plate 40, figs. 3a, b.

Nodosaria protumida SCHWAGER, *Novara*-Exped., Geol. Theil., vol. 2, 1886, p. 227, pl. 6, fig. 59.

Vaginulina protumida CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 81, pl. 30, fig. 1.

This species occurred at *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean off east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.).

Schwager described this species from the Pliocene of Kar Nicobar and I have previously recorded it off Japan.

The figured specimen is somewhat different from more typical ones in the aperture, but has the characteristic form and ornamentation.

Vaginulina protumida—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E...	494	° F. 41.2	fine.gy.s.....	

VAGINULINA PATENS H. B. Brady.

Vaginulina patens H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 533, pl. 67, figs. 15, 16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 80, pl. 32, figs. 7, 8.

Brady described this species from a *Challenger* station in 95 fathoms (174 meters) off the Philippines, and from Torres Strait. I have had specimens very typical in character from two *Albatross* stations, D5198, in 220 fathoms (403 meters), off western Bohol, bottom temperature 53.9° F. (12.1° C.) and D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte, bottom temperature 52.8° F. (11.5° C.).

Vaginulina patens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12652	U.S.N.M.	1	D5198...	° ' " ° ' " 9 40 50 N.; 123 39 45 E...	220	° F. 53.9	gn.m.....	Rare.
13355 13356	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy.s.,m.....	Rare.

Subfamily POLYMORPHININAE.

Genus POLYMORPHINA d'Orbigny, 1826.

POLYMORPHINA REGINA H. B. Brady, Parker, and Jones.

Polymorphina regina H. B. BRADY, PARKER, and JONES, Trans. Linn. Soc., London, vol. 27, 1870, p. 241, pl. 41, figs. 32a, b.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 571, pl. 73, figs. 11–13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 91, pl. 41, figs. 6, 7.

At three stations specimens of this species occurred, at the first station in considerable numbers. Brady records it from off Hongkong. It is a distinctive species. The following are the stations at which it has occurred: D5152, Sulu Archipelago, Tawi Tawi Group, 34 fathoms (62 meters); D5311, China Sea, vicinity Hongkong, 88 fathoms (161 meters); D5580, vicinity of Darvel Bay, Borneo, 162 fathoms (297 meters), bottom temperature 55.8° F. (13.2° C.).

Elsewhere this species is known from New South Wales, at Port Jackson, 6 fathoms (11 meters), Sydney Heads, and Port Stephens; Curtis Strait, Queensland; Bass Strait, 38 fathoms (70 meters); Storm Bay, Tasmania; Torres Strait, 155 fathoms (283 meters); off Ki Islands, 129 fathoms (236 meters); off Admiralty Islands, 17 fathoms (31 meters), and off Honolulu, Hawaiian Islands, 40 fathoms (73 meters) (Brady); off Kerguelen (Egger); Malay Archipelago (Millett); Altona Bay, Victoria, Australia (Chapman); off Ceylon (Dakin); Kerimba Archipelago, east coast of Africa (Heron-Allen and Earland); and east coast of Australia, 465 fathoms (850 meters) (Sidebottom). Chapman has recorded it as a fossil from Australia.

Polymorphina regina—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12620	U.S.N.M.	2	D5152...	5 22 55 N.; 120 15 45 E..	34	° F.	wh. s.	Abundant.
12619	U.S.N.M.	2	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., sh.	Rare.
12618	} U.S.N.M.	2	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.	Rare.
12621								

POLYMORPHINA PROBLEMA d'Orbigny.

Plate 54, figs. 3, 4.

Polymorphina (Guttulina) problema D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826 p. 266, No. 14; Modèles, 1826, No. 61.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 568, pl. 72, fig. 20, pl. 73, fig. 1.

At the following six stations this species has occurred but in few numbers; the stations are for the most part in shallow water: D5143 vicinity of Jolo, 19 fathoms (35 meters); D5144, vicinity of Jolo, 19 fathoms (35 meters); D5152, Sulu Archipelago, Tawi Tawi Group, 34 fathoms (62 meters); D5178, vicinity of Romblon, 78 fathoms (143 meters); D5318, China Sea, vicinity of Formosa, 340 fathoms (622 meters); from exterior of pearl oyster, Jolo Jolo, Philippine Islands.

Polymorphina problema—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12613	U.S.N.M.	1	D5143...	6 05 50 N.; 121 02 15 E..	19	° F.	co. s.	Rare.
14998	U.S.N.M.	1	D5144...	6 05 50 N.; 121 02 15 E..	19	co. s.	Rare.
12610	U.S.N.M.	2	D5152...	5 22 55 N.; 120 15 45 E..	34	wh. s.	Rare.
12612	} U.S.N.M.	7	D5178...	12 43 00 N.; 122 06 15 E..	78	fne. s.	Few.
12615								
12611	U.S.N.M.	1	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.	Rare.
14999	U.S.N.M.	1	Exterior pearl oyster, Jolo Jolo, P. I.	Rare.

POLYMORPHINA PROBLEMA d'Orbigny, var. INDICA, new variety.

Plate 52, figs. 1, 2; plate 53, figs. 1a, b.

Description.—Test large, broad, chambers comparatively few and thick-walled; sutures distinct but only slightly depressed; earlier chambers and the posterior half of the later ones with fine longitudinal costae, apertural end broadly rounded; aperture radiate.

Length up to 2.2 mm.

Distribution.—Type specimen (Cat. No. 16056, U.S.N.M.) from Albatross station D5579, vicinity of Darvel Bay, Borneo, in 175 fathoms (320 meters).

Several specimens were dredged at this station, all of this large robust variety, with partially ornamented chambers.

Polymorphina problema, var. *indica*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
16056	U.S.N.M.	1	D5579...	° ' " ° ' " 4 54 15 N.; 119 19 52 E..	175	° F. 55.3	fine.s., Co....	Rare.

POLYMORPHINA SORORIA Reuss.

Polymorphina (Guttulina) sororia REUSS, Bull. Acad. Roy. Belg., ser. 3, vol. 15, 1863, p. 151, pl. 2, figs. 25–29.

Polymorphina sororia H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 562, pl. 71, figs. 15, 16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 88, pl. 30, fig. 3.

The only station at which specimens of this species were found is D5268, between Verde Island Passage and Batangas Bay, in 170 fathoms (311 meters).

Polymorphina sororia—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12626	U.S.N.M.	2	D5268...	° ' " ° ' " 13 42 00 N.; 120 57 15 E..	170	° F.	s., p.....	Rare.

POLYMORPHINA LACTEA (Walker and Jacob).

"*Serpula tenuis ovalis laevis*" WALKER and BOYS, Test. Min., 1784, p. 2, pl. 1, fig. 5.

"*Polymorpha subcordiformia vel oviformia*" SOLDANI, Testaceographia, vol. 1, pt. 2, 1791, p. 114, pl. 112, figs. 11, *nn.* etc.

Serpula lactea WALKER and JACOB, Adam's Essays, ed. 2, 1798, p. 634, pl. 24, fig. 4. *Vermiculum lacteum* MONTAGU, Test. Brit., 1803, p. 522.

Polymorphina lactea MAGILLIVRAY, Moll. Aberd., 1843, p. 320.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 559, pl. 71, fig. 11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 84, pl. 34, fig. 8.

The specimen belonging to this species was obtained from a single station, D5311, China Sea, vicinity of Hongkong, 88 fathoms (161 meters).

Polymorphina lactea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12665	U.S.N.M.	1	D5311...	° ' " ° ' " 21 33 00 N.; 116 15 00 E..	88	° F.	crs. s., sh.....	Rare.

POLYMORPHINA LACTEA, var. OBLONGA Williamson.

Plate 52, fig. 4.

Polymorphina lactea, var. *oblonga* WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 71, pl. 6, figs. 149, 149a.

There is a single specimen which is very close to this variety as figured by Williamson. It is from *Albatross* station D5601, Gulf of Tomini, Celebes, in 765 fathoms, (1,399 meters).

Polymorphina lactea, var. *oblonga*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
		1	D5601...	° ' " ° ' " 1 13 10 N.; 125 17 05 E...	765	° F.	s., glob., ptr.	Rare.

POLYMORPHINA COMPRESSA d'Orbigny.

Polymorpha "subovalia" SOLDANI, Testaceographia, vol. 1, pt. 2, 1791, p. 114, figs. F. I., etc.

Polymorphina compressa d'ORBIGNY, Foram. Foss. Vienne, 1846, p. 233, pl. 12, figs. 32-34.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 565, pl. 72, figs. 9-11.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 319, pl. 67, fig. 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 89, pl. 40, fig. 3.

From two stations specimens of this species were obtained: D5311, China Sea, vicinity of Hongkong, 88 fathoms (161 meters), and D5315, China Sea, vicinity of Formosa, 148 fathoms (271 meters), bottom temperature 54.4° F. (12.4° C.). The specimens were typical but rare.

Polymorphina compressa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12617	U.S.N.M.	1	D5311...	° ' " ° ' " 21 33 00 N.; 116 15 00 E..	88	° F.	crs. s., sh....	Rare.
12616	U.S.N.M.	1	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh.....	Rare.

POLYMORPHINA GIBBA d'Orbigny.

"*Polymorphina Subcordiformia* vel *Oviformia*" SOLDANI, Testaceographia, vol. 1, pt. 2, 1791, p. 114, pl. 113, figs. ZZ, C, etc.

Polymorphina (Globulina) gibba D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 226, No. 20; Modèles, 1826, No. 63.

Polymorphina gibba H. B. BRADY, PARKER, and JONES (part), Trans. Linn. Soc. London, vol. 27, 1870, p. 216, pl. 39, figs. 2a-d.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 561, pl. 71, figs. 12a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 85, pl. 41, fig. 4.

This species occurred at but two stations, D5121, east coast of Mindoro, 108 fathoms (198 meters), and D5141, vicinity of Jolo, 29 fathoms (53 meters). Single specimens were obtained at each station.

Polymorphina gibba—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
12628	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E..	108	° F.	dk. gn. m....	Rare.
12627	U.S.N.M.	1	D5141...	6 09 00 N.; 120 58 00 E..	29	co. s.....	Rare.

POLYMORPHINA ELEGANTISSIMA Parker and Jones.

Plate 54, figs. 1, 2.

Polymorphina elegantissima PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 438.—H. B. BRADY, PARKER, and JONES, Trans. Linn. Soc., vol. 27, 1870, p. 231, pl. 40, figs. 15a-c.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 566, pl. 72, figs. 12-15.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 319, pl. 67, fig. 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 90, pl. 38, fig. 1.

This species is by far the most abundant in the region, both in the number of stations at which it was obtained as well as in its individuals at different stations. It is especially abundant at many stations less than 100 fathoms in depth, ranging from 17 to 88 fathoms (31 to 161 meters), averaging 37 fathoms (68 meters). At the following stations it occurred at more than 100 fathoms (183 meters): D5315, China Sea, vicinity of Formosa, 148 fathoms (271 meters), bottom temperature 54.4° F. (12.4° C.); D5580, vicinity of Darvel Bay, Borneo, 162 fathoms (297 meters), bottom temperature 55.8° F. (13.2° C.); D5268, Verde Island Passage and Batangas Bay, 170 fathoms (311 meters), bottom temperature not given; D5282, China Sea, vicinity of southern Luzon, 248 fathoms (454 meters), bottom temperature 47.4° F. (8.5° C.); D5569, north of Tawi Tawi, 303 fathoms (554 meters), bottom temperature 52.3° F. (11.2° C.); D5172, vicinity of Jolo, 318 fathoms (582 meters), bottom temperature not given; D5318, China Sea, vicinity of Formosa, 340 fathoms (622 meters), bottom temperature not given.

This is very clearly a species of tropical coral reefs. A peculiar specimen of *Polymorphina* species is shown on plate 52, figure 5.

Polymorphina elegantissima—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15268	U.S.N.M.	4	D5133...	Island off Panabutan Point, N. 52° E., 1.5 miles.	38	° F. 79.5	gn. m., s....	Few.
			D5139...	6 06 00 N.; 121 02 30 E..	20		co. s.....	Few.
13329	} U.S.N.M.	4	D5143...	6 05 50 N.; 121 02 15 E..	19		co. s.....	Rare.
13327								
15262	U.S.N.M.	2	D5145...	6 04 30 N.; 120 59 30 E..	23		co. s., sh.....	Few.
14993	U.S.N.M.	1	D5146...	5 46 40 N.; 120 48 50 E..	24		co. s., sh.....	Rare.
15263	U.S.N.M.	3	D5148...	5 35 40 N.; 120 47 30 E..	17		co. s.....	Few.
15264	U.S.N.M.	10	D5151...	5 24 40 N.; 120 27 15 E..	24		co. s., sh.....	Abundant.
13328	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E..	34		wh. s.....	Common.
			D5153...	5 18 10 N.; 120 02 55 E..	49		co. s., sh.....	Frequent.
15265	U.S.N.M.	3	D5172...	6 03 15 N.; 120 35 30 E..	318		fine. s., sh.....	Common.
12622	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E..	78		fine. s.....	Abundant.
12623	U.S.N.M.	10+	D5179...	12 38 15 N.; 122 12 30 E..	37		hrd. s.....	Abundant.
			D5268...	13 42 00 N.; 120 57 15 E..	170		s., p.....	
15266	U.S.N.M.	1	D5282...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. gy. s.....	Rare.
12624	U.S.N.M.	5	D5311...	21 33 00 N.; 116 15 00 E..	88		crs. s., sh.....	Frequent.
15267	U.S.N.M.	3	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh.....	Few.
12625	U.S.N.M.	6	D5318...	21 32 00 N.; 117 46 00 E..	340		s., br. Co.....	Abundant.
12666	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E..	303	52.3	co. s.....	Rare.
14994	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.....	Rare.
14995	U.S.N.M.	2		Jolo Jolo, Philippine Islands				Rare.
14996	U.S.N.M.	1		Tara Island, Philippine Islands.				Rare.

POLYMORPHINA OBLONGA d'Orbigny.

Plate 52, fig. 3.

Polymorphina oblonga d'ORBIGNY, Foram. Foss. Vienne, 1846, p. 232, pl. 12, figs. 29-31.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 569, pl. 73, figs. 2-4.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 319, pl. 67, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 88, pl. 37, fig. 6.

Next to *Polymorphina elegantissima* this species has occurred at more stations than any other species of the genus. With the exception of a single station, D5151, Sulu Archipelago, Tawi Tawi group, 24 fathoms (44 meters), the stations are all in depths from 148 to 500 fathoms (271 to 914 meters), the average depth 306 fathoms (560 meters). The bottom temperatures were given range from 41.1° to 54.4° F. (5° to 12.4° C.), average 47.9° F. (8.8° C.). Specimens at most of the stations were rare.

Polymorphina oblonga—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12604	U.S.N.M.	1	D5151...	5 24 40 N.; 120 27 15 E..	24		co. s., sh.....	Rare.
12607	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170		s., p.....	Few.
12603	U.S.N.M.	2	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh.....	Rare.
12605	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E..	340		s., br. Co.....	Rare.
12608	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500		gn. m.....	Rare.
14997	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s.....	Rare.
12606	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Rare.
12609	U.S.N.M.	2	D5586...	4 06 50 N.; 118 47 20 E..	347	44	gy. m.....	Rare.

Subfamily UVIGERININAE.

Genus UVIGERINA d'Orbigny, 1826.

UVIGERINA PIGMEA d'Orbigny.

Plate 55, fig. 1.

"*Polymorpha Pineiformia*" SOLDANI, Testaceographia, vol. 1, pt. 2, 1791, pl. 130, figs. *ss, tt*.

Uvigerina pigmea D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 269, pl. 12, figs. 8, 9; Modelès, 1826, No. 67.

Uvigerina pygmaea D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 190, pl. 11, figs. 25, 26.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 575, pl. 74, figs. 11-14.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 96, pl. 42, fig. 1; pl. 44, fig. 5.

Although there are apparently several forms or species in the region with coarsely costate tests no attempt has been made here to separate them. They have occurred at numerous stations, depths varying from 88 to 1,449 fathoms (161 to 2,595 meters), and the bottom temperatures were given from 39.2° to 52.8° F. (4° to 11.5° C.). Many of the specimens were like those figured by Brady, without name in the *Challenger* report.

Uvigerina pigmea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12403	U.S.N.M.	8	D5112...	13 48 22 N.; 120 47 25 E..	177	52.4	dk. gn. m...	Frequent.
12435	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s....	Rare.
12436	U.S.N.M.	3	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Few.
			D5216...	12 52 00 N.; 123 23 30 E..	215	gn. m.....	Frequent.
13118	U.S.N.M.	4	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.....	Few.
			D5235...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Rare.
13373	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s. p.....	Few.
			D5269...	13 39 50 N.; 120 58 30 E..	220	fine. s., p....	Frequent.
			D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....
14975	U.S.N.M.	5	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Few.
14976	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 25 E..	88	co. s.....	Rare.
14985	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Rare.
14984	U.S.N.M.	1	D5660...	5 36 30 S.; 120 49 00 E..	692	39.2	gy. m., s....	Rare.
14986	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Few.
14906	U.S.N.M.	1	Nero 843.	15 21 00 N.; 122 01 15 E..	1,449	gn. m.....	Rare.

UVIGERINA TENUISTRATA Reuss.

Plate 55, fig. 2.

Uvigerina tenuistriata REUSS, Sitz. Akad. Wiss. Wien, vol. 52, 1870, p. 485.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 574, pl. 74, figs. 4-7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 95, pl. 42, fig. 4.

Although this species occurred at numerous stations in the Philippines proper, it occurred but twice in the region to the south—D5590, vicinity of Sibuko Bay, Borneo, 310 fathoms (567 meters), bottom temperature 44.3° F. (6.8° C.), and D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F. (8.6° C.). The

other stations vary in depth from 78 to 604 fathoms (143 to 1,105 meters), average about 275 fathoms (503 meters). The bottom temperatures vary from 41.2° to 62.4° F. (5.1° to 16.8° C.) with the average above 50° F. (10° C.).

Uvigerina tenuistriata—material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15274	U.S.N.M.	3	D5112...	13 48 22 N.; 120 47 25 E...	177	62.4	dk. gn. m...	Frequent.
15275	U.S.N.M.	2	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m...	Few.
13117	U.S.N.M.	6	D5178...	12 43 00 N.; 122 03 15 E...	78	fine. s.....	Few.
12386	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m...	Rare.
12385	U.S.N.M.	1	D5211...	11 51 35 N.; 124 14 00 E...	155	56.6	gn. m., s...	Rare.
14963	U.S.N.M.	1	D5215...	12 31 30 N.; 123 35 24 E...	604	50.5	gn. m.....	Rare.
15276	U.S.N.M.	2	D5222...	13 38 30 N.; 121 42 45 E...	195	52.8	gn. m.....	Rare.
14964	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s...	Few.
12389	U.S.N.M.	1	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fine. s., m., sh.	Rare.
15277	U.S.N.M.	1	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m. s...	Rare.
			D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co...	Few.
			D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.....	Few.
12391	U.S.N.M.	1	D5420...	9 49 35 N.; 123 45 00 E...	127	59.0	Rare.
12387	U.S.N.M.	5	D5590...	4 10 59 N.; 118 39 35 E...	310	44.3	gy. m., s...	Frequent.
12388	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Rare.

UVIGERINA SCHWAGERI H. B. Brady.

Plate 55, figs. 3-5 a, b.

Uvigerina schwageri H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 575, pl. 74, figs. 8-10.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 97, pl. 37, figs. 3, 4.

Brady described this species from the Philippines, and also recorded it from the Fiji Islands and Torres Straits. In the wealth of material examined from the Philippines and adjacent waters it has proved to be by far the most common species in the region. With one exception all these records are from the region of the Philippine Archipelago. The other station is D5621, between Gillolo and Makyan Islands, 298 fathoms (545 meters), where it was apparently rare. The other stations range in depth from 61 to 742 fathoms (112 to 1,357 meters), with the average depth 185 fathoms (338 meters). The bottom temperatures range from 52.8° to 63.6° F. (11.5° to 17.5° C.), with a single station 43.7° F. (6.5° C.). The average temperature, excluding the single station referred to, is 56.1° F. (13.4° C.). Both the average depth and average bottom temperature show perhaps somewhat of the reasons for the limited distribution of this species. Apparently the region south of Japan, where so many species reach their northern limits, is too cold for this species, for it was not noted there. It is a very distinctive species, with its ivory-white, shining test and peculiar shape. It was often found in large numbers, as well as at a large number of stations.

Uvigerina schwageri—material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12398	U.S.N.M.	2	D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s....	Few.
14948	U.S.N.M.	10+	D5121...	13 27 20 N.; 121 17 45 E..	108	dk. gn. m....	Few.
14949	U.S.N.M.	10+	D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m....	Common.
12400	U.S.N.M.	5	D5178...	12 43 00 N.; 122 06 15 E..	78	fn. s....	Frequent.
12392	U.S.N.M.	3	D5243...	10 10 00 N.; 125 04 15 E..	554	53.8	gy. s., m....	Frequent.
			D5261...	6 50 55 N.; 126 14 35 E..	218	63.6	gy. m....	Rare.
			D5272...	12 30 55 N.; 121 34 24 E..	145	s. m....	Common.
				14 00 00 N.; 120 22 30 E..	118	57.4	m., sh., co. s.	Common.
12397	U.S.N.M.	10+	D5277...	13 56 55 N.; 120 13 45 E..	80	58.6	fine s....	Frequent.
12393	U.S.N.M.	6	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., sh....	Common.
14950	U.S.N.M.	2	D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s....	Common.
12394	U.S.N.M.	4	D5315...	21 40 00 N.; 116 58 00 E..	148	54.4	s., sh....	Few.
14951	U.S.N.M.	10+	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Frequent.
14952	U.S.N.M.	4	D5369...	13 48 00 N.; 121 43 00 E..	106	bk. s....	Common.
12399	U.S.N.M.	10+	D5370...	13 44 15 N.; 121 42 30 E..	159	54.3	sft. m....	Abundant.
14953	U.S.N.M.	10+	D5374...	13 46 45 N.; 121 45 08 E..	190	gy. m....	Rare.
12396	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 25 E..	88	co. s....	Frequent.
14954	U.S.N.M.	6	D5408...	10 40 15 N.; 124 15 00 E..	159	55.4	gn. m....	Rare.
14955	U.S.N.M.	1	D5412...	10 09 15 N.; 123 52 00 E..	162	54.8	gn. m....	Few.
14956	U.S.N.M.	1	D5420...	9 49 35 N.; 123 45 00 E..	127	59.0	Few.
14957	U.S.N.M.	2	D5481...	10 27 30 N.; 125 17 10 E..	61	s., sh., g....	Frequent.
			D5523...	8 48 44 N.; 123 27 35 E..	Few.
12395	U.S.N.M.	3	D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	gn. m....	Common.
14958	U.S.N.M.	5	D5538...	9 08 15 N.; 123 23 20 E..	256	gn. m., s....	Few.
14959	U.S.N.M.	3	D5621...	0 15 00 N.; 127 24 35 E..	298	gy. and bk. s.	Rare.
14960	U.S.N.M.	1		Tara Island, Philippine Islands.	Rare.
14961	U.S.N.M.	3			

UVIGERINA ANGULOSA Willamson.

Uvigerina angulosa WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 67, pl. 5, fig. 140.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 576, pl. 74, figs. 15–18.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 98, pl. 44, fig. 4.

This species is very rare in the region as far as the records show, having been found but three times. They are as follows: D5543, northern Mindanao and vicinity, 162 fathoms (297 meters), bottom temperature 54.5° F. (12.5° C.); D5575, north of Tawi Tawi, 315 fathoms (580 meters), bottom temperature 52.3° F. (11.2° C.); and D5637, Bouro Island (south) and vicinity, 700 fathoms (1,280 meters).

Uvigerina angulosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12415	U.S.N.M.	1	D5543...	8 47 15 N.; 123 35 00 E..	162	54.5	fine s., brk. sh.	Rare.
12413	U.S.N.M.	1	D5575...	5 28 30 N.; 120 02 27 E..	315	52.3	Co., s....	Rare.
12414	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m....	Rare.

UVIGERINA STRIATA d'Orbigny.

Plate 54, fig. 6.

Uvigerina striata D'ORBIGNY, Amér. Mérid., vol. 5, pt. 5, 1839, "Foraminifères," p. 53, pl. 7, fig. 16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 94, pl. 43, fig. 5.

A few specimens which seem to be this species have occurred in comparatively deep water at three stations: D5126, 742 fathoms

(1,357 meters), Sulu Sea, off southern Panay; D5300, China Sea, off southern Luzon, 265 fathoms (485 meters); and D5318, 340 fathoms (622 meters), China Sea, off Formosa. It has been recorded in the Pacific from off the Hawaiian Islands.

Uvigerina striata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14962	U. S. N. M.	1	D5126...	° ' " ° ' "		° F.		
13108	U. S. N. M.	1	D5300...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m ...	Rare.
12409	} U. S. N. M.	2	D5318...	20 31 00 N.; 115 49 00 E..	265	gy. m., s	Rare.
13120								

UVIGERINA SELSEYENSIS Heron-Allen and Earland.

Uvigerina selseyensis HERON-ALLEN and EARLAND, Journ. Roy. Micr. Soc., 1909 p. 437, pl. 18, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 93, pl. 42, fig. 5.

From a single station this species was found, and the specimen even more like the typical than that I have figured previously. It is from D5575, north of Tawi Tawi, 315 fathoms (580 meters), bottom temperature 52.3° F. (11.2° C.). The depth agrees very well with that from which I have recorded the species previously in the North Pacific.

Uvigerina selseyensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
		1	D5575...	° ' " ° ' "		° F.		
				5 28 30 N.; 120 02 27 E..	315	52.3	co. s.....	Rare.

UVIGERINA BRUNNENSIS Karrer.

Uvigerina brunnensis KARRER, Abh. k. geol. Reichs., vol. 9, 1877, p. 385, pl. 16b, fig. 49.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 577, pl. 75, figs. 4, 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 97, pl. 43, fig. 2.

Specimens which were very much like the *Challenger* figures of this species were found at the following few stations: D5115, Balayan Bay and Verde Island Passage, 340 fathoms (622 meters); D5523, northern Mindanao and vicinity; D5606, Gulf of Tomini, Celebes, 834 fathoms (1,525 meters); D5609, north of Celebes, 1,092 fathoms (1,998 meters), bottom temperature 36.3° F. (2.3° C.); D5626, between Gillolo and Kayoa Islands, 265 fathoms (485 meters); and D5652, Gulf of Boni, 525 fathoms (960 meters), bottom temperature 41.2° F. (5.1° C.).

Uvigerina brunnensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12407	U.S.N.M.	9	D5115...	13 37 11 N.; 120 43 40 E..	340	Frequent.
			D5523...	8 48 44 N.; 123 27 35 E..	
12411	U.S.N.M.	2	D5606...	0 16 28 N.; 121 33 30 E..	834	gn. m.	Rare.
12408	U.S.N.M.	2	D5609...	0 11 00 S.; 121 16 00 E..	1,092	36.3	gn. m.	Rare.
12406	U.S.N.M.	1	D5626...	0 07 30 N.; 127 29 00 E..	265	gy. m., fnc.s.	Rare.
12410	U.S.N.M.	2	D5652...	4 35 00 S.; 121 23 06 E..	525	41.2	gn. m.	Rare.

UVIGERINA ACULEATA d'Orbigny.

Plate 55, fig. 6.

Uvigerina aculeata D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 191, pl. 11, figs. 27, 28.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 578, pl. 75, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 100, pl. 43, fig. 4.

Next to *Uvigerina schwageri* this is the most common species in the region, being found at a large number of stations well scattered throughout the area covered by the dredgings. The depths range from 234 to 805 fathoms (428 to 1,472 meters), and the bottom temperatures from 41.1° to 56.4° F. (4.5° to 13.5° C.). The average depth is 440 fathoms (805 meters), and the average bottom temperature 48.4° F. (9.1° C.).

The specimens are excellent ones and at many stations are abundant, often the whole test covered with the long spinose growth shown in the figures given by Brady in the *Challenger* Report.

Uvigerina aculeata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s.	Rare.
14971	U.S.N.M.	2	D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sit. gn. m.	Rare.
14972	U.S.N.M.	1	D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	sit. gn. m.	Few.
12424	U.S.N.M.	5	D5201...	10 10 00 N.; 125 04 15 E..	554	53.8	gy. s., m.	Frequent.
12425	U.S.N.M.		D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s.	Few.
14973	U.S.N.M.	10+	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Abundant.
12423	U.S.N.M.		D5260...	12 25 35 N.; 121 31 35 E..	234	51.4	gn. m., s.	Frequent.
12426	U.S.N.M.	5	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s.	Few.
14974	U.S.N.M.		D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.	Few.
			D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co., s.	Few.
15271	U.S.N.M.	2	D5300...	9 38 30 N.; 121 11 00 E..	508	49.8	gy. m., co. s.	Few.
15272	U.S.N.M.	2	D5445...	12 44 42 N.; 124 59 50 E..	883	44.3	gn. m., s.	Few.
12412	U.S.N.M.	3	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.	Rare.
			D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.	Rare.
14977	U.S.N.M.	1	D5512...	8 16 02 N.; 123 58 26 E..	445	52.5	gy. m., fnc.s.	Few.
14978	U.S.N.M.	2	D5526...	9 12 45 N.; 123 43 30 E..	805	52.3	gn. m., glob.	Rare.
14979	U.S.N.M.	1	D5537...	9 11 09 N.; 123 23 00 E..	254	53.5	gn. m.	Few.
14980	U.S.N.M.	2	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.	Few.
15273	U.S.N.M.	4	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.	Frequent.
12422	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gy. m., s.	Frequent.
12427	U.S.N.M.	5	D5650...	4 53 45 S.; 121 29 00 E..	540	gn. m.	Rare.
14981	U.S.N.M.	1	D5652...	4 35 00 S.; 121 23 06 E..	525	41.2	gn. m.	Few.
14982	U.S.N.M.	3	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.	Few.
12428	U.S.N.M.	2						
14983	U.S.N.M.	2						

UVIGERINA ASPERULA Czjzek.

Plate 54, fig. 5; plate 55, fig. 8.

Uvigerina asperula CZJZEK, Haidinger's Nat., Abh. 2, 1848, p. 146, pl. 13, figs. 14, 15.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 578, pl. 75, figs. 6-8.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 320, pl. 68, fig. 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 101, pl. 43, fig. 1.

This species is almost as frequent in its occurrence as *Uvigerina aculeata* and has a similar distribution in the region. The range of depth is the same, 234 to 901 fathoms (428 to 1,648 meters), the average 445 fathoms (814 meters) being slightly more. The bottom temperatures range from 38.2° to 56.4° F. (3.4° to 13.5° C.) with the average 46.5° F. (8.0° C.). This is slightly lower also than for *U. aculeata*. One station was from 24 fathoms (44 meters).

Uvigerina asperula—Material examined.

Cat. No.	Coll. of—	No. of specimns.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
14965	U.S.N.M.	2	D5126...	10 34 45 N.;	121 47 30 E..	742	49.5	sft. gn. m....	Rare.
12416	U.S.N.M.	2	D5201...	10 10 00 N.;	125 04 15 E..	554	52.8	gy. s., m....	Few.
			D5210...	11 49 55 N.;	124 28 05 E..	393	43.7	gn. m., s....	Rare.
12417	U.S.N.M.	2	D5236...	8 50 45 N.;	126 26 52 E..	494	41.2	fne. gy. s....	Few.
12418	U.S.N.M.	3	D5259...	11 57 30 N.;	121 42 15 E..	312	49.3	gy. m., glob..	Few.
14966	U.S.N.M.	4	D5260...	12 25 35 N.;	121 31 35 E..	234	51.4	gn. m., s....	Few.
			D5300...	20 31 00 N.;	115 49 00 E..	265	gy. m., s....	Few.
			D5348...	10 57 45 N.;	118 38 15 E..	375	56.4	co. s.....	Few.
12420	U.S.N.M.	1	D5445...	12 44 42 N.;	124 59 50 E..	383	44.3	gn. m., s....	Rare.
15269	U.S.N.M.	2	D5526...	9 12 45 N.;	123 45 30 E..	805	52.3	gn. m., glob..	Few.
			D5537...	9 11 00 N.;	123 23 00 E..	254	gn. m.....	Few.
15270	U.S.N.M.	1	D5601...	1 13 10 N.;	125 17 05 E..	765	s. glob., ptr..	Few.
14967	U.S.N.M.	3	D5618...	0 37 00 N.;	127 15 00 E..	417	gy. m.....	Few.
12419	U.S.N.M.	10+	D5640...	4 27 00 S.;	122 55 40 E..	24	s., brk. sh....	Abundant.
			D5648...	5 35 00 S.;	122 20 00 E..	559	39.2	gn. m.....	Few.
14968	U.S.N.M.	2	D5650...	4 53 45 S.;	121 23 00 E..	540	gn. m.....	Few.
12421	U.S.N.M.	2	D5652...	4 35 00 S.;	121 23 06 E..	525	41.2	gn. m.....	Few.
14969	U.S.N.M.	3	D5666...	2 54 30 S.;	118 47 00 E..	272	47.5	gn. m.....	Few.
			D5668...	2 28 15 S.;	118 49 00 E..	901	38.2	gy. m.....	Rare.
14970	U.S.N.M.	2	H4937...	5 32 50 N.;	120 49 10 E..	385	38.2	gy. m.....	Few.

UVIGERINA AMPULLACEA H. B. Brady.

Plate 55, fig. 7.

Uvigerina asperula CZJZEK, var. *ampullacea* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 579, pl. 75, figs. 10, 11.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 320, pl. 68, fig. 5.

Uvigerina ampullacea EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 313, pl. 9, fig. 37.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 102, pl. 42, fig. 3.

Although less common than some of the other species this has occurred at too large a number of stations to list separately. They range in depth from 265 to 885 fathoms (485 to 1,618 meters), average 550 fathoms (1,005 meters), and bottom temperatures from 38.2° to 56.4° F. (3.4° to 13.5° C.), the average 45.6° F. (7.5° C.).

In the region this species has occasionally occurred with *U. asperula*, but more often not, and on the whole they seem well distinguished from one another.

Uvigerina ampullacea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
14987	U. S. N. M.	3	D5126...	10 34 45 N.;	121 47 30 E..	742	49.5	sft. gn. m.	Rare.
12430	U. S. N. M.	1	D52365...	6 04 20 N.;	120 59 20 E..	494	41.2	fine gy. s.	Rare.
12433	U. S. N. M.	3	D5259...	11 57 30 N.;	121 42 15 E..	312	49.3	gy. m., glob.	Frequent.
14988	U. S. N. M.	9	D5284...	13 42 05 N.;	120 30 45 E..	422	42.3	gy. m., s.	Few.
12429	U. S. N. M.	1	D5300...	20 31 00 N.;	115 49 00 E..	265	co. s.	Few.
14989	U. S. N. M.	2	D5348...	10 57 45 N.;	118 38 15 E..	375	56.4	co. s.	Few.
12431	U. S. N. M.	2	D5349...	10 54 00 N.;	118 26 20 E..	730	40.6	co. s.	Few.
			D5525...	9 12 45 N.;	123 45 30 E..	805	52.3	gn. m., glob.	Few.
12434	U. S. N. M.	1	D5572...	5 31 26 N.;	120 09 45 E..	394	52.3	s.	Rare.
14990	U. S. N. M.	1	D5538...	3 47 15 S.;	126 23 40 E..	517	fine gy. s.	Rare.
14991	U. S. N. M.	2	D5648...	5 35 00 S.;	122 20 00 E..	559	39.2	gn. m.	Rare.
12432	U. S. N. M.	1	D5660...	5 36 30 S.;	120 49 00 E..	692	39.2	gy. m., s.	Few.
14992	U. S. N. M.	6	H4937...	5 32 50 S.;	120 49 10 E..	885	38.2	gy. m.	Frequent.

UVIGERINA INTERRUPTA H. B. Brady.

Uvigerina interrupta H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 60, pl. 8, figs. 17, 18; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 580, pl. 75, figs. 12-14.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 103, pl. 44, fig. 1.

At only five stations, four of which were in the southern part of the region, this species has occurred sparingly. The depths are rather less than recorded for the North Pacific. The stations are as follows: D5185, between Panay and Negros, 638 fathoms (1,167 meters), bottom temperature 49.8° F. (9.8° C.); D5624, between Gilolo and Makyan Islands, 288 fathoms (527 meters); D5637, Pitt Passage, 700 fathoms (1,280 meters); D5650, Gulf of Boni, 540 fathoms (988 meters), bottom temperature 40.1° F. (4.5° C.); D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F. (8.6° C.).

Uvigerina interrupta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
12404	U. S. N. M.	7	D5185...	10 05 45 N.;	122 18 30 E..	638	49.8	gn. m.	Frequent.
12405	U. S. N. M.	1	D5624...	0 12 15 N.;	127 29 30 E..	288	fine s., m.	Rare.
12401	U. S. N. M.	1	D5637...	3 53 20 S.;	126 48 00 E..	700	gy. m.	Rare.
			D5650...	4 53 45 S.;	121 29 00 E..	540	40.1	gn. m.	Rare.
12402	U. S. N. M.	1	D5666...	2 54 30 S.;	118 47 00 E..	272	47.5	gn. m.	Rare.

UVIGERINA CANARIENSIS d'Orbigny.

"Testae pineiformes minusculae" SOLDANI, Testaceographia, vol. 2, 1798, p. 8, pl. 4, figs. E, F, G, H.

Uvigerina nodosa, var. β , d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 269, No. 3.

Uvigerina canariensis d'ORBIGNY, Foram. Canaries, 1839, p. 138, pl. 1, figs. 25-27.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 573, pl. 74, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 92, pl. 42, fig. 6.

The only station at which this species was noted is D5300, China Sea, vicinity of southern Luzon, in 265 fathoms (485 meters).

Uvigerina canariensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5300...	° ' " ° ' " 20 31 00 N.; 115 49 00 E...	265	° F.	gy. m., s....	

UVIGERINA SCROBICULATA, new species.

Description.—Test elongate, polygonal, early chambers generally triserial in a three-sided test; the angles rounded, latter half of the test triserial and tapering toward the apertural end; last-formed chamber and sometimes the one preceding somewhat distinct from the general mass of the test, triangular at the base, and broadest thence tapering rapidly to the aperture, which is rounded with a slight tubular neck and a very slightly developed phialine lip; sutures distinct but not greatly depressed, surface smooth except for a finely scrobiculate condition, due to numerous irregular pittings of the surface; color yellowish- or brownish-white.

Length up to 1.5 mm.

Distribution.—Type-specimen (Cat. No. 16057, U.S.N.M.), from *Albatross* station D5333, in 310 fathoms (567 meters), Mindoro Strait, where the species is common.

This has some of the characters of *U. angulosa* and *U. schwageri* but is distinct from either. The almost fusiform shape, broadest in the middle and tapering toward either end with the generally triangular section, both of the whole test and of the chambers with the finely scrobiculate surface, will distinguish it.




FIG. 10.—UVIGERINA SCROBICULATA, NEW SPECIES. $\times 50$. a, b, TWO VIEWS FROM DIFFERENT ANGLES. FROM STATION D5333.

Uvigerina scrobiculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
16057	U.S.N.M.	1	D5333...	° ' " ° ' " 12 26 30 N.; 120 37 45 E...	310	° F.	s.....	Common.

Genus SIPHOGENERINA Schlumberger, 1883.

SIPHOGENERINA COLUMELLARIS (H. B. Brady).

Plate 56, fig. 1.

Sagrina columellaris H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 64; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 581, pl. 75, figs. 15–17.

Siphogenerina columellaris EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 316, pl. 9, figs. 28, 31, 33.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 104, pl. 47, figs 2, 3.

Typical specimens of this species were obtained from material dredged at the following stations: D5133, Sulu Sea off western Mindanao, 38 fathoms (70 meters); D5172, vicinity of Jolo, 318 fathoms (582 meters); D5218, between Burias and Luzon, 20 fathoms (37 meters); D5236, Pacific Ocean east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5268, Verde Island Passage and Batangas Bay, 170 fathoms (311 meters); D5469, east coast Luzon, San Bernardino Strait to San Miguel Bay, 500 fathoms (914 meters); D5572, north of Tawi Tawi, 334 fathoms (611 meters); bottom temperature 52.3° F. (11.2° C.); D5638, Bouro Island (south) and vicinity, 517 fathoms (946 meters).

Siphogenerina columellaris—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5133...	° ' " ° ' " Island off Panabutan Point N. 52° E., 1.50 miles.	38	° F.	gn. m., s....	
13112	U.S.N.M.	2	D5172...	6 03 15 N.; 120 35 30 E...	318	fine. s., sh....	Rare.
13111	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s....	Rare.
13110	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Rare.
13109	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p....	Rare.
11313	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.	Rare.
			D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s....	
14929	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E...	517	fine. gy. s....	Rare.

SIPHOGENERINA BIFRONS (H. B. Brady).

Plate 56, figs. 2, 3.

Sagrina bifrons H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 64; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 582, pl. 75, figs. 18-20.

Siphogenerina (Sagrina) bifrons EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 317, pl. 9, figs. 25, 26, 29.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 105, pl. 45, figs. 1, 2, 5-7.

This species occurred at a number of stations in the region as follows: D5121, east coast of Mindoro, 108 fathoms (198 meters); D5138, vicinity of Jolo, 19 fathoms (35 meters); D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5242, Pujada Bay and vicinity, 191 fathoms (350 meters), bottom temperature 64.1° F. (17.8° C.); D5281, China Sea, vicinity southern Luzon, 201 fathoms (368 meters), bottom temperature 50.4° F. (10.2° C.); D5318, China Sea, vicinity of Formosa, 340 fathoms (622 meters); D5445, east coast of Luzon, 383 fathoms (699 meters), bottom temperature 44.3° F. (6.8° C.); D5538, between Negros and Siquijor, 256 fathoms (468 meters), bottom temperature 53.3° F. (11.8° C.); D5630, south of Patiente Strait, 569 fathoms (1,040 meters); D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F. (8.6° C.).

It will be seen that the depths agree well with those already published from the North Pacific, and it is also interesting that the bottom temperatures are comparatively high, as was the case in the stations noted off Japan. With these added stations we now know considerable about the distribution of this species, which was described by Brady from a single *Challenger* station off Japan. Its range can now be well filled in from off southern Japan to Australia, including the Malay Archipelago, and probably occurs widely distributed among the East Indian and Oceanic Islands. Chapman records it from off Funafuti.

Siphogenerina bifrons—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13141	U.S.N.M.	10	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Common.
14938	U.S.N.M.	1	D5242...	6 51 53 N.; 126 14 10 E...	191	64.1	sft.gy.m....	Rare.
13140	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk.gy.s....	Rare.
14939	U.S.N.M.	3	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co....	Rare.
14940	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s....	Rare.
13133	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s....	Rare.
13137	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m....	Rare.
13139	U.S.N.M.	10	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Common.
13143	U.S.N.M.	1	D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m...	Rare.
12751	U.S.N.M.	8	D5138...	6 06 00 N.; 120 58 50 E...	19	s., Co.....	Common.
13138	U.S.N.M.	1	N e r o	Rare.
14907	U.S.N.M.	1	849.	Rare.

SIPHOGENERINA BIFRONS (H. B. Brady), var. STRIATULA Cushman.

Plate 56, fig. 4.

Siphogenerina bifrons (H. B. BRADY), var. *striatula* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 662.

Description.—Differing from the typical in having the surface with numerous longitudinal striations, rather more elongate, and the central indented portion deeper and more defined.

At the following stations the variety occurred, sometimes with the typical but usually much more numerous: H4898, Sulu Sea off western Mindanao, 221 fathoms (405 meters); D5110, China Sea, off southern Luzon, 135 fathoms (247 meters), bottom temperature 59° F. (15° C.); D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5268, Verde Island Passage and Batangas Bay, 170 fathoms (311 meters); D5445, east coast of Luzon, San Bernardino Strait to San Miguel Bay, 383 fathoms (699 meters), bottom temperature 44.3° F. (6.8° C.); D5523, northern Mindanao and vicinity; D5621, between Gillolo and Makyan Islands, 298 fathoms (545 meters); D5636, Pitt Passage, 1,262 fathoms (2,309

meters); D5638, Bouro Island (south) and vicinity, 517 fathoms (946 meters); D5666, Macassar Strait, 272 fathoms (498 meters), bottom temperature 47.5° F. (8.6° C.).

Both microspheric and megalospheric forms occur, the megalospheric more common. At the only station at which a record was kept 11 were megalospheric and two were microspheric.

Siphogenerina bifrons, var. *striatula*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
				° ' " ° ' "		° F.		
14941	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E...	135	59.0	dk. gy. m...	Rare.
9144	} U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m...	Rare.
11642								
13134	U.S.N.M.	2	} D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Few.
14942	U.S.N.M.	3						
14943	U.S.N.M.	10+		D5268...	13 42 00 N.; 120 57 15 E...	170	p., s., sh....
13142	U.S.N.M.	3	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s....	Few.
14944	U.S.N.M.	2	D5523...	8 48 44 N.; 123 27 35 E...	gy. m., fne. s	Rare.
13135	U.S.N.M.	1	D5621...	0 15 00 N.; 127 24 35 E...	298	gy. and bk. s	Rare.
13136	U.S.N.M.	1	D5638...	1 55 00 S.; 127 42 30 E...	1,262	gy. m., fne. s	Rare.
14945	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E...	517	fne. gy. s....	Rare.
14946	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Rare.
14924	U.S.N.M.	10+	H4898...	7 43 45 N.; 122 03 45 E...	221	gy. m., glob.	Common.

SIPHOGENERINA DIMORPHA (Parker and Jones).

Plate 56, fig. 8.

Uvigerina (Sagrina) dimorpha PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 420, pl. 18, fig. 18.

Sagrina dimorpha H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 582, pl. 76, figs. 1-3.

Siphogenerina dimorpha EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 317, pl. 9, fig. 30.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 106, pl. 45, figs. 3, 4.

This species was the least common of any of the genus, occurring as single specimens or at most few. The following records are all that were noted: D5300, China Sea, vicinity of southern Luzon, 265 fathoms (485 meters); D5601, Gulf of Tomini, Celebes, 765 fathoms (1,399 meters); and D5666, Macassar Strait, 272 fathoms (498 meters). The last station had a bottom temperature of 47.5° F. (8.6° C.), the others not being recorded.

Siphogenerina dimorpha—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° P.		
13114	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s....	Rare.
13116	U.S.N.M.	1	D5601...	1 13 10 N.; 125 17 05 E...	765	s., glob., ptr	Rare.
13115	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Rare.

SIPHOGENERINA STRIATA (Schwager).

Plate 56, fig. 5.

Dimorphina striata SCHWAGER, *Novara-Exped.*, Geol. Theil., vol. 2, 1866, p. 251, pl. 7, fig. 99.

Sagrina striata SCHWAGER, *Boll. R. Com. geol. Ital.*, vol. 8, 1877, p. 25, pl., fig. 35.—H. B. BRADY, *Rep. Voy. Challenger*, Zoology, vol. 9, 1884, p. 584, pl. 75, figs. 25, 26.

Siphogenerina (Sagrina) striata EGGER?, *Abh. kön. bay. Akad. Wiss. München*, Cl. II, vol. 18, 1893, p. 316, pl. 9, figs. 32, 34, 35, 64, 65.—CUSHMAN, *Bull.* 71, U. S. Nat. Mus., pt. 3, 1913, p. 107, pl. 47, figs. 4, 5.

Specimens have occurred at numerous stations, at some of which they could be called common. The species seems to be different from either form of *S. raphanus*, the costae being less independent of the chambers than in *S. raphanus* and the sutures marking well-defined limits even in the costae. The neck is invariably short or wanting, no lip being present. It was found at the following stations: H4898, Sulu Sea off western Mindanao, 221 fathoms (405 meters); D5178, vicinity of Romblon, 78 fathoms (143 meters); D5201, Sogod Bay, southern Leyte Island, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5236, Pacific Ocean, east coast Mindanao, 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.); D5242, Pujada Bay and vicinity, 191 fathoms (350 meters), bottom temperature 64.1° F. (17.8° C.); D5268, Verde Island Passage and Batangas Bay, 170 fathoms (311 meters); D5318 China Sea, vicinity of Formosa, 340 fathoms (622 meters); D5443, east coast of Luzon, San Bernardino Strait to San Miguel Bay, 241 fathoms (441 meters), bottom temperature 51.3° F. (10.7° C.); D5575, north of Tawi Tawi, 315 fathoms (580 meters), bottom temperature 52.3° F. (11.2° C.).

Siphogenerina striata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13124	U.S.N.M.	8	D5178...	12 43 00 N.; 122 06 15 E..	78	fne. s.....	Common.
13123	} U.S.N.M.	6	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	Common.
12750								
14927	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s....	Rare.
14928	U.S.N.M.	1	D5242...	6 51 53 N.; 126 14 10 E..	191	64.1	sit. gy. m.....	Rare.
12749	U.S.N.M.	3	D5268...	13 42 00 N.; 120 57 15 E..	170	p., s., sh.....	Few.
13122	U.S.N.M.	3	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.....	Few.
13121	U.S.N.M.	1	D5443...	12 43 05 N.; 125 01 00 E..	241	51.3	co. s., sh.....	Rare.
14923	U.S.N.M.	1	D5575...	5 28 30 N.; 120 02 27 E..	315	52.3	Co., s.....	Rare.
			H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob..	Rare.

SIPHOGENERINA RAPHANUS (Parker and Jones).

Plate 56, fig. 7.

Uvigerina (Sagrina) raphanus PARKER and JONES, *Philos. Trans.*, vol. 155, 1865, p. 364, pl. 18, figs. 16, 17.

Sagrina raphanus H. B. BRADY, *Rep. Voy. Challenger*, Zoology, vol. 9, 1884, p. 585, pl. 75, figs. 21–24.

Siphogenerina (Sagrina) raphanus EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 317, pl. 9, fig. 36.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 108, pl. 46, figs. 1-5.

This species is the most common of the genus in the region, occurring at numerous stations well scattered over the region, and from Apra Bay, Guam. In depth the stations range from 78 to 554 fathoms (143 to 1,012 meters), and the bottom temperatures from 47.5° F. to 59° F. (8.6° C. to 15° C.). All but one station had a bottom temperature above 51° F. (10.5° C.). In its typical form this species has a tapering shape, acute at the initial end, with a narrow neck and well-developed phialine lip, with a few strong costae, independent of the sutures.

Siphogenerina raphanus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° " " ° "		° F.		
13128	U.S.N.M.	8	D5178...	12 43 00 N.; 122 06 15 E...	78		fine.s.....	Common.
14930	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	53.8	gy.s.,m.....	Rare.
12745	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E...	170		s.,p.....	Rare.
14931	U.S.N.M.	3	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine.s.....	Few.
14932	U.S.N.M.	2	D5313...	21 30 00 N.; 116 43 00 E...	150	53.6	s.....	Few.
14933	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E...	340		s.,br.Co.....	Few.
13125	U.S.N.M.	1	D5420...	9 49 35 N.; 123 45 00 E...	127	59.0		Few.
14934	U.S.N.M.	1	D5443...	12 43 05 N.; 125 01 00 E...	241	51.3	co.s.,sh.....	Rare.
			D5523...	8 48 44 N.; 123 27 35 E...				Rare.
14935	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn.m.,s.....	Rare.
12747	U.S.N.M.	1	D5551...	5 54 48 N.; 120 44 24 E...	193	53.3	fine.s.....	Few.
14936								
14937	U.S.N.M.	1	D5565...	5 51 42 N.; 120 30 30 E...	243	52.3	s.,ptr.,sh.....	Rare.
			D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.....	Few.
13126	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn.m.....	Few.

SIPHOGENERINA RAPHANUS (Parker and Jones), var. COSTULATA Cushman.

Plate 56, fig. 6.

Siphogenerina raphanus (PARKER and JONES), var. *costulata* CUSHMAN, Proc U. S. Nat. Mus., vol. 51, 1917, p. 662.

This variety differs from the typical in its much more cylindrical form, its few very definite costae, and the very short, wide neck without a lip. This variety does not always occur with the typical, but in all the material seen there appear to be no intermediate forms. Both microspheric and megalospheric forms were noted. The variety occurred at the following stations: From the exterior of pearl oyster, Jolo Jolo; D5134, Sulu Archipelago, near Basilan Island, 25 fathoms (46 meters); D5201, Sogod Bay, southern Leyte, 554 fathoms (1,012 meters), bottom temperature 52.8° F. (11.5° C.); D5575, north of Tawi Tawi, 315 fathoms, bottom temperature 52.3° F.; D5142, vicinity of Jolo, 21 fathoms (39 meters); D5143, vicinity of Jolo, 19 fathoms (35 meters); D5469, east coast of Luzon, San Bernardino Strait to San Miguel Bay, 500 fathoms (914 meters); D5546, Jolo Island and vicinity, 138 fathoms (252 meters), bottom temperature

58.3° F. (14.6° C.); D5572, north of Tawi Tawi, 334 fathoms (611 meters), bottom temperature 52.3° F. (11.2° C.); D5580, vicinity of Darvel Bay, Borneo, 162 fathoms (297 meters), bottom temperature 55.8° F. (13.2° C.); D5636, Pitt Passage, 1,262 fathoms (2,309 meters).

Siphogenerina raphanus, var. *costulata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13132	U.S.N.M.	1	D5134...	6 44 45 N.; 121 48 00 E...	25	fine s.....	Rare.
9145	U.S.N.M.	1	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s., sh.....	Rare.
			D5142...	6 06 10 N.; 121 02 40 E...	21	gy. s., m.....	Few.
13130	U.S.N.M.	5	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gn. m.....	Few.
13131	U.S.N.M.	7	D5469...	13 36 48 N.; 123 38 24 E...	500	fine s.....	Few.
			D5546...	6 06 48 N.; 121 20 32 E...	138	53.3	Co., s.....	
13129	U.S.N.M.	4	D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	br. s., Co.....	
			D5575...	5 28 30 N.; 120 02 27 E...	315	52.3	gy. m., fine s.....	
			D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	
14947	U.S.N.M.	1	D5636...	1 55 00 S.; 127 12 30 E...	1,262	Rare.
				Exterior pearl oyster, Jolo Jolo, P. I.	

Subfamily RAMULININAE.

Genus RAMULINA Rupert-Jones, 1875.

RAMULINA GLOBULIFERA H. B. Brady.

Ramulina globulifera H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 58, pl. 8, figs. 32, 33; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 587, pl. 76, figs. 22-28.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 321, pl. 68, fig. 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 3, 1913, p. 110, pl. 39, fig. 1.

Brady recorded this species from a single *Challenger* station off the Philippines in 95 fathoms (174 meters). At *Albatross* station D5259, off northwestern Panay, in 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.), this species was abundant. The only other record of its occurrence in the region is D5537, in 254 fathoms (465 meters), between Negros and Siquijor, bottom temperature 53.5° F. (11.8° C.).

Ramulina globulifera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12757	U.S.N.M.	10+	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob..	Common.
			D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.....	

Family CHILOSTOMELLIDAE.

Genus CHILOSTOMELLA Reuss, 1850.

CHILOSTOMELLA OVOIDEA Reuss.

Chilostomella ovoidea REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1850, p. 380, pl. 48, fig. 12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 436, pl. 55, figs. 12-23.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 2, pl. 1, figs. 1-5.

Although never noted in any considerable numbers, this species occurred at numerous stations, mostly within the Philippine Archipelago, as follows: China Sea, off southern Luzon; Tanon Strait, east coast of Negros; Sogod Bay, southern Leyte; east of Masbate Island; between Burias and Luzon; Palawan Passage; Dupon Bay, Leyte; between Cebu and Bohol; and Macassar Strait.

Bottom temperatures where given range from 44° F. to 59° F. (6.6° C. to 15° C.), with the average of 52.9° F. (11.6° C.). Depths range from 127 to 1,008 fathoms (233 to 1,844 meters), but two stations being over 375 fathoms (686 meters); one of 554 fathoms (1,012 meters), the other 1,008 fathoms (1,844 meters).

The specimens were of the typical form, ovoid in side view and the test pellucid.

Length up to 1 mm.

Chilostomella ovoidea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13380	U.S.N.M.	1	D5112...	13 48 22 N.; 120 47 25 E...	177	52.4	dk. gn. m...	Rare.
13374	U.S.N.M.	1	D5113...	13 51 30 N.; 120 50 30 E...	159	52.8	dk. gn. m...	Rare.
13377	U.S.N.M.	3	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s. m...	Few.
14367	U.S.N.M.	1	D5211...	11 51 35 N.; 124 14 00 E...	155	56.6	gn. m., s....	Few.
12840								
12845	U.S.N.M.	5	D5211...	11 51 35 N.; 124 14 00 E...	155	56.6	gn. m., s....	Few.
13375								
13375	U.S.N.M.	1	D5216...	12 52 00 N.; 123 23 30 E...	215	51.9	gn. m.....	Few.
12841	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 15 E...	195	52.8	gn. m.....	Rare.
14368	U.S.N.M.	3	D5239...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
13379	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.....	Few.
14370	U.S.N.M.	1	D5260...	12 25 35 N.; 121 31 35 E...	234	51.4	gn. m., s....	Rare.
13376	U.S.N.M.	1	D5404...	10 50 00 N.; 124 26 18 E...	190	55.4	m.....	Rare.
13378	U.S.N.M.	1	D5420...	9 49 35 N.; 123 45 00 E...	127	59.0	Rare.
12842	U.S.N.M.	1	D5546...	6 06 48 N.; 121 20 32 E...	138	58.3	fne. co. s....	Rare.
14365	U.S.N.M.	2	D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.....	Rare.
12843	U.S.N.M.	1	D5665...	4 27 00 S.; 118 44 00 E...	1,008	m.....	Rare.
14369	U.S.N.M.	2	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fne. s., glob.	Rare.
14366	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Rare.

CHILOSTOMELLA GRANDIS Cushman.

Plate 57, figs. 5a-c.

Chilostomella grandis CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 662.

Description.—Test broadly elliptical in side view; in end view circular; two chambers visible from the exterior; wall thick and opaque,

smooth; aperture in end view semicircular with a flange-like truncated lip.

Length up to 4 mm. and more.

Distribution.—Type-specimens (U.S.N.M. No. 9146) from *Albatross* station D5449, in 300 fathoms (549 meters), east coast of Luzon between San Bernardino Strait and San Miguel Bay. It also occurred at D5469, in 500 fathoms (914 meters), in the same region; D5300, in 265 fathoms (485 meters), China Sea, vicinity of southern Luzon; and D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.).

This species is nearly four times the size of typical *Chilostomella ovoidea*, and differs in the broadly elliptical outline of the side view, the very thick, opaque test, and in the appearance of the lip, which in direct side view is practically a straight line, while in typical *C. ovoidea* it is usually a depressed curve.

Chilostomella grandis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12844	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Common.
12846	U.S.N.M.	2	D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s....	Rare.
9146	U.S.N.M.	1	D5449...	13 21 36 N.; 124 00 30 E...	300	Rare.
14371	U.S.N.M.	3						
12847	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Rare.

Genus ALLOMORPHINA Reuss, 1850.

ALLOMORPHINA TRIGONA Reuss.

Allomorphina trigona REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1850, p. 380, pl. 48, figs. 14a-e.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 438, pl. 55, figs. 24-26.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 3, pl. 1, figs. 6-8.

Typical material of this species has occurred at the following six stations: D5112, in 177 fathoms (324 meters), China Sea off southern Luzon, bottom temperature 52.4° F. (11.3° C.); D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte, bottom temperature 52.8° F. (11.5° C.); D5467, in 480 fathoms (878 meters), east coast of Luzon, between San Bernardino Strait and San Miguel Bay; D5529, in 441 fathoms (812 meters), between Siquijor and Bohol Islands, bottom temperature 53° F. (11.7° C.); D5625, in 230 fathoms (420 meters), between Gillolo and Kayoa Islands; and D5654, in 805 fathoms (1,472 meters), Gulf of Boni.

Scattered specimens seem to be the rule as far as material from these stations show.

Allomorphina trigona—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12836	U.S.N.M.	1	D5112...	13 48 22 N.; 120 47 25 E..	177	52.4	dk. gn. m...	Rare.
12837	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m...	Rare.
12838	U.S.N.M.	1	D5467...	13 35 27 N.; 123 37 18 E..	480	gy. m.	Rare.
12835	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob.	Rare.
12839	U.S.N.M.	1	D5625...	0 07 09 N.; 127 29 00 E..	230	gy. m., fnc. s.	Rare.
14372	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E..	505	Rare.

Family GLOBIGERINIDAE.

Genus GLOBIGERINA d'Orbigny, 1826.

GLOBIGERINA BULLOIDES d'Orbigny.

Globigerina bulloides D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 277, No. 1; Modèles, 1826, No. 17 and No. 76.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 593, pl. 77; pl. 79, figs. 3-7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 5, pl. 2, figs. 7-9; pl. 9.

This species was found at 34 stations in the area, some of them in the protected waters of the Archipelago. It was frequent in the China Sea and occasional in the Sulu Sea; Gulf of Tomini, Celebes; between Gillolo and Makyan Islands, and between Gillolo and Kayao Islands; and rare at a few points in protected waters of the Archipelago, being noted as frequent at one station off Masbate. These stations range in depth from 10 to 765 fathoms (18 to 1,399 meters), the average depth being 247 fathoms (452 meters), and the bottom temperature ranges from 40.6° F. to 63.6° F (4.7° C. to 17.5° C.), with the average 52.7° F. (11.4° C.).

The distribution of this and various other species of the family seems to be due very largely to ocean currents. The current from the east flows against the eastern side of the islands, then turns and is swept back again. The currents undoubtedly carry some species into the waters of the Archipelago, but in comparatively few numbers, and the only stations at which they are found in any quantity are on the eastern coasts of Luzon and Mindanao. The southern current sweeping into the China Sea also carries certain species into that region also.

Globigerina bulloides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
			D5097...	14 19 15 N.	120 33 52 E.	30	gy. m., s., sh.	Few.
			D5106...	14 23 55 N.	120 32 33 E.	37	gy. m.	Few.
			D5109...	14 03 45 N.	120 16 30 E.	10	co.	Few.
			D5112...	13 48 22 N.	120 47 25 E.	177	52.4	dk. gn. m.	Few.
			D5113...	13 51 30 N.	120 50 30 E.	159	dk. gn. m.	Few.
			D5120...	13 45 30 N.	120 30 15 E.	393	43.7	gn. m., s.	Frequent.
			D5126...	10 34 15 N.	121 47 30 E.	742	49.5	sft. gn. m.	Few.
			D5146...	5 46 40 N.	120 48 50 E.	24	co. s., sh.	Few.
			D5192...	11 09 15 N.	123 50 00 E.	32	gn. m., s.	Few.
12524	U.S.N.M.	5	D5211...	11 51 35 N.	124 14 00 E.	155	56.6	gn. m., s.	Few.
12494	U.S.N.M.	8	D5213...	12 15 00 N.	123 57 30 E.	80	s., m., sh.	Common.
			D5214...	12 25 18 N.	123 37 15 E.	218	51.4	gn. m.	Few.
			D5216...	12 52 00 N.	123 23 30 E.	215	50.2	gn. m.	Few.
			D5217...	13 20 00 N.	123 14 15 E.	105	63.1	crs. gy. s.	Frequent.
			D5222...	13 38 30 N.	121 42 45 E.	195	52.8	gn. m.	Few.
			D5243...	6 50 55 N.	126 14 35 E.	218	63.6	gy. m.	Few.
			D5268...	13 42 00 N.	120 57 15 E.	170	s., p.	Few.
			D5349...	10 54 00 N.	118 26 20 E.	730	40.6	co. s.	Few.
			D5381...	13 14 15 N.	122 44 25 E.	88	co. s.	Few.
12522	U.S.N.M.	6	D5443...	12 43 05 N.	125 01 00 E.	241	51.3	co. s., sh.	Few.
			D5446...	12 43 51 N.	124 59 18 E.	300	gn. m.	Few.
			D5469...	13 36 48 N.	123 38 24 E.	500	gn. m.	Few.
			D5512...	8 16 02 N.	123 58 26 E.	445	52.8	gy. m., fine s.	Few.
			D5542...	8 48 30 N.	123 35 30 E.	200	54.3	fine s., brk. sh.	Few.
			D5543...	8 47 15 N.	123 35 00 E.	162	54.5	s.	Few.
			D5546...	6 06 48 N.	121 20 32 E.	138	58.3	fine. co. s.	Few.
			D5549...	6 01 15 N.	120 44 20 E.	263	52.3	s., glob., for.	Few.
			D5551...	5 54 48 N.	120 44 24 E.	192	53.3	fine. s.	Few.
			D5576...	5 25 56 N.	120 03 39 E.	277	53.3	s.	Few.
			D5601...	1 13 10 N.	125 17 05 E.	765	s., glob., ptr.	Few.
			D5621...	0 15 00 N.	127 24 35 E.	298	gy. bk. s.	Few.
			D5623...	0 16 30 N.	127 30 00 E.	272	fine. s., m.	Few.
			D5625...	0 07 00 N.	127 28 00 E.	230	gy. m., fine s.	Few.
			D5626...	0 07 30 N.	127 29 00 E.	265	gy. m., fine s.	Few.

GLOBIGERINA BULLOIDES d'Orbigny, var. TRILOBA Reuss.

Globigerina triloba REUSS, Denkschr. Acad. Wiss. Wien, vol. 1, 1850, p. 374, pl. 47, fig. 11.

Globigerina bulloides D'ORBIGNY, var. *triloba* H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 595, pl. 79, figs. 1, 2; pl. 81, figs. 2, 3.

Description.—Variety, with the three last-formed chambers making up the entire visible test.

Distribution.—This variety seems to be very rare in the region, occurring at Albatross station D5445, in 383 fathoms (699 meters) east coast of Luzon, bottom temperature 44.3° F. (6.8° C.); D5636, in 1,262 fathoms (2,309 meters), Pitt Passage; and D5638, the neighboring station, 517 fathoms (946 meters).

Globigerina bulloides, var. *triloba*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
			D5445...	12 44 42 N.	124 59 50 E.	383	44.3	gn. m., s.	Rare.
12525	U.S.N.M.	D5636...	10 55 00 S.	127 42 30 E.	1,262	gy. m., fine s.	
12526	U.S.N.M.	1	D5638...	3 47 15 S.	126 23 40 E.	517	fine. gy. s.	Rare.

GLOBIGERINA CRETACEA d'Orbigny.

Globigerina cretacea D'ORBIGNY, Mém. Soc. Géol. France, ser. 1, vol. 4, 1840, p. 34, pl. 3, figs. 12-14.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 596, pl. 82, figs. 10a-c (?).—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 7.

A form, which it does not seem clear whether to place under this species or under *G. dubia* Egger, is not uncommon in the outside waters: China Sea; Flores Sea; Macassar Strait; Gulf of Tomini, Celebes; off Bouro Island; off Gillolo, etc.; and at numerous stations in the protected waters of the Archipelago. These range in depth from 24 to 1,570 fathoms (44 to 2,871 meters), with the average depth 320 fathoms (586 meters); the bottom temperatures ranging from 39.2° to 63.6° F. (4.0° C. to 17.5° C.), with the average 51.2° F. (10.6° C.).

Globigerina cretacea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
			D5096...	14 20 23 N.; 120 34 15 E..	28	gy. m., s., sh.	Few.
			D5097...	14 19 15 N.; 120 33 52 E..	30	gy. m., s., sh.	Rare.
			D5106...	14 23 55 N.; 120 32 33 E..	37	gy. m.	Few.
			D5112...	13 48 22 N.; 120 47 25 E..	177	dk. gn. m.	Common.
			D5113...	13 51 30 N.; 120 50 30 E..	159	dk. gn. m.	Frequent.
			D5114...	13 36 11 N.; 120 45 26 E..	340	fne. s.	Frequent.
			D5115...	13 37 11 N.; 120 43 40 E..	340	Few.
			D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s.	Few.
			D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m.	Few.
			D5146...	6 04 30 N.; 120 59 30 E..	24	co. s., sh.	Few.
			D5152...	5 22 55 N.; 120 15 45 E..	34	wh. s.	Common.
			D5153...	5 18 10 N.; 120 02 55 E..	49	co. s., sh.	Common.
14379	U.S.N.M.	10	D5172...	6 03 15 N.; 120 35 30 E..	318	fne. s., sh.	Frequent.
			D5178...	12 43 00 N.; 122 06 15 E..	78	fne. s.	Few.
			D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	gn. m.	Few.
			D5187...	9 16 45 N.; 123 21 15 E..	225	53.6	sft. gn. m.	Few.
			D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.	Frequent.
			D5203...	9 58 00 N.; 125 07 40 E..	775	52.9	gn. m.	Frequent.
			D5206...	11 31 40 N.; 124 42 40 E..	32	gn. m.	Frequent.
			D5212...	12 04 15 N.; 124 04 36 E..	108	59.9	gy. s., m.	Few.
			D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s.	Few.
			D5219...	13 21 00 N.; 122 18 45 E..	530	50.8	gn. m.	Few.
14378	U.S.N.M.	10	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.	Frequent.
			D5243...	6 50 55 N.; 126 14 35 E..	218	63.6	gy. m.	Few.
14377	U.S.N.M.	8	D5247...	7 02 00 N.; 125 38 45 E..	135	m. s.	Rare.
			D5249...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Frequent.
			D5290...	12 25 35 N.; 121 31 35 E..	234	51.4	gn. m., s.	Few.
			D5291...	12 30 55 N.; 121 34 24 E..	145	s. m.	Few.
			D5298...	13 42 00 N.; 120 57 15 E..	170	s. p.	Frequent.
			D5292...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. gy. s.	Few.
			D5284...	13 42 05 N.; 120 30 45 E..	422	42.3	gy. m., glob.	Few.
			D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s.	Few.
			D5311...	21 35 00 N.; 116 15 00 E..	88	crs. s., sh.	Few.
			D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s.	Frequent.
			D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.	Few.
			D5349...	10 54 00 N.; 118 26 20 E..	730	40.6	co. s.	Few.
			D5364...	13 48 30 N.; 120 43 45 E..	106	bk. s.	Rare.
			D5377...	13 26 00 N.; 122 19 00 E..	400	49.6	sft. gn. m.	Few.
			D5381...	13 14 15 N.; 122 44 25 E..	88	co. s.	Few.
14376	U.S.N.M.	10	D5430...	9 49 40 N.; 119 03 20 E..	464	50.0	glob. oz.	Frequent.
			D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s.	Few.
			D5481...	10 27 30 N.; 125 17 10 E..	61	s., sh., g.	Few.
			D5512...	8 16 02 N.; 123 58 26 E..	445	52.8	gy. m., fne. s.	Few.
14373	U.S.N.M.	6	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn. m., glob.	Few.
			D5542...	8 48 30 N.; 120 35 30 E..	200	54.3	fne. s., brk. sh.	Few.
			D5543...	8 47 15 N.; 123 35 00 E..	162	54.5	s.	Few.
			D5548...	6 00 20 N.; 120 45 35 E..	232	53.5	s., brk. sh.	Rare.
			D5549...	6 01 15 N.; 120 44 20 E..	263	52.3	s., glob., for.	Few.
			D5551...	5 54 48 N.; 120 44 24 E..	193	53.3	fne. s.	Few.
			D5565...	5 51 42 N.; 120 30 30 E..	213	52.3	s., ptr. sh.	Few.

Globigerina cretacea—Material examined—Continued.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "				
			D5571...	5 30 45 N.; 120 07 57 E..	340	52.3	s., sh.....	Few.
			D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Rare.
			D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.....	Few.
			D5601...	1 13 10 N.; 125 17 05 E..	765	s., glob., ptr.....	Few.
			D5622...	0 19 20 N.; 127 28 30 E..	275	gy. m.....	Few.
			D5523...	0 16 30 N.; 127 30 00 E..	272	fine. s., m.....	Few.
			D5625...	0 07 00 N.; 127 28 00 E..	230	gy. m., fine. s.....	Frequent.
			D5626...	0 07 30 N.; 127 29 00 E..	265	gy. m., fine. s.....	Few.
			D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	Few.
14374	U.S.N.M.	4	D5638...	3 47 15 S.; 126 23 40 E..	517	fine. gy. s.....	Frequent.
			D5650...	4 53 45 S.; 121 29 00 E..	510	40.1	gn. m.....	Few.
			D5660...	5 36 30 S.; 120 49 00 E..	692	39.2	gy. m., s.....	Few.
			D5665...	4 27 00 S.; 118 44 00 E..	1,003	m.....	Few.
14375	U.S.N.M.	3	H4897...	7 46 00 N.; 122 00 00 E..	1,570	gy. m.....	Common.

GLOBIGERINA DIGITATA H. B. Brady.

Globigerina digitata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 72; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 599, pl. 80, figs. 6-10; pl. 82, figs. 6, 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 7, pl. 14, figs. 1-3.

Specimens of this interesting species have been found in few numbers from 14 *Albatross* stations in the region. These are in the Sulu Sea; between Panay and Negros; Tanon Strait, east coast of Negros; Gulf of Davao; off northwestern Panay; China Sea, off southern Luzon; Palawan Passage; off Jolo Island; and Flores Sea.

These stations range in depth from 135 to 885 fathoms (247 to 1,618 meters), the average depth being 495 fathoms (905 meters); bottom temperatures range from 38.2° to 62.8° F. (3.4° to 17.1° C.), average 47.5° F. (8.6° C.).

In Palawan Passage specimens at one station were common, but usually they were represented at any one station by very few specimens. At one station in Flores Sea specimens were found with the chambers branched at the tips.

Globigerina digitata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "				
12515	U.S.N.M.	1	D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m....	Rare.
			D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	gn. m.....	Few.
14382	U.S.N.M.	1	D5191...	10 29 45 N.; 123 31 15 E..	258	62.8	gn. m.....	Few.
12514	U.S.N.M.	1	D5247...	07 02 00 N.; 125 38 45 E..	135	m.....	Rare.
14380	U.S.N.M.	4	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Few.
12513	U.S.N.M.	2	D5284...	13 42 05 N.; 120 30 45 E..	422	42.3	gy. m., glob.	Few.
12507 12512	U.S.N.M.	2	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s.....	Few.
12508	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Few.
12509	U.S.N.M.	10	D5349...	10 54 00 N.; 118 26 20 E..	730	40.6	co. s.....	Common.
12511	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob.	Rare.
			D5549...	6 01 15 N.; 120 44 20 E..	263	52.3	s., glob., for.....	Rare.
14381	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Rare.
12510	U.S.N.M.	4	D5660...	5 36 30 S.; 120 49 00 E..	692	39.2	gy. m., s.....	Few.
14903	U.S.N.M.	2	H4937...	5 32 50 N.; 120 49 10 E..	885	38.2	gy. m.....	Few.

GLOBIGERINA INFLATA d'Orbigny.

Globigerina inflata D'ORBIGNY, in Baker, Webb and Berthelot, Hist. Nat. Îles Canaries, vol. 2, pt. 2, 1839, "Foraminifères," p. 134, pl. 2, figs. 7-9.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 601, pl. 79, figs. 8-10.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 8, pl. 4, figs. 4-8.

At eight stations specimens of this species occurred as follows: *Albatross* D5126, Sulu Sea, southern Panay, 742 fathoms (1,357 meters) bottom temperature 49.5° F. (9.8° C.); D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte, bottom temperature 52.8° F. (11.5° C.); D5152, in 34 fathoms (62 meters), Sulu Archipelago, Tawi Tawi Group; D5178, in 78 fathoms (143 meters), off Romblon; D5211, east of Masbate Island, 155 fathoms (283 meters), bottom temperature 56.6° F. (13.6° C.); D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.); and D5369, in 106 fathoms (194 meters), off Marinduque Island, and D5618, Molucca Passage, 417 fathoms (763 meters).

Globigerina inflata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
12531	U.S.N.M.	1	D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m....	Rare.
12529	U.S.N.M.	8	D5152...	5 22 55 N.; 120 15 45 E..	34	wh. s.....	Frequent.
12530	U.S.N.M.	2	D5178...	12 43 00 N.; 122 06 15 E..	78	fine s.....	Rare.
12527	U.S.N.M.	5	D5211...	11 51 35 N.; 124 14 00 E..	155	56.6	gn. m., s.....	Few.
			D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	
			D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s....	
			D5369...	13 48 00 N.; 121 43 00 E..	106	bk. s.....	
12528	U.S.N.M.	7	D5618...	00 37 00 N.; 127 15 00 E..	417	gy. m.....	Frequent.

GLOBIGERINA RUBRA d'Orbigny.

Globigerina rubra D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 94, pl. 4, figs. 12-14.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 602, pl. 79, figs. 11-16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 9, pl. 3, figs. 6-9.

Specimens of this species occurred at several stations in the area: Off northern Cebu; east of Masbate; off southeastern Mindoro; Verde Island Passage; Palawan Passage; Ragay Gulf; east coast of Luzon; and off Jolo Island.

These stations vary in depth from 32 to 730 fathoms (59 to 1,335 meters) with the average depth 237 fathoms (434 meters). Bottom temperatures from 40.6° F. to 56.6° F. (4.7° C. to 13.6° C.).

Few specimens showed the reddish color, but had the characteristic form of the species.

Globigerina rubra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12517	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.	Few.
			D5211...	11 51 35 N.; 124 14 00 E..	155	56.6	gn. m., s.	Few.
12520	U.S.N.M.	1	D5261...	12 30 55 N.; 121 34 24 E..	145	s., m.	Rare.
12516	U.S.N.M.	1	D5267...	13 42 20 N.; 120 58 25 E..	170	p., s., sh.	Rare.
			D5349...	10 54 00 N.; 118 26 20 E..	730	40.6	co. s.,	Rare.
12518	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 25 E..	88	co. s.,	Few.
12519	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s.	Rare.
12521	U.S.N.M.	1	D5551...	5 54 48 N.; 120 44 24 E..	193	53.3	inc. s.,	Rare.

GLOBIGERINA CONGLOBATA H. B. Brady.

Globigerina conglobata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 72; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 603, pl. 80, figs. 1-5; pl. 82, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 10, pl. 3, figs. 3-5; pl. 10, figs. 1, 6.

At 58 stations record was made of this species in my examination of the material. These are very largely in the China Sea; the east coasts of Luzon and Mindanao; off Jolo and Tawi Tawi; Darvel Bay, Borneo; Gulf of Tomimi, Celebes; between Gillolo and Makayan; off Bouro Island; Molucca Sea; Gulf of Boni; Flores Sea; with scattered stations in the protected waters of the Archipelago.

The stations range in depth from 18 to 1,570 fathoms (33 to 2,871 meters), the average depth being 324 fathoms (593 meters); the average bottom temperature, 50.5° F. (10.2° C.).

Globigerina conglobata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12500	U.S.N.M.	5	D5100...	14 17 15 N.; 120 32 50 E..	35	gy. s.	Few.
			D5110...	13 59 20 N.; 120 75 45 E..	135	59	dk. gy. m.	Few.
			D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s.	Few.
			D5121...	13 27 20 N.; 121 17 45 E..	108	dk. gn. m.	Few.
			D5152...	5 22 55 N.; 120 15 45 E..	24	co. s., sh.	Frequent.
			D5153...	5 18 10 N.; 120 02 55 E..	49	co. s., sh.	Few.
			D5156...	5 12 50 N.; 119 55 55 E..	18	fine. s., sh.	Rare.
			D5172...	6 03 15 N.; 120 35 30 E..	318	fine. s., sh.	Few.
			D5178...	12 43 00 N.; 122 06 15 E..	78	fine. s.,	Few.
			D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	gn. m.	Few.
			D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.,	Few.
			D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.	Frequent.
			D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s.	Few.
12498	U.S.N.M.	10+	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.	Frequent.
			D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.	Common.
			D5247...	7 02 00 N.; 125 38 45 E..	135	m.	Few.
			D5255...	7 03 00 N.; 125 39 00 E..	100	sft.	Few.
			D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Few.
			D5267...	13 42 20 N.; 120 58 25 E..	170	p., s., sh.	Few.
			D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.	Few.

Globigerina conglobata—Material examined—Continued.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12501	U.S.N.M.	4	D5300	20 31 00 N., 115 49 00 E..	265	gy. m., s.....	Few.
			D5301	20 37 00 N., 115 43 00 E..	208	50.5	gy. m., s.....	Rare.
			D5311	21 33 00 N., 116 15 00 E..	88	crs. s., sh.....	Rare.
			D5313	21 30 00 N., 116 43 00 E..	150	53.6	Few.
			D5315	21 40 00 N., 116 58 00 E..	148	54.4	s., sh.....	Few.
			D5338	11 33 45 N., 119 24 45 E..	43	co. s., m.....	Few.
			D5342	10 56 55 N., 119 17 24 E..	145	gy. m.....	Rare.
			D5348	10 57 45 N., 118 38 15 E..	375	56.4	co. s.....	Few.
			D5349	10 54 00 N., 118 26 20 E..	730	40.6	co. s.....	Few.
			D5370	13 44 15 N., 121 42 30 E..	159	54.3	sft. m.....	Few.
			D5381	13 14 15 N., 122 44 25 E..	88	co. s.....	Few.
			D5443	12 43 05 N., 125 01 00 E..	241	51.3	co. s., sh.....	Rare.
			D5445	12 44 42 N., 124 59 50 E..	381.	44.3	gn. m., s.....	Few.
			D5446	12 43 51 N., 124 59 18 E..	300	gn. m.....	Few.
12499	U.S.N.M.	2	D5449	13 21 36 N., 124 00 30 E..	300	Few.
			D5459	13 10 21 N., 123 59 54 E..	201	Rare.
			D5467	13 35 27 N., 123 37 18 E..	480	gy. m.....	Few.
			D5470	13 37 30 N., 123 41 09 E..	560	m.....	Rare.
			D5481	10 27 30 N., 125 17 10 E..	61	s., sh., g.....	Few.
			D5512	8 16 02 N., 123 58 26 E..	445	52.8	gy. m., fne. s.	Few.
			D5529	9 23 45 N., 123 39 30 E..	441	53	gy. m., glob.	Few.
			D5538	9 08 15 N., 123 23 20 E..	256	53.3	gn. m., s.....	Few.
			D5542	8 48 30 N., 123 35 30 E..	200	54.3	fne. s., brk. sh.	Few.
			D5543	8 47 15 N., 123 35 00 E..	162	54.5	s.....	Few.
			D5548	6 00 20 N., 120 45 35 E..	232	53.5	s., brk. sh.....	Few.
			D5548	6 01 15 N., 120 44 20 E..	263	52.3	s., glob., for.	Few.
			D5571	5 30 45 N., 120 07 57 E..	340	52.3	s., sh.....	Few.
			D5580	4 52 45 N., 119 06 45 E..	162	55.8	br. s., co.....	Rare.
			D5612	0 38 00 S., 121 45 40 E..	750	Few.
			D5621	0 15 00 N., 127 24 45 E..	298	gy., bk. s.....	Few.
			D5627	0 06 00 N., 127 26 00 E..	22	m.....	Few.
D5630	0 56 30 S., 128 05 00 E..	569	co. s., m.....	Rare.			
D5637	3 53 20 S., 126 48 00 E..	700	gy. m.....	Few.			
D5639	3 54 50 S., 123 27 20 E..	1,560	gy. m.....	Few.			
D5650	4 53 45 S., 121 29 00 E..	540	40.1	gn. m.....	Few.			
D5660	5 36 39 S., 120 49 30 E..	692	39.2	gy. m., s.....	Few.			
H4897	7 46 00 N., 122 00 00 E..	1,570	gy. m., glob.	Few.			
H4937	5 32 50 N., 120 49 10 E..	885	38.2	gy. m.....	Few.			
14908	U.S.N.M.	1	Nero 849.	15 16 00 N., 121 40 00 E..	737	gn. m.....	Rare.
9950	U.S.N.M.	1						

GLOBIGERINA SACCULIFERA H. B. Brady.

Globigerina helicina CARPENTER (not *G. helicina* d'Orbigny), Introd. Foram., 1862, pl. 12, fig. 11.

Globigerina sacculifera H. B. BRADY, Geol. Mag., Dec. 2, vol. 4, 1877, p. 535; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 604, pl. 80, figs. 11–17; pl. 82, fig. 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 11, pl. 2, figs. 4–6; pl. 5, pl. 10, fig. 4.

I have noted the occurrence of this species in material from nearly a hundred stations in the area. These stations cover practically the entire area which was worked over by the *Albatross* in this area. They range in depth from 20 to 1,570 fathoms (37 to 2,871 meters); average depth, 286 fathoms (523 meters). The average bottom temperature is 53.3° F. (11.8° C.).

As a rule specimens were most common in the areas already mentioned for other specimens.

Globigerina sacculifera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
12502	U.S.N.M.	10+	D5096...	14 20 23 N.	120 34 15 E.	28	° F.	gy. m., s., sh.	Few.
			D5097...	14 19 15 N.	120 33 52 E.	30	gy. m., s., sh.	Rare.
			D5100...	14 17 15 N.	120 32 40 E.	35	gy. s.	Common.
			D5106...	14 23 55 N.	120 32 33 E.	37	gy. m.	Few.
12504	U.S.N.M.	10	D5110...	13 59 20 N.	120 75 45 E.	135	59.0	dk. gy. m.	Common.
			D5112...	13 48 22 N.	120 47 25 E.	177	52.4	dk. gy. m.	Frequent.
			D5113...	13 51 30 N.	120 50 30 E.	159	dk. gn. m.	Few.
			D5114...	13 36 11 N.	120 45 26 E.	340	fine. s.	Frequent.
			D5115...	13 37 11 N.	120 43 40 E.	340	Frequent.
			D5120...	13 45 30 N.	120 30 15 E.	393	43.7	gn. m., s.	Few.
			D5121...	13 27 20 N.	121 17 45 E.	108	dk. gn. m.
			D5126...	10 34 45 N.	121 47 30 E.	742	49.5	sft. gn. m.	Few.
			D5146...	5 46 40 N.	120 48 50 E.	24	co. s., sh.	Few.
			D5152...	5 22 55 N.	120 15 45 E.	24	co. s., sh.	Frequent.
			D5153...	5 18 10 N.	120 02 55 E.	49	co. s., sh.	Few.
			D5172...	6 03 15 N.	120 35 30 E.	318	fine. s., sh.	Common.
D5178...	12 43 00 N.	122 06 15 E.	78	fine. s.	Common.			
D5179...	12 38 15 N.	122 12 30 E.	37	75.7	hrd. s.	Few.			
D5185...	10 05 45 N.	122 18 30 E.	638	49.8	gn. m.	Frequent.			
D5189...	9 56 30 N.	123 15 00 E.	300	62.8	gn. m.	Few.			
12505	U.S.N.M.	10	D5191...	10 29 45 N.	123 31 15 E.	258	62.8	gn. m.	Common.
			D5192...	11 09 15 N.	123 50 00 E.	32	gn. s.	Few.
			D5201...	10 10 00 N.	125 04 15 E.	554	52.8	gy. s., m.	Frequent.
			D5206...	11 31 40 N.	124 42 40 E.	32	gn. m.	Few.
			D5211...	11 51 35 N.	124 14 00 E.	155	56.6	gn. m., s.	Frequent.
			D5212...	12 04 15 N.	124 04 36 E.	108	59.9	gy. s., m.	Few.
			D5215...	12 31 30 N.	123 35 24 E.	604	50.5	gn. m.	Few.
			D5216...	12 52 00 N.	123 23 30 E.	200	50.2	Frequent.
			D5217...	13 20 00 N.	123 14 15 E.	105	63.1	crs. gy. s.	Frequent.
			D5218...	13 11 15 N.	123 02 45 E.	20	crs. s.	Few.
			D5219...	13 21 00 N.	122 18 45 E.	530	50.8	gn. m.	Few.
			D5220...	13 38 00 N.	121 58 00 E.	50	sft. gn. m.	Frequent.
12503	U.S.N.M.	10	D5222...	13 38 30 N.	121 42 45 E.	195	52.8	gn. m.	Few.
			D5236...	8 50 45 N.	126 26 52 E.	494	41.2	fine. gy. s.	Few.
			D5241...	6 50 45 N.	126 14 38 E.	215	sft. gy. m.	Few.
			D5243...	6 50 55 N.	126 14 35 E.	218	63.6	gy. m.	Common.
			D5247...	7 02 00 N.	125 38 45 E.	135	m.	Few.
			D5257...	7 22 12 N.	124 12 15 E.	28	m.	Few.
			D5259...	11 57 30 N.	121 42 15 E.	312	49.3	gy. m., glob.	Common.
			D5260...	12 25 35 N.	121 31 35 E.	234	51.4	gn. m., s.	Few.
			D5261...	12 30 55 N.	121 34 24 E.	145	s., m.	Few.
			D5267...	13 42 20 N.	120 58 25 E.	170	p., s., sh.	Few.
			D5268...	13 42 00 N.	120 57 15 E.	170	s. p.	Common.
			D5277...	13 56 55 N.	120 13 45 E.	80	58.6	fine. s.	Few.
D5281...	13 52 45 N.	120 25 00 E.	201	50.4	dk. gy. s.	Few.			
D5282...	13 53 00 N.	120 26 45 E.	248	47.4	dk. gy. s.	Frequent.			
D5291...	13 29 40 N.	121 00 45 E.	173	51.5	fine. bk. s.	Frequent.			
D5300...	20 31 00 N.	115 49 00 E.	265	gy. m., s.	Few.			
D5313...	21 30 00 N.	116 43 00 E.	150	53.6	s.	Few.			
D5315...	21 40 00 N.	116 58 00 E.	148	54.6	s., sh.	Few.			
D5338...	11 33 45 N.	119 26 00 E.	43	co. s., m.	Few.			
D5342...	10 56 55 N.	119 17 24 E.	145	gy. m.	Rare.			
D5348...	10 57 45 N.	118 38 15 E.	375	56.4	co. s.	Common.			
D5349...	10 54 00 N.	118 26 20 E.	730	40.6	co. s.	Common.			
D5370...	13 44 15 N.	121 42 30 E.	159	54.3	sft. m.	Few.			
D5374...	13 46 45 N.	121 35 08 E.	190	gy. m.	Common.			
D5381...	13 14 15 N.	122 44 25 E.	88	co. s.	Few.			
D5394...	12 00 30 N.	124 05 36 E.	153	gn. m.	Few.			
D5423...	9 13 00 N.	118 51 15 E.	1,105	49.4	gy. m.	Few.			
D5430...	9 49 40 N.	119 03 20 E.	464	50.0	glob. oz.	Rare.			
D5449...	13 21 36 N.	124 00 30 E.	300	Few.			
D5467...	13 45 27 N.	123 37 18 E.	480	gy. m.	Few.			
D5481...	10 27 30 N.	125 17 10 E.	61	s., sh., g.	Few.			
D5512...	8 16 02 N.	123 58 26 E.	445	52.8	gy. m., fine. s.	Few.			
D5526...	9 12 45 N.	123 45 30 E.	805	52.3	gn. m., glob.	Frequent.			
D5529...	9 23 45 N.	123 39 30 E.	441	53.0	gy. m., glob.	Few.			
D5538...	9 08 15 N.	123 23 20 E.	256	53.3	gn. m., s.	Common.			
D5546...	6 06 48 N.	121 20 32 E.	138	58.3	ine. co. s.	Few.			
D5549...	6 01 15 N.	120 44 20 E.	263	52.3	s., glob., for.	Few.			
D5571...	5 30 45 N.	120 07 57 E.	340	52.3	s., sh.	Few.			
D5576...	5 25 56 N.	120 03 39 E.	277	53.3	s.	Few.			
D5579...	4 54 15 N.	109 09 52 E.	175	55.3	ine. s., Co.	Rare.			
D5601...	1 03 10 N.	125 17 05 E.	765	s. glob., ptr.	Few.			
D5618...	0 37 00 N.	127 15 00 E.	417	gy. m.	Few.			
D5625...	0 07 00 N.	127 28 00 E.	230	gy. m., fine. s.	Few.			
D5627...	0 06 00 N.	127 26 00 E.	22	m.	Few.			
D5630...	0 56 30 S.	128 05 00 E.	569	co. s., m.	Few.			
D5637...	3 53 20 S.	126 48 00 E.	700	gy. m.	Few.			
D5638...	3 47 15 S.	126 23 40 E.	517	ine. gy. s.	Few.			
D5660...	5 36 30 S.	120 49 00 E.	692	39.2	gy. m., s.	Few.			
H4897...	7 46 00 N.	122 00 00 E.	1,570	gy. m., glob.	Common.			
9971	U.S.N.M.	1	H4914...	13 38 05 N.	120 33 00 E.	464	46.5	gy. m., s.	Rare.
9956.	U.S.N.M.	1	Nero 849	15 16 00 N.	121 40 00 E.	737	gn. m.	Rare.

GLOBIGERINA AEQUILATERALIS H. B. Brady.

Cassidulina globulosa (part) EGGER, Neues Jahrb. für Min., 1857, p. 296, pl. 11, fig. 4.
Globigerina aequilateralis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 71;
 Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 605, pl. 80, figs. 18-21.—CUSHMAN,
 Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 12, pl. 2, figs. 1-3; pl. 10, fig. 5.

There seem to be two distinct forms of this species in the material which I have examined—one, which may be taken as typical, in which the development is planospiral, with the coiling so loose that it is possible to see between the coils in side view, that is, the coils are not really involute; the other, the variety described below. Such specimens are figured by Brady in the *Challenger* Report and may be taken as the typical form of the species.

This typical form was far less common than the following one, being found at but 15 stations in the area, but well scattered: China Sea; Sulu Sea; east coast of Luzon, etc., in the Pacific; Sibuko Bay, Borneo; between Gilolo and Makyan Islands; south of Patiente Strait; off Bouro Island, and at a very few stations in the protected waters of the Archipelago; off Batangas Bay, between Marinduque and Luzon; and east of Masbate. The depth of these stations ranges from 26 to 569 fathoms (48 to 1,040 meters), with the average depth 235 fathoms (430 meters). The bottom temperatures range from 44.3° F. to 59° F. (6.8° C. to 15° C.).




FIG. 11.—*GLOBIGERINA AEQUILATERALIS*, VAR. *INVOLUTA* CUSHMAN. X 50. a, SIDE VIEW; b, FROM AN OBLIQUE ANGLE; c, APERTURAL VIEW. FROM STATION D5255.

Globigerina aequilateralis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5113...	13 51 30 N.; 120 50 30 E...	159	° F.	dk. gn. m....	Few.
			D5132...	Island off Panabutan Point, N. 15 W. 0.3 mi.	26		gn. m., s....	Rare.
			D5211...	11 51 35 N.; 124 14 00 E...	155	56.6	gn. m., s....	Few.
			D5216...	12 52 00 N.; 123 23 30 E...	200	50.2	gn. m....	Few.
			D5222...	13 38 30 N.; 121 42 45 E...	195	52.8	gn. m....	Few.
			D5268...	13 42 00 N.; 120 57 15 E...	170		s., p.	Few.
12535	U.S.N.M.	1	D5291...	13 29 40 N.; 121 00 45 E...	173	51.5	fne. bk. s....	Rare.
			D5445...	12 44 42 N.; 124 59 50 E...	323	44.3	gn. m., s....	Rare.
12533	U.S.N.M.	2	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.	Few.
			D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Few.
			D5625...	0 07 00 N.; 127 28 00 E...	230		gy. m., fne s.	Few.
12532	U.S.N.M.	2	D5630...	0 56 30 S.; 128 05 00 E...	569		co. s., m.	Rare.
			D5638...	3 47 15 S.; 126 23 40 E...	517		fne. gy. s....	Few.
			D5696...	14 20 23 N.; 120 34 15 E...	28		gy. m., s., sh.	Few.
12534	U.S.N.M.	6	D5110...	13 59 20 N.; 120 75 45 E...	135	59.0	dk. gy. m....	Few.
9969	U.S.N.M.	2	H4914...	13 38 05 N.; 120 33 00 E...	464	46.5	gy. m., s....	Few.

GLOBIGERINA AEQUILATERALIS H. B. Brady, var. INVOLUTA Cushman.

Globigerina aequilateralis H. B. BRADY, var. *involuta* CUSHMAN, Proc. U. S. Nat Mus., vol. 51, 1917, p. 662.

Description.—Variety differing from the typical in being much more closely coiled, although planospiral, the last-formed coil decidedly overlapping the previous coils in side view; visible aperture instead of semicircular in the typical in the much broader; whole test less delicately formed.

Distribution.—Type specimen (U.S.N.M. No. 9147) from *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, off eastern coast of Mindanao, bottom temperature, 41.2° F. (5.1° C.). This variety has occurred at 56 other stations well distributed over the area from northern Luzon and the China Sea through the Archipelago, Sulu Sea to Flores Sea. The depths range from 22 to 1,570 fathoms (40 to 2,871 meters), with an average of 359 fathoms (656 meters), the average bottom temperature being 51.4° F. (10.8° C.).

This variety has been distinctive in this mass of material and it seems worthy of separation.

Globigerina aequilateralis, var. *involuta*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				• / " • / "		° F.		
			D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s.....	Few.
			D5106...	14 23 55 N.; 120 32 33 E...	37	gy. m.....	Rare.
			D5112...	13 48 22 N.; 120 47 25 E...	177	52.4	dk. gn. m.....	Frequent.
			D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m.....	Few.
			D5126...	10 34 45 N.; 121 47 30 E...	742	49.5	sft. gn. m.....	Frequent.
			D5146...	5 46 40 N.; 120 48 50 E...	24	co. s., sh.....	Few.
			D5153...	5 18 10 N.; 120 02 55 E...	49	co. s., sh.....	Few.
			D5172...	6 03 15 N.; 120 35 30 E...	318	fne. s., sh.....	Few.
			D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Few.
14384	U.S.N.M.	2	D5184...	10 18 30 N.; 122 23 30 E...	565	49.8	gn. m.....	Few.
14385	U.S.N.M.	8	D5187...	9 16 45 N.; 123 21 15 E...	638	49.8	gn. m.....	Few.
			D5189	} 9 56 30 N.; 123 15 00 E...	225	53.6	sft. gn. m.....	Frequent.
			D5191					
			D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Few.
			D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Frequent.
			D5203...	9 58 00 N.; 125 07 40 E...	775	52.9	gn. m.....	Few.
			D5213...	12 15 00 N.; 123 57 30 E...	80	s., m., sh.....	Few.
			D5214...	12 25 19 N.; 122 37 15 E...	218	51.4	gn. m.....	Few.
			D5215...	12 31 30 N.; 123 35 24 E...	604	50.5	gn. m.....	Few.
			D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.....	Few.
			D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m.....	Few.
9147	U.S.N.M.		D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s.....	Few.
			D5243...	6 50 55 N.; 126 14 35 E...	218	63.6	gy. m.....	Few.
14387	U.S.N.M.	3	D5255...	7 03 00 N.; 125 39 00 E...	100	sft. m.....	Few.
			D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.....	Few.
			D5260...	12 25 35 N.; 121 31 35 E...	234	51.4	g., m., s.....	Few.
			D5261...	12 30 55 N.; 121 34 24 E...	145	s., m.....	Few.
			D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s.....	Few.
			D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.....	Few.
			D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s.....	Few.
			D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Rare.
			D5338...	11 33 45 N.; 119 24 45 E...	43	co. s., m.....	Rare.
			D5342...	10 56 55 N.; 119 17 24 E...	145	gy. m.....	Rare.
			D5349...	10 54 00 N.; 118 26 20 E...	730	40.6	co. s.....	Few.
14386	U.S.N.M.	3	D5374...	13 46 45 N.; 121 35 08 E...	190	gy. m.....	Few.
			D5381...	13 14 15 N.; 122 44 25 E...	88	co. s.....	Few.
			D5430...	9 49 40 N.; 119 03 20 E...	464	50.0	glob. oz.....	Rare.
			D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.....	Rare.
			D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m.....	Few.
			D5523...	8 48 44 N.; 123 27 35 E...	Rare.
			D5526...	9 12 45 N.; 123 45 30 E...	805	52.3	gn. m., glob.....	Few.
			D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gy. m., glob.....	Few.
			D5538...	9 03 15 N.; 123 23 20 E...	256	53.3	gn. m., s.....	Few.

Globigerina aequilateralis, var. *involuta*—Material examined—Continued.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				* ' " * ' "				
			D5546 ..	6 06 48 N.; 121 20 32 E...	138	58.3	fne.co.s....	Few.
			D5565...	5 51 42 N.; 120 30 30 E...	243	52.3	s., ptr., sh...	Rare.
			D5571...	5 30 45 N.; 120 07 57 E...	340	52.3	s., sh.....	Few.
			D5601...	1 13 10 N.; 125 17 05 E...	765	s., glob., ptr.	Few.
			D5612...	0 38 00 S.; 121 45 40 E...	750	Few.
			D5622...	0 19 20 N.; 127 28 30 E...	275	gy.m.....	Rare.
			D5626...	0 07 30 N.; 127 29 00 E...	265	gy.m., fne.s.	Few.
			D5627...	0 06 00 N.; 127 26 00 E...	22	m.....	Few.
			D5632...	1 00 00 S.; 127 50 00 E...	845	Few.
			D5637...	3 53 20 S.; 126 48 00 E...	700	gy.m.....	Few.
			D5638...	3 47 15 S.; 126 23 40 E...	517	fne.gy.s....	Few.
			D5660...	5 36 30 S.; 120 49 00 E...	692	39.2	gy. m., s....	Few.
			D5665...	4 27 00 S.; 118 44 00 E...	1,008	m.....	Few.
			H4897...	7 46 00 N.; 122 00 00 E...	1,570	gy. m., glob.	Frequent.
			H4937...	5 32 50 S.; 120 49 10 E...	885	38.2	gy.m.....	Rare.

GLOBIGERINA CONGLOMERATA Schwager.

Globigerina conglomerata SCHWAGER, *Novara-Exped.*, Geol. Theil., pt. 2, 1866, p. 255, p. 7, fig. 113.

Description.—Test globose, final whorl of three or four chambers, usually the fourth overlapping the first, chambers very high, much inflated; spire somewhat raised, but not greatly so; umbilical region open widely; wall coarsely perforate.

Diameter up to 1 mm.

Distribution.—Specimens of this species were best developed at *Albatross* station D5613, in 752 fathoms (1,375 meters), Gulf of Tomini, Celebes.

This material in ventral view is very much like the original figure, and Brady remarks that his material obtained from Schwager showed "an approach toward the general form of *Globigerina conglobata*," which our material decidedly does, but in form only. At this station the species stood out very clearly from all others.

Globigerina conglomerata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14383	U.S.N.M.	4	D5613...	0 42 00 S.; 121 44 00 E...	752	* F.	gy.m.....	Few.

Genus ORBULINA d'Orbigny, 1839.

ORBULINA UNIVERSA d'Orbigny.

Orbulina universa D'ORBIGNY, in De la Sagra, *Hist. Fis. Pol. Nat. Cuba*, 1839, "Foraminifères," p. 3, pl. 1, fig. 1.—H. B. BRADY, *Rep. Voy. Challenger, Zoology*, vol. 9, 1884, p. 608, pl. 78; pl. 81, figs. 8-26; pl. 82, figs. 1-3.—CUSHMAN, *Bull.* 71, U. S. Nat. Mus., pt. 4, 1914, p. 14, pl. 6; pl. 7; pl. 11, fig. 3.

Strangely enough, *Orbulina universa*, which is one of the commonest of Foraminifera, does not seem to be at all well represented

in the Philippine material I have had. To be sure, typical Globigerina-ooze deposits are not the common bottom in the region, but *Globigerina* is not nearly so restricted in the area. It has occurred at several stations—in the Flores Sea; Sulu Sea; Sogod Bay, southern Leyte; Pacific Ocean, east coast of Mindanao; off southeastern Mindoro; between Marinduque and Luzon; and east of Masbate Island. The stations range in depth from 20 to 885 fathoms (37 to 1,618 meters). The average depth is 376 fathoms (689 meters). The bottom temperatures where given range from 38.2° to 63.1° F. (3.4° C. to 17.2° C.), the average being 51.2° F. (10.7° C.).

Orbulina universa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12496	U.S.N.M.	1	D5126...	10 34 45 N.; 121 47 30 E..	742	49.5	sft. gn. m.	Rare.
12497	U.S.N.M.		D5172...	6 03 15 N.; 120 35 30 E..	318	fne. s., sh.	Few.
12693	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.	Rare.
			D5203...	9 58 00 N.; 125 07 40 E..	775	52.9	gn. m.	Rare.
		2	D5211...	11 51 35 N.; 124 14 00 E..	155	56.6	gn. m., s.	Rare.
		2	D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s.	Few.
			D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.	Frequent.
12671	U.S.N.M.	10+	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.	Abundant.
			D5222...	13 38 30 N.; 121 42 45 E..	195	gn. m.	Few.
12692	U.S.N.M.	6	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s.	Few.
			D5260...	12 25 35 N.; 121 31 35 E..	234	51.4	gn. m., s.	Rare.
14904	U.S.N.M.	2	H4937...	5 32 50 S.; 120 49 10 E..	885	38.2	gy. m., s.	Rare.

Genus CANDEINA d'Orbigny, 1839.

CANDEINA NITIDA d'Orbigny.

Candeina nitida D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 108, pl. 2, figs. 27, 28.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 622, pl. 82, figs. 13–20.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 16, pl. 11, fig. 1.

A single station seems to be the only record for this species—*Albatross* D5638, in 517 fathoms (946 meters), off Bouro Island.

Candeina nitida—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12763	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E..	517	fne. gy. s.	Rare.

CANDEINA NITIDA d'Orbigny, var. TRILOBA, new variety.

Plate 57, figs. 1a–c.

Description.—Variety differing from the typical in that the last three chambers compose the whole external surface of the test, the spire being enveloped.

Distribution.—Type specimen (U.S.N.M. No. 14388) and others from *Albatross* station D5191, in 258 fathoms (472 meters), Tanon Strait, east coast of Negros.

This is a peculiar modification which is thus placed, but which has a rougher, more punctate surface than typical *C. nitida*, and yet has a series of pores along the junctions of the chambers.

Candema nitida, var. *triloba*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14388	U.S.N.M.	1	D5191...	° ' " ° ' "	258	° F. 62.8	gn. m.	Rare.
			D5613...	10 29 45 N.; 123 31 15 E.. 0 42 00 S.; 121 44 00 E..	752	gy. m.	

Genus SPHAEROIDINA d'Orbigny, 1826.

SPHAEROIDINA BULLOIDES d'Orbigny.

Sphaeroidina bulloides D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 267, No. 1; Modèles, 1826, No. 65.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 620, pl. 84, figs. 1-7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 18, pl. 10, fig. 7; pl. 12, fig. 1.

Like *Orbulina universa* this species seems to be limited in the region, while the other species of the genus is very much more common. It occurred at the following stations: D5178, in 78 fathoms (143 meters), vicinity of Romblon; D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.); and D5638, in 517 fathoms (946 meters), off Bouro Island.

Sphaeroidina bulloides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12753	U.S.N.M.	4	D5178...	° ' " ° ' "	78	° F.	fine. s.	Few.
12755	U.S.N.M.	10+	D5236...	12 43 00 N.; 122 06 15 E.. 8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.	Frequent.
12754	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E..	517	fine. gy. s.	Rare.

SPHAEROIDINA DEHISCENS Parker and Jones.

Sphaeroidina dehiscens PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 369, pl. 19, figs. 5a, b.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 621, pl. 84, figs. 8-11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 19, pl. 10, fig. 2; pl. 13, fig. 1.

Globigerina seminulina SCHWAGER, *Novara*-Exped., Geol. Theil., vol. 2, 1886, p. 256, pl. 7, fig. 112.

In contrast with the preceding species this one has been noted at 43 stations in the region. It is rather interesting, however, to note that but very few of these are really among the islands of the Archipelago proper, but, except in the Sulu Sea, the records are mainly from the China Sea; Gulf of Davao; Palawan Passage; Gulf of Tomini, Celebes; Molucca Passage; between Gillolo and Makyan and

between Gillolo and Kayao; Patiente Strait; Pitt Passage; off Bouru Island; Molucca Sea; and Gulf of Boni. Besides these regions, most of which are outside the Philippine Archipelago proper, the species has occurred on the Pacific side of Luzon and Mindanao; Verde Island Passage; between Burias and Luzon; and in Sogod Bay, southern Leyte.

These stations range in depth from 80 to 1,570 fathoms (146 to 2,871 meters), with the average 428 fathoms (783 meters). The average bottom temperature was 51.9° F. (11° C.); but this is too high, as the bottom temperature is not given for many of the deeper stations.

The specimens are very typical, but only at *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, could it be called abundant. There specimens were very abundant and well developed. The bottom temperature there, however, was 41.2° F.

Sphaeroidina dehiscens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
13177	U.S.N.M.	10+	D5110...	13 59 29 N.: 120 75 45 E...	135	59.0	dk. gy. m...	Common.
			D5113...	13 51 30 N.: 120 50 30 E...	159	dk. gn. m...	Few.
			D5114...	13 36 11 N.: 120 45 26 E...	340	fne. s.	Few.
			D5115...	13 37 11 N.: 120 43 40 E...	340	Few.
			D5121...	13 27 20 N.: 121 17 45 E...	108	dk. gn. m...	Several.
			D5172...	6 03 15 N.: 120 35 30 E...	318	fne. s., sh.	Frequent.
			D5201...	10 10 00 N.: 125 04 15 E...	554	52.8	gy. s., m...	Few.
			D5212...	12 04 15 N.: 124 04 36 E...	108	59.9	gy. s., m...	Few.
			D5217...	13 20 09 N.: 123 14 15 E...	105	63.1	crs. gy. s.	Few.
13176	U.S.N.M.	10+	D5236...	8 50 45 N.: 126 26 52 E...	494	41.2	fne. gy. s.	Abundant.
			D5247...	7 02 00 N.: 125 38 45 E...	135	m.	Few.
			D5267...	13 42 20 N.: 120 58 25 E...	p. s., sh.	Rare.
			D5268...	13 42 00 N.: 120 57 15 E...	170	s. p.	Few.
			D5277...	13 56 55 N.: 120 13 45 E...	80	58.6	fne. s.	Rare.
			D5281...	13 52 45 N.: 120 25 00 E...	201	50.4	dk. gy. s.	Rare.
			D5282...	13 63 00 N.: 120 26 45 E...	248	47.4	dk. gy. s.	Few.
13223	U.S.N.M.	2	D5300...	20 31 00 N.: 115 49 00 E...	265	gy. m., s.	Rare.
			D5301...	20 37 00 N.: 115 43 00 E...	208	50.5	gy. m., s.	Rare.
			D5313...	21 30 00 N.: 116 43 00 E...	150	53.6	s.	Few.
			D5348...	10 57 45 N.: 118 38 15 E...	300	Rare.
			D5428...	9 13 00 N.: 118 51 15 E...	1,105	49.7	gy. m.	Few.
			D5443...	12 43 05 N.: 125 01 00 E...	241	51.3	co. s., sh.	Rare.
			D5446...	12 43 51 N.: 124 59 18 E...	300	gn. m.	Frequent.
			D5449...	13 21 36 N.: 124 00 30 E...	300	Rare.
			D5542...	8 48 30 N.: 123 35 30 E...	200	54.3	fne. s., brk.sh	Rare.
13175	U.S.N.M.	2	D5543...	8 47 15 N.: 123 35 00 E...	162	54.5	Few.
			D5551...	5 54 48 N.: 120 44 24 E...	193	53.3	fne. s.	Rare.
			D5570...	5 32 15 N.: 120 12 57 E...	330	52.3	fne. s., glob.	Rare.
			D5612...	0 38 00 S.: 121 45 40 E...	750	Frequent.
			D5614...	0 31 00 N.: 125 58 45 E...	1,100	gy. m., s., glob	Few.
			D5618...	0 37 00 N.: 127 15 00 E...	417	gy. m.	Few.
			D5621...	0 15 00 N.: 127 24 35 E...	298	gy. and bk. s.	Frequent.
			D5623...	0 16 30 N.: 127 30 00 E...	272	fne. s., m...	Rare.
			D5625...	0 07 00 N.: 127 28 00 E...	230	gy. m., fne. s.	Rare.
			D5626...	0 07 30 N.: 127 29 00 E...	215	gy. m., fne. s.	Rare.
12752	U.S.N.M.	5	D5630...	0 56 30 S.: 128 05 00 E...	569	co. s. m.	Frequent.
			D5636...	1 55 00 S.: 127 42 30 E...	1,262	gy. m., fne. s.	Few.
			D5637...	3 53 20 S.: 126 48 00 E...	700	gy. m.	Rare.
			D5638...	3 47 15 S.: 126 23 40 E...	517	fne. gy. s.	Few.
			D5639...	3 54 50 S.: 123 27 20 E...	1,560	gy. m.	Frequent.
			D5650...	4 53 45 S.: 121 29 00 E...	540	40.1	gn. m.	Few.
			H4897...	7 46 00 N.: 122 03 00 E...	1,570	gy. m., glob.	Few.
			H4937...	5 32 50 S.: 120 49 10 E...	885	38.2	gy. m.	Few.
9966	U.S.N.M.	1	Nero 849	15 16 00 N.: 121 40 00 E...	737	gn. m.	Rare.

Genus PULLENIA Parker and Jones, 1862.

PULLENIA SPHAEROIDES (d'Orbigny).

Nonionina sphaeroides D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 293, No. 1.

Pullenia sphaeroides PARKER and JONES, in Carpenter, Parker, and Jones, *Introd. Foram.*, 1862, p. 184, pl. 12, fig. 12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 615, pl. 84, figs. 12, 13, text figure 18, p. 616.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 20, pl. 11, fig. 2.

The only record for this species in the region is *Albatross* station D5639, Molucca Sea, in 1,560 fathoms (2,853 meters). This is the deepest station dredged by the *Albatross*. This specimen is very typical, but larger than usual.

Pullenia sphaeroides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13174	U.S.N.M.	1	D5639...	° ' " ° ' "	1,560	° F.	gy. m.	Rare.
				3 54 50 S.; 123 27 20 E..				

PULLENIA QUINQUELOBA (Reuss).

Nonionina quinqueloba REUSS, Zeitschr. deutsch. geol. Ges., vol. 3, 1851, p. 47, pl. 5, figs. 31a, b.

Pullenia quinqueloba H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 617, pl. 84, figs. 14, 15.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 21, pl. 13, fig. 2.

Single very typical specimens occur at two stations—D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms, (903 meters), and D5526, between Siquijor and Bohol Islands, 805 fathoms (1,472 meters).

All of the records for this species elsewhere are in comparatively deep water, and it never seems to occur in any considerable numbers.

Pullenia quinqueloba—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15354	U.S.N.M.	1	D5236...	° ' " ° ' "	494	° F.		Rare.
13119	U.S.N.M.	1	D5526...	8 50 45 N.; 126 26 52 E..	805	41.2	fine. gy. s....	Rare.
				9 12 45 N.; 123 45 30 E..		52.3	gn. m., glob.	

PULLENIA OBLIQUOCULATA Parker and Jones.

Pullenia obliquoculata PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 368, pl. 19, figs. 4a, b.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 618, pl. 84, figs. 16–20.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 22, pl. 10, fig. 3; pl. 12, figs. 2, 3.

This is by far the commonest species of the genus in the area. I have noted its occurrence in material from over a hundred stations

in the area, but, as in the case of *Sphaeroidina dehiscens*, the great majority of the stations are in the China Sea, off the Pacific coast of the islands, or the greater bodies of water to the south; off Bouru, Celebes, Gillolo, Patiente Strait, Pitt Passage, Molucca Sea, Buton Strait, Gulf of Boni, Flores Sea, and Macassar Strait. At the few stations inside the Archipelago the specimens were few, while in the deeper waters outside they are often very common.

The range of depth of these stations ranges from 10 to 1,570 fathoms (18 to 2,871 meters), the average being 363 fathoms (664 meters), and the average bottom temperature for 51 stations at which the bottom temperature was given was 51.8° F. (10.9° C.). This is much too high, however, for at most of the deeper stations the temperature was not given.

Pullenia obliquiloculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13172	U.S.N.M.	10+	D5096	14 20 23 N.; 120 34 15 E.	28	gy. m., s., sh	Few.
			D5100	14 17 15 N.; 120 32 40 E.	35	gy. s.	Few.
			D5109	14 03 45 N.; 120 16 30 E.	10	co.	Few.
			D5110	13 59 20 N.; 120 75 45 E.	135	59.0	dk. gy. m.	Common.
			D5112	13 48 22 N.; 120 47 25 E.	177	52.4	dk. gn. m.	Frequent.
			D5113	13 51 30 N.; 120 50 30 E.	159	dk. gn. m.	Frequent.
			D5114	13 36 11 N.; 120 45 26 E.	340	fine. s.	Common.
			D5115	13 37 11 N.; 120 43 40 E.	340	Common.
			D5120	13 45 30 N.; 120 30 15 E.	393	43.7	gn. m., s.	Few.
			D5121	13 27 20 N.; 121 17 45 E.	108	dk. gn. m.	Few.
			D5126	10 34 45 N.; 121 47 30 E.	742	49.5	sft. gn. m.	Few.
			D5132	Id. off Panabutan Pt. N. 15, W. 0.3 mi.	26	gn. m., s.	Few.
			D5153	5 18 10 N.; 120 02 55 E.	49	co. s., sh.	Frequent.
			D5172	6 03 15 N.; 120 35 30 E.	318	fine. s., sh.	Common.
			D5178	12 43 00 N.; 122 06 15 E.	78	fine. s.	Common.
			D5179	12 38 15 N.; 122 12 30 E.	37	75.7	hrd. s.	Few.
			D5184	10 18 30 N.; 122 23 30 E.	565	49.8	gn. m.	Few.
			D5185	10 05 45 N.; 122 18 30 E.	638	49.8	gn. m.	Frequent.
			D5187	9 16 45 N.; 123 21 15 E.	225	53.6	sft. gn. m.	Frequent.
			D5189	9 56 30 N.; 123 15 00 E.	300	62.8	gn. m.	Few.
			D5191	10 29 45 N.; 123 31 15 E.	258	62.8	gn. m.	Common.
			D5192	11 19 15 N.; 123 50 00 E.	32	gn. s.	Few.
			D5201	10 10 00 N.; 125 04 15 E.	554	52.8	gy. s., m.	Common.
			D5203	9 58 00 N.; 125 07 40 E.	775	52.9	gn. m.	Common.
			D5206	11 31 40 N.; 124 42 40 E.	32	gn. m.	Few.
			D5211	11 51 35 N.; 124 14 00 E.	155	56.6	gn. m., s.	Few.
			D5215	12 31 30 N.; 123 35 24 E.	604	50.5	gn. m.	Few.
			D5216	12 52 00 N.; 123 23 30 E.	200	50.2	Few.
			D5217	13 20 00 N.; 123 14 15 E.	105	63.1	crs. gy. s.	Frequent.
			D5219	13 21 00 N.; 122 18 45 E.	530	50.8	gn. m.	Common.
			D5220	13 38 00 N.; 121 58 00 E.	50	sft. gn. m.	Few.
			D5222	13 38 30 N.; 121 42 45 E.	195	52.8	gn. m.	Few.
			D5257	7 22 12 N.; 124 12 15 E.	28	m.	Few.
			D5259	11 57 30 N.; 121 42 15 E.	312	49.3	gy. m., glob.	Abundant.
			D5260	12 25 35 N.; 121 31 35 E.	234	51.4	gn. m., s.	Few.
			D5261	12 30 55 N.; 121 34 24 E.	145	s., m.	Few.
			D5267	13 42 20 N.; 120 58 25 E.	170	p., s., sh.	Rare.
			D5268	13 42 00 N.; 120 57 15 E.	170	s., p.	Common.
			D5282	13 53 00 N.; 120 26 45 E.	248	47.4	dk. gy. s.	Few.
			D5284	13 42 05 N.; 120 30 45 E.	422	42.3	gy. m., glob.	Few.
			D5291	13 29 40 N.; 121 00 45 E.	173	51.5	fine. bk. s.	Few.
			D5300	20 31 00 N.; 115 49 00 E.	265	gy. m., s.	Common.
			D5311	21 33 00 N.; 116 15 00 E.	88	crs. s., sh.	Rare.
			D5313	21 30 00 N.; 116 43 00 E.	150	53.6	s.	Frequent.
			D5315	21 40 00 N.; 116 58 00 E.	148	54.4	s., sh.	Few.
			D5342	10 56 55 N.; 119 17 24 E.	14	gy. m.	Few.
			D5348	10 57 45 N.; 118 38 15 E.	375	56.4	co. s.	Few.
			D5349	10 54 00 N.; 118 26 20 E.	730	40.6	co. s.	Common.
			D5370	13 44 15 N.; 121 42 30 E.	159	54.3	sft. m.	Few.
			D5377	13 26 00 N.; 122 19 00 E.	240	54.3	crs. s.	Rare.
			D5381	13 14 15 N.; 122 44 25 E.	88	co. s.	Few.
12759	U.S.N.M.	10+	D5300	20 31 00 N.; 115 49 00 E.	265	gy. m., s.	Common.
			D5311	21 33 00 N.; 116 15 00 E.	88	crs. s., sh.	Rare.
			D5313	21 30 00 N.; 116 43 00 E.	150	53.6	s.	Frequent.
			D5315	21 40 00 N.; 116 58 00 E.	148	54.4	s., sh.	Few.
			D5342	10 56 55 N.; 119 17 24 E.	14	gy. m.	Few.
			D5348	10 57 45 N.; 118 38 15 E.	375	56.4	co. s.	Few.
			D5349	10 54 00 N.; 118 26 20 E.	730	40.6	co. s.	Common.
			D5370	13 44 15 N.; 121 42 30 E.	159	54.3	sft. m.	Few.
			D5377	13 26 00 N.; 122 19 00 E.	240	54.3	crs. s.	Rare.
			D5381	13 14 15 N.; 122 44 25 E.	88	co. s.	Few.

Pullenia obliquiloculata—Material examined—Continued.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
			D5394...	12 00 30 N.	124 05 36 E.	153	° F.	gn. m.	Few.
			D5428...	9 13 00 N.	118 51 15 E.	1,105	49.4	gy. m.	Few.
			D5430...	9 49 40 N.	119 03 20 E.	464	50.0	glob. oz.	Rare.
			D5443...	12 43 05 N.	125 01 00 E.	241	51.3	co. s., sh.	Frequent.
			D5445...	12 44 42 N.	124 59 50 E.	383	44.3	gn. m., s.	Few.
			D5449...	13 21 36 N.	124 00 30 E.	300	Few.
			D5459...	13 10 21 N.	123 59 54 E.	201	Few.
			D5467...	13 35 27 N.	123 37 18 E.	480	gy. m.	Few.
			D5469...	13 36 48 N.	123 38 24 E.	500	gn. m.	Few.
			D5470...	13 37 30 N.	123 41 09 E.	560	m.	Few.
			D5481...	10 27 30 N.	125 17 10 E.	61	s., sh., g.	Few.
			D5512...	8 16 02 N.	123 58 26 E.	445	52.8	gy. m., fine s.	Frequent.
			D5523...	8 48 44 N.	123 27 35 E.	Few.
			D5526...	9 12 45 N.	123 45 30 E.	805	52.3	gn. m., glob.	Common.
			D5529...	9 23 45 N.	123 39 30 E.	441	53.0	gy. m., glob.	Frequent.
			D5542...	8 48 30 N.	123 35 30 E.	200	54.3	fine s., brk. sh.	Few.
			D5543...	8 47 15 N.	123 35 00 E.	162	54.5	s.	Few.
			D5546...	6 06 48 N.	121 20 32 E.	138	58.3	fine co., s.	Few.
			D5548...	6 00 20 N.	120 45 35 E.	232	53.5	s., brk. sh.	Few.
			D5549...	6 01 15 N.	120 44 20 E.	263	52.3	s., glob. for.	Few.
			D5551...	5 54 48 N.	120 44 24 E.	193	53.3	fine, s.	Few.
			D5565...	5 51 42 N.	120 30 30 E.	243	52.3	s., ptr., sh.	Few.
			D5570...	5 32 15 N.	120 12 57 E.	330	52.3	fine s., glob.	Few.
			D5571...	5 30 45 N.	120 07 57 E.	340	52.3	s., sh.	Few.
13173	U.S.N.M.	4	D5572...	5 31 26 N.	120 09 45 E.	334	52.3	s.	Common.
			D5576...	5 25 56 N.	120 08 39 E.	277	53.3	s.	Few.
			D5579...	4 54 15 N.	119 09 52 E.	175	55.3	fine s., Co.	Few.
			D5580...	4 52 45 N.	119 06 45 E.	162	55.8	br. s., Co.	Common.
			D5590...	4 10 50 N.	118 39 35 E.	310	44.3	gn. m., s.	Few.
			D5601...	1 13 10 N.	125 17 05 E.	765	s., glob. ptr.	Common.
			D5612...	0 38 00 S.	121 45 40 E.	750	Few.
			D5614...	0 31 00 N.	125 58 45 E.	1,100	gy. m., s., glob.	Frequent.
			D5618...	0 37 00 N.	127 15 00 E.	417	gy. m.	Few.
13171	U.S.N.M.	7	D5620...	0 21 30 N.	127 16 45 E.	358	gy. m.	Few.
			D5621...	0 15 00 N.	127 24 35 E.	298	gy. and bk. s.	Frequent.
			D5622...	0 19 20 N.	127 28 30 E.	275	gy. m.	Frequent.
			D5623...	0 16 30 N.	127 30 00 E.	272	fine s., m.	Few.
			D5625...	0 07 00 N.	127 28 00 E.	260	gy. m., fine s.	Common.
			D5626...	0 07 30 N.	127 29 00 E.	235	gy. m., fine s.	Few.
			D5627...	0 06 00 N.	127 26 00 E.	22	Few.
			D5630...	0 56 30 S.	128 05 00 E.	569	co. s., m.	Common.
			D5632...	1 00 00 S.	127 50 00 E.	845	Few.
			D5636...	1 55 00 S.	127 42 30 E.	1,262	gy. m., fine s.	Frequent.
			D5637...	3 53 20 S.	126 48 00 E.	700	gy. m.	Few.
			D5638...	3 47 15 S.	126 23 40 E.	517	fine gy. s.	Frequent.
			D5639...	3 54 50 S.	123 27 20 E.	1,560	gy. m.	Frequent.
			D5648...	5 35 00 S.	122 20 00 E.	559	gn. m.	Rare.
			D5650...	4 53 45 S.	121 29 00 E.	540	40.1	gn. m.	Few.
			D5652...	4 35 00 S.	121 23 06 E.	525	41.2	gn. m.	Few.
			D5660...	5 36 30 S.	120 49 00 E.	692	39.2	gy. m., s.	Frequent.
			D5665...	4 27 00 S.	118 44 00 E.	1,008	m.	Common.
9972	U.S.N.M.	2	H4914...	13 38 05 N.	120 33 00 E.	464	46.5	gy. m., s.	Few.
			H4897...	7 46 00 N.	122 03 00 E.	1,570	gy. m., glob.	Common.
			H4937...	5 32 50 S.	120 49 10 E.	885	38.2	gy. m.	Few.

Family ROTALIIDAE.

Subfamily SPIRILLININAE.

Genus SPIRILLINA Ehrenberg, 1841.

SPIRILLINA LIMBATA H. B. Brady.

Spirillina limbata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 278, pl. 8, figs. 26a, b; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 632, pl. 85, figs. 18-21.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 5, pl. 2, figs. 1, 2, fig. 5 (in text).

The only specimens in the material which can be referred to the typical form of this species are from *Albatross* station D5179, off

Romblon, in 37 fathoms (68 meters). The material here noted has a tendency to be tuberculate on the underside, as in the second variety here recorded. Brady did not figure the underside of this species, but the description infers that it is smooth.

Spirillina limbata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14406	U.S.N.M.	1	D5179...	° / ' ' ° / ' ' 12 38 15 N.; 122 12 30 E...	37	° F. 75.7	hrd. s.	Rare.

SPIRILLINA LIMBATA H. B. Brady, var. *DENTICULATA* H. B. Brady.

Spirillina limbata H. B. BRADY, var. *denticulata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 632, pl. 85, fig. 17.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 5, pl. 3, figs. 1, 2.

This variety, although apparently much more common than the typical, occurred, as far as the material shows, at one station—D5179, off Romblon, in 37 fathoms (68 meters), with both the typical and the following variety. It is undoubtedly common in the region, although material was not saved.

Spirillina limbata, var. *denticulata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14407	U.S.N.M.	1	D5179...	° / ' ' ° / ' ' 12 38 15 N.; 122 12 30 E...	37	° F. 75.7	hrd. s.	Rare.

SPIRILLINA LIMBATA H. B. Brady, var. *TUBERCULO-LIMBATA* Chapman.

Spirillina tuberculo-limbata CHAPMAN, Journ. Linn. Soc. London, Zoology, vol. 28, 1900, p. 11, pl. 1, figs. 8a-c.

Spirillina limbata H. B. BRADY, var. *papillosa* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 6, pl. 2, fig. 4.

From an examination of the Philippine material it seems that Chapman's species, described from Funafuti, and the variety I described from the Hawaiian Islands, are probably the same. The occurrence of this variety with the typical seems to show that it is but a varietal form, as the differences are largely in ornamentation and the number of coils of the test.

It has occurred at D5178, in 78 fathoms (143 meters), and D5179 in 37 fathoms (68 meters), both stations off Romblon. The bottom temperature at the shallower station is 75.7° F., (24.2° C.).

These records indicate that this variety is widely distributed in the Indo-Pacific region.

Spirillina limbata, var. *tuberculo-limbata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14409	U.S.N.M.	1	D5178...	° ' " ° ' "	78	° F.	fne. s.	Rare.
14408	U.S.N.M.	1	D5179...	12 43 00 N.; 122 06 15 E.. 12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.	Rare.

Subfamily ROTALINAE.

Genus PATELLINA Williamson, 1858.

PATELLINA CORRUGATA Williamson.

Patellina corrugata WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 46, pl. 3, figs. 86-89.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 634, pl. 86, figs. 1-7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 9, pl. 7, fig. 1.

A single specimen of this species was found at *Albatross* station, D5551, in 193 fathoms (353 meters), bottom temperature 53.3° F. (11.8° C.), near Jolo Island. This is the only station at which the species was noted in all the abundant material from the Philippine Expedition, and it seems to be decidedly rare in this region. The *Challenger* did not obtain the species from the Philippine stations dredged during that expedition. Millett records it as "rather scarce" in the Malay Archipelago. The only records for the North Pacific outside this region are two records of a single specimen from each station from Laysan and Chatham Islands, recorded by Rhumbler.

The single Philippine specimen was a very typical and well-developed one.

Patellina corrugata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15668	U.S.N.M.	1	D5551...	° ' " ° ' "	193	° F. 53.3	fne. s.	Rare.

Genus DISCORBIS Lamarck, 1804.

This genus is one which is either in a state of flux with a few very variable species or else is represented by many species. Heron-Allen and Earland have given much attention to this problem in

their Clare Island work.¹⁹ A comparison of the figures of the *Challenger* Report with the original figures of the species to which Brady referred his material will show that in most cases the choice of names was hardly wise. So other writers, following the *Challenger* determinations and adding their own variant views, have created a complexity of nomenclature that it is very difficult to unravel. However, the figures of the *Challenger* Report are very accurate, and a study of the originals with notes as to their occurrence would help much. In the Philippine material the genus has been very poorly represented, probably because the specimens are usually small and were lost in the original washing. Therefore, it is impossible with the meagre material to add greatly to the clearing up of the problems waiting to be solved. In certain of the species reference to the *Challenger* figures is given rather than describing them as new.

DISCORBIS SUBOBTUSA, new species.

Plate 70, figs. 2a-c.

Description.—Test circular in outline, periphery rounded, chambers about six or seven in the last-formed coil; sutures oblique, distinct, flush with the surface, ventral side more rounded; sutures nearly at right angles to the periphery, ending in slight, blunt projections in the middle, slightly umbilicate; aperture narrow, the elongate base of the last-formed chamber extending about halfway from the umbilicus to the periphery; wall smooth and slightly polished, punctate.

Diameter about 1.5 mm.

Distribution.—Type specimen (Cat. No. 15356 U. S. N. M.) from *Albatross* station D5236, off the east coast of Mindanao, in 494 fathoms (903 meters).

This is a large species, different from any others in the region. It is not found elsewhere.

Discorbis subobtusa.—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15356	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	°F. 41.2	fine gy. s.....	Rare.

DISCORBIS VILARDEBOANA (?) d'Orbigny.

The material from *Albatross* D5144, off Jolo, in 19 fathoms (35 meters), is almost exactly like the figure given by Brady in plate 86, figure 12, of the *Challenger* Report. It is evidently not the same as

¹⁹ The Foraminifera of the Clare Island District, County Mayo, Ireland, Proc. Royal Irish Acad., vol. 31, pt. 64, 1913, pp. 110-141.

d'Orbigny's *Rosalina vilardeboana*, as a comparison of the two will at once show.

From another station off Jolo, D5546, in 138 fathoms (252 meters), is a specimen which might almost have served for the original of Brady's plate 86, figure 9. This form is referred to *D. mediterraneensis* d'Orbigny by Heron-Allen and Earland, but does not seem the same as their figured specimens from Clare Island.

Discorbis vilardeboana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14399	U.S.N.M.	1	D5144...	6 05 50 N.; 121 02 15 E...	19	°F.	co. s.....	Rare.
14400	U.S.N.M.	1	D5546...	6 06 48 N.; 121 20 32 E...	138	58.3	fine. co. s.....	Rare.

DISCORBIS ORBICULARIS (Terquem).

Rosalina orbicularis TERQUEM, Anim. Plage Dunkerque, 1876, p. 75, pl. 9, figs. 4a, 4b.

Discorbis orbicularis BERTHELIN, Foram. Bourgneuf et Pornichet, 1878, p. 39, No. 63.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 16, pl. 11, fig. 1, figs. 18a-c (in text).

Discorbina orbicularis H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 647, pl. 88, figs. 4-8.

Specimens which are close to this species were obtained from two stations—D5139, in 20 fathoms (37 meters), and D5141, in 29 fathoms (53 meters), both off Jolo.

Discorbis orbicularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14402	U.S.N.M.	2	D5139...	6 06 00 N.; 121 02 30 E...	20	°F.	co. s.....	Rare.
14403	U.S.N.M.	1	D5141...	6 09 00 N.; 120 58 00 E...	29	co. s.....	Rare.

DISCORBIS BERTHELOTI (d'Orbigny).

Plate 59, figs. 1a-c.

Rosalina bertheloti D'ORBIGNY, in Barker, Webb, and Berthelot, Hist. Nat. Îles Canaries, vol. 2, pt. 2, "Foraminifères," 1839, p. 135, pl. 1, figs. 28-30.

Discorbina bertheloti H. B. BRADY, Trans. Linn. Soc. London, vol. 24, 1864, p. 469, pl. 48, figs. 10a, b; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 650, pl. 89, figs. 10-12.

Discorbis bertheloti CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 20, pl. 7, fig. 3, fig. 23 (in text).

Rare specimens referred to this species were obtained at the following stations: D5164, in 18 fathoms (33 meters), off Tawi Tawi; D5179,

in 37 fathoms (68 meters), off Romblon; D5220, in 50 fathoms (91 meters), between Marinduque and Luzon; D5268, in 170 fathoms (311 meters), Verde Island Passage; D5551, in 193 fathoms (353 meters), off Jolo; and D5576, north of Tawi Tawi, 277 fathoms (506 meters); bottom temperature 53.3° F. (11.8° C.).

Discorbis bertheloti—Material examined.

Cat. No.	Coll. of—	No. of specim.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14396	U.S.N.M.	1	D5164...	5 01 40 N.; 119 52 20 E...	18	gn. m.....	Rare.
14395	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Rare.
14398	U.S.N.M.	1	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m.....	Rare.
			D5268...	13 42 00 N.; 120 57 15 E...	170	p. s., sh.....	
14397	U.S.N.M.	1	D5551...	5 54 48 N.; 120 44 24 E...	193	53.3	fne. s.....	Rare.
15702	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.....	Rare.

DISCORBIS ALLOMORPHINOIDES (Reuss).

Plate 58, figs. 1a-c.

Valvulina allomorphinoides REUSS, Sitz. Akad. Wiss. Wien, vol. 40, 1860, p. 223, pl. 11, fig. 6.

Discorbina allomorphinoides H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 654, pl. 91, figs. 5, 8.

Discorbis allomorphinoides CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 21, pl. 9, fig. 1, figs. 25a-c (in text).

Rare specimens occurred in material from two stations, D5261, in 145 fathoms (265 meters), off southeastern Mindoro, and D5381, in 88 fathoms (161 meters), in Ragay Gulf, Luzon.

Discorbis allomorphinoides—Material examined.

Cat. No.	Coll. of—	No. of specim.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14404	U.S.N.M.	2	D5261...	12 30 55 N.; 121 34 24 E...	145	s., m.....	Rare.
14405	U.S.N.M.	2	D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.....	Rare.

DISCORBIS RUGGSA (d'Orbigny).

Plate 60, figs. 2a-c.

Rosalina rugosa d'ORBIGNY, Foram. Amér. Mérid., 1839, p. 42, pl. 2, figs. 12-14.

Discorbina rugosa H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 652, pl. 87, figs. 3a-c; pl. 91, figs. 4a-c.

A specimen referable to this species occurred at *Albatross* station D5144, off Jolo in 19 fathoms (35 meters). It is close to Brady's figure. The species seems to be largely recorded from the Indo-Pacific in comparatively shallow water. A few specimens were obtained at D5236, in 494 fathoms (903 meters), off Mindanao.

Discorbis rugosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14410	U.S.N.M.	1	D5144... D5236...	° ' " ° ' " 6 05 50 N.; 121 02 15 E.... 8 50 45 N.; 126 26 52 E....	19 494	° F. 41.2	co. s..... fine. gy. s.....	Rare.

DISCORBIS, species (?).

Plate 60, figs. 1a-c.

Discorbina saulcii H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 653, pl. 91, figs. 6a-c (not *Rosalina sauleyi* d'Orbigny, 1839).

It is very difficult to understand how Brady could have referred his material figured in the *Challenger* Report to the species figured by d'Orbigny. Brady refers to this material from the Gulf of Scala Nova, off Smyrna, and *Challenger* material from off Tahiti.

Sidebottom has figured material from the Bay of Palermo, Sicily²⁰ which he refers to this species. He mentions (p. 26) that the material from Sicily agrees with material he has from the coast of Algiers in 80 fathoms (146 meters). He also mentions the Bay of Eleusis, Greece.

As figured, the Mediterranean material has no definite carinate border, is not so rounded as the *Challenger* figure and has a different appearance on the umbilical side.

Therefore, as the matter stands, I would venture to predict that the true *D. sauleyi* of d'Orbigny may be confined to the west coast of South America and adjacent regions, while there are two other distinct species—that figured by Sidebottom, probably a Mediterranean one, widely spread in that region; and a third—that figured by Brady, which is characteristic of the Indo-Pacific. To which of these two the material referred by Heron-Allen and Earland to this species from the east coast of Africa may belong²¹ it would be interesting to know.

This species is best developed at *Albatross* D5236, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.), off Mindanao, and D5201 in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte. It is worth noting that the *Challenger* material off Tahiti was from depths of 420 and 620 fathoms (768 and 1,134 meters), while the Mediterranean records are from less than 100 fathoms (183 meters).

²⁰ Mem. and Proc. Manchester Lit. and Philos. Soc., vol. 54 (No. 16), 1910, pl. 3, figs. 11a-c.

²¹ Trans. Zool. Soc. London, vol 20, 1915, p. 696.

Discorbis, species (?)—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
			D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	
			D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	

Genus CYMBALOPORA Hagenow, 1850.

CYMBALOPORA POEYI (d'Orbigny).

Plate 59, figs. 2a-c.

Rotalia squamosa D'ORBIGNY (*nomen nudum*), Ann. Sci. Nat., vol. 7, 1826, p. 272, No. 8.

Rosalina poeyi D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 92, pl. 3, figs. 18-20.

Cymbalopora poeyi CARPENTER, Introd. Foram., 1862, p. 215, pl. 13, figs. 10-12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 636, pl. 102, figs. 13a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 24, pl. 10, fig. 1; pl. 14, fig. 5.

The typical form of the species has occurred at a number of stations in the Philippine area and to the south, the depths ranging from 12 to 554 fathoms (22 to 1,012 meters), average depth 318 fathoms (582 meters), and the bottom temperatures from 44.3° F. to 63.1° F. (6.8° C. to 17.2° C.). The most of the stations are in the Archipelago proper, but it was also obtained from station D5590, in 310 fathoms (567 meters), bottom temperature 44.3° F. (6.8° C.) in Sibuko Bay, Borneo.

Cymbalopora poeyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
11922	U.S.N.M.	1	D5156...	5 12 55 N.; 119 55 55 E..	18	fine. s., sh....	Rare.
		1	D5160...	5 12 40 N.; 119 55 10 E..	12	s.....	Rare.
11921	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m....	Rare.
		1	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.....	Rare.
11920	U.S.N.M.	1	D5217...	13 00 20 N.; 123 14 15 E..	105	63.1	crs. gy. s....	Rare.
11918	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Rare.
11919	U.S.N.M.	2	D5423...	9 38 30 N.; 121 11 00 E..	508	49.8	gy. m.....	Rare.
11917	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Rare.
		1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.....	Rare.
11743	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s....	Rare.

CYMBALOPORA POEYI (d'Orbigny), var. BRADYI Cushman.

Cymbalopora poeyi, var. H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 637, pl. 102, fig. 14.

Cymbalopora poeyi (d'Orbigny), var. *bradyi* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 25, pl. 10, fig. 2; pl. 14, fig. 2.

This variety differs from the typical form of the species in its much more compressed form and the much more open arrangement of the chambers on the ventral side.

Specimens of this very distinct variety were obtained at four Philippine stations: *Albatross* D5164, in 18 fathoms (33 meters), Sulu Archipelago; D5191, in 258 fathoms (471 meters), Tanon Strait, east coast of Negros Island, bottom temperature 62.8° F. (17.1° C.); D5217, in 105 fathoms (193 meters), between Burias and Luzon, bottom temperature 63.1° F. (17.2° C.); and D5348, in 375 fathoms (686 meters), Palawan Passage, bottom temperature 56.4° F. (13.5° C.). The other North Pacific record is the type station in 392 fathoms (717 meters), off the Hawaiian Islands.

Cymbalopora peoyi, var. *braydi*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15751	U.S.N.M.	2	D5164...	5 01 40 N.; 119 52 20 E..	18	°F.	gn. m.	Rare.
15752	U.S.N.M.	1	D5191...	10 29 45 N.; 123 31 15 E..	258	62.8	gn. m.	Rare.
11923	U.S.N.M.	1	D5217...	13 00 20 N.; 123 14 15 E..	105	63.1	crs. gy. s.	Rare.
15753	U.S.N.M.	3	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	Co., s.	Few.

Genus TRETOMPHALUS Moebius, 1880.

TRETOMPHALUS BULLOIDES (d'Orbigny).

Plate 59, figs. 3a-c.

Rosalina bulloides D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 104, pl. 3, figs. 2-5.

Cymbalopora bulloides CARPENTER, Introd. Foram., 1862, p. 216.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 638, pl. 102, figs. 7-12, text figs. 20a-c.—EARLAND, Journ. Quekett Micr. Club, ser. 2, vol. 8, 1902, pp. 309-322, pl. 16.

Tretomphalus bulloides MOEBIUS, Beitr. Meeresfauna Insel. Mauritius, 1880, p. 98, pl. 10, figs. 6-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 26, pl. 14, figs. 3, 4, text figs. 29a-c.

This is a pelagic species, at least in part, and its distribution by depth may or may not have a definite significance compared with a truly bottom-living species. It has occurred at six stations in the area—D5192, in 32 fathoms (59 meters), off northern Cebu Island; D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.); D5243, in 218 fathoms (399 meters), Pujada Bay, bottom temperature 63.6° F. (17.5° C.); D5257, off southern Mindanao, eastern Illana Bay; D5551, in 193 fathoms (353 meters), vicinity of Jolo Island, bottom temperature 53.3° F. (11.8° C.); and D5576, in 277 fathoms (506 meters), north of Tawi Tawi, bottom temperature 53.3° F. (11.8° C.).

In the *Challenger* Report Brady records this species as pelagic at Zamboanga, Philippine Islands. In the North Pacific it also occurs about the Hawaiian Islands and about Laysan.

Earland has shown the complicated structure of this species and the peculiar adaptation of the "float chamber." It is found in enormous numbers at certain times from the observations of several collectors. Records show that it is well scattered through the Indo-Pacific region, eastward to the Hawaiian Islands, south to Australia, and west to the coast of Africa. It is also recorded from the Mediterranean and from the West Indies.

Tretomphalus bulloides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15749	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	° F.	gn. s.	Rare.
14813	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s.	Rare.
			D5243...	6 50 55 N.; 126 14 35 E...	218	63.6	gy. m.	
15750	U.S.N.M.	1	D5257...	7 22 12 N.; 124 12 15 E...	28		m.	Rare.
11924	U.S.N.M.	1	D5551...	5 54 48 N.; 120 44 24 E...	193	53.3	fine. s.	Rare.
			D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.	

Genus *PLANORBULINA* d'Orbigny, 1826.

PLANORBULINA LARVATA Parker and Jones.

Planorbulina vulgaris D'ORBIGNY, var. *larvata* PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 5, 1860, p. 294.

Planorbulina larvata PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 379, pl. 19, figs. 3a, b.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 658, pl. 92, figs. 5, 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 27, pl. 8, fig. 2.

This species seems to be very clearly a tropical one, and is, as might be expected, a very characteristic species of the Philippine Archipelago, especially in comparatively shallow waters. Outside the Archipelago proper it was noted in the material from the Gulf of Boni; Buton Strait; Sibuko Bay and Darvel Bay, Borneo. The numerous stations from the Archipelago give it a rather general distribution among the islands. The depths range from 17 to 805 fathoms (31 to 1,472 meters), with the average depth 177 fathoms (324 meters). Bottom temperatures where given range from 38.3° to 58.6° F. (3.5° to 14.7° C.), with the average 49° F. (9.5° C.). These temperatures are really too low, as the bottom temperatures are not given in many cases from the shallow-water stations.

From the great mass of material available it would seem as though this species clearly displaced *P. mediterranensis* in the Philippine region, as this latter species was not found at all.

Planorbulina larvata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11947	U.S.N.M.	10+	D5132...	Island off Panabutan Point, N. 15° W., 0.30 mile.	26	° F.	gn. m., s....	Common.
11943	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	gn. m., s....	Common.
11939	U.S.N.M.	10+	D5142...	6 06 10 N.; 121 02 40 E...	21	co. s., sh....	Common.
11945	U.S.N.M.	10+	D5145...	6 04 30 N.; 120 59 30 E...	23	eo. s., sh....	Common.
11946	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Common.
11938	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E...	73	fine. s.....	Common.
11940	} U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Common.
11941		10+			554	52.8	gy. s., m....	Common.
11954	U.S.N.M.	10+	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Common.
11952	} U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Common.
11956		10+		
11741	} U.S.N.M.	5	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s....	Common.
11953		4		
11948	U.S.N.M.	6	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine. s.....	Common.
11951	U.S.N.M.	7	D5336...	11 37 45 N.; 119 46 00 E...	46	s., m.....	Common.
11950	U.S.N.M.	8	D5381...	13 14 15 N.; 122 44 45 E...	83	co. s.....	Common.
11949	U.S.N.M.	5	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.....	Few.
11742	U.S.N.M.	5	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fine. s., Co....	Few.
11955	U.S.N.M.	8	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co....	Common.
11937	U.S.N.M.	4	D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fine. gy. s....	Few.
11958	U.S.N.M.	10+	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh....	Common.
11944	U.S.N.M.	3	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Few.
11942	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3	Rare.
14401	U.S.N.M.	3	D5148...	5 35 40 N.; 120 47 30 E...	17	co. s.....	Rare.

PLANORBULINA ACERVALIS H. B. Brady.

Planorbulina acervalis H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 657, pl. 92, fig. 4.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 328, pl. 72, fig. 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 29, pl. 14, fig. 3.

Compared with *P. larvata* this species is rare, being found at but few stations and then represented by very few specimens. It occurred at the following stations: D5191, in 258 fathoms (472 meters), Tanon Strait, east coast of Negros, bottom temperature 62.8° F. (17.1° C.); D5255, in 100 fathoms (183 meters), Gulf of Davao; D5419, in 175 fathoms (320 meters), between Cebu and Bohol, bottom temperature 54.5° F. (12.5° C.); D5469, in 500 fathoms (914 meters), east coast of Luzon; and D5546, in 138 fathoms (252 meters), near Jolo Island, bottom temperature 58.3° F. (14.6° C.).

Brady's notes in regard to the distribution of this species contains the following sentence:

They frequent shallow water, and are most plentiful on bottoms of less than 20 or 30 fathoms (37 or 55 meters), but are sometimes found at much greater depths.

The Philippine material seemed to be represented in the shallow water almost entirely by *P. larvata*, and the few records for *P. acervalis* are all from comparatively deep waters.

Planorbulina acervalis.—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
11959	U.S.N.M.	2	D5191...	10	29 45 N.; 123 31 15 E...	258	° F. 62.8	gn. m.....	Rare.
11960				7	03 00 N.; 125 39 00 E....	100	sft. m.....	Rare.
11962	U.S.N.M.	1	D5255...	9	58 30 N.; 123 46 00 E....	175	54.5	gn. m.....	Rare.
11961	U.S.N.M.	1	D5419...	13	36 48 N.; 123 38 24 E....	500	gn. m.....	Rare.
11963	U.S.N.M.	1	D5469...	6	06 48 N.; 121 20 32 E....	138	58.3	fine. co. s....	Rare.
14112	U.S.N.M.	1	D5546...	6	06 48 N.; 121 20 32 E....	138	58.3	fine. co. s....	Rare.

Genus TRUNCATULINA d'Orbigny, 1826.

TRUNCATULINA REFULGENS (Montfort).

Plate 63, figs. 1a-c.

"*Hammonia Balanus* seu *Balanoidea*" SOLDANI, Testaceographia, vol. 1, pt. 1, 1789, p. 58, pl. 46, figs. *nn, oo*.

Cibicides refulgens MONTFORT, Conch. Syst., vol. 1, 1808, p. 122.

Truncatulina refulgens d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 279, pl. 13, figs. 8-11; Modèles, 1826, No. 77.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 659, pl. 92, figs. 7-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 30, pl. 12, fig. 2, text fig. 33.

Typical specimens of this species were obtained in material from numerous stations, ranging in depth from 32 to 901 fathoms (59 to 1,648 meters), the average depth being 268 fathoms (491 meters); bottom temperatures range from 38.2° to 75.7° F. (3.4° to 24.2° C.), the average being 54.1° F. (12.2° C.).

These stations cover a wide area in the Archipelago—off southern Luzon; off Jolo; off Romblon; off northern Cebu; Sogod Bay, southern Leyte; between Burias and Luzon; east coast of Mindanao; Verde Island Passage; between Masbate and Leyte; between Negros and Siquijor; north of Tawi Tawi and outside to the southward off Darvel Bay, Borneo; between Gillolo and Makyan Islands; and in Macassar Strait.

Truncatulina refulgens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
13218	U.S.N.M.	2	D5469...	13	36 48 N.; 123 38 24 E....	500	° F.	gn. m.....	Rare.
14792	U.S.N.M.	1	D5110...	13	59 20 N.; 120 75 45 E....	135	59.0	dk. gy. m.....	Rare.
			D5172...	6	03 15 N.; 120 35 30 E....	318	fine. s., sh....	Rare.
14794	U.S.N.M.	2	D5236...	8	50 45 N.; 126 26 52 E....	494	41.2	fine. gy. s....	Rare.
			D5192...	11	09 15 N.; 123 50 00 E....	32	gn. s.....	Rare.
14793	U.S.N.M.	6	D5268...	13	42 00 N.; 120 58 25 E....	170	p., s., sh....	Few.
15720	U.S.N.M.	1	D5178...	12	43 00 N.; 122 06 15 E....	78	fine. s.....	Few.
			D5179...	12	38 15 N.; 122 12 30 E....	37	75.7	hrd. s.....	Few.
14790	U.S.N.M.	2	D5269...	13	39 50 N.; 120 59 30 E....	220	fine. s., p....	Rare.
			D5201...	10	10 00 N.; 125 04 15 E....	554	52.8	gy. s., m.....	Rare.
14791	U.S.N.M.	1	D5538...	13	56 55 N.; 120 13 45 E....	80	58.6	fine. s.....	Rare.
			D5217...	13	09 20 N.; 123 14 15 E....	105	63.1	crs. gy. s....	Rare.
14893	U.S.N.M.	1	D5282...	13	03 15 N.; 123 23 20 E....	256	53.3	gn. m., s....	Rare.
			D5282...	13	03 00 N.; 120 26 45 E....	248	47.4	dk. gy. s....	Rare.
14890	U.S.N.M.	1	D5574...	5	30 45 N.; 120 07 57 E....	340	Rare.
			D5398...	11	35 12 N.; 124 13 48 E....	114	gn. m.....	Rare.
14891	U.S.N.M.	1	D5621...	0	15 00 N.; 127 24 35 E....	298	gy. and bk. s.	Rare.
			D5571...	5	30 45 N.; 120 07 57 E....	340	52.3	s., sh....	Rare.
14773	U.S.N.M.	1	D5623...	0	16 30 N.; 127 30 00 E....	272	fine. s., m.....	Rare.
			D5572...	5	31 26 N.; 120 09 45 E....	334	52.3	s.....	Rare.
14774	U.S.N.M.	1	D5668...	2	28 15 S.; 118 49 00 E....	901	38.2	gy. m.....	Rare.
			D5588...	4	52 45 N.; 119 06 45 E....	162	55.8	br. s., Co....	Rare.
15832	U.S.N.M.	1	D5546...	6	06 48 N.; 121 20 32 E....	138	58.3	fine. co. s....	Rare.

TRUNCATULINA LOBATULA (Walker and Jacob).

Plate 63, figs. 2a-c.

"*Nautilus spiralis lobatus*, etc." WALKER and BOYS, Test. Min., 1784, p. 20, pl. 3, fig. 71.

"*Hammonia tuberculata*, etc." SOLDANI, Testaceographia, vol. 1, pt. 1, 1789, p. 58, pl. 45, figs. *ii, kk, ll, mm*.

Nautilus lobatulus WALKER and JACOB, Adam's Essays, Kanmacher's ed., 1798, p. 642, pl. 14, fig. 36.

Serpula lobatula MONTAGU, Test. Brit., 1803, p. 515, Suppl., p. 160.

Truncatulina lobatula D'ORBIGNY, in Barker, Webb, and Berthelot, Hist. Nat. Îles Canaries, vol. 2, pt. 2, "Foraminifères," 1839, p. 134, pl. 2, figs. 22-24.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 660, pl. 92, fig. 10; pl. 93, figs. 1, 4, 5; pl. 95, figs. 4, 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 31, pl. 15, fig. 1, text fig. 34.

The occurrence of this species in the Philippine material seems to indicate that it is not common in warm waters compared to its abundance in cooler shallow waters. It is worthy of note that the bottom temperatures of the Philippine stations at which this species occurred and where temperatures are given range from 41.2° to 44.3° F. (5.1° to 6.8° C.), which for this material are decidedly low temperatures. The depths range from 20 to 750 fathoms (37 to 1,370 meters), with the average 282 fathoms. In the Archipelago the species occurred near Basilan Island, in the Sulu Archipelago; vicinity of Jolo; off northern Cebu; and east coast of Mindanao. Except at the last locality, where it was frequent, the species is comparatively few in numbers. To the southward it occurred at a few stations off Sibuko Bay, Borneo; Gulf of Tomini, Celebes; and Flores Sea.

Truncatulina lobatula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14892	U.S.N.M.	3	D5139...	6 06 00 N.; 121 02 30 E...	20	° F.	eo. s.....	Few.
14894	U.S.N.M.	1	D5141...	6 09 00 N.; 120 58 00 E...	29		co. s.....	Rare.
14895	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s.....	Common.
15721	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 45 E...	88		co. s.....	Rare.
14896	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E...	750		Rare.
14896	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.....	Rare.

TRUNCATULINA TENUIMARGO H. B. BRADY.

Truncatulina tenuimargo H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 62, pl. 93, figs. 2, 3.—EGGER, Abh. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 379, pl. 16, figs. 7-9.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 257, pl. 6, figs. 20 and 21.—CHAPMAN, Trans. New Zealand Inst., vol. 38, 1905, p. 102.

This interesting species, which is now known from several stations in the Indo-Pacific, has occurred at two Philippine stations—D5201, Sogod Bay, southern Leyte, 554 fathoms (1,012 meters), and D5277, China Sea, off southern Luzon, 80 fathoms (146 meters).

These are the usual flattish form figured by Brady. None of the specimens are like those figured by Sidebottom.

Truncatulina tenuimargo—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15727	U.S.N.M.	3	D5201...	° ' " ° ' "		° F.		
15728	U.S.N.M.	2	D5277...	10 10 00 N.; 125 04 15 E.. 13 56 55 N.; 120 13 45 E..	554 80	52.8 59.6	gy. s., m.... fine. s., m....	Few. Rare.

TRUNCATULINA VARIABILIS d'Orbigny.

Plate 65, fig. 2.

"Testae hammoniformes, plano-cochleatae, tuberosae articulatae, etc." SOLDANI, Testaceographia, vol. 1, pt. 1, 1789, pp. 77-80, pls. 70-92.

Truncatulina variabilis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 279, No. 8; in Barker, Webb, and Berthelot, Hist. Nat. Îsles Canaries, vol. 2, pt. 2, "Foraminifères," 1839, p. 135, pl. 2, fig. 29.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 661, pl. 93, figs. 6, 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 33, text fig. 35.

Rare specimens referable to this species were obtained at the following stations: D5110, in 135 fathoms (247 meters), China Sea off southern Luzon, bottom temperature 59° F. (15° C.); D5212, in 108 fathoms (198 meters), east of Masbate Island, bottom temperature 59.9° F. (15.5° C.); D5217, in 105 fathoms (193 meters), between Burias and Luzon, bottom temperature 63.1° F. (17.2° C.); D5272, China Sea off southern Luzon, 118 fathoms (203 meters); and D5546, in 138 fathoms (252 meters), vicinity of Jolo Island, bottom temperature 58.3° F. (14.6° C.).

Truncatulina variabilis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14419	U.S.N.M.	1	D5110...	° ' " ° ' "		° F.		
14418	U.S.N.M.	1	D5212...	13 59 20 N.; 120 75 45 E.. 12 04 15 N.; 124 04 36 E..	135 108	59.0 59.9	dk. gy. m.... gy. s., m....	Rare. Rare.
14417	U.S.N.M.	1	D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s....	Rare.
15736	U.S.N.M.	2	D5546...	6 06 48 N.; 121 20 32 E.. 14 00 00 N.; 120 22 30 E..	138 118	58.3 57.4	fine. co. s.... m., sh., co. s.	Rare. Rare.

TRUNCATULINA WUELLERSTORFI (Schwager).

Plate 64, figs. 1a-c.

Anomalina wuellerstorfi SCHWAGER, Novara-Exped., geol. Theil., vol. 2, 1866, p. 258, pl. 7, figs. 105, 107.

Truncatulina wuellerstorfi H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 662, pl. 93, figs. 8, 9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 34, pl. 12, fig. 3, text fig. 36.

Throughout the region, both in the Archipelago and in the region to the south, this species occurs, but usually is represented by but few specimens at any station. The stations range in depth from 88 to 1,560 fathoms (161 to 2,853 meters), with the average depth 490 fathoms (897 meters), and the bottom temperatures from 39.2° to 56.4° F. (4° to 13.5° C.), the average being 46.8° F. (8.2° C.).

In the region to the southward of the Archipelago proper the species occurs in Sibuko Bay, Borneo; Gulf of Tomini; Molucca Passage; south of Bouro Island; Molucca Sea; Gulf of Boni; Flores Sea; and Macassar Strait.

Truncatulina wuellerstorfi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14783	U.S.N.M.	8	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Common.
15833	U.S.N.M.	3	D5222...	3 38 30 N.; 121 42 45 E...	195	52.8	gn. m.	Few.
14782	U.S.N.M.	2	D5261...	12 30 55 N.; 121 34 24 E...	145	s., m.	Rare.
14766	U.S.N.M.	1	D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh....	Rare.
14767	U.S.N.M.	3	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.	Rare.
14768	U.S.N.M.	2	D5349...	10 54 00 N.; 118 26 20 E...	730	40.6	Co., s.	Rare.
15739	U.S.N.M.	2	D5465...	13 39 42 N.; 123 40 39 E...	500	gn. m.	Rare.
14769	U.S.N.M.	3	D5526...	9 12 45 N.; 123 45 30 E...	805	52.3	gn. m., glob.	Rare.
14770	U.S.N.M.	2	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s....	Rare.
14771	U.S.N.M.	3	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.	Rare.
14772	U.S.N.M.	2	D5660...	5 36 30 S.; 120 49 00 E...	692	39.2	gy. m., s....	Rare.
15738	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.	Rare.

TRUNCATULINA HAIDINGERII (d'Orbigny).

Plate 64, figs. 3a-c.

Rotalina haidingerii D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 154, pl. 8, figs. 7-9.

Planorbulina haidingerii H. B. BRADY, Trans. Linn. Soc. London, vol. 24, 1864, p. 469, pl. 48, fig. 11.

Truncatulina haidingerii REUSS, Sitz. kais. Akad. Wiss. Wien, vol. 55, 1867, p. 28.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 663, pl. 95, figs. 7a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 35, pl. 13, fig. 5; pl. 28, fig. 1, text fig. 37.

This species has been noted at but few stations, all but one (Darvel Bay, Borneo, D5580) in the Archipelago proper. In depth these stations range from 23 to 334 fathoms (42 to 611 meters), the average depth being 193 fathoms (353 meters). Bottom temperatures range from 50.5° to 63.1° F. (10.2° to 17.2° C.).

The stations at which this occurs are near Jolo, D5145; between Burias and Luzon, D5216, 5217; Verde Island Passage and Batangas Bay, D5268, 5269; China Sea, D5301; east coast of Luzon, D5446; and north of Tawi Tawi, D5572.

At nearly all the stations several specimens occurred, and at one or two the species was frequent.

Truncatulina haidingerii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
			D5145...	6 04 30 N.; 120 59 30 E...	23	Co., s., sh.....	
			D5216...	12 52 00 N.; 123 23 30 E...	215	51.9	gn. m.....	
14814	U.S.N.M.	4	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.....	Few.
14818	U.S.N.M.	5	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Few.
14421	U.S.N.M.	4	D5269...	13 39 50 N.; 120 59 30 E...	220	fine s., p.....	Few.
14815	U.S.N.M.	1	D5446...	12 43 51 N.; 124 59 18 E...	300	gn. m.....	Rare.
			D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s.....	
			D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.....	
14422	U.S.N.M.	3	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co....	Few.

TRUNCATULINA AKNERIANA (d'Orbigny).

Plate 63, figs. 3a-c.

Rotalina akneriana d'ORBIGNY, Foram. Foss. Vienne, 1846, p. 156, pl. 8, figs. 13-15.

Truncatulina akneriana REUSS, Denkschr. Akad. Wiss. Wien, vol. 25, 1866, p. 160.

No. 6.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 663, pl. 94, figs. 8a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 35, pl. 16, fig. 3, text fig. 38.

Specimens referable to this species were obtained in material from but three stations, D5138, in 380 fathoms (700 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 43° F. (6.1° C.); D5268, 170 fathoms (311 meters), Verde Island Passage; and D5445, east coast of Luzon, 383 fathoms (699 meters), bottom temperature 44.3° F. (6.8° C.).

In the North Pacific material this species occurred only from the Hawaiian Islands. Apparently it is not common as far as the North Pacific is concerned.

Truncatulina akneriana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14423	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E...	380	43.0	s., Co.....	Rare.
14425	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Rare.
15352	U.S.N.M.	3						
14424	U.S.N.M.	2	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.....	Rare.

TRUNCATULINA UNGERIANA (d'Orbigny).

Plate 65, figs. 3a-c.

Rotalina ungeriana D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 157, pl. 8, figs. 16-18.*Planorbulina ungeriana* H. B. BRADY, Trans. Linn. Soc. London, vol. 24, 1864, p. 469, pl. 48, fig. 12.*Truncatulina ungeriana* REUSS, Denkschr. Akad. Wiss. Wien, vol. 25, 1865, p. 161.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 664, pl. 94, figs. 9a-d.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 36, pl. 17, fig. 2, text fig. 39.

This species seems to be well distributed both in the Archipelago and in the region to the south. The stations range in depth from 24 to 834 fathoms (44 to 525 meters), the average depth being 422 fathoms (775 meters). The bottom temperatures range from 41.2° to 73.8° F. (5.1° to 23.2° C.), with the average 57.5° F. (14.2° C.).




FIG. 12.—TRUNCATULINA UNGERIANA, VAR. ORNATA, NEW VARIETY. $\times 100$. a, DORSAL VIEW; b, VENTRAL VIEW. FROM STATION D5623.

The bottom temperatures range from 41.2° to 73.8° F. (5.1° to 23.2° C.), with the average 57.5° F. (14.2° C.).

Truncatulina ungeriana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14779	U.S.N.M.	7	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Few.
14780	U.S.N.M.	5	D5333...	12 26 30 N.; 120 37 45 E...	310	73.8	s.....	Few.
14781	U.S.N.M.	1	D5606...	0 16 28 N.; 121 33 30 E...	834	gn. m.....	Rare.
14420	U.S.N.M.	1	D5640...	4 27 00 S.; 122 55 40 E...	24	s. brk. sh....	Rare.
14910	U.S.N.M.	3	Nero 849.	15 16 00 N.; 121 40 00 E...	737	gn. m.....	Rare.

TRUNCATULINA UNGERIANA (d'Orbigny), var. ORNATA, new variety.

Description.—Variety differing from the typical in the thickened character of the sutures and the periphery, causing a raised ornamentation.

Distribution.—Type specimen (Cat. No. 15723, U.S.N.M.) from *Albatross* station D5623, between Gillolo and Makyan Islands, in 272 fathoms (498 meters).

Specimens also occur at the neighboring stations, D5622, in 275 fathoms (503 meters), and D5537, between Negros and Siquijor, in 254 fathoms (465 meters) and two others.

Truncatulina ungeriana, var. *ornata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15724	U.S.N.M.	3	D5112...	° ' " ° ' "		° F.		
			D5112...	13 48 22 N.; 120 47 25 E...	159	dk. gn. m....	Few.
			D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	
			D5622...	Makyan Island (N.E.), N., 66° W.	275	fine s., m....	
15723	U.S.N.M.	1	D5623...	0 16 30 N.; 127 30 00 E...	272	fine s., m....	Rare.
15725	U.S.N.M.	3	D5666...	2 54 30 S.; 118 47 00 E...	367	41.7	gy. s., m....	Few.

TRUNCATULINA TENERA H. B. Brady.

Plate 64, figs. 2a-c.

Truncatulina tenera H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol 9, 1884, p. 665, pl. 95, figs. 11a-c.—EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 402, pl. 16, figs. 45-47.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 37, pl. 15, fig. 2.

Single specimens have occurred at the following five stations: D5215, in 604 fathoms (1,105 meters), east of Masbate Island, bottom temperature 50.5° F. (10.2° C.); D5216, in 215 fathoms (393 meters); between Burias and Luzon, bottom temperature 51.9° F. (11° C.); D5348, in 375 fathoms (686 meters), Palawan Passage, bottom temperature 56.4° F. (13.5° C.); D5612, in 750 fathoms (1,370 meters), and D5613, in 752 fathoms (1,375 meters), both in the Gulf of Tomini, Celebes.

The specimens are typical. The only other North Pacific record for the species is off Japan.

Truncatulina tenera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14775	U.S.N.M.	1	D5215...	° ' " ° ' "		° F.		
			D5215...	12 31 30 N.; 123 35 24 E...	604	50.5	gn. m.	Rare.
			D5216...	12 52 00 N.; 123 23 30 E...	215	51.9	gn. m.	Rare.
14776	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.	Rare.
14777	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E...	750	Rare.
14778	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.	Rare.

TRUNCATULINA PRAECINCTA (Karrer).

Rotalia praecincta KARRER, Sitz. Akad. Wiss. Wien, vol. 58, 1868, p. 189, pl. 5, fig. 7.

Truncatulina praecincta H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 667, pl. 95, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 39, pl. 26, fig. 2, text fig. 42.

Brady recorded this species from a *Challenger* station, off the Philippines, in 95 fathoms (174 meters). The other North Pacific stations are

from near the Hawaiian Islands and off Japan at one station. In the Philippine material I have had the species from about a hundred stations, mostly within the Archipelago, but it has also occurred in the vicinity of Darvel Bay and Sibuko Bay, Borneo; Molucca Passage; between Gillolo and Kayoa Islands; Buton Strait; and Gulf of Boni.

The whole number of stations range in depth from 6 to 976 fathoms (11 to 1,785 meters), with the average depth 238 fathoms (436 meters). Bottom temperatures range, where given, from 40.1° to 76.3° F. (4.5° to 24.6° C.), with the average 52.7° F. (11.4° C.). This is from about half the stations, the bottom temperatures from the others, usually the shallow-water ones, not being given; therefore the average is probably much too low.

At some of the stations specimens are very abundant and attain a much larger size than in the colder waters. Some of the specimens measured over 2 mm. in diameter. This is the most common species of the genus in this area, and is evidently at home in warm, comparatively shallow waters.

Truncatulina praeincta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15732	U.S.N.M.	7	D5096...	14 20 23 N.; 120 34 15 E..	28	gy. m., s....	Few.
14786	U.S.N.M.	10+	D5113...	13 51 30 N.; 120 50 30 E..	159	dk. gn. m....	Common.
14787	U.S.N.M.	10	D5121...	13 27 20 N.; 121 17 45 E..	108	dk. gn. m....	Common.
14784	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Rare.
14785	U.S.N.M.	10+	D5261...	12 30 55 N.; 121 34 24 E..	145	s., m.....	Common.
14788	U.S.N.M.	5	D5278...	14 00 10 N.; 120 17 15 E..	102	59.6	fine. s., m., sh	Few.
14789	U.S.N.M.	2	D5381...	13 14 15 N.; 122 44 45 E..	88	co. s.....	Rare.
15733	U.S.N.M.	2	D5412...
14889	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.....	Rare.
14888	U.S.N.M.	10+	D5523...	8 48 44 N.; 123 27 35 E..	Common.
15734	U.S.N.M.	6	D5537...
14887	U.S.N.M.	10+	D5567...	5 48 00 N.; 120 33 45 E..	268	52	fine. s.....	Common.
14797	U.S.N.M.	2	D5579...	4 54 15 N.; 119 09 52 E..	175	55.3	co. s.....	Rare.
14820	U.S.N.M.	5	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s....	Few.
14796	U.S.N.M.	10+	D5642...	4 31 40 S.; 122 49 42 E..	37	gy. m.....	Common.
14795	U.S.N.M.	1	D5650...	4 33 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Rare.

TRUNCATULINA MARGARITIFERA H. B. Brady.

Plate 65, figs. 1a-c; plate 74, figs. 1a-c.

Truncatulina margaritifera H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 66; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 667, pl. 96, figs. 2a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 40, pl. 17, fig. 1, text fig. 43.

Brady records this species from 95 fathoms (174 meters) in material dredged by the *Challenger* in the Philippines. Except for two stations from the southern end of Japan these are the only North Pacific records.

It is interesting to note that the species has occurred frequently in the Philippine region, but all the recorded stations are from the Archipelago proper, except two stations off Darvel Bay, Borneo,

from each of which single specimens were obtained. The stations range in depth from 25 to 241 fathoms (46 to 441 meters), with the average 93 fathoms (167 meters) and the bottom temperatures were given from 51.3° to 76.3° F. (10.7° to 24.6° C.), with the average 57.1° F. (13.9° C.).

At two stations—D5133 in 38 fathoms (70 meters), Sulu Sea, off western Mindanao, and D5220 in 50 fathoms (91 meters), between Marinduque and Luzon—it was abundant. At other stations it was frequent or represented by few specimens.

Specimens seem to be rather larger than the measurements given for this species, running up to over 2 mm. in well-developed specimens.

This is a very strikingly ornamented species, of large size, and excellently figured by Brady. For that reason the lack of records is significant. Brady's original records were from off the Philippines in 95 fathoms (174 meters); off the New Hebrides, 125 fathoms (229 meters); and the coast of Korea, 10 to 50 fathoms (18 to 91 meters). In the North Pacific work I added two records for the species off Japan in 66 and 39 fathoms (121 and 71 meters). The Philippine records complete what is known of its distribution.

It is not recorded by Millett from the Malay Archipelago, by Dakin from Ceylon, by Heron-Allen and Earland from Kerimba, by any writers from Australia, nor by Egger from any of his material.

Its fine development in the Philippines shows that it is a characteristic species, and we must come to the conclusion that it is a species of comparatively limited distribution.

Truncatulina margaritifera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				" " " " " "		° F.		
14897	U.S.N.M.	4	D5100...	14 17 15 N.; 120 32 40 E..	35	gy. s.....	Few.
14899	U.S.N.M.	1	D5134...	6 44 45 N.; 121 48 00 E..	25	fine. s.....	Rare.
14898	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	gn. m., s....	Common.
14901	U.S.N.M.	7	D5210...	11 49 55 N.; 124 28 05 E..	50	76.3	fine. gy. s....	Common.
14900	U.S.N.M.	10+	D5213...	12 15 00 N.; 123 57 30 E..	80	s., m., sh...	Common.
14798	U.S.N.M.	10+	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m...	Common.
14799	U.S.N.M.	10+	D5313...	21 30 00 N.; 116 43 00 E..	150	53.6	s.....	Common.
14800					43	Co., s., m....	Rare.
14426	U.S.N.M.	1	D5338...	11 33 45 N.; 119 24 45 E..	43
14801	U.S.N.M.	3	D5443...	12 43 05 N.; 125 01 00 E..	241	51.3	co. s., sh....	Few.
14802	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co....	Rare.
15735	U.S.N.M.	3	D5315...	21 40 00 N.; 116 58 00 E..	143	54.4	s., sh.....	Few.

TRUNCATULINA CULTER (Parker and Jones).

Plate 62, figs. 4a-c.

Planorbulina culter PARKER and JONES, Philos. Trans., 1865, vol. 155, pl. 27, figs. 1, a, b.

Anomalina bengalensis SCHWAGER, Novara-Exped., geol. Theil, vol. 2, 1866, p. 259, pl. 7, fig. 3.

Truncatulina culter H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 668, pl. 96, figs. 3, a, b, c.

The only record for this species is a specimen from *Albatross* station D5638, off Bouro Island, in 517 fathoms (946 meters). It is very typical.

Truncatulina culter—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15729	U.S.N.M.	1	D5638...	° ' " ° ' " 3 47 15 S.; 126 23 40 E..	517	° F.	fne. gy. s....	Rare.

TRUNCATULINA HUMILIS H. B. Brady.

Truncatulina humilis H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 665, pl. 94, figs. 7, a, b, c.

There are a few specimens from *Albatross* station D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

These seem to be the same as the specimens figured by Brady, but are much larger than the measurements given by him.

Truncatulina humilis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15726	U.S.N.M.	3	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E....	494	° F. 41.2	fne. gy. s....	Few.

TRUNCATULINA ROSTRATA H. B. Brady.

Truncatulina rostrata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 65; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 668, pl. 94, figs. 6a-c.

There are three specimens of this rare species described by Brady from the *Challenger* material, off the northern shores of New Guinea in Humboldt Bay, 37 fathoms (68 meters), and near the Admiralty Islands in 16 to 25 fathoms (29 to 46 meters); also off Tongatabu, Friendly Islands, in 18 fathoms (33 meters). The Philippine material is from *Albatross* station D5218, between Burias and Luzon, in 20 fathoms (37 meters). These specimens are typical.

Truncatulina rostrata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15730	U.S.N.M.	3	D5218...	° ' " ° ' " 13 11 15 N.; 123 02 45 E....	20	° F.	crs. s.....	Rare.

Genus SIPHONINA Reuss, 1849.

SIPHONINA RETICULATA (Czjzek).

Plate 60, figs. 3a-c.

Rotalina reticulata CZJZEK, Haidinger's Nat. Abh., vol. 2, 1848, p. 145, pl. 13, figs. 7-9.

Siphonina reticulata BRONN, Lethaea Geognostica, ed. 3, 1853-1856, p. 227, pl. 35 (?), figs. 23a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 43, pl. 16, fig. 4; pl. 28, fig. 3, text fig. 48.

Truncatulina reticulata H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 669, pl. 96, figs. 5-8.

The only records for this species are D5120, 393 fathoms (719 meters), Verde Island Passage; D5217, 105 fathoms (193 meters), between Burias and Luzon; and D5242, 191 fathoms (350 meters), Pujada Bay. The only other North Pacific records I have had are from two *Albatross* stations off Japan.

Siphonina reticulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14822	U.S.N.M.	1	D5120...	° ' " ° ' "	393	° F. 43.7	gn. m., s....	Rare.
15715	U.S.N.M.	1	D5217...	13 45 30 N.; 120 30 15 E...	105	63.1	crs. gy. s....	Rare.
15716	U.S.N.M.	1	D5242...	13 00 20 N.; 123 14 15 E... 6 51 53 N.; 126 14 10 E...	191	64.1	sft. gy. m....	Rare.

Genus ANOMALINA d'Orbigny, 1826.

ANOMALINA ARIMINENSIS (d'Orbigny).

Planulina ariminensis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 280, pl. 14, figs. 1-3; Modèles, 1826, No. 49.

Anomalina ariminensis H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 674, pl. 93, figs. 10, 11.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 335, pl. 79, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 44, pl. 19, fig. 1, text figs. 49a-c.

Apparently this is a very rare species in this region, its occurrence having been noted but twice in all the material examined—D5192, in 32 fathoms (59 meters), off northern Cebu Island, and D5268, in 170 fathoms (311 meters), Verde Island Passage. The bottom temperature is not given in either case.

Anomalina ariminensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14414	U.S.N.M.	1	D5192...	° ' " ° ' "	32	° F.	gn. s.	Rare.
14413	U.S.N.M.	1	D5268...	11 09 15 N.; 123 50 00 E... 13 42 00 N.; 120 57 15 E...	170	p., s., sh	Rare.

ANOMALINA GROSSERUGOSA (Gümbel).

Plate 62, fig. 3.

Truncatulina grosserugosa GÜMBEL, Abh. kais. bay. Akad. Wiss., vol. 10, 1868, p. 660, pl. 2, fig. 104.

Anomalina grosserugosa H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 673, pl. 94, figs. 4, 5.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 335, pl. 78, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 45, pl. 20, fig. 1.

Although occurring at the largest number of stations in the region this species is not so abundant in point of numbers as is *A. polymorpha*. *A. grosserugosa* occurs at a large number of stations, nearly half of which are in the region to the south, including Sibuko Bay, Borneo; Gulf of Tomini, Celebes; between Gillolo and Makyan Islands; south of Bouro Island; Gulf of Boni; Flores Sea; and Macassar Strait. In the Archipelago proper the depths are somewhat less. Taking in all the stations in the whole area the range in depth is from 17 to 765 fathoms (31 to 1,399 meters), with the average 335 fathoms (613 meters). The bottom temperature ranges from 39.2° to 75.7° F. (4° to 24.2° C.), average 50.4° F. (10.2° C.).

Anomalina grosserugosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
12669	U.S.N.M.	1	D5172...	60 03 15 N.; 120 35 30 E...	318	fine. s., sh...	Rare.
12668	U.S.N.M.	5	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Few.
11994	U.S.N.M.	5	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s....	Few.
11993	U.S.N.M.	4	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
11977	U.S.N.M.	4	D5300...	20 31 00 N.; 115 49 00 E...	265	gy. m., s....	Few.
11979		1	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s....	Few.
15718	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Few.
15719	U.S.N.M.	1	D5660...	5 36 30 S.; 120 49 00 E...	692	39.2	gy. m., s....	Few.
15722	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Few.

ANOMALINA AMMONOIDES (Reuss).

Plate 61, figs. 1a-c.

Rosalina ammonoides REUSS, Geogn. Skizze Böhmen, vol. 2, 1884, p. 214; Verstein böhm. Kreide, vol. 1, 1845-1846, p. 36, pl. 8, fig. 53; pl. 13, fig. 66.

Planorbulina ammonoides JONES, Geologist, vol. 6, 1863, p. 294, pl. 15, figs. 7, 8.

Discorbis ammonoides REUSS, Sitz. Akad. Wiss. Wien, vol. 52, 1865, p. 456, No. 5.

Anomalina ammonoides H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 672, pl. 94, figs. 2, 3.—H. B. BRADY, PARKER, and JONES, Trans. Zool. Soc., vol. 12, 1888, p. 228, pl. 45, figs. 20-22.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 335, pl. 78, fig. 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 46, pl. 19, fig. 2.

Anomalina ammonoides has been noted in the material from 8 stations, all within the Archipelago proper. These stations range in depth from 254 to 805 fathoms (465 to 1,472 meters), the average depth being 387 fathoms (708 meters). Where given, the range in bottom temperature is from 41.2° to 56.4° F. (5.1° to 13.5° C.), the average being 49.3° F. (9.6° C.). At a single station—D5348

in Palawan Passage—it was frequent, occurring at the other stations as scattered or single specimens.

Anomalina ammonoides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11971	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s.	Rare.
11970	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Rare.
11968	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s.	Rare.
11969	U.S.N.M.	2						
11965	U.S.N.M.	1	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.	Rare.
11966	U.S.N.M.	4	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.	Frequent.
11967	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.	Rare.
11972	U.S.N.M.	2	D5526...	9 12 45 N.; 123 45 30 E..	505	52.3	gn. m., glob.	Rare.
11973	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.	Rare.

ANOMALINA POLYMORPHA Costa.

Plate 61, figs. 3a-c.

Anomalina polymorpha COSTA, Atti Accad. Pont., vol. 7, 1856, p. 252, pl. 21, figs. 7, 9.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 676, pl. 97, figs. 3-7.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 336, pl. 79, fig. 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 47, pl. 19, figs. 3, 4.

This is the commonest species of the genus in the Philippine region, occurring at nearly as many stations as *A. grosserugosa*, and at certain stations being one of the most abundant species in point of number of individuals. This is especially true of station D5236, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.), off the east coast of Mindanao, where it occurs very abundantly; and again at station D5586, in 347 fathoms (635 meters), bottom temperature 44.0° F. (6.6° C.) in Sibuko Bay, Borneo, where it was common. At other stations it occurred as scattered individuals. The stations range in depth from 49 to 730 fathoms (90 to 1,335 meters), the average depth being 352 fathoms (644 meters). The bottom temperatures range from 40.1° to 73.8° F. (4.5° to 23.2° C.), the average being 48.2° F. (8.9° C.).

Anomalina polymorpha—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11987	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.	Rare.
11981	U.S.N.M.	7	D5236...	8 50 45 N.; 126 26 45 E..	494	41.2	fine gy. s.	Abundant.
11989	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E..	201	50.4	dk. gy. s.	Rare.
11992	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.	Rare.
11990	U.S.N.M.	2	D5333...	12 26 30 N.; 120 37 45 E..	310	73.8	s.	Rare.
11982	U.S.N.M.	2	D5446...	12 43 51 N.; 124 59 18 E..	300	gn. m.	Rare.
11986	U.S.N.M.	2	D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.	Rare.
11991	U.S.N.M.	2	D5469...	13 36 48 N.; 123 38 24 E..	500	m.	Rare.
11983	U.S.N.M.	1	D5571...	5 30 45 N.; 120 07 57 E..	340	52.3	s., sh.	Rare.
11985	U.S.N.M.	1	D5574...	5 30 45 N.; 120 07 57 E..	340	Rare.
11980	U.S.N.M.	4	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.	Few.
11988	U.S.N.M.	1	D5590...	4 10 60 N.; 118 39 35 E..	310	44.3	gn. m., s.	Rare.
11984	U.S.N.M.	4	D5650...	4 35 45 S.; 121 29 00 E..	540	40.1	gn. m.	Few.

ANOMALINA POLYMORPHA Costa, var. CERVICORNIS Cushman.

Plate 62, figs. 1a, b.

Anomalina polymorpha, var. *cervicornis* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 662.

Description.—Test differing from the typical by having a marginal crown of short, stout, more or less branching or bifid spines on the angular margin of the chamber.

Distribution.—Type specimen (U.S.N.M. No. 14415) from *Albatross* station D5236, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.), Pacific Ocean, east coast of Mindanao.

This variety shows a rather remarkable departure from the usual ornamentation of the species.

Anomalina polymorpha, var. *cervicornis*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14415	U.S.N.M.	1	D5236...	8° 50' 45" N.; 126° 26' 52" E..	494	° F. 41.2	fne.gy.s....	Rare.

ANOMALINA POLYMORPHA Costa, var. SIPHONIFERA, Cushman.

Plate 62, figs. 2a, b.

Anomalina polymorpha, var. *siphonifera* CUSHMAN, Proc. U. S. Nat. Mus., vol. 51, 1917, p. 662.

Description.—Test differing from the typical by having short, tubular projections from the exterior, rather indefinitely placed; wall otherwise smooth.

Distribution.—Type specimen (U.S.N.M. No. 14416) from *Albatross* station D5236, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.), Pacific Ocean, east coast of Mindanao.

This variety is rather unique among this group, reminding one somewhat of the tubulations seen in *Textularia siphonifera*.

Anomalina polymorpha, var. *siphonifera*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14416	U.S.N.M.	2	D5236...	8° 50' 45" N.; 126° 26' 52" E..	494	° F. 41.2	fne.gy.s....	Rare.

ANOMALINA CORONATA Parker and Jones.

Plate 61, figs. 2a-c.

Anomalina coronata PARKER and JONES, Ann. Mag. Nat. Hist., ser. 2, vol. 19, 1857, p. 294, pl. 10, figs. 15, 16.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 675, pl. 97, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 47, pl. 18, fig. 5.

Specimens referable to this species occurred at the following stations: D5236, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.), east coast of Mindanao; D5318, in 340 fathoms (622 meters), China Sea vicinity of Formosa; D5445, in 383 fathoms (699 meters), bottom temperature 44.3° F. (6.8° C.); and D5446, in 300 fathoms (549 meters) east coast of Luzon.

Anomalina coronata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11974	U. S. N. M.	4	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s.....	Few.
11976	U. S. N. M.	2	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co.....	Rare.
11975	U. S. N. M.	2	D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s.....	Rare.
11995	U. S. N. M.	2	D5446...	12 43 51 N.; 124 59 18 E..	300	gn. m.....	Rare.

Genus PULVINULINA Parker and Jones, 1862.

PULVINULINA REPANDA (Fichtel and Moll).

Nautilus repandus FICHEL and MOLL, Test. Micr., 1798, p. 35, pl. 3, figs. a-d.

Rotalia repanda PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 5, 1860, p. 175, No. 25.

Pulvinulina repanda PARKER and JONES, in Carpenter, Parker, and Jones, Introd. Foram., 1862, p. 311.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 684, pl. 104, figs. 18a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 50, pl. 24, fig. 3.

It is interesting to note that in the *Challenger* Report this species is characterized as a shallow-water species. That seems to be true usually, but in the Philippines the only specimens which seem referable to this species are from rather deep water, and it was not found in the material from less than 100 fathoms (183 meters). The stations include the following localities: Verde Island Passage; between Burias and Luzon; Pacific Ocean, east coast of Mindanao; Gulf of Davao; north of Tawi Tawi; Sibuko Bay, Borneo; Gulf of Tomini, Celebes; Molucca Sea.

The range in depth of these stations is from 100 to 1,560 fathoms (183 to 2,853 meters); the average depth 461 fathoms (843 meters). Bottom temperatures are not given in all cases, but those that are

range from 43.0° to 63.1° F. (6.1° to 17.2° C.), average 50.8° F. (10.4° C.).

Pulvinulina repanda—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12367	U.S.N.M.	4	D5217...	13 00 20 N.; 123 14 15 E..	105	° F. 63.1	crs. gy. s.....	Few.
14448	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.....	Rare.
12366	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E..	750	Rare.
12365	U.S.N.M.	2	D5639...	3 54 50 S.; 123 27 20 E..	1,560	gy. m.....	Rare.
		1	D5238...	7 34 45 N.; 126 38 15 E..	380	43.0	gn. m.....	Rare.
		1	D5255...	7 03 00 N.; 125 39 00 E..	100	sft. m.....	Rare.
			D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Few.
			D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Few.

PULVINULINA CONCENTRICA Parker and Jones.

Plate 68, figs. 4a-c.

Pulvinulina concentrica (Parker and Jones, MS.) H. B. BRADY, Trans. Linn. Soc., London, vol. 24, 1864, p. 470, pl. 48, fig. 14.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 686, pl. 105, figs. 1a-b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 51, pl. 28, fig. 4.

This is a very characteristic species of wide distribution, but usually not occurring in any considerable numbers. The Philippine material is very typical. All of the stations are within the Archipelago proper. They include the following localities: East coast of Luzon; Verde Island Passage; between Burias and Luzon; off Romblon; and numerous stations in the Sulu Sea about Jolo and Tawi Tawi. The range of depth at the stations is from 23 to 500 fathoms (42 to 914 meters), the average depth 204 fathoms (373 meters). Bottom temperatures are given at but five of the numerous stations, as follows: 50.4°, 52.3°, 52.3°, 63.1°, and 75.7° F. (10.2°, 11.2°, 11.2°, 17.2°, and 24.2° C.).

Pulvinulina concentrica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12890	U.S.N.M.	1	D5145...	6 04 30 N.; 120 59 30 E..	23	° F.	co. s., sh.....	Rare.
			D5151...	5 24 40 N.; 120 27 15 E..	24	co. s., sh.....	Few.
12887	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E..	318	fine s., sh.....	Few.
12889	U.S.N.M.	8	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Common.
		1	D5217...	13 00 20 N.; 123 14 15 E..	105	63.1	crs. gy. s.....	Rare.
12888	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.....	Rare.
12891	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.....	Rare.
14438	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E..	303	52.3	co. s.....	Rare.
11623	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s.....	Rare.

PULVINULINA PUNCTULATA (d'Orbigny).

Plate 68, figs. 1a-c.

Rotalia punctulata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 273, No. 25; Modèles, 1826, No. 12.

Pulvinulina repanda, var. *punctulata* PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 394, pl. 14, figs. 12, 13.

Pulvinulina punctulata PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 3, vol. 16, 1865, p. 20, pl. 3, fig. 82.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 685, pl. 104, figs. 17a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 52, pl. 24, fig. 1, text figs. 54a-c.

Material referable to this species seems to be very rare in the Philippine region, occurring at but five stations: D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast Mindanao, bottom temperature 41.2° F. (5.1° C.); D5296, China Sea, off southern Luzon, 210 fathoms (384 meters); D5333, in 310 fathoms (567 meters), Mindoro Strait, bottom temperature 73.8° F. (23.2° C.); D5478, between Samar and Leyte, 57 fathoms (104 meters); and D5642, in 37 fathoms (68 meters), Buton Strait. Specimens were not frequent at any of the stations.

Pulvinulina punctulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12376	U.S.N.M.	4	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s.....	Few.
12375	U.S.N.M.	1	D5333...	12 26 30 N.; 120 37 45 E..	310	73.8	s.....	Rare.
11627	U.S.N.M.	1	D5642...	4 31 40 S.; 122 49 42 E..	37	gy. m.....	Rare.
15709	U.S.N.M.	8	D5478...	10 46 24 N.; 125 16 30 E..	57	sh.....	Few.
15710	U.S.N.M.	1	D5296...	13 40 09 N.; 120 57 45 E..	210	m., s.....	Rare.

PULVINULINA CONCAMERATA (Montagu).

Serpula concamerata MONTAGU, Test. Brit., Suppl., 1808, p. 160.

Rotalina concamerata WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 52, pl. 4, figs. 102, 103.

Pulvinulina repanda, var. *concamerata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 685, pl. 104, figs. 19a-c.

Pulvinulina concamerata CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 52, pl. 25, fig. 1.

This rather strikingly ornamented species has occurred at five stations: D5217, east of Masbate Island, 105 fathoms (193 meters); D5238, Pacific Ocean, east coast of Mindanao, 380 fathoms (700 meters); D5255, Gulf of Davao, 100 fathoms (183 meters); D5268, in 170 fathoms (311 meters), Verde Island Passage; and D5481 in 61 fathoms (112 meters), between Samar and Leyte, vicinity of Surigao Strait. Specimens were rare at all these stations.

Pulvinulina concamerata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12392	U.S.N.M.	1	D5217...	13 00 20 N.; 123 14 15 E..	105	° F. 63.1	crs. gy. s.....	Rare.
12368	U.S.N.M.	1	D5238...	7 34 45 N.; 126 38 15 E..	380	43.0	gn. m.....	Rare.
12369	U.S.N.M.	1	D5255...	7 03 00 N.; 125 39 00 E..	100	sft. m.....	Rare.
12370	U.S.N.M.	8	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Few.
12393			D5481...	10 27 30 N.; 125 17 10 E..	61	s., sh., Co....	Rare.

PULVINULINA AURICULA (Fichtel and Moll).

Plate 69, figs. 3a-c.

Nautilus auricula, var. *a*, FICHEL and MOLL, Test. Micr., 1803, p. 108, pl. 20, figs. a-c.

Pulvinulina auricula PARKER and JONES, Philos. Trans.; vol. 155, 1865, p. 393.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 688, pl. 106, figs. 5a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 53, pl. 22, fig. 1.

This species was recorded by Brady off the Philippines in 95 fathoms (174 meters). It is one of the characteristic species in the dredgings from below 100 fathoms (183 meters), the range being 32 to 1,105 fathoms (59 to 2,021 meters), but one station being less than 100 fathoms (183 meters), and the average 331 fathoms (606 meters). Bottom temperatures range from 43.3° to 59.6° F. (6.1° to 15.3° C.), average 50.0° F. (10° C.). The stations include a wide range of localities: China Sea off Formosa and eastern and southern Luzon; between Marinduque and Luzon; off northwestern Panay; off northern Cebu; between Cebu and Bohol; Sogod Bay, southern Leyte; Sulu Sea, off Jolo and Tawi Tawi; off eastern Palawan; and to the south at several stations in Sibuko Bay, Borneo and Macassar Strait. Specimens do not usually occur in very considerable numbers at any of the stations but are large and well-developed.

Pulvinulina auricula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12646	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E..	554	° F. 52.8	gy. s., m.....	Rare.
12886	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Rare.
12645	U.S.N.M.	1	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Rare.
15745								
12648	U.S.N.M.	5	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Few.
14920	U.S.N.M.	1	D5419...	9 58 30 N.; 123 46 00 E..	175	54.5	gn. m.....	Rare.
11630	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.....	Rare.
12649	U.S.N.M.	1	D5566...	5 52 12 N.; 120 31 00 E..	244	52.5	fine. s., sh....	Rare.
14451	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Rare.
12344	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.....	Rare.
14450	U.S.N.M.	10	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s....	Common.
12647	U.S.N.M.	1	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Rare.
15744	U.S.N.M.	3	D5112...	13 48 22 N.; 120 47 25 E..	177	52.4	dk. gn. m....	Few.
15756	U.S.N.M.	3	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Few.

PULVINULINA OBLONGA (Williamson).

Rotalina oblonga WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 51, pl. 4, figs. 98-100.

Pulvinulina oblonga H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 688, pl. 106, figs. 4a-c.

A specimen which it seems can be referred to this species was found at a single station in the area—D5220, between Marinduque and Luzon, 50 fathoms (91 meters). It was very rare at this station, however.

Pulvinulina oblonga—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15746	U.S.N.M.	1	D5220...	° ' " ° ' " 13 38 00 N.; 121 58 00 E..	50	° F.	sft. gn. m ...	Rare.

PULVINULINA SCABRA H. B. Brady.

Plate 58, figs. 3a, b.

Pulvinulina oblonga (WILLIAMSON), var. *scabra* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 689, pl. 106, figs. 8a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 53, pl. 27, fig. 5.

From an examination of a considerable series of this form it seems that it is worthy of specific rank. Typical *P. oblonga* seems to be rare as far as the material I have had shows, although Brady apparently had material from the Philippines in 95 fathoms (174 meters) which he referred to it. On the other hand, *P. scabra*, while not frequent, does appear at a number of stations and at some in very considerable numbers. The specimens are very constant in their characters and agree very closely with those of Brady.

Sidebottom refers to specimens from off Cebu,²² and speaks of the fact that all his specimens showed the earlier chambers to be flat or concave, while in the figures given by Brady they seem to be somewhat convex. In the Philippine material I have seen they are flattened and then become convex in later chambers. The ventral side in one form is very much like that given by Brady, except that it is rather more roughened, and is thereby easily distinguished from *Discorbis ventricosa*, which occasionally occurs at the same station. In the latter species also the ventral side is much more convex and the umbilical region has a large cavity. The scabrous condition in the two is also very different in its characters.

Pulvinulina scabra occurs at stations represented by the following localities: China Sea, off southern Luzon; between Marinduque and Luzon; off Romblon; and Sulu Sea, off western Mindanao. It was not found in the region to the south of the Archipelago.

²² Mem. and Proc. Manchester Lit. and Philos. Soc., vol. 53, No. 21, 1909, p. 6.

The stations range in depth from 10 to 50 fathoms (18 to 91 meters), the average being 34 fathoms (62 meters). It is therefore in this region a species of comparatively shallow water. Brady's records were 95 fathoms (174 meters), Philippines, and 17 to 155 fathoms (31 to 283 meters) off New Guinea. In the Atlantic, however, Brady records it from much deeper water—435 fathoms (796 meters), off Bermuda, and 1,000 fathoms (1,829 meters) west of the Azores. It will be interesting to see if these are really identical forms in all their characters with the more shallow-water Indo-Pacific species.

Pulvinulina scabra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11916	U.S.N.M.	8	D5096...	14 20 23 N.; 120 34 15 E..	28	°F.	gy. m., s., sh.	Common.
12374	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E..	10	Co.	Rare.
12894	U.S.N.M.	4	D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	gn. m., s.	Few.
14455	U.S.N.M.	5	D5179...	12 33 15 N.; 122 12 30 E..	37	75.7	hrd. s.	Few.
15703	U.S.N.M.	6	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.	Frequent.
15704	U.S.N.M.	10	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.	Frequent.

PULVINULINA PHILIPPINENSIS, new species.

Plate 58, figs. 2a-c.

Pulvinulina hauerii (part) H. B. BRADY (not d'Orbigny 1846), Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 690, pl. 106, fig. 7 (not fig. 6).

Description.—Test rounded, composed of about three coils, the outer one consisting of five chambers, very rapidly increasing in size, the last-formed chamber in the adult making up nearly half the volume of the test; periphery very broadly rounded; dorsal and ventral sides slightly rounded; sutures distinct, but not deep on the dorsal side, somewhat deeper on the ventral side; slightly umbilicate; wall smooth, very thin, punctate.

Diameter 0.4 to 0.6 mm.

Distribution.—Type specimen (Cat. No. 15760, U.S.N.M.) from *Albatross* station D5268, Verde Island Passage, 170 fathoms (311 meters).

It also occurs at the following stations: D5178, off Romblon, 78 fathoms (143 meters); D5381, Ragay Gulf, Luzon, 88 fathoms (161 meters); and D5565, between Jolo and Tawi, Tawi, 243 fathoms (445 meters).

Brady refers this species to *P. hauerii* d'Orbigny, but comparison with that figured by Brady shows that the two are very distinct. Brady records it from seven stations in the South Pacific. It is very constant in its characters and seems to be a well-defined species of this general region.

Pulvinulina philippinensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bot- tom tem- pera- ture.	Character of bottom.	Abundance.
						° F.		
15761	U.S.N.M.	3	D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.	Few.
15760	U.S.N.M.	2	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.	Few.
15712	U.S.N.M.	4	D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.	Few.
15762	U.S.N.M.	2	D5565...	5 51 42 N.; 120 30 3 E...	243	52.3	s., ptr., sh ..	Few.

PULVINULINA INDICA, new species.

Pulvinulina hauerii H. B. BRADY (part) (not d'Orbigny 1846), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 690, pl. 106, fig. 6 (not fig. 7).

Description.—Test somewhat longer than broad, composed of about two whorls, the last one having six chambers, usually gradually increasing in length as added; periphery broadly rounded, dorsal side convex, as is also the ventral side; sutures on both sides deeply depressed; ventral side not umbilicate, but the ventral portion of the last-formed chamber having an irregularly oval area of clear shell material; surface smooth, thin, and punctate; aperture a narrow arcuate slit at the edge of the peripheral side of the last-formed chamber.

Diameter 0.50 to 0.75 mm.

Distribution.—Type specimen (Cat. No. 15759, U.S.N.M.) from *Albatross* station D5219, between Marinduque and Luzon, in 530 fathoms (969 meters).

There are also specimens from D5134, Sulu Archipelago, near Basilan Island, in 34 fathoms (62 meters); D5143, off Jolo, in 19 fathoms (35 meters); D5145, at a neighboring station, in 23 fathoms (42 meters); and D5178, off Romblon, in 73 fathoms (133 meters).

This species, like the preceding, *P. philippinensis*, is confused by Brady with *P. hauerii* d'Orbigny. A comparison of the places at which these two occur shows that *P. indica* is found in much more shallow water than *P. philippinensis*, and the two do not occur together at the same stations; but they are probably both species characteristic of this general region. Brady's figures of both of these are very careful drawings and show well the characteristics of these two species.

Pulvinulina indica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bot- tom tem- pera- ture.	Character of bottom.	Abundance.
						° F.		
15788	U.S.N.M.	5	D5134...	6 44 12 N.; 121 46 55 E...	34	76.2	gy. s.	Few.
15754	U.S.N.M.	1	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.	Rare.
15755	U.S.N.M.	3	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh ..	Few.
15747	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	73	fine. s.	Rare.
15759	U.S.N.M.	1	D5219...	13 21 00 N.; 122 18 45 E...	530	50.8	gn. m.	Rare.

PULVINULINA BERTHELOTIANA d'Orbigny, var. SUBORNATA, new variety.

Plate 70, fig. 1.

Description.—Test differing from the typical in the character of the ornamentation, the sutures being thickened on the dorsal side, the periphery with a thickened rim, and basal angle of each chamber with a raised oval development.

Distribution.—Type specimen (Cat. No. 15355, U.S.N.M.) from *Albatross* station D5301, China Sea off Hongkong, in 208 fathoms (381 meters).

There are also specimens from nine other stations, ranging in depth from 6 to 208 fathoms (11 to 381 meters), the average being 106 fathoms (194 meters). Bottom temperatures are given at six stations, ranging from 50.5° to 76.3° F. (10.2° to 24.6° C.), the average temperature being 58.9° F. (14.9° C.).

All these stations are in the Archipelago.

Pulvinulina berthelotiana, var. *subornata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15355	U.S.N.M.	4	D5301...	20 37 00 N.; 115 43 00 E...	208	50.5	gy. m., s....	Few.
15353	U.S.N.M.	8	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Common.
			D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fne. gy. s....	Few.
			D5217...	13 00 20 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Frequent.
			D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m....	Few.
			D5331...	15 36 45 N.; 119 47 45 E...	178	54.7	s., sh., m....	Few.
			D5338...	11 33 45 N.; 119 24 45 E...	43	Co., s., m....	Frequent.
			D5344...	10 50 40 N.; 119 22 32 E...	6	m.....	Few.
			D5398...	11 35 12 N.; 124 13 48 E...	114	gn. m.....	Few.
			D5412...	10 09 15 N.; 123 52 00 E...	162	54.8	gn. m.....	Few.

PULVINULINA MENARDII (d'Orbigny).

Plate 66, figs. 1a-c.

Rotalia menardii D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 273, No. 26; Modèles, 1826, No. 10.

Pulvinulina menardii OWEN, Journ. Linn. Soc., London, vol. 9, 1867, p. 148, pl. 5, fig. 6.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, 1884, p. 690, pl. 103, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 54, pl. 22, fig. 2.

This common species of Globigerina-ooze is abundant, being found at more than 100 of the stations examined. As might be expected the species is most common around the outer coasts of the Archipelago, but is also common in the Sulu Sea and found less abundantly in the more protected waters among the islands. To the southward it occurs at nearly all the stations; Darvel and Sibuko Bays, Borneo; Gulf of Tomini, Celebes; Molucca Passage; between Gillolo and Makyan Islands; between Gillolo and Kayoa; south of Patiente Strait; off Bouro Island; Molucca Sea; Buton Strait; Gulf of Boni; Flores Sea; and Macassar Strait.

The depths of these stations range from 15 to 1,560 fathoms (27 to 2,853 meters) with the average 269 fathoms (493 meters), although depth probably means little with a more or less pelagic species. The bottom temperatures are given at but 56 stations, ranging from 36.3° to 75.7° F. (2.3° to 24.2° C.), with the average 51.1° F. (10.6° C.).

Pulvinulina menardii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12371	U.S.N.M.	10+	D5126...	10 34 45 N.: 121 47 30 E..	742	49.5	sft. gn. m...	Common.
12373	U.S.N.M.	8	D5201...	10 10 00 N.: 125 04 15 E..	554	52.8	gy. s., m....	Few.
12324	U.S.N.M.	7	D5236...	8 50 45 N.: 126 26 52 E..	494	41.2	fine. gy. s....	Common.
12895	U.S.N.M.	6	D5281...	13 52 45 N.: 120 25 00 E..	201	50.4	dk. gy. s....	Few.
12331	U.S.N.M.	5	D5300...	20 31 00 N.: 115 49 00 E..	265	gy. m., s....	Few.
12330	U.S.N.M.	1	D5338...	11 33 45 N.: 119 24 25 E..	43	Co., s., m....	Common.
14446	U.S.N.M.	10+	D5381...	13 14 15 N.: 122 44 45 E..	88	co. s.....	Common.
12329	U.S.N.M.	7	D5571...	5 30 45 N.: 120 07 57 E..	340	52.3	s., sh.....	Few.
12332	U.S.N.M.	4	D5627...	0 06 00 N.: 127 26 00 E..	22	m.....	Common.
11629	U.S.N.M.	9	D5630...	0 56 30 S.: 128 05 00 E..	569	co. s., m....	Common.
14445	U.S.N.M.	10+	D5637...	3 53 20 S.: 126 48 00 E..	700	gy. m.....	Common.
12327	U.S.N.M.	10	D5639...	3 54 50 S.: 123 27 20 E..	1,560	gy. m.....	Common.
12372	U.S.N.M.	6	D5654...	3 42 00 S.: 120 45 50 E..	805	38.3	Co., s.....	Common.
12333	U.S.N.M.	1	D5446...	12 43 51 N.: 124 59 18 E..	300	gn. m.....	Rare.
14447	U.S.N.M.	8	D5660...	5 36 30 S.: 120 49 00 E..	692	39.2	gy. m., s....	Rare.
12334	U.S.N.M.	1	D5666...	2 54 30 S.: 118 47 00 E..	272	47.5	gn. m.....	Rare.
12335	U.S.N.M.	1	D5110...	13 59 20 S.: 120 75 45 E..	135	59.0	dk. gy. m....	Few.
12336	U.S.N.M.	1	D5112...	13 48 22 N.: 120 47 25 E..	177	52.4	dk. gn. m....	Few.
12337	U.S.N.M.	1	D5113...	13 51 30 N.: 120 50 30 E..	159	dk. gn. m....	Few.
12338	U.S.N.M.	1	D5114...	13 36 11 N.: 120 45 26 E..	340	fine. s.....	Few.
12339	U.S.N.M.	1	D5115...	13 37 11 N.: 120 43 40 E..	340	Frequent.
15743	U.S.N.M.	1	D5120...	13 45 30 N.: 120 30 15 E..	393	43.7	gn. m., s....	Few.
15748	U.S.N.M.	1	D5134...	6 44 45 N.: 121 48 00 E..	108	dk. gn. m....	Few.
15757	U.S.N.M.	2	D5153...	5 18 10 N.: 120 02 55 E..	25	fine. s.....	Few.
			D5172...	6 03 15 N.: 120 35 30 E..	49	co. s., sh....	Few.
			D5178...	12 43 00 N.: 122 06 15 E..	318	fine. s., sh....	Frequent.
			D5179...	12 38 15 N.: 122 12 30 E..	78	fine. s.....	Frequent.
			D5184...	10 18 30 N.: 122 23 30 E..	37	75.7	hrd. s.....	Frequent.
			D5185...	10 05 45 N.: 122 18 30 E..	565	49.8	gn. m.....	Few.
			D5187...	9 16 45 N.: 123 21 15 E..	638	49.8	gn. m.....	Frequent.
			D5191...	9 16 45 N.: 123 21 15 E..	225	53.6	sft. gn. m....	Few.
			D5203...	10 29 45 N.: 123 31 15 E..	258	62.8	gn. m.....	Common.
			D5206...	9 58 00 N.: 125 07 40 E..	775	52.9	gn. m.....	Few.
			D5209...	11 31 40 N.: 124 42 40 E..	32	gn. m.....	Frequent.
			D5211...	11 51 35 N.: 124 14 00 E..	155	56.6	gn. m., s....	Few.
			D5214...	12 25 18 N.: 123 37 15 E..	218	51.4	gn. m., s....	Few.
			D5215...	12 31 30 N.: 123 35 24 E..	604	50.5	gn. m., s....	Few.
			D5216...	12 52 00 N.: 123 23 30 E..	215	51.9	gn. m., s....	Frequent.
			D5217...	13 00 20 N.: 123 14 15 E..	105	63.1	crs. gy. s....	Frequent.
			D5218...	13 11 15 N.: 123 02 45 E..	20	crs. s.....	Frequent.
			D5219...	13 21 00 N.: 122 18 45 E..	530	50.8	gn. m.....	Frequent.
			D5220...	13 38 00 N.: 121 58 00 E..	50	sft. gn. m....	Few.
			D5222...	13 38 30 N.: 121 42 15 E..	195	52.8	gn. m.....	Common.
			D5243...	6 50 55 N.: 126 14 35 E..	218	63.6	gy. m.....	Common.
			D5247...	7 02 00 N.: 125 33 45 E..	135	m.....	Few.
			D5259...	11 57 30 N.: 121 42 15 E..	312	49.3	gy. m.....	Frequent.
			D5260...	12 25 35 N.: 121 31 35 E..	234	51.4	gn. m., s....	Frequent.
			D5261...	12 30 55 N.: 121 30 50 E..	145	s., m.....	Frequent.
			D5268...	13 42 00 N.: 120 57 15 E..	170	s., p.....	Frequent.
			D5272...	14 00 00 N.: 120 22 30 E..	118	57.4	m., sh.....	Few.
			D5276...	13 49 15 N.: 120 14 45 E..	18	sh., p., s....	Frequent.
			D5282...	13 53 00 N.: 120 26 45 E..	248	47.4	dk. gy. s....	Frequent.
			D5313...	21 30 00 N.: 116 43 00 E..	150	53.6	s.....	Few.
			D5315...	21 40 00 N.: 116 58 00 E..	148	54.4	s., sh.....	Few.
			D5318...	21 32 00 N.: 117 46 00 E..	340	s., br. Co....	Few.
			D5331...	15 36 45 N.: 119 47 45 E..	178	54.7	s., sh., m....	Few.
			D5333...	12 26 30 N.: 120 37 45 E..	310	73.8	s.....	Few.
			D5342...	10 56 55 N.: 119 17 24 E..	14-15	gy. m.....	Few.
			D5349...	10 54 00 N.: 118 26 20 E..	730	40.6	Co., s.....	Frequent.
			D5369...	13 48 00 N.: 121 43 00 E..	106	bk. s.....	Few.
			D5370...	13 44 15 N.: 121 42 30 E..	159	54.3	sft. m.....	Few.
			D5371...	13 49 40 N.: 121 40 15 E..	83	gn. m.....	Few.
			D5374...	13 46 45 N.: 121 35 08 E..	190	gy. m.....	Frequent.
			D5388...	12 51 30 N.: 123 26 15 E..	226	51.4	sft. gn. m....	Few.

Pulvinulina menardii—Material examined—Continued.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "				
			D5394...	12 00 30 N.: 124 05 36 E...	153	gn. m.	Few.
			D5419...	9 58 30 N.: 123 46 00 E...	175	54.5	gn. m.	Few.
			D5428...	9 13 00 N.: 118 51 15 E...	1,105	49.7	gy. m.	Several.
			D5429...	9 41 30 N.: 118 80 22 E...	766	gn. m.	Few.
			D5430...	9 49 40 N.: 119 03 20 E...	464	50.0	glob. oz.	Few.
	U.S.N.M.	1	D5439...	15 58 15 N.: 119 40 20 E...	940	36.7	gn. m.	Few.
			D5445...	12 44 42 N.: 124 59 50 E...	383	44.3	gn. m., s.	Few.
			D5449...	13 21 36 N.: 124 00 30 E...	730	40.6	Co., s.	Few.
	U.S.N.M.	1	D5459...	13 10 21 N.: 123 59 54 E...	201	Few.
	U.S.N.M.	1	D5467...	13 25 27 N.: 123 37 18 E...	480	gy. m.	Few.
	U.S.N.M.	1	D5468...	13 35 39 N.: 123 40 28 E...	569	gn. m.	Few.
			D5481...	10 27 30 N.: 125 17 10 E...	61	s., sh., Co.	Few.
	U.S.N.M.	1	D5493...	9 04 00 N.: 125 20 00 E...	478	52.1	gn. m.	Few.
			D5523...	8 48 44 N.: 123 27 35 E...	Frequent.
			D5526...	9 12 45 N.: 123 45 30 E...	305	52.3	gn. m.	Frequent.
			D5529...	9 23 45 N.: 123 39 30 E...	441	53.0	gy. m.	Frequent.
			D5538...	9 08 15 N.: 123 23 20 E...	256	53.3	gn. m., s.	Frequent.
			D5542...	8 48 30 N.: 123 35 30 E...	200	54.3	fine. s.	Few.
			D5543...	8 47 15 N.: 123 35 00 E...	162	54.6	s.	Frequent.
			D5546...	6 06 48 N.: 121 20 32 E...	138	68.3	fine. co. s.	Few.
			D5551...	5 54 48 N.: 120 44 24 E...	193	53.3	fine. s.	Few.
			D5580...	4 52 45 N.: 119 08 45 E...	162	55.8	br. s., Co.	Few.
			D5590...	4 10 50 N.: 118 39 35 E...	310	44.3	gn. m., s.	Few.
			D5601...	1 13 10 N.: 125 17 05 E...	765	s., glob.	Common.
			D5609...	0 11 00 N.: 121 16 00 E...	1,092	36.3	gn. m.	Few.
			D5606...	0 16 28 N.: 121 33 30 E...	834	gn. m.	Common.
			D5612...	0 38 00 S.: 121 45 40 E...	750	Common.
			D5614...	0 31 00 N.: 125 58 45 E...	1,100	gy. m., s.	Common.
	U.S.N.M.	1	D5619...	0 35 00 N.: 127 14 40 E...	435	fine. gy. s.	Few.
			D5621...	0 15 00 N.: 127 24 35 E...	298	gy. and bk. s.	Few.
			D5622...	Makyan Island (N. E.), N. 66° W.	275	gy. m.	Frequent.
			D5623...	0 16 30 N.: 127 30 00 E...	272	fine. s., m.	Frequent.
	U.S.N.M.	1	D5625...	0 07 00 N.: 127 28 00 E...	230	gy. m., s.	Rare.
			D5632...	1 00 00 S.: 127 60 00 E...	845	Frequent.
			D5638...	3 47 15 S.: 126 23 40 E...	517	fine. gy. s.	Frequent.
			D5640...	4 27 00 S.: 122 55 40 E...	24	Few.
	U.S.N.M.	1	D5648...	5 35 00 S.: 122 20 00 E...	559	39.2	m.	Few.
	U.S.N.M.	1	D5650...	4 53 45 S.: 121 20 00 E...	540	40.1	gn. m.	Few.
	U.S.N.M.	1	D5652...	4 35 00 S.: 121 23 06 E...	525	41.2	gn. m.	Few.
			D5665...	4 27 00 S.: 118 44 00 E...	1,008	m.	Few.

PULVINULINA MENARDII (d'Orbigny), var. FIMBRIATA H. B. Brady.

Plate 66, figs. 2a-c.

Pulvinulina menardii (D'ORBIGNY), var. *fimbriata* H. B. BRADY, Rep. Voy. *Chalenger*, Zoology, vol. 9, 1884, p. 691, pl. 103, figs. 3a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 55.

This variety seems to be rare in the North Pacific, and is equally so in the region covered. It does not seem to occur in the Archipelago, but rather poorly developed specimens were taken at D5638 in 517 fathoms (946 meters), off Bouro Island, and D5639, in 1,560 fathoms (2,853 meters), Molucca Sea; but in neither case are the specimens at all typical, but tend toward the variety.

Pulvinulina menardii, var. *fimbriata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "				
12326	U.S.N.M.	1	D5638...	3 47 15 S.; 126 23 40 E....	517	°F.	fine. gy. s.	Rare.
		1	D5639...	3 54 50 S.; 123 27 20 E....	1,560	gy. m.	Rare.

PULVINULINA LATERALIS (Terquem).

Plate 69, figs. 2a-c.

Rosalina lateralis TERQUEM, Mém. Soc. Géol. France, ser. 3, vol. 1; Mém. 3, 1878, p. 25, pl. 2, figs. 11a-c.

Pulvinulina lateralis H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 689, pl. 106, figs. 2, 3.—EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 413, pl. 18, figs. 48-50.—MILLETT, Journ. Roy. Micr. Soc., 1904, p. 497.—CUSHMAN, Proc. Boston Soc. Nat. Hist., vol. 34, 1908, p. 31, pl. 5, figs. 11, 12.—SIDEBOTTOM, Mem. Proc. Manchester Lit. and Philos. Soc., vol. 53, No. 21, 1909, p. 5, pl. 2, fig. 6, pl. 13, figs. 1, 2 (?).—HERON-ALLEN and EARLAND, Trans. Zool. London, vol. 20, pt. 17, 1915, p. 714, pl. 53, figs. 6-11.

Description.—Test usually somewhat ovate, later chambers becoming rapidly larger and longer on the ventral side, the last-formed chamber often covering at least half the surface of the test; dorsally sutures somewhat limbate, surface generally flattened, ventrally sutures simple, depressed, the surface tumid, finely punctate except the last-formed chamber, which is covered with large circular pores, which are often bordered with a somewhat thicker rim of shell material, these pores serving as the aperture.

Test up to 2.5 mm. in diameter.

The records for *P. lateralis* are rather scattered, largely limited to the Indo-Pacific region. It is, however, recorded from numerous areas by Sidebottom and Heron-Allen and Earland had it from the Kerimba Archipelago off the east coast of Africa.

In the Philippine material it has occurred at 11 stations, ranging in depth from 12 to 162 fathoms (22 to 297 meters), average depth 44 fathoms (80 meters). Six of the stations are in the Sulu Sea off Jolo and Tawi Tawi; the others, off Romblon, two stations; between Burias and Luzon; between Marinduque and Luzon; and off Darvel Bay, Borneo. At the two stations off Romblon specimens were fairly frequent in their occurrence; at most other stations rather few.

Pulvinulina lateralis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12377	U. S. N. M.	3	D5133...	6 06 00 N.; 120 58 50 E...	19	s., co.....	Few.
12379	U. S. N. M.	4	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Few.
12378	U. S. N. M.	5	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Few.
11624	U. S. N. M.	1	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co.....	Few.
	U. S. N. M.		D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.....	Few.
	U. S. N. M.		D5148...	5 35 40 N.; 120 47 30 E...	17	co. s.....	Few.
	U. S. N. M.		D5151...	5 24 40 N.; 120 27 15 E...	24	co. s., sh.....	Few.
	U. S. N. M.		D5153...	5 18 10 N.; 120 02 55 E...	49	co. s., sh.....	Few.
	U. S. N. M.	1	D5160...	5 12 40 N.; 119 55 10 E...	12	s.....	Rare.
	U. S. N. M.		D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.....	Frequent.
	U. S. N. M.	1	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m.....	Rare.

PULVINULINA TUMIDA H. B. Brady.

Plate 66, figs. 3a-c.

Pulvinulina menardii D'ORBIGNY, var. *tumida* H. B. BRADY, Geol. Mag., vol. 4, 1877, p. 294; Quart. Journ. Micr. Sci., vol. 19, 1879, p. 80.

Pulvinulina tumida H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 692, pl. 103, figs. 4-6.—EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 414, pl. 17, figs. 4-6, 35-37, 44.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 329, pl. 73, fig. 5.—MILLETT, Journ. Roy. Micr. Soc., 1904, p. 499.—CUSHMAN, Proc. Boston Soc. Nat. Hist., vol. 34, 1908, p. 31; Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 56, pl. 22, fig. 3.

This species, characteristic of *Globigerina*-ooze, has occurred at 26 stations in the area. Most of these are in comparatively deep water. About half of these stations are in the Archipelago proper; the others to the south in the following localities: Gulf of Tomini, Celebes; Molucca Passage, between Gillolo and Makyan; between Gillolo and Kayoa; south of Patiente Strait; south of Bouro Island; Gulf of Boni; Macassar Strait; and Molucca Sea. These stations range in depth from 32 to 1,560 fathoms (59 to 2,853 meters), the average being 437 fathoms (800 meters); bottom temperatures are given at but 10 stations, ranging from 38.2° to 63.6° F. (3.4° to 17.5° C.), the average temperature being 50.3° F. (10.1° C.).

Pulvinulina tumida—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms	Bottom temperature.	Character of bottom.	Abundance.
12350	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318	fne. s., sh.	Rare.
			D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Few.
12349	U.S.N.M.	9	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy., s.	Common.
14441	U.S.N.M.	2	D5243...	6 50 55 N.; 126 14 35 E...	218	63.6	gy. m.	Rare.
11628	U.S.N.M.	5	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.	Few.
12348	} U.S.N.M.	10+	D5639...	3 54 20 S.; 123 27 20 E...	1,560	gy. m.	Common.
14442								
15711	U.S.N.M.	1	D5671...	1 05 00 S.; 118 56 00 E...	960	38.2	gy. m.	Rare.
			D5217...	13 00 20 N.; 123 14 15 E...	105	63.1	crs. gy. s.	Few.
	} U.S.N.M.	1	D5241...	6 50 45 N.; 126 14 38 E...	215	sit. gy. m.	Few.
			D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.	Few.
			D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.	Few.
			D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.	Few.
			D5446...	12 43 51 N.; 124 59 18 E...	300	gn. m.	Few.
	U.S.N.M.	1	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh. g.	Few.
	U.S.N.M.	1	D5542...	8 48 30 N.; 123 35 30 E...	200	54.3	fne. s., brk. sh.	Few.
	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.	Few.
	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.	Few.
			D5612...	0 38 00 S.; 121 45 40 E...	750	Few.
	U.S.N.M.	1	D5619...	0 35 00 N.; 127 14 40 E...	435	fne. gy. s., m.	Few.
			D5621...	0 15 00 N.; 127 24 35 E...	298	gy. and bk. s.	Few.
			D5626...	0 07 30 N.; 127 29 00 E...	265	gy. m., fne. s.	Few.
	U.S.N.M.	1	D5632...	1 00 00 S.; 127 50 00 E...	845	Few.
			D5638...	3 47 15 S.; 126 23 40 E...	517	fne. gy. s.	Few.
			D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Few.
			D5665...	4 27 00 S.; 118 44 00 E...	1,008	m.	Few.
			D5543...	8 47 15 N.; 123 35 00 E...	162	54.5	s.	Few.

PULVINULINA CANARIENSIS (d'Orbigny).

Plate 67, figs. 1a-c.

Rotalina canariensis D'ORBIGNY in Barker, Webb, and Berthelot, Hist. Nat. Îles Canaries, vol. 2, pt. 2, 1839, "Foraminifères," p. 130, pl. 1, figs. 34-36.

Pulvinulina canariensis OWEN, Journ. Linn. Soc. London, vol. 9, 1867, p. 148, pl. 5, fig. 21.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 692, pl. 103, figs. 8-10.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 56, pl. 23, fig. 1, text figs. 55a-c.

Rather strangely this species has occurred rarely in the region to the southward, but not at all in the Archipelago as far as the collections show. Material is typical, but found only as single specimens or in few numbers. It was found at the following stations: D5606, in 834 fathoms (1,525 meters), Gulf of Tomini, Celebes; D5621, in 298 fathoms (545 meters), between Gillolo and Makyan Islands; D5638, in 517 fathoms (946 meters), off Bouro Island; and D5650, in 540 fathoms (988 meters), Gulf of Boni.

Pulvinulina canariensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14433	U.S.N.M.	1	D5606...	° ' " ° ' "	834	° F.	gn. m.	Rare.
14432	U.S.N.M.	1	D5621...	0 16 28 N.; 121 33 30 E..	298	gy. and bk. s.	Rare.
			D5638...	0 15 00 N.; 127 24 35 E..	517	fine. gy. s.	
			D5650...	3 47 15 S.; 126 23 40 E..	540	40.1	gn. m.	
				4 53 45 S.; 121 29 00 E..				

PULVINULINA CRASSA (d'Orbigny).

Plate 67, figs. 3a-c.

Rotalina crassa D'ORBIGNY, Mém. Soc. Géol. France, vol. 4, 1840, p. 32, pl. 3, figs. 7-8.

Pulvinulina crassa OWEN, Journ. Linn. Soc. London, vol. 9, 1867, p. 148, pl. 5, figs. 8(?), 9.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 694, pl. 103, figs. 11-12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 58, pl. 27, fig. 1.

While this species has been taken as pelagic, the main records of its occurrence are from fairly deep water. It has been noted as single specimens at but two stations: D5201, in 554 fathoms (1,012 meters), Sogod Bay, southern Leyte; and D5580, in 162 fathoms (297 meters), Darvel Bay, Borneo.

Pulvinulina crassa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12694	U.S.N.M.	1	D5201...	° ' " ° ' "	554	° F.	gy. s. and m.	Rare.
11622	U.S.N.M.	1	D5580...	10 10 00 N.; 125 04 15 E..	162	52.8	br. s., Co....	Rare.
				4 52 45 N.; 119 06 45 E..		55.8		

PULVINULINA TRUNCATULINOIDES (d'Orbigny).

Plate 67, figs. 2a-c.

Rotalina truncatulinoides D'ORBIGNY in Barker, Webb, and Berthelot, Hist. Nat. Îles Canaries, vol. 2, pt. 2, 1839, "Foraminifères," p. 132, pl. 2, figs. 25-27.

Pulvinulina truncatulinoides PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 398, pl. 16, figs. 41-43.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 59, pl. 23, fig. 4, text figs. 57a-c.

Rotalina micheliniana D'ORBIGNY, Mém. Soc. Géol. France, vol. 4, 1840, p. 31, pl. 3, figs. 1-3.

Pulvinulina micheliniana OWEN, Journ. Linn. Soc. London, vol. 9, 1867, p. 148, pl. 5, fig. 17.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 694, pl. 104, figs. 1-2.

Although fairly well distributed the species was recorded as occurring at but one station outside the Philippines proper—at D5665, in 1,008 fathoms (1,844 meters), Macassar Strait. In the Archipelago it occurs on both the southwestern and eastern coasts of Luzon; Verde Island Passage; east coast of Mindanao and Gulf of Davao; Sulu Sea and Palawan Passage; thus not being recorded except in waters either on the outer side of the Archipelago or in the direct connections with interior waters. Stations range in depth from 49 to 1,008 fathoms (90 to 1,844 meters), with the average 397 fathoms (727 meters). Bottom temperatures range from 40.6° to 56.4° F. (4.7° to 13.5° C.), the average 46.5° F. (8.0° C.).

Pulvinulina truncatulinoides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12345	U.S.N.M.	2	D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s.....	Rare.
12344	U.S.N.M.	4	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.....	Rare.
11915	U.S.N.M.	2	D5247...	7 02 00 N.; 125 38 45 E..	135	m.	Rare.
11625 12338	U.S.N.M.	10+	D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s.....	Common.
12346	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	Co., s.....	Rare.
12347	U.S.N.M.	4	D5349...	10 54 00 N.; 118 26 20 E..	730	40.6	Co., s.....	Rare.
14444	U.S.N.M.	3	D5446...	12 43 51 N.; 124 59 18 E..	300	gn. m.	Rare.
			D5665...	4 27 00 S.; 118 44 00 E..	1,008	m.	Rare.
			D5153...	5 18 10 N.; 120 02 55 E..	49	co. s., sh....	Few.
			D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s.....	Few.

PULVINULINA UMBONATA (Reuss).

Plate 71, figs. 1a-c.

Rotalina umbonata REUSS, Zeitschr. deutsch. geol. Ges., vol. 3, 1851, p. 75, pl. 5, figs. 35a-c.

Pulvinulina umbonata REUSS, Denkschr. Akad. Wiss. Wien, vol. 25, 1866, p. 206.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 695, pl. 105, figs. 2a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 60, pl. 27, fig. 2.

This species seems to be rare in the region, having been noted at but three stations, all near Luzon, as follows: D5219, in 530 fathoms (969 meters), between Marinduque and Luzon; D5468, in 569 fathoms

(1,040 meters); and D5469, in 500 fathoms (914 meters), both on the east coast of Luzon.

Pulvinulina umbonata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12355	U.S.N.M.	1	D5219...	° ' " ° ' "	530	° F. 50.8	gn. m.	Rare.
12354	U.S.N.M.	2	D5468...	13 21 00 N.; 122 18 45 E..	569	gn. m.	Rare.
14453	U.S.N.M.	1	D5469...	13 35 39 N.; 123 40 28 E..	500	gn. m.	Rare.
				13 36 48 N.; 123 38 24 E..				

PULVINULINA EXIGUA H. B. Brady.

Plate 68, figs. 3a-c.

Pulvinulina exigua H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 696, pl. 103, figs. 13, 14.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 60, pl. 23, fig. 5.

This seems to be a rare species for the region as far as the material shows, possibly owing to the fact that it seems to be more characteristic of deeper waters. The only station at which material was recorded is D5349 in 730 fathoms (1,335 meters), Palawan Passage, bottom temperature 40.6° F. (4.7° C.). In material from this station several typical specimens were found.

Pulvinulina exigua—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14431	U.S.N.M.	3	D5349...	° ' " ° ' "	730	° F. 40.6	Co., s.	Rare.
				10 54 00 N.; 118 26 20 E..				

PULVINULINA PAUPERATA (Parker and Jones).

Plate 68, fig. 2.

Pulvinulina repanda, var. *menardii*, subvariety *pauperata* PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 395, pl. 16, figs. 50, 51.

Pulvinulina pauperata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 696, pl. 104, figs. 3-11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 61, pl. 23, figs. 2, 3, text figs. 58a, b.

Although this species was found at 10 stations, but two of these are in the Philippines, or rather on their borders—D5236 in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, and D5439, in 940 fathoms (1,719 meters), off the west coast of Luzon. To the southward it has occurred at three stations off Sibuko Bay, Borneo; in the Gulf of Tomini, Celebes; south of Patiente Strait; Molucca Sea; and Macassar Strait. The depths range from 260 to 1,560 fathoms (476 to 2,853 meters), with the average 699 fathoms (1,278 meters). Bottom temperatures are given at six stations, rang-

ing from 36.7° F to 44.0° F. (2.6° C. to 6.6° C.), with the average 40.3° F. (4.6° C.). From the known records of this species it is a deep-water form, and would hardly be expected inside the Archipelago unless in the Sulu Sea.

At D5236 a peculiar modification of the species occurs, where the broad carina instead of being thin and translucent, as in typical specimens, often becomes encrusted with fine sand grains, even to such an extent that it becomes in some cases thicker than the central chambers. This character was not noted at any of the other stations.

Pulvinulina pauperata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15714	} U.S.N.M.	5	} D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s....	Few.
12336		6						
11626	U.S.N.M.	3	D5439...	15 58 15 N.; 119 40 20 E..	940	36.7	gn. m.	Few.
14434	U.S.N.M.	5	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.	Few.
12337	U.S.N.M.	2	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.	Rare.
11907	U.S.N.M.	4	D5591...	4 11 48 N.; 118 38 20 E..	260	Few.
11906	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E..	750	Rare.
11965	U.S.N.M.	5	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m.	Few.
11904	} U.S.N.M.	1	} D5639...	3 54 50 S.; 123 27 20 E..	1,560	gy. m.	Rare.
15713		2						
11903	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.	Rare.
11908	U.S.N.M.	1	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.	Rare.

PULVINULINA SCHREIBERSII (d'Orbigny).

Rotalina schreibersii D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 154, pl. 8, figs. 4-6.

Pulvinulina schreibersii PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 393.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 697, pl.

115, figs. la-c.—CHAPMAN, Trans. New Zealand Inst., vol. 38, 1905, p. 106.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 62, fig. 59 (in text).

From the available records this species is largely limited to the Indo-Pacific and Mediterranean regions. Brady records it from seven Challenger stations from the south Pacific, ranging in depth from 3 to 155 fathoms (5 to 283 meters).

Pulvinulina schreibersii.—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
			D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m. ...	Few.
			D5132...	Island off Panabutan Point, N; 15° W., 3 mile.	26	gn. m., s.	Frequent.
			D5133...	Island off Panabutan Point, N; 52° E., 1.5 miles.	38	gn. m., s.	Few.
11914	} U.S.N.M.	2	D5134...	6 44 45 N.; 121 40 00 E..	25	fine s.	Frequent.
				D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s.
12343	U.S.N.M.	6	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Frequent.
14437	U.S.N.M.	10	D5261...	12 30 55 N.; 121 34 24 E..	145	s., m.	Common.
14435	U.S.N.M.	4	D5272...	14 00 00 N.; 120 22 30 E..	118	57.4	m., sh., co. s.	Common.
14436	} U.S.N.M.	1	D5278...	14 00 10 N.; 120 17 15 E..	102	59.6	fine s., m., sh.	Few.
				D5311...	21 33 00 N.; 116 15 00 E..	88	co. s., sh.

PULVINULINA PROCERA H. B. Brady.

Plate 69, figs. 1a-c.

Pulvinulina procera H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 661; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 698, pl. 105, figs. 7a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 62, pl. 24, fig. 2; pl. 25, fig. 2.

Brady described this species from three *Challenger* stations off Kandavu, Fiji Islands, 210 fathoms (384 meters), Torres Strait, off Raine Island, 155 fathoms (283 meters), and off Cape York, 3 to 11 fathoms (5 to 20 meters). I have had the species previously from off the Hawaiian Islands and off southern Japan. It is evidently a tropical species of the Indo-Pacific, in comparatively shallow water. It has occurred in the Philippine material in the following localities: China Sea, off southern Luzon; off western Samar; China Sea, off Hongkong; Darvel Bay, Borneo; vicinity of Jolo; vicinity of Romblon; off northern Cebu; between Burias and Luzon; Pacific Ocean, east coast of Mindanao; China Sea, off Formosa; Ragay Gulf, Luzon; between Samar and Leyte; and north of Tawi Tawi. It has, however, not been common at any of these stations, and was not abundant in any of the shallow-water stations where other species of this genus are very abundant.

Pulvinulina procera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11910	U.S.N.M.	1	D5097...	14 19 15 N.; 120 33 52 E.	30	gy. m., s., sh.	Rare.
11909	U.S.N.M.	2	D5100.....	14 17 15 N.; 120 32 40 E.	35	gy. s.....	Rare.
			D5172.....	6 03 15 N.; 120 35 30 E.	318	fine s., sh.	Few.
			D5178.....	12 43 00 N.; 122 06 15 E.	78	fine s.....	Few.
			D5192.....	11 09 15 N.; 123 50 00 E.	32	gn. s., m.	Few.
11912	U.S.N.M.	3	D5210.....	11 49 55 N.; 124 28 05 E.	50	76.3	fine gy. s.....	Few.
			D5217.....	13 20 00 N.; 123 14 15 E.	105	63.1	crs. gy. s.....	Few.
11913	U.S.N.M.	4	D5236.....	8 50 45 N.; 126 26 52 E.	494	41.2	fine gy. s.....	Few.
11911	U.S.N.M.	2	D5276.....	13 49 15 N.; 120 14 45 E.	18	sh., p., s.....	Few.
			D5277.....	13 56 55 N.; 120 13 45 E.	80	58.6	fine s.....	Few.
12335	U.S.N.M.	1	D5311.....	21 33 00 N.; 116 15 00 E.	88	crs. sh.....	Rare.
			D5313.....	21 30 00 N.; 116 43 00 E.	150	53.6	s.....	Few.
			D5318.....	21 32 00 N.; 117 46 00 E.	340	s., br. Co	Few.
14452	U.S.N.M.	1	D5424.....	9 37 05 N.; 121 12 37 E.	340	50.4	co. s.....	Rare.
12353	U.S.N.M.	2	D5580.....	4 52 45 N.; 119 06 45 E.	162	55.8	br. s., Co.....	Rare.
			D5381.....	13 14 15 N.; 122 44 45 E.	88	co. s.....	Frequent.
			D5443.....	12 43 05 N.; 125 01 00 E.	211	51.3	co. s., sh.....	Few.
15740	U.S.N.M.	2	D5469.....	13 36 48 N.; 123 38 24 E.	500	gn. m.....	Few.
		1	D5478.....	10 46 24 N.; 125 16 30 E.	57	sh.....	Rare.
			D5481.....	10 27 30 N.; 125 17 10 E.	61	s., sh., g.....	Few.
15741	U.S.N.M.	1	D5509.....	5 33 15 N.; 120 15 30 E.	303	52.3	co. s.....	Rare.
			D5570.....	5 32 15 N.; 120 12 57 E.	330	52.3	fine s., glob.....	Few.
			D5574.....	5 30 45 N.; 120 07 57 E.	Few.
		1	D5576.....	5 25 56 N.; 120 03 39 E.	277	53.3	s.....	Rare.

PULVINULINA ELEGANS (d'Orbigny).

Rotalia (*Turbinulina*) *elegans* D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 276, No. 54.

Pulvinulina elegans JONES and PARKER, Geologist, vol. 7, 1864, p. 88.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 699, pl. 105, figs. 4-6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 63, pl. 26, fig. 3.

In the *Challenger* Report Brady practically unites *P. elegans* and *P. partschiana*, the former more common in shallow water, the

latter in deeper water. In the Philippine material I have examined, *P. elegans* was noted at 76 stations, ranging in depth from 24 to 1,092 fathoms (44 to 1,998 meters), averaging 365 fathoms (668 meters), while *P. partschiana* occurred at but 11 stations, ranging from 32 to 742 fathoms (59 to 1,357 meters), with the average 301 fathoms (551 meters). As a matter of interest the records were compared for the two species and both species were recorded from but two stations in common. This would indicate that, as far as this area is concerned, the two species seem to be rather distinct.

In this connection it seems worthy of note that in their Kerimba Archipelago work²³ Heron-Allen and Earland record as noteworthy the occurrence of *P. partschiana* in very shallow water, and in the same region they had a single specimen of *P. elegans*, and that is noted as a "weak specimen."

The distribution of *P. elegans* covers the whole area dredged, both through the Archipelago and all the region to the southward in which dredgings were made.

Pulvinulina elegans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12358	U.S.N.M.	7	D5110...	13 59 20 N.; 120 75 45 E.	135	59.0	dk. gy. m...	Frequent.
			D5151...	5 24 40 N.; 120 27 15 E.	24	co. s., sh....	Few.
			D5153...	5 18 10 N.; 120 02 55 E.	49	co. s., sh....	Few.
			D5172...	6 03 15 N.; 120 35 30 E.	318	fine s., sh....	Few.
			D5203...	12 43 00 N.; 122 06 15 E.	78	fine s., sh....	Frequent.
			D5179...	12 38 15 N.; 122 12 30 E.	37	75.7	hrd. s.	Frequent.
			D5203...	9 58 00 N.; 125 07 40 E.	775	52.9	gn. m.	Few.
			D5217...	13 00 20 N.; 123 14 15 E.	105	63.1	crs. gy. s....	Few.
			D5219...	13 21 00 N.; 122 18 45 E.
			12356	U.S.N.M.	6	D5236...	8 50 45 N.; 126 26 52 E.	530
D5243...	6 50 55 N.; 126 14 35 E.	494				41.2	fine gy. s....	Common.
D5244...	6 52 05 N.; 126 14 15 E.	218				63.6	gy. m.	Few.
D5258...	10 27 45 N.; 122 12 30 E.	171				gy. m.	Few.
D5230...	12 25 35 N.; 121 33 35 E.	234				51.4	gn. m., s....	Few.
U.S.N.M.	2	D5281...		12 30 55 N.; 121 34 24 E.	145	s., m.	Few.
		D5238...		13 42 00 N.; 120 57 15 E.	170	s., p.	Frequent.
		D5277...		13 56 55 N.; 120 13 45 E.	80	58.6	fine s.	Few.
		D5278...		14 00 10 N.; 120 17 15 E.	120	59.6	fine s., m...	Few.
		D5282...		13 53 00 N.; 120 26 45 E.	248	47.4	dk. gy. s....	Few.
14454	U.S.N.M.	10	D5300...	20 31 00 N.; 115 49 00 E.	265	gy. m., s....	Common.
			D5301...	20 37 00 N.; 115 43 00 E.	208	50.5	gy. m., s....	Rare.
	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E.	150	53.6	s.	Few.
			D5318...	21 32 00 N.; 117 46 00 E.	340	s. br. Co....	Few.
	U.S.N.M.	1	D5331...	15 36 45 N.; 119 47 45 E.	178	54.7	s., sh. m....	Rare.
			D5370...	13 44 15 N.; 121 42 30 E.	159	54.3	sft. m.	Few.
			D5374...	13 46 45 N.; 121 35 08 E.	190	gy. m.	Few.
	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 45 E.	88	co. s.	Few.
			D5398...	11 35 12 N.; 124 13 43 E.	114	gn. m.	Frequent.
	U.S.N.M.	1	D5410...	10 28 45 N.; 124 05 30 E.	385	gn. m.	Rare.
D5412...			10 09 15 N.; 123 52 00 E.	162	54.8	gn. m.	Rare.	
U.S.N.M.	1	D5423...	9 38 30 N.; 121 11 00 E.	508	49.8	gy. m., co. s.	Few.	
		D5449...	15 58 15 N.; 119 40 20 E.	940	36.7	gn. m.	Rare.	
U.S.N.M.	10+	D5445...	12 44 42 N.; 124 59 50 E.	383	44.3	gn. m., s....	Few.	
		D5449...	13 21 36 N.; 124 00 30 E.	730	40.6	co. s.	Few.	
12363	U.S.N.M.	10+	D5460...	13 32 30 N.; 123 37 18 E.	480	gy. m.	Common.
			D5465...	13 39 42 N.; 123 40 39 E.	500	gy. m.	Common.
12364	U.S.N.M.	6	D5467...	13 35 27 N.; 123 37 18 E.	480	gy. m.	Few.
			D5469...	13 36 48 N.; 123 38 24 E.	500	gn. m.	Frequent.
			D5470...	13 37 30 N.; 123 41 09 E.	560	m.	Few.
	U.S.N.M.	1	D5487...	10 02 45 N.; 125 05 33 E.	732	53.3	gn. m.	Rare.
			D5523...	8 48 44 N.; 123 27 35 E.	Few.
			D5529...	9 23 45 N.; 123 39 30 E.	441	53.0	gy. m.	Few.
D5537...	9 11 00 N.; 123 23 00 E.	251	53.5	gn. m.	Few.			

²³ Trans. Zool. Soc. London, vol. 20, pt. 17, 1915, p. 717.

Pulvinulina elegans—Material examined—Continued.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
			D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gn. m., s....	Few.
			D5542...	8 48 30 N.; 123 35 30 E..	200	54.3	fine. s.....	Few.
	U.S.N.M.	1	D5543...	8 47 15 N.; 123 35 00 E..	162	54.5	s.....	Rare.
			D5548...	6 00 20 N.; 120 45 35 E..	232	53.5	s., brk. sh....	Frequent.
	U.S.N.M.	1	D5566...	5 52 12 N.; 120 31 00 E..	244	52.5	fine. sh., s....	Few.
			D5567...	5 48 00 N.; 120 33 45 E..	298	52.0	fine. s.....	Rare.
			D5569...	5 32 15 N.; 120 15 30 E..	303	52.3	co. s.....	Frequent.
			D5570...	5 32 15 N.; 120 12 57 E..	330	52.3	fine. s.....	Few.
			D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s.....	Few.
			D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Few.
	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co....	Few.
12361			D5582...	4 19 54 N.; 118 58 38 E..	890	38.3	gy. m.....	Rare.
			D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Few.
	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.....	Few.
			D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s....	Few.
			D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Few.
12359	U.S.N.M.	2	D5606...	0 16 28 N.; 121 33 30 E..	834	gn. m.....	Few.
	U.S.N.M.	1	D5609...	0 11 00 S.; 121 16 00 E..	1,092	36.3	gn. m.....	Few.
			D5612...	0 38 00 S.; 121 45 40 E..	750	Few.
			D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.....	Common.
			D5621...	0 15 00 N.; 127 24 35 E..	298	gy. s.....	Few.
	U.S.N.M.	1	D5623...	0 16 30 N.; 127 30 00 E..	272	fine. s.....	Few.
12360	U.S.N.M.	2	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m....	Few.
	U.S.N.M.	1	D5632...	1 00 00 S.; 127 50 00 E..	845	Few.
	U.S.N.M.	2	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	Few.
12362	U.S.N.M.	1	D5640...	4 27 00 S.; 122 50 44 E..	24	s., brk. sh....	Few.
			D5648...	5 35 00 S.; 122 20 00 E..	559	39.2	gn. m.....	Few.
			D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Frequent.
			D5652...	4 35 00 S.; 121 23 06 E..	525	41.2	gn. m.....	Few.
12357	U.S.N.M.	6	D5654...	3 42 00 S.; 120 45 50 E..	805	38.3	Few.
15742	U.S.N.M.	1	D5660...	5 36 30 S.; 120 49 00 E..	692	39.2	gy. m., s....	Few.
			D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.....	Frequent.
14911	U.S.N.M.	1	Nero 849	15 16 00 N.; 121 40 00 E..	737	gn. m.....	Rare.

PULVINULINA PARTSCHIANA (d'Orbigny).

Rotalina partschiana D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 153, pl. 7, figs. 28-30, pl. 8, figs. 1-3.

Pulvinulina partschiana REUSS, Sitz. Akad. Wiss. Wien, vol. 62, 1870, p. 36.—VON SCHLICHT, Foram. Pietzpuhl, 1870, pl. 20, figs. 23-25, 29-31.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 699, pl. 105, figs. 3a-c, woodcut, fig. 21.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 64, text figs. 60a-c.

Mention has already been made of the distribution of this species in comparison with *P. elegans*. Not mentioned there, however, is the peculiar fact that *P. partschiana*, as far as our material seems to show, is limited to the area of the Archipelago and not found in the material to the southward, where *P. elegans* is fairly common.

Pulvinulina partschiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12339	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E..	393	43.7	gn. m., s....	Rare.
12340	U.S.N.M.	1	D5126...	10 34 54 N.; 121 47 30 E..	742	49.5	sft. gn. m....	Rare.
12341	U.S.N.M.	4	D5187...	9 16 45 N.; 123 21 15 E..	225	53.6	sft. gn. m....	Few.
12352	U.S.N.M.	4	D5217...	13 00 20 N.; 123 14 15 E..	105	63.1	crs. gy. s....	Few.
12342	U.S.N.M.	2	D5247...	7 02 00 N.; 125 38 45 E..	135	m.....	Rare.
12384	U.S.N.M.	4	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Few.
14440	U.S.N.M.	2	D5260...	12 25 35 N.; 121 31 35 E..	234	51.4	gn. m., s....	Few.
14439	U.S.N.M.	6	D5370...	13 44 15 N.; 121 42 30 E..	159	54.3	sft. m.....	Common.
14902	U.S.N.M.	2	Nero 849	15 16 00 N.; 121 40 00 E..	737	gn. m.....	Rare.

Genus ROTALIA Lamarck, 1804.

ROTALIA BECCARII (Linnaeus).

Plate 70, figs. 3a-c.

Nautilus beccarii LINNAEUS, Syst. Nat., ed. 12, 1767, p. 1162; ed. 13 (Gmelin's), 1788, p. 3370, No. 4.

Rotalia (Turbinulina) beccarii D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 275, No. 40; Modèles, 1826, No. 74.

Rotalia beccarii PARKER and JONES, Philos. Trans., vol. 155, 1865, p. 388, pl. 16, figs. 29, 30.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 704, pl. 107, figs. 2, 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 67, pl. 30, fig. 3.

Although the number of stations for this species is large, there are no stations among them from the region south of the Philippines themselves. It occurred in general in the following localities: Off Formosa; China Sea, off western and southern Luzon; Baylayan Bay and Verde Island Passage; between Marinduque and Luzon; between Burias and Luzon; off Romblon; east of Masbate; east coast of Luzon; off northwestern Panay; between Panay and Negros; Tanon Strait, east coast of Negros; between Siquijor and Bohol; Sogod Bay, southern Leyte; eastern Illana Bay, southern Mindanao; off Jolo; off Tawi Tawi; off eastern Palawan and Palawan Passage. These localities cover the area fairly well. The range in depth is from 12 to 805 fathoms (22 to 1,472 meters), with a few specimens from Little Harbor, near Manila, in a few feet of water; the average depth of all stations, 318 fathoms (582 meters). Bottom temperatures range from 40.6° to 63.1° F. (4.70° to 17.2° C.), with the average 51.3° F. (10.7° C.).

The specimens as a rule were rather small, with little or no raised surface ornament, and the usual brownish tinge to the translucent test.

Rotalia beccarii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / ° ° / °		° F.		
14863	U.S.N.M.	1	D5112...	13 48 22 N.; 120 47 25 E...	177	52.4	dk. gn. m....	Few.
14864	U.S.N.M.	1	D5114...	13 36 11 N.; 120 45 26 E...	340	fine. s.....	Rare.
		1	D5120...	13 45 30 N.; 120 30 15 E...	393	43.7	gn. m., s.....	Rare.
		1	D5139...	6 06 00 N.; 121 02 30 E...	20	co. s.....	Rare.
14865	U.S.N.M.	2	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.....	Rare.
		1	D5160...	5 12 40 N.; 119 55 10 E...	12	s.....	Rare.
14866	U.S.N.M.	4	D5219...	13 21 00 N.; 122 18 45 E...	530	50.8	gn. m.....	Rare.
			D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.....	Rare.
14867	U.S.N.M.	3	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
			D5185...	10 05 45 N.; 122 18 30 E...	638	49.8	gn. m.....	Rare.
14868	U.S.N.M.	1	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co.....	Rare.
14428	U.S.N.M.	3	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Few.
14869	U.S.N.M.	5	D5526...	9 12 45 N.; 123 45 30 E...	805	52.3	gn. m., glob.	Few.
14870	U.S.N.M.	9	Cebu.....
	U.S.N.M.	1	D5191...	10 29 45 N.; 123 31 15 E...	258	62.8	gn. m.....	Rare.
			D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Few.
	U.S.N.M.	1	D5215...	12 31 30 N.; 123 35 24 E...	604	50.5	gn. m.....	Rare.
			D5222...	13 38 30 N.; 121 42 15 E...	195	52.8	gn. m.....	Few.
			D5257...	7 22 12 N.; 124 12 15 E...	28	m.....	Few.
			D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Few.
	U.S.N.M.	1	D5331...	15 36 45 N.; 119 47 45 E...	178	54.7	s. sh., m.....	Rare.
	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E...	730	40.6	co. s.....	Rare.
			D5430...	9 49 40 N.; 119 03 20 E...	464	50.0	glob. ooze.....	Few.
	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.....	Rare.

ROTALIA BROECKHIANA Karrer.

Plate 72, figs. 1a-c.

Rotalia broeckhiana KARRER, in Drasche, Geol. Insel Luzon, 1878, p. 98, pl. 5, fig. 26.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 705, pl. 107, figs. 4a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 68, pl. 27, fig. 4; pl. 30, fig. 2, text figs. 61a-c.

Although of wide distribution in the area this species is represented at but 13 stations, and at most of these by very few specimens. The specimens, however, are very typical, both in size and other specific characters.

The stations include the following localities: China Sea, off Formosa; Verde Island Passage and Batangas Bay; Mindoro Strait; between Siquijor and Bohol; east coast of Mindanao; off Jolo; Palawan Passage; and in the region to the south, off Bouro Island and Gulf of Tomini, Celebes.

In depth these stations range from 220 to 805 fathoms (403 to 1,472 meters), with the average depth 492 fathoms (901 meters). Bottom temperatures range from 41.2° to 73.8° F. (5.1° to 23.2° C.), with the average 52.0° F. (11.1° C.).

Rotalia broeckhiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14871	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318	fne. s., sh.....	Rare.
14872	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s.....	Rare.
		1	D5269...	13 39 50 N.; 120 59 36 E...	220	fne. s., p.....	Rare.
14873	U.S.N.M.	4	D5333...	12 26 30 N.; 120 37 45 E...	310	73.8	s.....	Rare.
		1	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co.....	Rare.
14874	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.....	Rare.
		1	D5333...	12 26 30 N.; 120 37 45 E...	310	73.8	s.....	Rare.
14875	U.S.N.M.	1	D5421...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Rare.
		1	D5423...	9 38 30 N.; 121 11 00 E...	508	49.8	gy. m.....	Rare.
14876	U.S.N.M.	1	D5612...	0 38 00 S.; 121 45 40 E...	750	Rare.
		1	D5449...	13 21 36 N.; 124 00 30 E...	730	40.6	Co., s.....	Rare.
14877	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Rare.
		1	D5526...	9 12 45 N.; 123 45 30 E...	805	52.3	gn. m.....	Rare.

ROTALIA ORBICULARIS d'Orbigny.

Rotalia (Gyroidina) orbicularis d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 278, No. 1; Modèles, 1826, No. 13.

Rotalia orbicularis H. B. BRADY, Trans. Linn. Soc. London, vol. 24, 1864, p. 470, pl. 48, fig. 16; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 706, pl. 107, fig. 5; pl. 115, fig. 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 68, pl. 29, fig. 3; text figs. 62a-c.

Typical specimens of this species seem to be rare. The fact that later authors have departed more or less widely from the typical form described by d'Orbigny has been noted by Heron-Allen and Earland.²⁴ At only two stations have specimens been found which have seemed

²⁴ Trans. Zool. Soc. London, vol. 20, pt. 17, 1915, p. 718.

really identical with this species—D5217, in 105 fathoms (193 meters), between Burias and Luzon, bottom temperature, 63.1° F. (17.2° C.); and D5260, in 234 fathoms (428 meters), off southeastern Mindoro, bottom temperature 51.4° F. (10.8° C.). At each of these stations a few specimens were found.

Rotalia orbicularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14430	U.S.N.M.	2	D5217...	° ' " ° ' "	105	° F. 63.1	crs. gy. s.....	Rare.
14429	U.S.N.M.	4	D5260...	13 20 00 N.; 123 14 15 E... 12 25 35 N.; 121 31 35 E...	234	51.4	gn.m.,s.....	Few.

ROTALIA SCHROETERIANA Parker and Jones.

Plate 73, figs. 1a-c.

Faujasina, species, WILLIAMSON, Trans. Micr. Soc., London, ser. 2, vol. 1, 1853, p. 87, pl. 10.

Rotalia schroeteriana (Parker and Jones, MS), CARPENTER, Introd. Foram., p. 213, pl. 13, figs. 7-9.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 707, pl. 115, figs. 7a-c.

Brady gives the following note in the *Challenger* Report in regard to this species:

Though somewhat local in distribution, it is by no means rare amongst the islands of the Eastern Archipelago, at depths of less than 50 or 60 fathoms (91 or 110 meters).

The only records for this species in the Philippine region are in the Archipelago proper, mostly in shallow water.

Rotalia schroeteriana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14833	U.S.N.M.	10+	D5313...	° ' " ° ' "	150	° F. 53.6	s.....	Common.
14834	U.S.N.M.	3	D5336...	21 30 00 N.; 116 43 00 E...	46	s.,m.....	Few.
14835	U.S.N.M.	10+	D5338...	11 37 45 N.; 119 46 00 E...	43	co.s.,m.....	Common.
14836	U.S.N.M.	10+	D5342...	11 33 45 N.; 119 24 45 E...	14-25	gy.m.....	Common.
14837	U.S.N.M.	10+	D5348...	10 56 55 N.; 119 17 24 E...	375	56.4	co.s.....	Common.
14838	U.S.N.M.	7	D5426...	10 67 45 N.; 118 38 15 E...	27	inc.gy.s.....	Common.
12879	U.S.N.M.	2	9 12 00 N.; 118 28 00 E... Manila Bay.....	Few.

ROTALIA PAPILOSA H. B. Brady.

Plate 72, figs. 3a, b.

Rotalia papillosa H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 708, pl. 106, figs. 9a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 70, pl. 31, fig. 1.

From available records this species is characteristic of the Indo-Pacific region, and in the Philippines it is certainly one of the most

abundant species. The ornamentation is beautifully sculptured, and varies considerably in strength but very little in its main characters. The list of stations at which it occurs would be a recapitulation of the main localities at which dredging was done, from Formosa on the north through the large part of the Archipelago. Besides this general area it occurs at two stations to the south—D5580, in 162 fathoms (297 meters), Darvel Bay, Borneo, bottom temperature 55.8° F. (13.2° C.); and D5585, in 476 fathoms (871 meters), Sibuko Bay, Borneo, bottom temperature 41.1° F. (5° C.). The general distribution should be modified, however, to this extent, that the species is characteristic of the protected waters among the islands of the Archipelago and on the western border along the China Sea, but is not represented at the stations on the eastern coast bordered by the open Pacific. The range in depth is from 18 to 476 fathoms (33 to 871 meters), the average depth 118 fathoms (203 meters), the bottom temperatures ranging from 41.1° F. to 55.8° F. (5° C. to 13.2° C.), average 51.2° F. (10.7° C.).

Rotalia papillosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14846	U.S.N.M.	10+	D5096...	14 20 23 N.; 120 34 15 E...	28	° F.	gy. m., s., sh.	Common.
14847	U.S.N.M.	10	D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s.....	Common.
14850	U.S.N.M.	10+	D5132...	Island off Panabutan Point, N., 15° W., E., 0.30 mile.	26	gn. m., s.....	Common.
14851	U.S.N.M.	9	D5134...	6 44 45 N.; 121 48 00 E...	25	fne. s.....	Common.
14852	U.S.N.M.	4	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.....	Few.
14853	U.S.N.M.	10+	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fne. s.....	Common.
14854	U.S.N.M.	8	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Common.
14855	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Rare.
14857	U.S.N.M.	10	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Common.
14859	U.S.N.M.	9	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Common.
14861	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E...	150	53.6	s.....	Rare.
14867	U.S.N.M.	2	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Rare.
14832	U.S.N.M.	10+	D5642...	4 31 40 S.; 122 49 42 E...	37	gy. m.....	Common.
			D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co.....	
			D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.....	

ROTALIA PAPILOSA H. B. Brady, var. COMPRESSIUSCULA H. B. Brady.

Plate 72, figs. 2a-c.

Rotalia papillosa H. B. BRADY, var. *compressiuscula* H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 708, pl. 107, figs. 1a-c; pl. 108, figs. 1a-c.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 70, pl. 30, fig. 1.

The variety seems to be slightly more restricted in its range than the typical form, not being recorded to the southward of the Archipelago proper, although well distributed among the islands and in the China Sea. The range of depth is from 16 to 560 fathoms (29 to 1,024 meters), the average depth being 179 fathoms (327 meters). Bottom temperatures range from 50.5° F. to 63.1° F. (10.2° C. to 17.2° C.), the average being 61.1° F. (16.1° C.).

At some of the stations specimens occurred in very considerable numbers, at a few really abundant. Among the stations from

which this variety was described by Brady is one *Challenger* station in the Philippines in 95 fathoms (174 meters.) Although the typical form and the variety both occur commonly in shallow water they were most abundant in the Philippine material from 100 to 125 fathoms (183 to 229 meters).

Rotalia papillosa, var. *compressiuscula*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "		° F.		
14848	U.S.N.M.	10+	D5113	13 51 30 N.; 120 50 30 E.	159	dk. gn. m...	Common.
14849	U.S.N.M.	10+	D5121	13 27 20 N.; 121 17 45 E.	108	dk. gn. m...	Common.
14856	U.S.N.M.	10+	D5247	7 02 00 N.; 125 28 45 E.	135	m.....	Common.
14860	U.S.N.M.	10+	D5268	13 42 00 N.; 120 57 15 E.	170	s. p.....	Common.
14853	U.S.N.M.	7	D5272	14 00 00 N.; 120 22 30 E.	113	57.4	m, sh., co. s.	Few.
14862	U.S.N.M.	8	D5313	21 30 00 N.; 116 43 00 E.	159	53.6	s.....	Few.
14828	U.S.N.M.	7	D5385	13 24 50 N.; 123 03 70 E.	32.7	62.4	gy. m.....	Few.
14829	U.S.N.M.	9	D5394	12 00 30 N.; 124 05 36 E.	153	gn. m.....	Common.
		1	D5469	13 36 43 N.; 123 38 24 E.	500	gn. m.....	Rare.
14830	U.S.N.M.	2	D5470	13 37 30 N.; 123 41 09 E.	560	m.....	Rare.
14831	U.S.N.M.	7	D5576	5 25 55 N.; 120 03 39 E.	277	53.3	s.....	Few.
			D5110	13 59 20 N.; 120 75 45 E.	135	59	dk. gy. m...	Frequent.
			D5134	6 44 45 N.; 121 48 00 E.	25	fine. s.....	Few.
			D5156	5 12 50 N.; 119 55 55 E.	18	fine. s.....	Frequent.
			D3161	5 10 15 N.; 119 53 00 E.	16	fine. s.....	Few.
			D5178	12 43 00 N.; 122 06 15 E.	78	fine. s.....	Few.
			D5217	13 59 20 N.; 123 14 15 E.	105	63.1	crs. gy. s...	Frequent.
			D5220	13 38 00 N.; 121 58 00 E.	50	sft. gn. m...	Frequent.
			D5255	7 03 00 N.; 125 39 00 E.	100	sft. m.....	Few.
			D5261	12 30 55 N.; 121 34 24 E.	145	s. m.....	Few.
			D5269	13 39 50 N.; 120 59 30 E.	220	fine. s., p...	Frequent.
			D5301	20 37 00 N.; 115 43 00 E.	208	50.5	gv. m., s...	Few.
			D5370	13 44 15 N.; 121 42 30 E.	159	54.3	sft. m.....	Few.
			D5374	13 46 44 N.; 121 35 08 E.	190	gv. m.....	Few.
		1	D5410	10 28 45 N.; 124 05 30 E.	385	gn. m.....	Rare.
			D5412	10 09 15 N.; 123 52 00 E.	162	54.8	gn. m.....	Frequent.
		1	D5419	9 58 30 N.; 123 46 00 E.	175	54.5	gn. m.....	Rare.
			D5420	9 49 35 N.; 123 45 00 E.	127	59	Few.
		1	D5443	12 43 51 N.; 124 58 50 E.	241	51.3	co. s., sh...	Few.

ROTALIA SOLDANII d'Orbigny.

Plate 71, figs. 2a-c.

Rotalia (Gyroidina) soldanii D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 278, No. 5; Modèles, 1826, No. 36.

Rotalia soldanii HANTKEN, Mitt. Jahrb. ung. geol. Anstalt., 1875, p. 80, pl. 9, figs. 7a-c.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 706, pl. 107, figs. 6, 7.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 71, pl. 29, fig. 1; pl. 31, fig. 4.

In the Philippine region this species occurs at widely scattered stations, usually not in any considerable numbers. It also keeps to its usual condition of being found in comparatively deep water, the range here being from 78 to 1,560 fathoms (143 to 2,853 meters), the last in the Molucca Sea, average 530 fathoms (969 meters). Bottom temperatures range from 41.2° F. to 73.8° F. (5.1° C. to 23.2° C.), average 52.7° F. (11.4° C.). The localities include Mindoro Strait; off Romblon; between Siquijor and Bohol; between Negros and Siquijor; between Leyte and Mindanao; Sogod Bay, southern Leyte; Pacific Ocean, east coast of Mindanao; Gulf of Davao, Mindanao; north of Tawi Tawi; off Palawan; Darvel Bay, Borneo; Sibuko Bay, Borneo; Molucca Sea; and Macassar Strait.

Rotalia soldanii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14839	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
14841	U.S.N.M.	10	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Common.
14840	U.S.N.M.	1	D5247...	7 02 00 N.; 125 38 45 E...	135	m.....	Rare.
14842	U.S.N.M.	1	D5333...	12 26 30 N.; 120 37 45 E...	310	73.8	s.....	Rare.
14816	U.S.N.M.	1	D5526...	9 12 45 N.; 123 45 30 E...	805	52.3	gn. m., glob.	Rare.
14817	U.S.N.M.	1	D5575...	5 28 30 N.; 120 02 27 E...	315	52.3	co. s.....	Rare.
14427	U.S.N.M.	5	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.....	Few.
			D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.....	Few.
			D5430...	9 49 40 N.; 119 03 20 E...	464	50.0	glob. ooze...	Few.
		1	D5495...	9 06 30 N.; 125 00 20 E...	976	52.3	gy. m., s....	Rare.
		1	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s....	Rare.
		1	D5530...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co...	Rare.
			D5392...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.....	Few.
			D5665...	4 27 00 S.; 118 44 00 E...	1,008	m.....	Few.

ROTALIA CALCAR (d'Orbigny).

Plate 71, figs. 3a, b.

Calcarina calcar D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 276, No. 1; in De la Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 93, pl. 5, figs. 22-24.
Rotalia calcar H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 709, pl. 108, fig. 3, fig. 4?—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 69, pl. 28, fig. 2; pl. 29, fig. 2.

This species, which is characteristic of shallow water of coral-reef regions, is abundant under similar conditions in the Philippine Archipelago. Some specimens resemble those figured by Brady; others have longer extensions on the margins of the chambers and become more like *Calcarina*. It is usually associated with *Calcarina*, and there is some question whether or not it should be included in that genus.

Rotalia calcar—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14878	U.S.N.M.	1	D5096...	14 20 23 N.; 120 34 15 E...	28	gy. m., s., sh	Rare.
14879	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Rare.
14880	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Rare.
14843	U.S.N.M.	2	D5238...	7 34 45 N.; 126 38 15 E...	380	43	gn. m.....	Rare.
14844	U.S.N.M.	2	D5268...	13 42 00 N.; 120 57 15 E...	170	p., s., sh....	Rare.
14845	U.S.N.M.	1	Cebu	Rare.
15705	U.S.N.M.	1	TaraId	Rare.
15706	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co. s.....	Rare.
15707	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.....	Rare.
15708	U.S.N.M.	1	D5370...	13 44 15 N.; 121 42 30 E...	159	54.3	sft. m.....	Rare.

Genus CALCARINA d'Orbigny, 1826.

Nautilus (part) LINNAEUS, Syst. Nat., ed. 13 (Gmelin's), 1788, p. 3371.

Calcarina D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 276 (type, *Nautilus spengleri* (Linnaeus)).

Description.—Test rotaliform throughout, chambers numerous, close-coiled, biconvex; periphery with radiating spines; chambers of the last-formed coil visible on the ventral side, those of the dorsal

side partially hidden by the supplementary skeleton which is more or less developed in all the species; aperture typically a row of small openings along the inner margin of the apertural face; canal system well-developed.

This genus is represented by several recent species, which are confined to the shallow water of the tropics and the Mediterranean. In the Philippine material four species are found, some of them represented by abundant specimens.

A study of the three genera—*Calcarina*, *Baculogypsina*, and *Siderolites*—as well as *Tinoporos* Montfort, has shown very definite relationships. In an earlier short paper I discussed these genera. It seems that the genus *Tinoporos* Montfort is too indefinite to be of value, the species *baculatus* Montfort being now probably recognizable, but as he refers in his generic description to certain characters which belong more closely to what must be, by the study of the characters, now called *Baculogypsina sphaerulata*, the whole genus should be eliminated.

The type species of *Calcarina* is *Nautilus spengleri* Gmelin. This species is definitely taken as the genotype by d'Orbigny. This is a well-defined species, as will be shown. Of the other species, *C. defranceii*, described by d'Orbigny, is recognizable, and also *C. hispida*, described by H. B. Brady. The fourth species here called *C. baculatus* Montfort makes up the series.

Of the other genera, *Siderolites* Lamarck, of which the type is *S. calcitrapoides* Lamarck from the chalk of Maestricht, seems to have a very definite place. Material from the type locality was studied and shows that the species is much more complex than most of the species of *Calcarina*, and should be used in a definite way for *S. tetraedra* Gumbel, which occurs in the Philippine collections.

Baculogypsina, erected by Sacco in 1893 on account of the uncertainty of the identity of Montfort's *Tinoporos*, is here used for *Baculogypsina sphaerulata* (Parker and Jones), which is the species of the *Challenger* report and other works usually passing as *Tinoporos* Montfort.

Calcarina has the simplest development of the three and is least specialized, apparently going back geologically to the Cretaceous. The following species occur:

CALCARINA SPENGLERI (Gmelin).

Plate 75, fig. 1.

Nautilus spengleri LINNÆUS, Syst. Nat., ed. 13 (Gmelin's), 1788, p. 3371, No. 10.

Calcarina spengleri D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 276, No. 4.—H. B.

BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 712, pl. 108, figs. 5, 7.—

CUSHMAN, U. S. Nat. Mus., pt. 5, 1915, p. 72, pl. 31, fig. 2.

Description.—Test lenticular, biconvex, chambers numerous, in a close-coiled trochoid spire of three or four volutions, each with several chambers, but the sutures not usually all distinguishable on the dorsal side except the last few; wall covered by a supplementary skeleton, the surface with raised tubercles of solid, translucent material, espe-

cially in the central part of the disk at either side; peripheral border broadly rounded with a series of blunt, subcylindrical, spinose processes, three to six or more in number, usually five or six; smooth except for the channels of the supplementary canals; aperture a row of rounded pores along the inner margin of the chamber.

Diameter up to 3 mm. or more.

The species is very common in the Philippine collection, especially in warm shallow waters, where it is sometimes very abundant. The depths at which specimens were found range from 16 to 1,092 fathoms (29 to 1,998 meters), with the average depth 173 fathoms (316 meters). Bottom temperatures, where given, range from 51.5° F. to 59° F. (10.8° C. to 15° C.), the average temperature being 51.5° F. (10.8 C.).

The stations include the following localities: China Sea, off southern Luzon; Sulu Sea, off western Mindanao; vicinity of Jolo; off northern Cebu; eastern Palawan and vicinity; northern Mindanao and vicinity; Gulf of Tomini, Celebes; south of Patiente Strait; Binang Pool, Subin Bay; and off Cebu.

Linnaeus is usually given as the author of the species, but it apparently was not named until Gmelin's edition, and according to present usage should be given Gmelin's name. There seems to be no question as to what is meant by the Spengler figure of "Ammonshorn" in 1781 which is referred to by Gmelin. In his 1826 work d'Orbigny gives *Tinoporus baculatus* Montfort and *Siderolites calcitrapoides* Lamarck as synonyms of *Calcarina spengleri*, but both seem to be other things, as will be discussed later. The *Challenger* figures of this species seem to be excellent.

Calcarina spengleri—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
12762	U.S.N.M.	7	Cebu	° F.	Common.
12761	U.S.N.M.	3	Sulu Id.	Few.
14511	U.S.N.M.	1	D5105...	14 43 55 N.; 120 12 50 E...	25	Few.
14512	U.S.N.M.	10+	D5106...	4 23 55 N.; 120 32 33 E...	37	gy. m.	Common.
14513	U.S.N.M.	5	D5110...	13 59 20 N.; 120 75 45 E...	135	59.0	dk. gy. m.	Common.
14514	U.S.N.M.	10+	D5113...	13 51 30 N.; 120 50 30 E...	159	dk. gn. m.	Common.
14515	U.S.N.M.	2	D5132...	Island off Panabutan Point N.; 15° W., 0.3 mile.	26	gn. m., s.	Few.
14516	U.S.N.M.	3	D5133...	Island off Panabutan Point N.; 52° E., 1.5 miles.	38	gn. m., s.	Rare.
14517	U.S.N.M.	1	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.	Rare.
14518	U.S.N.M.	1	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.	Rare.
14519	U.S.N.M.	4	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.	Rare.
14520	U.S.N.M.	4	P5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Few.
14521	U.S.N.M.	1	P5161...	5 10 15 N.; 119 53 00 E...	16	fine s., blk. sp.	Few.
14737	U.S.N.M.	10	P5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.	Common.
14522	U.S.N.M.	1	P5338...	11 33 45 N.; 119 24 45 E...	43	co., s., m.	Rare.
14533	U.S.N.M.	1	P5429...	9 41 30 N.; 118 50 22 E...	766	gn. m.	Rare.
14534	U.S.N.M.	5	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., co.	Common.
14535	U.S.N.M.	5	P5523...	8 48 44 N.; 123 27 35 E...	Common.
14536	U.S.N.M.	1	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fine s., Co.	Few.
14537	U.S.N.M.	1	P5609...	0 11 00 S.; 121 16 00 E...	1,092	36.3	gn. m.	Rare.
14538	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.	Rare.
14539	U.S.N.M.	8	Binang Pool, Subin Bay, P. I.	Common.
14738	U.S.N.M.	9	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.	Common.
14739	U.S.N.M.	1	D5291...	13 29 40 N.; 121 00 45 E...	173	51.5	fine bk. s.	Rare.

CALCARINA DEFRANCI *d'Orbigny*.

Plate 75, fig. 2.

Calcarina defrancii D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 276, pl. 13, figs. 5-7.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 714, pl. 108, figs. 6a-c.—MILLETT, Journ. Roy. Micr. Soc., 1904, p. 598.—FORNASINI, Mem. Accad. Sci. Bologna, ser. 6, vol. 5, 1908, p. 7, pl. 3, figs. 3, 4.

Calcarina spengleri, var., CARPENTER, PARKER, and JONES, Introd. Foram., 1862, p. 217, pl. 14, fig. 2, figs. 34 A-G in text.—PARKER and JONES, Ann. Mag. Nat. Hist., ser. 3, vol. 12, 1863, p. 439, No. 6.

Description.—Test lenticular, biconvex, chambers numerous in a close-coiled, flattened, trochoid spire of about three volutions, each with several chambers; the sutures distinct on the ventral side, but much obscured or obsolete on the dorsal side, the spire, however, usually distinct; wall, especially on the dorsal side, covered by a supplementary skeleton, the surface with spinose tubercles; peripheral border bluntly angled, with a series of elongate spines, generally cylindrical; the outer ends blunt, typically one spine from the outer end of each chamber, channeled longitudinally; the neighboring spines of the different whorls often fusing near the base and the free ends continuing their direction make an apparent bifurcating spine; aperture at the base of the last-formed chamber.

Diameter up to 2.5 mm. or more.

The species is very common at a few stations, most of which are in shallow, quiet water, and it is evident that its best development is in such localities. D'Orbigny's original specimens were from the Red Sea. Brady had his best specimens from off the Admiralty Islands in 15 to 25 fathoms (27 to 46 meters). Millett gives the following:

In the Malay Archipelago this is the most abundant species of the genus.

Fornasini in figuring the tracings of the "planchés inédites" gives more bizarre forms than the *Challenger* or d'Orbigny's originals, and figures one specimen with decidedly forked tips to the spines. Carpenter also figures similar specimens as the "Philippine variety of *Calcarina*," and in his text clearly shows that *C. defrancii* is much more abundant in the Philippines than in the Mediterranean.

In the Philippine collections this has occurred at the following localities: China Sea, off southern Luzon; Sulu Sea, off western Mindanao; Sulu Archipelago; Tawi Tawi Group, between Panay and Negros; Tanon Strait; Pujuda Bay and vicinity; Gulf of Davao; Palawan Passage; Ragay Gulf; Luzon, between Burias and Luzon; Buton Strait; and Binang Pool, Subin Bay.

The range in depth is from 18 to 565 fathoms (33 to 1,033 meters), and the bottom temperatures, where given, range from 46.2° F. to 79.5° F. (7.8° C. to 26.3° C.).

It is evident that in the Philippine and Malay regions this is a very abundant species.

Calcarina defrancii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14491	U.S.N.M.	2	D5097...	14 19 15 N.; 120 33 52 E...	30	gy. m., s., sh.	Rare.
14492	U.S.N.M.	2	D5105...	14 43 55 N.; 120 12 50 E...	25	Rare.
14493	U.S.N.M.	6	D5132...	Island off Panabutan Point, N.; 15° W., 0.3 mile.	26	gn. m., s....	Few.
14494	U.S.N.M.	10+	D5133...	Island off Panabutan Point N.; 52° E. 1.5 miles.	38	79.5	gn. m., s....	Common.
14495	U.S.N.M.	10	D5134...	6 44 45 N.; 121 48 00 E...	25	fine s.....	Common.
14496	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Rare.
14497	U.S.N.M.	3	D5184...	10 18 30 N.; 122 23 30 E...	565	49.8	gn. m.....	Rare.
14498	U.S.N.M.	2	D5191...	10 29 45 N.; 123 31 15 E...	258	62.8	gn. m.....	Rare.
15374	U.S.N.M.	2	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m....	Rare.
14499	U.S.N.M.	5	D5244...	6 52 05 N.; 126 14 15 E...	171	gy. m.....	Few.
14500	U.S.N.M.	1	D5250...	7 05 07 N.; 125 39 45 E...	23	Co., s.....	Rare.
14501	U.S.N.M.	5	D5255...	7 03 00 N.; 125 39 00 E...	100	sft. m.....	Few.
14502	U.S.N.M.	6	D5276...	13 49 15 N.; 120 14 15 E...	18	sh., p., s....	Few.
14503	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s....	Rare.
14505	U.S.N.M.	1	D5338...	11 33 45 N.; 119 24 45 E...	43	Co., s., m....	Rare.
14506	U.S.N.M.	2	D5358...	6 06 40 N.; 118 18 15 E...	39	m.....	Rare.
14507	U.S.N.M.	2	D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.....	Rare.
14504	U.S.N.M.	1	D5388...	12 51 30 N.; 123 26 15 E...	226	51.4	sft. gn. m....	Rare.
14508	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Rare.
14509	U.S.N.M.	3	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh.	Few.
14510	U.S.N.M.	6	Binang Pool, Subin Bay, P. I.	Few.

CALCARINA BACULATUS (?) (Montfort).

Plate 75, fig. 3.

Tinoporos baculatus MONTFORT (part), Conch. Syst., vol. 1, 1808, p. 146.*Calcarina baculatus* (?) (MONTFORT), Bull. 100, U. S. Nat. Mus., vol. 1, pt. 6, 1919, p. 364, pl. 44, fig. 3.

Description.—Test lenticular, biconvex, chambers numerous, in a closely coiled test, periphery rounded; supplementary skeleton well-developed, dorsal and ventral surfaces with numerous small knobs or bosses with finer spines between; periphery with three or four short, cylindrical, truncate spines, placed at equidistant points on the surface; surface of the spines covered with spinose projections.

Diameter up to 2 mm. or slightly more.

Montfort's localities for this species are the East Indies, Arabian Sea, and the Adriatic. He figures a rounded test with three cylindrical, truncate spinose projections at three of four equidistant points of the circumference, the exterior of the test black with white points on both the body and spines. This has been referred to *Tinoporos baculatus* with the description which in the remarks given makes note of the color as "blanche, flambée, et teintée de jaune." This, which is evidently not descriptive of the figured specimen, probably refers to the *Tinoporos* of most authors, which is here referred to as *Baculogypsina sphaerulatus*. The figured specimen, however, is evidently a *Calcarina*, as at one Philippine station especially (D5134, Sulu Archipelago, near Basilan Island), specimens were found exactly like the crude figure of Montfort, dark brown and black with spines appearing as light spots and the spinose projections as in the figure

and at equidistant points on the periphery. A study of the young and the development of older specimens of the same type from this locality show it to be a *Calcarina*. The name *baculatus* may be used probably to distinguish this species, although as to the generic portion it is very questionable as to its clearness and therefore its availability.

In its development it is seen that this species of *Calcarina* has a considerable development of supplementary skeleton, which, as in *C. hispida*, hides most of the earlier portion of the test.

Specimens referred to in this species were found in the Philippine collections from Jolo Jolo, Sulu Archipelago, vicinity of Siasi; Tawi Tawi Group; off Romblon; off eastern Panay; between Burias and Luzon; Pacific Ocean; east coast of Mindanao; Gulf of Davao; China Sea, off southern Luzon; Palawan Passage; Marinduque Island and vicinity; east coast of Luzon; north of Tawi Tawi; and Sibuko Bay, Borneo.

These are mostly in the southern portion of the Archipelago in shallow, warm water.

This species should be carefully distinguished and its occurrence recorded that more may be known of its actual distribution.

Calcarina baculatus (?)—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14458	U.S.N.M.	10+	D5134...	6 44 45 N.; 121 48 00 E...	25	° F.	fne. s.....	Common.
14459	U.S.N.M.	10+	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.....	Common.
14460	U.S.N.M.	10+	D5138...	6 06 00 N.; 120 58 50 E...	19	s., Co.....	Common.
14461	U.S.N.M.	7	D5139...	6 06 00 N.; 120 02 30 E...	20	co. s.....	Common.
14462	U.S.N.M.	10+	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Common.
14463	U.S.N.M.	10+	D5144...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Common.
14464	U.S.N.M.	5	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.....	Common.
14465	U.S.N.M.	4	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.....	Few.
14466	U.S.N.M.	10+	D5148...	5 35 40 N.; 120 47 30 E...	17	co. s.....	Common.
14467	U.S.N.M.	3	D5149...	5 33 00 N.; 120 42 10 E...	10	co. sh.....	Few.
14468	U.S.N.M.	1	D5151...	5 24 40 N.; 120 27 15 E...	24	co. s., sh.....	Rare.
14469	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Common.
14470	U.S.N.M.	10+	D5154...	5 14 50 N.; 119 58 45 E...	12	co. s.....	Common.
14471	U.S.N.M.	10+	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.....	Common.
14472	U.S.N.M.	10+	D5160...	5 12 40 N.; 119 55 10 E...	12	s.....	Common.
14473	U.S.N.M.	1	D5161...	5 10 15 N.; 119 53 00 E...	16	fne. s.....	Rare.
14474	U.S.N.M.	5	D5178...	12 43 00 N.; 122 06 15 E...	78	fne. s.....	Few.
14475	U.S.N.M.	6	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Few.
14476	U.S.N.M.	6	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fne. s.....	Few.
14477	U.S.N.M.	2	D5182...	11 30 40 N.; 123 23 20 E...	24	fne. s., m.....	Few.
14478	U.S.N.M.	8	D5218...	13 11 15 N.; 123 02 45 E...	20	co. s.....	Few.
14479	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s.....	Common.
14480	U.S.N.M.	10+	D5250...	7 05 07 N.; 125 39 45 E...	23	co. s.....	Common.
14481	U.S.N.M.	4	D5255...	7 03 00 N.; 125 39 00 E...	100	sft. m.....	Few.
14482	U.S.N.M.	6	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Few.
14483	U.S.N.M.	10+	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fne. s., m., sh.....	Common.
14484	U.S.N.M.	7	D5342...	10 56 55 N.; 119 17 24 E...	14-25	gy. m.....	Common.
14485	U.S.N.M.	1	D5371...	13 49 40 N.; 121 40 15 E...	83	gn. m.....	Rare.
14486	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E...	500	gy. m.....	Rare.
14487	U.S.N.M.	2	D5569...	5 03 15 N.; 120 15 30 E...	303	52.3	co. s.....	Rare.
14488	U.S.N.M.	6	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	fne. s.....	Few.
14489	U.S.N.M.	10+	D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fne. gy. s., gy. m.	Common.
14490	U.S.N.M.	1	D5592...	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.....	Rare.

CALCARINA HISPIDA H. B. Brady.

Plate 75, fig. 4.

Calcarina spengleri, hispid variety, CARPENTER, Philos. Trans., vol. 150, 1860, p. 551, pl. 19, figs. 8-11; pl. 20, figs. 6, 8; Introd. Foram., 1862, pl. 14, figs. 6, 7.

Calcarina hispida H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 713, pl. 108, figs. 8, 9.—LISTER, Philos. Trans., vol. 186, 1895, p. 437, pl. 8, figs. 34-37.—MILLETT, Journ. Roy. Micr. Soc., 1904, p. 597.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 72, pl. 29, figs. 4, 5; pl. 31, fig. 3.

Calcarina calcar, var. *hispida*, CARTER, Ann. Mag. Nat. Hist., ser. 5, vol. 5, 1880, p. 453.

Calcarina quoyi D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 276, No. 6.—FORNASINI, Mem. Accad. Sci. Bologna, ser. 6, vol. 5, 1908, p. 7, pl. 3, figs. 8, 9.

Description.—Test lenticular, biconvex, chambers numerous, in a close-coiled, generally flattened trochoid spire, of about three volutions, each with several chambers; the sutures indistinct, as well as the line of the spire, due to the great development of a supplementary skeleton, only the last two or three chambers being at all distinguishable from either side; as formed these are rounded with a thin, finely perforate wall; the general surface of the test very rough and hispid, with short flattened spines, projecting radially and often covering the spines as well; periphery of the test with a few spines which are usually somewhat flattened and short, widest at the base; aperture a series of pores at the base of the last-formed chamber.

Diameter up to 2 mm.

Brady originally described this species from among the islands of the Pacific, one station at 155 fathoms (283 meters), the remainder ranging from 3 fathoms (5 meters) or less to 37 fathoms (68 meters). Millett gives it from the Malay Archipelago with the note:

The examples are neither numerous or large.

In the Philippines it occurs at the following localities: China Sea, off southern Luzon; vicinity of Jolo, Sulu Archipelago; Tawi Tawi Group; off Romblon; between Panay and Negros; Sogod Bay; southern Leyte; between Marinduque and Luzon; Gulf of Davao; between Bohol and Cebu; east coast of Luzon; Darvel Bay, Borneo; and Binang Pool, Subin Bay.

The newly added chambers in this species are added directly on top of the hispid secondary skeleton, and this is produced so rapidly that rarely more than two chambers in the series retain their original surface before being so covered.

From the evidence of the "planchés inédites" d'Orbigny's *Calcarina quoyi* seems to be the same as *C. hispida* H. B. Brady; but as the former could not have been identified until the publication of the figures by Fornasini in 1907, it must remain a synonym of *C. hispida*.

Calcarina hispida—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14530	U.S.N.M.	7	D5113...	13 51 30 N.; 120 50 30 E..	159	dk. gn. m.	Common.
14531	U.S.N.M.	8	D5136...	6 04 20 N.; 120 59 20 E..	22	s., sh.	Common.
14532	U.S.N.M.	2	D5143...	6 05 50 N.; 121 02 15 E..	19	co. s.	Few.
14533	U.S.N.M.	2	D5144...	6 05 50 N.; 121 02 15 E..	19	co. s., sh.	Few.
14534	U.S.N.M.	10+	D5145...	6 04 30 N.; 120 59 30 E..	23	co. s., sh.	Common.
14535	U.S.N.M.	3	D5161...	5 10 15 N.; 119 53 00 E..	16	fne. s., blk. sp.	Few.
14536	U.S.N.M.	10+	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.	Common.
14537	U.S.N.M.	10+	D5181...	11 36 40 N.; 123 26 35 E..	26	m., fne. s.	Common.
14539	U.S.N.M.	10+	D5184...	10 18 30 N.; 122 23 30 E..	565	49.8	gn. m.	Common.
14540	U.S.N.M.	4	D5191...	10 29 45 N.; 123 31 15 E..	258	62.8	gn. m.	Few.
14541	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.	Rare.
14542	U.S.N.M.	5	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.	Few.
14543	U.S.N.M.	1	D5250...	7 05 07 N.; 125 39 45 E..	23	co. s.	Rare.
14538	U.S.N.M.	10+	D5282...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. gy. s.	Common.
14544	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 45 E..	88	co. s.	Rare.
14545	U.S.N.M.	2	D5412...	10 09 15 N.; 123 52 00 E..	162	54.8	gn. m.	Rare.
14546	U.S.N.M.	1	D5459...	13 10 21 N.; 123 59 54 E..	201	Rare.
14547	U.S.N.M.	2	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.	Rare.
14548	U.S.N.M.	10+	Binang Pool, Subin Bay, P. I.	Common.
14549	U.S.N.M.	3	Jolo Jolo, P. I.	Few.

Genus *SIDEROLITES* Lamarck, 1801.

Siderolites LAMARCK, Syst. Anim. sans Vert., 1801, p. 376 (type, *S. calcitrapoides* Lamarck).

From a study of material from Maestricht, the type locality for this genus, and the type species, it is evident that the genus *Siderolites* can be used in the present case. It is a test much like *Calcarina* but with a more accelerated development. The early chambers are close-coiled, rotaliform, then later with numerous large, irregular chambers, coarsely punctate, between which as support to this loose tissue are large bosses, starting near the center and growing radially to the periphery, made of solid shelly material, not very evident in well-preserved specimens, but becoming very distinct when erosion takes place; the periphery usually with three or four blunt spines, not in one place, but extending radially; apertures in the young, as in *Calcarina*, but in the adult formed by the coarse perforations of the outer chambers.

This is the genus to which it seems the *Calcarina tetraedra* of Gümbel should be referred, as well as the common Philippine species which seems to be identical with Gümbel's species. The genus goes back at least to the Eocene, and in the present oceans seems to be limited so far as is known to the general Philippine region.

Carpenter²⁵ figures well the characteristic structure of this genus, which he refers to as "Philippine variety" of *Tinoporus baculatus*.

²⁵ Introd. Foram., 1862, pl. 15, fig. 10.

SIDEROLITES TETRAEDRA (Gümbel).

Plate 75, fig. 5; plate 76, figs. 1-5.

Calcarina tetraedra GÜMBEL, Abh. bay. Akad. Wiss., vol. 10, 1868 (1870), p. 656, pl. 2, figs. 97a, b.

Siderolites tetraedra CUSHMAN, Bull. 100, U. S. Nat. Mus., vol. 1, pt. 6, 1919, pl. 44, fig. 5; pl. 45, figs. 1-5.

Description.—Test in the young, biconvex, rotaliform, composed of two or more volutions, with several chambers in each; later the supplementary skeleton greatly developed and the exterior with irregularly placed chambers, subglobular in shape and with coarsely perforated walls; between the chambers, solid radial portions giving support to the remainder of the test, and at four or five (usually four) points thick tapering, bluntly pointed spinose projections are placed, having their initiation in the early portion of the test, in the adult these often covered nearly to the tips with the supplementary skeleton, in such cases the whole test instead of spherical being concave and polyhedral; aperture in the young at the base of the chamber, in the adult formed by the large perforations of the chamber wall.

Diameter up to 6 mm.

In the Philippine collections this species is often very abundant in all its stages. It is most common in warm, shallow water. It has occurred at the following localities: Jolo Jolo; off Romblon; between Burias and Luzon; China Sea, off southern Luzon; Malam-paya Sound; Palawan Island; north of Tawi Tawi; and Darvel Bay, Borneo.

There seems to be little to separate this species from that described from the Eocene by Gümbel. The four tapering spines are very characteristic. Various stages in the development of the species are shown on plate 76, from the early *Calcarina* stage through the development of a few of the irregular chambers and gradually the covering of the surface with these. An eroded specimen is shown in figure 4, where the chambers of the exterior have been partially broken away leaving the heavier solid supporting portions as projections.

Carpenter figures this species as a "Philippine variety" of *Tinoporos baculatus*,²⁶ and makes notes of its differences from the "Australian variety" (*Baculogypsina sphaerulatus*). He showed the differences in the structure of the interior also.

From what I have seen of this species it is very common in the Philippines, but rare elsewhere, while the following genus and species is rare in the Philippines but very abundant in the southern region about Australia especially.

In its development *Siderolites* stands between *Calcarina* and *Baculogypsina*, and geologically also its first occurrence is between the two as far as is known.

²⁶ *Introd. Foram.*, 1862, pl. 15, figs. 8-10.

Siderolites tetraedra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14823	U.S.N.M.	4	D5142	6 06 10 N.; 121 02 40 E...	21	co. s., sh.	Few.
14824	U.S.N.M.	10+	D5149	5 33 00 N.; 120 42 10 E...	10	co., sh.	Common.
14825	U.S.N.M.	10+	D5154	5 14 50 N.; 119 58 45 E...	12	co. s.	Common.
14826	U.S.N.M.	10+	D5179	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.	Common.
14803	U.S.N.M.	7	D5181	11 36 40 N.; 123 26 35 E...	26	m., fine. s.	Common.
14804	U.S.N.M.	10+	D5218	13 11 15 N.; 123 02 45 E...	20	crs. s.	Common.
14805	U.S.N.M.	10+	D5276	13 49 15 N.; 120 14 15 E...	18	sh., p., s.	Common.
14806								
14807	U.S.N.M.	10+	D5282	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.	Common.
14808	U.S.N.M.	7	D5342	10 56 55 N.; 119 17 24 E...	14-25	gy. m.	Common.
14810	U.S.N.M.	9	D5571	5 30 45 N.; 120 07 57 E...	340	52.3	s., sh.	Common.
14811	U.S.N.M.	2	D5589	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co.	Rare.
14812	U.S.N.M.	3	D5589	4 12 10 N.; 118 38 08 E...	260	45.7	fine. gy. s.	Rare.
15670	U.S.N.M.	2	D5136	6 04 20 N.; 120 59 20 E...	22	s., sh.	Rare.
15671	U.S.N.M.	1	D5145	6 04 30 N.; 120 59 30 E...	23	co. s., sh.	Rare.
15672	U.S.N.M.	1	D5147	5 41 40 N.; 120 47 10 E...	21	co. s., sh.	Rare.
15673	U.S.N.M.	1	D5152	5 22 55 N.; 120 15 45 E...	34	wh. s.	Rare.
15674	U.S.N.M.	10+	D5172	6 03 15 N.; 120 35 30 E...	318	fine. s., sh.	Common.
15675	U.S.N.M.	2	D5220	13 38 00 N.; 121 58 00 E...	50	sft. gn. m.	Rare.
15676	U.S.N.M.	1	D5255	7 03 00 N.; 125 39 00 E...	100	sft. m.	Rare.
15677	U.S.N.M.	1	D5281	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s.	Rare.
15678	U.S.N.M.	1	D5381	13 14 15 N.; 122 44 45 E...	88	co. s.	Rare.

Genus *BACULOGYPSINA* Sacco, 1893.

Description.—Test in the very young, rotaliform, later irregular, with numerous small finely punctate chambers, with four to eight or even more sharp tapering spines; supplementary skeleton greatly developed, at the surface, when well preserved, with bosses of clear shell material united with surrounding ones by radial connecting portions of the same sort of material, making a reticulate marking standing out slightly above the surface.

This genus represents the single living species which is found in enormous numbers in the southern islands, especially off Australia, but is rare in the Philippines.

In its development *Baculogypsina* is more accelerated than *Calcarina* and *Siderolites*, and evidently represents a higher and later development in the same series.

BACULOGYPSINA *SPHAERULATUS* (Parker and Jones).

Plate 75, fig. 6.

Baculogypsina sphaerulatus CUSHMAN, Bull. 100, U. S. Nat. Mus., vol. 1, pt. 6, 1919, pl. 44, fig. 6.

Description.—Test generally biconvex, the periphery somewhat acute, with four to eight or more rather sharp short spines, surface covered with a reticulated pattern of slightly projecting bosses, with connecting lines of clear shell material; early development consisting of one volution or slightly more of numerous chambers coiled as in *Calcarina*, after which the supplementary chambers are filled in rapidly on the surface, and the network of bosses and connecting rods built up; the radial spines are always smooth except for the slight channeling of the surface; color in fresh material often orange or yellow.

Diameter up to 3 mm.

From all the Philippine collections examined this has been found but a few times, and then in small numbers. It is abundant farther south, especially in the South Pacific islands and off Australia, from where I have had abundant material. Millett records it as very rare in the Malay Archipelago collections, finding it at but one station. This accords with the results I have had from the Philippines and with Carpenter, who speaks of it as—

found in greatest size and abundance in the Australian and Polynesian Seas, occurring chiefly in the shelly sands of rather shallow water; and * * * does not appear to range far beyond those regions.

Genus GYPSINA Carter, 1877.

GYPSINA GLOBULUS (Reuss).

Ceripora globulus REUSS, Haidinger's Nat. Abh., vol. 2, 1847, p. 33, pl. 5, fig. 7.
Gypsina globulus H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 717, pl. 101, fig. 8.—CHAPMAN, Proc. Zool. Soc., London, 1895, p. 44; Journ. Linn. Soc. London, Zoology, vol. 28, 1900, p. 198.—HERON-ALLEN and EARLAND, Proc. Roy. Irish Acad., vol. 31, pt. 64, 1913, p. 140, pl. 13, fig. 10; Trans. Zool. Soc. London, vol. 20, 1915, p. 727.

Description.—Test spherical or generally globular; surface regularly pitted with a more or less regular polygonal network; earliest chambers spiral, followed quickly by a mass of chambers covering the entire exterior; outer chamber walls finely perforated, forming the aperture of the test; color white.

Diameter up to 4 mm.

Specimens have not been at all abundant in the Philippine collections. There are records from five stations from the following localities: Sulu Archipelago, off Tawi Tawi; off Jolo; off Romblon; and off eastern Panay. These range in depth from 24 to 318 fathoms (44 to 582 meters), three of them in less than 35 fathoms (66 meters).

Gypsina globulus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
12433	U.S.N.M.	9	D5151...	5 24 40 N.; 120 27 15 E...	24	co. s., sh....	Few.
12492	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Common.
15763	U.S.N.M.	2	D5172...	6 03 15 N.; 120 35 30 E...	318	fine. s., sh....	Rare.
15763	U.S.N.M.	2	D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.....	Few.
12495	U.S.N.M.	1	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fine. s....	Few.

Genus CARPENTERIA Gray, 1858.

CARPENTERIA UTRICULARIS (Carter).

Plate 73, figs. 4, 5.

Polytremia utricularis CARTER, Ann. Mag. Nat. Hist., ser. 4, vol. 17, 1876, p. 211, pl. 13, figs. 11–17.

Carpenteria utricularis CARTER, Ann. Mag. Nat. Hist., ser. 4, vol. 20, 1877, p. 176.—
H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 678, pl. 99, figs. 6, 7; pl. 100, figs. 1-4.

This species, which was found by the *Challenger* in its best development off the Admiralty Islands, and which are allied in their faunal characters with that of the Philippines, has occurred at two stations in the Sulu Archipelago, D5146, off Siasi, in 24 fathoms (44 meters), and D5154, off Tawi Tawi, in 12 fathoms (22 meters). The specimens were not common.

Carpenteria utricularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5146...	° ' " ° ' "	24	° F.	co. s., sh.....	Few.
			D5154...	5 46 40 N.; 120 48 50 E..	12	co. s.....	Few.
				5 14 50 N.; 119 58 45 E..				

CARPENTERIA PROTEIFORMIS Goës.

Plate 73, figs. 2, 3.

Carpenteria balaniformis, var. *proteiformis* GOËS, Kongl. Svensk. Vet. Akad. Handl., vol. 19, 1882, p. 94, pl. 6, figs. 208-214; pl. 7, figs. 215-219.

Carpenteria proteiformis H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 679, pl. 97, figs. 8-14.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 49, pl. 20, fig. 2; pl. 21, fig. 1.

This species seems to be a common one in the Philippine material. It is a characteristic species of the coral-reef regions, and is also found to considerable depth. There seem to be two forms which are characterized by the ornamentation of the test. One of these has the test with coarse reticulate ornamentation and large intermediate portions; the other is very thinly perforate. This latter form occasionally has a secondary thinning of the wall, developing large papillae.

Similar specimens are found in the region about New Zealand, and the two may represent distinct species.

This species has been found at 25 stations in the area, abundant at some of them; these range in depth from 18 to 890 fathoms (33 to 1,628 meters), the average depth being 190 fathoms (348 meters).

Bottom temperatures range from 46.2° F. to 75.7° F. (7.8° C. to 24.2° C.), with the average 54.6° F. (12.5° C.). These stations are mostly in the region of the Archipelago, although it is common in the Sulu region and also to the southward in Darvel Bay, Borneo, and Buton Strait.

Carpenteria proteiformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11929	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N.; 52° E., 1.50 miles.	38	° F.	gn. m., s....	Common.
15696	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Frequent.
11933	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Rare.
11930	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Common.
11927								
11936	U.S.N.M.	7	D5213...	12 15 00 N.; 123 57 30 E...	80	s., m., sh....	Rare.
11934	U.S.N.M.	1	D5276...	13 49 15 N.; 120 14 15 E...	18	sh., p., s....	Few.
11928	U.S.N.M.	4	D5277...	13 56 55 N.; 120 13 45 E...	80	53.6	fine s.....	Few.
11931	U.S.N.M.	10+	D5338...	11 37 45 N.; 119 46 00 E...	46	s., m.....	Common.
11925	U.S.N.M.	10+	D5338...	11 33 45 N.; 119 24 25 E...	43	Co., s., m....	Common.
15698	U.S.N.M.	8	D5478...	10 46 24 N.; 125 16 30 E...	57	sh.....	Common.
11925	U.S.N.M.	7	D5571...	5 30 45 N.; 120 07 57 E...	340	52.3	s., sh.....	Few.
15698	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Rare.
11964	U.S.N.M.	4	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fine s., Co...	Rare.
15701	U.S.N.M.	4	D5590...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co...	Rare.
13216	U.S.N.M.	1	D5582...	4 19 54 N.; 118 58 38 E...	890	33.3	gy. m., fine s.	Rare.
13217	U.S.N.M.	3	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh...	Rare.
11932	U.S.N.M.	3	D5642...	4 31 40 S.; 122 49 42 E...	37	gy. m.....	Rare.
D5097...			14 19 15 N.; 120 33 52 E...	30	gy. m., s....	Few.	
			D5132...	Island off Panabutan Point, N.; 15° W., 0.3 mile.	26	gr. m., s....	Frequent.
			D5178...	12 43 00 N.; 122 06 15 E...	78	fine s.....	Frequent.
15697	U.S.N.M.	1	D5255...	7 03 00 N.; 125 39 00 E...	100	sft. m.....	Rare.
15699	U.S.N.M.	1	D5426...	9 12 00 N.; 118 28 00 E...	27	fine gy. s....	Rare.
15700	U.S.N.M.	1	D5446...	12 43 51 N.; 124 59 18 E...	300	gn. m.....	Rare.
			D5574...	5 30 45 N.; 120 07 57 E...	340	Rare.

CARPENTERIA MONTICULARIS Carter.

Carpenteria monticularis CARTER, Ann. Mag. Nat. Hist., ser. 4, vol. 19, 1877, p. 211, pl. 13, figs. 9-12.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 677, pl. 99, figs. 1-5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 48, pl. 20, fig. 3.

The specimens are very typical but not common in the area. Some of these are fine and large and agree very closely with the specimens figured by Brady (*Challenger*, pl. 99, figs. 1-5).

These occurred at the following stations: D5154, off Tawi Tawi, 12 fathoms (22 meters); D5192, off northern Cebu, 32 fathoms (59 meters); D5217, between Burias and Luzon, 105 fathoms (193 meters); and D5269, Verde Island Passage, 220 fathoms (403 meters).

Carpenteria monticularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
15692	U.S.N.M.	2	D5154...	5 14 50 N.; 119 58 45 E...	12	co. s.....	Rare.
			D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	
	U.S.N.M.	1	D5217...	13 00 20 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Rare.
15693			D5269...	13 39 50 N.; 120 59 30 E...	220	fine s., p....	

CARPENTERIA RHAPHIDODENDRON Moebius.

Carpenteria raphidodendron MOEBIUS, Beitr. Meeresfauna Insel. Mauritius, 1880, p. 81, pl. 5, figs. 6-10; pl. 6, figs. 1-6.

Ramulina herdmanni DAKIN, Ceylon Pearl Oyster Fisheries, Suppl. Rept., pt. 5, 1906, p. 236, pl., figs. 1-6, text figs. p. 227.

This species, which seems to be rare in the Indo-Pacific region about coral reefs, is recorded by Chapman from Funafuti, and occurs in the Philippine region from *Albatross* station D5154, Sulu Archipelago, off Tawi Tawi, in 12 fathoms (22 meters), and is very common at D5357, north of Balabac Strait, 68 fathoms (125 meters). This is a very peculiar species, at first not resembling the foraminifera, but apparently belonging to this genus. Some of the largest specimens measure nearly 20 mm. in height. Figures given by Chapman show well the characters of its form and arrangement of chambers. The big rootlike offset chambers are characteristic in all the large specimens, apparently having a definite use in supporting the rest of the column. This is probably the same as the species described by Dakin²⁷ from off Ceylon, as a comparison of his figures will show.

Carpenteria raphidodendron—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15694	U.S.N.M.	2	D5154...	5 14 50 N.; 119 58 45 E...	12	° F.	Co. s.....	Rare.
15695	U.S.N.M.	10+	D5357...	8 06 00 N.; 117 17 10 E...	68	Co., s.....	Frequent.

Genus POLYTREMA Risso, 1826.

POLYTREMA MINIACEUM (Gmelin).

Millepora miniacea LINNÆUS, Syst. Nat., ed. 13 (Gmelin's), vol. 1, pt. 6, 1788, p. 3784, No. 6.

Polytrema miniaceum BLAINVILLE, Dict. Sci. Nat., vol. 42, 1826, Atlas, Zooph., vol. 1, p. 17.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 721, pl. 100, figs. 5-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 5, 1915, p. 75, pl. 18, fig. 6; pl. 20, fig. 4.—HICKSON, Trans. Linn. Soc. London, Zoology, vol. 14, 1911, p. 453, pl. 30, fig. 1; pl. 31, fig. 8; pl. 32, figs. 18, 23, 27, 31.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 728.

This species, which is characteristic of coral-reef regions, is very common, attached to shell fragments and other objects in shallow water, except in the southern part of the Archipelago. Although probably widely distributed in the region it has been recorded and material saved from the following stations: D5146, Sulu Archipelago, off Siasi, 24 fathoms (44 meters); D5179, off Romblon, 37 fathoms (68 meters); D5217, between Burias and Luzon, abundant at 105 fathoms (193 meters); and D5357, north of Balabac Strait, 68 fathoms

²⁷ Ceylon Pearl Oyster Fisheries, Suppl. Rep., pt. 5, 1906.

(125 meters). Some of the specimens are nearly white in color, but most of them are of the reddish color characteristic of the species.

Polytrema miniaceum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15689	U.S.N.M.	2	D5146...	5 46 40 N.; 120 48 50 E...	21	co. s., sh.....	Rare.
15764	U.S.N.M.	5	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Few.
15690	U.S.N.M.	10+	D5217...	13 21 00 N.; 123 14 15 E...	105	63.1	ers. gy. s.....	Frequent.
15691	U.S.N.M.	3	D5357...	8 06 00 N.; 117 17 10 E...	68	Co., s.....	Rare.

Genus HOMOTREMA Hickson, 1911.

This genus has been separated by Hickson from *Polytrema* on the basis of the structure of the test. The Philippine material carries out the points of structure which he discusses in the fullest manner.

The following is the generic description given by Hickson:

The surface is marked by clearly defined areolae about 0.1 mm. in diameter, perforated by a large number of small foramina, 0.001 mm. in diameter. The boundaries of the areolae are solid, and there are no pillar pores. Below the surface there may be seen a number of chambers communicating with one another by large open passages and bounded by solid walls. There are no hollow pillars and no foramina except those on the outer walls of the superficial chambers.

HOMOTREMA RUBRUM (Lamarck).

Millepora rubra LAMARCK, Anim. sans Vert., vol. 2, 1816, p. 202.

Polytrema rubra CARPENTER, PARKER, and JONES, Introd. Foram., 1862, p. 235, pl. 13, figs. 18–20.

Homotrema rubrum HICKSON, Trans. Linn. Soc. London, Zoology, vol. 14, 1911, p. 454, pl. 30, fig. 2; pl. 31, fig. 9; pl. 32, figs. 19, 22, 28.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 729.

The Philippine material, although occurring at a few stations as far as the available material shows, is very characteristic, having the form and the deep red color as well as the finer structure of the species, as suggested by Hickson. The records include the following four stations: D5109, China Sea off southern Luzon, 10 fathoms (18 meters); D5136, vicinity of Jolo, 22 fathoms (40 meters); D5146, Sulu Archipelago off Siasi, 24 fathoms (44 meters); and D5179, off Romblon, 37 fathoms (68 meters).

Homotrema rubrum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15686	U.S.N.M.	2	D5109...	14 03 45 N.; 120 16 30 E...	10	Co.....	Rare.
15687	U.S.N.M.	10+	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.....	Frequent.
15688	U.S.N.M.	8	D5146...	5 46 40 N.; 120 48 50 E...	24	co. s., sh.....	Frequent.
			D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Frequent.

Genus SPORADOTREMA Hickson, 1911.

The original description by Hickson is as follows:

The surface of the stem, and, in many cases, of the proximal parts of the branches as well, are not marked by areolae at all. The foramina are scattered irregularly on the surface and are of relatively large size. There are no pillar pores. Below the surface there may be seen a number of chambers communicating with one another by large open passages and bounded by solid walls. There are no hollow pillars and no foramina except those on the outer walls of the superficial chambers.

SPORADOTREMA CYLINDRICUM (Carter).

Polytrema cylindrica CARTER, Ann. Mag. Nat. Hist., ser. 5, vol. 5, 1880, p. 441, pl. 18, figs. 1a-g.

Sporadotrema cylindricum HICKSON, Trans. Linn. Soc. London, Zoology, vol. 14, 1911, p. 454, pl. 30, figs. 3, 4, 6, 7; pl. 31, figs. 10, 13-16; pl. 32, figs. 20, 21, 24, 29, 32, 33.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 729.

The Philippine specimens of this species show the orange-yellow color characteristic of this species, as described and figured by Hickson. As far as the records show the species is not common and a few specimens have been found at the following stations: D5146, Sulu Archipelago off Siasi, 24 fathoms (44 meters); D5179, off Romblon, 37 fathoms (68 meters); and D5357, north of Balabac Strait, 68 fathoms (125 meters).

This genus can be easily distinguished through the generic description given by Hickson preceding this.

Sporadotrema cylindricum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
15683	U.S.N.M.	1	D5146...	5 46 40 N.; 120 48 50 E..	24	co. s., sh.....	Rare.
15684	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Rare.
15685	U.S.N.M.	2	D5357...	8 06 00 N.; 117 17 10 E..	68	Co., s.....	Rare.

Family NUMMULITIDAE.

Genus NONIONINA d'Orbigny, 1826.

NONIONINA UMBLICATULA (Montagu.)

Plate 74, figs. 3a, b.

Nautilus umblicatulus MONTAGU, Test. Brit., 1803, p. 191.

Nonionina umblicatula PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 4, vol. 8, 1871, p. 242, pl. 12, fig. 157.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 726, pl. 109, figs. 8, 9.—CUSHMAN, Bull. 71, U.S. Nat. Mus., pt. 4, 1914, p. 24, pl. 17, fig. 1.

Rather typical specimens of this species occurred at 13 stations in the region, ranging in depth from 105 to 500 fathoms (193 to 914

meters), the average depth being 302 fathoms (550 meters). Bottom temperatures are given for 8 of these stations, ranging from 41.2° F. to 63.1° F. (5.1° C. to 17.2° C.), the average temperature being 51.1° F. (10.6° C.).

Besides the few stations in the Archipelago this has also occurred off Darvel Bay, between Gillolo and Makyan Islands, and in Macassar Strait.

Nonionina umbilicatulula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bot- tom tem- per- ature.	Character of bottom.	Abundance.
						° F.		
15528	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E..	318	fne. s., sh....	Rare.
15529	U.S.N.M.	1	D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s....	Rare.
15530	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s....	Rare.
15531	U.S.N.M.	8	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Frequent.
15532	U.S.N.M.	6	D5309...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Frequent.
15533	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Rare.
15534	U.S.N.M.	3	D5445...	12 44 42 N.; 124 59 50 E..	383	44.3	gn. m., s....	Rare.
15535	U.S.N.M.	3	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.(net)..	Few.
15536	U.S.N.M.	6	D5565...	5 51 42 N.; 120 30 30 E..	243	52.3	s., ptr. sh....	Frequent.
15537	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Rare.
15538	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	bk. s., Co....	Rare.
15539	U.S.N.M.	2	D5622...	Makyan Island, N. 9° E..	275	gy. m.....	Rare.
15540	U.S.N.M.	2	D5667...	2 56 00 S.; 118 47 30 E..	367	41.7	gy. s., m....	Rare.

NONIONINA POMPILIOIDES (Fichtel and Moll).

Nautilus pompilioides FICHEL and MOLL, Test. Micr., 1803, p. 31, pl. 2, figs. a-e.

Nonionina pompilioides PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 3, vol. 16, 1865, p. 18, pl. 3, fig. 98; ser. 4, vol. 8, 1871, p. 246, pl. 12, fig. 158.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 727, pl. 109, figs. 10, 11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 25, pl. 17, fig. 2.

The only station at which this species is recorded is D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

Nonionina pompilioides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bot- tom tem- per- ature.	Character of bottom.	Abundance.
						° F.		
15525	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s....	Rare.

NONIONINA BOUEANA d'Orbigny.

Nonionina boueana D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 108, pl. 5, figs. 11, 12.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 729, pl. 109, figs. 12, 13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 28, pl. 16, fig. 1.

Rare specimens occur at the following stations: D5217, between Burias and Luzon, 105 fathoms (193 meters), D5201, China Sea, off

Hongkong, 554 fathoms (1,012 meters); D5570, north of Tawi Tawi, 330 fathoms (604 meters); and in shallow water in Little Harbor, south of Luneta, Manila Harbor.

The specimens are characteristic.

Nonionina boueana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15526	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
15766	U.S.N.M.	1	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Rare.
15527	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fine. s., glob..	Rare.
15767	U.S.N.M.	1	Manila Bay, P. I.....	Rare.

Genus POLYSTOMELLA Lamarck, 1822.

POLYSTOMELLA SUBNODOSA (Münster).

Robulina subnodosa MÜNSTER, Neues Jahrb. für Min., 1838, p. 391, pl. 3, fig. 61.

Polystomella subnodosa REUSS, Sitz. Akad. Wiss. Wien, vol. 18, 1855, p. 240, pl. 4, figs. 51a, b.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 734, pl. 110, figs. 1a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 32, pl. 14, fig. 8.

Very characteristic specimens of this species have been noted at 13 stations in the area, ranging in depth from 12 to 138 fathoms (22 to 252 meters), the average 41 fathoms (75 meters). The bottom temperatures at three stations ranged from 52.4° F. to 63.1° F. (11.3° C. to 17.2° C.); the average temperature 57.9° F. (14.5° C.).

All these specimens are in the region of the Archipelago, the most of them about the Sulu Sea, in comparatively shallow water. It does not seem to be common, however, at any of these stations, and is not nearly as common as the other two species of this genus in the same general area.

Polystomella subnodosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15541	U.S.N.M.	2	D5109...	14 03 45 N.; 120 16 30 E...	12	Co.....	Rare.
15542	U.S.N.M.	1	D5139...	6 06 00 N.; 121 02 30 E...	20	co. s.....	Rare.
15543	U.S.N.M.	1	D5144...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Rare.
15544	U.S.N.M.	1	D5145...	6 04 30 N.; 120 59 30 E...	23	co. sh.....	Rare.
15545	U.S.N.M.	1	D5148...	5 35 40 N.; 120 47 30 E...	17	co. s.....	Rare.
15546	U.S.N.M.	7	D5156...	5 12 50 N.; 119 55 55 E...	18	fine. s., sh...	Frequent.
15547	U.S.N.M.	3	D5160...	5 12 40 N.; 119 55 10 E...	12	s.....	Rare.
15548	U.S.N.M.	2	D5161...	5 10 15 N.; 119 53 00 E...	16	fine. s., blk.sp.	Rare.
15549	U.S.N.M.	10+	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Common.
15550	U.S.N.M.	10+	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Frequent.
15551	U.S.N.M.	1	D5220...	13 38 00 N.; 121 58 00 E...	50	52.4	gn. m.....	Rare.
15552	U.S.N.M.	1	D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh....	Rare.
15553	U.S.N.M.	2	D5546...	6 06 48 N.; 121 28 32 E...	138	58.3	fine. co. s....	Rare.

POLYSTOMELLA CRISPA (Linnaeus).

Nautilus crispus LINNAEUS, Syst. Nat., ed. 12, 1767, p. 1162.

Polystomella crisa LAMARCK, Anim. sans Vert., vol. 7, 1822, p. 625, No. 1.—H. B.

BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 736, pl. 110, figs. 6, 7.—

CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 32, pl. 18, fig. 1.

This is one of the species which is common in shallow water of the tropics, especially in the Indo-Pacific. It has been found at 36 stations in the area, ranging in depth from 10 to 554 fathoms (18 to 1,012 meters), with the average depth 122 fathoms (222 meters). Bottom temperatures are given at eight stations, ranging from 50° F. to 75.7° F. (10.2° C. to 24.2° C.), the average temperature 57.5° F. (14.2° C.). All but one of these stations are in the Philippines proper, except D5640, Buton Strait, and this is also in shallow water, the depth being 24 fathoms (44 meters).

Those stations at which it occurred in greatest abundance are all under 20 fathoms (37 meters), showing that the general habitat is in comparatively shallow water.

Polystomella crisa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
15554	U.S.N.M.	5	D5136...	6 04 20 N.	120 59 20 E.	22	° F.	s., sh.	Few.
15555	U.S.N.M.	10+	D5138...	6 06 00 N.	120 58 50 E.	19		s., Co	Common.
15556	U.S.N.M.	5	D5139...	6 06 00 N.	121 02 30 E.	20		co. s.	Common.
15557	U.S.N.M.	1	D5141...	6 09 00 N.	120 58 00 E.	29		co. s.	Rare.
15558	U.S.N.M.	6	D5142...	6 06 10 N.	121 02 40 E.	21		co. sh.	Few.
15559	U.S.N.M.	6	D5143...	6 05 50 N.	121 02 15 E.	19		co. sh.	Few.
15560	U.S.N.M.	10	D5144...	6 05 50 N.	121 02 15 E.	19		co. sh.	Common.
15561	U.S.N.M.	5	D5145...	6 04 30 N.	120 59 30 F.	23		co. sh.	Common.
15562	U.S.N.M.	10+	D5148...	5 35 40 N.	120 47 30 E.	17		co. sh.	Common.
15563	U.S.N.M.	6	D5151...	5 24 40 N.	120 27 15 E.	24		co. sh.	Common.
15564	U.S.N.M.	10+	D5154...	5 14 50 N.	119 58 45 E.	12		co. s.	Common.
15565	U.S.N.M.	9	D5156...	5 12 50 N.	119 55 55 E.	18		fne. s., sh.	Common.
15566	U.S.N.M.	6	D5159...	5 11 50 N.	119 54 00 E.	10		co. s.	Common.
15567	U.S.N.M.	2	D5160...	5 12 40 N.	119 55 10 E.	12		s.	Rare.
15568	U.S.N.M.	8	D5161...	5 10 15 N.	119 53 00 E.	16		fne. s.	Common.
15569	U.S.N.M.	8	D5172...	6 03 15 N.	120 35 30 E.	318		fne. s., sh.	Common.
15570	U.S.N.M.	5	D5178...	12 43 00 N.	122 06 15 E.	78		fne. s., sh.	Common.
15571	U.S.N.M.	10+	D5179...	12 38 15 N.	122 12 38 E.	37	75.7	hrd. s.	Common.
15572	U.S.N.M.	4	D5181...	11 36 40 N.	123 26 35 E.	26		m., fne. s.	Common.
15573	U.S.N.M.	6	D5192...	11 09 15 N.	123 50 00 E.	32		gn. s.	Common.
15574	U.S.N.M.	1	D5201...	10 10 00 N.	125 04 15 E.	554	52.8	gy. s., m.	Rare.
15575	U.S.N.M.	4	D5250...	7 05 07 N.	125 39 45 E.	23		co. s.	Few.
15576	U.S.N.M.	4	D5311...	21 33 00 N.	116 15 00 E.	88		crs. s.	Few.
15377	U.S.N.M.	10+	D5348...	10 57 45 N.	118 38 15 E.	375	56.4	co. s.	Common.
15578	U.S.N.M.	3	D5358...	6 06 40 N.	118 18 15 E.	39		m.	Few.
15579	U.S.N.M.	5	D5368...	13 35 30 N.	121 43 00 E.	181		gy. m.	Few.
15580	U.S.N.M.	3	D5380...	13 02 45 N.	122 29 00 E.				Few.
15581	U.S.N.M.	1	D5385...	13 24 50 N.	123 03 70 E.	327	62.4	gy. m.	Few.
18582	U.S.N.M.	1	D5430...	9 49 40 N.	119 03 20 E.	464	50.0	glob. ooze.	Few.
15583	U.S.N.M.	3	D5469...	13 36 48 N.	123 38 24 E.	500			Few.
15584	U.S.N.M.	3	D5481...	10 27 30 N.	125 17 10 E.	61		s., sh.	Few.
15585	U.S.N.M.	1	D5571...	5 30 45 N.	120 07 57 E.	340	52.3	s., sh.	Few.
15586	U.S.N.M.	3	D5576...	5 25 56 N.	120 03 39 E.	277	53.3	s.	Few.
15587	U.S.N.M.	1	D5640...	4 27 00 S.	122 55 40 E.	24		s., blk. sh.	Few.
13345	U.S.N.M.	2	D5580...	4 52 45 N.	119 06 45 E.	162	55.8	br. s., Co.	Few.
13341	U.S.N.M.	1		Cagayan, P. I.					Few.

POLYSTOMELLA CRATICULATA (Fichtel and Moll).

Nautilus craticulatus FICHEL and MOLL, Test. Micr., 1803, p. 51, pl. 5, figs. h-k.

Polystomella craticulata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 284, No. 3.—

H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 739, pl. 110, figs.

16, 17.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 34, pl. 19, fig. 4.

This is the most common species of the genus and one of the most common of the foraminifera in the whole region. It is distinctively characteristic of tropical shallow water and reaches exceptionally fine development in this region. It occurred at 55 stations, ranging in depth from 10 to 565 fathoms (18 to 1,038 meters), the average depth 95 fathoms (174 meters). Bottom temperatures are given at but 12 stations, ranging from 47.4° F. to 75.7° F. (8.5° C. to 24.2° C.), the average temperature 57.1° F. (13.9° C.).

It is much more abundant at the stations in depths of 25 fathoms (46 meters) or less, and apparently is found in very shallow water.

Polystomella craticulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
15388	U. S. N. M.	1	D5096...	14 20 23 N.;	120 34 15 E.	28	gy. m., s., sh.	Rare.
15389	U. S. N. M.	10+	D5106...	14 23 55 N.;	120 32 33 E.	37	gy. m.	Common.
15390	U. S. N. M.	8	D5109...	14 03 45 N.;	120 16 30 E.	10	Co.	Common.
15391	U. S. N. M.	1	D5110...	13 59 29 N.;	120 75 45 E.	135	59	dk. gy. m.	Rare.
15392	U. S. N. M.	10	D5113...	13 51 30 N.;	120 50 30 E.	159	dk. gn. m.	Common.
15393	U. S. N. M.	10+	D5132...	Island off Panabutan Point, N. 15° W., 0.3 m.		26	gn. m., s.	Common.
15594	U. S. N. M.	10+	D5133...	Island off Panabutan Point, N. 52° E., 1.5 m.		38	gn. m., s.	Common.
15595	U. S. N. M.	10	D5134...	6 44 45 N.;	121 48 00 E.	25	fne. s.	Common.
15596	U. S. N. M.	9	D5136...	6 04 20 N.;	120 59 20 E.	22	s., sh.	Common.
15597	U. S. N. M.	10+	D5138...	6 06 00 N.;	120 58 50 E.	19	s., Co.	Common.
15598	U. S. N. M.	2	D5139...	6 06 00 N.;	121 02 30 E.	20	co. s.	Rare.
15599	U. S. N. M.	4	D5141...	6 09 00 N.;	120 58 00 E.	29	co. s.	Rare.
15600	U. S. N. M.	10+	D5142...	6 06 10 N.;	121 02 40 E.	21	co. s., sh.	Common.
15601	U. S. N. M.	10+	D5143...	6 05 50 N.;	121 02 15 E.	19	co. s.	Common.
15602	U. S. N. M.	10+	D5144...	6 05 50 N.;	121 02 15 E.	19	co. s.	Common.
15603	U. S. N. M.	10+	D5145...	6 04 30 N.;	120 59 30 E.	23	co. s., sh.	Common.
15604	U. S. N. M.	3	D5146...	5 46 40 N.;	120 48 50 E.	24	co. s., sh.	Few.
15605	U. S. N. M.	1	D5147...	5 41 40 N.;	120 47 10 E.	21	co. s., sh.	Rare.
15606	U. S. N. M.	10+	D5148...	5 35 40 N.;	120 47 30 E.	17	co. s.	Common.
15607	U. S. N. M.	9	D5151...	5 24 40 N.;	120 27 15 E.	24	co. s., sh.	Common.
15608	U. S. N. M.	5	D5152...	5 22 55 N.;	120 15 45 E.	34	wh. s.	Few.
15609	U. S. N. M.	1	D5153...	5 18 10 N.;	120 02 55 E.	49	co. s., sh.	Few.
15610	U. S. N. M.	10+	D5154...	5 14 50 N.;	119 58 45 E.	12	co. s.	Common.
15611	U. S. N. M.	10+	D5156...	5 12 50 N.;	119 55 55 E.	18	fne. s., sh.	Common.
15612	U. S. N. M.	10+	D5159...	5 11 50 N.;	119 54 00 E.	10	co. s.	Common.
15613	U. S. N. M.	5	D5160...	5 12 40 N.;	119 55 10 E.	12	s.	Common.
15614	U. S. N. M.	5	D5161...	5 10 15 N.;	119 53 00 E.	16	fne. s.	Common.
15615	U. S. N. M.	10+	D5178...	12 43 00 N.;	122 06 15 E.	78	fne. s.	Common.
15616	U. S. N. M.	10+	D5179...	12 38 15 N.;	122 12 30 E.	37	75.7	hrd. s.	Common.
15617	U. S. N. M.	10+	D5181...	11 36 40 N.;	123 26 35 E.	26	m., fne. s.	Common.
15618	U. S. N. M.	2	D5182...	11 30 40 N.;	123 23 20 E.	24	fne. s., m.	Rare.
15619	U. S. N. M.	2	D5184...	10 18 30 N.;	122 23 30 E.	565	49.8	gn. m.	Rare.
15620	U. S. N. M.	10+	D5192...	11 09 15 N.;	123 50 00 E.	32	gn. s.	Common.
15621	U. S. N. M.	4	D5201...	10 10 00 N.;	125 04 15 E.	554	52.8	fne. s., m.	Few.
15622	U. S. N. M.	10+	D5218...	13 11 15 N.;	123 02 45 E.	20	crs. s.	Common.
15623	U. S. N. M.	2	D5220...	13 38 00 N.;	121 00 58 E.	50	sft. gn. m.	Rare.
15624	U. S. N. M.	5	D5288...	13 42 00 N.;	120 58 25 E.	170	p., s., sh.	Rare.
15625	U. S. N. M.	10+	D5276...	13 49 15 N.;	120 14 45 E.	18	sh., p., s.	Common.
15626	U. S. N. M.	4	D5277...	13 56 55 N.;	120 13 45 E.	80	fne. s.	Few.
15627	U. S. N. M.	1	D5282...	13 53 00 N.;	120 26 45 E.	248	47.4	dk. gy. s.	Rare.
15628	U. S. N. M.	3	D5311...	21 33 00 N.;	116 15 00 E.	88	crs. s., sh.	Rare.
15629	U. S. N. M.	2	D5338...	11 33 45 N.;	119 24 45 E.	43	Co., s., m.	Rare.
15630	U. S. N. M.	10+	D5348...	10 57 45 N.;	118 38 15 E.	375	56.4	gn. m.	Common.
15631	U. S. N. M.	2	D5358...	6 06 40 N.;	118 18 15 E.	39	m.	Rare.
15632	U. S. N. M.	1	D5419...	9 58 30 N.;	123 46 00 E.	175	54.5	gn. m.	Rare.
15633	U. S. N. M.	1	D5426...	9 12 00 N.;	118 28 00 E.	27	fne. gy. s.	Rare.
15634	U. S. N. M.	1	D5481...	10 27 30 N.;	125 17 10 E.	61	s., sh., g.	Rare.
15635	U. S. N. M.	1	D5538...	9 08 15 N.;	123 23 20 E.	256	53.3	gn. m., s.	Rare.
15636	U. S. N. M.	3	D5569...	5 33 15 N.;	120 15 30 E.	303	52.3	co. s.	Rare.
15637	U. S. N. M.	2	D5574...	5 30 45 N.;	120 07 57 E.	340	52.3	s.	Rare.
15638	U. S. N. M.	1	D5576...	5 25 56 N.;	120 03 39 E.	277	53.3	s.	Rare.
15639	U. S. N. M.	1	D5580...	4 52 45 N.;	119 06 45 E.	162	55.8	br. s., Co.	Rare.
15640	U. S. N. M.	3	D5640...	4 27 00 S.;	122 55 40 E.	21	s., brk. sh.	Rare.
15641	U. S. N. M.	10+	Taeloban Anchorage, P. I.		Common.
13340	U. S. N. M.	10+	Cebu.		Common.
13344	U. S. N. M.	10+	Common.

Genus AMPHISTEGINA d'Orbigny, 1826.

A reference to the various figures and descriptions given by Brady and by the subsequent authors will show that the genus *Amphistegina* includes numerous things. Brady refers all the forms of the genus to *Amphistegina lessonii*. Heron-Allen and Earland in their Kerimba paper reference (p. 736) distinguish three forms and mention a fourth. Fornasini²⁸ has given figures of the various forms from the Planchés Inédites of d'Orbigny. A study of the Philippine material, which is abundant, has shown several forms which seem to be distinct. A comparison of these with the originals of d'Orbigny shows several interesting points. The original figures of d'Orbigny²⁹ show a form nearly equally biconvex, with a blunt periphery and very numerous chambers, usually with a secondary series of clear shell tissue between the suture lines. D'Orbigny's model, however (No. 98), shows a more convex form with an acutely carinate periphery and much fewer chambers. This form is close to that figured by Fornasini (pl. 2, fig. 1). If from these two distinct forms which d'Orbigny had under this name the original of the figure in the 1826 work is taken as a type, the typical *A. lessonii* as considered here will be a rather flattened biconvex test with a bluntly rounded periphery, very numerous narrow oblique chambers, and with the umbo very small but prominent, usually clear shell material. The sutures are almost always darker than the rest of the test, and between them are usually shorter lines or points, also clear shell material, showing dark against the lighter colored background. The second form, which in Fornasini's paper is figured (pl. 2, fig.), represents a variety named by d'Orbigny *madagascariensis*. This has few chambers, an acutely carinate border, very oblique sutures, and no papillae near the umbilical area. These two forms are very distinct in the Philippine specimens. A third very large form has very numerous chambers, the sutures nearly straight, and radiating from the center to the periphery. This is the form distinguished by Chapman and Heron-Allen and Earland as variety *radiata* (Fichtel and Moll).

The genus is most common in shallow, warm water, especially among the islands of the Archipelago.

AMPHISTEGINA LESSONII d'Orbigny.

Amphistegina lessonii D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 304, No. 3, pl. 17, figs. 1-4.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 740, pl. 111, fig. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 35.

The typical form of the species, as has already been noted, is a biconvex test with very numerous oblique chambers, the periphery bluntly rounded, the sutures oblique, very sharply angled near the periphery; the sutures themselves on the surface usually all clear shell material, showing darker against the nearly white opaque background formed by the lateral walls of the chamber; also between the sutures there are shorter lines of clear shell material and very often

²⁸ Rend. Acad. Sci. Bologna, vol. 7, 1903.

²⁹ Ann. Sci. Nat., vol. 7, 1826, pl. 17, figs. 1-4.

small dots of similar color; wall usually smooth except on the ventral side near the aperture where it is finely papillate.

Distribution.—This species is very common especially in warm, shallow waters. It is very frequent in waters less than 30 fathoms (55 meters) in depth, and makes up a considerable proportion of the sand on some of the beaches. It is most abundant in the region of the Sulu Archipelago, and is usually associated with *Heterostegina*, *Operculina*, *Alveolina*, *Orbitolites*, and other genera characteristic of this same type of habitat. The deeper records, of which there are comparatively few, are usually represented by very few specimens. It has occurred in the following localities: China Sea, off southern Luzon; Sulu Archipelago; near Basilan Island: off Jolo Jolo; off Siasi; off Tawi Tawi; off Romblon; off eastern Panay; off northern Cebu; Sogod Bay, southern Leyte; between Burias and Luzon; Gulf of Davao; Verde Island Passage; China Sea, off Hongkong; Palawan Passage; Ragay Gulf, Luzon; between Cebu and Bohol; east coast Luzon; between Samar and Leyte; between Negros and Siquijor; Darvel Bay, Borneo; Buton Strait; and Macassar Strait.

Amphistegina lessonii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15425	U.S.N.M.	10+	D5106...	14 23 55 N.; 120 32 33 E....	37	gy. m.	Common.
15426	U.S.N.M.	3	D5109....	14 03 45 N.; 120 16 30 E....	10	Co.	Few.
15427	U.S.N.M.	10+	D5110....	13 59 20 N.; 120 75 45 E....	135	59.0	dk. gy. m....	Common.
15428	U.S.N.M.	10+	D5134....	6 44 45 N.; 121 48 00 E....	25	fine. s.	Common.
15429	U.S.N.M.	10+	D5138....	6 06 00 N.; 120 58 50 E....	19	s., Co.	Common.
15430	U.S.N.M.	10+	D5139....	6 06 00 N.; 121 02 30 E....	20	crs., s.	Common.
15431	U.S.N.M.	10+	D5142....	6 06 10 N.; 121 02 40 E....	21	crs. s., sh....	Common.
15432	U.S.N.M.	10+	D5144....	6 05 50 N.; 121 02 15 E....	19	crs. s., sh....	Common.
15433	U.S.N.M.	7	D5145....	6 04 30 N.; 120 59 30 E....	23	crs. s., sh....	Common.
15434	U.S.N.M.	10+	D5146....	5 46 40 N.; 120 48 50 E....	24	crs. s., sh....	Common.
15435	U.S.N.M.	10+	D5148....	5 35 40 N.; 120 47 30 E....	17	crs. s., sh....	Common.
15436	U.S.N.M.	10+	D5152....	6 22 55 N.; 120 15 45 E....	34	wh. s.	Common.
15437	U.S.N.M.	7	D5153....	5 18 10 N.; 120 02 55 E....	49	crs. s., sh....	Common.
15438	U.S.N.M.	3	D5156....	5 12 50 N.; 119 55 55 E....	18	fine. s., sh....	Few.
15439	U.S.N.M.	10+	D5159....	5 11 50 N.; 119 54 00 E....	10	crs. s., sh....	Common.
15440	U.S.N.M.	5	D5161....	5 10 15 N.; 119 53 00 E....	16	fine. s.	Common.
15441	U.S.N.M.	10	D5172....	6 03 15 N.; 120 35 30 E....	318	fine. s., sh....	Common.
15442	U.S.N.M.	10	D5178....	12 43 00 N.; 122 06 15 E....	78	fine. s.	Common.
15443	U.S.N.M.	10+	D5179....	12 38 15 N.; 122 12 30 E....	37	75.7	hrd. s.	Common.
15444	U.S.N.M.	10+	D5181....	11 36 40 N.; 123 26 35 E....	26	m., fine. s....	Common.
15445	U.S.N.M.	2	D5182....	11 34 40 N.; 123 23 20 E....	24	fine. s., m....	Few.
15446	U.S.N.M.	10+	D5192....	11 09 15 N.; 123 50 00 E....	32	gn. s.	Common.
15447	U.S.N.M.	6	D5201....	10 10 00 N.; 125 04 15 E....	554	52.8	gy. m., s....	Common.
15448	U.S.N.M.	6	D5218....	13 11 15 N.; 123 02 45 E....	20	crs. s.	Common.
15449	U.S.N.M.	1	D5219....	13 21 00 N.; 122 18 45 E....	530	50.8	gn. m.	Rare.
15450	U.S.N.M.	10+	D5250....	7 05 07 N.; 125 39 45 E....	23	co. s.	Common.
15451	U.S.N.M.	5	D5268....	13 42 00 N.; 120 57 15 E....	170	s., p.	Common.
15452	U.S.N.M.	6	D5276....	13 49 15 N.; 120 14 45 E....	18	sh., p., s....	Common.
15453	U.S.N.M.	10+	D5282....	13 53 00 N.; 120 26 45 E....	248	47.4	dk. gy. s....	Common.
15455	U.S.N.M.	2	D5296....	13 40 09 N.; 120 57 45 E....	210	m., s.	Few.
15456	U.S.N.M.	1	D5311....	21 33 00 N.; 116 15 00 E....	88	crs. s., sh....	Few.
15457	U.S.N.M.	1	D5349....	10 54 00 N.; 118 26 20 E....	730	40.6	co. s.	Few.
15458	U.S.N.M.	2	D5381....	13 14 15 N.; 122 44 45 E....	88	co. s.	Few.
15459	U.S.N.M.	1	D5412....	10 09 15 N.; 123 52 00 E....	162	54.8	gn. m.	Few.
15460	U.S.N.M.	1	D5465....	13 39 42 N.; 123 40 39 E....	500	gy. m.	Few.
15461	U.S.N.M.	3	D5469....	13 36 48 N.; 123 38 24 E....	500	gn. m.	Few.
15462	U.S.N.M.	4	D5481....	10 27 30 N.; 125 17 10 E....	61	s., sh., g....	Few.
15463	U.S.N.M.	7	D5538....	9 08 15 N.; 123 23 20 E....	256	53.3	gn. m., s....	Common.
15464	U.S.N.M.	1	D5599....	5 33 15 N.; 120 15 30 E....	303	52.3	co. s.	Rare.
15465	U.S.N.M.	2	D5570....	5 32 15 N.; 120 12 57 E....	330	52.3	fine. s., glob....	Few.
15466	U.S.N.M.	3	D5571....	5 30 45 N.; 120 07 57 E....	340	52.3	s., sh.	Few.
15467	U.S.N.M.	3	D5580....	4 52 45 N.; 119 06 45 E....	162	55.8	br. s., Co....	Few.
15468	U.S.N.M.	3	D5640....	4 27 00 S.; 122 55 40 E....	24	s., brk. sh....	Few.
15454	U.S.N.M.	1	D5668....	2 28 15 S.; 118 49 00 E....	901	38.2	gy. m.	Few.

AMPHISTEGINA LESSONII d'Orbigny, var. MADAGASCARIENSIS d'Orbigny.

Amphistegina madagascariensis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 305, No. 5.—FORNASINI, Rend. Acad. Sci. Bologna, vol. 7, 1903, p. 144, pl. 2, fig. 5.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 736.

This form, which is here referred to d'Orbigny's *Amphistegina madagascariensis*, seems to be very distinct in the Philippine material and may be a valid species. It differs from the typical *A. lessonii* in the smaller number of chambers and the acutely carinate periphery, the unequal biconvex form, and the larger umbonal clear space. The sutures are more evenly curved and are usually depressed. In the Philippine material there seem to be no intermediate forms between these two. This variety is found under similar conditions as the typical, most abundant in warm shallow waters, and occurs with the typical, although in less numbers.

Amphistegina lessonii, var. *madagascariensis*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15469	U.S.N.M.	1	D5106...	14 23 55 N.; 120 32 33 E.	37	gy. m.	Rare.
15470	U.S.N.M.	2	D5109...	14 03 45 N.; 120 16 30 E.	10	Co	Rare.
15471	U.S.N.M.	2	D5110...	13 59 20 N.; 120 75 45 E.	135	59.0	dk. gy. m.	Rare.
15472	U.S.N.M.	5	D5113...	13 51 30 N.; 120 50 30 E.	159	dk. gr. m.	Few.
15473	U.S.N.M.	3	D5134...	6 44 45 N.; 121 48 00 E.	25	fine. s.	Few.
15474	U.S.N.M.	1	D5139...	6 06 00 N.; 121 02 30 E.	20	ers. s.	Rare.
15475	U.S.N.M.	7	D5146...	5 46 40 N.; 120 48 50 E.	24	ers. s., sh.	Few.
15476	U.S.N.M.	7	D5148...	5 35 40 N.; 120 47 30 E.	17	ers. s., sh.	Few.
15477	U.S.N.M.	7	D5154...	5 14 50 N.; 119 58 45 E.	12	co. s.	Few.
15478	U.S.N.M.	8	D5159...	5 11 50 N.; 119 54 00 E.	10	ers. s.	Common.
15479	U.S.N.M.	10+	D5160...	5 12 40 N.; 119 55 10 E.	12	s.	Common.
15480	U.S.N.M.	10+	D5179...	12 38 15 N.; 122 12 30 E.	37	75.7	hrd. sh.	Common.
15481	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E.	20	ers. s.	Rare.
15482	U.S.N.M.	4	D5250...	7 05 07 N.; 125 39 45 E.	23	co. s.	Rare.
15483	U.S.N.M.	4	D5276...	13 49 15 N.; 120 14 45 E.	18	sh., p., s.	Rare.
15484	U.S.N.M.	4	D5430...	9 49 40 N.; 119 03 20 E.	464	50.0	glob. ooze.	Rare.
15485	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E.	500	gy. m.	Rare.
15486	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E.	500	gr. m.	Rare.
15487	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E.	Rare.
15488	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E.	256	53.3	gn. m., s.	Rare.
15489	U.S.N.M.	2	D5572...	5 31 26 N.; 120 09 45 E.	334	52.3	s.	Rare.
15490	U.S.N.M.	10+	D5640...	4 27 00 S.; 122 55 40 E.	24	s., brk. sh.	Common.

AMPHISTEGINA LESSONII d'Orbigny, var. RADIATA (Fichtel and Moll).

Nautilus radiatus FICHTEL and MOLL, Test. Micr., 1798, p. 58, pl. 8, figs. 8, a-d.

Amphistegina radiata CHAPMAN, Proc. Zool. Soc. London, 1895, pp. 45-47, pl. 1, figs. 8-10, 12.

Amphistegina lessonii, var. *radiata* HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 736.

This large, many-chambered, biconvex form, which is distinguished by Chapman and also by Heron-Allen and Earland, is common at some of the stations. It is characterized by the straight, radiating suture lines which appear as transparent lines in the material of the test.

It has occurred at the following localities: China Sea, off southern Luzon; off Jolo Jolo; off Tawi Tawi; off Romblon; off eastern Panay; between Burias and Luzon; Gulf of Davao; Verde Island Passage; and Darvel Bay, Borneo.

Amphistegina lessonii, var. *radiata*.—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15404	U.S.N.M.	2	D5110	13 59 20 N.; 120 75 45 E.	135	59.0	dk. gy. m.	Rare.
15405	U.S.N.M.	10+	D5141	6 09 00 N.; 120 58 00 E.	29		co. s.	Common.
15406	U.S.N.M.	10+	D5142	6 06 10 N.; 121 02 40 E.	21		crs. s., sh.	Common.
15407	U.S.N.M.	6	D5144	6 05 50 N.; 121 02 15 E.	19		crs. s.	Few.
15408	U.S.N.M.	9	D5145	6 04 30 N.; 120 59 30 E.	23		crs. s., sh.	Common.
15409	U.S.N.M.	5	D5147	5 41 40 N.; 120 47 10 E.	21		co. s., sh.	Common.
15410	U.S.N.M.	10+	D5149	5 33 00 N.; 120 42 10 E.	10		co. sh.	Common.
15411	U.S.N.M.	10+	D5151	5 25 40 N.; 120 27 15 E.	24		crs. s., sh.	Common.
15412	U.S.N.M.	10	D5154	5 14 50 N.; 119 58 45 E.	12		crs. s., sh.	Common.
15413	U.S.N.M.	9	D5172	6 03 15 N.; 120 35 30 E.	318		fine s., sh.	Common.
15414	U.S.N.M.	1	D5179	12 38 15 N.; 122 12 30 E.	37	75.7	hrd. s.	Rare.
15415	U.S.N.M.	5	D5181	11 36 40 N.; 123 36 25 E.	26		m., fine s.	Few.
15416	U.S.N.M.	9	D5184	10 18 30 N.; 122 23 30 E.	565	49.8	gn. m.	Common.
15417	U.S.N.M.	10+	D5218	13 11 15 N.; 123 02 45 E.	20		crs. s.	Common.
15418	U.S.N.M.	7	D5250	7 05 07 N.; 125 39 45 E.	23		co. s.	Common.
15419	U.S.N.M.	10+	D5255	7 03 00 N.; 125 39 00 E.	100		sft. m.	Common.
15420	U.S.N.M.	10+	D5268	13 42 00 N.; 120 57 15 E.	170		s., p.	Common.
15421	U.S.N.M.	2	D5276	13 49 15 N.; 120 14 45 E.	18		s., p., sh.	Rare.
15422	U.S.N.M.	2	D5569	5 33 15 N.; 120 15 30 E.	303	52.3	crs. s.	Rare.
15423	U.S.N.M.	3	D5579	4 54 15 N.; 119 09 52 E.	175	55.3	fine s., Co.	Few.
15424	U.S.N.M.	4	D5580	4 52 45 N.; 119 06 45 E.	162	55.8	br. s., Co.	Few.

Genus OPERCULINA d'Orbigny, 1826.

Nautilus (part) GRONOVIVS, Zooph. Gron., 1781, p. 282.

Lenticulites (part) DEFRANCE, Dict. Sci. Nat., vol. 25, 1822, p. 453.

Operculina D'ORBIGNY (Type, *O. complanata* DeFrance), Ann. Sci. Nat., vol. 7, 1826, p. 281.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 742.—CARPENTER, Introd. Foram., 1862, p. 247.

Carpenter has given a very detailed account of the structure of this genus, and mentions especially its relations to *Heterostegina*, the relations which are very clear in a series of Philippine specimens studied. Among his material Carpenter had specimens from the Philippine region, and his notes on these, after studying the *Albatross* Philippine collection, are very interesting. There is, undoubtedly, much variation, as Carpenter mentions, but this seems to be along definite lines, and from a study of the several thousand specimens of available Philippine material the various forms seem to fall into definite categories.

In looking up the literature the first species to be described is *O. complanata* DeFrance. This species, which is known from d'Orbigny's figure,³⁰ and from the Modèles, No. 80, is a definite many-chambered, very flat, unornamented test, the coils rapidly increasing

³⁰ Ann. Sci. Nat., vol. 7, 1826, pl. 14, figs. 7-10.

in size, and the chambers very narrow and much curved. The type specimen is a fossil from the region of Bordeaux.

Although many writers have referred many things to this species, a study of the Philippine collection does not show anything which may be reasonably referred to this species as shown by d'Orbigny. Therefore, as far as the Philippine species are concerned, the name is not available. A study of the other species described since d'Orbigny's time shows several names, none of which can well be applied here. As a rule the various forms of *Operculina* have been divided by various authors into smooth forms, which are referred to *Operculina complanata*, and into those showing granular ornamentation, which are usually assigned to *Operculina granulosa* Leymerie, either as a species or variety of *O. complanata*. Such a treatment is a very simple way to dispose of the entire matter. However, the study of the thousands of Philippine specimens has shown that there is no such general and simple rule which can apply to all the specimens obtained. Also a study of the type figure and description of *O. granulosa* by Leymerie, which was a fossil from Europe, shows that it is a very close-coiled form, apparently with rows of beads on the sutures. Such a form is possibly known from a few stations in the Philippines, and for this the name *O. granulosa* is restricted. Of the other forms two may take their names from the species named by d'Orbigny, and later figured in Fornasini's publication of the "planchès inédites."³¹ The first of these to which d'Orbigny gave the name *O. gaimairdi*, was from "Rawack, Nouvelle-Hollande." That would be in a closely related area to the Philippine faunal region, and a form very close to this occurs abundantly at a number of stations with definite geographical limitations. Another group is similar to that figured by d'Orbigny and named by him *Assilina discoidalis*. This has a large protuberant center, with numerous granules, followed by raised sutures, the whole with a definite keel and the revolving whorls thickest near the middle. Such forms are very abundant at a few stations in the Sulu Archipelago, but are very rare elsewhere. The type station for this species is Rawack.

Carpenter also mentions and figures a form from the Philippines very similar to this. For such forms the name *O. discoidalis* can be used.

In studying the large collection of *Operculina* each station was taken separately, and in this way it was possible to recognize a number of minor forms or races which seem to have a definite geographical distribution. The genus is best developed in comparatively shallow water but some of the large forms reach considerable depths.

³¹ Boll. Soc. Geol. Ital., vol. 22, 1903, pl. 14, figs. 4, 8.

The relations of the different species and varieties will be discussed in the pages that follow.

OPERCULINA GAIMAIRD *d'*Orbigny.

Operculina gaimairdi D'ORBIGNY, Ann Sci. Nat., vol. 7, 1826, p. 281, No. 5.—
FORNASINI, Boll. Soc. Geol. Ital., vol. 22, 1903, pl. 14, fig. 4.

Description.—Test of medium size, much compressed, complanate, opaque; chambers 15 or less in the last-formed coil, curved, sutures distinct, and the revolving suture fairly well shown; central area somewhat enlarged in megalospheric specimens, the remainder of the test marked by chains of beads along the sutures, the rest of the chamber being smooth and unornamented.

Diameter up to 3 mm., sometimes more.

This species has occurred in the Philippines in the following localities: Vicinity of Jolo; Sulu Archipelago; vicinity of Siasi; Tawi Tawi Group; between Burias and Luzon; and in the Gulf of Davao. Bottom temperatures are not given at any of the stations at which it is recorded; the depths at which it is found range from 10 to 318 fathoms (18 to 582 meters).

This seems to be a very definite species in the area and limited to warm regions, as the table of distribution shows.

The whole test is opaque, the ornamentation of the sutures showing as darker lines on account of the transparency of that portion. There is some variation in the width of the coil, and the granules of the ornamentation may blend into confluent lines.

Operculina gaimairdi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15943	U.S.N.M.	10+	D5138...	6 06 00 N.; 120 58 50 E...	19	s., Co.....	Frequent.
15944	U.S.N.M.	10+	D5139...	6 06 00 N.; 121 02 30 E...	20	co. s.....	Frequent.
15945	U.S.N.M.	10+	D5141...	6 09 00 N.; 120 58 00 E...	29	co. s.....	Frequent.
15946	U.S.N.M.	10+	D5142...	6 06 10 N.; 121 02 40 E...	21	co. s., sh.....	Frequent.
15947	U.S.N.M.	10+	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Frequent.
15948	U.S.N.M.	10+	D5144...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Frequent.
15949	U.S.N.M.	10+	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.....	Frequent.
15950	U.S.N.M.	10+	D5146...	5 46 40 N.; 120 48 50 E...	24	co. s., sh.....	Frequent.
15951	U.S.N.M.	3	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.....	Few.
15952	U.S.N.M.	10	D5148...	5 35 40 N.; 120 47 30 E...	17	co. s.....	Frequent.
15953	U.S.N.M.	2	D5149...	5 33 00 N.; 120 42 10 E...	10	co. sh.....	Few.
15954	U.S.N.M.	10+	D5151...	5 24 40 N.; 120 27 15 E...	24	co. s., sh.....	Frequent.
15955	U.S.N.M.	10+	D5154...	5 14 50 N.; 119 58 45 E...	12	co. s.....	Frequent.
15956	U.S.N.M.	10+	D5156...	5 12 50 N.; 119 55 55 E...	18	fine s.....	Frequent.
15957	U.S.N.M.	10+	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.....	Frequent.
15958	U.S.N.M.	5	D5160...	5 12 40 N.; 119 55 10 E...	12	s.....	Rare.
15959	U.S.N.M.	10+	D5161...	5 10 15 N.; 119 53 00 E...	16	fine s.....	Frequent.
15960	U.S.N.M.	5	D5172...	6 03 15 N.; 120 35 30 E...	318	fine s.....	Frequent.
15961	U.S.N.M.	8	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Frequent.
15962	U.S.N.M.	3	D5250...	7 05 07 N.; 125 39 45 E...	23	co. s.....	Few.

OPERCULINA BARTSCHI, new species.

Description.—Test comparatively large, compressed except in the central region, coils rapidly widening; central portion thickened, biconvex, composed of several coiled chambers, and ornamented on the surface with numerous comparatively large bosses; chambers numerous, 20 to 25 in the last-formed coil; curved sutures distinct, not ornamented, but the chambers between with numerous comparatively large granulations, usually in somewhat definite lines across the test.

Diameter up to 8 mm.

Distribution.—Type specimen (Cat. No. 15848, U.S.N.M.) from *Albatross* station D5134 Sulu Archipelago, near Basilan Island, 25 fathoms (46 meters).

This has occurred at several stations in the area, as follows: China Sea, off southern Luzon; Sulu Sea, off western Mindanao; vicinity of Jolo; vicinity of Siasi; Tawi Tawi Group; off Romblon; between Panay and Negros; off northern Cebu; Sogod Bay, southern Leyte; Pacific Ocean, east coast of Mindanao; Gulf of Davao; Linapacan Strait; Palawan Passage; between Cebu and Boho; eastern Palawan and vicinity; east coast of Luzon; between Leyte and Mindanao; northern Mindanao and vicinity; between Negros and Siquijor; north of Tawi Tawi; and Buton Strait.




FIG. 13.—OPERCULINA BARTSCHI, NEW SPECIES. SIDE VIEW, $\times 15$. FROM STATION D5134.

This is evidently one of the varieties described by Carpenter from the Philippines. It seems to be one of the most common species of the *Albatross* collection from the Philippine area, and has two definite varieties, which are here indicated. It is closely related to the granular forms of *Heterostegina* which were found in this material.

It is named in honor of Dr. Paul Bartsch, of the United States National Museum, whose careful collecting while in charge of the Philippine Expedition has been responsible for such abundant series of specimens.

There is a local race of this species found at two stations, D5640 and D5642, both from Buton Strait. These have a thickened periphery and larger size than the typical form. The granules of the exterior are smaller than the area farther north.

Operculina bartschi—Material examined.

Cat. No.	Coll. of -	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15845	U.S.N.M.	6	D5096...	14 20 23 N.; 120 34 15 E..	28	° F.	gy. m., s.....	Few.
15846	U.S.N.M.	5	D5097...	14 19 15 N.; 120 33 52 E..	30	gy. m., s.....	Few.
15847	U.S.N.M.	1	D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	gn. m., s.....	Rare.
15848	U.S.N.M.	5	D5134...	6 44 45 N.; 121 48' 00" E..	25	fine. s.....	Few.
15849	U.S.N.M.	10+	D5141...	6 09 00 N.; 120 58 00 E..	29	co. s.....	Frequent.
15850	U.S.N.M.	10+	D5142...	6 06 10 N.; 121 02 40 E..	21	co. s., sh.....	Frequent.
15851	U.S.N.M.	10+	D5143...	6 05 50 N.; 121 02 15 E..	19	co. s.....	Frequent.
15852	U.S.N.M.	10	D5144...	6 05 50 N.; 121 02 15 E..	19	co. s.....	Frequent.
15853	U.S.N.M.	10+	D5145...	6 04 30 N.; 120 59 30 E..	23	co. s., sh.....	Frequent.
15854	U.S.N.M.	2	D5146...	5 46 40 N.; 120 48 50 E..	24	co. s., sh.....	Few.
15855	U.S.N.M.	1	D5147...	5 41 40 N.; 120 47 10 E..	21	co. s., sh.....	Rare.
15856	U.S.N.M.	4	D5164...	5 01 40 N.; 119 52 50 E..	18	gn. m.....	Rare.
15857	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Rare.
15858	U.S.N.M.	4	D5184...	10 18 30 N.; 122 23 30 E..	565	49.8	gn. m.....	Few.
15859	U.S.N.M.	4	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.....	Few.
15860	U.S.N.M.	4	D5201...	10 10 00 N.; 125 04 15 E..	534	52.8	fine. gy. s.....	Few.
15861	U.S.N.M.	4	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.....	Few.
15862	U.S.N.M.	1	D5250...	7 05 07 N.; 125 39 45 E..	23	co., s.....	Few.
15863	U.S.N.M.	4	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s.....	Few.
15864	U.S.N.M.	2	D5281...	13 52 45 N.; 120 25 00 E..	201	50.4	dk. gy. s.....	Few.
15865	U.S.N.M.	10+	D5336...	11 37 45 N.; 119 46 00 E..	46	s., m.....	Frequent.
15866	U.S.N.M.	10+	D5338...	11 33 45 N.; 119 24 45 E..	43	Co., s., m.....	Frequent.
15867	U.S.N.M.	10	D5339...	11 22 00 N.; 119 12 00 E..	52	m.....	Frequent.
15868	U.S.N.M.	1	D5349...	10 54 00 N.; 118 26 20 E..	730	40.6	Co., s.....	Rare.
15869	U.S.N.M.	1	D5419...	9 58 30 N.; 123 46 00 E..	175	54.5	gn. m.....	Rare.
15870	U.S.N.M.	4	D5426...	9 12 00 N.; 118 28 00 E..	27	fine. gy. s.....	Few.
15871	U.S.N.M.	7	D5428...	9 13 00 N.; 118 51 15 E..	1,105	49.7	gy. m.....	Few.
15872	U.S.N.M.	2	D5429...	9 41 30 N.; 118 50 22 E..	766	gn. m.....	Few.
15873	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 21 E..	500	gn. m.....	Rare.
15874	U.S.N.M.	2	D5487...	10 02 45 N.; 125 05 33 E..	732	32.3	gn. m.....	Rare.
15875	U.S.N.M.	2	D5493...	9 04 00 N.; 125 20 00 E..	478	52.1	gn. m.....	Rare.
15876	U.S.N.M.	2	D5495...	9 06 30 N.; 125 00 20 E..	976	52.3	gy. m.....	Rare.
15877	U.S.N.M.	3	D5512...	8 16 02 N.; 123 58 26 E..	445	52.8	gy. m.....	Rare.
15878	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E..	Rare.
15879	U.S.N.M.	7	D5538...	9 08 15 N.; 123 23 20 E..	254	53.5	gn. m.....	Few.
15880	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E..	330	52.3	fine. s., glob.....	Rare.
15881	U.S.N.M.	10+	D5640...	4 27 00 S.; 122 55 40 E..	24	s., brk. sh.....	Frequent.
15882	U.S.N.M.	2	D5642...	4 31 49 S.; 122 49 42 E..	37	gy. m.....	Rare.

OPERCULINA BARTSCHI, new species, var. PLANA, new variety.

Description.—Test different from the typical form of the species in the ornamentation of the surface, which, after the early development, is usually smooth, very thin, and translucent.

This has occurred at the following stations: China Sea, off southern Luzon; vicinity of Jolo; off northern Cebu; Sogod Bay, southern Leyte; between Marinduque and Luzon; Pacific Ocean, east coast of Mindanao; southern Mindanao; China Sea, off Formosa; Palawan Passage; between Samar and Leyte; between Leyte and Mindanao; northern Mindanao and vicinity; and between Negros and Siquijor.

This usually occurs with the typical form of the species, but is not as common. There is a peculiar race which was noted only at D5257, southern Mindanao, in 28 fathoms (51 meters).




FIG. 14.—OPERCULINA BARTSCHI, VAR. PLANA, NEW VARIETY. $\times 30$. SIDE VIEW. FROM STATION D5105.

Operculina bartschi, var. *plana*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / "	° / "				
15883	U.S.N.M.	10+	D5097...	14 19 15 N.;	120 33 52 E..	30	° F.	gy. m., s....	Frequent.
15884	U.S.N.M.	2	D5105...	14 43 55 N.;	120 12 50 E..	25	Few.
15885	U.S.N.M.	1	D5106...	14 23 55 N.;	120 32 33 E..	37	gy. m.....	Rare.
15886	U.S.N.M.	4	D5142...	6 06 10 N.;	121 02 40 E..	21	co. s., sh....	Few.
15887	U.S.N.M.	10+	D5192...	11 09 15 N.;	123 50 00 E..	32	gn. s.....	Frequent.
15888	U.S.N.M.	2	D5201...	10 10 00 N.;	125 04 15 E..	554	52.8	gy. s., m....	Few.
15889	U.S.N.M.	1	D5229...	13 38 00 N.;	121 58 00 E..	50	sft. gn. m....	Rare.
15890	U.S.N.M.	5	D5236...	8 50 45 N.;	126 26 52 E..	494	41.2	fine. gy. s....	Few.
15891	U.S.N.M.	10+	D5257...	7 22 12 N.;	124 12 15 E..	28	m.....	Frequent.
15892	U.S.N.M.	1	D5315...	21 40 00 N.;	116 58 00 E..	148	s., sh.....	Rare.
15893	U.S.N.M.	5	D5338...	11 33 45 N.;	119 24 45 E..	43	Co., s., m....	Few.
15894	U.S.N.M.	4	D5339...	11 22 00 N.;	119 12 00 E..	57	m.....	Few.
15895	U.S.N.M.	9	D5478...	10 46 24 N.;	125 16 30 E..	52	sh.....	Common.
15896	U.S.N.M.	1	D5487...	10 02 45 N.;	125 05 33 E..	732	32.3	gn. m.....	Rare.
15897	U.S.N.M.	7	D5493...	9 04 00 N.;	125 20 00 E..	478	52.1	gn. m.....	Frequent.
15898	U.S.N.M.	3	D5495...	9 06 30 N.;	125 00 20 E..	976	52.3	gy. m.....	Few.
15899	U.S.N.M.	6	D5512...	8 16 02 N.;	123 58 26 E..	445	52.8	gy. m.....	Few.
15900	U.S.N.M.	2	D5537...	9 11 00 N.;	123 23 00 E..	254	53.5	gn. m.....	Few.
15901	U.S.N.M.	4	D5538...	9 08 15 N.;	123 23 20 E..	254	53.5	gn. m.....	Few.

OPERCULINA BARTSCHI, new species, var. ORNATA, new variety.

Plate 74, figs. 2a, b.

Description.—Variety differing from the typical in the ornamentation, especially in the later portion of the test, where the granules, instead of being confined to the surface of the chambers themselves, appear only above the sutures.

This variety occurs with the typical and with the preceding variety, and at D5257, southern Mindanao, is a peculiar local race; another occurs at D5640, Buton Strait.

Operculina bartschi, var. *ornata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / "	° / "				
15902	U.S.N.M.	10+	D5097...	14 19 15 N.;	120 33 52 E...	30	° F.	gy. m., s....	Frequent.
15903	U.S.N.M.	5	D5106...	14 23 55 N.;	120 32 33 E...	37	gy. m.....	Few.
15904	U.S.N.M.	1	D5113...	13 51 30 N.;	120 50 30 E...	159	dk. gn. m....	Rare.
15905	U.S.N.M.	1	D5134...	6 44 45 N.;	121 48 00 E...	25	fine. s.....	Rare.
15906	U.S.N.M.	4	D5142...	6 06 10 N.;	121 02 40 E...	21	co. s., sh....	Few.
15907	U.S.N.M.	7	D5143...	6 05 50 N.;	121 02 15 E...	19	co. s., sh....	Few.
15908	U.S.N.M.	3	D5144...	6 05 50 N.;	121 02 15 E...	19	co. s., sh....	Few.
15909	U.S.N.M.	10+	D5145...	6 04 30 N.;	120 59 30 E...	23	co. s., sh....	Common.
15910	U.S.N.M.	9	D5192...	11 09 15 N.;	123 50 00 E...	32	gn. s.....	Common.
15911	U.S.N.M.	2	D5236...	8 50 45 N.;	126 26 52 E...	494	41.2	fine. gy. s....	Rare.
15912	U.S.N.M.	1	D5247...	7 02 00 N.;	125 38 45 E...	135	m.....	Rare.
15913	U.S.N.M.	10+	D5257...	7 22 12 N.;	124 12 15 E...	28	m.....	Frequent.
15914	U.S.N.M.	2	D5338...	11 33 45 N.;	119 24 45 E...	143	co. s., m....	Rare.
15915	U.S.N.M.	4	D5339...	11 22 00 N.;	119 12 00 E...	52	m.....	Few.
15916	U.S.N.M.	1	D5538...	9 08 15 N.;	123 23 20 E...	254	53.5	gn. m.....	Few.
15917	U.S.N.M.	6	D5640...	4 27 00 S.;	122 55 40 E...	24	s., brk. sh...	Few.

OPERCULINA PHILIPPINENSIS, new species.

Description.—Test compressed, not flaring, close-coiled, complanate; central area thickened by the proloculum only; thin

throughout, opaque; chambers 15 to 20 in the last-formed coil; surface of the chambers smooth; sutures ornamented by a row of raised tubercles, the sutures typical, nearly straight, then near the periphery bend back sharply to the outer edge.

Diameter 3-5 mm.

Distribution.—Type specimen (Cat. No. 15923, U.S.N.M.) from *Albatross* station D5134, Sulu Archipelago, near Basilan Island, 25 fathoms (46 meters).

This is a common species in the Philippine area, and while some of its specimens show relations with other species it is usually distinct.

At *Albatross* station D5428 there is a local race, differing from the typical in the larger size, and in the very prominent keel which in the adult is thicker than that part of the test which is adjacent to it; the rows of beads in the adult specimens usually become fused.




FIG. 15.—OPERCULINA PHILIPPINENSIS, NEW SPECIES. SIDE VIEW. $\times 15$. FROM STATION D5134.

Operculina philippinensis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15918	U.S.N.M.	6	D5097...	14 19 15 N.; 120 33 52 E...	28	° F.	gy. m., s....	Few.
15919	U.S.N.M.	6	D5106...	14 23 55 N.; 120 32 33 E...	37		gy. m.	Few.
15920	U.S.N.M.	2	D5110...	13 59 20 N.; 120 75 45 E...	135	59.0	dk. gy. m....	Few.
15921	U.S.N.M.	10	D5132...	Island off Panabutan Point, N.; 15° W., 0.3 mile.	26		gn. m., s....	Frequent.
15922?	U.S.N.M.	10+	D5133...	Island off Panabutan Point N.; 52° E., 1.50 miles.	38		gn. m., s....	Frequent.
15923	U.S.N.M.	10+	D5134...	6 44 45 N.; 121 48 00 E...	25		fine. s....	Frequent.
15924	U.S.N.M.	2	D5145...	6 04 30 N.; 120 59 30 E...	23		co. s., sh....	Few.
15925	U.S.N.M.	10+	D5164...	5 01 40 N.; 119 52 20 E...	18		gn. m.	Frequent.
15926	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E...	78		fine. s....	Frequent.
15927	U.S.N.M.	6	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s....	Frequent.
15928	U.S.N.M.	10+	D5184...	10 18 30 N.; 122 23 30 E...	565	49.8	gn. m.	Frequent.
15929	U.S.N.M.	3	D5191...	10 29 45 N.; 123 31 15 E...	258	62.8	gn. m.	Few.
15930	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32		gn. s....	Frequent.
15931	U.S.N.M.	10+	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	fine. gy. s....	Frequent.
15932	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Few.
15933	U.S.N.M.	2	D5272...	14 00 00 N.; 120 22 30 E...	118	57.4	m., co. s....	Rare.
15934	U.S.N.M.	10+	D5276...	13 49 15 N.; 120 14 45 E...	18		sh., p., s....	Frequent.
15935	U.S.N.M.	5	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine. s....	Few.
15936	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s....	Rare.
15937	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E...	150	53.6	s....	Rare.
15938	U.S.N.M.	10+	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s....	Frequent.
15939	U.S.N.M.	10+	D5358...	6 06 40 N.; 118 18 15 E...			co. s....	Frequent.
15940	U.S.N.M.	7	D5426...	9 12 00 N.; 118 28 00 E...	27		fine. gy. s....	Frequent.
15941	U.S.N.M.	4	D5428...	9 13 00 N.; 118 51 15 E...	1,105	49.7	gy. m.	Few.
15942	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3		Rare.

OPERCULINA DISCOIDALIS (d'Orbigny).

Assilina discoidalis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 296, No. 1; Modèles, 1826, No. 88.—FORNASINI, Boll. Soc. Geol. Ital., vol. 22, 1903, pl. 14, fig. 8.

Description.—Test close-coiled throughout; central portion with a definite large proloculum, and the region next to it with numerous large bosses; three or four coils, with numerous chambers, the center part of each coil being thicker than the edges, leaving a deep-channelled spiral where the coils come together; the periphery with a

definite smooth edge, keeled, and finely striate; spiral lines of the early portion ornamented by rows of beadlike protuberances, which in later growth may become confluent.

It has occurred at 10 stations in the Philippine area, as follows: Sulu Sea, off western Mindanao; Sulu Archipelago, near Basilan Island; vicinity of Jolo, Sulu Archipelago; Tawi Tawi Group; off northern Cebu; off eastern Mindoro; and Jolo Sea.

This species was described by d'Orbigny from "la mer du Sud à Rawack." It is very abundant at three adjacent stations—D5132, D5133, D5134, Sulu Sea, off western Mindanao. It is evidently one of the species that Carpenter had from the Philippines, and can be recognized in his description. There is considerable variation in the thickness of the test and the following variety may be distinguished.

Operculina discoidalis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15963	U.S.N.M.	10+	D5132...	° " " ° " "	26	° F.	gn. m., s.....	Frequent.
15964	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N.; 15 W., 0.3 mile.	38		gn. m., s.....	Frequent.
15965	U.S.N.M.	10+	D5134...	6 44 45 N.; 121 48 00 E.	25		fne. s.....	Frequent.
15966	U.S.N.M.	8	D5136...	6 04 20 N.; 120 59 20 E.	22		s., sh.....	Frequent.
15967	U.S.N.M.	1	D5142...	6 06 10 N.; 121 02 40 E.	21		co. s., sh.....	Rare.
15968	U.S.N.M.	4	D5151...	5 24 40 N.; 120 27 15 E.	24		co. s., sh.....	Few.
15969	U.S.N.M.	5	D5156...	5 12 50 N.; 119 55 55 E.	18		fne. s.....	Few.
15970	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E.	32		gn. s.....	Few.
15971	U.S.N.M.	4	D5268...	13 42 00 N.; 120 57 15 E.	170		s., p.....	Few.
15972	U.S.N.M.	2	D5358...	6 06 40 N.; 118 18 15 E.			co., s.....	Few.

OPERCULINA DISCOIDALIS (d'Orbigny), var. INVOLUTA, new variety.

Description.—Differing from the typical in that the test, instead of being complanate, is more or less involute. The test is usually more smooth than the type form.




FIG. 16.—OPERCULINA DISCOIDALIS, VAR. INVOLUTA, NEW VARIETY. SIDE VIEW, $\times 20$. FROM STATION D5172.

This occurs at 12 stations in the area: Balayan Bay and Verde Island Passage; Sulu Sea, off western Mindanao; vicinity of Jolo; Sulu Archipelago; Tawi Tawi Group; Sogod Bay; southern Leyte; Gulf of Davao; China Sea, off southern Luzon; and Sibuko Bay, Borneo.

There are a few intermediate forms which seem to show that this involute variety should be placed with *O. discoidalis* rather than with any of the other species of the region. Some of the specimens are involute from the beginning; others become so after an early, more open development.

*Operculina discoidal*s, var. *involuta*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15973	U.S.N.M.	10+	D5113...	13 51 30 N.; 120 50 30 E.	159	° F.	dk. gn. m.	Frequent.
15974	U.S.N.M.	3	D5114...	13 36 11 N.; 120 42 28 E.	340	fine. s.	Few.
15975	U.S.N.M.	3	D5133...	Island off Panabutan Point, N.; 52 E., 1.5 miles.	38	gn. m., s.	Few.
15976	U.S.N.M.	10+	D5136...	6 04 20 N.; 120 59 29 E.	22	s., sh.	Frequent.
15977	U.S.N.M.	2	D5144...	6 05 50 N.; 121 02 15 E.	19	co. s., sh.	Rare.
15978	U.S.N.M.	10+	D5156...	5 12 50 N.; 119 55 55 E.	18	fine. s.	Frequent.
15979	U.S.N.M.	5	D5159...	5 11 50 N.; 119 54 00 E.	10	co. s.	Few.
15980	U.S.N.M.	1	D5172...	6 03 15 N.; 119 53 00 E.	16	fine. s.	Rare.
15981	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E.	554	52.8	gy. s., m.	Rare.
15982	U.S.N.M.	1	D5255...	7 03 00 N.; 125 39 00 E.	100	sft. m.	Rare.
15983	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E.	310	44.3	gn. m., s.	Rare.
15984	U.S.N.M.	1	D5121...	13 27 29 N.; 121 17 45 E.	108	dk. gn. m.	Rare.

OPERCULINA GRANULOSA Leymerie.

Operculina granulosa LEYMERIE, Mém. Soc. géol. France, ser. 2, vol. 1, 1846, p. 359, pl. 13, figs. 12, a, b.

To this species described by Leymerie from the Eocene of France are referred specimens from Tacloban Anchorage, where they occurred in great numbers, and from D5276, China Sea, vicinity of southern Luzon, 18 fathoms (33 meters). This is a close-coiled form, with a fairly large proloculum, very numerous chambers, larger size, with a definite keel; the others with the sutures ornamented by lines of tubercles, in the last development becoming solid lines; the test composed of usually about two and one-half coils.

These are referred with a question to Leymerie's species and are the only ones in the area which seem like that figured by the author, and it is doubtful if they really are the same.

Operculina granulosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15986	U.S.N.M.	10+	D5276...	13 49 15 N.; 120 14 45 E.	18	° F.	sh., p., s.	Frequent.
15985	U.S.N.M.	10+		Tacloban Anchorage, P. I.				Frequent.

OPERCULINA ELEGANS, new species.

Plate 97, fig. 3.

Description.—Test biconvex, smooth, and polished, translucent, usually of a brown color; somewhat involute; sutures of clear shell material, but not ornamented; periphery bluntly rounded.

Distribution.—Type specimen (Cat. No. 15988, U.S.N.M.) from *Albatross* station D5579, off Darvel Bay, Borneo.

This seems to be much different from any of the other species described, and probably represents a localized species in this particular region.

Operculina elegans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15988	U.S.N.M.	4	D5579...	4 54 18 N.; 119 09 52 E..	175	° F. 55.3	fine s., Co....	Few.
15987	U.S.N.M.	2	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co....	Few.

OPERCULINA AMMONOIDES (Gronovius).

Nautilus ammonoides GRONOVIVS, Zooph. Gron., 1781, p. 282, pl. 19, figs. 5, 6.

Operculina ammonoides PARKER and JONES, Introd. Foram., 1862, Appendix, p. 310.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 745, pl. 112, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 37, pl. 14, fig. 7.—HERON-ALLEN and EARLAND, Trans. Zool. Soc., London, vol. 20, 1915, p. 737.

This species is recorded by Brady in the Philippines in 95 fathoms (174 meters), and by Heron-Allen and Earland as "not uncommon in some dredgings we have from off Cebu (Philippine Islands), but the specimens are all very small compared with our British records."

All but one of the stations at which the specimens were found in the Philippine collection are from more than 100 fathoms (183 meters), and, except for this station, they range from 138 to 805 fathoms (252 to 1,472 meters); bottom temperatures are not low, and, except for two at 42.3° F. and 47.5° F. (5.7° C. and 8.6° C.), they range from 50.5° F. to 58.3° F. (10.2° C. to 14.6° C.).

The usual habitat for this species is comparatively deep and cold water, yet at several stations numerous specimens were found. They are all very typical and are not of large size, as has already been noted by Heron-Allen and Earland in their material from off Cebu.

Operculina ammonoides—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15834	U.S.N.M.	8	D5112...	13 48 22 N.; 120 47 25 E..	177	° F. 52.4	dk. gn. m....	Frequent.
15835	U.S.N.M.	1	D5114...	13 36 11 N.; 120 45 26 E..	340	fine s.	Rare.
15836	U.S.N.M.	5	D5115...	13 37 11 N.; 120 43 40 E..	340	Few.
15837	U.S.N.M.	1	D5160...	5 12 40 N.; 119 55 10 E..	12	s.	Rare.
15838	U.S.N.M.	3	D5215...	12 31 30 N.; 123 35 24 E..	604	50.5	gn. m.	Few.
15839	U.S.N.M.	10	D5216...	12 52 00 N.; 123 23 30 E..	215	51.9	gn. m.	Common.
15840	U.S.N.M.	3	D5222...	13 38 30 N.; 121 42 15 E..	195	52.8	gn. m.	Few.
15841	U.S.N.M.	1	D5284...	13 42 05 N.; 120 30 45 E..	422	42.3	gy. m., glob.	Rare.
15842	U.S.N.M.	1	D5526...	9 12 45 N.; 123 45 30 E..	805	52.3	gn. m., glob.	Rare.
15843	U.S.N.M.	1	D5546...	6 06 48 N.; 121 20 32 E..	138	58.3	fine co. s....	Rare.
15844	U.S.N.M.	2	D5666...	2 54 30 S.; 118 47 00 E..	272	47.5	gn. m.	Rare.

OPERCULINA VENOSA (Fichtel and Moll).

Nautilus venosus FICHEL and MOLL, Test. Micr., 1798, p. 59, pl. 8, figs. e-h.

Nummulites venosa CHAPMAN, Proc. Zool. Soc. London, 1895, p. 47.

Operculina venosa CHAPMAN, Proc. Roy. Soc. Victoria, vol. 26, 1913, p. 173.

Amphistegina cumingii CARPENTER, Philos. Trans., 1859, p. 32, pl. 5, figs. 13-17.

Nummulites cumingii H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 749, pl. 112, figs. 11-13; woodcut, fig. 22.—BAGG, Proc. U. S. Nat. Mus., vol.

34, 1908, p. 166.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 4, 1914, p. 39, pl. 14, fig. 6.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915,

p. 739.

There is a well-marked development of this species in certain parts of the areas, especially in shallow warm waters among the islands. At some stations it is very abundant and shows well the development of the species. Brady's figures in the *Challenger* report illustrate the early stages of this species, but the Philippine specimens show later characters: These consist of a broadening of the test and the development of a thin flangelike portion at the periphery. The edge of the test is marked by the development of a round carina, which is finely longitudinally striate, similar to what is found in other species of *Operculina*. Many of the specimens have the sutural lines broader than those figured by Brady, and of a very clear shell material, through which the supplementary canals can be seen as white lines.

It has occurred at the following localities: China Sea, off southern Luzon; Sulu Archipelago, near Basilan Island; vicinity of Jolo; vicinity of Siasi; Tawi Tawi Group; off Romblon; off northern Cebu; Sogod Bay, southern Leyte; and between Burias and Luzon.

Brady records this species in 95 fathoms (174 meters) from the Philippine Islands from dredgings of the *Challenger*. It seems to be limited to the Indo-Pacific region.

There are certain large specimens which seem to be microspheric, in which the flangelike portion is greatly developed, which grow to a size nearly 15 mm. in diameter.

Chapman (1913, above) mentions that "in my own cabinet there is a series of tests of this form from the East Indian Archipelago, which shows extreme modifications, ending with the typical *Operculina*." Also, according to Chapman, "*Operculina venosa* is found on parts of the Australian coast at the present day, in the neighborhood of the Great Barrier Reef."

If this species, as it certainly seems, is an *Operculina* instead of a species of *Nummulites*, it means that there are to-day no living species of the latter genus which suddenly became so abundant in the Eocene. As other genera of that period represented by complex tests, such as *Orthophragmina* and *Lepidocyclina*, have become extinct, it is not strange that *Nummulites* may also have died out.

Operculina venosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15642	U.S.N.M.	10	D5106...	14 23 55 N.; 120 32 33 E...	37	gy. m.	Common.
15643	U.S.N.M.	10	D5131...	6 44 45 N.; 121 48 00 E...	25	fine. s.	Common.
15679	U.S.N.M.	10+	D5138...	6 06 00 N.; 120 58 50 E...	19	s., co.	Common.
15644	U.S.N.M.	1	D5139...	6 06 00 N.; 121 02 30 E...	20	co. s.	Rare.
15645	U.S.N.M.	10+	D5141...	6 09 00 N.; 120 58 00 E...	29	co. s.	Common.
15646	U.S.N.M.	10+	D5142...	6 06 10 N.; 121 02 40 E...	21	co. s.	Common.
15647	U.S.N.M.	10+	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.	Common.
15648	U.S.N.M.	7	D5144...	6 05 50 N.; 121 02 15 E...	19	co. s.	Common.
15649	U.S.N.M.	10+	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.	Common.
15650	U.S.N.M.	3	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.	Few.
15651	U.S.N.M.	1	D5148...	5 35 40 N.; 120 47 30 E...	21	co. s., sh.	Few.
15652	U.S.N.M.	5	D5149...	5 33 00 N.; 120 42 10 E...	10	co. s., sh.	Few.
15653	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.	Few.
15654	U.S.N.M.	9	D5154...	5 14 50 N.; 119 58 45 E...	12	co. s.	Common.
15655	U.S.N.M.	7	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.	Common.
15656	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.	Rare.
15657	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.	Rare.
15658	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Rare.
15659	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.	Rare.
15660	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.	Rare.

Genus *HETEROSTEGINA* d'Orbigny, 1826.

This genus, which is comparatively abundant in certain parts of the area, especially in the region of the Sulu Archipelago, is not represented by the usual species of the genus, but by two other forms, one of which is very abundant and grows to a large size.

HETEROSTEGINA *DEPRESSA* d'Orbigny, var. *TUBERCALATA* Moebius.

Heterostegina tubercalata MOEBIUS, Beitr. Meeresfauna Insel Mauritius, 1880, p. 107, pl. 12, figs. 3-7.

This variety differs from the typical form of the species in that the earlier portion has numerous fine tubercles scattered over the surface; usually also the central portion is slightly thickened, the remaining part of the test being flat and complanate. None of the specimens show the typical smooth form which is characteristic of the typical form of the species. The types of d'Orbigny's *H. depressa* were from the Island of St. Helena.

This variety was especially abundant at *Albatross* station D5152, in 34 fathoms (62 meters), off Tawi Tawi group, Sulu Archipelago. It has also occurred in the following localities: Sogod Bay, southern Leyte; China Sea, off southern Luzon; and between Negros and Siquijor. It is only abundant in shallow water, the deeper stations being represented usually by single specimens. Moebius described this form from the shallow water of Mauritius.

Heterostegina depressa, var. *tubercalata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15491	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E...	34	° F.	wh. s.	Common.
15492	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.	Rare.
15493	U.S.N.M.	1	D5276...	13 49 15 N.; 120 14 45 E...	18	47.4	sh., p., s.	Rare.
15494	U.S.N.M.	1	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s.	Rare.
15495	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.	Rare.

HETEROSTEGINA SUBORBICULARIS d'Orbigny.

Heterostegina suborbicularis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 305.—
FORNASINI, Boll. Soc. Geol. Ital., vol. 22, 1903, pl. 14, fig. 6.

This species, not described by d'Orbigny, but later figured in his *planchés inédites* as published by Fornasini, seems to be the commonest in this region. Its whole appearance is very different from that of *H. depressa*, it being much thicker, strongly biconvex in the central portion, then later developing a compressed, broadly flaring portion which is thin near the periphery. The entire test is smooth and the subdivisions do not appear until a considerable amount of the test is built. These then appear near the periphery and gradually develop across the chambers of the test. The sutures in the early portion of the test are nearly straight, and remind one of those in *Operculina venosa*, which the early part of the test strongly resembles. The peripheral edge of the test is acute, except in very large specimens, where the border becomes thickened and somewhat carinate. Specimens attain a growth of 8 mm. or more. The species is abundant in warm, shallow waters, where it often makes up a large proportion of the bottom.

D'Orbigny described this species from the Hawaiian Islands as variety A, and variety B from the Marianna Islands, and from Port Jackson, Australia; therefore it is evidently a species of Indo-Pacific regions. Alcoholic specimens which were collected at D5179, off Romblon, show a green color, probably due to large numbers of symbiotic algae which occur in the test. Heron-Allen and Earland also note this character in *H. depressa* from the Kerimba Archipelago.

This species has occurred in the following localities: China Sea, off southern Luzon; Sulu Archipelago, near Basilan Island; vicinity of Jolo; vicinity of Siasi; near Tawi Tawi; off Romblon; between Burias and Luzon; Gulf of Davao; and Buton Strait.

Heterostegina suborbicularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
15496	U.S.N.M.	10+	D5106...	14 23 55 N.	120 32 33 E.	37	gy. m.....	Common.
15497	U.S.N.M.	1	D5109...	14 03 45 N.	120 16 30 E.	10	co.....	Rare.
15498	U.S.N.M.	1	D5134...	6 44 45 N.	121 48 00 E.	25	fine s.....	Rare.
15499	U.S.N.M.	8	D5138...	6 06 00 N.	121 02 30 E.	20	co. s.....	Few.
15500	U.S.N.M.	7	D5139...	6 06 00 N.	121 02 30 E.	20	co. s.....	Few.
15501	U.S.N.M.	7	D5141...	6 09 00 N.	120 58 00 E.	29	co. s.....	Few.
15502	U.S.N.M.	10+	D5142...	6 06 10 N.	121 02 40 E.	21	co. s.....	Common.
15503	U.S.N.M.	2	D5143...	6 05 50 N.	121 02 15 E.	19	co. s.....	Rare.
15504	U.S.N.M.	2	D5144...	6 05 50 N.	121 02 15 E.	19	co. s.....	Rare.
15505	U.S.N.M.	2	D5145...	6 04 30 N.	120 59 30 E.	23	co. s., sh.....	Rare.
15506	U.S.N.M.	2	D5147...	5 41 40 N.	120 47 10 E.	21	co. s., sh.....	Rare.
15507	U.S.N.M.	3	D5148...	5 35 40 N.	120 47 30 E.	17	co. s.....	Few.
15508	U.S.N.M.	5	D5149...	5 33 00 N.	120 42 10 E.	10	co., sh.....	Few.
15509	U.S.N.M.	4	D5151...	5 24 40 N.	120 27 10 E.	24	co. s., sh.....	Few.
15510	U.S.N.M.	1	D5152...	5 22 55 N.	120 15 45 E.	34	wh. s.....	Few.
15511	U.S.N.M.	4	D5154...	5 14 50 N.	119 58 45 E.	49	co. s., sh.....	Few.
15512	U.S.N.M.	10+	D5179...	12 38 15 N.	122 12 30 E.	37	75.7	hrd. s.....	Frequent.
15513	U.S.N.M.	1	D5184...	10 18 30 N.	122 23 30 E.	565	49.8	gn. m.....	Rare.
15514	U.S.N.M.	4	D5198...	9 40 50 N.	123 39 45 E.	220	53.9	gn. m.....	Few.
15515	U.S.N.M.	10+	D5218...	13 11 15 N.	123 02 45 E.	20	crs. s.....	Frequent.
15516	U.S.N.M.	3	D5250...	7 05 07 N.	125 39 45 E.	23	co. s.....	Few.
15517	U.S.N.M.	5	D5255...	7 03 00 N.	125 39 00 E.	100	sft. m.....	Rare.
15518	U.S.N.M.	1	D5276...	13 49 15 N.	120 14 45 E.	18	sh. p., s.....	Rare.
15519	U.S.N.M.	3	D5296...	13 40 09 N.	120 57 45 E.	210	m., s.....	Rare.
15520	U.S.N.M.	1	D5529...	9 23 45 N.	123 39 30 E.	441	53	gy. m., glob.	Rare.
15521	U.S.N.M.	1	D5572...	5 31 26 N.	120 09 45 E.	334	52.3	s.....	Rare.
15522	U.S.N.M.	4	D5579...	4 54 15 N.	119 09 52 E.	175	55.3	fine s., Co.	Rare.
15523	U.S.N.M.	5	D5580...	4 52 45 N.	119 06 45 E.	162	55.8	br. s., Co.	Few.
15524	U.S.N.M.	1	D5640...	4 27 60 S.	122 55 40 E.	24	s., brk. sh.....	Rare.

Genus CYCLOCLYPEUS Carpenter, 1856.

CYCLOCLYPEUS GUEMBELIANUS H. B. Brady.

Cycloclypeus CARPENTER, Philos. Trans., vol. 146, 1856, p. 555, pl. 30, figs. 1 and 3; Introd. Foram., 1862, p. 292.

Cycloclypeus guembelianus H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 66; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 751, pl. 111, figs. 8a, b (young form).

Cycloclypeus carpenteri H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 67; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 751.—LISTER, Philos. Trans., vol. 186B, 1895, pp. 437, 438, pl. 9, figs. 52-54.—CHAPMAN, Journ. Linn. Soc., vol. 28, 1900, p. 22, pl. 2, figs. 6, 7; pl. 3, figs. 1-5.

Brady gave this species a name in the *Challenger* Report, but gives almost no description. He mentions, however, that "some of these specimens are almost unique amongst discoidal Foraminifera in the point of size, presenting a diameter of 2½ inches (63 mm.)." Carpenter's genus was founded on this large species, which comes from "a considerable depth of water off the coast of Borneo." The *Challenger* material gave little to add to the knowledge of this species. However, it has been found occasionally since that time, as the above references show. The *Albatross* dredged this species in great quantity at D5179, off Romblon, in 37 fathoms (68 meters), and some of these specimens were preserved in a large-mouthed fruit jar, in which was a note saying that other specimens in the same dredging were too

large to be placed in the jar. The largest specimens fully equaled those described by Brady, and it is therefore the largest of the living foraminifera and only equaled in size by a few fossil species of *Lepidocyclina*. The test is very thin and fragile and easily becomes broken in the dredging.

It has occurred in large quantities at D5184, between Panay and Negros, 565 fathoms (1,033 meters); D5356, North Balabac Strait, 58 fathoms (106 meters); and D5640, Buton Strait, 24 fathoms (44 meters). It is evidently not a widely distributed species, and probably represents one of those older types which are now only found living in this particular region.

At station D5179, where the species was so abundant, there are specimens which have a secondary portion, due to a secondary outgrowth from near the center, so the specimens become composed of three almost equal portions, one of these, however, usually at a greater angle, so that the original plane of the surface can be seen.

Cycloclypeus guembelianus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15680	U.S.N.M.	10+	D5179...	12 38 15 N.; 122 12 30 E..	78	75.7	fine s.....	Abundant.
15765	U.S.N.M.	1	D5184...	10 18 30 N.; 122 23 30 E..	565	49.8	gn. m.....	Common.
15681	U.S.N.M.	2	D5356...	8 06 40 N.; 117 18 45 E..	58	s., sh.....	Common.
15682	U.S.N.M.	10+	D5640...	4 27 00 S.; 122 55 40 E..	24	s., brk. s....	Abundant.

Family MILIOLIDAE.

Subfamily CORNUSPIRININAE.

Genus CORNUSPIRA Schultze, 1854.

CORNUSPIRA FOLIACEA (Philippi).

Plate 77, fig. 1.

Orbis foliaceus PHILIPPI, Enum. Moll. Siciliae, vol. 2, 1844, p. 147, pl. 24, fig. 26.
Cornuspira foliacea CARPENTER, PARKER, and JONES, Introd. Foram., 1862, p. 68, pl. 5, fig. 16.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 199, pl. 11, figs. 5-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 24 pl. 1, fig. 1; pl. 2, fig. 1.

Specimens have occurred well scattered over the Archipelago proper, east and west coasts of Luzon, northern and eastern Mindanao, off Cebu and in Jolo Sea, depths ranging from 32 to 500 fathoms (59 to 914 meters), and bottom temperatures from 41.2° to 52.3° F. (5.1° to 11.2° C.). Outside the Archipelago specimens were obtained from D5613 in 752 fathoms (1,375 meters), Gulf of Tomini, Celebes; and D5668, in 901 fathoms (1,648 meters), Macassar Strait, bottom temperature 38.2° F. (3.4° C.).

The only station at which specimens were at all common is D5236 in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao.

Taken as a whole the specimens are very typical in size and general characters, at least for warm, comparatively shallow water. None of the specimens has a greater diameter than 3 mm. The species reaches its greatest development both in size and the amount of its spread in colder waters. The finest specimens I have ever seen were from the Arctic, while those of the colder parts of the North Atlantic reach large dimensions.

In the general faunal area it is known from the Hawaiian Islands (Bagg, Cushman) in the east to the Kerimba Archipelago, off south eastern Africa (Heron-Allen and Earland), and Mauritius (Egger) from southern Japan (Cushman) in the north to Raine Island, and Kandavu (Brady). In nearly all cases it is noted that the specimens from this region are of comparatively small size and rather close coiled. I noted this in the Philippine material, and it is confirmed by the following notes: ³²

Although tolerably abundant and widely diffused, the specimens are feeble, none of them exhibiting the rapid increase in size of the outer convolution which is characteristic of the species.

Heron-Allen and Earland: ³³

All the specimens are small and of the true Philippi type; none of the broad divergent type * * * were found at Kerimba.

It may be that Costa's name *ammonitiformis* should be applied to this smaller and more closely coiled form, and *foliacea* Philippi applied to the larger, more flaring form from colder and deeper water elsewhere.

Cornuspira foliacea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		°F.		
10718	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.	Few.
10719	U.S.N.M.	3	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.	Common.
10726	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E..	752	gy. m.	Rare.
10727	U.S.N.M.	1	D5668...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.	Rare.
10720	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	co. s.	Few.
10721	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.	Rare.
10722	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E..	500	gy. m.	Few.
10723	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E..	Rare.
10724	U.S.N.M.	1	D5549...	6 01 15 N.; 120 44 20 E..	263	52.3	s., glob., for.	Rare.
10725	U.S.N.M.	1	D5575...	5 28 30 N.; 120 02 27 E..	315	52.3	Co., s.	Rare.

CORNUSPIRA FOLIACEA (Philippi), var. EXPANSA Chapman.

Plate 77, fig. 2.

Cornuspira carinata COSTA, var. *expansa* CHAPMAN, Biol. Res. *Endeavour*, vol. 3, pt. 1, 1915, p. 12, pl. 1, fig. 3.

³² Millett, Journ. Roy. Micr. Soc., 1898, p. 612.

³³ Trans. Zool. Soc., vol. 20, 1915, p. 592.

Description.—Test differing from the typical in the character of the peripheral border, which is definitely thickened with a rounded periphery.

Chapman described this form as a variety of *C. carinata*, but in the Philippines it has seemed to be much more closely allied to *C. foliacea*. It occurs but once at the same station with *C. carinata*. Chapman's original notes are as follows:

In this variety the tests partake of the characters of both *C. carinata* Costa, sp., and *C. foliacea* Philippi, sp. The test, although increasing rapidly in width, as in *C. foliacea*, is, however, depressed on the sides, and even tends to become concave, as in typical examples of *C. carinata*. The larger specimen of the varietal form has a diameter of 1.23 mm.

Forty miles south of Cape Wiles (South Australia, lat. 35° 35' S.; long. 135° 15' E.) 100 fathoms. Two examples, identical in varietal form.

The variety has occurred in the *Albatross* Philippine material at seven stations, ranging in depth from 135 to 700 fathoms (247 to 1,280 meters), the average 415 fathoms (759 meters). Bottom temperatures given at four of these stations range from 49.3° F. to 59° F. (9.6° C. to 15° C.), the average 53.7° F. (12.1° C.). The localities include China Sea, off southern Luzon; off northwestern Panay; east coast of Luzon; between Siquijor and Bohol Islands; between Negros and Siquijor; and in the region southward it has occurred south of Patiente Strait and off Bouro Island.

This probably is a widely spread species in the Indo-Pacific region in water of 100 fathoms (183 meters), or more from the various records now known.

Cornuspira foliacea, var. *expansa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10728	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E. .	135	59.0	dk. gn. m. . .	Few.
10729	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E. .	312	49.3	gy. m., glob.	Few.
10730	U.S.N.M.	2	D5469...	13 36 48 N.; 123 38 24 E. .	500	gn. m. (net).	Few.
10731	U.S.N.M.	1	D5529...	9 23 45 N.; 123 39 30 E. .	441	53.0	gy. m., glob.	Few.
10732	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E. .	254	53.5	gn. m.	Few.
10733	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E. .	569	co. s., m.	Few.
10734	U.S.N.M.	1	D5637...	3 53 20 S.; 126 48 00 E. .	700	gy. m.	Few.

CORNUSPIRA INVOLVENS Reuss.

Plate 77, figs. 3, 4.

Operculina involvens REUSS, Denkschr. Akad. Wiss. Wien, vol. 1, 1849, p. 370, pl. 45, fig. 20.

Cornuspira involvens REUSS, Sitz. Akad. Wiss. Wien, vol. 48. 1863 (1864), p. 39, pl. 1, fig. 2.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 200, pl. 11, figs. 1-3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 25, pl. 1, fig. 2; pl. 2, fig. 2.

This is by far the most common species of the genus in the region, being widely distributed, but nowhere common. All the stations

but one are within the Archipelago, and range in depth from 78 to 569 fathoms (143 to 1,040 meters); bottom temperatures range from 41.2° to 59.6° F. (5.1 to 15.3° C.). These stations embrace the following general localities: Off Romblon; western Bohol; Panay; Verde Island Passage; off southern Luzon; eastern Luzon; off Mindanao; between Siquijor and Bohol; Negros and Siquijor; and Sulu Sea.

A few specimens were found at D5590, Sibuko Bay, Borneo, in 310 fathoms (567 meters), bottom temperature 44.3° F. (6.8° C.).

Heron-Allen and Earland³⁴ note that in their material from the Kerinba Archipelago "two or three distinct types occur" and give notes on the various forms. It probably indicates that if this and other widely distributed species are closely studied various forms may be distinguished.

Cornuspira involvens—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
10738	U.S.N.M.	1	D5120...	13 45 30 N.	120 30 15 E...	393	43.7	gn. m., s.....	Rare.
10739	U.S.N.M.	1	D5178...	12 43 00 N.	122 06 15 E...	78	fine. s.....	Rare.
10740	U.S.N.M.	1	D5198...	9 40 50 N.	123 39 45 E...	220	53.9	gn. m.....	Rare.
10741	U.S.N.M.	1	D5236...	8 50 45 N.	126 26 52 E...	494	41.2	fine. gy. s.....	Few.
		2	D5259...	11 57 30 N.	121 42 15 E...	312	49.3	gy. m., glob.	Rare.
10742	U.S.N.M.	1	D5288...	13 42 00 N.	120 57 15 E...	170	s., p.....	Rare.
10743	U.S.N.M.	1	D5278...	14 00 10 N.	120 17 15 E...	102	59.6	fine. s., m., sh.	Rare.
10744	U.S.N.M.	1	D5468...	13 35 39 N.	123 40 48 E...	569	gn. m.....	Rare.
10745	U.S.N.M.	1	D5512...	8 16 02 N.	123 58 26 E...	445	52.8	gy. m., fine. s.	Rare.
10746	U.S.N.M.	1	D5523...	8 48 44 N.	123 27 35 E...	Rare.
10747	U.S.N.M.	1	D5529...	9 23 45 N.	123 39 30 E...	441	53.0	gy. m., glob.	Rare.
10748	U.S.N.M.	2	D5537...	9 11 00 N.	123 23 00 E...	254	53.5	gn. m.....	Few.
10749	U.S.N.M.	1	D5570...	5 32 15 N.	120 12 57 E...	330	52.3	fine. s., glob.	Rare.
10750	U.S.N.M.	1	D5576...	5 25 56 N.	120 03 39 E...	277	53.3	s.....	Rare.
10751	U.S.N.M.	2	D5590...	4 10 50 N.	118 39 35 E...	310	44.3	gn. m., s.....	Few.

CORNUSPIRA INVOLVENS Reuss, var. SUBSTRIATULA, new variety.

Plate 77, fig. 5.

Description.—Test similar to the typical in general characters, but the exterior, instead of smooth, is marked by numerous fine longitudinal striations.

Distribution.—Type specimen (U.S.N.M. No. 10752) from *Albatross* station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao, bottom temperature 41.2° F. (5.1° C.). Specimens were common at this station, while very few of the typical form occurred. The variety was also noted from the following stations: D5282, in 248 fathoms (454 meters), China Sea, off southern Luzon, bottom temperature 47.4° F. (8.5° C.); D5567 in 268 fathoms (491 meters), bottom temperature 52.0° F. (11.1° C.); and D5571,

³⁴ Trans. Zool. Soc., vol. 20, 1915, p. 593.

in 340 fathoms (622 meters), bottom temperature 52.3° F. (11.2° C.), both north of Tawi Tawi.

Cornuspira involvens, var. *substriatula*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10752	U.S.N.M.	10+	D5236...	° ' " ° ' "		° F.		
				8 50 45 N.; 126 26 52 E...	494	41.2	fne. gy. s....	Common.
10753	U.S.N.M.	2	D5282...	13 53 00 N.; 120 26 45 E...	248	47.4	dk. gy. s....	Few.
10754	U.S.N.M.	2	D5567...	5 48 00 N.; 120 33 45 E...	268	52.0	fne. s.....	Few.
10755	U.S.N.M.	1	D5571...	5 30 45 N.; 120 07 57 E...	340	52.3	s., sh.....	Rare.

CORNUSPIRA LACUNOSA H. B. Brady.

Plate 78, fig. 1.

Cornuspira lacunosa H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 202, pl. 113, fig. 21.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 26, pl. 2, fig. 3.

Specimens of this rare species seem to confirm what was noted in the North Pacific material that the type figured by Brady was an immature specimen. In the adults the latter portion becomes nearly smooth and the last-formed coil wider. The size of the Philippine specimens also is nearer that of the Japanese material of this species.

Single specimens were obtained from the following stations: D5178 in 78 fathoms (143 meters), off Romblon; D5259, in 312 fathoms (571 meters), off northwestern Panay, bottom temperature 49.3° F. (9.6° C.); and D5281 in 201 fathoms (368 meters), China Sea, off southern Luzon, bottom temperature 50.4° F. (10.2° C.).

These few specimens confirm what I have already noted at some length in regard to the adult form of this species and its relation to what was evidently the young described by Brady from off Raine Island, Torres Strait, 155 fathoms (283 meters). The other material is that which I had from southern Japan. This species, while evidently of widespread distribution in the Indo-Pacific area, can not be very common, from the scattered records and few specimens at each station. The Philippine specimens are all microspheric.

Cornuspira lacunosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10735	U.S.N.M.	1	D5178...	° ' " ° ' "		° F.		
				12 43 00 N.; 122 06 15 E...	78	fne. s.....	Rare.
10736	U.S.N.M.	1	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Rare.
10737	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s....	Rare.

CORNUSPIRA CARINATA (Costa).

Plate 77, fig. 6.

Operculina carinata COSTA, Atti. Accad. Pont., vol. 7, 1856, p. 209, pl. 17, figs. 15, a, b.

Cornuspira carinata SEGUENZA, Atti. Accad. Gioenia di Sci. Nat., vol. 18, 1862, p. 93.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 201, pl. 11, figs. 4a, b.—EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 247, pl. 3, figs. 16, 17.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 11.

Description.—Test close-coiled, not flaring, increasing regularly in diameter; tube somewhat compressed laterally, the sides convex; peripheral margin with a thin but sharply defined narrow keel; surface smooth, aperture broadly elliptical.

Diameter up to 3 mm.

This is not a common species in the Pacific from the available records, yet its occurrence in the Philippine region at several stations with the records from off Raine Island, Torres Strait, 155 fathoms (283 meters), rare (Brady), and off the east coast of Australia in 465 fathoms (850 meters) (Sidebottom), show that it is well distributed in the Indo-Pacific. Egger recorded it from western Australia, from Mauritius, and from near New Amsterdam, extending the distribution westward across the Indian Ocean.

In the *Albatross* Philippine dredgings there are records from ten stations, ranging in depth from 297 to 890 fathoms (543 to 1,628 meters). These are off Baylayan Bay; Sogod Bay, southern Leyte; off northwestern Panay; west coast of Luzon; off northern Mindanao; north of Tawi Tawi; and to the southward off Darvel Bay and Sibuko Bay, Borneo.

Cornuspira carinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
				• • • • •		• F.		
10706	U.S.N.M.	1	D5114...	13 36 11 N.; 120 45 26 E...	340	fine s.	Rare.
10715	U.S.N.M.	5	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
10707	U.S.N.M.	3	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Few.
10708	U.S.N.M.	1	D5429...	9 41 30 N.; 118 50 22 E...	766	gn. m.	Rare.
10709	U.S.N.M.	1	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.	Rare.
10710	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E...	Rare.
10711	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.	Rare.
10712	U.S.N.M.	1	D5582...	4 19 54 N.; 118 58 38 E...	890	38.3	gy. m., fine s.	Rare.
10713	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.	Rare.
10714	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Rare.

CORNUSPIRA CRASSISEPTA H. B. Brady.

Plate 84, fig. 1.

Cornuspira crassisepta H. B. BRADY, Proc. Roy. Soc. Edinburgh, vol. 11, 1882, p. 714; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 202, pl. 113, fig. 20.—HOWCHIN, Trans. Roy. Soc. South Australia, vol. 12, 1889, p. 4.—EGGER, Abh. kön. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 246, pl. 3, fig. 22.—CHAPMAN, Journ. Linn. Soc., Zoology, vol. 30, 1907, p. 22, pl. 2, fig. 45; p. 399.

Description.—"Test discoidal, biconcave, peripheral edge nearly square; convolutions very numerous and very narrow near the center; spinal septal wall thick, and marked externally by a raised limbate line.

"Diameter, one-fiftieth of an inch (0.5 mm.) or rather more."

Brady described this species from specimens dredged in the "warm area" of the Faroe Channel. It is known from the work of Howchin, and especially that of Chapman, to be "to be the most abundant of the *Cornuspirae* in the Tertiaries of southeastern Australia." Egger records it in recent dredgings from *Gazelle* stations off Mauritius and off western Australia. Therefore it is not surprising to find it present in the *Albatross* Philippine material.

The specimen figured by Brady has a broader and more irregular thickening than our specimens. The few I have had agree very closely with that figured by Chapman as noted in the reference above. Except for this difference the specimens are very much alike. They were microspheric, as were Brady's and Chapman's figured specimens. The squarely truncate peripheral margin is very evident.

The species was rare at two stations—D5236, off the east coast of Mindanao in 494 fathoms (903 meters), and D5348, Palawan Passage, in 375 fathoms (686 meters).

Cornuspira crassisepia—Materials examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10716	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	° F. 41.2	fine. gy. s....	Rare.
10717	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co., s.....	Rare.

Genus **OPHTHALMIDIUM** Zwingli and Kübler, 1870.

OPHTHALMIDIUM INCONSTANS (H. B. Brady).

Plate 78, figs. 2, 3.

Hauerina inconstans H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 54.

Ophthalmidium inconstans H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 189, pl. 12, figs. 5, 7, 8.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 28, pl. 3, figs. 1-4.

Except at one station, D5236, where specimens were fairly frequent, the species has occurred only as scattered specimens: Sulu Sea; Verde Island Passage; east coast Mindanao; off northwestern Panay; southern Luzon; Palawan Passage; also from Sibuko Bay, Borneo (stations D5585, 5586), and D5637 in 700 fathoms (1,280 meters), vicinity of Bouro Island. The depths in the Archipelago proper range from 201 to 494 fathoms (368 to 903 meters), the greater depth,

that of D5236, where the specimens were most frequent. Some of the specimens are fine and large but all were very typical.

In the Indo-Pacific region the species is known from off Funafuti (Chapman), off western Australia (Egger), and off the east coast of Australia (Sidebottom).

A part of the material referred by Sidebottom to *Spiroloculina nitida* d'Orbigny³⁵ seems to be *Ophthalmidium inconstans*, especially figure 6, which is very typical. This may be compared to the figure I have given³⁶ of the optical section.

Ophthalmidium inconstans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
			D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine gy. s....	Frequent.
			D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	
			D5586...	4 06 50 N.; 118 47 20 E..	415	42.3	Co., s.....	
			D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.....	

Genus SPIROLOCULINA d'Orbigny, 1826.

SPIROLOCULINA DEPRESSA d'Orbigny.

Plate 81, fig. 2; plate 100, figs. 4, 5.

Spiroloculina depressa d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 298.—WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 82, pl. 7, fig. 177.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 29, pl. 3, figs. 6–10.

Spiroloculina limbata H. B. BRADY (not d'Orbigny), Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 150, pl. 9, figs. 15–17.

This widely distributed species, characteristic of open ocean, is rare in the Philippine region. The four stations represented the following localities: China Sea, off Formosa; Sulu Sea, off western Mindanao; and between Burias and Luzon.

Spiroloculina depressa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
						° F.		
10955	U.S.N.M.	5	D5132...	Island off Panabutan Point, N. 15°; W., 0.3 m.	26	gn. m., s....	Frequent.
10956	U.S.N.M.	2	D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s....	Few.
10957	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Few.
10958	U.S.N.M.	1	D5388...	12 51 30 N.; 123 26 15 E..	226	51.4	sft. gn. m....	Few.

³⁵ Journ. Roy. Micr. Soc., 1918, pl. 1, figs. 5, 6 (not fig. 7).

³⁶ Bull. 71, U. S. Nat. Mus., pt. 6, 1917, pl. 3, fig. 3.

SPIROLOCULINA CANALICULATA d'Orbigny.

Plate 80, figs. 3 a, b.

Spiroloculina canaliculata d'ORBIGNY, *Foram. Foss. Vienne*, 1846, p. 269, pl. 16, figs. 10-12.—CUSHMAN, *Bull. 71, U. S. Nat. Mus.*, pt. 6, 1917, p. 30, pl. 4, figs. 1-3.

Spiroloculina impressa H. B. BRADY (not Terquem), *Rep. Voy. Challenger, Zoology*, vol. 9, 1884, p. 151, pl. 10, figs. 3, 4.

This species is characterized by the deeply concave periphery between the sharply extended sides of the chambers.

It occurs in characteristic form at 24 stations in the region, ranging in depth from 18 to 1,262 fathoms (33 to 2,309 meters), the average 204 fathoms (373 meters). Bottom temperatures for the 10 stations recorded range from 44.3° to 75.7° F. (6.8° to 24.2° C.), with the average slightly less than 57° F. (13.8° C.).

The stations at which it is recorded as common are in depths, for the most part, of less than 100 fathoms (183 meters). At the others the specimens were usually few.

The following general localities are represented: Sulu Sea, off western Mindanao; about the Tawi Tawi group and near Basilan; China Sea, off southern Luzon and off Formosa; Palawan Passage, between Leyte and Mindanao; off Romblon; off eastern Panay; off northern Cebu; Sogod Bay, southern Leyte; east of Masbate; between Burias and Luzon; between Marinduque and Luzon; between southeastern Mindoro and Verde Island Passage. These localities cover the Archipelago very well except the open Pacific coast. Southward a few specimens were found at stations in Darvel Bay and Sibuko Bay, Borneo, and from Pitt Passage.

Spiroloculina canaliculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10861	U.S.N.M.	10+	D5110...	° " " ° " "	135	59.0	dk. gy. m...	Common.
10862	U.S.N.M.	3	D5132...	13 59 20 N.; 120 75 45 E... Island off Panabutan N. 15° W.; 0.3 mile.	26	gn. m., s.....	Few.
10863	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N.; 52° E.; 1.5 miles.	38	gn. m., s.....	Common.
10864	U.S.N.M.	2	D5134...	6 44 45 N.; 121 48 00 E...	25	fine s.....	Few.
10865	U.S.N.M.	8	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Common.
10866	U.S.N.M.	3	D5153...	5 18 10 N.; 120 02 55 E...	49	co. s., sh.....	Few.
10867	U.S.N.M.	4	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Few.
10868	U.S.N.M.	5	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fine s.....	Few.
10869	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Few.
10870	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	53.8	gy. s., m.....	Few.
10871	U.S.N.M.	1	D5212...	12 04 15 N.; 124 04 36 E...	108	59.9	gy. s., m.....	Rare.
10872	U.S.N.M.	9	D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	ers. gy. s.....	Common.
10873	U.S.N.M.	6	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn., m.....	Common.
10874	U.S.N.M.	1	D5261...	12 30 55 N.; 121 34 24 E...	145	s., m.....	Few.
10875	U.S.N.M.	7	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Common.
10876	U.S.N.M.	2	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p. s.....	Few.
10877	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s.....	Rare.
10878	U.S.N.M.	3	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Few.
10879	U.S.N.M.	2	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co.....	Few.
10880	U.S.N.M.	3	D5238...	11 33 45 N.; 119 24 45 E...	43	co. s., m.....	Few.
10881	U.S.N.M.	1	D5495...	9 06 30 N.; 125 00 20 E...	976	52.3	gy. m.....	Few.
10882	U.S.N.M.	3	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co.....	Few.
10883	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.....	Few.
10884	U.S.N.M.	1	D5636...	1 55 00 S.; 127 42 30 E...	1,262	gy. m., fine s.	Few.

SPIROLOCULINA GRATELOUPI d'Orbigny.

Plate 78, figs. 4a, b; plate 100, fig. 3.

Spiroloculina grateloupi d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 298.—TERQUEM, Mém. Soc. géol. France, ser. 3, vol. 1, 1878, p. 52. pl. 5, figs. 5, 6.—WIESNER, Archiv. Prot., vol. 25, 1812, p. 208.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 31, pl. 4, figs. 4, 5.

Spiroloculina excavata H. B. BRADY (not d'Orbigny), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 151, pl. 9, figs. 5, 6.—HERON-ALLEN and EARLAND, Trans. Zool. Soc., London, vol. 20, 1915, p. 554.

The characters of this species include a deeply excavate central portion, great thickness at the periphery, elongate test; peripheral margin broadly convex, the last-formed chamber projecting at both ends of the test, the apertural end with a decided projecting neck and phialine lip.

Both megalospheric and microspheric forms occur, the latter reaching 3 mm. in length.

Brady's notes in the *Challenger* Report are instructive in regard to this species. Under *S. excavata* he says:

The most "excavated" specimens take the somewhat elongate contour represented in the drawings, and have the final segment extending a little beyond the rest of the shell. Such forms must be sought chiefly in the shallow water surrounding the coral islands of the Pacific.


Heron-Allen and Earland note in their material from the Kerimba Archipelago off southeastern Africa, under *S. excavata*.

The typical *S. excavata* has a nearly flat peripheral edge, but the bulk of our specimens are very rounded * * *. This round-edged variety occurs at many stations.

From these notes and an examination of the Philippine material and that from other tropical localities of the Pacific, it seems evident that this species is very well represented if not the dominant species of the genus in the Indo-Pacific. It certainly is different from *S. excavata*, and may be later separated from *S. grateloupi* d'Orbigny.

It is the most common *Spiroloculina* in the Philippine region, especially in shallow water, often occurring in great numbers and not showing any great amount of variation. The microspheric form is much larger and more excavate than the megalospheric.

It has been noted at 32 stations, all but one in the Archipelago or the China Sea, none of them off the Pacific coast in deeper water. It is especially abundant in the shallow water about the Sulu Sea. The only record in the deeper water to the southward is in Sibuko Bay, Borneo.



FIGS. 17, 18.—SPIROLOCULINA GRATELOUPI D'ORBIGNY. $\times 50$. END VIEWS SHOWING APERTURE AND TEETH. 17, FROM ALBATROSS STATION D5136. 18, FROM ALBATROSS D5136.

The depths range from 10 to 554 fathoms (18 to 1,012 meters), the average, however, being about 70 fathoms (128 meters). The bottom temperatures where given range from 43.3° to 75.7° F. (6.1° to 24.2° C.), with the average 58.3° F. (14.6° C.). This is too low, however, as it is only from eight of the deeper stations. Both sets of data show that the species is most at home in shallow, very warm waters.

Spiroloculina grateloupi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		* F.		
10906	U.S.N.M.	1	D5097...	14 19 15 N.; 120 33 52 E..	30	gy. m., sh...	Few.
10907	U.S.N.M.	2	D5100...	14 17 15 N.; 120 32 40 E..	35	gy. s.....	Few.
10908	U.S.N.M.	4	D5106...	14 23 55 N.; 120 32 33 E..	37	gy. m.....	Common.
10909	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m...	Few.
10910	U.S.N.M.	8	D5133...	Island off Panabutan Point, N. 52° E., 1.5 miles.	38	gn. m., s....	Common.
10911	U.S.N.M.	10	D5134...	6 44 45 N.; 121 48 00 E..	25	fine. s.....	Common.
10912	U.S.N.M.	2	D5136...	6 04 20 N.; 120 59 20 E..	22	s., sh.....	Few.
10913	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E..	19	s., Co.....	Few.
10914	U.S.N.M.	2	D5139...	6 06 00 N.; 121 02 30 E..	20	co. s.....	Few.
10915	U.S.N.M.	7	D5145...	6 04 36 N.; 120 59 30 E..	23	co. s., sh...	Common.
10916	U.S.N.M.	3	D5146...	5 46 40 N.; 120 48 50 E..	24	co. s., sh...	Few.
10917	U.S.N.M.	2	D5148...	5 35 40 N.; 120 47 30 E..	17	co. s.....	Few.
10918	U.S.N.M.	1	D5149...	5 33 00 N.; 120 42 10 E..	10	Co., sh.....	Few.
10919	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E..	34	oh. s.....	Common.
10920	U.S.N.M.	1	D5154...	5 14 50 N.; 119 58 45 E..	12	co. s.....	Few.
10921	U.S.N.M.	3	D5156...	5 12 50 N.; 119 55 55 E..	18	fine. s., sh...	Few.
			D5157...	5 12 30 N.; 119 55 50 E..	18	fine. s., sh...	Few.
10922	U.S.N.M.	9	D5159...	5 11 50 N.; 119 54 00 E..	10	co. s.....	Common.
10923	U.S.N.M.	4	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Few.
10924	U.S.N.M.	2	D5181...	11 36 40 N.; 123 26 35 E..	26	m., fine. s...	Frequent.
10925	U.S.N.M.	8	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.....	Frequent.
10926	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s. m....	Few.
10927	U.S.N.M.	1	D5212...	12 04 15 N.; 124 04 36 E..	108	59.9	gy. s. m....	Rare.
10928	U.S.N.M.	1	D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s...	Rare.
10929	U.S.N.M.	10+	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Common.
10930	U.S.N.M.	2	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m...	Few.
10931	U.S.N.M.	8	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s...	Common.
10932	U.S.N.M.	6	D5277...	13 56 55 N.; 120 13 45 E..	80	58.6	fine. s.....	Common.
10933	U.S.N.M.	6	D5478...	10 46 24 N.; 125 16 30 E..	57	sh.....	Common.
10934	U.S.N.M.	10	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Common.
10935	U.S.N.M.	7	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Common.
10936	U.S.N.M.	9		Jolo Jolo.....	Common.

SPIROLOCULINA GRATELOUPI d'Orbigny, var. INCISA, new variety.

Plate 78, fig. 5.

Description.—Variety differing from the typical in the adding of the chambers, the inner margin of the newly added chamber not being equal to the outer margin of the preceding chamber, thus leaving a distinct sharp ridge between each of two chambers as added. This leaves a series of incised areas with high areas alternating with them.

Distribution.—Type specimen (Cat. No. 10941, U.S.N.M.) from *Albatross* station D5220, between Marinduque and Luzon, in 50 fathoms (91 meters). It has also occurred at six other stations: Off Romblon; off western Samar; between Burias and Luzon; Palawan Passage; and off Marinduque.

These seven stations ranged in depth from 20 to 106 fathoms (37 to 194 meters), the average 64 fathoms (117 meters). The bottom temperature is given for only two of the stations, averaging nearly 70° F. (21.1° C.).

Spiroloculina grateloupi, var. *incisa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10937	U.S.N.M.	10+	D5178 ..	12 43 00 N.; 122 06 15 E....	78	fine.s.....	Common.
10938	U.S.N.M.	1	D5210...	11 49 55 N.; 124 28 05 E....	50	76.3	fine.gy.s.....	Rare.
10939	U.S.N.M.	3	D5217...	13 20 00 N.; 123 14 15 E....	105	63.1	crs.gy.s.....	Frequent
10940	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E....	20	crs.s.....	Rare.
10941	U.S.N.M.	10+	D5220...	13 38 00 N.; 121 58 00 E....	50	sft.gn.m.....	Common.
10942	U.S.N.M.	5	D5338...	11 33 45 N.; 119 24 45 E....	43	co.s.,m.....	Common.
10943	U.S.N.M.	1	D5369...	13 48 00 N.; 121 43 00 E....	106	bn.s.....	Rare.

SPIROLOCULINA ACUTIMARGO H. B. Brady.

Spiroloculina acutimargo H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 154, pl. 10, figs. 12-15.—EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 222, pl. 1, figs. 26-28.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 264.—BAGG, Proc. U. S. Nat. Mus., vol. 34, 1908, p. 119.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 557.—SIDE-BOTTOM, Journ. Roy. Micr. Soc., 1918, p. 5 (?).

A reference to the original figures of this species given by Brady in the *Challenger* Report shows that evidently more than one species is included. Figure 12 strongly suggests the species described by Schlumberger as *S. inaequilateralis*, but no end view is given on Brady's plate. Figure 14 seems to belong to *Massilina*. Of the other figures 13 and 15 may be taken to represent the species *S. acutimargo*.

Brady's localities also show different faunal areas. These, as given in the *Challenger* Report, are "off Bermuda, at a depth of 435 fathoms (796 meters); at four stations in the South Atlantic, 350 to 1,425 fathoms (640 to 777 meters); at three localities in the South Pacific, 15 to 255 fathoms (27 to 466 meters); it has also been found in shore sands collected on the east coast of Madagascar."

The South Pacific 15 to 255 fathoms (27 to 466 meters) and shore sands of Madagascar evidently represent a faunal area, and with no further data it would be a guess that the large specimen (fig. 15) of the *Challenger* Report came from some of these localities. This is strengthened by Heron-Allen and Earland, who remark on their specimen from the Kerimba Archipelago (p. 557):

A very finely developed specimen at station 11, exhibiting the characteristic lacunae at the extremities of the chambers.

Egger's figures are not clear; his records are off Kerguelen, New Amsterdam; West Australia; and the Fiji Islands. Chapman records it

from Funafuti. The material from western Australia Sidebottom definitely refers to *Challenger* figures 12 and 13. Millett gives no figure and simply says: "A few small specimens from both areas" (Malay Archipelago).

With this more or less indefinite recording and the fact that the specimens from about the British Isles do not seem related to the Indo-Pacific material it would seem that there quite likely may be several species now going under this name. I would limit the name to those specimens of the Indo-Pacific like figure 15 of the *Challenger* Report and possibly the figure 13, which may be an earlier stage of the same.

In the Philippine material it has occurred at but one station, D5242, off Pujada Bay, in 191 fathoms (350 meters).

Spiroloculina acutimargo—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10948	U.S.N.M.	1	D5242...	6 51 53 N.; 126 14 10 E..	191	° F. 64.1	sft. gy. m....	Rare.

SPIROLOCULINA TENUIMARGO Cushman.

Plate 84, fig. 3.

Spiroloculina tenuimargo CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 32, pl. 5, figs. 2, 3.

Description.—Test much compressed, planospiral; chambers numerous, usually more swollen toward the basal end, inflated, nearly circular in transverse section, the apertural end somewhat produced, surface smooth; periphery of the test with a distinct, rather broad, sharp, plate-like carina, in young specimens rather even, but in older ones more or less irregular; chambers closely adjacent to one another, not separated; apertural end with a slightly produced and slightly flaring lip; aperture round.

Length slightly less than 1 mm.

This species, originally described from off Guam and off Japan, has occurred at two stations in the Philippine area—D5349, Palawan Passage, in 730 fathoms (1,335 meters), and D5445, east coast of Luzon, in 383 fathoms (699 meters).

Egger's figures of *S. acutimargo* suggest that they may be *S. tenuimargo*.

Spiroloculina tenuimargo—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10963	U.S.N.M.	1	D5349...	10 54 00 N.; 118 29 00 E...	730	40.6	co. s.....	Rare.
10964	U.S.N.M.	1	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s....	Rare.

SPIROLOCULINA TENUISSIMA Reuss.

Plate 84, fig. 2.

Spiroloculina tenuissima REUSS, Sitz. Akad. Wiss. Wien, vol. 55, 1867, p. 71, pl. 1, fig. 11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 32.

Spiroloculina tenuis H. B. BRADY (not Czjzek), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 152, pl. 10, figs. 9, 10 (not 7, 8, 11).—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 556.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 5.

Description.—Test very thin, translucent, elongate, compressed, in front view tapering toward either end; chambers long and narrow, peripheral margin broadly rounded, chambers in transverse section, circular; apertural end produced into an elongate neck, aperture rounded, usually without distinct teeth; wall smooth.

Length usually less than 0.5 mm.

The records for this species are mostly from water of several hundred fathoms, and the four Philippine stations range from 295 to 554 fathoms (540 to 1,012 meters). The localities are off Balayan Bay, Tanon Strait; east coast of Negros; Sogod Bay, southern Leyte; and east of Mindoro.

Sidebottom definitely refers one of his two varieties to the *Challenger* Report (pl. 10, fig. 9), which would make it this species. His material was from 465 fathoms. (850 meters), off the east coast of Australia. Chapman records it off Funafuti, in deep water.

Spiroloculina tenuissima—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5120...	13 45 30 N.; 120 30 15 E...	393	43.7	gn. m., s....	Rare.
10970	U.S.N.M.	1	D5190...	10 08 15 N.; 123 16 45 E...	295	63.0	gn. m.	Rare.
10971	U.S.N.M.	2	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
10972	U.S.N.M.	1	D5227...	12 53 45 N.; 121 52 30 E...	322	gn. m.	Rare.

SPIROLOCULINA MILETTI Wiesner.

Plate 81, figs. 6, 7.

Spiroloculina nitida H. B. BRADY (not *S. nitida* d'Orbigny), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 149, pl. 9, figs. 9, 10.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 265, pl. 5, figs. 9–12.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 296, pl. 41, fig. 4.—BAGG, Proc. U. S. Nat. Mus., vol. 34, 1908, p. 119.

Spiroloculina milletti WIESNER, Archiv. Prot., vol. 25, 1912, p. 207.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 33, pl. 5, fig. 4.

This seems to be distinct from *S. nitida*, and is very evidently a tropical species. It has occurred in the Philippine material but once, in a collection marked "Taeloban Anchorage." It is known from other tropical parts of the Pacific and elsewhere.

Spiroloculina milletti—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10947	U.S.N.M.	2	Taeloban Anchorage.....	* F.	Rare.

SPIROLOCULINA PLANISSIMA (Lamarck).

Plate 80, figs. 5a, b.

Miliolites planulata, var. *planissima* LAMARCK, Anim. sans. Vert., vol. 7, 1822 p. 613, no. 4c.

Spiroloculina planissima WIESNER, Archiv. Prot., vol. 25, 1912, p. 209.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 556, pl. 41, figs. 1-5.

This is a fine large species, with the chambers slightly convex and smooth, the peripheral margin carinate, the apertural end extended in an elliptical neck with a bifid tooth.

Length up to 2 mm. or more.

Heron-Allen and Earland had this species from the Kerimba Archipelago, off the southeastern coast of Africa. It has occurred at four stations in the Philippine material, ranging in depth from 14 to 108 fathoms (25 to 198 meters), off Romblon; east of Masbate; Malampaya Sound, Palawan Island, and off eastern Palawan.

Spiroloculina planissima—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
				* , " * , "		* F.		
10959	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	fine.s.....	Rare.
10960	U.S.N.M.	1	D5212...	12 04 15 N.; 124 04 36 E..	108	59.9	gy.s.,m.....	Rare.
10961	U.S.N.M.	2	D5342...	10 56 55 N.; 119 17 24 E..	14-25	gy.m.....	Rare.
10962	U.S.N.M.	1	D5426...	9 12 00 N.; 118 28 00 E..	27	fine.gy.s.....	Few.

SPIROLOCULINA TENUISEPTATA H. B. Brady.

Plate 82, figs. 1-3.

Spiroloculina tenuiseptata H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 153, pl. 10, figs. 5, 6.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 265.—CHAPMAN, Journ. Linn. Soc. Zoology, vol. 28, 1901, p. 172.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 5, pl. 1, fig. 7 (not figs. 5, 6).

Description.—"Test complanate, elongate-oval; extremities tapering, subangular, peripheral edge square or rounded. Segments nu-

merous, narrow, arched, tubular; the successive convolutions separated by deep depressions on both sides of the test, the interspace being occupied by a thin, horizontal, shelly septum, which is sometimes wanting between the later chambers; aperture simple, circular.

Length, one twenty-second inch (1.2 mm.)."

Brady's material was from off the Ki Islands, 580 fathoms (1,061 meters); off Kandavu, Fiji, 610 fathoms (1,116 meters); and the Mediterranean, 1,200 fathoms (2,200 meters). Egger records the species from off West Africa, but his figure is difficult to make out. Millett records it from the Malay Archipelago, and Chapman from the lagoon at Funafuti. Sidebottom records it from off eastern Australia, 465 fathoms (850 meters), and plate 1, figure 7, certainly seems to be this species.

In the Philippine material dredged by the *Albatross* the species has occurred in considerable numbers at D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters), and at six other stations, the depths of all ranging from 230 to 565 fathoms (420 to 1,033 meters), with the average 409 fathoms (748 meters). Bottom temperatures of six of these stations range from 40.1° F. to 52.9° F. (4.5° C. to 11.6° C.), with the average 45.7° F. (7.6° C.). The localities represented, besides the one already mentioned, are: Tawi Tawi Group, Sulu Archipelago; off northwestern Panay; east coast of Luzon; and to the southward, Sibuko Bay, Borneo, and Gulf of Boni.

My friend, Mr. Sidebottom, has sent me some specimens from the coast of Ireland which were identified by Brady as this species. They are much smaller than the Philippine specimens, less elongate, and in one beautifully sectioned specimen there is no "shelly septum" between the chambers. After seeing this material I am inclined to think that the true *S. tenuiseptata* is largely an Indo-Pacific species.

One very characteristic feature in adults is the shape of the apertural end of the chamber, which becomes free and takes a different line of growth from the rest of the chambers, and also is apt to become shorter than the half volution.

Spiroloculina tenuiseptata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
10898	U.S.N.M.	1	D5162...	° ' " ° ' "	230	° F. 52.9	crs. s., brk. sh.	Rare.
10899	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fne. gy. s.....	Common.
10900	U.S.N.M.	1	D5238...	7 34 45 N.; 126 38 15 E..	350	43.0	gn. m.....	Few.
10901	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob..	Few.
10902	U.S.N.M.	1	D5460...	13 32 30 N.; 123 58 06 E..	565	gy. m.....	Rare.
10903	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.....	Rare.
10904	U.S.N.M.	1	D5650...	4 53 45 N.; 121 29 00 E..	540	40.1	gn. m.....	Rare.
10905	U.S.N.M.	1	H4897...	7 46 00 N.; 122 00 00 E..	1,570	gy. m., glob..	Rare.

SPIROLOCULINA CIRCULARIS, new species.

Plate 80, figs. 1, 2.

Description.—Test circular in face view, composed of a large number of chambers, narrow in face view; sides concave; periphery slightly convex, broad, last-formed chamber only projecting a little beyond the general circular contour of the whole test; sutures flush with the surface, but of clear shell material making them stand out as darker lines; apertural end with a broad, low aperture, the edges being slightly thickened and with a low tooth; surface smooth and polished.

Diameter up to 1.5 mm.

Distribution.—Type specimen (Cat. No. 10897, U. S. N. M.) from *Albatross* station D5259, off northwestern Panay, in 312 fathoms (571 meters), bottom temperature 49.3° F. (9.6° C.).

This is a beautiful species frequent at this station. Its circular shape, concave sides, broad periphery, sutures forming a series of concentric circles with the two zigzag lines when the chambers meet; very broad low chambers and aperture, and the surface smooth and polished, should easily distinguish this species. It was found nowhere else in the region, yet is frequent in the material from this one station.

Spiroloculina circularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10897	U. S. N. M.	4	D5259...	11 57 30 N.; 121 42 15 E..	312	° F. 49.3	gy. m., glob..	Frequent.

SPIROLOCULINA ORBIS, new species.

Plate 83, fig. 3.

Description.—Test subcircular in front view, thick, the periphery squarely truncate; chambers rather few; sides flat, except near the periphery, when there is a slight thickening; ends of the final chamber slightly extended, apertural end with a short neck and rounded aperture with a very slight collar-like lip; surface smooth but dull.

Length usually less than 1 mm.

Distribution.—Type specimen (Cat. No. 10951, U. S. N. M.) from *Albatross* station D5178, off Romblon, in 78 fathoms (143 meters), where it is frequent. The specimens show very little variation of any sort. It was also found in material from D5179, off Romblon, in 37 fathoms (68 meters); D5269, Verde Island Passage, in 220 fathoms (403 meters); and D5277, China Sea, off southern Luzon, in 80 fathoms (146 meters).

Spiroloculina orbis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10951	U.S.N.M.	10+	D5178	° ' " ° ' "		° F.		
				12 43 00 N.; 122 06 15 E...	78	fne. s.	Frequent.
10952	U.S.N.M.	1	D5179	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.	Rare.
10953	U.S.N.M.	1	D5269	13 39 50 N.; 120 59 30 E...	220		fne. s., p.	Rare.
10954	U.S.N.M.	1	D5277	13 56 55 N.; 120 13 45 E...	80	53.6	fne. s.	Few.

SPIROLOCULINA ROBUSTA H. B. Brady.

Plate 79, figs. 2a, b.

Spiroloculina robusta H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 150, pl. 9, figs. 7, 8.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 296, pl. 42, figs. 1, 2.—CHAPMAN, Journ. Linn. Soc., Zoology, vol. 28, 1901, p. 170; vol. 30, 1910, p. 396.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 33.

This is another species with a peculiar distribution so far as available records show. Brady described it from off Culebra Island, West Indies, in 390 fathoms (713 meters), and Flint records it from the Gulf of Mexico, 200 to 1,200 fathoms (366 to 8,366 meters). Chapman records it from off Funafuti in 200 and 1,485 fathoms (366 and 2,716 meters), in the lagoon, one specimen in 7½ fathoms (13.5 meters); and I have had it from the Western Pacific in 814 and 1,040 fathoms (11,476 and 1,902 meters). In the *Albatross* Philippine Expedition it was dredged at station D5650, in the Gulf of Boni, in 540 fathoms (988 meters).

Spiroloculina robusta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10950	U.S.N.M.	4	D5650	° ' " ° ' "		° F.		
				4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.	Frequent.

SPIROLOCULINA VALIDUS, new species.

Plate 79, figs. 3a, b.

Description.—Test large, composed of but few chambers, sub-circular or broadly elliptical in front view; periphery broadly rounded, inner portion depressed, apertural portion extended into a definite cylindrical or slightly tapering neck; aperture circular, surface without ornamentation, slightly granular, dull.

Length up to 2.5 mm.

Distribution.—Type specimen (Cat. No. 10973, U.S.N.M.) from *Albatross* station D5668, Macassar Strait, in 901 fathoms (1,648 meters). It also occurred at six other stations, ranging in depth

from 162 to 901 fathoms (297 to 1,648 meters), the average about 498 fathoms (911 meters). Bottom temperatures of three stations range from 38.2° F. to 55.8° F. (3.4° to 13.2° C.), with the average temperature 45° F. (7.2° C.). Other localities are east coast of Luzon; Darvel Bay and Sibuko Bay, Borneo; and off Bouro Island.

This species is close to *S. robusta* H. B. Brady, from which it differs mainly in the apertural form and characters. It may be that the material referred to *S. robusta* in the Indo-Pacific should belong to *S. validus*, and Brady's name be used only for the Atlantic material. A study of this will later be made to see if this may not be the fact.

Spiroloculina validus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
10974	U.S.N.M.	5	D5446...	12 43 51 N.; 124 59 18 E...	300	gn. m.....	Frequent.
10975	U.S.N.M.	3	D5460...	13 32 30 N.; 123 58 06 E...	565	gy. m.....	Frequent.
10976	U.S.N.M.	2	D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m.....	Rare.
10977	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co....	Rare.
10978	U.S.N.M.	1	D5585...	4 07 00 N.; 118 49 54 E...	476	41.1	gy. m.....	Rare.
10979	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Few.
10973	U.S.N.M.	6	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.....	Common.

SPIROLOCULINA COSTIFERA Cushman.

Plate 79, figs. 1a, b.

Spiroloculina costifera CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 34, pl. 6, figs. 1-3.

In the North Pacific this species was recorded from off Midway and the Hawaiian Islands. The only Philippine material is from *Albatross* station D5236, Pacific Ocean, off Mindanao, in 494 fathoms, (903 meters). A single specimen here is referred to this species, although the following variety is common.

Spiroloculina costifera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
10946	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Rare.

SPIROLOCULINA COSTIFERA Cushman, var. PLENA, new variety.

Plate 82, figs. 5-7.

Description.—Variety differing from the typical in the smooth walls, but with the same rounded form of chambers, excavate on the inner margins, and very rounded periphery.

Distribution.—Type specimen (Cat. No. 10980, U.S.N.M.) from Albatross station D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters), where the variety is very common.

In this variety the whole chamber is rounded, without the costae of the typical form. In adult specimens the chambers decrease in length and fail to make up the entire half revolution as in the typical.

Spiroloculina costifera, var. *plena*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10980 10981	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fine. gy. s....	Common.

SPIROLOCULINA SCROBICULATA, new species.

Plate 81, figs. 1a, b.

Description.—Test elliptical in face view, composed of numerous chambers, each circular in transverse section, sutures deep; apertural end continued into an elongate cylindrical neck with a slight everted lip; surface peculiarly scrobiculate almost irregularly reticulate with very fine markings, with low magnification appearing simply granular.

Length up to 1.25 mm.

Distribution.—Type specimen (Cat. No. 10944, U. S. N. M.) from Albatross station D5181, off eastern Panay, in 26 fathoms (48 meters). It has also occurred at D5342, Malampaya Sound, Palawan, in 14 to 25 fathoms (25 to 46 meters).

This is a very peculiarly ornamented species, and should be looked for elsewhere in shallow water of the Indo-Pacific region.

Spiroloculina scrobiculata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10944	U.S.N.M.	3	D5181...	11 36 40 N.; 123 35 35 E..	26	° F.	m., fine. s....	Few.
10945	U.S.N.M.	1	D5342...	10 56 55 N.; 119 17 24 E..	14-25	gy. m.	Few.

SPIROLOCULINA ELEGANS Cushman.

Plate 80, figs. 4a, b.

Spiroloculina elegans CUSHMAN, Carnegie Inst., Dept. of Marine Biology, vol. 9, 1918, p. 290, pl. 96, figs. 1a, 1b, 2.

Description.—Test one and a half as long as wide, much compressed, peripheral margin broadly rounded, apertural end somewhat exserted,

rounded; sutures distinctly depressed, wall ornamented with a regular pattern of elliptical depressions, arranged in rows longitudinally, the depressions of each row alternating regularly with those of the adjacent rows.

Length up to 1 mm. or slightly more.

The species was described from Murray Island, on the Great Barrier Reef of Australia, from shallow water. Its nearest relative seems to be *S. foveolata* Egger, from which it differs very much in shape and general appearance.

It was found in few numbers at five stations in 19 to 318 fathoms (35 to 582 meters); bottom temperature at the only station given, 63.1° F. (17.2° C.).

Spiroloculina elegans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10965	U.S.N.M.	2	D5143...	6 05 50 N.; 121 02 15 E..	19	° F.	co. s.....	Rare.
10966	U.S.N.M.	1	D5151...	5 24 40 N.; 120 27 15 E..	24	co. s., sh.....	Rare.
10967	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E..	318	fne. s., sh.....	Rare.
10968	U.S.N.M.	1	D5217...	13 20 00 N.; 123 14 15 E..	105	63.1	crs. gy. s.....	Rare.
10969	U.S.N.M.	1	Tacloban Anchorage, P. I.	Rare.

SPIROLOCULINA ANTILLARUM d'Orbigny.

Plate 81, figs. 4a, b; plate 83, fig. 4.?

Spiroloculina antillarum D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat. Cuba, "Foraminifères," 1839, p. 166, pl. 9, figs. 3, 4.—(?) H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 155, pl. 10, figs. 21, a, b.

Spiroloculina grata TERQUEM (?) and late authors.

A study of shallow-water West Indian material and comparison with d'Orbigny's figures show that our common typical species so characteristic of coral-reef regions should have the name *S. antillarum* d'Orbigny.

A comparison of the two references given by Brady in the Challenger Report shows that of the two, d'Orbigny's figure is more characteristic, not only of most of our coral-reef material, but actually nearer the *S. grata* as figured by Brady. On the other hand, Brady refers to *S. antillea* and figures a specimen which only remotely suggests the original of d'Orbigny. This specimen was from 350 fathoms (640 meters) off the coast of Brazil, while d'Orbigny's original was from "shore sands." The West Indian material which I collected in Jamaica shows clearly that the common shallow-water species of those waters is identical with the species figured and named by d'Orbigny *S. antillarum*. The species seems to be very widely distributed in the tropics, and even allowing that Terquem's Pliocene specimen from the Isle of Rhodes is the same, d'Orbigny's name

should have priority. It is one of the most unfortunate choices of Brady, made in the *Challenger* Report, which has been blindly followed by later writers.

As elsewhere in the tropics this species is common in the Philippines, especially in shallow water. The stations range in depth from 10 to 305 fathoms (18 to 558 meters), the average being close to 90 fathoms (165 meters). It is most common at the shallower-water stations. It is most frequent in the southern part of the Archipelago about the shores of the Sulu Sea and occurs at one station in Sibuko Bay, Borneo.

Spiroloculina antillarum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
10982	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E..	10	Co.....	Few.
10983	U.S.N.M.	1	D5136...	6 04 20 N.; 120 59 20 E..	22	s., sh.....	Few.
10984	U.S.N.M.	2	D5145...	6 04 30 N.; 120 59 30 E..	23	co. s., sh.....	Few.
10985	U.S.N.M.	1	D5146...	5 46 40 N.; 120 48 50 E..	24	co. s., sh.....	Few.
10986	U.S.N.M.	3	D5154...	5 14 50 N.; 119 58 45 E..	12	co. s.....	Few.
10987	U.S.N.M.	10	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Common.
10988	U.S.N.M.	10+	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Common.
10989	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Few.
10990	U.S.N.M.	2	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., sh.....	Few.
10991	U.S.N.M.	1	D5578...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Few.
10992	U.S.N.M.	2	D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.....	Few.
10993	U.S.N.M.	2	Jolo Jolo, P. I.	Few.

SPIROLOCULINA ANTILLARUM d'Orbigny, var. ANGULATA Cushman.

Plate 81, figs. 5a, b.

Spiroloculina grata H. B. BRADY (part), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, pl. 10, figs. 22, 23.

Spiroloculina nitida (striate variety) MILLETT, Journ. Roy. Micr. Soc., 1898, p. 266.

Spiroloculina grata, var. *angulata* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 36, pl. 7, fig. 5.

This variety differs from the typical form in having the chambers angular instead of circular in transverse section, and the costae parallel to the peripheral margin of the chamber instead of oblique, in extreme forms with the periphery carinate.

The type specimen of the variety is from Cebu, Philippines. It was noted at two other stations—D5160, in 12 fathoms (22 meters), Tawi Tawi Group, Sulu Archipelago, and D5218, in 20 fathoms (37 meters), between Burias and Luzon.

It is widely distributed and with the typical is found in shallow, tropical waters.

Spiroloculina artillarum, var. *angulata*.—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
10994	U.S.N.M.	1	D5160...	° ' " ° ' "	12	°F.		
10995	U.S.N.M.	2	D5218...	5 12 40 N.; 119 55 10 E... 13 11 15 N.; 123 02 45 E...	20	s..... crs. s.....	Few. Few.

SPIROLOCULINA (?) CONVEXIUSCULA H. B. Brady.

Plate 82, figs. 4a, b.

Spiroloculina (?) *convexiuscula* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 155, pl. 10, figs. 18-20.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 266.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 6.

Description.—Test small, compressed, nearly circular in side view, biconvex; early chambers obscured, the later chambers overlapping nearly to the center; chambers ornamented by longitudinal, rounded, more or less interrupted costae; peripheral margin carinate; aperture with a definite lip, rounded or elliptical.

Length usually less than 1 mm.

Brady's material was from two stations—off Raine Island, Torres Strait, 155 fathoms (283 meters), and off the Admiralty Islands, 16 to 25 fathoms (29 to 46 meters). Millett records it from the Malay Archipelago, "common and widespread over area 1 (eastern part), but rare in area 2 (west of Borneo)." Sidebottom mentions "a single specimen, which, appears to be typical," from 465 fathoms (850 meters), east coast of Australia.

Millett remarks as follows on this species:

Assigned by Brady, with some hesitation, to this genus, it is more probably an arrested form of *Articulina*. It is remarkable for its uniformity, specimens from all localities being almost identical in size and structure.

The aperture and embracing characters would seem to place this species with *Vertebralina*, but other characters suggest that it be left for the present under *Spiroloculina*.

In the Philippine material it has occurred at eight stations, common at two of them. The depths range from 10 to 393 fathoms (18 to 719 meters), with the average 76 fathoms (139 meters). The two bottom temperatures given are 75.7° F. (24.2° C.) and 43.7° F. (6.5° C.). The stations are: China Sea off southern Luzon; Jolo Jolo, Tawi Tawi Group; off Romblon and off eastern Panay.

It is very evidently a species of warm, shallow waters of the Indo-Pacific region.

Spiroloculina (?) *convexiuscula*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10885	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E...	10	°F.	Co	Few.
10886	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E...	393	43.7	gn. m., s.....	Few.
10887	U.S.N.M.	2	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Few.
10888	U.S.N.M.	1	D5154...	5 14 50 N.; 119 58 45 E...	12	co. s.....	Few.
10889	U.S.N.M.	10+	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Common.
10890	U.S.N.M.	1	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fine. s.....	Rare.
10891	U.S.N.M.	4	D5218...	13 11 15 N.; 123 02 45 E...	20	ers. s.....	Common.
10892	U.S.N.M.	2	Jolo Jolo, P. I.....	Rare.

SPIROLOCULINA ADVENA, new species.

Plate 83, figs. 1a-c.

Description.—Test inequilateral, one side nearly flat or slightly convex, the other deeply excavate in the middle, but with a high raised portion near the middle running lengthwise of the chamber, peripheral portion again flattened; whole test elliptical, the ends somewhat produced, apertural end slightly extended, aperture small, rounded; sutures distinct, wall smooth.

Length about 1 mm.

Distribution.—Type specimen (Cat. No. 10949, U. S. N. M.) from *Albatross* station D5178, off Romblon, in 78 fathoms (143 meters).

This would have been taken for some sort of abnormal specimen if found alone, but it occurs in the same form at this station in other specimens. It was never found elsewhere, however. Its nearest relative seems to be *S. affixa* Terquem.

Spiroloculina advena—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10949	U.S.N.M.	2	D5178...	12 43 00 N.; 122 06 15 E...	78	°F.	fine. s.....	Few.

SPIROLOCULINA AFFIXA Terquem.

Plate 83, figs. 2a-c.

Spiroloculina affixa TERQUEM, Mém. Soc. géol. France, ser. 3, vol. 1, 1878, p. 55, pl. 5 (10), figs. 13a-c.—HOWCHIN, Trans. Roy. Soc. South Australia, vol. 12, 1889, p. 2.—CHAPMAN, Journ. Linn. Soc. Zoology, vol. 30, 1907, p. 16, pl. 1, figs. 23-25.

Spiroloculina inaequilateralis SCHLUMBERGER, Mém. Soc. Zool. France, vol. 6, 1893, p. 201, pl. 4, figs. 84-86; fig. 3 (in text).—SIDEBOTTOM, Mem. Proc. Manchester Lit. and Philos. Soc., vol. 54, No. 16, 1910, p. 2, pl. 1, fig. 2.

Spiroloculina acutimargo H. B. BRADY (in part), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, pl. 10, fig. 12 (not figs. 13-15).

Description.—Test elongate, fusiform, the sides very different from one another, one side flattened, the other deeply concave; chambers with definite sutures which are flush with the surface; wall thin and smooth; peripheral face somewhat concave, peripheral margin somewhat carinate, apertural end subacute; aperture elliptical, compressed.

Length from 0.5 to 0.75 mm.

The distribution of this species includes two areas so far as is known—the Mediterranean, represented by the types which Terquem described, from the Pliocene of the Isle of Rhodes, and the recent specimens from the Gulf of Marseilles (Schlumberger) and the Bay of Palermo, Sicily (Sidebottom); the other the Pacific region. Here Howchin and Chapman have found the species in the Tertiary of Australia, and I have it in the recent material from the Philippines and from off New Zealand.

The Philippine specimens are from four *Albatross* stations off northern Cebu, 32 fathoms (59 meters); off southeastern Mindoro, 145 fathoms (265 meters); and China Sea, off Formosa, 148 and 340 fathoms (271 and 622 meters). The most typical specimens are those off New Zealand.

Spiroloculina affixa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10893	U.S.N.M.	1	D5192...	° ' " ° ' "	32	° F.	gn. s.	Rare.
10894	U.S.N.M.	1	D5261...	11 09 15 N.; 123 50 00 E..	145	s., m.....	Rare.
10895	U.S.N.M.	2	D5315...	12 30 55 N.; 121 34 24 E..	148	54.4	s., sh.....	Rare.
10896	U.S.N.M.	1	D5318...	21 40 00 N.; 116 58 00 E..	340	s., br. Co....	Rare.
				21 32 00 N.; 117 46 00 E..				

SPIROLOCULINA ARENARIA H. B. Brady.

Plate 81, figs. 3a, b.

Spiroloculina arenaria H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 153, pl. 8, fig. 12.—EGGER, Abh. kön. Bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 226, pl. 1, fig. 29.—DAKIN, Ceylon Pearl-Oyster Fisheries, Suppl. Rept., pt. 5, 1906.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 36.

One of Brady's *Challenger* stations for this species was in the Philippines in 95 fathoms (174 meters). The others are off Kandavu, Fiji Islands, 210 fathoms (384 meters), and off Raine Island, Torres Strait, 155 fathoms (283 meters). Dakin records it off Ceylon. Egger's records are off Mauritius and off western Australia.

In the Philippines species have occurred at five stations ranging in depth from 80 to 248 fathoms (146 to 454 meters), bottom tem-

peratures from 47.4° F. to 59° F. (8.5° C. to 15° C.). The localities are: China Sea, off southern Luzon; off western Bohol; and east of Masbate.

This is another species known only from the Philippines and the Indo-Pacific.

Spiroloculina arenaria—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10996	U.S.N.M.	10+	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m. . . .	Common.
			D5192...	9 40 50 N.; 123 39 45 E..	220	53.9	gy. m.	Rare.
			D5213...	12 15 00 N.; 123 57 30 E..	80	s. m., sh. . . .	Few.
			D5218...	13 52 45 N.; 120 25 00 E..	201	50.4	dk. gy. s. . . .	Rare.
			D5282...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. gy. s. . . .	Few.

Genus PLANISPIRINA Seguenza, 1880.

PLANISPIRINA CONTRARIA (d'Orbigny).

Plate 85, figs. 1a, b.

Biloculina contraria D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 266, pl. 16, figs. 4-6.

Planispirina contraria H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 195, pl. 11, figs. 10, 11, woodcut, figs. 5, a.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 611, pl. 13, fig. 12.

This species was not found in the material from the North Pacific which I examined, yet it has been found in the Philippine material at 20 stations and in considerable numbers at some of them. The specimens are large and well developed.

The localities represented are Pacific Ocean, east and north coasts of Mindanao; off northwestern Panay; east and west coasts of Luzon; between Negros and Siquijor; between Jolo and Tawi Tawi. To the southward it has occurred in Darvel Bay and Sibuko Bay, Borneo; south of Patiente Strait and the Gulf of Boni.

Depths of the stations range from 243 to 890 fathoms (445 to 1,628 meters), the average 445 fathoms (814 meters). The bottom temperatures of 11 stations range from 38.3° F. to 53.5° F. (3.5° C. to 11.8° C.) with the average 45.7° F. (7.6° C.).

The distribution of this species is peculiar. Millett had but a single specimen in the Malay Archipelago, and it does not seem to extend into the southern or eastern Pacific to any extent from the records. Brady records it from *Challenger* material, north of New Guinea, 1,075 fathoms (1,966 meters), and off Amboina, 1,425 fathoms (2,607 meters). The record from off Chile is in doubt, and the others are from the North Atlantic.

Planispirina contraria—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10768	U.S.N.M.	3	D5115...	13 37 11 N.; 120 43 40 E...	340	° F.	Few.
10769	U.S.N.M.	1	D5136...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s.....	Few.
10770	U.S.N.M.	6	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.....	Common.
10771	U.S.N.M.	3	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Few.
10772	U.S.N.M.	5	D5460...	13 32 30 N.; 123 58 06 E...	565	gy. m.....	Common.
10773	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E...	509	gy. m.....	Rare.
10774	U.S.N.M.	5	D5467...	13 35 27 N.; 123 37 18 E...	480	gy. m.....	Common.
10775	U.S.N.M.	3	D5469...	13 36 48 N.; 123 38 24 E...	509	gn. m.....	Few.
10776	U.S.N.M.	1	D5470...	13 37 30 N.; 123 41 09 E...	560	m.....	Few.
10777	U.S.N.M.	1	D5523...	8 48 44 N.; 123 27 35 E...	Few.
10778	U.S.N.M.	3	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.....	Few.
10779	U.S.N.M.	1	D5565...	5 51 42 N.; 120 30 30 E...	243	52.3	s., ptr. sh.....	Few.
10780	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.....	Rare.
10781	U.S.N.M.	4	D5582...	4 19 54 N.; 118 58 39 E...	890	38.3	gy. m., fine. s.....	Common.
10782	U.S.N.M.	1	D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fine. gy. s.....	Few.
10783	U.S.N.M.	1	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.....	Rare.
10784	U.S.N.M.	1	D5591...	4 11 48 N.; 118 38 20 E...	260	Rare.
10785	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m.....	Rare.
10786	U.S.N.M.	1	D5654...	3 42 00 S.; 120 45 50 E...	805	38.3	Rare.
10787	U.S.N.M.	1	D5658...	3 32 40 S.; 120 31 30 E...	510	41.2	gy. m.....	Rare.

PLANISPIRINA cf. STRIATA Sidebottom.

Planispirina striata SIDEBOTTOM, Mem. Proc. Manchester Lit. and Philos. Soc., vol. 48, No. 5, 1904, p. 21, pl. 5, figs. 12-14, text figure 9.

Material resembling this species described by Sidebottom from off Delos in the Grecian Archipelago occurred at *Albatross* station D5218, between Burias and Luzon, in 20 fathoms (37 meters).

Planispirina cf. striata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10803	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E...	20	° F.	crs. s.....	Rare.

PLANISPIRINA SPHAERA (d'Orbigny).

Biloculina sphaera D'ORBIGNY, Voy. Amér. Mérid., 1839, "Foraminifères," p. 66, pl. 8, figs. 13-16.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 141, pl. 2, figs. 4a, b.

Planispirina sphaera SCHLUMBERGER, Mém. Soc. Zool. France, 1891, p. 190, text figs. 45, 46.—CHAPMAN, Journ. Linn. Soc. Zoology, vol. 30, 1910, p. 399.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 37, pl. 19, fig. 1.

This is a widely-distributed species especially found in deep water. Specimens were rare at three stations—D5428, off eastern Palawan, in 1,105 fathoms (2,021 meters); D5470, east coast of Luzon, 560 fathoms (1,024 meters); and D5668, Macassar Strait, 901 fathoms (1,648 meters). Chapman records it from off Funafuti in 1,050 fathoms (1,920 meters).

Planispirina sphaera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13589	U.S.N.M.	1	D5428...	9 13 00 N.; 118 51 15 E...	1,105	° F. 49.7	gy. m.	Rare.
13588	U.S.N.M.	1	D5470...	13 37 30 N.; 123 41 09 E...	560	m.	Rare.
13590	U.S.N.M.	1	D5668...	2 28 15 S.; 118 49 00 E...	901	38.2	gy. m.	Rare.

Genus VERTEBRALINA d'Orbigny, 1826.

VERTEBRALINA STRIATA d'Orbigny.

Vertebralina striata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 283, No. 1; Modèles, 1826, No. 81.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 187, pl. 12, figs. 14–16.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 38, pl. 22, figs. 3, 4.

This is a species characteristic of coral-reef conditions. Millett did not find it well developed in his material from the Malay Archipelago, but I found it fairly common in material from about the Hawaiian Islands.

The only Philippine record is a single specimen in shore material from Tataan Island. It is typical.

Vertebralina striata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13581	U.S.N.M.	1	Tataan Island, P. I.	° F.	Rare.

VERTEBRALINA STRIATA d'Orbigny, var. RETICULOSA, new variety.

Plate 97, fig. 4.

Description.—Variety differing from the typical in the ornamentation of the last few chambers, which, instead of being “the invariable surface ornament of delicate parallel striae,” as in the typical, has certain of the lines stronger than others in usually alternating areas, giving a very peculiar reticulose appearance.

This variety occurs in the Mediterranean, where it is noted by Sidebottom from Delos “* * * some of them boldly striated and pitted as in *V. insignis*.”³⁷ I am grateful to Mr. Sidebottom for Delos material in which this variety occurs. Heron-Allen and Earland³⁸ also note under *V. striata*, “at Stns. 10 and ? B specimens with a reticulate instead of a striate surface were observed.” The variety, therefore, has a wide distribution in warm waters.

The type (U.S.N.M. No. 13591) is from *Albatross* station D5160, Sulu Archipelago, Tawi Tawi Group, 12 fathoms (22 meters). This is figured here and, while not so definitely reticulose as the Mediterranean specimens, has carried the peculiar ornamentation still further in its development.

³⁷ Mem. Proc. Manchester Lit. and Philos. Soc., vol. 48, 1904, No. 5, p. 18.

³⁸ Trans. Zool. Soc. London, vol. 20, 1915, p. 587.

Vertebralina striata, var. *reticulosa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13591	U.S.N.M.	1	D5160...	5 12 40 N.; 119 55 10 E...	12	° F.	s.....	Rare.

Genus *NODOBACULARIA* Rhumbler, 1895.*NODOBACULARIA TIBIA* Jones and Parker.

Plate 85, figs. 2, 3.

Nodobacularia tibia JONES and PARKER, Quart. Journ. Geol. Soc., vol. 16, 1860, p. 455, pl. 20, figs. 48–51.—H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 52, pl. 8, figs. 1, 2; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 135, pl. 1, figs. 1–4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 39, pl. 8, figs. 1, 2.

There are two sorts of specimens in the material, one of these evidently the megalospheric form, in which the proloculum is large and followed by a long, straight chamber, sometimes constricted ones. The other is evidently the microspheric form, the proloculum and initial end usually wanting, and represented by a series of gradually enlarging subcylindrical chambers.

Brady records this species from 95 fathoms (174 meters) from the Philippines in the *Challenger* material.

The only material I have had has come from the following stations: D5201, Sogod Bay, southern Leyte, in 554 fathoms (1,012 meters), and D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (1,086 meters).

Nodobacularia tibia—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	
			D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	

Subfamily *QUINQUELOCULINÆ*.Genus *QUINQUELOCULINA* d'Orbigny, 1826.

This genus contains probably more specimens of various species than any other in the entire Philippine foraminifera. The species are numerous and many of them very definite, both in their characters and their geographical distribution.

They seem to fall into three different groups: First, those which occur in the deeper water and which are found in other regions and represent widely distributed species. These are usually few in number, as far as specimens occur, and are widely scattered in the region where sufficient depth is obtained, especially on the ocean side of the islands, and in the region southward where the *Albatross* dredged. The

second group of species is that represented by Indo-Pacific species, which occur characteristically in the South Pacific, about Australia, northward to southern Japan, westward to the Hawaiian Islands, and usually eastward through the Indian Ocean to the coast of Africa. Such species are often widely distributed in shallow water, often occurring in great numbers. The third group are those which seem to be characteristic of this region, and may either represent local species or more widely spread species which have not been recorded elsewhere. These include some of the largest and most characteristic species of the region.

QUINQUELOCULINA SEMINULUM (Linnaeus).

Plate 88, figs. 4a-c.

Serpula seminulum LINNAEUS, Syst. Nat., ed. 12, 1767, p. 1264, No. 791.

Quinqueloculina seminulum D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 303, No. 44.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 44, pl. 11, fig. 2, fig. 29 (in text).

Miliolina seminulum WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 85, pl. 7, figs. 183-185.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 157, pl. 5, figs. 6a-c.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, 1915, vol. 20, p. 569, pl. 42, fig. 31.


There are numerous specimens which can be referred to this species, but they are not common at any station, and are found, as a rule, in the region of the Sulu Sea.

Specimens have occurred at 13 stations, ranging in depth from 18 to 159 fathoms (35 to 291 meters), the average depth being 40 fathoms (73 meters). These localities represent the following: China Sea, off southern Luzon; Sulu Sea, off western Mindanao; near Basilian Island; off Jolo; and off Tawi Tawi.

It has also occurred rarely off northern Cebu and in Buton Strait.

Specimens are referred to this species from almost all parts of the world, and, being

smooth and with few distinguishing characters, many species or varieties have been placed under this name. Heron-Allen and Earland record it from the Kerimba Archipelago, where, according to them, it shows a great amount of variation. The largest specimens from the Philippine region, when fully grown, measure



FIGS. 19, 20.—QUINQUELOCULINA SEMINULUM (LINNAEUS). $\times 20$. FROM ALBATROSS STATION D5152. 19, SPECIMEN WITH THE TIP OF THE TOOTH BECOMING BIFID. 20, SPECIMEN WITH THE TOOTH FORKED AND MUCH BROADENED TOWARD THE TIP.

3 mm. in length, are smooth, with a glossy surface, and are slightly angled.

Quinqueloculina seminulum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13857	U.S.N.M.	1	D5143...	13 51 30 N.; 120 50 30 E...	159 ° F.	dk. gn. m. ..	Few.
13859	U.S.N.M.	1	D5132...	Island off Panabutan Point, N. 15°; W., 0.3 mile.	36	gn. m., s....	Rare.
13858	U.S.N.M.	3	D5133...	Island off Panabutan Point, N. 52°; E., 1.5 miles!	38	79.5	gn. m., s....	Few.
13860	U.S.N.M.	3	D5134...	6 44 45 N.; 121 48 00 E...	25	fne. s.....	Few.
13861	U.S.N.M.	1	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.....	Rare.
13862	U.S.N.M.	4	D5141...	6 09 00 N.; 120 58 00 E...	29	co. s.....	Few.
13863	U.S.N.M.	1	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Rare.
13864	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Rare.
13865	U.S.N.M.	1	D5164...	5 01 40 N.; 119 52 20 E...	18	gn. m.....	Rare.
13866	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Rare.
13867	U.S.N.M.	4	D5642...	4 31 40 S.; 122 49 42 E...	37	gy. m.....	Rare.
11631	U.S.N.M.	1	Cebu.....	Rare.
12881	U.S.N.M.	1	Cebu.....	Rare.

QUINQUELOCULINA VULGARIS d'Orbigny.

Plate 87, figs. 1a-c.

Quinqueloculina vulgaris D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 302, No. 33.—SCHLUMBERGER, Mém. Soc. Zool. France, 1893, p. 207, text figs. 13, 14, pl. 2, figs. 65, 66.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 46, pl. 11, fig. 3.

Miliolina vulgaris CHAPMAN, Trans. New Zealand Inst., vol. 38, 1905, p. 81.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 569.

This species, which is now known from this general region, from the Hawaiian Islands, south to New Zealand, and westward to the Kerimba Archipelago off southeastern Africa, has been found to be a common species in the Philippine region. Specimens have occurred at 35 stations, ranging in depth from 10 to 805 fathoms (18 to 1,472 meters), the average depth being 236 fathoms (432 meters). Bottom temperatures are given at 18 stations, ranging from 43.7° F. to 79.5° F. (6.5° C. to 26.3° C.), the average depth being 53.9° F. (12.1° C.). There does not seem to be any particular depth at which it is particularly abundant.




FIG. 21.—QUINQUELOCULINA VULGARIS D'ORBIGNY. X 50. FROM ALBATROSS STATION D5216. OPPOSITE SIDES OF SAME SPECIMEN.

There is no great variation in its general characters, and on the whole it forms a fairly well-defined species in the material. Most of the stations are in the Archipelago itself, but there are a few in the

deeper water in the regions to the south. It has been rare at the stations in the southern area where it is found.

The localities in the Archipelago are as follows: Tara Island; China Sea, off southern Luzon; Balayan Bay; Sulu Sea, off western Mindanao; off eastern Panay; between Panay and Negros; off northern Cebu; Sogod Bay, southern Leyte; off western Samar; east of Masbate; between Bohol and Leyte; Pacific Ocean, east coast of Mindanao; off northeastern Mindoro; Verde Island Passage; Malampaya Sound, Palawan Island; Palawan Passage; between Samar and Leyte; off northern Mindanao; between Siquijor and Bohol Islands; north of Tawi Tawi; and to the southward in Darvel Bay, Borneo; Gulf of Tomini, Celebes; and south of Patiente Strait.

Quinqueloculina vulgaris—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13636	U.S.N.M.	2	D5096...	14 20 23 N.; 120 34 15 E...	28	gy. m., s., sh.	Rare.
13640	U.S.N.M.	4	D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s.....	Few.
13641	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E...	10	Co.....	Rare.
13659	U.S.N.M.	1	D5120...	13 45 30 N.; 120 30 15 E...	393	43.7	gn. m., s.....	Rare.
13643	U.S.N.M.	10	D5133...	Island off Panabutan Point, N. 52°; E. 1.5 miles.	38	79.5	gn. m., s.....	Common.
13645	U.S.N.M.	10+	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fine. s.....	Common.
13644	U.S.N.M.	1	D5182...	11 30 40 N.; 123 23 20 E...	24	fine. s., m.....	Rare.
13646	U.S.N.M.	1	D5185...	10 05 45 N.; 122 18 30 E...	638	49.8	gn. m.....	Rare.
13648	U.S.N.M.	6	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Common.
13649	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Rare.
13647	U.S.N.M.	5	D5206...	11 31 40 N.; 124 42 40 E...	32	gn. m.....	Few.
13650	U.S.N.M.	1	D5213...	12 15 00 N.; 123 57 30 E...	80	s., m., sh.....	Rare.
13651	U.S.N.M.	4	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m.....	Few.
13652	U.S.N.M.	1	D5230...	10 01 50 N.; 124 42 30 E...	118	57.6	gy. s.....	Rare.
13655	U.S.N.M.	2	D5236...	8 50 45 N.; 126 56 52 E...	494	41.2	fine. gy. s.....	Rare.
13653	U.S.N.M.	2	D5260...	12 25 35 N.; 121 31 35 E...	234	51.4	gn. m., s.....	Rare.
13637	U.S.N.M.	10+	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Common.
13638	U.S.N.M.	2	D5269...	13 39 50 N.; 120 59 30 E...	220	fine. s., p.....	Rare.
13639	U.S.N.M.	3	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fine. s., m., sh	Few.
13642	U.S.N.M.	5	D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh.....	Few.
13660	U.S.N.M.	1	D5342...	10 56 55 N.; 119 17 24 E...	14-25	gy. m.....	Rare.
13661	U.S.N.M.	1	D5343...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.....	Rare.
13662	U.S.N.M.	4	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh., g.....	Few.
13663	U.S.N.M.	1	D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy. m., fine. s	Rare.
13664	U.S.N.M.	2	D5526...	9 12 45 N.; 123 45 30 E...	805	53.3	gn. m., glob.	Rare.
13665	U.S.N.M.	10+	D5529...	9 23 45 N.; 123 39 30 E...	441	53	gy. m., glob.	Common.
13666	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.....	Rare.
13668	U.S.N.M.	1	D5576...	5 28 30 N.; 120 02 27 E...	315	52.3	Co., s.....	Rare.
13669	U.S.N.M.	2	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.....	Rare.
13667	U.S.N.M.	2	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fine. s., Co...	Rare.
13654	U.S.N.M.	3	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co...	Rare.
13656	U.S.N.M.	1	D5601...	1 13 10 N.; 125 17 05 E...	765	s., glob., ptr.	Rare.
13657	U.S.N.M.	1	D5612...	0 35 00 S.; 121 45 40 E...	750	Rare.
13658	U.S.N.M.	3	D5630...	0 56 30 S.; 128 05 00 E...	569	Co., s., m...	Rare.
13670	U.S.N.M.	2	Tara Island, P. I.....	Rare.

QUINQUELOCULINA LAMARCKIANA d'Orbigny.

Plate 87, figs. 2, 3a-c.


Quinqueloculina lamarckiana D'ORBIGNY, in De La Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 189, pl. 11, figs. 14, 15.

This species seems to be very common in the region, occurring at 81 stations, ranging in depth from 12 to 1,560 fathoms (22 to 2,853 meters), the average being 215 fathoms (393 meters). Bottom temperatures are given at 32 of these stations, ranging from 41.1° to 76.3° F. (5.0° to 24.6° C.), the average 53.8° F. (12.1 C.).

These stations cover the Archipelago rather well, and to the south occur in the following localities: Sibuko Bay, Borneo; Gulf of Tomini, Celebes; Molucca Passage; between Gillolo and Makyan Island; between Gillolo and Kayoa; Molucca Sea; and Buton Strait.

The species may³ be distinguished by the very highly polished angular test, the aperture flush with no² definite neck, and a single, simple tooth.

Quinqueloculina lamarckiana—Material examined.



FIGS. 22, 23.—QUINQUELOCULINA LAMARCKIANA D'ORBIGNY. X 50. FROM ALBATROSS STATION D5220. 22, APERTURAL VIEW OF YOUNG SPECIMEN WITH SIMPLE ELONGATE TOOTH. 23, ADULT SPECIMEN WHICH HAS DEVELOPED A BIFID TOOTH AND A MORE ROUNDED APERTURE.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11030	U.S.N.M.	6	D5097	14 19 15 N.; 120 33 52 E.	30	gy. m., s. sh.	Common.
11031	U.S.N.M.	10	D5100	14 17 15 N.; 120 32 40 E.	35	gy. s.	Common.
11032	U.S.N.M.	2	D5105	14 43 55 N.; 120 12 50 E.	25	co. s.	Few.
11033	U.S.N.M.	1	D5106	14 23 55 N.; 120 32 33 E.	37	gy. m.	Rare.
11034	U.S.N.M.	1	D5121	13 27 20 N.; 121 17 45 E.	108	dk. gn. m.	Rare.
11035	U.S.N.M.	10+	D5132	Island off Panabutan Point, N. 15°, W. 0.30 mile.	26	gn. m., s.	Common.
11036	U.S.N.M.	10+	D5133	Island off Panabutan Point, N. 52°, E. 1.50 miles.	38	gn. m., s.	Common.
11037	U.S.N.M.	3	D5134	6 44 45 N.; 121 48 00 E.	25	fine. s.	Few.
11038	U.S.N.M.	2	D5136	6 04 20 N.; 120 69 20 E.	22	s., sh.	Few.
11039	U.S.N.M.	1	D5141	6 09 00 N.; 120 58 00 E.	29	co. s.	Rare.
11040	U.S.N.M.	5	D5142	6 06 10 N.; 121 02 40 E.	21	co. s., sh.	Common.
11041	U.S.N.M.	2	D5143	6 05 50 N.; 121 02 15 E.	19	co. s.	Common.
11042	U.S.N.M.	1	D5144	6 05 50 N.; 121 02 15 E.	19	co. s., sh.	Rare.
11043	U.S.N.M.	10+	D5145	6 04 30 N.; 120 59 30 E.	23	co. s., sh.	Common.
11046	U.S.N.M.	2	D5151	5 24 40 N.; 120 27 15 E.	24	co. s., sh.	Rare.
11044	U.S.N.M.	1	D5154	5 14 50 N.; 119 53 45 E.	12	s.	Rare.
11045	U.S.N.M.	1	D5160	5 12 40 N.; 119 65 10 E.	12	s.	Rare.
11047	U.S.N.M.	3	D5162	5 10 00 N.; 119 47 30 E.	230	52.9	crs. s., brk. sh.	Rare.
11048	U.S.N.M.	3	D5172	6 93 15 N.; 120 35 30 E.	318	fine. s., sh.	Rare.
11049	U.S.N.M.	10+	D5178	12 43 00 N.; 122 06 15 E.	78	fine. s.	Common.
11050	U.S.N.M.	6	D5197	11 09 15 N.; 123 60 00 E.	32	gn. s.	Common.
11051	U.S.N.M.	2	D5198	9 40 50 N.; 123 39 45 E.	220	53.9	gn. m.	Few.
11052	U.S.N.M.	10+	D5201	10 10 00 N.; 125 04 15 E.	554	52.8	gy. s., m.	Common.
11053	U.S.N.M.	1	D5206	11 31 40 N.; 124 42 40 E.	32	gn. m.	Rare.
11064	U.S.N.M.	6	D5210	11 49 55 N.; 124 28 05 E.	50	76.3	fine. gy. s.	Common.
			D5211	11 51 35 N.; 124 14 00 E.	155	56.6	gn. m., s.	Few.
11055	U.S.N.M.	5	D5212	12 04 15 N.; 124 04 36 E.	108	59.9	gy. s., m.	Few.

Quinqueloculina lamarckiana—Material examined—Continued.

Cat. No.	Call of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
11056	U.S.N.M.	4	D5213	12 15 00 N.; 123 57 30 E.	80		s., m., sh.	Few.
11057	U.S.N.M.	1	D5214	12 25 18 N.; 123 37 15 E.	218	51.4	gn. m.	Few.
11058	U.S.N.M.	10+	D5217	13 20 00 N.; 123 14 15 E.	105	63.1	crs. gy. s.	Common.
11059	U.S.N.M.	10+	D5220	13 38 00 N.; 121 58 00 E.	50		sft. gn. m.	Abundant.
11060	U.S.N.M.	1	D5227	12 53 45 N.; 121 52 30 E.	322		gn. m.	Rare.
11061	U.S.N.M.	1	D5255	7 03 00 N.; 125 39 00 E.	100		sft. m.	Rare.
11062	U.S.N.M.	10+	D5259	11 57 30 N.; 121 42 15 E.	312	49.3	gy. m., glob.	Common.
11063	U.S.N.M.	3	D5261	12 30 55 N.; 121 34 24 E.	145		s., m.	Rare.
11064	U.S.N.M.	2	D5276	13 49 15 N.; 120 14 45 E.	18		sh., p., s.	Rare.
			D5277	13 56 55 N.; 120 13 45 E.	80	58.6	fine. s.	Few.
11065	U.S.N.M.	1	D5301	20 37 00 N.; 115 43 00 E.	208	50.5	gy. m., s.	Rare.
11066	U.S.N.M.	5	D5277	13 56 55 N.; 120 13 45 E.	80	58.6	fine. s.	Few.
11067	U.S.N.M.	10+	D5311	21 33 00 N.; 116 15 00 E.	88		crs. s., sh.	Common.
11068	U.S.N.M.	10+	D5313	21 30 00 N.; 116 43 00 E.	150	53.6	s.	Common.
11069	U.S.N.M.	10+	D5315	21 40 00 N.; 116 58 00 E.	148	54.4	s., sh.	Common.
11070	U.S.N.M.	5	D5338	11 33 45 N.; 119 24 45 E.	43		co. s., sh.	Few.
11071	U.S.N.M.	3	D5342	10 56 55 N.; 119 17 24 E.	14-25		gy. m.	Few.
11072	U.S.N.M.	7	D5348	10 57 45 N.; 118 38 15 E.	375	56.4	co. s.	Few.
11073	U.S.N.M.	10	D5358	6 06 40 N.; 118 18 15 E.	39		m.	Common.
11074	U.S.N.M.	6	D5369	13 48 00 N.; 121 43 00 E.	106		bk. s.	Common.
11075	U.S.N.M.	6	D5370	13 44 15 N.; 121 42 30 E.	159	54.3	sft. m.	Common.
11076	U.S.N.M.	8	D5381	13 14 15 N.; 122 44 45 E.	88		co. s.	Common.
11077	U.S.N.M.	1	D5385	13 24 50 N.; 123 03 70 E.	327	62.4	gy. m.	Rare.
11078	U.S.N.M.	2	D5398	11 35 12 N.; 124 13 48 E.	114		gn. m.	Rare.
11079	U.S.N.M.	1	D5408	10 40 15 N.; 124 15 00 E.	159	55.4	gn. m.	Rare.
11080	U.S.N.M.	1	D5419	9 58 30 N.; 123 46 00 E.	175	54.5	gn. m.	Rare.
11081	U.S.N.M.	1	D5420	9 49 35 N.; 123 45 00 E.	127	59.0		Rare.
11082	U.S.N.M.	3	D5423	9 38 30 N.; 121 11 00 E.	508	49.8	gy. m., co. s.	Few.
11083	U.S.N.M.	2	D5424	9 37 05 N.; 121 12 37 E.	340	50.4	co. s.	Few.
11084	U.S.N.M.	1	D5425	9 37 45 N.; 121 11 00 E.	495	49.4	gy. m., co. s.	Rare.
11085	U.S.N.M.	2	D5428	9 13 00 N.; 118 51 15 E.	1,105	49.7	gy. m.	Rare.
11086	U.S.N.M.	1	D5429	9 41 30 N.; 118 50 22 E.	766		gn. m.	Rare.
11087	U.S.N.M.	4	D5443	12 43 05 N.; 125 01 00 E.	241	51.3	co. s., sh.	Rare.
11088	U.S.N.M.	1	D5446	12 43 51 N.; 124 59 18 E.	300		gn. m.	Rare.
11089	U.S.N.M.	1	D5469	13 36 48 N.; 123 38 24 E.	500		gn. m. (net).	Rare.
11090	U.S.N.M.	2	D5487	10 02 45 N.; 125 05 33 E.	732	52.1	gn. m.	Rare.
11091	U.S.N.M.	1	D5523	8 48 44 N.; 123 27 35 E.				Rare.
11092	U.S.N.M.	1	D5537	9 11 00 N.; 123 23 00 E.	254	53.5	gn. m.	Rare.
11093	U.S.N.M.	2	D5538	9 08 15 N.; 123 23 20 E.	256	53.5	gn. m., s.	Rare.
11094	U.S.N.M.	3	D5543	8 47 15 N.; 123 35 00 E.	162	54.5	s.	Rare.
11095	U.S.N.M.	1	D5565	5 41 42 N.; 120 30 30 E.	243	52.3	s., ptr., sh.	Rare.
11096	U.S.N.M.	1	D5569	5 33 15 N.; 120 15 30 E.	303	52.3	co. s.	Rare.
11097	U.S.N.M.	4	D5585	4 07 00 N.; 118 49 54 E.	476	41.1	gy. m.	Rare.
11098	U.S.N.M.	2	D5586	4 06 50 N.; 118 47 20 E.	347	44.0	gy. m.	Rare.
11099	U.S.N.M.	7	D5590	4 10 50 N.; 118 39 35 E.	310	44.3	gn. m., s.	Few.
11100	U.S.N.M.	4	D5613	0 42 00 S.; 121 44 00 E.	752		gy. m.	Few.
13573	U.S.N.M.	2	D5571	5 30 45 N.; 120 07 57 E.	334	52.3	s.	Rare.
13570	U.S.N.M.	4	D5618	0 37 00 N.; 127 15 00 E.	417		gy. m.	Few.
13571	U.S.N.M.	1	D5621	0 15 00 N.; 127 24 35 E.	298		gy. and bk. s.	Few.
13569	U.S.N.M.	1	D5627	0 06 00 N.; 127 26 00 E.	22		m.	Few.
			D5639	3 54 50 S.; 123 27 20 E.	1,560		gy. m.	Few.
13572	U.S.N.M.	2	D5640	4 27 00 S.; 122 55 40 E.	24		s., brk. sh.	Few.
			H4898	7 43 45 N.; 122 03 45 E.	221		gy. m., glob.	Rare.
13574	U.S.N.M.	6		Tara Island, P. I.				Few.
13575	U.S.N.M.	2		Tayabas Gulf.				Rare.

QUINQUELOCULINA VENUSTA KARRER.

Plate 91, figs. 2a-c.

Quinqueloculina venusta KARRER, Sitz. Akad. Wiss. Wien, vol. 57, 1868, p. 147, pl. 2, fig. 6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 45, pl. 11, fig. 1.
Miliolina venusta H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 162, pl. 5, figs. 5, 7.

This seems to be a rare species in the Philippine region. It has occurred at seven stations, ranging in depth from 21 to 340 fathoms (39 to 622 meters), the average depth being 117 fathoms (201 meters).

Bottom temperatures are given at only three stations, ranging from 52.3° to 59.0° F. (11.2° to 15.0° C.), the average temperature 54.9° F. (12.7° C.).

These stations include the following localities: Sulu Archipelago, off Siasi Island; off northern Cebu; China Sea, off Hongkong; between Cebu and Bohol; between Samar and Leyte; and north of Tawi Tawi.

Quinqueloculina venusta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13911	U.S.N.M.	1	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.....	Rare.
13912	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Rare.
13913	U.S.N.M.	2	D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh.....	Rare.
13917	U.S.N.M.	1	D5313...	21 30 00 N.; 116 43 00 E...	150	53.6	s.....	Rare.
13914	U.S.N.M.	2	D5420...	9 49 35 N.; 123 45 00 E...	127	59.0	Rare.
13915	U.S.N.M.	1	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh., g.....	Rare.
13916	U.S.N.M.	1	D5571...	5 30 45 N.; 120 07 57 E...	340	52.3	s., sh.....	Rare.

QUINQUELOCULINA PROCERA (Goës).

Plate 88, figs. 3a-c.

Miliolina procera GOËS, Bull. Mus. Comp. Zoöl., vol. 29, 1896, p. 82, pl. 7, figs.

7-9.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 6, pl. 1, figs. 11-14.

Quinqueloculina procera CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 45, pl. 19, fig. 2.

At a single station, D5236, Pacific Ocean, off the east coast of Mindanao, in 494 fathoms (903 meters), this species is very abundant and typical. It was not seen elsewhere in the region.

Sidebottom records this form from off the east coast of Australia, in 465 fathoms (850 meters), which is close to the Philippine station in depth.

The original material of Goës was from off the Galapagos Island, and I have recorded it from *Albatross* D3346, off the west coast of the United States. From these scattered stations it would seem to be widely distributed in comparatively deep water in the Pacific Ocean.

Quinqueloculina procera—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13837	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Abundant.

QUINQUELOCULINA APICULA, new species.

Plate 88, figs. 1, 2.

Description.—Test somewhat elongate, with five visible chambers, the initial end of each chamber with a long, rounded, somewhat pointed spine, which may be one-half the length of the chamber; periphery bluntly rounded, surface smooth; apertural end slightly truncate without a definite tooth; aperture lunar.

Length up to 2.2 mm.

Distribution.—Type specimen (Cat. No. 13868, U.S.N.M.) from Albatross station D5236, in 494 fathoms (903 meters), Pacific Ocean, east coast of Mindanao.

This species, with test very peculiar, and long basal spine, which appear often at both ends of the test, is not uncommon at this one station, where so many other peculiar species have been taken.

Quinqueloculina apicula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13868	U.S.N.M.	3	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E...	494	° F. 41.2	fne. gy. s....	Few.

QUINQUELOCULINA LIMBATA d'Orbigny.

Quinqueloculina limbata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 302, No. 20.—

FORNASINI, Mem. Accad. Sci. Bologna, ser. 6, vol. 2, 1905, p. 66, pl. 3, fig. 9.

Miliolina limbata HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 577, pl. 44, figs. 5-8.

D'Orbigny's original specimens are from the Red Sea, and this seems to be a species with Indo-Pacific distribution. Heron-Allen and Earland had it from the Kerimba Archipelago, and it has occurred at two stations, D5313, China Sea off Hongkong, 150 fathoms (275 meters), and D5369, off Marinduque Island, 106 fathoms (194 meters), in the Philippine material.

The specimens have the highly polished surface characteristic of this species.

Quinqueloculina limbata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13839	U.S.N.M.	10+	D5313...	° ' " ° ' " 21 30 00 N.; 116 43 00 E..	150	° F. 53.6	s.....	Common.
13840	U.S.N.M.	2	D5369...	° ' " ° ' " 13 48 00 N.; 121 43 00 E..	106	bk. s.....	Rare.

QUINQUELOCULINA CRASSA d'Orbigny, var. SUBCUNEATA, new variety.

Plate 89, figs. 4a-c.

Miliolina crassa HERON-ALLEN and EARLAND (part) (not d'Orbigny), Trans. Zool. Soc. London, vol. 20, 1915, p. 572, pl. 42, fig. 41 (not 37-40).

Description.—Test like the typical in ornamentation, but short and the chambers wedge-shaped, almost sharp at the peripheral angles.

Distribution.—Type specimen (Cat. No. 13847, U.S.N.M.) from Albatross station D5179, off Romblon.

This is evidently the variety figured by Heron-Allen and Earland as "small acute-edged variety," as above noted. It is found at several stations. All the Philippine material which can be referred to this species is of this variety.

The material has occurred at 13 stations, ranging in depth from 12 to 221 fathoms (22 to 405 meters), the average being 62 fathoms (114 meters). The bottom temperature was given but for a single station, 75.7° F. (24.2° C.), where the depth was 37 fathoms (68 meters).

The localities are as follows: Vicinity of Romblon; Sulu Archipelago; Tawi Tawi Group; off Jolo; off western Mindanao; Verde Island Passage; off western Bohol; and between Burias and Luzon.

Quinqueloculina crassa, var. *subcuneata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13841	U.S.N.M.	3	D5138...	6 06 00 N.; 120 58 50 E..	19	s., Co.....	Few.
13842	U.S.N.M.	1	D5142...	6 06 10 N.; 121 02 40 E..	21	co. s., sh.....	Rare.
13843	U.S.N.M.	1	D5144...	6 05 50 N.; 121 02 15 E..	19	Rare.
13844	U.S.N.M.	1	D5151...	5 24 40 N.; 120 27 15 E..	24	co. s., sh.....	Rare.
13845	U.S.N.M.	1	D5158...	5 12 00 N.; 119 54 30 E..	12	crs. s., sh.....	Rare.
13846	U.S.N.M.	2	D5178...	12 43 00 N.; 122 06 15 E..	78	fine s.....	Rare.
13847	U.S.N.M.	8	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Common.
13849	U.S.N.M.	1	D5250...	9 31 50 N.; 124 02 05 E..	Rare.
13848	U.S.N.M.	3	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Rare.
13850	U.S.N.M.	2	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Rare.
13851	U.S.N.M.	3	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., h.....	Rare.
13852	U.S.N.M.	1	D5353...	6 06 40 N.; 118 18 15 E..	39	m.....	Rare.
13853	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob..	Rare.

QUINQUELOCULINA RECTA, new species.

Plate 84, figs. 5a-c.

Description.—Test elongate, three to four times as long as wide surface ornamented by a few strong costae, rounded, peripheral face, and whole chamber rounded in section, the periphery of the last-formed chamber in side view forming a straight line for a large part of its length; apertural end but slightly extended, no neck being

formed; the area slightly back from the aperture smooth, the aperture itself with a slightly everted lip, tooth simple.

Length 1 to 1.25 mm.

Distribution.—Type specimen (Cat. No. 13836, U.S.N.M.) from *Albatross* station D5570, north of Tawi Tawi, 330 fathoms (604 meters). It also occurs off Romblon and off Jolo.

These stations are all, except the type station, in comparatively shallow water, ranging in depth from 19 to 78 fathoms (35 to 143 meters). Bottom temperature in shallow water, 75.7° F. (24.2° C.); in deeper water, 52.3° F. (11.2° C.).

Quinqueloculina recta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13832	U.S.N.M.	1	D5143...	6 05 50 N.; 121 02 15 E..	19	co. s.	Rare.
13834	U.S.N.M.	1	D5151...	5 24 40 N.; 120 27 15 E..	24	co. s., sh.	Rare.
13835	U.S.N.M.	3	D5178...	12 43 00 N.; 122 06 15 E..	78	fine s.	Few.
13833	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.	Rare.
13836	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E..	330	52.3	fine s., glob.	Rare.

QUINQUELOCULINA DISPARILIS d'Orbigny.

Plate 86, figs. 1a-c.

Quinqueloculina disparilis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 302, No. 21.—SCHLUMBERGER, Mém. Soc. Zool. France, vol. 6, 1893, p. 212, pl. 2, figs. 55-57, figs. 21, 22 (in text).

Mioliolina disparilis WIESNER, Archiv. Prot., vol. 25, 1912, p. 216.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 577.

In its typical form this is an elongate test, with the chambers having the outer peripheral face broadly rounded, but flattened and ornamented with a number of fine longitudinal costae, closely set, the sides meeting the peripheral face nearly at right angles, making a definite angle at either side; the aperture with a simple and slightly bifid tooth. This species in the Philippine regions reaches a length of 1.5 mm. or more. The specimen I figured from the North Pacific³⁹ does not belong to this species. In the Philippines very typical specimens occurred at nine stations, ranging in depth from 27 to 303 fathoms (50 to 554 meters), the average depth being 133 fathoms (243 meters). Bottom temperatures are given for three stations, ranging from 52.3° F. to 55.8° F. (11.2° to 13.2° C.), with the average temperature 53.8° F. (12.1° C.).

This species includes the following localities: Off Romblon; between Marinduque and Luzon; Jolo Sea; eastern Palawan; north of Tawi Tawi; and to the south, one station in Darvel Bay, Borneo.

³⁹ Bull. 71, U. S. Nat. Mus., pt. 6, 1917, pl. 14, fig. 1.

At the first of these stations, in 78 fathoms (143 meters), the species was common, and occurred in few numbers at the others. D'Orbigny's type specimens came from the Mediterranean, as did those of Schlumberger and Wiesner. Heron-Allen and Earland record it from the Kerimba Archipelago, and our specimens extend its range considerably.

Quinqueloculina disparilis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
13713	U.S.N.M.	1	D5175...	9 21 00 N.; 121 37 45 E..	° F.	Rare.
13714	} U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E..	78	fine s.	Common.
13633								
13715	U.S.N.M.	1	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.	Rare.
13716	U.S.N.M.	4	D5358...	6 06 40 N.; 118 18 15 E..	39	m.	Few.
13717	U.S.N.M.	1	D5426...	9 12 00 N.; 118 28 00 E..	27	fine gy. s.	Rare.
13718	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E..	303	52.3	co. s.	Raw.
13719	U.S.N.M.	3	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.	Raw.
13720	U.S.N.M.	1	D5580...	4 52 45 N.; 119 06 45 E..	162	55.8	br. s., Co.	Rare.
13721	U.S.N.M.	8	D5642...	4 31 40 S.; 122 42 36 E..	215	gn. m.	Common.

QUINQUELOCULINA BOUEANA d'Orbigny.

Plate 89, fig. 5.

Quinqueloculina boucana D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 293, pl. 19, figs. 7-9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 50, pl. 15, fig. 2.

Miliolina boueana H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 173, pl. 7, figs. 13a-c.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 581.

A very few, mostly triloculine, specimens are placed under this name for want of a better one, as the material is very meager. Heron-Allen and Earland record the species from the Kerimba Archipelago. The Philippine stations are D5148, Sulu Archipelago, off Siasi, 17 fathoms (31 meters); D5181, off eastern Panay, 26 fathoms (48 meters); and D5358, Jolo Sea, 39 fathoms (71 meters).

Quinqueloculina boueana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13630	U.S.N.M.	1	D5148...	5 35 40 N.; 120 47 30 E..	17	co. s.	Rare.
13631	U.S.N.M.	1	D5181...	11 36 40 N.; 123 26 35 E..	26	m., fine s.	Rare.
13632	U.S.N.M.	1	D5358...	6 06 40 N.; 118 18 15 E..	39	m.	Rare.

QUINQUELOCULINA BICORNIS (Walker and Jacob), var. ELEGANS Williamson.

Miliolina bicornis, var. *elegans* WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 88, pl. 7, fig. 195.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 171, pl. 6, figs. 9-11, 12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 48.

Numerous specimens, with the slender neck and more slender form characteristic of this variety, were found at *Albatross* station D5338,

Palawan Passage, in 43 fathoms (78 meters). This variety has been noted from the Hawaiian Islands.

Quinqueloculina bicornis var. *elegans*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13833	U.S.N.M.	7	D5338...	° ' " ° ' " 11 33 45 N.; 119 24 45 E..	43	° F.	co. s., m....	Common.

QUINQUELOCULINA CURTA Cushman.

Plate 100, figs. 1, 2.

Miliolina cuvieriana MILLETT (not d'Orbigny), Journ. Roy. Micr. Soc., 1898, p. 505, pl. 12, figs. 2a, b.

Quinqueloculina disparilis D'ORBIGNY, var. *curta* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 49, pl. 14, fig. 2; fig. 30 (in text).

This species, figured by Millett from the Malay Archipelago and which I described as a variety of *Q. disparilis* from southern Japan, is very common, often abundant, in the Philippine region, where it reaches a large size. The test is characterized by broad peripheral angles, which are marked by several coarse costae. The sides of the chambers are either unornamented or with fine wrinkle-like markings. The younger specimens, as is usual in many species, are shorter than the adult, and later become somewhat longer than broad. The test is a dull cream-white color. The aperture has a thickened wall, and the tooth development is very large and complex. The stages in development of this complex aperture are very interesting.

Length up to 3 mm.

The megalospheric form has a large proloculum and a second cornuspiralike chamber, which is ornamented with several costae, as shown in the figure. From a study of d'Orbigny's original description and figures, together with the recent material which I obtained in the shallow water of the West Indies, this possibly may be *Q. cuvieriana*. It is a species with sharp angles, often carinate, with slight, partially developed, costae at either side. D'Orbigny's figure of this species is very accurate when compared with shallow-water West Indian material, and evidently represents a definite species in that region. The material figured by Millett under this name is undoubtedly the same as that which I had from southern Japan, and which is now found to be abundant in the Philippine region. The species is characteristic of shallow water; specimens become few and rare as depth greater than 100 fathoms is reached. All the stations of this species are in the Archipelago except one. These include the following localities: China Sea, off southern Luzon;

Sulu Sea, off southern Panay; near Basilan Island; off Jolo; off Tawi Tawi; off Romblon; off eastern Panay; off northern Cebu; off western Samar; east of Masbate; off southeastern Mindoro; Palawan Passage; Malampaya Sound, Palawan Island; Ragay Gulf, Luzon; between Cebu and Bohol; and one station to the south in Darvel Bay, Borneo. There are 35 stations at which material is found. These range in depth from 18 to 400 fathoms (33 to 732 meters), the average depth 95 fathoms (174 meters). The bottom temperature is given at nine stations, ranging from 49.6° to 76.3°F. (19.8° to 24.6° C.), the average temperature being 55.3° F. (12.9° C.). As these are from some of the deeper stations, the average temperature for the entire number would probably be considerably higher. This is, then, a species of comparatively shallow, warm waters of the tropics, and should be found to be common in the warm area to the south under similar conditions. At some of the stations, especially in the China Sea off southern Luzon, and in parts of the Sulu region, it occurs very abundantly, making up a considerable proportion of the bottom samples.

Quinqueloculina curta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10997	U.S.N.M.	10+	D5096...	14 20 23 N.; 120 34 15 E...	28	° F.	gn. m., sh., s.	Common.
10998	U.S.N.M.	10+	D5097...	14 19 15 N.; 120 33 52 E...	30	gy. m., sh., s.	Abundant.
10999	U.S.N.M.	10+	D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s.....	Abundant.
11000	U.S.N.M.	8	D5105...	14 43 55 N.; 120 12 50 E...	25	Common.
11001	U.S.N.M.	10+	D5132...	Island off Panabutan Point, N. 15°; W., 0.30 mile.	26	gn. m., s....	Abundant.
11002	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N. 52°; E., 1.50 miles.	38	gn. m., s....	Abundant.
11003	U.S.N.M.	10+	D5134...	6 44 45 N.; 121 48 00 E...	25	fne. s.....	Abundant.
11004	U.S.N.M.	10+	D5143...	6 05 50 N.; 121 02 15 E...	19	co. s.....	Common.
11005	U.S.N.M.	10+	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Common.
11006	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E...	78	fne. s.....	Common.
11007	U.S.N.M.	6	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fne. s....	Common.
11008	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Common.
11009	U.S.N.M.	3	D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fne. gy. s....	Few.
11010	U.S.N.M.	4	D5213...	12 15 00 N.; 123 57 30 E...	80	s., m., sh....	Few.
11011	U.S.N.M.	7	D5261...	12 30 55 N.; 121 34 24 E...	145	s., m.....	Common.
11012	U.S.N.M.	10+	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s....	Common.
11013	U.S.N.M.	7	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fne. s.....	Common.
11014	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E...	201	50.4	dk. gy. s....	Rare.
11015	U.S.N.M.	3	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Few.
11016	U.S.N.M.	6	D5336...	11 37 45 N.; 119 46 00 E...	46	s., m.....	Common.
11017	U.S.N.M.	10+	D5338...	11 33 45 N.; 119 24 45 E...	43	co. s., m....	Common.
11018	U.S.N.M.	10+	D5339...	11 22 00 N.; 119 12 00 E...	52	m.....	Common.
11019	U.S.N.M.	3	D5342...	10 55 55 N.; 119 17 24 E...	14-25	gy. m.....	Few.
11020	U.S.N.M.	3	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.....	Few.
11021	U.S.N.M.	3	D5358...	6 06 40 N.; 118 18 15 E...	39	m.....	Few.
11022	U.S.N.M.	10+	D5369...	13 48 00 N.; 121 43 00 E...	106	bk. s.....	Few.
11023	U.S.N.M.	1	D5377...	13 26 00 N.; 122 19 00 E...	400	49.6	sft. gn. m....	Rare.
11024	U.S.N.M.	6	D5381...	13 14 15 N.; 122 44 25 E...	88	co. s.....	Few.
11025	U.S.N.M.	2	D5382...	13 15 20 N.; 122 45 30 E...	128	m.....	Few.
11026	U.S.N.M.	1	D5419...	9 58 30 N.; 123 46 00 E...	175	54.4	gy. m., s....	Few.
11027	U.S.N.M.	10+	D5426...	9 12 00 N.; 118 28 00 E...	27	fne. gy. s....	Common.
11028	U.S.N.M.	1	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fne. s., glob..	Rare.
11029	U.S.N.M.	2	D5579...	4 54 15 N.; 119 09 52 E...	175	55.3	fne. s., Co...	Rare.

QUINQUELOCULINA BICARINATA d'Orbigny.

Plate 86, figs. 2, 3; plate 100, fig. 7.

Quinqueloculina bicarinata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 302, No. 35.—
TERQUEM, Mém. Soc. géol. France, ser. 3, vol. 1, 1878, pl. vii (xii), figs. 10a-c.

In the Philippine material there is a very widely distributed species which is characterized by two somewhat rounded carinae at the outer border of each chamber. These are apparently identical with the *Q. bicarinata* d'Orbigny, as figured by Terquem. It forms one of the characteristic species both in the Archipelago and southward in shallow water. It is always short and broad, the central part especially bicarinate, but the carinae often coalescing toward either end. It is occasionally slightly striate, but it is not the same as *Q. cuvieriana*. It was recorded at 41 stations in the area, ranging in depth from 10 to 1,262 fathoms (18 to 2,309 meters), the average depth 104 fathoms (191 meters). Bottom temperatures are given for six stations, all of them in the deeper water. These range from 52.3° to 56.4 F. (11.2° to 13.5° C.).

The stations for this species range well over the Archipelago and in the Sulu Sea, and to the southward in Darvel Bay, Borneo; Pitt Passage; and Buton Strait.

Quinqueloculina bicarinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13568	U.S.N.M.	3	D5097	14 19 15 N.; 120 33 52 E.	30	gy. m., s., sh.	Few.
13527	U.S.N.M.	6	D5100	14 17 15 N.; 120 32 40 E.	35	gy. s.	Common.
13528	U.S.N.M.	1	D5105	14 43 55 N.; 120 12 50 E.	25	Rare.
13529	U.S.N.M.	7	D5106	12 23 55 N.; 120 32 33 E.	37	gy. m.	Common.
13530	U.S.N.M.	10+	D5134	6 44 45 N.; 121 48 00 E.	25	fine s.	Common.
13531	U.S.N.M.	4	D5136	6 04 20 N.; 120 59 20 E.	22	s., sh.	Common.
13532	U.S.N.M.	8	D5137	6 04 25 N.; 120 58 30 E.	20	s., sh.	Common.
13533	U.S.N.M.	2	D5138	6 06 00 N.; 120 58 50 E.	19	s., Co.	Few.
13534	U.S.N.M.	6	D5139	6 06 00 N.; 121 02 30 E.	20	co. s.	Common.
13535	U.S.N.M.	4	D5141	6 09 00 N.; 120 58 00 E.	29	co. s.	Few.
13536	U.S.N.M.	10	D5142	6 06 10 N.; 121 02 40 E.	21	co. s., sh.	Common.
13537	U.S.N.M.	9	D5143	6 05 50 N.; 121 02 15 E.	19	co. s.	Common.
13538	U.S.N.M.	6	D5144	6 05 50 N.; 121 02 15 E.	19	co. s.	Few.
13539	U.S.N.M.	10	D5145	6 04 30 N.; 120 59 30 E.	Common.
13540	U.S.N.M.	6	D5146	5 46 40 N.; 120 48 50 E.	24	co. s.	Common.
13541	U.S.N.M.	2	D5147	5 41 40 N.; 120 47 10 E.	21	co. s.	Rare.
13542	U.S.N.M.	10+	D5148	5 35 40 N.; 120 47 30 E.	17	co. s.	Common.
13543	U.S.N.M.	10+	D5153	5 18 10 N.; 120 02 55 E.	49	co. s., sh.	Common.
13544	U.S.N.M.	7	D5154	5 14 50 N.; 119 58 45 E.	12	co. s., sh.	Few.
13545	U.S.N.M.	5	D5156	5 12 50 N.; 119 55 55 E.	18	fine s., sh.	Common.
13546	U.S.N.M.	1	D5159	5 11 50 N.; 119 54 00 E.	10	co. s.	Rare.
13547	U.S.N.M.	4	D5162	5 10 00 N.; 119 47 30 E.	230	52.9	ers. s., brk. sh.	Few.
13548	U.S.N.M.	10+	D5164	5 01 40 N.; 119 52 50 E.	18	gn. m.	Common.
13549	U.S.N.M.	9	D5192	11 09 15 N.; 123 50 00 E.	32	gn. s.	Common.
13550	U.S.N.M.	1	D5218	13 11 15 N.; 123 02 45 E.	20	ers. s.	Rare.
13551	U.S.N.M.	10+	D5220	13 38 00 N.; 121 58 00 E.	50	sft. gn. m.	Common.
13552	U.S.N.M.	3	D5276	13 49 15 N.; 120 14 45 E.	18	sh., p., s.	Few.
13553	U.S.N.M.	1	D5336	11 37 45 N.; 119 46 00 E.	46	s., m.	Rare.
13554	U.S.N.M.	5	D5338	11 33 45 N.; 119 24 45 E.	43	Co. s. m.	Common.
13555	U.S.N.M.	8	D5342	10 56 55 N.; 119 17 24 E.	14-25	gy. m.	Common.
13556	U.S.N.M.	6	D5348	10 57 45 N.; 118 38 15 E.	375	56.4	co. s.	Common.
13557	U.S.N.M.	4	D5358	6 06 40 N.; 118 18 15 E.	39	m.	Common.
13558	U.S.N.M.	1	D5381	13 14 15 N.; 122 44 45 E.	85	co. s.	Rare.
13559	U.S.N.M.	9	D5481	10 27 30 N.; 125 17 10 E.	61	s., sh., g.	Common.
13560	U.S.N.M.	1	D5549	6 01 15 N.; 120 44 20 E.	263	52.3	s., glob., for.	Rare.
13561	U.S.N.M.	1	D5572	5 31 26 N.; 120 09 45 E.	334	52.3	s.	Rare.
13562	U.S.N.M.	2	D5574	5 30 45 N.; 120 07 57 E.	340	Rare.
13563	U.S.N.M.	3	D5576	5 25 56 N.; 120 03 39 E.	277	53.3	s.	Rare.
13564	U.S.N.M.	1	D5579	4 54 15 N.; 119 09 52 E.	175	55.3	fine s., Co.	Rare.
13567	U.S.N.M.	2	D5636	1 55 00 S.; 127 42 30 E.	1,262	gy. m., fine s.	Rare.
13566	U.S.N.M.	2	D5640	4 27 00 S.; 122 55 40 E.	24	s., brk. sh.	Rare.
13565	U.S.N.M.	4	D5642	4 31 40 S.; 122 49 42 E.	37	gy. m.	Rare.

QUINQUELOCULINA FERUSSACII d'Orbigny.

Plate 90, figs. 2a-c.

Quinqueloculina ferussacii D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 301, No.

18.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 50, pl. 19, figs. 3, 4.

Miliolina ferussacii H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p.

175, pl. 113, figs. 17, a, b.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 507, pl.

12, figs. 6, 7.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol.

20, 1915, p. 578.

Specimens with the projecting slender neck and the acute marginal carinae typical of this species were found at a few stations, but never in any quantity. Of the seven stations at which this species is recorded the depth ranges from 18 to 148 fathoms (33 to 271 meters), the average being 61 fathoms (112 meters). Bottom temperature was given for the deepest station 54.4° F. (12.4° C.); none of the others are given. Therefore the average would be much above this figure. The localities are as follows: China Sea, off southern Luzon; Sulu Archipelago, off Tawi Tawi; vicinity of Romblon; off eastern Panay; off northern Cebu; Gulf of Davao; and China Sea, off Formosa. Specimens were frequent at two of the shallower stations and rare at the others. Heron-Allen and Earland record it "not widely distributed, but very fine and typical examples occur at several stations, though not in any numbers." They also mention that the form figured by Brady⁴⁰ "represents a type which is of much more frequent occurrence (than d'Orbigny's type) in tropical gatherings, having long, narrow chambers with little curvature, produced tubular neck, and strong, acute, marginal, facial carinae." This evidently is the same as the form that we have had from the Philippines.

Quinqueloculina ferussacii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13904	U.S.N.M.	3	D5096...	14 20 23 N.; 120 34 15 E...	25	gy. m., s., sh.	Rare.
13905	U.S.N.M.	1	D5164...	5 01 40 N.; 119 52 20 E...	18	gn. m.	Rare.
13906	U.S.N.M.	8	D5181...	11 36 40 N.; 123 26 35 E...	26	m, fne. s.	Frequent.
13907	U.S.N.M.	3	D5178...	12 43 00 N.; 122 06 15 E...	75	fne. s.	Rare.
13908	U.S.N.M.	6	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Frequent.
13909	U.S.N.M.	1	D5255...	7 03 00 N.; 125 39 00 E...	100	sft. m.	Rare.
13910	U.S.N.M.	3	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.	Rare.

QUINQUELOCULINA RUGOSA d'Orbigny.

Plate 100, figs. 6a-c.

Quinqueloculina rugosa D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 302, No. 24.—

SCHLUMBERGER, Mém. Soc. Zool. France, 1893, p. 210, text figs. 18, 19, pl. 4,

figs. 91-93.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 53.

This species, of which abundant specimens appear at some stations, seems to be well characterized. Schlumberger revived this

⁴⁰ *Challenger*, pl. 113, fig. 17.

species of d'Orbigny's, described in 1826, for recent material of his from the Gulf of Marseilles. I have referred to it in a recent paper (Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 53), tropical and subtropical material from the North Pacific, from the Hawaiian Islands, Japan, Hongkong, and Manila Bay. Its surface has a peculiarly individual character, rugose, not from accretions of extraneous matter, but from what seems an eroded character; the test, especially the outer wall, being conspicuously pitted and rugose. This character persists at various ages, and in the whole series of adults from numerous stations, in fact, it is one of the most characteristic species of the Philippine material. Heron-Allen and Earland in their Kerimba paper⁴ refer to this species as related to *Q. contorta*:

Almost universally distributed, abundant, and attaining very fine proportions at some stations, especially at stations 9, 12, and ?X. There is, as usual, a considerable number of specimens passing into *M. sclerotica*, the essential difference between these two species being the nature of the superficial test. At station 1 the individuals all closely resemble the figure of *Quinqueloculina rugosa* given by Fornasini from the "Planchés inédites" (F. 1905, Mem. Accad. Sci. Bologna, ser. 6, vol. 2, pl. 3, fig. 13) but they have a porcellaneous texture, although, like all other specimens of *M. contorta*, the surface is matt. The nature of the shell of *Q. rugosa* must remain purely speculative, the species having its origin in a *nomen nudum*, the "Planché inédite" giving practically no guide to the texture. Schlumberger has identified d'Orbigny's specific name *rugosa* with specimens from Marseille (Mém. Soc. Zool. p. 68, pl. 4, (not 2), figs. 91-93, text figs. 18, 19), which differ considerably from the "Planché inédite" in external characteristics, but which are described as having a rough surface ("Tête d'apparence rugose"); they appear, therefore, to be a form of *M. sclerotica*.



FIG. 24.—QUINQUELOCLINA RUGOSA D'ORBIGNY. APERTURE AND UNDIVIDED SIMPLE TOOTH OF EARLY STAGE. X 150, FROM ALBATROSS STATION D5164.

However, in the Philippine material *Q. contorta* occurs at several stations, but in all cases of depths from 200 to 300 fathoms (366 to 549 meters) or more, and while it has the matt surface it is never, in this material, to be confused with what I have here called *Q. rugosa*.

Moreover, the two are not found together at any station—a fact which seems rather remarkable. *Q. rugosa* is more at home apparently in shallow water, for it is there that specimens were most abundant and of the largest size. It is not related to *Q. sclerotica* Karrer, as the rugose character is not due in *Q. rugosa* to accretion of other material, as in *Q. sclerotica*, but results from the peculiar character of the original wall, as already noted. It would seem to be a species characteristic of comparatively shallow warm waters of the Mediterranean and Indo-Pacific regions. Wiesner notes the relations of *Q. rugosa* and *Q. contorta* in the material from the Adriatic.

The Philippine records include 16 stations, ranging in depth from 18 to 375 fathoms (33 to 686 meters), the average depth being 55

⁴ Trans. Zool. Soc. London, 1915, vol. 20, p. 576.

fathoms (100 meters). Bottom temperatures are given at but three stations, ranging from 56.4° F. (13.5° C.) at the deepest station to 79.5° F. (26.3° C.) at 38 fathoms (70 meters). The average temperature for all stations undoubtedly would be around 75° F. (23.8° C.). These stations include the following localities: Jolo Sea; Palawan Passage; China Sea off southern Luzon; between Marinduque and Luzon; off northern Cebu; northeastern Panay; vicinity of Romblon; Sulu Sea; Tawi Tawi Group, and off western Mindanao. It did not occur at all in the regions to the south, probably because of the depth of water.

Quinqueloculina rugosa—material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13701	U.S.N.M.	2	D5096...	14 20 23 N.; 120 34 15 E..	28	° F.	gy. m., s., sh	Frequent.
13702	U.S.N.M.	7	D5100...	14 17 15 N.; 120 32 40 E..	35	gy. s.	Frequent.
13703	U.S.N.M.	2	D5105...	14 43 55 N.; 120 12 50 E..	25	Few.
13704	U.S.N.M.	4	D5106...	14 23 55 N.; 120 32 33 E..	37	gy. m.	Few.
13705	U.S.N.M.	10+	D5132...	Island off Panabutan Pt., N.; 15° W. 0.3 mile.	26	gn. m., s.	Common.
13706	U.S.N.M.	7	D5133...	Island off Panabutan Pt., N.; 52° E. 1.5 miles.	38	79.5	gn. m., s.	Common.
13707	U.S.N.M.	10+	D5164...	5 01 40 N.; 119 52 20 E..	18	gn. m.	Common.
13708	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.	Rare.
13709	U.S.N.M.	9	D5181...	11 36 40 N.; 123 26 35 E..	26	fine. m., s.	Common.
13710	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.	Few.
13696	U.S.N.M.	6	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.	Few.
13697	U.S.N.M.	4	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s.	Few.
13711	U.S.N.M.	1	D5333...	11 33 45 N.; 119 24 45 E..	43	co. s., m.	Few.
13712	U.S.N.M.	6	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	Frequent.
13698	U.S.N.M.	5	D5358...	6 06 40 N.; 118 18 15 E..	39	m.	Frequent.
13699	U.S.N.M.	2	D5640...	4 27 00 S.; 122 55 40 E..	24	s., brk. sh.	Rare.

QUINQUELOCULINA POLYGONA d'Orbigny.

Plate 90, figs. 3a-c.

Quinqueloculina polygona d'ORBIGNY, in De La Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 198, pl. 12, figs. 21-23.

There are a few specimens which seem to be very close to this West Indian species described by d'Orbigny. This is placed by Heron-Allen and Earland as a synonym of *Q. ferussacii*, but in the Philippines the two seem to be distinct. In the West Indies this is a well-characterized species.

Specimens are rare at a few stations. They are decidedly concave on the periphery between the two keels, the lateral faces at right angles to the periphery, or even concave, while *Q. ferussacii* has extended single keels unless in the last-formed chamber.

The localities include the following: Sulu Archipelago, off Tawi Tawi; off eastern Panay; between Marinduque and Luzon; and the Palawan Passage.

They range in depth from 10 to 375 fathoms (18 to 686 meters) or less.

Quinqueloculina polygona—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
				° ' " ° ' "		° F.		
13831	U.S.N.M.	2	D5159...	50 11 50 N.; 119 54 00 E..	10	co. s.	Rare.
13827	U.S.N.M.	1	D5161...	5 10 15 N.; 119 53 00 E..	16	fine. bk. s.	Rare.
13828	U.S.N.M.	1	D5181...	11 36 49 N.; 123 26 35 E..	26	m., fine. s.	Rare.
13829	U.S.N.M.	1	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.	Rare.
13830	U.S.N.M.	1	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	Co., s.	Rare.

QUINQUELOCULINA BRADYANA Cushman.

Miliolina undosa H. B. BRADY (not *Quinqueloculina undosa* Karrer), Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 176, pl. 6, figs. 6-8.

Quinqueloculina bradyana CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 52, pl. 18, fig. 2.

There are a very few specimens which seem referable to this species. They are not at all alike and may represent more than one form.

Specimens were noted at seven stations, ranging in depth from 10 to 318 fathoms (18 to 582 meters), the average depth being 93 fathoms (170 meters). These were mostly from shallow water about the islands in the Sulu Archipelago, off southern Luzon, and between Burias and Luzon.

Quinqueloculina bradyana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
10804	U.S.N.M.	2	D5109...	14 03 45 N.; 120 16 30 E..	10	Co.	Rare.
10805	U.S.N.M.	1	D5154...	5 14 50 N.; 119 58 45 E..	12	co. s.	Rare.
10806	U.S.N.M.	2	D5160...	5 12 40 N.; 119 55 10 E..	12	s.	Rare.
10807	U.S.N.M.	2	D5161...	5 10 15 M.; 119 53 00 E..	16	fine. s., blk. sp.	Few.
10808	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E..	318	fine. s., sh.	Rare.
10809	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E..	29	crs. s.	Rare.
10810	U.S.N.M.	4	D5549...	6 01 15 N.; 120 44 20 E..	263	52.3	s., glob., f.	Rare.

QUINQUELOCULINA CONTORTA d'Orbigny.

Plate 90, figs. 1a-c.

Quinqueloculina contorta D'ORBIGNY, Foram. Foss. Vienne, 1846, p. 298, pl. 20, figs. 4-6.

Miliolina contorta BRADY, Journ. Roy. Micr. Soc., London, 1887, p. 881.—HALKYARD, Trans. Ann. Rep. Manchester Micr. Soc., 1889, p. 60, pl. 1, fig. 4.—SIDEBOTTOM, Mem. Lit. Philos. Soc., Manchester, pt. 1, 1904, p. 13, pl. 4, figs. 7-9.—EARLAND, Journ. Quekett Micr. Club, 1905, p. 195.—HERON-ALLEN and EARLAND, Proc. Roy. Irish Acad., 1913, p. 30; Trans. Zool. Soc. London, vol. 20, 1915, p. 576.

Heron-Allen and Earland have suggested that this species and *Q. rugosa* may be the same species. In the Philippine material the two seem to be very different. Specimens referred to *Q. contorta* have occurred at six stations, ranging in depth from 200 to 340 fathoms (366 to 622 meters), the average 267 fathoms (489 meters). Bottom temperatures are very uniform. Those given for five sta-

tions range from between 52.2° F. to 54.3° F. (11.2° C. to 12.3° C.), the average being 52.7° F. (11.4° C.).

No specimens referred to this species occurred at less than 200 fathoms, while most of the specimens of *Q. rugosa* occurred in much shallower water. The surface in typical *Q. contorta* is smooth and matt, and is very different from the pitted surface of *Q. rugosa*. The stations for *Q. contorta* are in the following localities: Off northern Mindanao; off Jolo; between Jolo and Tawi Tawi: north of Tawi Tawi; and off western Mindanao, all of the stations but the first being in the Sulu Sea, where dredgings were made in comparatively deep water. Heron-Allen and Earland record it as abundant in the Kerimba Archipelago.

Quinqueloculina contorta—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13722	U. S. N. M.	1	D5542...	8 48 30 N.; 123 35 30 E..	200	54.3	fine s., brk. sh.	Few.
13723	U. S. N. M.	1	D5549...	6 01 15 N.; 120 44 20 E..	263	52.3	s., glob., for..	Few.
13724	U. S. N. M.	2	D5565...	5 51 42 N.; 120 30 30 E..	243	52.3	s., ptr. sh....	Few.
			D5571...	5 30 45 N.; 120 07 57 E..	349	52.3	s., sh.....	Few.
13725	U. S. N. M.	1	D5572...	5 31 26 N.; 120 09 45 E..	334	52.3	s.....	Few.
			H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob..	Few.

QUINQUELOCULINA SCULPTURATA, new species.

Plate 91, figs. 3, 4.

Description.—Test elongate, chambers with the outer face broadly convex, sides concave, the angle between the peripheral and lateral faces sharp, chambers much the broadest near the initial end, tapering toward the apertural end; peripheral face in the young with deep oblique chambers and ridges, in the later development becoming a coarse reticulation; apertural end very slightly produced, with a single, long tooth.

Length up to 1 mm.

Quinqueloculina sculpturata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13626	U. S. N. M.	1	D5143...	6 05 50 N.; 121 02 15 E..	19	co. s.....	Rare.
13629	U. S. N. M.	1	D5144...	6 05 50 N.; 121 02 15 E..	19	co. s.....	Rare.
13628	U. S. N. M.	2	D5145...	6 04 30 N.; 120 59 30 E..	23	co. sh.....	Rare.
13627	U. S. N. M.	2	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Rare.
				Jolo Jolo, P. I.....	Rare.

Distribution.—Type specimen (Cat. No. 13626, U. S. N. M.) from *Albatross* station D5143, off Jolo, in 19 fathoms (35 meters). It has also occurred at the two neighboring stations, D5144 and D5145, in 19 and 23 fathoms (35 and 42 meters), and D5218, between Burias and Luzon, in 20 fathoms (37 meters).

This is similar in some respects to one of Heron-Allen and Earland's figures of *Q. kerimbatica*.⁴²

Both the young and adult are very different from any of the other reticulate forms of the Philippine material.

QUINQUELOCULINA RETICULATA (d'Orbigny).

- Triloculina reticulata* D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 299, No. 9.
Quinqueloculina reticulata KARRER, Sitz. Akad. Wiss. Wien, vol. 44, 1861, p. 449, pl. 2, fig. 5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 55, pl. 16, figs. 1-3.
Miliolina reticulata PARKER, JONES, and BRADY, Ann. Mag. Nat. Hist., ser. 4, vol. 8, 1871, p. 249, pl. 8, fig. 18.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 177, pl. 9, figs. 2-4.—SCHLUMBERGER, Mém. Soc. Zool. France, vol. 6, 1893, p. 214, text fig. 25, pl. 2, fig. 62.

To this species are referred several specimens, which are found at a number of stations, with periphery and reticulations at the sides somewhat like *Q. sagra* d'Orbigny but not bicarinate. A study of the West Indian shallow-water material, together with d'Orbigny's Cuban Monograph, shows that *Q. reticulata* in the West Indies is a very definite species and similar in general characters to that referred to here, but not including the various reticulate forms or species which are referred to this name by various authors.

Typical material occurred at 15 stations, ranging in depth from 17 to 303 fathoms (31 to 554 meters), the average being 76 fathoms (139 meters). Bottom temperatures are given at two stations; these are 52.3° F. (11.2° C.), and 53.3° F. (11.8° C.); as both represent the deepest stations in the series, the average for the shallower water stations would probably bring this up to 75° F. (23.8° C.) at least. All but one of the stations are in the Archipelago proper, most of them in the Sulu region. It has occurred in the Sulu Archipelago, off Jolo, off Siasi, and off Tawi Tawi; also off northern Cebu, between Burias and Luzon; Verde Island Passage; and to the southward in Buton Strait, where it occurred at 24 fathoms (44 meters), showing that the species is probably widely distributed there if shallow-water material were obtainable.

Quinqueloculina reticulata—Material examined.


Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13683	U.S.N.M.	1	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.....	Rare.
13684	U.S.N.M.	10+	D5137...	6 04 25 N.; 120 58 30 E...	20	s., sh.....	Common.
13685	U.S.N.M.	2	D5141...	6 09 00 N.; 120 58 00 E...	29	co. s.....	Few.
13686	U.S.N.M.	8	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.....	Common.
13687	U.S.N.M.	4	D5146...	5 46 49 N.; 120 48 50 E...	24	co. s., sh.....	Few.
13688	U.S.N.M.	3	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.....	Few.
13689	U.S.N.M.	2	D5148...	5 35 40 N.; 120 47 30 E...	17	co. s.....	Few.
13690	U.S.N.M.	4	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Few.
13691	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Rare.
13692	U.S.N.M.	5	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Few.
13693	U.S.N.M.	1	D5269...	13 39 50 N.; 129 59 30 E...	220	fine s., p.....	Rare.
13700	U.S.N.M.	2	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.....	Rare.
13694	U.S.N.M.	2	D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.....	Rare.
13695	U.S.N.M.	5	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh.....	Few.
				Tara Island, P. I.....				

⁴² Trans. Zool. Soc., London, vol. 20, 1915, pl. 43, fig. 18

QUINQUELOCULINA ALBATROSSI, new species.

Plate 85, figs. 4a-c.

Description.—Test large, quinqueloeculine throughout, elongate, oval; in the young the chambers with rounded angles on the periphery, later becoming convexly rounded, ornamentation consisting of very numerous fine reticulations, regularly arranged in the young specimens, the reticulations gradually being limited to the sides of the chambers and the peripheral face becoming smooth, and finally the entire chamber smooth; the reticulations show in places near the inner margin or not at all; aperture rounded slightly, if at all, above



FIGS. 25-33.—QUINQUELOCULINA ALBATROSSI, NEW SPECIES. $\times 50$. VARIOUS STAGES IN THE DEVELOPMENT OF THE APERTURAL CHARACTERS. 25, A SIMPLE TOOTH IN A GENERALLY ROUNDED APERTURE. 26-29, A BIFID TOOTH, THE APERTURE GRADUALLY ASSUMING AN IRREGULAR FORM BY THE DEVELOPMENT OF PROCESSES AND REENTRANTS. 30, ONE ARM OF THE BIFID TOOTH BROADENING, THE OTHER ALREADY JOINED TO THE SIDE OF THE APERTURE. 31, 32, STAGES IN THE BROADENING OF THE TOOTH AFTER THE CENTRAL OPENING IS ENTIRELY SURROUNDED. 33, A LATER STAGE WHERE THE SIDES HAVE FUSED INTO THE PROCESSES LEAVING A CENTRAL APERTURE AND SIX SUPPLEMENTARY ONES ABOUT IT.

the base of the preceding chamber; the margin with a thickened lip, and in the adult a bifid tooth.

Length up to nearly 3 mm.

Distribution.—Type specimen (Cat. No. 13877, U. S. N. M.) from *Albatross* station D5133, Sulu Sea, off western Mindanao, 38 fathoms (70 meters).

Specimens have been found in the material from the following general localities: China Sea, off southern Luzon; Sulu Sea, off western Mindanao, near Basilan Island, off Jolo, off Tawi Tawi; off Romblon; off northern Cebu; off western Samar; Gulf of Davao; China Sea, off Formosa; between Cebu and Bohol; between Samar and Leyte; between Negros and Siquijor; Tara Island; and to the south at two stations in Buton Strait.

These stations range in depth from 18 to 500 fathoms (33 to 914 meters), the average 87.8 fathoms. Bottom temperatures are given but for eight stations, ranging from 52.9° F. to 79.5° F. (11.6° C. to

26.3° C.); the average temperature, 63.2° F. (17.3° C.). As these are mostly from the deeper stations, the average temperature would probably be around 75° F. (23.8° C.). Altogether it has occurred at 31 stations, and at many of them abundantly; therefore it is one of the characteristic species of the region. This is one of the fine large species dredged by the *Albatross*, and is abundant in the Philippine Archipelago, and probably in the region southward in shallow water. It is very distinctive at all ages, and can hardly be confused with any other reticulate species, as the very fine sculpture resembling *Spiroloculina elegans* are not so much reticulations as regular lozenge-shaped and elliptical depressions of the surface, in many of the specimens the wall between each series of depressions being as wide as the depressions themselves. The progressive smoothness of the peripheral margin and the retention of the sculpture proximally is just the opposite condition of *Q. kerimbatica*, var. *philippinensis*. Some of the large smooth adults would not be thought related to the reticulate group but for the large series of stages and by cutting back to the earlier chambers.

Quinqueloculina albatrossi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13871	U.S.N.M.	5	D5096...	14 20 23 N.; 120 34 15 E...	28	gy. m., s. sh.	Frequent.
13872	U.S.N.M.	10+	D5097...	14 19 15 N.; 120 33 52 E...	30	gy. m., s. sh.	Common.
13873	U.S.N.M.	10+	D5105...	14 43 55 N.; 120 12 50 E...	25	Common.
13874	U.S.N.M.	10+	D5106...	14 23 55 N.; 120 32 33 E...	37	gy. m.	Common.
13875	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E...	135	59.0	dk. gy. m.	Rare.
13876	U.S.N.M.	10+	D5132...	Island off Panabutan Point, N.; 15° W., 0.30 miles.	26	gn. m., s.	Abundant.
13877	} U.S.N.M.	10+	D5133...	Island off Panabutan Point, N.; 52° E., 1.50 miles.	38	79.5	gn. m., s.	Common.
13878								
13879	U.S.N.M.	3	D5134...	6 44 45 N.; 121 48 00 E...	25	fine s.	Few.
13880	U.S.N.M.	10+	D5137...	6 04 25 N.; 120 58 30 E...	20	s., sh.	Common.
13881	U.S.N.M.	10+	D5142...	6 06 10 N.; 121 02 40 E...	21	co. s., sh.	Common.
13882	U.S.N.M.	2	D5144...	6 05 50 N.; 121 02 15 E...	19	co. s.	Rare.
13883	U.S.N.M.	8	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.	Frequent.
13884	U.S.N.M.	1	D5147...	5 41 40 N.; 120 47 10 E...	21	co. s., sh.	Rare.
13885	U.S.N.M.	10+	D5151...	5 24 40 N.; 120 27 15 E...	24	co. s., sh.	Common.
13886	U.S.N.M.	2	D5152...	5 22 55 N.; 120 15 45 E...	34	whit. s.	Few.
13887	U.S.N.M.	4	D5162...	5 10 00 N.; 119 47 30 E...	230	52.9	crs. s., brk. sh.	Few.
13888	U.S.N.M.	4	D5172...	6 03 15 N.; 120 35 30 E...	318	fine s., sh.	Few.
13890	U.S.N.M.	8	D5179...	12 38 15 N.; 122 12 39 E...	37	75.7	hrd. s.	Few.
13889	U.S.N.M.	5	D5178...	12 43 00 N.; 122 06 15 E...	78	fine s.	Few.
13891	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Common.
13892	U.S.N.M.	4	D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fine gy. s.	Few.
13893	U.S.N.M.	10+	D5259...	7 05 07 N.; 125 39 45 E...	23	co. s.	Common.
13894	U.S.N.M.	7	D5255...	7 03 00 N.; 125 39 00 E...	100	sh. m.	Few.
13895	U.S.N.M.	5	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p. s.	Few.
13896	U.S.N.M.	1	D5315...	21 40 00 N.; 116 53 00 E...	148	54.4	s., sh.	Rare.
13897	U.S.N.M.	1	D5419...	9 53 30 N.; 123 46 00 E...	175	54.5	gn. m.	Rare.
13898	U.S.N.M.	3	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.	Few.
13899	U.S.N.M.	10+	D5478...	10 46 24 N.; 125 16 30 E...	57	sh.	Common.
13900	U.S.N.M.	1	D5538...	9 08 15 N.; 123 23 20 E...	256	53.3	gn. m., s.	Rare.
13901	U.S.N.M.	10	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh.	Common.
13902	U.S.N.M.	3	D5942...	4 31 40 S.; 122 49 42 E...	37	gy. m.	Few.
13903	U.S.N.M.	3		Tara Island, P. I.	Few.

QUINQUELOCULINA KERIMBATICA (Heron-Allen and Earland).

Miliolina parkeri MILLETT, Journ. Roy. Micr. Soc., 1898, p. 507, pl. 12, fig. 4.

Miliolina kerimbatica HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 574, pl. 43, figs. 13-23.

The following is the original description:

Test free, quinqueloculine. The walls of the chambers thick, irregularly furrowed in all directions with broad, deeply gouged-out channels, running obliquely and irregularly across the face of each chamber and generally connecting with a deeper straight furrow excavated down the peripheral edge. This straight peripheral furrow, when exposed on an earlier chamber in the center of the test by the quinqueloculine arrangement of the shell, affords a very striking appearance by contrast with the transverse furrows on the surface of the surrounding chambers. Aperture large and furnished with a prominent tooth. The oral end is usually but slightly produced, but in some individuals the aperture is situated on a produced neck. The ridges between the furrows are flat on the top—i. e., the furrows are cleanly gouged out of the shell substance.

The typical form of this species with irregular, excavate peripheral faces, and with flat-topped ridges, is very rare in the Philippine material, being replaced by varieties showing very constant characters.

The typical form occurred at the following stations: Tara Island, in shallow water, and D5143, in 19 fathoms (35 meters), off Jolo.

The types, of which the above is the original description, were from the Kerimba Archipelago, off southeastern Africa, described by Heron-Allen and Earland.⁴³

Quinqueloculina kerimbatica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13486	U.S.N.M.	1	D5143...	6° 05' 59" N.; 121° 02' 15" E...	19	° F.	-	Rare.
13485	U.S.N.M.	1	Tara Island, P. I.	co. s.	Rare.

QUINQUELOCULINA KERIMBATICA, var. RETICULO-STRIATA, new variety.

Plate 89, figs. 1a-c.

Description.—Test with peripheral faces rather squarely truncate and ornamented by longitudinal narrow costae, the sides of the chambers with oblique areolae; the apertural end contracted and terminating in a short cylindrical neck, and thin philaline lip with a bifid tooth.

Length 2 mm.

Description.—Type specimen (Cat. No. 13488, U.S.N.M.) from Albatross station D5276, in 18 fathoms (33 meters), China Sea, off southern Luzon.

This variety is larger than variety *philippinensis*, and is closer to the typical form of the species; but the peripheral face of the chamber has sharp, usually very even costae, and the sides are reticulated

⁴³ Trans. Zool. Soc., London, vol. 20, 1915, p. 574.

much as in *Q. parkeri*. It is not as common as the following variety, but has occurred at eight stations in the region, ranging in depth from 14 to 260 fathoms (25 to 476 meters).

The following localities are represented: Malampaya Sound, Palawan Island; Palawan Passage; Sulu Archipelago, off Tawi Tawi; off Jolo; China Sea, off southern Luzon; and one station to the south in Sibuko Bay, Borneo.

Quinqueloculina kerimbatica, var. *reticulo-striata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13487	U.S.N.M.	1	D5151...	5 24 40 N.; 120 27 15 E...	24	° F.	co. s., sp.....	Rare.
13493	U.S.N.M.	1	D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fne. gy. s.....	Rare.
13488	U.S.N.M.	10+	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Common.
13489	U.S.N.M.	8	D5338...	11 33 45 N.; 119 24 45 E...	43	Co., s., m.....	Common
13490	U.S.N.M.	1	D5339...	11 22 00 N.; 119 12 00 E...	52	m.....	Rare.
13491	U.S.N.M.	2	D5342...	10 56 55 N.; 119 17 24 E...	14-25	gy. m.....	Few.
13492	U.S.N.M.	2	D5358...	6 06 40 N.; 118 18 15 E...	39	m.....	Common.
13494	U.S.N.M.	1	D5589...	4 12 10 N.; 118 38 08 E...	260	45.7	fne. gy. s., gy. m.	Rare.

QUINQUELOCULINA KERIMBATICA, var. PHILIPPINENSIS, new variety.

Plate 89, figs. 2, 3.

Description.—Test elongate, quinqueloculine; peripheral margin of the chambers broadly rounded, sides convex, surface ornamented; the peripheral face with regular reticulations with thin walls between, large but very regular, sides either with the reticulate pattern entire, or becoming obsolescent or even wanting, especially in older specimens; apertural end of the chamber projecting and contracted to form an exsert cylindrical neck without ornamentation, with a thin border, not thickened, but with a phialine lip and a simple bifid tooth.

Length seldom exceeding 1 mm.

Description.—Type specimen (Cat. No. 13510, U. S. N. M.) from Albatross station D5159, Sulu Archipelago, Tawi Tawi Group, in 10 fathoms (18 meters).

It has occurred at numerous stations, all but one in the Philippine Archipelago. There are 32 stations, ranging in depth from 10 to 500 fathoms (18 to 914 meters), the average depth being 62 fathoms (114 meters). Bottom temperatures are given for only two stations—one in 375 fathoms, 56.4° F. (13.5° C.), the other in 50 fathoms (914 meters), 76.3° F. (24.6° C.). Therefore the average, including the various shallow-water stations, would be close to the latter figure. The species is abundant in shallow water and becomes less common as depths increase above 50 fathoms (91 meters).

These stations include the following localities: Tacloban Anchorage; China Sea, off southern Luzon; Sulu Sea, off western Mindanao,

near Basilan Island, off Tawi Tawi; off Romblon; off eastern Panay; off northern Cebu; off western Samar; between Burias and Luzon; between Marinduque and Luzon; Palawan Passage; Malampaya Sound, Palawan Island; off eastern Palawan; between Samar and Leyte; and to the south in Buton Strait.

This variety differs from the typical form in the greater regularity of the ornamentation, and in the peripheral margin which is not striate but reticulate. The progressive appearance of the smooth area on the periphery is just the opposite of the development in *Q. albatrossi*.

While there is a considerable variation in the amount of surface covered by the reticulate ornamentation, the form of the test, the produced neck without the reticulations, and the thin phialine lip, all mark it at once in a mixed lot of material, and these characters remain very constant. While here placed as a variety of *Q. kerimbatica*, it may be specifically different. At any rate it appears to replace *Q. kerimbatica* almost entirely in the Philippine region.



FIG. 34.—QUINQUELOCULINA KERIMBATICA HERON-ALLEN AND EARLAND, VAR. PHILIPPINENSIS, APERTURAL VIEW. $\times 50$, FROM ALBATROSS STATION D5164.

Quinqueloculina kerimbatica, var. *philippinensis*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13496	U.S.N.M.	10+	D5096...	14 20 23 N.; 120 34 15 E...	28	gy. m., s....	Common.
13497	U.S.N.M.	10+	D5097...	14 19 15 N.; 120 33 52 E...	30	gy. sh. s....	Common.
13499	U.S.N.M.	10+	D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s....	Common.
13500	U.S.N.M.	8	D5105...	14 43 55 N.; 120 12 50 E...	25	fne. s., sh....	Common.
13502	U.S.N.M.	7	D5106...	14 23 55 N.; 120 32 33 E...	37	gy. m....	Common.
13503	U.S.N.M.	3	D5113...	13 51 30 N.; 120 50 30 E...	159	dk. gn. m....	Few.
13504	U.S.N.M.	10+	D5132...	Island off Panabutan Point, N. 15 W°; 0.30 mile.	26	gn. m., s....	Common.
13505	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N. 15° W.; 1.50 miles.	38	gn. m., s....	Common.
13506	U.S.N.M.	10+	D5134...	6 44 45 N.; 121 48 00 E...	25	fne. s....	Common.
13507	U.S.N.M.	10+	D5152...	5 25 55 N.; 120 15 45 E...	34	wh. s....	Common.
13508	U.S.N.M.	2	D5153...	5 18 10 N.; 120 02 55 E...	49	co. s., sh....	Few.
13509	U.S.N.M.	10+	D5156...	5 12 50 N.; 119 55 55 E...	18	fne. s., sh....	Common.
13510	U.S.N.M.	10	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s....	Common.
13511	U.S.N.M.	10+	D5161...	5 10 15 N.; 119 53 00 E...	16	fne. s....	Common.
13512	U.S.N.M.	10+	D5164...	5 01 40 N.; 119 52 20 E...	18	gn. m....	Common.
13498	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E...	78	fne. s....	Common.
13501	U.S.N.M.	10+	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fne. s....	Common.
13514	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s....	Common.
13515	U.S.N.M.	7	D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fne. gy. s....	Common.
13517	U.S.N.M.	2	D5218...	13 11 15 N.; 123 02 45 E...	20	cr. s....	Few.
13516	U.S.N.M.	10+	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m....	Common.
13526	U.S.N.M.	2	D5278...	13 49 15 N.; 120 14 45 E...	18	sh., p., s....	Rare.
13518	U.S.N.M.	1	D5336...	11 37 45 N.; 119 46 00 E...	46	s., m....	Rare.
13519	U.S.N.M.	2	D5338...	11 33 45 N.; 119 24 45 E...	43	co. s., s., m....	Few.
13520	U.S.N.M.	1	D5342...	10 56 55 N.; 119 17 24 E...	14-25	gy. m....	Rare.
13522	U.S.N.M.	8	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s....	Common.
13521	U.S.N.M.	2	D5423...	12 12 00 N.; 118 28 00 E...	27	fne. gy. s....	Few.
13523	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m. (net).	Rare.
13524	U.S.N.M.	1	D5478...	10 46 24 N.; 125 16 30 E...	57	sh....	Rare.
13513	U.S.N.M.	6	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh., g....	Few.
13525	U.S.N.M.	1	D5642...	4 31 40 S.; 122 49 42 E...	37	gy. m....	Rare.
13495	U.S.N.M.	10+	Tacloban Anchorage, P. I.	Abundant.

QUINQUELOCULINA PARKERI (H. B. Brady).

Plate 86, figs. 4a-c.

"*Quinqueloculina* with oblique ridges" PARKER, Trans. Micr. Soc. London, vol. 6, 1858, p. 53, pl. 5, fig. 10.

Miliolina parkeri H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 46; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 177, pl. 7, fig. 14.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, 1915, vol. 20, p. 574.

Quinqueloculina parkeri CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 50, pl. 15, fig. 3.

This species, which seems to be a typical one of coral-reef regions, is fairly common in the Philippine area. All 28 stations at which it is recorded are in the Archipelago proper, none of the records being from the regions to the south. This is probably because those stations represent deeper water stations, and it is probably found in shallower water, as recorded by Millett and Chapman. The stations range in depth from 10 to 318 fathoms (18 to 582 meters), with the average depth 45 fathoms (82 meters). Bottom temperatures are given but for three stations, ranging from 52.9° F. to 75.7° F. (11.6° C. to 26.3° C.), the average temperature being 62.3° F. (16.7° C.). As the stations at which bottom temperatures are lacking are mostly those in the shallower water, the average bottom temperature would probably be nearer the maximum given here. From the records the species seems to be commonest in waters less than 30 fathoms (55 meters), and less frequent at depths greater than these. There is some variation, but on the whole the specimens are rather typical in their characters. From the available records this seems to be a species ranging from the Hawaiian Islands on the east to the Kerimba Archipelago and the Red Sea on the west.

Quinqueloculina parkeri—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10811	U.S.N.M.	8	D5106...	14 23 55 N.; 120 32 33 E...	37	° F.	gy. m.....	Common.
10812	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E...	10		co.....	Rare.
10813	U.S.N.M.	1	D5133...	Island off Panabutan Point N; 52° E., 1.5 miles	38		gn. m., s.....	Rare.
10814	U.S.N.M.	2	D5136...	6 01 20 N.; 120 59 20 E...	22		s., sh.....	Rare.
10815	U.S.N.M.	4	D5137...	6 04 25 N.; 120 58 30 E...	20		s., sh.....	Common.
10816	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E...	19		s., Co.....	Few.
10817	U.S.N.M.	1	D5139...	6 06 00 N.; 121 02 30 E...	20		co. s.....	Rare.
10818	U.S.N.M.	4	D5142...	6 06 10 N.; 121 02 40 E...	21		co. s., sh.....	Few.
10819	U.S.N.M.	5	D5143...	6 05 50 N.; 121 02 15 E...	19		co. s.....	Frequent.
10820	U.S.N.M.	2	D5144...	6 05 50 N.; 121 02 15 E...	19		co. s.....	Rare.
10821	U.S.N.M.	7	D5145...	6 01 30 N.; 120 59 30 E...	23		co. s., sh.....	Frequent.
10822	U.S.N.M.	3	D5148...	5 35 40 N.; 120 47 30 E...	17		co. s.....	Frequent.
10823	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34		wht. s.....	Rare.
10824	U.S.N.M.	1	D5153...	5 18 10 N.; 120 02 55 E...	49		co. s., sh.....	Rare.
10825	U.S.N.M.	9	D5154...	5 14 50 N.; 119 58 45 E...	12		co. s.....	Frequent.
10826	U.S.N.M.	8	D5159...	5 11 50 N.; 119 54 00 E...	10		co. s.....	Frequent.
10827	U.S.N.M.	3	D5160...	5 12 40 N.; 119 55 10 E...	12		s.....	Rare.
10828	U.S.N.M.	3	D5162...	5 10 00 N.; 119 47 30 E...	230	52.9	crs. s., brk. sh	Few.
10829	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318		fne. s., sh.....	Rare.
10830	U.S.N.M.	4	D5178...	12 43 00 N.; 122 06 15 E...	78		fne. s.....	Few.
10831	U.S.N.M.	2	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Few.
10832	U.S.N.M.	3	D5192...	11 09 15 N.; 123 00 00 E...	32		gn. s.....	Few.
10833	U.S.N.M.	6	D5218...	13 11 15 N.; 123 02 45 E...	20		crs. s.....	Common.
10834	U.S.N.M.	1	D5220...	13 38 00 N.; 121 58 00 E...	50		sft. gn. m.....	Rare.
10835	U.S.N.M.	1	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fne. s., m., sh.	Rare.
10836	U.S.N.M.	1	Tara Island	Rare.
10837	U.S.N.M.	5	Binang Pool, Subin Bay	Frequent.
10838	U.S.N.M.	2	Shell sand from beach, Midway Island.	Frequent.

QUINQUELOCULINA AGGLUTINANS d'Orbigny.

Plate 91, figs. 1a-c.

Quinqueloculina agglutinans D'ORBIGNY, in De la Sagra, Hist. Fis. Pol. Nat., Cuba, 1839, "Foraminifères," p. 168, pl. 12, figs. 11-13 (not of later authors).

From a study of West Indian material from shallow water and a comparison with d'Orbigny's figures it is clear that the true *Q. agglutinans* is an arenaceous species with a squarish edge to the peripheral portion. A very similar form occurs in the Philippine material at several stations. It is close to the forms assigned by Heron-Allen and Earland to *Q. sclerotica*.

Material of this species occurred at nine stations, ranging in depth from 10 to 88 fathoms (18 to 161 meters), with the average 31 fathoms (57 meters). The only station at which bottom temperatures were given was 75.7° F. (24.2° C.).

These stations are all in and about the Archipelago; China Sea, off southern Luzon; off Jolo; Sulu Archipelago, Tawi Tawi Group; off Romblon; between Burias and Luzon; and Ragay Gulf, Luzon.

Quinqueloculina agglutinans—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13605	U.S.N.M.	10+	D5106...	14 23 55 N.; 120 32 33 E...	37	gy. m.....	Abundant.
13612	U.S.N.M.	1	D5139...	6 06 00 N.; 121 02 30 E...	20	co. s.....	Rare.
13609	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Rare.
13635	U.S.N.M.	5	D5156...	5 12 50 N.; 119 55 55 E...	18	fine. s., sh.....	Frequent.
13606	U.S.N.M.	5	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.....	Frequent.
13608	U.S.N.M.	1	D5161...	5 10 15 N.; 119 53 00 E...	16	fine. s., blk sp.....	Rare.
13610	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Rare.
13611	U.S.N.M.	8	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Frequent.
13607	U.S.N.M.	1	D5381...	13 14 15 N.; 122 44 45 E...	88	co. s.....	Rare.

QUINQUELOCULINA SCLEROTICA Karrer.

Quinqueloculina sclerotica KARRER, Sitz. Akad. Wiss. Wien, vol. 43, 1868, Abth. 1, pp. 111-193, pls. 1-5.

Miliolina sclerotica BALKWILL and MILLETT, Journ. Micr., vol. 3, 1884, p. 24, pl. 1, fig. 2.—SIDEBOTTOM, Mem. Proc. Lit. Philos. Soc. Manchester, pt. 1, 1904, p. 14.—HERON-ALLEN and EARLAND, Journ. Roy. Micr. Soc., 1911, p. 304; Proc. Roy. Irish Acad., vol. 31, pt. 64, p. 30; Trans. Zool. Soc. London, vol. 20, 1915, p. 577.

To this species are referred numerous specimens from the Philippine dredgings which occur in comparatively shallow water, and which are not the same as *Q. agglutinans* d'Orbigny. I am not sure that this material should be referred to *Q. sclerotica* Karrer. It has occurred at 14 stations, ranging in depth from 10 to 312 fathoms (18 to 571 meters), the average depth being 49 fathoms (90 meters). Bottom temperatures were given at two stations, 49.3° F. and 75.7° F. (9.6° C. and 24.2° C.); the warmer representing 37 fathoms (68 meters)

and the colder 312 fathoms (571 meters). Therefore, as this species occurs in shallow water, the average temperatures would probably be about the maximum given. Heron-Allen and Earland record it from the Kerimba Archipelago, "often abundant and presenting a considerable range of external form." The localities in the Philippine area are as follows: Off Jolo; off Romblon; off Tawi Tawi; between Samar and Leyte; off eastern Panay; off northwestern Panay; China Sea, off southern Luzon; off northern Cebu; Malam-paya Sound, Palawan Island; Sulu Archipelago; and between Marinduque and Luzon.

Quinqueloculina sclerotica—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bot- tom tem- pera- ture.	Character of bottom.	Abundance.
						° F.		
13594	U.S.N.M.	1	D5145...	6 04 30 N.; 120 59 30 E...	23	co. s., sh.	Rare.
13595	U.S.N.M.	2	D5154...	5 14 50 N.; 119 53 45 E...	12	co. s.	Few.
13671	U.S.N.M.	6	D5105...	14 17 15 N.; 120 32 40 E...	35	gy. s.	Few.
13592	U.S.N.M.	4	D5105...	14 43 55 N.; 120 12 50 E...	25	Few.
13593	U.S.N.M.	1	D5136...	6 04 20 N.; 120 59 20 E...	22	s., sh.	Rare.
13596	U.S.N.M.	6	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.	Few.
13597	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.	Rare.
13598	U.S.N.M.	10+	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fine s.	Common.
13599	U.S.N.M.	3	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Few.
13600	U.S.N.M.	10+	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m. ...	Abundant.
13604	U.S.N.M.	4	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob..	Few.
13601	U.S.N.M.	10+	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.	Common.
13602	U.S.N.M.	4	D5342...	10 56 55 N.; 119 17 24 E...	14-25	gy. m.	Few.
13603	U.S.N.M.	4	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh., g.	Few.

QUINQUELOCULINA FUSCA Brady.

Plate 84, figs. 6a-c.

Quinqueloculina fusca BRADY, Ann. Mag. Nat. Hist., 1870, p. 286, pl. 2, fig. 2.

Miliolina fusca BRADY, Journ. Roy. Micr. Soc., London, 1887, p. 883.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 576.

There are specimens which have the characteristic reddish color, reminding one of the color of many *Astrorhizidae* and *Lituolidae*. These occurred at six stations, ranging in depth from 20 to 159 fathoms (37 to 291 meters), the average depth being 92 fathoms (169 meters).

These stations include the following localities: Tara Island, off Romblon; China Sea, off Hongkong; Malampaya Sound, Palawan Island; off Marinduque Island; and to the southward in Buton Strait.

Heron-Allen and Earland record this species from the Kerimba Archipelago as "very sparingly distributed."

Specimens were frequent off Tara Island and common off Hongkong, but less common at the other stations.

Quinqueloculina fusca—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13613	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78	fine s.	Few.
13614	U.S.N.M.	6	D5311...	21 33 00 N.; 116 15 00 E...	88	crs. s., sh.	Common.
13615	U.S.N.M.	4	D5342...	10 56 55 N.; 119 17 24 E...	14-25	gy. m.	Frequent.
13616	U.S.N.M.	1	D5370...	13 44 15 N.; 121 42 30 E...	159	54.3	sft. m.	Rare.
13617	U.S.N.M.	1	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh.	Rare.
13618	U.S.N.M.	3	Tara Island, P. I.	Frequent.

QUINQUELOCULINA CRASSATINA (H. B. Brady).

Plate 84, figs. 4a-c.

Miliolina incrassata H. B. BRADY, Quart. Journ. Micr. Soc., vol. 21, 1881, p. 46.*Miliolina crassatina* H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 180, pl. 8, figs. 5a, b.

Brady described this species from off East Moncoeur Island, Bass Strait, in 138 fathoms (252 meters). It is evidently a rare species and possibly limited to the Pacific region.

Specimens very similar to that figured by Brady have occurred at two stations—D5178, off Romblon, in 78 fathoms (143 meters), and D5640, Buton Strait, in 24 fathoms (44 meters). Bottom temperatures are not given in either case.

Quinqueloculina crassatina—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13869	U.S.N.M.	2	D5178...	12 43 00 N.; 122 06 15 E...	78	fine s.	Rare.
13870	U.S.N.M.	1	D5640...	4 27 00 S.; 122 55 40 E...	24	s., brk. sh.	Rare.

QUINQUELOCULINA ALVEOLINIFORMIS (H. B. Brady).

Miliolina alveoliniformis H. B. BRADY, Quart. Journ. Micr. Soc., vol. 19, 1879; Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 181, pl. 8, figs. 15-20.—EGGER, Abh. kön. bay. Akad. Wiss. München, Cl. II, vol. 18, 1893, p. 232, pl. 2, figs. 17-19.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 510.—CHAPMAN, Journ. Linn. Soc. Zoology, vol. 28, 1901, p. 177; vol. 30, 1910, p. 398.—HERON-ALLEN and EARLAND, Trans. Zool. Soc., vol. 20, 1915, p. 581.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 43.

The records for this species are from the Indo-Pacific, mostly in shallow water. In the Philippines there are 10 records from stations ranging from 10 to 375 fathoms (18 to 686 meters), with the average 60 fathoms (110 meters). The two stations at which it was really common are in 10 and 20 fathoms (18 and 37 meters). The bottom temperature of the two stations given are 56.4° F. and 75° F. (13.5° C.

and 23.8° C.), but the lower of these is from the deepest station, 375 fathoms (686 meters), where the specimen was evidently accidental, and the next deepest one 37 fathoms (68 meters). All the other stations are in still shallower water, with consequent higher bottom temperatures.

The most of the stations are from the shallow water about the Sulu Sea, with a few other localities scattered about the Archipelago. It was not found in the region to the southward, largely on account of the deeper water in which the dredgings in that area were made.

Quinqueloculina alveoliniformis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance
				° ' " ° ' "		° F.		
13073	U.S.N.M.	3	Apra Bay, Guam.....	Few
13673	U.S.N.M.	3	D5134...	6 44 45 N.; 121 48 00 E..	25	fine s.....	Few.
13676	U.S.N.M.	3	D5154...	5 14 50 N.; 119 58 45 E..	12	co. s.....	Few.
13672	U.S.N.M.	10+	D5159...	5 11 50 N.; 119 54 00 E..	10	co. s.....	Abundant.
13674	U.S.N.M.	2	D5160...	5 12 40 N.; 119 55 10 E..	12	s.....	Few.
13675	U.S.N.M.	3	D5179...	12 38 15 N.; 122 12 39 E..	37	75.7	hrd. s.....	Few.
13677	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.....	Rare.
13678	U.S.N.M.	8	D5218...	13 11 15 N.; 123 02 45 E..	29	crs. s.....	Common.
13679	U.S.N.M.	3	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s.....	Few.
13689	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Few.
13681	U.S.N.M.	10+	Apra Bay, Guam.....	Abundant.
13682	U.S.N.M.	1	Jolo Jolo, P. I.....	Rare.

Genus MASSILINA Schlumberger, 1893.

This genus includes species in which the early development is quinqueloculine, the later development becoming very much like *Spiroloculina*.

To this genus are referred those species whose early development is typical quinqueloculine, some of which have been previously referred to *Spiroloculina*.

MASSILINA SECANS, var. RETICULATA Heron-Allen and Earland.

Massilina secans, var. *reticulata* HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 582, pl. 45, figs. 1-4.

Specimens which have the general characters of this variety described by Heron-Allen and Earland from the Kerimba Archipelago, were obtained at seven stations, ranging in depth from 10 to 88 fathoms (18 to 161 meters), the average being 30.1 fathoms (55 meters). Bottom temperatures were not given for one of these stations, but would probably be high in such shallow water.

The stations which include it are as follows: Sulu Archipelago, Tawi Tawi, Jolo, and off western Mindanao; China Sea, off southern Luzon; Ragay Gulf, Luzon; and Tacloban Anchorage.

Massilina secans, var. *reticulata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13739	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E..	10	° F.		Rare.
13740	U.S.N.M.	1	D5133...	Island off Panabutan Point, N. 52°; E. 1.50 miles	38		Co. gn. m., s.	Rare.
13741	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E..	19		s., Co.	Rare.
13738	U.S.N.M.	1	D5159...	5 11 50 N.; 119 54 00 E..	10		co. s.	Rare.
13736	U.S.N.M.	2	D5161...	5 10 15 N.; 119 53 00 E..	16		fine s., blk. sp.	Rare.
13737	U.S.N.M.	2	D5381...	13 14 15 N.; 122 44 45 E..	88		co. s.	Rare.
13743	U.S.N.M.	1	Tacloban Anchorage, P. I.			Rare.

MASSILINA SECANS, var. *tenuistriata* Heron-Allen and Earland.

Massilina secans, var. *tenuistriata* HERON-ALLEN and EARLAND, Trans. Zool. Soc., London, vol. 20, 1915, p. 582, pl. 44, figs. 28-31.

A few specimens from the *Albatross* station D5465, east coast of Luzon, in 500 fathoms (914 meters), probably represent this variety described by Heron-Allen and Earland from the Kerimba Archipelago.

Massilina secans, var. *tenuistriata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13742	U.S.N.M.	1	D5465...	13 39 42 N.; 123 49 39 E..	500	° F.	gy. m. (m. b.)	Rare.

MASSILINA CRENATA (Karrer).

Spiroloculina crenata KARRER, Sitz. Akad. Wiss. Wien, vol. 57, 1868, p. 135, pl. 1, fig. 9.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 156, pl. 10, figs. 24-26.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 265.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 557, pl. 41, fig. 68. *Massilina crenata* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 57, pl. 20, fig. 2.

Specimens are rare in the Philippine material, due perhaps to the comparative coarseness of many of the available samples.

It has occurred rarely in the following localities: Sulu Archipelago, off Jolo, off Tawi Tawi; two stations off Romblon; and the Pacific Ocean, off the east coast of Mindanao.

These stations range in depth from 23 to 249 fathoms (42 to 458 meters).

It is recorded from the Malay Archipelago by Millett, and from the Kerimba Archipelago by Heron-Allen and Earland. All but one of Brady's records are from the Indo-Pacific.

Heron-Allen and Earland figure specimens which, although larger than the spiroloculine form, do not show that stage, and suggest that these may be the microspheric form, which, growing larger, does not show the adult characters.

Massilina crenata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13727	U.S.N.M.	1	D5144...	6 04 30 N.; 120 59 30 E...	23	° F.	co. s., sh.....	Rare.
13728	U.S.N.M.	2	D5152...	5 22 55 N.; 120 15 45 E...	34	wh. s.....	Rare.
13729	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78	fne. s.....	Rare.
13730	U.S.N.M.	2	D5179...	12 38 15 N.; 122 12 30 E...	37	75.7	hrd. s.....	Rare.
13731	U.S.N.M.	1	D5237...	8 09 06 N.; 126 31 45 E...	249	46.4	gn. m.....	Rare.

MASSILINA DURRANDII Millett.

Plate 94, figs. 4a-c.

Miliolina durrandii MILLETT, Journ. Roy. Micr. Soc., 1898, p. 268, pl. 6, figs. 7-10.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 565, pl. 42, figs. 11-16.

This species was originally described by Millett from the Malay Archipelago, and is recorded by Heron-Allen and Earland from the Kerimba Archipelago, where it is "generally distributed and often quite common." They also record it from the coasts of Burmah, Queensland, Java, and Macassar, as well as from Tahiti. It therefore has a widespread tropical distribution in the Indo-Pacific in shallow water.

The few Philippine specimens are all of 50 fathoms (91 meters) of water or less, including the following stations: Tacloban Anchorage; China Sea, off southern Luzon; off eastern Panay; and between Marinduque and Luzon.

Massilina durrandii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13732	U.S.N.M.	1	D5100...	14 17 15 N.; 120 32 40 E...	35	° F.	gy. s.....	Rare.
13733	U.S.N.M.	1	D5181...	11 36 40 N.; 123 26 35 E...	26	m., fne. s.....	Rare.
13734	U.S.N.M.	1	D5220...	13 38 00 N.; 121 53 00 E...	50	sft. gn. m.....	Rare.
13735	U.S.N.M.	1	Tacloban Anchorage, P. I.	Rare.

MASSILINA ARENARIA (H. B. Brady).

Plate 94, figs. 3a-b.

Spiroloculina arenaria H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 153, pl. 8, fig. 12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 36.

Massilina arenaria DAKIN, Ceylon Pearl-Oyster Fisheries Suppl. Rep., pt. 5, 1906, p. 231.

From large series dredged at several stations, showing the young of this species, it is evident that it is a species of *Massilina*, instead of *Spiroloculina*. The finely granular condition of the surface is very

characteristic, obscuring the sutural lines. This is as characteristic of the early quinqueloculine stage as of the adult. One of Brady's original stations for this species was from the Philippines, in 95 fathoms (174 meters). The others were off Kandavu, Fiji Islands, in 210 fathoms (384 meters), and off Raine Island, Torres Strait, in 185 fathoms (338 meters). Dakin records this from off Ceylon.

At station D5236 there is a very large elongate form, which may possibly be the microspheric form of this species. In the Philippine material this species has occurred at 31 stations, ranging in depth from 14 to 569 fathoms (25 to 1,040 meters); bottom temperatures range from 40.1° F. to 76.3° F. (14.5° C. to 24.6° C.).

Besides the stations scattered through the Archipelago, it has occurred in the regions to the south in Sibuko Bay, Borneo, Gulf of Boni, and south of Patiente Strait.

Massilina arenaria—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14600	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E..	135	59.0	dk. gy. m....	Rare.
14601	U.S.N.M.	6	D5132...	Island off Panabutan Point, N.; 15° W., 0.3 mile.	26	gn. m., s.....	Common.
14602	U.S.N.M.	10	D5133...	Island off Panabutan Point, N.; 52° E., 1.5 miles.	38	gn. m., s.....	Common.
14603	U.S.N.M.	3	D5178...	12 43 00 N.; 122 06 15 E..	78	fine. s.....	Few.
14604	U.S.N.M.	10	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.....	Common.
14605	U.S.N.M.	10+	D5210...	11 49 55 N.; 124 28 05 E..	50	76.3	fine. gy. s....	Common.
14606	U.S.N.M.	7	D5213...	12 15 00 N.; 123 57 30 E..	80	s., m., sh....	Common.
14697	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.	Rare.
14608	U.S.N.M.	10+	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s....	Common.
14609	U.S.N.M.	1	D5247...	7 02 00 N.; 125 38 45 E..	135	m.	Rare.
14610	U.S.N.M.	1	D5261...	12 30 55 N.; 121 34 24 E..	145	s., m.	Rare.
14611	U.S.N.M.	10+	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.	Common.
14612	U.S.N.M.	2	D5269...	13 39 50 N.; 120 59 30 E..	220	fine. s., p....	Rare.
14613	U.S.N.M.	10	D5278...	14 00 10 N.; 120 17 15 E..	102	59.6	fine. s., m., sh.	Common.
14614	U.S.N.M.	1	D5281...	13 52 45 N.; 120 25 00 E..	201	50.4	dk. gy. s....	Rare.
14615	U.S.N.M.	2	D5282...	13 53 00 N.; 120 26 45 E..	248	47.4	dk. dy. s....	Rare.
14616	U.S.N.M.	6	D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Rare.
14617	U.S.N.M.	10+	D5338...	11 33 45 N.; 119 24 45 E..	43	fine. co. s., m.	Common.
14618	U.S.N.M.	9	D5342...	10 56 55 N.; 119 17 24 E..	14-25	gy. m.	Common.
14619	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Rare.
14620	U.S.N.M.	1	D5358...	6 06 40 N.; 118 18 15 E..	39	m.	Rare.
14621	U.S.N.M.	1	D5424...	9 37 05 N.; 121 12 37 E..	340	50.4	g., s.....	Rare.
14622	U.S.N.M.	2	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m. (net).	Rare.
14623	U.S.N.M.	1	D5548...	6 00 20 N.; 120 45 35 E..	232	53.5	s., brk., sh..	Rare.
14624	U.S.N.M.	6	D5549...	6 01 15 N.; 120 44 20 E..	263	52.3	s., glob., for.	Rare.
14625	U.S.N.M.	2	D5555...	5 41 42 N.; 120 30 30 E..	243	52.3	s., ptr., sh..	Rare.
14626	U.S.N.M.	1	D4585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.	Rare.
14627	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E..	347	43.0	gy. m.	Rare.
14628	U.S.N.M.	10+	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s....	Common.
14629	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m....	Rare.
14630	U.S.N.M.	5	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.	Common.

MASSILINA ASPERULA (Karrer).

Spiroloculina asperula KARRER, Sitz. Akad. Wiss. Wien, vol. 57, 1868, p. 136, pl. 1, fig. 10.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 152, pl. 8, figs. 13, 14 (11?).

Brady's records for this species include one station in the Philippines, in 95 fathoms (174 meters); also off the Admiralty Islands, 68

to 125 fathoms (125 to 229 meters), Ki Islands, 129 fathoms (236 meters), and Humboldt Bay, New Guinea, 37 fathoms (68 meters).

Specimens much like those figured by Brady occur at three Philippine stations—D5178, off Romblon, 78 fathoms (143 meters), D5220, between Marinduque and Luzon, 50 fathoms (91 meters), and D5348, Palawan Passage, 375 fathoms (686 meters).

These seem to be the smaller megalospheric form of this species.

Massilina asperula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13744	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	fine s.....	Rare.
13745	U.S.N.M.	1	D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.....	Rare.
			D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.....	Rare.

Genus ARTICULINA d'Orbigny, 1826.

ARTICULINA SAGRA d'Orbigny.

Articulina sagra D'ORBIGNY, in De La Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 160, pl. 9, figs. 23-26.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 184, pl. 12, figs. 22-24.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 585, pl. 45, figs. 22-25.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 59, pl. 22, figs. 7, 8.

The only record for this species in the Philippine material is *Albatross* station D5311, China Sea, off Hongkong, in 88 fathoms (161 meters). This consists of the usual early chambers, with a single nodosarian one.

It is recorded by Millett from the Malay region, and it seems strange that it is not more frequent in the Philippine dredgings.

Articulina sagra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13582	U.S.N.M.	1	D5311...	21 23 00 N.; 116 15 00 E..	88	crs. s., sh....	Rare.

Genus SIGMOILINA Schlumberger, 1887.

SIGMOILINA SIGMOIDEA (H. B. Brady).

Planispirina sigmoidea H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 197, pl. 2, figs. 1-3, woodcut fig. 5c.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 10.

Sigmoilina sigmoidea CHAPMAN, Biol. Res. *Endeavour*, vol. 3, pt. 1, 1915, p. 317.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 61, pl. 24, figs. 2, 3.

All the records for this species in the *Albatross* Philippine material are from stations in and about the Archipelago, none having oc-

curred in the region to the southward. At the 12 stations recorded the depths range from 218 to 554 fathoms (399 to 1,012 meters), with the average 372 fathoms (681 meters). Bottom temperatures of the nine stations given range from 41.2° F. to 56.4° F. (15.1° C. to 13.5° C.), the average being 50.4° F. (10.2° C.).

The localities include the following: Off Jolo; Sogod Bay, southern Leyte; east of Masbate; Pacific Ocean, east coast of Mindanao; China Sea, off southern Luzon; east coast of Luzon; Palawan Passage; northern Mindanao and vicinity; north of Tawi Tawi; and Sulu Sea, off western Mindanao.

Sigmoilina sigmoidea—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° / " ° / "		° F.		
14687	U. S. N. M.	1	D5172...	6 03 15 N.; 120 35 30 E...	318	fine s., sh....	Rare.
14688	U. S. N. M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Rare.
14689	U. S. N. M.	2	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m.....	Rare.
14690	U. S. N. M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine gy. s....	Few.
14691	U. S. N. M.	1	D5284...	13 42 05 N.; 120 30 45 E...	422	42.3	gy. m., glob..	Rare.
14692	U. S. N. M.	3	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	Co., s.....	Few.
14693	U. S. N. M.	1	D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Rare.
14694	U. S. N. M.	2	D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy. m., fine s.	Rare.
14695	U. S. N. M.	1	D5549...	6 01 15 N.; 120 44 20 E...	263	52.3	s., glob., f....	Rare.
14696	U. S. N. M.	1	D5570...	5 32 15 N.; 120 12 57 E...	330	52.3	fine s., glob..	Rare.
14697	U. S. N. M.	2	D5572...	5 31 26 N.; 120 09 45 E...	334	52.3	s.....	Few.
14698	U. S. N. M.	1	H4898...	7 43 45 N.; 122 03 45 E...	221	gy. m., glob..	Rare.

SIGMOILINA SCHLUMBERGERI Silvestri.

Planispirina celata H. B. BRADY (not Costa), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 197, pl. 8, figs. 1-4.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 10.

Sigmoilina celata CUSHMAN (not Costa), Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 61, pl. 24, fig. 1.

Sigmoilina schlumbergeri SILVESTRI, Mem. dell. Pont. Acc. Nuovi-Lincei, vol. 22, 1904, p. 267.—CHAPMAN, Biol. Res. *Endeavour*, vol. 3, pt. 1, 1915, p. 317.

This is one of the common species of the region, especially in water of some depth. It is recorded at 38 *Albatross* stations in the Archipelago and the region to the southward. The depths of the stations range from 32 to 1,570 fathoms (59 to 2,871 meters), with the average 425 fathoms (781 meters). Bottom temperatures range from 39.2° F. to 64.5° F. (4° C. to 18° C.), with the average 50.2° F. (10.1° C.).

The 31 stations in the Archipelago are well scattered and in addition it is found to the southward in Sibuko Bay, Borneo; Gulf of Tomini, Celebes; Pitt Passage; Molucca Sea; Gulf of Boni; Flores Sea; and Macassar Strait.

It is a widely distributed species at moderate depths.

Sigmoilina schlumbergeri—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14701	U. S. N. M.	2	D5120...	13 45 30 N.; 120 30 15 E...	393	43.7	gn. m., s.....	Few.
14702	U. S. N. M.	1	D5122...	13 21 30 N.; 120 30 33 E...	220	gn. m.....	Rare.
14703	U. S. N. M.	1	D5185...	10 05 45 N.; 122 18 30 E...	638	49.8	gn. m.....	Rare.
14704	U. S. N. M.	10+	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Common.
14705	U. S. N. M.	2	D5206...	11 31 40 N.; 124 42 40 E...	32	gn. m.....	Few.
14706	U. S. N. M.	1	D5211...	11 51 35 N.; 124 14 00 E...	155	56.6	gn. m., s.....	Rare.
14707	U. S. N. M.	2	D5214...	12 25 18 N.; 123 37 15 E...	218	51.4	gn. m.....	Frequent.
14708	U. S. N. M.	1	D5219...	13 21 00 N.; 122 18 45 E...	530	50.8	gn. m.....	Rare.
14709	U. S. N. M.	1	D5227...	12 53 45 N.; 121 52 30 E...	322	gn. m.....	Rare.
14710	U. S. N. M.	9	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s.....	Common.
14711	U. S. N. M.	10+	D5242...	6 51 53 N.; 126 14 10 E...	191	64.1	sft. gy. m.....	Common.
14712	U. S. N. M.	10+	D5243...	6 50 55 N.; 126 14 35 E...	218	63.6	gn. m.....	Common.
14713	U. S. N. M.	10+	D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob..	Common.
14714	U. S. N. M.	2	D5260...	12 25 35 N.; 121 31 35 E...	234	51.4	gn. m., s.....	Rare.
14715	U. S. N. M.	6	D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine. s.....	Common.
14716	U. S. N. M.	2	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co.....	Few.
14717	U. S. N. M.	1	D5349...	10 54 00 N.; 118 26 20 E...	730	40.6	Co., s.....	Few.
14718	U. S. N. M.	1	D5410...	10 28 45 N.; 124 05 30 E...	385	gn. m.....	Rare.
14719	U. S. N. M.	2	D5423...	9 38 30 N.; 121 11 00 E...	508	49.8	gy. m., co. s.....	Rare.
14720	U. S. N. M.	4	D5430...	9 49 40 N.; 119 03 20 E...	464	50.0	glob. oz.....	Few.
14721	U. S. N. M.	3	D5438...	15 54 42 N.; 119 44 42 E...	297	46.2	gn. m.....	Few.
14722	U. S. N. M.	7	D5445...	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.....	Frequent.
14723	U. S. N. M.	2	D5512...	8 16 02 N.; 123 58 26 E...	445	52.8	gy. m., fine. s.....	Few.
14724	U. S. N. M.	3	D5523...	8 48 44 N.; 127 23 35 E...	Few.
14725	U. S. N. M.	4	D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gy. m., glob..	Few.
14726	U. S. N. M.	2	D5537...	9 11 00 N.; 123 23 00 E...	254	53.5	gn. m.....	Few.
14727	U. S. N. M.	1	D5538...	9 09 15 N.; 123 23 20 E...	256	53.3	gn. m., s.....	Rare.
14728	U. S. N. M.	1	D5542...	8 48 30 N.; 123 35 30 E...	200	54.3	fine. s., brk. sh.	Rare.
14729	U. S. N. M.	2	D5543...	8 47 15 N.; 123 35 00 E...	162	64.5	s.....	Rare.
14730	U. S. N. M.	10+	D5590...	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.....	Common.
14731	U. S. N. M.	1	D5612...	0 38 00 S.; 121 45 40 E...	750	Rare.
14732	U. S. N. M.	1	D5636...	1 55 00 S.; 127 42 30 E...	1,262	gy. m., fine. s.	Rare.
14733	U. S. N. M.	2	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.....	Rare.
14734	U. S. N. M.	1	D5652...	4 53 00 S.; 121 23 06 E...	525	41.2	gn. m.....	Rare.
14735	U. S. N. M.	1	D5660...	5 36 30 S.; 120 49 00 E...	692	39.2	gy. m., s.....	Rare.
14736	U. S. N. M.	9	D5666...	2 54 30 S.; 118 47 00 E...	272	47.5	gn. m.....	Common
14699	U. S. N. M.	2	D4897...	7 46 00 S.; 122 00 00 E...	1,570	gy. m., glob..	Few.
14700	U. S. N. M.	1	H4895...	7 43 45 S.; 122 03 45 E...	221	gy. m., glob..	Rare.

Genus HAUERINA d'Orbigny, 1846.

HAUERINA BRADYI Cushman.

Hauerina compressa H. B. BRADY (not *H. compressa* d'Orbigny) Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 190, pl. 11, figs. 12, 13.—MILLET, Journ. Roy. Micr. Soc., 1898, p. 610, pl. 13, fig. 11.—RHUMBLER, Zool. Jahrb., Abt. Syst., vol. 24, 1906, p. 52, pl. 3, fig. 39.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 588.

Hauerina bradyi CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 62, pl. 23, fig. 2.

As far as the material shows this is a very rare species in the region.

Records of its occurrence are from but two stations, D5161, Sulu Archipelago, off Tawi Tawi, in 16 fathoms (29 meters), and D5218, between Burias and Luzon, in 20 fathoms (37 meters). The species was not common at either station.

The species is of the form which I have separated⁴⁴ from d'Orbigny's *H. compressa*. Heron-Allen and Earland have the species from the Kerimba region, and note that the form figured by Brady

⁴⁴ Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 62.

which I have separated is different from some of the other material. The Kerimba specimens seem to be different from those figured by Brady, according to Heron-Allen and Earland. However, they note that the specimens "have walls, all of such extreme tenuity that they are of a delicate blue, opalescent tint." This is also characteristic of the Philippine specimens, which are very delicately colored in a similar way.

The available records for this species range from the Hawaiian Islands to the coast of Africa and to the Mediterranean.

Hauerina bradyi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13579	U.S.N.M.	1	D5161...	° ' " ° ' "	16	*F.	fne.s., blk.sp.	Rare.
13580	U.S.N.M.	1	D5218...	5 10 15 N.; 119 53 00 E... 13 11 15 N.; 123 02 45 E...	20	crs. s.	Few.

HAUERINA FRAGILISSIMA (H. B. Brady).

Spiroloculina fragilissima H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 149, pl. 9, figs. 12-14.

Hauerina fragilissima MILLETT, Journ. Roy. Micr. Soc., 1898, p. 610, pl. 13, figs. 8-10.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 587, pl. 46, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 64, pl. 24, fig. 4.

Specimens were not common, being found at but two stations, D5181, off eastern Panay, in 26 fathoms (48 meters), and D5218, between Burias and Luzon, in 20 fathoms (37 meters).

The records for this species seem to be all from the Indo-Pacific region, being recorded by Brady from off Tahiti in 420 and 620 fathoms (768 and 1,134 meters); off Kandavu, Fiji Islands, in 225 fathoms (412 meters); and several stations off New Guinea in 3 to 25 fathoms (5 to 46 meters). He records it as "a rare coral-reef species." The other records are Millett's, from the Malay region, and my own, from the Hawaiian Islands. Dakin records a single specimen from off Ceylon. Heron-Allen and Earland report occasional specimens from the Kerimba region.

Hauerina fragilissima—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13576	U.S.N.M.	1	D5181...	° ' " ° ' "	26	*F.	m., fne. s.	Rare.
13577	U.S.N.M.	3	D5218...	11 36 40 N.; 123 26 36 E... 13 11 15 N.; 123 02 45 E...	20	crs. s.	Frequent.

HAUERINA ORNATISSIMA (Karrer).

Quinqueloculina ornatissima KARRER, Sitz. Akad. Wiss. Wien, vol. 58, 1868, p. 151, pl. 3, fig. 2.—H. B. BRADY, Quart. Journ. Micr. Sci., vol. 16, 1876, p. 406.

Hauerina ornatissima H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 192, pl. 7, figs. 15–22.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 610.—RHUMBLER, Zool. Jahrb., Abt. Syst., vol. 24, 1906, p. 53.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 590.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 63, pl. 23, figs. 1, 5.

This species has occurred in the material from but one station, D5178, off Point Origon, Romblon, in 78 fathoms (143 meters).

I can not explain its absence in so much material that should, it seems, be just the habitat for this tropical Indo-Pacific species.

Millett records it from the Malay region, Chapman has had it from Funafuti, Dakin records it from off Ceylon, and Heron-Allen and Earland record it from the Kerimba region.

Hauerina ornatissima—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13578	U.S.N.M.	1	D5178...	° ' " ° ' " 12 43 00 N.; 122 06 15 E...	78	° F.	fne. s.	Rare.

Genus TRILOCULINA d'Orbigny, 1826.**TRILOCULINA TRIGONULA (Lamarck).**

Miliolites trigonula LAMARCK, Ann. du Mus., vol. 5, 1804, p. 351, No. 3.

Triloculina trigonula D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 299, No. 1, pl. 16, figs. 5–9; Modèles, 1826, No. 93.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 65, pl. 25, fig. 3.

Miliolina trigonula WILLIAMSON, Recent Foraminifera of Great Britain, 1858, p. 83, pl. 7, figs. 180–182.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 164, pl. 3, figs. 14–16.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 561.

This species is not abundant in this area, compared to that of *T. tricarinata*.

There are records from but seven stations, ranging in depth from 12 to 318 fathoms (22 to 582 meters), the average depth being 98 fathoms (179 meters).

These include the following localities: Sulu Archipelago, off Siasi, off Jolo, off Tawi Tawi; between Bohol and Leyte; between Masbate and Leyte; between Burias and Luzon; and China Sea, off Hongkong.

Specimens are common at but one of these stations—D5218, between Burias and Luzon, in 20 fathoms (37 meters).

Triloculina trigonula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14680	U.S.N.M.	1	D5148...	5 35 45 N.; 120 47 30 E..	17	° F.	co. s.	Rare.
14685	U.S.N.M.	6	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.	Common.
14681	U.S.N.M.	1	D5230...	10 01 50 N.; 124 42 30 E..	118	57.6	gy. s.	Rare.
14684	U.S.N.M.	3	D5311...	21 33 00 N.; 116 15 00 E..	88	crs. s., sh.	Rare.
14682	U.S.N.M.	1	D5398...	11 35 12 N.; 124 13 48 E..	114	gn. m.	Rare.
14686	U.S.N.M.	1	D5154...	5 14 50 N.; 119 58 45 E..	12	co. s.	Rare.
14683	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E..	318	fine. s., sh.	Rare.

TRILOCULINA TRIGONULA (Lamarck), var. MULTISTRIATA, new variety.

Description.—Variety differing from the typical in having the surface of the test with many fine striae.

Distribution.—Type specimen (Cat. No. 13810, U.S.N.M.) from *Albatross* station D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters).

This form has occurred only at this station. It seems to represent a modified form of this species.

Triloculina trigonula, var. *multistriata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13810	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fine. gy. s.	Rare.

TRILOCULINA ADVENA, new species.

Description.—Test large, triangular, angles bluntly rounded, sides slightly inflated; sutures distinct, surface smooth; aperture very elongate, narrow, crescentiform, without a neck; tooth single, narrow, plate-like, nearly filling but not rising above the rim of the aperture.

Length of largest specimen about 2 mm.

Distribution.—Type specimen (Cat. No. 13796, U.S.N.M.) from *Albatross* station D5438, west coast of Luzon, in 297 fathoms (543 meters). Other records of this species are: D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters); D5331, off western Luzon, 178 fathoms (329 meters); and D5572, north of Tawi Tawi, in 334 fathoms (611 meters). Bottom temperatures of these stations range from 41.2° F. to 54.7° F. (5.1°C. to 12.5° C.).

This differs from *Triloculina trigonula* in its aperture, which is very elongate, and filled with a narrow thin lip. It is apparently

rather rare, being found at but four stations. The correlation of apertural characters and form of test with rounded angles is very constant.

Triloculina advena—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13797	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.....	Few.
13798	U.S.N.M.	1	D5331 ..	15 36 45 N.; 119 47 45 E..	178	54.7	s., sh., m.....	Rare.
13796	U.S.N.M.	3	D5438...	15 54 42 N.; 119 44 42 E..	297	46.2	gn. m.....	Few.
13799	U.S.N.M.	1	D5572...	5 31 26 N.; 120 09 45 E..	334	54.3	s.....	Rare.

TRILOCULINA TRICARINATA d'Orbigny.


Triloculina tricarinata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 299, No. 7.—

H. B. BRADY, Trans. Linn. Soc. London, vol. 24, 1864, p. 446, pl. 48, fig. 3.—

CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 66, pl. 25, figs. 1, 2.

Miliolina tricarinata H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 165, pl. 3, fig. 17a, b.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 7.

This species is typical and very common in the area. There are two distinct varieties, one elongate and very rare, and the other



FIGS. 35, 36.—TRILOCULINA TRICARINATA D'ORBIGNY. 35, APERTURAL VIEW OF BROAD, PECULIARLY SHAPED SPECIMEN. $\times 65$. FROM ALBATROSS STATION D5160. 36, APERTURAL VIEW OF A MUCH LARGER SPECIMEN WHERE THE OUTER FACE HAS A DOUBLE ANGLE. $\times 35$. FROM ALBATROSS STATION D5236.

shorter and very common. The sides are usually concave in the larger species and the aperture often cruciform. In the young there are two types of aperture—one slightly prolonged, rounded, with a single or usually bifid tooth, and more elongated form of test; the shorter test with the aperture flush with the surface and with no neck. The short variety, with rotund inflated sides but sharp keeled, may be known as var. *convexa*, new variety.

The records for the region include 50 stations, ranging in depth from 10 to 901 fathoms (18 to 1,648 meters), the average depth being 231 fathoms (423 meters). Bottom temperatures are given

for 25 stations; these range from 38.2° F. (3.4° C.) in the deepest water to 76.3° F. (24.6° C.) in shallow water, the average being 52.9° F. (11.6° C.). The stations are well scattered in the region of the Archipelago, and also occur in the region to the southward in Sibuko Bay, Borneo; Gulf of Boni; Macassar Strait; Molucca Passage; and Darvel Bay, Borneo.

Triloculina tricarinata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				* / " * / "		* F.		
14551	U.S.N.M.	3	D5096...	14 20 23 N., 120 34 15 E...	28	gy.m.,s.,sh..	Few.
14551	U.S.N.M.	1	D5100...	14 17 15 N., 120 32 40 E...	35	gy.s.....	Rare.
14552	U.S.N.M.	1	D5106...	14 23 55 N., 120 32 33 E...	37	gy.m.....	Rare.
14553	U.S.N.M.	1	D5112...	13 48 22 N., 120 47 25 E...	177	52.4	dk. gn. m....	Few.
14554	U.S.N.M.	5	D5126...	10 34 45 N., 121 47 30 E...	742	49.5	sft. gn. m....	Common (very large).
14555	U.S.N.M.	1	D5139...	6 06 00 N., 121 02 30 E...	20	co.s.....	Rare.
14556	U.S.N.M.	2	D5148...	5 35 40 N., 120 47 30 E...	17	co.s.....	Rare.
14557	U.S.N.M.	3	D5152...	5 22 55 N., 120 15 45 E...	34	wht.s.....	Common.
14558	U.S.N.M.	6	D5159...	5 11 50 N., 119 54 00 E...	10	co.s.....	Few.
14559	U.S.N.M.	1	D5160...	5 12 40 N., 119 55 10 E...	12	s.....	Rare.
14560	U.S.N.M.	4	D5164...	5 01 40 N., 119 52 20 E...	18	gn.m.....	Common.
14561	U.S.N.M.	1	D5172...	6 03 15 N., 120 35 30 E...	318	fne.s.,sh....	Rare.
14562	U.S.N.M.	8	D5178...	12 43 00 N., 122 06 15 E...	78	fne.s.....	Common.
14563	U.S.N.M.	8	D5179...	12 38 15 N., 122 12 30 E...	37	75.7	hrd.s.....	Common.
14564	U.S.N.M.	10+	D5181...	11 36 48 N., 123 26 35 E...	26	m.,fne.s....	Common.
14565	U.S.N.M.	2	D5182...	11 30 40 N., 123 23 20 E...	24	fne.s.,m....	Few.
			D5189...	9 56 30 N., 123 15 00 E...	300	62.8	gn.m.....	Common.
14566	U.S.N.M.	8	D5192...	11 09 15 N., 123 50 00 E...	32	gn.s.....	Few.
14567	U.S.N.M.	1	D5201...	10 10 00 N., 125 04 15 E...	554	52.3	gy.s.,m....	Rare.
14568	U.S.N.M.	6	D5206...	11 31 40 N., 124 42 40 E...	32	gn.m.....	Common.
14569	U.S.N.M.	2	D5210...	11 49 55 N., 124 28 05 E...	50	76.3	fne.gy.s....	Rare.
14570	U.S.N.M.	2	D5211...	11 51 35 N., 124 14 00 E...	155	56.6	gn.m.,s....	Rare.
14571	U.S.N.M.	1	D5212...	12 04 15 N., 124 04 36 E...	108	59.9	gy.s.,m....	Rare.
14572	U.S.N.M.	2	D5214...	12 25 18 N., 123 37 15 E...	218	51.4	gn.m.....	Rare.
14573	U.S.N.M.	6	D5217...	13 20 00 N., 123 14 15 E...	105	63.1	crs.gy.s....	Few.
14574	U.S.N.M.	10+	D5236...	8 50 45 N., 126 26 52 E...	494	41.2	fne.gy.s....	Common (very large).
14575	U.S.N.M.	1	D5259...	11 57 30 N., 121 42 15 E...	312	49.3	gy.m.,glob..	Few.
14576	U.S.N.M.	5	D5268...	13 42 00 N., 120 57 15 E...	170	s.,p.....	Few.
14577	U.S.N.M.	1	D5276...	13 49 15 N., 120 14 45 E...	18	sh.,p.,s....	Rare.
14578	U.S.N.M.	10+	D5284...	13 42 05 N., 120 30 45 E...	422	42.3	gy.m.,glob..	Common.
14579	U.S.N.M.	2	D5311...	21 33 00 N., 116 15 00 E...	88	crs.s.,sh....	Rare.
14580	U.S.N.M.	1	D5315...	21 40 00 N., 116 58 00 E...	148	54.4	s.,sh.....	Rare.
14581	U.S.N.M.	2	D5338...	11 33 45 N., 119 24 45 E...	43	Co.,s.,m....	Rare.
14582	U.S.N.M.	6	D5342...	10 56 55 N., 119 17 24 E...	14-25	gy.m.....	Common.
14583	U.S.N.M.	2	D5348...	10 57 45 N., 118 38 15 E...	375	56.4	Co.,s.....	Few.
14584	U.S.N.M.	1	D5358...	6 06 40 N., 118 18 15 E...	39	m.....	Rare.
14585	U.S.N.M.	3	D5429...	9 41 30 N., 118 50 22 E...	766	gn.m.....	Common (very large).
14586	U.S.N.M.	1	D5469...	13 36 48 N., 123 38 24 E...	500	gn.m.(net)..	Rare.
14587	U.S.N.M.	6	D5481...	10 27 30 N., 125 17 10 E...	61	s.,sh.,g....	Common.
14588	U.S.N.M.	3	D5512...	8 16 02 N., 123 58 26 E...	445	52.8	gy.m.,fne.s..	Common.
14589	U.S.N.M.	1	D5548...	6 00 20 N., 120 45 35 E...	232	53.5	g.,brk.sh....	Rare.
14590	U.S.N.M.	2	D5567...	5 48 00 N., 120 33 45 E...	268	52.0	fne.s.....	Few.
14591	U.S.N.M.	1	D5370...	5 32 15 N., 120 12 57 E...	330	52.3	fne.s.,glob..	Rare.
14592	U.S.N.M.	1	D5376...	5 25 56 N., 120 03 39 E...	277	53.3	s.....	Rare.
14593	U.S.N.M.	1	D5380...	4 52 45 N., 119 06 45 E...	162	55.8	br.s.,Co....	Rare.
14594	U.S.N.M.	1	D5582...	4 19 54 N., 118 58 38 E...	890	38.3	gy.m.,fne.s..	Rare.
14595	U.S.N.M.	2	D5590...	4 10 50 N., 118 39 35 E...	310	44.3	gn.m.,s....	Few.
14596	U.S.N.M.	1	D5618...	0 37 00 N., 127 15 00 E...	417	gy.m.....	Rare.
14597	U.S.N.M.	5	D5650...	4 53 45 S., 121 29 00 E...	540	40.1	gn.m.....	Frequent.
14598	U.S.N.M.	2	D5668...	2 28 15 S., 118 49 00 E...	901	38.2	gy.m.....	Few.
14599	U.S.N.M.	1		Tara Island, P. I.....	Rare.

TRILOCULINA TRICARINATA d'Orbigny, var. CONVEXA, new variety.

Description.—Variety differing from the typical in the shorter test and the convex sides.



FIG. 37.—TRILOCULINA TRICARINATA D'ORBIGNY. APERTURAL VIEW, $\times 50$. FROM ALBATROSS STATION D5236.

This is constant in its characters at several stations as follows: D5133, Sulu Sea off western Mindanao; D5144, vicinity of Jolo, D5145, the neighboring station; D5268, Verde Island Passage; D5277, China Sea, off southern Luzon; and D5576, north of Tawi Tawi.

With two exceptions, D5268 and D5576, the stations are in less than 100 fathoms (183 meters), ranging from 19 to 80 fathoms (35 to 146 meters). The bottom temperature was recorded in two cases only, D5277, 58.6° F. (14.7° C.), and D5576, 53.5° F. (11.8° C.).

Triloculina tricarinata, var. *convexa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13802	U.S.N.M.	3	D5133...	Island off Panabutan Pt. N. 52°; E., 1.50 miles.	38	° F.	gn. m., s.	Few.
13803	U.S.N.M.	2	D5144...	6 05 50 N.; 121 02 15 E.	19	co. s.	Few.
13804	U.S.N.M.	1	D5145...	6 04 30 N.; 120 59 30 E.	23	co. s., sh.	Rare.
13805	U.S.N.M.	2	D5268...	13 42 00 N.; 120 57 15 E.	170	s., p.	Few.
13806	U.S.N.M.	3	D5277...	13 56 55 N.; 120 13 45 E.	80	58.6	fine s.	Few.
13807	U.S.N.M.	1	D5576...	5 25 56 N.; 120 03 29 E.	277	53.3	s.	Few.

TRILOCULINA TRICARINATA, var. PLICATA Terquem.

Triloculina plicata TERQUEM, Mém. Soc. géol. France, ser. 3, vol. 1, p. 61, pl. 6 (11), fig. 2.

Miliolina tricarinata, var. *plicata* HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 562, pl. 41, figs. 17-22.

There are rare specimens at two stations—D5160, Sulu Archipelago, off Tawi Tawi, 12 fathoms (22 meters), and D5236, Pacific Ocean, off the east coast of Mindanao, 494 fathoms (903 meters), which can be referred to this species.

As figured by Terquem and Heron-Allen and Earland, this is rather short, chambers with very decided extended edges, and the chambers often separated by deep clefts.

Terquem's material was from the Pliocene, of the Isle of Rhodes; Heron-Allen and Earland's from the Kerimba Archipelago.

Triloculina tricarinata, var. *plicata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13809	U.S.N.M.	1	D5160...	5 12 40 N.; 119° 55 10' E....	12	s.....	Rare.
13808	U.S.N.M.	1	D5236...	8 50' 45 N.; 126° 26 52 E....	494	41.2	fine. gy. s....	Rare.

TRILOCULINA BERTHELINIANA (H. B. Brady).

Miliolina bertheliniana H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 166, pl. 114, fig. 2.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 563, pl. 41, figs. 32-35.

Miliolina tricarinata, reticulated variety, MILLETT, Journ. Roy. Micr. Soc., 1898, p. 503, pl. 11, fig. 12.

Miliolina tricarinata, var. *bertheliniana* CHAPMAN, Journ. Linn. Soc. London (Zoology), vol. 28, 1902, p. 174.

Heron-Allen and Earland, in their Kerimba paper, record this species from Cebu in 45 to 120 fathoms (82 to 220 meters) and note that it "is probably of wide distribution in tropical shallow seas."

Brady describes the species from the following localities: Off Ascension Island, 7 fathoms (13 meters); off Calpentyn, Ceylon, 2 fathoms (4 meters); and in shore sands from near Tamatavé, Madagascar; and near Port Elizabeth, Algoa Bay.

Millett records it from the Malay region as very rare, having but a single specimen from but one station.

Chapman records it from Funafuti and Heron-Allen and Earland from the Kerimba Archipelago.

All the Philippine material is of the short form which Heron-Allen and Earland mention in their paper, none of the longer form being present.

There are also specimens from the nine stations which have the quinqueloculine form and very thin reticulate ornamentation, like the triloculine forms, and may be simply an early quinqueloculine stage.

This species ranges in depth from very shallow water to 318 fathoms (582 meters), representing the following localities: Binang Pool, Subin Bay; Tara Island; Sulu region, off Jolo, off Tawi Tawi, near Basilan; China Sea, off southern Luzon; and between Burias and Luzon.

Triloculina bertheliniana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13793	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E....	135	59	drk. gy. m....	Rare.
13791	U.S.N.M.	1	D5134...	6 44 45 N.; 121 48 00 E....	25	fine. s.....	Rare.
13748	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E....	19	s., Co.....	Rare.
13789	U.S.N.M.	1	D5153...	5 18 10 N.; 120 02 55 E....	49	co. s., sh....	Rare.
13749	U.S.N.M.	1	D5162...	5 10 00 N.; 119 47 30 E....	230	52.9	rs. s. brk. sh.	Rare.
13750	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E....	318	fine. s., sh....	Rare.
13790	U.S.N.M.	3	D5218...	13 11 15 N.; 123 02 45 E....	20	crs. s.....	Rare.
13751	U.S.N.M.	1	Binang Pool, Subin Bay, P. I.	Rare.
13752	U.S.N.M.	1	Tara Island, P. I.	Rare.

TRILOCULINA TERQUEMIANA (H. B. Brady).

Miliolina terquemiana H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 166, pl. 114, figs. 1a, b.—DAKIN, Ceylon Pearl-Oyster Fisheries Suppl. Rep., pt. 5, 1906, p. 230, pl. figs. 9, 10.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 563, pl. 41, figs. 29-31.

Triloculina terquemiana CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 72, pl. 27, fig. 2.

Miliolina tricarinata, striate variety, MILLETT, Journ. Roy. Micr. Soc., 1898, p. 503, pl. 11, figs. 10, 11.

Miliolina tricarinata, var. *terquemiana* CHAPMAN, Journ. Linn. Soc. Zool., vol. 23, 1900, p. 174.

The only specimens which can be referred to this species are like the inflated form figured by Heron-Allen and Earland from the Kerimba Archipelago. From the records this species seems to be decidedly Indo-Pacific in its range.

The only station at which it occurred is D5154, Sulu Archipelago, Tawi Tawi Group, in 12 fathoms (22 meters).

Triloculina terquemiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15791	U.S.N.M.	1	D5154...	5 14 50 N.; 119 58 45 E....	12	° F.	co. s.	Rare.

TRILOCULINA INSIGNIS (H. B. Brady).

Miliolina insignis H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 45; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 165, pl. 4, figs. 8, 10.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 562.

Triloculina insignis CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 72, pl. 27, fig. 3.

There are rare specimens from *Albatross* station D5141, off Jolo, 29 fathoms (52 meters), and D5236, Pacific Ocean, east coast of Mindanao, 494 fathoms (903 meters), which may be referred to this species.

There seems to be a mixture of forms in the material described by Brady. In the *Challenger* Report (pl. 4, fig. 8) is evidently a triloculine form of *Biloculina comata* Brady, as shown by the form of the aperture. Some such specimens are often found where *B. comata* is abundant. Brady has a single North Pacific record from deep water for this species. This may have been the form of *B. comata* which he figures. Flint⁴⁵ figures a form (pl. 45, fig. 2) and has it from the same station from which he reports *B. comata*, and the two probably represent the same species.

⁴⁵ Ann. Rep. U. S. Nat. Mus., 1897 (1899).

Returning to the *Challenger* Report (pl. 4, fig. 10) refers to a different species and may be taken as the type of *T. insignis*. This is a striate form of the *T. trigonula* group, and apparently found in shallow water tropical regions. It may be suggested that the form figured by Heron-Allen and Earland from Kerimba (pl. 41, figs. 29, 31) is the inflated form of *T. terquemiana*, and may really be this species, which is very similar to *Challenger* (pl. 4, fig. 10), noted above.

Triloculina insignis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13825	U.S.N.M.	1	D5141...	• • " • • "	29	*F.	co. s.....	Rare.
13826	U.S.N.M.	1	D5236...	6 09 00 N.; 120 58 00 E..	494	41.2	fine. gy. s....	Rare.
15777	U.S.N.M.	1	D5425...	8 50 45 N.; 126 26 52 E..	495	49.4	gy. m., co. s.	Rare.

TRILOCULINA OBLONGA (Montagu).

Plate 92, figs. 3a-c.

Vermiculum oblongum MONTAGU, Test. Brit., 1903, p. 522, pl. 14, fig. 9.

Triloculina oblonga D'ORBIGNY, Ann. Sci. Mat., vol. 7, 1826, p. 300, No. 16.—

CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 69, pl. 26, fig. 3.

Miliolina oblonga H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 160, pl. 5, figs. 4a, b.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 566.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 7, pl. 1, figs. 15, 16.

This usually common species occurs at but five stations, and is not common at any of them. These range in depth from 61 to 494 fathoms (112 to 903 meters); bottom temperatures range from 41.2° F. to 56.4° F. (5.1° C. to 13.5° C.).

These few stations are east of Masbate Island; Pacific Ocean, east coast of Mindanao; Palawan Passage; between Samar and Leyte; and off Jolo.

Triloculina oblonga—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13824	U.S.N.M.	1	D5214...	• • " • • "	218	*F.	gn. m.....	Rare.
			D5236...	12 25 18 N.; 123 37 15 E..	494	51.4	fine. gy. s....	Rare.
			D5348...	8 50 45 N.; 126 26 52 E..	375	56.4	co. s.....	Rare.
13822	U.S.N.M.	1	D5481...	10 57 00 N.; 118 38 15 E..	61	s., sh., g.....	Rare.
13823	U.S.N.M.	4	D5549...	10 27 30 N.; 125 17 10 E..	263	52.3	s., glob., for..	Few.

TRILOCULINA ROTUNDA d'Orbigny.

Triloculina rotunda D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 299, No. 4.—SCHLUMBERGER, Mém. Soc. Zool. France, vol. 6, 1893, p. 64, pl. 1, figs. 48-50; figs. 11, 12 (in text).

Miliolina rotunda MILLETT, Journ. Roy. Micr. Soc., 1898, p. 267, pl. 5, figs. 15, 16.—SIDEBOTTOM, Mem. Proc. Manchester Lit. and Philos. Soc., vol. 48, No. 5, 1905, p. 8.—HERON-ALLEN and EARLAND, Proc. Roy. Irish Acad., vol. 31, pt. 64, 1913, p. 25.—WIESNER, Archiv. Prot., vol. 25, 1912, p. 225.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 568, pl. 42, figs. 27-30.

I have records of this species from six stations, ranging in depth from 10 to 569 fathoms (18 to 1,040 meters), the average depth 36 fathoms. Bottom temperatures are given at but two stations, 41.2° F. (5.1° C.) and 56.4° F. (13.5° C.). Of these stations the one at 10 fathoms (18 meters) is the only one in shallow water, the next deepest station being in 298 fathoms (545 meters).

The localities are China Sea, off southern Luzon; Pacific Ocean, east coast of Mindanao; Palawan Passage; and to the southward, between Gillolo and Makyan Island; south of Patiente Strait; and Molucca Passage.

Triloculina rotundata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13316	U.S.N.M.	1	D5109...	14 03 45 N.; 120 16 30 E...	10	Co.	Rare.
13317	U.S.N.M.	4	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Few.
13319	U.S.N.M.	2	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co. s.	Few.
13320	U.S.N.M.	1	D5318...	0 37 00 N.; 127 15 00 E...	417	gy. m.	Rare.
13318	U.S.N.M.	2	D5621...	0 15 00 N.; 127 24 35 E...	298	gy. and bk. s. (m. b.)	Few.
13321	U.S.N.M.	1	D5630...	0 56 30 S.; 128 05 00 E...	569	co. s., m....	Rare.

TRILOCULINA VALVULARIS Reuss.

Triloculina valvularis REUSS, Zeitsch. deutsch. geol. Ges., vol. 3, 1851, p. 85, pl. 7, fig. 56.

Miliolina valvularis H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 161, pl. 4, figs. 4, 5.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 501, pl. 11, figs. 5-7.—FLINT, Rep. U. S. Nat. Mus., 1897 (1899), p. 290, pl. 44, fig. 4.—CHAPMAN, Journ. Linn. Soc., vol. 28, 1900, p. 172.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 599.

There are only two stations for this species—D5333, Mindoro Strait, in 310 fathoms (567 meters), where it is very common, and D5650, Gulf of Boni, 540 fathoms (988 meters), where it is rare.

This species is also found off New Zealand, and may be different from that of Reuss.

Heron-Allen and Earland record it from Kerimba, and note that their material was very variable, but that the triloculine form was most abundant.

Triloculina valvularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13800	U.S.N.M.	10	D5333...	° ' " ° ' "	310	° F. 73.8	s.....	Common.
13801	U.S.N.M.	1	D5650...	12 26 30 N.; 120 37 45 E.. 4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Rare.

TRILOCULINA SUBORBICULARIS d'Orbigny.

Miliolina suborbicularis HERON-ALLEN and EARLAND, Journ. Roy. Micr. Soc., 1911, p. 304; Trans. Zool. Soc. London, vol. 20, 1915, p. 560.

Quinqueloculina suborbicularis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 302, No. 29.—SCHLUMBERGER, Mém. Soc. Zool., France, 1893, p. 215, text figs. 26–28; pl. 2, figs. 63–64; pl. 3, fig. 67.

Triloculina suborbicularis D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 300, No. 12.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 70, pl. 21, fig. 3.

There are single, very typical, specimens from three stations—D5154, Tawi Tawi Group, Sulu Archipelago; D5162, Tawi Tawi Group, Sulu Archipelago; and D5179, off Romblon.

In these specimens the two last-formed chambers are almost opposite one another, and they have the fine striations characteristic of this species.

Triloculina suborbicularis—Material examined.

Cat. No.	Coll. of—	No. of specimens	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13854	U.S.N.M.	1	D5154...	° ' " ° ' "	12	° F.	co. s.....	Rare.
13855	U.S.N.M.	1	D5179...	5 14 50 N.; 119 58 45 E.. 5 10 00 N.; 119 47 30 E..	230	52.9	crs. s., bk. sh.	Rare.
13856	U.S.N.M.	1	D5162...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Rare.

TRILOCULINA CUNEATA Karrer.

Plate 92, figs. 4a-c.

Triloculina cuneata KARRER, Sitz. Akad. Wiss. Wien, vol. 55, Abteil. 1, 1867, p. 359, pl. 2, fig. 8.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 71, pl. 17, fig. 1.

Miliolina cuneata RHUMBLER, Zool. Jahrb., Abteil. Syst., vol. 24, 1907, p. 42, pl. 3, fig. 22.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 569.

There are single specimens from five stations. These are China Sea, off southern Luzon; between Burias and Luzon; between Samar and Leyte; and in shallow water off Jolo Jolo.

All these stations are in shallow water, ranging in depth from 18 to 61 fathoms (33 to 112 meters).

Rhumbler records it from off Laysan Island. Many of the specimens, according to variable information, are biloculine in character, showing that this is a species which is tending to become possibly more related to *Biloculina* than *Triloculina* in its later characters.

Triloculina cuneata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13812	U.S.N.M.	1	D5096...	14 20 23 N.; 120 34 15 E...	28	gy. m., s., sh.	Rare.
13813	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E...	20	crs. s.	Rare.
13814	U.S.N.M.	1	D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.	Rare.
13815	U.S.N.M.	1	D5481...	10 27 30 N.; 125 17 10 E...	61	s., sh., g.	Rare.
13811	U.S.N.M.	1	Exterior pearl oyster, Jolo Jolo, P. I.	Rare.

TRILOCULINA CUNEATA Karrer, var. INCISA, new variety.

Plate 93, figs. 1a-c.

Description.—Test differing from the typical largely in the character of the surface, which is ornamented by numerous short, sharply incised markings lengthwise of the test.

Specimen from shallow water from Siasi Island, Philippines.

Triloculina cuneata, var. *incisa*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13795	U.S.N.M.	1	Siasi Id, P. I.	Shallow water.	Rare.

TRILOCULINA CIRCULARIS Bornemann.

Plate 92, figs. 1, 2.

Triloculina circularis BORNEMANN, Zeitschr. deutsch. geol. Ges., vol. 7, 1855, p. 349, pl. 19, fig. 4.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 67, pl. 25, fig. 4; pl. 26, fig. 1.

Miliolina circularis H. B. BRADY, Rep. Voy. Challenger, Zoology vol. 9, 1884, p. 169, pl. 4, figs. 3 a-c; pl. 5, figs. 13, 14?—CHAPMAN, Trans. New Zealand Inst., vol. 38, 1905, p. 81.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 557.—SIDEBOTTOM, Journ. Roy. Micr. Soc., 1918, p. 8.

There is very typical material from 15 stations in the area, ranging in depth from 20 to 554 fathoms (37 to 1,012 meters), the average depth being 209 fathoms (383 meters); bottom temperature is given at eight stations, ranging from 52.3° F. to 75.7° F. (11.2° C. to 24.2° C.), the average temperature 58° F. (14.4° C.).

The specimens are of the form described by Bornemann, and are similar to those found in other parts of the Indo-Pacific region.

Triloculina circularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13774	U.S.N.M.	1	D5139...	6 06 00 N.; 121 02 30 E..	20	co. s.....	Rare.
13775	U.S.N.M.	1	D5142...	6 06 10 N.; 121 02 40 E..	21	co. s., sh.....	Rare.
13776	U.S.N.M.	1	D5153...	5 18 10 N.; 120 02 55 E..	49	co. s., sh.....	Rare.
13777	U.S.N.M.	3	D5178...	12 43 99 N.; 122 06 15 E..	78	fine. s.....	Rare.
13778	U.S.N.M.	7	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Common.
13779	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	Rare.
13780	U.S.N.M.	1	D5230...	10 01 50 N.; 124 42 30 E..	118	57.6	gy. s.....	Rare.
13781	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.(net)..	Rare.
13782	U.S.N.M.	6	D5546...	6 06 48 N.; 121 20 32 E..	138	58.3	fine. co. s.....	Frequent.
13783	U.S.N.M.	2	D5549...	6 01 15 N.; 120 44 20 E..	263	62.3	s., glob. for..	Few.
13784	U.S.N.M.	2	D5572...	5 31 26 N.; 120 69 45 E..	334	52.3	Rare.
13785	U.S.N.M.	1	D5575...	5 28 30 N.; 120 02 27 E..	315	52.3	Co., s.....	Rare.
13786	U.S.N.M.	3	D5576...	5 25 56 N.; 120 03 39 E..	277	53.3	s.....	Few.
13788	U.S.N.M.	1	Exterior pearl oyster, Jolo Jolo, P. I.	Rare.
13787	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob..	Rare.

TRILOCULINA LINNEIANA d'Orbigny.

Triloculina linneiana d'ORBIGNY, in De La Sagra, Hist. Fis. Pol. Nat. Cuba, 1839, "Foraminifères," p. 153, pl. 9, figs. 11, 13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 72, pl. 27, fig. 4.

Miliolina linnaeana H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 174, pl. 6, figs. 15–20.

This species, as might be expected, is very typical and finely developed at many stations, the most abundant and finest specimens being found in comparatively shallow water.

Of the 21 stations at which it is recorded, ranging in depth from 10 to 500 fathoms (18 to 914 meters), the average depth is 166 fathoms (303 meters). It is most numerous and best developed about the islands of the Sulu Archipelago in shallow water, usually under 20 fathoms (37 meters). It has also occurred in Verde Island Passage; China Sea, off southern Luzon; east of Masbate; between Burias and Luzon; and off Romblon.

Triloculina linneiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13753	U.S.N.M.	8	D5137...	6 04 25 N.; 120 58 30 E..	20	*F.	s., sh.....	Common.
13754	U.S.N.M.	7	D5138...	6 06 00 N.; 128 58 50 E..	19	s., Co.....	Common.
13755	U.S.N.M.	3	D5139...	6 06 00 N.; 121 02 30 E..	20	co. s.....	Few.
13757	U.S.N.M.	2	D5141...	6 09 00 N.; 120 58 00 E..	29	co. s.....	Few.
13758	U.S.N.M.	6	D5142...	6 06 10 N.; 121 02 40 E..	21	co. s., sh.....	Few.
13759	U.S.N.M.	10+	D5143...	6 05 50 N.; 121 02 15 E..	19	co. s.....	Common.
13756	U.S.N.M.	3	D5144...	6 05 50 N.; 121 02 15 E..	19	co. s.....	Rare.
13760	U.S.N.M.	10+	D5148...	5 35 40 N.; 120 47 30 E..	17	co. s.....	Common.
13761	U.S.N.M.	1	D5149...	5 33 00 N.; 120 42 10 E..	10	Co., sh.....	Few.
13762	U.S.N.M.	10	D5151...	5 24 40 N.; 120 27 15 E..	24	s., sh.....	Common.
13763	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E..	34	wh. t. s.....	Rare.
13764	U.S.N.M.	6	D5154...	5 14 50 N.; 119 58 45 E..	12	co. s.....	Common.
13767	U.S.N.M.	2	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Few.
13770	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.....	Rare.
13771	U.S.N.M.	1	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s.....	Rare.
13768	U.S.N.M.	1	D5212...	12 04 15 N.; 124 04 36 E..	108	59.9	gy. s., m.....	Rare.
13769	U.S.N.M.	6	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Common.
13766	U.S.N.M.	3	D5162...	5 10 00 N.; 119 47 30 E..	230	52.9	crs. s., brk. sh.	Few.
13765	U.S.N.M.	1	D5159...	5 11 50 N.; 119 54 00 E..	10	co. s.....	Rare.
13772	U.S.N.M.	1	D5469...	13 36 48 N.; 123 38 24 E..	500	gn. m.(net)..	Rare.
13773	U.S.N.M.	1	D5571...	5 30 45 N.; 120 07 57 E..	340	52.3	s., sh.....	Rare.

TRILOCULINA RUPERTIANA (H. B. Brady).

Plate 93, figs. 2a-c.

Miliolina rupertiana H. B. BRADY, Quart. Journ. Micr. Sci., vol. 19, 1879, p. 46; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 178, pl. 7, figs. 7-12.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 269, pl. 6, fig. 13.—DAKIN, Ceylon Pearl-Oyster Fisheries, Suppl. Rep., pt. 5, 1906, p. 230.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 565.

The only material of this species is a very fine large specimen, 3 mm. in length, from the station D5134, Sulu Archipelago, near Basilan Island, in 25 fathoms (46 meters). Millett records his material from the Malay region at several stations, but in few numbers. The *Challenger* records include several stations about the islands on the south shores of New Guinea, and west of Torres Strait in 6 to 28 fathoms (11 to 51 meters), "and even here the species is very rare."

Other records given by Brady are the northwest coast of Ceylon, 2 fathoms (4 meters), north of Suez, 15 to 20 fathoms (27 to 37 meters), and in shore sands, near Tamatavé, on the east coast of Madagascar. At the last station it was plentiful.

Heron-Allen and Earland record it from the Kerimba Archipelago, and note that it is "extremely common at Perim in the Red Sea," and that it also occurs on the Queensland coast. Dakin gives it as "frequent off Ceylon."

From the records it is evident the species is more characteristic of the Indian Ocean and the coast of Africa than in the Pacific part of this region, and is evidently most common in very shallow water. It resembles in its highly specialized ornamentation certain of the Eocene species of western Europe and North America.

Triloculina rupertiana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13794	U.S.N.M.	1	D5134...	6 44 45 N.; 121 48 00 E...	25	°F.	fine s.....	Rare.

TRILOCULINA TORTUOSA, new species.

Quinqueloculina agglutinans CUSHMAN (not d'Orbigny), Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 42, pl. 9, fig. 1.

Description.—Test triloculine, in the adult condition elongate, slender, the chambers somewhat twisted; walls made of coarse fragments on the exterior; peripheral end produced into a cylindrical neck, usually smooth; aperture circular; lip thin, slightly expanded, with a slender bifid tooth.

Length 1 to 1.1 mm.

Distribution.—Type specimen (Cat. No. 13622, U.S.N.M.) from *Abatross* station D5192, off northern Cebu, in 32 fathoms (59 meters).

Other localities represent 7 stations, ranging in depth from 18 to 80 fathoms (33 to 146 meters), the average depth being 34 fathoms (62 meters). Bottom temperatures are given at 2 stations, ranging from 58.6° F. to 75.7° F. (14.7° C. to 24.2° C.). These localities are China Sea, off southern Luzon; off Romblon; off eastern Panay; and between Burias and Luzon.

This species is a peculiar, and, in the Philippine material, a very distinctive one. The peculiar twisted character, long slender form, with the extended smooth neck, and especially very coarse material of the test, all make it conspicuous among the other arenaceous species that occur with it. It is found in comparatively shallow water.

Triloculina tortuosa—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13619	U.S.N.M.	2	D5096...	14 20 23 N.; 120 34 15 E..	28	gy. m., sh., s.	Rare.
13620	U.S.N.M.	1	D5097...	14 19 15 N.; 120 33 52 E..	30	gy. m., s., sh.	Few.
13621	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.....	Rare.
13622	U.S.N.M.	6	D5181...	11 36 40 N.; 123 26 35 E..	26	m., fine s.....	Common.
13622	U.S.N.M.	5	D5192...	11 09 15 N.; 123 50 00 E..	32	gn. s.....	Common.
13623	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Rare.
13624	U.S.N.M.	3	D5276...	13 49 15 N.; 120 14 45 E..	18	sh., p., s.....	Few.
13625	U.S.N.M.	1	D5277...	13 56 55 N.; 120 13 45 E..	80	58.6	fine s.....	Rare.

Genus FLINTINA, new genus.

Description.—Test free, in its earliest stages quinqueloculine, followed very early by a triloculine development, which is maintained in all the later chambers, which become planospiral, usually taking three to make a complete cycle; wall imperforated, porcellaneous; aperture with a large orifice, with a thickened border, and large complex tooth.

Type of the genus.—*Flintina bradyana*, new species.

This genus, while somewhat like *Planospirina*, differs from that genus in its development. *Planospirina* has no quinqueloculine or triloculine stages, but has a planospiral test from the beginning. *Flintina*, on the other hand, has a derivative from *Triloculina* through the quinqueloculine stages.

It is named in honor of the late Dr. James M. Flint, whose excellent work on the *Albatross* Foraminifera is well known.

FLINTINA BARTSCHI, new species.

Description.—Test generally triangular; last-formed whorl of three chambers, polygonal in transverse section, the apertural face flat or slightly concave, the lateral faces generally somewhat concave with a sharp angle between the apertural and lateral faces; in the last-formed chamber the lateral face is somewhat convex, the chamber

broadest near the middle, tapering toward either end, with a slightly-thickened cylindrical neck; the surface of the test smooth dull matt, cream white.

Diameter 1.6 mm.

Distribution.—Type specimen (Cat. No. 10801, U. S. N. M.) from *Albatross* station D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters), bottom temperature 41.2° F. (5.1° C.).

This is a peculiar but very well-characterized species, with its triangular form, matt surface, and sharp angles.

It is named for Dr. Paul Bartsch, of the United States National Museum, who was in charge of the Philippine collecting of the *Albatross* during the early part of her Philippine cruise.

Flintina bartschi—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10801	U.S.N.M.	1	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fne. gy. s....	Rare.

FLINTINA TRIQUETRA (H. B. Brady).

Plate 94, figs. 1a-c.

Miliolina triquetra H. B. BRADY, Quart. Journ. Micr. Soc., vol. 19, 1879, p. 54; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 181, pl. 8, figs. 8-10.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 581, pl. 44, figs. 22, 23.

This species, with a planospiral character in the adult, with three chambers making up a whorl, belongs to *Flintina*, as here described. It is rare at *Albatross* station D5214, east of Masbate Island, in 218 fathoms (399 meters), the bottom temperature, 51.4° F. (10.8° C.).

Heron-Allen and Earland record is as "very finely developed off Cebu, Philippine Islands; 120 fathoms (220 meters)."

Brady's original records are from three *Challenger* stations—Bass Strait, 38 fathoms (70 meters); Torres Strait, 155 fathoms (283 meters); and Humboldt Bay, New Guinea, 37 fathoms (68 meters).

It is a rare species from the records, largely limited to Australia and the East Indian region, and found also in the Kerimba Archipelago, where Heron-Allen and Earland record two specimens.

Flintina triquetra—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10802	U.S.N.M.	2	D5214...	° ' " ° ' " 12 25 18 N.; 123 37 15 E..	218	° F. 51.4	gn. m.	Rare.

FLINTINA BRADYANA, new species.


Plate 94, fig. 2.

Miliolina fichteliana H. B. BRADY (not d'Orbigny 1839), Rep. Voy. *Challenger*, Zoology, vol. 9. 1884, p. 169, pl. 4, fig. 9.

Description.—Test triloculine, after a typical early quinqueloculine development, then in the adult planospiral; wall highly polished, with numerous fine longitudinal and slightly oblique costae; aperture large with a thickened lip and large tooth, complex in the adult.

Diameter up to 3 mm.

Distribution.—Type specimen (Cat. No. 10851, U. S. N. M.) from *Albatross* station D5192, off northern Cebu, in 32 fathoms (59 meters).



FIGS. 38-40.—FLINTINA BRADYANA, NEW SPECIES. 38, SPECIMEN WITH SIMPLE BIFID TOOTH. $\times 100$. FROM ALBATROSS STATION D5141. 39, SPECIMEN IN LATER STAGE WITH BROADENED TOOTH. $\times 50$. FROM ALBATROSS STATION D5164. 40, LARGER SPECIMEN WITH REDUCED COSTAE, THE LAST-FORMED CHAMBER SMOOTH, THE TOOTH SENDING OUT PROCESSES TOWARD THE WALL AT THE SIDES OF THE APERTURE. $\times 50$.

This is an abundant species in the Philippines at numerous stations, and is evidently the species which Brady figures, although only a partially developed specimen is figured by him. In the development of the tooth of this species, figures of which are given in the plates, there is first the typical single tooth of *Quinqueloculina*, followed by a bifid tooth characteristic of *Triloculina*; then is developed a large rounded plate-like portion, which becomes united to the border of the aperture by several projections, and later the circular plate itself becomes perforated in the center and serves largely as the aperture in the adult. This is one of the most complex developments of the aperture in the family and is very similar to that found in *Quinqueloculina albatrossi*.

F. bradyana has occurred at 22 stations; one of these at San Luis, Apra Bay, Guam, in shallow water; the others, except D5613, Gulf of Tomini, Celebes, are in the Archipelago and the following locali-

ties: China Sea, off southern Luzon; Sulu Sea, off southwestern Mindanao; near Basilan Island; off Tawi Tawi; off Jolo; off Romblon; off northern Cebu; between Marinduque and Luzon; Palawan Passage; and Malam-paya Sound, Palawan Island.




These stations range in depth from 14 to 752 fathoms (25 to 1,375 meters), the average depth being 100 fathoms (183 meters). Bottom temperatures are given at but 3 stations, ranging from 56.4° F. to 59.6° F. (13.5° C. to 15.3° C.). Most of the stations at which this is really abundant are less than 50 fathoms (91 meters), and most of them around 30 fathoms (55 meters) in depth.

It is probably widely distributed in this general region from its occurrence at Guam, and has probably been referred to the name given by Brady in the *Chal-*

lenger report, but the study of the original material is necessary to determine this.

Flintina bradyana—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10840	U.S.N.M.	10+	D5097...	14 19 15 N.; 120 33 52 E...	30		gy. m., s., sh.	Common.
10841	U.S.N.M.	6	D5100...	14 17 15 N.; 120 32 40 E...	35		gy. s.	Common.
10842	U.S.N.M.	4	D5106...	14 23 55 N.; 120 32 33 E...	37		gy. m.	Common.
10843	U.S.N.M.	1	D5110...	13 59 20 N.; 120 75 45 E...	135	59.0	dk. gy. m.	Rare.
10844	U.S.N.M.	2	D5113...	13 51 30 N.; 120 50 30 E...	159		dk. gn. m.	Rare.
10845	U.S.N.M.	10+	D5132...	Island off Panabutan Pt. N.; 15° W., 0.30 mile.	26		gn. m., s.	Common.
10846	U.S.N.M.	10+	D5133...	Island off Panabutan Pt. N.; 52° E., 1.50 miles.	38		gn. m., s.	Common.
10847	U.S.N.M.	1	D5134...	6 44 45 N.; 121 48 00 E...	25		gy. s.	Rare.
10848	U.S.N.M.	7	D5152...	5 22 55 N.; 120 15 45 E...	34		wh. s.	Common.
10849	U.S.N.M.	10+	D5164...	5 01 40 N.; 119 52 20 E...	18		gn. m.	Common.
10850	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E...	78		fine. s.	Rare.
10851	U.S.N.M.	5	D5192...	11 09 15 N.; 123 50 00 E...	32		gn. s.	Common.
10852	U.S.N.M.	4	D5220...	13 38 00 N.; 121 58 00 E...	50		sft. gn. m.	Common.
10853	U.S.N.M.	10+	D5278...	14 00 10 N.; 120 17 15 E...	102	59.6	fine. s., m., sh.	Common.
10854	U.S.N.M.	10+	D5338...	11 33 45 N.; 119 24 45 E...	43		Co., s., sh.	Common.
10855	U.S.N.M.	2	D5339...	11 22 00 N.; 119 12 00 E...	52		m.	Few.
10856	U.S.N.M.	2	D5342...	10 56 55 N.; 119 17 24 E...	14-25		gy. m.	Few.
10857	U.S.N.M.	5	D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co., s.	Few.
10858	U.S.N.M.	1	D5358...	6 06 40 N.; 118 18 15 E...	39		m.	Rare.
10859	U.S.N.M.	2	D5426...	9 12 00 N.; 118 28 00 E...	27		fine. gy. s.	Rare.
10860	U.S.N.M.	2	D5613...	0 42 00 S.; 121 44 00 E...	752		gy. m.	Rare.
10839	U.S.N.M.	3		San Luis, Apra Bay, Guam.				Rare.



FIGS. 41-44.—*FLINTINA BRADYANA*, NEW SPECIES. $\times 50$. FROM ALBATROSS STATION D5133. 41a, SIDE VIEW; b, APERTURE VIEW OF YOUNG SPECIMEN WITH SIMPLE TOOTH. 42, APERTURE OF OLDER SPECIMEN WITH BIFID TOOTH. 43, APERTURE OF A STILL OLDER SPECIMEN WITH THE PROCESSES OF THE TOOTH BEGINNING TO EXTEND IN AND THE OPENING TO HAVE ANGLES. 44, APERTURE OF AN ADULT WHERE THE ARMS HAVE MET TO FORM A CENTRAL RING, ADDITIONAL ARMS EXTENDED FROM EITHER SIDE OF THE APERTURE AT THE ANGLES AND SECONDARY ARMS BEING GIVEN OFF ON THE SIDES OF THE RING.

Genus *BILOCULINA* d'Orbigny, 1826.*BILOCULINA* *DEPRESSA* d'Orbigny.


Plate 96, figs. 2a, b.

Biloculina depressa D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 298, No. 7.—
H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 145, pl. 2,
figs. 12, 16, 17; pl. 3, figs. 1, 2.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6,
1917, p. 74, pl. 28, figs. 1, 2.

This is one of the most common species of the genus, occurring usually in the deeper water, as is shown elsewhere. It has occurred at 44 stations in the material, ranging in depth from 78 to 1,560 fathoms (143 to 2,853 meters), the average 475 fathoms (869 meters). Bottom temperatures are given at 27 stations, ranging from 36.3° F. to 59.6° F. (2.3° C. to 15.3° C), the average being 48.5° F. (19.1° C.).

Specimens are not common at any of the stations. They were found mostly in the deeper water of the Sulu Sea, China Sea, and between the islands; also in the region to the south, in Sibuko Bay, Borneo; Gulf of Tomini, Celebes; Molucca; between Gillolo and Makyan Island; off Bouro Island; Molucca Sea; Flores Sea; Gulf of Boni; and Macassar Strait.

This is a common species of deep and cooler waters in all the oceans.

Biloculina depressa—Material examined.

FIGS. 45-47.—*BILOCULINA* *DEPRESSA* D'ORBIGNY. 45, APERTURE OF YOUNG SPECIMEN, WITH ELLIPTICAL APERTURE AND ELONGATE TOOTH. $\times 50$. FROM ALBATROSS STATION D5236. 46, APERTURE OF A LATER STAGE SHOWING THE GREATER ELONGATION OF THE APERTURE AND TOOTH. $\times 100$. FROM ALBATROSS STATION D5201. 47, APERTURE OF AN ADULT WITH VERY NARROW CURVED APERTURE, LARGELY OCCUPIED BY THE SOMEWHAT EXPANDED TOOTH. $\times 40$. FROM ALBATROSS STATION D5236.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14632	U.S.N.M.	1	D5112...	13 48 22 N.; 120 47 25 E..	177	52.4	dk. gn. m....	Few.
15784	U.S.N.M.	1	D5178...	12 43 00 N.; 122 06 15 E..	78	fine. s.	Rare.
14631	U.S.N.M.	1	D5185...	10 05 45 N.; 122 18 30 E..	638	49.8	gn. m.	Rare.
15785	U.S.N.M.	1	D5193...	9 40 50 N.; 123 39 45 E..	220	53.9	gn. m.	Rare.
14633	U.S.N.M.	4	D5201...	10 10 00 N.; 125 04 15 E..	554	53.8	gy. s., m....	Few.
15786	U.S.N.M.	1	D5214...	12 25 18 N.; 123 37 15 E..	218	51.4	gn. m.	Few.
15787	U.S.N.M.	2	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	inc. gy. s.	Few.
14634	U.S.N.M.	10+	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m.	Frequent.
15788	U.S.N.M.	1	D5260...	12 25 35 N.; 121 31 35 E..	234	51.4	gu. m., s....	Rare.
14635	U.S.N.M.	8	D5268...	13 42 00 N.; 120 57 15 E..	170	s., p.	Frequent.
			D5278...	14 00 10 N.; 120 17 15 E..	102	59.6	fine. s., m., sh.	Few.
			D5300...	20 31 00 N.; 115 49 00 E..	265	gy. m., s....	Few.
			D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Few.
			D5348...	10 57 45 N.; 118 38 15 E..	375	56.4	co. s.	Few.
14636	U.S.N.M.	3	D5349...	10 54 00 N.; 118 26 20 E..	730	46.0	co. s.	Few.

Biliculina depressa—Material examined—Continued.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "				
14638	U.S.N.M.	3	D5419...	9 58 30 N.; 123 46 00 E..	175	54.5	gn. m.	Few.
			D5423...	9 38 30 N.; 121 11 00 E..	508	49.8	gy. m., co. s..	Few.
			D5425...	9 37 45 N.; 121 11 00 E..	495	49.4	gy. m., co. s..	Few.
14639	U.S.N.M.	1	D5443...	12 43 05 N.; 125 01 00 E..	241	51.3	co. s., sh.	Few.
			D5447...	13 28 00 N.; 123 46 18 E..	310	45.3	gn. m.	Few.
			D5467...	13 35 27 N.; 123 37 13 E..	480	gy. m.	Few.
			D5468...	13 35 39 N.; 123 40 28 E..	569	gn. m.	Few.
			D5470...	13 37 30 N.; 123 41 09 E..	560	m.	Few.
			D5495...	9 06 30 N.; 125 00 20 E..	967	52.3	gy. m.	Few.
			D5523...	8 48 44 N.; 123 27 35 E..	Rare.
			D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob..	Frequent.
14637	U.S.N.M.	1	D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.	Rare.
			D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gn. m., s.	Frequent.
14640	U.S.N.M.	10+	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.	Frequent.
			D5586...	4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.	Few.
14641	U.S.N.M.	10+	D5590...	4 10 50 N.; 118 39 35 E..	310	44.3	gn. m., s.	Frequent.
			D5591...	4 11 48 N.; 118 38 20 E..	260	Few.
			D5592...	4 12 44 N.; 118 27 44 E..	305	43.3	gn. m.	Few.
14642	U.S.N.M.	1	D5601...	1 13 10 N.; 125 17 05 E..	765	s., glob., ptr.	Few.
			D5609...	0 41 00 S.; 121 16 00 E..	1,092	36.3	gn. m.	Few.
			D5613...	0 02 00 S.; 121 44 00 E..	752	gy. m.	Few.
			D5618...	0 37 00 N.; 127 15 00 E..	417	gy. m.	Few.
			D5621...	0 15 00 N.; 127 24 35 E..	293	gy. & bk. s. (m. b.).	Few.
14643	U.S.N.M.	3	D5637...	3 53 20 S.; 126 48 00 E..	700	gy. m.	Few.
			D5639...	3 54 50 S.; 123 27 20 E..	1,560	gy. m.	Few.
			D5660...	5 36 30 S.; 120 49 00 E..	692	gy. m., s.	Few.
14644	U.S.N.M.	1	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.	Frequent.
15789	U.S.N.M.	10+	D5663...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.	Few.
14645	U.S.N.M.	3	H4893...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob..	Few.

BILOCULINA MURRHYNIA Schwager.

Biliculina murrhyna SCHWAGER, *Novara-Exped.*, Geol. Thiel., vol. 2, 1866, p. 203, pl. 4, figs. 5a-c.—SCHLUMBERGER, *Mém. Soc. Zool. France*, vol. 4, 1891, p. 165, pl. 9, figs. 52-54, text figs. 8, 9.—CUSHMAN, *Bull. 71, U. S. Nat. Mus.*, pt. 6, 1917, p. 75, pl. 28, fig. 3; pl. 29, fig. 1.

Biliculina depressa D'ORBIGNY, var. *murrhyna* H. B. BRADY, *Rep. Voy. Challenger*, *Zoology*, vol. 9, 1884, p. 146, pl. 2, figs. 10, 11.

This species is represented by a few specimens at but six stations, all but one, D5446, east coast of Luzon, in the region to the south, where it occurs in Sibuko Bay, Borneo; south of Patiente Strait; Moluca Sea; and the Gulf of Boni. These stations are all in comparatively deep water, ranging from 300 to 1,560 fathoms (549 to 2,853 meters). Bottom temperatures are given at but three stations—38.3° F., 38.7° F., and 44° F. (3.5° C., 3.6° C., and 6.6° C.). This is a species which occurs in deep water in most parts of the ocean.

Biliculina murrhyna—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13473	U.S.N.M.	1	D5446...	12 43 51 N.; 124 59 18 E....	300	gn. m.	Rare.
13475	U.S.N.M.	1	D5586...	4 06 50 N.; 118 47 20 E....	347	44.0	gy. m.	Few.
			D5632...	1 00 00 S.; 127 50 00 E....	845	Few.
13474	U.S.N.M.	1	D5639...	3 54 00 S.; 123 27 20 E....	1,560	gy. m.	Few.
13472	U.S.N.M.	1	D5652...	4 35 00 S.; 121 23 06 E....	525	38.7	gn. m.	Rare.
			D5654...	3 42 00 S.; 120 45 50 E....	805	38.3	Rare.

BILOCULINA SERRATA L. W. Bailey.

Plate 95, figs. 3a, b.

Biloculina serrata L. W. BAILEY, Boston Journ. Nat. Hist., vol. 7, 1862, p. 350, pl. 8, fig. E.—SCHLUMBERGER, Feuille Jeun. Nat., vol. 13, 1883, p. 106, pl. 3, fig. 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 75, pl. 29, fig. 2.

Biloculina depressa D'ORBIGNY, var. *serrata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 146, pl. 2, fig. 3.

This species is very rare as far as the records show, which is similar to what I found in most of the North Pacific material.⁴⁶

The only two stations from which it was found are D5439, west coast of Luzon, in 940 fathoms (1,719 meters); and D5639, Molucca Sea, 1,560 fathoms (2,853 meters).

This seems to be much more common in the Atlantic than in the Pacific.

Biloculina serrata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13463	U. S. N. M.	1	D5439...	15 58 15 N.; 119 40 20 E...	940	36.7	gn. m.....	Rare.
13464	U. S. N. M.	1	D5639...	3 54 50 S.; 123 27 20 E...	1,560	gy. m.....	Few.

BILOCULINA SARSHI Schlumberger.

Plate 97, figs. 1a-c.

Biloculina ringens H. B. BRADY (not *B. ringens* Lamarck), Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 142, pl. 2, fig. 7.

Biloculina sarshi SCHLUMBERGER, Mém. Soc. Zool. France, vol. 4, 1881, p. 166, pl. 9, figs. 55-59, text figs. 10, 11.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 76, pl. 30, fig. 2.





FIG. 48.—BILOCULINA SARSHI SCHLUMBERGER. $\times 50$. FROM ALBATROSS STATION D5220. a, FRONT VIEW; b, SIDE VIEW; c, APERTURAL VIEW OF SAME SPECIMEN.

⁴⁶ Bull. 71, U. S. Nat. Mus. pt. 6, 1917, p. 76.

All the stations for this species are in the Archipelago, where it occurs at 24 stations, ranging in depth from 26 to 500 fathoms (48 to



FIGS. 49, 50.—*Biloculina sarsii* SCHLUMBERGER. 49, APERTURE OF EARLY STAGE WITH A BROAD, ROUNDED TOOTH AND AN ELLIPTICAL APERTURE. $\times 80$. FROM ALBATROSS STATION D5178. 50, APERTURE OF AN ADULT SPECIMEN WITH VERY BROAD APERTURE, THE TOOTH VERY NARROW AND ELONGATE WITH THE ANGLES PRODUCED AT EACH END. $\times 100$. FROM ALBATROSS STATION D5133.

914 meters), the average being 137 fathoms (250 meters); bottom temperatures are given in few cases, the average being 59° F. (15° C.).

The stations are well scattered throughout the Archipelago.

Biloculina sarsii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14646	U.S.N.M.	1	D5097...	14 19 15 N.; 120 33 52 E...	30	gy. m., s., sh	Few.
			D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s.....	Frequent.
			D5110...	13 59 20 N.; 120 75 45 E...	135	59	dk. gy. m....	Frequent.
			D5121...	13 27 20 N.; 121 17 45 E...	108	dk. gn. m....	Few.
			D5132...	Island off Panabutan Point, N. 15 W., E., 0.30 mile.	26	gn. m., s....	Few.
14647	U.S.N.M.	10+	D5133...	Island off Panabutan Point N. 52 E., 1.50 miles.	38	gn. m., s....	Common.
14648	U.S.N.M.	10+	D5178...	12 43 00 N.; 122 06 15 E...	78	fine. s.....	Common.
15775	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.....	Rare.
			D5210...	11 49 55 N.; 124 28 05 E...	50	76.3	fine. gy. s....	Few.
			D5217...	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s....	Rare.
14649	U.S.N.M.	5	D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m....	Frequent.
15774	U.S.N.M.	10+	D5255...	7 03 00 N.; 125 39 00 E...	100	sft. m.....	Few.
14650	U.S.N.M.	1	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.....	Few.
			D5277...	13 56 55 N.; 120 13 45 E...	80	58.6	fine. s.....	Few.
			D5296...	13 40 09 N.; 120 57 45 E...	210	m., s.....	Few.
14651	U.S.N.M.	2	D5315...	21 40 00 N.; 116 58 00 E...	148	54.4	s., sh.....	Few.
			D5338...	11 33 45 N.; 119 24 45 E...	43	co. s., m....	Few.
14652	U.S.N.M.	2	D5370...	13 44 15 N.; 121 42 30 E...	159	54.3	sft. m.....	Few.
			D5394...	12 00 30 N.; 124 05 36 E...	153	gn. m.....	Few.
14653	U.S.N.M.	1	D5398...	11 35 12 N.; 124 13 48 E...	114	gn. m.....	Few.
14654	U.S.N.M.	1	D5465...	13 39 42 N.; 123 40 39 E...	500	gy. m.....	Rare.
14655	U.S.N.M.	1	D5569...	5 33 15 N.; 120 15 30 E...	303	52.3	co. s.....	Few.
			D5574...	5 30 45 N.; 120 07 57 E...	340	Few.
			D5576...	5 25 56 N.; 120 03 39 E...	277	53.3	s.....	Frequent.

Biloculina vespertilio Schlumberger.

Plate 95, figs. 5a, b.

Biloculina ringens H. B. BRADY (not Lamarck), Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 142, pl. 2, fig. 8.

Biloculina vespertilio SCHLUMBERGER, Mém. Soc. Zool. France, vol. 4, 1891, p. 174, pl. 10, figs. 74-76, text figs. 20-22.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 77, pl. 30, fig. 1; figs. 37-39 (in text).

Very fine specimens of this species occurred at *Albatross* stations D5236, east coast of Mindanao, 494 fathoms (903 meters), and D5277, China Sea, off southern Luzon, 80 fathoms (146 meters). These are the only Philippine stations; the other stations being in the area further south, where it occurs off Bouro Island; Gulf of Tomini, Celebes, and the Gulf of Boni.

Biloculina vesperilio—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13461	U.S.N.M.	1	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine. gy. s....	Few.
13462	U.S.N.M.	1	D5277...	13 56 50 N.; 120 13 45 E...	80	58.6	fine. s.....	Few.
15790	U.S.N.M.	1	D5609...	0 11 00 S.; 121 16 00 E...	1,092	36.3	gn. m.....	Rare.
			D5637...	3 53 20 S.; 126 48 00 E...	700	gy. m.....	Few.
			D5650...	4 53 45 S.; 121 29 00 E...	540	40.1	gn. m.....	Few.
			D5658...	3 32 40 S.; 120 31 30 E...	510	41.2	gy. m.....	Rare.

BILOCULINA ELONGATA d'Orbigny.

Plate 95, figs. 4a-b.

Biloculina elongata D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 298, No. 4.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 144, pl. 2, fig. 9.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 78, pl. 31, fig. 1, fig. 40 (in text).

Typical material of this species occurred at but seven stations and only a few specimens at each. These stations range in depth from 10 to 752 fathoms (18 to 1,375 meters), the average depth 350 (640 meters); except at one station, at 10 fathoms (18 meters), the least depth is 135 (247 meters), so most of the material is from deep water. Bottom temperatures are given but for three stations, ranging from 43° to 49.8° F. (6.1° to 9.8° C.).

The localities for this species include the Sulu Archipelago, off Siasi, off Jolo; Pacific Ocean, east coast of Mindanao; Gulf of Davao; northwestern Panay; and to the southward in the Gulf of Tomini, Celebes.

Biloculina elongata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15768	U.S.N.M.	1	D5149...	5 33 00 N.; 120 42 10 E...	10	Co., sh.....	Few.
15769	U.S.N.M.	2	D5238...	7 34 45 N.; 126 38 15 E...	380	43.0	gn. m.....	Rare.
			D5247...	7 02 00 N.; 125 38 45 E...	135	m.....	Rare.
			D5259...	11 57 30 N.; 121 42 15 E...	312	49.3	gy. m., glob.	Few.
13454	U.S.N.M.	1	D5423...	9 38 30 N.; 121 11 00 E...	508	49.8	gy. m., co. s.	Few.
13455	U.S.N.M.	1	D5449...	13 21 36 N.; 124 00 30 E...	300	Few.
13453	U.S.N.M.	1	D5613...	0 42 00 S.; 121 44 00 E...	752	gy. m.....	Few.

BILOCULINA GLOBULUS Bornemann.

Plate 95, figs. 2a, b.

Biloculina globulus BORNEMANN, Zeitschr. deutsch. geol. Ges., vol. 7, 1855, p. 349, pl. 19, fig. 3.—SCHLUMBERGER, Mém. Soc. Zool. France, vol. 4, 1891, p. 188, pl. 12, figs. 97-100, figs. in text, 42-44.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 78, pl. 31, fig. 2.

A few specimens occurred at 11 different stations in the region. These ranged in depth from 32 to 752 fathoms (59 to 1,375 meters), the average depth being 373 fathoms (682 meters). Bottom temperatures at four of these stations ranged from 44° F. to 73.8° F. (6.6° C. to 23.2° C.), the average being 57.5° F. (14.2° C.). Most of the stations are from 300 to 500 fathoms (549 to 914 meters).

These are off northern Cebu; Verde Island Passage; China Sea, off Formosa; Mindoro Strait; Palawan Passage; east coast of Luzon, and to the southward in Darvel and Sibuko Bays, Borneo; Gulf of Tomini, Celebes; and south of Patiente Strait.

Biloculina globulus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13480	U. S. N. M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	gn. s.	Few.
13477	U. S. N. M.	2	D5268...	13 42 00 N.; 120 57 15 E...	170	s., p.	Few.
13476	U. S. N. M.	1	D5318...	21 32 00 N.; 117 46 00 E...	340	s., br. Co ..	Few.
13484	U. S. N. M.	1	D5333...	12 26 30 N.; 120 37 45 E...	310	73.8	s.	Few.
			D5348...	10 57 45 N.; 118 38 15 E...	375	56.4	co., s.	Few.
			D5167...	13 35 27 N.; 123 37 18 E...	480	gy. m.	Few.
13483	U. S. N. M.	2	D5168...	13 35 39 N.; 123 40 28 E...	569	gn. m.	Few.
13478	U. S. N. M.	1	D5580...	4 52 45 N.; 119 06 45 E...	162	55.8	br. s., Co ..	Few.
13475	U. S. N. M.	1	D5586...	4 06 50 N.; 118 47 20 E...	347	44.0	gy. m.	Few.
13479	U. S. N. M.	1	D5613...	0 42 00 N.; 121 44 00 E...	752	gy. m.	Few.
13481	U. S. N. M.	1	D5630...	0 56 30 S.; 128 05 00 E ..	569	co. s., m.	Few.

BILOCULINA ANOMALA Schlumberger.

Plate 96, figs. 1a-c.

Biloculina anomala SCHLUMBERGER, Mém. Soc. Zool. France, vol. 4, 1891, p. 182, pl. 11, figs. 84-86; pl. 12, fig. 101; text figs. 32-34.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 79, pl. 32, fig. 1.

There are but three records for this species; D5201, Sogod Bay, southern Leyte, in 554 fathoms (1,012 meters); D5622, between Gillolo and Makyan, in 275 fathoms (503 meters); and H4898, Sulu Sea, off western Mindanao, 221 fathoms (405 meters).

Schlumberger's material was from the Gulf of Marseilles, and I have recorded this species in the North Pacific, off the Hawaiian Islands.

From these records it is probably found in the general Indo-Pacific region in comparatively deep water.

Biloculina anomala—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13457	U. S. N. M.	1	D5201...	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m....	Rare.
			D5622...	0 19 20 N.; 127 28 30 E...	275	gy. m.	Rare.
15776	U. S. N. M.	1	D1898...	7 43 45 N.; 122 03 45 E...	221	gy. m. glob..	Rare.

BILOCULINA LUCERNULA Schwager.

Plate 97, figs. 2a-c; pl. 98, figs. 1a-c.

Biloculina lucernula SCHWAGER, *Novara-Exped.*, Geol. Theil., vol. 2, 1866, p. 202, pl. 4, figs. 14a, c, 17a, b.—SCHLUMBERGER, *Mém. Soc. Zool. France*, vol. 4, 1891, p. 185, pl. 12, figs. 90-96; text figs. 37-41.—CUSHMAN, *Bull. U. S. Nat. Mus.*, pt. 6, 1917, p. 79, pl. 32, fig. 2.

Biloculina tubulosa H. B. BRADY (not Costa), *Rep. Voy. Challenger*, Zoology, vol. 9, 1884, p. 147, pl. 3, figs. 6, 14.

There is but one station for the typical form of this species—D5630, south of Patiente Strait, in 569 fathoms (1,040 meters). The triloculine form occurs at D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters).

This species, described by Schwager from Kar Nicobar Island, is evidently widely spread in the Indo-Pacific, and probably elsewhere although it has not been common in this material.

I have had it in the Pacific from the western coast of America and the Hawaiian Islands.

Biloculina lucernula—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15771	U. S. N. M.	5	D5236...	8 50 45 N.; 126 26 52 E...	494	41.2	fine gy. s....	Few.
13458	U. S. N. M.	1	D5630...	0 56 30 S.; 128 05 00 E ..	569	co. s., m.....	Rare.

BILOCULINA PISUM Schlumberger.

Biloculina pisum SCHLUMBERGER, *Mém. Soc. Zool. France*, vol. 4, 1891, p. 569, pl. 11, figs. 81-83; fig. 31 (in text).—CHAPMAN, *Trans. New Zealand Inst.*, vol. 38, 1905, p. 80.

Specimens, apparently this species, occur at a single station D5121, east coast of Mindoro, in 108 fathoms (198 meters).

Schlumberger described this from the Mediterranean; Chapman records it from off Great Barrier Island, New Zealand; and I have had it from the Poor Knights Islands, in the same general region.

Bilocolina pisum—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15772	U.S.N.M.	1	D5121...	° ' " ° ' " N.; 121 17 45 E..	108	° F.	dk. gn. m....	Few.

BILOCULINA DENTICULATA (H. B. Brady).

Plate 98, figs. 3a, b.

Bilocolina ringens (Lamarck), var. *denticulata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 143, pl. 3, figs. 4, 5.

Bilocolina denticulata CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 80, pl. 33, fig. 1.

This species, which was described by Brady as a variety of *Bilocolina ringens*, I placed as a definite species (reference above).

It is evidently a shallow-water, tropical species, in its best development in the Indo-Pacific region, where it occurs from the Hawaiian Islands, west to the Indian Ocean.

In the Philippines it has occurred at 11 stations, all in shallow water, ranging in depth from 10 to 100 fathoms (18 to 183 meters), the average depth 33 fathoms (60 meters). The bottom temperature is given at but one of these stations, that from 37 fathoms (68 meters), where it is 75.7° F. (24.2°C.), showing that all the records for this species, except perhaps that one at 100 fathoms (183 meters), are in warm waters.

At the one station D5179, off Romblon in 37 fathoms (68 meters), it is very abundant.

Bilocolina denticulata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13466	U.S.N.M.	10+	D5134 ..	° ' " ° ' " N.; 121 48 00 E..	25	° F.	fne. s.	Common.
15781	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E..	19	s., Co	Rare.
			D5141...	6 09 00 N.; 120 58 00 E..	29	co. s	Rare.
13468	U.S.N.M.	9	D5159...	5 11 50 N.; 119 54 00 E..	10	co. s	Common.
15782	U.S.N.M.	1	D5161...	5 10 15 N.; 119 53 00 E..	16	fne. s.	Rare.
13470	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd. s.	Abundant.
13465	U.S.N.M.	1	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s	Common.
15783	U.S.N.M.	1	D5255...	7 03 00 N.; 125 39 00 E..	100	sft. m	Rare.
13469	U.S.N.M.	1	D5312...	10 56 55 N.; 119 17 24 E..	14-25	gy. m.	Rare.
13467	U.S.N.M.	1	Jolo Jolo, P. I.	Frequent.
13157	U.S.N.M.	3	Apra Bay, Guam.	Few.

BILOCULINA DENTICULATA (H. B. Brady), var. STRIOLATA H. B. Brady.

Plate 98, figs. 2a-c.

Biloculina ringens (Lamarck), var. *striolata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 143, pl. 3, figs. 7, 8.—MILLETT, Journ. Roy. Micr. Soc., 1898, p. 262, pl. 5, fig. 8.

Biloculina denticulata, var. *striolata* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 80, pl. 33, figs. 2, 3.

This variety is much less common than the typical form of the species, occurring at but five stations in the region D5139, off Jolo in 20 fathoms (37 meters); D5141, the neighboring station, 29 fathoms (53 meters); D5159, off Tawi Tawi in 10 fathoms; D5179, off Romblon in 37 fathoms (168 meters); and from the exterior of a pearl oyster from Jolo Jolo.

Evidently the variety is like the typical form in being most at home in shallow, warm water.

Biloculina denticulata, var. *striolata*—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13471	U.S.N.M.	10+	D5139...	6 06 00 N.; 121 02 30 E..	20	° F.	co.s.....	Few.
			D5141...	6 09 00 N.; 120 58 00 E..	29		co.s.....	Common.
			D5159...	50 11 50 N.; 119 54 00 E..	10		co.s.....	Rare.
			D5179...	12 38 15 N.; 122 12 30 E..	37	75.7	hrd.s.....	Few.
				Jolo Jolo, P. I.				

BILOCULINA COMATA H. B. Brady.

Plate 96, figs. 3a, b.

Biloculina comata H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 45; Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 144, pl. 3, figs. 9a, b.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 81, pl. 34, fig. 1.

As known elsewhere this species occurs only in waters of considerable depth. Specimens occurred at 21 stations, ranging in depth from 50 to 901 fathoms (91 to 1,648 meters), the average being 375 fathoms (686 meters). Bottom temperatures range from 38.2° to 59.6° F. (3.4° to 15.3° C.), the average being 51° F (10.5° C.).

The only stations at which this species is at all frequent range from 256 to 494 fathoms (468 to 903 meters).

These localities include stations in the Philippine region and about the Sulu Sea, off western Mindanao, and off Tawi Tawi; Sogod Bay,



FIG. 51.—BILOCULINA COMATA H. B. BRADY. $\times 100$. APERTURE OF SPECIMEN FROM ALBATROSS STATION D5236.

southern Leyte, between Marinduque and Luzon; Pacific Ocean, east coast of Mindanao; off northwestern Panay, China Sea, off southern Luzon and Formosa; between Siquijor and Bohol; between Negros and Siquijor; and several stations to the southward in Darvel




FIG. 52.—*Biloculina comata* H. B. BEADY. OUTLINE OF THE CHAMBERS OF A PARTIALLY BROKEN SPECIMEN, SHOWING THE EARLY QUINQUELOCULINE STAGE WITH NEARLY SMOOTH CHAMBERS. $\times 100$. FROM ALBATROSS STATION D5236.

and Sibuko Bays, Borneo; south of Patiente Strait; Gulf of Boni; and Macassar Strait.

Some of the stations show the triloculine form which has already been referred to under *Triloculina insignis*, and which has been confused by various writers under that name.

Biloculina comata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
13447	U.S.N.M.	1	D5201...	10 10 00 N.; 125 04 15 E..	554	52.8	gy. s., m.....	Rare.
			D5220...	13 38 00 N.; 121 58 00 E..	50	sft. gn. m.....	Few.
13442	U.S.N.M.	1	D5222...	13 38 30 N.; 121 42 45 E..	195	52.8	gn. m.....	Rare.
15778	U.S.N.M.	10	D5236...	8 50 45 N.; 126 26 52 E..	494	41.2	fine. gy. s.....	Frequent.
13450	U.S.N.M.	2	D5259...	11 57 30 N.; 121 42 15 E..	312	49.3	gy. m., glob.	Few.
			D5278...	14 00 10 N.; 120 17 15 E..	102	59.6	fine. s., m., sh.	Few.
			D5318...	21 32 00 N.; 117 46 00 E..	340	s., br. Co....	Few.
			D5537...	9 11 00 N.; 123 23 00 E..	254	53.5	gn. m.....	Frequent.
13451	U.S.N.M.	5	D5529...	9 23 45 N.; 123 39 30 E..	441	53.0	gy. m., glob.	Frequent.
			D5538...	9 08 15 N.; 123 23 20 E..	256	53.3	gn. m., s.....	Frequent.
13446	U.S.N.M.	3	D5567...	5 48 00 N.; 120 33 45 E..	268	52.0	fine. s.....	Few.
13443	U.S.N.M.	2	D5663...	2 28 15 S.; 118 49 00 E..	901	38.2	gy. m.....	Few.
13448	U.S.N.M.	2	D5650...	4 53 45 S.; 121 29 00 E..	540	40.1	gn. m.....	Few.
			D5630...	0 56 30 S.; 128 05 00 E..	569	co. s., m.....	Few.
13452	U.S.N.M.	4	D5591...	4 11 48 N.; 118 38 20 E..	260	Few.
13441	U.S.N.M.	1	D5582...	4 19 54 N.; 118 58 38 E..	890	38.3	gy. m., fine. s.	Few.
15779	U.S.N.M.	6	D5585...	4 07 00 N.; 118 49 54 E..	476	41.1	gy. m.....	Common.
13445	U.S.N.M.	5		4 06 50 N.; 118 47 20 E..	347	44.0	gy. m.....	Few.
13444	U.S.N.M.	2		4 12 10 N.; 118 38 08 E..	260	45.7	fine. gy. s., gy. m.	Few.
13449	U.S.N.M.	1	D5425...	9 37 45 N.; 121 11 00 E	495	49.4	gy. m., co. s.	Rare.
				(triloculine form).				
15780	U.S.N.M.	1	H4898...	7 43 45 N.; 122 03 45 E..	221	gy. m., glob.	Rare.

BILOCULINA MILETTII Cushman.

Miliolina durrandii MILLETT (part), Journ. Roy. Micr. Soc., 1898, p. 268, pl. 6, figs. 8-10 (not figure 7).—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 565, pl. 42, figs. 11-16.

Biloculina millettii CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 81, pl. 34, figs. 4, 5.

This species, described from the Malay region by Millett, and which I also found in Hongkong Harbor, has occurred in the Philippine region at one station. It is evidently characteristic of this rather limited region.

Heron-Allen and Earland record this in their paper from the Kerimba Archipelago, and also note its occurrence "from the coasts of Burma, Queensland, Java, and Macassar in the eastern seas; and in the Pacific is very abundant, and typical at Tahiti."

This species seems to be without an apertural tooth, none having occurred in our specimen, and Heron-Allen and Earland mention the same characteristic.

Biloculina millettii—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15773	U.S.N.M.	1	D5210...	° ' " ° ' " 11 49 55 N.; 124 28 05 E..	50	° F. 76.3	fine. gy. s....	Rare.

BILOCULINA IRREGULARIS d'Orbigny.

Plate 95, figs. 1a-b.

Biloculina irregularis D'ORBIGNY, Foram. Amér. Mérid., 1839, "Foraminifères," p. 67, pl. 8, figs. 22-24.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 140, pl. 1, figs. 17, 18.

This rather widely distributed species in comparatively deep water is recorded by Brady from three South Pacific stations, ranging in depth from 610 to 1,070 fathoms (1,116 meters to 1,957 meters), off Fiji, off Tahiti, and north of New Guinea. Numerous specimens have occurred at a single station in the Philippine dredgings—D5236, Pacific Ocean, east coast of Mindanao, in 494 fathoms (903 meters); bottom temperature, 41.2° F. (5.1° C.).

Biloculina irregularis—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15770	U.S.N.M.	5	D5236...	° ' " ° ' " 8 50 45 N.; 126 26 52 E..	494	° F. 41.2	fine. gy. s....	Few.

Genus *NEVILLINA* Sidebottom, 1905.

Nevillina SIDEBOTTOM (Type *N. coronata* (Millett)), Mem. Proc. Manchester Lit. Philos. Soc., vol. 49, pt. 2, No. 11, 1905, p. 1.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 83.

This very interesting genus has been discussed at some length in the references given above.

NEVILLINA CORONATA (Millett).

Plate 99, figs. 1a, b.

Biloculina coronata MILLETT, Journ. Roy. Micr. Soc., 1898, p. 263, pl. 6, figs. 6a-c.

Nevillina coronata SIDEBOTTOM, Mem. Proc. Manchester Lit. Philos. Soc., vol. 49, pt. 2, No. 11, 1905, p. 1, figs. 1-8.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 84, pl. 35, figs. 2-5.

Some of the relationships of this very interesting genus and species have already been noted⁴⁷. The species was originally discovered and described by Millett from a solitary specimen from "Asjahan, northeast coast of Sumatra," depth not given. Millett's specimen was evidently not adult, but still in the biloculine stage. Later Sidebottom described the genus *Nevillina* from more adequate material from two stations—Port Blair, east side of Andaman Island, at entrance to the port, between Ross Island and the main island in 16 fathoms (29 meters); and Sulu roadstead, 12 fathoms (22 meters). These two stations with that of Millett show that the species must have a wide distribution in the Indo-Pacific.

In the Philippine material it is very rare. A single apertural end of a fully developed specimen is from D5142, off Jolo, in 21 fathoms (39 meters); and two specimens, also fully mature and very typical, from D5160, off Tinakta Island, Sulu Archipelago, Tawi Tawi Group, in 12 fathoms (22 meters).

When sufficient material of this species can be obtained the early development of the microspheric form should be studied and compared with the sections figured by Schlumberger of the genera *Idalina* and *Periloculina*.

This is evidently one of the most interesting Foraminifera of the region, representing a species with Cretaceous affinities still existing in the Indo-Pacific region.

Nevillina coronata—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
13584	U.S.N.M.	1	D5142...	6 06 10 N.; 121 02 40 E..	21	° F.	co. s., sh....	Rare.
13585	U.S.N.M.	2	D5160...	5 12 40 N.; 119 55 10 E..	12	s.....	Rare.

⁴⁷ Cushman, 1917, pp. 83-84.

Genus *PENEROPLIS* Montfort, 1808.

This genus occurs in numerous forms in the area, and the same arrangement as that adopted by Heron-Allen and Earland in their paper on the Kerimba Archipelago is here adopted. They, having made an extended study of the earlier figures and descriptions, have given a very valuable series of notes on the systematic arrangement of the forms of this genus. Their paper is here referred to so that their disposition of the species may be followed.

PENEROPLIS PERTUSUS (Forskålt).

Nautilus pertusus FORSKÅL, Descr. Anim., 1775, p. 125, No. 65.

Peneroplis pertusus JONES, PARKER, and H. B. BRADY, Foram. Crag., 1865, p. 19.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 204, pl. 13, figs. 16, 17.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 601.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 86, pl. 36, fig. 1; pl. 37, figs. 1, 2, 6.

This typical form of the species is not the most abundant in the region. There are records from four stations: Sulu Archipelago; off Tawi Tawi; between Burias and Luzon; and Midway Island. It undoubtedly occurs, as do the other forms, in shallow water throughout the region.

Peneroplis pertusus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
14660	U.S.N.M.	3	D5154...	5 14 50 N.; 119 58 45 E..	12	co. s.....	Few.
14661	U.S.N.M.	1	D5159...	5 11 50 N.; 119 54 00 E..	10	co. s.....	Few.
14662	U.S.N.M.	10+	D5218...	13 11 15 N.; 123 02 45 E..	20	crs. s.....	Common.
14663	U.S.N.M.	2	Midway Island.....	Few.

PENEROPLIS PLANATUS (Fichtel and Moll.)

Nautilus (Lituus) arietinus (part) BATSCH, Conch. des Seesandes, 1791, p. 4, pl. 6, figs. 15a, b.

Nautilus planatus, var. β FICHEL and MOLL, Test. Micr., 1803, p. 91, pl. 16, figs. 1d, e, f.

Peneroplis planatus D'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 285, No. 1.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 204, pl. 13, fig. 15.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 601.

Peneroplis pertusus, var. *planatus* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 87, pl. 37, fig. 3.

This is the most common form, being very abundant at a few of the shallow-water stations, and occurring in lesser numbers than others. The stations range in depth from 12 to 78 fathoms (22 to 143 meters), the average being about 30 fathoms (55 meters). Bottom temperatures are not given.

Most of the stations are in the shallow, warm waters about the islands of the Sulu Sea.

Peneroplis planatus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14668	U.S.N.M.	5	D5133...	Island of Panabutan Point N. 52° E., 1.50 miles.	38	° F.	gn. m., s.....	Common.
				" " " " "				
14669	U.S.N.M.	1	D5139...	6 06 00 N.; 121 02 30 E...	20		co. s.....	Few.
14670	U.S.N.M.	1	D5152...	5 22 55 N.; 120 15 45 E...	34		wh. s.....	Few.
14671	U.S.N.M.	3	D5160...	5 12 40 N.; 119 55 10 E...	12		s.....	Few.
14672	U.S.N.M.	1	D5161...	5 10 15 N.; 119 53 00 E...	15		fine s.....	Few.
14673	U.S.N.M.	2	D5164...	5 01 40 N.; 119 52 20 E...	18		gn. m.....	Few.
14674	U.S.N.M.	1	D5178...	12 43 00 N., 122 06 15 E...	78		ine. s.....	Few.
14675	U.S.N.M.	2	D5192...	11 09 15 N.; 123 50 00 E...	32		gn. s.....	Few.
14676	U.S.N.M.	1	D5206...	11 31 40 N.; 124 42 40 E...	32		gn. m.....	Few.
14677	U.S.N.M.	10+	D5218...	13 11 15 N.; 123 02 45 E...	20		crs. s.....	Common.
14678	U.S.N.M.	1		Binang Pool, Subin Bay, P. I.				Few.
14679	U.S.N.M.	3		Tacloban Anchorage, P. I.				Common.

PENEROPLIS CARINATUS d'Orbigny.

Peneroplis carinatus D'ORBIGNY, Foram. Amér. Mérid., 1839, "Foraminifères," p. 33, pl. 3, figs. 7, 8.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 205, pl. 13, fig. 14.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 602.

Peneroplis pertusus, var. *carinatus* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 87, pl. 37, fig. 4.

This species is common at D5159, off Tawi Tawi, in 10 fathoms (18 meters).

Peneroplis carinatus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
4658	U.S.N.M.	10+	D5159...	5 11 50 N.; 119 54 00 E...	10	° F.	co. s.....	Common.

PENEROPLIS ARIETINUS (Batsch).

Nautilus (Lituus) arietinus (part) BATSCH, Conch. des Seesandes, 1791, p. 4, pl. 6, fig. 15c.

Peneroplis arietinus PARKER, JONES, and H. B. BRADY, Ann. Mag. Nat. Hist., ser. 3, vol. 16, 1865, p. 26, pl. 1, fig. 18.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 204, pl. 13, figs. 18, 19, 22.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 602.

Peneroplis pertusus, var. *arietinus* CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 88, pl. 36, fig. 2; pl. 37, fig. 5.

The elongate form is common at some stations, the depths at which it was recorded being between 10 and 20 fathoms (18 and 37 meters), mostly among the islands in the Sulu region.

Peneroplis arietinus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14664	U.S.N.M.	4	D5154...	5 14 50 N.; 119 58 45 E...	12	° F.	co. s.....	Common.
14665	U.S.N.M.	5	D5139...	5 11 50 N.; 119 54 00 E...	10		co. s.....	Few.
14666	U.S.N.M.	5	D5218...	13 11 15 N.; 123 02 45 E...	20		crs. s.....	Common.
14667	U.S.N.M.	1	Midway Island.....	Rare.
13156	U.S.N.M.	1	Manila Bay.....	Rare.

PENEROPLIS CYLINDRACEUS (Lamarck).

Nautilus acicularis BATSCH, Conch. des Seesandes, 1791, p. 4, pl. 6, figs. 16 a, b.
Spirolinities cylindracea LAMARCK, Ann. du Mus., vol. 5, 1804, p. 245; vol. 8, 1806, pl. 62, fig. 16.

Peneroplis cylindraceus H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 205, pl. 13, figs. 20, 21.—HERON-ALLEN and EARLAND, Trans. Zool. Soc. London, vol. 20, 1915, p. 662.

This elongate cylindrical species has occurred at *Albatross* station D5192, off northern Cebu, in 32 fathoms (59 meters).

Peneroplis cylindraceus—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
14659	U.S.N.M.	1	D5192...	11 09 15 N.; 123 50 00 E...	32	° F.	gn. s.....	Few.

Genus ORBICULINA Lamarck, 1816.

ORBICULINA ADUNCA (Fichtel and Moll).

Nautilus aduncus FICHEL and MOLL, Test. Micr., 1803, p. 115, pl. 23.

Orbiculina adunca LAMARCK, Tabl. Encycl. et Meth., 1816, pl. 468, figs. 2 a-c.—
 H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 209, pl. 14, figs. 1-13.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1917, p. 91, pl. 37, figs. 7, 8.

I have found this species at but two stations in the Philippine region—D5158 and D5164, Sulu Archipelago, off Tawi Tawi, in 12 and 18 fathoms (22 and 33 meters). The species is rare at both stations.

Brady records this from about the various East Indian Islands, but I have failed to find it in the North Pacific except at one station. It did not occur in the material I had from about the Hawaiian Islands, and it seems to be very rare in the Indo-Pacific, being replaced by various species of *Orbitolites*. Dakin records it as rare from the Gulf of Mannar, Ceylon. Heron-Allen and Earland do not record it from the Kerimba Archipelago.

Orbiculina adunca—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
			D5158...	° ' " ° ' "	12	°F.	crs. s., sh.....	Rare.
			D5164...	5 12 00 N.; 119 54 30 E... 5 01 40 N.; 119 52 20 E...	18	gn. m.....	Rare.

Genus **ORBITOLITES** Lamarck.**ORBITOLITES MARGINALIS** (Lamarck).

Orbitolites marginalis LAMARCK, Hist. Nat. Anim. sans Vert., vol. 2, 1816, p. 196, No. 1.

Orbitolites marginalis CARPENTER, Philos. Trans., 1856, p. 192, pl. 9, figs. 1-4.—
H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 214, pl. 15, figs. 1-5.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1915, p. 92, pl. 38, figs. 1, 2.

In the reference last referred to I have figured and described the young stages of the three common species of *Orbitolites*. Heron-Allen and Earland in their paper, "The Kerimba Archipelago Foraminifera," note the difficulties of distinguishing this species and the following (*O. duplex*) simply on the basis of the single or double series of marginal pores. The same holds true of the Philippine material, where both species often occur together in various stages of development. In such cases the early development is a much more accurate character to distinguish the species, especially in the megalospheric form. As noted in the above reference on the North Pacific, *O. marginalis* has a series of several Orbiculine chambers with single apertures and before divisions occur in the chambers has developed to a decidedly crozier-shaped test. Before true annular chambers are developed the test attains a considerable size. Throughout the test there is a single layer of chambers unless toward the edge there may be a tendency toward *O. duplex* and a double series of pores developed.

O. duplex, on the other hand, has a much more accelerated development as there figured, and the crozier-shape is attained almost at once and annular chambers are developed very early. The double tier of chambers also is developed early, but their apertures may for some time all be in a single series in the middle of the border.

As also noted by Heron-Allen and Earland, *O. duplex* is much more prone to exhibit excess developments of shell material in the form of extra plates than *O. marginalis*.

Where living material was preserved all three species show a decided greenish color, due to the presence of symbiotic algae. These were perhaps most marked in *O. complanata*.

As a rule the chamber walls of *O. marginalis* are thinner than in the other species giving a characteristic bluish tinge to the test not usually noticed in either of the other species.

O. marginalis was less common than *O. duplex*, but more so than *O. complanata* except in very shallow water. It is abundant at many of the stations usually in shallow water but in a few cases found at more than 500 fathoms (914 meters).

It seems to be most common among the islands rather than on the open outer shores. To the southward it was found between Gillolo and Kayoa.

Orbitolites marginalis—Material examined.

Cat. No.	Coll. of—	No. of specimans.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
15792	U.S.N.M.	1	D5097...	14 19 15 N.; 120 33 52 E...	30	° F.	gy. m., s., sh	Few.
15793	U.S.N.M.	1	D5100...	14 17 15 N.; 120 32 40 E...	35	gy. s.....	Few.
15794	U.S.N.M.	1	D5105...	14 43 55 N.; 120 12 50 E...	25	Few.
15795	U.S.N.M.	5	D5132...	Island off Panabutan Point, N. 15° W., 0.30 mile.	26	gn. m., s.....	Few.
15796	U.S.N.M.	10+	D5133...	Island off Panabutan Point, N. 52° E., 1.50 miles.	38	gn. m., s.....	Abundant.
15797	U.S.N.M.	1	D5138...	6 06 00 N.; 120 58 50 E...	19	s., Co.....	Few.
15798	U.S.N.M.	10+	D5159...	5 11 50 N.; 119 54 00 E...	10	co. s.....	Common.
15799	U.S.N.M.	4	D5160...	5 12 40 N.; 119 55 10 E...	12	s.....	Few.
			D5191...	10 29 45 N.; 123 31 15 E...	258	62.8	gr. m.....	Few.
			D5218...	13 11 15 N.; 123 02 45 E...	20	ers. s.....	Few.
			D5220...	13 38 00 N.; 121 58 00 E...	50	sft. gn. m.....	Few.
15800	U.S.N.M.	1	D5244...	6 52 05 N.; 126 14 15 E...	171	gy. m.....	Few.
15801	U.S.N.M.	1	D5257...	7 22 12 N.; 124 12 15 E...	28	m.....	Few.
			D5276...	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Few.
			D5382...	13 15 20 N.; 122 45 30 E...	128	m.....	Few.
			D5388...	12 51 30 N.; 123 26 15 E...	226	51.4	sft. gn. m.....	Few.
			D5408...	10 40 15 N.; 124 15 00 E...	159	55.4	gn. m.....	Few.
			D5419...	9 58 30 N.; 123 46 00 E...	175	54.5	gn. m.....	Few.
			D5424...	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Few.
15802	U.S.N.M.	3	D5529...	9 23 45 N.; 123 39 30 E...	441	53.0	gy. m., glob.	Few.
			D5465...	13 39 42 N.; 123 40 39 E...	500	gn. m.....	Few.
			D5469...	13 36 48 N.; 123 38 24 E...	500	gn. m.....	Few.
			D5170...	13 37 30 N.; 123 41 09 E...	540	m.....	Few.
			D5527...	0 03 00 N.; 127 26 00 E...	22	m.....	Few.
15804	U.S.N.M.	1	Tayabas Gulf, P. I.	Few.
15803	U.S.N.M.	1	Siaast Island, P. I.	Few.

ORBITOLITES DUPLEX Carpenter.

Orbitolites, duplex type CARPENTER, Philos. Trans., 1856, p. 120, pl. 5, fig. 10; pl. 9, fig. 10.

Orbitolites duplex CARPENTER, Rep. Challenger, "Orbitolites," 1883, p. 25, pl. 3, figs. 8-14; pl. 4, figs. 6-10; pl. 5, figs. 1-10.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 216, pl. 16, fig. 7.—CUSMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1916, p. 94, pl. 38, figs. 3, 4; pl. 39, fig. 1.

This species is represented at a greater number of stations than either of the others as far as our material shows. Depths range from 10 to 732 fathoms (18 to 1,339 meters). It is most common in fairly shallow water among the islands of the Archipelago, but occurred in Darvel and Sibuko Bays, Borneo. Most of the stations were at depths

of less than 100 fathoms (183 meters); eight between 100 and 200 fathoms (183 and 366 meters); five between 200 and 300 fathoms (366 and 549 meters); six between 300 and 400 fathoms (549 and 732 meters); five above 400 fathoms (732 meters). At the deeper stations the species was usually represented by few specimens.

Numerous specimens formed of fused individuals were found, two or more with the earlier annular chambers having fused into a complete annular growth.

Orbitolites duplex—Material examined.

Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' " ° ' "		° F.		
15805	U.S.N.M.	1	D5106	14 23 55 N.; 120 32 33 E...	37	gv. m.....	Few.
15806	U.S.N.M.	4	D5110	13 59 20 N.; 129 75 45 E...	135	59.0	dk. gy. m.....	Few.
15807	U.S.N.M.	1	D5148	5 35 40 N.; 120 47 30 E...	17	co. s.....	Few.
15808	U.S.N.M.	1	D5147
15809	U.S.N.M.	4	D5149	5 33 00 N.; 120 42 10 E...	10	Co., sh.....	Few.
15810	U.S.N.M.	3	D5154	5 14 50 N.; 119 58 45 E...	12	co. s.....	Few.
15811	U.S.N.M.	3	D5159	5 11 50 N.; 119 54 00 E...	10	co. s.....	Common.
15812	U.S.N.M.	1	D5160	5 12 40 N.; 119 55 10 E...	12	s.....	Few.
15813	U.S.N.M.	1	D5161	5 10 15 N.; 119 53 00 E...	16	fne. s.....	Few.
			D5201	10 10 00 N.; 125 04 15 E...	554	52.8	gy. s., m.....	Few.
15814	U.S.N.M.	1	D5206	11 31 40 N.; 124 42 40 E...	32	gn. m.....	Few.
15815	U.S.N.M.	1	D5214	12 25 18 N.; 123 37 15 E...	218	gn. m.....	Few.
15816	U.S.N.M.	2	D5217	13 20 00 N.; 123 14 15 E...	105	63.1	crs. gy. s.....	Few.
15817	U.S.N.M.	1	D5218	13 11 15 N.; 123 02 45 E...	20	crs. s.....	Few.
			D5220	13 38 00 N.; 121 53 00 E...	50	sft. gn. m.....	Few.
15818	U.S.N.M.	1	D5259	11 57 39 N.; 121 42 15 E...	312	49.3	gy. m., glob.....	Few.
15819	U.S.N.M.	1	D5261	12 30 55 N.; 121 34 24 E...	145	s., m.....	Few.
			D5266	13 49 15 N.; 120 14 45 E...	18	sh., p., s.....	Common.
			D5277	13 56 55 N.; 120 13 45 E...	80	58.6	fne. s.....	Few.
			D5278	14 00 10 N.; 120 17 15 E...	102	59.6	fne. s., m., sh.....	Few.
			D5338	11 33 45 N.; 119 24 45 E...	43	Co., s., m.....	Few.
			D5369	13 48 00 N.; 121 43 00 E...	106	bk. s.....	Few.
			D5374	13 46 45 N.; 121 35 08 E...	190	gy. m.....	Few.
			D5381	13 14 15 N.; 122 44 25 E...	88	co. s.....	Few.
			D5388	12 51 30 N.; 123 26 15 E...	225	51.4	sft. gn. m.....	Few.
			D5424	9 37 05 N.; 121 12 37 E...	340	50.4	co. s.....	Few.
15820	U.S.N.M.	1	D5426	9 12 00 N.; 118 28 00 E...	27	fne. gy. s.....	Few.
			D5438	15 54 32 N.; 119 44 42 E...	297	46.2	gn. m.....	Few.
			D5459	13 10 21 N.; 123 59 54 E...	201	Few.
			D5445	12 44 42 N.; 124 59 50 E...	383	44.3	gn. m., s.....	Few.
			D5467	13 35 27 N.; 123 37 18 E...	480	gn. m.....	Few.
			D5469	13 36 43 N.; 123 38 24 E...	500	gn. m.....	Few.
			D5487	10 02 45 N.; 125 05 33 E...	732	52.3	gn. m.....	Few.
			D5567	5 48 00 N.; 120 33 45 E...	238	52.0	fne. s.....	Few.
			D5570	5 32 15 N.; 120 12 57 E...	330	52.3	fne. s., glob.....	Few.
15821	U.S.N.M.	5		Binang Pool, Subin Bay	Few.
15822	U.S.N.M.	2		Siasi Island	Few.
			D5579	4 54 15 N.; 119 09 52 E...	175	55.3	fne. s., Co.....	Few.
			D5580	4 52 45 N.; 119 06 45 E...	162	55.8	fne. s., Co.....	Common.
			D5589	4 12 10 N.; 118 38 08 E...	260	45.7	fne. gy. s.....	Few.
				gy. m.....
			D5590	4 10 50 N.; 118 39 35 E...	310	44.3	gn. m., s.....	Few.
			D5592	4 12 44 N.; 118 27 44 E...	305	43.3	gn. m.....	Few.

ORBITOLITES COMPLANATA Lamarck.

Orbitolites complanata LAMARCK, Syst. Anim. sans Vert., 1801, p. 376.—H. B. BRADY, Rep. Voy. Challenger, Zoology, vol. 9, 1884, p. 218, pl. 16, figs. 1-6; pl. 17, figs. 1-6.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1916, p. 95, pl. 39, fig. 2.

Specimens of this species were abundant at many stations and at some reaching a very considerable size. Specimens of the micro-

spheric form were found with the outer annuli more or less broken and thin, filled with numerous megalospheric young. These large specimens were of the flattened form noted by Heron-Allen and Earland from the Kerimba Archipelago. Specimens composed of fused individuals are occasionally found also. The range of depths is from 12 to 766 fathoms (22 to 1,400 meters), but two stations being greater in depth than 500 fathoms (914 meters).

Orbitolites complanata—Material examined.

Cst. No.	Coll. of—	No. of specimens.	Station.	Locality.		Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
				° ' "	° ' "				
15824	U. S. N. M.	1	D5134...	6 44 45 N.	121 48 00 E.	25	fne. s.	Few.
15825	U. S. N. M.	6	D5152...	5 22 55 N.	120 15 45 E.	34	wh. s.	Common.
15826	U. S. N. M.	8	D5154...	5 14 50 N.	119 58 45 E.	12	co. s.	Common.
15827	U. S. N. M.	1	D5161...	5 10 15 N.	119 53 00 E.	16	fne. s.	Few.
			D5191...	10 29 45 N.	123 31 15 E.	258	62.8	gn. m.	Few.
15828	U. S. N. M.	1	D5201...	10 10 00 N.	125 04 15 E.	554	52.8	gy. s., m.	Few.
			D5220...	13 38 00 N.	121 58 00 E.	50	sft. gn. m.	Few.
15829	U. S. N. M.	3	D5272...	14 00 00 N.	120 22 30 E.	118	57.4	m., sh., co. s.	Few.
			D5276...	13 49 15 N.	120 14 45 E.	18	sh., p., s.	Few.
			D5374...	13 46 45 N.	121 35 08 E.	190	gy. m.	Few.
			D5381...	13 14 15 N.	122 44 25 E.	88	sm. l. co. s.	Few.
			D5429...	9 41 30 N.	118 50 22 E.	766	gn. m.	Few.
			D5567...	5 48 00 N.	120 33 45 E.	268	62.0	fne. s.	Few.
			D5569...	5 33 15 N.	120 15 30 E.	303	52.3	co. s.	Few.
			D5571...	5 30 45 N.	120 07 57 E.	340	52.3	s., sh.	Few.
15830	U. S. N. M.	1	Siasi Strait, P. I.	Rare.
13155	U. S. N. M.	2	Canagan Island, P. I.	Rare.
15831	U. S. N. M.	2	Tara Island, P. I.	Rare.

ORBITOLITES COMPLANATA Lamarck, var. PLICATA (Dana).

Marginopora vertebralis BLAINVILLE, var. *plicata* DANA, U. S. Exploring Expedition, vol. 7, 1846, p. 706, pl. 60, fig. 9.

Orbitolites laciniatus H. B. BRADY, Quart. Journ. Micr. Sci., vol. 21, 1881, p. 47.

Orbitolites complanata LAMARCK, var. *laciniata* H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 220, pl. 16, figs. 8-11.

Heron-Allen and Earland note the identity of Dana's variety with that of Brady.⁴⁸

It occurs at a few stations in shallow water with the outer annuli containing megalospheric young.

Genus ALVEOLINA d'Orbigny, 1826.

ALVEOLINA BOSCHII (Defrance).

Plate 99, figs. 2-5.

"Alveolites" DEFRANCE, Dict. Sci. Nat., vol. 1, 1816, p. 557.

Oryzaria boschii DEFRANCE, Dict. Sci. Nat., vol. 16, 1820, p. 104.

Alveolina boschii d'ORBIGNY, Ann. Sci. Nat., vol. 7, 1826, p. 306, No. 5, Modèles, 1826, No. 50.—H. B. BRADY, Rep. Voy. *Challenger*, Zoology, vol. 9, 1884, p. 222, pl. 17, figs. 7-12.—CHAPMAN, Journ. Roy. Micr. Soc., 1903, p. 151-153, pls. 2, 3.—CUSHMAN, Bull. 71, U. S. Nat. Mus., pt. 6, 1916, p. 98, pl. 39, fig. 30.

One of the interesting facts of distribution noted in the study of the Philippine material is the abundance of *A. boschii* and the entire

⁴⁸ Trans. Zool. Soc. London, vol. 20, 1915, p. 606.

absence as far as our material shows of *A. melo*. Both species occur off the Hawaiian Islands in shallow water. In the Malay region Millett notes the rarity of *A. melo*, occurring at but two stations, while *A. boscii* was more or less scattered over the area. In the Kerimba Archipelago, off the east coast of Africa, Heron-Allen and Earland note the abundance of *A. melo* and the scarcity in comparison of *A. boscii*. In the Philippine material *A. boscii* is, at the shallow-water stations, especially on the eastern side of the Sulu Sea, very abundant, while I have made no records of *A. melo* in all the material examined. With the data given by Millett it would seem that in its distribution from the Hawaiian Islands to the Red Sea, it does not extend far northward in the Malayan and Philippine regions.

Worthy of special remark is the finding of beautiful microspheric specimens of *A. boscii*, confirming in all details Chapman's observations on the two forms from the Great Barrier Reef of Australia. The two forms are here figured.

The species was not found outside the region of the eastern border of the Sulu Sea except at D5192, in 32 fathoms (59 meters), off northern Cebu, and D5218, in 20 fathoms (37 meters), between Burias and Luzon. This may be due in part to the fact that shallow-water collecting was most extensively carried out in this region. In the Sulu Sea, off western Mindanao, Busilan, Siasi, Jolo, and Tawi Tawi, it occurred in great abundance in material dredged from 10 to 38 fathoms (18 to 70 meters). At one station, D5153, in 49 fathoms (90 meters), off Tawi Tawi, the specimens were smaller; and at D5172 a specimen was obtained in 318 fathoms (582 meters)—an especially deep record for this species.

It was not found in the region to the south, largely for the reason that the material from that region is from much deeper water.

Alveolina boscii—Material examined.


Cat. No.	Coll. of—	No. of specimens.	Station.	Locality.	Depth in fathoms.	Bottom temperature.	Character of bottom.	Abundance.
10788	U.S.N.M.	10+	D5133...	Island off Panabutan Pt., ° N, 52° E., 1.50 miles.	38	° F.	gn. m., s.....	Abundant.
10789	U.S.N.M.	3	D5134...	6 44 45 N.; 121 48 00 E....	25	fine. s.....	Few.
10790	U.S.N.M.	10+	D5148...	5 35 40 N.; 120 47 30 E....	17	co. s.....	Common.
10791	U.S.N.M.	1	D5153...	5 18 10 N.; 120 02 55 E....	49	co. s., sh.....	Rare.
10792	U.S.N.M.	10+	D5154...	5 14 50 N.; 119 53 45 E....	12	co. s.....	Common.
10793	U.S.N.M.	1	D5158...	5 12 00 N.; 119 54 30 E....	12	crs. s., sh.....	Few.
10794	U.S.N.M.	10+	D5159...	5 11 50 N.; 119 54 00 E....	10	co. s.....	Abundant.
10795	U.S.N.M.	10+	D5160...	5 12 40 N.; 119 55 10 E....	12	s.....	Abundant.
10796	U.S.N.M.	1	D5161...	5 10 15 N.; 119 53 00 E....	16	fine. s.....	Few.
10797	U.S.N.M.	1	D5172...	6 03 15 N.; 120 35 30 E....	318	fine. s., sh.....	Rare.
10798	U.S.N.M.	1	D5179...	12 38 15 N.; 122 12 30 E....	37	75.7	hrd. s.....	Rare.
10800	U.S.N.M.	10+	D5192...	11 09 15 N.; 123 50 00 E....	32	gn. s.....	Common.
10799	U.S.N.M.	10+	D5218...	13 11 15 N.; 123 02 45 E....	20	crs. s.....	Common

EXPLANATION OF PLATES.

EXPLANATION OF PLATES.


PLATE 1.

- FIG. 1. *Astrorhiza granulosa*. × 13.
2. *Rhabdammina abyssorum*. × 13.
3. *Rhabdammina abyssorum*, var. *radiata*. × 7.
4. *Rhabdammina linearis*. × 13.
5. *Rhabdammina irregularis*. × 7.
6. *Marsipella gigantea*. × 7.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 490.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 491

PLATE 2.

- FIG. 1. *Bathysiphon filiformis*. $\times 7$.
2. *Bathysiphon rufus*. $\times 7$.
3. *Bathysiphon rufescens*. $\times 12$. *a*, side view; *b*, end view.
4. *Bathysiphon flavidus*, var. *giganteus*. $\times 3$. *a*, side view; *b*, end view.
5. *Bathysiphon papyraceus*. $\times 3$. *a*, side view; *b*, end view.
6. *Rhizammia indivisa*. $\times 27$.
7. *Psammospaera parva*. $\times 66$.
8. *Saccammia sphaerica*, var. *catenulata*. $\times 30$.


PLATE 3.

- FIG. 1. *Tholosina bulla*—hispid form. $\times 33$.
2. *Tholosina bulla*—smooth form. $\times 33$.
3. *Thurammia papyracea*. $\times 33$.
4. *Hyperammia friabilis*. $\times 7$.
5. *Hyperammia elongata*. $\times 13$.
6. *Jaculella obtusa*. $\times 13$.
7. *Jaculella acuta*. $\times 13$.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 492.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.
FOR EXPLANATION OF PLATE SEE PAGE 493.

PLATE 4.

- FIG. 1. *Dendrophya attenuata*. × 8. a, side view; b, end view.
2. *Tolypanmina vagans*. × 25.
3. *Tolypanmina vagans*. × 30.
4. *Tholosina bulla* attached to *Khabdammina irregularis*. × 10.
5. *Saccorhiza ramosa*. × 30.

PLATE 5.

- FIG. 1. *Ammodiscus incertus*. $\times 10$, microspheric form. *a*, side view; *b*, apertural view.
2. *Ammodiscus vicetus*. $\times 10$. Megalospheric form.
3. *Ammodiscus incertus*, var. *discoideus*. $\times 10$. *a*, side view, *b*, apertural view. Microspheric form.
4. *Ammodiscus evsetius*. $\times 13$.



1a



4



1




3a



2



3



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 495.

PLATE 6.


- FIG. 1. *Ammolagena clavata*, microspheric form attached to *Cyclammma*. $\times 7$.
2. *Ammolagena clavata*, megalospheric form with secondary chamber. $\times 7$.
3. *Ammolagena clavata*, megalospheric form attached to *Ammodiscus*. $\times 25$.
4. *Ammolagena clavata*, microspheric form attached to *Clavulina*. $\times 30$.
5. *Aschemonella catenata*. $\times 25$.
6. *Reophax scorpiurus*. $\times 30$.

PLATE 7.

- FIG. 1. *Tolypanmina vagans*. × 15.
2. *Tolypanmina vagans*. × 15.



1



2

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 496



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 497.

PLATE 8.

FIG. 1. *Saquinia dicaricans*. $\times 15$.

182152-20-33

PLATE 9.


FIG. 1. *Sagenina frondescens*, attached to shell. $\times 15$.

2, 3. *Hormosira ovaliformis*. $\times 15$.

4-6. *Ammosphaeroidina grandis*. $\times 7$.




1



2

3



4




5



6

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 498.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 499.


PLATE 10.

Figs. 1, 2. *Reophae nodulosus*. × 15.

3, 4. *Ammolagena clavata*. × 15.


PLATE 11.

- FIG. 1. *Reophax spiculotestus*. × 20.
2. *Haplophragmoides grandiformis*. × 30.
3. *Ammobaculites reophaciformis*. × 30.
4. *Trochammmina globigeriniformis*. × 15, ventral view.
5. *Trochammmina globigeriniformis*. × 15, dorsal view.
6. *Sphaerammina ovalis*. × 15, side view.
7. *Sphaerammina ovalis*. × 15, apertural view.
8. *Sphaerammina ovalis*. × 15, side view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 500.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 501.

PLATE 12.

- FIG. 1. *Reophax pilulifer*. $\times 60$.
2. *Reophax distans*. $\times 30$.
3. *Reophax bacillaris*. $\times 30$.
4. *Reophax dentaliniformis*. $\times 30$.
5. *Reophax nodulosus*. $\times 15$.
6. *Reophax guttifer*. $\times 30$.
7. *Reophax bilocularis*. $\times 30$.


PLATE 13.

- FIG. 1. *Reophax cylindricus*. $\times 15$.
2. *Reophax spiculotestus*. $\times 25$.
3. *Reophax horridus*. $\times 30$.
4. *Reophax findens*. $\times 30$. *a*, front view; *b*, side view.
5. *Hormosina globulifera*. $\times 25$.
6. *Hormosina oricula*. $\times 15$.
7. *Hormosina normani*. $\times 30$.
8. *Haplostiche dubia*. $\times 7$. Megalospheric form, *a*, side view; *b*, apertural view.
9. *Haplostiche dubia*. $\times 7$. Microspheric form, *a*, side view; *b*, apertural view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 502.



1



3



4



2b



5b



2a



5

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 503.

PLATE 14.

- FIG. 1. *Sagenina frondescens*. $\times 15$.
2. *Reophac agglutinatus*. $\times 15$. *a*, front view; *b*, apertural view.
3. *Ammobaculites reophaciformis*. $\times 40$.
4. *Ammobaculites cassis*. $\times 25$.
5. *Sphaerammina ovalis*. $\times 50$. *a*, front view; *b*, apertural view.


PLATE 15.

- FIG. 1. *Haplophragmoïdes subglobosum*. $\times 25$. *a*, apertural view; *b*, side view.
2. *Haplophragmoïdes ringens*. $\times 60$.
3. *Haplophragmoïdes sphaeriloculum*. $\times 30$. *a*, side view; *b*, apertural view.
4. *Trochammina turbinata*. $\times 20$.
5. *Hormosina carpenteri*. $\times 15$.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 504.




1b



1a



2a



2b




5



3b



3a



4a



4b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 505.

PLATE 16.


- FIG. 1. *Cyclammima cancellata*. $\times 18$. *a*, side view; *b*, apertural view.
2. *Cyclammima compressa*. $\times 18$. *a*, side view; *b*, apertural view.
3. *Cyclammima pauciloculata*. $\times 18$. *a* side view; *b*, apertural view.
4. *Cyclammima pusilla*. $\times 18$. *a*, side view; *b*, apertural view.
5. *Ammosphaeroidina grandis*. $\times 15$.

PLATE 17.

- FIG. 1. *Cribrostomoides bradyi*. $\times 25$. *a*, side view; *b*, apertural view.
2. *Trochammina squamata*, attached to *Ammodiscus*. $\times 30$.
3. *Ammobaculites calcareum*. $\times 18$.
4. *Ammobaculites agglutinans*. $\times 25$.
5. *Ammobaculites cylindricus*. $\times 25$.



1b



1a



2



3




4



5

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 506.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 507.

PLATE 18.

- FIG. 1. *Marsipella gigantea*. $\times 15$. Front view.
2. *Marsipella gigantea*. $\times 15$. Apertural view.
3. *Reophax horridus*. $\times 30$. Front view.
4. *Reophax horridus*. $\times 30$. Apertural view.
5. *Haliphysema catenulata*. $\times 15$. Front view.
6. *Haliphysema catenulata*. $\times 15$. Apertural view.
7. *Dendrophrya ramosa*. $\times 15$. Front view.
8. *Dendrophrya ramosa*. $\times 15$. Apertural view.
9. *Hormosira ovaliformis*. $\times 15$. Front view.
10. *Hormosira ovaliformis*. $\times 15$. Apertural view.
11. *Ammosphaerulina adhaerens*. $\times 30$. Interior view.
12. *Ammosphaerulina adhaerens*. $\times 30$. Exterior view.

PLATE 19.

- FIG. 1. *Ammobaculites pseudospirale*. × 18. H4898.
2. *Ammobaculites pseudospirale*. × 18. H4898.
3. *Reophar aduncus*. × 21. D5478.
4. *Clavulina parisiensis*. × 21. D5201.
5. *Bolivina limbata*. × 30. D5178.
6. *Textularia hauerii*. × 24. D5276.
7. *Textularia foliacea*. × 30. D5145. *a*, front view; *b*, side view.



1



4




2



3



5



6




7a



7b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 508.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 509.

PLATE 20.

- FIG. 1. *Spiroplecta bulbosa*. $\times 85$.
2. *Textularia sagittula*. $\times 15$.
3, 4. *Textularia sagittula*. Abnormal specimens. $\times 30$.
5. *Textularia sagittula*, var. *atrata*. $\times 35$.
6. *Textularia sagittula*, var. *fistulosa*. $\times 35$.
7. *Textularia gramei*. $\times 35$.
8. *Textularia agglutinans*. $\times 30$. a, front view; b, apertural view.


PLATE 21.

- FIG. 1. *Textularia stricta*. $\times 15$.
2. *Textularia abbreviata*. $\times 35$. *a*, front view; *b*, apertural view.
3. *Textularia goëssii*. $\times 35$.
4. *Textularia siphonifera*. $\times 75$. *a*, front view; *b*, apertural view; specimen with bifurcating rows of pores.
5. *Textularia siphonifera*. $\times 35$. *a*, front view; *b*, side view.
6. *Textularia siphonifera*. $\times 35$, microspheric specimen with twisted early portion.
7. *Textularia siphonifera*. $\times 35$, specimen with very large openings, two fused.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 510.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 511.

PLATE 22.

FIG. 1. *Textularia porrecta*. $\times 35$.

2. *Textularia transversaria*. $\times 75$. *a*, front view; *b*, side view; *c*, apertural view.

3. *Textularia vertebralis*. $\times 18$. *a*, front view; *c*, apertural view.

4. *Textularia flintii*. $\times 35$.

5. *Textularia pseudocarinata*. $\times 75$.

6. *Trochammina squamata*. $\times 75$. *a*, dorsal view; *b*, ventral view.


PLATE 23.

- FIG. 1. *Tectularia crassisepta*. $\times 75$.
2. *Tectularia quadrilatera*. $\times 75$. *a*, front view; *b*, side view.
3. *Tectularia rugosa*. $\times 35$.
4. *Tectularia rugosa*. $\times 35$. *a*, front view; *b*, apertural view.
5. *Tectularia catenata*. $\times 75$. D5551.
6. *Tectularia aspera*. $\times 35$. D5121.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 512.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 513.

PLATE 24.


- FIG. 1. *Textularia vertebralis*. $\times 23$. *a*, front view; *b*, apertural view.
2. *Textularia semialata*. $\times 50$. Megalospheric form.
3. *Textularia semialata*. $\times 50$. Microspheric form. *a*, front view; *b*, apertural view.
4. *Textularia immensa*. $\times 9$. *a*, front view; *b*, apertural view.
5. *Textularia exarata*. $\times 20$. *a*, front view; *b*, apertural view.

PLATE 25.

- FIG. 1. *Textularia trochus*. $\times 35$. *a*, front view; *b*, ventral view; *c*, dorsal view.
2. *Textularia conica*. $\times 35$. *a*, front view; *b*, ventral view.
3. *Bigenerina pennatula*. $\times 35$. *a*, front view; *b*, apertural view.



1a



2a



1c




2b



3b




1b



3a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 514.



1b



1a



2



3



5



4



6



7

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 515.

PLATE 26.

- FIG. 1. *Bigenerina caprea*. $\times 25$. *a*, front view; *b*, apertural view.
2. *Bigenerina nodosacia*. $\times 35$.
3. *Bolivina amygdaliformis*. $\times 50$.
4. *Bolivina karriana*. $\times 35$.
5. *Bolivina punctata*. $\times 35$.
6. *Bolivina dilatata*. $\times 35$.
7. *Bolivina compacta*. $\times 35$.


PLATE 27.

- FIG. 1. *Bolivina karreviana*, var. *carinata*. $\times 50$.
2. *Bolivina hantkeniana*. $\times 35$.
3. *Bolivina sculpturata*. $\times 50$. *a*, front view; *b*, apertural view.
4. *Verneuilina bradyi*. $\times 75$.
5. *Verneuilina spinulosa*. $\times 75$.
6. *Verneuilina affra*. $\times 35$.
7. *Valvulina conica*. $\times 75$.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 516.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 517.

PLATE 28.

- FIG. 1. *Valvulina fusca*. $\times 60$. *a*, dorsal view; *b*, ventral view.
2. *Tritaxia tricarinata*. $\times 25$. *a*, front view; *b*, apertural view.
3. *Tritaxia indiscreta*. $\times 35$.
4. *Tritaxilina capitata*. $\times 25$. *a*, front view; *b*, apertural view.
5. *Gaudryina scalra*. $\times 35$.


PLATE 29.

- FIG. 1. *Gaudryina flintii*. × 20.
2. *Gaudryina quadrangularis*. × 20.
3. *Gaudryina bradyi*. × 75.
4, 5. *Gaudryina paupercula*. × 35.
6. *Gaudryina baccaia*. × 75.
7. *Gaudryina apicularis*. × 75.
8. *Gaudryina pseudofiliformis*. × 75.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 518.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 519.

PLATE 30.

- FIG. 1. *Gaudryina robusta*. $\times 18$. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Clavulina primacra*. $\times 25$. *a*, front view; *b*, apertural view.
3. *Clavulina primacra*. $\times 25$. Young form still in biserial stage.
4. *Gaudryina attenuata*. $\times 25$.
5, 6. *Clavulina primacra*. $\times 25$. Apertures of biserial stage.
7. *Clavulina primacra*. $\times 25$. Aperture of adult.
8. *Clavulina rotundata*. $\times 25$. *a*, front view; *b*, apertural view.

PLATE 31.

- FIG. 1. *Clavulina communis*. × 35.
2. *Clavulina difformis*. × 50. *a*, front view; *b*, apertural view.
3. *Clavulina angularis*. × 35.
4. *Clavulina bradyi*. × 35.
5. *Bulimina aculeata*. × 75.
6. *Bulimina inflata*. × 75.
7. *Cassidulina lacvigata*. × 75.
8. *Bulimina pupoides*. × 75.
9. *Bulimicula contraria*. × 75.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 520.



3



4



1



6



5



2

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 521.

PLATE 32.

- FIG. 1. *Vernuilina polystropha*, side view. $\times 20$.
2. *Cassidulina subglobosa*, front view. $\times 35$.
3. *Virgulina cornuta*, rear view. $\times 50$.
4. *Virgulina cornuta*, side view. $\times 50$.
5. *Virgulina cornuta*, front view. $\times 50$.
6. *Virgulina cornuta*, apertural view. $\times 50$.


PLATE 33.

- FIG. 1. *Nodosaria (Glandulina) lacvigata*. $\times 50$. D5178.
2. *Nodosaria (Glandulina) lacvigata*, var. *striatula*. $\times 60$. D5236.
3. *Nodosaria pyrula*. $\times 33$. D5201.
4. *Nodosaria pyrula*. $\times 12$. D5201. *a*, apertural view; *b*, front view.
5. *Nodosaria pyrula*, with large proloculum. $\times 66$. D5259.
6, 7. *Nodosaria pyrula*, var. *semirugosa*. $\times 12$.
8. *Nodosaria pyrula*, var. *longi-costata*. $\times 12$. D5388.
9. *Nodosaria pyrula*, var. *longi-costata*. $\times 33$.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 522.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 523.

PLATE 34.

- FIG. 1. *Nodosaria annulata*. × 66.
2 *Nodosaria annulata*. × 66. D5626. *a*, apertural view; *b*, front view.
3. *Nodosaria insecta*. × 33. D5236.
4. *Nodosaria radícula*. × 33. D5586.
5. *Nodosaria soluta*. × 10. D5385. *a*, apertural view; *b*, front view.
6. *Nodosaria soluta*. × 10.
7. *Nodosaria communis*. × 33. D5236.
8. *Nodosaria consobrina*, var. *maciata*. × 20.
9. *Nodosaria filiformis*. × 20.


PLATE 35.

- FIG. 1. *Nodosaria consobrina*, var. *emaciata*. $\times 33$.
2. *Nodosaria japonica*. $\times 10$.
3. *Nodosaria antennula*. $\times 16$. D5236.
4. *Nodosaria subpolygona*. $\times 20$. D5318. *a*, apertural view; *b*, front view, megalospheric.
5. *Nodosaria subpolygona*. $\times 20$. D5318. Micro-spheric.
6. *Nodosaria scalaris*. $\times 66$.
7. *Nodosaria catenulata*. $\times 33$.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 524.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 525.

PLATE 36.

- FIG. 1. *Nodosaria subterrenata*. $\times 33$. D5236.
2. *Nodosaria subscalaris*. $\times 16$. D5178. *a*, apertural view; *b*, front view.
3. *Nodosaria subscalaris*, var. *paucicostata*. $\times 22$. D5133. *a*, apertural view; *b*, front view.
4. *Nodosaria subscalaris*, var. *paucicostata*. $\times 22$. D5133.
5. *Nodosaria milloti*. $\times 33$. D5281.
6. *Nodosaria lepidula*. $\times 33$. D5236.
7. *Nodosaria lepidula*, var. *hispidula*. $\times 33$. D5236.
8. *Nodosaria substriatula*. $\times 33$. D5123. Adult with apertural end broken.
9. *Nodosaria substriatula*. $\times 66$. D5112. Two-chambered stage.
10. *Nodosaria pauciloculata*. $\times 22$. D5201.
11. *Nodosaria pauciloculata*. $\times 22$. D5201.
12. *Nodosaria pauciloculata*. $\times 50$. D5371.


PLATE 37.

- FIG. 1. *Nodosaria sublineata*. × 22. D5586.
2. *Nodosaria perversa*. × 45. D5311.
3. *Nodosaria subperversa*. × 33. D5236.
4. *Nodosaria laevicostata*. × 66. D5178.
5. *Nodosaria raphanus*. × 16. *a*, apertural view; *b*, front view.
6. *Nodosaria raphanus*. × 16. D5272.
7. *Nodosaria raphanus*. × 16.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 526.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 527.

PLATE 38.

- FIG. 1. *Nodosaria obliqua*. × 33.
2. *Nodosaria vertebralis*. × 25.
3. *Nodosaria vertebralis*. × 20.
4. *Nodosaria spirostriolata*. × 12. D5236.
5. *Nodosaria hirsuta*. × 22. D 5201. *a*, apertural view; *b* front view.
6. *Nodosaria hirsuta*. × 20.
7. *Nodosaria hirsuta*, var. *aculeata*. × 33.
8. *Nodosaria hirsuta*, var. *aculeata*. × 33. D5259.


PLATE 39.

- FIG. 1. *Fronicularia philippinensis*. × 20. D5425.
2. *Fronicularia philippinensis*. × 20. D5429.
3. *Fronicularia philippinensis*. × 20. D5201.
4. *Fronicularia plicata* (?). × 12. D5584.
5. *Fronicularia annularis*. × 20. D5382.
6. *Fronicularia annularis*, var. *longistriata*. × 33. D5110.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 628.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 529.

PLATE 40.

- FIG. 1. *Vaginulina acicula*. D5613. *a*, $\times 30$; *b*, initial end. $\times 100$.
2. *Nodosaria vertebralis*. $\times 25$. H4898.
3. *Vaginulina protumida*. $\times 25$. D5236. *a*, apertural view; *b*, front view.
4. *Fronicularia plicata*. $\times 20$. D5301.
5. *Fronicularia inaequalis*. $\times 66$. D5123.
6. *Fronicularia inaequalis*. $\times 33$. D5652.


PLATE 41.

- FIG. 1. *Marginulina glabra*. × 66.
2. *Marginulina striatula*. × 66.
3. *Vaginulina legumen*. × 33.
4. *Vaginulina legumen*, var. *elegans*. × 33.
5. *Marginulina costata*, megalospheric form. × 33.
6. *Marginulina costata*, megalospheric form. × 22.
7. *Marginulina costata*, microspheric form. × 22.
8. *Marginulina costata*, microspheric form. × 33.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 530.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 531

PLATE 42.

- FIG. 1. *Vaginulina spinigera*. $\times 33$.
2. *Vaginulina margaritifera*. $\times 25$.
3. *Vaginulina bradyi*. $\times 25$.
4. *Vaginulina bradyi*. $\times 25$. *a*, apertural view; *b*, front view.
5. *Vaginulina linearis*, microspheric form. $\times 50$. D5569.
6. *Vaginulina linearis*, microspheric form. $\times 33$. D5172.
7. *Vaginulina linearis*, microspheric form. $\times 50$. D5569.
8. *Vaginulina linearis*, megalospheric form. $\times 33$. D5133.

PLATE 43.


- FIG. 1. *Lingulina grandis*. *a*, front view, $\times 12$; *b*, apertural view, $\times 16$.
2. *Lingulina costata*. *a*, front view, $\times 26$; *b*, apertural view, $\times 26$.
3. *Cristellaria rotulata*, var. *umbonata*. $\times 20$.




1b



2b



1a



2a




3

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 532.



1



2



3



4a



4b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 533.

PLATE 44.

- FIG. 1. *Cristellaria orbicularis*, var. *subearinata*. $\times 12$.
2. *Cristellaria orbicularis*, var. *papillata*. $\times 12$. D5152.
3. *Cristellaria orbicularis*, var. *subumbonata*. $\times 12$. D5259.
4. *Cristellaria convergens*. a, side view, $\times 26$; b, edge view, $\times 26$.

PLATE 45.

- FIG. 1. *Cristellaria gibba*. $\times 20$. *a*, side view; *b*, edge view.
2. *Cristellaria calcar*. $\times 33$. *a*, side view; *b*, edge view.
3. *Cristellaria calcarata*. $\times 26$.
4. *Cristellaria echinata*. $\times 23$. *a*, side view; *b*, edge view. D5423.




1a



1b




2b



4a



4b



3



2a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 534.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 535.

PLATE 46.


- FIG. 1. *Cristellaria echinata*. $\times 23$. *a*, side view; *b*, edge view. D5374.
2. *Cristellaria expansa*. $\times 26$. *a*, side view; *b*, edge view. D5467.
3. *Cristellaria expansa*, var. *planulata*. $\times 16$. *a*, side view; *b*, edge view.
4. *Cristellaria costata*. $\times 33$. D5590.

PLATE 47.

FIG. 1. *Cristellaria costata*. × 33.

2, 3. *Cristellaria costata*, var. *multicosta*. × 26.

4, 5. *Cristellaria costata*, var. *subdecorata*. × 26.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 536.



1a



2a



1b



2b



3a



3c



3b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 537.

PLATE 48.

- FIG. 1. *Cristellaria bradyi*. $\times 13$. *a*, side view; *b*, edge view. D5313.
2. *Cristellaria cassinoides*. $\times 26$. *a*, side view; *b*, edge view. D5381.
3. *Cristellaria dorso-costata*. $\times 13$. *a*, side view; *b*, edge view; *c*, apertural view. D5268.

PLATE 49.

- FIG. 1. *Cristellaria costata*, var. *sublaevis*. $\times 26$. D5374.
2. *Cristellaria cassis*. $\times 26$. *a*, side view; *b*, edge view.
3. *Cristellaria paucicostata*. $\times 40$. *a*, side view; *b*, edge view. D5586.
4. *Cristellaria tumido-costata*. $\times 23$. *a*, side view; *b*, edge view. D5570.




FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 538.




1a



2




1b



3a



3b



4



5

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 539.

PLATE 50.

- FIG. 1. *Cristellaria tumulo-costata*, var. *labyrinthica*. × 23. *a*, side view; *b*, edge view. D5567.
2. *Cristellaria tenuis*. × 50.
3. *Cristellaria tricarinella*. × 23. *a*, side view; *b*, edge view. D5301.
4, 5. *Cristellaria crepidula*. × 33.


PLATE 51.

- FIG. 1. *Cristellaria helicinoïdes*. $\times 26$. *a*, side view; *b*, reverse view. D 5301.
2. *Cristellaria italica*. $\times 26$.
3. *Cristellaria italica*, var. *acuto-carinata*. $\times 26$. *a*, side view; *b*, end view.
D5523.
4. *Cristellaria helicina*. $\times 16$. *a*, side view; *b*, edge view. D5272.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 540.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 541.

PLATE 52.

- FIG. 1, 2. *Polymorphina problema*, var. *indica*. × 20. D5579.
3. *Polymorphina oblonga*. × 50.
4. *Polymorphina lactea*, var. *oblonga*. × 66. D5601.
5. *Polymorphina*, species. × 50.
6. *Lagena apiculata*. × 33.
7. *Nodosaria substriatula*. × 33. D5123.
8. *Nodosaria substriatula*. × 100, showing details of neck and aperture.
9. *Nodosaria substriatula*. × 100, aperture.

PLATE 53.

- FIG. 1. *Polymorphina problema*, var. *indica*, short form. $\times 23$. D5579. *a*, side view; *b*, apertural view.
2. *Marginulina philippinensis*. $\times 33$. D5217.
3. *Marginulina philippinensis*. $\times 13$. D5220. *a*, side view; *b*, edge view.



1a



1b



2




3b



3a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 512



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 543.

PLATE 54.

- FIG. 1. *Polymorphina elegantissima*. × 66.
2. *Polymorphina elegantissima*. × 66.
3. *Polymorphina problema*. × 66. a, apertural view; b, front view.
4. *Polymorphina problema*. × 66.
5. *Urigerina asperula*. × 66.
6. *Urigerina striata*. × 66.


PLATE 55.

- FIG. 1. *Uvigerina pigma*. × 33.
2. *Uvigerina tenuistriata*. × 66.
3. *Uvigerina schwageri*. × 66.
4. *Uvigerina schwageri*. × 66.
5. *Uvigerina schwageri*. × 40. *a*, apertural view; *b*, front view
6. *Uvigerina aculeata*. × 66.
7. *Uvigerina ampullacea*. × 66.
8. *Uvigerina asperula*. × 66.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 544.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 545.

PLATE 56.

- FIG. 1. *Siphogenerina columellaris*. × 66.
2. *Siphogenerina bifrons*, twinned specimen. × 66.
3. *Siphogenerina bifrons*. × 66.
4. *Siphogenerina bifrons*, var. *striatula*. × 66.
5. *Siphogenerina striata*. × 66.
6. *Siphogenerina raphanus*, var. *costulata*. × 66. *a*, apertural view; *b*, front view.
7. *Siphogenerina raphanus*. × 66.
8. *Siphogenerina dimorpha*. × 66.

PLATE 57.

- FIG. 1. *Candiana nitida*, var. *triloba*. $\times 26$. D5191. *a-c*, three views.
2. *Cristellaria articulata* (?) $\times 26$. *a*, side view; *b*, apertural view.
3. *Cristellaria articulata* (?). $\times 26$, with broad cristellarian base.
4. *Cristellaria articulata* (?). $\times 46$. D5110. Very elongate, irregular specimen.
5. *Chilostomella grandis*. $\times 20$. D5449. *a*, front view; *b*, side view; *c*, apertural view.



1a



1b



1c



2a



2b



4



3



5a




5c




5b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 546.




1a




1b



3b



1c



2b



3a



2c



2a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.




FOR EXPLANATION OF PLATE SEE PAGE 547.

PLATE 58.

1. *Discorbis altomorphinoides*. × 66. D5261. *a*, dorsal view; *b*, ventral view; *c*, side view.
2. *Pulvinulina philippinensis*. √ 66. *a*, dorsal view; *b*, ventral view; *c*, side view.
3. *Pulvinulina scabra*. × 66. D5220. *a*, ventral view; *b*, side view.


PLATE 59.

- FIG. 1. *Discorbis bertheloti*. $\times 25$. D5268. *a*, dorsal view; *b*, ventral view; *c*, side view.
2. *Cymbalopora poeyi*. $\times 50$. D5469. *a*, dorsal view; *b*, ventral view; *c*, side view.
3. *Tretomphalus bulloides*. $\times 50$. D5576. *a*, dorsal view; *b*, ventral view; *c*, side view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 548.




FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.
FOR EXPLANATION OF PLATE SEE PAGE 549.

PLATE 60.

- FIG. 1. *Discorbis*, species. $\times 70$. D5201. *a*, dorsal view; *b*, ventral view; *c*, side view.
2. *Discorbis rugosa*. $\times 35$. D5236. *a*, dorsal view; *b*, ventral view; *c*, side view.
3. *Siphonina reticulata*. $\times 70$. D5178. *a*, dorsal view; *b*, ventral view; *c*, side view.

PLATE 61.


- FIG. 1. *Anomalina ammonoides*. $\times 66$. 5259. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Anomalina coronata*. $\times 33$. D5236. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Anomalina polymorpha*. $\times 33$. D5318. *a*, ventral view; *b*, dorsal view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.
FOR EXPLANATION OF PLATE SEE PAGE 550.




1a




1b



3



2a



2b



4a



4c



4b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 551.

PLATE 62.

- FIG. 1. *Anomalina polymorpha*, var. *cervicornis*. $\times 33$. D5236. *a*, ventral view
b, dorsal view.
2. *Anomalina polymorpha*, var. *siphonifera*. $\times 33$. D5236. *a*, ventral view
b, dorsal view.
3. *Anomalina grosserugosa*. $\times 33$. D5236.
4. *Truncatulina culter*. $\times 66$. D5635. *a*, ventral view; *b*, dorsal view; *c*, aper-
tural view.

PLATE 63.

- FIG. 1. *Truncatulina refulgens*. × 66. D5469. *a*, ventral view; *b*, dorsal view
c, apertural view.
2. *Truncatulina lobatula*. × 66. D5590. *a*, ventral view; *b*, dorsal view;
c, apertural view.
3. *Truncatulina akneriana*. × 66. D5445. *a*, ventral view; *b*, dorsal view;
c, apertural view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 552.



1a



1c




1b



2b




2c



2a



3a



3c



3b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 553.

PLATE 64.

- FIG. 1. *Truncatulina wuellerstorfi*. $\times 33$. D5445. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Truncatulina tenera*. $\times 66$. D5613. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Truncatulina haidingerii*. $\times 66$. D5589. *a*, ventral view; *b*, dorsal view; *c*, apertural view.

PLATE 65.

- FIG. 1. *Truncatulina margaritifera*. × 33. D5338. *a*, ventral view; *b*, dorsal view;
c, apertural view.
2. *Truncatulina variabilis*. × 33. D5546.
3. *Truncatulina ungeriana*. × 33. D5640. *a*, ventral view; *b*, dorsal view;
c, apertural view.




1b




1c



1a



2



3a




3c



3b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 564.




1a




1c



1b




2a



2c




2b



3b



3c



3a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 555.

PLATE 66.

- FIG. 1. *Pulvinulina menardii*. $\times 33$. D5571. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Pulvinulina menardii*, var. *fimbriata*. $\times 66$. D5638. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Pulvinulina tumida*. $\times 33$. D5637. *a*, ventral view; *b*, dorsal view; *c*, apertural view.

PLATE 67.

- FIG. 1. *Pulvinulina canariensis*. $\times 66$. D5606. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Pulvinulina truncatulinoides*. $\times 66$. D5300. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Pulvinulina crassa*. $\times 66$. D5580. *a*, ventral view; *b*, dorsal view; *c*, apertural view.




1b




1c




1a



2c




2b



2a



3b




3c



3a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 556.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 557.

PLATE 68.

- FIG. 1. *Pulvinulina punctulata*. $\times 33$. D5642. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Pulvinulina pauperata*. $\times 22$. D5439.
3. *Pulvinulina exigua*. $\times 66$. D5439. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
4. *Pulvinulina concentrica*. $\times 66$. D5572. *a*, ventral view; *b*, dorsal view; *c*, apertural view.


PLATE 69.

- FIG. 1. *Pulvinulina procera*. × 33. D5580. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Pulvinulina lateralis*. × 33. D5580. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Pulvinulina auricula*. × 66. D5465. *a*, ventral view; *b*, dorsal view; *c*, apertural view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 558.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 559.

PLATE 70.

- FIG. 1. *Pulvinulina berthelotiana*, var. *subornata*. $\times 33$. D5301. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Discorbis subobtusa*. $\times 66$. D5381. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Rotalia beccarii*. $\times 66$. D5469. *a*, ventral view; *b*, dorsal view; *c*, apertural view.

PLATE 71.

- FIG. 1. *Pulvinulina umbonata*. $\times 66$. D5468. *a*, ventral view; *b*, dorsal view; *c*, apertural view. "
2. *Rotalia soldanii*. $\times 33$. D5639. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Rotalia calcar*. $\times 50$. D5630. *a*, ventral view; *b*, dorsal view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 560.



1b



1a




1c




2b



2a



2c



3a



3b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 561.

PLATE 72.

- FIG. 1. *Rotalia broeckhiana*. $\times 50$. D5637. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
2. *Rotalia papillosa*, var. *compressiuscula*. $\times 33$. *a*, ventral view; *b*, dorsal view; *c*, apertural view.
3. *Rotalia papillosa*. $\times 33$. *a*, dorsal view; *b*, apertural view.

PLATE 73.


- FIG. 1. *Rotalia schroeteriana*. × 12. D5338. *a*, dorsal view; *b*, ventral view; *c*, side view.
2. *Carpenteria proteiformis*. × 15.
3. *Carpenteria proteiformis*. × 15.
4. *Carpenteria utricularis*. × 15.
5. *Carpenteria utricularis*. × 15.



1a



1b



1c



2



3



4



5

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.
FOR EXPLANATION OF PLATE SEE PAGE 562.



1a




1b



1c



2b



2a



3b



3a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 563.

PLATE 74.

- FIG. 1. *Truncatulina margaritifera*. $\times 23$. *a*, dorsal view; *b*, ventral view; *c*, apertural view.
2. *Operculina bartschi*, var. *ornata*. $\times 10$. *a*, side view; *b*, apertural view.
3. *Nonionina umbilicatula*. $\times 65$. D5236. *a*, side view; *b*, apertural view.


PLATE 75.

- FIG. 1. *Calcarina splengleri*, ventral view. $\times 20$.
2. *Calcarina defranci*, ventral view. $\times 20$.
3. *Calcarina baculatus* (?), ventral view. $\times 20$.
4. *Calcarina hispida*, ventral view. $\times 15$.
5. *Siderolites tetraedra*. $\times 15$.
6. *Baculogypsina sphaerulatus*. $\times 20$.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 564.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 565.

PLATE 76.

FIGS. 1-5. *Siderolites tetradra*.

1. Young, ventral view. $\times 45$.
2. Later stage, chambers invading dorsal side. $\times 40$. *a*, ventral side; *b*, dorsal side.
3. Still later stage where dorsal side is partly covered by the invading chambers. $\times 35$. *a*, ventral view; *b*, dorsal view.
4. Old eroded specimen, in which the chambers are largely broken away, leaving the raised bosses (*c*) as projections from the center. $\times 18$.
5. Specimens in which the chambers now cover even the spines. $\times 15$.

PLATE 77.

- FIG. 1. *Cornuspira foliacea*. × 20. D5424.
2. *Cornuspira foliacea*, var. *expansa*. × 20. D5259.
3. *Cornuspira involvens*. × 20. D5198.
4. *Cornuspira involvens*. × 20. D5256.
5. *Cornuspira involvens*, var. *striatula*. × 30. D5282.
6. *Cornuspira carinata*. × 20. D5523.



1



2



3



4




5



6

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.
FOR EXPLANATION OF PLATE SEE PAGE 566.



1




2



3



4a



4b



5

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 567.

PLATE 78.

- FIG. 1. *Cornuspira lacunosa*. $\times 20$. D5259.
2. *Ophthalmidium inconstans*. $\times 20$. Adult.
3. *Ophthalmidium inconstans*. $\times 50$. Young.
4. *Spiroloculina grateloupi*. $\times 20$. D5179. *a*, front view; *b*, apertural view.
5. *Spiroloculina grateloupi*, var. *incisa*. $\times 20$. D5369.

PLATE 79.


- FIG. 1. *Spiroloculina costifera*. × 18. D5236. *a*, front view; *b*, apertural view.
2. *Spiroloculina robusta*. × 18. D5650. *a*, front view; *b*, apertural view.
3. *Spiroloculina validus*. × 18. D5668. *a*, front view; *b*, apertural view.



1a



2b



1b



2a



3a



3b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 568.



1b



3b




1a




3a




4b



2



4a



5b



5a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 569.

PLATE 80.


- FIG. 1. *Spiroloculina circularis*. $\times 23$. D5259. *a*, front view; *b*, apertural view.
2. *Spiroloculina circularis*. $\times 66$. D5259. Aperture.
3. *Spiroloculina canaliculata*. $\times 23$. D5268. *a*, front view; *b*, apertural view.
4. *Spiroloculina elegans*. $\times 23$. D5143. *a*, front view; *b*, apertural view.
5. *Spiroloculina planissima*. $\times 66$. D5342. *a*, front view; *b*, apertural view.

PLATE 81.

- FIG. 1. *Spiroloculina scrobiculata*. $\times 30$. D5181. *a*, front view; *b*, apertural view.
2. *Spiroloculina depressa*. $\times 30$. D5217. *a*, front view; *b*, apertural view.
3. *Spiroloculina arcuaria*. $\times 30$. D5110. *a*, front view; *b*, apertural view.
4. *Spiroloculina antillarum*. $\times 30$. D5218. *a*, front view; *b*, apertural view.
5. *Spiroloculina antillarum*, var. *angulata*. $\times 30$. D5218. *a*, front view; *b*,
apertural view.
6, 7. *Spiroloculina millettii*. $\times 30$. Tacloban Anchorage.




1b



2b



1a



3b



2a



3a



4b



4a



6



7




5b



5a

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 570.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 571

PLATE 82.

- FIG. 1. *Spiroloculina tenuiscriptata*. × 27. D5238. *a*, face view; *b*, apertural view.
2. *Spiroloculina tenuiscriptata*. × 21. D5236. Starting to uncoil.
3. *Spiroloculina tenuiscriptata*. × 21. D5236. Later stage.
4. *Spiroloculina convexiuscula*. × 21. *a*, face view; *b*, apertural view.
5-7. *Spiroloculina costifera*, var. *plena*, new variety. × 18. D5236. Various stages in the development.


PLATE 83.

- FIG. 1. *Spiroloculina advena*. $\times 30$. D5178. *a*, convex side; *b*, concave side; *c*,
apertural view.
2. *Spiroloculina affixa*. $\times 48$. D5192. *a*, convex side; *b*, concave side; *c*, aper-
tural view.
3. *Spiroloculina orbis*. $\times 30$. D5178.
4. *Spiroloculina antillarum*, variety. $\times 30$. D5218.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 572.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 573.

PLATE 84.

- FIG. 1. *Cornuspira crassisepta*. $\times 56$.
2. *Spiroloculina tenuissima*. $\times 82$. D5227.
3. *Spiroloculina tenuimargo*. $\times 82$. D5349. Abnormal specimen.
4. *Quinqueloculina crassatina*. $\times 26$. D5178. *a*, side view; *b*, reverse side; *c*,
apertural view.
5. *Quinqueloculina recta*. $\times 26$. D5570. *a*, side view; *b*, reverse side; *c*, aper-
tural view.
6. *Quinqueloculina fusca*. $\times 26$. Tara Island, Philippine Islands. *a*, side
view; *b*, reverse side; *c*, apertural view.


PLATE 85.

- FIG. 1. *Planispirina contraria*. $\times 23$. D5582. *a*, side view; *b*, apertural view.
2. *Nodobacularia tibia*. $\times 23$. Specimen with the proloculum broken open, showing interior.
3. *Nodobacularia tibia*. $\times 23$. Exterior of another specimen.
4. *Quinqueloculina albatrossi*. $\times 23$. D5133. *a*, side view; *b*, opposite side; *c*, apertural view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 574.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 575.

PLATE 86.

- FIG. 1. *Quinqueloculina disparilis*. $\times 22$. D5178. *a*, side view; *b*, reverse side; *c*, apertural view.
2. *Quinqueloculina bicarinata*. $\times 30$. D5134. *a*, side view; *b*, reverse side; *c*, apertural view.
3. *Quinqueloculina bicarinata*. $\times 116$. D5143. Aperture of early stage with simple tooth.
4. *Quinqueloculina parkeri*. $\times 30$. D5160. *a*, side view; *b*, reverse side; *c*, apertural view.


PLATE 87.

- FIG. 1. *Quinqueloculina vulgaris*. $\times 23$. D5227. *a*, side view; *b*, opposite side; *c*, apertural view.
2. *Quinqueloculina lamarckiana*. $\times 23$. D5201. *a*, side view; *b*, opposite side; *c*, apertural view.
3. *Quinqueloculina lamarckiana*. $\times 20$. D5220. *a*, side view; *b*, opposite side; *c*, apertural view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 576.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 577.

PLATE 88.

- FIG. 1. *Quinqueloculina apicula*. $\times 18$. D5236. *a*, side view; *b*, opposite side
c, apertural view.
2. *Quinqueloculina apicula*. $\times 18$. D5236. Outline of specimen with the
spine greatly developed.
3. *Quinqueloculina procera*. $\times 12$. D5236. *a*, side view; *b*, opposite side;
c, apertural view.
4. *Quinqueloculina seminulum*. $\times 21$. *a*, side view; *b*, opposite side; *c*,
apertural view.


PLATE 89.

- FIG. 1. *Quinqueloculina kerimbatica*, var. *reticulo-striata*. $\times 26$. D5143. *a*, side view; *b*, reverse side; *c*, apertural view.
2. *Quinqueloculina kerimbatica*, var. *philippinensis*. $\times 26$. D5192. Four-chambered stage. *a*, side view; *b*, reverse side; *c*, apertural view.
3. *Quinqueloculina kerimbatica*, var. *philippinensis*. $\times 26$.
4. *Quinqueloculina crassa*, var. *subcuneata*. $\times 26$. D5151. *a*, side view; *b*, reverse view; *c*, apertural view.
5. *Quinqueloculina bancana*. $\times 26$. D5358.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 578.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 579.

PLATE 90.

- FIG. 1. *Quinqueloculina contorta*. $\times 66$. D5572. *a*, side view; *b*, reverse side;
c, apertural view.
2. *Quinqueloculina ferussacii*. $\times 20$. D5164. *a*, side view; *b*, reverse side;
c, apertural view.
3. *Quinqueloculina polygona*. $\times 66$. D5181. *a*, side view; *b*, reverse side;
c, apertural view.


PLATE 91

- FIG. 1. *Quinqueloculina agglutinans*. × 23. D5106. *a*, side view; *b*, reverse side; *c*, apertural view.
2. *Quinqueloculina venusta*. × 23. D5147. *a*, side view; *b*, reverse side; *c*, apertural view.
3. *Quinqueloculina sculpturata*. × 23. D5145. *a*, side view; *b*, reverse side; *c*, apertural view.
4. *Quinqueloculina sculpturata*. × 23. D5143.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 550.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 581.

PLATE 92.

- FIG. 1. *Triloculina circularis*. $\times 39$. D5236. *a*, side view; *b*, opposite side; *c*, apertural view.
2. *Triloculina circularis*. $\times 75$. D5236. Aperture with tooth.
3. *Triloculina oblonga*. $\times 39$. D5214. *a*, side view; *b*, opposite side; *c*, apertural view.
4. *Triloculina cuneata*. $\times 75$. *a*, side view; *b*, opposite side; *c*, apertural view.

PLATE 93.

- FIG. 1. *Triloculina cuneata*, var. *incisa*. $\times 26$. Siasi Island. *a*, side view; *b*, opposite side; *c*, apertural view.
2. *Triloculina rupertiana*. *a*, side view; *b*, opposite side; *c*, edge view; *d*, apertural view; $\times 13$. *e*, portion of surface, $\times 80$. D5134.



1a




1c




1b



2d



2e



2a



2c



2b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 582.



1a



1b



1c




3b



2




3a



4a



4c



4b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 583.

PLATE 94.

- FIG. 1. *Flintina triquetra*. × 10. D5236. *a* and *b*, opposite sides; *c*, apertural view.
2. *Flintina bradyana*. × 20. D5134.
3. *Massilina arenaria*. × 10. D5236. *a*, side view; *b*, apertural view.
4. *Massilina durrandii*. × 20. *a* and *b*, opposite sides; *c*, apertural view.




PLATE 95.

- FIG. 1. *Biloculina irregularis*. $\times 30$. *a*, front view; *b*, side view.
2. *Biloculina globulus*. $\times 30$. D5580. *a*, front view; *b*, apertural view.
3. *Biloculina serrata*. $\times 20$. D5639. *a*, front view; *b*, apertural view.
4. *Biloculina elongata*. $\times 20$. D5613. *a*, front view; *b*, apertural view.
5. *Biloculina vespertilio*. $\times 15$. D5277. *a*, front view; *b*, apertural view.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 584.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 585.

PLATE 96.


- FIG. 1. *Biloculina anomala*. $\times 50$. D5622. *a*, front view; *b*, side view; *c*, apertural view.
2. *Biloculina depressa*. $\times 10$. D5236. *a*, front view; *b*, side view.
3. *Biloculina comata*. $\times 17$. D5567. *a*, front view; *b*, apertural view.

PLATE 97.

- FIG. 1. *Biloculina sarsii*. $\times 24$. D5315. *a*, front view; *b*, side view; *c*, apertural view.
2. *Biloculina lucernula*. $\times 18$. D5236. Slightly triloculine form. *a*, side view; *b*, opposite side; *c*, apertural view.
3. *Operculina elegans*. Side view, $\times 20$. From station D5579.
4. *Vertebralina striata*, var. *reticulosa*. Side view, $\times 60$. From station D5160.



1a




1b



1c



2a



2b



2c




3




4

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.


FOR EXPLANATION OF PLATE SEE PAGE 586.



1a




1b



1c



2a



2b



2c



3a



3b

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 587.

PLATE 98.


- FIG. 1. *Biloculina lucernula*. $\times 24$. D5630. *a*, front view; *b*, side view; *c*, apertural view.
2. *Biloculina denticulata*, var. *striolata*. $\times 18$. D5139. *a*, front view; *b*, side view; *c*, apertural view.
3. *Biloculina denticulata*, var. $\times 18$. D5179. *a*, front view; *b*, apertural view.

PLATE 99.

- FIG. 1. *Nevillina coronata*. × 12. D5160. *a*, front view; *b*, basal view.
2. *Alveolina boscii*. × 8. Exterior of megalospheric form.
3. *Alveolina boscii*. × 12. Transverse section of megalospheric form.
4. *Alveolina boscii*. × 12. Transverse section of microspheric form.
5. *Alveolina boscii*. × 5. Exterior of microspheric form.



1a




1b




5



2




4



3

FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 588.



FORAMINIFERA OF THE PHILIPPINE AND ADJACENT SEAS.

FOR EXPLANATION OF PLATE SEE PAGE 589.

PLATE 100.

- FIG. 1. *Quinqueloculina curta?* × 33. D5134. 2-chambered stage. *a*, apertural view; *b*, side view.
2. *Quinqueloculina curta?* × 33. D5134. Central chambers by transmitted light.
3. *Spiroloculina grateloupi*. × 165. D5159. Central chambers by transmitted light.
4. *Spiroloculina depressa*. × 165. D5220. Central chambers by transmitted light.
5. *Spiroloculina depressa*. × 165. D5220. Central chambers by transmitted light.
6. *Quinqueloculina rugosa*. × 33. *a*, side view; *b*, opposite side; *c*, apertural view.
7. *Quinqueloculina bicarinata*. × 33. *a*, side view; *b*, opposite side; *c*, apertural view.

INDEX.

	Page.		Page.
abbreviata, Textularia	122	Ammobaculites cassis	91
abyssorum, radiata, Rhabdammina	37	cylindricus	92
Rhabdammina	36	pseudospirale	94
Rhabdopleura	39	reophaciformis	92
acaulis, Bolivina	129	tenuimargo	90
acervalis, Planorbulina	311	Ammochilostoma	97
acicula, Vaginulina	262	ringens	82
acicularis, Nautilus	483	rotundata	97
aculeata, Bulimina	161	Ammodiscinae	61
presli	161	Ammodiscus	62
Cristellaria	235	exsertus	64
Nodosaria hirsuta	214	incertus	62
Uvigerina	273	discoideus	63
acuta, Jaculella	59	tenuis	62
acuticosta, Lagena	180	Ammoglobigerina bulloides	285
acutimargo, Spiroloculina	398, 410	Ammolagena	61
acuto-carinata, Cristellaria italica	253	clavata	61
adhaerens, Ammosphaerulina	100	ammonoides, Anomalina	323
adunea, Orbiculina	483	Discorbis	323
Reophax	70	Nautilus	382
aduncus, Nautilus	483	Operculina	382
Reophax	70	Planorbulina	323
advena, Spiroloculina	410	Rosalina	323
Triloculina	453	Ammosphaeroidina	98
aenariensis, Bolivina	137	grandis	98
Brizalina	137	sphaeroidiniformis	98
aequilateralis, Globigerina	293	Ammosphaerulina	100
involuta	294	adhaerens	100
affinis, Bulimina	165	Amphistegina	370
affixa, Spiroloculina	410	cumingii	383
Verneuilina	142	lessonii	370
agglutinans, Ammobaculites	89	madagascariensis	372
fistula, Textularia	107	radiata	372
Haplophragmium	89, 92	radiata	372
nodosaria, Textularia	125	Amphorina gracilis	175
porrecta, Textularia	109	gracillima	175
Quinqueloculina	441, 464	ampullacea, Uvigerina	274
Spirolina	89, 92	asperula	274
Textularia	113	amygdalaeformis, Bolivina	133
agglutinata, Astrorhiza	36	angularis, Clavulina	155
agglutinatus, Reophax	73	Valvulina	155
akneriana, Rotalina	316	angulata, Spiroloculina antillarum	408
Truncatulina	316	grata	408
alata, Bolivina	129	angulosa, Astrorhiza	35
beyrichi	129	Uvigerina	271
Vulvulina	129	annularis, Frondicularia	217
albatrossi, Quinqueloculina	435	longistriata, Frondicularia	218
algaeformis, Rhizammina	46	annulata, Glandulina	189
Allomorphina	281	Nodosaria	189
trigona	284	radicula	189
allomorphinoides, Discorbina	306	anomala, Biloculina	474
Discorbis	306	Anomalina ammonoides	323
Valvulina	306	ariminensis	322
alternans, Pleurostomella	138	bengalensis	320
alveolata, substriata, Lagena	182	coronata	326
Alveolina	487	grosserugosa	323
boscii	487	polymorpha	324
alveoliniformis, Miliolina	443	cervicornis	325
Quinqueloculina	443	siphonifera	325
Ammobaculites	89	wuellerstorfi	314
agglutinans	89	antennula, Nodosaria	197
calcareum	90	antillarum, angulata, Spiroloculina	408

	Page.		Page.
antillarum, Spiroloculina	407	bertheliniana, Miliolina	757
aperturalis, Textularia	111	triearinata	457
apicula, Quinqueloculina	422	Triiloculina	457
apicularis, Gaudryina	151	Bertheloti, Discorbina	305
apiculata, Lagena	176	Discorbis	305
sulcata	180	Rosalina	305
Oolina	176	berthelotiana, subornata, Pulvinulina	333
arenacea, Bigenerina	126	beyrichi, alata, Bolivina	129
arenaria, Massilina	446	Bolivina	128
Spiroloculina	411, 446	bicarinata, Fissurina	184
arictinus, Nautilus (Lituus)	481, 482	Lagena	184
Peneroplis	482	Quinqueloculina	428
pertusus	482	bicornis, elegans, Miliolina	425
ariminensis, Anomalina	322	Quinqueloculina	425
Planulina	322	bifrons, Sagrina	277
articulata, Cristellaria	222	Siphogenerina	277
Robulina	222	(Sagrina)	277
Articulina	448	striatula, Siphogenerina	278
sagra	448	Bigenerina	125
Aschemonella	64	arenacea	126
catenata	64	capreolus	127
Aschemonellinae	64	nodosaria	125
aspera, Textularia	112	pennatula	127
asperula, ampullacea, Uvigerina	274	bilocularis, Reophax	74
Massilina	447	Biloculina	469
Spiroloculina	447	anomala	474
Uvigerina	274	comata	477
Assilina discoidalis	379	contraria	412
Astrorhiza	35	denticulata	476
agglutinata	36	striolata	477
angulosa	35	depressa	469
catenata	64	elongata	473
granulosa	35	globulus	474
Astrorhizidae	35	irregularis	479
Astrorhizinae	35	lucermula	475
atrata, Textularia sagittula	103	milletii	479
attenuata, Dendrophrya	57	murrhyna	470
Gaudryina	152	pisum	475
auricula, Nautilus	329	ringens	471, 472
Pulvinulina	329	denticulata	476
auriculata, substriata, Lagena	182	striolata	477
austriaca, Valvulina	144	sarsii	471
triangularis	144	serrata	471
baccata, Gaudryina	150	tubulosa	475
bacillaris, Reophax	67	vespertilio	472
baculatus, Calcarina	353	Bolivina	128
Tinoporos	353	acaulis	129
Baculogypsina	359	aenariensis	137
sphaerulatus	359	alata	129
balaniformis, proteriformis, Carpenteria	361	amygdalaeformis	133
bartschi, Flintina	465	beyrichi	128
Operculina	376	alata	129
ornata, Operculina	378	compacta	137
plana, Operculina	377	costata	134
Bathysiphon	41	dilatata	128
filiformis	41	hantkeniana	132
flavidum	44	karreriana	131
flavidus	44	carinata	132
giganteus	41	limbata	135
papyraceus	45	nobilis	130
rufum	42	punctata	136
rufus	42	pusilla	138
rufuscens	43	pygmaea	136
beccarii, Nautilus	345	quadrilatera	125
Rotalia	345	robusta	129
(Turbinulina)	345	compacta	137
bengalensis, Anomalina	320	schwageriana	130

	Page.		Page.
Bolivina sculpturata.....	134	calcar, Nautilus.....	231, 233
semicostata.....	134	Robulina.....	231
subangularis.....	135	Rotalia.....	350
boscii, Alveolina.....	487	calcarata, Cristellaria.....	232
Oryzaria.....	487	calcareum, Ammobaculites.....	89
boucaua, Miliolina.....	425	Haplophragmium.....	89
Nonionina.....	366	Calcarina.....	350
Quinqueloculina.....	425	baculatus.....	354
bradyana, Flutina.....	467	calcar.....	350
Quinqueloculina.....	432	var. hispida.....	356
bradyi, Clavulina.....	155	defrancii.....	353
Cribrostomoides.....	83	hispida.....	356
Cristellaria.....	243	quoyi.....	356
Cyclammina.....	88	spengleri.....	351, 356
Cymbalopora poeyi.....	308	var.....	353
Gaudryina.....	149	tetraedra.....	358
Hauerina.....	450	canaliculata, Spiroloculina.....	395
Vaginulina.....	260	canariensis, Haplophragmium.....	79
Vermenulina.....	111	Haplophragmoides.....	79
Brizalina aenariensis.....	137	Litua.....	79
broeckhiana, Rotalia.....	316	nautiloidea.....	79
brunckenthalii, Vaginulina.....	260	Nonionina.....	79
brunnensis, Uvigerina.....	272	Placopsilina.....	79
buchiana, Bulimina.....	160	Pulvinulina.....	338
presli.....	160	Rotalia.....	338
bulbosa, Spiroplecta.....	102	Uvigerina.....	275
Bulimina.....	159	cancellata, Cyclammina.....	84, 85
aculeata.....	161	candeiana, Textularia.....	109
affinis.....	165	sagittula.....	109
buchiana.....	160	Candeina.....	296
elegans, exilis.....	159	nitida.....	296
elegantissima.....	167, 168	triloba.....	296
elongata.....	165	caperata, Clavulina.....	153
inflata.....	160	Tritaxia.....	153
marginata.....	159	Tritaxilina.....	153
ovata.....	164	capreolus, Bigenerina.....	127
polystropha.....	139	Grammostomum.....	127
presli, aculeata.....	161	Valvulina.....	127
buchiana.....	160	carinata, Bolivina karreriana.....	132
elegantissima.....	167	Cornuspira.....	392
marginata.....	159	Operculina.....	392
pupoides.....	161	Textularia.....	121
marginata.....	159	carinatus, Peneroplis.....	482
spinulosa.....	161	pertusus.....	482
pyrula.....	162	carpenteri, Cycloclypeus.....	386
perversa.....	163	Hormosina.....	77
spinescens.....	164	Carpenteria.....	360
subaffinis.....	165	balaniformis, proteiformis.....	361
subteres.....	107	monticularis.....	362
Buliminella.....	167	proteiformis.....	361
contraria.....	167	raphidodendron.....	363
elegantissima.....	168	utricularis.....	360
subteres.....	167	Cassidulina.....	171
Bulimininae.....	159	crassa.....	172
bulia, Placopsilina.....	51	globulosa.....	293
Tholosina.....	51	laevigata.....	171
bulloides, Ammoglobigerina.....	285	subglobosa.....	171
Cymbalopora.....	309	Cassidulininae.....	171
Globigerina.....	96, 285	cassinoides, Cristellaria.....	244
Rosalina.....	309	cassis, Ammobaculites.....	91
Sphaeroidina.....	297	Cristellaria.....	244
Tretomphalus.....	309	Haplophragmium.....	91
triloba, Globigerina.....	286	Litua.....	91
calcar, Calcarina.....	350	Nautilus.....	244
var. hispida.....	356	castanea, Thurammina papillata.....	52
Cristellaria.....	231		

	Page.		Page.
catenata, Aschemonella.....	64	concamerata, Rotalina.....	328
Astrorhiza.....	64	Serpula.....	328
Textularia.....	112	concaua, Textularia.....	119
catenulata, Haliphysema.....	58	concauum, Plecanium.....	119
Nodosaria.....	198	concentrica, Pulvinulina.....	327
Saccamina sphaerica.....	48	conglobata, Globigerina.....	290
celata, Plauispirina.....	449	conglomerata, Globigerina.....	295
Sigmoidina.....	449	conica, Textularia.....	123
cenomana, Lituola.....	95	Valvulina.....	142
(Placopsilina).....	95	triangularis.....	142
Placopsilina.....	95	consobrina, emaciata, Nodosaria.....	195
Ceriodora globulus.....	360	contorta, Miliolina.....	432
cervicornis, Anomalina polymorpha.....	325	Quinqueloculina.....	432
chilostoma, Gaudryina.....	150	contraria, Biloculina.....	412
pupoides.....	150	Bulliminella.....	167
Textularia.....	150	Planispirina.....	412
Chilostomella.....	283	Rotalina.....	167
chilostomella, Gaudryina pupoides.....	150	convergens, Cristellaria.....	228
Chilostomella, grandis.....	283	convexa, Triloculina tricarinata.....	456
ovoidea.....	283	convexiuscula, Spiroloculina.....	409
Chilostomellidae.....	283	Cornuspira.....	387
Chrysalidina.....	144	carinata.....	392
dimorpha.....	144	crassisepta.....	392
Cibicides refulgens.....	312	foliacea.....	387
circularis, Miliolina.....	462	expansa.....	388
Spiroloculina.....	403	invovens.....	389
Triloculina.....	462	substriatula.....	389
clavata, Ammolagena.....	61	lacunosa.....	391
Lagena.....	174	Cornuspirininae.....	387
Oolina.....	174	cornuta, Virgulina.....	170
Trochammina irregularis.....	61	coronata, Anomalina.....	326
Webbina.....	61	Nevillina.....	480
Webbinella.....	61	corrugata, Patellina.....	303
Clavulina.....	145	costata, Bolivina.....	134
angularis.....	155	Cristellaria.....	239, 243
bradyi.....	155	Lenticulina.....	239
caperata.....	153	Lingulina.....	214
communis.....	154	Marginulina.....	256
cylindrica.....	155	multicosta, Cristellaria.....	240
difformis.....	156	Robulina.....	239
indiscreta.....	145	subdecorata, Cristellaria.....	241
parisiensis.....	156	sublaevis, Cristellaria.....	242
primaeva.....	158	costatus, Nautilus.....	239
rotundata.....	157	(Orthoceras).....	256
columellaris, Sagrina.....	276	Spincterules.....	239
Siphogenerina.....	276	costifera, plena, Spiroloculina.....	405
comata, Biloculina.....	478	Spiroloculina.....	405
communis, Clavulina.....	154	costulata, Nodosaria.....	187
Dentalina.....	192	stipitata.....	187
Nodosaria.....	193	Siphogenerina raphanus.....	251
(Dentalina).....	192	crassa, Cassidulina.....	172
Verneulina.....	154	Miliolina.....	420
compacta, Bolivina.....	137	Pulvinulina.....	338
robusta.....	137	Rotalina.....	338
complanata, lacinata, Orbitolites.....	487	subeuneata, Quinqueloculina.....	423
Operculina.....	373	crassatina, Miliolina.....	443
Orbitolites.....	486	Quinqueloculina.....	443
plicata, Orbitolites.....	487	crassisepta, Cornuspira.....	392
compressa, Cristellaria.....	247	Textularia.....	121
Cyclammima.....	85	craticulata, Polystomella.....	368
Hauerina.....	450	craticulatus, Nautilus.....	368
Nouria.....	102	crenata, Massilina.....	445
Polymorphina.....	266	Spiroloculina.....	445
compressiuscula, Rotalia papillosa.....	348	crepidula, Cristellaria.....	248
compressum, Haplophragmium.....	80	Nautilus.....	248
concamerata, Pulvinulina.....	328	cretacea, Globigerina.....	287
repanda.....	328	Cribrostomoides.....	83

	Page.		Page.
<i>Cribrostomoides bradyi</i>	83	<i>Cyclammina bradyi</i>	88
<i>crispa</i> , <i>Polystomella</i>	368	<i>cancellata</i>	84
<i>crispus</i> , <i>Nautilus</i>	368	<i>compressa</i>	85
<i>Cristellaria</i>	220	<i>pauciloculata</i>	86
<i>aculeata</i>	235	<i>pusilla</i>	87
<i>articulata</i>	222	<i>Cyeloclypeus</i>	386
<i>bradyi</i>	243	<i>carpenteri</i>	386
<i>calcar</i>	231	<i>guembeliaanus</i>	386
<i>calcarata</i>	232	<i>cyllindracea</i> , <i>Spirolinites</i>	483
<i>cassinoides</i>	244	<i>cyllindraceus</i> , <i>Peneroplis</i>	483
<i>cassis</i>	244	<i>cyllindrica</i> , <i>Clavulina</i>	155
<i>compressa</i>	247	<i>Marsipella</i>	41
<i>convergens</i>	228	<i>Polytrema</i>	365
<i>costata</i>	239, 243	<i>Reophax</i>	70
<i>multicosta</i>	240	<i>cyllindricum</i> , <i>Sporadotrema</i>	365
<i>subdecorata</i>	241	<i>cyllindricus</i> , <i>Ammobaculites</i>	92
<i>sublaevis</i>	242	<i>Reophax</i>	70
<i>crepidula</i>	248	<i>Cymbalopora</i>	308
<i>cultrata</i>	220, 238	<i>bulloides</i>	309
<i>decorata</i>	221	<i>poeyi</i>	308
<i>denticulifera</i>	238	<i>bradyi</i>	308
<i>dorso-costata</i>	247	<i>var</i>	308
<i>echinata</i>	233	<i>decorata</i> , <i>Cristellaria cultrata</i>	221
<i>expansa</i>	236	<i>defrancii</i> , <i>Calcarina</i>	353
<i>planulata</i>	237	<i>dehiscens</i> , <i>Sphaeroidina</i>	297
<i>gemmata</i>	239	<i>Dendrophrya</i>	56
<i>gibba</i>	228	<i>attenuata</i>	57
<i>helicina</i>	250	<i>ramosa</i>	56
<i>helicinoides</i>	251	<i>robusta</i>	57
<i>italica</i>	252	<i>Dentalina communis</i>	192
<i>acuto-carinata</i>	253	<i>Nodosaria</i>	192
<i>latifrons</i>	254	<i>enaciata</i>	195
<i>mamilligera</i>	235	<i>legumeni</i> , <i>linearis</i>	261
<i>orbicularis</i>	224	<i>mucronata</i>	195
<i>papillata</i>	226	<i>obliqua</i> , <i>Nodosaria</i>	195, 210
<i>subcarinata</i>	225	<i>roemeri</i>	193
<i>subumbonata</i>	226	<i>soluta</i>	192
<i>papillosa</i>	234	<i>subnodosa</i>	139
<i>paucicostata</i>	248	<i>dentaliniformis</i> , <i>Reophax</i>	68
<i>perprocera</i>	250	<i>denticulata</i> , <i>Biloculina</i>	476
<i>reniformis</i>	221	<i>ringens</i>	476
<i>Cristellaria</i> (<i>Robulina</i>) <i>echinata</i>	233	<i>Spirillina limbata</i>	302
<i>rotulata</i>	223	<i>striolata</i> , <i>Biloculina</i>	477
<i>umbonata</i>	223	<i>denticulifera</i> , <i>Cristellaria</i>	238
(<i>Saracenaria</i>) <i>italica</i>	252	<i>depressa</i> , <i>Biloculina</i>	489
<i>schloenbachi</i>	249	<i>Spiroloculina</i>	394
<i>submamilligera</i>	235	<i>tubercalata</i> , <i>Heterostegina</i>	364
<i>tenuis</i>	250	<i>diaphana</i> , <i>Iridia</i>	47
<i>tricarinnella</i>	230	<i>diffugiformis</i> , <i>testacea</i> , <i>Reophax</i>	50
<i>spinipes</i>	230	<i>Proteonina</i>	49
<i>tumido-costata</i>	245	<i>Reophax</i>	49
<i>labyrinthica</i>	245	<i>digitata</i> , <i>Globigerina</i>	288
<i>variabilis</i>	229	<i>dilatata</i> , <i>Bolivina</i>	128
<i>vortex</i>	227	<i>dimorpha</i> , <i>Chrysaladina</i>	144
<i>culter</i> , <i>Planorbulina</i>	320	<i>Sagrina</i>	279
<i>Truncatulina</i>	320	<i>Siphogenerina</i>	279
<i>cultrata</i> , <i>Cristellaria</i>	220, 228	<i>Uvigerina</i> (<i>Sagrina</i>).....	279
<i>decorata</i> , <i>Cristellaria</i>	221	<i>Dimorphina striata</i>	280
<i>Robulina</i>	220	<i>discoidalis</i> , <i>Assilina</i>	379
<i>cultratus</i> , <i>Robulus</i>	220	<i>involuta</i> , <i>Operculina</i>	380
<i>eumingii</i> , <i>Amphistegina</i>	383	<i>Operculina</i>	379
<i>Nummulites</i>	383	<i>discoideus</i> , <i>Ammodiscus incertus</i>	63
<i>emneata</i> , <i>incisa</i> , <i>Triloculina</i>	462	<i>Discorbina allomorphinoides</i>	306
<i>Miliolina</i>	461	<i>bertheloti</i>	305
<i>Triloculina</i>	461	<i>orbicularis</i>	305
<i>eurta</i> , <i>Quinqueloculina</i>	426	<i>rugosa</i>	306
<i>cuvieriana</i> , <i>Miliolina</i>	426	<i>saulei</i>	307
<i>Cyclammina</i>	84	<i>Discorbis</i>	308

	Page.		Page.
<i>Discorbis allomorphinoides</i>	206	<i>Fissurina laevigata</i>	174
<i>ammonoides</i>	323	<i>staphyllearia</i>	181
<i>bertheloti</i>	305	<i>fistula</i> , <i>Textularia agglutinans</i>	107
<i>orbicularis</i>	305	<i>fistulosa</i> , <i>Spiroplecta sagittula</i>	107
<i>rugosa</i>	306	<i>Textularia sagittula</i>	104
<i>sp.</i>	307	<i>flavidum</i> , <i>Bathysiphon</i>	44
<i>subobtusata</i>	304	<i>flavidus</i> , <i>Bathysiphon</i>	44
<i>vilardeboana</i>	304	<i>giganteus</i> , <i>Bathysiphon</i>	44
<i>discreta</i> , <i>Rhabdammina</i>	39	<i>flintii</i> , <i>Gaudryina</i>	146
<i>disparilis</i> , <i>Miliolina</i>	424	<i>Textularia</i>	113
<i>Quinqueloculina</i>	424	<i>Flintina</i>	465
<i>distans</i> , <i>Reophax</i>	66	<i>bartschi</i>	465
<i>divaricans</i> , <i>Sagenina</i>	60	<i>bradyana</i>	467
<i>dorso-costata</i> , <i>Cristellaria</i>	247	<i>triquetra</i>	466
<i>dubia</i> , <i>Haplostiche</i>	78	<i>foliacea</i> , <i>Cornuspira</i>	387
<i>Lituola</i>	78	<i>expansa</i> , <i>Cornuspira</i>	388
<i>Nodosaria</i>	78	<i>Textularia</i>	117
<i>duplex</i> , <i>Orbitolites</i>	185	<i>foliaceus</i> , <i>Orbis</i>	387
<i>durrandii</i> , <i>Massilina</i>	146	<i>formosa</i> , <i>Lagena</i>	183
<i>Miliolina</i>	446, 479	<i>fragillisma</i> , <i>Hauerina</i>	451
<i>echinata</i> , <i>Cristellaria (Robulina)</i>	233	<i>Spiroloculina</i>	451
<i>Robulina</i>	233	<i>friabilis</i> , <i>Hyperammina</i>	54
<i>Ehrenbergina</i>	172	<i>frondescens</i> , <i>Sagenella</i>	60
<i>serrata</i>	172	<i>Sagenina</i>	60
<i>elegans</i> , <i>exilis</i> , <i>Bulimina</i>	159	<i>Fronicularia</i>	216
<i>Miliolina bicornis</i>	425	<i>annularis</i>	217
<i>Operculina</i>	381	<i>longistriata</i>	218
<i>Pulvinulina</i>	342	<i>inaequalis</i>	216
<i>Quinqueloculina bicornis</i>	425	<i>philippinensis</i>	216
<i>Rotalia (Turbinulina)</i>	342	<i>plicata</i>	219
<i>Spiroloculina</i>	406	<i>fusca</i> , <i>Miliolina</i>	442
<i>Vaginulina</i>	258	<i>Psammospaera</i>	46
<i>legumen</i>	258	<i>Quinqueloculina</i>	442
<i>elegantissima</i> , <i>Bulimina</i>	167, 168	<i>Rotalina</i>	143
<i>presli</i>	167	<i>Valvulina</i>	143
<i>Buliminella</i>	167, 168	<i>fusiformis</i> , <i>Proteonina</i>	49
<i>Polymorphina</i>	267	<i>Reophax</i>	49
<i>elongata</i> , <i>Biloculina</i>	473	<i>gaimairdi</i> , <i>Operculina</i>	375
<i>Formosina</i>	76	<i>Gaudryina</i>	146
<i>Hyperammina</i>	53, 54	<i>apicularis</i>	151
<i>Lagena</i>	175	<i>attenuata</i>	152
<i>Miliola</i>	175	<i>baccata</i>	150
<i>emaciata</i> , <i>Dentalina</i>	195	<i>bradyi</i>	149
<i>Nodosaria consobrina</i>	195	<i>chilostoma</i>	150
<i>emaciatum</i> , <i>Haplophragmium</i>	80	<i>filiformis</i>	151
<i>Haplophragmoides</i>	80	<i>flintii</i>	146
<i>Entosolenia marginata</i>	182	<i>paupercula</i>	148
<i>lagenoides</i>	183	<i>pseudofiliformis</i>	151
<i>excavata</i> , <i>Spiroloculina</i>	120, 396	<i>pupoides</i>	149
<i>Textularia</i>	120	<i>chilostoma</i>	150
<i>exigua</i> , <i>Pulvinulina</i>	340	<i>chilostomella</i>	150
<i>exilis</i> , <i>Bulimina elegans</i>	388	<i>quadrangularis</i>	147
<i>expansa</i> , <i>Cornuspira foliacea</i>	388	<i>robusta</i>	152
<i>Cristellaria</i>	236	<i>rugosa</i>	146
<i>planulata</i> , <i>Cristellaria</i>	237	<i>scabra</i>	146
<i>exsertus</i> , <i>Ammolagena</i>	64	<i>siphonella</i>	151
<i>Faujasina</i>	347	<i>subrotundata</i>	146
<i>ferussacii</i> , <i>miliolina</i>	429	<i>triangularis</i>	148
<i>Quinqueloculina</i>	429	<i>gemmata</i> , <i>Cristellaria</i>	239
<i>filiformis</i> , <i>Bathysiphon</i>	41	<i>gibba</i> , <i>Cristellaria</i>	228
<i>Gaudryina</i>	151	<i>Polymorphina</i>	267
<i>Nodosaria</i>	194	(<i>Globulina</i>).....	267
<i>fimbriata</i> , <i>Pulvinulina menardii</i>	335	<i>gigantea</i> , <i>Marsipella</i>	40
<i>findens</i> , <i>Lituola</i>	71	<i>giganteus</i> , <i>Bathysiphon flavidus</i>	44
<i>Reophax</i>	71	<i>Girvanella vagans</i>	55
<i>Fissurina bicarinata</i>	184	<i>glabra</i> , <i>Marginulina</i>	254

	Page.		Page.
Glandulina annulata.....	185	Haliphysema.....	58
laevigata.....	185	catenulata.....	58
Nodosaria.....	185	hantkeniana, Bolivina.....	132
striatula, Nodosaria.....	186	Haplophragmium agglutinans.....	89, 92
rotundata.....	185	calcareum.....	89, 90, 92
Nodosaria.....	185	canariensis.....	79
Globigerina.....	285	cassis.....	91
aequilateralis.....	293	compressum.....	80
involuta.....	294	emaciatum.....	80
bulloides.....	96, 285	globigeriniforme.....	96
triloba.....	286	latidorsatum.....	81, 83
conglobata.....	290	pseudospirale.....	94
conglomerata.....	295	ringens.....	81
cretacea.....	287	scitulum.....	80
digitata.....	288	sphaeroidiniformis.....	98
helicina.....	291	subglobosum.....	81
inflata.....	289	tenuimargo.....	90
rubra.....	289	turbinatum.....	96
sacculifera.....	291	Haplophragmoides.....	79
triloba.....	286	canariensis.....	79
Globigerinidae.....	285	emaciatum.....	80
globigeriniformis, Haplophragmium.....	96	grandiformis.....	82
Lituola (Haplophragmium).....	96	ringens.....	81
nautiloidea.....	96	scitulum.....	80
Trochammina.....	96	sphaeriloculum.....	83
globosa, Lagena.....	173	subglobosum.....	81
globosum, Vermiculium.....	173	Haplostiche.....	78
globulifera, Hormosina.....	74	dubia.....	78
Ramulina.....	282	soldanii.....	78
(Globulina) gibba, Polymorphina.....	267	harrisii, Nouria.....	101
globulosa, Cassidulina.....	293	hauerii, Pulvinulina.....	331, 332
globulus, Biloculina.....	474	Textularia.....	105
Ceriopora.....	360	Hauerina.....	450
Gypsina.....	360	bradyi.....	450
goësi, Textularia.....	113	compressa.....	450
gracilis, Amphorina.....	175	fragilissima.....	451
Lagena.....	175, 181	inconstans.....	393
gracillima, Amphorina.....	175	ornatissima.....	452
Lagena.....	175	helicina, Cristellaria.....	250
gramen, Textularia.....	105	Globigerina.....	291
Grammostomum capreolus.....	127	helicinoides, Cristellaria.....	251
grandiformis, Haplophragmoides.....	82	herdmani, Ramulina.....	363
grandis, Ammosphaeroidina.....	98	Heterostegina.....	384
Chilostomella.....	283	depressa, tuberculata.....	384
Lingulina.....	215	suborbicularis.....	385
granulosa, Astrorhiza.....	35	tuberculata.....	384
Marsipella.....	35	hexagona, scalariformis, Lagena.....	177
Operculina.....	381	hirsuta, aculeata, Nodosaria.....	211
grata, angulata, Spiroloculina.....	408	Nodosaria.....	213
Spiroloculina.....	407	hispidula, Calcarina.....	356
grateloupi, incisa, Spiroloculina.....	397	Lagena.....	176
Spiroloculina.....	396	Nodosaria.....	213
grosserugosa, Anomalina.....	323	sublineata, Nodosaria.....	207
Truncatulina.....	323	tubulata, Lagena.....	177
guembelianus, Cyclolepeus.....	386	hispidula, Nodosaria lepidula.....	203
guttifera, Reophax.....	69	Homotrema.....	364
(Guttulina) problema, Polymorphina.....	264	rubrum.....	364
sororia, Polymorphina.....	265	Hormosina.....	74
Gypsina.....	360	carpenteri.....	77
globulus.....	360	elongata.....	76
(Gyroidina) orbicularis, Rotalia.....	346	globulifera.....	74
soldanii, Rotalia.....	349	monile.....	76
haidingeri, Lagena.....	178	normani.....	75
striata.....	178	ovaliformis.....	76
Oolina.....	178	ovicula.....	75
Planorbulina.....	315	hormosinoides, Reophax nodulosus.....	69
Rotalina.....	315	horrida, Reophax.....	73
Truncatulina.....	315		

	Page.		Page.
horridus, Reophax.....	73	lacteum, Vermiculum.....	265
humilis, Truncatulina.....	321	lacunosa, Cornuspira.....	391
Hyperammina.....	53	laeve, Vermiculum.....	173
elongata.....	53, 54	laevicostata, Nodosaria.....	209
friabilis.....	54	leivicostata, Cassidulina.....	209
ramosa.....	54	laevigata, Cassidulina.....	171
vagans.....	55	Fissurina.....	174
Hyperammininae.....	53	Glandulina.....	185
immensa, Textularia.....	118	Lagena.....	174
impressa, Spiroloeculina.....	395	Nodosaria (Glandulina).....	185
inaequalis, Frondicularia.....	216	striatula, Nodosaria (Glandulina).....	186
inaequilateralis, involuta, Globigerina.....	294	laevis, Lagena.....	173
Spiroloeculina.....	410	Lagena.....	173
incerta, Operculina.....	62	acuticosta.....	180
incertus, Ammodiscus.....	62	alveolata, substriata.....	182
discoidens, Ammodiscus.....	63	apiculata.....	176
incisa, Spiroloeculina grateloupi.....	397	auriculata, substriata.....	182
Triloeculina cuneata.....	462	bicaeinata.....	184
inconstans, Hauerina.....	393	clavata.....	174
Ophthalmidium.....	393	elongata.....	175
incrassata, Miliolina.....	443	formosa.....	183
indica, Polymorphina problema.....	265	globosa.....	173
Pulvinulina.....	332	gracilis.....	181
indiscreta, Clavulina.....	145	gracillima.....	175
Tritaxia.....	145	haidingeri.....	178
indivisa, Rhizammina.....	45	hexagona, scalariformis.....	177
inflata, Bolimina.....	160	hispidula.....	176
Globigerina.....	289	tubulata.....	177
Textularia solita.....	111	laevigata.....	174
inflexa, Nodosaria.....	191	laevis.....	173
insecta, Nodosaria.....	189	lagenoides.....	183
insignis, Miliolina.....	458	lamellata.....	184
Triloeculina.....	458	marginata.....	182
insolita, Nodosaria.....	191	(Serpula).....	182
interrupta, Uvigerina.....	275	semistriata.....	179
involuta, Globigerina inaequilateralis.....	294	staphyllearia.....	171
Operculina discoidalis.....	380	striata.....	177, 178
invovens, Cornuspira.....	389	haidingeri.....	178
Operculina.....	389	semistriata.....	179
substriatula, Cornuspira.....	389	strumosa.....	178
Iridia.....	47	sulcata.....	179
diaphana.....	47	apiculata.....	180
irregularis, Biloculina.....	479	Serpula.....	179
clavata, Trochammina.....	61	Lagenidae.....	173
Rhabdammina.....	38	Lageninae.....	173
Trochammina.....	61	lagenoides, Entosolenia marginata.....	183
italica, acuto-carinata, Cristellaria.....	253	Lagena.....	183
Cristellara (Saracenaria).....	252	lamarekiana, Quinqueloculina.....	418
Cristellaria.....	252	lamellata, Lagena.....	184
Saracenaria.....	252	larvata, Planorbulina.....	310
Jaculella.....	59	vulgaris.....	310
acuta.....	59	lateralis, Pulvinulina.....	336
obtusa.....	59	Rosalina.....	336
japonica, Nodosaria.....	196	latidorsata, Nonionina.....	81
jugosa, Textularia.....	119	latidorsatum, Haplophragmium.....	81, 83
karreriana, Bolivina.....	131	latifrons, Cristellaria.....	254
carinata, Bolivina.....	132	legumen, elegans, Vaginulina.....	258
kerimbatica, Miliolina.....	437	linearis, Dentalina.....	261
philippinensis, Quinqueloculina.....	437	Nautilus.....	257
Quinqueloculina.....	437	Technitella.....	50
reticulo-striata, Quinqueloculina.....	437	Vagiulina.....	257
jabyrinthica, Cristellaria tumido-costata.....	246	Lenticulina costata.....	239
laciniata, Orbitolites complanata.....	487	Lenticulites.....	373
laciniatus, Orbitolites.....	487	rotulata.....	223
lactea, oblonga, Polymorphina.....	266	lepidula, hispidula, Nodosaria.....	203
Polymorphina.....	265	Nodosaria.....	203
Serpula.....	265		

	Page.		Page.
lessonii, Amphistegina.....	370	Marsipella.....	40
madagascariensis, Amphistegina.....	372	cylindrica.....	41
radiata, Amphistegina.....	372	gigantea.....	40
limbata, Bolivina.....	134	granulosa.....	35
denticulata, Spirillina.....	302	Massilina.....	444
Miliolina.....	422	arenaria.....	446
Quinqueloculina.....	422	asperula.....	447
Spirillina.....	301	crenata.....	445
Spiroloculina.....	394	durrandii.....	446
tuberculo-limbata, Spirillina.....	302	secans, reticulata.....	444
linearis, Dentalina legumen.....	261	tenuistriata.....	445
Nautilus.....	261	menardii, fimbriata, Pulvinulina.....	335
Rhabdammina.....	40	Pulvinulina.....	333
Vaginulina.....	261	Rotalia.....	333
Lingulina.....	214	tumida, Pulvinulina.....	337
costata.....	214	michelimana, Pulvinulina.....	339
grandis.....	215	Rotalina.....	339
linuacana, Miliolina.....	463	Miliola elongata.....	175
Triloculina.....	463	Milioliidae.....	387
lituiformis, Lituotuba.....	88	Miliolina alveoliniformis.....	443
Trochammina.....	88	bertheliniana.....	457
Lituola.....	71	bicornis, elegans.....	425
canariensis.....	79	boueana.....	425
cassis.....	91	circularis.....	462
cenomana.....	95	contorta.....	432
dubia.....	78	crassa.....	423
findens.....	71	crassatina.....	443
(Haplophragmium) globigeriniformis.....	96	euneata.....	461
nautiloidea, canariensis.....	79	euvieriana.....	426
globigeriniformis.....	96	disparilis.....	424
(Placopsilina) cenomana.....	95	durrandii.....	479
soldanii.....	78	ferussacii.....	429
subglobosa.....	81	fusea.....	442
Lituoliidae.....	64	incrassata.....	443
Lituotuba.....	88	insignis.....	458
lituiformis.....	88	kerimbatia.....	437
(Lituus) arietinus, Nautilus.....	482	limbata.....	422
lobatula, Serpula.....	313	linuacana.....	463
Truncatulina.....	313	oblonga.....	459
lobatulus, Nautilus.....	313	parkeri.....	437, 440
longicostata, Nodosaria pyrula.....	188	procera.....	421
longistriata, Frondicularia annularis.....	218	reticulata.....	434
luernula, Biloculina.....	475	rotunda.....	460
luzonensis, Nodosaria pauciloculata.....	206	rupertiana.....	464
madagascariensis, Amphistegina lessonii.....	372	scleroica.....	441
mamilligera, Cristellaria.....	235	seminulum.....	416
margaritifera, Truncatulina.....	319	suborbicularis.....	461
Vaginulina.....	259	terquemiana.....	458
margaritifera, Nautilus (Orthoecras).....	259	tricarinata.....	454, 457, 458
marginalis, Orbitolites.....	484	bertheliniana.....	457
Orbitolites.....	484	plicata.....	456
marginata, Bulimina.....	159	terquemiana.....	458
presli.....	159	trigonula.....	452
pupoides.....	159	triquetra.....	466
Entosolenia.....	182	undosa.....	432
Lagena.....	182	valvularis.....	460
lagenoides, Entosolenia.....	183	venusta.....	420
Serpula (Lagena).....	182	vulgaris.....	417
marginatum, Vermiculum.....	182	Miliolites planulata, planissima.....	401
Marginopora vertebralis, plicata.....	487	trigonula.....	452
Margulinina.....	254, 259	Millepora miniacea.....	363
costata.....	256	rubra.....	364
glabra.....	254	millettii, Biloculina.....	479
philippinensis.....	257	Nodosaria.....	202
raphanus.....	256	Spiroloculina.....	400
striatula.....	255	miniacea, Millepora.....	363
tenuis.....	250	miniaceum, Polytrema.....	363

	Page.		Page.
monile, Hormosina	76	Nodosaria (Glandulina) laevigata	185
monticularis, Carpenteria	362	striatula	186
mucronata, Dentalina	195	rotundata	185
Nodosaria	195	hirsuta	213
multicosta, Cristellaria costata	240	aculeata	214
multistriata, Triloculina trigonula	453	hispidula	213
murrhyna, Biloculina	470	sublineata	207
nautiloidea, canariensis, Lituola	79	inflexa	191
globigeriniformis, Lituola	96	insecta	189
Nautilus	350, 373	insolita	191
acicularis	483	japonica	196
aduncus	483	laevicostata	209
ammonoides	382	lepidula	203
auricula, var	329	hispidula	203
becarii	345	milletii	202
calcar	231, 233	mucronata	195
cassis	244	nodosa	139
costatus	239	obliqua	210
craticulatus	368	pauciloculata	205
crepidula	248	luzonensis	205
crispus	368	perversa	208
legumen	257	protumida	262
linearis	261	pyrula	187
(Lituus) arietius	482	longi-costata	188
lobatulus	313	semirugosa	187
obliquus	210	radicula	190
(Orthoceras) costatus	256	annulata	189
margaritiferus	259	raphanus	210
pennatula	127	roemeri	193
scalaris	199	scalaris	199
vertebralis	211	separans	200
papillosus	234	semirugosa	187
pertusus	481	simplex	186
plabatus	482	soluta	192
pompilioides	366	spirostriolata	212
radiatus	372	stipitata, costulata	187
radicula	190	subcanaliculata	204
raphanus	210	sublineata	207
repandus	326	subperversa	208
spengleri	351	subpolygona	198
umbilicatus	365	subscalaris	201
venosus	383	pauci-costata	201
vortex	227	substriatula	204
Nevillina	480	subtenuata	213
coronata	480	Textularia agglutinans	125
nitida, Candeina	296	vertebralis	211
Spiroloculina	408	Nodosariinae	185
triloba, Candeina	296	nodulosus, Hormosinoidea, Reophax	69
nobilis, Bolivina	130	Reophax	68
Nodobacularia	415	Nonionina	365
tibia	415	boucana	366
nodosa, Nodosaria	139	canariensis	79
Uvigerina	275	latidorsata	81
Nodosaria	185	pompilioides	366
annulata	189	quinquecloba	366
antennula	197	umbilicatulata	365
Bigenerina	125	normani, Hormosina	75
catenulata	198	Nouria	101
communis	192	compressa	102
conobryna, emaciata	195	harrisii	101
costulata	187	Nummulites cumingii	383
stipitata	187	venosa	383
(Dentalina) communis	192	Nummulitidae	365
obliqua	210	obliqua, Nodosaria	210
dubia	78	(Dentalina)	195, 210
filiformis	194	obliquiloculata, Pullenia	299
(Glandulina) laevigata	185	obliquus, Nautilus	210
		oblonga, Miliolina	430

	Page.		Page.
oblonga, Polymorphina.....	268	ovata, Bulimina.....	164
laetea.....	266	ovicula, Hormosina.....	75
Pulvinulina.....	330	ovoidea, Chilostomella.....	283
Rotalina.....	330	papillata, castanea, Thurammina.....	52
seabra, Pulvinulina.....	330	Cristellaria orbicularis.....	226
Triloculina.....	459	papillosa, compressiuscula, Rotalia.....	347
oblongum, Vermieulum.....	459	Cristellaria.....	234
obtusa, Jaculella.....	59	Rotalia.....	347
Oolina apiculata.....	176	papillosus, Nautilus.....	234
clavata.....	174	papyracea, Thurammina.....	52
haidingeri.....	178	papyraceus, Bathysiphon.....	45
strata.....	177	parkeri, Miliolina.....	437, 440
Operculina.....	373	Quinqueloculina.....	440
ammonoides.....	382	parisiensis, Clavulina.....	156
bartschi.....	376	Valvulina.....	156
ornata.....	378	partschiana, Pulvinulina.....	344
plana.....	377	parva, Psammospaera.....	47
carinata.....	392	Patellina.....	303
complanata.....	373	corrugata.....	303
discoidalis.....	379	patens, Vaginulina.....	263
involuta.....	380	paucicostata, Cristellaria.....	248
elegans.....	384	Nodosaria subscalaris.....	201
gaimairdi.....	375	pauciloculata, Cyclammina.....	86
granulosa.....	381	Iuzonensis, Nodosaria.....	206
incerta.....	62	Nodosaria.....	205
involvens.....	389	Virgulina.....	168
philippinensis.....	378	pauperata, Pulvinulina.....	340
venosa.....	383	pauperula, Gaudryina.....	148
Ophthalmidium.....	393	Peneroplis.....	481
inconstans.....	393	arictinus.....	482
orbicularis, Cristellaria.....	224	carinatus.....	482
Discorbina.....	305	cylindraceus.....	483
Discorbis.....	305	pertusus.....	481
papillata, Cristellaria.....	226	arictinus.....	482
Robulina.....	224	carinatus.....	482
Rosalina.....	305	planatus.....	481
Rotalia.....	346	planatus.....	481
(Gyroïdina).....	346	pennatula, Bigenerina.....	127
subcarinata, Cristellaria.....	225	Nautilus (Orthoceras).....	127
subumbonata, Cristellaria.....	226	perprocera, Cristellaria.....	250
Orbiculina.....	483	pertusus, arictinus, Peneroplis.....	482
adunca.....	483	carinatus, Peneroplis.....	482
Orbis foliaceus.....	387	Nautilus.....	481
orbis, Spiroloculina.....	403	planatus, Peneroplis.....	481
Orbitolites.....	481	perversa, Bulimina pyrula.....	163
complanata.....	486	Nodosaria.....	208
laciniata.....	487	philippinensis, Frondicularia.....	216
plicata.....	487	Marginulina.....	257
duplex.....	485	Operculina.....	378
laciniatus.....	487	Pulvinulina.....	331
marginalis.....	484	Quinqueloculina kerimbatica.....	438
Orbulina.....	295	pigmaea, Uvigerina.....	269
universa.....	295	pilulifer, Reophax.....	66
Orbulites marginalis.....	484	pilulifera, Reophax.....	66
ornata, Operculina bartschi.....	378	pisum, Biloculina.....	475
Truncatulina ungeriana.....	317	Placopsilina.....	95
ornatissima, Hauerina.....	452	bulla.....	51
Quinqueloculina.....	442	canariensis.....	79
Orthocera raphanus.....	210	enomana.....	95
(Orthoceras) eostatus, Nautilus.....	256	(Placopsilina) enomana, Litola.....	95
margariferus, Nautilus.....	259	plana, Operculina bartschi.....	377
pennatula, Nautilus.....	127	planatus, Nautilus.....	481
scalaris, Nautilus.....	199	Peneroplis.....	481
vertebralis, Nautilus.....	211	pertusus.....	481
Oryzaria boscii.....	487	Planispirina.....	412
ovaliformis, Hormosina.....	76	celata.....	449
ovalis, Sphaerammina.....	100	ef. striata.....	413

	Page.		Page.
Planispirina contraria.....	412	presli, aculeata, Bulimina.....	161
sigmoidea.....	448	buchiana, Bulimina.....	160
sphaera.....	413	elegantissima, Bulimina.....	167
planissima, Miliolites planulata.....	401	marginata, Bulimina.....	159
Spiroloeculina.....	401	primaeva, Clavulina.....	158
Planorbulina.....	310	problema, indica, Polymorphina.....	265
acervalis.....	311	Polymorphina.....	264
ammonoides.....	323	(Guttulina).....	264
culter.....	320	procera, Miliolina.....	421
haidingerii.....	315	Pulvinulina.....	342
larvata.....	310	Quinqueloculina.....	421
ungeriana.....	317	propinqua, Verneuilina.....	140, 142
vulgaris, larvata.....	310	proteiformis, Carpenteria.....	361
planulata, Cristellaria expansa.....	237	balaniformis.....	361
planissima, Miliolites.....	401	Proteonina.....	49
Planulina ariminensis.....	322	difflugiformis.....	49
Plecanium concavum.....	119	fusiformis.....	49
rugosa.....	114	pseudospirale.....	94
plena, Spiroloeculina costifera.....	405	testicea.....	50
Pleurostomella.....	138	proteus, Trochammina.....	79, 95
alternans.....	138	Trochamminoides.....	79, 95
subnodosa.....	139	protumida, Nodosaria.....	262
plicata, Frondicularia.....	219	Vaginulina.....	262
Marginopora vertebralis.....	487	Psammospaera.....	46
Orbitolites complanata.....	487	fusca.....	46
Triloculina tricarinata.....	456	parva.....	47
pocyi, bradyi, Cymbalopora.....	308	pseudobacillaris, Reophax.....	71
Cymbalopora.....	308	pseudocarinata, Textularia.....	121
Rosalina.....	308	pseudofiliformis, Gaudryina.....	151
polygona, Quinqueloculina.....	431	pseudospirale, Ammobaculites.....	94
polymorpha, Anomalina.....	324	Haplophragmium.....	94
cervicornis, Anomalina.....	325	Proteonina.....	94
siphonifera, Anomalina.....	325	Pullenia.....	299
Polymorphina.....	263	obliquiloculata.....	299
compressa.....	268	quinqueloba.....	299
elegantissima.....	267	sphaeroides.....	299
gibba.....	267	Pulvinulina.....	326
(Globulina) gibba.....	267	auricula.....	329
(Guttulina) problema.....	267	berthlotiana, subornata.....	333
sororia.....	265	canariensis.....	338
lactea.....	265	concamerata.....	328
oblonga.....	266	concentrica.....	327
oblonga.....	26 ⁸	crassa.....	338
problema.....	264	elegans.....	342
indica.....	265	exigua.....	340
regina.....	263	hauerii.....	331, 332
sororia.....	265	indica.....	332
Polymorphininae.....	263	lateralis.....	336
Polymorphum sagittula.....	103	menardii.....	333
Polystomella.....	367	fimbriata.....	335
eraticulata.....	368	tumida.....	337
crispa.....	368	micheliniana.....	331
subnodosa.....	367	oblonga.....	330
vortex.....	227	scabra.....	330
polystropha, Bulimina.....	139	partschiana.....	344
Verneuilina.....	139	pauperata.....	340
Polytrema.....	363	philippinensis.....	331
cylindrica.....	365	procera.....	342
miniaceum.....	363	punctulata.....	328
rubra.....	664	repanda.....	326
utricularis.....	360	concamerata.....	328
pompilioides, Nautilus.....	366	punctulata.....	328
Nonionina.....	366	scabra.....	330
porrecta, Textularia.....	109	schreibersii.....	341
agglutinans.....	109	truncatulinoides.....	339
praecincta, Rotalia.....	318	tumida.....	337
Truncatulina.....	318	umbonata.....	339

	Page.		Page.
punctata, Bolivina.....	136	radicula, Nautilus.....	190
Pulvinulina.....	328	Nodosaria.....	190
repanda.....	328	ramosa, Dendrophrya.....	56
Rotalia.....	328	robusta, Dendrophrya.....	57
pupoides, Bulimina.....	161	Hyperammima.....	54
chilostoma, Gaudryina.....	150	Saccorhiza.....	54
chilostomella.....	150	Ramulina.....	282
Gaudryina.....	149	globulifera.....	282
marginata, Bulimina.....	159	herdmani.....	363
spinulosa, Bulimina.....	161	Ramulininae.....	282
pusilla, Bolivina.....	138	raphanus, costulata, Siphogenerina.....	281
Cyclammima.....	87	Marginulina.....	256
pygmaea, Bolivina.....	136	Nautilus.....	210
Uvigerina.....	269	Nodosaria.....	210
Verneuilina.....	141	Orthocera.....	210
pyrula, Bulimina.....	162	Siphogenerina.....	280
longi-costata, Nodosaria.....	188	Uvigerina (Sagrina).....	280
Nodosaria.....	187	raphidodendron, Carpenteria.....	363
perversa, Bulimina.....	163	recta, Quinqueloculina.....	423
semirugosa, Nodosaria.....	187	refulgens, Cibicides.....	312
spinescens, Bulmina.....	164	Truncatulina.....	312
quadrangularis, Gaudryina.....	147	regina, Polymorphina.....	263
quadrilatera, Bolivina.....	125	reniformis, Cristellaria.....	221
Textularia.....	125	reophaciformis, Ammobaculites.....	92
quinqueloba, Nonionina.....	299	Reophacinae.....	65
Pullemia.....	299	Reophax.....	65
Quinqueloculina.....	415	adunca.....	70
agglutinans.....	441, 464	aduncus.....	70
albatrossi.....	435	agglutinatus.....	73
alveoliniformis.....	443	baeillaris.....	67
apicula.....	422	bilocularis.....	74
bicarinata.....	428	cylindrica.....	70
bicornis, elegans.....	425	cylindricus.....	70
boucana.....	425	dentalimiformis.....	68
bradyana.....	432	ditflugiformis.....	49
contorta.....	432	testacea.....	50
crassa, subcuneata.....	423	distanis.....	66
crassatina.....	443	findens.....	71
curta.....	426	fusiformis.....	49
disparilis.....	424	guttifer.....	69
curta.....	426	guttifera.....	69
ferussaci.....	429	hormosinoides.....	69
lusca.....	442	horrida.....	73
kerimbatica.....	437	horridus.....	73
philippinensis.....	438	nodulosus.....	68
reticulo-striata.....	437	pilulifer.....	66
lamarckiana.....	418	pilulifera.....	66
limbata.....	422	pseudobaeillaris.....	71
ornatissima.....	452	scorpiurus.....	65
parkeri.....	440	spiculotestus.....	72
polygona.....	431	repanda, concamerata, Pulvinulina.....	328
procera.....	421	punctulata, Pulvinulina.....	328
recta.....	423	Pulvinulina.....	326
reticulata.....	434	Rotalia.....	326
rugosa.....	429	repandus, Nautilus.....	326
sclerotica.....	441	reticulata, Massilina secans.....	444
sculpturata.....	433	Miliolina.....	434
seminulum.....	416	Quinqueloculina.....	434
suborbicularis.....	461	Rotalina.....	322
venusta.....	420	Siphonina.....	322
vulgaris.....	417	Triloculina.....	434
quoyi, Alcarina.....	356	Truncatulina.....	322
radiata, Amphistegina.....	372	reticulosa, Vertebralina striata.....	414
lessonii.....	372	reticulo-striata, Quinqueloculina kerimbatica.....	437
Rhabdammina abyssorum.....	37	Rhabdammina.....	36
radiatus, Nautilus.....	372	abyssorum.....	36
radicula, annulata, Nodosaria.....	189	radiata.....	37

	Page.		Page.
Rhabdammina discreta	39	Rotalina fusca	143
irregularis	38	haidingerii	315
linearis	40	micheliniiana	339
Rhabdogonium tricarinatum	219	oblonga	330
Rhabdopleura	39	reticulata	332
abyssorum	39	schreibersii	341
rhaphtodendron, Carpenteria	363	umbonata	339
Rhizammina	45	ungariana	317
algaeformis	46	Rotalinae	303
indivisa	45	rotulata, Cristellaria	223
ringens, Ammochilostoma	82	Lenticulites	223
Biloculina	471, 472	umbonata, Cristellaria	223
denticulata, Biloculina	476	rotunda, Miliolina	460
Haplophragmium	81	Triloculina	460
Haplophragmoides	84	rotundata, Ammochilostoma	97
striolata, Biloculina	477	Clavulina	157
Trochammina	81	Glandulina	185
Robulina articulata	222	Nodosaria (Glandulina)	185
calcar	231	rubra, Globigerina	289
costata	239	Millepora	364
cultrata	220	Polytrema	364
echinata	233	rubrum, Homotrema	364
Cristellaria	233	rufescens, Bathysiphon	43
orbicularis	224	rufum, Bathysiphon	42
subnodosa	367	rufus, Bathysiphon	42
vortex	227	rupertiana, Miliolina	464
Robulus cultratus	220	Triloculina	464
robusta, Bolivina	129	rugosa, Discorbina	306
compacta, Bolivina	137	Discorbis	306
Dendrophrya ramosa	56	Gaudryina	146
Gaudryina	152	Plecanium	114
Spiroloculina	404	Quinqueloculina	429
roemeri, Dentalina	193	Rosalina	306
Nodosaria	193	Textularia	113, 114
Rosalina ammonoides	323	Saccammina	48
bertheloti	305	sphaerica	48
bulloides	309	catenulata	48
lateralis	336	Saccamininae	46
orbicularis	305	Saccorbiza	54
poeyi	308	ramosa	54
rugosa	306	sacculifera, Globigerina	291
rostrata, Truncatulina	321	Sagenella frondescens	60
Rotalia	345	Sagenina	60
beccarii	345	divaricans	60
broeckhiana	346	frondescens	60
calcar	346	sagittula, atrata, Textularia	103
(Gyroidiina) orbicularis	346	candeiana, Textularia	109
soldanii	349	fistulosa, Spiroplecta	107
menardii	333	Textularia	104
orbicularis	346	Polymorphum	103
papillosa	347	Textularia	103
compressiuscula	347	var	113
praecineta	318	sagra, Articulina	448
punctulata	328	Sagrina bifrons	277
repanda	326	Siphogenerina	277
schroeteriana	347	columellaris	276
soldanii	349	dimorpha	279
squamosa	308	Uvigerina	279
truncatulinoidea	339	raphanus, Uvigerina	280
(Turbinulina) beccarii	345	striata	280
elegans	342	Siphogenerina	280
Rotaliidae	301	Saracenaria italica	252
Rotalina akneriana	316	Cristellaria	252
canariensis	338	sarsii, Biloculina	471
concamerata	328	saulei, Discorbina	307
contraria	167	scabra, Gaudryina	146
crassa	338	Pulvinulina	330

	Page.		Page.
scabra, Pulvinulina oblonga.....	330	soldanii, Haplostiche.....	78
scalariformis, Lagena hexagona.....	177	Lituola.....	78
scalaris, Nautilus (Orthoceras).....	199	Rotalia.....	349
Nodosaria.....	199	(Gyroïdina).....	349
separans, Nodosaria.....	200, 202	solita, inflata, Textularia.....	111
schloenbachi, Cristellaria.....	249	soluta, Dentalina.....	192
schlumbergeri, Sigmoidina.....	449	Nodosaria.....	192
schreibersiana, Virgulina.....	169	sororia, Polymorphina.....	265
schreibersii, Pulvinulina.....	341	(Guttulina).....	265
schroeteriana, Rotalia.....	347	spengleri, Calcarina.....	351, 336
schwageri, Uvigerina.....	270	var.....	353
schwageriana, Bolivina.....	130	Nautilus.....	351
scitulum, Haplophragmoides.....	80	sphaera, Biloculina.....	413
sclerotica, Miliolina.....	441	Planispirina.....	413
Quinqueloculina.....	441	Sphaerammina.....	99
scorpiurus, Reophax.....	65	ovalis.....	100
serobiculata, Spiroloculina.....	406	sphaerica, catenulata, Saccammina.....	48
Uvigerina.....	276	Saccammina.....	48
sculpturata, Bolivina.....	134	sphaeriloculum, Haplophragmoides.....	83
Quinqueloculina.....	433	sphaeroides, Pullenia.....	299
seceans, reticulata, Massilina.....	441	Nonionina.....	299
tenuistriata, Massilina.....	445	Sphaeroidina.....	297
selseyensis, Uvigerina.....	272	bulloides.....	297
semialata, Textularia.....	116	dehiscens.....	297
semicostata, Bolivina.....	134	sphaeroidiaiformis, Ammosphaeroidina.....	98
seminulum, Miliolina.....	416	Haplophragmium.....	98
Quinqueloculina.....	416	sphaerulatus, Baculogypsina.....	359
Serpula.....	416	spiculotestus, Reophax.....	72
semirugosa, Nodosaria.....	187	Spiniterules costatus.....	239
pyrula.....	187	spinescens, Bulimina pyrula.....	164
semistriata, Lagena.....	179	spinigera, Vaginulina.....	259
striata.....	179	spinipes, Cristellaria tricarinata.....	230
separans, Nodosaria scalaris.....	200, 202	spinulosa, Bulimina pupoides.....	164
Serpula concamerata.....	328	Verneulina.....	141
lactea.....	265	Spirillina.....	301
(Lagena) marginata.....	182	limbata.....	301
sulcata.....	179	denticulata.....	302
lobatula.....	313	tuberculo-limbata.....	302
seminulum.....	416	tuberculo-limbata.....	302
Serpulella vagans.....	55	Spirillininae.....	301
serrata, Biloculina.....	471	Spirolina.....	89
Ehrenbergina.....	172	agglutinans.....	89, 92
Siderolites.....	357	Spirolinites cylindracea.....	483
tetraedra.....	358	Spiroloculina.....	394
sigmoidea, Planispirina.....	448	acutimargo.....	398, 410
Sigmoidina.....	448	advena.....	410
Sigmoidina.....	448	affixa.....	410
celata.....	449	antillarum.....	407
schlumbergeri.....	449	angulata.....	408
sigmoidea.....	448	arenaria.....	411
simplex, Nodosaria.....	186	asperula.....	447
Siphogenerina.....	276	canaliculata.....	395
bifrons.....	277	circularis.....	403
striatula.....	278	convexiuscula.....	409
columellaris.....	276	costifera.....	405
dimorpha.....	279	plena.....	405
raphanus.....	280	crenata.....	445
costulata.....	281	depressa.....	394
(Sagrina) bifrons.....	277	elegans.....	406
striata.....	280	excavata.....	396
striata.....	280	fragilissima.....	451
siphonella, Gaudryina.....	151	grata.....	407
Siphonina.....	322	angulata.....	408
reticulata.....	322	gratcloupi.....	396
siphonifera, Anomalina polymorpha.....	325	incisa.....	397
Spiroplecta.....	115	impressa.....	395
Textularia.....	115	inaequilateralis.....	410

	Page.		Page.
<i>Spiroloculina limbata</i>	394	<i>subobtusata, Discorbis</i>	304
<i>millettii</i>	400	<i>subornata, Pulvinulina berthelotiana</i>	333
<i>nitida</i>	408	<i>suborbicularis, Heterostegina</i>	385
<i>orbis</i>	403	<i>Miliolina</i>	461
<i>planissima</i>	401	<i>Quinqueloculina</i>	461
<i>robusta</i>	404	<i>Triloculina</i>	461
<i>serobiculata</i>	406	<i>subperversa, Nodosaria</i>	208
<i>tenuimargo</i>	398	<i>subpolygona, Nodosaria</i>	198
<i>tenuis</i>	400	<i>subrotundata, Gaudryina</i>	146
<i>tenuiseptata</i>	401	<i>subscalaris, Nodosaria</i>	201
<i>tenuissima</i>	400	<i>pauci-costata, Nodosaria</i>	201
<i>validus</i>	404	<i>subsquammosa, Virgulina</i>	168
<i>Spiroplecta</i>	102	<i>substriata, Lagena alveolata</i>	182
<i>bulbosa</i>	102	<i>auriculata</i>	182
<i>sagittula, fistulosa</i>	107	<i>substriatula, Cornuspira involvens</i>	390
<i>siphonifera</i>	115	<i>Nodosaria</i>	204
<i>spengleri, Calcarina</i>	351	<i>subteres, Bulimina</i>	167
<i>Nautilus</i>	351	<i>Buliminella</i>	167
<i>Spiroplectinae</i>	102	<i>subtertuata, Nodosaria</i>	213
<i>spirostriolata, Nodosaria</i>	212	<i>subumbonata, Cristellaria orbicularis</i>	226
<i>Sporadotrema</i>	365	<i>suleata, Lagena</i>	179
<i>cylindricum</i>	365	<i>Serpula (Lagena)</i>	179
<i>squamata, Trochammina</i>	95	<i>apiculata, Lagena</i>	180
<i>squammosa, Rotalia</i>	308	<i>Technitella</i>	50
<i>staphyllearia, Fissurina</i>	181	<i>legumen</i>	50
<i>Lagena</i>	181	<i>tenera, Truncatulina</i>	318
<i>stipitata, costulata, Nodosaria</i>	187	<i>tenuimargo, Ammobaculites</i>	90
<i>striata, Dimorphina</i>	280	<i>Haplophragmium</i>	90
<i>haidingeri, Lagena</i>	178	<i>Spiroloculina</i>	399
<i>Lagena</i>	177, 178	<i>Truncatulina</i>	313
<i>Oolina</i>	177	<i>tenuis, Ammodiscus</i>	62
<i>Planispirina</i>	413	<i>Cristellaria</i>	250
<i>reticulosa, Vertebralina</i>	414	<i>Margulinina</i>	250
<i>Sagrina</i>	280	<i>Spiroloculina</i>	400
<i>semistriata, Lagena</i>	179	<i>tenuiseptata, Spiroloculina</i>	401
<i>Siphogenerina</i>	280	<i>tenuissima, Spiroloculina</i>	400
<i>(Sagrina)</i>	280	<i>tennistriata, Massilina secans</i>	445
<i>strumosa, Lagena</i>	178	<i>Uvigerina</i>	269
<i>Uvigerina</i>	271	<i>terquemiana, Miliolina</i>	458
<i>Vertebralina</i>	413	<i>tricarinata</i>	458
<i>striatula, Marginulina</i>	255	<i>Triloculina</i>	458
<i>Nodosaria (Glandulina) laevigata</i> ..	186	<i>testacea, Proteonina</i>	50
<i>Siphogenerina bifrons</i>	278	<i>Reophax diffugiformis</i>	50
<i>stricta, Textularia</i>	107	<i>tetraedra, Calcarina</i>	358
<i>striolata, Biloculina denticulata</i>	477	<i>Siderolites</i>	358
<i>ringens</i>	477	<i>Textilaria chilostoma</i>	150
<i>strumosa, Lagena</i>	178	<i>Textularia</i>	103
<i>striata</i>	178	<i>abbreviata</i>	122
<i>subaffinis, Bulimina</i>	166	<i>agglutinans</i>	106, 113
<i>subangularis, Bolivina</i>	134	<i>fistula</i>	107
<i>subcanaliculata, Nodosaria</i>	204	<i>nodosaria</i>	125
<i>subcarinata, Cristellaria orbicularis</i>	225	<i>porrecta</i>	109
<i>subcuneata, Quinqueloculina crassa</i>	423	<i>aperturalis</i>	111
<i>subdecorata, Cristellaria costata</i>	241	<i>aspera</i>	112
<i>subdepressa, Virgulina</i>	170	<i>candeiana</i>	109
<i>subglobosa, Cassidulina</i>	171	<i>carinata</i>	121
<i>Lituola</i>	81	<i>catenata</i>	112
<i>subglobosum, Haplophragmium</i>	81	<i>concaea</i>	149
<i>Haplophragmoides</i>	81	<i>conica</i>	123
<i>sublaevis, Cristellaria costata</i>	242	<i>crassisepta</i>	121
<i>sublineata, Nodosaria</i>	207	<i>excavata</i>	120
<i>hispida</i>	207	<i>fluitii</i>	113
<i>submamilligera, Cristellaria</i>	235	<i>foliacea</i>	117
<i>subnodosa, Dentalina</i>	139	<i>goësi</i>	113
<i>Pleurostomella</i>	139	<i>gramen</i>	105
<i>Polystomella</i>	367	<i>hauerii</i>	105
<i>Robulina</i>	367	<i>immensa</i>	118

	Page.		Page.
Textularia jugosa.....	119	Trioculina reticulata.....	434
porrecta.....	109	rotunda.....	460
psendocarinata.....	121	rupertiana.....	464
quadrilatera.....	125	suborbicularis.....	461
rugosa.....	113, 114	terquemiana.....	458
sagittula.....	103	tortuosa.....	464
atrata.....	103	tricarinata.....	454
caudeiana.....	109	convexa.....	456
fistulosa.....	104	plicata.....	456
semialata.....	116	trigonula.....	452
siphonifera.....	115	multistriata.....	453
solita, inflata.....	111	valvularis.....	460
stricta.....	107	Triplesia.....	219
transversaria.....	120	tricarinata.....	219
tricarinata.....	145	triquetra, Flintina.....	466
triquetra.....	112	Miliolina.....	466
trochus.....	124	Textularia.....	142
(Verneulina) triquetra.....	112	(Verneulina).....	142
vertebralis.....	110	Verneulina.....	142
Textulariidae.....	102	Tritaxia.....	153
Textulariinae.....	103	caperata.....	153
Tholosina.....	51	indiscreta.....	145
bulla.....	51	tricarinata.....	145
Thurammia.....	52	Tritaxilina.....	153
papillata, castanea.....	52	caperata.....	153
papyracea.....	52	Trochammia.....	95
tibia, Nodobaularia.....	415	globigeriniformis.....	96
Tinoporos baculatus.....	354	irregularis.....	61
Tolypammia.....	55	clavata.....	61
vagans.....	55	lituiformis.....	88
tortuosa, Triloculina.....	464	proteus.....	79, 95
transversaria, Textularia.....	120	ringens.....	81
Tretomphalus.....	309	squamata.....	95
bulloides.....	309	trullisata.....	88
triangularis, austriaca, Valvulina.....	141	turbinata.....	96
conica, Valvulina.....	142	turbinatum.....	96
Gaudryina.....	148	Trochammininae.....	79
Valvulina.....	142	Trochamminoides.....	79
tricarinata, bertheliniana, Miliolina.....	457	proteus.....	79
convexa, Triloculina.....	456	trochus, Textularia.....	113, 124
Miliolina.....	454, 457, 458	trullisata, Trochammia.....	88
plicata, Triloculina.....	456	Truncatulina.....	312
terquemiana, Miliolina.....	458	akneriana.....	316
Textularia.....	145	culter.....	320
Triloculina.....	454	grosserugosa.....	323
Triplesia.....	219	haidingerii.....	315
Tritaxia.....	145	humilis.....	321
Vaginulina.....	219	lobatula.....	313
tricarinatum, Rhabdogonium.....	219	margaritifera.....	319
tricarinnella, Cristellaria.....	230	praecincta.....	318
spinipes, Cristellaria.....	230	refulgens.....	312
trigona, Allomorphina.....	284	reticulata.....	322
trigonula, Miliolites.....	452	rostrata.....	321
multistriata, Triloculina.....	453	tenera.....	318
Triloculina.....	452	tenuimargo.....	313
triloba, Candaina nitida.....	296	ungeriana.....	317
globigerina.....	286	ornata.....	317
bulloides.....	286	variabilis.....	314
Triloculina.....	452	wuellerstorfi.....	314
advena.....	453	truncatulinoidea, Pulvinulina.....	339
bertheliniana.....	457	Rotalina.....	339
circularis.....	462	tuberculata, Heterostegina.....	384
cuneata.....	461	depressa.....	384
incisa.....	462	tuberculo-limbata, Spirillina.....	302
insignis.....	458	limbata.....	302
linneiana.....	463	tubulata, Lagena hispida.....	177
oblonga.....	459	tubulosa, Biloculina.....	475

	Page.		Page.
tumida, Pulvinulina.....	337	Valvulina.....	142
menardii.....	337	allomorfinoides.....	306
tumido-costata, Cristellaria.....	245	angularis.....	155
labyrinthica, Cristellaria.....	246	austriaca.....	144
turbinata, Trochammina.....	96	capreolus.....	127
turbinatum, Haplophragmium.....	96	conica.....	142
Trochammina.....	96	fusa.....	143
(Turbinulina) beccarii, Rotalia.....	345	parisiensis.....	156
elegans, Rotalia.....	342	triangularis.....	142, 155
umbilicatulata, Nonionna.....	365	austriaca.....	144
umbilicatulata, Nautilus.....	365	conica.....	142
umbonata, Cristellaria rotulata.....	223	variabilis, Cristellaria.....	229
Pulvinulina.....	339	Truncatulina.....	314
Rotalina.....	339	venosa, Nimmulites.....	383
undosa, Miliolina.....	432	Operculina.....	383
ungarica, ornata, Truncatulina.....	317	venosus, Nautilus.....	383
Planorbulina.....	317	venusta, Miliolina.....	420
Rotalina.....	317	Quinqueloculina.....	420
Truncatulina.....	317	Vermiculum globosum.....	173
universa, Orbulina.....	295	lacteum.....	265
utricularis, Carpenteria.....	360	laeve.....	173
Polytrema.....	360	marginatum.....	182
Uvigerina.....	269	oblongum.....	459
aculeata.....	273	Verneuilina.....	139
ampullacea.....	274	affixa.....	142
angulosa.....	271	bradyi.....	141
asperula.....	274	communis.....	154
ampullacea.....	274	polystropha.....	139
brunnensis.....	272	propinqua.....	140
canariensis.....	275	pygmaea.....	141
interrupta.....	275	spinulosa.....	141
nodosa.....	275	triquetra.....	142
pigmea.....	269	Textularia.....	142
pygmaea.....	269	Verneuilinae.....	139
(Sagrina) dimorpha.....	279	Vertebrulina.....	414
raphanus.....	280	striata.....	414
schwageri.....	270	reticulosa.....	414
serobiculata.....	276	vertebralis, Nautilus (Orthoceras).....	211
selseyensis.....	272	Nodosaria.....	211
striata.....	271	plicata, Marginopora.....	487
tenuistriata.....	269	Textularia.....	110
Uvigerinae.....	269	respertilio, Biloculina.....	472
vagans, Givanelia.....	55	vildeboana, Discorbis.....	304
Hyperammina.....	55	Virgulina.....	168
Serpulella.....	55	cornuta.....	168
Tolypammina.....	55	pauciloculata.....	168
Vaginulina.....	257	schreibersiana.....	168
acicula.....	262	subdepressa.....	168
bradyi.....	260	subsquamosa.....	168
brukenenthalii.....	260	vortex, Cristellaria.....	227
elegans.....	258	Nautilus.....	227
legumen.....	257	Polystomella.....	227
elegans.....	258	Robulina.....	227
linearis.....	261	vulgaris, larvata, Planorbulina.....	310
margaritifera.....	259	Miliolina.....	417
patens.....	263	Quinqueloculina.....	417
protumida.....	262	Vulvulina alata.....	129
spinigera.....	259	Webbina clavata.....	61
tricarinata.....	219	Webbinella clavata.....	61
validus, Spiroloculina.....	404	wuellerstorfi, Anomalina.....	314
valvularis, Miliolina.....	460	Truncatulina.....	314
Triloculina.....	460		

SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01421 1601