

# Limited photosynthetic plasticity in the leaf-succulent CAM plant *Agave angustifolia* grown at different temperatures

Joseph A. M. Holtum<sup>A,B,C</sup> and Klaus Winter<sup>B</sup>

<sup>A</sup>Tropical Biology, James Cook University, Townsville, Qld 4811, Australia.

<sup>B</sup>Smithsonian Tropical Research Institute, PO Box 0843-03092, Balboa, Ancon, Republic of Panama.

<sup>C</sup>Corresponding author. Email: joseph.holtum@jcu.edu.au

**Abstract.** In *Agave angustifolia* Haw., a leaf-succulent constitutive crassulacean acid metabolism (CAM) plant of tropical Panama, we tested whether nocturnal CO<sub>2</sub> uptake and growth were reduced at night temperatures above 20°C. Unlike some CAM model species from habitats with pronounced day-night temperature variations, in *A. angustifolia* temperature affected little the relative contributions of CAM and C<sub>3</sub> photosynthesis to growth. In plants grown under 12 h light/dark regimes of 25/17, 30/22 and 35/27°C, biomass increased with temperature. Maintaining day temperature at 35°C and reducing night temperature from 27 to 17°C markedly lowered growth, a reduction partially reversed when roots were heated to 27°C. Across all treatments, whole-shoot δ<sup>13</sup>C values ranged between –14.6 and –13.2 ‰, indicating a stable proportion of CO<sub>2</sub> was fixed at night, between 75 and 83%. Nocturnal acidification reflected growth, varying between 339 and 393 μmol H<sup>+</sup> g<sup>-1</sup> fresh mass and 63–87 μmol H<sup>+</sup> cm<sup>-2</sup>. In outdoor open-top chambers, warming the air 3°C above ambient at night did not reduce biomass accumulation. The persistence of a high capacity for nocturnal CO<sub>2</sub> fixation at the expense of a limited capacity for switching between C<sub>3</sub> and CAM probably makes this *Agave*, and others like it, potential species for biomass production in seasonally-dry landscapes.

**Additional keywords:** biofuel, C<sub>3</sub> photosynthesis, climate change, crassulacean acid metabolism, open-top chamber.

Received 30 September 2013, accepted 19 February 2014, published online 28 April 2014

## Introduction

Crassulacean acid metabolism (CAM) in terrestrial plants is a water-use efficient photosynthetic adaptation that assists plants to occupy habitats subject to periodic water stress (Osmond 1978; Winter 1985). CAM is water-use efficient because CO<sub>2</sub> is assimilated at night when temperatures and rates of evaporation are at daily minima (Winter *et al.* 2005).

In addition to fixing CO<sub>2</sub> during the night, many species with CAM can also incorporate atmospheric CO<sub>2</sub> during the light via C<sub>3</sub> photosynthesis, a pathway that is less water-use efficient than CAM. It is not uncommon for CAM species with a capacity for light fixation to adjust the proportions of carbon fixed at night and during the day in response to the availability of water (Winter 1985; Borland *et al.* 2011; Dodd *et al.* 2002) such that CO<sub>2</sub> fixation in the light increases when water is available and decreases when water is scarce. The ability to regulate CO<sub>2</sub> uptake during the dark and the light thus enables many CAM species to maintain some daily carbon gain as they reduce water-loss, whereas C<sub>3</sub> and C<sub>4</sub> plants may not be able to maintain a positive carbon balance under similar stressful conditions.

Evaporative water loss increases with the difference in vapour pressure between the leaf and the atmosphere, and thus responds to increasing temperature. It is therefore not surprising that in CAM species the proportion of carbon fixed at night and during the day responds to temperature (De Vries 1884; Kluge and Ting

1978). In CAM plants, the temperature responses of CO<sub>2</sub> fixation during the night and the day are not the same (Nobel 1988; Yamori *et al.* 2014), as might be expected when two biochemically distinct processes are involved. CO<sub>2</sub> uptake in the dark is enhanced when daytime temperature maxima are typically above 30°C, and night maxima are below 20°C. As a result, night temperatures significantly above 20°C are not only widely considered suboptimal for nocturnal CO<sub>2</sub> uptake (Neales 1973a, 1973b; Kluge and Ting 1978; Neales *et al.* 1980; Medina and Osmond 1981) but, by extrapolation, suboptimal for growth. The night-time temperature responses are at odds with the expression of pronounced CAM in the many species (both epiphytes and terrestrial plants) that inhabit the tropics (Wong and Hew 1976; Holtum and Winter 1999; Crayn *et al.* 2004; Silvera *et al.* 2005) where night temperatures are commonly around 20–25°C.

Early studies proposed that the mechanistic basis for reduced nocturnal CO<sub>2</sub> uptake at higher night temperatures is a product of differing thermal optima of malate synthesising and consuming reactions and of increased respiration (Brandon 1967; Kaplan *et al.* 1976) but these ideas have not been further pursued experimentally. Similarly, there is little experimental evidence that the negative effects of high night temperatures on CO<sub>2</sub> uptake in the dark that have been observed by several authors ultimately adversely affect growth, even amongst the extensive

*Agave* and cactus research published by Nobel and his colleagues (see Nobel 1988 for a review).

Here we explored the relationship between growth,  $\delta^{13}\text{C}$  (as indicator of the relative contributions of light and dark fixation to carbon gain), and day and night temperatures in *Agave angustifolia* Haw., a constitutive strong-CAM species from the tropics and subtropics of Central America and Mexico. *A. angustifolia* is a putative wild ancestor of the domesticated agaves (Gentry 1982; Colunga-García Marín *et al.* 1999), *Agave fourcroydes* Lem. (henequen) which is used for fibre production, and *Agave tequilana* F.A.C.Weber, the source of tequila. Both *A. fourcroydes* and *A. tequilana* have been proposed as biofuel feedstock crops for seasonally dry landscapes in tropical northern Australia and elsewhere (Chambers and Holtum 2010; Holtum *et al.* 2011; Owen and Griffiths 2013). In a warming world containing roughly 400 million ha of abandoned agricultural land much of which is in the semiarid tropics and sub-tropics (Campbell *et al.* 2008), it could be expected that the area suitable for growing appropriate *Agave* species for biofuels might be expanding. However, if the expression of CAM in *Agave* is progressively reduced with increasing night temperature, and the expression of daytime  $\text{CO}_2$  uptake is reduced as well, then growing CAM *Agave* for biomass may become a less attractive proposition.

Here we demonstrate that *A. angustifolia* accumulates biomass more rapidly at a relatively high day/night temperature regime of 37/27°C than it does at 25/17°C. At a high daytime temperature of 35°C, growth increased when night temperature was increased from 17 to 27°C. In addition, the proportional contribution of light and dark  $\text{CO}_2$  uptake to net carbon gain, measured using  $\delta^{13}\text{C}$  values as a proxy (Winter and Holtum 2002), responded little to temperatures to which *A. angustifolia* might be exposed in its natural habitat. We suggest that this lack of photosynthetic plasticity may be a desirable trait when growing constitutive strong CAM plants for biomass production in seasonally-dry environments in the tropics and subtropics.

## Materials and methods

### Plant material

*Agave angustifolia* Haw. was collected from Playa Majagual, Panama (8°43'N, 79°45'E) and maintained outdoors in forest topsoil at the Smithsonian Tropical Research Institute, Santa Cruz Experimental Research Facility, Gamboa, Republic of Panama (9°07'N, 79°42'W).

### Experiment using growth chambers

Bulbils of  $0.68 \pm 0.09$  (s.e.) g DM with leaf areas of  $33.04 \pm 7.13$  cm<sup>2</sup> were planted in 6 L pots (16 cm internal diameter) containing 80% forest topsoil, 20% sand and 2 g Osmocote Plus fertiliser (Scotts-Sierra Horticultural Products, OH, USA). The plants ( $n = 6$  per treatment) were grown from 1/7/2012 until 24/9/2012 inside five controlled-environment chambers (Model GC15, Environmental Growth Chambers, OH, USA) operating under 12 h light/12 h dark cycles of 25/17°C, 30/22°C, 35/27°C, 35/17°C, and 35/17°C and maintaining soil temperature at  $\geq 27^\circ\text{C}$  (Table 1). Photon flux density was  $500 \mu\text{mol m}^{-2} \text{s}^{-1}$  at the top of the plants.

Air temperatures inside growth chambers, measured using copper-constantan fine wire thermocouples, were essentially identical to chamber set-temperatures and assumed their new value within 10 min of light/dark and dark/light transitions.

Soil temperatures within all pots reflected the day/night air temperatures except for pronounced lags as they gradually heated or cooled following switchovers between temperature regimes (Fig. 1). For the treatment in which roots were heated, pots containing plants were placed in a water-bath equipped with a heating thermostat (Model Alpha, Lauda, NJ, USA) set to 27°C. Soil temperature was measured in each pot using two copper-constantan thermocouples embedded in silicone sealant.

### Open-top chamber experiment


*Agave angustifolia* was grown outdoors in six open-top chambers between 1 November 2012 and 25 June 2013 at the Santa Cruz Experimental Research Facility. Each chamber

**Table 1. Growth chamber environments**

Leaf temperatures of *Agave angustifolia* were measured in the middle of the light period and in the dark (1 h before onset of the light) using an infrared thermometer (MT6; Raytek, CA, USA), the readings of which were verified using fine wire copper-constantan thermocouples. Values are means of six measurements  $\pm$  s.e.

Air temperature (°C)		Relative humidity (%)		Leaf temperature (°C)	
Light	Dark	Light	Dark	Light	Dark
25	17	56	76	27.4 $\pm$ 0.4	15.1 $\pm$ 0.1
30	22	40	57	32.8 $\pm$ 0.3	20.5 $\pm$ 0.0
35	27	29	41	38.5 $\pm$ 0.2	26.1 $\pm$ 0.1
35	17	29	77	37.5 $\pm$ 0.3	15.6 $\pm$ 0.0
35 <sup>A</sup>	17 <sup>A</sup>	29	77	38.2 $\pm$ 0.4	17.7 $\pm$ 0.2

<sup>A</sup>In this treatment, the roots were heated to achieve 27°C during the dark. Heating was maintained during the light.



**Fig. 1.** A 24 h day-night cycle of soil temperatures in pots containing *Agave angustifolia* plants exposed in growth cabinets to 12 h day/night temperature regimes of 25/17°C (1), 30/22°C (2), 35/27°C (3), 35/17°C (4) and 35/17°C plus roots warmed to achieve 27°C (5). Open bar denotes light period, closed bar denotes dark period. Measurements were at 5 min intervals.

contained six 19 L containers each with one bulbil planted in forest soil. At the onset of the experiment, the average weight of bulbils was  $0.99 \pm 0.10$  g DM and the average area of bulbil leaves was  $41.38 \pm 3.20$  cm<sup>2</sup>. The chambers and environmental control within the chambers are described by Cheesman and Winter (2013). In three chambers night temperature was maintained at 3°C above the ambient night temperature. Pots received ambient rainfall with additional daily watering as required to maintain soils at close to field capacity.

At harvest, soil was washed by hand from the roots. Leaf area was measured using a leaf area meter (Model LI-3100, Li-Cor, Lincoln, NE, USA). Roots, leaves and stems were oven-dried at 70°C.

#### Titrateable acidity

From mature leaves of each plant, two discs, each of 0.9 cm diameter, were excised using a cork borer at the end of the light and dark periods and were frozen in liquid nitrogen. Organic acids were extracted by sequentially boiling samples in 50% methanol and in water for 5 min. Extracts were cooled to room temperature and titrated with 10 mM KOH to pH 6.5 (Holtum *et al.* 2004).

#### Stable isotope analysis

The  $\delta^{13}\text{C}$  values of finely-ground homogeneous powder from the pooled dried leaves of whole plants were measured using an isotope ratio mass spectrometer (Delta V; Thermo Fisher Scientific, Ottawa, ON, Canada) in the Stable Isotope Laboratory of the Smithsonian Tropical Research Institute (Cernusak *et al.* 2011; Winter *et al.* 2014).


## Results

Whole-plant mean dry biomass recorded at the two higher day/night temperature treatments, 30/22 and 35/27°C, was significantly greater than that of plants grown at 25/17°C (Fig. 2a), with plants at 35/27°C, the warmest regime, having a mean biomass that was 27% greater than for plants at 25/17°C, the coolest regime.

Compared with 35/27°C, maintaining the day temperature at 35°C and reducing the night temperature to 17°C lowered mean biomass from 8.8 to 5.9 g plant<sup>-1</sup>, a significant drop of 33%. In contrast, the mean biomass of plants at 25/17°C and 35/17°C was similar.

The response pattern of leaf area to temperature was qualitatively similar to the biomass pattern but overall amplified (Fig. 2b). Compared with the mean leaf area of plants at 25/17°C, leaf area was 25% greater at 30/22°C and 49% greater at 35/27°C. As with biomass, the mean leaf area of plants grown at 35/17°C was 32% less than for plants grown at 35/27°C. Leaf area ratio (LAR) increased with temperature (Fig. 2c) whereas leaf mass/area (LMA), a measure of leaf density, remained constant or decreased slightly (Fig. 2d).

Maintaining roots at 27°C during a 35/17°C air temperature regime increased leaf area and LAR significantly (Fig. 2b, c), whereas LMA decreased slightly (Fig. 2d). The heating of roots was accompanied by an associated 2.1°C increase in leaf temperature at night (Table 1), presumably the product of



**Fig. 2.** Biomass (a), leaf area (b), leaf area ratio (leaf area per whole-plant dry mass) (c), leaf mass/area (leaf dry mass per unit leaf area) (d) and root mass ratio (root dry mass per whole-plant dry mass) (e) of *Agave angustifolia* plants exposed to 12 h day/night temperature regimes of 25/17°C, 30/22°C, 35/27°C, 35/17°C, and 35/17°C plus roots warmed to 27°C. Bars indicate 95% confidence levels of the means ( $n=6$  plants). Letters within columns indicate statistical significance, categories not sharing a letter differ significantly at  $P<0.05$  (Tukey's HSD).

thermal dissipation from H<sub>2</sub>O in the circulating bath used to heat the roots.

Nocturnal acid accumulation ( $\Delta\text{H}^+$ ) ranged between 339 and 393  $\mu\text{mol H}^+ \text{g}^{-1}$  FM and 63–87  $\mu\text{mol H}^+ \text{cm}^{-2}$  across treatments (Table 2). Acid levels were similar in plants grown at 25/17, 30/22 and 35/27°C, with the largest mean acid accumulation measured for plants at 30/22°C. In plants grown at 35°C during the light, mean acidification values were not significantly affected when the night temperature

**Table 2.** Nocturnal increases in leaf tissue acidity of *Agave angustifolia* grown under five 12 h light/12 h dark temperature regimes

Samples were taken from mature leaves at the end of the light and end of the dark periods. Values are means  $\pm$  s.e. ( $n=6$  plants). Means followed by the same letter do not differ (Tukey's HSD test,  $P < 0.05$ )

Titratable acidity	Light/dark temperature ( $^{\circ}\text{C}$ )				
	25/17	30/22	35/27	35/17	35/17 + root warming <sup>A</sup>
	$\mu\text{mol g}^{-1}$ fresh mass				
Light	34 $\pm$ 4	13 $\pm$ 1	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0
Dark	399 $\pm$ 7	406 $\pm$ 10	371 $\pm$ 5	343 $\pm$ 8	340 $\pm$ 12
$\Delta$	364 $\pm$ 10ab	393 $\pm$ 10a	371 $\pm$ 5ab	343 $\pm$ 8b	339 $\pm$ 12b
	$\mu\text{mol cm}^{-2}$				
Light	8 $\pm$ 4	3 $\pm$ 3	0 $\pm$ 2	0 $\pm$ 3	0 $\pm$ 2
Dark	88 $\pm$ 4	90 $\pm$ 3	75 $\pm$ 2	67 $\pm$ 3	63 $\pm$ 2
$\Delta$	80 $\pm$ 4ab	87 $\pm$ 3a	75 $\pm$ 2abc	67 $\pm$ 3bc	63 $\pm$ 2c

<sup>A</sup>In this treatment, the roots were heated to achieve 27 $^{\circ}\text{C}$  during the dark. Heating was maintained during the light.

**Table 3.**  $\delta^{13}\text{C}$  values of pooled leaves of *Agave angustifolia* plants grown under five 12 h light/12 h dark temperature regimes

Values are means  $\pm$  s.e. ( $n=6$  plants). Means followed by the same letter do not differ (Tukey's HSD test,  $P < 0.05$ )

	Light/dark temperature ( $^{\circ}\text{C}$ )				
	25/17	30/22	35/27	35/17	35/17 + root warming <sup>A</sup>
$\delta^{13}\text{C}$ ( $\text{‰}$ )	-14.6 $\pm$ 0.1a	-14.5 $\pm$ 0.0a	-14.4 $\pm$ 0.1a	-13.2 $\pm$ 0.1b	-13.2 $\pm$ 0.1b

<sup>A</sup>In this treatment, the roots were heated to achieve 27 $^{\circ}\text{C}$  during the dark. Heating was maintained during the light.

was reduced from 27 to 17 $^{\circ}\text{C}$  or when roots were warmed to 27 $^{\circ}\text{C}$  during 17 $^{\circ}\text{C}$  nights.

As an indicator of the proportional contribution of  $\text{CO}_2$  uptake in the dark and the light to shoot carbon gain,  $\delta^{13}\text{C}$  values differed by only 1.4  $\text{‰}$  across all treatments (Table 3). Plants cultured under day/night regimes of 25/17, 30/22 and 35/27 $^{\circ}\text{C}$  had similar  $\delta^{13}\text{C}$  values of between -14.4 and -14.6  $\text{‰}$ . The least negative  $\delta^{13}\text{C}$  value, -13.2  $\text{‰}$ , were observed for plants at 35/17 $^{\circ}\text{C}$  irrespective of whether the roots were heated.

For well-watered plants grown in open-top chambers under ambient light, elevating night temperature by 3 $^{\circ}\text{C}$  did not affect plant dry mass or leaf area (Table 4).

## Discussion

The growth response of the constitutive strong CAM plant *A. angustifolia* to a range of day-night temperature regimes contrasted with an often-cited generalisation that high day temperatures combined with low night temperatures favour CAM and enhance growth (Kluge and Ting 1978; Owen and Griffiths 2013). In *A. angustifolia*, biomass was greater for plants grown at day temperatures above 25 $^{\circ}\text{C}$  and night temperatures above 17 $^{\circ}\text{C}$ . Nocturnal accumulation of titratable acidity was similar for plants grown at night temperatures of 17 and 27 $^{\circ}\text{C}$ .

Biomass accumulation, leaf area and LAR increased as day and night temperatures were increased from 25/17 to 35/27 $^{\circ}\text{C}$ , whereas unchanged  $\delta^{13}\text{C}$  values indicated that the relative contribution of nocturnal  $\text{CO}_2$  uptake to carbon gain remained around 75% (using the calibration of Winter and Holtum 2002). In general agreement with the CAM-type  $\delta^{13}\text{C}$  values, nocturnal acidification was substantial under the three treatments.

**Table 4.** Dry biomass and leaf area of whole plants of *Agave angustifolia* grown in six open-top chambers under ambient conditions or ambient days with nights elevated by 3 $^{\circ}\text{C}$  at the Smithsonian Tropical Research Institute, Santa Cruz Experimental Research Facility, Gamboa, Republic of Panama (9 $^{\circ}$ 07'N, 79 $^{\circ}$ 42'W)

Significance of difference of the means ( $n=3$ ) was tested using a Students  $t$ -test (2-tailed, equal variance). n.s. = not significant,  $P \geq 0.05$

Parameter	Temperature treatment		Test of significance
	Ambient	Night temperature elevated by 3 $^{\circ}\text{C}$	
Dry mass (g)	36.2 $\pm$ 0.4	42.9 $\pm$ 5.7	n.s. ( $P=0.321$ )
Leaf area ( $\text{cm}^2$ )	713 $\pm$ 12	886 $\pm$ 153	n.s. ( $P=0.323$ )

As growth is driven by overall assimilation rate and LAR, and nocturnal  $\text{CO}_2$  assimilation remained relatively constant, the principal contributor to increased growth between 25/17, 30/22 and 35/27 $^{\circ}\text{C}$  appears to have been increased LAR.

Compared with the highest temperature regime, 35/27 $^{\circ}\text{C}$ , maintaining the day temperature at 35 $^{\circ}\text{C}$  and reducing the night temperature to 17 $^{\circ}\text{C}$  markedly reduced biomass accumulation. As LAR was similar for both treatments it follows that the overall rates of  $\text{CO}_2$  uptake were lower in plants under 17 $^{\circ}\text{C}$  nights, a conclusion supported by marginally lower nocturnal acidification at 17 $^{\circ}\text{C}$ .  $\delta^{13}\text{C}$  values suggest that over the life of the plants under 27 and 17 $^{\circ}\text{C}$  nights, 76 and 83% of the carbon, that is, similar proportions, were obtained in the dark respectively.

Biomass accumulation was similar between plants subjected to 17 $^{\circ}\text{C}$  nights and days of 25 or 35 $^{\circ}\text{C}$  but LAR was greater at



the higher day temperature. As was deduced following the comparison of plant growth at 35/27 and 35/17°C, it appears that the overall rate of CO<sub>2</sub> uptake per unit area was lower in the 35/17°C plants than in the 25/17°C plants, a deduction consistent with the lower nocturnal acidification in plants at 35/17°C.

In the 35/17°C treatment, warming the soil to 27°C increased total leaf area and LAR but biomass, acidification and δ<sup>13</sup>C values remained constant. Interpretation of the effects on growth of soil heating were complicated by an associated increase of nocturnal leaf temperature by 2.1°C (Table 1), as explained in 'Materials and Methods'. Stimulation of growth following root heating has previously been reported for the C<sub>3</sub> species, cotton, possibly the consequence of temperature shifting source–sink relationships (Königer and Winter 1993).

Warming of *A. angustifolia* grown outdoors by 3°C during the night did not adversely affect growth. In fact, higher means for whole-plant dry mass (19% increase) and leaf area (24% increase) of heated plants, although not statistically different, suggest that a positive statistically significant correlation between night temperature and growth might well be observed in a more extensively replicated experiment. Significantly increased growth rates have been reported for the C<sub>3</sub> pioneer tree species, *Ficus insipida* and *Ochroma pyramidale*, maintained at 3°C above ambient at night (Cheesman and Winter 2013).

The perception that CAM activity in general is stimulated by pronounced differences in day and night temperature stems in the main from studies of species of *Kalanchoë*, *Agave* and *Opuntia* from habitats where cool nights and hot days predominate (Queiroz 1966; Neales 1973a; Szarek and Ting 1974; Kluge and Ting 1978; Nobel and McDaniel 1988), although broadly similar observations have been reported for *Ananas comosus*, a tropical species that has been modified by breeding for desirable agronomic characters (Neales 1973b; Neales *et al.* 1980; Zhu *et al.* 1999). One might expect any requirement for substantial day/night temperature differences to be less evident in species like *A. angustifolia* that are native to the humid tropics where nights are typically warm and day/night temperature fluctuations are less pronounced (Milburn *et al.* 1968; Wong and Hew 1976; Griffiths and Smith 1983; Winter 1985; Holtum and Winter 1999; Crayn *et al.* 2004; Silvera *et al.* 2005). Nobel and Hartsock (1978, 1981) demonstrated that CO<sub>2</sub> exchange by *Agave* species can rapidly acclimate to new temperature regimes but the implications of these short-term responses for growth are not clear.

On the basis that nocturnal CO<sub>2</sub> uptake is reflected in growth, minimum night-time temperatures have been used to form the temperature index components of environmental productivity indexes (EPI) designed to predict the productivities of various *Agave* and *Opuntia* in the field (Nobel 1984, 1985, 1988, 1989, 1991; Nobel and Meyer 1985; Nobel and Quero 1986; Nobel and Valenzuela 1987; Pimienta-Barrios *et al.* 2001). Similarly, a model that integrates EPI estimates with soil water retention characters and GIS methodology to predict potential productivity of *Agave* species in Australia, apportioned to minimum night-time temperatures 95% of the effects of temperature on productivity in *A. fourcroydes* (Owen and

Griffiths 2013). The observations on the growth of *A. angustifolia* reported here suggest that EPI modelling based on minimum temperatures may be a too simplistic approach, particularly for tropical CAM species.

The responses of growth in *A. angustifolia* to day and night temperatures will not only reflect the effects of temperature on the CAM cycle but also the effects of temperature on other cell processes. Diel patterns of leaf and cladode expansion have been observed in CAM *M. crystallinum*, and in CAM perennials such as *Opuntia engelmannii*, *O. oricola* and *Kalanchoë beharensis* (Gouws *et al.* 2005) but the interaction between the demands of CAM cells for carbon skeletons and the demands by the rest of the plant for carbon skeletons and energy are not well understood, even for long-studied models such as the annual *Mesembryanthemum crystallinum* and perennial *Kalanchoë* spp. (Borland and Dodd 2002; Antony and Borland 2009). It has been proposed that the substantial requirements for carbon skeletons in CAM cells limits the export of carbohydrates for growth during acidification at night, and during deacidification in the light (Gouws *et al.* 2005; Haider *et al.* 2012). Growth could be particularly sensitive to temperature when carbon export for growth is expected to be most evident – during the first and last phases of the light, and perhaps late at night when the accumulation of malic acid is close to completion and turgor pressures are high.

Photosynthetic plasticity, which is the ability to switch between C<sub>3</sub> and CAM photosynthesis, contributes to the ecological success of some CAM species, particularly annuals, trees and hemi-epiphytes in seasonally-dry habitats (Zotz and Winter 1994a, 1994b; Winter and Holtum 2014). These photosynthetically highly flexible species often exhibit reduced rates of growth when CAM is the predominate photosynthetic pathway. In contrast, many open-field CAM perennials such as *A. angustifolia* and its close relatives, *A. fourcroydes* and *A. tequilana*, combine strong CAM with substantial growth rates. In these species, the persistence of CAM throughout large parts of the dry season can enable appreciable rates of growth in concert with high water-use efficiency (Nobel 1985, 1988, 1991; Nobel and Valenzuela 1987). It is this ability to exhibit year-round water-use efficient growth in seasonally-dry conditions that enables agaves to accumulate biomass at sizeable annualised rates and has resulted in them being considered as biofuel feedstock crops in regions where growth of traditional food and bioenergy crops is suboptimal (Nobel 1988; Borland *et al.* 2009; Davis *et al.* 2011; Holtum *et al.* 2011).

In conclusion, consistent with its tropical occurrence, *A. angustifolia* successfully grew at high day/night temperatures, and low night temperatures combined with high day temperatures adversely affected growth. In all situations, CAM was the principal contributor to carbon gain, suggesting limited photosynthetic plasticity in this leaf-succulent constitutive CAM species.

There is substantial literature on the effects of temperature on daily CO<sub>2</sub> exchange patterns of CAM plants. What is clearly needed is research that extends these studies to temperature effects on growth, particularly of species native to habitats with contrasting temperature regimes.

## Acknowledgements

The authors acknowledge the contributions of J Aranda, who oversaw the open-top chamber experiment, M. Garcia who maintained data-logging equipment, and A Virgo who assisted with an illustration. The research was supported by funds from the Smithsonian Tropical Research Institute. JAMH was supported by the School of Marine and Tropical Biology, James Cook University.

## References

- Antony E, Borland AM (2009) The role and regulation of sugar transporters in plants with crassulacean acid metabolism. *Progress in Botany* **70**, 127–143. doi:10.1007/978-3-540-68421-3\_6
- Borland AM, Dodd AN (2002) Carbohydrate partitioning in crassulacean acid metabolism plants: reconciling potential conflicts of interest. *Functional Plant Biology* **29**, 707–716. doi:10.1071/PP01221
- Borland AM, Griffiths H, Hartwell J, Smith JAC (2009) Exploiting the potential of plants with crassulacean acid metabolism for bioenergy production on marginal lands. *Journal of Experimental Botany* **60**, 2879–2896. doi:10.1093/jxb/erp118
- Borland AM, Zambrano VAB, Ceusters J, Shorrocks K (2011) The photosynthetic plasticity of crassulacean acid metabolism: an evolutionary innovation for sustainable productivity in a changing world. *New Phytologist* **191**, 619–633. doi:10.1111/j.1469-8137.2011.03781.x
- Brandon PC (1967) Temperature features of enzymes affecting crassulacean acid metabolism. *Plant Physiology* **42**, 977–984. doi:10.1104/pp.42.7.977
- Campbell JE, Lobell DB, Genova RC, Field CB (2008) The global potential of bioenergy on abandoned agriculture lands. *Environmental Science & Technology* **42**, 5791–5794. doi:10.1021/es800052w
- Cermusak L, Winter K, Martinez C, Correa E, Aranda J, Garcia M, Jaramillo C, Turner BL (2011) Responses of legume versus nonlegume tropical tree seedlings to elevated CO<sub>2</sub> concentration. *Plant Physiology* **157**, 372–385. doi:10.1104/pp.111.182436
- Chambers D, Holtum JAM (2010) 'Feasibility of *Agave* as a feedstock for biofuel production in Australia.' (Rural Industries Research and Development Corporation: Canberra)
- Cheesman A, Winter K (2013) Elevated night-time temperatures increase growth in seedlings of two tropical pioneer tree species. *New Phytologist* **197**, 1185–1192. doi:10.1111/nph.12098
- Colunga-García Marín P, Coello-Coello J, Eguiarte LE, Piñero D (1999) Isozymatic variation and phylogenetic relationships between henequén (*Agave fourcroydes*) and its wild ancestor *A. angustifolia* (Agavaceae). *American Journal of Botany* **86**, 115–123. doi:10.2307/2656960
- Crayn DM, Winter K, Smith JAC (2004) Multiple origins of crassulacean acid metabolism and the epiphytic habit in the Neotropical family Bromeliaceae. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 3703–3708. doi:10.1073/pnas.0400366101
- Davis SC, Dohleman FG, Long SP (2011) The global potential for *Agave* as a biofuel feedstock. *Global Change Biology - Bioenergy* **3**, 68–78. doi:10.1111/j.1757-1707.2010.01077.x
- De Vries H (1884) Über die periodische Säurebildung der Fettpflanzen. *Botanische Zeitung* **42**, 339–344.
- Dodd AN, Borland AM, Haslam RP, Griffiths H, Maxwell K (2002) Crassulacean acid metabolism: plastic, fantastic. *Journal of Experimental Botany* **53**, 569–580. doi:10.1093/jexbot/53.369.569
- Gentry HS (1982) 'Agaves of continental North America.' (University of Arizona Press: Tucson, AR, USA)
- Gouws LM, Osmond CB, Schurr U, Walter A (2005) Distinctive diel growth cycles in leaves and cladodes of CAM plants: differences from C<sub>3</sub> plants and putative interactions with substrate availability, turgor and cytoplasmic pH. *Functional Plant Biology* **32**, 421–428. doi:10.1071/FP05074
- Griffiths HG, Smith JAC (1983) Photosynthetic pathways in the Bromeliaceae of Trinidad: relations between life-forms, habitat preference and occurrence of CAM. *Oecologia* **60**, 176–184.
- Haider MS, Barnes JD, Cushman JC, Borland AM (2012) A CAM- and starch-deficient mutant of the facultative CAM species *Mesembryanthemum crystallinum* reconciles sink demands by repartitioning carbon during acclimation to salinity. *Journal of Experimental Botany* **63**, 1985–1996. doi:10.1093/jxb/err412
- Holtum JAM, Winter K (1999) Degrees of crassulacean acid metabolism in tropical epiphytic and lithophytic ferns. *Australian Journal of Plant Physiology* **26**, 749–757. doi:10.1071/PP99001
- Holtum JAM, Aranda J, Virgo A, Winter K (2004) δ<sup>13</sup>C values and crassulacean acid metabolism in *Clusia* species from Panama. *Trees* **18**, 658–668. doi:10.1007/s00468-004-0342-y
- Holtum JAM, Chambers D, Morgan T, Tan DY (2011) *Agave* as a biofuel feedstock in Australia. *Global Change Biology - Bioenergy* **3**, 58–67. doi:10.1111/j.1757-1707.2010.01083.x
- Kaplan A, Gale J, Poljakoff-Mayber A (1976) Resolution of net dark fixation of carbon dioxide into its respiration and gross fixation components in *Bryophyllum daigremontianum*. *Journal of Experimental Botany* **27**, 220–230. doi:10.1093/jxb/27.2.220
- Kluge M, Ting IP (1978) 'Crassulacean acid metabolism. Analysis of an ecological adaptation.' (Springer-Verlag: Berlin)
- Königer M, Winter K (1993) Growth and photosynthesis of *Gossypium hirsutum* L. at high photon flux densities: effects of soil temperatures and nocturnal air temperatures. *Agronomie* **13**, 423–431. doi:10.1051/agro:19930507
- Medina E, Osmond CB (1981) Temperature dependence of dark CO<sub>2</sub> fixation and acid accumulation in *Kalanchoe daigremontiana*. *Australian Journal of Plant Physiology* **8**, 641–649.
- Milburn TR, Pearson DJ, Ndegwe NA (1968) Crassulacean acid metabolism under natural tropical conditions. *New Phytologist* **67**, 883–897. doi:10.1111/j.1469-8137.1968.tb06401.x
- Neales TF (1973a) The effect of night temperature on CO<sub>2</sub> assimilation, transpiration, and water use efficiency in *Agave americana* L. *Australian Journal of Biological Sciences* **26**, 705–714.
- Neales TF (1973b) The effect of night temperature on the assimilation of carbon dioxide by mature pineapple plants, *Ananas comosus* (L.) Merr. *Australian Journal of Biological Sciences* **26**, 539–546.
- Neales TF, Sale PJM, Meyer CP (1980) Carbon dioxide assimilation by pineapple plants, *Ananas comosus* (L.) Merr. Effects of variation of the day/night temperature regime. *Australian Journal of Plant Physiology* **7**, 375–385. doi:10.1071/PP9800375
- Nobel PS (1984) Productivity of *Agave deserti*: measurement by dry weight and monthly prediction using physiological responses to environmental parameters. *Oecologia* **64**, 1–7. doi:10.1007/BF00377535
- Nobel PS (1985) PAR, water, and temperature limitations on the productivity of cultivated *Agave fourcroydes* (henequen). *Journal of Applied Ecology* **22**, 157–173. doi:10.2307/2403334
- Nobel PS (1988) 'Environmental biology of agaves and cacti.' (Cambridge University Press: Cambridge)
- Nobel PS (1989) A nutrient index quantifying productivity of agaves and cacti. *Journal of Applied Ecology* **26**, 635–645. doi:10.2307/2404088
- Nobel PS (1991) Environmental productivity indices and productivity for *Opuntia ficus-indica* under current and elevated atmospheric CO<sub>2</sub> levels. *Plant, Cell & Environment* **14**, 637–646. doi:10.1111/j.1365-3040.1991.tb01536.x
- Nobel PS, Hartsocck TL (1978) Resistance analysis of nocturnal carbon dioxide uptake by a crassulacean acid metabolism plant, *Agave deserti*. *Plant Physiology* **61**, 510–514. doi:10.1104/pp.61.4.510
- Nobel PS, Hartsocck TL (1981) Shifts in the optimal temperature for nocturnal CO<sub>2</sub> uptake caused by changes in growth temperature for

- cacti and agaves. *Physiologia Plantarum* **53**, 523–527. doi:10.1111/j.1399-3054.1981.tb02744.x
- Nobel PS, McDaniell RG (1988) Low-temperature tolerances, nocturnal acid accumulation, and biomass increases for seven species of *Agave*. *Journal of Arid Environments* **15**, 147–155.
- Nobel PS, Meyer SE (1985) Field productivity of a CAM plant, *Agave salmiana*, estimated using daily acidity changes under various environmental conditions. *Physiologia Plantarum* **65**, 397–404. doi:10.1111/j.1399-3054.1985.tb08663.x
- Nobel PS, Quero E (1986) Environmental productivity indices for a Chihuahuan Desert CAM plant, *Agave lechuguilla*. *Ecology* **67**, 1–11. doi:10.2307/1938497
- Nobel PS, Valenzuela AG (1987) Environmental responses and productivity of the CAM plant, *Agave tequilana*. *Agricultural and Forest Meteorology* **39**, 319–334. doi:10.1016/0168-1923(87)90024-4
- Osmond CB (1978) Crassulacean acid metabolism: a curiosity in context. *Annual Review of Plant Physiology* **29**, 379–414. doi:10.1146/annurev.pp.29.060178.002115
- Owen NA, Griffiths H (2013) Marginal land bioethanol yield potential of four crassulacean acid metabolism candidates (*Agave fourcroydes*, *Agave salmiana*, *Agave tequilana* and *Opuntia ficus-indica*) in Australia. *Global Change Biology - Bioenergy* doi:10.1111/gcbb.12094
- Pimienta-Barrios E, Robles-Murguía C, Nobel PS (2001) Net CO<sub>2</sub> uptake for *Agave tequilana* in a warm and a temperate environment. *Biotropica* **33**, 312–318.
- Queiroz O (1966) Sur le métabolisme acide des Crassulacées.II. Action à long terme de la température de jour sur les variations de la teneur en acide malique en jours courts. *Physiologie Vegetale* **4**, 323–339.
- Silvera K, Santiago LS, Winter K (2005) Distribution of crassulacean acid metabolism in orchids of Panama: evidence of selection for weak and strong modes. *Functional Plant Biology* **32**, 397–407. doi:10.1071/FP04179
- Szarek SR, Ting IP (1974) Seasonal patterns of acid metabolism and gas exchange in *Opuntia basilaris*. *Plant Physiology* **54**, 76–81. doi:10.1104/pp.54.1.76
- Winter K (1985) Crassulacean acid metabolism. In 'Photosynthetic mechanisms and the environment'. (Eds J Barber, NR Baker) pp. 329–387. (Elsevier: Amsterdam)
- Winter K, Holtum JAM (2002) How closely do the  $\delta^{13}\text{C}$  values of crassulacean acid metabolism plants reflect the proportion of CO<sub>2</sub> fixed during day and night? *Plant Physiology* **129**, 1843–1851. doi:10.1104/pp.002915
- Winter K, Holtum JAM (2014) Facultative crassulacean acid metabolism (CAM) plants: powerful tools for unravelling the functional elements of CAM photosynthesis. *Journal of Experimental Botany* doi:10.1093/jxb/eru063
- Winter K, Aranda J, Holtum JAM (2005) Carbon isotope composition and water use efficiency in plants with crassulacean acid metabolism. *Functional Plant Biology* **32**, 381–388. doi:10.1071/FP04123
- Winter K, García M, Holtum JAM (2014) Nocturnal versus diurnal CO<sub>2</sub> uptake: how flexible is *Agave angustifolia*? *Journal of Experimental Botany* doi:10.1093/jxb/eru097
- Wong SC, Hew CS (1976) Diffusive resistance, titratable acidity and CO<sub>2</sub> fixation in two tropical epiphytic ferns. *American Fern Journal* **66**, 121–124. doi:10.2307/1546463
- Yamori W, Hikosaka K, Way DA (2014) Temperature response of photosynthesis in C<sub>3</sub>, C<sub>4</sub>, and CAM plants: temperature acclimation and temperature adaptation. *Photosynthesis Research* **119**, 101–117. doi:10.1007/s11120-013-9874-6
- Zhu J, Goldstein G, Bartholomew DP (1999) Gas exchange and carbon isotope composition of *Ananas comosus* in response to elevated CO<sub>2</sub> and temperature. *Plant, Cell & Environment* **22**, 999–1007. doi:10.1046/j.1365-3040.1999.00451.x
- Zotz G, Winter K (1994a) Annual carbon balance and nitrogen-use efficiency in tropical C<sub>3</sub> and CAM epiphytes. *New Phytologist* **126**, 481–492. doi:10.1111/j.1469-8137.1994.tb04245.x
- Zotz G, Winter K (1994b) A one-year study on carbon, water and nutrient relationships in a tropical C<sub>3</sub>-CAM hemi-epiphyte, *Clusia uvitana* Pittier. *New Phytologist* **127**, 45–60. doi:10.1111/j.1469-8137.1994.tb04258.x