

Interannual variation in land-use intensity enhances grassland multidiversity

Eric Allan^{a,1}, Oliver Bossdorf^{a,b}, Carsten F. Dormann^c, Daniel Prati^a, Martin M. Gossner^{d,e}, Teja Tschardt^f, Nico Blüthgen^g, Michaela Bellach^h, Klaus Birkhoferⁱ, Steffen Boch^{a,j}, Stefan Böhm^{i,k}, Carmen Börschig^f, Antonis Chatzinotas^l, Sabina Christ^m, Rolf Danielⁿ, Tim Diekötter^o, Christiane Fischer^{n,p}, Thomas Friedl^q, Karin Glaser^l, Christine Hallmann^q, Ladislav Hodac^q, Norbert Hölzel^r, Kirsten Jung^k, Alexandra Maria Klein^s, Valentin H. Klaus^r, Till Kleinebecker^r, Jochen Krauss^h, Markus Lange^{d,t}, E. Kathryn Morris^{u,v}, Jörg Müller^w, Heiko Nackeⁿ, Esther Pašalić^d, Matthias C. Rillig^{v,x}, Christoph Rothenwöhrer^f, Peter Schall^y, Christoph Scherber^f, Waltraud Schulze^{z,aa}, Stephanie A. Socher^a, Juliane Steckel^h, Ingolf Steffan-Dewenter^h, Manfred Türke^{d,e}, Christiane N. Weiner^h, Michael Werner^h, Catrin Westphal^f, Volkmar Wolters^o, Tesfaye Wubet^{bb,cc}, Sonja Gockel^e, Martin Gorke^p, Andreas Hemp^{dd}, Swen C. Renner^{k,ee}, Ingo Schöning^t, Simone Pfeiffer^p, Birgitta König-Ries^{ff}, François Buscot^{bb,cc,gg}, Karl Eduard Linsenmair^h, Ernst-Detlef Schulze^t, Wolfgang W. Weisser^{d,e}, and Markus Fischer^{a,p,w}

^aInstitute of Plant Sciences, University of Bern, CH-3013 Bern, Switzerland; ^bPlant Evolutionary Ecology, University of Tübingen, D-72076 Tübingen, Germany; ^cBiometry and Environmental System Analysis, University of Freiburg, D-79106 Freiburg, Germany; ^dInstitute of Ecology, Friedrich-Schiller University Jena, D-07743 Jena, Germany; ^eTerrestrial Ecology Research Group, Department of Ecology and Ecosystem Management, Center for Food and Life Sciences Weihenstephan, Technische Universität München, D-85354 Freising, Germany; ^fAgroecology, Department of Crop Sciences, Georg-August University of Göttingen, D-37077 Göttingen, Germany; ^gEcological Networks, Biology, Technische Universität Darmstadt, D-64287 Darmstadt, Germany; ^hDepartment of Animal Ecology and Tropical Biology, Biocentre, University of Würzburg, D-97074 Würzburg, Germany; ⁱBiodiversity and Conservation Science, Lund University, SE-22362 Lund, Sweden; ^jBotanical Garden, University of Bern, CH-3013 Bern, Switzerland; ^kInstitute of Experimental Ecology, University of Ulm, D-89069 Ulm, Germany; ^lDepartment of Environmental Microbiology, Helmholtz Centre for Environmental Research-UFZ, D-04318 Leipzig, Germany; ^mDepartment of Diagnostics, Fraunhofer Institute for Cell Therapy and Immunology, D-04103 Leipzig, Germany; ⁿDepartment of Genomic and Applied Microbiology and Göttingen Genomics Laboratory, Institute of Microbiology and Genetics, Georg-August University of Göttingen, D-37077 Göttingen, Germany; ^oDepartment of Animal Ecology, Justus Liebig University, D-35392 Giessen, Germany; ^pSenckenberg Gesellschaft für Naturforschung, Biodiversity and Climate Research Centre, D-60325 Frankfurt, Germany; ^qAlbrecht-von-Haller Institute for Plant Sciences, Department of Experimental Phycology and Culture Collection of Algae in Göttingen, Georg-August University of Göttingen, D-37073 Göttingen, Germany; ^rInstitute of Landscape Ecology, University of Münster, D-48149 Münster, Germany; ^sEcosystem Functions, Institute of Ecology, Leuphana University of Lüneburg, D-21335 Lüneburg, Germany; ^tMax Planck Institute for Biogeochemistry, D-07745 Jena, Germany; ^uDepartment of Biology, Xavier University, Cincinnati, OH 45207; ^vPlant Ecology, Institut für Biologie, Freie Universität Berlin, D-14195 Berlin, Germany; ^wBiodiversity Research/Systematic Botany, University of Potsdam, D-14469 Potsdam, Germany; ^xBerlin-Brandenburg Institute of Advanced Biodiversity Research, D-14195 Berlin, Germany; ^yDepartment of Silviculture and Forest Ecology of the Temperate Zones, Georg-August University of Göttingen, D-37077 Göttingen, Germany; ^zMax Planck Institute for Molecular Plant Physiology, D-14476 Potsdam, Germany; ^{aa}Department of Plant Systems Biology, University of Hohenheim, D-70593 Stuttgart, Germany; ^{bb}Department of Soil Ecology, UFZ-Helmholtz Centre for Environmental Research, D-06120 Halle (Saale), Germany; ^{cc}German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, D-04103 Leipzig, Germany; ^{dd}Department of Plant Systematics, University of Bayreuth, D-95445 Bayreuth, Germany; ^{ee}Smithsonian Conservation Biology Institute, National Zoological Park, Front Royal, VA 22630; ^{ff}University of Jena, D-07743 Jena, Germany; and ^{gg}Institute of Biology, Chair of Soil Ecology, University of Leipzig, D-04103 Leipzig, Germany

Edited by Osvaldo E. Sala, Arizona State University, Tempe, AZ, and accepted by the Editorial Board November 20, 2013 (received for review July 2, 2013)

Although temporal heterogeneity is a well-accepted driver of biodiversity, effects of interannual variation in land-use intensity (LUI) have not been addressed yet. Additionally, responses to land use can differ greatly among different organisms; therefore, overall effects of land-use on total local biodiversity are hardly known. To test for effects of LUI (quantified as the combined intensity of fertilization, grazing, and mowing) and interannual variation in LUI (SD in LUI across time), we introduce a unique measure of whole-ecosystem biodiversity, multidiversity. This synthesizes individual diversity measures across up to 49 taxonomic groups of plants, animals, fungi, and bacteria from 150 grasslands. Multidiversity declined with increasing LUI among grasslands, particularly for rarer species and aboveground organisms, whereas common species and belowground groups were less sensitive. However, a high level of interannual variation in LUI increased overall multidiversity at low LUI and was even more beneficial for rarer species because it slowed the rate at which the multidiversity of rare species declined with increasing LUI. In more intensively managed grasslands, the diversity of rarer species was, on average, 18% of the maximum diversity across all grasslands when LUI was static over time but increased to 31% of the maximum when LUI changed maximally over time. In addition to decreasing overall LUI, we suggest varying LUI across years as a complementary strategy to promote biodiversity conservation.

biodiversity loss | agricultural grasslands | Biodiversity Exploratories

Ample theoretical and empirical work has shown that temporal heterogeneity can promote biodiversity by creating niches that allow species with different responses to the environment to coexist stably (1, 2). Among the processes currently eroding

Significance

Land-use intensification is a major threat to biodiversity. So far, however, studies on biodiversity impacts of land-use intensity (LUI) have been limited to a single or few groups of organisms and have not considered temporal variation in LUI. Therefore, we examined total ecosystem biodiversity in grasslands varying in LUI with a newly developed index called multidiversity, which integrates the species richness of 49 different organism groups ranging from bacteria to birds. Multidiversity declined strongly with increasing LUI, but changing LUI across years increased multidiversity, particularly of rarer species. We conclude that encouraging farmers to change the intensity of their land use over time could be an important strategy to maintain high biodiversity in grasslands.

Author contributions: E.A., O.B., and M.F. designed research; D.P., M.M.G., T.T., N.B., M.B., K.B., S. Boch, S. Böhm, C.B., A.C., S.C., R.D., T.D., C.F., T.F., K.G., C.H., L.H., N.H., K.J., A.M.K., V.H.K., T.K., J.K., M.L., E.K.M., J.M., H.N., E.P., M.C.R., C.R., P.S., C.S., W.S., S.A.S., J.S., I.S.-D., M.T., C.N.W., M.W., C.W., V.W., T.W., S.C.R., F.B., W.W.W., and M.F. contributed data; E.A. analyzed data; E.A., O.B., C.F.D., D.P., M.M.G., T.T., N.B., and M.F. wrote the paper; F.B., K.E.L., E.-D.S., W.W.W., and M.F. designed and established the Biodiversity Exploratories; and S.G., M.G., A.H., S.C.R., I.S., S.P., and B.K.-R. maintained the infrastructure of the Biodiversity Exploratories.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. O.E.S. is a guest editor invited by the Editorial Board.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: eric.allan@ips.unibe.ch.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1312213111/-DCSupplemental.

biodiversity, land-use intensification is one of the most important (3–5), with likely feedbacks on ecosystem functioning (6). However, previous studies of land-use effects have only considered how changes in mean intensity affect biodiversity (7–9) and have neglected the question of whether interannual variation in land-use intensity (LUI) could also have an impact on biodiversity. In grasslands, farmers frequently change animal stocking densities, fertilizer application, or mowing frequencies across years (10), meaning that temporal heterogeneity in LUI can be high. Spatial heterogeneity of land use can promote biodiversity (11, 12), and we might hypothesize that interannual variation in LUI also has positive effects on biodiversity (“land-use variation hypothesis”). Changing land use across years could even mitigate some of the negative effects of high management intensity, and therefore help to develop strategies that resolve the tradeoff between high agricultural production and biodiversity conservation.

Interannual variation in LUI might generally promote biodiversity, but effects could differ between rarer and more common species or between aboveground and belowground groups. Locally abundant (common) species are often generalists and less sensitive to land use (13), whereas less abundant (rarer) species may be more sensitive to LUI because they have more specialized habitat requirements or smaller populations (14, 15). Although linkages between aboveground and belowground diversity are expected (16, 17), in grasslands, LUI and interannual variation in intensity may have more direct effects on aboveground diversity, whereas belowground diversity may be more sensitive to general soil conditions (18).

One of the main challenges when studying land-use effects on biodiversity is that responses of different taxonomic groups of organisms can differ greatly (4), making it difficult to assess overall land-use effects. One promising approach would be a synthetic index of total ecosystem biodiversity that integrates information on a wide diversity of groups of organisms and allows us to identify the conditions that simultaneously maximize the diversity of most groups. Here, we introduce and apply an index of “multidiversity,” which computes the average scaled species richness per taxonomic group. Species richness values for each group were scaled to the maximum value observed for that group across all of the grasslands, so that groups differing in the total number of species were weighted equally. The approach is conceptually similar to indices of multifunctionality used in biodiversity-ecosystem functioning research (6, 19) and to the World Wildlife Fund’s living planet index, which quantifies the overall state of biodiversity at the global scale (20). We apply the multidiversity index to a large set of biodiversity data from 150 grasslands to examine how changes in the mean and interannual variance of land use affect multidiversity and to what degree land-use effects differ between aboveground and belowground, as well as between locally rare vs. common, organisms.

We addressed these questions with the unique set of comprehensive biodiversity and land-use data of our German Biodiversity Exploratories project (21). We investigated land-use effects, first using a set of 150 plots on which the species richness of 18 taxonomic groups were measured and then using a subset of 27 plots on which 49 taxonomic groups were surveyed (Fig. 1). We modeled the response of multidiversity to an integrated measure of LUI (10), which is the sum of the standardized intensities of grazing (duration and type of grazing animals), mowing (number of cuts per year), and fertilization [kilograms of nitrogen (N) per hectare]. The shape of the relationship between LUI and biodiversity has important management implications; for instance, if the relationship is saturating rather than linear, this would suggest that large losses of biodiversity occur even at modest levels of intensification (9). We therefore fitted a series of models (Table S1) that differed in the shape of the relationship




Fig. 1. Organism groups used to calculate multidiversity. Of 49 taxonomic groups surveyed on 27 grasslands, 45 are shown on the tree. Eighteen groups were measured on all 150 grasslands, and 16 of them are shown in green on the tree. The groups WS3 and TM7 are candidate bacterial phyla. In addition to the groups shown on the tree, our analyses included lichens (150 plots), Homoptera (now considered paraphyletic but used as a group for convenience, 150 plots), viruses (27 plots), and a fungal subphylum *incertae sedis* (27 plots), which could not be placed on the tree. The tree was created based on National Center for Biotechnology Information taxonomy (www.ncbi.nlm.nih.gov), and therefore shows relationships among groups but without true branch lengths.

between LUI and multidiversity, whether they contained interannual variation in LUI [SD in LUI (LUI_{sd})] and whether they modeled LUI as an integrated index or fitted individual land-use components separately. We used Akaike’s information criterion corrected for small sample size (AICc) to select the best-fitting models (95% confidence set; *Methods*).

Results and Discussion

Our analysis of the species richness of 18 taxa across 150 grasslands showed a clear negative response of multidiversity to increasing LUI (Fig. 24). Multidiversity followed a negative asymptotic exponential relationship, which indicates that intensification of land use will have particularly negative consequences for biodiversity in extensively managed grasslands (9). The asymptote of the curve was 0.3, indicating that species richness declined to 30% of the maximum across taxonomic groups. Although there were many “losers” (13) under land-use intensification, some (mostly animal) groups were hardly affected: Diptera, arbuscular mycorrhizal fungi, and bats did not decline with increasing LUI (Fig. 3). In contrast, plants and lichens, as well as Orthoptera, Araneae, and Lepidoptera, all declined strongly. In general, animal groups showed a wider range of responses to LUI than did plant groups; however, calculating multidiversity for plants and animals separately revealed that the overall response was the same for both groups (all best models were asymptotic exponentials), although LUI explained less of the variation in animal multidiversity (pseudo- R^2 of 0.2 for animals and 0.7 for plants; Fig. S1 A and B and Table S2) because of the more variable responses of the animals. Modeling




Fig. 2. Effect of LUI on multidiversity of 18 taxonomic groups of all species (A), relatively rare species (i.e., the 90% of species with the lowest total abundance) (B), and relatively common species (i.e., the 10% of species with the highest total abundance across plots) (C). Lines show model fits for different values of LUI_{sd} , from $LUI_{sd} = 0$ to the maximum interannual land-use change observed in any plot, $LUI_{sd} = 1$. In all cases, model predictions were calculated using multimodel averaging across all models in the 95% confidence set and were averaged across regions (more details are provided in *Methods*). In C, not all models in the 95% confidence set could be used for multimodel averaging because it would not be possible to average across different types of models (e.g., those with the compound LUI and those with individual components). AICc weights of models that could be used for multimodel averaging summed to 78%.

LUI using the integrated index proved better than modeling it using the individual intensities of grazing, mowing, and fertilization. Of the individual components, however, grazing and mowing seemed to be more important than fertilization in driving declines in multidiversity (*SI Methods* and *Fig. S2*). High rates of fertilization mostly occur in frequently mown grasslands (10), but these results suggest that fertilization may not have as many negative effects as high mowing intensity. As with all nonexperimental studies, it is impossible to identify LUI categorically as the driver of these differences in multidiversity among grasslands; however, great care was taken during plot selection to

minimize confounding between environmental variables and LUI (21). Loss of multidiversity could be driven by effects on abundance, or there could be effects on species richness per se. Land-use intensification might cause a reduction in abundance, and this, in turn, could cause a loss of species richness. Alternatively, higher LUI could reduce species richness more than would be expected based on changing abundance, which would be an effect on species richness per se. Further analyses (*SI Methods* and *Figs. S3* and *S4*) suggest that both processes are important: The relative importance varies among groups, but the overall effect on multidiversity seems principally driven by an effect on species richness per se. LUI also reduced the evenness of species abundances, but the effects on evenness were less pronounced than those on species richness (*Fig. S1C* and *Table S2*), supporting the idea that richness and evenness may show different responses (22) and that LUI has smaller effects on abundance and principally reduces species numbers.

Interannual changes in LUI were beneficial for biodiversity (*Fig. 2A*), supporting our land-use variation hypothesis. In the analysis of 18 groups of organisms in 150 grasslands, all of the




Fig. 3. Response to LUI of scaled species richness for the 18 taxonomic groups measured on 150 plots. For each group, the predictions from the best-fit model are shown (*Methods* and *Table S2*). Where temporal variation in LUI (LUI_{sd}) appeared in the model, model fits are shown for different values of LUI_{sd} . (A–R) The groups are sorted in order of their response to LUI, from the group showing the strongest decline (A) to the strongest increase (R). For each group, changes in species richness with increasing LUI are shown beside the name of the group on the graph and were calculated as the difference between the relative species richness predicted by the model at minimum LUI ($LUI = 0.5$) and the predicted species richness at maximum LUI ($LUI = 3.5$). Model predictions were evaluated at the mean LUI_{sd} and were averaged across regions. AMF, arbuscular mycorrhizal fungi.

pattern because individual responses differed in magnitude and shape (Fig. 3). Earlier, taxonomically less comprehensive studies have also reported idiosyncratic land-use responses of different groups of organisms (28, 29). However, our study, which covers a large number of groups of organisms, including taxa not normally assessed in environmental monitoring, shows that there is a consistent pattern in a comprehensive measure of the biodiversity of the ecosystem. The multidiversity index could facilitate efforts to identify areas of high conservation priority or to assess the effectiveness of restoration efforts. In our analysis, we assumed that all of the taxonomic groups were of equal importance. However, future uses of the index could consider weighting the groups differently. For instance, groups could be weighted by their conservation relevance, by their importance in providing certain ecosystem services, by their phylogenetic distinctiveness, or by aesthetic/cultural value.

Using the newly developed multidiversity index and our uniquely comprehensive grassland biodiversity data, we not only provide strong support for the importance of extensively managed grasslands for nature conservation (30) but, importantly, show that increased interannual variation in LUI maintains higher biodiversity and slows the rate at which rarer species are lost with increasing LUI. Varying LUI across time might also promote ecosystem service delivery if higher plant diversity increases forage production (31) or higher pollinator diversity promotes pollination of surrounding crops (32). One way of ameliorating the adverse effect of land-use intensification on biodiversity could therefore be to encourage farmers to alter the intensity of their land use somewhat across years. This could contribute to reconciling the need to produce high levels of biomass in grasslands with the maintenance of biodiversity.

Methods

Study Design. The study grasslands are located in three regions in Germany and are part of the Biodiversity Exploratories project (www.biodiversity-exploratories.de). The study regions are (i) the United Nations Educational, Scientific, and Cultural Organization (UNESCO) Biosphere Area of Schwäbische Alb in southwestern Germany, (ii) the National Park of Hainich-Dün and the surrounding area in central Germany, and (iii) the UNESCO Biosphere Reserve of Schorfheide-Chorin in northeastern Germany. The three regions differ in climate, geology, and topography and are representative of large parts of Central Europe, spanning a range of almost 3 °C in mean annual temperature and 500–1,000 mm of precipitation [details are provided by Fischer et al. (21)]. Grasslands in all three regions span a similar gradient in land-use intensity (LUI) (21). Because natural grasslands, those not requiring management to prevent succession to forest, are almost absent from Western and Central Europe, the land-use gradient is from seminatural to intensively managed grasslands. In each region, 50 permanent grassland plots (50 × 50 m) were established (150 in total) along a gradient of increasing LUI. A smaller number of plots, spanning the same range in LUI, on which more labor-intensive measurements could be carried out were also established: There were nine of these in each exploratory region (27 in total). All of the plots had been grassland for at least 20 y before the start of the project.

Land Use. Land use in these grasslands comprises fertilization, mowing, and grazing at different intensities. Land use was quantified based on a questionnaire submitted to farmers and landowners each year from 2006 to 2008 (10, 21). Grasslands could be grazed by cattle, horses, or sheep, and farmers reported the number of animals and the duration of grazing in each plot. Farmers were also asked about the number of mowing events per year (from one to three cuts) and the amount of N in fertilizer (organic and inorganic) added to the grassland.

Land use was quantified using a compound index of LUI (10), which does not suffer information loss due to categorization and makes different management types comparable. LUI integrates the intensity of fertilization (F), the mowing frequency (M), and the intensity of grazing (G) for each grassland plot. Grazing livestock were translated into livestock units weighted for their impact on grasslands (21). For each plot, an individual LUI component (F , M , or G) was standardized relative to its mean across all three regions and across all 3 y (details are provided in *SI Methods*). The compound

LUI is the sum of the three standardized components. The minimum LUI of 0.5 could be produced by mowing every 2 y, fertilizing at the rate of 6 kg of $N\cdot ha^{-1}\cdot y^{-1}$, or grazing one cow (>2 y old) per hectare for 30 d (or one sheep per hectare for the whole year). An intermediate LUI of 1.5 would equate to around two cuts per year, the addition of 60 kg of $N\cdot ha^{-1}\cdot y^{-1}$, or grazing one cow per hectare for most of the year (300 d). A high LUI of 3.0 could be produced by grazing three cows per hectare for most of the year (300 d) and fertilizing at the rate of 50 kg of $N\cdot ha^{-1}\cdot y^{-1}$ or by cutting three times and fertilizing with 130 kg of $N\cdot ha^{-1}\cdot y^{-1}$. For the analyses here, we used the average LUI across 3 y and the LUI_{sd} across 3 y (2006–2008). In addition to using the LUI (i.e., where all three types of land use are given equal weight), we tested the individual standardized land use components in our models to determine whether certain types of land use had a larger effect on biodiversity.

The intensity of land use in the grasslands changed considerably over time (10). We quantified this using the LUI_{sd} across the 3 y. This LUI_{sd} was uncorrelated with the mean LUI across the 3 y (Fig. S5). Because most of the data were collected in 2008 or 2009 (Table S4), LUI_{sd} calculates the change in land use in the years preceding data collection. We also calculated the SD in mowing, grazing, and fertilization intensity across the 3 y (*SI Methods*).

Species Richness Data. Data on the species richness of 18 taxonomic groups were collected with different standardized sampling methods on the 150 plots (Table S4). In some cases, more labor-intensive methods were used to sample the same groups on the subset of 27 plots; however, the intensity of sampling did not affect the results (Fig. S6A). Note that we use the term “species richness” throughout, although for the microbial and fungal groups, these are phylotypes and not necessarily true species.

Calculation of Multidiversity. We calculated multidiversity as the average proportional species richness across taxonomic groups. Species richness values were standardized for each taxonomic group by scaling them to the maximum observed value across all grasslands. Note that we could not simply sum species richness values to calculate multidiversity because this would have given higher weighting to species-rich groups. For instance, the bacterial groups had phylotype richness values of several thousand. However, we also conducted a range of sensitivity analyses to test other ways of calculating multidiversity (*SI Methods*). We used a different standardizations of species richness, we used a range of thresholds, and we calculated multidiversity using alternative taxonomic groupings (Fig. S6). The code used to calculate multidiversity is available at <https://github.com/eric-allan/multidiversity>.

We calculated multidiversity values for all 150 plots using 18 groups, and we also calculated multidiversity for the six plant groups (including lichens) and 11 animal groups separately. Furthermore, we calculated multidiversity based on Pielou's evenness index (J) rather than based on species richness. On the subset of 27 plots, we calculated multidiversity for the 17 above-ground groups and the 33 below-ground groups separately.

Additionally, we calculated multidiversity on all 150 plots for common and rare species separately. For each species, we calculated its total abundance across plots. Within each of 17 groups (we did not have data on the occurrence of each mycorrhizal fungal phylotype in each plot, so they were excluded from this analysis), we split the species into two categories: Common species were the top 10% in terms of total abundance, and the species we refer to as “rare” were the bottom 90% of species. Species abundances followed approximately lognormal distributions, so this split ensured that only abundant species were counted as common. Classifying rare species as the least abundant 50% of species or defining rare and common species separately for each region gave similar results (*SI Methods* and Table S3). Using a threshold in this way means that the species we classified as rare were only relatively rare (i.e., across the study plots) and not necessarily generally rare in the landscape. We analyzed multidiversity for the 17 groups of rare species and the 17 groups of common species separately.

Statistical Analysis. We first analyzed the response to LUI for each of the 18 taxonomic groups measured on all plots; all analyses were conducted with R version 2.15 (33). We used an approach similar to that of Scherber et al. (16) and scaled the species richness of each group between 0 and 1. We then fitted a series of models to estimate the shape of the response of each taxonomic group to LUI. We fitted “region” in all models to account for regional differences in species richness. We tested polynomial models with linear, quadratic, or cubic terms for LUI. These models test for a linear change in species richness with land use, a unimodal relationship, or a relationship with two turning points. We also used nonlinear regressions [fitted with the `gnls` function in `nlme` (34)] and tested for three further shapes of response: (i) negative exponential models, which model an

exponential decay of species richness with increasing LUI, asymptoting at 0; (ii) asymptotic exponential models in which species richness can asymptote at values greater than 0; and (iii) power law models, which allow a diversity of shapes of response to be modeled. In the nonlinear regressions, we modeled the intercept of each model as a function of region, which is therefore equivalent to fitting region as a categorical factor in the polynomial regressions. We also fitted a null model with only a main effect for region.

To model the influence of temporal changes in LUI, we fitted the same models as above but with covariates for LUI_{sd} . For the linear regressions (linear, quadratic, or cubic), we either fitted a main effect only for LUI_{sd} or an interaction between LUI_{sd} and all other parameters. For the nonlinear regressions, we modeled all of the possible combinations of each of the individual parameters as a function of LUI_{sd} . This resulted in 25 different models (Table S1). For each taxonomic group, the best-fit model was selected using AICc (35).

To analyze the shape of the relationship between LUI and multidiversity, we used the same series of models as above; in addition, we tested models with each of the three land-use components (fertilization, grazing, and mowing) fitted individually. We did not use models with individual land-use components for the analysis of individual diversities because we wanted to be able to compare the response of the different taxonomic groups to the same measure of LUI. In the analysis of multidiversity, we fitted all possible combinations of linear or linear and quadratic terms for each land-use component, while obeying the principle of marginality (36). All models contained a main effect for region. We did not conduct nonlinear regressions for the different land-use components

because models could not include nonlinear terms for more than one land-use component at a time (all 76 models are listed in Table S1).

To model multidiversity on the subset of 27 plots with fewer degrees of freedom, we used a reduced set of models. We did not include linear models with interactions between LUI_{sd} and LUI or nonlinear models with more than one parameter modeled as a function of LUI_{sd} (67 models).

We calculated AICc weights for each model: These weight the explanatory power of each model relative to the others tested. For the analysis of multidiversity, we present the models that account for 95% of the AICc weights as the set of best models (95% confidence set). We also calculated parameters for the relationships shown in Figs. 2 and 4A with multimodel averaging (35), using the MuMIn package in R (37). This averages the parameters across all models in the 95% confidence set, weighing each value by the model's AICc weight. We further calculated the square of the Pearson correlation coefficient between observed and model fitted values (pseudo- R^2). Although this measure may not be appropriate for nonlinear models, it conveys an idea of the goodness of fit.

ACKNOWLEDGMENTS. We thank Fabrice Grassein and Madalin Parepa for comments on the manuscript and Jens Nieschulze and Michael Owonibi for maintenance of the database. This work was funded by the Deutsche Forschungsgemeinschaft Priority Program 1374 "Infrastructure-Biodiversity Exploratories." Fieldwork permits were given by the responsible state environmental offices of Baden-Württemberg, Thüringen, and Brandenburg (according to §72 Brandenburgisches Naturschutzgesetz).

- Adler PB, HilleRisLambers J, Kyriakidis PC, Guan Q, Levine JM (2006) Climate variability has a stabilizing effect on the coexistence of prairie grasses. *Proc Natl Acad Sci USA* 103(34):12793–12798.
- Chesson P (2000) Mechanisms of maintenance of species diversity. *Annu Rev Ecol Syst* 31:343–366.
- Foley JA, et al. (2005) Global consequences of land use. *Science* 309(5734):570–574.
- Flynn DFB, et al. (2009) Loss of functional diversity under land use intensification across multiple taxa. *Ecol Lett* 12(1):22–33.
- Laliberté E, et al. (2010) Land-use intensification reduces functional redundancy and response diversity in plant communities. *Ecol Lett* 13(1):76–86.
- Isbell F, et al. (2011) High plant diversity is needed to maintain ecosystem services. *Nature* 477(7363):199–202.
- Clough Y, et al. (2011) Combining high biodiversity with high yields in tropical agroforests. *Proc Natl Acad Sci USA* 108(20):8311–8316.
- Geiger F, et al. (2010) Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic Appl Ecol* 11(2):97–105.
- Kleijn D, et al. (2009) On the relationship between farmland biodiversity and land-use intensity in Europe. *Proc Biol Sci* 276(1658):903–909.
- Blüthgen N, et al. (2012) A quantitative index of land-use intensity in grasslands: Integrating mowing, grazing and fertilization. *Basic Appl Ecol* 13(3):207–220.
- Benton TG, Vickery JA, Wilson JD (2003) Farmland biodiversity: Is habitat heterogeneity the key? *Trends Ecol Evol* 18(4):182–188.
- Fuhlendorf SD, Engle DM (2001) Restoring heterogeneity on rangelands: Ecosystem management based on evolutionary grazing patterns. *Bioscience* 51(8):625–632.
- McKinney ML, Lockwood JL (1999) Biotic homogenization: A few winners replacing many losers in the next mass extinction. *Trends Ecol Evol* 14(11):450–453.
- Fischer M, Stöcklin J (1997) Local extinctions of plants in remnants of extensively used calcareous grasslands 1950–1985. *Conserv Biol* 11(3):727–737.
- Lavergne S, Thuiller W, Molina J, Debussche M (2005) Environmental and human factors influencing rare plant local occurrence, extinction and persistence: A 115-year study in the Mediterranean region. *J Biogeogr* 32(5):799–811.
- Scherber C, et al. (2010) Bottom-up effects of plant diversity on multitrophic interactions in a biodiversity experiment. *Nature* 468(7323):553–556.
- De Deyn GB, Van der Putten WH (2005) Linking aboveground and belowground diversity. *Trends Ecol Evol* 20(11):625–633.
- Bardgett RD, Bowman WD, Kaufmann R, Schmidt SK (2005) A temporal approach to linking aboveground and belowground ecology. *Trends Ecol Evol* 20(11):634–641.
- Maestre FT, et al. (2012) Plant species richness and ecosystem multifunctionality in global drylands. *Science* 335(6065):214–218.
- Loh J, et al. (2005) The Living Planet Index: Using species population time series to track trends in biodiversity. *Philos Trans R Soc Lond B Biol Sci* 360(1454):289–295.
- Fischer M, et al. (2010) Implementing large-scale and long-term functional biodiversity research: The Biodiversity Exploratories. *Basic Appl Ecol* 11(6):473–485.
- Wilsey BJ, Chalcraft DR, Bowles CM, Willig MR (2005) Relationships among indices suggest that richness is an incomplete surrogate for grassland biodiversity. *Ecology* 86(5):1178–1184.
- Thies C, Steffan-Dewenter I, Tscharntke T (2008) Interannual landscape changes influence plant-herbivore-parasitoid interactions. *Agric Ecosyst Environ* 125(1–4):266–268.
- Isbell F, Tilman D, Polasky S, Binder S, Hawthorne P (2013) Low biodiversity state persists two decades after cessation of nutrient enrichment. *Ecol Lett* 16(4):454–460.
- Phalan B, Onial M, Balmford A, Green RE (2011) Reconciling food production and biodiversity conservation: Land sharing and land sparing compared. *Science* 333(6047):1289–1291.
- Tscharntke T, et al. (2012) Landscape moderation of biodiversity patterns and processes—Eight hypotheses. *Biol Rev Camb Philos Soc* 87(3):661–685.
- Rodrigues JLM, et al. (2013) Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. *Proc Natl Acad Sci USA* 110(3):988–993.
- Billetter R, et al. (2008) Indicators for biodiversity in agricultural landscapes: A pan-European study. *J Appl Ecol* 45(1):141–150.
- Dormann CF, et al. (2007) Effects of landscape structure and land-use intensity on similarity of plant and animal communities. *Glob Ecol Biogeogr* 16(6):774–787.
- Krauss J, et al. (2010) Habitat fragmentation causes immediate and time-delayed biodiversity loss at different trophic levels. *Ecol Lett* 13(5):597–605.
- Isbell F, et al. (2013) Nutrient enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *Proc Natl Acad Sci USA* 110(29):11911–11916.
- Kremen C, Williams NM, Thorp RW (2002) Crop pollination from native bees at risk from agricultural intensification. *Proc Natl Acad Sci USA* 99(26):16812–16816.
- R Core Team (2012) *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna).
- Pinheiro J, Bates D, DebRoy S, Sarkar D; R Development Core Team (2012) *lme4: Linear and Nonlinear Mixed Effects Models*.
- Burnham K, Anderson D (2002) *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach* (Springer, Berlin).
- Nelder JA (1977) A reformulation of linear models. *J R Stat Soc Ser A Stat Soc* 140(1):48–77.
- Bartoň K (2012) *MuMIn: Multi-Model Inference*. R Package Version 1.