



## Short Communication

# Field surveys in Western Panama indicate populations of *Atelopus varius* frogs are persisting in regions where *Batrachochytrium dendrobatidis* is now enzootic

<sup>1</sup>Rachel Perez, <sup>2,6</sup>Corinne L. Richards-Zawacki, <sup>3</sup>Alexander R. Krohn, <sup>2</sup>Matthew Robak, <sup>4</sup>Edgardo J. Griffith, <sup>4</sup>Heidi Ross, <sup>5</sup>Brian Gratwicke, <sup>6</sup>Roberto Ibáñez, and <sup>1,\*</sup>Jamie Voyles

<sup>1</sup>Department of Biology, New Mexico Institute of Mining and Technology, Socorro, New Mexico 87801, USA <sup>2</sup>Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, Louisiana 70118, USA <sup>3</sup>Department of Environmental Science Policy and Management, University of California, Berkeley, Berkeley, California 94720, USA <sup>4</sup>El Valle Amphibian Conservation Center, El Valle, PANAMÁ <sup>5</sup>Smithsonian Conservation Biology Institute, Washington D.C., USA <sup>6</sup>Smithsonian Tropical Research Institute, P.O. Box 0843-03092, Balboa, Ancón, PANAMÁ

**Key words.** Chytridiomycosis, Central America, Harlequin frog, golden frog, threatened species, emerging infectious disease, chytridiomycosis, *Batrachochytrium dendrobatidis*

**Citation:** Perez R, Richards-Zawacki CL, Krohn AR, Robak M, Griffith EJ, Ross H, Gratwicke B, Ibáñez R, Voyles J. 2014. Field surveys in Western Panama indicate populations of *Atelopus varius* frogs are persisting in regions where *Batrachochytrium dendrobatidis* is now enzootic. *Amphibian & Reptile Conservation* 8(2) [General Section]: 30–35 (e85).

**Copyright:** © 2014 Perez et al. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits unrestricted use for non-commercial and education purposes only, in any medium, provided the original author and the official and authorized publication sources are recognized and properly credited. The official and authorized publication credit sources, which will be duly enforced, are as follows: official journal title *Amphibian & Reptile Conservation*; official journal website <[amphibian-reptile-conservation.org](http://amphibian-reptile-conservation.org)>.

**Received:** 18 June 2014; **Accepted:** 25 August 2014; **Published:** 11 November 2014

The stunning Harlequin frogs of the genus *Atelopus*, once common, are now among the most imperiled of all amphibian species (La Marca et al. 2005; Zippel et al. 2006). Of 88 described *Atelopus* species in Central and South America, 65 (74%) are Critically Endangered (La Marca et al. 2005). The most pressing threat to these frogs is chytridiomycosis, a fungal disease caused by *Batrachochytrium dendrobatidis* (hereafter “*Bd*”) and associated with die-offs of amphibians around the world (Berger et al. 1998; Longcore et al. 1999; Kilpatrick et al. 2010). In Western Panama, an epidemic wave spreading from west to east caused mass mortality events, resulting in catastrophic losses in amphibian diversity (Lips et al. 2006; Brem and Lips 2008; Woodhams et al. 2008; Kilburn et al. 2010), including declines in three *Atelopus* species: *A. varius*, *A. zeteki*, and *A. chiriquiensis*.

Relatively few studies have been conducted on the amphibian communities of Western Panama following chytridiomycosis outbreaks. Those few investigations that have focused on understanding community composition where *Bd* is now enzootic report differential survival among host species (e.g., Brem and Lips 2008), with some species putatively driven to local extinction (Gagliardo et al. 2008). In particular, *Atelopus* species were considered to be highly vulnerable to disease-induced extinction for multiple reasons. First, *A. varius* was used as an “indicator species” to monitor declines


and thus helped to document *Bd* invasion and characterize *Bd*-related amphibian losses (Brem and Lips 2008). Second, *A. zeteki* has been repeatedly tested in controlled laboratory infection experiments and found to be highly susceptible to lethal chytridiomycosis (Bustamente et al. 2010; DiRenzo et al. 2014; Ellison et al. 2014). Third, recent immunogenetics research suggests that *A. zeteki* adaptive immune responses are suppressed by *Bd* (Ellison et al. 2014). Thus, *Atelopus* species have become important focal species in the study of chytridiomycosis dynamics and have also provided motivation for progressive conservation action (Gagliardo et al. 2008).

In October 2012, we revisited study sites that were surveyed for *Atelopus* from 2001 to 2004 (Richards and Knowles 2007) and we established new study sites in hopes of discovering extant populations of Critically Endangered *Atelopus* species. Using a measuring tape, a 200 m transect was developed and marked with flagging tape every 10 m. For our surveys, 2–3 observers walked these transects slowly, searching for amphibians according to established visual encounter survey protocols (e.g., Lips 1999). We captured all post-metamorphic amphibians we encountered using a fresh pair of gloves or inverted plastic bag to minimize transmission or infection and followed strict field hygiene protocols (Phillot et al. 2010). We noted the time and location of capture, identified the species, sex and age class, and measured


**Correspondence.** Email: [jamie.voyles@gmail.com](mailto:jamie.voyles@gmail.com) (Corresponding author, Jamie Voyles); tel: (720) 883-2341; fax: (208) 885-7905.

snout-to-vent length and body mass. We also collected skin swab samples for all amphibians using standardized swabbing techniques (Hyatt et al. 2007). We preserved the skin swab samples (by freezing at  $-20\text{ }^{\circ}\text{C}$ ) to test for *Bd* infection using quantitative polymerase chain reaction (qPCR; Boyle et al. 2005; Hyatt et al. 2007). For the qPCR assay, we analyzed all samples in triplicate with an internal positive control (Hyatt et al. 2007) and used a dilution set of plasmid standards (obtained from Pisces Molecular, Boulder, Colorado) to quantify pathogen load. We converted plasmid copy numbers to zoospore copy numbers using the line of best fit ( $r^2 > 0.999$ ) from a linear regression of  $\log(\text{plasmids})$  vs.  $\log(\text{zoospores})$  ( $t_4 = 210.6$ ,  $P < 0.0001$ ) that we obtained by running the plasmid standard set alongside a series of standards containing known quantities of zoospores (obtained from Alex Hyatt, Australian Animal Health Laboratory). If one of three replicate wells turned up positive, we checked Cycle Threshold (Ct) value to determine whether non-amplification in two of three wells could have been caused by a low-level infection (near the detection threshold) and verified that the qPCR was not inhibited (IPC amplified normally). In cases of inhibition or Ct values far from the detection threshold, we re-ran and considered them positive if *Bd* was detected in any of the three re-run wells.

We surveyed 16 field sites from 2012 to 2013, 10 of which were sites where *Atelopus* species were found in 2004 (Richards and Knowles 2007), prior to the chytridiomycosis epidemic. The remaining six sites were chosen based on other biologists' sightings of *Atelopus*



**Figure 1.** A female Harlequin frog, *Atelopus varius*. This species, classified as Critically Endangered by IUCN, has been found in small numbers in the mountains of Western Panama.



**Figure 2.** A pair of *Atelopus varius* in amplexus, found in the mountains of Western Panama.

## *Atelopus varius* populations persisting after chytridiomycosis

**Table 1.** Summary of infection prevalence of *Batrachochytrium dendrobatidis* from amphibians found at three sites where *Atelopus varius* still persist.

Site (Elevation)	Species	Prevalence	N	Lower 95%	Upper 95%
Highland 1 (735 m)	<i>Atelopus varius</i>	0%	0/1	0	0.975
	<i>Colostethus flotator</i>	0%	0/1	0	0.975
	<i>Craugastor bransfordii</i>	0%	0/1	0	0.975
	<i>Craugastor crassidigitus</i>	0%	0/1	0	0.975
	<i>Craugastor fitzingeri</i>	0%	0/1	0	0.975
	<i>Lithobates warszewitschii</i>	60%	3/5	0.147	0.947
	<i>Pristimantis cerasinus</i>	100%	1/1	0.025	1
	<i>Pristimantis cruentus</i>	0%	0/2	0	0.842
	<i>Pristimantis museosus</i>	0%	0/1	0	0.975
	<i>Pristimantis ridens</i>	0%	0/1	0	0.975
	<i>Sachatamia albomaculata</i>	25%	2/8	0.0715	0.591
	<i>Terahyla spinosa</i>	0%	0/1	0	0.975
	Glass frog metamorph	0%	0/1	0	0.975
Highland 2 (521 m)	<i>Atelopus varius</i>	0%	0/6	0	0.459
	<i>Colostethus flotator</i>	0%	0/7	0	0.41
	<i>Colostethus panamensis</i>	25%	1/4	0.073	0.524
	<i>Pristimantis ridens</i>	0/1	0/1	0	0.975
	<i>Rhaebo haemitticus</i>	42%	1/5	0.057	0.437
	<i>Silverstoneia flotator</i>	33%	1/3	0.008	0.906
	<i>Smilisca</i> spp.	100%	3/3	0.292	1
Lowland 1 (0 m)	<i>Atelopus varius</i>	0%	0/1	0	0.975
	<i>Craugastor bransfordii</i>	0%	0/2	0	0.842
	<i>Craugastor crassidigitus</i>	0%	0/1	0	0.975
	<i>Craugastor longirostris</i>	0%	0/1	0	0.975
	<i>Craugastor fitzingeri</i>	0%	0/2	0	0.842
	<i>Dendrobates auratus</i>	0%	0/3	0	0.708
	<i>Dendrobates minnutus</i>	0%	0/1	0	0.975
	<i>Diasperous</i> spp.	0%	0/3	0	0.708
	<i>Pristimantis caryophyllaceus</i>	0%	0/1	0	0.975
	<i>Rhinella alata</i>	0%	0/10	0	0.308
	<i>Silverstoneia flotator</i>	0%	0/1	0	0.975

(e.g., Hertz et al. 2012) or predicted habitat suitability in species distribution models. We found persisting populations of *A. varius* at three of 16 (18.7%) field sites (Fig. 1, Table 1). At one site, we found one juvenile *A. varius*, five adult males, and two adult females, including one pair in amplexus (Fig. 2). We found individual adult males at each of the other two respective sites (Table 1). We have intentionally only provided general site information, rather than precise site coordinates, due to the risk of illegal animal collections.

We confirmed that *Bd* is present in two of these three populations based on detection of *Bd* on skin swabs from other species (e.g., *Lithobates warszewitschii* and *Sachatamia albomaculata*), but none of the *Atelopus* samples were *Bd* positive on these transects (Table 1). These sites vary in elevation from 45 to 750 m and all three are in areas where fungal epidemics were associated with mas-

sive amphibian declines from 2004 to 2006 (Lips et al. 2006; Brem and Lips 2008; Kilburn et al. 2010). We did not detect *A. zeteki* or *A. chiriquiensis* at any of our study sites. Although survey efforts for these species are still underway, the absence of these species is concerning because we know that they previously had restricted ranges (Zippel et al. 2006) and at least *A. zeteki* is known to be highly susceptible to chytridiomycosis in laboratory infection experiments (Bustamente et al. 2010; DiRenzo et al. 2014; Ellison et al. 2014).

We found that *A. varius* is persisting in multiple sites following a chytridiomycosis outbreak in western Panama. Furthermore, our positive qPCR results suggest that these populations have survived despite the presence of *Bd*. Prior to this study, Hertz et al. (2012) was the only study to document sightings of *A. varius* in the wild in Panama since 2006. Those observations were made in

2009 at a site in Santa Fe National Park. These populations may be persisting for a wide range of biotic (e.g., changes in host behavioral, innate or acquired immune responses, anti-*Bd* microbial communities) or abiotic (e.g., environmental/thermal conditions) reasons. However, because there have been few coordinated efforts to locate new populations, resurvey historical localities, or test for *Bd* infections, the question of how these populations have persisted—and whether any other *Atelopus* populations have survived—remains to be unraveled. We believe that the lack of post-decline survey effort has not been so much an oversight, but likely a consequence of the enormity of the challenge of monitoring these species while simultaneously establishing conservation programs to abate the threat of chytridiomycosis to entire amphibian communities.

Post-decline surveys are critical for conservation of *Atelopus* species, as well as for other neotropical amphibians. Documenting rediscovered species is critically important for informing conservation and management initiatives (Minteer et al. 2014) and, in this case, could be accomplished with photographs, rather than collecting the individuals. Moreover, understanding the variables that permit some populations to persist while others die out will be critical to conservation, especially since several species are being bred in captivity (e.g., *A. varius* and *A. zeteki*) with the expectation of one day returning them to the wild (Gagliardo et al. 2008; Zippel et al. 2011). Our discovery of extant populations of *A. varius* in *Bd*-enzootic areas underscores the importance of continued monitoring for species presumed to be “extinct in the wild,” even after long periods without any sightings.

**Acknowledgments.**—Disney Worldwide Wildlife Fund, Association of Zoos and Aquariums (AZA) Conservation Endowment Fund, Riverbanks Conservation Support Fund, Bay and Paul Foundations, Minnesota Zoo Ulysses S. Seal Conservation Grant, Maryland Zoo Conservation Fund, the Chicago Board of Trade Endangered Species Fund, The AZA Amphibian Taxon Advisory Group, Stone Center for Latin American Studies, and Project Golden Frog funded this research. We thank the Smithsonian Tropical Research Institute for allowing access to their facilities. We thank Maggie Unkefer, Clifford Richardson, Kelly Cruz and the Cruz family, Daniel Medina, Karina Klonoski, and Simone Des Roches for their support. This work was conducted under IACUC NMT: 2013-1, IACUC Tulane: 0453, STRI IACUC: 2012-0901-2015, ANAM SE/AH-4-12, SE/AH-1-13, SE/AH-1-12, and SE/AH-4-13.

## Literature Cited

- Berger L, Speare R, Daszak P, Green DE, Cunningham AA, Goggin CL, Parkes H. 1998. Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Sciences of the United States of America* 95(15): 9031–9036.
- Brem FM, Lips KR. 2008. *Batrachochytrium dendrobatidis* infection patterns among Panamanian amphibian species, habitats and elevations during epizootic and enzootic stages. *Diseases of Aquatic Organisms* 81(3): 189–202.
- Bustamante HM, Livo LJ, Carey C. 2010. Effects of temperature and hydric environment on survival of the Panamanian Golden Frog infected with a pathogenic chytrid fungus. *Integrative Zoology* 5(2): 143–153.
- DiRenzo GV, Langhammer PF, Zamudio KR, Lips KR. 2014. Fungal Infection Intensity and Zoospore Output of *Atelopus zeteki*, a Potential Acute Chytrid Super-shedder. *PloS One* 9(3): e93356.
- Ellison AR, Savage AE, DiRenzo GV, Langhammer P, Lips KR, Zamudio KR. 2014. Fighting a Losing Battle: Vigorous Immune Response Countered by Pathogen Suppression of Host Defenses in the Chytridiomycosis-Susceptible Frog *Atelopus zeteki*. *G3: Genes Genomes Genetics*: g3–114.
- Gagliardo R, Crump P, Griffith E, Mendelson J, Ross H, Zippel K. 2008. The principles of rapid response for amphibian conservation, using the programmes in Panama as an example. *International Zoo Yearbook* 42(1): 125–135.
- Hertz A, Lotzkat S, Carrizo A, Ponce M, Köhler G, Streit B. 2012. Field notes on findings of threatened amphibian species in the central mountain range of western Panama. *Amphibian & Reptile Conservation* 6(2): 9–30 (e46).
- IUCN. 2013. Guidelines for using the IUCN Red List categories and criteria. Available: [www.iucnredlist.org](http://www.iucnredlist.org) [Accessed: 09 March 2014].
- Kilburn VL, Ibáñez R, Sanjurjo O, Bermingham E, Suraci JP, Green DM. 2010. Ubiquity of the pathogenic chytrid fungus, *Batrachochytrium dendrobatidis*, in anuran communities in Panama. *EcoHealth* 7(4): 537–548.
- Kilpatrick AM, Briggs CJ, Daszak P. 2010. The ecology and impact of chytridiomycosis: an emerging disease of amphibians. *Trends in Ecology & Evolution* 25(2): 109–118.
- La Marca E, Lips KR, Lötters S, Puschendorf R, Ibáñez R, Rueda-Almonacid JV, Young BE. 2005. Catastrophic population declines and extinctions in Neotropical harlequin frogs (Bufonidae: *Atelopus*). *Biotropica* 37(2): 190–201.
- Lips KR, Brem F, Brenes R, Reeve JD, Alford RA, Voyles J, Collins JP. 2006. Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proceedings of the National Academy of Sciences of the United States of America* 103(9): 3165–3170.
- Longcore JE, Pessier AP, Nichols DK. 1999. *Batrachochytrium dendrobatidis* gen. et sp. nov., a chytrid

- pathogenic to amphibians. *Mycologia* 91(2): 219–227.
- Mendelson JR, Lips KR, Gagliardo RW, Rabb GB, Collins JP, Diffendorfer JE, Brodie ED. 2006. Biodiversity-Confronting amphibian declines and extinctions. *Science* 313(5783): 48.
- Minteer BA, Collins JP, Love KE, Puschendorf R. 2014. Avoiding (Re) extinction. *Science* 344(6181): 260–261.
- Richards CL, Knowles LL. 2007. Tests of phenotypic and genetic concordance and their application to the conservation of Panamanian golden frogs (Anura, Bufonidae). *Molecular Ecology* 16(15): 3119–3133.
- Woodhams DC, Alford RA, Briggs CJ, Johnson M, Rollins-Smith LA. 2008. Life-history trade-offs influence disease in changing climates: strategies of an amphibian pathogen. *Ecology* 89(6): 1627–1639.
- Zipell KC, Ibáñez R, Lindquist ED, Richards CL, Jaramillo CA, Griffith EJ. 2006. Implicaciones en la conservación de las ranas doradas de Panamá, asociadas con su revisión taxonomica. *Herpetotropicos* 3(1). Available: <http://erevistas.saber.ula.ve/index.php/herpetotropicos/article/view/643> [Accessed: 25 October 2014].
- Zippel K, Johnson K, Gagliardo R, Gibson R, McFadden M, Browne R, Martinez C, Townsend E. 2011. The Amphibian Ark: A global community for ex situ conservation of amphibians. *Herpetological Conservation and Biology* 6(3): 340–352.



**Rachel Perez** is a M.S. student in the department of biology at New Mexico Tech. She received her B.S. at University of California, Riverside. Her main research interests are in the areas of epidemiology, ecoimmunology, and amphibian conservation. She is currently investigating innate defenses and microhabitat conditions of neotropical amphibians.



**Corinne Richards-Zawacki** is a professor in the Ecology and Evolutionary Biology department at Tulane University. Her research lies at the intersection of ecology and evolutionary biology in that she approaches questions about how changes in climate and habitat shape population and community processes in a way that explicitly considers their evolutionary implications. The questions she asks address (1) the effects of climate and landscape changes on species distributions and diversity, (2) how reproductive isolation evolves during speciation, and (3) how climate and host/pathogen evolution shape the dynamics of wildlife diseases. She is passionate about amphibian conservation and has 12 years of experience working on conservation-oriented projects in Panama. Much of her work in Panama has focused on the Critically Endangered Panamanian golden frogs. Her lab has also published studies related to the captive management of amphibian species threatened by *Bd*.



**Alexander Krohn** is a current Ph.D. candidate at the University of California, Berkeley, in the department of Environmental, Science, Policy and Management. His dissertation focuses on the convergent evolution of melanism in desert reptiles, but he has been interested in herpetology, tropical ecology, and conservation since he was in middle school.



**Matthew Robak** is broadly interested in amphibian conservation. He is currently a Ph.D. candidate at Tulane University where he is researching how differences in temperature affect amphibians' immune responses to *Batrachochytrium dendrobatidis* exposure.



**Edgardo J. Griffith** is a world-renowned leader in amphibian conservation in Panama and one of the founders of the El Valle Amphibian Conservation Center (EVACC). Edgardo is one of the pioneers that has proven that multi-species *ex situ* amphibian conservation is the only option we have at the moment to fight the dramatic amphibian declines in Panama. His work has been featured in several conservation books, documentaries, and peer-reviewed scientific papers. Edgardo has 14 years working with the Panamanian amphibians, including the Panamanian golden frog *in situ* and breeding them *ex situ*.



**Heidi Ross** is the project director at the El Valle Amphibian Conservation Center (EVACC). Heidi and her husband were awarded the San Diego Medal of International Conservation in 2012 for their work at EVACC. Heidi has 10 years of experience *ex situ* experience with Panamanian amphibians leading to the breeding of nine of the 12 priority species in the EVACC collection, including the Panamanian golden frog.



**Brian Gratwicke** is a conservation biologist that leads the amphibian conservation program at the National Zoo (Washington, DC, USA). Brian also leads the Panama Amphibian Rescue and Conservation Project. Previous conservation experience includes work on freshwater ecology in Africa, tropical marine ecology in the Caribbean, and tiger conservation efforts in Asia. He has published more than 25 peer-reviewed papers and book chapters and was a contributor to *Hotspots Revisited*. Brian received a Ph.D. in zoology from Oxford University. He obtained his bachelor's and master's degrees in zoology and fisheries ecology, respectively, from the University of Zimbabwe.



**Roberto Ibáñez** is the in-country Director of the Panama Amphibian Rescue and Conservation Project, based at the Smithsonian Tropical Research Institute. He has been part-time Associate Professor at the Universidad de Panamá (1996–2014), and professor during de Panama Field Study Semester of McGill University (2003–2013). He received his B.S. in Biology from the Universidad de Panama, and his Master's degree and Ph.D. in Zoology from the Ecology and Evolutionary Biology Department at the University of Connecticut. He has studied the amphibians and reptiles of Panama for more than 30 years. He has published more than 30 peer-reviewed papers, more than 20 notes, and a book guide to the amphibians of the lowlands of central Panama. He is a distinguished researcher of the national research system of Panama.



**Jamie Voyles** is currently an Assistant Professor at New Mexico Tech working on emerging infectious diseases in wildlife. She conducts chytridiomycosis research in Central America and in California. She is a member of multiple working groups investigating disease-related amphibian declines. She is actively involved in conservation initiatives, such as Amphibian Rescue and Conservation Project, and contributes to [amphibianrescue.org](http://amphibianrescue.org) and [AmphibiaWeb](http://AmphibiaWeb).