

SMITHSONIAN MISCELLANEOUS COLLECTIONS

PART OF VOLUME LIII

CAMBRIAN GEOLOGY AND PALEONTOLOGY

No. 5.—CAMBRIAN SECTIONS OF THE
CORDILLERAN AREA

WITH TEN PLATES

BY

CHARLES D. WALCOTT



No. 1812

CITY OF WASHINGTON
PUBLISHED BY THE SMITHSONIAN INSTITUTION
DECEMBER 10, 1908

CAMBRIAN GEOLOGY AND PALEONTOLOGY

No. 5.—CAMBRIAN SECTIONS OF THE CORDILLERAN AREA

By CHARLES D. WALCOTT

(WITH TEN PLATES)

CONTENTS

	Page
Introduction.....	167
Correlation of sections.....	168
House Range section, Utah.....	173
Waucoba Springs section, California.....	185
Barrel Spring section, Nevada.....	188
Blacksmith Fork section, Utah.....	190
Dearborn River section, Montana.....	200
Mount Bosworth section, British Columbia.....	204
Bibliography	218
Index	221

ILLUSTRATIONS

Plate 13. Map of central portion of House Range, Utah.....	172-173
Plate 14. West face of House Range south of Marjum Pass.....	173
Plate 15. Northeast face of House Range south of Marjum Pass; ridge east and southeast of Wheeler Amphitheater, House Range	178-179
Plate 16. North side of Dome Canyon, House Range.....	182
Plate 17. West face of House Range, below Tatow Knob.....	184
Plate 18. Cleavage of quartzitic sandstones, Deep Spring Valley, California.....	186
Plate 19. Sherbrooke Ridge on Mount Bosworth, British Columbia.....	207
Plate 20. Ridge north of Castle Mountain, Alberta; profile of southeast front of Castle Mountain.....	209
Plate 21. Mount Stephen, British Columbia.....	210
Plate 22. Profile of mountains surrounding Lake Louise, Alberta.....	216

INTRODUCTION

My first study of a great section of Paleozoic rocks of the western side of North America was that of the Grand Canyon of the Colorado River, Arizona. In this section the Cambrian strata extend down to the horizon of the central portion of the Middle Cambrian (Acadian) where the Cambrian rests unconformably on the pre-Cambrian formations.¹

¹ See American Jour. Sci., 3d ser., xxvi, 1883, pp. 437-442.

The second section studied was that of the Eureka District of central Nevada in 1880-1881, the results of which were incorporated in Monographs VIII and XX of the U. S. Geological Survey. This section includes the upper portion of the Lower Cambrian (Georgian), the Middle Cambrian (Acadian), and the Upper Cambrian (Saratogan). The studies of the Cambrian strata were afterward continued in the Cordilleran area from time to time as opportunity offered. These included the Highland Range section of Nevada and the Big Cottonwood section of the Wasatch Mountains (see Bulletin U. S. Geol. Survey, No. 30, 1886, pp. 33 and 38). The great House Range section of central western Utah was studied and measured in 1905, the Blacksmith Fork section of the Wasatch Mountains in 1906, and the Mount Bosworth section of British Columbia in 1907. The last three sections are included in this paper.

The strata of the Lower Cambrian (Georgian) are apparently well developed in the Big Cottonwood section of Utah, and the upper portion in the House Range, Eureka, and Highland Range sections, but it was not until the sections of the Lower Cambrian (Georgian) formations of western Nevada and southeastern California were examined that the fauna was found well developed. These sections are incorporated in this paper.

ILLUSTRATIONS.—In order that geologists and paleontologists who have not had an opportunity to see the sections may get an idea of the completeness of the exposures of the strata in the Cordilleran area, photographs are introduced in connection with the House Range and Mount Bosworth sections.

The map of the House Range gives the localities and names used in the section.

CORRELATION OF SECTIONS

The object of this preliminary correlation is to show in a broad way the interrelations of the strata and faunas in the North American Cordilleran area west of the great continental land area of Lower and much of Middle Cambrian time. The margin of this area was as far westward as the present position of the main range of the Wasatch Mountains in the vicinity of Salt Lake, Utah; from this point the shoreline trended gradually south-southwest to southwestern Utah and into southeastern Nevada. To the north of Salt Lake the trend of the early Cambrian shoreline was north-northeast to western Wyoming, and thence north into Montana (see Dearborn River section). It passed westward of the Belt Mountain

uplift, and thence north into Alberta, east of the Rocky Mountain front, where all traces of it are lost beneath the covering of Tertiary and Cretaceous rocks. In the vicinity of the international boundary (49th parallel) an uplift of pre-Cambrian (Beltian) strata appears to have largely prevented Cambrian sedimentation in northwestern Montana and northern Idaho. The faunas of the sections to the north in British Columbia and to the south in Utah clearly prove that the seas in which they lived were connected, but how or where we do not know.

In the following diagram the general relations of the sections are shown:

Table Showing Stratigraphic Position in the Cambrian System of Five of the Sections Described

Ordovician	+ 285			+		+
Upper Cambrian (Saratogan)	Utah, 3,315 feet.	300		Utah, 1,227 feet.		3,590 feet.
Middle Cambrian (Acadian)	House Range, 4,417 feet.	(?) (?)		Blacksmith Fork, 5,420 feet.		British Columbia, 4,963 feet.
Lower Cambrian (Georgian)	1,500 feet. Total, 9,232 +	Waucoba, California, 5,670 + +		Big Cottonwood, Utah, 12,000 feet.		Mount Bosworth, 3,800 feet.
			225			

The House Range section, supplemented by the Lower Cambrian sections of western Nevada and southeastern California, 230 miles (370.07 km.) west-southwest, gives a total of over 13,000 feet (3962 m.) of strata with Cambrian faunas throughout. If the Big Cottonwood section, 140 miles (225.26 km.) to the northeast of the

House Range, is found to have Cambrian fossils to its base, there will be over 19,000 feet of Cambrian strata in Utah. I think it quite probable that the quartzitic sandstones and siliceous shales of the Big Cottonwood section were being deposited as near-shore sediments while the calcareous, argillaceous, and arenaceous muds were accumulating at the same time 350 miles (563.15 km.) to the southwest.

The Upper and Middle Cambrian formations of the House Range section are much like those of the Blacksmith Fork and Mount Bosworth sections. From the top down the correlation of the various sections is as follows:


Correlation Table of Stratigraphic Sections


	House Range	Silver Peak	Big Cottonwood	Blacksmith Fork	Dearborn River	Mount Bosworth
Upper Cambrian (Saratogan)	Notch Peak, 1,490	Emigrant, 300	No Upper Cambrian	St. Charles, 1,227	(?)	Sherbrooke, 1,375
	Orr, 1,825					Paget, 360
Middle	Weeks, 1,390	(?)		Nouuan, 1,041	Limestone, 1,320	Eldon, 2,728
	Marjum, 1,102					
Cambrian	Wheeler, 570	(?)		Blacksmith, 570	Shale, 150	Stephen, 640
	Swasey, 340					
(Acadian)	Dome, 355	(?)		Ute, 729	Limestone, 130	Cathedral, 1,595
	Howell, 435					
Lower Cambrian (Georgian)	Spence, 20	Silver Peak, 5,670 +	Limestone, 75	Spence, 30	Shale, 210	Mt. Whyte, 390
	Langston(?), 205		Shale, 150	Langston, 498		
	Pioche, 125		Shale, 100	Brigham, 1,232 +	Shale, 190	
	Prospect Mountain, 1,375 +		Prospect Mountain, 11,750 +		Sandstone, 150	St. Piran, 2,705
					The sandstone rests unconformably on the pre-Cambrian	Lake Louise, 105
						Fairview, 600 +

The numerals indicate the thickness of each formation in feet. The only horizons definitely correlated by strongly marked and similar faunas are the Pioche and Mount Whyte; Spence and Stephen; Notch Peak, St. Charles, and Sherbrooke.

There are many partial sections that supplement various portions of the three great sections. These I wish to utilize in connection with the study of the Cambrian trilobites of the Cordilleran area, as our present knowledge of the vertical range and distribution of the trilobites is too limited and inaccurate to be more than of value in general and broad correlations. It is also true that many of the great limestone beds now considered as almost without fossils will be found in their extension away from the three great sections to contain a well-marked fauna.


In closing this brief review, I wish to call attention to the close relationship between the great Cambrian section of the Province of Shantung, China, and the Cordilleran sections. The thickness of the strata is very much less, but the general character and stratigraphic succession of the Cambrian faunas is very much the same. This will be discussed in the introduction to a paper on the Cambrian faunas of China, upon which I am now at work.





HOUSE RANGE, MILLARD COUNTY, UTAH


CENTRAL PORTION



Contour interval 200ft
 Datum is mean sea level

Contours below "A-B" have been added to Map to show approximate relations of Notch Peak and Orr Ridge to the rest of the range

TOPOGRAPHY BY W. D. JOHNSON, 1901



HOUSE RANGE, MILLARD COUNTY, UTAH CENTRAL PORTION




Contour interval 500'

Datum is in mean sea level

Contours below 'A' B' have been added to Map to show approximate relations
of Notch Peak and Orr Ridge to crest of the range

TOPOGRAPHY BY W. D. JOHNSON, 1901



WEST FACE OF NOTCH PEAK, HOUSE RANGE, SOUTH OF MARJUM PASS, UTAH

The summit of Notch Peak is capped with Ordovician limestone, and an intrusive mass of granite porphyry is intruded into the Cambrian beds on the north slope of the peak (left side). The Notch Peak formation is beautifully shown in the 1,400-foot cliff exposed in the dark canyon just beneath the peak. (See Plate 15, Figure 1, for opposite side of Notch Peak.)

HOUSE RANGE SECTION

In order to locate the various points referred to, the accompanying map has been prepared, under the direction of Mr. L. D. Burling, from a manuscript topographic map made by Mr. W. D. Johnson of the U. S. Geological Survey (see plate 13).

LOCALITY.—West and east of Antelope Springs and east-southeast and south of Marjum Pass, House Range, Millard County, Utah. Sawtooth Range is a name given locally to the House Range south of Marjum Pass.

The section begins at the top, 285 feet below the summit of Notch Peak, the highest point (8,828 feet) on the House Range south of Marjum Pass.

The top of the peak is formed of 285 feet of Ordovician limestone, which is a banded, thin-bedded, bluish gray and purplish limestone containing near the top a distinct fauna:

Obolus (Westonia) notchensis Walcott [1908*d*, p. 69].

Eoorthis coloradoensis (Meek) [1870, p. 425].

Raphistoma sp., etc.

The strike of the upper beds is north 20° east (magnetic); dip, 12° south.

The line of the section extends down the northeast slope of Notch Peak and thence to a high ridge east of the area of eruptive granite on the northwest slope of Notch Peak; thence north to Marjum Pass. It is then carried on the line of the upper beds of the Wheeler formation to a point southeast of Antelope Springs; thence west to Dome Pass and (on the north side of Dome Canyon) to the Lower Cambrian quartzitic sandstones that pass beneath the quaternary of the White Valley at the western foot of the House Range.

ORDOVICIAN

	Feet
Limestone resting conformably on the Cambrian.....	285

UPPER CAMBRIAN

NOTCH PEAK FORMATION:

The Notch Peak formation [Walcott, 1908*a*, p. 9] is exposed on the east and southeast slopes and ridges of Notch Peak. Feet

- 1a. Gray, arenaceous limestone in thick layers and bands of thin layers. Irregular nodules and thin layers of dark gray chert, weathering dark brown, occur at irregular intervals for 350 feet below the summit. Thin, cherty layers, one-half to one-eighth inch thick, also occur occasionally below..... 640

Ord.		Feet	Me- ters	
		250	76	Thin-bedded limestone
Upper Cambrian	Notch Peak lime- stone	1490	454	Massive bedded, arenaceous limestone with a few thin beds
	Orr limestones and shale	375	114	Thin bedded limestones with two bands of arenaceous shale
		84	26	
		206	63	
235		72		
	925	282		
Middle Cambrian	Weeks limestone	1390	424	Thin bedded and shaly lime- stones
	Marjum limestone	1102	336	Thin bedded limestones
	Wheeler shales	570	174	Calcareous shales
	Swasey limestone	152 188	46 57	Arenaceous limestone above, with thin bedded limestone below
	Dome limestone	355	108	Massive bedded, arenaceous limestone
	Howell limestone	435	133	Thick and thin bedded limestone
	Spence shale	< 20	6	Argillaceous shale
	Langston (?) limestone	205	62	Bluish gray arenaceous limestone
	Pioche shale	125	38	Arenaceous shale
Lower Cambrian	Prospect Mountain sandstones	1375	419	Brown quartzitic sandstone

FIG. 6.—House Range Section.

NOTCH PEAK FORMATION (continued):

1a (continued):

Feet

*Fauna:**Lingulella isse* (Walcott) [1905, p. 330].*Dicellosephalus* ? sp. ?

A drift boulder found 2.5 miles from the peak, and on its eastern drainage slope, and similar in its lithological appearance to the gray, arenaceous limestone of this horizon, contained the following fossils:

Eoorthis coloradoensis (Meek) [1870, p. 425].*Schizambon typicalis* Walcott [1884, p. 70].*Agraulos*.*Solenopleura*.*Illænurus*.

Another drift boulder was found near this with slightly different fauna.

Crepicephalus.*Ptychoparia*.

- 1b. Shaly, dark gray to bluish gray, arenaceous limestone, with small dark concretions in some layers..... 90.
No fossils observed.
- 1c. Gray, siliceous limestone in layers of varying thickness, 4 inches to 2 feet, banded with dark cherty layers and purer arenaceous limestone. The chert takes the form of flattened nodules and very thin irregular layers..... 340
- 1d. Shaly and thin-bedded, bluish gray, arenaceous limestone.... 65.
- 1e. Gray, siliceous limestone in layers 2 inches to 2 feet thick. In the lower part of this limestone, where it is not metamorphosed, it is dove-colored and in layers 6 inches to 3 feet thick. There are occasional occurrences of gray, cherty matter, as flattened nodules, and thin layers that weather a dark brown 355.

Fauna (about 120 to 150 feet from the base):*Obolus tetonensis leda* Walcott [1908d, p. 63].

Fragments of the free cheek of a trilobite.

Total of Notch Peak formation..... 1,490

ORR FORMATION:

The section is carried along the strike of the exposed strata two miles east to the west side of Orr Ridge, where the rocks of the Orr formation [Walcott, 1908a, p. 10] are unmetamorphosed and present the following characters:

- 1a. Bluish gray to gray, compact limestone in layers 1 inch to 2 feet thick. On weathering the thicker layers break down into thin, irregular layers, which form a talus of angular fragments 375.

ORR FORMATION (continued):

- | | Feet |
|---|------|
| 1a (continued): | |
| <i>Fauna:</i> | |
| Fragments of trilobites. | |
| 1b. Sandy and siliceous, bluish and drab-colored shales, with interbedded bands of dark, bluish gray limestone 6 inches to 2 feet thick | 84 |
| <i>Fauna:</i> | |
| Section of crinoid column. | |
| <i>Lingulella manticula</i> (White) [1874, p. 9]. | |
| <i>Lingulella isse</i> (Walcott) [1905, p. 330]. | |
| <i>Obolus rotundatus</i> (Walcott) [1898, p. 415]. | |
| <i>Ptychaspis</i> . | |
| <i>Anomocare</i> . | |
| 1c. Lead-colored, finely oölitic, and arenaceous limestone in layers 4 inches to 2 feet thick that are obscurely banded by thin strips of light and dark gray color..... | 91 |
| <i>Fauna:</i> | |
| Fragments of trilobites. | |
| 1d. Bluish gray, compact limestone in layers 2 inches to 4 feet thick that break down into irregular, thin layers on weathering... | 115 |
| <i>Fauna</i> (near base): | |
| Fragments of trilobites. | |
| <i>Linnarssonella modesta</i> Walcott [1908d, p. 90]. | |
| <i>Linnarssonella nitens</i> Walcott [1908d, p. 91]. | |
| <i>Solenopleura</i> . | |
| 1e. Dirty brown and bluish black, arenaceous shales, with thin nodules of gray, fossiliferous limestone in some horizons; also a few layers of bluish gray limestone 4 inches to 8 inches thick | 235 |
| <i>Fauna</i> (near the top): | |
| <i>Linnarssonella modesta</i> Walcott [1908d, p. 90]. | |
| <i>Lingulella isse</i> (Walcott) [1905, p. 330]. | |
| <i>Ptychoparia?</i> | |
| <i>Solenopleura</i> . | |
| <i>Fauna</i> (near the base): | |
| <i>Micromitra</i> (<i>Paterina</i>) <i>crenistria</i> ? (Walcott) [1897, p. 713]. | |
| <i>Obolus mcconnelli pelias</i> (Walcott) [1905, p. 330]. | |
| <i>Lingulella desiderata</i> (Walcott) [1898, p. 399]. | |
| <i>Lingulella isse</i> (Walcott) [1905, p. 330]. | |
| <i>Linnarssonella transversa</i> Walcott [1908d, p. 92]. | |
| <i>Agnostus</i> . | |
| <i>Crepiccephalus</i> . | |
| 2a. Gray, slightly arenaceous limestone in layers 2 to 6 feet thick, weathering lead gray. (Cliff-forming beds.)..... | 590 |

ORR FORMATION (continued):

	Feet
2a (continued):	
Fauna (at base):	
<i>Lingulella desiderata</i> (Walcott) [1898, p. 399].	
<i>Acrotreta idahoensis</i> Walcott [1902, p. 587].	
<i>Crepicephalus texanus</i> (Shumard) [1861, p. 218].	
<i>Bathyuriscus</i> .	
<i>Illænurus</i> ??	
Fauna (275 feet above base):	
<i>Agraulos</i> .	
<i>Crepicephalus texanus</i> (Shumard) [1861, p. 218].	
<i>Illænurus</i> .	
2b. Gray limestone and dark gray chert in alternating layers, one-half to 2 inches thick. The irregular cherty layers weather in relief as dark brown bands and the limestone as lead-colored bands, which give a very characteristic banded appearance to the cliff.....	170
2c. Gray, arenaceous limestone in massive beds that usually break up, on weathering, into irregular layers one-fourth to 4 inches thick. The upper 20 feet form a more massive, solid bed than the layers below.....	165
Fauna:	
Traces of trilobites and brachiopods.	
Total of Orr formation.....	1,825
Total Upper Cambrian.....	3,315

MIDDLE CAMBRIAN

WEEKS FORMATION:

The Weeks formation [Walcott, 1908a, p. 10] is exposed at Weeks Canyon (see pl. 13) from beneath the massive limestone on the south side of the canyon to the top of the cliffs on the south side of Marjum Pass. Average dip, 12°; strike, north 20° east (magnetic).

	Feet
1a. Thin-bedded limestones in layers 1 to 4 inches thick. The limestone is mainly fine-grained, dark gray, weathering lead-colored, except on bedding planes, where it is usually more or less pinkish colored.....	245
Fauna:	
Fragments of trilobites and brachiopods of the fauna in shaly limestone in 1b.	
1b. Shaly limestone, usually dark gray, with pinkish tinge in some layers and on the surfaces; sometimes buff yellow on weathering. The shales vary from one-eighth to 1 inch thick. This is a marked band in some sections and is arbitrarily separated from the shaly beds below.....	285

WEEKS FORMATION (continued):

1b (continued):

Feet

Fauna:

The fauna ranges through about 100 feet of the lower portion of this division.

Obolus (Fordinia) perfectus Walcott [1908d, p. 65].

Agnostus (2 species).

Ptychoparia.

Crepicephalus texanus (Shumard) [1861, p. 218].

Anomocare.

Bathyriscus.

Asaphiscus minor, new species.

-The fauna is much like that of 1c. Its most characteristic trilobite is *Asaphiscus minor*, new species.

- 1c. Shaly, bluish gray to dark gray limestone in layers one-eighth to 1 inch thick, with occasional layers 2 to 6 inches thick; 25 feet from the top a band of layers of arenaceous, dirty gray, finely oölitic limestone, 3 feet thick, occurs, and a second similar band 38 feet below.

170

Fauna:

The fauna is rich in numbers of specimens and quite varied. The best specimens occur on the surface of the shaly layers in the lower portion of the division.

Lingulella isse (Walcott) [1905, p. 330].

Obolus (Fordinia) perfectus Walcott [1908d, p. 65].

Acrotreta ophirensis Walcott [1902, p. 591].

Acrotreta ophirensis descendens Walcott [1908d, p. 94].

Hyalithes.

Agnostus (several species).

Ptychoparia (several species).

Crepicephalus texanus (Shumard) [1861, p. 218].

Solenopleura.

Asaphiscus minor, new species.

- 1d. Reddish tinted, more or less arenaceous, shaly limestone.

30

Fauna:

Same as 1c, but not abundant.

- 1e. Shaly, bluish gray to dark gray limestone, similar to 1c.

270

Fauna:

Same as that of 1c.

- 1f. Evenly bedded, bluish gray to dark gray, fine-grained limestone, in layers 2 to 16 inches thick, with shaly limestone partings

330

Fauna:

A few traces of *Agnostus* and *Ptychoparia* similar to those above.

- 1g. Calcareous shales with thin layers of limestone.

60

Total thickness of Weeks formation. 1,390



Fig. 1. VIEW FROM THE NORTHEAST OF THE EAST

The rounded hills of the foreground are eroded in the Wheeler Cambrian limestones of the Weeks, Orr, and Notch Peak formations, by 285 feet of Ordovician limestone.




Fig. 2. PANORAMIC VIEW OF RIDGE

Looking across Wheeler Amphitheater, House Range. The Wheeler shale extends to the base of the low f on the slopes of the mountain on the left side of the illustration.



HOUSE RANGE SOUTH OF MARJUM PASS, UTAH

limestone forms the long horizontal cliff, and back of this the
continue on up to near the summit of Notch Peak, which is capped

VOL. 53, PL. 15—CONTINUED



EAST OF ANTELOPE SPRINGS

Marjum formation to the summit of the ridge. The best known fossil localities in the Marjum formation



Fig. 1. VIEW FROM THE NORTHEAST OF THE EASTERN END OF THE HOUSE RANGE SOUTH OF MARJUM PASS, UTAH

The rounded hills of the foreground are eroded in the Wheeler shale. The Marjum limestone forms the long horizontal cliff, and back of this the Cambrian limestones of the Weeks, Orr, and Notch Peak formations, and black, continue on up to near the summit of Notch Peak, which is capped by 285 feet of Ordovician limestone.



Fig. 2. PANORAMIC VIEW OF RIDGE RUNNING SOUTHEAST OF ANTELOPE SPRINGS

Looking across Wheeler Amphitheater, House Range. The Wheeler shale extends to the base of the low part of the ridge. The best known fossil localities in the Marjum formation are on the slopes of the mountain on the left side of the illustration.

MARJUM FORMATION:

The Marjum formation [Walcott, 1908a, p. 10] is exposed in the cliffs southeast of Marjum Pass and in the ridge east of Wheeler Amphitheater.

Feet

1a. Gray, more or less thin-bedded limestone that weathers to a dark lead-gray color and breaks down into angular fragments one-half to 2 inches thick.

Flattened cherty nodules and thin, irregular cherty layers occur at intervals 305

Fauna (in upper 100 feet):

- Obolus mcconnelli pelias* (Walcott) [1905, p. 330].
- Obolus (Fordinia) gilberti* Walcott [1908d, p. 65]
- Obolus (Fordinia) perfectus* Walcott [1908d, p. 65].
- Acrotreta bellatula* Walcott [1908d, p. 93].
- Acrotreta marjumensis* Walcott [1908d, p. 94].
- Acrotreta* cf. *sagittalis* Salter [1866, p. 285].
- Agnostus* (4 species).

Fauna (central portion):

- Micromitra sculptilis* Meek [1873, p. 479].
- Lingulella arguta* (Walcott) [1898, p. 396].
- Dicellomus prolificus* Walcott [1908d, p. 77].
- Acrotreta attenuata* Meek [1873, p. 463].
- Acrotreta bellatula* Walcott [1908d, p. 93].
- Agnostus*.
- Ptychoparia*.
- Anomocare*.

Fauna (near base):

- Micromitra (Iphidella) pannula ophirensis* (Walcott) [1905, p. 306].
- Obolus mcconnelli pelias* (Walcott) [1905, p. 330].
- Obolus rotundatus* (Walcott) [1898, p. 415].
- Hyalithes*.
- Ptychoparia*.
- Anomocare*.

1b. Alternating bands of dark, blue-gray, compact limestone in massive layers that break up into thin irregular layers; and gray arenaceous limestone in layers 1 to 8 inches thick.

Feet

- 1. Gray limestone 35
- 2. Blue-gray limestone 7
- 3. Gray arenaceous limestone..... 95
- 4. Blue-gray limestone 12
- 5. Gray arenaceous limestone..... 90
- 6. Blue-gray limestone 8

247

Fauna:

- Ptychoparia*, sp. undt.

1c. Dark and light-gray, thin-bedded limestone, more or less arenaceous

250

MARJUM FORMATION (continued):

1c (continued):

Feet

Fauna (near top):

Acrotreta pyxidicula White [1874, p. 9].*Agnostus*.*Ptychoparia* like *P. kingi* (Meek) [1870, p. 63].

Fauna (in central portion, though ranging through 100 to 150 feet of the thin-bedded shaly limestone):

Obolus mcconnelli pelias (Walcott) [1905, p. 330].*Lingulella arguta* (Walcott) [1898, p. 396].*Acrothele subsidua* (White) [1874, p. 6].*Acrotreta ophirensis*? Walcott [1902, p. 591].*Eoorthis thyone* Walcott [1908d, p. 105].*Nisusia (Jamesella) nantes* (Walcott) [1905, p. 283].*Nisusia (Jamesella) spencei* (Walcott) [1905, p. 285].*Hyaloliths*.*Agnostus* (2 species).*Ptychoparia* (3 species).*Solenopleura*.*Oxtenella tyra*, new genus and new species.*Neolenus inflatus* Walcott [1908b, p. 30].*Neolenus intermedius* Walcott [1908b, p. 34].*Neolenus intermedius pugio* Walcott [1908b, p. 35].*Neolenus superbus* Walcott [1908b, p. 36].*Ogygopsis*?

1d. Gray, shaly limestone, passing below into shales, interbedded in the shaly limestone, and at 75 feet from the top into drab argillaceous shales

105

Fauna:

Micromitra (Iphidella) pannula ophirensis (Walcott) [1905, p. 306].*Micromitra sculptilis* Meek [1873, p. 479].*Obolus mcconnelli pelias* (Walcott) [1905, p. 330].*Obolus rotundatus* (Walcott) [1898, p. 415].*Lingulella arguta* (Walcott) [1898, p. 396].*Acrotreta attenuata* Meek [1873, p. 463].*Acrotreta ophirensis* Walcott [1902, p. 591].*Acrothele subsidua* (White) [1874, p. 6].*Acrothele subsidua laevis*, new variety.*Eoorthis remnicha* (N. H. Winchell) [1886, p. 317].*Eoorthis thyone* Walcott [1908d, p. 105].*Syntrophia unxia* Walcott [1908d, p. 105].*Agnostus* (3 species).*Ptychoparia*.*Oxtenella tyra*, new genus and new species.*Neolenus inflatus* Walcott [1908b, p. 30].*Neolenus intermedius* Walcott [1908b, p. 34].*Neolenus superbus* Walcott [1908b, p. 36].*Ogygopsis*?

MARJUM FORMATION (continued):

1c. Dark, bluish gray limestone in thick beds that break up on weathering into thin, irregular layers one-half to 2 inches thick 195

Fauna:

Linnarssonella sp.
Agnostus.
Ptychoparia.
Ogygopsis.

Total thickness of Marjum formation..... 1,102

WHEELER FORMATION:

The Wheeler formation [Walcott, 1908a, p. 10] is exposed at Marjum Pass, but the type locality is in Wheeler Amphitheater, southeast of Antelope Springs. The section was measured south from the ridge south of the lower springs of Antelope Springs. Feet

1. Alternating bands of thin, shaly limestone and calcareous shale, with shale gradually increasing and predominating toward the lower portion. At 405 feet from top a band of blue-gray, hard limestone, in layers one-eighth to 2 inches thick, occurs. At 473 feet another band, and below an occasional thin layer..... 570

Fauna:

Acrothele subsidua (White) [1874, p. 6].
Agnostus bidens Meek [1873, p. 463].
Asaphiscus wheeleri Meek [1873, p. 485].
Ptychoparia kingi (Meek) [1870, p. 63].

These species occur in great numbers at 230 feet to 350 feet from the base. Many hundred trilobites, entire and backed by "cone-in-cone," have been picked up on the surface of the clay, resulting from the disintegration of the shales.

Obolus mcconnelli pelias Walcott and *Acrotreta attenuata* Meek occur more rarely.

SWASEY FORMATION:

The section of the Swasey formation [Walcott, 1908a, p. 11] is exposed on the southwest ridge of Swasey Peak. Feet

1a. Oölitic and arenaceous limestone in massive layers near the top. Below, dark bluish gray limestone is occasionally interbedded, and gradually it becomes the principal rock; it breaks up on weathering into irregular, shaly layers one-half to 3 inches thick..... 152

Fauna (near the top):

Platyceras.
Zacanthoides.

Fauna (near the base):

Scenella.
Zacanthoides.
Ptychoparia.
Dorypyge.

SWASEY FORMATION (continued):

	Feet
1b. Drab and reddish argillaceous shales, with interbedded, thin layers of fossiliferous limestone.....	63
1c. Dark, bluish gray limestone in massive layers that break up into irregular, shaly layers one-fourth to 2 inches thick....	17
1d. Calcareous and argillaceous shales with thin layers of gray limestone	102

Fauna:

Micromitra (Paterina) labradorica utahensis (Walcott)
[1905, p. 306].

Lingulella arguta (Walcott) [1898, p. 396].

Ptychoparia (2 species).

1e. Bluish gray limestone in layers 4 to 10 inches thick, with numerous concretions from one-eighth to 1 inch in diameter, in a few layers.....	6
---	---

Fauna:

Obolus (Westonia) ella (Hall and Whitfield) [1877, p. 232].

Ptychoparia (3 species).

Total of Swasey formation.....	340
--------------------------------	-----

DOME LIMESTONE:

The section of the Dome limestone [Walcott, 1908a, p. 11] is exposed in the central portion of Dome Canyon and adjoining cliffs. Feet

Massive bedded, cliff-forming, gray, siliceous limestone, with small specks of calcite. One hundred feet from the top, and for 50 feet below, occasional layers 15 inches to 2 feet thick, of brownish yellow, arenaceous limestone, occur.....	355
---	-----

HOWELL FORMATION:

The section of the Howell formation [Walcott, 1908a, p. 11] is exposed on the west face of the House Range at Howell Mountain. Feet

1a. Bluish black limestone in massive layers that break up on weathering into irregular, thin layers.....	50
---	----

Fauna (in shaly bed at top of 1a):

Micromitra (Iphidella) pannula (White) [1874, p. 6].

Acrotreta cf. *ophirensis* Walcott [1902, p. 591].

Ptychoparia.

1b. Gray, siliceous limestone.....	8
1c. Bluish black limestone, similar to 1a.....	105
1d. Pinkish colored, argillaceous shale with interbedded, thin layers of limestone.....	10

Fauna:

Micromitra (Iphidella) pannula (White) [1874, p. 6].

Obolus (Westonia) ella (Hall and Whitfield) [1877, p. 232].


Acrotreta cf. *ophirensis* Walcott [1902, p. 591].

Scenella.

Hyalithes.

Zacanthoides.

Bathyriscus.



VIEW OF THE NORTH SIDE OF DOME CANYON BELOW DOME PASS, HOUSE RANGE

The dark Swasey limestone forms the dark peak at the left. This rests on the light gray cliff of Dome limestone, below which the Howell formation breaks down. A mass of the Dome limestone has been displaced in the foreground by a fault between it and the cliff. The dome from which the canyon formation takes its name is shown at the right.

HOWELL FORMATION (continued):

	Feet
1e. Gray, siliceous limestone in layers 2 to 10 inches thick.....	70
1f. Bluish black limestone in massive layers, breaking up into thin layers on weathering.....	102

Fauna:

Ptychoparia.
Bathyuriscus.

1g. Gray, siliceous limestone in thick beds.....	90
--	----

Total of Howell formation.....	435
--------------------------------	-----

Spence shale:

The Spence shale [Walcott, 1908a, p. 8] is exposed on the east side of Dome Canyon a little above where it bends to the westward. Feet

1. Pinkish, argillaceous shale.....	20
-------------------------------------	----

Fauna:

Micromitra (Iphidella) pannula (White) [1874, p. 6].
Obolus (Westonia) ella (Hall and Whitfield) [1877, p. 232].
Lingulella dubia (Walcott) [1898, p. 401].
Acrothele subsidua (White) [1874, p. 6].
Hyalithes billingsi Walcott [1886, p. 134].
Ptychoparia piochensis Walcott [1886, p. 201].
Ptychoparia sp.
Zacanthoides typicalis (Walcott) [1886, p. 183].
Bathyuriscus productus (Hall and Whitfield) [1877, p. 244].

LANGSTON (?) FORMATION:

The section of the beds which are doubtfully placed in the Langston formation [Walcott, 1908a, p. 8] was measured at the same locality as the Spence shale. Feet

1a. Massive bedded, bluish gray, arenaceous limestone, with irregular partings of buff-colored arenaceous limestone. The latter penetrates the layers of limestone in the most irregular manner and frequently surrounds small, irregular nodules of the bluish gray limestone.....	170
---	-----

Fauna:

Billingsella, sp. undt.
Platyceras.
Hyalithes.
Leperditia.
Ptychoparia.
Zacanthoides.
Dorypyge?

1b. Brown, buff weathering, arenaceous limestone in thick layers; almost sandstone in places.....	35
---	----

Total of Langston (?) formation.....	205
--------------------------------------	-----

Total Middle Cambrian.....	4,417
----------------------------	-------

LOWER CAMBRIAN

PIOCHE FORMATION:

The Pioche formation [Walcott, 1908a, p. 11] is exposed at the westward bend of Dome Canyon. Feet

- i. Arenaceous and siliceous shaly layers, with some thicker layers of quartzitic sandstone..... 125

Fauna:

- Annelid trails.
Trilobite tracks (*Cruziana*).

Southwest of Pioche, Nevada, on the Panaca Road, this formation contains the following fauna:

- Eocystites* ? *longidactylus* Walcott [1886, p. 94].
Obolus (*Westonia*) *ella* (Hall and Whitefield) [1877, p. 232].
Micromitra (*Iphidella*) *pannula* (White) [1874, p. 6].
Acrothele subsidua (White) [1874, p. 6].
Acrothele subsidua hera Walcott [1908d, p. 87].
Acrothele spurri Walcott [1908d, p. 86].
Acrotreta primæva Walcott [1902, p. 593].
Billingsella highlandensis (Walcott) [1886, p. 119].
Hyolithes billingsi Walcott [1886, p. 134].
Olenellus gilberti Meek [1874, p. 7].
Zacanthoides levis (Walcott) [1886, p. 187].
Crepicephalus augusta Walcott [1886, p. 208].
Crepicephalus liliana Walcott [1886, p. 207].

PROSPECT MOUNTAIN FORMATION:¹

The Prospect Mountain formation [see Walcott, 1908a, p. 12] is exposed on the west slope and foothills of the House Range north and south of Dome Canyon. Feet


- i. Gray and brownish quartzitic sandstone in layers 4 inches to three feet in thickness..... 1,375+

Total Lower Cambrian..... 1,500+

RÉSUMÉ, HOUSE RANGE SECTION

UPPER CAMBRIAN:	Feet	Feet
Notch Peak formation.....	1,490	
Orr formation	1,825	
Total		3,315


¹ As the result of conference with Mr. Arnold Hague, the following formation names are given for formations in the Eureka section (see Walcott, 1884, p. 284): Eldorado limestone replaces Prospect Mountain limestone; Dunderberg shale replaces Hamburg shale, the name Hamburg being retained for the Hamburg limestone.



The Prospect Mountain formation forms the lowest beds and above, the Pioche shale of the Lower Cambrian, then in turn the limestones of the Langston and Howell formations separated by a narrow band of Spence shale, and above, the Dome and Swasey limestones. Tatow is a word of Indian derivation signifying "nipple", and the topographic feature to which it refers is locally known as "Molli's Nipple."

SILVER PEAK GROUP (continued):

1d (continued):	Feet
Fauna (105 feet from the base):	
<i>Salterella</i> .	
<i>Holmia weeksi</i> , new species.	
Total of 1.....	1,040
2a. Dark, siliceous, indurated shales, shaly sandstone and quartzitic sandstone in alternating layers.....	35
Fauna:	
Annelid trails.	
<i>Crusiana</i> .	
2b. Buff, drab, and bluish gray arenaceous limestone alternating in layers and bands.....	120
2c. Gray and dirty brown sandstones, with bands of light gray quartzitic sandstones	125
2d. Gray limestone, becoming arenaceous and passing into gray and dirty brown sandstone.....	105
Fauna:	
Traces of fragments of trilobites on the surface of the sandstone.	
2e. Gray and dirty brown, compact sandstone in layers from 2 inches to 2 or 3 feet in thickness. In the lower portion of the strata are layers of massive bedded, gray quartzitic sandstone. Small concretions 1 to 2 millimeters in diameter are very abundant in many of the upper layers.....	365
2f. Hard, brown and gray, shaly sandstones, with an occasional irregular, thin layer of bluish gray limestone.....	155
2g. Bluish gray arenaceous limestone in thick layers.....	25
2h. Greenish-colored arenaceous shale.....	120
2i. Alternating bands of arenaceous shale and massive bedded, gray, quartzitic sandstones.....	430
Fauna:	
<i>Scolithus</i> occurs abundantly in many of the quartzitic sandstones.	
Fauna (50 feet from the top):	
<i>Salterella</i> .	
<i>Olenellus</i> (fragments).	
2j. Gray quartzitic sandstones in layers 8 inches to 3 feet in thickness, passing below 35 feet into buff to yellowish shale with greenish buff bands, with some interbedded, gray, quartzitic sandstones	485
Fauna (in quartzitic layers):	
<i>Scolithus</i> like <i>S. linearis</i> Haldeman [see Walcott, 1891, p. 603].	
Fauna (in lower portion):	
<i>Obolella</i> , sp. undt.	
<i>Holmia rowei</i> , new species.	



LOWER CAMBRIAN QUARTZITES

Showing vertical cleavage in massive layers and interbedded thin layers without cleavage. Soldiers Canyon, above Deep Spring Valley, White Mountain range, Inyo County, California.

SILVER PEAK GROUP (continued):

2j (continued):

Feet

Near Resting Springs, in the Kingston Range, and at about this horizon, Mr. R. B. Rowe collected the following:

Billingsella highlandensis (Walcott) [1886, p. 119].

Holmia weeksi, new species.

Olenellus fremonti, new species.

2k. Gray and brownish gray quartzitic sandstones in layers 6 inches to 3 feet in thickness..... 790

Fauna:

Annelid trails on the surface of the layers.

Total of 2 2,755

3a. At summit a band of bluish gray limestone, with sandstones and occasional layers of thin-bedded limestone below. At 290 feet down a band of arenaceous limestone 50 feet thick occurs. Below this, brown sandstone and sandy shales, with interbedded thin layers of limestone in the lower 100 feet 650

Fauna (430 feet from the base):

Numerous fragments of *Olenellus*.

3b. Argillaceous and sandy shale, with a few thin beds of limestone 200

Fauna (160 feet from base):

Obolella, sp. undt.

Trematobolus excelsis Walcott [1908d, p. 80].

3c. Alternating arenaceous limestones, shales, and dirty brown sandstones that break into angular blocks and fragments... 575

Fauna (275 feet from base):

Archæocyathus is very abundant.

3d. Shaly indurated sandstones, with a few thicker layers of almost quartzitic sandstone..... 450+

Fauna (on west slope of hill just east of the summit, where the Saline Valley wagon road passes down the slope toward Waucoba Springs):

Annelid trails.

Cruziana.

Trematobolus excelsis Walcott [1908d, p. 80].

Holmia rowei, new species.

Fauna (on the east side of the hill and in lower portion of 3d):

Archæocyathus.

Ethmophyllum gracile Meek [1868, p. 62].

Mickwitzia occidens Walcott [1908d, p. 54].

Obolella, sp. undt.

SILVER PEAK GROUP (continued):

3d (continued):

Feet

Trematobolus excelsis Walcott [1908d, p. 80].

Hyalithes sp.

Holmia rowei, new species.

Total of 3. 1,875+

Total of section. 5,670+

In this section the genus *Olenellus* is found extending through 4,900 feet of strata and its lower limit is unknown.

RÉSUMÉ, WAUCOBA SPRINGS SECTION

	Feet	Feet
1a. Limestone	525	
1b. Limestone	115	
1c. Limestone	60	
1d. Arenaceous limestone	340	
	—————	1,040
2a. Shales	35	
2b. Arenaceous limestone	120	
2c. Sandstone	125	
2d. Arenaceous limestone	105	
2e. Sandstone	365	
2f. Sandstone	155	
2g. Arenaceous limestone	25	
2h. Shale	120	
2i. Shale and sandstone.	430	
2j. Sandstone and shale.	485	
2k. Quartzitic sandstone	790	
	—————	2,755
3a. Shales, limestone, and sandstone.	650	
3b. Shaly sandstone.	200	
3c. Arenaceous limestone and shaly sandstone.	575	
3d. Hard sandstones.	450+	
	—————	1,875+
	—————	5,670+
Total		5,670+

BARREL SPRING SECTION

A section of Lower Cambrian strata studied by Mr. F. B. Weeks near Barrel Spring, 16 miles south of the town of Silver Peak, Nevada, is much like that east of Waucoba Springs, and has about the same fauna at various horizons in it.

- | | |
|---|-----|
| 1. Massive blue mottled limestone, with 50 feet of sandy limestone in the middle of the series. | 737 |
|---|-----|

Feet

Fauna:

Archaeocyathus and allied forms occur throughout this limestone.

	Feet
2. Sandy shales succeeded by coarse, thin, fine sandstone, with buff limestones at top.....	206
<i>Fauna</i> (in limestone):	
<i>Micromitra</i> (<i>Paterina</i>) <i>prospectensis</i> (Walcott) [1884, p. 19].	
<i>Nisusia</i> (<i>Jamesella</i>) <i>amii</i> Walcott [1905, p. 252].	
<i>Scenella</i> , sp.	
<i>Agraulos</i> ?	
<i>Olenellus gilberti</i> Meek [1874, p. 7].	
3. Green calcareous shale, arenaceous at top.....	390
<i>Fauna</i> :	
<i>Archæocyathus</i> ?	
<i>Kutorgina cingulata</i> (Billings) [1861, p. 8].	
<i>Kutorgina perugata</i> Walcott [1905, p. 310].	
<i>Siphonotreta</i> ? <i>dubia</i> , new species.	
<i>Acrotreta claytoni</i> Walcott [1902, p. 583].	
<i>Acrothole spurri</i> ? Walcott [1908d, p. 86].	
<i>Swantonina weeksi</i> Walcott [1905, p. 297].	
<i>Swantonina</i> ? sp.	
<i>Stenotheca</i> cf. <i>elongata</i> Walcott [1884, p. 23].	
<i>Stenotheca</i> cf. <i>rugosa</i> (Hall) [1847, p. 306].	
<i>Salterella</i> .	
<i>Ptychoparia</i> sp.	
<i>Holmia rowei</i> , new species.	
<i>Holmia weeksi</i> , new species.	
4. Massive blue mottled limestone.....	49
5. Mainly green shales, some quartzitic shale, bands of limestone at top.....	580
6. Green calcareous shale, with bands of limestone at top.....	564
<i>Fauna</i> :	
<i>Salterella</i> sp.	
<i>Holmia weeksi</i> , new species.	
<i>Olenellus claytoni</i> , new species.	
7. Andesite mass	750
8. Massive blue mottled limestone.....	81
9. Green calcareous shales.....	238
10. Mostly thin-bedded blue and gray shaly quartzites.....	904
11. Siliceous limestones at base, then blue coral limestones.....	1,349
<i>Holmia weeksi</i> , new species.	
<i>Olenellus</i> , sp.	
12. Massive quartzites, shaly in places.....	222
<i>Fauna</i> :	
<i>Holmia rowei</i> , new species.	
<i>Holmia weeksi</i> , new species.	
13. Siliceous buff limestones.....	180
Total	6,250
Base unknown.	




FIG. 7.—Blacksmith Fork Section

BLACKSMITH FORK SECTION

LOCALITY.—Wasatch Mountains, between Ute and Logan Peaks, in Blacksmith Fork Canyon, east side of Cache Valley, and 12 to 16 miles east of Hyrum, in northern Utah.

This section is 230 miles northeast of the House Range section and north of the greater effect of the pre-Cambrian Uinta Mountain uplift and island. The character of the sediments derived from the Uinta area is shown by the continuation of the arenaceous deposits up to the middle of the Middle Cambrian (Acadian) time, whereas in the House Range section the arenaceous deposits cease before the Middle Cambrian fauna appears. It is not until after the Belt Mountain and Kintla (of the 49th parallel) uplifts to the north are passed that the order of sedimentation, as shown in the Mount Bosworth section, is again of the type of that of the House Range section.

ORDOVICIAN

Feet

- i. Dark, bluish black and gray limestone. In the basal bed immediately above the Cambrian a fine fauna occurs. The limestone is of the same character as that of the Upper Cambrian for 190 feet below and, except for the change in the fauna, there is no break in the section. One of the characters common to the Cambrian and the superjacent Ordovician is the presence in most layers of flattened concretionary nodules and stringers from a minute size up to 6 or 8 cm. or more in diameter; the large ones rarely exceed 3 to 10 mm. in thickness.

Fauna:

Eoorthis coloradoensis (Meek) [1870, p. 425].

Syntrophia nudina Walcott [1905, p. 292].

Orthoceras.

Endoceras.

Fragments of trilobites.

UPPER CAMBRIAN

ST. CHARLES FORMATION [Walcott, 1908a, p. 6]:

1. Dark bluish gray and gray limestone in layers varying from 1 to 20 inches in thickness. Many of the layers are almost made up of flattened concretions varying from a minute size to 6 or 8 cm.

190

Fauna (25 feet below the top):

Lingulella manticula (White) [1874, p. 9].

Eoorthis coloradoensis (Meek) [1870, p. 425].

Syntrophia nudina Walcott [1905, p. 292].

Dicelloccephalus.

ST. CHARLES FORMATION (continued):

1 (continued):

feet

Fauna (105 to 125 feet below the top):

- Schizambon typicalis* Walcott [1884, p. 70].
- Eoorthis coloradoensis* (Meek) [1870, p. 425].
- Eoorthis newberryi* Walcott [1908d, p. 105].
- Syntrophia nudina* Walcott [1905, p. 292].
- Solenopleura*.
- Menocephalus*.
- Illænurus*.

Fauna (20 to 30 feet above base):

- Lingulella (Lingulepis) acuminata* (Conrad) [1839, p. 64].
- Eoorthis coloradoensis* (Meek) [1870, p. 425].
- Eoorthis newberryi* Walcott [1908d, p. 105].
- Agnostus*.
- Solenopleura*.
- Menocephalus*.
- Asaphus* ?

Fragments of fossils occur throughout.

- 2a. Massive bedded, dark lead-gray, arenaceous, cliff-forming limestone, becoming thinner bedded in the lower 50 feet. 195
- 2b. Massive bedded, gray, arenaceous limestone with occasional irregular cherty layers which extend down 85 feet, and just below this the dark, arenaceous limestone is almost made up of round concretions 2 to 4 mm. in diameter for a thickness of about 15 feet. 100
- 2c. Gray, siliceous, and arenaceous limestone in layers one-half inch to 6 inches thick, occurring in massive bands. Light gray chert fills large and small annelid borings, and it also occurs as flattened stringers in the line of the bedding and in the layers. 85

Fauna (34 feet from the base):

- Obolus (Westonia) iphis*, new species.
- Lingulella desiderata* (Walcott) [1898, p. 399].

- 2d. Massive bedded, arenaceous limestone, forming broken cliffs. A few cherty nodules occur near the top and the lower 50 feet has many irregular, oval cherty nodules and stringers of chert coincident with the bedding. 397

Total of 2. 777

- 3. Bedded, bluish gray fossiliferous limestone. 94

Fauna (upper part):

- Acrotreta* sp.
- Anomocare*.

Fauna (near base):

- Obolus*, sp. undt.
- Lingulella manticula* (White) [1874, p. 9].
- Billingsella coloradoensis* (Shumard) [1860, p. 627].
- Agnostus*.
- Ptychoparia*.

ST. CHARLES FORMATION (continued):

	Feet
3 (continued):	
<i>Fauna</i> (a mixture of the faunas at the base and at the top):	
<i>Obolus discoideus</i> (Hall and Whitfield) [1877, p. 205].	
<i>Obolus</i> ? sp. undt.	
<i>Lingulella manticula</i> (White) [1874, p. 9].	
<i>Billingsella coloradoensis</i> (Shumard) [1860, p. 627].	
<i>Huencella lesleyi</i> Walcott [1908 <i>d</i> , p. 110].	
<i>Hyalolithes</i> .	
<i>Cyrtolites</i> .	
<i>Agnostus</i> .	
<i>Ptychoparia</i> .	
<i>Anomocare</i> .	
4. Bedded, light gray sandstone, followed below by dirty brown sandstone, and toward the base shaly and thin-bedded sandstone	166
Strike, north 20° east (magnetic); dip, 25° west.	
<i>Fauna</i> (in upper 20 feet):	
<i>Obolus discoideus</i> (Hall and Whitfield) [1877, p. 205].	
<i>Obolus</i> (<i>Fordinia</i>) <i>bellulus</i> (Walcott) [1905, p. 323].	
<i>Acrotreta idahocensis alta</i> Walcott [1902, p. 588].	
<i>Billingsella coloradoensis</i> (Shumard) [1860, p. 627].	
<i>Fauna</i> (near the base):	
<i>Lingulella</i> (<i>Lingulepis</i>) <i>acuminata</i> (Conrad) [1839, p. 64].	
Total Upper Cambrian.....	1,227

MIDDLE CAMBRIAN

NOUNAN FORMATION [Walcott, 1908*a*, p. 6]:

1 <i>a</i> . Light-gray, arenaceous limestone.....	12
1 <i>b</i> . Lead-colored, arenaceous limestone.....	40
1 <i>c</i> . Light-gray, arenaceous limestone.....	85
1 <i>d</i> . Dark lead-gray, arenaceous limestone.....	87
1 <i>e</i> . Shaly and thin-bedded arenaceous limestone with intercalated reddish brown sandy layers.....	15
1 <i>f</i> . Light-gray, arenaceous limestone.....	18
1 <i>g</i> . Dark lead-gray, arenaceous limestone.....	198
1 <i>h</i> . Light-gray, arenaceous limestone.....	494
1 <i>i</i> . Dark lead-gray, arenaceous limestone, with numerous irregular annelid borings filled with light-gray, arenaceous limestone..	56
1 <i>j</i> . Massive bedded, arenaceous, cherty limestone.....	8
1 <i>k</i> . Bluish gray, cherty, more or less arenaceous limestone in thick bands that break up into thin layers on weathering.....	28
Total of 1	1,041

Fauna:

A few traces of fossils occur in the lower 28 feet and large annelid borings occur in many of the arenaceous limestones. If in a dark rock, the irregular borings are filled with lighter-colored rock, and in the light-gray rock by darker rock.

BLOOMINGTON FORMATION [Walcott, 1908a, p. 7] :

	Feet
1a. Thin-bedded, bluish gray, compact limestone with interbedded thick layers of gray limestone.....	22
<i>Fauna:</i>	
<i>Protospongia</i> (spicules).	
<i>Obolus mcconnelli pelias</i> (Walcott) [1905, p. 330].	
<i>Obolus</i> (<i>Westonia</i>) <i>wasatchensis</i> Walcott [1908d, p. 69].	
<i>Lingulella desiderata</i> (Walcott) [1898, p. 399].	
<i>Hyalithes</i> .	
<i>Agnostus</i> .	
<i>Ptychoparia</i> .	
1b. Greenish argillaceous shale.....	12
1c. Gray, coarse-grained limestone.....	13
Strike, north 20° east; dip, 20° west (magnetic).	
<i>Fauna:</i>	
<i>Hyalithes</i> .	
<i>Ptychoparia</i> .	
1d. Greenish argillaceous and sandy shale.....	147
<i>Fauna</i> (at base) :	
<i>Hyalithes</i> (fragments).	
<i>Agnostus</i> .	
<i>Ptychoparia</i> .	
1e. Gray, coarse-grained limestone.....	4
<i>Fauna:</i>	
<i>Micromitra sculptilis</i> (Meek) [1873, p. 479].	
<i>Hyalithes</i> (abundant).	
<i>Ptychoparia</i> .	
<i>Agraulos</i> .	
1f. Greenish argillaceous and sandy shale.....	22
Total of 1.....	
	220
2a. Bluish gray limestones, with small concretions and small nodules of calcite scattered through the layers, which range from an inch to 6 inches or more in thickness.....	380
<i>Fauna:</i>	
Fragments of fossils.	
2b. Massive bedded, gray limestone that forms a low cliff and breaks down readily on gentle slopes.....	132
<i>Fauna:</i>	
<i>Ptychoparia</i> . } Same as in 1e.	
<i>Agraulos</i> . }	
2c. Bluish gray limestone, with small concretions and small nodules of calcite scattered through the layers; a limestone similar to 2a.....	290

BLOOMINGTON FORMATION (continued):

	Feet
2c (continued):	
<i>Fauna:</i>	
<i>Hyalithes.</i>	
<i>Agraulos.</i>	
2d. Greenish argillaceous shale.....	39
<i>Fauna:</i>	
<i>Obolus (Westonia) wasatchensis</i> Walcott [1908d, p. 69].	
<i>Agraulos.</i>	
<i>Ptychoparia.</i>	
At this horizon in Wasatch Canyon, 5 miles north of Brigham, <i>Acrothele subsidua</i> (White) [1874, p. 6] occurs.	
2e. Bluish gray, thin-bedded limestone.....	182
2f. Arenaceous, steel-gray limestone.....	22
2g. Bluish gray limestone, with small concretions and small nodules of calcite scattered irregularly through the layers...	55
<i>Fauna:</i>	
<i>Micromitra sculptilis</i> (Meek) [1873, p. 479].	
<i>Ptychoparia.</i>	
<i>Dorypyge.</i>	
Total of 2.....	1,100
Total of Bloomington formation.....	1,320

BLACKSMITH FORMATION [Walcott, 1908a, p. 7]:

1a. Dark lead-gray, arenaceous limestone.....	195
1b. Arenaceous, steel gray, cliff-forming limestone, in the lower portion passing gradually into a dove-gray, compact lime- stone that weathers to a light-gray color. The layers vary in thickness from 4 inches to 2.5 feet.....	375
<i>Fauna:</i>	
Fragments of a small trilobite (<i>Ptychoparia</i> ?).	
Annelid borings.	
Total of Blacksmith formation.....	570

UTE FORMATION [Walcott, 1908a, p. 7]:

1a. Bluish gray, compact, thin-bedded limestone, with large irregu- lar annelid borings in the upper part filled with steel-gray, arenaceous limestone similar to the beds above. Below the limestone is purer and more uniformly gray and in layers that tend to form low cliffs on the steeper slopes.....	290
<i>Fauna</i> (in upper part):	
<i>Micromitra (Paterina) labradorica utahensis</i> (Walcott) [1905, p. 306].	
<i>Billingsella</i> , sp. undt.	
<i>Hyalithes.</i>	
<i>Agraulos.</i>	

UTE FORMATION (continued):

1a (continued):

Feet

Ptychoparia subcoronata (Hall and Whitfield) [1877, p. 237].

Dorypyge ? quadriceps (Hall and Whitfield) [1877, p. 240].

130 feet below the top of 1a a large trilobite is indicated by a head and tail.

1b. Gray, arenaceous limestone in thin layers, with occasional bands of layers 4 to 10 inches thick, often oölitic, and with interformational conglomerate and flattened concretions....

135

Fauna (in the upper 5 feet):

Scenella.

Ptychoparia subcoronata (Hall and Whitfield) [1877, p. 237].

Dorypyge ? quadriceps (Hall and Whitfield) [1877, p. 240].

Fauna (in layers 70 to 80 feet below the top):

Micromitra (Paterina) labradorica utahensis (Walcott) [1905, p. 306].

Obolus mcconnelli (Walcott) [1889, p. 441].

Billingsella coloradoensis (Shumard) [1860, p. 627].

Nisusia (Jamesella) nantes (Walcott) [1905, p. 283].

Eoorthis zeno Walcott [1908d, p. 106].

Syntrophia cambria Walcott [1908d, p. 106].

Hyalithes.

Scenella.

Zacanthoides.

Ptychoparia subcoronata (Hall and Whitfield) [1877, p. 237].

Dorypyge ? quadriceps (Hall and Whitfield) [1877, p. 240].

1c. Gray limestone, with numerous concretions one-fourth to one-half inches in diameter. A few thin layers of interformational conglomerate and some shaly limestone.....

58

Total of 1.....

483

2a. Gray, fine-grained, calcareous and argillaceous shaly beds.....

38

Fauna:

Micromitra (Paterina) labradorica utahensis (Walcott) [1905, p. 306].

Obolus (Westonia) ella (Hall and Whitfield) [1877, p. 232].

Acrothele cf. turneri Walcott [1908d, p. 87].

Isoxys cf. argentica (Walcott) [1886, p. 146].

Ptychoparia.

2b. Bluish gray to blue-black, fine-grained, thin-bedded limestone.

57

Fauna:

Obolus ?

Ptychoparia.

UTE FORMATION (continued):

Feet

2c. Greenish argillaceous and calcareous shale, weathering buff...	51
2d. Thin-bedded, grayish-blue limestone.....	36
2e. Gray, oölitic limestone in layers 3 to 14 inches thick.....	24

Fauna:

- Micromitra (Paterina) stuarti* Walcott [1908d, p. 58].
- Micromitra (Paterina) superba* (Walcott) [1897, p. 711].
- Hyalolithes*.
- Ptychoparia a*.
- Ptychoparia b*.
- Dorypyge* (fragment).

2f. Greenish argillaceous and sandy shale.....	18
--	----

Fauna:

- Obolus mcconnelli* (Walcott) [1889, p. 441].
- Micromitra (Paterina) superba* (Walcott) [1897, p. 711].
- Ptychoparia*, sp. undt.

2g. Bluish gray, thin-bedded limestone.....	22
---	----

Strike, north 30° (magnetic); dip, 12° northwest.

Fauna (near base):

- Micromitra (Paterina) superba* (Walcott) [1897, p. 711].
- Hyalolithes*.
- Ptychoparia* (small heads).

Total of 2.....	246
-----------------	-----

Spence shale [Walcott, 1908a, p. 8]:

1. Greenish argillaceous and sandy shale.....	30
---	----

Fauna:

- Micromitra (Iphidella) pannula* (White) [1874, p. 6].
- Obolus (Westonia) ella* (Hall and Whitfield) [1877, p. 232].
- Hyalolithes*.
- Orthotheca major* Walcott [1908c, p. 246, pl. I, fig. 11].
- Leperditia*.
- Ptychoparia*.
- Bathyriscus productus* (Hall and Whitfield) [1877, p. 244].

At Wasatch Canyon, 5 miles north of Brigham, Utah, the following were found at this horizon:

- Eocystites ? longidactylus* Walcott [1886, p. 94].
- Micromitra (Iphidella) pannula* (White) [1874, p. 6].
- Obolus (Westonia) ella* (Hall and Whitfield) [1877, p. 232].
- Lingulella desiderata* (Walcott) [1898, p. 399].
- Acrothele subsidua* (White) [1874, p. 6].
- Agnostus*.
- Ptychoparia piochensis* Walcott [1886, p. 201].
- Zacanthoides idahoensis* Walcott [1908b, p. 26].

Spence shale (continued):

I (continued):

- Neolenus a.*
- Neolenus b.*
- Bathyriscus howelli* Walcott [1886, p. 216].
- Bathyriscus productus* (Hall and Whitfield) [1877, p. 244].
- Ogygopsis.*

LANGSTON FORMATION [Walcott, 1908a, p. 8]:

- | | |
|--|----|
| 1a. Massive bedded, bluish gray limestone, passing downward into gray, arenaceous limestone with many round concretions, one-fourth to three-fourths of an inch in diameter..... | 64 |
|--|----|

Fauna:

- Obolus (Westonia) ella* (Hall and Whitfield) [1877, p. 232].
- Zacanthoides* sp.
- Bathyriscus productus* (Hall and Whitfield) [1877, p. 244]?
- Neolenus* ?

- | | |
|--|----|
| 1b. Massive bedded, bluish gray limestone that breaks up into layers 2 to 8 inches thick on weathering and with many round concretions | 44 |
|--|----|

Fauna:

- Ptychoparia.*
- Bathyriscus productus* (Hall and Whitfield) [1877, p. 244].

In the section two miles southeast of Malade, Idaho, a section which is 60 miles northwest of Blacksmith Fork, the fauna at this horizon is large and finely preserved in compact, bluish gray limestones. It includes:

- Micromitra haydeni* Walcott [1908d, p. 55].
- Micromitra (Iphidella) pannula* (White) [1874, p. 6].
- Micromitra (Iphidella) pannula ophirensis* (Walcott) [1905, p. 306].
- Lingulella desiderata* (Walcott) [1898, p. 399].
- Lingulella helena* (Walcott) [1898, p. 406].
- Lingulella isse* (Walcott) [1905, p. 330].
- Acrotreta idahoensis sulcata* Walcott [1902, p. 588].
- Acrotreta pyxidicula* White [1874, p. 9].
- Acrotreta* ?
- Acrothele artemis* Walcott [1908d, p. 82].
- Acrothele subsidua* (White) [1874, p. 6].
- Acrothele subsidua*, var.
- Acrothyra minor* Walcott [1905, p. 303].
- Billingsella coloradoensis* (Shumard) [1860, p. 627].
- Hyalolithes.*

LANGSTON FORMATION (continued):

Feet

- Orthotheca.*
- Stenotheca.*
- Platyceras.*
- Agnostus.*
- Microdiscus.*
- Solenopleura.*
- Ptychoparia* (2 species).
- Oryctocephalus.*
- Dorypyge* (2 species).
- Neolenus* (2 species).
- Asaphiscus.*
- Ogygopsis?*

- 2. Massive bedded, dark, arenaceous limestone, passing at about 150 feet down into a calcareous sandstone, and then a gray sandstone 390

Total of Langston formation..... 498

BRIGHAM FORMATION [Walcott, 1908a, p. 8]:

- 1a. Quartzitic sandstone, gray, greenish, gray brownish, dirty gray, all weathering reddish dirty brown in layers 3 inches to 3 feet in thickness..... 28
- 1b. Greenish, hard, sandy shale..... 4

Fauna:

- Annelid trails.
- Trilobite tracks.

- 1c. Same as 1a (estimated)..... 1,200+

Total of Brigham formation..... 1,232

Total of Middle Cambrian¹..... 5,420+

RÉSUMÉ, BLACKSMITH FORK SECTION

UPPER CAMBRIAN:

ST. CHARLES FORMATION:

	Feet	Feet
1. Fossiliferous limestone	190	
2. Arenaceous limestone	777	
3. Fossiliferous limestone	94	
4. Shaly and thin-bedded sandstones.....	166	
	—	1,227

MIDDLE CAMBRIAN:

NOUNAN FORMATION:

- 1. Arenaceous limestone

BLOOMINGTON FORMATION:

- 1. Limestone and shales..... 220
 - 2. Thin-bedded limestone
- 1,100
-
- 1,320

¹The line of separation between the Middle and Lower Cambrian occurs somewhere in the Brigham formation, and this thickness (5,420 feet) likely includes several hundred feet of Lower Cambrian beds.

RÉSUMÉ, BLACKSMITH FORK SECTION (continued):

	Feet
BLACKSMITH FORMATION:	
1. Arenaceous limestone	570
UTE FORMATION:	
1. Thin-bedded limestone	483
2. Limestone and shales.....	246
	729
<i>Spence shale</i>	30
LANGSTON FORMATION:	
1. Massive limestone	108
2. Arenaceous limestone	390
	498
BRIGHAM FORMATION:	
1. Quartzitic sandstones (estimated).....	1,232+
Total Middle Cambrian ¹	5,420+
Total section	6,647+

DEARBORN RIVER SECTION

LOCALITY.—North fork of the Dearborn River, south-southeast and south of Mount Dearborn, Lewis and Clark Forest Reserve, Montana.

The base of the section is 4 miles above Walker's ranch at the mouth of the canyon. The summit is the top of Mount Dearborn.

CARBONIFEROUS

	Feet
1. Brown, thin-bedded sandstone.....	135
2a. Massive bedded, light gray, siliceous limestone, forming a high cliff and breaking into talus slopes of small angular fragments	1,970
Fossils: Noted <i>Zaphrentis</i> and <i>Syringopora</i> in abundance.	
2b. Massive bedded dark gray, siliceous limestone breaking into fragments; 275 feet from the top there is a thin bed of shaly limestone	425
2c. Thin-bedded, steel-gray, buff, and gray weathering limestone..	725
Very few fossils were observed. A large <i>Spirifer</i> and specimens of a large <i>Productus</i> were noted about 400 feet from the top.	
Total Carboniferous	3,255

SILURIAN (?)

The strata referred to the Silurian are arenaceous limestones in which no fossils were observed. It is not improbable that the upper portion of them may be of Devonian age, or possibly Lower Carboniferous.

¹ See footnote on page 199.

	Feet
3a. Massive bedded, gray, arenaceous cliff-forming limestones. . . .	675
3b. Thin-bedded, gray limestone and sandstone, with small lentiles of light-gray limestone. The irregular arenaceous portions weather buff and form a buff band where the rock is in the cliffs	75
3c. Massive bedded, light gray, arenaceous limestone, with thinner bedded, purer limestones at intervals. On the mountain slopes the massive beds form cliffs, and the thin-bedded more shaly portion forms slopes.	245
3d. Massive bedded, light gray, arenaceous limestone, with somewhat purer, slightly banded limestones toward the top.	175
3e. Massive bedded light gray, banded limestones, that break up into thin and often shaly layers on exposure to the weather.	51
3f. Massive bedded, light gray, arenaceous limestones.	164
Total of Silurian?	
	1,385

The line drawn between the Cambrian and the Silurian is based largely on the change in the character of the limestone, from the coarse, gray, arenaceous limestone to a much purer, gray limestone, and the occurrence, about 150 feet from the top, of fragments of a species of *Ptychoparia*.

CAMBRIAN

LIMESTONE:

	Feet
1a. Massive bedded, hard, gray and bluish gray limestones that break up into thin, irregular layers on exposure to the weather. The color of many thin layers and the thick layers on their bedding planes is yellow to buff. The upper 100 feet contain massive dove-colored limestones and near the top a few feet of siliceous limestone.	550
Fragments of a species of <i>Ptychoparia</i> were noted about 150 feet below the summit.	
1b. Greenish and gray, thin-bedded limestone, with some arenaceous shale and thin layers of greenish sandstone in the shale. Numerous annelid trails and fragments of trilobites occur throughout.	90
1c. Massive bedded, gray limestone that breaks up into thin, irregular layers, in very much the same manner as the Pilgrim limestone, but is usually more massive. It is quite arenaceous near the central portions, where it is more massive bedded for a short distance.	680
One hundred and seventy feet from the top there is a band of thin-bedded oölitic limestone, in which fragments of trilobites are numerous; also a small <i>Obolus</i> -like shell. Oölitic limestone, interbedded with irregular, thin-bedded, bluish gray limestones, occurs in the lower 170 feet.	
Total of limestone.	
	1,320

SHALE:

- | | |
|--|------|
| | Feet |
| 2. Thin-bedded limestones, with partings of greenish, argillaceous, and arenaceous shale. Sometimes the shale and at other times the limestone predominates..... | 150 |

LIMESTONE:

- | | |
|--|-----|
| 3. Massive bedded, gray limestone, similar to the Meagher limestone, except that it is of a lighter gray color near the top.. Annelid trails are abundant and fragments of trilobites. | 130 |
|--|-----|

SHALE:

- | | |
|--|-----|
| 4. Thin-bedded limestones, with partings of greenish, argillaceous, and arenaceous shale, the limestones predominating. It breaks down readily on the slopes and forms a sloping terrace | 210 |
|--|-----|

LIMESTONE:

5. Massive bedded, fine-grained, gray limestone that breaks up on weathering into thin layers from a quarter of an inch to two inches in thickness. They have a very irregular surface, marked by a thin, buff-colored deposit that fills the annelid burrows and trails, and also occurs as irregular blotches on the surface.

This belt of limestone is divided into five thick beds that may be distinguished for miles in the cliffs. The two lower are usually broken down.....

Annelid trails are abundant and numerous fragments of trilobites.

55

SHALE:

- | | |
|--|-----|
| 6. Greenish purple and dark gray, argillaceous shales, with thin layers of sandstone and arenaceous shale at irregular intervals | 190 |
|--|-----|

Shale No. 6 is in the same stratigraphic position as the Wolsey shale [Weed, 1900, p. 285] of the Little Belt Mountains section, and the sandstone beneath corresponds stratigraphically to the Flat-head sandstone [Peale, 1893, p. 20] in the same section. The fauna of shale No. 6 on Scapegoat and Gordon mountains, localities west of the Dearborn River section, is, however, entirely unlike that of the Middle Cambrian Wolsey shale, and includes the following species:

Micromitra (Iphidella) pannula (White) [1874, p. 6].

Obolus (Westonia) ella (Hall and Whitfield) [1877, p. 232].

Acrothele colleni, new species.

Acrothele panderi, new species.

Wimanella simplex Walcott [1908d, p. 101].

Olenopsis ? sp.

Ptychoparia, sp.

Albertella helena Walcott [1908b, p. 19].

Vanuxemella contracta, new genus and new species.

Bathyriscus productus ? (Hall and Whitfield) [1877, p. 244].

Bathyriscus ?.

This fauna is strikingly similar to that occurring in the drift blocks which are believed to have come from the lower portion of the Mount Whyte formation of the Mount Bosworth section [see page 214]. At the localities in question neither fauna contains *Olenellus*, but that genus occurs so generally in the Mount Whyte formation, both above and below the Albertella horizon, that the entire formation is placed in the Lower Cambrian. This correlation places shale No. 6 and sandstone No. 7 in the Lower Cambrian.

SANDSTONE:

	Feet
7a. Thin-bedded sandstones, with partings of dark arenaceous shale Many varieties of annelid trails and tracks of trilobites occur on the surface of the sandstone.	70
7b. Massive bedded, coarse, more or less cross-bedded, light gray sandstone, with a few thin layers of fine quartzitic con- glomerate	80
Total of sandstone.....	150

RÉSUMÉ, DEARBORN RIVER SECTION

	Feet
1. Limestone.....	1,320
2. Shale.....	150
3. Limestone.....	130
4. Shale.....	210
5. Limestone.....	55
6. Shale.....	190
7. Sandstone.....	150
Total of Cambrian.....	2,205

Beneath the Cambrian sandstone the Empire shales of the Belt Terrane of the Algonkian occur with apparently the same strike and dip as the base of the sandstone. Traced on the strike, however, they appear to be unconformably beneath the sandstone.

MOUNT BOSWORTH SECTION

Mount Bosworth section, north of Hector, British Columbia, on the Continental Divide, north of the Canadian Pacific Railway.

The summit of the section is on the west spur (Sherbrooke ridge) of Mount Bosworth overlooking Sherbrooke Lake. The highest beds are on the south summit of the ridge, and from their lithologic character and the finding of obscure fossils that suggest *Ophileta* of the Lower Ordovician the upper 110 feet of strata are tentatively referred to the Ordovician system. The strata near the summit are much broken up owing to a fault line that crosses the ridge.

ORDOVICIAN

- | | Feet |
|---|------|
| 1. Massive bedded gray and bluish gray arenaceous limestone, with thin layers, irregular stringers, and nodules of dark chert | 110 |

UPPER CAMBRIAN

SHERBROOKE FORMATION [Walcott, 1908a, p. 2]:

- | | |
|---|-----|
| 1. Massive bedded, bluish gray limestone, with some cherty matter in the form of small nodules and stringers; also irregular partings and fillings of annelid borings by gray dolomitic limestone weathering buff. | 175 |
|---|-----|

Fauna:

Annelid borings and trails.
Fragments of undeterminable trilobites.

- | | |
|---|-----|
| 2a. Gray oölitic limestone in thick layers, with bluish banded limestone intercalated at irregular intervals. The banded appearance of the nonoölitic layers is owing to the buff weathering of the thin dolomitic layers. | 190 |
|---|-----|

Fauna (Upper Cambrian facies):

Crepicephalus.
Pterocephalus?
Ptychoparia.

- | | |
|--|-----|
| 2b. Greenish drab and gray siliceous shales with interbedded oölitic limestone in bands of layers from 6 inches to 4 feet thick; also a few bands of thick-bedded, bluish gray limestone that breaks up into shaly limestone on weathering. | 335 |
|--|-----|

Fauna (in green shales near summit):

Lingulella isse (Walcott) [1905, p. 330].

Fauna (in oölitic layers):

Agnostus, sp. undt.
Illænurus.
Ptychoparia.

SHERBROOKE FORMATION (continued):

	Feet
2c. Gray oölitic limestone, with thin bands of interbedded shaly, blue gray limestone. Gray, dolomitic, buff-weathering, flattened nodules, stringers, and thin layers of limestone occur in a very irregular manner.....	65

Fauna:

Illænenurus.

Agnostus.

Ptychoparia.

Bathyrurus-like pygidia.

Total of 2..... 590

3. Arenaceous, dolomitic, steel gray limestone weathering light gray and buff gray.....	610
---	-----

The line of demarcation between 3 and the bluish gray limestones below is irregular. The gray beds of 3 extend along the cliff and abruptly change to bluish gray. In the upper 100 feet of 3 irregular masses of bluish gray limestone occur like great lentiles, as though they were cores left in the general alteration (dolomitization) of the strata.

Total of Sherbrooke formation..... 1,375

PAGET FORMATION [Walcott, 1908a, p.3]:

1. Massive bedded, dark bluish gray limestone forming base of cliff on the west side of the amphitheater on the west slope of Mount Bosworth and, with 3 of Sherbrooke formation, the upper cliffs of Paget Peak and Mount Daly.....	60
2. Massive beds of oölitic limestone, with irregular, interbedded bands of green siliceous shale. Thin layers, irregular stringers, and nodules of gray buff weathering dolomite occur in the oölitic limestones.....	300+

The base of 2 is covered by talus slope on line of the section. It is well exposed on the southeast face of Mount Daly and Paget Peak. The thickness is placed at 300 feet, which I think is less than the total thickness. Over 200 feet was measured.

Fauna:

Hyalithes.

Agnostus.

Crepicephalus.


Total of Paget formation..... 360+

BOSWORTH FORMATION [Walcott, 1908a, p.3]:

1. Massive bedded, gray, and bluish gray arenaceous dolomitic limestone. Several bands of steel gray, yellowish buff weathering bands of strata occur in the lower half of 1.....	600+
---	------

		Feet	Me- ters	
Ordovician				Shaly and thin bedded limestone
Upper Cambrian	Sherbrooke	365	111	Gray limestone
		400	122	Thin bedded limestone
		610	186	Arenaceous limestone
	Paget	360	109	Gray and oolitic limestone
		600	183	Arenaceous limestone
	Bosworth	1255	383	Thin bedded limestone
Middle Cambrian	Eldon	410	125	Gray and bluish gray limestone
		110	34	Thin bedded limestone
		265	82	Siliceous limestone
		< 95	29	Thin bedded limestone
		1845	562	Arenaceous limestone

FIG. 8.—Mt. Bosworth Section (continued on following page)



EASTERN SIDE OF SHERBROOKE RIDGE

The summit of the ridge at (a) is formed of strata tentatively referred to the Ordovician. Below, the Sherbrooke and Paget formations extend down to the foot of the cliffs. Buff Point and Red Knob expose the Bosworth formation nearly down to the great Eldon siliceous limestone. A fault brings the base of the Paget formation up about 500 feet.




FIG. 9.—Mt. Bosworth Section (continued)

NOTE.—The thickness of the St. Piran, Lake Louise, and Fairview formations is taken from the Lake Louise section.

BOSWORTH FORMATION (continued):

1 (continued):	Feet
This formation forms the base of the high cliffs on the south-east face of Mount Daly and Paget Peak.	
The lower portion of 1 was measured and the upper parts estimated. The thickness given is probably 100 feet or more less than the actual thickness.	
2a. Shaly and thin-bedded, gray and dove-colored, compact fine-grained dolomitic limestone weathering buff and light gray. Thicker layers occur in bands from 1 to 6 feet thick.....	422
2b. Greenish siliceous shale with thin interbedded layers of siliceous, compact, gray limestone.....	48
<i>Fauna:</i>	
At about this horizon in the Castle Mountain section 20 miles southeast of Mount Bosworth small trilobite heads of the genera <i>Ptychoparia</i> and <i>Solenopleura</i> occur in a band of gray and bluish black limestone, and just below fragments of a species of <i>Obolus</i> .	
2c. Limestones similar to 2a.....	517
Total of 2.....	987
3. Variable arenaceous shales with alternating bands of color—greenish, deep red, buff, yellow, and gray. Numerous mud cracks and ripple-marks occur on many of the layers.....	268
Total of Bosworth formation.....	1,855
Total Upper Cambrian.....	3,590

NOTE.—The 1,855 feet of strata included in 1, 2, and 3 remind me, in lithologic character and appearance, of strata of the upper portions of the Cambrian Belt Terrane of Montana. No traces of life were observed and the shaly, banded character of the beds is very striking.

MIDDLE CAMBRIAN

ELDON FORMATION [Walcott, 1908a, p.3]:

1a. Irregularly bedded, gray, siliceous and arenaceous limestone in thick layers above and thin layers below; at 192 feet from the base a bed of bluish black limestone is fossiliferous. Above the fossiliferous bed the strata become more massive, arenaceous, steel gray in color, weathering to a light gray...	410
<i>Fauna</i> (192 feet above the base):	
<i>Agnostus</i> , sp.	
<i>Ptychoparia</i> , 2 species.	
<i>Bathyriscus</i> -like pygidium.	
1b. Light and dark gray, thin-bedded, arenaceous limestone, weathering to a light-gray color.....	110
1c. Massive bedded, siliceous, fine-grained, compact, dark bluish gray limestone.....	197



Fig. 1. NORTH RIDGE OF CASTLE MOUNTAIN

Showing the Eldon formation in the cliffs above the lake and the Bosworth formation in the snow-covered points above the cliff line.




Fig. 2. PROFILE OF SOUTHEAST FRONT OF CASTLE MOUNTAIN, OPPOSITE ELDON

The upper cliff is formed of the siliceous limestone of the Eldon formation; the terrace with snow on it the Stephen formation, and the lower cliff and slope the Cathedral formation. These formations are finely exposed on Mount Bosworth, but not so as to get good photographs of them.

ELDON FORMATION (continued):

1c (continued):

Feet

Two yellowish buff weathering bands of limestone 2 to 3 feet thick stand out in color on the face of cliffs.

Fauna (near the summit):

Billingsella ?

Neolenus-like pygidium.

1d. Massive bedded limestone much like that of 1c.....	71
Total of 1.....	788
2. Thin-bedded, bluish gray limestone with irregular layers and stringers of gray, buff weathering, dolomitic limestone.....	95
At 24 feet from the base a shaly, bluish gray, siliceous limestone about two feet thick is interbedded.	
<i>Fauna</i> (in shaly limestone):	
<i>Obolus membranaceous</i> Walcott [1908d, p. 61].	
<i>Lingulella</i> sp.	
<i>Isoxys argentca</i> (Walcott) [1886, p. 146].	
<i>Ptychoparia</i> , 2 species.	
3. Massive bedded dark gray arenaceous limestone.....	190
4. Massive bedded, cliff-forming, light gray arenaceous limestone. At several horizons bands of thinner layers from a few feet up to 30 feet in thickness occur. One of these 480 feet from the base forms a slight terrace.....	1,655

Fauna:

In the Mount Stephen section seven miles southwest of Mount Bosworth, at a horizon about 700 feet above the base of this limestone, the following fossils have been recognized:

Protospongia (spicules).

Lingulella cf. *isse* (Walcott) [1905, p. 330].

Hyalithes sp.

Agnostus cf. *montis* Matthew [1899, p. 43].

Zacanthoides spinosus Walcott [1884, p. 63].

Ptychoparia sp.

Bathyuriscus sp.

Ogygopsis sp.

Total of Eldon formation..... 2,728

STEPHEN FORMATION [Walcott, 1908a, p. 3]:

1. Thin-bedded, dark gray and bluish black limestone.....	315
---	-----

Fauna:

Micromitra (*Paterina*) *stissingensis* (Dwight) [1889, p. 145].

Obolus mconnelli (Walcott) [1889, p. 441].

STEPHEN FORMATION (continued):

I (continued):

Nisusia alberta (Walcott) [1889, p. 442], var.

Hyolithes carinatus Matthew [1899, p. 42].

Agnostus sp.

Agraulos sp.

Menocephalus sp.

Ptychoparia, 3 species.

Neolenus sp.

Bathyriscus sp.

At Mount Stephen, about seven miles southwest of Mount Bosworth, a siliceous shale occurs at the summit of the Stephen formation in which an unusually rich fauna occurs. This shale is not well developed on Mount Bosworth.

OGYGOPHIS SHALE.—This term is applied to the local development of arenaceous and calcareous shale at the summit of the Stephen formation on the northwest slope of Mount Stephen. The shale band (lentile) has a maximum thickness of about 150 feet. It thins out to the northeast and is faulted out to the southwest. At its maximum thickness, 2,800 feet above Field, it carries immense numbers of trilobites, especially *Ogygopsis klotzi* (Rominger), *Bathyriscus rotundatus* (Rominger), *Neolenus serratus* (Rominger), *Zacanthoides spinosus* (Walcott), and, in addition, sponges, cystids, brachiopods, pteropods, and gasteropods. The shale is less rich in fossils one-fourth of a mile northeast on the strike; also to the northwest. Lentiles of gray quartzitic sandstone and siliceous, gray limestone occur in the shale, and the entire shale band appears to be a lentile between the thin-bedded blue limestones and the superjacent massive, arenaceous limestone formation. There is no trace of the *Ogygopsis* shale on Mount Bosworth 6 miles northeast, at the same horizon, or at Castle Mountain, 20 miles east-southeast.

There is a sharp anticline, with a northeast-southwest axis, in the shale and the thin-bedded limestones beneath, on the northwest slope of Mount Stephen. The southeast limb is crushed and the beds are largely faulted out against the massive arenaceous limestone before reaching the amphitheater at the head of Field Brook. On the northwest limb the shales are unaltered and slope down the side of the mountain for 1,800 feet, thus affording a great exposure of the shale and contained fossils.

Fauna:

1. *Hyolithellus flagellum* (Matthew) [1899, p. 40].
2. *Hyolithellus annulatus* (Matthew) [1899, p. 42].
3. *Orthotheca corrugata* Matthew [1899, p. 42].
4. *Orthotheca major* Walcott [1908c, p. 246, pl. I, fig. 11].
5. *Hyolithes*, sp.
6. *Hyolithes carinatus* Matthew [1899, p. 42].
7. *Stenotheca wheeleri* Walcott [1908c, p. 245, pl. I, fig. 7].
8. *Platyceras romingeri* Walcott [1889, p. 442].
9. *Platyceras bellianus* Walcott [1908c, p. 246, pl. I, fig. 13].
10. *Acrotreta depressa* (Walcott) [1889, p. 441].
11. *Micromitra (Iphidella) pannula* (White) [1874, p. 6].
12. *Obolus mcconnelli* (Walcott) [1889, p. 441].
13. *Nisusia alberta* Walcott [1889, p. 442].
14. *Philhedra columbiana* (Walcott) [1889, p. 441].



MOUNT STEPHEN, BRITISH COLUMBIA, FROM THE NORTH

Near the base at (a) the Mount Whyte formation rests on the St. Piran quartzitic sandstones. The great Cathedral arenaceous limestone forms the north shoulder of the mountain up to (b), where the 800 feet of the Stephen formation breaks the profile. Above this the massive beds of the Eldon formation extend to the summit of the peak. The section shown in the profile is over 5,800 feet in thickness. At *xx* on the slope the great fossil beds of the Stephen formation are finely exposed.

STEPHEN FORMATION (continued):

Ogygopsis shale (continued):	Feet
15. <i>Scenella varians</i> Walcott [1886, p. 127].	
16. <i>Anomalocaris canadensis</i> Whiteaves [1892, p. 207].	
17. <i>Anomalocaris</i> ? <i>whiteavesi</i> Walcott [1908c, p. 246, pl. II, figs. 2, 2a, 4, 6, and 6a].	
18. <i>Anomalocaris</i> ?? <i>acutangula</i> Walcott [1908c, p. 247, pl. II, fig. 5].	
19. <i>Agnostus montis</i> Matthew [1899, p. 43].	
20. <i>Dorypyge</i> (<i>Kootenia</i>) <i>dawsoni</i> (Walcott) [1889, p. 446].	
21. <i>Bathyriscus rotundatus</i> (Rominger) [1887, p. 16].	
22. <i>Bathyriscus pupa</i> Matthew [1899, p. 51] probably = 23.	
23. <i>Bathyriscus occidentalis</i> (Matthew) [1899, p. 49].	
24. <i>Bathyriscus ornatus</i> Walcott [1908b, p. 39].	
25. <i>Karlia stephenensis</i> Walcott [1889, p. 445].	
<i>Corynexochus romingeri</i> Matthew [1899, p. 47] = 25.	
26. <i>Neolenus serratus</i> (Rominger) [1887, p. 13].	
<i>Neolenus granulatus</i> Matthew [1899, p. 55] = 26.	
27. <i>Ogygopsis klotzi</i> (Rominger) [1887, p. 12].	
28. <i>Oryctocephalus reynoldsi</i> Reed [1899, p. 359].	
<i>Oryctocephalus walkeri</i> Matthew [1899] = 28.	
29. <i>Burlingia hectori</i> Walcott [1908b, p. 15].	
30. <i>Ptychoparia cordillerae</i> (Rominger) [1887, p. 17].	
<i>Conocephalites</i> cf. <i>perseus</i> Hall, Matthew [1899, p. 46] = 30.	
31. <i>Ptychoparia palliseri</i> Walcott [1908c, p. 247, pl. III, fig. 6].	
32. <i>Zacanthoides spinosus</i> (Walcott) [1884, p. 63].	
2a. Greenish siliceous shale	23
Fauna:	
<i>Obolus</i> (<i>Westonia</i>) <i>ella</i> ? (Hall and Whitfield) [1877, p. 232].	
2b. Thick-bedded, bluish gray limestone, breaking up into thin layers one-half to 3 inches thick on weathering	22
Fauna:	
<i>Micromitra</i> (<i>Paterina</i>) <i>stissingensis</i> (Dwight) [1889, p. 145].	
<i>Nisusia alberta</i> Walcott [1889, p. 442], var.	
2c. Greenish siliceous shale	70
2d. Alternating bluish gray, bedded, compact limestone, siliceous and arenaceous shale, mostly shale below	210
Total 2	325
Fauna:	
<i>Cruziana</i> .	
<i>Micromitra</i> (<i>Iphidella</i>) <i>pannula</i> (White) [1874, p. 6].	
<i>Obolus</i> (<i>Westonia</i>) <i>ella</i> (Hall and Whitfield) [1877, p. 232].	
<i>Hyalolithes</i> .	
<i>Leperditia</i> .	
<i>Ptychoparia</i> .	
<i>Bathyriscus</i> .	

STEPHEN FORMATION (continued):

2d (continued):

Feet

On Mount Stephen, at a horizon 150 feet from the base of this limestone, the fauna includes:

Micromitra (Iphidella) pannula (White) [1874, p. 6].

Billingsella marion Walcott [1908d, p. 102].

Hyalithes.

Microdiscus.

Ptychoparia.

CATHEDRAL FORMATION [Walcott, 1908a, p. 4]:

1a. Thin-bedded gray to lead-gray, arenaceous limestones, weathering buff gray to dull light gray.....	404
1b. Massive bedded, steel-gray weathering, light gray, arenaceous limestone. In some localities thinner layers appear at various horizons and large lentiles of dark lead-gray-colored beds occur very irregularly.....	682
1c. Similar to 1a. Annelid borings and trails occur in and on some of the layers.....	126
1d. Similar to 1b.....	83
1e. Thin-bedded, lead-gray to blue-gray, thin-bedded (layers 1 inch to 4 inches thick) arenaceous limestone.....	25
1f. Alternating thin and massive bedded, arenaceous, steel-gray limestone weathering light gray.....	275
Total of 1.....	1,595

LOWER CAMBRIAN

MOUNT WHYTE FORMATION [Walcott, 1908a, p. 4]:

The line between the Middle and Lower Cambrian is placed at this horizon on account of the presence in the Mount Stephen section of *Olenellus* in the limestone 116 feet below the massive arenaceous limestone belt represented by 1f in the Cathedral formation of the Mount Bosworth section.

1a. Thin-bedded, bluish gray, slightly arenaceous limestone.....	120
<i>Fauna:</i>	
Numerous annelid trails and borings.	
1b. Gray oölitic limestone in layers 3 to 6 inches thick.....	44
<i>Fauna</i> (4 feet from base):	
<i>Nisusia (Jamesella) lowi</i> Walcott [1908d, p. 98].	
<i>Microdiscus</i> , sp. undt.	
<i>Agraulos</i> sp.	
<i>Ptychoparia</i> sp.	

At Castle Mountain fragments of a species of *Bornemannia* (new genus allied to *Zacanthoides*) occur at about this horizon.

MOUNT WHYTE FORMATION (continued):

1b (continued):

Feet

In the Mount Stephen section the following species occur at a horizon near the top of this limestone:

- Nisusia (Jamesella) lowi* Walcott [1908d, p. 98].
Stenotheca elongata Walcott [1884, p. 23], var.
Scenella varians Walcott [1886, p. 127].
Platyceras, new species.
Hyolithes billingsi Walcott [1886, p. 134].
Ptychoparia sp.
Crepicephalus, new species.
Protypus, new species.
Albertella, sp. undt.

About 50 feet down in the Mount Stephen section in a gray, siliceous shale the following species occur:

- Cystid plates.
Micromitra (Paterina), sp. undt.
Acrotreta sagittalis taconica (Walcott) [1887, p. 189].
Nisusia (Jamesella) lowi Walcott [1908d, p. 98].
Hyolithes (fragment).
Hyolithellus cf. *micans* Billings [1872, p. 215].
Scenella varians Walcott [1886, p. 127].
Olenellus (fragments of thoracic segments).

- 1c. Massive layers made up of banded bluish gray limestone and sandstone in layers one-half inch to 2 inches thick. 60

Fauna:

Agraulos, sp. undt.

Total of 1. 224

On Mount Stephen, at a horizon near the top of this bed of limestone, there was found:

- Acrothele colleni*, new species.
Acrotreta sagittalis taconica (Walcott) [1887, p. 189].
Scenella varians Walcott [1886, p. 127].
Stenotheca elongata Walcott [1884, p. 23], var.
Albertella, sp. undt.
Olenellus (fragments).
Bathyriscus, sp. undt.

Near the base on Mount Stephen:

- Micromitra (Paterina) labradorica* (Billings) [1861, p. 6], var.
Micromitra (Iphidella) pannula (White) [1874, p. 6].
Acrotreta sagittalis taconica (Walcott) [1887, p. 189].
Bornemannia prima, new genus and new species.
Ptychoparia, 3 species.

2. Gray and brownish gray sandstone in thin and massive layers. 31

MOUNT WHYTE FORMATION (continued):

2 (continued):

Feet

Fauna:

Hyalithes.*Agraulos*.

On Mount Stephen, at this horizon, there were found:

Microdiscus, sp. undt.*Olcuellus*, sp. undt. (fragments).*Ptychoparia*, sp. undt.*Protypus*, sp. undt.

3. Siliceous shale with a few interbedded thin layers of compact, hard, gray sandstone.....

115

In the Lake Agnes section 5 miles southeast of Mount Bosworth, the fauna of about this horizon includes:

Micromitra (Paterina) wapta Walcott [1908d, p. 59].*Obolus parvus* Walcott [1908d, p. 61].*Hyalithes billingsi* Walcott [1886, p. 134].*Olenopsis agnes*, new species.*Ptychoparia*, 3 species.*Albertella*, sp. undt.*Bathyriscus*.

On the south slope of Mount Bosworth two drift blocks of siliceous shale, supposed to be from this horizon, were found, from which the following species were collected:

Micromitra (Paterina) wapta Walcott [1908d, p. 59].*Obolus parvus* Walcott [1908d, p. 61].*Acrothelc colleni*, new species.*Wimanella simplex* Walcott [1908d, p. 101].*Agraulos*, sp.*Ptychoparia*, sp.*Bornemannia*, sp.*Albertella bosworthi* Walcott [1908b, p. 22].*Albertella helena* Walcott [1908b, p. 19].*Vanuxemella contracta*, new genus and new species.*Bathyriscus*, sp. a.

On Mount Stephen, at about the same horizon, the following were found:

Hyalithes billingsi Walcott [1886, p. 134].*Scenella varians* Walcott [1886, p. 127].*Olenopsis agnes*, new species.*Bornemannia prima*, new genus and new species.

4. Interbedded layers of gray fossiliferous limestone and greenish gray siliceous shale.....

20

Fauna:

Nisusia festinata (Billings) [1861, p. 10].*Scenella varians* Walcott [1886, p. 127].*Hyalithellus*.

MOUNT WHYTE FORMATION (continued):

Feet

4 (continued):

Ptychoparia.*Agraulos*.*Protypus fieldensis*, new species.*Olenellus canadensis*, new species.

At this horizon on Mount Stephen the following were found:

Micromitra (Iphidella) pannula (White) [1874, p. 6].*Acrotreta sagittalis taconica* (Walcott) [1887, p. 189].*Kutorgina cingulata* (Billings) [1861, p. 8].*Kutorgina*, sp. undt.*Nisusia festinata* (Billings) [1861, p. 10].*Hyalolithes billingsi* Walcott [1886, p. 134].*Scenella varians* Walcott [1886, p. 127].*Protypus*, new species.*Agraulos*, sp. undt.*Ptychoparia*, 3 sp. undt.*Olenellus canadensis*, new species.

BOW RIVER GROUP

ST. PIRAN FORMATION [Walcott, 1908a, p. 4]:

1a. Siliceous and arenaceous greenish-colored shales in layers 1 to 3 inches in thickness, interbedded in shaly and thin-bedded gray and brownish gray sandstone, with a thick layer of compact, gray sandstone near the top. 68

1b. Irregularly bedded brownish, dirty gray, and occasionally purplish-colored sandstones, more or less compact and quartzitic and in massive and thin layers that break down readily on slopes 310

Fauna:

Annelid trails and borings (*Scolithus*).*Hyalolithes*.*Olenellus canadensis*?, new species.*Ptychoparia* (2 species).

1c. Massive bedded, compact, light gray and pinkish quartzitic sandstones 125

Fauna:

Annelid trails and borings (*Scolithus*).*Hyalolithes*.*Olenellus canadensis*?, new species (fragments).

The general dip of the strata is to the northwest 20°; strike, north 30° east. The section is continuous, with the exception of the displacement between the Paget and Bosworth formations of the Upper Cambrian. This does not cut out any considerable thickness of strata, as is proven by the unbroken section in the cliffs of Mount Daly three miles to the north.

In the Lakes Louise and Agnes section, about five miles southeast of Mount

Bosworth, the total thickness of the St. Piran formation is 2,705 feet. Below the St. Piran the following section occurs:

LAKE LOUISE FORMATION [Walcott, 1908a, p. 5]:

	Feet
1. Compact, gray, siliceous shale.....	105

Fauna:

Annelid trails.

Cruziana.

Micromitra (Iphidella) louise Walcott [1908d, p. 56].

FAIRVIEW FORMATION [Walcott, 1908a, p. 5]:

1. Thin and thick layers of gray, quartzitic, brownish weathering, compact sandstones (estimated).....	600+
This formation is much thicker to the southeast.	

RÉSUMÉ, MOUNT BOSWORTH SECTION

UPPER CAMBRIAN.

SHERBROOKE FORMATION:

	Feet	Feet
1. Gray, partly cherty limestones.....	175	
2. Oölitic limestones and shaly band.....	590	
3. Arenaceous dolomitic limestone.....	610	
Total	1,375	

PAGET FORMATION:

1. Massive bedded bluish gray limestone.....	60	
2. Oölitic limestone with bands of shale.....	300+	
Total	360+	

BOSWORTH FORMATION:

1. Gray, arenaceous, dolomitic limestone.....	600+	
2. Shaly and thin-bedded, dolomitic limestones with two bands of shale.....	987	
3. Shales	268	
Total	1,855+	
Total Upper Cambrian.....	3,590+	

MIDDLE CAMBRIAN.

ELDON FORMATION:

1. Siliceous and arenaceous limestone.....	788	
2. Bluish gray limestone.....	95	
3 and 4. Arenaceous limestone	1,845	
Total	2,728	



PROFILE VIEW OF RIDGES SOUTHEAST (IN THE DISTANCE) AND WEST (FOREGROUND) OF LAKE LOUISE

The distant profile shows the Fairview formation at the base of Fairview Mountain. The Lake Louise formation is at (a), and from (a) to about (b) the St. Piran quartzitic sandstones. The Mount Whyte and Cathedral formations form the summits of the distant peaks. The quartzitic sandstones of the St. Piran formation are well shown on the ridge in the foreground.

RÉSUMÉ, MOUNT BOSWORTH SECTION (continued):

STEPHEN FORMATION:

	Feet	Feet
1. Thin-bedded, dark and bluish gray limestone.....	315	
2. Alternating limestones and shales.....	325	
	<hr/>	
Total		640

CATHEDRAL FORMATION:

1. Arenaceous dolomitic limestone.....	1,595	
	<hr/> <hr/>	
Total Middle Cambrian.....		4,963

LOWER CAMBRIAN.

MOUNT WHYTE FORMATION:

1. Thin-bedded limestones	224	
2. Sandstone	31	
3. Siliceous shale	115	
4. Gray limestone	20	
	<hr/>	
Total		390

ST. PIRAN FORMATION:

1. Sandy shales and quartzitic sandstones as exposed at Lake Agnes	2,705	
---	-------	--

LAKE LOUISE FORMATION:

1. Compact siliceous shale as exposed at Lake Louise.....	105	
---	-----	--

FAIRVIEW FORMATION:

1. Quartzitic sandstones as exposed at Lake Louise.....	600+	
	<hr/>	
Total Lower Cambrian.....		3,800+
		<hr/> <hr/>

Upper Cambrian	3,590+
Middle Cambrian	4,963
Lower Cambrian	3,800+
	<hr/>
Total thickness of sections examined.....	12,353+

Below the section of the quartzitic sandstones on Fairview Mountain there is, in the Bow River valley, a considerable, but unknown, thickness of sandstones and siliceous shales that have not been examined or measured.

BIBLIOGRAPHY.

BILLINGS, E.

1861. Geological Survey of Canada, Paleozoic Fossils, I, 1861 (November), pp. 1-24.
1872. The Canadian Naturalist and Quarterly Journal of Science, new (2d) series, VI, No. 2, 1872, pp. 213-222: On some new species of Paleozoic Fossils.

CONRAD, T. A.

1839. Third Annual Report New York State Survey (Printed as New York State Assembly Document, No. 275), 1839, (February 27), pp. 57-66: Second Annual Report of T. A. Conrad.

DWIGHT, W. B.

1889. American Journal of Science, 3d series, XXXVIII, 1889 (August), pp. 139-153: Recent explorations in the Wappinger Valley limestones and other formations of Dutchess Co., N. Y.

HALL, J.

1847. Natural History of New York, Paleontology, I, 1847: 4to, Albany, N. Y.

HALL, J., and WHITFIELD, R. P.

1877. United States Geological Exploration of the Fortieth Parallel, IV, 1877; Pt. 2, Paleontology, pp. 198-302.

MATTHEW, G. F.

1899. Proceedings and Transactions of the Royal Society of Canada for 1899, 2d series, V, 1899, Sec. 4, No. 2, pp. 39-66: Studies on Cambrian Faunas, No. 3.—Upper Cambrian Fauna of Mt. Stephen, British Columbia.—The Trilobites and Worms.

MEEK, F. B.

1868. American Journal of Science and Arts, 2d series, XLV, 1868 (January), pp. 62-64: Preliminary notice of a remarkable new genus of Corals, probably typical of a new family.
1870. Proceedings of the American Philosophical Society held at Philadelphia, XI, 1870, No. 84, pp. 425-431: A preliminary list of Fossils, collected by Dr. Hayden in Colorado, New Mexico, and California, with descriptions of new species.
1873. Sixth Annual Report of the United States Geological Survey of Montana, Idaho, Wyoming, and Utah for 1872, 1873, pp. 429-518: Preliminary Paleontological Report.

PEALE, A. C.

1893. Bulletin United States Geological Survey, No. 110, 1893: The Paleozoic section in the vicinity of Three Forks, Montana.

REED, F. R. C.

1899. Geological Magazine, Decade IV, Vol. VI, 1899 (August), pp. 358-361: A new trilobite from Mount Stephen, Field, British Columbia.

ROMINGER, C.

1887. Proceedings of the Academy of Natural Sciences of Philadelphia, 1887 (February 22), pp. 12-19: Description of Primordial fossils from Mount Stephens, N. W. Territory of Canada.

SALTER, J. W.

1866. Report of the 35th Meeting of the British Association for the Advancement of Science, held at Birmingham, September, 1865, pp. 284-286: Notes on the Sections and Fossils in a paper on the *Lingula*-flags by H. Hicks.

SHUMARD, B. F.

1860. Transactions of the Academy of Science of St. Louis for 1856-1860, I, 1860, pp. 624-627: Descriptions of five new species of Gasteropoda from the Coal Measures and a Brachiopod from the Potsdam sandstone of Texas.
1861. The American Journal of Science and Arts, 2d series, XXXII, 1861 (September), pp. 213-221: The Primordial Zone of Texas, with descriptions of new fossils.

TURNER, H. W.

1902. American Geologist, XXIX, 1902, pp. 261-272: A sketch of the historical geology of Esmeralda County, Nevada.

WALCOTT, C. D.

1883. The American Journal of Science, 3d series, XXVI, 1883 (December), pp. 437-442, 484: Pre-Carboniferous Strata in the Grand Canyon of the Colorado, Arizona.
1884. Monograph United States Geological Survey, VIII, 1884: Paleontology of the Eureka District, Nevada.
1886. Bulletin United States Geological Survey, No. 30, 1886: Second contribution to studies on the Cambrian Faunas of North America.
1887. American Journal of Science, 3d series, XXXIV, 1887 (September), pp. 187-199: Fauna of the "Upper Taconic" of Emmons, in Washington Co., N. Y.
1889. Proceedings United States National Museum for 1888, XI, 1889 (September 3), pp. 441-446: Description of new genera and species of fossils from the Middle Cambrian.
1891. Tenth Annual Report United States Geological Survey, 1891, pp. 509-774: The Fauna of the Lower Cambrian or *Olenellus* Zone.
1897. Proceedings United States National Museum, XIX, 1897 (August 27), pp. 707-718: Cambrian Brachiopoda: Genera *Iphidia* and *Yorkia*, with descriptions of new species of each, and of the genus *Acrothele*.
1898. Proceedings United States National Museum, XXI, 1898 (November 19), pp. 385-420: Cambrian Brachiopoda: *Obolus* and *Lingulella*, with description of new species.
1902. Proceedings United States National Museum, XXV, 1902 (November 3), pp. 577-612: Cambrian Brachiopoda: *Acrotreta*; *Linnarssonella*; *Obolus*; with descriptions of new species.
1905. Proceedings United States National Museum, XXVIII, 1905 (February 17), pp. 227-337: Cambrian Brachiopoda, with descriptions of new Genera and Species.
- 1908a. Smithsonian Miscellaneous Collections, LIII, Cambrian Geology and Paleontology, No. 1, 1908 (April 18), pp. 1-12: Nomenclature of some Cambrian Cordilleran Formations.
- 1908b. Smithsonian Miscellaneous Collections, LIII, Cambrian Geology and Paleontology, No. 2, 1908 (April 25), pp. 13-52: Cambrian trilobites.

WALCOTT, C. D. (continued) :

1908*c*. The Canadian Alpine Journal, I, No. 2, 1908, pp. 232-248: Mount Stephen Rocks and Fossils.

1908*d*. Smithsonian Miscellaneous Collections, LIII, Cambrian Geology and Paleontology, No. 3, 1908 (June), pp. 53-124: Cambrian Brachiopoda; descriptions of new genera and species.

WEED, W. H.

1900. Twentieth Annual Report United States Geological Survey for 1898-1899, Pt. III, 1900, pp. 271-461: Geology of the Little Belt Mountains, Montana, with notes on the mineral deposits of the Neihart, Barker, Yogo, and other districts.

WHITE, C. A.

1874. Geographical and Geological Explorations and Surveys West of the One Hundredth Meridian, Preliminary report upon Invertebrate Fossils, 1874 (December), pp. 5-27.

WHITEAVES, J. F.

1892. Canadian Record of Science, V, 1892, pp. 207-208: Description of a new genus and species of Phyllocarid Crustacea from the Middle Cambrian of Mount Stephens, British Columbia.

WINCHELL, N. H.

1886. Fourteenth Annual Report of the Geological and Natural History Survey of Minnesota for 1885, 1886, pp. 313-318: New Species of Fossils.

INDEX.

	Page
<i>Acrothele artemis</i> Walcott.....	198
<i>colleti</i> , new species.....	203, 213, 214
<i>panderi</i> , new species.....	203, 214
<i>spurri</i> Walcott.....	184, 189
<i>subsidia</i> (White).....	180, 181, 183, 195, 197
<i>subsidia hera</i> Walcott.....	184
<i>subsidia</i> var.	198
cf. <i>turneri</i> Walcott.....	196
<i>Acrothyra minor</i> Walcott.....	198
<i>Acrotreta attenuata</i> Meek.....	179, 180, 181
<i>bellatula</i> Walcott.....	179
<i>claytoni</i> Walcott.....	189
<i>depressa</i> (Walcott).....	210
<i>idahoensis</i> Walcott.....	177
<i>idahoensis alta</i> Walcott.....	193
<i>idahoensis sulcata</i> Walcott.....	198
<i>marjumensis</i> Walcott.....	179
<i>ophirensis</i> Walcott.....	178, 180, 182
<i>ophirensis descendens</i> Walcott.....	178
<i>primaeva</i> Walcott.....	184
<i>pyvidicula</i> White.....	180, 198
cf. <i>sagittalis</i> Salter.....	179
<i>sagittalis taconica</i> (Walcott).....	213, 215
sp. undt.	192, 198
<i>acuminata</i> , see <i>Lingulella</i> (<i>Lingulepis</i>).	
<i>acutangula</i> , see <i>Anomalocaris</i> .	
<i>agnes</i> , see <i>Olenopsis</i> .	
<i>Agnostus bidens</i> Meek.....	181
cf. <i>montis</i> Matthew.....	209, 211
sp. undt.	176, 178, 180, 192, 193, 194, 197, 199, 204, 205, 208, 210
<i>Agraulos</i>	175, 177, 189, 194, 195, 210, 212, 213, 214, 215
<i>alberta</i> , see <i>Nisusia</i> .	
Alberta, boundary of Cambrian land area in.....	169
<i>Albertella bosworthi</i> Walcott.....	214
<i>helena</i> Walcott.....	203, 214
sp. undt.	213, 214
<i>Albertella</i> fauna, in Montana and British Columbia, stratigraphic position of, discussed.....	203
<i>amii</i> , see <i>Nisusia</i> (<i>Jamesella</i>).	
<i>annulatus</i> , see <i>Hyolithellus</i> .	
<i>Anomalocaris</i> ?? <i>acutangula</i> Walcott.....	211
<i>canadensis</i> Whiteaves.....	211
? <i>whiteavesi</i> Walcott.....	211
<i>Anomocare</i>	176, 178, 179, 192, 193
<i>Archæocyathus</i>	187, 189
<i>argentea</i> , see <i>Isoxys</i> .	
<i>arguta</i> , see <i>Lingulella</i> .	
<i>artemis</i> , see <i>Acrothele</i> .	

	Page
<i>Asaphiscus minor</i> , new species.....	178
<i>checleri</i> Meek	181
sp. undt.	199
<i>Asaphus</i> ?	192
<i>attenuata</i> , see <i>Acrotreta</i> .	
<i>augusta</i> , see <i>Crepiccephalus</i> .	
Barrel Spring section, Nevada, described.....	188-189
<i>Bathyriscus howelli</i> Walcott.....	198
<i>occidentalis</i> (Matthew)	211
<i>ornatus</i> Walcott	211
<i>productus</i> (Hall and Whitfield).....	183, 197, 198, 203
<i>pupa</i> Matthew	211
<i>rotundatus</i> (Rominger)	211
sp. undt.	177, 178, 182, 183, 203, 208, 209, 210, 211, 213, 214
<i>Bathyrurus</i> , sp. undt.....	205
<i>bellatula</i> , see <i>Acrotreta</i> .	
<i>bellianus</i> , see <i>Platycceras</i> .	
<i>bellulus</i> , see <i>Obolus</i> (<i>Fordinia</i>).	
Belt Mountain uplift, mentioned.....	168, 191
Belt Terrane, Montana, mentioned.....	203, 208
Bibliography	218-220
<i>bidens</i> , see <i>Agnostus</i> .	
Big Cottonwood Canyon section, Utah, correlation.....	171
stratigraphic position of.....	169
Big Cottonwood Canyon sediments, Utah, probable nature of.....	170
<i>Billingsella coloradoensis</i> (Shumard).....	192, 193, 196, 198
<i>highlandensis</i> (Walcott)	184, 187
<i>marion</i> Walcott	212
sp. undt.	183, 195, 209
<i>billingsi</i> , see <i>Hyalithes</i> .	
Blacksmith Fork section, Utah, correlation.....	171
described	191-200
graphic representation of	199
résumé of	199-200
stratigraphic position	169
Blacksmith formation, Blacksmith Fork.....	171, 195
Bloomington formation, Blacksmith Fork.....	171, 194-195
<i>Bornemannia prima</i> , new genus and new species.....	213, 214
sp. undt.	214
Bosworth formation, Castle Mountain, view showing.....	pl. 20, figs. 1 and 2
Bosworth formation, Mount Bosworth.....	171, 205, 208, pl. 19
Bosworth and Paget formations, Mount Bosworth, break between.....	215
Bosworth section, see Mount Bosworth.	
<i>bosworthi</i> , see <i>Albertella</i> .	
Bow River Group, Mount Bosworth.....	215
Bow River Valley, Alberta, sediments in.....	217
Brigham formation, Blacksmith Fork.....	171, 199
British Columbia and Utah, connection between sections in.....	169
Burling, L. D., mentioned.....	173
<i>Burlingia hectori</i> Walcott.....	211
<i>cambrina</i> , see <i>Syntrophia</i> .	

	Page
Cambrian land area, extent and relations.....	168, 169
Cambrian (Lower) of Montana and British Columbia, compared.....	203
Cambrian sections of China and Cordilleran area, compared.....	172
<i>canadensis</i> , see <i>Anomalocaris</i> and <i>Olenellus</i> .	
Carboniferous rocks, section of on Dearborn River.....	200
<i>carinatus</i> , see <i>Hyolithes</i> .	
Castle Mountain, Alberta, fossils in.....	208, 212, 214
views of	pl. 20
Cathedral formation, Castle Mountain, view showing.....	pl. 20, fig. 2
Cathedral formation, Mount Bosworth.....	171, 212
Cathedral formation, Mount Stephen, view showing.....	pl. 21
Cathedral formation near Lake Louise, view showing.....	pl. 22
China, comparison of Cordilleran sections with sections in.....	172
<i>cingulata</i> , see <i>Kutorgina</i> .	
<i>claytoni</i> , see <i>Acrotreta</i> and <i>Olenellus</i> .	
<i>colleui</i> , see <i>Acrothelc</i> .	
<i>coloradoensis</i> , see <i>Billingsella</i> and <i>Eoorthis</i> .	
<i>columbiana</i> , see <i>Philhedra</i> .	
<i>Conocephalus</i> cf. <i>perseus</i> Hall, Matthew.....	211
<i>contracta</i> , see <i>Vanuxemella</i> .	
<i>cordillera</i> , see <i>Ptychoparia</i> .	
Cordilleran land area in Cambrian time.....	168, 169
Cordilleran sections compared with those in China.....	172
<i>corrugata</i> , see <i>Orthotheca</i> .	
<i>Corynexochus romingeri</i> Matthew.....	211
<i>crenistria</i> , see <i>Micromitra</i> (<i>Paterina</i>).	
<i>Crepicephalus augusta</i> Walcott.....	184
<i>liliana</i> Walcott.....	184
<i>texanus</i> (Shumard)	177, 178
sp. undt.....	175, 176, 204, 205, 213
<i>Cruziana</i>	184, 186, 187, 211, 216
<i>Cyrtolites</i>	193
<i>dawsoni</i> , see <i>Dorypyge</i> (<i>Kootenia</i>).	
Dearborn River section, Montana, correlation.....	171
described	200-203
résumé of	202-203
Deep Spring Valley, California, view of quartzite in.....	pl. 18
<i>depressa</i> , see <i>Acrotreta</i> .	
<i>desiderata</i> , see <i>Lingulella</i> .	
<i>Dicelloccephalus</i>	175, 191
<i>Dicellomus prolificus</i> Walcott.....	179
<i>discoideus</i> , see <i>Obolus</i> .	
Dome Canyon, House Range, view of.....	pl. 16
Dome formation, House Range.....	171, 182, pls. 16 and 17
<i>Dorypyge quadriceps</i> (Hall and Whitfield).....	196
(<i>Kootenia</i>) <i>dawsoni</i> (Walcott).....	211
sp. undt.	181, 183, 195, 197, 199
<i>dubia</i> , see <i>Lingulella</i> and <i>Siphonotreta</i> .	
Dunderberg shale, new formation name proposed.....	184
Eldon formation, Castle Mountain, views showing.....	pl. 20, figs. 1 and 2

	Page
Eldon formation, Mount Bosworth.....	171, 208-209
Eldon formation, Mount Stephen, view showing.....	pl. 21
Eldorado limestone, new formation name proposed.....	184
<i>ella</i> , see <i>Obolus (Westonia)</i> .	
<i>elongata</i> , see <i>Stenotheca</i> .	
Empire shales, Dearborn River, mentioned.....	203
<i>Endoceras</i>	189
<i>Eocystites ? longidactylus</i> Walcott.....	184, 197
<i>Eoorthis coloradoensis</i> (Meek).....	173, 175, 191, 192
<i>newberryi</i> Walcott	192
<i>remnicha</i> (N. H. Winchell).....	180
<i>thyone</i> Walcott	180
<i>zeno</i> Walcott	196
<i>Ethmophyllum gracile</i> Meek.....	187
Eureka District section, Nevada, new formation names proposed for....	184
<i>excelsis</i> , see <i>Trematobolus</i> .	
Fairview formation, near Lake Louise.....	171, 216, pl. 22
<i>festinata</i> , see <i>Nisusia</i> .	
<i>fieldensis</i> , see <i>Protypus</i> .	
<i>flagellum</i> , see <i>Hyolithellus</i> .	
Flathead sandstone, Little Belt Mountains.....	203
(<i>Fordinia</i>), see <i>Obolus (Fordinia)</i> .	
<i>fremonti</i> , see <i>Olenellus</i> .	
<i>gilberti</i> , see <i>Obolus (Fordinia)</i> and <i>Olenellus</i> .	
Gordon Mountain, discussion of <i>Albertella</i> fauna on.....	203
<i>gracile</i> , see <i>Ethmophyllum</i> .	
<i>granulatus</i> , see <i>Neolenus</i> .	
Hague, A., mentioned.....	184
Hamburgh limestone, old formation name retained.....	184
Hamburgh shale, Dunderberg shale proposed for.....	184
<i>haydeni</i> , see <i>Micromitra</i> .	
<i>hectori</i> , see <i>Burlingia</i> .	
<i>helena</i> , see <i>Albertella</i> and <i>Lingulella</i> .	
<i>highlandensis</i> , see <i>Billingsella</i> .	
<i>Holmia rowei</i> , new species.....	186, 187, 188, 189
<i>wecksi</i> , new species.....	186, 187, 189
House Range, Utah, map of.....	pl. 13
views of.....	pls. 14, 15, 16 and 17
House Range section, Utah, correlation.....	171
described	173-185
graphic representation of	174
stratigraphic position	169
résumé of	184-185
Howell formation, House Range.....	171, 182-183, pls. 16 and 17
<i>howelli</i> , see <i>Bathyriscus</i> .	198
<i>Huenella lesleyi</i> Walcott.....	193
<i>Hyolithellus annalatus</i> (Matthew).....	210
<i>flagellum</i> (Matthew)	210
<i>micans</i> Billings	213
sp. undt.	214

	Page
<i>Hyolithes billingsi</i> Walcott.....	183, 184, 213, 214, 215
<i>carinatus</i> Matthew	210
sp. undt.	178, 180, 182, 183, 188, 193, 194, 195, 196, 197, 198, 205, 209, 210, 211, 212, 213, 214, 215
<i>idahoensis</i> , see <i>Acrotreta</i> and <i>Zacanthoides</i> .	
<i>idahoensis alta</i> , see <i>Acrotreta</i> .	
<i>idahoensis sulcata</i> , see <i>Acrotreta</i> .	
<i>Illænurus</i>	175, 177, 192, 204, 205
<i>inflatus</i> , see <i>Neolenus</i> .	
<i>intermedius</i> , see <i>Neolenus</i> .	
<i>intermedius pugio</i> , see <i>Neolenus</i> .	
(<i>Iphidella</i>), see <i>Micromitra</i> (<i>Iphidella</i>).	
<i>iphis</i> , see <i>Obolus</i> (<i>Westonia</i>).	
<i>Isoxys</i> cf. <i>argentea</i> (Walcott).....	196, 209
<i>isse</i> , see <i>Lingulella</i> .	
(<i>Jamesella</i>), see <i>Nisusia</i> (<i>Jamesella</i>).	
Johnson, W. D., mentioned.....	173
<i>Karlia stephensis</i> Walcott	211
<i>kingi</i> , see <i>Ptychoparia</i> .	
Kintla uplift, mentioned	191
(<i>Kootenia</i>), see <i>Dorypyge</i> (<i>Kootenia</i>).	
<i>Kutorgina cingulata</i> (Billings).....	189, 215
<i>perugata</i> Walcott	189
sp. undt.	215
<i>labradorica</i> , see <i>Micromitra</i> (<i>Paterina</i>).	
<i>labradorica utahensis</i> , see <i>Micromitra</i> (<i>Paterina</i>).	
Lake Agnes, Alberta, fossils near.....	214
Lake Louise, view of mountains surrounding.....	pl. 22
Lake Louise formation, near Lake Louise.....	171, 216, pl. 22
Lake Louise section, Alberta, résumé of lower part.....	217
Langston formation, Blacksmith Fork.....	171, 198-199
Langston (?) formation, House Range.....	171, 183, pl. 17
<i>Leperditia</i>	183, 197, 211
<i>lesleyi</i> , see <i>Huenella</i> .	
<i>levis</i> , see <i>Zacanthoides</i> .	
<i>liliana</i> , see <i>Crepicephalus</i> .	
<i>linearis</i> , see <i>Scolithus</i> .	
<i>Lingulella arguta</i> (Walcott).....	179, 180, 182
<i>desiderata</i> (Walcott)	176, 177, 192, 194, 197, 198
<i>dubia</i> (Walcott)	183
<i>helena</i> (Walcott)	198
<i>isse</i> (Walcott).....	175, 176, 178, 198, 204, 209
<i>manticula</i> (White).....	176, 191, 192, 193
sp. undt.	209
(<i>Lingulepis</i>) <i>acuminata</i> (Conrad)	192, 193
<i>Linnarssonella modesta</i> Walcott.....	176
<i>nitens</i> Walcott	176
<i>transversa</i> Walcott	176
sp. undt.	181
Little Belt Mountains, discussion of horizons in.....	203

<i>longidactylus</i> , see <i>Ēocystites</i> .	
<i>louise</i> , see <i>Micromitra</i> (<i>Iphidella</i>).	
<i>lowi</i> , see <i>Nisusia</i> (<i>Jamesella</i>).	
<i>major</i> , see <i>Orthotheca</i> .	
Malade, Idaho, fossils near.....	198
<i>manticula</i> , see <i>Lingulella</i> .	
<i>marion</i> , see <i>Billingsella</i> .	
Marjum formation, House Range.....	171, 179-181, pl. 15, figs. 1 and 2
<i>marjumensis</i> , see <i>Acrotreta</i> .	
<i>mcconnelli</i> , see <i>Obolus</i> .	
<i>mcconnelli pelias</i> , see <i>Obolus</i> .	
<i>membranaceous</i> , see <i>Obolus</i> .	
<i>Menocephalus</i>	192, 210
<i>micans</i> , see <i>Hyalithellus</i> .	
<i>Mickwitzia occidentis</i> Walcott	187
<i>Microdiscus</i>	199, 212, 214
<i>Micromitra haydeni</i> Walcott	198
<i>sculptilis</i> Meek	179, 180, 194, 195
<i>stuarti</i> Walcott	197
(<i>Iphidella</i>) <i>louise</i> Walcott	216
<i>pannula</i> (White).....	182, 183, 184, 197, 198, 203, 210, 211, 212, 213, 215
<i>pannula ophiurensis</i> (Walcott).....	180, 198
(<i>Paterina</i>) <i>crenistria</i> ? (Walcott).....	176
<i>labradorica</i> (Billings)	213
<i>labradorica utahensis</i> (Walcott).....	182, 195, 196
<i>prospectensis</i> (Walcott)	189
<i>stissingensis</i> (Dwight)	209, 211
<i>superba</i> (Walcott)	197
<i>wapta</i> Walcott	214
sp. undt.	213
<i>minor</i> , see <i>Acrothyra</i> and <i>Asaphiscus</i> .	
<i>modesta</i> , see <i>Linnarssonella</i> .	
Montana, boundary of Cambrian land area in.....	168
<i>montis</i> , see <i>Agnostus</i> .	
Mount Bosworth section, British Columbia, correlation.....	171
described	204-217
discussion of <i>Albertella</i> fauna in.....	203
graphic representation of.....	206-207
résumé of	216-217
stratigraphic position	169
Mount Bosworth, view of Sherbrooke ridge on.....	pl. 19
Mount Daly, British Columbia, mentioned.....	205, 208, 215
Mount Fairview, view of.....	pl. 22
Mount Stephen, British Columbia, fossils on.....	209, 210, 211, 212, 213, 214, 215
view of	pl. 21
Mt. Whyte formation, stratigraphic position of, discussed.....	203
Mt. Whyte formation, Mount Bosworth.....	171, 203, 212-215
Mt. Whyte formation, near Lake Louise, view showing.....	pl. 22
Mt. Whyte formation, on Mt. Stephen, view showing.....	pl. 21
<i>nautcs</i> , see <i>Nisusia</i> (<i>Jamesella</i>).	

	Page
<i>Neolenus granulatus</i> (Matthew).....	211
<i>inflatus</i> Walcott	180
<i>intermedius</i> Walcott	180
<i>intermedius pugio</i> Walcott	180
<i>serratus</i> (Rominger)	211
<i>superbus</i> Walcott	180
sp. undt.	198, 199, 209, 210
Nevada, boundary of Cambrian land area in.....	168
<i>newberryi</i> , see <i>Eoorthis</i> .	
<i>Nisusia alberta</i> (Walcott).....	210, 211
<i>festinata</i> (Billings).....	214, 215
(<i>Jamesella</i>) <i>amii</i> Walcott	180
<i>lowi</i> Walcott	212, 213
<i>nautes</i> (Walcott)	180, 196
<i>spencei</i> (Walcott)	180
<i>nitens</i> , see <i>Linnarssonella</i> .	
Notch Peak formation, House Range.....	171, 173-175, pl. 14
<i>notchensis</i> , see <i>Obolus</i> (<i>Westonia</i>).	
Nounan formation, Blacksmith Fork.....	171, 193
<i>nundina</i> , see <i>Syntrophia</i> .	
<i>Obolella</i> , sp. undt.	186, 187
<i>Obolus discoidens</i> (Hall and Whitfield).....	193
<i>meconnelli</i> (Walcott).....	196, 197, 209, 210
<i>meconnelli pelias</i> (Walcott).....	176, 179, 180, 181, 194
<i>membranaceous</i> Walcott	209
<i>parvus</i> Walcott	214
<i>rotundatus</i> (Walcott)	176, 180
<i>tetonensis leda</i> Walcott	175
(<i>Fordinia</i>) <i>bellulus</i> (Walcott).....	193
<i>gilberti</i> Walcott	179
<i>perfectus</i> Walcott	178, 179
(<i>Westonia</i>) <i>ella</i> (Hall and Whitfield) ..	182, 183, 184, 196, 197, 198, 203, 211
<i>iphis</i> , new species.....	192
<i>notchensis</i> Walcott ..	173
<i>wasatchensis</i> Walcott	195
sp. undt.	192, 193, 196, 208
<i>occidens</i> , see <i>Mickwitzia</i> .	
<i>occidentalis</i> , see <i>Bathyriscus</i> .	
<i>Ogygopsis klotzi</i> (Rominger).....	211
sp. undt.	180, 181, 198, 199, 209
Ogygopsis shale, Mount Stephen, notes on.....	210-211
<i>Olenellus canadensis</i> , new species.....	215
<i>claytoni</i> , new species.....	139
<i>fremonti</i> , new species.....	187
<i>gilberti</i> Meek	184, 189
sp. undt.	186, 187, 189, 203, 213, 214
<i>Olenopsis agnes</i> , new species.....	214
? sp. undt.	203
<i>Ophileta</i>	204
<i>ophirensis</i> , see <i>Acrotreta</i> .	

	Page
<i>ophirensis descendens</i> , see <i>Acrotreta</i> .	
Ordovician rocks, sections of.....	173, 191, 204
<i>ornatus</i> , see <i>Bathyuriscus</i> .	
Orr formation, House Range.....	171, 175-177, pl. 15, fig. 1
<i>Orthoceras</i>	189
<i>Orthotheca corrugata</i> Matthew	210
<i>major</i> Walcott	197, 210
sp. undt.	199
<i>Oryctocephalus reynoldsi</i> Reed.....	211
<i>walkeri</i> Matthew	211
sp. undt.	199
<i>Owenella typa</i> , new genus and new species.....	180
Paget formation, Mount Bosworth.....	171, 205, pl. 19
Paget and Bosworth formations, Mount Bosworth, break between.....	215
<i>palliseri</i> , see <i>Ptychoparia</i> .	
<i>panderi</i> , see <i>Acrothele</i> .	
<i>pannula</i> , see <i>Micromitra (Iphidella)</i> .	
<i>pannula ophirensis</i> , see <i>Micromitra (Iphidella)</i> .	
<i>parvus</i> , see <i>Obolus</i> .	
(<i>Paterina</i>), see <i>Micromitra (Paterina)</i> .	
<i>perfectus</i> , see <i>Obolus (Fordinia)</i> .	
<i>perscus</i> , see <i>Conocephalus</i> .	
<i>perugata</i> , see <i>Kutorgina</i> .	
<i>Philhedra columbiana</i> (Walcott).....	210
Pioche formation, House Range.....	171, 184, pl. 17
<i>piochensis</i> , see <i>Ptychoparia</i> .	
<i>Platyceras bellianus</i> Walcott	210
<i>romingeri</i> Walcott	210
sp. undt.	181, 183, 199, 213
<i>prima</i> , see <i>Bornemannia</i> .	
<i>primava</i> , see <i>Acrotreta</i> .	
<i>Productus</i> , sp. undt.....	200
<i>productus</i> , see <i>Bathyuriscus</i> .	
<i>prolificus</i> , see <i>Dicellomus</i> .	
Prospect Mountain formation, House Range.....	171, 184, pl. 17
Prospect Mountain limestone, Eldorado limestone proposed for.....	184
Prospect Mountain sandstone, old formation name retained.....	184
<i>prospectensis</i> , see <i>Micromitra (Paterina)</i> .	
<i>Protospongia (spicules)</i>	194, 209
<i>Protypus fieldensis</i> , new species.....	215
new species	213, 215
sp. undt.	214
<i>Pterocephalus</i> ?	204
<i>Ptychaspis</i>	176
<i>Ptychoparia cordillera</i> (Rominger).....	211
<i>kingi</i> (Meek)	180, 181
<i>palliseri</i> Walcott	211
<i>piochensis</i> Walcott	183, 197
<i>subcoronata</i> (Hall and Whitfield).....	196
sp. undt.	175, 176, 178, 179, 180, 181, 182, 183, 189, 192, 193, 194, 195, 196, 197, 198, 199, 201, 204, 205, 208, 209, 210, 211, 212, 213, 214, 215.

<i>pupa</i> , see <i>Bathyriscus</i> .	
<i>pyxidicula</i> , see <i>Acrotreta</i> .	
<i>quadriceps</i> , see <i>Dorypyge</i> .	
<i>Raphistoma</i> sp.	173
<i>remnicha</i> , see <i>Eoorthis</i> .	
Resting Springs, California, fossils at.....	187
<i>reynoldsi</i> , see <i>Oryctocephalus</i> .	
<i>romingeri</i> , see <i>Corynexochus</i> and <i>Platyceras</i> .	
<i>rotundatus</i> , see <i>Bathyriscus</i> and <i>Obolus</i> .	
<i>rowei</i> , see <i>Holmia</i> .	
<i>rugosa</i> , see <i>Stenothecca</i> .	
<i>sagittalis</i> , see <i>Acrotreta</i> .	
<i>sagittalis taconica</i> , see <i>Acrotreta</i> .	
St. Charles formation, Blacksmith Fork.....	171, 191-193
St. Piran formation, near Lake Louise.....	171, 207, pl. 22
St. Piran formation, Mount Bosworth.....	215
<i>Salterella</i>	186, 189
Scapegoat Mountain, discussion of Albertella fauna on.....	203
<i>Scenella varians</i> Walcott	211, 213, 214, 215
sp. undt.	181, 182, 189, 196
<i>Schizambon typicalis</i> Walcott	175, 192
<i>Scolithus linearis</i> Haldeman.....	186
sp. undt.	186, 215
<i>sculptilis</i> , see <i>Micromitra</i> .	
<i>serratus</i> , see <i>Neolemus</i> .	
Sherbrooke formation, Mount Bosworth.....	171, 204-205, pl. 19
Sherbrooke ridge, view of.....	pl. 19
Silurian ? rocks, section of on Dearborn River.....	200-201
Silver Peak Group, California.....	185-188
Silver Peak section, Nevada, correlation.....	171
<i>simplex</i> , see <i>Wimanella</i> .	
<i>Siphonotreta ? dubia</i> , new species.....	189
Spence shale, Blacksmith Fork.....	171, 197-198
Spence shale, House Range.....	171, 183, pl. 17
<i>spencei</i> , see <i>Nisusia (Jamesella)</i> .	
<i>spinosus</i> , see <i>Zacanthoides</i> .	
<i>Spirifer</i>	200
<i>spurri</i> , see <i>Acrothele</i> .	
<i>Solenopleura</i>	175, 176, 178, 180, 192, 199, 208
<i>Stenothecca elongata</i> Walcott	189, 213
cf. <i>rugosa</i> (Hall).....	189
<i>wheeleri</i> Walcott	210
sp. undt.	199
Stephen formation, Castle Mountain, view showing.....	pl. 20, fig. 2
Stephen formation, Mount Bosworth.....	171, 209-212
Stephen formation, Castle Mountain, view showing.....	pl. 20, fig. 2
<i>stephenensis</i> , see <i>Karlia</i> .	
<i>stissingensis</i> , see <i>Micromitra (Paterina)</i> .	
<i>stuarti</i> , see <i>Micromitra (Paterina)</i> .	
<i>subcoronata</i> , see <i>Ptychoparia</i> .	

<i>subsida</i> , see <i>Acrothele</i> .	
<i>subsida hera</i> , see <i>Acrothele</i> .	
<i>superba</i> , see <i>Micromitra (Paterina)</i> .	
<i>superbus</i> , see <i>Neolenus</i> .	
<i>Swanton</i> <i>weeksii</i> Walcott	189
? sp. undt.	189
Swasey formation, House Range.....	171, 181-182, pls. 16 and 17
<i>Syntrophia cambria</i> Walcott	196
<i>nundina</i> Walcott	189, 191, 192
<i>unxia</i>	180
<i>Syringopora</i>	200
<i>tetonensis leda</i> , see <i>Obolus</i> .	
<i>texanus</i> , see <i>Crepicephalus</i> .	
<i>thyone</i> , see <i>Eoorthis</i> .	
<i>transversa</i> , see <i>Linnarssonella</i> .	
<i>Trematobolus excelsis</i> Walcott	187, 188
Turner, H. W., mentioned.....	185
<i>turneri</i> , see <i>Acrothele</i> .	
<i>tyba</i> , see <i>Owenella</i> .	
<i>typicalis</i> , see <i>Schizambon</i> and <i>Zacanthoides</i> .	
Uinta Mountain uplift, mentioned.....	191
<i>unxia</i> , see <i>Syntrophia</i> .	
Utah, boundary of Cambrian land area in.....	168
Utah and British Columbia, connection between sections in.....	169
Ute formation, Blacksmith Fork.....	171, 195-198
<i>Vanuxemella contracta</i> , new genus and new species.....	203, 214
<i>varians</i> , see <i>Scenella</i> .	
<i>walkeri</i> , see <i>Oryctocephalus</i> .	
<i>wapta</i> , see <i>Micromitra (Paterina)</i>	214
Wasatch Canyon, Box Elder County, Utah, fossils in.....	195, 197
<i>wasatchensis</i> , see <i>Obolus (Westonia)</i> .	
Waucoba Springs section, California, described.....	185-188
stratigraphic position	169
Weeks, F. B., mentioned.....	188
Weeks formation, House Range.....	171, 177-178, pl. 15, fig. 1
<i>weeksii</i> , see <i>Holmia</i> and <i>Swanton</i> .	
(<i>Westonia</i>), see <i>Obolus (Westonia)</i> .	
Wheeler Amphitheater, House Range, view of.....	pl. 15, fig. 2
Wheeler formation, House Range.....	171, 181, pl. 15, fig. 2
<i>wheeleri</i> , see <i>Asaphiscus</i> and <i>Stenothecca</i> .	
<i>whiteavesi</i> , see <i>Anomalocaris</i> .	
<i>Wimanella simplex</i> Walcott.....	203, 214
Wolsey shale, Little Belt Mountains.....	203
Wyoming, boundary of Cambrian land area in.....	168
<i>Zacanthoides idahoensis</i> Walcott	197
<i>levis</i> (Walcott)	184
<i>spinousus</i> Walcott	209, 211
<i>typicalis</i> (Walcott)	183
sp. undt.	181, 182, 183, 196, 198
<i>Zaphrentis</i>	200
<i>zeno</i> , see <i>Eoorthis</i> .	