SMITHSONIAN MISCELLANEOUS COLLECTIONS

VOLUME 73, NUMBER 2

OPINIONS RENDERED BY THE INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

OPINIONS 78 TO 81







CITY OF WASHINGTON

PUBLISHED BY THE SMITHSONIAN INSTITUTION
FEBRUARY 9, 1924



SMITHSONIAN MISCELLANEOUS COLLECTIONS

VOLUME 73, NUMBER 2

OPINIONS RENDERED BY THE INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

OPINIONS 78 TO 81



(Publication 2747)

CITY OF WASHINGTON

PUBLISHED BY THE SMITHSONIAN INSTITUTION

FEBRUARY 9, 1924

The Lord Gastimore (Press BALTIMORE, MD., U. S. A.

OPINIONS RENDERED BY THE INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

OPINIONS 78 TO 81

OPINION 78

Case of Dermacentor andersoni vs. Dermacentor venustus

SUMMARY.—On basis of the premises presented, the Commission is of the Opinion that *Dermacentor venustus* dates from Marx in Neumann, 1897, type specimen Collection Marx No. 122 (U. S. National Museum), from *Ovis aries*, Texas, and that *Dermacentor andersoni* dates from Stiles, 1908, holotype U. S. P. H. & M. H. S. 9467, from Woodman, Montana.

STATEMENT OF CASE.—This case has been submitted to the Commission by W. Dwight Pierce in the following letter, W. Dwight Pierce to Stiles:

Feb. 18, 1920: The recent publication of Wolbach's excellent monograph on Rocky Mountain Spotted Fever, again brings critically before the medical profession the confusion as to the name of the spotted fever tick. In order that we may get at this thing right and forever legally settle this name I appeal to the International Commission to give us a definite ruling on the proper name of the Rocky Mountain Spotted Fever Tick. In order that this ruling may be based on absolutely fair and just premises I would request that statements be requested of Dr. C. W. Stiles, Mr. Nathan Banks, Mr. F. C. Bishopp, and Dr. Nuttall, and others if necessary, these statements to be used as briefs and to be published with the ruling. My personal conclusions are as follows:

I. That there is no question whatever that Dermacentor andersoni Stiles (1905) refers to the Rocky Mountain Spotted Fever Tick.

2. That there is debatable ground as to whether *D. venustus* Banks (1908) is conspecific and refers to the fever tick.

3. The first reference I find to *D. venustus* Marx mss. is in Neumann (1897) as a synonym of *D. reticulatus* Fabricius, undescribed.

4. Dermacentor andersoni Stiles was described as the fever tick, in 1905, (U. S. Treas. Dept., Hyg. Lab., Bull. 20, pp. 1-119) and the description strengthened in 1908 and 1910.

5. In 1908 Banks drew up the description, as a new species, of *D. venustus* (Marx mss.), from the Marx material, which was subsequently examined by Stiles, and found to consist of three lots of material of at least two species. Stiles definitely picked from Bank's type material Marx No. 122 as type of

the species D. venustus. This was Texas material. Since both Marx and Banks confused more than one species and neither designated an individual type from the material, Stiles' designation is valid.

6. In 1910 Stiles differentiated between the two species *D. andersoni* and *D. venustus*, using the designated type individuals as basis of his differentiation.

7. It therefore appears to me that *D. andersoni* not only is definitely the fever tick, but that it antedates *D. venustus* Banks, which may have originally had specimens of the fever tick contained within its series, but which when typically defined according to our laws of nomenclature is a very different species, with a range extraterritorial to the fever area.

8. The entire medical profession would welcome a final legal decision on this name at the earliest possible moment.

In accordance with Pierce's suggestion, the Secretary has invited Mr. Banks, Mr. Bishopp, and Doctor Nuttall to submit statements. No reply has been received from Nuttall.

Banks submits the following letter:

Cambridge, Mass., April 29, 1920: As far as I am concerned there is no "question" as to the name of the Rocky Mt. Spotted Fever Tick, and no decisions of any committee can alter facts. D. venustus was published in 1908, D. andersoni a few months later. All previous references to either name had nothing to do with the matter, as there was no description till that time. D. andersoni of 1905 was not referred to as the fever tick but as the tick that did not carry the disease.

Type label was placed on a certain vial of *D. venustus* at time of publication and anyone who examined the collection of the Bureau of Entomology would have found it.

Later attempts at limitation of the name cannot alter the facts.

Bishopp submits the following:

Dallas, Texas, May 1, 1920: I am enclosing herewith a statement on this subject which I drew up in 1912, which I believe sets forth my viewpoint in a rather concise way.

THE CORRECT NAME OF THE ROCKY MOUNTAIN SPOTTED FEVER TICK

By F. C. BISHOPP

There is considerable confusion regarding the correct scientific name of the tick which transmits Rocky Mountain spotted fever. As the several statements which have been made upon this question do not seem to have cleared the matter up, it seems best to briefly review the situation and show the exact status of the question.

Labels bearing the name *D. venustus* n. sp. were placed by Marx in vials containing specimens of ticks from Soldier, Idaho, Las Cruces, N. M., and Texas (on sheep). All of these specimens were deposited in the U. S. National Museum. No manuscript notes or drawings were left with this material.

After the death of Dr. Marx, these specimens together with other material from the Marx collection, were sent to Prof. L. G. Neumann for study. In

1897 Neumann, after studying this material, considered it the same as the European D. reticulatus, the manuscript name D. renustus being cited in identifying the specimens from the United States.

In 1905 Stiles used the name andersoni for material from Montana, concluding that the species did not transmit the disease known as Rocky Mountain spotted fever. This was a nomen nudum as it was unaccompanied by a description or by a specific indication. In June, 1908, Banks, after studying all of the Marx material, described the species, using the Marx name D. venustus. He used one of the males from Soldier, Idaho, as the type for his species. By doing this Banks repudiated Neumann's placing the species as a synonym of reticulatus. The name venustus cannot date from 1897 because Neumann did not specifically differentiate this species from his reticulatus, but confused it with his material. Banks, by describing this species in 1908, gave it a standing in nomenclature as a distinct species. In July, 1908, Stiles, after studying part of the Marx material exclusive of Banks' type of D. venustus, briefly described specimens from Montana under the name D. andersoni. Subsequently, Aug. 1910—(Taxonomic Value of the Microscopic Structure of the Stigmal Plates in the Tick Genus, Dermacentor, Bull, No. 62, Hygienic Laboratory), Stiles applied the name D. venustus to the Texas material which was contained in the Marx collection, and designated this as the type of the species. He stated that the New Mexico material could not be positively identified and that the Idaho specimen was not sufficient to base a determination upon. In this publication he fully described certain Montana material under the name D. andersoni.

On Oct. 29, 1910, in the JAMA, Stiles reiterates Banks' statement that Neumann was incorrect in placing D. venustus, Marx's manuscript, as a synonym of D. reticulatus but claimed that venustus should date from Neumann, 1897. In the last paragraph of this statement he says "Were the premise correct that Marx's specimens from Texas and New Mexico are identical with the specimens from Montana, D. venustus would of necessity be the correct name for the Rocky Mountain spotted fever tick, but this premise is erroneous and the name venustus must be applied to the species containing the original specimens designated under this name." We must take exception to the last portion of this sentence, as a part of the material labeled D, remustus by Marx (specimens from Soldier, Idaho), is identical with the form found in Montana and called D. andersoni by Stiles. One of these males from Soldier, Idaho, was designated as type of D. venustus by Banks. A careful comparison of this type specimen with Stiles' type of D. andersoni shows the two species to be identical and there is no question that this is the form which conveys Rocky Mountain spotted fever. Hence D. andersoni is a synonym of D. venustus, and if Stiles is correct in his belief that the specimens from "Texas on sheep" are specifically different from D. venustus of Montana, this species requires another name.

Stiles submits the following statement to the Commission:

- I. IN SUMMARY, I submit to the Commission the following points:
 - a. Under the International Rules, the name *D. venustus* dates from Marx in Neumann, 1807a, 365. (Art. 25; Opinion of *Halicampus grayi* 1856, ruled upon in Opinion No. 53.)

- b. It would require, under the By-Laws, a two-thirds vote of the Commission to reverse Opinion 53 in the case of D. venustus.
- c. As the original publication of *D. venustus* 1897 mentioned only two localities (New Mexico and Texas), only these two localities and no other come into consideration as type locality. (Not covered by the International Rules but in harmony with Zoological practice.)
- d. The only original specimens of Marx's *D. venustus* mentioned by Neumann in 1897 have been found and identified, and only these come into consideration as type specimens. (Not covered by International Rules, but in harmony with Zoological practice.)
- e. Marx No. 122, from Texas, host *Ovis aries*, is the first and the only originally published specimen publicly or privately designated as type specimen and this must remain type specimen. (Not covered by the International Rules, but in harmony with Zoological practice.)
- f. D. venustus n. sp. Banks, 1908, is antedated by D. venustus 1897, hence is a homonym, hence is to be suppressed. (Art. 35.)
- g. It is generally admitted (by Banks, Bishopp, Stiles, etc.) that D. venustus n. sp. Banks, 1908, is specifically identical with D. andersoni Stiles, (1905) 1908, but evidence is not lacking that it also contains Marx's specimens 120 from New Mexico and 122 from sheep in Texas. The only specimen of D. venustus 1908 known to have the label of "type" in Banks' handwriting is in the U. S. National Museum (Marx No. 10) and although Banks specifically states that his type belongs in the collection of the Bureau of Entomology, the Museum specimens can be taken as Banks' type until evidence of error is presented; this specimen seems to be specifically identical with D. andersoni [but as it is a single specimen, it has not been mounted]. Accordingly, D. venustus Banks, 1908, (nec Marx, 1897) is synonymous with D. andersoni Stiles (1905) 1908.
- h. Under the International Rules D. andersoni is the earliest available name for the Rocky Mountain Spotted Fever Tick, hence (Art. 25, 35) it is the valid name.
- i. As a matter of propriety, I will refrain from utilizing my Commissioner's right of vote on this case, since it involves a name proposed by myself, but I obligate myself to accept the decision of the Commission as determined by the By-Laws.
- j. The following documents are submitted to the reviewing Commissioner (Stejneger) either in original or in copy, in connection with this case.

 Banks, 1908.—Revision of the Ixodoidea < Tech. Series, No. 15, Bureau of Entomology.
 - 1910.—The Scientific Name of the Spotted Fever Tick < JAMA, v. 55 (18), 1574-1575.
 - ? 1908.—Undated letter, Banks to Stiles regarding type specimen of D. venustus.
 - NEUMANN, 1897a.—Revision de la famille des Ixodidés. (2e mémoire) < Mém. Soc. Zool. France, Par., v. 10 (3-4), pp. 324-420.
 - STILES, 1905f. —A Zoological Investigation, etc., < Bull. 20, Hyg. Lab.
 1907. —[Transcript of Minutes, Ent. Soc. Wash., Jan. 10, 1907,
 pp. 10-11, giving Secretary's abstract of Stiles' paper
 on stigmal plates of the genus *Dermacentor*.]

1908m.—The common tick (Dermacentor andersoni) of the Bitter Root valley < Pub. Health Rep., U. S. Pub. Health & Mar.-Hosp. Serv., Wash., v. 23 (27), p. 949.</p>

1908. —Copy of letter, Stiles to Banks, June 10.

1909. —Copy of letter, Stiles to Banks, Mar. 19.

1909. —Copy of letter, Stiles to Banks, Oct. 23.

1910. —The taxonomic value of the microscopic structure of the stigmal plates in the tick genus *Dermacentor* < Bull. 62, Hyg. Lab.

1911. -Letter, Stiles to Banks, Feb. 20.

2. The first actual publication of the name *Dermacentor venustus* occurs in Neumann (1897a, 365) who examined specimens of ticks from the Marx collection, and determined them as *Dermacentor reticulatus*. His original reads as follows:

"D'Amérique, j'en ai 2 femelles originaires du Mont Diablo, en Californie (Coll. de l'Acad. des sciences de Californie). La Collection du Départ. of Agriculture de Washington et celle de la Smithsonian Institution en contiennent plusieurs mâles et femelles recueillis aussi en Californie, sur le Daim, et étiquetés par G. Marx D. occidentalis. D'autres proviennent du Texas et du Nouveau-Mexique et sont étiquetés D. venustus. Je rapporte aussi à la même espèce 9 mâles et 1 femelle, jeunes, à patine blanche encore peu marquée, à coloration génèrale brun foncé, provenant de Las Paz (?) et appartenant au Muséum de Berlin."

3. Accordingly, *D. venustus* was first published as a synonym of *D. reticulatus* and the original publication clearly cites Texas [Marx 122] and New Mexico [Marx 120] as the first published, hence type localities, unless it can be shown that Marx designated some other specimens from some other locality as type specimens.

4. The first point which arises is whether or not the manuscript or label name *D. venustus* received nomenclatorial status in this publication by Neumann. The answer to this question is found in three opinions already issued by the Commission, namely, Opinions Nos. 1, 4, and 53.

5. Status of a Manuscript Name published in Synonymy.—Article 25 of the Code reads:

"The valid name of a genus or species can be only that name under which it was first designated, on the condition:

(a) That this name was published and accompanied by an indication, or a definition, or a description; and

(b) That the author has applied the principles of binary nomenclature."

6. As Neumann (1897a) is both binary and binomial, the decision reverts to "(a)." This point has been discussed in several opinions, thus:

7. Opinion I states: "The word *indication* in Art. 25a is to be construed as follows: (A) with regard to specific names, an indication is (I) a bibliographic reference, or (2) a published figure (illustration), or (3) a definite citation of an earlier name for which a new name is proposed."

8. Opinion 4 states: "Manuscript names acquire standing in nomenclature when printed in connection with the provisions of Art. 25, and the question as to their validity is not influenced by the fact whether such names are accepted or rejected by the author responsible for their publication."

9. Opinion 53 covers a case identical with the one at issue, namely the status of "Halicampus grayi Kp. British Museum," published as synonym of "Halicampus conspicillatus," corresponding exactly to Dermacentor venustus, Collection Marx, U. S. Nat. Mus., published as synonym of D. reticulatus. In Opinion 53, written by Stejneger and Stiles, concurred in by 9 Commissioners, dissented from by 2 Commissioners, Halicampus grayi 1856 was recognized under Art. 25 and Opinion 4 as published and hence as available and was given precedence over H. koilomatodon (about 1865).

10. According to the By-Laws of the Commission, an Opinion cannot be reversed by less than a two-thirds vote. Opinion 53 has never before come up for reversal and unless a two-thirds vote now obtains against Opinion 53, D. venustus must be accepted as available from the date of 1897.

- 11. As D. venustus Marx in Neumann, 1897, is under Opinion 53 clearly to be accepted as a published and available name, and not as a nomen nudum, it remains to enquire into its validity. Two possibilities present themselves, namely,
 - a. Is *D. venustus* a synonym of *D. reticulatus*, as assumed by Neumann? If Neumann's view is sustained, the name *D. venustus* is clearly not valid for *D. reticulatus* unless it be shown that no earlier name for this species is available. But even then, as a synonym of *D. reticulatus* it would preclude its (*venustus*) later use for any other species.
 - b. Is *D. venustus* Marx in Neumann distinct from *D. reticulatus?* In other words, should *D. reticulatus* as defined by Neumann be sub-divided? All authors now agree that it should be, and that certain American (Marx) specimens of *D. reticulatus* (*D. venustus*) represent a distinct species.
- 12. Under this latter premise it is necessary to determine if possible the type specimen and the type locality of *D. venustus* Marx in Neumann.
- 13. Obviously, the type locality can be only the originally published locality and the type specimens can be only the originally published specimens. Fortunately, Neumann has given definite information as to the locality, namely, the United States of North America and he specifically cites two States, namely, Texas and New Mexico. Fortunately, it is possible to identify the original specimens also, on basis of the following data:
- 14. When Neumann returned the Marx material to the U. S. National Museum I borrowed the specimens. The exact date when these came into my hands does not appear to be recorded in my notes. There were three bottles which contained the name D. venustus on labels, namely, Marx No. 120, one male, from New Mexico; No. 121, one male from Soldier, Idaho, host, Mountain Goat; and No. 122, 3 males, I female, from Texas, host, Ovis aries. [See below, under Stiles, 1910.] It seems obvious that Nos. 120 and 122 represent the Marx material, and the only specimens of Marx's D. venustus mentioned by Neumann, 1897a, hence, only these two are available as type material. Later Stiles (1910, 44-46) definitely published Marx No. 122 as the type specimen. This is the first (and so far as I know, the only) publication of the Museum number of the type.
- 15. From copies of correspondence in my files it is clear that I returned Marx 122 to the U. S. National Museum accompanied by a letter dated March 19, 1909; and that I returned Marx 120 and 121 to the U. S. National Museum accompanied by a letter dated February 20, 1911.

16. My letter files also show that in answer to a letter from me dated Oct. 23, 1909, asking where the types of *D. parumapertus marginatus* and *D. nigrolineatus* were deposited, Mr. Banks replied (in an undated letter) that the type of *D. p. marginatus* was in his private collection, "the type of *Derm. venustus* in Bur[eau] Entom[ology] Coll[ection]," that of *D. nigrolineatus* in the Mus. Comp. Zool., Harvard, "cotypes or paratypes of *D. nitens* in Marx Coll., U. S. N. Mus." and of "*D. parumapertus* and *D. occidentalis*, also Marx coll., at least paratypes." It will be observed that this statement (namely, that the type of *D. venustus* is in the collection of the Bureau of Entomology [no mention of Marx collection]) is in harmony with Mr. Banks' statement of April 29, 1920. The Marx collection has at no time been the property of the Bureau of Entomology.

17. On Dec. 6, 1920, in the presence of Prof. H. E. Ewing, of the Bureau of Entomology, I examined three bottles of ticks at the U. S. National Museum, as follows: Marx 121 and 122 (see *supra*). Also a bottle containing the label "No. 10. *Dermacentor venustus* Marx Idaho Coll. Marx." This bottle also contains a paper with the word "type" written in a handwriting identified by Professor Ewing as that of Banks. The Marx label is in a different handwriting from that of Marx 121 and 122. This Marx 10 is not Marx 120.

18. Here is, accordingly, a bottle attributed to the Marx Collection which I had never seen prior to Dec. 6, 1920. It contains no label written either by Marx, by Neumann, or by E. A. Schwartz (who went over the Marx collection after Marx's death). Schwartz identifies the Marx label as probably written by C. V. Piper. That this specimen is not available as type specimen of *D. venustus* Marx in Neumann follows from the fact that Neumann (1897a) did not refer to any specimens from Idaho.

19. The fact that Banks twice states that the type of *D. venustus* is in the Bureau of Entomology Collection while the specimen with the label "No. 10, Coll. Marx," contains a slip of paper bearing the word "type" in Banks' handwriting is not, therefore, of special importance so far as the date 1897 is concerned, but comes into consideration in connection with the date 1908.

20. Banks (1908, 46-47, 55, pl. 8, figs. 4, 5, 7) described *Dermacentor venustus* n. sp. Banks. In addition to the specific description, which is clearly influenced chiefly by material from the Northwest, Banks states:

"Specimens come from various places in the West; Olympia, Yakima, Klikitat Valley, and Grand Coulee, Wash.; Fort Collins and Boulder, Colo.; Pecos and Las Cruces, N. Mex.; Bozeman, Mont.; Bridger Basin, Utah; Soldier, Idaho, and Texas (on sheep).

"This species is quite common in the Northwest. It has been included in D. occidentalis, by Neumann, but was separated out by Doctor Marx in manuscript under the name I have adopted. It is larger than D. occidentalis, with more red and less white in the coloring, and differs in many minor points of structure, as size of porose areas, size of hind coxae in male, etc. This is the species supposed to be concerned in the transmission of spotted fever in Montana."

21. It will be noticed that Banks cites specimens from "Pecos and Las Cruces, N. M." and "Texas (on sheep)" and that he says it was separated out from D. occidentalis "by Doctor Marx in manuscript under the name I have adopted." Banks does not cite the museum number of the type specimen.

- 22. The status of *D. venustus* n. sp. 1908 and its type specimen must be determined. Theoretically, three possibilities are present, namely:
 - a. D. venustus n. sp. Banks, 1908, might be identical with D. venustus Marx in Neumann, 1897; or
 - b. D. venustus n. sp. Banks, 1908, might represent a new species; or
 - c. D. venustus n. sp. Banks, 1908, might be D. venustus 1897 plus another species.
- 23. Is D. venustus n. sp. Banks, 1908, identical with D. venustus Marx in Neumann, 1897? Banks distinctly states that he adopts the name from Marx's manuscript. Neither Bishopp nor I have been able to find this manuscript, so possibly reference is made to the labels in the bottles. Banks quotes among the localities, "Las Cruces, New Mexico," "Soldier, Idaho," and "Texas (on sheep)." These three localities are in harmony with the Marx specimens Nos. 120, 121, 122. The presumption therefore would seem to be that Banks examined these three specimens. I am in a position to state that these three specimens, with drawings of No. 122, and with my manuscript giving No. 122 as type of D. venustus were placed on a table in my laboratory in front of Mr. Banks for examination prior to the publication of his paper. Bishopp (see supra) states that Banks studied "all of the Marx material" and this would seem to include Marx 120, 121, and 122. Banks, however, (1910, JAMA, 1574-1575) states that he never studied Marx 120 and 122 (namely the specimens published by me in 1910 as D. venustus). If Banks' D. venustus is identical with Marx's D. venustus as published in Neumann, the species should be attributed to Marx.
- 24. Is D. venustus n. sp. Banks, 1908, distinct from D. venustus Marx in Neumann, 1897? If this represents the correct status of facts, then D. venustus Banks, 1908, is a homonym of D. venustus 1897 and therefore cannot be used as a valid name.
- 25. Does *D. venustus* n. sp. Banks, 1908, include *D. venustus* Marx in Neumann, 1897, plus some other species? If this be the status of affairs, it is clear that such portion of *D. venustus* of Banks, 1908, as agrees with *D. venustus* 1897 should be allocated to *D. venustus* 1897 and that the remaining portion should be known under some other name.
- 26. It would appear, therefore, that the crux of the problem lies in establishing the type specimen of *D. venustus* of Banks, 1908. The evidence at my disposal, bearing on this point, is as follows:
- 27. Banks has twice stated in letters that the type of his *D. venustus* of 1908 is in the Collection of the Bureau of Entomolgy. He has also stated in a letter that "type label was placed on a certain vial of *D. venustus* at time of publication." Bishopp states that Banks "used one of the males from Soldier, Idaho, as type for his species." In the presence of Professor Ewing, Dec. 6, 1920, I established the fact that there is in the National Museum a specimen marked "Coll. Marx, *Dermacentor venustus* Marx Idaho," and that the bottle contains a label, identified by Ewing as in Banks' handwriting, reading "type."
- 28. The Marx specimen from "Soldier, Idaho," No. 121, was in my laboratory at the time Banks visited me in order to examine Marx's specimens, and it is not the specimen containing Banks' label "type." Banks (1910, JAMA, 1574-1575) states that his *D. venustus* 1908 is identical with my *D. andersoni*, and this view is in harmony with the specimen which bears Banks' label

"type." How and whether this specimen changed from the Bureau of Entomology Collection to the Marx Collection is as yet not clear.

29. Judged from the specimen containing Banks' label "type," D. venustus n. sp. Banks, 1908, falls, therefore, as a homonym of D, venustus Marx in Neumann, 1897, and it is either a synonym or it is not a synonym. To determine this latter point, it is necessary to examine Stiles (1910) who reexamined the specimens (Marx 120 and 121 from New Mexico and Texas) of D. venustus Marx published by Neumann, 1897. Specimen 122 (mentioned by Neumann) and selected by Stiles as type is specifically distinct from the specimen which bears Banks' label as representing the type of D. venustus Banks, 1908. As this was the first selection of any specimen of the Marx-Neumann (1897) material as type, and as the Idaho material was not available as type, since it was not mentioned by Neumann (although Marx 121 from a mountain goat, at Soldier, Idaho, was examined by him), a comparison of the type specimens in question, namely, Marx 120 (type of D. venustus Marx in Neumann, 1897, as published by Stiles, 1910) with Marx No. 10 (type of D. venustus Banks, 1908, according to the label in Banks' handwriting, but not entirely in harmony with his correspondence) appears therefore to settle the question that nomenclatorially D. venustus 1908 is not absolutely (from point of view of type specimen) synonymous with D, venustus 1897. Accordingly, the name D. venustus n. sp. Banks, 1908, drops as a homonym.

30. It next becomes necessary to enquire into the valid name for the species represented by *D. venustus* n. sp. Banks, 1908 (*nec* Marx in Neumann, 1897) incriminated as vector of Rocky Mountain Spotted Fever.

31. The systematic history of this tick is indeed complicated, owing to the difficulties connected with specific determinations. It has been studied by Marx, Neumann, Banks, and Stiles, all four of whom were fairly familiar with the group. These specialists confused the species with: D. occidentalis, D. venustus, D. electus, and D. reticulatus. These various species were not all clearly and definitely defined from each other until 1010, although all four of the authors just mentioned, and other authors also, had at various times determined a number of specimens correctly.

32. Anderson collected in the Bitter Root Valley some ticks which Wilson & Chowning and Anderson had incriminated as the vector of the Rocky Mountain Spotted Fever. Stiles (in Anderson, 1903, 21) made a provisional determination of this material as *Dermacentor reticulatus*.

33. Stiles (1904 1(m), 1649 (363)) obtained from the Bitter Root Valley a considerable amount of tick material which agreed with the tick which Wilson & Chowning (1902, 1903, 1904) and Anderson (1903) had incriminated as the vector of Rocky Mountain Spotted Fever. Stiles states:

"6. The tick most common in the valley is a dermacentor which is very closely allied to *D. reticulatus*. The data now at my disposal indicates, however, that it represents a distinct species."

"7. These ticks are common on horses, cattle, and dogs, and more or less frequent on man, but there is nothing to indicate that a hibernating animal is necessary for their development; in fact, indications (seasonal distribution) are not entirely lacking that the spermophile forms a more or less accidental host for this species."

34. Later, Stiles (1905f, 7, 22, 24) in discussing his negative results as to the piroplasmic nature of the Rocky Mountain Spotted Fever, uses the new name "Dermacentor andersoni" in referring to this tick which Wilson & Chowning (1902, 1903, 1904) and Anderson (1903) had incriminated as vector of the supposed Piroplasma hominis. Zoological characters are not cited and so far as this article is concerned, the name Dermacentor andersoni rests solely upon the geographic distribution of the tick and the earlier claims that this arachnoid is the vector of the disease.

35. Later, Stiles (1907, 10-12) presented to the Entomological Society of Washington drawings of *D. andersoni*, *D. venustus*, *D. occidentalis*, etc., demonstrating the differential characters on which the species in question are recognizable, but these names were not published in the Secretary's minutes

of the meeting. Mr. Banks was present and discussed the paper.

36. After the meeting, Mr. Banks asked to examine some of the specimens and was invited to do so. For this purpose he visited my laboratory (exact date unknown, but between Jan. 10, 1907 and June 6, 1908). I placed before him the manuscript, drawings, and specimens, and a miscroscope; he used his own hand lens. Among the specimens placed before him were "Marx·120, 121, 122." Mr. Banks examined some of the drawings and specimens; as he was received as a guest he was free to do this.

37. Upon the publication of *D. venustus* n. sp. Banks, 1908, Stiles, in the hope of forestalling further confusion, published (1908m, 949) a short note giving

some of the more important differential characters.

38. Later, Stiles (1910, 36-46) published his delayed manuscript, describing and figuring in detail *D. andersoni* Stiles (type No. 9467, from Woodman, Mont.) (giving *D. venustus* pars of Banks, 1908, as synonym) and *D. venustus* Marx, 1897, in Neumann, 1897 (type Marx 122 from Texas) giving *D. venustus* pars of Banks, 1908, as synonym).

DISCUSSION.—The present case, to my mind, is much less complicated than the argument submitted would indicate.

The facts appear to be as follows:

I. In 1897 G. Neumann (Mem. Soc. Zool. France, vol. 10, pp. 324-420) published a "Révision de la famille des Ixodidés," in which under the specific heading of Dermacentor reticulatus (Fabricius), up to that time known only from the Old World, he says on p. 365: "La Collection du Départ, of Agriculture de Washington et celle de la Smithsonian Institution en [i. e., D. reticulatus] contiennent plusieurs mâles et femelles receuilles aussi en Californie, sur le Daim, et étiquetés par G. Marx D. occidentalis. D'autres proviennent de Texas et du Noveau-Mexique et sont étiquetés D. venustus." There is no further reference to these specimens, and this is the first published reference to Dermacentor venustus. Although there is no description, the name is not a nomem nudum, since according to Opinion 53 it has a nomenclatorial status that cannot be ignored. The case is absolutely comparable, though not quite identical, with that of Halicampus grayi, quoted only in synonymy as being in the British Museum, but not described, regarding which Opinion 53 says that "there can be no question but that Halicampus grayi has been published in connection with a bibliographic reference, and in connection with a description, and on this account the name must be considered as dating from 1856." As Opinion 53 is in force and consequently is part of the Code, it is clear that *Dermacentor venustus* as a published and available specific name dates from 1897. But it is also unidentifiable from the published data then available. Dr. Neumann himself apparently thought it the same as *reticulatus*, but he gives no data by which it can be determined from his publication whether he was right or wrong. The reference to certain localities can have no bearing, nor is there any indication that he referred to actual type specimens. Marx's type specimens may have been examined, or they may not, as far as contemporaneous published evidence is concerned.

The next appearance of the name in any publication is in 1908 when Banks (A Revision of the Ixodoidea, or Ticks, of the United States, June 6, 1908, p. 46, pl. 8, figs. 4, 5, 7) described Dermacentor venustus as a new species without reference to Marx's manuscript name of 1897 in Neumann. He mentions neither a type specimen, nor does he give any single type locality. He says: "Specimens come from various places in the West: Olympia, Yakima, Klikitat Valley, and Grand Coulee, Wash.; Fort Collins and Boulder, Colo.; Pecos and Las Cruces, N. Mex.; Bozeman, Mont.; Bridger Basin, Utah; Soldier, Idaho; and Texas (on sheep)." On page 48, under D. occidentalis, he says: "Neumann first considered D. occidentalis and D. venustus of Marx as identical with the European D. reticulatus. When he described D. occidentalis, Neumann included with it D. venustus of the Marx manuscript. However, I have restricted the name to the form to which Marx applied it." This last sentence is not strictly correct. When Neumann described D. reticulatus occidentalis, which was done in January, 1905 (Arch. Parasitol., Paris, vol. 9, no. 2, p. 235), he did not mention D. venustus at all; he only recognized several of and of collected on "le Daim," California, and labeled D. occidentalis by G. Marx, as a distinguishable subspecies [variété] of the species D. reticulatus, in other words, in 1905 he recognized his species D. reticulatus of 1807, as a complex one including still the material which Marx had labeled D. venustus, and with the right of the first reviser he separated out and fixed the name of D. occidentalis. But he did nothing to D. venustus; he still kept it in the synonymy of D. reticulatus. Banks, however, in 1908, accepted Neumann's action as first reviser, as far as D. occidentalis is concerned (recognizing it however as full species), but went a step further and exercised his right as next reviser to segregate Marx's D. venustus out of the complex D. reticulatus of Neumann 1897. In the

D. venustus thus restricted, Banks included specimens from Washington, Colorado, New Mexico, Montana, Utah, Idaho, and Texas. No type locality, nor type is mentioned, as stated before. In the absence of definite type designation the presumption in 1908 is, therefore, that the D. venustus of 1908 and the one of 1897 are identical.

Later in the same year Dr. Stiles (Weekly Pub. Health Rep., vol. 23, pt. 2, nos. 27 to 52, July 3, 1908m, p. 949) briefly indicated that Banks' D. venustus of 1908 was still a specific complex, separating out from it, and for the first time diagnosing, the specimens from Montana as Dermacentor andersoni [D. andersoni Stiles 1905, nomen nudum]. Incidentally he also mentioned D. venustus as an allied species from Texas, but gave no characters and mentioned no type.

Up to that time there had been no published mention of type specimen or of the names having been tied down to any particular specimens, except in the case of *D. occidentalis*.

No further revision and subdivision of the complex took place until August, 1910, when Stiles' paper entitled "the taxonomic value of the microscopic structure of the stigmal plates in the tick genus *Dermacentor*" was published (Hyg. Lab. U. S. Publ. Health Mar. Hosp. Serv.). In this he undertook a final revision of the specific complex *D. reticulatus* as presented by Neumann in 1897. In this revision he described fully and figured *D. venustus* designating "Marx 122 in U. S. National Museum. Host, Sheep (*Ovis aries*) in Texas" as the type (holotype). As the final reviser of a complex group embracing specimens from a large number of localities, some of which had been variously named, he exercised *his* right to select the type for such components as had not already been so designated.

The case of Dermacentor andersoni seems to be simpler still.

Specimens of this form do not appear to have been known by Neumann in 1897, at least he does not mention Montana specimens as being among the material examined by him, and *D. andersoni* is consequently not involved in the revision of Neumann's *D. reticulatus* (of 1897). The name appears before 1908 only as a nomen nudum and consequently does not concern us until that year when it is briefly characterized by Stiles (Weekly Publ. Health Rep., vol. 23, pt. 2, Nos. 27 to 52, July 3, 1908m, p. 949) and said to be based on specimens from Montana. Specimens from the latter State were first mentioned by Banks in June, 1908, and by him included in his complex *D. venustus*. In 1910, a definite type specimen of *D. andersoni* was published by Stiles, viz., U. S. P. H. & M. H. S. 9467. This specimen is from Woodman, Montana; host, Equus caballus.

The subsequent discussion between Banks and Stiles as to what specimens in the museums were actually designated as types of *D. venustus*, but which had never been so designated in any publication, seems to me irrelevant.

The published record of the two forms and their gradual fixation nomenclatorially by the various revisers may be briefly summarized as follows:

D. VENUSTUS

1897. Component of the complex D. reticulatus Neumann (no type designation).
1908. Component of the complex D. venustus Banks (no type designation).
1910. Segregated from D. venustus Banks 1908 and type designated by Stiles:
Marx No. 122.

D. ANDERSONI

1908. June. Montana specimens (not named) included in the complex D. venustus Banks (no type designation).

1910. August. Type designated by Stiles.

I am, therefore, of the opinion that the answer to Dr. W. Dwight Pierce's communication should be:

- I. That the Commission as such is incompetent to express an Opinion as to the name of the spotted fever tick. It can only take cognizance of the systematic names which have been applied to the various forms mentioned by him, and decide as to their applicability under the Code as disclosed by the records before the Commission.
- 2. On basis of these records it appears that, assuming the taxonomic distinctness of these forms,
 - a. The name *Dermacentor venustus* Marx in Neumann 1897 belongs to a form with the specimen Marx No. 122, from Texas as holotype.
 - b. The name *Dermacentor andersoni* Stiles 1908 belongs to a form with specimen U. S. P. H. & M. H.S. 9467, from Woodman, Montana, as the holotype.

Opinion written by Stejneger.

Opinion concurred in by 11 Commissioners: Allen, Apstein, Bather, Loennberg, Handlirsch, Hoyle, D. S. Jordan, K. Jordan, Monticelli, Skinner, and Stejneger.

Opinion dissented from by two Commissioners: Horváth and Kolbe.

Horváth states: "Je n'accepte que la seconde partie de la proposition, celle qui se rapporte au nom de *Dermacentor andersoni* Stiles, 1908. En ce qui concerne la première partie de la proposition, l'auteur de *Dermacentor venustus* est, à mon avis, incontestablement Banks qui en a publié en 1908 la première description. *D. venustus* Marx in

Neumann 1897 est un nomen nudum, puisque ni Marx, ni Neumann n'en ont donné une description. Le principe statué par l'Opinion 4 et appliqué dans l'Opinion 53 est inadmissible et doit être rejeté comme tout-à-fait contraire aux lois fondamentales de la nomenclature zoologique."

Monticelli states: "I cannot agree with the *first* point of the opinion of Stejneger from which, according to my judgment, a contradiction results,

"As the Commission must, on the basis of the conclusions of the relator, determine the nomenclature of the two species of *Dermacentor* (as results from the *second* point of the same conclusions by the wide discussion of the case presented for the examination of the Commission), I think that the Commission cannot declare its incompetence to express an opinion on Dr. Pierce's question. I think, therefore, that the Commission could well give its opinion on the specific name of the species of *Dermacentor* which transmits 'spotted fever' to man.

"Because, having fixed the two specific names, *Dermacentor venustus* Marx, 1897, and *Dermacentor andersoni* Stiles, 1908, and having identified with these names all the other names that different authors have attributed to the ticks of 'spotted fever,' it seems to me that—from the elimination of the names by which the relator has arrived at the *second* point of his conclusions—the specific name of the *Dermacentor* that gives spotted fever logically should result.

"It only remains to identify which of the two species of *Dermacentor* is the intermediate host of the parasite of 'spotted fever.'

"2. I agree, however, to the second point of the conclusions of the relator."

Not voting, two Commissioners: Hartert, Stiles.

OPINION 79

Case of Lamarck's (1801a) Système des Animaux sans Vertèbres

SUMMARY.—" Rigidly construed," Lamarck's (1801a) Système des Animaux sans Vertèbres is not to be accepted as designation of type species.

STATEMENT OF CASE.—Dr. J. Chester Bradley has submitted to the Commission the following question:

Is the Système des Animaux sans Vertèbres of Lamarck, 1801 to be accepted as designating types of genera?

In the work cited, Lamarck, after the description of each genus, cites at least one species, frequently two, which would at first sight appear to be mere examples and not construable as designated types.

But on p. viii he states: "Pour faire connoître d'une manière certaine les genres dont je donne ici les caractères, j'ai cité sous chacun d'eux une espèce connue, ou très-rarement plusieurs, et j'y ai joint quelques synonymes que je puis certifier; cela suffit pour me faire entendre."

This work was not accepted by Rohwer of nor by Viereck in their careful attempts to fix the types of the genera of sawflies and of ichneumonwasps. It has been accepted in a paper by Morice & Durrant, but these authors accept several works that clearly do not fix generic types in the sense of the Code.

Discussion.—In another Opinion (No. 81, on Cinex) the Commission has not interpreted this book by Lamarck as fixing types, and no new evidence is now presented which appears to warrant the reversal of this interpretation. In the view of the Commission, Lamarck cites a "known species, or very rarely several" as examples, in order to illustrate the genera, but rigidly construed, he does not fix the types.

¹Lamarck, Jean Baptiste Pierre Antoin de Monet *chevalier de*. Système des animaux sans vertèbres, ou Tableau général des classes, des ordres et des genérès de ces animaux Par J. B. Lamarck Paris, Deterville, An ix—1801, viii, 452, p. fold. tables, 20 cm.

² Rohwer, Sievert Allen II. The genotypes of the sawflies and woodwasps, or the superfamily *Tenthredinoidea*. By S. A. Rohwer Washington, 1916, < Technical series, No. 20, part H., U. S. Department of Agriculture, Bureau of Entomology.

⁸ Viereck, Henry Lorenz . . . Type species of the genera of ichneumonflies. By Henry L. Viereck 1914, < Smithsonian Institution, U. S. National Museum, Bulletin 83.

^{&#}x27;Morice, F. D. & John Hartley Durrant. The authorship and first publication of the "Jurinean" genera of Hymenoptera: Being a reprint of a long lost work by Panzer, with a translation into English, and introduction, and bibliographical and critical notes < Trans. Ent. Soc. Lond. 1914: 339-436.

This interpretation is supported by an examination of Lamarck's (1816b) Hist. Nat. des Anim. sans Vertéb., in which he does not even cite certain species mentioned in 1801. For instance, in 1801, p. 293, he cites only *P. rufipes* under *Pentatoma*; if he had intended this as type designation, he would, presumably, have cited this species under *Pentatoma* in 1816b, 492-494, but he does not do so; he stated that *Pentatoma* contains a large number of species, of which he cites three: acuminata, baccarum and prasina.

Opinion written by Stiles.

Opinion concurred in by 14 Commissioners: Allen, Apstein, Bather, Handlirsch, Hartert, Hoyle, Jordan (D. S.), Jordan (K.), Kolbe, Loennberg, Monticelli, Skinner, Stejneger, and Stiles.

Opinion dissented from by two Commissioners: Horváth, Dautzenberg.

Not voting, two Commissioners: Roule, Simon.

Dautzenberg says: "A l'époque où Lamarck a publié son Système des Animaux sans Vertèbres, on n'attachait pas à la fixation des types des genres l'importance ni la précision que nous lui attribuons aujourd'hui. En désignant pour chaque genre une espèce connue, accompagnée de reférences, 'afin de se faire bien entendre,' Lamarck a certainement voulu désigner ce que nous appelons aujourd'hui des types, aussi ne verrais-je aucun inconvénient en ce qui concerne les mollusques, à adopter pour types les espèces citées connue exemples dans le Système des An. sans vert., car il ne s'agit, en somme que de deux mots différents, mais qui ont exactement la mème signification."

17

OPINION 80

Suspension of Rules in the Case of Holothuria and Physalia

SUMMARY.—The Echinoderm genus Holothuria Linn., 1767, restr. Bruguière, 1791, type H. tremula 1767 = H. tubulosa 1790, and the Siphonophorae genus Physalia Lamarck, 1801, type P. pelagica 1801 = Holothuria physalis 1758, are hereby placed in the Official List of Generic Names.

STATEMENT OF CASE.—Pages 49 to 57 of Opinion 76 are accepted as statement of Case.

Discussion.—Pages 49 to 57 of Opinion 76 are accepted as Discussion.

The fact that the suspension of the rules was under consideration for these names was duly published as follows: Science, 1917, v. 45, Feb. 2, p. 113; Nature, Lond., v. 98, 1916, Sept. 21, p. 49; Monit. Zool. Ital., 1917, v. 28 (11), p. 183.

The Secretary recommends that the Commission adopt the following action:

- (1) Suspend the rules in the case of the generic names *Holothuria* and Physalia;
- (2) Permanently reject Holothuria Linnaeus, 1758, type H. physalis 1758;
- (3) Validate Physalia Lamarck, 1801, type P. pelagica 1801 (syn. H. physalis 1758);
- (4) Accept Holothuria as dating from Linn., 1767a (type H. tremula 1767=H. tubulosa 1790) as restricted by Bruguière, 1791, and despite the publication of Holothuria Linn., 1758 (rejected);
- (5) This suspension is not to be construed as invalidating any specific name.

The grounds for said suspension are:

- (a) In the judgment of the Commission, the strict application of the Règles to the names Holothuria and Physalia "will clearly result in greater confusion than uniformity";
- (b) The cases involve a transfer of generic names, almost universally accepted in the sense given above since 1701 (for Holothuria), and since 1801 (for Physalia), to genera in other groups in connection with which they have been used during more than 100 years by only a very few authors. Important supergeneric names, also of long standing, are involved.
- (6) The Commission places on the Official List of Generic Names the name Holothuria Linn., 1767, type H. tremula 1767 = H. tubulosa

1790, as the correct name for a genus of Sea Cucumbers, and the generic name *Physalia* Lamarck, 1801, type *P. pelagica* 1801 = *Holothuria physalia* 1758, as the correct generic name for the Portuguese Man-of-War.

Opinion written by Stiles.

Opinion concurred in by 12 Commissioners: Apstein, Bather, Handlirsch, Horvath, Hoyle, Jordan (D. S.), Jordan (K.), Kolbe, Loennberg, Skinner, Stejneger, Stiles.

Opinion dissented from by no Commissioners.

Not voting, three Commissioners: Dautzenberg, Hartert, Monticelli.

OPINION 81

THE GENOTYPE OF CIMEX, ACANTHIA, CLINOCORIS, AND KLINOPHILOS

SUMMARY,—On basis of the premises before the Commission, the common bedbug of Europe, Cimcx lectularius, is the genotype for Cimcx 1758, Acanthia 1775, Clinocoris 1829, and Klinophilos 1899 (Clinophilus 1903), and its proper technical designation under the Rules is Cimcx lectularius. Cimex Linn., 1758, type C. lectularius is hereby placed in the Official List of generic names.

Presentation of Case.—Dr. W. Dwight Pierce has submitted the following case for opinion. (Additions by the Secretary are marked *):

The scientific name of the bedbug has proved one of the most confusing problems in entomological nomenclature. It appears to the writer that the proper name should be *Clinocoris lectularius* Linnaeus, as accepted by Girault, Kirkaldy, and Reuter, and used in some medical text books (Castellani and Chalmers).

In American literature it also passes under the generic names Cimex and Acanthia.

In 1758 Linnaeus (Syst. Nat., 10th edit., p. 441) described *Cimex* with 85 species, of which *lectularius* was first and *stockerus* second. The genus was described as having four wings, but *lectularius* is wingless and does not agree with the generic description. No type is designated by Linnaeus.

Dr. C. W. Stiles in 1907 (Proc. Ent. Soc. Wash., vol. 8, p. 67, 68) considers that *lectularius* must be considered type because of Linnaeus' rule to select the commonest and most medicinal species as type of his genera. Such a method of selection, it seems to me, would be valid if there were no definite designations of type preceding Dr. Stiles' paper. The evidence presented below is against the acceptance of Dr. Stiles' designation.

In 1775, Fabricius (Syst. Ent. p. 696) discusses Cimex, and includes 167 species with "stockerus" Linnaeus as the first species, and he describes (p. 693) Acanthia with 15 species, of which (Cimex) lectularius Linnaeus (= Acanthia lectularia) is first. This action by Fabricius definitely removes lectularius from Cimex. (* No type was designated.—C. W. S.)

In 1789, Oliver (Encycl. Meth., vol. 4, Intr., p. 25) reversed Fabricius' division of genera, and called *Acanthia* Fabricius "Cimex" (Punaise), and called Cimex Fabricius "Pentatoma." From this date begins the confusion.

In 1797, Latreille ((* 1796a,) Précis des Caractères, p. 85) in discussing Acanthia says, "Je ne rapporte à ce genre que les espèces de Fab. que l'on trouve ordinairement aux bords des eaux. Les autres appartiennent aux Genres Coré et Lygé." (* Latreille (1796a, 83) cites "Cimex Linn. Punaise s. Pentatoma, Oliv." No type is selected, no species mentioned.—C. W. S.)

Kirkaldy in 1899 (The Entomologist, vol. 32, p. 219) considers Latreille's remarks to definitely limit the genus *Acanthia* to *littoralis* and its allies. Accepting this interpretation of Latreille's action, we must concede that *lectularia* was definitely eliminated from *Acanthia* in 1797.

(* Cuvier, 1798a, 574-575 (Tableau élémentaire de l'histoire naturelle des animaux) says:

Les punaises (Cimex)

On les divise en

- a. Acanthies (Mentions only "C. lectularius.")
- b. Punaises proprement dites. (Cimex Fabr.) (4 sp., lineatus, haemorrhoidalis, oleraceus, ornatus.)
 - c. Corées. (Corcus Fabr.). (Mentions only marginatus.)
 - d. Lygées. (Lygaeus Fabr.). (2 sp.)
 - e. Gerres. (Gerris Fabr.)
 - f. Hydrometre. (Hydrométra Latr.) (Mentions only stagnorum.)
 - g. Reduves. (Reduvius.) (Mentions only personatus.)

(* The question arises whether the expression, "Punaises proprement dites (Cimex Fabr.)," when 4 species are cited, constitutes a restriction that affects the type designation. The Secretary is inclined to the view that even if this point were conceded, the type is not designated thereby either for Acanthia or for Cimex, and that while it might have been better under the circumstances to follow this division subsequent to 1798, we cannot alter the fact that this course was not uniformly followed. We must take the facts as they exist, not as they should or might have been.)

For the next few years we find the species in ever shifting positions, none of which can really be accepted if we view elimination as a legitimate process in limiting a generic concept.

Schellenburg in 1800 (Cimicum Helvetiae Genus, pp. 5, 6, 15, 16) in a monograph of the Cimicidae has both genera *Cimex* and *Acanthia*, and places *lectularia* in *Acanthia* (* but does not designate types).

(* Lamarck, 1801a (Syst. anim. sans vertèbres, pp. 293-294) adopts Cimex Linn. as genus, which he divides as follows:

"Corps ovale ou arrondi. (Acanth. Fab.)

"Cimex lectularius, Lin. Acanthia lectu—(p. 294) laria. Fab. Ent. 4, p. 67. Geoff. ins. 1, p. 434, n. 1. La punaise des lits.

"Cet insecte incommode et puant, n'a ni ailes ni élytres par un avortement qui se perpétue, et propage dans un état qui réssemble à celui de larve. Néanmoins sa classe et son genre sont determinés par la considération de ses congénères.

"Corps oblong, un peu étroit. (Ligaei, Fab.)

"Cimex equestris. Lin. Ligaeus equestris. Fab. ent. 4, p. 147. Climex. Geoff. ins. 1, p. 442, no. 14.")

(* On page viii, Lamarck says: "Pour faire connoître d'une mannière certaine les genres dont je donne ici les caractères, j'ai cité sous chacun d'eux une espèce connue, ou très rarement plusiers, et j'y ai joint quelques synonymes que je puis certifier; cela suffit pour me faire entendre.")

(* Thus while Lamarck clearly intended C. lectularius to be considered as a Cimex, he recognized two subgroups (Acanthia and Lygaeus), placing C. lectularius in the subgroup Acanthia. If his remarks on page viii (see above) are to be interpreted as definite designation of genotypes for the genera in which only one species is cited, it would appear that lectularius is here designated type of Acanthia. Since, however, he did not name one of his subgroups as Cimex s. str., it would appear that either Acanthia or Lygaeus should be interpreted as the typical subgroup, hence as Cimex s. str., hence also that

either *lectularius* or *equestris* should be type of *Cimex*. As this point is not definite from the context, it must be concluded that "rigidly construed" (Art. 30g), Lamarck did not here designate type for *Cimex*.)

(*Linnaeus (1802, Turton Ed., Syst. Nat., pp. 608-702) divides Cimex into six groups (cf. subgenera) as follows: I. Cimex (Aeanthia) which includes lectularius, littoralis and many other species; 2. Cimex (Cimex) which includes bidens and many other species; and four other groups which do not influence the present problem, namely, 3. Cimex (Lygacus); 4. Cimex (Gerris); 5. Cimex (Miris); and 6. Cimex (Reduvius). Types are not cited for these groups, but is is to be noticed that both lectularius and littoralis are placed in Aeanthia, and it is clear that a typical subgenus Cimex has been created, but as no type is designated this seems to leave the subject in the same status as did "Punaises proprement dites. (Cimex Fabr.)" of Cuvier, 1798a. So far as Aeanthia is concerned, the status of affairs has reverted to that which existed in 1775.)

Fabricius in 1803 (Syst. Rhyng., p. 112-113, 155-179) treats both Acanthia and Cimex and limits Acanthia to lectularia and hemiptera. Kirkaldy (1899, The Entomologist, vol. 32, p. 220) is very positive in asserting that Fabricius in this work designates bidens L. as type of Cimex. It is true that bidens is the first Linnaean species included in the Fabrician concept of Cimex, but I cannot find a positive designation.¹

Latreille in 1804 (Hist. Nat. Crust. et Ins. p. 237, 240-244, 254-255) definitely states ² on p. 237 that he reversed the Fabrician decisions and makes *lectularius* type of "punaise," which is his common name for Cimex, and on page 254-255 limits Cimex to lectularius. He places in Acanthia, zosterae, littoralis and four other species.

(*Dumeril, 1806, 264 (Zool. analytique) appears definitely to designate *lectularius* as type (by monotypy) of *Cimex*. The passage in question reads:

"2. Les punaises (cimex, Linne; acanthia Fab.) ont le corps ovale, très applati, cinq articles aux antennes, et le corcelet en croissant recevant la tête. On n'en a encore observé qu'une seule espèce, qui attaque pendant la nuit l'homme et certains oiseaux, en particulier les hirondelles.)

(* Dumeril (1806, 262) adopts Acanthia for species, not mentioned by name, which live on banks of bodies of water, on bark of trees, and on fruits.)

(* Latreille, 1807 (Gen. Crust. et insect), p. 136 mentions only C. lectularius under Cimex, and cites (p. 142) A. maculata, Lygaeus saltatorius, Salda littoralis, S. zosterae, and S. striata, under Acanthia.)

Latreille in 1810a (Consid. Gen., p. 433) in the list which is considered as designating types by an Opinion (* No. 11) of the International Commission, designates lectularia as type of Acanthia, thus contradicting his positive state-

¹ (* Fabricius, 1803, 112, cites *lectularia* (chef de file) and *hemiptera* as belonging to *Acanthia*, and p. 155-170 he cites 123 species (without type designation (See Art. 30r) for *Cimex*); (bidens is chef de file).—C. W. S.)

²P. 237: "Il nous a paru plus convenable de restituer à cet insecte le nom sous lequel il est généralement connu, et de le faire servir de type au genre punaise (* Cimex, p. 254), dont il est jusqu'à présent la seule espèce bien connue."

ments of 1797, 1804, and even on previous pages in the same book. He refers Cimex to Pentatoma.¹

(*On p. 434 he cites "Lygaeus saltatorius" as type of "Acanthic" (Acanthia).)

(* Lamarck 1816b, 501-503, clearly designates lectularius as type for Cimex, for though he cites two species (lectularius and hirundinis) the second (hirundinis) is not an original (1758) species, and he says "Par les nombreuses distinctions etablies, le genre: punaise (Cimex) se trouve presque (cf. hirundinis) reduit à la seule espèce (lectularius) qu'on eut souhaite ne jamais connaître." Under Acanthia he includes maculata, littoralis, and zosterae, but without type designation.)

Fallen in 1818 (Cimices Sveciae, p. 17, 27) has 18 species in Cimex and limits Acanthia to lectularia. (* Not a type designation—C. W.S.)

In 1825 Saint Fageau and Serville (Encycl. Meth., vol. 10, p. 250-251) follow Olivier in placing *lectularius* as the only (* positive) species in *Cimex*.

Fallen in 1829 (Hem. Svec., p. 140, 142) limits Acanthia to lectularia but suggests Clinocoris² as a better generic name. This is the first time that lectularia has had a bona-fide location since 1797. (* Fallen includes bidens and 17 other species in Cimex.—C. W.S.)

(* The publication by Fallen, 1829, brings up a very complicated combination of nomenclatorial possibilities.)

(*(a). It is clear that *Clinocoris* ($\hat{\eta}$ κλίνη a couch; $\hat{\delta}$ κόριs, a bug) 1829 is *Acanthia* (ἀκαιθίαs, a prickly thing) renamed, hence (Art. 30f, rule) "the type of either, when established, becomes ipso facto type of the other.")

(*(b). The first definite type designation for *Acanthia* was *Lygaeus saltatorius* (by Latreille, 1810a, 434), but as this was not an original species for *Acanthia* it is not available as type.)

1904: A. lectularia is apparently accepted as type by Kirkaldy, 1904, Nature, 465; 1905; and by Reuter, 1908, Ent. mon. Mag. 27.

1912: Cimex lectularius is definitely designated as type by Castellani & Chalmers, 1913, 637 and 1920, 763.

1917: C. lectularius is definitely accepted as type by Van Duzee, 1917, 285.

(* The only species (See dissenting view by Stejneger in Discussion) which can possibly come into theoretical consideration as genotype both of Acanthia and of Clinocoris are: A. lectularia and A. clavicornis; all theoretical arguments are in favor of accepting lectularia which is the only one of the two species which has ever been definitely cited by name in connection with Clinocoris and which is the first and only species ever designated as type of Clinocoris. Accordingly, unless it can be shown that clavicornis has been designated type of Acanthia, lectularia remains type of Clinocoris and therefore type of Acanthia also.)

¹ (* Latreille, 1810a, p. 257 says: "G. 324, Punaise, Cimex." and on p. 433 he says: "Punaise, Acanthia lectularia." Thus lectularius is designated type of Cimex.)

² Acanthia renamed. "Nomen generis ab $a\kappa ar\theta \acute{a}$ (spina) desumsit Cel. Fabricius, verisimiliter propter punctionem insecti. Forsitan convenientius judicabitur nomen Clinocoris (Germanice Bettwanze). I. A. lectularia.

(* Curtis, 1835 (Brit. Ent. vol. 12, pl. 548, 569) says: 548: "Acanthia Type of the Genus, Cimex littoralis Linn." and 569: "Cimex Type of the Genus, Cimex lectularius Linn.")

(*Westwood, 1840, vol. 2, Synopsis, p. 110, designates saltatoria Linn. as type of Acanthia , and p. 120 C. lectularius as type of Cimex; but saltatoria

is not cited as an original species by Fabricius in 1775.)

In 1843 Amyot & Serville (Hist. Nat. Ins. Hemiptères, p. 310-313) give a good discussion of the case in hand, stating that Fabricius by dividing Cimex into three genera definitely removed lectularius to Acanthia. They attribute all our present difficulties to Olivier's (1789) arbitrary reversal of the Fabrician genera calling Acanthia Fabr. "Cimex," and Cimex L., Fabr. "Pentatoma." They further recite Latreille's reversals of opinion in 1797 and later, first accepting Acanthia for lectularia and later Cimex. They treat Acanthia with only lectularia.

(*Reuter (Wien, Ent. Zeitung, 1882, 301-306) discusses the case in detail and accepts *lectularius* as type of *Cimex*; on basis of Fabricius (1803) he accepts *littoralis* as type of *Acanthia*. He argues that Fabricius (1803) definitely designated types by his method of comparison (chef de file).)

In 1899, Kirkaldy (The Entomologist, p. 219) overlooking *Clinocoris*, and considering the bedbug without a generic name, proposed *Klinophilos* (* tod. *Cimcx lectularius*, and he took *bidens* Linn., as type of *Cimcx.*—C. W. S.).

(*Blanford (1903, Nature, 200) changes Klinophilos to Clinophilus and adopts lectularius as type of Cimex on basis of the Linnaean rules. Kirkaldy (1904, Nature, 465), replying to Blanford, claims that (on basis of elimination) lectularius is excluded from being taken as type of Cimex and that Latreille (1707) restricted Acanthia to "littoralis and its congeners"; Kirkaldy accepts Clinocoris, instead of his Klinophilos, for the bedbug. Blanford (1904, Nature, 464), replies that the generic name was taken from a species in the Linnaean genus that was called Cimex in classical Latin. The only species that can be clearly identified with the Latin name appears to be C. lectularius L. and he accepts this as type of Cimex on basis of the Linnaean rules.)

In 1905, Kirkaldy (The Entomologist, vol. 38, p. 76, 78) withdrew Klino-

philos, accepting Clinocoris, and gave further proof on pp. 304-306.

In 1908, Reuter (Ent. mon. Mag., vol. 44, p. 27) reviewed the situation and agreed with Kirkaldy (1899) that littoralis should be type of Acanthia, bidens type of Cimex, and lectularius of Clinocoris.

Kirkaldy, 1909 (Cat. Hemiptera (Heteroptera) vol. 1, p. xxvi-xxviii), again insists that Fabricius 1803 named bidens as type of Cimex, but says that Latreille 1804 named (zosterac Latr.) = saltatorius L. as the type of Acanthia.

(* Apstein, 1915a, 158, (Nomina Conservanda) designates lectularius as

type of Cimex.)

(*Van Duzee (1917, Catalog. Hemipt., 285) accepts lectularius as type of Cimex on basis of Lamarck (1801a, 293), Latreille (1810a, 257, 433), Laport (1832, 51) and Westwood (1840), all of whom he quotes as "names lectularius type." He also accepts lectularia as type of Acanthia on basis of Fabr. (1803,

¹Reuter quotes (in part erroneously) Kirkaldy, 1899, p. 219, as follows: "I therefore see no alternative to adopting the name Acanthia for littoralis (*& c." in original of K. but omitted by R.—C. W. S.) as Kirkaldy has already done in his monograph of the palaearctic species."

112). The Secretary does not accept Laport (1832, 51) and Fabr. (1803, 112) as definite type designation.)

As I see the synonymy at present, it may be summarized as follows:

1. Cimex Linnaeus 1758, type bidens L. selected according to Kirkaldy by Fabricius 1803, but at least by Kirkaldy 1899. The genus is limited by removal of Acanthia Fabricius 1775 thus taking away lectularius. Accepted as above by Reuter 1908.

2. Acanthia Fabricius 1775, type littoralis L. selected by Latreille 1804 according to Reuter 1908. The genus was limited to exclude lectularia by

Latreille 1707.

3. Clinocoris Fallen 1829—monotype lectularia L. The genus is offered as substitute for Acanthia Fabricius 1803, Fallen 1829 (not Fabricius 1775, Latreille 1797). Accepted by Kirkaldy 1899, 1905, 1909; Reuter 1908; Girault, 1905.

Synonyms:

(a) Acanthia Schellenberg, 1800; Fabricius, 1803, type by elimination lectularius; Latreille, type by designation, 1810; Fallen monotype, 1818; Fallen monotype, 1829; Douglass and Scott 1865.

(b) Cimex Latreille, 1804, type by designation lectularius; Stiles, 1907

(designation); E. Saunders, 1892; Lethierry & Severin, 1896.

(c) Klinophilos Kirkaldy, 1899, type by original designation lectularius.

Discussion.—The case submitted is one more to be added to the many cases of generic confusion due to the fact that so many authors have been content with division of genera, but have ignored the principle of genotype fixation. If authors had followed the Linnæan code in this case, and had, in accordance with said code,¹ adopted *C. lectularius* as type of *Cimex* the confusion would have been automatically avoided.

The premises have been set forth by Dr. Pierce in the "Presentation of Case." In company with Dr. Pierce the Secretary has verified the references, but his interpretation of certain of the citations differs somewhat from that presented by Dr. Pierce. This case of nomenclatures has been discussed in more or less detail by a considerable number of authors and their views seem to be hopelessly at variance. No opinion the Commission adopts can count upon universal approval since so many complications, giving rise to different views, come into consideration. One principle develops in the case (see *Clinocoris*) which has never been before the Commission heretofore, which seems to be an entirely new principle, and yet one which seems to be clearly covered by the rules.

In addition to the literature cited by Dr. Pierce, the Secretary has consulted a number of other references which are briefly summarized

¹ The particular Linnæan rule in question reads "Si genus receptum, secundum jus naturae et artis, in plura dirimi debet, tum nomen antea commune manebit vulgatissimæ et officinali plantæ."

or cited herewith. As the Secretary sees the points at issue, they involve four generic names (Cimex, Acanthia, Clinocoris, and Klinophilos) and may be summarized as follows:

I. Cimex Linn., 1758a: Two species (lectularius and bidens) have been selected as type.

A. In the original publication the type is not determined under Art. 30—

(a) Original designation, (b) Use of typicus or typus, (c) Monotypy, or (d) Absolute tautonymy.

B. Neither species thus far designated as type (*lectularius* and *bidens*) is excluded under Art. 30(e).

C. No complication arises under Art. 30(f), renaming of genus.

D. In case of doubt, Recommendations (h to t), the following points are to be held in mind under Art. 30:

1758: C. lectularius (Cimex of Pliney) is on the preferred list under (h) the Linnaean rule, (n) best described, best figured, best known, and easily obtained species, (p) parasitic on man, (q) probably actually studied by author, (t) page precedence.

1775: *C. lectularius* would not be on the preferred list because (k) elimination by: Fabricius, 1775, 693; 1787, 280; 1794, 67; 1803, 112.—Cuvier, 1798.—Schellenberg, 1800, 15.—Turton, 1802.—Fallen, 1818, 19; 1829.—Burmeister, 1837a, 596.—Amyot & Serville, 1843.—Douglass & Scott, 1868, 278.—Claus, 1885a.—Leunis, 1886a.—R. Blanchard, 1890a, 473.—Railliet, 1895a, 820.—Kirkaldy, 1899; 1904, 465; 1905.—Reuter, 1908, 27.—And many others.

A. bidens seems to be on the preferred list under (k) because it remained in Cimex after A. lectularia was eliminated (1775) and (o) De Candolle's rule.

Apparently neither A. lectularia nor A. bidens has preference, one over the other, under (i) Virtual tautonymy, (j) non-exotic, (1) sexually mature vs. larvae, (m) name communis, etc., (s) Linnaeus did not declare in favor of the first species rule.

1803: C. bidens is on the preferred list under (r) as chef de file by Fabricius, 1803, 155.

E. "Rigidly construed" (Art. 30g) the following references are to be interpreted as citation of illustrative or characteristic species rather than as selection of type, or at best are debatable.

1764: C. lectularius by: Brunnich, 1764, 82 (see also p. 56).—Olivier, 1789, 25.—Lamarck, 1801a, 293.—Latreille, 1804, 254; 1807, 136.—St. Fagean & Serville, 1825.—DeLaporte, 1832, 51.—Stal, 1873, 104.—And many others.

1834: C. juniperinus by: Burmeister, 1837a, 597.

F. "Rigidly construed" (Art. 30g) the following references are undebatably definite designations of genotypes:

1804: *C. lectularius* by: Latreille, 1804, 254; 1810a, 257, 433.—Dumeril, 1806, 264.—Lamarck, 1816b, 502.—Curtis, 1835, 569.—Westwood, 1840, 120.—Pascoe, 1868, 94.—Reuter, 1882, 301.—Blanford, 1903, 200; 1904, 464.—Stiles, 1907, 67.—Apstein, 1915a, 158.—Van Duzee, 1917, 285.

1899: C. bidens by: Kirkaldy, 1899, 220; 1909, xxviii (on basis of Fabr. 1803), 4.—Reuter 1908.

G. Conclusion.—C. lectularius was the first original species definitely designated (1804) as type of Cimex in harmony with Art. 30 and this designation is not subject to change.

2. Acanthia Fabr. 1775: Four species (A. saltatoria, A. littoralis, A. sosterae, and A. lectularia) have been selected as type.

A. In the original publication, the type is not determined under Art, 30 (a, b, c, d).

B. Under Art. 30 (e. a), A. saltatoria is definitely excluded as type since it was not an original species. A. zosterae is not cited as an original species, and it was further considered later to be a synonym of saltatoria; accordingly, A. zosterae is definitely excluded as type.

C. A distinct complication arises because of the renaming of genus. Acanthia was renamed Clinocoris in 1829, hence under Art. 30(f) the type of either, when established, becomes, ipso facto, type of the other. As a natural result, no species which is excluded as type of one of these genera can come into consideration as type of the other, and as A. littoralis was definitely excluded from Clinocoris by the founder of the generic name, this species cannot (under Art. 30e, a) become type of Clinocoris, hence (Art. 30f),

dating with 1829 it is definitely excluded from consideration in selecting (Art. 30g) the type of *Acanthia*.

D. In case of doubt, the following points are to be held in mind:

1775: A. lectularia is on the preferred list under (h) the Linnean rule, (n) best known, etc., (p) parasitic on man, (q) probably actually studied by author, and (t) page precedence.

1789: A. lectularia would not be on the preferred list because of (k) elimination by: Olivier, 1789, 25.— Dumeril, 1806, 262.—Latreille, 1804; 1807; 1810a.— Lamarck, 1816b, 502.—St. Fagean & Serville, 1825.— DeLaporte, 1832, 51.—Curtis, 1835.—Westwood, 1840.—Stal, 1873, 104.—Reuter, 1882, 301; 1908, 27.— Kirkaldy, 1809; 1904; 1905.—Blanford, 1903; 1904.— Stiles, 1907.—Apstein, 1917a.—Van Duzee, 1917.— And many others.

1803: A. littoralis would not be on the preferred list because of (k) elimination by: Fabricius, 1803, 115, to Salda.—Fallen, 1829, 71.

A. littoralis seems to be on the preferred list under (o) DeCandolle's rule.

Apparently neither A. lectularia nor A. littoralis is on the preferred list under (i) Virtual tautonymy, (1) Sexually mature vs. larvae, (m) name communis, etc., (s) Fabricius did not declare in favor of the first species rule.

1803: A. lectularia is on the preferred list under (r) as chef de file by Fabricius, 1803, 112.

E. "Rigidly construed" (Art. 30g) the following references, are to be interpreted as citation of illustrative or characteristic species rather than selection of type, or at best are debatable.

1796: A. littoralis group by: Latreille, 1796a, 185; 1804, 240.—Dumeril, 1806.—Lamarck, 1816b, 508.—Kirkaldy, 1904, 465.

1798: A. lectularia by: Cuvier, 1798a, 574.—Schellenberg, 1800. 15.—Lamarck, 1801a, 293.—Fallen, 1818, 17, 27; ? 1829, 140.—Burmeister, 1837a, 596—Amyot & Serville, 1843, 310.—Douglass & Scott, 1868, 278.—Claus, 1885a.—Leunis, 1886a.—Knauer, 1887a,

339.—R. Blanchard, 1890a, 473.—Railliet, 1895a, 820.—And many others.

1832: A. saltatoria by: DeLaporte, 1832, 52.

F. "Rigidly construed" (Art. 30g) the following references are undebatably definite designations of genotypes.

1810: A. saltatoria by: Latreille, 1810a, 259, 434.—Westwood, 1840, 119.—Kirkaldy, 1909, xxviii (on basis of Latreille, 1804).

1835: A. littoralis by: Curtis, 1835, 548.—Reuter, 1882, 301 (on basis of Fabr. 1803); 1908, 26-27 (on basis of Kirkaldy, 1899, 218).

1868: A. zosterae by: Pascoe, 1868, 94-95 (on basis of Latr. 1802; 1804).—Kirkaldy, 1909, xxviii (so. saltatorius) (on basis of Latreille, 1804) (chef de file of Salda by Fabr., 1803, 113).

1917: A. lectularia by: Van Duzee, 1917, 285 (on basis of Fabr., 1803, 112).

G. CONCLUSION: A. lectularia is type because it is the first and only original species (Art. 30e, α) of both Acanthia and Clinocoris which has been validly designated as type either of Acanthia or of Clinocoris (see C).

3. Clinocoris (Petersson? in) Fallen, 1829, Acanthia Fabricius renamed hence both must have the same genotype. C. lectularius is the only species which has been definitely designated as type.

A. On basis of the original publication it is possibly a debatable point but very doubtful whether the type is determined under (a) original designation, but it is not determined under (b, c, or d).

B. C. lectularius is available under Art. 30 (e).

C. Complications arise under Art. 30 (f) as Clinocoris is Acanthia renamed. The following 7 of the 15 original species of Acanthia are definitely excluded (under 30 e, a) from consideration as type of Clinocoris, since Fallen (1829) himself definitely excluded them by not including them in Clinocoris and by classifying them elsewhere: A. betulae (in Aradus), A. cardui (in Tingis), A. corticalis (in Aradus), A. laevis (in Aradus), A. littoralis (in Salda), A. pyri (in Tingis), A. rugosa (in Aradus).

C'. Commissioner Stejneger holds another view as follows: The fact brought out by Dr. Stiles in the rewritten Opinion, that Fallen, in 1829, simultaneously with suggesting Clinocoris as a substitute for Acanthia, placed A. littoralis

of Fabricius in another genus, *Salda*, can have no influence on Curtis's right, in 1835, to designate it as type of *Acanthia* Fabricius.

As shown above, Acanthia, up to the year 1829, had not any valid type designation, and was consequently still polytypic. Fallen in this year did not alter the status of Acanthia; he only mentioned lectularia as one of the species, but gave a substitute name, Clinocoris. Consequently, Clinocoris at that date was equally polytypic, and must share the fate of Acanthia. It now appears that on the same occasion he also relegated Acanthia littoralis to another genus, Salda. The question then arises: Does this action of Fallen in placing A. littoralis in another genus nullify Curtis' explicit designation, in 1835, of littoralis as the type of Acanthia? Is there anything in Code Art. 30 which makes this action of Curtis invalid? These questions, it seems to me, have already been answered in Opinion 62 which specifically provides that Article 30 does not even exclude type species of other genera from consideration in the subsequent selection of the type of a given genus. The fact that Fallen removed littoralis to another genus, Salda, consequently does not bar its designation by Curtis in 1835, since even if he had made it the type of Salda (and so he may have done for all I know) that fact would not have invalidated the designation of littoralis as type of Acanthia. Fallen, in 1829, did not make a new genus Clinocoris, he only suggested a new name for an old genus, and this substitute name must ipso facto have the same designated type. If littoralis is the type of Salda, Salda also becomes a synonym of Acanthia.

D. In case of doubt, the following points are to be held in mind under Recommendations (h to t) of Art. 30:

1829: C. lectularius is on the preferred list under (h, n, p, q, and t).

1829: *C. lectularius* (known as κόρις by Aristophanes; κόρις ἀπὸ κλίνης by Discorides), is to be selected ("unless such preference is strongly contraindicated by other factors") under (i) Virtual tautonymy: ἡ κλίνη, a couch; *lectulus*, a little bed; ὁ κόρις, a bug.

? 1829: Acanthia lectularia by Monotypy, by Fallen, 1829, 141. This is open to debate. Certain it is that this is the species which Fallen had especially in mind. A difference of opinion seems, however, inevitable, as

theoretical arguments exist on both sides. Hence, rigidly construed, this designation or alleged designation might perhaps best be tabled.

1829: C. lectularius is on the preferred list under (j) as a non-exotic species, when compared with the following 6 of the 8 remaining original species (not mentioned above in C) of Acanthia; A. crassipes (Dresden); A. lunata (India); A. rhombea (Africa); A. scrrata (hab. unknown); A. serratulae (England); A. umbraculata (Hafniae).

1829: Acanthia clavicornis, the one remaining original species of Acanthia which comes into theoretical competition has nothing (under Art. 30) to give it preference over A. lectularia.

E. "Rigidly construed" (Art. 30g), it is not clear that Girault (1905, 61, 117) designates the genotype.

F. "Rigidly construed" (Art. 30g), the following references are undebatably definite designations of genotype.

1904: *C. lectularius* by: Kirkaldy, 1904, 465; 1905.—Reuter, 1908, 27.—Castellani & Chalmers, 1913, 637; 1920, 763.—Van Duzee, 1917, 285.

G. CONCLUSION.—C. lectularius was the first and only original species of *Clinocoris* definitely designated as type of *Clinocoris* in harmony with Art. 30 and this designation is not subject to change.

4. Klinophilos Kirkaldy, 1899=Clinophilus Blanford, 1903. 1899: lectularius type by monotypy (Art. 30c).

As soon as one departs from the foregoing citations to which the Rules can be strictly applied one encounters citations that are subject to interpretations that are diametrically opposed to each other and one becomes involved in the uncertainties of elimination, retransfer, and recliminations, and in the vagaries involved in the citation of a single species as example.

Accordingly, the Secretary recommends that the Commission adopt as its Opinion the following:

1. On basis of the premises before the Commission, the common bed-bug of Europe, Cimex lectularius Linn., 1758, is genotype for Cimex Linn., 1758, Acanthia Fabr., 1775, Clinocoris Petersson or Fallen, 1829, and Klinophilos Kirkaldy, 1899 (= Clinophilus Blanford, 1903), and its proper designation under the rules is Cimex lectularius.

2. Cimex Linn., 1758, type C. lectularius, is hereby placed in the Official List of generic names.

Commissioner Stejneger presents the following dissenting conclusion which is presented for vote as alternative Opinion:

I am therefore constrained to maintain that my original conclusions were correct as formulated in my first vote to the effect:

(1) That lectularius Linn., 1758, is the type of Cimex; (2) that Klinophilus of Kirkaldy, 1899, is a synonym of Cimex with the same type; (3) that Acanthia of Fabricius, 1775, has for type Cimex littoralis; (4) that Clinocoris of Fallen, 1829, is a synonym of Acanthia with the same type.

Opinion written by Stiles.

Opinion as written by Stiles concurred in by ten Commissioners: Allen, Handlirsch, Hartert, Horvath, Hoyle, Jordan (D. S.), Kolbe, Monticelli, Skinner, Stiles.

Opinion as modified by Stejneger (but accepting *lectularius* as type of *Cimex*) concurred in by one (or two?) Commissioners: Stejneger, ?Bather.

Opinion dissented from by one Commissioner: Jordan (K.).

Not voting on opinion as now written (see, however, footnote, p. 31) five Commissioners: Apstein (accepts *lectularius* as type of *Cimex*), R. Blanchard (deceased; prior to death he accepted *lectularius* as type of *Cimex*) Dautzenberg (accepts *lectularius* as type of *Cimex*), Roule, Simon.

The essential point is that 14 Commissioners have concurred in accepting *lectularius* as type of *Cimex* as against one Commissioner who dissents from this view.

Bather adds: "I do not accept Stiles' argument, p. 26, C. I am doubtful as to the validity of all of Stejneger's remarks, p. 28, C'. I incline to think that this is a case in which one should frankly give up argument and decide either on ground of practical convenience or by drawing lots. From first to last an amount of time must have been wasted on this bed-bug enough to decide the fate of six alleged murderers. Is it worth while?"

Handlirsch adds: "Wenn Cimex in dem Sinne 'lectularius' beibehalten wird und Salda für littoralis etc., so fällt endlich der Name

¹The Opinion as written in Circular Letter No. 36 was:

Concurred in by 14 Commissioners: Allen, Apstein, Bather, Blanchard, Dautzenberg, Handlirsch, Hartert, Horvath (part), Hoyle, Jordan (D.S.), Monticelli, Skinner, Stejneger (part), Stiles.

Dissented from (in part) by 2 Commissioners: Horvath, Stejneger. Not voting, 4 Commissioners: Jordan (K), Kolbe, Roule, Simon.

Acanthia, der so viel Confusion verursacht hat, und alle Zweifel sind endgültig beseitigt. Das its ja schliesslich doch die Hauptsache."

Hoyle adds: "On reading this re-statement of the case, the following points occur to me: (1) That the action of Linné in placing 'lectularius' as first species in 'Cimex,' taken in conjunction with his method of selecting types is almost sufficient to make 'lectularius' the type of 'Cimex,' though perhaps it does not justify the phrase 'rigidly construed.' (2) However this may be, it seems to me that Latreille (1804) definitely makes 'lectularius' the type of 'Cimex' and this action overrules any preceding subdivisions and eliminations. I, therefore, see no reason to reverse my previous opinion."

Jordan (D. S.) adds: "I should have taken Stejneger's view, but not insistently as the case is excessively complex."

Jordan (K.) adds: "I. As a matter of principle the original diagnosis of a genus should be considered first guide in determining the type species of the genus. If the original author, by the wording of his diagnosis, indicates from which kind of species the diagnosis is taken, this indication has priority over all subsequent ones. E. g., Hübner describes his genus Heraclia (Lepid.) as having 'glossy green black' forerings, and places into this genus three species, of which two agree with the description, while the third does not. Obviously, the type of the genus is one of the 'glossy green black' species. Similarly, Cimex is diagnosed by Linnaeus as having four wings; his conception of a true Cimex, therefore, was a four-winged insect. The bed-bug does not conform with this conception. Therefore, I cannot accept lectularius as type of Cimex. But something might be said in favor of discarding priority (or suspending the rules) in this important case."

"II. Acanthia Fabr., 1775, was based on a number of species inclusive of the bed-bug. The diagnosis of the genus seems to cover all species, being very general (and faulty). In 1794 Fabricius gave a fuller diagnosis of Acanthia, stating 'elytris coriaceis, planis, apice membranaceis longitudine abdominis. . .', but he, nevertheless, leaves lectularius in this Acanthia. Latreille in 1797 limits Acanthia to the species found near water. Both Fabricius in 1794 and Latreille in 1797 place the bed-bug outside the concept of true Acanthia, and I submit that from 1794 lectularius had no valid generic name.

"III. In 1803 Fabricius reversed his conception of 1794 and restricted Acanthia to the bed-bugs. He was not entitled to do so. This concept of 1803 and not the Acanthia Fabr., 1775, was renamed Clinocoris by Fallen in 1829. I consider Clinocoris to be the first valid generic term for lectularius."