

SMITHSONIAN MISCELLANEOUS COLLECTIONS

VOLUME 99, NUMBER 8

CHECK-LIST OF THE TERRESTRIAL AND  
FRESH-WATER ISOPODA  
OF OCEANIA

BY

HAROLD GORDON JACKSON

Department of Zoology, Birkbeck College,  
University of London



(PUBLICATION 3593)

CITY OF WASHINGTON  
PUBLISHED BY THE SMITHSONIAN INSTITUTION  
JULY 23, 1941



SMITHSONIAN MISCELLANEOUS COLLECTIONS  
VOLUME 99, NUMBER 8

CHECK-LIST OF THE TERRESTRIAL AND  
FRESH-WATER ISOPODA  
OF OCEANIA

BY

HAROLD GORDON JACKSON

Department of Zoology, Birkbeck College,  
University of London



(PUBLICATION 3593)

CITY OF WASHINGTON  
PUBLISHED BY THE SMITHSONIAN INSTITUTION  
JULY 23, 1941

The Lord Baltimore Press  
BALTIMORE, MD., U. S. A.

# CHECK-LIST OF THE TERRESTRIAL AND FRESH-WATER ISOPODA OF OCEANIA \*

By HAROLD GORDON JACKSON

*Department of Zoology, Birkbeck College,  
University of London*

## INTRODUCTION

The geographical classification of the Pacific island groups used in this check-list is that employed by the Bernice P. Bishop Museum, and is as follows:

Polynesia: Austral, Chatham, Cook, Easter, Ellice, Hawaii (Hawaii to Kure), Kermadec, Line (Palmyra to Jarvis), Manihiki, Marquesas, New Zealand, Phoenix, Samoa, Society, Tokelau (Union), Tonga, Tuamotu.

Melanesia: Admiralty, Bismarck Archipelago, Fiji, Loyalty, New Caledonia, New Hebrides, Santa Cruz, Solomon, Trobriand.

Micronesia: Bonin, Caroline, Gilbert, Marianas (Guam), Marshall, Palau, Volcano, Wake.

Records from Juan Fernandez and the Galápagos Islands have not been included. (For these see Van Name, 1936, pp. 40 and 41.)

References are, as a rule, restricted to Oceania, but a reference or synonym which seemed in danger of being overlooked is included even though it ranged wider.

The types of the genera dealt with below have been nominated in relatively few cases and, in case of doubt, I have chosen the earliest described species which has been most frequently recorded, or which has page preference in the original description. The type locality is given in each case as nearly as the original description allows.

Such a list as this must, perforce, contain a number of species of doubtful validity, as many of the earlier descriptions, especially of the Armadilline genera, do not suffice to identify the original species. Anyone who works on Pacific species must acknowledge his debt to Verhoeff, but I cannot follow him in all his subdivisions of *Philoscia*.

---

\* Acknowledgment is made of contributions toward the cost of publishing this paper from the Publication Committee of the University of London, the Royal Society of London, the British Association for the Advancement of Science, and the Pacific Entomological Survey.

and *Ligia* and have reduced many of his genera to subgenera in these groups. I agree with Van Name (1936) that the division of *Philoscia* into a number of full genera "involves losing sight of important resemblances and relationships in the effort to emphasize small differences," and I consider that the same applies to *Ligia*. There is no doubt, however, that large numbers of forms which still fall under the genera *Spherillo* and *Cubaris* must be further subdivided or reduced by synonymy.

The history of the nomenclature of these forms must be briefly considered in order to explain the divisions adopted in the following pages. Those which were known at the time were placed by Brandt (1833) in three genera: *Armadillo* (*A. officinalis*), *Cubaris* (*C. murina*, *cinercea*, and *brunnea*), and *Diploexochus* (*D. echinatus*). Budde-Lund (1885 and 1904) grouped all under *Armadillo* and in 1904 resurrected *Spherillo* Dana for a great army of new but related forms. However, many authors, rejecting both these names, made use of *Cubaris* for the whole group.

It is obvious that such a large assemblage of forms showing considerable diversity of character under their common likeness of structure cannot continue to be known under one or two all-embracing generic names, and the first step in subdividing them must be to go back to Brandt's three clearly defined genera. Here we are fortunately on firm ground (provided that the name *Armadillo* is retained<sup>1</sup>), as the types are well known and undoubtedly are "good" species, but if confusion is to be avoided it is essential that the genera be closely and strictly founded on the named types (for example, Budde-Lund defined *Diploexochus* very much wider than the type and has been followed by other authors).

Probably only one of Brandt's genera (*Cubaris*) is a genuine Pacific genus, but there is a great assemblage of related forms which Budde-Lund lumped under the one genus *Spherillo*.

This complex of forms has yet to be completely sorted, but Verhoeff (1926) is responsible for the first serious attempt to impose order and it seems probable his genera will stand. Unfortunately he has not defined them in relation to the tentative sections (Budde-Lund, XII, Wahrberg, XVI) which were already set up. It is clear that his genus *Sphaerillo* (which is a homonym of *Spherillo* Dana according to Article 35) contains the species placed by Budde-Lund under section XIII of *Spherillo* (Wahrberg's section XVI), and it would only intensify confusion to disagree with this now. If we seek

<sup>1</sup> See Jackson, 1933b, p. 148 for further discussion.

for a type for this section it appears that we must nominate *Spherillo danae* Heller 1868, which Heller wrongly attributed to Dana's genus. Of the remaining sections of *Spherillo*, VIII is now divided between *Australiodillo* Verhoeff and *Hawaiiodillo* Verhoeff, X is *Merulana* Budde-Lund, and XI (Wahrberg XIII) is *Merulanella* Verhoeff. The remaining sections await analysis, although most of them are probably entitled to generic rank.

Verhoeff's remaining genera would appear to fall outside the already described species of Budde-Lund's composite genus *Spherillo*, but his *Nesodillo* (by Verhoeff's admission, 1938) contained *Cubaris murina* Brandt (*Armadillo murinus* Budde-Lund) and the genera are therefore synonyms. The genus *Cubaris* is thus clearly defined by Brandt and Verhoeff and must be used only in this restricted sense. The rejection of *Nesodillo* is fortunate in another way, as it avoids any confusion with the same author's *Mesodillo*.

Past and present workers on the assemblage of terrestrial isopods under consideration have so consistently neglected to apply the International Rules of Nomenclature that it is too late to do more than protest formally and, in the interest of clarity, accept the status quo. On these grounds it is suggested that *Sphaerillo* Verhoeff be retained as the generic name of the forms included under Budde-Lund's section XIII (Wahrberg XVI) and that *Spherillo* should be allowed to die out, as its species are absorbed into new or already existing genera.

The origin of the land isopod fauna of Oceania is a problem upon which speculation may easily outrun the available data, but the following facts are of interest. About 155 species have been recorded from Pacific Islands and of these 139 are peculiar to this area. Of the 16 recorded from elsewhere 6 are cosmopolitan, 6 are also recorded from Malaysia and the Indian Ocean and 4 from South America. Of the latter, 1 (*Trichoniscus magellanicus*) is Antarctic, 2 are semimarine (*Ligia novaezealandiae* and *Doto bucculenta*) and present no problem of transport, and the remaining 1 (*Rhyscotus ortonedae*) is known by a single specimen recorded from Samoa and is an obvious introduction. From the fact that records of the *Ligia* and *Doto* are numerous from New Zealand and its islands and solitary from Chile it is reasonably probable that the former locality is the home of both. If they had been natives of Chile they might be expected to be more widely spread along the American coast.

The credentials of the western fauna are more respectable. *Philoscia angusticauda* is recorded from Borneo and Hawaii; *Cubaris murina*, although first recorded from Brazil, is widely distributed in Malaysia

and the Indian Ocean; *Spherillo testudinalis* is from the Bay of Bengal; *Philoscia truncata* is from Malaysia; and *Trichoniscus verrucosus* is from the Crozets (South Indian Ocean). Surprisingly few are shared with Australia, but it is not surprising that the Polynesian area shows no relationship with Hawaii, which is relatively poor in terrestrial isopods.<sup>2</sup>

Of this large residue of species more than half belong to the Armadillinae and about 50 of these to the conglobating forms included by Budde-Lund under the name "Spherillo." These forms may be considered to be the characteristic woodlice of the whole area, and it is significant that such a large assemblage of forms with short bushy penicilli on the maxillula are grouped together in the Pacific basin, even if the fact does not justify Budde-Lund's creation of a subfamily to contain them.

The only other group of species which approaches the Armadillinae in numbers is that comprised under the general name "Philoscia," of which there are 28 recorded.

In spite of the richness of species in Oceania, there are few records of endemic genera. Each group shows considerable uniformity, and specific and even generic differences are founded on minute points of no great moment. It would be reasonable therefore to postulate a recent origin for the majority of the woodlice of Oceania.

The terrestrial isopods, in spite of this wide distribution, would seem to be singularly ill-provided with means of dispersal. They are dependent on the maintenance of a high humidity, they have small powers of locomotion and must await the accidental carriage of a gravid female for colonizing distant places. Even the smallest are too large and unprotected to be carried easily by birds, like seeds or small eggs, except for very short distances, but transportation by hurricanes, perhaps with portions of plants, is by no means out of the question. The powers of distribution possessed by wind storms are shown, in this area, in records which "are marvellous to the point almost of embarrassment" (Gulick).<sup>3</sup> Usually terrestrial isopods must rely on driftwood or the cooperation of man in spreading over the remoter parts of the earth.

Between the groups of islands driftwood is probably the most usual means of dispersal, which can take place in any direction in spite of the prevailing trade wind. The British Admiralty sailing instructions takes particular notice of "singular interruptions to the trade

<sup>2</sup> Verhoeff, however, records *Ligia hawaiiensis* from Fiji (1938).

<sup>3</sup> Gulick, A., Quart. Rev. Biol., vol. 7, p. 415, 1932.

wind, unknown currents and strong squalls" among these islands, which might assist the random distribution of drift and these conditions have presumably persisted *mutatis mutandis* for the greater part of the Tertiary period, prior to which we have neither knowledge of the existence of terrestrial isopods nor reliable information on the conformation of land in the Pacific. There is certainly no confirmation of the existence of the NW. to SE. land chains, which are admittedly geologically possible, in the present distribution of the Pacific woodlice, or any indication that they traveled there by land.

It is significant in this connection that a large proportion of woodlice from the Pacific Islands belong to the Armadillinae—that is to say to a group which is strikingly adapted to resist desiccation but with very poor powers of locomotion. They may well have survived transportation by natural means from early times, and it may be assumed that the great variety of species of *Philoscia*, which, with their lightly calcified integument, exposed ventral surface, and low birth rate are ill-formed for rigorous conditions, have appeared with the free introduction of plants characteristic of man's activities, which gives the woodlouse a means of conveyance under conditions of the requisite high humidity.

It is believed to be highly probable that Polynesia was peopled by races which came from the west via India and Malaysia, and they may have been responsible for the woodlice with western affinities. Later dispersals in both directions were possibly made by the seventeenth- and eighteenth-century explorers of the Pacific and later still by the copra trade, which followed the same general route as the earliest settlers.

Speculation on the distribution of the Oniscoidea is inevitably checked by our ignorance of the rate at which modification may have taken place and, though the ancestral form of the "Spherillo" group may have had its origin in the land masses on the western border of the Pacific basin bounded by the line New Zealand—Tonga—Fiji—New Hebrides—Solomon and have spread east and west from island to island by natural means of dispersal, it is also within the bounds of possibility that they have not preceded man. Nevertheless, it must not be forgotten that two of the most modified forms, *Tridentodillo* and *Echinodillo*, come from the most isolated group of islands, the Marquesas, at considerable altitudes. I have suggested elsewhere (1933b) that these may be gerontic forms preserved from extinction by their inaccessible habitat.

The "Philoscias" more probably appeared with the earliest settlers in much more recent times, although *Cerberoides* from Rapa (Aus-

trals), whose affinities are with the Oniscinae, can scarcely be accounted for in that way. Its three subgenera provide a beautiful example of adaptive radiation, which, one feels, must have had more leisure to work itself out than the advent of man could have provided.

Finally, there is a group of sub-Antarctic forms—*Actaecia*, *Scypho-niscus*, *Scyphax*, and *Decto*—distinctly lower in the evolutionary scale, which have made their appearance in the extreme south and which contribute nothing to the wider problem.

To sum up: The woodlice of Oceania for the most part consist of forms belonging to the Armadillid or Oniscid type of structure. The Porcellionidae and the more terrestrial families of the Atracheata are almost unrepresented except by obvious recent introductions. These statements are almost equally applicable to the great land masses bordering the Pacific, but the fauna of Oceania is less rich than they in characteristic genera. Endemic species are certainly numerous, but the range of differences does not encourage the belief that they have been separated from one another for great periods of time, but rather that variations have occurred freely, after random population of the islands, under conditions which encourage their appearance and survival.

The assumption of land connections between the islands is gratuitous and would seem to provide more time than is necessary to account for the comparatively small range of differences observed, while the assumption that man is responsible would seem to provide too little.

There remains random distribution by hurricanes and other accidental means, which is probably adequate to explain the present distribution and to account for the greater individuality of more remote islands such as the Marquesas and the Australs.

## Suborder ONISCOIDEA

### Superfamily ATRACHEATA

#### Family LIGIIDAE

##### Genus LIGIA Fabricius, 1792

(Type, *Oniscus oceanicus* Linné)

*Ligia (Megaligia) exotica* ROUX: Roux, 1828, pl. 13; Budde-Lund, 1879, p. 8; 1885, p. 266; Dollfus, 1893, p. 24; Richardson, 1905, p. 676 (*Ligyda*); Jackson, 1922, p. 693; Panning, 1924, p. 196; Verhoeff, 1926, p. 348 (*Megaligia*); Jackson, 1927b, p. 11; Verhoeff, 1928a, p. 116; Van Name, 1936, p. 48 (q.v. for further references).

Synonyms: *L. gaudichaudi* Milne-Edwards, 1840, p. 157.

*L. grandis* Perty, 1830-34, p. 212.

? *L. malleata* Pfeffer, 1889, p. 36 (Jackson, 1922, p. 701).

Hawaii (Oahu); Samoa (Tutuila); New Zealand.

(Type locality, Mediterranean.)

**Ligia (Megaligia) hawaiensis** DANA: Dana, 1853, p. 740; Budde-Lund, 1885, p. 271; Jackson, 1922, p. 696; Verhoeff, 1928a, p. 116 (*Megaligia*); Jackson, 1933b, p. 149; Van Name, 1936, p. 52; Verhoeff, 1938, p. 12.

Hawaii (Oahu and Kauai); Fiji (Viti Levu).

(Type locality, Hawaiian Islands.)

**Ligia (Megaligia) perkinsi** (DOLLFUS): Dollfus, 1900, p. 525 (*Geoligia*); Jackson, 1922, p. 699; Chilton, 1922, p. 83; 1924a, p. 894; Verhoeff, 1926, p. 354; Jackson, 1927a, p. 134; 1927b, pp. 3 and 11.

Hawaii (Hawaii, Kauai); Samoa (Namua).

(Type locality, Hawaiian Islands.)

**Ligia (Nesoligia) novaezealandiae** DANA: Dana, 1853, p. 739 (*Lygia novaezealandiae*); Miers, 1876b, p. 103; Filhol, 1885, p. 444; Budde-Lund, 1885, p. 271; Thomson and Chilton, 1886, p. 157; Chilton, 1901, p. 106; 1910b, p. 287; 1911, p. 568; Jackson, 1922, p. 697; Chilton, 1924b, p. 287; Panning, 1924, p. 196; Verhoeff, 1926, p. 348 (*Nesoligia*); Barnard, 1932, p. 185; Van Name, 1936, pp. 54, 57 (? = *L. litigiosa* Wahrberg, 1922b, p. 277).

Synonyms: *L. quadrata*, Thomson, 1879a, p. 232; Filhol, 1885, p. 444.

*L. porteri*, Maccagno, 1931, p. 151.

*L. cursor* Budde-Lund (see Jackson, 1922, p. 697).

Stewart Island; New Zealand; Sunday Island, Victoria; Chile.

(Type locality, Bay of Islands, New Zealand.)

**Ligia (Euryligia) latissima** VERHOEFF: Verhoeff, 1926, p. 349.

(Type locality, New Caledonia.)

**Ligia rugosa** JACKSON: Jackson, 1938, p. 172.

Tuamotus (Mangareva); Australs (Marotiri).

(Type locality, Tuamotus [Mangareva].)

**Ligia philoscoides** JACKSON: Jackson, 1938, p. 173.

(Type locality, Australs [Rapa].)

**Ligia pallida** JACKSON: Jackson, 1938, p. 175.

(Type locality, Christmas Island.)

**Ligia vitiensis** DANA: Dana, 1853, p. 741; Budde-Lund, 1885, p. 271; Stebbing, 1900b, p. 646; Jackson, 1922, p. 696 (*L. hawaiensis*); 1933b, p. 149; 1938, p. 172.

Fiji; New Guinea; Marquesas (Eiao); Tuamotus (Mangareva).

(Type locality, Fiji.)

#### Genus STYLONISCUS Dana, 1858

(Type, *Styloiniscus longistylis* Dana)

**Styloiniscus longistylis** DANA: Dana, 1853, p. 737; Budde-Lund, 1885, p. 271.

(Type locality, Tongatabu [Friendly].)

## Family TRICHONISCIDAE

## Genus TRICHONISCUS Brandt, 1833

(Type, *Trichoniscus pusillus* Brandt)

Synonym: *Philougria* Bate and Westwood, 1868.

**Trichoniscus (Megatrichoniscus) thomsoni** CHILTON: Chilton, 1886, p. 159 (*Philygria*); 1901, p. 118; Budde-Lund, 1906, p. 83; Chilton, 1909a, p. 661; Wahrberg, 1922a, p. 79; Stephensen, 1927, p. 371; Jackson, 1938, p. 176 (*Megatrichoniscus*).

Auckland Islands; New Zealand; Australs (Rapa).

(Type locality, South Island, New Zealand.)

**Trichoniscus (Megatrichoniscus) magellanicus** (DANA): Dana, 1853, p. 736 (*Styloniscus*); Budde-Lund, 1885, p. 271 (*Styloniscus*); Stebbing, 1900a, p. 566 (*Trichoniscus*); Chilton, 1901, p. 106 (*Styloniscus*); Budde-Lund, 1906, p. 83; Chilton, 1909a, pp. 602 and 661; 1910b, p. 287; 1915b, p. 453; Wahrberg, 1922a, p. 76; Stephensen, 1927, p. 370; Van Name, 1936, p. 82. New Zealand.

(Type locality, Tierra del Fuego, Nassau Bay.)

**Trichoniscus phormianus** CHILTON: Chilton, 1901, p. 115.

Synonym: *Philongria rosca* Chilton, 1883a, p. 73; 1883c, p. 149; Thomson and Chilton, 1886, p. 157.

New Zealand.

(Type locality, Canterbury, New Zealand.)

**Trichoniscus otakensis** CHILTON: Chilton, 1901, p. 117; 1910b, p. 287; ? Wahrberg, 1922a, p. 76; Chilton, 1925a, p. 319; Stephensen, 1927, p. 370.

Synonym as for *T. phormianus*.

South Island, New Zealand; Chatham Island.

(Type locality, South Island, New Zealand.)

**Trichoniscus commensalis** CHILTON: Chilton, 1910a, p. 191.

New Zealand, "North Island and wider" (from ants' nests).

(Type locality, New Plymouth, New Zealand.)

**Trichoniscus verrucosus** BUDDE-LUND: Budde-Lund, 1906, p. 79; Chilton, 1909a, p. 602; 1910b, p. 287; 1915b, p. 453 (? = *T. magellanicus*); Wahrberg, 1922a, p. 72; Van Name, 1936, p. 83.

Sub-Antarctic islands of New Zealand and Marion Island.

(Type locality, Crozets.)

**Trichoniscus kermadecensis** CHILTON: Chilton, 1911, p. 569.

Kermadec Islands.

(Type locality, Sunday Island, Kermadec Islands.)

## Genus NOTONISCUS Chilton, 1915

(Type, *Haplophthalmus helmsii* Chilton)

**Notoniscus helmsii** (CHILTON): Chilton, 1901, p. 119 (*Haplophthalmus*); 1915a, p. 418.

New Zealand.

(Type locality, Greymouth, New Zealand.)

**Notoniscus australis** (CHILTON): Chilton, 1909a, p. 662 (*Haplophthalmus*); 1915a, p. 421.  
 (Type locality, Campbell Island.)

## Superfamily HYPOTRACHEATA

### Family TYLIDAE

**Genus TYLOS** Audouin and Savigny, 1827

(Type, *Tylos latreillei* Audouin and Savigny)

**Tylos neozelandicus** CHILTON: Chilton, 1901, p. 120; Budde-Lund, 1906, p. 78; Chilton, 1910b, p. 288.  
 New Zealand.  
 (Type locality, Wellington, New Zealand.)

## Superfamily PLEUROTRACHEATA

### Family SCYPHACIDAE (=DETONIDAE)

**Genus SCYPHAX** Dana, 1853

(Type, *Scyphax ornatus* Dana)

**Scyphax ornatus** DANA: Dana, 1853, p. 734; Miers, 1876b, p. 101; Budde-Lund, 1879, p. 2; 1885, p. 233; Filhol, 1885, p. 443; Thomson and Chilton, 1886, p. 158; Chilton, 1901, p. 123.

Synonyms: *Scyphax intermedius* Miers: Miers, 1876a, p. 227; 1876b, p. 102; Thomson and Chilton, 1886, p. 158.

*Philoscia violacea* Filhol, 1885, p. 445.

"On sandy shores in the North Island and also from Westport. Not found in the south of South Island." (Chilton, 1901.)

(Type locality, New Zealand.)

**Scyphax setiger** BUDDE-LUND: Budde-Lund, 1885, p. 232.

(Type locality, New Caledonia.)

**Genus DETO** Guérin, 1836

(Type, *Deto echinata* Guérin)

**Deto (Vinetta) aucklandiae** (G. M. THOMSON): Thomson, 1879b, p. 249 (*Actaecia*); Budde-Lund, 1885, p. 239 (*Actaecia*); Filhol, 1885, p. 443 (*Actaecia*); Chilton, 1901, p. 126 (*Scyphax* ?); 1906b, p. 273; Budde-Lund, 1906, p. 87; Chilton, 1909a, p. 666; 1910b, p. 288; 1915b, p. 445; Stephensen, 1927, p. 371.

Synonyms: *Deto magnifica* Budde-Lund, 1906, p. 86.

*Deto robusta* Budde-Lund, 1906, p. 87.

(Type locality, Ewing Island in Auckland Islands group.)

**Deto (Vinetta) bucculenta** (NICOLET): Nicolet, 1849, p. 267 (*Oniscus* ♂); Budde-Lund, 1885, p. 206 (*Oniscus*); Chilton, 1915b, p. 449.  
 Synonyms: *Oniscus tuberculatus* (?) Nicolet, 1849, p. 268; Budde-Lund, 1885, p. 206.  
*Oniscus novae-zealandiae* Filhol, 1885, p. 441; Budde-Lund, 1906, p. 87.  
*Scyphax* (?) *aucklandiae* Chilton, 1901, p. 126.  
*Deto novae-zealandiae* Chilton, 1906b, p. 273; Budde-Lund, 1906, p. 87; Chilton, 1909a, p. 667; 1910b, p. 288.  
 Chatham Island, Stewart Island, Wellington, New Zealand.  
 (Type locality, Valparaiso Bay, South America.)

#### Genus SCYPHONISCUS Chilton, 1901

(Type, *Scyphoniscus waitatensis* Chilton)

**Scyphoniscus waitatensis** CHILTON: Chilton, 1901, p. 128; 1910b, p. 288.  
 (Type locality, Blueskin Bay, Otago, New Zealand.)

**Scyphoniscus magnus** CHILTON: Chilton, 1909a, p. 665.  
 (Type locality, Auckland and Campbell Islands.)

#### Genus ACTAECIA Dana, 1853

(*Cylloma* Budde-Lund, 1885)

(Type, *Actaccia euchroa* Dana)

**Actaecia euchroa** DANA: Dana, 1853, p. 735; Miers, 1876b, p. 102; Thomson, 1879b, p. 249; Filhol, 1885, p. 443; Thomson and Chilton, 1886, p. 158; Thomson, 1893, p. 56 (12 in reprint); Chilton, 1901, p. 130; 1910b, p. 288.  
 Synonym: *Armadilloniscus euchroa* Budde-Lund, 1885, p. 239.  
 New Zealand.  
 (Type locality, Bay of Islands, New Zealand.)

**Actaecia opihensis** CHILTON: Chilton, 1901, p. 132; 1910b, p. 288.  
 (Type locality, Timaru, New Zealand.)

#### Family ONISCIDAE

##### Subfamily ONISCINAE

#### Genus PHILOSCIA Latreille, 1804

(Type, *Oniscus muscorum* Scopoli)

**Philoscia australis** SEARLE: Searle, 1914, p. 365.  
 (Type locality, Marquesas [Nukuhiva].)

**Philoscia lifuensis** STEBBING: Stebbing, 1900b, p. 648; Budde-Lund, 1912, p. 386  
 (? *Setaphora*); Verhoeff, 1926, p. 356, ? = *P.* (*Chactophiloscia*) *rouxi*  
 Verhoeff.  
 (Type locality, Lifu, Loyalty Islands.)

**Philoscia novae-zealandiae** FILHOL: Filhol, 1885, p. 441; Chilton, 1901, p. 138; 1910b, p. 289.

(Type localities, New Zealand and Stewart Island.)

**Philoscia oliveri** CHILTON: Chilton, 1911, p. 570.

(Type locality, Sunday Island, Kermadec Islands.)

**Philoscia persona** JACKSON: Jackson, 1938, p. 178.

(Type locality, Australs [South East Islet, Marotiri].)

**Philoscia pubescens** (DANA): Dana, 1853, p. 730 (*Oniscus*) ; Miers, 1876b, p. 99 (*Oniscus*) ; Filhol, 1885, p. 440 (*Oniscus*) ; Budde-Lund, 1885, p. 223; Thomson and Chilton, 1886, p. 158 (*Oniscus*) ; Chilton, 1901, p. 136; Budde-Lund, 1904, p. 43; 1906, p. 71; Chilton, 1910b, p. 289; Barnard, 1932, p. 244.

Auckland.

(Type locality, Whykare River, Auckland, New Zealand.)

**Philoscia societatus** MACCAGNO: Maccagno, 1932, p. 4.

(Type locality, Society Islands [Moorea].)

**Philoscia squamosa** JACKSON: Jackson, 1938, p. 180; var. *setosa* Jackson, 1938, p. 181.

(Type locality, Australs [South East Islet, Marotiri].)

**Philoscia (Chaetophiloscia) rouxi** VERHOEFF: Verhoeff, 1926, p. 345 (*Laevo-philoscia*) ; Verhoeff, 1928b, p. 218 (*Laevophiloscia* = *Chaetophiloscia*) ? = *P. lisiensis* Stebbing q. v.

Loyalty Islands.

(Type locality, Maré, Loyalty Islands.)

**Philoscia (Heroldia) appressa** VERHOEFF: Verhoeff, 1926, p. 342.

New Caledonia.

(Type locality, Tchalabel, New Caledonia.)

**Philoscia (Heroldia) canalensis** VERHOEFF: Verhoeff, 1926, p. 342.

New Caledonia.

(Type locality, Panié Mountain, New Caledonia.)

**Philoscia (Heroldia) digitifera** VERHOEFF: Verhoeff, 1926, p. 338; 1926, p. 339 (var. *marmorata*).

New Caledonia.

(Type locality, Humboldt Mountain, New Caledonia.)

**Philoscia (Heroldia) humboldtii** VERHOEFF: Verhoeff, 1926, p. 340.

New Caledonia.

(Type locality, Humboldt Mountain, New Caledonia.)

**Philoscia (Heroldia) monticola** VERHOEFF: Verhoeff, 1926, p. 344; 1926, p. 345 (subsp. *paniensis*).

New Caledonia.

(Type locality, Ignambi Mountain, New Caledonia.)

**Philoscia (Heroldia) reticulata** VERHOEFF: Verhoeff, 1926, p. 339.

New Caledonia.

(Type locality, Ignambi Mountain, New Caledonia.)

**Philoscia (Nesoniscus) longicornis** VERHOEFF: Verhoeff, 1926, p. 329.

New Caledonia.

(Type locality, Ignambi Mountain, New Caledonia.)

**Philoscia (Nesoniscus) noduligerus** VERHOEFF: Verhoeff, 1926, p. 330; 1926, p. 331 (subsp. *minor*).

New Caledonia.

(Type locality, Ignambi and Panié Mountains, New Caledonia.)

**Philoscia (Oroscia) paniensis** VERHOEFF: Verhoeff, 1926, p. 328.

New Caledonia.

(Type locality, Panié, New Caledonia.)

**Philoscia (Oroscia) squamuligera** VERHOEFF: Verhoeff, 1926, p. 325.

New Caledonia.

(Type locality, Ngoi Tal and Humboldt Mountain, New Caledonia.)

**Philoscia (Paraphiloscia) brevicornis** BUDDE-LUND: Budde-Lund, 1912, p. 374 (*Pseudophiloscia*).

(Type locality, Auckland, New Zealand.)

**Philoscia (Paraphiloscia) fragilis** (BUDDE-LUND): Budde-Lund, 1904, p. 43 (*Pseudophiloscia*).

Synonym: *Philoscia pubescens* Chilton, 1901, p. 136, *ex parte*.

Auckland and Howick, New Zealand.

(Type locality, Howick, New Zealand.)

**Philoscia (Paraphiloscia) gracilis** BUDDE-LUND: Budde-Lund, 1879, p. 2; 1885, p. 220; Stebbing, 1900b, p. 647; Budde-Lund, 1904, p. 42 (*Pseudophiloscia*); 1912, p. 372 (*Pseudophiloscia*); *ibid.* note by Stebbing, p. 372 (*Paraphiloscia*); Jackson, 1927b, p. 8; 1938, p. 178 (var. *australis*).

Lifu, Loyalty Islands ?; Samoan Islands; Australs (Tubuai).

(Type locality, Upolu, Samoan Islands.)

**Philoscia (Paraphiloscia) stenosoma** STEBBING: Stebbing, 1900b, p. 648.

(Type locality, New Britain.)

**Philoscia (Plymophiloscia) montana** VERHOEFF: Verhoeff, 1926, p. 334.

New Caledonia.

(Type locality, Canala and Humboldt Mountains, New Caledonia.)

**Philoscia (Setaphora) angusticauda** BUDDE-LUND: Budde-Lund, 1885, p. 216; Dollfus, 1900, p. 525; Budde-Lund, 1912, p. 386 (*Setaphora*); Arcangeli, 1927, p. 259; Jackson, 1936, p. 81.

Hawaii (Honolulu, Oahu).

(Type locality, Borneo.)

**Philoscia (Setaphora ?) fasciata** JACKSON: Jackson, 1933a, p. 87; 1933b, p. 151; 1938, p. 181.

Marquesas (Tahuata, Nukuhiva, Hivaoa, Fatuhiva, Uahuka); Tuamotus (Mangareva, Henderson); Pitcairn; Australs (Tubuai, Rapa, Rurutu, Raivavae); Society Islands (Tahiti, Moorea).

(Type locality, Marquesas.)

**Philoscia (Setaphora) truncata** DOLLFUS: Dollfus, 1898, p. 376; Stebbing, 1900b, p. 647; Budde-Lund, 1912, p. 386 (? *Setaphora*) ; Arcangeli, 1927, p. 260; Herold, 1931, p. 383; Jackson, 1933b, p. 150; 1938, p. 177.

New Britain; Marquesas (Fatuiva, Nukuhiva, Uapou); Tuamotus (Mangareva); Australs (Tubuai, Rapa); Society Islands (Tahiti, Moorea, Huahine, Raiatea).

(Type locality, Celebes and Flores.)

**Philoscia (Wahrbergia) sarasini** VERHOEFF: Verhoeff, 1926, p. 333.

New Caledonia.

(Type locality, Ignambi and Canala Mountains, New Caledonia.)

#### Genus **ONISCUS** Linnaeus, 1758

(Type, *Oniscus asellus* Linné)

**Oniscus cookii** FILHOL: Filhol, 1885, p. 442; Chilton, 1901, p. 136 (? = "Oniscus kenepurensis"); 1910b, p. 288. "Sous les pierres sur la portion ouest de l'île du milieu de la Nouvelle-Zélande" (Filhol).<sup>4</sup>

#### Genus **PHALLONISCUS** Budde-Lund, 1908

(Type, *Oniscus punctatus* Thomson)

**Phalloniscus punctatus** (THOMSON): Thomson, 1879a, p. 232 (*Oniscus*); Budde-Lund, 1885, p. 206 (*Oniscus*); Filhol, 1885, p. 440 (*Oniscus*); Thomson and Chilton, 1886, p. 158 (*Oniscus*); Chilton, 1901, p. 133 (*Oniscus*); 1906b, p. 273 (*Oniscus*); 1909a, p. 668 (*Oniscus*); 1910b, p. 288 (*Oniscus*); Wahrberg, 1922a, p. 91 (*Phalloniscus*); Chilton, 1925a, p. 319; Verhoeff, 1926, p. 331; Stephensen, 1927, p. 372.

New Zealand; Chatham Island; Auckland Islands.

(Type locality, New Zealand.)

**Phalloniscus kenepurensis** (CHILTON): Chilton, 1901, p. 135 (*Oniscus*); 1910b, p. 288 (*Oniscus*); Wahrberg, 1922a, p. 86 (*Phalloniscus*); Verhoeff, 1926, p. 331.

(Type locality, Kenepuru, New Zealand.)

#### Genus **ALLONISCUS** Dana, 1854

(Type, *Alloniscus perconvexus* Dana)

**Alloniscus oahuensis** BUDDE-LUND: Budde-Lund, 1879, p. 1; 1885, p. 225; Jackson, 1933b, p. 153; 1938, p. 181.

Synonym: *Alloniscus brevis* Budde-Lund, 1885, p. 226; Jackson, 1927b, p. 6 (see Jackson, 1933b, p. 154).

Samoa (Tutuila, Upolu); Marquesas (Hivaoa, Mohotani, Uahuka); Hawaii (Oahu); Tuamotus (Mangareva); Australs (Tubuai, Raivavae); Society Islands (Tahiti); Fanning Island.

(Type locality, Oahu, Hawaiian Islands.)

<sup>4</sup> This is almost certainly not an "*Oniscus*." Chilton's suggestion is probable, and in any case it is likely to be a "*Phalloniscus*."

**Alloniscus floresianus** DOLLFUS: Dollfus, 1898, p. 374; 1900, p. 524.  
 Hawaii (Lanai).  
 (Type locality, Flores.)

Genus **CERBEROIDES** Jackson, 1938

(Type, *Cerberoides pilosus* Jackson)

**Cerberoides (Philoscodillo) pilosus** JACKSON: Jackson, 1938, p. 183.  
 (Type locality, Australs [Rapa Island].)

**Cerberoides (Oniscomorphus) bicornis** JACKSON: Jackson, 1938, p. 184.  
 (Type locality, Australs [Rapa Island].)

**Cerberoides (Congloboniscus) brevicauda** JACKSON: Jackson, 1938, p. 187.  
 (Type locality, Australs [Rapa Island].)

Subfamily **PORCELLIONIINAE**

Genus **PORCELLIO** Latreille, 1804

(Type, *Porcellio scaber* Latreille)

**Porcellio (Euporcellio) scaber** LATREILLE: Latreille, 1804, p. 45; Dollfus, 1900, p. 524; Chilton, 1901, p. 139; 1905, p. 429; Jackson, 1938, p. 189.

Synonym: *Porcellio graniger* Miers, 1876a, p. 226; 1876b, p. 99; Filhol, 1885, p. 443; Thomson and Chilton, 1886, p. 158; Thomson, 1892, p. 48. For further references and synonymy see Van Name, 1936, p. 226.

New Zealand; Hawaii (Kona); Australs (Rapa).  
 (Type locality, Europe.)

**Porcellio (Mesoporcellio) laevis** LATREILLE: Latreille, 1804, p. 46; Dollfus, 1890b, p. 4; 1900, p. 524; Chilton, 1905, p. 430; Wahrberg, 1922a, p. 182; Verhoeff, 1926, p. 321; Jackson, 1933b, p. 155; 1938, p. 187. For further references and synonymy see Van Name, 1936, p. 229.

Hawaii (Lanai, Oahu); New Caledonia; Marquesas (Taiohae); Tuamotus (Mangareva).  
 (Type locality, Europe.)

Genus **PORCELLIONIDES** Miers, 1877 (= **METOPONORTHUS** Budde-Lund, 1885)

(Type, *Porcellio pruinosis* Brandt, 1833)

**Porcellionides pruinosis** (BRANDT): Brandt, 1833, p. 181 (*Porcellio*); Chilton, 1901, p. 141; 1905, p. 431; 1906a, p. 64; 1911, p. 571; Searle, 1914, p. 366; Wahrberg, 1922a, p. 148; Verhoeff, 1926, p. 321 (except Searle, all the above *Metoponorthus*); Jackson, 1933b, p. 155; 1938, p. 190; Verhoeff, 1938, p. 12 (*Metoponorthus*).

Synonyms: *Porcellio zealandicus* White, 1847, p. 99; Miers, 1876a, p. 226; 1876b, p. 100.

*Porcellio neo-secalanicus* Thomson and Chilton, 1886, p. 158; Chilton, 1901, p. 141.

For further references and synonymy see Van Name, 1936, p. 238.  
 New Zealand; Norfolk Island; Kermadec Islands (Sunday Island); Society Islands (Tahiti); Marquesas (Hivaoa, Mohotani Eiao, Fatuuku, Nuku-hiva); Tuamotus (Fakarava); Fanning Island; Fiji (Viti Levu).  
 (Type locality, Europe.)

**Genus NAGARA Budde-Lund, 1908**

(Type, *Porcellio cristatus* Dollfus, 1889)

**Nagara (Nagara) insularum** VERHOEFF: Verhoeff, 1926, p. 315.  
 (Type locality, Loyalty Islands.)

**Nagara (Heminagara) tahitiensis** JACKSON: Jackson, 1933a, p. 87; 1938, p. 189.  
 Society Islands (Tahiti, Raiatea); Tuamotus (Mangareva); Australs (Tu-buai, Maria); Christmas Island?  
 (Type locality, Society Islands [Tahiti].)

**Genus TRICHLORHINA Budde-Lund, 1908**

(Type, *Bathytrapa thermophila* Dollfus)

**Trichorhina squamata** VERHOEFF: Verhoeff, 1926, p. 319.  
 (Type locality, Loyalty Islands.)

Subfamily RHYSOTINAE

**Genus RHYSOTUS Budde-Lund, 1885**

(Type, *Rhysotus turgifrons* Budde-Lund)

**Rhysotus ortonedae** BUDDE-LUND: Budde-Lund, 1908, p. 299; Jackson, 1927b, p. 6; Van Name, 1936, p. 266.  
 Samoa (Upolu).  
 (Type locality, Ecuador.)

Subfamily ARMADILLINAE

**Genus ACANTHODILLO Verhoeff, 1926**

(Type, *Acanthodillo erinaceus* Verhoeff)

**Acanthodillo erinaceus** VERHOEFF: Verhoeff, 1926, p. 267.  
 (Type locality, New Caledonia.)

**Genus ARMADILLO Brandt, 1833 amend. Verhoeff, 1926**

(Type, *Armadillo officinalis* Duméril)

**Armadillo purpurescens** (BUDDE-LUND): Budde-Lund, 1912, p. 371 (*Spherillo*); Arcangeli, 1933, p. 32.

Synonyms: *Cubaris officinalis* Stebbing (not Duméril); Stebbing, 1900b, p. 655.

? *Armadillo rouxi* Verhoeff, 1926, p. 311.

New Caledonia.  
 (Type locality, Isle of Pines, New Caledonia.)

**Armadillo (Pseudosphaerillo) insularis** ARCANGELI: Arcangeli, 1933, p. 25.  
 (Type locality, Samoa [Tutuila].)

**Armadillo (Pseudosphaerillo) rouxi** VERHOEFF: Verhoeff, 1926, pp. 311 and 357.  
 Synonym: *Cubaris officinalis* Stebbing (not Duméril); Stebbing, 1900b, p. 655.  
 (Type locality, New Caledonia.)

**Genus CUBARIS Brandt, 1833<sup>5</sup> (= NESODILLO Verhoeff, 1926)**

(Type, *Cubaris cinerea* or *murina* Brandt)

**Cubaris bocki** (VERHOEFF): Verhoeff, 1926, p. 9 (*Nesodillo*).  
 (Type locality, Gilbert Islands [Aranuka].)

**Cubaris canalensis** (VERHOEFF): Verhoeff, 1926, p. 282 (*Nesodillo*).  
 (Type locality, New Caledonia.)

**Cubaris galbineus** (ESCHSCHOLTZ): Eschscholtz, 1823, p. 112 (*Armadillo*);  
 Budde-Lund, 1885, p. 39 (*Armadillo*); 1904, p. 120 (? *Sphaerillo*).  
 (Type locality, "Guahm" Island.)

**Cubaris incisus** (VERHOEFF): Verhoeff, 1926, p. 284 (*Nesodillo*).  
 (Type locality, New Caledonia.)

**Cubaris javanensis** (DOLLFUS): Dollfus, 1889, p. 91 (*Armadillo*); 1890b, p. 3  
 (*Armadillo*); Budde-Lund, 1904, p. 120 (*Armadillo*).  
 Hawaii (Oahu).  
 (Type locality, Java.)

**Cubaris lacustris** (VERHOEFF): Verhoeff, 1926, p. 286 (*Nesodillo*).  
 (Type locality, New Caledonia.)

**Cubaris lifuensis** STEBBING: Stebbing, 1900b, p. 653.  
 (Type locality, Lifu, Loyalty Islands.)

**Cubaris longicornis** (VERHOEFF): Verhoeff, 1926, p. 283 (*Nesodillo*).  
 (Type locality, New Caledonia.)

**Cubaris lundi** STEBBING: Stebbing, 1900b, p. 655. (Perhaps a *Sphaerillo*.)  
 (Type locality, New Britain.)

**Cubaris milleri** CHILTON: Chilton, 1917, p. 327. (Probably a new genus related  
 to *Sphaerillo*.)  
 (Type locality, Wellington, New Zealand.)

**Cubaris murina** BRANDT: Brandt, 1833, p. 190; Budde-Lund, 1885, p. 27 (*Arma-*  
*dillo*); 1904, p. 119 (*Armadillo*); Jackson, 1933a, p. 90; 1933b, p. 157;  
 1938, p. 192; Verhoeff, 1926, p. 8.  
 Synonym: *Nesodillo medius* Verhoeff, 1926,<sup>6</sup> p. 287.

<sup>5</sup> All species of *Cubaris* Brandt listed below belong to the genus as restricted by Verhoeff (= *Nesodillo* Verhoeff) except *C. lundi* Stebbing, *C. milleri* Chilton, and *C. suteri* Chilton, which are related to *Sphaerillo* and will ultimately find a home elsewhere.

<sup>6</sup> Verhoeff (1938) considers that *C. murina* and *C. medius* are different, but nearly related, species. An examination of numerous specimens of the former in

For further references and synonymy see Van Name, 1936, p. 387.

Hawaii (Oahu); Marquesas (Nukuhiva, Hivaoa, Fatuhiva, Uahuka, Uapou, Mohotani, Eiao); Tuamotus (Mangareva); Australs (Rapa, Raivavae); Society Islands (Tahiti, Moorea); Fanning; New Caledonia.  
(Type locality, Brazil.)

**Cubaris pacificus** (VERHOEFF): Verhoeff, 1926, p. 284 (*Nesodillo*).  
(Type locality, New Caledonia.)

**Cubaris plasticus** (VERHOEFF): Verhoeff, 1926, p. 290 (*Nesodillo*).  
(Type locality, New Caledonia.)

**Cubaris pronyensis** (VERHOEFF): Verhoeff, 1926, p. 289 (*Nesodillo*). *C. pronyensis* var. *erythrocephalus* Verhoeff, 1926, p. 289 (*Nesodillo*).  
(Type locality, New Caledonia.)

**Cubaris sarasini** (VERHOEFF): Verhoeff, 1926, p. 280 (*Nesodillo*). *C. sarasini* var. *marmoratus* Verhoeff, 1926, p. 280 (*Nesodillo*).

Synonym: ? *Cubaris lifuensis* Stebbing, 1900b, p. 653; Verhoeff, 1926, p. 357 ("lifouensis").  
(Type locality, New Caledonia.)

**Cubaris suteri** CHILTON: Chilton, 1915a, p. 425. (Probably a *Sphaerillo*.)  
(Type locality, Auckland, New Zealand)

#### Genus ECHINODILLO Jackson, 1933

(Type, *Echinodillo montanum* Jackson)

**Echinodillo montanum** JACKSON: Jackson, 1933b, p. 159.  
(Type locality, Marquesas [Uahuka].)

#### Genus EMYDODILLO Verhoeff, 1926

(Type, *Emydodillo testudo* Verhoeff)

**Emydodillo testudo** VERHOEFF: Verhoeff, 1926, p. 264.  
(Type locality, New Caledonia.)

#### Genus HAWAIODILLO Verhoeff, 1926

(Type, ? *Armadillo perkinsi* Dollfus)

**Hawaiodillo frontalis** (BUDDE-LUND): Budde-Lund, 1904, p. 67 (*Spherillo*); Verhoeff, 1926, p. 256 (*Hawaiodillo*).  
Synonym: *Armadillo Danac* Dollfus, 1900, p. 522.  
(Type locality, Hawaii [Kauai].)

**Hawaiodillo perkinsi** (DOLLFUS): Dollfus, 1900, p. 522 (*Armadillo*); Budde-Lund, 1904, p. 67 (*Spherillo*); Verhoeff, 1926, p. 256 (*Hawaiodillo*).  
(Type locality, Hawaii [Maui].)

---

the British Museum (Natural History) from widely spread localities suggests that the points of difference cited are not sufficiently constant to justify their separation.

**Hawaiiodillo sharpi** (DOLLFUS): Dollfus, 1900, p. 523 (*Armadillo*) ; Budde-Lund, 1904, p. 67 (*Sphaerillo*) ; Verhoeff, 1926, p. 256 (*Hawaiiodillo*).  
(Type locality, Hawaii [Kauai].)

**Genus MELANESILLO Verhoeff, 1938**

(Type, *Sphaerillo hebridarum* Verhoeff)

**Melanesillo hebridarum** (VERHOEFF): Verhoeff, 1926, p. 297 (*Sphaerillo*) ; Herold, 1931, p. 321 (*Lobodillo*) ; Verhoeff, 1938, p. 1 (*Melanesillo*).  
(Type localities, New Hebrides; Banks Islands.)

**Melanesillo bocki** VERHOEFF: Verhoeff, 1938, p. 4.  
(Type locality, Marshall Islands [Jaluit].)

**Melanesillo scamnorum** VERHOEFF: Verhoeff, 1938, p. 4.  
(Type locality, Fiji [Viti Levu].)

**Genus MERULANA Budde-Lund, 1913**

(Type, *Armadillo rugosus* Budde-Lund)

**Merulana canaliculatus** (BUDDE-LUND): Budde-Lund, 1904, p. 74 (*Sphaerillo*) ; Chilton, 1910b, p. 290 (*Cubaris*) ; Budde-Lund, 1913, p. 65 (*Merulana*) ; Chilton, 1925a, p. 319.

Synonym: *Armadillo speciosus* Chilton (not Dana), 1906b, p. 273.  
(Type locality, Chatham Islands.)

**Merulana chathamensis** (BUDDE-LUND): Budde-Lund, 1904, p. 75 (*Sphaerillo*) ; Chilton, 1910b, p. 290 (*Cubaris*) ; Budde-Lund, 1913, p. 65.

Synonyms: *Armadillo speciosus* Chilton, 1901, p. 146 (not Dana) ; Budde-Lund, 1904, p. 75.  
(Type locality, Chatham Islands.)

**Merulana exilis** (BUDDE-LUND): Budde-Lund, 1885, p. 288 (*Armadillo*) ; 1904, p. 76 (*Sphaerillo*).  
(Type locality, New Caledonia [Noumea].)

**Merulana noduligera** VERHOEFF: Verhoeff, 1926, p. 306.  
(Type localities, New Caledonia, Loyalty Islands.)

**Merulana translucida** (BUDDE-LUND): Budde-Lund, 1885, p. 290 (*Armadillo*) ; 1904, p. 76 (*Sphaerillo*) ; Verhoeff, 1926, p. 304 (*Merulana*). *M. translucida* var. *translucida* Verhoeff, 1926, p. 306. *M. translucida* var. *gracilior* Verhoeff, 1926, p. 306.

Synonym: ? *Pyrgoniscus cinctutus* Kinahan, 1859, p. 199.  
(Type localities, New Caledonia, Loyalty Islands.)

**Genus MERULANELLA Verhoeff, 1926**

(Type, *Merulanella carinata* Verhoeff)

**Merulanella carinata** VERHOEFF: Verhoeff, 1926, p. 308.  
(Type locality, New Caledonia.)

**Merulanella dollfusi** (STEBBING) : Stebbing, 1900b, p. 654 (*Cubaris*) ; Verhoeff, 1926, p. 357.

(Type locality, Loyalty Islands.)

**Merulanella wahrbergi** VERHOEFF : Verhoeff, 1926, p. 309.

(Type locality, New Caledonia.)

**Genus MESODILLO** Verhoeff, 1926

(Type, *Mesodillo eremitus* Verhoeff)

**Mesodillo eremitus** VERHOEFF : Verhoeff, 1926, p. 294.

(Type locality, New Caledonia.)

**Genus OCHEtodillo** Verhoeff, 1926

(Type, *Ochetodillo sulcatus* Verhoeff)

**Ochetodillo sulcatus** VERHOEFF : Verhoeff, 1926, p. 271.

(Type locality, Humboldt Mountains, New Caledonia.)

**Genus ORODILLO** Verhoeff, 1926

(Type, *Orodillo collaris* Verhoeff)

**Orodillo collaris** VERHOEFF : Verhoeff, 1926, p. 291.

(Type locality, New Caledonia.)

**Genus SCHISMADILLO** Verhoeff, 1926

(Type, *Schismadillo rouxi* Verhoeff)

**Schismadillo rouxi** VERHOEFF : Verhoeff, 1926, p. 272.

(Type locality, Canala Mountains, New Caledonia.)

**Genus SPHERILLO** Dana, 1853<sup>7</sup>

(Type, ? *Spherillo vitiensis* Dana)

*Spherillo* Budde-Lund, 1904

*Sphaerillo* Verhoeff (ex parte), 1926

(Type, *Spherillo danae* Heller)

**Spherillo albospinosus** (DOLLFUS) : Dollfus, 1900, p. 521 (*Armadillo*) ; Budde-Lund, 1904, p. 54 (*Spherillo*).

Hawaii (Oahu, Kauai).

(Type locality, Hawaii [Oahu].)

<sup>7</sup> All species which fall under *Sphaerillo* Verhoeff are distinguished by that spelling from those which were included under *Spherillo* Dana by Budde-Lund and will ultimately find a place in other new or existing genera. (See introduction.)

**Spherillo ambitiosus** (BUDDE-LUND): Budde-Lund, 1879, p. 7 (*Armadillo*); 1885, p. 34 (*Armadillo*); Chilton, 1901, p. 144 (*Armadillo*); Budde-Lund, 1904, p. 63 (*Spherillo*); Chilton, 1910b, p. 289 (*Cubaris*).

(Type locality, New Zealand.)

"Common over North Island and in South Island down west coast as far as Daggs Sound." (Chilton.)

**Spherillo bipunctatus** BUDDE-LUND: Budde-Lund, 1904, p. 62; Chilton, 1910b, p. 289 (*Cubaris*).

(Type locality, Lyttelton, New Zealand.)

"One male specimen." (Budde-Lund.)

**Sphaerillo brevis** (BUDDE-LUND): Budde-Lund, 1904, p. 93 (*Spherillo*); Chilton, 1910b, p. 290 (*Cubaris*).

(Type locality, Auckland.)

"One specimen." (Budde-Lund.)

**Spherillo carinulatus** BUDDE-LUND: Budde-Lund, 1904, p. 54.

Synonym: *Armadillo albospinosus* Dollfus, 1900, p. 521, ♀.

(Type locality, Hawaii [Kauai].)

**Sphaerillo danae** (HELLER): Heller, 1865, p. 134 (*Spherillo*); Budde-Lund, 1885, p. 39 (*Armadillo*); Filhol, 1885, p. 440 (*Spherillo*); Thomson and Chilton, 1886, p. 159 (*Spherillo*); Chilton, 1901, p. 145 (*Armadillo*); Budde-Lund, 1904, p. 94 (*Spherillo*); Chilton, 1910b, p. 290 (*Cubaris*).

Synonym: *Armadillo inconspicuus* Miers, 1876a, p. 225; 1876b; p. 95; Filhol, 1885, p. 439; Thomson and Chilton, 1886, p. 159.

New Zealand, North Island, more rare in South; Kapiti Island, Auckland.

(Type locality, Auckland.)

**Spherillo dispersus** BUDDE-LUND: Budde-Lund, 1904, p. 70.

(Type locality, Bismarck Archipelago?)

**Spherillo erinaceus** (BUDDE-LUND): Budde-Lund, 1879, p. 7; 1885, p. 36 (*Arma-*

*dillo*); 1904, p. 54 (*Spherillo*).

(Type locality, Samoa [Upolu].)

**Sphaerillo (Sphaerillo) fissus** VERHOEFF: Verhoeff, 1926, p. 299.

(Type locality, New Caledonia.)

**Spherillo hamiltoni** (CHILTON): Chilton, 1901, p. 148 (*Armadillo*); Budde-

Lund, 1904, p. 54 (*Spherillo*); Chilton, 1910b, p. 289 (*Cubaris*).

(Type locality, New Zealand.)

**Spherillo hawaiiensis** DANA: Dana, 1853, p. 722; Budde-Lund, 1885, p. 39 (*Armadillo*); Dollfus, 1900, p. 521 (*Armadillo*); Budde-Lund, 1904, p. 57 (*Spherillo*).

Synonyms: *Armadillo bidens* Budde-Lund, 1885, p. 18 (♂).

*Armadillo australis* Budde-Lund, 1885, p. 33 (♀).

Hawaii (Oahu, Lanai, Kauai, Molokai).

(Type locality, "Sandwich Islands.")

**Sphaerillo (Xestodillo) lifouensis** VERHOEFF: Verhoeff, 1926, p. 301; *S. lifouen-*

*sis* var. *ambrymensis*, Verhoeff, 1926, p. 302 (New Hebrides.)

(Type locality, Loyalty Islands.)

**Spherillo macmahoni** (CHILTON): Chilton, 1901, p. 149 (*Armadillo*) ; Budde-Lund, 1904, p. 56 (*Spherillo*) ; Chilton, 1910b, p. 289 (*Cubaris*). (Probably a *Sphaerillo*.)

(Type locality, Marlborough, New Zealand.)

**Spherillo marginatus** BUDDE-LUND: Budde-Lund, 1904, p. 65; Chilton, 1910b, p. 289 (*Cubaris*).

(Type locality, Auckland, New Zealand.)

"One female specimen."

**Spherillo melanurus** (DOLLFUS): Dollfus, 1887, p. 1 (*Armadillo*) ; Budde-Lund, 1904, p. 94 (*Spherillo*).

(Type locality, New Caledonia.)

**Spherillo monolinus** DANA: Dana, 1853, p. 719; Heller, 1865, p. 135; Budde-Lund, 1885, p. 39 (*Armadillo*) ; Filhol, 1885, p. 440; Thomson and Chilton, 1886, p. 159 (*Spherillo*) ; Chilton, 1901, p. 148 (*Armadillo*) ; Budde-Lund, 1904, p. 68 (*Spherillo*) ; Chilton, 1910b, p. 290 (*Cubaris*).

Synonym: *Armadillo Aucklandicus* Budde-Lund, 1885, p. 40; 1904, p. 69 (*Spherillo*).

New Zealand, "Whykare River near Bay of Islands."

(Type locality, Auckland, New Zealand.)

**Sphaerillo (Sphaerillo) montivagus** (BUDDE-LUND): Budde-Lund, 1879, p. 7 (*Armadillo*) ; 1885, p. 35 (*Armadillo*) ; 1904, p. 89 (*Spherillo*) ; Jackson, 1933a, p. 90; 1933b, p. 155; 1938, p. 190.

Samoa (Upolu); Tuamotus (Mangareva, Henderson); Australs (Rapa, Raivavae, Tubuai); Pitcairn; Society Islands (Tahiti); Tonga; Marquesas (Hivaaoa, Uahuka, Uapou); Bismarck Archipelago (Ruk).

(Type locality, Samoa [Upolu].)

**Sphaerillo (Xestodillo) marquesarum** (JACKSON): Jackson, 1933a, p. 90 (*Spherillo*) ; 1933b, p. 156; 1938, p. 191: var. *australis* Jackson, 1938, p. 191.

Marquesas (Uapou); Society Islands (Tahiti, Moorea); Tuamotus (Mangareva, Henderson); Pitcairn; Australs (Raivavae, Rurutu, Rimatara).

(Type locality, Marquesas [Uapou].)

**Sphaerillo pictus** (HELLER): Heller, 1865, p. 135 (*Spherillo*) ; Budde-Lund, 1885, p. 40 (*Armadillo*) ; 1904, p. 95 (*Spherillo*).

(Type locality, "Taiti" Island.)

**Sphaerillo (Xestodillo) politus** VERHOEFF: Verhoeff, 1926, p. 302.

(Type locality, New Caledonia.)

**Sphaerillo (Sphaerillo) pygmaeus** VERHOEFF: Verhoeff, 1926, p. 296; Jackson, 1933b, p. 156.

New Caledonia; Marquesas (Nukuhiva).

(Type locality, New Caledonia.)

**Spherillo ruficornis** BUDDE-LUND: Budde-Lund, 1885, p. 283 (*Armadillo*) ; 1904, p. 86 (*Spherillo*).

(Type locality, New Caledonia.)

**Spherillo rufomarginatus** BUDDE-LUND: Budde-Lund, 1904, p. 64; Chilton, 1910b, p. 289 (*Cubaris*).

(Type locality, New Zealand.)

"One female specimen was taken at Taranga [? Tauranga]."

**Spherillo rugulosus** (MIERS): Miers, 1876a, p. 225 (*Cubaris*); 1876b, p. 96 (*Cubaris*); Chilton, 1883a, p. 73 (*Cubaris*); Budde-Lund, 1885, p. 40 (*Armadillo*); Thomson and Chilton, 1886, p. 158 (*Cubaris*); Chilton, 1901, p. 147 (*Armadillo*); Budde-Lund, 1904, p. 65 (*Spherillo*); Chilton, 1909a, p. 668 (*Cubaris*); 1910b, p. 290 (*Cubaris*); Stephensen, 1927, p. 372 (*Cubaris*).

New Zealand, "very abundant on South Island" (Chilton); Auckland and Campbell Islands.

(Type locality, New Zealand.)

**Sphaerillo setaceus** (BUDDE-LUND): Budde-Lund, 1904, p. 89 (*Spherillo*); Chilton, 1910b, p. 290 (*Cubaris*).

(Type locality, Auckland.)

One specimen.

**Sphaerillo societatis** (MACCAGNO): Maccagno, 1932, p. 1 (*Spherillo*).

(Type locality, Society Islands [Moorea].)

**Spherillo speciosus** (DANA): Dana, 1853, p. 718 (*Armadillo*); Miers, 1876b, p. 95 (*Armadillo*); Budde-Lund, 1885, p. 39 (*Armadillo*); Thomson and Chilton, 1886, p. 159 (*Armadillo*); not Chilton, 1901, p. 146; Budde-Lund, 1904, p. 69 (*Spherillo*); Chilton, 1910b, p. 290 (*Cubaris*). See *Merulana chathamensis*.

(Type locality, New Zealand, near Bay of Islands.)

**Spherillo spicatus** JACKSON: Jackson, 1927b, p. 4.

(Type locality, Samoa [Upolu].)

**Spherillo spinosus** DANA: Dana, 1853, p. 723; Budde-Lund, 1885, p. 39 (*Armadillo*); Thomson and Chilton, 1886, p. 159 (*Spherillo*); Chilton, 1901, p. 150 (*Armadillo*); Budde-Lund, 1904, p. 54 (*Spherillo*); Chilton, 1910b, p. 289 (*Cubaris*); Verhoeff, 1926, p. 274 (? *Schismadillo*).

(Type locality, New Zealand, near Bay of Islands.)

**Spherillo squamatus** BUDDE-LUND: Budde-Lund, 1904, p. 61; Chilton, 1910b, p. 289 (*Cubaris*).

(Type locality, New Zealand.)

"One female from Lyttelton, near Christchurch."

**Spherillo tarangensis** BUDDE-LUND: Budde-Lund, 1904, p. 67; Chilton, 1910b, p. 290 (*Cubaris*).

New Zealand: Taranga [? Tauranga] and Lyttelton.

(Type locality, Taranga and Lyttelton, New Zealand.)

**Spherillo testudinalis** (BUDDE-LUND): Budde-Lund, 1879, p. 7 (*Armadillo*); 1885, p. 29 (*Armadillo*); 1904, p. 80 (*Spherillo*); 1908, p. 269; 1912, p. 377 (*Cubaris*); Searle, 1914, p. 367; Jackson, 1927b, p. 3; 1933b, p. 156; 1938, p. 190.

Synonyms: *Armadillo samoensis* Budde-Lund, 1885, p. 30; 1904, p. 80.  
*Armadillo tongensis* Budde-Lund, 1885, p. 284; 1904, p. 80.  
*? Armadillidium pacificum* Borradaile, 1900, p. 796; Searle, 1914,  
p. 369.

Samoa (Upolu); Tonga; Bismarck Archipelago (Ruk, Matapi); Cook Islands (Rarotonga); Caroline Islands; Marianas; Ellice Islands; Fiji; Marquesas (Nukuhiva); Tuamotus (Mangareva); Australs (Raivavae, Tubuai); Society Islands (Tahiti, Raiatea, Moorea, Tahaa, Mehetia [Maitea]).

(Type locality, Nicobar Island, Bay of Bengal.)

**Spherillo vitiensis** DANA: Dana, 1853, p. 721; Budde-Lund, 1885, p. 39 (*Armadillo*); not Dollfus, 1890b, p. 2 (*Armadillo*); Budde-Lund, 1904, p. 69 (*Spherillo*).

(Type locality, Fiji [Vanua Levu].)

**Sphaerillo (Xestodillo) zebicolor** (STEBBING): Stebbing, 1900b, p. 656 (*Cutbaris*); Verhoeff, 1926, p. 300 (*Sphaerillo*).

(Type localities, New Caledonia, Loyalty Islands.)

#### Genus TRIDENTODILLO Jackson, 1933

(Type, *Tridentodillo squamosus* Jackson)

**Tridentodillo squamosus** JACKSON: Jackson, 1933b, p. 161.

(Type locality, Marquesas [Nukuhiva].)

#### Subfamily ARMADILLIDINAE

##### Genus ARMADILLIDIUM Brandt, 1830

(Type, *Armadillo vulgare* Latreille.)

**Armadillidium vulgare** (Latreille): Latreille, 1804, p. 48 (*Armadillo*); Chilton, 1901, p. 142; 1905, p. 431.

For further references and synonymy see Budde-Lund, 1885, p. 67, and Van Name, 1936, p. 276.

New Zealand.

(Type locality, Europe.)

## Suborder PHREATOICIDEA

### Family PHREATOICIDAE

#### Genus PHREATOICUS Chilton, 1882

(Type, *Phreatoicus typicus* Chilton, 1882)

**Phreatoicus typicus** CHILTON: Chilton, 1883b, p. 87; Thomson and Chilton, 1886, p. 151; Stebbing, 1888, pp. 543 and 587; 1893, p. 388; Chilton, 1894, p. 196; Sheppard, 1927, p. 109.

New Zealand.

(Type locality, Eyrerton and Ashburton, New Zealand [in wells].)

**Phreatoicus kirkii** CHILTON: Chilton, 1906c, p. 275; Sheppard, 1927, p. 110.  
 (Type locality, "In a fresh water lagoon on Ruapuke Island.")  
 Var. *dunedensis* Chilton, 1906c, p. 276.  
 (Type locality, in streams, Mosgiel and Woodhaugh, near Dunedin, New Zealand.)

**Phreatoicus assimilis** CHILTON: Chilton, 1894, p. 186; Sheppard, 1927, p. 111.  
 (Type locality, in wells at Winchester, near Canterbury, New Zealand.)

## Suborder VALVIFERA

### Family IDOTHEIDAE

#### Subfamily MESIDOTEINAE

##### Genus AUSTRIDOTEA Nicholls, 1937

(Type, *Austridotea annectens* Nicholls, 1937)

**Austridotea (Austridotea) annectens** NICHOLLS: Nicholls, 1937, p. 118.  
 (Type locality, Stewart Island.)

**Austridotea (Austridotea) benhami** NICHOLLS: Nicholls, 1937, p. 123.  
 Synonyms: *Idotea lacustris* var.  $\beta$ : Chilton, 1891, p. 131; 1892, p. 263; 1909a, p. 658.  
*Pentidotea lacustris* (part): Collinge, 1916, p. 154.  
 Otago Peninsula, New Zealand, Campbell Island ?.  
 (Type locality, "mountain streams around Port Chalmers and Dunedin," Otago Peninsula, New Zealand.)

**Austridotea (Notidotea) lacustris** (THOMSON)<sup>8</sup>; Thomson, 1879c, p. 251 (*Idotea*); Miers, 1881, p. 39 (*Idotea*); Thomson and Chilton, 1886, p. 156 (*Idotea*); Chilton, 1890, p. 194 (*Idotea*); *I. lacustris* var.  $\alpha$  Chilton, 1892, p. 263; Chilton, 1909a, p. 658; Collinge, 1916, p. 154 (*Pentidotea*); Van Name, 1936, p. 451 (*Pentidotea*); Nicholls, 1937, p. 125 (*Notidotea*).  
 Dunedin, Waitati, Mihiwaka, New Zealand.  
 (Type locality, Dunedin, New Zealand.)

## Suborder FLABELLIFERA

### Family ANTHRIDAE

##### Genus CRUREGENS Chilton, 1882

(Type, *Cruregens fontanus* Chilton, 1882)

**Cruregens fontanus** CHILTON: Chilton, 1882, p. 175; 1883b, p. 88; Thomson and Chilton, 1886, p. 152; Chilton, 1894, p. 211.  
 Eyrerton, North Canterbury; Leeston; Winchester, South Canterbury (in wells).  
 (Type locality, Eyrerton, North Canterbury, New Zealand.)

---

<sup>8</sup>"*I. lacustris* G. M. T. is in all probability not restricted to, but merely tolerant of, fresh water." Nicholls, 1937.

## Family SPIHAEROMIDAE

## Subfamily SPHAEROMINAE

## Genus PARAVIREIA Chilton, 1925

(Type, *Paravireia typicus* Chilton)

Paravireia typicus CHILTON: Chilton, 1925b, p. 321.

(Type locality, Chatham Islands.)

## REFERENCES

ARCANGELI, A.

1927. Isopodi terrestri raccolti nell'Estremo Oriente. Boll. Lab. Zool., Portici, vol. 20, pp. 211-269, figs. 1-22.  
 1933. Isopodi terrestre delle Isole Samoa. . . . Boll. Mus. Zool. Anat. Comp. Torino, vol. 43, pp. 25-33, 4 pls.

BARNARD, K. H.

1932. Contributions to the crustacean fauna of South Africa. No. 11. Terrestrial Isopoda. Ann. South African Mus., vol. 30, pt. 2, pp. 179-388, figs. 1-80.

BATE, C. SPENCE, and WESTWOOD, J. O.

1868. A history of the British sessile-eyed Crustacea, vol. 2, lvi + 536 pp. London.

BORRADAILE, L. A.

1900. On some crustaceans from the South Pacific. Pt. 5, Arthrostracans and barnacles. Proc. Zool. Soc. London, pp. 795-799, pl. 51.

BRANDT, J. F.

1833. Conspectus Monographiae Crustaceorum Oniscodorum Latreillii. Bull. Soc. Imp. Nat. Moscou, vol. 6, pp. 171-193, pl. 4. (Also published with separate paging.)

BUdde-LUND, G.

1879. Prospectus generum specierumque Crustaceorum Isopodum terrestrium, pp. 1-10. Copenhagen.

1885. Crustacea Isopoda Terrestria per Familias et Genera et Species descripta, pp. 1-319. Håvniae.

1904. A revision of "Crustacea Isopoda Terrestria" with additions and illustrations. 2. Spherilloninae. 3. Armadillo. Pp. 33-144, pls. 6-10. Copenhagen.

1906. Die Landisopoden der Deutschen Südpolar-Expedition 1901-1903. Mit Diagnosen verwandter Arten. Deutsche Südpolar-Expedition 1901-1903, bd. 9, heft 2, pp. 69-92, pls. 3-4.

1908. Isopoda von Madagaskar und Ostafrika. In Voeltzkow, Reise in Ostafrika in den Jahren 1903-1905, bd. 2, pp. 265-308, pls. 12-18.

1912. Terrestrial Isopoda, particularly considered in relation to the distribution of the southern Indo-Pacific species. In Report Percy Sladen Trust Exped. Trans. Linn. Soc. London (Zool.), vol. 15, pp. 367-394, pls. 20-22. (Revised and footnotes added by Rev. T. R. R. Stebbing.)

1913. Über einige Oniscoideen von Australien, nachgelassenes Fragment. Jahrb. Hamburg Wiss. Anst., vol. 30, beiheft 2, pp. 65-72, figs. 1-8, 1 pl.

## CHILTON, CHARLES

1882. On some subterranean Crustacea. *Trans. New Zealand Inst.*, vol. 14 (1881), pp. 174-180, pls. 10, 11.
- 1883a. Further additions to our knowledge of the New Zealand Crustacea. *Trans. New Zealand Inst.*, vol. 15 (1882), pp. 69-86, pls. 1-3.
- 1883b. Notes on, and a new species of subterranean Crustacea. *Trans. New Zealand Inst.*, vol. 15 (1882), pp. 87-92, pl. 4.
- 1883c. Additions to the isopoden fauna of New Zealand. *Trans. New Zealand Inst.*, vol. 15 (1882), pp. 145-150, 1 pl.
1886. A new species of *Philygria*. *Trans. New Zealand Inst.*, vol. 18 (1885), pp. 159-161, pl. 5.
1890. Revision of the New Zealand Idoteidae. *Trans. New Zealand Inst.*, vol. 22 (1889), pp. 189-204.
1891. General notes. *New Zealand Journ. Sci. (new issue)*, vol. 1, No. 3, pp. 131-132.
1892. Notes on some New Zealand Amphipoda and Isopoda. *Trans. New Zealand Inst.*, vol. 24 (1891), pp. 258-269.
1894. The subterranean Crustacea of New Zealand: with some general remarks on the fauna of caves and wells. *Trans. Linn. Soc. London (Zool.)*, vol. 6, pt. 2, pp. 163-284, pls. 16-23.
1901. The terrestrial Isopoda of New Zealand. *Trans. Linn. Soc. London (Zool.)*, vol. 8, pt. 4, pp. 99-152, pls. 11-16.
1905. Notes on the distribution of some species of terrestrial Isopoda introduced into Australasia. *Ann. Mag. Nat. Hist.*, vol. 16, ser. 7, pp. 428-432.
- 1906a. Note on the occurrence of *Metoponorthus pruinosis*, Brandt, in New Zealand. *Trans. New Zealand Inst.*, vol. 38 (1905), pp. 64-65.
- 1906b. List of Crustacea from the Chatham Islands. *Trans. New Zealand Inst.*, vol. 38 (1905), pp. 269-273.
- 1906c. Description of a species of *Phreatoicus* from the surface waters of New Zealand. *Trans. New Zealand Inst.*, vol. 38 (1905), pp. 274-276.
- 1909a. The Crustacea of the Subantarctic Islands of New Zealand. The Subantarctic Islands of New Zealand, vol. 2, art. 26, pp. 601-671, figs. 1-19. Wellington, New Zealand.
- 1909b. The biological relations of the Subantarctic Islands of New Zealand: summary of results. The Subantarctic Islands of New Zealand, vol. 2, art. 35, pp. 793-807. Wellington, New Zealand.
- 1910a. On an isopod inhabiting ants' nests in New Zealand. *Trans. New Zealand Inst.*, vol. 42 (1909), pp. 190-192.
- 1910b. Additions to the terrestrial Isopoda of New Zealand. *Trans. New Zealand Inst.*, vol. 42 (1909), pp. 286-291.
1911. The Crustacea of the Kermadec Islands. *Trans. New Zealand Inst.*, vol. 43 (1910), pp. 544-573, figs. 1-4.
- 1915a. Some terrestrial Isopoda from New Zealand and Tasmania, with description of a new genus. *Journ. Linn. Soc. London (Zool.)*, vol. 32, pp. 417-427, pls. 36-37.
- 1915b. *Deto*, a subantarctic genus of terrestrial Isopoda. *Journ. Linn. Soc. London (Zool.)*, vol. 32, pp. 435-456, pls. 39, 40.

1917. A new tuberculate terrestrial isopod from New Zealand. Ann. Mag. Nat. Hist., vol. 19, ser. 8, pp. 327-329, pl. 13.
1922. Note on the isopod known as *Geoligia perkinsi* Dollfus (Crust.). Proc. Hawaiian Ent. Soc., vol. 1, No. 1, pp. 83-86.
- 1924a. Fauna of the Chilka Lake. Tanaidacea and Isopoda. Mem. Indian Mus., vol. 5, No. 12, pp. 875-895, figs. 1-10, pl. 60.
- 1924b. Occurrence in South America of the shore isopod *Ligia novae-zealandiae* Dana. New Zealand Journ. Sci. and Techn., vol. 6, Nos. 5 and 6, pp. 287-288.
- 1925a. Some Amphipoda and Isopoda from the Chatham Islands. Rec. Canterbury Mus., vol. 2, pp. 317-320.
- 1925b. A new genus of Isopoda (Family Sphaeromidae). Rec. Canterbury Mus., vol. 2, pp. 321-326, pls. 45-47.

## COLLINGE, W. E.

1916. Some remarks upon the structure and generic position of *Idotea lacustris* Thomson. Journ. Zool. Res., London, vol. 1, pp. 153-157, pl. 6.

## CUVIER, G.

1829. Le règne animal, édit. 2, tom. 4, par P. A. Latreille. Paris.
1849. Le règne animal, édition 3 accompagnée de planches gravées. Vol. 8, Les Crustacés par Milne-Edwards. Paris.

## DANA, J. D.

1853. Isopoda. In United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842, under the command of Charles Wilkes, U.S.N., vol. 14, Crustacea, pt. 2, pp. 696-805. Atlas, pls. 46-53 (1855).
1856. Catalogue and descriptions of Crustacea collected in California by Dr. John Le Conte. Proc. Philadelphia Acad. Nat. Sci., vol. 7, pp. 175-177.

## DESMAREST, A. G.

1825. Considérations générales sur la classe des crustacés. Pp. xix + 446, pls. 1-56. Paris.

## DOLLFUS, ADRIEN

1887. Diagnoses d'espèces nouvelles . . . des Armadilliens. Bull. Soc. Études Sci., vol. 9, pp. 1-7.
1889. Sur quelques isopodes du Musée de Leyde. Notes Leyden Mus., vol. 11, pp. 91-94, pl. 5.
- 1890a. Note au sujet des isopodes terrestres du *Challenger*. Bull. Soc. Études Sci. Paris, vol. 12, p. 71.
- 1890b. Isopodes terrestres du *Challenger*. Bull. Soc. Études Sci. Paris, vol. 12, pp. 63-70, 2 pls. (Also reprinted, paged 1-8.)
1893. Sur la distribution du genre *Ligia* Fabr. Feuille Natural., vol. 24, No. 3, pp. 23-26, figs. 1-4.
1898. Isopodes Terrestres des Indes Neerlandaises. M. Weber, Zool. Ergebni. einer Reise in Niederländisch-OstIndien, vol. 4, pp. 357-382, 3 pls. Leiden.
1900. Crustacea Isopoda. Fauna Hawaïensis, vol. 2, pt. 5, pp. 521-526, pl. 20 (vol. 2, 1899-1910 was issued in 1913).

## ESCHSCHOLTZ, F. FRID

1823. Animalia tetracera et myriapoda exotica. Mem. Soc. Imp. Nat. Moscou, vol. 6, pt. 2, pp. 111-114.

## FABRICIUS, J. C.

1793. Entomologia systematica emendata et aucta. Vol. 2, viii + 519 pp. Hafniae.  
1798. Supplementum entomologiae systematicae. Hafniae.

## FILHOL, H.

1885. Mission de l'île Campbell. Rec. Mémoires . . . , vol. 3, pt. 2, Crustacea, chap. 7, pp. 349-516, pls. 38-55. Paris (Inst. France, Acad. Sci.).

## GUÉRIN, F. E.

1836. *Doto*. Mag. Zool., vol. 6, classe 7, pl. 14 (rectius 21).

## HELLER, C.

- 1865.<sup>9</sup> Crustaceen in Reise der Oesterreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859. Zoologischer Theil., bd. 2, abt. 3, pp. 280, 25 pls. Vienna.

## HEROLD, W.

1931. Land-Isopoden von den Sunda-Inseln. Arch. Hydrobiol., suppl. bd. 9, heft 2, pp. 306-393, figs. 1-193.

## JACKSON, HAROLD GORDON

1922. A revision of the isopod genus *Ligia* (Fabricius). Proc. Zool. Soc. London, pp. 683-703, pls. 1, 2.

- 1927a. A new subgenus of *Ligia*, with further observations on the genus. Ann. Mag. Nat. Hist., vol. 19, ser. 9, pp. 129-136, pl. 2.

- 1927b. Isopoda terrestria. In Insects of Samoa. Brit. Mus. (Nat. Hist.), pt. 8, fasc. 1, pp. 1-11, pls. 1, 2.

- 1933a. *Porcellio (Heminagara) tahitiensis*, new subgenus and species, and other Tahitian terrestrial isopods. Bull. Bishop Mus., Honolulu, No. 113, pp. 87-90, fig. 1. (Pacific Ent. Surv. Publ. 6, art. 17.)

- 1933b. Marquesan terrestrial Isopoda. Bull. Bishop Mus., Honolulu, No. 114, pp. 145-162, figs. 1-6. (Pacific Ent. Surv. Publ. 7, art. 10.)

1936. Terrestrial isopods from Malaysia. Bull. Raffles Mus., Singapore, No. 12, pp. 77-87, figs. 1-4.

1938. Terrestrial isopods of southeastern Polynesia. Occ. Papers B. P. Bishop Mus., vol. 14, No. 10, pp. 167-192, figs. 1-10. (Mangarevan Exped. Publ. 26.)

## KINAHAN, J. R.

1859. On the genus *Platyarthrus* Brandt; with notices of allied undescribed genera. Proc. Dublin Univ. Zool.-Bot. Assoc., vol. 1, pt. 2, pp. 188-201.

## KOCH, C. L.

1847. System der Myriapoden mit den Verzeichnissen und Berichtigungen zu Deutschlands Crustacean, Myriapoden, und Arachniden. Regensburg.

<sup>9</sup> "The year 1865 is appended to the several plates, but the publication of the volume took place in 1866." Spence Bate, *Challenger Macrura*, 1888, p. 196.

## LATREILLE, P. A.

1804. Histoire naturelle des crustacés et des insectes. Vol. 7, 413 pp., pls. 58-66.  
 1806. Genera crustáceorum et insectorum . . . Vol. I. Parisiis et Argentorati.

## MACCAGNO, T.

1931. *Ligia porteri* Macc., nuova specie di isopodo terrestre del Cile. Boll. Zool. Napoli, vol. 2, pp. 151-157, pl. 3.  
 1932. Isopodi Terrestri delle Isole della Società. Boll. Mus. Zool. Anat. Comp. Torino, vol. 42, No. 20, pp. 1-7, 20 figs.

## MIERS, EDWARD J.

- 1876a. Description of some new species of Crustacea, chiefly from New Zealand. Ann. Mag. Nat. Hist., vol. 17, ser. 4, pp. 218-229.  
 1876b. Oniscoidea. In Catalogue of the stalk- and sessile-eyed Crustacea of New Zealand, pp. 94-103.  
 1877a. Crustacea. In Gunther, Account of the zoological collection made during the visit of H.M.S. *Peterel* to the Galapagos Islands. Proc. Zool. Soc. London, pp. 73-75, 1 pl.  
 1877b. On a collection of Crustacea, Decapoda, and Isopoda chiefly from South America, with descriptions of new genera and species. Proc. Zool. Soc. London, pp. 653-679, pls. 66-69.  
 1881. Revision of the Idoteidae, a family of sessile-eyed Crustacea. Journ. Linn. Soc. (Zool.), vol. 16, No. 89, pp. 1-88, pls. 1-3.

## MILNE-EDWARDS, H.

1840. Isopoda. In Histoire Naturelle des Crustacés, vol. 3, pp. 115-283, pls. 31-33. Paris.

## NICHOLLS, G. E.

1937. On the freshwater Idoteidae of New Zealand. Ann. Mag. Nat. Hist., vol. 19, ser. 10, pp. 113-136, 18 figs.

## NICOLET, H.

1849. Isopodes. In Gay, Historia fisica y política de Chile, Zool., vol. 3, pp. 256-287; atlas, pls. 3, 4, Crust. (1854).

## PANNING, A.

1924. Isopoda. In Michaelsen, Beiträge . . . Land-v. Süsswasserfauna Deutsch-Südwestafrikas, vol. 2, lief. 3, pp. 169-201, figs. 1-11.

## PERTY, M.

- 1830-34. Delectus animalium articulatorum quae in itinere per Braziliam annis 1817-1823 collegerunt J. B. de Spix et C. F. Ph. de Martius, pp. 1-224, pls. 1-40. Monachii. (Isopods pp. 211, 212, pl. 40.)

## PFEFFER, G.

1889. Ubersicht der von Herrn Dr. Franz Stuhlmann in Ägypten, auf Sansibar . . . gesammelten . . . Krebse. Jahrb. Hamburg Wiss. Anst., vol. 6, pt. 2 (1888), pp. 1-36.

## RICHARDSON, HARRIET

1905. A monograph on the Isopods of North America. U. S. Nat. Mus. Bull. 54, liii + 727 pp., figs. 1-740.

## ROUX, POLYDORE

1828. Crustacés de la méditerranée et de son littoral. Pp. 1-176, pls. 1-45. Marseille.

## SAVIGNY, J. C.

1827. Description de l'Egypte. Histoire naturelle Zoologie. Planches par J. C. Savigny. Vol. 12, Crust., pp. 1-13. Paris, 1826. Texte. Explication sommaire des planches de Crustacés de l'Egypte et de la Syrie publiées par J. C. Savigny et par V. Audouin, vol. 22, p. 249. Paris, 1827.

## SEARLE, H. RICHARDSON

1914. Isopoda. *Albatross* Exped. 1899-1900 and 1904-1905. Bull. Mus. Comp. Zool. Harvard, vol. 58, No. 8, pp. 361-372, figs. 1-16.

## SHEPPARD, EDITH M.

1927. Revision of the family Phreatoicidae (Crustacea), with a description of two new species. Proc. Zool. Soc. London, pp. 81-124, figs. 1-7.

## STEBBING, T. R. R.

1888. Report on the Amphipoda. . . . Rep. Sci. Res. H.M.S. *Challenger*, Zool., vol. 29, pp. 543 and 587.

1893. A history of Crustacea. xvii + 466 pp. New York.

- 1900a. On some crustaceans from the Falkland Islands collected by Mr. Rupert Vallentin. Proc. Zool. Soc. London, pp. 517-568, pls. 36-39.

- 1900b. On Crustacea brought by Dr. Willey from the South Seas. Willey's Zool. Results, pt. 5, pp. 605-690, pls. 64-74.

## STEPHENSEN, K.

1927. Crustacea from the Auckland and Campbell Islands. Vidensk. Medd. Dansk. Naturh. Foren., vol. 83, pp. 289-390, figs. 1-33.

## STIMPSON, WILLIAM

1857. Crustacea and Echinodermata of the Pacific shores of North America. Journ. Boston Soc. Nat. Hist., vol. 6, pp. 444-532, pls. 18-23.

## THOMSON, G. M.

- 1879a. New Zealand Crustacea, with descriptions of new species. Trans. New Zealand Inst., vol. 11 (1878), pp. 230-247, pl. 10.

- 1879b. Description of new crustacean from the Auckland Islands. Trans. New Zealand Inst., vol. 11 (1878), pp. 249-250.

- 1879c. Description of a new species of isopodous crustacean (*Idotea*). Trans. New Zealand Inst., vol. 11 (1878), pp. 250, 251.

1892. Notes on Tasmanian Crustacea, with descriptions of new species. Proc. Roy. Soc. Tasmania, pp. 45-76, pls. 1-6 (reprint, pp. 1-32).

## THOMSON, G. M., and CHILTON, CHARLES.

1886. Critical list of the Crustacea Malacostraca of New Zealand. Trans. New Zealand Inst. Zool., vol. 18 (1885), pp. 141-159.

## VAN NAME, WILLARD G.

1936. The American land and fresh-water isopod Crustacea. Amer. Mus. Nat. Hist. Bull. 71, vii + 535 pp., figs. 1-323.

## VERHOEFF, K. W.

1926. Isopoda Terrestria von Neu-Caledonien und den Loyalty-Inseln. Nova Caledonia, Zool., vol. 4, liv. 2, pp. 243-366, figs. 1-141.

- 1928a. Über einige Isopoden der zoologischen Staatssammlung in München.

38. Isopoden-Aufsatz. Zool. Anz., vol. 76, heft 1/2, pp. 25-36, figs. 1-16; heft 3/6, pp. 113-123, figs. 17-31.

1928b. Isopoden aus Formosa. 39. Isopoden-Aufsatz. Mitt. Zool. Mus. Berlin, bd. 14, heft 2, pp. 199-226, figs. 1-31.

1938. Über einige polynesische Oniscoideen von Prof. Sixten Bock's Pazifik-Expedition 1917-1918. Ark. Zool., vol. 30A, No. 16, pp. 1-14, figs. 1-21.

WAHRBERG, RAGNAR

1922a. Results of Dr. E. Mjöberg's Swedish scientific expeditions to Australia, 1910-1913, No. 30. Terrestre Isopoden aus Australien. Ark. Zool., vol. 15, No. 1, pp. 1-298, figs. 1-78.

1922b. Einige terrestre Isopoden von den Juan Fernandez Inseln. Skottsberg, C. The Natural History of Juan Fernandez and Easter Island, vol. 3, pp. 277-288, 4 text figs.

WHITE, A.

1847. List of the specimens of Crustacea in the collection of the British Museum. London.

## INDEX

Generic and subgeneric names are in capitals.  
Synonyms are in italics.

- ACANTHODILLO, 15  
*ACTAECIA*, 9  
**ACTAECIA**, 10  
*albospinosus*, *Armadillo*, 20  
albospinosus, *Spherillo*, 19  
**ALLONISCUS**, 13, 14  
*ambitiosus*, *Spherillo*, 20  
*angusticauda*, *Philoscia* (*Setaphora*), 12  
*annectens*, *Austridotea*, 24  
*annectens*, *Austridotea* (*Austridotea*), 24  
*appressa*, *Philoscia* (*Heroldia*), 11  
**ARMADILLIDIUM**, 23  
**ARMADILLIDIUM**, 23  
**ARMADILLO**, 15, 16, 18, 23  
**ARMADILLO**, 16, 17, 18, 19, 20, 21, 22, 23  
**ARMADILLONISCUS**, 10  
*asselus*, *Oniscus*, 13  
*assimilis*, *Phreatoicus*, 24  
*auklandiae*, *Deto* (*Vinetta*), 9  
*auklandiae*, *Scyphax*, 10  
*Aucklandicus*, *Armadillo*, 21  
*australis*, *Armadillo*, 20  
*australis*, *Notoniscus*, 9  
*australis*, *Philoscia*, 10  
**AUSTRIDOTEA**, 24
- BATHYTROPA, 15  
*benhami*, *Austridotea* (*Austridotea*), 24  
*bicornis*, *Cerberoides* (*Oniscomorphus*), 14  
*bidens*, *Armadillo*, 20  
*bipunctatus*, *Spherillo*, 20  
*bocki*, *Cubaris*, 16  
*bocki*, *Melanesillo*, 18  
*brevicauda*, *Cerberoides* (*Congloboniscus*), 14  
*brevicornis*, *Philoscia* (*Paraphiloscia*), 12  
*brevis*, *Alloniscus*, 13  
*brevis*, *Spherillo*, 20  
*bucculenta*, *Deto* (*Vinetta*), 10
- canalensis*, *Cubaris*, 16  
*canalensis*, *Philoscia* (*Heroldia*), 11  
*canaliculatus*, *Merulana*, 18  
*carinata*, *Merulanella*, 18  
*carinulatus*, *Spherillo*, 20  
**CERBEROIDES**, 14  
**CHAETOPHIOSCIA**, 11  
*chathamensis*, *Merulana*, 18, 22  
*cinctetus*, *?Pyrgoniscus*, 18  
*cinerea*, *Cubaris*, 16  
*collaris*, *Orodillo*, 19  
*commensalis*, *Trichoniscus*, 8  
**CONGLOBONISCUS**, 14  
*cookii*, *Oniscus*, 13  
*cristatus*, *Porcellio*, 15  
**CRUREGENS**, 24  
**CUBARIS**, 15, 16, 18, 19, 20, 21, 22, 23  
**CUBARIS**, 16, 17  
*cursor*, *Ligia*, 7  
**CYLLOMA**, 10
- Danae*, *Armadillo*, 17  
*danae*, *Sphaerillo*, 20  
*danae*, *Spherillo*, 19  
**DET**, 9, 10  
*digitifera*, *Philoscia* (*Heroldia*), 11  
*dispersus*, *Spherillo*, 20  
*dollfusi*, *Merulanella*, 19
- echinata*, *Deto*, 9  
**ECHINODILLO**, 17  
**EMYDODILLO**, 17  
*eremitus*, *Mesodillo*, 19  
*erinaceus*, *Acanthodillo*, 15  
*erinaceus*, *Spherillo*, 20  
*euchroa*, *Actaecia*, 10  
*euchroa*, *Armadilloniscus*, 10

- EUPORCELLIO, 14  
 EURYLIGIA, 7  
*exilis*, Merulana, 18  
*exotica*, Ligia (Megaligia), 6
- fasciata*, Philoscia (Setaphora ?), 12  
*fissus*, Sphaerillo (Sphaerillo), 20  
*floresianus*, Alloniscus, 14  
*fontanus*, Cruregens, 24  
*fragilis*, Philoscia (Paraphiloscia), 12  
*frontalis*, Hawaiodillo, 17
- galbineus*, Cubaris, 16  
*gaudichaudi*, Ligia, 7  
**GEOLIGIA**, 7  
*gracilis*, Philoscia (Paraphiloscia), 12  
*grandis*, Ligia, 7  
*graniger*, Porcellio, 14
- hamiltoni*, Spherillo, 20  
**HAPLOPHTHALMUS**, 8  
*hawaiensis*, Ligia, 7  
*hawaiensis*, Ligia (Megaligia), 7  
*hawaiensis*, Spherillo, 20  
**HAWAIODILLO**, 17, 18  
*hebridarum*, Melanesillo, 18  
*hebridarum*, Sphaerillo, 18  
*helmsii*, Haplophthalmus, 8  
*helmsii*, Notoniscus, 8  
**HEMINAGARA**, 15  
**HEROLDIA**, 11  
*humboldtii*, Philoscia (Heroldia), 11
- IDOTEA**, 24  
*incisus*, Cubaris, 16  
*inconspicuus*, Armadillo, 20  
*insularis*, Armadillo (Pseudosphaerillo), 16  
*insularum*, Nagara (Nagara), 15  
*intermedius*, Scyphax, 9
- javanensis*, Cubaris, 16
- “*kencpurensis*, Oniscus,” 13  
*kenepurensis*, Phallonuscus, 13  
*kermadecensis*, Trichoniscus, 8  
*kirki*, Phreatoicus, 24
- lacustris*, Austridotea (Notidotea), 24  
*lacustris*, Cubaris, 16  
*lacustris*, Idotea, 24  
*lacustris*, Pentidotea, 24  
*laevis*, Porcellio (Mesoporcellio), 14  
**LAEVOPHILOSIA**, 11  
*latissima*, Ligia (Euryligia), 7  
*latreillei*, Tylos, 9  
*“lifouensis”*, Cubaris, 17  
*lifouensis*, Sphaerillo (Xestodillo), 20  
*lifuensis*, Cubaris, 16  
*lifuensis*, ?Cubaris, 17  
*lifuensis*, Philoscia, 10  
*lifuensis*, Philoscia, 11  
**LIGIA**, 6, 7  
**LIGYDA**, 6  
*litigiosa*, Ligia, 7  
**LOBODILLO**, 18  
*longicornis*, Cubaris, 16  
*longicornis*, Philoscia (Nesoniscus), 12  
*longistylis*, Styloniscus, 7  
*lundi*, Cubaris, 16  
**LYGIA**, 7
- macmahoni*, Spherillo, 21  
*magellanicus*, Trichoniscus (Mega-trichoniscus), 8  
*magnifica*, Doto, 9  
*magnus*, Scyphoniscus, 10  
*malleata*, Ligia, 7  
*marginatus*, Spherillo, 21  
*marquesarum*, Sphaerillo (Xestodillo), 21  
*medius*, Nesodillo, 16  
**MEGALIGIA**, 6, 7  
**MEGATRICHONISCUS**, 8  
**MELANESILLO**, 18  
*melanurus*, Spherillo, 21  
**MERULANA**, 18  
**MERULANELLA**, 18, 19  
**MESODILLO**, 19  
**MESOPORCELLIO**, 14  
**METOPONORTHUS**, 14  
*milleri*, Cubaris, 16  
*monolinus*, Spherillo, 21  
*montana*, Philoscia (Plymophiloscia), 12  
*montanum*, Echinodillo, 17

- monticola, Philoscia (Heroldia), 11  
 montivagus, Sphaerillo, 21  
 murina, Cubaris, 16  
 muscorum, Oniscus, 10
- NAGARA, 15  
*neo-zelandicus*, *Porcellio*, 14  
*neozelandicus*, Tylus, 9  
*NESODILLO*, 16, 17  
*NESOLIGIA*, 7  
*NESONISCUS*, 12  
*noduligera*, Merulana, 18  
*noduligerus*, Philoscia (Nesoniscus), 12
- NOTIDOTEA, 24  
*NOTONISCUS*, 8, 9  
*novae-zelandiae*, *Doto*, 10  
*novaezealandiae*, Ligia (Nesoligia), 7  
*novae-zelandiae*, *Oniscus*, 10  
*novae-zelandiae*, Philoscia, 11  
*novizelandiac*, Lygia, 7
- oahuensis*, Alloniscus, 13  
*oceanicus*, Oniscus, 6  
*OCHETODILLO*, 19  
*officinalis*, Armadillo, 15  
*officinalis*, Cubaris, 15, 16  
*oliveri*, Philoscia, 11  
*ONISCOMORPHUS*, 14  
*ONISCUS*, 6, 10, 13  
*ONISCUS*, 10, 11, 13  
*opinensis*, Actaecia, 10  
*ornatus*, Scyphax, 9  
*ORODILLO*, 19  
*OROSCIA*, 12  
*ortonedae*, Rhyscotus, 15  
*otakensis*, Trichoniscus, 8
- pacificum*, ?*Armadillidium*, 23  
*pacificus*, Cubaris, 17  
*pallida*, Ligia, 7  
*paniensis*, Philoscia (Oroscia), 12  
*PARAPHIOSCIA*, 12  
*PARAVIREIA*, 25  
*PENTIDOTEA*, 24  
*perconvexus*, Alloniscus, 13  
*perkinsi*, Armadillo, 17  
*perkinsi*, Hawaiodillo, 17  
*perkinsi*, Ligia (Megaligia), 7
- persona, Philoscia, 11  
*PHALLONISCUS*, 13  
*PHILONGRIA*, 8  
*PHILOSCIA*, 9, 12  
*PHILOSCIA*, 10, 11, 12, 13  
*PHILOSCODILLO*, 14  
*philoscoides*, Ligia, 7  
*PHILOUGRIA*, 8  
*PHILYGRIA*, 8  
*phormianus*, Trichoniscus, 8  
*PHREATOICUS*, 23, 24  
*pictus*, Sphaerillo, 21  
*pilosus*, Cerberoides, 14  
*pilosus*, Cerberoides (Philoscodillo), 14  
*plasticus*, Cubaris, 17  
*PLYMOPHIOSCIA*, 12  
*politus*, Sphaerillo (Xestodillo), 21  
*PORCELLIO*, 14  
*PORCELLIO*, 14  
*PORCELLIONIDES*, 14  
*porteri*, Ligia, 7  
*pronyensis*, Cubaris, 17  
*pruinosus*, Porcellio, 14  
*pruinosus*, Porcellionides, 14  
*PSEUDOPHIOSCIA*, 12  
*PSEUDOSPHEARILLO*, 16  
*pubescens*, Philoscia, 11  
*pubescens*, Philoscia, 12  
*punctatus*, Oniscus, 13  
*punctatus*, Phalloniscus, 13  
*purpurescens*, Armadillo, 15  
*pusillus*, Trichoniscus, 8  
*pygmaeus*, Sphaerillo (Sphaerillo), 21  
*PYRGONISCUS*, 18
- quadrata*, Ligia, 7
- reticulata, Philoscia (Heroldia), 11  
*RHYSCOTUS*, 15  
*robusta*, *Doto*, 9  
*rosea*, Philongria, 8  
*rouxi*, Armadillo, 15  
*rouxi*, Armadillo (Pseudosphaerillo), 16  
*rouxi*, Philoscia (Chaetophiloscia), 10  
*rouxi*, Philoscia (Chaetophiloscia), 11  
*rouxi*, Schismadillo, 19  
*ruficornis*, Spherillo, 21  
*rufomarginatus*, Spherillo, 22

- rugosa, *Ligia*, 7  
rugosus, *Armadillo*, 18  
rugulosus, *Spherillo*, 22
- samoensis*, *Armadillo*, 23  
*sarasini*, *Cubaris*, 17  
*sarasini*, *Philoscia* (*Wahrbergia*), 13  
*scaber*, *Porcellio*, 14  
*scaber*, *Porcellio* (*Euporcellio*), 14  
*scamnorum*, *Melanesillo*, 18
- SCHISMADILLO*, 19  
*SCHISMADILLO*, 22
- SCYPHAX*, 9  
*SCYPHAX*, 10  
*SCYPHONISCUS*, 10  
*setaceus*, *Sphaerillo*, 22
- SETAPHORA*, 10, 12, 13  
*SEТАPHORA*, 12, 13  
*setiger*, *Scyphax*, 9  
*sharpī*, *Hawaiodillo*, 18  
*societatis*, *Sphaerillo*, 22  
*societatus*, *Philoscia*, 11  
*speciosus*, *Armadillo*, 8  
*speciosus*, *Spherillo*, 22
- SPHAERILLO*, 16, 17, 18, 19, 20, 21, 22, 23  
*SPHERILLO*, 15, 16, 17, 18, 20  
*SPHERILLO*, 19, 20, 21, 22, 23  
*spicatus*, *Spherillo*, 22  
*spinosis*, *Spherillo*, 22  
*squamata*, *Trichorhina*, 15  
*squamatus*, *Spherillo*, 22  
*squamosa*, *Philoscia*, 11  
*squamosus*, *Tridentodillo*, 23  
*squamuligera*, *Philoscia* (*Oroscia*), 12  
*stenosoma*, *Philoscia* (*Paraphiloscia*), 12
- STYLONISCUS*, 7
- STYLONISCUS*, 8  
*sulcatus*, *Ochetodillo*, 19  
*suteri*, *Cubaris*, 17
- tahitiensis*, *Nagara* (*Heminagara*), 15  
*taragensis*, *Spherillo*, 22  
*testudinalis*, *Spherillo*, 22  
*testudo*, *Emydodillo*, 17  
*thermophila*, *Bathytropa*, 15  
*thomsoni*, *Trichoniscus* (*Megatrichoniscus*), 8
- tongensis*, *Armadillo*, 23  
*translucida*, *Merulana*, 18
- TRICHONISCUS*, 8
- TRIC HORHINA*, 15
- TRIDENTODILLO*, 23
- truncata*, *Philoscia* (*Setaphora*), 13  
*tuberculatus*, *Oniscus*, 10  
*turgifrons*, *Rhyscotus*, 15
- TYLOS*, 9
- typicus*, *Paravireia*, 25  
*typicus*, *Phreatoicus*, 23
- verrucosus*, *Trichoniscus*, 8
- VINETTA*, 9, 10
- violacea*, *Philoscia*, 9
- vitiensis*, *Ligia*, 7  
*vitiensis*, *Spherillo*, 19, 23
- vulgare*, *Armadillidium*, 23  
*vulgare*, *Armadillo*, 23
- wahrbergi*, *Merulanella*, 19
- WAHRBERGIA*, 13
- waitatensis*, *Scyphoniscus*, 10
- XESTODILLO*, 20, 21, 23
- zealandicus*, *Porcellio*, 14
- zebricolor*, *Sphaerillo* (*Xestodillo*), 23