



A. S. Hitchcock

SMITHSONIAN INSTITUTION
UNITED STATES NATIONAL MUSEUM

CONTRIBUTIONS

FROM THE

UNITED STATES NATIONAL HERBARIUM

VOLUME VIII, PART 2

ECONOMIC PLANTS OF PORTO RICO

By O. F. COOK and G. N. COLLINS



WASHINGTON
GOVERNMENT PRINTING OFFICE

1903

NOTE.

ISSUED JUNE 27, 1903.

P R E F A C E .

One influence of the extension of American jurisdiction over Porto Rico and the Philippine Islands, in the year 1898, was to create a demand for information about the vegetation of those islands. All contained a tropical flora, and American botanists had devoted little attention to the study of tropical plants. The information demanded regarding the Porto Rican flora was largely of a popular and commercial character and was not at all to be satisfied by the existing and available technical works on the West Indian flora. In the course of their investigations on tropical agricultural plants in Porto Rico, Mr. O. F. Cook, botanist in charge of investigations in tropical agriculture, Department of Agriculture, and his assistant, Mr. G. N. Collins, have collated a large amount of information on the economic plants of that island such as is not suitable for incorporation in their detailed reports on staple and special agricultural plants, which are published by the United States Department of Agriculture. This information has been brought together in the present paper on Economic Plants of Porto Rico. Fortunately the plant names in use among the people of Porto Rico are applied with approximate precision, so that by adding the botanical identification of a plant to its Porto Rican popular name we have in most instances a direct key to its botanical name and position. The use of illustrations reproduced from good photographs conveys in a simple and direct way information which no amount of technical description could replace. Availing themselves of personal observation, the facts published by others, a study of collections, the use of photographs, and the intimate personal knowledge the Porto Ricans have of their useful plants, the authors have produced a work which will be of great value to botanists and to all other persons seeking information about the common and useful plants of Porto Rico.

FREDERICK V. COVILLE,
Curator of Botany.

CONTENTS.

	Page.
Introduction.....	57
Economic plants imperfectly known.....	58
Materials included and arrangement of list.....	58
Misplaced common names.....	59
References to botanical literature.....	60
Books on Porto Rican plants.....	60
Personal field work, acknowledgements, and desideratum.....	62
Catalogue of economic plants of Porto Rico.....	63

ILLUSTRATIONS.

PLATES.

PLATE XIII. Porto Rico Royal Palm (<i>Roystonea borinquena</i>).....	Frontispiece.
	Facing page.
XIV. Okra (<i>Abelmoschus esculentus</i>) showing healthy and diseased fruits. From Santurce.....	63
XV. Acacia amarilla (<i>Albizzia lebbek</i>). Near Juana Diaz.....	70
XVI. <i>Alpinia nutans</i> . At Caguas.....	73
XVII. <i>Anthurium acule</i>	82
XVIII. <i>Antigonon leptopus</i> . Coamo Springs.....	82
XIX. Flamboyant blanco (<i>Bauhinia kappleri</i>). Coamo Springs.....	90
XX. Ucar tree (<i>Bueida buceras</i>). Coamo Springs.....	96
XXI. Almacigo (<i>Bursera simaruba</i>). Coamo Springs.....	97
XXII. Algodon de seda (<i>Calotropis procera</i>). Road near Guayama.....	101
XXIII. Palo de burro (<i>Capparis cynophallophora</i>), fruit and leaves.....	104
XXIV. Ceiba tree (<i>Ceiba pentandra</i>). Near Ponce.....	111
XXV. Stem and edible fruit of <i>Cereus</i> sp. At Guayama.....	112
XXVI. Icaco (<i>Chrysobalanus icaco</i>).....	114
XXVII. Péndulo colorado (<i>Citharexylum quadrangulare</i>). Road between Ponce and Adjuntas.....	116
XXVIII. Cupey (<i>Clusia rosea</i>). Near Yauco.....	119
XXIX. Job's tears (<i>Coix lachryma-jobi</i>).....	122
XXX. Moral tree (<i>Cordia macrophylla</i>). Coamo Springs.....	127
XXXI. Cerezas (<i>Cordia nitida</i>).....	127
XXXII. Calabash tree (<i>Crescentia cujete</i>). Near Rio Piedras.....	129
XXXIII. Concombre (<i>Cucumis anguria</i>).....	130
XXXIV. Virgin forest of tabonuco (<i>Daeryodes hexandra</i>). At Isolina.....	132
XXXV. Jaguey (<i>Ficus lentiginosa</i>). Road between Ponce and Adjuntas.....	145

	Facing page.
PLATE XXXVI. Guácima (<i>Guazuma guazuma</i>). Coamo Springs	155
XXXVII. <i>Heckeria peltata</i> inflorescence and center of leaf	158
XXXVIII. <i>Heckeria umbellata</i> inflorescence and center of leaf	158
XXXIX. Talantala (<i>Herpetica alata</i>). At Cataño	159
XL. Lira (<i>Hibiscus schizopetalus</i>). Garden of the Captain-General. Rio Piedras	160
XLI. Pods of algarrobo (<i>Hymenaea courbaril</i>). Ponce Market	164
XLII. Physic nut (<i>Jatropha curcas</i>). Cataño	171
XLIII. Flowers, seeds, and seedlings of mangle blanco (<i>Laguncularia racemosa</i>). Cataño	173
XLIV. Molinillo (<i>Leonotis nepetalifolia</i>)	174
XLV. China berry (<i>Melia azedarach</i>). Coamo Springs	190
XLVI. Cundeamor (<i>Momordica charantia</i>)	193
XLVII. Cork wood (<i>Ochroma lagopus</i>). Near Ponce	205
XLVIII. Spineless Opuntia. Near Guayama	206
XLIX. Para grass (<i>Panicum molle</i>)	211
L. Emajagua (<i>Paritium filiaecum</i>)	212
LI. <i>Pentarraphia albiflora</i>	215
LII. Tachuelo (<i>Pictetia aristata</i>)	218
LIII. Tachuelo (<i>Pictetia aristata</i>), showing aristate leaves	218
LIV. Flame tree (<i>Poinciana regia</i>). In the Plaza, Caguas	223
LV. Berengena cimarrona (<i>Solanum torquatum</i>)	243
LVI. Anacagüitas (<i>Stereulia carthagenensis</i>). Near Ponce	245
LVII. Almendra (<i>Terminalia catappa</i>)	250
LVIII. Santa Maria (<i>Thespesia populnea</i>). Near Ponce	253
LIX. Santa Maria (<i>Thespesia populnea</i>), showing fruit	253
LX. Vincetoxicum sp	262

TEXT FIGURES.

	Page.
Fig. 12. Lawn of <i>Meibomia triflora</i> . Coamo Springs	189
13. Almendra, fruit of <i>Terminalia catappa</i> . Cataño	250

ECONOMIC PLANTS OF PORTO RICO.

By O. F. COOK and G. N. COLLINS.

INTRODUCTION.

Very few tropical countries have been provided with satisfactory manuals of botany by means of which the resident or the traveler can identify the plants and secure access to information regarding their uses and other noteworthy characteristics.

There is, as yet, no manual of the botany of Porto Rico nor any handbook of tropical cultivated plants which can be recommended to those who seek an introduction to the indigenous flora of the island and the economic species grown for food, ornament, and other purposes, or escaped and intermingled with the native vegetation. Fortunately, however, clues to the identity of a large proportion of the plants of Porto Rico are readily obtainable through the medium of the common names. The natives of Porto Rico have been noted by many travelers as having a larger series of popular names and as using them with more precision than the inhabitants of other parts of the American tropics. Certain it is that many species have received separate names in Porto Rico which in Cuba are not distinguished except by botanists. This fact is to be connected, perhaps, with the existence of a large rural population, which has preserved many names obtained from their Indian ancestors and the Spanish settlers with whom these amalgamated. The names are in many cases entirely local, not known outside the island, and quite different from those applied in Mexico and Cuba. It is accordingly believed that a list of Porto Rican names of economic plants, although necessarily imperfect, would be of special convenience at the present juncture when the native population is beginning the study of English and while numerous Americans are struggling to comprehend the language, products, and vegetation of a new and interesting country.

Some plants have several common names which may be confined to different districts, and in some cases the same names are applied to different plants in different places. Of course this popular knowledge lacks scientific accuracy and frequently breaks down when similar species are in question. The identification of single individuals

should always be verified by cumulative testimony and a consensus of the most intelligent opinion obtainable.

ECONOMIC PLANTS IMPERFECTLY KNOWN.

Owing partly to the fact that some of the botanists who have written most extensively on the West Indian flora have carefully avoided reference to the uses of the plants they studied, our knowledge of the economic plants of the West Indies is still in a very fragmentary condition. Moreover, a considerable amount of the existing literature is now out of date and unreliable on account of the uncertainty attaching to many of the names used by the earlier semipopular writers. The British West Indies have received the most attention, and the French islands are second in the amount of printed information. On the Cuban flora, as well, a considerable body of literature exists, in comparison with the extreme paucity of scientific information regarding Porto Rico. The island has not attracted, to any considerable extent, the interest of European or American naturalists, and the local prosecution of study received only negative encouragement.

Although the area is small, the topography is so complicated and the climatic and other conditions so varied that anything like an exhaustive knowledge of the wild plants will be very difficult to obtain. In the case of the cultivated economic species the same difficulty of extreme localization is apparent. Plant introduction has been carried on only in an extremely desultory manner. Transportation is difficult, and the numerous towns have had connection, for much of the time, only by sea, so that the dissemination even of successful species has been slow and difficult.

It is manifestly impossible to enter to best advantage upon the task of improving agricultural conditions in Porto Rico without more adequate knowledge of the nature and results of past experiments, and it would be useless and wasteful to reintroduce species already growing in the island except where superior varieties can be secured.

MATERIALS INCLUDED AND ARRANGEMENT OF LIST.

This paper includes miscellaneous information on the principal cultivated plants of Porto Rico, brief notes on many of the minor economic plants, and a list of all the native names of plants which have thus far been recorded from the island, with references to the scientific names of the species to which they are applied as far as these have been determined. As there are no botanical publications in either English or Spanish which give an even approximately complete treatment of the flora, it is believed that the present list of names and the brief notes accompanying will be found of use both to visitors and to residents of the island.

Owing to the fact that the same name is often applied to similar though botanically distinct species, and that various names are used

for the same species in different parts of the island, it has seemed best in the preparation of this list not to separate the information from the native name with which it was collected except where the botanical identity of the plants has been established, even though a guess might have been made with some probability of correctness. By the system of cross references it will, however, be possible for those who make use of the index to find all the available information, wherever classified.

MISPLACED COMMON NAMES.

Although sufficiently reliable to be of much assistance to strangers who desire to secure acquaintance with the local flora, the application of names in Porto Rico is in many cases misleading, well-known terms being used in the most novel relations. Thus the "níspero" of Porto Rico is not the loquat or Japanese medlar (*Eriobotrya japonica*) as in Spain, but the sapodilla (*Achras sapota*). Quite a list of such misapplied names might be made, but a few of the more prominent will suffice to show how unreliable are inferences drawn from verbal or even from written information which is not based on actual contact with the objects. Thus in Porto Rico the—

- Almendra (almond) is *Terminalia catappa*.
- Ciruela (plum) is *Spondias purpurea*.
- Castaña (chestnut) is *Artocarpus communis*, the bread fruit.
- Níspero (loquat) is *Achras sapota*, the sapodilla.
- Nuez moscado (nutmeg) is *Nectandra* sp.
- Pimienta (pepper or allspice) is *Amomis caryophyllata*, the bay-rum tree.
- Reseda (mignonette) is *Lawsonia alba*, the henna.
- Almacigo (mastic) is *Bursera simaruba*, the West Indian birch.
- Cereza (cherry) is *Cordia collococca* and other species.

Many plants also have different local names. Those supplied by Captain Hansard are mostly from the vicinity of Luquillo in the northeast corner of the island, and although they are names of trees several are not known to be used in other parts.

The English names of economic plants have also been included in this index, as far as practicable, in order to render the more certain the application of the Porto Rican designations for those who may have had previous experience in other tropical countries. The list of "colonial names" appended to Grisebach's flora has been drawn upon for this purpose to a limited extent, but is generally very unsatisfactory for the reason that the locality where a name is used is not given.

The names, English and Spanish, of some of the better known minor economic plants not yet recorded by botanists as growing in Porto Rico have been included in instances where these plants are known to have been introduced, and especially when they have become naturalized, in neighboring islands. It is extremely probable that some of these are already planted locally in Porto Rico, and as to those

which are not it is desirable to know the probabilities of their success and whence they may be most conveniently obtained.

REFERENCES TO BOTANICAL LITERATURE.

There is as yet nothing like a complete list of the plants of Porto Rico, and most of the literature is not even indexed. It has therefore seemed desirable, as a matter of convenience for those who may wish to become more familiar with the flora of the island, to add references to descriptions of many of the species enumerated below, although in the present paper no attempt has been made to follow the method or arrangement of formal botanical literature. Nor should it be inferred that the author cited is in all cases responsible for the statements accompanying a specific reference, although this is generally true where very short notes are given. The more recent literature of Urban has been cited in preference to the older writings in order to avoid multiplication of references, and because Urban generally gives complete synonymy from which the opinions of other writers can be traced.

BOOKS ON PORTO RICAN PLANTS.

The botany of Porto Rico is far from complete, and very little of it has been written in the English language. Descriptions of plants found in Porto Rico are scattered through a vast number of publications which can not be enumerated here, but brief notices of the more important papers may not be out of place. But two authors have attempted a connected sketch of the Porto Rican flora, and the efforts of these not only remain incomplete in that they do not cover the entire series of families of flowering plants, but the lists are also partial and local, as the writers themselves realized.

The first of these sketches was that of Don Domingo Bello y Espinosa.⁶ This consists almost entirely of a list of plants occurring, as explained by the author, in the triangular area included between Aquadilla, Lares, and Guanica. The prefatory note states that the writer resided in Porto Rico for thirty years, presumably at Mayaguez, and that his botanical studies were carried on in the intervals of a busy professional life. Considering this fact and the other difficulties under which he must have labored, the result is certainly most creditable, and the field observations and notes represent a distinct contribution to our knowledge of the flora of the area covered, although the adjustment of Bello's systematic treatment has given his successors considerable trouble. Many popular names are given, but no economic notes are included, although the useful species

⁶ Domingo Bello y Espinosa. Apuntes para la flora de Puerto-Rico. Anales de la Sociedad española de historia natural, tomo X, pp. 230-304, 1881; tomo XII, pp. 103-130, 1883.

are not omitted from their places in the natural sequence. Several new species are described, and the probable existence of others is often suggested.

The second of these sketches and the most important contribution to the botany of Porto Rico is the *Flora* projected by Dr. A. Stahl, of Bayamon, but unfortunately only partially published. Although of German paternity Dr. Stahl is a native of Porto Rico, and his scientific achievements are the result of tireless energy and perseverance under the most discouraging circumstances. Not only did he receive no public or official assistance or encouragement in the self-imposed task of making known the flora of the island, but his scientific labors drew upon him instead the suspicions of the government, and resulted in imprisonment and banishment on three or four occasions, without the satisfaction of even an alleged reason for his arbitrary ill treatment. In spite of public indifference and official animosity six parts of the *Flora* were issued, at the expense of the author, having been prepared in the intervals of his professional life as a physician. Publication ceased in 1888, and Dr. Stahl no longer hopes to continue the work. In addition to his botanical studies Dr. Stahl has written and made collections bearing upon medical, archaeological, and zoological subjects, and has well earned the distinction of being the first resident Porto Rican naturalist. The common names of Dr. Stahl's *Flora* appear also in Professor De la Maza's *Diccionario*,^b in which other names not yet reported from Porto Rico have been found and verified.

Further acknowledgement is due to Capt. Arthur C. Hansard, a retired English officer, for several years resident in the northeastern part of the island. A list of native names previously published by him in the *San Juan News* was carefully copied and the author then most obligingly revised and extended it, submitting as well to many questions which his extended residence and previous travels in other tropical countries rendered him particularly qualified to answer.

Apparently unknown to subsequent writers, a much more extended list containing the popular names of nearly 200 Porto Rican woods, with data regarding density, size, color, texture, and uses had appeared as a catalogue of a collection exhibited at the Agricultural Exhibition held in Madrid in 1857. This list has been safely preserved and securely hidden away in a bulky quarto volume entitled *Memoria sobre los Productos de la Agricultura española reunidos en la Exposición general de 1857*, published in Madrid between 1859 and 1861, and containing over 1,200 pages. The Porto Rican woods occur on pages 470 to 477 and are interjected among other material with which there is little apparent connection. Neither is there any statement as to the authorship or origin of the list, nor is it surprising that Maza,

^a Stahl, A. *Estudios para la flora de Puerto-Rico*. 1884-1888.

^b Manuel Gómez de la Maza. *Diccionario botánico de los nombres vulgares cubanos y puerto-riqueños*.

Hill, and other authors have overlooked it in making their compilations. The spelling of many names is different from that found elsewhere and there are some evident typographical errors, such as "diametro en pies," which must be intended for inches, as well as impossible specific gravities probably due to careless proof reading. On the whole, however, this anonymous contribution is an important one, and certainly represents a large amount of work. Many of the names are not yet identifiable botanically and the data are in such cases placed with the Porto Rican designation. Credit is given to "Exp. 1857," since a personal reference to the author is impossible.

For data respecting the characters and uses of woods we have drawn very freely upon Grosourdy's rare and valuable work on West Indian botany."

The native names used in the publications of Mr. Robert T. Hill, as well as those appearing in other recent works upon Porto Rico, have also received consideration, with the intention of rendering as definite as possible the information which has been collected.

PERSONAL FIELD WORK, ACKNOWLEDGMENTS, AND DESIDERATUM.

The writers made two trips to Porto Rico, one in 1899, the other in 1901, which covered the different types of floral and agricultural areas of the island. The specimen numbers cited in the catalogue refer to the collections made on these visits.

In the labor of compiling data from these and other sources we have received much friendly assistance from Messrs. Henry E. Baum, Edgar Brown, and William R. Beattie, of the Office of Botanical Investigations, Department of Agriculture, to whom our thanks are tendered herewith.

Not all the native names of woods and other plants have yet been identified botanically, and authentic flowering or fruiting specimens of such will be welcomed by the authors. Such specimens for identification should be of reasonable size—that is, 12 or 15 inches in length—and should be pressed and dried before mailing, otherwise they may arrive in a decayed condition and utterly worthless for study. With correspondents who will take the trouble to prepare good specimens of economic plants and information concerning their local names and uses, arrangements will be made for the transportation of packages without cost to the sender.

"Grosourdy, René de, M. D. *El médico botánico criollo*. 1864. 4 vols.



OKRA (*ABELMOSCHUS ESCULENTUS*), SHOWING HEALTHY AND DISEASED FRUITS.

CATALOGUE OF ECONOMIC PLANTS OF PORTO RICO.

Abacá. See *Musa textilis*.

Abejuelo.

A name given in Grosourdy, evidently the same as Abeyuelo and Abelluelo, with which the dimensions agree. The above author describes the wood as rose-colored, strong, and breaking with a vertical fracture. Commonly used for shelving. (Grosourdy, 2: 356.)

Abelluelo. See *Colubrina ferruginosa*.

Abelmoschus abelmoschus. ALGALIA.

Family Malvaceae: an annual or biennial shrub, 1 to 2 meters high, cultivated for the flowers and for the seeds, which have a strong odor of musk. At Cataño this was called "caballo asi." (Stahl, 2: 87, as *Abelmoschus moschatus*.)

Abelmoschus esculentus. GUINGAMBO. OKRA. PLATE XIV.

A woody annual or biennial, 1 to 3 meters high, cultivated for its large fruit capsules, known as "okra" and "gumbo." Bello notices two varieties, one pyramidal, and the other cylindrical, the latter being more delicate. (Stahl, 2: 86.)

Abeyuelo.

A tree from the northern parts of the island; height, 40 to 50 feet (12 to 15 meters); diameter, 12 to 14 inches (30 to 35 centimeters). Wood rose-colored, hard; specific gravity, 0.903; used in cabinet-making. (Exp. 1857.) Probably this is intended for the same name as abelluelo.

Abispillo.

A forest tree reported by Captain Hansard from the vicinity of Luquillo. Wood not used; specific gravity, 0.75. Perhaps the same as "avipillo" of Sintenis. *Maypea caribaea*, from Fajardo; avispillo is said to be *Turpinia paniculata*. A tree from all parts of the islands; height, 40 to 50 feet (12 to 15 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood chestnut-brown, hard; specific gravity, 0.601; used for lumber. (Exp. 1857.)

Grosourdy describes it as a tree abundant in all parts of the island, 40 to 50 feet (12 to 15 meters) high, 12 to 24 inches (30 to 60 centimeters) in diameter. The wood is said to be brittle, breaking with an oblique fracture; the heart brown; the sapwood white; used for flooring. (Grosourdy, 2: 356.)

Abrojo. See *Tribulus cistoides* and *T. maximus*.

Abrus precatorius. PERONIA.

A climbing, leguminous shrub; pubescent, biennial; partial to sandy soil; flowers in summer and autumn. The small, bright-red beans are extremely hard and are used as beads for rosaries. A highly toxic albuminoid, derived from this species, has been described under the name abrine. (Stahl, 3: 51.)

Abutilon leiospermum. PICHANA MOCHA.

Family Malvaceae: a pubescent shrub, common on the south side of the island. (Stahl, 2: 73.)

Abutilon pedunculare. PICHANA PEDUNCULAR.

Not found by Stahl on the north side of the island: reported by Bello from the south side.

Abutilon periplocifolium. See *Wissadula rostrata*.**Abutilon umbellatum.** MALVAVISCO CIMARRON.

From the vicinity of Coamo (Sintenis). The common name was heard at Santurce.

Acacia amarilla. See *Albizzia lebbek*.**Acacia angustiloba.**

A leguminous plant, reported from Porto Rico by Bello, with the common name "cojobilla," but Stahl thinks the identification erroneous, and would refer Bello's plant to *Piptadenia peregrina*.

Acacia arabica. GUM ARABIC TREE.

A native of the drier districts of Egypt, the Soudan, and East Africa. Cultivated in Jamaica and other islands of the West Indies and, according to Grisebach, naturalized in Antigua. It might be worth while to plant this species in the dry southwestern part of Porto Rico.

Acacia berteriana. ACACIA SILVESTRE.

According to Stahl, a small tree of 3 to 5 meters, preferring the summits of hills and shady situations. According to Grisebach, this species is abundant in some localities in Jamaica. Bello gives the common name "cojoba" for this species. (Stahl, 3: 150.)

Acacia farnesiana. AROMO.

A small tree introduced from the Old World: cultivated and now growing spontaneously in many parts of the West Indies, including Porto Rico. In the Mediterranean region it is extensively grown for the fragrant flowers, which are used in the manufacture of perfumery.

The wood is very heavy and solid, reddish-brown in color, and very handsomely grained. It is nearly all heart and has a specific gravity of 1.117. This wood can be used to good advantage wherever strength and durability are required. It would appear that it would also be suitable for furniture. (Grosourdy, 2: 362.)

Acacia lebbek. See *Albizzia lebbek*.**Acacia nudiflora.** ACACIA NUDOSA.

Stahl considers the occurrence of this species in Porto Rico doubtful, but it is known from St. Thomas and the neighboring Virgin Islands; also from Haiti. (Stahl, 3: 147.)

Acacia nudosa. See *Acacia nudiflora*.**Acacia palida.** See *Leucaena glauca*.**Acacia puertorriqueña.** See *Calliandra portoricensis*.**Acacia sarmentosa.** ACACIA ZARZA.

A climbing, spiny shrub, widely distributed in the West Indies and South America. By some considered synonymous with *A. riparia*. (Stahl, 3: 148.)

Acacia sylvestre. See *Acacia berteriana*.**Acacia zarza.** See *Acacia sarmentosa*.**Acajou.** See *Guarea swartzii*; also *Swietenia mahagoni*.**Acaju.** See *Anacardium occidentale*.

This and the preceding are different spellings of the same word, evidently of French origin, in which language it is more properly applied to *Swietenia mahagoni*, but is also used with *Anacardium occidentale* and *Cedrela glaziovii*.

Acalypha bisetosa.

Family Euphorbiaceae; found by Sintenis, growing wild in the mountains near Yabucoa. Other species are cultivated as ornamentals.

Acana. See *Labourdonnaisia albescens*.

A tree growing on hills; has great girth and height. The wood has a specific gravity of 1.09 and makes fair boards, good for boat building. It has a dry, white fruit, similar in taste, but not in form, to that of the algarrobo. Also called "jacana." (Hansard.)

Acanthotrichilia triacantha. See *Trichilia triacantha*.

Acayu. See *Anacardium occidentale*.

A form of the name "acaju."

Aceitillo. See *Simaruba tulae*.

A wood of this name, the locality not given, is described as strong and durable; very well suited to all kinds of coarse carpenter work where great strength is needed. (Grosourdy, 2: 356.)

Aceitillo labrado.

A tree from the northwest part of the island; height, 35 to 40 feet (10 to 12 meters); diameter, 35 to 40 inches (87 to 100 centimeters). Wood light yellow, hard; specific gravity, 0.947; used in cabinetmaking. (Exp. 1857.) (Grosourdy, 2: 356.)

Aceitillo liso.

A tree from the northwestern part of the island; height, 50 to 55 feet (15 to 17 meters); diameter, 35 to 40 inches (87 to 100 centimeters). Wood light yellow, hard; specific gravity, 0.789; used in making furniture. (Exp. 1857.) Grosourdy, 2: 356.)

Aceituna. See *Symplocos martinicensis*.

A tree abundant in the eastern part of the island; height, 35 to 40 feet (10 to 12 meters); furnishing a soft, light-colored wood, breaking with an oblique fracture; suitable for framework in houses. (Grosourdy, 2: 357.)

The European olive, *Olea europaea*, to which this name is properly applied, is not known to occur in Porto Rico.

Aceituna blanca. See *Symplocos martinicensis*.

A tree from the eastern part of the island; height, 12 to 15 feet (4 to 5 meters); diameter, 16 to 18 inches (40 to 45 centimeters). Wood white, rather hard; specific gravity, 0.646; used for weatherboards. (Exp. 1857.)

Aceituna cimarrona. See *Symplocos martinicensis*.

Aceituna macho.

A tree from the eastern part of the island; height, 35 to 40 feet (10 to 12 meters); diameter, 16 to 18 inches (40 to 45 centimeters). Wood, light-reddish, rather hard; specific gravity, 0.656; used for weatherboards. (Exp. 1857.)

Aceitunillo.

A tree reported by Captain Hansard from the northeastern part of the island; specific gravity of wood, 0.84. In Cuba this name is applied to *Alectoaricon punctatum*, a cultivated euphorbiaceous plant, native of Chile.

Achicoria de cabro. See *Erechtites hieracifolia*.

Achiote. See *Bixa orellana*.

A tree from all parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 8 to 9 inches (20 to 23 centimeters). Wood, white and very soft; specific gravity, 0.404; not used. (Exp. 1857.)

Achote. See *Bixa orellana*.

A shortening of Achiote, of questionable correctness.

Achras sapota. SAPODILLA. NÍSPERO.

A handsome tree belonging to the family Sapotaceae, commonly cultivated in Porto Rico and supposed to have been introduced from Venezuela. The applica-

tion of the name "nispero" to this species, though erroneous and misleading to strangers, is universal in Porto Rico, and has led Mr. Hill and possibly other writers to report the loquat as abundant, the latter fruit bearing the name nispero in Spain and in Spanish lexicons.

The sapodilla tree grows to a height of 25 or 30 feet (7 to 9 meters) and has shining dark green leaves. The fruit is somewhat larger than a hen's egg, with a rather rough, brownish skin, suggesting an Irish potato with earth adhering. It has a yellowish pulp and 4 or 5 black seeds. The wood is said to be soft, although Captain Hansard gives the specific gravity as 1.02. This tree furnishes the material imported in recent years in large quantities from Mexico under the name of "chicle," or "gum chicle," and used mostly in the manufacture of chewing gum. It is obtained by the evaporation of the milky juice secured by tapping the trees or by pressure from the fruit.

The sapodilla is one of the tropical fruits which, like the mango and avocado pear, may have a future in the American market. Experiments have been made by the Division of Pomology with sapodillas raised on the Florida Keys and they were found, contrary to expectations, to stand shipment well. The skin, while not thick or hard, does not seem particularly susceptible to decay, and it is believed that with proper care in harvesting and packing sapodillas could be kept in good condition for ten days or two weeks, to say nothing of the possibilities of cold storage. The fruit is also believed by some of the fruit experts to be one which is likely to become popular if it can be furnished in sufficient quantities to give it a regular place in the market.

The question of varieties of sapodillas has not received attention as yet, but should be given careful consideration if planting for export is undertaken. The testimony of travelers in Mexico and other countries where the fruit abounds, indicates that the natural differences in quality between the unselected seedlings are very appreciable so that a search for superior varieties might be richly repaid.

The fact that in the culture of the sapodilla the production of "chicle" could be made an alternative resource, when the fruit could not be properly marketed, would give the industry an advantage over many lines of fruit growing.

Grosourdy describes this tree as furnishing a very good wood, of good weight, more than ordinarily hard, compact in texture, and fine-grained. The color is light red with darker stripes. It is as strong and as good as "balata" or "pargo" and might replace these if it grew wild and was more abundant. Specific gravity, 1.021. (Grosourdy, 2: 413.)

Bello gives "ausubo" as the common name of a synonym of this species.

Acisanthera quadrata. CAMACEY DE CHARCOS.

Family Melastomaceae; a slender, somewhat woody herb growing in sandy overflowed situations. (Stahl, 4: 119.)

Acnistus arborescens. GALAN ARBOREO.

Family Solanaceae; a shrub or low tree, smooth, and much branched; grows in waste places and mountainous localities to an altitude of 500 feet (155 meters) or more; flowers in March. (Stahl, 4: 142.)

Grosourdy gives "palo de gallina" as the common name of this tree and describes it as a wild tree from Porto Rico, Trinidad, and as far as Caracas, 15 to 25 feet high (5 to 8 meters), with a trunk 4 to 6 inches (10 to 15 centimeters) in diameter. It furnishes a soft wood, dirty white in color, that breaks with a vertical fracture. Its most common uses are as fuel and for the framework of rude houses. He states that the common name is derived from the fact that the fruit kills chickens if eaten by them. (Grosourdy, 2: 406.)

Acrista monticola. PALMA DE SIERRA.

Family Arecaceae; among the mountains between Cayey and Guayama many summits are covered with the palma de sierra, probably in places which have

never been cleared. A few of the palms follow down the steeper uncultivated ravines. From a distance the crowns suggest royal palms, but a closer view renders the difference apparent. There is also no suggestion of the bulging trunk of *Roystonea*. In height the palma de sierra probably does not exceed the royal palm.

Acrocomia. COROZO PALM.

A genus of palms distributed through tropical America from Mexico and Cuba to Paraguay. All the species are of stocky, compact growth with a dense crown of numerous leaves. The trunk as well as the leaf stalks is usually armed with sharp spines, sometimes several inches long.

The economic value of *Acrocomia* lies mostly in the nut-like seeds, an inch or more in diameter, which are constructed on the principle of the cocoanut. The outer fibrous husk is oily and edible in some species, and there is a nearly solid kernel, from which oil is extracted for food as well as for soap making and other purposes for which the cocoanut and African oil palm are used. In Paraguay the collection of palm kernels from a species of *Acrocomia* has become a considerable industry and large quantities are exported to Europe. In order to free the nuts from the fibrous husk they are fed to cattle, after which they are cracked. The Porto Rican species has larger seeds than the Paraguayan and the husks are not known to be edible. The natives seem also to make little use of the meat except to eat it occasionally before fully ripe, when it tastes much like cocoanut. While the kernels, if extracted, could probably be marketed, it is possible that the labor required could find better remuneration in other work, though the subject might be worthy of investigation in districts where the palms are abundant.

The outer shell of the trunk of the corozo palm is extremely hard, but seems to be used in Porto Rico only for the manufacture of walking sticks. When well made these are both curious and beautiful. The general color is a deep rich reddish brown, irregularly nerved with the lighter color of the compacted fibrous strands.

From the leaves of *Acrocomia* a fiber of great strength and fineness has been obtained in some of the British West Indies, but it seems not to have been applied to any civilized purpose or to have become a regular article of commerce.

Acrocomia aculeata.

A species from Martinique, referred by Martius to *A. sclerocarpa*. It has not been noticed in Porto Rico.

Acrocomia media. COROZO.

Although totally different on close inspection this palm has a superficial resemblance to the royal palm, which often deceives travelers. The similarity lies mostly in the two facts that both the royal and "corozo" palms are more robust and stiffly erect than the cocoanut, and that the leaf divisions instead of lying horizontal and in one plane are tilted at different angles to the midrib, thus giving the foliage seen in the mass a somewhat unkempt appearance in comparison with the cocoanut.

In distinguishing the corozo palm from the royal palm when seen at a distance so great that the spines of the one and the columnar green leaf sheaths of the other can not be seen, recourse may be had to the following facts. The leaf crown of the corozo palm is much rounder, thicker, and more compact than that of the royal palm, since it contains many more leaves, and these persist much longer. The royal palm can also be known by the unopened leaves which project straight upward like flag poles or lightning rods, while in *Acrocomia* the leaves open as they are pushed out and seldom offer a suggestion of the spire-like effect.

Adansonia digitata.

Family Bombacaceae: the African baobab tree has been introduced into gardens in the West Indies, but is not known to have become naturalized.

Adelfa. See *Nerium oleander*.

Adelfa sencilla. See *Nerium oleander*.

The name means single oleander.

Adenanthera pavonina. PALO DE PERONIAS.

Also called "coralitos;" a leguminous tree, 5 meters high, introduced from India as an ornamental, and naturalized in some localities. (Stahl, 3: 135.)

Grosourdy gives the common name "coral," and describes it as a wild tree of the hills, 30 or 40 feet (9 to 12 meters) high, 12 to 14 inches (30 to 35 centimeters) in diameter, furnishing a rather strong reddish wood. Used for beams and boards of country houses and frequently turned. It also serves as a dyestuff. (Grosourdy, 2: 377.)

Adiantum fragile.

Sold in the market at Ponce for making an infusion used in diseases of women.

Aegiphila martinicensis. CAPAILLO.

Family Verbenaceae: a smooth shrub growing in thickets in the foothills: flowering in summer and autumn. (Stahl, 6: 222.) Reported from Yabucao. (Sintenis.)

Aeria attenuata. LLUME.

Family Arecaceae: the so-called "llume" palm is a most striking ornament of the rugged limestone hills from Vega Baja to Manati and Arecibo. At a sufficient distance the slender trunk is no longer visible and the crown of leaves appears as if suspended in mid-air, while at closer range it does not seem possible that so slender a shaft can maintain itself. This very slenderness, with the attending flexibility, is, however, an element of strength, since it permits the trees to bend before the wind while the leaves diminish the resistance by straightening out as in the coconut. The hurricane of August, 1899, seemed to have done little damage to these tallest of Porto Rico palms, many of which project for more than half their height above everything standing about them. As the trees of the rather sparse forest growth of these hills are commonly from 12 to 18 meters tall, the llume palms must often attain upwards of 30 meters.

Aeschynomene americana. YERBA ROSARIO.

A leguminous shrubby herb affecting roadsides and dry localities, flowering in autumn. The pod has but 5 to 8 joints, instead of 9 to 15 as in *A. sensitiva*. Our specimens were obtained at Carolina and Cayey.

Aeschynomene portoricensis.

A new species, named by Urban, from the shores of Lake Tortugero near Manati.

Aeschynomene sensitiva. YERBA ROSARIO.

An herb or biennial shrub growing in localities subject to inundation; flowers in August. (Stahl, 3: 37.)

Agati grandiflora. GALLITO.

A small, ornamental, leguminous tree, or large shrub, 3 to 5 meters high, said by Stahl to be a biennial. The large pendulous flowers are either white or pink. At Coamo Springs this was called "baculo." A native of India, cultivated in gardens. (Stahl, 3: 33.)

Agave.

Agaves or century plants are not uncommon in Porto Rico, and are plentiful on the dry hillsides near Guayanilla. What appeared to be several distinct species were observed, but specific determinations are so difficult and uncertain in this group that no attempt was made to classify them.

In the Ponce market pieces of the stems in the form of flat sticks about 10 inches long were on sale for use as razor strops.

Agave sisalana. SISAL HEMP.

Among the suggestions made for new cultures is that of sisal hemp. This is the product of an agave or century plant, and is now largely imported from Mexico. It is, however, improbable that any large area of Porto Rico could be profitably devoted to growing this plant, the culture of which finds a place in land which is too dry and rocky for ordinary crops, and is consequently of so little value that it can be given up to a rather slow-growing plant like the agave. Three or four years are required before the cutting of the leaves and the extraction of the fiber may be begun, and the business can be carried on to the best advantage when estates are large, those of Yucatan being from 500 to 20,000 acres or more in extent. It would be difficult to secure such tracts of suitable land in Porto Rico, especially at prices which would justify attempting the establishment of this industry. In the southwestern part of the island, between Ponce and Yauco, there is some unused rocky land which might produce sisal hemp. However, it is doubtful whether any land suitable even for the rough cultivations sufficing for this plant could not be more profitably employed for other purposes. Most of the island is certainly too valuable to be planted to agaves, and, on the other hand, these will not produce fiber in paying quantities in rich or moist soil.

Ageratum conyzoides. MENTRASTO.

An annual composite herb cultivated as a flower. (Stahl, 5: 110.)

Agitacalillo.

A wood known to Captain Hansard; specific gravity 0.79.

Agrio de guinea. See *Hibiscus sabdariffa*.

Agripalma. See *Leonurus sibiricus*.

Aguacate. See *Persea gratissima*.

Aguacatillo. See *Meliosma obtusifolia* and *M. herbertii*.

A tree from all parts of the island; height, 80 to 90 feet (24 to 27 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood white and soft; specific gravity 0.632; little used. (Exp. 1857.)

Aguardiente.

A general name for spirituous liquors; not so much used in Porto Rico as in some Spanish countries, "ron" (rum) being much more common.

Aguinaldo.

This name was applied at Cataño to a large lavender-flowered ornamental species of *Solanum*, perhaps *S. seaforthianum* (no. 972).

Aguinaldo amarillo. See *Ipomoea umbellata*.

Aguinaldo azul. See *Convolvulus pentanthus*.

Aguinaldo blanco. See *Convolvulus nodiflorus*.

Aguinaldo blanco de costa. See *Convolvulus jamaicensis*.

Aguinaldo peludo. See *Jacquemontia tamnifolia*.

Ahorca caballo. See *Heteropteris bellonis*.

Aiphanes.

A genus of palms recently described from South America, and reduced by Martius to a synonym of *Martinezia*, one species of which, *Martinezia corallina*, comes from Martinique and may prove to be rather closely related to the coyure palm of Porto Rico. On finding that the genus *Martinezia* could not include the Porto Rican species, the use of *Aiphanes* was considered, with a negative conclusion, the reasons for which appear below:

The genus *Aiphanes* was established by Willdenow on *A. aculeata*, a spiny

palm from the mountains about Caracas. The trunk is said to be erect, 30 feet (9 meters) high, subcylindrical, and very spiny. The fronds are about $\frac{5}{8}$ feet long, with four pairs of remote, broad, cuneate, praemorse pinnæ, strongly whitish-pubescent on the under side. The petiole is also beset with spines. Spathe acuminate at both ends, aculeate on the outside, smooth within, opening longitudinally; spadix $1\frac{1}{2}$ feet (45 centimeters) long, composed of cylindrical spikes placed opposite. Flowers hermaphrodite; calyx trifid, its divisions acute; petals acuminate; filaments 6, subulate; anthers rounded; style as long as the stamens; stigma trifid. Drupe globose, the fleshy farinaceous pulp rather tasteless, though edible; nut hard, of the size of a musket ball, unilocular, black, furrowed with a large number of greyish grooves, of which three are always much larger than the others; the kernel is white, very sweet, and very good to eat.

Aiphanes grows in the ravines and forests of the high mountains of the district of Caucagua, province of Caracas, Venezuela, and requires a fertile, somewhat moist soil. It flowers and fruits in July.

From the above it appears that Aiphanes is a genus quite different from that of the spiny palm collected at Vega Baja, and it seems to approach some of the South American forms described under *Bactris* much more closely than it resembles the Porto Rican tree.

Aipim. See *Manihot palmata*.

Ajenjo cimarron. See *Parthenium hysterophorus*.

Aji. See *Capsicum*.

A general name for red peppers, a favorite condiment and vegetable in Spanish countries. Numerous varieties are known and distinguished by common names in Mexico and Cuba, but the subject has not yet received attention in Porto Rico. The large "sweet" peppers are called "aji dulce" in Porto Rico, and belong to the botanical species *Capsicum annuum*, while the small, more pungent varieties belong to *C. frutescens*.

Ají caballero.

Near Toa Alta this name was used for a species of *Capsicum* supposed to be *C. frutescens*. (No. 894.)

Ají dulce. See *Capsicum annuum*.

The Spanish name for the large "sweet" peppers.

Ají picante. See *Capsicum*.

"Pungent peppers" are mostly small varieties of *Capsicum frutescens* or *C. baccatum*.

Ajo. See *Allium sativum*.

Ajonjolí. See *Sesamum orientale*.

Akee. See *Blighia sapida*.

Alambrillo. See *Rajania cordata*.

This is the Cuban name of this plant: the Porto Rican designation is unknown.

Albahaca. See *Ocimum basilicum*.

Also applied at Rio Piedras to a *Scutellaria*, probably *S. pilosa* or *S. coerulescens*.

Albahaca cimarrona. See *Ocimum micranthum* and *O. americanum*.

Albizzia lebbek. ACACIA AMARILLA.

PLATE XV.

A beautiful leguminous shade tree, native in the East Indies, but now extensively introduced in tropical and semitropical countries. It endures drought and thrives in rocky limestone soils, and would thus appear to be well fitted for the south side of Porto Rico, where it already exists in small numbers, as well as at Mayaguez and San Juan.

Alcachofa. See *Cynara scolymus*.



ACACIA AMARILLA ALBIZZIA LEBBEK . NEAR JUANA DIAZ.

Alchornea latifolia.

Family Euphorbiaceae; known from Lares and Peñuelas. (Sintenis.)

Alchorneopsis portoricensis. PALO DE GALLINA.

A euphorbiaceous tree of 10 to 15 meters, known only from the Sierra Luquillo.

Alder, West Indian. See *Conocarpus erecta*.**Alectra brasiliensis.** YERBA DE HIERRO.

Family Scrophulariaceae; a herbaceous annual, growing in ditches and inundated places. (Stahl, 6: 228.)

Alelí. See *Plumeria rubra*.**Alelí amarillo.** See *Plumeria tenorii*.**Alelí cimarron.** See *Plumeria obtusa*, *P. portoricensis*, and *P. alba*.**Aleluya roja de Guinea.** See *Hibiscus sabdariffa*.

The name signifies red sorrel of Guinea.

Alfalfa. See *Medicago sativa*.**Algalia.** See *Abelmoschus abelmoschus*.**Algarrobo.** See *Hymenaea courbaril*.

According to Stahl this name is applied only to *Hymenaea courbaril*, while Hill states that it is also used for *Pithecolobium saman*. The true "algarrobo" or "carob" of southern Europe is *Ceratonia siliqua*, "St. John's bread," the pods of which are valuable as fodder. Captain Hansard gives the specific gravity of "algarrobo" as 1.06, and believes the tree to be indigenous since large trees are found in the mountains in apparently undisturbed forests. He states that it is used especially for wagon wheels, and also believes that it is the same as the "carob" or "gum animae" of the British Islands.

Algarrobo amarillo. See *Hymenaea courbaril*.

This variety is described as a very beautiful tree, large and robust, reaching 90 feet (27 meters) in height and with a straight trunk 24 to 30 feet (7 to 9 meters) long, with a diameter of 6 feet (1.8 meters), or sometimes much more. Martins is said to have seen Algarroba trees in Brazil whose circumference was such that it took 15 Indians with arms extended to encompass one of them. The wood is very hard, yellowish with green veins, breaking vertically. The grain is fine and dense. The wood is much appreciated as well for its strength and durability as for its beauty. Specific gravity, 0.929. It is utilized for all sorts of fine carpenter and cabinet work. It constitutes an important branch of the exports to Europe, where it is much appreciated for fine furniture. In the Antilles it is used commonly in the making of sugar mills, whose cylinders and cogs are made of this material. It is, in fact, one of the best woods of the country and one of the most used. (Grosourdy, 2: 359.)

Reported from all parts of the island; specific gravity, 0.592. (Exp. 1857.)

Algarrobo colorado. See *Hymenaea courbaril*.

According to Grosourdy (2: 360), this is a variety differing from Algarrobo amarillo in having its wood rather light reddish brown, with much darker veins of varying sizes; susceptible also of a high polish. Used for the same purposes as the other.

In the list of woods at the exposition in 1857 this variety is described as a tree 40 to 50 feet (12 to 15 meters) high, from all parts of the island; specific gravity, 0.784.

Algodon. See *Gossypium*.

"Algodon" is the Spanish for cotton, and is applied not only to the genuine cotton but to other related or similar species, as below:

Algodon de seda. See *Catalpa procera*.

Algodon de yuca. See *Gossypium juniphaefolium*.

Algodon rojo. See *Gossypium barbadense*.

Said to be the variety which has been described botanically as *Gossypium purpurascens*.

Algodon yuca. See *Gossypium barbadense*.

Said by Maza to be the variety described by Bello as *Gossypium juniphaefolium*.

Algodoncillo. See *Anguria plumieriana* and *Aselepias curassavica*.

Alilaila.

A tree from all parts of the island; height, 35 to 40 feet (11 or 12 meters); diameter, 5 to 6 inches (12 to 15 centimeters). Wood pinkish, rather hard; specific gravity, 0.803; used in making agricultural implements. (Exp. 1857.) This is probably the same as "lilaila," a name applied to *Melia azedarach*.

Aljonjoli. See *Sesamum orientale*.

Allamanda cathartica. CANARIO.

Family Apocynaceae; an ornamental shrub or small tree introduced from South America. (Stahl, 6: 67.)

Alligator apple. See *Anona palustris*.

Alligator pear. See *Persca gratissima*.

Allium cepa. ONION. CEBOLLA.

Onions are occasionally cultivated in gardens, but the market supply is imported. An American correspondent states that only the Bermuda variety does well. The growing of Bermuda onions is one of the openings suggested for export gardening in Porto Rico and commercial experiments will probably be tried. The seed from Bermuda onions is all grown in Teneriffe.

Allium sativum. GARLIC. AJO.

Allophylus occidentalis. QUIEBRA HACHAS.

An indigenous tree of the family Sapindaceae. (Stahl, 2: 159, as *Schmidelia occidentalis*.)

Alloplectus ambiguus.

Family Gesneriaceae; a new species to replace the *Alloplectus cristatus*, which does not represent the genuine species of that name. Two varieties, *chlorosepalus* and *erythrosepalus*, are also named and reported from numerous localities. (Urban, Symb. 1: 408.)

Alloplectus cristatus. TIBEY DE CRESTA.

Family Gesneriaceae; a rather fleshy herb covered with a reddish pubescence; grows adhering to rocks in the higher mountains, flowering in summer. Urban includes this under his *A. ambiguus*. (Stahl, 6: 261.)

Allspice.

The occurrence of the true allspice, *Pimenta officinalis*, in Porto Rico has not been established, and the reports of Hill and others are probably to be taken as applying to *Amomis caryophyllata*, the tree yielding the oil of bay used in compounding bay rum.

Almacigo. See *Bursera simaruba*.

In Spain this name is applied to the true mastic tree, *Pistacia lentiscus*.

Almacigo blanco.

A tree from all parts of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood light-colored, soft; specific gravity, 0.271; not used. (Exp. 1857.)



ALPINIA NUTANS. AT CAGUAS.

Almacigo rojo.

A tree from all parts of the island; height, 10 to 12 feet (3 to 4 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood pinkish, soft; specific gravity, 0.274; not used. (Exp. 1857.)

Almendra.

The fruit of *Terminalia catappa*.

Almendrillo. See *Prunus occidentalis*.

A tree from all parts of the island; height, 35 to 40 feet (11 to 13 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood light-colored, rather hard; specific gravity, 0.742; not used. (Exp. 1857.)

Almendro. See *Terminalia catappa*.

In Cuba this name is applied to several species of *Haemocharis*, handsome trees of the tea family, Theaceae. In Porto Rico, however, it seems that "almendro" and "almendron" both refer to *Terminalia catappa*, although the account of the collection of Porto Rican woods of the exposition of 1857 gives different data under "almendro" and "almendron."

A tree from all parts of the island; height, 40 to 50 feet (13 to 15 meters); diameter, 20 to 22 inches (50 to 55 centimeters). Wood light-colored, rather hard; specific gravity, 0.644; little used. (Exp. 1857.)

Almendrón. See *Terminalia catappa*.

A tree from the southern part of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood cinnamon-color, hard; specific gravity, 0.922; used in cabinetmaking. (Exp. 1857.)

According to Bello the name "almendron" is also applied to *Prunus occidentalis*.

Almidon.

The Spanish word for starch, in the West Indies commonly made from "cassava" (*Manihot utilissima*).

Almond.

The true almond, *Amygdalus amygdalus*, is closely related to the peach, and does not grow in Porto Rico, but the name "almendron" is applied instead to *Terminalia catappa*, the fruit of which is called "almendra" and is commonly believed to be the genuine article.

Aloe.

This name is sometimes incorrectly applied to the various species of *Agave* or "century plants," which are native of Mexico, South America, and the West Indies. The genus *Aloe* belongs to the Liliaceae and is of African origin. Some of the species have been introduced into cultivation, and *Aloe vera*, in particular is now widely spread throughout the more arid tropical and subtropical regions. As early as 1596 specimens were taken to England from Barbados and the plant was long supposed to be native in that island, receiving the name *Aloe barbadensis*. It is also naturalized, according to Grisebach, in Jamaica, Antigua, and St. Vincent, and very probably in Porto Rico, although not yet reported.

Aloe-wood. See *Cordia sebestena*.**Alpinia nutans.**

PLATE XVI.

Family Zingiberaceae; native in East Indies, but now widely cultivated for the beauty of its flowers; apparently naturalized in Porto Rico. Collected near Bayamon by Sintenis. Our specimen was obtained near Cayey.

Altea. See *Nepsera aquatica*.**Alternanthera repens.** ARDOSNA JARDIN.

Family Amarantaceae; specimens from Cataño (no. 989). The flowers are straw-colored.

Alysicarpus vaginalis. YERBA DE CONTRABANDO.

A small creeping, shrubby leguminous plant with single leaflets and clusters of subcylindrical pods breaking up into single sections, of which the outer surface is irregularly reticulate. Reported from a single station, Boca Vieja, near the mouth of the Bayamon River. Our specimens were obtained in two localities near Ponce. (Stahl, 3: 158.)

Amaranthus paniculatus. PIGWEED. BLEDO BLANCO.

Collected by Sintenis at Guanica.

Amaranthus tristis. PIGWEED.

A common pigweed very similar to our well-known garden pest, *Amaranthus retroflexus*.

Ambrosia. ARTEMISA.

A specimen of *Ambrosia* (no. 966) collected at Cataño appears to be the same as Sintenis no. 196b, from Bayamon, but a specific identification has not been made.

American nutmeg. See *Monodora myristica*.**Ammannia humilis.** YERBA DE CANCER.

Family Lythraceae; this species and *A. latifolia* share the common name and are annual herbs growing in ditches and overflowed places. *A. coccinea*, variety *purpurea*, is reported from Guanica. (Stahl, 4: 128.)

Amor sin zelo.

An ornamental apocynaceous plant, evidently closely related to *Allamanda*, and having similar large yellow flowers, but not yet identified.

Amphilophium paniculatum. LIANA DE CUELLO.

Family Bignoniaceae; a hexagonal vine, scaly-pubescent and much branched; it is found both near the sea and in the mountains; flowers in winter. (Stahl, 6: 182.)

Amomis caryophyllata. BAYBERRY TREE. BAY RUM TREE. WILD CINNAMON. AUSU.

Variouly reported by Bello, Krug, Stahl, and Sintenis as "ausu," "auzu," "guayavita," "limoncillo," "malagueta," and "pimienta malagueta." A myrtaceous tree, from the dried leaves of which is obtained by distillation with water an essential oil, called "bay oil" or "oil of bay," the most important ingredient of bay rum. Only a pint and a half of oil is said to be required for the medication of 100 gallons of rum. The latter should be of good quality and strength. If below 18 or 19 proof it will not properly incorporate the oil. Large quantities of dried leaves of this species are imported from the West Indies, notably from the island of Dominica. They are generally put up into bales of about 200 pounds in weight. It is not known that any leaves have been shipped from Porto Rico, but in 1895, 95 gallons of bay oil, valued at \$1,390, and 12,544 gallons of bay rum, valued at \$6,414, were exported. The trees occur in all parts of the island and are said to be abundant in some districts on the south side. The industry is worthy of investigation as one probably capable of profitable expansion, although not likely to become of more than minor and local importance.

In the fresh condition the leaves of this tree have the taste and odor of lemon, whence the propriety of the name "limoncillo" or "little lemon." Although more common in Porto Rico as a shrub, this species is said to grow to a height of 35 or 40 feet (11 or 12 meters) and to attain a diameter of a foot (30 centimeters) or more; the wood is light-colored, mottled, very hard and heavy. Captain Hansard reports that the tree is cultivated at Maunabo and at other points for the sake of the leaves.

There are probably several other myrtaceous trees which share with this the

name "pimiento," but the present is probably that which Hill and others have supposed to be allspice. (Stahl, 4: 72.)

A tree 45 to 50 feet (14 to 16 meters) high, the straight, rather long trunk 15 to 24 inches (35 to 55 centimeters) in diameter. Furnishes a moderately hard and heavy wood, fine and compact in texture. The sapwood is very light red with darker lines, while the heart is brownish red, brown, or on account of the knots, almost black. It is susceptible of a very high polish. Specific gravity, 0.909. It is one of the best and most valued woods of these countries, very strong and durable, suitable for carpenters and cabinetwork, and it is exported to some extent. The bark is rough and ash-colored, and peels after the manner of the sycamore. (Grosourdy, 2: 398.)

Amyris balsamifera. ROSEWOOD.

A rutaceous shrub or small tree of 2 to 8 meters. Extends from Florida to Cuba, Jamaica, and South America. Known in Porto Rico only from the vicinity of Guanica. (Urban, Add. 3: 292.)

Amyris elemifera. TORCHWOOD. CUABILLA.

A rutaceous shrub or small tree, 2 to 8 meters in height, extending from Florida to Trinidad and known in Porto Rico in littoral thickets near Bayamon and Ponce. Native names are as yet unknown, and the above are from Florida and Cuba.

Amyris maritima. TÉA.

A shrub or small tree from 5 to 10 meters in height, growing in thickets near the sea: Cangrejos, Fajardo, Coamo, Ponce, Guanica, and Salinas de Cabo Rojo. (Urban, Add. 3: 292.)

Amyris silvatica. TÉA.

A wild tree whose height reaches 25 or 30 feet (8 or 9 meters) and the diameter of its trunk 5 to 8 inches (12 to 20 centimeters). Furnishes an aromatic wood, strong, solid, whitish in color, breaking with a vertical fracture.

On account of its resinous quality, splinters of this wood are used by the country people as torches for fishing and to light their huts. It is also suitable for furniture. (Grosourdy, 2: 379.)

Anacagüitas. See *Sterculia carthaginensis*.

Anacahuita.

Probably the same as "anacagüitas."

In Cuba a Mexican boraginaceous tree, *Cordia boissieri*, is known by this name, and the wood, yielding an oil supposed to be a specific for consumption, was formerly exported to Europe from Mexico, but experiments in Germany failed to show the presence of any medicinal virtues.

Flowers called by this name were sold in the market of Ponce for making a tea for colds, etc.

Anacardium occidentale. CASHEW. PAJUIL.

There is also a considerable variety of common names, such as "cajon," "acaju," and "marañón."

This tropical tree gives its name to the family Anacardiaceae, to which our poison ivy and sumac also belong. It is a handsome, quick-growing species, and bears fruit while still very young, sometimes when but two years old. Ultimately it reaches a height of 30 or 40 feet (9 to 13 meters) and has a close-grained, strong, and durable wood, useful for boat building and other purposes, and said to weigh 38 lbs. per cubic foot. It has large, entire oval leaves, not at all resembling those of its relatives mentioned above, but it shares with the poison ivy the possession of an acrid substance strongly irritant to the human epidermis and the mucous membranes. The poisonous material is not, however, spread through the plant, but is mostly concentrated in the rather soft shell of the nut, which is borne upon a pear-shaped, red or yellow fleshy receptacle 2 to 4 inches (5 to 10 centimeters) long.

This receptacle is edible and quite harmless when ripe, having a very agreeable, subacid taste in the raw state, and being also stewed or cooked; a distilled liquor, said to be of good quality, has also been made from it, and in Porto Rico it is used as an ingredient of a peculiar beverage which has been compared to punch. The nut is comparable in size to a rather thick peanut, is kidney-shaped, or distinctly curved near the middle, and contains a single large kernel of white, firm flesh of fine texture and a delicate, very pleasant, nutty flavor. No attempt should, however, be made to eat it in the raw state, on account of the poisonous juice of the shell, which must be driven off by the heat, so that roasting is an absolute necessity. In the fresh state the kernel also contains a slight amount of the irritant oil and has painful effects upon the lips and tongue.

The process of roasting the nuts is quite simple, but skill and experience are required to avoid burning on the one hand and insufficient heating on the other. The fumes driven off during the operation are also poisonous, and where cashew nuts are roasted in large quantities for export the workers suffer from blistered faces and hands, accompanied by painful swellings and other toxic effects of inhaling and coming in contact with the poisonous vapors. These difficulties might be obviated, it would seem, by a proper arrangement of the roasting ovens and by providing a system of drafts for carrying up the fumes.

After the roasting is completed the nuts are shelled and sorted and packed in sealed tin cans for shipment. In this way they keep fresh for a considerable period without the necessity of any other preliminary treatment or of the addition of preservatives. The trade in cashew nuts is still very small, but may be expected to increase. Mr. David G. Fairchild, agricultural explorer of the United States Department of Agriculture, became acquainted with the cashew in Jamaica and took a very enthusiastic view of its value and commercial possibilities, as the following extract from a letter will show:

"I assure you that there is no nut so delicious as the roasted cashew. Burnt almonds are flat in comparison. Huyler could, I believe, make a fine thing of it, and a big market could be created. Through William Fawcett, director of the gardens here, you can secure a large quantity of seeds for planting, I believe. The sooner groves are started the sooner the nuts can be pushed on the market. For Porto Rico and Hawaii, I believe, it would be a good thing. This fruit is bound to go. Porto Rico, Cuba, and Hawaii should be encouraged and steps be taken to secure a collection of the best varieties. Look out for Jamaica as a competitor. Mr. S. T. Scharschmidt, of Mandeville, Jamaica, has made an oil from the fruit, superior to olive oil for table use. There are wonderful prospects for such a fruit. It can be shipped to Boston from here easily without cold storage."

From a sample sent home the roasted nuts were pronounced excellent by all who tasted them. Probably they are better when fresh, though no tendency to become rancid was apparent. In British India the cashew has now been introduced very generally, and has found a much larger variety of uses than in its native home in the West Indies and South America. East Indian information has been compiled from numerous sources in Watt's *Economic Products of India*, and is, with the omission of some repetitions, as follows:

"The bark of this plant yields a gum which occurs in large stalactitic pieces; it is yellow or reddish, and only slightly soluble in water. It is obnoxious to insects. The astringent juice issuing from incisions in the bark is in demand as an ink, and is used by native workmen as a flux for soldering metals. The bark may be used for tanning. The pericarp gives an oil called cardol, which is very astringent, and is used by the Andamanese to tan or color fishing nets, so as to preserve them. Dr. Dymock informs me that this oil is called dik in Goa, where it is much used as a tar for boats and nets.

"From this plant two distinct oils are obtained. (1) The kernels when pressed yield a light-yellow bland oil, very nutritious, the finest quality in every

respect, equal to almond oil and considered superior to olive oil. The yield is about 40 per cent. The kernels are so extensively eaten in India, however, that it is almost impossible that a trade could at present be done in this oil. Samples of this fixed oil, and information as to methods of preparation and extent of trade, are much required. The kernels have been once or twice exported to Europe under the name of 'cassia nuts.'

"(2) *Cardole or Cashew-apple oil*.—This is prepared from the pericarp or shell of the nut. It is black, acrid, and powerfully vesicating. In the Andamans it is used to color and preserve fishing lines. It is an effective preventive against white ants in carved woodwork, books, etc. The yield is 29½ per cent.

"The medicinal uses of this plant are many. The acrid oil is used as an anaesthetic in leprosy, and as a blister in warts, corns, and ulcers. Between the laminae of the shell of the kernel there is a black caustic fluid, which contains an acrid oily principle, cardol, and a peculiar acid, anacardic acid. It possesses powerful rubefacient and vesicant properties. The spirit distilled from the expressed juices of the fruit may be used as a stimulant. Fruit eaten as a remedy for scurvy. The juice of the nut is used as a substitute for iodine locally. The oil I have used with benefit in the anaesthetic variety of leprosy. The oil obtained from the shell by maceration in spirit is the very best application for cracks of the feet so common with natives. The oil is efficacious, when faintly brushed, as a local stimulant in psoriasis."

Along the Adjuntas road we noticed trees of *Anacardium* from which the bark of the trunk had been largely hacked off. On inquiring it was learned that a decoction made of it was held in high esteem as an astringent remedy for diarrhea.

In the bulletins of the botanic gardens of Jamaica and Trinidad this tree has also received attention, and some additional notes from these sources are appended:

"A gum is obtained from the trunk. It is subastringent and highly unpalatable to insects. It consists principally of arabine and dextrine, both soluble in water, with a minor insoluble portion, probably bassarine. It forms a strong, yellowish mucilage with water. In South America it is largely used by bookbinders; it is occasionally imported from that country into England and possesses the same commercial value as the common and inferior sorts of Arabic and Senegal gums.

"The fruit of *Anacardium* is well known to West Indians, who, besides eating it in the fresh state, make conserves of it in various ways. Though sweet, it is at the same time very astringent and said to be useful in cases of dysentery and diarrhea. Many and various are the effects with which this tree, its fruit, bark, leaves, and seeds are credited, and if all were true, it would indeed be one of the wonders of nature. It is said to possess aphrodisiacal properties, the leaves to be capable of producing drunkenness, the nuts or seeds when roasted to excite the faculties, especially memory, so much so that a confection made therefrom has been called 'confection des sages,' and the oil from the nut is said to be equal to that made from the finest olives, while an acrid oil is produced from the epidermis of the nut, which is said by Barham in *Hortus Americanus*, 1794, to 'cure herpes and cancerous and malignant ulcers abounding in rotten flesh; it also kills worms in ulcers and chigoes; it takes away freckles and liver spots, but it draws blisters and therefore must be cautiously made use of. It has been observed that poor dropsical slaves that have had the liberty to go into a cashew walk and eat what cashews they pleased, and of the roasted nuts, have been recovered.'"

Another writer brings an indictment against the West Indian fair sex for using it as a cosmetic when they have become tanned by exposure to the sun. He describes the process as follows:

"They take a nut, scrape off the outside skin, and rub their faces with the

exposed oily surface. The face swells and blackens, but ultimately the tan skin peels off; and although the process necessitates rigid retirement for a fortnight, at the end of that time they emerge with a new skin and complexion as fair as a babe. It is also to be stated that the oil is found to be useful in leprosy, and that the fruit is a cure for the disorder which causes the patient to become what is known as a dirt-eater.

"The tree produces a beautiful clear gum, which makes a fine varnish, and is said to possess a special virtue in preserving woods from insects. The sap of the tree, like its congener, *Semecarpus auacardium*, produces an indelible stain on linen. Professor Lindley makes Martius responsible for a statement that the nut has a wonderful effect upon chronic inflammations of the eyes, especially such as are of scrofulous nature, when simply borne or carried by the persons affected. It is also a common belief in Trinidad that a necklace or collar of cashew nuts has a remarkable curative effect upon some of the diseases of the canine race, especially for coughs or distemper."

The wood is very hard and strong. It is much appreciated for making the hubs of carriage wheels, yokes, and other farm utensils. Its principal use in Porto Rico is for charcoal and fuel; specific gravity, 0.488. (Grosourdy, 2:404.)

Aname.

In the market of Ponce large nut-like seeds said to be derived from a tree were on sale. They are ground up and put into rum for the sake of the rather pleasant pungent flavor, and are also supposed to have medicinal properties.

Anamu. See *Paromia typhalea*.

In the market of Ponce material with this common name was on sale as a medicine. It is boiled in water and the decoction used for the relief of pain in the stomach.

Ananassa ananas. PINEAPPLE. PIÑA.

The pineapple is the one fruit in the production of which Porto Rico already has a good reputation, one of the largest and best varieties being known as the "Porto Rican." There can be little doubt that natural conditions, entirely favorable to this culture, exist in many parts of the island, but it is not on this account to be inferred that our markets will be flooded at once with cheap fruit from Porto Rico. In order to furnish a regular commercial supply it will first be necessary to bring into existence an organized productive industry. The wild and half-wild pineapples brought in by the rural natives and sold in the town markets would not go far in supplying an export trade, and while many individual high-grade fruits are secured the quality is, of course, extremely uneven and areas of one variety have seldom been planted. An initial difficulty, as in Florida, will probably arise in the scarcity of cuttings of good varieties. In Florida this is by far the largest item of expense in new plantings, even in comparison with the frost sheds which cost between \$300 and \$500 per acre. But cuttings of good varieties bring from 10 to 20 cents apiece, and although this means between \$2,000 and \$3,000 per acre for plants, the grower has his returns in disposing of his own suckers at similar prices. Of course it is not to be expected that this rate can be indefinitely maintained. Such prices are possible only while the industry is rapidly expanding, and as soon as the normal demands of the market are once filled there will come an inevitable and probably a very rapid decline in the valuation of cuttings.

Florida pineapple growers have found, however, that the partial shade of the frost sheds exerts a most beneficial influence upon the size and quality of the fruit, and the use of sheds for shade purposes is now being extended far below the frost line. The inference is natural that the production of high-grade fruits in Porto Rico will require the introduction of the Florida methods of culture,

especially since the small importations which have already been made from Porto Rico were badly damaged by sun scald.

It is not impossible, however, that situations might be found in Porto Rico where the injury from this cause would be very slight, and there is a further probability that if shade proves to be necessary it can be supplied more cheaply by means of some of the many leguminous shade trees, which will also assist in maintaining the fertility of the land. We did not notice any plantings which seemed to have been made with reference to shade, although it is well known in the Tropics that partial protection from the sun encourages the growth of large pineapple fruits.

Serious competition on the part of Porto Rico with the well-developed pineapple industry of Florida is not to be expected for a series of years, though the very favorable natural conditions render it certain that an increasing quantity of this fruit will be sent to the American market. The Florida supply does not, however, meet the popular demand, even at prices far in advance of the actual cost of raising and marketing the fruit, and at more reasonable rates there is little doubt that the present consumption would be quickly doubled or trebled. There is, accordingly, ample room for the expansion of pineapple growing as an agricultural industry both in Florida and in Porto Rico.

Local conditions and varieties.—The pineapple season at Ponce is from April to July, after which time none are obtainable. Small-sized fruits of fine flavor and texture are preferred, and command a higher price than the large ones for which the island is particularly noted. On the north side of the island pineapples are said to be obtainable from April to September. Those at higher altitudes ripen first, while lower down the crop is not ready for market until July, though the fruits are of better quality.

Sandy and gravelly loams are the preferable soils for fruit production in Porto Rico, although the plants apparently thrive in all situations. Limestone soils not subject to too great drought are also desirable. To facilitate clean culture the plants are set 3 feet (90 centimeters) apart in rows 5 feet (150 centimeters) apart. The same piece of ground remains productive from four to eight years, dependent upon soil and culture. For shipment the fruits are cut before the commencement of the ripening process, and it is desirable to make a clean cut and leave an inch (25 millimeters) or more of the stem to avoid decay and prevent drying out. With proper packing and ventilation pineapples will not suffer at all from the brief sea voyage, and the shipping rates will probably be as low as or lower than from Florida. The transportation difficulties arising on shore will not, however, be so easy of adjustment in the near future, except in the immediate vicinity of the ports, and will undoubtedly retard the development of the pineapple business.

The custom so prevalent in Porto Rico of placing the sliced fruit in salt water before eating is evidently very old, as Acosta says of the inhabitants of the West Indies, "They eat it being cut in morsels and steeped a while in water and salt."

There is also a belief, more or less widespread, in Porto Rico that pineapples and coffee should not be partaken of at the same meal.

An insect enemy.—On both sides of the island young pineapples were found to be infested by a mealy-bug, which is said to prevent normal growth and finally to rot the fruit. The same insects are also to be found in great numbers on the Bromelia "piñuelas," or wild pines, a related plant found everywhere in Porto Rico. To be rid of the mealy-bugs it may be found necessary to destroy all the Bromelias in the neighborhood of pineapple farms. It is not likely that the ants would carry the mealy-bugs any great distance, though a separation of several rods would probably be required. An experiment with kerosene emulsion for driving away the insects which had already settled was reported as quite successful.

Andira inermis. CABBAGE TREE. MOCA.

A leguminous tree called "cabbage tree" or "cabbage bark tree" in Jamaica on account of its disagreeable odor. It is generally distributed in Porto Rico, and is sometimes used for coffee shade, but is considered inferior to the two species of *Inga* on account of its slow growth. The bark, in the form of a powder or a decoction, is said in Jamaica to have value as a cathartic and vermifuge, but large doses are dangerous, causing vomiting, delirium, and even death. The wood is said to be hard and durable, having a specific gravity of 0.88, and is susceptible of a high polish. The pods are fleshy, about the size of a horse chestnut, and contain but a single seed. The floors of the caves of Aguas Buenas, Porto Rico, are in places covered with the seeds of this species, which are carried in by bats for the sake of the inclosing pulp. These seeds germinate in the caves, sending up slender white sprouts 2 or 3 feet (60 to 90 centimeters) high.

Grosourdy (2: 395) reports *A. racemosa* in Porto Rico with the same common name. As subsequent writers report only one species, the determination is probably in error. He says of the wood: "It is rather hard, and its color varies greatly, specimens varying from yellow to reddish flesh-color, brown, or black having been seen. For this reason two varieties are called 'moca colorada' and 'moca amarilla.' Both are, however, produced in the same tree. The wood is also variously marked and sometimes produces a very pleasing effect. Specific gravity, 0.748. It is used to considerable advantage in making the hubs of wheels. Made into boards it is suitable for flooring and all sorts of carpenter work. It was formerly used in Brazil in the construction of boats. In Porto Rico its most common use is for the framework of country houses. It is imported into Europe and used to make canes and umbrella and parasol handles, and for turned parts of cabinet work."

Anemia.

In the woods through which the road winds to the west of the Guajataca River, near Quebradillas, we found *Anemia adiantifolia* and the little species usually called *A. aurita*. In the latter the fertile frond is fertile throughout, and it should be associated with *Coptophyllum*, although that genus was established by Gardner with the obviously mistaken idea that in the typical *Anemia* two fertile fronds are united with a sterile, a theory of fasciation violently improbable on morphological grounds. All the requirements of the phenomena are much better met by observing the simple fact that in *Anemia* proper only the lowest pair of pinnules is fertile, while in *Coptophyllum auritum* and its congeners we have the ordinary case of completely dimorphic fronds. That the fertile frond of *Anemia* should have the fibro-vascular system modified in accordance with the requirements of its double function may be viewed as an adaptation and is not necessarily an anomaly.

Angela. See *Moringa moringa*.

Angelica tree. See *Sciadophyllum brownii*.

Angelón. See *Angelonia angustifolia*.

Angelonia angustifolia. ANGELÓN.

Family Scrophulariaceae: an annual herb, cultivated in gardens as an ornamental. The same may be said of *A. salicariifolia*. (Stahl, 6: 234, 235.)

Anguria plumieriana. ALGODONCILLO.

Family Cucurbitaceae: a pubescent vine, rare on the north side of the island and confined to mountainous districts made inaccessible by rain during the winter season, the time of flowering. (Stahl, 4: 176.)

Aniba bracteata.

Family Lauraceae; has oblong leaves 12 to 18 inches (30 to 45 centimeters) long; known from Yabucoa. (Sintenis.)

Añil. See *Indigofera anil*.

This is the Spanish for indigo and is applied also to other indigo yielding species.

Añil cenizo. See *Cracca cinerea*.

Añil falso. See *Cracca aniloides*.

Añil racimillo. See *Cracca leptostachya* and *C. purpurea*.

Anistus arborescens. See *Acnistus arborescens*.

Anoda hastata. VIOLETA.

Family Malvaceae; an annual herb native in Porto Rico, but also introduced from the continent. (Stahl, 2: 59.)

Anón. See *Anona squamosa* and *A. dolabripetala*.

Anona dolabripetala. ANÓN.

Reported from Porto Rico by Bello, but not known to Stahl, who seems to consider it somewhat doubtful. Bello's locality was Furnias, according to Stahl, but this is not so stated in Bello's paper. This species is also given as a synonym of *Rollinia longifolia*.

Anona montana. GUANÁBANO CIMARRON.

An indigenous shrub or tree of mountain districts; not common. The wood is fibrous and of poor quality. The fruit is globose with soft spines from the middle of round or polygonal areas. It is not edible. (Stahl, 2: 24.)

Anona muricata. SOURSOP. GUANÁBANO.

Family Anonaceae; a small tree. The fruit called the "soursop" is well known throughout the tropics. It has a rough green skin, inclosing a soft and juicy white pulp of an agreeable, slightly acid flavor, in which are embedded the large brown seeds. In the mature condition the weight may reach 2 pounds or more.

Grosourdy (2: 387) says there are several varieties. The wood which is not used is soft and light-colored, and breaks with a horizontal fracture. Specific gravity, 0.397.

An infusion of the leaves is said to be a remedy for dysentery.

Anona palustris. ALLIGATOR APPLE. CAYUR.

An indigenous shrub of 2 to 5 meters, affecting swampy localities. According to Dr. Stahl this species has astringent properties. The fruit is not edible. Bello calls this "corazon cimarron" as well as "cayurz." (Stahl, 2: 25.)

A small tree 24 to 30 feet (8 to 10 meters) high, the trunk reaching 10 or 12 inches (25 to 30 centimeters) in diameter. The wood is very light and soft, fibrous in texture, gray or light brown somewhat tinged with green, lustrous and sometimes somewhat marbled, but with long exposure to the air it turns a dirty white. Specific gravity of the roots, 0.175. Used in the country for rafts. The wood from roots that have grown in bogs is extremely soft and porous, and is used in Cuba to sharpen razors.

Called also "guanábano cimarron" and "corcho." (Grosourdy, 2: 364.)

Anona reticulata. BULLOCK'S HEART. CORAZON. CUSTARD APPLE.

A wild tree not very abundant, 25 to 30 feet (8 to 10 meters) high, rather dense. The trunk is ordinarily straight, and 10 to 14 inches (25 to 35 centimeters) in diameter. Furnishes a light wood, rather soft, but resistant. Fibrous in texture, moderately close-grained. The wood is handsome in appearance but is not used. (Grosourdy, 2: 377.) These dimensions are certainly much larger than any seen in the island.

The fruit is of a more regular oval shape than that of the soursop, and the skin is much more nearly smooth, being marked merely with small slightly raised areas. The flesh is yellow, and is generally considered inferior to that of the soursop.

Anona squamosa. SWEETSOP. ANÓN. CHERIMOLIA.

A low tree or shrub widely cultivated throughout the tropics, though not considered the equal of the soursop. According to Bello, the name "chirimoya" is applied to this species in the western part of the island. (Stahl, 2: 26.)

Anthacanthus spinosus. ESPINOSA.

Family Acanthaceae; a shrub growing in thickets on the north coast, near Cangujos, and also known from Humacao. (Stahl, 6: 246.)

Antherylium rhorii. ROSA DE CIENEGA.

Family Lythraceae; a much branched tree, 3 meters high, found on the south coast; also reported from the island of St. Thomas. (Stahl, 4: 123.)

Anthurium acaule.

PLATE XVII.

Family Araceae; a herbaceous plant of no economic value sent in by all collectors.

Antigonon leptopus.

PLATE XVIII.

An ornamental climbing vine belonging to the family Polygonaceae. The clusters of showy pink flowers sometimes cover large areas and suggest the appearance of Bougainvillea, for which Antigonon is probably often mistaken. The beautiful pink color of the latter is, however, very different and much more beautiful than the rather unpleasantly conspicuous purplish tint of Bougainvillea. Antigonon is probably also superior as a rapid grower and a more effective climber. The flower clusters also serve as tendrils and shorten themselves or "take in the slack" by bending at the joints in a zigzag form. The specimen photographed was taken at Coamo Springs, but the plant was seen at numerous places and has been collected by Sintenis at Fajardo. It is supposed to be a native of western Mexico.

In Cuba the species is called "coralillo" or "coralillo rosado," but no common name has been recorded from Porto Rico. Another Mexican species, *A. cinerascens* is reported as introduced into St. Thomas.

Antirrhoea.

A genus of rubiaceous shrubs. Several species have been collected by Sintenis in different parts of Porto Rico.

Antirrhoea acutata.

From Guanica. A tree 5 to 8 meters in height from the southwestern part of the island; Ponce, Guanica, Cabo Rojo. (Urban, Symb. 1: 439.)

Antirrhoea coriacea. QUINA.

Also called "boje." A rubiaceous tree attaining 20 meters in height; known in the vicinity of Utuado and from several other localities. It has been placed by Stahl and other writers under *Stenostomum*, *Exostemma*, and other genera. (Urban, Symb. 1: 436.)

Antirrhoea lucida. PALO LLORÓN.

Family Rubiaceae; a shrub of 3 meters; locality not known. (Stahl, 5: 59) as *Stenostomum lucidum*.

Antirrhoea obtusifolia. TORTUGUILLO.

A rubiaceous tree 8 to 15 meters in height, recently described as a new species from the mountains of Luquillo and Yabucoa. (Urban, Symb. 1: 435.)

Antirrhoea resinosa. AQUILON.

A tree 10 meters high; the branches covered with a sticky powder. Found in mountain districts; wood employed in building. (Stahl, 5: 58, as *Stenostomum resinosum*.)

Antirrhoea sintenisii.

Another new species from the vicinity of Manati. (Urban, Symb. 1: 438.)



ANTHURIUM ACAULE.



ANTIGONON LEPTOPUS. COAMO SPRINGS.

Aquilon. See *Langeria resinosa*.

According to Captain Hansard the specific gravity of wood of "aquilon" is 0.88.

Arachis hypogea. PEANUT. MANÍ.

Porto Rican peanuts are of very small size, and not many are grown. They are hawked upon the streets by boys, much as with us.

Aralia arborea. See *Gilibertia arborea*.

Araucaria excelsa. NORFOLK ISLAND PINE.

This beautiful coniferous tree is a conspicuous object in gardens and parks in the Canary Islands, but is either very scarce or entirely absent in Porto Rico. That it would flourish if introduced is indicated by the fact that it has been found to succeed well in Trinidad.

Árbol de la cera. See *Sapium sebiferum*.

Árbol del pan. See *Artocarpus communis*.

Árbol del sebo. See *Sapium sebiferum*.

Ardisia."

A large genus of shrubs, or small trees, of the family Myrsinaceae. Many of them are fine ornamentals, and a considerable number of species appear in horticultural trade. Several of the Porto Rican species are little known, as yet, and should be canvassed with reference to their desirability for introduction for the above purposes.

Ardisia coriacea. MAMEYUELO.

A tree 3 to 4 meters high found in the mountains and in waste places. Also reported from Panama and the island of St. Thomas. (Stahl, 6: 38.)

Ardisia crenulata. See *Parathesis crenulata*.

Ardisia glauciflora. MAMEYUELO.

A tree of 5 to 8 meters, with leaves 8 or 9 inches (20 to 23 centimeters) long, leathery and finely veined. Like *A. sintenisii* and *A. yunquensis* (shrubs of 3 to 4 meters) known only from the Luquillo Mountains. (Urban, Symb. 1: 382.)

Ardisia guadalupensis.

Collected by Sintenis about Coamo and near Rincon.

Ardisia pendula. JACANILLO.

A tree of 5 to 10 meters from the vicinity of Cayey (Aibonito, Lares, and Manati).

Ardisia purpurascens.

Native names: "Quebra hacha," "mameyulo," and "hacanilla." A shrub, or small tree, 10 meters or less in height. A variety *corymbifera* is known from Utuado and Yabucoa; a variety *genuina* from Barranquitas.

Ardisia sintenisii.

Known only from the Sierra de Luquillo.

Ardosna jardin. See *Alternanthera repens*.

Areca catechu. BETEL PALM.

Family Arecaceae: in the western end of the island the betel palm of the Malay region has been sparingly introduced. A few were seen in gardens about Mayaguez and others in and near San Sebastian. So far as we were able to learn, the people do not know the name or nature of this introduced species, which is apparently planted only as an ornament or a curiosity. The form is not unpleasing, but the extremely deep, somber green of the foliage seems almost unnatural and imparts a suggestion of artificiality.

"The generic name *Ardisia* (Swartz, 1788) is antedated by *Leacorea* Aubl. (1775), but as few of the species have been transferred to this genus, they are here retained under their commonly accepted designations.

Arenga saccharifera. GOMATU.

An East India palm cultivated in Jamaica, but not known to have been introduced into Porto Rico. Called "cabo negro" in the Philippines.

Argemone mexicana. CARDOSANTO.

Family Papaveraceae; an herb, native of Mexico. The seeds possess acrid, narcotic, and purgative properties, and are employed as a substitute for ipecacuanha. The yellow juice of the plant is said to be used in ophthalmia. (Stahl, 2: 33.)

Argithamnia candicans.

Family Euphorbiaceae; from Guanica.

Argithamnia fasciculata.

From Guanica (Sintenis).

Aristolochia oblongata.

Family Aristolochiaceae; this and other species of *Aristolochia* are twining vines, with very peculiar pipe-shaped flowers. There are numerous species in the West Indies and probably several will be found in Porto Rico. "Contrayerva" is given as local name for this genus by Grisebach.

Arnotta. See *Bixa orellana*.

A name used in some of the British Islands. Anatto is the preferable form.

Aroma. See *Lasianthus lanceolatus*.**Aromo.** See *Acacia farnesiana*.

At Coamo Springs this name was used for *Cleome spinosa*, but near Ponce the more correct application was made.

Arrabidaea chica.

A bignoniaceous shrub, with large, flat, leathery pods; the variety *thyrsoides* has been collected on calcareous mountains near Bayamon.

Arracacha. See *Arracacia esculenta*.**Arracacia esculenta.** ARRACACHA.

An umbelliferous garden vegetable presumably introduced from Venezuela by the Spanish refugees, who also brought the culture of cacao. The "arracacha" seems to be grown somewhat extensively in the mountains behind Ponce, where large quantities appeared in the market. It requires a very equable climate, and will not endure extremes either of heat or cold. As its botanical affinity might indicate, the flavor suggests that of the carrot or parsnip, but it is said to be superior to either, and some Europeans become very fond of it.

In Venezuela and Colombia the culture of "arracacha" has been long an industry of considerable importance, and attempts have been made to introduce it into Europe, but without success. The following account of the Venezuelan culture is by Mr. David G. Fairchild, agricultural explorer of this Department, who visited Venezuela in 1898.

"The culture of the native 'apio' in Venezuela is an extensive one. There is no question that it competes with the potato for a place as the most important food plant among the peones of the interior elevated regions. It is also highly esteemed by the Europeans living in Venezuela, and is met with on the hotel tables as a constituent of certain vegetable soups, resembling somewhat carrots and passing as such among the less observant. I consider it very much superior to carrots. In general it is served boiled, precisely as parsnips are treated. So-called *buñuelos* or fritters are prepared from the grated roots by frying them in lard. Cut in thin slices and fried it is delicious, and as a soufflé it is said to be unexcelled.

"The culture of this vegetable is confined to the higher altitudes, where the mean annual temperature is 58 to 60 degrees F., the 'tierra fria' of the Venezuelans, and carried on on a small scale by the peones, small cultivators and

Italians, which latter, by the way gave the name 'apio' (celery) to the plant because of its superficial resemblance to the Italian apio.

"The plant thrives best in good rich soil, though growing successfully in barren sandy soil. It resembles superficially celery. The plant is propagated from root cuttings almost exclusively, the enlarged leaf bases forming small 'heads,' which are cut from the edible roots and planted, one in each hill, just as onions are planted. In the course of eleven months a mass of edible roots the size of a man's head has been formed, and a dozen or more of the 'heads,' which can be used for planting the following season, after being removed from the roots and allowed partially to dry. How long these heads can be kept alive without planting is a question I have not been able to answer. According to Jose A. Diaz, El Agricultor, Venezolano o' Lecciones de Agricultura Practica Nacionae, J. I. Caracas, 1877, pages 67-71. Señor Vargas has published^a an article on the culture of arracacha. In this article, which is quoted at length by Diaz, the statement is made that the plant originated in the provinces of Santa Fe and Caracas; and that now its culture in Colombia is as common as that of the potato in England. It matures in three or four months sufficiently to be of value in the kitchen, although a longer period allows much larger roots to develop. The best variety comes from Lipacon, near Norte de Santa Fe de Bogota, in Colombia. Specimens of the plant were imported into Europe in 1824, and in Liverpool a specimen flowered. These plants were, however, imported from Trinidad, and not from Venezuela. It is probable that those from the high regions of Colombia and Venezuela will succeed better than those from the warm regions of Trinidad. Humboldt and Bonpland mentioned having found in New Granada, near Teindala, province of Pasto, at 700 meters, a wild plant called 'sacharacacha' (a name similar to aracacha). This is possibly a variety of the Venezuelan plant. For trial in America the sprouts should be planted as soon as possible in the spring and given a long season to develop in. They should be tested in all the southern experiment stations, and may be even worthy of trial in the northern ones. The question of keeping the sprouts over winter may be a difficult one to answer, as the plant rarely sets seed."

Arrayan. See *Myrica cerifera*, also *Myrsine coriacea* and *M. floribunda*.

In the market of Ponce leaves of "arrayan" were on sale as a remedy for rheumatism. The application is made in the form of a tincture with rum.

Arrowhead. See *Sagittaria acutifolia*.

Arrowroot. See *Maranta arundinacea*.

Artemisa. See *Ambrosia*.

Artocarpus communis. BREADFRUIT. ARBOL DE PAN.

Family Moraceae; in Porto Rico the breadfruit is commonly called "castaña," the Spanish name for the chestnut. This doubtless results from the fact that the seeds are the part generally used, and these bear considerable resemblance to large Spanish chestnuts. They are frequently sold in the market in a germinating condition (see photograph) and are boiled for a few moments before eating. Seedless varieties of the breadfruit also exist, but it was not ascertained that such have been introduced into Porto Rico. In some regions the name breadfruit is restricted to the large seedless form, while those which produce seed are called "bread nut." This name is also applied to other species of *Artocarpus*. One of these is the "jack fruit," which has been introduced from the East Indies into Jamaica. It bears an oval fruit, 18 inches (45 centimeters) or more long, weighing 30 or 40 pounds (13 to 18 kilograms). If this is not already known in Porto Rico steps should be taken to secure it.

^aAnnals of Botany (Sims & König), Vol. I, p. 400.

Furnishes a wood yellowish gray in color; rather light and soft, but strong, resistant, and elastic. It is highly appreciated for furniture and for building houses. Specific gravity, 0.495. Also called "palo de pan." (Grosourdy, 2: 406.)

The breadfruit grows about Ponce only in valleys and moist places, and is far less common than on the north side of the island. Along the Rio Portugas, near the baths of Quinitana, several trees were noticed which had had their trunks and larger branches for 20 feet (6 meters) up and more severely hacked, doubtless to induce fertility.

A substance somewhat resembling rubber may be made by boiling the milky mucilaginous juice of the breadfruit tree with cocoanut oil. The resulting material is said to be tough, durable, and waterproof, and is called "canoe gum," because used to close the seams of canoes and wooden utensils. Before hardening, it also serves as birdlime.

Artocarpus integrifolia. JACK FRUIT.

"I am strongly in favor of the jack as the tree best suited for providing shade for fields of coffee. In the first place, its presence, so far from being prejudicial, seems to be actually beneficial to the coffee plant; next, it is a subsoil feeder; then it produces a fruit much valued by the natives; its timber is also valuable, whether for cabinetmaking or building purposes; and, finally, it flourishes best precisely in those conditions where its shade is most required. Known to botanists as the *Artocarpus integrifolia*, the Jack grows to a large size; it resembles and belongs to the same family as the bread fruit tree. The timber, when newly cut, is of a light yellow color, possesses a beautiful grain, and is capable of a high polish, not greatly inferior to that of the mahogany or satinwood, both of which it resembles to a certain extent in grain and color after having been polished. The fruit is as large as a pumpkin and weighs from 20 to 30 pounds (9 to 13 kilograms), containing from 200 to 300 seeds, which, though somewhat unpleasant in smell when raw, are converted by being roasted or boiled into a wholesome farinaceous food, always much appreciated by the coolies.

"The jack is said to bear transplanting badly, and it will therefore be necessary to deposit two or three of the seeds a couple of inches below the surface wherever a tree is desired to grow. The most healthy of the plants can afterwards be selected."—Hull.

Asclepias curassavica. ALGODONCILLO. BLOODFLOWER.

Also called "platanillo." An indigenous milkweed appearing in Urban's Symbolæ as a variety of *A. nivea*. In some of the West Indies it is used as an emetic in substitution for the true ipecac, which is derived from a South American plant of the family Rubiaceæ. A new variety (*concolor*) has recently been described by Krug and Urban, distinguished by having the corolla yellow. (Urban, Symb. 1: 389.)

Asclepias nivea. PLATANILLO BLANCO.

An indigenous milkweed 1½ to 3 feet (0.5 to 1 meter) in height.

Asimina blainii.

An anonaceous shrub or tree from Utuado.

Assaea fascicularis. See *Henriettella fascicularis*.

Astromeda. See *Lagerstroemia indica*.

Atelandra. See *Meliosma*.

An indigenous genus of two species doubtfully referred by its author, Bello, to the Myrsinaceæ. The name is preoccupied in the Labiatae and would need to be changed. Urban refers the species to *Meliosma*.

Atelandra laurina. See *Meliosma herbertii*.

Atelandra obtusifolia. See *Meliosma obtusifolia*.

Atriplex cristata.

Family Chenopodiaceae; a native saltbush; known from the vicinity of Guanica.

Attalea.

Sintenis collected about houses near Victoria, Agnadilla, leaves and young fruit of a palm determined by Drude as belonging to the subtribe Attaleeae. *Attalea cohune*, native in Honduras, is cultivated in Jamaica. The seeds are said to be extremely hard and are turned into buttons and other small articles like those of the African wine palm *Raphia*.

Ausu. See *Amomis caryophyllata*.

The name is also applied to *Myrcia pagani* and probably to other Myrtaceae.

A tree from the northeast part of the island; height, 100 to 110 feet; diameter, 10 to 12 inches. Wood purplish, very hard; specific gravity 1.203; used in cabinet-making. (Exp. 1857.)

Grosourdy (2: 363) describes this as 35 to 40 feet (11 to 13 meters) high with a trunk 10 to 12 inches in diameter. The wood is very strong, the heart pale, the exterior portions white. Breaks with a vertical fracture, and is used for shelving.

Ausubo.

Stahl gives this name for *Bumelia nigra*, which in the Index Kewensis stands as a synonym of *Dipholis salicifolia*. According to Hill and others it is used for *Sideroxylon mastichodendron*, while Bello says *Sapota sideroxylon*, a synonym of *Achras sapota*.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 50 to 55 inches (1.2 to 1.3 meters). Wood red, very hard; specific gravity, 1.158; used in housebuilding. (Exp. 1857.)

The tree known to Captain Hansard as "ausubo" has leaves not unlike those of *Ficus elastica*, but somewhat smaller and thinner; when mature they turn red. The fruit somewhat resembles the sapodilla.

"The wood is clear, dull reddish-brown; moderately conspicuous ducts, in short, detached, long and short chains (single line of cells) evenly diffused; chains usually between two medullary rays; medullary rays very numerous, minute, indistinct; wood fibers slightly interlaced and appearing straight-grained; resembles somewhat a fine-grained teak. It is the chief and the most used timber on the island, being noted for its great durability. It is used in the making of wagon wheels, which are turned out by machinery in Ponce, and small stocks of it were noticed in several towns. It is close-grained and beautiful in color, and should be utilized for veneering. It would make most excellent furniture." (Hill and Sudworth.)

Under this name Grosourdy (2: 363) refers to *Acana* and adds that the military authorities of San Juan use this wood for the spokes of wheels, axles of carriages, etc. Some of its particular uses are for beams in houses, windows, doors, etc.

Auzu. See *Amomis caryophyllata*.

Averrhoa bilimbi. BIMBLING. BILIMBI.

Avicennia nitida. MANGLE BLANCO. BLACK MANGROVE.

Family Verbenaceae; a shrub or tree growing in tidal swamps; the wood is considered valuable for its peculiar durability in damp situations, and is accordingly used for foundations and drains; under ordinary circumstances it decays quickly; the bark is used for tanning. Bello calls this "mangle bobo." (Stahl, 6: 220.)

This is called both "black" and "white" mangrove. The former perhaps because of the dark-colored bark, the latter because of the tomentose leaves. Also called "olive mangrove."

Avicennia officinalis.

Reported by Grosourdy under the name "mangle blanco."

Avipillo. See *Majepoa caribaea*.

Avispillo. See *Turpinia paniculata*.

Avocado pear. See *Persea gratissima*.

Avocate. See *Persea gratissima*.

Azafran.

A tree from all parts of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 20 to 25 inches (50 to 63 centimeters); wood, yellowish, hard; specific gravity, 0.628; used in building houses. Also called "laurel puero." (Exp. 1857.)

Azota-caballo. See *Malpighia coccigera* and *Duranta plumieri*.

Azufaifo. See *Zizyphus reticulata*.

Babeiro. See *Echites circinalis* and *E. umbellata*.

Babeiro amarillo. See *Echites andrewsii*.

Bacopa chamaedryoides. YERBA DE CULEBRA AMARILLA.

Family Scrophulariaceae; a decumbent annual, found in ditches and moist places. (Stahl, 6 : 231, as *Herpestis chamaedryoides*.)

Bacopa monniera. YERBA DE CULEBRA.

Herb found in the same kind of places as the last. *B. repens* and one other species receive the same common name.

Bactris pavoniana.

A small palm reported from Porto Rico by Grisebach, but with no more definite locality. The pinnae are narrow and grass-like a quarter or half an inch apart, and there are more than 30 pairs, 12 to 18 inches (30 to 45 centimeters) long, from 2 to 4 lines (4 to 8 millimeters) broad, but remaining united toward the end of the leaf, and here 6 to 8 lines (12 to 16 millimeters) broad; prickles scattered, the longest 2 inches (5 centimeters).

Baculo. See *Aguti grandiflora*.

Badiera de Santo Domingo. See *Badiera domingensis*.

Badiera domingensis. BADIERA DE SANTO DOMINGO.

Family Polygalaceae. Bello reports this from Guanajibo; not found by Stahl. (Stahl, 2 : 54.)

Badinjan. See *Solanum melongena*.

Bahama grass. See *Capriola dactyloa*.

Ballata tree. See *Mimusops*.

In Dominica this name is said to be applied to *Bumelia retusa*.

Balsam apple. See *Momordica charantia* and *M. zeylanica*.

Balsam fig, Balsam tree. See *Clusia rosea*.

Balsamillo. See *Coccocypselum repens*.

Bálsamo. See *Psychotria undata*.

Bálsamo amarillo. See *Hamelia lutea* and *H. patens*.

Bálsamo colorado. See *Hamelia patens*.

Bamboo.

In Porto Rico the bamboo is apparently of rather recent introduction, and is neither so common nor so widely distributed as might be expected. A single species is growing at numerous points along the military road, but was nowhere seen in large quantities, and seems not to be known about Guayama. Neither is the bamboo much in use for domestic purposes, although the moist climate is very suitable for growing it and the difficulty of securing wood in some localities should make its value very considerable for miscellaneous farm purposes. It seems most desirable that a selection of the larger and more useful species be introduced from Japan, China, and the East Indies.

An important use of this plant from the agricultural standpoint is in making pots for seedlings. These will last for a year or two, and greatly decrease the trouble and expense of raising and transplanting seedlings of all kinds. The pots are made by simply sawing off the ripe stems an inch (25 millimeters) below the joint and 6 inches (15 centimeters) or more above it. On transplanting, the pot can be split down the side without injury to the roots of the young plant or simply buried in the ground, where it will soon decompose.

In using bamboo for fences in the Trinidad gardens, Superintendent Hart states that it has been found desirable to soak the split stems in water for ten or twelve days in order to dissolve out the sugar and gummy matters which otherwise cause them to ferment and invite the attacks of insects. After removal from the water the stems are to be dried as soon as possible, and will then be as durable as other kinds of wood.

Several species of the genus *Bambusa* and allied woody grasses have been introduced into the West Indies and have escaped from cultivation. The name bamboo is also applied in the British Islands to some of the large native grasses, such as *Arthrostyidium* and *Chusquea*.

Banana. See *Musa*.

Banara portoricensis. PALO DE RAMON.

A shrub or small tree of 3 to 8 meters; family Flacourtiaceae; an indigenous and peculiar recently described species from the vicinity of Adjuntas. (Urban, Symb. 1 : 370.)

Baobab. See *Adansonia digitata*.

Barbasco. See *Jacquinia armillaris*, *J. aristata*, and *Canella alba*.

Barbieria polyphylla. ENREDADERA.

Family Viciaceae. A handsome shrub with long, red flowers. (Stahl, 3 : 86, 160.)

Bardana mayor. See *Xanthium macrocarpum* and *X. canadense*.

Bariaco.

A wild tree reaching a height of 30 feet (9 meters), with a trunk diameter of 18 inches (45 centimeters). It furnishes a wood much appreciated in the country for carpenter work, used especially in house building; also for furniture and certain musical instruments. It is strong and very resistant, fibrous in texture, with a close, fine grain, reddish gray in color, variously marked and marbled. (Grosourdy, 2 : 364.) This author refers this to *Trichilia moschoxylum*. A species that does not appear in Index Kewensis; perhaps it should be *T. moschata*, a Jamaican species.

Basil. See *Ocimum basilicum*.

Basora prieta. See *Waltheria indica* and *Cordia ulmifolia*.

Basote. See *Chenopodium ambrosioides*.

Bastardia bivalvis. ESCORA BABOSA.

Family Malvaceae. (Stahl, 2 : 69, as *Sida viscosa*.)

Batata. See *Ipomoea batatas*.

Batata blanca. See *Ipomoea batatas*.

Batatilla blanca. See *Ipomoea pentaphylla* and *I. quinquefolia*.

Batatilla carnosa. See *Ipomoea carnea*.

Batatilla ventruda. See *Ipomoea ventricosa*.

Batatilla de zaeta. See *Ipomoea sericantha*.

Batatillo.

Recorded by Bello for *Ipomoea caruosa*.

Bauhinia amarilla. See *Bauhinia kappleri*.

Bauhinia kappleri. FLAMBOYANT BLANCO, BAUHINIA AMARILLA.

PLATE XIX.

Also called "varital." A curious and beautiful leguminous tree, 10 to 15 meters high, cultivated for ornament and shade, and also becoming wild in open places. Supposed to be native in French Guiana. *Bauhinia krugii* is now considered by Urban to be a synonym of this species. (Urban, Symb. 1: 315.)

Bauhinia krugii. See *Bauhinia kappleri*.

Bauhinia porrecta.

A leguminous tree about 4 meters high, cultivated for its conspicuous flowers. (Stahl, 3: 127.)

Bayberry. See *Myrica cerifera*.

Bayberry tree. See *Amomis caryophyllata*.

Bayleaf. See *Amomis caryophyllata*.

Bay rum. See *Amomis caryophyllata*.

Bay rum tree. See *Amomis caryophyllata*.

Bay tree. See *Amomis caryophyllata*.

Bejuco blanco. See *Bignonia acquinocialis*.

Bejuco colorado. See *Rudolphia colubilis*.

Bejuco de agua. See *Vitis caribaea*.

Bejuco de berac. See *Chiococca racemosa*.

Bejuco de caro. See *Vitis sicyoides*.

Bejuco de clavo. See *Bignonia caryophyllata*.

Bejuco de conchitas. See *Citoria ternatea*.

Bejuco de corrales. See *Serjania lucida*.

Bejuco de costilla. See *Paullinia pinnata*.

Bejuco de garrote. See *Rourea glabra*.

Bejuco de gloria. See *Ipomoea cathartica*.

Bejuco de miel. See *Salmea eupatoria*.

Bejuco de mato. See *Dioeclea reflexa*.

Bejuco de mona. See *Cissampelos pareira* and *C. microcarpa*.

Bejuco de palma. See *Marcgravia rectiflora jacquini* and *M. umbellata* and *M. sintenis*.

Bejuco de paralejo.

A name applied to numerous species of **Malpighiaceae**, such as *Stigmaphyllon* (synonym *Stigmatophyllon*) *floribundum*, *S. periplocifolium* and *S. puberum*; also *Heteropteris chrysophylla*, as well as *Triopteris* and *Tetrapteris*.

Bejuco de paralejo rosado. See *Heteropteris purpurea*.

Bejuco de peo. See *Lasianthus lanceolatus*.

Bejuco de puerco. See *Ipomoea setifera* and *I. fastigiata*.

At Santurce this common name was also applied to *Thunbergia alata*. Bello reports it for *Ipomoea bona-nox*.

Bejuco de puerco blanco. See *Ipomoea ciliolata*.

Bejuco de puerco de costa. See *Ipomoea carnososa*.

Bejuco de puerco de playa. See *Ipomoea biloba*.

Bejuco de rana. See *Marcgravia rectiflora jacquini*.

Bejuco de San Juan. See *Forsteronia corymbosa*.

Bejuco de sopla. See *Securidaca virgata*.

Bejuco de vaca. See *Ipomoea bona-nox*.



FLAMBOYANT BLANCO (BAUHINIA KAPPLERI). COAMO SPRINGS.

Bejuco inglés. See *Capparis jamaicensis*.

Bejuco prieto. See *Paullinia pinnata*, *Hippocratea ovata*, and *Rudolphia volubilis*.

Bejuquillo de puerco. See *Ipomoea triloba*.

Bellorita. See *Erigeron bellioides*.

Ben. See *Moringa moringa*.

Ben nut. See *Moringa moringa*.

Berengena. See *Solanum melongena*.

Berengena cimarrona. See *Solanum torvum* and *S. inclusum*.

Berengena de marimbo. See *Solanum mammosum*.

Berengena de paloma. See *Solanum virgatum*, *S. callicarpifolium* and *S. lentum*.

Berengena de playa. See *Solanum persicaefolium*

Berengena jamaiquiña. See *Solanum jamaicense*.

Bergamota. See *Citrus medica*.

Berica.

This name is noted as being used near Toa Alta for a shrub said to produce an edible fruit (no. 893) probably a species of *Eugenia*. At Vega Baja "berica" or "perica cimarron" was a species of *Acalypha* (no. 1035).

• **Berros.** See *Rorippa officinalis*.

Bertholletia excelsa. BRAZIL NUT.

A large tree belonging to the family Lecythidaceae, and yielding the Brazil or Para nuts of commerce. A tree 100 to 150 feet high, distributed throughout northeastern South America to the Island of Trinidad. It has been cultivated in Jamaica and probably in other islands.

Berugillo.

A wood not much used; specific gravity 0.73; abundant in the hills. (Hansard.)

Betel palm. See *Areca catechu*.

Bichy. See *Cola acuminata*.

Bidens bipinnata. MANZANILLA NEGRA.

An annual composite weed. (Stahl, 5: 128.)

Bidens coreopsidis. See *Bidens rubifolia*.

Bidens pilosa. MANZANILLA DEL PAIS.

An annual weed; found in all parts of the island. It seems to be more often called "margarita." (Stahl, 5: 127.)

Bidens rubifolia. MANZANILLA TREPADORA.

Reported as *B. coreopsidis*. (Stahl, 5: 129.)

Bignonia aequinoctialis. LIANA DE LA SIERRA.

Family Bignoniaceae; a shrubby climber. According to Bello this is called "bejuco blanco." (Stahl, 6: 177.)

Bignonia caryophyllea. BEJUCO DE CLAVO.

A vine with large aromatic roots which are used to color and impart a pleasant odor to rum. (Stahl, 6: 180.)

Bignonia odorata. LIANA FRAGRANTE.

A woody climber growing in waste places on the coast. (Stahl, 6: 178.)

Bignonia unguis. See *Bignonia unguis-cati*.

Bignonia unguis-cati. LIANA UÑADA.

A shrubby climber with sharp claw-like tendrils. Specimens were collected at Guayanillo by Sintenis. Bello calls this "pegapola." (Stahl, 6: 179.)

Bija. See *Bixa orellana*.

More often called Achiote.

Bilimbi. See *Arerrhoa bilimbi*.

Bimbling. See *Arerrhoa bilimbi*.

Birch (West Indian). See *Bursera simaruba*.

Biriji. See *Eugenia monticola* and *E. poiretii*.

Bitter wood. See *Picrasma excelsa*.

Bixa orellana. ANATTO. ACHIOTE.

Family Bixaceae. The achiote or anatto is a small tree of handsome appearance, with large, clean, cordate leaves and numerous pink flowers, followed by bur-like pods, at first green, but changing to deep red and becoming an inch in diameter. These contain the seeds, the arillus or fleshy covering of which is bright orange in color and constitutes the anatto of commerce. On drying, the arillus becomes dull orange. Quantities of dried seeds are to be found in Porto Rican markets for domestic use in soup and in coloring rice and other dishes yellow. In the English colonies the coloring matter is removed while fresh, and then dried and compacted into cakes, in which form it is exported to the United States for manufacture into butter-color. Supposedly for this purpose 726,269 pounds were imported into the United States in 1899, valued at \$34,827 but recent analyses of butter colors show that they consist largely of analine dyes.

The wood is nearly white, porous, light, and very soft, with a specific gravity of 0.399. It has no use whatever. When polished the wood is slightly yellow, or sometimes with a tendency toward red, and very uniform in color, although it retains narrow lines of a slightly darker color, which correspond to the concentric zones to be seen in a cross section. (Grosourdy, 2: 357.)

Black bead. See *Pithecolobium unguis-cati*.

Black mangrove. See *Avicennia nitida*.

Blechum brownei. YERBA DE PAPAGAYO.

Family Acanthaceae; an indigenous, annual herb, affecting dry situations and flowering continuously. (Stahl, 6: 244.)

Bleeding heart. See *Colocasia esculenta*.

Bledo blanco. See *Amaranthus paniculatus*.

Blero. See *Mesosphaerum capitatum*.

Blighia sapida. AKEE.

A beautiful tree of the family Sapindaceae, introduced from Africa to the West Indies by the same Captain Bligh who introduced the breadfruit, above which it is said to be valued by the people of Jamaica as a richly flavored and wholesome food. The bright yellow, fleshy arillus is the part eaten. In some islands, however, the akee is avoided as poisonous, though the existence of any deleterious principle has yet to be demonstrated, and it is now suggested by Superintendent Hart, of the Trinidad Gardens, that the cause of injury in the reported cases of poisoning has been the use of decayed fruits, which seems the more probable in view of the recent discovery of the extremely poisonous nature of some of the products of organic decomposition. The following caution should be accordingly heeded by any who may wish to experiment with this fruit:

“ It is prepared in Jamaica in many ways, a favorite method being to stew in milk and afterwards to brown in a frying pan with butter. Again it may be boiled or even baked. It is also commonly eaten boiled and mixed with salt fish, onions, and tomatoes as a breakfast dish. Although so commonly used it can not be denied that cases of poisoning have followed the consumption of it as an article

of food, but so little cause is there to fear its effects that when in season it is to be found generally on the tables of both rich and poor throughout Jamaica. The cases in which the fruit has produced poisonous effects are generally found to be such as show a careless preparation of the fruit for food. The arillus when in good condition is of a wholesome, nutty flavor, and should be quite firm and not easily broken. If, however, it is kept too long it readily becomes soft and decays, and such decayed parts are doubtless poisonous as any other decayed vegetable or animal matter and quite as likely to cause irritant poisoning. Care should therefore be exercised by the housewife in preparing such a delicacy for table, and if such is taken it may be eaten without the slightest fear, and with the satisfaction of having partaken of one of the finest *bonne bouches* that come to a West Indian table. The writer speaks of it from experience of some twenty years, during which time he has missed no opportunity of securing a share of it when in season."

Blood flower. See *Asclepias curassavica*.

Bocagea laurifolia.

Family Anonaceae; from Guajataca and Sierra de Lares.

Bocconia frutescens. PALO DE PAN CIMARRON.

Family Papaveraceae; an herb with a disagreeable odor. (Stahl, 2: 34.)

Boehmeria cylindrica.

An urticaceous plant of interest chiefly because related to the next species; known from Aybonito.

Boehmeria nivea. RAMIE.

Family Urticaceae; the now celebrated ramie plant or Chinese hemp, the cultivation of which has been so widely advocated. Notwithstanding the numerous attempts at commercial utilization of the fine fiber produced by this species, practical success has not yet been attained, the satisfactory processes and machinery for degumming the fiber being too complicated and expensive to permit ramie to compete in the manufacture of cheap fabrics. The Porto Rican specimens were found by Sintenis in cultivation, near Cabo Rojo. It is not known whether any local use has been made of the fiber.

Boerhavia paniculata. TOSTON.

Family Nyctaginaceae; this species and *B. hirsuta* are common tropical weeds known from Porto Rico, but without recorded local names, "toston" being given as Cuban only. (Urban, 3: 307, 308.)

Boerhavia scandens. PEGA POLLO.

A shrubby herb of about 4 feet (1.2 meters), widely distributed throughout the American tropics. (Urban, Add. 3: 309.)

Boje. See *Antirrhoea coriacea*.

A wild tree not infrequently met with in the northeast part of the island. Not more than 45 or 50 feet (14 or 15 meters) in height, with a straight trunk 18 to 24 inches (45 to 60 centimeters) in diameter. It furnishes a heavy durable wood, very strong and much appreciated for carpenter work, furniture, framework of houses, etc., yellow in color, brittle, breaking horizontally. Also called "quina" and referred to *Contarea coccinea*, a species that does not appear in Index Kewensis. (Grosourdy, 2: 367.)

Bombija. See *Cardiospermum halicacabum*.

Boniato.

A tree from all parts of the island; height 10 to 12 feet (3 to 3.7 meters); diameter 12 to 15 inches. Wood, dark, streaked with yellow; hard; specific gravity, 0.049 (1.049?); used in cabinetmaking. (Exp. 1857.)

Borreria parviflora. See *Spermatocoe ocymoides*.

Borreria podocephala. See *Spermacoce verticillata*.

Borreria arborescens. CLAVELON DE PLAYA.

A biennial composite shrub. (Stahl, 5: 134.)

Boton blanco. See *Spermacoce verticillata*.

Boton blanco de arenales. See *Spermacoce verticillata*.

Botoncillo. See *Spermacoce oeymoides*.

Boton de cadete. See *Leonotis nepetifolia*.

Boton de oro. See *Clerodendron aculeatum*.

Bottle gourd. See *Lagenaria vulgaris*.

Bougainvillea spectabilis. TRINITARIA.

A showy climbing shrub, belonging to the family Nyctaginaceae. Its striking appearance is due to the large purplish pink bracts, the true flowers being yellowish and quite small. The color is extremely conspicuous, and to some it appears painfully "loud." In Porto Rico a quite different plant, *Antigonum leptopus*, producing similar large masses of brilliant color, though of a more pleasing shade of pink, has been mistaken for Bougainvillea by travelers and popular writers.

Bourreria domingensis. ROBLE GUAYO. BURERIA.

Family Boraginaceae; a shrub or small tree reported by Stahl from the vicinity of Quebradillas. (Stahl, 6: 99.)

Bourreria succulenta. ROBLE GUAYO. BURERIA.

A small indigenous tree growing in the forests near the coast. It is said to strikingly resemble *Cordia elliptica* in both habit and inflorescence. (Stahl, 6: 97.)

Boussingaultia baselloides. MADEIRA VINE. SUELDA CON SUELDA.

Family Chenopodiaceae; a slender vine commonly cultivated and trained over arbors. Supposed at Cataño to have medicinal value in rheumatism.

Brachyramphus caribaeus. See *Lactuca intybaea*.

Bradburya laurifolia. CONCHITA LAURIFOLIA.

A leguminous plant reported by De Candolle as occurring in pastures in Porto Rico; not known by Stahl. (Stahl, 3: 64.)

Bradburya plumieri. CONCHITA DE PLUMIER.

A wild vine. (Stahl, 3: 60.)

Bradburya pubescens. CONCHITA PELUDA.

A wild annual vine growing in waste places. (Stahl, 3: 61.)

Bradburya virginiana. CONCHITA VIRGINIA.

A slender, wild, annual vine growing in all parts of the island. (Stahl, 3: 62.)

Brasil.

A tree from all parts of the island; height, 25 to 30 feet (8 to 10 meters); diameter, 5 to 6 inches (12 to 15 centimeters). Wood, dark, hard; specific gravity, 1.099; used in making a dye. (Exp. 1857.)

Brasil de costa.

A tree from the south part of the island; height, 25 to 30 feet (8 to 10 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood, yellow, hard; specific gravity, 0.671; used in making planks and joists. (Exp. 1857.)

Grosourdy refers this to *Caesalpinia cristata*, a species that does not appear in Index Kewensis. He may possibly refer to *C. crista*, a West Indian species.

A tree 30 feet (10 meters) high, with a naked, crooked trunk, 25 inches (62 centimeters) in diameter. The wood is rather compact, heavy, hard, and strong. In color much darker than *C. bijuga*, and used for the same purposes. Used also in Porto Rico for boards and beams in country houses. (Grosourdy, 2: 367.)

Brasilete.

Enumerated by Hill among trees used for dyeing and tanning. In Cuba the name is applied to *Caesalpinia pectinata* and other related species. In Stahl's Flora "palo de brasilete" is *Ecastophyllum monetaria*, a leguminous shrub, 6 to 12 feet (2 to 4 meters) high. Grosourdy refers it to *Caesalpinia bijuga*.

Brassica juncea. MOSTAZA.

An annual cruciferous weed, sometimes cultivated and used in medicine as a substitute for mustard. Bello reports this common name for *Sinapis brassicata*. (Stahl, 2: 184.)

Brassica napa. TURNIP.

The turnip is very little cultivated in Porto Rico. In the market at San Juan very small ones were selling at four for 1 cent.

Brassica oleracea. CABBAGE. COL.

The cabbage is a temperate vegetable very popular in tropical countries, and generally cultivated on a small scale in private gardens, in spite of the fact that only indifferent success is attained. The plants mostly fail to form heads or produce only a small and loose cluster of leaves. Under very favorable circumstances and by using good seed of early varieties it is occasionally possible to attain somewhat better results, but at the time of our visit cabbages from the United States were selling at half a peso (30 cents) apiece, no competition being afforded by the insignificant home-grown vegetable. With better care and better seed the latter can undoubtedly be improved, but the importation of American cabbages at lower prices may be expected to greatly increase.

For cultivation in the tropical garden the variety of cabbage called collards is generally more satisfactory than the sorts usually grown in the North. This variety does not head, but the leaves are more delicate than the outer leaves of cabbage and make very acceptable and wholesome "greens." It will flourish where cabbages can be produced only with difficulty or not at all, with the additional advantage of growing as a perennial and thus remaining available for food during most of the year.

Brazil nut. See *Bertholletia excelsa*.

Bread fruit. See *Artocarpus communis*.

Bretonica. See *Chaptalia integrifolia* and *Valerianodes*.

Bretonica afelpada. See *Melochia tomentosa*.

Bretonica aserrada. See *Melochia serrata*.

Bretonica peluda. See *Melochia hirsuta*.

Bretonica piramidal. See *Melochia pyramidata* and *M. nodiflora*.

Bretonica prieta. See *Melochia nodiflora*.

Brigueta. See *Ilex dioica*.

Brigueta naranjo. See *Ilex macoucoua*.

Bromelia pinguin. PINGUIN. PIÑUELA.

A wild relative of the pineapple, found everywhere at lower elevations in Porto Rico, and often planted for hedges. The margins of the leaves have sharp, cutting spines, mostly curved upward, but often reversed so that even the most casual contact is likely to result in a wound. It was Bromelia which added so much to the difficulty of operations about Santiago, and a hedgerow of it is indeed a formidable obstacle for men or horses.

The pulp inclosing the seeds has a pleasant, slightly acid taste somewhat suggesting the pineapple; it is also used as an anthelmintic. Like those of the true pineapple and most other members of the present family, the leaves contain a fine fiber which might be utilized if obtainable in good condition and in sufficient

quantity. In the present species, however, the fibers have been found to have a gummy coating which increases the difficulty of preparation, so that experiments in commercial production have not been successful. Other South American Bromeliaceae have cleaner fibers which have been shipped to Europe in considerable quantities, but the industry has remained on a rather precarious basis.

In the discussion of the pineapple, attention has been called to the fact that in Porto Rico Bromelia is the host plant of a mealy bug which also infests the pineapple, to which it is probably carried by the ants. If the pest should prove serious it may be found necessary to exterminate Bromelia in regions devoted to pineapple culture.

Bruja. See *Bryophyllum calycinum*.

Brunfelsia densifolia.

Family Solanaceae; an indigenous tree, 8 to 10 meters in height, from the vicinity of Maricao. (Urban, Symb. 1: 402.)

Secured by Sintenis near Maricao.

Brunfelsia lactea. VEGA BLANCA.

A shrub, sometimes reaching 6 meters; reported from Luquillo, Naguabo, and Cayey. (Urban, Symb. 1: 401.)

Brunfelsia portoricensis.

A shrub of 3 meters; known only from the Sierra de Luquillo. (Urban, Symb. 1: 400.)

Bryophyllum calycinum. BRUJA.

Family Crassulaceae. In Cuba this plant has several names—"hoja bruja," "immortal," "polipo herbacea," "siempreviva," and "vivora,"—all referring to its great tenacity of life, small pieces and even single leaves sending out new growth when suspended in a moist atmosphere. Buds arise from the notches of the leaves. It is this plant, a common tropical weed, which recently received the distinction of a letter from a consular officer which was published by the State Department and resulted in numerous inquiries from all parts of the country. It has long been known in cultivation, on account of its extreme hardiness, but seldom flowers outside the Tropics, where it seems to thrive best when growing in crevices of bare or slightly shaded rocks; in such situations it flowers abundantly in the dry season. (Stahl, 4: 191.)

Bucago. See *Erythrina glauca*.

Bucaro. See *Erythrina micropteryx* and *E. umbrosa*.

According to Captain Hansard the "bucare" of Porto Rico is the same as the tree called "madre de cacao" used for shade in the cacao growing districts of Venezuela and Trinidad, because of its extremely rapid growth, as much as 10 feet (3 meters) per year. It is also said to be called "immortelle" in the French islands. The South American species is supposed to be *E. umbrosa*, but according to Urban this is not known from the islands north of Trinidad, its place being taken by *E. micropteryx*, which is also found in Porto Rico. Hucar, jucar, ucar, and ucare are supposed to be variations of bucare, but another quite different tree (*Bucida buceras*) was pointed out to us under those names.

Buchnera elongata. ESPIGA DE SAN ANTONIO.

Family Scrophulariaceae; a stiff, scarcely branched herb, growing on inundated sandy soil. (Stahl, 6: 229.)

Bucida buceras. UCAR.

PLATE XX.

Family Combretaceae; a large tree noted at several points on the south side of the island; specimens from Coamo Springs (No. 714). (Stahl, 4: 140.)

A wild tree abundant in all parts of this and other islands of the West Indies; 24 to 80 feet high, with a straight trunk 4 to 5 inches in diameter. It furnishes a



UCAR TREE BUCIDA BUCERAS . COAMO SPRINGS.



ALMACIGO (BURSERA SIMARUBA). COAMO SPRINGS.

yellowish white wood, very strong and flexible, commonly used for shelves in houses and for mallets and wooden cogs; also used for knees in boat building. Native name "hucar blanco." (Grosourdy, 2:390.)

Under the native name "ucar," on page 362, the same author describes the wood of this species as olive-colored, more or less dark, with stripes and undulations, and adds that from its compact and fine-grained texture it would appear suitable for polishing. It has, however, the defect of cracking easily. Specific gravity, 1.009.

Buenoceo.

Spelling doubtful; applied at Caguinas to *Thunbergia erecta*.

Buenos dias. See *Hibiscus bifurcatus*.

Bullock's heart. See *Anona reticulata*.

Bully tree.

A name used in the British West Indies, according to Grisebach, for *Sapota sideroxyton*, *Myrsine lacta*, and several species of *Dipholis*.

Bumelia krugii.

Family Sapotaceae; a recently described species from the vicinity of Guanica.

Bunchosia glandulifera.

Family Malpighiaceae; from Sierra de Jabucoa and Añasco, in cultivation.

Bunchosia glandulosa.

From Guanica, Peñon, and near Ponce. A variety (*latifolia*) is known from Fajardo and Peñuelas. *Bunchosia nitida* is reported from Guanica.

Bureria. See *Bourreria*.

Burro. See *Capparis portoricensis*.

Burro blanco. See *Capparis portoricensis*.

Bursera gummifera. See *Bursera simaruba*.

Bursera simaruba. ALMACIGO.

PLATE XXI.

This species, better known as *Bursera gummifera*, is one of the commonest and most characteristic trees of Porto Rico. It is native in the forests of the limestone foothills, where it grows to considerable size, but is more conspicuous as a shade tree extensively planted along the improved roads, especially on the south side of the island. It may be recognized at once by the thin reddish bark, which loosens in shreds and sheets like that of some species of cherry and birch, and is on this account called "West Indian birch" in some of the British islands. As in case of the jobo tree (*Spondias lutea*) cuttings root readily, even when large stakes are used, rendering these species very easy of propagation, and accordingly popular for fence posts and similar uses requiring permanence. As a shade tree *Bursera* is seldom a success, the top being too small and irregular. Even when still young it fruits abundantly. The wood is soft and of little value.

Bursera simaruba is the largest tree of the thin forest growth which covers the dry limestone hills of the south side of the island. This is composed mostly of slender Leguminosae—*Pictetia*, *Pithecolobium unguis-cati*, and several others.

In some islands a somewhat aromatic sweetish balsam, supposed to have medicinal properties, is obtained from this species. According to Engler this is dried down to a sort of resin and goes into export trade as chibou, cahibou, or gomart, but it is not stated where these names are used. The leaves, bark, buds, and roots are also believed to have medicinal properties.

Button tree. See *Conocarpus erecta*.

Buxus vahlii.

Family Buxaceae; from Rincon, in thickets near the shore.

Byrsonima crassifolia.

Family Malpighiaceae; a tree collected near Mayaguez by Mr. Heller.

Byrsonima lucida. PALO DE DONCELLA.

Also called "sangra de doncella," according to Stahl, who considers the Porto Rican form a distinct variety or species, which he names *B. portoricensis*, but this suggestion seems not to have been noticed by other botanists. The plant is an indigenous shrub, or small tree, belonging to the Malpighiaceae, and having the branches and inflorescence covered with a yellow powder. It grows close to the shore to the east of San Juan, and blossoms in the winter. The timber is durable, but not of sufficient diameter to render it important.

Byrsonima portoricensis. See *Byrsonima lucida*.

Byrsonima spicata. MARICAO.

A tree 30 to 40 feet (9 to 13 meters) high. The bark is used for tanning. The yellow, acid berries are said to be eaten, although rather astringent. (Stahl, 2: 140.) It furnishes a wood of good weight and ordinary durability. This is compact, uniform, fine-grained, and somewhat fibrous in texture, and cinnamon-brown in color, with narrow parallel light stripes. The wood is strong and much appreciated; used in the country for boards and framework of houses, etc. Specific gravity, 0.812. Native name, "doncella." (Grosourdy, 2: 372.)

Caballo asi. See *Abelmoschus abelmoschus*.

Cabalonga. See *Theretia neriifolia*.

Cabbage. See *Brassica oleracea*.

Cabbage-bark. See *Audira inermis*.

Cabbage palm.

The "palm cabbage" is obtainable from many different palm trees, and consists of the delicate tissues of the newly-formed, unexpanded leaves. In the West Indies the name is more properly applied to *Oreodora oleracea*, a native of Barbados and other southern islands, though apparently extending to Porto Rico if specimens collected by Sintenis in the Luquillo Mountains were from native trees and are properly determined.

Cabbage tree. See *Audira inermis*.

Cabellos de Angel. See *Clematis dioica*.

Cabambreña.

A tree from all parts of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 8 to 10 inches (20 to 25 centimeters). Wood white, soft; specific gravity, 0.799; not used. (Exp. 1857.) Perhaps this is an error for "calambreña."

Cabo de hacha. See *Trichilia hirta* and *T. simplicifolia*.

Cabo negro. See *Arenga saccharifera*.

Cabra.

A tree from all parts of the island; height, 25 to 35 feet (8 to 11 meters); diameter, 8 to 10 inches (20 to 25 centimeters). Wood light red, soft; specific gravity, 0.597; used for weather boards. (Exp. 1857.) (Grosourdy, 2: 368.)

Cacaillo. See *Sloanea berteriana* and *Meliosma obtusifolia*.

Cacao. See *Theobroma cacao*.

Cacao bobo. See *Meliosma obtusifolia* and *M. herbertii*.

Cacao cimarron.

According to Captain Hansard a tree of great height, not resembling the true cacao. It has large, coarse, brittle leaves. The wood has a specific gravity of 1.08, and is used for firewood and walking sticks.

Cacao de monte.

A tree from the interior of the island; height, 40 to 44 feet (12 to 14 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood light-colored, rather hard; specific gravity, 0.964; not used. Also called "cacao macho." (Exp. 1857.)

Cacao hembra.

A tree from the interior of the island; height, 25 to 30 feet (8 to 10 meters); diameter, 8 to 10 inches (20 to 25 centimeters). Wood light-colored, rather hard; specific gravity, 0.817; little used. (Exp. 1857.)

Cacao macho. See *Cacao de monte*.

Cacao motillo. See *Sloanea berteriana*.

Cacao rojo. See *Sarcomphalus reticulatus*.

Cacao roseta. See *Sloanea berteriana*.

A tree from the interior of the island; height, 35 feet (11 meters); diameter, 30 to 40 inches (75 to 100 centimeters). Wood white, soft; specific gravity, 0.488; not used. (Exp. 1857.)

Cachimbo. See *Palicourea crocea*.

Cachimbo verde. See *Palicourea umbellata*.

Cadillo. See *Corchorus hirsutus* and *Urena lobata*; also see under "caillo."

Cadillo altea. See *Triumfetta altheoides* and *T. semitriloba*.

Cadillo anaranjado. See *Pavonia coccinea*.

Cadillo de ciénega. See *Pavonia racemosa*.

Cadillo espinoso. See *Pavonia spinifera*.

Cadillo pata de perro. See *Urena sinuata*.

Cadillo pequeque. See *Pavonia typhalea*.

Cadungo amargo. See *Lagenaria vulgaris*.

Caesalpinia bijuga. BRASILETE.

A small tree found on the coast, 9 to 15 feet (3 to 5 meters) high, with a trunk 18 inches (45 centimeters) in diameter. The wood is very compact, very hard, dark reddish orange in color, and breaks vertically with difficulty. The bark is rather thick. It is utilized in Europe as a dye. It serves also for turning and to make violin cases. (Grosourdy, 2: 367.)

Caesalpinia bonduc. MATO AZUL.

Bello reports the common name, "matos de playa," for this species. (Stahl, 3: 120).

Caesalpinia bonducella. MATO AMARILLO.

A much-branched leguminous shrub, 3 to 4 meters high, frequenting the coast. (Stahl, 3: 121.) The seeds of this species were on sale in the market of Ponce as a remedy for piles. See notes under "mato colorado" and "tortera."

Caesalpinia coriaria. DIVI-DIVI.

A leguminous tree, the pods of which are very rich in tanning and dying material, and constitute a considerable article of export from the West Indies and South America. The species is not included in Stahl's Flora, but is mentioned by Hill, though without anything to show that he established its existence in Porto Rico. Whether already introduced or not, it is probable that the planting of it on a somewhat extensive scale is worthy of serious consideration.

Caesalpinia gilliesii.

Reported from Guanica.

Caesalpinia sepiaria. ZARZA DE CERCAS.

Of shrubby growth; used for fences in Cuba. (Stahl, 3: 122.)

Café. See *Coffea arabica*.

Café de la India. See *Murraya exotica*.

Caffeillo. See *Drypetes alba* and *D. glauca*.

Evidently another spelling of "cafetillo." According to Captain Hansard caffeillo is a wood having a specific gravity of 0.85 and is not much used. The flowers are fragrant.

Cafetillo. See *Casearia nitida* and *Faramaea odoratissima*.

A tree from the interior of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 15 inches (37 centimeters). Wood yellowish, hard; specific gravity 0.840; used for lumber. (Grosourdy, 2: 368.) (Exp. 1857.)

Caillo.

Probably intended for "cadillo." At Santurce *Corchorus hirsutus* and *Xanthium canadense*, and at Quebrada Arenas *Triumfetta semitriloba* were called "caillo."

Caimitillo. See *Chrysophyllum monopyrenum* and *C. glabrum*.

A tree from all parts of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood, white, hard; specific gravity, 0.012 (1.012?); used in making laths. (Exp. 1857.)

Captain Hansard says that the wood is used for shingles and barrel staves. The tree is similar to the star-apple or caimito, but the fruit is only half the size.

Grosourdy gives this as the common name in Porto Rico and Cuba of *Chrysophyllum microphyllum* and *C. oliviforme*. Both of these names are treated as synonyms of *C. monopyrenum* and *C. caimito*. He describes it as a tree growing wild and reaching 40 to 45 feet (12 to 14 meters) in height, with a trunk 18 to 20 inches (45 to 50 centimeters) in diameter. Furnishes a wood of good weight and more than ordinary durability, rather compact and fibrous in texture and rather fine-grained. Dirty white in color or sometimes somewhat chocolate. The wood is used ordinarily for shingles and bowls. It might also be used for carpenter work. (Grosourdy, 2: 368.)

Caimito. See *Chrysophyllum caimito*.

A tree from all parts of the island; height 30 to 35 feet (9 to 11 meters); diameter 10 to 12 inches (25 to 30 centimeters). Wood, violet; hard; specific gravity, 0.194; little used. (Exp. 1857.)

Cainito. See *Chrysophyllum cainito*.

According to Captain Hansard the specific gravity is 0.85.

Cajanus bicolor. See *Cajanus cajan*.

Cajanus cajan. GANDUL.

Family Leguminosae; perennial shrub growing from 3 to 10 feet (1 to 3 meters) high, but treated as an annual in cultivation. Bello calls *Cajanus bicolor* "gandures." (Stahl, 3: 55.)

Cajou. See *Anacardium occidentale*.

Cakile aequalis. See *Cakile maritima*.

Cakile maritima. MOSTACILLA DEL MAR.

A seaside herbaceous plant related to mustard; the common name means "sea-mustard." (Stahl, 2: 37.)

Calabash nutmeg. See *Monodora myristica*.

Calabash tree. See *Crescentia cujete*.

Calabaza. See *Cucurbita pepo*.

Caladium. CARA DE CABALLO.

An ornamental *Caladium* was called by this common name, meaning "horse face," at Cataño.

Caladium esculentum. See *Colocasia esculenta*.

Calambrena. See *Coccolobis nivea*.



ALGODON DE SEDA CALOTROPIS PROCERA. ROAD NEAR GUAYAMA.

Caldo santo. See *Argemone mexicana*.

The local mispronunciation of cardosanto.

Calliandra portoricensis. ACACIA PUERTORIQUEÑA.

A white-flowered leguminous shrub, rare on the north side of the island. Our specimen (no. 481) was collected along the road between Cayey and Guayama. (Stahl, 3: 156.)

Calliandra saman. See *Pithecolobium saman*.

Callicarpa ampla. CAPA ROSA.

A verbenaceous shrub or small tree; specimens were collected by Sintenis at Utuado. Grisebach used the name *Petitia domingensis*. (Stahl, 6: 221.)

Calocarpum mammosum.

Family Sapotaceae, the variety *occideum* is identified by Pierre from Utuado (Sintenis). In Engler this genus is treated under *Lucuma*.

Calonyction megalocarpon. See *Ipomea bona-nox*.

Calophyllum calaba. PALO DE MARIA.

Family Clusiaceae; a tree 60 feet (18 meters) high. The oil pressed from the seeds is said to be used in lamps. Specimens are known from Manati and Rinçon. Bello calls this species "marias." (Stahl, 3: 125). Grosourdy (2: 403) varies the native name to "maria" and describes the plant as a handsome wild tree with a dense top, 55 to 60 feet (17 to 18 meters) high, and a long straight trunk 3 feet (1 meter) or more in diameter. The wood is lasting, rather light, soft, fibrous in texture, reddish in color, rather lighter than mahogany. Specific gravity, 0.655. This wood is highly appreciated for carpenter work and the trunks, when large, serve to make canoes. Planted in rows this tree is sometimes used in parks. If the lower limbs are cut off the tree will make a straight growth, This operation should be repeated every seven years, and each time a good top should be left.

Calopogonium caeruleum. JÍCAMA.

A leguminous vine known from Mexico, Peru, and Brazil. (Stahl, 3: 69, as *Stenolobium caeruleum*.)

Calopogonium orthocarpium.

A small vine with hairy pods; cabo rojo.

Calotropis procera. ALGODON DE SEDA.

PLATE XXII.

Also called "mata de seda." Family Asclepiadaceae; a shrub or small tree said to attain a height of 5 meters in favorable situations. The fibers of the seeds are twisted into thread by some of the African tribes; the bark is also used as a medicine, and from the milk an inferior rubber or rubber adulterant has been obtained. This species is native in the tropics of Africa and Asia, but has been introduced and widely disseminated in America. According to Stahl (6: 86) the height is but 6 feet. Specimens planted along the lower part of the Guayama road are 12 to 15 feet (4 to 5 meters) high, but are bushy rather than tree-like. The leaves are white or grayish with a heavy bloom, and its whole appearance is that of a large milkweed. From the Cape Verde Islands the silk of this milkweed is exported in small quantities as "kapok," the more common source of which is the ceiba or "tree cotton."

Calyptranthes chytraculia. See *Calyptranthes sintenisii*.

Calyptranthes krugii.

Family Myrtaceae; specimens were collected by Sintenis in the vicinity of Luquillo.

Calyptranthes sintenisii. HOJA MENUDA.

Native names: "Limoncillo" and "limoncillo de monte." A myrtaceous shrub or small tree indigenous in Porto Rico, Santo Domingo, and Martinique. Reported from Luquillo and Anasco. (Urban, Add. 2: 116.)

This is probably the "limoncillo" which Grosourdy identifies as *Calyptroanthus paniculata*, a Peruvian species not likely to occur in Porto Rico. A wild tree 50 or 60 feet (15 or 18 meters) high, with a trunk 18 inches (45 centimeters) in diameter. The crushed leaves have the odor of lemons. The wood is very hard, flexible, and violet-colored. It breaks with difficulty with a vertical fracture. Its common use is for shelving, door cases, etc. Specific gravity, 1.112. (Grosourdy, 2: 394.)

Calyptrosidium sintenisii.

A small tree, 15 to 25 feet (5 to 8 meters) high, related to the guava; known only from the forests along the crest of El Yunque.

Camacey. See *Tamouea prasina*.

Camacey-almendro. See *Tamouea integrifolia*.

Camacey blanco. See *Tamouea fothergilla*.

Camacey colorado. See *Tamouea impetiolearis*.

Camacey de charcos. See *Acisanthera quadrata*.

Camacey de ciénega. See *Nepsera aqualica*.

Camacey de paloma. See *Tamouea lacrigata* and *Sagraea fascicularis*.

Camacey-Mariana. See *Rhexia mariana*.

Camacey negligente. See *Clidemia neglecta*.

Camacey peludo. See *Clidemia hirta*.

Camacey racemoso. See *Tamouea racemosa*.

Camacey simple. See *Henriettella fascicularis*.

Camacey-Tomaso. See *Tamouea thomasiana*.

Camandula. See *Coix lachryma*.

Camasey.

Presumably an erroneous spelling of camacey.

Cambrón. See *Cascaria ramiflora* and *Randia aculeata*.

Cambures.

A variety of banana mentioned by Hill.

Cambustera. See *Quamoelit quamoelit*.

Cambutera. See *Quamoelit quamoelit* and *Q. coccinea*.

Cambutera de costa. See *Ipomea filiformis*.

Camocuy blanco.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood, white, rather hard; specific gravity, 0.807; used in building houses. Also known as "camocuy puz." (Exp. 1857.)

Camocuy colorado.

A tree from all parts of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood, pinkish, very soft; specific gravity, 0.80; used in making houses. (Exp. 1857.)

Camocuy prieto.

A tree from all parts of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood, white, rather hard; specific gravity, 0.804; used in making houses. (Exp. 1857.)

Camocuy puz. See *Camocuy blanco*.

Campana de Paris. See *Datura suaveolens*.

Campanilla. See *Sphenoclea zeylanica*.

Campeachy wood. See *Haematoxylon campechianum*.

Caña de azucar. See *Saccharum officinarum*.

Caña de Castilla.

In Cuba applied to the grass *Cyperium saccharoides*.

Caña de la India.

Also called wild cane, "caña cimarron." The juice is squeezed out and used as an emetic. In the market of Ponce the roots of "caña india" were on sale for use in the preparation of a refreshing drink.

Cañafistolo. See *Cassia fistula* (another spelling of Cañafistulo.)

A tree from the interior of the island; height, 60 to 70 feet; diameter, 3 to 9 inches. Wood red, hard; specific gravity, 1.193; little used. (Exp. 1857.)

Cañafistula.

The fruit of *Cassia fistula*.

Cañafistula cimarrona. See *Cassia grandis*.

Cañafistulo. See *Cassia fistula*.

Canario. See *Allamanda cathartica*.

Canavalia ensiformis. SWORD BEAN.

A leguminous woody vine, bearing long pods and white beans. In Africa the young pods and beans are eaten.

Canavalia obtusifolia. MATO DE LA PLAYA.

A woody vine. (Stahl, 3: 81.)

Canavalia rusiosperma. MATO COLORADO.

A large leguminous climbing vine, attaining a thickness of 10 cm. and ascending the highest forest trees; from Maricao and Quebradillas. (Urban, Symb. 1: 474.) The seeds of this species are red, and are probably the ones used with those of *Caesalpinia bonducella* as a magical remedy for piles. See note under that species, also that under "tortera."

Candle wood. See *Oreopanax capitatum*.

Canela.

Supposed by Captain Hansard to be cinnamon, but neither *Cinnamomum zeylanicum*, the true cinnamon, nor *C. cassia*, the Chinese cinnamon, are known from the island. The specific gravity, as given by Captain Hansard, is 0.87.

A tree from the interior of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 10 to 12 inches (25 to 30 centimeters); wood dark, hard; specific gravity, 0.956; used in making country houses. (Exp. 1857.)

A wild tree, 35 to 40 feet (11 to 12 meters) high, with a trunk 12 inches (30 centimeters) in diameter. The wood is dark-colored, rather hard, and used in country houses. Specific gravity, 0.515. (Grosourdy, 3: 369.)

Canelilla.

A tree from the interior of the island; height, 35 to 40 feet (10 to 12 meters); diameter, 10 to 12 inches (25 to 30 centimeters); wood greenish, rather hard; specific gravity, 0.897; used in making country houses. (Exp. 1857.) A wood of this name is noted by Captain Hansard with a specific gravity of 0.60.

Canella alba. BARBASCO. WILD CINNAMON.

Family Canellaceae; a tree 10 to 15 feet (3 to 5 meters) high. The pale, orange-colored bark has an aromatic odor and is used as a tonic. Specimens were collected by Sintenis at Rincon. (Stahl, 2: 127.)

Canelo.

A tree from the interior of the island; height, 15 to 20 feet (5 to 6 meters); diameter, 10 to 12 inches (25 to 30 centimeters); wood dark, rather hard; specific gravity, 0.874; used in making country houses. (Exp. 1857.)

Canna.

Several species of *Canna* grow wild in Porto Rico; whether they are indigenous or have escaped from cultivation is not evident. At Cataño the name "maraga" was applied to *C. edulis* or *C. indica*.

Capá blanca. See *Petitia domingensis* and *Cordia alba*.

Color, clear light brown. Very numerous ducts, occurring singly, and evenly diffused between the medullary rays. The latter numerous, in distinct parallel lines. Structurally similar on the radial section to American beech, the medullary rays giving a beautiful, fine-mottled, satiny appearance. Tangential section unattractive. Straight-grained and suitable for interior finish. Used for rollers in coffee-hulling mills. (Hill and Sudworth.)

According to Hill "capá blanca" is *Cordia alba*, but this identification is not seconded, and Stahl gives *Petitia domingensis* as above. Possibly the *Cordia* identification was inferred from the fact that "capá prieta" is *Cordia gerascanthus*.

A tree from the interior of the island; height, 65 to 70 feet (19 to 21 meters); diameter, 30 to 35 inches (75 to 87 centimeters). Wood white, very hard; specific gravity, 0.955; used in boat building. (Exp. 1857.)

Captain Hansard gives the specific gravity as 0.89.

Capá prieta. See *Cordia gerascanthus* and *C. gerascanthoides*.

A tree from the interior of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood light-colored, very hard; specific gravity, 0.899; used in boat building. (Exp. 1857.)

Captain Hansard gives the specific gravity as 0.75.

Color, rich light brown, with darker streaks and mottlings. Very numerous ducts, occurring singly or irregularly, aggregated in twos and threes between the medullary rays. Medullary rays numerous, indistinct, straight, parallel lines. Wood fibers interlaced, but wood appearing to be straight-grained. Radial section structurally similar to capá blanca. Tangential section somewhat similar to dark heartwood of American elm. Handsome wood for interior finish. Used for flooring. (Hill and Sudworth.)

Capá rosa. See *Callicarpa ampla*.

Capá sabanera.

A tree from the interior of the island; height, 55 to 60 feet (16 to 18 meters); diameter, 12 to 15 inches (30 to 40 centimeters). Wood dark, very hard; specific gravity, 0.978; used in boat building. (Exp. 1857.)

Described by Grosourdy as a wild tree, reaching 30 feet (9 meters) in height, with a trunk 12 inches (40 centimeters) in diameter and furnishing a wood of good weight, hard and strong, dark in color and used in the construction of country houses and for fences. (Grosourdy, 2: 370.)

Capacillo.

According to Captain Hansard this tree grows with "yaiti" on the calcareous foot hills and does not become large. The wood is very hard and has a specific gravity of 0.82, but is used for little except fire-wood. Perhaps the same as "capaillo."

Capaillo. See *Aegiphila martinicensis*.

Capitana. See *Dendropremon caribaeus*.

Capitaneja. See *Verbesina alata*.

Capparis cynophallophora. PALO DE BURRO.

PLATE XXIII.

Family Capparidaceae; a shrub or tree 8 to 25 feet (1.5 to 8 meters) high, found along the coast. Collected by Sintenis in the vicinity of Guanica; said to be a specific for dropsy. (Stahl, 2: 44.)



PALO DE BURRO (*CAPPARIS CYNOPHALLOPHORA*): FRUIT AND LEAVES.

Bello reports seven varieties of this species from the southwestern part of the island. (Bello, p. 236.)

Capparis frondosa. PALO DE BURRO.

A shrub 3 or 4 meters high, reported from Sabana Grande and Manati. (Stahl, 2: 47.)

Capparis jamaicensis. BEJUCO INGLES.

A shrub 6 to 10 feet (2 to 3 meters) high. (Stahl, 2: 185.)

Capparis latifolia. PALO DE BURRO.

Very similar to *C. cynophallophora*, but with the flowers in racemes. (Stahl, 2: 186.)

Capparis portoricensis. BURRO BLANCO.

A tree of 15 to 20 meters, known from Coamo, Peñuelas, San German, and Guanica. (Urban, Symb. 1: 309.)

Capparis verrucosa. PALO DE BURRO.

A tree 5 to 6 meters high, the largest of the Port Rican species. (Stahl, 2: 45.)

Capraria biflora. GOAT WEED. Té.

Family Scrophulariaceae. A small shrub, less than 4 feet (1.3 meters) high. Bello calls this "té del pais." (Stahl, 6: 227.)

Capriola dactylon. BERMUDA GRASS.

Also called Bahama grass in the West Indies. Now widely distributed in tropical and subtropical regions. A useful grass for sandy soils, but sometimes becoming a very bad weed in cultivated land.

Capsicum annuum. PIMIENTO.

Family Solanaceae; native of South America. The fruit furnishes the chilies of commerce. (Stahl, 6: 123.)

Capsicum baccatum. AJÍ PICANTE.

Reported by Bello.

Capsicum frutescens. AJÍ.

Fruits used in the preparation of Cayenne pepper. (Stahl, 6: 124.)

A specimen apparently belonging to this species was called "ají caballero" near Toa Alta.

Cara de caballo.

This name is applied to a variegated ornamental Caladium, probably *C. bicolor*.

Caracho. See *Lagenaria vulgaris*.

One of the names applied to the grooved gourd used as a musical instrument.

Caracolillo. See *Sabineca punicea*.

A wood stated by Captain Hansard to be common on hills. Not much used, but furnishes good firewood. Specific gravity, 1.20.

Caracolillo labrado.

Described by Grosourdy as a wild tree, 60 feet (18 meters) in height, with a trunk 3 feet (1 meter) in diameter. It furnishes a striped, yellow wood. In the form usually met with, the markings are in arches or spirals, from which fact the name is said to have arisen. The wood is very good for building purposes and is used quite extensively for shelves, etc.; its principal use, however, is in the manufacture of furniture. (Grosourdy, 2: 371.)

Caracolillo liso.

A tree from the western part of the island; height 50 to 65 feet (15 to 20 meters); diameter 30 to 35 inches (75 to 90 centimeters). Wood yellow, hard; specific gravity, 0.943; used in making furniture. (Exp. 1857.)

Thought by Grosourdy to be a variety of caracolillo labrado, which it closely resembles in height and aspect, as well as in the color and uses of its wood. (Grosourdy, 2: 371.)

Caracolillo listado.

A tree from the western part of the island; height, 15 to 20 feet (5 or 6 meters); diameter, 30 to 35 inches (75 to 90 centimeters). Wood yellow, streaked; specific gravity, 0.952; used in making furniture. (Exp. 1857.)

Cardenala azul. See *Lobelia cliffortiana*.

Cardiospermum halicacabum. FAROLITOS.

Family Sapindaceae; an annual, climbing herb, known as heart seed, winter cherry, and heart pea. At Rio Piedras a specimen (no. 1004) was obtained, under the name "bombija." (Stahl, 2: 153.)

Cardo. See *Carduus mexicanus*.

Cardosanto. See *Argemone mexicana*.

Carduus mexicanus. CARDO.

A native thistle, not abundant. (Urban, Symb. 1: 470, as *Cirsium mexicanum*.)

Cariaquillo. See *Lantana camara* and *L. crocea*.

Cariaquillo de Santa Maria. See *Lantana involucrata*.

Cariaquito blanco. See *Melanthera deltoidea*.

Carica papaya. See PAPAWE. PAYAYA.

Family in Porto Rico commonly called "lechosa," meaning "milky," or "papaya lechosa." This useful species is everywhere cultivated at low elevations; it has also escaped from cultivation and now grows spontaneously in waste places.

Caro. See *Vitis sicyoides*.

According to the list of the Exposition of 1857, there are two trees of this name, both occurring in all parts of the island. One has a height of 30 to 35 feet (9 to 11 meters); diameter 19 to 20 inches (48 to 50 centimeters). Wood light-colored, hard; specific gravity, 0.980; used in building houses. Also called "chupagallo." The other has a height of 35 to 40 feet (10 to 12 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood yellow, hard; specific gravity, 0.873; used in making furniture. Also called "chino."

Caro agrio. See *Vitis acida*.

Caro de tres hojas. See *Vitis trifoliata*.

Carrasco. See *Comocladia*.

Carubio. See *Fagara monophylla*.

Caryocar nuciferum. SOUARI NUT.

A tree belonging to the family Caryocaraceae. It is a native of Guiana, but has been cultivated in several of the West Indies. Three other species of *Caryocar*—*C. glabrum*, *C. barbinerve*, and *C. intermedium*—have been reported as growing in Martinique, but all are probably introduced.

Both the flowers and the fruit are very large, the latter several inches in diameter and inclosing from 2 to 4 hard-shelled nuts, the size of a hen's egg, but somewhat kidney-shaped, and with rounded warts on the outside. These are described as having a very pleasant flavor and also as yielding an oil to pressure. The tree attains a height of 100 feet (30 meters) or more, and the timber is of good quality, and very durable. Its chief use has been in the line of shipbuilding.

In his "Tropische Agrikultur" Semler expresses a very high opinion of the valuable qualities of the souari nut, considering it in fact the finest of all nuts, and the one most likely to prove successful and remunerative in culture. The taste is said to be similar to the almond, but sweeter, and the small quantities sent to the European markets have found ready sale at good prices. Successful

attempts at culture have already been made on the island of St. Vincent, whence seed could probably be obtained.

Caryophyllus aromaticus. CLOVE.

The true clove, a native of the Malay region, has been introduced into several of the West India islands and in some cases has escaped from cultivation, but this is not known to have happened in Porto Rico. In the British island of Grenada quite extensive experiments have been made with the cultivation of the cloves, but the results have not been as favorable as with nutmegs, not because the trees do not thrive, but because the market is overstocked with the product of Zanzibar and the East Indies. Whenever the practice of intelligent mixed farming shall have been introduced into Porto Rico, the clove may serve as a minor resource. For culture on a large scale it does not appear for the present to be especially promising, in view of the results of the Grenada experiment.

Carusillo.

A native grass valued for pasturage in the region of Yanco, and considered sweeter and more nutritious than either Guinea or Para grass.

Cascabelillo alado. See *Crotalaria stipularis*.

Cascabelillo axilar. See *Crotalaria lotifolia*.

Cascabelillo grande. See *Crotalaria retusa*.

Cascabelillo vacío. See *Crotalaria incana*.

Casearia bicolor. TALANTRON.

Also called "cotorrerillo." A tree of 15 meters, belonging to the family Flacourtiaceae. Known only from the vicinity of Utuado.

Casearia hirta.

A shrub or tree, from river banks near Sabana Grande.

Casearia nitida. CAFETILLO.

A shrub 6 to 15 feet (2 to 5 meters) high. (Stahl, 4: 33.)

Casearia parvifolia. GIA MANSA.

A shrub about 15 feet high. From Peñuelas and Utuado in mountains and waste places. (Stahl, 4: 31.)

Casearia ramiflora. CAMBRON.

A shrub or tree 6 to 20 feet high. From Utuado and Manati, in thickets. Bello reports the common name for *C. spinosa*. (Stahl, 4: 30.)

Casearia samyda. COTORRERILLO.

Reported by Bello from Utuado.

Casearia spinosa. See *Casearia ramiflora*.

Casearia stipularis. GIA VERDE.

A shrub or small tree of from 3 to 4 meters; also called "rabojunco." (Stahl, 4: 27.)

Casearia sylvestris. SARNA DE PERRO.

A shrub or tree attaining 35 feet (11 meters), common in mountain woods and waste places. (Stahl, 4: 26.)

Cashaw. See *Anacardium occidentale*.

The more usual and preferable form is cashew. In Jamaica the name "cashaw" is said to be applied to *Prosopis juliflora*.

Cashew. See *Anacardium occidentale*.

Cassada. See *Manihot*.

Cassareep. See *Manihot utilissima*.

Cassava. See *Manihot*.

Cassia alata. See *Herpetica alata*

Cassia bacillaris. SEN DE PALILLOS.

A trailing herb growing in waste places both in the lowlands and in the mountains. (Stahl, 3: 110.)

Cassia bicapsularis. SEN DEL PAIS.

Probably an introduced species; known in Porto Rico only from the vicinity of Fajardo, but distributed elsewhere in tropical America and in Madeira. (Stahl, 3: 109.)

Cassia biflora. RETAMA.

A fine-leaved species with numerous yellow flowers; reported by Bello from the vicinity of Cabo Rojo. (Stahl, 3: 114.)

Cassia diphylla. See *Chamaecrista diphylla*.**Cassia eglandulosa.** See *Cassia emarginata*.**Cassia emarginata.**

With rather long, straight, somewhat turgid pods. (Stahl, 3: 114.)

Cassia fistula. CAÑAFISTULO.

A tree sparingly introduced in Porto Rico. It is said to be a native of upper Egypt and is valued for the sake of the purgative pulp in which the seeds are embedded. The cylindrical pods are 15 to 20 inches long. (Stahl, 3: 105.)

In some countries the bark is said to be used for tanning.

This tree must have been very early introduced into the West Indies, as Acosta states that the fleet in which he returned to Spain in 1587 carried 45 hundred-weight of the fruit for medicinal purposes.

A low tree common in all the Antilles and on the American continent; about 20 feet high with a trunk 9 or 10 inches in diameter. Furnishes a strong wood, reddish in color, of ordinary weight and hardness, but without use. (Grosourdy, 2: 369.)

Cassia grandis. CAÑAFISTULA CIMARRONA.

A tree attaining the height of 10 meters, said by Stahl to be found in Porto Rico in the wild state, though rarely; it is also native on the Isthmus of Panama and other parts of tropical America. The pulp of the pods is said to be used as a substitute for that of *Cassia fistula*. The flowers are rose-colored and appear in April in large lateral racemes. (Stahl, 3: 106.)

A fine tree 45 to 54 feet (14 to 17 meters) high with a dense top and trunk 12 to 18 feet (4 to 6 meters) long and 12 to 15 inches (30 to 37 centimeters) in diameter. Furnishes a very handsome wood, very strong and very resistant, of ordinary weight and hardness, fibrous in texture and fine-grained; specific gravity 0.816. Although not at present utilized, Grosourdy considers this wood very suitable for carpenter work and perhaps for cabinetwork. (Grosourdy, 2: 370.)

Cassia glandulosa. See *Chamaecrista glandulosa*.**Cassia laevigata.** SEN DEL PAIS.

Shrubby; leaves very similar to those of *C. occidentalis*. Reported from Adjuntas. (Stahl, 3: 108.)

Cassia obtusifolia. See *Cassia tora*.**Cassia occidentalis.** HEDIONDA.

This is the common tropical weed which has been variously reported from the West Indies and from Florida as a coffee substitute. Even in Porto Rico this idea has long been prevalent, and the seeds are regularly on sale in the markets, and small quantities have been exported. The usual price is 12 centeros (7.2 cents) per pound. Analysis has failed to find caffeine or any similar alkaloid, and the "coffee" must be ranked with that made from burnt peas and the like. Tonic and febrifugal properties have been ascribed to this drink, and it has also been

used as a coffee adulterant. An attempt was made to introduce it into Europe from Africa as "fedegosa," "negro coffee," or "cafe de Soudan." (Stahl, 3: 111.)

Cassia polyphylla.

Known from Salinas de Cabo Rojo and Coamo.

Cassia tora.

A weedy rank herb with long falcate linear pods. (Stahl, 3: 115.)

Cassipourea elliptica. PALO DE OREJA.

Family Rhizophoraceae. A tree or shrub 3 to 9 meters high. (Stahl, 4: 142.)

Castalia. WATER LILY. FLOR DE AGUA.

Stahl describes under the synonym *Nymphaea* three indigenous species, *N. ampla*, *N. minor*, and *N. rudgeana*. Bello gives the name "yerba de hicotea" for *N. ampla*, with the explanation that tortoises (*tortugas*) are in Porto Rico called "hicotea."

Castaño.

In Spain the name of the chestnut, *Castanea sativa*, but in Porto Rico and Cuba applied to the breadfruit tree, doubtless on account of the fact that the seeds somewhat resemble chestnuts in size, shape, and color.

Castor bean. See *Ricinus*.

Casuarina.

The Casuarinas about San Juan are few and spindling, and do not make the vigorous growth apparent in the streets of Key West, but on the drier south side of the island they seem to flourish much better, and some are supposed to have been introduced from that island, although in reality natives of Australia.

Catalpa longisiliqua. ROBLE.

A wild tree rather abundant in the Antilles and on the Continent, 40 to 50 feet (12 to 15 meters) high, with a trunk 15 to 30 inches (37 to 75 centimeters) in diameter. It furnishes a wood very hard, very strong, and very good for rough carpenter work, carts, etc. (Grosourdy, 2: 382; also 411.)

Catesbaea melanocarpa.

Family Rubiaceae; a species recently described from Porto Rico, St. Croix, and Antigua, formerly confused with *C. parviflora*. A shrub from the south side of the island. (Urban, Symb. 1: 427.)

Catesbaea parviflora.

A spiny shrub 6 feet (2 meters) high, known from Guanica (summit of mountain called "El Maniel"), and from near Guayanilla. (Urban, Symb. 1: 428.)

Catire.

A vine sold in the market of Ponce as an ingredient of a refreshing drink.

Cat-tail. See *Typha domingensis*.

Cautiva.

A bignoniaceous tree was called by this name at Santurce, but our specimen (no. 81) has not yet been identified.

Cayaponia americana.

Family Cucurbitaceae. A variety *subintegrifolia* has been described from near Maunabo. (Stahl, 4: 172.)

Cayaponia racemosa. COLOQUINTILLA.

An annual vine growing in waste places. Cogniaux has named a variety *plumieri*. (Stahl, 4: 170, as *Cionandra racemosa*.)

Cayenne pepper. See *Capsicum*.

Cayur. See *Annona palustris*.

Ceanothus reclinatus. See *Colubrina reclinata*.

Cebada. See *Hordeum*.

Cebolla. See *Allium cepa*.

Ceboruquillo. See *Thouinia tomentosa*.

A tree from the western part of the island: height 70 to 80 feet (21 to 24 meters); diameter 10 to 12 inches (25 to 28 centimeters). Wood, light-colored, hard; specific gravity, 0.857; used for packing boxes. (Exp. 1857.)

Cecropia obtusa.

Collected by Sintenis near Bayamon.

Cecropia peltata. TRUMPET TREE. YAGRUMO HEMBRA.

The name "yagrumo hembra" has reference to the fact that this tree is popularly supposed to be the female of *Didymopanax morototoni*, called "yagrumo macho." The stems of the present species are hollow; it grows very rapidly and although attaining a height of 50 feet (15 meters) with a diameter of a foot (25 millimeters) or more, it seems rather like an overgrown herb than a real tree. The trunk and branches are partitioned at the nodes or former points of attachment of the very large peltately divided leaves.

The juice of *Cecropia* contains india rubber, but the quantity obtainable by the usual process of tapping is small and the product has not become of commercial importance.

Cedar. See *Cedrela odorata*.

Cedrela odorata. WEST INDIAN CEDAR. CEDRO.

Family Meliaceae. A large tree; wood of a brown color and fragrant odor, known variously as West Indian, Spanish, Jamaica, or Honduras cedar: used for furniture, moth-proof chests, shingles, canoes, and other articles that have to withstand the weather; also for cigar boxes. (Stahl, 2: 168.)

Color, pale reddish brown. Large ducts few, occurring singly, irregularly grouped, and scattered. Medullary rays few and indistinct. Wood fibers interlaced, the wood splitting irregularly. Very similar to the mahogany of commerce. Probably *Cedrela odorata*, the well-known cigar-box wood of commerce. It is no longer abundant in Porto Rico, and is now largely imported from Santo Domingo, costing \$150 per 1,000 feet. It still grows in Aguadilla and near Aibonito, Juana Diaz, Cayey, and Luquillo. (Hill and Sudworth.)

Grosourdy distinguishes three varieties of the wood—cedro blanco, cedro hembra colorado, and cedro macho—but states that the trees are identical.

The tree is 60, 90, or 100 feet (18, 27, or 30 meters) high, with a long straight trunk $4\frac{1}{2}$ to 6 feet (1.5 to 2 meters) or more in diameter. Planted this tree would be a source of wealth, and already plantations exist in Cuba, an example that should be followed in the other Antilles, for it not only grows with greater rapidity than mahogany, but is also much less delicate and will prosper in all localities. The tree, however, prefers a clean light soil on the tops of hills or on plains. It is not injured by moisture, except when the soil is submerged. It is propagated by seeds or cuttings, and reaches its growth in forty years.

This precious wood of agreeable odor is very lasting. It is heavier, more compact, and darker colored than mahogany, which the variety called "cedro macho" much resembles: the variety known as "cedro hembra" is much lighter, less compact, softer, and much lighter in color. The first-named variety is used for furniture, carpenter work, and the interior of houses, and for the latter is preferred to mahogany, being softer and more easy to work, also having the advantage of never being attacked by insects. It is also used for windows, balconies, etc. The military authorities use it for boxes, powder vessels, etc.

The second variety is fit for the same purposes, and is furthermore used extensively in making cigar boxes. The specific gravity is 0.474 and 0.461. (Grosourdy, 2: 374.)



CEIBA TREE (CEIBA PENTANDRA). NEAR PONCE.

Cedro blanco. See *Cedrela odorata*.

Cedro hembra. See *Cedrela odorata*.

Cedro hembra colorado. See *Cedrela odorata*.

Cedro macho.

A tree from all parts of the island; height, 50 feet (15 meters); diameter, 55 to 60 inches (133 to 150 centimeters). Wood, blackish, hard; specific gravity, 0.878; used in building houses. (Exp. 1857.)

Ceiba pentandra. SILK COTTON. CEIBA.

PLATE XXIV.

Family, Bombacaceae; a high tree, 20 to 30 meters, with spreading, horizontal branches. The cotton covering the seeds is used in making beds and pillows. The wood is white and soft, but in Porto Rico is considered desirable for minor building purposes, such as inside partitions. The leaves are palmately divided, like those of the horse-chestnut, and the tree is sometimes called the "five-leaved silk cotton," to distinguish it from *Ochroma* and others which produce a silky fiber. The name "Bombay ceiba" is also used, perhaps as a corruption of *Bombax*, formerly the generic name.

The ceiba tree is much more common on the south side of the island, but nearly all have had the larger branches twisted off by the hurricane, the wood being soft and brittle. The trunks of young trees of this species are beset with large, conical spines, but in age these fall away and are to be found only on the branches. The trunk is further transformed by the growth of prominent wings or buttresses, sometimes 2 feet (0.6 meter) wide, while but a few inches thick. In west Africa, where this tree is larger and more luxuriant, pieces of these supporting wings are sawed out and used as doors of native houses.

A specimen near Ponce measured 36 meters in circumference at 4 feet from the ground, by following the sinuosities of the trunk.

Grosourdy (2:375) says of the wood:

"It is white, soft, very light, but rather strong and resists water moderately well, for which reason it is used to make canoes, although for this purpose it is not as durable as is generally supposed. The bottoms of such boats are sometimes coated with the fat from turtles to render them more durable, but when so treated the bottom is likely to crumble suddenly. The wood is also used to make tubs and basins. Specific gravity, 0.520."

The fiber surrounding the seeds of this and related species is the "kapok" of commerce, and is exported in considerable quantities from the west coast of Africa.

Ceibilla.

A tree from all parts of the island; height, 40 to 45 feet (13 to 14 meters); diameter, 40 to 45 inches (100 to 115 centimeters). Wood, white, soft; specific gravity, 0.642; not used. (Exp. 1857.)

Cenizo. See *Chenopodium album*, *Fagara martinicensis*, and *Tetrazygia claeagnoides*.

A tree from all parts of the island; height, 90 to 100 feet (27 to 30 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood, yellowish, soft; specific gravity, 0.741; not used. (Exp. 1857.)

According to Captain Hansard the specific gravity is 0.74 and the wood was formerly used for hogsheds.

Under this name Grosourdy refers to a number of species of *Zanthoxylum* and describes it as a large, dense tree, one kind, "massa de hercules," reaching 15 to 20 meters in height, with a straight, rather long, spiny trunk from 24 to 30 inches (60 to 75 centimeters) in diameter. The wood is light-colored and very handsome. It is fibrous in texture, rather hard, light, and strong. Used in the con-

struction of country houses. Specific gravity, 0.380. The bark in some species is spineless. (Grosourdy, 2: 363.)

Centella asiatica. YERBA DE CLAVO.

An herbaceous annual, growing in moist places. (Stahl, 4: 194.)

Cephaelis triplocephala. TRESCAPEZAS.

Family Rubiaceae; an herb. (Stahl, 5: 80.)

Cera. See *Sapium sebiferum*.

Cerasus occidentalis. See *Prunus occidentalis*.

Cereipa.

Under this name flat, oily seeds with a reticulated shell, said to be useful in the form of a liniment made with rum, and a bark used as an ingredient of a refreshing drink, were sold in the market of Ponce.

Cerero. See *Myrica cerifera*.

Cereus.

PLATE XXV.

A large genus belonging to the family Cactaceae.

Numerous species occur in Porto Rico, particularly about Coamo and in the vicinity of Guayanilla. One species collected near Guayama bore large edible fruits with a delicate and very refreshing flavor. This species might well deserve cultivation.

Cereus grandiflorus. REINA DE LAS FLORES.

Family Cactaceae; a fleshy plant, with large, fragrant, white flowers, found in all the Antilles. (Stahl, 4: 188.)

Cereus quadricostatus.

From vicinity of Guayanilla.

Cereus swartzii. SEBUCAN.

A fleshy plant, 3 meters high, found on the southern coast of the island. (Stahl, 4: 187.)

Cereus triangularis. PITAJAYA.

A fleshy plant, 1 to 3 meters high, found on rocks and decayed trees. (Stahl, 4: 187.)

Cerezas. See *Cordia collococca*, *C. nitida*, and *Malpighia coccigera*.

Cerezas blancas. See *Cordia alba*.

Cerezas cimarronas. See *Cordia elliptica*.

Cerezo.

"Cherry tree;" according to Captain Hansard the specific gravity is 0.61.

Cerezo comun, cerezo de la tierra. See *Cicca disticha*.

Cerillo.

A specimen (no. 22) not yet identified was obtained at Lecheria under this name, not previously recorded for a Porto Rican plant. In Cuba the name is said to be used for *Ecostemma caribaeum*, a rubiaceous tree, also called "managua de costa," for which "cuero de sapo" is given by Stahl. "Cerillo de costa" and "cerillo de loma" are also in use in Cuba for other plants.

Cestrum laurifolium. GALAN DEL MONTE.


Family Solanaceae; a shrub 3 to 4 meters high, found in waste places. (Stahl, 4: 141.)

Cestrum nocturnum. DAMA DE NOCHE.

Family Solanaceae; an ornamental shrub cultivated in gardens. It has white flowers, very fragrant at night. (Stahl, 4: 140.)

Chamaecrista diphylla. SEN DE DOS HOJAS.

A low shrubby herb differing from the other species of this genus and *Cassia* in having two obovate leaflets and very large stipules. (Stahl, 3: 116, as *Cassia diphylla*.)



STEM AND EDIBLE FRUIT OF *CEREUS* SP. AT GUAYAMA.

Chamaecrista glandulosa. TAMARINDILLO.

A rather small species, with large flowers similar to *C. fasciculata*, of the eastern United States. The leaves are much like those of the tamarind, whence the common name. (Stahl, 3: 117, as *Cassia glandulosa*.)

Chamaecrista portoricensis. RETAMA.

The variety *callosa* is said to be called "retama," and the variety *granulata* "escobilla," but it is not probable that any such precision actually exists. The plant is a small shrub from the limestone hills about Guayanilla, Guanica, and Cabo Rojo. (Urban. Symb. 1: 317, under *Cassia*.)

Chamico blanco. See *Datura metel*.

Chamico morado. See *Datura stramonium*.

Chamisco. See *Datura stramonium*.

Chamiso. See *Dodonaea viscosa*.

Chaptalia nutans. BRETONICA.

An annual composite pubescent herb growing in ravines and steep places, preferably moist and sandy. (Stahl, 5: 152, as *Leria nutans*.)

Chayote. See *Sechium edule*.

Chenopodium album. PIGWEED. CENIZO.

Family Chenopodiaceae; the familiar pigweed of our gardens.

Chenopodium ambrosioides. BASOTE.

Collected at Cataño. (No. 991.)

Cherimolia. See *Anona squamosa*.

Cherimoyer. See *Anona squamosa*.

Cherry.

The genuine cherry does not thrive in the Tropics, but in Porto Rico the Spanish name for cherries, "cerezas," is applied to the fruits of species of *Cordia* and *Malpighia*.

Cherry, Jamaica. See *Ficus pedunculata*.

Chiagari. See *Cuphea micrantha*.

Chicarron. See *Comocladia*.

Chicharo.

A name applied to a considerable variety of smaller leguminous vines.

Chicharron. See *Reynosia uncinata*.

Chicoria. See *Sonchus oleraceus* and *S. oleraceus*.

Chicoria azul. See *Lactuca intybaea*.

Chicoria cimarrona. See *Lobelia assurgens*.

China. See *Citrus aurantium*.

China dulce. See *Citrus aurantium*.

Chino.

A tree from all parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood yellow, hard; specific gravity, 0.873; used in making furniture. Also called "caro," and presumably not the orange tree. (Exp. 1857.)

Chiococca racemosa. BEJUCO DE BERAC.

Family Rubiaceae; a shrub 2 to 3 meters high, biennial; found in waste places and at bases of mountains. (Stahl, 5: 62.)

Chione glabra. MARTIN AVILA.

Family Rubiaceae; a tree 10 meters high; used for lumber. (Stahl, 5: 57.)

Chirimoya. See *Anona squamosa*.

Chirivita. See *Erigeron spathulatus*.

Chlorophora tinctoria. FUSTIC.

Family Moraceae; a tree reported from Guanica. A variety *xanthoxylon* is also known from Bayamon. The yellow wood furnishes a yellow dye; also used for brown and green shades. Fustic to the value of \$121,665 was imported in 1899, according to the Treasury reports.

A tree of slow growth that reaches a height of 50 feet (15 meters), and with a trunk $1\frac{1}{2}$ to 2 feet (0.5 to 0.6 meter) in diameter. Furnishes an elegant wood of a handsome yellow or orange color, with time and exposure to the air turning slowly lighter until it becomes a dirty yellow, which is improved by varnishing when used for furniture. Specific gravity, 0.711. In the countries where it grows the wood is used for shelving in the country houses, for the spokes of carriage and wagon wheels, and wherever strength and resistance are necessary. (Grosourdy, 2: 383.)

The native name "mora" is given by this author.

Chocho. See *Sechium edule*.

Chocolate. See *Theobroma cacao*.

Chrysobalanus icaco. ICACO.

PLATE XXVI.

Family Rosaceae; shrub 2 meters high, found on dry beaches; known as "cocoaplum." Fruits about the size of a plum, used for preserves. The variety *pello-carpus* is reported from Manati. Bello spells the common name "hicaco." (Stahl, 4: 64.)

Chrysophyllum cainito. STAR APPLE. CAINITO.

Family Sapotaceae; a tree 10 to 15 meters high, with fruit about the size of an apple; edible, and said to be very constipating when eaten with orange; Bello and others sometimes write the common name "cainito." (Stahl, 6: 45.)

Very similar to the species referred to under "cainitillo," which should probably be referred to this species. The wood is darker, violet-colored, and rather coarse. Not used, but thought by Grosourdy to be suitable to the same purposes as cainitillo. Specific gravity, 0.889. (Grosourdy, 2: 369.)

Chrysophyllum glabrum. LECHECILLO.

A tree or shrub 4 to 5 meters high. (Stahl, 6: 48.)

Chrysophyllum krugii.

A recently described species collected by Sintenis between Coamo and Aibonito.

Chrysophyllum monopyrenum. CAIMITILLO.

A tree 3 to 5 meters high, bearing an edible fruit called in Jamaica "damson plum." Bello reports the common name "teta de burro" for *C. oliviforme*. (Stahl, 6: 46.)

Furnishes a very strong and solid flesh-colored wood that is not used except to make charcoal for forges, for which it is highly valued. (Grosourdy, 2: 415.)

Chrysophyllum oliviforme. See *Chrysophyllum monopyrenum*.

Chupa gallo. See *Cinnamodendron macranthum*.

Chupagallo.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 19 to 20 inches (47 to 50 centimeters). Wood light-colored, hard; specific gravity, 0.980; used in building houses. Also known as "caro." (Exp. 1857.)

Cicca disticha. CEREZO COMUN.

A tree 21 to 24 feet in height, with a trunk 6 to 8 inches (15 to 20 centimeters) in diameter. Furnishes, after polishing, a very pretty wood, having as yet no application. The wood is of ordinary weight, fibrous in texture, rather hard, fine-



ICACO ('CHRYSOBALANUS ICACO').

grained, but not very lasting. Specific gravity, 0.577. Also called "cerezo de la tierra." (Grosourdy, 2: 376.)

Cicca disticha is given as a synonym of *Phyllanthus distichus*, an Indian species.

Cicer arietinum. GARBANZO.

The extensive use of garbanzos or chick-peas by the Porto Ricans, and doubtless also by other Spanish-Americans, suggests the desirability of cultivating them in the United States for export to these tropical countries. Moreover, they are undoubtedly a palatable and nutritious dish which, if properly introduced, should find a place on the American table.

We could not learn that any "garbanzos" were grown in the island or that any experiments in this direction were ever tried.

Cidron. See *Lippia nodiflora*.

Cieneguillo. See *Daphnopsis philippiana*, *Eugenia confusa*, *Comidesia Lindeniana* and *Myrcia deflexa*.

Cienigilla.

A tree with very hard white wood; grows to no great size; similar to ash; not uncommon in the mountains. (Hansard.)

Cieniguillo. See *Myrcia ferruginea* and *M. deflexa*.

Doubtless the same name as cieneguillo.

Cinchona. QUININE.

Experiments have been tried with cinchona in Jamaica, but the market is now well supplied by Java and the British West Indies, so that planting in Porto Rico does not seem advisable from the commercial standpoint.

Cinnamodendron macranthum. CHUPAGALLO.

A tree belonging to the family Canellaceae; 10 meters high, with greenish white, orange-perfumed flowers. From Sierra de Luquillo. (Urban, Add. 1: 9.)

Cinnamomum camphora. CAMPHOR TREE.

The true camphor tree of Japan and China has been successfully planted in California and in the warmer parts of our Gulf States, since it is able to withstand light frosts. It makes a beautiful and in every way desirable shade tree in addition to its value for the extraction of camphor, for which some have calculated that its culture would be a paying business. It seems entirely probable that the camphor tree would grow well in Porto Rico, and the conditions are probably favorable for the production of the gum, in which respect the culture of camphor in North Africa has failed, perhaps owing to the too great heat and dryness of the climate.

The present species is not the only one yielding camphor; *Blumea balsamifera*, a composite herb native in China, and *Dryobalanops aromatica*, a large tree of the family Dipterocarpaceae, native in Borneo and neighboring islands, also yield substances closely similar to camphor, and even more highly valued.

Cinnamomum zeylanicum. CINNAMON.

The cinnamon tree is native in Ceylon and other parts of the East Indies, and has been introduced into the American tropics, though not known to have reached Porto Rico. It might thrive in a few situations on the south side of the island, but a dry, sandy soil is considered necessary for the production of bark of good quality, and Porto Rico has little land of that kind. *Cinnamomum cassia* and other species of the genus are more hardy and produce commercial barks used as substitutes and adulterants of the true cinnamon, to which they are inferior.

Cinnamon. See *Cinnamomum*.

Cinnamon, wild. See *Canella alba* and *Anomis caryophyllata*.

Cionandra racemosa. See *Cayaponia racemosa*.

Cipil.

A tree from the interior of the island; height 45 to 50 feet (13 to 15 meters); diameter 12 to 15 inches (30 to 37 centimeters). Wood white, very hard; specific gravity, 0.960; used in boat building. Also called "pino." (Exp. 1857.)

Ciralillo. See *Meliosma obtusifolia*.

Cirila. See *Cyrilla antillana*.

Ciruela del país. See *Spondias purpurea*.

Cissampelos microcarpa. BEJUCO DE MONA.

Reported by Bello from the western part of the island.

Cissampelos pareira. BEJUCO DE MONA.

Family Menispermaceae; an annual herb. The "pareira brava" of the pharmacopœia is a product of this plant; said to be an antidote for poison, and valuable in medicine. (Stahl, 2: 31.)

Cissus acida. See *Vitis acida*.

Cissus sicyoides. See *Vitis sicyoides*.

Cissus trifoliata. See *Vitis trifoliata*.

Citharexylum quadrangulare. PÉNDULO COLORADO. PLATE XXVII.

Family Verbenaceae; a tree 20 to 60 feet (6 to 18 meters) high, with red wood; useful for furniture and building. Found in mountain districts. Sinteris' specimen from Cayey (no. 3321) has the leaves broader and more nearly elliptical and may be distinct from his 3723. Bello gives the common names "pendola" and "palo de guitarra." (Stahl, 6: 211.)

Grosourdy (2: 409), under the native name "péndula," describes this species as a wild tree reaching 50 feet (15 meters) in height, with a moderately straight trunk 18 inches (45 centimeters) in diameter. Furnishes a rather strong, straw-colored wood that breaks with an oblique fracture. It is used for beams, windows, doors, etc. The natives make their guitars of this wood. Specific gravity, 0.753 and 0.722.

Citron. See *Citrus medica*.

Citrullus vulgaris. WATERMELON. SANDÍA. PATILLA.

Watermelons seldom reach marketable quality in humid tropical countries. A few were seen in the market of San Juan, but none of them would have been considered fit to eat in the United States. Watermelon growing is scarcely to be recommended as a profitable business in Porto Rico, though with properly selected conditions and careful cultivation fairly good fruit can probably be raised, as reported by one of the Department's correspondents. Muskmelons will probably succeed somewhat better.

Citrus.

Grosourdy (2: 394) says of the lemon and orange that they furnish a wood very strong, flexible, compact, and of good durability and resistance. It is, however, more an object of curiosity than utility, although it is used for moldings and inlaid work of fine furniture. The military authorities of San Juan use it for the handles of tools. He gives the specific gravity of lemon as 0.833 and that of the orange as 0.815.

Citrus aurantium. SWEET ORANGE. CHINA.

"China" is the name under which the common sweet orange is known in Porto Rico, the Spanish name for the orange, "naranja," being applied to the sour orange.

This fruit may be distinguished from the sour orange not only by the taste but by the smoother and more even surface, the thinner and more closely fitting rind, and the solid center.

Until recently the orange in Porto Rico has been propagated only by seeds, and



PÉNDULO COLORADO · CITHAREXYLUM QUADRANGULARE . ROAD BETWEEN PONCE AND
ADJUNTAS.

as might be expected under such circumstances the fruit is anything but uniform in character. The majority of those offered in the markets are of such low grade that they would hardly have been placed before a public accustomed to a fruit of good quality. As a rule, the fruits are very juicy but lack texture and flavor. Specimens of a high-grade orange were, however, secured at San Juan, and fruit specialists at Washington were of the opinion that this variety might even profitably be used for propagation by budding.

According to Captain Hansard, a weed called "cojitre" placed around the orange tree sweetens the fruit. This is believed to be a leguminous plant and suggests the possibility of its use as a substitute for the nitrogenous manure which is said to be necessary for the best results with this fruit.

Ten years are thought to be necessary for a tree to reach full bearing. It is claimed that a single tree will sometimes produce several thousand oranges.

"Naranja china" and "china dulce" are names sometimes applied to the sweet orange.

Citrus bigaradia. SOUR ORANGE. NARANJA.

The most common of the citrous fruits is perhaps the sour orange, or "naranja," as it is universally called in Porto Rico. The ordinary Porto Rican sour orange is believed to belong to this species, and is not the Seville orange, as has been reported.

In Florida the sour orange is used for stocks in all plantings on moist land, for the reason that it resists the foot rot, or *mal di goma*, and other diseases which affect other varieties when planted on heavy soils.

Citrus decumana. POMELO. TORONJA.

Pomelo seems now to be the name preferred for the fruit otherwise known as grape fruit. The shaddock is thought to be a slightly different horticultural variety.

Pomeloes were seen growing near Bayamon, but as the fruit was immature the quality could not be judged. Although not so sensitive to frosts as the lime or lemon, the pomelo requires tropical conditions for properly maturing the fruit, and the increasing popularity of this fruit will warrant careful experiments in Porto Rico.

Citrus hystrix acida. LIME. LIMA.

The lime is referred by different authors to both *C. medica* and *C. limetta*, as well as to the present species.

The limes of Porto Rico seem to be quite normal and of good quality.

The lime endures shipment well if properly packed, and as the quality of the Porto Rican fruit is good it is perhaps the most eligible fruit with which to open a trade.

In the British island of Montserrat the production of limes for the extraction of lime juice has been made quite an industry. About 1,400 acres are planted to limes, over 1,200 belonging to a single estate. The annual total of income from the industry averages over \$30,000, and other islands contribute large quantities.

Citrus limonum. LEMON. LIMON.

The Porto Rican lemon seems to be somewhat intermediate between the genuine lemon and the rough lemon of Florida and Jamaica.

Commercially it can not be expected to compete in the market with the ordinary lemon, but for eating and possibly for the preparation of unsweetened lemonade, it might become popular with those who are partial to acid fruits.

Citrus medica. CITRON. TORONJA.

To this species are referred by some authors not only the citron, but the lemon, lime, bergamot orange, and the sweet lemon, or "lemon dulce."

A specimen of citron was secured from a street vender in San Juan. He stated that considerable quantities are grown in the vicinity of Trujillo Alto, but there was no opportunity of investigating that locality. Large amounts of citron peel are imported into the United States from Mediterranean countries, notably from Corsica, and experiments in the culture of the trees are being made in Florida and California. The possibility of success in Porto Rico is worthy of investigation. Although none of the citrons now in commerce are grown in tropical countries, there are said to be varieties of excellent quality in Hindostan and the Malay Peninsula. The Porto Rican specimen observed was of medium size, but the skin was too thin and too loose in texture and apparently deficient in flavor. Some or all of these defects may, of course, be due to the absence of proper culture.

Citrus vulgaris. See *Citrus aurantium*.

Clavel. See *Cosmos caudatus*.

Clavelillo. See *Stemmodontia caruosa*.

At Toa Alto and Cataño this name was also used for a species of Spermaceae not yet identified.

Clavellina. See *Poinciana pulcherrima*.

Clavelon de playa. See *Borrchia arborescens*.

Clematis dioica. CABELLOS DE ANGEL.

Family Ranunculaceae; a shrubby climber, found in mountains, flowering in December. Called "virgin's bower," "rudding withe," and "traveler's joy" in Jamaica. From vicinity of Lares. (Stahl, 2: 19.)

Cleome pentaphylla. See *Gynandropsis pentaphylla*.

Cleome pungens. See *Cleome spinosa*.

Cleome speciosa. See *Gynandropsis speciosa*.

Cleome spinosa. VOLATINES PUNZANTES. SAMBO.

Family Capparidaceae; an annual thorny herb, 1 meter high; thrives in a dry soil. Specimens were obtained at Coamo Springs (no. 720), where the name "aromo" is used. Bello reports the name "jasmin del río" for *Cleome pungens*. (Stahl, 2: 41.)

Clerodendron aculeatum. BOTON DE ORO.

Family Verbenaceae; a shrub, 5 to 12 feet (1.5 to 3.5 meters) high. From Fajardo. (Stahl, 6: 218.) Reported by Bello.

Clerodendron fragrans. FLOR DE MUERTO.

A biennial shrub, 1 meter high. Bello reports the common name "jasmin hediondo." (Stahl, 6: 219.)

Cleyera albopunctata.

Family Theaceae; from vicinity of Maricas.

Clidemia hirta. CAMACEY PELUDO.

Family Melastomaceae; a common shrub, 3 to 8 feet (1 to 2.5 meters) high. (Stahl, 4: 95.)

Clidemia neglecta. CAMACEY NEGLIGENTE.

A shrub, 3 to 8 feet (1 to 2.5 meters) high. (Stahl, 4: 97.)

Clitoria arborescens. CONCHITA ESBELTA.

Family Leguminosae; a high climber, growing in dry places. (Stahl, 3: 59.)

Clitoria glycinoides. CONCHITA BLANCA.

An annual wild vine, growing in all parts of the island. According to Stahl the flowers are white, and according to Grisebach purple-variegated. Our specimens (nos. 299 and 373) were noted in life to be creamy white, with purplish lines on the banner; on drying they became distinctly yellow. (Stahl, 3: 58.)



CUPEY (CLUSIA ROSEA). NEAR YAUCO.

Clitoria ternatea. BEJUCO DE CONCHITAS.

A woody annual vine, with handsome blue and white flowers; a native of the East Indies, cultivated in gardens as an ornamental, and frequently escaped in waste places, including dry rocky hillsides. Our specimens were collected near Jobo. (Stahl, 3: 56.)

Clove. See *Eugenia aromatica*.

Clusia acuminata. CUPEILLO.

Family Guttiferae. (Stahl, 2: 124.)

Clusia cartilaginosa.

From Cayey; Adjuntas.

Clusia gundlachi. CUPEY DE ALTURA.

A shrub, 3 to 5 meters high; grows above 1,000 feet (300 meters) altitude. The resin is used for rheumatism. From Mariacao and Luquillo. (Stahl, 2: 122.)

Clusia krigiana. CUPEY.

A shrub or small tree of 3 to 15 meters, reported from Sierra de Luquillo (El Yunque) and from the vicinity of Utuado.

Clusia rosea. CUPEY.

PLATE XXVIII.

A wild tree or shrub 3 to 10 meters high. The resin is useful in rheumatism. (Stahl, 2: 121.)

A wild tree that grows usually as a parasite on other trees, which are often killed by it. The roots spread downward over the trunk until the ground is reached, when other roots are formed, and the tree finally comes to be supported by its own trunk. When grown entirely in the ground, the trunk will not be so tall but thicker and denser. The diameter is from 18 to 24 inches (45 to 60 centimeters), and it sometimes reaches a height of 30 or 40 feet (9 to 12 meters). The wood is reddish and is used only as fuel. Specific gravity, 0.876—(Grosourdy, 2: 378). Sometimes called "balsam fig" or "balsam tree." "L. cat. de ..."

Cnicus portoricensis. See *Carduus mexicanus*.

Cobano. See *Stahlia maritima*.

Cóbana negra.

Makes pretty walking sticks, perfectly black, very similar to ebony, a smoky but not a jet black; takes a high polish.

Cobano.

A tree from the eastern parts of the island; height 50 to 60 feet (15 to 18 meters), diameter 25 to 30 inches (60 to 75 centimeters). Wood very dark and very hard; specific gravity, 1.227; used in boat building. (Exp. 1857.) Said by Stahl to be the same as "cojobana," a name which he applies to *Pithecolobium filicifolium* and to *Piptadenia peregrina*.

Coca. See *Erythroxylon coca*.

Cocaine. See *Erythroxylon coca*.

Coccocypselum repens. BALSAMILLO.

Family Rubiaceae; a creeping tufted annual; stems 18 to 20 inches (45 to 50 centimeters) long, growing in mountains. Bello records the common name "yerba de guava." (Stahl, 5: 45.)

Coccolobis.

A large West Indian genus of polygonaceous shrubs and trees, many of which bear edible fruits known as "sea grapes" or "uvas del mar," although other more specialized names are used in Porto Rico where the genus is represented by numerous species.

Coccolobis diversifolia. PALO BOBO.

A shrub of 15 feet (5 meters) or less, known throughout the West Indies.

Coccolobis excoriata.

This tree attains a considerable size and is looked upon as a fine timber wood; grows in mountains. (Sunan.)

Coccolobis grandifolia. MORALON.

A large tree of the mountain forests.

Coccolobis kunthiana.

From Utuado.

Coccolobis laurifolia.

Native names: "Uvillo," "glateado," and "uverillo." See note under *Coccolobis*. From Cabo Rojo.

Coccolobis nivea. CALAMBREÑAS.

A branching tree 20 to 30 feet (6 to 9 meters) in height, with papery leaves, yellowish flowers, and small white edible berries. Specimens were obtained at Coamo Springs (no. 737) and near Ponce (no. 805), the latter with red berries.

Coccolobis obtusifolia. UVILLO.

A shrub of 10 to 12 feet (3 to 4 meters), known from the Bahamas, Santo Domingo, Porto Rico, and the Danish Islands.

Coccolobis pirifolia.

From Bayamon.

Coccolobis rugosa. ORTEGON.

A species with deeply wrinkled leaves 18 inches (45 centimeters) across; collected by Sintenis at Cayey. Also reported by Eggers from the region of El Yunque as *C. macrophylla*. Said to be the most conspicuous tree in some regions, up to an altitude of 2,000 feet, especially when the purple flowering spikes, a yard long, are displayed. A large tree, said to be abundant in Porto Rico, and furnishing an economic wood.

Coccolobis uvifera. SEA GRAPE. UVERO.

Also called "uva del mar," and in Jamaica "pigeon wood." A tree of medium or large size growing near the sea. The berries are described by Grisebach as dark blue and $\frac{1}{2}$ to $\frac{3}{8}$ of an inch (12 to 16 millimeters) in length. The wood is esteemed in Jamaica for cabinet work. The plants are propagated by cuttings and the trees grow to a large size.

Coco. See *Cocos nucifera*.

Cocoa plum. See *Chrysobalanus icaco*.

Cocoe. See *Colocasia esculenta*.

Cocorrón. See *Elaeodendron xylocarpum*.

A tree reported by Grosourdy as occurring in the eastern part of the island, 30 or 35 feet (9 to 11 meters) high, with a trunk 1 foot or more (30 centimeters) in diameter. It furnishes a yellow wood of medium durability, having a vertical fracture, used in the construction of country houses. (Grosourdy, 2: 376.)

Cocorroncito. *Elaeodendron attenuatum*.

Cocos aculeata. See *Acrocomia aculeata*.

Cocos mauritia.

A name given by Hill for the "palma moriche." There is no such name or synonym in the Index Kewensis, and the reference is doubtless to *Mauritia flexuosa*, which is also given by Hill as the botanical name of "moriche."

Cocos nucifera.

The rather extensive use of green coconuts for drinking purposes in Porto Rico suggests the question whether the extraction of fiber from the husks might not be undertaken to advantage. This fiber is a regular and important article of export from the East Indies to Europe, under the name of coir, and is used in

the manufacture of cordage, bags, brooms, brushes, nets, matting, and other coarse fabrics, and for numerous minor purposes. In some parts of India unripe cocoanuts are used for this purpose, the fiber being finer and more flexible than that of the mature nuts. For the extraction of the fiber the husks are split open upon a sharp stake or crowbar and then soaked in tanks or pits dug in the banks of tidal rivers. The labor of beating out the fibers from the rotted husks is considerable, and the people of Porto Rico could hardly have competed in this with the Hindoos, particularly since the latter have all the experience, but successful machinery has been invented and is now coming into general use, so that when power plants have been established in Porto Rico the possibilities of the extraction of coir will be worthy of further investigation. Very little coir is imported into the United States in the unmanufactured condition, so that plans for any extensive trade must have reference to the European market.

The other important cocoanut export of India and the South Sea Islands is copra, the dried meat of the nut. It is imported into Europe and America for the sake of the cocoanut oil, which is extracted from it to the amount of 30 to 60 per cent of its weight, depending upon quality and the degree of dryness. The Treasury reports for 1899 show importations of:

Copra	2,071,037 pounds, valued at \$109,421
Cocoanut oil.....	21,794,633 pounds, valued at \$1,034,710

Something over another million dollars was paid for cocoanuts in the shell and in other forms.

Cocoanut oil is used extensively for food, in the Far East as a substitute for butter, lard, or olive oil, and is also burned for illumination. In Europe and the United States it is employed mostly in the manufacture of candles and soaps, but in the latter form it has the disadvantage of having a disagreeable smell. At cool temperatures, 40° to 60°, it becomes solid, but in the Tropics it remains fluid. The "cake," left after the oil has been extracted, is used quite after the manner of cottonseed. Copra is used much more extensively in Europe than in this country, but recent statistics are not available. The trade is, however, known to have suffered in recent years from the competition of the African palm oil—secured from *Elæis guineensis*, which is superior to cocoanut oil for the purpose of soap making—and has probably been further affected by our cottonseed oil.

As a minor local use of the cocoanut, may be mentioned the employment of the roots as one of the ingredients of the numerous refreshing drinks so popular in Porto Rico.

Grosourdy (2: 495) states that the wood is used in the same manner as that of the royal palm, and some parts are more highly prized, being harder, handsomer, and more suitable for fine work: specific gravity 0.501 and 0.865.

Cocotero. See *Cocos nucifera*.

This is the name for the cocoanut tree, but it is little used, both the tree and the nut being commonly called coco.

Coco wood. See *Inga vera*.

Coffea arabica. COFFEE. CAFÉ.

Coffee is the most important product of Porto Rico, and its culture is too extensive a subject to be more than touched on here. The question of shade has been made the subject of a special publication. (See Bulletin No. 25, Division of Botany, United States Department of Agriculture.)

The most careless and wasteful methods are practiced in the culture of this important crop. No attention is paid to the selection of seedlings, most of the new plants being secured from seeds that have germinated under the trees in the old plantations. It is estimated that by proper methods of cultivation the yield from the land now devoted to coffee could be doubled or tripled.

Coffee is a very popular beverage on the island. As usually prepared it is very black, due, in part, to the custom of roasting the beans with sugar. This is sometimes added when the roasting is nearly completed and is supposed to coat the beans with a thin layer of burnt sugar, which helps to retain the aroma.

Coffee. See *Coffea arabica*.

Cohune nut. See *Attalea cohune*.

Coitre. See *Commelina*.

Coix lachryma-jobi. JOB'S TEARS. CAMANDULA. PLATE XXIX.

A grass with large oval, very hard seeds, smooth and polished, and of a light pearly-gray color; they are sometimes strung for beads or rosaries. This species is not uncommon in waste places in Porto Rico.

Cojoba. See *Copaifera hymenacifolia* and *Acacia berteriana*.

According to Captain Hansard cojoba wood is used for posts and fencing; specific gravity, 0.60.

A wild tree found sparingly in the eastern part of the island, 40 to 50 feet (12 to 15 meters) high, 2 feet (60 centimeters) in diameter; furnishes a wood that is flesh-colored, very hard, breaking vertically with great difficulty; used in boat building and for fences, in which case it strikes root; also called "palo de hierro," and "quebra hacha." (Grosourdy, 2: 376.)

Cojoba blanca.

A tree from the eastern part of the island; height, 50 to 60 feet (15 to 18 meters); diameter, 12 to 15 inches. Wood white, very hard; specific gravity, 0.928; used in boat building. (Exp. 1857.)

Cojoba negra.

A tree from the eastern part of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood dark, very hard; specific gravity, 0.994; used in boat building. (Exp. 1857.)

Cojobana. See *Piptadenia peregrina* and *Pithecolobium jilicifolium*.

Grosourdy (2: 376) refers this name and "cobana" to *Swietenia mahagouii*.

Cojobilla. See *Piptadenia peregrina*.

Bello reports this name under *Acacia angustiloba*.

Cola. See *Cola acuminata*.

Cola acuminata. COLA.

The cola tree was introduced into the West Indies from West Africa during slave trading times for the benefit of the slaves. It has become naturalized in Trinidad and probably in other islands.

Coladilla. See *Polypodium polypodioides*.

Coleus amboinicus. See *Coleus aromaticus*.

Coleus aromaticus.

Family Labiatae. Bello reports the name "oregano de Espana" for *Coleus amboinicus*. (Stahl, 6: 170.)

Collards. See *Brassica oleracea*.

Colocasia esculenta. TARO. YAUTIA MALANGA. BLEEDING HEART.

Family Araceae; the "taro" of Polynesia.

Elsewhere about the Caribbean Sea the same plant has a most confusing variety of native names, "coco" in Jamaica, "eddo" in Barbados, "tannia" in Trinidad, "taya" in the French islands, "oto" on the Isthmus of Panama, and "tiquisquis" in Nicaragua. European travelers and residents are commonly not aware of the differences between these closely similar plants, and generally apply these names indiscriminately to both *Colocasia* and *Xanthosma*, and some botanists have even failed to appreciate the differences.



JOB'S TEARS (COIX LACHRYMA-JOBI).

The species of *Colocasia* can be distinguished from those of *Xanthosma* by the peltate leaves.

Mr. W. M. Armstrong, of Honolulu, states that he found the Porto Rican taro, or yautia, invariably inferior to the Hawaiian, the introduction of which he considers would be an important service for the agricultural population of Porto Rico. Stock should be secured from Hilo through the experiment station authorities.

Mr. Armstrong also states that the Hawaiian taro has been found superior to that of Fiji and Samoa by Mr. Alexander, who has investigated the subject. Its wholesome and delicate quality recommend it to resident Americans, and many who have returned to California now import taro for household use.

Recently the Chinese have taken to raising another variety of taro, or taro-like plant, in California, particularly at Bakersfield. This is said to be propagated from tuberous lateral shoots, not like the taro in which the leafy cut-off ends are planted. Chinese taro is also imported to California from Canton and Hongkong. This is by some thought to be somewhat inferior to the Hawaiian, but the difference is not great.

Coloquintilla. See *Cayaponia racemosa*.

Colubrina ferruginosa. SNAKE WOOD. ABELLUELLO.

Family Rhamnaceae; a tree 15 to 20 meters high, found in mountains. Stahl states that this species is also called "quitaran." (Stahl, 4: 45.)

Colubrina reclinata. MABI.

A shrub or low tree found in the mountains; 15 to 20 meters high. From near Guanica. (Stahl, 4: 46.)

Called "palo mabi" by Grosourdy (2: 366), who says the wood is strong and durable and is used in Cuba for building. Specific gravity, 0.947.

Both this and the following species are said to be used to make the drink called "mabi." (*Ceanothus reclinatus*.)

Columnea hispida. See *Columnea tulae*.

Columnea tulae. TIBEY PARASITO.

Family Gesneriaceae: an annual 50 centimeters high, found in high mountains. Stahl's *C. hispida* is found by Urban not to be Swartz's species of that name and is redescribed as variety *flava* of *C. tulae*, the flowers being yellow instead of red. (Urban, Symb. 1: 410.)

Commelina. COITRE.

Several species of *Commelina* are common in Porto Rico and all seem to be called "coitre."

Comocladia.

A West Indian genus belonging to the family Anacardiaceae and having a poisonous juice similar to that of poison ivy and other species of *Rhus*. All the species are shrubs or trees with compound leaves and nearly all have the margins of the leaflets armed with pointed teeth into which the veins are produced.

This peculiarity, while not unique, will assist in the recognition of the species with which those who have occasion to visit West Indian forests will do well to become acquainted. In Cuba species of *Comocladia* are called "guao," in Porto Rico "carrasco," and in Santo Domingo "chicarron." In his last revision of the genus Professor Engler, of Berlin, recognized nine species. While several of these are represented in the National Herbarium, there are also several new forms, including three from Porto Rico, making ten species available for study.

Comocladia acuminata. CARRASCO.

A shrub 2 to 3 meters high; found at the base of the mountains; said to be poisonous.

Comocladia ilicifolia. CARRASCO.

A shrub 2 or 3 meters high found on the south coast of the island. (Stahl, 4: 59.)

Conchita blanca. See *Clitoria glycinoides*.

Conchita de Plumier. See *Centrosema plumieri*.

Conchita esbelta. See *Clitoria arborescens*.

Conchita laurifolia. See *Centrosema laurifolia*.

Conchita peluda. See *Centrosema pubescens*.

Conchita Virginia. See *Centrosema virginianum*.

Concombre. See *Cucumis anguria*.

This common name is applied at San Juan; at Ponce *Cucumis anguria* is called *pepino* or *pepinito*.

Condalia ferrea. See *Scuta ferrea*

Congo pea. See *Cajanus cajan*.

Congo.

A variety of banana. See *Musa*.

Conocarpus erectus. MANGLE BOTON.

Family Combretaceae; a shrub 5 meters high, growing in mangrove swamps. In Jamaica called "alder" or "button tree," and esteemed as fire wood. (Stahl, 4: 137.)

A wild tree abundant on the coasts, 20 to 25 feet (6 to 8 meters) high, 9 to 10 inches (22 to 35 centimeters) in diameter. It furnishes a wood very hard and heavy, fibrous in texture, fine-grained, and compact. When freshly worked it is gray tinted with light chocolate with narrow undulous light lines; after it has been worked for some time it becomes darker. For a long time the wood keeps the marshy odor of the swamps. Specific gravity, 1.009. Used in building boats and barges, also for shelving, being very durable when grown on dry ground. It is used also to make charcoal for forges. (Grosourdy, 2: 399.)

Conradia. See *Pentarrhaphia reticulata*.

Conradia reticulata. See *Pentarrhaphia reticulata*.

Contrayerva. See *Aristolochia*.

Convolvulus jamaicensis. AGUINALDO BLANCA DE COSTA.

Family Convolvulaceae; a wild vine growing in savannas and along the coast. (Stahl, 6: 169.)

Convolvulus nodiflorus. AGUINALDO BLANCO.

A creeping plant, found in waste places on the south coast of the island. (Stahl, 6: 170.)

Convolvulus pentanthus. AGUINALDO AZUL.

Wild in waste places and arid regions. (Stahl, 6: 168.)

Copaifera hymenaeifolia. COJOBA.

Baron Eggers expresses in the Kew Bulletin for 1890 the opinion that this species is confined to Cuba. It is enumerated by Hill among Porto Rican forest trees.

Copaifera officinalis. PALO DE ACEITE.

A wild tree, rather large and dense, which at first sight slightly resembles the walnut of Europe. It is 40 to 50 feet (12 to 15 meters) high with a rather straight trunk 12 to 18 inches (30 to 45 centimeters) in diameter. It furnishes a good wood, rather hard, very handsome, of good weight, fibrous in texture, but rather compact and fine-grained. The general color is brownish or sometimes about the color of dead leaves beautified with black markings and resembling walnut. The

shavings have the odor of copaiba balsam. The wood is used for building and general carpenter work, for making wheels of carts and carriages, for turning and for adorning the interior of houses, also for furniture. Specific gravity 0.777. (Grosourdy, 2: 378.)

Copaiva. See *Copaifera*.

Copillo. See *Cordia globosa*.

Coral.

A tree from the western parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood pinkish, hard; specific gravity 1.247; used in house building. (Exp. 1857.)

Coral bean. See *Erythrina corallodendron*.

Coralillo. See *Antigonon leptopus*.

Coralillo rosado. See *Antigonon leptopus*.

Coralitos. See *Adenantha paronina*.

Corazon. See *Anona reticulata*.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 12 inches (30 centimeters). Wood yellow, soft; specific gravity, 0.650; little used. (Exp. 1857.)

According to Captain Hansard this is the "bull's heart" or "sweet sop" of the British Islands. The specific gravity of the wood is 0.65. According to Grisebach the name "sweet sop" is applied both to *Anona squamosa* and *A. sericea*, while in the Treasury of Botany the "bullock's heart" of the West Indies is said to be *A. reticulata*.

Corazon cimarron. See *Anona palustris* and *Rollinia multiflora*.

Corcho. See *Anona palustris*.

A cork tree, with very soft wood and umbrella-like leaves; very rare.

Corcho blanco.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood white, very soft; specific gravity, 0.555; not used. (Exp. 1857.)

Corcho prieto.

A tree from all parts of the islands; height, 30 to 35 feet (9 to 11 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood light, soft; specific gravity, 0.558; not used. (Exp. 1857.)

Corchorus hirsutus. MALVA TÉ DE LA PLAYA.

Family Tiliaceae: a shrub 2 to 6 feet (0.6 to 2 meters) high, found along the sea-coast. A specimen collected at Santurce (no. 293), and there called "cadillo," agrees fairly well with a sheet from St. Croix in the National Herbarium, but is much less hirsute than material from other parts of the West Indies and South America. A variety, *fortipes*, was obtained by Sintenis at Aguadilla, who also secured the typical form at Guanica. (Stahl, 2: 117.)

Corchorus hirtus. MALVA TÉ.

A woody herb. (Stahl, 2: 115.)

Corchorus siliquosus. MALVA TÉ. BROOMWEED.

A woody shrub, 1 to 3 feet (0.3 to 1 meter) high, frequenting stony and sandy places; used by the natives of the West Indies for making brooms; the leaves used for tea. (Stahl, 2: 116.)

Cordia.

A large genus of boraginaceous trees yielding medicinal drugs, edible fruits, and valuable timber. In the characters of habit, leaves, and inflorescence the species of *Cordia* are extremely diverse, and there also appears to be considerable confusion as to the identity of closely allied species.

Cordia alba. CEREZAS BLANCAS.

Family Boraginaceae. According to Grisebach this is a tree 30 feet (9 meters) high. Stahl says that it is a shrub 9 to 12 feet (3 to 4 meters) high with slender branches trailing over the surrounding vegetation. From Peñuelas.

Grosourdy gives the native name "capa blanco" and agrees with Grisebach in the dimensions, describing it as a tree 30 to 45 feet (9 to 14 meters) high with a rather dense top. The trunk is moderately long, about 3 feet (1 meter) in diameter, and rather crooked. The wood is very hard and strong, dark yellow in color, breaking with an oblique fracture. It is very lasting and highly esteemed. Extensively employed by the military authorities of San Juan for gun carriages, hubs of wheels, carpenter benches, vises, etc. It is also used in making fine furniture and in the construction of buildings and for farm utensils. (Grosourdy, 2: 370.)

Cordia bellonis.

A shrub from the vicinity of Maricao. (Urban, Symb. 1: 393.)

Cordia borinquensis. MUÑECA.

Also called "palo de Muñeca." A tree attaining 20 meters, recently described as a new species; reported from numerous localities. (Urban, 1: 390.) From Isabon, Sierra de Nagnabo, and near Utuado.

Cordia collococca. CEREZAS.

A tree 5 meters high, with white bark; wood soft, brittle, and little used; found in all parts of the island except the central part and the north coast. In Jamaica this tree is said to reach a height of 50 feet (15 meters). The fruit is eaten by fowls. (Stahl, 6: 271.)

Cordia elliptica. CEREZAS CIMARRONAS.

A forest tree 30 feet (9 meters) high; more abundant on the south side of the island. (Stahl, 6: 93.)

Cordia gerascanthus. CAPÁ PRIETA.

A tree 30 to 60 feet (9 to 18 meters) high; wood used for furniture; found in mountainous regions. In Jamaica the wood of this tree is called "prince wood" and is considered one of the best woods. The small saplings make good hoops for hogsheads. It resembles the elm and is called Spanish elm. (Stahl, 6: 91.)

Grosourdy gives the native name "palo de rosa," and reports it also from Vieques.

A wild tree 30 feet high with a trunk 18 inches in diameter. Furnishes a rose-colored wood that breaks with a vertical fracture. Employed in the construction of crude houses and in building boats. It is also much appreciated for making furniture. The name "palo de rosa" has been applied to various different woods taken from South America to Europe, one of which is the "palissandre" of the French; another (*Amyris balsamifera*), also called rosewood, candle wood, and sweetwood, is very little used. The English rosewood is *Amyris montana*, which is also called yellow candle and yellow sander in Jamaica. (Grosourdy, 2: 407.) On page 307 this author reports the same species under the native name "capa prieta," and after giving slightly different dimensions, supplies the following data:

A wild tree which is 40 feet (12 meters) in height with a trunk 21 to 27 feet (7 to 9 meters) long and 10 inches in diameter. Found commonly in mountainous localities. The wood is of ordinary weight, very strong, and frequently very elastic, the color of dead leaves, sometimes gray, fibrous in texture, rather fine-grained. Specific gravity, 0.574. The wood is much appreciated and was employed by the military authorities of San Juan in making windlasses, gun carriages, arms for fencing, etc. Also used for doors, venetian blinds, cots, bungs of barrels, the sweeps of sugar mills, and ordinary furniture.

Cordia gerascanthoides. CAPÁ PRIETA.

A tree that reaches 60 feet (18 meters) in height and whose trunk is 18 inches (45 centimeters) in diameter. Furnishes a wood that is much appreciated, for it



MORAL TREE (CORDIA MACROPHYLLA). COAMO SPRINGS.



CEREZAS (CORDIA NITIDA).

is at the same time strong and somewhat flexible, elastic, hard, more than ordinarily heavy, fibrous in texture, with a fine, compact grain. The heart is very hard, chestnut-colored, ornamented with lighter lines that are sometimes reddish. The outside wood is gray to light yellow, with narrow lines of a darker color. The wood is susceptible of a high polish, very beautiful when varnished. Specific gravity, 0.754 or 0.752. The military authorities of San Juan used this wood for axles, gun carriages, etc. Also made into doors, venetian blinds, beds, carriages, beehives, etc. Bees are attracted to its fragrant flowers, and cattle and hogs to the fruit. (Grosourdy, 2: 364.)

Cordia globosa. COPILLO.

According to Grisebach, a shrub 6 to 10 feet (2 to 3 meters) high, found in dry soil along the coast. According to Stahl a low shrub 4 inches (10 centimeters) high in salt marshes. From Cabo Rojo. (Stahl, 6: 96.)

Cordia macrophylla. MORAL.

PLATE XXX.

A forest tree 8 meters high, found, according to Grisebach, in all inland woods. Wood little used; in Jamaica it is claimed to be a good timber wood. (Stahl, 6: 92.)

Cordia nitida. CEREZAS.

PLATE XXXI.

A small tree along the road between Ponce and Adjuntas. The flowers are greenish white and the berries cherry red. The pulp of the latter is used for gluing musical instruments.

Cordia rupicola.

A shrub of 4 or 5 meters, from the southwestern part of the island. From vicinity of Guanica. (Urban, 1: 392.)

Cordia sebestena. VOMITEL COLORADO. ALOE-WOOD.

A shrub 3 to 4 meters high, found wild on the eastern coast. Cultivated in gardens for its beautiful flowers. (Stahl, 6: 96.)

A variety *brachycalyx* has recently been described by Urban, who states that the genuine *sebestena* is not known from the island except in cultivation. The additional popular name "San Bartolome" is also given. (Urban, Symb. 1: 389.)

Cordia ulmifolia. PALO DE PERICO.

A shrub 6 to 15 feet (2 to 5 meters) high, found in rocky waste places. Bello records the name "basora prieta" for this species. (Stahl, 6: 94.)

The variety *ovata* is reported from Peñuelas by Urban on the basis of specimens secured by Sintenis.

Cordobancillo. See *Rondeletia arborescens* and *R. laevigata*, and *R. inermis*.

Corkwood. See *Ochroma lagopus* and *Anona palustris*.

Cornutia obtvata.

Family Verbenaceae; an indigenous tree attaining a height of 10 meters; known only from the vicinity of Barranquitas.

Cororron.

A tree from the eastern parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood yellow, rather hard; specific gravity, 0.862; used in building country houses. (Exp. 1857.)

Corozo. See *Acrocomia media*.

Corozo del Orinoco. See *Martinezia caryotaefolia*.

A palm of the family Cocaceae. This name was unknown to Captain Hansard.

Corynella paucifolia. RETAMA.

A leguminous shrub, attaining 2 meters in height, with blue flowers. Known from Yauco and San German.

Coscarron.

According to Captain Hansard the specific gravity is 0.90.

Cosmos caudatus. MARGARITA.

Family Compositae; a common weed in open places. Near Toa Baja this was called "claveles," and at Coamo "piquete." (Stahl, 5: 130.)

Cotona.

A tree from all parts of the island; height, 25 to 30 feet (7 to 9 meters); diameter, 8 to 10 inches (20 to 25 centimeters). Wood hard, yellow; specific gravity, 1.194; used in building country houses. (Exp. 1857.)

Captain Hansard gives a specific gravity of 0.90 for this tree.

Cotorra.

A wild tree, produced in abundance over the entire island, 30 to 35 feet (9 to 11 meters) high, with a trunk 8 to 10 inches (20 to 25 centimeters) in diameter. It furnishes a moderately hard wood, canary-yellow in color, breaking with a vertical fracture. Used in the framework of country houses, for fences and other uses of this class. (Grosourdy, 2: 378.)

Cotorrera. See *Heliotropium indicum*.

Cotorrera de agua. See *Heliotropium inundatum*.

Cotorrera de la playa. See *Heliotropium curassavicum*.

Cotorrerilla. See *Heliotropium parviflorum*.

Cotorrerillo. See *Casuarina bicolor* and *C. sumyda*.

Cotorrillo. See *Elephantopus tomentosus*.

Cotton.

Cotton nankeen. See *Gossypium religiosum*.

Cotton tree. See *Ceiba pentandra*.

Courbaril. See *Hymenaea courbaril*.

Cowhage. See *Mucuna pruriens*.

Cowitch. See *Mucuna pruriens*.

Coyure. See *Curima colophylla*.

In his account of this palm Mr. Hill has probably confused the royal palm with the llume. There is nothing to indicate reference to the small, spiny palm to which the name is in reality applied, at least in the vicinity of Vega Baja.

Cracca aniloides. AÑIL FALSO.

Family Viciaceae; an herbaceous annual, 40 centimeters high; flowers in November; reported from the beach of the east coast. (Stahl, 3: 28, as *Tephrosia aniloides*.)

Cracca cinerea. AÑIL CENIZO.

A woody herb, 40 centimeters high, found along the beach; flowers all the year. Common throughout tropical America. Specimens collected along the seashore at Cataño, where the plant is called simply añil, and is used as a blue dye for cloth. (Stahl, 3: 26, as *Tephrosia cinerea*.)

Cracca leptostachya. AÑIL RACIMILLO.

An herbaceous annual, 50 centimeters high, growing along the sandy shore, between Arecibo and Aguadilla. Common to the tropics of America and Africa. (Stahl, 3: 27, as *Tephrosia leptostachya*.)

Cracca purpurea. AÑIL RACIMILLO.

Craniolaria annua. ESCORZONERA.

Family Martyniaceae; an herb a meter high. The large fleshy roots are used as a substitute for the European "escorzonera." (Stahl, 6: 264.)

Crape myrtle. See *Lagerstroemia indica*.

Cremanium amygdalinum. See *Tamonea integrifolia*.



CALABASH TREE (*CRESCENTIA CUJETE*). NEAR RIO PIEDRAS.

Crescentia cucurbitina. HIGÜERILLA.

Family Bignoniaceae. From Aguada. (Stahl, 6: 184.)

Crescentia cujete. HIGÜERA. CALABASH. PLATE XXXII.

A tree, according to Lunan, growing about 20 feet (6 meters) high. The fruit is used for receptacles of various kinds, while the juice acts as a purge and is said to cause abortion in cattle. The pulp with other ingredients forms an excellent cough remedy. The wood is used in Jamaica for shafts, saddles, chairs, and other articles that require a tough, flexible wood. (Stahl, 6: 183.)

In Mexico and other Spanish-American countries the common name is "jicara."

Crescentia microcarpa. HIGÜERITO.

(Stahl, 6: 184.)

Cresta de gallo blanco. See *Teramnus uncinatus*.

Critonia dalea. See *Eupatorium dalea*.

Critonia parviflora.

The variety *portoricensis* is reported from Maricao.

Crotalaria incana. CASCABELILLO VACÍO.

An herbaceous, pubescent annual with trifoliate leaves, growing in dry, rocky places. From Cayey, along the road. (Stahl, 3: 24.)

Crotalaria lotifolia. CASCABELILLO AXILAR.

A slender, small-leaved shrub, 75 centimeters high; found at San Juan, Fajardo, Coamo, and Guanica. (Stahl, 3: 25.)

Crotalaria retusa. MATRACA.

A common leguminous weed with showy yellow and maroon flowers; frequent in open waste places. In India this species is cultivated for its fiber. A near relative, *C. juncea*, is the sunn hemp plant of India, a shrub 10 feet or more in height, extensively grown for its fiber, which is said to be better than jute. It was introduced by the Department some years since for planting in the Southern States, but the fiber did not prove to be of good quality except in Florida. An experiment in Porto Rico is desirable, particularly since *C. juncea* is considered of value in India for forage as well as for fiber. According to Stahl the common name of *C. retusa* is "cascabelillo grande," while Bello calls it "sonajuelas."

Crotalaria stipularia. CASCABELILLO ALADO.

An herb 50 centimeters high. (Stahl, 3: 22.)

Croton.

Family Euphorbiaceae; numerous species of shrubs of this genus are known in Porto Rico. *C. discolor* is reported from Ponce.

Croton flavus rigidus.

This plant is said to be used in baths, for nervous weakness. From Guanica.

Croton humilis.

A shrub 1 to 2 feet (30 to 60 centimeters) high. Called "seaside balsam" in Jamaica. From vicinity of Guanica.

Croton lucidus.

A shrub 5 to 6 feet (1.5 to 2 meters) high. From Salinas de Cabo Rojo.

Cryptomeria.

Like most tropical countries, Porto Rico is deficient in coniferous trees. It is not, however, impossible that *Cryptomeria japonica*, the largest and perhaps the most beautiful of Japanese evergreens, might be found to flourish at the higher altitudes. This species, while it will endure frost, thrives also in mild, moist climates, and has been very successfully cultivated in the Azores for its valuable timber.

Cryptostegia grandiflora.

An East Indian asclepiadaceous vine; has been introduced into Jamaica. It is said to produce rubber of good quality, but the latex is not abundant and collection is slow and difficult.

Cuabilla. See *Amyris elemifera*.

A Cuban name not yet recorded in Porto Rico.

Cuba bark. See *Paritium elatum*.

Cucubano.

A tree from all parts of the island; height, 35 to 40 feet (11 or 12 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood rather hard, ash-colored; specific gravity, 0.888; used in building country houses. (Exp, 1857.) (Grosourdy, 2: 379). Captain Hansard says that it is mostly of small size, the saplings, measuring about 3 inches (7 centimeters), being used in hut building; specific gravity, 0.84.

Cucumber. See *Cucumis anguria* and *C. sativus*.

Cucumis anguria. CONCOMBRE. ANGURIA. PEPINO. PLATE XXXIII.

Family Cucurbitaceae. In Cuba called "pepino cimarron;" an annual cucurbitaceous vine with small oval spiny fruits, which are used as a substitute for the common cucumber and are commonly sold in the markets. (Stahl, 4: 178.)

Cucumis citrullus. See *Citrullus vulgaris*.

Cucumis melo. MELON.

The introduced muskmelon. (Stahl, 4: 179.)

Muskmelons are not an uncommon fruit in the markets of Porto Rico, and several varieties were observed. All were, however, lacking in flavor.

Cucuracey.

A tree from the eastern parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 8 to 10 inches (20 to 25 centimeters). Wood rather hard, pinkish; specific gravity, 0.897; used in building houses in the country. (Exp. 1857.)

Cucurbita pepo. SQUASH. CALABAZA.

Several varieties of squashes were seen in the markets of San Juan and Ponce, but mostly of small size.

Cuenta de oro.

"Golden bead;" a large tree with orange-yellow fruit; not yet identified.

Cuernecillo. See *Morongia distachya* and *Helicteres jamaicensis*.

Cuero de sapo. See *Ecostemma caribaeum*, *Ilex dioica*, *I. nitida*, and *I. urbaniana*.

"Frog's skin;" a tree of small size; soft bark; used in hut building; plentiful in the mountains; not known in lowlands. Captain Hansard gives a specific gravity of 0.89 for this.

A tree from all parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 12 to 15 inches (30 to 45 centimeters). Wood yellow, rather hard; specific gravity, 1.883; not used. (Exp. 1857.)

Used for fuel in the country districts. (Grosourdy, 2: 380.)

Culantrillo. See *Scoparia dulcis*.

Culantrillo de Pozo.


No. 231. An unidentified species of *Adiantum*.

Culantro del monte. See *Eryugium foetidum*.

Cundeamor. See *Momordica charantia*.

Cupania americana. GUARA.

Family Sapindaceae; a forest tree, 5 meters high. The soft wood takes a high polish. (Stahl, 2: 156.)



CONCOMBRE (CUCUMIS ANGUIRIA).

Cupania sapida. See *Blighia sapida*.

Cupania tomentosa. See *Cupania americana*.

Cupei. See *Clusia krugiana*.

Cupeillo. See *Clusia krugiana* and *C. acuminata*.

Cupey. See *Clusia rosea*.

A tree from all parts of the island: height, 20 to 30 feet (6 to 9 meters); diameter, 6 to 8 inches (15 to 20 centimeters); wood yellow, rather hard; specific gravity, 1.058; kills trees to which it attaches itself. (Exp. 1857.)

Cupey colorado.

A tree from the interior parts of the island: height, 50 to 60 feet (15 to 18 meters); diameter, 15 to 20 inches (37 to 50 centimeters); wood hard, red; specific gravity, 0.874; used in building houses. (Exp. 1857.)

Cupey de altura. See *Clusia gundlachi*.

Cuphea micrantha. CHILAGARI.

Family Lythraceae: small shrub, 30 centimeters high, growing along the coast. The same common name is also applied to *Cuphea parsonsia*, a smaller species, only 15 centimeters high. (Stahl, 4: 125.)

Curbaril. See *Hymenaea courbaril*.

Curcuma longa. TURMERIC. YUQUILLO.

An introduced zingiberaceous plant running wild in Jamaica, and probably also in Porto Rico. The roots are used as an ingredient in curry powders, and also produce a yellow dye much used in India. The yellow principle is not developed until the roots mature, and in the younger state they are used much like arrow-root, starch being extracted from them.

Curia. See *Dianthera pectoralis*.

Curima calophylla. COYURE.

Family Arecaceae: Curima appeared to be especially abundant about Bayamon, but is probably rather generally distributed in the limestone hills of the island, perhaps also on other soils. A few trees were seen along the road between Utuado and Lares, and numerous others between Isolina and Manati. Sintenis collected specimens of what is apparently the same species near Juncos and Hato Grande, and young specimens at Maricao.

As far as Porto Rico is concerned, this palm is very easily recognized by means of the curiously truncate leaf divisions, the outer margins of which appear as though accidentally injured or eaten away by caterpillars. This feature is, however, shared with numerous other West Indian and South American palms, though apparently only one, the so-called *grigri* palm of Martinique, can be referred to the present genus with confidence.

Hill's statement that the coyure palm is taller than the royal palm does not apply to this species, but probably has reference to the llume palm (*Aeria attenuata*). Neither species is common nor much used. *Aeria* was pointed out to us from the distance as "coyure," but when close at hand all agreed that that name belongs to the present spiny species.

Cuscuta americana. DODDER. FIDEOS.

A slender parasitic vine, reported by Bello.

The large dodder of Porto Rico not only covers large masses of vegetation with its coarse yellow threads, but it sometimes invades and covers trees of considerable size.

Custard apple. See *Annona reticulata*.

Cycas revoluta.

Occasional in garden—growing to large size in the open air.

Cynara scolymus. ARTICHOKE. ALCACHOFA.

A few specimens of the genuine European artichoke were seen near Santurce, apparently escaped from cultivation.

Cynodon dactylon. See *Capriola dactylon*.**Cynometra portoricensis.**

Family Cassiaceae; a leguminous tree, 15 meters in height, recently described from the vicinity of Rincon. (Urban, Symb. 1: 312.)

Cyphomandra betacea. TREE TOMATO.

A plant related to the tomato and at first placed in the genus *Solanum*. It grows rapidly to a height of 10 or 12 feet (3 to 4 meters), becoming a veritable shrub or small tree. The heart-shaped flowers are sometimes a foot in length. The fruits are borne in clusters, and are oval in shape, or somewhat pointed; the color is purplish or reddish, and the taste is similar to that of the ordinary tomato, though some have reported a gooseberry flavor in the ripe, raw fruit, and if stewed after the skin and seeds have been removed a resemblance to apricot is said to be developed. It has become very popular for tarts and pastries; also for jellies and preserves. The suggestion which is considered under the eggplant of grafting on *Solanum mammosum* might also find application in the present instance.

Cyphomandra is a native of the Andes of Peru and neighboring regions of tropical America, but has been successfully introduced into Jamaica, Ceylon, and other mountainous regions of the Tropics, and at many places is considered a valuable addition to the list of garden vegetables. It would in all probability thrive in Porto Rico, and as it can be propagated readily from seed its introduction offers no special difficulties.

Cypress vine. See *Quamoclit quamoclit*.**Cyrilla antillana.** CIRILA.

Family Cyrillaceae; a tree found on the south coast. From the vicinity of Utuado and at Santa Ysabel. (Stahl, 6: 34.)

Dacryodes excelsa.

A tree of the family Burseraceae; reported from near Maricao.

Dacryodes hexandra. TABONUCO.

PLATE XXXIV.

A tree exuding a resin used for torches; found in mountain districts. Grisebach gives it as 10 to 15 feet high; Stahl as 75 feet high. The trunks are used for lumber. (Stahl, 4: 53.)

Near Isolina there still exists a considerable area of virgin forest composed largely of what was taken to be this species. The trees are much higher than even the figures given by Stahl, with very smooth, straight, white trunks, often scarred near the base where they have been cut to secure the resin. The wood has a beautiful color and is said to be free from the attacks of insects.

Dalbergia ecastaphyllum. MARAY-MARAY.

A leguminous shrub, 3 to 4 meters high, found in thickets along the coast.

Dalbergia monetaria. PALO DE BRASILETE.

The red wood is said to furnish a resin resembling dragon's blood. Known from Utuado as *Hecastaphyllum monetaria*. (Stahl, 3: 97.)

Dalechampia scandens.

Family Euphorbiaceae; a twining shrub; from Guanica.

Dama de noche. See *Cestrum nocturnum* and *Murraya exotica*.**Dama juana.**

Captain Hansard gives the specific gravity of 0.82 for the wood of this species.

Daphnopsis caribaea.

Family Thymelaeaceae; a tree 20 to 30 feet (6 to 9 meters) high; known from Cayey.



VIRGIN FOREST OF TABONUCO · DACRYODES HEXANDRA · AT ISOLINA

Daphnopsis philippiana. CIENEGUILLO.

Also called "emajagna" and "emajagua brava." A shrub or small tree of the family Thymelaeaceae, evidently generally distributed in forest districts, being reported from Luquillo, Naguabo, Juncos, Aibonito, Barranquitas, Penuelas, Utuado, and Jayuya.

Date. See *Phoenix dactylifera*.

Date palm. See *Phoenix dactylifera*.

Date plum. See *Diospyros tetrasperma*.

Another species, *Diospyros ebenaster*, is known from Porto Rico.

Datil. See *Phoenix dactylifera*.

Datura fastuosa. ESTRAMONIO.

Family Solanaceae; an herbaceous annual, 1 meter high. The seeds are said to possess poisonous properties. (Stahl, 6: 119.)

Datura metel. CHAMICO BLANCO.

An herbaceous annual weed, 1 meter high, frequenting rocky places along the coast. (Stahl, 6: 117.)

Datura stramonium. JIMSON WEED. ESTRAMONIO.

Also called "peo de fraile," "chamico morado," "chamisco," and "thorn apple." A solanaceous herb 3 to 4 feet (1 to 1.3 meters) high. The seeds are poisonous and are said to have medicinal value as a narcotic. (Stahl, 6: 115, 118.)

Datura suaveolens. CAMPANA DE PARIS.

A shrub 3 to 4 meters high, found along river banks. (Stahl, 6: 120.)

Datura tatula. See *Datura stramonium*.

Demajagua.

This name is probably intended for emajagua: at least that is its use in Cuba and South America.

Demajagua blanca.

A tree from all parts of the island; height, 10 to 12 feet (3 to 4 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood white, soft; specific gravity .696; bark used for making ropes. (Exp. 1857.)

Demajagua colorado.

A tree from all parts of the island; height, 10 to 12 feet (3 to 4 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood white, soft; specific gravity .698; bark used for making ropes. (Exp. 1857.)

Dendropremon bicolor. PARASITA.

Also called "hicaquillo del combro." A parasitic shrub of the family Loranthaceae. It is recorded as growing upon the orange and the custard apple.

Dendropremon caribaeus. HICAQUILLO. CAPITANA.

A parasitic shrub, attacking numerous trees, both wild and cultivated.

Desmanthus plenus. See *Neptunia plena*.

Desmanthus virgatus. DESMANTO RAYADO.

Family Mimosaceae; a woody herb 4 to 6 feet (1.3 to 2 meters) high, sometimes decumbent; found along the coast at Fajardo. (Stahl, 3: 140.)

Desmanto amarillo. See *Neptunia plena*.

Desmanto rayado. See *Desmanthus virgatus*.

Desmodium. See *Meibomia*.

Dianthera pectoralis. CURIA.

Family Acanthaceae; an herbaceous plant 40 centimeters high; cultivated in gardens, and wild in dry places. A sirup made from this plant is said to be beneficial for pulmonary disorders. Called also "justicia balsam" in Jamaica. (Stahl, 6: 248.)

Didymopanax micans. YAGRUME.

An indigenous araliaceous tree known from the vicinity of Adjuntas, Cabo Rojo, and Aguada. (Urban, Symb. 1: 204.)

Didymopanax morototoni. YAGRUME.

Variouly called "grayume" or "grayumo," "yagrume" or "yagrumo," "yagrume macho," and "pana cimarrona." A tree of 5 to 20 meters, probably the more common and widely distributed species in Porto Rico. "Yagrumo macho" or "male yagrumo" alludes to the fact that this tree is believed to be the male of *Cecropia*, which is called "yagrumo hembra."

Diffenbachia seguine.

Family Araceae; a very succulent aroid from swampy meadows near Juncos. It grows from 3 to 6 feet (1 to 2 meters) high.

Diego de noche.

In Cuba *Allionia incarnata*.

Dioclea reflexa. MATO.

Also called "bejuco de mato." A leguminous vine climbing to the tops of the highest trees. Reported from Mannabo, Yabucoa, and Luquillo. (Urban, Symb. 1: 473.)

Diodia maritima.

A rubiaceous herb. (Stahl, 5: 159, as *Diodia radicans*.)

Diodia radicans. See *Diodia maritima*.**Diodia rigida.** *Diodia rigida*.

Family Rubiaceae; an herbaceous annual; frequents dry clay soil; 20 to 30 centimeters high. (Stahl, 5: 84.)

Diodia sarmentosa. *Diodia sarmentosa*.

A woody herb growing among shrubs in arid soil; 1 meter high; from Maricao. (Stahl, 5: 86.)

Diodia scandens. DIODIA TREPADORA.

A woody herb, 1 meter high, found in dry soil. (Stahl, 5: 85.)

Diodia trepadora. See *Diodia scandens*.**Dioscorea.** YAM. ÑAME.

The true yams of the Tropics belong to a distinct botanical family, the Dioscoreaceae, related to the Smilax family and the Lily family, but having no affinity to the sweet potato, to some varieties of which the name is commonly applied in the South. Nearly all of the very numerous species form more or less thickened roots, and many of them have been introduced into cultivation, mostly in the Tropics. Some years since the Chinese yam, *Dioscorea batatas*, which is hardy in temperate climates, was advocated for general culture in Europe and the United States, but its habit of forming its tuberous roots 3 feet in the ground rendered it very difficult to handle by ordinary methods of agriculture. The tropical species have the same tendency, but usually not to so pronounced a degree, and in most tropical countries the yam is a more or less important article of diet. In some it is merely accessory to cassava, while in others it takes first place as a root crop. It seems certain that in Porto Rico they have received no such careful attention as in the British West Indies, and the following abridgment of a report on "Yams in the West Indies," recently furnished by Mr. David G. Fairchild, Agricultural Explorer of the Section of Seed and Plant Introduction, United States Department of Agriculture, will probably be of use to those interested in agriculture in Porto Rico.

It is important, of course, to plant the best sorts, but with so many species and varieties in cultivation it is difficult to tell the different forms apart or to know which is most desirable. At the Trinidad Botanic Gardens a series of experiments

have been conducted, on which a report has been made during the past year, and as this supplements the information furnished by Mr. Fairchild, and represents what is now known about the subject in that island, it seems best to include it here:

The experiments in connection with this class of plants have been continued at St. Clair, and the improvement of the quality of the tubers grown in the new ground is so marked as to be patent to the veriest novice.

The kinds grown were as follows:

Yellow yam	<i>Dioscorea cayenensis.</i>
White yam	<i>Dioscorea alata.</i>
Negro yam	<i>Dioscorea alata</i> , var.
Red yam	<i>Dioscorea alata</i> , var.
Horn yam	<i>Dioscorea alata</i> , var.
Water yam	<i>Dioscorea alata</i> , var.
Chinese yam	<i>Dioscorea batatas.</i>
Afoo yam	<i>Dioscorea lutea.</i>
Cush-cush, or Indian yam	<i>Dioscorea trifida.</i>

Dioscorea bulbifera is so evidently inferior that its culture has not been continued. The "devil yam," which was taken to be a very coarse variety of *D. alata*, has also been discarded as useless. The "negro yam" is by far the best of the white yams grown. The "yellow yam" is also a good table kind, very sweet and wholesome and much liked by American and European visitors.

The "cush-cush," or "Indian yam," also known as "yampee," gives but very small tubers, and there are three or more varieties—red cush-cush, white cush-cush, and Demerara cush-cush. The last-mentioned kind has larger tubers than the others, testiculate in form, and generally of excellent quality. It has been found that the return made by varieties of this species is much less than from any other, but it is probably superior to any for the table of the better classes, although it would not be profitable to grow where a cheap supply of food stuff was required.

The "Chinese yam," as I know it, is one of the best. Well grown and well ripened, it is of fine quality for table, and probably comes nearer in flavor to a good potato than any other variety. The "afoo yam" is the most inferior of all. This name, however, is sometimes applied to the "yellow yam," so that one may get the worst as well as the best yams under the same name, so little is local nomenclature to be trusted.

The yield of yams given in former reports has been more than maintained. White yam gave over 20 pounds weight to a plant; Negro yam, 17 pounds; Chinese yam, 19 pounds; yellow yam, 12 pounds; and cush-cush, 7 pounds. Yams grown from aerial tubers, i. e., tubers which develop on the vines, gave 19 pounds per root. White yam yielded at the rate of 13 tons to an acre. If 25 per cent is deducted from this for waste and for planting again, there is still a crop of 10 tons per acre to be registered as the crop for the year of 1898.

The importance of similar and more extended experiments in Porto Rico is obvious, both for the sake of stimulating progress in the direction of mixed farming and greater local production of food stuffs, and because it is probable that among so many botanical species of different origins special adaptations to soils and climates will be found to exist, a knowledge of which will be of the utmost practical importance in utilizing these valuable food plants. The excellence of the crop will also determine the possibility of opening a trade with the United States, as suggested by Mr. Fairchild. It has been found that properly packed yams may be shipped from the West Indies to New York and interior points and arrive in good condition.

Dioscorea alata.

This species seems to be one of the commonest in Porto Rico, and may be recognized by noting that the vine has thin ridges or wings on the stem.

In Jamaica this species is said to be particularly useful on account of the fact that it bears well on shallow or marly soils, while other kinds require deep mold; it is also thought to keep better than the other species.

Dioscorea chondrocarpa.

A wild yam, growing in forests near Yabucoa. (Sintenis.)

Dioscorea multiflora.

Collected by Sintenis in the Sierra de Naguabo.

Dioscorea pilosiuscula.

Another wild species from forests in the Sierra de Naguabo.

Diospyros ebenaster. GUAYABOTA.

Family Ebenaceae; a native forest tree reported by Stahl (6: 284) from Toa Alta; identified by Urban (Add. 1: 43).

Dipholis montana. VARITAL. TABLONCILLO.

Family Sapotaceae; a tree 5 to 6 meters high in mountain woods. (Stahl, 6: 55).

Dipholis nigra. BUMELIA NIGRA. AUSBBO.

A large tree, 15 to 20 meters high, found in all mountain woods. The hard, red wood is considered one of the best of the numerous kinds produced in the island, and is used for a large variety of purposes. (Stahl, 6: 56.)

Dipholis salicifolia.

A large, fragrant tree, growing in dry calcareous soil; reported from near Guanica.

Diplochita fothergilla. See *Tamonea fothergilla*.**Distictis lactiflora.** VIUDA.

Family Bignoniaceae; a tendril-bearing vine with the reticulate veins of the leaves very prominent. Our specimen (no. 694) was obtained near Coamo. The flowers are white, with the base of the corolla canary-yellow.

Distreptus spicatus. See *Elephantopus spicatus*.**Divi-divi.** See *Caesalpinia coriaria*.

A species also treated by some botanists as *Lebidibia coriaria*.

Dodonaea viscosa. CHAMISO.

Family Sapindaceae; a tree 3 to 4 meters high, found along the coast. (Stahl, 2: 161.)

Dolicholus minimus. FRIJOLILLO PEQUEÑO.

Family Viciaceae; a reddish pubescent vine found in all parts of the island, flowering nearly all the year, especially in winter; appears to be common to tropical countries generally. (Stahl, 3: 52, as *Rhynchosia minima*.)

Dolicholus phaseoloides. FRIJOLILLO COLORADO.

A vine seen only once in fruit by Stahl in the high hills of Bayamon. An indigenous species with short pods and broad, closely veined leaves. Found also in the other Antilles, Panama, Brazil, and the Galapagos Islands. (Stahl, 3: 53, as *Rhynchosia phaseoloides*.)

Dolichos lablab. FRIJOLES CABALLEROS.

Family Viciaceae; a woolly biennial cultivated in gardens for its edible beans. (Stahl, 3: 74.)

Dolichos monachalis. See *Vigna catjang*.**Dominicos.**

A variety of banana, mentioned by Hill.

Doncella. See *Byrsonima spicata*.

A tree from all parts of the island: height, 35 to 40 feet (11 or 12 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood pink, hard; specific gravity, 1.117; used in building houses. (Exp. 1857.)

Down tree. See *Ochroma lagopus*.**Drepanocarpus lunatus.** PALO DE HOZ.

A leguminous tree, 3 to 5 meters high; found in borders of swamps near the sea. (Stahl, 3: 94.)

Drymaria cordata. YERBA DE ESTRELLA.

Family Caryophyllaceae; an herbaceous reclining annual, 15 centimeters high, frequenting moist, shady places. (Stahl, 2: 56.)

Drypetes alba. CAFFEILLO.

Family Euphorbiaceae. A native tree reaching a height of 40 feet (12 meters); reported from mountain forests, Sierra de Luquillo, and also between Adjuntas and Ponce. (Urban, Add. 1: 71.)

Drypetes glauca. VARITAL.

Also called "caffeyillo" and "palo blanco." A euphorbiaceous shrub generally distributed in the native forests: Luquillo, Yabucoa, Penuelas, Adjuntas, Jayuya, Maricao. *Drypetes lateriflora* is known from Bayamon. (Urban, Add. 1: 73.)

Drypetes ilicifolia. ENCINILLO.

An indigenous shrub, 10 to 15 feet (3 to 4.5 meters) high, found by Stahl (4: 37) on the calcareous rocky hills along the north coast, between Cangregos and Loiza, and supposed to be a species of *Ilex*; secured by Gundlach at Arecibo. (Urban, Add. 1: 68.)

Duranta plumieri. LLEVIA.

Family Verbenaceae, a shrub 6 to 15 feet (2 to 4.5 meters) high; in moist, sunny places, also cultivated in gardens; collected at Manabo. According to Bello this is also called "azote-caballo" and "lila." (Stahl, 6: 212.)

Earth nut. See *Arachis hypogea*.

Ecastaphyllum brownei. See *Dalbergia ecastaphyllum*.

Ecastaphyllum monetaria. See *Dalbergia monetaria*.

Echites circinalis. BABEIRO.

Family Apocynaceae. (Stahl, 6: 79.)

Echites andrewsii. BABEIRO AMARILLO.

A woody climber, 3 to 4 meters high, reported from Manati. (Stahl, 6: 82, as *E. neriantra*.)

Echites paludosa.

An apocynaceous vine, native in several of the Bahamas, Cuba, Haiti, Jamaica, and Panama. According to Grisebach's Flora (p. 406) this species is a source of rubber in Jamaica.

Echites umbellata. BABEIRO.

A woody herb, 3 to 5 meters high, occurring in shady waste places. (Stahl, 6: 79.)

Eclipta alba. See *Eclipta erecta*.

Eclipta blanca. See *Eclipta erecta*.

Eclipta erecta.

Family Compositae. (Stahl, 5: 132, as *Eclipta alba*.)

Eddoe. See *Colocasia esculenta*.

Eggplant. See *Solanum melongena*.

Egyptian privet. See *Lawsonia inermis*.

Ehretia bourreria.

Family Boraginaceae; a branching tree, 8 to 14 feet (2.5 to 4.5 meters) high; it bears small edible berries, and in Jamaica is called "currant tree." Known from Yabucoa.

Elaeis guineensis. AFRICAN OIL PALM.

The African oil palm, *Elaeis guineensis*, furnishes the chief article of export and is the principal basis of the commerce of West Africa. Two products are exported—the oil, derived from the reddish exterior of the pulp of the fruit, and the kernel,

secured by cracking the inclosed nut after the pulp has been removed. The fruit is egg-shaped, and from 1 to 1½ inches (25 to 37 millimeters) in length. The deep red skin is thin, like that of the date. The oily pulp underneath is interspersed with tough fibers, but has a pleasant taste, and is frequently eaten after being slightly roasted. The oil is obtained in quantities by beating the nuts in wooden mortars and then skimming off from boiling water, also by less desirable methods in which the nuts are rotted. This is said to increase the yield of oil, but gives it a rancid taste unpleasant to the civilized palate, but not injuring the oil for export. In Europe the kernels are used for the extraction, under pressure, of another oil quite similar to that of the cocoanut and used for like purposes.

Elaeodendron attenuatum. COCORRONCITO.

Family Celastraceae; a low tree or shrub, 2 meters high, found along the sea-coast. (Stahl, 4: 20.)

Elaeodendron xylocarpum. COCORRON.

A low tree or shrub, 2 to 3 meters high, found along the seacoast; from Guanica; a variety of this is reported from near Mayaguez. (Stahl, 4: 18.)

Elephantopus mollis. See *Elephantopus scaber*.

Elephantopus scaber. LENGUA DE VACA.

An herbaceous composite weed, 1 to 3 feet (30 to 90 centimeters) high; reputed to have medicinal value in Jamaica and the French islands. (Stahl, 5: 105.)

Elephantopus spicatus. YERBA DE BURRO.

A woody annual, 50 centimeters high; along roadsides and in dry pastures. (Stahl, 5: 106.)

Elephantopus tomentosus. COTORRILLO.

Specimens from Vega Baja (no. 1023).

Eleusine indica.

An annual, low, spreading grass; collected at Fajardo.

Eleutheranthera ovata. OGIERA.

An herbaceous annual, 50 centimeters high, found in sandy, stony places, (Stahl, 5: 126.)

Elm, Spanish. See *Cordia gerascanthoides*.

Emajagua. See *Paritium tiliaceum*.

Emajagua brava. See *Daphnopsis philippiana*.

Emajagua de sierra. See *Daphnopsis philippiana*.

Emajagua excelsa. See *Hibiscus elatus*.

Emilia sonchifolia. EMILIA.

An herbaceous composite annual; found along roads and in dry pastures. (Stahl, 5: 151.)

Encinillo. See *Drypetes ilicifolia*.

Enernaseada.

A tree from all parts of the island; height, 25 to 30 feet (7 to 9 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood yellow, hard; specific gravity 0.783; fruit tree. (Exp. 1857.)

Enredadera. See *Barbieria polyphylla*.

Enrubio.

A wild tree, not common, reaching a height of 25 or 30 feet (7 to 9 meters), with a trunk 10 inches (25 centimeters) in diameter. Its wood is very hard, with a red heart and white sapwood. Has no application (Grosourdy, 2: 382). Specific gravity 0.988 (Exp. 1857). Grosourdy refers this to *Zanthoxylum lanceolatum*, a species given in Index Kewensis as a synonym of *Z. clava-herculis*, which has since been referred to *Fagara martinicensis*.

Entada polyphylla.

Family Mimosaceae; a perennial vine; from Fajardo.

Erechtites hieracifolia. ACHICORIA DE CABRA (GOAT'S CHICORY).

An herbaceous erect annual composite, 50 centimeters high; grows in pastures. (Stahl, 5: 150.)

Erigeron bellioides. BELLORITA.

An herbaceous annual composite, 12 centimeters high; found in stony moist places. (Stahl, 5: 119.)

Erigeron canadensis. PASCUETA.

An herbaceous erect annual, 50 centimeters high; grows in pastures. (Stahl 5: 158.)

Erigeron rivularis. PASCUETA AMARILLA.

Described as *E. canadensis*, for which it was at first mistaken. (Stahl, 5: 121.)

Erigeron spathulatus. CHIRIVITA.

An herbaceous annual, 30 centimeters high; found in dry pastures. (Stahl 5: 120.)

Eriobotrya japonica. LOQUAT. NÍSPERO.

"The loquat is planted along the roadsides on many of the estates in the Wynaad and elsewhere, and coffee appears to thrive well under it; but, so far as I am aware, the wood is not of any value, which at once places it at a disadvantage in competing with the jack. The loquat yields a pleasant fruit, in size and appearance much like the yellow plum."—Hull.

Eriodendron anfractuosum. See *Ceiba pentandra*.**Erithalis fruticosa.** IAYAJABICO.

Family Rubiaceae; a shrub 3 to 4 meters high, found on the beaches and cliffs near the sea. (Stahl, 5: 61.)

Ernodea de playa. See *Ernodea litoralis*.**Ernodea litoralis.** ERNODEA DE PLAYA.

A decumbent rubiaceous shrub; from Cabo Rojo. (Stahl, 5: 83.)

Eryngium foetidum. CULANTRO DEL MONTE.

An ill-smelling umbelliferous herb, 1 foot (30 centimeters) in height, found near cultivated places. "Fit weed," as this plant is called in Jamaica, is reputed to have curative value in cases of snake bite, epileptic fits, etc. Specimens were collected along the military road beyond Rio Piedras; used in that locality for seasoning food. (Stahl, 4: 196.)

Erythrina corallodendron. PIÑON ESPINOSO.

A leguminous tree or shrub, 5 meters high, growing on limestone hills. Used for hedges in Jamaica, where it is called the "red bean tree." (Stahl, 3: 89.)

Erythrina glauca. BUCAGO.

A leguminous tree, 10 meters high, found near rivers. (Stahl, 3: 88.)

Erythrina micropteryx. BUCARE.

Also known as "palo de boyo." A leguminous tree of 15 to 20 meters beset with short, conical spines; flowers red. Cultivated as a shade tree for coffee and reported from numerous localities in Porto Rico. It is a native of the lower Andes of Peru. (Urban, Symb. 1: 327.)

Erythrina umbrosa. BUCARE. MADRE DE CACAO.

Also called "bois immortelle" and "madre de cacao" in South America. Captain Hansard claims to have introduced seed of this species from Trinidad, but considered the seedlings the same as the tree already common in Porto Rico and now supposed to be *E. micropteryx*. This species is said to be the most popular shade tree for the cacao plantations of the mainland, and is reported by

Hill and Stahl (?) as a coffee shade tree in Porto Rico. According to Urban, the true *E. umbrosa* is not yet known from any of the West Indies, such references pertaining to *E. micropteryx*. (Urban, Symb. 1: 327.)

Erythroxyton brevipes. JIBA.

Family Erythroxylaceae; a shrub, 2 to 3 meters high, found in sandy places near the sea. Urban reports the form *intermedia* from Fajardo. (Stahl, 2: 136.)

Erythroxyton havanense. See *Erythroxyton ovatum*.

Erythroxyton obovatum. JIBA.

A shrubby tree, 12 feet (4 meters) high, on limestone hills near the sea at Veja-baja and Aricebo. (Stahl, 2: 138.)

Escambron. See *Pisonia aculeata*.

Escambron blanco. See *Clerodendron aculeatum*.

A simple-leaved spiny plant, collected about the large cacti near Guayanilla.

Escambron colorado.

A name applied to *Pithecolobium unguis-cati* at Guayanilla.

Escoba. See *Sida carpinifolia*.

Escoba acorazanada. See *Sida cordifolia*.

Escoba babosa. See *Bastardia bivalvis*.

Escoba blanca. See *Sida carpinifolia*.

Escoba colorada. See *Sida rhombifolia*.

At Cataño this name was also applied to a small species of *Croton* (no. 960).

Escoba de Jamaica. See *Sida jamaicensis*.

Escoba dulce. See *Sida ulmifolia*.

Escoba pestañosa. See *Sida ciliaris*.

Escoba purpurina. See *Sida purpurea*.

Escoba tendida. See *Sida supina*.

Escobillon. See *Pterocaulon virgatum*.

Escobita dulce. See *Sida ulmifolia*.

Escorzonera. See *Craniolaria annua* and *Martynia diandra*.

Espejuelo. See *Sarcomphalus reticulatus*.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood yellow, very hard; specific gravity, 1.299; used for cabinetwork. (Exp. 1857.)

The strong yellow wood is used for shelves and fences. (Grosourdy, 2: 382.)

Espejuelo amarillo.

A tree from the northeastern part of the island; height, 40 to 50 feet (12 to 15 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood yellow, very hard; specific gravity, 1.050; used for cabinetwork. (Exp. 1857.) (Grosourdy, 2: 382.)

Espejuelo bobo.

A thick-barked tree, not of very large size; found occasionally in the hills; specific gravity, 1.08. (Hansard.)

Espiga de San Antonia. See *Buchnera elongata*.

Espinillo.

A tree from all parts of the island; height, 35 to 40 feet; diameter, 12 to 15 inches. Wood greenish, very hard; specific gravity, 1.114; used for cabinetwork. Also called "huso." (Exp. 1857.) (Grosourdy, 2: 382.)

Captain Hansard gives this wood, which indicates that the tree occurs in the Luquillo region; specific gravity, 1.10. A spiny tree, similar to *Randia aculeata*, bears this name in the vicinity of Vega Baja, but our specimen (no. 1028) is sterile and has not been identified.

Espino. See *Fagara martinicensis* and *F. monophylla*, *Zanthoxylum clava-herculis*, *Z. acuminatum* and *Z. caribaeum*.

According to Captain Hansard, this tree has a specific gravity of 7.69. He also calls it the "box thorn." The wood is light yellowish. The trunk has large, conical spines, bark turning from gray to white. This suggests the possibility that *Hura crepitans*, generally called "avilla," may be intended.

Espino amarillo.

A tree from all parts of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 15 to 20 inches. Wood yellow, soft; specific gravity, 0.556; used for lumber. (Exp. 1857.)

Grosourdy (2: 382) refers this to the genus *Zanthoxylon*.

Espino blanco.

A tree from all parts of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 15 to 20 inches (37 to 45 centimeters). Wood white, soft; specific gravity, 0.642; used for lumber. (Exp. 1857.)

Grosourdy (2: 382) also refers this to the genus *Zanthoxylon*, and suggests that it may be identical with the "agua blanca" of Cuba.

Espino rubial. See *Fagara caribaea*, *F. martinicensis*, *F. monophylla*, *F. pterota*, and *Z. ochroarylum*.

A tree from all parts of the island; height, 40 to 45 feet (12 to 14 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood yellow, hard; specific gravity, 0.664; used for lumber. (Exp. 1857.)

Espino rubio.

Very scarce and not much used; specific gravity, 0.68, according to Captain Hansard.

Espinosa. See *Anthacanthus spinosus*.

Esponja. See *Luffa aegyptiaca*.

Espuela de galan. See *Scolosanthus grandiflorus*.

Estramonio. PEO DE FRAILE. See *Datura stramonium* and *D. fastuosa*.

Estrapajo. See *Luffa aegyptiaca*.

Eugenia aeruginea. GUASAVARA.

An indigenous myrtaceous tree sometimes 60 feet in height, known from Bayamon, Juncos, Yabucoa, Guanica, Mayaguez, and Arecibo. A related species of equal size, *E. eggersii*, is reported from several points between Luquillo and Penuelas. (Urban, Add. 2: 141.)

Eugenia aromatica. PALO DE CLAVO.

A tree 25 feet (7 meters) in height, with a rather straight trunk 4 or 5 feet (120 to 150 centimeters) in diameter, that furnishes a wood of good weight and harder than the average; fibrous and compact with a fine grain. It is not used, and when dry has neither odor nor taste. Specific gravity, 0.897. (Grosourdy, 2: 405.) The native name suggests that of the clove, *Caryophyllus aromaticus*.

Eugenia biflora ludibunda. PITANGUEIRA.

Also called "hoja menuda." A myrtaceous shrub or small tree, not reaching 20 feet in height. Indigenous in the thickets of the limestone hills near the coast. Reported from the vicinity of San Juan, Juncos, Fajardo, Manati, Morovis, and Dorado. The variety *lancea* comes from Fajardo and Camuy. (Urban, Add. 2: 149.)

Eugenia buxifolia. HOJA MENUDA.

A myrtaceous shrub or small tree, extending from Florida to Cuba, Santo Domingo, Jamaica, and Porto Rico; known in Porto Rico from the vicinity of Guanica and Cabo Rojo. A similar but somewhat larger species, *E. avillaris*, is

known from Bayamon and several localities in the western part of the island, but no common name is recorded. (Urban, Add. 2: 157.)

Eugenia caryophyllata. See *Caryophyllus aromaticus*.

Eugenia confusa. CIENEGUILLO.

A myrtaceous tree 60 feet (18 meters) in height from the Florida Keys, Bahamas, Jamaica, Virgin Islands, Antigua, Guadeloupe, and Dominica. In Porto Rico it is known only from near Maricao in the forests of Mount Alegrillo. *E. maricao*, 25 feet (7.5 meters), has a similar distribution, and is known from littoral woods near Ponce. (Urban, Add. 2: 161, 162.)

Eugenia cordata sintenisii. MURTA.

An indigenous myrtaceous shrub or small tree 30 feet (9 meters) high, reported from hillsides near Aibonito and Cayey.

Eugenia costata. See *Eugenia serrasuela*.

Cultivated in gardens at Yabucoa.

Eugenia flavo-virens. GUAYABILLO DE COSTA.

Family Myrtaceae; a shrub 2 meters high; grows on the coast. (Stahl, 4: 86.)

Eugenia floribunda. MURTA.

An indigenous myrtaceous tree attaining a height of 50 feet (15 meters). Reported from Cayey, Lares, and Quebradillas. A related species, *E. cordata*, also called murta, is represented in Porto Rico by a subspecies, *sintenisii*, a small tree of 30 feet (9 meters) reported from hillsides near Aibonito and Cayey. (Urban, Add. 2: 173-175.)

Eugenia jambos. See *Jambosa jambos*.

Eugenia ligustrina. PALO DE MURTA.

Also known as "hoja menuda." A myrtaceous shrub or small tree, 18 feet (5 meters) high, distributed from the Bahamas to Trinidad; reported in Porto Rico from Fajardo, Coamo, Guanica, and Cabo Rojo, in woods along the shore and in river valleys.

Eugenia malaccensis. OTAHEITE APPLE.

The "Otahete" or Malay apple has been introduced and become established in Jamaica and several other islands of the West Indies, but it is not known to exist in Porto Rico. The same may be said of *E. javanica*, the "Malacca apple," and *Syzygium jambolanum*, the Java or jack plum.

Eugenia monticola. BIRIJI.

Also called "hoja menuda." An indigenous myrtaceous shrub or small tree sometimes attaining 30 feet (9 meters). Apparently generally distributed in forests and thickets throughout the island, Bayamon, Juncos, Fajardo, Cayey, Coamo, Isabella, and Quebradillas. (Urban, Add. 2: 154.)

Eugenia poiretii. HOJA MENUDA.

Also called "biriji;" a shrub 3 to 4 meters high on sandy beaches; also reported from the vicinity of Boamo and from Juncos. (Stahl, 4: 79.)

Eugenia portoricensis. PETANGUEIRA.

Urban describes this species as a variety of *Eugenia pseudopsidium*. (Stahl, 4: 82.)

Eugenia procera. HOJA MENUDA.

A small indigenous tree of general distribution from Cuba to Barbados. Known in Porto Rico from the vicinity of Maunabo. (Urban, Add. 2: 163.)

Eugenia pseudopsidium portoricensis. QUIEBRA HACHA.

A small myrtaceous tree, under 25 feet (7.5 meters) in height. Indigenous in numerous localities, Bayamon, Yabucoa, Aibonito, Coamo, Mayaguez, Aguada, Lares, Quebradilla, and Manati. (Urban, Add. 2: 164.) Variety *genuina* of this

species is not known from Porto Rico, and the Porto Rican form is admitted by Urban to be easily distinguished by several characters, there seems little warrant for setting aside De Candolle's species, *E. portoricensis*. According to Stahl, *E. pseudopsidium* is called "guayabillo" and is a tree or shrub 3 to 4 meters high, found rarely in mountains and waste places. (Stahl, 4: 84.)

Eugenia sintenisii Kr. & Urb.

A striking species, with nearly round, heavily veined leaves. (Urban. Add. 2: 168.)

Eugenia serrasuela. SERRASUELA.

An indigenous myrtaceous tree of doubtful specific identity. Reported by Bello only from the vicinity of Anones as *Eugenia costata*. (Urban. Add. 2: 170.)

Eugenia stahlii.

Called "guayabota" and "limoncillo." A tree 60 feet (18 meters) in height, indigenous and peculiar to Porto Rico, Luquillo, Adjuntas, Penuelas. A smaller closely related species, *E. sintenisii*, is known only from El Yunque and from the Sierra de Naguabo. (Urban, 2: 168.)

Eugenia tetrasperma. GUASABARA.

Tree or shrub 5 meters high; native in the forests. (Stahl, 4: 80.)

Eupatorium dalea. GUERRERO.

A composite shrub 6 to 10 feet (2 to 3 meters) high; a rum made from the leaves and flowers is used in medicine. (Stahl, 5: 114, as *Critonia dalea*.)

Eupatorium macrophyllum. TURMA TORO.

An erect, herbaceous annual, 1 meter high, frequenting mountain shades. (Stahl, 5: 109.)

Eupatorium odoratum. SANTA MARIA.

A woody herb, 1 to 2 meters high, annual, found in waste places. (Stahl, 5: 111.)

Eupatorium polyodon. SANTA MARIA.

A recently described shrub of 2 to 6 meters, from Bayamon, Cayey, Aibonito, Maricao, and Mayaguez. It is *Eupatorium triste* of Stahl, which was not the true species of De Candolle. (Urban. Symb. 1: 462.)

Eupatorium portoricense. GUERRERO.

A shrub or tree, from 2 to 5 meters high. It grows in the mountains as well as in the thickets along the seashore, and has been introduced into cultivation. (Urban, Symb. 1: 459.)

Eupatorium resinifluum. GUERRERO.

A shrubby species attaining only 1 or 2 meters; known from Adjuntas and Maricao. (Urban, Symb. 1: 462.)

Eupatorium triste. See *Eupatorium polyodon*.

Eupatorium villosum. OREGANILLO.

A shrub 1 to 4 meters high. The variety *dolicholepsis* is known from several localities in the north and northwestern parts of the island. (Urban, Symb. 1: 462.)

Euphorbia.

A large genus of the Euphorbiaceae or spurge family, represented by numerous species in Porto Rico. These are mostly small herbs or low shrubs of no economic value.

Euphorbia buxifolia.

Family Euphorbiaceae: shrubby, 1 foot high (30 centimeters), frequenting the rocky seashore near Guinaca.

Euphorbia heterophylla. MARAVILLA.

Cultivated at Juana Diaz as an ornamental. The base of the upper leaves are light red, as in *Poinsettia*.

Euphorbia portoricensis.

A very small plant with subreniform pubescent leaves; recently named by Urban on specimens from near Cabo Rojo.

Euterpe.

A genus nearly related to that of the royal palm, *Roystonea*. An unnamed species is reported by Eggers as growing gregariously on El Yunque, at an altitude of 1,500 to 3,000 feet. This is doubtless the species known in the north-eastern part of the island as "palma de la sierra," which is said to be very similar to the royal palm, if not indeed identical with *Roystonea oleracea*, the so-called "cabbage palm."

Exostemma caribaeum. CUERO DE SAPO.

A rubiaceous shrub. (Stahl, 5: 48.)

Exostemma floribundum. See *Antirrhoea coriacea*.**Fagara.**

A genus of Rutaceae, including many species of West Indian trees formerly referred to *Zanthoxylum*; several of these are indigenous in Porto Rico.

Fagara caribaea. ESPINO RUBIAL.

An indigenous shrub or tree, 7 to 20 meters high. Reported from the western part of the island, at Peñuelas, Sabana, Guanica, Cabo Rojo, Rincon, and Aguadilla. (Urban, Add. 3: 251.)

Fagara flava. SATINWOOD.

Indigenous; 3 to 10 meters; known also from the Florida Keys and from several of the smaller Antilles. (Urban, Add. 3: 258.)

Fagara martinicensis.

Also called "espino," "espino rubial," and "cenizo." An indigenous tree of 5 to 20 meters. Of general distribution from Cuba to Trinidad, and known from mountain forests in numerous localities in Porto Rico. The root is considered medicinal. (Urban, Add. 3: 252.)

Furnishes a wood moderately hard, white in color, that breaks with a vertical fracture and is used in the construction of houses in the country. The bark contains a yellow dye. (Grosourdy, 2: 375.)

Fagara microphylla. See *Zanthoxylum fragodes*.**Fagara monophylla.**

Variouly named "carubio," "mapurito," "rubia," and "espino rubial." A tree of 2 to 10 meters, indigenous in forests and thickets in coast or foothill regions throughout the island. (Urban, Add. 3: 256.)

Fagara pterota. ESPINO RUBIAL.


A wild tree, produced in abundance over nearly the entire island, 40 to 50 feet (12 to 15 meters) high and 12 inches (30 centimeters) in diameter. The wood is rather strong and solid, greenish-yellow or smuff-colored; used for doors, floors, windows, etc. The military authorities of San Juan used it for making chests. It is also used in ships' bulwarks. (Grosourdy, 2: 382.)

Fagara spinifex. NIARAGATO.

A shrub or small tree of 1 to 5 meters. Reported from San Ildefonso, near Coamo; also from Guanica and the river Juey. (Urban, Add. 3: 265.)

Fagara trifoliata.

A shrub or small tree, 2 to 8 meters, known from the vicinity of Guanica. The distribution of the species extends southward to Trinidad. (Urban, Add. 3: 272.)



JAGUEY (FICUS LENTIGINOSA). ROAD BETWEEN PONCE AND ADJUNTAS.

Falsa belladona. See *Solanum seaforthianum*.

Falso guaco. See *Willughbaeya scandens*.

Falso marrubio. See *Maecosphaerum polystachyum*.

Faramea odoratissima. PALO DE TORO.

Family Rubiaceae; a tree or shrub 3 meters high, found in mountains and waste places; flowers fragrant. The name "cafetillo" is recorded by Bello. (Stahl, 5: 66.)

Farolitos. See *Cardiospermum halicacabum*.

Fevillea cordifolia. SECUA.

Also called "pepita amarga," and in Jamaica "antidote cocoon" and "sabo." Family Cucurbitaceae, a woody biennial creeper. In Jamaica an oil has been extracted from the seeds and used in lamps. The seeds are also used as an antidote in poisoning and as a remedy for dropsy.

Ficus arbutifolia.

Family Moraceae; said to be cultivated along the roadside at Arroyo (Sintenis).

Along the entire road from Aguadilla to Arecibo two species of fig were common. This part of the country offers much waste land in addition to the roadsides, which might be planted in rubber or any similar crop.

Ficus carica. FIG. HIGO.

A few fig trees were seen near Santurce, but they were not in a thriving condition, and the probabilities of success on the moist north coast seem to be slight.

Ficus crassinervis.

Near Naguabo.

Ficus lentiginosa. JAGUEY.

PLATE XXXV.

The fruits are light green, spotted with dull red, and are much larger than those of *F. pedunculata*. Both species are planted for shade along the road between Ponce and Adjuntas.

Ficus pedunculata.

According to Grisebach, this species is called "Jamaica cherry." It has been reported from south Florida and many of the West Indian islands, but was originally described from St. Christopher.

Ficus portoricensis.

A native fig tree attaining a height of 10 meters; from Yabucoa and Manati. (Urban, Symb. 1: 472.)

Fideos. See *Cuscuta americana*.

Fig. See *Ficus carica*.

Fit weed. See *Eryngium foetidum*.

Flamboyán. See *Poinciana regia*.

Flamboyant boanco. See *Bauhinia kappleri*.

Fleurya estuans. PICA-PICA.

A common weed about Santurce; belongs to the nettle family with stinging hairs.

Flor de agua. See *Castalia*.

Flor de muerto. See *Clerodendron fragrans*.

Forestiera segregata.

Family Oleaceae; a shrub or small tree; native in Florida and the northern Antilles, including Porto Rico.

Forsteronia corymbosa. BEJUCO DE SAN JUAN.

Reported by Bello.

Forsteronia floribunda.

Family Apocynaceae; a trailing shrub said to yield rubber.

Framboyant. See *Poinciana regia*.

A local mispronunciation of flamboyant.

Frangula sphaerocarpa.

Family Rhamnaceae; a shrub or small tree from near Lares. (Grisebach, 99.)

French Jasmine. See *Calotropis procera*.**Frijol cimarrón.** See *Vigna capensis*.**Frijol marrullero.** See *Vigna glabra*.**Frijoles.** See *Vigna catjang*.

In Central America and Mexico this name is applied to the common black bean. *Phaseolus vulgaris*.

Frijoles caballeros. See *Dolichos lablab*.**Frijolillo.** See *Dolicholus reticulatus*.**Frijolillo colorado.** See *Dolicholus phaseoloides*.**Frijolillo de ñame.** See *Pachyrrhizus angulatus*.**Frijolillo pequeño.** See *Dolicholus minimus*.**Fruta de pan.**

A tree from all parts of the island; height, 30 to 40 feet (9 to 12 meters); diameter, 15 to 20 inches (37 to 45 centimeters); wood white, very soft; specific gravity 0.36; not used. (Exp. 1857.)

Furcraea gigantea. MAURITIUS HEMP.

Family Amaryllidaceae. A large agave-like plant native in Mexico, but now generally known as Mauritius hemp from the fact that its culture was taken up on a large scale in Zanzibar. Fourcroya will grow on dry sandy land where few other crops of value can be raised, but as in the case of sisal hemp there is little probability that it would be desirable to plant extensively in Porto Rico. The species is already introduced into the island and is occasionally planted as an ornamental instead of agave.

Fustic. See *Machura tinctoria* and *Zanthoxylum*.**Gaita.** See *Hypelate paniculata*.

A tree from the interior of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 16 to 18 inches (40 to 45 centimeters). Wood white, rather hard; specific gravity, 0.753; used for cabinet work. (Exp. 1857.)

Galactia.

A genus of leguminous creeping or twining vines. Stahl describes *Galactia berteriana* and *G. filiformis*. Sintenis collected the former near Bananquitas and the latter near Cabo Rojo. (Stahl, 3: 66-67.)

Galactia tomentosa.

A recently described species known from between Aibonito and Coamo. (Urban, Symb. 1: 472.)

Galan arbóreo. See *Acnistus arborescens*.**Galan del monte.** See *Cestrum laurifolium*.**Gallina.**

"Chicken wood." Much used for machete handles. A very hard, yellow wood. (Hansard.)

Gallito. See *Agati grandiflora*.**Gandul.** See *Cajanus cajan*.**Gangolin.** See *Ilex sideroxyloides portoricensis*.

Garcinia mangostana. MANGOSTEEN.

Various attempts have been made to introduce this noble fruit into the West Indies, and a tree supposed to be the largest in those islands has been growing for many years in the botanic gardens of Trinidad. It began to fruit in 1875 and has continued to do so at irregular intervals. Seedlings raised from it have proved, however, to be very weakly and have not reached maturity, a failure due, in the opinion of Superintendent Hart, to the lack of cross fertilization. Additional trees have recently been secured from the East in the hope of meeting this deficiency and making possible the more rapid multiplication of the species, for which the natural conditions are considered entirely favorable.

Mr. F. L. Cervantes, of Habana, has recently called our attention to the fact the former existence of the mangosteen in Cuba was reported by the French botanist Descourtilz. It was introduced from the East Indies by a wealthy French planter of Santiago de Cuba, reached maturity, and fruited, but is now believed to have become extinct. It is also reported that the mangosteen is grown in the Philippines, especially at Jolo, whence it is shipped in some quantity to Manila.

Garden egg.

A Jamaica name for the egg plant, *Solanum melongena*.

Gardenia florida. CAPE JASMINE.

Family Rubiaceae; introduced as an ornamental; reported from Yabucoa.

Garlic. See *Allium sativum*.**Garracho.** See *Quararibea turbinata*.**Garrocho.**

A tree from all parts of the island; height, 30 to 38 feet (9 to 12 meters); diameter, 6 to 8 inches (15 to 20 centimeters). Wood white, rather hard; specific gravity, 0.713; not used. (Exp. 1857.)

Gateado.

A tree from the interior of the island; height, 18 to 20 feet; diameter, 10 to 13 inches (25 to 33 centimeters). Wood striped red, hard; specific gravity, 1.164; used for cabinet work. (Exp. 1857.)

Grosourdy (2: 384) refers this wood to *Brosimum guianensis*. He gives the height as 20 to 30 feet (6 to 9 meters), and says that the samples from Porto Rico are red, striped with black, and would seem to be susceptible of a fine polish.

Gato. See *Helicteres jamaicensis*.**Gen gen.**

Captain Hansard gives for this wood a specific gravity of 0.68. It has been used for making the wooden pans for gold seekers in the Luquillo mountains.

Gengibre. See *Zingiber officinale*.**Gengibre amargo.**

The "bitter ginger" is probably *Zingiber zerumbet*. It has large coarse roots, and is used as an external remedy in the form of a tincture made with rum.

Gengibrillo.

Enumerated by Hill among the "trees" used for dyeing and tanning. In Ponce we found this name applied to small, fleshy rootstocks having a yellow, bitterish juice; possibly they may be those of some species of *Curcuma*. The juice is said to be used for dyeing and marking handkerchiefs. The dealers obtain their supplies from the hills, where the plant is believed to grow wild.

Genipa americana. JAGUA.

A rubiaceous forest tree native in South America and the West Indies. It should not be confused with the genip tree *Melicocca bijuga*, of family Sapin-

daceae, which is called "genipe" in Porto Rico. The fruit of genipa is from 4 to 6 inches (10 to 15 centimeters) in length and oval in shape. The skin is somewhat leathery in texture and of a russet color, irregularly mottled with grayish. The section shows it to be two-celled. The rind and the pulp, in which the seeds are embedded, are used in the preparation of one of the refreshing drinks which are such a feature in Porto Rico. The "jagua" is also occasionally eaten, but can scarcely be recommended. It has somewhat the flabby texture and much of the taste of dried apples; according to Schumann, it contains large quantities of tannic acid.

It reaches a height of 50 feet (15 meters), with a long, rather straight trunk, from 15 to 20 inches (37 to 45 centimeters) in diameter. It furnishes a rather soft, light wood that is, however, strong and resistant, flexible, and fibrous in texture. The color is uniformly whitish, tinged with a light reddish brown. It resembles the ash and possesses also its properties. It is very suitable for boxes and for any work where strength and elasticity are needed. The rims of sieves and shoe lasts are made of this wood. The military authorities of San Juan used it for ammunition chests and for hoops to barrels. Specific gravity, 0.873. (Grosourdy, 2: 391.)

Geno. See *Lonchocarpus glaucifolius*.

Geno geno. See *Lonchocarpus violaceus*.

Captain Hansard says that the wood of this tree has a specific gravity of 0.89.

Geófila. See *Geophila reniformis*.

Geo geo.

A tree from all parts of the island; height, 25 to 40 feet (7 to 12 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood dark, soft; specific gravity, 0.511; not used. (Exp. 1857.)

Geophila reniformis. GEOFILA.

Family Rubiaceae; a creeping herb frequenting forest shades in moist places. (Stahl, 5: 81.)

Gerardia domingensis. YERBA VERONICA.

Family Scrophulariaceae; an herbaceous annual in sandy inundated places. (Stahl, 6: 233.)

Gesneria citrina.

Family Gesneriaceae; a new species from Utuado. (Urban, Symb. 1: 477.)

Gesneria portoricensis. YERBA PARRERA.

Reported by Bello, who gives De Candolle as the author of the species, which does not, however, appear in the Index Kewensis.

Gia mansa. See *Casearia parvifolia*.

Gia verde. See *Casearia stipularis*.

Gilibertia arborea. VIBONA.

Also known as "pana," "palo cachumba," and "muñeca." A tree of 4 to 20 meters, belonging to the family Araliaceae. Reported from forests in numerous localities in Porto Rico. (Urban, Symb. 1: 20; as *Sciadophyllum jacquini*; Stahl, 4: 197.)

Gilibertia laurifolia.

Native names: "Palo de vaca," "palo de gangulin," "palo cachumba" and "vibona." A tree of 8 to 20 meters known from native forests near Aibonito, Jayuya, Barranquitas, Peñuelas, and Adjuntas. (Urban, Symb. 1: 203.)

Ginger. See *Zingiber officinale*.

Ginoria rohrii.

Family Lythraceae; from Naguabo, Guayacan, and Guanica.

Glateado. See *Coccolobis laurifolia*.

Gnaphalium americanum. See *Gnaphalium purpureum*.

Gnaphalium purpureum. YERBA LANUDA.

An erect, herbaceous, annual composite on the borders of cane fields and in ditches; 25 centimeters high. (Stahl, 5: 147.)

Goat weed. See *Capraria biflora*.

Gomatu palm. See *Arenga saccharifera*.

Gomidesia lindeniana. CIENEGUILLO.

An indigenous, myrtaceous shrub or small tree 30 feet (9 meters) high, reported from forests between Bayamon and Aguas Buenas; also from Aibonito, Barranquitas, Adjuntas, and Lares.

Gomphya nitida. See *Ouratea nitida*.

Gonfia lustrosa. See *Ouratea nitida*.

Gongoli.

A tree from all parts of the island: height, 35 to 40 feet (11 to 12 meters); diameter, 10 to 12 inches. Specific gravity, 1.088. (Exp. 1857.)

Furnishes a strong, compact, flesh-colored wood that breaks with an oblique fracture. Its common use is for the framework of houses and for fences. (Grosourdy, 2: 384.)

Gonolobus. See *Vincetoxicum*.

Gonzalagunia spicata. RABO DE RATON.

Family Rubiaceae; an erect or somewhat trailing woody annual or biennial, 1 meter high, common in fence rows and in waste places partially shaded. The white or slightly pinkish flowers are all turned to the upper side of the long curved raceme. The fruit is a whitish berry, but turns black on drying and appears capsular. (Stahl, 5: 43, as *Gonzalea spicata*.)

Gonzalea spicata. See *Gonzalagunia spicata*.

Gossypium barbadense. COTTON. ALGODON.

Cotton is not raised in Porto Rico as a commercial crop, and indeed none is to be seen about the country except now and then a shrub in a dooryard.

The cotton plant is also one of many, of tropical origin, which have been selected and improved in subtropical or temperate regions and which tend to return to their original habits when taken back to the more humid tropics. The cottons of tropical countries, of which there are several species, are perennials, and scarcely begin to bear during a period sufficient for ripening a crop in the Southern States. There is, accordingly, little fear that any tropical country will successfully compete with warm temperate regions like Egypt and our Southern States, where the crop can be brought to early and even maturity so that the the necessary labor can be applied with advantage and economy.

The name "algodon rojo" (red cotton) is said to be the variety described botanically as *Gossypium purpurascens*.

Gossypium janiphaefolium. ALGODON DE YUCA.

A species of cotton described by Bello from the vicinity of Cabo Rojo, called "algodon de yuca" or "cassava cotton," on account of the similarity of the leaves to those of *Manihot*. Stahl reduces the species to *Gossypium herbaceum*. (Stahl, 2: 96.)

Gossypium purpurascens. See *Gossypium barbadense*.

Gouania domingensis.

Family Rhamnaceae; a shrubby vine, growing also in Jamaica, where it is called "chaw stick" and used for toothbrushes, etc. It yields a valuable stomachic drug, according to Grisebach. A variety *aptera* is reported by Urban from Coamo and Aibonito.

Gouania tomentosa.

Reported from near Aguada.

Gout tea. See *Cordia globosa*.

Graciosa. See *Oldenlandia glomerata*.

Grana.

Given in Maza's Diccionario as the Porto Rican name of the royal palm, but apparently little used, "palma de yagua" being the prevalent designation.

Granada. See *Punica granatum*.

Granadilla. See *Passiflora quadrangularis*.

Granadilla tree.

In Jamaica and Cuba the names "granadilla tree" and "granadilla" are applied to *Brya ebenus*, a small leguminous tree yielding the so-called "American ebony." This species is not yet reported from Porto Rico. Captain Hansard describes the "granadilla tree" as very large, with small leaves and spreading branches. Nothing will grow underneath it, and it is therefore unwelcome in coffee plantations. The yellow wood is not considered of marketable value, although it will make fairly good boards. The specific gravity is 1.11. The "granadilla" fruit is from a vine *Passiflora quadrangularis*, and has nothing to do with the tree.

Granadillo.

A tree from all parts of the island; height, 40 to 45 feet (13 to 14 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood, yellow, hard; specific gravity, 0.917; used for cabinet work. (Exp. 1857.)

This is also enumerated by Hill among trees used for dyeing and tanning. See also the note under *Granadilla tree*.

Grape. See *Vitis vinifera*.

Grape fruit. See *Citrus decumana*.

Grape tree. See *Coccolobis*.

Grasilla. See *Utricularia subulata*.

Grayume. See *Didymopanax morototoni*.

Grayumo. See *Didymopanax morototoni*.

Captain Hansard states that this tree is supposed to have two distinct sexes, the female being hollow, while the male is solid and is sometimes sawed into white narrow boards, although he has seen some 14 inches wide. Owing to the softness and lightness of the wood, it is used only for inside work. The tree is large, with parasol-shaped leaves, and the name is sometimes erroneously pronounced "yau-grumo."

Ground cherry. See *Physalis*.

Ground nut. See *Arachis hypogaea*.

Guaba. See *Inga vera*.

A most useful tree for shade of coffee; according to Captain Hansard the specific gravity of the wood is 0.64.

Guabara.

The wood is given a specific gravity of 0.63 by Captain Hansard.

Guácima del Norte. See *Guazuma ulmifolia*.

Guácima del Sur. See *Guazuma tomentosa*.

Guaco. See *Willughbaeya odoratissima* and *W. cordifolia*.

Guaco de cabra. See *Willughbaeya porosa*.

Guajacum officinale. LIGNUM VITAE. GUAYACAN.

Family Zygophyllaceae; a tree 5 meters high, wood called "lignum vitae." The wood of this tree is very hard, cross-grained, and of high specific gravity. It

takes a fine polish and is used in making rollers for casters and various utensils. The resin exuded from the stem called "guaiacum" is said to have the property of changing color when brought in contact with various substances, also used as a stimulant in medicine. (Stahl, 2: 173.)

A wild tree met with on the coasts, reaching a height of 45 or 50 feet (14 or 15 meters), with a trunk 12 to 15 inches (30 to 37 centimeters) in diameter. It furnishes a very hard wood, very fine-grained and compact. The sapwood which is almost as hard as the heart is light yellow, while the heart is brown mixed with green with darker stripes. It is a valuable wood and its exportation forms a branch of commerce. It cuts well with the saw and in spite of its fibrous texture is much worked. It can also be split, especially when recently cut, at which time the wood is much softer and more easily worked. It becomes much harder on exposure to the atmosphere and is then very difficult to work.

It is used extensively for machinery and in cabinet work and for turned articles, and in all works where strength and hardness are needed, such as wooden cogs, mortars, hubs of wheels, etc. It is susceptible of a high polish. The green wood has a specific gravity of 1.080; the dry, 1.354 to 1.360. (Grosourdy, 2: 389.)

It is believed in Porto Rico that coffee powdered with a mortar and pestle of "guayacan" is far superior to that ground in a wooden mill.

Guajacum sanctum. GUAYACANCILLO.

Family Zygothylaceae; a tree said to have been abundant formerly on the coast hills of the south side of the island, but now nearly exterminated as far as large trees are concerned. The wood is considered valuable, and a resin or balsam exuded from it is medicinal. (Stahl, 2: 174.)

According to Engler and Prantl the wood of this species is also utilized in commerce as *lignum vitae*.

Guajanilla amarilla.

A wild tree produced in abundance in the southern part of the island. Its height is about 40 feet (12 meters), and its trunk 12 to 15 inches (30 to 37 centimeters) in diameter. The wood is strong, dark yellow, breaking with a vertical fracture. Its most common use is for shelving in houses. (Grosourdy, 2: 386.)

Guajanilla prieta.

Said to be a variety of guajanilla amarilla, distinguished only by having its dark yellow wood streaked with brown. (Grosourdy, 2: 386.)

Guajanillo.

Captain Hansard gives the specific gravity as 0.80 and states that it is the same as "caracolillo."

Guajona.

A large tree occurring rarely in the mountains of the Luquillo region; the wood makes good boards. The name means gypsy. (Hansard.)

Gualteria americana. See *Waltheria indica*.

Guamá. See *Inga laurina*.

Specific gravity, 0.70 (Hansard). Used for firewood and as a coffee shade. The fruit is a broad, bean-like pod with edible, rather sweet-tasting, fluffy pulp. Abundant in foothills.

Guanábana.

Soursop. The fruit of *Anona muricata*.

Guanabancilla cimarrona. See *Vincocharicum*.

Guanabanillo.

A tree from all parts of the island; height, 20 to 25 feet (6 to 7.5 meters); diameter, 10 to 12 inches. Wood light, soft; specific gravity, 0.728; not used. (Exp. 1857.)

Guanábano. See *Anona muricata*.

A tree from all parts of the island; height, 25 to 30 feet (7.5 to 9 meters); diameter, 12 to 15 inches. Wood, light, soft; specific gravity, 0.688; not used. (Exp. 1857.)

Captain Hansard gives this wood a specific gravity of 0.44.

Guanábano cimarron. See *Anona montana* and *A. palustris*.

Guanávano. See *Anona muricata*.

A less correct form of "guanábano."

Guanaguao.

A tree from all parts of the island; height, 60 to 70 feet (18 to 21 meters); diameter, 60 to 65 inches (150 to 160 centimeters). Wood flesh-colored, soft; specific gravity, 0.741; not used. (Exp. 1857.)

Guano. See *Ochroma lagopus*.

Silk-cotton tree with long, fluffy pods sold in market; grows profusely in British islands.

A tree from all parts of the island; height, 25 to 30 feet (7.5 to 9 meters); diameter, 18 to 20 inches (45 to 50 centimeters). Wood, yellow very soft; specific gravity, 0.231; little used. (Exp. 1857.)

Guansa.

A tree from all parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood ash-colored, rather hard; specific gravity, 0.729; not used. (Exp. 1857.)

Guao. See *Comocladia*.

According to Engler this name is applied to *Comocladia ilicifolia*, an anacardiaceous tree, also known as "carrasco" in Porto Rico and other West Indian islands. Hill's reference (p. 34) is evidently an error for "guano," the popular name of *Ochroma lagopus*, the West Indian cork-wood tree, which furnishes a downy fiber.

Guara. See *Cupania americana*.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood white, soft; specific gravity, 0.596; not used. (Exp. 1857.)

Captain Hansard gives a specific gravity of 0.70.

Guaraguadillo. See *Guarea humilis*.

Guaraguao. See *Guarea swartzii*, *G. trichilioides*, and *Trichilia spondiodes*.

A tree occasionally found, but not plentiful, in the mountains of the northeastern part of the island. The wood is red and makes good boards, similar to mahogany; specific gravity, 0.69 (Hansard).

Light reddish brown, streaked with lighter and darker shades. Large ducts numerous, occurring singly or in irregular loose groups of two to four. Medullary rays numerous, but very indistinct. An exceedingly cross-grained, porous wood, somewhat similar in color to cedro. Suitable for a cabinet wood. (Hill and Sudworth, p. 29.)

Grosourdy (2: 387) refers this variety to both *Guarea trichilioides* and *Trichilia moscata* and describes it as an abundant wild tree, 70 feet (21 meters) in height, whose long, straight trunk reaches a diameter of 6 or 6½ feet (about 2 meters). It furnishes a rather hard wood of good weight, dense and close-grained, dark flesh-colored, and breaking with an oblique fracture. It is much appreciated for the construction of strong wagons and carriages and farm implements and in many branches of carpentry work.

Guaraguao cimarron. See *Trichilia haitiensis*.

Guarea humilis. GUARAGUADILLO.

Family Meliaceae: a tree or shrub 5 meters high from Barranquitas and Adjuntas, Luquillo, and Naguabo. (Stahl, 2: 168.)

Guarea swartzii. GUARAGUAO.

A large forest tree, 15 to 20 meters high. Wood like cedar in color, but harder; used for furniture. (Stahl, 2: 166.)

This species is not now recognized as genuine, the Porto Rican form representing probably *G. trichilioides*.

The common name "acajou" is sometimes applied to this species.

Guarea trichilioides.

A tree having a musk-like perfume.

Guarema. See *Picramnia pentandra*.

A tree from the northeastern part of the island: height, 30 to 35 feet (9 to 11 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood dark, hard; specific gravity, 1.213; used in building houses. (Exp. 1857.) (Grosourdy, 2: 387.)

Guasa.

A tree from the eastern part of the island; height, 30 to 35 feet: diameter, 10 to 12 inches. Wood white, soft; specific gravity, 0.890; used in building houses. (Exp. 1857.)

Grosourdy says (2: 387), in addition, that the wood is resistant and breaks with a vertical fracture.

Guasábara. See *Eugenia tetrasperma*.

A tree from all parts of the island: height, 55 to 60 feet (15 to 18 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood flesh-colored, hard; specific gravity, 0.947; used for cabinet work. (Exp. 1857.)

Specific gravity 0.87 according to Captain Hansard.

Guasabarillo.

Perhaps the same as "guasábara," but with the specific gravity given as 0.78.

Guasávara. See *Eugenia aeruginosa* and *E. eggertii*.

Another spelling of Guasabara; said by Grosourdy to be an abundant wild tree, 60 feet (18 meters) in height and 12 to 18 inches (30 to 37 centimeters) in diameter. The wood is strong, compact, and flesh-colored, and breaks obliquely. Its most common use is for fences and for fuel. (Grosourdy, 2: 387.)

Guásima. See *Guazuma guazuma*.

A tree from all parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood white, soft; specific gravity, 0.741; not used. (Exp. 1857.)

Captain Hansard gives a specific gravity of 0.35 under this name.

Guasimilla.

A wild tree, without doubt a variety of Guacima, 35 feet (11 meters) high and 12 inches (30 centimeters) in diameter. The wood is white, soft, and breaks with a vertical fracture. Its most common use is for framing country houses. (Grosourdy, 2: 387.)

A tree from the eastern part of the island; specific gravity, 0.508. (Exp. 1857.)

Guatapaná.

A wild tree, 24 to 30 feet (7 to 9 meters) high, with a trunk 12 inches (30 centimeters) in diameter. The wood is dark-colored, hard, and resistant; used for shelving and in boat building. (Grosourdy, 2: 388.)

A tree from the southern part of the island; specific gravity, 1.294. (Exp. 1857.)

Guauro. See *Vincetoxicum pubescens*.**Guava.**

The English name of the well-known tropical fruit *Psidium guajava*, the Spanish name of which is "guayava." In Porto Rico the name guava is applied

instead to a species of *Inga*, used as a shade tree for coffee. Captain Hansard claims to have been informed by Professor Hart, of the Trinidad Botanic Gardens, that the tree called guava in Porto Rico is not the genuine *Inga vera*, but our specimens seem to agree reasonably well with material from the other West Indies and from South America.

Guayaba. See *Psidium guajava*.

Guayabacison.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 15 to 18 inches (37 to 45 centimeters). Wood pinkish, hard; specific gravity, 1.029; used in building houses. (Exp. 1857.)

Guayabacon. See *Eugenia aeruginea*, *Myrica diraricata*, and *M. leptoclada*.

A tree of the coasts, reaching 30 feet (9 meters) in height by 18 inches (45 centimeters) in diameter. The rose-colored wood is compact and strong, and difficult to break. It is used for beams and rafters in country houses. (Grosourdy, 2: 388.)

Guayabillo. See *Eugenia pseudopsidium*.

Guayabillo de costa. See *Eugenia flavo-virens*.

Guayabo.

A tree from all parts of the island; height, 12 to 15 feet; diameter, 6 to 8 inches. Wood flesh-colored, hard; specific gravity, 0.811; fruit tree. (Exp. 1857.)

Guayabota. See *Diospyrus ebenaster* and *Eugenia stahlii*.

A hard wood used in peasant houses for posts; flowers like *Gardenia*. Specific gravity of the wood, 0.66. (Hansard.)

Said by Grosourdy to be a variety of "guayabota rojo," with a lighter-colored wood. (Grosourdy, 2: 388.)

Guayabota blanca.

A tree from all parts of the island; height, 20 to 30 feet (6 to 9 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood pinkish, soft; specific gravity, 0.686; not used. (Exp. 1857.)

Guayabota nispero. See *Maba sintenisii*.

Guayabota roja.

A rather abundant wild tree, reaching 30 feet (9 meters) in height, and 12 inches (30 centimeters) in diameter. The wood is red, rather soft, and not very durable, breaking with an oblique fracture. Its common use is for the framework of country houses. (Grosourdy, 2: 388.)

A tree from all parts of the island. Specific gravity, 0.667. (Exp. 1857.)

Guayacan. See *Guajacum officinale*.

Also called erroneously "guayavacan" by the peasants and others who have learned from them.

A tree from the southeastern part of the island; height, 40 to 45 feet; diameter, 12 to 15 inches. Wood greenish, hard; specific gravity, 0.972; used in making furniture. (Exp. 1857.)

Captain Hansard considers this the genuine *lignum-vitae* and gives the specific gravity as 1.16.

Heartwood dull yellowish-brown, with dark olive-brown streaks; sapwood pale yellow, with brownish areas. Minute ducts occurring singly and evenly scattered. Medullary rays very small and indistinct. Smoothed surface, oily to the touch. Exceedingly hard, brittle, and difficult to cut. Resembles *Guajacum officinale*, and is probably *G. sanctum*. It grows in comparative abundance in the entire mountain chain and on the southern coast of the island, producing a wood which is very solid and resistant. On this account it is much sought after in the ship-



GUÁCIMA ('GUAZUMA GUAZUMA'). COAMO SPRINGS.

yards for blocks and pulleys, for spokes and tires, and many other things which require great strength. Water boiled with this wood is an antivenereal specific and cures ulcers produced by this disease. The resin from the "guayacan," *lignum-vitae*, is highly valued for gout. The Caribbeans apply it to other local ailments, and foreigners extract it freely on the southern coast equally with the "ucar," "expinillo," "palo de Maria," "palo de Brazil" and others, for dyes and coloring matter. (Hill and Sudworth, 27.)

Guayacancillo. See *Guajacum sanctum*.

Grosourdy (2: 389) refers this to *G. verticale* and suggests that it is possibly only a variety of *G. officinalis*.

Guayacan de vera.

A name applied at Yauco to guayacan wood with white and yellow spots.

Guayaco.

Reported by Hill among resin-yielding trees. Perhaps "guayacan" was intended.

Guayánilla amarilla.

A tree from all parts of the island; height, 40 to 45 feet; diameter, 12 to 15 inches. Wood yellow, hard; specific gravity, 1.088; used for cabinetwork. (Exp. 1857.)

Guayánilla prieta.

A tree from the interior of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood dark yellow, hard; specific gravity, 1.116; not used. (Exp. 1857.)

Guayarote. See *Meliosma obtusifolia*.

Guayava. See *Psidium guayava*.

Guayava pera. See *Psidium guayava*.

A name applied to the pear-shaped varieties of the guava.

Guayavacon. See *Myrcia leptoclada* and *Trichilia hirta*.

Guayavita. See *Amomis caryophyllata*.

Guazuma guazuma. GUÁCIMA DEL NORTE.

PLATE XXXVI.

A common tree belonging to the family Sterculiaceae; Stahl ascribes qualities similar to those noted under *G. tomentosa*, and gives the name "guácima del norte." (Stahl, 2: 105.)

A wild tree very common throughout the Antilles and also on the continent. It reaches 30 to 40 feet (9 to 12 meters) in height, with a rather straight trunk 15 to 18 inches (37 to 45 centimeters) in diameter. A variety of the mountains with rather small leaves is somewhat larger and taller. Furnishes a rather light wood that is undoubtedly resistant, fibrous in texture, light gray in color, slightly mixed with rose or flesh color in unequal stripes that produce a handsome appearance. This wood has no heart, and some specimens are reddish-gray mottled almost uniformly. It has no use, but would appear suitable for fences, the interior of houses, etc. It serves for fuel, and the light charcoal is good for making powder. Specific gravity, 0.552. The common name is sometimes spelled "guázima" and "guásima." (Grosourdy, 2: 385.)

Guazuma tomentosa. GUÁCIMA DEL SUR.

A tree common on the southern coast of the island. (Stahl, 2: 106.) In Jamaica is said to be a tree 20 to 25 feet (6 to 8 meters) high; foliage and fruit eaten by cattle. The light wood is used for making sugar hogsheads.

Guazuma ulmifolia. See *Guazuma guazuma*.

Guerrero. See *Eupatorium dalea*, *E. portoricense*, and *E. resinifluum*.

Guettarda krugii.

A recently described rubiaceous tree, attaining a height of 10 meters; known from the vicinity of Guanica. (Urban, Symb. 1: 431.)

Guettarda laevis.

A tree 15 to 20 meters high from the primeval forests of the mountains of Luquillo and Naguabo. (Urban, Symb. 1: 433.)

Guettarda ovalifolia.

A tree 8 to 10 meters in height, recently described from the region of Cayey, Adjuntas, and Maricao. (Urban, Symb. 1: 432.)

Guettarda pungens.

The midrib of the small leathery leaves of this species is produced into a sharp spine. Known from Maricao. (Sintenis.)

Guettarda scabra. PALO DE CUCUBANO.

A tree 3 to 5 meters high, growing on mountains and in waste places. (Stahl, 5: 56.)

Guichara. See *Lagenaria vulgaris*.

The grooved musical instrument made from the gourd; also called "guira" and "caracho."

Guilandina bonduc. See *Caesalpinia bonduc*.**Guilandina bonducella.** See *Caesalpinia bonducella*.**Guinea grass.** See *Panicum maximum*.**Guinea-hen's weed.** See *Peliveria alliacea*.**Guinea pepper.** See *Capsicum*.**Guineos.**

A variety of banana.

Guingambo. See *Abelmoschus esculentus*.**Guira.** See *Lagenaria vulgaris*.

One of the names applied to the grooved gourd used as a musical instrument.

Guitaran.

A low tree growing on the coast, reaching 18 to 20 feet (5 to 6 meters) in height, with a trunk 10 or 12 inches (25 to 30 centimeters) in diameter. The wood is rose-colored and breaks easily with a vertical fracture. Has no application. (Grosourdy, 2: 389.)

A tree from the interior of the island; height, 18 to 20 feet; diameter, 10 to 12 inches. Wood pinkish, soft; specific gravity, 0.863; used for cabinetwork. (Exp. 1857.)

Gum arabic. See *Acacia arabica*.**Guyabacoa.** See *Rhedia portoricensis*.**Guyacan.** Probably an error for Guayacan.**Guzmannia sintenisii.**

Family Bromeliaceae; from Sierra de Luquillo.

Gynandropsis speciosa. VOLATINES PRECIOSOS.

Family Capparidaceae; an herbaceous annual cultivated in gardens; 1 meter high. (Stahl, 2: 42.)

Gynerium saccharoides. CAÑA DE CASTILLA.**Habas.** See *Phaseolus lunatus*.**Habichuela.** See *Phaseolus vulgaris*.**Habichuela cimarrona.** See *Phaseolus adenanthus* and *Phaseolus lanceolatus*.**Habichuela parada.** See *Phaseolus semierectus*.

Hacana. See *Lucuma multiflora*.

A tree from all parts of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 8 to 10 inches. Wood white, rather hard; specific gravity, 0.924; not used. This is perhaps the same as "acana" and "jacana." (Exp. 1857.)

Hacanilla. See *Ardisia purpurascens*.

This name was perhaps intended for "jacanillo," applied to *Ardisia pendula*.

Hachuelo. See *Pictetia*.

"Color, rich dark yellowish-brown, with streaks and mottlings of light yellow-brown. Ducts minute, very numerous, occurring singly, and irregularly diffused between the medullary rays or often interrupting the rays. Medullary rays very numerous, indistinct. Wood fibers interlaced, but appearing rather straight grained on the finished surface. Valuable for cabinetwork." (Hill and Sudworth.)

Hachuelo.

An erroneous name given by Hill for the calabash tree, *Crescentia cujete*, which seems also to have been confused by him with "tachuelo" or *Pictetia*.

Haematoxylon campechianum. PALO DE CAMPECHE.

Family Cassiaceae; a medium-sized tree yielding the logwood of commerce. (Stahl, 3: 104.)

A wild tree reaching 30 to 45 feet (9 to 14 meters) high, but with short trunk seldom attaining 6 or 9 feet (2 to 3 meters); very crooked and irregular, prolonged into large branches rather long and straight. The diameter of the trunk is 18 to 24 inches (45 to 60 centimeters); of the branches 8 to 10 inches (20 to 25 centimeters). The wood is very hard and heavy, compact in texture, dark purple with darker stripes, frequently tinted with orange, especially toward the outside, which is lighter, variegated with orange. Without doubt at the fall of the tree the wood is of one color, a reddish white, and takes on its beautiful color little by little after exposure to the air. Specific gravity, 0.992.

The wood is not only used as a dye, but can be utilized in turning. (Grosourdy, 2: 369.)

Haemocharis portoricensis. NIÑO DE COTA.

An indigenous tree belonging to the Theaceae and closely related botanically to the tea shrub. It is reported from the primeval forests of the mountain of Luquillo and Naguabo at an altitude of 1,000 meters.

Haenianthus ovatus.

An oleaceous shrub described from specimens collected on El Yunque. (Urban, Add. 1: 58.) Also from the mountain of Luquillo.

Hamelia lutea. BÁLSAMO AMARILLO.

Family Rubiaceae: an annual herbaceous shrub, 1 meter high, found along the foothills and shaded woody places. (Stahl, 5: 40.)

Hamelia patens. BÁLSAMO COLORADO.

An erect biennial shrub, 1 meter high, in stony places in the mountains. Our specimens (No. 619) are from the vicinity of Coamo. (Stahl, 5: 41.)

Hat palm. See *Inodes causiarrum*.

Haya. See *Ocandora laurifolia*.

A tree from the interior of the island; height, 30 to 35 feet; diameter, 8 to 10 inches; specific gravity, 0.705. (Exp. 1857.)

The wood is dirty white, flexible, and breaks with a vertical fracture. Its common use is for oars and the frames of country houses. (Grosourdy, 2: 389.)

Haya blanca. See *Ocandora laurifolia*.

Haya prieta. See *Ocandora virgata*.

Hebeclinium macrophyllum. See *Eupatorium macrophyllum*.

Hecastophyllum brownei. See *Dalbergia hecastophyllum*.

Hecastophyllum monetaria. See *Dalbergia monetaria*.

Heckeria peltata.

PLATE XXXVII.

Family Piperaceae; a large, succulent herb with peltate, heart-shaped leaves.

Heckeria umbellata.

PLATE XXXVIII.

A species appearing very similar to the preceding but very different on close inspection. The stems and petioles have prominent ridges fringed with rows of long hairs; the internodes and peduncles are shorter, the petioles not expanded at base, and the leaves are not peltate but have an open sinus. Both species were collected along the Military Road a short distance south of Rio Piedras.

Hedionda. See *Cassia occidentalis*.

In the market of Ponce roots called "hedionda" were being sold by an herb dealer as a remedy for flatulence.

Hediondilla. See *Leucaena glauca*.

Hedwigia balsamifera. See *Tetragastris balsamifera*.

Hedychium coronarium.

Family Zingiberaceae; reported from Cayey and Maricao. Also collected at Adjuntas.

Hedyosmum arborescens.

Family Chloranthaceae; a shrub 12 to 16 feet (4 to 5 meters) high, in mountain woods. A related species is called "headache weed" in Jamaica. Reported from Sierra de Luquillo.

Hedyotis glomerata. See *Oldenlandia glomerata*.

Heliconia.

A species of *Heliconia* is reported by Hill, apparently on the authority of Eggers, in a list of trees of the mountains. This genus is quite closely related to the banana, but the fruit is a dry capsule with three seeds or less.

Heliconia bihai.

A musaceous tree 10 or 12 feet (3 or 4 meters) high, reported from Bayamon, in forests on the mountains. Called bastard plantain in Jamaica.

Helicteres jamaicensis. CUERNECILLO.

Family Sterculiaceae; a shrub 2 to 3 meters high, in sands of the coast. Jacquin calls this "huevo de gato" (Stahl, 2: 102), while Bello gives simply "gato."

Heliophytum indicum. See *Heliotropium indicum*.

Heliophytum parviflorum. See *Heliotropium parviflorum*.

Heliotropium curassavicum. COTORRERA DE LA PLAYA.

Family Boraginaceae; a woody herb 50 centimeters high, growing in tufts in marshy ground along the seashore. (Stahl, 6: 106.)

Heliotropium indicum. COTORRERA.

An herbaceous, erect annual, 50 centimeters high; common. A decoction of this plant is said to act as a diuretic. (Stahl, 6: 108.) At Cataño this was called "yerba de cotorra," a translation of which would be "parrot grass," while at Ponce it is termed "yerba de culebra."

Heliotropium inundatum. COTORRERA DE AGUA.

Said to be a woody herb, found in wet places along the seashore. (Stahl, 6: 107.)

Heliotropium parviflorum. COTORRERILLA.

A woody herb 50 centimeters high, growing in stony places. (Stahl, 6: 109, as *Heliophytum parviflorum*.)

Heliotropium peruvianum. HELIOTROPO.

A beautiful, fragrant species, cultivated in gardens. (Stahl, 6: 108.)



HECKERIA PELTATA: INFLORESCENCE AND CENTER OF LEAF.



HECKERIA UMBELLATA; INFLORESCENCE AND CENTER OF LEAF.



TALANTALA (HERPETICA ALATA). AT CATAÑO.

Heliotropo. See *Heliotropium peruvianum*.

Henna. See *Larsonia inermis*.

Henriettella fascicularis. CAMACEY SIMPLE.

Family Melastomaceae: a shrub 3 to 4 meters high; grows on mountains above 500 feet elevation. (Stahl, 4: 114.)

Herbeclinium macrophyllum. See *Eupatorium macrophyllum* and *Bacopa chamaedryoides*.

Hernandia sonora.

Family Hernandiaceae: a tree with large peltate leaves, the juice of which is said to be a powerful depilatory; reported from near Rincon and Yabucoa.

Herpestis monniera. See *Bacopa monniera*.

Herpetica alata. TALANTALA.

PLATE XXXIX.

Family Cassiaceae: a shrubby plant 4 to 8 feet high, also called ringworm shrub in Jamaica on account of its reputed healing properties for that disorder. Bello gives the common name as "talantro." (Stahl, 3: 107, as *Cassia alata*.)

Heteropteris bellonis. AHORCA CARALLO.

An indigenous, climbing shrub of the family Malpighiaceae. From Sabana Grande, Maricao, and Lares. Appears in Stahl's "Flora" as *Stigmatophyllum chrysophyllum*.

Heteropteris chrysophylla. BEJUCO DE PARALEJO.

A vine found by Stahl at Humacao. (Stahl, 2: 147.)

Heteropteris laurifolia. PARALEJO VELLUDO.

A shrub 2 to 3 meters high; collected by Heller near Mayaguez. (Stahl, 2: 149, as *Heteropteris pubiflora*.)

Heteropteris pubiflora. See *Heteropteris laurifolia*.

Heteropteris purpurea. BEJUCO DE PARALEJO ROSADO.

A low, woody climber among shrubs. (Stahl, 2: 148.)

Hibiscus abelmoschus. See *Abelmoschus abelmoschus*.

Hibiscus bifurcatus. BUENOS DIAS.

An herbaceous, woody annual, 1 to 2 meters high, found in shady waste places. (Stahl, 2: 91.)

Hibiscus elatus. EMAJAGUA EXCELSA.

A spreading tree growing to be 60 feet high and 8 feet through; used for lumber; does not corrode the nails; very durable; the bark is said to make good rope. (Stahl, 2: 95, as *Paritium elatum*.)

Hibiscus esculentus. See *Abelmoschus esculentus*.

Hibiscus mutabilis. MARAVILLA.

A shrub 3 to 4 meters high, cultivated in gardens for its flowers. (Stahl, 2: 89.)

Hibiscus radiatus. PAVONA ENCENDIDA.

A woody shrub, 1 meter high, cultivated in gardens. (Stahl, 2: 90.)

Hibiscus rosa-sinensis. PAVONA.

A shrub 3 to 4 meters high, cultivated in gardens for its beautiful flowers. Introduced from Asia. At Santurce this species was called "mapola." (Stahl, 2: 88.)

Hibiscus sabdariffa. JAMAICA SORREL. VIÑA.

A shrubby annual or biennial, valued for the fleshy red calyx, which has a sharp but not unpleasant acid taste. The jellies and sauces made from it have been compared to those obtained from cranberries. In British India this plant is the basis of a considerable agricultural industry, it being grown there both for the calyx and for a fiber extracted from the stems by a process of retting. For

this purpose the crop is cut while in flower, dried, made into bundles, and soaked in water for fifteen to twenty days. It is then possible to wash out a strong, silky fiber known in commerce as Rozelle hemp, considered by some to be the equal of jute, although little or none is exported to Europe. The natives use it mostly for cordage and coarser textile purposes. The seeds are employed in various medicinal decoctions and compounds on account of supposed demulcent, diuretic, and tonic properties, and the calices are also made into a refreshing acid drink for fever patients and convalescents.

The calices and young capsules may be preserved by drying, and in this condition are sometimes met with in tropical markets. The leaves are also an ingredient in certain dishes and the seeds are fed to cattle. Analyses show considerable quantities of tartaric and malic acid, but no citric, and the Jamaica sorrel thus supplements the lime in the direction of the qualities of our temperate fruits. It is becoming increasingly popular with the people of Florida, and is now commonly cultivated in the southern part of that State. Americans in Porto Rico would do well to become acquainted with it. The species has been reported by Bello with the above native name as growing in country districts in the western part of the island, but is probably to be found occasionally elsewhere. Also called "aleluya roja de Guinea" and "agrio de Guinea."

Hibiscus schizopetalus. LIRA.

PLATE XL.

A shrub 6 to 8 feet (2 to 2.5 meters) high, of very slender trailing habit; cultivated for its very peculiar pink flowers. The petals are split into a fringe of narrow divisions and the stamen tube and style are extremely long.

Hibiscus tiliaceus. See *Paritium tiliaceum*.

Hibiscus tomentosus. MALVAVISCO APELPADO.

An herbaceous, woody annual; grows in swamps between Cataño and Palo Seco; 1 to 2 meters high. (Stahl, 2: 92.)

Hicaco. See *Chrysobalanus icaco*.

Hicaquillo. See *Dendroprenon caribaeus* and *Loranthus americanus*.

Hicaquillo del combro. See *Dendroprenon bicolor*.

Hierba. See *Yerba*.

Hieronymia alchornoides.

An euphorbiaceous tree, reported from Sierra de Lares.

Higo. See *Ficus carica*.

Higrofila de Puerto Rico. See *Hypophylla portoricensis*.

Higüera. See *Crescentia cujete*.

Hansard gives this wood a specific gravity of 0.51 and doubtfully refers it to the fig tree.

Higüerilla. See *Crescentia cucurbitina*.

Higuerillo. See *Vitex divaricata*.

A tree from all parts of the island; height, 25 to 30 feet; diameter, 10 to 12 inches. Wood ash-colored, hard; specific gravity 0.805; used for cabinet work. (Exp. 1857.)

Captain Hansard gives 0.61 as the specific gravity of this wood.

Said by Grosourdy (2: 390) to break with a vertical fracture, and to be used for shelves, boards, framework of houses, etc.

Higuereta cimarrona. See *Jatropha gossypifolia*.

Higüerito. See *Crescentia microcarpa*.

Higüerito de Sierra. See *Schlegelia acillaris* and *Schlegelia brachyantha portoricensis*.



LIRA (HIBISCUS SCHIZOPETALUS).

Higuero.

A tree from all parts of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood light, soft; specific gravity 0.638; not used. (Exp. 1857.)

Higuillo. See *Plumbago scandens*.

A tree from all parts of the island; height, 15 to 20 feet (5 to 6 meters); diameter, 5 to 6 inches. Wood white, soft; specific gravity, 0.689; not used. (Exp. 1857.)

Higuillo amargo. See *Piper citrifolium*.**Hippocratea ovata.** BEJUCO PRIETO.

Family Hippocrateaceae; a woody climber. (Stahl, 4: 39.)

Hippocratea scandens.

A woody climber; reported from Sierra de Luquillo.

Hippomane mancinella. MANCHINEEL. MANZANILLO.

Reported by Hill as a tree of the coast region. A member of the Euphorbiaceae and related to the tallow tree, *Sapium sebiferum*. The wood makes handsome furniture and takes a high polish. The abundant milky juice is extremely poisonous to most persons, although some are not affected by it. In the eye it is said to cause temporary and even permanent blindness. Its deleterious properties have, however, been greatly exaggerated, to the effect that sleeping in its shade is considered fatal. The fruit has somewhat the appearance of a small apple, whence the Spanish name. The acrid juice attacks the mouth and prevents one eating a fatal quantity. Hill's account may be quoted as well illustrating the popular belief, page 38:

"The alleged effects of the guao tree upon man and animal have just been related, but more serious still are said to be the results of resting in the shade of the manzanillo or manchineel. This tree spreads throughout the seacoasts and river banks; is covered with regular leaves; its stem almost two feet in circumference; its bark smooth and delicate; the flowers rose-tinted. It is loaded with small, apple-like fruit of a pleasant appearance and odor; the leaves resemble those of the pear tree; the entire tree is full of a milky juice, which exudes in the heat of the sun. According to current belief, as frequently related by the writer, the incautious traveler, attracted by the beautiful appearance of the manzanillo, reposes in its shade, soon finds himself poisoned, and if lacteal substance falls upon him from the leaves, or if he touches the leaves he suffers as from the application of some blistering substance. It is also said that a fish which eats the fruit becomes infected, the gills becoming yellow and black, and one who eats the fish in this state is said to fall into a profound lethargy, with a general relaxation of all the limbs, according to the quantity of the fish he has eaten. This effect, according to popular belief, continues twenty-four hours, and frequently results in death. The use of brandy or other spirituous liquors or the drinking of sea water is considered beneficial as an antidote."

Furnishes a valuable wood for fine furniture, as easy to work as mahogany, whose properties and durability it possesses. It is of ordinary weight and hardness, fine-grained, and somewhat fibrous in texture. Its color is a handsome yellowish gray tinged with brown, with brown or black lines. This wood should not be worked until thoroughly dried and seasoned. Specific gravity, 0.674. (Grosourdy, 2: 401.)

Hirtella.

A genus, according to Stahl, of two Porto Rican species, *H. triandra* and *H. rugosa*, both bearing the common name "tete de burro." They are shrubs or small trees and are related to *Chrysobalanus*, which Stahl considers the type of a distinct family of Leguminosae, although the prevailing tendency is to place it in the Rosaceae.

Hog plum. See *Spondias lutea*.

Hog weed. See *Boerhavia*.

Hoja de yagua.

The leaf base of the royal palm.

Hoja menuda. See *Calyptrocalyx sintenisii*, *Eugenia biflora ludibunda*, *E. buxifolia*, *E. ligustrina*, *E. monticola*, *E. poiretii*, *E. procera*, *Myrcia coriacea*, *M. paniculata*, and *M. splendens*.

A tree from all parts of the island; height, 12 to 15 feet (3.5 to 4.5 meters); diameter, 8 to 10 inches. Wood, ash colored, hard; specific gravity, 0.822; used for stakes. (Exp. 1857.)

Grosourdy adds (2: 390) that the wood is flexible, fibrous, and moderately resistant. Used also to make charcoal.

Homalium racemosum. TOSTADO.

Family Flacourtiaceae; a tree 5 to 8 meters high, found on mountains and in waste places. (Stahl, 4: 161.)

Hordeum vulgare. CEBADA.

Barley is not raised in the island, but is imported in small quantities, mostly for making soups and for barley water, which is used as a drink.

Horse bean. See *Canavalia gladiata*.

Horseradish tree. See *Moringa pterygosperma*.

Hortegon.

Hansard gives this wood a specific gravity of 1.25. It varies in color from deep red to black and is very durable and hard. The leaf is large. When making boundaries a piece 2 or 3 feet long is put in the ground and lasts indefinitely.

Hortegon labrado.

A wild tree not commonly met with, and then only in the northeast part of the island. It reaches a height of 60 feet, and the diameter of the long straight trunk is not over 2 feet. It furnishes a very coarse, flesh-colored wood of the greatest durability. It breaks with difficulty with a vertical fracture. Its common use is for shelving in houses. (Grosourdy, 2: 390.)

Specific gravity, 1.207. (Exp. 1857.)

Hortegon prieto.

A tree from all parts of the island; height, 75 to 80 feet; diameter, 20 to 25 inches. Wood dark, very hard; specific gravity, 1.064; used for cabinet work. (Exp. 1857.)

Grosourdy says this is but a variety of "hortegon labrado." The only difference being the brown-colored wood. It is used in the same way. (Grosourdy, 2: 390.)

Hortones.

A variety of banana or plantain mentioned by Hill, but not described.

Hoya carnos.

An ornamental plant of the milkweed family, introduced from the Malay region and now naturalized in several of the West Indies.

Hucar.

"Bois immortelle," good for coffee and cacao shades; specific gravity, 1.06 (Hansard). See also "bucar" and "ucar."

Probably refers to *Erythrina poeppigiana*.

Hucar amarillo.

A tree from all parts of the island; height, 80 to 85 feet (24 to 25 meters); diameter, 50 to 55 inches (125 to 137 centimeters). Wood yellowish, very hard; specific gravity, 1.080; used in boat building. (Exp. 1857.)

Captain Hansard calls this "bois immortelle" and says that it is good for coffee and cacao shades; specific gravity given at 1.07.

Hucar blanco. See *Bucida buceras*.

A tree from all parts of the island; height, 80 to 85 feet (24 to 25 meters); diameter, 45 to 80 inches (112 to 200 centimeters). Wood white, very hard; specific gravity, 1.093; used in boat building. (Exp. 1857.)

Hucar colorado.

Bois immortelle, good for coffee and cacao shades. Specific gravity, 0.93. (Captain Hansard.)

Hucar prieto.

A tree from all parts of the island; height, 70 to 75 feet (21 to 23 meters); diameter, 40 to 45 inches (100 to 112 centimeters). Wood dark, very hard; specific gravity, 1.073; used in boat building. (Exp. 1857.)

Grosourdy (2: 390) says this is a variety of "hucar blanco" (*Bucida buceras*) differing only in having the wood dark gray in color.

Hucarillo.

A tree from the eastern part of the island; height, 40 to 45 feet (12 to 14 meters); diameter, 20 to 24 inches (50 to 60 centimeters). Wood white, hard; specific gravity, 0.943; used in building houses. (Exp. 1857.)

Captain Hansard gives a specific gravity of 0.80 for wood of this tree.

Huesillo. See *Maypea domingensis*.

Hueso blanco. See *Maypea domingensis*.

Hueso prieto. See *Ilex nitida*.

Huevo de gato. See *Helicteres jamaicensis*.

Hura crepitans. HAVILLA.

This or a related species is not unfrequently planted as a shade tree along roads, particularly between San German and Mayaguez. It reaches a rather large size (2½ feet—75 centimeters—in diameter) and has a spreading habit, but the wood is soft and brittle so that it suffered much damage from the hurricane. The trunk suggests that of the *Ceiba*, having prominent roots, large stout spines, and a similar light gray color. The habit of the branches is different, however, and the leaves are simple and cordate instead of palmately compound. They are also peculiar in having unusually numerous white veins. *Hura* belongs to the Euphorbiaceae, but differs from most of the family in having a seed capsule of numerous divisions, somewhat resembling, exteriorly, the fruit of *Malva* on a large scale. The flowers are also peculiar, the staminate being on a spike suggesting the Piperaceae or aroids; the pistillate flower is naked, with a large spreading trumpet-shaped, lacerate stigma. The seeds are loosened and rattle in their separate compartments, perhaps suggesting the name "sand-box tree," used in the English colonies. The name "monkey's dinner-bell" is also applied in allusion to the fact that the ripe fruit splits with a loud report.

The slightly milky juice is acrid and irritant, while the seeds are sometimes used as a purgative. An oil expressed from them is said to be less nauseous than castor oil and to be effective in smaller doses. The fruits of the Porto Rican trees were much less prominently ribbed than those figured in botanical works; possibly it represents a distinct species.

Grosourdy says of the wood that it is rather light and soft, fibrous in texture, and resistant to the action of water. It is nearly white with stripes of light brown of the same color as the heart. There is a variety, light brown in color, that is highly valued because canoes made of it are much stronger and last much longer than those made of the other varieties. Specific gravity, 0.465. It is used also

for interior work in houses. In working this wood care should be taken that the dust does not enter the eyes or nose, as it is very irritating and produces considerable inflammation. It is sufficient that the face be covered with a gauze. The native name is here spelled "javillo." (Grosourdy, 2: 392.)

Huso.

A tree from all parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood greenish, very hard; specific gravity 1.114; used for cabinetwork. Also called "espinillo." (Exp. 1857.)

Color, light yellow, with irregular, thin, yellowish-brown streaks. Conspicuous ducts in short, detached chains (1 to 2 rows of cells) evenly diffused. Medullary rays minute, but distinct. Wood fibers strongly interlaced (crossgrained). Tough, uncleavable wood. Used for hubs. (Hill and Sudworth.)

Huso amarillo.

Specific gravity 0.88, according to Captain Hansard.

Huso blanco.

Captain Hansard says that this wood has a specific gravity of 0.89.

Huso colorado.

Specific gravity, according to Captain Hansard, is 1.30.

Hydrocotyle.

This is a genus of small umbelliferous herbs, growing in moist, shady places, and usually creeping by means of long rhizomes.

Hydrocotyle asiatica. See *Centella asiatica*.

Hydrocotyle hirsuta. YERBA DE CUARTO.

Hydrocotyle spicata. See *Hydrocotyle hirsuta*.

Hydrocotyle umbellata. YERBA DE CUARTO.

A creeping herb found in rivulets and swamps. Used by Brazilian natives to produce vomiting; also said to be an antidote for poison. Kills sheep if they feed upon it.

Hygrophila portoricensis. HIGROFILA DE PUERTO RICO.

Family Acanthaceae; an herbaceous annual somewhat fleshy, found in ponds and on sides of ravines. Peculiar to Porto Rico. (Stahl, 4: 243.)

Hymenaea courbaril. ALGARROBO.

PLATE XLI.

Also called "courbaril" and in Jamaica "locust tree." It belongs to the family Cassiaceae and is a large tree 15 or more meters high. The fruit is a short, thick, and very hard pod, the seeds of which are embedded in a dry, sweet pulp, eaten by children. The pods frequently appear in the markets. This tree is said to produce the "gum animae" of commerce, which is thought to be a finer varnish than shellac.

Specific gravity of the wood, 1.06. (Hansard.) Jacquin says that the wood is very hard and tough and is used for cogged wheels in sugar mills. Lunan attributes a variety of medicinal virtues to the different parts of this tree. (Stahl, 3: 129.)

Grosourdy describes two varieties, which he calls "algarrobo amarillo" and "algarrobo colorado."

Hypelate paniculata. GAITA.

Family Sapindaceae; a low or middle-sized tree. (Stahl, 2: 161.)

Hyptis. See *Mesosphaerum*.

Iaya jabico. See *Erithalis fruticosa*.

Ibatia maritima. POPON.

A densely pubescent, climbing, asclepiadaceous plant preferring forests and thickets near the coast.



PODS OF ALGARROBA : HYMENAEA COURBARIL . PONCE MARKET.

Icaco. See *Chrysobalanus icaco*.

Icaque. See *Chrysobalanus icaco*.

Ileucedran.

Enumerated by Hill among forest trees yielding timber and fuel, but the name is admitted to be doubtful and seems not to have been printed elsewhere.

Ilex.

A large genus of Aquifoliaceae, including many West Indian shrubs and trees; several of the latter are called by the natives "cuero de sapo," but *I. macoucoua* and *I. ridlaci* have no local names reported.

Ilex dioica. CUERO DE SAPO.

A shrub, 3 to 5 meters high, found in the mountains and waste places; also called "brigueta." (Stahl, 4: 36.)

Ilex macoucoua. BRIGUETA NARANJO.

A shrub, 3 to 5 meters high, found in waste places and in the mountains. (Stahl, 4: 35.)

Ilex nitida.

According to Urban (Add. 1: 35) variously known in Porto Rico as "cuero de sapo," "palo de hueso," and "hueso prieto." Reported from Bayamon, Sierra de Luquillo, Hato Grande, Cayey, Aibonito, Adjuntas, and Lares, in "coffee forests." Also known from Jamaica, Montserrat, and Martinique.

Ilex sideroxyloides. GONGOLIN.

A large tree reported from the Luquillo Mountains. There are two varieties, *portoricensis* and *occidentalis*. (Urban, Add. 1: 32.)

Ilex urbaniana. CUERO DE SAPO.

A tree reported from primeval forest at Maneyes, near Utuado. (Urban, Add. 1: 33.)

Ilysanthes gratioloides. See *Ilysanthes riparia*.

Ilysanthes riparia. YERBA GRACIOSA.

Family Scrophulariaceae; a procumbent herb growing in wet places. (Stahl, 6: 236.)

India rubber vine.

See note under "rubber vine."

Indian almond. See *Terminalia catappa*.

Indian corn. See *Zea mays*.

Indian creeper. See *Quamoclit quamoclit*.

Indian pink. See *Quamoclit quamoclit*.

Indian shot. See *Canna*.

Indian sorrel. See *Hibiscus sabdariffa*.

Indigo. See *Indigofera anil*.

Indigo berry. See *Randia aculeata*.

Indigofera anil and I. tinctoria. INDIGO. AÑIL.

An annual or biennial leguminous shrub very common as a weed in waste places at low elevations in Porto Rico. Many species of plants are now known to produce indigo, but the species of *Indigofera* are preferred. In India *I. tinctoria* is generally cultivated, while in the American tropics *I. anil* is believed to have been much more commonly used. Careful comparative experiments regarding the value of these and other indigo-yielding plants seem not to have been made. Many of these could undoubtedly be grown successfully in Porto Rico, but unless improvements in the methods of manufacturing can be made it is not likely that there could be successful competition with the East Indies, whence

the principal supply now comes. Prices vary at present from 60 cents to \$1.20 per pound, depending upon quality. The competition of an artificial coal-tar product called alizarin, which threatened at one time to entirely replace indigo, is not feared so much as formerly, it having been found that indigo dyes are preferable for many purposes.

This crop was cultivated on a considerable scale in our Southern States during the eighteenth century, about 1,500,000 pounds being exported in 1794. The conditions were, however, not sufficiently favorable to enable the planters to withstand the competition of tropical countries where three crops can be cut each year, and production gradually ceased. At present we import indigo to the value of about \$1,600,000 annually. Cultivation and manufacture are simple processes and have been well summed up in the following paragraph, extracted from a circular letter compiled by Mr. Lyster H. Dewey, of the Division of Botany. Although these operations are not complicated, much skill and experience are necessary in order to decide how long to continue the various stages of fermentation and aeration:

"Indigo is cultivated on sandy soil. The seeds are sown in spring either in drills or hills or broadcast. When sown in hills or drills it is cultivated or hoed. It is ready to cut from 45 to 60 days after seeding. The indigo is extracted by placing the plants, either fresh or after they have dried like hay, in a large tank, where they are covered with water. They are allowed to remain in the water about twenty-four hours, or a shorter time if the water is heated. It should not be boiled. The water is then drawn off into another tank, where it is stirred for at least three or four hours. Sometimes limewater is added during the stirring process to aid in precipitating the indigo. This is either allowed to settle in the tank where it is stirred or, more frequently, drained off into a third tank. The settling requires about twenty-four hours. When the indigo has settled to the bottom, leaving the water above practically clear, the water is carefully drawn off and the indigo is taken out and dried on cloths. When it is dried to about the consistency of molding clay it is usually made up into balls, and these are dried in the shade. When thoroughly dried and hard they are packed for market. The yield of indigo varies from 50 to 140 pounds per acre for each cutting. The average yield in Venezuela is about 112 pounds, but in the United States it was considerably less."

When planted in rows, these are about 2 feet (60 centimeters) apart, and 15 pounds (7 kilograms) of seed are used per acre. Blossoming is sometimes delayed for twelve weeks, and in some countries exposure of the cut indigo to the sun is considered detrimental. The average yield has been given as 300 pounds (142 kilograms) per acre. Although belonging to the Leguminosae, the indigo plant is said to rapidly exhaust the soil, so that planting on the same land can not be indefinitely repeated. Alternation with the pigeon-pea (*Cajanus*) has been advised.

In different countries there are also considerable differences in the methods of extraction, and in some parts of India the precipitate is boiled before being finally pressed and dried. This method is now defended by the best authorities as an improvement over that formerly followed in the United States and still generally prevalent in the American tropics.

The important difference in the market prices quoted above shows that quality is important in this product as in every other, and while indigo can not, perhaps, be recommended to the general farmer in Porto Rico, it is by no means certain that a properly organized plantation with a well-equipped and intelligently directed factory would not be a profitable enterprise. The production of indigo of high quality on a small scale is seldom accomplished, because of the absence of the most favorable conditions and best appliances; but in a large enterprise these can be profitably provided.

Another objection to indigo has been the injurious effect of the manufacturing processes upon the health of the operatives; but it would probably not be difficult at this time to provide machinery for stirring the tanks of fermenting indigo and also for handling and sacking the finished product.

Inga laurina. GUAMÁ.

Family Mimosaceae; a tree probably introduced, 30 to 50 feet (9 to 15 meters) high. The leaves usually with four leaflets. The flower clusters are longer and the stamens much shorter than in other members of this genus.

As a shade tree for coffee this species is second in importance only to the "guava" (*I. vera*). Specimens growing at Isolina were found in some cases at least to have tubercles on the roots.

According to Guerin, this tree is one of those employed in the cacao culture of Guadeloupe in making hedges or windbreaks which are planted across the direction of the prevailing winds at distances of 100 meters. Such hedges are used at elevations where permanent shade is thought not to be required, but where protection from the wind is necessary. In Venezuela and Colombia this is one of the favorite species for coffee shade. As a possible objection to it may be noted the report that it is susceptible to the disease called "mancha de hierro," or "iron rust," which has been very destructive to coffee, especially in overshaded plantations.

In Colombia, Saenz recommends the planting of this species at 10 meters where the temperature does not exceed 21° C.; in hotter regions a distance of 10 meters is advised. (Bull. 25, Division of Botany, U. S. Dept. Agr., p. 62.)

Reported from Utuado.

Inga vera. GUAVA.

A tree 30 to 50 feet (9 to 15 meters) high, producing round clusters of white flowers with very long stamens. As a shade tree for coffee this seems to be the favorite species in Porto Rico, with the related *Inga laurina*, or guamá, second. Both grow much less rapidly than the bucare (*Erythrina*), but are less liable to injury by hurricanes. In well-kept plantations the lower branches are trimmed out and the upper form an even layer of rather open foliage. But even where this amount of care is taken the shade is usually still too thick for the coffee to make normally vigorous growth, and the crop would probably be improved by the removal of alternate trees, so as to increase the distances to 30 or 40 feet.

The name of this tree is likely to be confused by the visitor with that of *Psidium guajava*, the well-known tropical fruit called guava in English, but by the Porto Ricans more correctly termed guayava. (Bull. 25, Division of Botany, U. S. Dept. Agr., p. 63.)

Tubercles were also observed on the roots of this species at Isolina.

Grosourdy (2: 386) describes the wood as strong and resistant, used only for the making of charcoal and for fuel. He gives the specific gravity as 0.505. This author is certainly in error in giving this species the native name of "guamá."

Ink berry. See *Randia aculeata*.

Inodes causiarum. PORTO RICAN HAT PALM. YARAY.

Family Sabalaceae. The most conspicuous difference between the present genus and Sabal is the fact that the former produces an upright trunk while the latter has only what might be called an underground rootstock. Although such a distinction is quite artificial, both groups of species begin life with a creeping axis which becomes erect in one and remains horizontal in the other. A much more important difference is to be found in the leaves which in *Inodes* have secured strength by the development of a midrib, a tendency early abandoned by Sabal, in which the midrib is rudimentary and the middle of the leaf is the weakest part. The leaves of Sabal are adapted for standing erect, and avoid

resistance to the wind by being split down the middle. The leaves of *Inodes*, which are held horizontal from an erect axis, have attained the unique adaptation of a decurved midrib which braces the sloping sides of the leaf and effectively prevents the breaking above the ligule common in some of the species of *Thrinax*. It is true that the palm-leaf hats manufactured in large quantities in Porto Rico are made from the present species. The center of the hat industry is at Joyua, a small village on the western coast of the island some miles southwest of Mayaguez and west of Cabo Rojo. Here many hundreds of the palms are growing along the shore in a narrow belt of coral sand.

Ipecac.

See note under *Asclepias curassavica*.

***Ipomoea acetosifolia*.** See *Ipomoea carnosae*.

***Ipomoea batatas*.** SWEET POTATO. BATATA.

In Porto Rico the sweet potato is found in all sorts of places, particularly at low elevations, often appearing quite wild, though in a country where such general and yet desultory agriculture has been carried on it would be very difficult to demonstrate the indigenous character of the plant. It seems, however, that the potato was cultivated in the West Indies before the advent of Europeans, but that does not necessarily involve an indigenous origin, since it was probably already spread throughout tropical America. Moreover, like the cocoanut, it seems to have crossed the Pacific and reached China in the second or third century. Some writers have believed in the Asiatic or Malayan origin of the plant, and have claimed that batata is a Malay name, though this is not the prevailing idea, the present tendency being to consider not only batata but also "camote," the Philippine name as of American origin, and probably introduced from Mexico by the Spaniards.

Little attention is paid in Porto Rico to the planting of pure or high-grade stock, and propagation is carried on exclusively from the vines, pieces a foot long being buried for half their length in the ground. When regular culture is attempted the cuttings are placed a foot apart in rows 2 feet apart. The weeds are sometimes kept down and the rows hilled up, but often neither of these matters is attended to. The crop is not dug at any regular time, but individual roots are discovered by probing and taken out without otherwise disturbing the vine, which is permitted to grow indefinitely.

Numerous varieties are grown in every patch, to judge from the very different leaf forms. The red sorts have lobed leaves, while those of the white kinds (*batata blanca*) are simply cordate or nearly circular. For good crops rich land is necessary, and the red clay soils seem to be very suitable. Four tons per acre is quoted as a good crop, but a second crop may be secured a month or six weeks later. In the eastern end of the island and in Vieques there is said, for some unexplained reason, to be a scarcity of sweet potatoes, so that with the regular demand a local market is open to those who might undertake this culture on a commercial scale. At the time of our visit retail prices were 3 or 4 centavos per pound in the city markets.

***Ipomoea biloba*.** BEJUCO DE PUERCO DE PLAYA.

(Stahl, 6: 160.)

***Ipomoea bona-nox*.** BEJUCO DE VACA.

An annual ornamental vine growing to great length and bearing many large flowers. Bello gives the name "bejuco de puerco" for *Calonyction megalocarpum*. (Stahl, 6: 148.)

***Ipomoea carnea*.** BATATILLA CARNOSA.

A large vine found along the coast. (Stahl, 6: 277.)

Ipomoea carnos. BEJUCO DE PUERCO DE COSTA.

A climbing vine growing in the sands along the coast. According to Bello this species is called "batatillo" in the western part of the island. (Stahl, 6: 161.)

Ipomoea cathartica. BEJUCO DE GLORIA.

A vine found in waste places along the coast. (Stahl, 6: 166.)

Ipomoea ciliolata. BEJUCO DE PUERCO BLANCO.

A climber occurring in all parts of the island. (Stahl, 6: 153.)

Ipomoea coccinea. See *Quamoclit coccinea*.

Ipomoea dissecta. NOYO.

A climber, cultivated in gardens; its leaves have an odor of prussic acid. (Stahl, 6: 150.)

Ipomoea fastigiata. BEJUCO DE PUERCO.

A twining vine found in waste places. The tuberous roots are called wild potatoes in Jamaica. (Stahl, 6: 156.)

Ipomoea filiformis. CAMBUTERA DE COSTA.

A much-branched vine found on the coast plains. (Stahl, 6: 163.)

Ipomoea pentaphylla. BATATILLA BLANCA.

A twining climber found in waste places. (Stahl, 6: 151.)

Ipomoea pes-caprae. See *Ipomoea biloba*.

Ipomoea quamoclit. See *Quamoclit quamoclit*.

Ipomoea quinquefolia. BATATILLA BLANCA.

A climber found in the southern part of the island. (Stahl, 6: 153.)

Ipomoea sericantha. BATATILLA DE ZAETA.

A creeping vine, in sands between Vega Baja and Manati. (Stahl, 6: 162.)

Ipomoea setifera. BEJUCO DE PUERCO.

A woody vine found in all parts of the island. (Stahl, 6: 152.)

Ipomoea triloba. BEJUQUILLO DE PUERCO.

A vine 3 to 4 feet (about 1 meter) long, found in hedges, flowering in winter. (Stahl, 6: 157.)

Ipomoea umbellata.* AGUINALDO AMARILLO.

A twining vine found in different parts of the island. (Stahl, 6: 159.)

Ipomoea ventricosa. BATATILLA VENTRUDA.

(Stahl, 6: 277.)

Iresine argentata.

Family Amarantaceae; reported from Juya.

Isotoma longiflora. TIBEY BLANCO.

Family Campanulaceae; an herbaceous annual growing in dry, rocky localities. Near Toa Alta this species was called simply "tibey." According to Dr. Stahl, it is reputed to be poisonous, but no direct evidence is given. (Stahl, 6: 27.)

Ixora ferrea. PALO DE HIERRO.

Family Rubiaceae; a tree or shrub 3 meters high, found on mountain slopes and waste places. (Stahl, 5: 63.)

Jaboncilla.

Captain Hansard calls this "soapwort;" specific gravity 0.63. Probably the same as "jaboncillo."

Jaboncillo. See *Sapindus saponaria* and *S. marginalis*.

Jacana. See *Lucuma multiflora*.

A wild tree, 40 feet (12 meters) high, with a trunk 2 feet (60 centimeters) in diameter. Furnishes a rather strong, light-colored, flexible wood, used only for fuel. (Grosourdy, 2: 391.)

Specific gravity 0.920. (Exp. 1857.) See also note under *Acana*.

Jacanillo. See *Ardisia pendula*.

Jack plum. See *Syzygium jambolanum*

Jack tree. See *Artocarpus integrifolia*.

Jacquemontia tamnifolia. AGUINALDO PELUDO.

Family Convolvulaceae; a twining herb found in dry places. (Stahl, 6: 171.)

Jacquinia armillaris. BARBASCO.

Family Myrsinaceae; a shrub 3 to 4 meters high; grows along the coast. (Stahl, 4: 40.)

Jacquinia aristata. BARBASCO.

(Stahl, 4: 41.)

Jacquinia berterii.

An indigenous tree of 6 to 8 meters; two subspecies, *portoricensis* and *retusa*, are described by Urban, both known only from the southwestern part of Porto Rico. (Urban, Symb. 1: 378.)

Jacquinia umbellata.

An indigenous shrub, maintained by Urban as distinct from *J. aristata*, to which it was referred by Grisebach, known from Coama, Yauco, Maricao, and from between Sabana Grande and Guanica.

Jagua. See *Genipa americana*.

A tree from all parts of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood white, hard; specific gravity 0.886; used in making packing boxes. (Exp. 1857.)

Captain Hansard gives the specific gravity as 0.80 and says that it is a large tree, yielding good white boards; also a large oval fruit.

Jaguey.

A tree from all parts of the island; height, 40 to 45 feet (14 to 15 meters); diameter, 55 to 60 inches (130 to 150 centimeters). Wood ash-colored, soft; specific gravity, 0.485; used in making fishing canoes. (Exp. 1857.)

In Cuba this name is applied to *Ficus membranacea*, a species not known to occur in Porto Rico.

What is evidently another tree is described by Grosourdy under this name, and also called "mato palo," which see.

Jamaica cherry. See *Ficus pedunculata*.

Jamaica bitterwood. See *Picrasma excelsa*.

Jamaica mignonette tree. See *Lawsonia inermis*.

Jamaica nutmeg. See *Monodora myristica*, the calabash nutmeg.

Jamaica sorrel. See *Hibiscus sabdariffa*.

Jambolin. See *Syzygium jambolanum*.


Jambosa caryophyllus. See *Caryophyllus aromaticus*.

Jambosa jambos. ROSE APPLE. POMA ROSA.

A myrtaceous shrub or small tree native in the Malay region but now introduced in all parts of the Tropics and escaped from cultivation in many countries. In Porto Rico it is abundant in thickets, in waste places, and furnishes the material from which most of the larger baskets are made. Considerable amounts are also used for hoops of the large sugar casks. It yields also an edible fruit which has a pleasant odor like roses, whence the name. The long and narrow leaves resemble those of the peach and oleander. (Stahl, 4: 87, as *Eugenia jambos*).

Grosourdy (2: 409) says that this tree reaches 40 feet in height and that the trunk attains a diameter of 15 to 18 inches. He also states that the wood is used to make charcoal for forges.

Jambosa vulgaris. See *Jambosa jambos*.



PHYSIC NUT (JATROPHA CURCAS). CATAÑO.

Japanese medlar. See *Eriobotrya japonica*.

Jasmin. See *Jasminum pubescens* and *J. officinale*.

Jasmin del rio. See *Cleome spinosa*.

Jasmin hediondo. See *Clerodendron fragrans*.

Jasminum officinale. JASMIN.

Family Oleaceae; a climbing shrub, cultivated in gardens. Introduced from Asia. (Stahl, 6: 63.)

Jasminum pubescens. JASMIN.

A biennial shrub, introduced from India, and cultivated in gardens. Reported from Sierra de Naguabo and near Rio Blanco, about dwellings; apparently spontaneous. Noted by us at Caguaitas and at Juana Diaz. (Stahl, 6: 62.)

Jatropha curcas. TARTAGO.

PLATE XLII.

Family Euphorbiaceae; reported from near Peñuelas. This is a shrub about 8 feet high, and is called "physic nut" in Jamaica, where the seeds are used as a purgative. The juice of the leaves is said to act as a resolvent. The black beans are similar to those of the castor-oil plant; they are ground up, boiled in water, and used as a remedy for diarrhea.

Jatropha gossypifolia. HIGUERETA CIMARRONA.

Occasional as a weed in waste places, particularly in sand banks along water courses. The common name was noted at Coamo Springs, and, doubtless, alludes to the similarity to the castor bean, *Ricinus communis*.

Java plum. See *Syzygium jambolanum*.

Javillo. See *Hura crepitans*.

Jaya.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood, yellow, hard; specific gravity, 0.778; not used. (Exp. 1857.)

Jerusalem thorn. See *Parkinsonia aculeata*.

Jiba. See *Erythroxylon brevipes* and other species of the same genus.

Jicama. See *Calopogonium caeruleum*.

Jimson weed. See *Datura stramonium*.

Jobo. See *Spondias lutea*.

Job's tears. See *Coix lachryma-jobi*.

Juan caliente. See *Rourea glabra* and *R. frutescens*.

Jucare.

Given by Hill as a variation of the name "ucar."

Juglans cinerea. NUEZ.

A tree growing wild in the interior of the Greater Antilles and on the Continent. It reaches a height of 40 to 45 feet (14 to 15 meters), with a straight, rather long trunk 18 to 24 inches (45 to 60 centimeters) in diameter. Furnishes a good, though rather porous wood, with the properties of the European walnut; is dark red in color, and breaks with a vertical fracture. It is employed for boards and timbers in housebuilding. (Grosourdy, 2: 403.) See also note under *Nues*.

This species, the common butternut of the United States, is not known to occur in Porto Rico, and this author's identification is probably in error.

Juglans insularis.

According to the Kew Bulletin for 1894, specimens collected by Sintenis near Adjuntas and distributed from Berlin as *Juglans jamaicensis* belongs to *J. insularis*, a species previously known from Cuba.

Juso.

A hard wood with a specific gravity of 1.12; small trees, making good posts. (Hansard.) Probably the same as "huso" or "hueso."

Jussiaea acuminata. See *Jussiaea linifolia*.

Jussiaea hirta. See *Jussiaea peruviana*.

Jussiaea linifolium. YERBA DE CLAVO AGUADA.

Family Onagraceae; a woody, erect plant, growing in hedges and along edges of ponds. (Stahl, 4: 132, as *Jussiaea acuminata*.)

Jussiaea palustris. YERBA DE CLAVO PALUSTRE.

An erect annual, 60 centimeters high, found in inundated plains. (Stahl, 4: 133).

Jussiaea peruviana. YERBA DE CLAVO PELUDA.

An herbaceous, woody annual, found in inundated places: 1 meter high. (Stahl, 4: 134, as *Jussiaea hirta*.)

Jussiaea pilosa.

An erect, woody herb in wet places: the variety *glabra* is known from near Manati.

Jussiaea repens. YERBA DE CLAVO ACUATICO.

A fleshy aquatic herb, sometimes floating, found in marshes. A great variety of medicinal properties are attributed to this by the natives of Jamaica, where it is called "primrose willow." (Stahl, 4: 129.)

Jussiaea suffruticosa. YERBA DE CLAVO.

An herbaceous, woody annual or biennial, common in wet places: 1 meter high; reported from near Manati. (Stahl, 4: 131.)

Kallstroemia maxima.

Family Zygophyllaceae; a leafy herbaceous plant. Secured by Sintenis in cultivated grounds near Manati.

Lablab vulgaris. See *Dolichos lablab*.

Labourdonnaisia albescens. ACANA.

A sapotaceous tree with hard wood.

Lache prieta.

A tree from the interior of the island; height, 25 to 30 feet (8 to 9 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood pink, hard; specific gravity, 0.993; used in building houses. (Exp. 1857.)

Lactuca floridana.

A coarse herbaceous composite collected by Sintenis on a coffee farm near Cayey.

Lactuca intybacea. CHICORIA AZUL.

An erect much-branched annual composite. (Stahl, 5: 155.)

Lactuca sativa. LETTUCE.

The lettuce offered in the markets in Porto Rico seemed to be of fairly good quality; better than could be raised in most tropical countries. Even at sea level, near Santurce, lettuce seemed to thrive moderately well, but it could probably be made to do better in the mountains. The seed is, of course, imported.

Lagenaria vulgaris. GOURD. MARIMBO.

Also called "amargo," "marimbo," and "cadungo amargo." Family Cucurbitaceae. This vine yields the gourds from which the so-called "guira," "guichara" or "caracho" is made. In Cuba the vine is said to be called "guiro cimarron," while "guira cimarrona" is the calabash tree *Crescentia*.

The young gourds, 6 to 10 inches long, are eaten. Two of the latter size were offered for 10 cents in the San Juan market, June 14.



MANGLE BLANCO (*LAGUNCULARIA RACEMOSA*); FLOWERS, SEEDS, AND SEEDLINGS.
CATAÑO.

Lagerstroemia indica. ASTROMEDA. CRAPE MYRTLE.

Family Lythraceae; an ornamental shrub, native in China and extensively cultivated in greenhouses in Europe and America, or, southward, in the open air. In the West Indies it is frequent in gardens: reported from Sierra de Luquillo and from Peñuelas, about dwellings.

Lagerstroemia reginae.

An East Indian tree valued for its timber, which is extremely durable in water. Introduced into Jamaica according to Grisebach.

Laguncularia racemosa. MANGLE BLANCO.

PLATE XLIII.

Family Combretaceae; a shrub or tree 5 to 8 meters high; grows in tide-water swamps and is often mistaken for the true mangrove, *Rhizophora*. It is abundant immediately behind Cataño. (Stahl, 4: 136.)

Lancewood.

A name applied in the British West Indies to species of *Oxandra* and *Cananga*.

Lantana camara. CARIAQUILLO.

Family Verbenaceae; a spiny shrub, 1 meter high, found in rocky places; reported from Guanica, Yanco, and Mayaguez. (Stahl, 6: 214.)

Lantana crocea. CARIAQUILLO.

A shrub slightly longer than *L. camara*. (Stahl, 6: 215.)

Lantana involucrata. CARIAQUILLO DE SANTA MARIA.

A shrub, 1 meter high, found in waste places; flowers fragrant. At Cataño this is called simply "Santa Maria." (Stahl, 6: 215.)

Lasianthus lanceolatus. AROMA.

Also called "bejuco de peo" and "mata de peo." A rubiaceous shrub from the eastern and southern parts of the island. (Urban, Symb. 1: 449.)

Lasianthus moralesii.

Family Rubiaceae; reported from Naguabo.

Laugeria resinosa. AQUILON.

Family Rubiaceae; a shrub, 8 feet (2.5 meters) high, on mountains.

Laurel amarillo.

Perhaps the same as "laurel sabino." The wood is yellow, makes good boards, stands water well, and is pretty for furniture; specific gravity, 0.96. (Hansard.)

Laurel blanco.

A tree from all parts of the island; height, 40 to 45 feet (12 to 14 meters); diameter, 20 to 25 inches (50 to 62 centimeters); wood, white, hard; specific gravity, 0.606; used in building houses. (Exp. 1857.)

Captain Hansard gives a specific gravity of 0.84 and says it is a white wood used for furniture.

Grosourdy (2: 393) includes both "laurel prieto" and "laurel blanco" under the following description:

A wild tree, 60 to 80 feet (18 to 24 meters) high, with a long, straight trunk 6 to 8 inches (15 to 20 centimeters) in diameter. It furnishes a wood very strong and resistant, rather light, fibrous in texture, neither hard nor flexible, and moderately fine-grained. The wood exhales an agreeable aromatic odor. The color is a grayish yellow or very light cinnamon. Specific gravity, 0.658. Used on the continent to make umbrellas and in building boats. It is also suitable for doors, windows, etc.

Laurel prieto.

A tree from all parts of the island; height, 40 to 45 feet (12 to 14 meters); diameter, 20 to 25 inches (50 to 62 centimeters); wood, dark, hard; specific gravity, 0.572; used in building houses. (Exp. 1857.) (See also *Laurel blanco*.)

Laurel puero.

A tree from all parts of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 20 to 25 inches (50 to 62 centimeters); wood, yellowish, hard; specific gravity, 0.628; used in building houses. Also called "azafran." (Exp. 1857.)

Perhaps this should be "laurel puerco."

Laurel sabino. See *Magnolia splendens*; also *Talauma*.

Color, clear olive-brown. Ducts occurring singly, evenly diffused in one or two irregular lines, between the indistinct medullary rays. A straight-grained wood similar in color to, but finer-grained than the heart of the tulip and cucumber tree of the United States. (Hill and Sudworth, p. 28.)

Laurel savino.

A tree from all parts of the island; height, 40 to 45 feet (12 to 14 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood, greenish, hard; specific gravity, 0.750; used in building houses. Probably another name for "laurel sabino." (Exp. 1857.)

Lawsonia alba. See *Lawsonia inermis*.

(Stahl, 4: 122.)

Lawsonia inermis. HENNA. RESEDA.

Family Lythraceae: a glabrous shrub, 2 meters high, with white flowers; naturalized from Egypt and Syria in Jamaica and Antigua. Yields a yellow dye, much used by the Egyptians and Arabs as a cosmetic for the palms of the hands, finger nails, and hair, also for the tails of horses, and for coloring cloth a dull red. "Egyptian privet" and "Jamaica mignonette tree" are additional names. Cultivated in gardens in Porto Rico. Specimens have been collected near Guanica. It will grow without irrigation in extremely dry situations and has been suggested as a hedge plant. (Stahl, 4: 122, as *L. alba*.)

Lechecillo. See *Chrysophyllum glabrum*.

A wild tree, 40 feet (12 meters) high, with edible fruit. The trunk 12 inches (30 centimeters) in diameter. The wood is rose-colored, hard, and breaks with a vertical fracture. Its common use is for shelving in houses. It is used also for making a fine charcoal for forges. (Grosourdy, 2: 394.) Specific gravity, 0.883. (Exp. 1857.)

Leche prieta.

A wild tree from the interior of the island, 20 feet (6 meters) high, with a trunk 27 inches (67 centimeters) in diameter. It furnishes a compact, rose-colored wood that is very difficult to break. Its common use is for shelving in houses. (Grosourdy, 2: 394.)

Lechicillo.

Probably another spelling of "lechecillo."

A small tree, abundant on mountain slopes near streams. The fruit is eaten by the natives, while the wood is very soft and not used. The specific gravity, according to Captain Hansard, is 0.79.

Lechoncillo.

Captain Hansard gives this wood a specific gravity of 0.80. It is probably the same as lechicillo.

Lechosa. See *Carica papaya*.**Lemon.** See *Citrus limonum*.**Lengua de vaca.** See *Elephantopus scaber* and *Sausserieria* sp.**Leonotis nepetaefolia.** MOLINILLO.

PLATE XLIV.

A herbaceous annual of the mint family, 15 to 30 inches (37 to 75 centimeters) high; the large bur-like flower clusters are very peculiar, and when covered with



MOLINILLO · LEONOTIS NEPETAEFOLIA.

the yellow flowers the plant is quite ornamental. Bello gives the names " boton de cadete " and " quina del pasto. " (Stahl, 6: 198.)

At Arecibo this plant was said to be poisonous.

Leonurus sibiricus. AGRIPALMA.

A herbaceous annual weed of the mint family. (Stahl, 6: 197.)

Lepidium virginicum. MASTUERZO.

This common North American cruciferous weed is not uncommon in some parts of Porto Rico. (Stahl, 2: 37.)

Leria nutans. See *Chaptalia nutans*.

Leucaena glauca. ACACIA PALIDA. HEDIONDILLA.

Family Mimosaceae; a shrub 3 meters high. Valued for its wood, edible pods, and seeds. (Stahl, 3: 146.)

A small tree, with characteristic clusters of long, thin pods, growing in open waste places, particularly in dry localities. It is said to withstand drought when all other vegetation withers, and is then sometimes eaten by domestic animals with bad results, especially in the case of horses. The skin and cuticular appendages are affected, the hair, including the mane and tail, fall out, and if the eating of the plant continues, even the hoofs drop off. It has been suggested that *Leucaena* may thus become valuable as a homeopathic remedy for baldness.

Liana de cuello. See *Amphilophium paniculatum*.

Liana de la sierra. See *Bignonia acquinocialis*.

Liana fragante. See *Bignonia odorata*.

Liana unada. See *Bignonia unguis-cati*.

Libi-dibi. See *Caesalpinia coriaria*.

Lichi. See *Litchi chinensis*.

Lignum-vitae. See *Guajacum officinale*.

Lila. See *Duranta plumieri*.

Lilaila. See *Melia azedarach*.

Lima. See *Citrus hystrix*.

Lima bean. See *Phaseolus lunatus*.

Lime. See *Citrus hystrix*.

Limon. See *Citrus limonum*.

A tree from all parts of the island; height, 30 to 35 feet; diameter, 8 to 10 inches. Wood, yellow, hard; specific gravity, 0.984; not used. (Exp. 1857.)

Limoncillo. See *Amomis caryophyllata*.

A name applied to several myrtaceous shrubs and trees: *Amomis caryophyllata*, the bay rum tree, *Calyptranthes sintenisii*, *Eugenia stahlia*, and doubtless to other related species.

A tree from all parts of the island; height, 45 to 50 feet (13 to 15 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood, purplish, hard; specific gravity, 1.236; used for cabinet work. (Exp. 1857.)

Limoncillo de monte. See *Calyptranthes sintenisii*.

Limon dulce. See *Sweet lemon*.

Linociera compacta. See *Mayepea compacta*.

Lippia micromera. OREGANO DEL PAIS.

Family Verbenaceae; reported by Bello.

Lippia nodiflora. CIDRON.

This plant is known as an introduced weed in our Southern States and the West Indies generally, but Mr. David G. Fairchild, Agricultural Explorer of this Department, recently reports that it is considered of value in Egypt as a lawn plant

where all grasses or other plants tried for this purpose have failed. In Porto Rico coarse grasses, such as "guinea" and "para" grass, are abundant, but lawns are uncommon, and perhaps *Lippia* may be found of use. Mr. Fairchild's report on the subject is accordingly added:

According to Ascherson and Schweinfurth the *Lippia* is a native of Egypt. It has probably been used for lawn purposes for a great many years.

It is a low, creeping plant of the *Verbena* family, with broad, flat, obovate leaves of a deep green color. The creeping stems throw out roots wherever they come in contact with the earth, and form thick mats of herbage. It is well known that in regions with climatic conditions similar to those of Egypt, grass lawns are generally very difficult to maintain. Although there are several substitutes for lawn grasses, none that I have seen are as good as *Lippia*. Owing to its rapid growth, the plant can be mown closely, and to a layman the lawn effects resemble closely those produced by English lawn grasses.

In order to plant a lawn with *Lippia* the ground is prepared as it would be for the reception of grass seed. A mass of old *Lippia* is dug from some neighboring lawn or field. The native gardener cuts off or breaks off two or three long cuttings of the plant, makes a hole with a pointed stick in the soft earth, thrusts the cuttings, doubled up, into the hole and packs the earth securely about them. These cuttings are placed about 4 to 6 inches apart, quite irregularly over the field. They are given plenty of water, being sprinkled every day until well started. In winter, in Egypt, the lawns made of this *Lippia* are watered every four to five days, while in summer they are kept green by daily waterings. Every twenty days the lawns are gone over with a scythe, and in this way kept quite closely mown. There is no evident reason why a lawn mower would not answer the purpose better than a scythe.

Lawns of *Lippia* will last five to six years without renewing. Whenever a patch gets old or is injured by the shade of some tree, it is very easily repaired by setting new cuttings.

While, according to the statement of Mr. Colombo, of the Gizeh Gardens, no grasses form in Cairo a real sod, this plant produces a permanent sod lasting five to six years.

The *Lippia* deserves a thorough trial as a lawn plant in Southern California, Arizona, Texas, and Florida. Just what degree of hardiness it will show remains to be seen. It is not exposed to a temperature below freezing here in Cairo, except at extremely long intervals. Whether it is injured then or not I have been unable to ascertain. Although, during the hottest part of the summer, the lawns of *Lippia* wear a much less vigorous look than they do in winter, yet, from the fact that they are able to withstand the extreme heat and dryness of the Egyptian summer, it is evident that the plant is well suited for hot, dry climates. It is to be hoped this will prove a valuable new lawn plant for the parks and gardens of the South.

It is well to note that this plant is already quite commonly introduced, especially in the Southern States. It occurs in low, moist situations, from North Carolina to Florida, Texas, and Missouri, and is also present in California. So far as known it has not been utilized as a lawn plant in this country, although it is recognized as having some value as a sand-binder on the South Atlantic and Gulf coasts.

Lippia reptans. CIDRÓN.

(Stahl 6: 210.)

Licorice weed. See *Scoparia dulcis*.

Lira. See *Hibiscus schizopetalus*.

Litchi. See *Litchi chinensis*.

Litchi chinensis. LITCHI.

An important fruit in Southern China, British India, and the Malay Peninsula. The tree is described as a handsome evergreen, and, although it can be successfully propagated only by layering, it is reasonably hardy when once established. Writing from the standpoint of the northwest provinces of India, Dr. Bonavia, the fruit expert, says:

Here, then, is a fruit tree which resists the heaviest rains and stands the hottest winds, and also the frosts of these provinces. Moreover, it bears annually an abundant crop of fine, well-flavored, and aromatic fruit, which can readily be sent to distant markets without injury. Instead of being planted by the one or two,

it should be planted by the thousand. From all I know of the hardiness and fruitfulness of this remarkable tree, I feel confident that if any individual (or company), possessing the necessary capital, were to plant an extensive orchard of litchi trees, say where canal water would be easily obtained, or where well water is within easy reach, he would very probably make a good life-long business of it.

On this testimony the editor of Watt's Dictionary of the Economic Products of India comments as follows:

This result has been abundantly attained in Bengal, and, although statistics of the extent of the trade can not be given, it may be said that in the lower provinces the litchi tree is almost coextensively cultivated with the mango. It comes into season a little before that fruit, and in the larger cities, such as Calcutta, is sold in every fruit dealer's shop, the streets for a month or six weeks being literally bestrewn with the rind and large seeds, rejected by the wayside consumers. The fruit, to be enjoyed, should, however, be eaten as soon after being plucked as possible. When fresh the great bunches look like bright, pinkish strawberries, but they rapidly lose their bloom and assume a dirty, brownish color. The dried fruit, as sold in Europe, bears no possible resemblance to the deliciously bitter-sweet pulp of the fresh litchi.

The fruit is nearly round, and about an inch and a half (37 millimeters) in diameter. The edible portion is the sweet, semitransparent, jelly-like pulp or aril which covers the seed, and the whole is inclosed in a thin reddish or brownish brittle shell, which is rough with warty protuberances. The Chinese dry the fruit, which then becomes blackish, and in this state it may often be seen in London fruit shops. The fresh fruit has a very pleasant acid flavor, and is much liked both by natives and Europeans in this country.

With such definite and authoritative commendation, it would seem that the litchi is one of the fruits most likely to succeed in Porto Rico and likely to be of value when once generally introduced. The difficulty of propagation by cuttings is of course an obstacle, but it could easily be obviated by a few enterprising nurserymen. Little seems to have been written on the results of experiments which have been made with this tree in other West Indian Islands. The name "lichi" is mentioned in Maza's Diccionario of Cuban common names, and the species has also been recorded among the plants growing in the Castleton Gardens in Jamaica, but nothing has been found to indicate that any attempt has been made at extensive propagation for general purposes. Quite a large specimen of this species exists in one of the greenhouses of the Department of Agriculture.

The dried fruits described above are an article of considerable importance among the Chinese in this country, and are served in most of the Chinese restaurants. Boxes containing about a quart of these dried fruits were selling in one of the Washington fruit stores at 50 cents apiece.

Although these may bear no resemblance to the fresh fruit, they are by no means unpalatable. In appearance and taste they are not unlike raisins.

Llagrume. See *Didymopanax morototoni*.

Llanten. See *Plantago major*.

Llume. See *Aeria attenuata*.

Lluvia. See *Duranta plumieri* and *Russellia juncea*.

Lobelia.

Family Campanulaceae; a species of *Lobelia* is reported by Hill in a list of trees of the mountains. The species known from Porto Rico are all herbs except *L. acuminata* (*Tupa acuminata*), which grows to a height of about 10 feet.

Lobelia acuminata. TIBEY. TUPA.

A shrub 2 meters high, found in waste places and on limestone mountains. (Stahl, 8: 29, as *Tupa acuminata*.)

Lobelia assurgens. CHICORIA CIMARRONA.

An herbaceous perennial, 3 to 4 feet (about 1 meter) high, growing on mountains. Urban treats *Tupa assurgens* of Grisebach and Stahl as variety *portoricensis* under *Lobelia assurgens*. (Urban, Symb. 1: 454.)

Lobelia cliffortiana. CARDENALA AZUL.

An herbaceous, erect annual, found in rocky places. (Stahl, 6: 28.)

Lobelia portoricensis.

A shrub, 2 to 8 meters in height; known from Luquillo and Adjuntas. (Urban, Symb. 1: 453.)

Locust tree. See *Hymenaea courbaril*.

Logwood. See *Haematoxylon campechianum*.

Lombricera. See *Spigelia anthelmia*.

Lonchocarpus glaucifolius. GENO.

A leguminous tree of 5 to 8 meters, known only from Porto Rico, Rincon, Aguada, Aguadilla, and Quebradillas. A specimen from Aguada shows elliptical, finely veined leaves and flat, bright russet pods. (Urban, Symb. 1: 326.)

Lonchocarpus latifolius. PALO HEDIONDO.

A tree, 10 meters high, along the coast. Hard wood, used for furniture. Reported from Manati and from Bayamon as "palo seco." (Stahl, 3: 91.)

Lonchocarpus violaceus. GENO GENO.

A tree or shrub, 5 meters high, found on the west coast. (Stahl, 3: 92.)

Lonicera. HONEYSUCKLE.

One or more species are cultivated in gardens.

Loniciera. See *Linociera* and *Magepea*.

Loquat. See *Eriobotrya japonica*.

Loranthus americanus. HICQUILLO.

Reported by Bello. Urban describes this as a new species, *Dendropremon bicolor*.

Loranthus portoricensis. See note under *Psychotria pendula*.

Lucuma mammosa. SAPOTE. MAMEY SAPOTE.

Family Sapotaceae: a tree, 10 meters high; yields the "marmalade fruit;" also called the "vegetable egg." Grisebach spells the native name "mamme sapota." (Stahl, 6: 51.)

Grosourdy (2: 398) says of the wood that it is fine-grained and compact, ordinarily hard, and of good weight. In color it resembles mahogany, but is redder and more uniform, though somewhat mottled with darker-colored spots. Specific gravity, 0.588. Its most common use is for shelving, but it would appear suitable for carpentry and cabinetwork. The fresh bark has the odor of bitter almonds.

This fruit is rare in Porto Rico, and the plant is probably introduced. The common names are, however, often incorrectly applied to the fruit of *Mammea americana*, which it somewhat resembles in external appearance. The smooth seeds with a roughened segment are, however, entirely unlike those of the latter species, which are rough over the entire surface, with the flesh closely adhering. In Central America and Mexico this is a very common wild fruit, and in times of want it serves to some extent as a food. The flavor is mild, by some considered insipid, and it will probably never become popular among people of temperate regions.

As the common name "marmalade fruit" implies, it is made into a marmalade, and this is not unlike good apple butter.

It was this fruit that kept Cortez and his army alive on their famous march from Mexico City to Honduras.

Lucuma multiflora. JACANA.

A large tree, 8 to 10 meters high, yielding a useful wood. Bello gives the common name as "hacana." (Stahl, 6: 50.)

Lucuma stahlia.

Reported from near Manati.

Lucuma urbani.

Reported from near Guanica.

Luffa acutangula. SPONGE CUCUMBER. ESPONJA.

Reported by Bello; also called "estropajo." A species of *Luffa*, probably this or *L. aegyptiaca*, was very common in Porto Rico, climbing over bushes and fences. It is not known whether the matted fibers of the fruit are used or not, but its increasing employment for the toilet and for other purposes, such as the linings of hats and the porous soles of slippers, is rendering it quite an article of export from Japan. In the younger state the fruits may be eaten as cucumbers or pickled.

Luffa cylindrica. See *Luffa aegyptiaca*.**Lycopersicon cerasiforme.** TOMATO AMARILLO.

Family Solanaceae.

Lycopersicon esculentum. See *Lycopersicon lycopersicum*.

(Stahl, 6: 126.)

Lycopersicon humboldtii. TOMATE.

(Stahl, 6: 125.)

Lycopersicon lycopersicum. TOMATO. TOMATO GRANDE.

Tomatoes, like many other vegetables that have been developed in temperate regions, are with difficulty made to succeed in the Tropics. Good tomatoes are, however, produced in Porto Rico, and with the use of fertilizers fruits are reported from the north side of the island weighing as high as 11 ounces.

Maba inconstans. NEGRA LORA.

Family Ebenaceae. (Stahl, 6: 60.)

Maba sintenisii. TABEIBA.

Also called "guayabota nispero;" a native forest tree, about 30 feet (9 meters) high, reported from the vicinity of Lares. (Urban, Add. 1: 43.)

Mabi. See *Colubrina reclinata* and *C. ferruginosa*.

A drink by this name is commonly sold in the markets and peddled on the streets. It is usually taken as a "refresco," and is also reputed to be a remedy for indigestion. The bark from which it is made is also offered in the markets.

Mabie. See *Colubrina reclinata*.

Said to be the St. Lucia form of the word "mabi."

Macachaira. See *Manihot palmata*.**Macaw bush.** See *Solanum mammosum*.**Macaw palm.** See *Acrocomia fusiformis*.

According to Grisebach the "mackaw tree" is *Acrocomia sclerocarpa* (= *A. aculeata*), while the "great mackaw tree" is *A. lasiospatha* (= *A. fusiformis*).

Maclura tinctoria. See *Chlorophora tinctoria*.**Madre de cacao.** See *Erythrina umbrosa*.

This name is not known to be used in Porto Rico, but is of South American origin, and has been referred to by Hill and other writers.

Maga. See *Thespesia grandiflora*.

A tree from the interior of the island; height, 45 to 50 feet (14 to 15 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood light, hard; specific gravity, 0.972; used in building houses. (Exp. 1857.)

Color, rich chocolate brown. Conspicuous ducts few, occurring singly, and irregularly scattered between the medullary rays; mostly distinct or occasionally two or three loosely grouped. Medullary rays comparatively few, distinct, but inconspicuous. Wood fibers slightly cross-grained, the smoothed surface appearing straight-grained. Quarter and radially cut medullary rays have a satiny

appearance. The rich color and attractive grain of this wood should make it valuable for cabinetwork. (Hill and Sudworth.)

This name is also given by Hill among those of trees planted for shading coffee. We could not learn that the magar is ever so used; perhaps an error for "moca."

Magar. See *Thespesia grandiflora*.

Magnolia. See *Talauma plumieri*.

Magnolia portoricensis. MAURICIO.

A handsome indigenous tree discovered by Bello, differing from *M. splendens* in having much broader and more rounded leaves. It is known only from the western half of the island—Utuado, Jayuya, Adjuntas, Guayanilla, Peñuelas, and Lares. (Urban, Symb. 1: 306).

Magnolia splendens. SABINO.

Also called "laurel sabino." A species recently described by Urban from the primeval forests of the Sierra de Luquillo and Sierra de Naguabo. Said to be one of the most beautiful species, similar in habit to *Talauma plumieri*, for which it was probably mistaken by Eggers in the account of his visit to Porto Rico, as quoted from Nature by Hill. The leaves are said to be used by the natives as a condiment, like those of *Laurus nobilis* of Europe.

Mago.

A tree found in all parts of the island; height, 30 to 40 feet (9 to 12 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood pinkish, soft; specific gravity, 0.791; not used. (Exp. 1857.)

Mahagua. See *Paritium tiliaceum*.

An erroneous spelling of "majagua."

Mahoe.

A name applied in the British West Indies to various malvaceous trees, such as *Paritium*, *Hibiscus*, and *Thespesia*.

Mahogany. See *Swietenia mahagoni*.

Mais, Maize. See *Zea mays*.

Majagua. See *Paritium tiliaceum*.

A soft wood, used for laths in hut building, the bark used for fastening. All ropes made here are from the majagua bark. (Hansard.)

Majagua quemona. See *Pisonia obtusata*.

Malachra alceifolia.

Family Malvaceae; variety *rotundifolia* reported from near Aibonito, along roads.

Malachra capitata. MALVA DE CABALLO.

An herbaceous annual, becoming woody with age.

Commonly seen about dwellings and moist places. A characteristic species with light-yellow flowers. We found it cultivated in a dooryard near Coamo, where a decoction was supposed to have medicinal properties in external application. In this locality the plant was known simply as "malva," as also stated by Bello. (Stahl, 2: 74.)

Malachra ciliata. See *Malachra urens*.

Malachra cordata. MALVA ACORAZONADA.

Family Malvaceae. (Stahl, 2: 77.)

Malachra palmata. MALVA DE CABALLO.

An herbaceous annual, found in all parts of the island. (Stahl, 2: 76.)

Malachra radiata. MALVA BLANCA.

A woody, annual herb, 1 meter high, found in stony places. (Stahl, 2: 75.)

Malachra urens. MALVA PESTAÑOSA.

(Stahl, 2: 77, as *Malachra ciliata*.)

Malagueta. See *Annonis caryophyllata*.

Malanea macrophylla. PALO LLORON.

Family Rubiaceae; a shrub 1 meter high, found among brambles. (Stahl, 5: 93.)

Malojillo. See *Panicum molle*.

In Cuba the present name does not appear to be used; Mazo gives "yerba para," "yerba del para," "yerba del paral," and "yerba de rarana."

Malonillo. See *Poinciana regia*.

According to Captain Hansard this name is extensively used in the northeastern part of the island instead of "flamboyant."

Malpighia coccigera. AZOTA-CABALLO.

A shrub 3 to 4 meters high; cultivated for its agreeable fruits, which resemble the common cherry and make good tarts and jelly. (Stahl, 2: 143.)

Malpighia urens. PALO BRONCO.

A shrub 4 meters high, along the seacoast; fruit and bark are used as astringents. The hairs on the under side of the leaves pierce the skin and cause severe pain. (Stahl, 2: 144.)

Malta arborea.

A tree from the northern part of the island; height, 12 to 15 feet (4 to 5 meters); diameter, 8 to 9 inches (20 to 22 centimeters). Wood light, very hard; specific gravity, 0.977; fruit tree. (Exp. 1857.)

Malta de mata.

A tree from the northern part of the island; height, 8 to 10 feet (2.5 to 3 meters); diameter, 4 to 6 inches (10 to 15 centimeters). Wood light, very hard; specific gravity, 0.890; fruit tree. (Exp. 1857.)

Maluco. See *Spilanthes acmella*.

Malva. See *Malachra capitata*.

A root called malva was being sold by an herb dealer in the market of Ponce as a remedy for pulmonary diseases.

Malva acorazonada. See *Malachra cordata*.

Malvabisco. A form of the name "malvavisco."

Malva blanca. See *Malachra radiata*.

Malva de caballo. See *Malachra capitata* and *M. palmata*.

Malva pestañosa. See *Malachra urens*.

Malvastrum spicatum. MALVAVISCO.

Family Malvaceae; a woody herb, found on south coast; rare on the north coast. (Stahl, 2: 78.)

Malva té. See *Corchorus hirtus* and *C. siliquosus*.

Malva té de la playa. See *Corchorus hirsutus*.

Malvavisco. See *Malvastrum spicatum*.

Malvavisco afelpado. See *Hibiscus tomentosus*.

Malvavisco cimarron. See *Abutilon umbellatum*.

Malvavisco purpureo. See *Hibiscus phoeniceus*.

Mamey. See *Mammea americana*.

A tree from all parts of the island; height, 40 to 45 feet (12 to 14 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood pinkish, hard; specific gravity, 0.978; fruit tree. (Exp. 1857.) The gum is used as incense.

Mamey del cura. See *Ternstroemia elliptica* and *T. stahlii*.

Mamey sapote. See *Lucuma mammosa*.

Mameyuelo. See *Ardisia coriacea*, *A. glauciflora*, and *A. purpurascens*.

A tree from all parts of the island; height, from 40 to 45 feet (13 to 14 meters); diameter, 9 to 10 inches (23 to 25 centimeters). Wood white, hard; specific gravity, 0.879; used in building houses and cabinetmaking. (Exp. 1857.)

Mamie apple. See *Mammea americana*.

Mammea americana. MAMMEE. MAMEY. MAMIE APPLE.

Family Clusiaceae; a handsome tree, 40 to 60 feet (12 to 18 meters) high; fruit large, with a pleasant taste and aromatic smell, eaten raw or preserved in sugar. Wood valued for timber; gum used by natives in extracting chigoes, etc. (Lunan.)

From between Fajardo and Ceiba. (Stahl. 2: 124.)

See note under *Lucuma mammosa*.

Mammee. See *Mammea americana*.

Mammee apple. See *Mammea americana*.

Mammee sapota. See *Lucuma mammosa*.

Mamoncillo. See *Melicocca bijuga*.

Manchineel. See *Hippomane mancinella*.

Mandioca. See *Manihot*.

Mangifera indica. MANGO.

Family Anacardiaceae; a fruit tree native in South Asia or the Malay Archipelago and introduced into America about the beginning of the eighteenth century.

It not infrequently happens that those who are visiting a tropical country for the first time confuse the mangrove and the mango. The former is a tree which forms tide-water swamps along the coasts and about the mouths of rivers in nearly all humid tropical countries, while the mango is an edible fruit, even more widely distributed.

In Porto Rico the name mangrove or "mangle" is applied to three or four different trees which grow in the coast swamps, while the mango is the best known and most abundant of fruits. Whether associated with ideas of unwholesomeness through the above confusion or other unreliable information, many of the army officers in Cuba and Porto Rico manifested a violent prejudice against the mango, and some even issued orders strictly forbidding the soldiers, under any circumstances, to eat it. The antimango crusade has even gone to the extent of cutting down mango trees in public grounds.

The only excuse for this mistaken policy seems to lie in the report that many soldiers were made sick by them when the armies were first landed, and this on being sifted a little resolves itself into the proposition that a hearty meal of green mangoes, followed by a large quantity of bad rum, made the hungry and fatigued soldiers very sick. Before the mango is finally condemned it might be well to consider what the results would have been with green apples.

In reality the mango is known throughout the Tropics as a delicious and wholesome fruit, comparable in quality and value with the apple or the orange, although entirely different from either in texture and flavor. Some varieties have, especially when unripe, a distinct flavor and odor of turpentine, which seems objectionable at first, but soon comes to add charm to the high qualities of the new favorite. It has been said that one should eat mangoes only in a bath tub, on account of keeping the yellowish pulp from spreading over the countenance and person, and this might prevent the fruit becoming popular with the most fastidious. This objection, however, applies only to the poorer varieties, such as at present exist in

Porto Rico. The finer, asexually propagated varieties are quite as manageable as the peaches and pears of our own climate.

Several of the best Indian varieties have already been introduced into Florida and the West Indies, and when these are produced in sufficient quantities to be placed regularly in the Northern markets the mango is almost sure to become a popular fruit.

The drier portions of Porto Rico are admirably adapted for growing mangoes, and the absence of all danger from frost places this island at a decided advantage over Florida, where the best varieties are now being propagated on an extensive scale by several of the more enterprising fruit growers. It has been found possible to ship even the better varieties from India to London, consequently there need be no fear as to the shipping qualities.

For a discussion of the possibilities and requirements of the mango in Porto Rico, see Bulletin 28, Bureau of Plant Industry, U. S. Department of Agriculture.

The wood is light, soft, fibrous in texture, and not very compact, brownish gray with little spots and irregular lines of brown. After varnishing this wood resembles light-colored walnut. It is resistant and elastic, and is used for the same purposes as the ash of Europe—for gunstocks, etc. It is not, however, used as much as it deserves. Specific gravity, 0.864. (Grosourdy, 2: 400.) Specific gravity also given as 0.738. (Exp. 1857.)

Mangle blanco. See *Avicennia nitida* and *Laguncularia racemosa*.

Grosourdy (2: 399) gives this common name to both *Avicennia tomentosa* (= *A. officinalis*) and *Laguncularia racemosa*, and describes the wood as being hard and heavy, fibrous and compact in texture, and gray in color with rather brown undulations, which produce a very pleasing aspect. Used for the small timbers of houses, but its most common use is in making a good charcoal and for fuel. The specific gravity of *Avicennia* is 0.867; of *Laguncularia*, 0.860.

Mangle bobo.

A tree from near the seashore; height, 25 to 30 feet (8 to 9 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood, dark, hard; specific gravity, 1.124; used for cabinetwork. Also called "mangle prieto." (Exp. 1857.)

A good firewood; on the higher elevations of El Yunque this grows as mere saplings, while in the lowlands it reaches 8 or 10 inches (20 to 25 centimeters) in diameter. (Hansard.)

Bello gives this common name with *Avicennia nitida*, but this is certainly not the tree reported by Captain Hansard.

Mangle botanico.

A tree from the beaches of the island; height, 20 to 25 feet (6 to 8 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood, silver-gray, hard; specific gravity, 1.072; used in making knees of boats. (Exp. 1857.)

Mangle boton. See *Conocarpus erectus*.

Mangle cobo.

A tree from the beaches of the island; height, 18 to 20 feet (5.5 to 6.5 meters); diameter, 9 to 10 inches (22 to 25 centimeters). Wood, white, soft; specific gravity, 0.840; not used. (Exp. 1857.)

Mangle colorado. See *Rhizophora mangle*.

A tree from the beaches of the island; height, 18 to 20 feet (5.5 to 6.5 meters); diameter, 9 to 10 inches (22 to 25 centimeters). Wood, pinkish, hard; specific gravity, 0.752; bark used in tanning leather. (Exp. 1857.)

Mangle prieto.

A tree from the beaches of the island; height, 25 to 30 feet (8 or 9 meters); diameter, 10 to 12 inches (22 to 25 centimeters). Wood, dark, hard; specific gravity, 1.124; used for cabinetwork. Also called "mangle bobo." (Exp. 1857.)

Mango. See *Mangifera indica*.

Mangrove. See *Rhizophora mangle*.

Mangrove, Black. See *Avicennia nitida*.

Mangrove, Olive. See *Avicennia nitida*.

Mangrove, White. See *Laguncularia racemosa* and *Avicennia nitida*.

Maní. See *Arachis hypogaea*.

Manihot aipi.

Considered as a variety of *Manihot palmata*.

Manihot cassava. CASSAVA. YUCA.

Family Euphorbiaceae; shrubby plants, 6 to 8 feet (1 to 1.5 meters) high; cultivated for the thick, fleshy roots, which are very rich in starch.

Cassava is a less important article of food in Porto Rico than in most other parts of the American tropics. Its place is to a great extent taken by "yantia" and "taro." The principal uses to which cassava is put in Porto Rico are the making of starch and a sort of bread baked in thin, semicircular loaves or cakes. This latter custom is undoubtedly indigenous, and curiously enough finds its counterpart among the so-called Caribs of Guatemala, a negro people of uncertain origin, with whom cassava is by far the most important food.

Forms with both 5 and 7 leaflets were observed, and it was inferred that both the sweet and the bitter varieties exist, although nothing was heard of the poisonous properties of any of the cassava grown in the island, but as none is eaten raw or used as fodder, such might easily exist and fail to be in evidence.

Captain Hansard's notes on culture are as follows: "Hot climate, coast region; plow the soil; cut stem in 6-inch lengths; plant 5 feet apart, leaving about 1 inch out of the ground; plant on a dry day. It will begin to grow in two weeks, and in three months will cover the ground. Will be ready to dig in nine months. Spoils quickly out of the ground, but keeps until dug. To prepare meal, tubers are pulled, washed, grated, put in bags, and the juice pressed out. The fibers are sifted out and the meal dried by a fire, but should not be scorched. Proper drying and heating draws off the juice."

Manihot palmata aipi. SWEET CASSAVA. YUCA.

According to Engler and Prantl *M. palmata aipi* in Brazil is called "aipim," "macacheira," or "sweet mandioca."

Sweet cassava, as a fodder crop, has received considerable attention, but in Florida, where the most extensive experiments have been tried, velvet beans have been found to produce cheaper fodder with less labor, and with the added advantage that the soil is thereby enriched.

As a starch producer the outlook is perhaps more promising, cassava starch being of a very fine quality. Several factories are in operation in Florida, but it is still too early to report on the success of the enterprise.

Manihot utilissima. CASSAVA. YUCA.

This is the so-called "bitter cassava," and usually contains sufficient hydrocyanic acid in the roots to render them poisonous if eaten before taking precautions to remove or destroy it. This can be accomplished by cooking or by expressing the juice, after which it is a very wholesome food.

Cassareep is the juice of this plant boiled down to a gummy consistency. It is used in the British West Indies to preserve fresh meat or in the dish called "pepper pot." It is also exported to England for making sauce.

An antidote used by the Indians for the poison of this plant consists of red peppers steeped in pure rum.

Manioc. See *Manihot*.

Manzanilla.

A name in use for several composite weeds, the most common of which is *Bidens leucantha*. Stahl proposes to distinguish the different species according to the following list:

Manzanilla cimarrona. See *Stemmodontia carnosu* and *Bidens leucantha*

Manzanilla de costa. See *Stemmodontia buphthalmoides*.

Manzanilla de la playa. See *Stemmodontia carnosu*.

Manzanilla de monte. See *Stemmodontia reticulata*.

Manzanilla del pais. See *Bidens pilosa*.

Manzanilla negra. See *Bidens bipinnata*.

Manzanilla trepadora. See *Bidens rubifolia*.

Manzanillo. See *Hippomane mancinella*.

Mapola. See *Hibiscus rosa-sinensis*.

In the Philippines and Guam this name is applied to *H. mutabilis*.

Mapurito. See *Fagara monophylla*.

Maraga.

Perhaps the same as "maraca" or "matraca;" applied at Cataño to a species of *Canna*.

Marañon. See *Anacardium occidentale*.

Maranta arundinacea. ARROWROOT.

Family Marantaceae; an herbaceous plant with tubers from which arrowroot is extracted. Said to be called arrowroot from its property of counteracting the poison put on arrow tips by the natives and other acrid poison, as that of spiders. Called touola by the natives of Jamaica.

The following recommendations are from Captain Hansard:

"Elevation not important; rows 3 feet (90 centimeters) apart, 12 inches (30 centimeters) in a row. Cut off flowers to throw strength into the tubers. Plant in April or early May; returns will be had in a little less than twelve months. Twenty per cent starch may sometimes be secured. Crop seven hundredweight (329 kilograms) per acre, but double this has been obtained by high cultivation. In preparation all skin and discolored parts are removed from the rhizomes. After rewashing the rhizomes are reduced to pulp in a mortar or against a revolving cylinder grater. Cleanness is very necessary. In St. Vincent glass-roofed houses and the very cleanest machinery are used."

Maravedi. See *Myginda rhacoma* and *M. pallens*.

Maravilla. See *Hibiscus mutabilis*, *Mirabilis jalapa*, and *Euphorbia heterophylla*.

Maray-maray. See *Dalbergia ecastaphyllum*.

Marcgravia rectiflora jacquini. BEJUCO DE PALMA.

An indigenous trailing or climbing shrub known also as bejuco de rana. According to Urban, *Marcgravia umbellata* of Grisebach's Flora includes this species, but the genuine *umbellata* does not occur in Porto Rico.

Marcgravia sintenisii. BEJUCO DE PALMA.

A species peculiar to Porto Rico and said to be very distinct from *M. rectiflora*. It is a climber and ascends large trees. Reported from the Sierra de Luquillo, Sierra de Naguabo, Cayey, and Adjuntas. (Urban, Add. 3: 208.)

Marcgravia umbellata. PEGA PALMA.

A trailing shrub clinging to trees by adventitious roots, called "bejuco de palma," by Bello. (Stahl, 2: 129.)

Margarita. See *Cosmos caudatus* and *Bidens pilosa*.

Mari-Lopez. See *Turnera ulmifolia* and *T. pumilea*.

Maria.

A tree from all parts of the island; height, 45 to 50 feet (13 to 15 meters); diameter, 25 to 30 inches (62 to 75 centimeters). Wood pinkish, hard; specific gravity, 0.800; used in building houses. (Exp. 1857.)

Enumerated by Hill among trees yielding resin. Perhaps the same as "palo de Maria."

According to Captain Hansard this species yields the balsam of Tolu. The extract is not made in Porto Rico, but extensively used by the druggists. Specific gravity of wood, 0.91.

Marias. See *Calophyllum calaba*.

Marialva elliptica. See *Toromita elliptica*.

Maricao. See *Byrsonima spicata*.

A tree from all parts of the island; height, 40 to 45 feet (13 to 15 meters); diameter, 25 to 30 inches (62 to 75 centimeters). Wood dark, hard; specific gravity, 0.730; used in building houses. (Exp. 1857.) (Grosourdy, 2: 401.)

Captain Hansard gives the specific gravity at 0.78, and says that it is a good firewood.

Marimbo. See *Lagenaria vulgaris*.

Marimbo is the name applied to a Central American musical instrument resembling a xylophone, which in former times had gourds hung beneath the keys. In some parts of Africa the name is given to a kind of drum.

Marimbo amargo. See *Lagenaria vulgaris*.

Marlierea dussii.

A small indigenous myrtaceous tree, known in Porto Rico only from the neighborhood of Adjuntas. In Guadeloupe the same species is said to grow as a shrub of 15 feet (5 meters) or less, half the size it attains in Porto Rico. *M. sintenisii*, an allied species, is known only from Luquillo. (Urban, Add. 108.)

Marmelluelo.

Captain Hansard mentions this wood as having a specific gravity of 1.02.

Martinezia.

Specimens of the flowers and fruit supposed by Dr. Drude to belong to *Martinezia* or *Bactris* (Sintenis no. 2500) were collected in the forest of Monte Hormigas near Jauco. The spathe is unarmed; the branches are long, slender, and attenuate. This may be the species collected by us at Vega Baja. There are no leaves with this specimen. Another specimen (Sintenis no. 2628) from Hato Grande, on slopes toward Monte Gregorio, has a much larger and thicker inflorescence with the main axis beset with slender black spines from one-quarter to 1 inch (6 to 25 millimeters) in length. The branches are much coarser than in the other specimen and the pinnae have irregularly crowded plications at base, showing a tendency to the grouped condition found in the South American species. There are no spines on the pinnae, and very scattering slender ones on the median carina of the rachis.

Marrubio. See *Mesosphaerum suaveolens* and *M. spicigerum*

Marrubio blanco. See *Mesosphaerum lantanaefolium*.

Marrubio-boton. See *Mesosphaerum capitatum*.

Marrubio oscuro. See *Mesosphaerum atrorubens*.

Marsypianthes hyptoides. ORTELA.

An herbaceous suberect annual of the mint family, found in sandy places. (Stahl, 6: 191.)

Martin avila. See *Chione glabra*.

Martynia diandra. UÑA DE GATO.

Family Martyniaceae; an annual herb found on the south coast. Bello gives the common name "escorzonera." (Stahl, 6: 265.)

Marunguay. See *Zamia erosa* and *Z. portoricensis*.

Masa. See *Hedwigia balsamifera*.

A tree from the eastern part of the island; height, 35 to 40 feet (11 or 12 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood ash-colored, hard; specific gravity, 0.827; used in building houses. (Exp. 1857.)

Mastic.

The true mastic tree, *Pistacia lentiscus*, is a native of the Mediterranean region, and is not known to have been introduced into the West Indies, where the name is applied to *Bursera simaruba*, also called the West Indian mastic.

Mastuerzo. See *Lepidium virginicum*.

Mata de niguas. See *Tournefortia hirsutissima*.

Mata de peo. See *Lasianthus lanceolatus*.

Mata de seda. See *Calotropis procera*.

Mata-gallinas. See *Solanum caribaeum*.

Matayba domingensis. RATON.

A tree of 10 to 15 meters; family Sapindaceae. Reported from Adjuntas, Utuado, and Sierra de Luquillo. *M. apetala* is also reported from Utuado and Luquillo. (Urban, Symb. 1: 351.)

Matillo.

A tree from all parts of the island; height, 25 to 30 feet (8 to 9 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood red, rather hard; specific gravity, 0.613; used for lumber. (Exp. 1857.) (Grosourdy, 2: 401.)

Mato. See *Dioclea reflexa*.

Mato amarillo. See *Caesalpinia bonducella*.

Mato azul. See *Caesalpinia bonduc*.

Mato blanco. See *Caesalpinia bonduc*.

Mato colorado. See *Canavalia rusiosperma*.

Red sea beans on sale by an herb dealer in the market of Ponce as a remedy for piles, used in combination with the gray beans of *Caesalpinia bonducella*, as described under "tortera."

Mato de la playa. See *Canavalia obtusifolia*.

Mato palo.

A moderately abundant wild tree, 40 to 50 feet (12 to 15 meters) high, with a rather long, straight trunk 6 inches (15 centimeters) in diameter. It furnishes a rather light soft wood, fibrous in texture, soft and fine-grained. Yellow, or sometimes light cinnamon-colored, with numerous parallel lines, giving it a very beautiful aspect. Its most common use is in making small boats, tubs, and wash basins. Specific gravity 0.450. Also called "gaguey" and referred to the genus *Ficus*. (Grosourdy, 2: 392.)

Matos. See *Ormosia krugii* and *Mucuna urens*.

Matos colorados. See *Canavalia ensiformis*.

Matos de playa. See *Caesalpinia bonduc*.

Matraca.

The Spanish word for rattle; applied in Porto Rico to rattles made from the calabash tree, *Crescentia cujete*, and also to *Crotalaria retusa*, the ripe seeds of which become loosened and rattle in the pod.

Mauricia. See *Mauritia flexuosa*.

Mauricio. See *Magnolia portoricensis*.

Mauritia flexuosa.

A palm mentioned and described by Hill under the names "moriche" and "palma moriche." The species is a tall South American fan-palm, native in the valleys of the Orinoco and Amazon. It grows in great abundance and forms extensive forests in level regions subject to annual overflow, and furnishes a large part of the material of subsistence and native arts of whole tribes of Indians. It is not, however, known from Porto Rico or, as a native, from any part of the West Indies. According to Maza it and *Mauritia setifera* (probably an error for *setigera*, a species native in Trinidad) are cultivated in Cuba and called "palma mauricia." As a useful tree which will flourish in swampy places this species might be of some utility in Porto Rico.

Mauritia setifera and Mauritia setigera.

See note under *Mauritia flexuosa*.

Mauritius hemp. See *Furcraea gigantea*.**Maya.**

A name applied to the fruits of the wild piñuela, *Bromelia pinguin*, used as a remedy for worms.

Maypea caribaea. AVIPILLO.

Family Oleaceae; a tree reported as collected by Sintenis at Fajardo—perhaps the same as Abispillo of Captain Hansard's list.

Maypea compacta. PALO DE HUESO.

A tree 10 to 12 meters high. White, hard wood used for heavy timbers. (Stahl, 6: 60, as *Linociera compacta*.)

Maypea domingensis.

Variouly named "palo de hueso," "huesillo," "hueso blanco." A native tree reported from Bayamon, Caguas, Sierra de Luquillo, Utuado, and Quebradillas. It is the same as Stahl's *Linociera compacta*. An allied species, *M. arilliflora*, comes from mountain forests at "El Manil." (Urban, Add. 1: 61.)

Maza.

Not used for special purposes; a common wood in the hills; specific gravity, 0.87. (Hansard.)

Medicago sativa. ALFALFA.

Alfalfa has not been found successful in humid tropical countries. A substitute has been suggested in phasemy (*Phaseolus semierectus*).

Medlar.

Reported by Hill among "forest trees yielding fruits." The reference is, however, to be understood as applying to *Achras sapota*, the sapote or sapodilla, neither the true medlar, *Mespilus germanica*, nor the Japanese medlar or loquat, *Eriobotrya japonica*, being known to the Porto Rican public.

Meibomia. ZARZABACOA.

A large genus of leguminous herbs, formerly known under the name *Desmodium*. Several of the species are useful forage and soiling plants.

Meibomia adscendens. ZARZABACOA GALANA.

A creeping woody annual found in dry pastures. The flowers are pale lavender or whitish. (Stahl, 3: 42, as *Desmodium adscendens*.)

Meibomia axillaris. ZARZABACOA DE MONTE.

A creeping or trailing annual, frequenting shady places in mountain districts. This species is too diffuse in habit and too woody in structure to be likely to prove of economic value.

Meibomia barbata. ZARZABACOA PELUDA.

A shrubby species found in dry pastures. Known from Lares. (Stahl, 4: 45, as *Desmodium barbatum*.)

Meibomia incana. ZARZABACOA COMUN.

An annual or biennial shrub found in pastures. The variety *angustifolia* is reported from Rincon. The general habit of this species seems not unlike that of *M. axillaris*, and economic utility is improbable. (Stahl, 3: 43, as *Desmodium incanum*.)

Meibomia mollis. ZARZABACOA DULCE.

Found on the west coast. Also known from Juana Diaz. (Stahl, 3: 49, as *Desmodium molle*.)

Meibomia scorpiurus. ZARZABACOA COLA DE ESCORPION.

An herb 50 centimeters high, prostrate, common in arid pastures and stony places. (Stahl, 3: 46, as *Desmodium scorpiurus*.)




FIG. 12. Lawn of *Meibomia triflora*. Coamo Springs.

Meibomia spiralis. ZARZABACOA ESPIRAL.

A slender species attaining a height of 2 feet. A form much more delicate and leafy than usual was collected at Coamo Springs (No. 716). This might be of interest if the plant should be found of economic utility. In the Index Kewensis the present species is treated as a synonym of *Desmodium neomexicanum*, but specimens in the National Herbarium do not indicate close affinity.

Meibomia tortuosa. FLORIDA BEGGAR-WEED. ZARZABACOA TORCIDA.

A shrub, 1 meter high, found along roadsides and in moist mountainous places. This is the species now somewhat extensively cultivated in our Southern States as a forage and soiling crop. From its habit and general characteristics it would seem that the form mentioned under *Meibomia spiralis* might be preferable, at least from the forage standpoint. (Stahl, 3: 48, as *Desmodium tortuosum*.)

Meibomia triflora. ZARZABACOA DE TRES FLORES.

FIGURE 12.

A small herbaceous annual, found in dry places along the coast. This species was collected in a dry, exposed situation at Coamo Springs, where it covered the

otherwise bare ground with a dense mat of green, the small leaflets giving a very pleasing effect like a white clover lawn and suggesting a possible use for this plant. (Stahl, 3: 41, as *Desmodium triflorum*.)

Meladillo. See *Plumbago scandens*.

Melanthera deltoidea. CARIAQUITO BLANCO.

A perennial composite herb 1 meter high, found in waste places and on the lower slopes of mountains. (Stahl, 5: 140.)

Melia azedarach. LILAILA.

PLATE XLV.

A meliaceous tree or shrub commonly known as "China berry" or "pride of China," introduced from the Old World. It grows 5 or more meters high and is found as an ornamental in gardens, etc. The bark of the roots is said to be an efficacious vermifuge, while the pulp of the fruit is considered poisonous. (Stahl, 2: 164.)

The wood is rose-colored, rather hard, breaking vertically. Used by the country people in making tool handles. (Grosourdy, 2: 360.)

This author spells the native name "alilaila."

The china-berry is very commonly planted in Porto Rico as a shade or rather as an ornamental tree. The Porto Rican tree may, however, be a different species or variety, since it has not at all the umbrella-like habit found in the United States and in the Canary Islands. It grows with great rapidity, sometimes sending up a central stem 20 feet (6 meters) high in a year and a half. At a height of about 10 feet (3 meters) lateral branches from 4 to 6 feet (1.2 to 1.9 meters) long are emitted, while those of the upper part remain short, giving the tree a narrowly pyramidal form. The slender top often bends over or is broken off. The foliage is rather loose and open; flowers and fruit are produced in great abundance. The extremely rapid growth might give this species a limited use in coffee culture, particularly where it is desired to protect without loss of time an exposed slope; and the numerous slender branches of the tree would also enable the shade to be removed gradually. Considerable shelter from winds and a moderate degree of shade could be secured by planting the china-berry from 2 to 4 rods apart.

Melia sempervirens. See *Melia azedarach*.

Melicocca bijuga. GENIP TREE. GENIPE.

A tree belonging to the family Sapindaceae, believed by some botanists to be native in the West Indies, by others thought to have been introduced from South America. It produces clusters of oval fruits about an inch in length. The pulpy covering of the seed is edible and has a pleasant flavor, but the fine fibers compact themselves against the mucous membranes and cause a very disagreeable sensation. In the Bulletin of the Jamaica Botanical Department it is stated that this material has caused the death of children by forming a coating over the lining of the stomach.

The tree is vigorous and handsome and has been recommended for planting along roads and avenues for shade. The timber is said to be hard and heavy.

Radlkofer gives the following collection of names for this tree: Honigboere, mamon, mamoncillo, knippa, knepier, guenepe, and genep tree. The Porto Rican pronunciation heard by us was, however, distinctly "genepe," not "guenepe," or "quenepa."

Meliosma herbertii.

Family Sabiaceae; shares with *M. obtusifolia* the names "cacao bobo" and "aguacatillo." A tree generally distributed in Porto Rico and the adjacent islands as far south as Grenada. Of medium size, with alternate, exstipulate, elliptical, or obovate acute leaves. (Urban, Symb. 1: 55.)

Meliosma obtusifolia. GUAYROTE.

Also called "arroyo," "aguacatillo," "cacao bobo," "cacaillo," "ciralillo," "serillos." A tree belonging to the family Sabiaceae, generally distributed



CHINA BERRY (MELIA AZEDARACH).

throughout the island. It yields a valuable wood, and from the multiplicity of common names is evidently in use locally, though all the names are probably not confined in application to this species. Reported from near Adjuntas and Sierra de Naguabo. Differs from *M. herbertii* in having the leaves obtuse and the fruits very small, only half the size of the others. (Urban, Symb. 1: 513; Stahl 6: 43, as *Atelandra obtusifolia*.)

Melocactus communis. MELON DE COSTA.

Family Cactaceae; a fleshy plant 25 to 30 centimeters in diameter, found in dry places and cultivated. (Stahl, 4: 184.)

Melochia hirsuta. BRETONICA PELUDA.

Family Sterculiaceae; an herbaceous, erect, shrubby annual, densely covered with soft hairs. (Stahl, 2: 109.)

Melochia nodiflora. BRETONICA PYRAMIDAL. BRETONICA PRIETA.

A woody annual, 50 centimeters high, found in rocky places. (Stahl, 2: 110.)

Melochia pyramidata. BRETONICA PYRAMIDAL.

A woody herb, 50 centimeters high, in sandy and stony pastures along the south coast. (Stahl, 2: 107.)

Melochia serrata. BRETONICA ASERRADA.

According to Grisebach a shrub about 2 feet high, covered with long, soft hairs. (Stahl, 2: 109.)

Melochia tomentosa. BRETONICA AFELPADA.

A shrub 1 meter high, in sandy coastal regions on the south side of the islands. (Stahl, 2: 108.)

Melon. See *Cucumis melo*.

Melon de costa. See *Melocactus communis*.

Melothria fluminensis. PEPINILLO.

Family Cucurbitaceae; a climber 5 or 6 feet (1.5 to 1.9 meters) long, found in hedges and waste places. Fruit eaten, pickled when green; also good when fully ripe. (Stahl, 4: 177.)

Mentha piperita. PEPPERMINT.

A plant named by Briquet as variety *citrata* of this species was collected in the borders of primeval forests near Adjuntas.

Mentrasto. See *Ageratum conyzoides*.

Mentzelia aspera. PEGADORA.

Family Loasaceae; a hairy herb with triangular dentate leaves and deep buff flowers. Our specimens were obtained to the southward of Coamo, not far from the military road.

Mesosphaerum atrorubens. MARRUBIO OSCURO.

Family Labiatae; an herbaceous annual, 40 centimeters high, in dry pastures. (Stahl, 4: 194, as *Hyptis atrorubens*.)

Mesosphaerum capitatum. MARRUBIO-BOTON.

An herbaceous annual weed of the mint family. At Cataño it was called "blero." (Stahl, 4: 192, as *Hyptis capitata*.)

Mesosphaerum lantanaefolium. MARRUBIO BLANCO.

An herbaceous reclining annual. (Stahl, 6: 193, as *Hyptis lantanaefolia*.)

Mesosphaerum polystachyum. FALSO MARRUBIO.

An herbaceous climber, 1 to 2 meters high. (Stahl, 6: 195, as *Hyptis polystachya*.)

Mesosphaerum spicigerum. MARRUBIO.

Reported by Bello.

Mesosphaerum suaveolens. MARRUBIO.

An herbaceous climber, 1 meter high; contains an oil with a strong odor. Grows in sandy soil. (Stahl, 6: 196, as *Hyptis suaveolens*.)

Mesquit. See *Prosopis juliflora*.

Metastelma.

A genus of climbing milkweeds represented in Porto Rico by numerous species. (Urban, Symb. 1: 245, a monograph.)

Miconia. See *Tamonea*.

Micropholis chrysophylloides.

Family Sapotaceae; reported from near Adjuntas.

Micropholis garciniaefolia.

From Sierra de Luquillo.

Micropholis portoricensis curvata.

From near Adjuntas.

Micropholis portoricensis mesnaefolia.

From near Peñuelas.

Micropholis urbani.

From Sierra de Naguabo.

Mignonette tree. See *Lawsonia inermis*.

Mikania. See *Willughbaeya*.

Millet. See *Panicum*.

Millo. See *Andropogon*.

The seeds of a millet were on sale in small quantities in the market of Ponce, and are said to be used in the preparation of a sort of flour and in sweets.

Mimosa ceratonia. ZARZA.

Family Mimosaceae; a vine-like shrub, 2 to 5 meters high, found in waste places. (Stahl, 3: 143.)

Mimosa pudica. MORIVIVI. SENSITIVE PLANT.

A branching perennial, 2 to 5 feet (0.6 to 1.3 meters) high. The sensitive plant of greenhouses; an extremely common weed along roadsides and in waste places generally. (Stahl, 3: 142.)

Mimusops globosa. SAPOTE DE COSTA.

Family Sapotaceae; a shrub, 3 to 4 meters high, growing in shaded situations along the sea coast. (Stahl, 6: 54.)

Mimusops pleeana.

From Manati.

Mimusops riedliana.

From Yabucoa.

Mirabilis jalapa. FOUR O'CLOCK. MARAVILLA.

The well-known garden flower said to be called "siciliana" in Porto Rico, where it is cultivated and has also become naturalized as a weed in open places. It is a native of Mexico. (Urban, Add. 3: 304.)

Mistletoe.

The true mistletoe, *Viscum album* of Europe, does not occur in West Indies, but there are numerous related parasites belonging to the family Loranthaceae and now referred to the genera *Dendropremon*, *Dendrophthora*, *Phoradendron*, and others.

Moca. See *Andira inermis*.

Moca acatera.

A tree from the northeastern part of the island: height, 25 to 30 feet (7 to 9 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood yellowish, hard; specific gravity, 0.993; used for cabinetwork. (Exp. 1857.)



CUNDEAMOR (MOMORDICA CHARANTIA).

Moca amarilla.

A tree from the northeastern part of the island; height, 25 to 30 feet (7 to 9 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood red, hard; specific gravity, 0.904; used for cabinetwork. (Exp. 1857.)

Moca negra.

A tree from the eastern part of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood dark, hard; specific gravity, 0.811; used for cabinetwork. (Exp. 1857.)

Molinillo. See *Leonotis nepetaefolia*.

Momordica.

A genus of Cucurbitaceae. The species are graceful and delicate climbing vines, bearing rather small edible fruits.

Momordica charantia. CUNDEAMOR.

PLATE XLVI.

This species is supposed to be the most common and generally distributed, perhaps because confused with *M. zeylanica*, which in Porto Rico, at least, is much more abundant. The present species was not noticed by us, but has been collected by Sintenis at Fajardo. It differs from *M. zeylanica* in the much larger and longer fruit, the surface of which is irregularly covered with rounded warts and tubercles. The seeds are also much larger, and the leaves have a very different cut, as may be seen by reference to the figure.

Both species are sometimes called "balsam apple."

Momordica zeylanica. CUNDEAMOR.

This species is one of the commonest and most widely distributed plants in Porto Rico, climbing over fences and other vegetation in open places. Many botanists have confused it with *M. charantia*, but it may be recognized readily by the different leaf pattern, the small oval fruits with rows of sharp spines, and the small seeds.

Monkey apple. See *Anona palustris*.

Monodora myristica. CALABASH NUTMEG.

Also called "American nutmeg" and "Jamaica nutmeg." An anonaceous tree, supposed to have been introduced from West Africa. Its occurrence in Porto Rico seems doubtful, a species of *Nectandra* having, in some cases at least, been mistaken for it.

Grosourdy (2: 382) reports it from all parts of the island, and gives the height of the tree as reaching 25 or 30 feet (8 or 9 meters), with a diameter of 10 inches (25 centimeters). The wood, he says, is yellow, rather strong, but without application. His identification was perhaps based on the native name "nuez moscada del pais," in which case the description probably refers to a species of *Nectandra*.

Mora. See *Chlorophora tinctoria*.

Hill & Sudworth (p. 27) describe a wood of this name as follows:

Color, bright orange brown, probably darkening with age and exposure. Marked on the radial section with parallel light and dark streaks. Large ducts arranged in wavy, irregular lines; lines also irregularly broken into crescent-shaped groups. Medullary rays distinct, rather coarse. Radically cut and polished surface satiny. Similar in general appearance to osage orange. Largely used for fellies.

Hansard gives the specific gravity as 0.86. Perhaps the same as "moral."

Mora amarilla.

A tree from the eastern part of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood, yellowish, hard; specific gravity, 0.807; used for cabinetmaking. (Exp. 1857.)

Moradilla azul. See *Salvia occidentalis*.

Moradilla encarnada. See *Salvia coccinea*.

Moral. See *Cordia macrophylla*.

A tree from all parts of the island; height, 45 to 50 feet (13 to 15 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood light, soft; specific gravity, 0.611; not used. (Exp. 1857.) (Grosourdy, 2: 402.)

Captain Hansard thought this might be the mulberry, *Morus nigra*, the leaf being similar, though the fruit is different. Specific gravity, 0.57.

At Coamo Springs another species of *Cordia* is known under this name. It is quite different from *C. macrophylla* and seems to have affinity with *C. collococca*.

Moralon. See *Coccolobis grandifolia*.

Moriche. See *Mauritia flexuosa*.

Moringa moringa. HORSE-RADISH TREE. BEN.

Family Moringaceae; a tree 3 to 4 meters high, native in India; reported as introduced and escaped from cultivation in Cuba and probably also in Porto Rico. At Cataño and also at Isabella it was called "angela" or angel tree. It yields the so-called oil of ben, used for fine lubricating purposes. The root has the taste and odor of horse-radish. Sintenis collected a specimen distributed as *M. oleifera* in a garden at Mayaguez. (Stahl, 2: 43.)

A specimen 20 feet (6 meters) high, with a trunk 8 or 10 inches (20 or 25 centimeters) in diameter, was seen between Yauco and Guayanilla.

Mori-vivi. See *Mimosa pudica*.

Other fine-leaved leguminous plants, such as *Aeschynomene americana* are also sometimes called mori-vivi or mori-vivi bobo.

Morongia distachya. CUERNECILLO.

A much-branched, trailing, spiny shrub reported by Stahl from the south side of the island. (Stahl, 3: 136, as *Schrankia distachya*.)

Morongia leptoclada. SALCILLA.

Collected at Coamo Springs (No. 726). A less spiny species than *distachya*, according to the specimens in the National Herbarium.

Morus. MULBERRY.

No mulberry trees were noticed in Porto Rico, but in Cuba experiments are said to have shown that both the mulberry and the silkworm thrive extremely well.

Mostacilla del mar. See *Cakile maritima*.

Mostaza. See *Brassica juncea*.

Motillo. See *Sloanea berteriana*.

Mouriria.

Family Melastomaceae; a species thought to be related to *Mouriria spathulata* was collected at Hatillo by Sintenis. The lower side of the leaves is yellow.

Mouriria domingensis. MURTA.

A cultivated shrub, 3 to 5 meters high. The agreeable fruits are eaten raw and are used to color liquors. (Stahl, 4: 91.)

Mucuna altissima.

Family Viciaceae; a leguminous vine, related to the velvet bean; known from Yabucoa; also from the vicinity of Ponce. (Sintenis.)

Mucuna pruriens. COWHAGE. PICA-PICA.

The pods of this leguminous plant, which is a close relative to the velvet bean, are covered with a dense growth of fine, rather stiff, poisonous hairs, said to be extremely irritating when brought in contact with the more sensitive parts of the skin. (Stahl, 3: 84.)

Mucuna urens. OJO DE BUEY.

A woody, biennial vine, growing in waste places. Bello gives "matos" as the common name of this species. (Stahl, 3: 85.)

Mucuna utilis. VELVET BEAN.

The velvet bean has been tried at a few points in Porto Rico, and vigorous growth is reported. In all probability it will be found one of many leguminous plants of value for restoring exhausted soils.

Mugwort. See *Parthenium hysterophorus*.

Mulberry. See *Morus*.

Multa.

Grosourdy (2: 402) refers this to the genus *Eugenia*, and describes it as a wild fruit tree, 15 to 20 feet (4 to 6 meters) high, with rather hard, resistant wood, the only use of which is in making charcoal.

A sample of fruit called "multas" in the San Juan market was not unlike those of the genus *Eugenia*. The pulp, though very thin, is of a pleasant taste, not unlike that of gooseberries.

The name "murta," given by Urban as the common name of *Eugenia floribunda* and *E. cordata sintensisii* and by Stahl for *Mouriria domingensis*, may in one or more cases be another spelling of "multa," misunderstood for the Spanish word for "myrtle."

The word "multa" is given in Velásquez's Spanish dictionary as the name of a fruit tree in Porto Rico.

According to Captain Hansard this name is applied to a wood with a specific gravity of 1.06, found in Vieques, and used for coloring rum.

Muñeca. See *Cordia borinquensis* and *Gilibertia arborea*.

A tree from the eastern part of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood, ash-colored, soft; specific gravity, 0.951; used for lumber. (Exp. 1857.) (Grosourdy, 2: 403.)

A soft, light, inferior wood, found occasionally in the hills; specific gravity, 0.59. (Hansard.)

Muñeca de costas.

A tree from the northern part of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters.) Wood, yellowish, rather hard; specific gravity, 0.923. (Exp. 1857.) Used for fuel. (Grosourdy, 2: 403.)

Murraya exotica. CAFÉ DE LA INDIA.

Family Rutaceae; a shrub, 2 to 3 meters high, native in the East Indies, and naturalized in the West Indies and on the continent. It is cultivated in gardens for its abundant and fragrant flowers and for its agreeable, small, reddish tart fruits, which much resemble coffee berries, although in reality the plant is closely related to the orange. (Stahl, 2: 135.) Near Carolina this plant was called "dama de noche," a name supposed to belong to *Cestrum nocturnum*.

The rather straight trunk is sometimes as thick as a man's thigh, but is ordinarily smaller. It furnishes a wood very similar to the boxwood of Europe, which it might very well replace. The wood is durable, yellowish white with brownish heart, fine-grained and dense, strong and solid. Very good for turned articles and susceptible of a polish. Specific gravity, 0.979. (Grosourdy, 2: 366.)

Murta. See *Eugenia floribunda*, *E. cordata*, *E. sintensisii*, and *Mouriria domingensis*.

Musa. BANANA. GUINEO.

The culture of bananas for export is one of the industries which could be added to the resources of Porto Rico with every hope of success. Although furnishing, perhaps, the most important food crop of the island the banana is seldom treated to any approximation of regular cultivation. To a large extent, the growing of bananas has gone hand in hand with the planting of coffee, for the shading of which large numbers of bananas are required. Extensive fields, which appear to

be devoted to bananas, are usually found, on careful inspection, to conceal spindle coffee plants. So many bananas are produced in this incidental manner that there has been little necessity for special cultivation to supply the local markets. Neither has any sustained attempt been made to maintain the quality or productiveness of any of the varieties, so that a general deterioration has probably taken place, instead of the rather careful though apparently desultory selection to which bananas are subjected in many tropical countries. As the edible bananas are seedless, all propagation is by cuttings of buds or branches which arise from root stocks. It is thus usually possible for natives to select good stock of any desired sort when setting out new plants, and therefore in most countries the bearing qualities, at least, are maintained. But in Porto Rico there has been comparatively little incentive to care in special planting. Our failure to find any marketable bananas seemed at first very strange, until the local conditions were understood. Moreover, the deficiency is further explainable by the fact that the banana is extremely responsive to good treatment, and really first-class fruit can be produced only under favorable conditions, requiring little attention to the enrichment of the soil. As such care is seldom forthcoming in Porto Rico it is doubtful whether high-grade stock of any of the local varieties could now be secured in the island.

Choice of varieties.—Accordingly, if the commercial planting of bananas is to be undertaken in Porto Rico it will be necessary either to make a careful choice of stock, which owing to special care has retained its vigor and productiveness, or to make new importations from other parts of the West Indies or from Central America. The Martinique or Jamaica banana, the variety almost exclusively imported into the United States, is nowhere particularly esteemed in the Tropics except for export purposes, and comparatively few are raised in Porto Rico, so that it is extremely doubtful whether plantations of any size could be stocked without some loss of time, unavoidable on account of the rather slow propagation of the superior varieties of the banana. Of course it would be useless to plant an indiscriminate mixture of varieties, the produce of which would be quite unmarketable abroad. At the same time it is proper to consider the question whether in view of the comparatively ready access to the American market it will not be wise to plant in export quantities some one of the numerous varieties superior to that now known in the trade. The value of the common sort lies in its shipping qualities, which result from the fact that it shares some of the attributes of the plantain in being of rather coarse texture, with a tough skin and a large quantity of mealy outer coating of the flesh, which gives the dry sensation and, when not thoroughly ripened, the astringent taste. Really fine varieties are so superior in flavor and texture that once regularly introduced into the trade they would certainly secure the popular preference and command special prices. It is true that the more delicate texture of most of the high-grade varieties would probably render it necessary to use somewhat greater care in shipping. But methods of packing and ventilation are now so far improved that some of the better varieties could undoubtedly be brought to our markets in time to reach the consumer in good condition. One of the requisites of the success of such an experiment should, however, be kept in mind. No matter how excellent the fruit may be, the public will not have a chance to assert a preference until it can be sent in quantities large enough to give it a place in the regular fruit trade. Many failures in opening new lines of agricultural activity have resulted from the fact that trial shipments were so small that the merchants would not take the trouble necessary to properly introduce an unknown product to the public.

As one instance of a superior banana which might be grown in Porto Rico may be mentioned a variety of the dwarf banana (*Musa cavendishii*), the same species which is cultivated in the Canary Islands for the English market. The Canary bananas are not, however, of particular excellence, having, if anything, even more

of the objectionable taste and mealy texture. The better variety is cultivated in Liberia, and even when ripe has a greenish rather than a yellow color. The skin is rather thin, there is no astringency, and the compact and yet delicate creamy texture and delicious flavor render the inferiority of our ordinary imported fruit distastefully evident to all who have the opportunity of comparison. The dwarf banana tree reaches a height of only 6 or 8 feet (2 or 2.5 meters), and the enormous bunches reach nearly to the ground and weigh in the neighborhood of 100 pounds. The individual bananas are of medium size and rather slender in comparison with those known in our markets.

Botanical knowledge of the banana.—It might naturally be expected that the botany of so important a plant as the banana would be in an advanced state of completeness; but quite the contrary is true, owing to the fact that economic plants are generally neglected by botanical students and collectors, especially when, as in the present instance, specimens are very difficult to preserve. No part of the banana plant is adapted to ordinary herbarium treatment, neither the enormous leaves nor the succulent flowers and fruits.

It is accordingly quite impossible to tell from books whether the superior African variety mentioned above is known in other parts of the world. In Porto Rico the names "congos" and "guineos" are applied to several varieties of banana, and there are also traditions that, like the cattle and Guinea grass, they were introduced from the Dark Continent. We had no personal opportunities of testing the variety called "enano," or "dwarf," which may prove to be the same as the African sort, though it is much more likely to have been derived from Canary Island stock. Bananas are not native in America, but several wild species with seeds are known in British India and the Malay region, where there are also large numbers of cultivated varieties, very few of which have been introduced into any part of tropical America. It would seem that the country which makes by far the largest banana trade in the world should enjoy the advantage of the best that could be afforded in the way of this now popular fruit, and that Porto Rico might realize brilliant results from the culture of high-grade varieties.

Porto Rican varieties.—Hundreds of varieties are known to exist in the East Indies, particularly in the Malay Archipelago and in the Philippines, but these are so confused and so poorly described that it is impossible to tell which of them has been introduced into Porto Rico, and even between the different West Indian islands the identity of varieties can not be asserted with confidence, even when they bear the same name. As an illustration of the confusion which may exist, even on the same island, our notes on Porto Rican varieties may serve a salutary purpose.

In addition to what was obtained from other sources, we had the advantage of the information gathered by two gentlemen who had given the subject considerable attention—Captain Hansard, of San Juan, formerly resident in the Luquillo region, and Mr. Mead, of Ponce and Peñuelas. The following alphabetical list of names, while probably incomplete and perhaps erroneous, may serve as a beginning in the collection of more complete information:

PORTO RICAN VARIETIES OF BANANAS.

Amarillo.—A name applied to the large plantain (plátano), especially when sliced and fried. (Ponce.)

Bulico.—See *Horse plantain*. (Ponce.)

Cambur.—A name given by Hill; not known to Captain Hansard. In Venezuela it seems to be a generic term for the bananas, as distinguished from the plantains.

Chinese.—See *Dwarf*.

Colorado.—See *Morado colorado*.

Commissario.—A name for plantains; thought to be the same as *mafafó*, or horse plantain.

Datil.—A name used about San Juan for a very small banana (fig. 77). Perhaps the same as niños. Although called manzanas by some of the hucksters, they had none of the apple flavor and were otherwise inferior.

Dominico.—Captain Hansard states that this name is applied in Porto Rico to the gigantes or Jamaica bananas, but the first use of it by Acosta had reference to a small, white, delicate species from Santo Domingo. In Venezuela the name "plátano dominico" is used for "the royal, or small-fruited, plantain." This is said to be "very similar to the small plantain in appearance and habit. The fruit, however, is smaller and the plant somewhat hardier; that is, it bears better the cold of the mountains."

Dwarf.—In the English-speaking tropics this name is applied to *Musa cavendishii*, also known as the Chinese banana. The name "enano," in use for a banana variety in Porto Rico, probably has reference to the above species, which is that cultivated in the Canary Islands.

Enano.—This name means "dwarf," and has reference to the tree rather than the fruit. It may prove to be *Musa cavendishii*, like the "platanos congos" at Ponce.

Fig.—A name used in the British Island for the Martinique or Jamaica banana, the variety commonly used in the United States.

Fig la rose.—Said to be the name used in Dominica for the Martinique or Jamaica banana.

Figue.—A name of the above in the French Islands.

Fotoko.—A variety said by Captain Hansard to be midway between a plantain and a banana, but not fit to eat uncooked. The name seems to be one of African origin.

Giant.—See *Gigante*.

Gigante.—According to Mr. Mead this is the same as the Martinique banana. The fruit grows very large, sometimes 10 inches (25 centimeters) long, but is slender and very angular. Eaten mostly cooked, especially with wine, and used for making "dulce" or preserves. In the San Juan market this name was used for a much smaller fruit, apparently derived from a form of *Musa cavendishii*.

Gros Michel.—See *Martinico*.

Guadeloupe.—See *Martinico*.

Guineo amarillo.—A name used in the San Juan market for a medium-sized banana which some of the dealers called "gigantes." The fruits seemed rather to resemble those of *Musa cavendishii*.

Guineo de manzana.—See *Manzana*.

Guineo de piña.—The smallest variety known to Captain Hansard, and not the same as datiles, thin-skinned; stem bright green; leaves broad; not so tall as the gigantes and less liable to have the leaves stripped by the wind. See also "Guineo de rosa."

Guineo de rosa.—A name used about Ponce for a variety also called guineos de piña. These were described as short, yellow, rather thick, and slightly angular. The taste is sweet and not astringent.

Guineo de tierra.—A very inferior sort of plantain. According to Captain Hansard vinegar is not made from the ordinary plantain as indicated by Hill, but from the present variety which is grown for that especial purpose.

Guineo indio.—A red banana grown about Ponce. The golden yellow flesh is said to be of very fine quality, much preferred by many Americans. Bunches over 3 feet (1 meter) long are sometimes secured under favorable conditions.

Higo.—See *Martinico*.

Horse plantain.—An English name for the mafafó or short plantain.

Horton.—According to Hill and Hansard this is the name for the plantain, presumably the long-fruited type. In Venezuela the common plantain is said to be called "plátano arton."

Jamaica.—A name used in the American trade for the large yellow bananas now practically the only variety brought to our markets. Jamaica is the center of the West Indian banana-growing industry, but the present variety is not confined to that island, and is not called "Jamaica" except by us. In the different islands it has many other names such as fig, figue, gigante, gros Michel, Guadeloupe, Martinique, etc.

Mafafó.—Apparently the most common Porto Rican name for the short angular plantain, sometimes called the horse-plantain by the English-speaking people. The mafafos are smaller, shorter, and thicker than the genuine plantains and the cross section is very angular. Quality and texture differ from the plátanos in being more mealy; the skin is extremely thick. This variety is peeled and toasted in the charcoal ashes and is then called "native bread;" it is frequently sold in the market. There are several other native names, such as bulicos, commissarios, and mata hambre, or "hunger-killer."

Manis.—This name seems to be unknown at San Juan, but Captain Hansard became acquainted with the variety in the vicinity of Luquillo, and declares it to be the best in the island, describing it as more substantial and yet with a more delicate flavor. There are at least three slightly different varieties. The individual fruits are small, only 4 or 5 inches in length, like the "datile" and "piña." A medium-sized tree with rather few leaves; the stem becomes superficially red with age. There is another similar variety, coarser and larger, and not tasting so well. Perhaps this is the "niño," a name unknown to Captain Hansard.

Manzano.—Specimens of this variety were kindly sent to us by Mr. Graham, of Ponce, under the name guineos de manzana. These were of a very bright yellow color, with a very thin and delicate skin, fine and soft texture, and very pleasant taste, with a distinct flavor of apple. An over-ripe apple is perhaps suggested, but the banana texture and flavor overcome any objection of that kind, and there is no mealiness or unpleasant aftertaste. This is, in short, an exceedingly fine fruit, but one which is looked upon as a delicacy rather than a regular product. In the unripe state the manzana is said to be particularly astringent, but this quality disappears with maturity. The tree is somewhat smaller than that of the niño and other varieties except the dwarf. The bunches of the manzana are generally small, seldom reaching 100 fruits.

Martinico.—According to Mr. Mead this name is used on the south side of the island for the variety more commonly known as "gigantes," the gros Michel or Jamaica banana.

Mata hambre.—See *Mafafo*.

Morado.—See *Morado colorado*.

Morado blanco.—Like the morado colorado in taste, form, texture, etc., but green, not becoming red, but yellow when ripe.

Morado colorado.—A very large, red variety. Captain Hansard has seen bunches of 100, of which the smallest fruit was 8½ inches (21 centimeters) in circumference, the largest over 9 inches (22 centimeters). The tree is the largest of any of the varieties and is on this account desirable for shade, but the fruit is not liked by the people of Porto Rico, hence there is no local market.

Niño.—Yellow, short, slender, slightly angular; taste very sweet, without acid or unpleasantness; trees about same size as the large varieties.

Piña.—See *Guineo de piña*.

Plantain.—In English this name is used either as a general term for all the banana-like fruits or, more commonly, it is restricted to the coarser varieties which are inedible without cooking.

Plátano.—Used alone, this name seems to refer in Porto Rico to the long plantain and not to the short plantain, or "mafafo."

The plátano is eaten only in the cooked condition, being generally sliced and fried, although often boiled or combined with other materials. The fruits are very often large, thick-skinned, and angular. They are generally cooked while still green, but if allowed to mature they become yellow and then black.

Plátano Congo.—Considered by Mr. Mead the finest of the Porto Rican bananas. Very slender and well flavored, eaten either raw or cooked; grows on a small tree and in large bunches. From the description this seems to be close to the African dwarf banana; but the fruit is said to become very yellow and the leaves to be entirely green, including the stems. These discrepancies seem to indicate at least a varietal difference, though botanical varieties of *Musa cavendishii* seem not to have been distinguished.

Plátano largo.—See *Plátano*.

St. Michael.—Probably an error for gros Michel.

Musa textilis. MANILA HEMP. ABACÁ.

The Philippine Islands are the home of this important fiber plant and the industry based on it. Various attempts have been made to introduce the culture of Manila hemp into other regions, thus far without success, probably because the process of extraction, though very simple, requires experience and skill.

Muskwood.

In the British West Indies this name is said to be used for *Mosehoerylum swartzii* and *Guarea swartzii*.

Myginda latifolia.

A branching celastraceous shrub, 3 to 4 feet (1 to 1.3 meters) high. Reported from Salinas de Cabo Rojo.

Myginda pallens. MARAVEDI.

A low shrub. (Stahl, 4: 22.)

Myginda rhacoma. MARAVEDI.

A low shrub from along the seacoast; 1 to 2 meters high. Reported from Rincon. (Stahl, 4: 21.)

Myginda rotundata.

Known from Guanica (Sintenis). This species is considered by Urban to be a monstrosity. It has clusters of very fine branches covered with small brown scales, apparently hexenbesens.

Myrcia coriacea. HOJA MENUDA.

Family Myrtaceae; a shrub 3 meters high; in waste places in dry regions. (Stahl, 4: 74.)

Myrcia deflexa. CIENEGUILLO.

An indigenous tree, sometimes exceeding 50 feet (15 meters), distributed throughout the mountainous districts, furnishing hard wood, also called "guayavacon." (Urban, Add. 2: 105; Stahl, 4: 76.)

Myrcia divaricata. See *Myrcia leptoclada*.**Myrcia ferruginea.** CIENEGUILLO.

A shrub 3 to 5 meters high, found in waste places. The red, hard wood is used to make canes. (Stahl, 4: 77.)

Myrcia leptoclada. GUAYAVACON.

An indigenous tree said to reach a height of 30 feet (9 meters). Known from Luquillo, Yabucoa, Maricao, and Añasco.

Myrcia pagani. AUSU.

An indigenous tree known only from sterile branches collected in forests in the Sierra de Lares.

Myrcia paniculata. HOJA MENUDA.

An indigenous tree reaching height of over 50 feet (15 meters), and furnishing a durable, flexible wood. Reported from Bayamon, Cayey, Aibonito, Rincon, Manati.

Myrcia sintenisii.

Has large elliptical leaves 7 inches long (17 centimeters); from Lares.

Myrcia sororia. See *Myrcia splendens*.**Myrcia splendens.** RAMA MENUDA.

Also called "hoja menuda." A shrub or small tree, indigenous in Porto Rico and found in forests and thickets in numerous localities from Luquillo to Mayaguez. *M. berberis* is known in Porto Rico only from the Luquillo region. (Urban, Add. 2: 101.)

Myrica cerifera. BAYBERRY. ARRAYAN.

The common bayberry of the Eastern States; extends to Cuba, Porto Rico, Santo Domingo, and Guadeloupe. In Porto Rico it has been collected in mountain forests near Utuado, Las Marias, and Mayaguez, as well as in humid meadows and sandy beaches near Bayamon.

The following account written about 1831 indicates that this tree must have been formerly of much greater importance than at present:

The "cerero" tree grows naturally in regions that are not extremely cold, and in humid temperate regions, particularly the plains and shores along the rivers; also in all regions that produce wheat, whether high or low. Its size is that of an ordinary olive tree or sometimes smaller. The fruit resembles the seeds of the pepper of Castile. Experience has shown that each large tree well fruited yields two "almudes." "

Smaller trees produce one almud and the poorer one-quarter. Annually they gather two crops; the first is more abundant than the second and lasts from March until May.

The season to collect the fruit is when it is white or ashy. Well dried in boxes

"An "almud" is a measure given as either about 8 or 50 pounds.

it can be kept a year without spoiling, and from each half almud you obtain a pound of wax more or less in the manner described below.

The collected fruit is broken up and spread on hides or boards in a well-ventilated place out of the sun. It is turned gently each day for thirty days, when they proceed to extract the wax; but the wax may be extracted as soon as the fruit is gathered, in which case the wax is too green in color, while after thirty days it is whiter. It is kept from the sun because it melts part of the fruit and wastes a part of its substance, but it has been found that by exposure to the sun for three days they get whiter wax.

Place on the fire a pot glazed on the interior, with a quantity of water, and when boiling introduce as much fruit as the pot will hold, and gather with a spoon the fat that rises to the surface of the water and allow it to cool until it coagulates, after which it is melted and filtered through a haircloth, with the result that the wax is clean and ready for use. The fruit remains in the boiling water until it has given up all its substance, which can be known when no oil rises to the surface, and then with fresh water and fruit the process is repeated.

This wax is brittle and no method for making it flexible has been discovered, so that it is worked with difficulty and the candles are liable to break. The wicks are fastened to strong sticks and bathed with the wax and guarded from the wind which will make them crooked, and while they are slime the bottom of each candle is pulled after each coating of wax to prevent their becoming crooked. (Cordova, *Memorias de Puerto Rico*, 1: 240. 1831-1833)

Myristica fragrans. NUTMEG.

The true nutmeg is not known to have been introduced into Porto Rico, but the Spanish name for it "nuez moscada" is in common use for the seeds of a species of *Nectandra*, which serve as a substitute.

See note under nutmeg.

Myrodia turbinata. See *Quararibea turbinata*.

Myrospermum frutescens.

Family Viciaceae; a tree or shrub reported from near Utuado.

Myroxylon buxifolium. ROSETA.

A bixaceous shrub or tree, reaching a height of 8 to 10 meters; known from the vicinity of Peñuelas and Mayaguez.

Myroxylon pachyphyllum. See *Xylosma pachyphyllum* and note under *X. schwaneckeanum*.

Myroxylon schwaneckeanum. PALO DE CANDELA. See *Xylosma schwaneckeanum*.

Also called "palo colorado." A tree related to the "achiote" (*Bixa orellana*). It is described as from 5 to 8 meters in height, native in the Luquillo Mountains. (Urban, *Add.* 1: 12.)

Myrsine coriacea. ARRAYAN.

A myrsinaceous shrub, 3 meters high found in mountains and waste places; collected by Sintenis at Adjuntas. (Stahl, 6: 35.)

Myrsine floribunda. ARRAYAN.

A shrub 3 meters high, found in the mountains and waste places. (Stahl, 6: 36.)

Myrsine guianensis.

A shrub reported from near Manati.

Myrtus acris. See *Anomis euryphyllata*.

Naguacatillo.

A tree about 60 feet (18 meters) in height, with light-green or greenish white wood, light and soft, but resistant. It is used in making boxes and tampions and for sheathing houses. Its fruit is used to feed hogs. (Grosourdy, 2: 357.) Referred to *Nectandra leucantha* and *Laurus borbonia*, the former a Brazilian species, the latter a synonym of *Persea borbonia*.

Naranja. See *Citrus aurantium* and *C. bigaradia*.

The Spanish general term for the orange. In Porto Rico, however, it is applied

only to the wild, sour orange, the sweet orange being called "china," or "naranja de china," or "china dulce."

Naranja china. See *Citrus aurantium*.

Naranja de china. See *Citrus aurantium*.

Naranjada.

A drink made from oranges. "orangeade."

Naranjillo.

A tree from all parts of the island; height, 25 to 30 feet (7 to 9 meters); diameter, 8 to 10 inches (20 to 25 centimeters). Wood, white, hard; specific gravity, 0.743; not used. (Exp. 1857.)

Naranjo. See *Naranja*.

The orange tree in Spanish is "naranjo" or "naranjero," while the fruit is "naranja."

A tree from all parts of the island; height, 20 to 25 feet (6 or 7 meters); diameter, 9 to 10 inches (21 to 25 centimeters). Wood white, hard; specific gravity, 0.784; fruit tree. (Exp. 1857.)

Captain Hansard gives a specific gravity of 0.48 for the wood of the bitter orange, but this seems too light for so hard a wood.

Naseberry tree. See *Sapota achras*.

Nasturtium officinale. See *Rorippa nasturtium*.

Nectandra. ENEMOSCO.

Family Lauraceae. In Porto Rico the seeds of a species of *Nectandra* have been confused with those of *Monodora myristica*, the "calabash nutmeg," and also with the true nutmeg, *Myristica moschata*.

See note under Nutmeg.

Nectandra leucantha.

See note under Nagnacatillo.

Neea buxifolia.

A nyctaginaceous shrub, reported from Fajardo.

Negra lora. MABA INCONSTANS.

Captain Hansard gives for this wood a specific gravity of 0.91.

A wild tree, 30 feet (9 meters) high, with a trunk 10 to 12 inches (25 to 30 centimeters) in diameter. The wood is flesh-colored, coarse, porous, and breaks with a vertical fracture. Its most common use is in building country houses and for fences. (Grosourdy, 2: 403.)

Negraz losa.

A tree from the interior of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood red, hard; specific gravity, 0.897; used in building houses. (Exp. 1857.) Probably the same as "negra lora."

Nephelium.

A genus of Sapindaceae, belonging to the Indian and Malayan region, and generally treated as including species which yield several important fruits, such as the "litchi," the "rambotan," and the "longan." *Litchi chinensis* has recently been recognized as generically distinct from *Nephelium*.

Nephelium litchi. See *Litchi chinensis*.

Nepsera aquatica. CAMACEY DE CIENEGA.

Family Melastomaceae; a small biennial shrub growing in clay or sandy soil subject to inundation. Of general distribution in tropical America. The generic name given by Stahl is *Nespera*, doubtless a typographical error. (Stahl, 4: 117.)

At Cataño this species is called "altea" and supposed to possess medicinal properties.

Neptunia plena. DESMANTO AMARILLO.

Family Mimosaceae; a slender herb, in sandy inundated places. (Stahl, 3: 141 as *Desmanthus plenus*.)

Nerium oleander. OLEANDER. ADELFA.

Family Apocynaceae. The common oleander is extensively planted in Porto Rico, as in other tropical countries, sometimes in the form of hedges. In some localities it has escaped from cultivation and grows apparently wild.

Nespera aquatica. See *Nepsera aquatica*.**Neurolaena lobata.** SEPI.

An annual herbaceous composite growing as a weed in waste places. (Stahl, 5: 128, as *Neurolema lobata*.)

Neurolema lobata. See *Neurolaena lobata*.

This is probably a typographical error on the part of Stahl.

Niaragato. See *Fagara spinifer* and *Zanthoxylum fragodes*.**Nicotiana tabacum.** TOBACCO.

Although only a small area is devoted to the culture of tobacco in Porto Rico, this crop ranks third among the exports of the island. The plant may be grown, indeed, in nearly all parts of the island, but there is only a limited area of the soil supposed to be adapted to the production of a marketable quality of leaf. The best tobacco, by general consent, is raised in the La Plata Valley above Cayey. Like nearly all the interior of the island the configuration of the country is extremely hilly, although, owing to the more friable nature of the soil, the slopes are not as steep as in many of the coffee-growing districts, and plowing can often be practiced. The underlying rock crumbles readily into small angular pieces, which, mingled with the finer superficial material, makes a loose, easily workable, deep-brownish soil. This is left in shallow furrows and ridges running along the slope. The rows are spaced at from 18 to 24 inches (45 to 60 centimeters), and the plants are set very close, only from 4 to 8 inches (10 to 20 centimeters) apart, there being no evidence of any serious attempt at regularity of distance.

The town of Cayey is the recognized center of the tobacco industry of the island, although the largest cigarette factories are located at Ponce and San Juan and many cigars are made in those and other cities and towns. In fact, cigar making is one of the largest of the domestic industries. All the men and boys smoke, of course, and many of the women are said to do so, although only those of the lower classes are seen indulging themselves in public. The home consumption of cigarettes must be enormous, since the Porto Rican, whether at work or at leisure, is seldom long without one, even though the cigar may be only an occasional indulgence. The Porto Rican cigarette is not, however, the medicated nuisance prevalent in the United States, but the general use of tobacco by the growing boys probably has its effect in increasing the debility of the population.

A peculiar type of black plug tobacco is also offered for sale in the market, in the form of large rolls variously and ingeniously coiled. Although fruits and vegetables are sold by weight in Porto Rico, this tobacco is retailed by linear measure.

No such careful methods of cultivation or selection have been practiced in Porto Rico as in Cuba, and the possibilities of the island will not be adequately known until careful experiments have been made under test conditions. It has been demonstrated by the experiments of Dr. Loew that the processes of curing are all important in the determination of the quality of tobacco. It has been stated, with what warrant is not known, that the Porto Rican growers are not acquainted with the Cuban methods and do not practice a uniform system of curing. If this be the case, it is not impossible that the introduction of the popular processes may render the Porto Rican crop of the very first quality.

In Cuba and other regions furnishing high-grade tobacco the crop is not only assorted by localities, but the leaves even from different parts of the same plant are separated into several classes, some of superlative quality. In Porto Rico the manufacturers uniformly declare that they handle only one grade of tobacco, and that the differences in the prices of their various classes of cigars represent merely the varying cost of labor and packing.

Nigua. See *Tournefortia hirsutissima*.

Several other species of *Tournefortia* are also known under this name, though Stahl proposes distinctive terms for the different species. A root called "nigua" was on sale in the market of Ponce as an ingredient of a refreshing drink.

Nigua de playa. See *Tournefortia guaphalodes*.

Nigua-enredadera. See *Tournefortia colubilis*.

Nigua fetida. See *Tournefortia foetidissima*.

Nigua-hoja-laurel. See *Tournefortia laurifolia*.

Nigua peluda. See *Tournefortia hirsutissima*.

Niguita. See *Tournefortia laccigata*.

Niño de cota. See *Haemocharis portoricensis*.

Nispero. See *Achras sapota*.

This is the Spanish name for the Japanese medlar or loquat (*Eriobotrya japonica*), but in Porto Rico it is universally applied to *Achras sapota*, elsewhere called "sapote" or "sapidilla." Captain Hansard gives a specific gravity of 1.02 for the "sapidilla" or "nispero" wood.

Nispero cimarron. See *Symplocos lanata*.

Nispero de España. See *Eriobotrya japonica*.

Nispero was originally the Spanish for *Mespilus germanica*, but was transferred to the loquat when that was introduced. In Porto Rico the use of "nispero" for the sapodilla has resulted in the explanatory "de España" for the Japanese species.

No-eye pea. See *Cajanus cajan*.

Nopalea coccinellifera. TUNA MANSA.

PLATE XLVII.

This is generally reported to be a spineless cactus on which the cochineal insect is raised. In the Canary Islands, however, where there is a considerable cochineal industry, the cacti employed are by no means spineless, and are apparently the plant usually designated by botanists as *Opuntia ficus-indica*. Nopalea is reported from Porto Rico by Stahl (4: 190) as *Opuntia coccinellifera*.

Norfolk Island Pine. See *Araucaria creelsa*.

Noyó. See *Ipomoea dissecta*.

Nues. (Probably a misspelling of "Nuez.")

A tree from the interior of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 20 to 25 inches (50 to 63 centimeters). Wood, red, hard; specific gravity, 0.654; used for making furniture. (Exp. 1857.)

Nuez. See *Juglans cinerea*.

The Spanish name for the fruit of the common walnut.

Nunguey.

On sale at the market of Ponce; said to be the root of a tree; used with sugar and water in the preparation of a refreshing drink.

Nutmeg.

Many attempts have been made to render the cultivation of the nutmeg a commercial industry in various parts of South America and the West Indies, but with uniformly discouraging results, excepting in the volcanic island of Grenada, at the extreme southern end of the Windward chain. Grenada lies out of the track of



CORK WOOD (*OCHROMA LAGOPUS*): NEAR PONCE.

the hurricanes, is not subject to the dry seasons, and appears to possess all the requirements for the successful growth of this spice. The geological and climatic conditions are, however, very different from those of Porto Rico, and while it is too early, perhaps, to say that the nutmeg can not be profitably cultivated in any part of that island, it is safe to say that the probabilities of success are slight and that extensive planting would be inadvisable until a careful investigation of any proposed locality has been made.

According to several published works the nutmeg is a common tree in Porto Rico, but these references are to be understood as applying to a species of *Nectandra*, the seeds of which are in common use and sold in the markets as "nuez moscado," or nutmeg, which the general public firmly believe them to be. The seeds may be distinguished at once from the genuine article by their smaller size and more slender shape and from the fact that the two halves readily fall apart.

Nymphaea. See *Castalia*.

Ochra. See *Abelmoschus esculentus*.

Ochroma lagopus. GUANO. CORKWOOD. PLATE XLVII.

Family Bombacaceae: a tree rising to the height of 60 or 70 feet (18 to 21 meters) and sometimes 6 feet in circumference. The wood is soft on account of the rapid growth of the tree, and is used for corks by fishermen, etc. The fiber enveloping the seeds is used to make pillows and mattresses, and may be made into garments. Reported from Manati. In Jamaica called "down tree." (Stahl, 2: 100.)

A wild tree, rather abundant, 30 to 40 feet (9 to 12 meters) high, 1 foot (30 centimeters) or more in diameter. The wood is white, stained with red, luminous in aspect, sometimes silky. It is very porous, and the lightest of all woods, lighter even than true cork, the specific gravity of which is 0.240, while the latter is 0.120; fibrous and very difficult to work, because of the fibers that impede the action of the tools. In Trinidad and other places it forms an article of commerce with fishermen, who use it in place of true cork on their nets. The Indians use it to make their canoes buoyant. Called "corcho." (Grosourdy, 2: 376.)

Ochroma is said to be shipped in quantities to Cuba. The bark and roots have medicinal properties. The fiber is chestnut-colored. It is said to keep free from insects. Sold at 20 cents per pound.

Ocimum americanum. ALBAHACA CIMARRONA.

An indigenous annual of the mint family. (Stahl, 6: 190.)

Ocimum basilicum. BASIL. ALBAHACA.

An herbaceous annual or biennial, cultivated in gardens, 40 centimeters high, said to grow spontaneously in Jamaica, where it is called by its English name, "basil." (Stahl, 6: 188.)

Ocimum micranthum. ALBAHACA CIMARRONA.

An erect woody biennial, 50 centimeters high, found in cultivated fields near forests. (Stahl, 6: 189.)

Ocotea moschata.

Family Lauraceae; reported from Adjuntas.

Ogiera ruderalis. See *Eleutheranthera orata*.

Ojo de buey. See *Mucuna urens*.

Okra. See *Abelmoschus esculentus*.

Oldenlandia corymbosa. YERBA OLDENLANDIA.

A slender rubiaceous herb; a weed in cane fields. (Stahl, 5: 53.)

Oldenlandia glomerata. GRACIOSA.

An herbaceous annual, 25 centimeters high. (Stahl, 5: 52, as *Hedyotis glomerata*.)

Oldenlandia herbacea. See *Oldenlandia corymbosa*.

Olea europaea. OLIVE. ACEITUNA.

Olive culture is one of the new industries which has been suggested for Porto Rico. The olive is native to the dry climate of the desert regions about the Mediterranean. It has also been successfully introduced into similiar situations in southern California and Arizona. In Florida, however, owing doubtless to the more moist atmosphere, the production of fruit has been found impossible, although the trees vegetate with apparent vigor. Under the more tropical conditions of Porto Rico, with a warmer and more moist atmosphere, there is still less probability of the production of fruit, to say nothing of securing it in commercial quantities and of superior quality, as would be necessary in order for it to become an export crop able to compete with established industries. An experiment in the driest localities on the south side of the island might be worth while from the scientific standpoint, but the general planting of olives in the hope of establishing an industry which might be substituted for those now in distress could result only in more complete disaster.

Oleander. See *Nerium oleander*.

Olive. See *Olea europea*.

Olive mangrove. See *Avicennia nitida*

Onion. See *Allium cepa*.

PLATE XLVIII.

Opuntia.

A large genus of Cactaceae, with numerous representatives in Porto Rico. About Guayanilla the members of this genus are very abundant, forming almost a forest in some places.

Opuntia coccinellifera. See *Nopalea coccinellifera*.

Opuntia decumbens. OTAGA.

A cactus abundant on the southern and eastern coasts. (Stahl, 4: 190.)

Opuntia guanica.

A small indigenous species recently described by Professor Schumann, of Berlin.

Opuntia repens.

Known from near Guanica.

Opuntia tuna. TUNA BRAVA.

The identity of this cactus seems to be very uncertain. Some botanists have confused it with the species usually called *Opuntia ficus-indica*.

Orange, Wild. See *Naranja*.

Mr. Hill says the wild or Seville orange in the country is an indigenous tree. Numerous seedling varieties of oranges and other citrus fruits have escaped from cultivation and are to be found in unexpected places in all parts of the island. No species of citrus is, however, supposed by botanists to be indigenous in Porto Rico or in any part of the American continents or islands.

Orchata.

A drink said to be made from the "sweetened and diluted milk of almonds." In view of the prevalent belief that the nuts of *Terminalia catappa* are "almond-drags," this statement should probably be understood as referring to that species rather than to imported nuts.

Oreganillo. See *Eupatorium villosum*.

Oregano.

Leaves sold in the market of Ponce; used for seasoning food.

Oregano brujo. See *Coltus aromaticus*.

Oregano de España. See *Coltus aromaticus*.

Oregano del pais. See *Lippia micromera*.

Oreodoxa. See *Roystonea*.



SPINLESS OPUNTIA. NEAR GUAYAMA.

Oreodoxa regia. See *Roystonea regia*.

Oreopanax capitatum. PALO CACHUMBA.

Family Araliaceae: a shrub, 4 meters high, on hills near Coamo. (Stahl, 4: 198, as *Aciadophyllum capitatum*.)

Ormosia dasycarpa. PALO DE MATOS.

Family Viciaceae: a tree, 10 meters high, in mountains. Wood of good quality. (Stahl, 3: 99.)

Ormosia krugii. PALO DE MATOS.

A leguminous tree of 10 to 25 meters, known only from Porto Rico and Dominica. Specimens in the National Herbarium, collected by Sintenis, near Juncos, show the wood to be light and pithy. The leaves are very large, about 20 inches (50 centimeters) long, and the pods are brown and leathery, containing from one to three red and black beans. Another specimen from Mount Francis, Sierra de las Piedras, has much smaller, closely veined leaves, more densely pubescent on the under side.

Orosne. See *Polygala paniculata*.

Orozuz. See *Scoparia dulcis*.

Ortegon. See *Coccolobis rugosa*.

Ortelá. See *Marsypianthes hyptoides*.

Ortiga.

An ingredient for a refreshing drink, on sale by a herb dealer in the market of Ponce. Said to be derived from a tree; perhaps ortegon.

Oryza sativa. RICE.

Very little rice is now grown in Porto Rico, and this is planted without irrigation in ordinary fields, such as we might use for wheat. A considerable quantity is, however, imported from the East Indies, largely by way of Germany and Spain. The question has been raised as to whether the low grade and broken rice produced in our Southern States might not be advantageously utilized in Porto Rico, and there seems to be every probability that considerable quantities would find a ready market. In the United States rice is seldom treated as a staple article of food, but is looked upon merely as an accessory dish or dessert, for which fancy prices are readily paid. This, however, excludes from the market all of the broken kernels and lower grades. The food value of these is, however, quite as great as that of the higher-priced article, and there is a large class in Porto Rico which might be expected to avail itself of a cheap but wholesome article of this kind.

Ossea fascicularis. See *Henriettella fascicularis*.

Otaga. See *Opuntia decumbens*.

Otaheite apple. See *Eugenia malaccensis*.

Otaheite gooseberry. See *Cicca disticha*.

Ouratea litoralis.

A shrub or small tree 2 to 5 meters high, belonging to the family Ochnaceae. Reported from thickets and sandy beaches, Cangrejitos, Bayamon, Fajardo, Yuba-coa, and Manati. (Urban, Symb. 1: 363.)

Ouratea nitida. GONFIA LUSTROSA.

A shrub or tree with the same habitat and dimensions as the preceding. (Stahl, 2: 120, as *Gomphia nitida*.)

Oxalis corniculata. VINAGRILLO.

Family Oxalidaceae: a herbaceous annual, preferring stony places, rather shaded and wet. (Stahl, 2: 169.)

Oxalis corymbosa. VINAGRILLO MORADO.

A herbaceous plant, cultivated in gardens. (Stahl, 2: 170, as *Oxalis martiana*.)

Oxalis frutescens. VINAGRILLO.

A low, tortuous shrub. (Stahl, 2: 170.)

Oxalis martiana. See *Oxalis corymbosa*.**Oxandra laurifolia.** HAYA BLANCA.

An indigenous anonaceous tree, reported by Bello; also called "yaya." It reaches a height of 30 feet (9 meters), and fruits in January. (Stahl, 2: 28.)

Oxandra virgata. HAYA PRIETA.

An indigenous tree, originally described from Cuba.

Pacholi.

The strongly aromatic roots of a grass, perhaps *Andropogon nardus*, were on sale in small bundles in the market of Ponce. They are put in chests to keep out moths.

Pachyrrhizus angulatus. YAM BEAN. FRIJOLILLO DE ÑAME.

A shrubby, leguminous vine, having thickened horizontal roots, several inches in diameter, and 6 to 8 feet (2 to 2.5 meters) long. Although edible, these roots are inferior to the true yam "cassava," or "taro," and when cooked they are said to resemble turnips in texture and taste. In some countries they form a resource in times of scarcity resulting from the failure of the usual crops. In Porto Rico this species was observed only in the vicinity of Rio Piedras, though it doubtless exists in other parts of the island. (Stahl, 3: 70.)

Pachyrrhizus tuberosus. YAM BEAN.

A leguminous vine with large edible roots; reported from Rincon

Pachyrrhizus angulatus. See *Pachyrrhizus angulatus*.**Pajuil.** See *Anacardium occidentale*.

A tree from all parts of the island; height, 12 to 15 feet (3.5 to 4 meters); diameter, 9 to 10 inches (22 to 25 millimeters). Wood pinkish, soft; specific gravity, 0.550; fruit tree. (Exp. 1857.)

Palm.

The palms thus far reported from Porto Rico are as follows:

Phoenix dactylifera (date palm), *Inodes causiaram* (Porto Rican hat palm), *Thrinax parviflora*, *Thrinax ponceana*, *Thrinaxoma alta*, *Thringis lara*, *Thringis latifrons*, *Acris attenuata* (Ilume), *Areca catechu* (Betel nut palm), *Roystonea borinquena* (Royal palm), *Acris monticola* (Palma de Sierra), *Bactris acanthophylla*, *Bactris paroniana*, *Curima calophylla* (coyure), *Acrocomia media* (corozo), *Cocos nucifera*, and *Cocops rivalis*.

A synopsis of the above species appeared in the Bulletin of the Torrey Botanical Club for October, 1901.

Palicourea barbinervis.

Family Rubiaceae; reported from Lares and Anones. Bello gives the common name "tafetán."

Palicourea coccinea.

Reported from Aibonito.

Palicourea crocea. CACHIMBO.

A biennial shrub, 1 to 2 meters high, found in waste places; known from Lares. (Stahl, 5: 77.)

Wood light-colored, soft; specific gravity, 0.482; not used. (Exp. 1857.)

Palicourea pavetta.

A shrub 6 feet (1.8 meters) high; known from Juncos, on Mount Goyo.

Palicourea umbellata. CACHIMBO VERDE.

A biennial shrub, 1 meter high, found at the base of mountains. (Stahl, 5: 78.)

Palma christi. See *Ricinus communis*.

Palma costa. See *Roystonea borinquena*.

A name applied to the royal palm in the vicinity of Coamo.

Palma de grana. See *Roystonea borinquena*.

Palma de la Sierra. See *Acrista monticola*.

Palma de los cerdos. See *Roystonea borinquena*.

This means "pig palm," and is one of the names applied to the royal palm on the south side of the island.

Palma mauricia. See *Mauritia flexuosa*.

A Cuban name, also said to be applied to *Mauritia setifera* (*setigera?*).

Palma moriche. See *Mauritia flexuosa*.

Palma real. See *Roystonea borinquena*.

Palmetto. See *Sabal* and *Thrinax*.

Palo blanco. See *Drypetes glauca*.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood white, soft; specific gravity, 0.866; used in building houses. (Exp. 1857).

Captain Hansard gives the specific gravity as 0.77. Used in hut building.

Palo bobo. See *Coccolobis diversifolia* and *Pisonia subcordata*.

A tree from all parts of the island; height, 30 to 35 feet (9 to 11 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood white, very soft; specific gravity, 0.531; not used. (Exp. 1857).

Captain Hansard gives the specific gravity at 0.54.

Palo bronco. See *Malpighia urens*.

Palo cachumba. See *Gilibertia arborea*, *G. laurifolia*, and *Oreopanax capitatum*.

Palo cano.

A tree from all parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood white, hard; specific gravity, 0.990; not used. (Exp. 1857).

Palo colorado. See *Myroxylon schumackerianum* and *Ternstroemia luquillensis*.

Palo de aceite. See *Copaifera officinalis*.

Palo de boyo. See *Erythrina micropteryx*.

Palo de Brasilete. See *Dalbergia monetaria*.

Palo de burro. See *Capparis cynophallophora*, *C. frondosa*, *C. latifolia*, and *C. verrucosa*.

Palo de cabra. See *Symplocos polyantha*.

Palo de Campeche. See *Haematoxylon campechianum*.

Palo de candela. See *Myroxylon schumackerianum*.

Palo de corcho. See *Pisonia obtusata*.

Palo de cucubano. See *Gnettarda scabra*.

Palo de doncella. See *Byrsonima lucida*.

Palo de galleria.

A tree from all parts of the island; height, 25 to 30 feet (8 to 9 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood white, soft; specific gravity, 0.409; used in building houses. (Exp. 1857.)

Palo de gallina. See *Alchorneopsis portoricensis*.

Palo de gangulin. See *Gilibertia laurifolia*.

Palo de garrocha. See *Quararibea turbinata*.

Palo de guitarra. See *Citharexylum quadrangulare*.

Palo de hierro. See *Leora ferrea*, *Sarcophalus reticulatus*, and *Scutia ferrea*.

Captain Hansard gives the specific gravity of this wood as 1.07.

Palo de hoz. See *Drepanocarpus lunatus*.

Palo de hueso. See *Ilex nitida*, *Linociera compacta*, and *Maypea domingensis*.

Palo de jaqueca. See *Thespesia populnea*.

Palo de Maria. See *Calophyllum calaba*.

Palo de matos. See *Ormosia dasycarpa* and *O. krugii*.

Palo de multa. See *Eugenia ligustrina*.

Palo de muñeca. See *Cordia borinquensis* and *Rauwolfia nitida*.

Palo de nispero cimarron. See *Symplocos lanata*.

Palo de oreja. See *Cassipourea elliptica*.

Palo de pan cimarron. See *Bocconia frutescens*.

Palo de perico. See *Cordia ulmifolia*.

Palo de peronias. See *Adenanthera parviflora*.

Palo de pollo. See *Pterocarpus draco*.

Palo de ramon. See *Bauera portoricensis*.

Palo de rayo. See *Parkinsonia aculeata*.

Palo de toro. See *Faramea odoratissima*.

Palo de vaca. See *Gilibertia laurifolia*.

Palo de geringa. See *Moringa moringa*.

Palo hediondo. See *Loucheocarpus latifolius*.

Palo hueso.

A tree from the northeastern part of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood olive green, hard; specific gravity 1.029; used in making shelves. (Exp. 1857.)

Groussourdy (2: 407) describes the wood as dirty white, and says that it is used for fuel and the framework of rude houses.

Palo llorón. See *Antirrhoea lucida* and *Malanea macrophylla*.

Palo moro. See *Psychotria*.

This name is applied to numerous species of this very large genus of Rubiaceae.

Palo pollo. See *Pterocarpus draco*.

A tree from the swampy parts of the island; height, 35 to 40 feet (11 to 12 meters); diameter, 20 to 25 inches (50 to 62 centimeters). Wood white, soft; specific gravity 0.614; not used. (Exp. 1857.)

Grows to a considerable height and attains great thickness; very irregular at base; wood used for handles for machetes and pans for gold miners; texture hard and solid.

Palo puerco.


Captain Hansard gives a specific gravity of 0.87 for this wood, but says that he does not know the tree.

Palo punz.

A tree from all parts of the island; height, 25 to 30 feet (8 to 9 meters); diameter, 10 to 12 inches (25 to 30 centimeters). Wood ash-colored, soft; specific gravity 0.725; used in building houses. Also called "palo santo." (Exp. 1857.)

Palo rosa.

A tree from the island of Vieques; height, 25 to 30 feet (7 to 9 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood pinkish, hard; specific gravity, 0.994; used in building houses. (Exp. 1857.)



PARA GRASS PANICUM MOLLE.

Palo santo. See *Palo puz*.

A wild tree 30 feet (9 meters) or more in height, with a trunk 8 to 12 inches (20 to 30 centimeters) in diameter. Furnishes a soft ash-colored wood that breaks with a vertical fracture. Used in building houses and fences. (Grosourdy, 2: 407.)

Captain Hansard gives 0.70 as the specific gravity of this wood.

Pana. See *Gilibertia arborea*.

Pana cimarrona. See *Didymopanax morototoni*.

Panales.

Small froth-like rolls made of sugar and white of egg. They are not for eating, but are dissolved in water to make a sweetish drink.

Panicum maximum. GUINEA GRASS. YERBA DE GUINEA.

The guinea grass is said to have been used as bedding in the slave ships, and to have become established in tropical America accidentally after floating ashore from the rubbish thrown overboard when the ships were unloaded. The origin of this account should be traced as a note on plant introduction.

Panicum molle. PARA GRASS. MALOJILLO. PLATE XLIX.

As a pasture grass this species is second only to the Guinea or "yerba de Guinea." It is grown in the swampy meadows of the north coast and wherever the ground is too moist for the latter species.

The Malojillo is deep green in color, decumbent at base, with very hairy joints. The flower stalk has fewer and shorter branches than the Guinea grass. Both species produce luxuriant crops, but both seem too coarse to represent the food value of the same amount of the temperate grasses.

Papa. See *Solanum tuberosum*.

Papaw. See *Carica papaya*.

Papaya. See *Carica papaya*.

This is a cultivated fruit, not a mountain forest tree as stated by Hill.

Papaya lechosa. See *Carica papaya*.

Para grass. See *Panicum molle*.

Paralejo velludo. See *Heteropteris laurifolia*.

Parasita. See *Dendropremou bicolor*.

Parathesis crenulata. See *Seca-garganta*.

Family Myrsinaceae; an erect shrub, 1 to 2 meters high, on the sides of ravines. Bello gives the common name "rasca garganta." (Stahl, 6: 37, as *Ardisia crenulata*.)

Parcha. See *Passiflora laurifolia*.

Parcha angosta. See *Passiflora suberosa*.

Parcha cimarrona. See *Passiflora maliformis*.

Parcha colorada. See *Passiflora rubra*.

Parcha de garjos. See *Passiflora serrata*.

Parcha multiflora. See *Passiflora multiflora*.

Parcha ojillosa. See *Passiflora tulae*.

Parcha palida. See *Passiflora suberosa*.

Parcha pelitiesa. See *Passiflora suberosa*.

Parcha yedra. See *Passiflora suberosa*.

Paritium elatum. See *Hibiscus elatus*.

Paritium tiliaceum. EMAJAGUA.

PLATE L.

A large shrub or small tree widely distributed throughout the warm regions of the Old and New worlds. It was in use throughout tropical America before the advent of Europeans, and is still one of the most important fiber plants for local and domestic uses. The bast furnishes a strong and flexible fiber, comparable to jute, and having the remarkable quality of becoming stronger by long maceration in water. Although this has been known for many years, and although the plant would grow readily and without cultivation in moist waste tracts common along the coasts of many tropical countries, there seems to have been no serious attempt at introducing the fiber to commerce or adapting it to civilized uses.

The oft-quoted experiment of Roxburgh, as reported by Watt, is as follows:

"A line broke when white with a weight of 41 pounds; after being tanned, with 62 pounds, and after having been tarred, with 61 pounds. A similar line macerated in water for one hundred and sixteen days broke when white with 40 pounds; tanned, 55 pounds, and tarred, 70 pounds. These observations are of great interest, for of the other fibers experimented with by Roxburgh, the majority were rotten after maceration, and no other fiber showed so marked an improvement for cordage purposes when tarred. English hemp and Indian grown hemp, treated in the same manner, were found to be rotten, and sunn-hemp broke with 65 pounds and jute with 60 pounds. The power of endurance under water is, therefore, a point of great importance.

"The fiber is readily separated from the green or unsteeped branches, the work of preparation being less tedious than applies to the other fiber-yielding plants of this genus. It appears to be well adapted for making ropes, mats, and possibly paper (*Cameron*). The fiber seems highly suitable for the paper trade, and immense quantities of it might be conveyed in boats to the paper mills of Bengal."

The last suggestion of paper making would seem to be worthy of investigation by our manufacturers in view of the increasing scarcity of raw materials for a product of good quality. Emajagua could be produced with extreme cheapness. Replanting would not be necessary, as sprouts come up abundantly from the stumps. Cultivation could be carried on after the manner of the basket osiers.

In Porto Rico nearly all the ropes in use are made from emajagua.

Recommended in Jamaica by Dr. Morris for windbreaks about cacao plantations.

Also called "Majagua."

Parkinsonia aculeata. JERUSALEM THORN. PALO DE RAYO.

A leguminous tree related to the Haematoxylon or logwood. It is supposed to be native in some parts of the American continent, but is now so generally cultivated that the original habitat is in doubt. (Stahl, 3: 119.)

Parra cimarrona. See *Vitis caribaea*.**Parthenium hysterophorus.** AJENJO CIMARRON.

An herbaceous annual composite, found in sandy places. In Jamaica this plant is called "mugwort" and Lunan gives a number of cases in which it was supposed to have curative value in healing different kinds of eruptive diseases. (Stahl, 5: 157.)

Pascueta. See *Erigeron canadensis*.**Pascueta amarilla.** See *Erigeron rivularis*.**Passiflora angustifolia.** See *Passiflora suberosa*.**Passiflora foetida.** TAGUA-TAGUA.

Family Passifloraceae; a shrubby vine found along roadsides. It has a disagreeable odor, but the flowers are beautiful and the fruit edible; called "love in a mist" in Jamaica. (Stahl, 4: 144.)

Passiflora hederaceae. See *Passiflora suberosa*.**Passiflora hirsuta.** See *Passiflora suberosa*.



EMAJAGUA (PARITIUM TILIACEUM).

Passiflora laurifolia. PARCHA.

A woody twining climber, cultivated in gardens; fruit tart and agreeable. (Stahl, 4: 147.)

Passiflora maliformis. PARCHA CIMARRONA.

A woody climber, found in waste places. Reported from Utuado. (Stahl, 4: 148.)

Passiflora multiflora. PARCHA MULTIFLORA.

The velvety-pubescent long-oval leaves seem characteristic among the Porto Rican species; collected by Sintenis at Manati.

Passiflora pallida. See *Passiflora suberosa*.**Passiflora quadrangularis.** GRANADILLA.

A twining vine, cultivated in gardens as an ornamental; used for covering arbors. Fruits agreeable. The pulp mixed with water and sugar makes a refreshing drink. (Stahl, 4: 145.)

Passiflora rubra. PARCHA COLORADA.

An herbaceous vine found in waste places. (Stahl, 4: 150.)

Passiflora serrata. PARCHA DE GARCOS.

A vine found in waste places and mountains. (Stahl, 4: 149.)

Passiflora suberosa.

According to the Kew Index, Stahl's species of *Passiflora angustiflora* "parcha angosta," *P. pallida* "parcha palida," *P. hirsuta* "parcha pelitiosa," and *P. hederacea* "parcha yedra" are included in *Passiflora suberosa*.

Grows to a height of 20 feet (6 meters). Reported from Yabuco and Guanica.

Passiflora tulae. PARCHA OJILLOSA.

An indigenous species recently described by Urban as new. It is *Murucuja ocellata* of Bello and *Passiflora murucuja* of Stahl, called "bull hoof" in Jamaica. (Stahl, 4: 151, as *Passiflora murucuja*.)

Pata de cabra.

These samples [pata de cabra and algarrobo] are so similar in detail of structure as to be from the same or closely related species. Color, rich blackish brown, irregularly mottled, and streaked with areas of pale reddish brown; sapwood (present in pata de caba), light brown. Ducts conspicuous, occurring singly or two together, between two medullary rays or often interrupting a ray; evenly diffused. Medullary rays conspicuous, visible to the naked eye, often disconnected. Wood fibers strongly interlaced, giving smoothed surface a "curled" appearance. Very attractive cabinet woods. Resemble forms of mesquit, especially *Prosopis odorata*. (Hill and Sudworth.)

Pata de gallina. See *Phoradendron chrysocarpum*.**Patata.** See *Solanum tuberosum*.**Patilla.** See *Citrullus vulgaris*.**Paullinia pinnata.** BEJUCO PRIETO.

Also called bejuco de costilla. Family Sapindaceae; a woody climber with compound leaves and a winged rhachis like that of *Inga vera*. It climbs by means of tendrils attached to the flower cluster. According to the Index Kewensis this name is a synonym of *Serjania curassavica*. (Stahl, 2: 155.)

Pavonia. See *Hibiscus rosa-sinensis*.**Pavonia coccinea.** CADILLO ANARANJADO.

Family Malvaceae; a woody herb, 1 meter high. (Stahl, 2: 85.)

Pavonia encendida. See *Hibiscus radiatus*.**Pavonia racemosa.** CADILLO DE CIÉNEGA.

A shrub 2 to 3 meters high, frequenting mangrove swamps. (Stahl, 2: 84.)

Pavonia spinifex. CADILLO ESPINOSO.

An annual or biennial shrub, 1 to 2 meters high, found on mountains and in shaded places. (Stahl, 2: 83.)

Pavonia typhalea. CADILLO PEQUEQUE.

A shrubby annual, growing in shaded wet places. The small white flowers are collected in a head and give an appearance quite different from related members of the Malvaceae. Specimens collected at El Rio were called "anamu" and were reputed a remedy for chigoes. (Stahl, 2: 81.)

Paw paw. See *Carica papaya*.**Peanut.** See *Arachis hypogea*.**Pectis humifusa.** YERBA DE SAN JUAN.

A woody annual composite from sandy seashores. (Stahl, 5: 145.)

Pectis serpyllifolia. See *Pectis humifusa*.**Pedilanthus tithymaloides.**

A peculiar fleshy Euphorbiaceous plant, with an irregular wing along the midrib on the lower side of the leaves. Frequently planted for hedges.

Pegadora. See *Meuzelia aspera*.**Pega-palma.** See *Maregravia umbellata*.**Pega-pega.**

This name meaning "stick-stick," like our "stick-tight," is probably applied to several seeds adapted for attaching themselves to one's clothes, as "pica-pica" is used for many spiny plants.

At Santurce *Priva echinata* and *Aeschynomene americana* were called "pega-pega."

Pega-palo. See *Bignonia unguis-cati*.**Pega pollo.** See *Boerhavia scandens*.**Pegoge.** See *Tabernaemontana citrifolia*.**Peireskia.**

Family Cactaceae; a thick, spiny shrub or small tree, valuable mostly as a stock on which to graft the various cacti. Known from specimens collected by Sintenis at Penuelas, where it was cultivated in gardens.

Penaejuelo.

A tree from the northeastern part of the island; height, 45 to 50 feet (13 to 15 meters); diameter, 15 to 20 inches (37 to 50 centimeters). Wood, ash-colored; rather hard; specific gravity, 0.810; used for lumber. (Exp. 1857.)

Pendejuelo.

Evidently the same as Penaejuelo.

A wild tree, 50 feet (15 meters) high, with a trunk reaching a diameter of 18 inches (45 centimeters). Furnishes an ash-colored wood that breaks with an oblique fracture. Its common use is for lumber used in building crude houses. (Gronourdy, 2: 408.)

Péndola. See *Citharexylum quadrangulare*.**Péndula.**

A tree from the eastern part of the island; height, 45 to 50 feet (13 to 15 meters); diameter 15 to 20 inches (37 to 50 centimeters). Wood, yellow, hard; specific gravity, 0.942; used for lumber. (Exp. 1857.)

A large tree, trunk rather short; wood very hard, making most excellent boards and posts. A very valuable wood with a specific gravity of 0.84. (Captain Hansard.)

Péndulo blanco. See *Vitex divaricata*.**Péndulo colorado.** See *Citharexylum quadrangulare*.



PENTARRHAPHIA ALBIFLORA.

Pentarrhaphia albiflora.

PLATE LI.

Family Gesneriaceae: this and *P. longiflora* seem to be very similar species; rather small shrubs growing in rocky places, and having the characteristic long-stalked asymmetric flowers, greenish yellow, mottled with dark reddish stripes and spots.

Pentarrhaphia longiflora. SPANISH TEA. See *Pentarrhaphia albiflora*.

Pentarrhaphia reticulata. CONRADIA.

An herb 8 to 12 centimeters high; found among rocks and in shady places. (Stahl, 6: 259.)

Peperomia.

A large genus of Piperaceae represented in Porto Rico by numerous species, mostly rather small and delicate herbs.

Peperomia portoricensis. YERBA DE GUAVA.

An indigenous herb from the mountain forests near Aybonito, Adjuntas, Utuado, and Maricao. (Urban. Symb. 1: 292.)

Pepinillo. See *Melothria fluminensis*.

Pepinito. See *Cucumis anguria*.

Pepino. See *Cucumis anguria*.

Pepino angolo. See *Sicana odorifera*.

Pepita amarga. See *Fevillea cordifolia*.

Pepper.

The red, Cayenne, Chile, or garden peppers belong to the genus *Capsicum* of the family Solanaceae, while the true black or white pepper is derived from species of *Piper*.

Peregil cimarron. See *Scrophularia micrantha*.

Perico cimarron.

This name was applied at Vega Baja to a species of *Acalypha* (No. 1035), still unidentified.

Periwinkle. See *Vinca rosea*.

Peronia. See *Abrus precatorius*.

Peronilas.

Bello gives this name for *Abrus precatorius*, but probably by error.

Persea carolinensis.

See note under "naguacatillo."

Persea gratissima. AGUACATE.

Variously known as "alligator-pear," "butter-pear," "avocate," "avocado," etc. Family Lauraceae. This well-known fruit of the Tropics has the external appearance of a large pear, but on being cut open is found to have a large central seed, which separates readily from the rather firm, somewhat buttery outer pulp, which is the part eaten. The "butter-pear," as it is sometimes called, is a salad fruit—if such a term may be used—being eaten with salt, vinegar, pepper, and other condiments. The pulp is scraped away from the outer hard skin with a spoon. Opinions differ greatly as to the value of this fruit; some are extremely fond of it, while others consider it quite insipid. Properly seasoned, it blends finely with dressings, and has recently been used as an ingredient of rich and elaborate salads. Treated with oil, salt, and tarragon vinegar, it has also been recommended cut in slices and served as a relish, or the "butter" mashed up with such a dressing may be applied to sandwiches. At present but a small quantity of this fruit is marketed in New York, but the demand is increasing and the prices are good, 20 cents apiece being an average figure, at which the business would certainly be very profitable. According to Semler, an oil extracted from

the alligator pear is used in large quantities in America in soap manufacture, but the seat of the oil industry is not stated.

There are a great many varieties of the alligator pear, but the forms already in Porto Rico compare well with those of other countries. In view of the difficulties experienced in shipping this fruit, an effort should be made to secure some of the thick-skinned varieties of Central America. Selection, or the introduction of new varieties that will extend the season, will also greatly improve the chances of making their culture profitable.

One danger in relying on alligator pears as a source of income arises from the brittle nature of the wood, the branches being easily broken by strong winds, only occasional trees having been left uninjured by the storm of August, 1901. The tree is rather handsome, but never reaches very large size, old specimens being 35 to 40 feet (10 to 12 meters) high and a foot or 18 inches (30 to 45 centimeters) in diameter. There is a wild species in Cuba, *P. sylvestris*, there called "*aguacate silvestre*."

It furnishes a rather soft wood, with a specific gravity of 0.643. The grain is very fine. The texture is fibrous and rather brittle. The color is reddish brown, or sometimes light brown, with poorly-marked lines or undulations, more visible in the white parts, while the heart is mottled with light, irregular, curved spots of different colors, disposed almost symmetrically, and giving to the whole a most beautiful aspect. This wood, which is not used at all, is very brittle, but might, in our opinion, be employed in cabinet work. The bark is rather thick, very rough, irregular, ash-colored. (Grosourdy, 2: 358.)

Petangueira. See *Eugenia portoricensis*.

Petitia domingensis. CAPÁ BLANCA.

Family Verbenaceae; a tree 10 meters high. The hard wood is good for fine work. Reported from Rancon and Utuado. (Stahl, 6: 223.)

Petiveria alliacea.

Family Phytolaccaceae; a slender weed very common in waste places but preferring partial shade. It has a very disagreeable odor, strongly suggesting onions or garlic, and is said to injure the milk of cows, which seems entirely probable.

Phasemy. See *Phaseolus semierectus*.

Phaseolus adenanthus. HABICHUELA CIMARRONA.

Family Viciaceae; a woody, climbing annual in hedges and at the base of mountains. (Stahl, 3: 76.)

Phaseolus lanceolatus. HABICHUELA CIMARRONA.

(Stahl, 3: 78.)

Phaseolus lunatus. HABAS. LIMA BEAN.

A twining climber, both cultivated and escaped. (Stahl, 3: 81.)

Phaseolus semierectus. PHASEMY. HABICHUELA PARADA.

An herbaceous, erect, branching annual 50 centimeters high, found in pastures. (Stahl, 3: 77.)

In British Guiana experiments with this plant have seemed to warrant the belief that it will prove to be a tropical substitute for alfalfa.

Phaseolus vulgaris. BEAN. HABICHUELA.

Several varieties of the common bean are imported into Porto Rico, but the one most favored for local cultivation is a rather large, red variety. A bean collected near Ponce (No. 796) has the pods warted along both edges. It was called "haba," but is not the Lima bean to which that name is more often applied.

Phoebe cubensis.

Family Lauraceae; the form *antillana* is reported from Bayamon.

Phoenix dactylifera. DATE. DATIL.

The date palm has been introduced into Porto Rico, and while the trees reach a considerable size, particularly on the drier southern side of the island, it is not known that any fruit of good quality has been produced. Specimens of a date palm were obtained by Sintenis at Puerto Real near Cabo Rojo, and it is in that region that the tree might be expected to thrive best in Porto Rico, and the results of experiments which have been made would be of interest.

Although the date palm grows with apparent vigor in Porto Rico it is not likely that it will ripen fruit of marketable quality, as the climate is too cool and too moist. The date reaches perfection only in desert regions, and while it can stand slight frosts it must have exceedingly hot weather during the ripening season. That the tree will flourish is no indication that it will produce good fruit. In the Canary Islands an indigenous species of date palm is extensively grown for the sake of the leaves, which are made into baskets and serve other domestic purposes, but the fruit is nearly worthless, and genuine date palms which have been introduced do not succeed.

As yet dates are not known to have been successfully raised in America except in the desert regions of Arizona, California, and Mexico. In the latter country the date was introduced very early and was spread to Lower and Upper California by the Spanish missionaries. Although, according to Forbes's "California," dates and figs were exported from Lower California as early as 1837, the industry has not been greatly developed, owing, perhaps, to the fact that only seedlings were planted. Of late it has been found that the date succeeds well in the extremely dry and hot regions of Arizona, and cuttings of superior varieties have been imported in order that fruit of uniform, marketable quality may be secured.

Pholacilia diversifolia. See *Trichilia simplicifolia*.**Phoradendron chrysocarpum.** PATA DE GALLINA.

A parasitic shrub of the family Loranthaceae. Bello gives the name "yerba capitana" for *Phoradendron berterianum*, which Urban refers to the present species. (Urban, Add. 4: 357.)

Phoradendron dichotomum. YERBA CAPITANA.

A parasitic shrub said to have been found on *Mangifera indica* in Haiti, but known elsewhere only from native trees.

Phoradendron quadrangulare. YERBA CAPITANA.

Also called "quasimilla de Canario." A parasitic shrub related to the mistletoe. It attacks numerous species of native trees.

Phthirusa emarginata.

Family Lauraceae; reported from Aybonito.

Phthirusa parvifolia.

From Guanica.

Phthirusa portoricensis. See *Psychotria pendula*.**Phyllanthus grandifolius.**

Family Euphorbiaceae; reported from Manati.

Phyllanthus nobilis antillanus.

From Fajardo and Aybonito.

Phyllocactus stenopetalus.

Family Cactaceae; cultivated in gardens at Pennelas. (Sintenis.)

Physalis angulata. SACABUCHE ANGULOSO.

Family Solanaceae; an herbaceous annual, 50 centimeters high, found in various parts of the island. The fruits are edible and in Jamaica are called winter cherries. The plant is one of the ground-cherry group. (Stahl, 6: 122.)

Physalis linkiana.

An herbaceous annual or weed; from near Guanica.

Physalis pubescens. SACABUCHE PELUDO.

An herbaceous annual, 60 centimeters high, common. This is the common cultivated ground cherry or strawberry tomato. (Stahl, 6: 121.)

At Arecibo the fruit of a species of *Physalis* was said to be used for stomach troubles, and the roots for toothache.

Physic nut. See *Jatropha curcas*.

Pica-pica. See *Mucuna pruriens* and *Fleurbaea aestuans*.

Picha de gato. See *Scolosanthus grandifolius*.

Pichana. See *Wissadula rostrata*.

Pichana mocha. See *Abutilon leiospermum*.

Pichana peduncular. See *Abutilon pedunculare*.

Picramnia pentandra. GUAREMA.

Family Simarubaceae; a shrub 10 to 15 feet (3 to 5 meters) high, from the vicinity of Cayey and Lares. The common name was recorded by Bello.

Picrasma excelsa. BITTERWOOD.

A small tree of the family Simarubaceae; found in Jamaica, Antigua, and several of the smaller islands. The bark and wood are extremely bitter and are now preferred to the genuine quassia from Guayana and Brazil. *Picrasma* is found in forests of low elevation near the sea.

Pictetia aristata. TACHUELO.

PLATES LII, LIII.

Family Viciaceae; a tree 6 to 8 meters high. The wood is extremely hard and heavy. Found in the southeastern coastal regions. (Stahl, 3: 50.)

Hill identifies "tachuelo" with "hachuelo," and with the calabash tree, (*Crescentia*). Our informants, including Captain Lugovine and Señor Besosa, agree in giving the name to an extremely hard and heavy wood, which endures indefinitely in the ground and is used in supporting houses. The leaves have a spine from the midrib. A photograph of a branch was taken at Penon and a "habitusbild" of a clump of shrubby growth about 20 feet (6 meters) high in an open pasture at Coamo Springs.

Pictetia squamata.

A specimen from Coamo is in the National Herbarium.

Pigeon pea. See *Cajanus cajan*.

Pigeon wood.

A name applied in the British West Indies to several trees belonging to the genera *Dipholis*, *Diospyros*, and *Coccolobis*.

Pigweed. See *Amaranthus paniculatus* and *A. tristis*.

Pilea ciliaris.^a

Family Urticaceae. This and *P. elegans*, broad-leaved species, are quite unlike *P. trianthemoides*. Known from Adjuntas and Sabana Grande.

Pilea trianthemoides. ARTILLERY PLANT. VERDOLAGUILLO.

A very fine-leaved, fern-like plant, common in most rocky situations where the ground is moist; also on roofs of houses in mountain districts. The stone water filters and storage jars for water are often overgrown with this species.

Pilocarpus racemosus.

An indigenous shrub or small tree, 2 to 5 meters high, belonging to the Rutaceae. Reported from Aguado, Agnadilla, and near Rio Grande.


Pimenta acris. See *Anomis caryophyllata*.

Pimenta officinalis. See *Pimenta pimenta*.

^aThe name *Adicea* Raf. antedates *Pilea* Lindl., but the species here referred to have not been transferred to that genus.



TACHUELO (PICTETIA ARISTATA)



TACHUELO ('PICTETIA ARISTATA'), SHOWING ARISTATE LEAVES.

Pimenta pimenta. ALLSPICE.

In Urban's revision of the Myrtaceae (Add. 2: 89) this species is not reported from Porto Rico, but the Porto Rican references of Stahl and other writers are applied to *Amomis caryophyllata*.

A tree belonging to the Myrtaceae and formerly placed in the genus *Eugenia*. Mr. Hill reports it as abundant on the south side of Porto Rico, in the region between Guayama, Coamo, and Ponce; but the species is not, in fact, known to occur in Porto Rico where the names "pimienta" and "pimienta malagueta" are applied to the bay-rum tree, *Amomis caryophyllata*. The allspice tree has been found in Cuba and Jamaica, also in Mexico and Central America, but not, as far as known, anywhere else in the West Indies. The allspice of commerce consists of the unripe fruits, which must be dried quickly before the flavor has time to deteriorate. Large quantities are shipped to Europe and the United States from Jamaica, where the tree is cultivated, or allowed to spread itself over waste lands or pastures. Hill explains the fact that no allspice has been exported from Porto Rico by saying that the fruits are allowed to drop from the trees and are not properly cured.

Pimenta vulgaris. See *Amomis caryophyllata*.

Pimento. See *Pimenta pimenta*.

Pimienjo.

Captain Hansard gives the specific gravity of this wood as 1.31.

Pimienta.

This name, which properly means "pepper," has been applied to *Amomis caryophyllata*, also in the combination "pimienta malagueta." In Cuba and Jamaica, where the true allspice tree (*Pimenta pimenta*) grows, the same names are used for it.

Pimienta malagueta. See *Amomis caryophyllata*.

Pimiento. See *Capsicum annum* and *Amomis caryophyllata*.

Pina.

Enumerated by Hill among "forest trees yielding fruits." There is nothing to indicate the identity of the plant alluded to.

Pindar. See *Arachis hypogea*.

Pineapple. See *Ananassa ananas*.

Pino.

A tree from the interior of the island; height, 45 to 50 feet (13 to 15 meters); diameter, 12 to 15 inches (30 to 37 centimeters). Wood white, very hard; specific gravity, 0.960; used in boat building. Also called "cipil." (Exp. 1857.)

Pinon espinoso. See *Erythrina corallodendron*.

Piñuela.

A name applied to numerous wild plants having a real or fancied resemblance to the pineapple. The numerous wild species of *Tillandsia* and other bromeliaceous plants are called "piñuelas," as are also orchids, both terrestrial and epiphytic. One of the latter, with large fleshy bulbs, furnishes a strong glue or varnish-like cement, used particularly for joining the rudely manufactured domestic guitars. It is quite rare, and each bulb is said to be valued at 25 centavos.

Pinzona calineoides.

Family Dilleniaceae. A climbing vine from Utuado.

Piper aduncum.

A species agreeing with other West-Indian material in the National Herbarium which was collected near Toa Baja (No. 259). The leaves are minutely but distinctly scabrous on both sides. A shrub 8 feet (2.5 meters) high.

Piper citrifolium. HIGUILLO AMARGO.

Collected at Atola Teja, between Bayamon and Toa Baja (No. 240). Very similar in general appearance to the form identified as *P. aduncum*, but agreeing with the present species in having the leaves smooth on both sides.

Piper macrophyllum.

Family Piperaceae; a shrub 12 to 20 feet (4 to 6 meters) high on rocky hills; known from Adjuntas. (Grisebach.)

Piper marginatum.

Known from Aguadilla.

Piper peltatum. See *Heckeria peltata*.**Piper umbellatum.** See *Heckeria umbellata*.**Piptadenia peregrina.** COJOBANA.

Also known as Cojobilla. Family Mimosaceae; a tree 5 meters high; older growth covered with spines; found at Bayamon. (Stahl, 3: 144-152.) See note under *Acacia angustiloba*.

Piptocarpha triflora.

A composite shrub, known from Sierra de las Piedras.

Piquete. See *Bidens leucantha* and *Cosmos caudatus*.**Piriqueta cistoides.** See *Piriqueta villosa*.**Piriqueta ovala.** See *Piriqueta ovata*.**Piriqueta ovata.** PIRIQUETA OVALA.

Family Turneraceae. (Stahl, 4: 159.)

Piriqueta villosa. PIRIQUETA.

An herbaceous erect annual, 40 to 50 centimeters high, in dry sandy soil. (Stahl, 4: 157.)

Pisonia aculeata. ESCAMBRON.

Also called "uña de gato," or "cat's claw." A spiny shrub belonging to the family Nyctaginaceae; reported from near Manati.

Pisonia obtusata. MAJAGUA QUEMONA.

Also called "palo de corcho." An indigenous shrub.

Pisonia subcordata. PALO BOBO.

An indigenous tree reaching a height of 20 meters. According to Urban two forms, *albida* and *gigantophylla*, have been found in Porto Rico; reported from Barranquitas. (Urban, Add. 3: 318.)

Pistia stratiotes. WATER-LETTUCE.

Family Araceae; a floating, stemless, stoloniferous herb, known from near Manati.

Pitajaya. See *Cereus triangularis*.**Pitangueira.** See *Eugenia biflora*.**Pithecolobium flicifolium.** COJOBANA.

Family Mimosaceae; a forest tree 10 meters high; leaves finely divided; spineless. (Stahl, 3: 137.)

Pithecolobium saman. SAMAN.

Some writers state that the name "algarrobo" is applied to this species. A large, spreading tree, 15 to 20 meters high, planted for shade in yards and public grounds. The trunk is inclined to be short and thick, while the branches are horizontal and extremely long. According to Stahl the wood is not very hard, with the heart of a handsome red color. On account of its lack of durability it is very little used. The flowers appear in spring and summer, and all the growing parts are slightly hairy. (Stahl, 3: 154.)