

First record of the box jellyfish *Tripedalia cystophora* (Cnidaria: Cubozoa: Tripedaliidae) in the Gulf of Mexico

Robert M. Lasley Jr*, Cheryl Lewis Ames, Robert Erdman, Sheri Parks and Allen G. Collins

(RML, SP) Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission, St. Petersburg, Florida, U.S.A.

(AC, CLA) Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC, U.S.A.

(CLA) Biological Sciences Graduate Program, University of Maryland, College Park, Maryland, U.S.A.

(RE) Department of Biological Sciences, Florida Gulf Coast University, Fort Myers, Florida, U.S.A.

(AGC) National Systematics Laboratory, NOAA Fisheries, National Museum of Natural History, Smithsonian Institution, Washington, DC, U.S.A.

Abstract.—Although observations of the box jellyfish *Tripedalia cystophora* have been widely documented since it was first recorded in Jamaica in 1897, to date there are no published reports of its occurrence in the Gulf of Mexico. Eighteen specimens of *Tripedalia cystophora* (Cubozoa: Tripedaliidae), 12 and 6, respectively, were collected from a mangrove waterway near Bonita Springs, Florida, in 2010 and a brackish canal in Englewood, Florida, in 2015. Additional records from Tampa Bay (2007–2015) and Everglades City (2015), Florida, were documented with photography. Within the past few years, new records of *T. cystophora* from other localities, including Hawai'i, northern Australia, and the east coast of Florida, have been reported. This study represents the first published report of the species from the Gulf of Mexico. Several other species of cubozoans have been reported from the Gulf region, but the presence of three pedalia at each corner of the swimming bell (12 tentacles total), its small size up to 13 mm, and the presence of sexually dimorphic gonads, markedly distinguishes it from other cubozoan taxa.

Keywords: box jellyfish, *Tripedalia cystophora*, Tripedaliidae, Cubozoa, Gulf of Mexico, distribution

Tripedalia cystophora Conant, 1897 is a small (maximum bell height ca. 12–13 mm) box jellyfish (Cnidaria: Cubozoa) that inhabits tropical and subtropical mangrove systems (Conant 1897, Kramp 1961, Ekins & Gershwin 2014). Since its description from Jamaica in 1897, the

species has been the subject of several phylogenetic and developmental studies (Werner 1973, Laska-Mehnert 1985, Bentlage et al. 2010, Gurska & Garm 2014). Its elaborate copulation strategies have been well documented (Werner 1971, 1973, Stewart 1996), making it an excellent cubozoan for laboratory and field studies examining paternity, sexual selection, and the potential involvement of visually-guided behavior related to mat-

* Corresponding author: Robert.Lasley@myfwc.com

DOI: 10.2988/0006-324X-129.Q2.164

ing. *T. cystophora* is one of three species in the family Tripedaliidae (Order: Carybdeida), along with *T. binata* Moore, 1988 and *Copula sivickisi* (Stiasny, 1926), all of which are sexually dimorphic and produce spermatophores (Lewis & Long 2005, Bentlage et al. 2010, Bentlage & Lewis 2012, Straehler-Pohl et al. 2014). While spermatophore transfer from male to female is well documented in *T. cystophora* and *C. sivickisi*, (Werner 1973, Lewis & Long 2005, Underwood et al. 2013), details remain unclear for *T. binata*. Furthermore, *T. cystophora* has long been used as a model for elucidating the biomechanics, physiology, and evolution of the complex lens-eyes characteristic of all box jellyfish (Nilsson et al. 2005, Garm et al. 2007, Garm & Mori 2009). The most recent study of this species used a genomics approach to investigate cubozoan opsin diversification (Liegertová et al. 2015). Despite extensive research on these various aspects of the species' biology and the description of two types of nematocysts (Orellana & Collins 2011), the venom components of the nematocysts are yet to be described. Unlike other box jellyfish that are notorious for painful and sometimes debilitating stings (Russell et al. 1984, Brinkman & Burnell 2009, Bentlage et al. 2010), there are no documented human stings by *T. cystophora* despite the extensive number of publications on this species. Furthermore, *T. cystophora* has never been listed among dangerous box jellyfish in the scientific literature (e.g. Williamson et al. 1996), suggesting that its sting is mild to humans.

The species has a widespread, pantropical to subtropical distribution. Within 40 years of the original description from Jamaica (Conant 1897), *T. cystophora* was reported from the eastern Pacific and Coral Triangle (Stiasny 1926, Thiel 1936). In the past 4 years, the reported range of the species has expanded to include records from the east coast of

Florida (Orellana & Collins 2011), northern Australia, (Ekins & Gershwin 2014), and Hawai'i (Crow et al. 2015). Ekins and Gershwin (2014) reviewed the geographic distribution of the species, listing various Atlantic, Pacific, and Indian ocean localities and, like prior authors, suggested that these reports represent recent introductions rather than oversight of earlier extensive surveys conducted in those localities. Our new record of the species is the first in the Gulf of Mexico Large Ecoregion, an area differing from neighboring regions (e.g., Caribbean) based on its distinct biota, genetic breaks between it and bordering regions, extensive freshwater drainage, currents, and other abiotic factors (Kumpf et al. 1999, Felder & Camp 2009, Fautin et al. 2010). Furthermore, the northernmost records for *T. cystophora*, presented herein, lie on the fringe between the Northern Gulf and Floridian subregions, which is the northern latitudinal limit of mangroves and a transition to oyster reef and saltmarsh dominated habitats (Fautin et al. 2010). Our report therefore likely represents the northernmost limit of *T. cystophora* in the Gulf of Mexico and perhaps the Atlantic Ocean.

Materials and Methods

The specimens were identified as *Tripedalia cystophora* based primarily on: 1) the presence of four groups of tentacular pedalia, one at each corner of the bell; 2) three pedalia per group; and 3) each pedaliu with one tentacle (Conant 1897).

In September 2010, numerous *T. cystophora* medusae were noticed swarming near red mangroves (*Rhizophora mangle*) in a waterway adjacent to Estero Bay near Bonita Springs, Florida. Twelve exemplars (bell height 5.5–13 mm), including males and females, were collected by a resident and delivered to the Florida Gulf Coast University's Vester Field Station in Bonita

Springs, where they were held in a small aquarium for observation. The samples were separated into three lots: 2 lots were prepared in 5% formalin/seawater solution and 1 lot in 95% ethanol for future molecular studies. Material was deposited into the National Museum of Natural History, Smithsonian Institution (USNM). The following year, a large number of medusae were seen swarming at the Vester Field Station in August 2011 (Lasley et al. 2016).

On July 15, 2015, six *T. cystophora* medusae (bell height 4.8–9.5 mm) were collected in a brackish canal parallel to East 3rd Street, Englewood, Florida (Fig. 1). The canal feeds into Rock Creek ca. 1.5 km from the creek's mouth at Lemon Bay. Specimens were caught near *Rhizophora mangle* roots roughly one hour before sunset (20h25; with low tide (–0.1 ft) at 20h47). One exemplar (Fig. 1A) was determined to be female based on the presence of large eggs and embryos circulating in the gastrovascular cavity. The other specimens appeared to be post-spawned adults given their mature size but lack of gonads, so the sex could not be verified (Fig. 1B–F). After live transport to the Fish and Wildlife Research Institute, St. Petersburg, Florida, each specimen was photographed (Figs. 1, 2), measured, preserved in 10% formalin/seawater solution, and deposited into the Institute's marine collection (FSBCI). Prior to preservation in formalin, one tentacle from each specimen was removed and stored in 95% EtOH for future molecular study. Preserved specimens from Englewood were examined using a Zeiss Discovery V8 SMZ-168 stereoscope with manual 8x zoom. Live color photos were taken using

a Nikon D7100 DSLR with 60 mm AF Micro-Nikkor f2.8D and the Nikon R1C1 strobe package. Specimens were photographed from above in a shallow aquarium with a black velvet background. Measurements of bell height and bell width are provided below under material examined in the following format (bell height × bell width).

Additional specimens were observed but not collected at Smacks Bayou (2007) and between Cockroach Bay and Anna Maria Island (2011), Tampa Bay, Florida (Sheri Parks pers. obs.); and Everglades City, Florida (2015; including identifiable females) (Anders Garm pers. comm.). Specimens from Smacks Bayou, St. Petersburg and Everglades City, Florida were photographed, the images providing reliable identification of the specimens as *T. cystophora* (Lasley et al. 2016).

Systematics

Phylum Cnidaria Verrill, 1865

Class Cubozoa Werner, 1973

Order Carybdeida Gegenbaur, 1857


Family Tripedaliidae Conant, 1897

Genus *Tripedalia* Conant, 1897

Tripedalia cystophora Conant, 1897

Material examined.—12 medusae: 1 gravid female (5.5 mm × 7.7 mm), gastrovascular cavity filled with planulae, suggesting recent (within days) fertilization, 1 unsexed (9.5 mm × 13 mm) in formalin (USNM 1204759); 3 unsexed in 95% ethanol (USNM 1204760); 4 mature females (8.33 × 12 mm – 10 × 13 mm), 2 mature males (9 × 12.6 mm), 1 unsexed (10 mm × 14 mm) in formalin (USNM

Fig. 1. *Tripedalia cystophora*. Lateral view of live specimens. A, ripe female (7.8 mm × 9.2 mm) (FSBCI 129558, 1); B, sex undetermined (9.5 mm × 11.4 mm) (FSBCI 129558, 2); C, sex undetermined (8.1 mm × 9.5 mm) (FSBCI 129558, 3); D, sex undetermined (4.8 mm × 6.5 mm) (FSBCI 129558, 4); E, sex undetermined (7.6 mm × 9.4 mm) (FSBCI 129558, 5); F, sex undetermined (5.2 mm × 7.9 mm) (FSBCI 129558, 6). →






Fig. 2. *Tripedalia cystophora*. A–C, ventral, lateral, and dorsal views, respectively of live specimen, sex undetermined (9.5 mm \times 11.4 mm) (FSBCI 129558, 2).

1205451) – collected from red mangroves near Bonita Springs, Florida, Gulf of Mexico (26°22'53"N 81°50'08"W), USA, coll. C. Ritter, 7 September 2010, daylight hours.

6 medusae: 1 female, 5 sex undetermined (4.8 \times 6.5 mm – 9.5 \times 11.4 mm) (FSBCI 129558, 1–6) – collected from red mangroves, in canal parallel to East 3rd Street, Englewood, Florida, Gulf of Mexico (26.941419, –82.335328), USA, coll. M. Vorkapic, R. Lasley, 15 July 2015.

Range.—*Tripedalia cystophora* occurs in tropical and subtropical mangrove habitats between 32°N and 28°S latitude. In the Atlantic, the species has been reported from Jamaica, Puerto Rico, the Bahamas, and the east coast of Brazil (Morandini & Marques 1997); the east coast of Florida (Orellana & Collins 2011); and the Gulf of Mexico (this report). Pacific Ocean specimens have been reported from Japan (Uchida 1970); Costa Rica (Rodríguez-Sáenz & Segura-Puertas 2009); northern Australia, Thailand, and Indonesia (Ekins & Gershwin 2014); and Hawai'i (Crow et al. 2015). Indian Ocean specimens have been reported from Aldabra Atoll, Seychelles (Ekins & Gershwin 2014).

Conclusions

Of the seven box jellyfish species reported from the Gulf of Mexico, including those in this study, three have been well documented as long-term inhabitants: 1) *Alatina alata* (Reynaud, 1830) (= *Carybdea alata*) (Larson et al. 1991, Lewis et al. 2013); 2) *Tamoya* sp. (*T.* cf. *haplonema*) Müller, 1859 (Phillips & Burke 1970, Collins et al. 2011); and 3) *Chiropsalmus quadrumanus* Müller, 1859 (Müller 1859, Guest 1959, Phillips & Burke 1970, Gómez Aguirre 1986). In addition to these well-documented taxa, *Alatina grandis* (Agassiz & Mayer, 1902) was reported (as *Carybdea* var. *alata*) a single time in the Gulf of Mexico (Graham 1998). Furthermore, Hedgepeth's (1954) species checklist of the Gulf of Mexico includes *Carybdea aurifera* Mayer, 1900, and newer checklists (Segura-Puertas et al. 2003, Segura-Puertas et al. 2009) include *Carybdea marsupialis* (Linnaeus, 1758). However, the former corresponds to collections by Mayer (1900) from Tortugas and the latter to reports by Suárez-Morales et al. (1999) from the western Caribbean Sea.

All species reported from the Gulf of Mexico, including *T. cystophora*, belong to

the order Carybdeida except *Chiropsalmus quadrumanus*. *Tripedalia cystophora* is easily distinguished from the other species of Carybdeida by the presence of three pedalia/tentacles at each interradial corner of the bell (versus one in other Gulf species of Carybdeida) (Fig. 2) (Conant 1897, Mayer 1910). *Chiropsalmus quadrumanus* (Chirodromida: Chiropsalmidae), superficially resembles *T. cystophora* in having multiple tentacles at each corner, but *C. quadrumanus* possesses only a single pedalum per corner that is divided into 5–9 branches, each giving rise to a single tentacle (Mayer 1910, Calder 2009).

Conant's 1897 original description of *T. cystophora* does not allude to any type specimens. Furthermore, the legendary F. S. Conant described *T. cystophora* as a species new to science while he was Director at the Johns Hopkins Marine Laboratory at Kingston Harbor, Jamaica (Conant 1897), but there is no indication that Conant deposited any cubozoan specimens into the affiliated Institute of Jamaica Museum (IJM) (Muka 2014) prior to his untimely death in 1897. A Johns Hopkins circulatory published in memory of Conant and his work (Brooks 1900) states that Conant "brought home specimens", but it does not say where they were deposited (Brooks 1900:2). No invertebrate zoology collection currently exists at Johns Hopkins University, bringing the existence of the specimens further into question. Nonetheless, given the completeness of Conant's original description and accompanying illustration of the specimens, there is no doubt about the species' identity as *T. cystophora*, negating the need to erect a neotype in accordance with ICZN Article 75.2 (ICZN). Finally, it has been suggested that the species has been extirpated from its type locality of Kingston, Jamaica (Ekins & Gershwin 2014).

This new record of *T. cystophora* extends the species' known range to the Gulf of Mexico as far north as Tampa Bay. Together with the movement of

mangrove systems into higher latitudes (Saintilan et al. 2014) and the possible increase in frequency of jellyfish blooms (Condon et al. 2013), the range expansion of *T. cystophora* and its possible link to climate change and human introduction is worthy of investigation. Molodtsova (2009) reviewed the long history of medusozoan studies in the Gulf. Despite being a small species that can easily blend in with mangrove infrastructure and murky water (Stewart 1996, Garm et al. 2012), the fact that *T. cystophora* was previously never reported in the Gulf of Mexico, despite decades of extensive medusozoan sampling, casts doubt on their prior occurrence in the region. Correspondingly, recent new records of the species in Australia (2014) and Hawaii (2015) are assumed to have resulted from human introductions (Ekins & Gershwin 2014, Crow et al. 2015), and the likelihood of anthropogenic influence and cryptogenesis (introduction from an obscure locality) was addressed in a report on the recent appearance of *Copula sivickisi*, another member of the Tripedaliidae, in the western Atlantic (Bennett et al. 2011), which had been considered strictly an Indo-Pacific species (Lewis et al. 2008). The recently published *T. cystophora* genome, along with future large-scale morphological and molecular studies, could serve as a tool to elucidate the mechanism of dispersal resulting in the apparent recent global range expansion of the species.

Acknowledgments

Thanks to Michael and Joy Vorkapic for collecting the Englewood specimens and notifying the Fish and Wildlife Research Institute. We also acknowledge Chris Ritter for collecting the Bonita Springs samples and notifying the Vester Field Station. Thanks also to Carol Davis for notifying the first author after receiving the report. Thank you to Geoff Keel,

Smithsonian National Museum of Natural History, Department of Invertebrate Zoology Collections Lab Manager, for assistance with cataloguing specimens. We thank also Jan Boyett and Bland Crowder for their help editing the manuscript.

Literature Cited

- Agassiz, A., & A. G. Mayer. 1902. Reports on the scientific results of the expedition to the tropical Pacific, in charge of Alexander Agassiz, by the U.S. Fish Commission steamer "Albatross," from August, 1899, to March, 1900, Commander Jefferson F. Moser, U.S.N., commanding. III. The Medusae. *Memoirs of the Museum of Comparative Zoology* (26):139–76.
- Bennett, C. L., M. D. Nicholson, & R. B. Erdman. 2011. First record of *Copula sivickisi* (Cnidaria: Cubozoa) in the tropical western Atlantic Ocean: an enigmatic occurrence from San Salvador Island, Bahamas. Pp. 33–37 in C. Tepper & R. Shaklee eds. *Proceedings of the 14th Symposium on the Natural History of the Bahamas*.
- Bentlage, B., P. Cartwright, A. A. Yanagihara, C. Lewis, G. S. Richards, & A. G. Collins. 2010. Evolution of box jellyfish (Cnidaria: Cubozoa), a group of highly toxic invertebrates. *Proceedings of the Royal Society B: Biological Sciences* 277(1680):493–501.
- Bentlage, B., & C. Lewis. 2012. An illustrated key and synopsis of the families and genera of carybdeid box jellyfishes (Cnidaria: Cubozoa: Carybdeida), with emphasis on the "Irukandji family" (Carukiidae). *Journal of Natural History* 46(41–42):2595–2620.
- Bentlage, B., A. Peterson, & P. Cartwright. 2009. Inferring distributions of chirodroid box-jellyfishes (Cnidaria: Cubozoa) in geographic and ecological space using ecological niche modeling. *Marine Ecology Program Series* 384:121–133.
- Brinkman, D. L., & J. N. Burnell. 2009. Biochemical and molecular characterisation of cubozoan protein toxins. *Toxicon* 54:1162–1173.
- Brooks, W. K. 1900. The expedition to Jamaica in the summer of 1897. Notes from the Biological Laboratory. John Hopkins University Circulars. In: Brooks WK (ed) *University Circulars*. The Johns Hopkins press, 352 pp.
- Calder DR. 2009. Cubozoan and scyphozoan jellyfishes of the Carolinian biogeographic province, southeastern USA. Ontario (Canada): Royal Ontario Museum.
- Collins, A. G., B. Bentlage, W. Gillan, T. H. Lynn, A. C. Morandini, & A. C. Marques. 2011. Naming the Bonaire banded box jelly, *Tamoya ohboya*, n. sp. (Cnidaria: Cubozoa: Carybdeida: Tamoyidae). *Zootaxa* 68:53–68.
- Conant, F. S. 1897. Notes on the Cubomedusae. *Johns Hopkins University Circ.* 132:8–10.
- Conant, F.S. 1898. The Cubomedusae. By Franklin Story Conant. A memorial volume. Memoirs from the biological laboratory of the Johns Hopkins university, IV, 1. The Johns Hopkins press, Baltimore
- Condon, R. H., C. M. Duarte, K. A. Pitt, K. L. Robinson, C. H. Lucas, K. R. Sutherland, H. W. Mianzan, M. Bogeberg, J. E. Purcell, M. B. Decker, & S. I. Uye. 2013. Recurrent jellyfish blooms are a consequence of global oscillations. *Proceedings of the National Academy of Sciences* 110(3): 1000–1005.
- Crow, G. L., L. M. Chiaverano, J. Crites, M. A. Khrarov, & B. S. Holland. 2015. Box jellyfish (Cubozoa: Carybdeida) in Hawaiian waters, and the first record of *Tripedalia cystophora* in Hawai'i. Pp. 93–108 in N. L. Evenhuis & J. T. Carlton, eds. *Lucius G. Eldredge III Memorial Volume: Tribute to a Polymath*. Bishop Museum Bulletin in Zoology. Bishop Museum Press: Honolulu, 177 pp.
- Ekins, M., & L. Gershwin. 2014. First record of the Caribbean box jellyfish *Tripedalia cystophora* in Australian waters. *Marine Biodiversity Records* 7:1–5.
- Felder, D. L., & D. K. Camp, eds. 2009. *Gulf of Mexico Origin, Waters, and Biota*. Vol. 1, Biodiversity. College Station, Tex.: Texas A&M University Press. 1384 pp.
- Fautin, D., P. Dalton, L. S. Incze, J. A. C. Leong, C. Pautzke, A. Rosenberg, P. Sandifer, G. Sedberry, J. W. Tunnell Jr, I. Abbott, R. E. Brainard, M. Brodeur, L. Eldredge, M. Feldman, F. Moretzsohn, P. S. Vroom, M. Wainstein, and N. Wolf. 2010. An overview of marine biodiversity in United States waters. *PLoS One* 5(8): e11914.
- Garm, A., J. Bielecki, R. Petie, & D. E. Nilsson. 2012. Opposite patterns of diurnal activity in the box jellyfish *Tripedalia cystophora* and *Copula sivickisi*. *The Biological Bulletin* 222(1):35–45.
- Garm, A., & S. Mori. 2009. Multiple photoreceptor systems control the swim pacemaker activity in box jellyfish. *The Journal of Experimental Biology* 212(24):3951–60.
- Garm, A., M. O'Connor, L. Parkefelt, & D. E. Nilsson. 2007. Visually guided obstacle avoidance in the box jellyfish *Tripedalia cystophora*

- and *Chiropsella bronzie*. The Journal of Experimental Biology 210(20):3616–23.
- Gegenbaur, C. 1857. Versuch eines Systemes der Medusen, mit Beschreibung neuer oder wenig gekannter Formen; zugleich ein Beitrag zur Kenntniss der Fauna des Mittelmeeres. Zeitschrift für wissenschaftliche Zoologie (8):202–273.
- Gómez-Aguirre, S. 1986. Ocurrencia de *Chiropsalmus quadrumanus* (Scyphozoa: Cubomedusae) en la Laguna de Términos. Anales del Instituto de Ciencias del Mar y Limnología 57(Serie Zoología 2):227–34.
- Graham, M. 1989. Short papers and notes: first report of *Carybdea alata* var. *grandis* (Reynaud 1830) (Cnidaria: Cubozoa) from the Gulf of Mexico. Gulf of Mexico Science 1:28–30.
- Guest, W. C. 1959. The occurrence of the jellyfish *Chiropsalmus quadrumanus* in Matagorda Bay, Texas. Bulletin of Marine Science 9(1):79–83.
- Gurska, D., & A. Garm. 2014. Cell proliferation in cubozoan jellyfish *Tripedalia cystophora* and *Alatina moseri*. PLoS One. 9(7): e102628. doi: 10.1371/journal.pone.0102628
- Hedgepeth, J. W. 1954. Scyphozoa. Fisheries Bulletin, U.S.A. 55: 277–278.
- International Commission on Zoological Nomenclature. Article 75.2. <http://www.nhm.ac.uk/hosted-sites/iczn/code/index.jsp?nfv=true&article=75>. Accessed 1 February 2016.
- Kramp, P. L. 1961. Synopsis of the medusae of the world. Journal of the Marine Biological Association of the United Kingdom 40:7–382.
- Kumpf H., K. Steidinger, & K. Sherman, eds. 1999. The Gulf of Mexico Large Marine Ecosystem: Assessment, Sustainability, and Management. Malden, Mass: Blackwell Science. 704 pp.
- Larson, R. J., C. E. Mills, & G. R. Harbison. 1991. Western Atlantic midwater hydrozoan and scyphozoan medusae: in situ studies using manned submersibles. Hydrobiologia 216(1):311–317.
- Laska-Mehnert, G. 1985. Cytologische Veränderungen während der Metamorphose des Cubopolypen *Tripedalia cystophora* (Cubozoa, Carybdeidae) in die Meduse. Helgoländer Meeresunters 39(1):129–64.
- Lasley R., C. L. Ames, R. Erdman, S. Parks, and A. Collins. 2016. Photo records of Gulf of Mexico *Tripedalia cystophora*. https://figshare.com/articles/Photo_records_of_Gulf_of_Mexico_Tripedalia_cystophora/3459626 (last accessed on 23 June 2016)
- Lewis, C., B. Bentlage, A. Yanagihara, W. Gillan, J. Van Blerk, D. P. Keil, A. E. Bely, & A. G. Collins. 2013. Redescription of *Alatina alata* (Reynaud, 1830) (Cnidaria: Cubozoa) from Bonaire, Dutch Caribbean. Zootaxa 3737(4):473–487.
- Lewis, C., S. Kubota, A. Esteves Migotto, & A. G. Collins. 2008. Sexually dimorphic cubomedusa *Carybdea sivickisi* (Cnidaria: Cubozoa) in Seto, Wakayama, Japan. Publications of the Seto Marine Biological Laboratory 40(5):1–8.
- Lewis, C., & T. A. F. Long. 2005. Courtship and reproduction in *Carybdea sivickisi* (Cnidaria: Cubozoa). Marine Biology 147(2):477–83.
- Liegertová, M., J. Pergner, I. Kozmiková, P. Fabian, A. R. Pombinho, H. Strnad, J. Pačes, Č. Vlček, P. Bartůněk, & Z. Kozmik. 2015. Cubozoan genome illuminates functional diversification of opsins and photoreceptor evolution. Scientific reports 5:1–18.
- Linnaeus, C. 1758. Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, Tomus I. Editio decima, reformata. pp. [1–4], 1–824. Holmiae. (Salvius). I. Editio dec. Holmiae. Laurentii Salvii.
- Mayer, A. G. 1900. Some medusae from the Tortugas, Florida. Bulletin of the Museum of Comparative Zoology Harvard 37:13–82.
- Moore, S. J. 1988. A new species of cubomedusan (Cubozoa: Cnidaria) from northern Australia. The Beagle: Records of the Museums and Art Galleries of the Northern Territory 5(1):1–4.
- Morandini, A. C., & A. C. Marques. 1997. “Morbakka” syndrome: first report of envenomation by Cubozoa (Cnidaria) in Brazil. Anals of VII Colacmar (Boletim de Resumos Expandidos VII Colacmar), vol. 2. Instituto Oceanográfico da Universidade de Sao Paulo, Santos, pp. 188–189.
- Muka, S. K. 2014. Working at water’s edge: life sciences at American Marine Stations, 1880–1930. Publicly Accessible Penn Dissertations. Paper 1379.
- Müller, F. 1859. Zwei neue Quallen von Santa Catharina. Abhandlungen der Naturforschenden Gesellschaft zu Halle Bd. 5:1–12.
- Nilsson, D. E., L. Gislén, M. M. Coates, C. Skogh, & A. Garm. 2005. Advanced optics in a jellyfish eye. Nature 435(7039):201–205.
- Orellana, E. R., & A. G. Collins. 2011. First report of the box jellyfish *Tripedalia cystophora* (Cubozoa: Tripedaliidae) in the continental USA, from Lake Wyman, Boca Raton, Florida. Marine Biodiversity Records 4:1–3.

- Phillips, P. J., & W. D. Burke. 1970. The occurrence of sea wasps (cubomedusae) in Mississippi Sound and the northern Gulf of Mexico. *Bulletin of Marine Science* 20(4):853–859.
- Reynaud, M. 1830. *Carybdea alata* n. sp. P. 95, pl. 33 in R. P. Lesson, ed., *Centurie Zoologique*. Levrault: Paris (France).
- Rodríguez-Sáenz K., & L. Segura-Puertas. 2009. Hydrozoa, Scyphozoa, and Cubozoa (Medusozoa). Pp. 143–149 in I. S. Wehrtmann & J. Cortés, ed., *Marine biodiversity of Costa Rica, Central America*. Dordrecht: Springer Netherlands, 533 pp.
- Russell, F. E., H. Gonzalez, S. B. Dobson, & J. A. Coats. 1984. Bibliography of venomous and poisonous marine animals and their toxins. *Attending Staff Association of LAC/USC Medical Center Los Angeles CA*.
- Saintilan, N., N. C. Wilson, K. Rogers, A. Rajkaran, & K. W. Krauss. 2014. Mangrove expansion and salt marsh decline at mangrove poleward limits. *Global Change Biology* 20(1):147–157.
- Segura-Puertas, L., E. Suárez-Morales, & L. Celis. 2003. A checklist of the medusae (Hydrozoa, Scyphozoa and Cubozoa) of Mexico. *Zootaxa* 15:1–15.
- Stewart, S. E. 1996. Field behavior of *Tripedalia cystophora* (class Cubozoa). *Marine and Freshwater Behaviour and Physiology* 27:175–88.
- Stiasny, G. 1926. Über einige Scyphomedusen von Puerto Galera, Mindoro (Philippinen). *Zoologische Mededelingen* 9:239–48.
- Straehler-Pohl I, A. Garm, & A. C. Morandini. 2014. Sexual dimorphism in Tripedaliidae (Conant 1897) (Cnidaria, Cubozoa, Carybdeida). *Zootaxa* 3785(4):533–49.
- Suárez-Morales, E., Segura-Puertas, L. & Gasca, R. (1999b) A survey of the reef-related medusa (Cnidaria) community in the western Caribbean Sea. *Gulf Research Reports* 11: 23–31.
- Thiel, M. 1936. Scyphomedusae: Cubomedusae. H.G. Bronns Klassen und Ordnungen des Tierreichs. Mitteilungen aus dem Museum für Naturkd. Berlin Zool Museum und Inst für Spez Zool. 2:179–308.
- Uchida, T. 1970. Revision of Japanese Cubomedusae. *Publications of the Seto Marine Biological Laboratory* 17(5):289–97.
- Underwood, A., S. Taylor, & J. E. Seymour. 2013. Range extension of the Cubozoan, *Tripedalia binata* Moore (Cnidaria : Carybdeida : Carybdeidea) from far north Queensland, Australia. *Memoirs of the Queensland Museum, Nature* 56(2):607–13.
- Verrill, A. E. 1865. Synopsis of the polyps and corals of the North Pacific Exploring Expedition under Commodore C. Ringgold and Captain John Rodgers, U. S. N. *Commun Essex Institute* 4:145–152.
- Werner, B. 1971. Life cycle of *Tripedalia cystophora* Conant (Cubomedusae). *Nature* 232(5312):582–583.
- Werner, B. 1973. Spermatozeugmen und Paarungsverhalten bei *Tripedalia cystophora* (Cubomedusae). *Marine Biology* 18(3):212–217.

Editor: Rick Hochberg