

13

Evaluating Indicators of Seagrass Stress to Light

Patrick D. Biber, Hans W. Paerl, Charles L. Gallegos, and W. Judson Kenworthy

CONTENTS

Introduction	193
Bio-Optical Modeling	194
Plant Physiology	195
Methods	196
Calibrating a Bio-Optical Model	196
Light-Stress Experiments	197
Results	198
Water Quality Monitoring; North River, North Carolina	198
Light Stress Experiment	200
Discussion	203
Water Quality Stress Indicators	203
Plant Physiology Indicators	204
Conclusions	205
Acknowledgments	206
References	207

Introduction

Estuaries and coastal waters are highly productive, ecologically and societally valuable ecosystems. They are under increasing stress from both anthropogenic factors, such as nutrient enrichment and sedimentation, and altered frequencies and intensities of natural disturbances arising across many scales, from inputs to a watershed to those wrought by global climate change. Seagrasses are often dominant primary producers that can play a central role in the stability, nursery function, biogeochemical cycling, and trophodynamics of coastal ecosystems and, as such, are important for sustaining a broad spectrum of organisms (Thayer et al., 1984; Hemminga and Duarte, 2000). For example, they stabilize sediments, which are easily resuspended if the plants are lost, resulting in increased and prolonged turbidity that reduces available light reaching the seafloor. For these reasons, seagrasses are widely recognized as “barometers” of estuarine water quality, being perhaps the most parsimonious integrator of estuarine water quality throughout the range of their current and historic distribution (Dennison et al., 1993). Thriving seagrass communities signal a productive, diverse, and biogeochemically and trophically well-coupled coastal ecosystem (Harlin and Thorne-Miller, 1981; Thayer et al., 1984; Fonseca et al., 1998; Hauxwell et al., 2001). Accordingly, the presence or absence of seagrass is a useful measure of estuarine condition, but reliance on presence/absence as an indicator implicitly requires significant degradation of estuarine water quality (Zimmerman et al., 1991; Short and Wyllie-Echeverria, 1996). By focusing on seagrass decline, we are restricted to detecting conditions when water quality is already so degraded that there is virtually no time for corrective actions. Therefore, early detection of sublethal stress thresholds in seagrass plants is crucial for effective conservation of this resource.

The role of seagrasses as indicators of estuarine condition, particularly decreased water clarity, was proposed in the early 1990s (Kenworthy and Haunert, 1991; Neckles, 1994). Dennison et al. (1993) concluded that seagrasses were potentially sensitive indicators of declining water quality because of their high light requirements (15 to 25% surface irradiance) compared to those of other aquatic primary producers, such as macroalgae and benthic microalgae, with much lower light requirements (Markager and Sand-Jensen, 1992, 1996; Agusti et al., 1994). To develop predictive indicators of estuarine suitability for seagrasses, both water-quality-driven stressors and the whole-plant integrated responses should be assessed. Potential predictive indicators need to respond clearly and reliably to abiotic factors that cause suboptimal seagrass growth (e.g., light limitation) and should come from a suite of approaches over a range of hierarchical levels.

Bio-Optical Modeling

Provided that all other environmental parameters are suitable, the light environment during the growing season is the most important abiotic factor determining survival of seagrasses (Moore et al., 1997; Batiuk et al., 2000; Dixon, 2000a). Light attenuation by the water column is a major variable related to seagrass decline (Dennison and Alberte, 1982, 1985; Bulthuis, 1983). Another is epiphytic light attenuation, which Dixon (2000b) found to be more important in the subtropical waters of Florida. Low light levels, below some minimum physiological requirement (typically 15 to 25% of incident surface light = I_0) may result in a loss of seagrasses. Light is attenuated down the water column resulting in less light available at the bottom (I_b) than at the surface (I_0) by factors including:

1. Turbidity, expressed as total suspended particulate matter (SPM)
2. Phytoplankton, which both absorb and scatter light, expressed in chlorophyll concentration (chl a)
3. Colored dissolved organic matter (CDOM) leaching from decaying vegetation
4. Macroalgae and epiphytic microalgae that grow on the seagrass, which are usually more problematic when eutrophication is taking place (Harlin and Thorne-Miller, 1981; Hauxwell et al., 2001)

Light attenuation (Figure 13.1A) can be expressed as PLW (percent light through water) or the combined effects of contributions due to turbidity, chlorophyll, and color, resulting in significantly reduced light with increasing depth. Additional light attenuation can occur at the leaf surface due to epiphyte fouling, PLL (percent light at leaf), which occurs primarily under heavily eutrophic conditions (Batiuk et al., 2000). However, this may not always be the case; for example, Dixon (2000b) found light attenuation due to epiphytes averaging 34% of all attenuation of light in lower Tampa Bay where chl a has an annual average that is less than 5 mg m^{-3} (not heavily eutrophic). One of the goals of our research has been to refine a bio-optical water quality model (Gallegos, 1994, 2001) by determining the importance of SPM, chl a , and CDOM on light attenuation to seagrasses in North Carolina during different seasons.

Graphically, the results of the bio-optical model are shown as in Figure 13.1B, where two components of light attenuation (chl a , TSS) are presented on the x and y axes. The third constituent, CDOM, can be plotted on the z axis on a three-dimensional (3D) plot to show all three components of attenuation. Median concentrations for one water quality sample are plotted on this graph and compared to a minimum light water quality requirement for a given depth (for simplicity, depicted as a line of constant attenuation), which is calculated using a radiative-transfer model and knowledge of seagrass species light requirements. Target minimum water clarity requirements for seagrass survival are found at the intersection of vectors perpendicular to the axes or the origin from the median sample concentration. The target concentrations in this figure suggest that both TSS and chl a need to be reduced to meet the minimum light requirements of this seagrass species.

Determining the minimum light requirements for a given species requires either monitoring and identifying the deep edge of existing seagrass beds within an estuary, or an experimental approach where the minimum light requirements for survival are determined over a period of time. An understanding of

Evaluating Indicators of Seagrass Stress to Light




FIGURE 13.1 (A) Conceptual diagram of light attenuation down the water column (PLW), and the relative contributions of turbidity, chlorophyll *a*, and color to attenuation. Additional attenuation can occur at the leaf surface due to epiphyte fouling (PLL), which occurs primarily under heavily eutrophic conditions. (B) Graphical representation of the bio-optical model. Components of light attenuation in the water column are presented along the axes. Median concentrations for one sample are plotted on this graph and compared to a minimum-light water quality requirement for a given depth (line of constant attenuation). Target minimum water clarity requirements for seagrass survival are found at the intersection of vectors perpendicular to the axes or the origin from the median sample concentration. The target concentrations in this figure suggest that both TSS and chl *a* need to be reduced to meet the minimum light requirements of this seagrass species.

the physiological processes occurring within plants subject to chronic light limitation is, therefore, important when applying the bio-optical model.

Plant Physiology

For a seagrass-dominated ecosystem to be judged sustainable, the seagrasses must be vital and capable of long-term persistence by growth and reproduction. We are focusing on a common set of photosynthetic and physiological metrics that measure seagrass condition and may provide *early warning* of plant stress or demise, before extensive loss of seagrass occurs.

A promising approach needing further investigation is chlorophyll fluorescence of photosystem II, also known as P680, where a molecule of water is split to form O_2 and a reductant in the light reaction of photosynthesis (Stryer, 1981). To measure this, we utilized a photosynthetic efficiency analyzer (PEA) that was developed to measure the health of crops (e.g., Critchley and Smilie, 1981; Havaux and Lannoye, 1983; Bowyer et al., 1991; Filiault and Stier, 1999), and because all plants utilize the same fundamental processes in photosynthesis, this instrument can also be used on marine macrophytes.

The PEA consists of a computer and a photo-emitter/sensor unit. Clips are supplied with the apparatus, which are attached to a plant leaf and serve to occlude all but a small area of the leaf needed for photosynthetic testing. A small area of the plant leaf is dark-adapted with the clip and then a shutter built into the clip is opened, exposing the leaf area under the clip to high-intensity light provided by nearly monochromatic light emitting diodes (LEDs) located in the photo-emitter/sensor unit. The chlorophyll in the dark-adapted area of the leaf fluoresces and the PEA measures this. The initial fluorescence (F_o) and the maximal fluorescence (F_m) are recorded and the difference between the maximal and initial fluorescence levels ($F_m - F_o$) is called the variable fluorescence (F_v). The PEA computer calculates the ratio F_v/F_m , or photosynthetic efficiency; the greater the fluorescence per unit incoming light, the higher the efficiency of the photosystem, which equates with a plant under low physiological stress. Conversely, low F_v/F_m indicates a plant under stress (Schulze and Caldwell, 1990; Krause and Weis, 1991; Rohacek and Bartak, 1999).

This approach has been used to measure acute stress in seagrasses, such as desiccation (Adams and Bates, 1994; Bjork et al., 1999), temperature, or salinity shifts (Ralph et al., 1998; Ralph, 1999), and even changes in ambient light over short time durations (Beer and Bjork, 2000; Major and Dunton, 2002). However, to our knowledge this technique has not been evaluated in seagrass plants subject to chronic stress, such as light limitation arising from reduced water clarity.

There are two components to our research that are discussed here:

1. Monitoring water quality to calibrate a bio-optical model and using this modeling approach to understand water quality changes that result in low-light stress
2. Determining seagrass photophysiological indicators as a measure of individual plant responses to light stress

Methods

Calibrating a Bio-Optical Model

To refine the bio-optical model as an indicator for seagrass habitat suitability in the mid-Atlantic region of the eastern coastal United States, we have initiated a monthly water-sampling program in North River, North Carolina, an estuary that contains sizable seagrass beds along a gradient in water clarity. We collect water samples at nine stations in North River, North Carolina (Figure 13.2). Additionally, at each station we profile water quality with YSI® 6600 multiparameter probes (temperature, salinity, dissolved oxygen, or DO, pH, turbidity, and chlorophyll fluorescence), as well as collecting light attenuation data using a LICOR® 4π sensor. The light data are used to calculate attenuation coefficients to compare with Secchi disk readings and laboratory-measured optical properties of the water samples. Water samples are analyzed for total absorption (referenced to pure water) and scattering coefficients using the methods described in Gallegos (1994, 2001).




FIGURE 13.2 Map of nine sampling locations in North River, a small estuary in the southern Pamlico Sound region of North Carolina. Seagrass beds are indicated by shaded polygons; 1 and 2 m depths are indicated by the dashed contour lines.

Light-Stress Experiments

A light gradient (Table 13.1) was created in a large indoor tank under controlled environmental (light, air temperature) conditions to determine minimum integrated light requirements for seagrass survival. A bank of overhead Halogen® lights of increasing wattage provided the different light treatments. Lights were placed on a timer to create a 12:12 light:dark (L:D) environment. Water was provided as a flow-through system and temperature was monitored at 30 min intervals at both ends of the light gradient. Water quality data were collected with a YSI 6000 multiparameter water quality probe. Two seagrass species were tested in 2003: *Zostera marina* (eelgrass) seedlings were grown for 20 weeks (27 January to 6 June 2003), and *Halodule wrightii* (shoalgrass) plants were grown for 16 weeks (8 July to 27 October 2003) in a range of light intensities (Table 13.1).

Two tubs (Rubbermaid® 2951) were planted with nine *Z. marina* seedlings or nine *H. wrightii* plants (3 × 3 arrangement) and placed into each of the seven light gradient treatments after allowing 10 days recovery from transplant stress. Initial morphological measurements were taken on the day the plants were placed into the light gradient tank and weekly thereafter until the experiment was terminated. Biweekly, mature leaves from a random subset of plants were collected to measure width and destructive analysis of tissue constituents, including chlorophyll extraction and tissue nutrient content (C:N ratio) using standard analytical methods.

For the weekly chlorophyll fluorescence measurements, the second leaf on the terminal apical shoot was clipped about half way along the adaxial side as per the recommendations of Durako and Kunzelman (2002). Dark adaptation of the clipped portion was 10 min, and the PEA recorded chlorophyll fluorescence from the leaf for 1 s. Typically, the maximum fluorescence in green plants occurs within the first second after exposure to the high-intensity LED lights (Srivastava et al., 1999).

TABLE 13.1
Seven Irradiance Treatments Used for Indoor Light Experiment^a

Instantaneous Flux ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	Integrated Daily Flux ($\text{mol m}^{-2} \text{d}^{-1}$)	Hsat (h)	Hcomp (h)
<i>Zostera marina</i>			
277.03	11.97	5.49	11.54
161.49	6.98	0.50	6.54
69.80	3.02	-3.46	2.58
33.29	1.44	-5.04	1.01
10.93	0.47	-6.01	0.04
4.21	0.18	-6.30	-0.25
0.00	0.00	-6.48	-0.43
<i>Halodule wrightii</i>			
248.20	10.72	4.24	10.29
95.91	4.14	-2.34	3.71
59.96	2.59	-3.89	2.16
31.11	1.34	-5.14	0.91
6.90	0.60	-5.88	0.16
2.96	0.26	-6.22	-0.18
0.00	0.00	-6.48	-0.43

^a Expressed as instantaneous irradiance, integrated daily irradiance for a 12-h day, number of hours (Hsat) that irradiance exceeded a saturation intensity ($150 \mu\text{mol m}^{-2} \text{s}^{-1}$), and number of hours (Hcomp) that irradiance exceeded a compensation intensity ($10 \mu\text{mol m}^{-2} \text{s}^{-1}$) where photosynthesis balances respiration. Negative numbers indicate insufficient light to meet physiological requirements.

Results

Water Quality Monitoring; North River, North Carolina

Temperature followed an approximate sinusoidal pattern during the year with minimum temperatures in February and maximum temperatures in July and August (Figure 13.3). Shallower upriver stations had more rapid temperature changes than the tidally influenced downstream stations, which reflected oceanic water temperatures. Salinity was higher and more stable in the downstream, ocean-influenced section of North River than the upstream portion, which was more influenced by terrestrial runoff from surrounding salt-marsh drainage streams. Salinity dropped after heavy rainfall events, which was especially evident during the wet spring and summer of 2003 (Figure 13.3); this year was the wettest on record with 2337 mm (92 in.) recorded, 50% greater than average (National Weather Service data).

Chlorophyll concentrations (mg m^{-3}) were lowest during the cold winter months and increased during the spring and summer when phytoplankton were more abundant (Figure 13.3). Highest values were observed at the farthest upstream station corresponding with increased nitrogen concentrations measured at this station. High chlorophyll concentrations were also recorded during storm events and co-occurred with elevated turbidity, likely due to resuspension of benthic microalgae (Figure 13.3). Turbidity was similar at all stations during a given sampling date. In general, there was a trend of increasing turbidity from the downstream to the upstream stations (Figure 13.3). Turbidity was lower in winter than other seasons. The highest turbidity values were recorded at shallow upstream sites during storms (e.g., September 2002).

The effect of chlorophyll and turbidity on measured light penetration to the bottom at each of the nine stations was expressed as an attenuation coefficient (K_d), derived from the negative exponential




FIGURE 13.3 Selected results from our monthly water quality monitoring program in North River. Time-space contour plots of temperature, salinity, chl *a* concentration, and turbidity (NTU) measured monthly from September 2002 to September 2003. Sites are numbered from downstream (1) to most upstream (9). Insets are same data plotted as 3D wireframe plots.




FIGURE 13.4 The effects of chlorophyll and turbidity on measured light attenuation with depth (K_d) at each of the nine stations. Filled symbols are samples collected during floodtide, open symbols are ebb tide samples. For seagrass survival K_d should not exceed 1.5, i.e., 22% of surface light reaches the bottom (dotted bar). Seagrass beds are found in North River between stations 2 and 6. The gray arrow indicates relative abundance of seagrasses.

Beer-Lambert function. Higher values of K_d indicate more light attenuation (i.e., more turbid water). K_d values increased with distance from the Beaufort Inlet at station 1 to the highest values found at station 9, where chl a and turbidity were also typically highest. During 2 months (September 2002 and March 2003), samples were collected during storm conditions (tropical storm and northeaster, respectively) with some of the highest attenuation values seen, indicating the importance of natural events in driving extreme values in this estuary. For seagrass survival to 1 m, K_d should not exceed 1.5 m^{-1} , i.e., 22% of surface light (I_0) reaching the bottom, as indicated by the dotted line (Figure 13.4). This condition was typically met at the stations near where seagrass are found in North River (stations 2 through 6). K_d values sufficient for seagrass survival were less frequently seen in the upstream stations.

Using our initial calibration values from the bio-optical model, monthly water quality conditions were plotted for North River sampling stations (Figure 13.5). Most samples fall in the region of acceptable light quantities (22% I_0) reaching 1 m depth, except for a few samples that were collected during storm conditions (September 2002 and March 2003). This indicates that water clarity in the North River is sufficient to support seagrasses to 1 m depth, and this agrees with observations made on the multidecadal persistence of seagrass beds in this estuary (since the early 1970s). Interestingly, the line of constant attenuation for the North River (solid line) allows much higher turbidity concentrations than the same light-at-depth requirement for Chesapeake Bay seagrasses (dashed line). The reason for this (based on initial results) is that per unit mass suspended sediments in the North River absorb and scatter much less light than those in the mesohaline Chesapeake Bay, due to differences in sediment properties related primarily to particle size (and perhaps also phytoplankton composition) between these regions.

Additionally, the North Carolina State estuarine water quality criteria of 40 mg m^{-3} chl a and 25 NTU turbidity (NC-DENR, 2003) are shown on the plot (Figure 13.5, black box). Almost all water quality samples from North River fall within the acceptable state criteria, and within the submerged aquatic vegetation (SAV) survival criteria based on the bio-optical model calibrated to North River conditions. There are combinations of turbidity and chlorophyll concentration permitted by state criteria that would not support seagrasses according to the bio-optical model (Figure 13.5, shaded triangle); interestingly, no water quality observations fell in this region.

The results from the bio-optical model can be corroborated by results from the permanent monitoring stations. Three locations in North River are designated as permanent stations, with continuous unattended water quality and light monitoring using YSI and LICOR sensors and data loggers. These locations are near stations 2, 6, and 9 in North River (Figure 13.2). Station 6 has been continuously monitored since September 2002. Integrated daily light ($\text{mol m}^{-2} \text{ d}^{-1}$) received at two sensor depths, 1 and 1.5 m, was




FIGURE 13.5 Median monthly water quality conditions plotted for North River, based on the bio-optical model. Symbol tints indicate season (gray shadow = winter, light gray = spring and fall, black = summer), symbol shape indicates location within North River. Additionally, the North Carolina State estuarine water-quality criteria of 40 mg m^{-3} chl *a* and 25 NTU are shown (rectangle outlined with thick lines). The bio-optical model was used to predict water quality criteria for survival of seagrass to 1 m depth in Chesapeake Bay (dashed line) and North River (solid line). The different slopes reflect regional differences in sediments and phytoplankton.

averaged for each month and the number of hours that irradiance exceeded the photosynthesis saturation intensity ($150 \mu\text{mol m}^{-2} \text{s}^{-1}$) for *Z. marina* (Dennison and Alberte, 1982) calculated (Table 13.2). Light available for photosynthesis was high in December to February (clear water) and June to July (strong solar irradiance) even though summertime water clarity was much reduced compared to the winter condition; i.e., stronger solar irradiance may compensate for higher chl *a* concentrations in the summer.

Light Stress Experiment

Comparing and contrasting the results of the light gradient experiments using *Z. marina* seedlings and *H. wrightii* plants showed similar responses by both species to light limitation stress. Both instantaneous and integrated (12 h) irradiance fluxes during the two experiments are given in Table 13.1, as well as integrated irradiance greater than the saturation irradiance ($\sim 150 \mu\text{mol}$) and compensation irradiance ($\sim 10 \mu\text{mol}$) for *Z. marina* (Dennison and Alberte, 1982) and *H. wrightii*.

For both species, all but the two greatest light treatments resulted in a deficit of light to saturate photosynthesis, ultimately resulting in mortality of many of the plants. This result indicates that irradiances consistently lower than the saturation threshold will ultimately result in plant death. Observations made during weekly measurements suggested that mortality in response to light limitation occurred from the base of the plant upward, with the meristem becoming nonviable about 1 to 2 weeks before the leaves were apparently dead (*Z. marina*) or broke off at the sheath (*H. wrightii*). This observed lag-time of 1 to 2 weeks may be critical in monitoring seagrass mortality under natural situations where light limitation is the cause of mortality.

During the *Z. marina* experiment, water temperatures increased from 10 to 28°C . The upper favorable temperature for this species is around 25°C (Thayer et al., 1984), which was attained in May (Figure 13.6A). This high-temperature stress resulted in reduced growth, as seen by a reduction in leaf area for the remainder of the experiment. In contrast, temperatures started to decline during the *H. wrightii*

TABLE 13.2

Light Data from 1- and 1.5-m-Deep Sensors Collected at Station 6 from 2000 to 2003^a

2002–2003 Month	mol m ⁻² d ⁻¹		Hsat	
	1 m	1.5 m	1 m	1.5 m
Sept	6.22	1.57	5.00	0.14
Oct	5.01	1.61	3.42	0.45
Nov	7.70	4.94	4.87	4.13
Dec	12.02	6.02	6.29	4.35
Jan	12.29	5.16	7.00	4.15
Feb	12.95	5.99	6.72	4.39
Mar	8.06	3.03	4.72	1.66
Apr	8.81	3.55	5.24	2.36
May	7.66	2.36	4.33	1.15
Jun	12.90	4.74	7.17	3.41
Jul	9.36	5.34	6.27	3.85
Aug	6.99	3.61	4.65	2.31
Sept	2.98	1.49	1.86	0.36
Mean	8.35	3.79	5.04	2.52

^a Expressed as mean daily integrated irradiance (mol m⁻² d⁻¹) each month, and the number of hours (Hsat) that irradiance exceeded a saturation intensity (150 μmol m⁻² s⁻¹). The growth period of *Z. marina* is from December to June, and the growth period for *H. wrightii* is from May to November. Both integrated and Hsat values are about twice as great at 1 m vs. 1.5 m depths.




FIGURE 13.6 Water temperatures recorded in the indoor tank during the two light gradient experiments. Water temperatures were recorded at 30 min time intervals in the highest light (tank 1) and lowest light (tank 6) treatments.

experiment (Figure 13.6B). They remained at 28°C until early September; then temperatures declined to 18°C by the end of the experiment.

Plant growth and photosynthesis responses for *Z. marina* and *H. wrightii* in the light gradient experiment showed some similarities as well as differences with respect to the minimum integrated light requirements for survival. Both single shoot *Z. marina* seedlings and two shoot *H. wrightii* plants were grown for the duration of their growth season. For *Z. marina*, no branching or spatial expansion was observed in any of the seven irradiance treatments. Of the 117 seedlings, 5 flowered (4.3%) all at irradiances of 33 μmol or higher (Figure 13.7A). In *H. wrightii* plants, branching or spatial expansion was observed in the highest irradiance treatment only (Figure 13.7B). Shoot loss and mortality were

observed at irradiances of 60 μmol or lower in this species. These integrated light thresholds indicate that tropical *H. wrightii* plants have somewhat higher minimum light requirements for survival than temperate *Z. marina* seedlings.

The length of the longest leaf on each surviving seedling was measured as a proxy for canopy height in both species. In *Z. marina*, leaf growth (linear extension) occurred only in the two highest light levels




FIGURE 13.7 Plant growth and photosynthesis responses for *Z. marina* and *H. wrightii* in the seven irradiance treatments (units of $\mu\text{mol m}^{-2} \text{s}^{-1}$) in the light gradient experiment. Selected variables shown are mean number of shoots per plant ($n = 18$), length of longest leaf measured as a proxy for canopy height, leaf area (single surface), and photosynthesis yield ratio (F_v/F_m).

(irradiance > saturation). Early mortality (leaf length = 0) was observed in the two lowest light levels with evidence of plant stress by weeks 3 and 5, respectively, about half the time to mortality (Figure 13.7C). In *H. wrightii*, leaf length was maintained only at the two highest light levels (irradiance $\geq 95 \mu\text{mol}$). Early mortality was observed in the lowest light level with evidence of plant stress by weeks 3 to 4, about half the time to mortality of all the plants in this treatment (Figure 13.7D).

Leaf area (single surface) was calculated for the strap-like leaves by multiplying maximum length by average width in both species. Width was measured on alternative sampling dates on a random subset of three to five plants with linear interpolation for nonsampled dates. Area was primarily influenced by leaf length, as width remained approximately constant within a given light treatment (Figure 13.7E and F).

Zostera marina seedling photosynthesis was measured with an OptiSciences® OS-30 PEA (Plant Efficiency Analyzer), and *H. wrightii* photosynthesis was measured with a Hansatech PEA that had increased sensitivity, an important consideration given the narrow leaves of this species. Photosynthetic yield, expressed as the ratio F_v/F_m , ranged from 0.6 to 0.75 in healthy *Z. marina* seedlings (Figure 13.7G) and from 0.65 to 0.78 in healthy *H. wrightii* plants (Figure 13.7H). In both species F_v/F_m fell below 0.5 in plants that were dead or dying. No significant difference in F_v/F_m was observed in those plants remaining alive across all seven light treatments, indicating acclimation of the photosynthetic apparatus to the ambient light field and temperature conditions. In summary, the yield ratio (F_v/F_m) is not a sensitive measure of chronic stress conditions, either light or temperature, to which tropical *H. wrightii* and temperate *Z. marina* can acclimate.

Discussion

Seagrass is an important estuarine habitat that is declining globally (Green and Short, 2003). Much of this decline can be attributed to decreased light availability to the plants because of reductions in water clarity due to degrading water quality (Dennison et al., 1993; Hauxwell et al., 2001). Seagrasses have relatively high light requirements compared to other marine primary producers and so are more susceptible to low light stress (Kenworthy and Hauxwell, 1991; Kenworthy and Fonseca, 1996; Dixon, 2000a). For this reason, they have been proposed as indicators of estuarine change (Dennison et al., 1993). However, it is desirable to have an indicator that provides an early warning of potential seagrass demise, rather than waiting until after the fact.

Water quality criteria, particularly those related to the optical water quality (i.e., water clarity) needed for the survival and growth of seagrasses, have been the subject of considerable research (Kenworthy and Hauxwell, 1991; Neckles, 1994; Gallegos and Kenworthy, 1996; Kenworthy and Fonseca, 1996; Longstaff et al., 1999; Zimmerman, 2003). A general conclusion of those workshops and research programs was that water column clarity needs to be greatly increased to provide light conditions suitable for the survival of most seagrasses (Kenworthy and Fonseca, 1996; Moore et al., 1996; Batiuk et al., 2000).

Water Quality Stress Indicators

We are developing a water quality stress indicator using a bio-optical model of water clarity that can identify harmful trends in water quality and make explicit what is causing the increased light attenuation. This will enable early preventative management actions to be taken to remediate water quality before it becomes critically limiting to seagrass survival, resulting in loss of this important habitat.

The bio-optical model is useful because it permits us to determine the relative contributions of the different water quality parameters to light attenuation at different positions in the estuary. This is a very important step in managing water quality as it makes explicit the relative contribution of suspended particles, phytoplankton, and color to light attenuation. From this knowledge, target reductions or target loading rates for one or more of these components of water quality can be set to achieve a desired water clarity. Comparing light attenuation calculated at the deep survival limit of seagrasses with water quality concentrations measured there allows the determination of ranges of water clarity that permit expansion

or cause contraction of the seagrass bed. Basing the model on inherent optical properties has the advantage that extrapolation beyond the range of water quality concentrations encountered during model development is possible, because the absorption and scattering coefficients are linearly related to the relevant water quality concentrations (Figure 13.1B). Such an exercise can be used to determine the availability of light at the edge of the seagrass bed in response to hypothetical scenarios, such as accelerated eutrophication resulting from increased nutrient loading in the watershed. This bio-optical model has already been calibrated to conditions typical for Chesapeake Bay, Maryland (Gallegos, 2001) and Indian River Lagoon, Florida (Gallegos and Kenworthy, 1996), and recently North River, North Carolina (Biber and Gallegos, unpubl. data), all estuaries with significant seagrass habitats.

Based on the initial results of the bio-optical model for North River seagrass habitats, we can conclude that, even though water quality appears to be adequate to sustain stable seagrass beds in this estuary, the long-term survival of these mostly shallow seagrass communities in North River is still uncertain, especially when considering the likelihood of sea-level rise projected to occur within the next century. More stringent water clarity criteria with respect to seagrass habitats should be adopted by the state to ensure seagrasses receive adequate light. Similar results have been found by researchers in Chesapeake Bay, where large declines in submerged aquatic vegetation (SAV) have been ongoing and are almost impossible to reverse (Orth and Moore, 1983; Dennison et al., 1993; Batiuk et al., 2000; U.S. EPA, 2003).

Seagrass communities in North Carolina present a unique situation at the overlap of temperate and tropical biogeographic regions resulting in the coexistence of *Z. marina* (temperate, winter dominant) and *H. wrightii* (tropical, summer dominant), allowing research on the competitive interactions between these two widely distributed seagrasses (Green and Short, 2003). Results are likely to be conservative as plants may already be experiencing some stress related to their presence at extremes of distribution. Further differences exist in that *Z. marina* reproduces sexually with annual seedling recruitment being a significant contribution to the population, whereas *H. wrightii* has never been observed to reproduce sexually in North Carolina, dispersing instead by clonal growth and asexual fragmentation (Thayer et al., 1984; Ferguson et al., 1993). These different reproductive strategies affect the dispersal ability and population recovery strategies after severe stress events, with potentially important repercussions in the face of possible global climate change scenarios. Because of the extreme importance of annual seedling recruitment to eelgrass population dynamics, we suggest that future research be focused on understanding the stress tolerance of this life stage, so that relevant differences in survival ability can be incorporated into management plans.

We suggest that the higher irradiance penetrating through the water column in the winter months, when water is clearer, may be a critical window of opportunity for seedling growth for *Z. marina*. Similarly for *H. wrightii*, high light levels, due to strong solar irradiances, in early summer may be critical for this species to grow rapidly early in the season, even though summer water clarity is much reduced compared to the winter condition. Similar critical windows of increased irradiance, due to higher water clarity, have been identified in the polyhaline Chesapeake Bay, from 1 March to 31 May, the early part of growing season for *Z. marina* (U.S. EPA, 2003). Management implications of this are that better water quality needs to be present during these critical periods.

Plant Physiology Indicators

Most current approaches do not address the integrated light requirements of seagrass, focusing instead on “instantaneous” measures of irradiance flux and seagrass photosynthetic rates, e.g., photosynthesis–irradiance curves (Dennison and Alberte, 1982, 1985; Goodman et al., 1995; Zimmerman et al., 1995; Bintz and Nixon, 2001). This approach implicitly ignores cumulative stress effects, so that important questions regarding how the duration of exposure or the frequency of exposure to a given level of environmental degradation might influence survival of the seagrasses are overlooked. Only recently has the frequency and duration of stressful conditions started to be investigated for the survival of seagrass (Dunton and Tomasko, 1994; Onuf, 1996; Moore et al., 1997). It is this whole-plant integrated response that is biologically and ecologically relevant to seagrass management. This is also the scale at which we can link water quality to seagrass photophysiology.

Recent technological advances and research into plant photosynthesis have resulted in the commercial production of instruments that can measure chlorophyll fluorescence to detect sublethal stress thresholds *in situ*, even underwater (e.g., Diving PAM, Walz, Germany). This technique has opened a new vista for seagrass physiologists, allowing rapid, nondestructive measurements of plant photosynthetic performance (e.g., Ralph and Burchett, 1995; Beer et al., 1998; Ralph et al., 1998; Bjork et al., 1999; Beer and Bjork, 2000; Durako and Kunzelman, 2002).

In our light-limitation experiments, seagrass chlorophyll induction kinetics were measured using a PEA. Contrary to expectation, the photosynthetic efficiency (F_v/F_m) indicated that both *Z. marina* and *H. wrightii* adapt their photosystem to the ambient light and temperature conditions within less than a week, as indicated by a near constant F_v/F_m ratio. Because of this adaptation, declines in the F_v/F_m ratio in low light treatments failed to precede losses in shoot number (see Figure 13.7A and G). F_v/F_m ratio alone is, therefore, unlikely to be a suitable tool to measure sublethal chronic stress responses in seagrass because of its invariance in the face of photosystem change. However, other ratios and indices derived from chlorophyll fluorescence induction kinetics that can be collected by PEA show more promise.

Chlorophyll fluorescence induction kinetics are extremely rich in terms of quality and quantity of different information (Strasser et al., 1995) and can only be measured by a PEA with a 1- μ s sampling rate or higher, but not a pulse amplitude modulated (PAM) fluorometer, which has only a 30- μ s sampling rate, as the lower sampling frequency misses the very early induction dynamics (Rohacek and Bartak, 1999). Because the induction kinetics are determined by both the physiological state of a plant and the ambient and past physical and chemical environmental conditions, they can be used to detect and predict sublethal stress thresholds not apparent in F_v/F_m . Strasser et al. (1995, 1996) have developed a method (JIP-test analysis) to further analyze the data on chlorophyll fluorescence induction kinetics gathered by PEA into ratios and indices of photosynthetic performance of the photosystem.

Upon illumination chlorophyll exhibits a fast fluorescence rise from an initial fluorescence intensity, F_o , to a maximal intensity, $F_p (= F_m)$. Between these two extremes, the fluorescence intensity usually shows two intermediate steps: F_j at about 2 ms, and F_i at about 30 ms (Strasser and Govindjee, 1992a,b; Strasser et al., 1995) followed by F_p at about 300 ms. The labeling of these steps follows an alphabetic order, from the initial (F_o) to the final (F_p) part of the transient. Based on the information inherent in the O-J-I-P fluorescence transient, a test has been developed (called "JIP-test" after the steps of the transient), which can be used as a tool for rapid screening of many samples providing additional information about the structure, conformation, and function of their photosynthetic apparatus (Strasser and Strasser, 1995; Strasser et al., 1996).

As seagrass leaves age, many of the JIP-test ratios and indices change (Figure 13.8). Standardized methods for collecting PEA data for JIP-test analysis from seagrasses have not yet been adopted, in part because this is a new approach, and in part because of lack of knowledge on the variability inherent in natural, healthy seagrass populations (Figure 13.9). To better understand seagrass photophysiology, PEA and JIP-test will need to be further investigated to determine the inherent variability of these ratios and indices in seagrass species under stress (e.g., light limitation). We will continue to test the responses of *Z. marina* and *H. wrightii* to chronic light stress using PEA and JIP-test analyses to determine whether this may provide potentially powerful early warning of imminent demise of seagrass plants.

The ability to combine both water quality stressors, i.e., the light environment linked explicitly to constituents of attenuation, with plant physiological responses to light stress forms the basis of an integrative indicator of water quality that will permit evaluation and determination of the suitability of water quality in an estuary for continued seagrass sustainability.

Conclusions

Once the optical properties of the bio-optical model are calibrated to regional conditions, the next step is to utilize the seagrass ecophysiological information to set light requirement thresholds for seagrass plant survival, growth, and reproduction. Each of these three thresholds will be successively greater than the previous one because of the increased cumulative light resource requirement by these plants to achieve the successive stages of development; i.e., resources (light) required for growth, a net increase




FIGURE 13.8 Results of PEA-derived JIP-test analyses on *H. wrightii* leaves of different ages, L1 = youngest, L3 = oldest, Brn and Blk denote dead leaves in increasing stages of decomposition. (A) Relationship between the relative number of active PSII reaction centers and the performance index. (B) Relationship between the probability that an electron will be transported beyond quinone A and the performance index.




FIGURE 13.9 Results of JIP-test on *Z. marina* plants at 2 months and 1 year of age. (A) Relationship between the relative number of active PSII reaction centers per unit area and the performance index. (B) Relationship between the probability that an electron will be transported beyond quinone A and the performance index.

in biomass, will be greater than those required for survival (maintenance metabolism or no net increase in mass) and similarly for growth to reproductive maturity. The final stage of this process will be to use the bio-optical model as a tool to forecast the results of possible altered water quality scenarios and determine the impacts of these changes on existing seagrass habitats.

Acknowledgments

Grateful thanks to the cadre of persons who were instrumental in assisting with data collection especially in the field, including Christine Addison, Craig Bonn, John Burke, John Brewer, Priscilla Delano, Teri Denault, Don Field, Nikki Fogarty, Tom Gallo, Kamille Hammerstrom, Kar Howe, Loretta Leist, Amit Malhotra, Ryan McQuinn, Joe Purifoy, Karen Rossignol, Stopher Slade, Amy Uhrin, Rich Weaver, Paula Whitfield, and Pam Wyrick. Additional support was provided by the Center for Coastal Fisheries and Habitat Research, National Centers for Coastal Ocean Science (NCCOS), National Ocean Service (NOS), National Oceanic and Atmospheric Administration (NOAA), Beaufort, North Carolina. Thanks to Glen Thursby for providing the tools to do the JIP-test analyses. This research was funded by EPA-STAR Grants (R82867701) to the ACE-INC EaGLes Center (www.aceinc.org) and (R82868401) to the Atlantic Slope Consortium EaGLes Center (www.asc.psu.edu).

References

- Adams, J. B. and G. C. Bates. 1994. The tolerance to desiccation of the submerged macrophytes *Ruppia cirrhosa* (Petagna) Grande and *Zostera capensis* Setchell. *Journal of Experimental Marine Biology and Ecology* 183:53–62.
- Agusti, E. S., S. Enriquez, H. Frost-Christensen, K. Sand-Jensen, and C. M. Duarte. 1994. Light-harvesting among photosynthetic organisms. *Functional Ecology* 8:273–279.
- Batiuk, R. A., P. Bergstrom, M. Kemp, E. Koch, L. Murray, J. C. Stevenson, R. Bartleson, V. Carter, N. B. Rybicki, J. M. Landwehr, C. Gallegos, L. Karrh, M. Naylor, D. Wilcox, K. A. Moore, S. Ailstock, and M. Teichberg. 2000. Chesapeake Bay Submerged Aquatic Vegetation Water Quality and Habitat-based Requirements and Restoration Targets: A Second Technical Synthesis. Chesapeake Bay Program, U.S. EPA, Washington, D.C., 231 pp.
- Beer, S. and M. Bjork. 2000. Measuring rates of photosynthesis of two tropical seagrass by pulse amplitude modulated (PAM) fluorometry. *Aquatic Botany* 66:69–76.
- Beer, S., B. Vilenkin, A. Weil, M. Veste, L. Susel, and A. Eshel. 1998. Measuring photosynthetic rates in seagrasses by pulse amplitude modulated (PAM) fluorometry. *Marine Ecology Progress Series* 174:293–300.
- Bintz, J. C. and S. W. Nixon. 2001. Responses of eelgrass, *Zostera marina*, seedlings to reduced light. *Marine Ecology Progress Series* 223:133–141.
- Bjork, M., J. Uku, A. Weil, and S. Beer. 1999. Photosynthetic tolerances of desiccation of tropical intertidal seagrasses. *Marine Ecology Progress Series* 191:121–126.
- Bowyer, J. R., P. Camilleri, and W. F. J. Vermaas. 1991. Photosystem II and its interaction with herbicides. In *Herbicides. Topics in Photosynthesis*, Vol. 10, N. R. Baker and M. P. Percival (eds.). Elsevier, Amsterdam, pp. 27–85.
- Bulthuis, D. A. 1983. Effects of in situ light reduction on density and growth of the seagrass *Heterozostera tamanica* in Western Port, Victoria, Australia. *Journal of Experimental Marine Biology and Ecology* 67:91–103.
- Critchley, C. and R. M. Smilie. 1981. Leaf chlorophyll fluorescence as an indicator of photoinhibition in *Cucumis sativus* L. *Australian Journal of Plant Physiology* 8:133–141.
- Dennison, W. C. and R. S. Alberte. 1982. Photosynthetic responses of *Zostera marina* to in situ manipulations of light intensity. *Oecologia* 55:137–144.
- Dennison, W. C. and R. S. Alberte. 1985. Role of daily light period in the depth distribution of *Zostera marina*. *Marine Ecology Progress Series* 25:51–61.
- Dennison, W. C., R. J. Orth, K. A. Moore, J. C. Stevenson, V. Carter, S. Kollar, P. W. Bergstrom, and R. A. Batiuk. 1993. Assessing water quality with submersed aquatic vegetation. *Bioscience* 43:86–94.
- Dixon, L. K. 2000a. Establishing light requirements for the seagrass *Thalassia testudinum*: an example from Tampa Bay, Florida. In *Seagrasses: Monitoring, Ecology, Physiology and Management*, S. A. Bortone (ed.). CRC Press, Boca Raton, FL, pp. 9–32.
- Dixon, L. K. 2000b. Light requirements of Tampa Bay Seagrasses: nutrient related issues still pending. In *Seagrass Management: It's Not Just Nutrients! Proceedings of a Symposium*, H. Greening (ed.). August 22–24, 2000. St. Petersburg, FL.
- Dunton, K. H. and D. A. Tomasko. 1994. In situ photosynthetic performance in the seagrass *Halodule wrightii* in a hypersaline subtropical lagoon. *Marine Ecology Progress Series* 107:281–293.
- Durako, M. J. and J. I. Kunzleman. 2002. Photosynthetic characteristics of *Thalassia testudinum* measured in situ by pulse-amplitude modulated (PAM) fluorometry: methodological and scale-based considerations. *Aquatic Botany* 73:173–185.
- Ferguson, R. L., B. T. Pawlak and L. L. Wood. 1993. Flowering of the seagrass *Halodule wrightii* in North Carolina. *Aquatic Botany* 46:91–98.
- Filiault, D. L. and J. C. Stier. 1999. The use of chlorophyll fluorescence in assessing the cold tolerance of three turfgrass species. *Wisconsin Turfgrass Research Report* 16:109–110.
- Fonseca, M. S., W. J. Kenworthy, and G. W. Thayer. 1998. Guidelines for the conservation and restoration of seagrasses in the United States and adjacent waters. NOAA Coastal Ocean Program Decision Analysis Series, No. 12. NOAA Coastal Ocean Office, Silver Spring, MD, 222 pp.

- Gallegos, C. L. 1994. Refining habitat requirements of submersed aquatic vegetation: role of optical models. *Estuaries* 17:187–199.
- Gallegos, C. L. 2001. Calculating optical water quality targets to restore and protect submersed aquatic vegetation: overcoming problems in partitioning the diffuse attenuation coefficient for photosynthetically active radiation. *Estuaries* 24:381–397.
- Gallegos, C. L. and J. W. Kenworthy. 1996. Seagrass depth limits in the Indian River Lagoon (Florida): application of an optical water quality model. *Estuarine and Coastal Shelf Science*, 42:267–288.
- Goodman, J. L., K. A. Moore, and W. C. Dennison. 1995. Photosynthetic responses of eelgrass, *Zostera marina*, to light and sediment sulfide in a shallow barrier island lagoon. *Aquatic Botany* 50:37–47.
- Green, E. P. and F. T. Short. 2003. *World Atlas of Seagrasses*. University of California Press, Berkeley, 320 pp.
- Harlin, M. M. and B. Thorne-Miller. 1981. Nutrient enrichment of seagrass beds in a Rhode Island coastal lagoon. *Marine Biology* 65:221–229.
- Hauxwell, J., J. Cebrian, C. Furlong, and I. Valiela. 2001. Macroalgal canopies contribute to eelgrass (*Zostera marina*) decline in temperate estuarine ecosystems. *Ecology* 82:1007–1022.
- Havaux, M. and R. Lannoye. 1983. Chlorophyll fluorescence induction: A sensitive indicator of water stress in maize plants. *Irrigation Science* 4:147–151.
- Hemminga, M. A. and C. M. Duarte. 2000. *Seagrass Ecology*. Cambridge University Press, Cambridge, U.K., 298 pp.
- Kenworthy, J. W. and M. S. Fonseca. 1996. Light requirements of seagrasses *Halodule wrightii* and *Syringodium filiforme* derived from the relationship between diffuse light attenuation and maximum depth distribution. *Estuaries* 19:740–750.
- Kenworthy, W. J. and D. E. Haunert. 1991. The Light Requirements of Seagrasses: Proceedings of a Workshop to Examine the Capability of Water Quality Criteria, Standards and Monitoring Programs to Protect Seagrasses. NOAA Technical Memorandum NMFS-SEFC-287, NOAA.
- Krause, G. H. and E. Weis. 1991. Chlorophyll fluorescence and photosynthesis: the basics. *Annual Review of Plant Physiology and Plant Molecular Biology* 42:313–349.
- Longstaff, B. J., N. R. Loneragan, M. J. O'Donohue, and W. C. Dennison. 1999. Effects of light deprivation on the survival and recovery of the seagrass *Halophila ovalis*. *Journal of Experimental Marine Biology and Ecology* 234:1–27.
- Major, K. M. and K. H. Dunton. 2002. Variations in light-harvesting characteristics of the seagrass *Thalassia testudinum*: evidence for photoacclimation. *Journal of Experimental Marine Biology and Ecology* 275:173–189.
- Markager, S. and K. Sand-Jensen. 1992. Light requirements and depth zonation of marine macroalgae. *Marine Ecology Progress Series* 88:83–92.
- Markager, S. and K. Sand-Jensen. 1996. Implications of thallus thickness for growth irradiance relationships of marine macroalgae. *European Journal of Phycology* 31:79–87.
- Moore, K. A., H. A. Neckles, and R. J. Orth. 1996. *Zostera marina* (eelgrass) growth and survival along a gradient of nutrients and turbidity in the lower Chesapeake Bay. *Marine Ecology Progress Series* 142:247–259.
- Moore, K. A., R. L. Wetzel, and R. J. Orth. 1997. Seasonal pulses of turbidity and their relations to eelgrass (*Zostera marina*) survival in an estuary. *Journal of Experimental Marine Biology and Ecology* 215:115–134.
- NC Department of Environment and Natural Resources. 2003. Classifications and Water Quality Standards. North Carolina Administrative Code: Section 15A NCAC 2B.0200.
- Neckles, H. A. 1994. Indicator Development: Seagrass Monitoring and Research in the Gulf of Mexico. EPA/620/R-94/029. U.S. EPA, ORD-ERL, Gulf Breeze, FL, 64 pp.
- Onuf, C. P. 1996. Seagrass responses to long-term light reduction by brown tide in upper Laguna Madre, Texas: distribution and biomass patterns. *Marine Ecology Progress Series* 138:219–231.
- Orth, R. J. and K. A. Moore. 1983. Chesapeake Bay: an unprecedented decline in submerged aquatic vegetation. *Science* 222:51–53.
- Ralph, P. J. 1999. Photosynthetic response of *Halophila ovalis* to combined environmental stress. *Aquatic Botany* 65:83–96.
- Ralph, P. J. and M. D. Burchett. 1995. Photosynthetic responses of the seagrass *Halophila ovalis* to high irradiance stress using chlorophyll *a* fluorescence. *Aquatic Botany* 51:55–66.

- Ralph, P. J., R. Gademann, and W. C. Dennison. 1998. *In situ* seagrass photosynthesis measured using a submersible, pulse-amplitude modulated fluorometer. *Marine Biology* 132:367–373.
- Rohacek, K. and M. Bartak. 1999. Techniques of modulated chlorophyll fluorescence: basic concepts, useful parameters, and some applications. *Photosynthetica* 37:339–363.
- Schulze, E. D. and M. M. Caldwell (eds.). 1990. *Ecophysiology of Photosynthesis*. Springer, Berlin, 432 pp.
- Short, F. T. and S. Wyllie-Echeverria. 1996. Natural and human-induced disturbance of seagrasses. *Environmental Conservation* 23:17–27.
- Srivastava, A., R. J. Strasser, and Govindjee. 1999. Greening of peas: parallel measurements of 77K emission spectra, OJIP chlorophyll *a* fluorescence transient, period four oscillation of the initial fluorescence level, delayed light emission and P700. *Photosynthetica* 37:365–392.
- Strasser, B. J. and R. J. Strasser. 1995. Measuring fast fluorescence transients to address environmental questions: the JIP-test. In *Photosynthesis: From Light to Biosphere*, Vol. 5, P. Mathis (ed.). Kluwer Academic, Dordrecht, pp. 977–980.
- Strasser, R. J. and Govindjee. 1992a. The Fo and the O-J-I-P fluorescence rise in higher plants and algae. In *Regulation of Chloroplast Biogenesis*, J. H. Argyroudi-Akoyunoglou (ed.). Plenum Press, New York, pp. 423–426.
- Strasser, R. J. and Govindjee. 1992b. On the O-J-I-P fluorescence transient in leaves and D1 mutants of *Chlamydomonas reinhardtii*. In *Research in Photosynthesis*, Vol. 4, N. Murata (ed.). Kluwer Academic, Dordrecht, pp. 29–32.
- Strasser, R. J., A. Srivastava, and Govindjee. 1995. Polyphasic chlorophyll *a* fluorescence transient in plants and cyanobacteria. *Photochemistry and Photobiology* 61:32–42.
- Strasser, R. J., P. Eggenberg, and B. J. Strasser. 1996. How to work without stress but with fluorescence. *Bulletin de la Société Royal des Sciences de Liège* 65:330–349.
- Stryer, L. 1981. *Biochemistry*, 2nd ed. W.H. Freeman, New York, 950 pp.
- Thayer, G. W., W. J. Kenworthy, and M. S. Fonseca. 1984. The Ecology of Eelgrass Meadows of the Atlantic Coast: A Community Profile. FWS/OBS-84/02. U.S. Fish and Wildlife Service, Washington, D.C., 147 pp.
- U.S. EPA. 2003. Ambient Water Quality Criteria for Dissolved Oxygen, Water Clarity and chlorophyll *a* for the Chesapeake Bay and Its Tidal Tributaries. EPA903-R-03-002. U.S. Environmental Protection Agency, National Service Center for Environmental Publications, Washington, D.C.
- Zimmerman, R. C. 2003. A bio-optical model of irradiance distribution and photosynthesis in seagrass canopies. *Limnology and Oceanography* 48:568–585.
- Zimmerman, R. C., J. L. Reguzzoni, S. Wyllie-Echeverria, M. Josselyn, and R. S. Alberte. 1991. Assessment of environmental suitability for growth of *Zostera marina* L. (eelgrass) in San Francisco Bay. *Aquatic Botany* 39:353–366.
- Zimmerman, R. C., J. L. Reguzzoni, and R. S. Alberte. 1995. Eelgrass, *Zostera marina*, transplants in San Francisco Bay: role of light availability on metabolism, growth and survival. *Aquatic Botany* 51:67–86.

