





Scientific note

**New record of the primnoid gorgonian  
*Primnoella delicatissima* Kükenthal, 1909 for Chilean waters**

(Anthozoa, Octocorallia, Primnoidae)

Verena Häussermann\*, Günter Försterra\* & Stephen Cairns\*\*

With 44 genera and approximately 275 species, the family Primnoidae constitutes the fourth largest octocoral family (Cairns & Bayer 2009), and also the family with the deepest living gorgonian species, which occur down to 5850 m (Cairns 2016a). Primnoids are typical deep-water species which are common on seamounts and deep-water coral banks. Due to a phenomenon called

deep-water emergence they can be found in diving depth in channels of Chilean Patagonia (Häussermann & Försterra 2009). In deep-water emergent species the depth ranges are enlarged; they occur from deep to shallow water (Försterra 2009). Eleven species of primnoids have been described from Chile (Van Ofwegen et al. 2009), five of which have been reported from shallow

**Fig. 1.** Sea whips from shallow water of Chilean Patagonia. **a.** *Primnoella chilensis*, entrance of Cahuelmo Fjord, Comau Fjord, 20 m; **b.** *Convexella magelhaenica*, Canal Pasaje, 12 m; **c.** *Primnoella delicatissima*, Caleta Tortuga, Chonos Archipelago, 12 m.

\* Verena Häussermann & Günter Försterra, Facultad de Recursos Naturales, Escuela de Ciencias del Mar, Pontificia Universidad Católica de Valparaíso, Avda. Brasil 2950, Valparaíso, Chile; and Huinay Scientific Field Station, Chile; e-mail: v.haussermann@gmail.com (corresponding author)

\*\* Stephen Cairns, Department of Invertebrate Zoology, National Museum of Natural History, W-329 Smithsonian Institution, P.O. Box 37012, Washington, D.C. 20560, U.S.A.

waters of Chilean Patagonia: *Primnoella chilensis* (Philippi, 1894), *Convexella magelhaenica* (Studer, 1879), *Thouarella koellikeri* (Wright & Studer, 1889), *Thouarella brucei* (Thomson & Richie, 1906) and one undescribed species of *Thouarella* (Van Ofwegen et al. 2009). Both *P. chilensis* and *Thouarella* spp. form so-called animal forests, providing habitat for other marine life (Försterra et al. in press). During the Huinay Fiordos (HF) expeditions, which hoped to inventory the marine life of Chilean Patagonia, some dozens of sea whips, which have the same colonial gross morphology and colour and thus under water resembled the common and widely distributed species *Primnoella chilensis* (41°S–54°S), have been collected and identified by SC. Most of these specimens belonged to the species *P. chilensis*, however, five specimens were identified as *Convexella magelhaenica* (12–27.5 m depth, 48°17'S–55°00'S; also see Van Ofwegen et al. 2009) and two recently collected specimens are *Primnoella delicatissima* Kükenthal, 1909. The latter specimens had been collected at Caleta Tortuga, Northern Patagonian Zone, at 13 m depth on April 4, 2014 (HF21, 45°19.518'S, 73°5.828'W) (USNM 1279400) and at Isla van der Meulen, Central Patagonian Zone, at 12 m depth on April 25, 2015 (HF24, 48°17.406'S, 74°20.150'W), respectively. *P. delicatissima* has been described from 55–103 m from off Brazil from Amapa to off Rio de Janeiro (Cairns 2006) and thus represents a new record for Chilean waters, although quite a disjunct distribution. The species was adequately described and figured by Cairns (2006), although in situ images were not provided.

The genus *Primnoella* consists of 11 species, occurring at depths of 8–1249 m in the western Atlantic, southwestern South America (including Chile), southeastern Australia, New Zealand, and Tasmania (Cairns 2016b). The genus has been recently reviewed by Cairns and Bayer (2009) and Cairns (2006, 2016b). Aside from *P. delicatissima*, two other species occur off Chile: *P. chilensis* (Philippi, 1894) and *P. scotiae* Thomson & Richie,

1906. The latter two species are easily distinguished by having many more scales in both the abaxial and outer lateral polyp rows, more polyps per whorl, and larger polyps.

**Acknowledgements.** Funding was provided through Fondocyt project nr 1131039 to VH and 1150843 to GF. This is publication nr 140 of Huinay Scientific Field Station.

## References

- Cairns, S. D. 2006. Studies on western Atlantic Octocorallia (Coelenterata: Anthozoa). Part 6. The genera *Primnoella* Gray, 1858; *Thouarella* Gray, 1870; *Dasystemella* Versluys, 1906. Proceedings of the Biological Society of Washington 119 (2): 161–194.
- 2016a. New abyssal Primnoidae (Anthozoa: Octocorallia) from the Clarion-Clipperton Fracture Zone, equatorial northeastern Pacific. Marine Biodiversity 46: 141–150.
- 2016b. The Marine Fauna of New Zealand: Primnoidae Octocorals (Anthozoa: Alcyonacea). Part 2. *Primnoella*, *Callozostrom*, *Metafannyella*, *Callogorgia*, *Fanelia* and other genera. NIWA Biodiversity Memoir 129: 1–129.
- & Bayer, F. M. 2009. A generic revision and phylogenetic analysis of the Primnoidea (Cnidaria: Octocorallia). Smithsonian Contributions to Zoology 629: 1–79.
- Försterra, G. 2009. Ecological and biogeographical aspects of the Chilean Fjord Region. In: Häussermann, V. & Försterra, G. (eds). Marine benthic fauna of Chilean Patagonia. Santiago de Chile (Nature in Focus).
- , Häussermann, V. & Laudien, J. in press. Animal forests in the Chilean fjords: discoveries, perspectives and threats in shallow and deep waters. In: Rossi, S. (ed.). Marine animal forests. Switzerland (Springer International Publishing).
- Häussermann, V. & Försterra, G. 2009 (eds). Marine benthic fauna of Chilean Patagonia. 1000 pp., Santiago de Chile (Nature in Focus).
- Kükenthal, W. 1909. Diagnosen neuer Alcyonarien (7. Mitteilung). Zoologischer Anzeiger 35(1/2): 46–53.
- Van Ofwegen, L., Breedy, O. & Cairns, S. 2009. Octocorals. In: Häussermann, V. & Försterra, G. 2009 (eds). Marine benthic fauna of Chilean Patagonia. 1000 pp., Santiago de Chile (Nature in Focus).

The annual subscription rate is 60.00 €, supplements are charged separately.  
For members of the "Freunde der Zoologischen Staatssammlung München" a reduced rate of 25.50 € applies.  
Journal exchange requests are welcome.  
Orders should be addressed to the  
Library of the Zoologische Staatssammlung München, Münchhausenstraße 21, 81247 München

### Author Guidelines

The journal considers manuscripts written in English, German, or French. Manuscripts should be submitted electronically as e-mail attachment to one of the managing editors (arthropods: Melzer@zsm.mwn.de; other invertebrates and vertebrates: Michael.Schroedl@zsm.mwn.de), or to spixiana@zsm.mwn.de. The organisation of the manuscript should be as follows: Title, Author(s), Address(es), Abstract (max 300 words), Text, Literature Cited, Appendix, Tables, Figure Captions, Figures. The text should be in Word or RTF format with a minimum of character and paragraph formatting (e.g. italics exclusively for genus and species names, and bold for headings). Identification keys should be formatted using tabulators. Avoid using blank spaces for adjusting text. Capitalization should be avoided except for abbreviations of depositories. The editors reserve the right to make minor textual changes that do not alter the author's meaning. The name, postal address, and e-mail of the corresponding author should be given on the first page. All papers must conform to the latest edition of the International Code of Zoological Nomenclature. The original author and date of a taxon name should be cited at least once (e.g. at first mention). If possible, representative material from contributions published in SPIXIANA should be deposited in the Zoologische Staatssammlung.

Digital photographs and other half-tone images should have high quality and a resolution of at least 400 dpi at the final image size. Line drawings should be scanned at a resolution of 1200 dpi as black and white (not greyscale) images and saved as TIFF (not JPG). Illustrations should be supplied in digital format, TIFF or PSD for raster images, EPS or Adobe Illustrator™ (together with externally linked images) for vector graphics. Labels and scale bars should be placed in separate layers (for PSD files) or added with a vector graphics program. Line-art and halftones must not be combined in a single grouped illustration. Printing of figures in colour is available at author's cost.

In-text citations should be in the form (Arratia et al. 1996, Glaw & Vences 2003, Meyer 2003). References should be listed at the end of the paper using the following format:

- Meyer, C. P. 2003. Molecular systematics of cowries (Gastropoda: Cypraeidae) and diversification patterns in the tropics. *Biological Journal of the Linnean Society* 79: 401-459.
- Glaw, F. & Vences, M. 2003. Introduction to Amphibians. Pp. 883-898 in: Goodman, S. M. & Benstead, J. P. (eds). *The Natural History of Madagascar*. Chicago and London (The University of Chicago Press).
- Arratia, G., Schultze, H.-P. & Viohl, G. 1996. *Mesozoic Fishes – Systematics and Paleoecology*. 604 pp., München (Pfeil).
- Noyes, J. S. 2003. Universal Chalcidoidea Database. World Wide Web electronic publication. [www.nhm.ac.uk/entomology/chalcidoids/](http://www.nhm.ac.uk/entomology/chalcidoids/) [accessed 05-Sep-2003]

The journal is published on a non-profit basis, editors and reviewers do not receive any payment. Authors will receive one copy of the issue with their article and the PDF of the article. Reprints are available at author's cost and must be ordered when proofs are returned.

## Contents

Schrödl, Michael: Winner of the Rolex Awards 2016: Congratulations to Vreni Häussermann!.....	145–146
Häussermann, Verena, Günter Försterra & Stephen Cairns: New record of the primnoid gorgonian <i>Primnoella delicatissima</i> Kükenthal, 1909 for Chilean waters (Anthozoa, Octocorallia, Primnoidae).....	147–148
Sikora, Bozena, Markus Unsöld & Maciej Skoracki: <i>Aulonastus paridus</i> spec. nov. – a parasite of two bird species of the genus <i>Melaniparus</i> in Kenya and Tanzania (Acari, Prostigmata, Syringophilidae).....	149–152
Ceseña, Feliza, Roland Meyer, Christian P. Mergl, Verena Häussermann, Günter Försterra, Kaitlin McConnell & Roland R. Melzer: Decapoda of the Huinay Fjords-expeditions to the Chilean fjords 2005–2014: Inventory, pictorial atlas and faunistic remarks (Crustacea, Malacostraca).....	153–198
Schrödl, Michael & Gerhard Haszprunar: Do we need Epitypes in Zoology?.....	199–201
Narayana, Sumantha & Deepak Apte: First record of <i>Epidendrium aureum</i> (Gastropoda, Epitoniidae) associated with <i>Tubastrea</i> corals (Scleractinia, Dendrophylliidae) from Lakshadweep, India (Gastropoda; Scleractinia).....	202
Sales, Licia, Vinicius Queiroz & Alvaro E. Migotto: Rediscovery of <i>Okenia impexa</i> Marcus 1957 on the Brazilian coast: the first image of a living specimen from Brazil (Mollusca, Nudibranchia).....	203–204
Scheers, Kevin: An updated checklist of the water beetles of Montenegro (Coleoptera, Hydradephaga).....	205–212
Hendrich, Lars & Michael Balke: A new epigeal <i>Paroster</i> Sharp, 1882 from coastal New South Wales, Australia (Coleoptera, Dytiscidae, Hydroporini).....	213–218
Bremer, Hans J.: Revision of the genus <i>Amarygmus</i> Dalman, 1823 and of related genera. Part LXXII. Amarygmmini of the Papuan faunal region: new subgenera of <i>Amarygmus</i> , new species of <i>Amarygmus</i> , a new synonym (Coleoptera, Tenebrionidae, Amarygmmini).....	219–246
Bremer, Hans J.: Species of <i>Corticeus</i> Piller & Mitterpacher from Peru (Coleoptera, Tenebrionidae, Hypophlaeini).....	247–251
Teruel, Rolando: Taxonomic identity of two <i>Pseudocellus</i> from Eastern Cuba (Arachnida, Ricinulei, Ricinoididae).....	252
Baehr, Martin: A new species of the genus <i>Tachyura</i> Motschulsky, subgenus <i>Tachyphanes</i> Jeannel from eastern Turkey (Coleoptera, Carabidae, Bembidiini, Tachyina).....	253–256
Baehr, Martin: Two new species of the genus <i>Adelotopus</i> Hope, 1834 from North Queensland, Australia. 15 <sup>th</sup> Supplement to the “Revision of the Pseudomorphinae of the Australian Region” (Coleoptera, Carabidae, Pseudomorphini).....	257–263
Glaw, Frank & Michael Franzen: On the maximum length of <i>Boa constrictor</i> (Serpentes, Boidae).....	264
Blick, Theo & Stefan Zaenker: Die österreichische Schneemücke, <i>Chionea austriaca</i> , erstmals in Deutschland (Diptera, Limoniidae).....	265–271
Prötzel, David, Kathrin Glaw, Julia Forster & Frank Glaw: Hibernation in tropical Madagascar? Unusual roosting sites of chameleons of the genus <i>Calumma</i> (Squamata, Chamaeleonidae).....	272
Ortiz, Antonio S., Rosa M. Rubio, Juan J. Guerrero, Manuel Garre & Axel Hausmann: Integrated taxonomy, phylogeography and conservation in the genus <i>Chelis</i> Rambur, [1866] in the Iberian Peninsula (Lepidoptera, Erebidae, Arctiinae).....	273–286
Hawlitshchek, Oliver, Cynthia Y. Wang-Claypool, Mark D. Scherz, Ludovic Montfort, Olivier Soumille & Frank Glaw: New size record of the snake genus <i>Liophidium</i> by the island endemic <i>L. mayottensis</i> (Squamata, Lamprophiidae).....	287–288