

# Arbuscular mycorrhizal fungal community composition is altered by long-term litter removal but not litter addition in a lowland tropical forest

Merlin Sheldrake<sup>1,2</sup>, Nicholas P. Rosenstock<sup>3,4</sup>, Daniel Revillini<sup>2,5</sup>, Pål Axel Olsson<sup>4</sup>, Scott Mangan<sup>2,6</sup>, Emma J. Sayer<sup>2,7</sup>, Håkan Wallander<sup>4</sup>, Benjamin L. Turner<sup>2</sup> and Edmund V. J. Tanner<sup>1</sup>

<sup>1</sup>Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK; <sup>2</sup>Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancón, Panama; <sup>3</sup>Center for Environmental and Climate Research, Lund University, Lund 22362, Sweden; <sup>4</sup>Department of Biology, Lund University, Lund 22362, Sweden; <sup>5</sup>Department of Biological Sciences, Northern Arizona University, PO BOX 5640, Flagstaff, AZ 86011, USA; <sup>6</sup>Department of Biology, Washington University in St Louis, St Louis, MO 63130, USA; <sup>7</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

## Summary

Author for correspondence:  
Merlin Sheldrake  
Tel: +44 207 794 9841  
Email: merlinsheldrake@gmail.com

Received: 20 September 2016  
Accepted: 14 November 2016

*New Phytologist* (2017)  
doi: 10.1111/nph.14384

**Key words:** 454-sequencing, arbuscular mycorrhizal (AM) fungi, litterfall, nutrient cycling, organic matter, tropical forest.

- Tropical forest productivity is sustained by the cycling of nutrients through decomposing organic matter. Arbuscular mycorrhizal (AM) fungi play a key role in the nutrition of tropical trees, yet there has been little experimental investigation into the role of AM fungi in nutrient cycling via decomposing organic material in tropical forests.
- We evaluated the responses of AM fungi in a long-term leaf litter addition and removal experiment in a tropical forest in Panama. We described AM fungal communities using 454-pyrosequencing, quantified the proportion of root length colonised by AM fungi using microscopy, and estimated AM fungal biomass using a lipid biomarker.
- AM fungal community composition was altered by litter removal but not litter addition. Root colonisation was substantially greater in the superficial organic layer compared with the mineral soil. Overall colonisation was lower in the litter removal treatment, which lacked an organic layer. There was no effect of litter manipulation on the concentration of the AM fungal lipid biomarker in the mineral soil.
- We hypothesise that reductions in organic matter brought about by litter removal may lead to AM fungi obtaining nutrients from recalcitrant organic or mineral sources in the soil, besides increasing fungal competition for progressively limited resources.

## Introduction

The productivity of most tropical forests is sustained by symbiotic associations between plants and arbuscular mycorrhizal (AM) fungi (Read, 1991; Alexander & Lee, 2005). AM fungi play crucial roles in nutrient cycling and are also major vectors of carbon (C) in the global C cycle (Johnson *et al.*, 2013). AM fungi obtain up to 20–30% of total plant photosynthates (Drigo *et al.*, 2010) and may enhance the decomposition of organic matter, releasing substantial quantities of CO<sub>2</sub> to the atmosphere through their respiration (Nottingham *et al.*, 2010).

Tropical forest growth currently constitutes the largest terrestrial sink for anthropogenic CO<sub>2</sub> (Oren *et al.*, 2001) and thus makes a substantial contribution to the regulation of the global climate system (Field *et al.*, 1998). Anticipating future effects of anthropogenic change on tropical forests demands a clearer understanding of how nutrient availability limits forest productivity and the roles of AM fungi in complex scenarios of nutrient limitation and co-limitation. Nonetheless, AM fungi are under-

investigated in tropical systems in general, and tropical forests in particular (Alexander & Selosse, 2009).

It is widely hypothesised that the symbiotic function of AM fungi is determined by the relative availability of C, nitrogen (N) and phosphorus (P; Johnson, 2010; Johnson *et al.*, 2013). This is based on evidence which shows that fertilisation with N and P can reduce AM fungal colonisation of roots (Johnson *et al.*, 2003), and that the relative amounts of N and P determine mycorrhizal symbiotic function (Johnson, 2010). In some cases this may cause AM fungi to behave less mutualistically (Johnson, 1993); where neither N or P is limited, the only limitation to fungal growth is the supply of plant C, meaning that fungal C demand can increase to the point where plant growth is depressed (Johnson, 2010).

Much current understanding concerning the function of AM fungal symbioses comes from studies that explore how variation in nutrient availability affects AM fungal characteristics (e.g. Treseder, 2004; Wurzbürger & Wright, 2015). Amongst these, nutrient addition experiments are one of the most widely used

approaches, particularly in field settings (Treseder, 2004). Nutrient addition is hypothesised to affect AM fungi either directly, by alleviating fungal nutrient limitation and thereby stimulating fungal growth (Treseder & Allen, 2002), or indirectly, by causing plants to reduce investment of carbohydrate in their AM fungal partners (Mosse & Phillips, 1971; Johnson, 2010).

Besides altering AM fungal biomass, nutrient addition may affect AM fungal community composition and diversity. Changes in community composition and diversity are likely to arise from differences in the functional properties of AM fungal taxa and their ability to compete with other fungi (AM or saprobe) for key resources (Hart & Reader, 2002; Maherali & Klironomos, 2007; Powell *et al.*, 2009). For instance, different AM fungal taxa can vary in the translocation of P (Ravnskov & Jakobsen, 1995) or N (Veresoglou *et al.*, 2012) to plant partners, carbon storage and demand (Pearson & Jakobsen, 1993), relative allocation to intra- and extra-radical biomass (Hart & Reader, 2002), and growth and life-history strategy (Hart & Reader, 2002; Maherali & Klironomos, 2007; Powell *et al.*, 2009). Furthermore, plant–AM fungal combinations perform differently in alternative settings, with a wide range of symbiotic outcomes (Klironomos, 2003; Powell *et al.*, 2009). Consequently, the advantage of AM fungal associations will vary according to the prevailing conditions and the ecological niche of the fungal partner. Evaluation of community parameters thus provides important information to supplement the aggregate metrics of root colonisation and concentration of the AM fungal biomarker lipid (a proxy for AM fungal biomass), which cannot distinguish between members of the AM fungal community.

In addition, AM fungal species that share a common evolutionary history may also share traits and ecological functions (Maherali & Klironomos, 2007; Powell *et al.*, 2009), and community data can thus be used to infer the ecological processes structuring AM fungal communities. Phylogenetically overdispersed communities (communities consisting of taxa that are less related to each other than expected by chance) are hypothesised to be structured by competition, preventing closely related and functionally similar taxa (those sharing a common niche) from co-occurring. By contrast, phylogenetically underdispersed (or clustered) communities are hypothesised to be structured by habitat filters; features of the environment that permit only the co-occurrence of species with specific traits or ecological tolerances, and which can cause taxa with similar traits to respond in similar ways to environmental pressures (Webb *et al.*, 2002; Maherali & Klironomos, 2007).

The great majority of nutrient addition studies apply inorganic fertilisers (e.g. see Treseder, 2004). These studies are useful in highlighting the roles of individual nutrients and simulating the effects of inorganic nutrient deposition. However, fertilisation treatments are artificial and do not mimic pathways of nutrient cycling under natural conditions (Sayer & Banin, 2016). Furthermore, the regulation of plant–AM fungal relationships is strongly dependent on the relative availability of different nutrients (Treseder & Allen, 2002; Johnson, 2010), whereas the addition of large quantities of one or more inorganic nutrients (e.g. N, P, potassium (K)) strongly distorts stoichiometric relationships, and

largely neglects the role of organic matter in nutrient cycling (Sayer & Banin, 2016).

Under natural conditions, nutrient cycling in forests occurs largely through litterfall, root death, root exudates, decomposition, and the growth and death of microorganisms (Attiwill & Adams, 1993; Leff *et al.*, 2012). It is via these processes that the regulatory processes governing plant–AM fungal exchange have evolved. Indeed, over large latitudinal gradients there is a strong relationship between leaf litter quality, the organic matter resulting from its degradation, and the predominant mycorrhizal type in a given bioregion (Read, 1991). Nonetheless, there have been few experimental investigations into the effects of leaf litter amendments on AM fungi in highly diverse tropical forests.

Although multiple lines of evidence suggest a key role for AM fungi in cycling nutrients via organic sources, the majority of studies investigating the effects of organic amendments on AM fungi have been conducted in experimental microcosms, and most have examined changes in biomass rather than community parameters (Hodge, 2014). These experiments demonstrate that AM fungal hyphae preferentially proliferate in organic substrates in experimental microcosms (Hodge & Fitter, 2010), are able to capture N from organic substrates (Leigh *et al.*, 2009), and can enhance the decomposition of organic material (Hodge, 2014). The few existing field studies show that organic matter additions in agricultural systems tend to increase AM fungal colonisation of plant roots and hyphal abundance in soils (Gryndler *et al.*, 2005; Gosling *et al.*, 2010). Furthermore, AM fungal hyphae can grow into decomposing leaf litter on tropical forest floors (Herrera *et al.*, 1978; Posada *et al.*, 2012; Camenzind & Rillig, 2013). Together, these studies strongly suggest that AM fungal hyphae are important in recycling nutrients from leaf litter. This is likely due to tightly coupled interactions between AM fungi and saprophytic fungi and bacteria (Herman *et al.*, 2012) given that AM fungi have not been shown to possess saprophytic capabilities (Hodge, 2014).

We investigated AM fungal responses to altered organic matter inputs in a lowland tropical forest in Panama using an existing long-term litter manipulation experiment in which 9 yr of litter removal and addition treatments have altered fine root biomass (Sayer *et al.*, 2006a), litter production, foliar and litter nutrient concentrations, and soil nutrient pools (Sayer & Tanner, 2010b; Vincent *et al.*, 2010). This platform provided a unique opportunity to evaluate the responses of AM fungal communities to changes in organic matter inputs in a well-studied lowland tropical forest setting.

We hypothesised that: litter addition would increase net AM fungal abundance, given the well-documented stimulatory effects of organic matter additions on AM fungal growth; litter removal would also increase net AM fungal abundance, given that plants may increase investment in AM fungi when nutrient availability is reduced (Johnson, 2010); that the addition or removal of organic matter would result in changes in the AM fungal community composition; and that litter manipulation would alter the ecological processes structuring AM fungal communities, and that this would be reflected in changes in the degree of relatedness (or phylogenetic structure), of AM fungal communities.

## Materials and Methods

### Site description and experimental design

The Gigante Litter Manipulation Experiment (GLMP) is located on the Gigante Peninsula (9°06' N, 79°54' W) within the Barro Colorado Nature Monument (BCNM) in Panama, Central America. Nearby Barro Colorado Island (BCI; c. 5 km from the study site) has a mean annual rainfall of 2600 mm, with a strong dry season between January and April and a mean annual temperature of 27°C (Leigh, 1999). Tree species composition and canopy height are characteristic of mature (>200-yr-old) secondary forest (Wright *et al.*, 2011) and the soils are classed as moderately acidic Oxisols (Dieter *et al.*, 2010; Turner & Wright, 2014), with low concentrations of available P and moderate concentrations of base cations (Turner & Wright, 2014). The GLMP consists of 15 45 × 45 m plots; starting in 2003, leaf litter from five plots was raked up once a month (litter removal treatment, L−), immediately added to five plots where it was distributed as evenly as possible (litter addition treatment, L+), and five plots were left undisturbed as controls (see Sayer & Tanner, 2010a,b; for details).

### Sampling

In May 2012, after 9 yr of treatments, we sampled at six points in the inner 30 × 30 m of each of the 15 experimental plots (a total of 30 samples per treatment); we selected sampling points at random using random number sheets to delineate point coordinates, with the provision that all points were separated by at least 3 m. At each sampling point, we collected the litter (Oi) and fermentation (Oe) horizons from a 78.5 cm<sup>2</sup> area, using a knife to cut around the edge of a metal disk (controls and L+ treatments only; the L− treatment lacked an organic horizon), and two cores from the mineral soil (0–10 cm depth) using a 5-cm diameter corer (all treatments). To prevent cross-contamination, we wiped down and flame-sterilised all equipment between samples, handled all samples with fresh latex gloves, and double-bagged samples in sealed Ziploc™ bags (Ziploc, USA). All samples were stored at 4°C and processed within 36 h of returning from the field. Root samples were obtained from one of the two cores per sampling point by washing away soil and organic matter under a continuous stream of filtered water over a sieve with a mesh size of 500 µm. We retained fine roots (≤1 mm in diameter) for further analysis, drying a subsample over silica gel for DNA extraction, and storing a second subsample in 70% ethanol for microscopic analysis. The remaining soil cores were sieved to remove stones and roots, composited to make one sample per plot, and thoroughly homogenised. 20 g subsamples for lipid analysis were frozen at −80°C for 12 h, lyophilised, and stored dry at −80°C until further processing.

Before lipid and nutrient extractions, an equal mass of each sample was pooled to make one composite sample per plot (a total of 15 samples). Before DNA extraction, the six root samples per plot were individually pulverised in a homogeniser (TissueLyser II; Qiagen, West Sussex, UK), and an equal mass of

each sample was pooled to make one composite sample per plot (a total of 15 samples). Microscopic analysis of root samples was performed on individual samples (total = 6 samples per plot, 90 samples in total).

### AM fungal abundance

We used the percentage of root length colonised as a measure of intraradical AM fungal abundance (McGonigle *et al.*, 1990). We soaked and rinsed the root samples with distilled water to remove the ethanol. Roots were then cleared by autoclaving in 5% KOH for 5–60 min; bleached in solution of ammonia in 3% H<sub>2</sub>O<sub>2</sub> for 15–60 min; acidified in 2% HCl for 30 min; and stained with 0.05% trypan blue (in a 1 : 1 : 1 solution of distilled water, glycerol and lactic acid) for 20 min at 60°C. The optimum clearing and bleaching time varied depending on the thickness and pigmentation of the roots. We quantified AM fungal colonisation by hyphae, vesicles and arbuscules using a compound light microscope at ×200 magnification, according to the method of McGonigle *et al.* (1990), with at least 100 intersections per sample, and one sample per core. AM fungal colonisation was expressed as the percentage fine root length colonised by AM fungal hyphae, vesicles or arbuscules.

We used the neutral lipid fatty acid (NLFA) 16:1ω5 as a biomarker for extra-radical AM fungal biomass. We performed lipid extraction and analysis according to Frostegård *et al.* (1993) with modifications (Nilsson *et al.*, 2007). Briefly, lipids extracted from 4 g lyophilised soil per plot were fractionated into neutral lipids, glycolipids and polar lipids on silica columns by successive elution with chloroform, acetone and methanol. Methyl nonadecanoate (FAME 19:0) was added as an internal standard, and neutral and polar fractions were converted to fatty acid methyl esters (FAMES) before analysis on a gas chromatograph with a flame ionisation detector and a 50 m HP5 capillary column (Hewlett Packard, Wilmington, DE, USA). The mean NLFA:PLFA ratio across all samples was 1.3, suggesting that NLFA 16:1ω5 is an effective AM fungal biomarker in these soils (Olsson, 1999).

### Soil chemistry

Measurement of inorganic N (NO<sub>3</sub><sup>−</sup> and NH<sub>4</sub><sup>+</sup>), resin-extractable P, organic P and pH was performed as described in Turner & Wright (2014). Analysis of total N and C was performed on air-dried soils by automated combustion and gas chromatography on a Thermo Flash EA1112 analyzer (CE Elantech, Lakewood, NJ, USA). Organic P was extracted in a mixture of 0.25 M NaOH and 0.05 M EDTA, and analysed as described by Turner (2008). Exchangeable cations were extracted in 0.1 M BaCl<sub>2</sub>, with detection by ICP-OES (Hendershot *et al.*, 2008), and effective base saturation (EBS) was calculated by dividing the cmol of positive charge kg per dry soil of exchangeable bases (calcium (Ca) + K + magnesium (Mg) + sodium (Na)) by that of the total cations (aluminium (Al) + Ca + iron (Fe) + K + Mg + Mn + Na; Hendershot *et al.*, 2008).

## DNA extraction and sequencing

We extracted DNA from 50 mg of pulverised root using MoBio PowerPlant DNA isolation kits according to the manufacturer's instructions (MoBio Laboratories Inc., Carlsbad, CA, USA).

We amplified the partial small subunit (SSU) region of 18S ribosomal DNA (*c.* 550 bp) with the universal eukaryotic primer NS31 (Simon *et al.*, 1992) and the AM fungal-specific primer AM1, which amplifies the major families of the Glomeromycota (Helgason *et al.*, 1998). We chose this primer set because it is widely represented in sequence databases, and because we wanted to facilitate comparisons with previous work using these primers. In addition, these primers have been demonstrated to have extremely low PCR bias against artificially assembled community templates (Cotton *et al.*, 2014). Before amplification, the primers were modified by the addition of the 454 pyrosequencing adaptors A and B, in addition to a 10 bp multiplex identifier (MID) on the forward primer (NS31). We conducted duplicate PCRs in 25 µl sample volume using Phire hot start II DNA polymerase (Life Technologies Ltd, Paisley, UK). Conditions were: 98°C for 1 min; 32 cycles of 98°C for 10 s and 72°C for 15 s; and a final extension phase of 72°C for 2 min.

We gel-purified the PCR products using MinElute PCR purification kits (Qiagen) and pooled the samples in equimolar concentrations, evaluating the concentration of DNA in the cleaned PCR products using Quant-iT PicoGreen dsDNA Assay Kit (Invitrogen, Life Technologies Ltd). Amplicon libraries were distributed on PicoTiter Plates and sequenced on an FLX Titanium system using Lib-L shotgun chemistry (Roche). No sequences were detected in the blanks included as negative controls at each of the extraction, PCR, gel purification and quantification steps.

## Bioinformatic analysis

All bioinformatic analyses were performed using the software mothur (Schloss *et al.*, 2009) unless stated otherwise. Sequence filtering was performed with the *sff.multiple* quality filtering protocol. Reads were removed from the dataset if they did not contain the 10 bp MID, had more than one error in the barcode sequence, more than two errors in the forward primer, or were shorter than 200 bp in length. After quality filtering and removal of barcode and primer sequences, clustering was performed using the algorithm Clustering 16S rRNA for operational taxonomic unit (OTU) prediction (CROP), an unsupervised Bayesian clustering method that forms clusters based on the organisation of sequences without setting a hard similarity cutoff (Hao *et al.*, 2011). To provide finer taxonomic resolution, we set the *i* and *u* parameters to 2% cluster difference rather than the conventional 3% because the SSU region has relatively low variation (Öpik *et al.*, 2013; Davison *et al.*, 2015). The centre sequence from each cluster was used as a representative sequence in subsequent analyses.

Sequence alignment was performed with the software MAFFT v.7.149b (Katoh *et al.*, 2002) using the L-INS-i algorithm (iterative refinement using local pairwise alignment) and the alignment

from Krüger *et al.* (2012) as a backbone. Alignments were improved with MUSCLE (Edgar, 2004) using the *-refine* option. Trees were built using RAxML v.8.0 (Stamatakis, 2014) with GTR GAMMA implementation and bootstrap values based on 1000 runs.

We used the Basic Local Alignment Search Tool (BLAST, Altschul *et al.*, 1990; minimum *e*-value  $10^{-30}$ ) on one representative sequence from each cluster iteratively against three databases in the following order of preference: sequences from Krüger *et al.* (2012); all virtual taxa (VT) from the MaarjAM AM fungal sequence database (<http://www.maarjam.botany.ut.ee>); and all 18S Glomeromycotan sequences from SILVA database. Non-Glomeromycotan clusters were removed when the highest blast match did not correspond to an AM fungal sequence in any of the three datasets.

Clusters were named based on matches to database entries at >97% covering a minimum of 80% of the query sequence. We used the generic names from Krüger *et al.* (2012), and VT numbers from the MaarjAM database. Where clusters did not match a VT at >97% we assigned a name based on the highest VT match and phylogeny (e.g. *Glomus\_OTU1*). We fused clusters based on matches to database sequences >97% and the tree topology obtained from RAxML. Clusters that occurred in fewer than two samples, and with fewer than five reads total were removed from the dataset. Raw sequence data were deposited in the International Nucleotide Sequence Database Sequence Read Archive (accession no. SRP076949).

## Statistical analysis

All statistical analyses were conducted in R v.3.1.2 (R Development Core Team, 2014).

**Multivariate analysis of AM fungal communities** We accounted for variation in the number of sequences between samples by using a variance stabilising (VS) transformation of the OTU table, implemented with the DESEQ2 package (Love *et al.*, 2014), according to McMurdie & Holmes (2014). This approach avoids the need for rarefying, which can result in data that misrepresent the original community (McMurdie & Holmes, 2014). All subsequent analysis was performed on the VS transformed OTU table, using the copy number of DNA sequences as a measure of relative abundance of each OTU.

In order to examine the effect of litter manipulation on AM fungal community composition, we used multivariate generalised linear models (M-GLMs) with negative binomial error structures using the *MVABUND* package (Wang *et al.*, 2012). M-GLMs provide a more robust way to analyse multivariate community data than distance-based approaches such as PERMANOVA (Warton *et al.*, 2015). We ascertained the degree to which individual OTUs were affected by litter manipulation using DESEQ2 (Anders & Huber, 2010), which estimates the effect size (as logarithmic fold-change) and reports *P*-values adjusted for multiple comparisons.

In order to visualise differences in AM fungal communities across litter manipulation treatments we used nonmetric

multidimensional scaling (NMDS) ordination, using the metaMDS function in the VEGAN package (Anderson, 2001; Oksanen *et al.*, 2010). Ordination was based on Bray–Curtis dissimilarity calculated from square-root transformed abundances. The range of data values was large, and a square root transformation was applied to improve the quality of the ordination by reducing the weighting of the most abundant OTUs (Oksanen *et al.*, 2010; Legendre & Legendre, 2012).

Soil physical characteristics were standardised to zero mean and unit variance, and fit to the NMDS ordinations (function `envfit` from the VEGAN package) with significance ascertained using 9999 permutations. Individual values of exchangeable cations were collapsed into the metric of effective base saturation (EBS). Organic phosphorus correlated closely with resin-extractable phosphorus ( $r^2 > 0.7$ ) and was omitted, because resin-extractable phosphorus better approximates the plant-available phosphorus fraction (Condit *et al.*, 2013).

**Community phylogenetic structure** We asked whether litter manipulation altered the degree of relatedness between taxa in AM fungal communities. We used two indices of community phylogenetic structure: Net relatedness index (NRI) and nearest taxa index (NTI; Webb, 2000). Positive values of these metrics indicate that taxa in a community are, on average, more closely related to each other than to members of the regional taxon pool (phylogenetically clustered), and negative values indicate that taxa in a community are less closely related (phylogenetically overdispersed). NRI is sensitive to tree-wide phylogenetic patterns, and NTI is sensitive to phylogenetic community patterns close to the tips of the phylogeny. Observed values of these metrics were compared with 10 000 null communities generated using the ‘independentswap’ algorithm, which maintains column and row totals and accounts for differences in community richness and taxon prevalence (Gotelli, 2000). Statistical significance of phylogenetic structure was ascertained using a two-tailed *t*-test. Community phylogenetic analysis was performed using the PICANTE package (Kembel *et al.*, 2010).

**Univariate analysis of AM fungal abundance and diversity, and soil physical characteristics** We analysed the effects of litter manipulation on the concentrations of NLFA 16:1ω5 in the soil, AM fungal colonisation of plant roots, AM fungal OTU richness and predominance, and metrics of phylogenetic community structure (NRI and NTI) using linear models having confirmed that all variables met the assumptions. Where the main effect of litter manipulation was significant, we performed Dunnett’s *post-hoc* analysis to compare each treatment with the controls.

In order to ascertain whether AM fungal colonisation of roots was greater in the mineral soil or organic layer, we built linear mixed effects models (using the LME4 package; Bates *et al.*, 2015). Models included ‘layer’ and ‘treatment’ as fixed effects, and ‘plot’ as a random effect. The significance of fixed effects was assessed by comparing nested models using parametric bootstrapping with 10 000 simulations, using the PBMODCOMP function from the PBKRTEST package (Halekoh & Højsgaard, 2014). Results are reported as significant at  $\alpha < 0.05$ .

## Results

### Soil chemistry

Soil nutrients were lower in litter removal compared with litter addition treatments for inorganic N; resin and organic P, pH, and extractable Ca, Mg and Mn (K was not significantly lower). Compared with the controls, the soils in the L– plots had lower concentrations of inorganic N, resin and organic P, Ca and Mg, whereas soils in the L+ plots had higher concentrations of resin P and Ca (Fig. 1; Supporting Information Table S1). A full discussion of the effects of litter manipulation on soil chemistry is provided in Sayer & Tanner (2010a,b) and Sayer *et al.* (2012).


### AM fungal abundance

There was no significant effect of litter manipulation on the proportion of root length colonised by any AM fungal structure in the mineral soil (total colonization, hyphae, vesicles or arbuscules), although for each of the structures there was a trend towards higher root colonisation in both litter removal and litter addition treatments compared with the controls (Fig. 2; total colonisation:  $F_{2,12} = 1.7$ ,  $P = 0.23$ ; hyphae:  $F_{2,12} = 1.4$ ,  $P = 0.29$ ; vesicles:  $F_{2,12} = 2.5$ ,  $P = 0.13$ ; arbuscules:  $F_{2,12} = 1.3$ ,  $P = 0.31$ ). In the control and litter addition plots, the proportion of root length colonised by all AM fungal structures was substantially greater in the superficial organic layer than in the mineral soil (significant ‘layer’ term; hyphae: likelihood-ratio test (LRT) = 50.0  $P < 0.001$ ; vesicles: LRT = 19.6,  $P < 0.001$ ; arbuscules: LRT = 28.6,  $P < 0.001$ ; all structures: LRT = 51.6,  $P < 0.001$ ; Fig. 2). Because root colonization was highest in the superficial organic layer, the overall abundance of AM fungi was lower in the litter removal treatment, which lacked this layer.

There was no effect of litter manipulation on AM fungal biomass in the mineral soil (concentration of NLFA 16:1ω5; Fig. S1), nor was AM fungal biomass correlated with any of the measured soil variables (soil pH:  $F_{1,13} < 0.001$ ,  $P = 0.98$ ; effective base saturation:  $F_{1,13} = 0.01$ ,  $P = 0.92$ , resin-extractable phosphorus:  $F_{1,13} = 0.12$ ,  $P = 0.74$ ; and inorganic nitrogen:  $F_{1,13} = 0.54$ ,  $P = 0.48$ ).

### AM fungal community composition and structure

Four AM fungal families were represented in the sequencing dataset (Acaulosporaceae, Archaeosporaceae, Gigasporaceae, Glomeraceae; Fig. 3), indicating reasonable taxonomic coverage of the Glomeromycota (based on the classification of Redecker *et al.*, 2013). No members of the Diversisporaceae, Paraglomeraceae, Geosiphonaceae, Ambisporaceae, Claroideosporaceae or Pacisporaceae were detected. Rarefaction curves for each sample indicated that sequencing intensity was sufficiently high to detect the majority of OTUs. Rarefaction curves pooled by experimental treatment approached asymptotes, indicating that sampling effort was sufficient to capture the range of AM fungal taxa across the sites (Fig. S2). A total of 10 197 sequences were retained after quality control, clustered into 72 OTUs, and 95.9% of all sequences matched Glomeromycota in the databases. Fifty-six



**Fig. 1** Effects of litter manipulation on soil physical characteristics. Values are means  $\pm$  Fisher's Least Significant Difference. Grey shaded regions represent control treatments. Litter treatments are significantly different from controls at  $\alpha < 0.05$  ( $n = 5$ ) where error bars do not overlap the grey shaded regions. Standard normal deviates are plotted to facilitate visual comparison of effect size. Aluminium (Al), manganese (Mn) and inorganic nitrogen (N inorg.) were log-transformed before analysis due to heteroscedasticity. N (inorg.), inorganic N; P (res.), resin extractable phosphorus (P); P (tot.), total P; TEB, total exchangeable bases; EBS, effective base saturation; L-, litter removal treatment; L+, litter addition treatment.

OTUs remained after blasting, filtering, merging and trimming (exclusion of OTUs with a total of five or less reads), representing a total of 8825 sequences. Each sample (one per plot) contained a mean of 18 OTUs (range: 11–24), and the mean number of sequences per sample was 588 (range: 237–1225; Table S2). A phylogenetic tree is provided in Fig. S3.

Overall AM fungal community composition was altered by litter removal but was not significantly affected by litter addition (multivariate GLM:  $Wald_{2,12} = 11.5$ ,  $P < 0.003$ ; treatment contrast for litter removal:  $Wald = 9.2$ ,  $P < 0.003$  and for litter addition:  $Wald = 5.9$ ,  $P = 0.24$ ; Fig. 4). There were no significant differences among treatments when the analysis was repeated at the levels of genus and family (multivariate GLM; genus:  $Wald_{2,12} = 3.9$ ,  $P = 0.24$ ; family:  $Wald_{2,12} = 2.1$ ,  $P = 0.66$ ; Fig. 3). In the analysis of individual OTUs (using the DESeq2 package), litter removal significantly ( $P < 0.05$ ) reduced the relative abundance of four OTUs and increased the relative abundance of three OTUs ( $P < 0.05$ ; Fig. 5). By contrast, litter addition significantly increased the relative abundance of two OTUs (Fig. 5; Table S3). All of the significantly affected OTUs were in the family Glomeraceae apart from a single OTU in the Acaulosporaceae, which had lower relative abundance in the litter removal treatment. Neither litter treatment altered the total number of AM fungal OTUs (richness; ANOVA:  $F_{2,12} = 0.15$ ,  $P = 0.86$ ), nor the proportional abundance of the dominant AM fungal taxon (predominance; ANOVA:  $F_{2,12} = 0.37$ ,  $P = 0.69$ ; Fig. S4). Of the variables fitted to the NMDS ordination, soil pH, effective base saturation (EBS), resin-extractable P, and inorganic N concentrations were significantly correlated with AM fungal community composition (Fig. 4).


### AM fungal community assembly

Litter manipulation moderately altered the degree of relatedness between taxa in AM fungal communities, as summarised by the

NRI. There was a greater likelihood of detecting closely related taxa in litter addition plots than litter removal plots (ANOVA:  $F_{2,12} = 4.02$ ,  $P = 0.05$ ; Fig. 6), although neither treatment differed significantly from controls. Furthermore, whereas neither treatment showed significant phylogenetic structure of AM fungal communities relative to null model distributions, the NRI was  $> 0$  in the litter addition treatment and  $< 0$  in the litter removal treatment, indicating a trend towards phylogenetic underdispersion (taxa more related to each other than expected by chance) in the litter addition treatment (Fig. 6) and phylogenetic overdispersion (taxa less related to each other than expected by chance) in the litter removal treatment. However, when we used the NTI, which is sensitive to patterns in relatedness close to the tips of the phylogeny, AM fungal communities were neither significantly structured relative to null distributions, nor affected by litter manipulation (ANOVA:  $F_{2,12} = 0.25$ ,  $P = 0.79$ ).

### Discussion


Litter removal altered arbuscular mycorrhizal (AM) fungal community composition (Fig. 4), indicating that inputs of organic matter are important in structuring AM fungal communities. Together with substantially greater AM fungal root colonisation in the superficial organic layer than the mineral soil (70% vs 30%, respectively; Fig. 2e–h), our findings suggest that AM fungi obtain a substantial part of their nutrition from decomposing organic matter in this lowland tropical forest. We observed a trend towards increased AM fungal colonisation of roots growing in the mineral soil both in litter addition and litter removal treatments relative to controls (Fig. 2), providing some support for our hypotheses that plants may increase investment in AM fungal associations in both litter addition and litter removal treatments.




**Fig. 2** Percent root length colonised by arbuscular mycorrhizal (AM) fungi (total colonisation, colonisation by hyphae, colonisation by vesicles and colonisation by arbuscules). Left-hand panels (a–d) show the effect of litter manipulation on AM fungal colonisation of roots in the mineral soil. Right-hand panels (e–h) compare colonisation in roots between the mineral soil ('soil') and superficial organic horizon ('organic') across control and litter addition treatments. L–, litter removal; C, control; L+, litter addition. In left hand panels (a–d) values are means  $\pm$  Fisher's Least Significant Difference, and nonoverlapping error bars indicate significance at  $\alpha < 0.05$  ( $n = 5$ ). In right-hand panels, values are means  $\pm$  95% confidence intervals obtained by parametric bootstrapping with 10 000 simulations.

### Litter removal


Litter removal may have altered AM fungal community composition by reducing nitrogen -availability, either via direct fungal N-limitation, or by altering plant N status, leading to changes in plant allocation to AM fungi. Litter removal reduced the amount and availability of soil inorganic N, and crucially, reduced N concentrations in leaf litter after 5 yr (Sayer & Tanner, 2010b; Sayer *et al.*, 2012), suggesting that N-availability to plants had decreased. N concentrations of AM fungal hyphae are substantially higher than those of plant tissues (Hodge *et al.*, 2010) and comparison of the carbon (C) : N ratios of plant and fungal tissues indicate that severe N-limitation may be more likely to



**Fig. 3** Mean proportional abundance of arbuscular mycorrhizal (AM) fungal (a) genera and (b) families in mixed root samples across litter manipulation treatments ( $n = 5$ ); L–, litter removal; C, control; L+, litter addition.



**Fig. 4** Nonmetric multidimensional scaling (NMDS) ordination plot showing changes in arbuscular mycorrhizal (AM) fungal community composition in long-term litter removal plots (circles), but not litter addition plots (triangles), compared with controls (squares) in a lowland tropical forest. Site scores are shown and ellipses describe 95% confidence areas. Arrows indicate the direction and degree of significant correlations between NMDS axes and soil physical characteristics ( $n = 5$ ). EBS, effective base saturation; P (resin), resin extractable phosphate; N (inorg.), inorganic N; L–, litter removal; C, control; L+, litter addition. Axes are scaled to half-change (HC) units, by which one HC unit describes a halving of community similarity.



**Fig. 5** Effect of litter addition (red) and removal (blue) on the relative abundance of individual arbuscular mycorrhizal (AM) fungal operational taxonomic units (OTUs). Significantly altered ( $P < 0.05$ ) OTUs are shown based on both adjusted and unadjusted  $P$ -values. The names of OTUs that are significantly affected by litter manipulation are emboldened. x-axis indicates the effect size as  $\log_2$  fold change, and error bars show SE. OTUs are arranged in order of decreasing rank abundance (more highly ranked OTUs are those that are more prevalent across all samples in the dataset). Significance was ascertained based on negative binomial Wald tests using standard maximum likelihood estimates for generalised linear models, as implemented in the DESeq2 package.


community composition by increasing AM fungal competition (both with other AM fungi and saprobes) for a more limited resource.

An alternative possibility is that changes in AM fungal community composition in the litter removal plots reflect niche separation arising from a shift in AM fungal phosphorus (P)-acquisition strategies. The availability of P is thought to limit many biological processes in lowland tropical forests (Vitousek & Sanford, 1986), and is a limiting nutrient in these forests (Wright *et al.*, 2011; Turner & Wright, 2014). As a large proportion of the P required for plant growth is cycled through leaf litter (Sayer & Tanner, 2010b), we would expect the litter removal treatment to affect plant P status. However, there was no reduction in leaf litter P in litter removal plots, nor a reduction in litterfall or plant productivity in the first 6 yr of litter manipulation (Sayer & Tanner, 2010b; Sayer *et al.*, 2012), indicating that trees in the litter removal plots were able to access sufficient P from alternative sources to maintain productivity and foliar P concentrations.

At least some of the additional P available to plants in the litter removal treatment was probably acquired from stable organic P pools in the mineral soil. Organic P in forests occurs in fresh organic matter (such as leaf litter), microbial biomass and

nonbiomass stable organic P (Vincent *et al.*, 2010). Under normal conditions, P is rapidly released from leaf litter via leaching (Schreeg *et al.*, 2013) or mineralisation (Richardson & Simpson, 2011) before being taken up directly by plants by mycorrhizal fungi (Herrera *et al.*, 1978). This results in 'direct' nutrient cycling by which nutrient losses through leaching might be minimised (Went & Stark, 1968). After 3 yr of litter removal, the stable organic P pool in the upper 2 cm of the mineral soil was reduced by 23%, whereas the overall inorganic P pool remained unchanged (Vincent *et al.*, 2010). Given that our study took place after 9 yr of litter removal, and the depletion of the stable organic P pool had conceivably continued, it is probable that additional P could also have been mobilised from recalcitrant mineral P stocks in the soil.

The role of AM fungi in P acquisition is well-known (Smith & Read, 2008), and it is likely that a shift in plants' primary source of P from decomposing litter to stable organic P and stocks of mineral P would involve a change in the primary function of plants' AM fungal associations. Although limited, there is evidence that different AM fungal species differ in their ability to acquire (Cavagnaro *et al.*, 2005) and transport P to plant hosts (Munkvold *et al.*, 2004), and that AM fungal taxa may benefit



**Fig. 6** Litter manipulation moderately altered the degree of relatedness between taxa in arbuscular mycorrhizal (AM) fungal communities when described using the metric of net relatedness index (NRI). Higher numeric values correspond to more closely related AM fungal communities. Values are means  $\pm$  Fisher's Least Significant Difference: nonoverlapping error bars indicate significance at  $\alpha < 0.05$  ( $n = 5$ ). Dotted lines indicate significance threshold of  $\alpha = 0.05$  derived from comparison with 10 000 null communities generated using the 'independentswap' algorithm. L-, litter removal; C, control; L+, litter addition.

plants to different degrees based on the type of soil P available (e.g. mineral vs organic; Reynolds *et al.*, 2005). Consequently, it is possible that the taxa with increased relative abundance in litter removal plots were mineral P specialists, and those with decreased relative abundance were litter specialists (Fig. 5). Nonetheless, it is striking that the dominant taxon and the relative abundances of most taxa in the litter removal treatment remained unchanged. Given the probable shift in plants' primary P source in the litter removal treatment, this would suggest that most of the AM fungal taxa observed at this site are readily able to adapt to the changed conditions. This is interesting in the light of studies of ectomycorrhizal fungi, which document wide differences in the ability of different taxa to mobilise and acquire P from different sources (Plassard *et al.*, 2011).

Other factors besides changes in nutrient availability could explain the shift in community composition observed in the litter removal treatment. Organic amendments such as leaf litter can affect a number of other soil properties besides nutrient availability, such as habitat space available for decomposers (Sayer, 2006). It is thus possible that AM fungal communities were affected by changes in the non-AM microbial community or soil fauna, which can impact AM fungal growth and function (Johnson *et al.*, 2005; Sayer *et al.*, 2006b; Gryndler *et al.*, 2008; Hodge, 2014), and which play a key role in AM fungal uptake of nutrients from leaf litter given the lack of documented saprophytic effects of AM fungi (Hodge, 2014). Previous studies at this site show no major changes in either temperature or soil water content among treatments (Sayer & Tanner, 2010a), and it is thus unlikely that these factors are responsible for the observed effects.

Soil pH was correlated with the nonmetric multidimensional scaling (NMDS) ordinations of AM fungal community shifts, and may have been responsible for some of the observed shifts in

community composition. However, studies documenting the effects of pH on AM fungi have largely reported a reduction in root colonisation and extra-radical hyphal biomass with decreasing pH (Wang *et al.*, 1993; Clark, 1997; van Aarle *et al.*, 2002) as well as reduced AM fungal  $\beta$  diversity (Dumbrell *et al.*, 2009), none of which were observed in this study.

Regardless of the mechanism underlying the shifts in AM fungal community composition, the trend towards more phylogenetically overdispersed (less closely related) AM fungal communities in the litter removal plots relative to the litter addition plots (Fig. 6) may reflect increasing competition between AM fungal taxa following litter removal. This is because more closely related AM fungal taxa tend to share functional traits (Maherali & Klironomos, 2007; Powell *et al.*, 2009), a phenomenon known as phylogenetic trait conservatism (Webb *et al.*, 2002). Consequently, phylogenetically overdispersed communities are thought to be structured more by competition than by habitat filtering, which reduces the likelihood that closely related and functionally similar taxa will co-occur (Webb *et al.*, 2002).

### Litter addition

AM fungal colonisation of roots was substantially higher in the organic horizons than the mineral soil in the control and litter addition treatments (70% vs 30%, respectively; Fig. 2e–h). This finding agrees with a sizeable body of evidence which shows that the addition of organic material may increase AM fungal colonisation of plant roots (Gryndler *et al.*, 2005, 2008; Gosling *et al.*, 2010), and AM fungal sporulation (Gosling *et al.*, 2010). Indeed, AM fungal hyphae proliferate in organic substrates (Hodge & Fitter, 2010), and grow into decomposing leaf litter in tropical forests (Herrera *et al.*, 1978; Posada *et al.*, 2012; Camenzind & Rillig, 2013). Together with the finding that fine roots proliferated into the organic horizons in the litter addition treatment (Sayer *et al.*, 2006a), our results suggest that AM fungi may represent important pathways for plant uptake of nutrients from sites of organic matter decomposition in this tropical forest. However, given that AM fungi lack substantial saprophytic capability (Hodge, 2014), it is unlikely that AM fungi themselves are actively involved in litter decomposition, but rather are able to efficiently acquire nutrients as they are released from decomposing organic matter by the action of saprobes.

Given much greater root colonisation by AM fungi in the organic horizons of the litter addition and control plots relative to the mineral soil, it is surprising that we observed no significant increase in root colonisation in the mineral soil of litter addition treatments relative to controls (Fig. 2a–d), where organic matter content is elevated relative to controls (Tanner *et al.*, 2016). It is possible that plant investment in AM fungi in litter addition plots is lower, due to the increases in soil fertility and tree nutrient status (indicated by marginal increases in litterfall and foliar N and P; Fig. 1; Sayer & Tanner, 2010b; Sayer *et al.*, 2012). This interpretation follows from the functional equilibrium hypothesis, by which plants allocate resources to the structures that are the most helpful in acquiring the most limiting nutrients (Johnson, 2010), and by which plants should reduce investment in AM fungal

associations when soil fertility increases because the carbon costs outweigh the nutritional benefits (Mosse & Phillips, 1971; Johnson, 2010). Reduced plant investment in AM fungi would counter the stimulatory effects of organic matter on AM fungal colonisation.

### Limitations of this study

We did not measure neutral lipid fatty acid (NLFA) in the superficial organic layer, or below 10 cm, so we were not able to determine if total AM biomass was affected by litter treatment. In addition, we did not characterise AM fungal communities from roots sampled from the superficial organic layer due to technical constraints. As such, we are unable to address the extra-radical responsiveness of AM fungi to increased inputs of organic matter, and directly address the selection of litter-specific AM fungal communities. Vertical stratification of ectomycorrhizal communities has been described in boreal forest podzols (Rosling *et al.*, 2003), and increased AM fungal colonisation of roots in the superficial organic layer could be hypothesised to reflect shifts in the structure and composition of AM fungal communities. This warrants further investigation. Finally, we made no direct measure of nutrient transfer, and our discussion of how leaf litter manipulation altered AM fungal function is thus necessarily speculative.

### Potential sequencing bias

AM fungal communities were strongly dominated by taxa in the Glomeraceae (Fig. 3), which was due in part to our choice of marker region because the small subunit (SSU) is biased towards Glomeraceae (Kohout *et al.*, 2014) and may underestimate diversity in some Diversisporales (Davison *et al.*, 2015). Indeed, a previous study in the Barro Colorado Nature Monument (BCNM) using Sanger sequencing and the same AM1/NS31 primer set likewise found a strong dominance of AM fungal species in the Glomeraceae (Husband *et al.*, 2002). Furthermore, a compilation of globally sampled AM fungal sequences obtained from the amplification of a similar SSU region (with the primers AML1/NS31) described a similar pattern: 79% of operational taxonomic units (OTUs) were from the order Glomerales (compared with 84% in this study), and 15% were from the Diversisporales (compared with 14% in this study; Öpik *et al.*, 2013). By contrast, a study in a montane forest in Ecuador using the ribosomal large subunit (LSU) region found their dataset dominated by the Diversisporales (Camenzind *et al.*, 2014).

We used the number of DNA sequences as a measure of relative abundance of OTUs. Although sequence abundance may reflect biases introduced through PCR and sequencing protocols, the NS31-AM1 primer set exhibited very low levels of PCR bias when used to amplify artificial community templates of known composition (Cotton *et al.*, 2014). This is possibly because of the consistent length (*c.* 1.5% variation) and GC content (*c.* 3% variation) of the amplified region across different AM fungal taxa (Helgason *et al.*, 1999), as variation in amplicon length and GC content are known to cause biases in PCR

(Ihrmark *et al.*, 2012), and may cause biases in the 454 sequencing process as well (Kausserud *et al.*, 2011). In any case, in a comparative analysis of our dataset using both quantitative and presence-absence approaches led to identical conclusions (Fig. S5).

### Concluding remarks

Our findings show that the presence of decomposing leaf litter is important both in structuring AM fungal communities and in determining the extent of root colonisation by AM fungi. Alterations in AM fungal community composition in response to litter removal may be due to a range of factors including the reduction of key nutrients supplied by decomposing leaf litter, notably N and P, changes in the action of saprobes, and changes in water availability and pH. We hypothesise that a reduction in the quantity of decomposing fresh organic matter brought about by litter removal may lead to AM fungi obtaining scarce nutrients such as P from recalcitrant organic or mineral sources in the soil. Our hypothesis helps to explain how trees were able to maintain their P-status despite the chronic removal of a major P input in this lowland tropical forest, and merits further investigation.

### Acknowledgements

We thank J. Grandez and J. Rodriguez for their assistance in the field. Dayana Agudo and Aleksandra Bielnicka assisted in the analysis of soils. I. Henderson provided advice on the sequencing work. A. Herre and E. Verbruggen provided discussion and comments on the manuscript. M.S. was funded by a Smithsonian Tropical Research Institute (STRI) predoctoral fellowship, a Cambridge Home and European Scholarship, the Department of Plant Sciences, Cambridge, and a Cambridge Philosophical Society travel grant. E.J.S. was supported by funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013), ERC Grant Agreement No. 307888. The Smithsonian Tropical Research Institute served as the base of operations and provided logistical support. The manuscript was improved by comments from four anonymous reviewers.

### Author contributions

M.S. and S.M. designed the study. E.V.J.T. and E.J.S. established the leaf litter manipulation experiment. M.S. and D.R. performed the lab and field work. P.A.O. and H.W. supported the lipid analysis. B.L.T. conducted the nutrient analysis. N.P.R. conducted the bioinformatic analysis. M.S. conducted the statistical analysis and wrote the manuscript with input from all authors.

### References

- van Aarle IM, Olsson PA, Soderstrom B. 2002. Arbuscular mycorrhizal fungi respond to the substrate pH of their extraradical mycelium by altered growth and root colonization. *New Phytologist* 155: 173–182.

- Alexander I, Lee S. 2005. Mycorrhizas and ecosystem processes in tropical rain forest: implications for diversity. In: Burslem D, Pinard M, Hartley S, eds. *Biotic interactions in the tropics*. Cambridge, UK: Biotic Interactions in the Tropics, 165–203.
- Alexander I, Selosse M-A. 2009. Mycorrhizas in tropical forests: a neglected research imperative. *New Phytologist* 182: 14–16.
- Altschul SF, Gish W, Miller W, Myers EW. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215: 403–410.
- Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biology* 11: R106.
- Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26: 32–46.
- Attiwill PM, Adams MA. 1993. Nutrient cycling in forests. *New Phytologist* 124: 561–582.
- Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67: 1–48.
- Camenzind T, Hempel S, Homeier J, Horn S, Velescu A, Wilcke W, Rillig MC. 2014. Nitrogen and phosphorus additions impact arbuscular mycorrhizal abundance and molecular diversity in a tropical montane forest. *Global Change Biology* 20: 3646–3659.
- Camenzind T, Rillig MC. 2013. Extraradical arbuscular mycorrhizal fungal hyphae in an organic tropical montane forest soil. *Soil Biology and Biochemistry* 64: 96–102.
- Cavagnaro TR, Smith FA, Smith SE, Jakobsen I. 2005. Functional diversity in arbuscular mycorrhizas: exploitation of soil patches with different phosphate enrichment differs among fungal species. *Plant, Cell & Environment* 28: 642–650.
- Clark RB. 1997. Arbuscular mycorrhizal adaptation, spore germination, root colonization, and host plant growth and mineral acquisition at low pH. *Plant Soil* 192: 15–22.
- Condit R, Engelbrecht BMJ, Pino D, Perez R, Turner BL. 2013. Species distributions in response to individual soil nutrients and seasonal drought across a community of tropical trees. *Proceedings of the National Academy of Sciences, USA* 110: 5064–5068.
- Cotton TEA, Dumbrell AJ, Helgason T. 2014. What goes in must come out: testing for biases in molecular analysis of arbuscular mycorrhizal fungal communities. *PLoS One* 9: e109234.
- Davison J, Moora M, Öpik M, Adhohleya A, Ainsaar L, Bâ A, Burla S, Diedhiou AG, Hiiesalu I, Jairus T *et al.* 2015. Global assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism. *Science* 349: 970–973.
- Dieter D, Elsenbeer H, Turner BL. 2010. Phosphorus fractionation in lowland tropical rainforest soils in central Panama. *Catena* 82: 118–125.
- Drigo B, Pijl AS, Duyts H, Kielak AM, Gamper HA, Houtekamer MJ, Boschker HTS, Bodelier PLE, Whiteley AS, van Veen JA *et al.* 2010. Shifting carbon flow from roots into associated microbial communities in response to elevated atmospheric CO<sub>2</sub>. *Proceedings of the National Academy of Sciences, USA* 107: 10 938–10 942.
- Dumbrell AJ, Nelson M, Helgason T, Dytham C, Fitter AH. 2009. Relative roles of niche and neutral processes in structuring a soil microbial community. *The ISME Journal* 4: 337–345.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797.
- Field C, Behrenfeld M, Randerson J, Falkowski P. 1998. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* 281: 237–240.
- Frostegård Å, Tunlid A, Bååth E. 1993. Phospholipid fatty acid composition, biomass, and activity of microbial communities from two soil types experimentally exposed to different heavy metals. *Applied and Environmental Microbiology* 59: 3605–3617.
- Gosling P, Ozaki A, Jones J, Turner M, Rayns F, Bending GD. 2010. Organic management of tilled agricultural soils results in a rapid increase in colonisation potential and spore populations of arbuscular mycorrhizal fungi. *Agriculture, Ecosystems & Environment* 139: 273–279.
- Gotelli NJ. 2000. Null model analysis of species co-occurrence patterns. *Ecology* 81: 2606–2621.
- Gryndler M, Hřselová H, Cajthaml T, Havránková M, Řezáčová V, Gryndlerová H, Larsen J. 2008. Influence of soil organic matter decomposition on arbuscular mycorrhizal fungi in terms of asymbiotic hyphal growth and root colonization. *Mycorrhiza* 19: 255–266.
- Gryndler M, Larsen J, Hřselová H, Řezáčová V, Gryndlerová H, Kubát J. 2005. Organic and mineral fertilization, respectively, increase and decrease the development of external mycelium of arbuscular mycorrhizal fungi in a long-term field experiment. *Mycorrhiza* 16: 159–166.
- Halekoh U, Højsgaard S. 2014. A kenward-roger approximation and parametric bootstrap methods for tests in linear mixed models—the R package pbrkrtest. *Journal of Statistical Software* 59: 1–32.
- Hao X, Jiang R, Chen T. 2011. Clustering 16S rRNA for OTU prediction: a method of unsupervised Bayesian clustering. *Bioinformatics* 27: 611–618.
- Hart MM, Reader RJ. 2002. Taxonomic basis for variation in the colonization strategy of arbuscular mycorrhizal fungi. *New Phytologist* 153: 335–344.
- Helgason T, Daniell TJ, Husband R, Fitter AH, Young JPW. 1998. Ploughing up the wood-wide web? *Nature* 394: 431.
- Helgason T, Fitter AH, Young JPW. 1999. Molecular diversity of arbuscular mycorrhizal fungi colonising *Hyacinthoides non-scripta* (bluebell) in a seminatural woodland. *Molecular Ecology* 8: 659–666.
- Hendershot WH, Lalonde H, Duquette M. 2008. Ion exchange and exchangeable cations. In: Carter MR, Gregorich E, eds. *Soil sampling and methods of analysis*. Boca Raton, FL, USA: Canadian Society of Soil Science and CRC Press, 173–178.
- Herman DJ, Firestone MK, Nuccio E, Hodge A. 2012. Interactions between an arbuscular mycorrhizal fungus and a soil microbial community mediating litter decomposition. *FEMS Microbiology Ecology* 80: 236–247.
- Herrera R, Merida T, Stark N, Jordan CF. 1978. Direct phosphorus transfer from leaf litter to roots. *Naturwissenschaften* 65: 208–209.
- Hodge A. 2014. Interactions between arbuscular mycorrhizal fungi and organic material substrates. *Advances in Applied Microbiology* 89: 47–99.
- Hodge A, Fitter AH. 2010. Substantial nitrogen acquisition by arbuscular mycorrhizal fungi from organic material has implications for N cycling. *Proceedings of the National Academy of Sciences, USA* 107: 13 754–13 759.
- Hodge A, Helgason T, Fitter AH. 2010. Nutritional ecology of arbuscular mycorrhizal fungi. *Fungal Ecology* 3: 267–273.
- Husband R, Herre EA, Turner SL, Gallery R, Young JPW. 2002. Molecular diversity of arbuscular mycorrhizal fungi and patterns of host association over time and space in a tropical forest. *Molecular Ecology* 11: 2669–2678.
- Ihrmark K, Bödeker ITM, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y, Stenlid J, Brandström Durling M, Clemmensen KE *et al.* 2012. New primers to amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiology Ecology* 82: 666–677.
- Johnson D, Krsek M, Wellington EMH, Stott AW, Cole L, Bardgett RD, Read DJ, Leake JR. 2005. Soil invertebrates disrupt carbon flow through fungal networks. *Science* 309: 1047.
- Johnson N. 1993. Can fertilization of soil select less mutualistic mycorrhizae? *Ecological Applications* 3: 749–757.
- Johnson NC. 2010. Resource stoichiometry elucidates the structure and function of arbuscular mycorrhizas across scales. *New Phytologist* 185: 631–647.
- Johnson NC, Angelard C, Sanders IR, Kiers ET. 2013. Predicting community and ecosystem outcomes of mycorrhizal responses to global change. *Ecology Letters* 16: 140–153.
- Johnson NC, Rowland D, Corkidi L, Egerton-Warburton L. 2003. Nitrogen enrichment alters mycorrhizal allocation at five mesic to semiarid grasslands. *Ecology* 84: 1895–1908.
- Katoh K, Misawa K, Kuma K-I, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30: 3059–3066.
- Kausserud H, Kumar S, Brysting AK, Nordén J, Carlsen T. 2011. High consistency between replicate 454 pyrosequencing analyses of ectomycorrhizal plant root samples. *Mycorrhiza* 22: 309–315.
- Kaye JP, Hart SC. 1997. Competition for nitrogen between plants and soil microorganisms. *Trends in Ecology & Evolution* 12: 139–143.
- Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, Blomberg SP, Webb CO. 2010. Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* 26: 1463–1464.

- Klironomos JN. 2003. Variation in plant response to native and exotic arbuscular mycorrhizal fungi. *Ecology* 84: 2292–2301.
- Kohout P, Sudová R, Janoušková M, Čtvrtlíková M, Hejda M, Pánková H, Slavíková R, Štajerová K, Vosátka M, Sýkorová Z. 2014. Comparison of commonly used primer sets for evaluating arbuscular mycorrhizal fungal communities: is there a universal solution? *Soil Biology and Biochemistry* 68: 482–493.
- Krüger M, Krüger C, Walker C, Stockinger H, Schüßler A. 2012. Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. *New Phytologist* 193: 970–984.
- Leff JW, Wieder WR, Taylor PG, Townsend AR, Nemergut DR, Grandy AS, Cleveland CC. 2012. Experimental litterfall manipulation drives large and rapid changes in soil carbon cycling in a wet tropical forest. *Global Change Biology* 18: 2969–2979.
- Legendre P, Legendre L. 2012. *Numerical ecology*. Elsevier Science. [www document] URL [http://store.elsevier.com/Numerical-Ecology/P\\_Legendre/ISBN-9780444538680/](http://store.elsevier.com/Numerical-Ecology/P_Legendre/ISBN-9780444538680/) [accessed 22 December 16].
- Leigh EG Jr. 1999. *Tropical forest ecology*. Oxford, UK: Oxford University Press.
- Leigh J, Hodge A, Fitter AH. 2009. Arbuscular mycorrhizal fungi can transfer substantial amounts of nitrogen to their host plant from organic material. *New Phytologist* 181: 199–207.
- Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* 15: 550.
- Maherali H, Klironomos JN. 2007. Influence of phylogeny on fungal community assembly and ecosystem functioning. *Science* 316: 1746–1748.
- McGonigle TP, Miller MH, Evans DG, Fairchild GL, Swan JA. 1990. A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. *New Phytologist* 115: 495–501.
- McMurdie PJ, Holmes S. 2014. Waste not, want not: why rarefying microbiome data is inadmissible. *PLoS Computational Biology* 10: e1003531.
- Mosse B, Phillips JM. 1971. The influence of phosphate and other nutrients on the development of vesicular-arbuscular mycorrhiza in culture. *Journal of General Microbiology* 69: 157–166.
- Munkvold L, Kjoller R, Vestberg M, Rosendahl S, Jakobsen I. 2004. High functional diversity within species of arbuscular mycorrhizal fungi. *New Phytologist* 164: 357–364.
- Nilsson LO, Bååth E, Falkengren-Grerup U, Wallander H. 2007. Growth of ectomycorrhizal mycelia and composition of soil microbial communities in oak forest soils along a nitrogen deposition gradient. *Oecologia* 153: 375–384.
- Nottingham AT, Turner BL, Winter K, Van Der Heijden MGA, Tanner EVJ. 2010. Arbuscular mycorrhizal mycelial respiration in a moist tropical forest. *New Phytologist* 186: 957–967.
- Oksanen J, Blanchet FG, Kindt R, Legendre P. 2010. *Multivariate analysis of ecological communities in R: vegan tutorial. R package version 1.17-0*. [WWW document] URL <http://vegan.r-forge.r-project.org> [accessed 21 December 2016].
- Olsson PA. 1999. Signature fatty acids provide tools for determination of the distribution and interactions of mycorrhizal fungi in soil. *FEMS Microbiology Ecology* 29: 303–310.
- Öpik M, Zobel M, Cantero JJ, Davison J, Facelli JM, Hiiesalu I, Jairus T, Kalwij JM, Koorem K, Leal ME *et al.* 2013. Global sampling of plant roots expands the described molecular diversity of arbuscular mycorrhizal fungi. *Mycorrhiza* 23: 411–430.
- Oren R, Ellsworth DS, Johnsen KH, Phillips N, Ewers BE, Maier C, Schäfer KV, McCarthy H, Hendrey G, McNulty SG *et al.* 2001. Soil fertility limits carbon sequestration by forest ecosystems in a CO<sub>2</sub>-enriched atmosphere. *Nature* 411: 469–472.
- Pearson JN, Jakobsen I. 1993. Symbiotic exchange of carbon and phosphorus between cucumber and three arbuscular mycorrhizal fungi. *New Phytologist* 124: 481–488.
- Plassard C, Louche J, Ali MA, Duchemin M, Legname E, Cloutier-Hurteau B. 2011. Diversity in phosphorus mobilisation and uptake in ectomycorrhizal fungi. *Annals of Forest Science* 68: 33–43.
- Posada RH, Madrinan S, Rivera EL. 2012. Relationships between the litter colonization by saprotrophic and arbuscular mycorrhizal fungi with depth in a tropical forest. *Fungal Biology* 116: 747–755.
- Powell JR, Parrent JL, Hart MM, Klironomos JN, Rillig MC, Maherali H. 2009. Phylogenetic trait conservatism and the evolution of functional trade-offs in arbuscular mycorrhizal fungi. *Proceedings of the Royal Society of London B* 276: 4237–4245.
- R Development Core Team. 2014. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ravnskov S, Jakobsen I. 1995. Functional compatibility in arbuscular mycorrhizas measured as hyphal P transport to the plant. *New Phytologist* 129: 611–618.
- Read DJ. 1991. Mycorrhizas in ecosystems. *Experientia* 47: 376–391.
- Redecker D, Schüßler A, Stockinger H, Stürmer SL, Morton JB, Walker C. 2013. An evidence-based consensus for the classification of arbuscular mycorrhizal fungi (Glomeromycota). *Mycorrhiza* 23: 515–531.
- Reynolds HL, Vogelsang KM, Hartley AE, Bever JD, Schultz PA. 2005. Variable responses of old-field perennials to arbuscular mycorrhizal fungi and phosphorus source. *Oecologia* 147: 348–358.
- Richardson AE, Simpson RJ. 2011. Soil microorganisms mediating phosphorus availability update on microbial phosphorus. *Plant Physiology* 156: 989–996.
- Rosling A, Landeweert R, Lindahl BD, Larsson KH, Kuyper TW, Taylor AFS, Finlay RD. 2003. Vertical distribution of ectomycorrhizal fungal taxa in a podzol soil profile. *New Phytologist* 159: 775–783.
- Sayer EJ. 2006. Using experimental manipulation to assess the roles of leaf litter in the functioning of forest ecosystems. *Biological Reviews* 81: 1–31.
- Sayer EJ, Banin LF. 2016. Tree nutrient status and nutrient cycling in tropical forest – lessons from fertilization experiments. In: Goldstein G, Santiago L, eds. *Tropical tree physiology*. Switzerland: Springer International Publishing, 275–297.
- Sayer EJ, Tanner EVJ. 2010a. A new approach to trenching experiments for measuring root–rhizosphere respiration in a lowland tropical forest. *Soil Biology and Biochemistry* 42: 347–352.
- Sayer EJ, Tanner EVJ. 2010b. Experimental investigation of the importance of litterfall in lowland semi-evergreen tropical forest nutrient cycling. *Journal of Ecology* 98: 1052–1062.
- Sayer EJ, Tanner EVJ, Cheesman AW. 2006a. Increased litterfall changes fine root distribution in a moist tropical forest. *Plant and Soil* 281: 5–13.
- Sayer EJ, Tanner EVJ, Lacey A. 2006b. Effects of litter manipulation on early-stage decomposition and meso-arthropod abundance in a tropical moist forest. *Forest Ecology and Management* 229: 285–293.
- Sayer EJ, Wright SJ, Tanner EVJ, Yavitt JB, Harms KE, Powers JS, Kaspari M, Garcia MN, Turner BL. 2012. Variable responses of lowland tropical forest nutrient status to fertilization and litter manipulation. *Ecosystems* 15: 387–400.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ *et al.* 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* 75: 7537–7541.
- Schreeg LA, Mack MC, Turner BL. 2013. Nutrient-specific solubility patterns of leaf litter across 41 lowland tropical woody species. *Ecology* 94: 94–105.
- Simon L, Lalonde M, Bruns TD. 1992. Specific amplification of 18S fungal ribosomal genes from vesicular-arbuscular endomycorrhizal fungi colonizing roots. *Applied and Environmental Microbiology* 58: 291–295.
- Smith SE, Read DJ. 2008. *Mycorrhizal symbiosis*. Cambridge, UK: Academic Press.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313.
- Tanner EVJ, Sheldrake MWA, Turner BL. 2016. Changes in soil carbon and nutrients following 6 years of litter removal and addition in a tropical semi-evergreen rain forest. *Biogeosciences* 13: 6183–6190.
- Treseder K. 2004. A meta-analysis of mycorrhizal responses to nitrogen, phosphorus, and atmospheric CO<sub>2</sub> in field studies. *New Phytologist* 164: 347–355.
- Treseder KK, Allen MF. 2002. Direct nitrogen and phosphorus limitation of arbuscular mycorrhizal fungi: a model and field test. *New Phytologist* 155: 507–515.

- Turner BL. 2008. Soil organic phosphorus in tropical forests: an assessment of the NaOH–EDTA extraction procedure for quantitative analysis by solution  $^{31}\text{P}$  NMR spectroscopy. *European Journal of Soil Science* 59: 453–466.
- Turner BL, Wright SJ. 2014. The response of microbial biomass and hydrolytic enzymes to a decade of nitrogen, phosphorus, and potassium addition in a lowland tropical rain forest. *Biogeochemistry* 117: 115–130.
- Veresoglou SD, Chen B, Rillig MC. 2012. Arbuscular mycorrhiza and soil nitrogen cycling. *Soil Biology and Biochemistry* 46: 53–62.
- Vincent AG, Turner BL, Tanner EVJ. 2010. Soil organic phosphorus dynamics following perturbation of litter cycling in a tropical moist forest. *European Journal of Soil Science* 61: 48–57.
- Vitousek PM, Sanford RL. 1986. Nutrient cycling in moist tropical forest. *Annual Review of Ecology and Systematics* 17: 137–167.
- Wang GM, Stribley DP, Tinker PB, Walker C. 1993. Effects of pH on arbuscular mycorrhiza. I. Field observations on the long-term liming experiments at Rothamsted and Woburn. *New Phytologist* 124: 465–472.
- Wang Y, Naumann U, Wright ST, Warton DI. 2012. mvabund – an R package for model-based analysis of multivariate abundance data. *Methods in Ecology and Evolution* 3: 471–474.
- Warton DI, Foster SD, De'ath G, Stoklosa J. 2015. Model-based thinking for community ecology. *Plant Ecology* 261: 669–682.
- Webb C. 2000. Exploring the phylogenetic structure of ecological communities: an example for rain forest trees. *The American Naturalist* 156: 145–155.
- Webb CO, Ackerly DD, McPeck MA, Donoghue MJ. 2002. Phylogenies and community ecology. *Annual Review of Ecology and Systematics* 33: 475–505.
- Went FW, Stark N. 1968. Mycorrhiza. *BioScience* 18: 1035–1039.
- Wright SJ, Yavitt JB, Wurzburger N, Turner BL, Tanner EVJ, Sayer EJ, Santiago LS, Kaspari M, Hedin LO, Harms KE *et al.* 2011. Potassium, phosphorus, or nitrogen limit root allocation, tree growth, or litter production in a lowland tropical forest. *Ecology* 92: 1616–1625.
- Wurzburger N, Wright SJ. 2015. Fine root responses to fertilization reveal multiple nutrient limitation in a lowland tropical forest. *Ecology* 96: 2137–2146.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information tab for this article:

**Fig. S1** Effect of litter manipulation on the levels of NLFA 16:1 $\omega$ 5 in the top 10 cm of forest soil.

**Fig. S2** Rarefaction curves pooled by experimental treatment and for each sample.

**Fig. S3** Maximum-likelihood phylogenetic tree of all OTUs detected in this study.

**Fig. S4** Effect of litter manipulation on AM fungal OTU richness and predominance.

**Fig. S5** Comparison of AM fungal communities described by the quantitative Bray–Curtis metric of dissimilarity, the Jaccard presence-absence based metric of dissimilarity, and correlation between the two.

**Table S1** Response of soil physical characteristics to 9 yr of litter removal and addition in a tropical forest

**Table S2** AM fungal OTUs altered by 9 yr of leaf litter addition and removal

**Table S3** Number of sequences per sample after blasting, filtering, merging and trimming

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



## About *New Phytologist*

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <28 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit [www.newphytologist.com](http://www.newphytologist.com) to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office ([np-centraloffice@lancaster.ac.uk](mailto:np-centraloffice@lancaster.ac.uk)) or, if it is more convenient, our USA Office ([np-usaoffice@lancaster.ac.uk](mailto:np-usaoffice@lancaster.ac.uk))
- For submission instructions, subscription and all the latest information visit [www.newphytologist.com](http://www.newphytologist.com)