













Vegetation demographics in Earth System Models: A review of progress and priorities

Rosie A. Fisher¹  | Charles D. Koven² | William R. L. Anderegg³  | Bradley O. Christoffersen⁴ | Michael C. Dietze⁵  | Caroline E. Farrior⁶ | Jennifer A. Holm² | George C. Hurtt⁷ | Ryan G. Knox²  | Peter J. Lawrence¹ | Jeremy W. Lichstein⁸ | Marcos Longo⁹  | Ashley M. Matheny¹⁰  | David Medvigy¹¹  | Helene C. Muller-Landau¹² | Thomas L. Powell²  | Shawn P. Serbin¹³  | Hisashi Sato¹⁴ | Jacquelyn K. Shuman¹ | Benjamin Smith¹⁵ | Anna T. Trugman¹⁶  | Toni Viskari¹² | Hans Verbeeck¹⁷ | Ensheng Weng¹⁸  | Chonggang Xu⁴ | Xiangtao Xu¹⁹ | Tao Zhang⁸  | Paul R. Moorcroft²⁰

¹National Center for Atmospheric Research, Boulder, CO, USA

²Lawrence Berkeley National Laboratory, Berkeley, CA, USA

³Department of Biology, University of Utah, Salt Lake City, UT, USA

⁴Los Alamos National Laboratory, Los Alamos, NM, USA

⁵Department of Earth and Environment, Boston University, Boston, MA, USA

⁶Department of Integrative Biology, University of Texas at Austin, Austin, TX, USA

⁷Department of Geographical Sciences, University of Maryland, College Park, MD, USA

⁸Department of Biology, University of Florida, Gainesville, FL, USA

⁹Embrapa Agricultural Informatics, Campinas, Brazil

¹⁰Department of Geological Sciences, Jackson School of Geosciences, University of Texas at Austin, Austin, TX, USA

¹¹Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA

¹²Smithsonian Tropical Research Institute, Panamá, Panamá

¹³Environmental and Climate Sciences Department, Brookhaven National Laboratory, Upton, NY, USA

¹⁴Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, Japan

¹⁵Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

¹⁶Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, NJ, USA

¹⁷Department of Applied Ecology and Environmental Biology, Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium

¹⁸Center for Climate Systems Research, Columbia University, New York, NY, USA

¹⁹Department of Geosciences, Princeton University, Princeton, NJ, USA

²⁰Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA

Correspondence

Rosie A. Fisher, National Center for Atmospheric Research, Boulder, CO, USA.
Email: rfisher@ucar.edu

Funding information

Division of Atmospheric and Geospace Sciences; Biological and Environmental Research; NASA Terrestrial Ecology, Grant/Award Number: NNX14AH65G; United States Department of Energy, Grant/Award

Abstract

Numerous current efforts seek to improve the representation of ecosystem ecology and vegetation demographic processes within Earth System Models (ESMs). These developments are widely viewed as an important step in developing greater realism in predictions of future ecosystem states and fluxes. Increased realism, however, leads to increased model complexity, with new features raising a suite of ecological questions that require empirical constraints. Here, we review the developments that

Number: DE-SC0012704; National Science Foundation; FAPESP, Grant/Award Number: 2015/07227-6; U.S. National Science Foundation Hydrological Science, Grant/Award Number: 1521238; USDA, Grant/Award Number: 11-JV-112423-059, 16-JV-11242306-050; US Department of Energy, Office of Science, Office of Biological and Environmental Research, Terrestrial Ecosystem Science (TES), Grant/Award Number: DE-SC0014363

permit the representation of plant demographics in ESMs, and identify issues raised by these developments that highlight important gaps in ecological understanding. These issues inevitably translate into uncertainty in model projections but also allow models to be applied to new processes and questions concerning the dynamics of real-world ecosystems. We argue that stronger and more innovative connections to data, across the range of scales considered, are required to address these gaps in understanding. The development of first-generation land surface models as a unifying framework for ecophysiological understanding stimulated much research into plant physiological traits and gas exchange. Constraining predictions at ecologically relevant spatial and temporal scales will require a similar investment of effort and intensified inter-disciplinary communication.

KEYWORDS

carbon cycle, demographics, dynamic global vegetation models, Earth System Model, ecosystem, vegetation

1 | INTRODUCTION

Ecological demographic processes govern terrestrial vegetation structure, and vegetation structure influences climatically important fluxes of carbon, energy, and water (Bonan, 2008). Better representation of vegetation demography in Earth System Models (ESMs) has repeatedly been identified as a critical step toward a more realistic representation of biologically mediated feedbacks in modeling future climates (Evans, 2012; Moorcroft, 2006; Moorcroft, Hurtt, & Pacala, 2001; Purves & Pacala, 2008; Thomas, Brookshire, & Gerber, 2015). Model-data comparison is greatly assisted by increasingly realistic model abstraction methods. Similarly, a greater range of data can be used for parameterization and initialization, and in some cases, models improvements can be directly linked to better simulation of biodiversity (Levine et al., 2016). These improvements are traded off against increasing complexity and computation expense.

Dynamic global vegetation models (DGVMs) are the components of land surface models (LSMs) that try to predict the global distribution of vegetation types from physiological principles (Cao & Woodward, 1998; Foley et al., 1996; Sitch et al., 2003; Woodward & Lomas, 2004). Traditionally, DGVMs represent plant communities using a single area-averaged representation of each plant functional type (PFT) for each climatic grid cell. This simplification and the resulting computational efficiency has allowed first generation DGVMs (hereafter g1DVMs) to be broadly adopted within ESMs (Arora & Boer, 2010; Bonan, Levis, Sitch, Vertenstein, & Oleson, 2003; Cox, 2001; Krinner et al., 2005).

This level of abstraction means, however, that g1DVMs do not capture many demographic processes considered important for the accurate prediction of ecosystem composition and function, including canopy gap formation, vertical light competition, competitive exclusion, and successional recovery from disturbance (Feeley et al., 2007; Hurtt, Moorcroft, Pacala, & Levin, 1998; Moorcroft et al., 2001; Smith, Prentice, & Sykes, 2001; Stark et al., 2012).

In contrast, forest gap (Bugmann, 2001; Dietze & Latimer, 2011) and “individual-based” models (IBMs) (Christoffersen et al., 2016; Fischer et al., 2016; Fyllas et al., 2014; Sato, Ito, & Kohyama, 2007; Shuman, Shugart, & Krankina, 2014; Smith et al., 2001) represent vegetation at the level of individual plants. IBMs represent spatial variability in the light environment and thereby simulate competitive exclusion, succession, and coexistence of tree species (Pacala et al., 1996; Smith et al., 2001). Simulation of individual trees in a spatially explicit, stochastic framework incurs a notable computational penalty, however. These challenges are typically addressed by limiting the spatial scope (Sakschewski et al., 2015), temporal frequency, and/or reduced sampling of the potential ensemble of model outcomes (Sato et al., 2007; Smith et al., 2014).

As a compromise between the abstraction of g1DVMs and the computational expense of IBMs, many groups have developed “cohort-based” models, whereby individual plants with similar properties (size, age, functional type) are grouped together (Haverd et al., 2013; Hurtt et al., 1998; Lischke, Zimmermann, Bolliger, Rickebusch, & Löffler, 2006; Medvigy, Wofsy, Munger, Hollinger, & Moorcroft, 2009; Moorcroft et al., 2001; Scherstjanoi, Kaplan, & Lischke, 2014; Smith et al., 2014; Weng et al., 2015). The cohort approach retains the dynamics of IBMs, with reduced computational cost, but removes stochastic processes that can enhance the representation of functional diversity (Fisher et al., 2010).

Herein we refer to both individual and cohort-based models as “vegetation demographic models” (VDMs). We define VDMs as a special class of DGVM, which include representation/tracking of multiple size-classes or individuals of the same PFT, which can encounter multiple light environments within a single climatic grid cell. We adopt this terminology since both individual and cohort models present similar opportunities and challenges as they are implemented within ESMs.

As in first-generation models the distributions of PFTs, and their associated traits, can be geographically and temporally “filtered” in

VDMs via the mechanisms of competition, differential recruitment and mortality. In VDMs, however, disturbance history and vertical light competition modulate interactions between plant traits and resource acquisition. Furthermore, it is typical (but not universally the case) that a priori constraints on distribution (climate envelopes) are removed (Fisher et al., 2015). Vegetation structure and distribution thus become entirely emergent model properties of ascribed plant functional traits and their interactions with abiotic environmental conditions. In addition, VDMs provide critical new opportunities for data-model integration owing to their higher fidelity representation of the structure of vegetation stands, as we will discuss in this review.

Several efforts to embed VDMs within ESMs are now coming to fruition, but understanding of their provenance, function and uncertainties remains specialized knowledge even within the land surface modeling community. In this review, we focus on those models currently resident in ESMs or regional atmospheric models. Many “offline” VDMs exist, with novel and beneficial approaches (e.g., Fyllas et al., 2014; Haverd et al., 2013; Pavlick, Drewry, Bohn, Reu, & Kleidon, 2013; Sakschewski et al., 2015; Scheiter, Langan, & Higgins, 2012; Scherstjanoi, Kaplan, Thürig, & Lischke, 2014). We focus on the particular challenges of large-scale implementations, coupled to atmospheric models, since this process imposes important boundary conditions on VDM functionality. For example, ESMs typically require land-atmosphere fluxes of carbon, water, and energy at hourly or sub-hourly timescales, and these must be in exact balance to prevent erroneous model drift. Implementation must be global in principle, imposing computational restrictions, and many other processes (hydrology, snow, lakes, urban areas, biogeochemical cycles, land-use change) must be simulated consistently.

Underpinning the implementation of VDMs in LSMs are a set of processes whose representations require significant modification (compared to first-generation DGVMs) to provide the appropriate function in the context of multi-layer, multi-PFT ecosystems. These include the partitioning of light and other resources between individuals or cohorts, the representation of ecophysiological processes involved in carbon and nutrient uptake, allocation, mortality, and recruitment within the newly resolved model dimensions, and the interpretations of land use, fire, and other disturbances. These new model structures pose several research challenges and opportunities.

In this review, which is the first to bring together expertise from such a diverse range of VDM groups, we:

1. summarize the state-of-the-art of VDM development,
2. discuss model features specific to VDMs and alternative assumptions currently used,
3. detail datasets available for validation and benchmarking, and
4. outline future code development and data collection strategies needed to better constrain these new model elements.

We hope to both motivate research aimed at informing the representation of plant ecology in ESMs and highlight gaps in basic ecological theory that are now at the front line of simulating the biosphere's role in the climate system. We argue that it is too early to assert that any one methodology is the ideal representation of plant demographics to use within ESMs. Rather, until relevant data and knowledge gaps are filled through concerted empirical and model-based research, we expect that an ensemble of techniques will allow for more robust predictions of likely trajectories of vegetation structural changes, their impacts on biogeochemistry and climate feedbacks (Koven et al., 2015; Sanderson, Knutti, & Caldwell, 2015).

2 | PROGRESS TO DATE

In this section we detail the ongoing progress of implementing vegetation demographic models (VDMs) inside ESMs on a loose continuum from individual- to cohort-based approaches (Table 1).

2.1 | SEIB-DGVM

The SEIB-DGVM (Spatially-Explicit Individual-Based Dynamic Global Vegetation Model, <http://seib-dgvm.com>) is an IBM, representing variability in light in both the vertical and horizontal dimensions. Following initial implementation at a global scale (Sato et al., 2007), SEIB-DGVM has been modified to represent plant population dynamics and biogeochemistry in south-east Asia (Sato, 2009), Africa (Sato & Ise, 2012), and Siberia (Sato, Kobayahi, & Delbart, 2010).

TABLE 1 Table of attributes of vegetation demographics models discussed in this paper

Model acronym	Name	Vegetation representation	Coupled to ESM?	Stochastic?	Canopy structure	Disturbance history patches?
SEIB	Spatially Explicit Individual-Based model	Individual	MIROC-ESM	Yes	Individuals	No
LPJ-GUESS	Lund-Potsdam-Jena General Ecosystem Simulator	Individual or Cohort	EC-Earth, RCA-GUESS	Yes (optional for some processes)	Flat-top	Yes
LM3-PPA	Perfect Plasticity Approximation	Cohort	GFDL-ESM	No	PPA	No
ED	Ecosystem Demography model	Cohort	RAMS	No	Flat-top	Yes
ED2	Ecosystem Demography model v2	Cohort	RAMS	No	Flat-top	Yes
CLM(ED)	Community Land Model with Ecosystem Demography	Cohort	CESM	No	PPA	Yes

SEIB simulates a 30 m × 30 m patch of forest, where individual trees establish, compete, and die. Each tree is composed of a cylindrical crown and trunk, plus fine roots. Tree crowns are horizontally sliced into 10 cm deep “disks,” for which photosynthesis is calculated separately with a daily physiological timestep. Leaf area is updated daily by turnover and growth. Crowns of different trees do not occupy the same physical space. To represent spatial plasticity, crowns are able to grow a given distance horizontally in response to light availability each year.

SEIB-DGVM is implemented within the MIROC-ESM (Watanabe et al., 2011). In the ESM context, SEIB-DGVM is run once for each grid cell, representing one particular integration of the stochastic forest, to allow global applications.

2.2 | LPJ-GUESS

The Lund-Potsdam-Jena General Ecosystem Simulator (Smith et al., 2001, 2014) is also an IBM, but with multiple patches accounting for stochastic heterogeneity in composition and structure arising from succession following stand-destroying disturbance. Both “individual” and more commonly used “cohort” modes are implemented. In the cohort mode, tree or shrub individuals of the same age and PFT within patch are grouped together and simulated as an average individual, scaled to patch level via cohort density. Multiple PFTs may occur within a single patch, and compete for light, water, and nitrogen. Photosynthesis, stomatal conductance, phenology, turnover, and allocation follow LPJ-DGVM (Sitch et al., 2003), with the addition (LPJ-GUESSv3.0 onward) of nitrogen cycling (Smith et al., 2014). The model includes the wildfire scheme of Thonicke, Venevsky, Sitch, and Cramer (2001), and a new representation of fire dynamics is in development.

LPJ-GUESS is coupled to the RCA4 regional climate model (Smith, Samuelsson, Wramneby, & Rummukainen, 2011; Wramneby, Smith, & Samuelsson, 2010). It also accounts for land cover dynamics and carbon cycling within the EC-EARTH ESM (Hazeleger et al., 2010; Weiss et al., 2014). Daily meteorological fields are input to LPJ-GUESS and adjustments in leaf area index for separate “high” and “low” vegetation tiles, averaged across patches, are returned to the land surface physics scheme. Impacts on energy and water exchange with the atmosphere are manifested via albedo, evapotranspiration, and surface roughness length. CO₂ is exchanged daily with the atmospheric transport model.

2.3 | LM3-PPA

The Geophysical Fluid Dynamics Laboratory (GFDL) Land Model 3 with the Perfect Plasticity Approximation (LM3-PPA) is a cohort-based VDM (Weng et al., 2015). The PPA assumes that tree crowns “perfectly” fill canopy gaps through phototropism (plasticity) (Strigul, Pristinski, Purves, Dushoff, & Pacala, 2008). Crowns thus self-organize into discrete canopy layers, within which all plants receive the same incoming radiation. The LM3-PPA model extends earlier work on simpler tractable PPA models (Farrior, Bohlman, Hubbell, & Pacala, 2016; Farrior, Dybzinski, Levin, & Pacala, 2013) to include prognostic energy, water, and carbon cycling. The simpler PPA

models allows ecosystem scale consequences of plant strategies to be rapidly predicted, allowing the properties of the complex model to be investigated in greater depth (Weng et al., 2015).

LM3-PPA successfully captured observed successional dynamics of one site in Eastern US temperate forest (Weng et al., 2015), and the changing relative abundances of deciduous and evergreen strategies over succession in three sites spanning temperate to boreal zones in North America (Weng, Farrior, Dybzinski, & Pacala, 2016). Coupling to the GFDL ESM for site-level simulations has been completed, and global implementation is currently in progress.

2.4 | Ecosystem Demography models

The Ecosystem Demography (ED) concept is also a cohort-based representation of vegetation dynamics (Hurt et al., 1998; Moorcroft et al., 2001). In contrast to the LM3-PPA, ED discretizes the simulated landscape into spatially implicit “patches” according to “age since last disturbance,” capturing the dynamic matrix of disturbance-recovery processes within a typical forest ecosystem in a deterministic manner (in contrast to LPJ-GUESS and SEIB). Within patches, individuals are grouped into cohorts by PFT and height class, and height-structured competition for light between cohorts drives successional dynamics. ED uses a patch fusion/fission scheme to track the landscape-scale age distribution resulting from disturbance. During fission, disturbance splits patch areas into undisturbed and disturbed fractions. During fusion, (to keep the number of patches from growing exponentially), patches that are similar in structure are merged. No subgrid geographic information is retained. Similar fusion/fission routines exist for cohorts. At least three derivatives of the original ED concept have emerged since its inception, including:

2.5 | ED

One implementation of the Ecosystem Demography concept (currently known solely as “ED”), was developed from (Moorcroft et al., 2001) applied to the U.S by Hurt et al. (2004) and Albani, Medvigy, Hurt, and Moorcroft (2006), and is now a global model (Fisk, 2015). Advances in this version have focused on the inclusion of land-use as driver of demography (Hurt et al., 2006), transient effects of tropical cyclones (Fisk et al., 2013), plant migration in response to climate change (Flanagan et al., 2016), and detailed use of vegetation structure to initialize and test ecosystem dynamics (Fisk, 2015; Hurt, Thomas, Fisk, Dubayah, & Sheldon, 2016; Hurt et al., 2004, 2010; Thomas, Hurt, Dubayah, & Schilz, 2008). This branch of ED has also been coupled to the RAMS mesoscale atmospheric model (Roy, Hurt, Weaver, & Pacala, 2003) and the GCAM integrated Assessment Model (Fisk, 2015) and also serves as base model of the NASA Carbon Monitoring System, and the NASA-GEDI mission (Dubayah et al., 2014).

2.6 | ED2

The Ecosystem Demography Model v2 (ED2) (Medvigy et al., 2009, <https://github.com/EDmodel/ED2>) also uses the scaling

concepts of Moorcroft et al. (2001), with numerous subsequent developments. In ED2, grid cells are further disaggregated by similar edaphic conditions. Heterogeneity in light environment and canopy structure is integrated into the biophysical scheme, giving rise to differentiated horizontal and vertical micro-environments within grid-cells that vary in temperature, humidity, soil moisture, and soil nutrient conditions. Recent developments include a plant hydrodynamic scheme, drought-deciduous phenology (Xu, Medvigy, Powers, Becknell, & Guan, 2016), nitrogen fixers, boreal-specific PFTs, dynamic soil organic layers, and trait-based recruitment (Trugman et al., 2016).

ED2 has been tested in boreal (Trugman et al., 2016), temperate (Antonarakis, Munger, & Moorcroft, 2014; Medvigy, Jeong, Clark, Skowronski, & Schäfer, 2013; Medvigy & Moorcroft, 2012; Medvigy et al., 2009), tropical (Kim et al., 2012; Levine et al., 2016; Xu et al., 2016; Zhang et al., 2015), tundra (Davidson et al., 2009), agricultural (Lokupitiya et al., 2016), and biofuel systems (LeBauer, Wang, Richter, Davidson, & Dietze, 2013). It has also been applied to ecosystems undergoing disturbance events such as fire, drought, elevated CO₂, land-use change, and insect defoliation (Medvigy, Clark, Skowronski, & Schäfer, 2012; Miller, Dietze, DeLucia, & Anderson-Teixeira, 2016; Trugman et al., 2016; Zhang et al., 2015). ED2 is coupled to the Regional Atmospheric Modeling System (RAMS) (Knox et al., 2015; Swann, Longo, Knox, Lee, & Moorcroft, 2015).

2.7 | CLM(ED)

CLM(ED) (Fisher et al., 2015) is a variant of the Community Land Model (CLM) (Lawrence et al., 2011; Oleson et al., 2013), integrating the ED concept within the architecture of the Community Earth System Model (Hurrell et al., 2013). CLM(ED) includes a merging of the ED and PPA concepts, allocating cohorts to canopy and understorey layers. It deviates from the standard PPA as it does not allocate canopy levels according to a definitive height threshold (z^* , Purves, Lichstein, Strigul, & Pacala, 2008; Strigul et al., 2008) and instead splits growing cohorts between canopy layers—the fraction of each cohort remaining in the canopy a continuous function of height (in principle increasing the probability of coexistence, Fisher et al., 2010). Canopy biophysics, hydrology, photosynthesis, and respiration all follow CLM4.5 (Oleson et al., 2013) subject to disaggregation into cohort-level fluxes. CLM(ED) includes new representations of phenology and carbon storage and a modified SPITFIRE fire model (Thonicke et al., 2010). CLM(ED) was applied regionally, focusing on the sensitivity of biome boundaries to plant trait representation (Fisher et al., 2015) and will be re-named FATES (Functionally Assembled Terrestrial Ecosystem Simulated) in future references.

3 | APPROACHES TO MODEL STRUCTURE AND PROCESS REPRESENTATION

Historically, demographic models (typically IBMs) have been distinct from models with detailed plant physiological representation. In the

VDMs discussed here, however, demographics (recruitment, growth, mortality) arise primarily as functions of physiological functions and so the two are intimately linked. Thus, design decisions in physiological algorithms have potentially critical impacts on the emergent population dynamics. In this section, we discuss the process modifications that are required when moving from a g1DVM model to a size-structured VDM. These include higher-order representation of competition for light, water and nutrients, demographic processes (recruitment, mortality) and disturbance (fire, land use). Our intention is to (i) illustrate the logic behind the inclusion of new model features, (ii) highlight process uncertainties that remain or emerge (by way of motivating new research themes), and (iii) provide context for the following discussion of model evaluation data.

3.1 | Competition for light

Land surface models calculate radiation partitioning and the within-canopy radiation regime using radiative transfer models (RTM). RTMs simulate the reflectance, interception, absorption, and transmission (into the ground) of shortwave radiation (0.3–2.5 microns) through a canopy comprised of scattering elements (leaves, wood, soil, and snow). Incoming radiation is typically partitioned into direct and diffuse streams. Interception of direct radiation by scattering elements results in reflected and transmitted fluxes of diffuse radiation. Upwards-reflected diffuse radiation affects leaves higher in the canopy, preventing a simple solution to the partitioning of energy. To resolve this, iterative methods calculate upwards and downwards diffuse fluxes until a solution is reached (Goudriaan, 1977; Norman, 1979). Other approaches use the “two-stream approximation” (Dickinson, 1983; Sellers, 1985), a system of two coupled ordinary differential equations which can be analytically solved (Liou, 2002) assuming a continuum with homogenous reflectance and transmission characteristics. The two-stream approximation is used for a single canopy divided into sun/shade fractions in CLM4.5 (Bonan et al., 2011) and as a set of canopy layers each with its own 2-stream solution for JULES (Mercado, Huntingford, Gash, Cox, & Jogireddy, 2007), ORCHIDEE-CAN (Naudts et al., 2015), ED2 (Medvigy et al., 2009) and CLM(SPA) (Bonan, Williams, Fisher, & Oleson, 2014).

In VDMs, the RTM is necessarily more complex than standard LSMs because of the need to have more than one PFT within a given vertical structure canopy, invalidating the homogeneity assumptions of the two-stream model. VDMs must further determine (i) how to partition leaves and stems of cohorts/individuals into discrete scattering elements within which there can be an assumption of homogeneity and (ii) how to arrange these scattering elements relative to each other, to discern the influence of plant height on radiation interception. The details of how these issues are resolved control the nature of the feedback between plant size and light acquisition, and thus are a pivotal component of any trait-filtering architecture (Fisher et al., 2010; Scheiter et al., 2012).

Solving these problems efficiently is a significant component of the implementation of VDMs in ESMs. Here, we summarize the




FIGURE 1 Organization of canopy schemes in four vegetation demographic models. Shades of yellow represent incident light levels, whereas shades of gray indicate alternative plant functional types (PFTs). Boxes represent cohorts as represented by ED & ED2, LM3-PPA, and CLM(ED). Dotted cohort boundaries denote cohorts that belong to the understory, all of which receive identical light levels, in the PPA schemes of the LM3-PPA and CLM(ED) models. Note that in the LM3-PPA there can be more than one understory layer, but in CLM(ED) there cannot. In the cohort-based schemes, horizontal positioning is for illustrative purposes only and not represented by the model, which is one-dimensional. Dotted lines in the CLM(ED) figure illustrate within-canopy leaf levels resolved by the radiation transfer scheme. In the LM3-PPA, “ z^* ” indicates the cohort height above which canopy/understory status is defined. In the CLM(ED), there is no “ z^* ” threshold, and larger cohorts in the understory may in principle be taller than the shorter cohorts in the canopy layer (reflecting imperfect competition processes, per Fisher et al., 2010). Note that for ED-derived models (ED, ED2, CLM(ED)), cohort organization is illustrated only for a single patch, though each model represents a multitude of patches having different ages since disturbance within a single site

status of the existing schemes, their advantage and disadvantages, and suggest ways in which these might be developed further. Alternative schemes are represented in Figure 1.

3.2 | Individual-based approach: SEIB

In SEIB-DGVM, each individual crown has an x-y location in space, and shading of trees by their neighbors is explicitly simulated. Direct and diffuse photosynthetically active radiation (PAR) are estimated for each crown disk. For diffuse PAR, all disks at the same height receive the same radiation, attenuated by the leaf area index (LAI) above each disk using Beer’s law (Goudriaan, 1977). For direct light, a “virtual cylinder” is calculated for each canopy disc. The cylinder extends South, at $0.86 \times$ the midday solar angle (Sato et al., 2007), and available PAR is attenuated (also using Beer’s law) by the leaves located within the cylinder. The grass layer is horizontally divided into 1×1 m cells, each of which receives PAR attenuated by the LAI above. Some simplifications are employed to efficiently simulate individual trees (daily timestep, static solar angle, few or no replicates). In contrast, cohort models (below) have a lower computational footprint, but must designate rules by which light is distributed to cohorts of differing height in the absence of direct spatial competition.

3.3 | Infinitely thin flat crowns: ED, ED2

Perhaps the most straightforward method for representing how cohort leaves are aligned with respect to incoming light is the “flat-top crown” idea; wherein the total leaf area of each cohort is conceptually distributed evenly across the entire canopy area of a patch (one infinitely thin layer). The cohort-layers are stacked vertically and the two-stream model is used to determine radiation absorbed by each layer at its midpoint. Each cohort thus is shaded by all taller cohorts.

The flat-top method is relatively straightforward to implement, but suffers from the biologically unrealistic outcome that marginally taller cohorts outcompete their neighbors in terms of light availability. This can lead to systematic growth biases (compared to observations) where the tallest trees grow too fast and next-tallest trees more slowly, making coexistence of multiple PFTs more difficult to achieve (Fisher et al., 2010). Furthermore, there is no representation of the effects of space on canopy structure (Farrion et al., 2016). In ED2, these negative effects have been partially mitigated by (i) the consideration of cohort crown area, which allows partial, rather than complete, shading among cohorts and, (ii) a cohort splitting algorithm that prevents a single cohort from accumulating a leaf area index above a predetermined maximum LAI threshold.

3.4 | Vertically overlapping crowns: LPJ-GUESS

LPJ-GUESS adopts an approach similar to ED2 but with three-dimensional crowns evenly distributed across the area (ca. 0.1 ha) of each stochastic patch, and uniformly distributed in the vertical dimension from ground level up to the current maximum height of each individual or cohort. A multilayer Beer's law integration partitions incoming PAR among cohorts by relating light absorbance of each layer to that layer's leaf area using a single fixed extinction coefficient (Smith et al., 2001). A herbaceous layer captures PAR not absorbed by the canopy.

3.5 | Perfect plasticity approximation: LM3-PPA

The PPA assumes that trees can forage for light in a "perfectly plastic" manner horizontally within a patch. Starting with the tallest tree, the crown area of each successively shorter tree is assigned to the canopy layer until the cumulative canopy crown area equals or exceeds the patch ground area. Once the "canopy" is filled with tree crowns, the next shorter trees inhabit the first understory layer and are shaded by the trees in the canopy. If the first understory layer is also full, then additional understory layers are created. All crowns in the same layer receive the same incoming radiation streams, and penetration of light through each crown is determined using Beer's law (Weng et al., 2015). The radiation streams penetrating the crowns of a layer are averaged before passing to the next lower layer. Light reflected by the soil can be absorbed by the leaves above. Physiologically-based PPA models (Farrion, et al., 2013; Farrion, Rodriguez-Iturbe, Dybzinski, Levin, & Pacala, 2015; Dybzinski, Farrion, Wolf, Reich, & Pacala, 2011; Weng et al., 2015) include a gap fraction parameter (η), whereby only $1-\eta$ of each layer can be filled. This allows more light to reach the understory, and thus more realistic understory behavior, but does not fundamentally change the PPA algorithm.

3.6 | Discretized PPA: CLM(ED)

The CLM(ED) follows a similar logic to the PPA, and resolves radiation streams between canopy layers, and also discretizes direct and diffuse fluxes into "leaf layers" within cohorts. To reduce computational intensity, cohorts are merged together for flux calculations, such that all leaves of each PFT are represented by a three-dimensional matrix of PFT, canopy layer, and leaf layer. An iterative, layered solution, following Norman, Perry, Fraser, and Mach, (1979), calculates equilibrium upwards and downwards radiation fluxes.

3.7 | Modifying extreme assumptions

The existing methodologies for the division of solar radiation in cohort models represent two extreme assumptions at either end of a continuum. The flat-top assumption implies that small differences in relative height lead to large changes in light availability (within a patch), whereas the PPA assumption means that

differences in height, however extreme, only affect light availability at the boundaries between canopy layers (canopy vs. understory). In reality, canopy trees all receive equivalent light from above, but the amount of lateral light they receive depends on their height relative to their neighbors. An ideal framework might include the capacity of the PPA to represent the impact of space on competition for overhead light, while adding change in lateral light availability with height among canopy trees. A spatially-implicit scheme that could capture both of these features would enhance the ability of VDMs to capture size-related variation in light availability, and thus presumably size-related variation in growth and survival.

3.8 | Water acquisition and its influence on plant physiology

In comparison with competition for light, competition for water is less well-understood, and model representations remain poorly developed, especially in terms of how water is distributed among competing plants, as well as in how water acquisition affects plant processes.

3.9 | Representing competition for water

3.9.1 | Shared vs. partitioned water resources

Models use contrasting assumptions of how water resources are divided horizontally within a grid-cell. Some VDMs represent a single "pool" of water from which all plants draw equally (CLM(ED), LM3-PPA, SEIB-DGVM, Figure 2b). Other models (ED2, LPJ-GUESS) divide water resources by patch (Figure 2a). Real ecosystems are unlikely to conform to either of these simplifications. In reality, whether water resources are shared across patches depends on the relative characteristic length scales of disturbance and of horizontal water redistribution (Jupp & Twiss, 2006). Where canopy gaps and thus patches are small, we might expect that water would be effectively shared between patches of different ages; in contrast, where disturbance events are larger-scale (blow-downs, fires) and patches accordingly larger, we might expect little or no such water sharing. No VDMs track length scales or arrangement of disturbance events by default, nor do they represent inter-patch water fluxes. The impact of this type of effect can be important for the surface energy balance (Shrestha, Arora, & Melton, 2016).

3.9.2 | Spatial aggregation of water resources

A rich literature exists on the spatial partitioning of water resources in semi-arid regions (Borgogno, D'Odorico, Laio, & Ridolfi, 2009; D'Odorico, Caylor, Okin, & Scanlon, 2007; Gilad, Shachak, & Meron, 2007; Meron, 2011; Meron, Gilad, von Hardenberg, Shachak, & Zarmi, 2004; Rodriguez-Iturbe, D'Odorico, Porporato, & Ridolfi, 1999; Scanlon, Caylor, Levin, & Rodriguez-Iturbe, 2007; van Wijk & Rodriguez-Iturbe, 2002), the focus of which is the tendency for soil




FIGURE 2 Illustration of unresolved belowground resource partitioning issues in vegetation demographic models. (a, b) show two alternative depictions of resource partitioning in an age-since-disturbance resolving (ED-type) model. In (a) resources (water/nutrients) are resolved for each age-since-disturbance patch, meaning that different extraction levels can affect resource availability over the successional gradient, a situation made more likely by large spatial-scale disturbances. In (b) all patches share a common pool, a situation more relevant to smaller (individual) scale disturbances. (c) Illustrates two mechanisms of water resource concentration in a semi-arid system, including greater infiltration rates near vegetation (differentially sized vertical arrows) and spatially extensive root systems that move water to the site of individual plants. Both allow greater local water availability than is possible using the grid cell mean soil moisture value. (d, e) illustrate a mechanism for size asymmetric in resource competition. In (d) two similar-sized root zones intersect, depleting the resource where they overlap. Since they are the same size, neither has an advantage. In (e) the smaller root system has most of its resource uptake soil volume depleted, whereas the larger root system is only mildly affected by the overlap, thus the larger root system gains an asymmetric advantage analogous to that of large trees in the forest canopy competing for light

moisture states to shift away from the mean due to vegetation-mediated positive feedback mechanisms (Figure 2c), including root foraging for water, and impacts of vegetation on infiltration and recruitment (Ivanov et al., 2010; Shachak et al., 2008). These mechanisms allow patchy vegetation to persist where the spatial mean moisture state might prevent viable vegetation growth. LSMs typically assume spatial homogeneity of moisture, leading in principle to underestimations of vegetation survival.

3.9.3 | Size symmetry of water competition

The degree to which competition for belowground resources is asymmetric with regard to plant size is unclear. VDMs typically allow partitioning of water between plants of differing root depth (Ivanov et al., 2012) but within a given soil volume assume perfectly symmetric competition. Schwinnig and Weiner (1998) argued that, where a large plant is in competition with a small plant, the fraction of the small plant's root zone affected by the resulting resource depletion is larger than the affected fraction of the root system of the large plant, (Figure 2d,e) suggesting the likelihood of size asymmetry, but the degree to which this is a dominant phenomenon remains unclear (DeMalach, Zaady, Weiner, & Kadmon, 2016; Schenk, 2006).

3.10 | Impacts of water on plant physiology

LSMs have used simplistic representations of the responses of plants to acquired soil moisture since their inception (McDowell et al., 2013). Recently, plant hydraulics theory (Sperry, Adler, Campbell, & Comstock, 1998), offline models (Bohrer et al., 2005; Gentile, Guérin, Uriarte, McDowell, & Pockman, 2015; Hickler, Prentice, Smith, Sykes, & Zaehle, 2006; Janott et al., 2011; McDowell et al., 2013; Williams, Bond, & Ryan, 2001), and datasets (Choat et al., 2012; Christoffersen et al., 2016; Maréchaux, Bartlett, Gaucher, Sack, & Chave, 2016) have improved substantially, giving rise to the potential for inclusion of "hydrodynamics" (prognosis of moisture states and fluxes within plants) in LSMs. These methods have shown promise for improving simulations of carbon, water, and energy fluxes, particularly during dry conditions (Bonan et al., 2014; Christoffersen et al., 2016; Fisher, Williams, Lobo do Vale, da Costa, & Meir, 2006; Fisher et al., 2007; Williams et al., 2001). Prediction of internal plant moisture status might also allow more realistic representation of drought deciduousness (Xu et al., 2016), sink limitations on growth (Fatichi, Leuzinger, & Körner, 2014), and stress-induced tree mortality (Anderegg et al., 2012).

Considerable effort is currently being expended on the implementation of such hydrodynamic schemes within VDMs (Christoffersen et al., 2016; Xu et al., 2016). Key challenges include (i) parameterization of hydraulic trait trade-offs and coordination across functional types and tissues, (ii) understanding the impact of segmentation of the hydraulic continuum, (iii) representing characteristic timescales of xylem embolism refilling (Mackay et al., 2015), (iv) linking stomatal responses to plant hydraulic states/fluxes (Bonan et al., 2014; Christoffersen et al., 2016; Sperry & Love, 2015; Sperry et al.,

2016), (v) integrating plant hydraulic status with existing growth and allocation schemes and demography, and (vi) integration with appropriate benchmarking data.

3.10.1 | Below-ground competition for nutrients

Nutrient cycling (nitrogen, rarely phosphorus) is now represented in several LSMs (Smith et al., 2014; Wang, Law, & Pak, 2010; Zaehle & Friend, 2010) yet uncertainties remain concerning the appropriate representation of many processes (Brzostek, Fisher, & Phillips, 2014; Xu et al., 2012; Zaehle & Dalmonech, 2011; Zaehle et al., 2014). VDMs inherit these uncertainties, and are subject to further structural degrees of freedom, derived from size/age structured representations of nutrient supply and demand. In principle, similar concerns of tiling impacts, aggregation, and asymmetric competition apply to nutrient as well as water uptake. One difference is that nutrients tend to be more abundant near the ground surface than at depth, thus may allow for more size-symmetric competition than for water. An advantage of VDMs is that they might better resolve some features of nutrient cycling that are difficult to include in typical LSMs, such as explicit representation of the successional status of nitrogen-fixing vs. nonfixing plants, and release of nutrient competition following disturbance. Of the models described here, ED2, LPJ-GUESS, and LM3-PPA have nutrient cycling schemes (Smith et al., 2014; Trugman et al., 2016; Weng et al., 2016).

3.11 | Vegetation demographics: recruitment & mortality

g1DVMs typically include representations of plant demography (recruitment, mortality; Sitch et al., 2003; Melton & Arora, 2016). Where, in g1DVMs, mortality and recruitment rates only impact mean vegetation biomass and PFT distributions, in size-structured VDMs, these processes also impact emergent forest structure and PFT composition, which in turn affect light competition and feedback on PFT filtering. There is thus a strong imperative to constrain demographic processes in VDMs.

3.12 | Mortality

Tree mortality is represented in g1DVMs as either a constant (in basic models) or as a function of various "proxies"—carbon balance, hydraulic stress, growth efficiency, plant traits, size, or age (McDowell et al., 2011, 2013). Persistent uncertainty about the major drivers of plant death means that a consensus on model structure is not yet justified. Some models now account for several additional sources of mortality, including biotic damage (Dietze & Matthes, 2014; Hicke et al., 2012; Jönsson, Schroeder, Lagergren, Anderbrandt, & Smith, 2012), atmospheric pollutants (Dietze & Moorcroft, 2011), wind damage (Lagergren, Jönsson, Blennow, & Smith, 2012), and herbivory (Pachzelt, Forrest, Rammig, Higgins, & Hickler, 2015), but relative importance of these various processes remains unclear. Representations of height structures and hydrodynamics in VDMs should

improve the fidelity of mortality proxy prediction, given that mortality can be strongly related to tree size (Bennett, McDowell, Allen, & Anderson-Teixeira, 2015; Lines, Coomes, & Purves, 2010; Muller-Landau et al., 2006).

The likelihood of increased climate-stress related mortality (Anderegg, Kane, & Anderegg, 2013) has motivated numerous experimental and observational studies in recent years (Anderegg et al., 2012, 2015; McDowell et al., 2008, 2011, 2013, 2015; Xu, McDowell, Sevanto, & Fisher, 2013; Zeppel, Adams, & Anderegg, 2011). One goal of this effort is to empirically relate hypothesized physiological proxies to rates of tree death at relevant scales.

At the scale of individual plants, mortality is a discrete process, occurring after some threshold of physiological stress is reached. If a model, however, predicted that all members of a given cohort died on the day that their average stress exceeded some threshold, that cohort would be extinguished across the whole landscape. Given ESM gridcells are often very large (>100 km resolution) this outcome would be ecologically unrealistic because of heterogeneity within the real population represented by that cohort. Cohort-based models therefore require empirical linkages between physiological proxies of death and mortality rates at the scale of model predictions. On a stand scale, the population represented by a cohort is heterogeneous due to variations in resource availability, genotype diversity, herbivory, and disease. Across a landscape, the population represented by a cohort might also encounter heterogeneity in soil texture, topography, aspect, microclimate, etc. Therefore we expect a looser connection between average physiological stress and landscape-scale mortality rates as the scale of prediction increases. In principle, the slope of the relationship between average physiological stress and landscape-scale mortality requires scale-dependent calibration (Figure 3).

3.13 | Recruitment

There is some evidence that establishment rates may be considerably more sensitive to environmental filters than selection of adult plants, thus, compositional shifts are as likely to be driven by changes in recruitment as by adult growth and mortality (Ibanez, Clark, & Dietze, 2008; Ibáñez et al., 2009). All demographic models represent plant recruitment processes (seed production, dispersal & germination), albeit simplistically. The rate of seed production is typically highly idealized; in ED-type models, it is a fixed fraction of net primary productivity (NPP) for plants that are in positive carbon balance (Moorcroft et al., 2001). Most VDMs do not consider dispersal among grid cells, given the complexities of this process (Nabel, 2015; Sato & Ise, 2012). To simulate germination, VDMs typically impose a minimum size threshold below which physiological processes and demography are not resolved and the emergence of new recruits is thus a phenomenological formulation (Farrion et al., 2013), modulated in some VDMs by climate envelopes (LPJ-GUESS, SEIB-DGVM), empirical proxies (forest-floor potential NPP, Smith et al., 2001), or plant traits and environmental conditions (Trugman et al., 2016).




FIGURE 3 Scale-dependence in extrapolating cohort-level mortality proxies to landscape-scale predictions of mortality. (a) illustrates a plausible multi-annual trajectory for mortality proxies throughout a chronic drought (solid line) and a hypothetical threshold whereby decline in this proxy is predictive of death (dashed line). (b) illustrates potential evolution of between mortality proxies and numbers of individuals through time (left to right). Long dashes represent a threshold-based mortality algorithm, whereby all individuals in a given cohort die in the same timestep. Short dashes illustrate a scaling from individual to landscape where there is a low level of heterogeneity across individuals, and the dotted line illustrates a condition with higher heterogeneity. In this case, parts of the grid cell (or genetic population) experience mortality at much lower degrees of average stress, and others are resilient under a given set of climatic drivers. As local mortality rates (gap scale) vary in comparison to landscape scale (entire forest), the slope of these lines requires calibration to a specific spatial scale

3.14 | Disturbance regimes: fire & land use

3.14.1 | Fire

Most ESMs contain representations of the impacts of fire, in g1DVMs, however, the impact of fire-induced tree mortality is simply to reduce the overall number density (individuals/m²) in the next timestep (Hantson et al., 2016). VDMs' tracking of size and age structure provides three opportunities to improve representation of fire-vegetation interactions. First, age-since-disturbance structured models can natively represent disturbance-recovery mosaics that

arise as a result of frequent fire regimes and fire-vegetation feedback processes (wind speed, flammability, recruitment) within those regimes. Second, vertical canopy structure can capture size-structured mortality resulting from fire events and thus represent the dynamics of the “fire-trap” in savanna-type ecosystems (Hoffmann & Solbrig, 2003; Hoffmann et al., 2012).

3.14.2 | Land use

Capturing the impact of human land-use and land-cover change on the carbon cycle, hydrology and other biogeophysical systems is a key application of ESMs (Brovkin et al., 2013; de Noblet-Ducoudré et al., 2012; Jones et al., 2011; Pongratz, Reick, Raddatz, & Clausen, 2009; Shevliakova et al., 2009). For the CMIP inter-comparison process, a single consolidated set of land-use transitions are specified (with carbon estimates from the “ED” model as described above, Hurtt et al., 2011), providing a matrix of transitions between land use classes (e.g., primary forest, secondary forest, pasture, cropland) through time (Lawrence et al., 2016). In traditional LSMs, land-use transitions must be translated into annual land-cover maps that specify the fraction of the land surface occupied by each PFT (Lawrence et al., 2012). A principal advantage of VDMs is that these land-use transitions can be directly implemented without the need for translation into PFT fractions, since they can explicitly simulate ecosystem disturbance and recovery (Shevliakova et al., 2009).

Representing human managed systems such as croplands, pasturelands, and plantation forests also requires the specification of transitions and management practices (e.g., harvest, grazing; Shevliakova et al., 2009; Lindeskog et al., 2013). Implementing standardized representations of these processes directly will emerge as a challenge as VDMs become more common elements of ESM structure. A further advantage of VDMs relates to the impacts of shifting cultivation. The impact of gross land use transitions has been estimated to generate emissions that are 15%–40% higher than the net transitions alone (Hansis, Davis, & Pongratz, 2015; Stocker, Feissli, Strassmann, Spahni, & Joos, 2014; Wilkenskjeld, Kloster, Pongratz, Raddatz, & Reick, 2014). This effect can be captured using age-since-disturbance mosaic approaches but is not directly possible with traditional LSMs.

4 | BENCHMARKING VDMs

4.1 | On the need for VDM-specific benchmarking data

Benchmarking and validation activities for LSMs have become increasingly numerous and sophisticated in recent years. These include comparisons against global or regional gridded data products (Luo et al., 2012), comparisons of relationships between two or more properties (emergent constraints), comparisons against intensively measured individual sites (including flux towers; Schwalm et al., 2010) and against manipulation experiments (de Kauwe et al., 2013;

Zaehle et al., 2014). The International Land Model Benchmarking Project (www.iLAMB.org) maintains a set of standard benchmarking products used for this purpose. Expanding this set of standardized data products to match the scope of VDMs will be a critical challenge in years to come (Hoffman et al., 2017). In this section, we describe potential metrics for benchmarking the novel aspects of VDMs described above. For some components, benchmarking datasets are already available, but for many they are scarce. We hope to illustrate potential platforms for future model-data integration made possible by the additional realism of VDM components.

Further to this, numerous model intercomparison projects in recent years have attempted to compare the outputs of large and complex Earth Systems Model components (including LSMs) against various types of benchmarking data. Vegetation demographics, in particular, are the emergent properties of a very large array of other simulated processes in VDMs, and so it is not clear that a straight intercomparison between the featured approaches would generate a clear comparison of how different methods for abstracting ecosystems into models compare. Given the lack of consensus on the parametric and structural approaches employed in the physiological and biophysical algorithms of all said models, it is notoriously difficult to assign differences in model performance to individual attributes, such as their demographic representation (c.f. Zaehle et al., 2014). This difficulty provides a motivation for assessing the skill of individual component parts (e.g., radiation transfer schemes, hydrodynamic representations, allocation, mortality, and recruitment models.). Illustrating that VDMs have these components in common might provide a framework for future more refined intercomparison studies (Table 1).

4.2 | Validation of radiation transfer and canopy organization

Radiation transfer models have two main components: The first is the underlying scheme; how radiation interacts with the scattering elements, reflectance and transmittance properties, and the treatment of diffuse radiation. Validating these representations is arguably an existing field of research, particularly in the realm of assessing canopy structure from remote sensing data (Smolander & Stenberg, 2005; Widlowski et al., 2007; Widlowski et al., 2015).

The second feature of VDM RTMs is the assumptions controlling the arrangement of scattering elements with respect to each other. For example: Is a discrete-layered PPA-like structure a good approximation of a forest canopy? How much do adjacent crowns affect each other's light interception? These questions might be addressed using detailed 3D observations on the arrangement of leaves and crowns in space, via high-density airborne or ground-based LiDAR (Detto, Asner, Muller-Landau, & Sonnentag, 2015; Stark et al., 2012, 2015), and then applying more complex 3D radiative transfer models (Morton et al., 2015) to assess how alternate VDM RTM structures perform. These exercises should be prioritized, since the impact of size on resource acquisition is such a fundamentally important process in determining ecological demographics.

4.3 | Validation of plant water use

Establishing credible boundary conditions (soil moisture, meteorology, vegetation structure) and appropriate validation data (sap flow, leaf water potential, gas exchange) is a challenging prerequisite for testing alternative hypotheses about the physics and physiology governing plant water utilization. The number of locations for which this validation is possible is small but growing. Such intensive ecosystem physiology observations have proven extremely valuable, however, illustrated by their repeated use in model validation exercises (Fisher et al., 2007; Joetzjer et al., 2014; Matheny et al., 2016; Plaut et al., 2012; Poyatos, Aguadé, Galiano, Mencuccini, & Martínez-Vilalta, 2013; Williams et al., 2001; Zeppel et al., 2008). Since plant water status is fundamentally linked to both height, (on account of gravitational effects) and canopy position (in relation to differing evaporation rates) vertically resolved models are critical to allowing direct model-data comparison exercises. Christofferson et al., for example, Christoffersen et al. (2016) illustrate the importance of canopy position in correctly simulating daytime leaf water potential at the Caxiuaana throughfall exclusion experiment in Amazonia, thus, g1DVMs with aggregated plant water status might be difficult to compare directly with plant hydraulics observations.

New datasets documenting stem water storage (Carrasco et al., 2015; Matheny et al., 2015), remotely sensed plant water status (Konings & Gentine, 2016) and solar induced fluorescence (Guanter, Köhler, Walther, & Zhang, 2016), also have the potential to provide additional metrics for evaluation of hydrodynamic model predictions.

4.4 | Validation of canopy structure

4.4.1 | Tests against plot-scale size structure data

The canopy structure (tree size frequency per plant type) predicted by a VDM can be validated using ecological census data from permanent sample plots. Predicted canopy structure is a high-level emergent property, however, and is influenced by radiation transfer, photosynthesis, respiration, allocation, and demographics (recruitment and mortality). Thus where discrepancies arise, it is difficult to diagnose the model specific errors that led to the poor predictive power.

Using the PPA, Fariior et al. (2016) circumvent this problem by collapsing the details of growth and demographic rates at a given location into constant rate parameters, which vary only with canopy status and PFT. This approach successfully captures the size distribution of a tropical rainforest, in particular, the observed shift in structure between understory trees (which approximate a power-law distribution) and canopy trees (which do not). This result highlights (i) the need to account for asymmetric availability of light across size classes (in contrast to West, Brown, & Enquist, 1999; Enquist, West, & Brown, 2009), (ii) that simulation of small-scale disturbances is critical in the tropics, necessitating a model inclusive of gap formation and (iii), that representation of the plant canopy as distinct strata (canopy and one or more understory layers) is a useful simplification.

4.4.2 | Tests against remote sensing of canopy structure

Existing Earth Observation products can detect phenological signals (Hansen et al., 2002) and vegetation stature (Lefsky et al., 2005; Simard, Pinto, Fisher, & Baccini, 2011). This allows a remote detection of “traditional” PFTs (defined by phenology and growth habit). As such, DVMs have historically been tested against these vegetation classification maps (Arora & Boer, 2006; Bonan et al., 2003; Fisher et al., 2015; Sitch et al., 2003). Emerging data products that observe high spectral resolution (i.e., “hyperspectral”) data streams can be used to discern the properties of plant surfaces, which themselves can be linked to leaf or canopy traits (Asner et al., 2012; Shiklomanov, Dietze, Viskari, Townsend, & Serbin, 2016; Singh, Serbin, McNeil, Kingdon, & Townsend, 2015). In principle, VDMs specified using PFTs that align with traits that can be detected using hyperspectral sensing (e.g., leaf nutrient and photosynthetic properties, moisture/temperature features, leaf thickness & venation) could be validated using these types of observation (Antonarakis et al., 2014; Asner et al., 2016; Serbin et al., 2015). Use of LIDAR to detect individual tree height and crown diameter (Barbier, Coueron, Proisy, Malhi, & Gastellu-Etchegorry, 2010; Garrity, Meyer, Maurer, Hardiman, & Bohrer, 2011; Hurtt et al., 2004, 2010, 2016; Jucker et al., 2017; Thomas et al., 2008) can be used to constrain model vegetation structure predictions.

4.5 | Validation of demographic rates (recruitment, growth, mortality)

4.5.1 | Tests against plot-scale demographic data

Recruitment, growth, and mortality rates can be estimated from repeated censuses at permanent sample plots (Lewis et al., 2004; Phillips et al., 2010). The direct use of plot data is hindered by the need to drive models with local climate data, however, and as such is typically limited to more intensively observed field sites (Powell et al., 2013). Emergent relationships, such as the change in mortality with environmental gradients (Phillips et al., 2010) other ecosystem properties such as NPP (Delbart et al., 2010) and regional extrapolation of mortality rates (Johnson et al., 2016; Lines et al., 2010) should also prove useful as benchmarks that a model might be expected to capture.

4.5.2 | Tests against remotely sensed demographic data

Recent developments in remote sensing-based disturbance detection, including a high-resolution (30 m × 30 m) global disturbance database (Hansen et al., 2013), provide opportunities to evaluate large tree mortality events at global scales. At smaller scales, (Garrity et al., 2013) illustrate the potential for detection of tree mortality using 1 m resolution QuickBird imagery. Hyperspectral and airborne LIDAR techniques will likely improve our ability to remotely detect tree mortality

rates (Eitel et al., 2016). Remaining limitations of these approaches include the fact that they primarily detect mortality of canopy trees (McDowell et al., 2015), and that issues related to return frequency, cloud cover, sensor lifetime impact the ability to detect the exact timing of mortality events, impeding attribution of their drivers.

5 | BROADER ISSUES CONCERNING THE INCLUSION OF VEGETATION DEMOGRAPHICS IN ESMS

5.1 | Plant trait information

The use of plant trait data for parameterization of LSMs in general (Reich, 2014; Reich, Rich, Lu, Wang, & Oleksyn, 2014; Verheijen et al., 2015) and VDMs in particular has been covered extensively elsewhere (Fisher et al., 2015; Fyllas et al., 2014; Pappas, Fatichi, & Burlando, 2016; Pavlick et al., 2013; Sakschewski et al., 2015; Scheiter et al., 2012). VDMs are primarily designed as “trait filtering” models, in that they predict the differential demographic performance (in terms of growth mortality and recruitment) from plant traits and environmental conditions, and thus in turn predict/filter the distribution of those traits across the landscape. The success of trait filtering approaches is linked to the fidelity with which trait trade-off surfaces are prescribed to the model (Scheiter et al., 2012). Designing balanced trade-offs is a particular concern with this approach. Specifically, allowing traits to vary such that one particular PFT gains a large growth advantage -without sufficient attendant cost- will result in the runaway dominance of that PFT, particularly given positive feedbacks between growth, resource acquisition, and reproductive success (Bohn et al., 2011; Fisher et al., 2010, 2015; Pacala & Tilman, 1994). Deriving balanced trade-off surfaces from plant trait datasets is also problematic if environmental variation affects plant trait expression. For example, Reich et al. (2014) find that leaf lifespan of needleleaf trees varies with temperature, breaking the more typical global correlation with leaf mass per unit area. Thus, geographical datasets can conflate the causes of trait variability, as can impacts of shade on trait plasticity (Keenan & Niinemets, 2016). We strongly advocate for detailed analyses of emerging trait databases to provide relevant trait relationships for VDM simulations (e.g., Christoffersen et al., 2016).

5.2 | Improving informatics

Working with VDMs presents notable challenges in informatics and statistics due to their complexity, input requirements and output dimensionality. Recently, model informatics systems have emerged to meet these challenges, including the Predictive Ecosystem Analyzer (PEcAn) (Dietze, 2014; Dietze, Lebauer, & Kooper, 2013; LeBauer et al., 2013). The primary goals of PEcAn are to reduce redundancy and improve reliability in the workflows associated with running, parameterizing, validating, calibrating, and reporting ecosystem models. PEcAn consists of a suite of open-source workflow and statistical tools (<https://github.com/PecanProject/pecan>) and a web interface,

(pecanproject.org). Tools include sensitivity analysis and variance decomposition, model-data assimilation, Bayesian calibration, as well as generation of common meteorological drivers, validation data streams, benchmarking, and visualizations. These common protocols allow analyses to be replicated across models, making inter-model comparison easier. Furthermore, PEcAn employs a database to track workflows across researchers and institutions, allowing robust provenance tracking. PEcAn currently supports ten different ecosystem models, including most of the models discussed in this paper (ED2, CLM(ED), LM3-PPA, LPJ-GUESS). PEcAn is an open community project, and is extensible for novel analyses and modules.

6 | SUMMARY

We describe the major modifications to traditional dynamic vegetation models that are necessary to allow structured representation of ecological demographic processes inside the architecture of Earth System Models. These developments open a number of avenues for better data-model integration, and highlight gaps in ecological observation and understanding that we hope could be a major focus of future scientific endeavor.

Priority areas for VDM development include:

1. Partitioning of solar radiation between competing plant canopies, and the physics of shading within and between individuals and cohorts.
2. Representation of plant hydrodynamics in models, and improving parameterizations and linkages to observations.
3. Distribution of below-ground resources (nutrients and water) between size classes, PFTs, and patches.
4. Representation of demographic processes (mortality and recruitment), scaling of reductionist physiological models of plant mortality across heterogeneous landscapes, calibration using emerging remote sensing products, and trait impacts on recruitment rates.
5. Better representation of land use processes and fire disturbance. For fire, this includes size-structured impacts of burning, as well as recovery and fire-vegetation feedback processes. For land use change, challenges include mapping transitions into clear impacts on ecosystem structure and management and global parameterization of heterogeneous anthropogenic impacts.

These foci integrate many potential avenues for novel model-data-fusion effort that are made plausible via the implementation of demographic models. We advocate for more intensive and innovative usage of ecological data streams in model validation and argue that the many different avenues for development of VDMs will benefit from coordinated approaches to these topics. The development of common, widely-available intensive (plot-scale) and distributed (network and remote-sensing scale) testbeds, accessible via commonly-used online tools (e.g., iLAMB, PEcAn) will provide the greatest chance of constraining future trajectories of the land biosphere in ESMS.

ACKNOWLEDGEMENTS

The outline for this manuscript was identified during a workshop focused on VDM implementation in ESMs held at the National Center for Atmospheric Research in January 2016. NCAR is sponsored by the National Science Foundation. CDK, BC, RK, JH, TP, JS, CX, & SPS were supported by the Next-Generation Ecosystem Experiments (NGEE Tropics) project that is supported by the Office of Biological and Environmental Research in the Department of Energy, Office of Science. TV and SPS were supported by NASA Terrestrial Ecology grant NNX14AH65G and through the United States Department of Energy contract No. DE-SC0012704 to Brookhaven National Laboratory. ATT was partially supported by a National Science Foundation Graduate scholarship. BS acknowledges support from the Strategic Research Fellowship Area MERGE. ML was funded by FAPESP (grant 2015/07227-6) AMM was supported by U.S. National Science Foundation Hydrological Science grant 1521238. GH acknowledges the support of NASA. JL and TZ were funded by USDA agreements 11-JV-112423-059 and 16-JV-11242306-050. DM acknowledges support from the US Department of Energy, Office of Science, Office of Biological and Environmental Research, Terrestrial Ecosystem Science (TES) Program under award number DE-SC0014363. HV was supported by the ERC starting grant 637643 (TREECLIMBERS).

ORCID

Rosie A. Fisher  <http://orcid.org/0000-0003-3260-9227>
 William R. L. Anderegg  <http://orcid.org/0000-0001-6551-3331>
 Michael C. Dietze  <http://orcid.org/0000-0002-2324-2518>
 Ryan G. Knox  <http://orcid.org/0000-0003-1140-3350>
 Marcos Longo  <http://orcid.org/0000-0001-5062-6245>
 Ashley M. Matheny  <http://orcid.org/0000-0002-9532-7131>
 David Medvigy  <http://orcid.org/0000-0002-3076-3071>
 Thomas L. Powell  <http://orcid.org/0000-0002-3516-7164>
 Shawn P. Serbin  <http://orcid.org/0000-0003-4136-8971>
 Anna T. Trugman  <http://orcid.org/0000-0002-7903-9711>
 Ensheng Weng  <http://orcid.org/0000-0002-1858-4847>
 Tao Zhang  <http://orcid.org/0000-0001-7135-1762>

REFERENCES

- Albani, M., Medvigy, D., Hurtt, G. C., & Moorcroft, P. R. (2006). The contributions of land-use change, CO₂ fertilization, and climate variability to the Eastern US carbon sink. *Global Change Biology*, 12, 2370–2390.
- Anderegg, W. R., Berry, J. A., Smith, D. D., Sperry, J. S., Anderegg, L. D., & Field, C. B. (2012). The roles of hydraulic and carbon stress in a widespread climate-induced forest die-off. *Proceedings of the National Academy of Sciences of the USA*, 109(1), 233–237.
- Anderegg, W. R. L., Flint, A., Huang, C.-Y., Flint, L., Berry, J. A., Davis Frank, W., ... Field, C. B. (2015). Tree mortality predicted from drought-induced vascular damage. *Nature Geoscience*, 8, 367–371.
- Anderegg, W. R., Kane, J. M., & Anderegg, L. D. (2013). Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature Climate Change*, 3(1), 30–36.
- Antonarakis, A. S., Munger, J. W., & Moorcroft, P. R. (2014). Imaging spectroscopy- and lidar-derived estimates of canopy composition and structure to improve predictions of forest carbon fluxes and ecosystem dynamics. *Geophysical Research Letters*, 41(7), 2535–2542.
- Arora, V. K., & Boer, G. J. (2006). Simulating competition and coexistence between plant functional types in a dynamic vegetation model. *Earth Interactions*, 10, 1–30.
- Arora, V. K., & Boer, G. J. (2010). Uncertainties in the 20th century carbon budget associated with land use change. *Global Change Biology*, 16, 3327–3348.
- Asner, G. P., Knapp, D. E., Boardman, J., Green, R. O., Kennedy-Bowdoin, T., Eastwood, M., ... Field, C. B. (2012). Carnegie Airborne Observatory-2: Increasing science data dimensionality via high-fidelity multi-sensor fusion. *Remote Sensing of the Environment*, 124, 454–465.
- Asner, G. P., Martin, R. E., Anderson, C. B., Kryston, K., Vaughn, N., Knapp, D. E., ... Tupayachi, R. (2016). Scale dependence of canopy trait distributions along a tropical forest elevation gradient. *New Phytologist*, <https://doi.org/10.1111/nph.14068>
- Barbier, N., Coutron, P., Proisy, C., Malhi, Y., & Gastellu-Etchegorry, J. P. (2010). The variation of apparent crown size and canopy heterogeneity across lowland Amazonian forests. *Global Ecology and Biogeography*, 19, 72–84.
- Bennett, A. C., McDowell, N. G., Allen, C. D., & Anderson-Teixeira, K. J. (2015). Larger trees suffer most during drought in forests worldwide. *Nature Plants*, 1, 15139.
- Bohn, K., Dyke, J. G., Pavlick, R., Reineking, B., Reu, B., & Kleidon, A. (2011). The relative importance of seed competition, resource competition and perturbations on community structure. *Biogeosciences*, 8, 1107–1120.
- Bohrer, G., Mourad, H., Laursen, T. A., Drewry, D., Avissar, R., Poggi, D., ... Katul, G. G. (2005). Finite element tree crown hydrodynamics model (FETCH) using porous media flow within branching elements: A new representation of tree hydrodynamics. *Water Resources Research*, 41, 11.
- Bonan, G. B. (2008). Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. *Science*, 320, 1444–1449.
- Bonan, G. B., Lawrence, P. J., Oleson, K. W., Levis, S., Jung, M., Reichstein, M., ... Swenson, S. C. (2011). Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data. *Journal of Geophysical Research: Biogeosciences*, 116(G2), G02014. <https://doi.org/10.1029/2010JG001593>
- Bonan, G. B., Levis, S., Sitch, S., Vertenstein, M., & Oleson, K. W. (2003). A dynamic global vegetation model for use with climate models: Concepts and description of simulated vegetation dynamics. *Global Change Biology*, 9, 1543–1566.
- Bonan, G. B., Williams, M., Fisher, R. A., & Oleson, K. W. (2014). Modeling stomatal conductance in the earth system: Linking leaf water-use efficiency and water transport along the soil–plant–atmosphere continuum. *Geoscientific Model Development*, 7, 2193–2222.
- Borgogno, F., D'Odorico, P., Laio, F., & Ridolfi, L. (2009). Mathematical models of vegetation pattern formation in ecohydrology. *Reviews of Geophysics*, 47(1), RG1005. <https://doi.org/10.1029/2007RG000256>
- Brovkin, V., Boysen, L., Arora, V. K., Boisier, J. P., Cadule, P., Chini, L., ... Hurtt, G. C. (2013). Effect of anthropogenic land-use and land-cover changes on climate and land carbon storage in CMIP5 projections for the twenty-first century. *Journal of Climate*, 26, 6859–6881.
- Brzostek, E. R., Fisher, J. B., & Phillips, R. P. (2014). Modeling the carbon cost of plant nitrogen acquisition: Mycorrhizal trade-offs and multi-path resistance uptake improve predictions of retranslocation. *Journal of Geophysical Research: Biogeosciences*, 119, 1684–1697.
- Bugmann, H. (2001). A review of forest gap models. *Climatic Change*, 51, 259–305.
- Cao, M., & Woodward, F. I. (1998). Dynamic responses of terrestrial ecosystem carbon cycling to global climate change. *Nature*, 393, 249–252.

- Carrasco, L. O., Bucci, S. J., Di Francescantonio, D., Lezcano, O. A., Campanello, P. I., Scholz, F. G., ... Holbrook, N. M. (2015). Water storage dynamics in the main stem of subtropical tree species differing in wood density, growth rate and life history traits. *Tree Physiology*, *35*, 354–365.
- Choat, B., Jansen, S., Brodribb, T. J., Cochard, H., Delzon, S., Bhaskar, R., ... Jacobsen, A. L. (2012). Global convergence in the vulnerability of forests to drought. *Nature*, *491*, 752–755.
- Christoffersen, B. O., Gloor, M., Fauset, S., Fyllas, N. M., Galbraith, D. R., Baker, T. R., ... Meir, P. (2016). Linking hydraulic traits to tropical forest function in a size-structured and trait-driven model (TFS v.1-Hydro). *Geoscientific Model Development*, *9*, 4227–4255.
- Cox, P. M. (2001). Description of the TRIFFID dynamic global vegetation model. *Hadley Centre Technical Note*, *24*, 1–16.
- Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., & Totterdell, I. J. (2000). Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature*, *408*, 184–187.
- Davidson, E., Lefebvre, P. A., Brando, P. M., Ray, D. M., Trumbore, S. E., Solorzano, L. A., ... Nepstad, D. C. (2009). Carbon inputs and water uptake in deep soils of an eastern Amazon forest. *Forest Science*, *57*, 51–58.
- de Kauwe, M. G., Medlyn, B. E., Zaehle, S., Walker, A. P., Dietze, M. C., Hickler, T., ... Smith, B. (2013). Forest water use and water use efficiency at elevated CO₂: A model-data intercomparison at two contrasting temperate forest FACE sites. *Global Change Biology*, *19*, 1759–1779.
- de Noblet-Ducoudré, N., Boisier, J.-P., Pitman, A., Bonan, G., Brovkin, V., Cruz, F., ... Lawrence, P. (2012). Determining robust impacts of land-use-induced land cover changes on surface climate over North America and Eurasia: Results from the first set of LUCID experiments. *Journal of Climate*, *25*, 3261–3281.
- Delbart, N., Ciais, P., Chave, J., Viovy, N., Malhi, Y., & Le Toan, T. (2010). Mortality as a key driver of the spatial distribution of aboveground biomass in Amazonian forest: Results from a dynamic vegetation model. *Biogeosciences*, *7*, 3027–3039.
- DeMalach, N., Zaady, E., Weiner, J., & Kadmon, R. (2016). Size asymmetry of resource competition and the structure of plant communities. *Journal of Ecology*, *104*, 899–910.
- Detto, M., Asner, G. P., Muller-Landau, H. C., & Sonnentag, O. (2015). Spatial variability in tropical forest leaf area density from multireturn lidar and modeling. *Journal of Geophysical Research: Biogeosciences*, *120*, 294–309.
- Dickinson, R. E. (1983). Land surface processes and climate-surface albedos and energy balance. *Advances in Geophysics*, *25*, 305–353.
- Dietze, M. C. (2014). Gaps in knowledge and data driving uncertainty in models of photosynthesis. *Photosynthesis Research*, *119*, 3–14.
- Dietze, M. C., & Latimer, A. M. (2011). Forest simulators. In A. Hastings & L. Gross (Eds.), *Invited chapter in: Sourcebook in theoretical ecology* (pp. 307–316). Berkeley, CA: University of California Press.
- Dietze, M. C., Lebauer, D. S., & Kooper, R. O. (2013). On improving the communication between models and data. *Plant, Cell & Environment*, *36*, 1575–1585.
- Dietze, M. C., & Matthes, J. H. (2014). A general ecophysiological framework for modelling the impact of pests and pathogens on forest ecosystems. *Ecology Letters*, *17*(11), 1418–1426.
- Dietze, M. C., & Moorcroft, P. R. (2011). Tree mortality in the eastern and central United States: Patterns and drivers. *Global Change Biology*, *17*, 3312–3326.
- D'Odorico, P., Caylor, K., Okin, G. S., & Scanlon, T. M. (2007). On soil moisture-vegetation feedbacks and their possible effects on the dynamics of dryland ecosystems. *Journal of Geophysical Research: Biogeosciences*, *112*(G4), G04010. <https://doi.org/10.1029/2006JG000379>
- Dubayah, R., Goetz, S. J., Blair, J. B., Fatoyinbo, T. E., Hansen, M., Healey, S. P., ... Swatantran, A. GEDI: The global ecosystem dynamics investigation. *American Geophysical Union, Fall Meeting*, abstract #U14A-07.
- Dybzinski, R., Farnier, C., Wolf, A., Reich, P. B., & Pacala, S. W. (2011). Evolutionarily stable strategy carbon allocation to foliage, wood, and fine roots in trees competing for light and nitrogen: An analytically tractable, individual-based model and quantitative comparisons to data. *American Naturalist*, *177*, 153–166.
- Eitel, J. U., Höfle, B., Vierling, L. A., Abellán, A., Asner, G. P., Deems, J. S., ... Mandlburger, G. (2016). Beyond 3-D: The new spectrum of lidar applications for earth and ecological sciences. *Remote Sensing of Environment*, *186*, 372–392.
- Enquist, B. J., West, G. B., & Brown, J. H. (2009). A general quantitative theory of forest structure and dynamics. *Proceedings of the National Academy of Sciences of the USA*, *106*, 7046–7051.
- Evans, M. R. (2012). Modelling ecological systems in a changing world. *Philosophical Transactions of the Royal Society London. B-Biology*, *36*, 181–190.
- Farnier, C. E., Bohlman, S. A., Hubbell, S., & Pacala, S. W. (2016). Dominance of the suppressed: Power-law size structure in tropical forests. *Science*, *8*, 155–157.
- Farnier, C. E., Dybzinski, R., Levin, S. A., & Pacala, S. W. (2013). Competition for water and light in closed-canopy forests: A tractable model of carbon allocation with implications for carbon sinks. *The American Naturalist*, *181*(3), 314–330.
- Farnier, C. E., Rodriguez-Iturbe, I., Dybzinski, R., Levin, S. A., & Pacala, S. W. (2015). Decreased water limitation under elevated CO₂ amplifies potential for forest carbon sinks. *Proceedings of the National Academy of Sciences of the USA*, *112*, 7213–7218.
- Faticchi, S., Leuzinger, S., & Körner, C. (2014). Moving beyond photosynthesis: From carbon source to sink-driven vegetation modeling. *New Phytologist*, *201*, 1086–1095.
- Feeley, K. J., Davies, S. J., Ashton, P. S., Bunyavejchewin, S., Supardi, M. N. N., Kassim, A. R., ... Chave, J. (2007). The role of gap phase processes in the biomass dynamics of tropical forests. *Proceedings of the Royal Society London B: Biological Sciences*, *274*, 2857–2864.
- Fischer, R., Bohn, F., de Paula, M. D., Dislich, C., Groeneveld, J., Gutiérrez, A. G., ... Pütz, S. (2016). Lessons learned from applying a forest gap model to understand ecosystem and carbon dynamics of complex tropical forests. *Ecological Modelling*, *326*, 124–133.
- Fisher, R. A., McDowell, N., Purves, D., Moorcroft, P., Sitch, S., Cox, P., ... Ian Woodward, F. (2010). Assessing uncertainties in a second-generation dynamic vegetation model caused by ecological scale limitations. *New Phytologist*, *187*, 666–681.
- Fisher, R. A., Muszala, S., Versteinstein, M., Lawrence, P., Xu, C., McDowell, N. G., ... Lawrence, D. (2015). Taking off the training wheels: The properties of a dynamic vegetation model without climate envelopes. *Geoscientific Model Development*, *8*, 3293–3357.
- Fisher, R. A., Williams, M., da Costa, A. L., Malhi, Y., da Costa, R. F., Almeida, S., & Meir, P. (2007). The response of an Eastern Amazonian rain forest to drought stress: Results and modelling analyses from a throughfall exclusion experiment. *Global Change Biology*, *13*, 2361–2378.
- Fisher, R. A., Williams, M., Lobo do Vale, R., da Costa, A. L., & Meir, P. (2006). Evidence from Amazonian forests is consistent with isohydric control of leaf water potential. *Plant, Cell & Environment*, *29*, 151–165.
- Fisk, J. P. (2015). Net effects of disturbance: Spatial, temporal, and societal dimensions of forest disturbance and recovery on terrestrial carbon balance. PhD Thesis, University of Maryland.
- Fisk, J. P., Hurtt, G. C., Chambers, J. Q., Zeng, H., Dolan, K. A., & Negrón-Juárez, R. I. (2013). The impacts of tropical cyclones on the net carbon balance of eastern US forests (1851–2000). *Environmental Research Letters*, *8*(4), 045017.
- Flanagan, S. A., Hurtt, G. C., Fisk, J. P., Sahajpal, R., Hansen, M. C., Dolan, K. A., ... Zhao, M. (2016). Potential vegetation and carbon redistribution in Northern North America from climate change. *Climate*, *4*, 2–13.

- Foley, J. A., Prentice, I. C., Ramankutty, N., Levis, S., Pollard, D., Sitch, S., & Haxeltine, A. (1996). An integrated biosphere model of land surface processes, terrestrial carbon balance, and vegetation dynamics. *Global Biogeochemical Cycles*, *10*, 603–628.
- Fyllas, N. M., Gloor, E., Mercado, L. M., Sitch, S., Quesada, C. A., Domingues, T. F., ... Higuchi, N. (2014). Analysing Amazonian forest productivity using a new individual and trait-based model (TFS v. 1). *Geoscientific Model Development*, *7*, 1251–1269.
- Garrity, S. R., Allen, C. D., Brumby, S. P., Gangodagamage, C., McDowell, N. G., & Cai, D. M. (2013). Quantifying tree mortality in a mixed species woodland using multitemporal high spatial resolution satellite imagery. *Remote Sensing of Environment*, *129*, 54–65.
- Garrity, S. R., Meyer, K., Maurer, K. D., Hardiman, B., & Bohrer, G. (2011). Estimating plot-level tree structure in a deciduous forest by combining allometric equations, spatial wavelet analysis and airborne LiDAR. *Remote Sensing Letters*, *3*, 443–451.
- Gentine, P., Guérin, M., Uriarte, M., McDowell, N. G., & Pockman, W. T. (2015). An allometry-based model of the survival strategies of hydraulic failure and carbon starvation. *Ecohydrology*, *9*, 529–546.
- Gilad, E., Shachak, M., & Meron, E. (2007). Dynamics and spatial organization of plant communities in water-limited systems. *Theoretical population biology*, *72*, 214–230.
- Goudriaan, J. (1977). *Crop micrometeorology: A simulation study*. Wageningen: PUDOC.
- Guanter, L., Köhler, P., Walther, S., & Zhang, Y. (2016). Recent advances in global monitoring of terrestrial sun-induced chlorophyll fluorescence. In *Geoscience and Remote Sensing Symposium (IGARSS), 2016 IEEE International* (pp. 1714–1716). IEEE.
- Hansen, M. C., DeFries, R. S., Townshend, J. R. G., Sohlberg, R., Dimiceli, C., & Carroll, M. (2002). Towards an operational MODIS continuous field of percent tree cover algorithm: Examples using AVHRR and MODIS data. *Remote Sensing of Environment*, *83*, 303–319.
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., ... Kommareddy, A. (2013). High-resolution global maps of 21st-century forest cover change. *Science*, *342*, 850–853.
- Hansis, E., Davis, S. J., & Pongratz, J. (2015). Relevance of methodological choices for accounting of land use change carbon fluxes. *Global Biogeochemical Cycles*, *29*, 1230–1246.
- Hantson, S., Arneith, A., Harrison, S. P., Kelley, D. I., Prentice, I. C., Rabin, S. S., ... Bachelet, D. (2016). The status and challenge of global fire modelling. *Biogeosciences*, *13*, 3359–3375.
- Haverd, V., Smith, B., Cook, G. D., Briggs, P. R., Nieradzki, L., Roxburgh, S. R., ... Canadell, J. G. (2013). A stand-alone tree demography and landscape structure module for Earth system models. *Geophysical Research Letters*, *40*, 5234–5239.
- Hazeleger, W., Severijns, C., Semmler, T., Ștefănescu, S., Yang, S., Wang, X., ... Bougeault, P. (2010). EC-Earth: A seamless earth system prediction approach in action. *Bulletin of the American Meteorological Society*, *91*, 1357–1363.
- Hicke, J. A., Allen, C. D., Desai, A., Dietze, M. C., Hall, R. J., Hogg, E. T., ... Vogelmann, J. (2012). The effects of biotic disturbances on carbon budgets of North American forests. *Global Change Biology*, *18*, 7–34.
- Hickler, T., Prentice, I. C., Smith, B., Sykes, M. T., & Zaehle, S. (2006). Implementing plant hydraulic architecture within the LPJ Dynamic Global Vegetation Model. *Global Ecology and Biogeography*, *15*, 567–577.
- Hoffman, F. M., Riley, W. J., Randerson, J. T., Keppel-Aleks, G., Lawrence, D. M., & Koven, C. D. (2017). *International Land Model Benchmarking (ILAMB) 2016 workshop report*. DOE/SC-0186, U.S. Department of Energy, Office of Science, Germantown, Maryland, USA.
- Hoffmann, W. A., Geiger, E. L., Gotsch, S. G., Rossatto, D. R., Silva, L. C., Lau, O. L., ... Franco, A. C. (2012). Ecological thresholds at the savanna-forest boundary: How plant traits, resources and fire govern the distribution of tropical biomes. *Ecology Letters*, *15*, 759–768.
- Hoffmann, W. A., & Solbrig, O. T. (2003). The role of topkill in the differential response of savanna woody species to fire. *Forest Ecology and Management*, *180*, 273–286.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., ... Lipscomb, W. H. (2013). The community earth system model: A framework for collaborative research. *Bulletin of the American Meteorological Society*, *94*, 1339–1360.
- Hurtt, G. C., Chini, L., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., ... Jones, C. D. (2011). Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Climatic Change*, *109*, 117. <https://doi.org/10.1007/s10584-011-0153-2>
- Hurtt, G. C., Dubayah, R., Drake, J., Moorcroft, P., Pacala, S. W., Blair, J. B., & Fearon, M. G. (2004). Beyond potential vegetation: Combining lidar data and a height-structured model for carbon studies. *Ecological Applications*, *14*, 873–883.
- Hurtt, G. C., Fisk, J., Thomas, R., Dubayah, R., Moorcroft, P., & Shugart, H. H. (2010). Linking models and data on vegetation structure. *Journal of Geophysical Research*, *115*, 1–11.
- Hurtt, G. C., Frolking, S., Fearon, M. G., Moore, B. III, Shevliakova, E., Malyshev, S., ... Houghton, R. A. (2006). The underpinnings of land-use history: Three centuries of global gridded land-use transitions, wood harvest activity, and resulting secondary lands. *Global Change Biology*, *12*, 1208–1229.
- Hurtt, G. C., Moorcroft, P. R., Pacala, S. W., & Levin, S. A. (1998). Terrestrial models and global change: Challenges for the future. *Global Change Biology*, *4*, 581–590.
- Hurtt, G. C., Thomas, R. Q., Fisk, J. P., Dubayah, R. O., & Sheldon, S. L. (2016). The Impact of Fine-Scale disturbances on the predictability of vegetation dynamics and carbon flux. *PLoS One*, *11*, e0152883.
- Ibanez, I., Clark, J. S., & Dietze, M. C. (2008). Evaluating the sources of potential migrant species: Implications under climate change. *Ecological Applications*, *18*, 1664–1678.
- Ibáñez, I., Silander, J. A., Wilson, A. M., LaFleur, N., Tanaka, N., & Tsuyama, I. (2009). Multivariate forecasts of potential distributions of invasive plant species. *Ecological Applications*, *19*, 359–375.
- Ivanov, V. Y., Fatichi, S., Jenerette, G. D., Espeleta, J. F., Troch, P. A., & Huxman, T. E. (2010). Hysteresis of soil moisture spatial heterogeneity and the “homogenizing” effect of vegetation. *Water Resources Research*, *1*(46), 9.
- Ivanov, V. Y., Hutrya, L. R., Wofsy, S. C., Munger, J. W., Saleska, S. R., de Oliveira, R. C., & de Camargo, P. B. (2012). Root niche separation can explain avoidance of seasonal drought stress and vulnerability of overstory trees to extended drought in a mature Amazonian forest. *Water Resources Research*, *48*, 12.
- Janott, M., Gayler, S., Gessler, A., Javaux, M., Klier, C., & Priesack, E. (2011). A one-dimensional model of water flow in soil-plant systems based on plant architecture. *Plant and Soil*, *341*, 233–256.
- Joetzer, E., Delire, C., Douville, H., Ciais, P., Decharme, B., Fisher, R., ... Meir, P. (2014). Predicting the response of the Amazon rainforest to persistent drought conditions under current and future climates: A major challenge for global land surface models. *Geoscientific Model Development*, *10*, 2933–2950.
- Johnson, M. O., Galbraith, D., Gloor, M., De Deurwaerder, H., Guimberteau, M., Rammig, A., ... Phillips, O. L. (2016). Variation in stem mortality rates determines patterns of above-ground biomass in Amazonian forests: Implications for dynamic global vegetation models. *Global Change Biology*, *22*(12), 3996–4013.
- Jones, C. D., Hughes, J. K., Bellouin, N., Hardiman, S. C., Jones, G. S., Knight, J., ... Boo, K. O. (2011). The HadGEM2-ES implementation of CMIP5 centennial simulations. *Geoscientific Model Development*, *4*, 543–570.
- Jönsson, A. M., Schroeder, L. M., Lagergren, F., Anderbrandt, O., & Smith, B. (2012). Guess the impact of Ips typographus—An ecosystem

- modelling approach for simulating bark beetle outbreaks. *Agricultural & Forest Meteorology*, 166–167, 188–200.
- Jucker, T., Caspersen, J., Chave, J., Antin, C., Barbier, N., Bongers, F., & Higgins, S. I. (2017). Allometric equations for integrating remote sensing imagery into forest monitoring programmes. *Global Change Biology*, 23, 177–190.
- Jupp, T. E., & Twiss, S. D. (2006). A physically motivated index of sub-grid-scale pattern. *Journal of Geophysical Research: Atmospheres*, 111 (D19).
- Keenan, T. F., & Niinemets, Ü. (2016). Global leaf trait estimates biased due to plasticity in the shade. *Nature Plants*, 3, 16201.
- Kim, Y., Knox, R. G., Longo, M., Medvigy, D., Hutya, L. R., Pyle, E. H., ... Moorcroft, P. R. (2012). Seasonal carbon dynamics and water fluxes in an Amazon rainforest. *Global Change Biology*, 18, 1322–1334.
- Knox, R. G., Longo, M., Swann, A. L. S., Zhang, K., Levine, N. M., Moorcroft, P. R., & Bras, R. L. (2015). Hydrometeorological effects of historical land-conversion in an ecosystem-atmosphere model of Northern South America. *Hydrological Earth System Science*, 19, 241–273.
- Konings, A. G., & Gentine, P. (2016). Global variations in ecosystem-scale isohydricity. *Global Change Biology*, 23, 891–905.
- Koven, C. D., Chambers, J. Q., Georgiou, K., Knox, R., Negron-Juarez, R., Riley, W. J., ... Jones, C. D. (2015). Controls on terrestrial carbon feedbacks by productivity vs. turnover in the CMIP5 Earth System Models. *Biogeosciences Discussions*, 12, 5211–5228.
- Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., ... Prentice, I. C. (2005). A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochemical Cycles*, 19(1), GB1015.
- Lagergren, F., Jönsson, A. M., Blennow, K., & Smith, B. (2012). Implementing storm damage in a dynamic vegetation model for regional applications in Sweden. *Ecological Modelling*, 247, 71–82.
- Lawrence, P. J., Feddema, J. J., Bonan, G. B., Meehl, G. A., O'Neill, B. C., Oleson, K. W., ... Thornton, P. E. (2012). Simulating the biogeochemical and biogeophysical impacts of transient land cover change and wood harvest in the Community Climate System Model (CCSM4) from 1850 to 2100. *Journal of Climate*, 25, 3071–3095.
- Lawrence, D. M., Hurr, G. C., Arneeth, A., Brovkin, V., Calvin, K. V., Jones, A. D., ... Seneviratne, S. I. (2016). The Land Use Model Intercomparison Project (LUMIP): Rationale and experimental design. *Geoscientific Model Development Discussion*, 9, 2973–2998.
- Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., ... Bonan, G. B. (2011). Parameterization improvements and functional and structural advances in version 4 of the Community Land Model. *Journal of Advances in Modeling Earth Systems*, 3(1), M03001.
- LeBauer, D. S., Wang, D., Richter, K. T., Davidson, C. C., & Dietze, M. C. (2013). Facilitating feedbacks between field measurements and ecosystem models. *Ecological Monographs*, 83, 133–154.
- Lefsky, M. A., Harding, D. J., Keller, M., Cohen, W. B., Carabajal, C. C., Del Bom Espirito-Santo, F., ... de Oliveira, R. (2005). Estimates of forest canopy height and aboveground biomass using ICESat. *Geophysical Research Letters*, 32, L22S02.
- Levine, N. M., Zhang, K., Longo, M., Baccini, A., Phillips, O. L., Lewis, S. L., ... Feldpausch, T. R. (2016). Ecosystem heterogeneity determines the ecological resilience of the Amazon to climate change. *Proceedings of the National Academy of Sciences of the USA*, 113, 793–797.
- Lewis, S. L., Phillips, O. L., Sheil, D., Vinceti, B., Baker, T. R., Brown, S., ... Lejoly, J. (2004). Tropical forest tree mortality, recruitment and turnover rates: Calculation, interpretation and comparison when census intervals vary. *Journal of Ecology*, 92, 929–944.
- Lindeskog, M., Arneeth, A., Bondeau, A., Waha, K., Seaquist, J., Olin, S., & Smith, B. (2013). Implications of accounting for land use in simulations of ecosystem services and carbon cycling in Africa. *Earth System Dynamics*, 4, 385–407.
- Lines, E. R., Coomes, D. A., & Purves, D. W. (2010). Influences of forest structure, climate and species composition on tree mortality across the eastern US. *PLoS One*, 5, p.e13212.
- Liou, K. N. (2002). *An introduction to atmospheric radiation*. New York: Academic Press.
- Lischke, H., Zimmermann, N. E., Bolliger, J., Rickebusch, S., & Löffler, T. J. (2006). TreeMig: A forest-landscape model for simulating spatio-temporal patterns from stand to landscape scale. *Ecological Modelling*, 199(4), 409–420.
- Lokupitiya, E., Denning, A. S., Schaefer, K., Ricciuto, D., Anderson, R., Arain, M. A., ... Ciais, P. (2016). Carbon and energy fluxes in cropland ecosystems: A model-data comparison. *Biogeochemistry*, 129, 53–76.
- Luo, Y. Q., Randerson, J. T., Abramowitz, G., Bacour, C., Blyth, E., Carvalhais, N., ... Friedlingstein, P. (2012). A framework for benchmarking land models. *Biogeosciences*, 9, 3857–3874.
- Mackay, D. S., Roberts, D. E., Ewers, B. E., Sperry, J. S., McDowell, N. G., & Pockman, W. T. (2015). Interdependence of chronic hydraulic dysfunction and canopy processes can improve integrated models of tree response to drought. *Water Resources Research*, 51, 6156–6176.
- Maréchal, I., Bartlett, M. K., Gaucher, P., Sack, L., & Chave, J. (2016). Causes of variation in leaf-level drought tolerance within an Amazonian forest. *Journal of Plant Hydraulics*, 3, e004.
- Matheny, A. M., Bohrer, G., Garrity, S. R., Morin, T. H., Howard, C. J., & Vogel, C. S. (2015). Observations of stem water storage in trees of opposing hydraulic strategies. *Ecosphere*, 6, 1–13.
- Matheny, A. M., Fiorella, R. P., Bohrer, G., Poulsen, C. J., Morin, T. H., Wunderlich, A., ... Curtis, P. S. (2016). Contrasting strategies of hydraulic control in two codominant temperate tree species. *Ecology*, 10, e1815. <https://doi.org/10.1002/eco.1815>
- McDowell, N. G., Beerling, D. J., Breshears, D. D., Fisher, R. A., Raffa, K. F., & Stitt, M. (2011). The interdependence of mechanisms underlying climate-driven vegetation mortality. *Trends in Ecology & Evolution*, 26, 523–532.
- McDowell, N. G., Coops, N. C., Beck, P., Chambers, J. Q., Gangogadage, C., Hicke, J. A., ... Meddens, A. J. (2015). Global satellite monitoring of climate-induced vegetation disturbances. *Trends in Plant Science*, 20, 114–123.
- McDowell, N. G., Fisher, R. A., Xu, C., Domec, J. C., Hölttä, T., Mackay, D. S., ... Limousin, J. M. (2013). Evaluating theories of drought-induced vegetation mortality using a multimodel-experiment framework. *New Phytologist*, 200, 304–321.
- McDowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., ... Yepez, E. A. (2008). Mechanisms of plant survival and mortality during drought: Why do some plants survive while others succumb to drought? *New Phytologist*, 178, 719–739.
- Medvigy, D., Clark, K. L., Skowronski, N. S., & Schäfer, K. V. R. (2012). Simulated impacts of insect defoliation on forest carbon dynamics. *Environmental Research Letters*, 7(4), 045703.
- Medvigy, D., Jeong, S. J., Clark, K. L., Skowronski, N. S., & Schäfer, K. V. (2013). Effects of seasonal variation of photosynthetic capacity on the carbon fluxes of a temperate deciduous forest. *Journal of Geophysical Research: Biogeosciences*, 118, 1703–1714.
- Medvigy, D., & Moorcroft, P. R. (2012). Predicting ecosystem dynamics at regional scales: An evaluation of a terrestrial biosphere model for the forests of northeastern North America. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 367, 222–235.
- Medvigy, D., Wofsy, S. C., Munger, J. W., Hollinger, D. Y., & Moorcroft, P. R. (2009). Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem Demography model version 2. *Journal of Geophysical Research: Biogeosciences*, 114(G1), G01002. <https://doi.org/10.1029/2008JG000812>
- Melton, J. R., & Arora, V. K. (2016). Competition between plant functional types in the Canadian Terrestrial Ecosystem Model (CTEM) v. 2.0. *Geoscientific Model Development*, 9, 323–361.

- Mercado, L. M., Huntingford, C., Gash, J. H., Cox, P. M., & Jöregreddy, V. (2007). Improving the representation of radiation interception and photosynthesis for climate model applications. *Tellus*, *59*, 553–565.
- Meron, E. (2011). Modeling dryland landscapes. *Mathematical Modelling of Natural Phenomena*, *6*, 163–187.
- Meron, E., Gilad, E., von Hardenberg, J., Shachak, M., & Zarmi, Y. (2004). Vegetation patterns along a rainfall gradient. *Chaos, Solitons & Fractals*, *19*, 367–376.
- Miller, A. D., Dietze, M. C., DeLucia, E. H., & Anderson-Teixeira, K. J. (2016). Alteration of forest succession and carbon cycling under elevated CO₂. *Global Change Biology*, *22*, 351–363.
- Moorcroft, P. R. (2006). How close are we to a predictive science of the biosphere? *Trends in Ecology & Evolution*, *21*, 400–407.
- Moorcroft, P. R., Hurtt, G. C., & Pacala, S. W. (2001). A method for scaling vegetation dynamics: The ecosystem demography model (ED). *Ecological Monographs*, *71*, 557–586.
- Morton, D. C., Rubio, J., Cook, B. D., Gastellu-Etchegorry, J. P., Longo, M., Choi, H., ... Keller, M. (2015). Amazon forest structure generates diurnal and seasonal variability in light utilization. *Biogeosciences Discussions*, *12*, 19043–19072.
- Muller-Landau, H. C., Condit, R. S., Chave, J., Thomas, S. C., Bohlman, S. A., Bunyavejchewin, S., ... Harms, K. E. (2006). Testing metabolic ecology theory for allometric scaling of tree size, growth and mortality in tropical forests. *Ecology Letters*, *9*, 575–588.
- Nabel, J. E. (2015). Upscaling with the dynamic two-layer classification concept (D2C): TreeMig-2L, an efficient implementation of the forest-landscape model TreeMig. *Geoscientific Model Development*, *8*, 3563–3577.
- Naudts, K., Ryder, J., McGrath, M. J., Otto, J., Chen, Y., Valade, A., ... Ghattas, J. (2015). A vertically discretised canopy description for ORCHIDEE (SVN r2290) and the modifications to the energy, water and carbon fluxes. *Geoscientific Model Development*, *8*, 2035–2065.
- Norman, J. M. (1979). Modelling the complete crop canopy. In B. J. Barfield, & J. F. Gerber (Eds.), *Modification of the aerial environment of plants*. St. Joseph, MI: American Society of Agricultural Engineers.
- Norman, J. M., Perry, S. G., Fraser, A. B., & Mach, W. (1979). Remote sensing of canopy structure. In *Proceedings of the Nth Conference on Agricultural and Forest Meteorology and the 4th Conference on Biometeorology*, 184–185.
- Oleson, K., Lawrence, D., Bonan, G., Drewniak, E., Huang, M., Koven, C., ... Yang, Z. (2013). Technical description of version 4.5 of the Community Land Model (CLM), NCAR Technical Note NCAR/TN-503+STR, 420 pp. <https://doi.org/10.5065/d6rr1w7m>
- Pacala, S. W., Canham, C. D., Saponara, J., Silander, J. A., Kobe, R. K., & Ribbens, E. (1996). Forest models defined by field measurements: Estimation, error analysis and dynamics. *Ecological Monographs*, *66*, 1–43.
- Pacala, S. W., & Tilman, D. (1994). Limiting similarity in mechanistic and spatial models of plant competition in heterogeneous environments. *American Naturalist*, *143*, 222–257.
- Pachzelt, A., Forrest, M., Rammig, A., Higgins, S. I., & Hickler, T. (2015). Potential impact of large ungulate grazers on African vegetation, carbon storage and fire regimes. *Global Ecology & Biogeography*, *24*, 991–1002.
- Pappas, C., Fatichi, S., & Burlando, P. (2016). Modeling terrestrial carbon and water dynamics across climatic gradients: Does plant trait diversity matter? *New Phytologist*, *209*, 137–151.
- Pavlick, R., Drewry, D. T., Bohn, K., Reu, B., & Kleidon, A. (2013). The Jena Diversity-Dynamic Global Vegetation Model (JeDi-DGVM): A diverse approach to representing terrestrial biogeography and biogeochemistry based on plant functional trade-offs. *Biogeosciences*, *10*, 4137–4177.
- Phillips, O. L., Van Der Heijden, G., Lewis, S. L., López-González, G., Aragão, L. E., Lloyd, J., ... Amaral, I. (2010). Drought–mortality relationships for tropical forests. *New Phytologist*, *187*, 631–646.
- Plaut, J. A., Yopez, E. A., Hill, J., Pangle, R., Sperry, J. S., Pockman, W. T., & McDowell, N. G. (2012). Hydraulic limits preceding mortality in a piñon–juniper woodland under experimental drought. *Plant, Cell & Environment*, *35*, 1601–1617.
- Pongratz, J., Reick, C. H., Raddatz, T., & Claussen, M. (2009). Effects of anthropogenic land cover change on the carbon cycle of the last millennium. *Global Biogeochemical Cycles*, *23*, GB4001. <https://doi.org/10.1029/2009GB003488>
- Powell, T. L., Galbraith, D. R., Christoffersen, B. O., Harper, A., Imbuzeiro, H., Rowland, L., ... Levine, N. M. (2013). Confronting model predictions of carbon fluxes with measurements of Amazon forests subjected to experimental drought. *New Phytologist*, *200*, 350–365.
- Poyatos, R., Aguadé, D., Galiano, L., Mencuccini, M., & Martínez-Vilalta, J. (2013). Drought induced defoliation and long periods of near-zero gas exchange play a key role in accentuating metabolic decline of Scots pine. *New Phytologist*, *200*, 388–401.
- Purves, D. W., Lichstein, J. W., Strigul, N. G., & Pacala, S. W. (2008). Predicting and understanding forest dynamics using a simple tractable model. *Proceedings of the National Academy of Sciences of the USA*, *105*, 17018–17022.
- Purves, D., & Pacala, S. (2008). Predictive models of forest dynamics. *Science*, *320*, 1452–1453.
- Reich, P. B. (2014). The world-wide ‘fast–slow’ plant economics spectrum: A traits manifesto. *Journal of Ecology*, *102*, 275–301.
- Reich, P. B., Rich, R. L., Lu, X., Wang, Y. P., & Oleksyn, J. (2014). Biogeographic variation in evergreen conifer needle longevity and impacts on boreal forest carbon cycle projections. *Proceedings of the National Academy of Sciences of the USA*, *111*, 13703–13708.
- Rodriguez-Iturbe, I., D’Odorico, P., Porporato, A., & Ridolfi, L. (1999). On the spatial and temporal links between vegetation, climate, and soil moisture. *Water Resources Research*, *35*, 3709–3722.
- Roy, S. B., Hurtt, G. C., Weaver, C. P., & Pacala, S. W. (2003). Impact of historical land cover change on the July climate of the United States. *Journal of Geophysical Research*, *108*(D24), 14. <https://doi.org/10.1029/2003JD003565>
- Sakschewski, B., Bloh, W., Boit, A., Rammig, A., Kattge, J., Poorter, L., ... Thonicke, K. (2015). Leaf and stem economics spectra drive diversity of functional plant traits in a dynamic global vegetation model. *Global Change Biology*, *21*, 2711–2725.
- Sanderson, B. M., Knutti, R., & Caldwell, P. (2015). A representative democracy to reduce interdependency in a multimodel ensemble. *Journal of Climate*, *28*, 5171–5194.
- Sato, H. (2009). Simulation of the vegetation structure and function in a Malaysian tropical rain forest using the individual-based dynamic vegetation model SEIB-DGVM. *Forest Ecology and Management*, *257*, 2277–2286.
- Sato, H., & Ise, T. (2012). Effect of plant dynamic processes on African vegetation responses to climate change: Analysis using the spatially explicit individual-based dynamic global vegetation model (SEIB-DGVM). *Journal of Geophysical Research-Biogeosciences*, *117*, G3017.
- Sato, H., Ito, A., & Kohyama, T. (2007). SEIB-DGVM: A new dynamic global vegetation model using a spatially explicit individual-based approach. *Ecological Modelling*, *200*, 279–307.
- Sato, H., Kobayahi, H., & Delbart, N. (2010). Simulation study of the vegetation structure and function in eastern Siberian larch forests using the individual-based vegetation model SEIB-DGVM. *Forest Ecology and Management*, *259*, 301–311.
- Scanlon, T. M., Caylor, K. K., Levin, S. A., & Rodriguez-Iturbe, I. (2007). Positive feedbacks promote power-law clustering of Kalahari vegetation. *Nature*, *449*, 209–212.
- Scheiter, S., Langan, L., & Higgins, S. I. (2012). Next-generation dynamic global vegetation models: Learning from community ecology. *New Phytologist*, *198*, 957–969.

- Schenk, H. J. (2006). Root competition: Beyond resource depletion. *Journal of Ecology*, *94*, 725–739.
- Scherstjanoi, M., Kaplan, J. O., & Lischke, H. (2014). Application of a computationally efficient method to approximate gap model results with a probabilistic approach. *Geoscientific Model Development*, *7*(4), 1543–1571.
- Scherstjanoi, M., Kaplan, J. O., Thürig, E., & Lischke, H. (2014). GAP-PARD: A computationally efficient method of approximating gap-scale disturbance in vegetation models. *Geoscientific Model Development*, *6*, 1517–1542.
- Schwalm, C. R., Williams, C. A., Schaefer, K., Anderson, R., Arain, M. A., Baker, I., ... Ciais, P. (2010). A model-data inter-comparison of CO₂ exchange across North America: Results from the North American Carbon Program site synthesis. *Journal of Geophysical Research: Biogeosciences*, *115*(G3), G00H05 <https://doi.org/10.1029/2009JG001229>.
- Schwinning, S., & Weiner, J. (1998). Mechanisms determining the degree of size asymmetry in competition among plants. *Oecologia*, *113*, 447–455.
- Sellers, P. J. (1985). Canopy reflectance, photosynthesis and transpiration. *International Institute of Remote Sensing*, *6*, 1335–1372.
- Serbin, S. P., Singh, A., Desai, A. R., Dubois, S. G., Jablonski, A. D., Kingdon, C. C., ... Townsend, P. A. (2015). Remotely estimating photosynthetic capacity, and its response to temperature, in vegetation canopies using imaging spectroscopy. *Remote Sensing of Environment*, *167*, 78–87.
- Shachak, M., Boeken, B., Groner, E., Kadmon, R., Lubin, Y., Meron, E., ... Ungar, E. D. (2008). Woody species as landscape modulators and their effect on biodiversity patterns. *BioScience*, *58*, 209–221.
- Shevliakova, E., Pacala, S. W., Malyshev, S., Hurtt, G. C., Milly, P. C. D., Caspersen, J. P., ... Crevoisier, C. (2009). Carbon cycling under 300 years of land use change: Importance of the secondary vegetation sink. *Global Biogeochemical Cycles*, *23*(GB2022). <https://doi.org/10.1029/2007GB003176>
- Shiklomanov, A. N., Dietze, M. C., Viskari, T., Townsend, P. A., & Serbin, S. P. (2016). Quantifying the influences of spectral resolution on uncertainty in leaf trait estimates through a Bayesian approach to RTM inversion. *Remote Sensing of Environment*, *183*, 226–238.
- Shrestha, R. K., Arora, V. K., & Melton, J. R. (2016). The sensitivity of simulated competition between different plant functional types to sub-grid-scale representation of vegetation in a land surface model. *Journal of Geophysical Research: Biogeosciences*, *121*, 809–828.
- Shuman, J. K., Shugart, H. H., & Krankina, O. N. (2014). Testing individual-based models of forest dynamics: Issues and an example from the boreal forests of Russia. *Ecological Modelling*, *10*, 102–110.
- Simard, M., Pinto, N., Fisher, J. B., & Baccini, A. (2011). Mapping forest canopy height globally with spaceborne lidar. *Journal of Geophysical Research: Biogeosciences*, *116*, G04021. <https://doi.org/10.1029/2011JG001708>
- Singh, A., Serbin, S. P., McNeil, B. E., Kingdon, C. C., & Townsend, P. A. (2015). Imaging spectroscopy algorithms for mapping canopy foliar chemical and morphological traits and their uncertainties. *Ecological Applications*, *25*, 2180–2197.
- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., ... Venevsky, S. (2003). Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ Dynamic Global Vegetation Model. *Global Change Biology*, *9*, 161–185.
- Smith, B., Prentice, I. C., & Sykes, M. T. (2001). Representation of vegetation dynamics in the modelling of terrestrial ecosystems: Comparing two contrasting approaches within European climate space. *Global Ecology & Biogeography*, *10*, 621–637.
- Smith, B., Samuelsson, P., Wramneby, A., & Rummukainen, M. (2011). A model of the coupled dynamics of climate, vegetation and terrestrial ecosystem biogeochemistry for regional applications. *Tellus*, *63A*, 87–106.
- Smith, B., Wärlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., & Zaehle, S. (2014). Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model. *Biogeosciences*, *11*, 2027–2054.
- Smolander, S., & Stenberg, P. (2005). Simple parameterizations of the radiation budget of uniform broadleaved and coniferous canopies. *Remote Sensing of Environment*, *94*, 355–363.
- Sperry, J. S., Adler, F. R., Campbell, G. S., & Comstock, J. P. (1998). Limitation of plant water use by rhizosphere and xylem conductance: Results from a model. *Plant, Cell & Environment*, *21*, 347–359.
- Sperry, J. S., & Love, D. M. (2015). What plant hydraulics can tell us about responses to climate-change droughts. *New Phytologist*, *207*, 14–27.
- Sperry, J. S., Venturas, M. D., Anderegg, W. R., Mencuccini, M., Mackay, D. S., Wang, Y., & Love, D. M. (2016). Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost. *Plant, Cell & Environment*, *40*, 816–830.
- Stark, S. C., Enquist, B. J., Saleska, S. R., Leitold, V., Schiatti, J., Longo, M., ... Oliveira, R. C. (2015). Linking canopy leaf area and light environments with tree size distributions to explain Amazon forest demography. *Ecology Letters*, *18*, 636–645.
- Stark, S. C., Leitold, V., Wu, J. L., Hunter, M. O., de Castilho, C. V., Costa, F. R., ... Keller, M. (2012). Amazon forest carbon dynamics predicted by profiles of canopy leaf area and light environment. *Ecology Letters*, *15*, 1406–1414.
- Stocker, B. D., Feissli, F., Strassmann, K. M., Spahni, R., & Joos, F. (2014). Past and future carbon fluxes from land use change, shifting cultivation and wood harvest. *Tellus Series B*, *66*(1), 23188. <https://doi.org/10.3402/tellusb.v66.23188>
- Strigul, N., Pristiniski, D., Purves, D., Dushoff, J., & Pacala, S. (2008). Scaling from trees to forests: Tractable macroscopic equations for forest dynamics. *Ecological Monographs*, *78*, 523–545.
- Swann, A. L. S., Longo, M., Knox, R. G., Lee, E., & Moorcroft, P. R. (2015). Future deforestation in the Amazon and consequences for South American climate. *Agricultural Forest Meteorology*, *214*, 12–24.
- Thomas, R. Q., Brookshire, E. N., & Gerber, S. (2015). Nitrogen limitation on land: How can it occur in Earth system models? *Global Change Biology*, *21*, 1777–1793.
- Thomas, R. Q., Hurtt, G. C., Dubayah, R., & Schilz, M. H. (2008). Using lidar data and a height-structured ecosystem model to estimate forest carbon stocks and fluxes over mountainous terrain. *Canadian Journal of Remote Sensing*, *34*(Suppl. 2), S351–S363.
- Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L., & Carmona-Moreno, C. (2010). The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: Results from a process-based model. *Biogeosciences*, *7*, 1991–2011.
- Thonicke, K., Venevsky, S., Sitch, S., & Cramer, W. (2001). The role of fire disturbance for global vegetation dynamics: Coupling fire into a Dynamic Global Vegetation Model. *Global Ecology and Biogeography*, *10*, 661–677.
- Trugman, A. T., Fenton, N. J., Bergeron, Y., Xu, X., Welp, L. R., & Medvigy, D. (2016). Climate, soil organic layer, and nitrogen jointly drive forest development after fire in the North American boreal zone. *Journal of Advances in Modeling the Earth System*, *8*, 1180–1209.
- van Wijk, M. T., & Rodriguez-Iturbe, I. (2002). Tree-grass competition in space and time: Insights from a simple cellular automata model based on ecohydrological dynamics. *Water Resources Research*, *38*(9), 1179. <https://doi.org/10.1029/2001WR000768>
- Verheijen, L. M., Aerts, R., Brovkin, V., Cavender-Bares, J., Cornelissen, J. H., Kattge, J., & Bodegom, P. M. (2015). Inclusion of ecologically based trait variation in plant functional types reduces the projected land carbon sink in an earth system model. *Global Change Biology*, *21*, 3074–3086.

- Wang, Y. P., Law, R. M., & Pak, B. (2010). A global model of carbon, nitrogen and phosphorus cycles for the terrestrial biosphere. *Biogeosciences*, 7, 2261–2282.
- Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., ... Ise, T. (2011). MIROC-ESM 2010: Model description and basic results of CMIP5-20c3m experiments. *Geoscientific Model Development*, 4, 845–872.
- Weiss, M., Miller, P. A., van den Hurk, B. J. J. M., van Noije, T., Stefanescu, S., Haarsma, R., ... Schurgers, G. (2014). Contribution of dynamic vegetation phenology to decadal climate predictability. *Journal of Climate*, 27, 8563–8577.
- Weng, E., Fariior, C. E., Dybzinski, R., & Pacala, S. W. (2016). Predicting vegetation type through physiological and environmental interactions with leaf traits: Evergreen and deciduous forests in an earth system modeling framework. *Global Change Biology*, 23, 2482–2498.
- Weng, E. S., Malyshev, S., Lichstein, J. W., Fariior, C. E., Dybzinski, R., Zhang, T., ... Pacala, S. W. (2015). Scaling from individual trees to forests in an Earth system modeling framework using a mathematically tractable model of height-structured competition. *Biogeosciences*, 12, 2655–2694.
- West, G. B., Brown, J. H., & Enquist, B. J. (1999). A general model for the structure and allometry of plant vascular systems. *Nature*, 400, 664–667.
- Widowski, J. L., Mio, C., Disney, M., Adams, J., Andredakis, I., Atzberger, C., ... Colombo, R. (2015). The fourth phase of the radiative transfer model intercomparison (RAMI) exercise: Actual canopy scenarios and conformity testing. *Remote Sensing of Environment*, 169, 418–437.
- Widowski, J. L., Taberner, M., Pinty, B., Bruniquel-Pinel, V., Disney, M., Fernandes, R., ... Leblanc, S. (2007). Third Radiation Transfer Model Intercomparison (RAMI) exercise: Documenting progress in canopy reflectance models. *Journal of Geophysical Research: Atmospheres*, 112 (D9), D09111. <https://doi.org/10.1029/2006JD007821>
- Wilkenskjeld, S., Kloster, S., Pongratz, J., Raddatz, T., & Reick, C. (2014). Comparing the influence of net and gross anthropogenic land-use and land-cover changes on the carbon cycle in the MPI-ESM. *Biogeosciences*, 11, 4817–4828.
- Williams, M., Bond, B. J., & Ryan, M. G. (2001). Evaluating different soil and plant hydraulic constraints on tree function using a model and sap flow data from ponderosa pine. *Plant, Cell & Environment*, 24, 679–690.
- Woodward, F. I., & Lomas, M. R. (2004). Vegetation dynamics—simulating responses to climatic change. *Biological Reviews*, 79, 643–670.
- Wramneby, A., Smith, B., & Samuelsson, P. (2010). Hotspots of vegetation-climate feedbacks under future greenhouse forcing in Europe. *Journal of Geophysical Research*, 115, D21119.
- Xu, C., Fisher, R., Wullschleger, S. D., Wilson, C. J., Cai, M., & McDowell, N. G. (2012). Toward a mechanistic modeling of nitrogen limitation on vegetation dynamics. *PLoS One*, 7, e37914.
- Xu, C., McDowell, N. G., Sevanto, S., & Fisher, R. A. (2013). Our limited ability to predict vegetation dynamics under water stress. *New Phytologist*, 200, 298–300.
- Xu, X., Medvigy, D., Powers, J. S., Becknell, J., & Guan, K. (2016). Diversity in plant hydraulic traits explains seasonal and inter-annual variations of vegetation dynamics in seasonally dry tropical forests. *New Phytologist*, 212, 80–95.
- Zaehle, S., & Dalmonech, D. (2011). Carbon-nitrogen interactions on land at global scales: Current understanding in modelling climate biosphere feedbacks. *Current Opinion in Environmental Stability*, 3, 311–320.
- Zaehle, S., & Friend, A. D. (2010). Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates. *Global Biogeochemical Cycles*, 24(1), GB1005. <https://doi.org/10.1029/2009GB003521>
- Zaehle, S., Medlyn, B. E., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hickler, T., ... Jain, A. (2014). Evaluation of 11 terrestrial carbon–nitrogen cycle models against observations from two temperate Free-Air CO₂ Enrichment Studies. *New Phytologist*, 202, 802–822.
- Zeppel, M. J., Adams, H. D., & Anderegg, W. R. (2011). Mechanistic causes of tree drought mortality: Recent results, unresolved questions and future research needs. *New Phytologist*, 192, 800–803.
- Zeppel, M., Macinnis-Ng, C., Palmer, A., Taylor, D., Whitley, R., Fuentes, S., ... Eamus, D. (2008). An analysis of the sensitivity of sap flux to soil and plant variables assessed for an Australian woodland using a soil–plant–atmosphere model. *Functional Plant Biology*, 35, 509–520.
- Zhang, K., Castanho, A. D. A., Galbraith, D. R., Moghim, S., Levine, N., Bras, R. L., ... Moorcroft, P. R. (2015). The fate of Amazonian ecosystems over the coming century arising from changes in climate, atmospheric CO₂ and land-use. *Global Change Biology*, 21, 2569–2587.

How to cite this article: Fisher RA, Koven CD, Anderegg WRL, et al. Vegetation demographics in Earth System Models: A review of progress and priorities. *Glob Change Biol.* 2018;24:35–54. <https://doi.org/10.1111/gcb.13910>