

ATOLL RESEARCH BULLETIN

NO. 442

CRUSTACEA DECAPODA OF FRENCH POLYNESIA

(ASTACIDEA, PALINURIDEA, ANOMURA, BRACHYURA)

BY

JOSEPH POUPIN

ISSUED BY
NATIONAL MUSEUM OF NATURAL HISTORY
SMITHSONIAN INSTITUTION
WASHINGTON, D.C., U.S.A.
APRIL 1996

ATOLE RESEARCH BULLETIN

NO. 443

CREATIVITY DEVELOPMENT OF REINACH FOR ANESTHESIA
(ANESTHESIOLOGY, PALINDROME, ANOMURA, BRAUCHIGUS)

BA

TOPICAL TOPIC

REVIEWED BY
NATIONAL MUSEUM OF NATURAL HISTORY
SMITHSONIAN INSTITUTION
WASHINGTON, D.C., U.S.A.
APRIL 1969

C O N T E N T S

SUMMARY	1
INTRODUCTION	2
HISTORICAL	3
CONVENTIONS	6
LIST OF THE SPECIES	8
INFRA-ORDER ASTACIDEA	8
FAMILY ENOPLOMETOPIDAE	8
INFRA-ORDER PALINURIDEA	9
FAMILY PALINURIDAE	9
FAMILY SYNAXIDAE	10
FAMILY SCYLLARIDAE	10
INFRA-ORDER ANOMURA	11
FAMILY COENOBITIDAE	11
FAMILY DIOGENIDAE	13
FAMILY PAGURIDAE	18
FAMILY GALATHEIDAE	19
FAMILY PORCELLANIDAE	20
FAMILY ALBUNEIDAE	22
FAMILY HIPPIDAE	23
INFRA-ORDER BRACHYURA	23
FAMILY DROMIIDAE	23
FAMILY DYNOMENIDAE	24
FAMILY RANINIDAE	24
FAMILY CALAPPIDAE	25
FAMILY LEUCOSIIDAE	26
FAMILY MAJIDAE	26
FAMILY PARTHENOPIDAE	28
FAMILY EUMEDONIDAE	29
FAMILY PORTUNIDAE	29
SUBFAMILY CATOPTRINAE	29
SUBFAMILY CAPHYRINAE	29
SUBFAMILY PORTUNINAE	30
SUBFAMILY PODOPHTHALMINAE	37
FAMILY XANTHIDAE	37
SUBFAMILY POLYDECTINAE	37
SUBFAMILY CYMOINAE	38
SUBFAMILY TRICHIINAE	38
SUBFAMILY LIOMERINAE	38
SUBFAMILY EUXANTHINAE	41
SUBFAMILY ACTAEINAE	42
SUBFAMILY ZOZIMINAE	45
SUBFAMILY XANTHINAE	48
SUBFAMILY PANCOPEINAE	51
SUBFAMILY KRAUSSIINAE	51
SUBFAMILY ETISINAE	51
SUBFAMILY CHLORODIINAE	53

CONTENTS

FAMILY TRAPEZIIDAE.....	57
FAMILY PILUMNIDAE	61
FAMILY CARPILIIDAE	62
FAMILY MENIPPIDAE	63
FAMILY GECARCINIDAE	65
FAMILY GRAPSIDAE	66
SUBFAMILY GRAPSINAE	66
SUBFAMILY VARUNINAE	69
SUBFAMILY SESARMINAE.....	69
SUBFAMILY PLAGUSIINAE.....	71
FAMILY PINNOTHERIDAE.....	72
FAMILY OCYPODIDAE.....	72
FAMILY CRYPTOCHIRIDAE	74
FAMILY HYMENOSOMATIDAE.....	75
FAMILY INCERTAE SEDIS	75
DISCUSSION	76
LITERATURE CITED	81
ACKNOWLEDGEMENT	95
APPENDICES.....	95
INDEX.....	101

CRISTACEA DECAPODA (DEPTHE) OF POLYNESIA


(ASTACOIDEA, MALINTIDAE, ANOMIADA, BRACHYURA)

and


with

and


</div




TUAMOTU



AUSTRAL



FRENCH POLYNESIA



CRUSTACEA DECAPODA OF FRENCH POLYNESIA

(ASTACIDEA, PALINURIDEA, ANOMURA, BRACHYURA)

BY

JOSEPH POUPIN

S U M M A R Y

From a bibliographic compilation and, to a lesser extent, from material collected in the field, 401 littoral and sublittoral decapods (Palinura, Anomura, Brachyura), are reported from French Polynesia. The Brachyura prevail, with 313 species, mainly Xanthidae (123 species), Portunidae (54 species), and Grapsidae (35 species). The Anomura are represented by 74 species, and the Palinura by only 14 species. The list of the deep species, *i.e.* living in depths of 100m or more, is updated. Ninety-two species are listed, making a total of 493 Polynesian species.

Amongst the material recently collected, 16 species are recorded for the first time in the area: *Calcinus guamensis*, *Calcinus imperialis*, *Dardanus australis*, *Dardanus brachyops*, *Albunea speciosa*, *Parthenope contrarius*, *Portunus macrophthalmus*, *Portunus orbitosinus*, *Thalamita danae*, *Thalamita macropus*, *Thalamita mitsiensis*, *Thalamita philippinensis*, *Quadrella maculosa*, *Planes cyaneus*, *Percnon guinotae*, and *Macrophthalmus serenei*. Moreover, after the examination of the type material, *Ruppelia granulosa* A. Milne Edwards, 1867, originally describe from the Marquesas, is here proposed as a junior synonym of *Lydia annulipes* (H. Milne Edwards, 1834).

Only 8 species, related to well defined species, are known solely from French Polynesia: *Parribacus holthuisi*, *Micropagurus polynesiensis*, *Nucia rosea*, *Nursia mimetica*, *Acanthophrys cristimanus*, *Lissocarcinus elegans*, *Ozius tricarinatus*, and *Macrophthalmus consobrinus*. For some of them, however, it is probable that their distributions extend at least to western Polynesia.

The French Polynesian fauna is typically Indo-West Pacific in its composition, with few endemic forms, and a low diversity compared to the Indo-Malaysian area. It includes, however, many more species than the Hawaiian fauna, possibly because the Polynesian islands are less isolated than the Hawaiian islands.

The Society, Tuamotu, and Gambier archipelagos have been well investigated, with numerous expeditions organised in these areas. In contrast, the Austral and Marquesas Islands, still remain poorly known. The French Polynesian fauna is more or less homogenous, with few regionally distinctive features. The single obvious exception is for the isolated southernmost islands, Rapa and Marotiri, subjected to a subtropical climate. In these islands, species that are very common elsewhere, are missing (*Coenobita*

Service Mixte de Surveillance Radiologique et Biologique, SMSRB, B.P. 208, 91311 Montlhéry Cedex, France, and Muséum national d'Histoire naturelle, Laboratoire de Zoologie des Arthropodes, 61 rue Buffon, 75005 Paris.

Manuscript received 8 December 1995; revised 28 March 1996

perlatus, *Birgus latro*, *Cardisoma carnifex*), and, on the contrary, at least one common species is still unknown in the northern part of French Polynesia (*Panulirus pascuensis*).

INTRODUCTION

What are the decapod crustacea known from French Polynesia? The answer to this, apparently simple question, would be very helpful for determinating the species collected during ecological studies. Moreover, from a biogeographical point of view, a check list of the species reaching this area, at the eastern limit of the Indo-West Pacific province, would be very interesting. The aim of this work therefore, is to answer this question by drawing up, mainly from a compilation of systematic and ecological studies, a list of the French Polynesian crustacea, the scope of the subject being restricted to littoral and sublittoral, Palinura (Astacidea and Palinuridae), Anomura, and Brachyura.

The check list given here has been mostly compiled from bibliographical records. In a first step, the most important works dealing with the French Polynesian crustacea have been consulted. They are the works by DANA (1852b, 1855), HELLER (1865), NOBILI (1907), RATHBUN (1907), BOONE (1934, 1935), HOLTHUIS (1953), FOREST & GUINOT (1961), and more recently, those by ODINETZ (1983), MONTEFORTE (1984), GUINOT (1985), MARQUET (1988), and PEYROT-CLAUSADE (1989). In a second step, the names of the species have been updated, for changes in the generic classification, or species reduced to synonymy, by looking through more general works, like the ones by HOLTHUIS (1991), SAKAI (1976), or SERÈNE (1984). This research has been completed by additional consultation of two bibliographical journals, the *Zoological Records* and the *Current Contents*, and by randomly looking through reprints available at the Muséum national d'Histoire naturelle, Paris. This last step was sometimes very fruitful, with additonal species mentioned in the area, often very discreetly.

For the most important families, the main works consulted during this research are the following:

The Palinuridae have been found in the recent catalogue of the *Marine lobsters of the world* by HOLTHUIS (1991), and in his revision of the Scyllaridae (HOLTHUIS, 1985).

The pagurids (Coenobitidae, Diogenidae, Paguridae) have been first searched through the work by NOBILI (1907) and the studies by FOREST, published between 1951 and 1956. Additional information has been found in the work of LEWINSOHN (1969), the report of RAHAYU (1988), and revisions of the genera, *Aniculus* (FOREST, 1984), *Calcinus* (MORGAN, 1991), *Catapaguroides* (DE SAINT LAURENT, 1968, 1970), *Clibanarius* (RAHAYU & FOREST, 1992), *Pagurixus* (MCLAUGHLIN & HAIG, 1984), and *Trizopagurus* (FOREST, 1995).

Except for NOBILI's (1907) work and, for a single species, BOONE's (1935) work, the few Galatheidae known from French Polynesia come from the ecological works by PEYROT-CLAUSADE (1977a, b, 1989), KROPP & BIRKELAND (1981), and ODINETZ (1983).

Almost all the Porcellanidae have been found in the works published by HAIG, between 1964 and 1992, HAIG & KROPP (1987), and KROPP (1983, 1986).

For the Brachyura, the beginning of the research has been greatly facilitated by the important studies of FOREST & GUINOT (1961), MONTEFORTE (1984), and GUINOT (1985). More information has been found in: MCLAY (1991, 1993), for the Dromiidae; GALIL & CLARK (1994), for the Calappidae of the

genus *Matuta*; GRIFFIN & TRANTER (1986), for the Majidae; STEPHENSON (1972, 1976), STEPHENSON & REES (1961, 1967), and MOOSA (1979), for the Portunidae; ODINETZ (1983, 1984a), and the works by GALIL, and co-authors, published between 1985 and 1990, for the Trapeziidae; SERÈNE (1984), for the Xanthidae, and CLARK & GALIL (1993) for the *Pilodius* xanthids; CROSNIER (1984), for the Carpiliidae and Menippidae; TÜRKAY (1973, 1974), for the Gecarcinidae; and SAKAI & TÜRKAY (1976), CRANE (1975), and BARNES (1977), for the Ocypodidae of the genera, *Ocypode*, *Uca*, and *Macrophthalmus*, respectively.

This bibliographical compilation has been completed, in a much more limited way, by the study of some specimens collected on the field, during the last few years. From the addition of this material, 16 species are recorded for the first time in French Polynesia.

The deep-water crustacea, *ie* collected from 100m and beyond, have already been listed in POUPIN (1996), with full references on origins of the collections, and depth ranges. A simple list is produced here, updated by inclusion of species described after the first compilation, or recently collected (*cf.* Appendices).

HISTORICAL

OLD VOYAGES: 1820-1900

At that time, the crustacea were collected during the exploring expeditions made around the world by large sailing vessels. DUPERREY, on board the *Coquille* (1822-1825), is one of the first to bring back some species from Tahiti and Bora Bora. They were studied by GUÉRIN-MÉNEVILLE (1829, 1838) who dedicated to DUPERREY a small ocypodid crab from Bora Bora, *Gelasimus Duperreyi* (now *Uca tetriconon*).

About 10 years later (1837-1840), DUMONT D'URVILLE, chief officier of DUPERREY on the *Coquille*, sailed again in French Polynesia, commanding the *Astrolabe* and the *Zélée*. His vessels visited the Gambier, Marquesas (Nuku Hiva), and Society Islands. JACQUINOT (1852), naturalist, commanding the *Zélée*, mentioned a dozen of species from the area, and described some from the Gambier Islands, including the small *Ocypode pallidula*, common on the white sandy beaches of Aukena island.


At the same time, the Americans, worried about participating, like the Europeans, in the discovery of remote marine areas, launched their first round the world campaign, the great *U.S. Exploring Expedition* (1838-1842). The squadron of 6 vessels, commanded by WILKES, left Norfolk in August, 1838. At least four vessels cruised in French Polynesia: the *Flying Fish*, *Peacock*, *Porpoise*, and the *Vincennes*. A great part of the collections from the Tuamotu Islands was lost during the wreck of the *Peacock*, on the banks of the Columbia river, however, the Polynesian material, about 10 Anomura and 50 Brachyura, studied by DANA (1851, 1852a-b, 1855), represents the most important collection from that area. DANA describes several new species, from Tahiti (*Phymodius monticulosus*, *Trapezia areolata*), and the Tuamotu Islands (*Globopilumnus globosus*, *Liomera tristis*, *Plagusia speciosa*, *Thalamita integra*, *Trapezia bella*).

Between 1857 and 1859, the Austrian frigate, *Novara*, put in at Tahiti, during her sea voyage around the world. HELLER (1862, 1865) studied the Crustacea of this campaign. He recorded 54 species from

Tahiti, and described several, for example the colourful *Calcinus nitidus*, and the small gecarcinid, *Epigrapsus politus*.

Limited collections were also made at Tahiti by the famous British H.M.S. *Challenger* (1873-1876). They are discreetly mentioned in the works of HENDERSON (1888), for the Anomura, MIERS (1886), for the Brachyura, and BANERJEE (1960), for the grapsid crabs.

This era ends with the voyages of the American ship *Albatross* (1899-1900, and 1900-1905), and new collections in the Society, Tuamotu, Gambier, and Marquesas Islands. RATHBUN (1907) studied the Brachyura collected by this vessel. She recorded 85 Polynesian species and described, for example, *Pachygrapsus fakaravensis*, a grapsid very common in the Tuamotu Islands, named after the large atoll of Fakarava.



French frigate *La Coquille* at anchor in Matavai bay, Tahiti (1823)

(Drawing by Jules-Louis LEJEUNE. Courtesy of HORIZON Magazine)

BEGINNING OF THE 20TH CENTURY: 1900-1967

The voyages around the world have ended and the collections are now made by people living in French Polynesia. The most striking in that respect is certainly SEURAT, the head of a small Zoological laboratory once established at Rikitea, Gambier Islands. Between 1902 and 1905 he gathered an important collection from the Gambier Islands, and also from the Tuamotu Islands, at Hao and Marutea South. This material was studied by NOBILI (1906, 1907) with more than 130 Polynesian species, belonging to the groups here concerned, and with the description of a score of new species, for example *Thalamita gatavakensis*, or *Thalamita seurati*. FOREST (1951), for *Calcinus seurati* and *Calcinus spicatus* spp. nov., STEPHENSON & REES (1961), for *Portunus guinotae* sp. nov., and FOREST & GUINOT (1961), in their study on the Polynesian Brachyura, have mentioned again the material collected by SEURAT.

Gilbert RANSON, of the Malacology department, Muséum national d'Histoire naturelle, Paris, also collected many specimens. In 1952, during a stay of several months, especially on the atoll of Hikueru, he gathered numerous scyllarids, pagurids, and crabs. The first two groups have been studied by FOREST (1953, 1954), with description of 5 new species, including *Parribacus holthuisi* and *Clibanarius ransonii*. The third is studied by FOREST & GUINOT (1961), who, in grouping RANSON and SEURAT material, and some smaller collections, such as the one made by CHABOIS, a teacher at the *Paul Gauguin* school, Papeete, have registered about 100 crabs, 21 as new records, with some new species, such as *Pilumnus ransonii*. The same year, MORRISON, as a part of the *Pacific Sciences Board's Coral Atoll Program*, sampled the most common Crustacea of Raroia atoll, and, to a less extent, of Pukapuka, Takume, and Tahiti. From that material HOLTHUIS (1953) produced a list of 70 species, and mentioned for the first time *Hippa ovalis*, from Tahiti.

During these years, a few expeditions, even if they no longer have the nature of great campaigns around the world, still visited the Polynesian Islands. For example, SENDLER (1923) recorded about 30 species from Makatea, Rimatara, and Tahiti, from the collections made during the *Hanseatischen Südsee-Expedition*. Some of them, like *Coenobita cavipes* or the gecarcinid *Discoplax longipes*, have never been collected since. In 1931, the yacht *Alva* explored the Marquesas (Nuku Hiva) and the Society Islands (Bora Bora, Raiatea, Tahiti). BOONE (1934, 1935) mentioned about 40 species collected during this cruise, and described two crabs, *Actaeomorpha alvae* and *Lissocarcinus elegans*.

Two important expeditions, at an interval of 10 years, mark the end of this period. In 1957, the Americans organised the *Smithsonian Bredin Expedition*, which visited the Society and the Tuamotu Islands. The portunids were studied by STEPHENSON & REES (1967) and STEPHENSON (1976), with about 30 species, including some new records like *Portunus iranjae*, *Thalamita corrugata*, or *Thalamita quadrilobata*. GALIL (1985) and, more recently, FOREST (1995), in their works on the genera *Tetraloides* and *Trizopagurus*, respectively, also studied the material of this expedition. In 1967, the boat *Pele*, during the *Marquesas Expedition*, visited the Marquesas, Tuamotu, Society, Gambier Islands, and, in the neighbourhood, the small island of Pitcairn (HARALD, 1967). The crustacea of this campaign, deposited in Washington and Perth Museums, have been studied by STEPHENSON (1976), with some fifteen portunids, and appear, more discreetly, in the studies by SERÈNE (1972), for *Palapedia marquesas* sp. nov., GALIL & LEWINSOHN (1985), for *Trapezia tigrina*, or HOLTHUIS (1985), for *Parribacus holthuisi*.

MODERN PERIOD.

In 1966, with the installation of the *Centre d'Expérimentation du Pacifique*, several scientific investigations were made, mainly on the atoll of Moruroa. Crustacea collected during these investigations are mentioned in some systematic works, such as DE SAINT LAURENT (1967), for *Catapaguroides fragilis*, or GUINOT (1979) for *Lophozozymus superbus*, and in ecological studies, for example in CHEVALIER *et al.* (1968), SALVAT & RENAUD-MORAND (1969), and LABOUTE & RICHER DE FORGES (1986). These latter, during the expedition of the old minesweeper *Paimpolaise*, in the south of French Polynesia (MacDonald bank), have made the first Polynesian record of *Panulirus pascuensis*, originally described from Easter island. From these different campaigns, unstudied collections are still deposited at the *Muséum national d'Histoire naturelle*, Paris, for example the crustacea collected by PLESSIS, from which we give here the first record of *Thalamita danae*. Others collections were made during the campaigns of the fishing boat *Marara*, used by the *Direction des Centres d'Expérimentations Nucléaires* for monitoring the marine environment, as a part of the radiological safety program in French Polynesia. Although they mainly

concerned the deep fauna (POUPIN, 1996), some of the most common littoral and sublittoral species were also collected, and were presented by POUPIGN (1994a), in a small illustrated document.

In 1971, the French *École Pratique des Hautes Études*, established its research center in French Polynesia. First located at Tahiti, it was afterwards transferred to Opunohu Bay, Moorea Island, and is now denominated *Centre de Recherche Insulaire et Observatoire de l'Environnement* (CRIODE). Several important works have been done by the students or researchers of this center. They are mostly ecological studies with, however, about 30 new records for the area. The cryptozoa was studied by PEYROT-CLAUSADE, at Moorea (1977, 1985), and Tikehau (1989). The study of this small fauna was completed by NAIM (1980) with a dozen of species associated with the algae, at Tiahura, Moorea. The crustacean associates of the coral *Pocillopora*, were studied by KROPP & BIRKELAND (1981), and by ODINETZ (1983, 1984a, b) who described two new *Trapezia* species, *Trapezia serenei* and *T. punctimanus*. MONTEFORTE (1984) in his *Contribution à la connaissance de la faune carcinologique de Polynésie française*, collected and studied more than 110 species, some of them, like *Calcinus minutus*, *Calappa calappa*, or *Esisus anaglyptus*, being new records. More recently, a dozen common species were recorded from the atoll of Nukutipipi by MERSCHARDT-SALVAT (1991), and the freshwater collections made by MARQUET (1988, 1991, 1993), have given two new grapsid records, *Varuna litterata*, from the Society Islands, and *Ptychognathus easteranus*, from the Austral and Marquesas Islands.

C O N V E N T I O N S

STUDIES INCLUDED

We have included only studies in which material from French Polynesia has been actually examined, excluding works where "French Polynesia" appears only in the "Distribution". These are followed by the indication, in parenthesis, of the island(s) where the material originated. Recent revisions and general syntheses, from which the names of the species are updated (changes in generic classification; synonymies), or useful in different aspects, have been added; these are followed by mention such as "Syn." or "Key", in parenthesis. Some works have been included, that do not mention new collections. These are: the important syntheses about French Polynesia, like FOREST & GUINOT (1962), followed by "Biogeography", or GUINOT (1985), followed by "List"; the ecological studies focusing on a particular island, such as DELESALLE (1985), for the atoll of Mataiva, or SALVAT & RICHARD (1985), for the atoll of Takapoto; general works dedicated to the fauna of French Polynesia, such as SEURAT (1934), CHABOIS L. & F. (1954), or more recently the *Encyclopédie de la Polynésie* (cf. CHARLEUX, 1986 and SALVAT, 1986a-c) and the books by PARDON (1992) and BONVALLOT *et al.* (1994). For these latter, however, only the species illustrated, usually in colour, have been cited.

No distinction has been made between systematic and ecological works. In her list of the Brachyura of French Polynesia, GUINOT (1985) has sometimes considered the record of a species doubtful (name followed by a ?), when it was known only from an ecological paper. As these works can easily be identified from the references, the reader will be able to judge for himself. It is clear, however, that the revision of these collections would be important, but it is often difficult, or even impossible, to retrieve the material.

LOCATIONS

The unit of location is the island (see map). When the name of a village, a particular locality, or a small islet on the reefal crown (*Motu* in Polynesian language), was indicated, the name of the corresponding island is mentioned with the following presentation: "Gatavake" = Mangareva, "Ohura" = Hao, or "Taiohae" = Nuku Hiva. The same presentation is adopted for corrections of obvious mistakes: "Tickahau" = Tikehau, "Timoe" = Temoe, or "Fakaina" = Fakahina. The Gambier Islands have a particular configuration, with 8 mountainous islands surrounded by a common reefal crown. In that particular case, we have considered as real islands three *Motu* of the external crown: Puaumu, Tarauru-Roa, and Vaiatekeue.

Sometimes, in the oldest works, the names of the islands were old names, no longer now in use. They are translated into modern names by using MOTTLER's (1986) work, and the following presentation: "*Carlshoff*" = Aratika, "*Clermont-Tonnerre*" = Reao, or "*Eimeo*" = Moorea.

When no particular location was specified, the reference is just followed by "French Polynesia".

CLASSIFICATION

The classification approximately follows BOWMAN & ABELE (1982), and, for the Xanthoidea, SERÈNE (1984). Subfamily ranks has been indicated only within the most important families: Portunidae, Xanthidae, and Grapsidae. The presentation has been clarified by ignoring subgeneric names in the check list. However, if they were used in the works cited, they appear in the references.

LITTORAL, SUBLITTORAL AND DEEP SPECIES

These three groups are here defined in the following way: littoral species are commonly collected on the reef, and in depth of few meters only; sublittoral species are collected from about 10m to 100m; and deep species are collected from 100m and deeper. It is sometimes difficult to classify the species according to these three groups, especially because our knowledge about the deep distribution of numerous species is often very limited, and will have to be revised in the future. As an example, the maximal depth known for some species has been increased, here, sometimes considerably, from collections made by traps. Some littoral forms have been found unusually deep (*Carpilius convexus*, 60m; *Charybdis paucidentata*, 100m). They are qualified as "Littoral to sublittoral" species. Sublittoral forms have been sometimes collected far beyond 100m (*Dardanus brachyops*, 110-300m; *Dromia wilsoni*, 190-350m; *Thalamita spinifera*, 42-200m), and, in contrast, deep forms have been found in less than 100m (*Palibythus magnificus*, 70-240m; *Scyllarus aurora*, 90-300m; *Alainodaeus rimatara*, 90-350m). They are qualified as "Sublittoral to deep" species. Because of these difficulties, 12 species included in this work were also listed with the deep species (POUPIN, 1996; see Appendices 1, species with a *).

It is important to realise that, in several cases, these classifications are questionable, and often only reflect the poor information that we have on that subject. For example, a species like *Oreotlos potanus*, known by a single specimen, has been included with the deep species, to within 1m (101m). Considering the limited accuracy of the measures at sea, *O. potanus* could have been reasonably considered as a sublittoral species.

OTHERS CONVENTIONS

When a "?" appears in front of the name of a species, it always means that the doubt on that name is ours. When it is an hesitation expressed in the work consulted, it is mentioned after the reference by "with a ?".

When a species has been reduced to synonymy, two cases are considered. If the species was not originally described from French Polynesia, the full name (author and date) appears at the end of the references, after "SYNONYMS". Otherwise, this information appears clearly with the reference, and is not repeated again. In both cases the origin of the synonymy is to be found in the work followed by "Syn.". The synonymies are restricted to French Polynesian species only.

As far as possible we have tried to avoid partial identifications. When a generic name is only available, the reference is not considered in the main list, but appears separately, in Appendices 2. Preliminaries identifications (*aff.* or *cf.*) have been retained only when the species referred to is not yet recorded from French Polynesia (for example, *Actaea* *aff. glandifera* in PEYROT-CLAUSADE, 1989: 111). Otherwise, they appear under the species referred to, after "RELEVANT MATERIAL" (see for example, *Lybia* *cf. caestifera* in MONTEFORTE, 1984: 171, under *Lybia caestifera*).

The following abbreviations are used: BM (Natural History Museum, London); CRIODE (Centre de Recherche Insulaire et Observatoire de l'Environnement, Moorea); MNHN (Muséum national d'Histoire naturelle, Paris); USNM (National Museum of Natural History, Washington).

L I S T O F T H E S P E C I E S

INFRA-ORDER ASTACIDEA

FAMILY ENOPLOMETOPIDAE

Enoplometopus holthuisi Gordon, 1968

DISTRIBUTION. — Tuamotu - Sublittoral.

REFERENCES. — *Enoplometopus holthuisi* - BONVALLOT *et al.*, 1994: 144-145, photograph (Tuamotu).

REMARK. — At least another *Enoplometopus* is present in French Polynesia (cf. *Enoplometopus* sp. nov. in POUPIN *et al.*, 1990: 16, pl. 3c).

INFRA-ORDER PALINURIDEA

FAMILY PALINURIDAE

Justitia longimanus (H. Milne Edwards, 1837)

DISTRIBUTION. — Austral (Rurutu); Society (Bora Bora, Raiatea, Tahiti); Tuamotu (Makemo, Maria, Tenarunga) - *Sublittoral to deep*.

REFERENCES. — *Justitia longimana* - POUPIN *et al.*, 1990: 16 (French Polynesia). — *Justitia longimanus* - POUPIN, 1994b: 46, fig. 3e', pl. 1d, 2d (Bora Bora, Tahiti, Tenarunga, Rurutu; 62-160m); 1996: in press (Bora Bora, Makemo, Maria, Raiatea, Rurutu, Tenarunga; 80-190m).

Panulirus homarus (Linné, 1758)

DISTRIBUTION. — Marquesas (Fatu Hiva, Nuku Hiva); Society (Tahiti).

REFERENCES. — *Panulirus homarus* - GORDON, 1953: 29, fig. 2b-d, 6, 7b (Marquesas "Hana Hevané" = Hanavave bay?, Fatu Hiva; Puerulus larvae only). — MICHEL, 1971: 467 (Marquesas; Phyllosom larvae only). — HOLTHUIS, 1991: 139, fig. 267-268 (Marquesas, with a ?; Syn.). — ? *Panulirus* (sic) *spinosus* (Edwards) - CANO, 1888: 179 (Tahiti) - NEW MATERIAL - February 1996, Coll. & det. J. POUPIN (Nuku Hiva) - SYNONYMS - *Palinurus spinosus* H. Milne Edwards, 1837 (with a ?, in HOLTHUIS, 1991: 139).

Panulirus longipes (A. Milne Edwards, 1868)

DISTRIBUTION. — Marquesas; Society (Tahiti); Tuamotu.

REFERENCES. — *Senex femoristriga* - ORTMANN, 1891: 23 (Tahiti). — *Panulirus longipes* - MICHEL, 1971: 467 (Marquesas, Tuamotu; Phyllosom larvae only). — *Panulirus longipes femoristriga* - HOLTHUIS, 1991: 146, fig. 277b, 278 (Syn.).

REMARK. — In the Indo-West Pacific HOLTHUIS (1991) recognises two subspecies: *Panulirus longipes*, the western form, distributed from Africa to Thailand, Taiwan, Indonesia, and Philippines; and *P. longipes femoristriga*, the eastern form, known from Japan, the Moluccas, New Guinea, New Caledonia, eastern Australia, and French Polynesia.

Panulirus pascuensis Reed, 1954

DISTRIBUTION. — Austral (MacDonald bank, Marotiri, Rapa) - *Littoral to sublittoral*.

REFERENCES. — *Panulirus pascuensis* - LABOUTE & RICHER DE FORGES, 1986: 7, 21, pl. 2c (MacDonald bank, Marotiri, Rapa; 40m). — SALVAT, 1986b: 70, photograph (MacDonald bank). — HOLTHUIS, 1991: 149, fig. 283-284 (Pitcairn, 500km south-east off the Gambier). — POUPIN, 1994a: 8 (after LABOUTE & RICHER DE FORGES).

REMARK. — LABOUTE & RICHER DE FORGES (1986: 18) also record *Panulirus polyphagus* (Herbst, 1793) in French Polynesia ("Iles hautes et atolls"). This species, which is only listed without material examined, is not reported from the area by HOLTHUIS (1991: 152). We therefore consider that this record is not valid.

Panulirus penicillatus (Olivier, 1791)

DISTRIBUTION. — Gambier; Marquesas; Society (Tahiti); Tuamotu (Makatea, Mataiva, Moruroa, Fakarava, Hao, Raroia, Taiaro, Takapoto).

REFERENCES. — *Panulirus penicillatus* - STIMPSON, 1860: 23 [92] (Tahiti). — BATE, 1888: 82, pl. 12-fig. 2 (Tahiti). — NOBILI, 1907: 366 (Hao). — BOONE, 1935: 67, pl. 17 (Tahiti). — SEURAT, 1934: 60 (Gambier, Tuamotu). — HOLTHUIS, 1953: 50 (Raroia). — CHABOIS L. & F., 1954: 89 (Tahiti). — MORRISON, 1954: 16 (Raroia). — MICHEL, 1971: 467 (Marquesas, Tuamotu; Phyllosom larvae). — CHEVALIER *et al.*, 1968: 92, 137 (Moruroa). — MONTEFORTE, 1984: 173, annex 1, tab. a (Makatea, Mataiva, Tahiti, Takapoto). — DELESALLE, 1985: 289, 293 (Mataiva). — SALVAT, 1986b: 70, photograph (French Polynesia). — BAGNIS & CHRISTIAN, 1983: 108 (Tuamotu). — HOLTHUIS, 1991: 151, fig. 285-286

(Tuamotu; Syn.). — PARDON, 1992: 83, photograph (Fakarava). — POUPIN, 1994a: 8, fig. 4 (Taiaro). — *Cancer theresae* Curtiss, 1938 ("Tautira" = Tahiti).

Panulirus versicolor (Latreille, 1804)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — ? *Panulirus fasciatus* - CANO, 1888: 179 (Tahiti; cf. Remark). — *Panulirus ornatus* - BOONE, 1935: 63, pl. 16 (Tahiti). — POUPIN, 1994a: 8 (French Polynesia; after BOONE, and erroneously after NOBILI, 1907 and GRUVEL, 1911) - Not *Palinurus ornatus* (Fabricius, 1798) (cf. Remark). — *Panulirus versicolor* - HOLTHUIS, 1946: 142, pl. 6-j, pl. 9-b, pl. 11-e,f,m (Tahiti); 1991: 156, fig. 293-294 (French Polynesia).

REMARK. — In HOLTHUIS (1991: 152) *Panulirus fasciatus* Fabricius, 1798 is a synonym of *P. polyphagus* (Herbst, 1793). However, CANO's (1888) reference to *P. fasciatus* would rather be *P. versicolor*, often recorded under *P. fasciatus* (HOLTHUIS, 1991: 152). Moreover, we observe that, in his catalogue, HOLTHUIS (1991) does not mention *P. polyphagus* from French Polynesia.

According HOLTHUIS (1946: 140, 142), BOONE's (1935) Tahitian record of *Panulirus ornatus* (Fabricius, 1798) would be erroneous, but it is not certain that his material really belongs to *P. versicolor*.

FAMILY SYNAXIDAE

Palibythus magnificus Davie, 1990

DISTRIBUTION. — Society (Tahiti) - Sublittoral to deep.

REFERENCES. — *Palinurellus wienechi* (sic) - ANONYMOUS, 1979: 6, 8, 11, not *Palinurellus wieneckii* (de Man, 1881) = *Palibythus magnificus* (Tahiti, 70-240m; material corresponding to the photographs examined and considered by DAVIE (1990: 686) as "almost definitely of this species"). — *Palibythus magnificus* Davie, 1990: 686, fig. 1a-b, 3a, c, 4a, 5a (Tahiti; but not Tuamotu). — POUPIN, 1996: in press (Tahiti, Tuamotu?).

REMARK. — The beautiful set of 10 specimens, collected in 1978 in front of Taravao, Tahiti, has disappeared. It remains only the photographs examined by DAVIE (1990).

Palinurellus wieneckii (De Man, 1881)

DISTRIBUTION. — Tuamotu - Sublittoral.

REFERENCES. — *Palinurellus wieneckii* - MICHEL, 1971: 460, fig. 1a-j, tab. 1 (Tuamotu; Puerulus larvae only). — HOLTHUIS, 1991: 170, fig. 315-316 (Tuamotu: larvae and juveniles; 9-27m).

REMARK. — The larvae, once attributed to this species, could in fact belong to *Palibythus magnificus*, afterwards collected in the area (cf. previous species).

FAMILY SCYLLARIDAE

Arctides regalis Holthuis, 1963

DISTRIBUTION. — Tuamotu.

REFERENCES. — *Arctides antipodarum* - MICHEL, 1971: 467 (Tuamotu; Phyllosom larvae) not *A. antipodarum* Holthuis, 1960 = *A. regalis*, with a doubt, *fide* HOLTHUIS (1991: 177). — *Arctides regalis*. — HOLTHUIS, 1991: 177, fig. 331-332 (Tuamotu; Syn.).

Parribacus antarcticus (Lund, 1793)

DISTRIBUTION. — Society (Maupiti, Tahiti); Tuamotu (Anaa, Manihi?, Moruroa, Raroia, Takapoto, Tureia).

REFERENCES. — *Scyllarus antarcticus* - OWEN, 1839: 86 ("Carysfort" = Tureia). — *Parribacus antarcticus* - SEURAT, 1934: 60 (Tuamotu). — HOLTHUIS, 1953: 50 (Raroia) *pro parte fide* HOLTHUIS (1985: 74); 1985: 73, fig. 21, 25a (Anaa,

Maupiti, Tahiti, Takapoto; Syn.). — CHABOIS L. & F., 1954: 89, unnumbered fig. (French Polynesia). — FOREST, 1954b: 345, fig. 26a (Tahiti). — CHEVALIER *et al.*, 1968: 92, 137 (Moruroa). — ? BABLET, 1972: 32, pl. 10 (French Polynesia). — *Parribacus ursus-major* - BOONE, 1935: 54, pl. 13 (Tahiti). — *Cancer barffi* Curtiss, 1938: 164 ("Tautira" = Tahiti). — ? "Tiane" - PARDON, 1992: 83, photograph (Manihi) (det. according to the photograph). — Not *Parribacus antarcticus* - STIMPSON, 1860: 92 [23] (Tahiti). — NOBILI, 1907: 366 (Hao, "Rikitea" = Mangareva). — HOLTHUIS, 1953: 50 (Raroia). *pro parte*. — MORRISON, 1954: 50 (Raroia) - All = *Parribacus holthuisi* Forest, 1954 *fide* HOLTHUIS (1985: 75) - SYNONYMS - *Parribacus ursus-major* (Herbst, 1793).

Parribacus holthuisi Forest, 1954

DISTRIBUTION. — Gambier (Mangareva); Society (Moorea, Tahiti); Tuamotu (Hao, Hikueru, Mataiva, Raroia, Takapoto).

REFERENCES. — *Parribacus antarcticus* - STIMPSON, 1860: 92 [23] (Tahiti). — NOBILI, 1907: 366 (Hao, "Rikitea" = Mangareva). — HOLTHUIS, 1953: 50 (Raroia) *pro parte*. — MORRISON, 1954: 50 (Raroia) - All, not *P. antarcticus* (Lund, 1793) = *P. holthuisi* *fide* HOLTHUIS (1985: 98). — *Parribacus holthuisi* Forest, 1954b: 346, fig. 25, 26b (Hikueru, Tahiti). — MONTEFORTE, 1984: 173, annex 1, tab. a (Mataiva, Moorea, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — HOLTHUIS, 1985: 98 (Hao, Hikueru, Mangareva, Tahiti, Raroia). — SALVAT, 1986b: 70, 71, photograph (French Polynesia).

Parribacus scarlatinus Holthuis, 1960

DISTRIBUTION. — Marquesas (Fatu Hiva).

REFERENCES. — *Parribacus scarlatinus* - MICHEL, 1971: 472 (Marquesas, Omoa bay = Fatu Hiva). — HOLTHUIS, 1985: 102, fig. 26; 1991: 215, fig. 411-412 (Marquesas).

Scyllarus aurora Holthuis, 1981

DISTRIBUTION. — Austral (Maria, Rurutu, Tubuai); Gambier; Marquesas (Fatu Hiva, Tahuata); Society (Maupiti, Moorea, Raiatea, Tupai); Tuamotu (Akiaki, Fangataufa, Hao, Makemo, Marutea South, Maria, Moruroa, Tuanake, Tureia, Vanavana) - *Sublittoral to deep*.

REFERENCES. — ? *Scyllarus* sp. IV & V - MICHEL, 1971: 467, tab. 3 (Marquesas, Tuamotu; larvae only). — *Scyllarus aurora* Holthuis, 1981: 847, fig. 1-2 (Tubuai; 200m). — MANACH & CARSIN, 1985: 473 (Moruroa and/or Fangataufa). — POUPIN, 1996: in press ("Common, 90-300m"; in the distribution, most of the islands are mentioned here for the first time).

INFRA-ORDER ANOMURA

FAMILY COENOBITIDAE

Birgus latro (Linné, 1767)

DISTRIBUTION. — Gambier (Temoe); Tuamotu (Amanu, Makatea, Marutea South, Matureivavao, Morane, Niau, Pukapuka, Raroia, Taiaro, Takapoto) - *Terrestrial*.

REFERENCES. — *Birgus latro* - DANA, 1852b: 474; 1855, pl. 30, fig. 5a-b (several islands in the Tuamotu; drawing of a specimen from "Honden" = Pukapuka). — NOBILI, 1907: 375 (Amanu). — SEURAT, 1904a: 242 (Marutea South, Temoe, "Moture-vavao" = Matureivavao); 1934: 51 (French Polynesia). — SENDLER, 1923: 44 (Makatea). — HOLTHUIS, 1953: 36 (Raroia). — MORRISON, 1954: 10 (Raroia). — FOREST, 1954a: 79; 1956a: 1073 (Niau). — CHABOIS L. & F., 1954: 92, unnumbered fig. (Makatea). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Takapoto). — DELESALLE, 1985: 288 (Mataiva). — CHARLEUX, 1986: 80, photograph (French Polynesia). — SALVAT &

RICHARD, 1985: 356 (Takapoto). — SALVAT, 1986b: 71; 1986c: 8-9, photograph (French Polynesia). — BONVALLOT *et al.*, 1994: 76, photograph (Tuamotu). — POUPIN, 1994a: 10, fig. 6, pl. 1h (Taiaro) - NEW MATERIAL - Coll. and det. J. POUPIN (Morane).

REMARK. — GIBSON-HILL (1948: 10) mentions this species from the Marquesas Islands, but it is doubtful that it really occurs in these Islands, where we have made several unsuccessful investigations.

Coenobita brevimanus Dana, 1852

DISTRIBUTION. — Society (Bora Bora, Tahiti); Tuamotu (Amanu, Hao, Hikueru, Makatea, Mataiva, Niau, Nukutipipi, Raroia, Takapoto, Taiaro) - Terrestrial.

REFERENCES. — *Cenobita clypeata* Latr. - HELLER, 1865: 82 (Tahiti). — SEURAT, 1934: 52 (Amanu, Hao). — *Coenobita clypeatus* (Herbst) - ORTMANN, 1892a: 316, pl. 12, fig 20 (Tahiti) not *C. clypeatus* (Herbst, 1794) = *C. hilgendorfi* Terao in TERAO (1913: 388). — *Coenobita clypeatus* Latr. - NOBILI, 1907: 373 (Amanu, "Ohura" = Hao). — SENDLER, 1923: 42 (Makatea, "Nian" = Niau). — *Coenobita hilgendorfi* Terao, 1913: 388 (Syn.; cf. Remark). — FOREST, 1954a: 77 (Hikueru; Syn.); 1956a: 1072 (Hikueru, Tahiti). — *Cenobita clypeatus* Latreille - BOONE, 1935: 40, pl. 9 (Bora Bora). — *Coenobita brevimanus* - HOLTHUIS, 1953: 36 (Raroia). — MORRISON, 1954: 10 (Raroia). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Takapoto). — SALVAT, 1986b: 72 (French Polynesia). — NAKASONE, 1988: 174 (Syn.). — MERSCHARDT-SALVAT, 1991: 40 (Nukutipipi). — SALVAT F. & B., 1992: 5 (Nukutipipi). — POUPIN, 1994a: 11, fig. 7, pl. 1c (Hikueru, Tahiti, Taiaro). — *Coenobita ollivieri* (sic) - CHARLEUX, 1986: 80-81, photograph (French Polynesia) not *C. olivieri* (Owen, 1839) = *C. brevimanus* (correction according to the photograph).

REMARK. — In TERAO (1913: 389) *Coenobita clypeatus* (Herbst, 1794) is different from *Coenobita clypeatus* (Latreille, 1826), and the name *Coenobita hilgendorfi* is proposed for LATREILLE's material. More recently, NAKASONE (1988) considers that TERAO's (1913) *C. hilgendorfi* is the same than *C. brevimanus* Dana, 1852, and states that, until 1955, DANA's species has been often referred to as, either *C. clypeatus*, or *C. hilgendorfi*.

Coenobita carnescens Dana 1851

DISTRIBUTION. — Tuamotu (Ahe and/or Manihi, Aratika, Kauehi, Pukapuka, Raraka) - Terrestrial.

REFERENCES. — *Cenobita carnescens* Dana, 1851: 272 (Paumotu); 1852b: 472; 1855, pl. 30, fig. 3a-b ("Carlshoff" = Aratika, "Honden" = Pukapuka, Raraka, "Vincennes" = Kauehi, "Waterland" = Ahe and/or Manihi). — POUPIN, 1994a: 9, fig. 5 (Text).

REMARK. — In NAKASONE (1988: 165) this species would be valid, although it was considered doubtful by BOUVIER (in ALCOCK, 1905: 193). According to the drawings provided by DANA, *Coenobita carnescens* could be in fact the juvenile form of *C. perlatus* (cf. POUPIN, 1994a: 12, pl. 1d-f).

Coenobita cavipes Stimpson, 1858

DISTRIBUTION. — Austral (Rimatara) - Terrestrial.

REFERENCES. — *Coenobita cavipes* - SENDLER, 1923: 43 (Rimatara). — MIYAKE, 1991: 116, fig. 3 (cited only for the illustration). — POUPIN, 1994a: 9 (Text).

Coenobita olivieri (Owen, 1839)

DISTRIBUTION. — Gambier (Tarau-Roa); Society (Tahiti) - Terrestrial.

REFERENCES. — *Coenobita olivieri* - NOBILI, 1907: 374 ("Tarawao, Papenoo" = Tahiti, Tarau-Roa; cf. Remark). — SEURAT, 1934: 52 (Tahiti, Gambier). — FOREST, 1956a: 1056 (French Polynesia). — POUPIN, 1994a: 14 (Tahiti; cf. Remark). — Not *Coenobita olivieri* - DANA, 1852b: 470 = *C. spinosus* H. Milne Edwards *fide* ORTMANN (1892a: 318). — BAGNIS & CHRISTIAN, 1983: 108, photograph (Tuamotu) = *C. perlatus* H. Milne Edwards *fide* POUPIN (1994a: 12).

REMARK. — A specimen attributed by NOBILI to *Coenobita olivieri* (Owen, 1839) has been examined previously (POUPIN, 1994a; Tahiti "Papenoo", MNHN Pg2111). It is very close to *Coenobita spinosus*, and ORTMANN's (1892a) opinion that *Coenobita olivieri* is only a variety of *C. spinosus*, could be justified. Examination of the type material is required to clarify that point.

Coenobita perlatus H. Milne Edwards, 1837

DISTRIBUTION. — Gambier (Mangareva, Tarauru-Roa, Temoe); Society (Moorea, Tahiti); Tuamotu (Amanu, Hao, Hikueru, Kaukura, Makatea, Marutea South, Mataiva, Moruroa, Nukutipipi, Raroia, Taiaro, Takapoto, Takume) - *Terrestrial*.

REFERENCES. — *Coenobita perlata* - SEURAT, 1904a: 238 (Mangareva, Tarauru-Roa, Temoe, Marutea South); 1904b: 95 (Marutea South); 1934: 51 (French Polynesia). — CHEVALIER *et al.*, 1968: 85, 137 (Moruroa). — *Coenobita perlatus* - NOBILI, 1907: 373 (Amanu, Hao, Kaukura, Marutea, Tarauru-Roa). — SENDLER, 1923: 43 (Makatea, "Uusuroa, Paumotu" = ?). — HOLTHUIS, 1953: 37 (Raroia). — MORRISON, 1954: 7 (Raroia). — CHABOIS L. & F., 1954: 93 ("Mataia" = Tahiti). — FOREST, 1954a: 78; 1956a: 1072 (Hikueru, Takume). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 6 (Moorea). — DELESALLE, 1985: 288, 289 (Mataiva). — SALVAT & RICHARD, 1985: 359, 360 (Takapoto). — CHARLEUX, 1986: 80-81, photograph (French Polynesia). — SALVAT, 1986b: 71 (French Polynesia). — MERSCHARDT-SALVAT, 1991: 40 (Nukutipipi). — PARDON, 1992: 83, photograph (Tuamotu). — SALVAT F. & B., 1992: 5 (Nukutipipi). — BONVALLOT *et al.*, 1994: 77, photograph (Tuamotu). — POUPIN, 1994a: 12, fig. 8, pl. 1d,f (Hao, Mangareva, Marutea, Taiaro). — *Coenobita rugosus* var. *granulatus* Bouvier - NOBILI 1907: 373 (Marutea, "Ohura" = Hao) not *C. rugosus* H. Milne Edwards, 1837 = *C. perlatus* fide FOREST (1954a: 78). — *Coenobita rugosus* - NOBILI, 1907: 373 (Hao, Kaukura) — SEURAT, 1934: 52 (NOBILI's material) - These two references, *pro parte* not *C. rugosus* H. Milne Edwards, 1837 = *C. perlatus* fide FOREST (1954a: 78). — *Coenobita olivieri* - BAGNIS & CHRISTIAN, 1983: 108, photograph (Tuamotu) not *C. olivieri* (Owen, 1839) = *C. perlatus* fide POUPIN (1994a: 12).

Coenobita rugosus H. Milne Edwards, 1837

DISTRIBUTION. — Austral (Raevavae); Society (Tahiti); Tuamotu (Hikueru, Kaukura, Raraka, Raroia, Takume) - *Terrestrial*.

REFERENCES. — *Cenobita rugosa* - DANA, 1852b: 471; 1855, pl. 30, fig. 1 (Raraka). — STIMPSON, 1858c: 245 [83]; 1907: 199 (Tahiti). — HELLER, 1865: 82 (Tahiti). — HENDERSON, 1888: 51 (Tahiti). — SEURAT, 1934: 52 (Kaukura) — *Coenobita rugosus* - ORTMANN, 1892a: 317, pl. 12, fig. 22 (Tahiti). — NOBILI, 1907: 373 (Kaukura, "Ohura" = Hao) *pro parte* fide FOREST (1954a: 78; some specimens are *C. perlatus*). — SENDLER, 1923: 42 (Tuamotu). — FOREST, 1954a: 78; 1956a: 1073 (Hikueru, Tahiti, Takume). — HOLTHUIS, 1953: 40 (Raroia). — MORRISON, 1954: 10 (Raroia). — POUPIN, 1994a: 13, fig. 9, pl. 1e (Raevavae, Tahiti). — Not *C. rugosus* (*pro parte*) and *C. rugosus* var. *granulosa* Bouvier - NOBILI, 1907: 373 = *C. perlatus* fide FOREST (1954a: 78).

Coenobita spinosus H. Milne Edwards, 1837

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Ahe and/or Manihi, Amanu, Niau, Nukutipipi, Reao) - *Terrestrial*.

REFERENCES. — *Cenobita olivieri* - DANA, 1852b: 470 ("Clermont Tonnerre" = Reao, Tahiti, "Waterland" = Ahe and/or Manihi) not *C. olivieri* (Owen, 1839) = *C. spinosus* fide ORTMANN (1892a: 318). — *Coenobita spinosus* - NOBILI, 1907: 374 (Amanu). — SENDLER, 1923: 43 ("Nian" = Niau). — SEURAT, 1934: 52 (Amanu). — FOREST, 1956a: 1056 (French Polynesia). — MERSCHARDT-SALVAT, 1991: 40 (Nukutipipi). — SALVAT F. & B., 1992: 5 (Nukutipipi). — POUPIN, 1994a: 14, fig. 10, pl. 1g (Amanu, Moorea, Tahiti).

FAMILY DIOGENIDAE

Aniculus aniculus (Fabricius, 1787)

DISTRIBUTION. — Gambier (Mangareva); Marquesas (Nuku Hiva); Society (Moorea, Tahiti); Tuamotu (Ahe and/or Manihi, Aratika, Fakahina, Hikueru, Makatea, Marokau, Marutea South, Mataiva, Moruroa, Nukutipipi, Rangiroa, Raraka, Raroia, Takapoto, Tikehau).

REFERENCES. — *Aniculus typicus* - DANA, 1852b: 461; 1855, pl. 29, fig. 1 ("Carlshoff" = Aratika, Raraka, "Waterland" = Ahe and/or Manihi). — CANO, 1888: 178 (Tahiti). — *Aniculus aniculus* - NOBILI, 1907: 371 (Marokau). — SEURAT, 1934: 61 (Tuamotu, Gambier). — BOONE, 1935: 36, pl. 8 (Tahiti). — HOLTHUIS, 1953: 41 (Raroia). — MORRISON,

1954: 13 (Raroia). — FOREST, 1953b: 561; 1956a: 1067 (Hikueru, Tahiti); 1984: 21, fig. 8, 16, 28-30, 35, 66, 68, 74, 76-85 (Fakahina, Hikueru, Mangareva, Marokau, Marutea South, "Matahiva and Tiahura" = Moorea, Moruroa, Rangiroa, Raroia, Tahiti, "Tawhae, Marquesas" = Taiohae at Nuku Hiva, "Tikahau atoll" = Tikehau; Syn.). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Moorea, Takapoto); 1987: 8 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — RAHAYU, 1988: 40 (French Polynesia). — MERSCHARDT-SALVAT, 1991: 40 (Nukutipipi). — SALVAT F. & B., 1992: 5 (Nukutipipi).

REMARK. — *Aniculus typicus*, proposed by DANA (1852c) in the place of *Pagurus aniculus* Fabricius, is no more a valid name (*cf.* FOREST, 1984: 21).

Aniculus maximus Edmonson, 1952

DISTRIBUTION. — Marquesas (Fatu Hiva, Nuku Hiva).

REFERENCES. — *Aniculus maximus* - FOREST, 1984: 61, fig. 14, 22, 59-61 (Fatu Hiva, Nuku Hiva). — SALVAT, 1986a: 6, 7, photograph (French Polynesia). — ? *Aniculus* sp. - SALVAT, 1986b: 71, photograph (French Polynesia; det. according to the photograph).

Calcinus elegans (H. Milne Edwards, 1836)

DISTRIBUTION. — Gambier (Kamaka, Makaroa, Mangareva); Society (Bora Bora, Tahiti); Tuamotu (Ahe and/or Manihi, Amanu, Apataki, Aratika, Hao, Hikueru, Marutea South, Mataiva, Raroia, Reao, Tagatau, Taiaro, Takume).

REFERENCES. — *Calcinus elegans* - DANA, 1852b: 458; 1855, pl. 28, fig. 10a-c ("Carlshoff" = Aratika, "Clermont Tonnerre" = Reao, "Waterland" = Ahe and/or Manihi). — HELLER, 1865: 88 (Tahiti). — NOBILI, 1907: 368 (Amanu, Hao, Kamaka, Makaroa, Marutea South, "Rikitea" = Mangareva, Tagatau, "Wakatihi" = ?). — BOONE, 1935: 23, pl. 3 (Bora Bora, Tahiti). — HOLTHUIS, 1953: 41 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST, 1953b: 555; 1956a: 1062 (Hikueru, Tahiti, Takume). — MONTEFORTE, 1984: 172, annex 1, tab. a (Mataiva). — RAHAYU, 1988: 10, 17 (French Polynesia). — POUPIN, 1994a: 15, fig. 11, pl. 2a (Hao, Taiaro).

Calcinus gaimardi (H. Milne Edwards, 1848)

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Calcinus gaimardi* - HELLER, 1865: 87 (Tahiti). — FOREST, 1953b: 555; 1956a: 1062 (Tahiti). — KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea). — MONTEFORTE, 1984: 172, annex 1, tab. a; 1987: 8 (Moorea). — RAHAYU, 1988: 20 (Tahiti).

Calcinus guamensis Wooster, 1984

DISTRIBUTION. — Marquesas (Fatu Hiva, Hiva Oa).

REFERENCES. — *Calcinus guamensis* - NEW MATERIAL - February 1996, Coll. J. POUPIN, det. J. POUPIN & J. FOREST (Fatu Hiva, Hiva Oa).

Calcinus imperialis Whitelegge, 1901

DISTRIBUTION. — Society (Tahiti); Tuamotu (Moruroa).

REFERENCES. — *Calcinus imperialis* - NEW MATERIAL - October 1995, Coll. (in coral *Pocillopora*) & det. J. POUPIN (Moruroa, Tahiti "Taravao"). — MORGAN, 1991: 882, fig. 21-23 (Syn.; Key).

Calcinus laevimanus (Randall, 1840)

DISTRIBUTION. — Gambier (Kamaka, Mangareva, Puaumu); Society (Moorea, Tahiti); Tuamotu (Ahe and/or Manihi, Aratika, Hao, Hikueru, Kauehi, Makatea, Marutea South, Mataiva, Raraka, Raroia, Tagatau, Taiaro, Takapoto, Takume).

REFERENCES. — *Calcinus tibicen* - DANA, 1852b: 457 ("Carlshoff" = Aratika, Raraka, "Vincennes" = Kauehi, "Waterland" = Ahe and/or Manihi) not *C. tibicen* (Herbst, 1791) = *C. laevimanus* fide MORGAN (1991: 888). — HELLER, 1865: 87 (Tahiti; *cf.* Remark). — HENDERSON, 1888: 61 (Tahiti; *cf.* Remark). — *Calcinus herbstii* - NOBILI, 1907: 368 *pro parte* fide FOREST (1951: 84) (Hao, Hikueru, Kamaka, Marutea, "Puaumu" = Puaumu, "Rikitea" =

Mangareva, Tagatau, Tahiti). — SENDLER, 1923: 42 (Makatea). — FOREST, 1951: 84 (NOBILI's material); 1953: 555; 1956a: 1062 (Hikueru, Tahiti, Takume). — *Calcinus herbstii* var. *lividus* Edw. - NOBILI, 1907: 369 (Marutea; cf. Remark). — *Calcinus laevimanus* - HOLTHUIS, 1953: 43 (Raroia). — MORRISON, 1954: 7 (Raroia). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Moorea, Takapoto); 1987: 8 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — RAHAYU, 1988: 10, 18, fig. 1-3 (French Polynesia). — POUPIN, 1994a: 16, fig. 12 (Hikueru, Taiaro) - SYNONYMS - *Calcinus herbstii* de Man, 1888; *Pagurus lividus* H. Milne Edwards, 1848.

REMARK. — *Calcinus tibicen* (Herbst, 1791) is an Atlantic species. MORGAN (1991: 888) has cited several works in which *C. laevimanus* is referred to as *C. tibicen*, *C. herbstii*, or *C. herbstii* var. *lividus*, but without the works by HELLER, HENDERSON or NOBILI. Although not verified, this material is here attributed to *C. laevimanus*.

Calcinus latens (Randall, 1840)

DISTRIBUTION. — Gambier (Vaiatekeue); Society (Moorea, Tahiti); Tuamotu (Fakahina, Hao, Hikueru, Makatea, Mataiva, Raroia, Taiaro, Takapoto, Takume).

REFERENCES. — *Calcinus latens* - HELLER, 1865: 88 (Tahiti). — NOBILI, 1907: 369 (Hao). — FOREST, 1951: 84, fig. 14-18 (French Polynesia); 1953b: 556 (Syn.); 1956a: 1062 (Hikueru, Tahiti, Takume). — HOLTHUIS, 1953: 44 (Raroia). — MORRISON, 1954: 7 (Raroia). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — KROPP & BIRKELAND, 1981: 630, tab. 5 (Takapoto). — MONTEFORTE, 1984: 172, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 8 (Moorea). — GALZIN & POINTIER, 1985: 100 (Moorea). — SALVAT & RICHARD, 1985: 358 (Takapoto). — RAHAYU, 1988: 10, 19 (French Polynesia). — POUPIN, 1994a: 17, fig. 13, pl. 2b (Takume, Taiaro). — *Calcinus herbstii* - NOBILI, 1907: 368 (Hao, "canal Waiatekene" = Vaiatekeue) pro parte not *C. herbstii* de Man, synonym of *C. laevimanus* = *C. latens* fide FOREST (1951: 84). — *Calcinus terrae-reginae* - NOBILI, 1907: 369 (Fakahina, Hao, Mangareva) - SYNONYMS - *Calcinus terrae-reginae* Haswell, 1882.

Calcinus minutus Buitendijk, 1937

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Calcinus minutus* - MONTEFORTE, 1984: 172, annex 1, tab. a (Moorea).

Calcinus nitidus Heller, 1865

DISTRIBUTION. — Society (Tahiti); Tuamotu (Takapoto).

REFERENCES. — *Calcinus nitidus* Heller, 1865: 89, pl. 7, fig. 4 (Tahiti). — DE MAN, 1890: 111 (Tahiti). — ORTMANN, 1892a: 293 (Tahiti). — FOREST, 1956b: 218, fig. 1-4 (Tahiti). — RAHAYU, 1988: 10 (French Polynesia). — POUPIN, 1994a: 18, fig. 14, pl. 2c (Tahiti, Takapoto).

Calcinus seurati Forest, 1951

DISTRIBUTION. — Gambier (Tarauru-Roa, Vaiatekeue); Society (Tahiti); Tuamotu (Hao, Hikueru, Fakahina, Raroia, Taiaro, Takume).

REFERENCES. — *Calcinus herbstii* - NOBILI, 1907: 368 (Hao, Fakahina, "Taraourouroa" = Tarauru-Roa, "chenal Waiatekene" = Vaiatekeue) pro parte not *C. herbstii* de Man, 1888, synonym of *C. laevimanus* = *C. seurati* fide FOREST (1951: 86). — *Calcinus seurati* Forest, 1951: 84, fig. 1, 3-4, 7-8 (NOBILI's material); 1953b: 556; 1956a: 1062 (Hikueru, Tahiti, Takume). — HOLTHUIS, 1953: 44 (Raroia, Takume). — MORRISON, 1954: 7 (Raroia). — RAHAYU, 1988: 19 (Hikueru, Tahiti, Takume). — POUPIN, 1994a: 19, fig. 15, pl. 2d (Hao, Taiaro).

Calcinus spicatus Forest, 1951

DISTRIBUTION. — Gambier (Vaiatekeue).

REFERENCES. — *Calcinus herbstii* - NOBILI, 1907: 368 ("chenal Waiatekene" = Vaiatekeue) pro parte not *C. herbstii* de Man, 1888, synonym of *C. laevimanus* = *C. spicatus* fide FOREST (1951: 84). — *Calcinus spicatus* Forest, 1951: 90, fig. 10-13 ("chenal Waiatekene" = Vaiatekeue). — RAHAYU, 1988: 21 (French Polynesia).

Ciliopagurus krempfi (Forest, 1952)

DISTRIBUTION. — Marquesas (Tahuata) - *Sublittoral*.

REFERENCES. — *Ciliopagurus krempfi* - FOREST, 1995: 59 fig. 10c, 11, 12c, 31d, 37g-h (Tahuata; 48m).

Ciliopagurus strigatus (Herbst, 1804)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Moorea, Tahiti).

REFERENCES. — *Pagurus strigatus* - ORTMANN, 1892a: 285 (Tahiti). — *Trizopagurus strigatus* - MONTEFORTE, 1984: 172, annex 1, tab. a (Moorea, Tahiti); 1987: 8 (Moorea). — *Ciliopagurus strigatus* - FOREST, 1995: 49, fig. 8a, 9, 10a, 12a, 31a-b, 37d (Moorea, Nuku Hiva, Tahiti).

Clibanarius corallinus (H. Milne Edwards, 1848)

DISTRIBUTION. — Society (Bora Bora, Tahiti); Tuamotu (Apataki, Hao, Hikueru, Mataiva, Moruroa, Raroia, Taiaro, Takapoto, Takume).

REFERENCES. — *Clibanarius corallinus* - HELLER, 1865: 89 (Tahiti; cf. Remark). — NOBILI, 1907: 367 (Apataki, Hao). — BOONE, 1935: 17, pl. 1 (Bora Bora). — HOLTHUIS, 1953: 45 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST, 1953a: 442; 1956a: 1057 (Hikueru, Tahiti, Takume). — MONTEFORTE, 1984: 172, annex 1, tab. a (Mataiva, Takapoto). — RAHAYU, 1988: 26, fig. 4-6 (Tahiti). — POUPIN, 1994a: 20, fig. 16 (Apataki, Moruroa, Taiaro).

REMARK. — HELLER (1862: 527) has also described a new *Clibanarius* from Tahiti, *Clibanarius semistriatus*. According to J. FOREST (Personal communication) this species is very doubtful and could be a *Pagurus* or a *Paguristes*.

Clibanarius eurysternus Hilgendorf, 1878

DISTRIBUTION. — French Polynesia.

REFERENCES. — *Clibanarius eurysternus* - RAHAYU, 1988: 10, 28 (French Polynesia). — RAHAYU & FOREST, 1992: 750 (Distribution only "Polynésie").

Clibanarius humilis (Dana, 1851)

DISTRIBUTION. — Gambier (Mangareva, Tarauru-Roa); Society (Tahiti); Tuamotu (Hikueru, Mataiva, Moruroa, Takume).

REFERENCES. — *Clibanarius aequabilis* - ? DANA, 1852b: 464; 1855, pl. 29, fig. 4a-f (Tahiti; cf. Remark). — NOBILI, 1907: 367 ("Rikitea" = Mangareva, Tarauru-Roa) not *C. aequabilis* Dana, 1852 = *C. humilis* fide FOREST (1953a: 443). — *Clibanarius humilis* - FOREST, 1953a: 443, fig. 1, 5; 1956a: 1057 (Hikueru, Tahiti, Takume). — MONTEFORTE, 1984: 172, annex 1, tab. a (Mataiva). — RAHAYU, 1988: 10, 27 (French Polynesia). — POUPIN, 1994a: 21, fig. 17 (Moruroa, Tahiti).

REMARK. — *Clibanarius aequabilis* Dana, 1852 is an Atlantic species. Concerning DANA's material, FOREST (1953a: 446) writes: "A quoi correspond le sp. *C. aequabilis* de Tahiti, figuré par Dana ? Il est souhaitable que le terme d'*aequabilis* ne soit plus utilisé pour les espèces de l'Indo-Pacifique."

Clibanarius ransoni Forest, 1953

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Clibanarius ransoni* Forest, 1953a: 446, fig. 2, 6; 1956a: 1059 (Tahiti). — RAHAYU, 1988: 30 (Tahiti).

Clibanarius rhabdodactylus Forest, 1953

DISTRIBUTION. — Tuamotu (Hao, Hikueru).

REFERENCES. — *Clibanarius zebra* - NOBILI, 1907: 367 (Hao), not *C. zebra* Dana, 1852 = *C. rhabdodactylus* fide RAHAYU & FOREST (1992: 777). — *Clibanarius zebra* var. *rhabdodactylus* Forest, 1953a: 448, fig. 3, 8; 1956a: 1059 (Hikueru). — RAHAYU, 1988: 29 (French Polynesia). — *Clibanarius rhabdodactylus* - RAHAYU & FOREST, 1992: 777 (Syn.).

Clibanarius striolatus Dana, 1852

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Clibanarius striolatus* - HELLER, 1865: 89 (Tahiti). — FOREST, 1953a: 448; 1956a: 1059 (Tahiti). — RAHAYU, 1988: 10, 27 (Tahiti).

Clibanarius zebra (Dana, 1852)

DISTRIBUTION. — Marquesas (Fatu Hiva, Nuku Hiva, Ua Pou).

REFERENCES. — *Clibanarius zebra* - FOREST, 1953a: 449, fig. 4 ("Taiohae" = Nuku Hiva) - NEW MATERIAL - February 1996, Coll. J. POUPIN, det. J. POUPIN & J. FOREST (Fatu Hiva, Nuku Hiva, Ua Pou).

REMARK. — FOREST (1953a) states that the specimens from Marquesas (Coll. Père Simon DELMAS) are typical of *C. zebra*, and distinctly larger than the *rhabdodactylus* variety.

Dardanus australis Forest & Morgan, 1991

DISTRIBUTION. — Austral (Rapa) - Sublittoral to deep.

REFERENCES. — *Dardanus australis* - NEW MATERIAL - March 1995, Coll. J. POUPIN, det. J. FOREST (Rapa, 70-115m).

Dardanus brachyops Forest, 1962

DISTRIBUTION. — Marquesas (Tahuata); Society (Bora Bora, Maupiti) - Sublittoral to deep.

REFERENCES. — *Dardanus brachyops* - POUPIN, 1996: in press (Bora Bora, Maupiti, Tahuata; 110-300m cf. Remark).

REMARK. — This species is still known only beyond 100m in French Polynesia, but it is merely sublittoral in FOREST (1962: 365; more than 33m to 80m).

Dardanus deformis (H. Milne Edwards, 1836)

DISTRIBUTION. — Gambier (Mangareva?); Society (Tahiti); Tuamotu (Hao, Hikueru, Matureivavao).

REFERENCES. — *Pagurus difformis* - HELLER, 1865: 86 (Tahiti). — HENDERSON, 1888: 57 (Tahiti). — *Pagurus deformis* - NOBILI, 1907: 370 (Hao, and Mangareva with a ?). — BOONE, 1935: 28, pl. 5 (Tahiti). — FOREST, 1953b: 556; 1956a: 1063 (Hikueru, Tahiti). — *Dardanus deformis* - HOLTHUIS, 1953: 47 (Raroia, Tahiti). — MORRISON, 1954: 7 (Raroia). — RAHAYU, 1988: 35 (French Polynesia) - NEW MATERIAL - Coll. PLESSIS (Matureivavao), coll. C. HILY (Tahiti), det. J. POUPIN.

Dardanus gemmatus (H. Milne Edwards, 1848)

DISTRIBUTION. — Austral (Maria); Marquesas (Ua Huka); Society (Moorea, Tahiti); Tuamotu (Manihi?, Taiaro) - Littoral to sublittoral (10-20m).

REFERENCES. — *Pagurus gemmatus* H. Milne Edwards, 1848: 60 (Marquesas). — STIMPSON, 1858c: 234 [72] (Marquesas; new material ?). — FOREST, 1953b: 557, fig. 10-11; 1956a: 1063 (Tahiti). — *Dardanus gemmatus* - MONTEFORTE, 1984: 172, annex 1, tab. a; 1987: 8 (Moorea). — RAHAYU, 1988: 36 (Tahiti). — POUPIN, 1994a: 22, fig. 18, pl. 2e (Maria, Tahiti, Taiaro, Ua Huka; 10-20m). — Without name - ? SALVAT, 1986a: 23, photograph (Tahiti). — ? PARDON, 1992: 80, photograph (Manihi) - Det. after the photographs.

Dardanus guttatus (Olivier, 1812)

DISTRIBUTION. — Tuamotu (Tureia).

REFERENCES. — *Pagurus guttatus* - OWEN, 1839: 82 ("Carysfort" = Tureia). — *Dardanus guttatus* - HAIG & BALL, 1988: 165 (Syn.).

Dardanus lagopodes (Forskål, 1775)

DISTRIBUTION. — Society (Maupiti?, Moorea, Tahiti, Tupai); Tuamotu (Hao, Hikueru, Moruroa).

REFERENCES. — *Pagurus euopsis* - NOBILI, 1907: 370 (Hao). — *Pagurus sanguinolentus* - FOREST, 1953b: 559, fig. 12-14 (Syn.); 1956a: 1064 (Hikueru, Tahiti). — *Dardanus lagopodes* - LEWINSOHN, 1969: 32 (Syn.). — MONTEFORTE, 1984: 172, annex 1, tab. a; 1987: 8 (Moorea). — RAHAYU, 1988: 32 (Tahiti, Tuamotu). — "Bernard l'hermite bigaré" - ? PARDON, 1992: 20, 21, double page photograph (Maupiti; det. according to the photograph) - NEW MATERIAL - Coll. and det. J. POUPIN (Moruroa, Tahiti, Tupai) - RELEVANT MATERIAL - *Dardanus* aff. *sanguinolentus* - SALVAT & RENAUD-MORNANT, 1969: 165 (Moruroa) - SYNONYMS - *Pagurus euopsis* Dana, 1852; *P. sanguinolentus* Quoy & Gaimard, 1824.

Dardanus megistos (Herbst, 1804)

DISTRIBUTION. — Society (Tahiti); Tuamotu (Hao, Raroia, Tagatau, Taiaro).

REFERENCES. — *Pagurus spinimanus* Edw. - DANA, 1852b: 452 (with a ?); 1855, pl. 28, fig. 5a-c (Tuamotu). — BOONE, 1935: 34, pl. 7 (Tahiti). — *Pagurus punctulatus* Olivier - HELLER, 1865: 87 (Tahiti). — NOBILI, 1907: 370 ("Ohura" = Hao, Tagatau). — *Pagurus megistos* - FOREST, 1953b: 559; 1956a: 1064 (Tahiti). — *Dardanus megistos* - HOLTHUIS, 1953: 49 (Raroia). — MORRISON, 1954: 7 (Raroia). — RAHAYU, 1988: 33 (French Polynesia). — POUPIN, 1994a: 23, fig. 19, pl. 2g (Tahiti, Taiaro). — Without name - ? SALVAT & RIVES, 1975: 57, full page photograph (French Polynesia; det. according to the photograph) - SYNONYMS - *Pagurus punctulatus* Olivier, 1811; *P. spinimanus* H. Milne Edwards, 1848.

Dardanus pedunculatus (Herbst, 1804)

DISTRIBUTION. — Austral (Tubuai); Tuamotu (Moruroa) - *Littoral to sublittoral*.

REFERENCES. — *Dardanus haani* - CHEVALIER et al., 1968: 119 (Moruroa). — SALVAT & RENAUD-MORNANT, 1969: 165, 176 (Moruroa). — *Dardanus pedunculatus* - RAHAYU, 1988: 10, 34 (French Polynesia). — POUPIN, 1994a: 24, fig. 20, pl. 2f (Tubuai; 65m) - SYNONYMS - *Dardanus haani* (Rathbun, 1902).

Dardanus scutellatus (H. Milne Edwards, 1848)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Pagurus scutellatus* - FOREST, 1953b: 560; 1956a: 1066 (Tahiti). — *Dardanus scutellatus* - RAHAYU, 1988: 36 (Tahiti).

Diogenes gardineri Alcock, 1905

DISTRIBUTION. — Gambier (Mangareva); Tuamotu (Marutea South).

REFERENCES. — *Diogenes gardineri* - NOBILI, 1907: 366 (Marutea South; "Rikitea" = Mangareva). — SEURAT, 1934: 61 (Mangareva, Marutea South). — FOREST, 1956a: 1056 (French Polynesia); 1957: 530 (NOBILI's material).

FAMILY PAGURIDAE

Catapaguroides fragilis (Melin, 1939)

DISTRIBUTION. — Tuamotu (Moruroa) - *Littoral to sublittoral*.

REFERENCES. — *Catapaguroides fragilis* - DE SAINT LAURENT, 1968: 940, fig. 26 with hesitations (cf. p. 941, note 1) (Moruroa; 40m). — CHEVALIER et al., 1968: 119, 138 (Moruroa).

Micropagurus polynesiensis (Nobili, 1906)

DISTRIBUTION. — Gambier (Mangareva); Tuamotu (Moruroa).

REFERENCES. — *Anapagurus polynesiensis* Nobili, 1906a: 260; 1907: 372, pl. 1, fig. 10. ("Rikitea" = Mangareva). — FOREST, 1956a: 1056 (French Polynesia). — CHEVALIER et al., 1968: 119 (Moruroa). — *Micropagurus polynesiensis* - HAIG & BALL, 1988: 184 (Syn.).

***Pagurixus anceps* (Forest, 1954)**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Hikueru).

REFERENCES. — *Eupagurus anceps* Forest, 1954a: 71, fig. 15-19; 1956a: 1067 (Hikueru, Tahiti). — ELDREDGE, 1967: 13 (Hikueru). — *Pagurixus anceps* - MC LAUGHLIN & HAIG, 1984: 135, fig. 5 (Hikueru; Syn.).

***Pagurixus laevimanus* (Ortmann, 1892)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Eupagurus laevimanus* Ortmann, 1892a: 302, pl. 12, fig. 13 (Tahiti). — *Pagurixus laevimanus* - MC LAUGHLIN & HAIG, 1984: 142, fig. 7 (Syn.).

***Pagurixus maorus* (Nobili, 1906)**

DISTRIBUTION. — Gambier (Mangareva).

REFERENCES. — *Eupagurus maorus* Nobili, 1906a: 259; 1907: 371, pl. 1, fig. 9 (Mangareva). — FOREST, 1954a: 73; 1956a: 1056 (French Polynesia). — *Pagurixus maorus* - MC LAUGHLIN & HAIG, 1984: 126, fig. 2 (Syn.). — KOMAI & ASAKURA, 1995: 341, 353 (Key).

***Trichopagurus trichophthalmus* (Forest, 1954)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Catapaguroides trichophthalmus* Forest, 1954a: 74, fig. 20-24, with a ? for the genus; 1956a: 1069 (Tahiti). — *Trichopagurus trichophthalmus* - DE SAINT LAURENT, 1970: 212, fig. 1-16 (Tahiti).

FAMILY GALATHEIDAE

***Coralliogalathea humilis* (Nobili, 1905)**

DISTRIBUTION. — Society (Moorea); Tuamotu (Hao, Tikehau).

REFERENCES. — *Galathea megalochira* Nobili, 1906a: 260; 1907: 376, pl. 1, fig. 12 (Hao). — *Coralliogalathea humilis* - LEWINSOHN, 1969: 117 (Syn.). — PEYROT-CLAUSADE, 1977a, annex of the species: 24 (Moorea); 1989: 113, 115 (Moorea, Tikehau). — KROPP & BIRKELAND, 1981: 629, tab. 5 (Moorea). — ? *Galathea himilis* (sic) - PEYROT-CLAUSADE, 1977b: 213 (Moorea).

***Galathea aculeata* Haswell, 1882**

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Takapoto).

REFERENCES. — *Galathea aculeata* - ODINETZ, 1983: 208 (Moorea, Tahiti, Takapoto). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti, Takapoto). — MONTEFORTE, 1987: 8 (Moorea).

***Galathea aff. amamiensis* Miyake & Baba, 1966**

DISTRIBUTION. — Society (Moorea) - Littoral to sublittoral.

REFERENCES. — *Galathea aff. amamiensis* - PEYROT-CLAUSADE, 1989: 115 (Moorea; 30m).

***Galathea affinis* Ortmann, 1892**

DISTRIBUTION. — Gambier (Mangareva); Society (Moorea); Tuamotu (Hao, Marutea South, Tikehau) - Littoral to sublittoral.

REFERENCES. — *Galathea affinis* - NOBILI, 1907: 375, pl. 1, fig. 11 (Marutea South, "Ohura" = Hao, "Rikitea and Gatavake" = Mangareva). — PEYROT-CLAUSADE, 1977a, annex of the species: 24; 1977b: 213; 1985: 462 (Moorea); 1989: 112, 115 (Moorea, Tikehau; 30m). — KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea).

***Galathea algae* Baba, 1969**

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau) - *Littoral to sublittoral*.

REFERENCES. — ***Galathea algae*** - PEYROT-CLAUSADE, 1989: 112, 115 (Moorea, Tikehau; 30m).

***Galathea latirostris* Dana, 1852**

DISTRIBUTION. — Society (Raiatea, Tahiti).

REFERENCES. — ***Galathea latirostris*** - BOONE, 1935: 50, pl. 12 (Raiatea, Tahiti).

REMARK. — The status of this species, originally described by DANA (1852b: 480; 1855, pl. 30, fig. 8) from the Fiji is doubtful. The examination of topotypic material, and the revision of all the references attributed to *Galathea latirostris* is necessary to establish its real identity (K. BABA, personal communication). *Galathea latirostris* Lenz, 1902, describes form Juan Fernandez must be attributed to *Phylladiorhynchus pusillus* (Henderson, 1885) (*cf.* BABA, 1991: 487), and must not be confounded with DANA's species.

***Phylladiorhynchus serrirostris* (Melin, 1939)**

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau) - *Littoral to sublittoral*.

REFERENCES. — ***Galathea serrirostris*** - PEYROT-CLAUSADE, 1977b: 213 (Moorea). — ***Phylladiorhynchus serrirostris*** - PEYROT-CLAUSADE, 1977a, annex of the species: 24 (Moorea); 1989: 112, 115 (Moorea, Tikehau; "...extremely abundant at 30m"). — KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea).

***Sadayoshia miyakei* Baba, 1969**

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau) - *Littoral to sublittoral*.

REFERENCES. — ***Sadayoshia miyakei*** - KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea). — PEYROT-CLAUSADE, 1989: 112, 115 (Moorea, Tikehau; 30m).

FAMILY PORCELLANIDAE

***Neopetrolisthes maculatus* (H. Milne Edwards, 1837)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — ***Neopetrolisthes oshimai* (sic)** - PARDON, 1992: 81 (Tahiti). — ***Neopetrolisthes maculatus*** - HAIG, 1979: 127 (Syn.).

REMARK. — According to HAIG (1979), *Neopetrolisthes ohshima* Miyake, 1937 is a synonym of this species. This small crustacea is commensal of a sea-anemone. PARDON (1992) illustrates a shrimp *Stenopus* captured in a sea-anemone. "*Neopetrolisthes oshimai*" is only indicated in the caption, but is not visible on the photograph.

***Pachycheles pisoides* (Heller, 1865)**

DISTRIBUTION. — Society; Tuamotu (Tikehau).

REFERENCES. — ***Pachycheles pisoides*** - HAIG, 1966: 290 (Tuamotu; with the mention that it is the first record for the area, but without details on the material examined); 1983: 284 (Distribution only "Society Islands"). — PEYROT-CLAUSADE, 1989: 113 (Tikehau).

***Pachycheles sculptus* (H. Milne Edwards, 1837)**

DISTRIBUTION. — Society (Moorea); Tuamotu.

REFERENCES. — ***Pachycheles sculptus*** - HAIG, 1966: 287 (Tuamotu; same remark than for the previous species); 1983: 284 (Distribution only "Tuamotu"); 1992: 310 (Syn.). — PEYROT-CLAUSADE, 1989: 115 (Moorea).

***Petrolisthes bispinosus* Borradaile, 1900**

DISTRIBUTION. — French Polynesia.

REFERENCES. — *Petrolisthes bispinosus* - HAIG & KROPP, 1987: 171, 172, fig. 1-2 (French Polynesia, only in summary and discussion).

***Petrolisthes borradailei* Kropp, 1983**

DISTRIBUTION. — Gambier (Mangareva); Marquesas (Nuku Hiva); Society (Huahine, Moorea, Tahiti); Tuamotu (Fakarava, Makemo, Moruroa, Rangiroa, Raroia, Taiaro, Tikehau).

REFERENCES. — *Petrolisthes rufescens* - NOBILI, 1907: 377 ("Rikitea" = Mangareva; cf. Remark under *P. rufescens*). — POUPIN, 1994a: 25, fig. 21, pl. 2h (Mangareva, Moruroa, Taiaro) not *P. rufescens* = *P. borradailei* fide KROPP personal communication. — *Petrolisthes borradailei* Kropp, 1983: 96, 106, fig. 3 ("Fakarava" = Fakarava, Huahine, "Makeno" = Makemo, Moorea, Nuku Hiva, Rangiroa, Raroia, "Pascua Pass" = ?, Tahiti, "Tikahua" = Tikehau).

***Petrolisthes coccineus* (Owen, 1839)**

DISTRIBUTION. — Society (Moorea); Tuamotu.

REFERENCES. — *Porcellana coccinea* Owen, 1839: 87, pl. 26, fig. 1-2 ("Low Islands" = Tuamotu). — *Petrolisthes coccineus* - PEYROT-CLAUSADE, 1977a, annex of the species: 24; 1977b: 213 (Moorea). — HAIG, 1983: 280; 1992: 313, fig. 9 (Syn.).

***Petrolisthes decacanthus* Ortmann, 1897**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Petrolisthes decacanthus* Ortmann, 1897a: 285, pl. 17, fig. 2 (Tahiti). — HAIG & KROPP, 1987: 176 (French Polynesia, in the distribution only; Syn.).

***Petrolisthes eldredgei* Haig & Kropp, 1987**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Raroia).

REFERENCES. — *Petrolisthes eldredgei* Haig & Kropp, 1987: 180, fig. 5-6 (Tahiti, Raroia).

***Petrolisthes elegans* Haig, 1981**

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Fakarava).

REFERENCES. — *Petrolisthes bosci* - NOBILI, 1907: 377 ("Sakarava" = Fakarava) not *P. bosci* (Audouin, 1826) = *P. elegans* sp. nov. in HAIG (1981: 266). — *Petrolisthes elegans* Haig, 1981: 266, fig. 2 (Tahiti, "Sakarava" = Fakarava). — PEYROT-CLAUSADE, 1989: 115 (Moorea).

***Petrolisthes lamarckii* (Leach, 1820)**

DISTRIBUTION. — Tuamotu (Raraka, Takaroa).

REFERENCES. — *Porcellana speciosa* Dana, 1852b: 417; 1855, pl. 26, fig. 8 (Raraka). — EVANS, 1967: 409 (Raraka; syntypes at the BM). — *Petrolisthes lamarckii* - HAIG, 1964: 362 (Takaroa); 1992: 315, fig. 11 (Syn.). — KROPP, 1983: 100, 106 (Syn.).

***Petrolisthes militaris* (Heller, 1862)**

DISTRIBUTION. — Tuamotu (Marutea South).

REFERENCES. — *Petrolisthes militaris* - NOBILI, 1907: 377, with a ? (Marutea South).

REMARK. — The presence of this species in French Polynesia is doubtful. NOBILI's determination is uncertain and, moreover, it has never been reported in the area by HAIG (1979: 122; 1982: 280; 1992: 316).

***Petrolisthes pubescens* Stimpson, 1858**

DISTRIBUTION. — Marquesas (Nuku Hiva).

REFERENCES. — *Petrolisthes pubescens* - KROPP, 1986: 456, fig. 2 ("Taiohae, Haka Paa, baie du Controleur" = Nuku Hiva).

***Petrolisthes rufescens* (Heller, 1861)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Porcellana rufescens* - HELLER, 1865: 76 (Tahiti). — Not *Petrolisthes rufescens* - NOBILI, 1907: 377 (cf. Remark). — POUPIN, 1994a: 25, fig. 21, pl. 2h = *P. borradalei* Kropp, 1983 fide KROPP personal communication.

REMARK. — *Petrolisthes rufescens* and *P. borradalei* are two very close species. The main difference concerned the posterior border of the cheliped carpus, strongly toothed in *P. borradalei*, more smooth in *P. rufescens*. We have confused these two species in a previous work (POUPIN, 1994a) and re-examination of NOBILI's material in Paris (♀ ov. 8x7.3, MNHN Ga96) show that it also belongs to *P. borradalei*. HELLER's reference remains the only record of *P. rufescens* in the pacific and could also belongs to *P. borradalei*.

***Petrolisthes scabriculus* (Dana, 1852)**

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Petrolisthes scabriculus* - PEYROT-CLAUSADE, 1977a, annex of the species: 25; 1977b: 213 (Moorea); 1989: 112, 115 (Moorea, Tikehau; 30m). — HAIG, 1979: 120 (Syn.).

***Petrolisthes tomentosus* (Dana, 1852)**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Raraka).

REFERENCES. — *Porcellana tomentosa* Dana, 1852b: 420; 1855, pl. 26, fig. 10 (Raraka). — *Petrolisthes tomentosus* - KROPP, 1986: 453, fig. 1 (Tahiti, Tahitian neotype at the USNM; Syn.).

***Porcellana mitra* Dana, 1852**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Porcellana mitra* - HELLER, 1865: 74, 265, ("Südsee" and Tahiti; cf. Remark).

REMARK. — In HELLER, Tahiti is not mentioned in the main text (p. 74), but only at the end of the work, in the part concerning the geographical distribution (p. 265). It is thus not certain that the location "Tahiti" is correct for this species.

***Porcellana monilifera* Dana, 1852**

DISTRIBUTION. — Tuamotu (Raraka?).

REFERENCES. — *Porcellana monilifera* Dana, 1852b: 413; 1855, pl. 26, fig. 3 (Raraka, with a ?).

FAMILY ALBUNEIDAE

***Albunea speciosa* Dana, 1852**

DISTRIBUTION. — Society (Moorea?).

REFERENCES. — *Albunea speciosa* - NEW MATERIAL - CRIODE collections, Moorea (without label), 1 ♂ 11x10.5, det. J. POUPIN.

REMARK. — This specimen agrees very well with DANA's species, illustrated by SERÈNE (1973). Formerly, *Albunea speciosa* was considered as endemic from Hawaii. SERÈNE (1973) has examined specimens from the type locality but mentions that the type material has disappeared. In 1973, THOMASSIN, has described *A. madagascariensis*, very close to

A. speciosa. By the shape of the ocular peduncle and the number of frontal spines, the specimen from Moorea do belongs to DANA's species.

FAMILY HIPPIDAE

Hippa adactyla Fabricius, 1787

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Tahiti) - *Brackish water*.

REFERENCES. — *Remipes testudinarius* - ? HELLER, 1865: 72 (Tahiti; reference not found in HAIG, 1970). — DE MAN, 1896: 466 ("Nuka-hiwa (Mus. Paris)" = Nuku Hiva). — *Hippa adactyla* - HOLTHUIS, 1953: 35 (Tahiti). — HAIG, 1970: 294 (Syn.); 1974: 179, fig. 2, pl. 6 (Marquesas, distribution only) - NEW MATERIAL - 1 ♂ 28x23.5, coll. C. HILY, det. J. POUPIN ("Arue" = Tahiti; confronted with DE MAN's material) - SYNONYMS - *Remipes testudinarius* Latreille, 1806.

Hippa ovalis (A. Milne Edwards, 1863)

DISTRIBUTION. — Society (Tahiti) - *Brackish water*.

REFERENCES. — *Hippa ovalis* - HOLTHUIS, 1953: 35 (Tahiti). — THOMASSIN, 1969: 154, fig. 7b, 8b, 9, pl. 6, fig. 1-8 (Syn.).

Hippa pacifica (Dana, 1852)

DISTRIBUTION. — Gambier (Mangareva); Society (Tahiti) - *Brackish water*.

REFERENCES. — *Remipes pacificus* - NOBILI, 1907: 378 (Mangareva). — SEURAT, 1934: 60 (Tahiti). — ? *Hippa* sp. - CHABOIS L. & F., 1954: 92, unnumbered fig. (French Polynesia; det. according to the shape of the front margin, on the figure). — *Hippa pacifica* - HAIG, 1974: 181, fig. 3 (Gambier, distribution only; Syn.).

REMARK. — NOBILI's material has been re-examined (MNHN Hi38, Mangareva, 1 ♀ ov. 22x17, coll. SEURAT 1905, det. NOBILI 1906) and is similar to other specimens attributed by DE MAN to this species (MNHN Hi107, Java sea; MNHN Hi108, Atjeh).

INFRA-ORDER BRACHYURA

FAMILY DROMIIDAE

Cryptodromia coronata Stimpson, 1859

DISTRIBUTION. — Gambier (Mangareva).

REFERENCES. — *Cryptodromia coronata* - NOBILI, 1907: 378 ("Rikitea" = Mangareva). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 448 (List). — *Cryptodromia* ? *coronata* - McLAY, 1993: 199 (Syn.; with the indication that the status of the species is uncertain).

Cryptodromia fallax (Lamarck, 1818)

DISTRIBUTION. — Tuamotu (Raroia).

REFERENCES. — *Cryptodromia canaliculata* - HOLTHUIS, 1953: 3 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 448 (List). — *Cryptodromia fallax* - McLAY, 1993: 206, fig. 18e (Syn.) - SYNONYMS - *Cryptodromia canaliculata* Stimpson, 1858.

Cryptodromiopsis tridens Borradaile, 1903

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Cryptodromiopsis tridens* - McLAY, 1991: 467, fig. 5a-d (Moorea, Tahiti).

Dromia wilsoni (Fulton & Grant, 1902)

DISTRIBUTION. — Austral (Raevavae); Marquesas (Tahuata); Tuamotu (Makemo, Takapoto) - Sublittoral to deep.

REFERENCES. — *Petalomera wilsoni* - McLAY, 1991: 470, fig. 6a-d, 7a-c, 8a-c (Makemo, Raevavae, Tahuata, Takapoto; 190-350m). — *Dromia wilsoni* - McLAY, 1993: 156, fig. 16e (Syn.). — POUPIN, 1996: in press (same material than McLAY).

REMARK. — *Dromia wilsoni* is mainly a sublittoral species, usually found within the first 100m, but it is also recorded up to 520m (cf. in McLAY, 1991: 475).

FAMILY DYNOMENIDAE

Dynomene hispida Desmaret, 1825

DISTRIBUTION. — Society (Moorea); Tuamotu (Marutea South, Tikehau).

REFERENCES. — *Dynomene hispida* - NOBILI, 1907: 378 ("Marutea-Vaitutaki" = Marutea South). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 25; 1977b: 212 (Moorea); 1989: 111, 114 (Moorea, Tikehau). — GUINOT, 1985: 448 (List).

Dynomene praedator A. Milne Edwards, 1879

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Dynomene praedator* - ORTMANN, 1892b: 534, pl. 26, fig. 3 (Tahiti). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 448 (List). — *Dynomene* sp. - NAIM, 1980a: 55, fide observation and personal communication of McLAY (Moorea; MNHN B20203). — *Dynomene sinense* - ODINETZ, 1983: 208 (Tahiti; MNHN B17090). — *Dynomene sinensis* (sic) - GUINOT, 1985: 448 (List; after ODINETZ) - These two references, not *D. sinense* Chen, 1979 = *D. praedator* fide observation and personal communication of McLAY.

Dynomene spinosa Rathbun, 1911

DISTRIBUTION. — Marquesas; Tuamotu (Raroia).

REFERENCES. — *Dynomene spinosa* - BALSS, 1935: 115 (Marquesas). — HOLTHUIS, 1953: 3 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"; Marquesas). — GUINOT, 1985: 448 (List).

FAMILY RANINIDAE

Notosceles chimonis Bourne, 1922

DISTRIBUTION. — Marquesas (Eiao) - Sublittoral to deep.

REFERENCES. — *Notosceles chimonis* - POUPIN, 1996: in press (Eiao; 54-101m).

REMARK. — Species known between 45-52 m (SERÈNE & UMALI, 1972), 75-90m (RIBES, 1989), and up to 450m (MONOD, 1975).

Ranina ranina (Linné, 1758)

DISTRIBUTION. — Marquesas (Eiao, Fatu Hiva, Nuku Hiva) - Sublittoral.

REFERENCES. — *Ranina ranina* - GUINOT, 1985: 449 (List; certainly after the following dry specimen: MNHN n°223 "Nouhiva" = Nuku Hiva) - NEW MATERIAL - Coll. and det. J. POUPIN (Eiao, Fatu Hiva; juveniles, 100m).

FAMILY CALAPPIDAE

Ashtoret lunaris (Forskål, 1775)

DISTRIBUTION. — Marquesas (Nuku Hiva).

REFERENCES. — *Matuta banksii* - RATHBUN, 1907: 68 (Nuku Hiva). — GUINOT, 1985: 453 (List). — *Ashtoret lunaris* - GALIL & CLARK, 1994: 5, fig. 1a-b, pl. 1a-b (Syn.) - SYNONYMS - *Matuta banksi* Leach, 1817.

Ashtoret picta (Hess, 1865)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Matuta picta* - POUPIN, 1994a: 27, fig. 23, pl. 3b (Tahiti). — *Ashtoret picta* - GALIL & CLARK, 1994: 18, fig. 3c-d, pl. 6a-b (Tahiti).

REMARK. — In the revision of the genus *Matuta* by GALIL & CLARK (1994), the location "Tahiti", that should appear under *Ashtoret picta*, is erroneously mentioned under *Ashtoret granulosa* (Miers, 1877), species still unknown from French Polynesia (GALIL, personal communication).

Calappa calappa (Linné, 1758)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Calappa calappa* - MONTEFORTE, 1984: 173, annex 1, tab. a (Tahiti; MONTEFORTE's material verified in the CRIODE collections, Moorea). — GUINOT, 1985: 449 (List).

Calappa hepatica (Linné, 1758)

DISTRIBUTION. — Gambier (Mangareva); Marquesas (Nuku Hiva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Hao, Hikueru, Marutea North?, Marutea South, Mataiva, Moruroa, Takapoto).

REFERENCES. — *Calappa hepatica* - NOBILI, 1907: 378 ("Gatavake" = Mangareva, Hao, Marutea = Marutea North?, Marutea South). — RATHBUN, 1907: 67 (Bora Bora). — PESTA, 1913: 37 (Tahiti). — BOONE, 1934: 32, pl. 8-10 (Nuku Hiva). — SEURAT, 1934: 59, 60 (Hao, Marutea South). — CHABOIS L. & F., 1954: 92, unnumbered fig. (French Polynesia). — FOREST & GUINOT, 1961: 11, fig. 1a-b, 2 (Hikueru; Syn.); 1962: 56 (Biogeography "Tahiti-Tuamotu; Marquesas"). — MONTEFORTE, 1984: 173, annex 1, tab. a, photograph p. 140c (Mataiva, Moorea, Tahiti, Takapoto); 1987: 8 (Moorea). — DELESALLE, 1985: 288 (Mataiva). — GUINOT, 1985: 449 (List). — POUPIN, 1994a: 26, fig. 22, pl. 3a (Hikueru, Moruroa, Tahiti). — *Calappa tuberculata* - HELLER, 1865: 69 (Tahiti) - SYNONYMS - *Calappa tuberculata* Fabricius, 1798.

Matuta victor (Fabricius, 1781)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Matuta victor* - HELLER, 1865: 69 (Tahiti). — GALIL & CLARK, 1994: 39, fig. 7a-b, pl. 13a-b (cf. Remark).

REMARK. — HELLER's reference is not mentioned in the revision by GALIL & CLARK (1994). These authors consider that the eastern distribution of this species is limited to the Fiji. Thus, the revision of HELLER's material would be important to confirm the presence of *Matuta victor* in French Polynesia.

FAMILY LEUCOSIIDAE

***Ebaliopsis erosa* (A. Milne Edwards, 1874)**

DISTRIBUTION. — Gambier; Tuamotu (Marutea South).

REFERENCES. — *Ebalia erosa* - NOBILI, 1907: 378 (Marutea). — *Ebaliopsis erosa* - FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — SERÈNE, 1977: 55, fig. 5-6 (Gambier). — GUINOT, 1985: 453 (List).

***Heteronucia venusta* Nobili, 1906**

DISTRIBUTION. — Society (Moorea); Tuamotu (Fakahina, Fakarava, Hao, Tikehau).

REFERENCES. — *Heteronucia venusta* Nobili, 1906a: 260; 1907: 379, pl. 5, fig. 14 ("Ohura" = Hao). — FOREST & GUINOT, 1961: 13, fig. 3a-b, 4, pl. 5, fig. 1-2 (Fakahina; Syn.); 1962: 56 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 25; 1977b: 212 (Moorea); 1989: 113 (Tikehau). — GUINOT, 1985: 453 (List). — *Nucia gelida* Rathbun, 1907: 68, pl. 5, fig. 4, pl. 9, fig. 2 (Fakarava). — GUINOT, 1985: 453 (List).

***Nucia rosea* Nobili, 1906**

DISTRIBUTION. — Gambier (Mangareva).

REFERENCES. — *Nucia rosea* Nobili, 1906a: 261; 1907: 381 (Mangareva). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List).

***Nursia mimetica* Nobili, 1906**

DISTRIBUTION. — Gambier (Mangareva).

REFERENCES. — *Nursia mimetica* Nobili, 1906a: 261; 1907: 380, pl. 5, fig. 13 (Mangareva). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List).

FAMILY MAJIDAE

***Acanthophrys cristimanus* A. Milne Edwards, 1865**

DISTRIBUTION. — Marquesas (Nuku Hiva).

REFERENCES. — *Acanthophrys cristimanus* A. Milne Edwards, 1865a: 141, pl. 5, fig. 3 (Nuku Hiva). — GRIFFIN & TRANTER, 1986: 105 (cf. Remark).

REMARK. — GRIFFIN & TRANTER (1986) have re-established the validity of the genus *Acanthophrys*, formerly transferred in *Hyastenus*, and *Acanthophrys cristimanus* is the type species of the genus.

***Camposcia retusa* Latreille, 1829**

DISTRIBUTION. — Tuamotu (Fangataufa) - Littoral to deep?

REFERENCES. — *Camposcia retusa* - GUINOT, 1985: 452, with a ? (List) - NEW MATERIAL - Coll. and det. J. POUPIN (Fangataufa; 220m, cf. Remark).

REMARK. — The presence of this species in French Polynesia, only inferred by GUINOT because of its large distribution, is here confirmed by one specimen, collected at an unusual depth.

***Cyclax suborbicularis* (Stimpson, 1858)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Cyclax suborbicularis* - FOREST & GUINOT, 1961: 15, fig. 5-6, 8 bis, 10, pl. 6, fig. 1-2 (Tahiti); 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List).

***Huenia proteus* de Haan, 1839**

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Huenia proteus* - KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea). — GUINOT, 1985: 452 (List).

***Hyastenus* aff. *borradailei* (Rathbun, 1907)**

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Hyastenus* aff. *borradaeilli* (sic) - PEYROT-CLAUSADE, 1989: 115 (Moorea).

***Menaethius monoceros* (Latreille, 1825)**

DISTRIBUTION. — Marquesas; Society (Moorea, Tahiti); Tuamotu (Fakarava, Hao, Raroia, Takapoto, Tikehau) - *Littoral to sublittoral* (30m).

REFERENCES. — *Menaethius tuberculatus* Dana, 1852b: 123; 1855, pl. 5, fig. 1a-c (Tuamotu). — *Menaethius monoceros* - NOBILI, 1907: 382 ("Ohura" = Hao). — RATHBUN, 1907: 64 (Fakarava). — HOLTHUIS, 1953: 4 (Raroia). — FOREST & GUINOT, 1961: 14, fig. 9a-b (Tahiti; Syn.); 1962: 56 (Biogeography "Tahiti-Tuamotu, Marquesas"). — PEYROT-CLAUSADE, 1977a, annex of the species: 25 (Moorea); 1977b: 212; 1989: 112, 115 (Moorea, Tikehau; 30m). — ODINETZ, 1983: 208 (Moorea, Tahiti, Takapoto). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti, Takapoto).

***Micippa margaritifera* Henderson, 1893**

DISTRIBUTION. — Tuamotu (Tikehau).

REFERENCES. — *Micippa* (sic) *margaritifera* - PEYROT-CLAUSADE, 1989: 112 (Tikehau).

***Micippa parca* Alcock, 1895**

DISTRIBUTION. — Tuamotu (Makemo).

REFERENCES. — *Lophomicippa limbata* Rathbun, 1907: 65, pl. 5, fig. 3, pl. 6, fig. 1, 1g (Makemo). — FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List). — *Micippa parca* - GRIFFIN & TRANTER, 1986: 277 (Syn.).

***Micippoides angustifrons* A. Milne Edwards, 1873**

DISTRIBUTION. — Tuamotu (Raroia).

REFERENCES. — *Micippoides angustifrons* - HOLTHUIS, 1953: 5 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List).

***Perinea tumida* Dana, 1851**

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Fakarava, Takapoto, Tikehau).

REFERENCES. — *Perinea tumida* - RATHBUN, 1907: 65 (Fakarava). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 25; 1977b: 212 (Moorea); 1989: 112, 115 (Moorea, Tikehau). — KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea, Takapoto). — ODINETZ, 1983: 208 (Moorea, Tahiti, Takapoto). — GUINOT, 1985: 453 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti, Takapoto).

***Schizophrys aspera* (H. Milne Edwards, 1834)**

DISTRIBUTION. — French Polynesia.

REFERENCES. — *Schizophrys aspera* - GUINOT, 1985: 453, with a ? (List).

REMARK. — Cited by GUINOT, only because of the large distribution of the species (see SAKAI, 1976, or DAI & YANG, 1991: Japon, Hawaii, ... Australia).

Simocarcinus obtusirostris (Miers, 1879)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Trigonothir obtusirostris* - FOREST & GUINOT, 1961: 14 (Tahiti); 1962: 56 (Biogeography "Tahiti-Tuamotu"). — *Simocarcinus obtusirostris* - GUINOT, 1985: 453 (List). — GRIFFIN & TRANTER, 1986: 98 (Syn.).

Tylocarcinus dumerili (H. Milne Edwards, 1834)

DISTRIBUTION. — Gambier (Mangareva); Tuamotu (Hao).

REFERENCES. — *Tylocarcinus gracilis* - NOBILI, 1907: 382 (Hao, Mangareva). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List). — *Tylocarcinus dumerili* - GRIFFIN & TRANTER, 1986: 197, fig. 67a-b (Syn.) - SYNONYMS - *Tylocarcinus gracilis* Miers, 1879.

FAMILY PARTHENOPIDAE

Actaeomorpha alvae Boone, 1934

DISTRIBUTION. — Society (Raiatea).

REFERENCES. — *Actaeomorpha alvae* Boone, 1934: 37 pl. 11 (Raiatea). — FOREST & GUINOT, 1962: 56 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966b: 759 (cf. Remark); 1985: 453 (List).

REMARK. — According to the drawing published by BOONE (1934), GUINOT (1966b) considers that this species could possibly be, either *Actaeomorpha erosa* Miers, 1878, or *A. punctata* Edmonson, 1935.

We keep here the genus *Actaeomorpha* in the Parthenopidae, but GUINOT (1966b, 1967), in her study of the genera *Aethra* (cf. hereafter *A. scruposa*), *Osachila*, *Hepatus*, *Hepatella* and *Actaeomorpha*, has modified this usual classification and placed these genera in a group *Incertainae sedis* Parthenoxystomata (cf. GUINOT's, 1985 List).

Aethra scruposa (Linné, 1764)

DISTRIBUTION. — Society.

REFERENCES. — *Aethra scruposa* - GUINOT, 1985: 453, with a ? (Society; listed only according to the large distribution of the species, without material from French Polynesia).

Daldorfia horrida (Linné, 1758)

DISTRIBUTION. — Gambier (Mangareva); Society (Tahiti); Tuamotu (Fangatau, Hao, Mataiva).

REFERENCES. — *Parthenope horrida* - NOBILI, 1907: 382 ("Fagatau" = Fangatau, Hao, Mangareva). — SEURAT, 1934: 60 ("Fagatau" = Fangatau, Gambier, Hao). — CHABOIS L. & F., 1954: 91, fig. 7 (Tahiti, Tuamotu). — FOREST & GUINOT, 1961: 26, fig. 14 (Tahiti); 1962: 58 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 174, annex 1, tab. a (Mataiva). — GUINOT, 1985: 453 (List). — *Daldorfia horrida* - SAKAI, 1976: 283, pl. 96, fig. 2, text-fig. 157 (Syn.).

Parthenope contrarius (Herbst, 1796)

DISTRIBUTION. — Marquesas (Eiao) - Sublittoral.

REFERENCES. — *Parthenope contrarius* - NEW MATERIAL - Coll. and det. J. POUPIN (Eiao; 42m).

REMARK. — This new material (1 ♀ and 3 juveniles, Marara st. D38) agrees very well with the description and the good photograph published by RATBHUN (1906: 885, pl. 17, fig. 1), under *Parthenope (Rhinolambrus) lamelligera* (White, 1847). According to SAKAI (1976: 273) WHITE's species is a synonym of *P. (Rhinolambrus) pelagicus* (Rüppell, 1830), but RATHBUN's material belongs to *P. contrarius*.

Parthenope hoplonotus (Adams & White, 1848)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Aulacolambrus hoplonotus* - FOREST & GUINOT, 1961: 26, fig. 12a-c, 13 (Tahiti); 1962: 58 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List). — *Parthenope (Aulacolambrus) hoplonotus* - SAKAI, 1976: 280 (Syn.).

FAMILY EUMEDONIDAE

Echinoecus pentagonus (A. Milne Edwards, 1879)

DISTRIBUTION. — Tuamotu (Hao, Raroia).

REFERENCES. — *Eumedon convictor* Bouvier & Seurat, 1905: 629 (Hao). — NOBILI, 1907: 382 (Hao). — SEURAT, 1934: 58 (Hao). — *Eumedonus convictor* - HOLTHUIS, 1953: 6 (Raroia). — MORRISON, 1954: 6 (Raroia) — *Echinoecus pentagonus* - SERÈNE et al., 1958: 152 (Syn.). — GUINOT, 1985: 453 (List). — *Eumedonus pentagonus* - FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu").

FAMILY PORTUNIDAE

SUBFAMILY CATOPTRINAE

Carupa tenuipes Dana, 1852

DISTRIBUTION. — Gambier (Akamaru); Society (Huahine, Maiao?, Moorea, Tahiti); Tuamotu (Makatea, Makemo, Marutea South, Pukapuka, Raroia, Tikehau).

REFERENCES. — *Carupa tenuipes* Dana, 1852a: 85 (Tuamotu); 1852b: 279; 1855, pl.17, fig. 4a-e ("Paumotu archipelago?"). — A. MILNE EDWARDS, 1861: 386 (Tuamotu; DANA's material). — STEPHENSON & REES, 1967: 5 (Huahine, Moorea, Maiai = Maiao?, "Tickahau" = Tikehau). — SAKAI, 1976: 325 (Syn.). — STEPHENSON, 1976: 12 (Pukapuka). — MONTEFORTE, 1984: 173, annex 1, tab. a (Makatea, Moorea); 1987: 8 (Moorea). — GUINOT, 1985: 449 (List). — *Carupa laeviuscula* Heller, 1862: 520 ("Tarti" = Tahiti); 1865: 27, pl.3, fig. 2 (Tahiti). — NOBILI, 1907: 386 (Akamaru, "Marutea-Vaitutaki" = Marutea South). — RATHBUN, 1907: 64 (Makemo). — HOLTHUIS, 1953: 9 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu") - NEW MATERIAL - coll. C. HILY, det. K. MOOSA (Tikehau).

Catoptrus nitidus A. Milne Edwards, 1870

DISTRIBUTION. — Marquesas (Eiao); Society (Huahine); Tuamotu (Makemo, Tikehau) - *Sublittoral*.

REFERENCES. — *Catoptrus nitidus* - RATHBUN, 1907: 60 (Makemo). — FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1972: 29 (Syn.). — GUINOT, 1985: 449 (List). — *Libystes truncatifrons* - STEPHENSON & REES, 1967: 6 ("Maroe" = Huahine, "Tickahau" = Tikehau) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao; 42m) - SYNONYMS - *Libystes truncatifrons* (de Man, 1887).

SUBFAMILY CAPHYRINAE

Caphyra rotundifrons (A. Milne Edwards, 1869)

DISTRIBUTION. — Society (Bora Bora, Tahiti).

REFERENCES. — *Caphyra rotundifrons* - RATHBUN, 1907: 60, pl. 1, fig. 4 (Tahiti). — FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON & REES, 1967: 7 (Bora Bora, "Mata Uta Papeete" = Tahiti). — GUINOT, 1985: 449 (List).

***Caphyra tridens* Richters, 1880**

DISTRIBUTION. — Gambier (Mangareva).

REFERENCES. — *Caphyra rotundifrons* var. *tridens* Richters - NOBILI, 1907: 386 ("Rikitea, Teone Kura" = Mangareva). — *Caphyra tridens* - CROSNIER, 1975: 747, fig. 3a-n (Mangareva).

***Lissocarcinus elegans* Boone, 1934**

DISTRIBUTION. — Society (Raiatea).

REFERENCES. — *Lissocarcinus elegans* Boone, 1934: 50, pl. 16 (Raiatea). — FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1972: 27 (Distribution; French Polynesia only). — GUINOT, 1985: 449 (List).

***Lissocarcinus laevis* Miers, 1886**

DISTRIBUTION. — Marquesas (Eiao, Hiva Oa, Nuku Hiva) - *Sublittoral*.

REFERENCES. — *Lissocarcinus laevis* - STEPHENSON, 1976: 12 (Nuku Hiva) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao, Hiva Oa; 42-53m).

***Lissocarcinus orbicularis* Dana, 1852**

DISTRIBUTION. — Society (Huahine, Moorea, Raiatea, Tahiti); Tuamotu (Hikueru, Moruroa, Tikehau).

REFERENCES. — *Lissocarcinus orbicularis* - FOREST & GUINOT, 1961: 27, fig. 15a-b, 16a-c (Hikueru); 1962: 58 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON & REES, 1967: 7 (Huahine, Raiatea, "Tikahau" = Tikehau). — STEPHENSON, 1976: 12 (Moorea). — CHEVALIER *et al.*, 1968: 112, 137 (Moruroa). — MONTEFORTE, 1984: 173, annex 1, tab. a (Tahiti). — GUINOT, 1985: 449 (List). — PEYROT-CLAUSADE, 1989: 113 (Tikehau).

SUBFAMILY PORTUNINAE

***Charybdis annulata* (Fabricius, 1798)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Goniosoma annulatum* (Fabricius) - ORTMANN, 1893a: 82 (Tahiti). — *Charybdis annulata* - FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — *Charybdis (Charybdis) annulata* - CROSNIER, 1962: 78, fig. 136-139, pl. 5, fig. 2 (Distribution only, Tahiti). — GUINOT, 1985: 449, with a ? (List).

***Charybdis erythrodactyla* (Lamarck, 1818)**

DISTRIBUTION. — Austral (Rurutu); Marquesas (Nuku Hiva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fangataufa, Hikueru, Makatea, Mataiva, Moruroa, Raroia, Taiaro, Takapoto).

REFERENCES. — *Goniosoma erythrodactylum* - A. MILNE EDWARDS, 1861: 369 (Marquesas). — DE MAN, 1889: 424 (Tahiti). — ORTMANN, 1893a: 81 (Marquesas). — *Charybdis erythrodactyla* - NOBILI, 1906b: 118, fig. 3 (Nuku Hiva). — SENDLER, 1923: 40 (Makatea). — BOONE, 1934: 57, pl. 18-19 (Tahiti). — LEENE, 1936: 117, fig. 1-5 (Marquesas, Makatea). — HOLTHUIS, 1953: 6 (Raroia). — MORRISON, 1954: 16 (Raroia). — FOREST & GUINOT, 1961: 30 (Hikueru); 1962: 58 (Biogeography "Tahiti-Tuamotu, Marquesas"). — GUINOT, 1966a: 48 (Raroia). — CHEVALIER *et al.*, 1968: 92, 137 (Fangataufa). — MONTEFORTE, 1984: 173, annex 1, tab. a, photograph p. 140(b) (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — SALVAT, 1986b: 72, photograph (French Polynesia). — *Charybdis (Goniosupradens) erythrodactyla* - LEENE, 1938: 134, fig. 77-80 (Marquesas). — STEPHENSON & REES, 1967: 13 (Bora Bora, Moorea). — STEPHENSON, 1976: 15 ("Taiohae" = Nuku Hiva). — GUINOT, 1985: 449 (List). — POUPIN 1994a: 29, fig. 25, pl. 3d (Rurutu, Taiaro) - NEW MATERIAL - Coll. and det. J. POUPIN (Moruroa).

***Charybdis hawaiensis* Edmonson, 1954**

DISTRIBUTION. — Tuamotu (Taiaro).

REFERENCES. — *Charybdis (Charybdis) hawaiensis* - POUPIN, 1994a: 28, fig. 24, pl. 3c (Taiaro).

Charybdis orientalis Dana, 1852

DISTRIBUTION. — Society.

REFERENCES. — *Charybdis (Charybdis) orientalis* - LEENE, 1938: 69 (Society; cf. Remark).

REMARK. — Although LEENE (1938) does not mention clearly the location "Society", she indicates that F. CHACE has examined, for her, in the USNM collections, a male from the Society Islands attributed to *C. orientalis* (confronted with the type specimen).

Charybdis paucidentata A. Milne Edwards, 1861

DISTRIBUTION. — Marquesas (Hiva Oa, Tahuata); Tuamotu (Taiaro) - *Littoral to sublittoral*.

REFERENCES. — *Charybdis (Gonioinfradens) paucidentata* - POUPIN, 1994a: 30, fig. 26, pl. 3e (Hiva Oa, Tahuata, Taiaro; 0-100m); 1996: in press (Hiva Oa, Tahuata).

Lupocycclus quinquedentatus Rathbun, 1906

DISTRIBUTION. — Austral (Maria, Rurutu); Marquesas (Nuku Hiva); Society (Bora Bora) - *Sublittoral to deep*.

REFERENCES. — *Lupocycclus quinquedentatus* - STEPHENSON, 1976: 15 ("Hatwata" = Haatuatua bay, Nuku Hiva) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Bora Bora, Maria, Rurutu; 80-110m).

Portunus alexandri (Rathbun, 1907)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Callinectes alexandri* Rathbun, 1907: 61, pl. 2, fig. 1, pl. 9, fig. 3, 3a-b (Tahiti). — GUINOT, 1985: 449 (List).

REMARK. — This portunid has been described from Tahiti, with a paratype from "Suva, Fidjis". According to STEPHENSON (1976: 13), it is in fact a non-identifiable *Portunus*.

Portunus dubius (Laurie, 1906)

DISTRIBUTION. — Marquesas (Eiao, Fatu Hiva, Hiva Oa, Nuku Hiva, Tahuata) - *Sublittoral to deep*.

REFERENCES. — *Portunus dubius* - STEPHENSON, 1976: 16 (Marquesas; numerous stations without precisions) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao, Fatu Hiva, Hiva Oa, Nuku Hiva, Tahuata; 42-140m).

Portunus granulatus (H. Milne-Edwards, 1834)

DISTRIBUTION. — Gambier (Mangareva); Marquesas (Nuku Hiva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fakarava, Raroia, Tikehau).

REFERENCES. — *Neptunus (Achelous) granulatus* - ORTMANN, 1893a: 72 (Tahiti). — NOBILI, 1907: 383 ("Rikitea" = Mangareva). — BOONE, 1934: 60, pl. 20 (Nuku Hiva). — SEURAT, 1934: 59 (Mangareva). — *Portunus (Achelous) granulatus* - RATHBUN, 1907: 60 (Bora Bora, Fakarava). — SENDLER, 1923: 40 (Tahiti). — *Portunus (Cycloachelous) granulatus* - HOLTHUIS, 1953: 6 (Raroia). — MORRISON, 1954: 7 (Raroia). — GUINOT, 1985: 449 (List). — *Portunus granulatus* - FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu, Marquesas"). — STEPHENSON & REES, 1967: 25 (Moorea, Tuamotu). — TÜRKAY, 1971: 127 (Tahiti). — STEPHENSON, 1976: 16 (Tahiti). — MONTEFORTE, 1984: 173, annex 1, tab. a (Moorea, Tahiti); 1987: 8 (Moorea) - NEW MATERIAL - Coll. C. HILY, det. K. MOOSA (Tahiti, Tikehau).

Portunus guinotae Stephenson & Rees, 1961

DISTRIBUTION. — Tuamotu (Marutea South).

REFERENCES. — *Portunus guinotae* Stephenson & Rees, 1961: 425, fig. 1b, d, g, 2d-f (Marutea South). — FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1972: 39 (Marutea South). — *Portunus (Xiphonectes) guinotae* - GUINOT, 1985: 449 (List).

Portunus iranjae Crosnier, 1962

DISTRIBUTION. — Marquesas (Eiao, Fatu Hiva, Hiva Oa, Tahuata); Society (Moorea) - *Littoral to sublittoral*.

REFERENCES. — *Portunus iranjae* - STEPHENSON & REES, 1967: 30 ("Papetoai bay" = Moorea). — STEPHENSON, 1976: 16 ("Marquesas expedition, st. THX" = Haava strait, between Tahuata and Hiva Oa, cf. HARALD, 1967). — *Portunus (Xiphonectes) iranjae* - GUINOT, 1985: 449 (List) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao, Fatu Hiva, Hiva Oa, Tahuata; 54m).

Portunus longispinosus (Dana, 1852)

DISTRIBUTION. — Marquesas (Hiva Oa, Tahuata); Tuamotu (Marutea South, Raroia).

REFERENCES. — *Neptunus (Hellenus) longispinosus* - NOBILI, 1907: 383 (Marutea South). — *Portunus (Hellenus) longispinosus* - HOLTHUIS, 1953: 7 (Raroia). — MORRISON, 1954: 8 (Raroia). — *Portunus longispinosus* - FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1976: 16 ("Marquesas expedition st. THX, haul 5" = Haava strait, between Tahuata and Hiva Oa, cf. HARALD, 1967). — *Portunus (Xiphonectes) longispinosus* - GUINOT, 1985: 449 (List).

REMARK. — This species can be confused with all the species belonging to the *longispinosus* complex, as *Portunus iranjae* and *P. macrophthalmus* (cf. STEPHENSON & REES, 1967; STEPHENSON, 1976; and NAGAI, 1981).

Portunus macrophthalmus Rathbun, 1906

DISTRIBUTION. — Marquesas (Eiao, Hiva Oa) - *Sublittoral*.

REFERENCES. — *Portunus macrophthalmus* - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao, Hiva Oa; 42-53m).

Portunus nipponensis Sakai, 1938

DISTRIBUTION. — Tuamotu (Moruroa) - *Sublittoral to deep*.

REFERENCES. — *Portunus nipponensis* - POUPIN et al. 1990: 17 (French Polynesia). — POUPIN, 1996: in press *pro parte* (Moruroa; cf. Remark).

REMARK. — This species is usually found in shallow waters (15-50m) but, in French Polynesia, it has been trapped up to 130m. Except for Moruroa, the localities mentioned in POUPIN (1996) concerned in fact a new species, related to *P. nipponensis*, but with a distinct male pleopod (MOOSA & CROSNIER, in study).

Portunus orbitosinus Rathbun, 1911

DISTRIBUTION. — Marquesas (Eiao) - *Sublittoral*.

REFERENCES. — *Portunus orbitosinus* - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao; 42m).

Portunus pelagicus (Linné, 1758)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Neptunus pelagicus* - HELLER, 1865: 27 (Tahiti). — *Portunus pelagicus* - FOREST & GUINOT, 1962: 58 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1972: 41 (Distribution only, Tahiti). — *Portunus (Portunus) pelagicus* - GUINOT, 1985: 449 (List).

REMARK. — Although this species has often been reported from Tahiti (BOONE, 1934; STEPHENSON & CAMPBELL, 1959; CROSNIER, 1962; STEPHENSON & REES, 1967; STEPHENSON, 1972; SAKAI, 1976; DAI & YANG, 1991), it seems that the single material examined from this locality is mentioned in HELLER (1865). It would thus be very interesting to check this reference.

Portunus sanguinolentus (Herbst, 1783)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Mataiva).

REFERENCES. — *Neptunus sanguinolentus* - CHABOIS L. & F., 1954: 91, fig. 9 (French Polynesia). — *Portunus sanguinolentus* - FOREST & GUINOT, 1961: 29, fig. 17a-b, 18 (Tahiti); 1962: 58 (Biogeography "Tahiti-Tuamotu"). —

MONTEFORTE, 1984: 173, annex 1, tab. a (Moorea). — SALVAT, 1986b: 72 (French Polynesia). — POUPIN, 1994a: 31, fig. 27, pl. 3f (Mataiva, Tahiti). — *Portunus sanguinolentus sanguinolentus* - STEPHENSON, 1976: 19 (Tahiti). — *Portunus (Portunus) sanguinolentus* - GUINOT, 1985: 449 (List).

Scylla serrata (Forskål, 1775)

DISTRIBUTION. — Society (Huahine, Raiatea, Tahiti, Tupai) - *Brackish to sea water*.

REFERENCES. — *Scylla serrata* - HELLER, 1865: 27 (Tahiti). — MIERS, 1886: 185 (Tahiti). — BOONE, 1934: 68, pl. 25-30 (Huahine, Tahiti). — SEURAT, 1934: 58 (Tahiti). — CHABOIS L. & F., 1954: 90, unnumbered fig. (Huahine, Raiatea). — FOREST & GUINOT, 1961: 27 (Tahiti); 1962: 58 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (Society); 1985: 449 (List). — BABLET, 1972: 32, pl. 11 (French Polynesia). — SALVAT, 1986b: 70, 72 (French Polynesia) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Raiatea, Tupai).

Thalamita admete (Herbst, 1803)

DISTRIBUTION. — Gambier (Mangareva); Marquesas?; Society (Bora Bora, Huahine, Moorea, Raiatea, Tahiti); Tuamotu (Fakarava, Hikueru, Kaukura, Makemo, Mataiva, Takapoto, Tikehau).

REFERENCES. — *Thalamita admete* - HELLER, 1865: 28 (Tahiti). — ORTMANN, 1893a: 83 (Tahiti). — NOBILI, 1907: 383 (Kaukura, Mangareva). — RATHBUN, 1907: 63 (Fakarava, Makemo). — FOREST & GUINOT, 1961: 30, fig. 19a-b (Hikueru, Tahiti); 1962: 58 (Biogeography "Tahiti-Tuamotu, Marquesas" with a ?). — STEPHENSON & REES, 1967: 56, fig. 20 (Bora Bora, Huahine, Moorea, Raiatea, Tahiti, "Tikahau" = Tikehau). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — MONTEFORTE, 1984: 173, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 8 (Moorea). — DELESALLE, 1985: 288 (Mataiva). — *Thalamita (Thalamita) admete* - GUINOT, 1985: 449 (List).

Thalamita bouvieri Nobili, 1906

DISTRIBUTION. — Gambier (Mangareva).

REFERENCES. — *Thalamita bouvieri* Nobili, 1906a: 262; 1907: 384, pl. 2, fig. 2 ("Rikitea" = Mangareva). — SEURAT, 1934: 60 (French Polynesia). — CROSNIER, 1962: 119, fig. 201-204, pl. 10, fig. 2 (Mangareva; syntypes). — FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — *Thalamita (Pseudothalamitopsis) bouvieri* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 47) - RELEVANT MATERIAL - *Thalamitoïdes* (sic) aff. *bouvieri* - PEYROT-CLAUSADE, 1989: 113 (Tikehau).

Thalamita chaptalii (Audouin, 1826)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Thalamita chaptalii* - FOREST & GUINOT, 1961: 34, fig. 21a-b (Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — *Thalamita (Neothalamita) chaptalii* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 43).

Thalamita coerulipes Jacquinot, 1852

DISTRIBUTION. — Gambier (Kamaka, Mangareva); Society (Bora Bora, Huahine, Tahiti); Tuamotu (Fakarava).

REFERENCES. — *Thalamita coerulipes* Jacquinot, 1852, pl. 5, fig. 6-10 (Mangareva). — JACQUINOT & LUCAS, 1853: 53 (Mangareva). — A. MILNE EDWARDS, 1861: 363 (Mangareva). — NOBILI, 1907: 383 (Kamaka) — RATHBUN, 1907: 63 (Fakarava, Society). — BOONE, 1934: 78, pl. 35 (Bora Bora). — FOREST & GUINOT, 1961: 32 (Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON & REES, 1967: 64 (Huahine). — POUPIN, 1994a: 32, fig. 28, pl. 3g (Kamaka, Mangareva, Tahiti, Tuamotu). — *Thalamita (Thalaminella) coerulipes* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 51).

Thalamita cooperi Borradaile, 1903

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Thalamita cooperi* - NAIM, 1980a: annex 1, tab. 3 (Moorea; material not found in MNHN).

***Thalamita corrugata* Stephenson & Rees, 1961**

DISTRIBUTION. — Tuamotu (Tikehau).

REFERENCES. — *Thalamita corrugata* - STEPHENSON & REES, 1967: 65, fig. 23 ("Tickahau" = Tikehau). — GUINOT, 1985: 449 (List).

***Thalamita crenata* (Latreille, 1829)**

DISTRIBUTION. — Marquesas; Society (Bora Bora, Maiao?, Moorea, Tahiti); Tuamotu (Mataiva, Takapoto, Tikehau).

REFERENCES. — *Thalamita crenata* - A. MILNE EDWARDS, 1861: 365 (Marquesas). — ORTMANN, 1893a: 86 (Marquesas). — RATHBUN, 1907: 62 (Bora Bora). — SEURAT, 1934: 59 (Marquesas). — FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu, Marquesas"). — STEPHENSON & REES, 1967: 66 ("Maiai" = Maiao?, "Tikahau" = Tikehau). — MONTEFORTE, 1984: 173, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 8 (Moorea). — SALVAT & RICHARD, 1985: 356, 362 (Takapoto) — *Thalamita (Thalaminella) crenata* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 51) - NEW MATERIAL - Coll. C. HILY, det. K. MOOSA (Tahiti, Tikehau).

***Thalamita dakini* Montgomery, 1931**

DISTRIBUTION. — Society (Bora Bora, Moorea); Tuamotu (Makatea, Mataiva, Takapoto).

REFERENCES. — *Thalamita dakini* - STEPHENSON & REES, 1967: 69 (Bora Bora, Moorea). — MONTEFORTE, 1984: 173, annex 1, tab a (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — *Thalamita (Thalamitopsis) dakini* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 47).

***Thalamita danae* Stimpson, 1858**

DISTRIBUTION. — French Polynesia.

REFERENCES. — *Thalamita danae* - NEW MATERIAL - Coll. PLESSIS, det. K. MOOSA (French Polynesia).

***Thalamita demani* Nobili, 1905**

DISTRIBUTION. — Marquesas (Nuku Hiva).

REFERENCES. — *Thalamita demani* - STEPHENSON, 1976: 20 ("Hatwata" = Haatuatua bay, Nuku Hiva).

***Thalamita edwardsi* Borradaile, 1900**

DISTRIBUTION. — Gambier (Mangareva, Temoe); Society (Tahiti).

REFERENCES. — *Thalamita admete* var. *Edwardsii* Borr. - NOBILI, 1907: 383 ("Rikitea" = Mangareva, "Timoe" = Temoe). — *Thalamita edwardsi* - FOREST & GUINOT, 1961: 32, fig. 20a-b (Tahiti); 1962: 58 (Biogeography "Tahiti-Tuamotu"). — *Thalamita edwardsi* - GUINOT, 1985: 449, with " = *T. admete* ?" (List; cf. Remark).

REMARK. — This species was formerly considered as a synonym of *Thalamita admete* by STEPHENSON & HUDSON (1957). This assertion is not followed by FOREST & GUINOT (1961), CROSNIER (1962) and DAI & YANG (1991).

***Thalamita gatavakensis* Nobili, 1906**

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Tahiti).

REFERENCES. — *Thalamita pilumnoides* var. *gatavakensis* Nobili, 1906a: 262; 1907: 384 ("Gatavake" = Mangareva). — *Thalamita pilumnoides* ssp. *gatavakensis* - FOREST & GUINOT, 1961: 34, fig. 22a-b, 23-25 (Mangareva); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — *Thalamita gatavakensis* - CROSNIER, 1962: 99, fig. 156a-c, e (Mangareva; syntypes). — STEPHENSON & REES, 1967: 75 (Bora Bora, Tahiti). — *Thalamita (Thalamita) gatavakensis* - GUINOT, 1985: 449 (List) - NEW MATERIAL - Coll. C. HILY, det. K. MOOSA (Tahiti).

***Thalamita gloriensis* Crosnier, 1962**

DISTRIBUTION. — Society (Huahine).

REFERENCES. — *Thalamita gloriensis* - STEPHENSON & REES, 1967: 76 (Huahine). — *Thalamita (Thalamita) gloriensis* - GUINOT, 1985: 449 (List).

Thalamita gracilipes (A. Milne Edwards, 1873)

DISTRIBUTION. — Society (Tahiti); Tuamotu (Moruroa?).

REFERENCES. — *Thalamita gracilipes* - STEPHENSON, 1976: 21 (Tahiti) - RELEVANT MATERIAL - *Thalamonyx* aff. *gracilipes* - SALVAT & RENAUD-MORNANT, 1969: 165 (Moruroa). — *Thalamita (Thalamonyx)* aff. *gracilipes* - GUINOT, 1985: 449 (List).

Thalamita integra Dana, 1852

DISTRIBUTION. — Gambier (Mangareva); Society (Tahiti); Tuamotu (Takapoto, Nukutipipi).

REFERENCES. — *Thalamita integra* Dana, 1852a: 85; 1852b: 281; 1855, pl. 17, fig. 6a-d (Tuamotu). — A. MILNE EDWARDS, 1861: 358 (Tahiti, Tuamotu). — NOBILI, 1907: 383 ("Gatavake" = Mangareva). — FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 173, annex 1, tab. a (Takapoto). — SALVAT & RICHARD, 1985: 350 (Takapoto). — MERSCHARDT-SALVAT, 1991: 89 (Nukutipipi). — *Thalamita (Thalamita) integra* - GUINOT, 1985: 449 (List).

Thalamita macropus Montgomery, 1931

DISTRIBUTION. — Austral (Neilson bank); Marquesas (Fatu Hiva) - *Sublittoral to deep*.

REFERENCES. — *Thalamita macropus* - NEW MATERIAL - Coll. B. RICHER DE FORGES and J. POUPIN, det. K. MOOSA (Neilson bank, Fatu Hiva; 49-100/130m).

Thalamita macrospinifera Rathbun, 1911

DISTRIBUTION. — Austral (Rurutu); Marquesas (Eiao, Hiva Oa); Society (Raiatea); Tuamotu (Makemo, Moruroa) - *Sublittoral to deep*.

REFERENCES. — *Thalamita macrospinifera* - POUPIN, 1996: in press (Makemo, Raiatea; 120-160m) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao, Hiva Oa, Makemo, Moruroa, Raiatea, Rurutu; 80-160m).

Thalamita minuscula Nobili, 1906

DISTRIBUTION. — Tuamotu (Kaukura, Vahitahi).

REFERENCES. — *Thalamita minuscula* Nobili, 1906a: 262; 1907: 386, pl. 1, fig. 15 (Kaukura, Vahitahi). — FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1972: 49 (List with "Only from Tuamotu Is."). — *Thalamita (Neothalamita) minuscula* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 43).

REMARK. — This very small species (maximum width, 4mm), never recorded since its description, could be the juvenile of another species.

Thalamita mitsiensis Crosnier, 1962

DISTRIBUTION. — Austral (Rurutu) - *Sublittoral*.

REFERENCES. — *Thalamita mitsiensis* - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Rurutu; 80m).

Thalamita philippinensis Stephenson & Rees, 1967

DISTRIBUTION. — Austral (Rurutu); Tuamotu (Moruroa) - *Sublittoral to deep*.

REFERENCES. — *Thalamita philippinensis* - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Moruroa, Rurutu; 95-130m).

Thalamita picta Stimpson, 1858

DISTRIBUTION. — Marquesas (Hiva Oa, Nuku Hiva, Tahuata); Gambier?; Society (Raiatea, Tahiti); Tuamotu (Fakarava, Makatea, Mataiva, Raroia, Takapoto).

REFERENCES. — *Goniosoma lineatum* A. Milne Edwards, 1861: 377, pl. 35, fig. 4 (Nuku Hiva) *fide* CROSNIER (1962: 138). — *Thalamita alcocki* - NOBILI, 1907: 384 ("Tagatau" = Gambier?). — *Thalamita gardineri* - RATHBUN, 1907: 63 (Fakarava, Makemo). — *Thalamita picta* - HOLTHUIS, 1953: 8 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1961: 33 (Tahiti; Syn.); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1972: 50 (Syn.); 1976: 23 ("Hoava Strait" = Haava strait, between Tahuata and Hiva Oa, *cf.* HARALD, 1967). — MONTEFORTE, 1984: 173, annex 1, tab. a (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — *Thalamita (Neothalaminella) picta* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 51) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Raiatea) - SYNONYMS - *Thalamita alcocki* de Man, 1902; *T. gardineri* Borradaile, 1902.

Thalamita pilumnoides Borradaile, 1903

DISTRIBUTION. — Society (Huahine, Moorea, Raiatea).

REFERENCES. — *Thalamita pilumnoides* - STEPHENSON & REES, 1967: 87, fig. 32 (Huahine, Moorea, Raiatea). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 212 (Moorea). — MONTEFORTE, 1984: 173, annex 1, tab. a; 1987: 8 (Moorea). — *Thalamita (Neothalamita) pilumnoides* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 43).

Thalamita prymna (Herbst, 1803)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Anaa).

REFERENCES. — *Thalamita prymna* - STEPHENSON, 1976: 23 (Anaa). — PEYROT-CLAUSADE, 1989: 115 (Moorea) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Tahiti).

Thalamita quadrilobata Miers, 1884

DISTRIBUTION. — Society (Bora Bora).

REFERENCES. — *Thalamita quadrilobata* - STEPHENSON & REES, 1967: 92 (Bora Bora). — *Thalamita (Pseudothalamitopsis) quadrilobata* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 47).

Thalamita seurati Nobili, 1906

DISTRIBUTION. — Tuamotu (Marutea South).

REFERENCES. — *Thalamita seurati* Nobili, 1906a: 262; 1907: 385, pl. 2, fig. 1 (Marutea). — SEURAT, 1934: 60 (French Polynesia). — FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON, 1972: 51 (List "Tuamotu"). — *Thalamita (Pseudothalamitopsis) seurati* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 47).

Thalamita spinifera Borradaile, 1903

DISTRIBUTION. — Austral (Raevavae, Tubuai); Marquesas (Eiao, Fatu Hiva, Nuku Hiva); Tuamotu (Makemo) - Sublittoral to deep.

REFERENCES. — *Thalamita spinifera* - STEPHENSON, 1976: 24 ("Marquesas Expedition, 40-80m, 18/ix/1967 to 1/x/1967" = Marquesas, *cf.* HARALD, 1967) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Eiao, Fatu Hiva, Makemo, Nuku Hiva, Raevavae, Tubuai; 42-200m).

Thalamita woodmasoni Alcock, 1899

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Thalamita woodmasoni* - FOREST & GUINOT, 1961: 33 (Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — *Thalamita (Pseudothalamitopsis) woodmasoni* - GUINOT, 1985: 449 (List; subgenus from MOOSA, 1979: 47).

Thalamitoides quadridens A. Milne Edwards, 1869

DISTRIBUTION. — Tuamotu (Moruroa, Rangiroa).

REFERENCES. — *Thalamitoides quadridens* - STEPHENSON, 1976: 26 (Rangiroa) - NEW MATERIAL - Coll. J. POUPIN, det. K. MOOSA (Moruroa).

SUBFAMILY PODOPHTHALMINAE

Podophthalmus vigil (Fabricius, 1798)

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Podophthalmus vigil* - FOREST & GUINOT, 1961: 36 (Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — STEPHENSON & REES, 1967: 104 ("Opunohu Bay" = Moorea). — MONTEFORTE, 1984: 173, annex 1, tab. a (Moorea). — GUINOT, 1985: 449 (List).

FAMILY XANTHIDAE

SUBFAMILY POLYDECTINAE.

Lybia caestifera (Alcock, 1897)

DISTRIBUTION. — Society (Tahiti); Tuamotu (Mataiva?).

REFERENCES. — *Lybia caestifera* - RATHBUN, 1907: 60 (Tahiti). — FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1976: 75 (Syn.; RATHBUN's reference with a ?); 1985: 452, with a ? (List) - RELEVANT MATERIAL - *Lybia cf. coestifera* (sic) - MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva).

Lybia plumosa Barnard, 1947

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau).

REFERENCES. — *Lybia leptochelis* - PEYROT-CLAUSADE, 1977a, annex of the species: 27 (Moorea) not *L. leptochelis* (Zehntner, 1894) = *L. plumosa* fide SERÈNE (1984: 29, 31). — *Lybia plumulosa* (sic) - PEYROT-CLAUSADE, 1989: 113 (Tikehau).

Lybia tessellata (Latreille, 1812)

DISTRIBUTION. — Marquesas; Society (Moorea, Tahiti); Tuamotu (Makatea, Mataiva, Rangiroa, Raroia).

REFERENCES. — *Melia tessellata* - FINNEGAR, 1931: 647 (Marquesas). — *Lybia tessellata* - HOLTHUIS, 1953: 23 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu, Marquesas"). — TAKEDA & MIYAKE, 1970: 15 ("Avatoru" = Rangiroa, Tahiti). — GUINOT, 1976: 70, fig. 17d, 18e, 19c, 20e-h, 22d, pl. 2, fig. 6 (Syn.); 1985: 452 (List). — MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti); 1987: 9 (Moorea).

Polydectus cupulifer (Latreille, 1812)

DISTRIBUTION. — Tuamotu (Raraka).

REFERENCES. — *Polydectus villosus* Dana, 1852a: 81; 1852b: 227; 1855, pl. 13, fig. 3a-e (Raraka). — *Polydectus cupulifer* - FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1976: 65 (Syn.); 1985: 452 (List). — SERÈNE, 1984: 24, fig. 1, pl. 1a (Syn.).

SUBFAMILY CYMOINAE

Cymo andreossyi (Audouin, 1826)

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Cymo andreossyi* - DANA, 1852b: 225; 1855, pl. 13, fig. 2a-b (Tahiti). — HELLER, 1865: 20 (Tahiti). — FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu"). — ODINETZ, 1983: 206 (Moorea, Tahiti). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti).

Cymo deplanatus A. Milne Edwards, 1873

DISTRIBUTION. — Tuamotu (Raroia).

REFERENCES. — *Cymo deplanatus* - HOLTHUIS, 1953: 18 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 450 (List).

Cymo melanodactylus de Haan, 1833

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fakarava, Hao, Hikueru, Moruroa).

REFERENCES. — *Cymo Andreossyi* var. *melanodactyla* - NOBILI, 1907: 397 ("Otepa" = Hao, Mangareva). — *Cymo melanodactylus* - RATHBUN, 1907: 53 (Bora Bora, Fakarava). — PESTA, 1913: 46 (Tahiti). — BOONE, 1934: 144, pl. 74 (Tahiti). — FOREST & GUINOT, 1961: 119 (Hikueru, "Rikitea" = Mangareva); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — ODINETZ, 1983: 206 (Tahiti). — SERÈNE, 1984: 34, fig. 8, pl. 2b (Syn.). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Tahiti). — POUPIN, 1994a: 33, fig. 29, pl. 3h (Mangareva, Moruroa).

Cymo quadrilobatus Miers, 1884

DISTRIBUTION. — Society (Moorea, Tahiti?).

REFERENCES. — *Cymo quadrilobatus* - ODINETZ, 1983: 206 (Moorea). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti).

SUBFAMILY TRICHIINAE

Banareia parvula (Krauss, 1843)

DISTRIBUTION. — Marquesas.

REFERENCES. — *Actaea parvula* - ODHNER, 1925: 35, 51, pl. 3, fig. 13 (Marquesas). — *Banareia parvula* - GUINOT, 1976: 179, with a ? for the genus (Syn.; see the considerations about the generic rank); 1985: 452 (List). — *Banareia parvula* - GARTH et al., 1987: 243 (cited only for the generic rank).

SUBFAMILY LIOMERINAE

Liomera bella (Dana, 1852)

DISTRIBUTION. — Gambier (Mangareva, Tarauru-Roa, Vaiatekeue); Society (Moorea, Tahiti); Tuamotu (Fakareva, Hikueru, Makatea, Makemo, Marutea South, Mataiva, Takapoto, Takaroa, Tikehau).

REFERENCES. — *Carpiloxanthus rugipes* - HELLER, 1865: 17 (Tahiti) not *Liomera rugipes* (Heller, 1861) = *L. bella* fide SERÈNE (1984: 65). — *Carpilodes rugatus* - NOBILI, 1907: 387 ("chenal Waiatekene = Vaiatekeue, "Rikitea" = Mangareva, "Waitutaki" = Marutea South). — RATHBUN, 1907: 37 (Makemo) - All, not *Liomera rugata* (H. Milne Edwards, 1834) = *L. bella* fide SERÈNE (1984: 61). — *Carpilodes vaillantianus* - NOBILI, 1907: 387. — *Carpilodes bellus* - ODHNER, 1925: 16, pl. 1, fig. 9 ("Eimeo" = Moorea, Makemo, Tahiti). — BUITENDIJK, 1960: 257, fig. 2b (Tahiti). — *Liomera bella* - FOREST & GUINOT, 1961: 38, fig. 26a-b (Hikueru, Tahiti, "Taraourou-roa" = Tarauru-Roa);

1962: 60 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 212 (Moorea); 1989: 111 (Tikehau). — MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti). — *Liomera (Liomera) bella* - SERÈNE, 1984: 60, fig. 21, pl. 5e (Syn.) - SYNONYMS - *Carpilodes vaillantianus* A. Milne Edwards, 1862.

Liomera cinctimana (White, 1847)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Moorea, Tahiti).

REFERENCES. — *Liomera lata* - HELLER, 1865: 9 (Tahiti). — *Liomera cinctimana* - ORTMANN, 1893b: 450, pl. 17, fig. 8 (Tahiti). — FOREST & GUINOT, 1961: 39, fig. 27a-b (Tahiti, "Taihoae" = Nuku Hiva); 1962: 60 (Biogeography "Tahiti-Tuamotu, Marquesas"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26 (Moorea). — MONTEFORTE, 1984: 171, annex 1, tab. a (Tahiti). — GUINOT, 1985: 450 (List). — *Carpilodes cinctimanus* - ODHNER, 1925: 14 (Tahiti). — *Liomera (Liomera) cinctimana* - SERÈNE, 1984: 57, fig. 17, pl. 5a (Syn.) - SYNONYMS - *Liomera lata* Dana, 1852.

Liomera laevis (A. Milne Edwards, 1873)

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Liomera laevis* - PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 212 (Moorea). — GUINOT, 1985: 451, with a ? (List).

Liomera laperousei Garth, 1985

DISTRIBUTION. — Austral (MacDonald bank) - *Littoral to sublittoral*.

REFERENCES. — *Liomera laperousei* - LABOUTE & RICHER DE FORGES, 1986: 21 (MacDonald bank, 40m; with hesitation: "semble être une femelle juvénile de *Liomera laperousei* Garth, 1985 décrit de l'île de Pâque").

Liomera monticulosa (A. Milne Edwards, 1873)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Marutea South) - *Littoral to sublittoral*.

REFERENCES. — *Carpilodes monticulosus* - NOBILI, 1907: 387 (Marutea South). — ODHNER, 1925: 21, pl. 1, fig. 18 (Tahiti, Marutea South). — *Liomera monticulosa* - FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1964: 11 (Syn.); 1985: 451 (List). — PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 212 (Moorea); 1989: 115 (Moorea; 30m). — *Liomera (Liomera) monticulosa* - SERÈNE, 1984: 64, fig. 24, pl. 6c (Syn.). — Not *Carpilodes monticulosus* - RATHBUN, 1907: 37 (Fakarava, Makemo) = *Liomera (Liomera) rugata* (H. Milne Edwards, 1834) *fide* SERÈNE (1984: 62).

Liomera pallida (Borradaile, 1900)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Carpilodes pallidus* - ODHNER, 1925: 20, pl. 1, fig. 17 (Tahiti). — *Liomera pallida* - FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 451, with a ? (List). — *Liomera (Liomera) pallida* - SERÈNE, 1984: 62, pl. 5f (Syn.).

Liomera rubra (A. Milne Edwards, 1865)

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Liomera rubra* - MONTEFORTE, 1984: 171, annex 1, tab. a; 1987: 9 (Moorea). — GUINOT, 1985: 451, with a ? (List). — *Liomera (Liomera) rubra* - SERÈNE, 1984: 65, fig. 26, pl. 6e-f, pl. 9f (Syn.).

Liomera rugata (H. Milne Edwards, 1834)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Moorea, Tahiti); Tuamotu (Fakarava, Makemo, Mataiva, Raroia, Tikehau).

REFERENCES. — *Carpilodes rugatus* - ORTMANN, 1893b: 468 (Tahiti). — ODHNER, 1925: 20, pl. 1, fig. 16 (Fakarava, Tahiti). — BOONE, 1934: 91, pl. 46 (Nuku Hiva). — HOLTHUIS, 1953: 13 (Raroia). — MORRISON, 1954: 16 (Raroia). — BUITENDIJK, 1960: 259, fig. 2d (Tahiti). — *Carpilodes monticulosus* - RATHBUN, 1907: 37 (Fakarava, Makemo) not *Liomera monticulosa* (A. Milne Edwards, 1873) = *L. rugata* fide SERÈNE (1984: 62). — *Liomera rugata* - FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva, Moorea); 1987: 9 (Moorea). — GUINOT, 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 113, 115 (Moorea, Tikehau). — *Liomera (Liomera) rugata* - SERÈNE, 1984: 62, fig. 22, pl. 6b (Syn.).

Liomera semigranosa De Man, 1888

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Liomera semigranulosa* (*sic*) - ODINETZ, 1983: 209 (Tahiti). — *Liomera semigranosa* - GUINOT, 1985: 451 (List). — *Liomera (Liomera) semigranosa* - SERÈNE, 1984: 63, pl. 7c, f (Syn.).

Liomera stimpsoni (A. Milne Edwards, 1865)

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Liomera stimpsoni* - PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 212; 1989: 115 (Moorea). — GUINOT, 1985: 451, with a ? (List).

Liomera tristis (Dana, 1852)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Fakarava).

REFERENCES. — *Carpilodes tristis* Dana, 1852a: 77; 1852b: 193; 1855, pl. 9, fig. 7a-d (Tuamotu with a ?). — HELLER, 1865: 17 (Tahiti). — A. MILNE EDWARDS, 1865b: 225 (Tuamotu). — DE MAN, 1890: 50 (Tahiti). — RATHBUN, 1907: 37 (Fakarava). — ODHNER, 1925: 12, pl. 1, fig. 1 ("Eimeo" = Moorea, Tahiti). — BUITENDIJK, 1960: 254, fig. 1c (Tahiti). — *Carpilodes granulatus* Heller, 1862: 520 (Tahiti). — *Liomera tristis* - FOREST & GUINOT, 1961: 38 (Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 451 (List). — *Liomera (Liomera) tristis* - SERÈNE, 1984: 59, fig. 19, pl. 5b (Syn.).

Liomera venosa (H. Milne Edwards, 1834)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Carpilodes venosus* - ORTMANN, 1893b: 467 (Tahiti). — ODHNER, 1925: 22, pl. 2, fig. 1 (Tahiti). — BUITENDIJK, 1960: 259, fig. 2e (Tahiti). — *Liomera venosa* - FOREST & GUINOT, 1962: 60 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 451 (List). — *Liomera (Liomera) venosa* - SERÈNE, 1984: 58, fig. 18, pl. 7d-e (Syn.).

Neoliomera demani Forest & Guinot, 1961

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hikueru).

REFERENCES. — *Neoliomera demani* Forest & Guinot, 1961: 80, fig. 76, 77bis, pl. 3, fig. 3-5 (Hikueru, Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — SERÈNE, 1984: 71, fig. 31, pl. 8b (Hikueru). — GUINOT, 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 115 (Moorea). — *Neoliomera pubescens* - ODHNER, 1925: 28, pl. 2, fig. 6, 6a-7 (Tahiti) not *Neoliomera pubescens* (H. Milne Edwards, 1834) = *N. demani* sp. nov. in FOREST & GUINOT (1961: 80).

Neoliomera insularis (White, 1847)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Neoliomera insularis* - SAKAI, 1976: 398 (Tahiti; cf. Remark). — GUINOT, 1985: 451 (List; after SAKAI).

REMARK. — The only reference in French Polynesia seems to be in SAKAI (1976) where "Tahiti" is cited without material examined from that island, and without older references for that location. Thus, the occurrence of this species in the French Polynesia still remains to be confirmed.

***Neoliomera pubescens* (H. Milne Edwards, 1834)**

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Neoliomera pubescens* - PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 212 (Moorea). — SERÈNE, 1984: 71, fig. 30, pl. 8a (Syn.; cf. Remark).

REMARK. — According to SERÈNE (1984) it is, in most of the works, necessary to check that the material attributed to *Neoliomera pubescens* has not been confounded with *N. demani* FOREST & GUINOT, 1961. He also mentions that *N. pubescens* is known, with certainty, only from Mauritius.

***Neoliomera richtersi* (De Man, 1889)**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Marutea South).

REFERENCES. — *Actaeodes richtersii* de Man 1889: 412, pl. 9, fig. 2; 1890: 51 (Tahiti). — *Liomera richtersi* - NOBILI, 1907: 387 ("Marutea-Vaitutaki" = Marutea South). — *Neoliomera richtersii* - ODHNER, 1925: 33, pl. 2, fig. 13 (Tahiti). — BUITENDIJK, 1960: 262 (Tahiti). — FOREST & GUINOT, 1961: 79, fig. 74 (Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1964: 47, fig. 17 (Tahiti); 1985: 451 (List). — SERÈNE, 1984: 70, fig. 28, pl. 8e (Tahiti; Syn.).

***Neoliomera variolosa* (A. Milne Edwards, 1873)**

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Neoliomera variolosa* - PEYROT-CLAUSADE, 1977b: 212, 220 (Moorea).

SUBFAMILY EUXANTHINAE

***Alainodaeus rimatara* Davie, 1993**

DISTRIBUTION. — Austral (Raevavae, Rimatara); Tuamotu (Akiaki, Fangataufa, Hao, Takapoto) - Sublittoral to deep.

REFERENCES. — *Alainodaeus rimatara* Davie, 1993: 519, fig. 6, pl. 6 (Akiaki, Fangataufa, Hao, Raevavae, Rimatara, Takapoto; 90-350m). — POUPIN, 1996: in press (same material).

***Euxanthus exsculptus* (Herbst, 1790)**

DISTRIBUTION. — Gambier (Mangareva); Society (Moorea, Tahiti).

REFERENCES. — *Euxanthus exsculptus* var. *rugosus* - NOBILI, 1907: 389 (Mangareva) not *Euxanthus rugosus* Miers, 1884 = *E. exsculptus* fide GUINOT-DUMORTIER (1960b: 170). — *Euxanthus exsculptus* - GUINOT-DUMORTIER, 1960b: 169, pl. 1, fig. 4, pl. 2, fig. 10, pl. 6, fig. 36-37, pl. 8, fig. 42-47 ("Rikitea" = Mangareva). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 170, annex 1, tab. a; 1987: 9 (Moorea). — SERÈNE, 1984: 86, fig. 48, pl. 11b (Tahiti). — GUINOT, 1985: 450 (List).

***Euxanthus sculptilis* Dana, 1852**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Euxanthus sculptilis* - BOONE, 1934: 107, pl. 57 (Tahiti). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — GUINOT-DUMORTIER, 1960b: 167, pl. 6, fig. 39, pl. 9, fig. 49 (Syn.); 1985: 450 (List).

***Medaeus grandis* Davie, 1993**

DISTRIBUTION. — Tuamotu (Hao, Moruroa) - Sublittoral to deep.

REFERENCES. — *Medaeus grandis* Davie, 1993: 526, fig. 8, pl. 8 (Hao, Moruroa; 90-210m). — POUPIN, 1996: in press (same material).

***Paramedaeus noelensis* (Ward, 1934)**

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Medaeus noelensis* - FOREST & GUINOT, 1961: 56, fig. 42-43, 44a-b, pl. 1, fig. 1 (Tahiti); 1962: 62 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a (Moorea, Tahiti); 1987: 9 (Moorea). — *Paramedaeus noelensis* - SERÈNE, 1984: 90, fig. 51, pl. 12f (Syn.). — GUINOT, 1985: 451 (List).

Paramedaeus simplex (A. Milne Edwards, 1873)

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Paramedaeus simplex* - PEYROT-CLAUSADE, 1977b: 212 (Moorea). — GUINOT, 1985: 451, with a ? (List).

SUBFAMILY ACTAEINAE

Actaea aff. *glandifera* Rathbun, 1914

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Actaea* aff. *glandifera* - PEYROT-CLAUSADE, 1989: 111, 114 (Moorea, Tikehau; 25m).

Actaea calculosa (H. Milne Edwards, 1834)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Actaea calculosa* - ODHNER, 1925: 52 (Tahiti). — FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu").

REMARK. — ODHNER mentions a Tahitian specimen in the collections of Hamburg. However, GUINOT (1976: 215-216) states that most of the references to *Actaea calculosa*, especially ODHNER (1925: 52), must be re-examined.

Actaea danae A. Milne Edwards, 1865

DISTRIBUTION. — Tuamotu (Raraka).

REFERENCES. — *Actaeodes areolatus* Dana, 1852a: 77; 1852b: 194; 1855, pl. 9, fig. 8a-d (Raraka). — *Actaea danae* - FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — (*Actaea*) *danae* - GUINOT, 1976: 247 (Syn.); 1985: 450, with "species inquirenda" (List).

Actaea polyacantha (Heller, 1861)

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Actaea polyacantha* - PEYROT-CLAUSADE, 1989: 114 (Moorea).

Actaeodes consobrinus (A. Milne Edwards, 1873)

DISTRIBUTION. — Marquesas; Society (Moorea) - *Littoral to sublittoral*.

REFERENCES. — *Actaea consobrina* - ODHNER, 1925: 67, pl. 4, fig. 14 (Marquesas). — *Actaeodes consobrinus* - GUINOT, 1976: 246, pl. 15, 5a (Syn.); 1985: 450, with a ? (List). — *Actaeodes consobrina* - PEYROT-CLAUSADE, 1989: 114 (Moorea; 30m). — Not *Actaea consobrina* - NOBILI, 1907: 390 = *Actaea ruppellioides* sp. nov. in ODHNER (1925: 47; cf. under *Pseudoliomera ruppellioides*).

Actaeodes hirsutissimus (Rüppell, 1830)

DISTRIBUTION. — Society (Bora Bora, Moorea, Raiatea, Tahiti); Tuamotu (Mataiva, Tikehau).

REFERENCES. — *Actaea hirsutissima* - HELLER, 1865: 9 (Tahiti). — RATHBUN, 1907: 42 (Bora Bora, Tahiti). — ODHNER, 1925: 69, pl. 4, fig. 13 (Tahiti). — BOONE, 1934: 124, pl. 66 (Raiatea). — FOREST & GUINOT, 1961: 78 (Tahiti); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1989: 111 (Tikehau). — *Actaeodes hirsutissimus* - GUINOT, 1976: 245, fig. 38e, pl. 15, fig. 2, 2a (Tahiti; Syn.); 1985: 450 (List). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213; 1985: 462 (Moorea). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva, Moorea, Tahiti); 1987: 8 (Moorea). — SERÈNE, 1984: 135 (Syn.).

***Actaeodes tomentosus* (H. Milne Edwards, 1834)**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Makatea).

REFERENCES. — *Actaeodes tomentosus* - HELLER, 1865: 17 (Tahiti). — GUINOT, 1976: 244, fig. 38d, 41c, pl. 15, fig. 1, 1a (Syn.); 1985: 450, with a ? (List). — SERÈNE, 1984: 134, 137 (Syn.). — *Actaea tomentosa* - SENDLER, 1923: 37 (Makatea). — FOREST & GUINOT, 1962: 66 (Biogeography "Tahiti-Tuamotu").

***Forestia depressa* (White, 1847)**

DISTRIBUTION. — Marquesas.

REFERENCES. — *Actaea depressa* - BALSS, 1935: 136 (Marquesas). — *Forestia depressa* - GUINOT, 1976: 262 (Syn.; with a ? for BALSS' reference); 1985: 450 (List; with a ? in front of "Marquesas"). — SERÈNE, 1984: 106 (Syn.; with a ? in front of BALSS' reference).

REMARK. — In GUINOT (1976) and SERÈNE (1984), at least one specimen attributed to *Actaea depressa*, by BALSS (1938: 54), would be in fact a *Forestia scabra* (Odhner, 1925).

***Forestia scabra* (Odhner, 1925)**

DISTRIBUTION. — Marquesas.

REFERENCES. — *Actaea scabra* - BALSS, 1935: 136 (Marquesas). — *Forestia scabra* - GUINOT, 1976: 263 (Syn.; with a ? in front of BALSS' reference); 1985: 450 (List; with a ? in front of "Marquesas").

***Gaillardiellus rueppelli* (Kraus, 1843)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Gaillardiellus rueppelli* - GARTH & KIM, 1983: 684 (Distribution only, Tahiti; cf. Remark). — GUINOT, 1985: 450 (List; presumably after the previous work).

REMARK. — The material examined by GARTH & KIM (1983: 685) was collected in the Philippines by the *Albatross* (1908-1909). "Tahiti", mentioned in the "Distribution" only, corresponds neither to material examined, nor to former references cited in this work.

***Gaillardiellus superciliaris* (Odhner, 1925)**

DISTRIBUTION. — Tuamotu (Raroia, Taiaro).

REFERENCES. — *Actaea superciliaris* - HOLTHUIS, 1953: 11 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — *Gaillardiellus superciliaris* - GUINOT, 1976: 257 (Syn.); 1985: 450 (List). — POUPIN, 1994a: 34, fig. 30, pl. 4a, with a ? (Taiaro).

***Paractaea excentrica* Guinot, 1969**

DISTRIBUTION. — Tuamotu (Marutea South).

REFERENCES. — *Paractaea excentrica* Guinot, 1969: 263, fig. 36 (Marutea South); 1985: 451 (List).

***Paractaea retusa* (Nobili, 1905)**

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Raroia).

REFERENCES. — *Actaea garretti* Rathbun, 1906: 852, pl. 9, fig. 8 (Society). — *Actaea rufopunctata* - HOLTHUIS, 1953: 11 (Raroia). — MORRISON, 1954: 16 (Raroia). — FOREST & GUINOT, 1961: 79, fig. 79a-b (Tahiti) - All, not *Paractaea rufopunctata* (H. Milne Edwards, 1834) = *Paractaea retusa* form *hippocrepica* nov. in GUINOT (1969: 256). — *Paractaea retusa* form *hippocrepica* - GUINOT, 1969: 256, fig. 30 (Tahiti, Raroia; Syn.); 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 115 (Moorea). — *Paractaea retusa* - MONTEFORTE, 1984: 171, annex 1, tab. a; 1987: 9 (Moorea).

REMARK. — GUINOT (1969: 255), when creating the new genus *Paractaea*, has examined the syntype of *garretti* from Gilbert Islands (but not the specimen from the Society Islands), and considers that RATHBUN's species belongs to *Paractaea retusa* (Nobili) form *garretti* (Rathbun).

***Paractaea rufopunctata* H. Milne Edwards, 1834**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Makemo, Marutea South).

REFERENCES. — *Actaea rufopunctata* - NOBILI, 1907: 392 (Marutea). — RATHBUN, 1907: 43 (Makemo, Tahiti). — ODHNER, 1925: 60 (Tahiti). — FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — *Paractaea rufopunctata* form *plumosa* - GUINOT, 1969: 248, fig. 21 (Marutea). — *Paractaea rufopunctata* - GUINOT, 1985: 451 (List).

***Paractaeopsis quadriareolatus* (Takeda & Miyake, 1968)**

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Paractaeopsis quadriareolatus* - SERÈNE, 1984: 127 (Syn.; gen. nov.). — *Paractaea quadriareolata* - PEYROT-CLAUSADE, 1989: 115 (Moorea).

***Paractaeopsis tumulosus* (Odhner, 1925)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Actaea tumulosa* Odhner, 1925: 61, pl. 4, fig. 10 (Tahiti). — FOREST & GUINOT, 1962: 66 (Biogeography "Tahiti-Tuamotu"). — *Paractaeopsis tumulosus* - SERÈNE, 1984: 127, fig. 74, pl. 17d (Syn.; gen. nov.). — *Paractaea tumulosa* - GUINOT, 1985: 451 (List).

***Psaumis cavipes* (Dana, 1852)**

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fakarava, Hao, Rangiroa, Raroia, Tikehau).

REFERENCES. — *Actaea cavipes* - NOBILI, 1907: 390 ("Ohura" = Hao, "Rikitea" = Mangareva). — RATHBUN, 1907: 44, pl. 1, fig. 2 (Bora Bora, Fakarava, Rangiroa). — ODHNER, 1925: 68 ("Eimeo" = Moorea, Tahiti). — BOONE, 1934: 128, pl. 68 (Tahiti). — HOLTHUIS, 1953: 10 (Raroia). — MORRISON, 1954: 16 (Raroia). — FOREST & GUINOT, 1961: 78 (Tahiti); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 212 (Moorea). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a (Moorea, Tahiti); 1987: 8 (Moorea). — *Psaumis cavipes* - ODINETZ, 1983: 209 (Moorea, Tahiti). — SERÈNE, 1984: 129, fig. 76, pl. 18f (Syn.). — GUINOT, 1985: 451 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti, Takapoto). — MONTEFORTE, 1987: 9 (Moorea). — PEYROT-CLAUSADE, 1989: 113 (Tikehau).

***Psaumis cellulosa* (Dana, 1852)**

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Psaumis cellulosa* - ODINETZ, 1983: 209 (Moorea, Tahiti). — GUINOT, 1985: 451 (List).

***Pseudoliomera granosimana* (A. Milne Edwards, 1865)**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Marutea South).

REFERENCES. — *Liomeria granosimana* - ORTMANN, 1893b: 451 (Tahiti). — NOBILI, 1907: 387 ("Marutea-Vaitutaki" = Marutea South). — *Pseudoliomera granosimana* - ODHNER, 1925: 79, fig. 5-6 (Tahiti). — FOREST & GUINOT, 1961: 39, fig. 28a-c, pl. 7, fig. 1-2 (Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — SERÈNE, 1984: 100, fig. 56, pl. 13a (Tahiti). — GUINOT, 1985: 451 (List).

***Pseudoliomera lata* (Borradaile, 1902)**

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Pseudoliomera lata* - KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea). — SERÈNE, 1984: 102 (Syn.). — PEYROT-CLAUSADE, 1989: 115 (Moorea). — (*Pseudoliomera*) *lata* - GUINOT, 1985: 451, with a ? (List). — Not *Actaea lata* - NOBILI, 1907: 392 ("Marutea, Vaitutaki" = Marutea South) = *A. ruppellioides* Odhner, 1925 *fide* GUINOT (1962: 237).

Pseudoliomera ruppellioides (Odhner, 1925)

DISTRIBUTION. — Tuamotu (Marutea South).

REFERENCES. — *Actaea consobrina* - NOBILI, 1907: 390 (Marutea South) not *Actaea consobrina* A. Milne Edwards, 1873 = *A. ruppellioides* sp. nov. in ODHNER (1925: 47). — *Actaea lata* - NOBILI, 1907: 392 ("Marutea, Vaitutaki" = Marutea South) not *Actaea lata* Borradaile, 1902 = *A. ruppellioides* Odhner *fide* GUINOT (1962: 237). — *Actaea ruppellioides* Odhner, 1925: 47, pl. 3, fig. 9 (Marutea South; NOBILI's material). — FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — (*Pseudoliomera*) *ruppellioides* - GUINOT, 1976: 203, 246 (Genus *Pseudoliomera* "ou à sa proximité"); 1985: 451 (List).

Pseudoliomera speciosa (Dana, 1852)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Actaea speciosa* - SENDLER, 1923: 38 (Tahiti). — ODHNER, 1925: 62 (Tahiti). — FOREST & GUINOT, 1962: 66 (Biogeography "Tahiti-Tuamotu"). — (*Pseudoliomera*) *speciosa* - GUINOT, 1976: 203, 243 (Genus uncertain); 1985: 451, with a ? (List).

Pseudoliomera variolosa (Borradaile, 1902)

DISTRIBUTION. — Society (Moorea); Tuamotu (Mataiva, Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Pseudoliomera variolosa* - KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva, Moorea; cf. Remark); 1987: 9 (Moorea). — PEYROT-CLAUSADE, 1989: 111, 115 (Moorea, Tikehau; 30m). — Aff. *Pseudoliomera variolosa* - PEYROT-CLAUSADE, 1977a, annex of the species: 27 (Moorea). — (*Pseudoliomera*) *variolosa* - GUINOT, 1985: 451 (List).

REMARK. — MONTEFORTE writes, *Pseudoliomera variolosa* (A. Milne Edwards, 1837 *sic*). Then, it could be *Neoliomera variolosa* (A. Milne Edwards, 1873), which is different from BORRADAILE's species (cf. SERÈNE, 1984: 66).

SUBFAMILY ZOZIMINAE

Atergatis floridus (Linné, 1767)

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Hao).

REFERENCES. — *Atergatis floridus* - DANA, 1852b: 159; 1855, pl. 7, fig. 4 (Society and Tuamotu). — HELLER, 1865: 8 (Tahiti). — NOBILI, 1907: 388 ("Ohura" = Hao). — SEURAT, 1934: 59 (Hao). — BUTENDIJK, 1960: 268 (Society). — FOREST & GUINOT, 1961: 41 (Tahiti; Syn.); 1962: 62 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 212 (Moorea). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a (Moorea, Tahiti); 1987: 8 (Moorea). — GUINOT, 1985: 450 (List). — POUPIN, 1994a: 35, fig. 31, pl. 4b (Mangareva, Tahiti). — *Atergatis ocyroe* - RATHBUN, 1907: 37 (Bora Bora). — SENDLER, 1923: 37 (Tahiti) - SYNONYMS - *Atergatis ocyroe* (Herbst, 1801).

Atergatopsis cf. germaini A. Milne Edwards, 1865

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Atergatopsis cf. germaini* - MONTEFORTE, 1984: 170, annex 1, tab. a; 1987: 9 (Moorea). — GUINOT, 1985: 450 (List).

Atergatopsis signatus (Adams & White, 1848)

DISTRIBUTION. — Tuamotu (Makatea, Mataiva, Raroia, Takapoto).

REFERENCES. — *Atergatopsis signatus* - HOLTHUIS, 1953: 12 (Raroia). — MORRISON, 1954: 16 (Raroia). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (Raroia); 1985: 450 (List). — MONTEFORTE, 1984: 170, annex 1, tab. a (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — SALVAT, 1986b: 72 (French Polynesia).

***Lophozozymus cristatus* A. Milne Edwards, 1867**

DISTRIBUTION. — Austral (Maria); Society.

REFERENCES. — *Lophozozymus cristatus* - BUITENDIJK, 1960: 292, fig. 7a (Society). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — POUPIN, 1994a: 36, fig. 32, pl. 4c (Maria).

***Lophozozymus dodone* (Herbst, 1801)**

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Atergatis elegans* Heller, 1862: 519; 1865: 7, pl. 1, fig. 3 (Tahiti). — *Lophozozymus dodone* - BALSS, 1938: 39 (Tahiti). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a (Moorea, Tahiti); 1987: 9 (Moorea). — SERÈNE, 1984: 171, pl. 24e (Syn.). — GUINOT, 1985: 451 (List). — Not *Lophozozymus dodone* - FOREST & GUINOT, 1961: 54, fig. 39a-b (Tahiti) = *Lophozozymus glaber* fide GUINOT (1979: 65).

***Lophozozymus edwardsi* Odhner, 1925**

DISTRIBUTION. — Gambier (Mangareva); Society (Moorea); Tuamotu (Marutea South).

REFERENCES. — *Lophozozymus superbus* - NOBILI, 1907: 388 (Mangareva, "Marutea Vaitutaki" = Marutea South) not *Lophozozymus superbus* A. Milne Edwards, 1873 = *L. edwardsi* fide FOREST & GUINOT (1961: 56) & GUINOT (1979: 63). — *Lophozozymus edwardsi* - FOREST & GUINOT, 1961: 56, fig. 41 (Mangareva); 1962: 62 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1979: 63 (Mangareva, Marutea South); 1985 (List). — MONTEFORTE, 1984: 171, annex 1, tab. a; 1987: 9 (Moorea).

***Lophozozymus glaber* Ortmann, 1893**

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Lophozozymus glaber* - GUINOT, 1979: 65, pl. 8, fig. 2, 2a (Tahiti). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — MONTEFORTE, 1984: 171, annex 1, tab. a; 1987: 9 (Moorea). — *Lophozozymus dodone* - FOREST & GUINOT, 1961: 54, fig. 39a-b (Tahiti) not *Lophozozymus dodone* (Herbst, 1801) = *L. glaber* fide GUINOT (1979: 65).

***Lophozozymus pictor* (Fabricius, 1798)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Lophozozymus octodentatus* - BOONE, 1934: 102, pl. 54-55 (Tahiti). — SAKAI, 1976: 407, pl. 146, fig. 3 (Syn., but without BOONE's reference) - SYNONYMS - *Lophozozymus octodentatus* (H. Milne Edwards, 1834).

REMARK. — BOONE has examined and illustrated a large male and a female from Tahiti. However, it seems that this material has never been re-examined (cf. BUITENDIJK, 1960, or SAKAI, 1976).

***Lophozozymus superbus* (Dana, 1852)**

DISTRIBUTION. — Tuamotu (Moruroa, Raraka, Raroia).

REFERENCES. — *Xantho superbus* Dana, 1852a: 74; 1852b: 167; 1855, pl. 8, fig. 5a-b (Raraka). — *Lophozozymus superbus* - HOLTHUIS, 1953: 23 (Raroia). — MORRISON, 1954: 13 (Raroia). — GUINOT, 1979: 63 (Moruroa); 1985: 451 (List). — *Lophozozymus incisus* - FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu") not *Lophozozymus incisus* (H. Milne Edwards, 1834) = *L. superbus* (Dana) (cf. Remark). — Not *Lophozozymus superbus* - NOBILI, 1907: 388 = *L. edwardsi* fide FOREST & GUINOT (1961: 56) & GUINOT (1979: 63).

REMARK. — FOREST & GUINOT (1962: 62) have recorded *Lophozozymus incisus* (H. Milne Edwards, 1834) in French Polynesia because it has formerly been considered as a synonym of *L. superbus* (Dana). This opinion has been changed later on (see for example HOLTHUIS, 1953, or GUINOT, 1979).

***Platypodia anaglypta* (Heller, 1861)**

DISTRIBUTION. — Society (Moorea); Tuamotu (Fakarava, Tikehau).

REFERENCES. — *Platypodia anaglypta* - RATHBUN, 1907: 38 (Fakarava). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 113, 115 (Moorea, Tikehau).

Platypodia granulosa (Rüppell, 1830)

DISTRIBUTION. — Society (Tahiti?); Tuamotu (Nukutipipi).

REFERENCES. — *Atergatis limbatus* - ? HELLER, 1865: 8 (Tahiti). — *Platypodia granulosa* - ? PESTA, 1913: 41 (Tahiti). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — SERÈNE, 1984: 159, 162 (Syn.). — GUINOT, 1985: 451 (List). — MERSCHARDT-SALVAT, 1991: 89 (Nukutipipi). — Not *Lophactea granulosa* - NOBILI, 1907: 388 ("Rikitea" = Mangareva, Marutea). — Not *Platypodia granulosa* - FOREST & GUINOT, 1961: 51 (Mangareva) - These two references = *Platypodia pseudogranulosa* sp. nov. in SERÈNE (1984: 159; cf. Remark) - SYNONYMS - *Atergatis limbatus* (H. Milne Edwards, 1834).

REMARK. — SERÈNE (1984: 159), for the description of *Platypodia pseudogranulosa*, closely related to *P. granulosa*, does not mention HELLER's (1865) and PESTA's (1913) references, neither under *P. granulosa*, nor under *P. pseudogranulosa*. Thus we place these two references under both species, with uncertainty.

Platypodia pseudogranulosa Serène, 1984

DISTRIBUTION. — Gambier (Mangareva); Society (Tahiti?); Tuamotu (Marutea South).

REFERENCES. — *Atergatis limbatus* - ? HELLER, 1865: 8 (Tahiti; cf. Remark under *P. granulosa*). — *Lophactaea granulosa* (Rüpp.) - NOBILI, 1907: 388 ("Rikitea" = Mangareva, Marutea). — *Platypodia granulosa* - ? PESTA, 1913: 41 (Tahiti; cf. Remark under *P. granulosa*). — FOREST & GUINOT, 1961: 51 (Mangareva) - NOBILI's and FOREST & GUINOT's references, not *Platypodia granulosa* (Rüppell, 1830) = *P. pseudogranulosa* sp. nov. in SERÈNE (1984: 159). — *Platypodia pseudogranulosa* Serène, 1984: 159, pl. 22d ("Rikitea" = Mangareva, Marutea; Syn.).

Platypodia semigranosa (Heller, 1861)

DISTRIBUTION. — Society (Moorea) - Littoral to sublittoral.

REFERENCES. — *Platypodia semigranosa* - SERÈNE, 1984: 160, fig. 95, pl. 22b (Syn.; with material collected by PEYROT-CLAUSADE at Madagascar). — PEYROT-CLAUSADE, 1989: 115 (Moorea; 22m).

Zozimus aeneus (Linné, 1758)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Tahiti); Tuamotu (Fakahina, Hikueru, Makatea, Makemo, Mataiva, Moruroa, Raroia, Taiaro, Takapoto).

REFERENCES. — *Zozymus aeneus* - DANA, 1852b: 192; 1855, pl. 10, fig. 3a (Tuamotu). — STIMPSON, 1858a: 32 [30]; 1907: 42 (Tahiti). — NOBILI, 1907: 388 (Fakahina). — SEURAT, 1934: 59 (Fakahina, Tahiti). — MONTEFORTE, 1984: 171, annex 1, tab. a, photograph p. 136b (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — SALVAT & RICHARD, 1985: 360 (Takapoto). — SALVAT, 1986b: 72 (French Polynesia). — *Zozimus aeneus* - RATHBUN, 1907: 38 (Makemo). — BOONE, 1934: 99, pl. 50-53 (Nuku Hiva, Tahiti). — HOLTHUIS, 1953: 27 (Raroia). — MORRISON, 1954: 16 (Raroia). — BUTTENDIJK, 1960: 284, fig. 6a (Society). — FOREST & GUINOT, 1961: 51 (Hikueru, Tahiti); 1962: 62 (Biogeography "Tahiti-Tuamotu, Marquesas"). — GUINOT, 1966a: 48 (Raroia); 1985: 451 (List). — BONVALLOT *et al.*, 1994: 140-141, photograph (Tuamotu). — POUPIN, 1994a: 37, fig. 33, pl. 4d (Tahiti, Taiaro). — ? *Zozimus* s. p. - SALVAT, 1986a: 19, photograph (French Polynesia; det. according to the photograph). — ? *Lophozozymus* sp. - BAGNIS & CHRISTIAN, 1983: 110, photograph (Tuamotu; det. according to the photograph) - NEW MATERIAL - Coll. and det. J. POUPIN (Moruroa).

Zozymodes pumilus (Jacquinot, 1852)

DISTRIBUTION. — Gambier?; Society (Tahiti); Tuamotu (Hikueru, Makemo).

REFERENCES. — *Xanthodius cristatus* - RATHBUN, 1907: 41 (Makemo). — *Zozymodes carinipes* - NOBILI, 1907: 388 ("Tagatau" = Gambier?) not *Zozymodes carinipes* Heller, 1861 synonym of *Z. xanthoides* (Krauss, 1843) = *Z. pumilus* fide FOREST & GUINOT (1961: 52). — *Zozymodes pumilus* - FOREST & GUINOT, 1961: 52, fig. 36a-b (Hikueru); 1962:

62 (Biogeography "Tahiti-Tuamotu"). — SERÈNE, 1984: 153, fig. 90, pl. 14e (Tahiti; Syn.). — GUINOT, 1985: 452 (List) - SYNONYMS - *Leptodius cristatus* Borradaile, 1902.

Zozymodes xanthoides (Krauss, 1843)

DISTRIBUTION. — Tuamotu (Takapoto, Tikehau).

REFERENCES. — *Zozymodes xanthoides* - MONTEFORTE, 1984: 171, annex 1, tab. a (Takapoto). — GUINOT, 1985: 452, with a ? (List). — *Zozymoïdes xanthoïdes* (sic) - PEYROT-CLAUSADE, 1989: 113 (Tikehau).

SUBFAMILY XANTHINAE

Lachnopus bidentatus (A. Milne Edwards, 1867)

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Xantho arcuatus* Heller, 1865: 11, pl. 2, fig. 1 (Tahiti). — *Lachnopus bidentatus* - FOREST & GUINOT, 1961: 42, fig. 29-30, 32bis, 33a-b, pl. 7, fig. 1-2 (Tahiti; Syn.); 1962: 62 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 212 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a; 1987: 9 (Moorea). — GUINOT, 1985: 250 (List).

Lachnopus ponapensis (Rathbun, 1907)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Xanthias ponapensis* Rathbun, 1907: 44, pl. 7, fig. 5, 5a (Tahiti). — *Paraxanthias ponapensis* - FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 451 (List). — *Lachnopus ponapensis* - SERÈNE, 1984: 203 (Key).

Lachnopus subacutus (Stimpson, 1858)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Mataiva).

REFERENCES. — *Lachnopus subacutus* - PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 212 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva, Moorea, Tahiti); 1987: 9 (Moorea). — SERÈNE, 1984: 204, fig. 122, pl. 29a (Syn.). — GUINOT, 1985: 450, with a ? (List).

Lachnopus tahitensis De Man, 1889

DISTRIBUTION. — Society (Tahiti); Tuamotu (Makatea, Raroia, Takapoto).

REFERENCES. — *Xantho (Lachnopus) tahitensis* de Man, 1889: 418, pl. 9, fig. 4, 4a; 1890: 52 (Tahiti). — *Lachnopus tahitensis* - HOLTHUIS, 1953: 22 (Raroia). — MORRISON, 1954: 16 (Raroia). — FOREST & GUINOT, 1961: 49 (Tahiti); 1962: 62 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (Raroia); 1985: 450 (List). — MONTEFORTE, 1984: 170, annex 1, tab. a (Makatea, Takapoto). — SERÈNE, 1984: 203, fig. 123, pl. 29d (Tahiti).

Leptodius davaoensis Ward, 1941

DISTRIBUTION. — Tuamotu (Hao, Hikueru, Mataiva, Moruroa, Takapoto).

REFERENCES. — *Leptodius leptodon* Forest & Guinot, 1961: 65, fig. 55-56, 59a-b, pl. 2, fig. 3 (Hikueru) *fide* TAKEDA (1980: 318); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva, Takapoto). — GUINOT, 1985: 450 (List). — *Leptodius exaratus* - NOBILI, 1907: 389 (Hao) not *Leptodius exaratus* (H. Milne Edwards, 1834) = *L. leptodon* nov. in FOREST & GUINOT (1961: 65). — *Leptodius davaoensis* - TAKEDA, 1980: 318 (Syn.). — POUPIN, 1994a: 38, fig. 34, pl. 4e (Hikueru, Moruroa) - RELEVANT MATERIAL - *Leptodius cf. davaoensis* - MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva). — GUINOT, 1985: 450 (List).

Leptodius exaratus (H. Milne Edwards, 1834)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society; Tuamotu (Raroia).

REFERENCES. — *Leptodius exaratus* - BOONE, 1934: 110, pl. 58 (Nuku Hiva). — FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu, ?Marquesas"). — SERÈNE, 1984: 183, fig. 106, pl. 26a (Syn.). — GUINOT, 1985: 450 (List). — *Xantho exaratus* - HOLTHUIS, 1953: 27 (Raroia). — MORRISON, 1954: 7 (Raroia). — BUITENDIJK, 1960: 331, fig. 9k-m (Society). — Not *Leptodius exaratus* (H. Milne Edwards) - NOBILI, 1907: 389 = *Leptodius leptodon* nov. in FOREST & GUINOT (1961: 65) synonym of *L. davaoensis* fide TAKEDA (1980: 318).

Leptodius gracilis (Dana, 1852)

DISTRIBUTION. — Gambier (Rikitea); Tuamotu (Hikueru, Moruroa, Rangiroa, Raroia).

REFERENCES. — *Leptodius gracilis* - NOBILI, 1907: 389 (Rikitea). — FOREST & GUINOT, 1961: 64, fig. 57, 58a-b, pl. 2, fig. 4 (Hikueru); 1962: 62 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 450 (List). — POUPIN, 1994a: 39, fig. 35, pl. 4f (Hikueru, Moruroa). — *Xantho gracilis* - HOLTHUIS, 1953: 27 (Raroia). — MORRISON, 1954: 7 (Raroia). — BUITENDIJK, 1960: 335 (Rangiroa).

Leptodius sanguineus (H. Milne Edwards, 1834)

DISTRIBUTION. — Gambier (Kamaka, Mangareva, Tarauru-Roa); Marquesas (Nuku Hiva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Ahe and/or Manihi, Fakarava, Makatea, Makemo, Marutea South, Mataiva, Rangiroa, Taiaro, Takapoto).

REFERENCES. — *Chlorodius sanguineus* - DANA, 1852a: 79; 1852b: 207; 1855, pl. 11, fig. 11a-d ("Waterland" = Ahe and/or Manihi). — *Leptodius sanguineus* - NOBILI, 1907: 389 ("Rikitea" = Mangareva, Kamaka, Marutea South). — RATHBUN, 1907: 39 (Bora Bora, Fakarava, Makatea, Makemo, Mangareva, Nuku Hiva, Rangiroa, Tahiti). — SENDLER, 1923: 37 (Makatea, Tahiti). — BOONE, 1934: 116, pl. 60-61 (Nuku Hiva). — FOREST & GUINOT, 1961: 63, fig. 50a-b ("Gatavake" = Mangareva, Tahiti, "Taraourou-roa" = Tarauru-Roa); 1962: 62 (Biogeography "Tahiti-Tuamotu, Marquesas"). — MONTEFORTE, 1984: 170, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — SERÈNE, 1984: 185 (Syn.). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List). — POUPIN, 1994a: 40, fig. 36, pl. 4g (Nuku Hiva, Tahiti, Taiaro). — *Xantho sanguineus* - BUITENDIJK, 1960: 323 (Nuku Hiva).

Lioxanthodes alcocki Calman, 1909

DISTRIBUTION. — Society (Moorea); Tuamotu (Makatea, Mataiva, Takapoto, Tikehau).

REFERENCES. — *Lioxanthodes alcocki* - MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Moorea, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 113 (Tikehau).

Macromedaeus crassimanus (A. Milne Edwards, 1867)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Xantho crassimanus* - BUITENDIJK, 1960: 318, fig. 9c-f (Tahiti). — *Leptodius crassimanus* - FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — *Macromedaeus crassimanus* - SERÈNE, 1984: 179, fig. 103, pl. 25b (Syn.). — GUINOT, 1985: 451 (List).

Macromedaeus distinguendus (de Haan, 1835)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Macromedaeus distinguendus* - KIM, 1973: 630 (Distribution only, Tahiti). — GUINOT, 1985: 451 (Society; List).

REMARK. — GUINOT (1985) has listed this species after FOREST & GUINOT (1961). These authors have actually examined some *Xantho distinguendus* de Haan, 1835 (p. 57, under *Medaeus noelensis* Ward, 1934), but they are from Hong Kong. It could be that GUINOT refers to KIM (1973), who has quoted "Tahiti" in the distribution of this species. However, this location concerns neither the material examined, nor one of the references cited by KIM under *Macromedaeus distinguendus*. The presence of this species in French Polynesia remains thus doubtful.

***Macromedaeus nudipes* (A. Milne Edwards, 1867)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Xantho nudipes* - FOREST & GUINOT, 1961: 59, fig. 47a-b. (Tahiti); 1962: 62 (Biogeography "Tahiti-Tuamotu"). — *Macromedaeus nudipes* - SERÈNE, 1984: 178, fig. 101, pl. 25a (Syn.). — GUINOT, 1985: 451 (List).

***Neoxanthops cavatus* (Rathbun, 1907)**

DISTRIBUTION. — Tuamotu (Fakarava).

REFERENCES. — *Cycloxoanthops cavatus* Rathbun, 1907: 41, pl. 5, fig. 8, pl. 6, fig. 3, 3a (Fakarava). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — *Neoxanthops cavatus* - SERÈNE, 1984: 212, fig. 128, pl. 29f (Syn.). — GUINOT, 1985: 451 (List).

***Paraxanthias notatus* (Dana, 1852)**

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Fakarava, Makatea, Makemo, Marutea North?, Marutea South, Mataiva, Takapoto, Tikehau).

REFERENCES. — *Xanthodes notatus* Dana, 1852a: 76 (Tahiti, Tuamotu); 1852b: 178; 1855, pl. 8, fig. 12a-b. (Society or Tuamotu). — *Xanthias notatus* - NOBILI, 1907: 392 (Fakarava, Makatea, Marutea = Marutea North?, Marutea South). — RATHBUN, 1907: 45 (Fakarava, Makemo). — *Paraxanthias notatus* - FOREST & GUINOT, 1961: 76, fig. 70a-b (Tahiti); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 113, 115 (Moorea, Tikehau).

***Xanthias canaliculatus* Rathbun, 1907**

DISTRIBUTION. — Society (Moorea); Tuamotu (Makemo).

REFERENCES. — *Xanthias canaliculatus* Rathbun, 1907: 45 (Makemo). — FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 212 (Moorea). — GUINOT, 1985: 451 (List).

***Xanthias lamarcki* (H. Milne Edwards, 1834)**

DISTRIBUTION. — Gambier (Tarauru-Roa, Temoe); Society (Bora Bora, Moorea, Raiatea, Tahiti); Tuamotu (Fakarava, Hikueru, Makatea, Makemo, Marutea South, Mataiva, Raroia, Takapoto, Tikehau).

REFERENCES. — *Xanthodes granoso-manus* Dana, 1852a: 75; 1852b: 175; 1855, pl. 8, fig. 10a-c (Society, Tuamotu). — *Xanthias lamarckii* - NOBILI, 1907: 393 (Hikueru, "Timoe" = Temoe). — RATHBUN, 1907: 44 (Bora Bora, Fakarava, Makemo, Tahiti). — SENDLER, 1923: 37 (Makatea). — BOONE, 1934: 131, pl. 70 (Raiatea, Tahiti). — HOLTHUIS, 1953: 26 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1961: 70, fig. 63, 66a-b (Hikueru, Marutea, Tahiti, "Taraourou-roa" = Tarauru-Roa); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 212 (Moorea); 1989: 113 (Tikehau). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — SERÈNE, 1984: 195, fig. 112, pl. 27b (Syn.). — DELESALLE, 1985: 305 (Mataiva). — GUINOT, 1985: 451 (List).

***Xanthias latifrons* (De Man, 1888)**

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Xanthias latifrons* - FOREST & GUINOT, 1961: 70, fig. 67a-b (Tahiti); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a; 1987: 9 (Moorea). — GUINOT, 1985: 451 (List).

***Xanthias nitidulus* (Dana, 1852)**

DISTRIBUTION. — Tuamotu (Marutea South).

REFERENCES. — *Xanthodes nitidulus* Dana, 1852a: 76; 1852b: 177; 1855, pl. 8, fig. 11a-c (Tuamotu). — *Xanthias nitidulus* - NOBILI, 1907: 392 (Marutea South).

REMARK. — FOREST & GUINOT (1961: 72) have placed these two references under *Xanthias tetaodon*, with a ? and this remark "Nous nous abstiendrons pour l'instant de tirer des conclusions définitives, mais il est probable que lorsqu'on disposera d'une série de *Xanthias tetaodon* de diverses tailles, l'on sera amené à désigner cette espèce sous le nom de *Xanthias nitidilus* (Dana)".

Xanthias punctatus (H. Milne Edwards, 1834)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Xanthias punctatus* - FOREST & GUINOT, 1961: 68, fig. 61, 65a-b (Tahiti); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 451 (List).

Xanthias tetaodon (Heller, 1865)

DISTRIBUTION. — Gambier (Mangareva); Society (Tahiti); Tuamotu (Hao, Hikueru, Makatea, Marutea South?, Mataiva, Raroia, Takapoto).

REFERENCES. — *Eudora tetaodon* Heller, 1865: 14, pl. 2, fig. 3 (Auckland = ? Tahiti in FOREST & GUINOT, 1961). — *Xantho (Eudora) tetaodon* - NOBILI, 1907: 389 (Hao). — *Juxtaxanthias tetaodon* - WARD, 1942: 92 (Mangareva). — HOLTHUIS, 1953: 22 (Raroia). — MORRISON, 1954: 16 (Raroia). — *Xanthias tetaodon* - FOREST & GUINOT, 1961: 72, fig. 61, 68a-c, 69bis (Hikueru, Tahiti); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (Raroia); 1985: 451 (List). — MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva).

REMARK. — ODHNER (1925) and FOREST & GUINOT (1961) consider that the type locality, Auckland, mentioned by HELLER (1865) for the description of *Eudora tetaodon*, is a mistake, and that it could very likely be Tahiti. Moreover, FOREST & GUINOT (1961) mention that this species could be a synonym of *Xanthias nitidilus* (Dana, 1852) (cf. above).

SUBFAMILY PANOPINEAE

Panopeus pacificus Edmondson, 1931

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Panopeus pacificus* - FOREST & GUINOT, 1961: 116, fig. 102, 103a-b, 104, 105a-b, pl. 4, fig. 3 (Tahiti); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List).

SUBFAMILY KRAUSSIINAE

Palapedia marquesas (Serène, 1972)

DISTRIBUTION. — Tuamotu (Anaa).

REFERENCES. — *Kraussia marquesas* Serène, 1972: 53, fig. 14-15, 23g, k (Anaa). — GUINOT, 1985: 449 (List). — *Palapedia marquesas* - NG, 1993: 141 (subfamily nov. and gen. nov.).

Palapedia rastripes (Müller, 1887)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Kraussia rastripes* - MONTEFORTE, 1984: 171, annex 1, tab. a (Tahiti). — GUINOT, 1985: 449 (List). — *Palapedia rastripes* - NG, 1993: 141 (subfamily nov. and gen. nov.).

SUBFAMILY ETISINAE

Etisus bifrontalis (Edmonson, 1935)

DISTRIBUTION. — Tuamotu (Hikueru).

REFERENCES. — *Etisodes electra* - NOBILI, 1907: 390 (Hikueru) *pro parte* not *Etisodus electra* (Herbst, 1801) = *Etisus aff. bifrontalis* fide GUINOT (1964: 56, 61; cf. Remark under *E. electra*). — *Etisus aff. bifrontalis* - GUINOT, 1964: 61 (Hikueru); 1985: 450 (List) = *E. bifrontalis* fide SERÈNE (1984: 230).

Etisus anaglyptus H. Milne Edwards, 1834

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Etisus anaglyptus* - MONTEFORTE, 1984: 170, annex 1, tab. a; 1987: 9 (Moorea). — GUINOT, 1985: 450 (List).

Etisus demani Odhner, 1925

DISTRIBUTION. — Society (Tahiti); Tuamotu (Hikueru?).

REFERENCES. — *Etisus demani* - MONTEFORTE, 1984: 170, annex 1, tab. a (Tahiti). — SERÈNE, 1984: 227, fig. 140, 143a, pl. 31f (Hikueru?; cf. Remark). — GUINOT, 1985: 450 (List).

REMARK. — The location "Hikueru" in SERÈNE (1984: 227) is only mentioned in the observations. It is not indicated under the material examined, and was not retrieved in the references cited by SERÈNE. It could be an erroneous reading in GUINOT (1964: 59), where "Hikueru" is cited under *Etisus frontalis* Dana, just beneath *E. demani* Odhner.

Etisus dentatus (Herbst, 1785)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Taiaro).

REFERENCES. — *Etisus dentatus* - BOONE, 1934: 119, pl. 62-63 (Tahiti). — FOREST & GUINOT, 1961: 86, fig. 80a-b (Tahiti); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 170, annex 1, tab. a (Moorea, Tahiti); 1987: 9 (Moorea). — GUINOT, 1985: 450 (List). — POUPIN, 1994a: 41, fig. 37, pl. 4h (Tahiti, Taiaro).

Etisus electra (Herbst, 1801)

DISTRIBUTION. — Gambier (Mangareva); Society (Tahiti); Tuamotu (Fakarava, Hikueru, Marutea South).

REFERENCES. — *Etisus rugosus* Jacquinot, 1852, pl. 4, fig. 2. — JACQUINOT & LUCAS, 1853: 33 (Mangareva). — *Etisodes electra* - NOBILI, 1907: 390 (Hikueru, "Rikitea" = Mangareva, Marutea South) *pro parte* cf. Remark. — RATHBUN, 1907: 42 (Fakarava). — FOREST & GUINOT, 1961: 89, fig. 82a-b (Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 450 (List). — *Etisus electra* - MONTEFORTE, 1984: 170, annex 1, tab. a (Tahiti). — SERÈNE, 1984: 228 (Syn.).

REMARK. — GUINOT (1964: 54) indicates that the material from Hikueru, attributed by NOBILI to *E. electra*, includes in fact three species: *E. electra*, *Etisus frontalis* (Dana, 1852), and *E. aff. bifrontalis* (Edmonson, 1935).

Etisus frontalis (Dana, 1852)

DISTRIBUTION. — Society (Moorea); Tuamotu (Hikueru).

REFERENCES. — *Etisodes electra* - NOBILI, 1907: 390 (Hikueru) *pro parte* not *Etisus electra* (Herbst, 1801) = *E. frontalis* fide GUINOT (1964: 54; cf. Remark under *E. electra*). — *Etisus frontalis* - GUINOT, 1964: 54 (Hikueru). — ODINETZ, 1983: 209 (Moorea). — SERÈNE, 1984: 229, fig. 139, pl. 31e (Syn.). — GUINOT: 1985: 450 (List). — *Etisodes frontalis* - GUINOT, 1985: 450 (List; cf. Remark).

REMARK. — In her list, GUINOT (1985: 450) mentions the two following species: "*Etisodes frontalis* (Dana, 1852) and *Etisus frontalis* Dana, 1852". It is obviously a mistake for a single species, described under *Etisodes frontalis* by DANA (1852b: 187).

Etisus laevimanus Randall, 1839

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora); Tuamotu (Mataiva).

REFERENCES. — *Etisus macrodactylus* - JACQUINOT, 1852, pl. 9, fig. 2 (Mangareva). — JACQUINOT & LUCAS, 1853: 30 (Mangareva). — *Etisus laevimanus* - NOBILI, 1907: 390 ("Rikitea" = Mangareva). — RATHBUN, 1907: 42 (Bora Bora). — FOREST & GUINOT, 1961: 88 (Mangareva); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 170,

annex 1, tab. a (Mataiva). — SERÈNE, 1984: 225 (Syn.). — GUINOT, 1985: 450 (List). — DELESALLE, 1985: 289 (Mataiva). — POUPIN, 1994a: 42, fig. 38, pl. 5a (Mangareva) - SYNONYMS - *Esisus macrodactylus* Bianconi, 1851.

Esisus punctatus Jacquinot, 1852

DISTRIBUTION. — Gambier (Mangareva).

REFERENCES. — *Esisus punctatus* Jacquinot, 1852, pl. 3, fig. 5. — JACQUINOT & LUCAS, 1853: 31 (Mangareva).

REMARK. — This *Esisus* has been figured by JACQUINOT, then described by JACQUINOT & LUCAS with this commentary "Cette espèce n'ayant pas été déposée au Muséum, c'est d'après la figure qui en a été donnée par MM. HOMBON et JACQUINOT que nous avons fait cette description". According to JACQUINOT & LUCAS, *Esisus punctatus* is related to *E. macrodactylus* Bianconi, 1851 (= *E. laevimanus* Randall, 1851) and to *E. anaglyptus* H. Milne Edwards, 1834.

Esisus splendidus Rathbun, 1906

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hikueru, Mataiva?, Raroia, Taiaro, Takapoto).

REFERENCES. — *Esisus (Etisodes) splendidus* - HOLTHUIS, 1953: 21 (Raroia). — MORRISON, 1954: 16 (Raroia). — *Esisus splendidus* - FOREST & GUINOT, 1961: 88, fig. 81a-c (Hikueru); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (Raroia); 1985: 450 (List). — MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva?, Moorea, Tahiti, Takapoto?). — SALVAT, 1986b: 72, photograph (French Polynesia). — BONVALLOT *et al.*, 1994: 145 (Tuamotu). — POUPIN, 1994a: 43, fig. 39, pl. 5b (Taiaro, Takapoto).

SUBFAMILY CHLORODIINAE

Chlorodiella barbata (Borradaile, 1900)

DISTRIBUTION. — Gambier (Mangareva); Society (Moorea, Tahiti); Tuamotu (Mataiva, Marutea South, Takapoto, Tikehau).

REFERENCES. — *Chlorodiella barbata* - FOREST & GUINOT, 1961: 96, fig. 93-94, 99a-b, 100 (Mangareva, Marutea South, Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213; 1985: 462 (Moorea); 1989: 112 (Tikehau). — NAIM, 1980a, annex 1, tab. 3; 1980b: 550 (Moorea). — THOMASSIN *et al.*, 1982: 394 (Moorea). — ODINETZ, 1983: 97 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GALZIN & POINTIER, 1985: 100 (Moorea). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti). — SALVAT & RICHARD, 1985: 360 (Takapoto).

Chlorodiella cytherea (Dana, 1852)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hikueru, Mataiva, Raraka, Takapoto, Tikehau).

REFERENCES. — *Chlorodius cytherea* Dana 1852a: 79; 1852b: 213; 1855, pl. 12, fig. 2a-c (Raraka, Tahiti). — *Chlorodiella cytherea* - FOREST & GUINOT, 1961: 95, fig. 90-92, 98a-b (Hikueru, Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213 (Moorea); 1989: 112 (Tikehau). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List).

Chlorodiella laevissima (Dana, 1852)

DISTRIBUTION. — Gambier (Mangareva); Marquesas (Nuku Hiva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fakarava, Hao, Makatea, Makemo, Mataiva, Rangiroa, Takapoto, Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Chlorodius laevissimus* - NOBILI, 1907: 393 ("Rikitea, Gatavake" = Mangareva, "Ohura" = Hao). — *Chlorodiella laevissima* - RATHBUN, 1907: 46 (Fakarava, Makemo, Rangiroa, Tahiti; 46m). — FOREST & GUINOT, 1961: 95, fig. 95-96, 101a-b (Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213; 1985: 462 (Moorea); 1989: 111, 114 (Moorea, Tikehau). — ODINETZ, 1983: 209 (Moorea, Tahiti). — MONTEFORTE, 1984: 170, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9

(Moorea). — SERÈNE, 1984: 260, fig. 171-172, pl. 36d-e (Syn.). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti). — *Chlorodiella laevissimus* - BOONE, 1934: 138, pl. 72 (Nuku Hiva, Tahiti).

Chlorodiella nigra (Forskål, 1775)

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fakarava, Hao, Makatea, Makemo, Marutea South, Mataiva, Nukutipipi, Rangiroa, Raraka, Takapoto, Tikehau).

REFERENCES. — *Chlorodius niger* - HELLER, 1865: 18 (Tahiti). — STIMPSON, 1858a: 33; 1907: 50 (Tahiti). — NOBILI, 1907: 393 ("Gatavake" = Mangareva, Hao, Marutea, "Tikahan" = Tikehau). — *Chlorodiella niger* - RATHBUN, 1907: 46 (Bora Bora, Fakarava, Makemo, Rangiroa). — SENDLER, 1923: 38 (Makatea). — SEURAT, 1934: 59 ("Gatavake" = Mangareva, Hao, Marutea South, "Tikahan" = Tikehau). — *Chlorodiella nigra* - FOREST & GUINOT, 1961: 95, fig. 87-89, 97a-b (Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213 (Moorea). — KROPP & BIRKELAND, 1981: 630, tab. 5 (Moorea, Takapoto). — THOMASSIN *et al.*, 1982: 394 (Moorea). — MONTEFORTE, 1984: 170, annex 1, tab. a, photograph p. 136a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — SERÈNE, 1984: 258, fig. 168, pl. 36b (Syn.). — DELESALLE, 1985: 288 (Mataiva). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti, Takapoto). — MERSCHARDT-SALVAT, 1991: 89 (Nukutipipi).

Garthiella aberrans (Rathbun, 1906)

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Pilodus aberrans* - PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 212 (Moorea); 1989: 111, 115 (Moorea, Tikehau; 30m). — GUINOT, 1985: 451 (List). — *Garthiella aberrans* - TITGEN, 1986: 57, fig. 1-2 (Syn.).

Liocarpilodes armiger (Nobili, 1906)

DISTRIBUTION. — Society (Moorea); Tuamotu (Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Liocarpilodes armiger* - PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213 (Moorea); 1989: 111, 115 (Moorea, Tikehau; 30m). — GUINOT, 1985: 450, with a ? (List).

Liocarpilodes harmsi (Balss, 1934)

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Pilodus harmsi* - PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 213 (Moorea). — *Liocarpilodes harmsi* - SERÈNE, 1984: 264 (Syn.).

Liocarpilodes integerrimus (Dana, 1852)

DISTRIBUTION. — Gambier (Mangareva); Society (Moorea, Tahiti); Tuamotu (Fakarava, Mataiva, Takapoto, Tikehau).

REFERENCES. — *Actumnus integerrimus* - RATHBUN, 1907: 56, pl. 1, fig. 12, pl. 8, fig. 3, 3a-b (Fakarava, Tahiti). — *Pilumnus margaritatus* - NOBILI, 1907: 398 ("Rikitea" = Mangareva) not *Pilumnus margaritatus* Ortmann, 1893 = *L. integerrimus* fide GUINOT (1964: 63). — *Liocarpilodes integerrimus* - FOREST & GUINOT, 1962: 64 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1964: 63, fig. 36a-b ("Rikitea" = Mangareva). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213; 1985: 462 (Moorea); 1989: 111, 115 (Moorea, Tikehau). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List).

Phymodius granulosus (De Man, 1888)

DISTRIBUTION. — Society (Moorea and/or Tahiti).

REFERENCES. — *Phymodius granulosus* - GUINOT, 1985: 451, with a ? (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti).

REMARK. — ODINETZ does not mention this material in her thesis (1983), but it is recorded in the Museum of Paris (MNHN B17071, coll. ODINETZ 1981 "Moorea, Tahiti, Takapoto, associé au *Pocillopora damicornis* et *P. elegans*", det. GUINOT).

Phymodius monticulosus (Dana, 1852)

DISTRIBUTION. — Gambier (Mangareva); Society (Moorea, Tahiti); Tuamotu (Fakahina, Fakarava, Mataiva, Marutea South, Takapoto).

REFERENCES. — *Chlorodius monticulosus* Dana, 1852a: 79; 1852b: 206; 1855, pl. 11, fig. 9a (Tahiti). — STIMPSON, 1858a: 31; 1907: 50 (Tahiti). — *Chlorodius Dehaanii* - HELLER, 1865: 19 (Tahiti) *pro parte*, some sp. attributed to *Phymodius ungulatus* in FOREST & GUINOT (1961: 106, 114), not *C. Dehaani*, synonym of *Phymodius granulatus* (Targioni Tozzetti, 1877) in SERÈNE (1984: 250). — *Cyclodius ornatus* - NOBILI, 1907: 397 (Fakahina, Marutea South). — RATHBUN, 1907: 51, pl. 5, fig. 5, pl. 7, fig. 8 (Fakarava, Tahiti). — *Phymodius monticulosus* - FOREST & GUINOT, 1961: 106, pl. 10, fig. 1-6 (Fakahina, Marutea South, Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — GUINOT, 1985: 451 (List). — POUPIN, 1994a: 44, fig. 40, pl. 5c (Fakahina, Mangareva) - SYNONYMS - *Cyclodius ornatus* Dana, 1852.

Phymodius nitidus (Dana, 1852)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Kaukura, Tikehau).

REFERENCES. — *Pilodius nitidus* - NOBILI, 1907: 393 (Kaukura). — *Chlorodopsis scabricula* - RATHBUN, 1907: 50, pl. 1, fig. 3, pl. 9, fig. 5 (Tahiti) not *Pilodius scabriculus* (Dana, 1852) = *Phymodius nitidus* fide FOREST & GUINOT (1961: 114). — *Phymodius nitidus* - FOREST & GUINOT, 1961: 114, pl. 15, fig. 1-4 (Kaukura, Tahiti); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a; 1987: 9 (Moorea). — GUINOT, 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 113 (Tikehau).

Phymodius ungulatus (H. Milne Edwards, 1834)

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Moorea, Raiatea, Tahiti); Tuamotu (Fakahina, Fakarava, Hikueru, Makemo, Mataiva, Rangiroa, Raroia, Takapoto, Tikehau).

REFERENCES. — *Chlorodius ungulatus* - DANA, 1852b: 205; 1855, pl. 11, fig. 8a-b (Tahiti). — *Chlorodius dehaanii* - HELLER, 1865: 19 (Tahiti) *pro parte fide* FOREST & GUINOT (1961: 110), not *C. Dehaani*, synonym of *Phymodius granulatus* (Targioni Tozzetti, 1877) in SERÈNE (1984: 250). — *Cyclodius gracilis* - NOBILI, 1907: 397 (Fakahina, "Rikitea" = Mangareva). — *Phymodius ungulatus* - RATHBUN, 1907: 46, pl. 3-4 (Bora Bora, Fakarava, Makemo, Rangiroa, Tahiti) *pro parte fide* FOREST & GUINOT (1961: 110; cf. Remark). — NOBILI, 1907: 393 ("Rikitea" = Mangareva). — BOONE, 1934: 140, pl. 73 (Raiatea, Tahiti) *pro parte fide* FOREST & GUINOT (1961: 110). — HOLTHUIS, 1953: 25 (Raroia). — FOREST & GUINOT, 1961: 110, fig. 86a-b, pl. 11, fig. 1-4, pl. 12, fig. 1-4, pl. 13, fig. 1-3, pl. 14, fig. 1-3 (Fakahina, Hikueru, Mangareva, Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1964: 74, fig. 38 (Tahiti); 1985: 451 (List). — PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 213 (Moorea); 1989: 113 (Tikehau). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — SERÈNE, 1984: 251, fig. 158, 161, pl. 35e (Tahiti; Syn.). — DELESALLE, 1985: 288 (Mataiva). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti, Takapoto). — SALVAT & RICHARD, 1985: 356 (Takapoto) - SYNONYMS - *Cyclodius gracilis* Dana, 1852.

REMARK. — FOREST & GUINOT (1961: 111) indicate that RATHBUN's material is only partially assignable to this species, but without mentioning the localities referring to the real *Phymodius ungulatus*. Thus, for this species, some of the islands mentioned by RATHBUN are doubtful.

Pilodius areolatus (H. Milne Edwards, 1834)

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Hikueru, Makemo, Marutea South, Mataiva, Raroia).

REFERENCES. — *Actaeodes affinis* Dana, 1852a: 78; 1852b: 197; 1855, pl. 11, fig. 3 (Tahiti, Tuamotu). — *Actaea affinis* - RATHBUN, 1907: 42 (Makemo). — *Chlorodopsis areolata* - NOBILI, 1907: 396, pl. 2, fig. 3 (Hikueru, "Rikitea")

= Mangareva, Marutea South). — HOLTHUIS, 1953: 15 (Raroia). — MORRISON, 1954: 13 (Raroia). — *Pilodius areolatus* - FOREST & GUINOT, 1961: 90 (Hikueru, Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva). — GUINOT, 1985: 451 (List). — CLARK & GALIL, 1993: 1125, fig. 1a-g, 31a, 40a, 44b (Bora Bora, "Maharepa, Afareaitu, Temae" = Moorea; Syn.).

Pilodius flavus Rathbun, 1893

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hao, Tikehau).

REFERENCES. — *Pilodius pubescens* - NOBILI, 1907: 395, with a ? ("Ohura" = Hao) not *Pilodius pubescens* Dana, 1852 = *P. flavus* fide CLARK & GALIL (1993: 1130, 1146). — *Pilodius flavus* - FOREST & GUINOT, 1961: 95 (Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 27 (Moorea); 1989: 111, 115 (Moorea, Tikehau). — GUINOT, 1985: 451 (List). — CLARK & GALIL, 1993: 1130, fig. 4a-g, 32b, 40d, 41a (Syn.).

Pilodius paumotensis Rathbun, 1907

DISTRIBUTION. — Society (Bora Bora); Tuamotu (Fakarava, Makemo, Marutea South, Tikehau).

REFERENCES. — *Chlorodopsis granulatus* - NOBILI, 1906: 396 (Marutea South) not *Pilodius granulatus* Stimpson, 1859 = *P. paumotensis* fide GUINOT (1962: 238). — *Pilodius paumotensis* Rathbun 1907: 52, pl. 8, fig. 2, 2a-b (Fakarava, Makemo). — FOREST & GUINOT, 1962: 66 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 451 (List). — PEYROT-CLAUSADE, 1989: 113 (Tikehau). — CLARK & GALIL, 1993: 1143, fig. 10a-g, 35b, 43a (Bora Bora, Fakarava, Makemo, Marutea South).

Pilodius pubescens Dana, 1852

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Pilodius pubescens* - GUINOT, 1985: 451 (List; probably after NOBILI, cf. under *P. flavus*). — CLARK & GALIL, 1993: 1146, fig. 12a-g, 36b, 43b ("Ternae" = Temae at Moorea, Society). — Not *Pilodius pubescens* - NOBILI, 1907: 395, with a ? = *P. flavus* in CLARK & GALIL (1993: 1130).

Pilodius pugil Dana, 1852

DISTRIBUTION. — Gambier (Mangareva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fakarava, Hikueru, Makatea, Makemo, Mataiva, Rangiroa, Takapoto, Tikehau).

REFERENCES. — *Chlorodopsis pugil* - NOBILI, 1907: 395 ("Rikitea" = Mangareva). — *Chlorodopsis spinipes* - RATHBUN, 1907: 50, pl. 2, fig. 5 (Bora Bora, Fakarava, Makemo, Rangiroa) not *Pilodius spinipes* Heller, 1861 = *P. pugil* fide CLARK & GALIL (1983: 1149). — *Pilumnus globosus* - BOONE, 1934: 152, pl. 78 (Tahiti) not *Globopilumnus globosus* (Dana, 1852) = *Pilodius pugil* with a ? fide SERÈNE & LUOM (1959: 320). — *Pilodius pugil* - FOREST & GUINOT, 1961: 91 (Hikueru, Mangareva); 1962: 64 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 213; 1985: 462 (Moorea); 1989: 113 (Tikehau). — MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289, 305 (Mataiva). — GUINOT, 1985: 451 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea, Tahiti, Takapoto). — SALVAT & RICHARD, 1985: 358, 360 (Takapoto). — CLARK & GALIL, 1993: 1149, fig. 13a-g, 37a, 43c (Hikueru, Mangareva, "Ternae, Tiahura and Afareaitu" = Moorea, Tahiti, Takapoto).

Pilodius scabriculus Dana, 1852

DISTRIBUTION. — Gambier (Temoe); Society (Bora Bora, Huahine, Maiao?, Moorea, Tahaa?, Tahiti); Tuamotu (Fakarava, Fakahina, Hao, Hikueru, Makatea, Makemo, Marutea North?, Marutea South, Mataiva, Raraka, Takapoto, Tikehau).

REFERENCES. — *Pilodius scabriculus* Dana, 1852a: 80; 1852b: 220; 1855, pl. 12, fig. 9 (Raraka). — NOBILI, 1907: 394 ("Fakaina" = Fakahina, Fakarava, Marutea, Marutea South, Temoe). — FOREST & GUINOT, 1961: 91, fig. 83a-b, 84, 86bis (Hikueru, Tahiti); 1962: 66 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977b: 213 (Moorea); 1989: 113 (Tikehau). — MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289, 305 (Mataiva). — GUINOT, 1985: 451 (List). — CLARK & GALIL, 1993: 1152, fig. 14a-g, 37b, 43d (Bora Bora, Fakahina, Hao, Hikueru, Huahine, Maiai =

Maiao?, Marutea, Moorea, Raiatea, Tahiti, Temoe, "Tickahau" = Tikehau, "Vaioreia" = Vaitoare?, Tahaa; Syn.). — *Chlorodopsis venusta* Rathbun, 1907: 49, pl. 1, fig. 5 (Fakarava, Makemo).

Tweedieia laysani (Rathbun, 1906)

DISTRIBUTION. — Tuamotu (Tikehau).

REFERENCES. — *Tweedieia laysani* - PEYROT-CLAUSADE, 1989: 112 (Tikehau).

Tweedieia odhneri (Gordon, 1934)

DISTRIBUTION. — Society (Moorea).

REFERENCES. — *Tweedieia odhneri* - PEYROT-CLAUSADE, 1977a, annex of the species: 27; 1977: 212 (Moorea). — GUINOT, 1985: 451, with a ? (List).

FAMILY TRAPEZIIDAE

Jonesius triunguiculatus (Borradaile, 1902)

DISTRIBUTION. — Society (Raiatea).

REFERENCES. — *Jonesius triunguiculatus* - GALIL & TAKEDA, 1986: 165, fig. 1-4 ("Tetaro" = Raiatea).

Quadrella lewinsohni Galil, 1986

DISTRIBUTION. — Marquesas (Tahuata).

REFERENCES. — *Quadrella* sp. - MONOD, 1979: 9, fig. 1-8 (Tahuata). — *Quadrella cyrenae* - SERÈNE, 1975: 510, fig. 3-4, pl. 1b', e' (Tahuata; MONOD's material) not *Q. cyrenae* Ward, 1942, synonym of *Q. maculosa* Alcock, 1898 = *Q. lewinsohni* nov. in GALIL (1986a: 285). — GUINOT, 1985: 452 (List). — *Quadrella lewinsohni* Galil, 1986a: 285, fig. 5a-b, 6 (Marquesas; MONOD's and SERÈNE's material).

Quadrella maculosa Alcock, 1898

DISTRIBUTION. — Marquesas (Fatu Hiva) - *Littoral to sublittoral*.

REFERENCES - *Quadrella maculosa* - NEW MATERIAL - Coll. J. POUPIN, det. B. GALIL (Fatu Hiva; 49m).

Tetralia cinctipes Paulson, 1875

DISTRIBUTION. — Austral (Rapa) - *Sublittoral*.

REFERENCES. — *Tetralia cinctipes* - GALIL, 1986b: 97, fig. 1-3 (Rapa; 90m).

Tetralia glaberrima (Herbst, 1790)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Raiatea, Tahiti); Tuamotu (Aratika, Fakarava, Hikueru, Makemo, Marutea South, Tikehau).

REFERENCES. — *Tetralia glaberrima* - DANA, 1852b: 262; 1855, pl. 16, fig. 3a-h ("Carlshoff" = Aratika, Tahiti). — ORTMANN, 1893b: 485 (Tahiti). — NOBILI, 1907: 404 (Marutea). — RATHBUN, 1907: 60 (Fakarava, Makemo, Tahiti). — BOONE, 1934: 174, pl. 89 (Raiatea, Tahiti). — SEURAT, 1934: 59 (Tahiti, Marutea South, Marquesas). — FOREST & GUINOT, 1961: 139 (Hikueru); 1962: 70 (Biogeography "Tahiti-Tuamotu, Marquesas"). — SERÈNE, 1984: 281 (Syn.; cf. Remark). — GUINOT, 1985: 452 (List). — PEYROT-CLAUSADE, 1989: 111 (Tikehau). — *Trapezia serratifrons* Jacquinot, 1852, pl. 4, fig. 20-23. — JACQUINOT & LUCAS, 1853: 47 (Nuku Hiva). — *Tetralia cavimana* - HELLER, 1865: 26 (Tahiti) - SYNONYMS - *Tetralia cavimana* Heller, 1861.

REMARK. — In SERÈNE (1984: 281) all the above references (except BOONE, 1934, and PEYROT-CLAUSADE, 1989) can be attributed either to *T. glaberrima* or to *T. heterodactyla* Heller, 1861. However, the specimens attributed to *cavimana*, and the specimens identified to *glaberrima* with a figure of the male pleopod, are *T. glaberrima* without hesitation.

Tetraloides nigrifrons (Dana, 1852)

DISTRIBUTION. — Society (Raiatea); Tuamotu (Makemo, Pukapuka, Tikehau).

REFERENCES. — *Tetralia nigrifrons* Dana, 1852a: 83; 1852b: 262; 1865, pl. 16, fig. 2a-d ("Honden" = Pukapuka). — *Tetraloides nigrifrons* - GALIL, 1985: 72, fig. 1-3 (Makemo, Tikehau, "Taoru" = Raiatea).

Trapezia areolata Dana, 1852

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hikueru, Takapoto).

REFERENCES. — *Trapezia areolata* Dana, 1852a: 83; 1852b: 259; 1855, pl. 15, fig. 8a (Tahiti). — CANO, 1888: 173 (Tahiti). — ORTMANN 1893b: 485 (Tahiti; cf. Remark). — FOREST & GUINOT, 1961: 135, fig. 133 (Hikueru). — ODINETZ, 1983: 31 (French Polynesia). — ODINETZ, 1984a: 443, fig. 3c, 4c (Moorea, Tahiti, Takapoto). — GALIL & LEWINSOHN, 1985a: 286, fig. 1, 3-4 (Tahiti). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea, Tahiti, Takapoto). — ? *Trapezia ferruginea areolata* - SEURAT, 1934: 59 (Tahiti). — *Trapezia reticulata* - KROPP & BIRKELAND, 1981: 629 (Moorea, Takapoto). — ODINETZ, 1983: 31, 205 photograph 3 (Moorea, Tahiti, Takapoto). — GUINOT, 1985: 452 (List; after ODINETZ, 1983) - All, not *Trapezia reticulata* Stimpson, 1858, synonym of *T. septata* Dana, 1852 = *T. areolata* fide ODINETZ (1984a: 443; does not mention KROPP & BIRKELAND, but it is probably the same material).

REMARK. — In GALIL & LEWINSOHN (1985a), ORTMANN's (1893b) work, in which *T. aerolata* is recorded from Tahiti, New Guinea, and Palau, is cited at the same time under *T. aerolata* and *T. septata*. It is probable that the two Tahitian specimens belong to *T. areolata*, and the other to *T. septata* (opt. cit.: 291).

Trapezia bella Dana, 1852

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Aratika, Hikueru, Mataiva, Takapoto).

REFERENCES. — *Trapezia bella* Dana, 1852a: 83; 1852b: 254; 1855, pl. 15, fig. 2 ("Carlshoff" = Aratika). — FOREST & GUINOT, 1961: 133, fig. 129-130, 135a-b (Hikueru); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — KROPP & BIRKELAND, 1981: 629 (Moorea, Takapoto). — ODINETZ, 1983: 206 (Tahiti, Takapoto). — SERÈNE, 1984: 278 (Syn.). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Tahiti?, Takapoto). — *Trapezia digitalis bella* - RATHBUN, 1907: 59 (Tahiti). — Not *Trapezia bella* Dana - NOBILI, 1907: 403 (Hao). — SEURAT, 1934: 59 (Hao; NOBILI's material) = *Trapezia speciosa* fide SERÈNE (1984: 278).

Trapezia cymodoce (Herbst, 1799)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Bora Bora, Moorea, Tahiti); Tuamotu (Fakarava, Makemo, Rangiroa, Takapoto).

REFERENCES. — *Trapezia dentata* - DANA, 1852b: 258; 1855, pl. 15, fig. 6a-d (Tahiti). — *Trapezia hirtipes* Jacquinot, 1852, pl. 4, fig. 14-16 (Nuku Hiva). — JACQUINOT & LUCAS, 1853: 44 (Nuku Hiva). — *Trapezia cymodoce dentata* - RATHBUN, 1907: 58 (Bora Bora, Fakarava, Makemo, Rangiroa; cf. Remark). — *Trapezia cymodoce ferruginea* - RATHBUN, 1907: 58 (Bora Bora, Fakarava, Makemo, Rangiroa; cf. Remark). — *Trapezia ferruginea dentata* - SEURAT, 1934: 59 (Tahiti). — *Trapezia cymodoce* - ORTMANN, 1897b: 203 (Tahiti). — SEURAT, 1934: 59 (Tahiti, Marquesas). — ODINETZ, 1983: 205, photograph 2 (Moorea, Takapoto, Tahiti); 1984a: 432, fig. 1-2 (Moorea, Tahiti, Takapoto; pro parte, some specimens would belong to *T. ferruginea*, cf. Remark under that species). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea?, Tahiti, Takapoto). — GALIL & CLARK, 1990: 378 (Syn.). — *Trapezia cymodoce* sp.1 - KROPP & BIRKELAND, 1981: 629 (Moorea, Takapoto) fide distinction in ODINETZ (1984b: 124). — ODINETZ, 1984b: 125 (Moorea, Tahiti, Takapoto). — Not *Trapezia cymodoce* - DANA, 1852b: 257; 1855, pl. 15, fig. 5a-i (Tahiti). — SENDLER, 1923: 39 (Tahiti) - The two references = *T. ferruginea* Latreille 1825 fide GALIL & CLARK (1990: 378) - SYNONYMS - *Trapezia dentata* (Macleay, 1838).

REMARK. — The material attributed to *T. cymodoce dentata* by RATHBUN (1907) has been partially re-examined by GALIL & CLARK (1990; sp. of Ellice islands "Funafuti reef") and belongs to *T. cymodoce*. Arbitrarily we also attribute to this species the material from French Polynesia. Concerning the *T. cymodoce ferruginea* also recorded by this author, from French Polynesia, Ellice, and Easter island, they would belong in part to that species (?Bora Bora, Fakarava, Makemo, Rangiroa), in part to *T. guttata* Rüppell (2 sp. from "Mohican reef" at Rangiroa) (cf. GALIL & CLARK, 1990: 380-382), and in part to *T. punctimanus* (sp. from Easter island; cf. ODINETZ 1984a: 446).

Trapezia digitalis Latreille, 1825

DISTRIBUTION. — Marquesas (Nuku Hiva); Tuamotu (Makatea, Takapoto).

REFERENCES. — *Trapezia fusca* Jacquinot, 1852, pl. 4, fig. 17-19. — JACQUINOT & LUCAS, 1853: 45 (Nuku Hiva). — *Trapezia digitalis* - SENDLER, 1923: 40 (Makatea). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu, Marquesas"). — ODINETZ, 1983: 31, 206 (Takapoto; cf. Remark). — SERÈNE, 1984: 277 (Syn.). — GUINOT, 1985: 452 (List).

REMARK. — Although ODINETZ (1983: 31) has indicated that her material was collected at Guam, one specimen in table 9 (p. 206) is recorded from Takapoto.

Trapezia ferruginea Latreille, 1825

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Bora Bora, Moorea, Raiatea, Tahiti); Tuamotu (Takapoto).

REFERENCES. — *Trapezia ferruginea* - DANA, 1852b: 260; 1855, pl. 16, fig. 1a-b (Tahiti, *pro parte*, see under *T. guttata*; Samoan specimens are *T. serenei* fide ODINETZ, 1984a: 440, 442). — BOONE, 1934: 171, pl. 88 (Bora Bora, Nuku Hiva, Raiatea). — SEURAT, 1934: 59 (Tahiti, Marquesas). — FOREST & GUINOT, 1961: 136, fig. 137a-b (Tahiti). — KROPP & BIRKELAND, 1981: 629 (Moorea). — SERÈNE, 1984: 273 (Syn.). — GUINOT, 1985: 452 (List). — PEYROT-CLAUSADE, 1989: 115 (Moorea). — *Trapezia cymodoce* - DANA, 1852b: 257; 1855, pl. 15, fig. 5a-i (Tahiti). — SENDLER, 1923: 39 (Tahiti) - The two references not *Trapezia cymodoce* (Herbst, 1799) = *T. ferruginea* fide GALIL & CLARK (1990: 380). — ODINETZ, 1984a: 432 (Moorea, Tahiti, Takapoto; *pro parte*, cf. Remark). — *Trapezia miniata* Jacquinot, 1852, pl. 4, fig. 10-13. — JACQUINOT & LUCAS, 1853: 43 (Nuku Hiva). — Not *Trapezia cymodoce ferruginea* - RATHBUN, 1907: 58 (cf. Remark under *T. cymodoce*).

REMARK. — *Trapezia ferruginea* Latreille, 1825 was proposed as a synonym of *T. cymodoce* (Herbst, 1799) in ODINETZ (1984a), but this proposition was not followed by GALIL & CLARK (1990).

Trapezia flavopunctata Eydoux & Souleyet, 1842

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hikueru).

REFERENCES. — *Trapezia flavopunctata* - ORTMANN, 1893b: 485 (Tahiti). — FOREST & GUINOT, 1961: 136, fig. 138a-b (Hikueru). — ODINETZ, 1983: 34, 205 (Tahiti; p. 205 = "*T. flavomaculata*" sic). — GALIL & LEWINSOHN, 1985b: 210 ("Papetoai bay" = Moorea, Tahiti). — GUINOT, 1985: 452 (List). — *Trapezia rufopunctata flavopunctata* - SEURAT, 1934: 59 (Tahiti).

Trapezia formosa Smith, 1869

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Mataiva, Takapoto).

REFERENCES. — *Trapezia formosa* - KROPP & BIRKELAND, 1981: 629 (Takapoto). — ODINETZ, 1983: 206, photograph 4 (Moorea, Takapoto, Tahiti). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea?, Takapoto, Tahiti).

REMARK. — According to P. CASTRO (personal communication), who has re-examined the material of these references, it could rather belongs to a new species.

Trapezia guttata Rüppell, 1830

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hikueru, Rangiroa, Takapoto, Tikehau).

REFERENCES. — *Trapezia guttata* - HELLER, 1865: 25 (Tahiti). — FOREST & GUINOT, 1961: 136, fig. 134, 139a-b (Hikueru, Tahiti); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — ODINETZ, 1983: 205, photograph 8 (Moorea, Tahiti, Takapoto); 1984a: 442 (Moorea, Tahiti). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea?, Tahiti, Takapoto). — PEYROT-CLAUSADE, 1989: 111 (Tikehau). — GALIL & CLARK, 1990: 381 (Syn.). — *Trapezia cymodoce ferruginea* - RATHBUN, 1907: 58 (only some specimens from Rangiroa) not *T. ferruginea* Latreille, 1825 = *T. guttata* fide GALIL & CLARK (1990: 381; cf. Remark under *T. cymodoce*). — *Trapezia ferruginea* - DANA, 1852b: 260; 1865, pl. 16, fig. 1b (Tahiti) *pro parte* not *T. ferruginea* Latreille, 1825 = *T. guttata* fide GALIL & CLARK (1990: 381, 382). — *Trapezia ferruginea guttata* - SEURAT, 1934: 59 (Tahiti). — *Trapezia*

davaoensis - KROPP & BIRKELAND, 1981: 629 (Moorea). — GUINOT, 1985: 452 (List) - SYNONYMS - *Trapezia davaoensis* Ward, 1941.

Trapezia punctimanus Odinetz, 1984

DISTRIBUTION. — Society (Tahiti); Tuamotu (Takapoto).

REFERENCES. — *Trapezia punctimanus* Odinetz, 1983: 35, 206 photograph 7 (Thesis; French Polynesia); 1984a: 445, fig. 3e, 4e (Tahiti, Takapoto). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Tahiti, Takapoto).

Trapezia rufopunctata (Herbst, 1801)

DISTRIBUTION. — Society (Moorea, Raiatea, Tahiti); Tuamotu (Makemo, Takapoto, Tikehau).

REFERENCES. — *Trapezia rufo-punctata* - DANA, 1852b: 255; 1855, pl. 15, fig. 3a-b (Tahiti). — *Trapezia rufopunctata* - RATHBUN, 1907: 57 (Makemo). — BOONE, 1934: 166, pl. 86 (Raiatea). — SEURAT, 1934: 59 (Tahiti, but not the Marquesas certainly cited after JACQUINOT & LUCAS; see below). — ODINETZ, 1983: 34 (Moorea, Tahiti, Takapoto). — GUINOT, 1985: 452 (List). — PEYROT-CLAUSADE, 1989: 112 (Tikehau). — Not *Trapezia rufo-punctata* - JACQUINOT, 1852, pl 4, fig. 8-9. — JACQUINOT & LUCAS, 1853: 41 (Nuku Hiva) = *T. tigrina* Eydoux & Souleyet, 1842 *fide* GALIL & LEWINSOHN (1985b: 166).

Trapezia septata Dana, 1852

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Trapezia ferruginea areolata* - SENDLER, 1923: 40 (Tahiti) not *T. areolata* Dana, 1852 = *T. septata* *fide* GALIL & LEWINSOHN (1985a: 288; cf. Remark).

REMARK. — Although SENDLER is cited under that species in GALIL & LEWINSOHN, these authors have not examined Polynesian material. They only mention that (p. 291) "T. septata seems to be more widely distributed and more common than *T. areolata*. Thus, specimens identified as *T. aerolata*, with the exception of those mentioned by CANO (1888), ORTMANN (1893) (part) and FOREST & GUINOT (1961), should rightly be named *T. septata*".

Trapezia serenei Odinetz, 1984

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Mataiva, Takapoto).

REFERENCES. — *Trapezia serenei* Odinetz, 1983: 34, 206, photograph 6; 1984a: 440, fig. 3b, 4b (Mataiva, Moorea, Tahiti, Takapoto). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea?, Tahiti, Takapoto). — SALVAT & RICHARD, 1985: 344 (Takapoto). — *Trapezia cymodoce* sp. 2 - KROPP & BIRKELAND, 1981: 629 (Moorea, Takapoto). — ODINETZ, 1984b: 125 (Moorea, Tahiti, Takapoto) - *Fide* distinction in ODINETZ (1984b: 124).

Trapezia speciosa Dana, 1852

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Aratika, Fakarava, Hao, Hikueru, Makemo, Marutea South, Mataiva, Takapoto).

REFERENCES. — *Trapezia speciosa* Dana, 1852a: 83; 1852b: 253; 1855, pl. 15, fig. 1 ("Carlshoff" = Aratika). — NOBILI, 1907: 403 (Marutea). — SEURAT, 1934: 59 (Marutea South). — FOREST & GUINOT, 1961: 133, fig. 131-132, 136a-b (Hikueru); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — KROPP & BIRKELAND, 1981: 629 (Moorea, Takapoto). — ODINETZ, 1983: 205 (Moorea, Tahiti, Takapoto). — SERÈNE, 1984: 278 (Syn.). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea, Tahiti, Takapoto). — *Trapezia digitalis speciosa* - RATHBUN, 1907: 59 (Fakarava, Makemo, Tahiti). — *Trapezia bella* - NOBILI, 1907: 403 (Hao) not *T. bella* Dana, 1852 = *T. speciosa* *fide* SERÈNE (1984: 278).

Trapezia tigrina Eydoux & Souleyet, 1842

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Moorea, Tahiti); Tuamotu (Anaa, Makemo?, Takapoto, Tikehau).

REFERENCES. — *Trapezia maculata* - DANA, 1852b: 256; 1855, pl. 15, fig. 4 (Tahiti) not *T. maculata* (MacLeay, 1838) = *T. tigrina* fide SERÈNE (1984: 275) and GALIL & LEWINSHON (1984: 166). — *Trapezia rufo-punctata* - JACQUINOT, 1852, pl 4, fig. 8-9. — JACQUINOT & LUCAS, 1853: 41 (Nuku Hiva) not *T. rufopunctata* (Herbst, 1799) = *T. tigrina* fide GALIL & LEWINSOHN (1984: 166). — *Trapezia rufopunctata* var. *maculata* - ORTMANN, 1893b: 484 (Tahiti). — *Trapezia cymodoce maculata* - RATHBUN, 1907: 59 (Makemo) - These two references, with a ?, not *T. maculata* (MacLeay, 1838) = *T. tigrina* fide GALIL & LEWINSOHN (1984: 167). — ? *Trapezia ferruginea maculata* - SEURAT, 1934: 59 (Tahiti). — *Trapezia wardi* - KROPP & BIRKELAND, 1981: 629 (Moorea, Takapoto). — GUINOT, 1985: 452 (List). — *Trapezia tigrina* - ODINETZ, 1983: 205, photograph 5 (Moorea, Tahiti, Takapoto). — GALIL & LEWINSOHN, 1984: 166, fig. 1 (Anaa, "Tikehae lagoon, Tuamotu" = Tikehau, Society; Syn.). — GUINOT, 1985: 452 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea?, Tahiti, Takapoto) - SYNONYMS - *Trapezia wardi* Serène, 1969.

FAMILY PILUMNIDAE

Actumnus asper (Rüppell, 1830)

DISTRIBUTION. — Gambier (Mangareva); Tuamotu (Marutea South).

REFERENCES. — *Actumnus bonnieri* - NOBILI, 1907: 400 ("Rikitea" = Mangareva, Marutea South). — *Actumnus asper* - BALSS, 1933: 36 (Syn.). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1964: 98 (Syn.); 1985: 452 (List) - SYNONYMS - *Actumnus bonnieri* Nobili, 1905.

Actumnus digitalis (Rathbun, 1907)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Platypodia digitalis* Rathbun, 1907: 38, pl. 1, fig. 6, pl. 9, fig. 4, 4a (Tahiti). — FOREST & GUINOT, 1962: 62 (Biogeography "Tahiti-Tuamotu"). — *Actumnus digitalis* - GUINOT, 1969: 225 (Syn.); 1985: 452 (List).

Actumnus globulus Heller, 1861

DISTRIBUTION. — Tuamotu (Hao, Hikueru).

REFERENCES. — *Actumnus globulus* - NOBILI, 1907: 400 ("Ohura" = Hao, Hikueru; cf. thereafter). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1969: 226 (writes about NOBILI's work: "par contre il est bien possible que les "globulus" polynésiens de NOBILI (1907, p. 50 sic) soient en fait des *digitalis*"); 1985: 452 (List).

Actumnus obesus Dana, 1852

DISTRIBUTION. — Marquesas; Society (Tahiti).

REFERENCES. — *Actumnus obesus* - BALSS, 1933: 37 (Marquesas). — BOONE, 1934: 154, pl. 79 (Tahiti). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu, Marquesas"). — GUINOT, 1985: 452 (List).

Actumnus setifer (De Haan, 1835)

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Actumnus tomentosus* Dana, 1852a; 1852b: 243; 1855, pl. 14, fig. 2a-c (Tahiti). — *Actumnus setifer* - ORTMANN, 1893b: 474 (Syn.). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List).

Pilumnus merodentatus Nobili, 1906

DISTRIBUTION. — Gambier (Mangareva); Marquesas; Society (Tahiti); Tuamotu (Mataiva?).

REFERENCES. — *Pilumnus merodentatus* Nobili, 1906a: 263; 1907: 399 ("Rikitea" = Mangareva). — SEURAT, 1934: 60 (French Polynesia). — FOREST & GUINOT, 1961: 132, fig. 128 (Mangareva; Syn.); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List). — *Pilumnus longicornis merodentatus* - BALSS, 1933: 16 (Mangareva,

NOBILI's material and also Tahiti, Marquesas) - RELEVANT MATERIAL - *Pilumnus cf. merodentatus* - MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva). — GUINOT, 1985: 452 (List).

Pilumnus parvulus Nobili, 1906

DISTRIBUTION. — Gambier (Mangareva); Tuamotu (Moruroa).

REFERENCES. — *Pilumnus parvulus* Nobili, 1906a: 263; 1907: 398 ("Gatavake, Rikitea, Tokaerero" = Mangareva; some sp. in pearl oyster, 25m). — SEURAT, 1934: 60 (French Polynesia). — FOREST & GUINOT, 1961: 130, fig. 126, pl. 27, fig. 1 (Mangareva); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List). — TAKEDA & MIYAKE, 1968: 6 (Key) - NEW MATERIAL - Coll. & det. J. POUPIN (Moruroa; 1sp. in pearl oyster *Pinctada margaritifera*).

Pilumnus ransoni Forest & Guinot, 1961

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Pilumnus ransoni* Forest & Guinot, 1961: 130, fig. 123-124, 127, pl. 4, fig. 1-2, pl. 17, fig. 2. (Tahiti); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List).

Pilumnus tahitensis De Man, 1890

DISTRIBUTION. — Society (Moorea, Raiatea, Tahiti); Tuamotu (Fakarava, Marutea South, Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Pilumnus tahitensis* de Man, 1890: 61, pl. 3, fig. 4 (Tahiti). — ORTMANN, 1893b: 437 (Tahiti). — NOBILI, 1907: 399 (Marutea South). — RATHBUN, 1907: 56 (Fakarava). — BALSS, 1933: 25 (Tahiti). — SEURAT, 1934: 60 (Marutea South). — FOREST & GUINOT, 1961: 129, fig. 125 (Raiatea, Tahiti); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 452 (List). — PEYROT-CLAUSADE, 1989: 111, 115 (Moorea, Tikehau; 30m).

FAMILY CARPILIIDAE

Carpilius convexus (Forskål, 1775)

DISTRIBUTION. — Austral (Rapa, Rurutu); Society (Moorea, Tahiti); Tuamotu (Hikueru, Makatea, Makemo, Mataiva, Moruroa, Taiaro, Takapoto, Takaroa, Raroia) - *Littoral to sublittoral*.

REFERENCES. — *Carpilius convexus* - STIMPSON, 1858a: 32; 1907: 37 (Tahiti). — RATHBUN, 1907: 37 (Makemo). — BOONE, 1934: 89, pl. 43-45 (Tahiti). — HOLTHUIS, 1953: 12 (Raroia). — CHABOIS L. & F., 1954: 91, fig. 2 (French Polynesia). — MORRISON, 1954: 16 (Raroia). — BUITENDIJK, 1960: 263 (Takaroa). — FOREST & GUINOT, 1961: 37 (Hikueru, Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (French Polynesia); 1985: 449 (List). — CHEVALIER *et al.*, 1968: 92, 138 (Moruroa). — MONTEFORTE, 1984: 170, annex 1, tab. a (Makatea, Mataiva, Moorea, Takapoto). — SERÈNE, 1984: 303, fig. 208-209 (Mataiva). — DELESALLE, 1985: 289 (Mataiva). — SALVAT & RICHARD, 1985: 362 (Takapoto). — SALVAT, 1986b: 72 (French Polynesia). — BONVALLOT *et al.*, 1994: 140, photograph (Tuamotu). — POUPIN, 1994a: 45, fig. 41, pl. 5d (Hikueru, Rapa, Rurutu, Tahiti, Taiaro; up to 60m).

Carpilius maculatus (Linné, 1758)

DISTRIBUTION. — Gambier; Society (Moorea, Tahiti); Tuamotu (Fakarava, Hao, Hikueru, Makatea, Mataiva, Moorea, Moruroa, Raraka, Raroia, Taiaro, Takapoto).

REFERENCES. — *Carpilius maculatus* - DANA, 1852b: 160 (Raraka). — STIMPSON, 1858a: 32; 1907: 37 (Tahiti). — HELLER, 1865: 9 (Tahiti). — NOBILI, 1907: 386 ("Ohura" = Hao). — RATHBUN, 1907: 37 (Fakarava, Tahiti). — PESTA, 1913: 39, pl. 3, fig. 4 (Tahiti, with a ?). — BOONE, 1934: 86, pl. 39-42 (Tahiti). — SEURAT, 1934: 60 (Gambier, Tuamotu). — HOLTHUIS, 1953: 12 (Raroia). — CHABOIS L. & F., 1954: 91, fig. 1 (French Polynesia). — MORRISON, 1954: 16 (Raroia). — BABLET, 1972: 32, pl. 11 (French Polynesia). — FOREST & GUINOT, 1961: 37 (Hikueru, Tahiti); 1962: 60 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (French Polynesia); 1985: 450 (List). — CHEVALIER *et al.*, 1968: 92, 138 (Moruroa). — CROSNIER, 1984: 302, fig. 208-209, pl. 44e (Mataiva). — MONTEFORTE, 1984: 170,

annex 1, tab. a, photograph p.140a (haut) (Makatea, Mataiva, Moorea, Tahiti, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — SALVAT, 1986b: 72, photograph (French Polynesia). — PARDON, 1992: 82, photograph (Tahiti). — BONVALLOT *et al.*, 1994: 141, photograph (Tuamotu). — POUPIN, 1994a: 46, fig. 42, pl. 5e (Mataiva, Tahiti, Taiaro).

FAMILY MENIPPIDAE

Dacryopilumnus eremita Nobili, 1906

DISTRIBUTION. — Gambier (Mangareva); Tuamotu (Amanu, Hao, Makatea, Marutea South, Mataiva).

REFERENCES. — *Dacryopilumnus eremita* Nobili, 1906a: 264; 1907: 400, pl. 2, fig. 4 (Amanu, Hao, "Rikitea" = Mangareva; gen. and sp. nov.). — FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu"). — CROSNIER, 1984: 313, fig. 240-241, pl. 47e (Mangareva, Marutea). — MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva). — GUINOT, 1985: 450 (List).

Domecia glabra Alcock, 1899

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hao, Tikehau) - *Littoral to sublittoral*.

REFERENCES. — *Domecia hispida* - NOBILI, 1907: 404 (Hao) not *Domecia hispida* Eydoux & Souleyet, 1842 = *D. glabra* fide FOREST & GUINOT (1961: 126). — *Domecia glabra* - FOREST & GUINOT, 1961: 126, fig. 115-116, 120-122, 124bis (Hao); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1962: 240, fig. 13a-b (Hao). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213 (Moorea); 1989: 111 (Tikehau; 30m). — GUINOT, 1985: 450 (List).

Domecia hispida Eydoux & Souleyet, 1842

DISTRIBUTION. — Gambier (Mangareva); Marquesas (Nuku Hiva); Society (Moorea, Tahiti); Tuamotu (Hikueru, Makemo, Takapoto, Tikehau).

REFERENCES. — *Domecia hispida* - DANA, 1852b: 251 (Tahiti). — RATHBUN, 1907: 60 (Makemo). — BOONE, 1934: 162, pl. 85 (Nuku Hiva). — FOREST & GUINOT, 1961: 126, fig. 117-119, 124bis, pl. 28, fig. 1 (Hikueru); 1962: 68 (Biogeography "Tahiti-Tuamotu, Marquesas"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 213 (Moorea); 1989: 111, 114 (Moorea, Tikehau). — NAIM, 1980a, annex 1, tab. 3 (Moorea). — KROPP & BIRKELAND, 1981: 629, tab. 5 (Moorea, Takapoto). — ODINETZ, 1983: 205 (Moorea, Tahiti, Takapoto). — GUINOT, 1985: 450 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti, Takapoto). — Not *Domecia hispida* - NOBILI, 1907: 404 (Hao) = *Domecia glabra* Alcock, 1899 fide FOREST & GUINOT (1961: 126).

Eriphia scabricula Dana, 1852

DISTRIBUTION. — Society (Tahiti); Tuamotu (Fakarava, Raroia).

REFERENCES. — *Eriphia scabricula* Dana, 1852a: 82; 1852b: 247; 1855, pl. 14, fig. 5a-b (Tahiti). — RATHBUN, 1907: 57 (Fakarava). — HOLTHUIS, 1953: 20 (Raroia). — MORRISON, 1954: 7 (Raroia). — FOREST & GUINOT, 1961: 123, fig. 113a-b, 114 (Tahiti); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 170, annex 1, tab. a (Tahiti). — GUINOT, 1985: 450 (List).

Eriphia sebana (Shaw & Nodder, 1803)

DISTRIBUTION. — Gambier (Mangareva, Tarauru-Roa); Society (Moorea, Tahiti); Tuamotu (Fakarava, Hikueru, Makatea, Makemo, Marutea South, Mataiva, Moruroa, Pukapuka, Rangiroa, Raroia, Taiaro, Takapoto, Tauere).

REFERENCES. — *Eriphia laevimana* Latr. - DANA, 1852b: 249; 1855, pl. 14, fig. 7a-c ("Honden" = Pukapuka, Society). — CANO, 1888: 171 (Tahiti). — NOBILI, 1907: 403 ("Gatavake" = Mangareva, Tarauru-Roa, Tauere). — *Eriphia sebana* - RATHBUN, 1907: 57 (Fakarava, Makatea, Makemo, "Manga Reva, Motus" = Mangareva?, Rangiroa). — SENDLER, 1923: 39 (Makatea). — HOLTHUIS, 1953: 20 (Raroia). — MORRISON, 1954: 7 (Raroia). — FOREST & GUINOT, 1961: 122, fig. 111a-b, 112 (Hikueru, Tahiti; Syn.). — 1962: 68 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 170, annex 1, tab. a, photograph 140a (bas) (Makatea, Mataiva, Moorea, Tahiti, Takapoto). — DELESALLE, 1985:

289 (Mataiva). — GUINOT, 1985: 450 (List). — SALVAT, 1986b: 72 (French Polynesia). — POUPIN, 1994a: 50, fig. 46, pl. 6a (Makatea, Marutea, Moruroa, Taiaro) - SYNONYMS - *Eriphia laevimana* Guérin, 1829-1844 in Latreille.

Globopilumnus globosus (Dana, 1852)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Ahe?, Makatea, Manihi?, Mataiva, Raraka, Raroia, Takapoto, Tikehau).

REFERENCES. — *Pilumnus globosus* Dana, 1852a: 81; 1852b: 236; 1855, pl. 13, fig. 10 (Raraka, Tahiti, "Waterland" = Ahe and/or Manihi). — RATHBUN, 1907: 56 (Tahiti). — *Pilumnus margaritatus* Ortmann, 1893b: 436 (Tahiti). — *Globopilumnus globosus* - BALSS, 1933: 7, pl. 1, fig. 1-2. (Tahiti) ? pro parte. — HOLTHUIS, 1953: 21 (Raroia). — GUINOT-DUMORTIER, 1960a: 99, fig. 1-2, 5-6 (Tahiti; Syn.). — FOREST & GUINOT, 1961: 121 (Tahiti); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 170, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List). — PEYROT-CLAUSADE, 1989: 113 (Tikehau). — Not *Pilumnus globosus* - NOBILI, 1907: 398 ("Marutea Vaitutaki") = *Liocarpilodes* sp. fide GUINOT-DUMORTIER (1960a: 100). — BOONE, 1934: 152, pl. 78 (Tahiti) = *Pilodius pugil* with a ? fide SERÈNE & LUOM (1959: 320).

Lydia annulipes (H. Milne Edwards, 1834)

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Tahiti); Tuamotu (Hikueru, Mataiva, Moruroa, Pukapuka, Raroia, Taiaro, Takapoto).

REFERENCES. — *Ruppellia annulipes* - DANA, 1852b: 246; 1855, pl. 14, fig. 4a-c (Tahiti). — *Lydia annulipes* - HOLTHUIS, 1953: 23 (Pukapuka, Raroia). — MORRISON, 1954: 7 (Raroia). — FOREST & GUINOT, 1961: 122, fig. 109a-b, 110 (Hikueru); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a (Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List). — POUPIN, 1994a: 47, fig. 43, pl. 5f (Hikueru, Moruroa, Nuku Hiva, Taiaro). — *Ruppellia granulosa* A. Milne Edwards, 1867: 279 (Marquesas; new synonymy, cf. Remark).

REMARK. — *Ruppellia granulosa* has been very rarely cited after its description. It is mentioned for the genus *Lydia* in SAKAI (1976: 477). It has been briefly described from a single specimen: "Cette espèce se distingue de *Ruppellia annulipes* par la profondeur des sillons qui limitent les lobules des régions. Ces lobules sont rugueux et granuleux. Les pattes antérieures sont également couvertes de grosses granulations peu élevées. Les autres caractères sont les mêmes que chez la *Ruppellia annulipes*". We have re-examined the type specimen (MNHN B9344, 1 ♂ 17x25) and, after its comparison with specimens of *L. annulipes* collected in the Marquesas and the Tuamotu (cf. in POUPIN, 1994a: 47), we consider that it is a junior synonym of this species.

Ozius hawaiensis Rathbun, 1902

DISTRIBUTION. — Marquesas (Nuku Hiva); Tuamotu (Fakarava, Makemo, Rangiroa).

REFERENCES. — *Ozius hawaiensis* - RATHBUN, 1907: 54 (Fakarava, Makemo, Nuku Hiva, Rangiroa). — FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu, Marquesas"). — GUINOT, 1985: 450 (List).

Ozius rugulosus Stimpson, 1858

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Tahiti).

REFERENCES. — *Ozius rugulosus* - HELLER, 1865: 22, pl. 3, fig. 1 (Tahiti). — PESTA, 1913: 47 (Tahiti). — FOREST & GUINOT, 1961: 121, fig. 107a-b, 108 (Tahiti); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 450 (List). — POUPIN, 1994a: 48, fig. 44, pl. 5g (Nuku Hiva, Tahiti).

Ozius tricarinatus Rathbun, 1907

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Tahiti).

REFERENCES. — *Ozius tricarinatus* Rathbun, 1907: 53, pl. 2, fig. 3 (Nuku Hiva, Tahiti). — FOREST & GUINOT, 1962: 68 (Biogeography "Tahiti-Tuamotu, Marquesas"). — GUINOT, 1985: 450 (List).

Ozius truncatus A. Milne Edwards, 1834

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Ozius lobatus* Heller, 1865: 21, pl. 2, fig. 4 (Tahiti) *fide* CHILTON & BENNETT (1929: 750).

Remark. — CHILTON & BENNETT consider, with doubt, that HELLER's species is the same as *Ozius truncatus*, but they do not mention "Tahiti", in the distribution of *O. truncatus*. It is possible, as often seen in HELLER's work, that this locality was mentioned by error.

Ozius tuberculatus H. Milne Edwards, 1834

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Tahiti).

REFERENCES. — *Ozius tuberculatus* - BOONE, 1934: 150, pl. 77 (Nuku Hiva). — POUPIN, 1994a: 49, fig. 45, pl. 5h (Nuku Hiva, Tahiti).

Pseudozius caystrus (Adams & White, 1848)

DISTRIBUTION. — Gambier (Kamaka, Tarauru-Roa); Society (Moorea, Tahiti); Tuamotu (Ahe and/or Manihi, Makatea, Mataiva, Moruroa, Raraka, Raroia, Taiaro, Takapoto).

REFERENCES. — *Pseudozius planus* Dana, 1852a: 81; 1852b: 233; 1855, pl. 13, fig. 6a-h (Raraka, "Waterland" = Ahe and/or Manihi). — EVANS, 1967: 409 ("Paumotu"; BM syntypes). — *Pseudozius caystrus* - NOBILI, 1907: 397 (Kamaka). — SENDLER, 1923: 38 (Makatea). — HOLTHUIS, 1953: 26 (Raroia). — MORRISON, 1954: 9 (Raroia). — FOREST & GUINOT, 1961: 125 (Tahiti, "Taraourou-roo" = Tarauru-Roa; Syn.); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 171, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 450 (List). — POUPIN, 1994a: 52, fig. 48, pl. 6c (Kamaka, Moruroa, Taiaro).

FAMILY GECARCINIDAE

Cardisoma carnifex (Herbst, 1794)

DISTRIBUTION. — Society (Bora Bora, Moorea, Raiatea, Tahiti, Tupai); Tuamotu (Ahe, Hao, Makatea, Mataiva, Nukutipipi, Pukarua, Rangiroa, Raroia, Taiaro, Takapoto, Tauere) - *Terrestrial*.

REFERENCES. — *Cardisoma obesum* Dana, 1851: 252; 1852b: 375; 1855, pl. 24, fig. 1 ("Peacock" = Ahe). — STIMPSON, 1858b: 100; 1907: 111 (Tahiti). — *Perigrapsus excelsus* Heller, 1862: 522; 1865: 50, pl. 5, fig. 1 (Tahiti). — *Cardisoma carnifex* - MIERS, 1886: 220 (Tahiti). — NOBILI, 1907: 407 (Bora Bora, Hao, Tahiti). — RATHBUN, 1907: 26 (Rangiroa, Tahiti). — SENDLER, 1923: 22 (Tahiti). — BOONE, 1934: 187, pl. 97-98 (Bora Bora). — SEURAT, 1934: 52 (Moorea, Tahiti, Tuamotu). — HOLTHUIS, 1953: 34 (Raroia). — CHABOIS L. & F., 1954: 92, unnumbered fig. (French Polynesia). — MORRISON, 1954: 2 (Raroia). — FOREST & GUINOT, 1961: 165 (Tahiti); 1962: 74 (Biogeography "Tahiti-Tuamotu"). — EDMONSON, 1962: 25 (Raiatea). — GUINOT, 1966a: 48 (French Polynesia); 1985: 454 (List). — TÜRKAY, 1973: 108 (Syn.). — SAKAI, 1976: 680 (Syn.). — MONTEFORTE, 1984: 174, annex 1, tab. a (Makatea, Mataiva, Moorea, Takapoto); 1987: 6 (Moorea). — DELESALLE, 1985: 288, 295 (Mataiva). — CHARLEUX, 1986: 80, photographs (French Polynesia). — BONVALLOT *et al.*, 1994: 78, photograph (Tuamotu). — BAGNIS & CHRISTIAN, 1983: 110-111, photograph (Tuamotu). — MERSCHARDT-SALVAT, 1991: 40 (Nukutipipi). — SALVAT F. & B., 1992: 5 (Nukutipipi). — POUPIN, 1994a: 53, fig. 49, pl. 6e (Pukarua, Tahiti, Taiaro, Tauere, Tupuai). — Without name - PARDON, 1992: 78, 79, photograph, double page (Tahiti, Papeete market) (det. according to the photograph).

Cardisoma hirtipes Dana, 1851

DISTRIBUTION. — Society (Tahiti) - *Terrestrial*.

REFERENCES. — *Cardisoma hirtipes* - HELLER, 1865: 35 (Tahiti). — TÜRKAY, 1974: 229, fig. 2, 12-13 (Tahiti). — GUINOT, 1985: 454, with a ? (List). — POUPIN, 1994a: 53 (Text).

***Cardisoma rotundum* Quoy & Gaimard, 1834**

DISTRIBUTION. — Marquesas (Nuku Hiva); Tuamotu (Tikehau) - *Terrestrial*.

REFERENCES. — *Cardisoma rotundum* - TÜRKAY, 1974: 234, fig. 1, 14 ("Tickahau-Atoll" = Tikehau). — POUPIN, 1994a: 54, fig. 50, pl. 6g (Nuku Hiva).

***Discoplax longipes* A. Milne Edwards, 1867**

DISTRIBUTION. — Tuamotu (Makatea).

REFERENCES. — *Discoplax longipes* - SENDLER, 1923: 23, pl. 20, 1a-b (Makatea). — FOREST & GUINOT, 1962: 74 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 454 (List).

***Epigrapsus politus* Heller, 1862**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Hikueru, Taiaro).

REFERENCES. — *Epigrapsus politus* Heller, 1862: 522 (Tahiti). — HOLTHUIS, 1953: 34 ("Taravao" = Tahiti). — FOREST & GUINOT, 1961: 162, fig. 176a-b (Hikueru; Syn.); 1962: 74 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Tahiti). — GUINOT, 1985: 454 (List). — POUPIN, 1994a: 55, fig. 51, pl. 6d (Hikueru). — *Nectograpsus politus* Heller, 1865: 57, pl. 5, fig. 3 (Tahiti). — Not *Epigrapsus politus* - NOBILI, 1907: 407 (Hikueru) = *Cyclograpsus integer* H. Milne Edwards, 1837 *fide* FOREST & GUINOT (1961: 162). — SEURAT, 1934: 58 (NOBILI's material).

FAMILY GRAPSIDAE

SUBFAMILY GRAPSINAE

***Geograpsus crinipes* (Dana, 1851)**

DISTRIBUTION. — Marquesas (Nuku Hiva); Society (Tahiti); Tuamotu (Makatea, Makemo, Mataiva, Moruroa, Pukapuka, Raroia, Taiaro, Takapoto).

REFERENCES. — *Geograpsus crinipes* - HELLER, 1865: 48 (Tahiti). — NOBILI, 1907: 404 (Pukapuka). — RATHBUN, 1907: 28 (Makemo). — SENDLER, 1923: 32 (Makatea). — SEURAT, 1934: 52 (Pukapuka). — HOLTHUIS, 1953: 29 (Raroia). — MORRISON, 1954: 9 (Raroia). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 453 (List). — SALVAT & RICHARD, 1985: 359, 360 (Takapoto). — POUPIN, 1994a: 57, fig. 53, pl. 6h (Moruroa, Nuku Hiva, Taiaro, Takapoto).

***Geograpsus grayi* (H. Milne Edwards, 1853)**

DISTRIBUTION. — Society (Tahiti); Tuamotu (Makatea, Nukutipipi, Raroia).

REFERENCES. — *Geograpsus grayi* - KINGSLEY, 1880c: 196 (Tahiti). — ORTMANN, 1894: 707 (Tahiti). — SENDLER, 1923: 32, pl. 21, fig. 6 (Makatea). — SEURAT, 1934: 52 (Tahiti). — HOLTHUIS, 1953: 30 (Raroia). — MORRISON, 1954: 11 (Raroia). — BANERDJEE, 1960: 159 (Tahiti). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea). — GUINOT, 1985: 453 (List). — MERSCHARDT-SALVAT, 1991: 40 (Nukutipipi). — SALVAT F. & B., 1992: 5 (Nukutipipi).

***Geograpsus stormi* De Man, 1895**

DISTRIBUTION. — Marquesas (Nuku Hiva).

REFERENCES. — *Geograpsus lividus stormi* de Man - RATHBUN, 1907: 29 (Nuku Hiva). — *Geograpsus stormi* - BANERDJEE, 1960: 167 (Syn.). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu, Marquesas"). — GUINOT, 1985: 453 (List). — POUPIN, 1994a: 58, fig. 54, pl. 7a (Nuku Hiva).

Grapsus depressus Heller, 1862

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Grapsus depressus* Heller 1862: 521 (Tahiti; to our knowledge this species has never been mentioned after its description).

Grapsus longitarsis Dana, 1851

DISTRIBUTION. — Society; Tuamotu (Fakarava, Hao, Hikueru, Makatea, Mataiva, Rangiroa, Raroia, Taiaro, Takapoto, Tikei).

REFERENCES. — *Grapsus longitarsis* Dana, 1851: 249; 1852b: 339; 1855, pl. 21, fig. 4a-d (Tuamotu). — RATHBUN, 1907: 28 (Fakarava, Rangiroa, Tikei). — HOLTHUIS, 1953: 31 (Raroia). — MORRISON, 1954: 7 (Raroia). — BANERJEE, 1960: 144, fig. 1b, 2h-n (Society, Raroia). — FOREST & GUINOT, 1961: 152, fig. 160a-b, 161, pl. 18, fig. 2 (Hikueru); 1962: 72 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 453 (List). — POUPIN, 1994a: 59, fig. 55, pl. 7b (Hao, Hikueru, Taiaro). — *Grapsus strigosus* - NOBILI, 1907: 404 (Hao) not *G. strigosus* (Herbst, 1799) synonym of *Grapsus albolineatus* Lamarck, 1818 in BANERJEE (1960: 147) = *G. longitarsis* fide FOREST & GUINOT (1961: 152).

Grapsus tenuicrustatus (Herbst, 1783)

DISTRIBUTION. — Gambier (Kamaka); Marquesas (Nuku Hiva); Society (Moorea, Tahiti); Tuamotu (Fakarava, Hao, Hikueru, Makatea, Makemo, Mataiva, Moruroa, Rangiroa, Raroia, Taiaro, Takapoto).

REFERENCES. — *Grapsus pictus* - DANA, 1852b: 336; 1855, pl. 21, fig. 1 (Tuamotu). — *Grapsus grapsus* - NOBILI, 1907: 404 (Hao, Kamaka) not *Grapsus grapsus* Linné, 1758 = *G. tenuicrustatus* fide FOREST & GUINOT (1961: 154). — *Grapsus grapsus tenuicrustatus* - RATHBUN, 1907: 27 (Fakarava, Makemo, Rangiroa). — *Grapsus gracilipes* - SENDLER, 1923: 31 (Makatea). — *Grapsus gracillimus* Sendler, 1923: 32, pl. 21, fig. 5 (Makatea). — *Grapsus tenuicrustatus* - HOLTHUIS, 1953: 31 (Raroia). — CHABOIS L. & F., 1954: 91, fig. 6 (French Polynesia). — MORRISON, 1954: 9 (Raroia). — BANERJEE, 1960: 134, fig. 1a, 2a, c-g (Raroia; Syn.). — FOREST & GUINOT, 1961: 154 (Hikueru; Syn.); 1962: 72 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 453 (List). — SALVAT & RICHARD, 1985: 360 (Takapoto). — PARDON, 1992: 82, photograph (Tuamotu). — POUPIN, 1994a: 60, fig. 56, pl. 7c (Nuku Hiva, Tahiti, Taiaro, Takapoto). — The following works refer also probably to this species: *Grapsus maculatus* - KINGSLEY, 1880c: 192 (Tahiti) = *G. grapsus* fide ORTMANN (1894: 703). — *Grapsus grapsus* - SENDLER, 1923: 30 (Makatea). — BOONE, 1934: 178, pl. 90 ("Anaho Bay" = Nuku Hiva). — CHEVALIER *et al.*, 1968: 95, 138 (Moruroa). — BAGNIS & CHRISTIAN, 1983: 112-113, photograph (Tuamotu). — *Grapsus albolineatus* - BONVALLOT *et al.*, 1994: 137, photograph, (Tuamotu; det. according to the photograph) - SYNONYMS - *Grapsus pictus* Latreille, 1802-1803; *Grapsus gracilipes* H. Milne Edwards, 1853; *Grapsus gracillimus* Sendler, 1923.

REMARK. — BANERJEE (1960: 139) mentions that *Grapsus grapsus* (Linné, 1758) is solely Atlantic. More recently, MANNING & HOLTHUIS (1981: 233) have also mentioned it from the Eastern Pacific.

Leptograpsus variegatus (Fabricius, 1793)

DISTRIBUTION. — Austral (Rapa); Marquesas.

REFERENCES. — *Leptograpsus variegatus* - DE MAN, 1890: 84 (Marquesas; with regard of *L. ansoni* H. Milne Edwards, 1853). — GRIFFIN, 1973: 461, fig. 1-6 (Syn.) - SYNONYMS - *Leptograpsus ansoni* H. Milne Edwards, 1853 - NEW MATERIAL - March 1995, Coll. and det. J. POUPIN (Rapa, Haurei bay, very common).

Metopograpsus messor (Forskål, 1775)

DISTRIBUTION. — Society (Bora Bora, Tahiti).

REFERENCES. — *Metopograpsus messor* - KINGSLEY, 1880c: 190 (Tahiti). — MIERS, 1886: 258 (Tahiti). — RATHBUN, 1907: 29 (Bora Bora). — PESTA, 1913: 61 (Tahiti). — GUINOT, 1985: 453 (List). — POUPIN, 1994a: 61 (Text).

Metopograpsus thukuhar (Owen, 1839)

DISTRIBUTION. — Austral (Tubuai); Gambier (Mangareva); Society (Moorea, Tahiti); Tuamotu (Hao, Taiaro).

REFERENCES. — *Metopograpsus thukuar* - STIMPSON, 1858b: 101; 1907: 114 [47] (Tahiti). — HELLER, 1865: 43 (Tahiti). — NOBILI, 1907: 404 (Hao, "Rikitea" = Mangareva). — SEURAT, 1934: 59 (Mangareva). — BANERJEE, 1960: 186, fig. 6f-g (Tahiti). — FOREST & GUINOT, 1961: 155, fig. 162, 167 (Tahiti); 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List). — MARQUET, 1988: 90, fig. 48, tab. 23; 1991: 130, tab. 1-2; 1993: tab. 1, 3 (Mangareva, Moorea, Tahiti, Tubuai). — POUPIN, 1994a: 61, fig. 57, pl. 7d (Hao, Mangareva, Tahiti, Taiaro).

Pachygrapsus fakaravensis Rathbun, 1907

DISTRIBUTION. — Tuamotu (Fakarava, Makatea, Mataiva, Taiaro, Takapoto).

REFERENCES. — *Pachygrapsus fakaravensis* Rathbun, 1907: 29, pl. 5, fig. 1, pl. 9, fig. 6, 6a (Fakarava). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Takapoto). — GUINOT, 1985: 453 (List). — POUPIN, 1994a: 62, fig. 58, pl. 7e (Taiaro).

Pachygrapsus minutus A. Milne Edwards, 1873

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Hikueru, Makatea, Mataiva, Takapoto, Tikehau).

REFERENCES. — *Pachygrapsus minutus* - FOREST & GUINOT, 1961: 155 (Hikueru, Tahiti); 1962: 72 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 25; 1977b: 213; 1985: 462 (Moorea); 1989: 113 (Tikehau). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 453 (List) - RELEVANT MATERIAL - *Pachygrapsus aff. minutus* - NAIM, 1980a: 55, annex 1, tab. 3 (Moorea; very small adult specimens, possibly of a new species). — GUINOT, 1985: 453 (List).

Pachygrapsus planifrons De Man, 1888

DISTRIBUTION. — Tuamotu (Fakarava, Raroia).

REFERENCES. — *Pachygrapsus planifrons* - HOLTHUIS, 1953: 31 (Raroia). — MORRISON, 1954: 7 (Raroia). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List). — *Pachygrapsus longipes* - RATHBUN, 1907: 30 (Fakarava) - SYNONYMS - *Pachygrapsus longipes* Rathbun, 1893 (in TESCH, 1918, p. 78; with uncertainty).

Pachygrapsus plicatus (H. Milne Edwards, 1837)

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Fakarava, Hikueru, Makatea, Makemo, Mataiva, Raroia, Taiaro, Takapoto).

REFERENCES. — *Pachygrapsus plicatus* - KINGLEY, 1880c: 200 (Tahiti). — RATHBUN, 1907: 29 (Fakarava, Makemo). — HOLTHUIS, 1953: 32 (Raroia). — MORRISON, 1954: 13 (Raroia). — FOREST & GUINOT, 1961: 154 (Hikueru); 1962: 72 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Moorea, Tahiti, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 453 (List). — SALVAT & RICHARD, 1985: 359 (Takapoto). — POUPIN, 1994a: 63, fig. 59, pl 7f (Hikueru, Taiaro).

Planes cyaneus Dana, 1851

DISTRIBUTION. — Austral (Neilson bank).

REFERENCES. — *Planes cyaneus* - NEW MATERIAL - April 1995, coll. J. POUPIN, det. A. CROSNIER (Neilson bank; on a drifting buoy with cirripeds).

SUBFAMILY VARUNINAE

***Hemigrapsus crenulatus* (H. Milne Edwards, 1837)**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Heterograpsus crenulatus* Guérin - NOBILI, 1907: 405 (Tahiti; in Paris, NOBILI's material is well recorded under *Hemigrapsus crenulatus* MNHN B12830). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — *Hemigrapsus crenulatus* (H. Milne Edwards) - BENNETT, 1964: 81 (Syn.). — GUINOT, 1985: 453 (List).

***Pseudograpsus albus* Stimpson, 1858**

DISTRIBUTION. — Tuamotu (Fakarava, Raroia).

REFERENCES. — *Pseudograpsus albus* - RATHBUN, 1907: 32 (Fakarava). — HOLTHUIS, 1953: 32 (Raroia). — MORRISON, 1954: 10 (Raroia). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List).

***Ptychognathus crassimanus* Finnegan, 1931**

DISTRIBUTION. — Marquesas - Freshwater.

REFERENCES. — *Ptychognathus crassimanus* Finnegan, 1931: 649 (Marquesas).

REMARK. — It seems that this species was never recorded after its description. In particular, it does not appear in the works by MARQUET (1988, 1991, 1993), who has only collected *Ptychognathus easteranus* (det. HOLTHUIS), in the Marquesas rivers.

***Ptychognathus easteranus* Rathbun, 1907**

DISTRIBUTION. — Austral (Rurutu); Marquesas (Hiva Oa) - Freshwater.

REFERENCES. — *Ptychognathus easteranus* - MARQUET, 1988: 90, fig. 48, tab. 23; 1991: 132, tab. 1-2; 1993: tab. 1, 3 (Hiva Oa, Rurutu).

***Ptychognathus intermedius* (de Man, 1879)**

DISTRIBUTION. — Society (Tahiti) - Freshwater.

REFERENCES. — *Ptychognathus intermedius* - ORTMANN, 1894: 711 (Tahiti). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List).

***Thalassograpus harpax* (Hilgendorf, 1892)**

DISTRIBUTION. — French Polynesia.

REFERENCES. — *Thalassograpus harpax* - GUINOT, 1985: 453 (List only; origin not found).

***Varuna litterata* (Fabricius, 1798)**

DISTRIBUTION. — Society (Moorea, Tahiti) - Fresh & Brackish water.

REFERENCES. — *Varuna litterata* - MARQUET, 1988: 90, fig. 48, tab. 3; 1991: 133, tab. 1-2; 1993: tab. 1, 3 (Moorea, Tahiti). — POUPIN 1994a: 67, fig. 63, pl. 8b (Tahiti).

SUBFAMILY SESARMINAE

***Chasmagnathus subquadratus* Dana, 1851**

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Chasmagnathus subquadratus* - ORTMANN, 1894: 728 (Tahiti). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List).

REMARK. — Species described by par DANA (1851: 251) from an uncertain locality "Novi-Zealandiae ? Novi-Hollandiae orientalis ?", not very often cited.

Cyclograpsus integer H. Milne Edwards, 1837

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Fakarava, Hikueru, Kaukura?, Mataiva, Raroia, Taiaro, Takapoto).

REFERENCES. — *Epigrapsus politus* - NOBILI, 1907: 407 (Hikueru, Kaukura) not *Epigrapsus politus* Heller, 1862 = *C. integer* fide FOREST & GUINOT (1961: 160). — *Cyclograpsus parvulus* - RATHBUN, 1907: 36 (Fakarava). — HOLTHUIS, 1953: 32 (Raroia). — MORRISON, 1954: 5 (Raroia). — *Cyclograpsus integer* - FOREST & GUINOT, 1961: 160, fig. 175a-c (Hikueru; Syn.); 1962: 74 (Biogeography "Tahiti-Tuamotu"). — MONTEFORTE, 1984: 172, annex 1, tab. a (Mataiva, Moorea, Tahiti, Takapoto). — GUINOT, 1985: 454 (List). — POUPIN, 1994a: 56, fig. 52, pl. 6f (Hikueru, Taiaro) - SYNONYMS - *Cyclograpsus parvulus* de Man, 1897.

Cyclograpsus longipes Stimpson, 1858

DISTRIBUTION. — Society (Tahiti); Tuamotu (Makemo, Marutea South, Raroia).

REFERENCES. — *Cyclograpsus longipes* - RATHBUN, 1907: 36 (Makemo, Tahiti). — HOLTHUIS, 1953: 32 (Raroia). — MORRISON, 1954: 5 (Raroia). — FOREST & GUINOT, 1961: 160 (Marutea South); 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 454 (List).

Labuanium trapezoideum (H. Milne Edwards, 1837)

DISTRIBUTION. — Society (Moorea, Raiatea, Tahiti) - Freshwater.

REFERENCES. — *Sesarma trapezoidea* - GUÉRIN-MÉNEVILLE, 1838: 14 (Tahiti). — SEURAT, 1934: 51 (Tahiti). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — MARQUET, 1988: 90, fig. 48, tab. 23; 1991: 133, tab. 1-2; 1993: tab. 1, 3 (Moorea, Tahiti). — *Sesarma (Sesarma) trapezoidea* - NOBILI, 1907: 405 (Tahiti). — EDMONSON, 1951: 237, fig. 33b (Raiatea, Tahiti). — FOREST & GUINOT, 1961: 157, fig. 164a-b, 165 (Tahiti). — *Sesarma (Sesarma) trapezoideum* - RATHBUN, 1907: 33 (Tahiti). — *Labuanium trapezoideum* - SERÈNE & SOH, 1970: 402, 406 (Syn.). — GUINOT, 1985: 454 (List).

REMARK. — *Labuanium rotundatum* (Hess, 1865) is also recorded in Polynesia by SAKAI (1976: 663; distribution only "Micronesia, Polynesia"; cited afterwards by GUINOT, 1985: 454). SAKAI must consider the Polynesia s.l., with about 10 states, including French Polynesia. It is doubtful that this species have been really collected in French Polynesia since we have not find any mention of it in TESCH (1917: 193), who gives a detailed distribution, or in SERÈNE & SOH (1970: 402, 406), when they have established the genus *Labuanium*. For the moment, it thus seems better to exclude it from the area.

Metasesarma rousseauxii granularis Heller, 1862

DISTRIBUTION. — Gambier (Tarauru-Roa); Society (Tahiti); Tuamotu (Hikueru).

REFERENCES. — *Metasesarma granularis* Heller, 1862: 522 (Tahiti). — *Metasesarma rugulosa* Heller, 1865: 65 (Tahiti; cf. Remark). — *Metasesarma rousseauxii* H. Milne Edwards - ? ORTMANN, 1894: 717 (Tahiti). — ? HOLTHUIS, 1953: 33 ("Taravao" = Tahiti). — *Sesarma (Metasesarma) rousseauxii* - NOBILI, 1907: 405 (Tarauru-Roa). — *Metasesarma rousseauxii granularis* - FOREST & GUINOT, 1961: 158, fig. 168, 169, 174a-b (Hikueru, "Papenoo" = Tahiti; Syn.); 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 454 (List).

REMARK. — FOREST & GUINOT (1961: 158) writes "*M. rousseauxii granularis*, décrit en 1862 de Tahiti par HELLER, qui, en 1865, substituait à ce nom, sans raison apparente, celui de *M. granulosa*, a été mis en synonymie avec *Metasesarma rousseauxii* H. Milne Edwards, 1853, par DE MAN (1889, p. 439)". The same authors recognise differences between H. MILNE EDWARDS' species and the specimens from Tahiti and Tuamotu, which they attribute to the subspecies *granularis* Heller. According to the location, ORTMANN's and HOLTHUIS' references should be also attributed to this subspecies.

Sarmatium crassum Dana, 1851

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Sarmatium crassum* - GUINOT, 1985: 454 (List; origin ?). — DAVIE, 1992: 81, fig. 1a, 2, 3a-c (Tahiti).

Sesarma angustifrons A. Milne Edwards, 1869

DISTRIBUTION. — Society (Moorea, Tahiti).

REFERENCES. — *Sesarma* (*Sesarma*) *angustifrons* - DE MAN, 1889: 432, pl. 10, fig. 10 (Tahiti). — NOBILI, 1907: 405 (Tahiti). — SEURAT, 1934: 51 (Tahiti). — *Sesarma angustifrons* - FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 454 (List). — MARQUET, 1988: 90, fig. 48, tab. 23; 1991: 133, tab. 1-2; 1993: tab. 1, 3 (Moorea, Tahiti).

Sesarma jacquinoti Ortmann, 1894

DISTRIBUTION. — Society (Tahiti).

REFERENCES. — *Sesarma jacquinoti* Ortmann, 1894: 718 (Tahiti). — FOREST & GUINOT, 1962: 72 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 454 (List).

SUBFAMILY PLAGUSIINAE

Percnon abbreviatum (Dana, 1851)

DISTRIBUTION. — Gambier (Mangareva); Society (Tahiti); Tuamotu (Raroia).

REFERENCES. — *Acanthopus abbreviatus* Dana, 1851: 252; 1852b: 373; 1855, pl. 23, fig. 11a-c (Tahiti). — *Percnon affinis* - NOBILI, 1907: 406 (Mangareva), *pro parte* not *P. affine* H. Milne Edwards, 1853 = *P. abbreviatum* fide FOREST & GUINOT (1961: 164). — *Percnon abbreviatum* - HOLTHUIS, 1953: 33 (Raroia). — MORRISON, 1954: 16 (Raroia). — EDMONSON, 1959: 195, fig. 25c, 26a-c (Syn.). — FOREST & GUINOT, 1961: 164 (Mangareva; Syn.); 1962: 74 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1966a: 48 (Raroia); 1985: 454 (List).

Percnon affine (H. Milne Edwards, 1853)

DISTRIBUTION. — Marquesas; Gambier (Mangareva); Society (Tahiti); Tuamotu (Makatea, Mataiva).

REFERENCES. — *Percnon affinis* - NOBILI, 1907: 406 ("Gatavake" = Mangareva), *pro parte* some sp. are *P. abbreviatum* — FOREST & GUINOT, 1961: 164 ("Gatavake" = Mangareva; Syn.); 1962: 74 (Biogeography "Tahiti-Tuamotu"). — *Percnon pilimanus* - BOONE, 1934: 181, pl. 92-94 (Tahiti). — FOREST & GUINOT, 1962: 74 (Biogeography "Tahiti-Tuamotu, Marquesas"). — *Percnon affine* - CROSNIER, 1965: 88 (Tuamotu; Syn.). — MONTEFORTE, 1984: 172, annex 1, tab. a, photograph p. 138a-bas (Makatea, Mataiva, Tahiti). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 454 (List) - SYNONYMS - *Percnon pilimanus* (A. Milne Edwards, 1873).

Percnon guinotae Crosnier, 1965

DISTRIBUTION. — Marquesas (Hiva Oa).

REFERENCES. — *Percnon guinotae* - NEW MATERIAL - February 1996, Coll. & det. J. POUPIN (Hiva Oa).

REMARK. — The presence of this species in French Polynesia was assumed in POUPIN (1994a: 64). It is confirmed here with a specimen collected in the Marquesas.

Percnon planissimum (Herbst, 1804)

DISTRIBUTION. — Gambier (Mangareva); Marquesas (Nuku Hiva); Society (Tahiti); Tuamotu (Fakarava, Hao, Makatea, Marutea South, Mataiva, Moruroa, Raraka, Taiaro, Tikehau).

REFERENCES. — *Acanthopus planissimus* - DANA, 1852b: 372 (Raraka, Tahiti). — HELLER, 1865: 51 (Tahiti). — *Acanthopus tenuifrons* H. Milne Edwards, 1853: 180 (Nuku Hiva). — *Percnon planissimum* - NOBILI, 1907: 406 (Hao, Mangareva, Marutea). — MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea, Mataiva, Tahiti). — PEYROT-CLAUSADE, 1989: 113 (Tikehau). — *Percnon planissimum* - RATHBUN, 1907: 37 (Fakarava). — PESTA, 1913: 64 (Tahiti). — EDMONDSON, 1959: 197, fig. 25c, 27a-c (Marquesas). — FOREST & GUINOT, 1961: 163 (Marutea South, Tahiti; Syn.);

1962: 74 (Biogeography "Tahiti-Tuamotu, Marquesas"). — CHEVALIER *et al.*, 1968: 95, 138 (Moruroa). — GUINOT, 1985: 454 (List). — POUPIN, 1994a: 64, fig. 60, pl. 8a (Mangareva, Taiaro).

Plagusia speciosa Dana, 1851

DISTRIBUTION. — Marquesas; Society (Tahiti); Tuamotu (Ahe and/or Manihi, Hao, Hikueru, Makatea, Makemo, Mataiva, Raroia, Taiaro, Takapoto).

REFERENCES. — *Plagusia speciosa* Dana, 1851: 252; 1852b: 369; 1865, pl. 23, fig. 9 ("Waterland" = Ahe and/or Manihi). — KINGSLEY, 1880c: 223 (Tahiti). — DE MAN, 1890: 89 (Tuamotu). — ORTMANN, 1894: 731 (Tuamotu). — NOBILI, 1907: 406 (Hao). — RATHBUN, 1907: 36 (Makemo). — SENDLER, 1923: 35 (Makatea). — BOONE, 1934: 185, pl. 95-96 (Tahiti). — HOLTHUIS, 1953: 34 (Raroia). — MORRISON, 1954: 16 (Raroia). — FOREST & GUINOT, 1961: 162, fig. 177a-c, 178 (Hao, Hikueru); 1962: 74 (Biogeography "Tahiti-Tuamotu, Marquesas"). — GUINOT, 1966a: 48 (Raroia); 1985: 454 (List). — MONTEFORTE, 1984: 172, annex 1, tab. a, photograph p. 138a-haut (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289 (Mataiva). — SALVAT & RICHARD, 1985: 362 (Takapoto). — POUPIN, 1994a: 65, fig. 61, pl. 7g (Hao, Hikueru, Taiaro).

REMARK. — *Plagusia immaculata* Lamarck, 1818 is erroneously cited from Tahiti and the Tuamotu by DAI & YANG (1991: 563). This error must come from a quick reading of EDMONSON (1959: 194), where "Tuamotus (type locality), Tahiti" is mentioned under *P. immaculata*, but for remarks concerning only *P. speciosa* Dana.

Plagusia tuberculata Lamarck, 1818

DISTRIBUTION. — Austral (Raevavae); Gambier (Kamaka, Mangareva); Marquesas (Nuku Hiva); Tuamotu (Makatea?).

REFERENCES. — *Plagusia squamosa* - NOBILI, 1907: 406 (Kamaka, Mangareva; inferred only: reference not retrieved in recent works, and material not found in Paris). — *Plagusia depressa tuberculata* - FOREST & GUINOT, 1962: 74 (Biogeography "Tahiti-Tuamotu"). — *Plagusia depressa* - ? MONTEFORTE, 1984: 172, annex 1, tab. a (Makatea) not *P. depressa* (Fabricius, 1775) = *P. tuberculata* cf. Remark. — *Plagusia tuberculata* - GUINOT, 1985: 454 (List). — POUPIN, 1994a: 66, fig. 62, pl. 7h (Nuku Hiva, Raevavae, Tuamotu) - SYNONYMS - *Plagusia squamosa* (Herbst, 1882) (cf. SAKAI, 1976: 676, then ALCOCK, 1900: 437).

REMARK. — *Plagusia depressa* (Fabricius, 1775) is a species of the tropical Atlantic (cf. DAWSON, 1987: 42) and cannot be MONTEFORTE's (1984) material which is more likely *P. tuberculata* (One specimen of this species is actually deposited in the CRIODE collections, Moorea, Coll. and det. MONTEFORTE).

FAMILY PINNOTHERIDAE

Pinnotherelia laevigata A. Milne Edwards & Lucas, 1843

DISTRIBUTION. — Marquesas (Nuku Hiva).

REFERENCES. — *Pinnotherelia laevigata* - RATHBUN, 1918: 181, fig. 115, pl. 39, fig. 1-3, pl. 40, fig. 1-2 (Marquesas "Tawhoe" = Taiohae, Nuku Hiva). — SCHMITT *et al.*, 1973: 125 (catalogue "Marquesas Islands").

FAMILY OCYPODIDAE

Macrophthalmus consobrinus Nobili, 1906

DISTRIBUTION. — Gambier (Mangareva) - Brackish water.

REFERENCES. — *Macrophthalmus consobrinus* Nobili, 1906a: 265; 1907: 408 (Mangareva). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List). — *Macrophthalmus parvimanus* - BARNES, 1977: 273, *pro parte*, only NOBILI's *consobrinus*, not *M. parvimanus* Guérin-Méneville, 1834 (cf. Remark).

REMARK. — According to BARNES (1977: 273) NOBILI's *Macrophthalmus consobrinus* is the same as *Macrophthalmus parvimanus* Guérin-Méneville, 1834. However, to check that point, a large sample of *M. consobrinus* has been recently

collected in the Gambier Islands, and it appears that NOBILI's species is valid, and closely related to *M. convexus* Stimpson, 1858 (POUPIN, in study).

Macrophthalmus convexus Stimpson, 1858

DISTRIBUTION. — Society (Bora Bora, Tahiti) - Brackish water.

REFERENCES. — *Macrophthalmus convexus* - ORTMANN, 1894: 745 (Tahiti). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List) - NEW MATERIAL - Coll. and det. J. POUPIN (Bora Bora).

Macrophthalmus serenei (Serène, 1983)

DISTRIBUTION. — Tuamotu (Tikehau).

REFERENCES. — *Macrophthalmus serenei* - NEW MATERIAL - Coll. C. HILY, det. J. POUPIN (Tikehau).

Ocypode ceratophthalma (Pallas, 1772)

DISTRIBUTION. — Gambier? (Mangareva); Society (Scilly, Tahiti); Tuamotu (Fakarava, Makatea, Makemo, Mataiva, Marutea South?, Rangiroa, Raroia, Taiaro, Takapoto, Tikehau).

REFERENCES. — *Ocypode urvillei* Guérin-Méneville, 1829: pl. 1, fig. 1, 1a-b; 1838: 9 (Tahiti). — OWEN, 1839: 80 ("Low Islands" = Tuamotu). — NOBILI, 1907: 407 *pro parte* (Marutea?, cf. Remark sous *O. pallidula*). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453, with a ? (List). — *Ocypode ceratophthalma* - STIMPSON, 1858b: 100 [46]; 1907: 108, pl. 12, fig. 2 (Tahiti). — ORTMANN, 1897a: 364 (Syn.). — RATHBUN, 1907: 26 (Fakarava, Makemo, Rangiroa). — SENDLER, 1923: 21 (Tahiti). — HOLTHUIS, 1953: 28 (Raroia). — MORRISON, 1954: 9 (Raroia). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — SAKAI & TÜRKAY, 1976: 86, fig. 13 (Syn.). — MONTEFORTE, 1984: 173, annex 1, tab. a (Makatea, Mataiva, Takapoto). — DELESALLE, 1985: 289, 303 (Mataiva). — GUINOT, 1985: 453 (List). — HARMELIN-VIVIEN, 1985: 239 (Tikehau). — SALVAT & RICHARD, 1985: 359 (Takapoto). — POUPIN, 1994a: 68, fig. 64, pl. 8c-d (Tahiti, Taiaro, Scilly). — *Ocypode cordimana* - KINGSLEY, 1880b: 186 (Tahiti) not *O. cordimana* Desmaret, 1825 = *O. urvillei*, synonym of *O. ceratophthalma*, fide ORTMANN (1897a: 366). — ? *Oxypode* (*sic*) - CHARLEUX, 1986: 81, photograph (French Polynesia), det. according to the photograph.

Ocypode cordimana Desmaret, 1825

DISTRIBUTION. — Society (Bora Bora, Tahiti).

REFERENCES. — *Ocypode cordimana* - BOONE, 1934: 191, pl. 99-100 (Bora Bora). — FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — GUINOT, 1985: 453 (List). — POUPIN, 1994a: 69, fig. 65, pl. 8e (Tahiti). — Not *Ocypode cordimana* - KINGSLEY, 1880b: 186 (Tahiti) *fide* ORTMANN (1897a: 366) = *O. urvillei*, synonym of *O. ceratophthalma*.

REMARK. — BOONE mentions this species in the Tuamotu ("Paumotus") after RATHBUN (1907: 26). In fact it is *O. ceratophthalma* that RATHBUN has cited from this archipelago.

Ocypode pallidula Jacquinot 1852

DISTRIBUTION. — Gambier (Aukena, Mangareva); Tuamotu (Marutea South?, Moruroa).

REFERENCES. — *Ocypode pallidula* Jacquinot, 1852, pl. 6, fig. 1a (Mangareva). — SAKAI & TÜRKAY, 1976: 87, fig. 14-15 ("Rikitea" = Mangareva; type material, Syn.). — JONES, 1988: 34 (Syn.). — POUPIN, 1994a: 70, fig. 66, pl. 8f (Aukena). — *Ocypode cordimana* (Junior) - JACQUINOT & LUCAS, 1853: 64 (Mangareva) not *O. cordimana* Desmaret, 1825 = *O. pallidula* fide SAKAI & TÜRKAY (1976: 87). — *Ocypode urvillei* - NOBILI, 1907: 407, *pro parte* ("Rikitea" = Mangareva, Marutea?). — SEURAT, 1934: 52 (Mangareva; NOBILI's material) - The two references, not *O. urvillei* Guérin-Méneville, 1829, synonym of *O. ceratophthalma* = *O. pallidula* (cf. Remark). — *Ocypode laevis* - CHEVALIER et al., 1968: 109 (Moruroa) - NEW MATERIAL - Coll. B. SALVAT, 1966, det. J. POUPIN (Moruroa) - SYNONYMS - *Ocypode laevis* Dana, 1852.

REMARK. — After ORTMANN (1897a: 366), *Ocypode pallidula*, was usually considered as the same as *O. urvillei* (= *O. ceratophthalma*). Its validity was re-established by SAKAI & TÜRKAY (1976). The material mentioned in NOBILI (1907)

and SEURAT (1934), under *O. urvillei* has been collected at Marutea South and Mangareva. The specimens from this second island are in fact *O. pallidula* (verification in the collections of Paris: 1 sp. labelled "Ocypoda urvillei Guér., Seurat coll. 1905, Bouvier dét., G. Nobili vérif. 1906", MNHN B11841, is a real *O. pallidula*). The specimens from Marutea South could reasonably be *O. cerathophtalma*, very common in the Tuamotu.

Uca chlorophtalmus crassipes (Adams & White, 1848)

DISTRIBUTION. — Austral (Raevavae); Marquesas; Society (Bora Bora, Maupiti, Raiatea, Tahiti) - Brackish water.

REFERENCES. — *Gelasimus latreillei* H. Milne Edwards, 1852: 114, pl. 4, fig. 20, 20a (Bora Bora). — *Gelasimus pulchellus* Stimpson, 1858b: 100 [46]; 1907: 107, pl. 15, fig. 1 (Tahiti). — *Gelasimus gaimardi* - HELLER, 1865: 38 (Tahiti). — *Uca chlorophtalmus* - NOBILI, 1907: 408 ("Taravao" = Tahiti). — *Uca gaimardi* - RATHBUN, 1907: 26 (Bora Bora, Tahiti). — HOLTHUIS, 1953: 29 ("Taravao" = Tahiti). — CRANE, 1957: 74, 78 (Bora Bora, Raiatea, Tahiti). — FOREST & GUINOT, 1961: 140, fig. 140-145, 153, 156a-b (Tahiti); 1962: 70 (Biogeography "Tahiti-Tuamotu"). — *Gelasimus (Uca) chlorophtalmus* - SEURAT, 1934: 60 (Tahiti). — *Uca (Amphiuca) chlorophtalmus crassipes* - CRANE, 1975: 98, 102, 599, fig. 13-14, 26c, 31c, 37h, 39a-b, 56c, 60 l-m, 68a-b, 81g, 83a, 99, pl. 15 a-f, 46b (Bora Bora, Raiatea, Tahiti, Marquesas p. 599; Syn.). — GUINOT, 1985: 453 (List). — POUPIN, 1994a: 71, fig. 67, pl. 8g (Maupiti, Raevavae, Tahiti) - SYNONYMS - *Uca gaimardi* H. Milne Edwards, 1852.

Uca tetragonon (Herbst, 1790)

DISTRIBUTION. — Austral (Rapa); Gambier (Mangareva); Society (Bora Bora, Raiatea, Tahiti); Tuamotu (Moruroa, Napuka, Raroia).

REFERENCES. — *Gelasimus duperreyi* Guérin-Méneville, 1829, pl. 1, fig. 2, 2a (Bora Bora). — *Gelasimus tetragonon* - GUÉRIN-MÉNEVILLE, 1838, pl. 1, fig. 2, 2a (Bora Bora) *pro parte*. — HELLER, 1865: 37 (Tahiti). — KINGSLEY, 1880a: 143, pl. 9, fig. 11 (Tahiti). — Miers, 1886: 243 (Tahiti). — DE MAN, 1891: 24, pl. 2, fig. 6 (Tahiti). — ORTMANN, 1894: 754 (Tahiti). — *Uca tetragonon* - NOBILI, 1907: 408 ("Rikitea, Gatavake" = Mangareva). — RATHBUN, 1907: 26 (Bora Bora). — HOLTHUIS, 1953: 29 (Raroia). — MORRISON, 1954: 8 (Raroia). — CRANE, 1957: 79 (Bora Bora). — *Gelasimus (Uca) tetragonon* - SEURAT, 1934: 59 (Mangareva). — *Uca tetragonum* - FOREST & GUINOT, 1962: 70 (Biogeography "Tahiti-Tuamotu"). — *Uca (Thalassuca) tetragonon* - CRANE, 1975: 77, 81, 596, fig. 37d, 63a-b, 81f, 82e, 99, pl. 13 (Bora Bora, Raiatea, Raroia, Tahiti; Syn.). — GUINOT, 1985: 453 (List). — POUPIN, 1994a: 72, fig. 68, pl. 8h (Mangareva, Moruroa, Napuka, Rapa).

REMARK. — FOREST & GUINOT (1962: 70) mention *U. dussumieri* (H. Milne Edwards) from French Polynesia, probably after ORTMANN (1894; *Gelasimus dussumieri*, Tahiti, p. 755). ORTMANN's reference is cited by CRANE (1975) for two subspecies: *Uca (Deltuca) dussumieri spinata* (specimens from Java and Singapore) and *Uca (Deltuca) dussumieri dussumieri* (specimens from the Philippines and Mindanao). ORTMANN's Tahitian *U. dussumieri* do not appear in CRANE, who clearly indicates (p. 437) that *Uca dussumieri* does not occur in French Polynesia. ORTMANN's "*U. dussumieri*", if they exist, must probably be, either *U. chlorophtalmus*, or *U. tetragonon*.

FAMILY CRYPTOCHIRIDAE

Cryptochirus coralliodytes Heller, 1861

DISTRIBUTION. — Tuamotu (Makatea, Marutea South, Marokau).

REFERENCES. — *Cryptochirus coralliodytes* - NOBILI, 1907: 409 (Marutea South, Marokau). — SENDLER, 1923: 41 (Makatea). — FOREST & GUINOT, 1962: 74 (Biogeography "Tahiti-Tuamotu"). — SEURAT, 1934: 60 (Marokau, Marutea South). — GUINOT, 1985: 454 (List). — KROPP, 1988: 873 (Revision of this species but without mention of French Polynesia).

Hapalocarcinus marsupialis Simpson, 1859

DISTRIBUTION. — Society (Moorea, Tahiti); Tuamotu (Takapoto).

REFERENCES. — *Hapalocarcinus marsupialis* - KROPP & BIRKELAND, 1981: 629, tab. 5 (Moorea, Takapoto). — ODINETZ, 1983: 29, 205 (Moorea, Tahiti, Takapoto). — GUINOT, 1985: 454 (List). — ODINETZ-COLLART & RICHER DE FORGES, 1985: 201 (Moorea and/or Tahiti).

FAMILY HYMENOSOMATIDAE

Elamena mathaei (Desmaret, 1825)

DISTRIBUTION. — Tuamotu (Tikehau).

REFERENCES. — *Elamena mathaei* - PEYROT-CLAUSADE, 1989: 113 (Tikehau).

FAMILY INCERTAE SEDIS

Daira perlata (Herbst, 1790)

DISTRIBUTION. — Society (Moorea, Raiatea, Tahiti); Tuamotu (Makatea, Mataiva, Hao, Hikueru, Raroia, Taiaro, Takapoto, Tikehau).

REFERENCES. — *Daira perlata* - NOBILI, 1907: 392 (Hao). — RATHBUN, 1907: 44 (Tahiti). — SENDLER, 1923: 38 (Makatea). — BOONE, 1934: 129, pl. 69 (Raiatea, Tahiti). — HOLTHUIS, 1953: 19 (Raroia). — CHABOIS L. & F., 1954: 91, fig. 4 (French Polynesia). — MORRISON, 1954: 15 (Raroia). — FOREST & GUINOT, 1961: 119 (Hikueru, Tahiti); 1962: 68 (Biogeography "Tahiti-Tuamotu"). — PEYROT-CLAUSADE, 1977a, annex of the species: 26; 1977b: 212 (Moorea); 1989: 112, 115 (Moorea, Tikehau). — MONTEFORTE, 1984: 170, annex 1, tab. a, photograph p. 137a (Makatea, Mataiva, Moorea, Takapoto); 1987: 9 (Moorea). — DELESALLE, 1985: 289, 305 (Mataiva). — GUINOT, 1985: 453 (List). — SALVAT & RICHARD, 1985: 362 (Takapoto). — SALVAT, 1986b: 72, photograph (French Polynesia). — POUPIN, 1994a: 51, fig. 47, pl. 6b (Hikueru, Taiaro).

REMARK. — This species has sometimes been classified in the Zalasiinae Serène, 1968 (cf. SAKAI, 1976: 513).

Parapleurophrycoides roseus Nobili, 1906

DISTRIBUTION. — Tuamotu (Marutea South).

REFERENCES. — *Parapleurophrycoides roseus* Nobili, 1906a: 264; 1907: 402, pl. 2, fig. 5 (Marutea; gen. nov. and sp. nov. described from a very small specimen, 1.3x1.7mm).

REMARK. — For this species, and the following, FOREST & GUINOT (1962: 41) write: "...nous les considérons comme des juvéniles difficilement identifiables."

Platyozius perpusillus Nobili, 1906

DISTRIBUTION. — Tuamotu (Hao).

REFERENCES. — *Platyozius perpusillus* Nobili, 1906a: 264; 1907: 401 (Hao; described from a very small specimen, 1.45x1.75mm).

REMARK. — In SAKAI (1976: 535), *Platyozius* Borradaile, 1902 is the same as *Eucrate* de Haan, 1835 (Goneplacidae).

DISCUSSION

NUMBER OF POLYNESIAN SPECIES

A total of 401 littoral or sublittoral species are recorded in this work. The number by infra-order and family is computed on table 1. The Brachyura clearly prevail, with 78% of the species, followed by the Anomura (18%), and the Palinura (4%).

Within the crabs, the Xanthidae account for 123 species, distributed in 5 main subfamilies: Liomerinae, Actaeinae, Zoziminae, Xanthinae, and Chlorodiinae. The Portunidae account for 54 species, including 6 recorded for the first time from determinations made by MOOSA and CROSNIER: *Portunus macrophthalmus*, *P. orbitosinus*, *Thalamita danae*, *T. macropus*, *T. mitsiensis*, and *T. philippinensis*; half of the sublittoral species belong to this family. Two other families are also well represented, the Grapsidae, with 35 species, and the Trapeziidae, with 20 species. From these two families, *Percnon guinotae*, *Planes cyaneus* and *Quadrella maculosa*, are recorded for the first time. With respect to the list presented 10 years ago by GUINOT (1985), excluding the species mentioned erroneously, or not fully determined, about 60 species are added to the Polynesian Brachyura, and approximately a hundred, if the deep species are included.

Within the Anomura, the Diogenidae account for 40% of the species with three main genera: *Calcinus*, *Clibanarius*, and *Dardanus*. *Calcinus guamensis*, *C. imperialis*, and *Dardanus australis*, are now recorded in French Polynesia. The porcellanids account for about 25% of the species, the single genus *Petrolisthes* representing 12 species out of 17. The Albuneidae are represented by only one species, *Albunea speciosa*, which was previously thought to be endemic from the Hawaiian islands.

Only 14 palinurids are recorded in French Polynesia, of which 2 only by larvae (*Palinurellus wieneckii* and *Arctides regalis*).

SPECIES ERRONEOUSLY RECORDED IN FRENCH POLYNESIA

Fourteen species have been erroneously recorded in French Polynesia. They are: *Panulirus ornatus* (cf. under *P. versicolor*), *Panulirus polyphagus* (cf. under *P. pascuensis*), *Dynomene sinense* (cf. under *D. praedator*), *Ashtoret granulosa* (cf. under *A. picta*), *Lophozozymus incisus* (cf. under *L. superbus*), *Labuanium rotundatum* (cf. under *L. trapezoideum*), *Plagusia immaculata* (cf. under *P. speciosa*), *Uca dussumieri* (cf. under *U. tetragonon*), *Pachygrapsus transversus*, *Dotilla fenestrata*, *Ocypode macrocera*, *Ocypode platytarsis*, and two freshwater crabs (Potamonidae).

Pachygrapsus transversus Gibbes, 1850 is recorded from Tahiti by KINGSLEY (1880c: 199) (Tahiti). According to HOLTHUIS & GOTTLIED (1958: 102) this record is obviously false, *P. transversus* being an Atlantic species. This conclusion is later supported by MANNING & HOLTHUIS (1981: 235), who report however the species in the Pacific, but only along the American coasts.

The ocypodid *Dotilla fenestrata* Hilgendorf, 1869, is listed by GUINOT (1985: 453) after KROPP & BIRKELAND (1981). It is probably a mistake, because this species is not mentioned in that work, and, to our knowledge, has never been reported, elsewhere, from French Polynesia. Two other ocypodids, with an uncertain status, are also erroneously reported from Tahiti, by HELLER (1865: 42): *Ocypode macrocera* (H. Milne Edwards, 1837) and *Ocypode platytarsis* (H. Milne Edwards, 1852) (see ORTMANN, 1897a: 362).

HELLER has mentioned two potamonids crabs in Tahiti: *Thelphusa wüllerstorfi*, described as a new species in 1862 (p. 520); and *Thelphusa leschenaudii* (H. Milne Edwards, 1853) (in HELLER, 1865: 32). RATHBUN (1904: 287) places these two references under a single species *Potamon (Potamon) hydrodromus* (Herbst, 1796) and writes (p. 289): "il est douteux que cette espèce ou quelqu'autre habite Tahiti". Since the recent works by MARQUET (1988, 1991, 1993), who has intensively sampled the freshwater Polynesian fauna, it is almost certain that the Potamonidae are not represented in French Polynesia.

IMPROVEMENT AND CORRECTION OF THIS LIST

This bibliographic compilation is of course tentative and certainly does not account for all the species living in French Polynesia. When new collections become available, other species will undoubtedly be added to the present list. Moreover, despite a careful research, it is possible that a few works, recording additional species, have passed undetected. Right now, several species listed here deserve a particular attention, either because their presence in French Polynesia remained to be confirmed, or because their identification, or taxonomic status, are uncertain.

Twelve species of this list could have been erroneously recorded from French Polynesia. They are known only by larvae (*Palinurellus wieneckii*, *Arctides regalis*), are mentioned with doubt, or in an ambiguous way (*Petrolisthes militaris*, *Porcellana mitra*, *Porcellana monilifera*, *Ozius truncatus*), appear only in a part "Distribution", the origin of the French Polynesian material remaining unknown (*Neoliomera insularis*, *Gaillardiellus rueppelli*, *Macromedaeus distinguendus*, *Thalassograpus harpax*), or, are cited from French Polynesia only because of the large geographic distribution of the species, without real collections in the field (*Schizophrys aspera*, *Aethra scruposa*).

For a score of species the revision of the material would be particularly interesting. They are *Coenobita cavipes*, *Petrolisthes rufescens*, *Dardanus guttatus*, *Matuta victor*, *Charybdis annulata*, *Portunus pelagicus*, *Lophozozymus pictor*, *Trapezia septata*, *Chasmagnathus subquadratus*, and *Cryptochirus coralliodytes*, corresponding to isolated, usually old references, never again cited in recent revisions; *Enoplometopus holthuisi* and *Neopetrolisthes maculatus*, cited only in non-taxonomic books; *Neoliomera pubescens*, *Actaea calculosa*, *Forestia depressa*, *Forestia scabra*, and *Actumnus globulus*, for which it is clearly indicated, in systematic studies, that the revision of this material is necessary; and *Trapezia formosa*, re-examined in Paris and perhaps belonging to a new species (P. CASTRO, personal communication). Furthermore, about 30 additional species, recorded in ecological works, with sometimes only provisional determinations, could be added to the above mentioned species (cf. for example, *Calcinus minutus*, *Galathea aff. amamiensis*, *Liomera laperousei*, *Paramedaeus simplex*, *Actaea aff. glandifera*, *Zozymodes xanthoides*...).

The status of 15 species is doubtful and must be revised. They are: *Coenobita carnescens* and *C. olivieri*, that could respectively be synonyms of *C. perlatus* and *C. spinosus*; *Galathea latirostris* and *Cryptodromia coronata*, two species whose exact identity remains to be defined; *Thalamita minuscula*, *Parapleurophrycoides roseus*, and *Platyozius perpusillus*, described from very small specimens which could only be the juveniles of more common species; *Xanthias tetaodon*, possibly a synonym of *X. nitidulus*; *Eitisus punctatus*, described only after drawings, the corresponding material being lost; and *Porcellana monilifera*, *Actaeomorpha alvae*, *Portunus alexandri*, *Grapsus depresso*, *Ptychognathus crassimanus*, and *Sesarma jacquinoti*, 6 species described from French Polynesia a long time ago, and never recorded since.

BIOGEOGRAPHY

For the Brachyura, FOREST & GUINOT (1962) have already established that the French Polynesian fauna is a part of the Indo-West Pacific fauna. Located at the eastern limit of this area, French Polynesia is characterised by a lower diversity than in the Indo-Malaysian area, considered as the origin from where the Indo-West Pacific fauna has extended. This assumption can be verified here for other groups. For the Astacidea and Palinuridea, HOLTHUIS (1991) records 27 western-pacific species (zone 71 = Malaysia, Indonesia, Philippines, New Guinea), collected within the first 100m, against only 14 in French Polynesia. For the genus *Clibanarius*, RAHAYU & FOREST (1992) report 20 Indonesian species, against only 7 in this work. For other diogenids the comparaison with the Indonesian fauna (in RAHAYU, 1992) reveal the following discrepancies: *Aniculus* 4 vs 2 species, *Calcinus*, 23 vs 10 species, *Dardanus* 10 vs 8 species, and *Diogenes* 18 vs 1 species.

In our list, 21 species are known only from French Polynesia. For most of them, it is doubtful that they are real endemic forms: 3 have been described from very small specimens which could be juveniles of other species (*Parapleurophrycoides roseus*, *Platozius perpusillus*, *Thalamita minuscula*); 2 are sublittoral to deep species recently described from material collected with difficulty by the use of deep traps (*Alainodaeus rimatara*, *Medaeus grandis*); and 8 are some of the doubtful species already mentioned (*Coenobita carnescens*, *Eitisus punctatus*, *Porcellana monilifera*, *Actaeomorpha alvae*, *Portunus alexandri*, *Grapsus depressus*, *Ptychognathus crassimanus*, and *Sesarma jacquinoti*). Concerning this last group, let us recall that *Ruppellia granulosa*, never recorded since its description from the Marquesas, is here proposed as a junior synonym of the *Lydia annulipes*, widely distributed in the Indo-West Pacific. The 8 remaining species, which could be true endemic forms, are the following: *Parribacus holthuisi*, *Micropagurus polynesiensis*, *Nucia rosea*, *Nursia mimetica*, *Lissocarcinus elegans*, *Acanthophrys cristimanus*, *Ozius tricarinatus*, and *Macrophthalmus consobrinus*. However, it is likely that some of them are distributed at least as far as the Western Polynesia, and that they will be recorded there when more collections are available. As an example, *Calcinus nitidus*, formerly considered as endemic from Tahiti, has been recently reported in the Samoa (POUPIN, 1994a).

The French Polynesian fauna could be related to the fauna of the Hawaiian islands, which are of similar origin and geomorphology. A comparison between the two areas remains difficult because no detailed list of the Hawaiian fauna is yet available. We however notice that 4 species, *Albunea speciosa*, *Charybdis hawaiensis*, *Panopeus pacificus* and *Sesarma angustifrons*, are still known only from these two areas. ELDREDGE & MILLER (1995) have recently published the number of Hawaiian species, by Infra-Order. The same calculation, made after our work, is compared with the data of these authors in table 2. The most obvious result is that the French Polynesian fauna is almost twice as rich as the Hawaiian fauna. Nevertheless, the fauna of the Hawaiian islands has been well studied, with some important works, like RATHBUN (1906) or EDMONSON (1959, 1962). This discrepancy could come from the as exhaustive as possible approach that we have adopted in our compilation. In particular we have included: about 30 species recorded only in ecological works, with sometimes incomplete or only preliminary determinations; 45 doubtful species (uncertainty about the taxonomic status or the effective presence in French Polynesia); 92 species belonging to the deep fauna, which has been intensively studied and collected recently. Yet, if these three groups are eliminated from the calculation, the result remains still clearly higher in French Polynesia (326 species vs 246). Thus, this observation would reveal a real difference between the two areas, the number of species being greater in French Polynesia. A similar result has been observed for the barnacles by NEWMAN (1986). This author explains the relatively low diversity of the Hawaiian islands by their great isolation, in particular if the low islands are excluded from the chart (see *opt. cit.*, fig. 2), and a settlement from the southern hemisphere, in part from French Polynesia.

Table 1 - Number of French Polynesian species, by Infra-Order and Family. The littoral and sublittoral species come from the above compilation. The deep species have been published in a previous work, updated here in Appendices 1 (The 12 sublittoral to deep species, listed in both works, are counted with the littoral and sublittoral species).

INFRA-ORDER	FAMILY	Littoral and sublittoral	Deep species (>100m)	TOTAL
ASTACIDEA & PALINURIDEA	Enoplometopidae	1	1	2
	Palinuridae	6	3	9
	Synaxidae	2		2
	Scyllaridae	5	1	6
<i>subtotal</i>		14	5	19
ANOMURA	Coenobitidae	8		8
	Diogenidae	31	6	37
	Paguridae	6	1	7
	Parapaguridae		10	10
	Chirostylidae		2	2
	Galatheidae	8	19	27
	Porcellanidae	17		17
	Albuneidae	1		1
	Hippidae	3		3
<i>subtotal</i>		74	39	113
BRACHYURA	Dromiidae	4	1	5
	Homolidae		8	8
	Latreillidae		1	1
	Dynomeneidae	3	1	4
	Raninidae	2	2	4
	Poupinidae		1	1
	Calappidae	5	2	7
	Leucosiidae	4	3	7
	Majidae	13	3	16
	Parthenopidae	5	2	7
	Eumedonidae	1		1
	Cancridae		1	1
	Geryonidae		2	2
	Goneplacidae		3	3
	Portunidae	54	1	55
	Xanthidae	123	14	137
	Trapeziidae	20		20
	Pilumnidae	9		9
	Carpiliidae	2		2
	Menippidae	13		13
	Gecarcinidae	5		5
	Grapsidae	35		35
	Pinnotheridae	1		1
	Ocypodidae	8		8
	Cryptochiridae	2		2
	Hymenosomatidae	1		1
	<i>Incertae Sedis</i>	3	3	6
<i>subtotal</i>		313	48	361
TOTAL		401	92	493

Table 2 - Total number of species in French Polynesia (this work, deep species included) and Hawaii (data of ELDREDGE & MILLER, 1995: 7).

	French Polynesia	Hawaii
Palinuridae & Astacidae	19	14
Anomura	113	43
Brachyura	361	189
Total	493	246

Few regional distinctions are observed within the French Polynesian Islands. The differences presented in table 3, where the number of species is calculated by archipelago, mainly indicate differences in the number of explorations, and amount of collections.

Table 3 - Number of species by archipelago. The number in parenthesis indicates the species known only from the corresponding islands.

ARCHIPELAGO	TOTAL
Austral	28 (7)
Gambier	79 (10)
Marquesas	82 (25)
Society	282 (103)
Tuamotu	226 (49)

The Society Islands, with Tahiti, almost inevitable during a stay in French Polynesia, is of course the best studied place. Nearly as many species are known from the Tuamotu Islands, which illustrates the importance of the collections made during the voyages of either the *US Exploring Expedition* (Ahe, Manihi, Rangiroa, Reao...) or the *Albatross* (Fakarava, Rangiroa...), and those made by SEURAT (Hao, Marutea South...), RANSON (Hikueru), or MORRISSON (Raroia). The atolls of this archipelago, without rivers, are of course not colonized by fresh or brackish water species such as: the Hippidae of the genus *Hippa*, the Grapsidae Varuninae (*Ptychognathus*, *Varuna*), the Grapsidae Sesarminae (*Labuanium*, *Sesarma*), and some Ocypodidae such as *Uca chlorophthalmus* (cf. POUPIN, 1994a: 71), *Macrophthalmus convexus* and *M. consobrinus* (POUPIN, in study).

At the southeastern part of Polynesia, the small archipelago of the Gambier Islands, despite its isolation, has been relatively well sampled, thanks to the collections made by SEURAT during his several years stay at Mangareva. In comparison, the Marquesas, far more extended, in the vicinity of the equator, remain poorly known. Affected by particular hydrological conditions, at the origin of the remarkable absence of a reef barrier, they, nonetheless, do not have any regional characteristics. The 25 species that, within French Polynesia, are still known only from these islands, are almost always very common in the Indo-West Pacific. The only exception seems to be the absence of the coconut crab (*Birgus latro*) whose presence was never verified during our frequent visits and inquiries.

With only 28 species, the Austral islands have been clearly less sampled than the others. The northern islands (Maria, Rimatara, Rurutu, Tubuai, and Raevavae) have a fauna similar to the rest of Polynesia, and a lot of common species, although not yet recorded here, have been observed during our

stays: *Panulirus penicillatus*, *Coenobita perlatus*, *Calcinus laevimanus*, *Pachygrapsus plicatus*, *Cardisoma carnifex*... and even, in the mouths of the rivers of Raevavae, the big portunid *Scylla serrata*, common in the high Society Islands. Far more south, at the southern limit of the tropical area, the island of Rapa, and the islets of Marotiri, are affected by the particular climatic conditions prevailing in that place. The perceptible decrease of the water temperature allows only a feeble growth of coral, without a barrier reef. Ashore, the vegetation is affected by a milder climate, the coconut tree being almost absent. This particular situation has an influence on the decapod fauna. Some species, very common elsewhere, have never been found after several visits and inquiries: the coenobite *Coenobita perlatus*, the coconut crab *Birgus latro* and the land crab *Cardisoma carnifex*. In contrast, at least one species, the lobster *Panulirus pascuensis*, has settled in these islands, whereas it is absent in the northern Polynesia.

LITERATURE CITED

- ALCOCK, A., 1900. — Material for a carcinological fauna of India, 6. The Brachyura Catometopa or Grapsidae. *Journal Asiatic Society of Bengal, Calcutta*, **69**, part 2 (3): 279-456.
- 1905. — Catalogue of the Indian decapod crustacea in the collection of the Indian Museum. Part II, Anomura, Fasciculus I, Pagurides. Calcutta: 1-197, pl. 1-16.
- ANONYMOUS, 1979. — Essai de pêche de fond à l'extérieur du récif. Pose de casiers par le *Tainui*. Association Territoire de la French Polynesia, Centre National d'Exploitation des Océans, Tahiti, Centre Océanologique du Pacifique COP/D: 1-14, fig. 1-7.
- BABA, K., 1991. — Crustacea Decapoda: *Alainius* gen. nov., *Leiogalathea* Baba, 1969, and *Phylladiorhynchus* Baba, 1969 (Galatheidae) from New Caledonia. In: A. CROSNIER (éd.), Résultats des campagnes MUSORSTOM, Volume 9. *Mémoires du Muséum national d'Histoire naturelle*, Paris (A), **152**: 479-491, fig. 1-5.
- BABLET, J. P., 1972. — Echinodermes et Crustacés. In: BABLET, J. P & O. CAYET (éds), Le Monde vivant des atolls, Chap IV. *Publication de la Society des Océanistes*, Musée de l'homme, Paris, **28**: 29-35, pl. 10-11.
- BAGNIS, R. & E. CHRISTIAN, 1983. — Guide sous-marin de Tahiti. *Les éditions du Pacifique*, 1-152, illustrated in colour.
- BALSS, H., 1933. — Beiträge zur Kenntnis der Gattung *Pilumnus* (Crustacea Decapoda) und verwandter Gattungen. *Capita Zoologica*, **4** (3): 1-47, fig. 1-7, pl. 1-7.
- 1935. — Brachyura of the Hamburg Museum Expedition to South-Western Australia, 1905. *Journal of the Royal Society of Western Australia*, **21**: 113-151, fig. 1-3, pl. 13.
- 1938. — Die Dekapoda Brachyura von Dr. Sixten BOCKS Pazifik-Expedition 1917-1918. *Göteborgs Kungl. Vetenskaps och Vitterhets Samhälles*, **5** (7): 1-85, fig. 1-18, pl. 1-2.
- BANERJEE, S. K., 1960. — Biological results of the *Snellius* Expedition. XVIII. The genera *Grapsus*, *Geograpsus*, and *Metopograpsus* (Crustacea Brachyura). *Temminckia*, **10**: 132-199, fig. 1-6.
- BARNES, R. S., 1977. — Concluding contributions towards a revision of, and a key to, the genus *Macrobrachium* (Crustacea, Brachyura). *Journal of Zoology*, London, **182**: 267-280, fig. 1-3.
- BATE, C. S., 1888. — Report on the Crustacea Macrura collected by the H.M.S. *Challenger* during the years 1873-76. *Report on the Scientific Results of the Voyage of H.M.S. Challenger, Zoology*, **24** (i-ix): 1-942, fig. 1-76; 1 vol. planches: pl. 1-150.

- BENNETT, E. W., 1964. — The marine fauna of New Zealand: Crustacea Brachyura. *New Zealand Department of Scientific and Industrial Research Bulletin*, **153** (22): 1-120, fig. 1-141.
- BONVALLOT, J., LABOUTE, P., ROUGERIE, F. & E. VIGNERON, 1994. — Les atolls des Tuamotu. *Editions de l'ORSTOM*, Paris: 1-296, illustrated in colour.
- BOONE, L., 1934. — Scientific Results of the World Cruise of the Yacht *Alva*, 1931, William K. VANDERBILT, Commanding. Crustacea: Stomatopoda and Brachyura. *Bulletin of the Vanderbilt Marine Museum*, Huntington, L.I. New York, USA, **5**: 1-210, pl. 1-109.
- BOONE, L., 1935. — *Ibid.* Crustacea: Anomura, Macrura, Euphausiacea, Isopoda, Amphipoda, and Echinodermata: Asteroidea and Echinoidea, **6**: 1-263, fig. 1-13, pl. 1-96.
- BOUVIER, E. L. & G. SEURAT, 1905. — *Eumodon convictor*, crabe commensal d'un oursin. *Compte Rendus Hebdomadaires des séances de l'Académie des Sciences*, Paris, **140**: 629-631.
- BOWMAN, T. E. & G. ABELE, 1982. — Classification of the recent crustacea. In: L. G. ABELE (ed.) *The Biology of Crustacea*, Academic Press, New York, **1** (1): 1-27.
- BUITENDIJK, A. M., 1960. — Biological results of the *Snellius* Expedition. XXI - Brachyura of the families Atelecyclidae and Xanthidae. *Temminckia*, **10**: 252-338, fig. 1-9.
- CANO, G., 1888. — Crostacei raccolti dalla R. Corvetta *Caracciolo* nel viaggio intorno al globo durante gli anni 1881-84. *Bollettino della società di Naturalisti in Napoli*, serie 1, **2** (2): 160-184, fig. 1-3.
- CHABOIS L. & F., 1954. — Petite Histoire Naturelle des Etablissements Français de l'Océanie. II. Zoologie. *Editions Paul Lechevalier*, Paris: 1-137, planches.
- CHARLEUX, M., 1986. — Les invertébrés: le tupa.. et les autres. In: C. GLEIZAL (éd.) *Encyclopédie de la Polynésie*, Tome 2, Flore et faune terrestres. C. GLEIZAL/Multipress: 80-81, illustré.
- CHEVALIER, J. P., DENIZOT, M., MOUGIN, J. L., PLESSIS, Y. & B. SALVAT, 1968. — Etude géomorphologique et bionomique de l'atoll de Moruroa (Tuamotu). *Cahiers du Pacifique*, **12**: 1-144, fig. 1-62, pl. 1-24.
- CHILTON, C. & E. W. BENNETT, 1929. — Contribution for a revision of the Crustacea Brachyura of New Zealand. *Transactions of the New Zealand Institute*, **59**: 731-778.
- CLARK, P. F. & B. S. GALIL, 1993. — A revision of the xanthid genus *Pilodius* Dana, 1851 (Crustacea, Brachyura, Xanthoidea). *Journal of Natural History*, **27**: 1119-1206, fig. 1-44.
- CRANE, J., 1957. — Basic patterns of display in fiddler crabs (Ocypodidae, Genus *Uca*). *Zoologica*, **42** (2): 69-82, fig. 1-4, pl. 1.
- 1975. — Fiddler Crabs of the World. Ocypodidae: Genus *Uca*. Princeton University Press: 1-736, fig. 1-101, pl. 1-50.
- CROSNIER, A., 1962. — Crustacés Décapodes Portunidae. *Faune de Madagascar*, **16**: 1-154, fig. 1-256, pl. 1-13.
- 1965. — Crustacés Décapodes Grapsidae et Ocypodidae. *Faune de Madagascar*, **18**: 1-143, fig. 1-260, pl. 1-11.
- 1975. — Sur les *Caphyra* (Crustacea, Decapoda, Portunidae) de l'océan Indien occidental et de la mer Rouge. *Bulletin du Muséum national d'Histoire naturelle*, 3ème série, n°304, Zoologie, **214**: 743-764, fig. 1-7.
- 1984. — Famille des Carpiliidae et des Menippidae. In: R. SERÈNE, 1984: Crustacés Décapodes Brachyoures de l'océan Indien occidental et de la mer Rouge. Xanthoidea: Xanthidae et Trapeziidae. *ORSTOM Collection Faune Tropicale*, **24**: 299-313, fig. 208-243, pl. 45-48.
- 1995. — *Pleurocolpus boileauai*, nouveau genre et une espèce nouvelle de Polynésie française (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle*, 4ème série (A), **17** (3-4): 245-251, fig. 1, pl. 1.

- CURTISS, A., 1938. — A short zoology of Tahiti in the Society islands. *Germantown*, i-xvi, 1-193 (not seen).
- DAI, A. & S. YANG, 1991. — Crabs of the China seas. China Ocean Press Beijing; Springer-Verlag Berlin, Heidelberg, New York, Tokyo (English Edition): 1-682, fig. 1-295, pl. 1-74.
- DANA, J. D., 1851: — *Conspectus Crustaceaorum quae in Orbis Terrarum circumnavigatione, Carolo WILKES e Classe Reipublicae Faederatae Duce, lexit et descriptis J. D. DANA. Proceedings of the Academy of Natural Sciences of Philadelphia*, 5: 247-254, 267-272.
- 1852a. — *Ibid.* 6: 73-86.
- 1852b. — Crustacea. United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842, Part I, 13: i-viii, 1-685.
- 1852c. — *Conspectus Crustaceaorum, ect.* Conspectus of the Crustacea of the Exploring Expedition under Capt. WILKES, U.S.N., including the Paguridae continued, the Megalopidae, and the Macroura. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 6: 6-28.
- 1855. — Crustacea. Atlas. 13: 1-27, pl. 1-96.
- DAVIE, P. J., 1990. — A new genus and species of marine crayfish, *Palibythus magnificus*, and new records of *Palinurellus* (Decapoda, Palinuridae) from the Pacific Ocean. *Invertebrate Taxonomy*, 4 (4): 685-695., fig. 1-5.
- 1992. — Revision of *Sarmatium* Dana (Crustacea, Brachyura, Sesarminae) with description of three new species. *Memoirs of the Queensland Museum*, 32 (1): 79-97, fig. 1-8.
- 1993. — Deepwater Xanthid crabs from French Polynesia (Crustacea, Decapoda, Xanthoidea). *Bulletin du Muséum national d'Histoire naturelle*, Paris, 4ème série (A), 1992 (1993), 14 (2): 501-561, fig. 1-12, pl. 1-13.
- 1995. — Two new species of *Nanocassiope* from the western Pacific (Crustacea, Decapoda, Xanthidae). *Bulletin du Muséum national d'Histoire naturelle*, Paris, 4ème série (A), 17 (1-2): 201-210, fig. 1-2.
- DAWSON, E. W., 1987. — A key to the world species of *Plagusia* (Crustacea: Brachyura), with a new record of *P. depressa tuberculata* Lamarck from New Zealand. *National Museum of New Zealand Records*, 3 (4): 37-45, fig. 1.
- DELESALLE, B., 1985. — Mataiva atoll, Tuamotu archipelago. In: B. DELESALLE, R. GALZIN & B. SALVAT (eds). *Proceedings of the Fifth International Coral Reef Congress*, Tahiti, 27 May - 1 June 1985, 1: 269-307, fig. 1-44.
- EDMONSON, C. H., 1951. — Some Central Pacific Crustaceans. *Occasional Papers of Bernice P. Bishop Museum*, Honolulu, Hawaii, 20 (13): 183-243, fig. 1-38.
- 1959. — Hawaiian Grapsidae. *Occasional Papers of Bernice P. Bishop Museum*, Honolulu, Hawaii, 22 (10): 153-202, fig. 1-27.
- 1962. — Hawaiian Crustacea: Gonoplacidae, Pinnotheridae, Cymopoliidae, Ocypodidae, and Gecarcinidae. *Ibid.*, 23 (1): 1-27, fig. 1-10.
- ELDREDGE, L. G., 1967. — Catalog of Invertebrate type specimens. *Pacific Scientific Information Center*, 21 p.
- EVANS, A. C., 1967. — Syntypes of Decapoda described by William STIMPSON and James DANA in the collections of the British Museum (Natural History). *Journal of Natural History*, 1: 399-411.
- FINNEGAN, S., 1931. — Report on the Brachyura collected in Central America, the Gorgona and Galapagos islands, by Dr. CROSSLAND on the St George Expedition to the Pacific, 1924-25. *Linnean Society's Journal, Zoology*, 37 (255): 607-673, fig. 1-6.
- FOREST, J., 1951. — Remark sur quelques Paguridae du genre *Calcinus* à propos de la description de deux espèces nouvelles de Polynésie Orientale: *Calcinus seurati* et *Calcinus spicatus*. *Extrait du Bulletin de la Société Zoologique de France*, 76 (1-2): 83-89, fig. 1-18.

- 1953a. — Crustacés Décapodes marcheurs des îles de Tahiti et des Tuamotu. I. Paguridea. *Bulletin du Muséum national d'Histoire naturelle*, Paris, 2ème série, **25** (5): 441-450, fig. 1-9.
- 1953b. — *Ibid.* I. Paguridea (suite), **25** (6): 555-561, fig. 10.
- 1954a. — *Ibid.* I. Paguridea (suite), **26** (1): 71-79, fig. 15-19.
- 1954b. — *Ibid.* II. Scyllaridea, **26** (3): 345-352, fig. 25-26.
- 1956a. — Crustacés décapodes Paguridae d'Océanie Française. *Proceedings of the eighth Pacific Science congress*, **3** (A): 1053-1076.
- 1956b. — Sur *Calcinus nitidus* Heller et *C. rosaceus* Heller (Crust. Paguridae). *Bulletin du Muséum national d'Histoire naturelle*, 2ème série, **28** (2): 218-227, fig. 1-8.
- 1957. — Les Pagures du Viet-Nam. I - Le genre *Diogenes* Dana. *Bulletin du Muséum national d'Histoire naturelle*, 2ème série, 1956 (1957), **28** (6): 524-532, fig. 1-15.
- 1962. — Sur un *Dardanus* des Hawaii et de Madagascar, *D. brachyops* sp. nov. (Crustacea, Paguridea, Diogenidae). *Bulletin du Muséum national d'Histoire naturelle*, 2ème série, **34** (5): 365-370, fig. 1-3.
- 1984. — Révision du genre *Aniculus* Decapoda Diogenidae. *Crustaceana*, supplément n° 8: 1-91, fig. 1-89.
- 1995. — Crustacea Decapoda Anomura: Révision du genre *Trizopagurus* Forest, 1952 (Diogenidae) avec l'établissement de deux genres nouveaux. In: A. CROSNIER (éd.), Résultats des campagnes MUSORSTOM volume 13. *Mémoires du Muséum national d'Histoire naturelle*, **163**: 9-149, fig. 1-30.
- FOREST, J. & D. GUINOT, 1961. — Crustacés Décapodes Brachyoures de Tahiti et des Tuamotu. In: Expédition française sur les récifs coralliens de la Nouvelle-Calédonie. Volume préliminaire, Editions de la Fondation Singer Polignac, Paris, IX-XI: 1-195, fig. 1-178, pl. 1-18.
- 1962. — Remark biogéographiques sur les crabes des archipels de la Society et des Tuamotu. *Cahiers du Pacifique*, **4**: 41-75, fig. 1, tab. 1-2.
- GALIL, B. S., 1985. — *Tetraloides*, a new genus of coral-inhabiting crabs. *Crustaceana*, **50**: 68-77.
- 1986a. — *Quadrella* (Brachyura, Xanthoidea, Trapeziidae), review and revision. *Journal of Crustacean Biology*, **6** (2): 275-293, fig. 1-8.
- 1986b. — On the identity of *Tetralia cinctipes* Paulson, 1875 (Decapoda, Brachyura). *Crustaceana*, **51** (1): 97-102, fig. 1-3.
- GALIL, B. S. & C. LEWINSOHN, 1984. — On the taxonomic status of *Trapezia tigrina* Eydoux & Souleyet, 1842 (Decapoda, Brachyura). *Crustaceana*, **46** (2): 166-175, fig. 1.
- 1985a. — On the taxonomic status of *Trapezia areolata* Dana, and *Trapezia septata* Dana (Decapoda, Brachyura). *Crustaceana*, **48** (3): 286-293, fig. 1-4.
- 1985b. — On the taxonomic status of *Trapezia rufopunctata* (Herbst) and *Trapezia flavopunctata* Eydoux & Souleyet (Decapoda, Brachyura). *Crustaceana*, **48** (2): 209-217, fig. 1-7.
- GALIL, B. S. & M. TAKEDA, 1986. — Ressurection of the genus *Jonesius* and establishment of a new genus: commensal crabs associated with corals from Indo-Pacific ocean. *Bulletin of the National Science Museum*, serie A, Zoology, Tokyo, **12** (4): 163-171, fig. 1-8.
- GALIL, B. S. & P. F. CLARK, 1990. — Crustacea Decapoda: Notes on trapeziid crabs from New Caledonia including description of two new species. In: A. CROSNIER (éd.), Résultats des campagnes MUSORSTOM, vol. 6. *Mémoires du Muséum national d'Histoire naturelle*, Paris (A), **145**: 369-388, fig. 1-6.
- 1994. — A revision of the genus *Matuta* Weber, 1795 (Crustacea, Brachyura, Calappidae). *Zoologische Verhandelingen*, Leiden, **294**: 1-55, fig. 1-7, pl. 1-14.

- GALZIN, R. & J. P. POINTIER, 1985. — Moorea island, Society archipelago. In: B. DELESALLE, R. GALZIN & B. SALVAT (eds.). *Proceedings of the Fifth International Coral Reef Congress*, Tahiti, 27 May - 1 June 1985, 1: 73-102, fig. 1-19.
- GARTH, J. S. & H. S. KIM, 1983. — Crabs of the family Xanthidae (Crustacea, Brachyura) from the Philippines islands and adjacent waters based largely on collections of the U.S. Fish Commission steamer *Albatross* in 1908-1909. *Journal of Natural History*, 17: 663-729, fig. 1-14.
- GARTH, J. S., HAIG, J. & J. W. KNUDSEN, 1987. — Crustacea Decapoda (Brachyura and Anomura) of Enewetak Atoll. In: DEVANEY, D. M., REESE, E. S., BURCH, B. L., & P. HELFRICH (eds.), *The Natural History of Enewetak Atoll. Volume II, chapter 23 - Biogeography and Systematics. United States Department of Energy, Office of Energy Research, Office of Health and Environmental Research, Ecological Research Division*: 235-261.
- GIBSON-HILL, M. A., 1948. — The Robber Crab. *The Malayan Nature Journal*, 3 (1): 10-14.
- GORDON, I., 1953. — On the Puerulus stage of some spiny lobsters (Palinuridae). *Bulletin of the British Museum, (Natural History) Zoology*, 2 (2): 17-42, fig. 1-9.
- GRIFFIN, D. J., 1973. — A revision of the two southern temperate shore crabs *Leptograpsus variegatus* (Fabricius) and *Plagusia chabrus* (Linnaeus) (Crustacea, Decapoda, Grapsidae). *Journal of the Royal Society of New Zealand*, 3 (3): 415-440, fig. 1-15.
- GRIFFIN, D. J. & H. A. TRANTER, 1986. — The Decapoda Brachyura of the *Siboga* Expedition. Part 8, Majidae. *Siboga Expedition, Monographie*, 39, C4, Livr. 148: 1-335, fig. 1-112, pl. 1-22.
- GUÉRIN-MÉNEVILLE, F. E., 1829-1830. — Atlas. In: *Voyage autour du monde, exécuté par ordre du roi, sur la corvette La Coquille, pendant les années 1822, 1823, 1824 et 1825. Arthus Bertrand, libraire éditeur, rue de Hauteville, n°23, Paris, crustacés*: pl. 1-5.
- 1838. — Crustacés, arachnides et insectes. *Ibid. Zoologie*, vol. 2, pt. 2, div. 1: xii + 319. Crustacés: 1-47.
- GUINOT, D., 1962. — Sur une collection de Crustacés Décapodes Brachyoures des îles Maldives et de la Mer Rouge (Expédition "Xarifa" 1957-1958). *Kieler Meeresforschungen*, Kiel, 18 (2): 231-244, fig. 1-17.
- 1964. — Crustacés décapodes brachyoures (Xanthidae) des campagnes de la *Calypso* en mer rouge (1952), dans le golfe persique, et à l'île Aldabra (1954). *Mémoires du Muséum National d'Histoire Naturelle, nouvelle série (A)*, 32 (1): 1-108, i-iii, fig. 1-57, pl. 1-12.
- 1966a. — Les crabes comestibles de l'Indo-Pacifique. Editions de la Fondation Singer-Polignac, Paris, deuxième volume préliminaire: 1-145, fig. 1-23, pl. 1-10.
- 1966b. — Recherches préliminaires sur les groupements naturels chez les crustacés décapodes brachyoures. I. Les affinités des genres *Aethra*, *Osachila*, *Hepatus*, *Hepatella* et *Actaeomorpha*. *Bulletin du Muséum national d'Histoire naturelle, 2ème série*, 38 (5): 744-762, fig. 1-24.
- 1967. — *Ibid. 38* (6), 1966 (1967): 828-845, fig. 25-41.
- 1969. — Sur divers Xanthidae notamment sur *Actaea* de Haan et *Paractaea* gen. nov. (Crustacea, Decapoda, Brachyura). *Cahiers du Pacifique*, 13: 1-267, fig. 1-36.
- 1976. — Constitution de quelques groupes naturels chez les crustacés décapodes brachyoures. I - La super-famille des Bellioidea, et trois sous-familles de Xanthidae (Polydectinae Dana, Trichiinae de Haan, Actaeinae Alcock). *Mémoires du Muséum national d'Histoire naturelle, nouvelle série (A)*, 97: 1-308, fig. 1-47, pl. 1-19.
- 1979. — Morphologie et phyllogénèse des brachyoures. *Mémoires du Muséum national d'Histoire naturelle*, Paris, (A), 112: 1-351, fig. 1-70, pl. 1-27.
- 1985. — Crustacea. In: G. Richard (ed.), French Polynesia coral reefs, fauna and flora. A first compendium of French Polynesian sea-dwellers. *Proceedings of the Fifth International Coral Reef Congress*, Tahiti 27 May - 1 June 1985, 1: 446-455.

- GUINOT, D. & B. RICHER DE FORGES, 1995. — Crustacea Decapoda Brachyura: Révision de la famille des Homolidae de Haan, 1839. In: A. CROSNIER (éd.), Résultats des campagnes MUSORSTOM volume 13. *Mémoires du Muséum national d'Histoire naturelle*, **163**: 283-517, fig. 1-76.
- GUINOT-DUMORTIER, D., 1960a. — Les espèces indo-pacifiques du genre *Globopilumnus* (Crustacea, Brachyura, Xanthidae). *Mémoires de l'Institut Scientifique de Madagascar*, série F, 1959 (1960), **3**: 97-119, fig. 1-14.
- 1960b. — Révision des genres *Euxanthus* Dana et *Hypocolpus* Rathbun (Crustacea, Decapoda, Brachyura). Remark sur les cavités sous-hépatiques et les coaptations des *Hypocolpus*. *Mémoires du Muséum national d'Histoire naturelle*, nouvelle série (A), **20** (2): 153-218, fig. 1-5, pl. 1-12.
- HAIG, J., 1964. — Porcellanid crabs from the Indo-West Pacific, Part I. *Videnskabelige meddelelser fra Dansk Naturhistorisk Forening*, **126**: 355-386, fig. 1-4.
- 1966. — A review of the indo-west Pacific species of genus *Pachycheles* (Porcellanidae, Anomura). *Proceedings of Symposium on Crustacea, India*, Part I: 285-294.
- 1970. — The status of *Remipes testudinarius* Latreille, and designation of a neotype for *Hippa adactyla* J. C. Fabricius (Decapoda, Hippidae). *Crustaceana*, **19** (3): 287-296.
- 1974. — A review of the australian crabs of family Hippidae (Crustacea, Decapoda, Anomura). *Memoirs of the Queensland Museum*, **71** (1): 175-189, fig. 1-5, pl. 6.
- 1979. — Expédition *Rumphius* II (1975), Crustacés parasites, commensaux, etc., V. Porcellanidae (Crustacea, Decapoda, Anomura). *Bulletin du Muséum national d'Histoire naturelle*, 4ème série, section A, **1** (1): 119-136, fig. 1-25.
- 1981. — Three new species of *Petrolisthes* (Decapoda, Anomura, Porcellanidae) from the indo-west Pacific. *Journal of Crustacean Biology*, **1** (2): 265-271, fig. 1-3.
- 1983. — Porcellanidae (Decapoda, Anomura) from the Seychelles, western Indian Ocean. *Crustaceana*, **45** (3): 279-289.
- 1992. — Hong Kong's porcellanid crabs. In: B. MORTON (ed.), The marine flora and fauna of Hong Kong and southern China, III. *Proceedings of the fourth international marine biological workshop* (Hong Kong 11-29 April 1989): 303-327, fig. 1-20.
- HAIG, J. & E. E. BALL, 1988. — Hermit crabs from north australian and eastern indonesian waters (Crustacea, Decapoda, Anomura, Paguroidea) collected during the 1975 ALPHA HELIX expedition. *Records of the Australian Museum*, **40** (3): 151-196, fig. 1-15.
- HAIG, J. & R. K. KROPP, 1987. — *Petrolisthes eldredgei*, a new porcellanid crab from the indo-west Pacific, with redescription of two related species. *Micronesica*, **20**: 171-186, fig. 1-6.
- HARALD, A. R., 1967. — The National Geographic Society Smithsonian-Bishop Museum Marquesas Expedition, August 15 - November 21, 1967. Mimeographed Report: 1-42.
- HARMELIN-VIVIEN, M., 1985. — Tikehau atoll, Tuamotu archipelago. In: B. DELESALLE, R. GALZIN & B. SALVAT (eds). *Proceedings of the Fifth International Coral Reef Congress*, Tahiti, 27 May - 1 June 1985, **1**: 211-256, fig. 1-74.
- HELLER, C., 1862. — Neue Crustaceen, gesammelt während der Weltumsegelung der k.k. Fregatte *Novara*. Zweiter vorläufiger Bericht. *Verhandlungen der kaiserlich-königlichen, Zoologisch-Botanischen Gesellschaft*, Wien, **12**: 519-528.
- 1865. — Die Crustaceen. Reise der öesterreichischen Fregatte *Novara* um die Erde in den Jahren 1857-1859 unter den Befehlen des Commodore B. von WÜLLERSTORF-URBAIR. *Zoologischer*, **2** (3): 1-280, pl. 1-25.
- HENDERSON, J. R., 1888. — Report on the Anomura collected by the H.M.S. *Challenger* during the years 1873-1876. *Report on the Scientific Results of the Voyage of H.M.S. Challenger*, Zoology, **27** (1): 1-211, pl. 1-21.

- HOLTHUIS, L. B., 1946. — Biological results of the *Snellius* expedition. XIV - The Decapoda, Macrura of the *Snellius* expedition. 1. The Stenopidae, Nephropsidae, Scyllaridae and Palinuridea. *Temminckia*, 7: 1-178, pl. 1-11.
- 1953. — Enumeration of the Decapod and Stomatopod Crustacea from Pacific coral islands. *Atoll Research Bulletin*, 24: 1-66.
 - 1981. — A new species of *Scyllarus* (Crustacea, Decapoda, Palinuridea) from the Pacific Ocean. *Bulletin du Muséum national d'Histoire naturelle*, 4ème série (A), 3 (3): 847-853, fig. 1-2.
 - 1985. — A revision of the family Scyllaridae (Crustacea, Decapoda, Macrura). I - Subfamily Ibacinae. *Zoologische Mededeelingen*, Leiden, 218: 1-130, fig. 1-27.
 - 1991. — Marine Lobsters of the world. An annotated and illustrated catalogue of species of interest to Fisheries known to date. *FAO Fisheries Synopsis*, 125 (13): 1-292, fig. 1-459.
 - 1993. — *Scyllarus rapanus*, a new species of locust lobster from the South Pacific (Crustacea, Decapoda, Scyllaridae). *Bulletin du Muséum national d'Histoire naturelle*, Paris, 4ème série (A), 15 (1-4): 179-186, fig. 1-3.
- HOLTHUIS, L. B. & E. GOTTLIEB, 1958. — An annotated list of the decapod crustacea of the mediterranean coast of Israel, with an appendix listing the Decapoda of the eastern mediterranean. *Bulletin of the Research Council of Israel*, 7b (1-2): 1-126, fig. 1-15.
- JACQUINOT, H., 1852. — In: HOMBON et JACQUINOT, Voyage au Pôle Sud et dans l'Océanie sur les corvettes l'*Astrolabe* et la *Zélée* pendant les années 1837-1838-1839-1840 sous le commandement de M. DUMONT D'URVILLE, Capitaine de vaisseau, publié par ordre du Gouvernement et sous la direction supérieure de M. JACQUINOT, Capitaine de vaisseau, Commandant de la *Zélée*. Atlas Crustacés, pl. 1-9.
- JACQUINOT, H. & H. LUCAS, 1853. — *Ibid.* Crustacés, 3: 1-107.
- JONES, D. S., 1988. — The occurrence of *Ocypode pallidula* Jacquinot (Decapoda, Brachyura) in Australia and the coral sea. *Crustaceana*, 54 (1): 33-38.
- KIM, H. S., 1973. — A catalogue of Anomura and Brachyura from Korea. In: Illustrated Encyclopedia of Fauna & Flora of Korea. Vol. 14, Anomura, Brachyura. Samwha Publishing Compagny: 1-694, fig. 1-265, pl. 1-112 (en coréen with a ?atalogue en anglais: 589-670).
- KINGSLEY, J. S., 1880a. — Carcinological notes, II - Revision of the Gelasimi. *Proceedings of the Academy of Natural Sciences of Philadelphia*: 135-155, pl. 9, fig. 1-15, pl. 10, fig. 16-34.
- 1880b. — Carcinological notes, III - Revision of the genus *Ocypoda*. *Ibid.*: 179-186.
 - 1880c. — Carcinological notes, IV - Synopsis of the Grapsidae. *Ibid.*: 187-224.
- KOMAI, T. & A. ASAKURA, 1995. — *Pagurixus nomurai*, new species, and additional record of *Pagurixus maorus* (Nobili, 1906), hermit crabs from Kume-Jima island, the Ryukyus, Japan (Decapoda, Anomura, Paguridae). *Journal of Crustacean Biology*, 15 (2): 341-354, fig. 1-6.
- KROPP, R. K., 1983. — Three new species of Porcellanidae (Crustacea, Anomura) from the Mariana islands and a discussion of Borradaile's *Petrolisthes lamarckii* complex. *Micronesica*, 19 (1-2): 91-106, fig. 1-3.
- 1986. — A neotype designation for *Petrolisthes tomentosus* (Dana), and description of *Petrolisthes heterochrous*, new species, from the Mariana islands (Anomura, Porcellanidae). *Proceeding of the Biological Society of Washington*, 99 (3): 452-463, fig. 1-3.
 - 1988. — The status of *Cryptochirus coralliodytes* Heller and *Lithoscaptus paradoxus* Milne Edwards (Brachyura, Cryptochiridae). *Ibid.*, 101 (4): 872-882, fig. 1-6.
- KROPP, R. K. & C. BIRKELAND, 1981. — Comparaison of Crustacean associates of *Pocillopora verrucosa* from a high island and an atoll. *Proceedings of the Fourth International Coral Reef Symposium*, Manila, 2: 627-632.

- LABOUTE, P. & B. RICHER DE FORGES, 1986. — Le volcan sous-marin MacDonald (Archipel des îles Austral). Nouvelles observations biologiques et géomorphologiques. *Notes et Documents d'océanographie de l'ORSTOM*, Tahiti, **29**: 1-31, pl. 1-4.
- LEENE, J. E., 1936. — Note on *Charybdis erythrodactyla* (Lam.), *Charybdis acutifrons* (De Man), and *Charybdis obtusifrons* nov. spec. *Zoologische Mededeelingen*, Leiden, **19**: 117-127, fig. 1-12.
- 1938. — The Portunidae of the *Siboga* expedition. VII - Brachygnatha, Portunidae. *Siboga Expedition, Monographie*, **39**, C3, livr. 131: 1-156, fig. 1-87.
- LEMAITRE, R., 1994. — Crustacea Decapoda: Deep-water hermit crabs (Parapaguridae) from French Polynesia with description of four new species. In: A. CROSNIER (ed.), *Résultats des campagnes MUSORSTOM, Volume 12. Mémoires du Muséum National d'Histoire naturelle*, **161**: 375-419, fig. 1-28.
- LEWINSOHN, C., 1969. — Die Anomuren des roten meeres (Crustacea, Decapoda, Paguridae, Galatheidea, Hippidae). *Zoologische Verhandelingen*, Leiden, **104**: 1-213, fig. 1-37, pl. 1-2.
- MAN, J. G. de, 1889. — Ueber einige neue oder seltene indopacifische Brachyuren. *Zoologische Jahrbücher, Abteilung für Systematik*, Jena, **4**: 409-452, pl. 9-10.
- 1890. — Carcinological studies in the Leyden Museum, N°4. *Notes from the Leyden Museum*, **12** (13): 49-126, pl. 3-6.
- 1891. — *Ibid.*, N°5, **13** (1): 1-61, pl. 1-4.
- 1896. — Bericht über die von Herrn Schiffscapitän STORM zu Atjeh, an den westlichen Küsten von Malakka, Borneo und Celebes sowie in der Java-See gesammelten Decapoden und Stomatopoden. *Zoologischen Jahrbüchern, Abtheilung für Systematik*, **9**: 459-514, pl. 33-34.
- MANACH, F. & J. L. CARSIN, 1985. — Pêche profonde sur la pente externe des atolls. *Proceedings of the Fifth International Coral Reef Congress*, Tahiti 27 May - 1 June 1985, **5**: 469-474.
- MANNING, R. B., 1993. — A new deep-sea crab, genus *Chaceon*, from the Austral Islands, southwestern Pacific Ocean (Decapoda, Geryonidae). *Crustacean Research*, **22**: 7-10, fig. 1-2.
- MANNING, R. B. & L. B. HOLTHUIS, 1981. — West african brachyuran crabs (Crustacea, Decapoda). *Smithsonian Contribution to Zoology*, **306**: i-xii, 1-379, fig. 1-88.
- MARQUET, G., 1988. — Les eaux intérieures de la French Polynesia. Principales caractéristiques physiques, chimiques et biologiques. *Thèse de Doctorat de l'Université Paris VI, spécialité Sciences de la Vie (Océanologie Biologique)*: 1-233, fig. 1-64.
- 1991. — Freshwater crustaceans of French Polynesia: Taxonomy, Distribution and Biomass (Decapoda). *Crustaceana*, **61** (2): 125-140, fig. 1.
- 1993. — Etude biogéographique de la faune d'eau douce de French Polynesia. *Biogeographica*, **69** (4): 157-170, fig. 1-3.
- MCLAUGHLIN, P. A. & J. HAIG, 1984. — A review of *Pagurixus* (Decapoda, Anomura, Paguridae) and description of new species. *Crustaceana*, **47** (2): 121-148, fig. 1-7.
- MCLAY, C. L., 1991. — A small collection of deep water sponge crabs (Brachyura, Dromiidae) from French Polynesia, including a new species of *Sphaerodromia* Alcock, 1899. *Bulletin du Muséum national d'Histoire naturelle*, Paris, 4ème série (A), **13** (3-4): 457-481, pl. 1-2.
- 1993. — Crustacea Decapoda: The sponge crabs (Dromiidae) of New Caledonia and the Philippines with a review of the genera. In: A. CROSNIER (éd.), *Résultats des campagnes MUSORSTOM, vol. 10. Mémoires du Muséum national d'Histoire naturelle*, Paris (A), **156**: 111-251, fig. 1-19.
- MERSCHARDT-SALVAT, F., 1991. — L'atoll de Nukutipipi (Tuamotu, French Polynesia) Géomorphologie et peuplements. *Thèse pour l'obtention du diplôme de l'Ecole Pratique des Hautes Etudes (Section des Sciences de la Vie et de la Terre)*, Laboratoire de Biologie Marine et de Malacologie, Paris: 1-153, fig. 1-21.

- MICHEL, A., 1971. — Note sur les Puerulus de Palinuridae et les larves phyllosomes de *Panulirus homarus* (L.). Key de détermination des larves phyllosomes récoltées dans le Pacifique équatorial et sud-tropical (Décapodes). *Cahiers de l'ORSTOM, Océanographie*, Nouméa Nouvelle-Calédonie, **9** (4): 459-473, fig. 1-6.
- MIERS, E. J., 1886. — Report on the Brachyura collected by H.M.S. *Challenger* during the years 1873-1876. *Report on the Scientific Results of the Voyage of H.M.S. Challenger, Zoology*, part 49, **17** (2): i-I, 1-362, pl. 1-29.
- MILNE EDWARDS, A., 1861. — Etudes Zoologiques sur les crustacés récents de la famille des Portuniens. *Archives du Muséum d'Histoire naturelle*, Paris, **10**: 309-428, pl. 28-38.
- 1865a. — Description de quelques crustacés nouveaux appartenant à la tribu des Maiens. *Annales de la Society Entomologique de France*, 4ème série, **5**: 133-147, pl. 3-5.
 - 1865b. — Etudes zoologiques sur les crustacés récents de la famille des Cancériens. *Nouvelles Archives du Muséum*, Paris, 1ère série, **1**: 177-308, pl. 11-19.
 - 1867. — Descriptions de quelques espèces nouvelles de crustacés brachyoures. *Annales de la Society Entomologique de France*, 4ème série, **7**: 263-288.
- MILNE EDWARDS, H., 1848. — Note sur quelques nouvelles espèces du genre Pagure. *Annales des Sciences Naturelles, Zoologie*, Paris, 3ème série, **10**: 59-64.
- 1852. — Observations sur les affinités zoologiques et la classification naturelle des crustacés. *Ibid.*, **18**: 109-166, pl. 3-4.
 - 1853. — Mémoire sur la famille des Ocypodiens. *Ibid.*, **20**: 163-228, pl. 6-11.
- MIYAKE, S., 1991. — Japanese Crustaceans Decapods and Stomatopods in color. Vol. I, Macrura, Anomura and Stomatopoda. Hoikusha Publishing Co., Ltd., Tsurumi-ku, Osaka, 538, Japan, 1-261, pl. 1-56.
- MONOD, T., 1975. — Sur quelques crustacés malacostracés de l'île de la Réunion. *Bulletin du Muséum national d'Histoire naturelle*, Paris, 3ème série, Zoologie, **319** (226): 1005-1033, fig. 1-118.
- 1979. — Crustacés associés à un Anthipathaire des îles Marquises. *Cahiers de l'Indo-Pacifique*, **1** (1): 1-23, fig. 80-85.
- MONTEFORTE, M., 1984. — Contribution à la connaissance de la faune carcinologique de French Polynesia. Inventaire faunistique, répartition bionomique et données quantitatives sur les Crustacés Décapodes Reptantia (Brachyura, Anomura, Macrura) et les Crustacés Stomatopodes habitant les complexes récifo-lagonaires de quelques îles hautes et atolls. *Thèse de l'Ecole Pratique des Hautes Etudes*, 3ème section: 1-196, fig. 1-33.
- 1987. — The decapod reptantia and stomatopod crustaceans of a typical high island coral reef complex in French Polynesia (Tiahura, Moorea island): zonation, community composition and trophic structure. *Atoll Research Bulletin*, **309**: 1-37, fig. 1-10.
- MOOSA, M. K., 1979. — Observations sur la systématique et la zoogéographie des crabes Portunidae indo-ouest-pacifiques. *Diplôme de l'Ecole Pratique des Hautes Etudes*, Paris, 6 juin 1979: 1-130.
- MORGAN, G. J., 1991. — A review of the hermit crab genus *Calcinus* Dana (Crustacea, Decapoda, Diogenidae) from Australia, with description of two new species. *Invertebrate Taxonomy*, **5**: 869-913, fig. 1-63.
- MORRISON, J. P., 1954. — Animal Ecology of the Raroia atoll, Tuamotu. Part 1 - Ecological notes on the mollusks and other animals of Raroia. *Atoll Research Bulletin*, **34**: 1-18.
- MOTTELER, L. S., 1986. — Pacific Island Names. A map and name guide to the new Pacific. *Bishop Museum Miscellaneous Publication*, **34**: 1-91.
- NAGAI, S., 1981. — Notes on *Portunus (Xiphonectes) longispinosus* (Dana) and some related species from Japan. *Nankiseibutu, The Nanki Biological Society*, **23** (1): 27-32, pl. 1-2.

- NAIM, O., 1980a. — Etude qualitative et quantitative de la faune mobile associée aux algues du lagon de Tiahura, île de Moorea, French Polynesia. *Thèse de 3ème cycle*, Université Pierre et Marie Curie, Paris VI: 1-105, fig. 1-40.
- 1980b. — Bilan qualitatif et quantitatif de la petite faune associée aux algues du lagon de Tiahura, île de Moorea, French Polynesia. *Comptes Rendus de l'Académie des Sciences*, Paris, 291, série D: 549-551.
- NAKASONE, Y., 1988. — Land hermit crabs from the Ryukyus, Japan, with a description of a new species from the Philippines (Crustacea, Decapoda, Coenobitidae). *Zoological Science*, 5 (1): 165-178, fig. 1-9.
- NEWMAN, W.A., 1986. — Origin of the Hawaiian marine fauna: Dispersal and vicariance as indicated by barnacles and other organisms. In: GORE, R. H. & K. L. HECK (eds), *Crustacean Biogeography. Crustacean Issues*, 4: 21-49, fig. 1-10.
- NG, P., 1993. — Kraussiinae, a new subfamily for the genera *Kraussia* Dana, 1852, *Palapedia*, new genus, and *Garthasia*, new genus (Crustacea, Decapoda, Brachyura, Xanthidae), with descriptions of two new species from Singapore and the Philippines. *Raffles Bulletin of Zoology*, 41 (1): 133-157, fig. 1-8.
- NOBILI, G., 1906a. — Diagnoses préliminaires de Crustacés Décapodes et Isopodes nouveaux recueillis par M. le Dr G. SEURAT aux îles Touamotou. *Bulletin du Muséum d'Histoire naturelle*, 12 (5): 256-270.
- 1906b. — Mission J. BONNIER et Ch. PÉREZ (Golfe Persique, 1901). Crustacés Décapodes et Stomatopodes. *Bulletin Scientifique de la France et de la Belgique*, 40: 13-159, fig. 1-3, pl. 2-7.
- 1907. — Ricerche sui Crostacei della Polinesia. Decapodi, Stomatopodi, Anisopodi e Isopodi. *Memori della Reale Accademia delle Scienze di Torino*, sér. 2, 57: 351-430, pl. 1-3.
- ODHNER, T., 1925. — Monographierte Gattungen der Krabbenfamilie Xanthidae. I. *Göteborgs Kungl. Vetenskaps- och Vitterhets-Samhälles Handlingar, Fjärde Följden*, 29 (1): 1-92, fig. 1-7, pl. 1-5.
- ODINETZ, O., 1983. — Ecologie et structure des peuplements de crustacés décapodes associés aux coraux du genre *Pocillopora* en French Polynesia et en Micronésie. *Thèse de 3ème cycle*, Université de Paris VI: 1-221, fig. 1-48, photo. 1-16.
- 1984a. — Révision des *Trapezia* du groupe *cymodoce-ferruginea* (Crustacea, Decapoda, Brachyura), avec des notes complémentaires concernant *T. serenei* Odinetz, 1983, et *T. punctimanus* Odinetz, 1983. *Bulletin du Muséum national d'Histoire naturelle*, Paris, 4ème série (A), 2: 431-452, fig. 1-4.
- 1984b. — L'éthologie au service de la systématique: l'exemple des *Trapezia* (Crustacés, Décapodes, Brachyoures). *Océanis*, 10 (1): 123-130, fig. 1-4.
- ODINETZ-COLLART, O. & B. RICHER DE FORGES, 1985. — Ecologie des crustacés décapodes associés aux *Pocillopora* en Polynésie et à Guam (Micronésie). *Proceedings of the Fifth International Coral Reef Congress*, Tahiti 27 May - 1 June 1985, 5: 197-203, fig. 1-4.
- ORTMANN, A., 1891. — Die Decapoden-Krebse des Strassburger Museums. III, Homaridae, Loricata und Thalassinidea. *Zoologischen Jahrbüchern, Abtheilung für Systematik*, Iena, 3: 1-58, pl. 1.
- 1892a. — *Ibid.* IV, Galatheidea und Paguridea, 4: 241-326, pl. 11-12.
- 1892b. — *Ibid.* V, Hippidea, Dromiidea und Oxystomata, 5: 532-588, pl. 26.
- 1893a. — *Ibid.* VI, Majoidea und Cancroidea, Section Portuninae, 7: 23-88, pl. 3.
- 1893b. — *Ibid.* VII, Cyclometopa, 7: 411-495, pl. 17.
- 1894. — *Ibid.* VIII, Catametopa, 8: 683-772, pl. 23.
- 1897a. — Carcinologische Studien. *Ibid.*, 10: 258-372, pl. 17.

- 1897b. — Die geographische Verbreitung der Decapoden-Familie Trapeziidae. *Ibid.*, **10**: 201-216.
- OWEN, R., 1839. — Crustacea. The Zoology of the captain BEECHEY's Voyage: 77-92, pl. 24-28.
- PARDON, D., 1992. — Tahiti entre ciel et mer. Editions du Pacifique, Tahiti & Editions Glénat, Grenoble: 1-143, illustrated in colour.
- PESTA, O., 1913. — Crustacea. I Theil - Decapoda Brachyura aus Samoa. In: K. RECHINGER, Botanische und Zoologische Ergebnisse einer Wissenschaftlichen Forschungreise nach den Samoainseln, dem Neuguinea-Archipel und den Salomonsinseln, März bis Dezember 1905. IV. *Denkschriften der Kaiserlichen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche klasse*, Wien, **88**: 36-65, fig. 1-5, pl. 3.
- PEYROT-CLAUSADE, M., 1977a. — Faune cavitaire mobile des platiers coralliens de la région de Tuléar (Madagascar). *Thèse de Doctorat en Sciences Naturelles*, Université d'Aix-Marseille II: 1-184, fig. 1-26.
- 1977b. — Décapodes brachyures et anomoures (à l'exception des Paguridae) de la cryptofaune de Tiahura, île de Moorea. *Cahiers du Pacifique*, **20**: 211-221, fig. 1-3.
- 1985. — Motile cryptofauna modifications related to coral degradations on Tiahura coral reef flat (Moorea, Polynesia). In: GABRIE, C. & M. HARMELIN VIVIEN (eds). *Proceedings of the Fifth International Coral Reef Congress*, Tahiti 27 May - 1 June 1985, **6**: 459-464, fig. 1-7.
- 1989. — Crab cryptofauna (Brachyura and Anomura) of Tikehau, Tuamotu Archipelago, French Polynesia. *Coral Reefs*, **8**: 109-117, fig. 1-5.
- POUPIN, J., 1994a. — Quelques crustacés décapodes communs de French Polynesia. *Rapport Scientifique du Service Mixte de Surveillance Radiologique et Biologique*: 1-86, fig. 1-68, pl. 1-8.
- 1994b. — The genus *Justitia* Holthuis, 1946, with the description of *J. chani* and *J. vericeli* spp. nov. (Crustacea, Decapoda, Palinuridae). *Journal of Taiwan Museum*, **47** (1): 37-56, fig. 1-4, pl. 1-2.
- 1995. — Etude des *Naxioides* du groupe *robillardii*, Miers, 1882, avec la description de deux nouvelles espèces de French Polynesia. *Journal of Natural History*, **29**: 85-109, fig. 1-10.
- 1996. — Recent contributions to the deep sea decapod Crustacea of French Polynesia. *Proceedings of International Senckenberg Symposium*, Frankfurt a.M., October 18-22, 1993 (in press).
- POUPIN J. & P. MC LAUGHLIN, 1996. — A new *Solitariopagurus* from French Polynesia. *Bulletin du Muséum national d'Histoire naturelle*, (in press).
- POUPIN, J., TAMARI, T. & A. VANDENBOOMGAERDE, 1990. — Pêches profondes aux casiers sur les pentes océaniques des îles de French Polynesia (N/O Marara - 1986/1989). *Notes et Documents d'Océanographie*, centre ORSTOM de Tahiti, **42**: 1-97, fig. 1-21, pl. 1-3.
- RAHAYU, D. L., 1988. — Les Pagures littoraux de Nouvelle-Calédonie et de French Polynesia: Taxonomie, Ecologie et Distribution géographique. *Rapport de D.E.A.*, Université Pierre et Marie Curie, Paris VI: 1-47, fig. 1-15.
- 1992. — Étude des pagures littoraux (Crustacés, Décapodes) d'Indonésie: systématique, écologie, et Biogeography. *Thèse de Doctorat de l'Université de Paris VI, Océanographie Biologique*: 1-232, fig. 1-26.
- RAHAYU D. L. & J. FOREST, 1992. — Le genre *Clibanarius* (Crustacea, Decapoda, Diogenidae) en Indonésie, avec la description de six espèces nouvelles. *Bulletin du Muséum national d'Histoire naturelle*, Paris, 4ème série (A), **14** (3-4): 745-779, fig. 1-7.
- RATHBUN, M. J., 1904. — Les crabes d'eau douce. *Nouvelles Archives du Muséum*, 4ème série, Paris, **6**: 225-312, fig. 1-37, pl. 9-14.

- 1906. — The brachyuran and macrura of the Hawaiian islands. *Bulletin of the United States Fish Commission*, **23** (3): 827-930, fig. 1-79, pl. 1-24.
- 1907. — Reports on the scientific results of the expedition to the Tropical Pacific, in charge of Alexander AGASSIZ, by the U.S. Fish Commission steamer *Albatross*, from August 1899, to March 1900, Commander Jefferson F. MOSER, U.S.N., commanding - IX. *ibid.* from October 1904, to March 1905, lieut.-commander L. M. GARRETT, U.S.N., commanding - X. The Brachyura. *Memoirs of the Museum of Comparative Zoology at Harvard College*, **35** (2): 23-74, pl. 1-9.
- 1918. — The grapsoid crabs of America. *Bulletin of the U.S. National Museum*, **97**: i-xxii, 1-461, fig. 1-172, pl. 1-161.
- RIBES, S., 1989. — Les Raninidae du sud-ouest de l'océan Indien (Crustacea, Decapoda, Brachyura). *Bulletin du Muséum national d'Histoire naturelle*, 4ème série, section A, **11** (4): 905-919, fig. 1-3, pl. 1-2.
- SAINT LAURENT, M. de, 1968. — Révision des genres *Catapaguroides* et *Cestopagurus* et description de quatre genres nouveaux. I. *Catapaguroides* A. Milne Edwards et Bouvier et *Decaphyllus* nov. gen. (Crustacés, Décapodes, Paguridae). *Bulletin du Muséum national d'Histoire naturelle*, Paris, 2ème série, (A), 1967 (1968), **39** (5): 923-954, fig. 1-32.
- 1970. — *Ibid.* V, *Trichopagurus* de Saint Laurent (Crustacés, Décapodes, Paguridae); VI, Conclusion, **42** (1): 210-222, fig. 1-16.
- SAINT LAURENT, M. de & J. POUPIN, 1996. — Crustacea, Anomura: Les espèces indo-ouest pacifiques du genre *Eumunida* Smith, 1880 (Chirostylidae): description de six espèces nouvelles. In: A. CROSNIER (éd.), Résultats des campagnes MUSORSTOM, volume 15. Mémoires du Muséum national d'Histoire naturelle, Paris (A), **168**: 337-385, fig. 1-13.
- SAKAI, T., 1976. — Crabs of Japan and the adjacent seas. Tokyo, Kodansha, Ldt, volume en anglais: i-xxix, 1-773, fig. 1-379; volume en japonais: 1-461; volume de planches, pl. 1-251.
- SAKAI, K. & M. TÜRKAY, 1976. — Bemerkungen zu einigen *Ocypode*-Arten (Crustacea, Decapoda). *Senckenbergiana biologica*, **57** (1/3): 81-96, fig. 1-16.
- 1982. — List der Brachyura des Okinoshima-Gebietes von Kochi, Japan, mit Angabe der vorläufigen Determinationen (det. K. SAKAI & M. TÜRKAY 1982), der Katalognummern und Fundorte. *Bulletin of Shikoku Women's University*, **2** (1): 1-4.
- SALVAT, B., 1986a. — Le littoral corallien. In: C. GLEIZAL (éd.) Encyclopédie de la Polynésie, Tome 3, Le monde marin. C. GLEIZAL/Multipress: 6-24, illustrated in colour.
- 1986b. — Crabes, langoustes, ti'an'e et varo. *Ibid.*: 70-72, illustrated in colour.
- 1986c. — Origine de la vie dans les îles. *Ibid.*, Tome 2, Flore et faune terrestres: 9-22, illustré.
- SALVAT, B. & J. RENAUD-MORNANT, 1969. — Etude écologique du macrobenthos et du meiobenthos d'un fond sableux du lagon de Moruroa (Tuamotu, Polynésie). *Cahiers du Pacifique*, **13**: 159-179, fig. 1-4.
- SALVAT, B. & G. RICHARD, 1985. — Takapoto atoll, Tuamotu archipelago. In: B. DELESALLE, R. GALZIN & B. SALVAT (eds). *Proceedings of the Fifth International Coral Reef Congress*, Tahiti, 27 May - 1 June 1985, **1**: 323-362, fig. 1-34.
- SALVAT, B. & C. RIVES, 1975. — Coquillages de Polynésie. Les éditions du Pacifique, Papeete, Tahiti: 1-391, fig. 1-446.
- SALVAT, F. & B. SALVAT, 1992. — Nukutipipi atoll, Tuamotu archipelago; geomorphology, land and marine flora and fauna and interrelationships. *Atoll Research Bulletin*, **357**: 1-43, fig. 1-9, pl. 1-25.
- SCHMITT, W. L., MCCAIN, J. C. & E. S. DAVIDSON, 1973. — Decapoda I, Brachyura I, Fam. Pinotheridae. In: GRUNER, H. E. & L. B. HOLTHUIS (eds), *Crustaceorum Catalogus*, Pars 3: 1-160.

- SENDLER, A., 1923. — Die Decapoden und Stomatopoden der Hanseatischen Südsee-Expedition. *Abhandlungen Herausgegeben von der Senckenbergischen Naturforschenden Gesellschaft*, **38**: 21-47, fig. 1-3, pl. 5-6.
- SERÈNE, R., 1972. — Observations on the Indo-Pacific species of *Kraussia* Dana 1852 (Decapoda, Brachyura). *Journal of the Royal Society of Western Australia*, **55** (2): 41-61, fig. 1-24.
- 1973. — A new species of Decapoda Hippidae: *Albunea mariellae* nov. sp. from the Banda sea. *Crustaceana*, **24** (3): 261-264, pl. 1-2.
- 1975. — Note additionnelle sur les espèces indo-pacifiques de *Quadrella* Dana, 1851 (Crustacea, Decapoda, Brachyura). *Bulletin de la Society Zoologique de France*, **100** (4): 509-521, fig. 1-13, pl. 1-2.
- 1977. — Crustacés hippidés et brachyoures des îles Seychelles (1ère partie). *Revue de Zoologie Africaine*, **91** (1): 45-68, fig. 1-43.
- 1984. — Crustacés décapodes brachyoures de l'océan Indien occidental et de la mer Rouge. Xanthoidea: Xanthidae et Trapeziidae. *ORSTOM Collection Faune Tropicale*, **24**: 1-349, fig. 1-243, pl. 1-47.
- SERÈNE, R. & N. V. LUOM, 1959. — Note additionnelle sur les espèces de *Chlorodopsis* (Brachyoures). *Annales de la Faculté des Sciences*, Saigon: 301-340, fig. 1-5, pl. 1-3.
- SERÈNE, R. & C. L. SOH, 1970. — New Indo-Pacific genera allied to *Sesarma* Say 1817 (Brachyura, Decapoda, Crustacea). *Treubia*, **27** (4): 387-416, pl. 1-8.
- SERÈNE, R. & A. F. UMALI, 1972. — The family Raninidae and other new and rare species of brachyuran decapods from the Philippines and adjacent regions. *The Philippine Journal of Science*, Manila, **99** (1-2): 21-105, fig. 1-131, pl. 1-9.
- SERÈNE, R., TRAN, V. D. & V. L. NGUYEN, 1958. — Eumedoninae du Viet-Nam (Crustacea). *Treubia*, **24**, part 2: 135-242, fig. 1-14, pl. 4-7.
- SEURAT, L. G., 1904a. — Observations biologiques sur les Cénobites (*Cenobita perlata*, Edwards). *Bulletin du Muséum d'Histoire naturelle*, **5**: 238-242.
- 1904b. — Lettre sur les travaux du laboratoire de Rikitea. *Bulletin du Muséum d'Histoire naturelle*, **10** (3): 94-95.
- 1934. — La faune et le peuplement de la French Polynesia. In: P. LECHEVALIER & fils (éd.), Contribution à l'étude du Peuplement zoologique et botanique des îles du Pacifique. *Society de Biogeography*, **4**: 41-74.
- STEPHENSON, W., 1972. — An annotated check list and key to the Indo-West Pacific swimming crabs (Crustacea, Decapoda, Portunidae). *Bulletin of the Royal Society of New Zealand*, **10**: 1-64.
- 1976. — Notes on Indo-West Pacific Portunids (Decapoda, Portunidae) in the Smithsonian Institution. *Crustaceana*, **31** (1): 11-26, fig. 1.
- STEPHENSON, W. & B. CAMPBELL, 1959. — The australian portunids (Crustacea, Portunidae). III, The genus *Portunus*. *Australian Journal of Marine and Freshwater Research*, **10** (1): 84-124, fig. 1-3, pl. 1-5.
- STEPHENSON, W. & J. HUDSON, 1957. — The Australian Portunids (Crustacea, Portunidae). I, The genus *Thalamita*. *Ibid.*, **8** (3): 312-368, fig. 1-5, pl. 1-10.
- STEPHENSON, W. & M. REES, 1961. — Sur deux nouveaux crustacés Portunidae indo-pacifiques. *Bulletin du Muséum national d'Histoire naturelle*, 2ème série, **33** (4): 421-427, fig. 1-2.
- 1967. — Some portunids crabs from the Pacific and Indian oceans in the collection of the Smithsonian Institution. *Proceedings of the United States National Museum*, Smithsonian Institution, Washington, **120** (3356): 1-114, fig. 1-38, pl. 1-9.

- STIMPSON, W., 1858a. — *Prodromus descriptionis animalium evertebratorum quæ in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladara Ringgold et Johanne Rodgers Ducibus, observavit et descriptis. Pars IV, Crustacea Cancroidea et Corystoidea. Proceedings of the Academy of Natural Sciences, Philadelphia*, **10**: 31-40 [29-37].
- 1858b. — *Ibid. Pars V, Crustacea Ocypodoidea*, **10**: 93-110 [39-56].
- 1858c. — *Ibid. Pars VII, Crustacea Anomoura*, **10**: 225-252 [63-90].
- 1860. — *Ibid. Pars VIII, Crustacea Macrura*, **10**: 91-116, [22-49].
- 1907. — Report on the Crustacea (Brachyura and Anomura) collected by the North Pacific Exploring Expedition, 1853-1856. *Smithsonian Miscellaneous Collections*, **49** (1717): 1-240, pl. 1-26.
- TAKEDA, M., 1980. — *Pilumnus planus* Edmonson and *Leptodius leptodon* Forest & Guinot as synonym of *Forestia depressa* (White) and *Leptodius davaoensis* Ward (Decapoda, Brachyura). *Crustaceana*, **39** (3): 318-320.
- TAKEDA, M. & S. MIYAKE, 1968. — Pilumnid crabs of the family Xanthidae from the West Pacific. I. Twenty-three species of the genus *Pilumnus*, with description of four new species. OHMU Occasional papers of Zoological Laboratory Faculty of Agriculture, Kyushu University, **1** (1): 1-58, fig. 1-13, pl. 1-4.
- 1970. — *Lybia edmondsoni* sp. nov., a new anemone crab from the Hawaiian islands. *Proceeding of the Japanese Society of Systematic Zoology*, **6**: 11-15, fig. 1-6.
- TERAO, A., 1913. — A catalogue of hermit-crabs found in Japan (Paguridae excluding Lithodidae), with descriptions of four new species. *Annotationes Zoologicae Japonenses*, **8** (2): 355-391, fig. 1-4.
- TESCH, J. J., 1917. — Synopsis of the genera *Sesarma*, *Metasesarma*, and *Clistocoeloma* with a key to the determination of the Indo-Pacific species. *Zoologische Mededeelingen*, Leiden, **3** (2-3): 127-260, pl. 15-17.
- 1918. — The Decapoda Brachyura of the *Siboga* expedition. I, Hymenosomatidae, Retroplumidae, Ocypodidae, Grapsidae, and Gecarcinidae. *Siboga Expedition, Monographie*, **39 c**: 1-148, pl. 1-6.
- THOMASSIN, B., 1969. — Identification, variabilité et écologie des Hippidae (Crustacea, Anomura) de la région de Tuléar, S.W. de Madagascar. *Recueil des Travaux de la Station Marine d'Endoume*, fascicule hors série, supplément n°9: 135-177, fig. 1-20.
- 1973. — *Albunea madagascariensis* n. sp., nouvelle espèce d'Hippidae (Decapoda, Anomura) des sables coralliens de la région de Tuléar (S.W. de Madagascar). *Crustaceana*, **24** (3): 265-274, fig. 1-2, pl. 1.
- THOMASSIN, B. A., JOUIN, C., RENAUD-MORNANT, J., RICHARD, G. & B. SALVAT, 1982. — Macrofauna and meiofauna in the coral sediments on the Tiahura reef complex, Moorea island (French Polynesia). *Tethys*, **10** (4): 392-397, fig. 1-5.
- TITGEN, R. H., 1986. — Hawaiian Xanthidae (Decapoda, Brachyura). II, Description of *Garthiella*, new genus, with a redescription of *G. aberrans* (Rathbun, 1906). *Proceedings of the Biological Society of Washington*, **99** (1): 56-60, fig. 1-2.
- TÜRKAY, M., 1971. — Die Portunidae des naturhistorischen Museums Genf, mit einem anhang über die typen von *Ovalipes ocellatus floridanus* Hay & Shore 1918 (Crustacea, Decapoda). *Archives des Sciences de Genève*, **24** (1): 111-143, fig. 1-3, pl. 1-6.
- 1973. — Zur synonymie von *Epigrapsus notatus* und *Cardisoma carnifex* (Crustacea, Decapoda). *Senckenbergiana Biologica*, **54** (1-3): 105-110, fig. 1-8.
- 1974. — Die Gecarcinidae Asiens und Ozeaniens (Crustacea, Decapoda). *Senckenbergiana Biologica*, **55** (4-6): 223-259, fig. 1-19.
- WARD, M., 1942. — Notes on the crustacea of the Desjardins Museum, Mauritius Institute, with descriptions of new genera and species. *The Mauritius Institute Bulletin*, **2** (2): 49-113, pl. 5-6.

WILLIAMS A. B., 1982. — Revision of the genus *Latreillia* Roux (Brachyura, Homoloidea). *Quadreni del Laboratorio di Tecnologia della Pesca*, 3 (2-5): 227-255, fig. 1-8.

A C K N O W L E D G E M E N T

The authorities of the *Service Mixte de Surveillance Radiologique et Biologique*, G. MARTIN and C. PAYEN, have permitted this research through the collaboration between their Institution and the *Laboratoire de Zoologie des Arthropodes* (Muséum national d'Histoire naturelle, Paris), where most of the literature was found. In the *Laboratoire de Zoologie des Arthropodes*, A. CROSNIER, J. FOREST, and D. GUINOT, have always assisted us in our research, and have contributed to improve this work by their corrections and advice. A.J. BRUCE has helped us for the english translation. Finally, the recent collections made in French Polynesia have been greatly facilitated by the whole crew of the *Marara*, and her last two commanding officers, R. AUDIGIER and M. BENARD. To all of them we wish to express our thanks.

A P P E N D I C E S

1 - DEEP SPECIES (from 100m, and deeper)

With the exception of the shrimps (Dendrobranchiata and Caridea), this list resume the work presented at the *International Senckenberg Symposium, Crustacea Decapoda*, Frankfurt, October, 1993 (POUPIN, 1996). Full references on the origin of the material, location and depth of the collections are available in that work. Some species, collected or described since the first compilation are added (in bold). For the new species the reader will find more information in the following works: CROSNIER (1995) for *Pleurocolpus boileaui* gen. & sp. nov., DAVIE (1995) for *Nanocassiope oblonga* nov., FOREST (1995) for the new genera *Ciliopagurus* and *Strigopagurus*, GUINOT & RICHER DE FORGES (1995) for the revision of the homolids and the new genus *Yaldwynopsis*, HOLTHUIS (1993) for *Scyllarus rapanus* nov., LEMAITRE (1994) for the genus *Sympagurus*, MANNING (1993) for *Chaceon australis* nov., POUPIN (1994b, 1995) for the genera *Justitia* and *Naxioides*, POUPIN & MCLAUGHLIN (1996) for *Solitariopagurus* sp. nov., and SAINT LAURENT, de & POUPIN (1996) for *Eumunida treguieri* nov.

Twelve sublittoral to deep species, followed by a "*" are common with the previous list (cf. Conventions).

INFRA-ORDER ASTACIDEA

FAMILY ENOPLOMETOPIDAE

Hoplometopus gracilipes de Saint Laurent, 1988

INFRA-ORDER PALINURIDEA

FAMILY PALINURIDAE

Justitia longimanus (H. Milne Edwards, 1837) **Justitia vericeli* Poupin, 1994*Palinustus unicornutus* Berry, 1979 (Coll. 1995, Fangataufa, 250m, det. POUPIN & CHAN).*Puerulus angulatus* (Bate, 1888)

FAMILY SYNAXIDAE

Palibythus magnificus Davie, 1990 *

FAMILY SCYLLARIDAE

Scyllarus aurora Holthuis, 1981 **Scyllarus rapanus* Holthuis, 1993

INFRA-ORDER ANOMURA

FAMILY DIOGENIDAE

Bathynarius albicinctus (Alcock, 1905)*Bathynarius pacificus* Forest, 1993*Ciliopagurus major* Forest, 1995*Ciliopagurus pacificus* Forest, 1995*Ciliopagurus plessisi* Forest, 1995*Dardanus australis* Forest & Morgan, 1991 * (Coll. 1995, Rapa 70-115m, det. FOREST).*Dardanus brachyops* Forest, 1962 **Strigopagurus pouponi* Forest, 1995

FAMILY PAGURIDAE

Solitariopagurus sp. nov. Poupin & McLaughlin, 1996

FAMILY PARAPAGURIDAE

Strobopagurus cf. *gracilipes* (A. Milne Edwards, 1891) = *S. cf. sibogae* in POUPIN 1996 *fide* LEMAITRE (1994: 378)*Sympagurus affinis* (Henderson, 1888)*Sympagurus boletifer* (de Saint Laurent, 1972)*Sympagurus bougainvillei* Lemaitre, 1994*Sympagurus dofleini* (Balss, 1912)*Sympagurus planimanus* (de Saint Laurent, 1972)*Sympagurus pouponi* Lemaitre, 1994*Sympagurus trispinosus* (Balss, 1911)*Sympagurus tuamotu* Lemaitre, 1994*Sympagurus wallisi* Lemaitre, 1994

FAMILY GALATHEIDAE

Leiogalathea laevirostris (Balss, 1913)*Munida amathea* Macpherson & de Saint Laurent, 1991*Munida ducoussoi* Macpherson & de Saint Laurent, 1991*Munida evarne* Macpherson & de Saint Laurent, 1991*Munida hystrix* Macpherson & de Saint Laurent, 1991*Munida lenticularis* Macpherson & de Saint Laurent, 1991*Munida longicheles* Macpherson & de Saint Laurent, 1991*Munida normani* Henderson, 1885*Munida ocellata* Macpherson & de Saint Laurent, 1991

- Munida pasithea* Macpherson & de Saint Laurent, 1991
Munida plexaura Macpherson & de Saint Laurent, 1991
Munida polynoe Macpherson & de Saint Laurent, 1991
Munida profunda Macpherson & de Saint Laurent, 1991
Munida pulchra Macpherson & de Saint Laurent, 1991
Munida rubella Macpherson & de Saint Laurent, 1991
Munida rubrovata Macpherson & de Saint Laurent, 1991
Munida sp. cf. *pilosimanus* Baba, 1969
Munida sp. cf. *solae* Baba, 1986
Sadayoshia aff. *edwardsii* Miers, 1884

FAMILY CHIROSTYLIIDAE

- Eumunida keijii* de Saint Laurent & Macpherson, 1990
Eumunida treguieri de Saint Laurent & Poupin, 1996

FAMILY LITHODIDAE

- Lithodes megacantha* Macpherson, 1991

INFRA-ORDER BRACHYURA

FAMILY DROMIIDAE

- Dromia wilsoni* (Fulton & Grant, 1902) *
Sphaerodromia ducoussoi McLay, 1991

FAMILY DYNOMENIDAE

- Dynomene tanensis* Yokoya, 1933 (Coll. 1995, Fangataufa, 310m, det. MCLAY).

FAMILY HOMOLIDAE

- Homola ikedai* Sakai, 1979
Homola orientalis s.l. Henderson, 1888
Homologenus broussei Guinot & Richer de Forges, 1981
Hypsophrys inflata Guinot & Richer de Forges, 1981
Hypsophrys aff. *murotoensis* Sakai, 1979
Hypsophrys personata Guinot & Richer de Forges, 1981
Moloha aff. *majora* Kubo, 1936
Yaldwynopsis aff. *spinimanus* Griffin, 1965

FAMILY LATREILLIIDAE

- Latreillia metanesa* Williams, 1982.

FAMILY RANINIDAE

- Notopoides latus* Henderson, 1888
Notosceles chimonis Bourne, 1922 *
Notosceles viaderi Ward, 1942

FAMILY POUPINIIDAE

- Poupinia hirsuta* Guinot, 1991

FAMILY CALAPPIDAE

- Calappa* aff. *hepatica* (Linné, 1758) = *Calappa* sp. nov. (GALL com. pers.)
Mursia hawaiensis Rathbun, 1893

FAMILY LEUCOSIIDAE

- Oreotlos encymus* Tan & Ng, 1993

Oreotlos potanus Tan & Ng, 1993
Randallia serenei Richer de Forges, 1983

FAMILY MAJIDAE

Cyrtomaia ihlei Guinot & Richer de Forges, 1982
Naxioides teatui Poupin, 1995
Naxioides vaitahu Poupin, 1995

FAMILY PARTHENOPIDAE

Parthenope (Platylambrus) poupini Garth, 1993
Parthenope (Platylambrus) stellata Rathbun, 1906

FAMILY CANCRIDAE

Platepistoma balssii (Zarenkov, 1990)

FAMILY GERYONIDAE

Chaceon australis Manning, 1993
Chaceon poupini Manning, 1992

FAMILY GONEPLACIDAE

Carcinoplax aff. cooki Rathbun, 1906
Carcinoplax aff. crosnieri Guinot & Richer de Forges, 1981
Carcinoplax aff. verdensis Rathbun, 1914

FAMILY PORTUNIDAE

Charybdis paucidentata A. Milne Edwards, 1861 *
Parathranites hexagonum Rathbun, 1906 (Coll. Eiao, Vanavana; 155-240m, det. Moosa)
Portunus nipponensis Sakai, 1938 *
Thalamita macrospinifera Rathbun, 1911 *

FAMILY XANTHIDAE

Alainodaeus akiaki Davie, 1993
Alainodaeus rimatara Davie, 1993 *
Banareia fatuhiva Davie, 1993
Demania garthi Guinot & Richer de Forges, 1981
Demania mortensenii (Odhner, 1925)
Epistocavea mururoa Davie, 1993
Euryozius danielae Davie, 1993
Hypocolpus mararae Crosnier, 1991
Lophozozymus bertoniciae Guinot & Richer de Forges, 1981
Medaeus grandis Davie, 1993 *
Meractaea tafai Davie, 1993
Meriola rufomaculata Davie, 1993
Nanocassiope oblonga Davie, 1995
Paraxanthodes polynesiensis Davie, 1993
Rata tuamotense Davie, 1993
Pleurocolpus boileau Crosnier, 1995

XANTHOIDEA INCERTAE SEDIS

Beuroisia manquenei Guinot & Richer de Forges, 1981
Mathildella maxima Guinot & Richer de Forges, 1981
Progeryon mararae Guinot & Richer de Forges, 1981

2 - PARTIAL IDENTIFICATIONS

These incomplete references were generally found in ecological works. To avoid partial identifications in the main list, they are presented separately hereafter, by alphabetical order. Four genera are cited for the first time in the area: *Glabropilumnus*, *Heteropanope*, *Libinia*, and *Tylodiplax*.

- Actaea* sp. : MONTEFORTE, 1984: 170, annex 1, tab. a (Moorea, Tahiti); 1987: 8 (Moorea).
- Actumnus* sp. : MONTEFORTE, 1984: 170, annex 1, tab. a (Makatea). — GUINOT, 1985: 452 (List).
- Dromia* sp. : ODINETZ, 1983: 208, with a ? (Tahiti). — GUINOT, 1985: 448 (List).
- Euxanthus* sp. : PEYROT-CLAUSADE, 1977a, annex of the species: 26 (Moorea), juvenile; 1977b: 212 (Moorea). — GUINOT, 1985: 450 (List).
- Glabropilumnus* sp. : MONTEFORTE, 1984: 170, annex 1, tab. a (Mataiva). — DELESALLE, 1985: 289 (Mataiva). — GUINOT, 1985: 452 (List).
- Heteropanope* sp. : PEYROT-CLAUSADE, 1989: 113 (Tikehau).
- Libinia* sp. : SENDLER, 1923: 40 (Tahiti).
- Lybia* sp. : PEYROT-CLAUSADE, 1977a, annex of the species: 27, juvenile (Moorea).
- Neoliomera* sp. : PEYROT-CLAUSADE, 1977a, annex of the species: 27, juvenile (Moorea).
- Pachygrapsus* sp. : HOLTHUIS, 1953: 32 (Raroia).
- Paramedaeus* sp. : PEYROT-CLAUSADE, 1989: 113 (Tikehau).
- Parthenope* sp. : MONTEFORTE, 1984: 174, annex 1, tab. a (Takapoto). — SALVAT & RICHARD, 1985: 350 (Takapoto).
- Petrolishtes* spp. : MONTEFORTE, 1984: 173. — PEYROT-CLAUSADE: 1977: 25.
- Phymodius* sp. : PEYROT-CLAUSADE, 1977a, annex of the species: 27, juvenile (Moorea).
- Pilodius* sp. : PEYROT-CLAUSADE, 1977a, annex of the species: 27, juvenile (Moorea).
- Pilumnus* spp. : MONTEFORTE, 1984: 171, annex 1, tab. a, photo p. 131a (Takapoto); 1987: 9 (Moorea). — SALVAT & RICHARD, 1985: 350 (Takapoto).
- Tylodiplax* sp. : THOMASSIN *et al.*, 1982: 393 (Moorea). — GUINOT, 1985: 453 (List).
- Xanthias* sp. : PEYROT-CLAUSADE, 1989: 112, 115 (Moorea, Tikehau).

INDEX

A

<i>abbreviatum, Percnon</i>	71
<i>abbreviatus, Acanthopus</i>	71
<i>aberrans, Garthiella, Pilodius</i>	54
<i>Acanthophrys cristimanus</i>	26; 78
<i>Acanthopus</i>	
<i>abbreviatus</i>	71
<i>planissimus</i>	71
<i>tenuifrons</i>	71
<i>Achelous, Neptunus, Portunus</i>	31
<i>Actaea</i>	
<i>affinis</i>	55
<i>calculosa</i>	42; 77
<i>cavipes</i>	44
<i>consobrina</i>	42; 45
<i>danae</i>	42
<i>depressa</i>	43
<i>garretti</i>	43
<i>glandifera</i>	42; 77
<i>hirsutissima</i>	42
<i>lata</i>	45
<i>parvula</i>	38
<i>polyacantha</i>	42
<i>rufopunctata</i>	43; 44
<i>rupPELLIOIDES</i>	42; 45
<i>scabra</i>	43
<i>sp.</i>	99
<i>superciliaris</i>	43
<i>tomentosa</i>	43
<i>tumulosa</i>	44
<i>Actaeodes</i>	
<i>affinis</i>	55
<i>areolatus</i>	42
<i>consobrinus</i>	42
<i>hirsutissimus</i>	42
<i>richtersii</i>	41
<i>tomentosus</i>	43
<i>Actaeomorpha</i>	
<i>alvae</i>	5; 28; 77; 78
<i>erosa</i>	28
<i>punctata</i>	28
<i>Actumnus</i>	
<i>asper</i>	61
<i>bonnieri</i>	61
<i>digitalis</i>	61
<i>globulus</i>	61; 77
<i>integerrimus</i>	54
<i>obesus</i>	61
<i>setifer</i>	61
<i>sp.</i>	99
<i>tomentosus</i>	61
<i>aculeata, Galathea</i>	19
<i>adactyla, Hippa</i>	23
<i>admete, Thalamita</i>	33
<i>aeneus, Zozimus</i>	47
<i>aequabilis, Clibanarius</i>	16
<i>Aethra scruposa</i>	28; 77

<i>affine, Percnon</i>	71
<i>affinis</i>	
<i>Actaea</i>	55
<i>Actaeodes</i>	55
<i>Galathea</i>	19
<i>Percnon</i>	71
<i>Sympagurus</i>	96
<i>akiaki, Alainodaeus</i>	98
<i>Alainodaeus</i>	
<i>akiaki</i>	98
<i>rimatara</i>	7; 41; 78; 98
<i>albicinctus, Bathynarius</i>	96
<i>albolineatus, Grapsus</i>	67
<i>Albunea speciosa</i>	22; 76; 78
<i>albus, Pseudograpsus</i>	69
<i>alcocki</i>	
<i>Lioxanthodes</i>	49
<i>Thalamita</i>	36
<i>alexandri</i>	
<i>Callinectes</i>	31
<i>Portunus</i>	31; 77; 78
<i>algae, Galathea</i>	20
<i>alvae, Actaeomorpha</i>	5; 28; 77; 78
<i>amamiensis, Galathea</i>	19; 77
<i>amathea, Munida</i>	96
<i>Amphiuca, Uca</i>	74
<i>anaglypta, Platypodia</i>	46
<i>anaglyptus, Etisus</i>	6; 52; 53
<i>anceps, Eupagurus, Pagurixus</i>	19
<i>andreossyi, Cymo</i>	38
<i>angulatus, Puerulus</i>	96
<i>angustifrons</i>	
<i>Micippoides</i>	27
<i>Sesarma</i>	71; 78
<i>Aniculus</i>	
<i>aniculus</i>	13
<i>maximus</i>	14
<i>sp.</i>	14
<i>typicus</i>	13
<i>aniculus, Aniculus, Pagurus</i>	13
<i>annulata</i>	
<i>Charybdis</i>	30; 77
<i>Goniosoma</i>	30
<i>annulipes</i>	
<i>Lydia</i>	64; 78
<i>Ruppellia</i>	64
<i>antarcticus</i>	
<i>Parribacus</i>	10; 11
<i>Scyllarus</i>	10
<i>antipodarum, Arctides</i>	10
<i>Arctides</i>	
<i>antipodarum</i>	10
<i>regalis</i>	10; 76; 77
<i>arcuatus, Xantho</i>	48
<i>areolata</i>	
<i>Chorodopsis</i>	55
<i>Trapezia</i>	3; 58; 60

<i>areolatus</i>	
<i>Actaeodes</i>	42
<i>Pilodius</i>	55
<i>armiger, Liocarpilodes</i>	54
<i>Ashtoret</i>	
<i>granulosa</i>	25; 76
<i>lunaris</i>	25
<i>picta</i>	25
<i>asper, Actumnus</i>	61
<i>aspera, Schizophrys</i>	27; 77
<i>Atergatis</i>	
<i>elegans</i>	46
<i>floridus</i>	45
<i>limbatus</i>	47
<i>ocyroe</i>	45
<i>Atergatopsis</i>	
<i>germaini</i>	45
<i>signatus</i>	45
<i>Aulacolambrus hoplonotus</i>	29
<i>aurora, Scyllarus</i>	7; 11; 96
<i>australis</i>	
<i>Chaceon</i>	95; 98
<i>Dardanus</i>	17; 76; 96

B	
<i>balssii, Platepistoma</i>	98
<i>Banareaia</i>	
<i>fatuhiva</i>	98
<i>parvula</i>	38
<i>banski, Matuta</i>	25
<i>barbata, Chlorodiella</i>	53
<i>barffi, Cancer</i>	11
<i>Bathynarius</i>	
<i>albicinctus</i>	96
<i>pacificus</i>	96
<i>bella</i>	
<i>Liomera</i>	38
<i>Trapezia</i>	3; 58; 60
<i>bellus, Actaeodes, Carpilodes</i>	38
<i>bertonciniae, Lophozozymus</i>	98
<i>Beuroisia manquenei</i>	98
<i>bidentatus, Lachnopus</i>	48
<i>bifrontalis, Etisus</i>	51; 52
<i>Birgus latro</i>	11; 80; 81
<i>bispinosus, Petrolisthes</i>	21
<i>boileaui, Pleurocolpus</i>	95; 98
<i>boletifer, Sympagurus</i>	96
<i>bonnieri, Actumnus</i>	61
<i>borradaillei</i>	
<i>Hyastenus</i>	27
<i>Petrolisthes</i>	21; 22
<i>bosci, Petrolisthes</i>	21
<i>bougainvillei, Sympagurus</i>	96
<i>bouvieri, Thalamita</i>	33
<i>brachyops, Dardanus</i>	7; 17; 96
<i>brevimanus, Coenobita</i>	12
<i>broussei, Homologenus</i>	97

C

<i>caestifera, Lybia</i>	37
<i>Calappa</i>	
<i>calappa</i>	6; 25
<i>hepatica</i>	25; 97
<i>sp. nov.</i>	97
<i>tuberculata</i>	25
<i>Calcinus</i>	
<i>elegans</i>	14
<i>gaimardi</i>	14
<i>guamensis</i>	14; 76
<i>herbstii</i>	14; 15
<i>imperialis</i>	14; 76
<i>laevimanus</i>	14; 81
<i>latens</i>	15
<i>minutus</i>	6; 15; 77
<i>nitidus</i>	4; 15; 78
<i>seurati</i>	4; 15
<i>spicatus</i>	4; 15
<i>terrae-reginae</i>	15
<i>tibicen</i>	14
<i>calculosa, Actaea</i>	42; 77
<i>Callinectes alexandri</i>	31
<i>Camposcia retusa</i>	26
<i>canaliculata, Cryptodromia</i>	23
<i>canaliculatus, Xanthias</i>	50
<i>Cancer</i>	
<i>barffi</i>	11
<i>theresae</i>	10
<i>Caphyra</i>	
<i>rotundifrons</i>	29
<i>tridens</i>	30
<i>Carcinoplax</i>	
<i>cooki</i>	98
<i>crosnieri</i>	98
<i>verdensis</i>	98
<i>Cardisoma</i>	
<i>carnifex</i>	65; 81
<i>hirtipes</i>	65
<i>obesum</i>	65
<i>rotundum</i>	66
<i>carinipes, Zozymodes</i>	47
<i>carnescens, Coenobita</i>	12; 77; 78
<i>carnifex, Cardisoma</i>	65; 81
<i>Carpilius</i>	
<i>convexus</i>	7; 62
<i>maculatus</i>	62
<i>Carpilodes</i>	
<i>bellus</i>	38
<i>cinctimanus</i>	39
<i>granulatus</i>	40
<i>monticulosus</i>	39; 40
<i>pallidus</i>	39
<i>rugatus</i>	38; 40
<i>tristis</i>	40
<i>vaillantianus</i>	38
<i>venosus</i>	40
<i>Carpiloxanthus rugipes</i>	38

<i>Carupa</i>		
<i>laeviuscula</i>	29
<i>tenuipes</i>	29
<i>Catapaguroides</i>		
<i>fragilis</i>	5; 18
<i>trichophthalmus</i>	19
<i>Catoptrus nitidus</i>	29
<i>cavatus, Cycloanthops, Neoxanthops</i>	50
<i>cavimana, Tetralia</i>	57
<i>cavipes</i>		
<i>Actaea</i>	44
<i>Coenobita</i>	5; 12; 77
<i>Psaumis</i>	44
<i>caystrus, Pseudozius</i>	65
<i>cellulosa, Psaumis</i>	44
<i>Cenobita</i>		
<i>carnescens</i>	12
<i>clypeata</i>	12
<i>olivieri</i>	13
<i>rugosa</i>	13
<i>cerathophthalma, Ocypode</i>	73
<i>Chaceon</i>		
<i>australis</i>	95; 98
<i>poupini</i>	98
<i>chaptali, Thalamita</i>	33
<i>Charybdis</i>		
<i>annulata</i>	30; 77
<i>erythrodactyla</i>	30
<i>hawaiensis</i>	30; 78
<i>orientalis</i>	31
<i>paucidentata</i>	7; 31; 98
<i>Chasmagnathus subquadratus</i>	69; 77
<i>chimmonis, Notosceles</i>	24; 97
<i>Chlorodiella</i>		
<i>barbata</i>	53
<i>cytherea</i>	53
<i>laevissima</i>	53
<i>laevissimus</i>	54
<i>nigra</i>	54
<i>Chlorodius</i>		
<i>cytherea</i>	53
<i>dehaani</i>	55
<i>laevissimus</i>	53
<i>monticulus</i>	55
<i>niger</i>	54
<i>sanguineus</i>	49
<i>ungulatus</i>	55
<i>Chlorodopsis</i>		
<i>areolata</i>	55
<i>granulatus</i>	56
<i>pugil</i>	56
<i>scabricula</i>	55
<i>spinipes</i>	56
<i>venusta</i>	57
<i>chlorophthalmus</i>		
<i>Amphiaca</i>	74
<i>Gelasimus</i>	74
<i>Uca</i>	74; 80
<i>Ciliopagurus</i>		
<i>krempfi</i>	16
<i>major</i>	96
<i>pacificus</i>	96
<i>plessisi</i>	96
<i>strigatus</i>	16
<i>cinctimana, Liomera</i>	39
<i>cinctimanus, Carpilodes</i>	39
<i>cinctipes, Tetralia</i>	57
<i>Clibanarius</i>		
<i>aequabilis</i>	16
<i>corallinus</i>	16
<i>eurysternus</i>	16
<i>humilis</i>	16
<i>ransonni</i>	5; 16
<i>rhabdodactylus</i>	16
<i>semistriatus</i>	16
<i>striolatus</i>	17
<i>zebra</i>	16; 17
<i>clypeata, Cenobita</i>	12
<i>clypeatus, Cenobita, Coenobita</i>	12
<i>coccinea, Porcellana</i>	21
<i>coccineus, Petrolisthes</i>	21
<i>Coenobita</i>		
<i>brevimanus</i>	12
<i>carnescens</i>	12; 77; 78
<i>cavipes</i>	5; 12; 77
<i>clypeatus</i>	12
<i>hilgendorfi</i>	12
<i>olivieri</i>	12; 13; 77
<i>perlata</i>	13
<i>perlatus</i>	12; 13; 77; 81
<i>rugosus</i>	13
<i>rugosus var. granulatus</i>	13
<i>spinosus</i>	12; 13; 77
<i>coerulipes, Thalamita</i>	33
<i>consobrina</i>		
<i>Actaea</i>	42; 45
<i>Actaeodes</i>	42
<i>consobrinus</i>		
<i>Actaeodes</i>	42
<i>Macrophthalmus</i>	72; 78; 80
<i>contrarius, Parthenope</i>	28
<i>convexus</i>		
<i>Carpilius</i>	7; 62
<i>Macrophthalmus</i>	73; 80
<i>convictor, Eumedon, Eumedonus</i>	29
<i>cooki, Carcinoplax</i>	98
<i>cooperi, Thalamita</i>	33
<i>corallinus, Clibanarius</i>	16
<i>coralliodytes, Cryptochirus</i>	74; 77
<i>Coralligalathea humilis</i>	19
<i>cordimana, Ocypode</i>	73
<i>coronata, Cryptodromia</i>	23; 77
<i>corrugata, Thalamita</i>	5; 34
<i>crassimanus</i>		
<i>Leptodius</i>	49
<i>Macromedaeus</i>	49
<i>Ptychognathus</i>	69; 77; 78
<i>Xantho</i>	49
<i>crassipes, Uca</i>	74
<i>crassum, Sarmatium</i>	70
<i>crenata, Thalamita</i>	34
<i>crenulatus, Hemigrapsus, Heterograpsus</i>	69
<i>crinipes, Geograpsus</i>	66
<i>cristatus</i>		
<i>Leptodius</i>	48
<i>Lophozozymus</i>	46
<i>Xanthodius</i>	47

<i>cristimanus</i>	
<i>Acanthophrys</i>	26; 78
<i>Hyastenus</i>	26
<i>crozieri, Carcinoplax</i>	98
<i>Cryptochirus coralliodytes</i>	74; 77
<i>Cryptodromia</i>	
<i>canaliculata</i>	23
<i>coronata</i>	23; 77
<i>fallax</i>	23
<i>Cryptodromiopsis tridens</i>	24
<i>cupulifer, Polydectus</i>	37
<i>cyaneus, Planes</i>	68; 76
<i>Cyclax suborbicularis</i>	26
<i>Cycloachelous, Portunus</i>	31
<i>Cyclodius</i>	
<i>gracilis</i>	55
<i>ornatus</i>	55
<i>Cyclograpus</i>	
<i>integer</i>	70
<i>longipes</i>	70
<i>parvulus</i>	70
<i>Cymo</i>	
<i>andreossyi</i>	38
<i>deplanatus</i>	38
<i>melanodactyla</i>	38
<i>melanodactylus</i>	38
<i>quadrilobatus</i>	38
<i>cymodoce, Trapezia</i>	58; 59; 60; 61
<i>cyrenae, Quadrella</i>	57
<i>Cyrtomaia ihlei</i>	98
<i>cytherea, Chlorodiella, Chlorodius</i>	53

D

<i>Dacryopilumnus eremita</i>	63
<i>Daira perlata</i>	75
<i>dakini, Thalamita</i>	34
<i>Daldorfia horrida</i>	28
<i>danae</i>	
<i>Actaea</i>	42
<i>Thalamita</i>	5; 34; 76
<i>danielae, Euryozius</i>	98
<i>Dardanus</i>	
<i>australis</i>	17; 76; 96
<i>brachyops</i>	7; 17; 96
<i>deformis</i>	17
<i>gemmatus</i>	17
<i>guttatus</i>	17; 77
<i>haani</i>	18
<i>lagopodes</i>	17
<i>megistos</i>	18
<i>pedunculatus</i>	18
<i>sanguinolentus</i>	18
<i>scutellatus</i>	18
<i>davaoensis</i>	
<i>Leptodius</i>	48
<i>Trapezia</i>	60
<i>decacanthus, Petrolisthes</i>	21
<i>deformis, Dardanus, Pagurus</i>	17
<i>dehaani, Chlorodius</i>	55
<i>Deltuca, Uca</i>	74

<i>demani</i>	
<i>Etisus</i>	52
<i>Neoliomera</i>	40; 41
<i>Thalamita</i>	34
<i>Demania</i>	
<i>garthi</i>	98
<i>mortenseni</i>	98
<i>dentata, Trapezia</i>	58
<i>dentatus, Etisus</i>	52
<i>deplanatus, Cymo</i>	38
<i>depressa</i>	
<i>Actaea</i>	43
<i>Forestia</i>	43; 77
<i>Plagusia</i>	72
<i>depressus, Grapsus</i>	67; 77; 78
<i>diformis, Pagurus</i>	17
<i>digitalis</i>	
<i>Actumnus</i>	61
<i>Platypodia</i>	61
<i>Trapezia</i>	59
<i>digitalis, Trapezia</i>	58; 60
<i>Diogenes gairdineri</i>	18
<i>Discoplax longipes</i>	66
<i>distinguendus</i>	
<i>Macromedaeus</i>	49; 77
<i>Xantho</i>	49
<i>dodone, Lophozozymus</i>	46
<i>dofleini, Sympagurus</i>	96
<i>Domecia</i>	
<i>glabra</i>	63
<i>hispida</i>	63
<i>Dotilla fenestra</i>	76
<i>Dromia</i>	
<i>sp.</i>	99
<i>wilsoni</i>	7; 24; 97
<i>dubius, Portunus</i>	31
<i>ducousoi</i>	
<i>Munida</i>	96
<i>Sphaerodromia</i>	97
<i>dumerilii, Tylocarcinus</i>	28
<i>duperreyi, Gelasimus</i>	3; 74
<i>dussumieri, Gelasimus, Uca</i>	74; 76
<i>Dynomene</i>	
<i>hispida</i>	24
<i>praedator</i>	24
<i>sinense</i>	24; 76
<i>sp.</i>	24
<i>spinosa</i>	24
<i>tanensis</i>	97
E	
<i>easteranus, Ptychognathus</i>	6; 69
<i>Ebalia erosa</i>	26
<i>Ebaliospis erosa</i>	26
<i>Echinoecus pentagonus</i>	29
<i>edwardsi</i>	
<i>Lophozozymus</i>	46
<i>Sadayoshia</i>	97
<i>Thalamita</i>	34
<i>Elamena, mathaei</i>	75

<i>eldredgei</i> , <i>Petrolisthes</i>	21
<i>electra</i>	
<i>Etisodes</i>	52
<i>Etisus</i>	52
<i>elegans</i>	
<i>Atergatis</i>	46
<i>Calcinus</i>	14
<i>Lissocarcinus</i>	5; 30; 78
<i>Petrolisthes</i>	21
<i>encymus</i> , <i>Oreotlos</i>	97
<i>Enoplometopus</i>	
<i>holthuisi</i>	8; 77
<i>sp. nov.</i>	8
<i>Epigrapsus politus</i>	4; 66; 70
<i>Epistocavea mururoa</i>	98
<i>eremita</i> , <i>Dacryopilumnus</i>	63
<i>Eriphia</i>	
<i>laevimana</i>	63
<i>scabricula</i>	63
<i>sebana</i>	63
<i>erosa</i>	
<i>Actaeomorpha</i>	28
<i>Ebalia</i>	26
<i>Ebaliopsis</i>	26
<i>erythrodactyla</i> , <i>Charybdis</i>	30
<i>erythrodactylum</i> , <i>Goniosoma</i>	30
<i>Etisodes</i>	
<i>electra</i>	52
<i>frontalis</i>	52
<i>splendidus</i>	53
<i>Etisus</i>	
<i>anaglyptus</i>	6; 52; 53
<i>bifrontalis</i>	51; 52
<i>demanii</i>	52
<i>dentatus</i>	52
<i>electra</i>	52
<i>frontalis</i>	52
<i>laevimanus</i>	52; 53
<i>macrodactylus</i>	52; 53
<i>punctatus</i>	53; 77; 78
<i>rugosus</i>	52
<i>splendidus</i>	53
<i>Eucrate</i>	75
<i>Eudora tetraodon</i>	51
<i>Eumedon convictor</i>	29
<i>Eumedonus</i>	
<i>convictor</i>	29
<i>pentagonus</i>	29
<i>Eumunida</i>	
<i>keijii</i>	97
<i>treguieri</i>	95; 97
<i>euopsis</i> , <i>Pagurus</i>	18
<i>Eupagurus</i>	
<i>anceps</i>	19
<i>laevimanus</i>	19
<i>maorus</i>	19
<i>Euryozius danielae</i>	98
<i>eurysternus</i> , <i>Clibanarius</i>	16
<i>Euxanthus</i>	
<i>exsculptus</i>	41
<i>rugosus</i>	41
<i>sculptilis</i>	41
<i>sp.</i>	99
<i>evarne</i> , <i>Munida</i>	96

<i>exaratus</i>	
<i>Leptodius</i>	48
<i>Xantho</i>	49
<i>excelsus</i> , <i>Perigrapsus</i>	65
<i>excentrica</i> , <i>Paractaea</i>	43
<i>exsculptus</i> , <i>Euxanthus</i>	41

F

<i>fakaravensis</i> , <i>Pachygrapsus</i>	4; 68
<i>fallax</i> , <i>Cryptodromia</i>	23
<i>fasciatus</i> , <i>Panulirus</i>	10
<i>fatuhiva</i> , <i>Banareia</i>	98
<i>femoristriga</i> , <i>Panulirus</i> , <i>Senex</i>	9
<i>fenestra</i> , <i>Dotilla</i>	76
<i>ferruginea</i> , <i>Trapezia</i>	58; 59; 60; 61
<i>flavopunctata</i> , <i>Trapezia</i>	59
<i>flavus</i> , <i>Pilodius</i>	56
<i>floridus</i> , <i>Atergatis</i>	45
<i>Forestia</i>	
<i>depressa</i>	43; 77
<i>scabra</i>	43; 77
<i>formosa</i> , <i>Trapezia</i>	59; 77
<i>fragilis</i> , <i>Catapaguroides</i>	5; 18
<i>frontalis</i>	
<i>Etisodes</i>	52
<i>Etisus</i>	52
<i>fusca</i> , <i>Trapezia</i>	59

G

<i>Gaillardiellus</i>	
<i>rueppelli</i>	43; 77
<i>superciliaris</i>	43
<i>gaimardi</i>	
<i>Calcinus</i>	14
<i>Gelasimus</i>	74
<i>Uca</i>	74
<i>gairdineri</i> , <i>Diogenes</i>	18
<i>Galathea</i>	
<i>aculeata</i>	19
<i>affinis</i>	19
<i>algae</i>	20
<i>amamiensis</i>	19; 77
<i>latirostris</i>	20; 77
<i>megalochira</i>	19
<i>serrirostris</i>	20
<i>gardineri</i> , <i>Thalamita</i>	36
<i>garretti</i> , <i>Actaea</i>	43
<i>garthi</i> , <i>Demania</i>	98
<i>Garthiella aberrans</i>	54
<i>gatavakensis</i> , <i>Thalamita</i>	4; 34
<i>Gelasimus</i>	
<i>chlorophthalmus</i>	74
<i>duperreyi</i>	3; 74
<i>dussumieri</i>	74
<i>gaimardi</i>	74
<i>latreillei</i>	74
<i>pulchellus</i>	74
<i>tetragonon</i>	74
<i>gelida</i> , <i>Nucia</i>	26

<i>gemmatus, Dardanus, Pagurus</i>	17
<i>Geograpsus</i>	
<i>crinipes</i>	66
<i>grayi</i>	66
<i>lividus</i>	66
<i>stormi</i>	66
<i>germaini, Atergatopsis</i>	45
<i>glaber, Lophozozymus</i>	46
<i>glaberrima, Tetralia</i>	57
<i>glabra, Domecia</i>	63
<i>Glabropilumnus sp.</i>	99
<i>glandifera, Actaea</i>	42; 77
<i>Globopilumnus globosus</i>	3; 64
<i>globosus</i>	
<i>Globopilumnus</i>	3; 64
<i>Pilumnus</i>	56; 64
<i>globulus, Actumnus</i>	61; 77
<i>gloriensis, Thalamita</i>	34
<i>Gonioinfradens, Charybdis</i>	31
<i>Goniosoma</i>	
<i>annulatum</i>	30
<i>erythrodactylum</i>	30
<i>lineatum</i>	36
<i>Goniosupradens, Charybdis</i>	30
<i>gracilipes</i>	
<i>Grapsus</i>	67
<i>Thalamita</i>	35
<i>gracilipes, Strobopagurus</i>	96
<i>gracilis</i>	
<i>Cyclodius</i>	55
<i>Leptodius</i>	49
<i>Tylocarcinus</i>	28
<i>Xantho</i>	49
<i>gracillimus, Grapsus</i>	67
<i>gracillipes, Hoplometopus</i>	95
<i>grandis, Medaeus</i>	41; 78; 98
<i>granosimana, Liomera, Pseudoliomera</i>	44
<i>granoso-manus, Xanthodes</i>	50
<i>granularis, Metasesarma</i>	70
<i>granulatus</i>	
<i>Carpilodes</i>	40
<i>Chlorodopsis</i>	56
<i>Coenobita</i>	13
<i>Neptunus</i>	31
<i>Phymodius</i>	55
<i>Portunus</i>	31
<i>granulosa</i>	
<i>Ashtoret</i>	25; 76
<i>Lophactea</i>	47
<i>Platypodia</i>	47
<i>Ruppellia</i>	64; 78
<i>granulosus, Phymodius</i>	54
<i>Grapsus</i>	
<i>albolineatus</i>	67
<i>depressus</i>	67; 77; 78
<i>gracilipes</i>	67
<i>gracillimus</i>	67
<i>grapsus</i>	67
<i>longitarsis</i>	67
<i>maculatus</i>	67
<i>pictus</i>	67
<i>strigosus</i>	67
<i>tenuicrustatus</i>	67
<i>grayi, Geograpsus</i>	66

<i>guamensis, Calcinus</i>	14; 76
<i>guinotae</i>	
<i>Percnon</i>	71; 76
<i>Portunus</i>	4; 31
<i>guttata, Trapezia</i>	58; 59
<i>guttatus</i>	
<i>Dardanus</i>	17; 77
<i>Pagurus</i>	17

H

<i>haani, Dardanus</i>	18
<i>Hapalocarcinus marsupialis</i>	74
<i>harmsi, Liocarpilodes, Pilodius</i>	54
<i>harpax, Thalassograpus</i>	69; 77
<i>hawaiensis</i>	
<i>Charybdis</i>	30; 78
<i>Mursia</i>	97
<i>Ozius</i>	64
<i>Hellenus, Portunus</i>	32
<i>Hemigrapsus crenulatus</i>	69
<i>Hepatella</i>	28
<i>hepatica, Calappa</i>	25; 97
<i>Hepatus</i>	28
<i>herbstii, Calcinus</i>	14; 15
<i>heterodactyla, Tetralia</i>	57
<i>Heterograpus crenulatus</i>	69
<i>Heteronucia venusta</i>	26
<i>Heteropanope sp.</i>	99
<i>hexagonum, Parathranites</i>	98
<i>hilgendorfi, Coenobita</i>	12
<i>Hippa</i>	
<i>adactyla</i>	23
<i>ovalis</i>	5; 23
<i>pacifica</i>	23
<i>sp.</i>	23
<i>hippocrepica, Paractaea</i>	43
<i>hirsuta, Poupinia</i>	97
<i>hirsutissima, Actaea</i>	42
<i>hirsutissimus, Actaeodes</i>	42
<i>hirtipes</i>	
<i>Cardisoma</i>	65
<i>Trapezia</i>	58
<i>hispida</i>	
<i>Domecia</i>	63
<i>Dynomene</i>	24
<i>holthuisi</i>	
<i>Enoplometopus</i>	8; 77
<i>Parribacus</i>	5; 11; 78
<i>homarus, Panulirus</i>	9
<i>Homola</i>	
<i>ikedai</i>	97
<i>orientalis</i>	97
<i>Homologenus broussei</i>	97
<i>Hoplometopus gracilipes</i>	95
<i>hoplonotus, Aulacolambrus, Parthenope</i>	28
<i>horrida, Daldorfia, Parthenope</i>	28
<i>Huenia proteus</i>	27
<i>humilis</i>	
<i>Clibanarius</i>	16
<i>Coralliogalathea</i>	19

Hyastenus

<i>borradalei</i>	27
<i>cristimanus</i>	26
<i>hydrodromus, Potamon</i>	77
<i>Hypocolpus mararae</i>	98
<i>Hypsophrys</i>	
<i>inflata</i>	97
<i>murotoensis</i>	97
<i>personata</i>	97
<i>hystrix, Munida</i>	96

I

<i>ihlei, Cyrtomaia</i>	98
<i>ikedai, Homola</i>	97
<i>immaculata, Plagusia</i>	72; 76
<i>imperialis, Calcinus</i>	14; 76
<i>incisus, Lophozozymus</i>	46; 76
<i>inflata, Hypsophrys</i>	97
<i>insularis, Neoliomera</i>	40; 77
<i>integer, Cyclograpsus</i>	70
<i>integerrimus, Actumnus, Liocarpilodes</i>	54
<i>integra, Thalamita</i>	3; 35
<i>intermedius, Ptychognathus</i>	69
<i>iranjae, Portunus</i>	5; 32

J

<i>jacquinoti, Sesarma</i>	71; 77; 78
<i>Jonesius triunguiculatus</i>	57
<i>Justitia</i>	
<i>longimanus</i>	9; 96
<i>vericeli</i>	96
<i>Juxtaxanthias tetraodon</i>	51

K

<i>keiji, Eumunida</i>	97
<i>Kraussia</i>	
<i>marquesas</i>	51
<i>rastripes</i>	51
<i>kremphi, Ciliopagurus</i>	16

L

<i>Labuanium</i>	
<i>rotundatum</i>	70; 76
<i>trapezoideum</i>	70
<i>Lachnopodus</i>	
<i>bidentatus</i>	48
<i>ponapensis</i>	48
<i>subacutus</i>	48
<i>tahitensis</i>	48
<i>laevigata, Pinnotherelia</i>	72
<i>laevimana, Eriphia</i>	63

laevimanus

<i>Calcinus</i>	14; 81
<i>Etisus</i>	52; 53
<i>Eupagurus</i>	19
<i>Pagurixus</i>	19
<i>laevirostris, Leiogalathea</i>	96
<i>laevis</i>	
<i>Liomera</i>	39
<i>Lissocarcinus</i>	30
<i>Ocypode</i>	73
<i>laevissima, Chlorodiella</i>	53
<i>laevissimus, Chlorodiella, Chlorodius</i>	53
<i>laeviuscula, Carupa</i>	29
<i>lagopodes, Dardanus</i>	17
<i>lamarcki</i>	
<i>Petrolisthes</i>	21
<i>Xanthias</i>	50
<i>lamelligera, Parthenope</i>	28
<i>laperousei, Liomera</i>	39; 77
<i>lata</i>	
<i>Actaea</i>	45
<i>Liomera</i>	39
<i>Pseudoliomera</i>	44
<i>latens, Calcinus</i>	15
<i>latifrons, Xanthias</i>	50
<i>latirostris, Galathea</i>	20; 77
<i>latreillei, Gelasimus</i>	74
<i>Latreillia metanesa</i>	97
<i>latro, Birgus</i>	11; 80; 81
<i>latus, Notopoides</i>	97
<i>laysani, Tweedieia</i>	57
<i>Leiogalathea laevirostris</i>	96
<i>lenticularis, Munida</i>	96
<i>leptochelis, Lybia</i>	37
<i>Leptodius</i>	
<i>crassimanus</i>	49
<i>cristatus</i>	48
<i>davaoensis</i>	48
<i>exaratus</i>	48
<i>gracilis</i>	49
<i>leptodon</i>	48
<i>sanguineus</i>	49
<i>leptodon, Leptodius</i>	48
<i>Leptograpsus variegatus</i>	67
<i>leschenaudii, Thelphusa</i>	77
<i>lewinsohni, Quadrella</i>	57
<i>Libinia sp.</i>	99
<i>Libystes truncatifrons</i>	29
<i>limbata, Lophomicippa</i>	27
<i>limbatus, Atergatis</i>	47
<i>lineatum, Goniosoma</i>	36
<i>Liocarpilodes</i>	
<i>armiger</i>	54
<i>harmsi</i>	54
<i>integerrimus</i>	54
<i>sp.</i>	64
<i>Liomera</i>	
<i>bella</i>	38
<i>cinctimana</i>	39
<i>granosimana</i>	44
<i>laevis</i>	39
<i>laperousei</i>	39; 77
<i>lata</i>	39
<i>monticulosa</i>	39

<i>pallida</i>	39	<i>nudipes</i>	50
<i>richtersi</i>	41	<i>Macrophthalmus</i>	
<i>rubra</i>	39	<i>consobrinus</i>	72; 78; 80
<i>rugata</i>	38; 39	<i>convexus</i>	73; 80
<i>rugipes</i>	38	<i>parvimanus</i>	72
<i>semigranosa</i>	40	<i>serenei</i>	73
<i>stimpsoni</i>	40	<i>macrophthalmus, Portunus</i>	32; 76
<i>tristis</i>	3; 40	<i>macropus, Thalamita</i>	35; 76
<i>venosa</i>	40	<i>macrospinifera, Thalamita</i>	35; 98
<i>Lioxanthodes alcocki</i>	49	<i>maculata, Trapezia</i>	61
<i>Lissocarcinus</i>		<i>maculatus</i>	
<i>elegans</i>	5; 30; 78	<i>Carpilius</i>	62
<i>laevis</i>	30	<i>Grapsus</i>	67
<i>orbicularis</i>	30	<i>Neopetrolisthes</i>	20; 77
<i>Lithodes megacantha</i>	97	<i>maculosa, Quadrella</i>	57; 76
<i>litterata, Varuna</i>	6; 69	<i>magnificus, Palibythus</i>	7; 10; 96
<i>lividus</i>		<i>major, Ciliopagurus</i>	96
<i>Calcinus</i>	15	<i>majora, Moloha</i>	97
<i>Geograpsus</i>	66	<i>manquenei, Beuroisia</i>	98
<i>Pagurus</i>	15	<i>maorus, Eupagurus, Pagurixus</i>	19
<i>lobatus, Ozius</i>	65	<i>mararae</i>	
<i>longicheles, Munida</i>	96	<i>Hypocolpus</i>	98
<i>longicornis, Pilumnus</i>	61	<i>Progeryon</i>	98
<i>longimanus, Justitia</i>	9; 96	<i>margaritatus, Pilumnus</i>	54; 64
<i>longipes</i>		<i>margaritifera, Micippa</i>	27
<i>Cyclograpsus</i>	70	<i>marquesas</i>	
<i>Discoplax</i>	66	<i>Kraussia</i>	51
<i>Pachygrapsus</i>	68	<i>Palapedia</i>	5; 51
<i>Panulirus</i>	9	<i>marsupialis, Hapalocarcinus</i>	74
<i>longispinosus, Neptunus, Portunus</i>	32	<i>mathaei, Elamena</i>	75
<i>longitarsis, Grapsus</i>	67	<i>Mathildella maxima</i>	98
<i>Lophactea granulosa</i>	47	<i>Matuta</i>	
<i>Lophomicippa limbata</i>	27	<i>banksi</i>	25
<i>Lophozomyrus</i>		<i>picta</i>	25
<i>bertonciniae</i>	98	<i>victor</i>	25; 77
<i>cristatus</i>	46	<i>maxima, Mathildella</i>	98
<i>dodone</i>	46	<i>maximus, Aniculus</i>	14
<i>edwardsi</i>	46	<i>Medaeus</i>	
<i>glaber</i>	46	<i>grandis</i>	41; 78; 98
<i>incisus</i>	46; 76	<i>noelensis</i>	42; 49
<i>octodentatus</i>	46	<i>megacantha, Lithodes</i>	97
<i>pictor</i>	46; 77	<i>megalochira, Galathea</i>	19
<i>sp.</i>	47	<i>megistos, Dardanus</i>	18
<i>superbus</i>	5; 46	<i>melanodactyla, Cymo</i>	38
<i>lunaris, Ashtoret</i>	25	<i>melanodactylus, Cymo</i>	38
<i>Lupocyclus quinquedentatus</i>	31	<i>Melia tessellata</i>	37
<i>Lybia</i>		<i>Menaethius</i>	
<i>caestifera</i>	37	<i>monoceros</i>	27
<i>leptochelis</i>	37	<i>tuberculatus</i>	27
<i>plumosa</i>	37	<i>Meractaea tafai</i>	98
<i>sp.</i>	99	<i>Meriola rufomaculata</i>	98
<i>tessellata</i>	37	<i>merodontatus, Pilumnus</i>	61
<i>Lydia annulipes</i>	64; 78	<i>messor, Metopograpsus</i>	67
<hr/>			
M		<i>metanesa, Latreillia</i>	97
<i>macrocera, Ocypode</i>	76	<i>Metasesarma</i>	
<i>macrodactylus, Etisus</i>	52; 53	<i>granularis</i>	70
<i>Macromedaeus</i>		<i>rousseauxii</i>	70
<i>crassimanus</i>	49	<i>rugulosa</i>	70
<i>distinguendus</i>	49; 77	<i>Metopograpsus</i>	
		<i>messor</i>	67
		<i>thukuhar</i>	68
		<i>Micippa</i>	
		<i>margaritifera</i>	27
		<i>parca</i>	27

<i>Micippoides angustifrons</i>	27
<i>Micropagurus polynesiensis</i>	18; 78
<i>militaris, Petrolisthes</i>	21; 77
<i>mimetica, Nursia</i>	26; 78
<i>miniata, Trapezia</i>	59
<i>minuscula, Thalamita</i>	35; 77; 78
<i>minutus</i>	
<i>Calcinus</i>	6; 15; 77
<i>Pachygrapsus</i>	68
<i>mitra, Porcellana</i>	22; 77
<i>mitsiensis, Thalamita</i>	35; 76
<i>miyakei, Sadayoshia</i>	20
<i>Moloha majora</i>	97
<i>monoceros, Menaethius</i>	27
<i>monolifera, Porcellana</i>	22; 77; 78
<i>monticulosa, Liomera</i>	39
<i>monticulosus</i>	
<i>Carpilodes</i>	39; 40
<i>Chlorodius</i>	55
<i>Phymodius</i>	3; 55
<i>mortenseni, Demania</i>	98
<i>Munida</i>	
<i>amathea</i>	96
<i>ducoussoi</i>	96
<i>evarne</i>	96
<i>hystrix</i>	96
<i>lenticularis</i>	96
<i>longicheles</i>	96
<i>normani</i>	96
<i>ocellata</i>	96
<i>pasithea</i>	97
<i>pilosimanus</i>	97
<i>plexaura</i>	97
<i>polynoe</i>	97
<i>profunda</i>	97
<i>pulchra</i>	97
<i>rubella</i>	97
<i>rubrovata</i>	97
<i>solae</i>	97
<i>murotoensis, Hypsophrys</i>	97
<i>Mursia hawaiensis</i>	97
<i>mururoa, Epistocavea</i>	98

N

<i>Nanocassiope oblonga</i>	95; 98
<i>Naxioides</i>	
<i>teatui</i>	98
<i>vaitahu</i>	98
<i>Nectograpsus politus</i>	66
<i>Neoliomera</i>	
<i>demari</i>	40; 41
<i>insularis</i>	40; 77
<i>pubescens</i>	40; 41; 77
<i>richtersi</i>	41
<i>sp.</i>	99
<i>variolosa</i>	41; 45
<i>Neopetrolisthes</i>	
<i>maculatus</i>	20; 77
<i>ohshimai</i>	20
<i>Neothalaminella, Thalamita</i>	36

<i>Neothalamita, Thalamita</i>	33; 35; 36
<i>Neoxanthops cavatus</i>	50
<i>Neptunus</i>	
<i>longispinosus</i>	32
<i>pelagicus</i>	32
<i>sanguinolentus</i>	32
<i>niger, Chlorodius</i>	54
<i>nigra, Chlorodiella</i>	54
<i>nigrifrons, Tetralia, Tetraloides</i>	58
<i>nipponensis, Portunus</i>	32; 98
<i>nitidulus</i>	
<i>Phymodius</i>	55
<i>Pilodius</i>	55
<i>Xanthias</i>	50; 77
<i>Xanthodes</i>	50
<i>nitidus</i>	
<i>Calcinus</i>	4; 15; 78
<i>Catoptrus</i>	29
<i>noelensis</i>	
<i>Medaeus</i>	41; 49
<i>Paramedaeus</i>	41
<i>normani, Munida</i>	96
<i>notatus, Paraxanthias, Xanthias, Xanthodes</i>	50
<i>Notopoides latus</i>	97
<i>Notosceles</i>	
<i>chimmonis</i>	24; 97
<i>viaderi</i>	97
<i>Nucia</i>	
<i>gelida</i>	26
<i>rosea</i>	26; 78
<i>nudipes, Macromedaeus, Xantho</i>	50
<i>Nursia mimetica</i>	26; 78

O

<i>obesum, Cardisoma</i>	65
<i>obesus, Actumnus</i>	61
<i>oblonga, Nanocassiope</i>	95; 98
<i>obtusirostris, Simocarcinus, Trigonothir</i>	28
<i>ocellata, Munida</i>	96
<i>octodentatus, Lophozozymus</i>	46
<i>Ocypode</i>	
<i>ceratophtalma</i>	73
<i>cordimana</i>	73
<i>laevis</i>	73
<i>macrocera</i>	76
<i>pallidula</i>	3; 73
<i>platytarsis</i>	76
<i>urvillei</i>	73
<i>ocyroe, Atergatis</i>	45
<i>odhneri, Tweedieia</i>	57
<i>ohshima, Neopetrolisthes</i>	20
<i>olivieri, Coenobita</i>	12; 13; 77
<i>orbicularis, Lissocarcinus</i>	30
<i>orbitosinus, Portunus</i>	32; 76
<i>Oreotlos</i>	
<i>encymus</i>	97
<i>potanus</i>	7; 98
<i>orientalis</i>	
<i>Charybdis</i>	31
<i>Homola</i>	97

<i>ornatus</i>	
<i>Cyclodius</i>	55
<i>Panulirus</i>	10; 76
<i>Osachila</i>	28
<i>ovalis, Hippa</i>	5; 23
<i>Ozius</i>	
<i>hawaiensis</i>	64
<i>lobatus</i>	65
<i>rugulosus</i>	64
<i>tricarinatus</i>	64; 78
<i>truncatus</i>	65; 77
<i>tuberculosus</i>	65
 P	
<i>Pachycheles</i>	
<i>pisoides</i>	20
<i>sculptus</i>	20
<i>Pachygrapsus</i>	
<i>fakaravensis</i>	4; 68
<i>longipes</i>	68
<i>minutus</i>	68
<i>planifrons</i>	68
<i>plicatus</i>	68; 81
<i>sp.</i>	99
<i>transversus</i>	76
<i>pacifica, Hippa</i>	23
<i>acificus</i>	
<i>Bathynarius</i>	96
<i>Ciliopagurus</i>	96
<i>Panopeus</i>	51; 78
<i>Remipes</i>	23
<i>Pagurixus</i>	
<i>anceps</i>	19
<i>laevimanus</i>	19
<i>maorus</i>	19
<i>Pagurus</i>	
<i>aniculus</i>	14
<i>deformis</i>	17
<i>diformis</i>	17
<i>euopsis</i>	18
<i>gemmaeus</i>	17
<i>guttatus</i>	17
<i>lividus</i>	15
<i>punctulatus</i>	18
<i>sanguinolentus</i>	18
<i>spinimanus</i>	18
<i>strigatus</i>	16
<i>Palapedia</i>	
<i>marquesas</i>	5; 51
<i>rastripes</i>	51
<i>Palibythus magnificus</i>	7; 10; 96
<i>Palinurellus wieneckii</i>	10; 76; 77
<i>Palinurus spinosus</i>	9
<i>Palinustus unicornutus</i>	96
<i>pallida, Liomera</i>	39
<i>pallidula, Ocypode</i>	3; 73
<i>pallidus, Carpilodes</i>	39
<i>Panopeus pacificus</i>	51; 78
<i>Panulirus</i>	
<i>fasciatus</i>	10

<i>homarus</i>	9
<i>longipes</i>	9
<i>ornatus</i>	10; 76
<i>pascuensis</i>	5; 9; 81
<i>penicillatus</i>	9; 81
<i>polyphagus</i>	9; 10; 76
<i>spinosus</i>	9
<i>versicolor</i>	10
<i>Paractaea</i>	
<i>excentrica</i>	43
<i>quadriareolata</i>	44
<i>retusa</i>	43
<i>rufopunctata</i>	44
<i>tumulosa</i>	44
<i>Paractaeopsis</i>	
<i>quadriareolatus</i>	44
<i>tumulosus</i>	44
<i>Paramedaeus</i>	
<i>noelensis</i>	41
<i>simplex</i>	42; 77
<i>sp.</i>	99
<i>Parapleurophrycoides roseus</i>	75; 77; 78
<i>Parathranites hexagonum</i>	98
<i>Paraxanthias notatus</i>	50
<i>Paraxanthodes polynesiensis</i>	98
<i>parca, Micippa</i>	27
<i>Parribacus</i>	
<i>antarcticus</i>	10; 11
<i>holthuiisi</i>	5; 11; 78
<i>scarlatinus</i>	11
<i>ursus-major</i>	11
<i>Parthenope</i>	
<i>contrarius</i>	28
<i>hoplonotus</i>	28
<i>horrida</i>	28
<i>lamelligera</i>	28
<i>pelagicus</i>	28
<i>poupini</i>	98
<i>sp.</i>	99
<i>stellata</i>	98
<i>parvimanus, Macrophthalmus</i>	72
<i>parvula, Actaea, Banareia</i>	38
<i>parvulus</i>	
<i>Cyclograpsus</i>	70
<i>Pilumnus</i>	62
<i>pascuensis, Panulirus</i>	5; 9; 81
<i>pasithea, Munida</i>	97
<i>paucidentata, Charybdis</i>	7; 31; 98
<i>paumotensis, Pilodius</i>	56
<i>pedunculatus, Dardanus</i>	18
<i>pelagicus</i>	
<i>Neptunus</i>	32
<i>Parthenope</i>	28
<i>Portunus</i>	32; 77
<i>penicillatus, Panulirus</i>	9; 81
<i>pentagonus, Echinoecus, Eumedonous</i>	29
<i>Percnon</i>	
<i>abbreviatum</i>	71
<i>affine</i>	71
<i>affinis</i>	71
<i>guinotae</i>	76
<i>pilimanus</i>	71
<i>planissimum</i>	71
<i>planissimus</i>	71

<i>Perigrapsus excelsus</i>	65	spp.	99
<i>Perinea tumida</i>	27	<i>tahitensis</i>	62
<i>perlata</i>		<i>Pinnotherelia laevigata</i>	72
<i>Coenobita</i>	13	<i>pisoides, Pachycheles</i>	20
<i>Daira</i>	75	<i>Plagusia</i>	
<i>perlatus, Coenobita</i>	12; 13; 77; 81	<i>depressa</i>	72
<i>perpusillus, Platyozius</i>	75; 77; 78	<i>immaculata</i>	72; 76
<i>personata, Hypsophrys</i>	97	<i>speciosa</i>	3; 72
<i>Petalomera wilsoni</i>	24	<i>squamosa</i>	72
<i>Petrolisthes</i>		<i>tuberculata</i>	72
<i>bispinosus</i>	21	<i>Planes cyaneus</i>	68; 76
<i>borradalei</i>	21; 22	<i>planifrons, Pachygrapsus</i>	68
<i>bosci</i>	21	<i>planimanus, Sympagurus</i>	96
<i>coccineus</i>	21	<i>planissimum, Percnon</i>	71
<i>decacanthus</i>	21	<i>planissimus, Acanthopus, Percnon</i>	71
<i>eldredgei</i>	21	<i>planus, Pseudozius</i>	65
<i>elegans</i>	21	<i>Platepistoma balssii</i>	98
<i>lamarckii</i>	21	<i>Platylambrus, Parthenope</i>	98
<i>militaris</i>	21; 77	<i>Platyozius perpusillus</i>	75; 77; 78
<i>pubescens</i>	22	<i>Platypodia</i>	
<i>rufescens</i>	21; 22; 77	<i>anaglypta</i>	46
<i>scabriculus</i>	22	<i>digitalis</i>	61
<i>sp.</i>	99	<i>granulosa</i>	47
<i>tomentosus</i>	22	<i>pseudogranulosa</i>	47
<i>philippinensis, Thalamita</i>	35; 76	<i>semigranosa</i>	47
<i>Phylladiorhynchus</i>		<i>platytarsis, Ocypode</i>	76
<i>pusillus</i>	20	<i>plessisi, Ciliopagurus</i>	96
<i>serrirostris</i>	20	<i>Pleurocolpus boileaui</i>	95; 98
<i>Phymodius</i>		<i>plexaura, Munida</i>	97
<i>granulatus</i>	55	<i>plicatus, Pachygrapsus</i>	68; 81
<i>granulosus</i>	54	<i>plumosa</i>	
<i>monticulosus</i>	3; 55	<i>Lybia</i>	37
<i>nitidulus</i>	55	<i>Paractaea</i>	44
<i>sp.</i>	99	<i>Podophthalmus vigil</i>	37
<i>ungulatus</i>	55	<i>politus</i>	
<i>picta</i>		<i>Epigrapsus</i>	4; 66; 70
<i>Ashtoret</i>	25	<i>Nectograpsus</i>	66
<i>Matuta</i>	25	<i>polyacantha, Actaea</i>	42
<i>Thalamita</i>	35	<i>Polydectus</i>	
<i>pictor, Lophozozymus</i>	46; 77	<i>cupulifer</i>	37
<i>pictus, Grapsus</i>	67	<i>villosus</i>	37
<i>pilimanus, Percnon</i>	71	<i>polynesiensis</i>	
<i>Pilodius</i>		<i>Micropagurus</i>	18; 78
<i>aberrans</i>	54	<i>Paraxanthodes</i>	98
<i>areolatus</i>	55	<i>polynoe, Munida</i>	97
<i>flavus</i>	56	<i>polyphagus, Panulirus</i>	9; 76
<i>harmsi</i>	54	<i>ponapensis, Lachnopodus, Paraxanthias, Xanthias</i>	48
<i>nitidulus</i>	55	<i>Porcellana</i>	
<i>paumotensis</i>	56	<i>coccinea</i>	21
<i>pubescens</i>	56	<i>mitra</i>	22; 77
<i>pugil</i>	56; 64	<i>monilifera</i>	22; 77; 78
<i>scabriculus</i>	55; 56	<i>rufescens</i>	22
<i>sp.</i>	99	<i>speciosa</i>	21
<i>spinipes</i>	56	<i>tomentosa</i>	22
<i>pilosimanus, Munida</i>	97	<i>Portunus</i>	
<i>pilumnoides, Thalamita</i>	36	<i>alexandri</i>	31; 77; 78
<i>Pilumnus</i>		<i>dubius</i>	31
<i>globosus</i>	56; 64	<i>granulatus</i>	31
<i>longicornis</i>	61	<i>guinotae</i>	4; 31
<i>margaritatus</i>	54; 64	<i>iranjae</i>	5; 32
<i>merodontatus</i>	61	<i>longispinosus</i>	32
<i>parvulus</i>	62	<i>macrophthalmus</i>	32; 76
<i>ransonni</i>	5; 62	<i>nipponensis</i>	32; 98
		<i>orbitosinus</i>	32; 76

<i>pelagicus</i>	32; 77
<i>sanguinolentus</i>	32
<i>Potamon hydrodromus</i>	77
<i>potanus, Oretlos</i>	7; 98
<i>poupini</i>	
<i>Chaceon</i>	98
<i>Parthenope</i>	98
<i>Strigopagurus</i>	96
<i>Sympagurus</i>	96
<i>Poupinia hirsuta</i>	97
<i>praedator, Dynomene</i>	24
<i>profunda, Munida</i>	97
<i>Progeryon mararae</i>	98
<i>proteus, Huenia</i>	27
<i>prymna, Thalamita</i>	36
<i>Psaumis</i>	
<i>cavipes</i>	44
<i>cellulosa</i>	44
<i>pseudogranulosa, Platypodia</i>	47
<i>Pseudograpsus albus</i>	69
<i>Pseudoliomera</i>	
<i>granosimana</i>	44
<i>lata</i>	44
<i>rupPELLIOIDES</i>	42; 45
<i>speciosa</i>	45
<i>variolosa</i>	45
<i>Pseudothalamitopsis, Thalamita</i>	33; 36
<i>Pseudozius</i>	
<i>caystrus</i>	65
<i>planus</i>	65
<i>Ptychognathus</i>	
<i>crassimanus</i>	69; 77; 78
<i>easteranus</i>	6; 69
<i>intermedius</i>	69
<i>pubescens</i>	
<i>Neoliomera</i>	40; 41; 77
<i>Petrolisthes</i>	22
<i>Pilodius</i>	56
<i>pugil</i>	
<i>Chlorodopsis</i>	56
<i>Pilodius</i>	56; 64
<i>pulchellus, Gelasimus</i>	74
<i>pulchra, Munida</i>	97
<i>pumilus, Zozymodes</i>	47
<i>punctata, Actaeomorpha</i>	28
<i>punctatus</i>	
<i>Etisus</i>	53; 77; 78
<i>Xanthias</i>	51
<i>punctimanus, Trapezia</i>	6; 58; 60
<i>punctulatus, Pagurus</i>	18
<i>pusillus, Phylladiorhynchus</i>	20

Q

<i>Quadrella</i>	
<i>cyrenae</i>	57
<i>lewinsohni</i>	57
<i>maculosa</i>	57; 76
<i>sp.</i>	57

<i>quadriareolata, Paractaea, Paractaeopsis</i>	44
<i>quadridens, Thalamita</i>	36
<i>quadrilobata, Thalamita</i>	5; 36
<i>quadrilobatus, Cymo</i>	38
<i>quinquedentatus, Lupocyclus</i>	31

R

<i>Randallia serenei</i>	98
<i>Ranina ranina</i>	24
<i>ransonii</i>	
<i>Clibanarius</i>	5; 16
<i>Pilumnus</i>	5; 62
<i>rapanus, Scyllarus</i>	95; 96
<i>rastripes, Kraussia, Palapedia</i>	51
<i>Rata tuamotense</i>	98
<i>regalis, Arctides</i>	10; 76; 77
<i>Remipes</i>	
<i>pacificus</i>	23
<i>testudinarius</i>	23
<i>reticulata, Trapezia</i>	58
<i>retusa</i>	
<i>Camposcia</i>	26
<i>Paractaea</i>	43
<i>rhabdodactylus, Clibanarius</i>	16
<i>Rhinolambrus, Parthenope</i>	28
<i>richtersi, Actaeodes, Liomera, Neoliomera</i>	41
<i>rimatara, Alainodaeus</i>	7; 41; 78; 98
<i>rosea, Nucia</i>	26; 78
<i>roseus, Parapleurophrycoides</i>	75; 77; 78
<i>rotundatum, Labuanium</i>	70; 76
<i>rotundifrons, Caphyra</i>	29
<i>rotundum, Cardisoma</i>	66
<i>rousseauxii, Metasesarma, Sesarma</i>	70
<i>rubella, Munida</i>	97
<i>rubra, Liomera</i>	39
<i>rubrovata, Munida</i>	97
<i>rueppelli, Gaillardiellus</i>	43; 77
<i>rufescens</i>	
<i>Petrolisthes</i>	21; 22; 77
<i>Porcellana</i>	22
<i>rufomaculata, Meriola</i>	98
<i>rufopunctata</i>	
<i>Actaea</i>	43; 44
<i>Paractaea</i>	43; 44
<i>Trapezia</i>	59; 60; 61
<i>rugata, Liomera</i>	38; 39
<i>rugatus, Carpilodes</i>	38; 40
<i>rugipes, Carpiloxanthus, Liomera</i>	38
<i>rugosa, Cenobita</i>	13
<i>rugosus</i>	
<i>Coenobita</i>	13
<i>Etisus</i>	52
<i>Euxanthus</i>	41
<i>rugulosa, Metasesarma</i>	70
<i>rugulosus, Ozius</i>	64
<i>Ruppellia</i>	
<i>annulipes</i>	64
<i>granulosa</i>	64; 78
<i>rupPELLIOIDES, Actaea, Pseudoliomera</i>	42; 45

S

<i>Sadayoshia</i>	
<i>edwardsii</i>	97
<i>miyakei</i>	20
<i>sanguineus, Chlorodius, Leptodius, Xantho</i>	49
<i>sanguinolentus</i>	
<i>Dardanus</i>	18
<i>Neptunus</i>	32
<i>Pagurus</i>	18
<i>Portunus</i>	32
<i>Sarmatium crassum</i>	70
<i>scabra</i>	
<i>Actaea</i>	43
<i>Forestia</i>	43; 77
<i>scabricula</i>	
<i>Chlorodopsis</i>	55
<i>Domecia</i>	63
<i>scabriculus</i>	
<i>Petrolisthes</i>	22
<i>Pilodius</i>	55; 56
<i>scarlatinus, Parribacus</i>	11
<i>Schizophrys aspera</i>	27; 77
<i>scruposa, Aethra</i>	28; 77
<i>sculptilis, Euxanthus</i>	41
<i>sculptus, Pachycheles</i>	20
<i>scutellatus, Dardanus, Pagurus</i>	18
<i>Scylla serrata</i>	33; 81
<i>Scyllarus</i>	
<i>antarcticus</i>	10
<i>aurora</i>	7; 11; 96
<i>rapanus</i>	95; 96
<i>sp.</i>	11
<i>sebana, Eriphia</i>	63
<i>semigranosa</i>	
<i>Liomera</i>	40
<i>Platypodia</i>	47
<i>semistriatus, Clibanarius</i>	16
<i>Senex femoristriga</i>	9
<i>septata, Trapezia</i>	58; 60; 77
<i>serenei</i>	
<i>Macrophthalmus</i>	73
<i>Randallia</i>	98
<i>Trapezia</i>	6; 59; 60
<i>serrata, Scylla</i>	33; 81
<i>serratifrons, Tetralia</i>	57
<i>serirostris, Galathea, Phylladiorhynchus</i>	20
<i>Sesarma</i>	
<i>angustifrons</i>	71; 78
<i>jacquinoti</i>	71; 77; 78
<i>rousseauxi</i>	70
<i>trapezoidea</i>	70
<i>setifer, Actumnus</i>	61
<i>seurati</i>	
<i>Calcinus</i>	4; 15
<i>Thalamita</i>	4; 36
<i>sibogae, Strobopagurus</i>	96
<i>signatus, Atergatopsis</i>	45
<i>Simocarcinus obtusirostris</i>	28
<i>simplex, Paramedaeus</i>	42; 77
<i>sinense, Dynomene</i>	24; 76
<i>solae, Munida</i>	97

Solitariopagurus sp. nov...... 95; 96
speciosa

<i>Actaea</i>	45
<i>Albunea</i>	22; 76; 78
<i>Plagusia</i>	3; 72
<i>Porcellana</i>	21
<i>Pseudoliomera</i>	45
<i>Trapezia</i>	58; 60
<i>Sphaerodromia ducoussoi</i>	97
<i>spicatus, Calcinus</i>	4; 15
<i>spinata, Uca</i>	74
<i>spinifera, Thalamita</i>	7; 36
<i>spinimanus</i>	
<i>Pagurus</i>	18
<i>Yaldwynopsis</i>	97
<i>spinipes, Chlorodopsis, Pilodius</i>	56
<i>spinosa, Dynomene</i>	24
<i>spinosus</i>	
<i>Coenobita</i>	12; 13; 77
<i>Palinurus</i>	9
<i>Panulirus</i>	9
<i>splendidus, Etisus</i>	53
<i>squamosa, Plagusia</i>	72
<i>stellata, Parthenope</i>	98
<i>stimpsoni, Liomera</i>	40
<i>stormi, Geograpsus</i>	66
<i>strigatus, Ciliopagurus, Pagurus, Trizopagurus</i>	16
<i>Strigopagurus poupinii</i>	96
<i>strigosus, Grapsus</i>	67
<i>striolatus, Clibanarius</i>	17
<i>Strobopagurus</i>	
<i>gracilipes</i>	96
<i>sibogae</i>	96
<i>subacutus, Lachnopodus</i>	48
<i>suborbicularis, Cyclax</i>	26
<i>subquadratus, Chasmagnathus</i>	69; 77
<i>superbus</i>	
<i>Lophozozymus</i>	5; 46
<i>Xantho</i>	46
<i>superciliaris, Actaea, Gailliellus</i>	43
<i>Sympagurus</i>	
<i>affinis</i>	96
<i>boletifer</i>	96
<i>bougainvillei</i>	96
<i>dofleini</i>	96
<i>planimanus</i>	96
<i>poupini</i>	96
<i>trispinosus</i>	96
<i>tuamotu</i>	96
<i>wallisi</i>	96

T

<i>tahitensis</i>	
<i>Lachnopodus, Xantho</i>	48
<i>Pilumnus</i>	62
<i>tanensis, Dynomene</i>	97
<i>teatui, Naxioides</i>	98
<i>tenuicrustatus, Grapsus</i>	67
<i>tenuifrons, Acanthopus</i>	71

<i>tenuipes</i> , <i>Carupa</i>	29	<i>Porcellana</i>	22
<i>terrae-reginae</i> , <i>Calcinus</i>	15	<i>tomentosus</i>	
<i>tessellata</i> , <i>Lybia</i> , <i>Melia</i>	37	<i>Actaeodes</i>	43
<i>testudinarius</i> , <i>Remipes</i>	23	<i>Actumnus</i>	61
<i>tetragonon</i> , <i>Gelasimus</i> , <i>Uca</i>	74	<i>Petrolisthes</i>	22
<i>Tetralia</i>		<i>transversus</i> , <i>Pachygrapsus</i>	76
<i>cavimana</i>	57		
<i>cinctipes</i>	57	<i>Trapezia</i>	
<i>glaberrima</i>	57	<i>areolata</i>	3; 58; 60
<i>heterodactyla</i>	57	<i>bella</i>	3; 58; 60
<i>nigrifrons</i>	58	<i>cymodoce</i>	58; 59; 60; 61
<i>serratifrons</i>	57	<i>davaoensis</i>	60
<i>Tetraloides nigrifrons</i>	58	<i>dentata</i>	58
<i>tetraodon</i>		<i>digitalis</i>	58; 59; 60
<i>Eudora</i>	51	<i>ferruginea</i>	58; 59; 60; 61
<i>Juxtaxanthias</i>	51	<i>flavopunctata</i>	59
<i>Xanthias</i>	51; 77	<i>formosa</i>	59; 77
<i>Xantho</i>	51	<i>fusca</i>	59
<i>Thalaminella</i> , <i>Thalamita</i>	33; 34	<i>guttata</i>	58; 59
<i>Thalamita</i>		<i>hirtipes</i>	58
<i>admete</i>	33	<i>maculata</i>	61
<i>alcocki</i>	36	<i>miniata</i>	59
<i>bouvieri</i>	33	<i>punctimanus</i>	6; 58; 60
<i>chaptali</i>	33	<i>reticulata</i>	58
<i>coerulipes</i>	33	<i>rufopunctata</i>	59; 60; 61
<i>cooperi</i>	33	<i>septata</i>	58; 60; 77
<i>corrugata</i>	5; 34	<i>serenei</i>	6; 59; 60
<i>crenata</i>	34	<i>speciosa</i>	58; 60
<i>dakini</i>	34	<i>tigrina</i>	5; 60
<i>danae</i>	5; 34; 76	<i>wardi</i>	61
<i>demanii</i>	34	<i>trapezoidea</i> , <i>Sesarma</i>	70
<i>edwardsi</i>	34	<i>trapezoideum</i> , <i>Labuanium</i>	70
<i>gardineri</i>	36	<i>treguieri</i> , <i>Eumunida</i>	95; 97
<i>gatavakensis</i>	4; 34	<i>tricarinatus</i> , <i>Ozius</i>	64; 78
<i>gloriensis</i>	34	<i>Trichopagurus trichophthalmus</i>	19
<i>gracilipes</i>	35	<i>trichophthalmus</i> , <i>Catapaguroides</i> , <i>Trichopagurus</i>	19
<i>integra</i>	3; 35	<i>tridens</i>	
<i>macropus</i>	35; 76	<i>Caphyra</i>	30
<i>macrospinifera</i>	35; 98	<i>Cryptodromiopsis</i>	24
<i>minuscula</i>	35; 77; 78	<i>trispinosus</i> , <i>Sympagurus</i>	96
<i>mitsiensis</i>	35; 76		
<i>philippinensis</i>	35; 76	<i>Carpilodes</i>	40
<i>picta</i>	35	<i>Liomera</i>	3; 40
<i>pilumnoides</i>	36	<i>triunguiculatus</i> , <i>Jonesius</i>	57
<i>prymna</i>	36	<i>Trizopagurus strigatus</i>	16
<i>quadrilobata</i>	5; 36	<i>truncatifrons</i> , <i>Libystes</i>	29
<i>seurati</i>	4; 36	<i>truncatus</i> , <i>Ozius</i>	65; 77
<i>spinifera</i>	7; 36	<i>tuamotense</i> , <i>Rata</i>	98
<i>woodmasoni</i>	36	<i>tuamotu</i> , <i>Sympagurus</i>	96
<i>Thalamitoides quadridens</i>	36		
<i>Thalamitopsis</i> , <i>Thalamita</i>	34	<i>tuberculata</i>	
<i>Thalamonyx gracilipes</i>	35	<i>Calappa</i>	25
<i>Thalamonyx</i> , <i>Thalamita</i>	35	<i>Plagusia</i>	72
<i>Thalassograpsus harpax</i>	69; 77	<i>tuberculatus</i> , <i>Menaethius</i>	27
<i>Thalassuca</i> , <i>Uca</i>	74	<i>tuberculosus</i> , <i>Ozius</i>	65
<i>Thelphusa</i>		<i>tumida</i> , <i>Perinea</i>	27
<i>leschenaultii</i>	77	<i>tumulosa</i> , <i>Actaea</i> , <i>Paractaea</i> , <i>Paractaeopsis</i>	44
<i>wüllerstorfi</i>	77		
<i>theresae</i> , <i>Cancer</i>	10	<i>Tweedieia</i>	
<i>thukuhar</i> , <i>Metopograpsus</i>	68	<i>laysani</i>	57
<i>tibicen</i> , <i>Calcinus</i>	14	<i>odhneri</i>	57
<i>tigrina</i> , <i>Trapezia</i>	5; 60	<i>Tylocarcinus</i>	
<i>tomentosa</i>		<i>dumerili</i>	28
<i>Actaea</i>	43	<i>gracilis</i>	28

U*Uca*

- chlorophthalmus* 74; 80
crassipes 74
dussumieri 74; 76
gaimardi 74
spinata 74
tetragonon 3; 74
ungulatus
 Chlorodius 55
 Phymodius 55
unicornutus, Palinustus 96
ursus-major, Parribacus 11
urvillei, Ocypode 73

V

- vallantianus, Carpilodes* 38
vaitahu, Naxioides 98
variegatus, Leptograpsus 67
variolosa
 Neoliomera 41; 45
 Pseudoliomera 45
Varuna litterata 6; 69
venosa, Liomera 40
venosus, Carpilodes 40
venusta
 Chlorodopsis 57
 Heteronucia 26
verdensis, Carcinoplax 98
vericeli, Justitia 96
versicolor, Panulirus 10
viaderi, Notosceles 97
victor, Matuta 25; 77
vigil, Podophthalmus 37
villosus, Polydectus 37

W

- wallisi, Sympagurus* 96
wardi, Trapezia 61
wieneckii, Palinurellus 10; 76; 77
wilsoni
 Dromia 7; 24; 97
 Petalomera 24

-
- woodmasoni, Thalamita* 36
wüllerstorfi, Thelphusa 77

X

- Xanthias*
 canaliculatus 50
 lamarcki 50
 latifrons 50
 nitidulus 50; 51; 77
 notatus 50
 ponapensis 48
 punctatus 51
 sp. 99
 tetraodon 51; 77

Xantho

- arcuatus* 48
crassimanus 49
exaratus 49
gracilis 49
sanguineus 49
superbus 46
tahitensis 48
tetraodon 51

Xanthodes

- granoso-manus* 50
nitidulus 50
notatus 50
Xanthodius cristatus 47
xanthoides, Zozymodes 47; 48; 77
Xiphonectes, Portunus 31; 32

Y

-
- Yaldwynopsis spinimanus* 97

Z

- zebra, Clibanarius* 16; 17
Zozimus
 aeneus 47
 sp. 47
Zozymodes
 carinipes 47
 pumilus 47
 xanthoides 47; 48; 77