

THE FISHES OF PORTO RICO,

BY

BARTON WARREN EVERMANN, Ph. D.,
Ichthyologist of the United States Fish Commission,

AND

MILLARD CALEB MARSH,
Assistant, United States Fish Commission.



THE FISHES OF PORTO RICO.

BY BARTON W. EVERMANN AND MILLARD C. MARSH.

INTRODUCTION.

In preparing this report upon the fishes of Porto Rico it seemed desirable to include not only those obtained by us, but to give diagnoses of all the families and genera and detailed descriptions of all the species now known from those waters. That the species may be readily determined we have given very full keys leading to the definite identification of each, including certain genera and species which, though not yet actually known from Porto Rican waters, may hereafter be found to occur there. The unrecorded genera and species are, however, not numbered in the keys. The keys to the families are also of the same character.

The keys and all the diagnoses of families and genera are adapted from "The Fishes of North and Middle America," by Jordan & Evermann, which work we have followed in the arrangement and sequence of genera and species.

The descriptions of all the species of which we have specimens have either been wholly from Porto Rican material or verified upon specimens from that island, and in the majority of cases the color descriptions are from living specimens.

The common names inclosed in quotation marks are the names by which the species are known in Porto Rico. The other common names are those by which the species are known at Key West or elsewhere.

It has not been the intention to give much synonymy. In every case reference is made to the original description of each species, to all original synonyms, to Poey's and Stahl's papers on Porto Rican fishes, and to Jordan & Evermann's "Fishes of North and Middle America." The locality named at the end of each reference is that from which the type of the species came.

Of the 49 colored plates, 33 were made by Mr. A. H. Baldwin, and 16 by Mr. C. B. Hudson. All those by Mr. Baldwin were painted on board the *Fish Hawk*, the fish being placed in an aquarium as soon as caught and the life colors gotten before they had undergone any appreciable change.

Among the fishes painted from life by Mr. Hudson at Key West, Fla., in the winter of 1897-98, are 16 species which occur in Porto Rico, and his paintings of those are used in this report. The fish were selected from those (usually a large number) brought in by the fishermen in the live-wells of their boats and transferred to a large aquarium on the pier only a few feet distant from the fishing boats, where they were kept in as nearly a normal condition as possible while being painted. We can not give too high praise to the artists, Mr. Hudson and Mr. Baldwin, for the care

and fidelity with which they have done their work. All who are familiar with the life-colors of the species and who have seen these paintings have been struck with their accuracy, both as to color and structural detail.

The lithographers, Messrs. Julius Bien & Co., have given faithful and satisfactory reproductions of the originals.

Besides the colored plates of 49 species, 115 drawings were to be reproduced, each 5.25 inches long, as cuts illustrating the text. But as these pages were being written all of these drawings were destroyed by fire in New York. Steps were at once taken to have the 28 new species redrawn. Some of the others can not be replaced, but 82 of them had recently been reproduced in the "Fishes of North and Middle America," by Jordan & Evermann, and through the courtesy of the U. S. National Museum the same blocks are used for illustrating this report, but we were thus compelled to have them appear 4.25 inches long instead of 5.25 inches, as originally intended.

Another serious loss was incurred by the burning of the model and training school at San Juan, July 1, resulting in the total destruction of a valuable collection of fishes made in the vicinity of San Juan by Mr. Oscar Riddle, and his notes upon them, both of which he was about sending for use in preparing the present paper.

As already stated, the investigations of the Fish Commission in Porto Rican waters, though extending over about forty-five days and really embracing not over thirty-eight days of actual work, resulted in adding to the known fish fauna of the island nearly 200 species, among which were 3 genera and 33 species which are new to science. The 3 new genera and 20 of the new species were described by the present writers in the Annual Report of the U. S. Fish Commission for 1899.¹

One of the new eels (*Aphthalmichthys caribbeus*), the type of which was collected by Mr. G. M. Gray, was described by Drs. Gill and Smith in Science for June 22, 1900.²

The following is a full list of the new genera and species resulting from these investigations. Those which appear for the first time in this report are indicated by the star. The number following each name is that which the type specimen bears on the register of the U. S. National Museum, in which all the types are deposited. Cotypes of species of which duplicate specimens were obtained are in the reserve series of the U. S. Fish Commission and in the Museum of Leland Stanford Junior University.

Species.	Type No.	Species.	Type No.
<i>Aphthalmichthys caribbeus</i>		<i>Bollmannia boqueronensis</i>	49366
<i>Sphagebranchus ophioneus</i> *.....	49526	<i>Microgobius mceki</i>	49367
<i>Lycodontis albimentis</i> *.....	49527	<i>Gillias</i>	
<i>Lycodontis jordani</i>	49358	<i>Gillias jordani</i>	49368
<i>Stolephorus lyolepis</i> *.....	49528	<i>Malacoctenus culebrae</i>	49369
<i>Stolephorus gilberti</i>	49359	<i>Malacoctenus moorci</i>	49370
<i>Stolephorus garmani</i>	49360	<i>Malacoctenus puertoricensis</i>	49371
<i>Apogon sellicauda</i> *.....	49529	<i>Auchenopterus albicaudus</i>	49373
<i>Mycteroperca bowersi</i> *.....	49530	<i>Auchenopterus fajardo</i>	49376
<i>Prionodes baldwini</i>	49361	<i>Auchenopterus rubescens</i>	49374
<i>Neomænis megalophtalmus</i> *.....	49531	<i>Auchenopterus cingulatus</i>	49375
<i>Calamus kendalli</i>	49362	<i>Auchenistius</i>	
<i>Doratonotus decoris</i>	49363	<i>Auchenistius stahli</i>	49372
<i>Scorpaena albifimbria</i> *.....	49532	<i>Coralliozetus</i>	
<i>Scorpaena bergii</i> *.....	49533	<i>Coralliozetus cardonæ</i>	49377
<i>Pontinus beanorum</i> *.....	49534	<i>Emblemaria pandionis</i> *.....	49535
<i>Sicydium caguaitæ</i>	49364	<i>Citharichthys arenaceus</i> *.....	49536
<i>Gobius bayamonensis</i>	49365	<i>Haliutichthys smithii</i> *.....	49537

¹ Description of new genera and species of fishes from Porto Rico. <Report U. S. Fish Com. 1899 (December 19), 351-362.

² The Moringuoid Eels in American Waters, by Theo. Gill and H. M. Smith. <Science, N. S., vol. XI, No. 286, pp. 973-974, June 22, 1900.

The total number of fishes now known from Porto Rican waters is 291 species. Prior to the investigations by the U. S. Fish Commission the number was 99. Of the 291 species, 263 were obtained by us, and of this number 33 species proved to be new, which is more than 12.5 per cent of the species collected by us, or 11.3 per cent of the entire known fish fauna of the island.

The 291 species represent 76 families and 165 genera, 3 of the latter being new. The majority of the families and genera are represented by but few species each. Among the families with largest representation are the grunts (*Hæmulidae*), with 17 species; the sea basses and groupers (*Serranidae*), with 16 species; the pampanos (*Carangidae*) and parrot-fishes (*Scaridae*), with 15 species each; and the blennies (*Blenniidae*) and gobies (*Gobiidae*), with 14 species each. The snappers (*Lutjanidae*) are represented by 11 species. The largest genera are *Sparisoma*, with 11 species; *Neomænis*, with 10; *Hæmulon* and *Stolephorus*, with 8 each; and *Caranx*, *Scorpena*, and *Auchenopterus*, with 5 each. Several families and many genera are represented by a single species each.

The food-fishes of the island are numerous as to species, but only fairly abundant as to individuals. The absence of extensive shoals or banks about the island, the comparatively limited area of shallow water about the river mouths, and the fewness and small size of the bays, are evidence that there can not be suitable feeding-grounds such as could support large numbers of commercial fishes. The number of species of fishes used as food is, however, large, as is shown in the chapter on the commercial fisheries of the island.

A comparison of the fish fauna of Porto Rico with that of Cuba and other neighboring regions will prove interesting. Poey records 499 species from Cuba, 301 of which have not been taken in Porto Rico. Of the 291 species known from Porto Rico, 198 are common to it and Cuba, while 93 of them are not known from Cuba.

The number of species known from the Florida Keys is almost exactly the same as the number recorded from Porto Rico. With few exceptions the food-fishes found at Key West, Habana, and Porto Rico are identical, and the same is true for Jamaica, from which Jordan & Rutter give a list of 197 species. Of these, 45 were not obtained by us, and of our 291 species, 139 were not listed by them from Jamaica.

The species common to Key West, Habana, Jamaica, and Porto Rico, however, show that those four regions belong to one fauna—the West Indian. It is true that at Key West are found a number of species which are not yet known from the West Indies, and which probably do not occur among them, but they are mostly stragglers or the peripheral species of the faunas of the Gulf and the coast of the South Atlantic States.

Another remarkable feature of the Porto Rican fish fauna is the paucity in the representation of the cyprinodonts. The family *Peciliidae* is a large one, and in most of our tropical waters it is represented by many species. From Florida 21 species of this family are recognized, but only 2 are known from Porto Rico. This great difference is doubtless largely due to the marked difference in the character of the environment. The *Peciliidae* are chiefly fishes of brackish water and mud bottom, conditions adequately met in Florida, but almost entirely absent from Porto Rico.

KEY TO THE FAMILIES OF FISHES REPRESENTED IN WEST INDIAN WATERS.¹

I.—VENTRAL FINS PRESENT, ABDOMINAL.

- A. Back with an adipose fin behind the single-rayed dorsal fin.
 B. Head with 4 to 8 long barbels about the mouth and nostrils.....SILURIDÆ.
 BB. Head without barbels as described above.
 C. Sides of body without photophores or luminous glands; no barbel at throat.
 D. Pseudobranchiæ present.
 E. Maxillary very narrow, rudimentary, or obsolete; hypocoracoids not divergent.....SYNODONTIDÆ.
 EE. Maxillary well developed, dilated behind; pectoral normal; hypocoracoids mostly divergent.....AULOPIDÆ.
 DD. Pseudobranchiæ absent.
 F. Pectoral undivided, subhumeral; pseudobranchiæ absent.....BENTHOSAURIDÆ.
 FF. Pectoral rays elongate, arranged in two groups.....BATHYPTEROIDIDÆ.
 CC. Sides of body with photophores more or less developed. Barbel at throat present, very long; body naked.
 ASTRONESTHIDÆ.
 G. Vertebral spines projecting through skin of back before dorsal fin; body short and deep, greatly compressed.
 STERNOPTYCHIDÆ.
 GG. Vertebral spines not exerted in front of dorsal.
 H. Pseudobranchiæ present.
 I. Premaxillaries forming entire margin of upper jaw; body scaly; opercular apparatus complete.
 J. Form elongate, the snout pointed, barracuda-like; photophores very small.....PARALEPIDIDÆ.
 JJ. Form oblong, the snout not much produced; photophores conspicuous.....MYCTOPHIDÆ.
 II. Premaxillaries not forming the whole margin of upper jaw, the maxillary entering into it; body naked; opercular apparatus incomplete.....MAUROLOCIDÆ.
 HH. Pseudobranchiæ absent; mouth large, with canine teeth; scales deciduous or wanting.....CHAULIODONTIDÆ.
 AA. Back without adipose fin.
 B. Dorsal fin single, made up of rays, and not preceded by a series of free spines or followed by finlets.
 C. Tail evidently strongly heterocercal; scales ganoid; no gular plate; dorsal fin short.....LEPISOSTEIDÆ.
 CC. Tail not evidently heterocercal.
 D. Tail tapering to a point, without caudal fin; anal fin very long, of about 200 rays; body scaly.....HALOSAURIDÆ.
 DD. Tail not tapering to a point; caudal fin developed.
 E. Body naked.
 F. Throat with a long barbel; no caudal filament; mouth large.
 G. Barbel free at tip.....STOMIATIDÆ.
 GG. Barbel connecting throat with symphysis of lower jaw.....MALACOSTEIDÆ.
 FF. Throat without barbel.
 H. Caudal fin with a long filament; body elongate; mouth very small.....FISTULARIDÆ.
 HH. Caudal fin without filament. Pectorals present.....RONDELETHIDÆ.
 EE. Body scaly.
 I. Head with a large divided luminous plate in place of eyes.....IPNOPIDÆ.
 II. Head with normally developed eyes.
 J. Anal fin with many spines; mouth toothless, sucker-like.....LIPOGENYIDÆ.
 JJ. Anal fin without distinct spines.
 K. Pectoral fins inserted high, near axis of body; lower pharyngeals united; lateral line along side of belly.
 L. Jaws each with long sharp teeth mixed with smaller ones.....ESOCIDÆ.
 LL. Jaws with small equal, conic or tricuspid teeth.
 M. Lower jaw more or less produced; teeth tricuspid.....HEMIRAMPHIDÆ.
 MM. Lower jaw a little produced; teeth conic; pectoral elongate, forming an organ of flight.....EXOCTIDÆ.
 KK. Pectoral fins inserted below axis of body; lower pharyngeals separate.
 N. Phosphorescent spots present; teeth unequal.....CHAULIODONTIDÆ.
 NN. Phosphorescent spots none.
 O. Head scaly, more or less.
 P. Maxillaries connate with premaxillaries; jaws long.....SYNODONTIDÆ.
 PP. Maxillaries distinct; upper jaw protractile, its margin formed by premaxillaries alone; no lateral line.....PECTIDÆ.
 OO. Head naked.
 Q. Dorsal fin inserted more or less before anal (rarely slightly behind it); shore fishes or river fishes, usually silvery in coloration and with skeleton firm; air-bladder well developed.
 R. Gular plate present, between branches of lower jaw; mouth large; teeth present, all pointed; axillary scales and sheaths large.....ELOPIDÆ.
 RR. Gular plate none.
 S. Lateral line well developed; mouth small, horizontal; posterior part of tongue and roof of mouth covered with coarse-paved teeth.....ALBULIDÆ.
 SS. Lateral line wanting; no gular plate.
 T. Mouth moderate, terminal, the maxillary of about 3 pieces; stomach not gizzard-like.....CLUPEIDÆ.
 TT. Mouth subinferior, very large, below a tapering, pig-like snout; maxillary very long.....ENGRAULIDÆ.
 QQ. Dorsal fin posterior, opposite anal; deep-sea fishes, of loose organization; mostly blackish in color; mouth small, with small pointed teeth; air-bladder wanting.....ALEPOCEPHALIDÆ.

¹ The lancelets, sharks, and rays are not included in this key.

- BB. Dorsal fin single, preceded by free spines. Body scaly; snout tubular.....AULOSTOMIDÆ.
 BBB. Dorsal fin composed of free spines; ventrals with 1 or 2 spines each; body elongate.....NOTACANTHIDÆ.
 BBBB. Dorsal fins 2, the anterior of spines only, the posterior chiefly of soft rays.
 U. Pectoral fin with 5 to 8 lowermost rays detached and filamentous.....POLYNEMIDÆ.
 UU. Pectoral fin entire.
 V. Teeth strong, unequal; lateral line present.....SPHYRENIDÆ.
 VV. Teeth small or wanting; lateral line obsolete.
 W. Dorsal spines 4, stout; anal spines 3.....MUGILIDÆ.
 WW. Dorsal spines 4 to 8, slender; anal spine single.....ATHERINIDÆ.

II.—VENTRAL FINS PRESENT, THORACIC OR SUBJUGULAR, NUMBER OF RAYS DEFINITELY 1, 5.

- A. Gill-openings in front of the pectoral fins.
 B. Body more or less scaly or armed with bony plates.
 C. Ventral fins completely united; gill-membranes joined to the isthmus; no lateral line.....GOBIDÆ.
 CC. Ventral fins separate.
 D. Suborbital with a bony stay, which extends across the cheek to or toward the preopercle; cheeks sometimes entirely mailed.
 E. Pectoral fin with 3 lower rays detached and free; head bony.....TRIGLIDÆ.
 EE. Pectoral fin with 2 lower rays detached and free; body mailed.....PERISTEDIIDÆ.
 EEE. Pectoral fin entire; slit behind fourth gill small or wanting.
 F. Dorsal spines 4; lips fringed; eyes superior.....URANOSCOPIDÆ.
 FF. Dorsal spines 8 to 17; anal spines 3; body scaly.....SCORPENIDÆ.
 DD. Suborbital stay wanting; cheeks not mailed.
 G. Spinous dorsal transformed into a sucking disk on top of head, composed of 8 to 30 transverse plates.....ECHENEIDIDÆ.
 GG. Spinous dorsal (if present) not transformed into a sucking disk.
 H. Dorsal spines all or nearly all disconnected from each other.
 I. Body elongate, spindle-shaped.....RACHYCENTRIDÆ.
 II. Body oblong or ovate, compressed.
 J. Caudal peduncle very slender, the fin widely forked; preopercle entire.....CARANGIDÆ.
 JJ. Caudal peduncle stoutish, the fin little forked. Gill-membranes broadly united to isthmus.....EPHIPPIDÆ.
 HH. Dorsal spines (if present) all, or most of them, connected by membrane.
 K. Pectoral fin with 4 to 9 lowermost rays detached and filiform.....POLYNEMIDÆ.
 KK. Pectoral fin entire.
 L. Dorsal and anal each with 1 or more detached finlets.
 M. Anal preceded by 2 free spines.....CARANGIDÆ.
 MM. Anal not preceded by 2 free spines.
 N. Caudal peduncle keeled.....SCOMBRIDÆ.
 NN. Caudal peduncle not keeled.....GEMPYLIDÆ.
 LL. Dorsal and anal without finlets.
 O. Lateral line armed posteriorly with a series of keeled plates; 2 free anal spines; gill-membranes free from isthmus.
 CARANGIDÆ.
 OO. Lateral line armed posteriorly with a sharp, movable, lancet-like spine, or with a few bony tubercles; scales small, rough; gill-membranes adherent to isthmus.....TEUTHIDIDÆ.
 OOO. Lateral line unarmed.
 P. Throat with 2 long barbels (placed just behind chin); dorsal fins 2.....MULLIDÆ.
 PP. Throat without long barbels.
 Q. Anal fin preceded by 2 free spines (these obsolete in very old, joined by membrane in very young).....CARANGIDÆ.
 QQ. Anal fin not preceded by free spines.
 R. Nostril single on each side; lateral line interrupted; lower pharyngeals united.
 S. Anal spines 2.....POMACENTRIDÆ.
 SS. Anal spines 3 to 11. Fresh-water fishes.....CICHLIDÆ.
 RR. Nostril double on each side.
 T. Lateral line extending to tip of middle rays of caudal.
 U. Anal spines 3, the second strong.
 V. Dorsal fins 2, separate; body elongate.....CENTROPOMIDÆ.
 VV. Dorsal fin continuous.....HEMULIDÆ.
 UU. Anal spines 1 or 2, the second large or small.....SCLENIDÆ.
 TT. Lateral line not extending beyond base of caudal fin.
 W. Gills 3.5, the slit behind the last very small or wanting.
 X. Mouth not vertical, lips not fringed; dorsal fin continuous, spines 8 to 18; scales cycloid; lower pharyngeals united.
 Y. Teeth in each side of each jaw united, forming a sort of beak.....SCARIDÆ.
 YY. Teeth distinct or nearly so, the anterior usually more or less canine.....LABRIDÆ.
 XX. Mouth nearly vertical, the lips with fleshy fringes; dorsal divided, the spinous part short, of about 4 spines; lower pharyngeals separate.....URANOSCOPIDÆ.
 WW. Gills 4, a long slit behind the fourth.
 Z. Teeth setiform, like the teeth of a brush; body elevated, longer than deep, the soft fins completely sealed.
 a. Dorsal fin continuous.....CHETODONTIDÆ.
 aa. Dorsal fin divided.....EPHIPPIDÆ.

- ZZ. Teeth not setiform.
- b. Body deeper than long, covered with rough scales; dorsal spines 8; anal spines 3; soft fins very long.CAPROIDE.
- bb. Body longer than deep.
- c. Gill-membranes broadly joined to isthmus; body long and low; no lateral line.GOBIIDE.
- cc. Gill-membranes free from isthmus or very nearly so.
- d. Premaxillaries excessively protractile, their basal process very long, in a groove at top of cranium.
- e. Teeth small; scales large, silvery; spines strong.GERRIDE.
- cc. Teeth none; spines slender.MORNIIDE.
- dd. Premaxillaries moderately protractile or not protractile.
- f. Lateral line incomplete or interrupted, running close to dorsal fin; dorsal spines very slender, continuous with the soft rays; body low, covered with small scales; anal fin very long.OPISTHOGNATHIDE.
- ff. Lateral line, if present, not as above.
- g. Anal fin much longer than dorsal; body much compressed, the belly prominent.PEMPHERIDIDE.
- gg. Anal fin not much, if any, longer than dorsal.
- h. Pseudobranchiæ wanting or covered by skin; dorsal fin of soft rays, only beginning as a crest on the head; caudal widely forked. Pelagic fishes.CORYPHENIDE.
- hh. Pseudobranchiæ developed.
- i. Spinous dorsal of 2 or 3 short spines only; anal without spines; scales small, smooth.SERRANIDE.
- ii. Spinous dorsal, if present, not as above.
- j. Dorsal fin continuous, the spines few, slender; maxillary usually with an enlarged tooth behind; nape sometimes with an adipose appendage; anal fin long, even.MALACANTHIDE.
- jj. Dorsal fin continuous or divided, not as above.
- k. Perch-like fishes, the caudal peduncle not very slender, the scales well developed, ctenoid or cycloid; the dorsal with distinct spines; the anal with at least 1 spine, its soft rays usually few.
- l. Maxillary not sheathed by the preorbital, or only partially covered by the edge of the latter; ventral with its accessory scale very small or wanting; pectoral without accessory scale; sheath at base of spinous dorsal little developed; vomer usually with teeth; opercle usually ending in a spine.
- m. Precaudal vertebrae with transverse processes from the third or fourth to the last; ribs all but the last 1 to 4 sessile, inserted on the centra behind the transverse processes; anal spines 3; species silvery in color, the dorsal deeply notched, with 10 spines; vertebrae 10+15=25.KUHLEIDE.
- mm. Precaudal vertebrae normal, anteriorly without transverse processes; all or most of the ribs inserted on the transverse processes when these are developed.
- n. Anal spines 2, rarely 3; vertebrae 24 or 25; dorsal fin divided. Marine fishes.CHEILODIPTERIDE.
- nn. Anal spines 3, never 2 nor 1; dorsal fin continuous or divided; vertebrae 24 to 35.
- o. Vomer, and usually palatines also, with teeth.
- p. Anal fin shorter than dorsal; head not everywhere covered with rough scales; postocular part of head not shortened.SERRANIDE.
- p. Anal fin scarcely shorter than dorsal and similar to it; head and body everywhere covered with rough scales; body deep, compressed, the posterior part of head shortened.PRIACANTHIDE.
- oo. Vomer without teeth; dorsal fin continuous; body deep, compressed.LOBOTIDE.
- ll. Maxillary slipping for most of its length under the edge of the preorbital, which forms a more or less distinct sheath; ventrals with an accessory scale; opercle without spines; maxillary without supplemental bone; anal spines 3, rarely 2.
- q. Fishes carnivorous; intestine of moderate length; teeth in jaws not all incisor-like; vertebrae usually 24 or 25.
- r. Vomer with teeth, these sometimes very small; maxillary long.LUTIANIDE.
- rr. Vomer without teeth; palatines and tongue toothless.
- s. Teeth on sides of jaws not molar; maxillaries formed essentially as in the *Serranidae*; preopercle mostly serrate.HEMULIDE.
- ss. Teeth on sides of jaws molar; maxillaries peculiar in form and in articulation; anterior teeth conical or else more or less incisor-like, preopercle entire.SPARIDE.
- qq. Fishes herbivorous; intestinal canal elongate; anterior teeth in jaws incisor-like; no molars or canines; premaxillaries moderately protractile.KYPHOSIDE.
- kk. Mackerel-like fishes, with the caudal peduncle usually very slender, the fin widely forked, the scales various, usually not ctenoid; the dorsal spines various; anal fin long.
- t. Scales firm, linear, parchment-like; body compressed; bones of head rough; dorsal spines few; mouth small.GRAMMICOLEPIDIDE.
- tt. Scales not linear, mostly cycloid.
- u. Dorsal fin very long, all the rays soft; skeleton soft.ICOSTEIDE.
- uu. Dorsal fin with 3 or more spines.
- v. Dorsal fin divided, the spines 6 to 12 in number.
- w. Scales weak, cycloid; jaws without canines.NOMEIDE.
- vv. Scales firm, each with a median ridge; no canines.STEINERGERIDE.
- ww. Dorsal spines 3 or 4, the fin not divided.
- BB. Body scaleless, smooth or armed with tubercles, prickles, or scattered bony plates.
- C. Breast with a sucking disk; gill-membrane free from the isthmus; no spinous dorsal.GOBIESOCIDE.
- CC. Breast without sucking disk.
- D. Gill-membranes broadly attached to the isthmus.
- E. Ventrals completely united.GOBIIDE.
- EE. Ventrals widely separated; body depressed; preopercle with a strong spine.CALLIONYMIDE.
- DD. Gill-membranes nearly or quite free from the isthmus.

- F. Anal preceded by 2 free spines (these lost with age; connected by membranes in the very young)CARANGIDÆ.
 FF. Anal without free spines.
 G. Dorsal and anal fins followed by finletsSCOMBRIDÆ.
 GG. Dorsal and anal without finlets; mouth very large, nearly horizontal, the teeth sharp; no pseudobranchiæ.
 CHIASMODONTIDÆ.
 AA. Gill-openings small, behind, above, or below the pectoral fins, which are more or less pedicellate.
 II. Gill-openings in or behind upper axil of pectoral; mouth smallOGCOEPHALIDÆ.
 III. Gill-openings in or behind lower axil of pectoral; mouth large.
 I. Head compressed; no pseudobranchiæANTENNARIIDÆ.
 II. Head depressed; pseudobranchiæ presentLOPHIDÆ.

III.—VENTRAL FINS PRESENT, THORACIC OR JUGULAR, NUMBER OF RAYS NOT DEFINITELY 1, 5.

- A. Eyes unsymmetrical, both on the same side of head.
 B. Eyes large, well separated; edge of preopercle usually evidentPLEURONECTIDÆ.
 BB. Eyes small, very close together; edge of preopercle hidden by skin; mouth very smallSOLEIDÆ.
 AA. Eyes symmetrical, one on each side of the head.
 C. Ventral fin with or without spine, the number of soft rays more than 5.
 D. Caudal fin wanting; scales spinousMACROURIDÆ.
 DD. Caudal fin well developed.
 E. Ventral rays about 15; dorsal fin single, elevatedLAMPRIDÆ.
 EE. Ventral rays 1, 6 to 1, 10; dorsal with spines.
 F. Chin with two long barbels behind symphysis; dorsal continuous, with 5 spinesPOLYMIXIDÆ.
 FF. Chin without barbels.
 G. Dorsal fin divided, the anterior part of a single slender spine; ventrals elongateBREGMACEROTIDÆ.
 GG. Dorsal fin divided, the anterior part of many spines.
 H. Body covered with firm serrated scales; anal spines 4; dorsal spines not elevatedHOLOCENTRIDÆ.
 HH. Body naked or covered with small scales, besides bony plates or wartsZEIDÆ.
 GGG. Dorsal fin continuous, its spines 2 to 8BERYCIDÆ.
 CC. Ventral fin with or without spine, the number of soft rays fewer than 5.
 I. Gill-opening before the pectoral fin.
 J. Anal fin present; caudal fin not directed upward.
 K. Upper jaw not prolonged into a sword.
 L. Dorsal fin with some spines or simple rays.
 M. Dorsal fin without soft rays, composed of spines onlyBLENNIIDÆ.
 MM. Dorsal fin with soft rays anteriorly, with spines posteriorly; gill-membranes joined to isthmusZOARCIDÆ.
 MMM. Dorsal fin of spines anteriorly, with soft rays posteriorly.
 N. Suborbital with a bony stay, extending across cheek, to or toward preopercle, the cheek sometimes entirely covered with a coat of mailCEPHALACANTHIDÆ.
 NN. Suborbital without bony stay.
 O. Dorsal spines 2 to 4 only; head very broad, depressed; gills 3; gill-membranes broadly united to the isthmus.
 P. Ventrals each a strong spine; teeth incisor-like; scales shagreen-likeTRIACANTHIDÆ.
 PP. Ventrals not reduced each to a single spineBATRACHOIDIDÆ.
 OO. Dorsal spines numerous; gills 4.
 Q. Gill-membranes separate, free from the isthmus.
 R. Body greatly elongate; lower jaw with a slit at base to permit free motion; lips not fringed.
 S. Soft dorsal and anal with a distinct lobe anteriorly, distinct from spinous partGEMPYLIDÆ.
 SS. Soft dorsal and anal without anterior lobe, continuous with spinous partLEPIDOPODIDÆ.
 RR. Body moderately elongate; opercles and lips fringed; eyes superiorDACTYLOSCOPIDÆ.
 QQ. Gill-membranes broadly united, attached to the isthmus or notBLENNIIDÆ.
 LL. Dorsal fins of soft rays only.
 T. Breast with a large sucking disk between ventral finsGOBIESOCIDÆ.
 TT. Breast without sucking disk.
 U. Lateral line and base of dorsal beset with prickles; skeleton very soft; body compressedICOSTEIDÆ.
 UU. Lateral line unarmed.
 V. Tail isocercal, the vertebral column pointed behind, the last vertebra very small; hypereoraoid not perforate; no pseudobranchiæ.
 W. Caudal fin presentGADIDÆ.
 WW. Caudal fin wantingMACROURIDÆ.
 VV. Tail not isocercal, truncate at base of caudal; hypereoraoid perforate.
 X. Gill-membranes joined to the isthmus; pseudobranchiæ presentZOARCIDÆ.
 XX. Gill-membranes free from the isthmus.
 Y. Ventral fins inserted below or before the eyes; pseudobranchiæ generally well developedOPHIDIIDÆ.
 YY. Ventral fins inserted below shoulder-girdle; no pseudobranchiæBROTULIDÆ.
 KK. Upper jaw prolonged into a bony sword; dorsal fin long and high; size largeISTIOPHORIDÆ.
 JJ. Anal fin wanting; caudal fin distorted or directed upward; body ribbon-like; ventral fins each of a few slender raysTRACHYPTERIDÆ.
 II. Gill-opening behind the pectoral fin.
 Z. Gill-opening above and behind pectoral; mouth small, lowOGCOEPHALIDÆ.
 ZZ. Gill-opening below and behind pectoral; mouth large, nearly verticalANTENNARIIDÆ.

IV.—VENTRAL FINS WHOLLY WANTING.

- A. Premaxillary and maxillary wanting or grown fast to palatines; body greatly elongate, eel-shaped, gill-openings restricted to the sides; scales minute or wanting; scapular arch not attached to the skull. Eels
- B. Gill-openings well developed, leading to large interbranchial slits; tongue present; opercles and oranchial bones well developed; scapular arch present.
- C. Skin covered with rudimentary embedded scales, usually linear in form, arranged in small groups, and placed obliquely at right angles to those of the neighboring groups; pectorals and vertical fins well developed, the latter confluent about the tail; lateral line present; posterior nostril in front of eyes; tongue with its margins free ANGUILLIDÆ.
- CC. Scales wholly wanting; eggs (so far as known) of moderate size, much as in ordinary fishes.
- D. Tip of tail with a more or less distinct fin, dorsal and anal fins confluent around it; tail sometimes ending in a long filament; coloration almost always plain, brownish, blackish, or silvery, the fins often black-margined.
- E. Posterior nostril without tube, situated entirely above the upper lip.
- F. Tongue broad, largely free anteriorly and on sides; vomerine teeth moderate.
- G. Pectoral fins well developed; body not excessively elongate; lower jaw not projecting; anterior nostril remote from eye LEPTOCEPHALIDÆ.
- FF. Tongue narrow, adnate to the floor of the mouth or only the tip slightly free; vomerine teeth well developed, sometimes enlarged.
- II. Jaws not attenuate and recurved at tip; gill-openings well separated; anterior nostril remote from eye.
- I. Pectoral fin small or wanting MORINGUIDÆ.
- II. Pectoral fin well developed, skin thick; skeleton firm; snout moderate; tail not ending in a filiform tip. MURENESOCIDÆ.
- HH. Jaws long and slender, tapering to a point, recurved at tip; nostrils large, both pairs close in front of eye; gill-openings convergent forward, separate or confluent; pectorals and verticel fins well developed; membranes of fins thin, not enveloping the rays; skeleton well developed. Deep-sea eels NEMICHTHYIDÆ.
- EE. Posterior nostril close to the edge of the upper lip; tongue more or less fully adnate to the floor of the mouth; teeth subequal MYRIDÆ.
- DD. Tip of tail without rays, projecting beyond the dorsal and anal fins (not filiform); posterior nostril on the edge of the upper lip; anterior nostril near tip of snout, usually in a small tube; tongue usually adnate to the floor of the mouth; coloration frequently variegated. OPHICHTHYIDÆ.
- BB. Gill-openings small, roundish, leading to restricted interbranchial slits; tongue wanting; pectoral fins (typically) wanting; opercles feebly developed; fourth gill-arch modified, strengthened, and supporting pharyngeal jaws. MURENIDÆ.
- AA. Premaxillary and maxillary present, often immovably united to rest of cranium.
- J. Gill-openings united in a single slit below throat; no pectoral fins; body eel-shaped SYMBRANCHIDÆ.
- JJ. Gill-openings not united in a longitudinal slit.
- K. Body eel-shaped, ending in a long filament, longer than rest of body; no anal or caudal fin STYLEPHORIDÆ.
- KK. Body not truly eel-shaped.
- L. Gill-membranes broadly united to the isthmus, restricting the gill-openings to the sides.
- M. Snout tubular, bearing the short, toothless mouth at the end; body mailed SYNGNATHIDÆ.
- MM. Snout not tubular.
- N. Dorsal fin single, of spines or undivided rays only BLENNIDÆ.
- NN. Dorsal fins 2, the anterior of spines, the posterior of soft rays; body short and deep.
- O. Spinous dorsal of 2 or 3 spines; scales rather large, rough or bony BALISTIDÆ.
- OO. Spinous dorsal of 1 or 2 spines; scales minute, rough, forming a velvety covering MONACANTHIDÆ.
- NNN. Dorsal fin continuous, of soft rays only.
- P. Teeth in each jaw confluent into 1.
- Q. Body compressed, rough MOLIDÆ.
- QQ. Body not compressed, spinous DIODONTIDÆ.
- PP. Teeth in each jaw confluent into 2.
- R. Back broadly rounded TETRAODONTIDÆ.
- RR. Back with a sharp median ridge CANTHIGASTERIDÆ.
- PPP. Teeth separate; body enveloped in a bony box OSTRACIDÆ.
- LL. Gill-membranes free from the isthmus.
- S. Vent at the throat; vertical fins confluent; body elongate, almost eel-shaped FIERASFERIDÆ.
- SS. Vent posterior, not at the throat.
- T. Caudal fin wanting; body naked, greatly elongate TRICHIURIDÆ.
- TT. Caudal fin present.
- U. Upper jaw prolonged into a sword; size very large XIPHIIDÆ.
- UU. Upper jaw not prolonged into a sword.
- V. Belly with a series of bony scutes along its edge; body much compressed CLUPEIDÆ.
- VV. Belly not armed with scutes.
- W. Body ovate, much compressed.
- X. Scales small, cycloid, silvery STROMATEIDÆ.
- XX. Scales wanting; caudal peduncle very slender ICOSTEIDÆ.
- WW. Body oblong or elongate, much longer than deep; gill-membranes broadly united; teeth present; dorsal fin of spines only, or the posterior half of soft rays, the anterior of spines BLENNIDÆ.

DESCRIPTIVE LIST OF FISHES KNOWN FROM PORTO RICO.

Family I. BRANCHIOSTOMIDÆ. The Lancelets.

Body elongate, lanceolate, compressed, naked, colorless; fins represented by a low fold extending along back, with usually a rudimentary fold below, which passes by vent to the abdominal pore. Mouth inferior, appearing as a longitudinal fissure, surrounded by conspicuous, rather stiff, cirri. Eye rudimentary. Liver reduced to a blind sac of the simple intestine.

Small, translucent creatures found embedded in sand on warm coasts throughout the world. Eight species are now recognized, referable to two or three genera, all very similar in appearance and habits. The numbers of the muscular impressions furnish the only characters thus far known by which the species can be distinguished.

- a. Gonads (reproductive structures) present on both sides of the median line; anal fin present, with traces of fin-rays; no caudal process BRANCHIOSTOMA, 1
- aa. Gonads present on right side only; anal fin without fin-rays or successive fin-ray chambers; a long caudal process or tail, about as long as head..... ASYMMETRON, 2

Genus 1. BRANCHIOSTOMA Costa. Amphioxus.

Lancelets with the gonads present on both sides of median line. Anal fin present, with traces of rays. Vertebral column not produced backward into a caudal process.

Six or seven species recognized, found in warm seas, usually buried in sand flats at no great depth. Very tenacious of life and enduring considerable mutilation. Of the three American species, only one is known from Porto Rico.

- a. Myocommata or muscular bands 58 to 64.
- b. Myocommata behind vent 11 to 13, the formula usually $35+14+12=61$ lanceolatum
- bb. Myocommata behind vent 7 to 10, the formula usually $35+14+9=58$ caribæum, 1



FIG. 1.—*Branchiostoma caribæum*.

1. *Branchiostoma caribæum* Sundevall. West Indian Lancelet.

Muscular bands (myocommata) usually $35 + 14 + 9 = 58$; gonads 22 to 26 on each side; tail short; extremities attenuate. Usual length 1.75 inches.

In shallow waters, buried in the sand, from Beaufort, N. C., to the mouth of the La Plata; abundant off the Carolina coast and in localities in Florida (Port Tampa), Jamaica, Brazil, etc.

In the white coral sand at the east end of the island of Porto Rico, at a depth of 10 to 15 fathoms, this interesting little species was fairly abundant. About 40 specimens were obtained at stations 6084, 6086, 6087, 6093, and 6097, mostly at station 6084. The first four of these stations are between Vieques and Culebra islands and the last is off Hucars about 5 miles. The bottom at all was of live coral and white sand. One specimen (station 6086) was obtained in the dredge, while all the others were caught in the tangle. In many cases the lancelets were still alive when picked out from the meshes of the tangle, though they had been subjected to considerable rough treatment. The majority do not exceed an inch in length, and the largest is but 1.25 inches long. These are considerably smaller than the numerous individuals collected by the *Fish Hawk* off Port Tampa, Fla., in 1898.

Branchiostoma caribæum Sundevall, Öfers. Vet. Akad. Forhandl., 12, 1853, St. Thomas; Jordan & Evermann, Fishes of North and Middle America, 3, 1896.

Genus 2. ASYMMETRON Andrews.

Gonads, or reproductive structures, developed on the right side only. Anal fin without fin rays or successive fin-ray chambers. A long caudal process. Otherwise as in *Branchiostoma*. One species.

2. *Asymmetron lucayanum* Andrews. *Bahama Lancelet*.

Right metapleuron continuous with median ventral (anal) fin, which passes to right of anus. Preoral hood extensive, the cirri united by the membrane throughout greater part of their length, and smooth, without sensory papillæ. Gonads on the right, 29, extending from fifteenth to forty-third myotomes inclusive. Myotome formula $44+9+13=66$. Length $\frac{1}{2}$ inch. Adult and young swimming at surface in the evening in June and July at Bemini and Nassau, Bahamas; also taken buried in calcareous sand. (Andrews.) Three specimens, somewhat less than 0.75 inches in length, taken in the dredge and tangle at Fish Hawk stations 6086 and 6093, off Culebra, in 14.75 and 15 fathoms respectively, and 6097, off Humacao, in 10 fathoms. These are decidedly larger than specimens in the U. S. National Museum collected by Dr. Andrews.

Asymmetron lucayanum Andrews, Studies Biol. Lab. Johns Hopkins Univ., V, 237, 1893, Bemini, Bahamas; Jordan & Evermann, l. c., 4, 1896.

Family II. GINGLYMOSTOMIDÆ. The Nurse Sharks.

Large sharks with general characters of *Scylliorhinidæ*, but with tail very long and more or less abruptly bent upward at its base, as in the *Galeidæ*. First dorsal above or behind ventrals, the second opposite or rather before anal; eyes very small, with small spiracles behind them; nostrils confluent with mouth; nasal valves on both sides forming a quadrangular flap in front of mouth, each being provided with a free cylindrical cirrus; an upper and lower lip, the latter not extending across symphysis; fourth and fifth gill-openings close together.

Large sharks of the warm seas; genera 3, species about 5.

Genus 3. GINGLYMOSTOMA Müller & Henle.

Characters of the genus included above.




FIG. 2.—*Ginglymostoma cirratum*.

3. *Ginglymostoma cirratum* (Gmelin). *Nurse Shark*; "*Gata*."

Head obtuse, depressed; nasal cirrus reaching lower lip; angles of fins obtusely rounded; tail forming nearly one-third of total length; skin very thick; uniform brownish, the young with small, scattered, round black spots.

A large shark of the warmer parts of the Western Hemisphere, abundant about coral reefs in the West Indies and on the west coast of Mexico, and occasionally on our South Atlantic coast. Length 6 to 10 feet. Not seen by us in Porto Rico, but included on the authority of Professor Poey.

Squalus cirratus Gmelin, Syst. Nat., I, 1492, 1788, American Seas; after Broussonet.

Squalus punctatus Bloch & Schneider, Syst. Ichth., 134, 1801, Cuba; after *Gata Hispanas* of Parra.

Squalus punctulatus Bloch & Schneider, l. c., 549, 1801, Cayenne; after *Squale pointille* of Lacépède.

Squalus argus Baneroff, Zool. Jour., V, 82, 1832-1834, West Indies.

Ginglymostoma fulvum Poey, Memorias, II, 342, 1861, Havana; Poey, Fauna Puerto-Riqueña, 349, 1881; Stahl, Fauna de Puerto Rico, 81 and 167, 1883.

Ginglymostoma caboverdianus Capello, Jour. Sci. Phys. Lisbon, 1867, 167, Cape Verde.

Ginglymostoma cirratum, Jordan & Evermann, l. c., 26, 1896.

Family III. GALEIDÆ. The Requiem Sharks.

Sharks with two dorsal fins, first short and high, entirely before ventrals; second comparatively small, opposite anal; no spines; gill-openings moderate, the last above base of pectoral; tail more or less bent upward from base of caudal fin; sides of tail not keeled; eyes with nictitating membranes; head not hammer-shaped, the snout being longitudinally produced, as usual among sharks. Spiracles small or obsolete. Oviviviparous.

A large family of 20 or more genera and about 60 species; found in all seas. The species are often closely related and difficult of determination. Of 11 genera recognized by Jordan & Evermann as occurring in American waters, only one is as yet known from Porto Rico.

GALEINÆ:

- a. Teeth flat and paved, without cusps or ridges; spiracles present; no pit at root of tail; labial folds well developed.
 b. Embryo not attached to uterus by a placenta; teeth very blunt MUSTELUS
 aa. Teeth more or less compressed, with entire or serrate sharp edges.

GALEORHININÆ:

- c. Spiracles present.
 d. Root of tail with conspicuous pit above; teeth all coarsely serrate, alike in both jaws and each with a deep notch on outer margin; caudal fin with a double notch GALEOCERDO

CARCHARHININÆ:

- cc. Spiracles obsolete; lower teeth narrower than upper teeth.
 e. Angle of mouth without groove or with merely a slight depression, which does not extend along either jaw.
 f. First dorsal inserted anteriorly, nearer ventrals than pectorals; embryo (so far as known) attached to uterus by a placenta.
 g. Teeth all serrate more or less, often entire in the very young CARCHARHINUS, 4
 gg. Teeth of upper jaw serrate at base only; lower teeth entire, erect HYPOPRION
 ggg. Teeth all entire at all ages, and nearly all erect APRIONODON
 ee. Angle of mouth provided with a more or less distinct groove which extends along one or both jaws; teeth entire, or very nearly so, more or less obliquely placed, their points turned away from median line; embryo (so far as known) with a placenta SCOLIODON

Genus 4. CARCHARHINUS Blainville.

Body rather robust; head broad and depressed; mouth inferior, with teeth in both jaws strongly serrated in adult, less so or entire in young, those in upper jaw broad or narrow, those below narrow, straight, and nearly erect. No spiracles. First dorsal large, not far behind pectorals; pectoral falcate; second dorsal small. Embryos attached by placenta to uterus, as in *Scoliodon*, *Triakis*, and *Galeus*.

Voracious sharks of the warm seas. Species very numerous and difficult of separation.

- a. Teeth in both jaws distinctly serrate in adult; serræ on lower teeth smaller; upper teeth rather broad, lower teeth narrower; snout not very acute.

PLATYFODON:

- b. Upper teeth oblique, deeply notched on outer margin; lower teeth narrow, scarcely or not notched.
 c. Pectoral very large, 3 times as long as broad, falciform, extending beyond base of first dorsal; color blue-gray *obscurus*
 cc. Pectoral shorter, not 3 times as long as broad, extending little if any beyond base of first dorsal.
 d. Length of snout from mouth little if any greater than width of mouth.
 e. Distance from end of base of first dorsal to ventrals less than length of base of first dorsal.
 f. Nasal flap without sharp lobe; second dorsal and anal nearly equal; color blue-black *falciformis*, 4
 ff. Nasal flap with an acute lobe.
 g. Snout not very short, its length from the mouth not notably less than the width of mouth; color yellowish-brown *acronotus*
 gg. Snout very short and blunt, its length from mouth but two-thirds width of mouth *perci*
 ee. Distance from end of base of first dorsal to ventrals greater than length of first dorsal; snout moderate *remotus*
 dd. Length of snout from mouth greater than width of mouth; first dorsal small *heleni*

CARCHARHINUS:

- bb. Upper teeth triangular, suberect, scarcely notched on the outer margin; lower teeth similar but much narrower.
 h. Snout moderate, its length from mouth about equal to width of mouth *milberti*
 hh. Snout very short, its length from mouth less than width of mouth.
 i. Pectoral long and falcate, reaching to posterior part of base of dorsal. Anterior margin of first dorsal convex, height of fin about equal to depth of body *lania*
 ii. Pectoral fin moderate, scarcely falcate, not reaching to end of base of dorsal; second dorsal not larger than anal; length of snout from mouth 1.5 times in breadth of mouth; upper teeth very broad *platyodon*

ISOGOMPHODON:

- aa. Teeth slightly serrated, similar in form in the two jaws, narrow, claviform, constricted at base; snout rather sharp.
 j. Snout moderate, its length from mouth not greater than breadth of mouth; teeth moderate; fins edged with black *limbatus*, 5
 jj. Snout very long and narrow, its length from mouth twice distance between nostrils; teeth small, about 48 in each jaw *oxyrhynchus*

4. *Carcharhinus falciformis* (Bibron). "*Cazon de Playa.*"

Snout moderately prolonged and acute; nostrils without lobe; first dorsal rather backward; second dorsal and anal opposite each other and of medium size; pectoral not twice as long as broad; upper teeth with a marked reentrant angle on outer border; two pores of nape well marked.

Color, blue-black, deeper than in any other species.

Two specimens, taken on a hook off the Morro at San Juan January 13, measured as follows:

Measurements.	No. 0298.		No. 0299.	
	<i>Ft.</i>	<i>In.</i>	<i>Ft.</i>	<i>In.</i>
Total length.....	3	3	7	4
Length to base of caudal.....	2	4	5	6
Tip of snout to first gill opening.....	8		
Length of caudal fin from pit.....	11.5		2	1
Length of pectoral fin.....	6		1	4
Tip of snout to mouth.....	3.5		5.5	
Width of mouth.....	3		7.5	
Snout to origin of dorsal.....	1	1.5	2	6
Origin of dorsal to caudal pit.....	1	4	

No. 0299 possessed the following characters: Snout gently and narrowly rounded; first dorsal far in advance of ventrals, its height 9 inches, its base 11 inches; pectoral falcate, inner rays little produced; second dorsal and anal opposite, equal in size; ventrals small, claspers 10 inches long; lower lobe of caudal about one-third length of upper, which is somewhat falcate.

Color, olivaceous, paler below.

Though we were constantly on the lookout for sharks and often had lines out trying for them, this is the only species obtained by us in Porto Rico, and only these two specimens were seen. They are probably more abundant at other seasons.

Carcharias falciformis Bibron, in Müller & Henle's Plagiostomen, 47, 1838, Cuba.

Squalus tiburo Poey, Memorias, II, 331, 1861, Havana.

Carchacinas falciformis, Jordan & Evermann, l. c., 36, 1896.

5. *Carcharhinus limbatus* (Müller & Henle). "*Cuçonetta.*"

Snout somewhat pointed in front, rather produced, distance between its extremity and mouth somewhat less than width of mouth; nostrils nearly midway between extremity of snout and mouth; teeth $\frac{25}{27}$ - $\frac{29}{30}$, similar in form in both jaws, erect, constricted, on a broad base, upper more distinctly serrated than lower; gill-openings wide, at least twice as wide as the small eye. Pectorals falciform, extending beyond end of dorsal, length of their upper margin nearly four times that of lower. First dorsal commencing very close behind axil of pectoral; origins of second dorsal and anal opposite each other, the bases being nearly equally long. Caudal fin long, its length equal to distance between origins of the two dorsal fins.

Color, gray, lower side of extremity of pectoral, extremities of second dorsal and anal, and of lower caudal lobe, black. (Günther).

Found in tropical seas, north to Florida; numerous specimens once taken at Woods Hole, Mass.; common in Brazil; used as food by the very poor. Not seen by us in Porto Rico, but recorded from there by Doctor Stahl.

Carcharias (Prionodon) limbatus Müller & Henle, Plagiostomen, 49, 1838, Martinique.

Isogomphodon maculipinnis Poey, Repertorio, I, 191, pl. 4, figs. 2 and 3, 1867, Cuba.

Carcharias unilcei Steindachner, Sitzb. Akad. Wiss. Wien 1867, 356, West Indies.

Carcharias microps Lowe, Proc. Zool. Soc. 1840, 38, Madeira.

Prionodon cucuzi Castlenau, Anim. Nou. Rares. Amer. Sud, Poiss., 99, 1855, Bahia.

Platypodon ? Poey, Fauna Puerto-Riqueña, 348, 1881.

Platypodon maculipinnis, Stahl, l. c., 81 and 167, 1883.

Carcharhinus limbatus, Jordan & Evermann, l. c., 40, 1896.

Family IV. SPHYRNIDÆ. The Hammer-head Sharks.

General characteristics of the *Galeidæ*, but with the head singularly formed, kidney-shaped, or hammer-shaped, from extension of its sides, the nostrils being anterior and the eyes on the sides of the hammer; mouth crescent-shaped, under hammer; teeth of both jaws similar, oblique, each with a notch on the outside near base; no spiracles; last gill-opening over pectoral; first dorsal and pectorals large, the dorsal nearer pectorals than ventrals; second dorsal and anal small; a pit at root of caudal; caudal fin with a single notch toward its tip, its lower lobe developed.

One genus, with 5 species, inhabiting most warm seas. Large sharks, known at once by the singular form of the head, which is not quite the same in any two species.

Genus 5. SPHYRNA Rafinesque.

Characters of the genus included above. In form of head there is a perfect gradation among the species from a narrow hammer, with the lobes three times as long as broad and deeply grooved along the anterior edge, to kidney shape, in which the anterior grooves are obsolete. Only one species of this genus is known from Porto Rico.

- a. Nostrils near eyes.
- b. Nostril with frontal groove short or obsolete; lateral extension of head moderate, so that the head is rather kidney-shaped than hammer-shaped.

RENICEPS:

c. Nostril with groove obsolete; anterior and lateral margins of head confluent into a semicircle..... *tiburø*

PLATYSQUALUS:

cc. Nostril with a short groove; anterior margin of head curved, not continuous with lateral edges..... *tudes*

SPHYRNA:

bb. Nostril with a well-developed groove, which extends along front of the hammer-shaped head, the anterior and posterior outlines of which are nearly parallel..... *zyggæna*, 6

6. *Sphyrna zygæna* (Linnaeus). Hammer-head Shark; "Cornuda."

Head truly hammer-shaped; width of head about twice its length; length of hinder margin of hammer nearly equal to its width near eye, prolonged into a groove which runs along nearly the whole front margin of head; first dorsal large, second quite small, smaller than anal; pectoral rather large. Color, gray. A large, voracious shark, reaching a length of 15 feet or more; found in all warm seas; occasional on our coasts from Cape Cod and from Point Concepcion southward. Not obtained by us, but included on the authority of Professor Poey.

Squalus zygæna Linnaeus, Syst. Nat., ed. X, 234, 1758, Europe; America.
Cestracion zygæna, Poey, Fauna Puerto-Riqueña, 348, 1881; Stahl, Fauna de Puerto Rico, 81 and 167, 1883.
Sphyrna zygæna, Jordan & Evermann, l. c., 45, 1896.



FIG. 3.—*Pristis pectinatus*.

Family V. PRISTIDÆ. The Saw-fishes.

Body elongate, depressed; pectoral fin moderate, front margin quite free, not extending to head; snout produced into a very long, thin, flat blade, armed with series of strong tooth-like processes placed in sockets along each edge; teeth in jaws minute, obtuse; gill-openings moderate, inferior; spiracles wide, behind eye; nostrils inferior, no tentacles; no nictitating membrane; dorsal fins large, without spine, the first nearly opposite ventrals. Caudal well developed, bent upward, a fold along each side of tail.

A single genus, with five or more species, inhabiting warm seas on sandy shores, sometimes ascending rivers.

Genus 6. PRISTIS Latham.

Characters of genus included above.

- a. Rostral teeth in 18 to 20 pairs; first dorsal chiefly before ventrals; caudal with a small lower lobe..... *perrotteii*
- aa. Rostral teeth in 24 to 32 pairs; first dorsal opposite ventrals; caudal without lower lobe..... *pectinatus*, 7

7. *Pristis pectinatus* Latham. *Common Saw-fish; "Pez Sierra."*

First dorsal over ventrals; second dorsal scarcely smaller than first; no lower caudal lobe. Saw with 24 to 32 pairs of teeth, the posterior farther apart than the anterior. Found in the tropical seas north to West Indies and Florida; abundant in the Gulf of Mexico, ascending the Lower Mississippi. The species probably reaches a length of 20 feet. Not obtained by us, but included on the authority of Professor Poey.

The saw-fish is abundant in Indian River, Florida, where it is permanently resident. It is held in much dread by the commercial fishermen on account of the damage which it does by becoming entangled in their nets. The larger ones tear or cut the nets so seriously as to render them valueless, while the smaller ones become entangled and can only be removed with great difficulty. Those seen in Indian River usually do not exceed 3 feet in length, the saw included, but very large ones are occasionally taken. It is said that one caught near Eau Gallie in October, 1895, was 12.5 feet long and weighed 425 pounds. The largest reported by fishermen were 16 or 17 feet long.

Some interesting information concerning the young of the saw-fish is recorded. Mr. F. B. Everett says that from a large female saw-fish he took a number of young which swam away when placed in the water. The "saws" were enveloped in a membrane which soon disappeared in specimens left to dry in the sun, and the teeth became visible. Mr. Stypmann, of Stuart, Fla., took 18 or 20 young from a saw-fish about the 1st of July. The "saws" were well developed, but they, like the teeth, were soft like leather. There is some variation in the number of teeth on the saw, and there is usually one more tooth on one side than on the other. From a large number counted it appears that the usual numbers are 25 and 26, respectively.

Another species of saw-fish (*Pristis perroteti* Valenciennes) was described in 1838 from the Senegal River, on the west coast of Africa. This is said to occur in the West Indies, and the saw-fish from the Pacific coast of America north to Mazatlan has been identified with it. The Pacific coast species, however, is now regarded as distinct, and has been named *Pristis zephyreus* Jordan & Starks. No direct comparison has been made between specimens from the west coast of Africa and the West Indies, and it may be that there is but the one species in the West Indies.

- Pristis pectinatus* Latham, Trans. Linn. Soc., II, 278, 1794, "in the ocean"; Poey, Fauna Puerto-Riqueña, 349, 1881; Stahl, l. c., 81 and 167, 1881; Jordan & Evermann, l. c., 60, 1896.
Pristis granulosa Bloch & Schneider, Syst. Ichth., 352, 1801, Havana; after Parra.
Pristis mississippiensis Rafinesque, Ichth. Ohl., 80, 1820, Lower Mississippi River
Pristis megalodon Duméril, Elasmobranches, 476, pl. 9, fig. 4, 1870, Cayenne.
Pristis acutirostris Duméril, l. c., 479, 1870, Martinique.

Family VI. DASYATIDÆ. The Sting Rays.

Disk usually more or less broad than long; pectoral fins uninterruptedly confluent in front, forming tip of snout; tail variously formed, usually whip-like, sometimes short and stout, sometimes bearing a single dorsal or caudal fin, but never with two dorsals; usually one or more vertical folds of skin on the tail, rarely a lateral fold. Tail generally armed with a large, sharp, retrorsely serrate spine on its upper surface toward the base; 2 or 3 spines occasionally present. Ventral fins not emarginate. Skin smooth or variously prickly or spinous, roughest in adult; no differentiated spines on pectorals in males, the sexes being similar. Mouth rather small; teeth small, paved, usually more or less pointed or tubercular. Nostrils close together; nasal valves forming a rectangular flap, which is joined to upper jaw by a narrow frenum. Spiracles large, placed close behind eyes. Skull not elevated, spiracles and eyes superior. Ovoviviparous.

Found in most warm seas, some of them in the fresh waters of the northern parts of South America. Genera about 10; species 50. The large jagged spine on the muscular tail is capable of inflicting a severe and even dangerous wound. Only two species of this family are as yet known from Porto Rico, though doubtless others will be found.

- a. Tail stout, provided with a rayed caudal fin; no dorsal fin; disk roundish; caudal spine strong UROLOPHUS
aa. Tail slender, without caudal fin; pelvis without sword-shaped process. (Marine species.)

DASYATINÆ:

- b. Tail whip-like, longer than disk, which is rhomboid or roundish; caudal spine strong..... DASYATIS, 7
bb. Tail very short, shorter than the very broad, transversely rhombic disk; caudal spine weak, often wanting. No trace of dorsal fin..... PTEROPLATEA

Genus 7. *DASYATIS* Rafinesque.

Disk oval, flat, with rounded angles. Tail very long and slender, whip-like, without fin, but often with 1 or 2 vertical membranous folds; a strong, serrated spine toward base of tail. Skin more or less spinous or prickly, rarely smooth. Teeth small, paved; a few papillæ usually present in mouth behind lower jaw.

This genus contains about 30 species; sting rays of large size abundant in warm seas. Many of the spinous species are nearly or quite smooth when young, becoming rough with age. Some of our species are yet imperfectly known and much of the synonymy is uncertain.

HEMISTRYGON:

- a. Tail with a keel or wing-like expansion below only; adult with stout bucklers on back and tail; tail rough, more than twice length of disk..... *centrura*

DASYATIS:

- aa. Tail with a narrow keel or expansion both above and below.
 b. Tail simply keeled above, with a wing-like expansion below.
 c. Shoulder with 3 series of tubercles; tail less than twice length of disk..... *hastata*, 8
 cc. Shoulder with fewer than 3 series of tubercles.
 d. Tail round, more than twice disk; back with strong tubercles; snout sharp..... *gymnura*
 bb. Tail more or less compressed, with a wing-like expansion above, a larger one below.
 e. Skin more or less prickly in adult, with a median series of tubercles on back..... *sabina*
 ee. Skin nearly or quite smooth in adult; median line of back not prickly or with but one spine..... *say*, 9

8. *Dasyatis hastata* (DeKay). *Sting Ray*; "*Raya*."

Disk quadrangular, about one-fourth wider than long; anterior margins nearly straight, meeting in a blunt angle on end of snout, curved near outer angle to meet slightly convex posterior margins; inner border convex; outer and hinder angles rounded. Ventrals almost entirely covered by pectorals, their hinder margins convex. Tail more than one and one-half times length of disk, low-keeled on upper side, a long, broad, membranous expansion below, roughened with small asperities; one or more serrated spines. Body smooth in young; very old with scattered, small asperities; a row of narrow compressed tubercles along middle of back and on base of tail; points of tubercles depressed and directed backward. On each shoulder, parallel with median row, is a shorter row, its length varying according to age. Mouth with 3 papillæ. Jaws with more curvature than those of *D. centrura* and less than those of *D. sabina*; young without tubercles. Color, bluish or olivaceous-brown, pale below.

West Indies to Brazil; north to Florida. One female obtained at Hucaree, whose length (of disk) is 11.25 inches; tail, 15.5 inches; width of disk, 14.5 inches; a row of 12 spines along middle line of back and a shorter lateral row of 2 or 3 spines on each side.

Trygon hastata DeKay, New York Fauna: Fishes, 373, pl. 65, fig. 214, 1842, Rhode Island.

Dasyatis hastata, Jordan & Evermann, l. c., 83, 1896.

9. *Dasyatis say* (Le Sueur). *Southern Sting Ray*; "*Raya*."

Disk quadrangular, one-sixth wider than long, anterior margins nearly straight, posterior and inner borders convex, outer and posterior angles rounded. Snout not protruding beyond lines of margins. Ventrals rounded. Tail strong, rather more than one and one-half times length of disk, with a strong serrated spine, bearing a short, low, cutaneous expansion behind spine on upper side, and a longer, little wider one below, ending nearly opposite. Upper jaw undulated, lower prominent in middle. Teeth small, smooth in young and females, sharp in adult males; 3 papillæ at bottom of mouth, and 1 at each side. Body and tail smooth.

Color, olive-brown in adult, reddish or yellowish in young; lower surface whitish.

This species closely resembles the European *D. pastinaca*. In *D. say* the anterior margins form a more blunt angle at end of snout, which is less prominent at apex; outer and posterior extremities of pectoral rounder, posterior margin more convex, disk broader toward ventrals, and tail longer; in *D. pastinaca* the lateral and hinder angles of pectoral and lateral angle of ventral are marked by blunted corners; a single small, rounded tubercle on middle of back.

This ray is found from Carolina to Brazil. It is quite common in Florida, and is occasionally taken as far northward as New York. Not obtained by us, but given on authority of Poey.

Raja say Le Sueur, Jour. Ac. Nat. Sci. Phila., I, 1817, 42, New Jersey.

? *Trygon sayi*, Poey, Fauna Puerto-Riqueña, 350, 1881; Stahl, l. c., 81 and 167, 1883.

Dasyatis say, Jordan & Evermann, l. c., 86, 1896.

Family VII. MYLIOBATIDÆ. The Eagle Rays.

Disk broad; pectoral fins not continued to end of snout, but ceasing on sides of head and reappearing in front of snout as one or two fleshy protuberances (cephalic fins), which are supported by fin rays. Tail very long and slender, whip-like, with a single dorsal fin near its root, behind which is usually a strong retrorsely serrated spine. Nasal valves forming a rectangular flap, with posterior margin free, attached by a frenum to upper jaw. Skull less depressed than usual among rays, its surface raised so that the eyes and spiracles are lateral in position. Teeth hexangular, large, flat, tessellated, middle ones usually broader than the others. Skin smooth; no differentiated spines on pectorals in males, the sexes being similar. Ventrals not emarginate. Ovoviviparous.

Only a single species of this family is yet known from Porto Rico.

AETOBATINÆ:

a. Teeth in a single series, very broad; muzzle entire..... AETOBATUS, 8

RHINOPTERINÆ:

aa. Teeth in several series, the middle series very broad.

b. Muzzle entire..... MYLIOBATUS

bb. Muzzle emarginate; cephalic fins below level of disk..... RHINOPTERA




FIG. 4.—*Actobatus narinari*, dorsal view.

Genus 8. AETOBATUS Blainville.

General form of *Myliobatis*. Muzzle entire. Teeth flat, broad, forming a single series corresponding to the middle series in *Myliobatis*, there being no small lateral teeth. Upper dental lamina straight, lower curved and projecting beyond upper. Free border of nasal valve deeply emarginate. Skin smooth. Found in tropical seas.

10. *Aetobatus narinari* (Euphrasen). *Spotted Sting Ray*; "Obispo"; "Chucho."

Disk nearly or quite twice as broad as long; tail very long, about 2.5 times length of disk; snout 7 in length of disk; distance from snout to eye 10 in width of disk; width of mouth 10 in length of disk. A long furrow in middle of interorbital space, deepest in front; spiracles obliquely placed.

General color of whole upper surface light chocolate-brown, everywhere covered with roundish or oblong pearly or bluish spots or blotches, largest about size of eye, smallest less than half as large; under surface milky-white except margin of snout, which is dark-gray; tail uniform chocolate-brown; iris yellowish-gray.




FIG. 5.—*Aetobatus narinari*, ventral view.

Found in tropical seas; north on our Atlantic coast to Virginia. One male seined at Culebra Island February 9. This is one of the largest and most interesting of the rays. It is not uncommon on our Florida coast, though the published records do not so indicate. It attains a length of 2 feet or more exclusive of the tail, which is two to three times the length of the disk.

Raja narinari Euphrasen, Vet. Ak. Nya. Handl., XI, 1790, 217, Brazil; after *Narinari* of Maregrave.

Myliobatis celtenke Rüppell, Neu. Wirb., 70, 1835, Red Sea.

Goniobatis macroptera McClelland, Calcutta Jour. Nat. Hist., I, 1841, 60, Bengal.

Actobatus narinari, Poey, Fauna Puerto-Riqueña, 349, 1881; Stahl, l. c., 81 and 167, 1883; Jordan & Evermann, l. c., 88, 1896.

Family VIII. ANGUILLIDÆ. The Truc Eels.

The true eels are characterized by their scaly skin in association with a conical head and a general resemblance to the congers.

Genus 9. ANGUILLA Shaw.

Body elongate, compressed behind, covered with embedded linear scales placed obliquely, some at right angles to others. Lateral line well developed. Head long, conical, moderately pointed, the rather small eye well forward and over angle of mouth. Teeth small, subequal, in bands on each jaw and a long patch on vomer. Tongue free at tip. Lips rather full, with free margin behind, attached by a frenum in front. Lower jaw projecting. Gill-openings rather small, slit-like, about as wide as base of pectorals and partly below them. Nostrils superior, well separated, the anterior with a slight tube. Vent close in front of anal. Dorsal inserted at some distance from head, confluent with anal around tail. Pectoral well developed.

Found in most warm seas (the eastern Pacific excepted), ascending streams, but mostly spawning in the sea. They often move for a considerable distance on land, in damp grass, and in this way pass waterfalls, dams, and other obstructions. It is thought that they spawn only in the sea, the female dying after having once produced ova. The females are larger than the males, paler in color, with smaller eyes and higher fins. Eels are among the most voracious of fishes. "On their hunting excursions they overturn alike huge and small stones, beneath which they find species of shrimp and crawfish, of which they are excessively fond. Their noses are poked into every imaginable hole in their search for food, to the terror of innumerable small fishes."

11. *Anguilla chrysypa* Rafinesque. "*Anguilla*"; *Common Eel*.

(PLATE I.)

Head 3.2 in trunk; depth 5.8; eye 1.8 in snout; snout 5.5 in head; interorbital 5.4.

Body elongate, compressed behind, covered with linear embedded scales obliquely placed in groups, the scales of which are at right angles to those of adjoining groups; lateral line well developed; head long, conic, pointed; eye situated over angle of mouth; lower jaw projecting; fine conic teeth in bands on each jaw and on vomer; the gill-openings are vertical slits in front of and partly below pectoral; dorsal inserted far back of head, but considerably in front of vent; anal commencing shortly behind vent, confluent with dorsal around the tail, forming a caudal fin.

Atlantic coast of the United States, and Mississippi Valley; abundant from Maine to Mexico, ascending all rivers south of Canada and east of the Rocky Mountains. Common in the West Indies and highly valued as a food-fish. It is caught in considerable numbers in Porto Rico in the small bamboo traps or "nasas" set in the small rivers.

Six examples, the smallest 4 inches long, the largest over 2 feet, from San Juan market and Bayamon River at Bayamon, the smallest one from the sea at Mayaguez.

Anguilla chrysypa Rafinesque, Amer. Month. Mag. and Crit. Rev. 1817, 120, Lake George; Hudson River; Lake Champlain; Jordan & Evermann, l. c., 348, 1896.

Anguilla blephura Rafinesque, Amer. Month. Mag. and Crit. Rev. 1817, 120, Long Island.

Anguilla laticauda Rafinesque, Amer. Month. Mag. and Crit. Rev. 1817, 445, Ohio River.

Anguilla aterrima Rafinesque, Ieh. Ohiensis, 78, 1820, Tennessee and Cumberland rivers.

Anguilla xanthomelas Rafinesque, Ieh. Ohiensis, 78, 1820, Ohio River.

Anguilla lutea Rafinesque, Ieh. Ohiensis, 78, 1820, Ohio River.

Muræna rostrata Le Sueur, Jour. Ac. Nat. Sci. Phila. 1821, 81, Cayuga Lake, New York.

Muræna bostoniensis Le Sueur, Jour. Ac. Nat. Sci. Phila. 1821, 81, Boston.

Muræna serpentina Le Sueur, Jour. Ac. Nat. Sci. Phila. 1821, 82, Newport, R. I.

Muræna macrocephala Le Sueur, Jour. Ac. Nat. Sci. Phila. 1821, 82, Saratoga, N. Y.

Muræna argentea Le Sueur, Jour. Ac. Nat. Sci. Phila. 1821, 82, Boston Bay.

Anguilla tenuirostris De Kay, Fishes New York, 310, 1842, New York.

Anguilla novaeortensis Kaup, Apodes, 43, fig. 33, 1856, New Orleans, La.

Anguilla punctatissima Kaup, Apodes, 41, 1856, Niagara River.

Anguilla cubana Kaup, Apodes, 44, 1856, Cuba.

Anguilla novaterrore Kaup, Apodes, 45, fig. 35, 1856, Newfoundland.

Anguilla texana Kaup, Apodes, 45, fig. 36, 1856, Texas.

Anguilla wabashensis Kaup, Apodes, 46, 1856, Wabash River.

Anguilla tyrannus Girard, U. S. and Mexican Bound. Surv., 75, 1859, Rio Grande.

Muræna anguilla, Poey, Fauna Puerto-Riqueña, 344, 1881.

Muræna cubana, Stahl, l. c., 166, 1883.

Family IX. LEPTOCEPHALIDÆ. The Conger Eels.

This family includes those eels which are scaleless, and have the tongue largely free in front; body moderately elongate, end of tail surrounded by a fin; posterior nostril remote from upper lip and near front of eye, and pectoral fins well developed. All the species are plainly colored, grayish or dusky above, silvery below, and the dorsal edged with black.

Genera 3, species about 15; in most warm seas, usually at moderate depths. Most of the species undergo a metamorphosis, the young being loosely organized and transparent, band-shaped, and with very small head. The body grows smaller with increased age, owing to compacting of the tissues.

- a. Vomerine teeth in bands, none of them canine-like; lips thick.
- b. Dorsal fin inserted at a point behind base of pectoral, but nearer pectoral than vent; head with inconspicuous mucous cavities; jaws with an outer series of close-set teeth, forming a cutting edge; tail about half longer than rest of body..... LEPTOCEPHALUS, 10
- bb. Dorsal fin beginning over the gill-opening; bones of front of head with large muciferous cavities; mouth rather small; jaws with bands of small teeth, the outer not forming a cutting edge; tail from half to two-thirds of total length..... CONGERMURENA
- aa. Vomerine teeth uniserial, some of them canine-like; maxillary teeth biserial; dorsal beginning above root of pectoral; cleft of mouth extending beyond middle of eye; tail very long and slender, about half longer than rest of body..... UROCONGER




FIG. 6.—*Leptocephalus conger*.

Genus 10. LEPTOCEPHALUS (Gronow) Scopoli.

Body formed as in *Anguilla*, skin scaleless. Head depressed above, anteriorly pointed. Lateral line present. Mouth wide, its cleft extending at least to below middle of eye. Teeth in outer series in each jaw equal and close-set, forming a cutting edge; no canines; band of vomerine teeth short. Tongue anteriorly free. Vertical fins well developed, confluent around tail; pectoral fins well developed; dorsal beginning close behind pectorals. Gill-openings rather large, low. Eyes well developed; posterior nostril near eye, the anterior near tip of snout, with short tube. Lower jaw not projecting. Skeleton differing in numerous respects from that of *Anguilla*. Vertebrae about 56+100.

This genus is found in most warm seas, and contains the well-known and widely distributed conger eel and three or four closely related species. The earliest generic name used for members of the group is *Leptocephalus*, based on a curious, elongate, transparent, band-like creature, with minute head and very small mouth, found in the waters of Europe, and known as *Leptocephalus morrissi*. This has been shown by Gill, Günther, and Faccioli to be the young and larval form of *Leptocephalus conger*. A number of the genera and species of the supposed family of *Leptocephalidæ* have been described, but there is no doubt that all of them are larvæ, some of eels, as *Conger*, *Congermurena*, *Oplisurus*, and *Nettastoma*, others of Isospondylous fishes, as *Albula*, *Elops*, *Alepocephalus*, *Stomias*, etc. (See Günther, Cat., VII, 136.) It is thought by Dr. Günther that the Leptocephalid forms are probably "individuals arrested in the development at a very early period of their life, yet continuing to grow to a certain size, without corresponding development of their internal organs, and perishing without having attained the characters of the perfect animal." The recent observations of Dr. Gilbert on the larvæ of *Albula*, *Elops*, and *Conger*, however, seem to point to the conclusion that these curious forms are normal young, and that the individuals grow smaller in size for a time with increased age, owing to the increasing compactness of the tissues.

- a. Dorsal beginning nearly opposite tip of pectoral; head about 1.8 in trunk..... *conger*, 12
- aa. Dorsal fin beginning above middle of pectoral; head about 1.6 in trunk..... *caudilimbatus*

12. *Leptocephalus conger* (Linnaeus). *Conger Eel; Congrio.*

Dorsal beginning opposite or just behind tip of pectoral; eye 1.5 in snout, 5 to 6 in head; snout 3.25 to 4.25 in head; gape extending nearly or quite to posterior margin of eye; head 1.8 to 1.86 in trunk; tail longer than rest of body; pectoral 3.5 in head; upper lip full, with conspicuous pores.

Ashy-gray or blackish; vertical fins with black margin; body sometimes (var. *niger*) entirely black.

Atlantic Ocean, generally common on both coasts, from Cape Cod to Brazil; also on coasts of Asia and Africa; almost cosmopolitan, but not found in the eastern Pacific. It reaches a length of 8 feet, and is an important food-fish, especially in Europe. Recorded by Dr. Stahl; not obtained by us in Porto Rico.

(a) LARVAL FORMS.

- Leptocephalus morrissi* Gmelin, Syst. Nat., 1150, 1788, Holyhead, England.
Ophidium pectucidum Couch, Lond. Mag. Nat. Hist., V, 1832, 313, 742, England.
Leptocephalus gracilis Storer, Mem. Amer. Acad., II, 524, 1839, Massachusetts.
Leptocephalus spatanzanii, candidissimus, etc., of European writers.

(b) ADULT FORMS.

- Muraena suprema margine pinnae dorsalis nigro* Artedi, Synon., 40, 2, 1738, Mediterranean.
Muraena conger Linnaeus, Syst. Nat., X, 245, 1758, Mediterranean Sea; based on Artedi.
Muraena nigra Risso, Ich. Nice, 93, 1810 (black variety), Nice.
Anguilla oceanica Mitchell, Jour. Ac. Nat. Sci. Phila. 1818, 407, off New York.
Conger verus Risso, Eur. MÉR., III, 201, 1826, Nice.
Conger vulgaris Cuvier, Règne Animal, ed. II, vol. 2, 350, 1829, France; Günther, Cat., VIII, 38, 1870.
Conger rubescens Ranzani, De Novis Spec. Pisc. Diss. Prima, 19, pl. V, fig. 5, 1838, Mediterranean.
Ophisoma obtusa Swainson, Fish., Rep., and Amph., II, 395, 1839, Sicily.
Conger orbignyanus Valenciennes, D'Orbigny, Voy. Am. MÉRID., Poiss., pl. 12, 2, 1839, South America.
Conger esulentus Poey, Memorias, II, 346, 1860, Cuba; Stahl, l. c., 80, 1883 (misprinted "Conges").
Leptocephalus conger, Jordan & Evermann, l. c., 354, 1896.

Family X. MURÆNESOCIDÆ.

Scaleless anguilloid eels, with posterior nostril not labial, tongue largely adnate, jaws not excessively elongate, end of tail surrounded by caudal fin, and pectoral fins well developed. None of these characters appears to have in itself great importance, but, according to Dr. Gill, in the genus *Muraenesox*, the only genus in which the osteology is well known, the characters are such as fully to justify family distinction.

Dr. Gill gives the following diagnosis of the *Muraenesocidæ*: "Enchelycephalous Apodals with the tongue not free, the branchiostegal membrane connecting the opposite sides below, the epipharyngeals reduced to one pair, and the hypopharyngeals linguiform and encroaching on the fourth branchial arch." To this should be added: Gill-openings rather wide; pectoral fins well developed; jaws of moderate length; vomer well armed.

Whether all these characters are found in the other genera commonly associated with the *Muraenesox* is not yet known. The family, as understood by us, seems divisible into two well-marked groups, which are, perhaps, as distinct from each other as from the *Echelidæ* or the *Congridæ*. The species of this family are not very numerous, and a large proportion are American. In general appearance and habits they approach the congers. All are plainly colored, and some descend to rather deep water. One genus represented in Porto Rico.

MURÆNESOCINÆ:

- a. Dorsal and anal fins well developed throughout, the dorsal beginning nearly above gill-opening; snout moderately produced; vomerine teeth very strong.
 b. Teeth in jaws in several series; gill-openings wide..... MURÆNESOX, 11
 bb. Teeth in jaws biserial, small; vomer with a series of long, pointed canines; tail about four times as long as rest of body; gill-openings narrow HOPLUNNIS

STILBISCINÆ:

- aa. Dorsal and anal fins very low anteriorly, developed chiefly on the tail.
 c. Tail about as long as rest of body; teeth moderate; dorsal beginning before the vent. Body moderately elongate, diameter more than one-thirtieth the length; dorsal beginning just before vent NEOCONGER
 cc. Tail short, little more than half as long as rest of body; teeth all uniserial, unequal, some of them canine-like; body very slender, whip-shaped.
 d. Dorsal fin beginning not far behind pectoral..... LEPTOCONGER
 dd. Dorsal fin beginning behind the vent..... STILBISCUS
 ddd. Dorsal fin beginning close behind the nape GORDIICHTHYS

Genus 11. *MURÆNESOX* McClelland.

Body robust. Dorsal and anal fins well developed, dorsal beginning nearly above gill-opening. Mouth large, teeth in jaws in several series, those of one series enlarged and depressed, forming long canines in front; vomer with several long series of teeth, the middle one of strong canines.

This genus contains numerous species, large, conger-like eels, some of which are found in all warm seas. They are remarkable for the strong armature of the vomer. One species known from Porto Rico.

13. *Murænesox savanna* (Cuvier).

Median series of teeth on vomer distinctly tricuspidate in the young, becoming entire with age, with nearly even surface; pectorals as long as maxillary, 2.66 in head; eye 2 in snout, which is 4.5 in head; dorsal inserted over gill-opening.

Brown above, silvery below; dorsal and anal edged with black.

Cuba to Rio Janeiro, not common; occasional in the Mediterranean Sea. Not obtained by us in Porto Rico, but included on authority of Poey.

Muræna savanna Cuvier, Règne Animal, ed. 2, vol. 2, 350, 1829, Martinique.

Conger brasiliensis Ranzani, Nov. Spec. Pisc. Diss. Prima, IV, 17, pl. 13, fig. 1, 1838, Brazil.

Congrus curvidens Richardson, Voy. Erebus and Terror, III, 184, no locality given.

Cynoponticus ferox Costa, Fauna Napoli, Pesc., pl. 28, 1854, Naples.

Conger limbatus Castlenau, Anim. Am. Sud. 83, pl. 43, fig. 3, 1855, Rio Janeiro.

Murænesox savanna, Poey, Fauna Puerto-Riqueña, 344, 1881; Stahl, l. c., 166, 1883; Jordan & Evermann, l. c., 360, 1896.

Family XI. MORINGUIDÆ. The Whip Eels.

Excessively elongate eels, with the abdominal cavity forming two-thirds of the length (the anus opening about the commencement of the last third) and the heart far behind gills. Body scaleless; cylindrical, with the trunk much longer than the tail; pectorals none or small; vertical fins but little developed, limited to the tail; posterior nostril in front of small eye; cleft of mouth narrow; teeth uniserial; gill-opening rather narrow, inferior.

Genus 13. *APHTHALMICHTHYS* Kaup.

Characters of the genus included above.

14. *Aphthalmichthys caribbeus* Gill & Smith.

Body exceedingly long and slender, the trunk of uniform depth throughout and not compressed, post-anal part of body slightly compressed. Head long, not conspicuously wider than body, its length contained 13 times in total length of fish, nearly 9 times in distance between snout and vent, and 4 times in tail. Anterior nostrils large and tubular, at end of snout; posterior nostrils large, immediately in front of eye. Length of snout one-seventh that of head. Gape equal to depth of body, about one-fourth length of head extending considerably beyond eye. Lower jaw projecting. Teeth rather large, canine, retrorse, uniserial, a single row of small teeth on vomer. Eye very small and rudimentary, little larger than the posterior nostril, and contained over 20 times in length of head and 3 times in snout. Interorbital narrow, about half length of snout. Depth of body contained 54 times in total length, 4.2 times in head. About the length of the head behind vent, a narrow, shallow groove extends backward on both the dorsal and ventral surfaces; in this groove the dorsal and anal fins exist as low ridges, and are entirely undeveloped until about length of head from end of body; they then become less than half a millimeter in height and are confluent around the blunt tail. The pectoral is a mere rudiment lying on the upper posterior edge of the gill-slit and is less than half the length of the latter. The branchial openings are nearly vertical slits, about twice length of eye, and separated from each other by a space equal to 1.5 times their length. The ventral opening is situated a little more than two-thirds the distance from snout to end of tail. A series of large circular pores along lateral line.

Color in life, a uniform grayish-olive, without markings. The type, 270 mm. long, collected among coral (*Porites*) at San Geronimo, Porto Rico, by Mr. George M. Gray.

The following table gives the measurements:

Measurements.	Milli- meters.	Measurements.	Milli- meters.
Length	270	Length of snout	3
Length of head	21	Interorbital	2.5
Length of head and trunk	185	Dorsal groove behind vent	21
Length of post-ventral region	85	Anal groove behind vent	21
Depth at gill-slit	5	Dorsal fin appears above surface, in front of end of tail	21
Depth at vent	4.5	Anal fin appears above surface, in front of end of tail about	16
Depth at dorsal origin	5		
Length of gape	5		

Aphthaliethys caribbeus Gill & Smith, Science, n. s., vol. XI, Nos. 286, 973, June 22, 1900, San Geronimo, Porto Rico.

Family XII. MYRIDÆ. The Worm Eels.

End of tail surrounded by the confluent vertical fins; posterior nostril in, or very near, upper lip, the tongue more or less fully adnate to the floor of the mouth. The species are usually of small size and plain colors, more or less worm-like in form, and inhabit sandy coasts in tropical seas. The genera have but few species each. They are intermediate in character between the *Ophichthyidæ* and the *Muraenesocidæ*. The osteology has not been carefully studied, but they will probably be found to be most nearly related to the latter family, if indeed the two should not be, as in Bleeker's arrangement, reunited with the *Leptocephalidæ*.

- a. Body elongate, subterete; pectoral present, sometimes minute; anterior nostril tubular; dorsal fin beginning behind head; teeth small.
- b. Dorsal fin beginning behind vent; no teeth on vomer; teeth mostly uniserial; body slender, terete..... AHLIA
- bb. Dorsal fin beginning before vent; vomer with teeth.
- c. Dorsal beginning at a point about midway between gill-opening and vent; pectoral very small; teeth subequal; body slender, terete; tail much longer than rest of body..... MYROPHIS
- aa. Body short, much compressed; pectoral almost invisible; mouth narrow; snout obtuse, depressed; vertical fins well developed, dorsal beginning behind gill-opening..... CHILORHINUS, 13

Genus 13. CHILORHINUS Lütken.

Body short, much compressed, especially posteriorly; mouth narrow, lower jaw slightly the shorter; snout depressed, obtuse, two rows of short conic or incisor-like teeth on each palatine and a group of similar teeth on nasals, back of which is a large tooth on anterior end of vomer; teeth on lower jaw triserial; pectoral fin very minute; vertical fins well developed, dorsal commencing behind gill-opening. One species known.

15. *Chilorhinus suensonii* Lütken. *Worm Eel*.

Head 5; depth 14.4; eye 8; snout 6.3; interorbital 5.6; cleft of mouth 3.6 in head; head and trunk a little less than half total length; dorsal fin beginning a short distance back of gill-opening, slightly nearer vent than tip of snout.

Body elongate, trunk slightly compressed anteriorly, much so posteriorly; head large, deeper and wider than body, its width somewhat greater than its depth, its upper surface convex; snout slightly tapering, rather broadly truncate; eye chiefly lateral, posterior border just above the angle of mouth; cleft of mouth horizontal, lower jaw included, lips fleshy, upper slightly overhanging the closed mouth, its outline somewhat irregular; a double series of short canine-like teeth on each palatine, similar teeth on nasals, behind which is a large tooth on vomer; teeth on lower jaw triserial; anterior nostril in a short tube well forward at edge of lip, directed downward; posterior nostril without tube, opening in lower part of upper lip just in front of eye, scarcely visible when mouth is closed; skin not firmly attached to body, more or less movable, especially full and loose about head, where it is gathered into longitudinal folds; dorsal and anal regular in outline, somewhat lower near their confluence; anal very slightly lower than dorsal, each with numerous rays; lateral line present, being a straight shallow groove running nearer dorsal outline anteriorly, containing a series of muciferous pores.

Color in spirits: Nearly everywhere uniform dark-brown; belly and ventral surface of head white.

Twenty-seven individuals, from 1.5 to 4.4 inches in length, were collected at Mayaguez, Ponce, Arroyo, Hucare, and Boqueron. Hitherto known only from St. Croix, Danish West Indies.

Chilorhinus suensonii Lütken, Vid. Med. Naturg. Foren. Kjöben., I, 1851, St. Croix; Jordan & Evermann, I. c. 372, 1896.

Family XIII. OPHICHTHYIDÆ. The Snake Eels.

This family includes those enchelycephalous eels which are scaleless and have the end of the tail projecting beyond dorsal and anal fins, and without rudiment of a caudal fin. Anterior nostrils placed in upper lip, opening downward; gill-openings not confluent; tongue more or less fully adnate to floor of mouth. Species usually moderate or small in size, very abundant in tropical seas, especially about coral reefs. The eggs are numerous, of moderate size, similar to those of ordinary fishes.

Of 12 genera of this family occurring in America only 4 are known to have species in Porto Rico.

- a. Body without traces of fins anywhere; teeth all small, conical; gill-openings near together, subinferior; anterior nostril tubular; tongue scarcely free in front; mouth small.
- b. Gill-slits inferior, converging forward SPHAGEBRANCHUS, 14
- bb. Gill-slits small, lateral, placed vertically VERMA
- aa. Body with distinct fins, at least on back.
- c. Anal fin wholly wanting; no pectoral fin; dorsal fin high, beginning on head; gill-openings subinferior, converging; anterior nostrils tubular; tongue slender, somewhat free in front LETHARCHUS
- cc. Anal fin well developed; anterior nostril usually in a short tube near tip of snout.
- d. Teeth blunt, mostly granular or molar; vomer with teeth; pectoral fins present, small.
- e. Dorsal rather high, beginning on head, before gill-opening MYRICHTHYS, 15
- cc. Dorsal fin beginning behind gill-opening, fin usually low PISODONOPHIS
- dd. Teeth all pointed, none of them molar; vomer with teeth.
- f. Dorsal fin beginning before nape, on anterior part of head; pectoral fin small or wanting.
- g. Pectoral fins wholly wanting; body compressed, dorsal fin high. CALLECHELYS
- gg. Pectoral fins small, but present; body elongate, subterete, dorsal fin moderate. BASCANICHTHYS
- ff. Dorsal fin beginning more or less behind gill-opening.
- h. Teeth subequal, with no elongate canines on jaws or vomer. OPHICHTHUS, 16
- hh. Teeth unequal, some of them long canines, either on vomer or on sides of one or both jaws; mouth large, snout short, and eyes more or less superior. MYSTRIOPHIS

Genus 14. SPHAGEBRANCHUS Bloch.

This genus contains several little-known species of small eels remarkable for showing no trace of fins in the adult stage. The snout projects beyond the small mouth, giving a shark-like profile, and the small teeth are mostly uniserial. Gill-slits inferior and converging. The name *Sphagebranchus* was based on a species which evidently belongs to the genus. It has, therefore, clear priority over *Ichthyapus* and *Apterichthys*.

This genus is the most simple in structure among the genera of *Ophichthyidæ*, as *Ophichthus* is probably the most specialized. Its loss of fins is doubtless due to degeneration, but it seems nearer the primitive type than *Brachysomophis* or *Ophichthus*. Only one species obtained in Porto Rico.

- a. Head 6 times in trunk *anguiformis*
- aa. Head little more than 4 times in trunk.
- b. Color brown, head mottled. *scelachops*
- bb. Color greenish, without mottlings *ophioneus*, 16

16. *Sphagebranchus ophioneus* Evermann & Marsh, new species.

Head 4.5 in distance from tip of snout to vent, 12.5 in total length; depth of trunk 48 times in total length, depth of head 36 times; distance from tip of snout to vent 2.8 in total length; snout 6 in head, mandible 2.5, interorbital 12.7.

Body cylindrical, tail tapering; longest diameter of head greater than that of body; head forward of occipital region long, slender, conic, and sharply pointed; eye minute; mouth inferior, the pointed snout far overhanging, tip of lower jaw considerably in front of eye; teeth numerous, directed backward, conic, sharp, depressible, uniserial in each jaw and on vomer; roof of mouth not entirely covered by lower jaw, exposing a triangular space in front of its tip, a single tooth exposed in front on median line when the mouth is closed; in the preserved specimen the tooth next behind on each side is also exposed, probably due to the contraction of the fleshy tip of lower jaw; a few fleshy papillæ at edge of upper lip on each side opposite tip of lower jaw; nostrils not tubular, the anterior on ventral side of fleshy tip of snout, the entrance guarded by a pair of fleshy papillæ; the posterior in upper lip just behind tip of lower jaw which covers nostril when mouth is closed; a row of muciferous pores on each ramus of mandible; scattered pores about snout, behind eyes and along upper lip, and a transverse row across head at occiput; lateral line distinct, complete, and continuous, commencing on side of head and extending to tip of tail, consisting of a slight elevation of the integument, the pores arranged at regular

intervals along its lower edge; gill-slits 5.4 in head, entirely inferior, converging anteriorly where they are very narrowly separated; skin over branchial chambers with longitudinal plications.

Color in spirits: Everywhere pale-green, without evident markings; traces of a faint yellowish area on side of head, in front of occiput, and on snout; tip of snout a little darker.

One specimen, the type (No. 49526 U.S.N.M.) 11.25 inches long, dredged in 4 fathoms at Fish Hawk Station 6065, off Mayaguez, Porto Rico, January 20.



FIG. 7.—*Sphagebranchus ophioneus*.

Genus 15. MYRICHTHYS Girard.

Teeth mostly blunt and molar; pectoral fins small; dorsal beginning on head before gill-opening; otherwise essentially as in *Ophichthus*. Coloration variegated. Species numerous, found in most tropical seas. Only one of the three American species known from Porto Rico.

- a.* Spots on body large, black, most of them with a distinct pale center, the ground-color paler *oculatus*, 17
aa. Spots on body large, round, nearly whitish in color, the ground-color dark *acuminatus*

17. *Myrichthys oculatus* (Kaup).

Head 4.2 in trunk (tip of snout to vent); eye 2.5 in snout; interorbital 6 in head, slightly shorter than snout; cleft of mouth 3.25 in head.

Body scaleless, very slender, tail tapering to a point, without caudal fin; dorsal and anal fins not continuous around it; dorsal low, commencing on head far in advance of gill-opening, extending almost to tip of tail; anal similar, shorter and lower; pectoral reduced to a thin, narrow, membranaceous flap immediately behind gill-opening; anterior profile of head straight, making an angle of 45° with the horizontal mouth; eye very small; nostril tubular, on side of tip of snout; lower jaw considerably the shorter, upper overhanging, exposing some of the teeth, which are all small, molar, and scarcely elevated.

Color in spirits: Body grayish, trunk paler below; side with two rows of about 35 large roundish black spots not sharply circumscribed, the center of each with a small pale spot, upper row close to dorsal fin, other row just below, the spots above alternating with those below; a large dark spot on the head in front of the dorsal; lower jaw and snout with numerous much smaller dark spots, some of these with a pale center; dorsal with dark-brown edge and many diffuse blotches of same; anal pale, unmarked.

Tropical Atlantic, Cuba to Surinam, and Cape Verde Islands. One specimen, nearly 2 feet long, collected at Hucares, Porto Rico.

Pisodonophis oculatus Kaup, Apodes, 22, 1856, Curaçao.

Ophisurus latimaculatus Poey, Repertorio, II, 252, pl. 3, fig. 1, 1867, Cuba.

Ophichthys latimaculatus, Poey, Fauna Puerto-Riqueña, 345, 1881.

Myrichthys oculatus, Jordan & Evermann, l. c., 376, 1896.

Genus 16. *OPHICHTHUS* Thunberg & Ahl.

This genus contains all the ophisuroid eels which have sharp teeth, no marked canines, well-developed pectoral fins, and dorsal inserted behind head. The species are very numerous in the tropical seas, and many attempts have been made to split the group into smaller genera. Notwithstanding the great differences when extremes are compared, these small genera can not be well defined. Only one of the ten American species is known from Porto Rico.

a. Teeth of upper jaw in two or three series.

b. Teeth of lower jaw uniserial, or nearly so; vomerine teeth in one series or slightly biserial in front.

CRYPTOPTERUS:

c. Coloration uniform, or nearly so; teeth of lower jaw not quite uniserial; tail half longer than rest of body..... *puncticeps*

OPHICHTHUS:

cc. Coloration not uniform; anterior teeth slightly enlarged; eye rather large, nearly median.

d. Sides of body with large, round, black spots; head with smaller ones; dorsal inserted opposite tip of pectoral..... *havanuensis*

dd. Sides of body with large, round, whitish spots; dorsal inserted behind tip of pectoral..... *retropinnis*

bb. Teeth of lower jaw in two to four series.

MURÆNOPSIS:

e. Vomerine teeth in one row; anterior teeth of jaws or vomer sometimes enlarged; teeth in both jaws biserial, those of inner series sometimes small and turned inward.

f. Dorsal beginning an eye's diameter behind tip of pectoral..... *guttifer*

ff. Dorsal beginning over or just before tip of pectoral..... *ocellatus*

SCYTALOPHIS:

ee. Vomerine teeth biserial throughout; teeth in both jaws biserial, subequal; no canines. Color plain-brownish.

g. Eye large, more than half length of snout.

h. Head rather short, 2.5 to 3 in trunk..... *gomesii*, 18

hh. Head long, 1.75 to 2.25 in trunk; pectoral a little longer than gape..... *magnioculis*

gg. Eye small, 2.5 in snout; gill-opening narrow; anterior nostril with long tube; pectoral longer than gape... *parilis*

18. *Ophichthus gomesii* (Castelnaud). *Sea Serpent.*

Head 2.8 in trunk; head and trunk 2 in tail; eye about equal to interorbital space, 1.5 in snout, which is 6 in head; pectoral 2.8 in head; teeth small, sharp and subequal, biserial on each jaw and on vomer; upper jaw the longer; pores on head and lower jaw; dorsal and anal very low.

Color in spirits: Everywhere brownish above, color laid on in a multitude of very small brown points somewhat in rows, on a yellow ground; ventral surface yellow with a few brown points; side of head and lower jaw with dusky spots; dorsal edge of pectoral with dusky points.

South Carolina to Rio Janeiro; generally common, especially about the Florida Keys and Cuba. One example 8.5 inches long collected at Mayaguez.

Ophisurus gomesii Castelnaud, Anim. Amér. Sud, 84, pl. 44, fig. 2, 1855, Rio Janeiro.

Ophisurus chrysops Poey, Memorias, II, 321, 1867, Havana.

Orydontichthys brachyurus Poey, Synopsis, 426, 1868, Havana.

Orydontichthys macrurus Poey, Anal. Soc. Hist. Nat. Esp., 1880, 254, Havana.

Ophichthus gomesii, Jordan & Evermann, l. c., 384, 1896.

Family XIV. MURÆNIDÆ. The Morays.

The *Murænidæ* represent the most degenerate type of eels so far as the skeleton is concerned, and are doubtless the farthest removed from the more typical fishes from which the eels have descended. The essential characters of the family are thus stated by Dr. Gill: "Colocephalous apodals with conic head, fully developed opercular apparatus, long and wide ethmoid, posterior maxillines, pauciserial teeth, roundish, lateral branchial apertures, diversiform vertical fins, pectoral fins (typically) suppressed, scaleless skin, restricted interbranchial slits, and very imperfect branchial skeleton, with the fourth branchial arch modified, strengthened, and supporting pharyngeal jaws."

The morays may be readily distinguished from other eels by their small round gill-openings and by absence of pectorals. The body and fins are covered by a thick leathery skin, the occipital region is elevated through the development of the strong muscles which move the lower jaw, and the jaws are usually narrow and armed with knife-like or else molar teeth. They inhabit tropical and subtropical waters, being especially abundant in crevices about coral reefs. Many of the species reach a large size, and all are voracious and pugnacious. Coloration usually strongly marked, the color cells being highly specialized. Only two of the eight American genera represented in Porto Rico.

- a. Vertical fins well developed, dorsal beginning before vent.
- b. Posterior nostril an oblong slit; anterior in a short tube; teeth all pointed; dorsal beginning above gill-opening; canine teeth strong; tail moderate..... ENCHELYCORE
- bb. Posterior nostril circular, with or without tube; tail moderate, not twice as long as trunk; body not excessively elongate.
- c. Teeth all, or nearly all, acute, none of those in jaws obtuse or molar-like.
- d. Anterior nostrils without tube; vomerine teeth in many series; lips with a free fold..... PYTHONICHTHYS
- dd. Anterior nostrils each with a long tube; vomerine teeth in one or two series; lips continuous with skin of head.
- e. Posterior nostrils without tube, margins sometimes slightly raised.
- f. Dorsal fin inserted behind head, over or behind the gill-opening..... RABULA
- ff. Dorsal fin inserted on head, considerably before gill-opening..... LYCODONTIS, 17
- cc. Posterior nostrils as well as anterior each in a conspicuous tube..... MURENA
- ce. Teeth mostly obtuse, molar-like; only anterior nostrils tubular; cleft of mouth rather short; dorsal beginning before gill-opening..... ECHIDNA, 18
- aa. Vertical fins rudimentary, confined to end of tail (often appreciable only on dissection, or altogether wanting); teeth rather small, pointed, subequal, in several series; posterior nostril round, with a short tube, or none, CHANOMURENA

Genus 17. LYCODONTIS McClelland.

This genus, as here understood, comprises the great bulk of the *Murena*, including all the species with sharp teeth, the body normally formed, anterior nostrils only tubular, and the dorsal fin beginning on head. *Priodonopsis* with serrated teeth has been recognized as a distinct genus by Bleeker, but the character in question disappears by degrees and seems not to be suitable for generic distinction. The morays of this genus are everywhere abundant in tropical seas, where some reach a great size. They are the most active and voracious of the eels, often showing much pugnacity. Most of them live in shallow water about rocks or reefs. Four species known from Porto Rico.

- a. Teeth all entire, with no serrations anywhere, and none of them with basal lobes.
- b. Body without small, round, bluish-white or yellow spots, the spots, if any, blackish or dull-grayish; dorsal without distinct paler margin, or with merely the very edge whitish.
- c. Dorsal with a distinct black margin; anal with a pale edge; teeth uniserial..... *vicius*
- cc. Dorsal without distinct, darker margin, its border colored nearly or quite like rest of fin.
- d. Body and tail covered with close-set dark points; tail longer than rest of body..... *virescens*
- dd. Body and tail not covered with close-set dark points.
- e. Color olivaceous or brownish, with conspicuous markings, marblings or spots darker than the ground-color; belly without distinct transverse lines, marked like the back and sides; tail slightly longer than rest of body.
- f. Dark markings forming narrow reticulations, never rounded spots; these reticulations dark-lilac in color, covering back and sides, some of them inclosing irregular polygons..... *polygonius*
- ff. Dark markings in the form of rounded spots, more or less confluent, sometimes obscuring the pale ground-color..... *moringa*, 19
- cc. Color dark-brown, dark-green, or blackish, either plain or with faint markings.
- g. Belly with black, wavy, transverse lines; no dark lines along dorsal fin..... *mordax*
- gg. Belly without black transverse lines; body nearly plain dark olive-brown.
- h. Dorsal and anal with dark longitudinal streaks; chin pale but not white..... *funbris*, 20
- hh. Dorsal and anal without dark longitudinal streaks; chin white..... *albimentis*, 21
- ccc. Color brownish-black with irregular pale-grayish spots of various sizes; margin of anal not pale; cleft of mouth less than half head..... *sancta-helenae*
- bb. Body with distinct small spots, blue, white, or yellow.
- i. Dorsal and anal without distinct colored margin; pale spots mostly smaller than eye.
- j. Teeth of upper jaw uniserial.
- k. Vomerine teeth uniserial; spots irregular, few, and scattered; dorsal colored like back.
- kk. Vomerine teeth biserial; entire body covered with small blue dots; dorsal with vertical bluish streaks.. *conspersus*
- jj. Teeth of upper jaw biserial; body with small yellow spots.
- l. Vomerine teeth uniserial, mostly small and rounded; color nearly uniform from head to tail; spots innumerable..... *miliavis*
- ll. Vomerine teeth biserial, small and rounded; color dark-brown, with yellow points excessively numerous. *elaboratus*
- ii. Dorsal with a blackish border, interrupted with white; anal with white markings; body with close-set, irregular pale spots.
- m. Body greenish, marbled with brownish, almost obscured by the ground-color..... *obscuratus*
- mm. Body rich-yellow or tawny; head and body covered with small, round, white spots..... *jordani*, 22
- PRIODONOPSIS:**
- aa. Teeth serrate, more or less.
- n. Color brown, with irregular light-yellowish spots irregularly placed; dorsal with large, dark spots on its edge, these sometimes obsolete, usually running together to form a continuous dark band; anal black-edged; teeth large, uniserial, the larger ones serrate; mouth nearly closing..... *ocellatus*

19. *Lycodontis moringa* (Cuvier). *Common Spotted Moray; Hamlet.*

Head 3.2; snout 6 in head; eye 1.5 in snout; cleft of mouth 2.5 in head; tail somewhat longer than trunk. Teeth strong and sharp; two or three long depressible teeth on vomer in front and a median row of small ones behind; teeth in jaws uniserial, long in front, those behind retrorse.

Color in spirits: Dark-brown, everywhere mottled and reticulated with paler or light yellow, which is properly the ground-color, but is nearly covered by the dark pigment; ventral surface of head pale with a few brown spots; fins colored like body.

West Indies, Pensacola to Rio Janeiro and St. Helena. Two young examples, about 7 inches long, from Culebra.

Muraena maculata nigra (the Black Moray), Catesby, Nat. Hist. Carolina, etc., pl. 21, 1738, Bahamas.

Muraena moringa Cuvier, Règne Animal, ed. II, vol. II, 352, 1829, Bahamas; after Catesby.

Gymnothorax rostratus Agassiz, Spix, Pisc. Bras., 91, pl. 50, a, 1829, Brazil; Stahl, l. c., 80, 1883.

Muraena moringua Richardson, Voy. Erebus and Terror, Fishes, 89, 1844, Jamaica.

Muraena punctata Gronow, Catalogue Fishes, 18, 1854, North America.

Marcenophis curvilincata Castelnau, Anim. Amér. Sud, Poiss., 81, pl. 42, fig. 2, 1855, Rio Janeiro.

Muraenophis caramura Castelnau, Anim. Nouv. Rares Amérique du Sud, 85, pl. 43, fig. 1, 1855, Bahia.

Gymnothorax flavoscriptus Poey, Enumeratio, 158, 1873, Cuba.

Gymnothorax picturatus Poey, Anal. Soc. Esp. Hist. Nat. 1880, 257, Cuba.

Lycodontis moringa, Jordan & Evermann, l. c., 395, 1896.




FIG. 8.—*Lycodontis moringa*.

20. *Lycodontis funebris* (Ranzani). *Black Moray; Morena Verde.*

Tail a little longer than head and trunk. Teeth uniserial in jaws in the adult; teeth on vomer uniserial (var.? *erebus*), or biserial (*funebris*); long, depressible canines on front of vomer; jaws not completely closing; eye 2 to 2.5 in snout, above middle of gape; cleft of mouth 2.5 in head; head 2.5 in trunk.

Dark olive-brown, nearly plain, paler on throat, sometimes with very faint darker marblings; dorsal and anal fins with dark lines running longitudinally; belly without black transverse lines.

Found in tropical America, on both coasts; the largest of our eels, reaching 5 or 6 feet or more; extremely ferocious; common from Florida Keys to Rio Janeiro and from Gulf of California to Panama. A single example, 4.12 inches long, obtained at Fajardo February 17, 1899.

Muraena maculata nigra et viridis, Catesby, Nat. Hist. Carolina, etc., pl. 20, 1738, Bahamas.

Gymnothorax funebris Ranzani, Nov. Comm. Ac. Sc. Bonon., IV, 76, 1840, Brazil.

Muraena lineopinnis Richardson, Voy. Erebus and Terror, Fishes, 89, 1844, Puerto Cabello.

Teniophis westphali Kaup, Aale Hamburg Mus., I, 1859.

Thysoidea aterrima Kaup, Aale Hamburg Mus., I, 22, 1859.

Muraena infernalis Poey, Memorias, II, 347 and 354, 1860, Cuba.

Thysoidea concolor Abbott, Proc. Ac. Nat. Sci. Phila. 1860, 479, Vera Cruz.

Muraena erebus Poey, Memorias, II, 426, 1861, Cuba.

Lycodontis funebris, Jordan & Evermann, l. c., 396, 1896.

21. *Lycodontis albimentis* Evermann & Marsh, new species.

Head 6.4; snout about 7; mandible 1.8; eye 2.3 in cleft of mouth; head and trunk somewhat compressed, tail considerably so; cleft of mouth a little less than half of head; jaws equal; teeth strong, conical, and sharp, in one rather irregular row in upper jaw, strongest in front; in lower jaw one row of similar teeth, becoming smaller posteriorly, those in front stronger than in upper jaw; eye rather large, slightly nearer tip of snout than angle of mouth, bordering the lip; upper lip with a row of very minute cirri along the edge, resembling teeth, visible only under a lens; a row of rather large pores around upper lip and a pair on top of snout; anterior nostril tubular, in the lip at angle of snout; posterior nostril without tube, just above front of eye; fins higher posteriorly.

Color in spirits: Everywhere uniform dark-brown, except lower part of head and edge of fins; upper lip and lower jaw perfectly white, the brown sharply demarcated from the colorless area, which begins just back of mandible, extending over lower jaw, past angle of mouth, along upper lip, under eye, and around snout; inside of mouth colorless; fins darker than body, narrowly pale-edged posteriorly.

One very small specimen, only 2 inches long, brought up by the tangle from coral bottom in 15 fathoms at Fish Hawk station 6093, off Culebra Island, February 8; possibly the young of a known species, but the sharply defined white area on the lower jaw is so prominent a character that we have thought it best to describe the species as new. Type No. 49527 U.S.N.M.




FIG. 9.—*Lycodontis albimentis*.

22. *Lycodontis jordani* Evermann & Marsh.

(PLATE 2.)

Head 7 in total length; depth about 14; eye 8 in head; snout 5; gape 2.2; interorbital a little less than snout. Teeth uniserial, strong, sharp, not close-set, all entire and without basal lobes; tail considerably longer than rest of body; gill-opening smaller than eye; snout rather pointed, lower jaw the shorter, the mouth capable of being completely closed. Dorsal fin high, much higher than anal; nasal tube long, about 3 in eye.

Color in life: Tawny-ochraceous, paler below; upper jaw gray; iris blue; longitudinal brown stripes on side of head in front of gill-opening; head and body covered with numerous small, round, white spots, those on head smallest; a series of larger ones along upper part of side, and one or two irregular series of large ones on side of belly; between these, on middle of side, the spots are smaller; dorsal with an irregular series of small white spots along the base, and another series of about 16 much larger, more quadrate spots of same color along edge of fin, some of the spots cutting the border, which is black; anal similarly spotted and with black border.

In alcohol the general color is grayish-black, yellowish below, the fawny-ochraceous or yellow becoming darker, almost black, and the white spots on body becoming yellowish.

This species seems to be related to *L. obscuratus* (Poey), but differs markedly from it in color. Only the type (No. 49358, U. S. N. M.), a specimen about 15 inches long, was obtained. This was collected at Mayaguez, January 20, 1899.

Lycodontis jordani Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 352, Mayaguez, Porto Rico.

Genus 18. ECHIDNA Forster.

This well-marked genus is distinguished from the other Morays by the blunt teeth. It contains 12 or 15 species, most of them belonging to the Western Pacific and representing the highest degree of specialization among the morays, as *Uropterygius* represents the extreme of degradation. The name *Echidna* was suggested for this group of eels long before its application by Cuvier to a genus of Australian monotremes. Only one species known from Porto Rico.

23. *Echidna catenata* (Bloch). *Morena*.

Head 3.7 in trunk; eye 2 in snout, which is 6 in head; cleft of mouth 3 in head; trunk a little longer than tail; teeth somewhat molar-like, not very blunt, chiefly uniserial, but reduced in size and biserial in rear of upper jaw; a median series on vomer.

Color in spirits: Ground-color pale-yellow, with some 30 heavy dark-brown transverse bars, some oblique, some branched, some connected, forming heavy reticulations; on ventral portion of trunk these bars are broken into definite roundish spots, these elongate in front of anal, forming a single median series; intermediate pale area of ground-color everywhere marked with dark-brown in spots of all sizes from punctulations to the size of pupil or larger; markings of body extend upon fins.

West Indies, Bermuda to Surinam, generally common. A very handsome eel. Mr. Gray obtained 3 fine specimens at San Geronimo; our collection contains one, of 6.5 inches, from Arroyo.

Muraena seu conger brasiliensis Seba, Thesaurus, II, 72, pl. 69, figs. 4 and 5, 1738, Brazil.

Gymnothorax catenatus Bloch, Aust. Fische, XII, 74, pl. 415, fig. 1, 1795, Coromandel, an error; Stahl, l. c., 166, 1883.

Muraenophis catenula Lacépède, Hist. Nat. Poiss., V, 628 and 641, 1803, Palmerston Island; after Bloch.

Echidna flavofasciata Poey, Repertorio, II, 264, 1868, Cuba.

Echidna fuscomaculata Poey, Repertorio, II, 263, 1868, Cuba.

Echidna catenata, Poey, Fauna Puerto-Riqueña, 345, 1881; Jordan & Evermann, l. c., 403, 1896.

Family XV. ELOPIDÆ. The Tarpons.

Body elongate, more or less compressed, covered with silvery, cycloid scales; head naked. Mouth broad, terminal, the lower jaw prominent. Premaxillaries not protractile, short, the maxillaries forming lateral margins of upper jaw; maxillary composed of about three pieces, extending backward beyond eye; an elongate bony plate between the branches of lower jaw (analogous to gular plate in *Amia*); bands of villiform teeth in both jaws and on vomer, palatines, pterygoids, tongue, and base of skull; no large teeth. Eye large, with an adipose eyelid. Opercular bones thin, with expanded membranaceous borders; a scaly occipital collar. Gill-membranes entirely separate, free from isthmus. Branchiostegals numerous (29 to 35). Gillrakers long and slender. Pseudobranchiæ present or absent. Belly not keeled nor serrated, rather broad and covered with ordinary scales. Lateral line present. Dorsal fin inserted over or slightly behind ventrals; caudal fin forked; no adipose fin; dorsal and anal depressible into a sheath of scales; pectorals and ventrals each with a long accessory scale. Parietal bones meeting along top of head. Pyloric caeca numerous.

Genera three, species about five, forming two well-marked subfamilies, both widely distributed in the tropical seas. Not much valued as food, the flesh being dry and bony.

MEGALOPINÆ:

a. Pseudobranchiæ none; body oblong, covered with large scales; anal fin larger than dorsal; last ray of dorsal produced in a long filament.

b. Dorsal fin inserted notably behind insertion of ventrals..... TARPON, 19


ELOPINÆ:

aa. Pseudobranchiæ large; body elongate, covered with small scales; anal fin smaller than dorsal; last ray of dorsal not produced in a filament..... ELOPS, 20

Genus 19. TARPON Jordan & Evermann. Grande Écaille.

Body oblong, compressed, covered with very large, thick, silvery, cycloid scales; belly narrow, but not carinated, its edge with ordinary scales. Mouth large, oblique, the lower jaw prominent; maxillary broad, extending beyond eye. Villiform teeth on jaws, vomer, palatines, tongue, sphenoid, and pterygoid bones. Eye very large, with an adipose eyelid. Lateral line nearly straight, its tubes radiating widely over surface of scales. Branchiostegals 23. Pseudobranchiæ wanting. Gillrakers long and slender. Dorsal fin short and high, inserted behind ventrals (over ventrals in *Megalops*), its last ray elongate and filamentous as in *Megalops*, *Dorosoma*, and *Opisthonema*; anal fin much longer than dorsal, falcate, its last ray produced; caudal widely forked; pectorals and ventrals rather long; anal with a sheath of scales; dorsal naked; caudal more or less scaly; a collar of large scales at the nape. Vertebrae about 57 (28 + 29). Size very large, the largest of the herring-like fishes.

The posterior insertion of the dorsal fin distinguishes the single species of *Tarpon* from the East Indian *Megalops cyprinoides*, a fish of similar habit, in which the dorsal is inserted above the ventrals.

FIG. 10.—*Tarpon atlanticus*.24. *Tarpon atlanticus* (Cuvier & Valenciennes).

Tarpon; *Tarpum*; *Grande Écaille*; *Silver King*; "*Sabalo*"; *Savanilla*; *Savalle*.

Head 4; depth 3.75; eye 4.45; snout 5; maxillary 1.65; mandible 1.70; interorbital 4.8; preorbital 11; D. 12; A. 19 or 20; pectoral 1.3; ventral 1.75; scales 5-42-5. Proportions in young of about 3 inches: Head 3.34; depth 4.7; eye 3.3; snout 4.75; maxillary 1.65; mandible 1.6; interorbital 4.6.

Body elongate, compressed, not elevated, with large cycloid scales; mouth large, maxillary reaching far beyond eye, lower jaw projecting; villiform teeth on jaws, vomer, palatines, tongue, sphenoid and pterygoid bones; lateral line straight with radiating tubes; pseudobranchiæ minute; last dorsal ray produced, the fin inserted behind ventrals; pectorals and ventrals each with an accessory scale. Dark above, silvery on sides.

Found from Long Island to Brazil; common on our southern coasts, especially about Florida; common about Porto Rico where it evidently breeds, as numerous immature individuals were taken at Hucars and Fajardo. The four examples from Hucars are from 7.5 to 11.5 inches long and were seined in a small brackish pool of dark-colored water, not over 5 feet deep, in the corner of a mangrove swamp, and at that time (February) entirely separated from the ocean by a narrow strip of land scarcely 25 feet wide. The 13 others are nearly all very young, of 2.25 to 3.25 inches, collected at Fajardo. No large individuals were seen. The young of the tarpon seems to be rare or wanting in collections. The U. S. National Museum contains one individual of about 9 inches. We do not know that any as small as those above mentioned have ever before been recorded.

The tarpon reaches a length of 5 or 6 feet and a weight of 30 to more than 300 pounds. The largest one recorded as taken on a hook weighed 209 pounds, and the largest taken with the harpoon weighed 383 pounds, if we may believe the record; but examples weighing over 100 pounds are not often seen. The silver king is the greatest of game-fishes. "An immense and active fish, preying eagerly on schools of small fry, in pursuit of which it ascends fresh-water streams quite a long distance." It is often dangerous to seine fishermen, leaping over or through their nets with great force.

Megalops atlanticus Cuvier & Valenciennes, Hist. Nat. Poiss., XIX, 398, 1846, Guadeloupe, Santo Domingo, Martinique, and Porto Rico; Poey, Fauna Puerto-Riqueña, 343, 1881; Stahl, l. c., 80 and 165, 1883.

Megalops elongatus Girard, Proc. Ac. Nat. Sci. Phila. 1858, 224, Long Island.

Tarpon atlanticus, Jordan & Evermann, l. c., 409, 1896.

Genus 20. *ELOPS* Linnæus.

Body elongate, covered with thin, small, silvery scales. Dorsal fin slightly behind ventrals, its last rays short, the fin depressible into a sheath of scales; anal fin smaller, similarly depressible; pectorals and ventrals moderate, each with a long accessory scale. Opercular bones thin, with expanded, membranaceous borders; a scaly occipital collar. Lateral line straight, its tubes simple. Pseudobranchiæ present, large. Vertebrae $43 + 29 = 72$.

Large fishes of the open seas, remarkable for the development of scaly sheaths. The young are ribbon-shaped and elongate, passing through a series of changes like those seen in *Albula*.

25. *Elops saurus* Linnæus.

"Piojo"; *Matajuelo Real*; *Chiro*; *Lisa Francesa*; *Ten-Pounder*; *John Mariiggle*; *Bony-fish*; *Big-eyed Herring*.

Head 4.3; depth 5 to 6; eye 5; snout 4.3; maxillary 1.6; mandible 1.5; interorbital 5.6; D. 20; A. 13; pectoral 1.8; ventral 2; caudal 0.8; scales 13-110-12.




FIG. 11.—*Elops saurus*.

Body very elongate, moderately compressed, scales small and thin, none on head; head small, pointed; mouth very large, the extremely long maxillary reaching far beyond eye, which has a well-developed adipose eyelid, sheathing the eye anteriorly and posteriorly; rather blunt, villiform teeth on jaws, vomer, and palatines and along lower edge of maxillary; jaws subequal; a pointed gular plate; dorsal and anal fins with well-developed basal sheaths of scales, that of dorsal large; ventral with a very large and pointed accessory scale; caudal lobes long and slender. Blue above, the sides silvery.

This species is abundant and widely distributed in the tropical seas. It is common in America north to the Carolinas and the Gulf of California. Probably not uncommon about Porto Rico, though seen by us only at Arecibo, where a specimen 15 inches long was obtained.

Elops saurus Linnæus, Syst. Nat., ed. XII, 518, 1766, Carolina; Jordan & Evermann, l. c., 410, 1896.

Argentina carolina Linnæus, Syst. Nat., ed. XII, 519, 1766, Carolina.

Argentina machnata Forskål, Descr. Anim., 68, 1775, Djidda, Arabia.

Mugilomorus anna-carolina Lacépède, Hist. Nat. Poiss., V, 398, 2803, South Carolina.

Elops incermis Mitchill, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 445, New York.

Elops capensis Smith, Zool. South Africa, 1845, pl. 7, Cape of Good Hope.

Elops purpurascens Richardson, Ichth. China, 311, 1846, China.

Family XVI. ALBULIDÆ. The Lady-fishes.

Body rather elongate, little compressed, covered with rather small, brilliantly silvery scales; head naked. Snout conic, subquadrangular, shaped like the snout of a pig, and overlapping the small, inferior, horizontal mouth. Maxillary rather strong, short, with a distinct supplemental bone, slipping under the membranaceous edge of the very broad preorbital; premaxillaries short, not protractile. Lateral margin of upper jaw formed by maxillaries; both jaws, vomer, and palatines with bands of villiform teeth; broad patches of coarse, blunt, paved teeth on the tongue behind and on the sphenoid and pterygoid bones. Eye large, median in head, with a bony ridge above it, and almost covered with an annular adipose eyelid. Opercle moderate, firm; preopercle with a broad, flat, membranaceous edge, which extends backward over the base of opercle. Pseudobranchiæ present. Gillrakers short, tubercle-like. Gill-membranes entirely separate, free from isthmus; branchiostegals about 14; a fold

of skin across gill-membranes anteriorly, its posterior free edge crenate; no gular plate. Lateral line present. Belly not carinate, flattish, covered with ordinary scales. Dorsal fin moderate, in front of ventrals, its membranes scaly; no adipose fin; anal very small; caudal widely forked. Pyloric caeca numerous. Parietal bones meeting along top of head. Vertebrae numerous, $42 + 28 = 70$.

A single species known, found in all warm seas. In this, and probably in related families, the young pass through a metamorphosis analogous to that seen in the conger eels. They are for a time elongate, band-shaped, with very small head and loose transparent tissues. From this condition they become gradually shorter and more compact, shrinking from 3 or $3\frac{1}{2}$ inches in length to 2 inches. According to Dr. Gilbert, this process, like that seen in various eels, is a normal one, through which all individuals pass. In the Gulf of California, where these fishes abound, these band-shaped young are often thrown by the waves on the beach in great masses.

Genus 21. ALBULA (Gronow) Bloch & Schneider.

The characters of this genus are included above with those of the family.




FIG. 12.—*Albula vulpes*.

26. *Albula vulpes* (Linnaeus). “*Macabi*”; “*Piojo*”; *Lady-fish*; *Bone-fish*; *Banana-fish*.

Head 3.4; depth 4.5; eye 7; snout 2.2; maxillary 3; interorbital 3.8; preorbital 5.6; D. 15; A. 8; pectoral 1.9; ventral 2.4; caudal 1.1; scales 9–70–6.

Body elongate, robust, the trunk covered with large shining scales with membranous edges; head large, naked, subconic; the snout pig-like, overhanging the inferior mouth, which is armed with villiform teeth; eye high in position, midway between tip of snout and edge of opercle, entirely covered, save for a circular central opening smaller than pupil, with an adipose eyelid; vertical fins scaled, the dorsal and anal very densely; caudal widely forked, upper lobe the longer; median line of back with one series of modified scales, which are smaller and narrower than those of body and have a long membranous appendage much narrower than the scale, this appendage alone exposed.

Color, bright-silvery, darker above; faint longitudinal dark or bluish streaks. A metamorphosis (as in *Elops*) takes place in the young, previous to which they do not resemble the adult.

Tropical seas, on sandy coasts, almost universally distributed and generally abundant, ranging northward on our coasts to San Diego and Massachusetts. Three examples in the collection, one 19 inches long, the others about 12 inches, from San Juan market and Culebra Island. Used as food to some extent, but not highly esteemed.

Umbarana, Maregrave, Hist. Bras., 1648, Brazil.

Vulpes bahamensis (the Bone-fish), Catesby, Nat. Hist. Carolinas, pl. II, fig. 1, 1737, Bahamas.

Esox vulpes Linnaeus, Syst. Nat., ed. X, 1758, 313, Bahamas; after Catesby.

Argentina glossodonta Forskål, Descr. Anim., 68, 1775, Djidda, Arabia.

Macabi, Parra, Dif. Piezas, Cuba, 88, pl. 35, fig. 1, 1787, Cuba; on *Umbarana* of Maregrave.

Synodus argenteus Bloch & Schneider, Syst. Ichth., 398, 1801, Asia.

Clupea brasiliensis Bloch & Schneider, Syst. Ichth., 427, 1801, Brazil.

Albula conorhynchus Bloch & Schneider, Syst. Ichth., 432, 1801, Antilles; after Gronow and Plumier; Poey, Fauna Puerto-Riqueña, 342, 1881; Stahl, l. c., 80 and 165, 1883.

Amia immaculata Bloch & Schneider, Syst. Ichth., 451, 1801, Central America; after Parra.

Butyrinus banana Lacépède, Hist. Nat. Poiss., V, 46, 1803, Isle de France.

- Clupca macrocephala* Lacépède, Hist. Nat. Poiss., V, 426, 1803, Martinique; on a drawing by Plumier.
Glossodus forskåli Agassiz, Spix, Pisc. Bras., 49, 1829, Bahia; called *Engraulis scrieus* and *Engraulis bahiensis* on the plates, 22 and 24.
Albula parra Cuvier & Valenciennes, Hist. Nat. Poiss., XIX, 339, 1846, Martinique; Bahia; Rio de Janeiro.
Albula gorensis Cuvier & Valenciennes, Hist. Nat. Poiss., XIX, 342, 1846, Gorea.
Albula neoguinaica Cuvier & Valenciennes, l. c., XIX, 350, 1846, New Guinea.
Albula seminuda Cuvier & Valenciennes, l. c., XIX, 351, 1846, New Guinea.
Albula erythrocheilos Cuvier & Valenciennes, l. c., XIX, 352, pl. 540, 1846, Friendly Islands.
Albula forsteri Cuvier & Valenciennes, l. c., XIX, 354, 1846, Tahiti.
Albula rostrata Gronow, Cat. Fishes, 189, 1854, American Ocean, etc.
Albula vulpes, Jordan & Evermann, l. c., 411, 1896.

Family XVII. CLUPEIDÆ. The Herrings.

Body oblong or elongate, more or less compressed, covered with cycloid or pectinated scales. Belly sometimes rounded, sometimes compressed, in which case it is often armed with bony serratures. Head naked, usually compressed. Mouth rather large, terminal, jaws about equal, the maxillaries forming the lateral margins of upper jaw, each composed of about three pieces. Premaxillaries not protractile; teeth mostly small, often feeble or wanting, variously arranged. Adipose eyelid present or absent. Gillrakers long and slender; gill-membranes not connected, free from isthmus. No gular plate. Gills 4, a slit behind the fourth. Branchiostegals usually few (6 to 15). Posterior lower part of opercular region often with an angular emargination, the tips of larger branchiostegals abruptly truncate. Pseudobranchiæ present. No lateral line. Dorsal fin median or somewhat posterior, rarely wanting. No adipose fin. Ventrals moderate or small. Anal usually rather long; caudal fin forked. Vertebrae 40 to 56.

The *Clupeidæ* comprise about 30 genera and 150 species, inhabiting all seas, usually swimming in immense schools; many species ascend fresh waters, and some remain there permanently. The northern and fresh-water species, as in many other families, differ from the tropical forms in having a larger number of vertebral segments.

- a. Belly rounded, covered with ordinary scales; supplemental bones of maxillary very narrow; anal fin short.
 b. Ventral small; teeth small, persistent, on jaws, vomer, palatines, pterygoids, and tongue.
 c. Scales of breast not forming a corselet.
 d. Species very small, with teeth minute; a silvery lateral band; dorsal short, of 11 to 16 rays; ventrals inserted nearly under its front..... JENKINSIA, 22
 dd. Species of moderate size, with moderate teeth; no silvery lateral band; dorsal long, of 18 to 20 developed rays; ventrals inserted much behind dorsal, much nearer base of caudal than tip of snout..... ETRUNEUS
 aa. Belly compressed, armed with bony serræ; supplemental bones of maxillary broad.
 e. Anal fin moderate, of 15 to 25 rays; dorsal inserted nearly opposite ventrals.
 f. Scales with their posterior margins entire and rounded; intestinal canal of moderate length.
 g. Last ray of dorsal not produced.
 h. Vertebrae about 50 in number (46 to 56); species of northern regions.
 i. Vomer with teeth; ventral scutes weak, ventrals below middle of dorsal; vertebrae 50 to 56..... CLUPEA
 ii. Vomer without teeth.
 j. Ventral scutes very weak, belly more or less rounded; vertebrae about 52; ventrals under middle of dorsal.
 CLUPANODON, 23
 hh. Vertebrae about 42 (40 to 44); tropical species with scales large and usually firmly attached; ventrals inserted under middle of dorsal; adipose eyelid obsolete..... SARDINELLA, 24
 gg. Last ray of dorsal produced in a long filament; scales large, not firmly attached; otherwise essentially as in *Scrdinella*..... OPISTHONEMA, 25
 ff. Scales with their posterior margins vertical, and pectinate or fluted; head very large; no teeth; intestines elongate; herbivorous..... BREVOORTIA
 PRISTIGASTERINÆ:
 cc. Anal fin very long, of more than 30 rays; dorsal fin inserted behind ventrals.
 k. Teeth not all villiform; both jaws with strong canines; ventrals present, very small..... CHIROCENTRODON
 kk. Teeth all villiform; no canines; ventral fins present..... ILISHA, 26

Genus 22. JENKINSIA Jordan & Evermann.

Very small species, closely allied to *Etruneus*, but with minute teeth and a silvery lateral band; dorsal with fewer than 18 rays; ventrals inserted below or just behind it. Two of the three known species are found in Porto Rico.

- a. Dorsal inserted midway between snout and base of caudal; dorsal rays 14; anal 15..... *lamprotaenia*, 27
 aa. Dorsal inserted nearer snout than base of caudal; dorsal rays 11; anal 17..... *stolifera*, 28

27. Jenkinsia lamprotænia (Gosse).

Head 3.75; depth 5.75; eye 3; snout 3; maxillary 2.3; mandible 2.2; interorbital 5; D. 14; A. 15.

Body long, slender, and compressed; head narrow; snout long, conical, as long as the large eye; mouth large, jaws subequal, maxillary narrow, reaching slightly past anterior border of eye; minute teeth on jaws, vomer, and palatines; origin of dorsal midway between tip of snout and base of caudal, slightly in front of insertion of ventrals. Length 2 to 3 inches.

Pale straw-color; a narrow dark stripe along median line of back, and a broad silvery lateral band nearly as broad as eye.

Known only from Jamaica and Porto Rico. One specimen, 2.5 inches long, obtained.

Clupea lamprotænia Gosse, Nat. Sojourn in Jamaica, 291, pl. 1, fig. 2, 1851, Jamaica.

Jenkinsia lamprotænia, Jordan & Evermann, l. c., 419, 1896.

28. Jenkinsia stolifera (Jordan & Gilbert).

Head 3.75; depth 5.5; eye 2.5; D. 11; A. 17; scales caducous, about 36.

Body elongate, slender, moderately compressed. Snout sharp, tapering; jaws equal; maxillary 2.5 in head, reaching slightly beyond front of eye. Teeth minute, evident in both jaws. Eye large. Dorsal high, inserted at a point slightly nearer snout than base of caudal; ventrals under fourth dorsal ray, nearly half head, and slightly shorter than pectoral.

Translucent green; side with a silvery band as in *Stolephorus*, one-fourth depth of body, a little broader than pupil; a double row of dots along back before dorsal and a single row behind; fins pale.

Found in the Gulf of Mexico from Key West to Yucatan. A small silvery fish very abundant in schools in the surf with *Stolephorus brownii*, a species it much resembles in form and coloration. Length 2 inches. Numerous specimens obtained near the shore of the reefs at Culebra Island. Most of these lack the silvery lateral band, which is replaced by a dark band, this condition appearing to be due to the formalin in which they were first preserved.

Dussumicria stolifera Jordan & Gilbert, Proc. U. S. N. M. 1884 (June 3), 25, Key West.

Jenkinsia stolifera, Jordan & Evermann, l. c., 419, 1896.

Genus 23. CLUPANODON Lacépède. **The Sardines.**

This genus is close to *Clupea*, which it resembles in the elongate form and weak ventral serratures. Vomer toothless; teeth in jaws mostly weak. Scales thin, deciduous. Adipose eyelid present. Gill-rakers very numerous. Species about 6; chiefly confined to the two temperate zones; all closely related to the European sardine, *Clupanodon pilchardus*, and agreeing with it in the rich and delicate flesh; less firm than that of related species, and much richer in oil. Species marine, not anadromous.

29. Clupanodon pseudohispanicus (Poey). *Sardina de España*; "Sardina."

Head 4; depth 3.75 to 4.5; eye 3.75; D. 16; A. 16; scales about 45. Vertebrae 46 to 48.

Body slender, little compressed, belly scarcely carinated, its scutes not prominent; mouth small, maxillary not quite reaching pupil, 2.6 in head; gillrakers very long, slender, and numerous, 30 to 40 below angle, the longest two-thirds eye. Lower jaw with a few feeble teeth; some minute teeth on tongue. Cheek much longer than deep, their depth below eye two-thirds diameter of eye. Adipose eyelid well developed. Opercle with very faint striae, preopercle with very few. Caudal well forked, lower lobe as long as head and a little longer than upper; ventrals inserted nearly below middle of dorsal, a little nearer base of caudal than tip of snout; pectoral 1.33 in head, a conspicuous sheath of scales at base. Intestine 1.5 times length of body.

Color, bluish with no distinct markings, sides golden and silvery; peritoneum dusky; opercle dusky within.

Found in the Gulf of Mexico from Pensacola and Tampa southward to Cuba, Jamaica, and Porto Rico; occasionally taken in numbers at Woods Hole, Mass., whither it is probably carried in the Gulf Stream. Called "sardina" in Porto Rico and "bang" in Jamaica. Length 8 inches. Specimens taken at Culebra Island.

Sardinia pseudohispanica Poey, Mem., II, 311, 1861, Cuba; Poey, Fauna Puerto-Riq., 343, 1881; Stahl, l. c., 80 and 165, 1883.

Clupanodon pseudohispanicus, Jordan & Evermann, l. c., 423, 1896.

Genus 24. **SARDINELLA** Cuvier & Valenciennes. Scaled Sardines.

Small herrings of the tropical seas, with the vertebrae in reduced number, about 40 to 44; scales large, usually firm and adherent, often crossed by vertical striae. Ventral scutes strong, 25 to 35 in number. Adipose eyelid obsolete. Lower jaw projecting; upper jaw somewhat emarginate; teeth weak. Ventrals inserted behind front of dorsal. Body compressed; cheek not deep; gillrakers long and numerous; otherwise essentially as in *Pomolobus*.

The genus *Sardinella*, as here understood, covers a wide diversity of forms and may be divisible into several genera when the anatomy of the species is better known.

- a. Side of body without distinct silvery lateral band; mouth moderate, teeth very small, but permanent over most of bones of mouth; scales large and usually firm.

SARDINELLA:

- b. Ventral scutes 33 to 35.
 c. Body slender, depth about 4.5 in length; a black opercular spot..... *anchovia*
 cc. Body rather deep, depth about 3.5 in length; no black opercular spot..... *clupcola*
 bb. Species imperfectly described.
 d. Snout and chin black..... *apicalis*
 dd. Snout and chin not black; a black humeral spot; side with dark streaks..... *bishopi*

HARENGULA:

- bbb. Ventral scutes 25 to 28; body short and deep, compressed, scales usually with vertical striae.
 e. Scales not very firm and little adherent, so that many are lost in preserved examples; each scale with four vertical wavy striae; ventral scutes about 15 + 10; depth 3.4 in length; eye 2.5 in head; no humeral spot..... *sardina*
 cc. Scales firm and closely adherent so that few, if any, are lost in preserved examples; usually a humeral spot.
 f. Body moderately elongate, ventral outline not strongly arched; depth 3.33 to 3.4 in length.
 g. Head long, 3.43 in length; eye 2.66 in head..... *macrophthalma*, 30
 ff. Body deep, ventral outline arched, forming an even curve from snout to vent; depth 2.75 to 3; eye 2.66 in head..... *humeralis*, 31
 aa. Side with a very distinct lateral silvery band; scales very firm, without vertical striae; mouth very small, almost vertical; teeth small, none on vomer; tip of snout, chin, and upper fins dusky..... *stolifera*

30. *Sardinella macrophthalma* (Ranzani). "Sardina."

Head 3.5; depth 3.4; eye 3; snout 3.4; maxillary 2.1; mandible 1.9; interorbital 4.1; D. 16; A. 17; pectoral 1.5; ventral 2.5; caudal 1.1; scales 40-10; scutes about 28.

Close to *S. humeralis* but differing in shape, chiefly in greater slenderness of body; the ventral outline is not so strongly arched, head not quite so heavy and more pointed, and in spirits the outlines of scales are less distinct. Humeral spot said (Jordan & Evermann) to be usually evident but sometimes wanting. In all our specimens it is absent. A dark, rather indistinct, longitudinal stripe from humeral region to base of upper lobe of caudal separated from the dark of the back by a pale stripe. Otherwise as in *S. humeralis*, with which it is taken.

West Indies, Cuba to Brazil; not common. Nine examples, 2.5 to 6 inches, from San Juan, Puerto Real, Culebra, and Fajardo.

Clupea macrophthalma Ranzani, Nov. Comm. Ac. Sci. Bonon., V, 320, 1842, Brazil.

Harengula maculosa Cuvier & Valenciennes, Hist. Nat. Poiss., XX, 292, 1847, Martinique.

Harengula jaguana Poey, Repertorio, I, 169, 1866, Jagua, near Cienfuegos, Cuba.

Sardinella macrophthalma, Jordan & Evermann, l. c., 430, 1896.

31. *Sardinella humeralis* (Cuvier & Valenciennes). "Sardina"; *Sardina Escamuda*; *White-bill*.

Head 3.5; depth 3; eye 2.7; snout 3.4; maxillary 2; mandible 2; interorbital 4.3; D. 16; A. 17 or 18; pectoral 1.4; ventral 2.2; caudal 1.2; scales 40-10; scutes 27 to 30. Resembling *Opisthonema*, but without the produced ray in dorsal, and with heavier head, larger eye and scales; fine teeth on jaws, palatines, and tongue; eye with a thin adipose eyelid; scales large, firm, with vertical striae.

Blue above, silvery below; a dusky humeral spot; some golden on opercle behind eye; chin and snout dusky.

Very common and much used as food; captured extensively with the cast-net. Eighty-nine examples, 3 to 5.5 inches, collected at San Juan, Palo Seco, Mayaguez, Puerto Real, Boqueron, Arroyo, Ilucares, and Culebra; one from San Geronimo.

Harengula humeralis Cuv. & Val., Hist. Nat. Poiss., XX, 293, 1847, Rio Janeiro; Bahia; Guadeloupe; Santo Domingo.

Alausa striata Cuvier & Valenciennes, Hist. Nat. Poiss., XX, 429, 1847, Guadeloupe; Bahia.

Harengula pensacole Goode & Bean, Proc. U. S. N. M. 1879 (Nov. 5), 152, Pensacola, Fla.

Sardinella humeralis, Jordan & Evermann, l. c., 431, 1896.

Genus 25. *OPISTHONEMA* Gill. Thread-herrings.

Characters essentially those of *Sardinella*, except that the last ray of the dorsal is produced in a long filament as in *Dorosoma*, *Megalops*, and *Tarpon*. Species few, American.

32. *Opisthonema oglinum* (Le Sueur). Thread Herring; *Machuelo*; *Cailleu-Tassart*; *Sprat*.

Head 4.7; depth 3.2; eye 3.6; snout 3.3; maxillary 2.5; mandible 2.1; interorbital 4; D. 17 to 19; A. 23; pectoral 1.4; ventral 2.7; caudal 0.6; scales 50-15; scutes about 31.

Body much compressed below, ventral outline trenchant, much more strongly arched than dorsal; head small; mouth small, the wide maxillary reaching about to front of pupil, the lower jaw barely projecting; opercle with a deep emargination at lower angle; adipose eyelid present; fins small, last dorsal ray greatly produced, reaching base of caudal in younger individuals, somewhat shorter in older ones; pectoral fitting into a shallow depression; ventral small, with an accessory scale nearly as long as fin; anal extremely low; caudal lobes long and slender, subequal; scales large, cycloid, rather loosely attached; lateral line absent.

Color in spirits: Bluish above, silvery below, back with interrupted brownish longitudinal stripes along middle of rows of scales; a dusky shoulder-spot, the young with a series of smaller spots behind, and black-tipped caudal lobes.

West Indies, north to the Carolinas, occasionally to Massachusetts; common; 47 examples, 3.25 to 9 inches, from San Juan Market, Palo Seco, Mayaguez, Guanica, Hucars, Isabel Segunda, and Fajardo; one from San Geronimo.

Clupea thrissa Broussonet, Ichthyologie, fasc. I, 1782, Carolina; Jamaica; not of Osbeck, 1757.

Megalops oglina Le Sueur, Jour. Ae. Nat. Sci. Phila., I, 1817, 359, Newport, R. I.

Megalops notata Le Sueur, Jour. Ac. Nat. Sci. Phila., I, 1817, 359, Guadeloupe.

Chatoëssus signifer De Kay, N. Y. Fauna: Fishes, 264, 1842, New York.

Chatoëssus eumorphus Gosse, Nat. Sojourn in Jamaica, 290, 1851, Jamaica.

Opisthonemus thrissa, Poey, Fauna, Puerto-Riqueña, 343, 1881; Stahl, l. c., 80 and 166, 1883.

Opisthonema oglinum, Jordan & Evermann, l. c., 432, 1896.

Genus 26. *ILISHA* Gray.

Body much compressed, thorax and abdomen strongly serrated. Scales moderate; lower jaw prominent; mouth moderate with rasp-like bands of minute teeth on jaws, palatines, pterygoids, and tongue; none on vomer. Anal fin very long; ventrals present, small, inserted before the small dorsal; upper ray of pectoral strong; caudal deeply forked.

Tropical coasts of America and Asia. Only one species known from Porto Rico.

a. Anal rays 40 to 45; anal placed behind dorsal.

b. Dorsal rays 19 or 20; eye 4 in head..... *flavipinnis*

bb. Dorsal rays 15; eye 3.5 in head..... *bleekeriana*, 33

33. *Ilisha bleekeriana* (Poey). *Manjua*; *Anchoa Pelada*.

Depth 5.66 in length with caudal; eye 3.5; D. 15; A. 43; scutes 25; ventral line strongly curved. Mouth wide, oblique. Anal placed behind dorsal; length of anal equal to its distance from posterior border of eye. Scales very caducous. Silvery.

Matanzas, Cuba; rare. Not obtained by us, but included on the authority of Poey and Stahl.

Pellona bleekeriana Poey, Repertorio, II, 242, 1867, Matanzas, Cuba; Poey, Fauna Puerto-Riqueña, 343, 1881; Stahl, l. c., 79 and 166, 1883.

Ilisha bleekeriana, Jordan & Evermann, l. c., 436, 1896.

Family XVIII. ENGRAULIDIDÆ. The Anchovies.

Body elongate, more or less compressed, covered with thin cycloid scales. Head compressed. Mouth extremely large, more or less oblique, usually overlapped by a pointed, compressed, pig-like snout. Gape very wide, maxillary very long and slender, formed of about three pieces, extending backward far behind eye, in some species beyond head. Premaxillaries not protractile, very small, firmly joined to maxillaries. Teeth usually small, sometimes obsolete, usually fine and even, in a single row

in each jaw; canines sometimes present. Eye large, well forward, without adipose eyelid. Preorbital narrow. Opercles thin and membranaceous. Gillrakers long and slender. Branchiostegals slender, 7 to 14 in number. Gill-membranes separate or joined, free from the isthmus. Pseudobranchiæ present. No lateral line. Belly rounded or weakly serrate. Fins various; dorsal usually short and median; no adipose fin; caudal forked.

Small, carnivorous shore fishes, usually swimming in large schools on sandy shores; abundant in all warm seas, occasionally entering rivers. Only one of the American genera known from Porto Rico.

- a. Teeth in jaws equally small, if present; no canines..... STOLEPHORUS, 27
- aa. Teeth in jaws unequal, some of them enlarged and canine-like..... LYCENGRULIS

Genus 27. STOLEPHORUS Lacépède. Silvery Anchovies.

Body oblong, compressed, covered with rather large, thin, deciduous scales. Belly rounded, or weakly compressed. Snout conical, compressed, projecting beyond the very large mouth. Maxillary narrow, little movable, usually formed of three pieces extending backward far behind eye to base of mandible or beyond, not beyond gill-opening. Premaxillaries very small. Teeth small, subequal, present at all ages, usually on jaws, vomer, palatines, and pterygoids. Anal fin moderate, free from caudal (its rays 12 to 40). No pectoral filaments. Dorsal inserted about midway of body, posterior to ventrals. Pectorals and ventrals each with a large axillary scale. Adipose eyelid obsolete. Vertebrae about 40 (40 to 42) in species examined. Flesh rather pale and dry, more or less translucent; bones firm. Pseudobranchiæ present. Branchiostegals 9 to 14. Gillrakers long and slender. Gill-membranes separate, free from the narrow isthmus.

Species about 50; small, carnivorous shore fishes, swimming in large schools on sandy shores of all warm seas, occasionally entering rivers; usually marked by a very broad, distinct, silvery band.

- a. Anal rays 12 to 14; body very slender, depth 6 in length; no distinct lateral band..... *miarchus*
- aa. Anal rays 15 to 17; body slender, compressed, depth 5 to 6 in length.
- b. Maxillary short, not quite reaching margin of preopercle; lateral band well defined, three-fourths eye. *per fasciatus*, 34
- bb. Maxillary of moderate length, reaching beyond preopercle nearly to gill-opening.
- c. Side with a well-defined lateral silvery band; belly not serrulate.
- d. Eye 4 in head, as long as snout; scales caducous..... *cubanus*, 35
- dd. Eye 3.33 in head, longer than snout; axillary sheaths very large..... *pertheccatus*
- aaa. Anal rays 19 to 24.
- e. Side with a distinct silvery lateral band, with well-defined edges.
- f. Maxillary long, reaching past root of mandible, nearly or quite to gill-opening; snout projecting considerably beyond lower jaw.
- g. Body moderately elongate, depth 4 to 4.75 in length of body.
- h. Silvery lateral band very sharply defined, as broad as eye or slightly narrower, not much narrowed anteriorly; eye large, 3.5 in head; belly serrulate; A. 20; gillrakers two-thirds eye..... *brownii*, 36
- gg. Body more elongate, depth about 5.25 in length of body; gillrakers as long as eye. A. 23 or 24; eye 4.4 in head..... *chirostomus*, 37
- ff. Maxillary short, not reaching root of mandible; eye small, not longer than snout, 4 in head; silvery stripe rather diffuse, half broader than eye; body little compressed, approaching *Engraulis mordax* in form.... *argyrophanus*
- ee. Side without distinct silvery band, or with a faint diffuse streak.
- i. Opercle short, distance from lower posterior angle of cheek to gill-opening much less than from same point forward to middle of eye; snout bluntish, not produced.
- j. Dorsal rays 12 to 14.
- k. Anal rays 18 to 20; depth 6.9..... *lyolepis*, 38
- kk. Anal rays 23; depth 3.3..... *garmani*, 39
- jj. Dorsal rays 15 or 16; anal 23 or 24.
- l. Depth 3.3; snout much projecting..... *gilberti*, 40
- aaaa. Anal rays about 30 (25 to 36).
- m. Silvery lateral band diffuse or obsolete; body much compressed; eye 3 to 3.5 in head.
- n. Gillrakers shorter than eye; lateral band narrow.
- o. Belly slightly serrulate; gillrakers two-thirds eye; scales caducous; anal rays 25 or 26..... *mitchilli*
- mm. Gillrakers long and slender, longer than eye; belly trenchant, not serrate; scales 40; lateral band broad and diffuse or obsolete; snout much projecting.
- p. Snout moderately pointed; minute teeth in both jaws; lateral band diffuse..... *clupcoïdes*
- aaaaa. Anal fin extremely long, its rays 37 or 38. Lateral band ill defined or obsolete; snout pointed, much projecting; insertion of dorsal nearer snout than base of caudal; subopercle with a flat triangular prominence..... *spinifer*
- ll. Depth 4.3; snout little projecting..... *pocyi*
- pp. Snout pointed; minute teeth in upper jaw only; lateral band obsolete..... *productus*, 41

34. *Stolephorus perfasciatus* (Poey).

Head 4; depth 6; eye 3.7; snout 4.4; maxillary 1.5; mandible 1.5; interorbital 4.2; D. 12 to 15; A. 14 to 16; scales about 40,-7. Body very slender, the back wide, the belly compressed to a rather trenchant keel with weak serrulations; snout shorter than eye, pointed, and considerably projecting; top of head with a slight median ridge; maxillary rounded at end, shorter than in the other species, reaching not quite to edge of preopercle; front of anal under last rays of dorsal.

Color much as in *S. brownii*, the lateral band nearly as wide as eye, in spirits its upper edge bordered with dark.

Florida Keys to Cuba, Porto Rico, and Jamaica; common; 15 specimens, 2.75 to 4 inches long, from Aguadilla.

Engraulis perfasciatus Poey, Memorias, II, 313, 1861, Cuba.

Stolephorus perfasciatus, Jordan & Evermann, l. c., 441, 1896.

35. *Stolephorus cubanus* (Poey).

Head 5 in length, with caudal; depth 6.66; eye 4; D. 14; A. 17. Allied to *S. brownii*, but with anal shorter. Body slender, compressed. Eye as long as snout. Maxillary with teeth, its tip extending beyond opercular border. Dorsal beginning midway between front of caudal and posterior edge of eye; pectoral not quite reaching ventral. Scales caducous. A silvery band one-fourth depth of body. Length 2.75 inches. Cuba and Porto Rico. Not seen by us; included on authority of Poey and Stahl.

Engraulis cubanus Poey, Synopsis, 420, 1868, Cuba; Poey, Fauna Puerto-Riqueña, 344, 1881; Stahl, l. c., 80 and 166, 1883.

Stolephorus cubanus, Jordan & Evermann, l. c., 442, 1896.

36. *Stolephorus brownii* (Gmelin). *Striped Anchovy; Manjua.*

Head 3.7; depth 4.9; eye 3.5; snout 4.6; maxillary 1.1; mandible 1.4; interorbital 4.3; D. 13 to 15; A. 20 to 23; pectoral 2; ventral 2.8; caudal 1.6; scales 38,-7. Body slender but little compressed, not elevated, the dorsal outline nearly straight, the belly compressed to a not very trenchant edge; scales deciduous; snout much projecting, shorter than eye, maxillary reaching nearly to gill-opening; teeth long and slender, on both maxillary and lower jaw; dorsal and anal with a low basal sheath; front of dorsal nearer base of caudal than snout.

Translucent in life, with a very distinct and well-defined silvery lateral band, usually narrower than eye; on the caudal peduncle just before the end of the band it is slightly narrower than in the rest of its course.

Cape Cod to Brazil. The most abundant of the American anchovies; abundant in Porto Rico; 33 examples, from Palo Seco, Mayaguez, Puerto Real, Hucares, Isabel Segunda, and Fajardo.

Piquitíngia, Maregrave, Hist. Bras., 159, 1648, Brazil.

Menidia, Browne, Hist. Jamaica, 441, 1756, Jamaica.

Atherina brownii Gmelin, Syst. Nat., 1397, 1788, Jamaica; after Browne.

Esox epsetus Bonnaterra, Tabl. Ichth., 175, 1788, Jamaica; after Browne.

Engraulis lemniscatus Cuvier, Règne Animal, ed. 2, vol. 2, 323, 1829, Brazil; after *Piquitíngia* of Maregrave.

Engraulis tricolor Agassiz, Pisc. Brasil., 51, 1829, Bahia; Pará.

Engraulis piquitíngia Agassiz, Pisc. Brasil., pl. 23, fig. 1, 1829, Bahia; Pará; same type as *tricolor*.

Argentina menidia Gronow, Cat., 141, 1854, Jamaica; after Browne.

Stolephorus hiuleus Goode & Bean, Proc. U. S. N. M. 1879 (March 25, 1880), 343, Clearwater Harbor, Fla.

Stolephorus brownii, Jordan & Evermann, l. c., 443, 1896.

37. *Stolephorus chærostomus* (Goode). *Hog-mouth Fry.*

Head 3.5; depth 5.3; eye 4.6; snout 4.6; D. 13; A. 23; scales 38. Snout projecting much beyond lower jaw, which just passes vertical from front of eye; eye as long as snout. Maxillary tapering, reaching gill-opening. Gillrakers 10 + 25, as long as eye. Dorsal inserted before middle of body; anal under middle of body; pectoral reaching front of ventrals; scales large.

Brownish, with a lateral silvery band, as broad as eye.

This species reaches a length of 2.75 inches. Heretofore known only from Hamilton Harbor, Bermuda Islands. Apparently common about Porto Rico, specimens having been obtained by us at Hucares, Puerto Real, and Fajardo.

Engraulis chærostomus Goode, Amer. Jour. Sci. Arts, August, 1874, 125, Bermudas.



Stolephorus chærostomus, Jordan & Evermann, l. c., 444, 1896.

38. *Stolephorus lyolepis* Evermann & Marsh, new species.

Head 4.3; depth 6.9; eye 4.6; snout 4.2; maxillary 1.4; mandible 1.4; interorbital 4.4; D. 12 to 14; A. 18 to 20. Body slender and compressed, the dorsal and ventral outlines nearly alike, little arched; caudal peduncle gently tapering; no serræ evident; snout not greatly produced; maxillary with free end rounded, reaching beyond root of mandible to edge of preopercle, armed with fine teeth; lower jaw with minute teeth, smaller than those of maxillary; anal inserted under last dorsal ray; scales deciduous.

Color in spirits: Flesh-color, without silvery band, a lateral longitudinal row of small black pigment dots, which are stellate under a lens, usually more numerous posteriorly; a single row of several larger, circular black spots across opercle, and a number scattered on top of head; a row of black spots at base of anal and a few small black blotches about base of caudal rays; peritoneum with rows of black dots, these showing faintly through the flesh on the side near ventral outline between pectoral and ventral.

This species is related to *S. curtus* and *S. poeyi*, both Pacific coast species, from each of which it differs apparently in the fewer anal rays. Thirty-seven specimens of nearly uniform size were collected at Culebra, February 10, of which the type (No. 49528, U. S. N. M.) is 1.5 inches long.

FIG. 13.—*Stolephorus lyolepis*.FIG. 14.—*Stolephorus garmani*.39. *Stolephorus garmani* Evermann & Marsh.

Head 3.2; depth 3.3; eye 3.5; snout 5.5; maxillary 1.7; mandible 1.7; interorbital 5; D. 14; A. 23; pectoral 2; ventral 3.5; caudal 1.3; scales 42,-9.

Body comparatively deep and strongly compressed; belly not strongly trenchant, without serrulations; snout thick, much projecting; maxillary reaching nearly to root of mandible, very finely and weakly serrate; eye large; tip of lower jaw reaching vertical from front of eye; distance from lower posterior angle of cheek to vertical from posterior margin of opercle much less than from same point to eye; dorsal inserted far in advance of anal, just behind insertion of ventrals, midway between anterior edge of pupil and base of caudal.

Color in spirits: Back dark to near the median line, below this somewhat reddish; rest of body

below a line from shoulder to upper base of caudal silvery; some golden on snout and behind eye; no lateral band.

This species has a general resemblance to *Stolephorus productus*, but is unquestionably distinct from it; the anal is much shorter and inserted farther back, the body is deeper, the eye larger, and the snout longer. It is very close to *Stolephorus gilberti*, differing chiefly in the larger eye, in the color of back, and in the somewhat less sharply compressed belly. One specimen, the type (No. 49360, U. S. N. M.), 4.5 inches long, collected at Puerto Real, January 27, 1899.

Stolephorus garmani Evermann & Marsh, Report U. S. F. C. 1899 (December 19, 1899), 352, Puerto Real, Porto Rico.

40. *Stolephorus gilberti* Evermann & Marsh.

Head 3.25; depth 3.4; eye 4; snout 6; maxillary 1.7; mandible 1.7; interorbital 4.9; D. 15; A. 23; pectoral 2.1; ventral 3.5; caudal 1.3; scales 42-9.

Body comparatively deep and strongly compressed, belly trenchant, without serrations; snout thick, much projecting; maxillary reaching nearly to root of mandible, scarcely serrate; eye moderate; tip of lower jaw reaching vertical from front of eye; distance from lower posterior angle of cheek to vertical from posterior margin of opercle much less than from same point to eye; dorsal inserted far in advance of anal, just behind insertion of ventrals, midway between anterior edge of eye and base of caudal.




FIG. 15.—*Stolephorus gilberti*.

Color in spirits: Back light-olivaceous with dark punctulations; rest of body below a line from shoulder to upper base of caudal, silvery; faint traces of golden behind eye; no lateral band.

This species is very close to *Stolephorus garmani*, differing chiefly in the much smaller eye, the more uniform color of the back, the somewhat more sharply compressed belly, and the more nearly entire maxillary. One specimen, the type (No. 49359, U. S. N. M.), 4.5 inches long, collected at Palo Seco, near San Juan, January 13, 1899, associated with *S. productus*, with which species both *S. gilberti* and *S. garmani* Evermann & Marsh are allied.

Stolephorus gilberti Evermann & Marsh, Report U. S. F. C. 1899 (December 19), 352, Palo Seco, Porto Rico.

41. *Stolephorus productus* (Poey). *Hechudo*; *Grubber Broad-head*.

Head 3.6; depth 3.7; eye 3.9; snout 7; maxillary 1.3; mandible 1.4; interorbital 4.6; D. 13; A. 31 to 33; pectoral 1.75; ventral 3.3; caudal 1.1; scales 43-8.

Body elongate, not elevated, strongly compressed, belly not rounded; head small, broadest above, the pointed snout projecting beyond mouth; eye large, much greater than length of snout, placed high and well forward; mouth very wide, maxillary very long and thin, reaching to or beyond root of mandible nearly to gill-opening, ending in a point; dentition very weak, edge of maxillary being minutely serrate, no teeth in lower jaw; fins small, pectoral inserted under gill-opening, ventral very small, anal long and low, caudal deeply forked; dorsal and anal with a basal sheath of scales; no lateral line; scales large, somewhat deciduous.

Silvery on sides, the back darker, no lateral band.

Known only from Cuba, Jamaica, and Porto Rico. One of the larger anchovies; rather common; 16 specimens of about 6 inches, from Palo Seco and Ponce.

Engraulis productus Poey, Repertorio, I, 380, 1866, Cuba.

Stolephorus productus, Jordan & Evermann, l. c., 447, 1896.

Family XIX. SYNODONTIDÆ. The Lizard-fishes.

Body oblong or elongate, little compressed, covered with cycloid scales, rarely naked. Mouth very wide, the entire margin of the upper jaw formed by the long and slender premaxillaries, closely adherent to which are the slender maxillaries, the latter mostly rudimental or obsolete, never widened at tip. Teeth mostly cardiform on both jaws, tongue, and palatines; canines rarely present; large teeth usually depressible. No barbels. Opercular bones usually thin, but complete. Gill-membranes separate, free from the isthmus. Branchiostegals usually numerous. Pseudobranchiæ present. Gill-rakers tubercular or obsolete. Lateral line present. Adipose fin present, rarely obsolete; dorsal fin short, of soft rays only; pectorals and ventrals present; anal fin moderate or long; caudal forked. Skeleton rather well ossified. Air-bladder small or wanting. Intestinal canal short. Sides sometimes with phosphorescent spots or photophores. Eggs inclosed in the sacs of the ovary and extruded through an oviduct.

Genera about 10; species about 40, mostly inhabiting shore waters, some of them descending to the depths.

a. Scales present, more or less adherent.

b. Teeth of premaxillary simple, compressed, not barbed, in one or two rows; a broad band of similar teeth on palate.

c. Vent about midway between base of caudal and axil of pectoral; head short, blunt, compressed.

TRACHINOCEPHALUS, 28

cc. Vent much nearer base of caudal than axil of pectoral; head depressed, with flat triangular snout . . . SYNODUS, 29

bb. Teeth of premaxillary in a very broad band, curved, unequal, and barbed at the end; a similar band on palatines.

BATHYSAURUS

aa. Scales very caducous or wanting; teeth in narrow bands; vent posterior BATHYLACO

Genus 28. TRACHINOCEPHALUS Gill.

This genus is closely related to *Synodus*, from which it differs chiefly in form and in the relative development of the fins. Body stout, head short, blunt, and compressed. Vent well formed, about midway between base of caudal and axil of pectoral, under tip of last dorsal ray. Teeth as in *Synodus*, but slender, smaller, and closely set. Lower jaw projecting.

Species few; shore fishes, widely diffused in tropical seas.

42. *Trachinocephalus myops* (Forster). *Ground Spearing; Lagarto.*

Head 3.4; depth 5; eye 6.4; maxillary 1.8; mandible 1.5; interorbital 10; D. 12; A. 14; pectoral 2.6; ventral 1.1; caudal 1.6; scales 4-58-7. Vertebrae 58; caeca 25. Snout very short, shorter than eye, lower jaw very slightly projecting; top of head scaleless, very rough; interorbital space with an abrupt longitudinal depression. Pectoral very short, ventrals long, reaching vent.

Color in spirits: Very light brown; 3 or 4 rather faint longitudinal lines above lateral line, these demarcating color stripes in life; traces of longitudinal dark lines below lateral line; back with 6 or 8 obscure dark vertical bars, barely crossing lateral line; an oblong dark-brown scapular blotch; a dark-brown band from eye across lower jaw; fins nearly plain.

One specimen, 7 inches long, from San Geronimo, collected by Mr. Geo. M. Gray.

Salmo myops Forster MS., Bloch & Schneider, Syst. Ichth., 421, 1801, St. Helena.

Osmerus lemniscatus Lacépède, Hist. Nat. Poiss., V, 236, 1803, Martinique; after Plmmer.

Saurus truncatus Agassiz, Pisc. Brasil., 82, 1829, Brazil.

? *Salmo trachinus* Schlegel, Fauna Japonica, Poissons, 231, 1842, Japan; the East Indian, Chinese, and Japanese form, *Trachinocephalus limbatus*, is little if at all different from *T. myops*.

Saurus brevirestris Poey, Memorias, II, 305, 1861, Cuba; erroneously stated to have 10 rays.

Trachinocephalus myops, Jordan & Evermann, l. c., 533, 1896.

Genus 29. *SYNODUS* (Gronow) Bloch & Schneider.

First superior pharyngeals cartilaginous; second without teeth; third and fourth separate, with teeth; lower pharyngeals separate. Body elongate, subterete. Head depressed, snout triangular, rather pointed. Interorbital region transversely concave. Mouth very wide; premaxillaries not protractile, very long and strong, more than half length of head; maxillaries closely connected with them, very small or obsolete; premaxillaries with one or two series of large, compressed, knife-shaped teeth, the inner and larger depressible; palatine teeth similar, smaller, in a single broad band; lower jaw with a band of rather large teeth, the inner and larger teeth depressible; a patch of strong depressible teeth on tongue in front, and a long row along hyoid bone; jaws nearly equal in front. Eye rather large, anterior; supraorbital forming a projection above eye. Pseudobranchiæ well developed. Gillrakers very small, spine-like. Gill-membranes slightly connected. Top of head naked; cheek and opercles scaled like body; body covered with rather small, adherent, cycloid scales; lateral line present; no luminous spots. Dorsal fin short, rather anterior; pectoral moderate, inserted high; ventrals anterior, not far behind pectoral, large, inner rays longer than outer; anal short; caudal narrow, forked. Vent posterior, much nearer base of caudal than axil of pectorals. Branchiostegals 12 to 16. Stomach with a long, blind sac and many pyloric cæca. Skeleton rather firm.

Voracious fishes of moderate size, inhabiting sandy bottoms at no great depth, in most warm seas. Species numerous; two known from Porto Rico.

- a. Scales large, 43 to 50 in lateral line, which has a blunt keel posteriorly.
 b. Tips of first rays of dorsal not reaching tips of last rays when depressed; snout short, broad; shoulder-girdle with a large black blotch, anal rays 10 or 11; scales 4-48-6. *intermedius*, 43
 bb. Tips of first rays of dorsal reaching tips of last rays when depressed.
 c. Shoulder-girdle with the black spot very small or wanting; snout rather pointed, 3.75 in head; ventrals 1.25 in head; anal rays 10 to 12. *poeyi*
 aa. Scales small, 58 to 68 in lateral line.
 d. Anal fin very short, its rays 8 only; pectoral fin short.
 e. Scales moderate, 58 in lateral line; snout 4 in head; shoulder-girdle slightly dusky; dorsal mottled. *synodus*
 dd. Anal fin moderate, its rays 10 to 13; shoulder-girdle chiefly yellowish.
 f. Snout very broad, broader than long; about 10 scales in a cross series from dorsal to ventral; jaws subequal; scales 4-60 to 64-5; tail with a slight keel. *saurus*
 ff. Snout not broader than long; more than 10 scales in a cross series from dorsal to ventral; lower jaw included; tail not keeled.
 g. About 4 rows of scales (6 counting obliquely) between lateral line and adipose fin; scales on cheek in 4 to 7 rows.
 h. Head vermiculate above, its length 4 to 4.25 in body; 7 rows of scales on cheek. *fætens*, 44

43. *Synodus intermedius* (Agassiz). *Sand Diver*.

Head 3.8; depth 7; eye 7; snout 3.8; premaxillary 1.6; mandible 1.4; interorbital 5.8; D. 11; A. 11; pectoral 2.1; ventral 1.1; caudal 1.7; scales 5-49-6. Body elongate, fusiform, slightly depressed, a slight keel on caudal peduncle; head rather blunt and heavy; scales behind eye, with tubes or muciferous canals; tips of first dorsal rays not reaching tips of last rays when depressed. Easily distinguishable from *S. fætens* by the less slender body, heavier head, larger scales, and color markings.

Color in spirits: Grayish, paler below, each row of scales with a pale longitudinal stripe, plainest on sides; about eight faint dark vertical bars, terminating at middle of side, a dark blotch between their ventral ends, plainest in the young; pectoral, caudal, and dorsal barred; shoulder-girdle with a black blotch near angle of opercle, which covers it.

This species is generally common and ranges from southern Florida to Brazil. It is apparently the most abundant species of *Synodus* in Porto Rico, but of little value as food. Five examples, from 7 to 12 inches in length, were collected at Puerto Real, Boqueron, and Culebra.

Saurus intermedius Agassiz, in Spix, *Piscium Brasil.*, 81, pl. XLIV, 1829, Brazil.

Saurus anolis Cuvier & Valenciennes, *Hist. Nat. Poiss.*, XXII, 483, 1849, Bahia; Martinique.

Synodus intermedius, Jordan & Evermann, l. c., 535, 1896.

44. *Synodus fætens* (Linnæus). *Lizard-fish; Galliwasp; Lagarto; Soap-fish.*

Head 4; depth 8; eye 8; snout 3.4; premaxillary 1.6; mandible 1.4; interorbital 5.5; D. 10; A. 12; pectorals 2.1; ventral 1.3; caudal 1.8; scales 6-60-7.

Body very slender, the caudal peduncle without trace of keel; interorbital space with radiating ridges, plainest in adult; opercles with 5 rows of scales, cheek with 7 rows.

Color in spirits: Grayish, obscurely mottled with darker on back, plain white below; no distinct longitudinal lines in adult; the young have the back more distinctly mottled, sometimes with regular bars, with diffuse regular blotches along lateral line and with more or less distinct longitudinal stripes; fins plain.

Three examples, 5.5 to 14 inches in length, from Puerto Real, Boqueron, and Isabel Segunda, and two young from Fish Hawk station 6063 in Mayaguez Harbor, in 75 fathoms.

Salmo falcens Linnæus, Syst. Nat., ed. XII, 513, 1766, South Carolina.
Osmerus albidus Lacépède, Hist. Nat. Poiss., V, 229, 1803, Carolina; after Linnæus.
Coregonus ruber Lacépède, Hist. Nat. Poiss., V, 263, 1803, Martinique; after Plumier.
Esox salmoncus Mitchill, Trans. Lit. and Phil. Soc., I, 1815, 442, New York.
Saurus longirostris Agassiz, Spix, Pisc. Brasil., pl. 43, 1829, Brazil.
Saurus mexicanus Cuvier, Règne Animal, ed. II, vol. 2, 314, 1829, Mexico.
Saurus spicianus Poey, Memorias, II, 304, 1861, Cuba.
Synodus falcens, Jordan & Evermann, l. c., 538, 1896.




FIG. 16.—*Synodus falcens*.

Family XX. AULOPIDÆ.

Allied to the *Synodontidæ*, but with the maxillary separate, well developed, and dilated behind. Hypocoracoids extended downward, as in many spiny-rayed fishes. Gillrakers mostly long and slender, needle-shaped. Eyes normal, large or small. No luminous spots; jaws without fang-like teeth. Dorsal fin moderate, nearly median in position; body elongate. Pectorals present, normal in form and position; adipose fin normally present. Pseudobranchiæ present.

Fishes of moderate depths, chiefly Atlantic, including, as here understood, about six species.

Genus 30. CHLOROPHTHALMUS Bonaparte.

Head elongate, body subterete, covered with moderate-sized, adherent, pectinate, or ctenoid scales arranged in straight, parallel, oblique lines. Mouth rather large, maxillary well developed, dilated behind, reaching to beyond front of orbit; lower jaw projecting. Teeth very small, sharp on jaws, vomer, and palatines, usually minute teeth on tongue. Eye very large. Dorsal short, inserted before middle of length of body; adipose fin small; anal short; caudal forked; pectorals and ventrals well developed, ventrals inserted under the dorsal and not far behind pectorals, none of the rays forming exerted filaments. Gill-openings wide. Branchiostegals 10. Pseudobranchiæ well developed. Gillrakers needle-shaped, rather numerous.

Color, silvery, with darker markings.

Deep-sea fishes, resembling smelt. Of four known species, only one is from Porto Rico.

- a. Eye 2 in head; scales 60 to 63..... *agassizii*
- aa. Eye 3 in head; scales 45 to 52.....
- b. Dorsal rays 11; depth 6.25 in length..... *chalybeius*, 44
- bb. Dorsal rays 8; depth 5.5 in length..... *truculentus*

45. *Chlorophthalmus chalybeius* (Goode).

Head 3; depth 6; eye 2.5; snout 3.5; interorbital 7; maxillary 2; mandible 1.8; D. 9; A. 6; scales 51; pectoral 1.5; ventrals 1.6; height of dorsal 1.4; of anal 2.5; length of caudal lobes 1.4; least depth of caudal peduncle about half that of body.

Body terete; head rather large; mouth large; maxillary long, broad at tip, reaching front of pupil; lower jaw somewhat projecting; eye very large; minute teeth on jaws, vomer, palatines, and tongue;

gillrakers long and slender; opercular flap long, reaching base of pectoral. Scales large, arranged in regular, oblique, transverse rows, overlapping in such a manner as to resemble plates, most of the scales apparently cycloid, but those of lateral line pectinate. Origin of caudal slightly nearer tip of snout than adipose fin and in front of base of ventrals; adipose fin over middle of anal whose distance from snout is three-fourths length of body; pectoral long, pointed, its tip not reaching tips of ventrals; ventrals long; anal opening between ventrals and near their base.

Color in alcohol: Pale; back and sides crossed by about 10 rather broad dark-brown bars, usually interrupted on the side, the broadest being just back of dorsal; a broad one crossing front of dorsal, and a narrow one at its last rays; a large quadrangular dark blotch on side above and anterior to anal fin; base of caudal dark; cheek, opercles, and top of head with some dark; snout pale; lower part of side, especially between pectoral and ventral, with fine black specks; a few above pectoral; breast and region between ventrals with numerous shining black specks; anal black; scales of belly with rows of similar black specks; bases of anal and ventrals similarly marked; inner ventral rays black; lower part of anterior dorsal rays black; base of caudal with numerous dark specks.

The single specimen (2.5 inches long) which we have agrees fairly well with the type with which we have compared it. The lower jaw, however, is less projecting in our specimen and the dorsal rays are one fewer. The known specimens of this interesting species have been taken as follows:

Fish Hawk Station.	U.S. Nat. Mus. No.	Lat. N.	Long. W.	Depth.	Albatross Station.	U.S. Nat. Mus. No.	Lat. N.	Long. W.	Depth.
		° ' "	° ' "	Fms.			° ' "	° ' "	Fms.
876	26092	Off Block	Island.	126	2420.....	43829	37 03 20	74 31 40	104
877					2424.....	43830	36 41 37	74 42 15	85
878					2425.....	43831	36 20 24	74 46 30	119
1043.....	28995	38 39 00	73 11 00	130	2536.....	43833	39 56 15	70 47 30	157
1038.....	28976	39 58 00	70 06 00	146	43834				
1108.....	31644	40 02 00	70 37 30	101	2537.....	43835	39 56 45	70 50 30	156
6070.....		Mayaguez Harbor.		220					

It will be observed that all the specimens previously taken were obtained in the Gulf Stream southeast of Rhode Island, in depths ranging from 85 to 157 fathoms. The specimen obtained by us in Mayaguez Harbor was taken from rocky bottom in the beam trawl and at the greatest depth yet known for the species.

Hyphalonedrus chalybeius Goode, Proc. U. S. N. M. 1880 (Feb. 16, 1881), 484, Gulf Stream, off Block Island.
Chlorophthalmus chalybeius, Jordan & Evermann, l. c., 542, 1896.

Family XXI. PECILIIDÆ. The Killi-fishes.

Body oblong or moderately elongate, compressed behind, depressed forward, covered with rather large cycloid scales, which are adherent and regularly arranged. Lateral line wanting or represented by a few imperfect pores. Head scaly, at least above. Mouth terminal, small, the lower jaw usually projecting; margin of the upper jaw formed by the premaxillaries only; premaxillaries strong, extremely protractile. Teeth incisor-like or villiform, sometimes present on vomer, but usually in jaws only; lower pharyngeals separate, with cardiform or rarely molar teeth; third upper pharyngeal enlarged, fourth wanting or united to third. Gill-membranes somewhat connected, free from isthmus; gillrakers very short, thick. Branchiostegals 4 to 6. Pseudobranchiæ none. Dorsal fin single, inserted posteriorly, of soft rays only, rarely with a single spine or a rudimentary spinous dorsal; caudal fin not forked; ventral fins abdominal, rarely wanting; pectoral fins inserted low; no adipose fin. Stomach siphonal without pyloric appendages. Air-bladder simple, often wanting. Basis cranii simple (*vide* Cope). Sexes usually unlike, the fins being largest in the males, but in some species the females are much larger in size. Many of the species are ovoviviparous or viviparous, the sexes very unlike, anal fin of male being developed as an intromittent organ, the young well developed at birth.

A large family of brackish or fresh-water fishes in southern Europe, Asia, Africa, and America, some of them occurring in bays and arms of the sea. They are mostly of small size, and the species are very difficult of determination. Only two species known from Porto Rico. This is easily understood when it is remembered that there is so little brackish water about this island.

- a. Intestinal canal comparatively short, little convoluted; teeth little movable; bones of the dentary firmly connected; lower jaw strong and usually projecting beyond upper; species chiefly carnivorous.
- b. Anal fin of male similar to that of female and not modified into an intromittent organ. Species oviparous.
- c. Teeth all pointed, none of them compressed or bicuspid or tricuspid.
- FUNDULINÆ:**
- d. Teeth in villiform bands or at least in more than one series.
- e. Air-bladder well developed (in all species examined); no caudal ocellus.
- f. Gill-openings not restricted above, the opercular angle free from shoulder-girdle; body oblong; dorsal various in size and insertion FUNDULUS, 31
- ff. Gill-openings restricted, the opercle from upper root of pectoral upward being adnate to shoulder-girdle, body short and deep ADINIA
- ec. Air-bladder wanting; a black ocellus at root of caudal, at least in males. RIVULUS
- dd. Teeth arranged in a single series; dorsal inserted in advance of anal; mouth oblique.
- g. Dorsal and anal fins short, each of 9 to 13 rays. LUCANIA
- gg. Dorsal and anal fins very long, each of more than 20 rays. GIRARDINICHTHYS
- CYPRINODONTINÆ:**
- cc. Teeth incisor-like, notched, bicuspid or tricuspid.
- h. Ventral fins well developed (occasionally lost through atrophy); gill-openings restricted; opercle adnate above to the shoulder-girdle.
- i. Incisors bicuspid, with a band of villiform teeth behind them; body rather deep, vertical fins moderately developed CHARACODON
- ii. Incisors tricuspid, in one row, with no villiform teeth behind them; body short and deep, compressed.
- j. Dorsal fin short, of 10 to 12 rays, first ray slender and rudimentary CYPRINODON
- jj. Dorsal fin very long, of 16 to 18 rays, first ray developed as a stout, grooved spine. JORDANELLA
- bb. Anal fin in the males placed well forward and modified into a sword-shaped intromittent organ teeth all pointed, arranged in bands. Species viviparous or ovoviviparous, the young of large size at birth.
- GAMBUSINÆ:**
- k. Eye normal, pupil not divided by a partition; dorsal inserted more or less behind front of anal.
- l. Jaws not produced into a beak, lower jaw prominent, longer than upper; male fish much smaller than female.
- m. Dorsal fin long, of 14 to 16 rays; anal short PSEUDOXIPHOPHORUS
- mm. Dorsal fin short, of 6 to 10 rays; anal short. GAMBUSIA
- ll. Jaws produced into a moderate beak, much as in *Labidesthes*; dorsal and anal short. BELONESOX
- ANABLEPINÆ:**
- kk. Eye divided into two portions by a horizontal cross partition; vertical fins short; body elongate ANABLEPS
- aa. Intestinal canal elongate, with numerous convolutions; dentary bones loosely joined; teeth movable; species chiefly mud-eating.
- GOODEINÆ:**
- n. Teeth incisor-like, all tricuspid, in one series, with a series of villiform teeth behind them; sexual characters unknown, the sexes probably alike; lower jaw projecting.
- o. Fins small; scales large; form oblong GOODEA
- PECILINÆ:**
- nn. Teeth all pointed; anal fin in male advanced and modified into an intromittent organ; lower jaw short and weak. Species ovoviviparous.
- p. Teeth in a single series; dorsal and anal both short; scales large.
- q. Dorsal fin inserted in advance of anal. PLATYPECILUS
- qq. Dorsal fin inserted more or less behind front of anal. HETERANDRIA
- pp. Teeth in more than one series.
- r. Dorsal inserted more or less behind anal; both fins very small. LEBISTES
- rr. Dorsal inserted over or in advance of anal, its rays much elevated in male.
- s. Dorsal fin short, of fewer than 12 rays.
- t. Teeth of inner series in both jaws trifid ACROPECILIA
- tt. Teeth of inner series in both jaws entire PECILIA, 32
- ss. Dorsal fin long, of 12 to 16 rays.
- u. Caudal fin normal, alike in both sexes, or with the lower angle merely sharp in the male MOLIENISIA
- uu. Caudal fin in males with its lower lobe much produced and sword-shaped, in adult as long as the rest of the body. XIPHOPHORUS

Genus 31. FUNDULUS Lacépède. Killi-fishes.

Body rather elongate, little elevated, compressed behind. Mouth moderate, lower jaw projecting. Jaws each with two or more series of pointed teeth, usually forming a narrow band. Bones of the mandible firmly united. Scales moderate. Gill-opening not restricted above, opercle with its margin not adnate to shoulder-girdle. Preopercle, preorbital, and mandible with mucous pores. Dorsal and anal fins similar, small or rather large, dorsal inserted either in front of, above, or behind front of anal; ventrals well developed. Air-bladder present. Sexes differing in color, size, and development

of fins, anal fin in the male normal. Intestinal canal short. First superior pharyngeal without teeth, second with teeth; third and fourth coossified, with teeth.

Species very numerous, mostly American, inhabiting fresh waters and arms of the sea. They are the largest in size of the cyprinodonts, and some of them are very brightly colored. They are oviparous and feed chiefly on animals. Some of them are bottom fishes, burying themselves in the mud of estuaries; others swim freely in river channels and bays; still others are "top minnows," surface swimmers, feeding on floating insects in swamps and streams.

- a. Species with dorsal fin moderate or rather large, of 11 to 17 rays, its insertion above or usually in front of insertion of first ray of anal; scales large or small. Free-swimming species, not feeding at the surface, some of them often burying themselves in the mud of bottoms in shallow water.
- b. Dorsal fin inserted before origin of anal; branchiostegals 5 or 6.

FUNDULUS:

- c. Scales large, 31 to 38 in a lengthwise series.
- d. Scales in lateral line 31 to 34, 10 to 12 in a cross series; body rather elongate, depth 4 to 4.5 in length; dorsal rays usually 12.
- e. Body without crossbands; each scale above, and especially posteriorly, with a vertical purplish spot in the center; dorsal with series of blackish dots; anal rays 13 *punctatus*
- ee. Body with crossbands light or dark, probably in both sexes; anal rays 9 to 11.
- f. Snout shortish, shorter than eye; a black spot on back before dorsal; head bluntish, 3.5 in length *pallidus*
- ff. Snout very long, 1.5 times length of eye in adult; no black spot before dorsal; male with an ocellated dorsal spot; head pointed and elongate, 3.33 in body; branchiostegals 6 *similis*
- dd. Scales in lateral line 35 to 38.
- g. Anal rays 10 to 12.
- h. Female with two or three black horizontal stripes; male with about 12 dark crossbars and a dorsal ocellus; head long; 3.75 in length; branchiostegals 6 *majalis*
- hh. Female plain or with dark crossbands only; no black horizontal stripes.
- i. Dorsal rays 10 or 11.
- j. Color greenish; both sexes usually with dusky or silvery crossbars and pearly spots, at least on the fins of the male; dorsal sometimes with an ocellus; scales 35 or 36.
- k. Form robust, the depth 3.66 in length; head 3.66; dorsal ocellus faint or wanting; males with many pearly spots; females nearly plain *heteroclitus*
- kk. Form rather slender, the depth 4 in length; head 3.25; dorsal ocellus on female very conspicuous; pearly spots fewer; female banded or spotted *ocellaris*
- jj. Color greenish; in spirits without bands or spots; body deep; tail slender; fins small *fonticola*, 46
- ii. Dorsal rays 12 to 14; olivaceous, with about 15 dark crossbands *bermuda*
- gg. Anal rays 16 or 17; dorsal rays 13 or 14; upper lip thick; scales 38-15 or 16.
- l. Form robust, the depth in adult about 3 in length; color uniform pale-brown; fins unspotted *robustus*
- ll. Form rather slender, the depth 4.5 in length; color uniform-brown, or slightly mottled on tail; fins unspotted; the anal of male black at base, yellow distally *labialis*
- cc. Scales comparatively small, 44 to 48 in lateral line; dorsal fin of 13 to 15 rays; anal rays 11 to 14 *alinia*
- bb. Dorsal fin inserted over or slightly behind front of anal fin; branchiostegals 4 or 5; bright-colored species with orange or brown spots, inhabiting mountain springs and brooks; scales about 35, 24 before dorsal; body with about 15 well-marked black crossbands, as in *Fundulus similis* or *F. cingulatus*, between which this species seems to find its place; fins plain. D. 9 or 10; A. 9 *funduloides*

ZYGONECTES:

- aa. Species with dorsal fin small, of 7 to 11 rays, its insertion distinctly behind front of anal fin; small species with large scales (29 to 40); surface swimmers, "top minnows," seeking insects at the surface of water.
- m. Anal rays 14; dorsal rays 8; scales 31-8; depth 5 in length; olive, the body plain; dorsal and anal with dark crossbands; base of caudal with round pale spots *dorii*
- mm. Anal rays 8 to 13.
- n. General coloration olivaceous, either plain or with pearly or orange spots paler than ground-color, or blackish spots not forming distinct series; no sharply defined black crossbars or longitudinal black stripes.
- o. Anal rays 12 or 13; body slender, depth 4 to 5 in length *jenkinsi*
- oo. Anal rays 8 to 11; scales rather large, about 33 to 35-10; depth about 4 in length; anal rays 8 to 11; head 3.25 to 3.5 in length *pulvcreus*
- nn. General coloration olivaceous, with a single black lateral stripe from head to tail; body stout, depth 3.33 in length; sides silvery, with black lateral shade above silvery part. D. 11; A. 11; scales 31-10 *melapleurus*

46. *Fundulus fonticola* Cuvier & Valenciennes.

D. 11; A. 12; B. 5; scales 37. Body plump, with long caudal peduncle. Head broad, little depressed; tail slenderer and body deeper than in *Fundulus heteroclitus*; dorsal inserted in front of anal; dorsal and caudal small and rounded, anal high and pointed, paired fins short. Teeth in broad bands, outer little enlarged. Uniform green, apparently without spots or band in spirits. (Color entirely lost in the original type.) Length 2 inches.

Known only from mountain springs in Porto Rico; here described from the original type, the only known example, as the other specimens possessed by Cuvier & Valenciennes belong to a species of *Gambusia*. This species was not obtained by us, but is recorded from Porto Rico by Poey.

Fundulus fonticola Cuvier & Valenciennes, Hist. Nat. Poiss., XVIII, 198, 1846, Porto Rico; Poey, Fauna Puerto-Riqueña, 342, 1881; Jordan & Evermann, l. c., 643, 1896.

Genus 32. PÆCILIA Bloch & Schneider.

Body oblong, often rather deep; mouth small, transverse, with weak jaws; teeth small, in narrow bands, the outer series in each jaw being usually enlarged, curved, movable, and with brown tips; lower jaw not prominent, bones movable. Scales large. Dorsal fin rather small, of 7 to 11 rays; anal fin short, in female nearly opposite dorsal, in males advanced and modified into a sword-shaped intromittent organ. Vertebrae about 28. Intestine long.

Numerous species, mud-eating and viviparous, inhabiting the West Indies, Mexico, and South America. The genus differs from *Mollinisia* only in the smaller size of the dorsal, which usually has 9 or 10 rays and is nearly opposite the anal in the female, but behind it in the male.

- a. Scales very large, 23 to 25 in a lengthwise series; coloration plain; dorsal and anal dotted.
- b. Dorsal rays 7; anal rays 8; coloration plain, fins dotted *vivipara*, 47
- bb. Dorsal rays 9; anal rays 8 *gilli*
- aa. Scales moderate, 28 to 32 in a lengthwise series.
- c. Base of caudal with a black ocellus *pavonina*
- cc. Base of caudal without distinct black ocellus.
- d. Dorsal and anal each with 10 or 11 rays; scales 30 to 32, -10; depth 3 to 3.33 in length; side with two rows of black dots *vittata*
- dd. Dorsal with 7 to 11 rays; anal rays 6 to 9.
- e. Body rather robust, depth in adult more than one-fourth length *mexicana*;
thermalis; *chisoyensis*; *petenensis*; *sphenops*; *dovii*; *boucardi*; *vandepolli*; *dominicensis*; *melanogaster*; *spilurus*
- ee. Body rather elongate, depth in adult not more than one-fourth length *longata*

47. *Pœcilia vivipara* (Bloch & Schneider).

Head 3.6; depth 3.5; eye 3.5; snout 4.2; interorbital 1.8; D. 7; A. 8; scales 25, -8.

Body stout, compressed posteriorly; head depressed, flat; mouth small, jaws weak, lower projecting; teeth small, in a narrow band in each jaw, outer series greatly enlarged, curved, movable, tips brown, those of inner series of short, blunt papillae; greatest width of body at pectorals about 1.2 in head; caudal peduncle greatly compressed 7 or 8 times in its least depth; origin of dorsal fin in female somewhat posterior to that of anal, a little nearer snout than tips of caudal fin; dorsal and anal fins small, the rays short; pectorals and ventrals short; scales large; intestinal canal long and convoluted.

Color, olivaceous, paler below; borders of scales brownish; a dark bar across anterior part of opercle, meeting its fellow below; dorsal fin with a broad black bar, plainest on posterior rays, sometimes not evident on anterior rays; caudal with many fine dark specks, faint traces of two dark crossbars; an irregular black blotch on base of upper caudal rays and a similar one on lower, these sometimes indistinct. The males differ somewhat from the females in color. The black at the base of the caudal is more distinct, and the body is crossed by some 10 or 12 indistinct brownish bars.

The male also differs from the female in having the anterior rays of the anal modified into a long, somewhat curved intromittent organ, the length of which is usually about equal to length of head; the anal is situated much farther forward, its origin being nearer tip of snout than base of caudal; the ventrals are also much lengthened, about 1.3 in head.

This little fish has been recorded from Brazil, Guiana, and Martinique. It was found near Ponce and Fajardo and at Arroyo and Hucare. At the latter place 422 specimens were obtained February 14 in a small brackish-water lagoon, where it was associated with *Dormitator maculatus* and the young of *Tarpon atlanticus*. They most frequented the shallow water along the edges of the lagoon, where they swarmed in great numbers and were easily caught in a small hand-net. Of the 422 examined only 34 were males. The largest females are about 2 inches in total length, the males about 1.44 inches. At Fajardo 71 were taken, of which only one was a male. These were evidently adults, some being 3 inches long and most of them over 2 inches. Some of the females contained well-developed young.

Pœcilia vivipara Bloch & Schneider, Syst. Ichth., 452, pl. 86, fig. 2, 1801, Surinam; Jordan & Evermann, l. c., 691, 1896.

Pœcilia surinamensis Cuvier & Valenciennes, Humboldt, Observ. Zool., II, 158, 1817, Surinam.

Pœcilia schneideri Cuvier & Valenciennes, Hist. Nat. Poiss., XVIII, 135, 1846, Surinam.

Family XXII. ESOCIDÆ. The Needle-fishes.

Body elongate, very slender, compressed or not, covered with small thin scales. Lateral line very low, running as a fold alongside of belly. Both jaws produced in a beak, the lower jaw the longer, very much the longer in the young, which resemble *Hemiramphus*; maxillaries grown fast to premaxillaries; each jaw with a band of small, sharp teeth, besides a series of longer, wide-set, sharp, conical teeth. No finlets. Dorsal fin opposite anal, both fins rather long. Air-bladder present. Lower pharyngeals united to form a long, slender, narrow plate, with flat surface, covered with small, pointed teeth; upper pharyngeals distinct, the third pair little enlarged, each with some 15 moderate, unequal pointed teeth (*Tylosurus marinus*); fourth pair well developed, with similar teeth, but without anterior processes. Vertebrae numerous, with zygapophyses. Ovary single.

Voracious, carnivorous fishes, bearing a superficial resemblance to the gar pikes; found in all warm seas, sometimes entering rivers.

This family contains 4 genera, only 2 of which (*Tylosurus* and *Athlenes*) are found in our waters, and only the first is represented in Porto Rico; the species are about 50, the majority of them American. Their habits are ordinarily much like those of the pike, but when startled they swim along the surface with extraordinary rapidity, often leaping above the water for short distances. When thus leaping the large species of the tropics are sources of danger to incautious fishermen, sometimes piercing the naked abdomen of the savages. Most of them are good food-fishes, but the green color of the bones of the larger species often causes them to be avoided, for no good reason.

a. Gillrakers none; no teeth on vomer; dorsal and anal elevated in front; caudal fin lunate.

b. Body subterete or slightly compressed, its breadth more than two-thirds its greatest depth..... TYLOSURUS, 33

bb. Body much compressed, its breadth not half its greatest depth..... ATHLENES

Genus 33. TYLOSURUS Cocco. The Hound-fishes.

Body elongate, very slender, not much compressed. Both jaws prolonged into a beak, lower jaw somewhat the longer, much the longer in young fishes, the very young resembling *Hemiramphus*. Each jaw armed with a band of small, sharp teeth, beside which is a series of longer, wide-set, sharp, conical, unequal teeth; no teeth on vomer or palatines. Scales small, thin; lateral line running along the side of the belly, becoming median on the tail. No finlets. Dorsal fin more or less elevated anteriorly; caudal fin short, unequally lunated or forked; pectorals moderate; ventrals small, inserted behind the middle of the body. Gillrakers obsolete. Bones usually more or less green.

This genus contains numerous species of comparatively large size. Voracious fishes, chiefly American, one species crossing to Europe, some of them entering rivers.

a. Mouth capable of being nearly or quite closed, upper jaw not conspicuously arched at base.

b. Caudal peduncle compressed, deeper than broad, without trace of keel along the lateral line; no fold of skin across preopercle; caudal subtruncate, the lower lobe somewhat produced; sides with a bluish-silvery band; species of small size, with scales and bones not green.

c. Scales comparatively large, about 85 before the dorsal fin, and about 7 or 8 rows on the cheek; body robust, depth about 5 in head; coloration pale, dorsal and caudal brick-red in life; lateral stripe narrow for its entire length; no scapular blotch..... *notatus*

cc. Scales small, 140 to 150 before dorsal fin, about 12 rows on the cheek; body slender; ventrals inserted at a point nearer cheeks than base of caudal; fins without red; lateral stripe broadened below dorsal fin.

d. Body very slender, depth 7 in head, which is 2.83 in body; eye moderate, 2.33 to 2.75 in postorbital part of head; no distinct notch in the temporal ridge; maxillary not entirely concealed by preorbital. D. 1, 15; A. 1, 17; scales in lateral line 225..... *timucu*, 48

dd. Body less slender, depth 6 in head, which is 2.9 in body; eye large, 2.2 in postorbital part of head; a distinct notch on temporal ridge elose behind eye; maxillary almost entirely concealed by the preorbital. D. 1, 15; A. 1, 17; scales in lateral line 200..... *curyops*

bb. Caudal peduncle very much depressed, wider than deep, but without trace of keel. Head 2.66 in length; eye 2.5 in postorbital part of head; maxillary nearly concealed by preorbital; body subterete; snout very nearly twice length of rest of head; brownish above, silvery below, a bluish lateral stripe edged below with black and yellowish; scales not very small. D. 16; A. 17..... *diplotenia*

bbb. Caudal peduncle more or less depressed, or at least with a more or less developed dermal keel along the lateral line; scales and bones more or less green.

e. Dorsal and anal fins short, each of 14 to 18 rays, anal larger than dorsal and beginning farther forward; last rays of dorsal and anal low; jaws slender, about twice as long as rest of head; no fold of skin across preopercle.

f. Eye very small, 4 to 6 in postorbital part of head; caudal keel sharp, color black; body and tail much depressed, *microps*

- ff.* Eye moderate, 2 to 3.25 in postorbital part of head.
- g.* Caudal fin forked; caudal keel sharp, broad, and conspicuous; top of head flat, striated, without median groove; base of upper jaw much depressed; maxillary entirely hidden by preorbital; teeth very small; ventral fin midway between eye and caudal; scales not very small. D. 1, 13; A. 1, 18 *ardeola*, 49
- gg.* Caudal fin unequally lunate, the emargination not deep, the lower rays moderately produced; scales very small; sides with a silvery lateral stripe; caudal keel not very conspicuous, not black; top of head with median groove; maxillary not entirely concealed by preorbital; ventral inserted midway between preopercle and base of caudal. Species of moderate size, with the scales and bones more or less green.
- h.* Eye moderate, 2.5 in postorbital part of head; pectorals not black posteriorly. D. 1, 15; A. 1, 17; lateral line 300; a dark bar on opercle *marinus*
- hh.* Eye small, 2.25 in postorbital part of head. D. 13 or 14; A. 15 or 16; pectoral pale *almeyda*
- cc.* Dorsal and anal fins long, each of 17 to 25 rays, last rays of dorsal fin more or less elevated in the young, becoming lower in adult; caudal keel rather strong, black; one or more folds of skin across edge of preopercle; caudal fin deeply emarginate or unequally forked. Ventrals inserted midway between base of caudal and middle of eye. Species of large size, with scales and bones green; no distinct lateral stripe.
- i.* Beak short and very strong, its length 1.5 to 1.3 times length of rest of head; body comparatively robust, depth more than one-fifth length of head.
- j.* Dorsal fin long, its rays 1, 21 to 1, 24; anal rays 1, 22 to 1, 24; insertion of dorsal almost opposite that of anal; snout longer, 1.66 to 1.83 length of rest of head; lateral line about 350 *raphidoma*, 50
- jj.* Allied to *T. raphidoma*, but insufficiently characterized *galeatus*
- ii.* Beak strong, but more elongate, about twice length of rest of head; dorsal beginning behind front of anal; greatest depth of body about two-thirds length of pectoral. D. 1, 23; A. 1, 21; lateral line 380; no lateral stripe *acus*
- aa.* Mouth not closing completely, upper jaw arched at base; lobes of dorsal and anal low, the last rays elevated; eye very large, 2.1 in head; scales small, green. D. 24; A. 22; a bluish lateral band *caribbaeus*

48. *Tylosurus timucu* (Walbaum). "Agujon."

Head 2.75; depth 7; eye moderate 2.33 to 2.75 in postorbital part of head; scales small, 225, about 150 before dorsal; D. 15; A. 17. Body very slender, subterete; caudal peduncle not keeled; ventrals inserted nearer cheeks than base of caudal. No distinct temporal notch; maxillary not entirely concealed. Scales and bones not green. Greenish; a silvery-bluish lateral band, widened below dorsal; no scapular spot. Length 1½ feet. Florida Keys to Brazil; not rare.

Found by us at San Juan, Palo Seco, and Hucares, and doubtless occurring everywhere about the island. By the natives all the species of *Tylosurus* are known as "agujon."

Timucu or *Peixe agulha*, Maregrave, Pisc. Brasil., 168, 1648, Brazil.

Esox timucu Walbaum, Artdi Piscium, III, 295, 1792, Brazil; after Maregrave.

Belone subtruncata Poey, Memorias, II, 295, 1861, Havana.

Belone depressa, Poey, Memorias, II, 296, 1861, Havana.

Tylosurus sagitta Jordan & Gilbert, Proc. U. S. N. M. 1884, 25, Key West.

Tylosurus timucu, Jordan & Evermann, l. c., 711, 1896.

49. *Tylosurus ardeola* (Cuvier & Valenciennes). "Agujon."

Head 3.75; depth 8 in head; eye 7; lower jaw from eye 4 in eye (upper jaw broken in our specimen); interorbital equal to eye; D. 13; A. 17. Body very slender; head broad and flat, much broader than in a specimen of *T. timucu* of same size; upper jaw depressed; preorbital completely covering maxillary; teeth small and weak; no fold of skin on opercle; caudal peduncle depressed, with a strong sharp keel; scales moderate, about 150 before dorsal; origin of dorsal decidedly behind that of anal; caudal unequally lunate, lower lobe the longer; ventrals midway between eye and caudal.

Our collection contains a specimen of *Tylosurus*, 13 inches long, obtained at Isabel Segunda, which we are not able to identify with certainty. It is much emaciated and starved in appearance, with very short jaws, which have probably been broken. It seems to agree with the brief description given by Cuvier & Valenciennes of their *Belone ardeola* and also of their *B. cigonella*, the type of which came from Porto Rico.

Belone ardeola Cuvier & Valenciennes, Hist. Nat. Poiss., XVIII, 425, 1846, Martinique.

Belone cigonella Cuv. & Val., Hist. Nat. Poiss., XVIII, 436, 1846, Porto Rico; Poey, Fauna Puerto-Riqueña, 337, 1881.

Tylosurus ardeola, Jordan & Evermann, l. c., 713, 1896.

50. *Tylosurus raphidoma* (Ranzani). "Agujon"; Hound-fish.

D. 21 to 24; A. 22 to 24; scales 350. Caudal keel rather strong, black; one or more folds of skin across edge of preopercle. Body robust, little compressed, its greatest breadth a little more than two-thirds greatest depth; caudal peduncle slightly depressed, a little broader than deep, with a slight black dermal keel. Head broad, interorbital space nearly two-thirds length of postorbital part of

head, with a broad, shallow, nearly naked median groove, which is wider behind and forks at nape. Supraorbital bones with radiating striae. Jaws unusually short, stiff, strong, rapidly tapering forward; large teeth of jaws very strong, knife-shaped. Upper jaw from eye about 1.75 times as long as the rest of head. Eye large, 7 in snout, 2.66 in postorbital part of head, and 1.8 in interorbital width. Maxillary entirely covered by preorbital. Cheek densely scaled; opercle mostly naked except along the anterior margin. Scales of body minute, especially above. Dorsal fin low posteriorly, height of its anterior lobe equaling that of anal, or length of postorbital part of head, its longest ray two-fifths the base of fin; last rays of dorsal and anal much elevated in young; caudal lunate, its lower lobe nearly half longer than upper; middle rays about as long as eye. Ventrals inserted midway between base of caudal and middle of eye, their length a little less than that of pectoral and equal to postorbital part of head; insertion of anal opposite that of dorsal.

Green, silvery below; no lateral stripe; pectoral and dorsal blackish; scales and bones green.

This species reaches a length of 3 to 5 feet, and is generally abundant in the West Indies from the Florida Keys to Brazil, the young occasionally reaching northward, having been reported by Dr. Bean from Ocean City, N. J. It is the most common species of the genus in Porto Rico, having been obtained at San Juan, Mayaguez, Ensenada del Boqueron, Isabel Segunda, and Culebra. It is a vigorous fish and sometimes dangerous in its leaps from the water.

Belone raphidoma Ranzani, Nov. Comm. Ac. Nat. Sci. Inst. Bonon., V, 1842, 359, pl. 37, fig. 1, Brazil.

Belone gerania Cuvier & Valenciennes, Hist. Nat. Poiss., XVIII, 437, 1846, Martinique.

Belone crassa Poey, Memorias, II, 291, 1861, Cuba; Poey, Fauna Puerto-Riqueña, 337, 1881; Stahl, l. c., 79 and 166, 1883.

Belone melanochira Poey, Memorias, II, 294, 1861, Havana.

Tylosurus gladius Bean, Proc. U.S.N.M. 1882, 239 and 430, Pensacola.

Tylosurus raphidoma, Jordan & Evermann, l. c., 715, 1896.




FIG. 17.—*Tylosurus raphidoma*.

Family XXIII. HEMIRAMPHIDÆ. The Balaos.

Body elongate, more or less compressed, covered with large cycloid scales; upper jaw short, lower jaw various, sometimes much produced, the toothed portion at base fitting against the toothed premaxillaries; teeth equal, mostly small and tricuspid; maxillaries ankylosed to premaxillaries. Gillrakers long. Caudal fin rounded, or forked; if forked, the lower lobe the longer. Anal fin modified in the viviparous species (*Zenarchopterus*), unmodified in the others and usually similar to dorsal; no finlets; air-bladder large, sometimes cellular. Third upper pharyngeal on each side much enlarged, solidly united with its fellow to form an oval plate, with slightly convex surface and covered with blunt tricuspid teeth; this is about as large as the united lower pharyngeals and fits into the cavity of the latter; fourth upper pharyngeal wanting or grown fast to third; lower pharyngeal large, thick, triangular, with concave surface. Vertebrae about 50. (Characters verified in *Hemiramphus browni*, *Hyporamphus roberti*, and *Chriodorus atherinoides*.)

Herbivorous fishes of warm seas; mostly shore species; a few pelagic. They feed chiefly on green algæ, and, like the related forms, swim at the surface, occasionally leaping into the air. Size rather small, about a foot in length.

- a. Lower jaw bluntish, not at all produced; teeth rather large; the pectorals and ventrals moderate; shore fishes..... CHRIODORUS
- aa. Lower jaw acute, longer than upper, or more or less produced; teeth small; species oviparous, anal fin in the male not modified, caudal fin unequally lunate.
- b. Lower jaw produced in a long pointed beak, usually longer than rest of head.
- c. Body moderately compressed; pectoral moderate; shore fishes.
- d. Air-bladder simple; sides of body more or less convex; ventrals inserted anteriorly, far in advance of dorsal..... HYPORAMPHUS, 34
- dd. Air-bladder cellular; sides of body nearly vertical and parallel; ventrals inserted posteriorly, not far before dorsal..... HEMIRAMPHUS, 35
- cc. Body very slender and compressed, more or less band-like; pectoral fins very long, ventral very short, inserted posteriorly; pelagic species..... EULEPTORAMPHUS

Genus 34. **HYPORHAMPHUS** Gill. The Half-beaks.

Body elongate, moderately compressed, sides of body not vertical, but more or less convex; the dorsal outline parallel with that of belly. Upper jaw short; lower jaw prolonged into a slender beak, bordered with membrane; the beak shorter in the young; premaxillaries forming a triangular plate, the teeth of which fit against the toothed portion of mandible; maxillaries joined to premaxillaries. Teeth feeble, mostly tricuspid. Gillrakers rather long. Head covered above with large, shield-like scales. Scales large, deciduous. No finlets; caudal more or less forked, lower lobe the longer; dorsal and anal similar, opposite each other, not modified in male; last ray of dorsal usually short; ventrals small, inserted well forward, nearly midway between opercle and base of caudal. Air-bladder large, simple, not cellular. Young with lower jaw short. Side in our species with a distinct silvery band, as in *Atherina*. Oviparous.

Species numerous in all warm seas, going in large schools, but usually remaining near the shore, feeding chiefly on green algæ.

- a. Length of mandible from tip of upper jaw less than rest of head in adult (longer in young); body rather stout; D. 15; A. 16 *unifasciatus*, 51
 aa. Length of mandible from tip of upper jaw not less than rest of head, at all ages, much greater in adult; body more slender. D. 14; A. 15..... *roberti*

51. *Hyporhamphus unifasciatus* (Ranzani). "Balaju"; *Escribano*.

Head 4.6; depth 6.8; eye 4; snout 2.8; maxillary 3.2; mandible 3; interorbital 3.6; D. 14 or 15; A. 14 to 16; pectoral 1.6; ventral 3.2; lower lobe of caudal 1.1; scales 52.

Body elongate, not greatly compressed, sides not parallel, back wider than belly; lower jaw produced into a long beak which, from tip of upper jaw, is shorter than rest of head, this character




FIG. 18.—*Hyporhamphus unifasciatus*.

separating the species from *H. roberti*, in which at all ages the mandible from tip of upper jaw is at least as long as head and in adults much longer; dorsal and anal fins completely scaled, the anterior rays produced but not falcate; lower lobe of caudal the longer; lateral line running very low, close to ventral fin, demarcating side from belly.

Color in life: Back pale-olive or greenish; scales with dark punctulations forming a streak near border; 3 narrow distinct black lines along middle of back from occiput to dorsal fin, the median one faintest; dorsal and anal pale, dusky-tipped; caudal pale, dark-edged; fleshy tip of beak red; a distinct silvery lateral band, about as wide as eye, from upper part of base of pectoral to base of caudal.

The balyhoo is very common in the West Indies, ranging from Key West to Rio de Janeiro, and is considerably used for food; it often leaps from the water and swims in schools, skimming along the surface, 43 examples, 4 to 10½ inches, from San Antonio Bridge, San Juan market, Boqueron, Ponce, Hucares, and Isabel Segunda.

Hemirhamphus unifasciatus Ranzani, Nov. Comm. Ac. Sci. Bonon., V, 1842, 326, Brazil.

? *Hemirhamphus picarti* Cuvier & Valenciennes, Hist. Nat. Poiss., XIX, 25, 1846, Algiers.

Hemirhamphus richardi Cuvier & Valenciennes, Hist. Nat. Poiss., XIX, 26, 1846, Antilles; Cayenne; Bahia; Rio de Janeiro

Hyporhamphus tricuspidatus Gill, Proc. Ac. Nat. Sci. Phila. 1859, 131, Barbados.

Hemirhamphus fasciatus Poey, Memorias, II, 299, 1861, Cuba; not of Bleeker.

Hemirhamphus poeyi Günther, Cat., VI, 262, 1866, Cuba; after Poey.

Hyporhamphus unifasciatus, Jordan & Evermann, I. c., 720, 1896.

Genus 35. **HEMIRAMPHUS** Cuvier. The Balaos

Body more robust than in *Hyporhamphus* and different in form, the sides of body being compressed and nearly vertical and parallel. Head and jaws as in *Hyporhamphus*. Dorsal longer than anal fin and inserted farther forward, its last ray more or less produced in American species. Ventral fins small and inserted well backward, much nearer base of caudal than gill-opening. Air-bladder cellular with many

partitions (in *H. browni*). Species probably numerous, but most of them have not been examined as to the characters which separate this genus from *Hyporhamphus*.

- a. Upper lobe of pectoral orange in life; length of pectoral scarcely greater than depth of body. D. 14; A. 12; scales 53 *brasiliensis*, 52
 aa. Upper lobe of caudal dull-bluish in life; scales rather smaller; length of pectoral one-fifth greater than depth of body *balao*

52. *Hemirhamphus brasiliensis* (Linnæus). "*Balaju*"; *Balao*; *Escribano*.

Head 4.2; depth 6.2; eye 3.8; snout 2.9; maxillary 3.6; mandible 3; interorbital 3.9; D. 13 or 14; A. 12 or 13; pectoral 1.5; ventral 3.1; lower lobe of caudal 1.1; scales 53.

Body much elongate, evenly compressed, the sides parallel, so that a cross section of the body is nearly rectangular; mandible produced into a very long beak, longer than head, and ending in a fleshy tip; upper jaw not produced. Lower lobe of caudal the longer.

Color in life: Back uniform dark-greenish, the scales very slightly paler on edges; top of head like back; upper lobe of caudal yellow, lower olivaceous, the inner edge of both lobes dark; green color of the back ceasing abruptly at level of middle of base of caudal and upper edge of base of pectoral; sides and under parts silvery white. The yellow on caudal and the orange on tip of beak are color markings that distinguish it from *Hyporhamphus unifasciatus* or *roberti*.

Very common; edible; its distribution and habits similar to *Hyporhamphus*; 14 specimens, 9 to 14 inches, from San Antonio Bridge, Aguadilla, Mayaguez, Boqueron, and Fajardo; 2 from San Geronimo.




FIG. 19.—*Hemirhamphus brasiliensis*.

Esox maxilla inferiore producta, Browne, Hist. Jamaica, 443, 1756, Jamaica.

Esox brasiliensis Linnaeus, Syst. Nat., ed. X, 314, 1758, Jamaica; after Browne.

Hemirhamphus marginatus Le Sueur, Jour. Ac. Nat. Sci. Phila., II, 1823, 135, Lesser Antilles; not of Forskål.

Hemirhamphus browni Cuvier & Valenciennes, XIX, 13, 1846, Guadeloupe; Martinique.

Hemirhamphus pleii Cuvier & Valenciennes, Hist. Nat. Poiss., XIX, 19, 1846, Martinique; Santo Domingo.

Macrogathus brevisrostris Gronow, Cat., 148, 1854, Jamaica; after Browne.

Hemirhamphus filamentosus Poey, Mem., II, 297, 1861, Cuba; Poey, Fauna Puerto-Riqueña, 337, 1881; Stahl, l. c., 166, 1883.

Hemirhamphus brasiliensis, Jordan & Evermann, l. c., 722, 1896.

Family XXIV. EXOCETIDÆ. The Flying-fishes.

Body oblong or elongate, covered with cycloid scales, which are rather deciduous. Lateral line running very low, along side of belly. Head more or less scaly, with vertical sides. Mouth moderate, terminal, the jaws not prolonged into a beak. Premaxillaries not protractile, hinged at base mesially; margin of upper jaw chiefly formed by premaxillaries, the short maxillaries entering the lateral margin; maxillary free from premaxillary, its edge slipping under front of preorbital. Dentition various, teeth small and weak. Dorsal fin without spines, inserted on posterior part of body, opposite anal and more or less similar to it; ventrals abdominal, of several soft rays, inserted posteriorly; pectoral fin inserted high, used as an organ of flight; shoulder-girdle and pectoral muscles very strong; caudal fin forked, lower lobe the longer. No finlets. Vent close in front of the anal. Nostrils large, double, near eye. Lower pharyngeals enlarged and fully united, forming a large, transversely concave plate, covered with large, close-set, blunt, tricuspid teeth; third upper pharyngeal greatly enlarged, not united with its fellow, both covered with large, blunt, tricuspid teeth; fourth superior pharyngeal wanting in adult (probably coossified with the third); (these characters verified on *Evocetus californicus*); vertebrae without zygapophyses. Gill-membranes not united, free from isthmus. Pseudobranchiæ hidden, glandular. Gillrakers various. Gills 4, a slit behind fourth. Air-bladder very large, not cellular, so far as known, and extending far backward among hæmapophyses of the caudal vertebrae. Vertebrae about 50. Intestinal canal simple, without cæca.

Carnivorous or herbivorous fishes; 4 genera and about 65 species, abounding in all warm seas, mostly pelagic, swimming near the surface, and skipping or sailing through the air, sometimes for considerable distances.

- a. Roof of mouth (vomer, palatines, pterygoids) and tongue provided with teeth; body not angular in outline (elliptical in cross section); pectoral fins moderate, not reaching beyond middle of dorsal fin, ventrals rather long, inserted behind middle of body; dorsal fin elevated; anal long, its base scarcely shorter than that of dorsal.

PAREXOCÆTUS, 36

- aa. Roof of mouth and tongue with fewer teeth or none (vomer and palatines toothed or not); body angular in outline (a cross section subquadrate); pectoral fins very long, their tips usually reaching nearly to base of caudal, lower jaw little prominent; snout short.

- b. Ventral fins inserted anteriorly, much nearer tip of snout than base of caudal, not used as organs of flight, their tips not reaching nearly to front of dorsal; anal fin long, its base nearly equal to that of dorsal. EXOCÆTUS

- bb. Ventral fins inserted posteriorly, usually nearer base of caudal than tip of snout, used as organs of flight, their tips reaching past middle of base of anal.

- c. Anal fin long, its base a little less than that of dorsal, its origin nearly opposite that of dorsal. EXONAUTES

- cc. Anal fin short, its base one-half to two-thirds that of dorsal, its origin behind that of dorsal. CYPSELURUS; 37

Genus 36. PAREXOCÆTUS Bleeker.

Body moderately elongate, elliptical in cross section. Snout short; lower jaw not produced. Roof of mouth (vomer, palatines, and pterygoids) fully provided with teeth; pectoral fins moderate, not reaching beyond middle of dorsal; ventrals long, inserted behind middle of body; anal fin about as long as dorsal; dorsal high.

Small flying-fishes of tropical coasts, widely distributed.

53. *Parexocætus mesogaster* (Bloch). *Volador; Flying-fish.*

Head 4.4 in length of body; depth 5; D. 12; A. 13; about 38 scales in lateral line, 5 rows of scales between lateral line and dorsal fin. Body elongate, compressed (not angulated), rather deep; width of body at base of pectorals 2 in head; head narrow, compressed, almost trenchant below; inter-orbital area flattish, about as wide as eye, 3 in head. Snout short, rather pointed, its length 4.25 in head; teeth on tongue and palatines; gillrakers numerous, long and slender; pectoral fins of moderate length, their length 1.66 to 2 in length of body, their tips reaching middle of base of dorsal fin; second ray of pectoral divided; dorsal fin very high, its longest rays about 0.16 longer than head; base of dorsal about 1.14 in length of head; tips of anterior rays of dorsal reaching beyond tips of posterior rays when fin is deflexed, almost to base of caudal fin; ventrals rather short, 4.75 in length of body, their tips reaching slightly past origin of anal fin; origin of ventrals midway between pupil and last caudal vertebra; anal fin opposite dorsal; lower lobe of caudal rather short, slightly longer than head.

Color, blue above, silvery below, pectoral (dusky in the young) becoming nearly white in the adult; color of ventrals very similar to pectoral, the duskiess in the young formed of fine blackish dots; upper half of anterior rays of dorsal fin black; anal fin with few small black dots, more numerous in the young; caudal dusky-reddish.

This species reaches a length of 7 inches. It is common in the tropical seas of both the East Indies and West Indies, and in the Hawaiian Islands. It ranges north in the Gulf Stream to Rhode Island and is the most common flying-fish of the Carolina region. The young often has one or two fleshy barbels on the tip of the lower jaw, these being fragile and easily destroyed.

During the voyage of the *Fish Hawk* to Porto Rico and return to Norfolk, flying-fish were seen nearly every day and were particularly abundant between Savannah and the Bahamas, in the Windward and Mona passages, and along the north coasts of Cuba, Santo Domingo, and Porto Rico. Scarcely a day passed when schools of twenty to a hundred or more did not follow along abreast of the ship. They were more numerous on bright days when there was considerable wind and the sea was somewhat rough. Apparently two or more species were seen, though the vast majority probably belong to this species which seems to come nearer to shore than any other. A specimen 5.5 inches long flew aboard the ship at Aguadilla January 18.

Exocætu mesogaster Bloch, Ichthyologia, pl. 399, 1795, Martinique; on a poor drawing by Plumier.

Exocætus orbignianus Cuvier & Valenciennes, Hist. Nat. Poiss., XIX, 131, 1846, Montevideo, based on a drawing.

Exocætus hillianus Gosse, Nat. Sojourn in Jamaica, II, pl. 1, fig. 1, 1851, Jamaica.

Exocætus gryllus Klunzinger, Fische des Rothen Meeres, 586, 1870, Red Sea.

Parexocætus mesogaster Jordan & Evermann l. c., 728, 1896.

Genus 37. *CYPSILURUS* Swainson.

Body elongate, broad above, somewhat compressed; head short, blunt, narrowed below; mouth small; jaws very short, about equal; chin without barbel; maxillaries not joined to premaxillaries; teeth very feeble or wanting; eyes large; gillrakers moderate; scales large, deciduous; no finlets; dorsal fin short, opposite anal; caudal widely forked, lower lobe the longer; pectoral fins very long, reaching past beginning of anal and serving as organs of flight, their great size enabling these fishes to sustain themselves in the air for some time; ventral fins large, posteriorly inserted, also used as organs of flight; air-bladder very large; no pyloric cæca.

Species numerous, in all warm seas, living mostly in the open water and swimming in large schools; largely cosmopolitan, and any of the forms may be expected to be found within our limits.

- a. Second ray of pectoral divided (first simple); third and fourth rays longest.
- b. Ventral fins inserted about midway between pupil and last caudal vertebra.
- c. Dorsal and anal fins without black markings; ventrals pale.
- d. Base of anal 1.66 in base of dorsal; pectoral 1.44 in length, reaching last ray of dorsal; ventrals 2.75 in body, reaching last ray of anal *heterurus*
- dd. Base of anal 2 in base of dorsal; pectoral 1.40 in length of body, the tip reaching end of dorsal fin; length of ventrals 2.89 in body, their tips nearly reaching last ray of anal..... *tutkeni*
- cc. Dorsal and anal fins marked with black; dorsal with one or more dark blotches; anal with a black spot on tips of third to sixth rays; ventrals black, with pale edgings and a white spot near base..... *furcatus*
- bb. Ventral fin inserted midway between posterior margin of preopercle and last caudal vertebra.
- e. Pectoral with posterior half rather abruptly black; anal white *nigricans*
- cc. Pectoral unicolor or nearly so, not abruptly black posteriorly.
- f. Dorsal fin slightly dusky, but without distinct markings; other fins faintly shaded, but without distinct black markings; pectoral reaching base of last anal ray; ventrals almost as far. D. 13; A. 10..... *lineatus*
- ff. Dorsal fin with a round, black blotch as large as eye on tips of middle rays; other fins all pale; pectoral reaching beyond tips of dorsal and anal. D. 12; A. 11..... *cyanopterus*
- bbb. Ventral fins inserted at a point midway between middle of opercle and last caudal vertebra (or between tip of snout and tip of upper lobe of caudal) *bahiensis*, 54
- aa. Second ray of pectoral simple (like the first); third ray divided.
- j. Snout more obtusely descending than in any other species, its length 4.5 in head..... *gibbifrons*

54. *Cypsilurus bahiensis* (Ranzani). *Volador*.

Head 4; depth 5; eye 3.1; snout 4.25; interorbital 3; D. 13; A. 9; scales about 50.

Body quadrate, stout, its width about two-thirds its depth; mouth small, snout short; eye large; top of head flat; pectoral very long, reaching last rays of dorsal and anal, second ray divided, third and fourth longest; ventrals long, longer than head, reaching nearly as far as tip of pectoral, their origin midway between last caudal vertebra and middle of opercle; dorsal considerably in front of anal, its base nearly double that of anal; caudal widely forked, lower lobe the longer.

Color, bluish, silvery above, silvery on sides, white below; side of head silvery; pectoral nearly uniform dusky, paler at base, and bluish-silvery outside; ventrals and anal pale; dorsal and caudal somewhat dusky.

Found in tropical seas, north to Cuba, and said to be one of the commonest species; reaches a length of 8 inches. One specimen flew aboard the ship off Mayaguez, January 20.

Exocetus bahiensis Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., V, 1842, 362, pl. 38, Bahía.

Exocetus vermiculatus Poey, Memorias, II, 300, 1861, Cuba.

Exocetus splanopterus Bleeker, "Nederl. Tydschr. Dierk., III, 1863, 113," Sumatra.

? *Exocetus parva* Poey, Synopsis, 385, 1868, Cuba; description insufficient; taken from an old drawing.

Exocetus bahiensis, Jordan & Evermann, l. c., 739, 1896.

Cypsilurus bahiensis, Jordan & Evermann, l. c., 2836, 1898.

Family XXV. AULOSTOMIDÆ. The Trumpet-fishes.

Body compressed, elongate, covered with small ctenoid scales. Lateral line continuous. Head long; mouth small, at end of a long, compressed tube. Lower jaw prominent, with a barbel at symphysis. Premaxillaries feeble, not protractile; maxillary broad, triangular, with a supplemental bone. Teeth minute, in bands on lower jaw and vomer. Branchiostegals 4; gills 4, a slit behind fourth. Pseudobranchiæ well developed. Gillrakers obsolete. Gill-membranes separate, free from isthmus. Air-bladder large. Spinous dorsal present, of 8 to 12 very slender free spines; the soft dorsal and

anal rather long, similar, posterior, with 23 to 28 rays each; caudal small, rhombic, the middle rays longest but not produced into a filament; ventrals abdominal, of 6 rays, all articulated; pectorals broad, rounded, the space in front of them scaly. First four vertebrae elongated. Two pyloric cæca.

A single genus with two species, found in tropical seas.

Genus 38. AULOSTOMUS Lacépède.

Characters of this genus included with those of the family.

a. Base of soft dorsal and anal not black; each fin with a black band parallel with its base.

b. Eye 2 to 2.5 in postorbital part of head; ground-color reddish; silvery lateral streaks, not all below lateral line.

bb. Eye 3.5 in postorbital part of head; ground-color brown; lateral silvery streaks all below lateral line..... *maculatus*, 55
cinereus

55. *Aulostomus maculatus* Valenciennes. *Trumpet-fish*; "*Trompetero*."

Head 3; eye 2 to 2.5 in postorbital part of head; D. x-23; A. 25; V. 6. Lower jaw prominent, keeled, with a small barbel at symphysis; premaxillary slender, maxillary broad; triangular patches of minute teeth on lower jaw, vomer, palatines, gill-arches, and pharyngeals. Intestinal canal short; two pyloric cæca.

Color, olivaceous, with one or two series of brown or blue dots along each side of the back; another irregular series from the preoperculum along each side of the belly to anal fin; three or four silvery lines on each side of abdomen, replaced on head by irregular oblique streaks; anterior part of dorsal and anal with a horizontal black band, parallel with base of fin but remote from it; caudal fin with usually two round black spots; ventral fins plain, spotted. (Günther.)




FIG. 20.—*Aulostomus maculatus*.

Found in the Caribbean Sea, north to southern Florida; rather common southward; apparently not common about Porto Rico, as only one small specimen was obtained. This was seined in Boqueron Bay January 26.

Aulostoma maculatum Valenciennes, in Cuvier's *Illst. Poissons*, pl. 92, fig. 2, about 1845.

Aulostoma coloratum Müller & Troschel, in Schomburgk's *Hist. Barbados*, 673, 1848, Barbados.

Aulostomus maculatus, Jordan & Evermann, *l. c.*, 754, 1896.

Family XXVI. FISTULARIIDÆ. The Cornet-fishes.

Body extremely elongate, much depressed, broader than deep. Scaleless, but having bony plates present on various parts of body, mostly covered by skin. Head very long, anterior bones of skull much produced, forming a long tube, which terminates in the narrow mouth; this tube formed by symplectic, proethmoid, metapterygoid, mesopterygoid, quadrate, palatines, vomer, and mesethmoid. Both jaws, and usually vomer and palatines also, with minute teeth; membrane uniting bones of tube below very lax, so that tube is capable of much dilation. Post-temporal coossified with the cranium. Branchiostegals 5 to 7; gills 4, a slit behind fourth. Gill-membranes separate, free from isthmus. Gillrakers obsolete. Basibranchial elements wanting. Pseudobranchiæ present. Air-bladder large. Spinous dorsal entirely absent; soft dorsal short, posterior, somewhat elevated; anal fin opposite it and similar; caudal fin forked, middle rays produced into a long filament; pectoral small, with a broad base, preceded by a smooth area as in the *Gasterosteidae*; pectoral ossicles 3; interclavicles greatly lengthened; supraclavicles very small; ventral fins very small, wide apart, abdominal (through partial atrophy of the girdle, by which they lose connection with the interclavicles), far in advance of the dorsal, composed of 6 soft rays. Pyloric cæca few; intestine short. Vertebrae very numerous (4 + 44 to 49 + 28 to 33), the first four very long.

Fishes of tropical seas, related to the sticklebacks in structure, but with prolonged snout and different ventral fins. A single genus, with three species.

Genus 39. *FISTULARIA* Linnæus. Trumpet-fishes.

Characters of the genus included above with those of the family.

- a.* Upper lateral edges of snout with few serrations or none; body with blue spots..... *tabacaria*, 56
aa. Upper lateral edges of snout sharply serrated; body with few blue spots or none..... *petimba*

56. *Fistularia tabacaria* (Linnæus). Trumpet-fish; Trompetero.

Head 2.8; mandible about 4 in snout; snout 3.75 in length of body. D. 14; A. 13. Mouth slightly oblique, lower jaw the longer, overlapping upper; snout much prolonged, tapering but little forward, its edges with fine serrations or none. Margin of orbit with sharp compressed points in front and behind. Reddish-brown above, variegated with numerous large, unequal, oblong, pale-blue spots on sides and back, arranged in series.

West Indies and the neighboring seas, generally common; occasional northward to Carolina and Florida, or even to Massachusetts. Reaches a length of 6 feet. One specimen obtained by Mr. Gray at San Geronimo; reported from Porto Rico by Poey and Stahl.

Petimbuaba, Maregrave, Hist. Brazil, 148, 1648, Brasil.

Fistularia tabacaria Linnæus, Syst. Nat., X, 312, 1758, America; Jordan & Evermann, l. c., 757, 1896.

Fistularia neoboracensis Mitchell, Trans. Lit. and Phil. Soc., I, 1815, 437, New York.

Autostome maregravii Castelnau, Anim. Nouv. Amér. Sud, 30, 1850, Bahia; Rio Janeiro.

Flagellaria fistularis Gronow, Cat. Fishes, ed. Gray, 146, 1854, American Ocean.

Solenostomus serratus, Poey, Fauna Puerto-Riqueña, 327, 1881.

Solenostomus tabacarius, Stahl, l. c., 79 and 165, 1883.

Family XXVII. SYNGNATHIDÆ. The Pipe-fishes.

Body elongate, usually slender, covered with bony plates which are firmly connected, forming a bony carapace. Head slender, snout long, tube-like, bearing short toothless jaws at end. Gill-openings reduced to a small aperture behind the upper part of the opercle. Tail long, prehensile or not, usually provided with a small caudal fin. Male fishes with an egg pouch, usually placed on under side of tail, sometimes on abdomen, commonly formed of two folds of skin which meet on median line. The eggs are received into this pouch and retained until some time after hatching, when the pouch opens, permitting the young to escape. Dorsal fin single, nearly median, of soft rays only; pectorals small, or wanting; ventrals none; anal fin minute, usually present.

a. Tail not prehensile, usually with a caudal fin; axis of head usually in line with axis of body.

b. Top of head with a slight carination, or none.

SYNGNATHINÆ:

c. Pectoral fins present; caudal present.

d. Male with egg pouch under tail, formed by lateral membranes which become connected along middle, forming a closed pouch.

e. Dorsal fin inserted over or just before vent..... SIPHOSTOMA, 40

dd. Male with egg pouch on abdomen; ridges of body prominent and distinct; caudal fin moderate... DORYRHAMPHUS

ee. Pectoral fins wanting; caudal wanting or rudimentary; male with ova attached to abdomen, without closed pouch; no adipose fin.

f. Back without peculiar tube..... SYNGNATHUS

ff. Back with a peculiar tube inclosed by the scutes, and extending for a distance before dorsal fin..... OSPHYOLAX

bb. Top of head strongly carinated..... CORYTHOICHTHYS, 41

HIPPOCAMPINÆ:

aa. Tail prehensile; caudal fin small; head shaped like that of a horse, placed at a large angle with axis of body; egg pouch at base of tail.

g. Body compressed; occiput with a narrow bony crest, surmounted by a coronet; shields with tubercles or spines.

HIPPOCAMPUS, 42

Genus 40. *SIPHOSTOMA* Rafinesque. The Pipe-fishes.

Body elongate, very slender, 6 or 7-angled, not compressed, tapering into a very long tail; dorsal keels of the trunk not continuous with those of tail. Head slender, tapering into a long, tube-like subterete snout, which bears the very short, toothless jaws at end. Humeral bones firmly united with "breast ring." Body covered with a series of bony, keeled, radiated plates, arranged in linear series. Dorsal fin distinct, rather short, inserted before or opposite the vent, which is near middle of body; caudal fin present, rather small; anal fin minute, close behind vent; pectorals developed, short and

rather broad. Male fishes with an egg pouch along under side of tail, formed by two cutaneous folds, and splitting lengthwise to release the young fishes.

Species very numerous, inhabiting all warm seas; abounding in bays among the seaweeds and entering rivers. The females in most species are deeper than the males, with more robust trunk, longer snout, and a more distinct ventral keel.

- a. Top of head with a slight carination or none; snout keeled or not; opercle without prominent ridge; base of dorsal not elevated.
- b. Dorsal moderate or long, its first ray in advance of vent; snout moderate or long; angles of body generally prominent.
- c. Dorsal covering 1 or 2 body rings.
- d. Dorsal covering about 6 or 7 caudal rings, rarely fewer.
- e. Caudal rings 36 to 41. *fistulatum*
- ee. Caudal rings 31 to 34.
- f. Dorsal 29 to 32, on 2+6 rings *mackayi*, 57
- ff. Dorsal 27, on 1+7 rings, rarely on 2+6. *floridæ*, 58
- eee. Caudal rings 24; body rings 19; dorsal rays 30, on 1+6 rings; snout long; body slender. *poeyi*
- ddd. Dorsal covering 4 or 5 caudal rings.
- g. Dorsal rays 24 to 32.
- h. Dorsal rays 29 to 32, on 1+9 rings; rings 17+35; snout 2 in head, head with keel *plagicum*
- hh. Dorsal rays 26 to 28, on 1.5+5 rings; rings 16+34; snout short, 2 in head; head with slight keel *roussau*
- hhh. Dorsal rays 23 to 25, on about 1+4 rings; rings usually 16+33; snout short, 2 in head *clucens*, 59
- gg. Dorsal very short, 18 rays, on 1+5 rings; rings 17+32; snout short *jonesi*, 60
- ddd. Dorsal very short, covering 3 caudal rings and 1.5 body rings, its rays 20 to 23; rings 17+27; snout very short; head keeled above. *brachycephalum*
- cc. Dorsal covering 3 body and 4 to 6 caudal rings.
- i. Rings 16 to 18+29 to 33.
- j. Snout rather short, not half length of head; dorsal fin very high; rings 18+33; dorsal 35, on 5+4 rings; belly in female with a black keel; sides with narrow vertical silvery streaks; dorsal spotted. *affine*
- jj. Snout rather long, more than half head.
- ii. Rings 20 to 21+36 to 38; dorsal 32 to 37, on 3+5 rings; belly flat or slightly concave; snout moderate. *louisianæ*
- iii. Rings 16 to 20+36 to 40; dorsal 36 to 40, on 5+4 or 4+5 rings. *fuscum*
- bb. Dorsal fin very short, its first ray not in advance of vent; rings 15+37 to 39; snout very short, less than two-fifths head; angles of body little marked, the form subterete. *crinigerum*

57. *Siphostoma mackayi* Swain & Meek.

Head 5.66 to 6.25 in total length. D. 29 to 32; rings 18+33 to 36. Snout compressed, 1.75 to 2 in head, its median line with a slight keel above and below, with smaller keels on each side; opercle not keeled. Dorsal somewhat higher than width of a body ring, its base about 1.33 in length of head, covering 2+6 rings. Pectoral higher than length of base; tail about 1.83 in total length of fish; body deep, its greatest depth in adult females equaling width of four body rings.

Color in spirits: Grayish or dark olive; males with about 14 dark-gray crossbars on side, broader than interspaces; body usually more or less spotted with small white spots; dorsal pale, usually dotted over with darker; caudal dusky, generally spotted with white; opercle usually with white bars.

Gulf of Mexico and West Indies; known from the Snapper Banks off Pensacola, Key West, and Cozumel, Yucatan; rather common. One specimen, a female, 7 inches long, taken at Mayaguez, January 19.

Siphostoma mackayi Swain & Meek, Proc. U. S. N. M. 1884, 239, Key West; Jordan & Evermann, l. c., 766, 1896.

58. *Siphostoma floridæ* Jordan & Gilbert.

Head 6 to 6.5 in total length. D. 27; rings 17 to 18+31 to 32. Snout rather short, about 1.66 in head; median line well keeled above and below, the ridge on both sides of median ridges above and below not so conspicuous. Occiput and opercle little keeled. Dorsal shorter than head, covering 1+6 to 7 rings, its height 5 times in base; caudal fin 2.5 in base of dorsal; pectoral slightly higher than length of base; tail longer than trunk, 1.83 in total length, caudal pouch covering about 18 rings.

Color in life: Dark green; side with gray specks and without dark band; tail with faint darker bars, broader than interspaces; sides of tail especially mesially, with many rough and oblong pale spots; snout mottled, especially on side; lower part of the opercle nearly plain; dorsal translucent, yellowish at base; caudal yellow, dusky at tip; anal plain.

Found on sandy shores from North Carolina to Texas (Beaufort, N. C., Corpus Christi, Tex.); rather common. Three specimens seined in Ensenada del Boqueron, January 26.

Siphostoma floridæ Jordan & Gilbert, Proc. U. S. N. M. 1882, 263, Pensacola, Fla.; Jordan & Evermann, l. c., 766, 1896.

59. *Siphostoma elucens* (Poey).

Head 7 in total length; depth 3.6 in head; snout 1.9 in head; body rings 17; caudal rings 32. D. 23, on 1 + 4 rings. Body rather stout; snout moderate; keels rather strong; pouch of male covering about 17 rings.

Color grayish-brown, back and sides more or less mottled with lighter and darker; lower half of snout with alternate bars of white and brown.

Known only from Havana and Porto Rico; one specimen, a male, 6 inches long, with pouch filled with eggs, obtained at San Antonio Bridge, January 12.

Syngnathus elucens Poey, Synopsis, 443, 1867, Havana.

? *Syngnathus flavirostris* Poey, Enumeratio, 178, 1875, Havana.

? *Syngnathus picturatus* Poey, Enumeratio, 178, 1875, Havana.

? *Syngnathus linea* Poey, Enumeratio, 178, 1875, Havana.

? *Syngnathus marmoratus* Poey, Enumeratio, 178, 1875, Havana.

? *Syngnathus ascendens* Poey, Enumeratio, 179, 1875, Havana.

Siphostoma elucens, Jordan & Evermann, l. c., 768, 1896.

60. *Siphostoma jonesi* (Günther).

Rings 17 + 32. Dorsal 18, on 1 + 5 rings. Head and snout short, the latter somewhat bent upward, shorter than the postorbital portion of the head, keeled above, but without spines in the median line. Interorbital space concave, crown and occiput with a median keel. Keels of body sharp but not spiny; lateral keel of the trunk passing into lower keel of tail; a ridge running across opercle but not reaching its end; tail twice as long as trunk without head; caudal very short.

Color: Blackish-brown; upper half of trunk with 3, upper half of tail with 7, equidistant white crossbands.

Bermudas and Porto Rico. Two small males, 2.75 and 3.50 inches long, respectively, dredged at station 6085, between Vieques and Culebra islands, in 14 fathoms, February 6. The larger has the pouch well filled with eggs and the other has a few eggs.

Syngnathus jonesi Günther, Ann. Mag. Nat. Hist., series 4, XIV, 8, 1874, Bermudas.

Siphostoma jonesi, Jordan & Evermann, l. c., 768, 1896.



FIG. 21.—*Corythroichthys cayorum*.

Genus 41. *CORYTHROICHTHYS* Kaup.

This genus differs from *Siphostoma* in the strong keel on top of head, the strong opercular ridges, the short, stout body with prominent angles, and the very short snout. Three species known.

- a. Top of head strongly carinated, keel usually extending from interorbital space to first body ring; body stout, usually with sharp angles and variegated coloration; head in line with axis of body; opercle with a prominent ridge.
- b. Caudal rings about 30; dorsal rays 23 to 27, snout short.
- c. Caudal rings 18 + 30; dorsal rays 23, on 1 + 4 rings; snout 2.4 in head, its color white; body with 12 irregular brown crossbands, and white markings..... *albirostre*
- bb. Caudal rings 25, body rings 20, dorsal rays 40, on 3 + 7 rings; snout long, more than half head; form of body undescribed..... *cayennense*
- bbb. Caudal rings 26, body rings 17; dorsal rays 21, on 1.5 + 3.5 rings; snout very short, 3.2 in head..... *cayorum*, 61

61. *Corythroichthys cayorum* Evermann & Kendall.

Head 8.6; depth 12.6; snout 3.2 in head; eye 4.33; D. 21 rays, on 1.5 + 3.5 rings; A. 3, on first caudal ring; C. 10; P. 10. Rings 17 + 26 = 43. Body short and stout; head short, snout very short; tail but little longer than head and trunk. Cranial ridges strong; a high, sharp keel on snout; occipital keel very high, its edges convex, notched near middle, not continuous with keel on snout; a strong supra-ocular ridge, beginning opposite posterior end of nasal keel and continuing backward, with one hiatus upon upper edge of opercle; just below this on opercle another longer but scarcely stronger ridge;

another short ridge on anterior part of opercle at level of lower part of eye; opercles very convex, as if swollen outward; keels on body and tail all strong, the two lateral keels on body terminating on third caudal ring; the two lateral keels on tail beginning on last body ring, thus overlapping body keels; median keel on side well developed, terminating on sixteenth body ring; ventral keels strong; abdominal keel very strong. Egg-sac on first 18 caudal rings.

Color, yellowish-brown, with darker punctulations; tip of snout white; cheek, throat, and under parts of snout white, crossed by about 7 or 8 irregular brownish bars extending downward and backward; opercles brown; fins pale.

This species is related to *C. albirostre*, of Heckel, differing from it chiefly in the shorter snout, smaller dorsal, and fewer rings. It is known only from Key West and Porto Rico. A fine female, 4 inches long, was seined on Cayo Jancudo, at Fajardo, February 17.

Corythoichthys cayorum Evermann & Kendall, Bull. U. S. F. C. 1897 (Feb. 7, 1898), 128, pl. 7, fig. 7, near Crawfish Bar, Key West, Fla.; Jordan & Evermann, l. c., 2838, 1898.

Genus 42. HIPPOCAMPUS Rafinesque. The Sea-horses.

Body strongly compressed, belly gibbous, tapering abruptly to a long, quadrangular, prehensile tail. Head with a distinct curved neck placed nearly at a right angle with direction of the body, surmounted by a compressed occipital crest, on top of which is an angular, star-shaped coronet; top and sides of head with spines. Physiognomy remarkably horse-like, like that of a conventional "knight" at chess. Body and tail covered with bony plates, forming rings, those on body each with 6 spines or tubercles, those of tail with 4. Pectoral fins present, short and broad; anal minute, usually present; dorsal fin moderate, opposite the vent. Egg-pouch in male a sac at base of tail, terminating near vent.

Species numerous in all warm seas. They attach themselves by their tails to seaweed and other floating substances, and are often carried to great distances by currents.

- a. Dorsal fin large, with 19 rays; rings 11 + 32 to 35; depth of body equals length of head; snout 2.33 in head; dorsal fin on 3.5 + 0 rings; body mottled, not dotted *hudsonius*
aa. Dorsal fin smaller, with 16 to 18 rays.
b. Dorsal fin with 17 or 18 rays, on 1 caudal ring; snout short, less than half length of head; light-blue spots on head and snout; head usually without filaments; size large *punctulatus*, 62
bb. Dorsal fin with 16 rays, on 4 + 0 rings; rings 12 + 31; snout longer than postocular distance; body unspotted; size moderate *stylifer*
aaa. Dorsal fin very small, with 12 rays, on 2 + 1 rings; rings 11 + 30; snout very short; body without white spots. . . *zosteræ*

62. *Hippocampus punctulatus* Guichenot. Sea-horse; Cabalito de Mar.

D. 18, covering $2\frac{1}{2} + 1$ rings; rings 12 + 30. Snout short, about 2.6 in head, or slightly shorter than postorbital part of head; eye 2 in snout; supraorbital spines divergent, each with a minute spine in front; coronet high, about 1.5 in snout; no filament on coronet, and only a few very short ones anywhere on head, none on body; spines of body strong, but blunt.

Color, dark-brown, fins pale; head with numerous minute white specks; body and tail with a few similar specks.

Found in the tropical parts of the Atlantic; common in the West Indies, Brazil, and western Africa, reaching occasionally northward in the Gulf Stream as far as Beaufort, N. C. (Jenkins.) One specimen, about 3 inches long, seined at Ponce, January 30.

Hippocampus punctulatus Guichenot, in Sagra's Cuba, Poiss., 174, pl. 5, fig. 2, 1860, Cuba; Poey, Fauna Puerto-Riqueña, 347, 1881; Stahl, l. c., 79 and 165, 1883; Jordan & Evermann, l. c., 777, 1896.

Hippocampus marginalis Heckel, in Kaup's Lophobr., 15, 1856, Mexico.

Hippocampus fascicularis Heckel, in Kaup's Lophobr., 15, 1856, Mexico.

Hippocampus longirostris Heckel, in Kaup's Lophobr., 12, 1856; not of Cuvier.

Hippocampus guttulatus Günther, Cat., VIII, 202, 1870; probably not of Cuvier.

? *Hippocampus kuda* Bleeker, Nat. Tyds. Ned. Ind., 111, 82, East Indies.

Family XXVIII. ATHERINIDÆ. The Silversides.

Body rather elongate, somewhat covered with scales of moderate or small size, which are usually, but not always, cycloid. No lateral line; some scales often with rudimentary mucous tubes. Cleft of the mouth moderate. Teeth small, on jaws and sometimes on vomer and palatines, rarely wanting. Premaxillaries protractile or not. Opercular bones without spines or serrature. Gill-openings wide,

the gill-membranes not connected, free from isthmus; gills 4, a slit behind fourth. Pseudobranchiæ present; gillrakers usually long and slender. Branchiostegals 5 or 6. Dorsal fins 2, well separated, the first of 3 to 8 slender flexible spines, second of soft rays; anal with a weak spine similar to soft dorsal, but usually larger; ventral fins small, abdominal, not far back, 1 small spine and 5 soft rays; pectorals moderate, inserted high. Air-bladder present. No pyloric cæca. Vertebrae numerous, usually about $23 + 23 = 46$; third and fourth superior pharyngeals coossified, with teeth.

This family comprises about 15 genera and 60 species of carnivorous fishes, mostly of small size, living in great schools near shore in temperate and tropical seas; a few species in fresh water; all the species have a silvery band along side; this is sometimes underlaid by black pigment. All which are large enough are highly valued as food; hence the common name of "fishes of the king" (pescados del rey, or pesce re, or peixe rey).

- a. Premaxillary narrow posteriorly, its edge nearly straight. Body little compressed, belly rounded; pectorals short; scales cycloid; vomer with teeth; first dorsal with 5 to 9 spines, inserted in front of the rather short anal; mouth short *ATHERINA*, 4
- aa. Premaxillary broad posteriorly, its edge strongly curved.
- b. Lower jaw strong, projecting beyond upper.
- c. Scales small, rough, in 70 series; teeth well developed; vomer usually with a few teeth; jaws long.... *LETHOSTOLE*
- cc. Scales large, smooth, in 38 to 50 series..... *CHIROSTOMA*
- bb. Lower jaw moderate, its tip included; vomer without teeth.
- d. Scales laciniate; dorsal and anal sealy *KIETLANDIA*
- dd. Scales cycloid; soft dorsal and anal mostly without scales..... *MENIDIA*




FIG. 22.—*Atherina stipes*.

Genus 43. ATHERINA (Artedi) Linnæus.

Body oblong, compressed. Mouth large, terminal, oblique; jaws about equal, their edges nearly straight; maxillary extending to front of eye. Premaxillaries narrow posteriorly, strongly protractile. Villiform teeth in bands on jaws, vomer, and palatines. Species numerous, mostly European.

- a. Anal fin rather short, of 10 to 16 rays.
- b. Scales large, 36 to 40; first dorsal with 5 or 6 spines.
- c. Head very broad, interorbital width about equal to the large eye, which is about 2.33 in head. Scales 36 to 41.
- d. Anal rays 1, 12, or 1, 13; head about 4 in length *stipes*, 63
- dd. Anal rays 1, 10 or 11; head 3.5 to 3.66 in length *laticeps*, 64
- cc. Head narrow and pointed, the large eye much greater than interorbital width; body slender and weak, depth 6 in length; anal 1, 12; scales 38 to 41 *aræa*, 65
- bb. Scales small, 45 to 52; body very slender; first dorsal long, with 7 or 8 spines.
- e. Anal rays 1, 11; eye 3 in head; scales 45 *harringtonensis*
- cc. Anal rays 1, 15; eye 2.5 in head; scales 52..... *carolina*
- aa. Anal fin longer, of about 20 rays; dorsal rays v-1, 15..... *microps*

63. *Atherina stipes* (Müller & Troschel).

Head 4; depth 5.3; eye 2.5; snout 4; maxillary 2.7; mandible 2.3; interorbital 3; D. v-1, 9; A. 1, 12 or 13; pectoral 1.6; ventral 2.3; caudal 1.2; scales 41.

Close to *A. laticeps*, but with top of head narrower and more pointed, eye smaller, lateral band much wider, about 4.5 in head, equal to length of snout. Known only from the Barbados and Porto Rico; apparently rare, only two specimens collected, about 2.5 inches long, these being taken with the abundant *A. laticeps* at Culebra.

Atherina stipes Müller & Troschel, in Schomburgk, Hist. Barb., 671, 1848, Barbados; Jordan & Evermann, l. c., 790, 1896.

64. *Atherina laticeps* Poey. *Cabezote*.

Head 3.5; depth 4.8; eye 2.3; snout 4; maxillary 2.1; mandible 2; interorbital 2.4; D. v-1, 9; A. 1, 10 or 11; pectoral 1.5; ventral 2; caudal 1.2; scales 38. Body elongate, somewhat compressed, especially posteriorly, very heavy forward; head very large, flat above, interorbital space very broad, equal to eye, which is very large; snout very short, wide and blunt; mouth wide, very oblique; maxillary narrow, reaching beyond front of eye; lower jaw the shorter; jaws, vomer, and palatines with villiform bands of very short teeth; spinous dorsal of extremely slender flexible spines; scales large, with a glazed or enamel-like appearance, as if highly polished, in alcoholic specimens when dry.

Color in life: Translucent green, silvery below with a well-defined silvery lateral band, below which is a series of dots along side; back with dark dots forming streaks along rows of scales; snout above with black dots; fins pale, nearly plain; a dusky shade at base of caudal. (Jordan & Evermann.)

Caribbean Sea, north to western Florida. A small and very common species; 290 examples, 1.5 to 3 inches in length, from Ponce and Culebra.

Atherina laticeps Poey, *Memorias*, II, 265, 1861, Havana; Jordan & Evermann, l. c., 790, 1896.

65. *Atherina aræa* Jordan & Gilbert.

Head 4.5; depth 6; eye 2.6; snout 4; maxillary 2.7; mandible 2.3; interorbital 3.1; D. vi-1, 9; A. 1, 12; pectoral 1.5; ventral 2.2; caudal 1.4; scales 41. Body more slender and head smaller and more pointed than in *A. laticeps* or *A. stipes*; maxillary scarcely reaching front of orbit.




FIG. 23.—*Atherina aræa*.

Color of *A. stipes*, belly dusky behind vent; base of anal dusky, the color continued along caudal peduncle in a series of black dots; no black dots on sides.

Gulf of Mexico to Key West and Cozumel. Uncommon in Porto Rico; only four specimens, 2.5 inches in length, taken with *A. laticeps*, at Culebra.

Atherina aræa Jordan & Gilbert, *Proc. U. S. N. M.* 1884, 27, Key West; Jordan & Evermann, l. c., 790, 1896.

Family XXIX. MUGILIDÆ. The Mulletts.

Body oblong, more or less compressed, covered with rather large cycloid scales; no lateral line, but the furrows often deepened on middle of each scale so as to form lateral streaks. Mouth small, jaws with small teeth, or none, teeth various in form; premaxillaries protractile. Gill-openings wide, membranes separate, free from isthmus. Branchiostegals 5 or 6. Gillrakers long and slender. Gills 4, a slit behind fourth. Pseudobranchiæ large. Two short dorsal fins, well separated, anterior with 4 stiff spines, last one of which is much shorter than the others; second dorsal longer than first, similar to the anal; anal spines 2 or 3, graduated; ventral fins abdominal, not far back, composed of 1 spine and 5 rays; caudal forked. Air-bladder large, simple. Intestinal canal long. Peritoneum usually black. Vertebrae 24.

The family comprises 8 or 10 genera and about 100 species, inhabiting the fresh waters and coasts of warm regions, feeding on organic matter contained in mud.

“In the genus *Mugil*, a considerable indigestible portion of the latter is swallowed, and in order to prevent larger bodies from passing into the stomach, or substances from passing through the gill-openings, these fishes have the organs of the pharynx modified into a filtering apparatus. They take in a quantity of sand or mud, and after having worked it for some time between the pharyngeal bones, they eject the roughest and indigestible portion of it. The upper pharyngeals have a rather irregular

form; they are slightly arched, the convexity being directed toward the pharyngeal cavity, tapering anteriorly, and broad posteriorly. They are coated with a thick, soft membrane, which reaches far beyond the margin of the bone and is studded all over with minute horny cilia. Each branchial arch is provided with a series of long gillrakers, which are laterally bent downward, each series closely fitting to the sides of the adjoining arch; they constitute together a sieve admirably adapted to permit a transit for the water, retaining at the same time every solid substance in the cavity of the pharynx."

Of the 5 genera occurring in American waters, only 2 are represented in Porto Rico.

MUGILINÆ:

- a. Stomach muscular, gizzard-like; teeth slender, usually having the form of cilia; lower jaw angular in front; species chiefly marine.
- b. Anal spines 3; teeth cilliform, flexible.
- c. Orbit with a well-developed adipose eyelid, covering part of iris; cilia in one or few series, slender; cleft of mouth chiefly anterior..... MUGIL, 41
- bb Anal spines 2, first soft ray simple but articulate; teeth distinct, in a few series, scarcely cilliform, often obsolete in lower jaw; lips thin; no adipose eyelid; preorbital serrate..... QUERIMANA

AGONOSTOMINÆ:

- aa. Stomach not gizzard-like; teeth not cilliform, lower jaw not angular in front; cleft of mouth lateral; fresh-water species, inhabiting chiefly mountain torrents in the Tropics.
- d. Teeth in villiform bands.
- c. Anal spines 2; teeth in bands on jaws and vomer; lower jaw without lamelliform folds..... AGONOSTOMUS, 45
- dd. Teeth coarse, broad, truncate incisors, with their free edges serrate; smaller teeth on vomer; none on palatines; head heavy, the blunt, tumid snout overhanging the small, inferior mouth; lower jaw forming a sharp soft edge..... JOTURUS

Genus 44. MUGIL (Artedi) Linnæus. The Mullet.

Body oblong, somewhat compressed, covered with large scales. Head large, convex, scaled on sides and above. Mouth small, subinferior, the lower jaw angulated. Jaws with one or a few series of short, flexible, cilliform teeth; no teeth on vomer or palatines. Eye large, with a large adipose eyelid, which is little developed in the young. Stomach muscular, like the gizzard of a fowl.

Species very numerous, living on mud and running in great schools along the shores and in brackish lagoons of all warm regions.

- a. Soft dorsal and anal fins almost naked; anal rays III, 8, rarely III, 7; sides with dark longitudinal stripes along rows of scales; caudal deeply forked, size large.
- b. Scales about 33 in longitudinal series; depth about 4.5 in length to base of caudal, teeth very minute; distance from tip of pectoral to front of dorsal about two-sevenths the length of pectoral; lips rather thin..... *brasiliensis*, 66
- bb. Scales about 41 in a longitudinal series; depth about 4 in length to base of caudal; teeth close-set, rather small; distance of tip of pectoral from front of dorsal about two-ninths the length of pectoral..... *cephalus*
- aa. Soft dorsal and anal fins scaled, sides without dark stripes along rows of scales; caudal less deeply forked; size smaller.
- c. Anal rays III, 9, scales 35 to 45 in a longitudinal series.
- d. Scales 42 to 45 in longitudinal series; teeth small..... *incilis*
- dd. Scales 35 to 38 in longitudinal series.
- c. Pectoral not nearly reaching origin of dorsal, distance from its tip to front of dorsal being in the adult one-sixth length of pectoral, teeth close-set, rather small, but distinctly visible without a lens; scales 38 or 39 in longitudinal series; dorsal less falcate..... *curma*, 67
- cc. Pectoral nearly reaching origin of dorsal; scales 35 or 36 in longitudinal series; bare space between dentary bones small..... *gaimardianus*
- cc. Anal rays III, 8; scales very large, about 33 in a longitudinal series; teeth wide-set, larger than in any other species except *setosus*, about as long as nostril; upper lip thick; pectoral not nearly reaching front of dorsal; size small..... *trichodon*, 68

66. *Mugil brasiliensis* (Agassiz). "Liza"; Mullet; Lebranco.

Head 4; depth 5; eye 5.75; snout 4.5; interorbital 2; D. IV-1, 8; A. III, 8; scales 35-12.

Body long and slender, more so than in any other of our species; snout broad, blunt, and evenly rounded; profile from snout to dorsal fin almost straight; ventral outline somewhat more convex; top of head nearly flat; upper lip thin; preorbital large, almost covering maxillary; eye hidden in front and behind by a broad adipose membrane; teeth minute; scales large, about 21 between tip of snout and origin of spinous dorsal; soft dorsal and anal almost naked, caudal covered with fine scales at base; margin of soft dorsal and anal very concave; caudal deeply forked, the lobes equal, about as long as head; pectoral moderate, one-sixth in head, not reaching origin of spinous dorsal by a distance greater than diameter of eye; ventrals slightly shorter than pectoral.

Color, dusky above, silvery below; a dusky streak along each row of scales; scales on side and opercle with dark punctulations; ventrals pale-yellowish; other fins dusky.

Found from Cuba to Patagonia; common throughout the West Indies and along the coast of Brazil, and very common in Havana market, where it is known as "lebrancho." It is the most abundant mullet seen in the markets of Porto Rico, and perhaps the most abundant species of the family about the island. It is an important food-fish and is held in high esteem. Length, 16 to 18 inches.

Mugil brasiliensis Agassiz, in Spix, Pisc. Brasil., 234, pl. 72, 1829, Atlantic Ocean off Brazil; the types in the museum at Munich; Poey, Fauna Puerto-Riqueña, 335, 1881; Jordan & Evermann, l. c., 810, 1896.

Mugil liza Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 83, 1836, Brazil, Porto Rico, Maracaibo, Surinam, and Martinique.

Mugil lebranchus Poey, Memorias, II, 260, pl. 18, fig. 3, 1861, Cuba; Poey, Fauna Puerto-Riqueña, 335, 1881, Stahl, l. c., 78 and 164, 1883.

67. *Mugil curema* Cuvier & Valenciennes. "Liza"; "Josea"; White Mullet.

Head 4 to 4.4; depth 3.9 to 4.5; eye 4; snout 3.5; D. IV-1, 8; A. III, 9; scales 38,-11; interorbital width 2.6. Body moderately elongate; snout rather narrow and pointed; upper profile curved very slightly from snout to dorsal fin; ventral outline more convex. Upper lip rather thick; preorbital narrow, nearly covering maxillary posteriorly; eye with broad adipose membrane in front and behind; teeth thick-set, small, but distinctly visible to naked eye; scales small, about 23 from origin of dorsal to tip of snout; soft dorsal, anal, and caudal densely scaled; soft dorsal and anal concave; pectoral short, not reaching origin of spinous dorsal, longer in young; caudal deeply forked, the lobes nearly twice length of middle rays.




FIG. 24.—*Mugil curema*.

Color, dark-olive above, with steel-blue reflections, silvery below; no dusky streaks along side; a rather small dark blotch on base of pectoral; spinous and soft dorsals and pectoral pale, with numerous small dark punctulations; caudal pale, yellowish at base, margin dark; anal and ventrals yellowish; side of head with two yellowish blotches.

This species measures a foot or less. Next to *M. brasiliensis* it is the most abundant mullet about Porto Rico. Specimens are in the collection from San Juan, Aguadilla, Mayaguez, Ensenada del Boqueron, Arroyo, Fajardo, Vieques, Culebra, and Caguas. It has a very wide range, being found from Cape Cod to Brazil, and Magdalena Bay to Chile, and is generally common on both coasts of America, particularly in the Tropics. Its occurrence in fresh water in the Rio Grande at Caguas, some 40 miles from salt water, is of interest. It is found there in some numbers, and is one of the most important food-fishes of the region.

Mugil curema Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 87, 1836, Brazil, Martinique, and Cuba; Jordan & Evermann, l. c., 813, 1896.

Mugil petrosus Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 89, 1836, Brazil, Surinam, Gulf of Mexico, and Cuba.

68. *Mugil trichodon* Poey. "Liza"; Fan-tail Mullet.

Head 4.2; depth 3.6; eye 4; snout 3.5; interorbital width 2.5; D. IV-1, 8; A. III, 8; scales 33,-11. Body robust, depth somewhat greater than in *Mugil curema*; snout rather narrow and pointed; dorsal and ventral outlines about equally curved; upper lip thicker than in any other American species; preorbital narrow, covering little of maxillary; eye covered in front or behind by a broad, adipose membrane; teeth wide-set, larger than in most other species, about as long as nostril, and plainly

visible in each jaw; scales large, about 21 between tip of snout and origin of dorsal fin; soft dorsal, anal, and caudal densely covered with fine scales; soft dorsal and anal concave; pectoral not reaching origin of dorsal; caudal broad, widely forked; caudal peduncle deep but thin.

Color, dusky-olive above with some bluish reflections, silvery below; no dusky streaks along rows of scales; a dark blotch on base of pectoral; dorsals and caudal pale, the former with very small dark punctulations; caudal margined with dark; anal and ventrals yellowish; pectoral pale, finely punctulate with brown.

This species reaches a length of 6 to 10 inches. It is found from Florida Keys to Brazil, and is abundant about Key West, but not about Cuba, and does not seem to be common about Porto Rico, the collection containing a single small specimen obtained at Puerto Real.

Mugil trichodon Poey, Ann. Lye. Nat. Hist. N. Y., XI, 1875, 66, pl. 8, figs. 4 to 8, Cuba; Jordan & Evermann, l. c., 816, 1896

Genus 45. AGONOSTOMUS Bennett. The Dajaos.

Fresh-water mullets, with cleft of mouth extending laterally about to front of eye. Small teeth in villiform bands in both jaws, and sometimes on vomer. Edge of lower lip rounded, not sharp. Stomach not gizzard-like. Anal spines usually 2, first soft ray slender and often taken for a spine.

This genus occurs in streams of mountainous regions in tropical countries. The American species constitute the subgenus *Dajaus*, characterized by the presence of teeth on the palatines.

- a. Interorbital space flat, narrow, 4 in head; head 3.66; depth 4.33 to 4.50; lips thin; scales 40 to 42, -13. *percooides*
 aa. Interorbital space convex, broad, 2.66 to 3 in head.
 b. Lips thin; body rather deep; maxillary to front of eye *monticola*, 69
 bb. Lips thick.
 c. Maxillary short, barely reaching front of eye, 4 in head; eye 5 in head; head 4.25 in length *nasutus*
 cc. Maxillary long, reaching posterior margin of pupil, 2.5 in head; eye very small, 6.5 in head; head 3.5 in length *microps*




FIG. 25.—*Agonostomus monticola*.

69. *Agonostomus monticola* (Bancroft). "Dajao."

Head 3.5; depth 3.8; eye 6; snout 3.4; maxillary 3; mandible 2.4; interorbital 2.9; D. iv-1, 8; A. iii, 9; pectoral 1.7; ventral 1.9; caudal 1.2; scales 42, -13.

Body elongate and compressed; back scarcely elevated; head very broad above and slightly convex; mouth moderate, upper lip broadened in front, maxillary reaching front of eye, lower jaw included; villiform teeth on each jaw, vomer, and palatines; eye much nearer tip of snout than edge of opercle; scales large, slightly ctenoid; dorsal fin as in *Mugil*, spinous dorsal with an appendage of 2 modified scales on each side of its base anteriorly; caudal large, moderately forked.

Color in life: Brownish above, scales very dark-edged on upper three-fifths of side; lower parts white; top of head dark, cheek and opercles white, with brassy shades; under part of head white; anal black; a black blotch at base of caudal, becoming faint or disappearing with age; dorsal spines dark; soft dorsal brassy at base, pale at tip; pectoral and ventral pale; anal yellowish, pale at tip, a black blotch near posterior border; caudal darker, yellowish at base; peritoneum black. In spirits not much different.

This species occurs in the fresh waters of the West Indies and eastern Mexico and is not uncommon. It is very abundant in the fresh-water streams of Porto Rico, and is much used as food; many were collected at Caguas in the Río Loiza and Río Caguaitas and in Río Bayamon at Bayamon, of all stages of growth from 1.5 to 11 inches; also common in the large spring at Aguadilla, from which Columbus is said to have supplied his ships with fresh water November 17, 1493.

Mugil monticola Baneroff, in Griffith's edition of Cuvier's Animal Kingdom, Fishes, 367, pl. 36, 1836.

Mugil irretitus Gosse, Nat. Sojourn Jamaica, 84, 1851, Jamaica.

Agonostomus monticola, Poey, Fauna Puerto-Riqueña, 335, 1881; Stahl, l. c., 164, 1883; Jordan & Evermann, l. c., 819, 1896.

Dajaus (*Agonostomus*) *monticola*, Stahl, l. c., 78, 1883.

Family XXX. SPHYRÆNIDÆ. The Barracudas.

Body elongate, subterete, covered with small cycloid scales; head very long, pointed, pike-like, scaly above and on sides; mouth horizontal, large; jaws elongate, the lower considerably projecting; upper jaw nonprotractile, its border formed by premaxillaries, behind which are the broad maxillaries; large, sharp teeth of unequal size on both jaws and on palatines, none on vomer; usually a very strong, sharp canine near tip of lower jaw. Opercular bones without spines or serratures. Gill-openings wide, gill-membranes not united, free from isthmus; gillrakers very short or obsolete. Branchiostegals 7; gills 4, a slit behind the fourth. Pseudobranchiæ well developed. Air-bladder large, bifurcate anteriorly; many pyloric cæca. Lateral line well developed, straight. Pectoral fin short, placed in or below line of axis of body; ventrals 1, 5, abdominal, in advance of middle of body; first dorsal over ventrals, of 5 rather stout spines; second dorsal remote from first dorsal, similar to anal and opposite to it; caudal fin forked. Vertebrae 24. First superior pharyngeal not present; second, third, and fourth separate, with teeth; lower pharyngeals separate.

A single genus of about 20 species; carnivorous, pike-like fishes, often of large size, active and voracious, inhabiting warm seas, many of them highly valued as food.




FIG. 26.—*Sphyræna barracuda*.

Genus 45. SPHYRÆNA (Artedi) Bloch & Schneider.

Characters of this genus included above.

- a. Scales large, 75 to 85 in lateral line; origin of first dorsal behind root of ventrals, over last third or fourth of pectoral; body compressed; lower jaw with fleshy tip; maxillary reaching past front of orbit; teeth large ... *barracuda*, 70
- aa. Scales moderate, 110 to 130 in lateral line; body subterete or compressed.
- b. Pectoral reaching front of spinous dorsal; maxillary reaching front of orbit; origin of spinous dorsal behind root of ventrals..... *guachancho*, 71
- bb. Pectoral not reaching front of first dorsal; maxillary not reaching front of orbit..... *picudilla*, 72
- aaa. Scales very small, 150 to 170 in lateral line; origin of spinous dorsal well behind tip of pectoral, before the vertical from root of ventrals; lower jaw with fleshy tip. Body slender, subterete..... *sphyræna*

70. *Sphyræna barracuda* (Walbaum). "*Picuda*."

Head 3.3; depth 7 (2 in head); eye 6.6; snout 2.1; maxillary 2.1; mandible 1.6; interorbital 5.1; D. v-i, 10; A. 1, 8; pectoral 2.7; ventral 3.3; caudal 1.3; scales 12-83-11.

Body elongate, slightly compressed, head cuboid back of eyes, snout conic; mouth very large, lower jaw strongly projecting, with a fleshy tip at symphysis; maxillary reaching past front of eye; teeth on jaws, premaxillaries, and palatines unequal, very strong, sharp, and compressed; much smaller teeth on edges of premaxillaries; scales of trunk large, cycloid; smaller scales on cheeks and opercles; dorsal fins 2, widely separated, origin of first midway between front of eye and origin of soft dorsal; its spines slender and flexible; soft dorsal and anal similar; caudal well forked, upper lobe slightly the longer

Color in spirits: Dark above, pale below, with silvery bluish reflections; dark longitudinal streaks along rows of scales above lateral line; young with dark, irregularly shaped blotches on middle of

side, these often extended as vertical bars, disappearing with age; usually from one to several small, very dark brown spots, sometimes black, scattered irregularly on side; ventral, anal, and soft dorsal chiefly black; top of head, upper edge of maxillary, and edge of opercle above, black.

The picuda is common in the West Indies, ranging north to Charleston and the Bermudas, and south to Brazil. It is a fierce and voracious fish, of good food qualities. Thirteen examples, from 6 to 16 inches in length, were collected at San Juan market, San Antonio Bridge, Mayaguez, Ensenada del Boqueron, Culebra, Hucares, and Fajardo.

Umbla minor marina, the Barracuda, Catesby, Fishes of Carolina, etc., pl. 1, 1731, Bahamas.

Picuda, Parra, Dif. Piezas, Hist. Nat. Cuba, 90, pl. 35, fig. 2, 1787, Havana.

Esox barracuda Walbaum, Artedi Piscium, III, 94, 1792, Bahamas; after Catesby.

Sphyræna becuna Lacépède, Hist. Nat. Poiss., V, pl. 9, fig. 3, 1803, Martinique; from a drawing made by Plumier.

Sphyræna picuda Poey, Fauna Puerto-Riqueña, 334, 1881; Stahl, l. c., 76 and 162, 1883; Jordan & Evermann, l. c., 823, 1896.

Sphyræna barracuda, Jordan & Evermann, l. c., 2841, 1898.

71. *Sphyræna guachancho* Cuvier & Valenciennes. *Guachanche; Guachanche Pélon.*

Head 3; depth 7; eye 5.4; snout 2.1; maxillary 2.2; mandible 1.6; interorbital 7.2; D. v-1, 9; A. 1, 8; pectoral 3; ventral 3.6; caudal 1.6; scales 115.

This species differs from *S. barracuda* in the much smaller scales, larger eye, longer and more slender head, and in the absence of color markings. The single specimen in the collection, 9 inches long, taken at Isabel Segunda, is nearly plain in spirits, save for fine dark longitudinal lines following the rows of scales above the lateral line. The back is of a faded brown color, the lower parts silvery white everywhere below lateral line; no regular bars or scattered spots.

The guachanche is a common West Indian species ranging north to Pensacola, and occasionally wandering in the Gulf Stream to Woods Hole. It does not reach as great size as the picuda.

Sphyræna guachancho Cuv. & Val., Hist. Nat. Poiss., III, 342, 1829, Havana; Jordan & Evermann, l. c., 824, 1896.

Sphyræna guntheri Haly, Ann. Mag. Nat. Hist., XV, 1875, 270, Colon.

72. *Sphyræna picudilla* Poey. *Picudilla.*

Head 3.16; depth 2.25 in head; eye large, about 5 in head, 1.5 times interorbital space. D. v-1, 9; A. 1, 9; scales 110. Body rather robust, subterete, covered with scales of moderate size; head rather large; maxillary rather small, about 2.6 in head, not reaching orbit. Jaw with fleshy tip, bluntly conical. Interorbital area flattish; median groove shallow, divided by a very indistinct median ridge; supraocular ridge bony, striate; preocular ridge rather prominent. Premaxillary teeth small, subconical; dentition as in *Sphyræna borealis*, but slightly weaker; position of spinous dorsal, in comparison with the ventrals, variable; distance from tip of snout to origin of spinous dorsal about 2.1 in body; pectoral not reaching spinous dorsal; space separating dorsals about 5.5 in body; second dorsal equal to and somewhat in advance of anal; cheek and opercles scaly; small embedded scales on upper part of head; scales on body moderate, uniform in size.

Color light-olive above; soft dorsal, anal, and ventral fins yellowish; spinous dorsal and pectorals darker; upper parts of preopercle and opercle each with a dark spot; top of head and tip of snout blackish.

S. picudilla is very closely allied to *S. borealis*. Its eye is, however, much larger (when specimens similar in size are compared), and the frontal groove is somewhat different. This species is found in the West Indies, on the coasts of Cuba, ranging southward to Bahia. It was not seen by us in Porto Rico, but is included on the authority of Dr. Stahl. Length 18 inches.

Sphyræna picudilla Poey, Memorias, II, 162, 1860, Havana; Stahl, l. c., 76 and 162, 1883; Jordan & Evermann, l. c., 824, 1896.

Family XXXI. POLYNEMIDÆ. The Thread-fishes.

Body oblong, compressed, and covered with rather large, loosely inserted, ctenoid scales. Lateral line continuous, continued on tail, usually forked, with a branch on each lobe. Head entirely scaly; snout more or less conical, projecting over mouth, which is rather large, inferior, with lateral cleft; premaxillary protractile, its basal process vertical; maxillary without supplemental bone, extending much beyond eye, which is anterior, lateral, rather large, with a well-developed, adipose eyelid. Villiform teeth on jaws, palatines, and sometimes on vomer. Pseudobranchiæ concealed. Branchi-

ostegals 7. Gill-membranes separate and free from isthmus. Gills 4, a slit behind fourth. Two separate dorsals, somewhat remote from each other, the first of 8 feeble but rather high spines, first and last spines very short, third longest; second dorsal equal to first in height but base somewhat longer, of soft rays only. Anal fin either similar to or much longer than soft dorsal; caudal fin rather long, widely forked. Second dorsal, anal, and caudal fins more or less covered with small scales; first three or four dorsal spines winged. Ventrals 1, 5, abdominal, but not far removed from pectorals; pectoral fins moderate, placed low, in two parts, the lower and anterior portion of several filiform articulated appendages, free from each other, used as organs of touch.

In the young the dorsal, caudal, and pectoral fins are dusky, the anal and ventral fins white; all the fins grow darker with age, pectorals usually becoming black, operculum blackish.

Bones of skull with a well-developed muciferous system, as in *Sciænidæ*. Basis cranii double, with muscular tube; post-temporal bifurcate; hypercoracoid with median foramen; superior pharyngeal bones 4. Pectoral actinosts divided; two of them normal, supporting pectoral fin, one longitudinal, without rays, and one a plate on the coracoid, supporting the pectoral filaments. Stomach caecal, with a few pyloric appendages. Air-bladder various, sometimes wanting. Vertebrae $10 + 14 = 24$.

- a. Anal fin much longer than soft dorsal, of about 30 rays; vomer without teeth; preoperculum entire; free filaments of pectoral longer than body..... POLYNEMUS
 aa. Anal fin not much longer than soft dorsal, of about 13 or 14 rays; vomer with teeth; preoperculum serrate; free filaments of pectoral mostly shorter than body..... POLYDACTYLUS, 47

Genus 47. POLYDACTYLUS Lacépède.

Anal fin not much longer than soft dorsal, of about 13 or 14 rays; vomer with teeth; preoperculum serrate; free filaments of pectoral mostly shorter than body. Teeth in villiform bands on both jaws, vomer, palatines, and pterygoids. Preopercle sharply serrated on its posterior margin, its angle with a scaly flap. Scales rather small, finely ctenoid. First dorsal with 7 or 8 feeble, rather high spines, first and last short. Soft dorsal and anal fins about equaling each other; pectoral filaments 3 to 9. Pyloric caeca in great numbers.

This genus is represented by numerous species occurring in the warm seas; but of four-American species only one is known from Porto Rico.

- a. Pectoral filaments 7..... *virginicus*, 73
 aa. Pectoral filaments 8 or 9..... *oceanus*

73. *Polydactylus virginicus* (Linnæus). "*Barbudo*"; *Barbu*.

Head 3.3; depth 3.25; eye 5; snout 4.8; maxillary 2.2; mandible 2; interorbital 4.2; D. VIII-1, 12; A. III, 13; pectoral 1.25; ventral 2.2; caudal 0.9; scales 7-60-10.

Body oblong, compressed, covered with large, loose, slightly ctenoid scales; the lateral line complete, nearly straight, bifurcating at base of caudal, the branches continued on tail; head completely scaled except about eye, which is entirely covered with an adipose eyelid; snout somewhat pig-like, overhanging the rather large mouth; maxillary reaching far beyond eye; lower jaw included, villiform teeth on jaws, vomer, and palatines; preopercle toothed; dorsal fins 2, well separated, the anterior of weak flexible spines, the first very short, third longest, 1.5 in head, the rest rapidly diminishing in length; pectoral with a large sheath of elongated scales above axil and with 7 long filaments, longer than head, inserted considerably in advance of base of pectoral; caudal deeply forked; all the fins more or less scaly.

Color in life: Whitish-olive above, dirty white below; spinous dorsal dark, soft dorsal and anal pale, with dark punctulations; pectoral with irregular black blotches, filaments white; ventrals dark, paler on margins. In spirits: Pale below, darker above; pectoral fin black; opercle with a dusty blotch in young.

The barbudo is an abundant and useful food-fish, found throughout the West Indies and north to Florida Keys, but not to Virginia. Twenty-three examples of all sizes up to 11 inches in length were collected at San Juan market, Palo Seco, Mayaguez, Puerto Real, Ponce, and Isabel Segunda.

Piracoaba, Maregrave, Hist. Nat. Brasil., 176, 1648, Brazil.

Polynemus virginicus Linnæus, Syst. Nat., ed. X, 317, 1758, America.

Polynemus mango Lacépède, Hist. Nat. Poiss., V, 413, 417, 418, 1803, America; after Linnæus.

Polydactylus plumieri Lacépède, Hist. Nat. Poiss., V, 419, 1803, Martinique; from a drawing by Plumier.

Polynemus americanus Cuvier & Valenciennes, Hist. Nat. Poiss., III, 393, 1829, Santo Domingo and Martinique.

Polynemus oligodon Günther, Cat., II, 322, 1860, Rio Janeiro.

Trichidion plumieri, Poey, Fauna Puerto-Riqueña, 334, 1881; Stahl, l. c., 78 and 164, 1883.

Polydactylus virginicus, Jordan & Evermann, l. c., 829, 1896.

Family XXXII. HOLOCENTRIDÆ. The Squirrel-fishes.

Body oblong or ovate, moderately compressed, covered with very strong ctenoid or spinous scales. Head with large muciferous cavities; eye lateral, very large; preorbital very narrow; mouth moderate, oblique; premaxillaries protractile; maxillary very large, with supplemental bone; bands of villiform teeth on jaws, vomer, and palatines. Opercular bones and membrane bones of head generally serrated or spinescent along their edges. Branchiostegals 8. Gill-membranes separate, free from isthmus. Gills 4, a slit behind fourth. Pseudobranchiæ present. Gillrakers moderate; no barbels. Sides of head scaly. Lateral line present. Dorsal fin very long, deeply divided, with about 11 strong spines depressible in a scaly groove; anal with 4 spines, the third longest and strongest; ventrals thoracic, with 1 spine and 7 rays; caudal deeply forked with sharp rudimentary rays or fulcra at base. Vertebrae about 27. Pyloric caeca 8 to 25. Air-bladder large, sometimes connected with organ of hearing. General color, red. Young with snout sharp and produced (constituting the nominal genera *Rhynchichthys*, *Rhamphoberyx*, and *Rhinoberyx* based on peculiarities of immature examples).

There are 4 genera and about 70 species of these gayly colored inhabitants of the tropical seas, abounding about coral reefs.

- a. Preopercle without conspicuous spine at its angle; air-bladder divided by a contraction, anterior part extending to the otocrane MYRIPRISTIS
- aa. Preopercle with a conspicuous spine.
- b. Suborbital arch armed with 3 long spines curved forward; scales undescribed..... PLECTRYPOPS
- bb. Suborbital arch simply serrated; scale moderate, 38 to 55.
- c. Mouth moderate, its length less than one-half head..... HOLOCENTRUS, 48
- cc. Mouth very large, considerably more than one-half head, lower jaw projecting..... FLAMMEO

Genus 48. HOLOCENTRUS (Gronow) Scopoli.

Body oblong, moderately compressed, ventral outline nearly straight, back a little elevated, tail very slender. Head compressed, narrowed forward. Operculum with a strong spine above, below which the edge is sharply serrated; a strong spine at angle of preopercle. Orbital ring, preorbital, preopercle, interopercle, subopercle, occiput, and shoulder-girdle with their edges sharply serrate. Mouth small, terminal, the lower jaw projecting in adult. In the young the snout is much produced. Maxillary broad, striate, with supplemental bone. Eye excessively large. Scales moderate, closely imbricated, posterior margin strongly spinous. Lateral line continuous. Dorsal deeply emarginate, spines usually 11, depressible in a groove; soft dorsal short and high; anal with 4 spines, first and second quite small, third very long and strong, fourth smaller; caudal widely forked, both lobes with rudimentary rays spine-like. Ventrals large, 1, 7, spine very strong.

The numerous species of this genus are remarkable for the development of sharp spines almost everywhere on the surface of the body. Of the nine known American species, two occur in Porto Rico.

- a. Preopercular* spine long, tapering, acute; third anal spine very long, more than half depth of body.
- b. Mouth moderate, lower jaw extending to below first one-third to one-half length of eye.
- c. Scales small, 48 to 55 in lateral line; upper lobe of caudal longer than lower, falcate; soft dorsal and anal elevated, pointed at tip, this character subject to variation..... *ascensionis*, 74
- cc. Scales moderate, about 45; depth of body 2.6 in length; membrane of front spines of dorsal black..... *siccifer*
- cd. Scales rather large, 38 to 42 in lateral line; caudal lobes nearly or quite equal.
- d. Maxillary extending to below middle of eye.
- e. Depth of body greater than length of head..... *coruscus*
- ee. Depth of body equal to length of head..... *brachypterus*
- dd. Maxillary extending to below first third of eye; dorsal with black markings..... *rexillarius*, 75
- bb. Mouth small, maxillary reaching to below first fourth of eye; upper lobe of caudal longer; cheek with a white blotch..... *osculus*
- aa. Preopercular spine short, flattish, notched at tip; third anal spine short, its length about one-third depth of body; soft dorsal and anal low, rounded..... *sancti-pauli*

74. *Holocentrus ascensionis* (Osbeck). "*Candil*"; Squirrel-fish.

(PLATE 3.)

Head 3.1; depth 3.3; eye 2.8; snout 4; maxillary 2.3; mandible 2; interorbital 2 in eye; D. XI, 15; A. IV, 10; scales 4-48-7, 6 before dorsal; caeca 25; vertebrae 11 + 16. Body considerably compressed,

*Little dependence can be placed on this key, as several species are imperfectly known and of doubtful validity.

back moderately elevated, ventral line nearly straight; mouth moderate, little oblique, maxillary reaching middle of eye; caudal peduncle long and slender, least width 2 in its depth. Fins all well developed, longest dorsal spine about 2 in head; third anal spine very strong, about 2 in head; longest dorsal rays 1.2 in head; longest anal ray 1.8; pectoral short, 1.6 in head; ventral long, reaching vent, 1.1 in head; caudal well forked, upper lobe the longer, about 1.1 in head; head very rugose; suborbital coarsely and irregularly dentate; opercle and preopercle strongly dentate, a large sharp spine at angle of preopercle, its free portion about 1.5 in eye; a broad flat spine at upper edge of opercle; scales coarse and strongly ctenoid.

Color in life: Chiefly bright rosy-red, paler below; shining longitudinal streaks along rows of scales; fins light-red, spinous dorsal largely golden-olive, its edge scarlet; head very red above, a white bar descending obliquely backward from eye. All these colors fade in alcohol, and the general color becomes a greenish-white, with steel-colored iridescence.

The candil frequents the rocks and reefs throughout the West Indies and is especially abundant in Cuba. It is common about Porto Rico, and attracts attention at once by reason of its bright color and the exceeding sharpness and completeness of its armature. It is reputed "malo" by the native, but is occasionally found in market. Length 1 to 2 feet. Specimens were obtained at San Juan, Aguadilla, Puerto Real, Arroyo, Hucares, Isabel Segunda, and San Geronimo.

Perca ascensionis Osbeck, *Iter Chinensis*, 388, 1771, Ascension Island.

Bodianus pentacanthus Bloch, *Aust. Fische*, IV, 40, pl. 225, 1790, Brazil.

Holocentrus sago Bloch, l. c., 61, pl. 232, 1790, Africa.

Sciæna rubra Bloch & Schneider, *Syst. Ichth.*, 82, 1801; after *Perca marina rubra* of Catesby.

Amphiprion matejuelo Bloch & Schneider, l. c., 206, 1801, Cuba; after *Matejuelo* of Parra.

Bodianus jaguar Lacépède, *Hist. Nat. Poiss.*, IV, 286, 1202, Brazil; after *Jaguaraca* of Maregrave.

Holocentrum longipinne Cuvier & Valenciennes, *Hist. Nat. Poiss.*, III, 185, 1829, Martinique, Santo Domingo, Porto Rico, St. Thomas, and Havana.

? *Holocentrus striatus* Gronow, *Cat. Fishes*, ed. Gray, 173, 1854, Antilles; name preoccupied.

? *Holocentrus rostratus* Gronow, l. c., 173, 1854, near the Equator; young example, not certainly identifiable.

Holocentrum matejuelo, Poey, *Fauna Puerto-Riqueña*, 322, 1881; Stahl l. c., 76 and 162, 1883.

Holocentrus ascensionis, Jordan & Evermann, l. c., 848, 1896.

75. *Holocentrus vexillarius* Poey.

Head 2.8; depth 2.6; eye 2.7; snout 4.4; maxillary 3; mandible 2.25; interorbital 3.5; D. x1, 13; A. 1v, 9; pectoral 1.4; ventral 1.6; caudal 1.5; scales 4-40-7.

Body oblong, back little elevated, anterior profile steep and nearly straight from tip of snout to above eye, where there is an abrupt angle, rest of profile to nape being low and straight; interorbital wide, slightly concave; top of head behind eyes with about 9 serrated bony ridges on each side; eye very large, considerably wider than interorbital space; snout short; maxillary reaching front of eye (to first third of eye in example 2 inches long); preorbital with a recurved spine in front, retrorse teeth back of this, finer and more numerous serrations continuing nearly halfway around eye; a pair of strong equal spines near upper angle of opercle; several much smaller spines above, of which 3 are somewhat enlarged, longest dorsal spines 2.1 in head; anterior rays of dorsal somewhat produced, 2 in head; scales at base of spinous dorsal with their serrations next the fin enlarged; ventrals not reaching vent; second anal spine longest and much the strongest, 1.5 in head; caudal well forked; lobes nearly equal.

Color in spirits: Faded-rosy, 6 narrow longitudinal very dark-purple bands between each 2 rows of scales on dorsal half of side; below, brown punctulations in same position and scattered more or less over whole body; axil of pectoral very dark-purple, the color sharply circumscribed; membrane of spinous dorsal white behind each spine and purple-black in front, except first 2 spines, the membranes of which are chiefly black; other fins pale.

Two examples, 2 and 4 inches in length, from Ponce and Guanica, are identical with specimens of *Holocentrum riparium* from Cuba in the U. S. National Museum. Following Jordan & Evermann, we include *H. riparium* in the synonymy of *H. vexillarius*, though these species may prove to be distinct.

Holocentrum vexillarium Poey, *Memorias*, II, 158, 1860, Cuba.

Holocentrum productum Poey, *Synopsis*, 300, 1868, Matanzas; based on a young individual 3 inches long.

Holocentrum riparium Poey, *Enumeratio*, 37, 1875, Cuba.

Holocentrus vexillarius, Jordan & Evermann, l. c., 852, 1896.

Family XXXIII. MULLIDÆ. The Surmulletts.

Body elongate, slightly compressed, covered with large scales, which are usually slightly ctenoid; lateral line continuous, the pores often branched; large scales on head; upper profile of head more or less parabolic. Mouth small, low, subterminal; teeth mostly small, variously placed; no canines, incisors, nor molars. Premaxillaries somewhat protractile; maxillaries thin, nearly as broad at base as at tip, without supplemental bone, partly hidden by broad preorbital. Preopercle entire or slightly serrate; opercle unarmed, or with a single spine. Eye moderate, placed high; branchiostegals 4; pseudobranchiæ present; 2 long unbranched barbels at throat, attached just behind symphysis of lower jaw. Dorsal fins 2, remote from each other, both short, the first of 6 to 8 rather high spines, which are depressible in a groove; anal short, similar to soft dorsal, with 1 or 2 small spines; ventrals thoracic, 1, 5. Air-bladder usually present, simple. Vertebrae $9 + 14 = 23$; stomach siphonal; pyloric cæca about 20.

Species about 40, referable to 5 closely related genera, found in all tropical seas, some species straying northward. Many of the species are highly valued as food, especially the European *Mullus barbatus* and *M. surmuletus*. The family is a very natural one and not closely related to any other. It bears some superficial likeness to the *Sciænidæ* and *Cheilodipteridæ*, but this may not show real affinity. The singular barbels appear also in *Polymixiidæ*, but in that family the ventral rays are numerous, as in *Berycidæ*. The small number (4) of the branchiostegals is found both in *Mullidæ* and *Polymixiidæ*.

- a. Teeth on lower jaw, vomer, and palatines; upper jaw toothless; bone which forms a downward hook over maxillary strongly developed; interorbital space flat and wide; opercle without spine MULLUS
 aa. Teeth on both jaws; vomer and palatines toothless; bone which forms a downward hook over maxillary moderately developed; interorbital space rather narrow; opercle ending in a single spine UPENEUS, 49

Genus 49. UPENEUS Cuvier. The Goat-fishes.

Body oblong, compressed; mouth moderate, nearly horizontal, low, jaws subequal; eye large high, posterior; opercle short, deep, with posterior spine; both jaws with rather strong, unequal teeth, in one or two series in each jaw; no teeth on vomer or palatines; lips well developed; bone which forms a hook over the maxillary less developed than in *Mullus*; interorbital space concave and narrow; opercle ending in one spine; barbels nearly as long as head; scales very large, somewhat ctenoid; lateral line continuous, its tubes ramifying on each scale; head covered with large scales; first dorsal with about 7 spines; anal with 2, the first very short; caudal fin forked.

Species numerous in the tropical seas; two known from Porto Rico.

- a. Teeth in both jaws uniserial (or irregularly biserial above); all teeth coarse and distinct; eye 4 in head; barbels 1.33 in head. Scales 31; depth 4 in length; side with three black blotches *maculatus*, 76
 aa. Teeth of both jaws biserial, at least in front.
 b. Dorsals and caudal with dark crossbands *parvus*, 77
 bb. Dorsals and caudal plain-yellow *martineus*, 78

76. *Upeneus maculatus* (Bloch). "Salmonete"; Red Goat-fish.

(PLATE 4.)

Head 3.2; depth 3.7; eye 4 to 5; snout 1.8; maxillary 3.2; mandible 2.4; interorbital 3.7; preorbital 3.33; D. VIII-1, 8; A. II, 7; pectoral 1.5; ventral 1.4; caudal 1.3; scales 3-30-5.

Body elongate, little compressed, tapering posteriorly to the long and slender caudal peduncle; anterior profile strongly arched, with an elevation in front of eye; snout very long; eye small, high and posterior; mouth small, maxillary not nearly reaching eye; teeth strong, uniserial, bluntly pointed, some of those in front of upper jaw bent variously sidewise and forward; throat with two long barbels reaching to preopercular margin or beyond. Scales large, very finely ctenoid. Dorsal fins 2, well separated, the spines slender and flexible; caudal deeply forked, lobes equal or upper very slightly the longer.

Color: In life, everywhere above red, merging into light-yellow on sides, becoming pale-greenish below; bluish oblique streaks and bars on head; several longitudinal rows of light-blue round spots, much smaller than pupil, on sides, the two rows above lateral line plainest; about four diffuse blotches of darker red than the surrounding ground-color on sides, the first just under and partly on anterior end of lateral line, second under first dorsal, third under front of second dorsal, fourth just behind second dorsal, all on or near lateral line; there may be an additional faint one on peduncle; spinous dorsal light-red near base, yellowish outwardly; soft dorsal mostly pale-bluish, especially near edge,

with some light-yellow on membrane and red on middle of rays; pectoral chiefly yellow, with red on rays; ventral pale-blue with streaks of red and yellow on first rays; anal pale-reddish, nearly pale; caudal pale-blue, lower base yellow, upper reddish and yellow; barbels pink near base, yellow outwardly; lips and throat pinkish. In spirits the specimens become pale below and on sides, dark above, the bright colors fading quickly; the red blotches along lateral line become dark, sometimes entirely fading. In young individuals some color occasionally persists in spirits, becoming pink.

This species is found at Key West and the Tortugas, through the West Indies to Brazil; in the West Indies it is known from Cuba, Jamaica, Porto Rico, and Martinique. It occurs in abundance in Porto Rico, where it is extensively used as food and considerably esteemed. It is one of the most common species that one sees carried about the streets by the local fishermen. Fifty-one individuals, from 2.25 to 9.5 inches long, collected at nearly every collecting station; also from San Geronimo.

Mullus maculatus Bloch, Ichthyologia, 348, 1793, Brazil.

Upeneus punctatus Cuvier & Valenciennes, Hist. Nat. Poiss., III, 482, 1829, Martinique.

Upeneus maculatus, Jordan & Evermann, l. c., 858, 1896.

77. *Upeneus parvus* Poey.

D. VII-1, 8; A. II, 6; scales 2.5-40-6. Snout short and decurved; maxillary reaching front of eye; barbels reaching angle of preopercle, on jaws only. Teeth conical, very small, on anterior part of jaws in two series; lateral teeth in a single series; all of the teeth obtusely conic and distinct from each other. Vermilion above, fading into white below; a yellow longitudinal band along side, with similar narrower streaks below; ventrals and anal yellow; other fins whitish, with dusky crossbands, 3 on first dorsal, 2 on second, and 5 on each caudal lobe. Known only from the type, which was obtained by Poey in Cuba, and from a specimen recorded from Porto Rico by Dr. Stahl.

Upeneus parvus Poey, Memorias, I, 226, 1851, Cuba.

Upeneoides parvus Stahl, l. c., 76 and 162, 1883.

78. *Upeneus martinicus* Cuvier & Valenciennes. *Yellow Goat-fish; Salmonete Amarilla.*

(PLATE 5.)

Head 3.3; depth 4; eye 3.45; snout 2.35; maxillary 3.1; mandible 2.6; interorbital 3.15; preorbital 6; D. VIII-1, 8; A. II, 6; pectoral 1.6; ventral 1.4; caudal 1.1; scales 2-37-6. Resembling *U. maculatus* in form, but with slightly larger eye, smaller scales, and weaker dentition arranged in more than one series.

Color in life: Body with shades of pale-blue and pink or pale-red, the latter chiefly above, the blue below; a straight yellow band from eye to base of upper caudal rays; a black vertical bar at base of caudal; head with yellow streaks and reddish patches; pectorals red, ventrals, anal, and caudal reddish near base, outer part yellow; dorsals yellow, plainest near tips. In spirits the colors fade, becoming dark above, pale below, fins all pale.

This species reaches the length of a foot. It is found in the West Indies north to Florida, and is known from Key West, Jamaica, Cuba, Martinique, and Porto Rico, but is less abundant in Porto Rico than *U. maculatus*. One example, 6.5 inches long, obtained at Palo Seco.

Upeneus martinicus Cuv. & Val., Hist. Nat. Poiss., III, 483, 1829, Martinique; Jordan & Evermann, l. c., 859, 1896.

Upeneus balteatus Cuvier & Valenciennes, Hist. Nat. Poiss., III, 484, 1829, Cuba; young.

Upeneus flavovittatus Poey, Memorias, I, 224, 1851, Cuba; adult.

Family XXXIV. SCOMBRIDÆ. The Mackerels.

Body elongate fusiform, not much compressed, covered with minute cycloid scales; scales anteriorly sometimes forming a corselet. Lateral line present, its course undulate. Head subconic, pointed anteriorly. Mouth rather large, with lateral cleft; premaxillaries not protractile; maxillary without supplemental bone; jaws with sharp teeth, large or small. Vomer and palatines toothed or not. Preopercle entire; opercle unarmed. In the very young the preopercle is armed with radiating spines, which are later absorbed and lost. Gill-openings very wide, membranes not united, free from isthmus. Gillrakers usually long. Pseudobranchiæ present, large. Gills 4, a slit behind fourth. Branchiostegals 7. Dorsal fins 2, the first of rather weak spines, depressible in a groove, second similar to the anal; elevated anterior lobe always distinct; anal spines weak; last rays of dorsal and anal detached and separate, forming in each case a series of finlets; caudal peduncle extremely slender, keeled, caudal lobes abruptly diverging, falcate, fin adapted for rapid motion; ventral fins well developed, thoracic, 1, 5.


Vertebrae in greater number than in the *Carangidae*, the number ranging from 31 to 66. First upper pharyngeal present, without teeth; second with teeth; third and fourth coossified, with teeth; lower pharyngeals separate. Stomach sac-shaped. Pyloric caeca numerous. Air-bladder small, sometimes absent. Coloration metallic, often brilliant, the prevailing shade steel-blue.

The *Scombridae* comprise about 12 genera and 60 species; of the 8 American genera only 2 are represented in Porto Rican waters. These fishes are inhabitants of the high seas, all having a wide range, and many of them being cosmopolitan. Most of them are valued as food-fishes, the flesh being firm and oily, but sometimes coarse.

- a. Dorsal spines 10 to 16, gills normal, the laminae not forming a network; teeth entire.
- b. Body scaleless, excepting about lateral line and corselet; abdominal vertebrae with their lower foramina enlarged and a portion between vertebrae proper and hamapophyses developed in form of a network or trellis.
- c. Dorsals well separated, interspace more than half head; corselet well developed; teeth small, some present on vomer, none on palatines; gillrakers long, slender, and numerous; pectorals rather high; vertebrae 39. AUXIS, 50
- cc. Dorsals contiguous, interspace more than 5 in head; palatine teeth villiform; pectorals low.
- d. Vomer toothless; dorsal spines 15 or 16; vertebrae 38 GYMNOSARDA
- bb. Body wholly covered with small scales, those on corselet and lateral line sometimes larger; dorsal spines 14 to 26; vertebrae normally formed, not as in *Auxis* and *Gymnosarda*.
- e. Teeth of jaws slender, subconical, little, if at all, compressed; gillrakers numerous; corselet distinct; pectoral inserted low.
- f. Vomer and palatines with villiform or sand-like teeth; body robust, not compressed; vertebrae 39 to 41.
- g. Pectoral short, not reaching much beyond tip of the moderate ventral; size enormous THUNNUS
- gg. Pectoral very long, ribbon-shaped, reaching much beyond front of anal; size moderate GERMUS
- ff. Vomer toothless; palatines with a single row of rather strong, conical teeth; body elongate, slightly compressed; vertebrae 50 to 54 SARDA
- ce. Teeth of jaws strong, subtriangular or knife-like, more or less compressed; villiform teeth on vomer and palatines; gillrakers comparatively few; corselet obscure; pectorals inserted near level of eye; dorsal spines 14 to 18; body elongate, compressed; head short; snout short; vertebrae 45 SCOMBEROMORUS, 51

ACANTHOCYBINE:

- aa. Dorsal spines about 25; gills with laminae forming a network, as in *Xiphias*; teeth large, compressed, serrated.
- h. Dorsal spines 24 to 26; body elongate, fusiform; snout long; vertebrae 32 + 34 = 66 ACANTHOCYBIUM

FIG. 27.—*Auxis thazard*.

Genus 50. AUXIS Cuvier. The Frigate Mackerels.

Body oblong, plump, mostly naked posteriorly, anteriorly covered with small scales, those of pectoral region enlarged, forming a corselet. Snout very short, conical, scarcely compressed. Mouth rather small, jaws equal. Teeth very small, mostly in a single series, on jaws and vomer only. Tail very slender, depressed, with a rather large keel on each side. First dorsal short, separated from second by a considerable interspace. Second dorsal and anal small, each with 7 or 8 finlets. Pectorals and ventrals small. No air-bladder. Branchiostegals 7. Pyloric caeca dendritical. Gillrakers very long and slender, numerous. Vertebrae 39 in number, peculiarly modified, essentially as in *Gymnosarda*.

One pelagic species, widely distributed.

79. *Auxis thazard* (Lacépède). "*Albacora*"; *Frigate Mackerel*.

Head 3.8; depth 4.4; eye 6; snout 3.7; mandible 2.2; interorbital 3.2; D. x-12-viii; A. 13-vii; pectoral 1.8; ventral 2.25; caudal 1.4.

Body elongate, robust, scaleless, save for a corselet of moderate and very small scales about

anterior part of trunk and along lateral line; caudal peduncle long and slender, with lateral keel; weak conical teeth in jaws only; first dorsal of flexible spines, the first three elevated, the rest graduated to last, which is very short; soft dorsal and anal small, somewhat falcate, followed by detached finlets; caudal widely forked.

Color in spirits: Bluish above, very deep purple, almost black, on upper part of head; oblique dark bars and stripes, somewhat as in *Scomber*, on sides of back; silvery white below; several large, regular, dark blotches, not sharply circumscribed, on the lower part of side of trunk anteriorly; pectoral and ventral purple, black on inner side, on outer side pectoral paler, ventral white.

This species is found in all warm seas, occasionally northward to Cape Cod. It is very erratic in its movements, swimming in large schools. It rarely reaches the coasts of the United States, but occasionally comes in immense numbers. It is a poor fish, of little value as food. It is probably not rare about Porto Rico; examples were seen at Arecibo, where one 15 inches long was obtained.

Scomber thazard Lacépède, Hist. Nat. Poiss., II, 9, 1802, between 6° and 7° S. lat., off coast of New Guinea.

Scomber rochei Risso, Ichth. Nice, 165, 1810, Nice.

Scomber bisus Rafinesque, Caratteri, etc., 45, 1810, Palermo.

Thynnus rocheanus Risso, Eur. Mérid., III, 417, 1827, Nice.

Auxis vulgaris Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 139, 1831, Mediterranean.

Auxis tapinosoma Bleeker, Fauna Japon., 408, 1854, Japan.

Auxis thynnoides Bleeker, Ternate, V, 301, 1855, Ternate.

Auxis thazard, Jordan & Evermann, l. c., 867, 1896.

Genus 51. SCOMBEROMORUS Lacépède.

Body elongate, wholly covered with rudimentary scales, which do not form a distinct corselet. Head pointed, comparatively short and small. Mouth wide, the strong teeth in the jaws more or less compressed or knife-shaped; villiform or sand-like teeth on the vomer and palatines; maxillary not concealed by preorbital. Gillrakers few. Caudal peduncle with a single keel. Spinous dorsal low, of 14 to 18 feeble spines. Soft dorsal and anal short, similar, somewhat elevated and falcate, each followed by 7 to 10 finlets; ventrals small; pectorals moderate, near level of eye. Air-bladder present. Vertebrae normally formed, 45 in number.

Fishes of the high seas; graceful in form and beautiful in color, and among the best of food-fishes.

- a. Soft dorsal inserted in advance of anal *maculatus*, 80
 aa. Soft dorsal inserted over anal.
 b. Body deep, the depth about 5 in length; teeth about 40 in each jaw *regalis*, 81
 bb. Body more slender, the depth about 6 in length *cavalla*, 82

80. *Scomberomorus maculatus* (Mitchill). "*Carita*"; *Spanish Mackerel*.

(PLATE 6.)

Head 4.5; depth 4.5; D. XVII-18-IX; A. II-17-IX; maxillary 1.8 in head; eye 4.75; pectoral 1.75; ventral 4.5; dorsal and anal lobes subequal, 2. Body elongate, its dorsal and ventral outlines equal; profile straight from snout to dorsal; head small and pointed; mouth large, oblique, jaws equal; maxillary reaching posterior margin of orbit; teeth large, compressed, and sharp, their formula being 24-24 to 32-32; gillrakers 2 + 11. Soft dorsal inserted in advance of anal a distance about equal to diameter of eye; lateral line undulating, with about 175 pores.

Color silvery, bluish above; sides with many elliptical spots of dull orange-color, two rows of these spots below lateral line and one row above; spinous dorsal white at base, black above; soft dorsal tinged with yellowish, its margins black; anal white; posterior side of pectoral black, anterior side yellowish with black borders; caudal blackish.

This species is found on both coasts of North America, appearing in large but very irregular schools in the Gulf of Mexico and along the Carolina coast; ranging north in the fall as far as Cape Ann, and south to Brazil. It is rare or unknown in Cuba, but is known from Jamaica and Porto Rico. It reaches a weight of 8 or 9 pounds and is one of the very best food-fishes in the United States.

Scomber maculatus Mitchill, Trans. Lit. & Philos. Soc. N. Y., I, 1815, 426, New York.

Scomberomorus maculatus, Jordan & Evermann, l. c., 874, 1896.

81. *Scomberomorus regalis* (Bloch). *Sierra*; *Pintado*.

Head 4.25; depth 5.2; eye 5.8; snout 2.5; maxillary 1.8; mandible 1.6; interorbital 3.5; D. xviii-14-ix; A. ii-15-viii; pectoral 1.9; ventral 4.2; caudal 1.1. Lateral line descending gradually from opposite last dorsal spines to caudal peduncle, slightly undulate; maxillary reaching posterior border of orbit.

Color in spirits: Silvery below, dark-blue above; side with a blackish longitudinal band from base of pectoral nearly to base of caudal, crossing lateral line under soft dorsal, somewhat interrupted posteriorly; below this a row of oblong dark spots forming an interrupted band; a few other faint spots at border of dark color of back; anterior portion of spinous dorsal black.

Cape Cod to Brazil; not very common on our south Atlantic coast, but abundant about Cuba; known also from Jamaica, Martinique, and Porto Rico. One 16 inches long from Puerto Real, and others seen.

Scomberomorus regalis Bloch, Ichthyol., pl. 333, 1795, Martinique; after a drawing by Plumier.

Scomberomorus plumieri Lacépède, Hist. Nat. Poiss., III, 292, 1806, Martinique; after Aubriet's copy of drawing by Plumier.

Cybitum acerrum Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 186, 1831, Cuba.

Cybitum regale, Poey, Fauna Puerto-Riqueña, 331, 1881; Stahl, l. c., 77 and 163, 1883.

Scomberomorus regalis, Jordan & Evermann, l. c., 875, 1896.




FIG. 28.—*Scomberomorus regalis*.

82. *Scomberomorus cavalla* (Cuvier & Valenciennes). *King-fish*; *Cero*; *Cavalla*; *Sierra*.

Head 5; depth 6; eye large; 2 in snout. D. xv-1, 15-viii; A. ii, 15-viii. Mouth large, maxillary reaching to below eye. Lateral line descending abruptly below second dorsal. Teeth triangular, strongly compressed, about 30 on each jaw. Pectoral 5 in body. Gillrakers very short; less than one-third diameter of eye, about 8 below angle.

Adult iron-gray, nearly or quite immaculate; young with sides of body marked with darker yellowish spots; spinous dorsal without black blotch anteriorly.

Found in the tropical Atlantic, in the open seas, coming in immense numbers to Florida Keys and Charleston, ranging north to Cape Cod and south to Africa and Brazil; very common on our south Atlantic coast, especially among the Florida Keys, the catch at Key West very large. One of the best food-fishes of the Florida coast. Its flesh is firm and of excellent flavor. It usually appears in large numbers from November until April, when it is caught by trolling. The usual weight is about 10 pounds, sometimes reaching 50 pounds. The largest of which there is any record dressed 52 pounds, and 40 pounds is not an unusual weight. Said to school at spawning time, which is believed to be late in the winter.

According to Mr. William H. Abbott, who studied the fisheries in 1891, the average weight of king-fish, as caught by the fishermen of Key West, is about 6 pounds. The larger, weighing from 15 pounds upward, are never as abundant as those weighing under 15. When the fisherman desires to catch large king-fish, he directs his course to the inshore grounds, lying in about 3 fathoms of water and from 1.5 to 3 miles from shore, where the water is muddy; and when small ones are desired, the fishing is done farther offshore along the edge of the Gulf Stream, where the water is much clearer. They are almost invariably found in two separate schools. The Florida spawning-grounds of the king-fish are "down the bay." The first of the winter a great many of the fish have large roes, but it is very seldom that one is taken that has a roe fully matured. If the weather has been very cold in the bay early in the fall, the king-fish will leave before they have spawned, and it is during such seasons that fish containing ripe spawn are most frequently taken.

On the coast of southern Florida, especially off Indian River, Biscayne Bay, and Key West, there is no fish, excepting possibly the tarpon, which affords more sport for the angler than the king-fish, nor is there any to whose season the angler and the commercial fishermen look forward with greater interest and anticipation of pleasure and profit.

Though not obtained by us in Porto Rico it doubtless occurs there, and we include it on the authority of Poey.

Cybium cavalla Cuvier, Règne Animal, ed. 2, II, 200, 1829, Brazil.

Cybium immaculatum Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 191, 1831; no locality given.

Cybium caballa, Poey, Fauna Puerto-Riqueña, 331, 1881.

Scomberomorus cavalla, Jordan & Evermann, l. c., 876, 1896.

Family XXXV. TRICHIURIDÆ. The Cutlas-fishes.

Body extremely elongate, band-shaped, naked, tapering to a point, the ventral fins imperfect or wanting, and the spinous and soft parts of the dorsal fin not differentiated. Mouth wide, jaws armed with strong unequal teeth. Premaxillaries not protractile. Pseudobranchiæ present. Gills 4, a slit behind fourth; gill-membranes separate, free from isthmus; lateral line present. Dorsal fin very long, low, usually continuous, rays all similar. Caudal fin absent; anal fin long and low, scarcely rising above surface of skin. Ventrals thoracic, rudimentary (*Eupleurogrammus*) or wanting. Vertebrae in greatly increased number, about 160. Air-bladder present. Pyloric caeca numerous.

Surface fishes of the tropical seas, comprising 2 genera and about 6 species; very close to the *Lepidopidae* and *Gempylidae*, differing from the former chiefly in the absence of a caudal fin, the last stage in the progressive reduction of parts seen in these groups.




FIG. 29.—*Trichiurus lepturus*.

Genus 52. TRICHIURUS Linnaeus. The Hair-tails.

Body extremely elongate, band-like, tail very slender, tapering to a fine point, without caudal fin. Head long, with very wide mouth, jaws armed with unequal and very strong teeth; upper jaw with about four long, strongly compressed barbed teeth; teeth on palatines, none on vomer. Lower jaw the longer, preorbital covering cleft of mouth posteriorly. Dorsal fin single, low, occupying whole of back, spines not distinguishable from soft rays; anal very long, its base more than half length of body; composed of detached spines, which are very short, nearly hidden in skin, anterior directed backward, posterior forward; ventral fins wanting; pectorals small. No scales. Lateral line decurved, concurrent with belly. Vertebrae 39 + 120; ribs excessively frail. Color, silvery.

Voracious fishes of the high seas; reaching a considerable size.

83. *Trichiurus lepturus* Linnaeus. "Machete"; Cutlas-fish; Scabbard-fish; Silver-fish; Sable; Savola.

Head 8; depth 15.5; eye 6.3 (2.1 in snout); snout 2.8; mandible 1.8; interorbital 6.8; D. about 135; A. about 100. Body scaleless, greatly elongate and compressed, band-like, tail tapering to a fine point, without caudal fin; jaws armed with very strong long teeth, larger ones of upper jaw barbed; lower jaw pointed; projecting preorbital with radiating striae; dorsal fin as long as back; anal fin of very short embedded spines, anterior directed backward, posterior forward; ventrals wanting. Color, bright silvery.

Found in warm seas, chiefly of the western Atlantic, north to Massachusetts; occasionally in Lower California (Streets); common in the West Indies, swimming near the surface, where it becomes easily benumbed with cold. It is probably not rare about Porto Rico. Two examples, 25 and 15 inches in length, were collected at Palo Seco and Mayaguez.

Trichiurus lepturus Linn., Syst. Nat., ed. X, 246, 1758, America; after *Lepturus* of Arfedi; Jordan & Evermann, l. c., 889, 1896.
Lepturus lepturus, Poey, Fauna Puerto-Riqueña, 333, 1881; Stahl, l. c., 162, 1883.

Family XXXVI. CARANGIDÆ. The Pampanos or Pompanos.

Body more or less compressed and often elevated, sometimes naked, or more usually covered with small, thin, cycloid scales. Head compressed, occipital keel prominent, usually trenchant. Mouth of varying size, dentition various, teeth generally small; premaxillaries usually protractile; maxillary with or without a supplemental bone; preopercle usually entire in adult, in the very young armed with three or more spines. Lateral line complete, anteriorly arched, posterior part straight, sometimes armed with bony plates. Dorsal fins more or less separated, the spinous part rather weak, the spines usually depressible in a groove; anal fin long, similar to soft dorsal, always preceded by two stiff spines, usually separate, but in young often more or less connected with fin or with each other, these sometimes disappear with old age, and sometimes the spinous dorsal also vanishes; often a procumbent spine before dorsal fin; ventral fins thoracic, well developed, 1, 5; caudal peduncle very slender, fin widely forked; pectoral fins narrow. Gill-openings very wide, membranes usually not united, free from isthmus. Gills 4, a slit behind last. Gillrakers usually long. Branchiostegals commonly 7. Air-bladder present, often bifurcate behind. Pseudobranchiæ large, present in all our genera, sometimes disappearing with age. Œsophagus unarmed. Pyloric caeca generally numerous. Vertebrae fewer than in the *Scombridae*, usually 10 + 14 = 24 in number. First superior pharyngeal without teeth; second, third, and fourth separate, with teeth; lower pharyngeals separate.

Coloration generally metallic and silvery or golden.

The *Carangidae* include 29 genera and about 200 species, abounding in warm seas, often moving northward in summer, like the *Scombridae*. They swim swiftly, often with the dorsal fin above the surface of the water. Most of the species are widely distributed, and nearly all are valued as food.

SCOMBEROIDINÆ:

- a. Premaxillaries not protractile (except in very young). Pectoral fins short, rounded; soft dorsal similar to anal, each much longer than abdomen; lateral line unarmed.
b. Maxillary without supplemental bone; no pterygoid teeth; scales linear, embedded OLIGOPLITES, 53
aa. Premaxillaries protractile.

SERIOLINÆ:

- c. Anal fin much shorter than soft dorsal, its base not longer than abdomen; pectoral fin short, not falcate; maxillary with a distinct supplemental bone.
d. Dorsal spines low and weak.
e. Dorsal and anal fins without finlets.
f. Membrane of dorsal spines disappearing with age..... NAUCRATES
ff. Membrane of dorsal spines persistent..... SERIOLA, 54
cc. Dorsal and anal fins each with a detached two-rayed finlet..... ELAGATIS
ce. Anal fin about as long as soft dorsal, its base longer than abdomen.
g. Maxillary with a supplemental bone; lateral line arched anteriorly, usually armed posteriorly; pectoral long, falcate.

CARANGINÆ:

- h. Dorsal outline more strongly envered than ventral outline.
i. Dorsal and anal each with a single detached finlet; body slender..... DECAPTERUS, 55
ii. Dorsal and anal without finlets.
j. Lateral line with well-developed scutes for its entire length; body elongate..... TRACHURUS
jj. Lateral line with scutes on its straight posterior portion only (these sometimes very few and small, especially in those species with body much compressed).
k. Shoulder-girdle with deep cross-furrow at its junction with isthmus, above which is a fleshy projection; body elongate..... TRACHUROPS, 56
kk. Shoulder-girdle normal, its surface even; body deeper.
l. Body oblong or more or less elevated, not as below.
m. Teeth of jaws in few series or in one series, unequal, or at least not forming villiform bands, outer series above usually enlarged, lower teeth usually uniserial.
n. Maxillary very narrow, its greatest width scarcely one-fourth eye; head small, lateral line strongly arched in front; teeth uniserial, those on palatines and vomer minute or obsolete..... HEMICARANX
mm. Maxillary broad; head rather large; vomer and palatines with teeth..... CARANX, 57

- mm.* Teeth of jaws equally small or wanting, forming villiform bands if present; maxillary broad; body compressed; spinous dorsal weak, usually disappearing with age.
- o.* Soft dorsal lobe very high, filamentous..... ALECTIS
- oo.* Soft dorsal lobe low..... HYNNIS
- ll.* Body broad-ovate, very strongly compressed, its outlines everywhere trenchant, anterior profile nearly vertical; scutes almost obsolete..... VOMER, 58
- jjj.* Lateral line without any scutes; body short and elevated, strongly compressed..... SELENE, 59
- CHLOROSCOMBRINÆ:
- hh.* Dorsal outline less strongly curved than ventral; body much compressed, its outlines everywhere trenchant; armature of lateral line obsolete, or nearly so..... CHLOROSCOMBRUS, 60
- TRACHINOTINÆ:
- gg.* Maxillary without supplemental bone; anal fin similar to soft dorsal, its base much longer than abdomen; tail unarmed; pectoral short, not falcate.
- p.* Forehead convex; teeth small or deciduous..... TRACHINOTUS, 61

Genus 53. OLIGOPLITES Gill. Leather-jackets.

Body compressed, oblong or lanceolate. Caudal peduncle slender, not keeled. Head short, compressed, acute. Occipital keel sharp. Mouth rather large, with small sharp teeth in bands on jaws, tongue, vomer, and palatines, none on pterygoids. Jaws about equal, upper not protractile, except in very young, in which it is movable as in other *Carangidae*; maxillary very narrow, without distinct supplemental bone. Gillrakers rather long. Scales small, linear, and extremely narrow, embedded in skin at different angles. Lateral line unarmed. Dorsal spines rather strong, 3 to 5 in number, nearly free in adult; second dorsal very long, its posterior rays pencillated and nearly or quite disconnected, forming finlets; anal rather longer than soft dorsal, much longer than abdomen, its last rays forming similar finlets; anal spines strong; ventral fins depressible in a groove; pectoral fins very short.

Species few, occurring in the tropical seas of America.

- a.* Maxillary not reaching posterior border of eye; lowest infraorbital bone usually narrower than one above it. *saurus*, 84
- aa.* Maxillary reaching beyond posterior border of eye; lowest infraorbital bone usually broader than one next above it. *salicis*

84. *Oligoplites saurus* (Bloch & Schneider). "*Zapatero*"; *Quiebra*; *Runner*; *Leather-jack*.

(PLATE 7.)

Head 4.8; depth 3.75; eye 3.6; snout 3.25; maxillary 1.9; mandible 1.7; interorbital 3; D. v-i, 20; A. ii-i, 20; pectoral 1.7; ventral 1.9; caudal 1.

Body elongate, lanceolate, and much compressed; back gently elevated; caudal peduncle very slender, somewhat higher than thick; head small and pointed; anterior profile slightly concave in occipital region; mouth wide, maxillary reaching nearly to posterior border of eye, jaws subequal or lower very slightly projecting; villiform teeth on vomer, palatines, and tongue; bands of larger teeth in jaws; head naked, body covered with very narrow, linear, embedded scales, placed at various angles, making a tough leathery integument; fins small, dorsal spines nearly free, soft dorsal and anal long and low, anterior rays somewhat elevated, others expanded and nearly disconnected from each other, forming finlets, as in the mackerel; ventrals depressible in a groove; caudal widely forked.

Color, blue above, silvery below, the demarcation fairly abrupt, corresponding with a line from anterior end of lateral line to base of upper caudal lobe; fins yellow.

The zapatero inhabits both coasts of tropical America, extends north to New York and Lower California, and is very common in the West Indies and along the coast of Florida. It is a very graceful and handsome species, very common, often leaping from the water. It is not much valued as food, being dry and bony. Forty-two, from 2.25 to 10.5 inches in length, were taken at Pato Seco, Mayaguez, Ponce, Hucars, and Fajardo; one at San Geronimo.

Scomber saurus Bloch & Schneider, Syst. Ichth., 321, 1801, Jamaica.
Centronotus argenteus Lacépède, Hist. Nat. Poiss., III, 316, 1802, Equatorial America.
Lichia quiebra Quoy & Gaimard, Voyage Freycinet, Zool., 365, 1824, Equatorial America.
Chorinemus saltans Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 393, 1831, Martinique, Brazil, and Santo Domingo.
Ologoplites inornatus Gill, Proc. Ac. Nat. Sci. Phila. 1863, 166, Panama.
Ologoplites occidentalis, Poey, Fauna Puerto-Riqueña, 332, 1881; Stahl, l. c., 77 and 163, 1883.
Ologoplites saurus, Jordan & Evermann, l. c., 898, 1896.

Genus 54. *SERIOLA* Cuvier. The Amber-fishes.

Body oblong, moderately compressed, not elevated. Occiput and breast not trenchant. Head usually more or less conical, not very blunt. Mouth comparatively large, with broad bands of villiform teeth on both jaws, tongue, vomer, and palatine; a broad, strong, supplemental maxillary bone; premaxillaries protractile. Scales small; lateral line scarcely arched, forming a keel on caudal peduncle, not armed with bony plates; sides of head with small scales. First dorsal with about 7 low spines, connected by membrane; second dorsal very long, elevated in front; anal similar to soft dorsal but not nearly so long, shorter than abdomen, preceded by 2 very small free spines, which disappear in old fishes; no finlets; ventral fins very long; pectoral short and broad. Gillrakers moderate.

Species of moderate or large size, often gracefully colored; most of them valued as food-fishes.

SERIOLA:

- a. Head longer than deep, profile not very steep. Dorsal and anal fins not falcate, height of their lobes less than half depth of body. Dorsal rays 30 to 38; species of large size, elongate, with 5 or 6 broad, dark crossbars when young, these becoming obsolete with age; a yellow lateral band; nuchal bar pale.
- b. Dorsal rays 36 to 38; dark bands on young very broad. *zonata*
- bb. Dorsal rays 30 to 34. Mouth rather large, maxillary reaching middle of pupil; dark bands on young broad.
- c. Body slender, depth 3.5 to 3.75 in length *lalandi*
- cc. Body deeper, depth 3 in length *dumeritii*

ZONICHTHYS:

- aa. Head deeper than long, anterior profile steep; no yellow longitudinal band; size small.
- d. Dorsal not falcate, its soft rays 32; young with about 12 narrow dark crossbars *fasciata*
- dd. Dorsal and anal falcate, their anterior lobes more than half depth of body; head deeper than long; body deep; dorsal rays 27 to 30.
- e. Nuchal band dark-brown or black *rivoliiana*
- cc. Nuchal band pale-yellowish *falcata*, 85

85. *Seriola falcata* Cuvier & Valenciennes. *Madregal*; *Rock Salmon*.

Head 3.8 (4.6 in total); depth 3.4 (4 in total). D. VII-1, 29; A. II-1, 21. Cæca 30. Body rather deep and compressed. Head somewhat longer than deep, not conical. Snout 2.75 in head; maxillary reaching front of pupil, 3.5 in head, its tip broad; eye large, 5.25 in head, 1.75 in snout. Occiput somewhat carinated. Interorbital space wide, convex. Caudal keel little developed. Dorsal high, somewhat falcate, its anterior lobe 1.4 in head, 2.33 in base of fin; pectoral 2 in head; ventrals 1.6; anal lobe 1.75; anal spines small.

Life coloration as follows! Grayish above, paler but hardly silvery below; fins blackish, pectoral pale; caudal not at all yellow; eye white; lining of opercle pale; a very obscure olivaceous band from eye to front of dorsal, scarcely visible in fresh specimens; preorbital and preopercle shaded with olive.

Found in the West Indies, north to Florida and the Carolinas. Not seen by us in Porto Rico, but included on the authority of Valenciennes and Poey.

Seriola falcata Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 210, pl. 515, 1833, Porto Rico, Jamaica, and Mexico; Poey, Fauna Puerto-Riqueña, 332, 1881; Jordan & Evermann, l. c., 905, 1896.

Seriola dubia Lowe, Proc. Zool. Soc. Lond., VII, 1839, 81, Madeira.

Seriola declivis Poey, Memorias, II, 230, 1860, Havana.

Seriola ligulata Poey, Memorias, II, 231, 1860, Cuba.

Genus 55. *DECAPTERUS* Bleeker. The Mackerel Scads.

Body elongate, little compressed, almost perfectly fusiform. Head short, pointed. Mouth rather small; jaws about equal, the dentition feeble; maxillary rather broad, with a supplementary bone. Premaxillaries protractile. Scales moderate, enlarged for the whole length of lateral line, but spinous and bony posteriorly only; second dorsal and anal each with a single detached finlet; free anal spines very strong; first dorsal well developed, persistent; pectoral comparatively short. Abdomen rather shorter than anal fin. Gillrakers long and slender.

Species numerous; only one known to occur in Porto Rico.

- a. Shields of lateral line numerous, 40 to 50 in number. *punctatus*, 86
- aa. Shields of lateral line few, 20 to 30 in number.
- b. Teeth minute on both jaws, vomer, palatines, and tongue; shields 22 to 28; depth 5 in length. *sanctæ-helenæ*
- bb. Teeth obsolete; caudal keel of 25 shields; depth 5.75 in length. *macarellus*

86. *Decapterus punctatus* (Agassiz). *Scad; Round Robin; Cigar-fish; Quia-quia.*

(PLATE 8.)

Head 4; depth 5.6; eye 3.3; snout 3; maxillary 3; mandible 2.3; interorbital 3.5; D. VIII-1, 30-1; A. II-1, 27-1; pectoral 1.4; ventral 2.1; caudal 1.7; scutes 39.

Body elongate, fusiform, scarcely compressed at all; caudal peduncle very short and slender; head small and pointed; jaws subequal, maxillary reaching front of eye; teeth of jaws weak, uniserial; adipose eyelid as in *T. crumenophthalmus*, but not covering so much of eye; lateral line scarcely arched, posterior portion with large rather sharply armed scutes, anterior with a series of ordinary scales which are not as easily deciduous as others. Shoulder-girdle with a small papilla of the integument (seen by raising gill-covers), somewhat similar to that in *T. crumenophthalmus*, but much smaller and higher in position. This species resembles the latter, but is readily recognized by the presence of the single detached finlet at end of soft dorsal and anal fins.

Color, bluish above, silvery below, with a dark opercular spot, its edge extending upon shoulder.

The scad ranges from Cape Cod to Brazil, but is only occasional northward. It is common on the coasts of Florida and in the West Indies. Nine examples collected, three of them each about 6½ inches in length, taken in the seine by native fishermen at Aguadilla, where, with *T. crumenophthalmus*, it was the principal part of the catch; the other six are the young, about 2 inches in length, taken in 220 fathoms, 9 miles from Mayaguez, by the beam-trawl. The species is commonly used as food.

Caranx punctatus Agassiz, Spix, Pisc. Bras., 108, 1829, Brazil.

Caranx suareus Risso in Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 33, 1833, Mediterranean.

Decapterus punctatus, Jordan & Evermann, l. c., 907, 1896.




FIG. 30.—*Trachurops crumenophthalmus*.

Genus 56. TRACHUROPS Gill.

This genus is close to *Caranx*, differing in the more elongate form and especially in the structure of the shoulder-girdle, which has a deep cross-furrow at its junction with the isthmus, with a fleshy projection above the furrow. Species few, found in all warm seas.

87. *Trachurops crumenophthalmus* (Bloch). *Goggler; Big-eyed Scad; Goggle-eye Jack; Chicharro.*

Head 3.2; depth 3.8; eye 3.2; snout 2.9; maxillary 2.5; mandible 2; interorbital 4.4; D. VIII-1, 25 or 26; A. II-1, 22; pectoral 1.2; ventral 2.1; scutes 35.

Body elongate; but little compressed; back not elevated; dorsal outline little curved; caudal peduncle subcylindrical, very slender and short; head rather large and pointed; mouth large, lower jaw projecting; teeth on vomer, palatines, tongue, and jaws all weak, uniserial on jaws; maxillary reaching about to front of pupil; eye very large, with an adipose eyelid covering two-thirds of eye, leaving a vertical elliptical area exposed opposite pupil; lateral line slightly arched anteriorly, covered posteriorly with large weak scutes, their spines not strong nor sharp; pectoral falcate; ventral small; soft dorsal and anal long and low, their anterior rays somewhat elevated; caudal widely forked. Just above junction of isthmus with shoulder-girdle is a large fleshy nipple-like projection.

F. C. B. 1900-9

Color, bluish above, silvery below; a faint dark opercular blotch; no striking colors.

This species reaches a length of 2 feet, and is found on both coasts of tropical America and in tropical seas generally. Common about Porto Rico; 38 examples, from 4 to 9 inches, collected at Palo Seco, Aguadilla, and Culebra; one from San Geronimo.

Scomber crumenophthalmus Bloch, Ichthyologia, pl. 343, 1793, Acara in Guinea.

Scomber plumieri Bloch, Ichthyologia, pl. 344, 1793, Antilles.

Scomber balantiophthalmus Bloch & Schneider, Syst. Ichth., 29, 1801, Guinea.

Caranx macropthalmus Agassiz in Spix, Pisc. Bras., 107, 1829, Brazil.

Trachurops brachychirus Gill, Proc. Ac. Nat. Sci. Phila. 1862, 261, Cape San Lucas.

Trachurops crumenophthalmus, Poey, Fauna Puerto-Riqueña, 331, 1881; Jordan & Evermann, l. c., 911, 1896.

Trachurops plumieri, Stahl, l. c., 77 and 163, 1888.

Genus 57. CARANX Lacépède. The Crevallés.

Body ovate or oblong, compressed; back sometimes considerably elevated, sometimes little arched. Head moderate or rather large, more or less compressed. Mouth moderate or large, oblique; maxillary broad, with a well-developed supplemental bone, extending to below eye. Premaxillaries protractile. Teeth developed in one or few series, unequal, or at least not in villiform bands. Villiform teeth usually present on vomer, palatines, and tongue, wanting or deciduous in some species. Gillrakers long. Eye large, with adipose eyelid. Dorsal spines rather low, connected; second dorsal long, usually elevated in front, both fins depressible in a groove. Anal fin similar to second dorsal and nearly as long, preceded by two rather strong spines, its base longer than abdomen. Caudal fin strongly forked, peduncle very slender. Ventral fins moderate; pectorals falcate; no finlets. Scales present, mostly very small. Lateral line with its posterior portion armed with strong bony plates, which grow larger on tail, each plate armed with a spine; a short dorsal branch of lateral line usually present. Preopercle entire in adult, serrate in young, usually with a membranous border.

Species very numerous, in all warm seas, most of them valued for food.

- a. Teeth on vomer and palatines persistent.
- b. Soft dorsal and anal low, not much elevated in front, little if at all falcate; teeth in jaws in one or few series, with no canines.
- c. Body slender, depth about 3.5 in length; color dark, chiefly bluish..... *ruber*, 88
- cc. Body deeper, depth about 2.8 in length; color pale, mostly golden..... *bartholomæi*, 89
- bb. Soft dorsal and anal much elevated in front and more or less falcate; upper teeth in a band, outer enlarged; lower teeth in one series; dorsal sheath of scales not greatly developed; soft dorsal and anal both rather short.
- d. Breast naked, except a small rhombic area before ventrals; two small canines in front of lower jaw; body robust, compressed.
- e. Opercular spot large; adult with a black spot on pectoral; pectoral 3 in length..... *hippos*, 90
- dd. Breast entirely covered with small scales.
- f. Body subfusiform, depth less than one-third the length; breast scaly; teeth of outer series small, not canine-like; a black opercular spot; no spot on pectoral; arch of lateral line about half straight part..... *crysos*, 91
- ff. Body oblong-ovate, depth more than one-third length; outer teeth rather strong, lower teeth not canine-like.
- g. General color silvery; vertical fins not all black..... *latus*, 92
- gg. General color brassy or blackish; vertical fins black; lower teeth not canine-like *lugubris*

URASPIS:

- aa. Teeth on vomer and palatines wanting or deciduous; teeth in jaw subequal, bluntish, in one or two rows; lateral line not strongly arched; soft dorsal and anal low; shields rather few.
- h. Body deep, compressed, back and belly arched; shields 24 to 30; second dorsal 1, 26; anal 1, 22; opercular spot present..... *guara*

88. *Caranx ruber* (Bloch). *Cibi Mancho*; *Carbonero*.

Head 3.5; depth 3.5; eye 5.4; snout 2.6; maxillary 2.6; mandible 2.25; interorbital 3; preorbital 9; D. VIII-1, 26; A. II-1, 23; pectoral 1; ventral 2.8; caudal 1.1; scutes about 30.

Body elongate, back scarcely elevated, caudal peduncle slender, much wider than deep, scutes forming a very strong keel; head pointed, symmetrical; maxillary nearly or quite reaching front of eye; teeth on vomer, palatines, tongue, and jaws, in villiform bands; no enlarged teeth; arch of lateral line moderate, somewhat shorter than straight part, which is about 2.5 in body; pectoral falcate, with very slender tip, reaching anal; anterior rays of soft dorsal and anal somewhat produced, those of dorsal 2.3 in head, of anal 2.7; caudal widely forked.

Color in spirits: Bluish above, pale on sides and below; parts of head with golden wash; lower lobe of caudal with a wide dusky lengthwise bar; dorsal somewhat dusky, other fins nearly pale.

West Indies. Four specimens, 6.5 to 14 inches in length, from Mayaguez, Ponce, and Culebra, taken in the seine. One of 5 inches from San Geronimo we refer with some doubt to this species.

Scomber ruber Bloch, Ichthyologia, pl. 342, 1793, St. Croix.

Caranx blochii Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 69, 1833, St. Croix.

Caranx iridimus Poey, Memorias, II, 226, 1860, Cuba.

Caranx ruber, Jordan & Evermann, l. c., 919, 1896.

89. *Caranx bartholomæi* (Cuvier & Valenciennes). *Yellow Jack; Cibi Amarillo.*

Head 3.3; depth 2.8; eye 4.8; snout 2.8; maxillary 2.4; mandible 2.1; interorbital 3; preorbital 8; D. viii-1, 26; A. ii-1, 22; pectoral 1; ventral 2.6; caudal 1.1; about 30 scutes. Close to *C. ruber*, but the body not so slender, and straight portion of lateral line longer, nearly as long as curved part, 2.8 in body; maxillary reaching past front of eye.

Bluish above, a golden wash everywhere.

West Indies, north to Florida and the Carolinas; common in Cuba, but not very common in Porto Rico. One specimen, 9.5 inches long, from San Juan market.

Caranx bartholomæi Cuv. & Val., Hist. Nat. Poiss., IX, 100, 1833, St. Bartholomew; Jordan & Evermann, l. c., 919, 1896.

Caranx cibi Poey, Memorias, II, 224, 1860, Cuba.

Caranx beani Jordan, Proc. U. S. N. M. 1880, 486, Beaufort, N. C.

90. *Caranx hippos* (Linnaeus). *Crevallé; Toro; Horse Crevallé; Cavally; Jack; Jiguayua.*

Head 3; depth 2.5; eye 3.8; snout 3.4; maxillary 2.3; mandible 2; interorbital 3.7; D. viii-1, 20; A. ii-1, 17; pectoral 1; ventral 2.3; caudal 1.1; scutes about 30.




FIG. 31.—*Caranx hippos*.

Body not greatly elongate, anterior profile very strongly arched, approximating vertical from front of eye to tip of snout in adult; mouth nearly horizontal, low; maxillary reaching posterior border of eye in adult, middle of eye in 5-inch individuals; teeth of upper jaw in a villiform band, with an outer row of considerably enlarged teeth, rather wide-set; one row of teeth in lower jaw, a pair of canines at symphysis; lateral line with a rather strong arch, shorter than straight part; breast naked, except a patch in front of ventrals. A large, very distinct jet-black spot on opercle; a fainter black spot on lower rays of pectoral, often wanting in young; axil of pectoral black; lobe of soft dorsal and upper surface of caudal peduncle dusky.

The most strongly marked of the species of *Caranx*, known at once from the strong arch of the head, the naked breast, and the opercular and pectoral spots. The collection contains only the young, 2.75 to 5.5 inches long, from Palo Seco. It does not appear to be common in Porto Rico, though it is of wide distribution, being found in all warm seas and generally abundant. It occurs on both coasts of tropical America and ranges as far north as Cape Cod and the Gulf of California. It bites voraciously and affords much sport to the angler, who takes it by trolling. It is taken at Key West both by trolling and by bottom fishing. It is very ravenous, swimming with great swiftness, and preying upon smaller fishes. It is said to attain a weight of 20 pounds, but those caught in Indian River, Florida, do not average over 3 pounds. As a food-fish it does not occupy a high rank. It was not seen in any of the Porto Rican markets, which, however, was probably due to its scarcity at that particular season.

- Scomber hippos* Linnæus, Syst. Nat., ed. XII, 494, 1766, Charleston, S. C.
Scomber carangus Bloch, Ichth., pl. 340, 1793, Antilles.
Caranx erythrus Laépède, Hist. Nat. Poiss., III, 68, 1802, South Carolina.
Caranx carangua Laépède, Hist. Nat. Poiss., III, 59 and 74, 1802, Martinique; on a drawing by Plumier.
Caranx daubentonii Laépède, Hist. Nat. Poiss., III, 72, 1802, Martinique; on a drawing by Plumier.
Caranx xanthopigus Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 109, 1833, Isle de France.
Caranx ekala Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 117, 1833, Vizagapatam.
Caranx antillarum Bennett, Whaling Voyage, II, 282, 1840, West Indies.
Caranx defensor De Kay, New York Fauna; Fishes, 120, 1842, New York.
Carangus csculentus Girard, U. S. Mex. Bound. Surv., 23, pl. 11, figs. 1-3, 1859, Brazos Santiago, Texas.
Caranx caninus Günther, Fishes Centr. Amer., 432, 1869, Panama.
Caranx hippos, Jordan & Evermann, l. e., 920, 1896.

91. *Caranx crysos* (Mitchill). "*Cojinuda*"; *Hard-tail*; *Runner*; *Jurel*; *Yellow Mackerel*; *Crevalle*.

(PLATE 9.)

Head 3.5; depth 3.2; eye 5.6; snout 3; maxillary 2.4; mandible 2.1; interorbital 3; preorbital 11; D. VIII-1, 24; A. II-1, 20; pectoral 0.8; ventral 2.3; caudal 0.95; about 45 scutes.

Body elongate, slightly elevated, not greatly compressed; maxillary reaching front of pupil or middle of eye; teeth in jaws conical and sharp, not very close-set, in one series in lower jaw, in two in upper, inner series of smaller teeth; no canines; lateral line more strongly arched anteriorly than




FIG. 32.—*Caranx crysos*.

in *C. ruber* or *C. bartholomæi*, arch rather more than half straight part; pectoral long and falcate, reaching anal or slightly beyond; anterior rays of soft dorsal and anal elevated and falcate. A distinct black blotch on opercle.

This species is found from Cape Cod to Brazil, is generally abundant, and is common farther to the north than any other of the species of *Caranx*. It is an important and well-known food-fish, fairly common. In Porto Rico only two specimens were collected, one 10.5 inches long from San Antonio Bridge, and one 17.5 inches long, which was taken on the hook at Isabel Segunda, displaying good game qualities. This is one of the best game-fishes of Porto Rico and can be taken by either still-fishing or trolling.

- Scomber crysos* Mitchill, Trans. Lit. & Phil. Soc. N. Y., I, 1815, 424, New York.
Caranx pisquetus Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 97, 1833, Santo Domingo, Cuba, and Brazil.
Trachurus squamosus Gronow, Cat. Fishes, ed. Gray, 125, 1854, Carolina.
Paratractus chrysos, Stahl, l. e., 77 and 163, 1883.
Caranx crysos, Jordan & Evermann, l. e., 921, 1896.

92. *Caranx latus* Agassiz. "*Jurel*"; *Xurel*; *Horse-eye Jack*.

Head 3.35; depth 2.7; eye 4; snout 3.5; maxillary 2.1; mandible 1.8; interorbital 3.2; D. VIII-1, 22; A. II-1, 18; pectoral 0.95; ventral 2.1; caudal 1.1; about 35 scutes. Form of *C. hippos*, but with anterior profile not quite so strongly arched, breast scaled; axillary and pectoral spots absent; opercular spot not so large nor distinct, scutes somewhat greater in number; adipose eyelid moderately developed; maxillary reaching past middle of eye.

Lobe of dorsal dusky, that of anal golden; the young with 4 or 5 dark vertical bars and a faint opercular spot, these sometimes absent; opercular spot is larger, deeper in color, and more persistent in young of *C. hippos*, and usually serves to separate readily small individuals of the two species.

C. latus is a very widely distributed species, occurring in all tropical seas, and very abundant in the West Indies, and by far the commonest species of *Caranx* in Porto Rico. The collection contains 137 specimens, varying in size from 3 to 9.5 inches. The poisonous character sometimes attributed to its flesh evidently does not attach to it in Porto Rico, where it may be found for sale in the markets. Our specimens were collected at Palo Seco, San Antonio Bridge, Aguadilla, Mayaguez, Culebra, Isabel Segunda, and Fajardo. Mr. Gray took two at San Geronimo.

Caranx latus Agassiz, Misc. Bras., 105, 1829, Brazil; Jordan & Evermann, l. c., 923, 1896.

Caranx lepturus Agassiz, Misc. Bras., 106, 1829, Brazil.

Scomber heberi Bennett, Fishes Ceylon, pl. 26, 1830, Ceylon.

Caranx fallax Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 95, 1833, Antilles: Brazil.

Caranx sen Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 105, 1833, Pondicherry.

Caranx forsteri Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 107, 1833, East Indies.

Caranx peroni Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 112, 1833, East Indies.

Caranx lessoni Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 113, 1833, Malabar.

Caranx belengeri Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 116, 1833, Malabar.

Caranx parasistes Richardson, Voyage Erebus and Terror, 136, 1844, Port Essington.

Caranx richardi Holbrook, Ichth. South Carolina, 96, pl. 13, fig. 1, 1860, South Carolina.

Caranx aureus Poey, Enumeratio, 76, 1875, Cuba.

Carangus fallax, Poey, Fauna Puerto-Riqueña, 331, 1881; Stahl, l. c., 77 and 163, 1883.




FIG. 33.—*Caranx latus*.

Genus 58. VOMER Cuvier & Valenciennes.

This genus is closely allied to *Caranx*, from which it differs only in its distortion of form, and in its weak teeth and very low fins. Body broad-ovate, very strongly compressed, all the outlines sharply trenchant. Head very gibbous above eyes, its anterior profile vertical; lateral line strongly arched, its posterior portion with very weak shields. Scales minute, rudimentary; the soft dorsal and anal extremely low, not falcate. Young much deeper in form than adult, all the fins higher, resembling *Selene*. Found in the warm seas.

a. Soft dorsal with about 25 rays; depth in adult less than half length..... *dorsalis*

aa. Soft dorsal with 21 or 22 rays.

b. Depth in adult about half length..... *setipinnis*

bb. Depth in adult much more than half length..... *gabonensis*, 93

93. *Vomer gabonensis* Guichenot. "*Corcobado*."

Head 2.75; depth 1.5; eye 3.8; snout 2.2; maxillary 2.5; mandible 2.2; interorbital 4.2; preorbital 2.6; D. VIII-1, 22; A. 1, 18; pectoral 1; caudal 1.1; scales minute.

Body ovate, scarcely longer than deep, very greatly compressed; occipital region greatly elevated, making height of body greatest above eye; anterior profile nearly vertical from occiput to eye,

opposite which it becomes concave; snout protruding; mouth oblique, maxillary extending opposite nostrils, not reaching front of eye; lower jaw entering profile; a single row of numerous weak teeth in each jaw; preorbital very broad; spinous dorsal weak, first four spines comparatively long, second about half diameter of eye, very slender, weak, and flexible, next four much shorter, stiff, pungent, and partially embedded; soft dorsal long and low, anterior rays somewhat elevated, not falcate; anal similar to soft dorsal, its rays somewhat more widely set, first one or two a little elevated; pectoral falcate, as long as head, reaching past middle of anal; ventrals minute, caudal peduncle short and very slender, the fin widely forked, its lobes equal; head and upper parts of body naked, minute persistent scales below; lateral line with minute rudimentary scutes on its straight portion, anterior part strongly arched, its chord a little shorter than straight part.




FIG. 34.—*Scene vomer*. Young.

Color in life: Rich silvery with an iridescence of steel-blue above and of pink below, with a light golden wash on lower parts.

This species seems to be the common *Vomer* of Porto Rico and distinct from *V. setipinnis*, which does not appear in our collection. It may be separated from the latter form by the much deeper body, larger eye, greater elevation of occipital region, and more nearly vertical anterior profile. The young of the two species resemble each other more closely. At a length of about 4 inches the difference between them in relation of depth to length becomes apparent, but before that size is attained *V. gabon-*

ensis may be distinguished by its larger eye, at least in some individuals of 2 or 3 inches in length. The ventral outline of *V. setipinnis* is somewhat more strongly curved than in *V. gabonensis*, while in the adult the reverse is true. The young of *V. gabonensis* has the caudal lobes more slender and somewhat longer than in the corresponding size of the other species, but this difference disappears in the adult. The species reaches a weight of a pound or more, is abundant and much used as food, being hawked about the streets upon strings attached to a stick, which is carried over the shoulders.

The corcobado is known from Africa and the West Indies. We collected 32 examples, 2.5 to 8 inches long, from San Juan market, Palo Seco, Arecibo, Aguadilla, Arroyo, and Isabel Segunda.

Argyrosus setipinnis var., Günther, Cat., II, 459, 1860, Fernando Po, Santo Domingo, Jamaica, and Bahia.

Vomer gabonensis Guichenot, Ann. Soc. Linn. Maine et Loire, 42, 1865, Gaboon; Jordan & Evermann, I. c. 934, 1896.

Genus 59. **SELENE** Lacépède. The Moon-fishes.

Body very closely compressed and much elevated; profile very oblique or nearly vertical; edges of body everywhere trenchant, especially anteriorly. Head short and very deep, opercle very short, and preorbital extremely deep; an abrupt angle at occipital region. Mouth rather small; premaxillaries protractile, fitting into a notch between the bases of maxillaries; maxillaries broad, each with supplemental bone. Tongue narrow, free. Teeth minute, on jaws, tongue, vomer, and palatines. Gillrakers long and slender. Spines of fins usually weak, more or less filamentous in young; free anal spines immovable, sometimes obsolete in the adult. Soft fins falcate, much elevated. No finlets. Head naked. Scales minute. Lateral line wholly unarmed. Coloration silvery.

Found in the tropical seas. Notwithstanding its extraordinary form this genus differs in no important regard from *Caranx*.




FIG. 35.—*Selene vomer*. Adult.

94. **Selene vomer** (Linnaeus). Moon-fish; Jorobado; Look-down.

Head 3; long dorsal rays 2; pectoral 2.75; long anal rays 2.66; depth 1.5, the young much deeper. D. VII-1, 23; A. II-1, 18. Anterior profile from tip of snout to occiput almost perfectly straight in adult. Diameter of eye, length of opercle, and distance from eye to profile about equal; eye 2 in maxillary, 2.5 in preorbital; mandibles very deep, dentary bones thin, approximate; one or two of the dorsal spines greatly elongate and filamentous in young, short in adult; ventrals variable in length, usually about as long as the eye in the adult, variously elongate in partly grown specimens.

Color, uniform silvery in adult.

Our observations of this species tend to confirm the correctness of Dr. Lütken's views (*Spolia Atlantica*, 139) as to the transformations incident to its growth.

A species of tropical America, occurring on both coasts, from Cape Cod to Brazil, and from Lower California to Peru. Very common southward on sandy shores, both in the Atlantic and Pacific. No specimens were obtained by us, but we include it on the authority of Poey and Stahl. It attains a weight of about 2 pounds and is a delicious pan-fish.

- Zeus vomer* Linn., Syst. Nat., ed. X, 1758, 266, America; after *Zeus cauda bifurca* of Linnæus, Mus. Adolph-Fred., I, 66.
Zeus gallus Linn., Syst. Nat., ed. X, 1758, 267, America; after *Zeus cauda bifurca* of Artedi; confused with *Alectis ciliaris*.
Zeus capillaris Mitchill, Trans. Lit. & Phil. Soc., I, 1815, 383 (young), New York.
Zeus rostratus Mitchill, l. c., 384, New York (young).
Zeus geometricus Mitchill, Amer. Monthly Mag., II, 1818, 245 (adult), New York.
Argyrosus triacanthus Swainson, Nat. Hist. Class'n Fish., 250, 1839, Brazil; after Spix & Agassiz, pl. 58 (young).
Argyrosus mauricei Swainson, l. c., 408, Brazil (adult).
Argyrosus mitchilli De Kay, N. Y. Fauna: Fishes, 126, 1842 (young), New York.
Argyrosus brevoortii Gill, Proc. Ac. Nat. Sci. Phila. 1863, 83, Panama (young).
Argyrosus pacificus Lockington, Proc. Ac. Nat. Sci. Phila. 1876, 84, Lower California; adult.
Selene argentea, Poey, Fauna Puerto-Riqueña, 332, 1881; Stahl, l. c., 77 and 164, 1883.
Selene vomer, Jordan & Evermann, l. c., 936, 1896.




FIG. 36.—*Chloroscomber chrysurus*.

Genus 60. **CHLOROSCOMBERUS** Girard. The Casabes.

Body oblong-ovate, closely compressed, but not elevated, the abdomen prominent anteriorly, its curve much greater than curve of back. Occiput and thoracic region trenchant. Caudal peduncle very narrow, fin widely forked. Scales small, smooth. Lateral line arched in front, unarmed, or with a few small plates. Head nearly naked. Preorbital low. Mouth rather small, oblique, lower jaw scarcely projecting; upper jaw protractile; maxillary broad, emarginate behind, with large supplemental bone. Jaws, vomer, and palatines with feeble teeth, mostly in a single series. First dorsal of feeble spines, connected by membrane; second dorsal and anal long and low, similar, much longer than the short abdomen. No finlets. Anal spines strong. Ventrals small; pectoral falcate. Gillrakers long.

An American genus, of small size, and little valued as food.

95. **Chloroscomber chrysurus** (Linnæus). *Casabe*; *Bumper*.

Head 3.75; depth 2.33; eye very large, longer than snout, about 3 in head; D. VIIII-1, 26; A. II-1, 26. Head rather deeper than long; opercles very short; snout short. Mouth very oblique; maxillary reaching anterior margin of eye. Chord of curved part of lateral line scarcely longer than head, 1.66 to 1.75 times in length of straight part. Lateral line wholly unarmed. Caudal peduncle longer than deep, its diameter less than that of eye; ventrals very small, fitting into a groove in which the vent is situated; pectoral long, falcate, one-third the length.

Greenish above, sides and below golden; a jet-black spot on caudal peduncle above; dark opercular and axillary spots; inside of mouth black; fins not bordered nor tipped with black.

This species ranges from Cape Cod to Brazil; it is very common on our south Atlantic coast and in Cuba, but is not valued as food, the flesh being thin and dry, the bones large. It is apparently not common in Porto Rico, specimens being obtained by us only at Mayaguez and Isabel Segunda.

Scomber chrysurus Linnaeus, Syst. Nat., XII, 494, 1766, Charleston, South Carolina.

Scomber chloris Bloch, Ichthyologia, pl. 339, 1793.

Micropteryx cosmopolita Agassiz, Spix, Pisc. Bras., 104, 1829, Brazil.

Scomber latus Gronow, Catalogue Fishes, ed. Gray, 127, 1854, Carolina.

Chloroscombrus caribbeus Girard, Mex. Bound. Surv., Zool., 21, pl. 9, fig. 6, 1859, St. Joseph Island, Texas.

Chloroscombrus chrysurus, Poey, Fauna Puerto-Riqueña, 332, 1881; Stahl, l. c., 77 and 163; 1883; Jordan & Evermann, l. c., 938, 1896.

Genus 61. TRACHINOTUS Lacépède. The Pámpanos or Pompanos.

Body compressed, moderately elevated, general outline ovate. Caudal peduncle short and rather slender. Abdomen not trenchant, shorter than anal fin. Head moderately compressed, very blunt, snout abruptly truncate. Mouth nearly horizontal, the maxillary reaching middle of eye; premaxillaries protractile; maxillary without distinct supplemental bone. Jaws, vomer, and palatines with bands of villiform teeth, which are deciduous with age. Preopercle entire in adult. Gillrakers short. Gill-membranes considerably united. Spinous dorsal represented by six rather low spines, which are connected by membrane in young, but are free in adult. In old examples the spines appear small on account of encroachments of flesh, and ultimately often disappear. Second dorsal long, elevated in front; anal opposite to it and similar in form and size; two stout, nearly free spines in front of anal, and one connected with the fin, these often disappearing with age. Scales small, smooth. Lateral line unarmed, little arched; no caudal keel.

"When extremely young the preoperculum is armed at the angle with three large spines, and smaller ones above and below. The spinous dorsal is developed as a perfect fin, and teeth are present on the jaws and palatine arch. In this stage the species has never been described by previous naturalists, and consequently has received no name, as the corresponding stage of *Nauclerates* (*Nauclerus*) has. At an early period the preopercular spines are absorbed into the substance of the preoperculum and disappear. The spinous dorsal and the teeth are still retained. In this condition it remains for some time, the spinous dorsal, however, gradually losing its relative size, while the soft vertical fins increase. In this stage the species belongs to the nominal genus *Doliodon* of Girard. At a later period the membrane connecting the dorsal spines has become obsolete, and the species then represents the genus *Trachinotus*, as understood by Cuvier & Valenciennes and others. Finally, in old age, the teeth of the jaws, palate, and pharyngeal bones have fallen out, and the lobes of the dorsal, anal, and caudal fins attain their greatest extension and become pointed. This final stage has been made known by Holbrook under the new generic name of *Bothrolæmus*" (Gill, Proc. Ac. Nat. Sci. Phila. 1862, 440). The pseudobranchiæ also disappear in old individuals.

Some species of *Trachinotus* (*carolinus*, etc.) are among the most highly valued of our food-fishes.

- a. Dorsal with 19 or 20 soft rays; anal with 17 to 19 soft rays.
- b. Body very much compressed; sides with narrow black crossbars; lobes of vertical fins elongate, reaching past middle of caudal fin in adult *glaucus*, 96
- bb. Body moderately compressed; sides without narrow black crossbars; lobes of vertical fins shorter, rarely reaching base of caudal; lobes of dorsal and anal usually blackish.
- c. Body broad-ovate; back arched; greatest depth at all ages nearly two-thirds length of body; profile from nostril to dorsal everywhere nearly evenly convex; axil without black spot *falcatus*, 97
- cc. Body oblong, the profile not strongly arched; depth in young and old 2 to 2.6 in length of body; dorsal lobes low *goodii*
- aaa. Dorsal with 25 to 27 soft rays; anal with 22 to 26 soft rays; body oblong, rather robust; greatest thickness 3 in greatest depth of body; depth less than half length; lobes of vertical fins short, not black; sides without dark crossbars.
- d. Dorsal with 25 soft rays; anal with 22 soft rays; profile from snout to procumbent spine evenly convex.
- e. Body very deep, depth about half length *argenteus*
- cc. Body moderately deep, depth about 2.5 in length *carolinus*, 98
- ddd. Dorsal with 27 soft rays; anal with 26 *cayennensis*

96. *Trachinotus glaucus* (Bloch). *Gaff-topsail*; *Pompano*; *Palometa*.

Head 4; depth 2; eye 3.6. D. vi-1, 19; A. ii-1, 18; pyloric cæca 13. Body elliptical, much compressed; snout blunt, subtruncate, vertical from mouth to horizontal from upper edge of the eye; profile from supraorbital to front of dorsal fin convex. Mouth nearly horizontal; maxillary nearly

reaching vertical from middle of eye, its length 3 in head; jaws without teeth in adult; dorsal spines separate in adult; dorsal and anal fins falcate, anterior soft rays reaching middle of caudal fin; dorsal lobe 1.5, anal 1.75 in length of body; ventrals reaching four-fifths distance to vent, their length 2.4 in head; caudal very deeply forked, its lobes nearly half length of body.

Color, bluish above, golden below; lobes of dorsal and anal very dark; rest of fins pale, with bluish edges; caudal bluish; pectorals golden and bluish; ventrals whitish. Body crossed by four black vertical bands, first under procumbent spine, second under third dorsal spine, third and fourth under soft dorsal; a black spot representing a fifth band on lateral line between last rays of dorsal and anal, this sometimes obsolete; the position of the bands appearing to be subject to slight variation.

Young individuals, 2 inches in total length, may be described as follows: Head 2.9; depth 2.6; eye 4; dorsal profile and general form as in adult; anterior dorsal and anal rays not produced; caudal less deeply forked. Color, bright-silvery, merging into metallic-bluish on back and yellowish on lower sides and belly; the four dark vertical bars very faint, but usually distinguishable with a lens, even in very small specimens; the first of these very short and under procumbent spine, second longer and under fourth spine, third and fourth under soft dorsal; the dark spot on posterior part of lateral line usually not evident; anterior rays of dorsal and anal and outer rays of caudal black.

Comparing young examples in our collection from Porto Rico with specimens of similar size from Woods Hole which have been identified as the young of *T. goodii*, it is found that they are difficult to distinguish. The general form, proportional measurements, fins, and general coloration are very much alike, but the black vertical bars are apparently not present on the Woods Hole specimens.




FIG. 37.—*Trachinotus glaucus*.

This species attains a length of a foot or more and is a handsome fish, though not highly esteemed as food. In Porto Rico, however, it is handled by the fishermen and seems to rank with the species of *Caranx* in food value. It is found in tropical America from Virginia to the Caribbean Sea; generally common from the Carolinas to Florida and in Porto Rico. Our collection contains examples from Mayaguez, Aguadilla, Ponce, and Isabel Segunda. Mr. Gray's collection has two from San Geronimo.

Chaetodon glaucus Bloch, Ichthyologia, pl. 210, 1787, Martinique; on a drawing by Plumier.

Trachynotus glaucus, Poey, Fauna Puerto-Riqueña, 333, 1881; Stahl, l. c., 77 and 164, 1883.

Trachinotus glaucus, Jordan & Evermann, l. c., 940, 1896.

97. *Trachinotus falcatus* (Linnaeus). *Round Pompano; Palometa; Permit.*

Head 3.6; depth 1.6; eye 3.9; snout 3.6; maxillary 3; mandible 2.4; interorbital 2.25; D. vi-r, 19; A. ii-r, 17; pectoral 1.4; ventral 2.7; caudal 0.8; scales very numerous.

Body ovate, compressed, back greatly elevated; profile of snout nearly vertical; mouth small, maxillary reaching nearly or quite to middle of eye; teeth deciduous, in villiform bands in younger individuals; anterior ray of dorsal and anal much produced, that of dorsal reaching beyond posterior end of fin, 2.5 in body; that of anal shorter, 3.3 in body; ventrals very small, reaching vent; caudal widely forked, the lobes long.

Color in life: Upper parts bluish-silvery, lower parts silvery; produced dorsal rays black outwardly, anal spines and produced rays reddish-orange, rays blackish at tips; inner edge of caudal lobes lemon, outer margin blackish; ventrals reddish.

T. falcatus is known at once from the other species of the genus by the form of the body and the elevation of dorsal and anal rays, which are longer than in any other of our Atlantic species except *T. glaucus*. Its range is from Cape Cod to Porto Rico and Brazil, common southward, apparently only the young occurring in the Gulf Stream as far north as Woods Hole. It does not seem to be at all common in the Indian River, Florida, though it occurs in some numbers, especially in the southern part of that region. It is not well known to the fishermen, only a few of whom had any name for it. These called it "permit," probably confusing it with the much larger species, *T. goodii*. It is regarded as a fair food-fish. The majority of the specimens seen in the Indian River weighed less than a pound each, and the largest about 3 pounds. In Porto Rico it is held in considerable esteem, ranking perhaps with most other species of the family. It appears to be abundant, as specimens were obtained by us at San Juan market, and at San Antonio Bridge, Palo Seco, Mayaguez, Puerto Real, Boqueron, Ponce, Arroyo, Hucares, and Isabel Segunda. It appeared to be common in most of the markets. Our Porto Rican specimens range from 1.4 to 7 inches in length.

Labrus falcatus Linnaeus, Syst. Nat., ed. X, 284, 1758, America.

Chaetodon rhomboides Bloch, Ichthyologia, pl. 209, 1787, Martinique; on a drawing by Plumier.

Trachinotus fuscus Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 410, 1831, Brazil.

Trachinotus spinosus De Kay, N. Y. Fauna: Fishes, 117, pl. 19, fig. 53, 1842, New York Harbor.

Trachinotus ovatus, Poey, Fauna Puerto-Riqueña, 333, 1881; Stahl, l. c., 77 and 164, 1883.

Trachinotus falcatus, Jordan & Evermann, l. c., 941, 1896.




FIG. 38.—*Trachinotus falcatus*.

98. *Trachinotus carolinus* (Linnaeus). Common Pompano.

(PLATE 10.)

Head 3.5; depth 2.33; eye 4.1; snout 3.7; maxillary 2.8; mandible 2.4; interorbital 2.7; D. VI-1, 24; A. II-1, 22; pectoral 1.4; ventral 2.2; caudal 9; scales very numerous. Body oblong, compressed, moderately elevated; caudal peduncle short, slender, and much compressed; mouth small, horizontal, jaws equal, maxillary reaching the middle of eye, the teeth in jaws in villiform bands in the young, finally deciduous; a procumbent spine before dorsal; spinous dorsal short and low, spines connected by membrane in young, becoming embedded in the flesh with age, membrane disappearing; anterior rays of soft dorsal and anal elevated and falcate, 1.4 in head in young of 5 inches.

Color, bluish above, silvery below with golden tinge; pectoral and anal light-orange, shaded with bluish; anterior lobe of soft dorsal dusky in young.

Found on the South Atlantic and Gulf coasts of United States, ranging north on sandy shores as far as Cape Cod; said to be rare or accidental in West Indies and Brazil, though we found the young, from 2.5 to 5 inches long, in considerable numbers at Palo Seco, Mayaguez, Ponce, and Isabel Segunda.

The pompano is perhaps the most delicious of all food-fishes. The richness, firmness, and delicacy

of flavor of its flesh render it superior not only to all other members of the family to which it belongs, but to any other fish. On the east and west coasts of Florida and about Key West it is held in the highest esteem and is most assiduously sought by the commercial fishermen, to whom it brings the highest price. It reaches a maximum length of nearly 2 feet and a weight of 6 or 8 pounds. The average weight of those taken in Indian River, Florida, probably does not now exceed 2 pounds, though before commercial fishing on that coast was so vigorously prosecuted the average weight was much greater; one was seen 21.5 inches long, weighing 4.5 pounds. At Key West the average weight is said to be about 1.5 pounds and the maximum 5 pounds.

The habits of this important fish have never been carefully studied, and its life-history therefore is not well known. It occurs in the Indian River throughout the year, but is said to be most abundant in the winter months. The best fishing is from January to early April. It is probably common in the summer months, but is not fished for then. It is most abundant about the inlets, playing in and out with the tides. It runs in bunches or schools and is easily influenced by changes in temperature, seeming to prefer rather warm water, and continued cold weather causes them to leave the river temporarily. At Key West the pompano is found only in the winter when the cold weather drives it south. It is fished for there with hook and line, while in Indian River nets are used.

The food of this fish seems to consist very largely of small bivalve shells and small crustaceans. They feed extensively about the inlets, in the surf outside, and are often seen feeding near shore. They are said to have the habit of digging in the mud or sand, which is probably done in their search for food. Not much definite information could be obtained concerning the abundance or the habits of the pompano in Porto Rico. The species was not found in any of the markets of the island nor were any large individuals seen. Our examples are smaller than any we have seen from Florida, but are about the size of those usually observed at Woods Hole in August and September, and from which they do not seem to differ.

The Porto Rican fishermen seem to recognize the species as an important food-fish, but not as being at any time abundant. The shore of the island is, in most places, unfavorable to this fish and it is probably not found in considerable numbers except on the more sandy, protected portions.

Gasterosteus carolinus Linnæus, Syst. Nat., ed. XII, 490, 1766, Carolina.

? *Trachynotus argenteus* Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 410, 1831, New York and Rio Janeiro.

Trachynotus cupreus Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 414, 1831, Martinique.

Trachynotus pompanus Cuvier & Valenciennes, Hist. Nat. Poiss., VIII, 415, 1831, Brazil and Charleston, S. C.

Trachinotus carolinus, Jordan & Evermann, l. c., 944, 1896.

Family XXXVII. STROMATEIDÆ. The Fiatolas.

Body compressed and more or less elevated, covered with small or minute cycloid scales. Profile anteriorly blunt and rounded. Mouth small. Premaxillaries not protractile. Dentition feeble; no teeth on vomer or palatines; pharyngeals little developed; œsophagus armed with numerous horny, barbed, or hooked teeth. Opercular bones smooth, not serrate. Gills 4, a slit behind fourth. Gill-membranes either separate and free from (*Stromateinæ*) or broadly joined to isthmus (*Stromateoidinæ*), restricting gill-openings to sides, as in *Chatodipterus*. Gillrakers rather long. Pseudobranchiæ present. Cheeks scaly. Preopercle entire or serrate. Lateral line well developed. Dorsal fin single, long, with the spines few or weak, often obsolete; anal fin long, similar to soft dorsal, usually with 3 small spines, which are often depressible in a fold of skin; ventrals thoracic, 1, 5 in the young, but reduced or altogether wanting in the adult; caudal fin well forked. Usually no air-bladder. Pyloric cæca commonly numerous. Vertebrae 30 to 36.

a. Dorsal and anal fins very high in front, anterior lobe falcate; body suborbicular..... PEPRILUS, 62

aa. Dorsal and anal fins moderately elevated in front, anterior lobe scarcely falcate; side with a series of large, wide-set pores above lateral line..... PORONOTUS

Genus 62. PEPRILUS* Cuvier. The Butter-fishes.

Body ovate or suborbicular, strongly compressed, tapering into a slender caudal peduncle, which is not keeled or shielded. Head short, compressed, profile obtuse. Mouth small, terminal, jaws subequal. Premaxillaries not protractile. Jaws each with single series of weak teeth. Scales very

*The name *Rhombus* Lacépède (1800), which has long been in use for this genus, is preoccupied by *Rhombus* Humphreys (1797), a genus of mollusks. The next available name is *Peprilus* of Cuvier (1817).

small, cycloid, silvery, loosely inserted, extending on vertical fins. Opercular bones entire. Gill-membranes separate, free from isthmus; gillrakers moderate. Lateral line continuous, concurrent with back. Dorsal fin long, more or less elevated in front, preceded by a few indistinct spines; usually one or more procumbent spines in front of dorsal and anal, each of these with a free point both anteriorly and posteriorly; anal fin similar to dorsal, or shorter, usually with three small spines; ventral fins wanting; a single small, sharp spine, attached to pubic bone, occupying the place of the ventrals; pectorals long and narrow; caudal widely forked.

There are but few species of this genus, mostly American. *Peprilus* differs from *Stromateus* chiefly in the prominence of the pelvic bone, which projects in a lamina beyond the skin. Species of *Stromateus* occur in Europe and South America, but none within our limits.

Only one of the two recognized species is known from Porto Rico.

- a. Dorsal rays III, 43 *paru*, 99
- aa. Dorsal rays about III, 39..... *xanthurus*




FIG. 39.—*Peprilus paru*.

99. *Peprilus paru* (Linnaeus). “*Palometa*”; *Harvest-fish*; *Puppy-fish*.

Head 3.6; depth 1.5; eye 3; snout 4.7; maxillary 3.5; mandible 3; interorbital 2.5; D. III, 42; A. II, 39; pectoral and caudal 2.5 in body; scales about 90.

Body nearly circular in outline, strongly compressed, with a very short and slender caudal peduncle; head small, not at all pointed, its anterior outline scarcely breaking the nearly even curve from above and below; mouth very small, oblique, lower jaw not projecting, maxillary reaching front of eye; teeth in jaws small and weak, uniserial; lateral line unarmed; dorsal and anal fins falcate, their anterior rays much produced, those of anal reaching base of caudal or beyond, those of dorsal shorter, reaching middle of fin when depressed; caudal lobes long, equal, the fin well forked; ventrals wanting, a small spine similar to anal spines in their place; scales somewhat deciduous.

Color in life: Pale-blue above, silvery on sides and below, rich bluish or purplish iridescence everywhere; inside of mouth and tongue inky-black, the jaws pale.

Atlantic coast of United States from Massachusetts south to Jamaica, Porto Rico, and Brazil.

This species reaches a length of 6 to 8 inches, a weight of a half pound or more, and is a highly valued pan-fish, particularly common along our south Atlantic coast. It is apparently not common in Porto Rico, as only two specimens were obtained, 6.5 and 7.5 inches long, respectively, from the San Juan market.

- Paru Brasiliense congener*, Sloane, Hist. Jamaica, 285, 1727, Jamaica.
Stromateus paru Linnæus, Syst. Nat., ed. X, 248, 1758, Jamaica; based on Sloane.
Chatodon atepidotus Linnæus, Syst. Nat., ed. XII, 460, 1766, Charleston.
Sternoptyx gardenii Bloch & Schneider, Syst. Ichth., 494, 1801, Charleston; after Linnæus.
Stromateus longipinnis Mitchell, Trans. Lit. & Philos. Soc. N. Y., I, 1815, 366, New York.
Rhombus paru, Jordan & Evermann, l. c., 966, 1896.
Peprilus paru, Jordan & Evermann, l. c., 3197, 1900.

Family XXXVIII. CHEILODIPTERIDÆ. The Cardinal-fishes.

Body oblong or elongate, sometimes compressed and elevated, covered with rather large scales, which are striated and ctenoid or sometimes cycloid; cheeks scaly; lateral line continuous; cleft of the mouth wide, oblique; villiform teeth on jaws and vomer, and sometimes on palatines; canines sometimes present (teeth wanting in *Brephostoma*); preopercle with a double ridge, its edge entire or slightly serrated; opercular spine little developed; lower pharyngeals separate, with sharp teeth; pseudobranchiæ present; branchiostegals 6 or 7. Dorsal fins well separated, the first with 6 to 9 rather strong spines; no dorsal sheath or furrow; anal fin short, usually with 2 spines, sometimes with 3 or 4; ventral fins thoracic, 1, 5, without axillary scale.

Small fishes of the tropics, especially abundant in the East Indies, some of them in fresh water, most of them in rather deep waters. Color often bright-red. Genera about 15; species about 130.

CHEILODIPTERINÆ:

- a. Body oblong, not greatly compressed nor greatly elongate; anal spines 2, rarely 3; teeth present in jaws at least.
 b. Anal fin with two spines; soft rays usually 8 or 9.
 c. Canines none; teeth all villiform; lateral line normal.
 d. Palatines with teeth.
 e. Scales large, 20 to 30 in lateral line.
 f. Preopercle with its posterior edge distinctly serrate, at least in young APOGON, 63
 ff. Preopercle with its ridges entire at all ages APOGONICHTHYS, 61
 cc. Scales small, 40 to 45 in lateral line GLOSSAMIA
 dd. Palatines toothless; teeth moderate; eye very large; body elongate; scales rather small; preopercle entire or nearly so EPIGONUS
 cc. Canines present in jaws. Dorsal spines 6; opercle unarmed; anal spines 2 CHEILODIPTERUS
 bb. Anal fin with 3 spines and 8 soft rays; no teeth on vomer or palatines; no canines; caudal rounded; opercles entire; body moderately elongate AMICHTHYS

SCOMBROPINÆ.

- aa. Body elongate, the form approaching that of the barracuda (*Sphyraena*); mouth large; anal spines 3 or 4.
 g. Teeth subequal, without distinct canines; preopercle serrate. D VII-1, 10; A. III, 7; scales small, about 60. SPHYRENOPS
 gg. Teeth unequal, jaws with long canines; preopercle entire or nearly so.
 h. Second dorsal and anal long, each of 12 to 14 soft rays; soft parts of vertical fins scaly; scales rather small, 45 to 50. SCOMBROPS
 hh. Second dorsal and anal short, each with 7 to 9 soft rays; vertical fins scarcely scaly; scales large, about 30. HYPOCLYDONIA

Genus 63. APOGON Lacépède. Kings of the Mulletts.

Body oblong, compressed, covered with large ctenoid scales. Lateral line continuous, with 20 to 30 scales. Head large; mouth wide, oblique, maxillary extending to below middle of large eye; villiform teeth on jaws, vomer, and palatines; no canine teeth; preopercle with a double ridge, the edge somewhat serrate, at least in the young, becoming entire with age in some species; opercle with spine behind. Gillrakers rather long. Dorsal spines 6 or 7, strong; second dorsal remote, short; anal with 2 spines and 8 or 9 soft rays, second much the longer, soft part similar to soft dorsal; pectorals and ventrals moderate; vertebrae 11 + 14 = 25. Warm seas; the species numerous.

- a. Base of caudal with a distinct round blackish blotch (rarely wanting).
 b. Base of soft dorsal without blackish saddle-like blotch.
 c. Opercle without dark spot; cheek without evident dark dots; soft dorsal much higher than spinous; scales 28 to 30 *imberbis*
 aa. Base of caudal without blackish blotch.

- d. Soft dorsal with a round black blotch below it and a similar one on caudal peduncle above; scales 26... *maculatus*
 dd. Soft dorsal and caudal peduncle without round black spots.
 e. Body with few black specks or none; a blackish bar between last rays of soft dorsal and anal; another on caudal peduncle..... *binotatus*
 ee. Body covered everywhere with blackish dots, like fly-specks; no other distinct markings; scales 25... *pigmentarius*
 aaa. A broad saddle-like dark blotch on caudal peduncle; scales 27..... *sellicauda*, 100

100. *Apogon sellicauda* Evermann & Marsh, new species.

Head 2.6; depth 3; eye 2.7; snout 4.75; maxillary 2; mandible 2.2; interorbital 4; D. vi-i, 9; A. ii, 7; scales 2-27-10. Body rather short; head large; caudal peduncle long, deep, and compressed; mouth large, little oblique, maxillary reaching past pupil; eye very large; preopercle finely serrate; scales large, finely ctenoid; lateral line complete, following curve of back. Fins moderate; second and third dorsal spines longest; first anal spine very short, second about equal to eye; pectoral long, 1.65 in head.

Color in life: Rich scarlet, nearly uniform; a jet-black spot as large as pupil on opercle and another of same size on side between lateral line and base of soft dorsal; a dark blotch from opercular spot to eye, beneath which are a few brownish specks; axil of pectoral somewhat dusky; top of head and snout with small black specks; caudal peduncle with a broad saddle-like dusky area close to base of caudal fin; a few small dark specks on dorsal, anal, and caudal fins.

In alcohol, the color fades to pale yellowish-white, the black spots and specks persisting.




FIG. 40.—*Apogon sellicauda*.

This interesting species is evidently close to *A. maculatus*, but differs in its much larger eye and in coloration. In four specimens of *A. maculatus* examined the eye was 3.25, 3.3, 3.5, and 3.5 in head, respectively. A specimen of the same size as the type of the present species has the eye 3.5 as against 2.7 in said type. The coloration is also very different. In *A. maculatus* there is a definite lateral black spot about the size of the pupil on the upper posterior part of the caudal peduncle; the opercle and postocular region in some specimens have a few minute dark specks sometimes forming an obscure opercular blotch; and the tips of the anterior dorsal and anal rays and the outer caudal rays are quite black. In *A. sellicauda*, instead of the definite black spot on the caudal peduncle, there is a broad dark saddle-like area completely surrounding the peduncle except, possibly, at the under edge; there is a definite black spot nearly as large as that under soft dorsal, on the middle part of the opercle, and from this a broad dark band runs forward to the eye. The tips of the caudal lobes and of the anal and soft dorsal are broken in the type, but they do not seem to have been black, as in *A. maculatus*.

This species is based upon a single specimen, 1.63 inches long, obtained February 11, 1899, on a coral reef at Culebra Island. Type No. 49529 U. S. N. M. (*sella*, saddle; *cauda*, tail.)

Genus 64. APOGONICHTHYS Bleeker.

This genus differs from *Apogon* only in having the preopercle entire at all ages; scales very large (20 to 26) and cycloid. Small species, similar in habit to those of *Apogon*, found in the tropical seas. The genus is scarcely distinct from *Apogon*.

- a. Scales in lateral line 21 to 23; body with many dark points.
 b. Ventrals short, not reaching vent; dorsal rays VI-1, 9 *alutus*, 101
 bb. Ventrals long, extending beyond anal; dorsal rays VII-1, 9 *stellatus*
 aa. Scales in lateral line 30; ventrals long, reaching beyond front of anal; body everywhere with black specks; dorsal VI-1, 9 *puncticulatus*




FIG. 41.—*Apogonichthys alutus*.

101. *Apogonichthys alutus* (Jordan & Gilbert).

Head 2.4; depth 2.75; eye 3; snout 4.75; maxillary 2; mandible 1.8; interorbital 4; D. VI-1, 9; A. 11, 9; scales 1-22-7. Body short and deep; head compressed, short and high, its height at occiput six-sevenths its length; snout short and blunt, less than interorbital width, about half diameter of orbit; mouth very oblique, maxillary reaching beyond pupil, but not to posterior margin of orbit; length of maxillary 1.75 in head; teeth in narrow villiform bands in each jaw, those on vomer and palatines minute; eye of moderate size, 2.8 in head; orbital rim elevated above and behind; interorbital width 3.33 in head, with a low median longitudinal ridge; both ridges of preopercle entire; opercle without spine; gillrakers slender, longest rather more than half diameter of orbit, 8 or 9 on anterior branch of outer arch. First dorsal low, of 6 rather weak spines, its base two-fifths length of head, and equal to greatest height of fin; second dorsal high, longest ray 1.5 in head. Anal similar to second dorsal; second anal spine half length of longest ray, which is contained 1.75 in head; caudal 1.33; ventrals not reaching vent, 1.66, and pectorals 1.75 in length of head.

Color, rusty-red, with silvery luster; sides of head little reddish; body and fins everywhere much soiled and freckled with dark points; first dorsal blackish, thickly punctate; second dorsal, anal, and caudal yellow, smutty, with dark points, posterior half of the caudal more dusky; ventrals smutty-yellow; pectorals colorless.

Found on the Snapper Banks off Pensacola and Tampa and at Porto Rico, where one specimen 2 inches long was seined in Guanica Bay January 28. Until now the only specimens known had been obtained from the stomachs of red snappers, *Neomenus aya*, which probably all came from depths of 20 to 50 fathoms. The finding of this species in shallow water near the shore is therefore an interesting addition to our knowledge of its habits.

Apogon alutus Jordan & Gilbert, Proc. U. S. N. M. 1882, 279, Snapper Banks off west coast of Florida.
Apogonichthys alutus, Jordan & Evermann, l. c., 1110, 1896.

Family XXXIX. CENTROPOMIDÆ. The Robalos.

This family is defined thus by Professor Gill (Proc. U. S. N. M. 1882, 484):

"Typical acanthopterygians with the postorbital portion of the skull longer than the oculo-rostral; the parietals behind the constriction continuous with the epiotics and transverse laminae arising from the supraoccipital crest, the three together forming a well differentiated posterior oblong pentagonal or hastiform area; the re-entering parietal sinus, with its anterior margin, produced forward nearest the opisthotics; the exoccipitals well developed and contiguous above the foramen magnum; the vertebrae in typical number (10 + 14) and longish; the anterior two partly coossified and the first with selliform apophyses extending backward and embracing the second vertebra; the vertebrae mostly with foveae or pits for the ribs and only with developed parapophyses for the posterior (6 to 10) pairs of ribs; the second neural spine suberect, and with laminiform extensions, which embrace the first; neuropophyses and neural spines of other vertebrae depressed at their bases continuous with the zygapophyses in front, and slightly curved upward at their tips; the hæmal spines resembling the neural."

Subocular laminae produced behind in a pointed process. External characters are the elongate body, with elevated back, straight abdomen, and angulated base of anal. Scales ctenoid, varying in size, lateral line conspicuous, extending on caudal fin, the tubes straight, confined to basal half of scale. Head depressed, pike-like, lower jaw projecting; villiform teeth, in bands, on jaws, vomer, and palatines; tongue smooth. Maxillary broad, truncate behind, with a strong supplemental bone. Pseudobranchiae present, small. Preopercle with a double ridge, posterior margin strongly serrated, with larger spines at angle; preorbital and suprascapula serrated, opercle without true spines. Gillrakers long. Dorsal fins well separated, first with 8 spines, first and second short, third and fourth longest; anal with 3 spines, second strong, third long and slender, these fins moving in scaly sheaths. Caudal forked. Ventrals large, 1, 5, inserted well behind pectorals, a scaly process at their base; pectorals narrow and rather pointed, upper rays longer than lower. Branchiostegals 7. Air-bladder well developed, simple, or with appendages anteriorly.

This family comprises about 15 species, all American and referred to one genus. They are game-fishes, excellent as food.

Genus 65. CENTROPOMUS Lacépède. The Robalos.

The characters of this genus are included in the above.

In American waters this genus contains 14 species, as recognized by Jordan & Evermann, only 2 of which are known from Porto Rico. All the species are of some value as food-fishes and some of them reach a large size. They occur on both coasts of tropical America and are fishes of shallow or moderate depths, two or three of them even ascending fresh-water streams for considerable distances.

- a. Preorbital entire or very faintly serrated; spines of fins moderate, none more than half head; lateral line black, with about 70 scales (pores); caudal fin short; air-bladder usually with recurved appendages at its anterior end; anal with 6 soft rays, the last cleft to base. Size large.
- b. Appendages to air-bladder short, shorter than eye; second anal spine projecting beyond third.... *undecimalis*, 162
- aa. Preorbital with well-developed retrorse teeth, especially posteriorly; air-bladder without appendages; spines longer. Size comparatively small.
- c. Lateral line in a narrow black streak.
- d. Scales moderate or large, 50 to 60 in lateral line; second anal spine very strong; ventrals more or less dusky, usually broadly tipped with black; anal rays III, 7, second spine a little shorter than third.
- e. Scales in lateral line 57 to 60; second anal spine very long, 1.25 to 1.60 in head; depth of body 3.66 in length..... *pedimacula*
- cc. Scales in lateral line 51 or 52; second anal spine 1.66 in head. Body more slender, the depth 3.75 in length.. *cuvieri*
- dd. Scales small, about 70 in lateral line; ventrals pale; second anal spine strong, equal to depth of body; anal rays III, 6..... *mexicanus*
- cc. Lateral line pale, not in a dark stripe; ventral fins yellowish, without black tip.
- f. Scales very small, about 80; sides of body parallel with each other; second anal spine longer than third, equal to depth of body; anal rays III, 7..... *parallelus*, 103
- ff. Scales small, 65; second anal spine very long, about equal to depth of body; third anal spine same length; angle of preopercle with about 6 long, comb-like teeth; anal rays III, 7..... *pectinatus*
- fff. Scales large, 49 to 53 in lateral line; eye moderate, about 6 in head; anal rays III, 6.
- g. Body moderately elongate, the depth 3.25 to 3.66 in length.
- h. Scales before dorsal small, 16 to 18 in number; maxillary reaching past front of pupil. Third dorsal spine half head; scales 51..... *armatus*
- hh. Scales before dorsal not crowded, 10 to 14 in number; maxillary barely reaching front of pupil; third dorsal spine less than half head; scales 7-53-11..... *affinis*
- gg. Body more elongate, depth 4 in length; the second anal spine excessively long, 1.4 times depth of the body; scales 53..... *ensiferus*

102. *Centropomus undecimalis* (Bloch). "Robalo"; Snook.

Head 3; depth 4.25; eye 7.75; snout 3.5; maxillary 2.5, reaching past middle of eye; interorbital equal to eye; D. VII or VIII-1, 10; A. III, 6; scales 9-75-12, about 18 before dorsal; gillrakers 4+9. Body stout, not much compressed; head long and pointed; mouth large; lower jaw strongly projecting; preorbital faintly or not at all serrate; preopercle and interopercle serrate; subopercular flap broad; cranial ridges prominent; caudal peduncle stout, its least width 2.5 in its least depths; fins moderate; first and second dorsal spines very short, first sometimes absent, third about 2 in head; first anal spine very short, second very strong, about 2.6 in head, usually a little longer than third, which is more slender; pectorals long, about 2 in head; ventrals long, slightly greater than the pectoral; scales moderate, closely imbricated, rather firm.

Color in life: Irregularly iridescent-green above; top of head grayish, cheeks silvery, with golden wash; lower jaw flesh-color, with some blue; lateral line darkest posteriorly; side silvery, with light purple iridescence; under parts white; fins all pale, ventrals darker at tips, caudal dark-edged.

Color in spirits: Greenish-olive, sides dull-silvery, lighter below; fins all pale except dorsals, which are dark; lateral line black.

A common food-fish in Porto Rico, where it reaches a length of 2 or 3 feet or more. Specimens were obtained at San Juan, Ponce, Palo Seco, Mayaguez, and Arroyo.

Sciæna undecimalis Bloch, Ichth., VI, 60, pl. 303, 1792, Jamaica.

Centropomus undecimradiatus Lacépède, Hist. Nat. Poiss., IV, 268, 1802, Jamaica; after Bloch.

Perca loubina Lacépède, Hist. Nat. Poiss., IV, 397, 1802, Cayenne.

Sphyræna aurcoriviridis Lacépède, Hist. Nat. Poiss., V, 324, 1803, Martinique.

Centropomus appendiculatus Poey, Memorias, II, 119, 1860, Havana and Cienfuegos.

Centropomus undecimalis, Poey, Fauna Puerto-Riqueña, 321, 1881; Stahl, l. c., 76 and 162, 1883; Jordan & Evermann, l. c., 1118, 1896.




FIG. 42.—*Centropomus undecimalis*.

103. *Centropomus parallelus* Poey. "Robalo."

Head 2.7; depth 3.33; eye 5.7; snout 3.4; maxillary 2.4, reaching past middle of eye; mandible 1.6; interorbital 7, less than eye; D. VIII-1, 10; A. III, 7; scales 12-80-14. Back considerably elevated, making body deeper and anterior profile more steep than in *C. undecimalis*. It differs also from that species in the much less projecting lower jaw, stronger serrations of preorbital and preopercle, smaller scales, and color of the lateral line, which is pale instead of black.

Our specimens differ slightly from Poey's description of *C. parallelus*: Second anal spine but two-thirds depth of body, third very nearly as long as second; otherwise identical with it.

Color, silvery in life, inclining to brassy-olive on back; in spirits, faint longitudinal streaks, pale below but brownish on back, traverse the rows of scales.

This is a small species, rarely exceeding a foot in length, found about Cuba, Santo Domingo, Porto Rico, and at Pernambuco. It enters rivers and lakes and may be found at long distances from salt water. In Porto Rico it ascends the larger streams well toward the interior of the island. Numerous examples from the Río Loiza were seen hawked about the streets of Caguas. One from this place is a foot in length. *C. undecimalis* also enters the rivers but does not ascend so far as does this one. Each species possesses some game qualities and both are sought by the local anglers. The best fishing is said to be in the lower portions of the Río de la Plata, Manatí, and the Río Grande de Arecibo.

Centropomus parallelus Poey, Memorias, II, 120, 1860, Havana and Cienfuegos; Jordan & Evermann, l. c., 1122, 1896.

Family XL. SERRANIDÆ. The Sea Basses.

Body oblong, more or less compressed, covered with adherent scales of moderate or small size, which are usually but not always ctenoid; dorsal and ventral outlines usually not perfectly corresponding. Mouth moderate or large, not very oblique, the premaxillary protractile, and broad maxillary usually not slipping for its whole length into a sheath formed by preorbital, which is usually narrow. Supplemental maxillary present or absent. Teeth all conical or pointed, in bands, present on jaws, vomer, and palatines. Gillrakers long or short, usually stiff, armed with teeth. Gills 4, a long slit behind fourth. Pseudobranchiæ present, large. Lower pharyngeals rather narrow, with pointed teeth, separate (except in *Centrogenys*). Gill-membranes separate, free from isthmus. Branchiostegals normally 7 (occasionally 6). Cheeks and opercles always scaly; preopercle with its margin more or less serrate, rarely entire; opercle usually ending in one or two flat spine-like points. Nostrils double. Lateral line single, not extending on caudal fin. Skull without cranial spines and usually without well-developed cavernous structure. No suborbital stay. Post-temporal normal. Second suborbital with an internal lamina supporting globe of eye; entopterygoid present; all or most of the ribs inserted on transverse processes when these are developed; anterior vertebrae without transverse processes. Dorsal spines usually stiff, 2 to 15 in number; soft dorsal with 10 to 30 rays; anal fin rather short, its soft rays 7 to 12, its spines, if present, always 3, in certain genera (*Grammistina*, *Rypticina*) altogether wanting. Ventrals thoracic, usually 1, 5 (1, 4, in *Plesiopina*), normally developed, without distinct axillary scale. Pectoral well developed, with narrow base, rays branched. Caudal peduncle stout, the fin variously formed. Vertebrae typically $10 + 14 = 24$, number sometimes increased, never more than 35. Air-bladder present, usually small, and adherent to wall of abdomen. Stomach cæcal, with few or many pyloric appendages; intestines short, as is usual in carnivorous fishes.

The *Serranidæ* include 60 to 70 genera and about 400 species; carnivorous fishes, chiefly marine, and found in all warm seas; several genera found in fresh waters, comprising most of the family of *Percidæ*, as understood by Günther and others, exclusive of those with imperfect pseudobranchiæ, those with one or two anal spines, those with the number of vertebrae increased, those in which the whole length of maxillary slips under preorbital, and those with anal fin many-rayed, and cranium shortened behind.

- a. Anal spines 3, well developed. Dorsal fin single, sometimes deeply divided.
- b. Maxillary with a distinct supplemental bone (rarely obscured by the skin); dorsal usually divided or deeply notched
- c. Inner teeth of jaws not depressible nor hinged.

LIOPROPOMINÆ:

- d. Soft dorsal longer than spinous part; dorsal deeply divided, spines 6 to 9 in number; preopercle entire; vertebrae $10 + 14 = 24$; lateral line arched anteriorly.
- e. Dorsal spines 9; caudal lunate LIOPROPOMA
- ee. Dorsal spines 6; caudal truncate CHORISTIUM

POLYPRIONINÆ:

- dd. Soft dorsal shorter than spinous part; vertebrae more than 24 (25 to 36); head without rugose dermal ossifications. Teeth all villiform, without canines; soft dorsal with 10 to 12 rays. Head armed with rough spinigerous crests, there being spinous projections above the eyes and a rough, bony ridge on opercle, with others on post-temporal; dorsal fin low, continuous; tongue with teeth; dorsal spines 11 or 12; soft dorsal scaly; caudal rounded; ventral not inserted before axil of pectoral; pyloric cæca numerous (about 70); vertebrae 27 POLYPRION

EPINEPHELINÆ:

- ee. Inner teeth of jaws depressible or hinged; canine teeth more or less distinct, in front of each jaw; scales small, firm, the top of head more or less scaly; lateral line running low (except in *Gonioplectrus*, etc.); supraoccipital crest usually more or less encroaching on the top of the skull, so as to leave no distinct smooth area at vertex (except in *Variola*); temporal crests usually distinct; gillrakers various, generally small and short. Dorsal rays VIII to XIV, 12 to 20, number of spines usually not 10; anal rays III, 7 to III, 12; ventral fins inserted more or less behind axil of pectoral; head unarmed, except for opercular spines and serræ on preopercle; soft dorsal scaly; scales of lateral line usually triangular and cycloid; vertebrae almost always $10 + 14 = 24$, rarely 26 or 27. Chiefly shore fishes, often of large size; all of them, so far as known, bisexual.
- ff. Pectoral unsymmetrical, its upper rays longest; dorsal spines 8; plectroid spine on preopercle single, very strong; a strong canine on middle of side of lower jaw; opercle with a long, knife-shaped spine; body rather deep; lateral line running high; jaws naked; scales small, firm, and rough; caudal rounded; soft dorsal rather short, of 12 or 13 rays GONIOPECTRUS
- fff. Pectoral rounded, symmetrical, its middle rays longest; canines usually distinct, in front of one or both jaws.
- gg. Frontals with a transverse ridge on posterior part in front of supraoccipital connecting the parietal crests; frontal bones without processes or longitudinal ridges on upper surface; dorsal spines always 9. Posterior process of premaxillary extending to between frontals; mandible without curved canines on its sides; caudal not forked; scales ctenoid PETROMETOPON, 66
- hh. Frontals without transverse ridge.
- ii. Dorsal spines 9, soft dorsal of moderate length and height, its rays 13 to 15; anal rays III, 7 or 8; skull and head essentially as in *Epinephelus*, snout not very short, frontal region flat or convex, supraoccipital crest continued forward over it; lateral crest short, low and diverging; mouth and teeth as in *Epinephelus* BODIANUS, 67

- hh. Dorsal spines 11 (rarely 10, never 9).
- i. Parietal crests not produced forward on frontals; frontals with process or knob on each side, behind interorbital area; premaxillary processes fitting into a cavity at anterior extremity of frontals, or into an emargination of these bones; anal rays III, 8, or III, 9.
- j. Scales of the lateral line normal, marked by radiating ridges.
- k. Cranium narrow above interorbital space, deeply concave; occipital crest meeting interorbital region. EPINEPHELUS, 68
- kk. Cranium very broad and flat above; interorbital little concave, occipital crest disappearing before reaching interorbital region GARRUPA
- jj. Scales of lateral line each with 4 to 6 strong radiating ridges; cranium short, extremely broad, and depressed between eyes; anterior profile of head a little concave; dorsal spines low; dorsal rays XI, 16 PROMICRODUS
- ii. Parietal crests produced forward on frontals.
- l. Frontals with a process or knob on each side behind interorbital area; premaxillaries fitting into a cavity at anterior extremity of frontals; anal rays III, 8, rarely III, 9.
- m. Preopercle with a single antrorse hook or spine near angle; supraoccipital and parietal crests not extending to between orbits; scales ctenoid ALPHESTES, 69
- mm. Preopercle without antrorse spine; supraoccipital and parietal crests extending to between orbits; scales smooth. DERMATOLEPIS
- ll. Frontals without processes on upper surface; parietal crests extending to between orbits; premaxillary processes not extending to frontals. Anal fin elongate, its rays III, 11, or III, 12 (very rarely III, 9 or III, 10) caudal fin lunate or truncate; dorsal rays XI, 16 to 18 MYCETOPERCA, 70
- bb. Maxillary without supplemental bone; canine teeth, if present, usually developed on side of lower jaw as well as in front; no depressible teeth; scales mostly ctenoid, including those of lateral line; tubes of lateral line straight or with an ascending tubule, covering most of length of scale. Temporal crests on cranium almost obsolete.

SERRANINÆ:

- n. Gillrakers comparatively short and wide apart; lateral line not running close to back (except in *Serranus*); dorsal rays X, 11 to 15; anal rays usually III, 7; supraoccipital crest not extending far forward on top of skull, a more or less distinct convex smooth area being left on vertex between supraoccipital and interorbital area; mouth not very oblique; vertebrae about 10 + 14 = 24. Chiefly shore fishes of olivaceous colors.
- o. Ventral fins inserted below, or more or less behind, axil of pectoral; branchiostegals 7.
- p. Dorsal without long filamentous spines, not more than one of its spines specially produced. Body short and deep, with elevated back, depth more than two-fifths length, usually nearly half; preopercle with a few antrorse serrae on its lower limb; top of supraoccipital crest very high, about as long as smooth area on vertex of cranium, which is well developed, as in *Serranus* and *Prionodes*. Top of head naked; dorsal rays usually X, 14. HYPOPLECTRUS
- oo. Ventral fins anterior, inserted more or less in advance of axil of pectoral, well separated; upper half of pectoral fin usually vertically truncate.
- q. Smooth area on top of cranium very short and small; the long supraoccipital crest encroaching on posterior border of cranium so that the latter in profile is not nearly vertical along occipital region. Caudal fin not lunate, rounded, or ending in 3 points, middle rays produced like outer ones. CENTROPRISTES
- qq. Smooth area on top of cranium very large, longer than the supraoccipital crest, which is low and short; posterior border of cranium at occipital region nearly vertical in profile. Caudal fin lunate or truncate.
- r. Branchiostegals 7; caudal fin forked or lunate; none of dorsal spines elongate.
- s. Preopercle with numerous strong diverging spines at its angle, these spines diverging from one or two centers; preorbital broader than maxillary, which is widest near its middle; scales rather large DIPLECTRUM, 71
- ss. Preopercle simply and rather finely serrate; preorbital narrow PRIONODES, 72
- rr. Branchiostegals 6; caudal fin truncate DULES, 73
- nn. Gillrakers (in American species) very long, slender, and close-set; lateral line running close to back; supraoccipital crest high; occiput with a short convex smooth area; canines strong; no depressible teeth; preorbital narrow; maxillary without supplemental bone or with a rudiment only.

ANTHINÆ:

- t. Lateral line complete and continuous, extending to base of caudal; ventral rays I, 5; dorsal rays IX to XII, 9 to 20; anal rays III, 7 to 10; pectoral rays branched; snout short, mostly convex in profile. Fishes of rather deep waters, chiefly bright-red in life.
- u. Dorsal spines 9, all low; soft rays about 19. Caudal fin deeply forked, lobes produced; scales small, ctenoid; ventrals long, inserted behind axil of pectoral; maxillary scaly; frontal region flattish, supraoccipital crest very prominent PARANTHIAS
- uu. Dorsal spines 10 or more; scales not very small; preopercle angular, with salient teeth at its angle; one or more dorsal spines sometimes filamentous; ventral fins long.
- v. Maxillary and frontal region naked; tongue mostly toothless; caudal fin lunate; parietals weak; posterior process of premaxillary reaching frontals. Ventrals inserted behind axil of pectoral; scales 50 to 60 HEMIANTHIAS
- vv. Maxillary scaly; top of head scaled to snout.
- w. Pterygoids toothless; tongue with few teeth or none; caudal forked ANTHIAS
- ww. Pterygoids with a large patch of teeth; tongue toothed; parietal crest strong, extending to above eye; posterior processes of premaxillary not reaching frontals; caudal truncate, with outer rays much produced. OCYANTHIAS
- tt. Lateral line interrupted, running close to back, beginning again on caudal peduncle.

GRAMMINÆ:

- z. Ventral rays 1, 5; preopercle serrate; caudal convex; scales rather large, somewhat ctenoid; dorsal spines 12. GRAMMA

RYPHTICINÆ:

- aa. Anal spines wanting; dorsal spines 2 to 4 only; preorbital narrow; no canine teeth; scales small, smooth, embedded. RYPHTICUS, 74

Genus 66. **PETROMETOPON** Gill. *Enjambres*.

Frontal bones with an anterior groove or excavation for reception of posterior processes of the premaxillaries, without processes on upper surface; a curved or angular ridge across posterior portion of frontals in front of supraoccipital, connecting parietal crests; supraoccipital and parietal crests not produced forward. Dorsal spines 9; anal rays mostly 11, 8; scales ctenoid; otherwise essentially as in *Epinephelus*.

Species rather few, mostly of small size, distinguished from *Bodianus* chiefly by the peculiarities of the frontal bones, the above account being taken from Boulenger, Cat., 1, 175.

104. Petrometopon cruentatus (Lacépède). "*Cabrilla*"; *Enjambre*; *Coney*; *Red Hind*.

Head 2.5; depth 2.8; eye 5.5; snout 4; maxillary 2; mandible 1.6; interorbital 6.7; preorbital 11; D. IX, 14; A. III, 8; pectoral 1.7; ventral 2.1; caudal 1.7; scales about 12-85 to 95-30.

Body not greatly elongate, somewhat compressed; caudal peduncle short; head moderate, pointed; mouth large, the large maxillary reaching beyond eye, lower jaw moderately projecting; teeth much as in *Epinephelus*, depressible teeth long and slender; outer row on sides of upper jaw enlarged but not depressible; canines in front of both jaws, the upper somewhat larger; preopercle convex, with very minute serrations; opercle ending in 3 flat, exposed spines, opercular flap thin and rounded or obtusely pointed. Scales ctenoid. Dorsal fin continuous, with a slight emargination; spines slender and sharp; caudal rounded or almost double-truncate; pectoral reaching considerably beyond tip of ventrals, which do not reach vent.

Color in life, reddish-gray with red spots nearly everywhere. Our single specimen, in spirits, is a rather dark brown, the spots appearing brown posteriorly and pale forward, largest and plainest on head; on anterior part of body they have faded almost entirely; the four larger very black spots along base of dorsal are still visible; fins spotted and barred; caudal edged with a moderate band of dark, outside of which is a very narrow pale edging.

This fish ranges from Florida and West Indies to Brazil; is common about Jamaica and Cuba, and not uncommon on the reefs about Key West, where it is called "coney." A handsome species of considerable importance as a food-fish; length about a foot. One specimen, 8.5 inches, from Isabel Segunda, obtained from the local fishermen.

Sparus cruentatus Lacépède, Hist. Nat. Poiss., IV, 157, pl. 4, fig. 1, 1803, Martinique; on a copy of a drawing by Plumier.

Serranus apiarius Poey, Memorias, II, 143, 1860, Havana.

Petrometopon cruentatus, Jordan & Evermann, l. c., 1141, 1896.

Genus 67. **BODIANUS** Bloch.

This genus is close to *Epinephelus*, from which it is separated mainly by the presence of 9 spines in dorsal fin instead of 11. In character of cranium the two genera differ little, the skull above having its bones thin and smooth, the angular ridge on posterior part of frontals being wanting, and parietal and supraoccipital crests not extending on frontals.

Species rather numerous in warm seas, of small size and bright color. Of seven American species and subspecies of *Bodianus* recognized by Jordan & Evermann two were obtained in Porto Rico.

BODIANUS:

- a. Scales etenoid; none of dorsal spines elevated.
 b. Caudal fin not lunate; head and body with few or many small, blue, dark-edged spots.
 c. Caudal fin rounded, the middle rays longest; snout with 1 or 2 blue stripes; back of tail without conspicuous black blotch; scales small; lateral line about 115 *teniops*
 cc. Caudal fin truncate, middle and outer rays about equal; snout without stripes; back of tail with 2 black spots; lower jaw with a black spot at tip; scales moderate; lateral line about 90.
 d. Ground-color lemon-yellow; blue spots few *fulvus*
 dd. Ground-color bright scarlet *ruber*, 105
 ddd. Ground-color brown *punctatus*, 106

MENEPIHORUS:

- bb. Caudal fin lunate, with prominent angles; body covered with blue spots.
 c. Body rather deep; preopercle evenly convex, without salient angle; mouth small, lower jaw much projecting; gillrakers slender, long, about $x + 20$; color earmine, head, back, and sides more or less covered with blue, dark-edged points; caudal tips black; some dark spots on maxillary and about eye; no black blotch on caudal peduncle or on tip of lower jaw *dubius*
 ee. Body more slender, depth 4 in total length; eye 5.5 in head *punctiferus*

105. *Bodianus ruber* (Bloch & Schneider). "Fino"; *Red Guatireve*; *Ouatilibi*.

Head 2.6; depth 3; eye 5.3; snout 3.8; maxillary 2.1; mandible 1.7; interorbital 6; preorbital 9; D. IX, 15; A. III, 9; pectoral 1.6; ventral 2.1; caudal 1.7; scales about 12-90 to 110-32.

Body elongate, not greatly compressed; mouth moderate; maxillary not extending beyond eye in our specimens; lower jaw projecting; teeth about as in *Petrometopon cruentatus*, depressible teeth smaller; preopercle with weak serrations, its outline convex, with a very slight emargination; opercle with three large, flat exposed spines, and a pointed opercular flap. Scales ctenoid. Dorsal spines slender and sharp, of nearly equal length, save the first and second, which are shorter; caudal truncate; second anal spine somewhat stronger than third, about 3.5 in head; pectoral pointed, reaching far past tip of ventral, 1.7 in head; ventral not reaching vent, 2.1 in head.

Color in life: Body and head rich rosy-red, darkest above, palest on belly; back and sides to lower level of pectoral, as well as head, with numerous small round blue spots, those on head largest; head with a few similar black or darkish spots; dorsal and anal fins blood-red, with black edge; caudal pale-red; pectoral orange-red; ventrals blood-red, with slight black border; inside of mouth flesh-color; two large black spots on tip of lower jaw and two similar black spots on dorsal side of caudal peduncle. In spirits the color is pale-yellowish, in some specimens brownish-gray; the spots mostly remain blue, but some change to gray and brown; caudal dusky, edged with two narrow bands, the inner dark, outer pale, these faded in some specimens.

Two specimens, 8 to 9 inches long, from Arroyo, February 4; called "fino" by the local fishermen. This good food-fish is fairly common throughout the West Indies to Brazil; usually in moderate depths.

Carauna, Maregrave, Hist. Brasil., 147, 1648, Brazil.

Guatireve, Parra, Descr. Dif. Piezas, Hist. Nat., pl. 5, fig. 1, 1787, Cuba.

Gymnocephalus ruber Bloch & Schneider, Syst. Ichth., 346, pl. 67, 1801, Brazil; on *Carauna* of Maregrave.

Serranus ouatilibi Cuvier & Valenciennes, Hist. Nat. Poiss., 71, 381, 1828, Havana.

Serranus carauna Cuvier & Valenciennes, Hist. Nat. Poiss., II, 384, 1828, Brazil.

Bodianus fulvus ruber, Jordan & Evermann, l. c., 1145, 1896.




FIG. 41.—*Bodianus punctatus*.

106. *Bodianus punctatus* (Linnaeus). *Nigger-fish*; *Black Guatireve*.

Not differing materially from the preceding, except in color. Color in life, brownish or blackish-olive; spots everywhere on sides and head, dark-blue with light-blue centers; dorsal fin dusky-olive, edged with darker, a few spots on its base; soft dorsal margined with whitish; caudal dusky-olive; anal and ventrals violaceous-black; pectorals olivaceous; the spots in spirits become brown, with gray centers. Length 8.5 to 12 inches.

This species is common in the West Indies, north to Florida. One specimen from Puerto Real and three from San Geronimo. *B. ruber* and *B. punctatus* have usually been regarded as subspecies of *B. fulvus*, but until the fact of intergradation is established it is best to treat them as distinct species.

Perca marina punctulata, Catesby, Hist. Carolinas, etc., pl. 7, 1743, Bahamas.

Perca punctata Linnaeus, Syst. Nat., X, 1758, 291, Bahamas; based on Catesby.

Perca punctulata Gmelin, Syst. Nat., 1315, 1788, Bahamas; after Catesby.

Enneacentrus punctulatus Poey, Fauna Puerto-Riqueña, 319, 1881; Stahl, l. c., 162, 1883.

Bodianus fulvus punctatus, Jordan & Evermann, l. c., 1146, 1896.

Genus 68. EPINEPHELUS Bloch. The Groupers.

Body stout, compressed, covered with small ctenoid scales, which are often somewhat embedded in the skin; scales of lateral line triangular, cycloid; soft parts of vertical fins generally more or less scaly. Cranium narrow above. Parietal crests not produced on frontals, which are without transverse ridge posteriorly; frontals with a process or knob on each side behind interorbital area; premaxillary processes fitting into a notch or cavity on anterior end of frontals. Preopercle moderately serrate behind, its lower limb entire, without distinct antrorse spine; opercle with two strong spines. Nostrils well separated. Mouth large; maxillary large with a well-developed supplemental bone, its surface usually with small scales. Canine teeth few, large in front of jaws; enlarged teeth of inner series of each jaw depressible. Gillrakers short and rather few. Dorsal spines usually 11, rarely 10, not filamentous, last ones somewhat shorter than middle ones. Anal spines 3, second usually the largest; number of soft rays 7 to 9. Caudal fin rounded or lunate. Pyloric caeca few (usually 10-20). Pectoral rounded, shortish, nearly symmetrical, of 15 to 20 rays. Ventrals moderate, inserted below pectorals, close together, each with a strong spine.

Species very numerous, most of them of large size, abounding in all tropical seas, where they are valuable food-fishes. This is the largest and most important genus of the *Serranidae*, and its species are most widely distributed.

I. SCHISTORUS:

Nostrils unequal, posterior much the larger, three times diameter of the anterior; preopercle with two or three small irregular teeth below its angle; pyloric caeca in increased number; head large.

a. Second and third anal spines about equal in length; color brownish, with about 8 darker crossbands; dark bands radiating from eye; a dark mustache above maxillary; a dark blotch on back of caudal peduncle... *mystacinus*

II. EPINEPHELUS:

Nostrils subequal, posterior scarcely larger than anterior; pyloric caeca in moderate number.

b. Second dorsal spine short, lower than third or fourth, fin not much notched; caudal rounded more or less; lower opercular spine inserted farther back than upper. Lateral teeth of lower jaw in more than two rows, at least in adult. Interorbital space of moderate width, its breadth more than half diameter of eye and 7 to 10 times in length of head. Dorsal spines 11; preopercle without distinct spinules on its lower limb.

c. Maxillary naked.

d. Lower jaw strongly projecting.

e. Body and head covered with red or orange spots (dusky in spirits and always darker than the ground-color); vertical fins without dark edge, their bases spotted like the body; body with large pale spots besides the orange spots; the young with large black blotches at base of dorsal; angle of preopercle not salient; form robust..... *adscensionis*, 107

ee. Body and head reddish-brown, adult nearly plain, young with darker spots; vertical fins broadly edged with dark brown. Body robust, depth 3 in length. D. XI, 15 or 16. Caudal fin convex behind; maxillary naked; dorsal spines low, subequal; interorbital moderate, 6.5 in head; preopercle with strong teeth at its angle, lower limb entire..... *guaza*

cc. Maxillary more or less scaly.

f. Preopercle with a more or less distinct salient angle, which is armed with teeth (these teeth occasionally undeveloped in *E. striatus*, which species may be known by presence of black points around eyes).

g. Body without orange or dark-brown spots; spots (if any) brownish or pearly, diffuse or irregular; vertical fins without broad black margin.

h. Caudal peduncle without black, saddle-like blotch above. Caudal fin truncate or emarginate when spread open, not convex behind; maxillary usually more or less scaly; vertical fins without broad edging of black; dorsal fin, or a part of it, distinctly edged with bright yellow; color of body uniform reddish-brown, a clear blue streak from eye to angle of preorbital; a faint dark mustache; no black spots anywhere; whole dorsal with a bright-yellow edging; anal and caudal without pale edging; caudal slightly lunate; maxillary scaly; dorsal rays XI, 14; lower jaw strongly projecting (as in *Epinephelus niveatus*, with which this species seems to agree very closely in all respects except color)..... *flavolimbatus*

hh. Caudal peduncle with a large quadrate saddle-like black blotch above (sometimes wanting in *E. niveatus*, especially in the young).

i. Eye not surrounded by dark points; sides brown, marked with large blotches of steel-blue, these more or less regularly arranged and not distinct on breast; no dark crossbars; lower jaw strongly projecting; caudal fin subtruncate, its angles rather acute; pyloric caeca rather numerous..... *niveatus*

ii. Eye surrounded by conspicuous dark-brown points; body with irregular dark crossbars; angle of preopercle little salient; third dorsal spine highest, 2.5 in head; scales moderate, about 100; caudal rounded; lower jaw little projecting; vertical fins in life broadly edged with yellow..... *striatus*, 108

gg. Body covered with small dark-orange or brown spots; lower jaw not prominent; interorbital space very narrow, not half diameter of eye. Vertical fins broadly edged with blue-black, their bases unspotted; body without pale spots, the orange spots rather small; body rather slender; size small..... *guttatus*, 109

ff. Preopercle without salient angle; body, head, and fins dark reddish-brown, profusely covered with small pearly-white stellate spots, body robust, lower jaw projecting; caudal subtruncate, with sharp angles. *drummond-hayi*

bb. Second dorsal spine elevated, not lower than third or fourth; caudal fin lunate; preopercular angle little salient, without enlarged teeth; interorbital width 7 in head..... *morio*, 110

107. *Epinephelus adscensionis* (Osbeck). *Rock-hind; Cabra Mora.*

(PLATE 11.)

Head 2.4; depth 3.2; eye 6; snout 4.3; maxillary 2.3; mandible 2; interorbital 6.7; preorbital 10; D. xi, 17; A. iii, 8; pectoral 2; ventral 2.5; caudal 2; scales about 100. Body oblong, robust, somewhat compressed posteriorly; head large, pointed; mouth large, maxillary reaching posterior border of eye, lower jaw strongly projecting; teeth in broad bands, canines in front of lower jaw smaller than those of upper; preopercular margin rounded, with a slight emargination, finely serrate; gillrakers short and thick, 9 + 18; middle opercular spine strong and prominent, lower much smaller and upper not evident without dissection; scales ctenoid; fins rather large, dorsal continuous, with a very shallow notch, spines strong; pectoral broad, middle rays longest, reaching beyond tips of ventrals, equal in length to caudal; ventrals not reaching vent; caudal truncate or gently rounded in younger, adult with margin wavy, due to extension of middle branches of the much-branched rays.

Color in life: Olivaceous-gray, with darker clouds; everywhere covered with round orange-brown or reddish spots, largest and reddest on breast, their centers more orange, borders brown; some scattered whitish blotches; five roundish, ill-defined black blotches along side of back, those under dorsal fin extending upon it and disappearing with age; mouth pale within, roof with red spots; dorsal and anal spotted similarly to body, without dark edge, edge of anal red on anterior rays; ventrals with orange spots; basal half of pectoral colored similarly to dorsal and anal, outer part plain; caudal plain olive with both pale and colored spots at base. In spirits the orange-brown spots become dull-brown.

This fish is widely distributed throughout the western Atlantic from southern Florida to Brazil and is also known from Ascension and St. Helena islands and Cape of Good Hope. At Key West it is very common about rocky elevations in moderately deep water, and is one of the most important food-fishes, reaching a length of 2 feet or more and a weight of 15 or 16 pounds, though the average weight of those brought to Key West does not exceed 2 or 3 pounds. It is caught with hook and line, will take any kind of bait, and exhibits good game qualities. In Porto Rico it is probably common, though we took only two specimens, 10 and 15 inches long, from Ponce and Isabel Segunda.

The rock-hind is one of the most beautiful of our tropical fishes, whether we consider the richness of its colors or the trimness of its form. It bears a strong general resemblance to the red-hind, but its spots are less thickly placed and are larger on some parts of the body.

Percu tota maculis Seba, Thesaurus, III, tab. 27.

Trachinus adscensionis Osbeck, Iter Chin., etc., 1757, English edition, 96, 1771, Ascension Island.

Trachinus punctatus Bonnaterre, Tableau Enyel. Method., 1788, 46; after Osbeck.

Percu stellio Walbaum, Artedi Piscium, 349, 1792; after Seba.

Percu maculata Bloch, Ichthyol., pl. 313, 1792, Martinique.

Trachinus osbeck Lacépède, Hist. Nat. Poiss., II, 364, 1800, Ascension Island; after Osbeck.

Sparus atlanticus Lacépède, l. c., IV, 158, pl. 5, fig. 1, 1803, Martinique; on a copy of a drawing by Plumier.

Serranus nigricaudus Cuvier & Valenciennes, Hist. Nat. Poiss., II, 375, 1828, Martinique.

Serranus pizanga Cuvier & Valenciennes, Hist. Nat. Poiss., II, 383, 1828, Brazil; after Maregrave.

Serranus aspersus Jenyns, Voyage of Beagle, Fishes, 6, 1842, Porto Praya, St. Jago, of the Cape Verde Islands.

Serranus impetiginosus Müller & Troschel, Schomburgk's Hist. Barb., 665, 1848, Barbados.

Serranus varius Bocourt, Ann. Sei. Nat. (5), X, 1868, 222, Gulf coast of Mexico.

Epinephelus punctatus, Poey, Fauna Puerto-Riqueña, 319, 1881; Stahl, l. c., 76 and 162, 1883.

Epinephelus adscensionis, Jordan & Evermann, l. c., 1152, 1896.

108. *Epinephelus striatus* (Bloch). "*Cherna*"; *Nassau Grouper; Hamlet; Cherna Criolla.*

(PLATE 12.)

Head 2.5; depth 3; eye 5.5; snout 4.2; maxillary 2.2; mandible 1.8; interorbital 7; preorbital 10; D. xi, 17; A. iii, 8; pectoral 1.8; ventral 2.1; caudal 1.8; scales about 110.

Body rather elongate, of moderate depth, dorsal outline regularly arched, back somewhat compressed; head pointed; mouth rather large, maxillary reaching posterior border of orbit, jaws nearly equal, or lower slightly projecting; teeth in jaws in bands with enlarged patches in front, a pair of canines in front of each jaw, lower pair the smaller; inner teeth of bands enlarged and depressible; preopercle with fine serrations, these enlarged near the angle; opercle ending in 3 flat spines, middle one strongest, upper one scarcely evident; a membranous opercular flap; gillrakers 8 + 15, counting rudiments; dorsal spines strong, fleshy prolongations of the membrane extending beyond their tips; soft dorsal higher than spinous portion; anal short, high, and rounded; caudal rounded; pectoral large, reaching tips of ventrals, middle rays longest; soft dorsal and anal little scaly; scales of body small, firm and strongly ctenoid.

Color in life: Pale-gray, paler below, side with five or six broad, brown, vertical bars, irregular in shape, the color darker above, extending upon dorsal fin; lower parts of these bars broken by pale blotches and cloudings; a band of same color from snout through eye to dorsal, where it joins its fellow; a narrow median band from snout backward, bifurcating opposite eye, the parallel branches extending to occiput and ceasing abruptly without joining the other band; a quadrate jet-black spot on upper surface of caudal peduncle; some obscure dark bars from eye downward and backward; numerous black points variously placed around and about eye; a narrow brown (in spirits) streak on edge of preorbital, overlapped by maxillary; lower part of side and under parts with whitish spots; dorsal yellow-edged; anal and caudal tipped with orange-yellow; ventrals blackish, tipped with faint yellow; pectorals light-orange or brownish, their bases dusky. The alcoholic specimens have the colors faded variously, some are nearly pale, some nearly uniform brown, and some retain the vertical bars; the square blotch on peduncle and the black points around eye persist in all cases and distinguish this well-marked species from its allies.

This excellent food-fish is found from the Florida Keys southward among the West Indies to Brazil; common about Porto Rico, numerous specimens obtained at San Antonio Bridge, Puerto Real, Ensenada del Boqueron, Ponce, and in the San Juan market; one from San Geronimo. A common and very important food-fish, reaching a weight of 50 pounds or more. The average weight of those brought to the Key West market is less than 10 pounds, while those seen in Porto Rico were still smaller. This grouper is taken with hook and line both about Porto Rico and Key West. It is said to be found at the latter place throughout the year.

Cherna, Parra, Dif. Piezas, Hist. Nat., 1787, 50, lam. 24, Havana.

Anthias striatus Bloch, Ichthyologia, IX, 109, 1792, Martinique.

Anthias cherna Bloch & Schneider, Syst. Ichth., 310, 1801, Cuba.

Sparus chrysomelanurus Lacépède, Hist. Nat. Poiss., IV, 160, 1803, Martinique.

Epinephelus striatus, Poey, Fauna Puerto-Riq., 319, 1881; Stahl, l. c., 76 and 162, 1883; Jordan & Evermann, l. c., 1157, 1896.

109. *Epinephelus guttatus* (Linnaeus). "*Mero Guajiro*"; *Cabrilla*; *Red-hind*.

(PLATE 13.)

Head 2.5; depth 3.2; eye 4.3; snout 4.3; maxillary 2.2; mandible 1.9; interorbital 7.8; preorbital 11; D. XI, 16; A. III, 8; pectoral 1.9; ventral 2.25; caudal 2; scales about 20-100 to 120-x.

Body elongate, somewhat compressed, not so heavy forward as in *E. adscensionis*; head pointed, mouth moderate, maxillary (which is finely scaled) reaching about to posterior border of eye, falling short in young; jaws subequal, lower barely projecting; canines of upper jaw the larger; interorbital space narrow, little more than half eye; preopercle finely serrate, more strongly at angle; a shallow emargination above angle; gillrakers not long, about 17 below angle, counting rudiments; scales ctenoid. Dorsal fin continuous, with a shallow emargination; caudal gently rounded.

Color: In life light yellowish-olive above, whitish or reddish below; three broad, oblique, obscure bands of olive running upward and backward on side; body everywhere with vivid scarlet spots, those above a little darker, those below sometimes with darker centers; some scarlet inside of mouth; soft parts of vertical fins olivaceous and reddish, broadly edged with black, especially dorsal; caudal palest; each with a very narrow white edge; pectoral light-yellow or reddish-orange, with scarlet spots; ventrals red, blackish at tips; inside of mouth red. Another specimen 10 inches long, from Arroyo, showed the following: Body dirty or yellowish-white, profusely covered with round brick-red spots about size of pupil, these darkest above and rosiest red below; these spots somewhat regularly in rows and covering head, lips, and lower jaw; inside of mouth red on sides; tongue white; dorsal olivaceous, with darker and paler areas, edge of membrane at tips of spines rich yellow, elsewhere black-edged; soft dorsal with broad black border narrowly edged with white; anal with six or eight red spots near base beyond which the fin is mottled, then dark like the soft dorsal; pectoral pale-red; ventral with a few pale-red spots, fin purplish-black at tip. In spirits, the oblique bands fade and the scarlet spots become brown.

This species resembles, at least in preserved specimens, *E. adscensionis*, from which it may be distinguished by the scales of maxillary and the broad black edgings of vertical fins. Its range is from the Carolinas southward through the Bahamas and West Indies to Brazil; it is known from Charleston, the Bahamas, Key West, the Bermudas, Havana, Jamaica, Porto Rico, and Martinique. It was obtained by us at Aguadilla, Puerto Real, Arroyo, Isabel Segunda, and Culebra, and by Mr. Gray at San Geronimo.

This is one of the smallest of the groupers, rarely attaining a greater length than 18 inches; it is, however, an important food-fish, and in the Havana market is one of the most abundant species. It

is also common at Key West. It is taken by the line fishermen at moderate depths near the coral reefs, and displays good game qualities. Anglers going to Key West or Porto Rico will find the red-hind one of the most beautiful and interesting of the game fishes.

Until recently this species was identified as *E. maculosus* (Cuvier & Valenciennes), but Dr. Günther, in a paper received since the colored plates of this report were printed, shows that the red-hind is evidently the *Perca guttata* of Linnaeus. The name on plate 13 should therefore be *Epinephelus guttatus* (Linnaeus).

Cugupuguacu brazil, the Hind, Catesby, Nat. Hist. Carolina, etc., pl. 11, 1743, Bahamas.

Cabrilla, Parra, Dif. Piezas, Hist. Nat. Cuba, 1787, Havana.

Perca guttata Linnaeus, Syst. Nat., ed. X, 292, 1758, Brazil?

Serranus catus Cuvier & Valenciennes, l. c., II, 373, 1828, Martinique.

Serranus arara Cuvier & Valenciennes, Hist. Nat. Poiss., II, 377, 1828, Havana.

? *Serranus angustifrons* Steindachner, Verh. Ges. Wien, XIV, 1864, 230, pl. VII, fig. 213, Cuba.

Epinephelus cubanus Poey, Repertorio, I, 202, 1867, Cuba.

Epinephelus maculosus, Jordan & Evermann, l. c., 1158, 1896, and 3197, 1900.

Epinephelus guttatus, Jordan & Evermann, l. c., 3197, 1900.

110. *Epinephelus morio* (Cuvier & Valenciennes).

"Cherna"; "Mero"; *Cherna Americana*; *Cherna de Yivera*; *Nigre*; *Red Grouper*; *Jaboncillo*.

(PLATE 14.)

Head 2.5; depth 2.7; eye 5.5; snout 4; maxillary 2.1; mandible 1.8; interorbital 7; preorbital 9; D. xi, 16; A. iii, 9; pectoral 2; ventral 2.1; caudal 1.7; scales 20-130 to 140-60.

Body deeper than in the other species of *Epinephelus*, and more compressed; head large, pointed, with a large mouth and projecting lower jaw; maxillary reaching past eye; 2 pairs of canines in front of each jaw, lower pair the smaller; preopercle finely serrate, teeth at the angle enlarged, a slight emargination above; dorsal fin with a moderate notch, the spines rather slender, but stiff and sharp, first less than half length of second, which is the longest, 2.5 in head; middle rays of soft dorsal highest; caudal lunate, upper rays produced slightly beyond lower; pectoral reaching beyond tip of ventral, latter not reaching vent.

Color in life: Olive-gray or olive-brown, clouded with pale-olive; lower part of head and breast usually salmon-color, with some red shades; irregular blotches of grayish-white over body; preorbital and adjacent regions with round points of dark orange-brown, becoming brown in spirits; inside of mouth posteriorly bright-orange; iris golden; vertical fins like body, soft parts with a broad edge of blue-black and narrow pale border; spinous dorsal black-edged; ventrals dusky; pectorals light-olive. In spirits, uniform light-brown with pale blotches. With age this species becomes more and more of a flesh red, especially below and on mouth, and the pale spots and blotches become less distinct.

An easily recognizable species, separated from all the others by the elevation of second dorsal spine. Found on the Atlantic coast of America from Virginia to Rio Janiero, and probably common about Porto Rico. Two fine specimens, each about a foot in length, obtained at Puerto Real and Isabel Segunda, where others were seen.

This is a very handsome fish, bearing some resemblance to the Nassau grouper, but the warm browns on the head and side are richer, while the general appearance is somewhat coarser. It is one of the largest and most important food-fishes of our tropical waters. It reaches a length of 2 to 3 feet and a weight of 20 to 25 pounds, or sometimes even 40 pounds. It is very abundant on the west coast of Florida in company with the red snapper. It is most abundant on the south Florida coast and is found throughout the year on the "grounds" at sea, and in the summer in some of the bays. It probably spawns in early spring in both places. Silas Stearns says the young are often seen in Pensacola Bay, where in June he obtained examples about an inch long. The red grouper is more of a bottom fish than the red snapper. It swims more slowly and seldom rises to the surface. It is very voracious, consuming enormous quantities of crustaceans and small fish. Large crabs in almost perfect condition are often found in their stomachs. On the "Snapper Banks" off the west coast of Florida it is caught by the red-snapper fishermen and in the same way, which is with hook and line, a piece of bone-fish or other fish being used as bait. It does not rank high as a game fish, its movements being slow, and when hooked it is hauled up almost as a dead weight. It will take almost any kind of bait. When red snappers were more abundant the red grouper did not find a ready sale in the Northern markets, though it has always been in good demand at Key West and Havana. The maximum weight of those taken at Key West now probably does not exceed 25 pounds and the average is only 8 to 15 pounds.

The red grouper is very tenacious of life, and will live several hours after being taken from the water, even though exposed to considerable heat. This is doubtless one reason why the Key West fishing fleet has preferred groupers for transportation to Cuba, since they are obliged to go a long way to market and through warm water, and the grouper bears the crowding and chafing in the wells of the smacks better than other species.

On the Florida coast this fish is known as the red grouper, or grouper, while in Cuba and Porto Rico it, together with one or two other groupers, is called *cherna*, *cherna de vivera*, or *jaboncillo*.

Serranus morio Cuvier & Valenciennes, Hist. Nat. Poiss., II, 285, 1828, New York and Santo Domingo.

Serranus crythrogaster De Kay, New York Fauna: Fishes, 21, pl. 19, 1842, Florida.

Serranus remotus Poey, Memorias, II, 140, 1860, Havana.

Epinephelus morio, Poey, Fauna Puerto-Riqueña, 319, 1881; Stahl, l. c., 76 and 162, 1883; Jordan & Evermann, l. c., 1160, 1896.

Genus 69. ALPHESTES Bloch & Schneider.

The genus *Alphestes* contains two species of small fishes which differ from *Epinephelus* proper in the presence of a strong antrorse spine on lower limb of preopercle. Frontal bones with an anterior excavation for reception of posterior processes of premaxillaries, a process or knob on each side of skull behind interorbital area; supraoccipital and parietal crests produced on frontals, but not extending to between orbits. Dorsal rays xi, 17 to 20; A. iii, 9. Only one of the two species known in Porto Rico.




FIG. 14.—*Alphestes chloropterus*.

111. *Alphestes chloropterus* (Cuvier & Valenciennes). "*Cherna*"; *Guaseta*.

Head 2.6; depth 2.7; eye 4.6; snout 6; maxillary 2.4; mandible 2; interorbital 6.8; D. xi, 18; A. iii, 9; pectoral 1.7; ventral 2; caudal 1.6; scales 12-75 to 85-32.

Body ovate and compressed, caudal peduncle short; head small and pointed, anterior profile depressed in occipital region; eye large, its diameter greater than length of short snout; mouth oblique, maxillary reaching posterior border of orbit or beyond; teeth conical and sharp, some of them depressible, in patches on front of jaws, in more than one row on sides; 2 pairs of weak canines on front of upper jaw; lower jaw slightly projecting; preopercle strongly convex, angle rounded, and with a strong, rather flat spine, pointing downward and curved slightly forward; above this are fine serrations along entire upper limb, decreasing in size upward; opercle with 3 flat spines, middle one strongest, upper and lower nearly concealed by scales; scales covering head and body, reduced in size at nape, occiput, and everywhere on head save opercles, where they are rather larger than on body; scales all cycloid save a distinctly ctenoid patch under or above pectoral; gillrakers short, about 15 below angle; dorsal fin continuous, with a shallow notch, spines strong and pungent, fourth and fifth highest, 2.5 in head; soft dorsal as high as spinous, about the fourteenth ray longest, 2.25 in head, fin pointed posteriorly, rounded in young; anal high, second spine strongest and about equal to third, 2.7 in head; caudal rounded; pectoral and ventral subequal, not reaching vent; in a small individual (4 inches) the pectoral reaches past ventral, which just reaches vent.

Color in life: Body yellowish-brown, paler below, upper part of side with about 7 longitudinal stripes of dark brown from head to tail, these becoming rows of round orange spots below; 6 dark, inconspicuous, vertical body bars; head with many smaller orange-brown spots; lower part of head and breast with pale-bluish spots; fins brownish; soft dorsal with oblique bars of black; spinous dorsal olive, blotched with brown; ventral olive, edged with darker; other fins obscurely barred; inside of mouth pale.

Found in the West Indies, Cuba to Brazil; also recorded from Africa and the Falkland Islands; not known from Florida; apparently not uncommon about Porto Rico. Two females, 7.5 and 9.5 inches long, respectively, and each with well-developed roe, were obtained in the San Juan market January 17. Another specimen, 4 inches long, was gotten at Mayaguez. The species reaches a length of a foot or more and is a food-fish of importance. The only local name heard for it was "cherna."

Plectropoma chloropteron Cuvier & Valenciennes, Hist. Nat. Poiss., II, 398, 1828, Santo Domingo and Martinique.

Plectropoma monacanthus Müller & Troschel, in Schomburgk's Hist. Barbados, 665, 1847, Barbados.

Alphestes afer, Jordan & Evermann, l. c., 1164, 1896.

Genus 70. MYCTEROPERCA Gill.

Cranium broad and transversely concave between eyes, its lateral crests very strong, nearly parallel with supraoccipital crest and extending much farther forward than latter, joining supraocular crest above eye, supraoccipital crest not extending on frontals; frontal bones without anterior concavity or notch for reception of premaxillaries, without processes on the upper surface; lower jaw strongly projecting; anal fin elongate, with 11 or 12 (in one species 9 or 10) soft rays; caudal lunate; spines of fins slender, none of them much elevated; scales small, mostly cycloid, those on lateral line simple; pyloric caeca few; gillrakers various; nostrils small, and subequal or with posterior enlarged. Otherwise essentially as in *Epinephelus*, from which genus *Mycteroperca* is well separated by the structure of the skull, and superficially by longer anal, larger mouth, and more elongate body.

Large, handsome fishes of the Tropics, mostly American, including some of the most valuable food-fishes in the Gulf of California and at least six of the valuable species of the Florida coast. Of the 20 or more American species only 2 are thus far known from Porto Rico. All are excellent for food, and those found where fisheries have become established are much sought.

- a. Nostrils subequal, well separated; scales on head cycloid.
- b. Gillrakers comparatively few and short, 8 to 20 below angle of arch.

TRISOTROPIS:

- c. Anal rays III, 11 or III, 12, the fin long.
- d. Anal fin not angulated, its outline more or less evenly rounded in adult as well as in young; soft parts of vertical fins edged with black in life.
- e. Angle of preopercle not salient, its teeth scarcely enlarged; gillrakers $x + 8$ to 10.
- f. Gillrakers very few and short, $x + 8$ developed (besides some rudiments); general color pale, bright red or grayish, with roundish spots or blotches of black or red darker than the ground-color; the blacker blotches along middle of sides, much larger and quadrate in young; red always present somewhere in life (fading in spirits); pectorals blackish, in the adult broadly tipped with orange-yellow; scales rather small (about 125); caudal lunate.
- g. General color gray, with red and black markings..... *venenosa*
- gg. General color scarlet, with red and black markings..... *apua*
- ff. Gillrakers rather slender, about $x + 10$ (besides several rudiments); caudal subtruncate; nostrils small.
- h. Scales not very small (about 110); color dark olive-green; sides of head and body with rivulations of dark-bluish around roundish dark-bronze spots, large or small (these markings subject to considerable variation, fading in spirits); sides with darker quadrate areas.
- i. Dark blotches on body rather large, often quadrate..... *bonaci*, 112
- ii. Dark spots on body very small, close-set, of a deep-bronze orange..... *xanthosticta*
- cc. Angle of preopercle more or less salient, its teeth somewhat enlarged, gillrakers more numerous, $x + 12$ to 14.
- j. Scales not very small (about 110); upper part of body dark-brown, lower half abruptly paler; a pale ring around caudal peduncle, behind which is a squarish dark blotch, smaller than eye, at base of upper rays of caudal; caudal deeply lunate; teeth strong..... *dimidiata*

PAREPINEPHELUS:

- bb. Gillrakers close-set, very long and slender, 25 to 35 below angle of arch.
- k. Caudal fin lunate, its angles more or less produced in adult, fin subtruncate in young; anal fin more or less angulate in adult, rounded in young; soft dorsal somewhat angular; scales rather large (lateral line 95); body rather deep, snout sharp; preopercle with a salient angle which is armed with larger teeth; dorsal spines low; gillrakers close-set, $x + 30$, longest 7.5 in head; ventrals not reaching to vent; color olive-gray, with darker reticulations around pale spots; fins not much darker on their edges; a dark mustache along maxillary; adult examples nearly uniform brown; not known to be red..... *rubra*
- aa. Nostrils very close together, posterior decidedly larger than anterior, and with a more or less distinct horizontal cross-septum within; scales on head cycloid.

MYCTEROPERCA:

- l. Gillrakers moderate, fewer in number, 6 to 18 below angle of arch.
 m. Second dorsal spine low, shorter than third, the third and fourth highest.
 n. Margin of anal fin posteriorly concave, its middle rays much exserted.
 o. Gillrakers rather numerous, 17 to 20 below angle of arch.
 p. Outer rays of caudal much produced, more than two-thirds length of head; preopercle with salient angle; canine teeth strong; scales small (140); gillrakers about 4 + 20. Color brownish, with small darker spots; vertical fins broadly edged with blackish.
 q. Upper canines directed strongly forward, lower backward; coloration obscure..... *falcata*
 qq. Upper canines nearly vertical; coloration paler and brighter..... *phenax*
 mm. Margin of anal fin not concave posteriorly, outline of fin rounded or slightly angular.
 r. Gillrakers rather few, x + 12; body without dark crossbars, covered with grayish reticulations around small round spots, these not evident on head; anal fin rounded; preopercle with a salient angle; form rather robust; anal fin not angulate.
 s. Scales very small, about 20-140-37; caudal not deeply lunate; eye small..... *bowersi*, 113
 ss. Scales larger, about 90 to 100 in lateral line; caudal deeply lunate; eye larger..... *calliura*
 rr. Gillrakers very few, short, and thick, about x + 6; body olive or (var. *camelopardalis*) bright red, with light and dark crossbars, these often becoming obsolete with age; head usually with distinct reticulations around yellowish spots; anal fin with angular margin, subtruncate posteriorly; preopercle without salient angle; scales rather small (lateral line 133); form rather robust.
 t. Ground-color dark-olive..... *tigris*
 tt. Ground-color bright-red..... *camelopardalis*

112. *Mycteroperca bonaci* (Poey). *Aguaji*; *Bonaci Acara*; *Black Grouper*.

Head 2.75; depth 3.25; eye moderate, 6 in head (young); D. XI, 16 to 18; A. III, 11 or 12; scales 18-120 to 125-50, pores 70 to 85. Body comparatively slender, a little more robust than in *M. microlepis*, its breadth 2.33 in its depth; head moderate, rather pointed, its anterior profile little curved; mouth rather large, maxillary reaching slightly beyond eye, 2.2 in head (in young), proportionately longer in adult; maxillary with cycloid scales. Teeth in rather narrow bands; two rather strong canines directed little forward in front of each jaw. Interorbital space slightly convex, its width 6 in head. Preopercle forming a regular curve without salient angle, emargination near angle very slight. Nostrils small, roundish, subequal; not very close together. Gillrakers few and long, 10 to 12, besides rudiments on lower part of anterior arch. Scales rather small, chiefly cycloid; dorsal spines comparatively slender and weak, outline of fin gently convex; tenth spine about as long as second; third and fourth spines longest, 3.33 in head; caudal fin truncate when spread open, its outer rays a very little produced, 1.6 in head; anal rather high and rounded, its longest rays 2.25 in head; pectoral reaching slightly beyond tips of ventrals, 1.7 in head. Ventrals short, not reaching vent. Pyloric caeca 15.

Color in life: Deep orange-brown, more olive on back, clouded above by paler or grayish; sides and belly marked everywhere by reticulations of pearly gray, which surround roundish or oblong spots of ground-color, pale streaks being largely horizontal on sides; sides of head similarly marked, the spots smaller, bronze-brown, reticulations decidedly bluish; 6 or 7 spots in a straight line between eye and preopercle, having nearly the diameter of the pupil; spots on body mostly covering 4 to 6 scales, all of them larger than a scale; dorsal olive-brown, somewhat mottled; caudal similar to dorsal, narrowly edged with whitish; anal similar with 2 or 3 rows of bluish spots, its tip blackish, with a narrow whitish edge; pectorals olive-brown, plain; ventrals blackish, rays bluish. Mouth not green, lips olive, barred with bluish; iris reddish. Here described from a specimen 11.5 inches long, from Key West. A large specimen, about 2.5 feet in length, seen at Key West, retained the same general coloration, the bronze spots and rivulations being distinct and not smaller than in young.

In spirits the orange-brown of the body is replaced by dark-brown, and the blue reticulations of the head by gray; all the markings become more faint.

This fish ranges from Florida to Brazil. It is common at Key West, where it is called black grouper. Probably not uncommon about Porto Rico, though seen by us only at Puerto Real. It is one of the important food-fishes of Key West and is probably held in equal esteem in Porto Rico. It reaches a length of 2 to 3 feet and a weight of 50 pounds. The larger individuals are caught with the hook, but the young are often seined along the shore.

- Serranus bonaci* Poey, Memorias, II, 129, 1860, Cuba.
Serranus brunneus Poey, Memorias, II, 131, 1860, Havana.
Serranus decimatis Poey, Memorias, II, 138, 1860, Cuba.
Serranus cyclopinatus Poey, Memorias, II, 353, 1861, Cuba.
Serranus latepictus Poey, Memorias, II, 353, 1861, Cuba.
Trisotropis aguaji Poey, Repertorio, 229, 1868, Havana.
Mycteroperca bonaci, Jordan & Evermann, l. c., 1174, 1896.

113. *Mycteroperca bowersi* Evermann & Marsh, new species.

“Rock-hind”; “Rock-fish”; “Mero Cabrilla.”

Head 2.8 measured from tip of upper jaw; depth 3.4; eye 7.5; snout 3.7; maxillary 2.1; mandible 1.7; interorbital 4.6; preorbital 8.4; scales about 20-140-37; D. xi, 16; A. iii, 11; P. 17; gillrakers x+10.

Body rather long, compressed, the dorsal and ventral outlines each gently and regularly arched; head long and pointed, the greatest width 2.25 in its length; snout long; mouth large, lower jaw strongly projecting and entering in profile of snout; maxillary reaching far beyond the orbit; supplemental maxillary bone evident; eye small, high up; preopercle very finely serrate, slightly concave at its angle; opercle ending in a long flat point upon base of which is a broad, flat spine; nostrils close together, posterior much the larger, with an anterior horizontal cross-partition at base. Fins rather low; origin of dorsal over end of opercular flap; first dorsal spine short, equal to diameter of orbit; second spine somewhat exceeding twice length of first; third, fourth, and fifth spines longest, subequal, about a third longer than second, or 3.35 in head; soft dorsal rounded, its longest rays about 3 in head; caudal slightly lunate, outer rays 1.7 in head; anal rounded, spines slender and weak, longest (third) 2.6 in head, longest ray 2.6; pectoral rounded, middle rays longest, 2.2 in head; ventral 2.4. Scales small, thin, and cycloid, those on head and nape very small and embedded.

Color in life: Body dark reddish-brown, covered with many small, round, blood-red spots, these also on head, lower jaw, and base of pectoral and anal, especially numerous on anal, and a few on spinous dorsal; soft dorsal mottled with white and black, bordered with a very narrow white edge inside of which is a broad black band; tip of caudal narrowly white, then a broad black band, rest of fin mottled with white and black; anal similar to soft dorsal, but with more red spots; pectoral crossed by two broad dark bars, outer end of fin yellow; inside of mouth pale red. In alcohol the general color becomes dark-grayish, paler below; red spots changing to black or dark-brown.




FIG. 45.—*Mycteroperca bowersi*.

Judging from descriptions this species appears to be related to *M. callura* Poey, but differs in a number of important particulars, among which are the smaller scales, smaller eye, and very different coloration. According to Poey, *M. callura*, in life, was dark brownish-olive; with rounded yellowish spots; fins dark-brown, darker on edges of vertical fins; eight narrow, dusky crossbands on body; caudal with a beautiful green crossband; pectoral yellowish toward the center, the posterior margin green.

Only a single specimen (type No. 49530, U.S.N.M.) of this important species was obtained. This is 21.5 inches long and was purchased from a Tortola fisherman at Culebra Island, February 10. It was called by him “rock-hind,” “rock-fish,” or “mero cabrilla,” and was caught, along with the tangs, angel-fish, and other species, in a fish-trap of the style in common use about Culebra Island. Though no others were seen, the species seems to be well known to the fishermen about Culebra and Vieques islands, and is held in high esteem as a valuable food-fish.

We name this species for the Hon. George M. Bowers, U. S. Commissioner of Fish and Fisheries.

Genus 71. **DIPLECTRUM** Holbrook. The Squirrel-fishes.

This genus is very close to *Prionodes*, from which it differs chiefly in the armature of preopercle, which is provided in adult with one or two clusters of strong, straight, divergent spines. Smooth area on top of head, as in *Serranus*, large, extending backward to a line connecting borders of preopercle; supraoccipital and parietal crests very short; preorbital broad; maxillary widest before its tip; profile of snout rounded; pectoral unsymmetrically rounded, its upper rays longest; ventrals inserted somewhat before axil of pectoral; dorsal spines slender, none of them much elevated; soft dorsal short, rays x, 12; anal rays m, 7; caudal lunate.

Species of small size and bright colors, all American; only one species known from Porto Rico, but *D. formosum* also doubtless occurs there.

HALIPERCA:

- a. Preopercle with a single center of divergence of the spinules about its angle (in adult as well as in young).
 b. Gillrakers moderate, about 10 below angle of arch; spines on produced portion of preopercle numerous, 8 to 20 in number; outline of spinous dorsal fin somewhat convex, so that the fin is more deeply notched than in *Diplectrum formosum*; jaws equal; vertex naked; opercle black within.
 c. Scales on cheek small and regularly placed, in about 10 rows; 22 scales before dorsal..... *radiale*, 114

DIPLECTRUM:

- aa. Preopercle with 2 clusters of divergent spines, one at angle, other higher (the 2 fascicles well separated in adult, but smaller and coalescent in young).
 d. Head and body marked with many interrupted blue lines; preorbital broad, more than twice width of maxillary; lower half of preopercle with strong, straight spines diverging from two centers; gillrakers short and small, x+14; 11 rows of scales on cheek; caudal deeply lunate, upper lobe the longer, sometimes ending in a long filament. Color brownish, silvery below; side with 7 or 8 longitudinal deep-blue lines and about as many dark crossbars, last bar forming a large black blotch at upper base of caudal; young with 2 broad, dusky longitudinal stripes, which become interrupted with age; 3 or 4 distinct blue stripes on sides of top of head; 2 across preorbital, the lower forked; fins with narrow, wavy bars of blue and pale yellow..... *formosum*

114. **Diplectrum radiale** (Quoy & Gaimard). *Aguavina*.

Head 2.8; depth 3.8; eye 3.75; snout 3.8; maxillary 2.1; mandible 1.9; interorbital 4.6; preorbital 9; D. x, 12; A. m, 7; pectoral 1.5; ventral 1.7; caudal 1.35; scales 7-10-18.

Body elongate, dorsal and ventral outlines nearly alike save for a depression at nape; head pointed, a large, smooth round area above, behind eyes; lower jaw barely projecting; teeth chiefly in one row on sides of jaws, in small patches in front; maxillary reaching posterior edge of pupil; eye large, placed high in head; about 9 rows of regularly placed scales on cheek; preopercle with a strongly salient angle composed of radiating spines; opercle with 7 or 8 rows of scales. Dorsal fin continuous, slightly notched, spines slender and very sharp; caudal lunate, upper lobe very slightly produced. Scales strongly ctenoid.

Color in spirits: Olive, paler below; many faint vertical dark bars wider than interspaces, plainest at middle of sides; 2 faint longitudinal dark bands, first from shoulder to soft dorsal, other from nearly same origin to base of caudal, where it forms a faint dark spot; some faint, pale streaks on preorbital and cheek. In life, the soft dorsal has blue spots encircled by darker blue rings; caudal barred with similar spots; body light-brown above, yellowish below, sides salmon-color, head and fins with red shadings.

Found on both coasts of tropical America, north to Havana and Guaymas; very common on the coast of Brazil and in the Gulf of California, usually in shallow bays; apparently not common in Porto Rico, where one young individual, 5 inches in length, was collected at Mayaguez.

Serranus radialis Quoy & Gaimard, Voyage Uranie, 316, 1824, Rio Janeiro.

Serranus bivittatus Cuvier & Valenciennes, Hist. Nat. Poiss., II, 241, 1828, Martinique.

Centropristes ayresi Steindachner, Ichth. Notizen, VII, 1, pl. 1, fig. 1, 1868, Santos, Brazil.

Diplectrum radiale, Jordan & Evermann, l. c., 1204, 1896.

Genus 72. **PRIONODES** Jenyns. **Serranos**.

Body oblong, moderately compressed, covered with moderate-sized ctenoid scales. Lateral line normal in direction, not running close to back. Cranium above with a very large convex smooth area, which is longer than the low supraorbital crest; supraoccipital and parietal crests short, extending to a line connecting borders of preopercle; posterior outline of cranium nearly vertical in profile; mouth large, maxillary not scaly and without supplemental bone; canines small, lateral; no depressible teeth in jaws; teeth always present on vomer and palatines. Gillrakers usually few. Branchiostegals 7. Dorsal with 10 rather slender spines either subequal or one of them much produced, the fin not

deeply notched, the soft portion short, of 11 to 13 rays, and nearly or quite destitute of scales; anal short, with slender spines; caudal lunate or truncate. Ventrals not very close together, inserted somewhat in advance of pectorals, as in *Centropristes* and *Diplectrum*. Vertebrae $10 + 14 = 24$.

Species of small size, probably all American, closely allied to the Old World genus *Serranus*, from which they are distinguished by the short, naked, soft dorsal, the anterior insertion of ventrals, and the smaller teeth. Of the 11 known American species only one has been found in Porto Rico. At least two other species (*P. phabe* and *P. tabacarius*) probably occur there.

PRIONODES:

- a. Scales large, 42 to 55 in the lateral line.
 b. Scales on cheek very large, in about 5 series; dorsal spines not quite equal, fourth longest, about twice length of ninth.
 c. Teeth unusually strong, 3 or 4 on each side in upper jaw as large as largest lateral teeth; jaws equal; preorbital extremely narrow, not one-third width of pupil; preopercle very sharply serrate; gillrakers short, slender, $x + 10$; caudal forked. Color plain-olivaceous; fins all pale; sides with about 6 faint dark crossbands, more or less confluent along lateral line *fuscus*
 bb. Scales on cheek moderate, in about 8 series; body oblong, heavy anteriorly; dorsal outline a little elevated, profile almost straight; lower jaw not projecting; canines small; mouth large; preorbital narrow; dorsal fin not notched.
 d. Eye large, equal to snout; gillrakers short, $x + 10$; caudal deeply lunate; pectoral long. Color light-brownish, with lighter blotches and faint dusky bars; a very conspicuous, sharply defined, vertical white bar extending upward on each side from just before vent; fins pale *phabe*
 dd. Eye very large, greater than snout; gillrakers short, $x + 6$; caudal truncate. Color, upper parts scarlet, lower parts bluish-white; a yellow lateral band, under which are 4 quadrate black blotches. *baldwini*, 115

MENTIPERCA:

- aa. Scales small, lateral line with 60 to 75 in its course.
 c. Lower jaw not very strongly projecting.
 f. Body covered with irregular, inky-black spots and bands. Body long and low, head low and sharp, lower jaw projecting; scales on cheek in 10 or 12 rows; teeth small; gillrakers very short, $x + 7$; dorsal low, not notched; caudal forked. Color brownish above, sides yellowish, everywhere above, below, and on fins covered with irregular inky-black spots, blotches, and bands, latter meeting around belly; pectorals and anal plain; a broad ring around base of caudal, and many irregular spots around bases of ventrals and pectorals; numerous black spots on dorsals and caudal, one of those on front of spinous dorsal very conspicuous. *tigrinus*
 ff. Body not covered with conspicuous inky-black spots and bands.
 g. Back with 3 or more large, conspicuous blotches of yellowish-white. Body more elongate than in related species; lower jaw slightly projecting; scales on cheek in 11 series; dorsal spines low, fifth the longest, 2.25 in head; caudal strongly lunate; pectoral and ventrals short. Color brownish-red above, with areas of light yellow on sides of back; one before dorsal; a large one and a small one below spinous dorsal; a large one below last rays of soft dorsal; one on back of tail; top of head with 2 pale cross-shades, one before, one behind eye; lower fins light-orange; caudal red, with 2 conspicuous longitudinal stripes of blackish-red; dorsal red-shaded, a maroon blotch on each part of it extending upward from a similar blotch on back *tabacarius*
 gg. Back without conspicuous blotches of yellowish-white.
 h. Coloration nearly uniform; body elongate; snout short and thick; caudal slightly lunate. Color brownish-yellow on back, orange on sides, and brighter or red on belly; no spots nor bands; upper part of head bluish, fins gray; caudal bordered above and below with brown *flavescens*
 cc. Lower jaw very strongly projecting; body elongate, moderately compressed; scales small, about 70; snout sharp, much longer than the large eye; preopercle finely denticulated; top of head with vertex naked; caudal deeply forked; skull depressed, with a single crest; dorsal spines moderate, third highest.
 i. Color clear brown with larger darker spots or bars on sides; fins pale, more or less tinged with orange. *luciopercaus*

115. *Prionodes baldwini* Evermann & Marsh.

(PLATE 15.)

Head 2.5; depth 3.2; eye 4; snout 4.6; maxillary 2.4; mandible 2; interorbital 7; D. x, 12; A. III, 7; pectoral 1.4; ventral 1.3; caudal 1.7; scales 4-12-12.

Body elongate, moderately compressed, not elevated, covered with ctenoid scales; dorsal and ventral outlines alike; head moderate, pointed, naked above and below; eye large, greater than length of snout, high in position; mouth terminal, slightly oblique, maxillary reaching middle of eye or somewhat beyond; gillrakers short, 6 developed on lower limb; teeth small, conical, and sharp, on vomer and palatines and in several series in each jaw, with weak canines in front and a few canine-like teeth on middle of side of lower jaw; cheek with about 7 rows of scales; preopercle finely serrate; opercle ending in 3 sharp, flat spines, middle one largest, a membranous pointed flap projecting beyond; fins all naked, dorsal continuous, with a slight emargination; spines slender and pungent, first 4 or 5 graduated, rest subequal, 3.2 in head, lower than soft rays which are contained about 2.5 in head; anal fin short, second spine longest and strongest, 3 in head, soft part high, fifth or sixth ray longest,

reaching almost to front of anal, 2 in head; ventral with second ray produced, reaching vent; caudal truncate, or with middle rays very slightly shorter, making the margin slightly concave.

Color in life: Dorsal half of head and trunk and all of caudal peduncle scarlet, ventral portion pale-blue, almost white; a yellow longitudinal band, nearly as wide as pupil, from preopercular margin straight across opercle and along body to lateral line under last dorsal rays; 4 quadrate or oblong black blotches just under this band, the first about under middle of spinous dorsal, second under last spines, third under first rays, fourth under last rays; from each of the first three of these blotches a square, well-defined yellow shade extends downward to belly or base of anal, a similar one from base of pectoral to ventral; 4 smaller black blotches at base of caudal, 2 others, somewhat larger than the last, just in front of them on caudal peduncle; a row of 9 round black dots on each side at base of dorsal fin, first one smallest, opposite membrane of first spine, the other 8 separated somewhat obscurely into pairs, the first pair under middle spines, second under last spines, third under first rays, fourth under last rays; 2 or 3 very small black dots on upper edge of caudal peduncle; 2 or 3 more in front of dorsal on median line, each accompanied by a similar one on either side; in some specimens a few scattering ones on top of head behind eyes, sometimes regularly arranged; a few dark-brown spots behind eye; various dark markings on side of head, without very definite pattern, but usually 2 oblique stripes on cheek, a heavy black blotch on interopercle and 2 on ramus of mandible, which, with their fellows of the other side, make distinct crossbars on lower side of head, usually extending across maxillary; chin and lower part of opercle with dark spots; lateral line white, with a few broken spots, comparatively faint, just below it; iris red, with an inner ring of white surrounding pupil; spinous dorsal pale, edge of membrane black, this color bordered below with faint yellow; soft dorsal pale, spotted throughout with light-orange, with a marginal band of same, outside of which is a very narrow pale-blue edge; ventral very pale-blue, produced ray somewhat yellow; anal pale-blue with some light orange on last rays; pectoral and caudal uniform pale-reddish, unmarked.

In spirits, all the red and yellow markings disappear, the dark persists, and additional markings are brought out as follows: Along anterior and upper part of the trunk and crossing the lateral line are dark-brown vertical bars, diffuse and running together, or separated and broken into round or quadrate blotches; in the middle part of the course of yellow longitudinal band appears a row of very small black points; spots on soft dorsal dusky; dark mottlings on caudal; upper and lower base of pectoral, and sometimes axil, dusky.

A beautiful and strongly marked species; 2 specimens dredged and 33 others, ranging in size from 0.55 to 2 inches, caught in the tangle, off Culebra and Vieques islands, on coral bottom, in depths of 15 and 16 fathoms. Known only from Porto Rico.

This interesting species was "named for Mr. Albertus H. Baldwin, the artist of the expedition, in recognition of his excellent drawings and paintings of American fishes."

Prionodes baldwini Evermann & Marsh, Report U. S. F. C. for 1899 (December 19), 353, Fish Hawk station 6093, off Culebra Island, 5.25 miles southwest of Culebritas Light-House, February 8, 1899, in 15 fathoms.

Genus 73. DULES Cuvier.

This genus is close to *Prionodes*, differing in having but 6 branchiostegals and in the truncate form of the caudal fin. In *D. auriga* the third dorsal spine is prolonged in a whip-like spine. There are 3 known species of the genus, the one described below and *D. subligarius*, which occurs on the South Atlantic coast of the United States, and *D. auriga*, from the coasts of Brazil and Uruguay.

- a. Third dorsal spine not longer than fourth, 3 in head; second anal spine considerably longer than third, about 2.25 in head; lower jaw little projecting; gillrakers short and few, 6 or 8 in number; jaws scaleless; soft dorsal with small scales; pectoral long, reaching anal. Color brown, with darker cross-shades; soft dorsal, anal, and caudal fins checkered with blackish on a white ground; a broad white area or bar before anal fin.
- b. Dorsal rays X, 13; head small, acuminate; pectoral fin finely barred with black and whitish, precisely like caudal fin; a very conspicuous inky-black blotch on front of soft dorsal (at least in the young), this being a continuation of one of the bars on body; a black ring about tail at base of caudal, before which are 6 or 7 dark bars, becoming progressively broader and fainter forward; lower parts of head with a conspicuous network of dark streaks..... *subligarius*
- bb. Dorsal rays X, 12; head less slender; pectoral red; inky blotch on soft dorsal small or obsolete; dusky bars on body distinct..... *dispilurus*, 116
- aa. Third dorsal spine in adult male greatly elevated, reaching past middle of soft dorsal, its length quite variable; second anal spine as long as third, 2.83 in head; lower jaw prominent; gillrakers 10 to 12 below arch; pectoral shorter than head; brownish lower parts with light and dark shades; fins clouded..... *auriga*

116. *Dules dispilurus* (Günther).

Head 2.4; depth 2.75; eye 4.2; snout 4.2; maxillary 2.3; mandible 1.9; interorbital 7; D. x, 12; A. iii, 7; pectoral 1.35; ventral 1.6; caudal 2; scales 5-44-12.

Body oblong, of moderate depth, considerably compressed; mouth large, slightly oblique, maxillary reaching center of large eye or beyond; teeth in villiform bands in each jaw, in lower becoming uniserial posteriorly; a few enlarged teeth in outer row of both jaws; vomer and palatines with narrow bands of villiform teeth, some of them enlarged; jaws subequal; preopercle evenly rounded, finely serrate above; opercle with 3 pungent spines, middle one longest; scales rather large, ctenoid; caudal fin truncate; second anal spine longest and more or less curved.

Color in spirits: Brownish, some specimens more olive; 5 or 6 crossbands of darker color on body, not very distinct, these extending on dorsal fin, where they are quite black and sharply defined—1 on front of spinous dorsal, 1 on last spines, and 1 on first dorsal rays at height of last dorsal spines, the bar from this one extending to about lateral line only; another at base of last dorsal rays. Between these inky-black blotches are much smaller spots not parts of the body bars; caudal peduncle with 1 or 2 vertical bars; an oval black spot, smaller than pupil, at base of upper caudal rays, and 1 at base of lower, with fainter markings between; caudal and anal mottled, ventrals dark, pectoral pale (red in life); many (about 15) longitudinal brownish or olive lines on body, formed of color on upper and lower edges of scales, rather wider than interspaces, and giving the fish its general color; a faint, wider, dark streak from occiput to upper edge of eye, and 1 from shoulder through eye to tip of snout. A creamy quadrate blotch from belly, in front of vent, extending upward about halfway to lateral line, its posterior border more sharply defined than anterior.

A strongly marked species heretofore known only from Trinidad and Jamaica. Five examples, one about 1.75 inches, the others 2.75. The smaller one was taken with the tangle, off Point Melones, in 7 fathoms; one of the others in the beam trawl, 9 miles from Mayaguez, in 220 fathoms, on rocky bottom; the others were taken at Mayaguez.

Centropristis dispilurus Günther, Proc. Zool. Soc. Lond., 1867, 99, Trinidad.

Dules dispilurus, Jordan & Evermann, l. c., 1219, 1896.

Genus 74. RYPTICUS Cuvier. The Soap-fishes.

Body oblong, compressed, covered with very small, smooth, embedded scales. Lateral line normal, head scaly. Mouth rather large, oblique, lower jaw the longer; maxillary with a supplemental bone, as in *Epinephelus*, with which this genus agrees in general osteology; smooth area on top of cranium very large, transversely convex, much longer than supraoccipital crest; interorbital area very narrow; parietal and supraoccipital bones short, with feeble crests, which do not extend on frontals; premaxillaries reaching frontals, which have a fossa in front; teeth all villiform, in bands on jaws, vomer, and palatines; preopercle crescent-shaped, without angle or serratures, but provided with 2 or 3 spinous hooks on the posterior margin; opercle with 2 or 3 spines; gillrakers short; branchiostegals 7. Dorsal fins separate, the first of 2 or 3 (rarely 4) small spines, second of many (about 25) soft rays; anal long, rounded, of soft rays only; caudal rounded; pectoral rounded, nearly symmetrical, of 17 rays; ventrals small, 1, 5, inserted slightly before pectorals, the spine short and strong. Vertebrae 10 + 14 = 24. Skeleton generally similar to that of *Epinephelus*.

Species about 8, all from the seas of tropical America.

RYPTICUS:

- a. Dorsal spines 2 or 3 (rarely 4).
- b. Preopercle with two spines only, lower scarcely the longer.
- c. Opercular spines 3, all well developed.
- d. Color not red, chiefly olivaceous; dorsal fins distinctly connected by membrane.
- e. Eye not longer than snout; pores in lateral line 85 to 90. *saponaceus*, 117
- ee. Eye longer than snout; pores in lateral line 67; brownish, with blackish spots and dots. *arcnatus*
- cc. Opercular spines 2, small, uppermost the smaller. *coriaceous*, 118

PROMICROPTERUS:

- bb. Preopercle with 3 spines *bistrispinus*, 119

117. *Rypticus saponaceus* (Bloch & Schneider). *Soap-fish*.

Head 3 to 3.33 in length; depth 2.6 to 3.25; D. III, 23 to 25; A. 16 or 17; scales 85 to 90 (pores). Body comparatively deep, young more slender; back elevated, snout rather pointed; lower jaw much projecting; anterior profile before eye little concave; eye 4.5 to 5 in head; maxillary reaching posterior edge of eye, 2.12 in head; preopercle with 2 straight spines behind; opercle with 3 spines, middle one largest and nearer upper than lower; first and second dorsal spines subequal, third smallest; dorsals slightly connected, ventrals very small, not half longer than eye; pectoral rounded; gillrakers very small and short, about 8 developed. Color dusky-brown, fins marked with blackish and usually with a narrow pale edge; sides generally with irregular pale spots; back and head usually immaculate.

West Indies, Pensacola to West Africa and Brazil; generally common. The best known and most widely distributed of the soap-fishes. Though not obtained by us in Porto Rico, we include it on the authority of Poey.

Jabonillo, Parra, Dif. Piezas Hist. Nat., 51, pl. 24, fig. 2, 1787, Havana.

Anthias saponaceus Bloch & Schneider, Syst. Ichth., 310, 1801, Havana; after Parra.

Rypticus microps Castlenau, Anim. Nouv. et Rares, 6, 1855, Bahia; after *Percu microps* Broussonet, a MS. name.

Rypticus saponarius, Poey, Fauna Puerto-Riqueña, 322, 1881; Stahl, l. c., 76 and 162, 1883.

Rypticus saponaceus, Jordan & Evermann, l. c., 1232, 1896.




FIG. 46.—*Rypticus bistrispinus*.

118. *Rypticus coriaceus* (Cope). *Black Soap-fish*.

Head 3.3; depth 3.5; eye 5; snout 5.8; maxillary 2.4; mandible 1.7; interorbital 11.5; scales 19–125–32; D. m–25; A. 15; pectoral 1.5 in head; ventral 2.8; caudal 1.3.

Body rather elongate, compressed, back little elevated; head pointed, snout small, pointed; mouth large, somewhat oblique, maxillary reaching beyond orbit, its fore end very broad, triangular; lower jaw strongly projecting; teeth rather strong; interorbital narrow; preopercular spines 2, the lower the larger; opercle with 2 obscure spines, partly covered with skin, the lower the stronger; caudal peduncle compressed, deep, least depth 2 in head, its least width about 6 in its depth. Scales small, more or less embedded, especially on nape and cheeks. Fins all rather large; dorsal well separated, spines short; last dorsal rays about 2 in head; last anal rays about 2.75 in head; caudal rounded, its middle rays about 1.5 in head; pectoral broadly rounded; ventrals short, 2 times eye.

Color, plain brown, darkest on back; under parts paler; a broad white line from tip of lower lip to occiput; vertical fins dark, edges black; pectoral with some dark; ventrals pale.

Two specimens obtained, one 4.75 inches long, at Mayaguez, and one 5.5 inches long at Hucares. This West Indian species has hitherto been known only from St. Martins and Jamaica.

Eleutheractis coriaceus Cope, Trans. Amer. Philos. Soc, 1870, 467, St. Martins.

Rypticus coriaceus, Jordan & Evermann, l. c., 1235, 1896.

119. *Rypticus bistrispinus* (Mitchill).

Head 3; depth 3.6; eye 4.5; snout 4.75; maxillary 2.1; mandible 1.8; interorbital 7; D. n, 26; A. 15. Body rather slender, depth considerably less than length of head; head compressed, snout pointed; mouth large, lower jaw strongly projecting, maxillary reaching posterior border of orbit, its exposed portion broadly triangular; interorbital very narrow; 3 strong, distinct, preopercular spines, nearly uniform in size, middle one sometimes slightly the largest; opercle with 3 flat spines, middle

one largest; opercle ending in a long flap; scales very minute, cycloid; entire head scaled; lateral line complete, arched above pectoral; dorsal fin continuous, spinous portion continuous with soft rays; second dorsal spine longer than first, about equal to diameter of orbit; last dorsal rays longest, about 2.5 in head; anal resembling last portion of dorsal, its longest ray 2.5 in head; pectoral broad and rounded, its length 1.9 in head; ventrals short, scarcely greater than orbit; caudal rounded.

Color in alcohol: Rich brownish or olivaceous above, paler below; upper half of head to level of lower part of eye dark-brown; lower half of head pale-yellowish, usually with scattered minute brown specks; tip of lower jaw brown; pale median line from tip of snout to nuchal region, growing less distinct posteriorly; whole body, and usually most of head, covered profusely with small rich brown spots, largest above pectoral; under jaw, throat, and breast often immaculate; fins usually unspotted.

South Atlantic coast of the United States and West Indies, usually in rather deep water; known from off Charleston, the Bahamas, Key West, Pensacola, Cuba, and Porto Rico, and occasionally taken off Newport, Rhode Island. Nineteen specimens obtained by the Porto Rico expedition from the vicinity of Culebra Island, some dredged in 12 to 15 fathoms, others found in shallow water about the coral reefs; length about 3 inches. These specimens agree fully with Mitchell's original description of this species and with the type of *R. pituitosus* Goode & Bean.

Bodianus bistrispinus Mitchell, Amer. Month. Mag. and Crit. Rev., II, Feb., 1818, 247, deep water in Bahama Straits.

Rhypticus maculatus Holbrook, Ichth. S. Carolina, ed. 1, 39, 1856, and ed. 2, 42, 1860, Cape Romain, South Carolina.

Rhypticus pituitosus Goode & Bean, Proc. U. S. N. M. 1879, 341, Key West, Florida.

Rypticus bistrispinus, Jordan & Evermann, l. c., 1233, 1896.

Family XLII. LOBOTIDÆ. The Triple-tails.

This family is thus defined by Dr. Gill:

"Percoidæ with an oblong, compressed body, equally developed above and below; a short snout and anterior eyes; edentulous palate; dorsal and anal with the soft portions equal and opposite, the former preceded by a much larger spinous portion, the latter with 3 spines; vertebrae 24, 12 abdominal and 12 caudal, the fifth to eleventh with short but gradually lengthening parapophyses projecting sideways and behind downward, and the twelfth with the parapophyses elongated, converging at their extremities and fitting into a groove of the first hæmal spine, the costiferous pits excavated obliquely in the developed parapophyses, and gradually ascending forward on the vertebrae, and finally on the neurapophyses; the skull with its frontal portion broad, expanded forward and outward, and entering into the posterior borders of the orbits, which are advanced far forward; the postfrontals elongated forward and underlying the frontals; ethmoid short, decurved, and expanded sideways."

This family contains a single species, a large fish closely allied to the *Serranidæ*, but lacking vomerine and palatine teeth, and with the fore part of the head very short. Its relations are decidedly with the *Serranidæ* and not with the *Hæmulidæ*, with which group, however, it agrees in the absence of teeth on the palate.

Genus 75. LOBOTES Cuvier.

Body oblong, compressed, and elevated, covered with moderate-sized, weakly ctenoid scales; profile of head concave, snout prominent; mouth moderate, oblique, with thick lips; upper jaw very protractile, the lower longer; maxillary without supplemental bone; jaws with narrow bands of villiform teeth, in front of which is a row of larger conical teeth directed backward; no teeth on vomer or palatines; preorbital narrower than eye; preopercle strongly serrate. Branchiostegals 6. Dorsal fin continuous, with 12 spines, which may be depressed into a shallow groove; soft rays of dorsal and anal fins elevated; anal spines graduated; bases of soft dorsal and anal thickened and scaly; caudal rounded. Air-bladder present. Pyloric cæca 3.

120. *Lobotes surinamensis* (Bloch). "*Sama*"; *Flasher*; *Triple-tail*; *Dormcur*.

Head 2.8; depth 2; eye 6.5; snout 4.1; maxillary 2.8; mandible 1.9; interorbital 3.5; D. XI, 1, 15; A. III, 11; pectoral 1.7; ventral 1.4; caudal 1.4; scales 5-48-15.

Resembling the serranoid genus *Alphestes* in form, but without teeth on vomer or palatines; scales large, firm, and ctenoid; anterior profile with a strong concavity at occiput; eye small; mouth small,

lower jaw projecting; teeth in a narrow villiform band in each jaw, with a single outer row of enlarged conical teeth; preopercle with a few strong spines, those at angle greatly enlarged; soft dorsal and anal high, nearly as large as caudal, the 3 fins suggesting a 3-lobed tail, whence the name "triple-tail."

Color in life: Body greenish, grayish, and yellowish, more or less mottled; fins mottled-grayish; soft dorsal and anal black-edged, caudal with yellowish margin, black inside; ventral pale, with black blotches; branchiostegals pale, with irregular black lines; our smaller specimen creamy-yellow, overlaid with dark-brown.

A fish of rather sluggish habits, found in most warm seas; north on our coast to Cape Cod; probably not uncommon about Porto Rico; not recorded from Key West, though known from the St. Johns and Indian rivers, Pensacola and Tampa. It is said to reach a length of 3 feet and a weight of 50 pounds, and is regarded as a very good food-fish. The collection contains two young examples, 6.5 and 8 inches in length. The smaller was taken by the seine at the mouth of the Río Bayamon at Palo Seco, in water very nearly fresh; the other was seined in San Juan Harbor between Palo Seco and Cataño.

Holocentrus surinamensis Bloch, Ichth., pl. 243, 1790, Surinam.

Bodianus triurus Mitchell, Trans. Lit. and Phil. Soc., I, 1815, 418, Powles Hook, New Jersey.

Lobotes crata Cuvier & Valenciennes, Hist. Nat. Poiss., V, 322, 1830, Pondicherry.

Lobotes furkharii Cuvier & Valenciennes, Hist. Nat. Poiss., V, 324, 1830, Malacca; on a drawing by Major Farkhar.

Lobotes somnolentus Cuvier & Valenciennes, Hist. Nat. Poiss., V, 324, 1830, Santo Domingo.

Lobotes auctororum Günther, Cat., I, 338, 1859, Cuba; Calcutta; China.

Lobotes surinamensis, Poey, Fauna Puerto-Riq., 329, 1881; Stahl, l. c., 77 and 163, 1883; Jordan & Evermann, l. c., 1235, 1896.




FIG. 47.—*Lobotes surinamensis*.

Family XLIII. PRIACANTHIDÆ. The Catalufas.

Body oblong or ovate, compressed, covered with small, firm, rough scales; all parts of body and head, even the snout and maxillaries, being densely scaly, each scale with a more or less developed plate on its posterior border, most developed in the young. Head deep. Mouth large, very oblique, lower jaw prominent. Villiform teeth on jaws, vomer, and palatines, none on tongue. Premaxillaries protractile. Maxillary broad, without supplemental bone, not slipping under the very narrow preorbital, which is usually serrate; no suborbital stay. Eye very large, forming about one-half length of side of head. Posterior nostril long, slit-like, close to eye. Preopercle more or less serrated, one or more strong spines at its angle; operculum very short, ending in two or three points behind; no barbels. Gill-membranes separate, free from isthmus. Pseudobranchiæ very large, extending along whole length of opercle. Postorbital part of head very short, opercle small. Gills 4, a slit behind the fourth. Gillrakers long. Branchiostegals 6. Lateral line continuous, not extending on

caudal. Dorsal fin continuous, x, 9 to 15, spines depressible in a groove; anal rays iii, 9 to 15, soft part long, similar to soft dorsal, spines strong; ventrals very large, thoracic, 1, 5, close together, in advance of base of pectoral, joined to belly by a membrane which incloses a groove; no axillary process; spine strong; pectoral small, pointed, not symmetrical, of 19 or 20 rays, upper longest; caudal fin truncate or lunate. Spines of fins generally rough, with small serræ. Air-bladder large. Pyloric cæca few. Vertebrae in reduced number, 9 or 10 + 13 = 22 or 23, first vertebrae being very small or absent; transverse process beginning on seventh (sixth) vertebra, last two precaudal bridged across; ribs attached to transverse processes; epipleurals absent on last three precaudal vertebrae. Supra-occipital crest very low, continued forward to over front of orbit, where it is joined by parietal crests; processes of premaxillaries moderate.

Carnivorous fishes of the tropical seas, chiefly in deep waters; mostly rose-colored in life.

- a. Scales very small, 80 to 100 in lateral line; body oblong, its depth not half its length; preopercle with a flat spine; dorsal and anal each with 12 to 15 soft rays..... PRIACANTHUS, 76
 aa. Scales large and very rough, 35 to 50 in lateral line; body ovate, its depth more than half its length; preopercle without spines; dorsal and anal each with 9 to 11 soft rays..... PSEUDOPRIACANTHUS

Genus 76. PRIACANTHUS Cuvier.

Scales very small, 80 to 100 in lateral line; body oblong, more than twice as long as deep; preopercle with a spine at angle; interorbital area externally transversely convex, cranium itself transversely concave, elevation being formed of flesh; a conspicuous foramen in interorbital area; lateral line extending upward and backward from upper angle of gill-opening toward second dorsal spine, below which it changes its course, following outline of back to end of dorsal fin, thence direct to middle of caudal; anal fin rather long, its rays about iii, 14; dorsal rays about x, 13.

Species rather numerous, in the tropical seas.

- a. Preopercular spine obsolete or nearly so; depth about equal to length of head; dorsal unspotted; dorsal rays x, 14; anal iii, 15..... *arenatus*, 121
 aa. Preopercular spine well developed; depth of body greater than length of head; dorsal spotted; dorsal rays x, 12 or x, 13, anal iii, 13 or 14..... *cruciatatus*, 122

121. *Priacanthus arenatus* (Cuvier & Valenciennes). "*Toro*"; "*Comico*"; *Catalufa*.

(PLATE 16.)

Head 3.25; depth 2.8; eye 2.7; snout 3; maxillary 2; mandible 1.6; interorbital 4.6; D. x, 14, rarely 13; A. iii, 15, rarely 16; pectoral 1.9; ventral 1.1; caudal 1.2; scales about 94.

Body oblong-ovate, compressed, covered with small, very firm, and slightly ctenoid scales; head almost entirely scaled; eye very large; mouth extremely oblique; lower jaw strong and prominent; maxillary very broad posteriorly, reaching slightly beyond front of eye; opercle and angle of preopercle each with a weak flat spine; dorsal spines slightly roughened; caudal slightly lunate.

Color in life: Body and fins nearly everywhere bright-red, ventrals and caudal deepest; base of pectoral yellow; ventral spine pale-blue, rays black-tipped, most of membrane dusky; soft dorsal and anal with a few small dusky spots, and faintly edged with dark; caudal distinctly dark-edged; upper edge of caudal peduncle dusky; a series of about 12 indistinct dark round blotches along and just above lateral line; inside of mouth red behind; iris chiefly bright-red, a narrow yellow circle about pupil.

This fish occurs in the tropical Atlantic south to Brazil, occasionally northward in the Gulf Stream to Newport and Woods Hole; also known from Jamaica, Key West, and Madeira; probably common about Porto Rico. Our collection contains six examples, 6 to 12 inches long, from Aguadilla, Mayaguez, and Arroyo. It does not usually exceed a foot or 15 inches in length, but is a food-fish of some importance. Its flesh is firm and flaky and of good flavor. Nothing is known as to its game qualities. The brilliant red color and large eye make this a very striking fish.

Catalufa, Parra, Dif. Piezas Hist. Nat., pl. 20, 1787, Havana.

Priacanthus arenatus Cuvier & Valenciennes, Hist. Nat. Poiss., III, 97, 1829, Brazil and Atlantic; Jordan & Evermann, l. c., 1237, 1896.

Priacanthus fulgens Lowe, Trans. Zool. Soc. Lond., II, 1839, 174, Madeira.

Priacanthus catalufa Poey, Proc. Ac. Nat. Sci. Phila. 1863, 182, Havana.

122. *Priacanthus cruentatus* (Lacépède). "Ojon"; *Catalufa*; "Ojudo"; *Big-eye*.

Head 3; depth 2.5; eye 2.6; snout 3.2; maxillary 2; mandible 1.6; interorbital 3.7; D. x, 13; A. iii, 14; pectoral 1.8; ventral 1.4; caudal 1.2; scales about 90. Body deeper than in *P. arenatus*, and preopercular spine stronger, curved, and serrate.

Color in life: Body silvery, washed with rosy; back with five or six rosy, saddle-like blotches extending on sides to below lateral line; under parts rosy; vertical fins with pale bases, brighter outwardly; caudal black-edged; pectoral and ventral rosy, ventral black-tipped.

This fish ranges from the West Indies to St. Helena and the Canaries, and is known from Cuba, Jamaica, and Porto Rico, but not yet recorded from the United States. Probably not common in Porto Rico. Our collection contains but a single specimen, 8 inches long, obtained in the San Juan market, though others were seen. It reaches a length of a foot or more, and is a good food-fish, common in the Havana market.

Labrus cruentatus Lacépède, Hist. Nat. Poiss., III. 522, 1800, Martinique.

Priacanthus cepedianus Desmarest, Prem. Déc. Ichthy., 9, pl. 1, 1823, Havana; Poey, Fauna Puerto-Riqueña, 322, 1881.

Stahl, l. c., 76 and 162, 1883.

Priacanthus cruentatus, Stahl, l. c., 76 and 162, 1883; Jordan & Evermann, l. c., 1238, 1896.

Family XLIV LUTIANIDÆ. The Snappers.

Body oblong or more or less elevated, covered with moderate-sized adherent scales, which are more or less strongly ctenoid or almost cycloid. Lateral line well developed, concurrent with back, not extending on caudal fin. Head large, crests on skull usually largely developed. No suborbital stay; mouth moderate or large, usually terminal, low, and horizontal. Premaxillaries moderately protractile, their spines not extending to occiput; maxillary long, without supplemental bone, for most of its length slipping under edge of preorbital, which forms a more or less distinct sheath, its form essentially as in the *Serranidæ*; teeth various, unequal, and sharp, never incisor-like, some of them sometimes molar; vomer and palatines usually with villiform teeth, these sometimes molar, sometimes very small, sometimes wanting; lower pharyngeals separate; gills 4, a slit behind fourth; pseudobranchiæ large; gillrakers moderate or long, slender; gill-membranes separate, free from isthmus. Preopercle serrate or entire; opercles without spines; sides of head usually scaly. Dorsal fin single, continuous, or deeply notched, sometimes divided into two fins, spines usually strong, depressible in a groove, heteracanthous, that is, alternating, one stronger on right side, other on left; spines 10 to 12 in number. Anal fin similar to soft dorsal and with 3 spines; ventral fins thoracic, the rays 1, 5, with a more or less distinct scale-like appendage at base; caudal fin usually more or less concave behind. Air-bladder present, usually simple. Intestinal canal short. Pyloric cæca few. Vertebrae usually $10 + 14 = 24$. No distinct tubercles from cranium for articulation of epipharyngeal bones; enlarged apophyses for articulation of palatine and preorbital bones; anterior 4 vertebrae without parapophyses.

The *Lutianidæ* comprise about 20 genera and some 250 species, chiefly inhabiting shores of warm regions, all active, carnivorous, and voracious, and all valued as food. The group is closely related to the *Serranidæ* on the one hand and to the *Hæmulidæ* on the other.

HOPLOPAGRINÆ:

a. Vomer with teeth.

b. Nostrils near together, placed just before eye, anterior not tubular; vomerine teeth villiform, the patch \wedge , \blacktriangle , or \circ -shaped; teeth in jaws all acute; no incisors or molars.

c. Palatines with teeth; teeth in jaws strong, more or less unequal.

LUTIANINÆ:

d. Interorbital area not flat nor separated from occipital region, median and lateral crests procurvent on it, and frontal narrowed forward; dorsal fin continuous, spines not separated by a notch from soft rays.

e. Prefrontals with articular facets arising from diverging V-shaped ridges; basi-sphenoid with an anterior lobiform extension; soft dorsal and anal scaly; dorsal spines 10 or 11 (in American species); tongue with teeth (at least in adult examples).

f. Fronto-occipital crest ceasing anteriorly far from front of frontal; prefrontal with posterior areas impressed, long, and cribriform; no pterygoid teeth; caudal fin lunate or forked; gillrakers rather few, shortish.

g. Top of head naked; an oblique band of scales on each side of nape; parietal crest not confluent with fronto-occipital crest, either fading away anteriorly or running into ocular rim; preopercle with a shallow notch or emargination only. NEOMENIS, 77

- ff.* Fronto-occipital crest continued forward along top of head to nearly opposite nostrils; prefrontals with posterior area short, excavated above and in front.
- h.* Gillrakers long and numerous, about 25; anal rather high, its rays III, 9; pterygoid teeth present (in the adult) in a narrow band; caudal fin very deeply forked..... OCYRUS, 78
- cc.* Prefrontals with articular facets developed from simple tubercles and not V-shaped; basi-sphenoid not lobigerous; canines small; soft rays of dorsal 10 or 11.
- i.* Prefrontals with posterior areas cribriform; pterygoids with a broad patch of teeth (in adult); hyoid bones and tongue with teeth; canines very small or obsolete; dorsal spines 12 (or 13); soft dorsal and anal somewhat sealed; top of head sealed to before middle of eye; gillrakers numerous..... RHOMBOPLITES, 79
- ii.* Prefrontals with posterior areas solid and somewhat tumid; pterygoids, hyoid bone, and tongue toothless; dorsal spines 10; soft dorsal and anal scaleless..... APSILUS
- ETELINÆ:
- dd.* Interorbital area flat, separated by a transverse line of demarcation from occipital, by which median as well as lateral crests are limited; frontals wide in front; tongue and pterygoids toothless; soft rays of dorsal 10 or 11.
- i.* Dorsal fin continuous; frontals not cavernous; supraorbital margin crenate; pterotic region much swollen outward and with bones thin and polished; preorbital moderate; frontals behind with funnel-shaped foramina; soft dorsal and anal scaleless; last rays of dorsal and anal produced..... APRION
- jj.* Dorsal nearly or quite divided into two fins by a deep notch; eye very large; preorbital very narrow.
- k.* Frontals not cavernous, simply normally perforate; supraorbital margins crenate; prefrontals behind, with funnel-shaped foramina; head naked above and on snout; soft dorsal and anal naked; peritoneum and lining of gill-cavity pale; caudal deeply forked; color crimson..... ETELIS, 80
- kk.* Frontals cavernous (like those of Sciaenoids), with longitudinal, osseous bars, leaving interspaces in front of transverse ridge and on each side near front; supraorbital margins smooth; prefrontals behind with simple foramina for olfactory nerves; head scaly above and on jaws and snout; soft dorsal and anal scaly at base; peritoneum and lining of gill-cavity black; caudal lunate. Deep-water species, blackish-purple in color... VERLUS
- DENTICINÆ:
- aa.* Vomer and palatines toothless; one or both jaws with strong canines; no molars; preopercle entire; dorsal continuous.
- l.* Dorsal spines 10; scales large, 50 in lateral line, those on cheek in 3 rows; mouth moderate, jaws subequal; fins usually with filaments..... NEMIPTERUS

Genus 77. NEOMÆNIS Girard. Snappers.

Body oblong, compressed, back somewhat elevated; head long, naked above, except for a broad oblique band of scales at nape; nostrils normally close together, neither with a tube; mouth large, jaws with bands of villiform teeth, besides which is usually an outer series of larger teeth in each jaw, and 2 to 4 stronger teeth or canines in front of upper jaw; vomer with villiform teeth; villiform teeth on palatines; usually one or more patches of teeth on tongue in the adult; no molar teeth; no teeth on pterygoids; preopercle without notch or with a shallow emargination; posterior limb of preopercle finely serrate; gillrakers rather few, shortish; soft rays of dorsal and anal scaly at base; dorsal spines 10 (rarely 11), continuous with soft rays; caudal lunate or forked; anal rays 7 to 9. Interorbital area not flat nor separated from occipital region, median and lateral crests procurrent on it, and frontal narrowed forward; fronto-occipital crest ceasing anteriorly far from front of frontal, usually behind eye; prefrontal with posterior areas impressed, long and cribriform; parietal crest not confluent with orbital rim, but nearly or quite joined anteriorly to fronto-occipital crest (in species examined); prefrontals with articular facets arising from diverging V-shaped ridges; basisphenoid with an anterior lobiform extension. Vertebrae $10 + 14 = 24$.

We follow Jordan & Evermann in separating the American pargos or snappers from the Old World genus *Lutjanus* on the following characters, distinctive so far as known: Parietal crest usually confluent anteriorly with orbital rim, never joined anteriorly to fronto-occipital crest; top of head naked; a more or less isolated band of scales extending obliquely on each side of nape; notch on preopercle for reception of knob of interopercle shallow and broad, sometimes obsolete, otherwise essentially as in *Lutjanus*.

Species very numerous, American; active predatory fishes, highly valued as food.

Among the food-fishes of Key West and the West Indies no fewer than 15 species belong to this genus, while on our Pacific coast are at least 5 more species, and all of them are excellent for food. Most of the species reach a good size and are objects of important fisheries.

- a.* Soft dorsal normally with 14 rays, rarely with 13.
- b.* Anal fin rounded, its middle rays less than half length of head; no black lateral spot.
- c.* Developed gillrakers 7 to 9, usually with few rudiments, if any; preorbital deep; caudal lunate. Shallow-water species, olivaceous in color, more or less marked by crossbands when young, often with a blue streak along preorbital.
- d.* Vomerine teeth forming a Λ or ∇ shaped patch, backward prolongation on median line very short or wanting; scales above lateral line in oblique series, which are not throughout parallel with it; body comparatively elongate, depth 3 to 3.5 in length, upper and lower canines very strong, lower considerably stronger than in other species; mouth very large, vertical fins dusky; size very large.

- e.* Maxillary 2.29 in head; preorbital 4.75 in head; maxillary reaching past middle of eye, about 2.33 in head; usually a black spot or shade at base of pectoral; head 2.75; depth 3; D. x, 14; A. III, 8; scales 7-50-12. . . . *cyanopterus*, 123
- dd.* Vomerine teeth forming an anchor-shaped patch, with a distinct backward prolongation on median line; second anal spine little, if any, shorter than third; upper canines strong, lower moderate or small.
- f.* Scales above lateral line arranged in series which are not throughout parallel with lateral line, being oblique and irregular, at least below second dorsal.
- g.* Body comparatively elongate, depth 2.75 to 3 in length; mouth large, maxillary 2.5 in head; scales 7 in an oblique series between dorsal and lateral line; pectoral short, not two-thirds length of head; soft dorsal, anal, and caudal blackish, tinged with wine-color, always becoming dusky in spirits *griscus*, 124
- gg.* Body comparatively deep, depth about 2.5 in length; snout long and pointed; mouth rather small, maxillary about 3 in head; pectoral long, more than two-thirds length of head; soft dorsal, anal, and caudal orange or yellow, becoming pale in spirits.
- h.* Scales moderate, about 9 in an oblique series from first dorsal to lateral line, about 55 vertical series above lateral line between gill-opening and base of caudal; lateral line with more than 45 pores; a whitish area below eye; blue streak along suborbital region usually not disappearing with age; scales 8-56-15. *joeu*, 125
- hh.* Scales unusually large, 5 or 6 in an oblique series from first dorsal to lateral line, about 45 vertical series above lateral line between gill-opening and base of caudal; lateral line with fewer than 40 pores; blue streak on suborbital region not permanent; scales 6-44-13. *apodus*, 126
- cc.* Developed gillrakers more numerous, about 10, with several rudiments before them (in *N. buccanella*; not examined in *N. lutjanoides*).
- i.* Caudal deeply forked; mouth small, maxillary reaching posterior nostril; preopercle slightly notched, little serrate; canines strong; tongue with teeth; soft dorsal and anal rounded; pectoral pointed, 4.5 in total length. Color brownish-green, with 6 brown crossbands; a broad greenish stripe from opercle to base of caudal. D. x, 14; A. III, 8. *lutjanoides*
- ii.* Caudal moderately forked; mouth large, maxillary reaching anterior edge of eye, 2.6 in head; preopercle serrate, serrae strong on angle; canines medium; vomerine teeth in an anchor-shaped patch; eye large; the base and axil of pectoral with a jet-black blotch; scales moderate, about 8 oblique series from the lateral line to first dorsal spine, about 63 vertical rows above lateral line; second anal-spine long, about 2.66 in head. Color crimson; caudal peduncle and caudal fin largely yellow; iris orange-red; no lateral blotch. Head 2.5; depth 2.8; D. x, 14; A. III, 8; scales 8-63-15. *buccanella*
- bb.* Anal fin angulated, its median rays produced, longest in adult, at least half head; body rather robust; upper canines rather long; lower small. Color more or less red, young with a black lateral blotch.
- j.* Scales above lateral line arranged in series, which are not throughout parallel with it; side with a black blotch, which usually disappears with age; anal fin bright red.
- k.* Teeth on vomer in an anchor-shaped patch, with a median backward prolongation; lingual teeth well developed; maxillary reaching edge of pupil, 2.5 in head; caudal edged with black.
- l.* Iris golden-yellow in life. Scales rather small, 9-72-10, about 50 pores in lateral line; body rather slender, depth 2.5 in length; second anal spine about 3.5 in head; gillrakers 9 below angle; eye large, 4.75 in head in adult; preorbital 5.75 in head. Color bright rose-red, with golden streaks. *vianus*, 127
- ll.* Iris rose-red. Scales rather large, 8-60-14; body robust, depth 2.6 in length; second anal spine about 4 in head; gillrakers about 8 below angle. Color rose-red, nearly uniform; size large. *aya*, 128
- kk.* Teeth in vomer in a Λ -shaped patch, without distinct prolongation on median line; lingual teeth very few or none; maxillary reaching edge of eye, 2.7 in head *analis*, 129
- aa.* Soft dorsal with 12 rays (rarely 13); body oblong, the back not greatly elevated; upper canines moderate, lower small or obsolete; scales above lateral line in very oblique series; anal fin low, its outline rounded.
- m.* Mouth moderate; maxillary 2.6 to 2.75 in head.
- n.* Caudal not deeply forked; gillrakers rather few (8 or 9 besides rudiments).
- o.* Pectoral short, 1.66 in head; teeth on vomer in an anchor-shaped patch. Color olivaceous, no black lateral blotch; lower jaw included. Head 2.75; depth 3; D. x, 12; A. III, 8; scales 8-51-x. (Hybrid, *griscus* + *synagris*?).
brachypterus
- oo.* Pectoral long, more than two-thirds length of head; color chiefly red; a large black lateral blotch; lower jaw slightly projecting.
- p.* Vomer with teeth in a Λ or ∇ shaped patch, the prolongation on median line short.
- q.* Eye large, 4.75 in head; back greatly elevated; pectoral long, 1.25 in head. *megalopthalmus*, 130
- qq.* Eye smaller, 5 in head; back less elevated; pectoral shorter, 1.5 in head. *synagris*, 131
- nn.* Caudal deeply forked; gillrakers rather numerous, about 10 on lower part of anterior arch; teeth on vomer in an anchor-shaped patch; body rather elongate, compressed; lower jaw projecting or not; eye small; scales small; lateral line with about 50 pores; anal spines graduated. Color reddish, with horizontal yellow streaks; no black lateral blotch. Head 3; depth 3; D. x, 13; A. III, 9; scales 9-53-15. *ambiguus*
- mm.* Mouth large; maxillary 2.4 in head; teeth on vomer in an anchor-shaped patch. *mahogoni*, 132

123. *Neomænis cyanopterus* (Cuvier & Valenciennes). *Cubera*.

Head 2.75; depth 3; eye rather small, 5.66 in head; D. x, 14; A. III, 8; scales (6) 7-50-12, 50 pores. Body elongate, rather robust, back little elevated; profile from snout to nape nearly straight; snout long, thick, rather acute in profile, 3 in head; interorbital space flattish or gently convex, 6.25 in head; occipital keel low; preorbital broad, 4.66 in head; mouth very large; maxillary reaching middle of eye, 2.33 in head. Canine teeth larger than in any other of the genus, especially those in lower jaw; upper

jaw with a narrow band of villiform teeth, outside of which is a series of strong, sharp teeth; 4 canines in front, 2 of them very long and strong, their length two-thirds diameter of eye; lower jaw with 5 or 6 very strong canine-like teeth on each side, largest little smaller than canines of upper jaw; a few villiform teeth in front of jaw; tongue with a large oblong patch of teeth, pointed behind, its length about twice its greatest width; vomer with a Λ -shaped patch of teeth, usually without backward prolongation on median line, but sometimes with a short median prolongation (\wedge -shaped), its length always less than width of patch in front; pterygoid and hyoid bones without teeth. Gillrakers rather short and thick, about one-third length of diameter of eye, about 8 on lower arch; no rudiments. Preopercle with posterior margin nearly vertical, emargination broad and shallow, the edge finely serrate above, teeth coarser just above angle, lower limb almost entire. Scales rather large, loosely attached; cheek with about 8 rows, 1 row on interopercle, 1 row on subopercle, and about 7 on opercle; temporal region with about 2 rows of large scales; tubes of lateral line simple; base of soft dorsal and anal scaly. Dorsal spines rather strong, outline of fin gently convex, fourth spine longest, 3.25 in head; tenth spine, 6 in head; anal spines strong, second spine stronger, slightly shorter than third, which is 5 in head; caudal little forked; pectoral about 1.4 in head.

Color, dusky-gray, paler below, belly sometimes tinged with reddish; membranes of dorsal, anal, and caudal grayish-black, anal and soft dorsal especially blackish; ventrals blackish at tip; pectoral plain olivaceous, base and inner margin dusky; head dusky above, without markings.

This fish attains a length of 2 to 4 feet; the specimen described (from Cuba) measured $17\frac{1}{2}$ inches. It is found in the West Indies and south to Brazil, and is rather common. It is a large, coarse fish, regarded as unwholesome by fishermen, but probably without sufficient cause. It is recorded by Cuvier & Valenciennes and by Poey from Porto Rico; not seen there by us.

Mesoprion cyanopterus Cuvier & Valenciennes, Hist. Nat. Poiss., II, 472, 1828, Brazil.

? *Mesoprion pargus* Cuvier & Valenciennes, Hist. Nat. Poiss., II, 473, 1828, Porto Rico.

Lutjanus dentatus A. Duméril, in Vaillant & Bocourt, Miss. Sci. au Mex., 125, 1881, Brazil.

Lutjanus pargus, Poey, Fauna Puerto-Riqueña, 320, 1881.

Neomænis cyanopterus, Jordan & Evermann, l. c., 1254, 1898.

124. *Neomænis griseus* (Linnæus). *Gray Snapper; Mangrove Snapper; Cabellerote; "Pargo Prieto."*

(PLATE 17.)

Head 2.75; depth 2.87 to 3.25; D. x, 14; A. 11, 8; scales (6) 7-50-12, 47 pores. Body comparatively elongate, back not strongly compressed, little elevated; profile almost straight from snout to nape, thence gently convex. Snout rather pointed, 3 in head. Eye rather small, 4.66 in head. Interorbital space gently convex, 6 in head; occipital keel little prominent; preorbital rather broad, 5.5 to 6.5 in head. Mouth large; jaws subequal; maxillary reaching front of pupil, 2.6 in head; upper jaw with a narrow band of villiform teeth, outside of which is a single series of enlarged teeth; 4 canines in front of upper jaw, 2 of them quite large—one-third diameter of eye; lower jaw with a very narrow band of villiform teeth in front of jaw only; outside of these a single row of teeth larger than outer teeth of upper jaw, becoming canine-like in adult; tongue with an oval patch of teeth, its width about half its length; vomer with an arrow-shaped patch of teeth, with backward prolongation on median line, its length about twice its width in front. Gillrakers rather short and thick, their length about one-third diameter of eye, about 8 on lower arch, with no rudimentary ones before them. Preopercle with its posterior margin nearly vertical, with rather broad and deep emargination; preopercle finely serrate above, teeth coarser at angle. Scales comparatively large, rows in horizontal series below lateral line, those above running parallel with lateral line until below soft dorsal, where they become slightly irregular and oblique; 7 rows of scales on cheek; an embedded row on interopercle; 1 row on subopercle, and 7 on opercle; temporal region with about 3 rows of large scales; top of head, snout, and jaws naked; base of soft dorsal and anal scaly; tubes of lateral line branched. Dorsal spines rather strong, the outline of fin gently convex; fourth spine longest, 2.5 in head, tenth spine 4 in head; margin of soft dorsal rounded; ninth and tenth rays longest, 1.33 length of first, and 1.6 last ray, 2.5 in head; caudal emarginate, upper lobe longest, 1.33 length of middle rays, which are 1.75 in head; anal fin high, its margin slightly angulate, middle rays longest, 2 times length of last ray, 2.16 in head, first ray reaching almost to tip of last ray, when the fin is depressed; second anal spine as long or slightly longer and stronger than third, 3.25 to 4 in head; ventrals 1.75 in head; pectoral shortish, scarcely reaching vent, 1.57 in head.

Color in life: Very dark green above, middle part of each scale brassy-black, its edge broadly pearly-whitish; below lateral line the duskiness of middle of scale passes into brassy, and below into bright coppery, belly and lower parts of head being more or less distinctly bright coppery-red; lower jaw grayish; no blue stripe below eye, except in very young; top of head blackish-olive; dorsal blackish, its margin darker and tinged with maroon-red; soft dorsal dusky, anteriorly slightly edged with whitish; caudal violaceous or maroon-black; anal wine-color, edged with whitish; pectoral pale flesh-color; ventrals whitish, faintly marked with reddish. Young with a blackish band from snout through eye to nape, very distinct in life; a blue streak below eye; spinous dorsal with a dark maroon-colored band along edge. An example, 10.5 inches long, from the San Juan market, was brownish-red on body; blue suborbital streak quite distinct and continuous; two blue streaks below and forward from eye, only one on right side; apparently a deep-water example. Fishes from deep water are much redder than those taken near shore. In no case is the caudal yellowish or of any pale shade.

The above description chiefly from a specimen 11 inches long from Key West but verified on specimens from Porto Rico. The variations in proportional measurements are not great.

Found in the West Indies and ranging from New Jersey to Brazil; very common along our South Atlantic and Gulf coasts, occasionally straying northward as far as New Jersey, being the northernmost in its range of any member of the genus in the Atlantic; everywhere generally known as gray snapper. In Florida and the Bahamas, where the coasts are lined by mangrove bushes, among which the young of this species abound, the name mangrove snapper comes into use. It inhabits water of varying depths, large specimens being often found near the shore, while others may be taken in waters of considerable depth in company with *Neomænis aya*. These latter individuals are much redder than those found in shoal water; their general color is paler and the body is a trifle less elongate; such correspond to the form named *Lutjanus stearnsi*.

All the snappers are game-fishes of considerable importance, and the gray snapper is one of the best. Its abundance and wide distribution, the ease with which it can be found at all seasons, together with the readiness and vigor with which it takes the hook and the fairly good fight which it makes, should cause this fish to be much sought after by anglers who visit our southern and tropical waters.

In Indian River, Florida, the mangrove snapper is regarded as a very good food-fish and is of considerable commercial importance, its average weight being about 2 pounds, the maximum about 6 or 7 pounds. At Key West, where it is the most abundant of the snappers, it attains a length of 3 feet and a weight of 18 pounds, though the average of those caught is 5 pounds or less. Here it is usually called gray snapper and is regarded as a warm-water fish, being most plentiful in shallow water in summer, but retiring to deeper water during winter, and always running in schools; it is said to spawn in July and August, usually on the shoals, the eggs being nonadhesive and separating from each other at spawning; it is caught with hook and line, sardines and pilchard being the usual bait.

About Porto Rico this is an important food-fish and is known as "pargo prieto." Specimens are in the collection from San Juan market, Puerto Real, Arroyo, and Isabel Segunda, and one from San Geronimo. It was one of the most common species in the San Juan market and was seen in all the other markets of the island. The largest seen weighed about 6 pounds.

Turdus pinnis branchialibus carcus (mangrove snapper), Catesby, Hist. Carolina, pl. 9, 1743, Bahamas.

Caballerote, Parra, Descr. Dif. Piezas, Hist. Nat., pl. 25, fig. 1, 1787, Havana.

Labrus griseus Linnaeus, Syst. Nat., ed. X, 283, 1758, Bahamas; after Catesby.

Sparus tetracanthus Bloch, Ichthyol., pl. 279, 1791, Martinique; on a drawing by Plumier.

Anthias caballerote Bloch & Schneider, Syst. Ichth., 310, 1801, Cuba; after Parra.

Bodianus vivancti Lacépède, Hist. Nat. Poiss., IV, pl. 4, fig. 3, 1803, Martinique; on a drawing by Plumier.

Mesoprius griseus Cuvier & Valenciennes, Hist. Nat. Poiss., II, 469, 1828, Santo Domingo; not after Linnaeus.

Lobotes emarginatus Baird & Girard, 9th Smith. Rept. 1855, 332, Beesley Point, N. J.

Lutjanus stearnsi Goode & Bean, Proc. U. S. N. M. 1878, 179, Pensacola, Fla.

Neomænis griseus, Jordan & Evermann, l. c., 1255, 1898.

125. *Neomænis jocu* (Bloch & Schneider). "Pargo Colorado"; Dog Snapper; Joca.

(PLATE 18.)

Head 2.57; depth 2.57; eye 4.75; snout 2.6; maxillary 2.57; mandible 5; interorbital 5.5; D. x, 13; A. III, 8; scales 8-50-16, about 45 pores. Body comparatively deep and compressed, back elevated; profile steep and almost straight from snout to nape, thence little convex; snout rather long and pointed; eye moderate; interorbital space narrow, gently convex; occipital keel moderate; preorbital

broad, 4.33 in head; mouth rather large, jaws subequal; maxillary reaching front of orbit; upper jaw with a narrow band of villiform teeth, outside of which is a single series of larger teeth; 4 canines in front of upper jaw, 2 of them very large, almost equaling in length diameter of pupil; lower jaw with a narrow villiform band in front only, and a series of larger teeth outside, largest on side of jaw almost canine-like; tongue with a single large oval patch of teeth, its length more than twice its width; teeth on vomer forming a broadly arrow-shaped patch with backward prolongation on median line twice length of width of anterior part. Gillrakers rather short and thick, longest about one-fourth diameter of eye, about 9 on lower part of arch, with no rudiments in front of them. Preopercle with its posterior margin slanting obliquely downward and forward, emargination very broad and shallow; preopercle finely serrate above, teeth coarser at angle, which is not salient. Scales moderate, smaller than in *N. griseus* or *N. apodus*, in nearly horizontal series below, and obliquely upward and backward above lateral line; about 7 or 8 rows of scales on cheek; 1 row on interopercle, 1 on subopercle, and 7 on opercle; about 3 rows of large scales on temporal region; top of head, snout, and jaws naked; tubes of lateral line branched; bases of soft dorsal and anal scaly; dorsal spines rather strong, outline of fin evenly curved, fourth and fifth spines longest, 2.6 in head; tenth spine 4 in head; margin of soft dorsal convex, middle rays longest, 2.83 in head; caudal little forked, upper lobe longest, 1.4 length of middle rays, 1.6 in head; margin of anal well rounded, middle rays about twice length of last ray; first ray reaching nearly to tip of last ray when fin is depressed; pectoral slightly falcate, reaching almost to front of anal, 1.33 in head; anal spines strong, second rather longest and strongest, not always reaching past tip of third, 3.4 in head.

Color of adult in life: Olivaceous above, paler below, much flushed, so that the general hue is everywhere coppery-red; sides of body with numerous narrow crossbars, rather faint, the light and dark of about equal width, or the pale narrower; scales of upper parts mesially bronzed; head coppery; especially above; broad whitish area from eye to angle of mouth, becoming rosy in spirits; an irregular line of small round, or oblong spots below the eye, from snout to angle of opercle; soft fins all plain light brick-red, anal somewhat orange, caudal more or less yellowish; spinous dorsal with a light orange band at base and edge, middle pearly; blue stripe below eye persisting longer than in any of the other species which possess it. Young, in life: Greenish-olive, head and breast flushed with bright coppery-red; base of each scale bright orange-yellow, this color more extensive than dark ground-color, so that the general hue of body, especially below and posteriorly, is rich golden-yellow; dusky spot on top of head; temporal region with a dusky shade; an undulating blue stripe below eye from snout to angle of opercle; a similar fainter streak below it; pectoral pale-red or light-orange; ventrals orange; other fins rich golden-yellow, front of anal and edge of spinous dorsal rich, clear, bright-orange.

The above description based upon specimens 10 and 12 inches long. The dog snapper bears some resemblance to the gray snapper, but is not quite so trim a fish, as may be seen by comparing plates 17 and 18. The body color is similar, but the fins are colored very differently.

This excellent food-fish is known from the West Indies south to Bahia and north to Florida Keys, occasionally straying north in the Gulf Stream to Woods Hole. It attains a weight of about 20 pounds, though the individuals usually seen are much smaller. At Key West it occurs in greatest numbers during fall and winter, but is not very common at any time; nor does it seem to be abundant in Porto Rico. A good many were seen in the market at San Juan where four, from 10 to 15 inches long, were obtained. Specimens were secured also at Palo Seco and Vieques Island. Our notes make no mention of it as having been seen elsewhere about the island.

Jocu, Parra, Descr. Dif. Piezas, Hist. Nat., I, pl. 25, fig. 2, 1787, Cuba.

Anthias jocu Bloch & Schneider, Syst. Ichth., 310, 1801, Cuba; after Parra.

Mesoprion litura Cuvier & Valenciennes Hist. Nat. Poiss., II, 467, 1828, Cayenne; St. Thomas.

Neomænis jocu, Jordan & Evermann, l. c., 1257, 1898.

126. *Neomænis apodus* (Walbaum). "*Pargo Amarilla*"; *Schoolmaster*; *Caji*.

(PLATE 19.)

Head 2.5; depth 2.83; eye 4.33; snout 2.67; maxillary 2.67; interorbital 4.83; preorbital 4.83; D. x, 14; A. iii, 8; scales 5-42 to 45-13, about 36 pores. Body comparatively deep, moderately compressed, back considerably elevated; profile almost straight from snout to nape, nuchal region rather convex; snout unusually long and pointed, its outline before eye a little depressed, its length 2.71 in head; eye moderate, 4.33 in head; interorbital space flattish or gently convex. 5.5 in head;

mouth large, maxillary reaching front of orbit, 2.6 in head; upper jaw with a narrow band of villiform teeth, outside of which is a single series of larger teeth; 4 canines in front of upper jaw, one of them on each side very large, almost as long as pupil; lower jaw with a narrow villiform band in front only, and an enlarged series outside, these largest on side of jaw, where some of them are somewhat canine-like; tongue with a single large oval patch of teeth, its length more than twice its width; teeth on vomer forming an arrow-shaped patch with backward prolongation on median line, length of which is twice the width of arrow-patch in front. Gillrakers rather short and thick, longest about one-third diameter of eye, about 9 on lower part of arch. Preopercle with its posterior margin directed somewhat obliquely forward, usually very weakly emarginate, finely serrate above, almost entire at angle. Scales large, decidedly larger than in *N. jocu*; series below lateral line almost horizontal; those above in rows parallel with lateral line, these becoming more or less irregular posteriorly and extending upward and backward below soft dorsal; about 7 rows of scales on cheek, 1 row on interopercle, 1 on subopercle, and 7 on opercle; temporal region with a few large scales in about 2 rows; base of soft dorsal and anal scaly; tubes of lateral line each with 4 or 5 branches. Dorsal spines strong, outline of fin not greatly convex, fourth spine longest, 2.66 in head, tenth spine 4 in head; margin of soft dorsal well rounded, middle rays longest, twice length of last, 2.75 in head; caudal not deeply forked, upper lobe longer, 1.5 length of middle rays, which are 2 in head; margin of anal well rounded, its middle rays twice length of last, 2.3 in head, first ray reaching about to middle of last when fin is depressed; anal spines strong, second longer than third, 3.33 in head; ventrals 2 in head; pectoral reaching to front of anal, 1.33 in head.

Color of young in life: Greenish, with about eight very narrow vertical paler bars on body; scales of lower part of side with central orange spots, forming faint streaks along rows of scales; belly pearly; head greenish; a blackish streak from snout through eye to nape; a narrow, sharply defined blue stripe below eye from snout to angle of opercle; no lateral spot; spinous dorsal edged with orange; ventrals, anal, and caudal pale orange-yellow; pectoral paler. Adult examples differ from the young in the vertical bars being fainter or obsolete, and in the absence, usually, of blue stripe below eye and dark stripe on temporal region; soft dorsal, anal, and caudal always yellow, of varying intensity, and the edge of spinous dorsal orange, not dusky; the whitish area below eye, very constant in *N. jocu*, is wanting in *N. apodus*.

A specimen, 9 inches long, from Arroyo, had the following colors: Upper parts reddish or purplish and brownish, with some yellow or brassy tints; lower sides and belly purplish-rosy; cheek pale-rosy; top of head dark-rosy; lower jaw white; fins all rich orange or gamboge-yellow, a triangle at tip of each dorsal membrane very rich; ventral rays whiter than membranes; side with about 12 faint brassy lines running backward and somewhat upward; inside of mouth pale.

The schoolmaster is one of the most richly colored of the snappers, and is, withal, a very interesting and attractive fish. It is known from the West Indies, south to Bahia and north to southern Florida, sometimes straying northward in the Gulf Stream to Woods Hole; known from Indian River and Key West, but not common at either place. It is one of the most abundant of the snappers occurring in Porto Rico. Numerous specimens are in the collection from San Juan, Palo Seco, Puerto Real, Ensenada del Boqueron, Ponce, Arroyo, Hueares, Vieques, and Culebra. It is evidently one of the most valued food-fishes of the island. None seen about Porto Rico would exceed a pound in weight. At Key West it is said to reach a weight of 7 or 8 pounds, though the average of those caught on the reefs does not exceed 3 pounds, while those caught in the "bay" do not often weigh more than one-third of a pound. It is said to take the hook readily and with considerable vigor.

? *Perca marina pinnis branchialibus carens* (Schoolmaster), Catesby, Hist. Carolina, etc., tab. 41, 1743, Bahamas; figure very poor, the pectoral fins omitted.

Caris, Parra, Descr. Dif. Piezas, Hist. Nat., pl. 8, fig. 2, 1787, Havana.

? *Perca apoda* ("Forster, Catal. of Anim., MS., 21," 1774; printed 1844), Walbaum, Artedi Piscium, 351, 1792, Bahamas; based on the schoolmaster of Catesby.

Sparus caris Bloch & Schneider, Ichthyol., 284, 1801, Havana; after Parra.

Bodianus striatus Bloch & Schneider, Syst. Ichth., 335, pl. 65, 1801, West Indies; misprinted *albostrigatus*, p. 237; called *B. fasciatus* on plate.

Lutjanus acutirostris Desmarest, Prem. Déc. Ichthyol., 12, pl. 3, 1823, Cuba.

Mesoprion cynodon Cuvier & Valenciennes, Hist. Nat. Poiss., II, 465, 1828, Martinique; Santo Domingo.

Mesoprion linea Cuvier & Valenciennes, Hist. Nat. Poiss., II, 468, 1828, Cuba; Santo Domingo.

Mesoprion flavescens Cuvier & Valenciennes, Hist. Nat. Poiss., II, 472, 1828, Martinique.

Neomacris apodus, Jordan & Evermann, I. c., 1258, 1898.

127. *Neomænis aya* (Bloch). *Red Snapper*; *Pargo Colorado*; *Pargo Guachinango*; *Acara Aya*.

(PLATE 20.)

Head 2.6; depth 2.6; D. x, 14; A. iii, 9; scales (7) 8-60-15, pores 46. Body rather deep, moderately compressed, back well elevated, profile steep, and almost straight from snout to nape. Snout rather pointed, 2.8 in head; eye moderate, 5.5 in head (larger in young). Interorbital space angulate or strongly convex, 5 in head; occipital keel strong; preorbital rather broad, 5 in head; mouth rather large, maxillary reaching front of orbit, 2.5 in head; upper jaw with a narrow band of villiform teeth, outside of which is a row of larger but comparatively small teeth; 4 canines in front, 2 (sometimes duplicate) of them larger, their length about one-third diameter of eye; lower jaw with a single row of rather small teeth, usually largest on side of jaw, where some of them are almost canine-like; within these is a very narrow band of villiform teeth in front of jaw only; tongue with a broad oval patch of teeth, scarcely twice as broad as long; in front of this patch is a small, irregular patch; vomer with a broadly arrow-shaped patch, with a rather short backward prolongation on median line, its length about equaling width of patch in front. Gillrakers moderate, their length about half diameter of eye, 8 on lower arch. Preopercle with its posterior margin about vertical, its emargination deep, its edge rather finely serrate above, coarser at angle, dentate on lower border. Scales rather large, rows horizontal below lateral line, rows above running backward and upward; 6 rows of scales on cheek, 1 on interopercle, 1 on subopercle, and 7 on opercle; bases of soft dorsal and anal scaly; pores of lateral line branched; temporal region with a broad band of scales, with a few scattering ones below it; top of head, snout, and jaws naked. Dorsal spines rather strong, outline of fin moderately convex, fourth and fifth spines longest, 2.8 in head; tenth spine about 4 in head; margin of soft dorsal nearly straight, fin pointed behind; middle rays little longer than first ray, 1.5 length of last, 3 in head; caudal lunate, upper lobe scarcely longer than lower, its length 1.4 times length of middle rays, which are 1.86 in head; margin of anal strongly angulate, middle rays reaching nearly to base of caudal, 2.5 length of last ray, 1.8 in head; first ray reaches about to middle of last ray when fin is depressed; anal spines strong, second scarcely as long as third, 4 in head; ventrals, 1.8 in head; pectoral reaching to front of anal fin, 1.2 in head.

Color in life: Deep rose-red, paler on throat; bluish streaks along rows of scales, above becoming fainter and disappearing with age; fins brick-red; dorsal bordered with orange, with narrow blackish edge; caudal narrowly edged with blackish; eye red; a large blackish blotch above lateral line and below front rays of soft dorsal in the young, this spot usually disappearing with age; axil of pectoral dusky.

A specimen 21 inches long taken at Aguadilla, and from which plate 20 was made, showed in life the colors faithfully portrayed by the artist. The following color notes were made upon this specimen at the time: Back and side down to lateral line rich rosy, scales edged with paler; side below lateral line and belly pale-rosy; a small black spot just above lateral line below second and third dorsal rays; dorsal fins pale-rosy on basal half, tips of rays and spines pale-yellowish; caudal red, lower lobe richest, upper yellowish on outer portion; anal rich-rosy on anterior outer part, rest of fin paler; ventral rosy; pectoral pale-rosy; upper part of head rose-color, side paler; some irregular blue lines or spots about eye; inside of mouth white. The general color of the Porto Rican red snappers seems to be a decidedly paler red than that of Florida examples and the black spot on the side appears to be more persistent. Otherwise the colors are essentially the same.

Found from Long Island to Brazil on rocky banks in rather deep water, especially abundant in the Gulf of Mexico, off Cape San Blas and about Yucatan, where the most important fisheries are located; also abundant on the banks off eastern Florida and Georgia. About Porto Rico it is said to be plentiful, though at the time of our visit it was seen only at Aguadilla, probably because the fishermen of that place go out toward Mona Island and fish in deep water. It reaches a length of 2 to 3 feet and a weight of 30 to 40 pounds, though the average weight of those caught on the Florida "Snapper Banks" is much less. Adams and Kendall give the weight of those examined on the banks from the Tortugas to the latitude of Charlotte Harbor at 5 to 20 pounds. The largest seen by them was 32 inches long; one 30 inches long weighed 18 pounds.

The red snapper is the most valuable food-fish of the genus in the waters of the United States. It is the object of the principal fishery of Pensacola, from which point it is shipped fresh to all important cities in the United States as far north as Boston, Chicago, and Minneapolis and west to Omaha and Denver. In Florida it is known everywhere as "red snapper," or to the Spanish-speaking people as

"pargo colorado." In Havana it is known as "pargo guachinango," or "Mexican snapper," because it was brought to that city from the coast of Mexico. In Porto Rico it is called "pargo colorado."

The red snapper is strictly carnivorous, feeding upon small fish, crabs, prawns, and mollusks. It is caught only with hand lines and in 15 to 50 fathoms of water. As the fish are hauled up from these considerable depths the rapid decrease in external pressure permits the air-bladder to expand and the stomach is often forced out into the mouth of the fish and the stomach contents thrown out upon the deck of the smack. Among the material thus ejected may frequently be seen specimens of small fish or crustaceans still alive and only very slightly injured. In this way have been obtained the types of several new fishes and good specimens of many rare species. The stomach of the specimen painted for this report contained one *Olivia* shell and the partially digested body of a *Trachurops crumeniophthalmus*.

There is considerable sport in catching the red snapper. The line used is the size of an ordinary chalk-line; at its lower end is a sinker weighing about 3.5 pounds, consisting of a piece of lead shaped like a frustum of a cone, with the lower end hollowed out and filled with a mixture of lard and wax, for the purpose of determining the character of the bottom, as some of the sand, coral, shells, mud, or whatever the bottom is composed of, will stick to the wax and be brought up with it. On the line a few feet above the lead are attached two short lines, with hooks, not quite long enough to reach the lead. These hooks are baited with pieces of meat, bone-fish, lady-fish, or some other species. When a fish strikes, a quick jerk is given to fasten it and then the line is hauled in hand over hand. The red snapper does not ordinarily make any fight, but comes up as a dead weight until near the surface of the water, when, getting sight of the boat and fishermen, he becomes frightened and begins darting to right and left in the most frantic manner. The sport then is very exciting and, if the snapper be large, assistance may have to be called in order to finish the fight and lift the fish on deck. It sometimes happens that a fish is gotten on each hook at the same time.

Acara aya Maregrave, Hist. Brasil., 167, 168, 1648, Brazil.

Bodianus aya Bloch, Ichthyol., 227, 1790, Brazil; after Maregrave.

Bodianus ruber Bloch & Schneider, Syst. Ichthy., 330, 1801, Brazil; based on Maregrave.

Mesoprion campechanus Poey, Memorias, II, 149, 1860, Campeche.

Lutjanus blackfordi Goode & Bean, Proc. U. S. N. M. 1878, 176, Pensacola.

Neomænis aya, Jordan & Evermann, l. c., 1264, 1898.

128. *Neomænis vivanus* (Cuvier & Valenciennes). *Pargo de lo Alto; Silk Snapper.*

Head 2.75; depth 3; D. x, 14; A. m, 8; scales (7) 8-72-17, 50 pores. Body rather slender, sub-elliptical, back not greatly elevated; profile very slightly convex from snout to nape, thence more arched; snout rather long and pointed, 3 in head; eye rather large, 4 in head; interorbital space slightly convex, 4.8 in head, occipital keel not very prominent; preorbital rather broad, 5.8 in head; mouth rather small; jaws subequal; maxillary reaching front of pupil, 2.5 in head; upper jaw with a narrow band of villiform teeth, outside of which is a single series of well-developed teeth; 4 moderate canines in front of jaw, the longest two about half diameter of pupil; lower jaw with a single series of rather large, unequal teeth, inside of which is a very narrow band of villiform teeth in front of jaw only; tongue with an oval patch of teeth, about twice as long as broad, in front of which is a roundish patch; no teeth on hyoid bone; pterygoids toothless; vomer with a broadly arrow-shaped patch of teeth, with a backward prolongation on median line somewhat longer than width of patch in front. Gillrakers slender, their length almost equal to half diameter of eye, about 11 developed below angle, in front of these about 5 rudiments. Preopercle with posterior limb slanting slightly downward and forward, with a broad and rather shallow emargination, its margin finely serrate above; coarser teeth at angle and on lower limb; posterior nostril oval. Scales very small, rows running obliquely upward and backward above lateral line, rows being almost horizontal; 7 rows of scales on cheek, 2 rows on interopercle, 1.5 rows on subopercle, and about 8 on opercle; temporal region with 1 row of large scales, behind which are smaller ones; top of head, snout, and jaws naked; bases of soft dorsal and anal scaly. Dorsal spines rather strong, outline of fin rather strongly convex and without deep emargination; fourth spine longest, 2.4 in head; tenth, 3.4 in head; margin of soft dorsal straightish, rounded behind, ninth ray longest, 1.33 length of first and two times last ray, 2.5 in head; caudal lunate, upper lobe slightly longer than lower, its length 1.5 times middle rays, which are 2 in head; margin of anal angulate, middle rays longest, 2 times length of last ray, 1.8 in head; first ray reaching almost to tip of last ray, when the fin is depressed; ventrals 1.66 in head; pectoral not quite reaching front of anal, 1.2 in head; second anal spine slightly longer than third, 3.25 in head,

Color in life: Bright rose-color, paler below, some narrow, undulating, light-golden streaks following rows of scales above lateral line; iris always bright-yellow (an important color-mark); mouth reddish within; traces of dark lateral spot in most specimens; dorsal rosy, its base pale, its edge yellow; caudal rosy, dusky behind, sometimes blood-red at tip; pectoral very pale yellow, ventrals and anal pale rosy, latter yellowish behind. The bright colors all fade and disappear in spirits. The scales of upper parts, in spirits, are marked with dark dots, which form streaks along rows of scales.

The above description verified on a specimen 14 inches long from Mayaguez.

West Indies; known from Cuba, St. Kitts, Martinique, and Porto Rico. A handsome species, rather common in the Havana markets, where it is known as pargo de lo alto. When fresh it may always be known by the bright-yellow color of the eye, a color which does not entirely fade in spirits.

The silk snapper was not common in the Porto Rican markets during our stay about the island, but Mr. Oscar Riddle says that it is quite common in the San Juan market at certain times. It is taken in the line fishery and is one of the most valued species. It is usually fished for in about 60 fathoms of water, and about a mile off Morro Castle. Five or 6 hooks are fastened with short snoods to a very tough, strong stick of native wood about 2 feet long, and this is weighted with heavy leads. In the center is fastened the strong hand-line with which it is lowered and raised. The hooks are usually baited with pieces of sardina (*Opisthonema oglinum*) or muniama (*Xystema cinereum*). While the fish are being raised to the surface they are very often attacked by sharks. Our collection contains two specimens, 9.5 and 14 inches long, obtained in the market at Mayaguez.

Mesoprion vivanus Cuvier & Valenciennes, Hist. Nat. Poiss., II, 454, 1828, Martinique.

Mesoprion profundus Poey, Memorias, II, 150, 1860, Cuba.

Lutjanus torridus Cope, Trans. Am. Philos. Soc. 1869, 468, St. Kitts.

Lutjanus profundus Poey, Fauna Puerto-Riqueña, 320, 1881; Stahl, l. c., 76 and 162, 1883.

Neomænis vivanus, Jordan & Evermann, l. c., 1262, 1898.

129. *Neomænis analis* (Cuvier & Valenciennes). Mutton-fish; Pargo Criollo; Pargo; "Sama."

(PLATE 21.)

Head 2.7; depth 2.9; eye 5.67; snout 2.25; maxillary 2.8; mandible 2.3; interorbital 4.43; preorbital 3.5; D. x, 14; A. m, 8; scales 10-70-18, about 51 pores. Body rather deep and compressed, back strongly elevated, profile steep and nearly straight from snout to nape; snout long and pointed; eye rather small; interorbital space gently convex; occipital keel moderate; preorbital very broad; mouth moderate, maxillary scarcely reaching front of orbit; upper jaw with a narrow band of villiform teeth, outside of which is a single series of larger but small teeth; 6 rather strong canines in front, 4 of them larger, about equaling in length half of diameter of pupil; lower jaw with a narrow villiform band in front only and a series of larger teeth outside; these unequal, largest on side of jaw, some of them almost canine-like; tongue with a single very small patch of teeth on its middle, this wanting in young examples; teeth on vomer forming a broadly Λ -shaped patch, without backward prolongation on median line. Gillrakers moderate, half length of diameter of eye, about 8 on lower arch, with no rudiments before them. Preopercle with its posterior margin almost straight, slanting gently downward and forward, notch broad and very shallow; edge of preopercle rather coarsely serrate, most so at angle; scales small, rows almost horizontal below lateral line, running backward and upward above; tubes of lateral lines branched; about 7 rows of scales on cheek; 1 row on interopercle, 1 on subopercle, and about 9 on opercle; temporal region with about 8 rows of scales, which become smaller posteriorly; bases of soft dorsal and anal scaly. Dorsal spines weak and slender, outline of fin not greatly curved, fourth spine longest, 2.66 in head, tenth spine 3.33 in head; margin of soft dorsal angulate, ninth ray longest, twice last and 1.5 times first ray, 2 in head; caudal well forked, upper lobe the longer, 1.6 length of middle rays, which are about 2.16 in head; anal angular, similar to soft dorsal, middle rays more elevated than in any other species, longest 2.25 length of last, 2 in head; first ray nearly reaching tip of last when fin is depressed; second and third anal spines rather strong, of equal length, 3.75 in head; ventrals 1.6 in head; pectoral reaching slightly past origin of anal, 1.3 in head.

Color in life: Dark olive-green above; many of the scales with pale-blue spots, these forming irregular oblique streaks upward and backward; similar stripes more regular and numerous on caudal peduncle and above anal. In old fishes these blue spots and streaks disappear; belly white, strongly tinted with brick-red; about 6 narrow, dusky, vertical bars, a little broader than interspaces and not well defined, between gill-opening and anal; head bronze-olive, darker above; a broad, undulating pearly

streak from snout below eye to upper edge of gill-opening; a narrow blue streak from eye to nostrils; iris fiery red; pectoral, caudal, anal, and ventrals brick-red, caudal narrowly margined with black and little bronzed above; dorsal reddish along the rays and tips of membranes, otherwise yellowish; distinct lateral blotch just above lateral line and below first soft ray on dorsal, about as large as pupil, smaller than in other species similarly marked and seldom disappearing with age; axil and bar across base of pectoral above pale or dusky olive. In spirits the markings become fainter, the lateral blotch and bluish streaks on head usually persisting.

The above description of color from a specimen 11 inches long taken at Key West; Porto Rican examples appear to be somewhat more brightly colored. A fine specimen, 14 inches long, taken at Arroyo February 14, was, in life, pale-rosy, richest on sides, pale below; back olive-green with rosy wash; bases of scales brownish, edge with a pale-blue crescent; cheek and opercle rosy; an irregular blue line under eye; some blue in front and behind eye; top of head dark-reddish; chin and belly nearly white; fins all rosy, dorsal edged with lemon; upper caudal lobe yellowish. A specimen 8 inches long from San Antonio Bridge had side with narrow blue stripes and broader yellowish ones; cheek with 2 or 3 pale-blue stripes; belly rosy; ventrals and anal rosy; caudal greenish with slight rosy wash and dark edge; small dark spot under anterior dorsal rays; mouth pale inside.

This species ranges from Pensacola and Key West southward among the West Indies to Brazil, straying northward in the Gulf Stream to Woods Hole. Apparently one of the most abundant and important food-fishes of Porto Rico. Our collections contain specimens from San Juan, Mayaguez, Puerto Real, Ensenada del Boqueron, Ponce, Arroyo, Hucars, Fajardo, Vieques, and Culebra.

At Key West this species is called mutton-fish or pargo, and is one of the most important fishes brought to that market. It is said to reach a maximum weight of 25 pounds, though examples weighing more than 15 or 18 pounds are not common. The average weight of those seen at Key West or in Porto Rico probably did not exceed 5 pounds. It is said to be the most important food-fish of Havana, being always abundant. Its flesh is fairly flavored, though not very delicate, but always healthful. About Key West it is found on rock bottom in 3 to 9 fathoms, and is caught with hook and line. They are quite gamy, taking the hook promptly and fighting well. They are found throughout the year, but are scarcest in July and August, which is their spawning time. They are said to school at spawning time; the eggs are nonadhesive and the size of a rice grain.

In Porto Rico this species is highly esteemed. It is called "sama" or "pargo criollo." It is usually taken in the fish-traps set in 5 to 20 fathoms, though considerable numbers of the smaller individuals are caught with the haul seines in shallow water along the shore.

Mesoprion analis Cuvier & Valenciennes, Hist. Nat. Poiss., II, 452, 1828, Santo Domingo.

Mesoprion sobra Cuvier & Valenciennes, Hist. Nat. Poiss., II, 453, 1828, Martinique.

Mesoprion isodon Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 443, 1833, Santo Domingo.

Mesoprion rosaceus Poey, Ann. Lyc. Nat. Hist. N. Y., IX, 1870, 317, Cuba.

Lutjanus analis, Poey, Fauna Puerto-Riqueña, 330, 1881; Stahl, l. c., 76 and 162, 1883.

Neomænis analis, Jordan & Evermann, l. c., 1265, 1898.

130. *Neomænis megalophthalmus* Evermann & Marsh, new species.

Head 2.67; depth 2.9; eye 4.75; snout 2.85; maxillary 2.6; mandible 2.1; interorbital 6.25; preorbital 4.8; D. x, 12; A. III, 8; scales 7-64-16, about 57 pores. Body oblong, compressed, back greatly elevated, anterior profile very steep and nearly straight or slightly concave from tip of snout to occiput, thence regularly convex to origin of dorsal fin; snout rather long and pointed; mouth large, maxillary reaching anterior third of pupil; lower jaw somewhat projecting; eye large; edge of preopercle nearly vertical, notch long and shallow, serrations obscure; teeth on vomer, palatines, and tongue, those on vomer in an A-shaped patch; patch on tongue long and narrow, truncate in front, narrowing behind; a narrow band of villiform teeth on upper jaw and 6 canines in front, one of these on each side much smaller than others; a row of moderately strong conic teeth in lower jaw; 10 gill-rakers on lower arm of first arch, longest about 2 in eye; scales moderate, those above lateral line in very oblique series; 6 rows each on cheek and opercle; temporal region with a broad band of scales arranged in several series; bases of soft dorsal and anal scaly; caudal well scaled; dorsal spines rather slender and weak, fourth to seventh longest, about 3.25 in head; second and third anal spines equal in length, second stronger, about equal to eye; caudal moderately forked, lobes equal, 1.6 in head; pectoral long, 1.25 in head, reaching origin of anal; ventrals about 2 in head.

Unfortunately, no life-color notes were taken of this species. In alcohol the type is olivaceous or grayish on back, with narrow pale-bluish lines following the rows of scales upward and backward; body below lateral line yellowish-white, with about 3 broad indistinct bluish lines near middle of side; a large black blotch, somewhat smaller than eye, just above lateral line and below first 4 dorsal rays, this spot smaller and less conspicuous than in *N. synagris*; preorbital bluish with traces of 2 narrow bronze lines running forward and downward from eye; fins all pale-lemon; iris yellow.

This species is related to *N. synagris*, but appears to differ in the larger eye, the greatly elevated back, longer pectoral, and in coloration. In *N. synagris* the upper caudal lobe is usually the longer, while in this species they are equal.

Type, No. 49531, U. S. N. M., 11.5 inches long, obtained at Puerto Real, Porto Rico, January 25, 1899, by Evermann & Marsh.




Fig. 48.—*Neomænis megalophthalmus*.

131. *Neomænis synagris* (Linnaeus). “Manchego”; “Manchera”; “Raiado”; Lane Snapper.

(PLATE 22.)

Head 2.65; depth 2.75 to 3; eye 4.5 to 5; snout 2.5 to 2.8; maxillary 2.4 to 2.7; mandible 2 to 2.2; interorbital 5 to 5.75; preorbital 4.5 to 5; D. x, 12; A. iii, 8; scales 7-64 to 68-15, about 50 pores. Body oblong, compressed, the back moderately elevated, profile almost straight from snout to nape; occipital keel little prominent; maxillary reaching front of orbit; upper jaw with a narrow band of villiform teeth, outside of which is a single series of enlarged teeth; 4 (rarely 6) rather small canines in front, 2 of them larger; lower jaw with villiform band in front only, single row of larger teeth nearly equal in size, none of them canines; tongue with a single oval patch, its length more than twice its width; vomer with a Λ or $-$ shaped patch of teeth, without backward prolongation on median line, or with only a very slight one. Gillrakers rather long, their length slightly more than half diameter of eye, about 5 + 9, and usually no rudiments before them. Preopercle with its posterior margin slanting downward and forward, the emargination broad and moderately deep; preopercle rather finely serrate above, with coarser teeth at angle. Scales rather small, rows almost horizontal below lateral line, above somewhat undulate, running upward and backward; tubes of lateral line simple; 6 rows of scales on the cheek, 1 row on interopercle, 1 on subopercle, and 6 on opercle; temporal region with a broad band of scales, arranged in several series; base of soft dorsal and anal scaly; dorsal spines rather weak and slender, outline of fin gently convex; fourth spine longest, 2.66 in head, tenth spine 3.6 in head; soft dorsal short, its margin somewhat angulated, eighth ray longest, twice length of last ray and 1.5 first, 2.4 in head; caudal moderately forked, upper lobe the longer, 1.5 length of middle rays, which are 2 in head; anal rather high, rounded in outline, its middle rays longest, 1.66 length of last ray, 2.57 in head, first ray reaching middle of last ray when fin is depressed; second anal spine stronger than third and of equal length, 3.66 in head; ventrals 1.75 in head; pectoral reaching front of anal, 1.25 in head.

Color in life: Rose-colored, silvery-tinged below, slightly olivaceous but not dark above; a large, round, maroon blotch, larger than eye, just above lateral line and below front of soft dorsal, always present; series of stripes of deep golden-yellow along sides; 3 on head, upper from snout through eye; about 10 on body, lower nearly straight and horizontal, upper undulating and irregular, extending upward and backward; belly white, its sides largely yellowish; lips red; maxillary partly yellow; tongue yellowish; iris fiery red; caudal deep blood-red; spinous dorsal nearly transparent, with a marginal and basal band of golden; soft dorsal light-red, edged with golden; ventrals and anal golden; pectoral pinkish. Young quite green above. Similarly striped Cuban specimens are generally duller, with yellow stripes decidedly coppery. In spirits, the bright colors fade, only the lateral blotch and the streaks on the head being persistent.

Two specimens, each 8 inches long, from the market at San Juan showed the following colors: Bluish-olive on back, becoming paler on side; belly white; narrow irregular bars on back running upward and backward; side with about 7 broad orange or brassy lines, 1 above lateral line, next 2 running into it, others wholly below it; a large round black blotch, as large as eye, between lateral line and front of soft dorsal; dorsal edged with red; caudal reddish. Another specimen, 9 inches long, from San Antonio Bridge was, in life, rich-rosy, with 8 or 9 narrow orange or brassy lines on side, 4 or 5 of which extend on head, fourth from below uniting with fifth just behind opercular flap; dorsal pale-rosy, margined with lemon; caudal rosy, with a narrow black border; anal white on last rays, lemon anteriorly; pectoral very pale-rosy; ventrals pale-lemon; lips somewhat rosy.

The lane snapper is found from southern Florida southward to Colon and Brazil, and is usually abundant throughout its range. On the east coast of Florida it is known from Indian River and Biscayne Bay, and on the west coast from Pensacola southward. It is abundant among the Florida Keys, and is known from the Bahamas, Cuba, Martinique, Jamaica, Santo Domingo, and Porto Rico. Wherever known it is a food-fish of importance. Its maximum weight is about 4 pounds, though the average of those brought to the Key West market is not over half a pound. The largest seen in Porto Rico was 14 inches long and weighed about 2 pounds. Next to *Neomænis apodus*, it is the most abundant snapper in Porto Rico, and was obtained in the markets of San Juan, Mayaguez, and Ponce, our collections containing specimens from the following localities: San Juan, Mayaguez, Puerto Real, Ensenada del Boqueron, Fish Hawk station 6070 in Mayaguez Harbor in 220 fathoms, Ponce, Arroyo, Hucares, Isabel Segunda, and Culebra.

Salpa purpureseens variegata (Lane Snapper) Catesby, Hist. Nat. Carolina, pl. 17, 1743, Bahamas.

Sparus syntagris Linnæus, Syst. Nat., X, 280, 1758, Bahamas; after Catesby.

Sparus vermicularis Bloch & Schneider, Syst. Ichth., 275, 1801, Martinique; on a drawing by Plumier.

Lutjanus aubrieti Desmarest, Prem. Déc. Ichth., 17, pl. 2, 1823, Cuba.

Mesoprion unnotatus Cuvier & Valenciennes, Hist. Nat. Poiss., II, 449, 1828, Santo Domingo; Martinique.

Neomænis syntagris, Jordan & Evermann, l. c., 1270, 1898.

132. *Neomænis mahogoni* (Cuvier & Valenciennes). "*Ojanco*"; *Mahogany Snapper*.

Head 2.5; depth 3; eye 3.75; snout 2.8; maxillary 2.3; mandible 1.9; interorbital 6; preorbital 6.5; D. x, 12; A. III, 8; scales 8-63-14, about 50 pores. Body elongate, strongly compressed, back somewhat elevated, profile almost straight or slightly concave from tip of snout to nape, thence gently convex; snout rather slender and pointed; eye large; interorbital space somewhat convex; mouth large, maxillary reaching front of pupil; lower jaw strongly projecting; upper jaw with a narrow band of villiform teeth, outside of which is a single series of enlarged but comparatively small teeth; 6 canines on front of upper jaw, 2 of them small; lower jaw with a single series of rather uniform small teeth, none of them canine-like; tongue with an oblong patch of teeth, tapering behind, its length more than twice its width; vomer with a broad, arrow-shaped patch of teeth, with backward prolongation on median line. Gillrakers moderate, about 10 developed on lower limb of arch, and 3 or 4 rudimentary ones; preopercle with its posterior margin almost vertical, broadly and rather deeply emarginate, very weakly or scarcely serrate above, angle projecting backward and armed with several rather coarse teeth, lower limb smooth. Scales rather small, those below lateral line somewhat larger, rows above lateral line running obliquely upward and backward, those below in almost straight horizontal series; cheek with 6 rows of scales, 1 row on interopercle, 1 on subopercle, and 7 on opercle; temporal region with a band of small scales, before and behind which is a series of larger ones; top of head, snout, and jaws naked; bases of soft dorsal and anal scaly. Dorsal spines rather weak and slender, outline of fin rather strongly convex, fourth spine longest, 2.57 in head; tenth spine 4 in head; margin of soft dorsal

very gently convex, the first and last rays slightly shorter than rest of fin, median rays 3.33 in head; caudal not deeply forked, upper lobe little longer than lower, its length 1.4 in middle rays, which are 2.12 in head, margin of anal little rounded, the middle rays 1.66 length of last ray, 3 in head, first ray reaching almost to tip of last ray when fin is depressed; anal spines small, second as long as third and stronger, 4.4 in head; ventrals 2.2 in head; pectoral scarcely reaching front of anal, 1.33 in head.

Color in life: Deep brown, silvery below, everywhere shaded with red, especially on head; eye scarlet; a large blackish blotch on side, chiefly above lateral line and below first rays of soft dorsal; maxillary yellow on the covered parts; narrow bronze streaks following rows of scales, these streaks distinct, chiefly above the lateral line; dorsal fin pale, edged with blood-red; caudal deeply red; anal, ventrals, and pectoral scarlet. Bright colors fade and disappear in spirits, leaving the back dark-gray, lower parts silvery, more or less flushed with red.

An inhabitant of the West Indies; known from Cuba, Jamaica, Martinique, and Porto Rico; not known to occur in Florida, but said to be not uncommon in the Havana markets, where it is called "ojanco," in allusion to the large eye. It was seen by us only at Ponce, where a single specimen, 9 inches long, was obtained. This species does not reach a large size.

Mesoprion mahogoni Cuvier & Valenciennes, Hist. Nat. Poiss., II, 447, 1828, Martinique.

Mesoprion ricardi Cuvier & Valenciennes, Hist. Nat. Poiss., II, 447, 1828, Martinique.

Mesoprion ojanco Poey, Memorias, II, 150, pl. 13, fig. 10, 1860, Cuba.

Lutjanus ojanco, Poey, Fauna Puerto-Riqueña, 321, 1881; Stahl, l. c., 76 and 162, 1883.

Neomænis mahogoni, Jordan & Evermann, l. c., 1272, 1898.

Genus 78. OCYURUS Gill. Rabirubias.

This genus is allied to *Neomænis*, from which it differs notably in the structure of the skull, especially in the forward extension of the fronto-occipital crest to the ethmoidal projection; prefrontals with posterior areas short and excavated above and in front. The single species shows numerous minor peculiarities, as the peculiar form of body, the large, well-forked caudal fin, and the small head, as well as an increased number of gillrakers and the presence (in the adult) of pterygoid teeth.

133. *Ocyurus chrysurus* (Bloch). "*Colirubia*"; *Rabirubia*; *Yellow-tail*.

(PLATE 23.)

Head 3; depth 3; D. x, 13; A. n1, 9; scales 7-65-15, 51 pores. Body elliptical, comparatively elongate, back little elevated; profile straight from tip of snout to nape, thence rather strongly arched; caudal peduncle long and slender; snout pointed, of a moderate length, 3 in head; eye small, 5 in head; interorbital space very convex, with sharp median keel, 4 in head; preorbital narrow, its least width 6.66 in head. Mouth small, oblique, lower jaw projecting; maxillary reaching very slightly beyond front of orbit, 2.71 in head; upper jaw with narrow band of villiform teeth, outside of which is a single series of larger teeth, 5 or 6 of those in front being somewhat canine-like, but small; lower jaw with a single series of moderately strong teeth, none of them large enough to be called canines; tongue with a large, oval patch of teeth, in front of which is a smaller but similar patch; teeth on vomer forming a broadly arrow-shaped patch, with backward prolongation on median line, which is nearly twice width of patch; a narrow band of pterygoid teeth behind patch on vomer, this not evident in young examples. Gillrakers rather long and slender, longest about half diameter of eye, about 8 + 21, none of them rudimentary. Preopercle with its posterior margin almost vertical, with a slight but distinct emargination above angle; serrations of the preopercle very feeble, teeth at angle scarcely enlarged; nostrils well separated, posterior slit-like; scales small, those above the lateral line arranged in very oblique series, those below in rows nearly horizontal; cheek with 5 or 6 rows of scales, about 2 rows on interopercle; temporal region with 2 or 3 series of large scales, before and behind which are many small scales; top of head, snout, and jaws naked; bases of soft dorsal and anal scaly. Dorsal spines rather long and slender, fin not deeply emarginate, fifth spine longest, 2.4 in head; tenth spine 3.75; soft dorsal and anal similar, their margins nearly straight, last rays slightly shortened, median rays about 3 in head; caudal fin long, very deeply forked, upper lobe longest, three times as long as middle rays, which are 2.5 in head; pectoral long and slender, reaching vent, 1.12 in head; ventrals 1.57 in head; anal spines rather weak, the third a third longer than second, 4 in head.

Color in life: Olivaceous above, rather pale, and somewhat violet-tinged; a number of large, irregular, deep yellow blotches on sides of back; a deep yellow stripe from tip of snout straight through eye to caudal peduncle, there broadening and including all of tail above lateral line and behind dorsal fin;

above this a pearly-purplish area; below it a flesh-colored or rosy area or band, 2 scales broad, then a succession of about 16 narrow streaks alternately flesh-colored and yellow, growing fainter progressively below; yellow on edges of scales, reddish on their middles; iris fiery-red; lower parts of head flesh-color with some yellow spots; maxillary mostly yellow; caudal deep yellow, its edges reddish; dorsal chiefly yellow; anal faintly yellow; ventrals and pectoral translucent.

In spirits, all markings fade, leaving fins yellowish, upper parts grayish, lower rosy-silvery.

Found from southern Florida to Brazil, generally abundant; known from Biscayne Bay, Key West, Cuba, Martinique, St. Kitts, Jamaica, Porto Rico, and Brazil; at Key West, where it is called "yellow-tail" or "rabirubia," it is even more abundant than the lane snapper, and is the principal fish served at the Key West hotels and boarding-houses in the fall. It is said to be plentiful throughout the year except during the winter, when unusual cold may drive them away. During the warmer weather they are found at a depth of 2 fathoms or more, usually in about 5 fathoms, and generally about shoals where there is some mud bottom. The spawning time in Florida is said to be in July, when they are found about the reefs from Miami to the Tortugas.

In Porto Rico this fish is called "colirubia," and is an abundant and important food species. It was seen by us at most of the places visited, and specimens are in the collection from Mayaguez, San Antonio Bridge, Puerto Real, Ensenada del Boqueron, Ponce, Arroyo, Hucares, Fajardo, Culebra, and Isabel Segunda. Mr. Gray's collection from San Geronimo also contains several.

This species attains a length of about 2 feet and a weight of 3 or 4 pounds or more, and is quite gamy. The average weight of those seen at Key West was probably not over a pound. Those seen in Porto Rico were somewhat larger. At Key West they are caught with hook, using sardine bait.

Acara pitamba, Marcgrave, Hist. Brasil., 155, 1648, Brazil.

Rabirubia, Parra, Descr. Dif. Piezas, Hist. Nat., pl. 20, fig. 1, 1787, Cuba.

Sparus chrysurus Bloch, Ichthyol., pl. 262, 1790, Brazil; after Marcgrave.

Anthias rabirubia Bloch & Schneider, Syst. Ichth., 309, 1801, Cuba; after Parra.

Sparus semiluna Lacépède, Hist. Nat. Poiss., IV, 141, 1803, Martinique; on a copy of a drawing by Plumier.

Mesoprion aurovittatus Agassiz, Spix, Pisc. Brasil., pl. 66, 1829, Brazil.

Ocyurus rijgersmaei Cope, Trans. Am. Phil. Soc. 1871, 468, St. Kitts.

Ocyurus Poey, Fauna Puerto-Riqueña, 321, 1881.

Ocyurus chrysurus, Jordan & Evermann, l. c., 1275, 1898.

Genus 79. RHOMBOPLITES Gill.

This genus is closely allied to *Neomænis*, but cranial peculiarities and extension of villiform teeth over pterygoid and hyoid bones well warrant generic separation. Form of vomerine patch of teeth is also somewhat peculiar. Prefrontals with articular facets developed from simple tubercles and not V-shaped, posterior areas eribriform; basisphenoid not lobigerous; pterygoid with a broad patch of teeth (in adult); hyoid bones and tongue with teeth; canines very small or obsolete; dorsal spines 12, soft rays 10 or 11; gillrakers slender and numerous. But one species known.

134. *Rhomboplites aurubens* (Cuvier & Valenciennes). *Cagon de lo Alto*.

Head 3; depth 3; eye 3.4; snout 3.9; maxillary 2.7; mandible 2.2; interorbital 4; preorbital 8.5; D. XII, 11; A. III, 8; scales 8-68-16, about 50 pores. Body elongate, irregularly elliptical, the back not greatly elevated, highest at nape; profile regularly and strongly convex from above eye to spinous dorsal; snout rather short and bluntish, its upper profile straight and steep; eye very large; interorbital space very convex; preorbital narrow; mouth small, oblique, lower jaw somewhat projecting; maxillary scaleless, reaching front of orbit; upper jaw with a broad band of villiform teeth, outside of which is a row of enlarged, but comparatively small teeth; no canines; lower jaw with one series somewhat stronger than outer teeth of upper jaw; inside of these is a rather broad villiform band of teeth in front of jaw only; tongue with a very broad irregularly ovate patch of teeth, its width almost as great as width of tongue, 1.5 in its length; in front of this patch is a large roundish patch of teeth; an oblong patch of teeth on hyoid bone; vomer with a rhomboid (◇-shaped) patch of teeth, forming almost a right angle in front, with a broadly wedge-shaped backward prolongation on median line, its length about twice its width; palatine band of teeth very wide; pterygoids with a large patch of teeth, these teeth undeveloped and covered by skin in young examples. Gillrakers numerous, longest about half diameter of eye, about 6 + 21. Preopercle with posterior margin almost straight and vertical, slightly emarginate, weakly serrate above, teeth coarser at angle and on lower border; posterior nostril larger, nearly round.

Scales very small, rows above lateral line running upward and backward, rows below rather wavy, almost horizontal; temporal region covered with small partially embedded scales, in 4 or 5 rows; cheek with 7 rows of scales; 4 rows on interopercle, 3 rows on subopercle, and 7 on opercle; snout, preorbital, and jaws naked; top of head scaly to near middle of eye; soft dorsal and anal with but few scales at base. Dorsal spines long and slender, fourth spine longest, 2.9 in head, length of spines thence gradually decreasing to twelfth spine, which is 3.33 in head; margin of soft dorsal truncate, its rays of subequal length, 4.5 in head; last ray slightly shorter; caudal deeply forked, upper lobe longer than lower, its length 1.75 times middle rays, which are 2 in head; upper lobe of caudal 1.25 in head; anal similar to soft dorsal, its rays 3.4 in head; second anal spine shorter than third, 5 in head; ventrals 2 in head; pectoral somewhat falcate, reaching opposite vent, 1.4 in head.

Color in life: Vermilion, paler below; faint brown lines running obliquely forward and downward from dorsal along rows of scales; side with narrow sinuous streaks of golden-yellow, some of them longitudinal, others oblique; dorsal rosy, its margin chiefly orange; anal pale at base, rosy at extremity; pectoral yellowish, ventrals rosy, caudal and iris vermilion; inside of mouth dusky.

The bright colors grow faint or disappear in spirits.




FIG. 49.—*Rhomboplites aurorubens*.

Charleston and Pensacola, south to Brazil; known from Pensacola, the "Snapper Banks," Cuba, Martinique, Santo Domingo, Jamaica, Porto Rico, and Brazil. Only a single specimen, 6 inches long, was obtained in Porto Rico. The species reaches a length of a foot and is a good food-fish.

Centropristis aurorubens Cuvier & Valenciennes, Hist. Nat. Poiss., III, 45, 1829, Brazil; Martinique; Santo Domingo.
Mesoprion elegans Poey, Memorias, II, 153, 1860, Cuba.
Aprion arionomus Jordan & Gilbert, Proc. U. S. N. M., 1883, 142, Pensacola; young with pterygoid teeth undeveloped.
Rhomboplites aurorubens, Jordan & Evermann, l. c., 1277, 1898.

Genus 80. ETELIS Cuvier & Valenciennes.

Body elongate, covered with large scales; eye very large; preorbital very narrow; mouth moderate, lower jaw projecting; canines in upper jaw only; no teeth on tongue or pterygoids; gillrakers long and slender. Dorsal fin deeply notched, rather short, its spines 10 in number, its soft rays not scaly; caudal very deeply forked; head naked above, skull with interorbital area flat, separated from occipital area by a transverse line, limiting median and lateral crests also; frontals wide in front, not cavernous, simply normally perforate; supraorbital margins crenate; periotic region little convex and with bones thick, unpolished; prefrontals behind, with funnel-shaped foramina.

The relationships of this interesting genus have been repeatedly misunderstood, but it belongs in the *Lutianidae* and it has no special affinity with *Anthias*, *Perca*, or *Serranus*. Its synonymy and relations have been well discussed by Dr. Gill. In spite of the difference in form of dorsal, its relations with *Aprion* are very close. The skulls in the two are almost identical, as has been pointed out by Gill and by Poey. Two American species are known.

a. Maxillary scaly; depth 3.5 or more in length..... *oculatus*, 135
 aa. Maxillary naked; depth less than 3 in length..... *aquilonaris*

135. *Etelis oculatus* (Cuvier & Valenciennes). "*Cachucho*."

Head 3.5; depth 3.8; eye 3.4; snout 3.1; maxillary 2.1; mandible 1.7; interorbital 3.8; D. x, 11; A. III, 8; pectoral 1.1; ventral 1.5; caudal 0.8; scales 5-51-11. Body elongate, fusiform, not greatly compressed, caudal peduncle rather long and slender; scales large, regular, minutely ctenoid; top of head above eyes broad and flat, naked, skin somewhat rugose; top of head back of eyes with 2 lateral patches of scales, separated from those of body by a groove; cheek and opercles scaly, save preopercle, which is naked, its edge finely toothed; opercle ending in 2 weak, flat spines; eye very large; lower jaw projecting, maxillary reaching past front of pupil; teeth of upper jaw in a villiform band with an outer enlarged row and 1 or 2 pairs of strong forward-pointing canines in front; lower jaw with a row of conical teeth on sides, villiform ones in front, without distinct canines; vomer with villiform teeth, palatines with a row of stronger conical ones; ventral accessory scale not well developed; dorsal deeply notched; caudal scaly, widely forked.

Color: In life, bright-red nearly everywhere; in spirits, pale, the red nowhere persisting.

A handsome, edible species. Length 2 to 3 feet. West Indies to Madeira; not yet known from Florida; in rather deep water; generally common on rocky bottom.

One specimen, nearly 2 feet long, taken at Mayaguez.

Serranus oculatus Cuvier & Valenciennes, Hist. Nat. Poiss., II, 266, 1828, Martinique.
Etelis oculatus, Jordan & Evermann, l. c., 1282, 1898.




FIG. 50.—*Etelis oculatus*.

Family XLV. HEMULIDÆ. The Grunters.

Body oblong, or more or less elevated, covered with moderate-sized, adherent scales, which are more or less strongly ctenoid or almost cycloid; lateral line well developed, concurrent with back, usually not extending on caudal fin; head large, crests on skull usually largely developed; no suborbital stay; mouth large or small, usually terminal, low, and horizontal; premaxillaries protractile, their spines not greatly produced backward; maxillary without supplemental bone, for most of its length slipping under edge of preorbital, which forms a more or less distinct sheath; preorbital usually broad; no barbels; teeth all pointed, none of them forming marked canines; no teeth on vomer, palatines, and tongue; lower pharyngeals separate, with pointed teeth; gills 4, a large slit behind fourth; pseudo-branchiæ large; gillrakers moderate; gill-membranes separate, free from isthmus; preopercle serrate or entire; opercle without spines; sides of head usually scaly; dorsal fin single, continuous or deeply notched, sometimes divided into 2 fins, spines usually strong, depressible in a groove, heteracanthous, that is, alternating, one stronger on right side, other on left, usually 10 to 12 in number; anal fin similar to soft dorsal, with 3 spines; ventral fins thoracic, rays 1, 5, with a more or less distinct scale-like appendage at base; caudal fin usually more or less concave behind; air-bladder present, usually simple; stomach cæcal; pyloric cæca few; vertebrae usually 10 + 14 = 24. Branchiostegals usually 6 or 7. Cranium with its muciferous system moderately developed or rudimentary. Intestinal canal short.

Carnivorous fishes of the warm seas, most of them valued as food. The family includes about 15 genera and nearly 150 species. It is very close to the *Lutianidæ* on the one hand and to the *Sparidæ* on the other, while some of its members show affinities with some *Sciænidæ* and *Serranidæ*.

- a. Chin with a central groove behind the symphysis of the lower jaw.
- b. Mouth more or less wide, the jaws scarlet posteriorly in life; soft parts of vertical fins densely scaly to their margins.
- c. Scales above lateral line arranged in very oblique series, not parallel with the lateral line.
- d. Jaws subequal, or the lower included; mouth little oblique; gillrakers comparatively few and short.
- e. Dorsal spines 12, rarely 11; scales large; gillrakers few and small (10 to 14 on lower part of anterior arch); frontal foramen a single or divided slit at the base of the high supraoccipital crest in front.
- f. Mouth moderate or large, its cleft more than one-third length of head; back more or less elevated; second anal spine strong, notably longer than third..... HÆMULON, 81
- ff. Mouth small, its cleft less than one-third length of head; body rather elongate; second anal spine small; back and sides with longitudinal yellow stripes; teeth weak; gillrakers rather few and small; snout very short, 2.66 in head; frontal foramina separate and placed some distance in front of the very low supraoccipital crest; premaxillary spine very short, 4.25 in head..... BRACHYGENYS
- cc. Dorsal spines 13; anal fin low; preorbital low; gillrakers in moderate or rather large numbers, 12 to 18 on lower part of arch; lower jaw not projecting; mouth little oblique; body comparatively elongate, the depth 2.75 to 3.5 in length; body with longitudinal yellowish stripes; scales rather small; frontal foramina long divided slits in front of supraoccipital crest; size small..... BATHYSTOMA, 82
- bb. Mouth more or less narrow, not scarlet within; soft fins naked or with scales on their basal parts.
- g. Anal fin short, its rays III, 7, to III, 10; dorsal fin more or less emarginate, its spines rather robust.
- h. Body ovate, back elevated; depth greater than length of head; outer teeth of upper jaw enlarged; lips thick; second anal spine strong; soft rays of dorsal and anal scaly at base..... ANISOTREMUS, 83
- hh. Body oblong, depth usually less than length of head; lips not very thick; scales large, those above lateral line in series mostly parallel with lateral line.
- i. Preopercle very sharply serrate, serræ at angle much enlarged, those below angle turned forward; outer teeth in both jaws considerably enlarged; soft rays of dorsal and anal more or less scaly; second anal spine enlarged..... CONODON, 84
- ii. Preopercle finely serrate, serræ at angle scarcely enlarged, those below not antrorse; teeth subequal, or outer in upper jaw somewhat enlarged; gillrakers very short and weak.
- j. Soft part of dorsal and anal with series of small scales on membranes behind each ray; anal spines small or moderate, second little, if any, longer or stronger than third; body oblong, not elevated; scales above lateral line parallel with back; dorsal spines 12, soft rays 15 or 16; outer teeth of upper jaw slightly enlarged..... BRACHYDEUTERUS, 85
- jj. Soft parts of dorsal and anal scaleless, except a low sheath at base; anal spines strong, second much longer and stronger than third; dorsal spines 12 or 13, the soft rays 11 to 14..... POMADASIS, 86
- gg. Anal fin long and low, its rays III, 10, to III, 13; dorsal fin low, usually not deeply emarginate; anal spines small; preopercle finely serrate or entire; outer teeth of jaws slightly enlarged; gillrakers moderate, rather slender.
- k. Dorsal spines 12 or 13.
- l. Scales of body without series of small accessory scales at base; soft dorsal and anal naked or somewhat scaly; mouth small; temporal crest, which rises from behind eye, very low and inconspicuous, upper edge below base of high supraoccipital crest, which originates over pupil..... ORTHOPRISTIS.
- aa. Chin with pores, but with no central groove at symphysis; preopercle finely serrate.
- m. Anterior profile concave above eye; snout gibbous; outer teeth in both jaws enlarged and blunt (appearance of *Anisotremus*); gillrakers small and slender; anal fin rather long, soft dorsal and anal scaleless..... GENYATREMUS.

Genus 81. HÆMULON Cuvier. The Grunts.

Body oblong, usually more or less elevated; mouth wide, maxillary long and curved, reaching to below eye, its tip extending to posterior end of preorbital; chin with a central groove behind symphysis; lower jaw included; gillrakers moderate; no teeth on vomer or palatines; teeth of jaws conical, outer series stronger, curved; lips and inside of mouth posteriorly commonly bright-red or scarlet in life; preopercle serrate, with no recurved hooks below; soft part of vertical fins completely covered with scales; scales above lateral line in series not parallel with it; a marked angle formed at junction of spinous and soft parts of dorsal; dorsal spines 12 or 11; second anal spine enlarged, generally larger and longer than third; caudal forked.

There are 12 or 13 species of *Hæmulon*, all of them American and all important food-fishes. They are of fair size, the flesh is firm and sweet, and they find a ready sale.

All the species have more or less of orange on inside of mouth, a trait of coloration not found in *Pomadasis*. The amount of redness is greatest in those species having the largest mouth. The young differ in proportion considerably from the adults. Besides the changes usual in other fishes, we may observe that in *Hæmulon* the young have the snout proportionately much shorter, so that the maxillary, although also much shorter in proportion, extends further back in comparison with the eye. Nearly all the species have, when young, at least two more or less sharply defined, dark, longitudinal stripes along the side, one or more along the top of the head, and a dark spot at the base of caudal. These markings persist longer in some species than in others, but traces of them at least may be found in the young of nearly all the species of *Hæmulon* and *Pomadasis*. In a few species these markings persist during life.

- a. Scales below lateral line anteriorly not especially enlarged.
 b. Scales above lateral line anteriorly not much enlarged.
 c. Maxillary 2.33 to 2.75 in head, not reaching center of eye (in adult).
 d. Back and sides without yellow or blue stripes; each scale above with a median blackish spot, these forming undulating lines (spots rarely obsolete in adult, obscure or wanting in young): maxillary 2.5 to 2.75 in head.
 e. Scales in a vertical row from first dorsal spine to lateral line, 7 or 8 (9 in oblique series).
 f. Mouth rather small, maxillary scarcely reaching to front of eye; back elevated; preorbital very deep, its least breadth greater than length of eye in adult, 4.12 to 4.66 in head in young; second anal spine not reaching to tip of last ray; snout long and pointed, 2.25 to 2.75 in head.
 g. Sides without dark bars; head unspotted; dorsal spines graduated; second anal spine, when depressed, reaching beyond tip of last spine..... *album*, 136
 ff. Mouth rather large, maxillary reaching front of pupil; back little elevated; preorbital rather narrow, its least width 5 in head; second anal spine reaching tip of last ray; snout rather long and pointed, 2.33 in head; back and sides with 4 or 5 black longitudinal streaks, which disappear only in very old examples *macrostomum*, 137
 ee. Scales in a vertical row from first dorsal spine to lateral line, 5 or 6.
 h. Series of scales from scapular scale extending backward to front of soft dorsal; snout rather long and pointed; mouth small; maxillary 2.75 to 3 in head; pectoral fin long, three-fourths length of head; black spots on sides coalescing in continuous stripes *bonariense*, 138
 hh. Series of scales from scapular scale not extending farther backward than the middle of spinous dorsal; snout shorter, not very acute; mouth larger, the maxillary about 2.5 in head; premaxillary processes about 3 in head; dark spots on scales not coalescent.
 i. Depth of body about 2.66 in length; pectoral fin short, less than two-thirds length of head; scales above lateral line scarcely enlarged. Head 3; depth 2.66; scales 6-50-14; D. XII, 17; A. III, 7..... *parra*, 139
 dd. Back and sides with distinct horizontal yellow stripes, fading but not disappearing in spirits; no black spots anywhere; maxillary 2.33 in head, reaching front of pupil *carbonarium*, 140
 cc. Maxillary nearly or quite half length of head, reaching center of eye in adult; no black spots or stripes anywhere in adult (except under angle of preopercle).
 j. Back and sides with continuous yellow stripes, which are horizontal and do not everywhere follow the direction of the rows of scales; ground-color bluish-gray; back with a well-defined blackish area from first dorsal spine to base of caudal, this color covering most of soft dorsal and middle of caudal lobes; body rather elongate; snout moderate; second anal spine 2.75 in head. Head 3..... *melanurum*
 jj. Back and sides of head and body with continuous blue stripes, horizontal and not everywhere following rows of scales; ground-color bright-yellow; second anal spine 2.5 in head. Head 2.75..... *sciurus*, 141
 bb. Scales above lateral line anteriorly much larger than other scales..... *plumieri*, 142
 aa. Scales below lateral line anteriorly much enlarged; head, back, and sides with continuous bright-yellow stripes, those below following direction of scales, and therefore extremely undulating for most part.. *flavolineatum*, 143

136. *Hæmulon album* Cuvier & Valenciennes. "Vieja"; Jallao; Margate-fish; Margaret Grunt.

(PLATE 24.)

Head 2.7; depth 2.5; eye 6; snout 2.3; maxillary 2.5; interorbital 4; preorbital 4.4; scales 8-51-16; D. XII, 16; A. III, 8; pectoral 1.6; ventral 1.8. Body comparatively deep, back elevated and compressed; anterior profile nearly straight, very slightly concave above eye; snout long and pointed; maxillary reaching front of eye; lower jaw included; teeth not large, those on sides of lower jaw more numerous and weaker than in *H. sciurus* or *H. plumieri*; interorbital space convex; preopercle finely serrate above angle; scales moderate, none of them much enlarged, in oblique series above lateral line, nearly horizontal below; soft fins scaly; dorsal spines moderate, fourth highest, 2.4 in head; caudal lobes subequal, 1.4 in head; second anal spine stronger and slightly longer than third, 3 in head.

Color in life, of adults: Pearly-white or rich olivaceous, especially above, where a few scales have very faint, dark spots at their bases; still fainter spots visible along scales of lower part of side; mouth orange within; lips and a faint blotch on each side of snout light yellow; a dusky shade under edge of preopercle (much more distinct in young); fins all light olive; soft dorsal somewhat dusky; head without stripes or spots. Young more distinctly spotted, spots small, round, blackish, each with a pearly edge; one under each scale of back and sides very distinct when the fish is alive, or after its scales are removed, but disappearing almost entirely with death. In life, a broad, dusky lateral band is also present, disappearing with death.

Found from Florida Keys to Brazil; known from Key West, Bahamas, Havana, Jamaica, Porto Rico, and St. Thomas; specimens obtained by us at Isabel Segunda.

The margate-fish is a common and important species at Key West. It reaches a weight of 8 or 10 pounds, the average being about 4 to 6 pounds. It is found in deep water, most abundantly on the reefs. It spawns early in summer, probably in July, on rock bottom, at which time it is

said to school. One intelligent fisherman says, however, that it does not school at Key West, though it does in the Bahamas. It is generally found on rock or barry bottom, around shoals. At night it comes into more shallow water to feed—crabs, crawfish, worms, etc., constituting the bulk of its food. The bait used for it is crawfish or crabs. Cold is said not to affect this fish to any great extent. It does not appear to be very common about Porto Rico, but is highly esteemed. The largest obtained by us was 8.75 inches long.

The origin of the common name of the "margate-fish" is not generally understood. It appears, however, to have been derived from Margate, a well-known seaport and watering-place in England. Some of the fishermen of the Bahamas came originally from Margate and applied the name to one of the fishes which they found in the Bahamas. Many of these fishermen (Conchs) have come to Key West and brought the name with them. The name Margate is, at Key West, sometimes corrupted into "Margat" and "Margaret," while in Biscayne Bay it is "Margat," "Market," or "Margarite."

Perca marina gibbosa (Margate-fish), Catesby, Nat. Hist. Carolinas, etc., 2, pl. 2, 1742, Bahamas.

Hæmulon album Cuvier & Valenciennes, Hist. Nat. Poiss., V, 241, 1830, St. Thomas; Jordan & Evermann, l. c., 1296, 1898.




FIG. 51.—*Hæmulon macrostomum*.

137. *Hæmulon macrostomum* Günther. *Gray Grunt; Striped Grunt; "Corocoro."*

Head 2.75 to 2.8; depth 2.9; eye 4.33 to 5; snout 2.33; maxillary 2; interorbital 3.6; preorbital about 5; D. xii, 16; A. iii, 8; scales 9-53-12; pectoral 1.4 in head; ventral 1.75 to 1.8. Body deep, compressed, back elevated in adult, profile nearly straight from tip of snout to occiput, thence gently curved to origin of dorsal; ventral outline straight; snout long and pointed, jaws subequal; mouth large, nearly horizontal, maxillary reaching middle of pupil. Teeth moderate, outer row in upper jaw and posterior teeth in both jaws considerably enlarged; preopercle finely serrate; gillrakers moderate; scales moderate, those above lateral line not enlarged, those below slightly enlarged on anterior part of body; scales above arranged in very oblique series, those below oblique anteriorly, becoming horizontal posteriorly. Dorsal spines strong, longest about 2.4 in head; soft dorsal high, longest rays 3 in head; caudal lobes subequal, lobes 1.5 in head; anal spines strong, second longest and strongest, 2.8 in head, not reaching past tip of last soft rays when depressed; soft anal high, its free edge concave, longest rays 2.5 in head and reaching beyond tips of last rays when depressed; caudal and soft parts of dorsal and anal densely and finely scaled; pectorals and ventrals with a few fine scales.

Color in life: Body dirty-silvery with about nine dark longitudinal streaks, plainest in young; a median stripe from snout to origin of dorsal; first and second lateral stripes extend from above eye to posterior end of soft dorsal, third begins on upper rim of orbit and extends to vertical of posterior end of soft dorsal where it joins fifth line; fourth line, which is usually indistinct, begins at eye, extends

across opercle, and finally disappears on middle of side; fifth extends from eye along middle of side, crossing lateral line, where it is joined by third, then extending on caudal peduncle above lateral line to base of caudal fin; the remaining lines are on lower part of side and are more or less broken and irregular; head dark grayish-purple; an inky-black spot on inner lower edge of opercle; lower jaw flesh-color, with numerous fine dark specks; dorsal, caudal, anal, and pectoral yellow, with dusky wash on bases; ventrals dark; inside of mouth flesh-color. In larger individuals the dark streaks persist, but are less distinct.

Found in the waters of southern Florida and West Indies; known from Clearwater Harbor, Key West, Florida Keys, Indian River Inlet, Jamaica, St. Thomas, and Porto Rico. Probably not common in Porto Rico, as we obtained specimens only at San Juan and Arroyo. The largest of these was 10.5 inches long. The life-colors in the above description were taken from a specimen 6.5 inches long, obtained at Arroyo.

Hæmulon macrostoma Günther, Cat., I, 308, 1859, Jamaica.

Hæmulon fremebundum Goode & Bean, Proc. U. S. N. M. 1879, 340, Clearwater Harbor, Florida.

Hæmulon macrostomum, Jordan & Evermann, l. c., 1296, 1898.

138. *Hæmulon bonariense* Cuvier & Valenciennes. "Ronco Prieto"; "Arrayado"; Black Grunt.

Head 2.8; depth 2.6; scales 5-44-10; D. XII, 16; A. III, 8. Body oblong, compressed, the back considerably elevated; head rather long; snout pointed, rather longer and sharper than in *H. parra*, anterior profile straight, or a little concave before eyes. Snout 2.66 in head (in young of 9 inches). Mouth rather small, smaller than in *H. parra*, maxillary barely reaching front of eye, its length 3 in head. Teeth of moderate size, outer and posterior somewhat enlarged. Eye moderate, 4.66 in head; interorbital space flattish, its width 4.25 in head; preorbital moderate, its least width 4.8 in head; preopercle moderately serrate. Gillrakers few and small, about 12 on lower part of arch. Scales larger than in *H. parra* or any other of the genus, those above and below lateral line about equal in size, those above arranged in series which are less oblique and more undulating than in related species, the series from scapular scale following direction of lateral line for about 10 scales, then turning abruptly, reaching base of last dorsal spine, or sometimes anterior part of soft dorsal; soft fins scaly as usual. Dorsal spines of moderate strength, fourth 2.6 in head; longest ray of soft dorsal 4 in head; caudal 1.66 in head; anal high, second spine and longest rays extending, when depressed, well beyond tip of last ray; longest soft ray 2.75 in head; second spine longer and stronger than third, 2.66 in head; pectorals long, 1.33 in head; ventrals 1.75.

Color in spirits, pearly-gray; center of each scale brownish-black, these coalescing and forming very sharply defined continuous undulating stripes; about 16 of these between front of dorsal and front of anal; sixth extending from scapular scale to last dorsal spine; base of caudal blackish; fins dusky.

This species is found in the West Indies south to Buenos Ayres; not very common. It is not common about Porto Rico, only 2 specimens being obtained, both in San Juan market. It reaches the length of a foot or less, and is of some value as a food-fish.

Hæmulon canna Cuvier & Valenciennes, Hist. Nat. Poiss., V, 233, 1830, Martinique; not of Agassiz, 1829.

Hæmulon bonariense Cuv. & Val., Hist. Nat. Poiss., V, 254, 1830, Buenos Ayres; Jordan & Evermann, l. c., 1297, 1898

Hæmulon notatum Poey, Memorias, II, 179, 1860, Cuba.

Hæmulon retrocurrans Poey, Repertorio, II, 236, 1868, Cuba.

Hæmulon continuum Poey, Enum., 46, 1875, Cuba; Poey, Fauna Puerto-Riqueña, 325, 1881; Stahl, l. c., 77 and 163, 1883.

139. *Hæmulon parra* (Desmarest). "Arrayado"; "Ronco"; Sailor's Choice.

Head 3; depth 3; eye 4.4; snout 2.4; interorbital 4; preorbital 4.8; D. XII, 17; A. III, 8; scales 6-54-12; pectoral 1.6; ventral 1.8. Body comparatively deep, back compressed and arched; anterior profile rather steep and convex, nearly straight from tip of snout to opposite front of eye, from which point to origin of dorsal fin, it is gently and regularly arched; back from origin of dorsal to caudal peduncle a uniform long curve; ventral line of body from tip of lower jaw to anal fin nearly straight; caudal peduncle compressed, its narrowest width 3 in its least depth. Mouth moderate but slightly oblique, gape 3 in head; lower jaw slightly shorter; maxillary reaching somewhat past front of eye. Teeth small, in villiform bands, those of outer series on sides of lower jaw somewhat enlarged; no antrorse teeth. Interorbital space convex; preopercle scarcely roughened, wholly without serra-

tions; gillrakers short and not very stout, about 15 on lower arm of arch. Squamation very complete, all soft fins densely and finely scaled; scales of body rather large, those above lateral line somewhat enlarged anteriorly, running in very oblique series, series from humeral scale terminating under eighth dorsal spine; scales below lateral line in more nearly horizontal series, those on anterior part of body somewhat enlarged; lateral line arched, approximately parallel to dorsal outline, but very gently approaching it posteriorly. Dorsal spines strong, fourth to sixth longest, fin, when depressed, being inclosed in a groove by the sheathing scales; first anal spine quite small, second large and strong, much stronger but scarcely longer than third, their points not reaching tips of last soft rays when depressed; free end of soft anal slightly emarginate; caudal lobes subequal, their length about 1.66 in head; pectoral rather short and broad, reaching slightly past tips of ventrals; ventrals short, their tips acutely rounded.

Color in spirits, dark-brown; center of each scale on upper parts of body dark-brown surrounded by silvery, free edge of scale paler or purplish-brown; lower part of side and under parts more silvery, but with numerous fine dark punctulations everywhere; head dark; fins all dark.

The range of this fish extends from southern Florida through the West Indies to Brazil; it is recorded from the Tortugas, Key West, Cards Sound, Marco, Lemon Bay, and Biscayne Bay in Florida, and from Havana, Jamaica, Porto Rico, and Brazil. It is abundant about Key West. It collects into schools in July and August, at which time it spawns on rocky bottom. It reaches a weight of 2 pounds,




FIG. 52.—*Hæmulon parra*.

the average being about half a pound, and is a valuable food-fish. The best fishing for this species is in summer. The single specimen obtained in Porto Rico is from Puerto Real, and is 10.5 inches long. It is probably not uncommon at the west end of the island.

Diabasis parra Desmarest, Prem. Déc. Ichth., 30, pl. 2, fig. 2, 1823, Havana.

Hæmulon caudimacula Cuvier, Règne Animal, ed. 2, vol. 2, 176, 1829, Brazil; Havana.

Hæmulon chromis Broussonet MS. in Cuvier & Valenciennes, Hist. Nat. Poiss., V, 242, 1830, Jamaica.

Hæmulon acutum Poey, Memorias, II, 180, 1860, Cuba.

Hæmulon serratum Poey, Memorias, II, 181, 1860, Cuba; Poey, Fauna Puerto-Riqueña, 325, 1881; Stahl, l. c., 77 and 163, 1883.

Hæmulon albidum Poey, Memorias, II, 181, 1860, Cuba.

Hæmulon parra, Jordan & Evermann, l. c., 1297, 1898.

140. *Hæmulon carbonarium* Poey. *Ronco Carbonero*.

Head 3; depth 2.8; eye large, 3.66 in head; D. XII, 16; A. III, 8; scales 7-55-14. Body oblong; back not greatly elevated, profile nearly straight or slightly convex from tip of snout to above eye, thence gibbous to front of dorsal; snout short, moderately pointed, its length 3.16 in head; mouth not very large; gape somewhat curved; maxillary extending nearly or quite to front of pupil, its length 2.5 in head; lower jaw rather included. Teeth strong, much as in *H. sciurus*, but a little shorter. Interorbital space flattish, 4 in head; preorbital moderate, its least breadth 6 in head; pre-

orbital finely but rather sharply serrate; gillrakers small, 9+14. Scales moderate, those below lateral line anteriorly moderately enlarged, their series nearly horizontal; series above lateral line very oblique. Dorsal spines slender and high, the fourth 1.87 in head; longest soft rays 3.5; upper caudal lobe a little longer than lower, 1.2 in head; longest anal rays 2.2 in head, their tips when depressed reaching beyond tip of last ray; second anal spine strong, 2 in head, its tip reaching when depressed about to tip of last soft ray; ventrals 1.5 in head; pectoral 1.3.

Color in life, light bluish-gray, much as in *H. plumieri*; body with 7 or 8 deep brassy-yellow stripes which are horizontal above, those below lateral line a little curved, following the rows of scales; stripes narrower than interspaces of ground-color; 3 stripes above lateral line, 3 or 4 below, the latter paler; little black under angle of preopercle; caudal blackish-yellowish at tip; soft dorsal, anal, and ventrals yellowish-gray, distal portion blackish; spinous dorsal bluish, deep yellow at base and edge; yellowish stripe along middle of fin; pectoral plain, yellowish bar across its base; mouth deep red, its angle dusky. In spirits, grayish, more or less shaded with dusky, the stripes rather faint orange-brown.

West Indies and the Bermudas, south to Brazil; very common at Hayana. Length about 10 inches. One 8 inches long in Mr. Gray's collection from San Geronimo; not seen by us in Porto Rico.

Hæmulon carbonarium Poey. Memorias, II, 176, 1860, Cuba; Jordan & Evermann, l. c., 1300, 1898.




FIG. 53.—*Hæmulon sciurus*.

141. *Hæmulon sciurus* (Shaw). "*Ronco Amarillo*"; *Boca Colorado*; "*Cachicata*"; *Yellow Grunt*.

Head 2.66 to 2.83; depth 2.83 to 3; eye 4 to 4.75; snout 2.25 to 2.33; maxillary 2 to 2.1; interorbital 3.5 to 4.33; preorbital 6; D. XII, 16; A. III, 8; scales 9-50 to 53-13 or 14; pectoral 1.5 to 1.43; ventral 1.75 to 1.86. Body oblong, back regularly and gently elevated, ventral outline nearly straight; snout rather long and pointed; mouth large, gape curved, maxillary usually reaching to vertical of front of pupil; lower jaw slightly included; teeth strong, upper jaw in front with about 3 strong canines on each side; front teeth of lower jaw rather strong; antrorse teeth of back part of each jaw rather strong. Scales moderate, those above lateral line not enlarged anteriorly, arranged in oblique series, those below slightly larger anteriorly, arranged in nearly horizontal series; lateral line gently arched in front. Dorsal spines moderate, fourth longest, about 2.5 in head; longest ray of soft dorsal about 4 in head; second anal spine strongest and somewhat longer than third, but not reaching tips of last soft rays when depressed; upper caudal lobe somewhat the longer; soft parts of dorsal, anal, and caudal densely scaled; pectoral and ventral less fully scaled; preopercle finely serrate; gillrakers short and slender, about 17 below angle of first arch.

General color in life: Yellowish, side with about 10 broad, brassy bands alternating with somewhat narrower pale-blue bands, fourth of which runs forward across upper edge of orbit, crosses forehead and joins its fellow from other side; cheek and snout with similar blue lines, the one on

middle of cheek forking below eye and inclosing an oblong area of ground-color; spinous dorsal pale-yellowish olive, bordered with orange; soft dorsal rusty-olivaceous, with orange border; caudal dusky at base, yellowish-olive on outer third; anal, greenish-yellow; pectoral and ventral, rusty-lemon or light-yellow; inside of mouth, except tips of jaws, blood-red.

The yellow grunt reaches a length of 18 inches and a weight of a pound or less and is a common and important food-fish from the Florida Keys to Brazil. It appears to be generally abundant about Porto Rico and was observed at most of the places visited. There are specimens in the collection from San Juan, Arroyo, Vieques Island, Culebra Island, Fajardo, Guanica, Puerto Real, and Ensenada del Boqueron; taken by Mr. Gray at San Geronimo. It has previously been recorded from Biscayne Bay, the Tortugas, Key West, and others of the Florida Keys, Cuba, Aspinwall, Bahia, and Jamaica.

At Key West this fish is known as "boar grunt" and is very plentiful, usually in schools on rock bottom. A Key West fisherman reports that he has often caught 500 to 600 in a single day; the best fishing is in August. The best bait is a long worm which the fishermen get from the stem of a tall grass which grows on the bars. Nothing could be learned concerning its spawning habits, except that it probably spawns in August. Cold is said to affect this species seriously.

Anthias formosus Bloch, Ichth., pl. 323, 1790, Antilles; not *Perca formosa* Linnæus, with which it has been identified; the latter is *Diplectrum formosum*.

Sparus sciurus Shaw, General Zoology, IV, pl. 64, 1803, Antilles; based on the description and figure of Bloch.

Hæmulon elegans Cuvier, Règne Animal, ed. 2, vol. 2, 175, 1829; no description; based on Bloch's figure.

? *Diabasis obliquatus* Bennett, Zool. Journ. London, V, 1835, 90, Jamaica.

? *Hæmulon similis* Castelnau, Anim. Nou. et Rares, II, 1855, Bahia.

Hæmulon luteum Poey, Memorias, II, 174, 1860, Cuba; Poey, Fauna Puerto-Riqueña, 325, 1881; Stahl, l. c., 77 and 163, 1883.

Hæmulon multilincatum Poey, Memorias, II, 178, 1860, Cuba; Stahl, l. c., 163, 1883.

Hæmulon hians Haly, Ann. Nat. Hist., XV, 1875, 268, Aspinwall.

Hæmulon sciurus, Jordan & Evermann, l. c., 1303, 1898.




FIG. 54.—*Hæmulon plumieri*.

142. *Hæmulon plumieri* (Lacépède).

"*Cachicata*"; "*Boca Colorado*"; *Ronco Ronco*; *Ronco Arará*; *Common Grunt*.

Head 2.5 to 2.8; depth 2.43 to 2.66; eye 4.5 to 5; snout 2.16; maxillary 2; interorbital 3.2 to 4; preorbital 5 to 5.8; D. XII, 15 or 16; A. III, 8 or 9; scales 6-51-17; pectoral 1.5 to 1.8; ventral 1.75 to 1.8. Body moderately elongate, back considerably elevated and compressed; head long, snout sharp and projecting; anterior profile more or less S-shaped, nearly straight from tip of snout to front of eye, there concave, thence gibbous to front of dorsal, this character more pronounced in adults; mouth very large, gape curved, maxillary nearly reaching middle of eye; jaws subequal, or lower slightly included; teeth strong, in rather broad bands, those of outer series enlarged; antrorse teeth of

posterior part of each jaw strong; interorbital space convex; preopercle finely serrate; gillrakers small and rather weak, about 12+15; scales rather large, those above lateral line very much enlarged anteriorly, arranged in irregular and very oblique series; those below lateral line but slightly enlarged anteriorly and in oblique series. Dorsal spines stout, third to fifth longest, about 2.5 in head, longest soft ray about 4.5; first anal spine very short, as usual, second long and strong, 2.4 in head, scarcely reaching tips of last soft rays when depressed and scarcely longer than third; soft portion of anal slightly concave, anterior rays longest, about 2.5 in head; upper caudal lobe the longer, 1.4 in head; pectoral long and somewhat falcate, about 1.33 in head; ventrals long, about 1.57 in head; caudal and soft parts of dorsal and anal densely and finely scaled, few scales on pectorals and ventrals.

Color in life: General color light bluish, series of scales each with a small brown or brassy spot, these forming narrow indistinct lines running upward and backward; above the lateral line the body of each scale is bluish, the border brownish olive; a brassy band along lateral line; back with some bronze; under parts whitish; about 12 narrow, irregular bright-blue lines on head, separated by broader brassy lines, these lines sometimes extending upon body; inner edge of maxillary orange; lower anterior edge of opercle yellow; inside of mouth red or deep yellow; lips dusky; dorsal grayish, with a narrow yellow edge on spinous portion; caudal plain gray; anal gray, tinged with yellow.

Color in spirits: Body pale-grayish with slight bluish iridescence on terminal borders of scales; bases of scales above lateral line brown; under parts paler; head purplish, blue lines persistent but duller, the brassy entirely faded; bright color of inside of mouth also faded.

The range of this species is from Cape Hatteras and Pensacola south to Brazil, on sandy shores. It is par excellence "the grunt" of our South Atlantic States and Florida. About Porto Rico it is one of the most abundant and valuable food-fishes. Numerous specimens were obtained at San Juan, Mayaguez, Puerto Real, Boqueron, Guanica, Ponce, Arroyo, Hucares, Isabel Segunda, and Culebra. It was one of the most common species in all the markets of the island. It is caught either in the fish-traps or with haul seines. In our seining about the island this was the species most frequently taken, particularly on sandy shores. At Key West it is the most abundant of all the food-fishes, and is caught all the year round, the best season being during the fall. Their spawning season is during August and September, at which time they gather up into schools on shoal, feathery, rock bottom, where they spawn. Each roe is from 1 to 2 inches in length. The eggs are said to be "gritty" to the touch and about the size of a No. 10 shot. When ripe they separate and flow freely from the fish. After spawning the schools break up and the fish scatter. They are so abundant, however, that they can usually be found in large numbers on suitable bottom, the best fishing being on rock bottom. These fish grow to about 18 inches in length, with a maximum weight of 4 pounds; but those over 2 pounds are rare, and the average weight does not exceed one-third of a pound.

Guabi coara brasiliensis, Maregrave, Hist. Brasil., 163, 1648, Brazil.

Perca marina capite striato (the Grunt), Catesby, Hist. Carolina, pl. 6, 1743, Bahamas, etc.

Labrus plumieri Lacépède, Hist. Nat. Poiss., III, 480, pl. 2, fig. 2, 1802, Martinique; on a copy of a drawing by Plumier.

Hæmulon formosum Cuvier, Règne Animal, ed. 2, II, 175, 1829, Martinique.

Hæmulon arcuatum Cuvier & Valenciennes, Hist. Nat. Poiss., IX, 481, 1833, Charleston, S. C.

Hæmulon arara Poey, Memorias, II, 177, 1860, Cuba.

Hæmulon subarcuatum Poey, Memorias, II, 419, 1860, Cuba; a specimen with blue bands on anterior half of head only.

Hæmulon plumieri, Jordan & Evermann, I. c., 1304, 1898.

143. *Hæmulon flavolineatum* (Desmarest). *Ronco Condado; French Grunt; Open-mouthed Grunt.*

Head 2.8; depth 3; eye 3; snout 3; maxillary 2.5; interorbital 3.5; scales 7-50-11; D. XII, 14; A. III, 8. Body oblong-ovate, compressed, back not much elevated, profile from tip of snout to nape nearly straight, thence gently arched to origin of dorsal; mouth large, nearly horizontal, gape curved, maxillary nearly reaching middle of pupil; teeth moderate, outer somewhat enlarged; antrorse teeth in posterior part of each jaw considerably enlarged, those of upper jaw canine-like. Scales large, those on anterior part of body below lateral line considerably enlarged; rows of scales above lateral line running very obliquely upward and backward, those below somewhat wavy, most of them forming a curve with convexity downward and backward. Fins all moderate; dorsal spines slender, rather weak, longest about 2 in head; second anal spine longest and strongest, about 1.8 in head, its tip, when depressed, extending beyond tip of last ray.

Color in life: Side with about 12 irregular brassy lines alternating with about same number of similar pale-bluish lines, upper brassy lines rather dark, middle ones brightest and broadest, the broadest

two uniting at gill-opening and running across opercle as one brassy line; anterior edge of opercle bright indigo-blue.

The above description based upon young individuals 2.5 to 4 inches long. The color of larger individuals, from Havana, has been described as follows: Light bluish-gray as ground color; a bronze-yellow spot on the upper part of each scale, these forming continuous undulating stripes on the whole body and head, wider than the interspaces of ground-color, and nearly straight on caudal peduncle, on anterior part of body below lateral line broader and very oblique; a horizontal golden-yellow stripe, crossing the others, runs alongside of back from occiput to last rays of soft dorsal; yellow around eye; yellow shades and streaks on cheek, not strongly marked as in *H. sciurus* and *H. plumieri*; yellow stripes on top of head; angle of mouth black, inside brick-red; a large black blotch under angle of preopercle; fins bright golden-yellow, the pectoral and spinous dorsal paler.

In spirits the ground-color becomes grayish and the stripes brownish or dusky.

The range of *H. flavolineatum* is from the Bermudas, the Florida Keys, and Tortugas south to Brazil. It is generally common in the West Indies and abundant about Porto Rico, usually on sandy shores. No adults were obtained, but numerous young from the following localities: San Antonio Bridge, Mayaguez, Puerto Real, Ensenada del Boqueron, Ponce, Arroyo, Hucares, and Culebra.

This well-marked species reaches a length of a foot and is a good food-fish.

Diabasis flavolineatus Desmarest, Prem. Décade Ichth., 35, pl. 2, fig. 1, 1823, Cuba.

Hemulon heterodon Cuvier, Règne Animal, ed. 2, vol. 2, 176, 1829, Cuba.

Hemulon xanthopteron Cuvier & Valenciennes, Hist. Nat. Poiss., V, 254, 1830, Martinique.

Hemulon flavolineatum, Jordan & Evermann, l. c., 1306, 1898.




FIG. 55.—*Bathystoma rimator*.

Genus 82. BATHYSTOMA Scudder.

This genus differs from *Hemulon* in having 13 dorsal spines; body rather elongate; gillrakers rather numerous, 12 to 18 on lower part of anterior arch; mouth moderate; scales small; frontal foramina long, divided slits in front of supraoccipital crest. Jaws red within.

- a. Mouth large, maxillary reaching middle of eye, its length about one-half head; gillrakers rather short and few; scales moderate, 50 to 56 in lateral line; second anal spine scarcely larger or longer than third, 2.75 or more in head.
- b. Body oblong, back moderately elevated, depth 2.75 to 3 in length *rimator*, 144
- bb. Body subfusiform, back little elevated, depth 3.33 to 3.5 in length *caeruleum*
- aa. Mouth rather small, maxillary not reaching to opposite middle of eye, its length not quite one-half head; gillrakers numerous, rather long; scales small, about 70 in lateral line; second anal spine notably longer and larger than third; body more elongate than in other species, depth about 3.5 in length *striatum*, 145

144. *Bathystoma rimator* (Jordan & Swain). *Tom-tate*; *Red-mouth Grunt*; *Cesar*.

Head 2.8; depth 3; eye 4.2; snout 2.5; maxillary 2; mandible 2.2; interorbital 3.7; preorbital 8; D. XIII, 15; A. III, 8; pectoral 1.5; ventral 2; scales 8-56-12. Body elongate, back somewhat elevated, anterior profile slightly convex; snout rather long, pointed; eye large; mouth large, maxillary reaching

pupil; teeth not strong, outer series enlarged; scales rather small, in oblique series above lateral line, horizontal below and slightly enlarged anteriorly. Dorsal spines slender, fourth 2.4 in head; second anal spine equaling third in length, but much the stronger, 3 in head.

Color in life: Silvery white, slightly bluish above, with iridescent reflections; edges of scales of body light-yellow, these forming continuous light-yellow lines, those below lateral line horizontal, those above very oblique; besides these a narrow continuous streak of light yellow above lateral line from head to end of soft dorsal, and another from eye to middle of caudal; head silvery yellowish above; inside of mouth red; no black under preopercle; traces of black blotch at base of caudal; fins colorless, lower slightly yellowish.

The range of this fish is from Cape Hatteras and Pensacola southward through the West Indies to Trinidad. It is abundant about Charleston, S. C., where it is one of the most common food-fishes; adults said to be uncommon about Pensacola and Key West, but at the latter place the young swarm everywhere about the wharves and shores. It is as yet not known from Cuba, and does not seem to be at all abundant in Porto Rico, as it was seen only at Aguadilla, Ensenada del Boqueron, and Culebra Island. The specimens obtained are 5 and 6 inches long.

Hæmulon rimator Jordan & Swain, Proc. U. S. N. M. 1884, 308, Charleston, Key West, and Pensacola.

Bathystoma rimator, Jordan & Evermann, l. c., 1308, 1898.

145. *Bathystoma striatum* (Linnaeus). *White Grunt*.

Head 2.8; depth 3.3; eye 3; snout 3.28; maxillary 3.16; mandible 2.75; interorbital 3.8; preorbital 8.3; scales 8-70-13; D. XII, 13, the longest spine about 2.2 in head; A. III, 7; pectoral 1.5 in head; ventrals 1.75; caudal 1.5. Body elongate, fusiform, back little elevated, anterior profile scarcely arched; head moderate, snout short and blunt; mouth small, maxillary reaching front of pupil; eye large; teeth small, outer series somewhat enlarged; interorbital wide, preorbital narrow; preopercle finely serrate; gillrakers numerous, long and slender, 7 + 20. Scales very small and crowded, those above lateral line in very oblique series, those below more nearly horizontal, none enlarged; soft parts of all fins densely scaled; spines slender and weak.

Color in alcohol: Pearly-gray, with 5 or 6 continuous brownish streaks (probably golden in life), 1 on median line from tip of snout to origin of dorsal, 1 diverging from snout and passing above eye and along side to soft dorsal; another from snout crosses upper part of eye and terminates near beginning of lateral line; a fourth passes through eye and along middle of side to base of caudal; another crosses opercle and base of pectoral.

This fish is known from the Bermudas, Key West, Cuba, Santo Domingo, and Porto Rico, but is apparently not common. It probably never exceeds a foot in length. We seined one specimen, 3.25 inches long, at San Antonio Bridge.

Percu striata Linnaeus, Syst. Nat., ed. X, 233, 1758, North America.

Grammistes trivittatus Bloch & Schneider, Syst. Ichth., 188, 1801, Brazil; on the description of Maregrave.

Serranus capcuna Lichtenstein, Abhandl. Berlin Akad. 1821, 288, Brazil; on the description of Maregrave.

Hæmulon quadrilineatum Cuvier & Valenciennes, Hist. Nat. Poiss., V, 238, pl. 120, 1830, Santo Domingo.

Hæmulon quinquelineatum Poy, Memorias, II, 419, 1860, Cuba.


Bathystoma striatum, Jordan & Evermann, l. c., 1310, 1898.

Genus 83. ANISOTREMUS Gill.

Body ovate, short, deep, and compressed; mouth rather small with thick lips, maxillary rather short; inside of mouth not red; teeth in jaws only, all pointed, those of outer series in upper jaw enlarged; chin with a median groove besides smaller pores. Dorsal spines strong; soft rays of dorsal and anal scaly at base, anal spines strong; caudal mostly lunate; scales large; lower pharyngeals broad, with coarse, blunt teeth.

This genus, which is closely related to *Hæmulon*, contains 12 species, valued as food-fishes, all from the shores of tropical America and only two of which have been taken in Porto Rico. All of the species undergo considerable change in form with age. The young are marked with two or three blackish, lengthwise stripes. These disappear with age, quickest in the brightly colored species, but persist a long time in species like *A. bilineatus* and *A. interruptus*, which agree in coloration with *Hæmulon parva* and related species.

- a. Scales above lateral line arranged in oblique series which are not parallel with it.
- b. Scales comparatively large, less than nine in a vertical series between first dorsal spine and lateral line; coloration olivaceous, adult nearly plain, young with two or more dusky lateral stripes which disappear with age; fins blackish.
- c. Scales 5 or 6-52-15 (lateral line with 49 pores); scales above lateral line on anterior part of body more or less enlarged, especially in adult.
- d. Scales above lateral line not much enlarged, about 9 in an oblique series from first dorsal spine to lateral line; profile of head anteriorly rounded..... *surinamensis*, 146
- cc. Scales 7 or 8-46-15 (lateral line with 51 pores); scales above lateral line anteriorly not especially enlarged... *bicolor*
- bb. Scales rather small, more than 9 in a vertical series between first dorsal spine and lateral line.
- c. Body not striped longitudinally with yellow or blue; preorbital narrow; gillrakers $x + 13$.
- f. Anterior part of body with a black vertical bar.
- g. Body with 2 lengthwise bands; humeral bar brownish; a dark caudal spot, and a spot on the back of the caudal peduncle..... *spleniatus*
- cc. Body with longitudinal stripes of blue or yellow or both; young with a black blotch at base of caudal; preorbital broad; gillrakers $x + 16$.
- h. Anterior part of body with 2 broad, dark crossbars, the one from nape obliquely forward through eye, other from front of dorsal downward; behind these a series of horizontal stripes alternately yellow and blue; pectoral longer than head; second anal and fourth dorsal spines nearly equal.
- i. Blue stripes on side as broad as a scale, each more than two-thirds width of golden-yellow interspaces and each very faintly edged with darker; vertical bands on head and shoulder jet-black..... *virginicus*, 147
- hh. Anterior part of body without dark crossbars, body sometimes plain yellowish, back usually violet with 4 or 5 yellow lines; silvery below; snout short, not longer than width of the eye; dorsal fin very deeply notched, with feeble spines; the second and third anal spines equal in length; the body a little more oblong than in *surinamensis*..... *serrula*

FIG. 56.—*Anisotremus surinamensis*.146. *Anisotremus surinamensis* (Bloch). *Pompon*.

Head 3.14; depth 2.2; eye 4.5; snout 2.6; maxillary 3; D. XII, 16; A. III, 8 or 9; scales 5-50-13. Body deep, back elevated, greatly compressed; profile steep, nearly straight from snout to above eye, a slight depression in front of nostril and another in interorbital space; profile from interorbital space to dorsal strongly arched in a broad curve. Head moderate; cheek deep; mouth rather small; jaws subequal, maxillary barely reaching front of orbit; ventral line of body nearly straight; caudal peduncle moderately long, its least depth equal to snout; teeth in several bands, outer enlarged and canine-like. Fourth dorsal spine strongest and longest, its length 2.2 in head; soft dorsal as well as anal, pectoral, ventrals, and caudal densely covered with minute scales; height of longest soft dorsal ray 3 in head; second anal spine very stout, its length equal to that of fourth dorsal spine; third anal spine broad at

base, but shorter than second; free edge of soft anal straight; dorsal and anal fins depressible in a scaly sheath; pectoral long and falcate, nearly reaching tip of ventrals, 1.2 in head; ventral shorter, 1.2 in pectoral; caudal well forked, the lobes about equal to ventral. Preopercle strongly but irregularly serrate. Scales of cheek in about 7 rows, those on opercle in about 8 rows, on interorbital and nape small and crowded; scales of back and sides arranged in oblique rows not parallel with lateral line, which is arched, following approximately contour of back. Gillrakers rather short, stiff, 13+19.

Color, grayish, darkest on anterior half of body, where each scale is dark-brown on its basal half, then with a white ellipse, narrow border darker, contrast between dark base and white ellipse very marked; owing to the irregular arrangement of scales the dark bases in some cases appear as spots; upper side of caudal peduncle brown, sides nearly plain white; snout and under parts of head lilac-brown; under parts of body rusty-brown; fins all dark brown, especially the soft parts of dorsal and anal.

The above description from a specimen 15 inches long, taken in Indian River, Florida.

The recorded range of this species is from the Indian River, Florida, and the Tortugas to Brazil; it is known from Surinam, Martinique, Porto Rico, Jamaica, and Cuba. It attains a length of 2 or 3 feet and is a good food-fish, though it is not sufficiently numerous anywhere to be of great importance. Only a single specimen, 2.4 inches long, seined at Ponce, was obtained in Porto Rico.

Lutjanus surinamensis Bloch, Ichth., pl. 253, 1791, Surinam.

Holocentrus gibbosus Lacépède, Hist. Nat. Poiss., IV, 344, 1803, Surinam.

Pristipoma bilineatum Cuvier & Valenciennes, Hist. Nat. Poiss., V, 271, 1830, Martinique.

Pristipoma melanopteron Cuvier & Valenciennes, Hist. Nat. Poiss., V, 273, 1830, Brazil.

Hemulon obtusum Poey, Memorias, II, 182, 1860, Havana.

Hemulon labridum Poey, Memorias, II, 419, 1861, Cuba.

Anisotremus surinamensis, Jordan & Evermann, l. c., 1318, 1898.




FIG. 57.—*Anisotremus virginicus*.

147. *Anisotremus virginicus* (Linnæus). "Sisi"; *Catalineta*; *Pork-fish*.

Head 3 to 3.2; depth 2 to 2.25; eye 3.75 to 4; snout 2.75; maxillary 3.25 to 3.5; interorbital 3; preorbital 4; scales 13-57 to 60-16; D. XII, 16; A. III, 9; pectoral a little longer than head; ventral 1.5; caudal lobe 1.2. Body ovate, back greatly elevated, anterior profile very steep, slightly convex along snout and over eye, very much arched on nape; highest point of body at origin of dorsal fin; mouth small, little oblique, maxillary extending to anterior edge of orbit; jaws subequal; outer row of teeth enlarged; gillrakers 11+11, short and not very stiff. Dorsal spines rather strong, longest (third) about 1.8 in head; dorsal rays short; second anal spine strongest and longest, about 2 in head; caudal well forked, lobes subequal; ventrals just reaching vent; pectoral long, somewhat falcate, reaching origin

of anal. Scales large, strongly ctenoid, those on nape and breast reduced in size; scales above lateral line in oblique series.

Color in life: Side with about 8 broad, lemon-yellow lines alternating with similar lines of dirty silvery, the upper 3 or 4 of these yellow lines branch anteriorly, the fifth extending on middle of caudal peduncle; belly silvery-white; a broad black bar from origin of spinous dorsal downward to base of pectoral, the black continuing on shoulder-girdle to near isthmus; another broad black bar from occiput through eye to angle of mouth; cheek metallic or brassy-greenish; top of head brassy; fins all orange-yellow, spinous dorsal, pectoral, and ventral dusted with brownish; scaly sheath at base of anal rich yellow. In alcohol, the longitudinal stripes become bluish and the yellow fades.

This handsome fish ranges from Florida to Brazil. It is known from Biscayne Bay, Key West, Santo Domingo, Jamaica, Porto Rico, Martinique, and St. Catharine Island, Brazil, and is the commonest species of the genus in the West Indies. About Key West it is said to school from June to August, which is the spawning season, and is found then about the shoals, but soon retires to deeper water. It spawns all through the channel about the shoals, and is then caught in greatest numbers. About a month after the spawning season immense numbers of young are seen on the shoals. It reaches a length of about a foot and a weight of 2 pounds. The average weight of those brought to market probably does not exceed one-third of a pound. It does not appear to be very common about Porto Rico as only five examples were obtained—one at Ponce, one at Mayaguez, and three at Arroyo, where it was called "sisi" by the fishermen. It is regarded as a good food-fish.

Guatucupa juba, Maregrave, Hist. Brasil., 148, 1648, Brazil.

Acara pinima, Maregrave, Hist. Brasil., 152, 1648, Brazil.

Sparus virginicus Linnaeus, Syst. Nat., X, 281, 1758, South America.

Sparus vittatus Bloch, Ichth., taf. 263, fig. 2, 1791, Brazil; after *Acara pinima* of Maregrave.

Perca juba Bloch, Ichth., taf. 308, fig. 2, 1791, Brazil; after *Guatucupa juba* of Maregrave.

Grammistes mauritii Bloch & Schneider, Syst. Ichth., 185, 1801, Brazil; after *Sparus vittatus* Bloch.

? *Pristipoma catharine* Cuvier & Valenciennes, Hist. Nat. Poiss., V, 269, 1830, St. Catharine Island, Brazil.

Pristipoma vodo Cuvier & Valenciennes, Hist. Nat. Poiss., V, 274, 1830, Brazil, Martinique, Porto Rico, and Santo Domingo.

Pristipoma acara pinima, Castlenau, Anim. Nouv. ou Rares, 8, 1856, Brazil.

Anisotremus virginicus, Poey, Fauna Puerto-Riqueña, 324, 1881; Jordan & Evermann, l. c., 1322, 1898.

Genus 84. CONODON Cuvier & Valenciennes.

This genus is close to *Pomadasis*, from which it is separated by the enlarged outer teeth and by the peculiar armature of the preopercle, which is very sharply serrate, serræ at angle enlarged, those before angle turned forward. Body oblong; soft rays of dorsal and anal more or less scaly; second anal spine large. Of the two species of this genus only one occurs in Porto Rico.

148. *Conodon nobilis* (Linnaeus). "*Burteado*."

Head 3.25; depth 3.25; eye 4.1; snout 3.4; maxillary 2.8; mandible 2.25; interorbital 5.1; preorbital 5.3; D. XI-1, 13; A. III, 7; pectoral 1.4; ventral 1.5; caudal 1.4; scales 5-51-11. Body oblong, back elevated and moderately compressed, depth equal to length of head; mouth small, maxillary falling short of front of eye; lower jaw projecting; teeth in front of both jaws much enlarged, conic, short, but very stout; pectoral with about 10 short, sharp serrations on upper limb, lowermost, at angle of preopercle, much the strongest; antrorse serrations on lower limb; dorsal spines strong, a deep notch between them and soft dorsal; second anal spine longest and strongest.

Color in life: Back grayish, becoming paler on sides; belly white, silvery; top of head grayish; side with 8 broad, vertical dark bars, broadest above, gradually narrowing and disappearing below; interspaces broader, with brassy wash; some evidence of 2 pale brassy stripes below lateral line; side of snout, head, and tip of lower jaw pale-lemon; fins all with some yellow, ventral brightest.

This species is found on the coast of Texas, West Indies, and Brazil, not uncommon on sandy shores; recorded from Jamaica, Martinique, Brazil, and Texas; not rare in Porto Rico, where specimens were obtained by us at Arecibo, Mayaguez, Palo Seco, Ponce, Arroyo, and Vieques. It reaches a length of about a foot, but our examples range from 2.75 to 4.5 inches.

Perca nobilis Linnaeus, Syst. Nat., ed. X, 191, 1758, North America.

Sciæna plumieri Bloch, Ichthyol., VI, 66, taf. 306, 1791, Martinique.

Sciæna coro Bloch, l. c., pl. 307, fig. 2, 1791, Brazil; after *Coro coro* Maregrave.

Conodon antillanus Cuvier & Valenciennes, Hist. Nat. Poiss., V, 156, 1830, Jamaica.

Pristipoma coro, Poey, Fauna Puerto-Riqueña, 324, 1881; Stahl, l. c., 77 and 163, 1883.

Conodon nobilis, Jordan & Evermann, l. c., 1324, 1898.

Genus 85. BRACHYDEUTERUS Gill. The Burritos.

Body oblong; scales large, those above in series parallel with lateral line; mouth small, outer teeth in jaws somewhat enlarged; inside of jaws not red. Anal spines small or moderate, second little, if any, longer or stronger than third, and lower than soft rays; soft dorsal and anal largely covered with small scales; dorsal spines 12; soft dorsal comparatively long, of 15 or 16 rays; otherwise essentially as in *Pomadasis*, fins smaller and more scaly. Only one of the four American species is known from Porto Rico.

149. *Brachydeuterus corvinæformis* (Steindachner).

Head 3 to 3.25; depth 3.25; eye 3.75 to 4 in head; snout 2.2 to 3; D. XII, 15; A. III, 7; scales 6-51-10. Upper profile regularly arched from snout to tail, highest point at origin of dorsal; lower profile nearly straight to base of anal; preorbital slightly less than eye; maxillary not reaching eye, 3.33 to 3.4 in head; anterior nostril oval, twice size of posterior; preopercle finely toothed, about 20 teeth on upper limb, increasing slightly in size toward angle; dorsal notched almost to base, fourth spine longest, 2.12 to 2.5 in head; longest soft ray (second) equal to or slightly less than longest spine; pectoral pointed, 1.25 to 1.4 in head; ventral broad, margin nearly straight, inner ray 1.25 in second, which is 2 in head; the first ray branched once, slightly filamentous, the other rays much branched; second and third anal spines about equal, 3 to 3.75 in head, relatively larger in smaller specimens, second stouter; first soft ray 2.5 in head; margin of anal slightly concave, last ray shorter than second spine; upper lobe of caudal longer, the difference more noticeable in smaller specimens, 1.33 to 1.5 in head, middle rays 1.75 in upper; pectoral scaly at base only, other fins, except spinous dorsal, more or less completely scaled, a narrow sheath of scales on sides of dorsal and anal, supplementary scales slightly developed in axil of ventrals; least depth of caudal peduncle 1.5 to 1.6 in length from below end of dorsal, its length being equal to middle caudal rays; lateral line and rows of scales above it concentric with back, scales below lateral line in horizontal rows.

Color in alcohol: Dark olive above, lower sides more or less silvery, larger specimens much the lighter, a dark line along each row of scales below lateral line, these rather indistinct in larger specimens; scales above lateral line with dark centers, these not forming distinct lines; a diffuse dark blotch on scapular region, very faint in the larger specimens; pectoral colorless, other fins punctate, margins very dark. There is considerable variation in the ground-color, larger specimens being distinctly silvery, some of the smaller heavily washed with olive and having scapular blotch more developed.

Our specimens differ from Steindachner's description in having a rather shorter maxillary, not reaching eye, narrower preorbital, and a blotch on scapular region. They are, however, probably not specifically distinct.

The range of this species is from the West Indies to Brazil; apparently common about Porto Rico, specimens being in the collection from Palo Seco, Aguadilla, Mayaguez, Arroyo, Hucares, and Isabel Segunda. It reaches a length of about a foot, and is a good pan-fish.

Hemulon corvinæforme Steindachner, Ichth. Notizen, VII, 16, 1868, Santos, Brazil.

Brachydeuterus corvinæformis, Jordan & Evermann, I. c., 1326, 1898.

Genus 86. POMADASIS Lacépède. The Burros.

Body oblong, somewhat compressed, back not much elevated; mouth rather small, terminal, low, lips thin; maxillary usually not extending to opposite eye, its tip not reaching posterior edge of broad preorbital; premaxillary protractile; teeth on jaws only, in villiform bands, subequal, or outer series in upper jaw more or less enlarged; no red on jaws; a central groove behind symphysis of lower jaw; cheek and opercles scaly; the preopercle rather distinctly serrate, serræ below not turned forward; suprascapular serrate; scales large, those above lateral line in series parallel with it; no small scales at base of the others; soft dorsal and anal fins naked, or with very few scales at the base; dorsal fin emarginate, spines strong, 12 or 13 in number, soft rays 11 to 14; anal fin of 7 or 8 soft rays, short, with second spine always very strong; gillrakers feeble, few in number; caudal lunate, forked.

This genus is composed of small shore fishes, some of its representatives being found in most tropical seas; several enter fresh waters, and perhaps belong to the brackish-water fauna. Numerous species are found on the west coast of Africa and about the Cape Verde Islands, but so far as known none enters European waters. Only three of the eight American species are known from Porto Rico.

- a. Dorsal spines XII; preorbital broad.
 b. Scales small, 65 in a longitudinal series; body elongate; maxillary extending a little beyond front of eye; anal spines strong, second three-fourths depth of body. Color, nearly plain; silvery below *productus*, 150
 aa. Dorsal spines XIII; preorbital narrow.
 c. Body moderately elongate, depth 2.66 to 3 in length *crocro*, 151
 cc. Body very long and low, compressed, back little elevated, depth about 3.6 in length..... *ramosus*, 152

150. *Pomadasis productus* (Poey).

Head 4 in total length with caudal; depth 4.33; eye 4; D. XII, 12; A. III, 7; scales 65. Body elongate, maxillary extending a little beyond front of eye; anal spines strong, second three-fourths depth of body. Color, nearly plain; silvery below. (Poey.)

Apparently closely allied to *P. bayanus* and differing from *P. ramosus*, if the descriptions can be trusted, in having but 12 dorsal spines. Recorded from Cuba and Porto Rico by Poey; not seen by us.

Pristipoma productum Poey, Memorias, II, 186, 1860, Havana; Poey, Fauna Puerto-Riqueña, 324, 1881.

Pomadasis productus, Jordan & Evermann, l. c., 1332, 1898.

151. *Pomadasis crocro* (Cuvier & Valenciennes). "Viejo."

Head 3; depth 2.66 to 3.5; eye 3.5 to 5; D. XIII, 11 or 12; A. III, 6 or 7; scales 6-54-16; preorbital 4 to 8; snout 2.8 to 3.66; maxillary 3 to 4; pectoral 1.16 to 1.5; fourth dorsal spine 1.8 to 2.25; second anal spine 1.5 to 2; soft dorsal 2 in spinous. Body elongate, compressed, back elevated, high at nape, anterior profile rather irregular, varying with age; a more or less distinct frontal depression above eye in old examples; mouth small, maxillary barely extending to anterior edge of orbit; lower jaw included; preopercle coarsely serrate, teeth wide apart; teeth small, outer scarcely enlarged; scales rather large, those above lateral line in parallel series; dorsal fin moderately notched; second anal spine very strong and long, reaching past tips of all rays; pectoral short, caudal slightly lunate.

Color in life: Grayish, with about 12 narrow interrupted stripes formed by pale-silvery quadrate spots, one on middle of each scale, plainest on middle of side; under parts white, washed with rusty; opercle brassy; top of head rusty; spinous dorsal darkish; ventrals and pectoral dusky; iris lemon-yellow; mouth scarlet within, pale flesh-color anteriorly.

West Indies, from Cuba to Brazil; generally common on sandy coasts. A specimen, 7.5 inches long, obtained in San Juan market, where others were seen. A good food-fish, though of small size.

Pristipoma crocro Cuvier & Valenciennes, Hist. Nat. Poiss., V, 264, 1830, Martinique.

Pristipoma cultriferum Poey, Memorias, II, 185, 1860, Havana; Poey, Fauna Puerto-Riqueña, 324, 1881; Stahl, l. c., 77 and 163, 1883.

Pomadasis crocro, Jordan & Evermann, l. c., 1333, 1898.

152. *Pomadasis ramosus* (Poey). "Ronco Blanco."

Head 3.15; depth 3.7; eye 4; snout 3.1; maxillary 2.7; mandible 2.25; interorbital 4.4; preorbital 6.6; D. XIII, 12; A. III, 7; pectoral 1.7; ventral 1.7; caudal 1.4; scales 6-57-13. Body long and low, back somewhat compressed and a little elevated, anterior profile nearly straight; head pointed, mouth not large, maxillary reaching past front of orbit, lower jaw included; teeth in villiform bands, outer row in upper jaw slightly enlarged; preopercle and suprascapular scale strongly serrate; dorsal fin continuous, well notched, spines very strong; second anal spine very greatly enlarged, as long as longest rays; bases of soft dorsal and anal with a sheath of small scales, that of soft dorsal low; caudal lunate; lateral line nearly straight save under soft dorsal.

Color in life: Silvery, bluish on back, slightly brassy on side, white below; pectoral pale-yellow; ventral pale; anal and soft dorsal darker, membrane of spinous dorsal dark-edged; caudal dark with black edge.

Found in the West Indies, south to Brazil. Two specimens, about 10 inches long, one from San Juan market, the other from the Rio Loiza near Caguas, more than 25 miles from the coast and in fresh water. Five young examples, each 2 to 3 inches long, were taken in the seine at Aguadilla. This species is said by the native fishermen to utter the grunting noise characteristic of the family. Though not large, it is nevertheless a good food-fish and is highly valued.

Pristipoma ramosum Poey, Memorias, II, 186, 1860, Cojimar River, Havana, Cuba.

Pomadasis ramosus, Jordan & Evermann, l. c., 1334, 1898.

Family XLVI. SPARIDÆ. The Porgies.

Body oblong, or more or less elevated, covered with rather large, adherent scales, which are never truly ctenoid. Lateral line well developed, concurrent with back, not extending on caudal fin. Head large, crests on skull usually largely developed. No suborbital stay. Mouth small, terminal, low, and horizontal. Premaxillaries little protracile; the maxillary short, peculiar in form and in articulation, without supplemental bone, for most of its length slipping under edge of the preorbital, which forms a more or less distinct sheath; preorbital usually broad; teeth strong, those in front of jaws conical, incisor-like or molar; lateral teeth of jaws always blunt and molar; no teeth on vomer or palatines; posterior nostril largest, usually more or less oblong or slit-like; lower pharyngeals separate; gills 4, a large slit behind the fourth; pseudobranchiæ large; gillrakers moderate; gill-membranes separate, free from the isthmus; preopercle entire or serrulate; opercle without spines; sides of head usually scaly; dorsal fin single, continuous, or deeply notched, spines usually strong, depressible in a groove; spines heteracanthous, that is, alternating, the one stronger on the right side, the other on the left, 10 to 13 in number; anal fin rather short, similar to soft dorsal, and with 3 spines; ventral fins thoracic, rays 1, 5, with a more or less distinct scale-like appendage at base; caudal fin usually more or less concave behind; air-bladder present, usually simple; pyloric cæca few; vertebrae usually 10 + 14 = 24; intestinal canal short.

Carnivorous shore fishes of the tropical seas, especially abundant in the Mediterranean, the Red Sea, and the West Indies.

- a. Teeth in front of jaws conical or incisor-like, not molar; dorsal fin continuous; posterior nostril oblong; preopercle entire.
- b. Second interhamal bone enlarged, hollowed anteriorly, or pen-shaped, receiving the posterior end of air-bladder in its anterior groove; posterior nostril slit-like.
- c. Front teeth narrow, compressed, forming lanceolate incisors; first spine-bearing interneural with an antrorse spine; temporal crest obsolete; lateral crest nowhere coalescing with supraoccipital crest; interorbital area flattish, with two low ridges; a small foramen in each of these above front of pupil; interorbital area much contracted anteriorly; a strongly projecting prefrontal process, which makes an acute angle with supraorbital.
- d. Frontal bones partly porous and gibbous; antrorse dorsal spine attached directly to interneural: third dorsal spine very long, longer than the head..... OTRYNTER
- dd. Frontal bones not gibbous nor porous; antrorse dorsal spine attached to interneural by a long process or spur; third dorsal spine about half head..... STENOTOMUS
- cc. Front teeth conical or canine-like; first spine-bearing interneural without antrorse spine; temporal crest very thin and high, joining lateral crest which forms part of margin of orbit above middle of eye, both crests coalescing with supraoccipital in cavernous anterior part of interorbital area; interorbital area somewhat contracted anteriorly; prefrontal process very strong, making an obtuse angle with supraorbital, this process forming a conspicuous knob above the long posterior nostril..... CALAMUS, 87
- bb. Second interhamal spine normal, not "pen-shaped."
- e. Front teeth conic, not compressed; no incisors; occipital crest coalescent with temporal crests; no antrorse spine on first interneural; dorsal spines usually 11 to 13.
- f. Anterior teeth in both jaws strong, decidedly canine-like; body more or less deep and compressed..... PAGRUS
- ee. Front teeth incisor-like; no canines.
- g. Incisors broad; molars in 2 to 4 series in each jaw.
- h. First spine-bearing interneural with an antrorse spine in front.
- i. Supraoccipital and temporal crests nowhere coalescent, interorbital area not swollen; frontal bone in interorbital area thin, concave in transverse section; temporal crest low, separated from supraoccipital crest by a flattish area which extends forward on each side of supraoccipital crest and to groove of premaxillary spines. Incisors conspicuously notched..... LAGODON
- ii. Supraoccipital and temporal crests coalescent anteriorly, both disappearing in the gibbous interorbital area; frontal bone between eyes transversely convex and more or less honeycombed; temporal crest separated from occipital crest by an excavated area bounded anteriorly by lateral crest, which merges into supraoccipital above eye. Incisors entire or with a shallow notch..... ARCHOSARGUS, 88
- hh. First spine-bearing interneural without antrorse spine above; skull essentially as in *Archosargus*, frontal bone more cavernous..... DIPLODUS

Genus 87. CALAMUS Swainson. Pez de Plumas.

Body oblong, compressed, back elevated; head large, preorbital deep; mouth small, teeth strong, those in front conical or pointed, those on sides molar; preopercle entire, posterior nostril slit-like; dorsal fin rather low, not much notched, soft rays low, not scaly; caudal well forked; anal spines small; pectoral rather long. Second interhamal bone enlarged, hollowed anteriorly, or pen-shaped, receiving posterior end of air-bladder in its anterior groove; first spine-bearing interneural without antrorse spine; temporal crest very thin and high, joining lateral crest which forms part of margin of orbit above middle

of eye, both crests coalescing with supraoccipital in cavernous anterior part of interorbital area; interorbital area somewhat contracted anteriorly; prefrontal process very strong, making an obtuse angle with supraorbital, this process forming a conspicuous knob above the long posterior nostril.

Shore fishes, remarkably distinguished by the structure of the interhæmal. This genus contains numerous very closely related species, all American and all valued as food-fishes.

CALAMUS:

- a. Scales comparatively small, 8 or 9-54 to 58-18 or 19; about 6 vertical rows of scales on base of preopercle, with about 12 scales entering into formation of lower margin; species of large size, with preorbital deep, pectoral fin long, and outer teeth strong.
- b. Body very deep, back elevated, depth in adult half length to base of caudal; outer teeth about $\frac{10-12}{12-10}$ in number, outer one on each side in one or both jaws sometimes enlarged, canine-like; sometimes directed forwards, especially in adult.
- c. Preorbital with reticulations of bluish ground-color around bronze spots; canines of upper jaw usually vertical, but sometimes, especially in old examples, directed more or less horizontally forward; body deeper than in other species, depth 1.9 to 2.25 in length; anterior profile not very steep, slightly curved; depth of preorbital less than half head *calamus*, 153
- cc. Preorbital region, snout, cheek, and opercles brassy, crossed by horizontal, wavy, non-reticulating lines of violet-blue, brightest on preorbital and snout; a sky-blue blotch behind eye over opercle, extending a short distance on body; outer canines of upper jaw directed horizontally forward, except in very young, these teeth longer than in *C. calamus*; anterior profile nearly straight and very steep to nape, then strongly convex.
- d. Eye about 4 in head; side with dark crossbands; blue lines on preorbital reticulating; iris dark..... *proridens*
- dd. Eye larger, about 3.5; side with bluish longitudinal lines; blue lines on preorbital not reticulating; iris yellow. *kendalli*, 154
- bb. Body more elongate, depth 2.17 to 2.75 in length.
- c. Upper jaw with a strong antrorse canine on each side, as in *C. proridens*; preorbital with blue, wavy stripes; preorbital deep; dorsal high; pectoral reaching front of anal; cheek with blue flexuous lines, anastomosing and forming rivulations; spinous dorsal edged with black *pennatula*
- ee. Upper jaw without antrorse canines, anterior teeth strong, $\frac{4-6}{6-8}$, one on each side of upper jaw more or less enlarged; body rather oblong, snout long and pointed, anterior profile forming a nearly even curve to front of dorsal. Color dull-brassy with little blue; a faint blue stripe below eye; preorbital dull-coppery, usually plain, sometimes faintly veined with bluish. Young, as in other species, with dark crossbands *bajonado*, 155

GRAMMATEUS:

- aa. Scales comparatively large, 6 or 7-45 to 52-13 or 14; about 5 vertical rows of scales on base of preopercle, with about 9 scales entering into the formation of lower margin; no antrorse canines.
- f. Pectoral fin long, about 3 in body.
- g. Scales of moderate size, 50 to 52 in the lateral line. Body very deep, the back elevated, depth about 2.2 to base of caudal; longest dorsal spine about half head.
- b. Canines moderate, about $\frac{8-10}{10}$; preorbital broad, its least width about 2.25 in head. Body moderately compressed, rather elongate, back only moderately elevated, anterior profile convex to eye, thence straight to point of snout. Color smutty-silvery, with dark crossbands; blotches on fins; no black axillary spot *leucosteus*
- gg. Scales large, about 46 (45 to 48) in lateral line; body rather elongate, depth about 2.25 in body; longest dorsal spine about 2.5 in head.
- i. Canine small, about $\frac{1}{2}$; eye large, 3 in head; dorsal spines xi; preorbital narrow, about equal to eye. Dorsal outline forming a comparatively regular arch, back being elevated, anterior profile steep and nearly straight. Color, plumbeous-gray, with a blue spot on each scale, preorbital with blue streaks; a blue streak below eye; a blue point in axil *macrops*
- ff. Pectoral fin short, about 3.5 in body.
- j. Dorsal outline forming a comparatively regular arch, anterior profile from snout to base of spinous dorsal evenly convex; back elevated, depth in adult about 2.17 in length; canines subequal, $\frac{8}{10}$; preorbital not very deep; pectoral shortish. Color, dull silvery, with pearly spots on scales of back; preorbital bluish, plain, or with pearly markings, without blue stripes; a faint pale streak below eye; axil with a small inky-black spot; crossbars on body usually persistent *penna*
- jj. Dorsal outline not forming a regular arch, anterior profile straight from base of spinous dorsal to nape, where a rather sharp angle is formed, thence straightish above eye, snout convex; body rather elongate, depth about 2.5 in length.
- k. Preorbital deep, nearly twice diameter of eye; canine teeth $\frac{8}{10}$. Body oblong, back little elevated, anterior profile unevenly curved, very convex before eye. Color, olivaceous, with dark bars or spots, centers of many scales pearly; 6 yellowish spots along lateral line; preorbital brownish, usually with dashes of golden-yellow; membrane of opercle orange; fins mostly barred or spotted *arctifrons*, 156
- kk. Preorbital not deep, pectoral short, 1.2 in head; dorsal fin low, longest spine about 3 in head; canines $\frac{8}{8}$, moderate, equal. Body little elevated, the anterior profile rather strongly convex, curve continuous from snout to middle of dorsal. Color, olivaceous, with darker crossbands; preorbital plain; a dark axillary spot; a blue subocular band *medius*

153. *Calamus calamus* (Cuvier & Valenciennes). "*Pluma*"; *Saucer-eye Porgy*.

Head 3.1; depth 2.2; eye 3.4; snout 1.6; maxillary 2.5; interorbital 3.3; preorbital 2.1; D. XII, 12; A. III, 10; pectoral 0.9; ventral 1.5; caudal 1.2; scales 8-54-16; teeth in the outer row in front of each jaw enlarged but without distinct canines. Anterior profile forms a regular curve from tip of snout to dorsal, save in a small specimen, which has a distinct depression above eye.

Color in spirits: Not markedly different from that of other examples of the species of *Calamus* in the collection.

Five examples, 3.5 to 8.75 inches long, collected at Arroyo, Mayaguez, and Boqueron. The range of this species is from the Florida Keys southward among the West Indies. It has been recorded from Martinique, Jamaica, Cuba, and various points in southern Florida. It attains a length of about a foot and a weight of a pound or more. Those seen in market probably do not average over half a pound in weight. It is an excellent food-fish and always commands a fair price. At Key West, and apparently in Porto Rico also, it is usually taken with hook and line and affords some sport as a game fish.




FIG. 58.—*Calamus calamus*.

Pagellus calamus Cuvier & Valenciennes, Hist. Nat. Poiss., VI, 206, pl. 152, 1830, Martinique and Santo Domingo.

Calamus megacephalus Swainson, Nat. Hist. Fish., I, 222, 1839, Martinique and Santo Domingo; after Cuvier & Valenciennes.

Pagellus orbitarius Poey, Memorias, II, 201, 1860, Havana.

Calamus macrops Jordan & Gilbert, Synopsis, 927, 1883, Garden Key, Florida.

Calamus calamus, Jordan & Evermann, I. c., 1349, 1898.

154. *Calamus kendalli* Evermann & Marsh. "*Pluma*."

Head 3.1; depth 2.1; eye 3.5; snout 1.5; maxillary 2.4; interorbital 3.5; preorbital 2.1; D. XII, 12; A. III, 10; pectoral 1; ventral 1.8; caudal 1.3; scales 7-53-16. Body deep, back strongly elevated, more so than in *C. bajonado*, but less than in *C. calamus* or *C. proridens*, anterior profile a nearly regular curve, lacking the abrupt nuchal elevation of those species; eye large, larger than in *C. proridens*; 7 or 8 rows of scales on cheek; teeth about as in *C. proridens*; molars in two or more rows on sides, those of inner row much the largest, those in front becoming more numerous and merging into cardiform teeth, the most anterior of which in each jaw are somewhat enlarged; in front of upper jaw are 2 much enlarged antrorse canines, curved slightly upward; highest dorsal spine 2.7 in head, second anal spine 4.6.

Color in spirits: Silvery, sides with bluish longitudinal lines following rows of scales, plainest above; a pale-blue line bordering orbit below; some blue lines on preorbital, not evidently reticulated

and not as numerous as in *C. proridens*; iris yellow; otherwise as in *C. proridens*, to which this species is very close.

Type, No. 49362, U. S. N. M., 10.5 inches long, collected at Mayaguez, January 20, 1899; 2 others, each 8.5 inches long, from Mayaguez and Arroyo, are more slender (depth 2.3 and 2.45 in length), but not differing in any other character.

This species is probably not common about Porto Rico, as it was obtained only at Mayaguez and Arroyo. It was not noticed in any of the markets. Like all other members of the genus, however, it is doubtless a good food-fish, and probably reaches a length of a foot or more.

Calamus kendalli Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 353, Mayaguez, Porto Rico.




FIG. 59.—*Calamus kendalli*.

155. *Calamus bajonado* (Bloch & Schneider). "*Pluma*"; *Jolt-head Porgy*; *Bajonado*.

(PLATE 25.)

Head 3.2; depth 2.3; eye 3.3; snout 1.7; maxillary 2.5; interorbital 3.3; preorbital 2.1; D. XII, 12; A. III, 10; pectoral 0.9; ventral 1.4; caudal 1.1; scales 7-54-17. Body less elevated than in *C. calamus* or *C. proridens*, snout more pointed, anterior profile not so steep, rising in a regular curve to front of dorsal; anterior teeth enlarged, becoming very strong in adult, one on each side on front of upper jaw usually enlarged and canine-like, but not antrorse.

Color in spirits: Dull olive, with blue luster; faint longitudinal lines of blue; a blue line bordering orbit below, and one above nostril, extending on forehead; obscure irregular blue lines on preorbital; caudal faintly and obscurely barred.

This porgy is found throughout the West Indies and north to southern Florida. It is the most abundant species of the genus and reaches the largest size. We have examined specimens 2 feet long and weighing 8 or 10 pounds. The average weight of those usually caught does not exceed 5 or 6 pounds. It frequents smooth rock bottom upon which it is said to spawn in July and August. It is one of the most common species about Porto Rico, as elsewhere, and is said to be found at all seasons. Because of its large size it is more important as a food-fish than any of its congeners, though its flesh is somewhat coarse. It is taken in the hook-and-line fishery and also in various fish-traps.

Our collections contain 5 examples, each 6 to 8 inches long, obtained at Puerto Real and Arroyo. Others were seen in the possession of the fishermen at those places.

Bajonado Parra, Dif. Piezas, Hist. Nat. Cuba, 13, lam. 8, 1787, Havana.

Sparus bajonado Bloch & Schneider, Syst. Ichth., 284, 1801, Havana; after Parra.

Pagellus caninus Poey, Memorias, II, 199, 1860, Havana.

Calamus plumatula Guichenot, Rev. Pagels, 119, Martinique.

Calamus bajonado, Poey, Fauna Puerto-Riqueña, 328, 1881; Stahl, l. c., 78 and 164, 1883; Jordan & Evermann, l. c., 1352, 1898.

156. *Calamus arctifrons* Goode & Bean. Grass Porgy; Shad Porgy.

Head 3.3; depth 2.3; eye 3.4; snout 1.9; maxillary 2.4; interorbital 3.5; preorbital 2.8; D. xii, 12; A. iii, 10; pectoral 1; ventral 1.4; caudal 1.1; scales 5-47-12. Back not greatly elevated, anterior profile not steep, a slight angle in front of eye; about 5 rows of scales on cheek; 8 to 10 teeth in front of each jaw about equally enlarged and canine-like.

Color in spirits: Olivaceous, bluish above the trunk, with 6 or 7 obscure dark vertical bars, with scattering dark spots between them; similar dark spots on opercle and cheek; snout and top of head dark; a broad dark bar from eye straight downward across cheek; pectoral pale, ventrals mostly dark, color plainest inside; vertical fins with dark shades.

The grass porgy has hitherto been known only from various places on the Florida coast from Pensacola and Biscayne Bay south to Key West, and has not until now been reported from the West Indies. The only example seen by us in Porto Rico is 7 inches long, and was seined at San Antonio Bridge. This is probably the smallest species of the genus, rarely exceeding a foot in length, but is a good food-fish wherever found in sufficient numbers. It is usually taken with haul seines.




FIG. 60.—*Calamus arctifrons*.

While the other porgies are usually found on hard bottom, the grass porgy frequents the more shallow water where there is an abundance of grass or other aquatic vegetation. The limited area of this sort of bottom about Porto Rico doubtless accounts for the scarcity of this species.

Calamus arctifrons Goode & Bean, Proc. U. S. N. M. 1882, 425, Pensacola; Jordan & Evermann, l. c., 1355, 1898.

Genus 88. ARCHOSARGUS Gill. Sheepsheads.

Body robust, short and deep, compressed, covered with large scales. Head deep, mouth moderate, jaws with broad incisors in front and coarse molars on sides; incisors entire or with a shallow notch; posterior nostril slit-like; opercle entire. Dorsal and anal spines strong, soft parts of fin short and rounded; a procumbent spine before dorsal; caudal forked. Gillrakers small. Supraoccipital and temporal crests coalescent anteriorly, both disappearing in the gibbous interorbital area; frontal bone between eyes transversely convex and more or less honeycombed; temporal crest separated from occipital crest by an excavated area, bounded anteriorly by lateral crest, which merges into supraoccipital above eye.

This genus, like *Lagodon*, *Stenotomus*, and *Otrynter*, which show the same character of the procumbent

dorsal spine, is confined to American waters. There are two color types in the genus, one group being made up of species with broad black crossbands, the other of species with golden streaks and inconspicuous crossbands, resembling the species of *Lagodon*. The common sheepshead (*A. probatocephalus*) is the most important member of this genus, but is not yet known from any point in the West Indies.

SALEMA:

- a. Occipital crest rather thin, its honeycomb structure not exposed. Species with streaks of steel-blue and golden, dark crossbands narrow, disappearing with age, about one-third the interspaces; a black humeral spot.
- b. Dorsal spines 13; incisors $\frac{3}{4}$ on each side, side of back with 8 or 9 golden streaks, which are narrower than the metallic-blue interspaces.
- c. Scales 9-48-15; pectoral fin not quite reaching second anal spine; body rather deep and compressed. Incisors $\frac{3}{4}$ on each side, entire, or with a shallow notch. Fifth dorsal spine highest, 2 to 2.5 in head; second anal spine strong, recurved, 2.5 in head. Olivaceous, silvery below, upper parts with golden longitudinal stripes alternating with bluish interspaces; humeral spot larger than eye..... *unimaculatus*, 157
- bb. Dorsal spines 12; incisors $\frac{3}{4}$ on each side. Profile with a slight depression above eye; second anal spine much longer than third. Color, grayish, belly white; 8 golden longitudinal bands; a black shoulder-spot..... *tridens*

ARCHOSARGUS:

- aa. Occipital crest broad, its honeycomb structure plainly exposed at upper margin; dorsal spines 12; species without blue or golden markings, but with about seven broad black crossbands crossing body; no distinct shoulder-spot. Body much compressed; dorsal outline strongly arched; ventral outline almost straight. Profile straight and steep anteriorly. Incisors $\frac{3}{4}$, entire or slightly emarginate, serrate in young; molars in 3 series above, in 2 below, those of inner series larger, those behind incisors very small. Highest dorsal spine 1.33 in head; second anal spine about twice in head, much longer than third.
- d. Incisors broad, their breadth about half their length. Scales 8-48-15..... *probatocephalus*
- dd. Incisors narrower, their breadth 2.5 in their length. Scales 7-44-14..... *aries*

157. *Archosargus unimaculatus* (Bloch). "*Chopa Amarilla*."

(PLATE 26.)

Head 3.4; depth 2.15; eye 4; snout 2.3; maxillary 3; mandible 3.1; interorbital 3; preorbital 4.1; D. xii, 10 to 12; A. iii, 10 or 11; pectoral 6; ventral 1.3; caudal 1; scales 8 or 9-45 to 50-14 to 16. Body ovate, deep, strongly compressed, anterior profile steep and convex; mouth small, maxillary not reaching front of eye; jaws equal, armed with strong, broad incisors, their edges sometimes notched, 3 on each side in upper jaw, 4 on each side in lower; 3 rows of molars of various sizes above, 2 rows below; scales moderate, regular and cycloid, an enlarged scapular scale; dorsal fin continuous, spines strong and sharp, graduated, middle ones highest, 2.5 in head, soft rays low, last longest, 3.6 in head, base of soft fin with a low sheath of small scales; second anal spine strongest, somewhat recurved, 2.5 in head, soft rays short and weak, with basal sheath; pectoral very long, slender, reaching front of anal; caudal well forked, lobes subequal.

Color in life: Pale bluish-silvery, bluish above, side with 9 to 12 narrow brassy lines, 5 above lateral line, somewhat irregular behind, those below broader, more regular, and fainter; head bluish, with brassy bars behind eye; under parts white, throat rusty; dorsal pale-bluish, brassy at base, tips of spines darker; pectoral pale-greenish, washed with rusty, its base with yellow blotch; ventral chiefly orange, inner ray white, tips of other rays white and violet; anal dirty-orange; caudal rusty, washed with olive, middle rays black at tip. In spirits there is a diffuse black shoulder-spot, usually smaller than eye, under lateral line several rows of scales back of its anterior end, faint or invisible in life; the young have 6 or 7 diffuse vertical bars, not observed in life.

Found among the Florida Keys and in the West Indies south to Rio Janeiro, occasionally as far north as South Carolina; recorded from Charleston, Key West, Cuba, Jamaica, and Brazil. It is an abundant and important food-fish in Porto Rico. Our collection contains specimens 4 to 9 inches long, from San Juan, Puerto Real, Guanica, Ponce, and Hucars. In the market at San Juan it was noted as one of the most common species. Though not reaching a large size, it is an excellent pan-fish.

Salma, Marcgrave, Hist. Brasil., 153, 1648, Brazil.

Bream, Browne, Jamaica, 446, No. 1, 1756, Jamaica.

Perca unimaculata Bloch, Ichthyologia, pl. 308, 1792, Brazil; on a figure by Prince Maurice.

Sparus saline Lacépède, Hist. Nat. Poiss., IV, 136, 1803, Brazil; based on *P. unimaculata* of Bloch.

Sargus humeri-maculatus Quoy & Gaimard, Voyage Freycinet, Zool., 297, 1825, Rio Janeiro.

Sargus flavolincatus Cuvier & Valenciennes, Hist. Nat. Poiss., VI, 60, 1830, Cuba.

Cynadus brama Gronow, Cat. Fishes, ed. Gray, 56, 1854, South Carolina.

Sargus caribaens Poey, Memorias, II, 197, 1860, Cuba; Fauna Puerto-Riqueña, 328, 1881.

Archosargus unimaculatus, Jordan & Evermann, l. c., 1359, 1898.

Family XLVII. GERRIDÆ. The Mojarras.

Body oblong or elevated, compressed, covered with large, smooth scales; lateral line continuous, concurrent with back; mouth moderate, extremely protractile, descending when protruded, spines of premaxillary extending to above eye, closing a deep groove in top of head; maxillary without supplemental bone, not slipping under very narrow preorbital, its surface silvery, like rest of head; base of mandible scaly, a slit between it and preorbital to permit its free motion; both jaws with slender, villiform teeth; no incisors, canines, nor molars; no teeth on vomer or palatines; preopercle entire or serrate; sides of head scaly; nostrils double, round; pseudobranchiæ concealed; gillrakers short, broad; gill-membranes separate, free from isthmus; dorsal fin single, continuous or deeply notched, spinous and soft portions about equally developed, with a scaly sheath along base; dorsal spines usually 9 or 10; anal usually with 3 spines; soft portion of fin similar to soft dorsal but shorter; ventral fins thoracic, 1, 5, rather close together, slightly behind pectorals; branchiostegals 6; lower pharyngeal bones close together, often appearing to be united, teeth blunt; air-bladder present; pyloric cæca rudimentary; vertebræ 10 + 14 = 24.

The *Gerridæ* comprise 6 or 8 genera and about 40 species. Four American genera are now recognized, and each is represented in Porto Rican waters by one or more species. They are carnivorous fishes, of moderate or small size, inhabiting the tropical seas, differing considerably in form and in development of spines; the intergradations are, however, very perfect, so that but for the osteological peculiarities of certain species all might be placed in one genus. Oviparous. The larger species of this family are used as food, though they do not seem to be highly esteemed anywhere. In Indian River, Florida, they are rarely used, but about Porto Rico they are in better demand.

- a. Dorsal fin continuous, deeply notched.
- b. Second interhæmal spine singularly developed, as a hollow cylinder, comparatively short and much expanded, posterior end of air-bladder entering its cavity; preopercle and preorbital entire, anal spines 3, second not much enlarged..... EUCINOSTOMUS, 89
- bb. Second interhæmal spine normally developed, not hollow, air-bladder not entering it.
- c. Second interhæmal spine very short, bluntish; anal spines 2, both small; preopercle and preorbital entire..... ULLEMA, 90
- cc. Second interhæmal spine long, spear-shaped; anal spines 2 or 3, second enlarged.
- d. Preopercle entire; second anal spine moderate..... XYSTEMA, 91
- dd. Preopercle serrate; second anal spine much enlarged..... GERRES, 92

Genus 89. EUCINOSTOMUS Baird & Girard. Mojarritas.

Interhæmal of second anal spine greatly modified, expanded into a hollow cylinder, into which posterior end of air-bladder enters. Preopercle and preorbital entire; body comparatively elongate, subelliptical in form; anal spines 3; second anal spine and fourth dorsal spine not greatly enlarged.

Species numerous, in warm seas, remarkable for the peculiar structure of the second interhæmal, which is formed somewhat as in *Calamus*, but much more modified than in that genus.

- a. Premaxillary groove wholly naked, linear or semioval, sometimes constricted at base, but never scaled. anal rays III, 7.
- b. Eye very large, its diameter much greater than length of snout, 2.66 in length of head. Exposed portion of maxillary small, triangular; premaxillary groove linear..... *dowi*
- bb. Eye moderate, usually more than 3 in head, its diameter about equal to length of snout. Exposed portion of maxillary triangular in front, oblong behind.
- c. Body elongate, back little elevated; greatest depth 3.25 to 3.5 in length..... *pseudogula*, 158
- cc. Body more compressed, deeper, back more elevated; greatest depth 2.66 in length.
- d. Snout blunt; eye large, scarcely 3 in head; second anal spine large, 2.66 to 3.3 in head. premaxillary groove linear..... *harcngulus*, 159
- aa. Premaxillary groove scaled in front, scales leaving a naked pit behind. Depth 2.4 in length; head 3 to 3.2 in length of body. Second anal spine about 3.5 in head..... *gula*, 160

158. *Eucinostomus pseudogula* Poey. *Mojarra*.

Head 3.2; depth 3.3; eye 3; snout 3.25; maxillary 3.3; mandible 2; interorbital 3.6; scales 5-49-9; D. IX, 10; A. III, 7. Body compressed, very slender, back very little elevated, the curve a gentle, regular one from nape to caudal peduncle; mouth moderate, nearly horizontal, maxillary reaching anterior edge of pupil, exposed portion triangular; preorbital and preopercle entire, eye large; premaxillary groove rather narrow and without scales; snout rather long and pointed. Fins moderate; dorsal spines

all slender, second not enlarged, 1.66 in head; second anal spine scarcely enlarged, slightly shorter than third, 3.5 in head, the base of fin 2 in head; caudal widely forked; pectoral rather long, scarcely reaching vent, 1.2 in head; ventrals short, 2.2 in head.

Color in life: Bright silvery, with bluish and purplish reflections; back somewhat mottled with dark; fins all pale except tip of spinous dorsal, which is black; axil of pectoral dusky; snout blackish.

Usual length, 4 or 5 inches. Known from the Bermudas and from Cuba to Brazil. Not common in Porto Rico, where four specimens were obtained, all from Mayaguez.

Eucinostomus pseudogula Poey, Enumeratio, 53, pl. 1, 1875, Havana; Jordan & Evermann, l. c., 1368, 1898.
Gerres jonesi Günther, Ann. Mag. Nat. Hist., III, 1879, 150 and 389, Bermudas.

159. *Eucinostomus harengulus* Goode & Bean. *Mojarra*.

Head 3; depth 2.7; eye 3.2; snout 3; maxillary 2.8; mandible 1.8; interorbital 4; scales 5-45-10; D. ix, 10; A. iii, 8. Body compressed, slender, back not much elevated; snout rather long and pointed; mouth horizontal, moderate, maxillary reaching anterior edge of pupil, its exposed portion triangular; preorbital and preopercle entire; premaxillary groove narrow, naked; eye rather large. Fins moderate; second dorsal spine slender, weak, 1.8 in head, shorter than the third; second anal spine short and weak, 3.2 in head, a little shorter than the third; caudal moderately forked, pectoral long and pointed, not quite reaching anal, 1.1 in head; ventrals short, not reaching vent, 1.9 in head.

Color in life: Silvery-white, the back with pale steel-blue iridescence; faint, shining, silvery longitudinal stripes along rows of scales; fins all pale except tips of dorsal spines, which are black.

E. harengulus is very close to *E. californiensis* and may not be really distinct. The principal difference we discover between our specimens and specimens of *E. californiensis* from the Pacific coast is that the second anal spine is somewhat larger in *E. harengulus*. Length 4 to 8 inches. It is found on the Atlantic coast of tropical America from Florida to Bahia; apparently the most abundant species of the family in Porto Rico, as is shown by numerous specimens from San Antonio Bridge and Palo Seco near San Juan, Mayaguez, Boqueron, Fajardo, Isabel Segunda, San Geronimo, and Culebra.

Eucinostomus harengulus Goode & Bean, Proc. U. S. N. M. 1879, 132, West Florida; Jordan & Evermann, l. c., 1368, 1898.

160. *Eucinostomus gula* (Cuvier & Valenciennes). *Mojarra*.

Head 3.2; depth 2.5; eye 3; snout 3.2; maxillary 3; mandible 2; interorbital 3; scales 4-45-9; D. ix, 10; A. iii, 8. Body compressed, rather short, back somewhat elevated; profile straight from tip of snout to occiput, thence arched to origin of dorsal; mouth rather small, horizontal, maxillary reaching anterior edge of pupil, exposed portion narrowly triangular; eye moderate; preorbital and preopercle entire; premaxillary groove scaled except a central, nearly circular pit. Fins moderate; dorsal spines all weak and slender, second and third subequal, 1.75 in head; second anal spine slightly stouter and shorter than third, 3 in head; caudal widely forked, lobes equal to head; pectoral long, reaching past vent, 1 in head; ventrals short, 1.75 in head.

Color, silvery-white, darker on back; fins all pale, front of spinous dorsal dark at tip.

Carolinus to Brazil, the young often taken at Woods Hole, Massachusetts. Common about Porto Rico, specimens having been obtained at San Antonio Bridge, Ensenada del Boqueron, Fajardo, Culebra, and San Geronimo. Length 4 to 5 inches.

Gerres gula Cuvier & Valenciennes, Hist. Nat. Poiss., VI, 464, 1830, Martinique.
Eucinostomus argenteus Baird & Girard, Ninth Smith. Report 1855, 345, Beesley Point, N. J.
Eucinostomus gulula Poey, Enumeratio, 54, pl. 2, 1875, Havana.
Diapterus homonymus Goode & Bean, Proc. U. S. N. M. 1879, 340, Clearwater Harbor, Florida.
Eucinostomus gula, Jordan & Evermann, l. c., 1370, 1898.

Genus 90. *ULÆMA* Jordan & Evermann.

This genus is close to *Eucinostomus*, from which it differs in the form of the second interhæmal, which is short, bluntish, and not hollowed out. The single known species is slender in form, with weak spines, the anal fin having but 2.

161. *Ulæma lefroyi* (Goode).

Head 3.2; depth 3.1; eye 3; snout 3.1; maxillary 3.1; mandible 2; interorbital 3.5; D. ix, 10; A. ii, 8; scales 5-46-10. Body long and slender, dorsal profile only slightly elevated; snout long and pointed; mouth moderate, maxillary reaching only to anterior border of eye, its exposed portion short and triangular; interorbital broad; premaxillary groove long, narrow, and naked; preorbital and preopercle entire; eye large. Fins small; dorsal spines all slender and weak, second slightly shorter than third, 1.9 in head; longest dorsal rays about 3.1 in head; second anal spine small and rather short, 4 in head; caudal fin long, lobes about equal to head; pectoral slender, 1.25 in head; ventrals short, reaching only halfway to anal, 2.1 in head.

Color, silvery, darker above, with bluish iridescence, white below; fine dusky punctulations everywhere, thickest on back; dorsal, anal, and caudal dusky, other fins pale; axil dusky; snout dusky.

Found in the West Indies, on sandy shores; known from the Bermudas, Cuba, Porto Rico, Key West, and Cedar Keys. Length 4 to 6 inches. A specimen, 5 inches long, was obtained at Culebra Island, February 9.

Diapterus lefroyi Goode, Amer. Journ. Sci. Arts 1874, 123, Bermudas.

Eucinostomus productus Poey, Enumeratio, 55, 1875, Havana.

Ulæma lefroyi, Jordan & Evermann, l. c., 1371, 1898.

Genus 91. **XYSTÆMA** Jordan & Evermann. **Mojarras Blancas; Muniamas.**

This genus differs from *Gerres* in having the preopercle entire. The body is compressed, but not greatly elevated, and the second anal and fourth dorsal spines are less enlarged than in *Gerres*. The second interhæmal is long and spear-shaped, not hollow and not receiving air-bladder, its structure as in *Gerres*. One species, widely distributed.




FIG. 61.—*Xystæma cinereum*.

162. *Xystæma cinereum* (Walbaum). *Mojarra; "Muniama."*

Head 3.2; depth 2.6; eye 3.4; snout 3; maxillary 2.6; mandible 2; interorbital 3.25; scales 6-45-11; D. ix, 10; A. iii, 7. Body compressed, elongate, dorsal profile moderately and regularly elevated, interorbital region slightly depressed; mouth moderate, maxillary extending to vertical of anterior edge of pupil, its exposed portion triangular in form and twice as long as wide, its length 5 in head; preorbital and preopercle entire; maxillary groove broadly ovate and without scales; gillrakers weak, 7 below angle; distance from tip of snout to origin of dorsal fin, 2.4 in body; second dorsal spine longest, 1.6 in head, not much longer than the others, all being weak and flexible; general outline of upper margin of dorsal fin falcate; second and third anal spines subequal, second not much enlarged, 2.6 in head; pectoral long and pointed, its length slightly greater than head; ventrals short, 2 in head; caudal deeply forked, lobes equal, 0.7 in head. Scales moderately firm, a sheath at dorsal and anal fins; caudal densely covered with minute scales.

Color in life: Light bluish-white above, silvery below, with about 6 faint, vertical, bluish bars.

Found on both coasts of tropical America, and in the West Indies, north to southern Florida and Lower California; generally common in waters of moderate depth; entering rivers. Length a foot or more. A good fish, of considerable importance, reaching a larger size than most species of the family. Common about Porto Rico, where we obtained specimens from San Juan, San Antonio Bridge, Ensenada del Boqueron, and Isabel Segunda. Common in the San Juan market.

Turdus cinereus peltatus (the Shad), Catesby, Nat. Hist. Carolinas, etc., 1731, Bahamas.

Mugil cinereus Walbaum, Artedi Piscium, 228, 1792, Bahamas; after Catesby.

Gerres aprion Cuvier, Règne Animal, ed. 2, vol. 2, 104, 1829, Bahamas; based on Catesby.

Gerres zebra Muller & Troschel, in Schomburgk, Hist. Barbados, 668, 1848, Barbados.

Gerres squamipinnis Günther, Cat. Fishes, 1, 349, 1859, Jamaica.

Eucinostomus aprion, Poey, Fauna Puerto-Riqueña, 328, 1881; Stahl, l. c., 77 and 163, 1883.

Xystema cinereum, Jordan & Evermann, l. c., 1372, 1898.

Genus 92. GERRES Cuvier. Mojarra.

Second interhæmal long and spear-shaped, not excavated and not receiving end of air-bladder; preopercle serrate; body elevated and more or less rhomboid in form; third or fourth dorsal spine and second anal spine more or less elevated. Species numerous.

a. Preorbital entire; no distinct dark streaks along rows of scales.

MOHARRA:

b. Anal spines 2 only, soft rays 9; second anal spine 1.75; premaxillary groove broad, scaleless; body deep.

rhombus, 163

DIAPTERUS:

bb. Anal spines 3, soft rays 8.

c. Premaxillary groove broad, oval, and covered with scales (these sometimes deciduous in poorly preserved specimens).
Anal III, 8; second dorsal spine 1.25 in head; second anal spine 1.6 in head; teeth rather long and slender.

olisthostomus, 164

GERRES:

aa. Preorbital serrate; a distinct dark streak along each row of scales on back and sides; body rhomboidal, with angular outlines; spines very strong; anal rays III, 8 or 9.

d. Spines moderate, second dorsal spine two-thirds to three-fourths length of head.

e. Pectoral as long as head, not reaching front of anal, 3 to 3.33 in body; scales 38; the longest dorsal spine 1.4 in head..... *brasiliensis*, 165

cc. Pectoral very long, one-sixth longer than head, 2.5 to 2.75 in body; the second dorsal spine longest, 1.5 in head; scales 35..... *embryx*

dd. Spines very high, second dorsal spine equaling head or longer; second anal spine about equal to length of head; lateral stripes very distinct, about 12 in number; depth of the body 2.17 in length; pectoral very long, 2.66 in body; scales 37..... *plumieri*, 166

163. *Gerres rhombus* Cuvier & Valenciennes. Mojarra.

Head 2.8; depth 2; eye 4; snout 3; maxillary 2.4; mandible 1.8; interorbital 3.4; scales 6-38-10; D. IX, 10; A. II, 9. Body much compressed, rhomboidal, back much elevated; profile evenly convex from dorsal to supraorbital, where there is a slight depression; snout somewhat pointed; mouth large, maxillary reaching vertical of posterior border of pupil, exposed portion oblong, its width about 3 in its length, which is 3.4 in head; preorbital and preopercle entire; gillrakers short and weak, 18 below angle; premaxillary groove broadly oval and without scales. Distance from tip of snout to origin of dorsal 2 in length; second and third dorsal spines longest, second strongest, 1.6 in head; dorsal rays short; second anal spine very strong, 1.9 in head; pectoral long, 1.1 in head, reaching origin of anal; ventrals rather long, reaching past vent, 1.6 in the head; caudal well forked, the lobes subequal, 1.1 in the head.

Color in life: Silvery-white with bluish reflections, paler below; snout dusky; no dark lines along rows of scales; margin of dorsal fin black; other fins pale, ventrals and anal somewhat dusky.

Found in the West Indies and along the Atlantic coast of tropical America; generally common. Numerous specimens from San Antonio Bridge, Mayaguez, and Palo Seco. Length 10 inches or less.

Gerres rhombus Cuvier & Valenciennes, Hist. Nat. Poiss., VI, 459, 1830, Martinique; Jordan & Evermann, l. c., 1374, 1898.

Moharra rhombus?, Poey, Fauna Puerto-Riqueña, 327, 1881; Stahl, l. c., 77 and 163, 1883.

164. *Gerres olisthostomus* Goode & Bean. *Mojarra*; *Irish Pompano*; *Mutton-fish*.

Head 3; depth 2.4; eye 3.4; snout 3.4; maxillary 2.6; mandible 1.9; interorbital 3.2; scales 6-39-11; D. ix, 10; A. iii, 8. Body compressed, rather long, rhomboid, more slender than in *G. brasiliensis* dorsal outline in a double curve from tip of snout to origin of dorsal, strongly convex from nape to dorsal fin; snout conical, bluntish; mouth little oblique, large, maxillary reaching anterior edge of orbit; exposed portion of maxillary oblong, length 2.75 times its width; eye moderate; preorbital entire; preopercle weakly serrate; premaxillary groove broad, oval, covered with small, deciduous scales, a naked, linear, median depression. Fins moderate; first dorsal spine very short, the second long and moderately strong, about 2 in head, others gradually shorter; second anal spine strong, 1.6 in head, slightly shorter than third; caudal deeply forked, lobes longer than head; pectoral as long as head; ventrals shorter, 1.4 in head; least depth of caudal peduncle 1.8 in head.

Color, silvery-olivaceous; scales with faint silvery streaks but no dark ones; fins mostly pale or yellowish, ventrals somewhat dusky.

Found in the West Indies, north to southern Florida, south to Brazil. Length a foot or less. Apparently not common in Porto Rico, as the collection contains but one specimen, obtained in the San Juan market. This is 8.5 inches long, and differs somewhat from Florida examples, the second dorsal spine being shorter and the body more slender.

Gerres olisthostoma Goode & Bean, Proc. U. S. N. M. 1882, 423, Indian River, Florida.

Gerres olisthostomus, Jordan & Evermann, I. c., 1376, 1898.




FIG. 62.—*Gerres olisthostomus*.

165. *Gerres brasiliensis* Cuvier & Valenciennes. *Mojarra*.

Head 3.4; depth 2.3; eye 3.4; snout 3.25; maxillary 2.75; mandible 2; interorbital 3.3; scales 6-36-17; D. ix, 10; A. iii, 8. Body compressed, rhomboidal, back very much elevated; profile nearly straight from tip of snout to origin of dorsal fin, slightly concave above eyes; snout conical, bluntish, less acute than in *G. lineatus*; mouth horizontal, rather large, maxillary reaching middle of pupil, its exposed portion oblong, about 2.75 times as long as wide; preorbital and preopercle serrate; eye rather large; premaxillary groove very broad, narrowed posteriorly, without scales. First dorsal spine very short, second very long and strong, 1.2 in head; third slightly shorter but much weaker; remaining dorsal spines short; dorsal rays short, about 2 in head; second anal spine very long and strong, 1.2 in head; third spine slender and slightly longer than second; caudal widely forked, middle rays about 3.5 in outer; pectoral long, 0.9 in head, reaching past vent; ventrals long, 1.15 in head; least depth of caudal peduncle 2.5 in head. Length a foot or less.

F. C. B. 1900-14

Color in life: Pale-silvery, with about six bronze or blackish stripes above lateral line, upper two ceasing at origin of dorsal, and about nine similar but fainter longitudinal stripes below lateral line, upper two of which unite at anterior end; fins all dark except pectoral, which is pale; dorsal fin black-edged; anterior edge of second dorsal spine black; axil dusky.

The range of this species extends from Cuba to Bahia. It is generally common, but our collection contains only two specimens, one from the San Juan market, the other from Puerto Real, 10 and 9 inches long, respectively.

Gerres brasiliannus Cuvier & Valenciennes, Hist. Nat. Poiss., VI, 458, 1830, Brazil; Poey, Fauna Puerto-Riqueña, 327, 1881; Jordan & Evermann, l. c., 1378, 1898.

166. *Gerres plumieri* Cuvier & Valenciennes. *Mojarra*.

Head 3; depth 2.16; eye rather large, 3 in head; snout 4 in head; scales 5-37-11; D. 1x, 10; A. 11, 8. Body compressed, rhomboidal in form, back very much elevated. Mouth rather large, maxillary extending slightly beyond vertical from anterior margin of pupil, its length 2.8 in head; exposed portion of maxillary oblong, its width 2.5 in length, which is 4.75 in length of head; preorbital and preopercle serrate; premaxillary groove broad and entirely free from scales; gillrakers small, weak, 13 below the angle; distance from tip of snout to dorsal fin equal to greatest depth of fish; upper margin of dorsal fin much concave; second dorsal spine very strong and long, its length equaling length of head; second anal spine stronger and slightly shorter than second dorsal spine, its tip reaching to vertical from base of caudal rays; third spine shorter and much weaker than second; pectoral fins reaching beyond front of anal, their length 2.66 in length of body; ventral fins reaching past vent, almost to front of anal, their length 3.4 in length of body. Length 10 inches.

Color, bluish-silvery above, silvery below; very distinct dark longitudinal lines along each row of scales; dorsal, caudal, and anal fins dusky; margin of dorsal fin black; a dark supraorbital spot; pectoral and ventral fins pale.

Found on the Atlantic coast of tropical America and among the West Indies; rather common; known from Havana, Porto Rico, Santo Domingo, Jamaica, Martinique, Indian River, Florida, Pernambuco, Bahia, Aspinwall, and Guatemala. Not seen by us in Porto Rico, but included here on the authority of Cuvier & Valenciennes and of Poey.

Gerres plumieri Cuvier & Valenciennes, Hist. Nat. Poiss., VI, 452, 1830, Antilles, Porto Rico; Poey, Fauna Puerto-Riqueña, 327, 1881; Jordan & Evermann, l. c., 1379, 1898.

Family XLVIII. KYPHOSIDÆ. The Rudder-fishes.

Herbivorous fishes, with incisor teeth only in front of jaws. Body oblong or elevated, with moderate or small scales, ctenoid or not. Mouth moderate, with incisor-like teeth in front of each jaw; no molars; teeth on vomer and palatines present or absent; premaxillaries moderately protractile; preorbital rather narrow, sheathing maxillary. Gillrakers moderate; pseudobranchiæ well developed; opercles entire. Gills 4, a slit behind fourth; gill-membranes separate, free from isthmus; dorsal fin continuous or divided, with 10 to 15 rather strong spines, soft dorsal naked or scaly; anal with 3 spines; ventrals thoracic, rays 1, 5, an accessory scale at base; caudal lunate or forked; pectoral fin with all its rays branched. Intestinal canal elongate, with few or many pyloric cæca. Air-bladder usually with 2 posterior horns. Vertebrae in ordinary or slightly increased number, 24 to 28. Post-temporal of normal percoid form, the stout forks not adnate to cranium.

Shore fishes, feeding largely on green or olive algae; chiefly of the Mediterranean Sea and the Pacific Ocean; most of them valued as food. Genera 20, species about 70.

Genus 93. KYPHOSUS Lacépède. Chopas.

Body elongate-ovate, regularly elliptical, moderately compressed; head short, with blunt snout; eye large; mouth small, horizontal; maxillary barely reaching front of eye; each jaw with a single series of rather narrow, obtusely lanceolate incisors, implanted with compressed conspicuous roots posteriorly; behind these a narrow band of villiform teeth; fine teeth on vomer, palatines, and tongue. Branchiostegals 7; gillrakers long. Preopercle obsolete serrate; preorbital narrow, covering but little of the maxillary. Squamation very complete, space between and about eyes being the only naked part; scales smallish, thick, ctenoid, 60 to 70 in lateral line, which is continuous; similar scales entirely covering soft parts of vertical fins and extending up on paired fins. Dorsal fin low, with about 11

spines, which are depressible in a groove of scales, fin continuous, but last spines low, so that a depression occurs between the two parts of fin, bases of spinous and soft parts about equal; soft dorsal rather low in front, not falcate, pointed behind; anal similar to soft dorsal, with 3 spines; caudal fin moderately forked; pectoral fin small, ventrals well behind them. Intestinal canal long. Pyloric cæca very numerous. Vertebrae 9 or 10+15 or 16=25.

This genus contains some 10 species, chiefly confined to the Pacific Ocean, and most of them found in the East Indies.

- a. Anal fin moderately elevated in front and rather short, its rays III, 11, longest ray 1.5 to 2 in base of soft part of fin. D. XI, 11 or 12. Teeth 35 to 40 in each jaw.
- b. Teeth rather narrow and subacute; maxillary short, barely reaching eye, about 3.5 in head.
- c. Scales moderate, 10-65-20; A. III, 13. Coloration bright-plumbeous, with many bright-yellow streaks on a plumbeous ground. Mouth and teeth not fully described. *incisor*, 167
- cc. Scales rather large, 10-55-16; depth 2.33 in length; head 3.75. Coloration dusky-gray, with about 25 gray streaks following rows of scales, those near middle of body broadest; a silvery streak along preorbital. D. XI, 12; A. III, 11. *sectatrix*, 168

167. *Kyphosus incisor* (Cuvier & Valenciennes). *Chopa Amarilla*.

Head 5 in total with caudal; depth 3; D. XI, 14; A. III, 13; scales 10-65, pores 20; anal a third higher than soft dorsal; caudal lunate; teeth as in *K. sectatrix*; preopercle scarcely serrate; scales of back much smaller than those of sides, as are also those of head, throat, and belly. Plumbeous, with yellow lines marking edges of scales; besides yellow streak across cheek to axil, another below eye ending on opercle; yellow streak before nostrils emphasized by black edgings; fins blue, deeper on soft part; base of pectoral with yellow scales.

Cuba (Poey) to Brazil and Canary Islands; a large species, reaching 2.5 to 3 feet in length, larger than *K. sectatrix*. Evidently distinct from *K. sectatrix*, with which it is confounded by Jordan & Fesler, its relations being with *K. analogus*. Not seen by us; reported by Dr. Stahl from Porto Rico.

? *Salema aurata* Bowdich, Excursion Madeira, 238, 1825, Bona Vista Island; description and figure very bad. D. X, 17; A. III, 14; body with light-orange stripes.

Pimblepterus incisor Cuvier & Valenciennes, Hist. Nat. Poiss., VII, 266, 1831, Brazil.

Pimblepterus flavolineatus, Stahl, l. c., 77 and 163, 1883.

Kyphosus incisor, Jordan & Evermann, 1386, 1898.

168. *Kyphosus sectatrix* (Linnæus). *Rudder-fish; Bermuda Chub; Chub; Chopa Blanca*.

Head 3.75; depth 2.12; D. XI, 12; A. III, 11; scales 10-55-16; vertebrae 9+16. Body ovate, somewhat compressed; longest dorsal spine one-fifth height of body, rather higher than soft dorsal and nearly equal to longest ray of anal; teeth 35 to 40 on each side; horizontal process of teeth not much longer than vertical; interorbital space 2.5 in head; top and sides of head finely scaled; interorbital region gibbous, below which point snout is truncate; preopercle weakly serrulate; gillrakers long; soft dorsal and anal very low; longest ray of anal 2.25 in head, longest spine 2.25; median dorsal spines highest; second anal spine highest; caudal well forked, lower lobe longer.

Color in life: Steel-gray, very slightly bluish, not much paler below; edges of each row of scales on back and sides slightly brassy, so that very faint yellowish stripes alternate with bluish ones of about equal width; a diffuse pale stripe below eye, a yellowish one above and below this; fins all dull-grayish; ventrals and anal somewhat blackish; edge of opercle slightly darker. Here described from Key West specimens.

K. sectatrix ranges from Cape Cod to the West Indies, crossing the ocean to the Canary Islands; accidental in the Mediterranean; once taken at Palermo by Prof. Pietro Doderlein. Not rare off our Atlantic coast, becoming rather common southward, especially at Key West. Long noted for its habit of following vessels, supposably for waste food thrown from them, hence called rudder-fish. It is apparently not common about Porto Rico, and was not obtained by us, but is recorded by Professor Poey and Dr. Stahl.

This fish reaches a length of 18 inches or more, though the average of those caught is much less. The average weight is 3 or 4 pounds and the maximum about 9 pounds. As a food-fish it is held in high esteem, its flesh being firm and of good flavor. At Key West it occurs in certain places in abundance and is found throughout the year. The particular bottom which it seeks is the shoals in

or near the channels where the water is 4 to 20 feet deep. It is one of the best and most interesting game-fishes of our southern waters. It is remarkable for the vigor and "rush" with which it takes the hook and for the mad dashes it makes to free itself. The fight which it makes is much like that one sometimes gets from an unusually large fresh-water sunfish, but is stronger, more prolonged, and more exciting, as the fish is larger and stronger.

Chub fishing at Key West, though not known to everyone who has gone to those waters to wet a line, is really one of the surest and most satisfactory ways of enjoyment the angler can find at that place. The chub is caught by still-fishing and the usual bait is pieces of the soft parts of the large spiny lobster (*Panulirus americanus*), which is very abundant about Key West. The fish swim at various depths, but usually not far above the bottom. The water is very clear and the fact that the angler can see the fish, as it rushes frantically in one direction and then in another to free itself, adds not a little to the excitement and interest of the sport.

Perca marina sactatrix (the Rudder-fish), Catesby, Nat. Hist. Car., 138, 1731, Carolina.

Perca saltatrix Linnæus, Syst. Nat., ed. X, 293, 1758 (misprint, incorrectly copied from Catesby, who called it *sactatrix*), Carolina.

Chatodon cyprinaceus (Broussonet) Gmelin, Syst. Nat., I, 1269, 1788, name only; on a specimen from the tropical Atlantic (coll. Parkinson, in voyage of Capt. Cook; described by Cuvier & Valenciennes, l. c., VIII, 263).

Pimblepterus bosquii Lacépède, Hist. Nat. Poiss., IV, 429, 1803, South Carolina.

Pimblepterus oblongior Cuvier & Valenciennes, Hist. Nat. Poiss., VII, 264, 1831, locality unknown; depth 3 in total length; 14 longitudinal streaks.

Pimblepterus bosci var. *sicula* Doderlein, Nat. Sicil., Ann. 11, fasc. 2, 1883, Palermo.

Pimblepterus boscii, Poey, Fauna Puerto-Riqueña, 330, 1881; Stahl, l. c., 77 and 163, 1883.

Kyphosus sactatrix, Jordan & Evermann, l. c., 1387, 1898.




FIG. 63.—*Kyphosus sactatrix*.

Family XLIX. SCIÆNIDÆ. The Croakers.

Body compressed, more or less elongate, covered with rather thin scales, which are usually more or less ctenoid. Lateral line continuous, usually more or less concurrent with back, extending on caudal fin. Head prominent, covered with scales; bones of the skull cavernous, muciferous system highly developed, surface of skull, when the flesh is removed, very uneven. Suborbital bones without backward projecting "stay." Chin usually with pores, sometimes with barbels. Mouth small or large, the teeth in one or more series, outer of which are sometimes enlarged; canines often present. No incisor nor molar teeth; no teeth on vomer, palatines, pterygoids, nor tongue. Maxillary without supplemental bone, slipping under free edge of preorbital, which is usually broad. Premaxillaries protractile, but not very freely movable. Nostrils double. Pseudobranchiae usually large, present in most of the genera. Gills 4, a slit behind fourth. Gillrakers present. Branchiostegals 7. Gill-membranes separate, free from isthmus. Lower pharyngeals separate or united, often enlarged, teeth conic or molar. Preopercle serrate or not. Opercle usually ending in 2 flat points. Dorsal fin deeply notched or divided into 2 fins, soft dorsal being the longer, spines depressible into a more or less perfect groove. Anal fin with 1 or 2 spines, never more than 2. Ventral fins thoracic, 1, 5, below

or behind pectorals. Pectoral fin normal. Caudal fin usually not forked. Ear bones very large. Pyloric caeca usually rather few. Air-bladder usually large and complicated (wanting in *Menticirrhus*).

An important family of 30 genera and about 150 species, found on sandy shores in all warm seas, a few being confined to fresh waters. None occurs in deep water and none among rocks. Many of them reach a large size, and nearly all are valued for food. All are carnivorous and some are of interest as game-fishes. Most of the species make a peculiar noise, called variously croaking, grunting, drumming, and snoring; this sound is supposed to be caused by forcing air from the air-bladder into one of the lateral horns.

OTOLITHINÆ:

- I. Vertebrae 14 or 15 + 10 or 11, abdominal portion of spinal column having always more vertebrae than caudal portion, anal fin being posterior in its insertion; body more or less elongate, mouth large, lower jaw projecting; preopercle with a crenulate, membranaceous border; snout without distinct pores or slits; no barbels, preorbital narrow; gillrakers slender, moderate, or rather long; anal fin with 1 or 2 very weak spines, second closely connected with first soft ray; scales small, smoothish.
- a. Anal fin long, of 15 to 21 soft rays, its length more than half that of soft dorsal; dorsal fins more or less separated; soft dorsal and anal fins closely scaled.
- b. Teeth large, very unequal; tip of upper jaw with one or two strong canines; enlarged teeth or canines on sides of lower jaw; anal fin one-fourth shorter than soft dorsal, 15 to 18 soft rays; dorsal fins well separated, interspace about equal to eye; soft dorsal of 21 rays; body compressed; scales rather small, cycloid. ISOPISTHYS
- aa. Anal fin moderate, or short, of 7 to 13 soft rays, its length less than one-half that of second dorsal; dorsal fins contiguous; lateral line arched in front.
- c. Canine teeth, if present, not lance-shaped, tapering from base to tip.
- d. Lower jaw without canines at its tip, some of its lateral teeth sometimes enlarged; tip of upper jaw usually with canines. CYNOSCION, 94
- cc. Canine teeth lance-shaped, widened toward tip, then abruptly pointed; canines of front of premaxillary largest; about 2 canines on front of lower jaw on each side; outer teeth of upper jaw enlarged, somewhat lance-shaped; outer teeth of lower jaw compressed; air-bladder with 2 horn-like processes; gillrakers moderate, slender; soft dorsal and anal fin scaly. SAGENICHTHYS
- II. Vertebrae 9 to 12 + 13 to 20, typically 10 + 14, number in abdominal part of body being always fewer than in caudal part; dorsal fins contiguous, soft dorsal being long, much longer than anal.
- c. Dorsal spines well separated, first dorsal spine attached to third or fourth interneural, not more than 2 of spine-bearing interneurals being placed between same pair of vertebrae; soft rays of dorsal fin usually 17 to 32 (37 to 40 in *Lonchirus*, 45 to 50 in *Sciænoides*); occipital crest not greatly elevated.

SCIÆNINÆ:

- f. Lower pharyngeals separate.
- g. Lower jaw without barbels.
- h. Caudal fin moderately scaly, its distal portion usually more or less naked, scales not numerous enough to give a thickened appearance to fin.
- i. Teeth well developed, permanent in both jaws.
- j. Lower pharyngeals rather narrow, their teeth conic and mostly sharp, none of them molar; outer teeth of upper jaw more or less enlarged.
- k. Gillrakers comparatively long and slender; mouth more or less oblique; anal fin usually (but not always) inserted posteriorly; preorbital usually narrow, flat; edge of snout above upper jaw with pores and slits little conspicuous or obsolete.
- l. Skull excessively cavernous, soft and spongy to touch, interorbital space very broad; eye very small; mouth large, oblique; preopercle with a broad membranaceous border, which is striated and fringed; scales small; spinous dorsal short and weak; anal spines weak; caudal fin pointed.
- m. Pseudobranchiæ present; teeth subequal, all villiform, in narrow bands; soft dorsal long, of 30 to 35 rays; anal fin rather long, soft dorsal and anal scaly; lower jaw projecting; vertebrae 10 + 14; gillrakers long and slender; air-bladder with two horns. NEBRIS
- ll. Skull firm, not excessively cavernous, interorbital space less broad; preorbital not turgid; soft dorsal of fewer than 30 rays.
- n. Scales of lateral line considerably enlarged, almost entirely concealed by smaller ones; anal fin small, inserted well forward, its first spine usually as near ventrals as caudal; caudal fin pointed, its peduncle long and slender; soft dorsal and anal scaly; scales small; preopercle without bony serræ; pseudobranchiæ small, often obsolete on one side. (Fluviatile species.) PLAGIOSCION
- mm. Scales of lateral line similar to the others, not concealed by smaller ones; anal fin inserted more or less posteriorly, first spine usually nearer caudal than ventrals; caudal peduncle rather short; pseudobranchiæ well developed.
- o. Head not very broad, the interorbital space not notably spongy nor deeply cavernous.
- p. Preopercle with its membranaceous edge entire, crenulate or eiliate, with no bony teeth; teeth in lower jaw in few series.
- q. Teeth very small, equal, uniserial or very nearly so; snout very short; cleft of mouth oblique or even vertical, lower jaw projecting. LARIMUS, 95
- qq. Teeth larger, more or less unequal, those of lower jaw in one or two series or in bands; cleft of mouth not vertical.
- r. Upper jaw with some of the teeth enlarged, forming canines; some canines in lower jaw; lower jaw projecting.

- rr. Upper jaw with narrow band of teeth, those of outer row more or less enlarged; no distinct canines...CORVULA, 97
- pp. Preopercle with its bony margin armed with sharp teeth or serræ.
- s. Preopercle with its lowermost spine directed abruptly downward; soft dorsal and anal fin moderately scaly; lower jaw without canines; second anal spine moderate or large BAIRDIELLA, 98
- oo. Head very broad above, interorbital space flattish, excessively cavernous, septa reduced to thin partitions; soft dorsal and anal fin usually densely scaly; second spine of dorsal usually thickened STELLIFER
- kk. Gillrakers comparatively short and thick, usually not longer than posterior nostril; anal fin inserted farther forward; snout above lower jaw with large pores, and with two more or less distinct slits on its edge, these sometimes obsolete; preorbital more or less broad; mouth more or less inferior.
- l. Preopercle with its bony margin armed with strong persistent spines which do not disappear with age; caudal fin not lunate, middle rays longer than lower OPHIOSCION, 99
- tt. Preopercle with its bony margin serrate in young, becoming entire with age; caudal fin truncate or lunate, middle rays not longest; slits and pores of upper jaws well developed..... SCLENOPS
- ii. Teeth very small, subequal, those in lower jaw wanting or deciduous; lower pharyngeals rather broad, with paved teeth; mouth small, inferior; snout as in *Sciæna*; preopercle entire; anal fin long, with about 12 soft rays; gillrakers shortish, rather slender LEIOSTOMUS
- gg. Lower jaw with one or more barbels, either at symphysis or on the rami; snout with slits and pores as in *Sciæna*; lower jaw included; preorbital broad; lower teeth in villiform bands; gillrakers more or less short.
- u. Pseudobranchiæ well developed; pectoral fin not elongate.
- vv. Lower jaw with slender barbels, usually several in number.
- w. Barbels mostly in a tuft at symphysis of lower jaw; mouth very small, inferior; gillrakers minute, thickish; dorsal spines X or XI; preopercle sharply but finely serrate; preorbital turgid and cavernous, more or less translucent; caudal fin rhombic. (Fluviatile species) PACHYPOPS
- www. Barbels chiefly lateral, along rami of lower jaw, usually none at symphysis; lower pharyngeals narrow, with sharp teeth; preopercle with its bony margin armed with strong teeth; D. X or XI; gillrakers short, thickish. MICROPOGON, 100
- vv. Lower jaw with a single thickish barbel at its tip.
- x. Air-bladder large; anal spines 2; back more or less elevated; preopercle with its bony margin crenate or serrate; pectoral short, shorter than ventrals. (Free-swimming species) UMBRINA, 101
- xx. Air-bladder none; anal spine single, weak; back not elevated; preopercle with its membranaceous edge crenulate; pectoral fins long, longer than ventrals. (Bottom fishes.) MENTICERRHUS, 102
- uu. Pseudobranchiæ obsolete; body long and low; caudal pointed; pectoral fin elongate; preopercle without bony serratures.
- y. Mandible without barbels along inner edge. Chin with two short barbels; soft dorsal with 30 to 40 rays. LONCHIURUS
- APLODINOTINE:
- ff. Lower pharyngeals very large, completely united, covered with coarse, blunt, paved teeth; lower jaw included; snout with slits and pores, as in *Sciæna*; gillrakers rather short.
- z. Lower jaw with numerous barbels along inner edge of rami; preopercle nearly entire POGONIAS

EQUITINE:

- ee. Dorsal spines close together, first spine attached to first interneural, and from 5 to 12 of spine-bearing interneurals wedged in between high occipital crest and neural spine of second vertebra on the one hand, and that of third vertebra on the other; occipital crest much elevated; soft dorsal very long, of 36 to 55 rays..... EQUUS, 103

Genus 94. CYNOSCION Gill. Weak-fishes.

Body elongate, little compressed, back not elevated. Head conical, rather pointed; mouth very large, terminal, not very oblique, lower jaw projecting, symphysis produced, angle at base of maxillary not prominent. Maxillary very broad. Teeth sharp, not closely set, in rather narrow bands; tip of lower jaw without canines; upper jaw with 2 long canines, 1 of which is sometimes obsolete; canines tapering from base to tip; lateral teeth of lower jaw larger than anterior. Preopercle with its membranaceous edge serrulate, bone entire. Lower pharyngeal bones separate, their teeth all pointed. Gillrakers strong, rather long. Vertebrae about 14 + 10 (instead of 10 + 14 as in *Sciænoideis* generally). Pseudobranchiæ well developed; dorsal spines slender, fins closely contiguous; anal spines 1 or 2, very feeble, soft rays 7 to 13; second dorsal long and low, more than twice length of anal; ventrals inserted below pectorals, pubic bone long and strong; caudal fin subtruncate or lunate.

Large fishes, chiefly of the waters of America, closely related to the Old World genus *Otolithus*, from which they are distinguished by the absence of canines in the lower jaw. All of them rank high as food-fishes; the flesh is rich, but in some species tender and easily torn; hence the popular name weak-fishes. The number of species of this genus known from American waters is about 20, among which are some of our most important food-fishes, including the common squeteague (*Cynoscion regalis*), which, though not exceeding 5 or 6 pounds in weight, is said occasionally to attain a weight of 30 pounds, and the spotted squeteague (*C. maculatum*), somewhat smaller and ranging further south. In the Gulf of California occurs another species (*C. macdonaldi*), which attains the enormous size of more than 170 pounds. Only one species of this genus is at present known from Porto Rico.

- a. Scales not very small, lateral line having 55 to 75 pores, number of transverse series ranging from 55 to 85, not much in excess of number of pores; head compressed, not truly conical; upper jaw with distinct canines, band of teeth in upper jaw rather narrow, lower teeth small and in few series in front, larger and uniserial on sides.
- b. Soft rays of dorsal and anal more or less closely scaled; gillrakers comparatively long and slender, 9 to 12 on lower part of arch, longest at least one-half diameter of eye.
- c. Soft dorsal of 20 to 23 rays.
- d. Caudal fin rhombic, middle rays considerably produced.
- e. Snout short, bluntish, 4.6 in head; mouth small, little oblique, canines quite small; color pale, with faint darker streaks; axil pale, pseudobranchiæ sometimes wanting. D. IX-1, 20; A. I, 8. *acoupa*
- cc. Soft dorsal of 26 to 29 rays; caudal fin subtruncate or double truncate, middle rays but slightly produced.
- f. Coloration nearly uniform silvery.
- g. Caudal truncate; body slender, depth more than 4 in length; snout short; maxillary not reaching beyond eye. D. X-1, 27; A. I, 11 *obliquatus*
- gg. Caudal weakly double-concave; body deep, depth 3.5 to 4 in length.
- h. Snout long, 3.75 in head, longer than eye. *jamaicensis*, 169
- hh. Snout long, 4.5 in head, shorter than eye. D. X-1, 27 to 29, A. II, 9 or 10. *nothus*
- ff. Coloration brownish-silvery above, with many dark-brown spots, arranged in undulating streaks; body more or less compressed; eye moderate, 5 to 7 in head; maxillary extending to below posterior margin of eye, 2.17 in head; canines large; color brownish-silvery, with iridescent reflections and marked with many small, rather irregular, dark-brown spots, some of which form undulating lines running upward and backward; upper fins dusky, lower yellowish.
- i. Snout not very sharp, about 4.25 (4 to 4.33) in head; gillrakers long and slender, usually 5 + 10 to 12 in number; membranes of soft dorsal and anal more or less closely scaly, scales readily deciduous *regalis*
- ii. Snout very sharp, 3.75 to 3.8 in length of head; gillrakers shorter, rather slender, 4 + 8 or 9 in number; membrane of soft dorsal and anal with very few scales, these readily deciduous *thalassinus*
- bb. Soft rays of dorsal and anal scaleless; gillrakers comparatively short and thickish, usually not longer than pupil, and but 6 to 8 on lower limb of arch.
- j. Coloration not uniform, grayish and silvery, back with distinct darker spots, lines, or reticulations; caudal fin truncate or slightly double-concave.
- k. Soft dorsal fin with conspicuous round black spots; back and sides covered with similar spots smaller than pupil, larger than those on fins; snout acute, much longer than eye; pectoral 2.25 in head. D. X-1, 25 to 27; A. II, 10. *nebulosus*
- aa. Scales comparatively small; number of pores in lateral line 70 to 90, and very much fewer than number of transverse rows, which is from 85 to 150; teeth of upper jaw in a rather broad band, 1 to 4 of them usually more or less canine-like, canines generally small and sometimes wholly disappearing with age; lateral teeth of lower jaw not much enlarged; gillrakers usually small and short.
- l. Caudal fin lunate or subtruncate; scales not very small; head more or less distinctly conical, not flattened above; soft dorsal with 21 to 23 rays.
- m. Soft dorsal fin with its lower portion covered with small, caducous scales. Body compressed; head compound-conic; canines small, both present; pectoral 2 in head; caudal weakly double-truncate. D. IX-1, 21 to 23; A. II, 10. *leiarachus*
- ll. Caudal fin rhombic or S-shaped, middle rays produced, upper lobe usually pointed; soft dorsal with 23 to 28 rays.
- n. Soft dorsal entirely naked; anal with a few scales; body long and low, spindle-shaped; head depressed above; mouth large; canines present, short and thick; eye small, caudal S-shaped, middle rays longest; pectoral 1.8 in head; scales very small. D. X-1, 28; A. I, 8. *virescens*
- mm. Soft dorsal and anal fins densely scaly throughout; teeth all small, canines moderate; scales very small; pectoral 1.75 in head; caudal S-shaped; color greenish, silvery below. D. XI-1, 23; A. II, 9 *microlepidotus*

169. *Cynoscion jamaicensis* (Vaillant & Bocourt). "Corvina"; Mongolar Drummer.

Head 3.3; depth 4; eye 4.7; snout 3.8; maxillary 2.1; mandible 1.7; interorbital 3.6; preorbital 15.5; D. X-1, 25; A. II, 10; pectoral 1.8; ventral 2; caudal 2.1; scales 11-76-8. Body elongate-elliptical, considerably compressed; head pointed; snout a little longer than eye; mouth large, lower jaw projecting; teeth in narrow bands, a few of the outer ones enlarged, especially on sides of lower jaw, where they are recurved and incurved; one strong recurved canine in front of upper jaw; lateral line decurved under last dorsal spines. Color in life, silvery-grayish, bluish above, whiter below; fins all pale except edge of soft dorsal, which is dark.

Known only from Jamaica and Porto Rico. Two specimens, about 11 inches long, from San Juan market, January 6, and two young ones, 5.5 and 4.5 inches, from Isabel Segunda, February 8, 1899. The dentition of the upper jaw exhibits an interesting condition. The single large canine tooth of the upper jaw is located at one side of the symphysis; on the other side is a tooth of similar appearance and nearly as large, but soft and flexible. Apparently there is structurally a pair of canines in the front of upper jaw, but only one becomes functional. Both specimens exhibit this peculiarity; in one the developed tooth is upon the right side of the symphysis, in the other on the left.

Otolithus jamaicensis Vaillant & Bocourt, Miss. Sci. au Mexique, Poissons, 156, 1874, Jamaica.
Cynoscion jamaicensis, Jordan & Evermann, I. c., 1406, 1898.

Genus 95. *LARIMUS* Cuvier & Valenciennes.

Body rather elongate, compressed; skull firm, not greatly cavernous; interorbital space rather narrow; preorbital flattish, not turgid; upper jaw with usual slits and pores little developed; no barbels; no canines; snout very short, mouth large, terminal, very oblique or even vertical, lower jaw projecting; teeth minute, equal, uniserial or partly biserial above; preopercle entire or nearly so, without bony teeth. Scales moderate, subequal. Pseudobranchiæ well developed. Fins essentially as in *Bairdiella*, second dorsal long, anal short, its spines moderate or small; fins not thickened by accessory scales. Gillrakers long and slender. Vertebrae 10+14. Silvery fishes, all American.

- a. Mouth more or less oblique, not quite vertical; upper part of body with dark streaks along rows of scales; profile slightly convex, a little oblique; maxillary extending to below front of orbit, 2 in head.
- b. Dorsal rays 27 to 30; mouth notably oblique.
- c. Upper parts with distinct dark streaks along rows of scales.
- d. Second anal spine 1.66 in head, reaching tips of soft rays; dark streaks on sides not very distinct; mouth very oblique; gill-cavity pale..... *breviceps*, 170
- bb. Dorsal rays 24 to 27; mouth still less oblique, snout more convex, profile descending forward.
- e. Color grayish, silvery with about 7 dark vertical crossbars; second anal spine short, 3.25 in head. Body heavy forward, much compressed, snout very short and blunt, 5.5 in head; mouth large, less oblique than in other species; tip of premaxillary on level of middle of pupil; maxillary 2 in head; gillrakers extremely elongate, as long as eye, 12+24; second anal spine short, one-fourth shorter than the first soft ray. D. x-1, 24 to 26; A. 11 5 or 6..... *fasciatus*

170. *Larimus breviceps* Cuvier & Valenciennes. "*Cabezón*"; "*Corbino Cabezón*."

Head 3.2; depth 3; eye 3.8; snout 5.1; maxillary 2.1; mandible 1.8; interorbital 4; preorbital 11; D. x-1, 27; A. 11, 6; pectoral 1.1; ventral 1.4; scales 6-50-9. Body oblong, heavy forward, back much compressed, caudal peduncle long and slender; scales thin and ctenoid; head short; mouth large, very oblique, approaching a perpendicular, maxillary reaching front of pupil or beyond, lower jaw heavy and prominent, its tip projecting; snout very short; preopercle unarmed; teeth small, in one row in each jaw; head scaled; gillrakers long and slender, 13+20; fins small, dorsal spines weak and flexible, soft dorsal long and low, anal very small, second spine much enlarged, reaching about to tip of longest soft rays; caudal double truncate.

Color in spirits: Pale, back dark-olivaceous; membrane of spinous dorsal dusky; caudal streaked with dusky; axil with some black. In life the fins showed the following colors: Dorsal darkish, caudal pale, yellow above and below; anal yellow next to spine, pale elsewhere; ventral yellow on outer half, other half pale; pectoral pale-lemon.

Found in the West Indies, and south to Brazil; known from Jamaica, St. Lucia, and Porto Rico. Length a foot or less. A good food-fish. Three adults, about 8.5 inches in length, from San Juan market, and five young, 3.5 to 5 inches, from Ponce and Isabel Segunda.

Larimus breviceps Cuvier & Valenciennes, Hist. Nat. Poissons, V, 146, 1830, Santo Domingo and Brazil; Jordan & Evermann, l. c., 1423, 1898.

Monosira stahli Poey, Fauna Puerto-Riqueña, 326, pl. 6, 1881, Porto Rico; Stahl, l. c., 163, 1883.

Larimus stahli, Eigenmann & Norris, Sobre alguns peixes de S. Paulo, Brazil, 361, 1900.

Genus 96. *ODONTOSCION* Gill.

This genus differs from *Larimus* mainly in the presence of canines, and may be described as a *Larimus* armed with canine teeth. It also approaches closely to *Bairdiella*, from which it differs in lacking the plectroid spine on the preopercle and also in the dentition, the group *Elattarchus* lying between the two, as does also the closely related group *Corvula*.

171. *Odontoscion dentex* (Cuvier & Valenciennes). *Corvina*.

Head 3; depth 3.4; eye 3.7; snout 4; maxillary 2.25; mandible 2; interorbital 4.6; preorbital 3.8 in eye; D. XI-1, 24; A. 11, 9; pectoral 1.8; caudal 1.4; scales 6-56-8. Body rather elongate, back compressed and slightly elevated; anterior profile nearly straight but not very steep; caudal peduncle long and much compressed; head moderate, snout short and blunt; eye large and placed high; teeth in a single row in each jaw, long and sharp, the pair in front of lower jaw enlarged; teeth of upper jaw similar to lower but smaller, largest in front but without distinct canines; preopercle entire; maxillary reaching middle of pupil; gillrakers about 7+15 (in individual 4 inches long); soft parts of

vertical fins scaly; caudal truncate; dorsal spines very slender, separated from rays; second anal spine very slender, 3 in head, much longer than first, which is much reduced.

Color in spirits: Silvery-bronze, with many dark points, under parts pale; darker longitudinal streaks along rows of scales; chin dusky, axil and base of pectoral black; vertical fins dusky, with dark points, paired fins pale.

Here described from the young, of which 15 examples were taken, from 3 to 4.75 inches in length, at Mayaguez, Puerto Real and Boqueron. Length about a foot. A food-fish of some importance. Known from the West Indies, and generally common; recorded by Jordan & Rutter from Jamaica.

Corvina dentex Cuvier & Valenciennes, Hist. Nat. Poiss., V, 139, 1830, Santo Domingo.

Odontoscion dentex, Jordan & Evermann, I. c., 1425, 1898.

Genus 97. *CORVULA* Jordan & Eigenmann.

This genus is closely allied to *Bairdiella* in nearly all respects, but with preopercle entire and unarmed, as in *Larimus*. The species differ considerably among themselves, and they form with *Larimus* and *Odontoscion* an almost continuous series. American.

- a. Body rather short and deep, depth 2.5 to 3.33 in length; distance from insertion of ventrals to first anal spine about equal to depth of body; color silvery, usually with dusky streaks along rows of scales.
- b. Dorsal rays x-1, 28; posterior rays of soft dorsal higher than anterior ones; dorsal outline strongly and regularly convex and elevated. Color, silvery-white, darker above; sides and back with rather distinct dark lines along scales; spinous dorsal, tips of ventrals, and anal dusky; upper part of head brownish; lower part of head, cheek, and breast with numerous rusty dots, base of soft dorsal and anal rusty..... *sialis*
- bb. Dorsal rays x to XII-1, 23 to 25; jaws equal; outer teeth above enlarged, lower teeth nearly uniserial; preopercle with flexible serræ; second anal spine, 3.66 in head; caudal fin subtruncate.
- c. Maxillary reaching middle of pupil, 2.33 in head; pectoral rather long..... *subæqualis*
- cc. Maxillary reaching beyond middle of pupil, 2.33 in head; pectoral very short; D. XI-1, 23; A. II, 8 or 9; color silvery, with very distinct dark longitudinal stripes..... *sanctæ-luciæ*, 172
- aa. Body rather elongate and compressed, depth 3.5 in length; distance from insertion of ventrals to first anal spine one-half greater than depth of body; coloration dusky, with conspicuous dark streaks along rows of scales. *batabana*, 173

172. *Corvula sanctæ-luciæ* Jordan.

Head 3.1; depth 3.2; eye 4.5; snout 4.3; maxillary 2.4; mandible 2; interorbital 4.4; preorbital 2.4 in eye; D. XI-1, 23; A. II, 9; pectoral 1.9; ventral 1.8; caudal 1.8; scales 6-47-8. Body oblong, back compressed posteriorly and a little elevated; head rather small and blunt; snout as long or a little longer than eye; eye large, about equal to interorbital space; mouth somewhat oblique, jaws equal, maxillary reaching middle of pupil; a band of fine teeth in upper jaw, outer a little enlarged; teeth in lower jaw in one row with a few scattering ones; preopercle with a finely crenulate membranous edge; gillrakers 8 + 19, counting rudiments, longest one-half eye; scales large and strongly ctenoid, rows above lateral line anteriorly parallel with it; below last dorsal spines these rows turn upward and become horizontal again under anterior part of soft dorsal; dorsal spines slender, finely pointed but not sharp; soft dorsal and anal scaly at base; caudal truncate, scaled to near edge; second anal spine slender, shorter than first rays.

Color in spirits: Pale, darker above, with a bluish luster; each row of scales save the few lowermost with a grayish-brown stripe, these most continuous above, where they follow the upturned rows of scales; many dark punctulations everywhere; axil dark; fins not notably colored, but with many punctulations.

A species known at once by the horizontal stripes and their characteristic bend upward under the notch separating the dorsals. Hitherto known only from the type from Port Castries, St. Lucia. This description is from our largest specimen, 7.75 inches long, from Isabel Segunda; about 20 other individuals from Isabel Segunda, Ponce, Mayaguez, Hucare, and Boqueron, ranging in length from 2.5 to 6.5 inches; smaller specimens have the eye 3.5 in head.

Corvula sanctæ-luciæ Jordan, Proc. U. S. N. M. 1889, 649, Port Castries, St. Lucia; Jordan & Evermann, I. c., 1429, 1898.

173. *Corvula batabana* (Poey). "*Barriga Blanca*."

Head 3.3; depth 3.2; eye 4; snout 4.2; maxillary 2.25; mandible 2; interorbital 3.9; preorbital 7.8; D. XI-1, 28; A. II, 8; pectoral and ventral 1.7; caudal 1.6; scales 7-45-7. Body oblong, back compressed posteriorly, belly long, distance from ventrals to anal greater than depth of body; head mod-

erate, profile nearly straight, a slight concavity above eye; mouth not small, maxillary reaching middle of eye or beyond; teeth of upper jaw in an enlarged outer row, behind which is a band of very small teeth; lower jaw with a single series similar to outer row in upper jaw; snout with a few small pores, chin with 5 larger ones; edges of scales with fine striæ, rows below lateral line bending slightly upward opposite origin of anal, most apparent in smaller individuals; lateral line curved, concurrent with outline of back, becoming straight on caudal peduncle; dorsal spines weak; soft parts of vertical fins densely scaled, a sheath of larger scales at their bases; caudal rounded, or double truncate.

Color in spirits: Body nearly everywhere lustrous grayish-brown; about 8 longitudinal stripes of darker brown following rows of scales below lateral line; above lateral line are interrupted stripes and scattered spots of same color; fins all about color of body.

Known only from Cuba and Porto Rico. Four specimens, 8 to 10 inches in length, were obtained by us at Puerto Real and Arroyo.

Johnius batavanus Poey, Memorias, II, 184, 1860, Batabano, Cuba; Poey, Fauna Puerto-Riqueña, 327, 1881; Stahl, l. c., 77 and 163, 1883.

Corvula batavana, Jordan & Evermann, l. c., 1430, 1898.

Genus 98. BAIRDIELLA Gill. Mademoiselles.

This genus is characterized by the oblique mouth, little cavernous skull, few rows of small teeth, slender gillrakers, and preopercle armed with a plectroid spine. It is certainly a very natural group, and worthy of recognition as a distinct genus, although its relationships with *Ophioscion* and especially with *Stellifer* are very close. The numerous species are all American, all small in size and silvery in coloration, and some of them are remarkable for the great size of second anal spine. In others this spine is quite small. These variations among species unquestionably closely allied show how slight is the systematic value to be attached to the size of this spine.

a. Teeth of lower jaw unequal, chiefly biserial; inner teeth more or less enlarged; preorbital narrow.

b. Second anal spine moderate, 2.33 in head, not so long as soft rays, not reaching tip of last ray when depressed. Color, silvery, punctate; fins yellow; depth 3 in length. D. x-1, 22; A. II, 10. *chrysur*

bb Second anal spine very long, two-thirds length of head, reaching beyond tip of last ray; base of anal oblique, forming an angle with ventral outline.

c. Mouth not quite terminal; preorbital narrow; dorsal rays x-1, 23; dorsal spines stiff, lower, highest 2 in head; second anal spine 1.4; pectorals 1.6. Color, soiled silvery; depth 3.14. *ronchus*, 174

174. *Bairdiella ronchus* (Cuvier & Valenciennes). *Ronco*; *Ground Drummer*.

Head 3.1; depth 3.2; eye 4.6; snout 4.1; maxillary 2.5; mandible 2.1; interorbital 4.5; preorbital 9.3; D. x-1, 24; A. II, 8; pectoral 1.6; ventral 1.5; caudal 1.5; scales 7-50-8. Body oblong and compressed, anterior profile straight and rather steep; ventral outline straight from chin to origin of anal; base of anal very oblique, forming a strong angle in ventral outline; caudal peduncle long, compressed; head small, snout rather pointed, a little longer than eye; mouth nearly horizontal, maxillary reaching past middle of eye (in individual of 4.25 inches nearly to posterior border of orbit), lower jaw included; teeth in upper jaw villiform, in a very narrow band, outside of which is a row of much larger teeth; in lower jaw a band of villiform teeth with a somewhat enlarged inner row; preopercle with strong serrations, largest at angle, lowermost being the downward-pointing plectroid spine characteristic of *Bairdiella*; dorsal spines moderately strong, sharp, fourth highest, 2 in head; first anal spine reduced, second very long and strong, reaching tips of first rays and beyond all the other rays, 1.6 in head; ventrals nearly to vent; caudal truncate.

Color in spirits: Dirty-grayish, darker above, paler below; some dark punctulations on lower part of side scattered along from snout to base of lower lobe of caudal; faint dark streaks along longitudinal and oblique rows of scales; paired fins pale; edge of soft dorsal, spinous dorsal, caudal, and part of anal dusky, with dark dots.

Found on Atlantic coast of tropical America; generally common in the West Indies and on the coast of Brazil; recorded from Jamaica, Cuba, Maracaibo, Surinam, and obtained by us at San Juan and Mayaguez, Porto Rico. Though not exceeding 8 or 10 inches in length, this is a good food-fish.

Corvina ronchus Cuvier & Valenciennes, Hist. Nat. Poiss., V, 107, 1830, Maracaibo and Surinam.

Bairdiella ronchus, Jordan & Evermann, l. c., 1436, 1898; Poey, Fauna Puerto-Riqueña, 326, 1881; Stahl, l. c., 77 and 163, 1883.

Genus 99. OPHIOSCION Gill.

This genus is composed of small species, nearly all American, allied to *Sciæna* (*Sciæna umbra* L.), but differing in the armature of the preopercle, its bony margin being at all ages armed with strong persistent serræ, the lowermost teeth not directed forward. The caudal fin in this group is never lunate; the soft dorsal and anal are scaly; teeth in bands; gillrakers rather short.

There are 7 or 8 American species of this genus, all except *O. adustus* being known only from our Pacific coast.

175. *Ophioscion adustus* (Agassiz).

Head 3.25; depth 3.3; eye 4.2; snout 3.2; maxillary 2.8; mandible 2.5; interorbital 3.7; pre-orbital 5.4; D. x1-1, 22; A. II, 7; pectoral 1.5; ventral 1.5; caudal 1.4; scales 6-56-10. Body compressed, the dorsal outline of trunk trenchant, ventral outline of body nearly straight, the dorsal outline evenly arched; caudal peduncle long and thin; top of head not trenchant, evenly convex; snout rather long and blunt, scarcely projecting beyond the small inferior mouth; several pores and clefts in fleshy tip of snout; chin with five pores, the middle one, at symphysis, smallest; lower jaw well included; maxillary reaching anterior edge of pupil or middle of eye; teeth in both jaws, in villiform bands, the outer row in upper jaw enlarged; top of head, cheeks, and opercles scaled; snout and under parts of head naked; eye moderate, nearer tip of snout than gill-opening; preopercular serræ rather blunt, 9 or 10 in number; dorsal fin deeply notched, the spines weak, flexible, sharp, with filamentous appendages projecting beyond their ends, the first very short, the third longest, the rest graduated to the tenth, which is shortest, the eleventh being somewhat longer, the twelfth much longer, pertaining to the soft portion of the fin, which is scaled on membrane between rays; ventrals rather long and slender, outer rays slightly filamentous, reaching about to vent; anal small but high, first spine very short, second very long and strong, 1.6 in head, longest ray about equaling second spine; caudal large, rounded, finely scaled.

Color in spirits: Everywhere brownish with a bluish iridescence, save under parts, which are pale; the brown color on lower part of side is broken up into spots, which are darker at center; faint longitudinal brown lines following rows of scales, plainest below, scarcely evident in some specimens; anal and the paired fins grayish-black; dorsal and caudal like body; one specimen has scarcely any brown color, is grayish above and white below.

Five specimens, 3 to 3.5 inches in length, from Vieques and Arroyo. These differ in certain respects from Agassiz's colored figure of *Sciæna adusta*—chiefly in the larger eye, in the shorter straight portion of lateral line, and in the length of the second anal spine, which equals the longest rays, but is only half as long in the drawing. Nevertheless, we believe that our specimens are the young of Agassiz's species. The artist was manifestly somewhat inaccurate in his reproduction, as in the number of soft dorsal rays, and certainly in the omission of the short first anal spine. Moreover, the drawing was based upon a specimen much larger than ours (10 inches). We think these considerations are sufficient to explain the differences. The description of *O. adustus* in Jordan & Evermann's Fishes of North and Middle America hardly applies to the *Sciæna adusta* of Agassiz, but is based upon a specimen from Pernambuco, which probably belongs to some other species.

Sciæna (Corvina) adusta Agassiz, Pisc. Brasil., 126, pl. 70, 1829, Montevideo.

Genus 100. MICROPOGON Cuvier & Valenciennes. Croakers.

Body moderately elongate, compressed, somewhat elevated; preopercle strongly serrate; teeth in villiform bands, outer row in upper jaw enlarged; lower jaw with a row of minute barbels on each side; gillrakers short, thickish; spinous dorsal rather short, of 10 or 11 stoutish spines; second anal spine moderate; caudal fin double-truncate; lower pharyngeals narrow, distinct, with sharp conical teeth; air-bladder with long horns. A well-marked genus, the species all American, allied to *Ophioscion* and *Sciænops*, but distinguished by the presence of barbels; species all closely related, similar in form, size, and color, and all of value as food-fishes.

a. Dorsal rays x-1, 28 to 30.

b. Scales comparatively small, about 9 in a vertical series between front of dorsal and lateral line, 12 in an oblique series; outer teeth of upper jaw evidently enlarged; dark spots on scales above lateral line not forming continuous stripes; scales 54..... *undulatus*

- bb. Scales larger, 7 in a vertical series from front of dorsal to lateral line, 9 or 10 in an oblique series; teeth of outer series in upper jaw scarcely enlarged; dark spots above lateral line forming continuous streaks nearly as wide as interspaces; short vertical bars extending across lateral line; many oblique lines above these; markings more regular, though less sharply defined, than in *M. undulatus*.
- c. Second anal spine moderate, 5 in head; eye small, 6 in head; scales 54 *furnieri*, 176

176. *Micropogon furnieri* (Desmarest). *Verrugato*; *White-mouth Drummer*.

Head 3.25; depth 3.7; eye 5.35; snout 3.25; maxillary 3.2; mandible 2.7; interorbital 3.8; preorbital 4.6; D. x-1, 30; A. 11, 8; pectoral 1.4; ventral 1.9; caudal 1.6; scales 7-51-9. Body elongate, with a rather long and slender caudal peduncle; back a little elevated, profile from the eye to the dorsal nearly straight, head not large, snout rather long and blunt; mouth small, low, little oblique, lower jaw included; teeth small, in a band in each jaw; preopercle serrate with 2 enlarged spines at angle; soft dorsal and anal naked; soft dorsal low, with a 1-rowed basal sheath of scales; caudal double-truncate, its basal half scaly; base of pectoral with a few scales; dorsal spines very slender, sharp; anal very small.

Color in spirits: Silvery-white, upper half with many oblique bluish bars a little wider than interspaces, anteriorly extending across lateral line and becoming wider; a dark opercular spot.

This species is close to *M. undulatus*, and is known from Cuba to Surinam; abundant on coast of Cuba; recorded by Jordan & Rutter from Jamaica; probably not rare in Porto Rico. Our collection contains two fine specimens, 7.5 and 10.5 inches long, obtained at San Juan and Arecibo. A good food-fish, reaching the length of a foot or more.

Umbrina furnieri Desmarest, Première Décade Ichth., 22, pl. 2, fig. 3, 1822, Havana.

? *Micropogon argenteus* Cuvier & Valenciennes, Hist. Nat. Poiss., V, 218, 1830, Surinam.

? *Micropogon undulatus*, Poey, Fauna Puerto-Riqueña, 325, 1881; Stahl, l. c., 77 and 163, 1883.

Micropogon furnieri, Jordan & Evermann, l. c., 1462, 1898.

Genus 101. UMBRINA Cuvier.

Body moderately elongate; back somewhat arched. Head oblong, with snout thick and protuberant; mouth almost horizontal, of moderate size; preoperculum with its bony margin finely serrate; lower jaw with a single thickish barbel. Teeth in villiform bands, outermost in upper jaw somewhat enlarged. Anterior dorsal with about 10 spines; anal fin with 2 spines, second not very small. Caudal lunate or truncate. Gillrakers normal, but short. Air-bladder well developed.

This genus contains a considerable number of species, most of them being American. It agrees with *Sciæna* in nearly all respects, excepting the presence at the chin of a short, thick barbel. A similar barbel is found in the genus *Menticirrhus*, but notwithstanding the fact that all European writers have confounded *Menticirrhus* with *Umbrina*, the two genera are not really very closely related.

a. Dorsal rays only x-1, 25? no crossbands? *broussonetii*

aa. Dorsal rays x-1, 26 to 29; serræ of preopercle slender, not notably flattened.

b. Body with about 9 dark vertical crossbands, besides narrow undulating streaks along rows of scales; second anal spine 2.3 in head; pectoral 1.66 *coroides*, 177

177. *Umbrina coroides* Cuvier & Valenciennes.

Head 3.5; depth 3.5; eye 4.25; snout 3.4; maxillary 3; mandible 2.9; interorbital 4; preorbital 4.3; D. x-1, 26; A. 11, 6; pectoral 1.7; ventral 1.7; caudal 1.4; scales 5-48-10. Body rather elongate, back elevated, anterior profile nearly straight save for a slight convexity opposite eye; ventral outline nearly straight; caudal peduncle rather long and slender; head small, snout blunt and overhanging the small inferior mouth; teeth in villiform bands in each jaw; maxillary reaching beyond front of pupil; chin with a short barbel; snout with pores and clefts; preopercle with 5 teeth; fins small; caudal truncate, second anal spine much the longer and stronger.

Color in spirits: Light below, light-olivaceous above, a silvery luster everywhere; 9 darker cross-bars ending about level of pectoral; faint longitudinal stripes of dark following rows of scales; some dark on dorsal, fins otherwise plain.

This interesting species ranges from southern Florida to the West Indies and Brazil; recorded from Indian River, Florida (as *U. broussonetii*), and Jamaica; not uncommon in Porto Rico; our collection contains six specimens, 4 to 6 inches long, from Aguadilla, Arroyo, Hucares, and Vieques.

Umbrina coroides Cuvier & Valenciennes, Hist. Nat. Poiss., V, 187, 1830, Brazil; Jordan & Evermann, l. c., 1466, 1898.

Genus 102. MENTICIRRHUS Gill. The Whittings.

Body comparatively elongate, little compressed; head long, subconic, bluntish snout considerably projecting beyond mouth; mouth small, horizontal, both jaws with bands of villiform teeth, outer teeth in upper jaw more or less enlarged; chin with a single stoutish barbel; preopercle with its membranaceous edge serrulate; gillrakers short and tubercular or obsolete; dorsal spines high, slender, 10 or 11 in number (13 in *Cirrimens*); second dorsal long and low; caudal fin with lower angle rounded, upper sharp; anal fin with a single weak spine; no air-bladder. Lower pharyngeals separate, teeth varying from sharp to very obtuse.

This genus is one of the most strongly marked in the family. It has been confounded by all European writers with *Umbrina*, with which it has not very much in common except the presence of the barbel at the chin. All the species are American, and all bottom fishes. The low, elongate body, large pectoral, and obsolete air-bladder are all characters related to this peculiarity of habit. The species are all valued as food-fishes.

MENTICIRRHUS:

- a. Dorsal spines usually 11; head not terete, depressed, with low snout.
- b. Gillrakers obsolete, reduced to tubercular prominences, covered with teeth similar to those on the other gill-arches, more developed in the young; lower pharyngeals narrow; the teeth villiform or cardiform, all of them acute or conical, none with rounded heads (molar); teeth in the outer series of upper jaw more or less enlarged; scales on breast large.
- c. Soft dorsal longer, its rays 1, 23 to 1, 25.
- d. Outer teeth of upper jaw decidedly enlarged; dorsal spines not much elevated, the longest usually not reaching front of soft dorsal, 1.5 to 1.66 in head. Coloration, grayish-silvery, the dark markings not pronounced and often obsolete.
- e. Dorsal rays x-1, 22 to 24; snout rather shorter and less pointed than in *M. americanus*, 3.5 in head; mouth smaller, the maxillary 3 in head. Coloration usually plain, sometimes very dark, otherwise as in *Menticirrhus americanus*..... *martinicensis*, 178
- ee. Dorsal rays x-1, 24 or 25; snout longer, 3.33 in head; maxillary reaching nearly to middle of eye 2.8 to 3 in head; eye small, 2 in snout; teeth villiform, in broad bands, the outer series of the upper jaw very much enlarged, larger than in the other species; ventrals short, 1.5 in pectoral; pectoral 1.25 in head; caudal with the broad rounded lower lobe longer than the acute upper; scales all ctenoid, those of the breast larger and regularly placed. Color, grayish-silvery, with obscure darker clouds along the back and sides, these marks forming dusky bars, running obliquely forward and downward to considerably below the lateral line, these often obsolete; the bar at the nape saddle-like; lining of gill-cavity dusky; pectoral yellowish, dusky at tip; an obscure dusky streak along the lower part of side running into lower lobe of caudal..... *americanus*
- dd. Outer teeth of upper jaw less enlarged; spinous dorsal elevated, the longest spine reaching past front of soft dorsal, its length 1.5 in head; coloration strongly marked, body scarcely silvery; eye small, 2.33 in snout, 2 in interorbital area, about 7 in head; snout long, bluntish, 3.3 in head; mouth large; maxillary reaching middle of eye, 2.8 in head; pectoral 1.14 in head. Color dusky-gray above, sometimes blackish, the back and sides with distinct dark oblique crossbands running downward and forward, the anterior one at the nape extending downward, meeting the second and thus forming a V-shaped blotch on each side; a dark lateral streak bounding the pale color of the belly, most distinct posteriorly and extending on lower lobe of caudal; inside of gill-cavity scarcely dusky; pectoral dark..... *saxatilis*

UMBRULA:

- bb. Gillrakers present, very short and rather slender; lower pharyngeals rather broad; some or most of teeth molar, that is, enlarged, with thickened rounded heads, the molar teeth covering at least anterior portion of bone; teeth in outer series of upper jaw scarcely larger than the others; scales on breast small..... *littoralis*

178. *Menticirrhus martinicensis* (Cuvier & Valenciennes). *Jewsharp Drummer*.

Head 3.3; depth 4; eye 7; snout 3.7; maxillary 3; mandible 2.8; interorbital 4.2; preorbital 5.3; D. x-1, 24; A. 1, 7; pectoral 1.4; ventral 2.2; caudal 1.6; scales 6-54-12. Body quite elongate, back considerably elevated, ventral outline nearly straight; head small, conical, upper profile gently convex; mouth not large, inferior, maxillary reaching past middle of eye; outer teeth of upper jaw considerably enlarged; snout projecting beyond mouth; spinous dorsal small, spines slender and pointed, but not sharp; soft dorsal very low; anal very small; pectoral large; caudal with an emargination in upper lobe, lower lobe the longer; lateral line almost straight except at ends, parallel with outline of back.

Coloration usually plain, sometimes very dark; back and sides usually with oblique dusky bars. Our specimens are pale below, darker above, and show few definite markings; there are faint oblique dark lines following rows of scales; tip of pectoral and ventrals and edge of anal nearly black.

Known from the West Indies to Patagonia; very common on the Brazilian coast, where it replaces the closely related *M. americanus*, from which it is not well separated; recorded from Jamaica, Martinique, and Rio Janeiro, and obtained by us in the San Juan market and at Palo Seco, Porto Rico, where it is probably not rare. It is a fair food-fish.

Umbrina martinicensis Cuvier & Valenciennes, Hist. Nat. Poiss., V, 186, 1830, Martinique.

Umbrina gracilis Cuvier & Valenciennes, Hist. Nat. Poiss., V, 189, 1830, Brazil.

Umbrina arenata Cuvier & Valenciennes, Hist. Nat. Poiss., V, 190, 1830, Brazil.

Umbrina januarua Steindachner, Ichth. Beitr., V, 122, 1876, Rio Janeiro.

Menticirrhus martinicensis, Jordan & Evermann, l. c., 1473, 1898.

Genus 103. *EQUES* Bloch. Ribbon-fishes.

Body oblong, compressed, back much elevated anteriorly, rapidly tapering to narrow caudal peduncle; mouth small, lower jaw included; preorbital wide; snout with slits and pores well developed; teeth all villiform, in broad bands, outer scarcely enlarged; preopercle with a fringed border and no bony serræ; scales small, irregular, with smaller ones intermixed, extending on soft fins; gillrakers few, short, and slender; dorsal fin very long, of 9 to 15 close-set spines and 36 to 55 soft rays; anterior interneurals closely wedged in behind occiput; anal small, its spine small; caudal rhombic; pyloric cæca few; vertebrae 10+15=25.

This genus is one of the most remarkable in the family in respect to form as well as to the coloration of its species.

- a. Dorsal rays x to XII-1, 36 to 46, first 4 to 6 of the interneurals wedged in between neurals of second and third vertebrae.
- b. Profile steep, but not vertical; distance from snout to first dorsal spine about equal to depth of body.
- c. Dorsal spines little elevated, not nearly as long as head; back arched; dorsal with 38 to 41 soft rays; color variously dusky or gray, with at least traces of about 7 lengthwise streaks; depth 2.6 to 2.75 in length *acuminatus*, 179
- cc. Dorsal spines elevated, longest 2.75 in length of body; soft parts of vertical fins with white spots; body robust, back much compressed, general form much as in *Eques acuminatus*, but caudal peduncle deeper and more compressed; pectoral and ventral short and equal, 1.14 in head. Color, dark-brown, a light bar in front of eye extending around chin, a second pale bar extending around head immediately behind eyes, a third extending from in front of dorsal over base of pectoral; a light bar along base of soft dorsal; a light bar extending from behind elevated portion of spinous dorsal downward, dividing into two, the branches running straight back, upper branch to beginning of last fourth of soft dorsal, lower branch to base of caudal; 2 or 3 light, undulating longitudinal bars below these; fins all dark brown, vertical fins with many whitish stellate spots. Head 3.75 in length; depth 3. D. XI or XII-1, 46. *punctatus*
- bb. Profile very steep. Body deepest below first dorsal spine, thence rapidly tapering to narrow caudal peduncle. Color, olivaceous, 3 dark-brown longitudinal bands along side, middle one from eye backward reaching tips of middle caudal rays. D. X-1, 37. *pulcher*
- aa. Dorsal rays XIV or XV-1, 53; about 9 interneurals wedged in between neurals of second and third vertebrae; profile almost vertical; body highly variegated, with ribbon-like oblique bands. *lancoletus*

179. *Eques acuminatus* (Bloch & Schneider). "*Berdugo*" or "*Bergudo*."

Head 3.1; depth 2.8; eye 3.8; snout 4.1; maxillary 2.6; mandible 2.4; interorbital 4.2; preorbital 5.2; D. X-1, 39; A. II, 7; pectoral 1.5; ventral 1.5; scales about 50. Body elongate, compressed posteriorly, back much elevated, ventral outline nearly straight; anterior profile straight from front of dorsal to upper part of blunt snout; mouth moderate, low, maxillary reaching middle of eye, lower jaw included; teeth in bands, outer row in upper jaw enlarged; soft fins all more or less scaly; gillrakers short, not pointed, 6+8.

Color in spirits: Everywhere dark brown, with pale longitudinal stripes, as follows: two from region of nape nearly confluent at middle of soft dorsal; one from a little above eye to last rays of soft dorsal; one from upper edge of eye to upper edge of caudal peduncle; two from near base of pectoral to base of caudal, the upper of these extending faintly across head to eye.

This species ranges from South Carolina to Brazil; it has been recorded from Key West, Tortugas, and Jamaica, and was obtained by us at Arroyo, Porto Rico, from which place we have two specimens, each 6 inches long. Valued as food.

Grammistes acuminatus Bloch & Schneider, Syst. Ichth., 184, 1801, no locality; after Seba.

Eques lancatus Cuvier & Valenciennes, Hist. Nat. Poiss., V, 169, 1830, Brazil.

Eques acuminatus, Jordan & Evermann, l. c., 1487, 1898.

Family L. POMACENTRIDÆ. The Demoiselles.

Body short, deep, compressed, covered with ctenoid scales of varying size; lateral line wanting posteriorly; mouth small, usually with rather strong teeth, either conic or incisor-like; vomer and palatines toothless; nostril single on each side, nearly round; preopercle with its posterior edge largely free, serrate or entire; preorbital sheathing the small maxillary; dorsal fin single, with numerous strong spines, spinous portion longer than soft, which is similar to soft anal, both fins scaly at base; anal spines 2; ventral fins thoracic, 1, 5, anterior rays longest, usually filamentous; a scaly appendage at base of ventral. Lower pharyngeals fully united; branchiostegals 5 to 7; gills 3.5, slit behind last gill very small or obsolete; gillrakers rather long and slender; no labyrinthiform appendage; air-bladder and pseudobranchiæ present; pyloric cæca 2 or 3; gill-membranes free from isthmus. Vertebrae 12+14=26.

Fishes of tropical seas, similar in mode of life to the *Chatodontidæ*, feeding on small marine animals and plants in the coral reefs. Genera 15; species about 180, most of them too small to be used as food. They are very active in life and the coloration is usually brilliant, sometimes changing much with age. The family shows strong affinities with the *Labridæ* in its gill-structures and pharyngeals. In other respects it approaches the *Kyphosidæ*, while the unique character of the simple nostril is shared with the *Cichlidæ* only, from ancestors of which group the *Pomacentridæ* are probably descended.

I. Scales large, 25 to 40 in lengthwise series.

POMACENTRINÆ:

- a. Teeth fixed, conical or incisor-like, covering nearly whole free edge of each jaw; carnivorous species.
- b. Teeth conical, in two to four series, outer enlarged and bluntish; preopercle entire; scales large; body oblong, depth 2.5 to 1.33 in length; lateral line wanting on tail..... CHROMIS
- bb. Teeth more or less flattened or incisor-like, in one or two series.
- c. Preopercle, and usually preorbital also, sharply serrate.
- d. Teeth entire, strictly uniserial in each jaw; preorbital not very deep, its edge not notched; snout scaly; lower jaw naked..... EUPOMACENTRUS, 104
- cc. Preopercle and preorbital strictly entire; snout naked.
- e. Teeth emarginate or Y-shaped..... ABUDEFDUF, 105

MICROSPATHODONTINÆ:

- aa. Teeth movable, incisor-like, in one row on front of each jaw; lower jaw weak, with teeth along its front only; preopercle and preorbital entire; snout scaled almost to lips; preorbital notched behind nostril; lower limb of preopercle scaled; soft dorsal and anal elevated; caudal deeply forked; soft anal rather long, of 14 or 15 rays; herbivorous species..... MICROSPATHODON

Genus 104. EUPOMACENTRUS Bleeker. Pescados Azules.

Body ovate, deep, and compressed, profile steep, usually rounded. Head moderate, nearly as deep as long, snout scaly, lower jaw naked. Mouth quite small, terminal, jaws equal; each jaw armed with a single close-set series of compressed, immovable teeth, which are truncate at tip. Gillrakers long; preopercle more or less serrate; preorbital serrate. Scales large, strongly ctenoid, lateral line running parallel with back to near end of dorsal fin, at which point it ceases. Dorsal fin continuous, with 12 or 13 low stout spines; membrane of spinous dorsal usually not deeply incised nor lobed; soft part more or less elevated, its last rays gradually shortened; lower limb of preopercle usually more or less scaly; preorbital narrow, without deep notch; anal fin similar to soft dorsal, with 2 spines, of which the second is much the larger; soft rays 12 to 16; dorsal spines with a sheath of large scales, membranes of both dorsal and anal covered high up with small scales; caudal fin more or less forked, lobes rounded; lower pharyngeals triangular; branchiostegals 5 or 6.

Species numerous in the tropical seas, chiefly American, and extremely variable in form and color, the brilliant coloration apparently dependent on surroundings. The species are little known and the classification of those found in the West Indies is not wholly satisfactory.

- a. Upper anterior profile of head arched.
- b. Depth of body moderate, 1.75 to 2.25 in length, without caudal.
- c. Lower posterior half of body dark, like anterior half; caudal fin mostly dusky.
- d. Pectoral fin not edged with white.
- e. Depth of body 1.75 in length of body (without caudal); side with faint cross-streaks..... *adustus*
- ee. Depth of body 2 to 2.25 in length.
- f. Opercle without distinct dark spot; caudal not tipped with orange.
- g. Anal without distinct blue spot in its posterior axil, except in young.
- h. Head with few if any small accessory scales.
- i. Base of pectoral without black spot or with but one..... *fuscus*, 180
- Base of pectoral with two black spots..... *dianæus*

- gg. Anal with a bluish spot at base of last ray; head and fins much spotted with blue *analis*, 181
 ff. Opercle with a distinct dark spot above; pectoral with a dark spot; tips of all fins orange *otophorus*
 cc. Lower posterior half of body unlike anterior part, being more or less abruptly bright-yellow; caudal fin bright-yellow; usually a blue spot at base of last ray of anal.
 j. Region below lateral line with many blue spots..... *leucostictus*, 182
 jj. Region below lateral line with few blue spots or none.
 k. Back with two round dusky spots on each side..... *flaviventer*
 bb. Depth of body about 2.75 in length; posterior half of body yellowish; fins with more or less yellowish..... *partitus*
 aa. Upper anterior profile of head straight, not arched; body and fins mostly dusky, with pale spots..... *planifrons*

180. *Eupomacentrus fuscus* (Cuvier & Valenciennes). *Brown Cockeye Pilot; Maria Mollé*

(PLATE 27.)

Head 3.5; depth 2.5; eye 3.4; snout 3.5; maxillary 3; D. XII, 15; A. II, 13; scales 2-28-9. Body compressed, anterior profile steep and evenly convex; dorsal and ventral outlines similar; mouth moderate, little oblique; maxillary reaching anterior border of orbit; jaws equal; teeth in a single series in each jaw, incisor-like, and closely placed; preopercle and preorbital serrate; gillrakers small, weak; scales large, ctenoid, reduced in size on head; snout, opercles, and cheek scaled; lateral line concurrent with back to below middle of soft dorsal, where it ceases. Fins rather large; longest dorsal spine about 2 in head, dorsal rays elevated, longest 1.2 in head; second anal spine 2 in head; soft anal similar to soft dorsal; caudal forked, lobes equal, about 1 in head; pectoral rather long, reaching vent; ventrals long and pointed, passing vent.

Color, dark-blue above, lighter on the sides and under parts; side of head and body with some yellowish, under parts with purple tinge; in some specimens the general color is dark-brown or nearly black; edges of scales dark, forming narrow dark vertical bands, but in most specimens these are entirely absent; dorsal and anal fins dark-bluish or olivaceous at base, the margin brighter blue; caudal dusky or pale, usually with some yellow at base; pectoral and ventrals greenish-olivaceous, no black spot at base of pectoral, no dorsal ocellus and no white spot at base of last anal ray; no bluish or whitish spots on head or body.

This species may be distinct from *E. analis*, or the latter may be simply an individual color variation from the typical form. The specimens which we refer to *E. fuscus* were obtained by us at San Juan, Aguadilla, and Ponce, and by Mr. Gray at San Geronimo.

Found in the West Indies south to Brazil and north to Key West. The least common of the three species of *Eupomacentrus* which we recognize from Porto Rico.

Pomacentrus fuscus Cuvier & Valenciennes, Hist. Nat. Poiss., V, 432, 1830, Brazil.

Pomacentrus variabilis Castelnau, Anim. Nouv. ou Rares, Poiss., 9, pl. 3, fig. 3, 1855, Bahia.

Pomacentrus atrocyaneus Poey, Memorias, II, 190, 1860, Havana.

Eupomacentrus fuscus, Jordan & Evermann, l. c., 1552, 1898.

181. *Eupomacentrus analis* (Poey).

Head 3.2 to 3.5; depth 2 to 2.4; snout 3.2 to 3.5; eye 3.2 to 3.5; maxillary 3.2 to 3.75; mandible 3.75 to 4.25; interorbital 3.25 to 3.66; preorbital 7; scales 3-28-8; D. XI, 15; A. II, 13. Body rather deep, strongly compressed, anterior profile steep and evenly convex; interorbital area strongly convex; eye usually a little greater than snout; mouth small, slightly oblique, jaws equal; teeth in single series, incisor-like and close-set; maxillary scarcely reaching vertical of anterior part of orbit; gillrakers slender, weak, and short, length less than diameter of pupil; preopercle strongly serrate; preorbital also serrate, but less distinctly so; scales large, firm, and strongly ctenoid; lateral line beginning at the upper edge of opercular opening running parallel with dorsal outline to beneath middle of base of soft dorsal, where it abruptly terminates; top of head and snout densely scaled, scales smaller than on side; opercle and preopercle scaled; chin naked; bases of dorsal and anal fins densely scaled; membranes of caudal fin covered with very small scales; dorsal spines strong and sharp, length of longest 1.8 to 2 in head; soft dorsal with its middle rays elevated, their length 1.2 to 1.4 in head; first anal spine very short, second much longer and stronger, its length 2 in head; soft anal similar to soft dorsal, its longest rays about 1.25 in head; pectoral broad, short, scarcely reaching vent, length about 1.2 in head; ventral longer, tips of outer rays passing vent, about equal to length of head; caudal forked, upper lobe the longer, usually somewhat longer than head.

Color in life very variable; back and upper part of side usually dark-blue or bluish, becoming somewhat paler on lower part of side and belly; head dark-blue; sometimes body is a dark brown, with scarcely any trace of blue; in other specimens the color is a sooty blue-black in life, under parts

scarcely paler; scales on anterior part of side sometimes with a long conspicuous bronze-olive spot; head, side, and bases of vertical fins usually with small sky-blue spots, these spots frequently in life inclining to black, in alcohol becoming at first more intense blue and later fading to white; the abundance of these spots varies greatly; in larger individuals, from 3 to 4 inches in length, the spots are greatly reduced in number, sometimes almost wholly absent; more abundant in individuals 2.5 inches or under in length, sometimes profusely so. Spots on anterior part of head sometimes take the shape of narrow lines or oblong blotches; a large blue blotch at bases of upper pectoral rays, becoming black in alcohol; a large black blotch on middle of anterior dorsal rays, sometimes quite round and surrounded by a narrow blue border; a small blue spot, fading to white in alcohol, near base of last anal rays.

Seventeen specimens, varying from 3 to 3.5 inches in length, collected at Ponce, February 1, were quite uniform in coloration; they each show a few white spots, some very faintly, on side and occasionally at base of anal; none noticeable on head or fins; a white spot present in every case on last anal ray at base; a single black spot at base of upper pectoral rays; dorsal ocellus showing distinctly on only three or four specimens; caudal usually pale, yellowish on two specimens. Seven specimens, 3 to 4 inches in length, collected at Culebra Island, February 11, were each profusely covered with pale-blue or whitish spots, most abundant on head, dorsal, and base of caudal and anal; spot at base of last anal ray always present; black spot at base of pectoral sometimes obscure; dorsal ocellus usually present; caudal pale or yellowish on three or four of the specimens. Nineteen specimens, somewhat smaller, collected at Fajardo, February 17, closely resemble the Culebra specimens; some of them, however, resemble more closely specimens from Ponce in the less numerous spots; the bluish-white spot at base of last anal ray is always present, and most of the specimens show the dorsal ocellus. Twenty other specimens from Fajardo, collected on same date and of smaller size, are much more profusely spotted; the dorsal ocellus is more distinct and the caudal fin more yellowish; the lower part of sides and the belly are paler. These specimens show a near approach to *E. leucostictus*. Four specimens from Mayaguez, each about 2.5 inches long, are profusely spotted like those from Culebra and Fajardo, but the general color is much darker; white spot on anal and black one on base of pectoral present; the dorsal ocellus is fairly distinct. Eleven specimens from Puerto Real, each 3.5 inches long, resemble closely those from Ponce in having but few spots; white spot on anal and black blotch at base of pectoral both present; dorsal ocellus evident in only six of eleven specimens; caudal peduncle and fin usually yellowish. Three small specimens from Hucaras are profusely covered with small blue specks and agree with those from Culebra.

In our collections from Porto Rico we have what we provisionally identify as three species of *Eupomacentrus*, namely, *E. fuscus*, *E. analis*, and *E. leucostictus*; but we are not at all sure that all three of these should not be united under one species—*E. fuscus*. The color variations and intergradations are most perplexing. In the field, fresh specimens sometimes show a most diverse coloration, but when brought into the laboratory these differences, to a very large extent, disappear. In life the caudal peduncle, caudal fin, and under parts are inclined to be more decidedly yellow than in alcoholic specimens, but this is not always true. In some instances, as, for example, the majority of specimens obtained at Ponce and Fajardo, the under parts are scarcely less dark than the upper. As a general rule the younger individuals are more profusely covered with brilliant sky-blue spots than are the older ones. As already stated, these spots appear in life as black or very dark-blue, but in alcohol they invariably fade first to sky-blue and finally to white or bluish-white; the spot near the base of last anal ray is usually bluish-white in life, fading to white in spirits.

The blotch on base of pectoral is sometimes bluish, but usually black. Some of the spots on anterior part of head are oblong or linear; the blotch on anterior soft rays of soft dorsal is usually black and irregular in form, though frequently it is beautifully ocellated, the border being a narrow line of bright sky-blue; this ocellus is always present in young examples, but frequently disappears in those of larger size. Except in a very few specimens, which we refer to *E. fuscus*, the spot near the base of the last anal ray is always present; except on certain specimens, which we refer to *E. leucostictus*, and whose colors agree closely with plate 28, there is no sharp contrast between the color of the upper and the lower parts, but where there is any difference at all it is through the gradual change from the dark of the back to the lighter of the ventral parts; except for the constant presence of the spot upon the anal fin, the specimens which we refer to *E. analis* would agree equally well with the prescriptions of *E. fuscus*; but as we have found no specimens without the anal spot, it seems best at present to regard the two forms as distinct. To ascertain the true relations of these species careful field studies in many different localities in the West Indies are necessary.

This species attains a length of 3 to 4 inches. Its proportional measurements are quite constant, as may be seen from the appended table:

Comparative measurements of 33 specimens of *Eupomacentrus analis*.

Locality.	No.	Length.	Head.	Depth.	Eye.	Snout.	Maxillary.	Mandible.	Interorbital.	Preorbital.	Scales.	Dorsal fin.			Anal fin.			Pectoral.	Ventral.	Caudal lobes.
												Formula.	Longest spine.	Longest ray.	Formula.	Longest spine.	Longest ray.			
Ponce	29	3.18	3.2	2.13	3.25	3.5	3.5	4.25	3.25	6	2-29-9	XII, 15	1.8	1.4	II, 13	2	1.33	1.33	1	
Do.	688	3.5	3.5	2.13	3.2	3.25	3.25	3.75	3.25	7	2-29-9	XII, 15	1.66	1.17	II, 13	2	1.2	1.2	1	.87
Do.	689	3.5	3.5	2.4	3.2	3.25	3.25	3.75	3.66	7	3-29-9	XII, 15 ¹⁾	1.25	1.25	II, 13	2	1.2	1.2	1.1	.87
Do.	690	3.25	3.4	2	3.2	3.25	3.25	3.25	3.5	7	3-29-9	XII, 15	1.8	1.25	II, 13	2	1.25	1.2	1	.87
Do.	691	3.36	3.5	2.13	3.2	3.5	3.5	4.25	3.75	7	3-29-9	XII, 15	1.8	1.4	II, 13	2	1.4	1.25	1.1	1
Do.	692	3.25	3.4	2.25	3.2	3.5	3.5	4	3.25	7	3-29-9	XII, 15	1.8	1.25	II, 13	2	1.5	1.2	1	.87
Do.	693	3.25	3.4	2.1	3.2	3.25	3.5	4	3.75	7	3-29-9	XII, 15	2	1.25	II, 13	2	1.4	1.2	1.1	1
Do.	694	3	3.25	2	3.2	3.5	3.5	4	3.5	7	3-29-9	XII, 15	2	1.25	II, 13	2	1.4	1.2	1	.87
Do.	695	3	3.4	2.4	3.2	3.4	3.4	4	3.5	7	3-29-9	XII, 15	2	1.25	II, 13	2	1.2	1.2	1	.87
Do.	696	3	3.4	2.33	3.2	3.4	3.5	4	3.25	7	2-28-9	XI, 15	2	1.2	II, 13	2	1.25	1.2	1	.87
Do.	697	3.5	3.5	2.25	3.5	3.5	3.5	4	3.5	7	3-28-9	XII, 15	2	1.2	II, 13	2	1.2	1.2	1	.87
Do.	698	3.25	3.25	2.25	3.25	3.5	3.75	4	3.5	7	3-28-9	XII, 15	2	1.2	II, 13	2	1.2	1.2	1	.87
Culebra	699	4	3.25	2.25	3.75	3.5	3.5	4	3	7	2-26-9	XII, 15	2	1.1	II, 13	1.9	1.2	1.1	1	.87
Do.	700	3	3.4	2.4	3	3.1	3.1	4	3	7	3-28-9	XII, 15	2	1	II, 13	2	1.2	1.17	1	.87
Do.	701	3.25	3.5	2.5	3.5	3.2	3.5	4	3	7	3-28-9	XII, 15	1.75	1.1	II, 13	2	1.1	1.1	1	.87
Do.	702	3	3.1	2.25	3.5	3.5	3.25	4	3.5	7	3-28-9	XII, 15	1.75	1.1	II, 13	2	1.2	1.17	1	.87
Do.	703	3	3.33	2.25	3	3.5	3.25	4	3	7	3-28-9	XII, 15	2	1.2	II, 13	2	1.4	1.25	1	.87
Do.	704	3	3.4	2.4	3.5	3.5	3.25	4	3.5	7	3-28-9	XII, 14	1.4	1.1	II, 13	1.75	1	1.25	1	.75
Do.	705	2.5	3	2.4	3.5	3.5	3.1	4	3.25	7	3-28-9	XII, 14	1.75	1	II, 13	2	1.2	1.25	1	.87
Fajardo	706	3.37	3.2	2.4	3.1	3.5	3.2	4	3.5	7	2-27-9	XII, 15	1.5	1	II, 13	1.4	1.2	1.2	1	.87
Do.	707	3.25	3.4	2.5	3.1	3.5	3	4	3.5	7	3-28-9	XII, 15	1.75	1.2	II, 13	1.75	1.2	1.2	1	.87
Do.	708	3.25	3.5	2.3	3.5	3.5	3.5	4	3.2	7	3-29-9	XII, 15	1.6	1	II, 13	1.75	1.2	1.2	1	.87
Do.	709	3.5	3.4	2.5	3.5	3.2	3.5	4	3.5	7	3-28-9	XII, 15	1.6	1	II, 13	1.75	1	1.2	1	.87
Do.	710	3	3.4	2.4	3.5	3.5	3	4	3.4	7	3-28-9	XII, 15	1.6	1	II, 13	1.75	1.2	1.3	1	.87
Puerto Real	748	3.5	3.4	2.1	3.4	3.7	3.5	4	3.4	7	3-28-9	XII, 15	1.75	1.1	II, 13	1.75	1.1	1.25	1	.87
Do.	749	3.5	3.4	2.4	3.5	3.7	3.6	4	3.5	7	3-29-9	XII, 15	1.75	1.1	II, 13	1.5	1.1	1.25	1	.87
Do.	750	3.5	3.2	2.4	3.5	3.5	3.6	4	3.5	7	3-29-9	XII, 15	1.75	1.1	II, 13	1.75	1.2	1.25	1	.87
Do.	751	3	3.1	2.2	3.3	3.5	3.5	4	3.5	7	3-28-9	XII, 15	1.75	1.25	II, 13	1.75	1.4	1.25	1	1
Do.	752	3.5	3.5	2.5	3.6	3.5	3.4	4	3.5	7	3-28-9	XII, 15	1.75	1	II, 13	1.75	1.2	1.25	1	.87
Do.	753	3.37	3.4	2.1	3.5	3.5	3.4	4	3.3	7	3-29-9	XII, 15	1.6	1.1	II, 13	2	1.2	1.25	1	.87
Do.	754	3.37	3.4	2.4	3.5	3.5	3.4	4	3.3	7	3-29-9	XII, 15	1.6	1.25	II, 13	1.75	1.2	1.25	1	1
Do.	755	3.5	3.2	2.2	3.5	3.5	3.5	4	3.3	7	3-29-9	XII, 15	1.75	1	II, 13	1.75	1.4	1.25	1	1
Do.	756	3.25	3.5	2	3.5	3.7	3.5	4	3.5	7	3-29-9	XII, 15	1.75	1.25	II, 13	2	1.2	1.25	1.4	1

¹ Broken.

West Indies north to Key West; hitherto regarded as only from Havana and Key West. Ninety per cent of the specimens of this genus obtained by us are referred to this species. Specimens were obtained at San Juan, Aguadilla, Mayaguez, Puerto Real, Guanica, Ponce, Hucares, Fajardo, and Culebra; it was exceedingly abundant at Culebra, Fajardo, and Ponce about the coral reefs and around the little islets, every haul of the seine securing several. At Fajardo they were particularly common among the algae on the west side of the small keys off the Playa.

Pomacentrus analis Poey, Synopsis, 327, 1867, Havana.

Eupomacentrus analis, Jordan & Evermann, l. c., 1554, 1898.

182. *Eupomacentrus leucostictus* (Müller & Troschel). *Beau Gregory*; *Cockeye Pilot*; *Black Pilot*.

(PLATE 28.)

Head 3.5; depth 2; eye 3; snout 3.75; maxillary 3.75; interorbital 3.75; preorbital 6.5; D. XII, 15; A. II, 13; scales 2-29-10. Body compressed; dorsal and ventral outlines similar; profile convex, but less steep than in *E. fuscus* or *E. analis*. Mouth moderate, maxillary reaching anterior part of orbit; jaws equal; teeth in a single band in each jaw, incisor-like and close-set. Preopercle and preorbital distinctly serrate; scales large, firm, strongly ctenoid, reduced in size on head; snout and sides of head completely scaled; lateral line beginning at upper end of gill-opening, arched, gradually approaching dorsal outline as it proceeds backward, ceasing under middle dorsal rays; fins moderate; longest dorsal spines nearly 2 in head; middle rays of soft dorsal elevated, about 1.4 in head; second anal spine long, strong, and gently curved, 2 in head; soft anal similar to soft dorsal, the rays somewhat longer; caudal forked, lobes about equal, upper sometimes slightly the longer; pectoral short, scarcely reaching vent; ventral short, reaching scarcely past vent.

Color: Top of head and back rich dark-blue, sides and under parts abruptly lemon-yellow, this yellow extending over side of head, entire side of body below lateral line and to some extent above it, and entire caudal peduncle and caudal fin; side of head and anterior part of body as well as top of head and dorsal region with numerous small pale-blue spots; similar spots on spinous dorsal and anterior soft dorsal rays, as well as a few on anal fin; a large black blotch on middle anterior dorsal rays, not ocellated. Posterior half of dorsal and anal yellow like caudal; pectoral pale-yellow; pale-blue spot near base of last anal ray; a small black blotch at base of pectoral. In life the pale-blue spots on top of head and upper part of side appear white by contrast with the darker-blue ground-color, those on side of body and head appearing bluer. Along the lateral line are sometimes present a few small, oblong dark spots. The coloration of this species is extremely variable; usually the colors are about as given above, but some specimens showed a gradual encroachment of the blue of the back upon the yellow of the sides, thereby approaching the general color-pattern of what we have called *E. analis*.

The extreme forms of this species and of *E. analis* are remarkably different and one would not hesitate to regard them as entirely different species, but individuals taken from different places and of various sizes, show very perplexing variations. It has, however, been impossible to show that these variations present actual intergradation, and for that reason we prefer, for the present, to recognize the species as distinct.

Found in the West Indies north to southern Florida. In Porto Rico we obtained specimens at San Juan, Puerto Real, Culebra, and Guanica. While fairly common, it is apparently much less abundant than *E. analis*.

Pomacentrus leucostictus Müller & Troschel, in Schomburgk's Exc. Barbados, 674, 1848, Barbados.

Pomacentrus caudalis Poey, Synopsis, 328, 1867, Havana.

Pomacentrus zanthurus Poey, Memorias, II, 190, 1860, Havana.

Pomacentrus dorsopunicans Poey, Synopsis, 328, 1867, Havana.

Eupomacentrus leucostictus, Jordan & Evermann, l. c., 155, 1898.

Genus 105. ABUDEFDUF Forskål. Pintados.

Body deep, compressed, covered with large ctenoid scales; snout without scales; preopercle and preorbital entire, lower limb of preopercle scaleless; 3 to 4 rows of scales between lateral line and dorsal; teeth compressed, fixed, more or less distinctly emarginate, in one series in each jaw, those below occupying most of free edge of jaw; jaws subequal. Dorsal usually with 13 spines, the last slightly shorter than median ones; branchiostegals 5 or 6; pyloric caeca 3. Lower pharyngeals triangular.

Species numerous, often brightly colored, about coral reefs in the tropical seas. We exclude from this genus all the species formerly referred to *Glyphidodon*, in which the teeth are in more than one series, and also those in which the snout is scaly or the lower pharyngeals not triangular. For these forms different generic names, *Hemiglyphidodon*, *Amblyglyphidodon*, *Glyphidodontops*, etc., have been defined by Bleeker. The genus *Stegastes* Jenyns (*imbricatus*) is also very close to *Abudefduf*, but it seems to have entire teeth, and the snout and fins are densely scaly. The genus *Nexilarius* is less closely related to *Abudefduf*.

GLYPHISODON:

a. Preopercle entire.

b. Preorbital very narrow, its least breadth less than pupil, even in adult; anterior profile of head nearly straight, snout rather acute; dorsal spines 13; anal with about 12 soft rays.

c. Scales about 4-30-11; green, with about 6 dark-blue or blackish crossbands; depth 1.75 to 2 in length; anal rays II, 12..... *saxatilis*, 183

EUSCHISTODUS:

bb. Preorbital broad, its least breadth not less than diameter of pupil, greater than pupil in adult; anterior profile of head more or less arched, snout low and blunt, projecting beyond the small mouth; anal rays II, 10; coloration dull.

d. Dorsal spines 13; scales 27 or 28.

e. Color in adult brownish, with green dots, not distinctly banded; teeth smaller, $\frac{2}{5}$ on each side..... *analogus*

dd. Dorsal spines 12; body with 5 dark crossbands, fainter than in *A. saxatilis*; scales about 25..... *taurus*

aa. Preopercle said to be coarsely serrate; dorsal spines 13; scales very large, 25; body with dark crossbands..... *rudis*

183. *Abudefduf saxatilis* (Linnaeus). *Cockeye Pilot*; *Demoiselle*; "*Chirivita*"; "*Pintado*."

Head 3.2; depth 1.8; eye 3.2; snout 3.5; interorbital 2.7; D. XIII, 13; A. II, 12; pectoral 0.9; ventral 1.0; caudal 0.8; scales 5-28-11.

Color in spirits: Chiefly silvery, somewhat darker above, top of head brown; side with five vertical

brown bars, largest as wide as eye, first from front of dorsal to base of pectoral, second downward from fourth, fifth, and sixth spines, third from ninth and tenth spines toward vent, fourth from last spines to middle of anal, fifth from soft dorsal across peduncle upon anal; sometimes a faint sixth bar upon base of caudal; first four not reaching belly; axil dark, front of base of pectoral black. Smaller individuals (2 inches and less) have, in spirits, ground-color nearly everywhere uniform light-brown, made up of small punctulations. In life the color is not much different; the interspaces are brassy-yellow, fading to white on side; head and fins pale-bluish.

This species, which reaches a length of 6 inches, is found in tropical America on both coasts; abundant in tide-pools and about coral reefs from Guaymas to Peru and Florida to Uruguay, but much less abundant than the species of *Eupomacentrus*. It is common about Porto Rico, but difficult to capture, as it is found chiefly where nets can not be used to advantage. We have specimens, from 0.75 to 5.5 inches, from San Juan, Puerto Real, and Culebra.

Chatodon cauda bifurca, fasciis S. albis, Linnæus, Mus. Adolph. Frederici, I, 64, "India."

Chatodon saxatilis Linnæus, Syst. Nat., ed. X, 276, 1758; after Mus. Ad. Fr.

Chatodon mauritii Bloch, Ichthyol., III, 213, pl. 109, 1785, Brazil; on a bad drawing by Prince Maurice

Chatodon marginatus Bloch, Ichthyol., III, 98, pl. 207, 1787, Martinique; on a drawing by Plumier.

Glyphisodon moucharra Lacépède, Hist. Nat. Poiss., IV, 542, 1803, Brazil, etc.; after various authors.

Chatodon sargoïdes Lacépède, Hist. Nat. Poiss., IV, 453, 1803, Martinique; on a drawing by Plumier.

Glyphidodon troschelii Gill, Proc. Ac. Nat. Sci. Phila. 1862, 150, Cape San Lucas.

Glyphidodon saxatilis Poey, Fauna Puerto-Riqueña, 336, 1881; Stahl, l. c. 77 and 164, 1883.

Abudcduf saxatilis, Jordan & Evermann, l. c., 1561, 1898.




FIG. 64.—*Abudcduf saxatilis*.

Family LI. LABRIDÆ. The Wrasse-fishes.

Body oblong or elongate, covered with cycloid scales; lateral line well developed, continuous or interrupted, often angularly bent. Mouth moderate, terminal; premaxillaries protractile; maxillaries without supplemental bone, slipping under the membranaceous edge of preorbital; anterior teeth in jaws usually very strong and canine-like; teeth of jaws separate or soldered together at base, not forming a continuous plate; no teeth on vomer or palatines; lower pharyngeals completely united into one bone, without median suture, this bone T-shaped or Y-shaped, its teeth conical or tubercular. Lips thick, longitudinally plicate. Nostrils round, with 2 openings on each side. Dorsal fin continuous, spinous part usually long, its spines rather slender, 3 to 20 in number; anal similar to soft dorsal, with 2 to

3 spines. Ventrals thoracic, 1, 5, inserted below pectorals. Branchiostegals 5 or 6; pseudobranchiae well developed; gills 3.5, slit behind last arch small or obsolete; gill-membranes somewhat connected, sometimes joined to narrow isthmus. Air-bladder present; no pyloric caeca.

This family comprises 60 genera and 450 species, chiefly of the tropical seas, living among rocks or kelp, many of them brilliantly colored and some valued as food-fish. Most of them feed upon mollusks, the dentition being adapted for crushing shells. As in most other large groups there is considerable difference of opinion as to the characters which should be used in dividing the Labroids into genera. The tendency with all recent writers has been toward a rather minute subdivision. The numbers of vertebrae seem to us to yield characters of the highest importance. Other characters not to be neglected can be drawn from the size of the scales, the numbers of the dorsal spines, and the dentition. The degree of squamation of the head seems to us to have an importance lower than that attributed to it by Bleeker and Günther, but it may be used for generic subdivision.

- a. Dorsal spines 8 or more, usually well distinguished from soft rays; anal spines 2 to 6.
- b. Vertebrae and dorsal spines not in greatly increased numbers; vertebrae 22 to 29; dorsal spines 8 to 13; anal spines 2 or 3; species of tropical or subtropical seas.
- c. Vertebrae 27 to 29 (so far as known); dorsal spines usually 12 (11 to 14); sides of head more or less scaly; preopercle serrulate or entire.

HARPINÆ:

- d. Anterior canines strong; lower pharyngeals large, with large, tubercular teeth; spinous dorsal not enveloped in scales; lower jaw naked; species mostly of large size and bright coloration, inhabiting semitropical seas.
- e. Dorsal spines about 14, the 3 or 4 anterior falcate, produced in long streamers; body deep and compressed, anterior profile steep; teeth uniserial; no posterior canine; cheek and opercles scaly; bases of soft dorsal and anal scaly; soft parts of vertical fins produced; scales moderate (40)..... LACHNOLAIMUS, 106
- ee. Dorsal spines 11 or 12 (rarely 13), none of them produced in filaments; cheek and opercles scaly; body oblong; back not greatly elevated.
- f. Soft dorsal and anal each with a scaly sheath at base; scales large (about 32); posterior canine present; soft dorsal and anal elevated, produced behind..... HARPE
- ff. Soft dorsal and anal without sheath of scales; preopercle serrulate (at least in young); soft dorsal and anal more or less falcate.
- g. Scales large, about 30; lower limb of preopercle scaly; posterior canine present; anterior canines $\frac{4}{4}$ DECODON

CLEPTICINÆ:

- dd. Anterior teeth small, bluntish, not canine-like; no posterior canine; mouth very small, terminal; snout short and blunt; dorsal and anal enveloped in scales, except produced tips of both fins; caudal deeply forked; dorsal spines almost hidden by series of scales; head everywhere closely scaled, except on lips and snout; scales of body large; preopercle serrulate; gillrakers slender, short; pectoral falcate; lower pharyngeals very small, Y-shaped, their teeth small, very blunt, and coalescent; vertebrae 10 + 17=27; dorsal spines, 12..... CLEPTICUS
- ee. Vertebrae 23 to 26; dorsal spines 8 or 9; anterior canines strong, 2 to 4 on each side in each jaw; head mostly naked; preopercle entire. Species of the Tropics, mostly of small size and bright coloration.

JULIDINÆ:

- h. Lateral line complete and continuous.
- i. Snout not tubiform; preopercle entire; teeth uniserial, none chisel-shaped.
- j. Cheeks and opercles naked.
- k. Scales large, 25 to 30 in lateral line; anal spines 2 or 3.
- l. Dorsal spines 9; dorsal enlarged, without scaly sheath; scales of breast not enlarged.
- m. Anterior canines all normal in position; lower pharyngeals T-shaped, with numerous teeth; anal spines 3.
- n. Posterior canine well developed on both sides; dorsal spines pungent; anterior canines $\frac{2}{4}$ IRIDIO, 107
- ll. Dorsal spines 8; no posterior canines; anterior canines $\frac{2}{2}$, normal in position; a low sheath of scales at base of dorsal; dorsal spines pungent; anal spines 3, never 2..... CHLORICHTHYS

XERICHTHYINÆ:

- hh. Lateral line interrupted posteriorly, beginning again on level of axis of body, on caudal peduncle; scales large, 20 to 30 in lateral line; dorsal spines 9; anal spines 3; anterior canines $\frac{2}{2}$.
- o. Posterior canine present; snout slender, anterior profile not convex; cheek and opercles scaly; dorsal spines pungent, the three anterior longer and with filamentous appendages; dorsal and anal with a scaly sheath; scales very large..... DORATONOTUS, 108
- oo. Posterior canine none; anterior profile more or less convex; head naked, except usually a few scales below eye; body more or less strongly compressed; ventrals thoracic, inserted below the pectorals.
- p. Scales very large, about 20 in lateral line, which is placed on first row of large scales below dorsal sheath; anterior dorsal spines not detached; head not trenchant above..... XYRULA
- pp. Scales large, about 26 in lateral line, which is placed on second row of large scales below dorsal sheath.
- q. First two dorsal spines joined by membrane to the others and inserted nearly above base of pectoral.
- r. Upper anterior profile of head not trenchant, curve of head not parabolic; cheek not very deep... NOVACULICHTHYS
- rr. Upper anterior profile of the head sharply trenchant, its curve parabolic; cheek very deep, eye near upper profile..... XYRICHTHYS

Genus 106. *LACHNOLAIMUS* Cuvier & Valenciennes.

Body strongly compressed, back sharp and elevated, profile long and steep. Snout sharp; mouth low, horizontal, jaws narrow; premaxillary slipping under membranaceous edge of the very broad preorbital, which is twice depth of eye. Teeth in front prominent, canine-like, in a single series; no posterior canines. Cheek and opercles with imbricate scales; scales of moderate size, thin, adherent; lateral line complete. Dorsal with 14 spines, first 3 strong, falcate, produced in long streamers in adult, membranes between these spines very low, filamentous tips longer than head; other spines all low, gradually shorter to the eleventh; second dorsal and anal much produced; caudal lobes falcate; third anal spine strong; pectoral and ventrals short.

This genus contains a single species, a large, showy fish of tropical America, remarkable for the long, streamer-like filaments on the dorsal spines.




FIG. 65.—*Lachnolaimus maximus*.

184. *Lachnolaimus maximus* (Walbaum). “*Capitan*”; *Perro Perro*; *Hog-fish*.

Head 3; depth 2.2; eye 5.4; snout 2.4; maxillary 2.1; mandible 2; interorbital 4.1; preorbital 4.4; D. XIV, 11; A. III, 10; pectoral 1.6; ventral 1.6; caudal 1; scales 8–38–14. Body deep and compressed, back elevated, anterior profile long and steep, nearly straight, slightly concave before eye; scales rather large and thin; premaxillaries very protractile; 4 strong canines in front of upper jaw, 1 pair in front of lower with 2 smaller conical teeth between them; sides of jaws with small, uniserial, bluntly conical teeth; first 3 dorsal spines produced in long filaments, reaching to end of fin, rest of spines low; soft dorsal with anterior rays somewhat elevated, nearly as long as head; caudal deeply lunate.

Color in life, brick-red, scales edged with reddish-yellow thus giving a cross-hatched appearance; a large black spot on last 4 dorsal rays and body at their base; caudal red, with 2 pale-brown crossbars and 2 spots of same on each tip, anal similar to caudal; ventral dark purplish-red; pectoral lemon; purplish lines radiating downward and forward from eye. In spirits, grayish, snout blackish; caudal with 2 faint vertical dark bars near base; base of last rays of soft dorsal with deep black spot extending upon body. The sexes differ in color and in size of mouth, and individuals vary much with age, but the species may be known at once from the long dorsal streamers.

“*El Capitan*” is found throughout the West Indies and north to Key West and the Bermudas, abundant about reefs and rocks. It has been recorded from various places in southern Florida, the Tortugas, Bahamas, St. Bartholomew, Cuba, Martinique, St. Thomas, Santo Domingo, Porto Rico, and Jamaica. Examples, 6 to 13 inches long, were obtained at Ensenada del Boqueron, Arroyo, and Isabel Segunda. Others were seen in the fishermen’s boats at each of these places and also at Culebra Island.

This is a large and showy fish, reaching a weight of 10 to 15 pounds, or even 20 pounds in the Bermudas. It changes much in appearance in the course of its growth, which accounts for the many specific names it has received. The large adult male is remarkable on account of a heavy black blotch

over the forehead and eyes. The name "hog-fish" refers to the swine-like appearance of the head, jaws, and teeth. Like all other members of the family, it feeds chiefly upon small fish and upon bottom mollusks and crustaceans. It is an important food-fish throughout its range and is one of the most common and attractive species seen in the wells of fishing boats at Key West. It is a favorite food-fish in Cuba, though at one time its sale was forbidden by law on account of the supposed poisonous character of its flesh. This opinion obtains to some extent in Porto Rico, but apparently the flesh of only the large individuals is believed to possess any deleterious properties. Whether there is any good reason for this belief is doubtful.

Suillus (the Great Hog-fish), Catesby, Nat. Hist. Carolina, etc., pl. 15, 1750, Bahamas.

Labrus maximus Walbaum, Artedi Piscium, 261, 1792.

Lachnolaimus suillus Cuvier, Règne Animal, cd. II, vol. 2, 257, 1829, Bahamas; after Catesby.

Lachnolaimus aigula Cuvier & Valenciennes, Hist. Nat. Poiss., XIII, 277, 1839, St. Bartholomew.

Lachnolaimus dux Cuvier & Valenciennes, Hist. Nat. Poiss., XIII, 285, Martinique.

Lachnolaimus caninus Cuvier & Valenciennes, Hist. Nat. Poiss., XIII, 288, 1839, St. Thomas and Santo Domingo.

Lachnolaimus psittacus Cuvier & Valenciennes, Hist. Nat. Poiss., XIII, 291, 1839, Porto Rico.

Lachnolaimus suillus Poey, Fauna Puerto-Ríqueña, 336, 1881.

Lachnolaimus maximus, Jordau & Evermann, l. c., 1579, 1898.


Genus 107. IRIDIO Jordan & Evermann. Doncellas.

Body oblong, compressed, not elevated, covered with large scales, there being 25 to 30 in course of lateral line, which is not interrupted, but abruptly bent posteriorly. Scales on breast rather smaller. Head naked, compressed, conic. Preopercle entire. Teeth large, upper jaw with 2 strong canines in front, none of them bent backward; lower jaw with 4 anterior canines, a posterior canine tooth directed forward on each side of upper jaw. Dorsal spines 9; anal spines 3, graduated; ventrals inserted under axil of pectoral. Gillrakers short and feeble; gill-membranes slightly joined to a narrow isthmus. Vertebrae 10 + 15 = 25.

There are numerous species of this genus, most of them brilliantly colored, abounding in kelp in the tropical seas. All of them are American. The genus is very close to the Old World *Halichares*, differing chiefly in the dentition and in the presence of 3 anal spines instead of 2.

- a. Caudal fin very slightly concave, truncate when spread open, outer rays longer than middle ones; body deep and compressed, depth about 2.75 in length; ventral fins filamentous, outer ray produced, more than twice as long as inner ray; scales before dorsal not crossing middle line, in about 5 series.
- b. Side below spinous dorsal without dark crossbar; general color bluish (male), or bronze (female), with many sky-blue spots, most distinct posteriorly; sky-blue spots and streaks on head; a stripe passing through upper part of eye; fins with blue stripes; a dark axillary spot; end of pectoral dusky..... *radiatus*
- bb. Side below spinous dorsal with a very broad, blackish crossbar..... *kirschii*, 186
- aa. Caudal fin rounded or subtruncate, outer rays not produced, shorter than middle rays.
- c. Scales before dorsal large, in 4 to 6 rows, not crossing median line; snout moderately pointed.
- d. Ventral fins with outer ray produced, more than twice length of inner.
- e. Side without conspicuous dark lateral band and with a distinct dark vertical bar, extending downward from spinous dorsal; axillary spot obscure; body rather elongate, depth about 3.75 in length; profile not steep; posterior canines rather small; head with black streaks and spots above; caudal sharply barred..... *garoti*
- cc. Side with a broad blue-black lateral band extending from eye to tip of caudal; back above this, dark brown or bluish; spinous dorsal with no conspicuous black spot; a dark-blue stripe from eye to nape; fins mostly blue-black with pale edgings; middle and base of caudal dusky; tip of pectoral dusky; profile rather steep; body rather robust, depth 3.33 in length..... *cyanocephalus*
- dd. Ventral fins with outer ray not produced, its length not more than twice that of inner rays; side with a dark lateral band; species of small size.
- f. Spinous dorsal with a conspicuous blue-black spot between fifth and seventh spines; body not very slender, depth 3.8 in length; a dark band from snout through eye to opercle, lateral band on side broader than eye and placed a little above opercular band, lateral band extending nearly to tip of caudal; no second dark band below it; a faint dark spot under last dorsal ray and one at base of pectoral above; 2 or 3 narrow bluish-white stripes across cheek; body and fins in life with bright colors which fade in alcohol..... *maculipinna*
- ff. Spinous dorsal pale, black spot very small or wanting; body slender, depth 4 in length; opercle with a conspicuous black spot; a blue-black band from snout through eye and across opercles to base of caudal, not extending on fin; a narrower and fainter band from lower base of pectoral to above anal, these bands growing fainter with age and sometimes disappearing, lower always wanting in adult; no axillary spot; no distinct bands across cheek; fins mostly pale, with bright red and blue colors in life, young and deep-water individuals often showing a black spot at base of caudal and sometimes a dark spot near middle of dorsal, with sometimes a larger one at base of its last ray; angles of caudal black in adult, lower pharyngeals T-shaped, the anterior limb very short..... *bivittatus*, 185
- aaa. Caudal fin double-concave, median portion convex, outer rays more or less produced in adult (fin rounded in young); scales before dorsal in 6 or 7 rows, not crossing median line; a blue-black spot close behind eye, sometimes obsolete in adult.

- g. Lateral line without blue-black spot; ventrals with outer rays scarcely filamentous, about reaching tips of pectorals.
- h. Tubes of pores of lateral line distinctly branched, branches usually 3 in number; body moderately slender, depth a little less than length of head and 3.75 to 4 in body; head 3.4.
- i. Eye large, 1.66 in snout.
- ii. Eye small, 2.66 in snout; brownish above, bright violet-red below; many blue spots above; a crescent on base of pectoral; caudal edged with violet and with convergent streaks of yellow..... *pocyi*
- hh. Tubes of pores of lateral line all simple or very nearly so, not trifid; body very slender, depth much less than length of head, 4.5 in body; head 3.4; snout very sharp, anterior profile of head straightish and not steep; snout 2.8 in head; eye 2 in snout; pectoral moderate, 1.6 in head; color in spirits, pale, unmarked, except for small black spot behind eye.
- j. Color in life, olive-green, bluish below; back with blue spots; a yellow band on side with vague outlines; posterior parts paler, with rows of blue spots; head with blue bands; dorsal and anal rosy, with blue spots..... *caudalis*
- jj. Color in life, olivaceous; a broad band-like area of orange mingled with violet spots along side backward from head to middle of body, lower edge of this band serrate; below this a pale-violet band, darker behind; still lower a yellow stripe; head olivaceous, marked with blue; preorbital scarlet, with 3 violet stripes; opercles bright red, with 3 violet stripes, postocular black spot in the uppermost; dorsal and anal orange and yellow, with blue spots; caudal with convergent bands of orange forming reticulations around blue spots..... *pictus*

FIG. 66.—*Iridio bivittatus*.

185. *Iridio bivittatus* (Bloch). *Slippery Dick*; *Doncella*.

Head 3.2; depth 4; eye 6.3; snout 3; interorbital 4.7; preorbital 4.7; D. ix, 11; A. iii, 12; pectoral 1.7; ventral 1.9; caudal 1.6; scales 2-28-8. Scales before dorsal reduced in size, in about 4 rows, not crossing median line; ventral fin with first ray moderately produced, about twice inner ray; caudal slightly rounded.

Color in spirits: Grayish, a purple band from snout through eye to base of caudal, much widened on opercle; a fainter band, more blue, from base of pectoral to above end of anal; an oblong purple-black spot on upper edge of opercle, bordered behind by blue; a blue band from nape to eye; chin with a blue transverse band. An example, 2.5 inches long, has longitudinal bands brown, traces of very faint vertical dark bars, and 2 small dark spots near base of front of soft dorsal.

This species is gorgeous in color, but of little value as food, the flesh being dry and insipid. It is found throughout the West Indies, north to Pensacola and Beaufort, N. C., and south to Brazil, and is usually very abundant along weedy shores and reefs. Specimens, 2.5 to 5.5 inches long, obtained at Mayaguez, Fajardo, and Culebra; 4 from San Geronimo.

Sparus radiatus Linnaeus, Syst. Nat., ed. XII, 472, 1766, Carolina; not *Labrus radiatus* L., ed. X.

Labrus bivittatus Bloch, Ichthyol., pl. 284, fig. 1, 1792; from a painting by Plumier, made at Martinique.

Labrus psittaculus Lacépède, Hist. Nat. Poiss., III, 522, 1800, Martinique; from a copy of Plumier's painting.

Julis humeralis Poey, Memorias, II, 212, 1860, Havana; adult.

Chærojulis grandisquamis Gill, Proc. Ac. Nat. Sci. Phila. 1863, 206, Beaufort, N. C.

Chærojulis arangoi Poey, Enumeration, 109, 1875, Havana; young.

PlatyGLOSSUS florcalis Jordan & Gilbert, Proc. U. S. N. M. 1882, 287, Pensacola; young.

Iridio bivittatus, Jordan & Evermann, l. c., 1595, 1898.

186. *Iridio kirschii* Jordan & Evermann.

Head 3.4; depth 4; eye 5.5; snout 3; D. ix, 12; A. iii, 11; scales 2-28-8. Body elongate, compressed; anterior profile gently convex; head rather long; snout pointed; mouth moderate, horizontal, maxillary not reaching vertical of eye by half its diameter; teeth strong, curved; opercular flap long, reaching past base of pectoral; caudal peduncle compressed, its least width 5 times in its least depth;

scales large, about 4 rows in front of dorsal fin, scales not extending across median line; tubes of pores of lateral line distinctly branched, branches usually 3 in number, lateral line running on third row of scales to near vertical of last dorsal ray, where it drops to fifth row. Fins moderate; longest dorsal spine about 3 in head, rays about 2.5; anal similar to soft dorsal; pectoral 1.5 in head; ventral with outer rays somewhat produced, about twice length of inner ray, or 1.75 in head; caudal fin almost truncate, slightly, if at all, double-concave.

Color in life: Side beautiful pale-green; scales of back and upper part of side pale-brown at base; under parts greenish-white; a rich green axillary spot and a few scales above it with rich-green edges; pectoral brick-red on base, followed by a narrow blue bar, then by a broader lemon-colored bar, the red bar on base extending on body about 3 scales below pectoral; cheek lemon-colored, a rosy bar from mouth to base of pectoral, another from mouth to eye, and 2 others from preopercle across opercle, a narrow, wavy, rosy line under eye, a somewhat paler bar from mouth downward upon lower jaw, and another forward on snout, meeting its fellow in front; a rich blue postocular spot, with a small yellow one above it; humeral region rich green; a small black spot at base of last dorsal ray; dorsal and anal fins lemon; numerous short rosy bars crossing dorsal, which is bordered by rosy; anal base rosy, then a narrow green stripe, followed by a very narrow pale border; caudal pale greenish-lemon, with rosy on middle rays; pectoral pale-greenish; ventral white with slight rosy wash; iris green and red, pupil black. In alcohol all the colors fade to a yellowish olivaceous, the postocular spot and the one at base of last dorsal ray becoming black.

Found in the West Indies, south to Bahia; recorded from Cuba, Jamaica, St. Croix, and Bahia. The species reaches a foot in length. Two specimens, each 4.25 inches long, in the collections from Porto Rico—one from San Antonio Bridge, the other from Fajardo. We have compared our specimens with the type of *L. kirschii* (No. 43303, U.S.N.M.) and find them to agree perfectly.

Iridio kirschii Jordan & Evermann, Check-List of Fishes of North and Middle America, 413, 1896, Bahia; name only; Jordan & Evermann, Fishes North and Middle America, 1598, 1898.

Genus 108. DORATONOTUS Günther.

Body compressed; head not compressed to an edge anteriorly, its profile in front straight or concave; preorbital not very deep; mouth rather wide; teeth in a single series, 2 large canines in front of each jaw; a posterior canine; cheek and opercles scaly; gill-membranes united, free from isthmus; scales large; lateral line interrupted behind, beginning again lower down; dorsal fin with 9 strong pungent spines, some of the anterior elevated, median spines short, so that the outline of fin is concave; caudal rounded. Colors brilliant. Size small. This genus contains two known species, among the most beautiful of the *Labridæ*, and the genus to which they belong is one of the best defined in the group.

a. Scales large 1.5-20-6.5; color grass-green, over entire body and caudal fin..... *megalepis*, 187
aa. Scales smaller, 1-26-6; caudal fin white, with 2 small reddish-brown spots..... *deccoris*, 188

187. *Doratonotus megalepis* Günther.

Head 2.75; depth 2.66; D. IX, 10; A. III, 9; scales $1\frac{1}{2}$ -20-6 $\frac{1}{2}$. Body much compressed, moderately elevated, its greatest width behind the head two-sevenths of its height; caudal peduncle short and deep, its length but little more than one-half its height; profile from dorsal to nape convex, carinated; occiput and supraorbital region depressed and flat, snout protruding, profile of top of head thus strongly concave. Snout slender, sharp, compressed, its length 3.2 in head; mouth wide; maxillary 4 in head; teeth growing gradually larger anteriorly, the 2 front teeth in each jaw distinctly the largest, canine-like, diverging, opposed to each other; a small but distinct posterior canine in upper jaw, none in lower; eye moderate, little wider than interorbital width, 5 in head; cheek with a single series of large scales, 4 in number; opercle covered with 5 or 6 similar scales; gill-membranes broadly united, free from isthmus. Dorsal spines robust and pungent, the first 3 with conspicuous filamentous appendages; first and second spines with their filaments about equal, 1.5 in head; without their filaments second spine is slightly the longer, equaling distance from end of snout to middle of eye, the fin rapidly descending to fourth spine, which is one-half as long as second, then gradually rising to ninth and highest, which is, however, shorter than the following soft rays; longest soft ray 1.66 in head; anal spines similar to those of dorsal fin, the longest about one-half head; caudal evenly convex, its longest ray 1.5 in head; ventrals short, about one-half length of head, an elongate scale between them at base; pectoral reaching beyond ventrals, but not to vent, 1.75 in head; membranes of vertical fins

with elongate scales on basal portion; lateral line following outline of back 1 scale beyond end of dorsal fin, thence interrupted and continued on 4 scales of middle of caudal peduncle.

Color in life: Very intense grass-green, about uniform over the body; head more yellowish, slightly paler below; opercle mesially a little darker; iris red, with a green ring; dorsal, anal, and caudal grass-green, mottled with light-orange; tips of lower spines green, of short ones orange; ventrals deep-green, membranes largely orange; pectoral light-yellowish.

Found in the West Indies north to Key West, rare. Length 2.75 inches. Here described from the type of *Doratonotus thalassinus*, obtained with a seine in eelgrass at Key West. The Porto Rican collections contain but a single specimen of this beautiful little species. It is 1.75 inches long and was seined among the algæ at Hucares.

Doratonotus megalepis Günther, Cat., IV, 125, 1862, St. Kitts; Jordan & Evermann, l. c., 1611, 1898.

Doratonotus thalassinus Jordan & Gilbert, Proc. U. S. N. M. 1884, 28, Key West.

188. *Doratonotus decoris* Evermann & Marsh.

(PLATE 29.)

Head 2.6; depth 3.4; eye 4; snout 3.5; maxillary 4; interorbital 4.6; D. ix, 10; A. iii, 9; pectoral 1.6; ventral 2.2; caudal 1.6; scales 1-26-6. Body moderately elongate, compressed throughout; back a little elevated, caudal peduncle deep and rather long; dorsal and ventral outlines nearly alike, dorsal somewhat more strongly arched; anterior profile not trenchant, almost straight from snout to front of dorsal, very slightly convex in front of dorsal and very slightly concave between eye and tip of snout; head pointed, interorbital space broad and flat; eye large, high in position, middle of pupil nearer tip of snout than end of opercle; snout long, somewhat longer than diameter of eye, moderately produced, lips broad in front, characteristically labroid; mouth not large, maxillary not reaching front of orbit, the jaws equal, armed with strong sharp teeth, about 4 canines in front of upper jaw, 2 in front of lower; teeth on sides of jaws also canine-like, smaller than those in front, but not distinctly different from them; a few smaller teeth behind the main row of large ones; vomer and palatines toothless; soft dorsal and anal each with a basal sheath of about two rows of large scales, that of dorsal extending over half the fin or more, that of anal lower, the fins otherwise naked; dorsal fin continuous, with a shallow notch, spines slender and pungent, second longer than first, following ones graduated to fifth, which is shortest, thence increasing in length to ninth, which is longest, 2.3 in head; soft dorsal with its middle rays highest, 2.2 in head; anal with three slender, sharp, graduated spines, third longest, 2.2 in head; soft part similar to soft dorsal, longest rays 2.3 in head; pectoral large, symmetrical, of 11 rays, middle ones longest, reaching past tip of ventral nearly to the vent; ventral moderate, pointed, reaching halfway to vent; caudal rounded; scales large, cycloid, lateral line on second row below dorsal, interrupted near end of dorsal and beginning again on row below, on caudal peduncle.

Color in life: Body chiefly green, darker green on back, lighter below; lower parts of head and breast light yellow; a broad white bar from eye obliquely across cheek and opercle, bordered above by an undulating maroon line and below by a similar but fainter line; a brown bar from eye to snout; 4 dusky spots near base of dorsal extending as fainter shades downward and slightly forward to or beyond lateral line, 1 from in front of dorsal, 2 under spinous dorsal, and 1 under soft rays; short pale-blue bars or spots on breast and about pectoral; iris blue, a pinkish border surrounding pupil; dorsal greenish, soft part with yellow shade, a pale-blue edging to whole fin, a maroon border to green color posteriorly just inside the pale-blue edge, a small dark spot on membrane between seventh and eighth rays and a blue spot on membrane of first spine; anal colored like soft dorsal, the maroon border extending from first spine to last ray inside the pale edging, the dark spot between sixth and seventh rays; ventral green near base, pale-blue outwardly, the green color bordered by maroon spots; pectoral plain pale-green; caudal very pale transparent blue, a wedge-shaped maroon spot on the 2 upper rays near tip and a corresponding one on the 2 lower rays, base of wedge on outer ray; base of caudal with a pale undulate vertical bar bordered in front by a black line. In spirits, pale-green, maroon markings faintly persistent, becoming dusky.

One specimen, the type (No. 49363, U. S. N. M.), 1.45 inches long, taken in the seine at Ponce, January 30, 1899. Two other specimens were seined in the algæ on a little sandy islet near Playa de Ponce, but they were inadvertently lost. This is one of the most beautiful of fishes and differs from *D. megalepis* chiefly in the smaller scales and the coloration.

Doratonotus decoris Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 354, Ponce, Porto Rico.

FAMILY LII. SCARIDÆ. The Parrot-fishes.

Body oblong, moderately compressed, covered with large cycloid scales as in the *Labridæ*. Mouth moderate, terminal. Teeth in jaws more or less coalescent, at least at base; lower pharyngeals much enlarged, united in a concave or spoon-shaped body, their teeth broadest transversely and truncate, arranged in mosaic; dorsal continuous, usually 1x, 10; A. 11, 9; 23 to 25 scales in lateral line; vertebrae about 11 + 14 = 25. Sexes similarly colored, the coloration almost always brilliant. Fin rays essentially the same throughout the group, the squamation varying little except on head.

The *Scaridæ* comprise 7 genera and about 110 species, herbivorous fishes of the tropical seas, especially abundant about coral reefs; often of large size, not much valued as food, the flesh being soft and pasty. The species in the various genera are very closely related, being distinguished chiefly by the coloration and the dentition, both series of characters being highly specialized. Jordan & Evermann recognize 5 genera of *Scaridæ* as inhabiting the waters of America north of Panama, of which only 3 are as yet known from Porto Rico.

SPARISOMATINÆ:

- a. Lower pharyngeal broader than long, flattish or basin-shaped; gill-membranes broadly joined to isthmus, not forming a fold across it; lateral line subcontinuous; scales about head few and large, those on cheek in one row; lower jaw projecting; teeth whitish or rosy.
- b. Dorsal spines flexible; teeth more or less distinct, at least anteriorly.
- c. Teeth in each jaw in few series, not imbricated or quincunx; lateral teeth of each jaw coalescent in a more or less continuous cutting edge, teeth more free anteriorly and not adnate to dental plate..... CRYPTOTOMUS
- bb. Dorsal spines stiff, pungent; teeth of upper jaw more or less coalescent.
- d. Teeth of each jaw chiefly coalescent, jaws divided by a rather indistinct median suture SPARISOMA, 109

SCARINÆ:

- aa. Lower pharyngeal spoon-shaped, much longer than broad; teeth of jaws fully coalesced, each jaw divided by a distinct median suture; gill-membranes forming a fold across isthmus; dorsal spines flexible; lateral line interrupted behind, beginning again lower down on caudal peduncle of tail; scales about head rather numerous, those on cheek in two or more series; lower jaw included.
- c. Teeth and jaws whitish or rosy in color SCARUS, 110
- α. Teeth and jaws blue or bluish-green PSEUDOSCARUS, 111

Genus 109. SPARISOMA Swainson. Viejas.

Lower pharyngeal broader than long, subhexagonal, its surface moderately concave or flattish; teeth in each jaw largely coalescent in adult, their tips more or less separate in young, the edge, especially of lower jaw, sometimes uneven; median suture in each jaw present, but not well defined; 1 to 4 radiating canines sometimes present on each side of upper jaw above its cutting edge; gill-membranes broadly united to isthmus; dorsal spines pungent; upper lip double for its entire length; lower jaw projecting beyond upper; lateral line not interrupted, passing gradually from its row of scales posteriorly to the series next below it; tubes of lateral line much branched; scales about head large, those on cheek in a single row, those on median line in front of dorsal 3 or 4 in number.

Species of rather small size, most of them American; some of them showily colored.

SPARISOMA:

- a. Upper jaw with one or more canines above its cutting edge (these occasionally obsolete on one or both sides); coloration often brilliant.
- b. Caudal truncate or slightly rounded, angles not acute.
- c. Posterior canines 2 to 4 on each side.
- d. Caudal fin with more or less of black on posterior margin, yellowish at base. Canines strong, 4 (rarely 3) on each side..... *xystrodon*, 189
- dd. Caudal fin without black in adult; 1 or 2 more or less distinct whitish bars across chin.
- e. Canines 3 or 4 on each side, radiating horizontally; axil with little or no blue, but with a dusky blotch partly hidden by fin; front steeper and less curved than in *S. hoplomystax*; body and fins mottled, but much less so than in *S. hoplomystax*.
- f. Canines 3 on each side; pores of lateral line with but 2 branches; sides of head much dotted with black; caudal barred..... *atomarium*
- ff. Canines 4 on each side; tubes of lateral line much branched; a distinct narrow streak of blue downward and forward from eye; caudal nearly plain dusky-olive; anal mottled..... *radians*
- cc. Canines 2 or 3 on each side; axillary region extensively deep-blue in life, this forming a large blotch around and on base of pectoral; a curved series of small white specks around the blue on base of pectoral; fins all mottled, anal with 3 darker areas; body with 3 faint pale lengthwise streaks, more or less obscure, 2 of these bounding a more or less interrupted dusky band from eye to base of caudal *hoplomystax*, 190
- cc. Posterior canine single on each side; body rather stout..... *niphobles*, 191

- bb.* Caudal fin simply lunate, outer rays more or less exerted, but not twice as long as inner rays and much shorter than head; canine single on each side (rarely obsolete or duplicated).
- g.* Head with a scarlet stripe from below eye to angle of mouth; a small scarlet streak behind eye; color, chiefly purplish brown; a round spot of yellow and black behind head, just below lateral line; fins chiefly red; angles of caudal black; axillary spot obscure *aurofrenatum*, 192
- gg.* Head without scarlet stripe.
- h.* Pectoral very long and sharp, 5 in body; body brownish, not striped; no axillary spot *oxybrachium*
- hh.* Pectoral moderate, less than one-fifth length.
- i.* Color, dark reddish-brown, with white mottlings; no yellow or black spot; belly abruptly red; fins mostly cherry red; axillary spot obsolete; body rather deep; scales large, their outlines well defined *abildgaardii*, 193
- ii.* Color, brownish, with 3 or 4 pale longitudinal streaks, upper running to a faint pale blotch on back of tail between 2 dark-brown blotches; caudal distinctly pale-edged behind and more distinctly barred than in *S. flavescens*; spot at base of pectoral brownish and very faint; about 4 small dusky blotches along base of dorsal, last one most distinct at base of last ray; caudal with many crossbars and blotches; snout dusky; chin with 1 or 2 whitish crossbars; caudal concave, with sharp angles; dorsal and anal mottled with brown; pectorals and ventrals plain; young with dark opercular blotch and dark points about eye *distinctum*
- bbb.* Caudal fin in adult deeply forked, upper lobe about as long as head and twice or more length of inner rays; caudal fin variegated.
- j.* Canines 4 to 6 on each side; pores of lateral line excessively branched, each with several (6 to 8) much divided branches. Color, bright greenish-blue (the sides sometimes with a blue band); caudal lobes blue, middle rays red; dorsal and anal red; pectoral yellowish, axillary spot large, black, edged with red *chrysopterum*, 194
- jj.* Canines 1 or 2 on each side; upper and lower caudal lobes greenish.
- k.* Opercle without black-and-yellow spot; pores of lateral line each with 4 or 5 nearly simple branches *lorito*, 195
- kk.* Opercle with an inky-black spot, in front of which is a golden spot; no spot at base of pectoral *viride*, 196
- EUSCARUS:
- aa.* Upper jaw never with posterior lateral canines; colors dull, usually mottled brown or greenish.
- l.* Caudal slightly rounded, angles not produced.
- m.* Scales of lateral line and some on nape and opercle black; dorsal spines stout; olive, vertical fins edged with violet; axil violet. *strigatum*
- ll.* Caudle lunate, or truncate with sharp angles (rounded in very young).
- n.* Caudal fin distinctly barred with irregular brown spots and markings.
- o.* Body without distinct pale longitudinal streaks above; caudal not evidently pale-edged; spot on base of pectoral blackish and distinct; no evident pale or dark blotches on back of tail.
- p.* Caudal lunate or subtruncate in adult, rounded in young. General color olivaceous or reddish-brown, clouded, and washed with cherry-red; lower fins mostly red; pectoral light-orange; chin pale, with whitish cross-band. *flavescens*, 197
- pp.* Caudal truncate, not at all lunate in adult, angles very slightly produced.
- q.* Color olivaceous or bluish-green, a whitish streak below mouth; a dark axillary spot usually present; a whitish band on caudal; fins dotted. *rubripinne*, 198
- mm.* Caudal fin not crossbarred.
- r.* Axillary spot black, very distinct; outer rays of caudal considerably produced, length of exerted part one-third to one-half that of head.
- s.* Caudal red, its outer rays green; axillary spot very distinct; body olivaceous, nearly plain-reddish below; some greenish-blue on head; a faint greenish streak running backward from angle of mouth *brachiale*, 199
- ss.* Caudal violaceous, its outer rays one-half head; a dark spot at base of pectoral; color dusky-red, scales of back and sides with red spots. *maschalespilos*
- rr.* Axillary spot faint or wanting; coloration uniform dark purplish-violet; 3 large scales on cheek; dorsal spines rather slender, but pungent; caudal emarginate; tubes of each scale of lateral line much ramified and extending over whole scale; teeth of moderate size, very distinct on edges of jaws *frondosum*

189. *Sparisoma xystrodon* Jordan & Swain. *Loro; Parrot-fish.*

Head 3.2; depth 3; eye 5; snout 2.75; interorbital width 3.8; preorbital 3.2; scales 1-25-5; D. IX, 10; A. 11; pectoral 1.5; ventral 1.9; middle caudal rays 1.75. Body oblong wedge-shaped, heavy forward; profile in a regular curve from snout to dorsal fin; upper jaw with 3 or 4 exerted canines on each side above cutting edge, largest in front of angle of mouth curved outward and backward, one next in front less curved, directed downward and slightly forward (sometimes backward), the pair at median suture small; upper lip covering most of upper jaw. Scales large and thin, a row of five on cheek; 4 scales in front of origin of dorsal fin; dorsal low, spines pungent, longest shorter than snout, rays longer, about 2 in head; anal similar to soft dorsal but lower; caudal slightly convex when spread, outer rays slightly shorter than middle ones; pectoral broad, reaching a little past tips of ventrals; ventrals short, not reaching anus.

Color in life: Bright olive-green above, paler below, the upper parts very much mottled, speckled with white and marbled with coppery-red; head similarly green, dotted with whitish above; a narrow ring of bright blue below eye, interrupted above; a narrow blue stripe from eye to angle of mouth; a blue spot behind eye; blue and coppery markings on opercle; lower part of head

light yellow, a blue band around margin of lower jaw; axil and a spot at base of pectoral in front above deep blue-black; dorsal orange flesh-color, the tip paler; caudal yellowish at the base, paler beyond, its posterior portion more or less black, a few whitish dots near base; anal light blue and reddish, the tip dusky; ventrals pale; pectoral light-yellowish; lining of opercle blackish; jaws pale. Some specimens are pearly-bluish rather than green above, livid below, blue on head paler, and the red of a light-yellowish carmine. Some highly colored specimens are greener, with belly bright yellow, brightest at throat; anal and caudal chiefly jet-black. (Jordan & Evermann, Key West specimens.)

Color in alcohol: Dark olive or grayish above, with slight green shade; lower parts paler; snout paler; blue line around eye and from eye to angle of mouth persistent; blue at base of pectoral usually changing to black; dorsal, anal, and caudal mottled; edge of anal and tip of caudal usually blackish.

This species can best be distinguished from *S. hoplomystax* and related species by the blue line about eye and from eye to snout, the dark edge of caudal fin, and the presence of 2 or 3 canines on each side of upper jaw. It is found in the West Indies and north to Key West, where it is very common in *Fucus* and eelgrass, along with *S. hoplomystax*. Length 4 to 7 inches. It is very common about Porto Rico, our collection containing specimens from San Juan, Aguadilla, Mayaguez, Puerto Real, Ensenada del Boqueron, Guanica, Ponce, Arroyo, Hucares, Fajardo, Culebra, Isabel Segunda, and San Geronimo. It was especially abundant at Ponce and Mayaguez.

Sparisoma rystrodon Jordan & Swain, Proc. U. S. N. M. 1884 (July 1, 1884), 99, Key West, Fla.; Jordan & Evermann, l. c., II, 1630, 1898.




FIG. 67.—*Sparisoma hoplomystax*.

190. *Sparisoma hoplomystax* (Cope).

Head 3; depth 2.8; eye 3.5; snout 2.8; interorbital 3.7; preorbital 5.4; D. ix, 10; A. ii, 9; pectoral 1.5; ventral 1.8; caudal 1.5; scales $1\frac{1}{2}$ –25–5. Body deep, the back considerably elevated; one strong posterior canine on each side of upper jaw, directed slightly backward, often a second canine in front of this and a small one directed downward on each side of front of upper jaw above the cutting edge and close to the median suture, these not evident in our specimen; caudal rounded.

Color in spirits: Grayish and olivaceous, caudal dark, with a few white mottlings and a narrow pale edge, the large scales at its base pale; dorsal and anal mottled with dark; side with about three longitudinal series of white spots smaller than pupil, one on row of scales above lateral line, another on row immediately below it, and the third two rows lower; about six spots in each row; some fainter and irregular white spots on side below.

This species ranges from Bahia through the West Indies north to Key West and Cape Florida; it has been recorded from Key West, St. Martins, and St. Lucia, but is apparently not common in Porto Rico. The collection contains but one individual, 2.75 inches long, from Mayaguez, of this type of coloration, and it seems to be identical with specimens identified as *Scarus radians* and *Sparisoma cyanoleucum* in the U. S. National Museum.

- Labrus radians* Castelnau, Anim. Nouv., etc., Amérique du Sud, 29, 1855; not *Scarus radians* Cuvier & Valenciennes.
Scarus radians Günther, Cat., IV, 211; Jordan & Gilbert, Synopsis, 906, 1888; not of Cuvier & Valenciennes.
Scarus hoplomystax Cope, Trans. Am. Philo. Soc. 1869, 462, St. Martins.
Sparisoma cyanolone Jordan & Swain, Proc. U. S. N. M. 1884, 98, Key West.
Sparisoma hoplomystax, Jordan & Evermann, l. c., 1632, 1898.

191. *Sparisoma niphobles* Jordan & Bollman. Loro; Parrot-fish.

Head 3; depth 2.67; eye 4.25; snout 3; D. ix, 10; A. ii, 9; scales 1-25-5. Body short and stout; snout short, obtuse; dorsal outline gently and regularly arched from tip of snout to origin of dorsal, thence in a regular descent to caudal peduncle; ventral outline less arched; a minute canine directed downward on front of upper jaw on each side close to median suture, and a larger, stronger lateral one, directed backward; lower jaw somewhat projecting, teeth coalesced except at tips; upper lip double for its entire length, covering most of upper jaw; cheek with one row of 5 large scales; lateral line complete, dropping from second to third row of scales under last dorsal rays, the tubes each with 3 to 5 branches; 4 scales on median line before dorsal. Origin of dorsal above base of pectoral, spines pungent, about equal to snout in length, rays a little longer; caudal slightly rounded, outer rays about 2 in head; anal similar to soft dorsal, its origin about under first dorsal ray; pectoral broad and short, about equal to snout and eye; ventral short, equal to snout and half eye.

Color in alcohol: Dirty gray, body mottled and speckled with whitish; an obscure whitish line, often not evident, from eye along lateral line to caudal, and a plainer, better-defined one from base of caudal to opercular flap, where it connects with a similar line from eye across opercle; in some specimens lower half of body abruptly paler, the dividing line running from lower edge of eye to the middle of caudal; base of pectoral and axil black, probably greenish in life; lower parts of side usually with more numerous white specks; under parts of the body paler; tip of lower jaw brown, followed by a white band, then another brown one, behind which is a series of 6 large white spots, middle ones largest, these spots sometimes confluent; subopercle and breast blotched with brown on bases of scales; dorsal with about 4 faint dark bars; anal similarly marked; caudal mottled, a pale line at tip. In life, grayish with washings of greenish and reddish; base of pectoral green.

This species is known only from the Bahamas, Cape Florida, Key West, and Porto Rico. It is quite abundant about Porto Rico, where numerous specimens were obtained from San Geronimo, Aguadilla, Mayaguez, Ensenada del Boqueron, Ponce, Hucars, and Fajardo. It was abundant at each of these places. Length 5 or 6 inches or less. Like most other species of this genus this fish frequents the patches of algae in shallow water. From *S. hoplomystax* (Cope), which it closely resembles, it may best be distinguished by the presence of but a single lateral canine and the numerous small white specks, which are diagnostic.

- Sparisoma niphobles* Jordan & Bollman, Proc. U. S. N. M. 1888 (Sept. 20), 551, Green Turtle Cay, Bahamas; Jordan & Evermann, l. c., II, 1633, 1898.

192. *Sparisoma aurofrenatum* (Cuvier & Valenciennes).

Head 3.1; depth 3.1; eye 5; snout 2.6; interorbital 4.5; preorbital 4.6; D. ix, 10; A. ii, 9; pectoral 1.4; ventral 1.7; caudal 1.5; scales 1½-25-5. One posterior canine (obsolete or broken on right side in our specimen); caudal fin lunate, upper lobe very slightly longer than lower.

Color in life: Whitish green above, slightly rosy on side, greenish below; an orange splotch bordered at upper anterior edge by a black blotch just below lateral line under third dorsal spine; eye red; a brick-red bar from mouth under eye to opercle, above this 2 small oblong spots of same; rest of head bluish, teeth white, dorsal pale yellowish-red; anal rich red with bluish border; caudal olive at base, then a broad, rich blood-red bar or crescent, then a pale, whitish terminal crescent; outer rays clear red, tips of fin black; pectoral very pale-rosy, the anterior ray dark; ventral pale-rosy, a blue blotch at base of pectoral. In spirits, dark above, pale green on sides and below; a pale-yellow stripe from angle of mouth to below posterior margin of eye, bordered below by black; a short, narrow bar of same color behind eye; a distinct black blotch on fourth and fifth scales of lateral line, and a much larger blotch of yellow below the black; upper ray and base of pectoral dark, front of base black; anal with a dark line at base and a dark edge; angles of caudal inky-black, blotch on lower lobe the larger. Easily recognizable, even in spirits, by the characteristic persistent markings.

A West Indian species, known from Havana, Sombrero Key, Santo Domingo, Jamaica, Porto Rico, St. Thomas, and St. Lucia. One specimen, 7.5 inches long, obtained by us at Arroyo.

Scarus aurofrenatus Cuvier & Valenciennes, Hist. Nat. Poiss., XIV, 191, 1839, Santo Domingo.

Scarus mincefrenatus Poey, Memorias, II, 279, 1860, Cuba.

Sparisoma aurofrenatum, Jordan & Evermann, l. c., 1634, 1898.

193. *Sparisoma abildgaardi* (Bloch). "Loro Colorado"; Red Parrot-fish.

(PLATE 30.)

Head 3; depth 2.7; eye 6.4; snout 2.1; interorbital 4.5; preorbital 3.4; D. IX, 10; A. II, 9; pectoral 1.3; ventral 1.6; caudal 1.3; scales 2-25-5. Dorsal and ventral outlines of body alike, snout without fleshy hump; caudal fin lunate, lobes equal; a single canine on each side of upper jaw above cutting edge (in one small specimen 2 on right side).

Color in life: Fins and lower parts below a line from tip of under jaw to base of caudal under the end of lateral line, red, edges of scales paler; body above grayish, edges of scales black, bases of those below lateral line rosy; brownish-red wavy stripes radiating from eye below; oblique pale-blue stripes on dorsal, tips of row of smaller scales at base of anal blue, and 3 blotches on upper and 3 on lower ray of caudal; a few scales at base of caudal with pale yellow; iris yellow, pupil bordered narrowly with red; membranous edge of opercle black.

This gorgeously colored parrot-fish is generally common from Cuba south to Brazil and has been recorded also from Martinique, Jamaica, Santo Domingo, and Havana. The collection from Porto Rico contains 2 specimens, 11 and 12.5 inches long, respectively, from Arroyo, where many others were seen. While it was not noticed elsewhere it is probably not rare about the island.

Vieja, Parra, Descr. Dif. Piezas, Hist. Nat., 58, pl. 28, fig. 2, 1787, Cuba.

Scarus abildgaardi Bloch, Ichth., pl. 259, 1791, America; from a specimen sent by Professor Abildgaard.

Scarus coccineus Bloch & Schneider, Syst. Ichth., 289, 1801, Cuba; after Parra.

Scarus aurcoruber Lacépède, Hist. Nat. Poiss., IV, 55, 163, 1803, Martinique; on a drawing by Plumier.

Scarus amplus Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., 324, taf. 5, pl. 25, 1842, Brazil (fide Guichenot; not seen by us).

Scarus erythrinoides Guichenot, Scarides Mus. Paris, 10, 1865, Santo Domingo.

Scarus oxybrachius Poey, Synopsis, 342, 1868, Cuba.

Sparisoma abildgaardi, Jordan & Evermann, l. c., 1635, 1898.

194. *Sparisoma chrysopterum* (Bloch & Schneider).

"Loro Verde"; "Cotoro Verde"; *Vieja*; Blue Parrot-fish.

Head 3.1; depth 3; eye 5.8; snout 2.1; interorbital 5.5; preorbital 3.2; D. IX, 10; A. II, 9; pectoral 1.5; ventral 1.9; caudal 1.2; scales 1½-25-5. Dorsal and ventral outlines nearly alike, dorsal somewhat more strongly arched; 4 to 6 strong canines on each side of upper jaw, those behind usually pointing outward or backward, those in front forward, sometimes one or two small ones at suture; pores of lateral line much branched, more so than in any other known Porto Rican species; caudal deeply lunate, outer rays much produced, upper lobe the longer, twice length of middle rays. Known from the other species by the canines, produced rays of the caudal, and the blue color.

Color in life: Rich greenish-blue or bluish-green above; edges of scales greener, their bases darker; under parts rich blue; head greenish-blue, somewhat mottled; lower jaw and breast sky-blue; dorsal pale rose; anal pale blue at base, then broadly pale brick-red, then a very narrow pale-blue border; caudal reddish at base, blue on outer rays extending to tips; abaft the red a pale crescent, then a broad blood-red one, and last a narrow bluish-green border; pectoral pale lemon, a black splotch at base above; ventral pale-greenish.

In spirits, body nearly everywhere a faded-blue, brighter on head and under pectoral; dorsal and pectoral pale; ventral chiefly pale, with traces of blue; anal soiled-grayish; middle of caudal pale, its edge dark, outer rays greenish; a black blotch on base of pectoral above.

This species is generally common from the West Indies to Brazil, and is known from Havana, Santo Domingo, Jamaica, Porto Rico, Martinique, St. Thomas, St. Croix, St. Kitts, Guadeloupe, and Bahia. It reaches a good size and is of some value as a food-fish. Two fine specimens, each a foot long, were obtained from Arroyo and Isabel Segunda, and others were seen at each of these places and at Culebra.

Vieja, Parra, Descr. Dif. Piezas, Hist. Nat., 58, pl. 28, fig. 4, 1787, Cuba.

Scarus chrysopterus Bloch & Schneider, Syst. Ichth., 286, pl. 57, 1801, American seas.

Scarus chloris Bloch & Schneider, Syst. Ichth., 289, 1801, Cuba; after Parra.

Scarus lateralis Poey, Memorias, II, 219, 1860, Cuba.

? *Scarus spinidens* Guichenot, Scarides, 15, 1865, Bahia.

Sparisoma chrysopterum, Jordan & Evermann, l. c., 1636, 1898.

195. *Sparisoma lorito* Jordan & Swain. "*Loro*"; *Parrot-fish*.

Head 3; depth 2.9; eye 5.3; snout 2.3; interorbital 5.2; preorbital 3.5; D. ix, 10; A. ii, 9; pectoral 1.5; ventral 1.8; caudal 1.3; scales $1\frac{1}{2}$ -25-5. Back moderately elevated, dorsal outline much more strongly arched than ventral; eye high in position; one posterior canine on each side of upper jaw, and one or more small ones near suture; caudal deeply lunate, upper lobe very slightly the longer.

Color in spirits: Uniform greenish; a very distinct deep-black blotch on base of upper rays of pectoral; outer rays of caudal darker than middle. Much resembling *S. brachiale* in spirits, but differing in the presence of canine teeth.

A West Indian species, known from Havana, Haiti, Jamaica, Porto Rico, Sombrero, and Barbados. Two specimens, each about 10 inches long, were obtained by us—one at Puerto Real, the other at Arroyo—and others were seen. It is of some value as a food-fish.

Sparisoma lorito Jordan & Swain, Proc. U. S. N. M. 1884 (July 1), 95, Havana; Jordan & Evermann, l. c., 1637, 1898.

196. *Sparisoma viride* (Bonnaterre). "*Loro Verde*"; "*Cotoro*"; *Dark-green Parrot-fish*.

Head 3; depth 2.6; eye 7.5; snout 2; interorbital 4; preorbital 3.1; D. ix, 10; A. ii, 9; pectoral 1.4; ventral 1.7; caudal 1.3; scales 2-25-5. Body rather deep, back elevated, profile nearly straight from snout to dorsal; caudal deeply lunate, outer rays considerably produced, upper lobe slightly the longer; 2 canines on each side of upper jaw.

Color in life: Rich bluish-green, edges of scales as well as their bases brownish; top of head light clear brown down to level of pupil, this brown patch extending from occiput to near tip of snout, which is green with a narrow brown border; a narrow brown bar backward from eye to humeral region, and a similar one from mouth backward to opercle, this connecting with the narrow line around each lip; space above this rich green, below paler green; upper edge of opercle with a small pale-lemon spot, below it a green one, then rest of opercle with a broad reddish-orange border; subopercle with a very narrow orange border; caudal peduncle with a large irregular orange blotch covering about 5 or 6 scales; dorsal pale-reddish, the sheathing scales at base very rich green; caudal dark green at base and on outer rays nearly to tips; next a crescent of rich orange, the horns extending to tips; lastly a broad crescent of paler green or bluish green; anal green at base, then a broad rich orange band, then a broad blue border; pectoral green on upper and lower rays, paler between; ventral rich blue-green on outer ray, rest pale-orange; teeth white, with slight bluish tinge.

In spirits: Pale-greenish, top of head grayish-brown, a band of same color behind eye; side of head blue, abruptly demarcated from color of top of head; a pale band from angle of mouth across cheek, then narrowing and curving upward across opercle; lips blue, their edges pale; membranous edge of opercle pale, a bright-yellow spot above; anal with blue margin and base, pale in middle; caudal with blue margin, widening on middle rays, a yellow crescent-shaped band in front of it, rest of fin greenish; pectoral bluish above and in front, pale-yellowish below and behind; scales at base of caudal pale yellow.

A West Indian species; known from the Bahamas, Sombrero Key, Jamaica, Porto Rico, St. Thomas, and St. Croix. Obtained by us at Arroyo and Culebra, where it is probably common. It attains a length of 2 feet or more and, though used for food, it is not of much value.

Pisces viridis bahamensis (the Parrot-fish), Catesby, Nat. Hist. Carolinas, etc., II, 29, pl. 29, 1738, Bahamas.

Scarus viridis Bonnaterre, Enc. Méth., X, 96, 1788, Bahamas; after Catesby.

Scarus catesbyi Lacépède, Hist. Nat. Poiss., IV, 16, 1803, Bahamas; after Catesby.

Scarus catesbyi Cuvier & Valenciennes, Hist. Nat. Poiss., XIV, 183, 1839, Bahamas.

Scarus melanotis Bleeker, Ichth. Notizen, I-X, 4, 1862, St. Croix.

Sparisoma viride, Jordan & Evermann, l. c., 1638, 1898.

197. *Sparisoma flavescens* (Bloch & Schneider). "*Vieja Colorada*"; *Mud Parrot-fish*.

Head 3.1; depth 2.7; eye 5.5; snout 2.4; interorbital 4; preorbital 4; D. ix, 10; A. ii, 9; scales 1-25-5. Body rather stout, compressed, heavy forward; profile evenly curved from tip of snout to dorsal fin, then in a long regular slope to caudal peduncle; mouth normal, gape not reaching eye by a distance greater than half diameter of orbit; jaws pale, lower jaw projecting; no canines; upper lip covering all of upper jaw; scales large and thin, a single row of 5 scales on cheek, 4 scales before origin of dorsal; tubes of lateral line dividing into 4 or 5 branches, covering most of scale; top of head with numerous small papillae, forming elevated lines posteriorly and behind eye; origin of dorsal over base of pectoral; longest dorsal spine about 2.5 in head; dorsal rays slightly longer; anal similar to soft

dorsal; pectoral rather long, reaching past tips of ventrals, rays about 1.4 in head; ventrals about 1.7 in head; caudal fin usually lunate, lobes pointed and projecting, especially in adult; in the young the fin is sometimes merely truncate, never rounded as in *S. rubripinne*.

Color of adult in life: Olivaceous, clouded with light and dark, and usually flushed with pinkish, especially below, edges of the scales more yellow-olive; scales of belly and lower parts light orange-red toward their bases, giving a decidedly reddish cast; dorsal mottled with different shades of olive; caudal creamy, mottled and barred with darker orange, markings more distinct on outer edge; ventrals and anal rich cherry-red, mottled and barred with brown; pectoral light orange-red, color formed by narrow orange cross-streaks on a paler ground; a light band across lower jaw, which is otherwise brown; a dusky or black blotch at base of pectoral; blackish blotches sometimes present on body at base of soft dorsal. In alcohol, the red and yellow fade and become pale and the general color is brownish, paler below; usually a dark blotch at base of soft dorsal and one on base of pectoral.

S. flavescens is close to *S. rubripinne*, from which it can be distinguished most readily by the difference in shape of caudal fin and its lighter coloration. In *S. rubripinne* the caudal is more or less rounded, while in *S. flavescens* it is lunate or truncate, tips of lobes always being pointed.

The range of this species extends from Biscayne Bay and Key West southward to Rio Janeiro. It is generally common, and is known from Cape Florida, Key West, Havana, Bahamas, Tortugas, St. Thomas, Jeremie, Haiti, Port au Prince, Jamaica, Porto Rico, St. Lucia, and Rio Janeiro. It is abundant in Porto Rico. The collection contains specimens from the following localities: San Juan, Mayaguez, Puerto Real, Guanica, Ponce, Ensenada del Boqueron, Hucares, Fajardo, Isabel Segunda, and Culcra Island. It rarely exceeds a foot in length. It is very abundant at Key West, swarming everywhere about the island in the eelgrass. At Havana it is apparently equally common, the numbers seen in the market exceeding that of all other species of the genus combined. Its flesh, although not unpleasant in flavor, is soft and rather poor. In the Havana market it is usually called *Vieja Colorado*, but the species of this group are seldom distinguished by the fishermen.

Vieja, Parra, Deser. Piezas Dif. Hist. Nat., 59, pl. 28, fig. 4, 1787, Cuba.

Scarus flavescens Bloch & Schneider, Syst. Ichth., 290, 1801, Cuba; after Parra.

Scarus squalidus Poey, Memorias, II, 218, 1860, Cuba.

Sparisoma flavescens, Jordan & Evermann, l. c., 1640, 1898.

198. *Sparisoma rubripinne* (Cuvier & Valenciennes). *Vieja*; *Loro*; *Parrot-fish*.

Head 3.3; depth 2.9; eye 5.5; snout 2.5; interorbital 3.7; preorbital 4; D. ix, 10; A. ii, 9; scales 1-25-5. Body stout, compressed; head large, somewhat rugose in the adult; mouth small, gape not reaching vertical of eye by a distance equal to half of its diameter; lower jaw projecting; no canines; suborbital with venules; pores of lateral line with 3 or 4 branches; 4 scales in front of dorsal. Fins moderate; dorsal spines pungent, longest about equal to snout; dorsal rays a little longer, about equal to those of anal; pectoral long, 1.3 in head; ventral shorter, about 1.9 in head; caudal truncate in adult, usually rounded in young, never lunate as in *S. flavescens*, outer rays about equal to snout and eye.

Color in life (a specimen 9 inches long): Head and side dirty mottled-brown; 3 brown bars on lower jaw; fins all mottled-brown and yellowish; belly white. Another specimen, 4 inches long, from San Juan, had the sides bluish-olivaceous; edges of scales darkest; white on belly; side of head with 2 irregular brownish bars; tip of lower jaw brownish, followed by a white stripe, then a broad brown bar followed by white; dorsal pale, mottled with brown, especially near base; caudal crossed by alternating bars of brown and paler; irregular brown blotches at base of caudal; anal rosy, blotched with brown; pectoral pale; ventrals pale-rosy, inner rays white; jaws white. Another specimen, 4 inches long, from Aguadilla, was described as grayish, paler below; chin with 2 broad brown bands; 4 black blotches on back, first on and at base of first dorsal spine, second on base of sixth and seventh rays, third on about eleventh and twelfth rays, and fourth on last ray but one, all these extending somewhat on sides; iris green, rest of eye silvery and brownish; dorsal and anal fins mottled reddish-brown; caudal brown, with 2 or 3 irregular cross-series of pale spots; ventrals reddish; pectoral pale. In alcohol, general color grayish-brown, paler below; head brown; chin with 2 or 3 broad brown bands separated by white ones; base of pectoral dark; a dark blotch on body at edge of opercle; 4 rather distinct dark blotches on back extending upon dorsal fin, which is mottled with dark and light; caudal dark, with 2 irregular cross-rows of white spots, more or less confluent in anterior row; extreme tip of tail with a pale border; anal barred and mottled like dorsal; pectoral and ventral pale; jaws pale.

A West Indian species (length 6 to 9 inches) closely resembling *S. flavescens*, from which it can best be distinguished by its rounded (or rarely truncate) caudal, and absence of whitish or snowy blotches on sides. It is known from Santo Domingo, Porto Rico, Jamaica, and Martinique. It is common about Porto Rico, the collection containing specimens from San Juan, Aguadilla, Mayaguez, Ensenada del Boqueron, Puerto Real, Guanica, Ponce, Arroyo, Hucares, and Fajardo.

Scarus rubripinnis Cuvier & Valenciennes, Hist. Nat. Poiss., XIV, 199, 1839, Santo Domingo.

Scarus virens Cuv. & Val., Hist. Nat. Poiss., XIV, 203, 1839, Porto Rico (Coll. Plée), and Martinique (Coll. Achard).

Scarus circumnotatus Poey, Memorias, II, 423, 1861, Havana.

Scarus truncatus Poey, Synopsis, 339, 1868, Havana; Poey, Fauna Puerto-Riqueña, 336, 1881; Stahl, l. c., 78 and 164, 1883.

? *Scarus emarginatum* Poey, Synopsis, 340, 1868, Havana.

Sparisoma rubripinne, Jordan & Evermann, l. c., 1640, 1898.

199. *Sparisoma brachiale* (Poey).

Head 3.2; depth 3.1; eye 5.6; snout 2.2; interorbital 5.3; preorbital 3.4; D. IX, 10; A. II, 9; pectoral 1.4; ventral 1.7; caudal 1.3; scales $1\frac{1}{2}$ -25-5. Body elongate, back moderately elevated, dorsal and ventral outlines much alike; eye about midway between tip of snout and end of opercle, very high in position; no canines; outer rays of caudal considerably produced, making fin rather strongly lunate, middle part truncate or slightly convex; lateral line interrupted near end of dorsal, beginning again on row below, the two ends usually overlapping.

Color in life: Rich red above, paler below, edges of scales white; fins all rosy-red; teeth white. In spirits, very pale greenish, dark olive in one specimen; in some cases with rosy flushes on side; a large jet-black spot at base of upper rays of pectoral, not extending into axil; no other distinct markings.

A West Indian species known only from Cuba, Jamaica, and Porto Rico. Seen by us at Aguadilla, Arroyo, and Isabel Segunda, where specimens 9 to 12.5 inches long were obtained; also collected by Mr. Gray at San Geronimo.

Scarus brachialis Poey, Memorias, II, 345, 1861, Cuba.

Scarus humeralis Poey, Memorias, II, 422, 1861, Havana; based on an old drawing; a black axillary spot, sown with white points.

Sparisoma brachiale, Jordan & Evermann, l. c., 1641, 1898.

Genus 110. SCARUS Forskål. Loros; Parrot-fishes.

Lower pharyngeals spoon-shaped, ovate-oblong, transversely concave; teeth in each jaw fully coalescent, appearing as tessellations on the surface; jaws with distinct median sutures; edges of jaw even, teeth whitish or rosy in color, never green. Upper pharyngeals each with 2 rows of teeth; gill-membranes scarcely united to the narrow isthmus, across which they form a broad fold; dorsal spines flexible, scarcely different from soft rays; upper lip laterally double, interior fold becoming very narrow or obsolete mesially; lower jaw included in the closed mouth; lateral line interrupted posteriorly, commencing again on next series of scales below; tubes of lateral line scarcely branched; scales on cheek in 2 to 4 rows; scales in front of dorsal on median line 6 to 8. D. IX, 10; A. II, 9 in all species; scales $2\frac{1}{2}$ -24-6. Body robust.

SCARUS:

- a. Upper jaw with from 1 to 4 posterior canines.
- b. Check with 2 or 3 rows of scales.
- c. Head with a longitudinal band; a yellow longitudinal stripe on body; outer rays of caudal not colored like inner; caudal subtruncate.
- d. Outer rays of caudal blackish or greenish, darker than median rays.
- e. Yellow stripe above pectoral about on a level with eye; outer rays of caudal deep greenish-blue; upper jaw with 1 posterior canine (rarely duplicated); 2.5 rows of scales on cheek; head with 2 bluish-green stripes, interspaces reddish or yellow; dorsal and anal each with 2 green bands and 1 orange one, anal having a roundish blue spot on membrane between every 2 rays. General color bluish-green mixed with orange. *punctulatus*
- cc. Yellow stripe above pectoral, mostly below level of eye; outer rays of caudal blackish, rest of caudal green; upper jaw with 2 posterior canines; 2 rows of scales on cheek, upper part of head dark-green, below eye bright yellowish-green, with bluish markings on opercle; dorsal bright-green at base; ventrals pale; base of pectoral with a blue-black mark. General color in life, bright-green; darker on back, paler below. *boltoni*
- dd. Outer rays of caudal orange, lighter than median rays, its edge blackish; yellow stripe above pectoral, below level of green stripes on head, which are nearly horizontal; upper jaw with 1 posterior canine (rarely duplicated); 2.5 rows of scales on cheek; head with 2 bluish-green stripes, interspaces reddish or yellow; dorsal and anal each with 2 green bands and 1 orange one, the latter without blue spots; basal band of dorsal not broken into green spots. General color bluish-green, mixed with orange. *teniopterus*
- cc. Head without longitudinal bands; posterior canines 2 to 4.

- f. Caudal truncate; 2 series of scales on cheek, and 2 scales on lower preopercular limb; canines 2 or 3 on each side. Color uniform violet-purple; vertical fins very dark *aracanga*
- ff. Caudal fin innate, the outer rays more or less produced; cheek with 2.5 or 3 rows of scales; posterior canines 3 or 4; color (dried skin) plain brownish, caudal in one specimen darker, or paler mesially, its border and angles dark *trispinosus*
- bb. Cheek with 4 rows of scales; angles of caudal more or less salient.
- g. Color dusky-olivaceous, some scales with a rosy blotch at base; dorsal edged with dusky; caudal dark, pale at base, and with pale shades, its angles little produced; opercle with blue blotches; canines 3; snout rather acute; 7 scales before dorsal *euzamila*
- gg. Color, dark sky-blue; jaws with bright colors *vetula*, 200
- ggg. Color brown, with 2 brown lateral bands *gnathodus*
- aa. Upper jaw without canines; 2.5 series of scales on cheek.
- h. Third (partial) row of scales of cheek of 3 or 4 scales, those of upper row little larger than those of second row.
- i. Caudal slightly rounded, its outer rays not produced.
- j. Side of body with 2 broad, dark longitudinal shades; sides of belly each with 3 sharply defined lines, each on a row of scales, these stripes running from breast to beyond front of ventrals and usually becoming faint or even obsolete in old individuals.
- k. Stripes on side of breast, if present, whitish. Color dark reddish-brown above, paler below; back dark; side with 2 dark parallel stripes of color of back, separated by paler interspaces, upper one extending backward from eye; snout above bluish-brown; a narrow whitish streak running from head along middle line of belly; a faint dark spot on base of pectoral; caudal pale orange-red, outer rays somewhat barred with brown; dorsal orange, edged with bluish; other fins nearly plain *croiccasis*, 201
- kk. Stripes on side of breast, if present, inky-blue. Color bright-green, olivaceous above, paler below, lower half of body becoming posteriorly more and more yellow, and on lower half of caudal peduncle bright light-yellow, this color being brightest above front of anal; longitudinal shades on side of body bright crimson, separated on head by a band of green; no spot on base of pectoral; caudal fin green, its lower half yellow; dorsal, anal, and pectoral green, at least at base; ventrals yellow *evermanni*
- jj. Sides of body without distinct, broad, darker stripes. Color brown, no bands or lines upon body or head; dorsal spotted with violet and edged above and below with yellow, like caudal; caudal without spots; yellow line near edge and another along base of dorsal *flavomarginatus*
- ii. Caudal truncate, its points slightly salient; side with a broad whitish band *acutus*
- hh. Third (partial) row of scales on cheek of 1 or 2 scales only; scales of upper row much larger than those of second row; caudal subtruncate, its outer rays more or less produced, becoming much elongate with age; adult with a fleshy hump above snout.
- l. Color bright-blue, young more or less shaded with reddish-brown; fins mostly blue. Size large *cervulus*, 202
- ll. Color dusky-olive; a pale-yellowish streak from upper part of eye to upper base of caudal *emblematicus*

200. *Scarus vetula* Bloch & Schneider. *Mud-fish; Vieja; Old Wife.*

(PLATE 31.)

Head 2.7; depth 2.7; eye 7; snout 2.3; interorbital 2.8; preorbital 4.2; D. IX, 10; A. II, 9; pectoral 1.5; ventral 2; caudal 1.3; scales 2-25-6. Dorsal and ventral outlines alike, head pointed; 2 posterior lateral canines in upper jaw; caudal margin slightly concave between outer produced rays.

Color in life: Scales of trunk bluish-green, their edges pink; head variegated, the ground-color in shades of grayish, pink, and blue; a wavy green stripe from slightly below angle of mouth obliquely upward to below posterior margin of eye, bordered on both sides with yellow; a similar shorter stripe on snout, similarly bordered; a bent green bar before eye and a very short one behind; lower lip and chin blue; iris yellow; outer half of dorsal fin abruptly green, the line of demarcation curving obliquely upward from base of first spine, thence nearly straight to tip of ninth ray; yellow below the green, shading into rosy on base of fin, which has oblique pale-blue bars, one near base of each interradial membrane, below which is an oblong deep-blue blotch; anal colored exactly like dorsal, but without oblique pale-blue bars; pectoral with its upper edge blue, below which is a band of pink spreading out over the base, rest of fin light-green; ventral with spine and first ray blue, second yellow, the rest pale, with rosy membranes; caudal with upper and lower produced rays blue, the two inner of the produced rays rosy at base, yellowish at ends, rest of fin green; several scales at base of caudal rosy.

This is one of the most gaudy of the parrot-fishes, the combination of colors being very extraordinary. It is thus far known only from Cuba and Porto Rico, where it is not uncommon. The specimen in our collection from which the colored drawing (plate 31) was made is 12.5 inches long and was obtained at Puerto Real. We have compared it with a specimen in the National Museum which has been identified with Poey's *S. superbus* and find that they agree perfectly. The identity of *S. superbus* with *S. vetula* seems certain, though there are some slight discrepancies in the descriptions.

Vieja, Parra, Descr. Dif. Piezas, Hist. Nat., 58, pl. 28, fig. 1, 1787, Havana.

Scarus vetula Bloch & Schneider, Syst. Ichth., 289, 1801, Cuba; after Parra; Jordan & Evermann, l. c., 1649, 1898.

Scarus superbus Poey, Memorias, II, 218, 1860, Cuba.

201. *Scarus croicensis* (Bloch). *Bullon*.

Head 2.9; depth 3.3; eye 4.8; snout 2.9; interorbital 3; preorbital 4.8; D. IX, 10; A. II, 9; pectoral 1.6; ventral 2; caudal 1.9; scales $1\frac{1}{2}$ -25-6. Body elongate, little elevated, dorsal and ventral outlines alike, jaws without canines; caudal fin slightly rounded.

Color in life: Brownish above, pale below; a broad band of brown from eye to base of caudal, a narrower one below this from base of pectoral to caudal; a pale-bluish line bordering upper band above and below; two or three white or pale-bluish longitudinal streaks on sides of abdomen.

A species of small size but great abundance, from southern Florida and the West Indies; known from Key West, the Tortugas, Bermuda, Havana, Porto Rico, St. Thomas, Jamaica, St. Lucia, Martinique, and St. Croix. Numerous specimens obtained by us at San Antonio Bridge, Boqueron, Mayaguez, Hucars, Fajardo, and Culebra; 8 by Mr. Gray at San Geronimo. It probably does not exceed 5 or 6 inches in length.

Scarus croicensis Bloch, Ichth., pl. 221, 1790, St. Croix; Jordan & Evermann, l. c., 1650, 1898.

Scarus insule-sanctæ-crucis Bloch & Schneider, Syst. Ichth., 312, pl. 62, fig. 2, 1801; after Gronow.

Scarus alternans Cuvier & Valenciennes, Hist. Nat. Poiss., IV, 200, 1839, Martinique.

Pseudoscarus lincolatus Poey, Repertorio, II, 239, 1868, Cuba.

202. *Scarus cœruleus* (Bloch). "*Loro*"; *Blue Parrot-fish*; "*Tumble-nose*."

(PLATE 32.)

Head 3; depth 3; eye 8; snout 2.3; interorbital 2.8; preorbital 3.7; D. IX, 10; A. II, 9; pectoral 1.6; ventral 1.8; caudal 1; scales 2-25-6. Body elongate, back scarcely elevated, but snout with a large fleshy hump so that head is not pointed, hump much more prominent in old individuals; anterior profile strongly concave above eye in a specimen 20 inches long, nearly straight or slightly convex in one of 14 inches; eye small; caudal truncate or slightly rounded between the produced outer rays; no canines; pores of lateral line little branched.

Color in life: Everywhere uniform turquoise-blue, bases of vertical fins much deeper blue; iris blue, pupil with a narrow brownish-yellow border; a brownish-yellow stripe bordering lower jaw; faint wine-colored shades on side of head below and behind eye; head, back, belly, and tips of fins greenish-blue.

The blue parrot-fish reaches a length of 2 or 3 feet and a weight of 12 to 20 pounds. It has the widest distribution of any member of the family, being found from Chesapeake Bay southward to Florida and throughout the West Indies. It has been recorded from St. George Island (Maryland), Cape Charles, Bahamas, Key West, Tortugas, Havana, Martinique, Jamaica, and Porto Rico. Two specimens, 14.5 and 20 inches long, were obtained by us at Culebra Island, and many others were seen in the boats of the St. Thomas, St. Croix, and Tortola fishermen who frequent the waters about Culebra and Vieques islands. Though evidently not held in high esteem, it is doubtless the most important food species of the parrot-fishes occurring about Porto Rico, due chiefly to its abundance and large size, and the catch seems to be always saved by the fishermen.

So far as known, none of the parrot-fishes ever takes the hook, and they are therefore usually caught in some sort of trap. The trap in common use about Porto Rico is shown on page 31. About Culebra Island these traps or baskets are set in 5 to 8 fathoms of water, and are usually baited with large chunks of cactus pulp, made white by trimming off the green outer part.

These large parrot-fish are taken in considerable numbers, along with hog-fish, doctor-fish (medicos), groupers, snappers, and grunts. The name "tumble-nose" is given by fishermen to unusually large blue parrot-fish, on account of the peculiar topography of the nose.

Novacula carulea (the Blue-fish), Catesby, Nat. Hist. Carolina, etc., 18, pl. 18, 1743, Bahamas.

Loro, Parra, Deser. Dif. Piezas, Hist. Nat., 57, pl. 27, fig. 1, 1787, Cuba.

Trompa, Parra, l. c., fig. 2.

Coryphæna carulea Bloch, Ausländische Fische, II, 120, pl. 176, 1786, in part, Bahamas; after Catesby and a figure by Aubriet, altered from a figure by Plumier.

Scarus loro Bloch & Schneider, Systema Ichthyol., 288, 1801; after *Loro* of Parra.

Scarus trilobatus Lacépède, Hist. Nat. Poiss., IV, 21, 1803, Martinique; on a drawing by Plumier.

? *Scarus holoqancos* Lacépède, Hist. Nat. Poiss., IV, 45, 1803, Martinique; on a copy by Aubriet of a drawing by Plumier; the copy colored entirely blue in order to represent this species; the original drawing probably intended for

Sparisoma chrysopterygum; the same copy by Aubriet, the original of Bloch's engraving of *Scarus caruleus*.

Scarus obtusus Poey, Memorias, II, 217, 1860, Cuba; adult.

Scarus nuchalis Poey, Memorias, II, 220, 1860, Cuba; young.

Scarus caruleus, Jordan & Evermann, l. c., 1652, 1898.

Genus 111. PSEUDOSCARUS Bleeker. Guacamaias.


This genus differs from *Scarus*, as here understood, chiefly in the deep green or blue color of its highly modified jaws and teeth. The species are mostly of large size and robust form. Five species of this genus are recognized by Jordan & Evermann as occurring in American waters north of Panama, only one of which is thus far known from Porto Rico.

PSEUDOSCARUS:

- a. Upper jaw with canines; caudal fin with angles much exerted, especially in adult; soft dorsal and anal ending in points; 2.5 rows of scales on cheek.
- b. Upper jaw usually with 1 posterior canine. Color bright-blue, edges of scales brownish; fins dark-brown, with green upon external border of ventrals, which are long and pointed; forehead with a fleshy hump in adult.
- c. Tubes of lateral line considerably branched *caelestinus*
- cc. Tubes of lateral line not branched *simplex*
- bb. Upper jaw with from 3 to 6 posterior canines; jaws very convex. Color green under pectoral and along side and posterior part of body; head, anterior and upper part of back, and belly grayish-yellow; dorsal and anal brown, spotted with green along their bases; pectorals and ventrals tinted with green; caudal grayish-yellow. Size large *pleianus*

LORO:

- aa. Upper jaw without posterior canines; teeth deep blue-green. Size large.
- d. Caudal deeply notched, angles much produced in adult (fin truncate or rounded in young); body moderately elongate; depth 2.5 to 3 in length; cheek with 2.5 rows of scales, those of upper row larger than those of second, 1 scale below second row. Color olive-green, with more or less ill-defined green markings on head; lower parts more or less reddish; vertical fins brownish-orange, all edged with deep-blue *guacamaia*, 203

FIG. 68.—*Pseudoscarus guacamaia*.203. *Pseudoscarus guacamaia* (Cuvier). *Guacamaia*; *Green Parrot-fish*.

Head 2.9; depth 2.7; eye 5; snout 2.8; interorbital 3; preorbital 5.5; D. ix, 10; A. ii, 9; pectoral 1.5; ventral 1.9; caudal 1.5; scales $1\frac{1}{2}$ –25–6. No canine teeth; lower jaw included in closed mouth; lateral line interrupted under last dorsal rays, beginning again two rows lower, its pores scarcely branched; caudal slightly rounded.

Color in life: Sides mottled blue and brown or with blue bars separated by irregular brown ones; under parts white, with numerous fine dark punctulations; head with a narrow pale greenish-blue bar under eye, extending upon lower jaw and fading to white, uniting with its fellow on chin; a short preocular blue bar; teeth green; dorsal and anal brick-reddish, with narrow blue border; a few indistinct bluish spots along base and through middle of fin; base of anal with pale-blue spots; caudal dirty-red with darkish border; ventral pale-rosy, anterior edge blue; pectoral pale. In spirits: Pale-greenish, darker above; teeth greenish-blue; the green stripes on head sometimes persisting; vertical fins dark-edged.

This species is known at once by the absence of canines and the blue teeth. It ranges from Florida south to Rio Janeiro; it has been recorded from St. Augustine, Key West, Havana, and Porto Rico. Numerous specimens, all young, from 1.75 to 5.88 inches long, were seized at San Antonio Bridge, Puerto Real, Boqueron, Culebra, and Fajardo. It is probably one of the most abundant of

the large parrot-fishes about Porto Rico, as it is also about Key West. It is not uncommon at Havana, where it is still known as "guacamaia," the name by which it was known to Parra more than a century ago. It reaches a length of 2 feet or more, and is of some value as food.

Guacamaia, Parra, Descr. Dif. Piezas, Hist. Nat., 54, pl. 26, 1787, Cuba.

Scarus guacamaia Cuvier, Règne Animal, ed. II, vol. 2, 265, 1829, Cuba; after Parra.

Scarus turchesius Cuvier & Valenciennes, Hist. Nat. Poiss., XIV, 181, 1839, Porto Rico.

Scarus rostratus Poey, Memorias, II, 221, 1860, Havana.

Pseudoscarus turchesius, Poey, Fauna Puerto-Riqueña, 337, 1881; Stahl, l. c., 78 and 164, 1883.

Pseudoscarus guacamaia, Jordan & Evermann, l. c., 1657, 1898.

FAMILY LIII. EPHIPPIDÆ. The Spade-Fishes.

Body compressed, usually greatly elevated, anterior profile steep, caudal peduncle short. Scales moderate or small, ctenoid, densely covering soft parts of vertical fins; lateral line present, following curve of back. Mouth small, terminal, horizontal; premaxillaries slightly protractile; maxillary short, without supplemental bone, partly slipping under the narrow preorbital; jaws with bands of slender, pointed, movable, brush-like teeth; nostrils double; preopercle very finely serrated or entire; gill-membranes broadly attached to isthmus, openings restricted to sides; branchiostegals 6 or 7; pyloric caeca few; gillrakers very short; pseudobranchiæ present. Dorsal fins 2, somewhat connected, the first of 8 to 11 spines, which are depressible in a groove; soft dorsal and anal fins anteriorly high, their bases thickened by scales; anal spines 3 or 4, short; caudal fin truncate or doubly concave; pectoral short, rays all branched; ventrals thoracic, normally 1, 5; sometimes rudimentary; a large accessory scale as in the *Sparidae*; air-bladder large, commonly bifurcate in front, and with 2 slender horns behind. Vertebrae 10 + 14 = 24. Post-temporal bifurcate, as usual among fishes, not joined to skull.

As here understood, the *Ephippidæ* comprise about 4 genera and 10 or 12 species, related to the *Chaetodontidæ* but showing important differences in the skeleton, which shows resemblances to both scombroid and sparoid forms. Shore fishes, mostly of large size, in warm seas, often valued as food.

The following diagnosis of this family is given by Dr. Gill: "Chaetodontoidea with a wide scaly isthmus extending from pectoral region to the chin and separating the branchial apertures; spinous partially differentiated from soft portion of dorsal; upper jaw scarcely protractile; ethmoid cariniform above (not sunk and concave) and vomer declivous (not projecting forward or retuse); parapophyses spiniform and posteriorly inclosing a hæmal canal, and post-temporal bones bifurcated."

Genus 112. CHÆTODIPTERUS Lacépède.

Body much elevated and compressed, its outline nearly orbicular, anterior profile nearly vertical. Scales small, 55 to 70 in course of lateral line. Jaws about equal; no teeth on vomer or palatines; teeth on jaws slender, somewhat movable; preopercle finely serrulate. Branchiostegals 6. Dorsal fins 2, somewhat connected, first usually of 9 spines, the third of which is elongate; anal spines 3, small, second the longest; ventral with large accessory scale. Pyloric caeca 4 to 6.

An American genus distinguished from the Asiatic *Ephippus*, by the very much smaller scales.

204. *Chaetodipterus faber* (Broussonet). "*Paguala*"; *Angel-fish*; *Spade-fish*.

(PLATE 33.)

Head 3.5; depth 1.1 to 1.75; eye 4.2; snout 2.6; maxillary 3.5; interorbital 2.6; preorbital 4.7; D. VIII-1, 20 to 22; A. III, 18; pectoral 1.8; ventral 1; caudal 0.8; scales about 60. Body nearly orbicular, greatly compressed, caudal peduncle short and slender; head small, not pointed; mouth very small, the equal jaws with broad brush-like bands of slender, very closely set, movable, pointed teeth; maxillary not reaching front of eye; profile of snout nearly vertical; spinous dorsal low, the third spine much enlarged, its membrane black; fourth spine enlarged, but shorter than third, widened and flattened, the other spines short and partially embedded; soft dorsal and anal densely covered throughout with very fine scales; the anterior rays much produced, those of dorsal 1.4 in body, of anal 2.2; bases of these fins with sheaths of scales intermediate in size between those of body and those of rest of fin; pectoral small, with fine scales on outer side; ventral long, falcate, densely scaled, the first ray produced and filamentous; caudal large, scaly at base, its margin slightly concave. The young have the third dorsal spine much more produced than in adult, reaching past middle of produced rays of

soft dorsal, the black membrane continued to tip; the anterior soft dorsal rays are much less produced in the young and the body is deeper than in the adult.

Color in spirits: Pearly-gray, with dark vertical bands; one from occiput through eye to lower margin of cheek; a second from front of dorsal to belly behind base of ventrals, extending slightly upon opercle; a third from under spinous dorsal downward, tapering out below middle of side; a fourth from last dorsal spines to front of anal; a fifth from middle of base of dorsal to near end of base of anal; sixth and last at base of caudal, faint; these bands plainest in young, almost or quite disappearing in old individuals; ventrals black.

The spade-fish ranges from Cape Cod to Rio Janeiro. It is occasionally taken at New York and a few have been caught in traps at Menemsha Bight near Woods Hole in August and September. It is not uncommon about the mouth of the Chesapeake and increases in abundance southward to Key West and Pensacola, at which latter place it is called "spade-fish," while at Key West, along the east coast of Florida, and on the Carolina coast it is known as the "angel-fish," a name which, according to Schœpf, appears to have been current for over a century at Beaufort, N. C., where it is also called "porgee" or "porgy," names which are also in use for this species among the New York fishermen. In the West Indies it is known from Cuba, Santo Domingo, Jamaica, Martinique, and Porto Rico. We obtained specimens from San Juan, Mayaguez, Puerto Real, Arroyo, and Isabel Segunda, and it is doubtless common everywhere about the island in suitable places. It has also been recorded from the coasts of Guatemala and Texas. It has been erroneously reported from the Pacific coast of America, particularly from San Diego, Cal., the Pacific coast species (*Chatodipterus zonatus*) being an entirely different fish.

Mr. Silas Stearns studied the habits of the spade-fish on the coasts of west Florida, Alabama, and Louisiana, where it is common. He says it is found throughout the summer and fall in the bays, about wharves, rock piles, and old wrecks, where crustaceans are abundant. In October and November large schools are seen along the sea-beaches, evidently leaving the coast for warmer water, at which time many are caught in haul seines. In that region they probably spawn in early summer, and the young are seen until October.

The spade-fish reaches a length of 2 or 3 feet and a weight of 20 pounds, although the average size of those caught is much less. Mr. Stearns gives 15 inches as the largest he has seen, and the average at not more than 8 inches. Much larger ones were examined by us at Key West and at Fort Pierce and Eden, on Indian River. The largest seen in Porto Rico was about 14 inches long. Very large examples, which have been described as *Ephippus gigas*, but which are evidently adults of unusual size of this species, have the occipital crest and anterior interhamals developed into large, thick, bony masses, which are quite remarkable in appearance.

The spade-fish has in the last twenty-five or thirty years come to be one of the most highly prized food-fishes, and is held in very high esteem by connoisseurs in Washington and New York, in the markets of which cities it is most abundant during the summer months.

- Chatodon faber* Broussonet, Ichth. Decas., I, V, pl. 4, 1782, Jamaica; Carolina; Society Islands.
Faber marinus fere quadratus (the Pilot-fish) Sloane, Nat. Hist. Jamaica, II, 290, pl. 251, 1793, Jamaica.
Zeus quadratus Gmelin, Syst. Nat., I, 1225, 1788, Jamaica; after Sloane.
Chatodon plumieri Bloch, Ichth.; pl. 211, 1793, Martinique; after Plumier.
Scæne quadrangularis Lacépède, Hist. Nat. Poiss., IV, 564, 1803, Jamaica; after Sloane.
Chatodon oriformis Mitchell, Trans. Lit. and Phil. Soc., I, 1815, 247, pl. 5, fig. 4, New York.
Ephippus gigas Cuvier, Règne Animal, ed. 2, II, 191, 1829, America; Stahl, l. c., 163, 1883.
Ephippus faber, Poyé, Fauna Puerto-Riqueña, 330, 1881; Stahl, l. c., 77 and 163, 1883.
Chatodipterus faber, Jordan & Evermann, l. c., 1668, 1898.

Family LIV. CHÆTODONTIDÆ. The Butterfly-fishes.

Body strongly compressed, elevated, suborbicular in outline, covered with moderate-sized or small scales, which are finely ciliated or nearly smooth; lateral line present, concurrent with back, not extending on caudal fin; mouth small, protractile, terminal; maxillary very short, irregular in form, divided into two by a longitudinal suture; upper part of skull solid, occipital crest strong; post-temporal firmly joined to skull, its form really trifurcate though appearing simple, interspaces between forks filled in by bone so that only a foramen is left; last bone of suborbital ring firmly joined to preoperculum; teeth brush-like or setiform, often extremely long, in narrow bands in jaws; no teeth on vomer or palatines; no canines, molars, or incisors; eyes lateral, of moderate size; branchiostegals 6 or 7;

pseudobranchiæ very large; air-bladder present. Gill-membranes more or less attached to isthmus; gillrakers very small. Dorsal fin single, continuous, its rays sometimes filamentous, its soft part as well as soft part of anal densely covered with small scales; anal similar to soft dorsal, with 3 or 4 spines; ventrals thoracic, 1, 5; caudal usually truncate. Vertebrae 10 + 14 = 24, the anterior abbreviated; insertion of ribs inferior; post-temporal usually reduced and not bifurcate.

Carnivorous fishes of the tropical seas, noted for their singular forms, bright colors, and great activity. The family comprises 8 to 10 genera and about 180 species, most of them belonging to *Chaetodon* and *Pomacanthus*. Their excessive quickness of sense and motion enables these fishes to maintain themselves in the struggle for existence in the close competition of the coral reefs, notwithstanding their bright colors. The young are very different from the adult, and pass through a stage termed *Tholichthys*, in which the membranes are greatly developed, forming collars and sheaths about head and neck.

CHÆTODONTINÆ:

- a. Preopercle unarmed; dorsal spines not graduated, some of median spines longer than last spines; scales comparatively large (young with the *Tholichthys* form).
 b. Snout (nasal, palatines, etc.) with premaxillaries, articular, and dentary bones much produced, beak-like; cleft of mouth with maxillaries, short; lateral line ceasing under soft dorsal.
 c. Dorsal spines 12 or 13; soft rays about 20 (19 to 23).
 d. Scales large; beak moderate. PROGNATHODES
 bb. Snout little if at all produced; dorsal spines usually 12 to 14, not graduated, some of middle ones highest; anal spines 3. CHÆTODON, 113

POMACANTHINÆ:

- aa. Preopercle armed at its angle with a very strong spine, which is sometimes grooved.
 c. Interopercle unarmed; vertical limb of preopercle above spine entire or nearly so; dorsal fin with 8 to 11 spines, its soft rays 23 to 32 POMACANTHUS, 114
 cc. Interopercle short and broad, armed with 1 to 4 strong spines; preopercle serrate or spinous; dorsal spines about 14, graduated, last one longest; scales rather small; isthmus very narrow.
 f. Vertical limb of preopercle simply serrate, with 10 to 30 small teeth; body oblong, rather robust. HOLACANTHUS, 115
 ff. Vertical limb of preopercle with 3 to 9 conspicuous spines; body ovate, much compressed. ANGELICHTHYS, 116

Genus 113. CHÆTODON (Artedi) Linnæus.

Body short, deep, very strongly compressed, especially above and behind; head small, compressed, almost everywhere scaly; mouth very small, terminal, jaws provided with long, slender, flexible, bristle-like teeth; vomer sometimes with teeth; preoperculum entire or nearly so, without spine. Dorsal fin single, continuous, not notched, spinous part longer than soft part, of about 13 spines, spines not graduated, some of middle ones being longer than the last; last rays of soft dorsal usually rapidly shortened, some of them occasionally filamentous (in East Indian species); caudal peduncle short, caudal fin fan-shaped; anal similar to soft dorsal, preceded by 3 strong spines. Body covered with rather large ctenoid scales, somewhat irregular in their arrangement; lateral line curved, high, parallel with back. Gill-openings rather narrow, membranes narrowly joined to isthmus; branchiostegals 6.

A very large genus of singular and beautiful fishes, abounding in the tropical seas, especially about volcanic rocks and coral reefs; most of them have the body crossed by transverse black bars. They are all very active, feeding on small animals.

a. Scales on trunk all subequal, their posterior margins regularly rounded. None of the rays of soft dorsal produced. CHÆTODONTOPS:

- b. Series of scales below axis of body running obliquely upward and backward, lowest becoming more or less horizontal.
 c. Base of soft dorsal with a large black spot, not ocellated *ocellatus*
 d. Humeral band absent.
 c. Body with a dark band between dorsal and anal; no caudal ocellus.
 f. Ocular band edged with yellowish or whitish above; soft dorsal and anal with much black *sedentarius*
 ff. Ocular band jet-black, not white-edged; soft dorsal and anal without black *aya*
 cc. Body without black crossbands, ocular band only present; an ocellus on caudal peduncle *atenuatus*

CHÆTODON:

- bb. Series of scales below axis of body extending downward and backward, forming an angle with those above, each series marked by a continuous black streak.
 g. Body without ocelli, crossed by dark bands. *striatus*, 205
 gg. Body with a large black ocellus below soft dorsal. *capistratus*, 206
 ggg. Body with 2 ocelli, a large one on caudal peduncle and a smaller one on first 8 or 9 soft rays of dorsal. *bricci*, 207

205. *Chaetodon striatus* Linnæus. "Mariposa"; Butterfly.

(PLATE 34.)

Head 3; depth 1.6; eye 3.2; snout 2.8; interorbital 3.25; D. XII, 20 or 21; A. III, 16 or 17; pectoral 1.3; ventral 1.25; caudal 1.7; scales 8-38 to 40-16. Body ovate, much compressed, covered with large ctenoid scales, the series of which above axis of body run obliquely upward and backward, those below downward and backward; head small, pointed, with much smaller scales, anterior profile slightly concave; snout somewhat produced; mouth very small, terminal, with brush-like teeth in jaws; dorsal fin continuous, spines graduated, strong and pungent, soft rays densely covered with small scales graduated in size, becoming smaller toward margin of fin; anal similar to dorsal, both angulated behind; caudal convex.

Color in life: Ground-color very light yellow, almost white; narrow longitudinal dark stripes between rows of scales visible through the brown crossbands; a black stripe about half width of eye from occiput through eye to lower side of head, not extending on breast; a second broad band, brown in color, wider above, where it is greater than width of eye, from first dorsal spines downward across side, meeting its fellow on belly in front of vent; another band, similar to last in color and size, from last dorsal spines across body and extending upon anal fin; another on soft dorsal, yellowish-brown, narrowly joined at its top to the preceding, extending downward on soft dorsal, caudal peduncle, and posterior part of base of anal, where it is joined to ventral end of preceding bar; first three of these bars slightly curved, their concavities backward; margins of soft dorsal and anal with a narrow band of black, bordered on its inner edge with brownish-yellow, on its outer edge by a narrow pale stripe, outside of this a yellow one; this same combination of color-bands extending vertically on middle of caudal, which is pale posteriorly; a black blotch on caudal peduncle; a rounded black blotch on anterior portion of soft dorsal, sometimes absent or merging with the surrounding color, very distinct and ocellated in young of about 2 inches; iris yellow, save where crossed by the black band; yellow shades on top of head; breast dusky; membrane of second, third, and fourth dorsal spines black at base, the color extending on base of spines; membrane of longest spines yellow outwardly; pectoral pale; ventral black outwardly, light-greenish near base, a little yellow on membrane of spine.

In spirits, the delicate color shading disappears but the main features of the markings persist.

A beautiful little fish, rarely over 6 or 7 inches in length, rather common from the West Indies to Brazil; recorded by Jordan & Rutter from Jamaica and fairly abundant in Porto Rico. Numerous specimens, 1.75 to 4.5 inches long, were seized at Mayaguez, Puerto Real, Ponce, and Arroyo.

Chaetodon striatus Linnæus, Syst. Nat., ed. X, 275, 1758, Indies; Jordan & Evermann, l. c., 1677, 1898.

206. *Chaetodon capistratus* Linnæus. "Mariposa"; Parché; Butterfly.

(PLATE 35.)

Head 2.9; depth 1.7; eye 2.9; snout 2.9; interorbital 3.5; D. XII or XIII, 19 or 20; A. III, 17; pectoral 1.2; ventral 1.25; caudal 1.4; scales about 40. General form of *Chaetodon striatus*, series of scales both above and below more oblique; those above upward and backward, those below downward and backward; very distinct blue stripes between rows of scales, sometimes zigzag or undulate, sharply marking their difference in direction; a black band, edged with white, from occiput across head through eye, as in *C. striatus*; a diffuse brownish vertical bar across body from first dorsal spines to below pectoral; a round jet-black ocellated spot, larger than the eye, on body under soft dorsal, the lateral line extending upon it, surrounded by an obscure brownish shade; soft dorsal and anal and basal portion of caudal with a narrow black stripe edged inwardly with brown and outwardly with yellow; basal portion of the anal yellowish, tip of fin bright yellow; spinous dorsal pale-blue, yellow-edged; outer portion of caudal pale blue; ventrals yellow; iris, snout, breast, and belly yellow. In spirits, the blue lines become brown, and the yellow shades pale.

One of the smallest and handsomest of the chaetodonts, probably never exceeding 4 or 5 inches in length. It is found from southern Florida southward through the West Indies, and is generally common. It is recorded from Key West, Havana, Jamaica, and Porto Rico. Numerous examples, each about 2 inches long, obtained by us at San Antonio Bridge, Mayaguez, Porto Real, Boqueron, Ilucars, and Fajardo; the most abundant chaetodont in Porto Rico.

Chaetodon capistratus Linnæus, Syst. Nat., ed. X, 275, 1758, Indies; Jordan & Evermann, l. c., 1677, 1898.

207. *Chaetodon bricei* H. M. Smith. "Mariposa"; Butterfly.

Head 2.5; depth 1.7; eye 2.8; snout 3.2; interorbital 3; D. XIII, 20; A. III, 17; pectoral 1.4; ventral 1.2; caudal 1.4; scales 6-40-17. Close to *C. capistratus*, but with 2 ocelli, the larger one, on body under soft dorsal, rather elliptical in shape, the smaller one, size of eye, directly above the larger, on anterior rays of soft dorsal; the brown band from spinous dorsal across body is deeper in color and its edges are more sharply defined than in *C. capistratus*, and the broad posterior brown band of body is relatively larger and more distinct, extending from soft dorsal to anal and involving both ocelli.

One example only in the collection, 1.25 inches in length, collected at Fajardo. Hitherto known only from Woods Hole, Mass., where 6 specimens were taken in 1897 and 42 in September, 1899, by the author of the species.

Chaetodon bricei H. M. Smith, Bull. U. S. F. C. 1897, 102, Woods Hole, Mass.; Jordan & Evermann, l. c., 1678, 1898.




FIG. 69.—*Pomacanthus arcuatus*.

Genus 114. **POMACANTHUS** Lacépède. Chirivitas.

Body much compressed and elevated, covered with small scales, among which smaller ones are distributed so that the series are not distinct; preopercle with a very strong spine at its angle, vertical limb entire in adult, usually serrulate in young; interopercle entire or nearly so; dorsal fin entirely scaly, with 8 to 11 graduated spines; soft dorsal usually much elevated in front; anal with 3 graduated spines; branchiostegals 6; air-bladder with 2 posterior horns; pyloric caeca numerous.

Found in the tropical seas; few species, chiefly American; the young brilliantly colored, the adult usually dull-colored. The species vary greatly with age and have been almost inextricably confused, as the age variations are much more striking than the specific distinctions. The number of dorsal spines is usually diagnostic.

- a.* Dorsal spines VIII to X, 29 to 32.
b. Scales in lateral line about 50 to 55; dorsal VIII or IX, 30 to 32; A. III, 24. Color of adult steel-gray or scarcely yellowish; young with 4 whitish crossbands..... *arcuatus*, 208
bb. Scales in lateral line 79 to 90; dorsal usually X, 29 or 30; anal III, 23 or 24. Color black in adult, with yellow mottlings; base of pectoral yellow; young with several yellowish crossbands..... *paru*

208. Pomacanthus arcuatus (Linnaeus). "*Mariposa*"; *Black Angel*; *Chiricita*; *Portugais*.

Head 4; depth 1.4; eye 4; snout 2.1; interorbital 3.1; preorbital 2.75; D. ix, 30 to 32; A. iii, 24; pectoral 1.1; ventral 0.75; caudal 0.9; scales 8-53-27. Body nearly orbicular, greatly compressed and elevated, covered with large scales interspersed with smaller ones, not in regular series; nape, head, and breast with fine scales; dorsal and anal densely scaled to margins and tips of produced rays; a strong flat spine at angle of preopercle, head otherwise unarmed, save for a few serrulations on upper limb of preopercle; dorsal and anal with a few of their anterior rays produced, reaching beyond truncate caudal.

Color in spirits: Grayish-olive, most of scales of body very dark brown with a pale edge, the color not on scales, but on membranous expansions of the integument lining the scales; caudal with a broad, pale edge; young much darker, sometimes nearly black, with white, arcuate, vertical bars, one from nape across head behind eye to breast, another from spinous dorsal across body behind pectoral to vent, a third from soft dorsal to anal, the ends extending on fins, one or more of these bars often absent; a median white line on top of head; a pale bar from angle of mouth downward; chin dusky, with a white spot at symphysis, sometimes chin and lower jaw pale; caudal with a pale bar at base, sometimes narrowly connected along outer rays with the pale marginal bar, thus surrounding a dark area in middle of fin; posterior margin of dorsal and anal very narrowly pale-edged.

The black angel is generally common in the West Indies and is occasionally taken as far north as New Jersey and as far south as Bahia. It is known from Key West, the Tortugas, Cuba, Jamaica, Porto Rico, and Martinique. Our specimens from Porto Rico are 5 to 12 inches long and were obtained at Puerto Real, Isabel Segunda, and Culebra. It is probably not uncommon in suitable places about the island. It reaches a length of 1 to 2 feet, and is of some value as food. At Key West it is found throughout the year, and is caught chiefly in traps, though it is frequently taken with spear or hook. The average weight of those taken at Key West does not exceed 2 or 3 pounds, and the largest rarely weigh more than 6 pounds.

Chatodon arcuatus Linnaeus, Syst. Nat., ed. X, India; from spec. Mus. Ad. Fr.; D. viii, 30; dusky with 5 dark bands.

Chatodon aureus Bloch, Ichthyol., pl. 193, fig. 273, 1758, Martinique; on a drawing by Plumier, the spines 9 in the original drawing; Poey, Fauna Puerto-Riqueña, 329, 1881.

Chatodon lutescens Bonnaterre, Encycl. Méth., 182, 1788, Jamaica; after Browne.

Chatodon littoricola Poey, Synopsis, 351, 1868, Cuba; black fins bordered with yellowish; fin rays not counted; Poey, Fauna Puerto-Riqueña, 329, 1881; Stahl, l. c., 77 and 164, 1883.

Pomacanthus balteatus Cuvier & Valenciennes, Hist. Nat. Poiss., VII, 208, 1831, Porto Rico (Coll. Plécé).

Pomacanthus cingulatus Cuv. & Val., Hist. Nat. Poiss., VII, 209, 1831, West Indies; probably sent by Plécé from Porto Rico.

Pomacanthus quinquecinctus Cuvier & Valenciennes, Hist. Nat. Poiss., VII, 210, 1831, West Indies; probably from Porto Rico.

Pomacanthus arcuatus, Jordan & Evermann, l. c., 1679, 1898.

Genus 115. HOLACANTHUS Lacépède. *Catalinetas*.

Body oblong, rather robust, back not greatly elevated nor compressed; scales rather small, roughish, often mixed with smaller ones. Vertical limb of preopercle with small equal serræ; a strong spine at angle of preopercle, this usually grooved; interopercle short, armed with 1 to 4 strong spines. Dorsal fin with 12 to 15 strong spines, which are usually graduated, increasing in height to the last; soft dorsal moderate, with 17 to 20 rays, usually not ending in streamers. Coloration usually brilliant and well defined, changes due to age less than in *Pomacanthus*.

Species numerous in all tropical seas, abounding about coral reefs.

209. Holacanthus tricolor (Bloch). "*Palmoneta*"; *Rock Beauty*; *Catalineta*; *Vaqueta de dos Colores*.

(PLATE 36.)

Head 3.4; depth 2.1; eye 4.4; snout 2.2; interorbital 3; preorbital 4.7; D. xiv, 18; A. iii, 18; pectoral 1.4; ventral 1.1; caudal 1.3; scales 8-48-24. Body oblong, little elevated and not greatly compressed; scales for the most part regularly imbricated, exposed portion with many fine parallel ridges ending as very fine sharp points, making the scales extremely ctenoid; mouth small, prominent; preorbital with an antrorse and a retrorse spine varying greatly with age; a long and strong spine at angle of preopercle, the upper limb with a row of short spines; 2 spines on lower limb, and about 4 on interopercle; soft dorsal and anal with one or more rays, the seventh or eighth, or adjoining ones, produced in fine filaments, the dorsal slightly the longer, reaching end of caudal; caudal convex, its upper and lower rays produced in slender filaments, the upper the longer.

Color in life: Caudal, pectoral, and ventral fins, and the forward parts in front of a line from fourth dorsal spine to near base of pectoral, thence to third anal spine, bright golden-yellow; rest of trunk and most of dorsal and anal deep black; reddish-orange on preopercular spine, edge of opercle, and horizontal margins of dorsal and anal, the vertical margins and produced rays of which are yellow; orange punctulations on caudal; lips pale-blue, becoming black in spirits; breast dusky; iris yellow, blue above and below.

The rock beauty is one of the most gorgeously colored of tropical fishes, the rich orange of the head, tail, and anterior third of trunk, and the soft, satiny-black of the rest of the body, together with the narrow red and orange borders of dorsal and anal fins, give the fish a most striking appearance. The line of demarcation between the colors is everywhere abrupt and clean cut, and the contrast is as great as it could well be.

This interesting fish is not uncommon in the West Indies, ranging south to Bahia and north to the Bermudas. It is known from Cuba, Bermuda, Jamaica, Porto Rico, St. Thomas, Guadaloupe, Trinidad, and Bahia, but has not been recorded from southern Florida or the Bahamas. Specimens were obtained by us at Arroyo and Isabel Segunda, but it was not seen elsewhere. This species reaches a length of a foot or more, and is regarded as a good food-fish. It frequents water of moderate depths about the coral reefs and is usually taken in the fish traps or baskets.

Catalineta, Parra, Descr. Dif. Piez., Hist. Nat. Cuba, 12 pl., VII, fig. 2, 1787, Cuba.

Chatodon tricolor Bloch, Ichth., pl. 426, 1795, Cuba.

Holacanthus tricolor, Jordan & Evermann, l. c., 1684, 1898.

Genus 116. ANGELICHTHYS Jordan & Evermann. Isabelitas.

This genus is separated from *Holacanthus* by the presence on the ascending limb of the preopercle of several stout graduated spines in addition to the large grooved spine at the angle. The soft dorsal and anal are much falcate and the preorbital is without spine; interopercle armed with 1 to 4 spines; scales rather large; body ovate, rather deep, and compressed. The known species are among the largest of the chatodonts and perhaps the most gaily colored of all. Species all American.

a. Spines on ascending limb of preopercle moderate, longest less than one-fourth length of large spine at angle.

b. Nape with a blue ocellus; soft dorsal and anal edged with dark-blue; depth 1.87 in length in adult..... *ciliaris*, 210

bb. Nape without distinct ocellus; no dark-blue edgings to soft dorsal and anal; body deep, the depth 1.6 in length in adult..... *isabelita*

210. *Angelicthys ciliaris* (Linnaeus). "*Mariposa*"; *Blue Angel-fish*; *Isabelita*.

(PLATE 37.)

Head 3.8; depth 1.8; eye 4.7; snout 2.6; interorbital 3; preorbital 3.8; D. XIV, 21; A. III, 21; pectoral 1.2; ventral 1; caudal 1.1; scales 47. Body oblong and compressed; head small, obtusely pointed; scales on body of various sizes, a large series of regularly imbricated ones, with many much smaller ones between, all strongly ctenoid, those of head much smaller than the large scales of body; preopercle with a very strong grooved spine at angle, upper limb with shorter strong spines, irregular in length; 2 spines on lower limb, about 3 on interopercle and 3 on lower margin of opercle, these sometimes reduced in number or entirely absent; preorbital ending in 3 diverging spines; spines of head subject to variation in number and size with age of specimen; middle rays of soft dorsal and anal much produced, reaching beyond caudal, the fins densely scaled.

Color in life: Ground-color blue, margins of scales yellowish; sides of head pale-yellow; branchiostegal membranes, pectoral, ventral, and caudal fins lemon-yellow; dorsal and anal reddish-orange, with bright-blue border, bases of soft portions shaded with blue of body, their last rays with a dark blotch, their produced tips yellow; pectoral with an olive base bordered anteriorly by a pale-blue stripe, posteriorly by a semicircular black line; nape with a large round deep-blue spot ocellated with pale-blue and containing pale-blue specks; iris yellow, with a blue upper and lower edge; preorbital spines pale-yellow, other spines of head blue. All the bright colors fade in spirits, leaving a ground-color of olive, the lighter edges of scales standing out prominently; ocellus persistent.

The variation in the color of this species is considerable. While the blue ground-color is usually sufficiently strong to justify the name "blue angel-fish," the yellow margins of the scales are sometimes so broad as to give the fish a decidedly yellow appearance, which accounts for the name "yellow angel-fish," by which the species is known at Key West.

This is a very handsome fish and, like the rock beauty, appears very striking when seen swimming leisurely in the clear water about the coral reefs. It is known from southern Florida through the West Indies to Brazil; it has been recorded from the Tortugas, Key West, the Bahamas, Cuba, Jamaica, the Bermudas, the Lesser Antilles, and Bahia. Specimens 9 to 13 inches long were obtained by us at Ponce and Culebra, and it is doubtless a common species about Porto Rico. At Key West it is probably the most common of the 4 angel-fishes occurring there. It attains a good size (1 to 2 feet in length) and is a fair food-fish. Like all similar fishes, it is usually caught by traps, though it sometimes takes a baited hook.

Angel-fish, Catesby, Nat. Hist. Carolina, etc., 1737.

Isabelita, Parra, Dif. Piezas, etc., 1787, Cuba.

Chetodon ciliaris Linnæus, Syst. Nat., ed. X, 276, 1758, Indies; in part.

Chetodon squamulosus Shaw, Naturalists' Miscellany, 275, 1789-1813; after Angel-fish of Catesby.

Chetodon parrae Bloch & Schneider, Syst. Ichth., 235, 1801, Cuba; after Isabelita of Parra.

Holacanthus cornutus Desmarest, Décade Ichthyologique, 44, pl. 3, fig. 3, 1823, Cuba.

Holacanthus formosus Castelnau, Anim. Nouv. ou Rares de l'Amér. du Sud, Poissons, 19, pl. 2, fig. 2, 1855, Bahia.

Angelichthys ciliaris, Jordan & Evermann, l. c., 1684, 1898.

Family LV. TEUTHIDIDÆ. The Surgeon-fishes.

Body oblong, compressed, and usually elevated, covered with very small scales; lateral line continuous. Tail armed with one or more spines or bony plates. Eye lateral, high up; preorbital very narrow and deep. Nostrils double. Mouth small, low; each jaw with a single series of narrow incisor-like teeth; vomer and palatines toothless; premaxillaries somewhat movable, but not protractile; maxillary short, closely united with premaxillary; gillrakers obsolete; pseudobranchiæ large; gills 4, a slit behind fourth; gill-membranes attached to isthmus, openings thus restricted to sides. A single dorsal fin, with strong spines, spinous part of fin shorter than soft part; anal fin similar to soft dorsal; pectoral moderate; ventral fins present, thoracic, mostly 1, 5, never 1, 4, 1. Pelvic bones long, narrow, curved, closely connected, evident through skin, as in *Balistidæ*, with which group the *Teuthididæ* have the closest affinities. Pyloric caeca rather few; air-bladder large; intestinal canal long. Vertebrae $9 + 13 = 22$. Posterior suborbital bones in close contact with preopercle; post-temporal immovably united with skull, apparently simple, but really trifurcate, with interspaces filled in with bone, the foramen not passing through it; interneural bones with transversely expanded buckler-like subcutaneous plates, which intervene between spines and limit their motion forward; epipleurals developed from ribs.

Herbivorous fishes of the tropical seas.

Genus 117. TEUTHIS Linnæus. The Tangs.

This genus includes those *Teuthididæ* which have the tail armed with a sharp, antrorse, lancet-like, movable spine; strong, fixed, incisor teeth; ventral rays 1, 5, and usually 9 spines in dorsal fin. The numerous species are found in all tropical seas. Herbivorous fishes, living about coral reefs; adult protected by the murderous caudal spine, which grows larger with age. Of the six species of this genus known to occur in America north of the Equator, three are known from Porto Rico.

a. Outline rhomboid, depth 1.5 in length; color brown, washed with deep blue *cœruleus*, 211

aa. Outline ovate, depth about 2 in length; color brown, never blue.

b. Caudal simply lunate *hepatus*, 212

bb. Caudal deeply emarginate. Upper lobe of caudal produced in a filament *bahianus*, 213

211. *Teuthis cœruleus* (Bloch & Schneider). "Médico"; Barbero; Blue Tang.

(PLATE 38.)

Head 3.4; depth 1.75; eye 4.5; snout 1.3; interorbital width 3; preorbital 1.67; D. IX, 26; A. III, 25. Body rhomboid, anterior profile subvertical, nearly straight, making an angle of about 60° with axis of body; dorsal outline from origin of dorsal fin to caudal peduncle a gentle convex curve; ventral outline a regular curve from snout to caudal peduncle; mouth slightly below axis of body; least depth of caudal peduncle 2 in snout. Dorsal fin moderate, longest spines about 2.4 in head, a little shorter than longest rays; longest anal rays about 2.75 in head, longest anal spine 2; pectoral long, slightly falcate, as long as head; ventrals 1.5 in head; caudal deeply lunate, its lobes subequal, middle rays about half length of outer, which are about a sixth longer than head.

Color, rich blue throughout, body with about 45 or 50 narrow longitudinal lines of lighter blue

or purplish; no blue lines on breast or head; no crossbars; dorsal crossed by alternating lines of rich blue and light brown, the edge broadly light blue; caudal blue, a subterminal brown border, then a border of rich blue; anal like dorsal; ventral rich blue; pectoral lighter blue; young with blue shades less evident. A specimen, 3.25 inches long, had sides olivaceous-blue, rich blue on belly; side with many narrow pale-blue, almost white, cross-lines; dorsal, ventral, anal, breast, and chin rich sky-blue; lips whitish; caudal olivaceous, bordered all round by blue; spine pale-yellowish.

The blue tang is generally common from the Bermudas and southern Florida to Brazil. It has been recorded from the Tortugas, Key West, Cuba, and Jamaica. It is common about Porto Rico, numerous specimens having been obtained at Aguadilla, Arroyo, and Culebra, but it is apparently less abundant than either *T. hepatus* or *T. bahianus*. It ordinarily does not reach a greater length than 8 or 10 inches, and is usually found in the algae in water of but a few feet in depth. The larger individuals are found in deeper water about or near the coral reefs.

Turdus rhomboïdalis, Catesby, Nat. Hist. Carolina, etc., 1742.

Acanthurus caeruleus Bloch & Schneider, Syst. Ichth., 214, 1801, Carolina, Havana, and Jamaica; based on Catesby Parra, and Browne.

Acanthurus broussonetii Desmarest, Prem. Déc. Ichth., 26, 1823, Cuba.

Acanthurus brevis Poey, Memorias, II, 207, 1860, Havana.

Aconurus caruleatus Poey, Enumeratio, 69, 1875, Cuba; young.

Teuthis caeruleus, Jordan & Evermann, l. c., 1691, 1898.

212. *Teuthis hepatus* Linnaeus. "Médico"; "Barbero"; Tang; Doctor-fish; Lanct-fish.

Head 3.5; depth 2; eye 3.5; snout 1.6; interorbital 3.3; preorbital 1.8; D. ix, 24 to 26; A. iii, 22 to 24; pectoral 1.2; ventral 1.4; caudal 1.1; scales numerous. Body ovate, back greatly elevated, anterior profile very steep and moderately convex; caudal lunate, lobes about equal.

Color in life: Dark olive-brown, sides with about 12 black vertical bars narrower than interspaces, plainer in young; a brownish stripe along base of dorsal; spinous dorsal with alternate stripes running upward and backward, of dark-blue and bronze-olive, the two colors of about equal width; soft dorsal with a bluish streak on anterior side of each ray, and a bronze stripe behind it; fins dark, often almost black. A young example, taken at San Juan, was olivaceous, with narrow vertical blue bars; spine pale, surrounded by rich blue; branchiostegals, breast, and belly pale-blue; anal bluish, especially anteriorly, its border deep blue; ventrals blue; caudal brownish, peduncle yellowish; eye blue and yellow; two or three small postocular orange spots. The vertical bars usually persist in spirits, but other markings fade. In the young the caudal is nearly truncate and without the pale edge of the ocean tang.

This species is the most abundant of the tangs, and is found from the Carolinas and southern Florida south to Brazil. It has been recorded from the Tortugas, Key West, Charleston, Havana, Jamaica, Martinique, and Bahia. About Porto Rico it is the most common of the tangs and was seen by us at nearly every place where collecting was done, particularly at San Juan, Mayaguez, Puerto Real, Ensenada del Boqueron, Guanica, Hucares, and Culebra. It reaches a foot in length, though the specimens obtained do not exceed 6 or 7 inches. It is of considerable importance as a food-fish.

Teuthis hepatus Linnaeus, Syst. Nat., ed. XII, 507, 1766, Carolina; after *Hepatus mucrone reflexo*, Gronow; Jordan & Evermann, l. c., 1691, 1898.

Chaetodon chirurgus Bloch, Ansl. Fisch., 99, pl. 208, No. 24, 1784, Martinique; on a drawing by Plumier.

Acanthurus phlebotomus Cuvier & Valenciennes, Hist. Nat. Poiss., X, 176, 1835, Martinique, Brazil, Havana, and New York; Poey, Repertorio, I, 256, 1867, and Poey, Synopsis, 245, fig. 7, 1868.

Aconurus fuscus Gronow, Cat. Fishes, ed. Gray, 119, 1854, Carolina; same type as *T. hepatus* Linnaeus.

Aconurus carneus Poey, Memorias, II, 207, 1860, Cuba; young.

213. *Teuthis bahianus* (Castelnau). "Médico"; Barbero; Ocean Tang.

Head 3.5; depth 2 to 2.4; eye 3 to 4.3; snout 1.75; interorbital 3.4; preorbital 1.67; D. ix, 25; A. iii, 23. Body ovate, anterior profile moderately convex, making an angle of about 45° with axis of body; ventral outline less arched; caudal fin deeply lunate, upper lobe much the longer, slender and often produced into a filament in adult, inner rays about half length of outer ones, which are longer than head.

Color in life: Dark bluish-brown, blotched with paler below; no transverse bars; brown, wavy, longitudinal brassy streaks on body; 8 dark lines on dorsal fin running parallel with its edge for its

whole length, separated by interspaces of similar width; margin of caudal fin bluish, with a violet base; no distinct dark crossbar at base of caudal. A young individual, less than 3 inches long, had sides bluish-brown, with numerous fine, wavy, darker brown longitudinal lines; dorsal, caudal, and anal fins bluish, with darker blue border, caudal palest; pectoral pale; ventrals darker blue; caudal spine bluish.

Though not the most abundant species of the genus, the ocean tang is the most valuable of those found in our waters. It occurs throughout the West Indies and on neighboring coasts of tropical America from southern Florida to Brazil. It has been recorded from Key West, Havana, and Bahia. It is the least common of the three species found at Key West, but about Porto Rico it is the second in abundance. Specimens are in the collections from San Juan, Aguadilla, Mayaguez, Ponce, Arroyo, Hucares, Fajardo, Isabel Segunda, Culebra, and San Geronimo. Large examples were observed in greatest numbers at Culebra, where this was one of the most common fishes seen in the fishermen's boats. A pot or trap basket lifted in our presence by a Tortola fisherman contained 30 good-sized examples of this species, and many others had been taken from the other pots.

The ocean tang apparently reaches the largest size of any of the three species (a foot or more), and is by far the most important as a food-fish. It is held in high esteem by the fishermen from Tortola, St. Croix, and St. Thomas who come to the Porto Rican waters. It is usually caught in the common native trap basket or pot, which is baited with large chunks of the white pulp of the cactus and set in 4 to 10 fathoms of water. Sometimes the fish are "grained," or speared, and occasionally they are taken with hook and line. This is one of the principal fishes that are corned (so that they will keep well for about two weeks), and taken chiefly to Santa Cruz, where they bring about \$5 a barrel.

Acanthurus bahianus Castelnau, Anim. Nouv. ou Rares de l'Amér. Sud, 24, pl. 11, fig. 1, 1855, Bahia.

Acanthurus tractus Poey, Mem., II, 208, 1860, Cuba; Poey, Fauna Puerto-Riqueña, 330, 1881; Stahl, l. c., 77 and 164, 1883.

Acronurus nigriculus Poey, Enumeratio, 69, 1875, Cuba; larval form.

Teuthis bahianus, Jordan & Evermann, l. c., 1693, 1898.




FIG. 70.—*Teuthis bahianus*.

Family LVI. BALISTIDÆ. The Trigger-fishes.

Body oblong, or ovate, moderately compressed, covered with rather large rough scales or scutes of varying form, scutes not forming an immovable carapace. Lateral line obscure or wanting. Mouth small, terminal, low; jaws short, each with about one series of separate incisor-like teeth; eye near occiput; preorbital very deep. Chin without barbel. Gill-openings small, slit-like, above or in front of pectoral fins, and not before eyes. Dorsal fins 2, anterior of 2 or 3 spines, first spine highest, very strong, second locking it in erection; second dorsal remote from first, of many soft rays; caudal fin rounded or forked; ventral fins wanting, their place occupied by a single stout thick spine at end of the very long, usually movable, pubic bone. Post-temporal short, simple, forks obliterated, bone grown solidly to skull, and without foramen. Vertebrae in reduced number (17).

Shore-fishes of the tropical seas, of rather large size, carnivorous or partly herbivorous. The family contains about 9 genera and 50 species. They are very rarely used as food, many of them being reputed as poisonous. According to Dr. Day:

"Eating the flesh of these fishes occasions in places symptoms of most virulent poisoning. Dr. Memier, at the Mauritius, considers that the poisonous flesh acts primarily on the nervous tissue of the stomach, occasioning violent spasms of that organ and shortly afterwards of all the muscles of the body. The frame becomes racked with spasms, the tongue thickened, the eye fixed, the breathing laborious, and the patient expires in a paroxysm of extreme suffering."

- a. Teeth unequal, oblique, each one deeply notched.
 - b. Gill-opening with a number of enlarged bony plates or scutes behind it; ventral flap movable, supported by a series of spines, more or less free at tip, and resembling fin rays; cheek entirely scaled, without naked grooves or patches; eye with a groove before it; scales rather small, 50 to 75.
 - c. Dorsal and anal fins falcate in adult; caudal lobes acuminate in adult; lateral line slender, undulate, more or less developed; scales of tail and posterior parts unarmed, similar to those on rest of body; ventral flap with slender, sharp spines; third dorsal spine little smaller than second and remote from it..... BALISTES, 118
 - bb. Gill-opening with only ordinary scales behind it; no enlarged plates or scutes; ventral flap scarcely movable, its surface scaled; lateral line obsolete; third dorsal spine small or wanting; vertical fins in adult more or less angulate or falcate.
 - d. Chin not projecting; cheek closely scaled; dorsal spines 3; scales of posterior parts unarmed or keeled.
- CANTHIDERMIS
- dd. Chin much projecting; cheek with 3 to 5 narrow parallel grooves; dorsal spines 2; scales of posterior parts more or less keeled..... XANTHICHTHYS
 - aa. Teeth even, incisor-like; scales of posterior parts more or less keeled; a groove before eye, enlarged scutes behind it; lateral line obsolete; third dorsal spine small or wanting; cheek entirely scaled, but marked by narrow grooves; enlarged scales present behind gill-opening; ventral flap scarcely movable, its surface scaled; vertical fins more or less angulated..... MELICHTHYS

Genus 118. BALISTES (Artedi) Linnæus. Trigger-fishes.

Body compressed, covered with thick, rough scales or plates of moderate size, 50 to 75 in a lengthwise series; a naked groove before eye below nostrils; lateral line more or less developed, very slender, undulate, conspicuous only when scales are dry, extending on cheek. Pelvic flap large, movable, supported by a series of slender, pungent spines. Caudal peduncle compressed, its scales unarmed, without spines or differentiated tubercles similar to those on rest of body. Gill-opening with enlarged bony scutes behind it; cheek entirely scaly, without naked patches or grooves. Both jaws with irregular, incisor-like teeth, usually 4 on each side in each jaw. First dorsal of 3 spines, anterior of which is much the largest, second acting as a trigger, locking first when erected; third nearly as large as second and remote from it; second dorsal and anal long, similar to each other, in adult always falcate or filamentous in front; caudal fin rounded, with outer rays much produced in adult; branchiostegals 6; vertebrae 7 + 10.

Species rather few, chiefly American; some of them straying to the Old World.

CAPRISCUS:

- a. Lateral line complete, beginning on lower part of cheek, thence extending upward to behind eye, thence backward to beyond first dorsal, thence abruptly downward to above anal, then upward and at last horizontally backward on caudal peduncle, line everywhere much undulated; lines of the two sides connected by a cross line at nape; dorsal fin falcate or filamentous; dorsal rays about III, 27; A. 25.
- b. Scales moderate, about 60 (50 to 65) in a lengthwise series.
- c. Body with few blue spots or none.
- d. Dorsal and anal with oblique dark bands of bluish spots; young clouded, and with vague, dusky blotches at base of dorsal; scales about 60..... *carolinensis*
- cc. Body covered with roundish blue-black spots; dorsal and anal similarly spotted..... *forcipatus*

BALISTES:

- aa. Lateral line incomplete, usually developed only on head, nape, and caudal peduncle; 37 scales between origin of dorsal and vent..... *vetula*, 214

214. Balistes vetula Linnæus. *Old Wife; Old Wench; Cochino; "Peje Puerco."*

(PLATE 39.)

Head 3; depth 1.8; D. III, 29; A. 27; scales 63. Median part of lateral line, from base of first dorsal to front of caudal peduncle wanting in adult, branch on cheek ceasing opposite gill-opening; cross-branch present; ventral flap well developed, with slender, sharp spines. Scales on head much smaller and more crowded than those on body; third dorsal spine rather shorter and weaker than second,

remote from it; caudal fin widely forked, lobes filamentous and about equal; dorsal in adult filamentous at tip; anal little elevated anteriorly.

Color in life: Olivaceous, yellowish below, especially on breast; lips green, bordered behind by a narrow blue line; a narrow blue line narrowly bordered by yellow, from blue line surrounding mouth backward and curving downward across lower side of cheek toward throat; a similar but broader band from above snout across cheek and parallel with first to below base of pectoral; a narrow blue line at base of pectoral; about 8 or 10 narrow, black, wavy lines very narrowly bordered with yellowish-green, radiating from eye; below eye a similar black line extending in an upward curve from above snout to base of pectoral; soft dorsal greenish at base, somewhat purplish above, with 2 or 3 irregular, broken, narrow blue lines near border, these sometimes being more numerous; anterior part of soft dorsal with a narrow blue line extending from base of first ray upward to height of about tenth ray; anal olivaceous with a broad subterminal blue band anteriorly convex and bordered by yellowish; tip of fin and prolonged external rays faint-purplish; caudal peduncle with a broad blue ring anteriorly followed by two narrow pale-blue lines, last at base of caudal fin; caudal greenish-olive, a narrow blue line following direction of first ray for nearly its entire length, and a broader blue subterminal band extending across fin from first to last ray; spinous dorsal purplish and greenish, anterior side of first spine blue, second and third spines pale; iris with an orange ring and radiating lines of white and blue. In alcohol most of these colors disappear, the broad blue lines on side of head changing to blackish.

This interesting fish is found throughout the West Indies and occasionally northward in the Gulf Stream to Woods Hole, Mass.; not uncommon at Key West; recorded by Jordan & Rutter from Jamaica, where it is called "Bessy Cerka"; recorded also from the Bahamas and Ascension Island; probably not uncommon about Porto Rico; the collection contains four excellent specimens from Arroyo, where other individuals were seen in the possession of local fishermen.

Guaperva Maregrave, Hist. Bras., 163, 1648, Brazil.

Turdus oculo radiato (the Old Wife), Catesby, Hist. Carol., pl. 22, 1725, Bahamas.

Balistes vetula Linnæus, Syst. Nat., ed. X, 329, 1758, Ascension Island; after *Balistes vetula* of Osbeck, Iter Chin ensis 294, 1757.

Balistes bellus Walbaum, Artedi Piscium, III, 467, 1792, West Indies; after Froyer.

Chaliosma velata Swainson, Class. Fishes, II, 325, 1839; probably a misprint for *vetula*.

Balistes equestris Gronow, Cat. Fishes, ed. Gray, 31, 1854, American seas.

Balistes vetula, Jordan & Evermann, l. c., 1703, 1898.

Family LVII. MONACANTHIDÆ. The File-fishes.

Body much compressed, covered with very small rough scales, forming a velvety covering; males sometimes with spines on caudal peduncle. Upper jaw with a double series of incisor-like teeth, 6 in outer and 4 in inner series; lower jaw with 6 similar teeth in a single series; first dorsal with a single strong spine and generally a rudimentary one behind it; second dorsal long, similar to anal; ventral fins reduced to a simple osseous, fixed or movable, small appendage at end of long pelvic bone; this appendage often rudimentary or entirely absent; no barbel; vertebrae 7+11 to 14=18 to 21.

Herbivorous shore-fishes of warm seas, closely allied to the *Balistida*, differing chiefly in having the first dorsal represented by a single spine, behind which is sometimes a rudiment; scales small, spinigerous, skin mostly rough-velvety. The family contains 6 or more genera and about 50 species, mostly small in size and not used as food, having little flesh and that of a bitterish taste.

a. Pubic bone with a small spine at its end; gill-opening short, nearly vertical; dorsal and anal moderate, each of fewer than 40 rays.

b. Pelvic spine fixed.

bb. Pelvic spine movable.

c. Dorsal spine armed with strong retrorse barbs, usually in two series..... MONACANTHUS, 120

d. Dorsal spine with about 4 series of small barbs..... PSEUDOMONACANTHUS

dd. Dorsal spine without barbs..... CANTHERINES, 119

aa. Pubic bone without spine at its end; gill-opening long, oblique; dorsal and anal long, each of 40 or more rays; dorsal spine without barbs, inserted above orbit..... ALUTERA, 121

Genus 119. CANTHERINES Swainson.

This genus differs from *Monacanthus* chiefly in the absence of barbs on the dorsal spine, which is long, strong, and placed over the front of the eye. Scales minute. Species few.

215. *Cantherines pullus* (Ranzani). *Lija Colorado*; "Peje Puerco."

Head 3.3; depth about 2; D. II, 35; A. 31. Body moderately elevated; snout moderately produced, upper profile slightly concave; posterior margin of eye directly above axil. Adults (12 inches long) with 2 to 6 pairs of strong recurved spines on each side of tail; caudal short; dorsal spine nearly straight, rather shorter than head, without barbs, serrulate in front, situated above front of eye; skin with a velvety appearance; scales minute.

Coloration variable, generally with a whitish spot behind the last dorsal ray, and several more or less distinct pale longitudinal bands along tail; head with undulated bluish streaks; body sometimes with scattered round light spots, each with a dark speck in center; young sometimes uniform silvery; color probably varying with surroundings. Specimens 7 inches long were, in life, dirty-brown, body sparsely covered with small reddish-brown spots; side of head with narrow brown lines and brown spots; dorsal and anal pale; caudal brown, crossed by a pale bar near middle, which is followed by a dark bar; tip of tail with a green border. In alcohol, the body and head become grayish or blackish; an irregular black blotch behind gill-opening; fins all pale, traces of bars on caudal remaining. In the young there are no spines on tail.

From the West Indies to coast of Brazil, and occasionally north to Florida. It has been recorded from the Florida Keys and the Tortugas; also from Cuba and Bahia. Probably not abundant in Porto Rico; two obtained at Arroyo, each 7 inches long; others seen. It reaches a weight of 6 pounds.

Lija colorada, Parra, Dif. Piczas, etc., pl. 23, 1787, Cuba.

Monacanthus pullus Ranzani, Nov. Comm. Act. Sci. Inst. Bonon., V, 4, pl. 1, 1842, Brazil.

Monacanthus pardalis, Günther, Cat., VIII, 230, 1870; in part, probably not of Rüppell.

Monacanthus ruppellii Castelnau, Anim. Nouv. Amér. Sud, Poiss., 97, pl. 47, fig. 2, Bahia.

Monacanthus macrocerus Hollard, Ann. Sci. Nat. 1854, 4th series, II, 327, pl. 12, fig. 1, 1854; Bahia, adult.

Monacanthus irroratus Poey, Memorias, II, 330, 1861, Cuba.

Monacanthus stratus Poey, Memorias, II, 329, 1861, Cuba.

Monacanthus parvaianus Poey, Proc. Ac. Nat. Sci. Phila. 1863, 185, Cuba; after *Lija Colorado* of Parra.

Monacanthus punctatus Poey, Synopsis, 437, 1868, Cuba.

Cantherines pullus, Jordan & Evermann, l. c., 1713, 1898.

Genus 120. MONACANTHUS Cuvier.

Body short and deep, very strongly compressed, covered with minute, rough scales. Mouth very small; upper jaw with a double series of incisor-like teeth, usually 6 in outer and 4 in inner series; lower jaw with about 6 incisors in a single series; teeth connivent, unequal; gill-opening a small slit, shorter than eye, nearly vertical below posterior part of eye and just in front of upper edge of pectoral. Dorsal spine large, armed with two series of retrorse barbs, no conspicuous filaments; second dorsal and anal fins similar to each other, of about 25 to 35 rays each; caudal fin moderate, rounded; pelvic bone with a blunt, movable spine, bone connected by a movable flap of varying size; side of tail often with a patch of spines, especially in males. Vertebrae 7 + 11 to 14 = 18 to 21.

Species very numerous, in warm seas; most of them small, lean fishes with leathery skin and bitter flesh, unsuited for food.

MONACANTHUS:

a. Ventral flap in adult greatly developed, extending much beyond ventral spine; adult with 2 or 3 pairs of recurved spines on caudal peduncle; young without these characters, similar to young of *Stephanolepis*.

b. D. I, 30; A. 30. Color very variable. *ciliatus*, 216

STEPHANOLEPIS:

aa. Ventral flap, even in adult, moderately developed, not reaching beyond pelvic spine; no recurved spines on caudal peduncle.

c. Dorsal and anal each with 30 to 32 soft rays.

d. Depth more than half length of body *hispidus*, 217

dd. Depth less than half length of body *spilonotus*

cc. Dorsal and anal each with about 27 soft rays. *oppositus*

216. *Monacanthus ciliatus* (Mitchill). *Leather-fish*; *Lija*; "Pez de Puerco."

Head 3.5; depth 1.75; young 1.5; D. I, 30; A. 30; scales very small, without median crest. Spines becoming longer on caudal peduncle, which has in addition 2 or 3 pairs of strong spines curved forward, these prominent only in adults; ventral flap longer than head, about one-third length of body. Scales on ventral flap developed as flat plates, with their free margins pectinate. Snout pointed, upper profile concave. Dorsal spine strong, nearly as long as head, armed behind with 2 rows of retrorse barbs; ventral spine small, rough.

Color varying very much with the surroundings of the fish, from dull olive-gray to the most vivid grass-green; markings not well defined and not very constant; green, with white cirri on sides; a whitish longitudinal cloud behind pectoral; a pale band downward and forward from eye; lower side of head with darker crossbands; dorsal and anal pinkish, with (usually 3) darker spots at base; ventral flap edged with scarlet; caudal greenish, mottled with darker and pale; some specimens show neither red nor green shades, and have vague, dusky, longitudinal stripes. A specimen 3 inches long, from Isabel Segunda, Vieques Island, in alcohol, is brownish-gray, dusky, with a large black clouded blotch from front of soft dorsal downward and forward to middle of side; body irregularly clouded with dusky elsewhere; top of head brownish-gray; below very dusky, becoming black on flap, which has a light submarginal stripe followed by a narrow white border; dorsal and anal translucent; a faint dark blotch below middle of soft dorsal, and one at front of anal; caudal barred with white and dark.

This species is known from the West Indies and southern Florida, and is very abundant about the Florida Keys, along with *M. hispidus*. It has been recorded also from Bahamas, Puerto Cabello (Cuba), Jamaica, and Porto Rico, where it is not uncommon. Our collection contains specimens from San Geronimo, Boqueron, Guanica, and Isabel Segunda, off Vieques Island, where a specimen was dredged in 16 fathoms at station 6092, and off St. Thomas, at station 6079, in 20 to 23 fathoms. Length




FIG. 71.—*Monacanthus ciliatus*.

3 to 8 inches. The young of this species and of *M. hispidus* are much alike, but *M. ciliatus* is always more elongate, and as it grows older the pelvic flap grows much larger and the armature of the tail more distinct.

Balistes ciliatus Mitchell, Amer. Monthly Mag. and Crit. Rev., March, 1818, 326, Bahama Straits.

Monacanthus occidentalis Günther, Cat., VIII, 237, 1870, Puerto Cabello, Cuba.

Monacanthus davidsoni Cope, Trans. Amer. Phil. Soc. Phila., XIV, 1870, 476, Florida Reef.

Monacanthus ciliatus, Jordan & Evermann, L. c., 1714, 1898.

217. *Monacanthus hispidus* (Linnaeus). Fool-fish; File-fish; Leather-fish; Horny Cong; Lija.

Head 3.4; depth 1.75; D. 1, 32; A. 32. Young slightly deeper (1.5) proportionally than adult. Body rather deep. Jaws subequal; eye large, about 3 in snout. Gill-opening about as long as eye, separated from eye by an interspace nearly equal to its length. Anterior profile slightly concave. Dorsal spine somewhat shorter than snout, more than half head, inserted above posterior part of eye, stout, rough, armed behind with 2 rows of retrorse barbs; first ray of soft dorsal often filamentous in adult (male?), its length varying from that of snout to that of depth of body (longest among specimens seen by us is one from Canary Islands); pectoral small. Pelvic bone long, ending in a short, blunt, movable spine, beyond which the abdominal flap does not extend. Scales minute, each with a crest of about 3 prickles, those on caudal peduncle villous, those on ventral flap larger, elongate; no naked areas; no recurved spines on tail. Length 10 inches.

Color, grass-green or olive; back and sides with faint, irregular whitish spots; head plain; spinous dorsal and caudal green; second dorsal and anal translucent; adult less variegated; dull olivaceous, mottled with dusky.

This species ranges from Cape Cod to Cuba, and is abundant on our South Atlantic coast and among the Florida Keys, and southward, through the West Indies to Brazil. It is recorded by Jordan & Rutter from Jamaica, where it is called "mingo"; known from Key West, the Pensacola snapper-grounds, Big Gasparilla, and the Anclote Sponge Kraals; apparently not common in Porto Rico, as the collection contains but a single small specimen seined at Boqueron and five larger ones from Puerto Real.

Balistes hispidus Linnæus, Syst. Nat., ed. XII, 405, 1766, Carolina.

Balistes broccus Mitchill, Trans. Lit. and Phil. Soc., I, 1815, 467, New York.

Monacanthus filamentosus Valenciennes, Îles Canaries, 95, 1836, Canaries; adult.

Monacanthus gallinula Valenciennes, l. c., 95, Canaries; young.

Monacanthus varius Ranzani, Nov. Comm. Bonon., V, 6, 1842, Brazil.

Monacanthus massachusettsensis De Kay, N. Y. Fauna: Fishes, 337, pl. 57, fig. 187, 1842, Massachusetts Bay.

Monacanthus setifer De Kay, N. Y. Fauna: Fishes, 337, pl. 59, fig. 194, 1842, New York Harbor; probably not of Bennett.

Monacanthus signifer Storer, Synopsis Fishes N. A., 497, 1846, Massachusetts; substitute for *setifer*, preoccupied.

Monacanthus hispidus, Jordan & Evermann, l. c., 1715, 1898.




FIG. 72.—*Monacanthus hispidus*.

Genus 121. ALUTERA Cuvier. File-fishes.

Body oblong or rather elongate, strongly compressed, covered with minute rough scales. Mouth and teeth essentially as in *Monacanthus*, but lower jaw more projecting, so that the lower teeth are directed obliquely upward and backward. Gill-opening an oblique slit, longer than eye, situated below and in advance of eye, its posterior end behind base of pectoral. Pelvic bone long, falcate, movable under skin, without spine at its extremity. Dorsal spine small, inserted over eye, rough, but without barbs; soft dorsal and anal long, each of 36 to 50 rays; caudal fin convex; pectoral small. Species numerous.

CERATACANTHUS:

a. Dorsal rays 1, 36.

b. Anal rays 38; coloration nearly uniform *schepffi*

bb. Anal rays 38; body covered with small, round dark-brown spots *punctata*

aa. Dorsal rays about 1, 46; anal rays about 50.

OSBECKIA:

c. Caudal fin elongate, with rounded angles. Coloration not uniform, head and body with irregular blue spots and lines, besides small round black spots; upper profile of snout concave *scripta*, 218

ALUTERA:

cc. Caudal fin short, subtruncate, with acute angles. Coloration uniform; upper profile of snout convex.... *monoceros*

218. *Alutera scripta* (Osbeck). Unicorn-fish; *Lija Trompa*.

Depth 3 to 3.25; D. 1, 44 to 48; A. 47 to 52; vertebrae 7+14. Body oblong, its depth being nearly equal to distance of hind margin of orbit from extremity of snout; snout produced, with upper profile concave; dorsal spine long and slender, above middle of orbit, about 1.25 in head; middle of gill-opening in advance of middle of eye; pectoral fin below posterior part of eye; caudal fin elongate, nearly as long as, or longer than, head, rounded; dorsal and anal fins low; ventral spine none. Head and body olivaceous, with irregular light-blue spots and curved streaks; besides these, numerous round black spots about as large as pupil; dorsal and anal yellowish; caudal reddish; skin finely velvety.

An inhabitant of tropical seas. Common in the West Indies, occasionally northward to South Carolina; also occasionally taken about the islands off the west coast of Mexico. Not seen by us in Porto Rico, but recorded from there by Professor Poey. Length 2 to 3 feet.

Unicornia pisces bahamensis (the Unicorn-fish) Catesby, Hist. Nat. Carolina, etc., II, pl. 19, 1737, Bahamas.

Balistes scriptus Osbeck, Iter Chin., I, 144, 1757, China.

Lija trompa Parra, Dif. Piezas Hist. Nat., 46, pl. 22, fig. 1.

Balistes laevis Bloch, Ichthyol., IX, 82, pl. 414, 1795, Morocco; Tranquebar.

Balistes ornatus Marion de Procé, Bull. Soc. Philom., 131, 1822.

Aluteres pareva Lesson, Voy. Coquille, Zool., 106, 1828.

Monacanthus proboscideus Ranzani, Nov. Comm. Ac. Sc. Inst. Bonon., 1842, 8, Brazil.

Aluteres venosus Hollard, Ann. Sc. Nat., IV, 1855, 14, pl. 1, fig. 3.

Alutera picturata Poey, Proc. Ac. Nat. Sci. Phila. 1863, 183, Cuba.

Alutera scripta, Poey, Fauna Puerto-Riqueña, 345, 1881; Jordan & Evermann, l. c., 1719, 1898.




FIG. 73.—*Alutera scripta*.

Family LVIII. OSTRACIIDÆ. The Trunk-fishes.

Body short, cuboid, triquetrous or pentagonal, covered by a carapace formed of firmly united polygonal bony patches; jaws, bases of fins, and caudal peduncle free and covered by smooth skin. Mouth small; each jaw with a single series of long, narrow teeth. Maxillaries and premaxillaries firmly united. Gill-opening a nearly vertical slit, below and behind eye. Dorsal fin single, short, without spine; anal short, similar to dorsal; caudal rounded; no ventral fins; vertebrae 14, the anterior 9 elongate, the last 5 extremely short; no ribs.

The *Ostraciidæ* comprise 3 genera and about 20 species, all of the tropical seas, living near bottom in shallow waters. The species are so singular in appearance and so easily preserved that they have been common in collections ever since the collecting of tropical curiosities began. The 4 American species were well known to Artdi and Linnæus. Goode says of these fishes:

“The locomotion of the trunk-fishes is very peculiar. The propelling force is exerted by the dorsal and anal fins, which have a half-rotary, sculling motion, resembling that of a screw propeller; the caudal fin acts as a rudder, save when it is needed for unusually rapid swimming, when it is used as in other fishes; the chief function of the broad pectorals seems to be that of forming a current of water through the gills, thus aiding respiration, which would otherwise be difficult on account of the narrowness and inflexibility of the branchial apertures. When taken from the water one of these fishes will live for two or three hours, all the time solemnly fanning its gills, and when restored to its native element seems none the worse for its experience, except that, on account of the air absorbed, it can not at once sink to the bottom.”

Genus 122. **LACTOPHRYS** Swainson. Three-angled Trunk-fishes.

Trunk-fishes with carapace 3-angled, ventral surface flat or concave, never carinate; carapace closed behind anal fin; carapace with or without frontal and abdominal spines; dorsal rays 9 or 10; caudal rays always 10. This genus contains 5 species, 4 of them American, and differs from the Old World genus *Ostracion* only in form of carapace. The median dorsal ridge of carapace is much more developed than the others, so that the body is 3-sided and 3-angled, instead of 4-sided and 4-angled, as in *Ostracion*. Although this character is a striking one, it is not one of high structural importance. Holland and Bleeker have discarded it as being of no real systematic value. All writers agree that the species of the group are most closely related, and that the relations of the species are closer than they appear. We think, with Dr. Goode, that the shape of the carapace affords "the most reliable guide in the arrangement of the species of the genus," and we think it not improper to accord generic distinction to the 3-angled species, as distinct from the more specialized 4-angled forms.

RHINESOMUS:

- a. Carapace without spines anywhere *triqueter*, 219
 aa. Carapace with distinct spines, at least on the ventral ridges behind.
 b. Frontal spines none.

CHAPINUS:


- c. Carapace closed behind dorsal fin; body everywhere with round, dark spots..... *bicaudalis*, 220

LACTOPHRYS:

- cc. Carapace open behind dorsal fin; body mottled with paler *trigonus*, 221

ACANTHOSTRACION:

- bb. Frontal region with 2 strong spines like horns..... *tricornis*, 222

FIG. 74.—*Lactophrys triqueter*.**219. *Lactophrys triqueter*** (Linnaeus). "*Chapin*"; Trunk-fish.

Head 3.5; height of side about 2; greatest ventral width 2.6; eye 3.4; snout 1.4; interorbital 1.7; least depth of caudal peduncle 4.1, its width in its depth 2.5; D. 10, its base 3.7 in head, the height 1.6; A. 10, its base equal to that of dorsal, its height a little less; P. 11 or 12, 1.6 in head. Body sharply 3-angled, sides rather double concave; dorsal carina strongly arched, more so than in *L. bicaudalis*, ending anteriorly above eye; ventral angles more convex and flaring than in *L. bicaudalis*; supraoccipital ridge less strong, continued backward along side in a low, flat ridge; profile from snout to eye somewhat concave; interorbital space concave; ventral surface slightly convex anteriorly, posteriorly with two low ridges, the surface between them and between each and the ventral angle shallowly concave; scales of sides hexagonal, in young with striae radiating from center to angles of each scale, in adults armed simply with tubercles, 9 or 10 plates in horizontal series from gill-opening to tail, 8 in median line of ventral surface, 8 between ventral keel and back; posterior dorsal scute unarmed; carapace closed behind dorsal and anal, width behind dorsal about 1.75 that behind anal; carapace without spines anywhere; branchial aperture oblique, its length about 2.5 in head, greater than eye; dorsal, anal, and caudal obtusely rounded; pectoral triangular, first rays longest.

Color in life: Body dirty white, upper third with small round white spots and some narrow black bars promiscuously arranged; head whiter and with whiter stellate spots; ventral angles with some

white spots; ventral surface plain pale; opercular opening bordered in front by brown, then black, and behind by black; base of pectoral and dorsal black, the fin pale yellow; caudal peduncle black with many round white spots; base of caudal similarly marked; caudal dirty white with numerous large irregular black spots. In spirits dark brown, thickly covered with circular spots of yellowish-white; ventral surface lighter and without spots; lips, bases of fins, border of branchial opening, and caudal peduncle black, or brown like ground-color of body; caudal peduncle with numerous small round white spots; caudal dusky, the tip black; other fins plain.

Found among the West Indies north to the Bermudas, Key West, and Pensacola; generally very common in the Tropics; common about Porto Rico, the collection containing specimens from Arroyo, Isabel Segunda, Culebra, and Puerto Real. A sluggish fish, reaching a foot in length, living on the bottom about reefs and among algae, feeding on minute animals of various kinds.

Ostracion polyodon inermis triquetus Linnaeus, *Adolphi-Frederici*, I, 60, 1754, India.

Ostracion triquetus Linnaeus, *Syst. Nat.*, ed. X, 330, 1758, India; after Mus. Ad. Fr.

Ostracion concatenatus Bloch, *Ichthyol.*, pl. 131, 1785, Martinique; on a painting by Plumier.

Lactophrys triquetus, Jordan & Evermann I. c., 1722, 1898.




FIG. 75.—*Lactophrys trigonus*.

220. *Lactophrys trigonus* (Linnaeus). "Chapin"; Common Trunk-fish.

Head 2.9; height of side 1.7; greatest ventral width 1.75; width between ventral spines 3.5; eye 2.8; snout 1.3; interorbital 1.8; D, 10, its base 4.25 in head, 1.5 in eye or 2 in height of fin; A, 10, its base and height about equal to dorsal; P, 2 in head. Body 3-angled, ventral angles strongly convex and strongly flaring; dorsal carina strongly and regularly arched, ending anteriorly above posterior border of orbit; supraoccipital ridge strong; interorbital space concave; profile from snout to eye nearly straight, a slight angle; carapace open behind dorsal, closed behind anal; caudal peduncle slender, its length 1.6 in head, its least depth 2.9 in its length, its least width 2 in its depth. Scales of sides hexagonal, covered with small tubercles.

Color, olive-gray, a faint blue spot in center of each of most of the scales; nostril in a yellow spot; boundaries of upper scutes blackish, of lower bluish; scutes behind gill-opening black, surrounding a white center, forming a large black blotch; a similar blotch on side between eye and base of caudal peduncle; caudal peduncle pale-olivaceous, with a few obscure white spots; fins all pale except caudal, which is somewhat dusky, especially at tip.

An inhabitant of the West Indies, north to the Bermudas and Florida, occasionally in the Gulf Stream to Woods Hole and Chesapeake Bay; common on the coast of Florida; recorded by Jordan & Rutter from Jamaica; apparently not common in Porto Rico, only one specimen, 3 inches long, having been obtained, at San Antonio Bridge. It reaches a length of about a foot.




FIG. 76.—*Lactophrys trigonus*; front view.

Ostracion triangulatus lenibis figurararum hexagonarum eminentibus, etc., Artedi, Genera, 56, 1738, Jamaica; seen by Artedi in the collection of Sir Hans Sloane and in the Naggs Head Inn, London.

Ostracion trigonus Linnaeus, Syst. Nat., ed. X, 330, 1758, India; after Artedi.

Ostracion yulei Storer, Bost. Journ. Nat. Hist., 1, 1837, 353, Holmes Hole, on Marthas Vineyard.

Lactophrys oviceps Kaup, Archiv Naturg. 1855, 218; specimens with 10 dorsal rays, Linnaeus having given by error "D. 14" in the original description of *O. trigonus*.

Ostracion undulatus Poey, Synopsis, 441, 1868, Havana.

Ostracion expansum Cope, Trans. Am. Phil. Soc. 1870, 474, figs. 9-10, St. Martins, West Indies.

Lactophrys trigonus, Poey, Fauna Puerto-Riqueña, 347, 1881; Stahl, l. c., 81 and 167, 1883; Jordan & Evermann, l. c., 1723, 1898.

221. *Lactophrys bicaudalis* (Linnaeus). "Chapin"; Spotted Trunk-fish.

(PLATE 40.)

Head 3.5; height of side 2.25; greatest ventral width 2.7; width between bases of ventral spines 3.6; eye 3; snout 1.3; interorbital 2.1; least depth of caudal peduncle 3.4; its width in its depth 3.2; D. 10, base of fin 4 in head, the height 2; A. 10, its base and height equal to dorsal; P. 12, 1.8 in head. Body sharply 3-angled, sides and ventral surface posteriorly somewhat concave; dorsal carina sharp, considerably arched, ending in front over middle of eye; supraocular ridges strong, a slight ridge from supraocular ridge along side to caudal peduncle; no spines on head; a sharp, stiff spine, 3.4 in head, on each side at posterior angle of ventral surface; carapace closed behind dorsal and anal fins, its width behind dorsal about twice that of strip behind anal; caudal peduncle about 1.4 to 1.6 in head; branchial aperture oblique, short, about 3.6 in head; body and head everywhere rough, tuberculate, tubercles strongest on body; hexagonal plates best defined on ventral surface.

Ground-color, light-yellowish, changing to whitish below; everywhere on head, body, caudal peduncle, and caudal covered with small round black spots, largest on middle of side, where they are somewhat smaller than pupil, those on the caudal peduncle small, larger on caudal fin, arranged in 5 to 7 irregular vertical bars; tip of snout rose, with a few small black spots above fleshy base of dorsal; pectoral with several black spots, few or none upon anal; pupil black, with orange border. In spirits. 3 to 5 irregular white blotches in center of hexagonal plates on low ridge behind eye.

An inhabitant of the West Indies, generally common from Cuba to Ascension Island; not known from Florida; recorded by Jordan & Rutter from Jamaica; next to *L. trigonus* the least common species of the genus about Porto Rico, specimens having been obtained only at Ensenada del Boqueron, Ponce, and Arroyo. It reaches a length of a foot or more.

Ostracion triangulatus tuberculatus hexagonis radiatis, etc., Artedi, Genera, 57, 1738, India.

Ostracion bicaudalis Linnaeus, Syst. Nat., X., 330, 1758, India.

Lactophrys bicaudalis, Jordan & Evermann, l. c., 1723, 1898.

222. *Lactophrys tricornis* (Linnaeus). Toro; Cow-fish.

Head 4.25; height of side 2.5; greatest ventral width 4.8; width between ventral spines 8.3; eye 2.7; snout 1.25; interorbital 2; D. 10, base equal to eye, height nearly equal to head; anal base 1.25 in eye, height 2 in head, caudal long, equal to distance from tip of snout to posterior edge of pectoral base; pectoral 1.2 in head; caudal peduncle long and slender, length a little exceeding that of head, least depth 2.75 in its length, least width about 3 in its least depth. Body 3-angled; dorsal carina gently arched, beginning over posterior part of pupil; ventral angles not flaring nor wide apart, each ending posteriorly in a stout spine; supraoccipital ridge moderate, with a stout spine in front; middle of side with a broad low ridge from eye to caudal peduncle; snout projecting, outline from snout to occipital spines concave below, then rising abruptly; interorbital very convex; gill-opening oblique, short, scarcely equal to eye; carapace closed behind dorsal and anal fins, the width of part behind dorsal about four times that behind anal, plate behind dorsal sometimes ending in a short spine, directed upward; hexagonal plates with small, blunt tubercles. Occasional specimens have a real spine on supracaudal plate, as figured by Lister. Of nine Porto Rican specimens examined by us two have this spine.

Color, brown, yellow, blue, or green, with irregular blue blotches, centers of scutes often lighter than margins. Young light gray, tinged with olive; belly white; head and carapace with round spots of light-blue, these sometimes forming more or less interrupted longitudinal stripes; about four of these

stripes on cheek; tail above with blue, brown-edged spots; dorsal olive, its base blackish; caudal olive, edged and mottled with light blue; anal similar; pectoral olive.

Found in the tropical Atlantic; very common from the Carolinas to Brazil. Recorded from several Florida localities and common in Jamaica and Porto Rico, specimens being at hand from Puerto Real, Ensenada del Boqueron, Ponce, Arroyo, and Isabel Segunda. The four species of trunk-fishes found in our waters have the same general habits. They are all slow and sluggish, and are found on sandy bottom where there are open or naked areas surrounded by patches of algae. They seem peculiarly susceptible to cold, and on the Florida coast large numbers are frequently washed up on the shore that have apparently been killed by a sudden fall in the temperature.

Piscis triangularis capilli cornutus cui c media cauda cutanea aculeus longus erigitus, Lister, in Willughby, Hist. Pisc., Appendix, 19, 1686; locality not given.

Ostracion triangulatus aculeis duobus in capite et unico longiore superne ad caudam, Artedi, Genera, part 3, 56, 1738; after Lister in Willughby.

Ostracion triangulatus duobus aculeis in fronte et totidem in imo ventre, Artedi, Genera, part 3, 56, 1738; specimens seen in London at the house of Mr. Lillja and in the Naggs Head Inn.

Ostracion tricornis Linnaeus, Syst. Nat., ed. X, 331, 1758; after Artedi.

Ostracion quadricornis Linnaeus, Syst. Nat., ed. X, 331, 1758; after Artedi.

Ostracion listeri Lacépède, Hist. Nat. Poiss., I, 468, 1798; after Willughby.

Ostracion seecornutus, Mitchill, Amer. Monthly Mag., II, 1818, 328, mouth of Mississippi River.

Ostracion maculatus Hollard, Ann. Sci. Nat. 1857, 149.

Ostracion guineensis Bleeker, Ned. Tydskr. Dierk, II, 298, Guinea.

Ostracion gronovii Bleeker, Ned. Tydskr. Dierk, II, 298.

Acanthostracion polygonius Poey, Enumeratio, 175, 1876, Cuba.

Acanthostracion quadricorne, Stahl, l. c., 167, 1883.

Lactophrys tricornis, Jordan & Evermann, l. c., 1724, 1898.




FIG. 77.—*Lactophrys tricornis*.

Family LIX. TETRAODONTIDÆ. The Puffers.

Body oblong or elongate, usually little compressed, sometimes very broad; head and snout broad; belly capable of great inflation; skin scaleless, usually more or less prickly, spines or prickles usually weak and movable, not rooted; in one genus (*Ephippion*) the skin is armed with bony scutes forming a sort of carapace, approaching that seen in *Ostracion*; teeth confluent, forming a sort of beak which in each jaw is divided by a median suture; maxillaries curved outward behind premaxillaries; lips full; nostrils various. Spinous dorsal and ventral fins wanting, fins composed of soft rays only; dorsal fin posterior, opposite and similar to anal; caudal fin distinct; no ventral fins, pelvic bone undeveloped; no ribs; pectoral fins short and broad, upper rays longest; caudal fin and its vertebrae normally developed. Medifrontals articulated with supraoccipital, postfrontals confined to sides, ethmoid more or less projecting in front of frontals; postfrontals extending outward as far as frontals; proethmoid short and narrow, little prominent to view above; vertebrae few, 7 or 8 + 9 to 13. Gill-openings small, placed close in front of pectorals; air-bladder present.

There are about 10 genera and 60 species of this family inhabiting warm seas and found on soft or sandy bottom where there is some vegetation. They are sluggish in movement and noted for their habit of filling the stomach with air. When disturbed they then float on the surface, belly upward. Not used as food, the flesh being ill-flavored and reputed poisonous. None of them over 12 or 15 inches in length.

TETRAODONTINÆ.

- a. Frontal bones expanded sidewise and forming lateral roofs of orbits, postfrontals limited to posterior portions. Species chiefly marine.
- b. Nostril on each side with two distinct openings; frontal region longer than broad.
- c. Dorsal and anal fins comparatively long, falcate, each of 12 to 15 rays; nostrils sessile, or nearly so, not forming a distinct papilla; mucous tubes on upper part of head and on sides of body very conspicuous.

LAGOCEPHALUS, 123

- cc. Dorsal and anal fins comparatively short, rounded, each of 6 to 8 rays; nostrils at summit of a hollow, simple (or lobed) papilla; mucous tubes inconspicuous..... SPHEROIDES, 124

COLOMESINÆ.


- aa. Frontal bones narrowed and excluded from orbit; postfrontals being elongated and projected forward and connected with prefrontals; dorsal and anal fins short, rounded; snout very obtuse; vertebrae $8 + 11 = 21$; nostrils (probably) as in *Spherooides*. Fluvial species..... COLOMESUS

Genus 123. LAGOCEPHALUS Swainson.

Body comparatively elongate; skin smooth or variously prickly, prickles most developed on abdomen; abdomen capable of very great inflation. Dorsal and anal rather long, falcate, 12 to 15 rays each; caudal lunate. Nostril without distinct papilla, each with two distinct openings; mucous tubes on upper part of head and on sides of body very conspicuous. Lower side of tail with a fold.

The fishes of this genus are chiefly tropical, *Lagocephalus lagocephalus* reaching the coasts of southern Europe. Vertebrae about $8 + 13 = 21$. The increased number of vertebrae and of rays in vertical fins mark a transition toward the allied family, *Chonerhinidae*, in which there are about 29 vertebrae, the dorsal rays about 35, the anal 30.

- a. Body elongate; head 3.25 in length; depth 4.25..... *lævigatus*, 223
- aa. Body stout; head 2.8 in length; depth 3.5..... *pachycephalus*

FIG. 78.—*Lagocephalus lævigatus*.223. *Lagocephalus lævigatus* (Linnaeus). Smooth Puffer; "Tamboril"; "Conejo."

Head 3.3; depth 4.3; eye 4.8; snout 2.1; interorbital 2.3; D. 13; A. 12; pectoral 2; caudal 1.25. Body elongate, robust, caudal peduncle long and slender, abdomen greatly distensible and set with short spines which are depressed when abdomen is deflated, skin elsewhere smooth; head large and blunt; upper jaw with two great broad confluent incisor teeth with a median suture; lower teeth much smaller; side of body separated from belly by a ridge or angle; lines of mucous pores around eye and on top of head armed with weak spines; caudal lunate.

Color in spirits: Purple above, nearly black; side dirty-silvery, belly white. The young have three broad, dark, transverse bands on back of trunk, one on peduncle, one through base of dorsal, and one opposite pectoral; similar narrower bands on top of head.

The smooth puffer is a large, sluggish fish, reaching a length of 2 feet, and of no value as food. It is found in bays and river mouths from Cape Cod to Brazil; it is common southward, but rather rare north of Cape Hatteras; recorded from New York, Chesapeake Bay, Charleston, Biscayne Bay, Key West, Pensacola, St. Johns River, Tortugas, Cuba, and Jamaica. Found by us at Palo Seco and Mayaguez, most of our specimens being young, only 2.5 to 4 inches long.

- Ostracion cathoplateco oblongus*, Artedi, Genera Piscium, genus 58, species 13, 1738; after *orbis lagocephalus*, Grew, etc.
Tetrodon levigatus Linnaeus, Syst. Nat., ed. XII, 411, 1766, Charleston, S. C.; Poey, Fauna Puerto-Riqueña, 366, 1881; Stahl, l. c., 81 and 167, 1883.
Tetrodon curvus Mitchell, Trans. Lit. and Philos. Soc., I, 1815, 474, New York; young.
Tetrodon mathematicus Mitchell, Trans. Lit. and Philos. Soc., I, 1815, 474, New York.
Tetrodon pachycephalus, Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., IV, 73, pl. II, fig. 2, 1840, Brazil.
Holacanthus melanothus Gronow, Syst. Nat., ed. Gray, 24, 1854, Carolina; based on *Tetrodon levigatus* of Linnaeus.
Tetrodon livcolatus Poey, Synopsis, 432, 1868, Cuba; young.
Lagocephalus levigatus, Jordan & Evermann, l. c., 1728, 1898.

Genus 124. SPHEROIDES Lacépède. Swell-fishes.

Body oblong, not elongate; skin variously prickly or smooth, sometimes with cirri. A single, short, simple nasal tube on each side, with 2 rather large openings near its tip. Dorsal and anal fins short, little falcate, of 6 to 8 rays each; caudal truncate or rounded, rarely slightly concave. Vertebrae 8+10=18. Frontal bones expanded sidewise and forming lateral roof of orbit, postfrontals limited to posterior portions.

Species numerous, in warm seas; largely American. Our species represent two well-marked subgenera, the extremes of which appear very different from each other so far as the skulls are concerned. Some typical species of *Spheroides* approach *Canthigaster* in narrowness of frontal area.

SPHEROIDES:

- a. Skull very narrow above, interorbital area more or less concave, 2.5 to 6.5 times in length of long snout, 5 to 12 times in head; sides of body usually with small dermal flaps.
b. Interorbital space flat or moderately concave; no dorsal flaps; back without curved cross-streaks; upper ray of caudal produced.
c. Sides of head and body always smooth except sometimes a strip behind pectoral; spines larger, higher, more stellate, wider apart than in *Spheroides maculatus*, irregularly placed and often wholly wanting; side usually with small dermal slips or flaps, especially in the young.
d. Lower part of side with a series of about 12 round black spots; caudal fin barred with black and white. *spengleri*, 224
dd. Lower part of side with a series of black blotches appearing as short vertical bars (never round); caudal fin not barred *nepheus*
cc. Sides of head and body always prickly, as is back from upper lip to base of dorsal; belly prickly, prickles all similar, small, 3-rooted, stiff and close-set, never obsolete; no cirri; back with dark spots; black blotches on sides forming short oblique crossbars, those behind pectoral most conspicuous; caudal nearly plain, darker at tip.
e. Color dark-brown, with black blotches; a series of about a dozen irregular black spots along under side. *maculatus*
ce. Color dark, olivaceous above; black blotches on lower part of sides in form of short, oblique crossbars. *marmoratus*, 225

CHEILICHTHYS:

- aa. Skull very broad above, interorbital space broad, flattish, or very little concave, its width more than half snout and 2.66 to 4 in head; sides without series of dark blotches bounding line of belly.
f. Caudal fin rounded or subtruncate; back and sides with many small irregular black spots; no series of larger blotches bounding edge of belly.
g. Dark shades on back, broad, appearing as ground-color and crossed by pale curved crossbars and streaks forming arcs of concentric circles, these sometimes broken by reticulations everywhere profusely spotted with black in adult; snout somewhat produced *testudineus*, 226
ff. Caudal fin lunate or truncate, with angles acute or produced.
h. Dorsal rays 8; body more or less prickly, above and below; color brownish above, vermiculated with paler; eye broader than interorbital space (in young of 4 inches) *trichocephalus*
hh. Dorsal rays 10; body (in adult) everywhere perfectly smooth; interorbital space broad, equal to snout and twice diameter of eye; brown, with darker spots above *pachygaster*

224. *Spheroides spengleri* (Bloch). Southern Puffer; Swell Toad; Tambor; "Tamboril."

Head 2.38; eye 5; snout 2; interorbital 7; D. 7. Head compressed, long; snout very long, profile from tip of snout to above eye rising gradually, slightly concave in front of eye; interorbital space narrow, concave, slightly wider than in *S. nepheus*; posterior part of body usually smooth, with a number of small white dermal flaps; anterior part of body usually covered with small stiff prickles; region back of eyes and between pectorals and entire under-parts from throat nearly to anal fin prickly, those underneath strongest; head wholly naked above, a few weak prickles on cheek and a few on posterior part of throat.

Color in alcohol: Dark grayish-brown above on head and body, thickly covered and mottled with black or brown spots and paler areas, spots often grouped, forming blotches; under parts white; tip of chin pale, behind and on sides of which is a broad dusky border; dark of side bordered below by a line of about 12 round black spots, varying somewhat in size, but averaging about size of pupil and iris; pectoral, dorsal, and anal pale; caudal rather distinctly barred, a broad dark bar at base, then a somewhat broader pale bar, next another broad darkish bar, tip of fin pale.

The current descriptions of *S. spengleri* have been based partly on specimens of *S. nephelus*, an entirely distinct species, which has been confused with the present one. The two species differ markedly in the very differently-shaped head and coloration. In *S. spengleri* the head is more slender, snout much longer, profile less steep and less concave, and the interorbital somewhat wider. The color of *S. nephelus* resembles that of *S. marmoratus*, as does also its shape; the spots along side in *S. spengleri* are usually nearly round, while in *S. nephelus* they are never round, but are oblong or short vertical bars invading the white of lower parts from the dark side; the spots in *S. spengleri* are always more distinct and the caudal fin is always barred, a character which may be regarded as diagnostic.

An inhabitant of the West Indies, north to Florida; recorded from Tortugas, Key West, Garden Key, Biscayne Bay, Big Gasparilla, and Tampa; also from Cuba and Martinique; common about Porto Rico, specimens having been obtained at San Antonio Bridge, Puerto Real, Fajardo, Culebra, and San Geronimo; most of the specimens are small, but the fish attains a length of about a foot.

Tetrodon spengleri Bloch, Ichthyologia, I, 135, pl. 144, 1782, East Indies.

Le Tetrodon plumier Lacépède, Hist. Nat. Poiss., I, 504, 1797, Martinique; on a drawing by Plumier.

Le Sphéroïde tuberculé Lacépède, Hist. Nat. Poiss., II, 1, 1798, Martinique; on a front-view drawing by Plumier.

Tetrodon plumieri Bloch & Schneider, Syst. Ichth., 508, 1801, Martinique; after Lacépède.

Spheroïdes tuberculatus Pilot edition of Lacépède, VI, 279, 1831, Martinique.

Tetrodon turgidus Poey, Synopsis, 432, 1868, Cuba; not of Mitchill.

Spheroïdes spengleri Jordan & Evermann, 1732, 1898; in part only; description confused with that of *S. nephelus*.




FIG. 79.—*Spheroïdes spengleri*.

225. *Spheroïdes marmoratus* (Ranzani). *Spiny-back Blow-fish*.

Head 2.75; depth 4; eye 4.5 in head; snout long, 1.66 in head; D. 7; A. 6; P. 14. Outline of head concave in front of eye; eye full and high, its distance above a line drawn from corner of mouth to upper base of pectoral equal to its longitudinal diameter. Interorbital space very narrow, grooved, its width equal to that of pupil. Nostrils at end of a tube, situated about equally distant from end of snout and posterior edge of eye. Gill-opening equal to base of pectoral, but higher. Length of caudal peduncle from anal 2 in head. Length of head equal to half of distance in front of dorsal. Posterior rays of dorsal 1.5 in longest, which are 2.5 in head. Pectoral very broad, folding fan-like, margins scalloped, broadly rounded, lowest ray 1.8 in upper, which is 2.75 in head. Caudal fin slightly longer than distance of its base from dorsal, its rays all of equal length, 1.57 in head. Prickles on ventral surface between chin and vent, extending on side of head in front of pectoral fin, on side behind pectoral fin to vertical from dorsal, above from nostrils to dorsal; only snout, axil of pectoral, and caudal peduncle naked. Lateral line very faint, extending obliquely upward from side of snout under eye, then backward, curving slightly downward under dorsal, most distinct on side of tail.

Color in alcohol: Above very dark-brown, with black blotches, sides lighter, with very pale reticulations, a series of about a dozen irregular black spots along lower side; below white; caudal slightly dusky, with no indications of bars; other fins colorless.

This species differs from *Spheroides spengleri* in the high and prominent eye, very narrow interorbital, strongly concave outline of snout, extensive distribution of prickles, and in color. It is found in the West Indies to Brazil; recorded from Brazil and Jamaica; common in Porto Rico, the collection containing specimens from San Antonio Bridge, Aguadilla, Mayaguez, and Hucares.

Tetodon marmoratus Ranzani, Nov. Comm. Ac. Sci. Inst., Bonon., IV, 1840, 72, pl. 10, fig. 1, Brazil.

Spheroides marmoratus, Jordan & Evermann, l. c., 1733, 1898.

226. *Spheroides testudineus* (Linnaeus). "*Tambaril*"; *Tambar*; *Globe-fish*; *Puffer*.

(PLATE 41.)

Head 3; snout moderately long, 2 in head; eye small, about 7.5 in head, nearer gill-opening than end of snout; interorbital width 4 in head; D. 8; A. 6; skin of back from nape to before dorsal fin covered with small, sparsely set prickles; belly from throat to anal with prickles which are rather large and closely set; axil usually prickly, these prickles rarely wanting or obscured; sides sometimes with cirri. Back dark-olivaceous, with whitish curved lines and streaks paler than the ground-color, these streaks usually arranged as follows: A circle or rhomb on middle of back, in front of dorsal fin, this surrounded by an ellipse, the ellipse sometimes broken up by cross-streaks; before this 3 or 4 cross-streaks extending downward and backward, one at nape and one behind eyes connected on median line; back and sides with many irregular, round, blackish spots of different sizes; a dark bar at base of pectoral; caudal dusky at base, then pale, posterior half blackish; skull not very broad, interorbital area somewhat concave, prefrontal grooves narrow. Length a foot or more.

An inhabitant of the West Indies, generally common; occasionally ascending rivers; ranging north in the Gulf Stream to Woods Hole; recorded from several points in Florida and apparently common in Porto Rico, the collections containing specimens from Palo Seco, Aguadilla, Mayaguez, Ensenada del Boqueron, Fajardo, Hucares, San Antonio Bridge, and Isabel Segunda; taken by Mr. Gray at San Geronimo; recorded also from Jamaica, Dominica, Puerto Cabello, and Brazil.

Ostracion oblongus glaber, Artedi, Genera Piscium, 60; after Clusius, Willughby, etc., Balk, Amoen. Acad. I, 591, 1749.

orbis laevis variegatus (the Globe-fish), Catesby, Nat. Hist. Carolina, pl. 28, 1743, Virginia.

Tetraodon testudineus Linnaeus, Syst. Nat., ed. X, 332, 1758; based on Balk and Artedi.

? *Tetraodon punctatus* Bloch & Schneider, Syst. Ichth., 506, 1801, Brazil.

Tetraodon geometricus Bloch & Schneider, Syst. Ichth., 508, 1801, Virginia; after Catesby.

Tetraodon ammocryptus Gosse, Nat. Sojourn Jamaica, 287, 1851, Jamaica.

Auchisomus reticularis (Kaup) Richardson, Voyage Herald, 161, pl. 31, 1854; not *Tetraodon reticularis* Bloch & Schneider, which is *Tetraodon testudineus* Bloch, not of Linnaeus.

Holocentrus leionothos Gronow, Syst. Nat., ed. Gray, 24, 1854, American ocean.

Tetraodon testudineus, Poey, Fauna Puerto-Riqueña, 347, 1881; Stahl, l. c., 81 and 167, 1883.

Spheroides testudineus, Jordan & Evermann, l. c., 1734, 1898.

Family LX. CANTHIGASTERIDÆ. The Sharp-nosed Puffers.

This family includes small puffers, similar in external appearance to the *Tetraodontide*, but with the snout sharp and the back more or less compressed or ridge-like. The skeletal characters by which the group is defined are thus given by Dr. Gill: Medifrontals separated from supraoccipital by intervention of sphenotics, which are connected and laterally expanded, but short; prosethmoid prominent above, enlarged and narrowed forward. Vertebrae about 8 + 10. Head compressed, with a projecting, attenuated snout; dorsal and anal short, few-rayed. Nostrils obsolete, imperforate.

One genus, with 15 species, found in the tropical seas; none over 5 inches in length.

Genus 125. CANTHIGASTER Swainson.

Characters of the genus included above.

227. *Canthigaster rostratus* (Bloch). *Sharp-nosed Puffer*.

Head 2.6; eye 3.6; snout 1.7; interorbital 3; D. 6; A. 8; caudal of 9 rays, 1.6 in head. A tetraodont with a produced and pointed snout, elevated and somewhat compressed back, and the belly and back with weak prickles.

Color in life: Back and one-third distance down side olivaceous, rest of body pale, demarcation rather abrupt, an oblong black blotch extending downward and forward from front of dorsal; about 12 narrow blue lines radiating from eye; snout with 6 or 8 similar lines; side of head with many small

blue spots, fainter ones scattered over body; a group of finger-like bars upward from base of anal, and another behind the first on caudal peduncle; fins pale, except the caudal, which has some orange on middle rays, upper and lower rays brown, their bases each with a black blotch. In spirits, much the same, the blue becoming brown.

A small and prettily colored puffer, not uncommon; found in the West Indies, north, in rather deep water, to the banks off Pensacola; also in the Madeiras and Bermudas. Seven examples, 1.75 to 2.75 inches long, taken in the seine at San Antonio Bridge and Fajardo, and 2 from San Geronimo.

Tetrodon rostratus Bloch, Ichthyol., I, pl. 146, 1782, India.

Tetrodon capistratus Lowe, Proc. Zool. Soc. London 1839, 90, Madeira.

Tetrodon ornatus Poey, Synopsis, 433, 1868, Havana.

Canthigaster rostratus, Jordan & Evermann, l. c., 1741, 1898.

Family LXI. DIODONTIDÆ. The Porcupine-fishes.

Body short, broad, depressed above. Belly moderately inflatable, covered everywhere except on lips and caudal peduncle with spines, which are usually 2-rooted or 3-rooted at their bony base. Caudal peduncle short and slender. Mouth moderate, terminal, each jaw covered with a bony plate like the beak of a bird, these not divided by a median suture. Nostrils on each side forming a small tentacle, usually with two openings. Eye rather large, gill-opening moderate, immediately in front of pectoral, which is short, broad, and rounded. Dorsal and anal fins short, similar to each other, rounded in form and placed posteriorly.

The *Diodontidæ* comprise about 6 genera and 15 species, sluggish fishes, living on the bottom among weeds and corals, in tropical seas. When disturbed, they swallow air and float belly upward on the water. Their capacity of inflation is very much less than that of the *Tetraodontidæ*, from which family they differ chiefly in the stronger armature and in having no division in the bony plate of either jaw. Rarely used as food, being generally regarded as poisonous. The species are mostly well known in collections, the singular form having attracted the attention of travelers in the earliest times.

- a. Dermal ossifications very small, each one 2-rooted, with fine flexible spine or hair-like bristle. Nasal tentacles present TRICHODIODON
- aa. Dermal ossifications mostly 2-rooted; spines rather slender, but stiff and erectile. Nasal tentacle simple, with 2 lateral openings..... DIODON, 126
- aaa. Dermal ossifications all or nearly all 3-rooted, each with a short, stiff, immovable spine. Nasal tentacle simple, with 2 openings CHILOMYCTERUS, 127
- aaaa. Dermal ossifications of flattish, papery, or cartilaginous plates with minute hair-like papillæ; nostril short, entire, with 2 lateral openings..... LYOSPHERA

Genus 126. DIODON Linnæus. Porcupine-fishes.

Body robust, belly moderately inflatable. Dermal spines strong, stiff, most of them 2-rooted and erectile, a few 3-rooted and therefore immovable; both jaws entire; nasal tube simple, with two lateral openings. Pectorals broad, their margin undulate, upper lobe longest; vertical fins rounded, dorsal and anal short, posteriorly inserted, similar to each other.

Found in tropical seas, the few species being very widely distributed.

- a. Spines terete.
- b. Frontal spines not as long as post-pectoral spines (in adults not half as long, about as long as eye); predorsal spines very short, 3-rooted, fixed or nearly so; 20 spines in a series between snout and dorsal; post-pectoral spines very much elongate, especially in adult, shorter in young; dorsal rays 15; anal 15; upper lobe of pectoral little longer than lower. Adult above everywhere covered with round black spots, these largest in front of dorsal, smallest on naked area about mouth; white below; fins profusely spotted with black; young with fewer spots, but never with large blotches..... *hystrix*, 228
- bb. Frontal spines long, usually longer than post-pectoral spines, about twice as long as eye in adult; predorsal spines not shortened, 2-rooted, erectile; 14 to 17 spines in a series between snout and dorsal; post-pectoral spines not especially elongate, their development variable; dorsal rays usually 12; anal 12; pectoral broader than long, its upper lobe pointed, lower lobe rounded. Body marked with black spots and blotches, irregular in size, usually a broad black bar from eye to eye, continued below eye as a narrow bar; a broad bar across occiput; a black blotch above each pectoral; a short bar in front of dorsal; another in which dorsal is inserted; a blotch behind pectoral, and many small spots and blotches on upper parts; fins with few spots, usually unmarked in the young *holacanthus*, 229
- aa. "Spines compressed laterally, short; 15 spines in a series between snout and dorsal; upper parts covered with round spots, those about pectoral sometimes confluent into a blotch; fins immaculate" *maculifer*

228. *Diodon hystrix* Linnaeus. *Porcupine-fish; Erizo; Puerco Espino; "Guanabano."*

Head 3; depth 3.5; D. 13 to 15; A. 13 to 15. Spines strong, dilated at base, with a pair of basal grooves; frontal spines not as long as post-pectoral spines (in adults not half as long, about as long as eye); post-pectoral spines longer than any others, especially in adult, usually about as long as pectoral fin, those of posterior part of back and tail short and broad, 3-rooted, and therefore not erectile; predorsal spines very short, 3-rooted, fixed, or nearly so; about 20 spines in a series between snout and dorsal; upper lobe of pectoral little longer than lower; upper and lower part of tail with 2 or 3 pairs of 3-rooted, immovable, recumbent spines. Adult above everywhere covered with small, round, black spots, these largest in front of dorsal, smallest on naked area about mouth; white below; fins all more or less spotted in adult, nearly plain in young.

This species attains a length of about 3 feet. An inhabitant of the tropical seas, everywhere common, north to Lower California, Florida, and Hawaiian Islands; abundant in collections, being stuffed and dried as a curiosity; not used as food: recorded from the Tortugas, Key West, Biscayne Bay, and Indian River in Florida, and by Jordan & Satter from Jamaica; not seen by us in Porto Rico, but recorded from there by Professor Poey and Dr. Stahl.

Orbis echinatus Rondelet, De Piscibus, 324, 1558, Northern Ocean.

Guamajacu guara Maregrave, Hist. Nat. Bras., 159, 1648, Brazil.

Ostracion conico oblongus, Artedi, Genera Piscium, 60, No. 19, 1738.

Erizo, Parra, Desc. Dif. Piczas, Hist. Nat. Cuba, 60, pl. 29, fig. 1, 1787, Havana.

Diodon hystrix Linnaeus, Syst. Nat., ed. X, 335, 1758, India; after Artedi.

Diodon atinga Bloch, Ichth., IV, 75, pl. 125, 1787; not of Linnaeus.

Le Diodon (Plumier) Lacépède, Hist. Nat. Poiss., II, 1 and 10, pl. 3, fig. 3, 1798, Martinique; on a drawing by Plumier.

Diodon brachiatus Bloch & Schneider, Syst. Ichth., 513, 1801, Cuba; after Parra, pl. 29, fig. 1.

Diodon punctatus, Cuvier, Mém. Mus. Hist. Nat., IV, 132, 1818, no locality.

Diodon echinus (Rafinesque) Bonaparte, Cat. Met. Pisc. Eur., 87, 1846, Mediterranean Sea; accidental.

Paradiodon hystrix, Poey, Fauna Puerto-Riqueña, l. c., 346, 1881; Stahl, l. c., 81 and 166, 1883.

Diodon hystrix Jordan & Evermann, l. c., 1745, 1898.




FIG. 80.—*Diodon hystrix*.

229. *Diodon holacanthus* Linnaeus.

D. 12; A. 12. Very similar to *Diodon hystrix*, but with frontal spines usually longer than spines behind pectoral, about twice as long as eye. Predorsal spines not shortened, 2-rooted, erectile; about 14 to 17 in a series between snout and dorsal; post-pectoral spines not especially elongate, but movable; pectoral broader than long, upper lobe pointed, lower lobe rounded.

Coloration much as in *Diodon hystrix*, but more variable, spots fewer and larger; usually a broad black bar from eye to eye, continued below eye as a narrow bar; a broad bar across occiput; a black blotch above each pectoral; a short bar in front of dorsal; another in which dorsal is inserted; a blotch behind pectoral, and many small spots and blotches on the upper parts; fins with few spots, mostly immaculate in young.

Found in all warm seas, north to Florida Keys, Lower California, and Hawaiian Islands, its range coinciding with that of *Diodon hystrix*, from which it may prove to be not distinct. The distinctions are generally evident in the adult, but young examples apparently intermediate are often found. Possibly the two are different sexes of the same species. Length 1 to 2 feet. The single specimen obtained by us was seined in Guanica Bay, January 28, and is 5 inches long.

- Crayracion*, Nos. 9 and 15, Klein, Historia Pisc., 19 and 20, pl. 3, fig. 6, 1740.
Ostracion oblongus holacanthus, Artedi, Genera, 60, No. 20.
Diodon holacanthus Linnaeus, Syst. Nat., ed. X, 335, 1758, India; based on Artedi; misprint for *holacanthus*.
Le Diodon tacheté, Lacépède, Hist. Nat. Poiss., II, 13, 1798, New Cythera.
Diodon spinosissimus Cuvier, Mém. Mus., IV, 134, 1818, no locality.
Diodon melanopsis Kaup, Wiegmann's Archiv 1855, 228.
Paradiodon quadrimaculatus, Bleeker, Atl. Ichth., Gymnod., pl. 8, fig. 2, 1865.
Diodon maculatus, var. *a*, Günther, Cat., VIII, 307, 1870; based on *Diodon tacheté* of Lacépède; St. Croix; Jamaica
 Panama; South America; Hawaiian Islands; China; Sooloo Sea; Indian Ocean.
Erizo Guanabana Parra, Desc. Dif. Piezas Hist. Nat. Cuba, 62, pl. 29, fig. 2, 1787, Havana.
Diodon tiberosus Shaw, Gen. Zool., V, pl. 2, 436, 1804; after *Diodon tacheté*, Lacépède.
Diodon novemmaculatus Cuvier, Mém. Mus. Hist. Nat., IV, 136, pl. 6, 1818, no locality.
Diodon sepmaculatus Cuvier, Mém. Mus. Hist. Nat., IV, 136, pl. 7, 1818, no locality.
Diodon quadrimaculatus Cuvier, Mém. Mus. Hist. Nat., IV, 137, pl. 6, 1818, Otaiti.
Diodon multimaculatus Cuvier, Mém. Mus. Hist. Nat., IV, 136, 1818, no locality.
Diodon holacanthus, Jordan & Evermann, l. c., 1746, 1898.

Genus 127. CHILOMYCTERUS Bibrón. Burr-fishes.

Body broad, depressed, moderately inflatable. Dermal spines short, stout, immovable, triangular, each with 3 roots; nasal tube simple, with 2 lateral openings; tube sometimes rounded, sometimes flattened, and with partition feeble and easily torn, so that tentacle appears divided; caudal peduncle short; fins small, formed as in *Diodon*; jaws without median suture. Species numerous, of smaller size than those of *Diodon*, spines broader and lower, their bases forming a coat of mail.

CYCLICHTHYS:

- a*. Nasal tentacle subcylindrical, not divided.
b. Fins unspotted; supraorbital spines 2, with generally a tentacle between them; a spine in middle of forehead.
c. Superciliary edge raised.
d. Upper parts greenish-black, with a series of undulating blackish stripes running from nape backward; a similar series between eyes and across face; an ocellated black spot above pectoral; a larger one behind pectoral; an ocellated spot on each side of dorsal, and an elongated spot behind each of ventral antennæ..... *schorff*
dd. Upper parts plain, without series of lines; spots as in *C. schorff*..... *spinosus*
ddd. Upper parts covered with black, hexagonal reticulations..... *antillarum*
cc. Superciliary edge not raised; upper parts with numerous black dots, some with bluish centers; a black spot in middle of nape; a large kidney-shaped spot above pectoral, and a subtriangular blotch before and along base of dorsal fin; a series of antennæ along lower part of side..... *antennatus*, 230

CHILOMYCTERUS:

- aa*. Nasal tentacle flattened, divided; fins spotted with black; supraorbital spines 3, feeble, none on forehead.
c. Supraocular cirrus well developed; upper parts densely covered with small, round, blackish spots; a large black blotch before and around dorsal, another on each side above gill-opening and pectoral; spines short, compressed, anterior root flat, longer than the others..... *atinga*

230. Chilomycterus antennatus (Cuvier). Spiny Puffer.

(PLATE 42.)

Spines strong, but short; 2 above orbit, 1 more or less prominent in the middle of forehead. Superciliary edge not raised; generally a tentacle between superciliary spines. Tentacles along lower part of side, 1 on each side and in advance of anal fin being especially developed. Tail spineless, but roots of one pair of spines reaching across behind dorsal fin.

A black spot in middle of nape; a large kidney-shaped spot above pectoral, and a subtriangular blotch before and along base of dorsal fin; generally a small black spot below eye; some or all of these spots edged with lighter; upper and lateral parts with numerous black dots, some with a bluish pupil; abdomen brown; fins unspotted.

In an example 3 inches long, obtained at Mayaguez, and from which plate 42 in this volume was made, the general color is a light-brown, somewhat darker on back and under parts; a large grass-green blotch on occiput, a similar oblong one above base of pectoral, and another on back across base of dorsal fin; back, head, and side with numerous small, roundish black specks; iris yellow, pupil pale-blue; fins all pale-blue or greenish-blue.

West Indies and ranging southward; recorded from St. Croix, Jamaica, Porto Rico, and Cape of Good Hope. Length 8 inches. One specimen seized at Mayaguez and one taken at San Geronimo.

- Diodon antennatus* Cuvier, Mém. Mus., IV, 131, pl. 7, 1818.
Chilomycterus puncticulatus Poey, Anal. Hist. Nat., 346, 1881, Porto Rico.
Chilomycterus antennatus, Jordan & Evermann, l. c., 1750, 1898.

Family LXII. SCORPÆNIDÆ. The Rock-fishes.

Body oblong, more or less compressed, head large and with one or more pairs of ridges above, which usually terminate in spines. Opercle usually with 2 spinous processes; preopercle with 4 or 5. Mouth terminal, usually large, with villiform teeth on jaws and vomer, and usually on palatines. Premaxillaries protractile; maxillary broad, without supplemental bone, not slipping under preorbital. Gill-openings wide, extending forward below; gill-membranes separate and free from isthmus, usually no slit behind fourth gill. Scales ctenoid or sometimes cycloid, usually well developed, sometimes nearly obsolete. Lateral line single, continuous, concurrent with back; a narrow bony stay extending backward from suborbital toward preopercle. Ventral fins thoracic, of normal percoid form, 1, 5, rays branched; dorsal fin continuous, sometimes so deeply notched as to divide it into 2 parts, with 8 to 16 rather strong spines and about as many soft rays; anal rather short, with 3 spines and 5 to 10 soft rays; soft rays in all fins branched, except some or all of rays of pectoral; pyloric caeca in moderate or small number (fewer than 12). Pseudobranchiæ large. Air-bladder usually present. Actinosts moderate, inserted on posterior edges of hypercoracoid and hypocoracoid; ribs borne on enlarged pleurapophyses, Post-temporal bifurcate, normally connected; myodome more or less developed.

The *Scorpænidæ* are a large and very interesting family of about 30 genera and more than 250 species inhabiting all seas, but especially abundant in the temperate parts of the Pacific Ocean, where they form a large proportion of the fish fauna. They are non-migratory fishes, living about rocks. Most of them are of large size and many of the species are used as food, though some of them, particularly species in the genus *Scorpæna*, are in some places regarded as being poisonous. Whether there is any good reason for this belief has not been demonstrated. Many of the species of this family are viviparous, the young being produced after reaching considerable size.

- a. Dorsal spines 12; vertebræ 10 + 14 = 24.
- b. Palatine teeth present; anal rays usually III, 5.
- c. Bones of head scarcely cavernous; occiput with 2 pairs of spines; scales ctenoid or provided with dermal flaps.
- d. Pectoral with some of its median rays more or less branched.
- e. Scales on top and sides of head ctenoid; cranium much as in *Sebastes*, the armature moderate..... HELICOLENUS
- ee. Scales on top and sides of head cycloid or wanting; cranium with many spines..... SCORPÆNA, 128
- dd. Pectoral rays all simple; head more or less scaly, the scales ctenoid..... PONTINUS, 129
- cc. Bones of head with large muciferous cavities; occiput with only 1 pair of spines; scales cycloid; pectoral rays 20 or more; head scaleless above; no groove at occiput; some of the pectoral rays branched..... SETARCHES


Genus 128. SCORPÆNA (Artedi) Linnæus. The Scorpion-fishes.

Body oblong, somewhat compressed. Head large, not much compressed, naked above, and more or less uneven with spinous ridges, often with dermal flaps. Mouth large, with bands of villiform teeth on jaws, vomer, and palatines. Scales mostly ctenoid, of moderate size, often with skinny flaps. Dorsal fin with 12 stout spines; anal with 3 spines, second commonly the longest; pectoral large, rounded, base usually procurvent, some or all of upper rays divided, lower simple; ventrals inserted behind pectorals. No air-bladder. Vertebræ 10 + 14 = 24. Species numerous in the tropical seas; fishes of singular forms and bright colors; the variation in squamation and armature very great, but, as in most similar cases, it is not easy to find definite characters for subdivision.

- a. Breast scaly.
- b. Occiput with a distinct quadrate pit.
- c. Supraocular tentacle less than twice diameter of orbit.
- d. Dorsal rays XII, 9.
- e. Top of head scaleless, 3 small spines on suborbital carina..... *agassizii*
- ee. Top of head not wholly scaleless, interorbital space incompletely scaled; suborbital carina with 6 spines. *cristulata*
- dd. Dorsal rays XII, 10.
- f. Anterior border of orbit with no distinct pit below it.
- g. Suborbital stay with 3 distinct spines; third anal spine longer and stronger than second.
- h. Axil pale, with small dark spots; body with a few large, diffuse dark spots *brasiliensis*, 231
- hh. Axil pale without spots; body and head with numerous small milky-white spots..... *albifimbria*, 232
- gg. Suborbital stay without spines; cheek more or less scaly; second anal spine longer and stronger than third. *bergii*, 233
- ff. Anterior border of eye with a distinct pit between it and suborbital stay.
- i. About 30 scales on lateral line, most of scales with dermal flaps; supraorbital flap large, longer than eye; axil black, with large white spots.
- j. Color rather pale or reddish; interorbital area narrow; occipital pit deep..... *plumieri*, 234
- cc. Supraocular tentacle more than twice diameter of eye; flaps on lateral line longer than eye; suborbital stay with a small spine near its center, another at its posterior end; axil gray with many small white spots .. *grandicornis*, 235
- bb. Occiput with only a very shallow depression or none.
- k. Pectoral with but 2 branched rays; no pit at occiput..... *incermis*.

231. *Scorpaena brasiliensis* Cuvier & Valenciennes.

Head 2.6; depth 2.66 to 3; orbit 4 in head; D. xii, 10; A. iii, 5 (5.5); transverse rows of scales (oblique) about 25 to 30, (vertical) 50 to 60; tubes 25 to 30. Body short, compressed, profile convex, depth of caudal peduncle a little less than 4 in head. Head compressed; interorbital space narrow, a little more than half orbit, about 7 in head, deeply concave, deepest between preocular spines, and with 2 marked longitudinal ridges on frontal bones, parallel with supraocular ridges; a deep pit at occiput, deepest behind, its anterior side sloping back from the base of coronal (tympanic) spines, its posterior side nearly vertical; preocular, supraocular, postocular, and coronal (tympanic) spines moderate, parietal and nuchal spines sharpest, their ridges thin and sharp, exoccipital spines and ridges present; spine between orbit and occipital small, not bifid. Preorbital very broad with 2 sharp spines, no pit between anterior inferior border of orbit and suborbital stay, latter low, with 2 or 3 small spines; uppermost preopercular spine much the longest, a small spine at its base, others very small or almost obsolete; opercular spines rather small, with not very prominent ridges; all ridges of head less prominent than in *Scorpaena plumieri*. Jaws equal, lower with small symphyseal knob; maxillary reaching posterior edge of pupil, about 2.25 in head; teeth on jaws, vomer, and palatines in rather narrow bands. Pseudobranchiae reaching down nearly to epihyal bone; gillrakers short, compressed, about 7 on anterior limb. Dorsal fin deeply notched, membrane reaching halfway up twelfth spine; dorsal spines slender, low, the longest equal to maxillary, about 2.25 in head; soft rays higher, about 1.75 in head; anal spines graduated, second a little the thickest, soft rays 1.6 in head; caudal truncate or very slightly rounded; pectoral reaching beyond origin of anal, a little shorter than head, its base not procurent, 10 lower rays slightly thickened, exerted and simple, upper rays (except

FIG. 81.—*Scorpaena brasiliensis*.

uppermost one) branched; ventrals reaching beyond vent, last soft ray attached to body for three-quarters of its length by a rather broad membrane. Supraocular flap long and slender, longer than orbit, a little more than 3 in head; preocular flap small; flaps at base of spines of preorbital and lower spines of preopercle, small ones on cheek, membrane of spinous dorsal, and on many of scales of body; larger flaps on lateral line and along base of dorsal fin. Scales large, rather smooth, with membranaceous edges; a few rudimentary scales on front and flap of opercle, on preopercle, and lower part of cheek; breast with small scales.

Color: Dusky-olivaceous or brownish, whitish below; a few large diffuse dark spots on side above, nearly as large as eye; posterior part of each scale darker, giving a slightly speckled appearance; axil pale, with small dark spots, which are also sparsely present along lower part of side; pectoral mottled, faintly banded, the lower part paler; spinous and soft dorsal and anal irregularly marbled; caudal with median and terminal blackish bands; ventrals dusky at tip; side of head dark with some small darker spots; snout, interorbital space, and tip of maxillary dark, faintly marbled; under side of head whitish or marbled with brownish; peritoneum white.

This species ranges from the South Atlantic and Gulf coasts of the United States to Brazil. It has been recorded from Charleston, S. C., Indian River, Biscayne Bay, Key West, Pensacola, Pensacola

Snapper Banks, Egmont Key, Boca Grande, Jamaica, and Rio Janeiro. The collections from Porto Rico contain but a single specimen, 2.75 inches long, taken in the seine at Palo Seco, January 16. It differs somewhat from larger examples in having the second anal spine stronger than third.

Scorpaena brasiliensis Cuvier & Valenciennes, Hist. Nat. Poiss., IV, 305, 1829, Brazil; Jordan & Evermann, l. c., 1842, 1898.
Scorpaena stearnsi Goode & Bean, Proc. U. S. N. M. 1882, 421, Pensacola, Fla.

232. *Scorpaena albifimbria* Evermann & Marsh, new species.

Head 2.1; depth 2.4; eye 3.3; snout 4.25; maxillary 2; mandible 1.9; interorbital 7; preorbital 6.5; scales 7-45-15, about 21 pores; D. XII, 10; A. III, 5; pectoral 19. Body very short; head heavy and broad, width 1.6 in its length; snout broad and short; profile rather evenly curved from tip of snout to origin of dorsal fin; occipital pit shallow but distinct; interorbital space rather broad, shallow; no pit below anterior part of eye; mouth large, maxillary reaching posterior border of eye; spines of head strong; supraocular ridge moderate, a strong preocular spine and two smaller supraocular ones; coronal, postocular, nuchal, parietal, and exoccipital spines strong; opercle with 2 spines, lower terminating a




FIG. 82.—*Scorpaena albifimbria*.

strong ridge; 4 preopercular spines, uppermost strong, with a small accessory spine on its base; suborbital stay prominent, with 3 evenly spaced, strong spines, in line with upper preopercular spine; nasal spines obscure; preorbital with 2 broad, blunt spines. Cephalic filaments mostly long, but slender, nasal pair short and broad; preocular pair long, greater than eye; supraocular pair slender, about equal to eye; a few small dermal flaps below suborbital stay, and a very broad one on lower edge of preorbital continuous with skin of the 2 preorbital spines. Scales small, scarcely ctenoid; scaling of head obscure; dermal flaps numerous, largest along lateral line. Fins large; longest dorsal spine 3 in head; longest ray 2.9; second anal spine 2.1, longer and somewhat stronger than third; longest anal ray 2.1; pectoral reaching origin of anal, 1.6; base broad, 2.5; third, fourth, fifth, and on one side sixth rays branched, others undivided; ventral 2.2; caudal 1.6.

Color in alcohol: Pale-rosy, with dark dustings on head, humeral region, back, and side; top and sides of head profusely covered with fine milky-white specks; similar spots on body, but less numerous; dermal flaps on head and body all milky-white; under part of head white, a large rosy blotch at base of fourth to seventh dorsal spines; dorsal rosy, richest on soft portion; anal pale, with some black on membrane between first and second spines; caudal pale; pectoral rosy, with dark dustings around margin; axil pale, without spots; ventrals pale, dusky at tips. In life the rosy color was doubtless much more evident.

This species seems quite distinct from any previously known species of *Scorpaena*. Its most characteristic features are the numerous milky-white specks and dermal flaps. The only specimen

obtained is the type (No. 49532, U.S.N.M.), 1.75 inches long, taken in the tangle February 8 at Fish Hawk station 6093, off Culebra Island, 5.25 miles southwest of Culebritas Light-house, in 15 fathoms, on live coral bottom.

Albus, white; *fimbria*, flap.

233. *Scorpæna bergii* Evermann & Marsh, new species.

Head 2.4; depth 2.7; eye 3.5; snout 4.8; maxillary 2.4; mandible 2; interorbital 6; preorbital 6.5; D. XI, 10; A. III, 5; pectoral 17; scales 9-38-15, about 22 pores. Body short and stout; head short and broad, width 1.4 in its length; snout short and broad; cephalic spines and ridges prominent; supraocular ridges strong, with 3 spines; interorbital groove deep, its 2 ridges very low; a pair of strong nasal spines; occipital pit deep, a deep transverse postocular groove crossing occipital pit; a small spine at each anterior angle of pit and a larger one at each posterior angle, back of which is, on each side, a smaller spine; a postocular and 2 small humeral spines; opercle with 2 strong flat spines and 2 short blunt ones below; preopercle with 5 blunt spines, upper one largest and with a small accessory spine on its base; no pit below anterior part of eye; suborbital stay strong, without any spines except terminal one at posterior end; preorbital with 3 blunt spines, 2 of them obscure; maxillary broad, reaching past pupil; cephalic filaments short; a pair of short nasal cirri, a small pair on anterior part of supraocular ridge and a larger pair on posterior part of same ridge; preorbital with




FIG. 83—*Scorpæna bergii*.

2 flat flaps, anterior the smaller; dermal flaps on body few and small, a series of 7 or 8 along lateral line; fins high; distance from snout to origin of dorsal 3 in body; first dorsal spine 1.5 in second, which equals eye; fourth spine longest, 2.6 in head; longest dorsal ray 2.75 in head; first anal spine 1.75 in second, which is 2.2 in head; third anal spine weaker and somewhat shorter than second; longest anal rays 2.5 in head, pectoral broad, its base 3 in head, length 1.6; tips of rays nearly reaching anal; ventral 2 in head, reaching past vent; caudal 1.8 in head.

Color in alcohol: Dark-grayish on head and body, under parts paler; a broad black bar crossing body between soft dorsal and anal and extending across these fins; caudal peduncle pale, with a narrow black bar at base of fin; dorsal grayish mottled with black and white, a large round black spot between third and seventh spines; soft dorsal with 2 or 3 broad black bars with pale interspaces; anal pale at base, with a few dark spots, then a pale band, next a broad black band continuous with that on body, then a broad white band, followed by another blackish bar, tip of fin white-edged; pectoral gray, crossed by 3 blackish bands; axil pale, with 5 or 6 large round brown spots; ventral pale.

This species seems related to *S. pamosa* Cramer, from Panama, but differs in many important respects, notably in the fewer pectoral rays, the greater length of the pectoral, and the color. The presence of but 11 dorsal spines instead of 12, the usual number, may be merely an individual variation

and of no specific value. It is known only from the type (No. 49533, U. S. N. M.), 3 inches long, obtained at Mayaguez, January 20, and one cotype (No. 813, U. S. F. C.), 2 inches long, seined at Culebra Island, February 11.

Named for Dr. Carlos Berg, director of the National Museum of Buenos Ayres, in recognition of his excellent work on South American fishes.

234. *Scorpæna plumieri* Bloch. *Rascacio*.

Head 2.25 to 2.5; depth 3; eye 5; snout 3.75; maxillary 2; mandible 2; interorbital 4.75; preorbital 5.5; D. XII, 10; A. III, 5; scales 8–40–16, about 25 pores. Body short and thick; head irregular in form, with many grooves and pits, and numerous fleshy flaps; a large, deep quadrate pit on occiput; a large, deep pit below anterior part of eye, between it and front of suborbital stay; supraocular ridge with 2 long flaps, posterior one broadest, each nearly twice length of eye; anterior nostril with a double fringed flap, about two-thirds length of eye; a pair of simple, slender tentacles on snout, each somewhat longer than eye; 3 or 4 flaps on edge of preorbital; numerous other smaller, shorter flaps along edges of opercle and elsewhere about head; parietal, nuchal and exoccipital spines present; suborbital stay with 3 or 4 sharp spines; preopercular and opercular spines moderate, bluntish, a smaller spine at base of upper preopercular spine; occiput with 2 pairs of spines; a few scales on preopercle and opercular flap, head otherwise scaleless; maxillary long and broad, reaching to posterior border of eye; lower jaw included; breast covered with small embedded scales; scales of body large, thin, firm, many of them with membranaceous flaps; lateral line with a series of fleshy flaps; dorsal low, longest spine 2.6 in head, longest ray 2.6; first anal spine short, second much stronger and somewhat longer than third, 2.5 in head; longest anal ray 2.2; pectoral broad, procurvent, reaching about to anus, 1.4 in head, upper rays branched, lower simple, tips free; ventrals reaching anus, 1.9 in head, equal to caudal.

Color, highly variegated and subject to much variation; sand-color, with 2 broad, blackish shades on body and 1 on head; belly purplish; lower parts of head finely speckled in all shades of light, dark, and pearly bluish; upper parts covered with whitish cirri profusely speckled; surface appearing as if covered with sand; eye with radiating dark spots; dorsal covered like body with some well-marked whitish spots; dark band of body passing on to soft dorsal; caudal variously mottled, with 3 pale and 3 black bands; anal whitish, variegated with reddish and black; ventrals similar, with more maroon-red; pectoral still more variegated, tip scarlet-shaded; inside of pectoral largely bright-yellow, then blackish, tinged with cherry-red; axil jet-black, with large, round, white spots, not fading in alcohol; lips barred with black and whitish; branchiostegal membranes and angle of mouth bright-yellow; peritoneum white.

The rascacio is found from southern Florida to Brazil; it is common among the Florida Keys and has been recorded from Clearwater Harbor, the Tortugas, and Key West. It was obtained by the Nutting expedition off Key West in 60 fathoms. It is known also from Jamaica, Cuba, and Martinique, and Porto Rican specimens were obtained by us at San Antonio Bridge, Mayaguez, Puerto Real, Guanica, Ponce, and Hucaras. It is, next to *S. grandicornis*, the most abundant species of the family in Porto Rican waters. It reaches a length of a foot or more.

The above description is based upon a specimen 10 inches long, from San Antonio Bridge.

Scorpæna plumieri Bloch, Nya, Handl. Stockh., X, 234, 1789, Martinique.

Scorpæna bufo Cuvier & Valenciennes, Hist. Nat. Poiss., IV, 306, 1829, Martinique; Poey, Fauna Puerto-Riqueña, 323, 1881.

Scorpæna rascacio Poey, Synopsis, 303, 1868, Havana.

Scorpæna plumieri, Jordan & Evermann, I. c., 1848, 1898.

235. *Scorpæna grandicornis* Cuvier & Valenciennes. *Rascacio*; *Lion-fish*.

Head 2.5; depth 2.6; eye 4; snout 4; maxillary 2.2; mandible 1.75; interorbital 5; preorbital 6.75; D. XII, 10; A. III, 5; P. 17; scales 9–40–15, about 28 pores. Body short, more compressed than in *S. plumieri*; head rough with many spines and ridges; a deep quadrate pit on occiput; a shallow suborbital groove but no pit at its anterior end; interorbital groove deep but narrow; suborbital stay not strong, with 1 small spine near its middle and 1 at its posterior end; supraocular ridge with 1 strong spine on posterior portion; 2 spines on side of nape at posterior angle of pit, the anterior with a fringed filament; below these, 3 other spines in an irregular row; opercle with 2 flat spines; preopercle with 3 flat spines, the one at angle strongest, and with a supplementary spine on its base; a pair of short nasal spines; supraocular filament long, broad, and fringed, more than twice length of

eye; a short filament on front of supraocular ridge and a small one on upper part of eye; anterior nostril with a short fringe; a pair of short filaments at tip of snout; 2 or 3 short, broad, dark flaps on cheek and others on various parts of head; dorsal high, longest spine 2 in head, equaling longest ray; first anal spine short, less than half second, which is slightly longer, but much stronger, than third, 1.9 in head, or equaling longest anal ray; pectoral broad, width at base 2.5 in head, length 1.2, upper rays but one divided, lower simple, their tips free, longest ray reaching past origin of anal; ventrals shorter, barely reaching origin of anal, 1.6 in head; caudal 1.5 in head. Scales smooth, thin, and firm; a series of very large dermal flaps along lateral line and smaller flaps scattered over body; a few embedded scales on cheek and opercle; breast scaled.

Color in life variable. Three examples, taken at San Antonio Bridge January 14, were described as follows: Body dark-brown, inclining to brick-red; belly pale-reddish; top and sides of head olive-brown; dorsal brown, soft portion palest; caudal reddish-brown, crossed by 2 narrow pinkish bars; anal brownish, a large black blotch near base of rays, then a broad whitish bar with reddish blotches, next a series of 5 black spots on rays, then brown, and last a pale border; pectoral brown with a




FIG. 84.—*Scorpæna grandicornis*.

narrow pale border and 2 irregular dark crossbars separating lighter-brown, central portion of fin with light wash of old gold; axil pale-grayish with numerous small white spots; similar but larger spots upon branchiostegals; lower jaw greenish, with series of white blotches; ventrals pale at base, then 3 oblong black blotches on middle rays, beyond which is pale-brown with pale border. Two other examples taken at same time had body brownish, belly rosy; head mottled-brownish; dorsal brown, soft part paler; caudal pale with 2 broad brown bars; ventrals rosy at base, dark at tips; pectoral yellowish-brown, with black on tips of upper rays and a black blotch at base of lower rays; ocular cirri brown.


The lion-fish occurs from the Florida Keys to Brazil, and has been recorded from Key West, Havana, Santo Domingo, Porto Rico, Jamaica, and Martinique. It is the most abundant species of the family about Porto Rico, our collections containing numerous specimens from San Juan, San Geronimo, Aguadilla, Mayaguez, Puerto Real, Ensenada del Boqueron, Guanica, Ponce, Hucares, and Fajardo. While *Scorpæna plumieri* is often found in water of considerable depth, this species frequents shallow water among algae, where, on account of its remarkable protective coloration and long alga-like filaments, which are also in the nature of protective mimicry, together with the rough armature of the head and its poison spines, it would seem extraordinarily well protected. It reaches the length of a foot or less and is of striking appearance, much dreaded by the fishermen, who pronounce it "muy malo" and can not be induced to touch it.

Scorpæna grandicornis Cuvier & Valenciennes, Hist. Nat. Poiss., IV, 309, 1829, Martinique, Porto Rico, Havana, Santo Domingo; Poey, Fauna Puerto-Riqueña, 323, 1881; Stahl, l. c., 78 and 164, 1883; Jordan & Evermann, l. c., 1850, 1898.

Genus 129. PONTINUS Poey.

This genus differs from *Scorpaena* chiefly in having the pectoral rays all simple and only their tips free; anal with 5 to 9 rays; suborbital keel composed of 3 or 4 distinct, differentiated spines, 2 prominent retrorse spines on each preorbital; no pit at occiput; scales ctenoid; cheek and opercles usually scaly; pectoral not procurent. The American species all have D. XII, 10; A. III, 5. Of the seven known American species only two were obtained in Porto Rico.

- a. Base of pectoral broad, fin fan-shaped.
 b. Snout naked above, as is interorbital space.
 c. Eleventh dorsal spine nearly as long as twelfth.
 d. Eye small, 4.1 in head *beanorum*, 236
 dd. Eye larger, 3.4 in head *macrolepis*, 237
 ee. Eleventh dorsal spine half as long as twelfth.
 e. Eye 5.5 in head; maxillary reaching two-thirds across eye; head 4 in total length; spinous dorsal low; pectoral pointed; supraorbital tentacle 5 in total length; carmine, without marblings *castor*
 ee. Eye 4 in head; maxillary reaching anterior third of orbit; spinous dorsal high; pectoral rounded; carmine with vertical rosy bands *pollux*
 bb. Snout fully scaled above; interorbital space with few scales, top of head otherwise entirely scaly *rathbuni*
 aa. Base of pectoral narrow.
 f. Head without filaments; nape and top of snout scaly; ventrals reaching vent. Pectoral rays 16 *longispinis*

FIG. 85.—*Pontinus beanorum*.236. *Pontinus beanorum* Evermann & Marsh, new species.

Head 2.5; depth 3.5; eye 4; snout 4; maxillary 2.4; mandible 2.25; interorbital 9; preorbital 6.2; D. XII, 10; A. III, 5; P. 16; scales 7-36-11, about 24 pores. Body short and stout, head heavy, its width 1.6 in its length; snout moderately long, broad; profile regularly curved from tip of snout to origin of dorsal; occiput scarcely depressed; spines of head much as in *P. macrolepis*, but smaller; a pair of small nasal spines; supraocular ridge with 3 spines, first at anterior end scarcely perceptible, the 2 others near posterior end, each short and weak; postocular, tympanic, parietal, and nuchal spines present but short; 1 paroccipital and 2 humeral spines present, all small; 2 stouter, flat spines on opercle, lower at end of a low ridge; 4 preopercular spines, upper largest and with a small accessory spine on its base; suborbital stay strong, with 4 spines, first small and indistinct, near anterior end, second under anterior edge of pupil somewhat stronger, third under posterior edge of eye and last at end of stay, third and fourth about equally strong; preorbital with 3 broad, blunt spines; no occipital pit; no pit below eye; interorbital narrow, a central groove, between which and each supraocular ridge there is a small sharp ridge; premaxillary broad, reaching middle of eye; nape, opercle, and cheek scaled, rest of head naked; scales of body moderate in size, firm, ctenoid; breast scaly; head almost without cirri, the only ones being a pair of small internasal filaments, a pair of small supraocular ones, and a cluster on

edge of suborbital; no dermal flaps on scales; teeth in villiform bands on jaws, vomer, and palatines; anterior nostril tubular, with a rather broad fringed flap; distance between anterior nostril and eye greater than interorbital width; distance from tip of snout to origin of dorsal twice length of maxillary; first dorsal spine 1.6 in second; fourth dorsal spine about 3.5 in head, penultimate spine 5 in head, last about 4, or about twice length of first spine; longest dorsal ray 2 in head; first anal spine very short, about 7.5 in head, second about 3 times length of first, 2.6 in head; third somewhat shorter and much weaker than second; pectoral base not so broad as in *P. macrolepis*, 3.75 in head, rays all simple, reaching origin of anal, 1.6 in head; ventrals shorter, their tips scarcely reaching vent, 1.6 in head; caudal 1.7.

Color in life: Back and top and sides of head pale-rosy, with irregular brownish markings; lower parts of head richer rosy; belly white, with some rosy; dorsal pale-rosy, with a series of large irregular brownish spots along outer border; soft dorsal with brownish spot; caudal pale-rosy, upper half with brown spots, lower half plain; anal rosy, dark at tip; pectoral and ventral rosy, blackish on outer portion.

This species resembles *P. macrolepis*, from which it differs in the much smaller eye, longer snout, fewer and smaller cephalic cirri, somewhat smoother head, narrower pectoral, and the very different coloration. The only example obtained is the type (No. 49534, U. S. N. M.), 5.5 inches long, taken in the beam trawl January 13, 1899, at Fish Hawk Station 6050, 1.25 miles northward from the entrance to San Juan Harbor, in 91 fathoms.

This interesting species is named in honor of Dr. Tarleton H. Bean, director of forestry and fisheries of the United States Commission at the Paris Exposition of 1900, and his brother, Mr. Barton A. Bean, acting curator of fishes in the U. S. National Museum, in recognition of their valuable services to American ichthyology.

237. *Pontinus macrolepis* Goode & Bean.

(PLATE 43.)

Head 2.2; depth 3.25; eye 3.4; snout 4.5; maxillary 2.4; mandible 2.2; interorbital 10; preorbital 9; scales 6-35-10, about 22 pores; D. xii, 10; A. iii, 5; P. 17. Body short; head large, its width half its length; snout short and broad; profile not much arched; occiput not depressed; interorbital groove deep, a slight ridge on each side at base of supraocular ridge; no pit under anterior part of eye; spines of head strong; a pair of moderate nasal spines, 1 at anterior end of supraocular ridge and 2 stronger ones near its posterior end; postocular, tympanic, nuchal, and parietal spines strong; 1 paroccipital and 2 humeral spines present; 2 flat, stout, opercular spines, and 4 on preopercle, upper one strongest and with an accessory spine on its base; suborbital stay prominent, with 3 strong spines; 2 strong spines on preorbital, each directed backward; scales strongly ctenoid, rubbing off easily; nape, opercles, and cheek scaled, rest of head naked; breast with small embedded scales; cephalic tentacles moderate, a short, broad nasal pair; a slender one at preocular spine, and a longer one, about 2 in eye, at supraocular spine; another small pair on nape and 2 small filaments on preorbital; distance from tip of snout to origin of dorsal twice length of maxillary; first dorsal spine about three-fifths of second, third longest, 2.4 in head; longest dorsal rays 2.5 in head; eleventh spine about 4.5 in head; twelfth 3.75; longest dorsal ray 2.5; first anal spine about 3 in second, which is 2.6 in head and longer and stronger than third; longest anal ray about 2.4; pectoral rays all simple, scarcely reaching anal, 1.5 in head; ventral shorter, 2 in head; caudal 2.1.

Color in life: Pale below, reddish above; body with about 6 vertical bars of deeper red reaching to lateral line or below; head chiefly red, iris gray, upper and lower edges red, pupil black; dorsal with 2 long rows of light-red blotches; pectoral with 3 V-shaped, vertical, light-red bars; bases of ventral and pectoral reddish; caudal with 3 curved, vertical, yellowish bars; a series of oblong reddish spots on caudal membrane at margin.

Hitherto known only from the type, a specimen 4.4 inches long, from off Yucatan. Our collection contains a single specimen 5.5 inches long, dredged by the *Fish Hawk* at station 6068, in 224 to 237 fathoms, in Mayaguez Harbor, 7.5 miles northwest from the Mayaguez custom-house.

Pontinus macrolepis Goode & Bean, *Oceanic Ichthyology*, 257, fig. 247, 1896, lat. 20° 59' 30" N., long. 86° 23' 45" W., at Albatross station 2354, off Yucatan, in 130 fathoms; Jordan & Evermann, l. c., 1856, 1898.

Family LXIII. TRIGLIDÆ. The Gurnards.

Body elongate, usually more or less fusiform, covered with scales or bony plates. Head externally bony, entirely cuirassed with rough, bony plates, some of which are armed with spines; eyes high; mouth terminal or subinferior; premaxillaries protractile; maxillary without supplemental bone, slipping under preorbital; teeth very small, in bands in jaws, and usually on vomer and palatines; gills 4, a large slit behind fourth; pseudobranchiæ present; gillrakers various; gill-membranes free from isthmus. Ventral fins thoracic, wide apart, separated by a flat area, their rays 1, 5. Spinous dorsal present, short; soft dorsal similar to anal, which is without spines; caudal narrow, few-rayed; pectoral large, with broad base, with 3 lower rays detached, forming feelers. These free rays are used chiefly in search for food, turning over stones, exploring shells, etc. Air-bladder present; pyloric cæca usually present, few in number.

Singular-looking fishes, found in all warm seas, comprising 5 genera and about 40 species, some of them in rather deep water, these red in color, the others living about rocks.

- a. Palatines with teeth.
- b. Dorsal spines low, the longest usually much shorter than head; scales moderate, 50 to 80 pores PRIONOTUS, 130
- bb. One or 2 of the dorsal spines greatly elevated, about as long as body; scales large, rough, the pores 40 .. BELLATOR
- aa. Palatines toothless; scales small.
- c. Lateral line without enlarged bony plates CHELIDONICHTHYS
- cc. Lateral line armed with a series of transverse bony plates TRIGLA

Genus 130. PRIONOTUS Lacépède.

Body subfusiform; profile of head descending to the broad depressed snout, which is much longer than the small eye; eyes close together, high up; surface of head entirely bony, bones rough with ridges and granulations; scales on head few or none; preopercle with 1 or 2 sharp spines at its angle; opercle with a sharp spine; nape with two strong spines; a spine on shoulder-girdle. Mouth rather broad; bands of small, almost granular, teeth on jaws, vomer, and palatines; gill-membranes nearly separate, free from isthmus; gillrakers rather long. Body covered with small, rough scales, which are not keeled; lateral line continuous; scales on breast very small. Dorsal fins distinct, first of 8 to 10 rather stout spines, third usually highest, but mostly shorter than head; anal fin similar to soft dorsal; pectoral fin with 3 lower anterior rays thickened, entirely free from each other and from fins; ventrals 1, 5, wide apart, with a flat space between them, inner rays longest. Pyloric cæca in moderate number; air-bladder generally with lateral muscles and divided into 2 lateral parts; vertebrae 10 or 11+15.

Species numerous, all but one being American, representing in America the Old World genus *Trigla*, some in deep water. They are well defined and easily recognized, but vary considerably with age, and are not easily thrown into subordinate groups. Most of the characters in the following analysis have been taken from adult individuals. Young examples in most cases differ from adults in the following respects, in addition to those characters which usually distinguish young fishes: The spines on the head are sharper, more conspicuous, and more compressed in the young, and some spines, especially those on side of head, disappear entirely with age. The interorbital space is more concave in the young. The pectoral fins are also much shorter. The gillrakers are longer in the young, and proportionately more slender, and some of the color markings—especially the darker cross-shades—are more conspicuous, while the spots on body and fins are less so.

Of the 21 recognized American species of the genus *Prionotus* 9 are known from the Pacific coast, 5 from the Atlantic coast of the United States, 5 (one of which is among those occurring on the Atlantic coast) from the Gulf of Mexico, while but 3 are known from the West Indies. The majority of the species are from rather deep water, and not many are taken in shore collecting.

- a. Mouth comparatively small, maxillary less than a third length of head; mandible usually not extending backward as far as vertical from front of eye; generally a more or less distinct cross-groove on top of head behind eye; black spot on spinous dorsal usually more or less distinct.
- b. Snout not distinctly birostrate, anterior profile usually not strongly concave.
- c. Pectoral fin long, reaching past front of anal.
- d. Pectoral fin not reaching base of caudal; gillrakers moderate, 8 or 10 developed; snout not strongly emarginate; no spine on cheek bone or edge of snout; dorsal spines 10.
- e. Body not very slender, depth 5 in length; head not very small, its length 3 in body; groove across top of head behind eye, very conspicuous; interorbital area moderately concave, rather broad, about equal to diameter of eye; bones of head comparatively smooth; preocular, postocular, occipital, and nuchal spines low, depressed; temporal ridge conspicuous, without spines. Dorsal spines low, second 2.12 in head, first moderately serrate;

- base of soft dorsal equal to distance from tip of snout to tip of humeral spine; caudal fin lunate, its outer ray one-eighth to one-fourth longer than inner; pectoral fin somewhat rounded, longest ray about the fifth; free rays of pectoral expanded toward tip, with decurrent membrane; scales rather large; about 58 pores. Body and fins nearly plain, mottled with darker, but without well-defined spots except dorsal ocellus; back with 4 obscure cross-blotches; 2 or 3 oblique pale streaks across spinous dorsal. Gill-membranes dusky. Young with head rougher, pectoral fins shorter, dark spots on body more distinct..... *carolinus*
- ee. Body very slender, depth about 6 in length; groove across top of head behind eye, conspicuous; interorbital area narrow, deeply concave, its width about two-thirds the diameter of eye; bones of head very smooth, striations very weak; spines on top of head (preocular, supraocular, occipital, and nuchal) short and sharp, not depressed; temporal ridge blunt, without spine.
- f. Pectoral short, reaching little past front of anal, not one-eighth length of body. Dorsal spines very high, second 1.75 in head, first moderately serrate; soft dorsal high, its base about one-fourth longer than head; caudal truncate; free rays of pectoral a little expanded at tip; 52 pores. Body covered with roundish bronze spots of various sizes; smaller bronze spots on head; both dorsals, caudal, and pectoral fins with similar bronze spots, these especially numerous and distinct on soft dorsal..... *scitulus*
- ff. Pectoral longer, reaching past middle of anal, more than one-half body; pores 62; color rose-red, not spotted; pectoral dusky..... *roseus*
- dd. Pectoral fin very long, reaching base of caudal, rays graduated; 50 pores in lateral line; gillrakers shortish, 1 + 6; body rather stout, depth 4 in length; palatine teeth few, feeble; caudal subtruncate; second dorsal spine longest, one-half length of head; first spine strongly serrated in front; preopercular spine with a smaller one at its base; head 2.5 in length. D. X-12; A. 11. Body with 4 faint crossbands; caudal with black tip and 2 paler cross-shades; spinous dorsal with small dark spots besides large one; soft dorsal plain; pectoral clouded..... *alatus*

PRIONOTUS:

- aa. Mouth comparatively large, maxillary 2 to 2.75 in length of head, mandible extending backward to opposite eye, or nearly so; usually no distinct cross-groove on top of head; free rays of pectoral tapering, not expanded at tip; black blotch on spinous dorsal diffuse, not ocellated, involving membranes of more than two spines.
- g. Preopercular spine without a distinct smaller spine at its base in front.
- h. Pectoral fin very long, reaching at least to beyond second third of soft dorsal.
- i. Head not plane above, interorbital space more or less concave; dorsal spines 10.
- j. Supraorbital cirrus wanting.
- k. Pectoral very long, reaching in adult beyond base of dorsal and anal; interorbital space moderately concave, its width about four-fifths length of eye; no cirrus above eye; distance from supraocular spine to nuchal scales about equal to eye; supraocular and nuchal spines low; occipital spines wanting; temporal ridge sharp, ending in a blunt spine; preorbital projecting, strongly serrate; a blunt spine on each side of snout, behind serrae of preorbital; a blunt spine behind this above angle of mouth; no spine on cheek bone in adult; upper opercular spine almost obsolete; bones of head rather strongly striate, not granulate; gillrakers short; mouth moderate; scales rather large, about 52 pores. D. X-11; A. 10. First dorsal spine not much shorter than second, which is 2.2 in head; caudal very slightly concave; ventrals reaching a little past vent; head 3; depth 5. Color nearly plain-brownish, with darker clouds; no distinct spots anywhere on body or fins; pectoral marbled with paler..... *rubio*
- jj. Supraorbital cirrus present, fringed. Interorbital space very deeply concave, its width about three-fifths length of eye; a fringed cirrus above eye; distance from supraocular spine to nuchal scales about one-half eye; occipital as well as nuchal spines distinct; temporal ridge with a small spine; no spine on cheek bone; bones of head with fine, sharply defined striae, but no granulations; upper opercular spine well developed; scales moderate (about 50 pores). D. VIII-13; A. 11. First dorsal spine longest, 1.5 in head; caudal subtruncate; pectoral nearly twice as long as head, reaching nearly to last rays of dorsal; ventrals about reaching to vent; head 3 in length; depth 4.5. Color crimson, nearly plain; caudal with 2 dark cross-shades..... *ophryas*
- hh. Pectoral fin short, not reaching beyond middle of dorsal; head much smoother than in any other species, bones of head faintly striate, with small granulations; cranial spines little developed; supraocular, occipital, and temporal spines wholly wanting, there being only 3 pairs of spines on head; mouth large, maxillary 2 in head. Gillrakers short and thick in adult, slender in young, about 10 developed; interorbital space concave, rather broad, its width in adult rather more than length of eye; first dorsal spine granulated; caudal slightly lunate; pectoral subtruncate, second ray longest, as long as head in adult; scales large, 48 pores in lateral line. Head large, 2.66 in length; depth 3.66. D. X-12; A. 11. Color crimson, with darker clouds and small spots; both dorsals with dark cross-streaks; head and pectoral fin conspicuously reticulated with blackish (in adult); anal plain, whitish; free rays of pectoral unspotted..... *stearnsi*
- gg. Preopercular spine with a distinct smaller one at base; gillrakers slender.
- l. Cheek bone without distinct spine at center of radiation; edge of preorbital granular-serrate, without distinct spine, serrae about 12 in number on each side; temporal ridges roughish, but without spines; bones of head with striae coarsely granular; mouth moderate, maxillary about 2.6 in head; head not very broad, spines above, except nuchal spines, not conspicuous; gillrakers long and slender, 15 to 20 developed; head 2.75 in length; depth about 4. D. X-12; A. 11. Coloration brownish; side with a very distinct dusky bronze band below lateral line and parallel with it, this becoming broken posteriorly into a series of roundish dark spots; some dark streaks and clouds below this stripe; fins with dark clouds, soft dorsal with 2 dark blotches which extend as bars on back; head with scattered dark spots; dusky area below eye.
- m. Pectoral with its rays each crossed by fine black bars, these especially distinct toward base of fin; free rays spotted; scales comparatively small, 10 + 1 + 23 in a vertical line from last dorsal spine to vent; interorbital area broad and almost flat, its width a little more than length of eye; first dorsal spine granulated; second spine 2.75 in head; pectoral about one-half length of body..... *strigatus*

- mm.* Pectoral fin with its rays all plain-blackish; free rays plain-dusky; scales larger, 8 + 1 + 21 in a vertical line from last dorsal spine to vent; interorbital space more deeply concave, its width in adult not quite length of eye; first dorsal spine nearly smooth; second spine 3 in head; pectoral a little more than one-half body..... *evolans*
- ll.* Cheek bone with a spine (small in adult, larger in young) at center of radiation, this rarely obsolete in old examples.
- n.* Spines on bones of head moderate, not knife-like; preorbital with a series of serræ and one or more bluntish spines.
- o.* First 3 dorsal spines little if at all serrate. Back obscurely spotted; dorsal and caudal fins spotted with brown, first dorsal with a black blotch, pectoral with obscure dark spots, and margined with blue..... *punctatus*, 238
- oo.* First 3 dorsal spines more or less serrate. Color brownish-yellow; spinous dorsal with a black blotch; pectoral with 2 longitudinal broad dark areas separated and surrounded by paler..... *beanii*
- nn.* Spines on bones of head elevated, knife-like; head very large, more than two-fifths length; temporal ridge with 2 bluntish spines; bones of head very sharply striate; young with 4 sharp, knife-like spines on side of cheek and snout, in a line before preocular spine, these nearly disappearing with age; maxillary about 2.33 in head; side without dark longitudinal stripe.
- p.* Pectoral fin moderate, about one-half body in adult, 2.5 in young; gillrakers slender in young, becoming shorter and thicker with age, about 10 developed on lower part of arch; head broad, spines on its upper surface very prominent, all of them more or less compressed and knife-like, especially in young. Second dorsal spine 2.5 in head; head 2.33; depth 4.33. D. x-12; A. 11. Body brownish, much mottled with grayish and dusky, and with 3 or 4 obscure dark crossbands; head and dorsal fins with many dark spots; caudal with 2 dusky shades; free rays of pectoral spotted..... *tribulus*

238. *Prionotus punctatus* (Bloch). *Gurnard*.

Head 2.8; depth 2.8; eye 6 in head. D. x-12; A. 12; about 50 pores in lateral line. Body stout; head large; preopercular spine with smaller one at its base; pectoral reaching past middle of anal, its length not quite one-half body; gillrakers rather long and slender, about 10 developed; maxillary 2.5 in head; a bluntish spine on edge of snout behind serræ; behind this 1 or 2 smaller ones, at least in young; no spine on cheek bone; groove behind eye evident; interorbital area rather narrow, concave; preocular, supraocular, occipital, and nuchal spines rather prominent; dorsal spines high, third 2.33 in head; first spine not serrate; mouth large, maxillary 2.5 to 2.4 in head, and reaching nearly to eye; a small spine on center of radiation of cheek and one before it.

Color, nearly plain; spinous dorsal with dark clouds and without black ocelli; pectoral dark, with some round brown spots above; caudal dark-barred; a whitish area on back between dorsals. Our description is taken from two small specimens collected (probably at Tuxpan) on the east coast of Mexico, by Mr. T. Salt; from specimens in museum at Paris, the types of Cuvier & Valenciennes, and from a specimen taken by the *Albatross* at Bahia. This species is certainly the *Prionotus punctatus* of Cuvier & Valenciennes, but it may not be the species figured by Plumier, to which Bloch has given the name of *Trigla punctata*. The figure of Plumier shows a bright-red body, with many small spots of a darker red, while red spots are scattered all over the fins, except spinous dorsal and ventrals. In general form and in the armature of head, so far as this is shown in the plate, Plumier's figure most resembles the present species, but the red color suggests a possibility that some of the deep-water species may have been intended. The present species corresponds better to the figure than any other yet known. Bloch's figure of *Trigla carolina*, which has been identified with *P. punctatus*, is almost certainly *P. tribulus*. (Jordan & Evermann.)

Found in the West Indies and on the coast of South America; not known from the coasts of the United States; not common in Porto Rico, specimens (all small) being at hand from Mayaguez, station 6059, off Mayaguez in 7 fathoms, Hucares, and Isabel Segunda.

Trigla punctata Bloch, Ichthyol., pl. 353, 1793, Martiniq; on a drawing by Plumier.

Prionotus punctatus, Poey, Fauna Puerto-Riqueña, 324, 1881; Stahl, l. c., 78 and 164; Jordan & Evermann, l. c., 2169, 1898.

Family LXIV. PERISTEDIIDÆ. The Deep-water Gurnards.

Body elongate, fusiform, covered with bony plates, each of which is armed with a strong spine; head bony; each preorbital produced into a long, flat process, which projects more or less beyond mouth; mouth small, inferior, like that of a sturgeon; teeth none; lower jaw provided with barbels; gill-membranes separate, narrowly joined to the isthmus anteriorly; gillrakers slender. Dorsal fin continuous or divided. Pectoral fin short, with 2 lowermost rays detached. Ventrals 1, 5, separated by a broad, flat area. Air-bladder simple. Pyloric cæca about 10. Color generally red.

Deep-sea fishes, comprising 2 or 3 genera and about 13 species, bearing some resemblance to young sturgeons.

- a.* Barbels at angle of mouth in large tufts of fringes..... PERISTEDION, 131
- aa.* Barbels at angle of mouth minute, simple or nearly so..... VULSICULUS

Genus 131. PERISTEDION Lacépède.

Barbels large, forming large fringed tufts at angles of mouth and on lower jaw. Dorsal fins 2; characters otherwise included above.

Of the ten or a dozen known species of this genus only four occur in American waters. The first of these (*P. miniatum* Goode) is known only from the type locality, which is in the Gulf Stream off Rhode Island; the second (*P. longispathum* Goode & Bean) is known from Blake station LVIII, off Havana, in 242 fathoms (type locality), Blake stations LXII and LXIII, each off Barbados and each in 209 fathoms, and from the following Albatross stations in the Gulf of Mexico: 2397 in 280 fathoms; 2376 in 324 fathoms, and 2358 in 222 fathoms; the third (*P. platycephalum* Goode & Bean) is known only from 2 specimens, the type from Blake station LX, off Barbados, in 123 fathoms, and one other from Blake station LIX, off Barbados, in 288 fathoms; and the fourth species (*P. gracile* Goode & Bean) is also known from but two specimens, the type collected by the Albatross at station 2401 in the Gulf of Mexico in 142 fathoms, and the single specimen obtained by us in Porto Rico. A fifth species (*P. truncatum* Günther) was dredged by the *Challenger* at station 122, off Pernambuco, Brazil, in 30 or 350 fathoms, or in some intermediate depth; and a single specimen of a species in the closely related genus *Vulsiculus* Jordan & Evermann (*V. imberbis* Poey) was obtained by Poey from the stomach of a barbudo (*Polymixia lowei* Günther) taken near Havana in deep water.

The other species of *Peristedion* are known from the Mediterranean, Amboina, Japan, Molucca, etc. It will be noticed that all are deep-water species and that but little is known about any of them.

- a. Body rather robust, depth 4.5 to 5 in length to base of caudal; head about 2.5 in length; barbels long.
- b. Length of preorbital extension about 3.5 in snout; color uniform crimson *miniatum*
- bb. Length of preorbital extension 2 in snout; color red, with a black blotch near tip of pectoral and black on caudal.
longispathum
- aa. Body slender, depth 6 to 6.5 in length to base of caudal; head 3 to 3.5; barbels moderate; fins mottled or blotched.
- c. Body very slender; preorbital process 2.25 in snout; color yellowish; a pearly lateral band; back dotted; fins mottled *gracile*; 239
- cc. Body much depressed; preorbital process 3 in snout; color red, mottled and blotched *platycephalum*

239. *Peristedion gracile* Goode & Bean. Deep-water Gurnard.

(PLATE 44.)

Head (measured from base of rostral prolongation to tip of opercular spine) 3; depth 6.5; width of head 6; eye 4; snout (without spine) 2.1; rostral prolongation or spine 3.6; D. VII-19; A. I, 19; branchiostegals 8; interorbital space deeply concave, 1.2 in eye. Body slender, heavy forward; profile from tip of snout to eyes straight or slightly concave; orbital ridges strong, high; rostral spine continued backward on each side in a sharp, thin ridge, ending in a blunt spine on lower part of opercle, a similar ridge diverging from it at angle of mouth and ending in a similar spine lower down; width between bases of rostral spines 1.5 in eye; opercular spine small, length of opercle and its spine scarcely equal to eye; jaws feeble and toothless; lower jaw with an irregular group of fringed barbels on each side, longest about twice length of eye; gillrakers 26 on first arch, the longest 2 in eye. Spinous dorsal originating above opercular spine, second spine longest, 2.5 in head, or equaling length of rostral spine; longest dorsal rays 1.5 in rostral spine; origin of anal under that of soft dorsal; longest anal ray equal to those of dorsal; base of anal more than 5 times as long as interorbital width; caudal somewhat forked, the lobes rounded; pectoral long, the lower rays detached, length of fin 1.6 in head; ventral 1.75 in head; bony plates of body strong, spines sharp, their surfaces rough, the number of spines in each series, beginning with the uppermost, about 26, 32, 26, and 27.

Color in life: Whitish, with pale-rosy wash, strongest on top of head and anterior part of body; side of head and body with faint greenish blotches; similar yellowish blotches in same region; tip of lower jaw red; dorsal fin pale with a broad rosy bar on membrane about at middle of height in spinous dorsal, somewhat narrower and higher up in soft dorsal; caudal pale at base, outer third rosy-red; anal pale; ventral pale; pectoral pale, crossed by about 5 series of rosy-red spots.

So rapidly do such fishes as this change color when brought up from considerable depths that we can never be sure that the colors they exhibit when we first behold them are really those which they possess in the depths which they inhabit; in fact we may be quite sure the colors are not the same, but whether the colors are more or less intense is difficult to determine. The specimen here described had a rather faded appearance when first seen, and very soon became still paler, and it is perhaps a safe inference that the natural color, particularly the red and green, was more pronounced.

This fish attains a length of 5 or 6 inches. Until now it has been known only from the type collected by the *Albatross* in the Gulf of Mexico. We obtained a single specimen, 5.5 inches long, dredged at station 6070, in Mayaguez Harbor, January 20, in 225 fathoms, on rocky bottom.

Peristedion gracile Goode & Bean, Oceanic Ichthyology, 473, pl. CXIV, fig. 387, 1896, Gulf of Mexico, in 142 fathoms, lat. 28° 82' 30" N., long. 85° 52' 30" W., at Albatross station 249; Jordan & Evermann, l. c., 2179, 1898.

Family LXV. CEPHALACANTHIDÆ. The Flying Gurnards.

Body elongate, subquadrangular, tapering behind; head very blunt, quadrangular, its surface almost entirely bony; nasals, preorbitals, suborbitals, and bones of top of head united into a shield; nuchal part of shield on each side produced backward in a bony ridge, ending in a strong spine, which reaches past front of dorsal; interocular space deeply concave; preorbitals forming a projecting roof above jaws; preopercle produced in a very long, rough spine; cheek and opercles with small scales; opercle smaller than eye; gill-openings narrow, vertical, separated by a very broad, scaly isthmus; pseudobranchiæ large; gillrakers minute; mouth small, lower jaw included; jaws with granular teeth; no teeth on vomer or palatines; scales bony, strongly keeled; 2 serrated, knife-like appendages at base of tail; first dorsal of 4 or 5 rather high flexible spines, first 1 or 2 spines nearly free from others; an immovable spine between dorsals; anal and second dorsal short, of slender rays; caudal small, lunate; pectoral fin divided to base into 2 parts, anterior portion about as long as head, of about 6 rays, closely connected; posterior and larger portion more than twice length of head, reaching nearly to caudal in adult (*Dactylopterus*), much shorter in young (*Cephalacanthus*), these rays very slender, simple, wide apart at tip; ventral rays 1, 4, long, fins pointed, their bases close together, the inner rays shortest; air-bladder with 2 lateral parts, each with a large muscle; pyloric cæca numerous; vertebrae 9 + 13 = 22. Inhabitants of the warm seas; the adult able to move in the air like the true flying-fish, but for shorter distances. One genus.




FIG. 86.—*Cephalacanthus volitans*.

Genus 132. CEPHALACANTHUS Lacépède.

Characters of the genus included above. Two species known, the following and the East Indian *Cephalacanthus spinarella*.

240. *Cephalacanthus volitans* (Linnaeus). *Flying Robin*; *Bat-fish*; *Volador*; *Murciélago*.

Head 4.33; depth 5.5; D. II-IV, 8; A. 6; P. 28 + 6. First 2 dorsal spines free, slightly connected by membrane at base; preopercular spine reaching beyond base of pectoral, not to end of occipital spine; pectoral reaching nearly to base of caudal in adult, very much shorter in young; in the young the spines of head are much longer.

Color, greenish-olive and brown above, of varying shades; below pale, marked irregularly with dusky and bright brick-red, varying to a salmon-yellow; pectoral fin mottled with bright-blue streaks near the base and blue spots and bars toward tip, the under side glaucous-blue, edged with darker; caudal fin with about 3 brownish-red bars; coloration extremely variable.

This handsome and singular fish attains a length of 12 inches. It is found in the Atlantic Ocean, on both coasts, and is abundant on the South Atlantic and Gulf coasts. It has been recorded from Tortugas, Pensacola, St. Augustine, Cuba, and Jamaica; not seen by us in Porto Rico, but recorded from there by Professor Poey and Dr. Stahl.

Pirabê Maregrave, Hist. Brasil., IV, 162, 1648, Brazil.

Milvus cirratus Sloane, Hist. Jamaica, II, 288, Jamaica.

Trigla digitis vicenis palmatis Artedi, Genera, 44, 1738, Mediterranean, etc.

Hirundo Catesby, Nat. Hist. Carolina, II, tab. 8, Bahamas.

Trigla volitans Linnæus, Syst. Nat., ed. X, i, 302, 1758; after Artedi: "Mari Mediterraneo, Oceano, Pelago inter tropicos, in Asia ad Cap. b. Spei. Sæpi agitata evolans ex aqua."

Trigla tentabunda Walbaum, Artedi, Piscium, III, 362, 1792; after *Cataphractus* Klein, Missus, which is after Catesby, Fishes of Carolina, IV, 44, taf. 14, f. 1.

Trigla fasciata Bloch & Schneider, Syst. Ichth., 16, tab. 3, f. 1, 1801; after *Corystion* Klein, Missus, IV, 45, taf. 14, f. 2, locality not stated.

Dactylopterus piraepeda Lacépède, Hist. Nat. Poiss., III, 326, 1802, Mediterranean and almost all warm seas.

Polynemus scrradiatus Mitchill, Trans. Lit. and Phil. Soc., I, 1815, pl. 4, f. 10, New York.

Dactylopterus volitans, Poey, Fauna Puerto-Riqueña, 323, 1881; Stahl, l. c., 78 and 164, 1883.

Cephalacanthus volitans, Jordan & Evermann, l. c., 2183, 1898.

Family LXVI. GOBIIDÆ. The Gobies.

Body oblong or elongate, naked or covered with ctenoid or cycloid scales. Dentition various, the teeth generally small; premaxillaries protractile; suborbital without bony stay. Skin of head continuous with covering of eyes. Opercle unarmed; preopercle unarmed or with a short spine; pseudobranchiæ present. Gills 4, a slit behind fourth; gill-membranes united to isthmus, gill-openings thus restricted to sides. No lateral line. Dorsal fins separate or connected, spinous dorsal least developed, of 2 to 8 flexible spines, rarely wanting; anal usually with a single weak spine, similar to soft dorsal; ventral fins close together, separate or fully united, each composed of a short spine and 5 (rarely 4) soft rays, inner rays longest; ventral fins, when united, form a sucking disk, a cross-fold of skin at their base completing the cup; caudal fin convex; anal papilla prominent. No pyloric caeca; usually no air-bladder.

Carnivorous fishes, mostly of small size, living on the bottoms near the shores in warm regions. Some inhabit fresh waters, and others live indiscriminately in either fresh or salt water. Many of them bury in the mud of estuaries. Few of them are large enough to be of much value as food. The family embraces about 80 genera and nearly 600 species. The species are for the most part easily recognized, but their arrangement in genera is extremely difficult. Until the many Asiatic forms are critically studied, any definition of the American genera must be tentative only.

a. Ventral fins separate; body scaly.

OXYMETOPONTINÆ:

b. Ventral rays, I, 4.

c. Forehead bluntly rounded, without sharp keel; tongue very slender, sharp; body elongate, compressed, covered with very small scales; head short, compressed, rather broad above, mouth oblique, lower jaw projecting; teeth in few series, some of them canine-like; isthmus narrow. Dorsals separate, the first of six slender spines; soft dorsal and anal elongate; caudal lanceolate IOGLOSSUS

ELEOTRIDINÆ:

bb. Ventral rays I, 5.

d. Vomer with a broad patch of villiform teeth; gill-openings extending forward to below posterior angle of mouth, isthmus thus very narrow; skull above with conspicuous elevated ridges, one of these bounding orbit above, orbital ridges connected posteriorly above by a strong cross ridge PHLYPNUS, 133

dd. Vomer without teeth; isthmus broad; gill-openings scarcely extending forward below to posterior angle of preopercle; skull without crests.

e. Body scaly, both anteriorly and posteriorly.

f. Lower pharyngeal teeth stiff and blunt; bones with an outer series of broad flexible lamelliform appendages, which are rudimentary gill-filaments; body short and elevated DORMITATOR, 134

ff. Lower pharyngeals normal, subtriangular, teeth stiff, villiform, no lamelliform appendages; scales of moderate or small size; body oblong or elongate.

g. Body moderately robust, depth 4 to 5.5 times in length to base of caudal; scales ctenoid; cranium without distinct median keel; a small supraoccipital crest.

h. Post-temporal bones little divergent, not inserted close together, distance between their insertions greater than moderate interorbital space, or 3.8 in length of head; top of skull little gibbous; lower pharyngeals narrower than in *Eleotris*; preopercle without spine, scales very small, about 110 in a longitudinal series. Vertebrae 11 + 13; teeth moderate, outer series on lower jaw enlarged GUAVINA, 135

- hh. Post-temporal bones very strongly divergent, their insertions close together, distance between them about two-thirds the narrow interorbital space, and less than one-seventh length of head; top of skull somewhat elevated and declivous; interorbital area somewhat convex transversely; lower pharyngeals rather broad, teeth bluntish preopercle with partly concealed spine directed downward and forward at its angle; scales moderate, 45 to 60 in a longitudinal series ELEOTRIS, 136
- gg. Body very slender, elongate, depth 8 to 9 times in length to base of caudal; scales very small, cycloid.
- i. Preopercle without spine; caudal without many accessory rays at base; post-temporal bones short, strongly divergent, distance between their insertions about equal to narrow interorbital space, or about one-sixth length of head; top of head with a strong median keel, which is highest on occipital region; no supraoccipital crest, mouth very oblique; teeth small EROTELIS
- aa. Ventral fins united.
- j. Dorsal fins separate, free from caudal.

SCYDINÆ:

- k. Ventral disk short, adnate to belly; body subcylindrical, covered with ctenoid scales; lips very thick; upper teeth mostly small and movable, lower fixed; dorsal spines 6. Teeth simple; no canines in front of lower jaw.

SCYDIUM, 137

GOBIINÆ:

- kk. Ventral disk free from the belly.
- l. Dorsal spines 4 to 8; eyes well developed.
- m. Teeth emarginate, uniserial, those of lower jaw nearly horizontal; dorsal spines 6; scales large, ctenoid; gill-openings moderate EVORTHODUS
- mm. Teeth simple.
- n. Body scaly, more or less.
- o. Maxillary normal, not prolonged behind rictus; skull of usual gobioid form, comparatively short and abruptly broadened behind orbits; occiput depressed; supraoccipital and temporal ridges continuous.
- p. Dorsal spines 6; scales evidently ctenoid; head naked (nape scaly as usual).
- q. Interorbital area anteriorly elevated, with a large foramen-like depression in front of eye; body short, compressed, formed much as in *Dormitator*; nape with a fleshy crest; scales large. Vertebrae 11 + 15. LOPHOGOBIOUS
- qq. Interorbital area not elevated in front; body more elongate; no fleshy nuchal crest; isthmus broad.
- r. Inner edge of the shoulder-girdle without fleshy cirri or papillæ; cranium anteriorly short; interorbital space narrower, grooved, with a low median ridge or none; median crest on cranium low.
- s. Body scaly anteriorly and posteriorly (sometimes a naked strip on back or belly). Vertebrae 12 + 16 to 10 + 15. GOBIUS, 138
- ss. Body entirely naked anteriorly, posterior half scaled; scales moderate or small. GARMANNIA
- rr. Inner edge of shoulder-girdle with 2 or 3 conspicuous dermal flaps; preorbital region very long; premaxillary and maxillary strong; interorbital groove with a conspicuous median crest; scales rather small (45 to 70). AWAOUS, 139
- pp. Dorsal spines 7 or 8 (very rarely 6, especially in *Eucyclogobius*).
- t. Scales large, ctenoid; shoulder-girdle without dermal flaps. Sides of head scaled; soft dorsal and anal rather short, of 11 to 14 rays each; deep-water species BOLLMANNIA, 140
- tt. Scales very small, cycloid or nearly so. Inner edge of shoulder-girdle without fleshy processes; head naked; body more or less compressed; mouth very oblique; teeth strong; interorbital groove with or without a median ridge. Vertebrae 11 + 15 or 16; soft dorsal and anal long, of 15 to 17 rays each. Body chiefly scaly, anteriorly as well as posteriorly MICROGOBIUS, 141
- nn. Body and head entirely naked. Dorsal spines 7 (rarely 6).
- u. Chin without barbels; mouth small, little oblique; body robust, soft dorsal and anal short GOBIOSOMA
- uu. Chin with a fringe of short barbels; mouth terminal, oblique; soft dorsal and anal very short BARBULIFER
- jj. Dorsal fin continuous, soft part and anal joined to base of caudal; eye minute; body elongate; scales minute or wanting; mouth very oblique, lower jaw projecting; gill-openings moderate.
- v. Dorsal rays vi, 16 to 23; anal rays 17 to 23. Teeth in a band, those of outer series being very strong; scales present.
- w. Body entirely scaled GOBIOIDES, 142
- ww. Anterior part of body naked CAYENNA

Genus 133. PHILYPNUS Cuvier & Valenciennes. Guavinas.

Body elongate, terete anteriorly, compressed behind. Head elongate, depressed above. Mouth large; lower jaw considerably projecting; teeth in jaws rather small, slender, recurved, outer scarcely enlarged; teeth on vomer villiform, in a broad, crescent-shaped patch; gill-openings extending forward to below posterior angle of mouth, isthmus very narrow. Scales moderate, ctenoid, covering most of head, 55 to 66 in a longitudinal series. Dorsal with 6 spines and 9 or 10 rays; anal rays i, 9 or 10; ventrals separate. No preopercular spine; insertion of post-temporals almost midway between occipital crest and edge of skull; parietals with a crest running from insertion of post-temporal forward to just behind eye, where they are connected by a thin, high, transverse crest; supraocular with a short, high crest, extending from above front of eye back to posterior edge of orbit, thence extending outward parallel with transverse crest, leaving a deep groove between them; bony projections before and behind eye prominent. Vertebrae 12 + 13 = 25; lower pharyngeals triangular, with slender teeth.

The fishes of this genus are the largest of the gobies, some reaching a length of 2 or 3 feet and being valued as food. Found in tropical rivers.

241. *Philypnus dormitor* (Lacépède). "*Guavina*."

Head 3; depth 5; eye 9.75; snout 3; maxillary 2.2; mandible 2.1; interorbital width 3.8; pre-orbital 9.5; D. vi-12; A. i, 10; scales about 61-23. Body long, subterete; head long, broad, depressed; snout long, lower jaw strongly projecting, the point a broad angle; mouth large, somewhat oblique, maxillary very long, reaching posterior border of orbit; eye nearly on level with top of head; cheeks full; teeth in each jaw in broad cardiform bands; vomerine teeth in a broad crescent-shaped patch; gillrakers 4 + 11, very short, covered with fine prickles or spines, especially on their posterior surface. Fins all large; dorsal fins separated by a space less than half diameter of orbit, spines flexible, longest about equal to snout and eye; last rays of second dorsal longest, somewhat longer than snout and eye, about 2 in head; caudal fin rounded, convex, the middle rays about 1.5 in head; last anal ray longest, a little shorter than last dorsal rays; pectoral broad, broadly pointed, with about 17 rays, middle ones longest, about 1.8 in head, reaching beyond tips of ventrals; ventrals 1, 5, about 2.2 in head, their tips reaching less than two-thirds distance to vent.

Color in life: Dark-yellowish or olive-brownish, back darkest, belly most yellow; sides with extensions of the dark of back; head dark, with a few small dark specks; arrangement of colors presenting a mottled appearance as on body; fins all mottled with brown and yellowish, no dark border to spinous dorsal; base of pectoral dark; ventrals paler.

This large goby is everywhere common in the fresh-water streams of the West Indies and the Atlantic shores of Mexico, Central America, and Surinam. It seems to be an abundant fish in all the larger streams of Porto Rico, specimens having been obtained by us from the Rio Loiza and Rio de Caguaitas near Caguas, and from the Rio Bayamon at Bayamon and near Palo Seco. It has been




Fig. 87.—*Philypnus dormitor*.

recorded from Havana, Martinique, Jamaica, Mexico, and Nicaragua. It reaches a length of 2 feet or more, and is one of the most important fresh-water food-fishes in Porto Rico and elsewhere in the West Indies. The examples obtained by us are 4 to 17 inches long.

Guavina, Parra, Descr. Dif. Piezas, Hist. Nat. Cuba, tab. 39, fig. 1, 1787, Havana.

Gobiomorus dormitor Lacépède, Hist. Nat. Poiss., II, 599, 1798, Martinique; from a drawing by Plumier.

Platycephalus dormitor Bloch, Syst. Ichth., pl. 12, 1790, Martinique; after Lacépède.

Batrachus guavina Bloch & Schneider, Syst. Ichth., 44, 1801; based on *Guavina* of Parra.

Eleotris longiceps Günther, Proc. Zool. Soc. Lond. 1864, 151, Nicaragua.

Eleotris dormitatrix Cuvier, Règne Animal, ed. II, vol. 2, 246, 1829, Antilles.

Philypnus dormitor, Poey, Fauna Puerto-Riqueña, 339, 1881; Stahl, l. c., 79 and 165, 1883.

Philypnus dormitor, Jordan & Evermann, l. c., 2194, 1898.

Genus 134. DORMITATOR Gill. Puñecas.

Body short, robust; head broad and flat above; mouth little oblique; maxillary reaching to anterior margin of orbit; lower jaw little projecting; no teeth on vomer; lower pharyngeal teeth stiff and blunt, the bones with an external series broad, flexible, lamelliform, these being rudimentary gill-filaments; scales large, ctenoid, 30 to 33 in a longitudinal series; skull much as in *Eleotris*; D. vii-1, 8; A. i, 9 or 10; no spine on preopercle; post-temporals inserted midway between occipital crest and edge of skull; supraoccipital crest low.

242. *Dormitator maculatus* (Bloch). *Masaguan; Mapiro.*

(PLATE 45.)

Head 3.5; depth 3.5; eye 5.75; snout 4.5; maxillary 3; mandible 3.2; interorbital 2; preorbital 6.5; scales 33, -12; D. VIII-9; A. I, 9; longest dorsal spine 1.9; longest dorsal ray 1.1; longest anal ray 1.2; pectoral 1.2; ventral 1.25; caudal 1. Body short and stout, heavy forward; head broad, flat above, interorbital space wide; caudal peduncle long, rather compressed, length 1.4 in head; profile gently and regularly arched from tip of snout to origin of dorsal, from which point body tapers to caudal peduncle. Mouth small, considerably oblique, upper lip on level with pupil; maxillary reaching vertical at front of orbit; lower jaw scarcely projecting; teeth on jaws short and weak, in villiform patches; no teeth on vomer. Fins all large; interspace between dorsals very narrow, less than orbit; soft dorsal and anal, when depressed, reaching beyond base of caudal, which is rounded. Scales large, smaller on head and belly, about 15 series from base of ventral to vent, 14 series across breast between bases of pectorals, and about 25 from tip of snout to origin of dorsal fin; origin of dorsal fin midway between tip of snout and posterior base of soft dorsal.

Color in life: Pale-greenish on head and back, lower sides and belly pale-bluish; sides of head and back mottled with darker; a narrow dark line from eye downward and forward to mouth, and 2 or 3 similar but broader bands from eye to posterior edge of opercle, these bands more or less broken; dorsal fins brownish, barred with narrower white bands, each with a narrow brick-red edge; anal banded with light brick-red and narrower white interspaces; edge of fin pale, with a subterminal dark bar; caudal, ventral, and pectoral pale; iris brick-red. In alcohol this fish becomes quite dark, almost black, the bright colors changing completely.

This description is from numerous specimens 6 to 8 inches long, but the species reaches a length of 1 to 2 feet and is a valued and important food-fish. It is found on both coasts of tropical America, ranging from South Carolina through the West Indies to Para, and from Cape San Lucas to Panama; usually abundant. On the coast of Florida it has been taken at Hillsboro River, Indian River Inlet, and on the Pensacola Snapper Banks. No adults were seen by us in Porto Rico, but we collected one young specimen at Mayaguez, and on February 15 we obtained 85 near Hucares from a small, shallow lagoon bordered with mangrove bushes, filled with decayed vegetation, and with the water strongly colored. These specimens varied in length from 3 to 8 inches.

Sciæna maculata Bloch, Syst. Ichth., pl. 299, fig. 2, 1790, West Indies.

Eleotris mugiloides Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 226, 1837, Martinique and Surinam.

Eleotris sima Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 232, 1837, Vera Cruz.

Eleotris latifrons Richardson, Voy. Sulphur, Fishes, 57, pl. 35, figs. 4 and 5, 1837, locality unknown, probably Pacific coast of Central America.

?*Eleotris grandisquama* Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 229, 1837, America.

Eleotris somnolentus Girard, Proc. Ac. Nat. Sci. Phila. 1858, 169, near mouth of Rio Grande, Texas.

Eleotris omocyanus Poey, Memorias, II, 269, Havana.

Dormitator microphthalmus Gill, Proc. Ac. Nat. Sci. Phila. 1863, 170, Panama.

Dormitator gundlachi Poey, Synopsis, 396, 1868, Cuba.

Dormitator lineatus Gill, Proc. Ac. Nat. Sci. Phila. 1863, 271, Savannah, Georgia.

Dormitator mugiloides, Poey, Fauna Puerto-Riqueña, 339, 1881; Stahl, l. c., 79 and 165, 1883.

Dormitator maculatus, Jordan & Evermann, l. c., 2196, 1898.

Genus 135. *GUAVINA* Bleeker. *Morons.*

This genus is allied to *Eleotris*, differing in having the post-temporal bones little divergent, not inserted close together, distance between their insertions greater than the moderate interorbital space, or 3.8 in length of head; top of skull little gibbous; lower pharyngeals narrower than in *Eleotris*; preopercle without spine; scales very small, ctenoid, about 110 in a longitudinal series. Vertebrae II + 13; teeth moderate, the outer series on lower jaw enlarged.

Inhabitants of the fresh waters of the West Indies and Brazil. Two species known—*Guavina brasiliensis* (Sauvage), from Bahia, and the following.

243. *Guavina guavina* (Cuvier & Valenciennes). "*Moron.*"

Head 3.5; depth 4.5; eye 6.5; snout 4; maxillary 2.25 to 2.5; mandible 2.5; interorbital 3 to 3.25; preorbital 7.2; scales about 100, -38; D. VII-12, the longest spine 2.5 to 3 in head, the longest soft ray about 2; A. I, 10, longest ray about 2 in head; pectoral 1.5; caudal 1.25 to 1.5. Body stoutish, oblong, heavy forward; head heavy, broad; mouth oblique, large, maxillaries reaching middle of

eye, lower jaw slightly projecting; isthmus very broad, gill-opening not reaching farther forward than vertical opercle; preopercle smooth with concealed spine; caudal peduncle compressed, least width 3.75 in its least depth, which is 2 in head; teeth in broad bands, outer ones of lower jaw somewhat enlarged; dorsal outline rising gently from snout to origin of spinous dorsal; ventral line relatively straight; scales very small, those on head embedded and cycloid, those on back, belly, and anterior part of sides cycloid, on posterior part of body ctenoid; lower portion of preocular and cheek naked, upper portions with small embedded scales. Fins moderate; origin of spinous dorsal nearer tip of snout than posterior edge of soft dorsal by a distance equal to diameter of orbit; distance between spinous soft dorsal slightly greater than diameter of orbit; dorsal spines flexible; dorsal and anal rays when depressed barely reaching rudimentary caudal rays; caudal fin regularly rounded; tips of ventrals reaching halfway to origin of anal fin; tips of pectorals extending slightly beyond tips of ventrals.

Color, dark, marbled with darker and lighter; under parts dirty-whitish; fins dark like body, bordered with white, especially the dorsals and anal.

The moron reaches a foot in length and is a good food-fish. It is found on the east coast of tropical America from Cuba to Rio Janeiro, in fresh and brackish water, and is generally common. Only two examples (6.5 and 8.5 inches long) were seen in Porto Rico by us; they were obtained in the San Juan market January 15, and probably came from the mouth of the Bayamon River.

Eleotris guavina Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 223, 1837, Martinique.

Guavina guavina, Jordan & Evermann, Fishes of North and Middle America, 2198, 1898.

Genus 136. ELEOTRIS (Gronow) Bloch & Schneider.

Body long and low, compressed behind. Head long, low, flattened above, without spines or crests, almost everywhere scaly. Mouth large, oblique, lower jaw projecting. Lower pharyngeals rather broad, teeth small, bluntish. Preopercle with a small concealed spine below, its tip hooked forward. Branchiostegals unarmed. Eyes small, high, anterior; isthmus broad. Post-temporal bones very strongly divergent, their insertions close together, the distance between them about two-thirds the narrow interorbital space, and less than one-seventh length of head; top of skull somewhat elevated and declivous; interorbital area slightly convex transversely; the dorsal fins well separated, first of 6 or 7 flexible spines; ventrals separate. Scales moderate, ctenoid, 45 to 62 in longitudinal series; vertebræ (*pisonis*) 11 + 15.

Found in tropical seas, entering fresh waters.

- a. Teeth subequal, those of inner or outer series enlarged.
- b. Cheek entirely scaled.
- c. Teeth of inner series of each jaw enlarged.
- d. Scales in a median series 40 to 51, in a cross series 12 to 20.
- e. Eye large, 5 to 6 in head; scales 40 to 44, -12 to 14 *amblyopsis*
- ee. Eye small, 8 in head; scales 51, -20 *abacurus*
- dd. Scales in a median series 57 to 66; in a cross series 18 to 24 *pisonis*. 244
- bb. Lower half of cheek naked; scales 61 *perniger*

244. *Eleotris pisonis* (Gmelin). "Moron."

Head 2.9; depth 3.75; eye 7; snout 5; maxillary 2.75; mandible 2.75; interorbital 3.6; preorbital 10; pectoral 1.4; ventral 2; caudal 1.4; D. vi-9, the longest spine 3 in head, the longest ray 2.5; A. i, 8, the longest ray 2.3; scales 57 to 63, -19 to 22; vertebræ 11 + 15. Body stout, not compressed; head broad, depressed, tapering forward; mouth large, oblique, maxillary reaching posterior border of pupil; eyes high up; isthmus moderately broad, gill-openings extending forward somewhat beyond angle of preopercle; preopercle with a strong concealed spine at angle, directed forward; no teeth on vomer or palatines; teeth on jaws in wide villiform bands, outer series somewhat enlarged; lower pharyngeals broad, triangular, without fringe, teeth bluntish; branchiostegals 6; gillrakers 2 + 9, very short; caudal peduncle compressed, its least width one-third its depth, which is 2 in head. Fins moderate; origin of spinous dorsal about midway between tip of snout and tip of dorsal rays; space separating dorsals scarcely more than half orbit; dorsal and anal rays not reaching base of caudal; ventrals inserted somewhat behind insertion of pectorals, their tips reaching halfway to origin of anal; pectoral reaching beyond ventrals. Scales small, crowded anteriorly, embedded on nape, head, and cheeks, ctenoid only on sides.

Color, dark-brown or blackish; sides with faint narrow dark lines alternating with narrow light ones; fins with dark wavy lines.

The range of this species is from Florida to Rio Janeiro; it is recorded from Pensacola, Rio Almendares (Cuba), Santo Domingo, Martinique, Surinam, and Rio Janeiro, and was found by us to be common in fresh and brackish waters of Porto Rico. In the Bayamon River 13 specimens were obtained January 12, varying in length from 3.5 to 6.25 inches. There is a single specimen, 1.5 inches long, from Mayaguez, January 20, and 3 small ones, from 1.31 to 2.06 inches long, from Arroyo. This species is known in Porto Rico only as "moron." It does not seem to be much used as food. It is apparently most common in the lower portions of rivers, and probably does not run to the headwaters.

Amore picuma, Marcgrave & Piso, Hist. Brasil., IV, 166, 1648, Brazil.

Eleotris capite plagioplateo, Gronow, Mus. Ichth., II, 168, 1757; after Marcgrave & Piso.

Gobius pisonis Gmelin, Syst. Nat., 1206, 1788; based on *Eleotris* of Gronow.

Eleotris gyrynus Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 220, pl. 356, 1837, Martinique, Santo Domingo and Surinam; Poey, Fauna Puerto-Riqueña, 339, 1881; Stahl, l. c., 79 and 165, 1883.

Eleotris (Culius) belizianus, Sauvage, Bull. Soc. Philom. Paris 1879, 55, Belize and Cayenne.

Eleotris pisonis, Jordan & Evermann, l. c., 2200, 1898.




FIG. 88.—*Eleotris pisonis*.

Genus 137. SICYDIUM Cuvier & Valenciennes.

Body subcylindrical, covered with rather small ctenoid scales; head oblong and broad, with cleft of mouth nearly horizontal; upper jaw prominent; snout obtusely rounded; lips very thick, lower jaw with a series of numerous slender horizontal teeth, of which sometimes only the extremities are visible; upper jaw with a single uniform series of numerous movable small teeth attached by ligament to edge of maxillary; behind this outer visible series lie numerous other parallel series of young teeth hidden in the gum, which succeed the former as they become worn out or broken; lower jaw with a series of widely-set conical teeth; teeth all simple, slender, distal half bent inward nearly at a right angle; eyes of moderate size; 2 dorsal fins, anterior with 6 (5 or 7) flexible spines; caudal quite free; ventrals united into a short cup-shaped disk; gill-openings of moderate width; 4 branchiostegals.

Found in the streams of the West Indies. Few species.

- a. Body usually well scaled.
- b. Scales large, about 56 in longitudinal series *punctatum*
- bb. Scales moderate, about 68..... *antillarum*
- bbb. Scales small, about 84.
- c. Body densely scaled; no dark vertical bars on body..... *caguitæ*, 245
- cc. Body less densely scaled; squamation sometimes quite incomplete; about 7 more or less distinct vertical bars on body..... *plumieri*, 246
- aa. Body nearly naked..... *vincente*

245. *Sicydium caguitæ* Evermann & Marsh.

Head 4.4; depth 4.8; eye 5.75; snout 2.5; maxillary 2; mandible 2.75; interorbital width 3; preorbital 3.5; D. VI-1, 10; A. I, 9; scales 83,-25; longest dorsal spine 1.5 in head, longest ray 2; longest anal spine 2 in head, longest ray 2; pectoral 1.1; ventral disc 1.75; caudal 1. Body rather stout, heavy forward; head large, broad; mouth large, its width 1.5 in head; lips very thick; maxillary not greatly produced; teeth simple, flexible; a median cleft in upper lip; pectoral somewhat shorter than head; dorsal spines without filaments, longest about 1.5 in depth of body; space between dorsals about equal to orbit; soft rays of dorsal and anal scarcely reaching base of caudal; ventrals united,

forming a cup-shaped disk, only about two-fifths posterior edge free from belly; caudal rounded. Scales very small, ctenoid, densely covering entire body except a broad strip on belly; posterior portion of nape with very fine scales; entire head naked.

Color, dark-brown or olivaceous on head, sides, and back; under parts pale; fins all pale, anal with a narrow darkish margin; caudal somewhat dark; no dark vertical bars on body and none at base of pectoral; no H-shaped figure at base of caudal.

This species is close to *S. plumieri*, from which it differs chiefly in color, the more complete squamation, the shorter pectoral, and the nonfilamentous character of dorsal spines. Known only from the type, 3.63 inches long, from the Rio de Caguas at Caguas, Porto Rico.

Sicydium caguizæ Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 355, Rio de Caguas, Caguas, Porto Rico.




FIG. 89.—*Sicydium caguizæ*.

246. *Sicydium plumieri* (Bloch). *Sirajo*.

Head 4 to 4.6; depth 4.5; eye 6 to 7 in head, 2 to 3 in interorbital width; D. VI-I, 10; A. I, 10; scales 84. Teeth in upper jaw long, slender, bent inward at right angles, only tips protruding from gums. Front teeth of lower jaw not larger than those behind; a single row of inconspicuous papillæ on gum beneath upper lip, a large median papilla above maxillary suture; a median cleft in upper lip. Pectoral longer than head; third, fourth, and fifth dorsal spines produced into long ribbons, the fourth, which is longest, being 2 to 3 times height of body. Body usually covered with small scales, reduced in size on neck and belly; frequently almost naked, scales present only on posterior part of body. Caudal deeply emarginate.

Color, olive or violet-brown, with about 7 more or less distinct dark vertical bars; a dark bar at base of pectoral; dorsal with irregular dark markings; anal fin with a dark marginal band, sometimes edged with white; an H-shaped figure on base of caudal fin, and a black bar on its posterior half.

An inhabitant of the fresh waters of the West Indies. Not obtained by us in Porto Rico, but recorded by Professor Poey.

Gobius plumieri Bloch, Ichth., 125, pl. 178, fig. 3, 1786, Martinique; on a drawing by Plumier.

Sicydium siragus Poey, Memorias, II, 278, 1861, Santiago de Cuba.

Sicydium plumieri Poey, Fauna Puerto-Riqueña, 338, 1881; Jordan & Evermann, l. c., 2206, 1898.

Genus 138. GOBIUS (Artedi) Linnæus. Gobies.

Body oblong or elongate, compressed behind. Head oblong, more or less depressed. Eyes high, anterior, close together; opercles unarmed. Mouth moderate. Teeth on jaws only, conical, in several series, those in outer row enlarged; no canines. Isthmus broad. Shoulder-girdle without fleshy flaps or papillæ. Skull depressed, abruptly widened behind eyes and without distinct median keel. Scales moderate, ctenoid, permanently covering body; cheek usually naked; belly generally scaly. Dorsal with 6 rather weak spines; pectoral well developed, upper rays sometimes very slender and silky; ventrals completely united, not adnate to belly; caudal fin usually obtuse.

The genus *Gobius*, as here understood, comprises a very large number of species more or less closely related to the European type of the genus, *Gobius niger*, and its American relative, *Gobius soporator*. An examination of skulls or skeletons of numerous European and American species shows a remarkable uniformity in most respects. The general form and structure of the cranium is the same

in all, the only differences being very minor ones in the height of certain crests. *Gobius oceanicus* seems most aberrant, but is difficult to separate generically on account of intermediate forms. Probably several of the many genera indicated by Bleeker will prove valid, but only a thorough study of skeletons can establish them. It is not unlikely that *Ctenogobius*, to which group most of our species belong, may be separable from *Gobius*.

GOBIUS:

- a. Upper rays of pectoral fin silk-like; i. e., short and very slender and flexible, free for nearly their whole length.
 b. Body robust, compressed posteriorly. *soporator*, 247

CTENOGOBIUS:

- aa. Upper rays of pectoral normal, not silk-like, similar to others.
 c. Scales large, 25 to 33. Color in life olivaceous, more or less spotted, never red.
 d. Dorsal soft rays 12 to 14; vertex and nape with a slight median fold of skin. Body long, not much compressed; head 3.5; eye 3 in head; a dark spot on first dorsal. *eigenmanni*
 dd. Dorsal soft rays 10 to 12.
 e. Caudal with 2 spots at its base; jaws unequal, lower slightly produced; body robust, compressed behind, depth 5 in total length; head 4.33; eye longer than snout, 3.5 in head; maxillary reaching pupil; teeth in a band, the outer enlarged and distant, the inner enlarged and bent backward. Brownish; a faint blue spot on each scale; six spots along middle of back; similar spots on scapular region and middle of side; a dark spot above opercle; blue dots on head; a straight blue line crossing cheek above and continued on opercle; dorsals faintly spotted. D. VI-10; A. 10. Scales 25, 7. *glaucofrænum*
 ee. Caudal plain or with but a single spot at its base.
 f. Dorsal spines low, highest little longer than head.
 g. Region from nape to dorsal entirely scaled.
 h. Pores on preopercle not very conspicuous; no canine teeth. Body subfusiform, little compressed; depth 4.5 in length; head blunt, 4 in length, rounded in profile; eye equal to snout, 4 in head. Mouth small, horizontal, lower jaw included; maxillary 3 in head, reaching to below eye. Teeth small, in bands in both jaws, the outer enlarged, those of upper jaw very slender. Scales large, ctenoid, those of nape and belly little reduced. Longest dorsal spine shorter than head. Caudal scarcely pointed, about as long as head. Color, whitish-gray, middle of side with 4 or 5 dark blotches, from each of which a narrow dark bar extends downward and forward; a large black blotch above pectoral, obsolete in female; a small black spot at base of caudal; a dark mark below eye; vertical fins barred. D. VI-12; A. 11 or 12. Scales 33. *stigmaturus*
 hh. Pores on preopercle very conspicuous; lower jaw with small canines. D. VI-1, 9; A. I, 9. *quadriporus*
 gg. Region between nape and dorsal with a narrow naked median strip. Body moderately elongate, subfusiform, depth 5.33 in length. Head large, not so blunt as in *G. bolcosoma*, 3.4 to 3.66 in length; anterior profile gently decurved; snout 3.33 to 3.5 in head; eye 4; mouth large, slightly oblique; maxillary extending to front of pupil, 2.33 in head. Teeth small, slender and curved, in moderate bands. Scales moderate, ctenoid, those in front much reduced in size; breast naked. Longest dorsal spine 1.5 in head. Caudal as long as head, somewhat pointed. Olivaceous, mottled with gray; about 5 rounded dark blotches along middle of side, last forming a spot at base of caudal; no dark spot on side of nape; some dark marks on head; vertical fins barred. D. VI-12; A. 13. Scales 33 to 35. *shufeldti*
 ggg. Region between nape and dorsal entirely naked.
 i. Highest rays of second dorsal little more than half head, none of them reaching base of caudal.
 j. Profile much decurved, skull rounded behind, without distinct median ridge; mouth horizontal. Body elongate, deepest below front of dorsal, tapering regularly backward, greatest depth 5.5 in length. Head short, blunt, profile anteriorly abruptly decurved, cheek somewhat swollen. Length of head 3.5 in body. Snout about equal to eye, 3.66 in head. Mouth horizontal, maxillary reaching to below pupil (in male); lower jaw included. Teeth in each jaw in a band, outer row of upper jaw large, recurved. Scales large, ctenoid, somewhat reduced anteriorly. Nape, breast, and belly naked. Dorsal spines about two-thirds of head. Caudal pointed, 2.66 to 3.5 in body. Color olivaceous, with numerous dark reticulations on back; 5 black spots along side, the last forming a spot on base of caudal, sometimes with V-shaped dark bars extending from them to dorsal; breast and sides of belly with numerous dark specks in male; a dark line between eyes; a dark line from eye to middle of premaxillary, some dark spots below eye, sometimes forming bars, sometimes a stripe; a large oblique spot above opercle, continued on opercle; a black spot at base of pectoral; dorsals and caudal barred, anal uniform dusky, ventrals and pectoral black in male, white in female. D. VI-11; A. 10 to 12. Scales 25 to 30. *bolcosoma*
 jj. Profile moderately decurved; eye longer than snout, 3.75 in head. Color yellowish, oblong dark blotches on middle of side; dorsal and caudal barred. Head 4; depth 6. D. VI-12; A. 10. *fasciatus*
 ii. Highest rays of second dorsal as long as head, the last reaching base of caudal. Body elongate, back not arched; depth 6 in length; head 4, not compressed, cheek tumid. Profile abruptly decurved, the snout 3.33 in head. Mouth large, nearly horizontal, maxillary reaching posterior edge of eye in males, middle of eye in females. Teeth in narrow bands in each jaw, outer somewhat enlarged, outer in some (males?) much enlarged above and recurved, enlarged teeth fixed, others movable. Scales large, ctenoid, reduced anteriorly; belly naked. Dorsal spines little filamentous, longest about equal to head; caudal 2.25 to 3 in body. Males dark-olive, with 4 oblong dark blotches along middle of side; a dark caudal spot; a black blotch larger than eye on each side of shoulder; dorsal spotted; caudal reddish above, dusky below; females with 5 oblong dark blotches on side the last on base of caudal; from each of middle blotches a V-shaped bar runs to back; a black shoulder blotch; a dark bar from eye to mouth; ventrals pale, with 2 dark streaks. D. VI-11; A. 12. Scales 30 (27 to 33) at least in males. *encæmus*

- ff.* Dorsal spines high, highest reaching past middle of second dorsal. Nape scaly. Body elongate, moderately compressed.
- k.* Depth 5 to 6 in length; profile little dcurved, skull flattish behind, much broader than in *Gobius bolcosoma*, with an evident median ridge; mouth very oblique, much larger than in *G. bolcosoma*; lower jaw thin and flat. Back slightly arched. Body a little deeper and rather less compressed than in *G. encaomus*, depth 5 to 6 in length. Head 4. Anterior profile moderately dcurved. Eye 3.33 in head. Mouth large, oblique; maxillary reaching to below pupil in both sexes. Teeth above uniserial, some of them enlarged and recurved; lower teeth in a narrow band; males sometimes with hindermost of outer series a strong, exserted, recurved canine; belly naked. Longest dorsal spine two-thirds head in females, elevated in males; soft dorsal elevated in males; caudal 3.5 in body. Color, light-greenish, sides of male with 5 or 6 narrow, straight, rather sharply defined whitish or yellowish crossbars, regularly placed; 4 dark bars, 3 below eye and 1 on opercle; a small dark spot behind and above opercle; vertical fins barred; female with a row of irregular dark spots connected by a dusky streak, and with pale crossbars obsolete. D. VI-12; A. 13. Scales 27..... *stigmaticus*
- kk.* Depth 4.66 in length; profile very obtuse anteriorly; maxillary extending beyond pupil, 2.6 in head. Teeth strong, uniserial, 4 shortish canines in lower jaw behind other teeth; upper teeth largest. D. VI-11; A. 10. *lyricus*, 248
- kkk.* Depth 4 in length; teeth short and thick, uniserial; yellowish, much mottled and blotched. D. VI-11; A. 11. Scales 30..... *garmani*
- cc.* Scales moderate or small, 40 to 90.
- l.* Soft dorsal and anal short, each of 10 to 14 rays; body more or less elongate.
- EUCTENOGOBIIUS:**
- m.* Caudal rounded, not much longer than head.
- n.* Scales 40; dorsal with 9 soft rays only; anal with 9; depth 6.5 in total length; head broad, flattish; snout short, dcurved; eye 4.25 in head, 1.33 in interorbital area, longer than snout; maxillary extending to below middle of eye. Some of dorsal spines produced in filaments, the third 1.5 times depth of body; caudal short, rounded. Two rows of ill-defined blotches on upper half of body; 2 rows of brownish spots on second dorsal, upper strongly marked..... *poeyi*
- nn.* Scales 50; dorsal and anal with 10 soft rays each; profile very oblique. Color dark-brown..... *badius*
- GOBIONELLUS:**
- mm.* Caudal lanceolate, much longer than head; lower jaw thin; usually a green spot on roof of mouth in life.
- o.* Body rather deep, depth about 5 in length.
- p.* Scales rather large, 39 to 42; body moderately elongate, compressed; depth 5.25; head 4. Head not compressed, cheek tumid, snout short, abruptly dcurved; mouth large, little oblique, jaws equal, maxillary 2.33 in head, reaching to below pupil; eye 5 in head; teeth above large, unequal, uniserial, some of them fixed, those below small, in a band. Scales anteriorly cycloid, becoming larger posteriorly, and etenoid; dorsal spines scarcely filamentous, none of them as high as body; caudal 2.25 in body. Light-olive, with dark-olive blotches; body and head with many conspicuous round spots of cream color, each surrounded by a dusky ring, these most distinct on head, all smaller than pupil; snout with dusky streaks; dorsals and caudal sharply barred; anal and ventrals dusky (in male); a small round spot at base of caudal. D. VI-11; A. 11..... *smaragdus*
- pp.* Scales comparatively small (53). Body elongate, compressed behind; head a little compressed, 3.75 in length; depth 5; eye 3.5 in head, shorter than rounded snout; maxillary reaching to below middle of eye; teeth small, outer a little enlarged; dorsal spines all shorter than head, not filamentous. Nape scaly, its scales much reduced in size; scales etenoid. Two violet stripes from eye to mouth; 8 or 9 violet bars on side; 3 or 4 bars on caudal; second dorsal spotted. D. VI-12; A. 11 or 12..... *strigatus*
- oo.* Body elongate, depth 6.5 to 9 in length; head 4.5; caudal very long.
- q.* Scales very small (60 to 90); caudal more than twice as long as head in adult. Body compressed, extremely elongate, depth 6 to 9 in length; head higher than wide, short, compressed, 4.5 to 5 in length; mouth wide, oblique; maxillary in adult reaching to below posterior border of eye. Lower jaw very thin and flat; teeth in both jaws small, subequal, those in upper jaw in a single series, those of lower in a narrow band; outer teeth somewhat movable. Scales anteriorly small, cycloid, embedded, those behind larger and etenoid; a few scales on upper anterior corner of opercle; dorsal fins high, some spines filamentous, longer than head. Caudal very long, filamentous, 2 to 2.66 in body. Light-olive; fins dusky in male; a round, black spot on side, a little larger than eye, below spinous dorsal; first dorsal spine with 2 or 3 black spots; a small dusky spot at base of caudal; emerald spot on tongue conspicuous, fading in spirits. D. VI-14; A. 14 or 15.
- r.* Only a few scales on opercle.
- s.* Scales large, about 60..... *hastatus*
- ss.* Scales smaller, about 71..... *bayamonensis*, 249
- rr.* Upper part of opercle with a large patch of scales..... *oceanicus*, 250

247. *Gobius soporator* Cuvier & Valenciennes. *Mapo*.

Head 3 to 3.3; depth 4 to 4.8; eye 4.5 to 5; snout 3.5 to 4; maxillary 2.5 to 3; mandible 2 to 2.3; interorbital 5 to 6; preorbital about 8.5; scales 38, -13; D. VI-10, the longest spine about 1.7 in head, the longest ray 1.8; A. 9, the longest ray 1.8; pectoral 1.3; ventral 1.5. Body stout, heavy forward, compressed posteriorly, the least width of caudal peduncle about 4 in its least depth; head broad, compressed blunt; mouth moderate, slightly oblique, the maxillary reaching middle of eye; isthmus very broad, gill-openings extending forward but slightly in front of base of pectoral; lips thick; teeth in

upper jaw in a broad band, those of outer series enlarged, the inner ones small; teeth of lower jaw similar but smaller. Entire body covered with large, ctenoid scales, smaller on nape and belly; head naked. Fins rather large; origin of spinous dorsal slightly nearer snout than base of last soft ray; space separating dorsals about 2 in eye; dorsal and anal rays not reaching caudal when depressed; pectoral broad, upper rays partially detached and silk-like; ventral disk rather free from belly, not reaching vent; caudal short, rounded.

Color, mottled or marbled-gray above and on sides and head, under parts white; back with about 4 dark areas extending down on sides; sides with irregular series of small white specks; dorsals and caudal barred with white and dark; anal light, with dark edge; other fins pale. Some specimens are considerably darker and have the fins darker, but the color pattern is essentially the same in all.

Found in the tropical seas; widely distributed and almost everywhere common, lurking among stones or on sand in shallow water, or in rock pools, moving very quickly when disturbed; north on our coasts to Carolina and Gulf of California. The commonest of all shore-fishes in tropical America. Length 3 to 5 inches. Among our species this seems to be the most nearly related to the European *Gobius niger*, and may, therefore, be held to represent the subgenus *Gobius*, if our other species be placed in different subgenera. Perhaps all the others will ultimately be removed from *Gobius*.

This is apparently the most abundant and generally distributed goby in Porto Rico, the collection containing numerous specimens from San Antonio Bridge, San Geronimo, Palo Seco, San Juan, Mayaguez, Puerto Real, Guanica, Ponce, Fajardo, Hucares, Isabel Segunda, and Culebra Island.

Gobius saporator Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 56, 1837, Martinique; Jordan & Evermann, l. c., 2216, 1898.

Gobius lineatus Jenyns, Zool. Voy. Beagle, 95, pl. 19, fig. 2, 1842, Galapagos Archipelago.

Gobius catulus Girard, Proc. Ac. Nat. Sci. Phila. 1858, 169, St. Joseph Island, Texas.

Gobius mapo Poey, Memorias, II, 277, 1861, Cuba.

Gobius lacertus Poey, Memorias, II, 278, 1861, Cuba.

Gobius andrei, Sauvage, Bull. Soc. Philom., ser. 7, IV, 44, 1880, Rio Guayas, Ecuador.

Gobius carolinensis Gill, Proc. Ac. Nat. Sci. Phila. 1863, 268, Charleston, S. C.

Gobius brunneus Poey, Synopsis, 393, 1868, Havana.

Gobius arundelii Garman, Proc. N. E. Zool. Club, I, June 9, 1899, Clipperton Island.

248. *Gobius lyricus* Girard.

Description of male: Head 4; depth 4.5 to 5; eye 4 to 5; snout 3.6; maxillary 2.3; mandible 2.3; interorbital 4 to 5; preorbital 5 to 6; scales 29-9; D. vi-11, the longest spine about 3 in head, the longest ray about 1; A. i, 10, the longest ray 1.2; pectoral about equal to head; ventrals 1.1; caudal about twice length of head. Body elongate, tapering; head heavy, body somewhat compressed; head short, broad; snout short, abruptly decurved; mouth rather small, somewhat oblique, maxillary reaching vertical of pupil; lower jaw included; eye small, high; isthmus broad, gill-openings not extending forward under opercles; teeth rather strong, in one series in each jaw; lower jaw with about 4 short canine-like teeth behind the other teeth; teeth of upper jaw larger than those of lower. Fins rather large; first dorsal with 2 or 3 filamentous spines reaching, when depressed, beyond base of second dorsal, or even to caudal; space separating dorsals very short, distance from snout to origin of first dorsal 3 in body; anal rays reaching caudal; caudal long and pointed; pectoral rather long, reaching origin of anal; ventrals rather short, disk not reaching vent. Scales large, cycloid, somewhat crowded anteriorly; nape scaled, head and breast naked.

Color, olivaceous or dark, with 6 or 7 somewhat regular dark crossbars, the 2 or 3 posterior ones broadest, body with other dark blotchings and irregular markings; head marbled with darker, jaws, opercles, and branchiostegals black; a black bar across isthmus; first dorsal mostly dusky translucent, somewhat barred; second dorsal and anal plain dusky; caudal dark-blue with 2 longitudinal stripes of bright red changing to rose or orange in alcohol; pectoral finely barred with blackish and pale; lower parts yellowish.

Female examples may be described as follows: Head 3.8; depth 4.9; eye 3.6 to 4; snout 4 to 4.5; maxillary 2.5; mandible about 3; interorbital 5; preorbital 5; scales 29-9. D. vi-11, the longest ray about 1.5 in head; A. i, 10, longest ray 1.7; pectoral 1; ventral 1.3; caudal 0.8. Form not markedly differing from that of male, caudal peduncle less slender. Color darker, head and body blotched with dark; about 4 large dark blotches across back, more distinct than in male; black blotch at base of caudal large; fins all barred with light and dark; no reddish bands on caudal; dorsal spines less filamentous.

Found in the Gulf of Mexico and along the South Atlantic coast from Galveston and Indian River,

south through the West Indies; generally common, but probably not in Porto Rico. Length 2 to 3 inches. The collection contains 5 males and 4 females from Fajardo and 2 males from Isabel Segunda.

Gobius lyricus Girard, Proc. Ac. Nat. Sci. Phila. 1858, 169, Brazos Santiago, Texas; Jordan & Evermann, l. c., 2224, 1898.

Gobius wurdemannii Girard, Proc. Ac. Nat. Sci. Phila. 1858, 169, Brazos Santiago, Texas.

Smaragdus costalesi Poey, Memorias, II, 280, 1861, Havana.

249. *Gobius bayamonensis* Evermann & Marsh.

Head 5.8; depth 6.4; eye 5; snout 3.2; maxillary 1.8; mandible 1.9; interorbital 7.6; preorbital 4.6; scales 74,-19, about 29 before dorsal; D. vi-14, longest spine about 0.7 in head, the longest ray 1.5; A. 15, longest ray 1.5; pectoral 0.9; ventrals 1; caudal very long and pointed. Body very long and slender; head long; caudal peduncle long; mouth very large, oblique; maxillary long, reaching past posterior border of orbit; only a few scales on opercle.

Color as in *G. oceanicus*, which this species closely resembles. The smaller (74 instead of 63 to 65), almost cycloid scales, fewer scales on opercle, longer head, larger mouth, longer maxillary, and longer and more slender body are differences which we can not reconcile with the descriptions of *G. oceanicus* or with numerous specimens of it which we have from Porto Rico.

This description is based on a single specimen 9 inches in length, No. 49365, U. S. N. M., bought in the San Juan market, January 14. It probably came from near the mouth of Bayamon River at Palo Seco, for which stream the species was named.

Gobius bayamonensis Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 355, San Juan market, Porto Rico.




FIG. 90.—*Gobius bayamonensis*.


250. *Gobius oceanicus* Pallas. *Esmeralda*; "Seti."

Head 5; depth 6 to 6.5; eye 5 to 5.5; snout about 3; maxillary 2 to 2.3; mandible 2; interorbital 7 to 8; preorbital 4 to 4.5; scales about 63,-17; D. vi-14, the longest spine variable in length, 0.7 to 1.5 in head, the longest ray about 1.5; A. 15, the longest ray 1.4 to 2 in head; pectoral 1.1; ventrals 1.2; caudal long and pointed. Body very elongate, tapering gradually to long caudal peduncle; head short; snout short; mouth large, oblique, lower jaw slightly projecting, maxillary reaching posterior border of pupil; interorbital very narrow, preorbital broad; isthmus broad, gill-openings not extending forward much beyond angle of opercle; teeth in narrow bands on jaws, outer slightly enlarged. Scales ctenoid, rather large, smaller and crowded anteriorly, about 26 before dorsal; nape scaled, breast and cheek naked, opercle with a few scales at top.

Color in alcohol: Pale-olivaceous; head, back, and upper part of side with fine dark punctulations; a large black blotch on side below spinous dorsal; middle of side with about 12 broad <-shaped black markings opening backward; an obscure dark blotch at base of caudal; opercle with a dark blotch; 3 or 4 small black spots on first dorsal spine; spinous dorsal dusky with a light and a dusky streak at base; soft dorsal dusky, with a light area between each two rays, anterior rays barred with light and dark; anal pale; pectoral somewhat dusky; ventrals dark, probably blue in life, with white border.

Found along the South Atlantic and Gulf coasts of the United States and southward through the West Indies to the South American coast; not rare; recorded from Cuba, Martinique, Surinam, and Cayenne. Apparently common in Porto Rico, as shown by the numerous specimens in the collection from San Juan market, Palo Seco, and Boqueron. Specimens were dredged at stations 6054, off San Juan Harbor, in 4 to 5 fathoms, and 6087, off Culebra Island, in 15 fathoms. Our largest examples are about 7.5 inches long. This species reaches a foot or more in length and is of considerable value as food.

- Gobius cauda longissima acuminata* Gronow, Zoophyl., 82, No. 227, pl. 4, fig. 4, 1763, locality unknown.
Gobius oceanicus Pallas, Spicilegia, VIII, 4, 1769, after Gronow; Jordan & Evermann, l. c., 2230, 1898.
Gobius lanccolatus Bloch, Fische Deutschlands, II, 8, pl. 38, fig. 1, 1783, Martinique.
Gobius buccalvus Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 119, 1837, Surinam, Cayenne, and Cuba.
Gobionellus lanccolatus, Poey, Fauna Puerto-Riqueña, 338, 1881; Stahl, l. c., 78 and 165, 1883.

FIG. 91.—*Gobius oceanicus*.

Genus 139. AWAOUS Steindachner.

Inner edge of shoulder-girdle with two or more conspicuous dermal flaps; preorbital region very long; premaxillary and maxillary strong; lips thick; scales rather small, ctenoid, 40 to 80 in a longitudinal series; interorbital groove with a conspicuous median crest; otherwise essentially as in *Gobius*. The species reach a large size and are confined to the fresh waters of the tropics of America and to the Hawaiian Islands. Asiatic species of similar habit have much larger scales and seem to form a distinct genus, *Rhinogobius* Gill. The physiognomy in each is peculiar, the snout being long and convex.

- a. Scales about 53, little crowded anteriorly, 21 before dorsal on nape; depth 5.66 in length; head 4; eyes placed high, interorbital area equal to diameter of eye; mouth horizontal; maxillary extending to middle of eye, 2.33 in head, lower jaw more flat than in *A. taiasica*; teeth small, in narrow bands, those of outer row above enlarged, some large teeth in band of lower jaw. D. vi-1, 12; A. i, 10. Uniform yellowish in spirits. *flavus*
aa. Scales 60 to 70, crowded anteriorly, about 30 scales before dorsal on nape; 21 scales between second dorsal and anal; head broader than high; body compressed posteriorly, rather depressed anteriorly; greatest depth 5.25 in length; head 3.25 in length. Olivaceous; a series of irregular, roundish blotches along middle of side; narrow dark streaks radiating from eye; a blackish streak running across upper margin of opercle and extending obliquely across base of upper pectoral rays; belly white; dorsal and caudal more or less distinctly barred with wavy blackish lines. D. vi-11; A. 11; scales about 65. *taiasica*, 251
aaa. Scales 76 to 82, 24 scales between second dorsal and anal; head as broad as high; depth of body 6.6 in length; head 4, flat above, snout more elongate, upper profile oblique, eye one-eighth of head, equaling interorbital area (in adult); mouth horizontal; lower jaw included; maxillary reaching to below anterior margin of eye; teeth of outer series enlarged; canine teeth none; scales ctenoid, those on nape and anterior part of body very small; head naked; dorsal fins lower than body, none of the spines produced; caudal rounded, 7 in length of body. Yellowish olive; back and sides reticulated with blackish; head, dorsal, caudal, and pectoral fins dotted with blackish, the spots forming streaks on second dorsal; six cross-series of dots on caudal; an irregular small blackish spot on upper part of root of pectoral. D. vi-11; A. 11; scales about 80. *mexicanus*

251. *Awaous taiasica* (Lichtenstein). "Guavina"; "Saga" or "Zaga."

Head 3.5; depth 5.4 to 6; eye 5.5 to 6; snout 2.2 to 2.4; maxillary 2.4 to 2.9; mandible 2.4 to 3; interorbital 6.5 to 8; preorbital 3.3 to 4; scales 66 to 71, -22; D. vi-11, longest spine about 2 in head, longest ray 2; A. i, 10, longest ray 2 to 3 in head; pectoral 1.6; ventral 1.7; caudal 1.5. Body rather slender, head heavy, somewhat depressed, tapering, its greatest width about 1.5 in its length; snout long; mouth moderate, very broad, its width about 2.5 in head; maxillary reaching anterior border of orbit; lips very heavy, upper with a broad flap; teeth of upper jaw in two series, those of anterior series much enlarged and recurved; teeth of lower jaw in a narrow band, outer scarcely enlarged; no teeth on vomer or palatines; isthmus very broad, gill-openings scarcely extending forward; branchiostegals 4; gillrakers 3 + 6, very short and soft; inner edge of shoulder-girdle with 2 or 3 short papillae. Posterior half of body compressed, least width of caudal peduncle about half its least height. Scales small, ctenoid, much reduced anteriorly, those on nape and in front of dorsal embedded and cycloid, about 30 series between dorsal and occiput; head naked; breast scaled. Fins rather large; origin of

spinous dorsal about midway between tip of snout and middle of soft dorsal; space between dorsals scarcely half orbit; dorsal and anal rays, when depressed, not reaching base of caudal; pectoral long, nearly reaching vent; ventral disk broad and free; caudal rounded.

Color, pale-olivaceous or yellowish, with a series of about 8 large black blotches along middle of side; back and head blotched and vermiculated with dark; under parts pale; dorsals and caudal with alternate bars of light and dark; pectoral dusky; ventrals and anal pale.

This goby reaches a length of a foot or more. It is extremely variable in form and coloration, as is the case with most widely distributed fresh-water fishes. It is found in the fresh waters of the West Indies and on both coasts of Mexico, south to Brazil; common in Cuba, in Sinaloa, and about La Paz in Lower California, thence southward to Panama. It is also common in the fresh and brackish waters of Porto Rico, specimens being obtained from the Bayamon River near Bayamon, the Rio de Caguaitas at Caguas, and from a brackish pool on Vieques Island just west of the town of Isabel Segunda. It is of some value as food.

Amore guacu, Maregrave, Hist. Brasil., 166, 1648, Brazil.

Gobius taitasica Lichtenstein, Berl. Abhandl., 273, 1822, Brazil; not *Taitasica* of Maregrave.

Gobius banana Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 103, 1837, Santo Domingo.

Gobius martinicus Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 105, 1837, Martinique.

Chonophorus bucculentus Poey, Memorias, II, 275, 1861, Cuba.

Rhinogobius contractus Poey, Mem., II, 424, 1861, Cuba; Poey, Fauna Puerto-Riq., 338, 1881; Stahl, l. c., 78 and 165, 1883.

Gobius dolichocephalus Cope, Trans. Amer. Philos. Soc. Phila. 1869, 403, near Orizaba, Mexico.

Euctenogobius latus O'Shaughnessy, Ann. Mag. Nat. Hist., series 4, XV, 1875, 146, Bahia.

Awaous taitasica, Jordan & Evermann, l. c., 2236, 1898.




FIG. 92.—*Bollmannia boqueronensis*.

Genus 140. *BOLLMANNIA* Jordan.

This genus differs from *Lepidogobius* in having no fleshy processes on inner edge of shoulder-girdle, the interorbital area of skull narrower and without trace of median keel, and in the very large ctenoid scales. From *Gobius* proper it is distinguished by the presence of 7 dorsal spines and by the presence of large scales on cheek. The genus *Bollmannia* has now five known species, *B. chlamydes*, *B. ocellata*, *B. macropoma*, *B. stigmatura*, and *B. boqueronensis*. The first of these was described in 1889 by Dr. Jordan from the west coast of Columbia and the next three by Dr. Gilbert in 1891 from the Gulf of California. Not until the last species was obtained by us off Porto Rico was the genus known to have any representative in the Atlantic. All the species are found only at considerable depths, and do not inhabit shoal water, as is the case with most other gobies.

252. *Bollmannia boqueronensis* Evermann & Marsh.

Head 4; depth 5.5; eye 3.5; snout 4.4; maxillary 2.2; mandible 2.5; interorbital width 3 in eye; preorbital 6; scales 27-8; D. VII-13, the longest spine 1.5 in head, the longest ray 1.2; A. 12, the longest ray 1.25 in head; pectoral 0.75; ventrals 1.1; caudal 0.4. Body long, slender, tapering; head short; snout blunt; mouth large, oblique; jaws subequal, maxillary reaching posterior border of pupil; isthmus narrow, gill-openings reaching forward to below preopercle; eyes large, high, close together, interorbital very narrow and without median keel; no fleshy process on inner edge of shoulder-girdle; teeth on jaws in narrow bands, those of outer series somewhat enlarged; opercle short, about 3 in

head. Fins moderate; origin of spinous dorsal slightly behind base of pectoral, its spines 7 in number, not filamentous; interspace between dorsals less than diameter of eye; soft rays of dorsal and anal reaching, when depressed, beyond base of caudal; caudal long and pointed, as in *Gobius oceanicus*; pectoral pointed, reaching beyond origin of anal; ventral disk moderate, free from belly, longest rays barely reaching origin of anal. Scales very large, weakly ctenoid; nape, cheeks, and breast scaled, scales somewhat smaller than on body, about 9 scales before dorsal.

Color, pale-olivaceous or straw-color, back and upper part of head with profuse fine dark punctulations; under parts pale, breast somewhat dusky; dorsal fins barred with white and dark, a large jet-black ocellus on posterior part of spinous dorsal; other fins pale, ventral disk somewhat dusky in front.

Known only from the type and four cotypes dredged by the *Fish Hawk* at station 6074, off Puerto Real, in 8.5 fathoms, January 25, 1899. It reaches a length of 2.75 inches. Type No. 49366, U. S. N. M.

Named from Ensenada del Boqueron, in which the type was obtained.

Bollmannia boqueronensis Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 356, Ensenada del Boqueron, Porto Rico.

Genus 141. MICROGOBIUS Poey.

Dorsal spines 7 or 8; scales very small, cycloid or weakly ctenoid, body scaled anteriorly as well as posteriorly, head naked, nape, belly and breast usually so. Inner edge of shoulder-girdle without fleshy processes; body more or less compressed; mouth large, very oblique; lower jaw conspicuous, teeth strong; interorbital groove with or without a median ridge. Vertebrae 11 + 15 or 16.

- a. Scales about 42. Body elongate, moderately compressed, depth 4 to 5 in length; head long and large, rather sharp in profile, 3 to 3.5 in body; eye longer than snout, 4 in head; mouth large, very oblique, lower jaw strongly projecting; maxillary 1.5 to 2.5 in head, extending to opposite middle of eye, or much beyond front of orbit; teeth in few series, the outer very long and slender, curved, the lower longest, none canine-like; scales small, some of them with short, thick teeth, those of anterior part of body not well developed; dorsal spines more or less filamentous, third and fourth or fourth and fifth sometimes with long filaments; caudal pointed, about as long as head. Grayish-olive, with rather sharply defined markings of darker brown overlaid with orange in life; head with a pale-bluish or gilt stripe from maxillary backward across suborbital region to upper edge of gill-opening; another pale gilt streak from snout along lower part of eye, another from angle of mouth upward and backward; rest of head dark; opercle with an oblique, blackish bar; top of head and nape with dark marbling surrounded by paler reticulations; back with a series of black cross-blotches mostly separated on median line; 2 narrower dark vertical bars behind pectoral; middle line of side posteriorly with longitudinally oblong black blotches; besides these, numerous other blotches not regularly arranged; first dorsal with 2 or 3 oblique black bands; second dorsal pale, with about 4 series of black dots; caudal spotted with black; pectoral yellowish; ventral black, its center yellowish (male); anal pale. D. VII-15; A. 16 or 17. *gulosus*
- aa. Scales 50 to 55.
- b. Scales 50, -14, cycloid; D. VII-15; body scarcely compressed. *eulcypis*
- bb. Scales 55, -12, strongly ctenoid; D. VII-17; body greatly compressed. *meeki*, 253
- aaa. Scales 65 or more.
- c. Caudal fin more than one-third (two-fifths) length of body. Scales very small, cycloid, deciduous. Body elongate, much compressed, highest in front of ventrals, tapering regularly to very narrow, short caudal peduncle; greatest depth 4.75 in length; head 3.5. Head compressed, much higher than wide; snout very short, acute, preorbital not as wide as pupil; mouth terminal, very wide and oblique; jaws equal; maxillary reaching vertical from middle of orbit, 2 in head. Outer series of teeth enlarged. Eye 3 in head. Dorsals closely contiguous; spines very slender, fifth slightly produced and filamentous; pectoral as long as head. Head and body translucent, overlaid by brilliant green luster, formed by minute, close-set green points; 3 conspicuous translucent bars wider than interspaces, crossing body close behind head; head with 2 brilliant narrow blue and green lines running obliquely across cheek below eye; dorsal whitish, with 2 or 3 lengthwise series of large reddish-brown spots; spinous dorsal blackish at base, upper caudal rays marked with red, lower portion of caudal and most of anal fin blackish, anal whitish at base, anterior rays tipped with white. In spirits, body dusted with dark points; 2 light crossbars toward head; lower part of caudal and anal black. D. VII-16; A. 15. *thalassinus*
- cc. Caudal fin less than one-third length of body. Scales small, cycloid, embedded. Body very much compressed, more or less elongate, greatest depth at ventrals 4 (female) to 6.5 (male) in length; head 3.5 to 4. Head much compressed, much deeper than wide. Snout very short, acute, anterior profile not deurved, not steep; preorbital not as wide as pupil; mouth very large, very oblique or almost vertical; maxillary extending to below pupil, 2 in head (in male), 2.25 (in female). Lower jaw projecting, teeth of outer series enlarged, recurved. Eye 3.25 to 4 in head. Dorsals contiguous, spines very fine, produced in filaments, third highest, a little longer than head; second dorsal and anal high. Head and nape naked. In the female the depth is greater, mouth less oblique, smaller; profile from spinous dorsal oblique. First dorsal spine highest, 3.33 in length. Ventrals much shorter than in males. Dark gray; female with a short bright-blue bar bordered by blackish above pectoral; a blotch of sky-blue and orange below eye; fins dusky, ventrals pale in female, dusky in males. Males with body plain bluish-gray. D. VII-17 to 20; A. 18 to 21; scales 68 to 70. *signatus*

253. *Microgobius meeki* Evermann & Marsh.

Head 3.75; depth 6; eye 3.5; snout 5.5; interorbital 7; preorbital 7; maxillary 2; mandible 1.5; scales 55,-12; D. VII-17; A. 16. Body slender, greatly compressed, tapering regularly from pectorals to caudal; head moderately heavy, interorbital space very narrow; eye large, high; mouth large, oblique; maxillary reaching posterior border of orbit; lower jaw projecting; teeth in bands in each jaw, outer series greatly enlarged and strongly recurved, those of lower jaw largest; isthmus rather narrow, gill-openings continuing forward; body densely scaled, scales strongly ctenoid, those anteriorly somewhat reduced; nape, breast, and entire head naked; origin of spinous dorsal from snout 3.5 in length; dorsals very close together; spines of first dorsal filamentous, exceeding head in length; soft dorsal and anal long, their bases about equal, about 2.5 in body, their last rays reaching past base of caudal when depressed; caudal pointed, its longest rays about equal to head; pectoral about equal to head, reaching origin of anal; ventrals united, almost reaching origin of anal.

Color, light-olivaceous, dusted over uniformly with fine dark punctulations; large dark shoulder-spot between base of pectoral and origin of spinous dorsal; a few indistinct dark areas on side of head; lower jaw dark at tip; an obscure dark blotch at base of caudal; fins all rather pale except ventrals, which are dark, perhaps bluish in life; caudal somewhat dusky; anal dark-edged.

This species seems related to *M. eulepis* Eigenmann & Eigenmann, described from Fortress Monroe, Va., but differs in the smaller and strongly ctenoid scales, greatly compressed body, and in coloration. It is described from a single specimen, 1.5 inches in length (No. 49367, U. S. N. M.), collected at Fish Hawk station 6087, in 15.25 fathoms, between Culebra and Vieques islands.

Microgobius meeki Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 356, between Vieques and Culebra islands, east of Porto Rico.




FIG. 93.—*Microgobius meeki*.

Genus 142. GOBIOIDES Lacépède. Barretos.

Body greatly elongate, compressed behind, scales very minute; head small; eyes very small; mouth large, oblique, lower jaw projecting; gill-openings moderate. Teeth in a band, those in outer series being very strong. Dorsal rays v to VII, 16 to 23; anal rays 17 to 23. Dorsal fin low, continuous, spines similar to soft rays, but more widely separated; soft dorsal and anal joined to base of caudal; ventrals 45, united in a disk which is formed much as in *Gobius*. No air-bladder; no pseudobranchiae. From *Tenioides* (= *Amblyopus*) the genus *Gobioides* is distinguished by the absence of barbels, the presence of scales, and by the much smaller number of rays in its vertical fins.

Found in the brackish waters of the Tropics, reaching a considerable size.

254. *Gobioides broussonnetii* Lacépède. Barreto.

Head 5.25 (young) to 7 (adult); caudal 3.5 to 5; eye small but evident, 7 to 10 in head; interorbital space 1 to 1.66 diameter of eye; D. VI, 16; A. 1, 16. Body elongate, mouth oblique, maxillary extending beyond eye; teeth in bands, outer series enlarged, shorter, and closer set than in *Gobioides peruanus*; scales twice as large as in *G. peruanus*, those on anterior part of body not imbricated, much

smaller than those on posterior part, which are elongate-oval in form. Violet bars extending downward and forward on upper part of body; sometimes a violet spot with a lighter or darker dot at end of bars; head marbled or spotted with dark violet or brown. (Steindachner.)

West Indies to Brazil; common southward, ascending rivers; once taken near New Orleans (Bean & Bean). Not seen by us in Porto Rico, but recorded from that island by Professor Poey. Length 20 inches or more.

Gobioides broussonnetii Lacépède, Hist. Nat. Poiss., II, 580, 1798, probably from Surinam, "given by Holland to France"; Jordan & Evermann, I. c., 2263, 1898.

Amblyopus brasiliensis Bloch & Schneider, Syst. Ichth., 69, 1801, Brazil; on drawing made by Prince Maurice.

Gobius oblongus Bloch & Schneider, Syst. Ichth., 548, 1801; based on Lacépède.

Gobioides barreto Poey, Memorias, II, 282, 1861, Cuba; Poey, Fauna Puerto-Riqueña, 338, 1881.

Amblyopus mexicanus O'Shaughnessy, Ann. Mag. Nat. Hist., series IV, vol. XV, 1875, 147, Mexico.

Family LXVII. ECHENEIDIDÆ. The Remoras.

Body fusiform, elongate, covered with minute, cycloid scales. Mouth wide, with villiform teeth on jaws, vomer, palatines, and usually on tongue. Premaxillaries not protractile. Lower jaw projecting beyond upper. Spinous dorsal modified into a sucking disk, which is placed on top of head and neck and is composed of a double series of transverse, movable, cartilaginous plates, serrated on their posterior or free edges. By means of this disk these fishes attach themselves to other fishes or to floating objects and are carried for great distances in the sea. Opercles unarmed. Pectoral fins placed high; ventral fins present, thoracic and close together, 1, 5; dorsal and anal fins long, without spines, opposite each other; caudal fin emarginate or rounded. Branchiostegals 7. Gills 4, slit behind fourth; gillrakers short; gill-membranes not united, free from isthmus. Pseudobranchiæ obsolete. Several pyloric appendages. No air-bladder. No finlets. No caudal keel. Vertebrae more than 10 + 14.

This family embraces about 4 genera and 10 species, found in all seas, all having a very wide range. The species of this group are apparently descended from a fossil genus, *Opisthomyzon*.

- a. Body very slender, vertebrae 14 + 16 = 30; ventrals narrowly adnate to abdomen; lower jaw produced in a flap; pectoral acute, with flexible rays. PITHEIRICHTHYS
- bb. Laminae 10 only ECHENEIS, 143
- aa. Body rather robust, vertebrae 12 + 15 = 27; ventrals broadly adnate to abdomen; lower jaw not produced; pectorals rounded. REMILEGIA
- c. Laminae 24 to 27. REMORA
- cc. Laminae 16 to 20. RHOMBOCHIRUS
- d. Pectoral rays soft and flexible. REMORA
- dd. Pectoral rays stiff and ossified. RHOMBOCHIRUS

Genus 143. ECHENEIS (Artedi) Linnæus.

Body comparatively elongate, vertebrae 14 + 16 = 30; disk long, of 20 to 28 laminae; pectoral pointed, its rays soft and flexible; soft dorsal and anal long, of 30 to 41 rays each; caudal lunate in adult, convex in young.

Species of wide distribution, attaching themselves mainly to sea turtles and large fishes.

- a. Disk of 22 to 26 laminae (rarely 21 or 28), its length less than one-fourth body *naucrates*, 255
- aa. Disk of 20 or 21 laminae, its length more than one-fourth body *naucrateoides*

255. *Echeneis naucrates* Linnæus. *Shark-sucker; Pega; Pegador; Sucking-fish.*

Head 5.25; depth 11 to 12; D. XXII to XXVIII (rarely XXI), 32 to 41; A. 31 to 38. Breadth between pectorals 7.5; disk 4 to 5 in body; eye 5 in head; snout 2.33; maxillary 3; from angle of mouth to tip of lower jaw 2.66; pectoral 1.4; ventrals 1.5; middle caudal rays 1.4; highest anal ray 2; highest dorsal ray 2.33; width of disk 2.5 in its length; base of dorsal 2.5, anal 2.5, in body. Body elongate, subterete, slender; lower jaw strongly projecting, the tip flexible; maxillary reaching nostril; teeth uniform in adult, young with series of small slender teeth in advance of others; gillrakers short and slender, about equal to pupil; vertical fins low; anal rays higher than dorsal anteriorly; pectoral reaching very slightly past tips of ventrals; origin of ventral spine under middle of pectoral base; inner rays of ventral fins narrowly adnate to abdomen; dorsal and anal commencing and ending opposite each other; caudal with middle rays produced in young, the fin becoming emarginate or lunate with age.

Color, brownish; belly dark, like back, as usual in this family; side with a broad stripe of darker, edged with whitish, extending through eye to snout; caudal black, its outer angles whitish; pectoral and ventrals black, sometimes bordered with pale; dorsal and anal broadly edged with white anteriorly; adult nearly uniform dark-brown, not paler below.

Widely distributed in the warm seas; common north to Cape Cod and occasionally to San Francisco, attaching itself to turtles and to large fishes. Very common in the Tropics, being found attached to sharks, groupers, or any other large fish, without regard to species. Few large sharks at Key West are without them. They are often caught with hook and line from the wharf, where they frequently forsake their host to take the bait. Lütken's remark that only *Remora remora* has been recorded from sharks is no longer true. Several writers have recognized 2 species of *Echeneis* proper—*naucrates*, with 22 to 26 laminae, the disk 4 to 5 in body, and *naucratoides* (= *albicauda* = *holbrookii* = *lineatus*), in which the disk is longer, 3.6 to 4 in body, but composed of fewer, 20 or 21, laminae. The latter form is rather common on our coast, the specimens from Key West above mentioned having 21.

Not obtained by us in Porto Rico, but recorded from that island by Professor Poey and Dr. Stahl.

Echeneis naucrates (misprint for *naucrates*) Linnaeus, Syst. Nat., ed. X, 261, 1758, "in Pelago Indico."

Echeneis albicauda Mitchill, Amer. Monthly Mag., II, 1817, 244, New York.

Echeneis lunata Bancroft, Proc. Comm. Zool. Soc., I, 134.

Echeneis australis Griffith, Cuvier Anim. Kingdom, 504, 1837.

? *Echeneis vittata* Lowe, Proc. Zool. Soc. Lond. 1839, 89, Madeira.

Leptecheneis naucrates, Poey, Fauna Puerto-Riqueña, 333, 1881; Stahl, l. c., 80 and 166, 1883.

Echeneis naucrates, Jordan & Evermann, l. c., 2269, 1898.



FIG. 94.—*Echeneis naucrates*.

Family LXVIII. MALACANTHIDÆ. The Blanquillos.

Body more or less elongate, fusiform or compressed. Head subconical, anterior profile usually convex; suborbital without bony stay; bones not greatly developed; cranial bones not cavernous; opercular bones mostly unarmed. Mouth rather terminal, little oblique; teeth rather strong; no teeth on vomer or palatines; premaxillary usually with a blunt posterior canine, somewhat as in the *Labridæ*; premaxillaries protractile; maxillary without supplemental bone, not slipping under edge of preorbital. Gills 4, a long slit behind fourth; pseudobranchiæ well developed; gill-membranes separate, or more or less united, often adherent to isthmus; lower pharyngeals separate. Scales small, ctenoid; lateral line present, complete, more or less concurrent with back; dorsal fin long and low, usually continuous, spinous portion always much less developed than soft portion, but never obsolete; anal fin very long, its spines feeble and few; caudal fin forked; tail diphyccercal; ventrals thoracic or subjugular, 1, 5, close together; pectoral fin not very broad, rays all branched; vertebrae in normal or slightly increased number (24 to 30). Pyloric cæca few or none.

Fishes of temperate and tropical seas, some of them reaching a large size. Genera about 6; species about 8 to 10, mostly American. The relationships of the family are obscure, and it may be that the genera here associated are not really closely allied.

MALACANTHINÆ:

a. Vertebrae 24; preopercle entire.

b. Soft dorsal and anal extremely long, each with more than 40 rays; form slender; scales very small.. MALACANTHUS

aa. Vertebrae more than 24; preopercle more or less serrate.

CAULOLATILINÆ:

c. Soft dorsal and anal moderate, each with 22 to 27 soft rays; preopercle serrate; scales rather small; form robust.

d. Upper jaw with posterior canines; dorsal spines graduated..... CAULOLATILUS, 144

LATILINÆ:

cc. Soft dorsal and anal short, each of 13 to 15 soft rays; preopercle denticulate; scales small; form robust.

e. Nape with a large adipose appendage; a fleshy prolongation on each side of labial fold, extending forward behind angle of mouth..... LOPHOLATILUS

Genus 144. CAULCLATILUS Gill. Blanquillos.

Body elongate, subfusiform, not strongly compressed, heavy forward, tapering to a rather slender caudal peduncle; profile of head strongly arched; mouth moderate, little oblique, jaws nearly equal; lips thick; maxillary narrow, not slipping under preorbital; teeth in villiform bands, preceded by a row of stronger acute teeth; posterior teeth in each jaw canine-like, directed forward; posterior canines of upper jaw largest; no teeth on vomer or palatines; preopercle pectinate, teeth nearly even; opercle with a blunt, flat spine; eyes large, lateral; gill-membranes slightly connected, forming a fold across isthmus, with which they are narrowly joined; branchiostegals 6; gillrakers short and stout; nostrils double, round, close together; scales small, firm, ctenoid; lateral line continuous, concurrent with the back; dorsal with 7 to 9 slender, pointed, graduated spines and 22 to 27 soft rays; anal similar to soft dorsal, with 1 or 2 small spines and more than 20 soft rays; caudal fin forked; ventral fins thoracic; no adipose appendage at nape; vertebrae $12 + 15 = 27$.

Large fishes of the warm seas of America; valued as food.

a. Scales small, about 125 in lateral line, about 50 in a transverse series.

b. Eye small, 6 in head; depth 3.5 in body; scales 13-120-35..... *microps*

aa. Scales larger, about 10-108-25..... *cyanops*, 256

256. *Caulolatilus cyanops* Poey. *Blanquillo*; *Tremba*.

Head 4 in total length; D. VII, 24; A. I, 22 (scales 10-108-25. Bean). Profile convex before eye, not ascending to nape; no scales on fins; soft rays little divided; caudal slightly lunate; the first caudal vertebra spoon-like, its cavity receiving air-bladder; vertebrae $12 + 15$; no pyloric caeca; stomach short; air-bladder large.

Color greenish above, a faint, broad, interrupted brown band above lateral line; some small brown spots above and below it; region below eye clear blue, not very different from color of belly; soft dorsal brown, paler at its base, edged with orange; spinous dorsal orange. (Poey.)

The blanquillo is related to the tile-fish, and is known only from the coasts of Cuba and Porto Rico, though it may be identical with *C. chrysops*, described from the coast of Brazil. It was not obtained by us in Porto Rico, but has been recorded from that island by Poey and Stahl.

When at Nassau, en route to Porto Rico, we caught with hook and line 2 fine examples of *Malacanthus plumieri* (Bloch), another relative of the tile-fish. This species is called "sand-fish" at Nassau, and its occurrence about Porto Rico may be confidently expected.

Caulolatilus cyanops Poey, Repertorio, I, 312, 1867, Cuba; Poey, Fauna Puerto-Riqueña, 334, 1881; Stahl, l. c., 78 and 164, 1883; Jordan & Evermann, l. c., 2278, 1898.

Family LXIX. DACTYLOSCOPIDÆ. The Sand Star-gazers.

Body oblong, low, compressed posteriorly, covered with moderate, cycloid, imbricated scales; lateral line complete, anteriorly running alongside of back, posteriorly median; head oblong, nearly plane above; eyes small, superior, well forward; suborbital bones enlarged, but without bony stay connecting with preopercle; nostrils double; opercles fringed; mouth nearly vertical; premaxillaries protractile, not forming entire edge of upper jaw; lips fringed as in *Uranoscopidae*; gill-openings very broad, membranes separated and free from isthmus, pseudobranchiæ present or obsolete. Dorsal fin very long, continuous or divided, several of anterior rays spinous; anal very long, commencing close behind vent, which is near breast; caudal diphycecal, free from dorsal and anal; pectoral variable, base broad and procurvent; ventrals jugular, 1, 3; vertebrae more than $10 + 14$; pyloric caeca none.

Small fishes living on sandy shores of tropical America, comprising 4 genera and about 10 species. The family is nearly related to the *Uranoscopidae*, of which group it seems to be a reduced or degenerate branch. Its relation with the Asiatic family *Leptoscopidae* are most intimate, the incomplete ventrals and simple pectoral rays of *Dactyloscopidae* being the chief distinctive features.

Genus 145. *DACTYLOSCOPUS* Gill.

Body moderately elongate, covered with rather large, cycloid scales; head cuboid, oblong, and nearly flat above; eyes small; interorbital space broad; mouth nearly vertical; lower jaw not dilated beneath nor emarginate in front, without barbels; no intralabial filament; teeth villiform, on jaws only; pseudobranchiæ very small or obsolete. Dorsal commencing at nape, with 6 to 12 slender spines, soft rays numerous; anal inserted behind dorsal; ventral rays, 1, 3.

a. Dorsal rays x to XII, 22 to 31; anal rays fewer than 35.

b. Soft dorsal with 28 to 31 soft rays; anal with 32 or 33; scales about 45.

c. Body rather slender, depth about 6 in length (7 with caudal); opercular fringe of 15 filaments *tridigitatus*, 257

cc. Body rather stout, depth 5.25 in length (6 in total with caudal); opercular fringe of 18 filaments.

d. Back not barred; head blotched and dotted *poeyi*

257. *Dactyloscopus tridigitatus* Gill.

Head 5 (in total) with caudal; depth 7; D. XII, 28; A. II, 32; P. 13; V. I, 3; scales $11 + 4 + 30 = 45$. Body slender, much compressed posteriorly; opercular fringe of 15 separate filaments. Origin of dorsal fin over lower angle of base of pectorals, or immediately before margin of operculum, its distance from snout to dorsal 5 in total length of body. Pseudobranchiæ very small (overlooked by Dr. Gill, but evident in living specimens).

In life, pale sand-color above, lower part whitish; above 12 narrow crossbands of whitish on the back, not extending down far on the side; head mottled above; fins all pale.

This species of star-gazer is found in the West Indies and the Barbados and north to southern Florida. It has been taken at Key West and Cape Florida, and a single example, 3 inches long, was seized by us in Ensenada del Boqueron, Porto Rico. It does not reach a length of more than a few inches, and is a shallow-water fish, frequenting the coral sands near shore, in which it is wont to bury itself, leaving only its eyes and mouth exposed, after the fashion of the flounders.

Dactyloscopus tridigitatus Gill, Proc. Ac. Nat. Sci. Phila. 1859, 132, Barbados; Jordan & Evermann, l. c., 2301, 1898.

Family LXX. GOBIESOCIDÆ. Cling-fishes.

Body rather elongate, tadpole-shaped, broad and depressed in front, covered by smooth, naked skin; mouth moderate; upper jaw protractile; teeth usually rather strong, anterior conical or incisor-like; posterior canines sometimes present; suborbital ring wanting; no bony stay from suborbital across the cheek; opercle reduced to a spine-like projection concealed in skin, behind angle of large preopercle, this spine sometimes obsolete; pseudobranchiæ small or wanting, the gills 3 or 2.5; gill-membranes broadly united, free or united to isthmus; dorsal fin on posterior part of body, opposite to anal and similar to it, both fins without spines; ventral fins wide apart, each with 1 concealed spine and 4 or 5 soft rays. Between and behind ventrals is a large sucking-disk, the ventrals usually forming part of it. This sucking-disk, which is wholly different in structure from that of *Cyclopterus* and *Liparis*, is thus described by Dr. Günther:

"The whole disk is exceedingly large, subcircular, longer than broad, its length being (often) one-third of the whole length of the fish. The central portion is formed merely by skin, which is separated from the pelvic or pubic bones by several layers of muscles. The peripheric portion is divided into an anterior and posterior part by a deep notch behind the ventrals. The anterior peripheric portion is formed by the ventral rays, the membrane between them, and a broad fringe which extends anteriorly from one ventral to the other. This fringe is a fold of the skin containing on one side the rudimentary ventral spine, but no cartilage. The posterior peripheric portion is suspended on each side on the coracoid, the upper bone of which is exceedingly broad, becoming a free, movable plate behind the pectoral. The lower bone of the coracoid is of triangular form, and supports a very broad fold of the skin, extending from one side to the other and containing a cartilage which runs through the whole of that fold. Fine processes of the cartilage are continued into the soft striated margin, in which the disk terminates posteriorly. The face of the disk is coated with a thick epidermis, like the sole of the foot in higher animals. The epidermis is divided into many polygonal plates. There are no such plates between the roots of the ventral fins." (Günther, Cat., III, 495.)

No air-bladder; intestines short; pyloric cæca few or none; skeleton firm; vertebræ 13 to $14 + 13$ to $22 = 26$ to 36.

Carnivorous fishes of small size and of no value as food, chiefly of the warm seas, usually living among loose stones between tide marks and clinging to them firmly by means of the adhesive disk. Their relations are obscure, but they are probably descended from allies or ancestors of the *Cottidae* or the *Batrachoididae*. Genera about 15; species 50. The principal genus is *Gobiesox*.

Genus 146. GOBIESOX Lacépède.

Body anteriorly very broad and depressed, posteriorly slender, covered with tough, smooth skin; opercle with a strong spine; head large, rounded in front; mouth terminal, crescent-shaped; lower jaw with a series of strong incisors in front, their edges rounded or truncate; upper jaw with a series of strong teeth, behind which are sometimes smaller teeth; no teeth on vomer or palatines; gills 3; gill-membranes broadly united under throat, not attached to isthmus; sucking disk large, posterior portion without anterior free margin. Dorsal and anal moderate, dorsal rays 6 to 12, anal rays 6 to 10. Vertebrae about 26, as far as known.

Species numerous, all American; mostly tropical, clinging to rocks near the shore.

Jordan & Evermann recognize 18 species of this genus in American waters; 5 of these are known only from our Pacific coast and the remaining 13 are all known only from the West Indies and the South Atlantic and Gulf coasts, though but one species has as yet been found in Porto Rico.

a. Dorsal fin moderate or short, its rays 6 to 11.

GOBIESOX:

b. Upper teeth in more than 1 series (character not verified in a few species); head broad.

c. Coloration in life chiefly olivaceous, with little red, sometimes banded with darker or paler.

d. Dorsal rays 12; anal rays 7..... *gyrinus*

dd. Dorsal rays 11; anal rays 6; fins black..... *nigripinnis*

ddd. Dorsal rays 9 or 10; anal rays 6..... *cephalus*

dddd. Dorsal rays 8; anal rays 6..... *tudes*, 258

ddddd. Dorsal rays 11; anal rays 10..... *strumosus*

ddddd. Dorsal rays 10; anal rays 8..... *virgatulus*

ee. Coloration in life chiefly bright-red, or else with red spots or bands, color not fading in spirits.

e. Color uniform red, unspotted, color not fading in spirits; dorsal rays 6 to 8; anal rays 6.

f. Lower jaw with short incisors on each side, followed by canines..... *macrophthalmus*

ff. Lower jaw with 2 horizontal incisors on each side, third horizontal tooth not incisor-like; no distinct canines.
crasinus

SICYASES:

bb. Upper teeth in a single series (character not verified in some species); dorsal and anal short.

g. Color chiefly red.

h. Body with dark crossbands and with spots of clear blue; body rather slender. D. 6 or 7; A. 6..... *rubiginosus*

hh. Body plain, light red; form rather slender..... *carneus*

gg. Color olivaceous or brownish, not red.

i. Dorsal rays 9; anal rays 6.

j. Color olivaceous, without bands..... *heeres*

jj. Color greenish, with 3 dark crossbands and many dots..... *punctulatus*

ii. Dorsal rays 7; anal rays 7; body with dark crossbands..... *fasciatus*

258. Gobiesox tudes Richardson.

Head 2.5; depth 4.66; width of head 2.5; D. 8; A. 6 in plate (5 in description, the first short ray apparently not counted by Richardson). Head very broad, as broad as long, abruptly truncate anteriorly; mouth large, maxillary reaching front of eye; lower jaw included; teeth entire; eye large, 4.75 in head, a little more than half interorbital width, 1.5 in snout. Distance from front of dorsal to caudal about equal to length of head; insertion of dorsal before vent; anal behind dorsal and much shorter than it; pectoral short.

This is a very little known species, which reaches a length of 5 inches. The type locality is not certainly known, but it has been supposed to be China. It is, however, certainly not Chinese, and is now shown to be West Indian. A single specimen, 1.25 inches long, collected on the reef at Culebra Island February 9, agrees very well with the above description by Richardson. There are but 8 dorsal and 6 anal rays. The color is uniform pale-yellowish, with a slight tinge of rosy on middle of back; fins all pale; eye very large, about 3 in head, much larger than in specimens of *G. strumosus* of same size, and the interorbital width is decidedly narrower. It seems to be a good species, distinct from *G. cephalus* and *G. nigripinnis*.

Gobiesox tudes Richardson, Voy. Sulphur, Fish., 103, pl. 46, figs. 1-3, 1845, habitat unknown; supposed to be China; Jordan & Evermann, l. c., 2333, 1898.

Family LXXI. BLENNIIDÆ. The Blennies.

Body oblong or elongate, naked or covered with moderate or small scales which are ctenoid or cycloid; lateral line variously developed, often wanting, often duplicated; mouth large or small, teeth various; gill-membranes free from isthmus or more or less attached to it; pseudobranchiæ present; ventrals jugular or subthoracic, of 1 spine and 1 to 3 soft rays, often wanting; dorsal fin of spines anteriorly, with or without soft rays; anal fin long, similar to the soft dorsal; caudal well developed. Vertebræ in moderate or large number, 30 to 80.

Carnivorous fishes of moderate or small size, mostly living near shore in tropical and temperate or arctic seas; most of them are carnivorous, the *Clinina*, so far as known, ovoviviparous, the others mostly oviparous. Genera about 80; species about 400; chiefly of rock pools and algae; some species in the lakes of Italy. Of the 57 genera of this family recognized as American about 17 are found on our South Atlantic coast, and species of each may at any time be discovered in Porto Rican waters. At present, however, only 7 of these 17 genera are known to have Porto Rican species.

- I. Tropical blennies, with vertebræ mostly in moderate number, usually fewer than 45; lateral line usually arched high above pectoral, if present; dorsal fin usually with at least one soft ray, none in some species of *Auchenopterus* and *Auchenistius*; anal spines little developed; ventral well developed, usually 1, 3.

a. Body sealy.

CLININÆ:

- b. Lateral line present, arched anteriorly over pectoral, becoming posteriorly median in position, or else obsolete; species ovoviviparous.

c. Scales etenoid, very rough, 33 to 40 in lateral line; dorsal divided into 3 fins. An ocular cirrus present.

GILLIAS, 147

cc. Scales eyeloid; dorsal fin not divided into 3 fins.

d. Dorsal with 6 to 20 soft rays.

e. Shoulder-girdle without upturned hook on its inner edge above. Maxillary normal, not greatly expanded. Anterior part of lateral line normally formed; usually a comb of filaments at nape. Palatines without teeth; scales moderate or small, 38 to 110 in lateral line.

f. Teeth in jaws in 1 row only; teeth usually on vomer, none on palatines; usually a comb of filaments at nape.

MALACOTENUS, 148

ff. Teeth in jaws in more than 1 row, a band of villiform teeth behind others; teeth on vomer, none on palatines. Body oblong, depth 3.5 to 4.5 in length; a filament above eye..... LABRISOMUS, 149

dd. Dorsal with 1 short soft ray only; scales large; teeth in jaws in more than 1 series; teeth on vomer, none on palatines.

g. Dorsal fin more or less deeply notched behind the third spine. First 3 dorsal spines stiff, wide set, not remote from rest of fin behind dorsal notch; anal spines short; body more elongate, snout less acute... AUCHENOPTERUS, 150

gg. Dorsal fin continuous, not notched..... PARACLINUS

bb. Lateral line absent..... AUCHENISTIUS, 151

aa. Body scaleless; species oviparous, so far as known.

h. Teeth comb-shaped, in a single row in each jaw, behind which are sometimes long canines; vomer and palatines usually toothless; lateral line usually single, with a strong arch anteriorly; dorsal fin long, continuous, or divided into 2 fins, anterior portion composed of spines, which are stiff or flexible; anal fin long, usually with 1 or 2 small spines; ventrals well developed, jugular, of 2 or 3 rays.

i. Teeth all fixed, attached to bone of jaw and not movable. Caudal fin rounded; teeth slender; gill-membranes not reduced to a small slit.

j. Gill-membranes free from isthmus, or at least forming a distinct fold across it.

k. Jaws one or both with a posterior fang-like canine, much longer than anterior teeth..... BLENNIUS

kk. Jaws without canines, teeth all equal..... SCARTELLA

jj. Gill-membranes broadly united to isthmus, gill-openings restricted to sides.

l. Jaws one or both with posterior fang-like canines..... HYPLEUROCHILUS

ll. Jaws without posterior canines; teeth equal; three articulated ventral rays.

m. Mouth small, maxillary extending scarcely beyond front of eye; head decurved in profile..... HYPSOBLENNIUS

mm. Mouth large, maxillary extending beyond vertical from middle of eye; head rather pointed in profile... CHASMODES

ii. Teeth of front of jaws all movable, implanted on skin of lips.

n. Vomer toothless. Jaws one or both with posterior fang-like canines.

o. Dorsal fin continuous..... RUPISCARTES

oo. Dorsal fin divided..... ENTOMACRODUS

nn. Vomer with a few teeth; posterior canines small..... SALARIICHTHYS

hh. Teeth unequal, not comb-like; body oblong or elongate, more or less eel-shaped, naked, or rarely with rudimentary scales; supraocular flap sometimes present. Gill-membranes united, free from isthmus; dorsal fin very long, sometimes divided into 2 fins, formed of flexible spines, which often pass gradually into soft rays; anal fin long; ventral fins thoracic or subjugular, usually not much, if any, before pectorals, composed of 2 soft rays each, spine rudimentary; caudal well developed, dorsal and anal usually more or less joined to it at base.

OPHIOBLENNINÆ:

p. Jaws with few teeth; a cirrus above eye and one above nostril; body scaleless and not eel-shaped; dorsal and anal free from caudal; dorsal fin notched.

q. Jaws each with 4 strong, hooked canines in front and a hooked posterior canine below; caudal fin forked.

OPHIOBLENNIUS

qq. Jaws each with 8 enlarged curved, conical teeth, not hooked, behind which is a single row of smaller ones; caudal fin not forked CORALLOZETUS, 152

pp. Jaws with numerous teeth, not as above; caudal fin not forked.

EMBLEMARINÆ:

r. Body not eel-shaped; dorsal and anal not joined to caudal; no scales; no cirri; no lateral line; ventrals before pectorals; teeth on palatines; caudal fin rounded. Dorsal fin very high, not notched, spines passing gradually into soft rays; jaws long, sharp at tip..... EMBLEMARIA, 153

Genus 147. *GILLIAS* Evermann & Marsh.

Body short and stout, tapering rapidly from the short, broad head to the short, compressed caudal peduncle; scales large, rough-ctenoid; lateral line complete, or nearly so, broken under last spines of middle dorsal; a broad, double-pointed tentacle above eye; dorsal fin divided into 3 parts, first of 3 short spines, second of 11 longer spines, and third of 7 rays.

This genus is closely related to *Emmancetes* Jordan & Evermann, differing in the presence of the orbital tentacle, the more complete development of the lateral line, and the larger scales.

Gillias Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 357 (*jordani*).




FIG. 95.—*Gillias jordani*.

259. *Gillias jordani* Evermann & Marsh.

Head 3.5; depth 4.3; eye 2.5; snout 3.5; maxillary 2.4; mandible 1.9; scales 2-33-3, 6 in transverse series; D. III-XII-7; A. II, 15; longest dorsal spine 1.8 in head, longest ray 1.6; longest anal ray 2.3; pectoral 0.8; ventral 1.3; caudal 1.3. Body short and stout, tapering rapidly to the short, compressed caudal peduncle; head short; snout short, blunt, concave in front of eyes; mouth small, slightly oblique, jaws equal; eye large, high up, interorbital width very narrow; a broad bifid orbital tentacle and a short nasal one; none on nape. Scales very large and rough-ctenoid; opercles and entire head rough; lateral line nearly complete, beginning immediately above base of pectoral at upper end of gill-opening and extending parallel with back to posterior part of middle dorsal fin (or for 12 scales), where there is a break, the line dropping down 3 scales, then continuing with one or two interruptions to base of caudal; belly and breast scaled; dorsals 3, first of 3 short, flexible spines, close to the second, which has 12 longer, rather stiffer, spines, separated from third by a space one-third diameter of eye; anal long and low, the membranes deeply notched between the rays; pectoral of 15 rays, broad and short, reaching posterior end of second dorsal; ventral of 2 slender rays.

Color in alcohol: Brown, body crossed by 4 broad blackish bars, one at origin of second dorsal, one under last spines of same fin, the third between second and third dorsals, and the fourth under third dorsal; an inky-black bar across caudal peduncle at base of caudal fin; head and under parts rusty; fins all barred with light and dark; caudal with a narrow light bar at base, then a black one, then a broader white one, followed by a much broader dark bar containing some white areas, the fin finally tipped with white.


Known only from Porto Rico. Two specimens of this well-marked and interesting species were obtained, the type, 1.5 inches long (No. 49368, U.S.N.M.), taken on the Cardona Light-house Reef, at Ponce, February 1, 1899, and another of about the same size taken at the same place the preceding day.

Gillias jordani Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 357, Cardona Light-house Reef, Ponce, Porto Rico.

Genus 148. MALACOTENUS Gill.

This genus is very close to *Labrisomus*, differing in dentition, the teeth in the jaws being in single series; vomer with a few teeth or with none, and none on palatines. The dorsal fin has usually a notch behind the fourth dorsal spine as well as at front of soft dorsal, but in some species the notch is obscure. Most of the species are not well known, and perhaps more than one genus is here included. About a dozen species are recognized, only four of which are as yet known from Porto Rico, though several others may occur there.

- a. Nape without filaments.
 b. Orbital filament present.
 c. D. XXI, 8; spinous dorsal not notched, first rays shortest; body elongate; snout pointed; scales large, about 38. *ocellatus*
 cc. D. XX, 12; spinous dorsal weakly notched; body rather robust *varius*
 bb. Orbital filament wanting; dorsal rays XXI, 11; spinous dorsal weakly notched; ventrals long *macropus*
 aa. Nape with a single filament; a tentacle above eye.
 d. D. XVIII, 9 *lugubris*
 dd. D. XXI, 8; scales 35 *culebræ*, 260
 ddd. D. XXI, 11; scales 45 *moorci*, 261
 aaa. Nape with a comb of slender filaments; spinous dorsal more or less notched, behind fourth or fifth spine.
 e. Orbital filament present. D. XVIII to XX, 10 to 12; vomer with teeth.
 f. Scales 43 or 44.
 g. Highest soft ray of dorsal 1.5 to 1.9 in head; dorsal without ocelli.
 h. A. II, 17; first 3 dorsal spines wider apart than the others, the first longest; anal edged with dark *gilli*
 hh. A. II, 19; first 3 dorsal spines not wider apart than the others, the second longer than the first; anal not edged with dark *puertoricensis*, 262
 gg. Soft rays of dorsal 1.2 in head; dorsal fin with 2 large black ocelli; ventral fins long, as long as head *bimaculatus*
 ff. Scales about 55; ventrals moderate, shorter than head *delalandi*, 263
 ee. Orbital filament wanting (no vomerine teeth?) *versicolor*
 aaaa. Nuchal and other filaments undescribed; a black ocellus on front of dorsal. D. XX, 11; scales 46 *biguttatus*

FIG. 96.—*Malacotenus culebræ*.260. *Malacotenus culebræ* Evermann & Marsh.

Head 3.35; depth 5; eye 4.2; snout 4.5; maxillary 2.2; mandible 1.8; interorbital 6.5; scales 2-35-11; D. XXI, 8; A. II, 18; pectoral 1.3; ventral 1.3; caudal 1.4. Body slender, compressed; head rather long, pointed, upper profile convex; mouth large, maxillary nearly reaching posterior border of orbit; lips thick, jaws equal; teeth very small, conical, a single row in each jaw; a single nasal, ocular, and nuchal filament; dorsal fin moderately high, originating above origin of lateral line, a shallow notch in front of last two dorsal spines, membrane free from caudal; anal origin under about tenth dorsal spine; caudal somewhat pointed; pectoral large, reaching anal; ventrals moderate, not reaching anus, of two rays, no spine evident; lateral line distinct throughout, running high anteriorly, where it is slightly curved, turning abruptly downward over origin of anal, thence median to base of caudal.

Color in spirits: Body everywhere mottled with dark brown, in somewhat regularly arranged blotches, a series of about nine of these at base of dorsal, barely extending upon fin; a similar series of much smaller ones at base of anal, not evident on all specimens; below the series at base of dorsal are two other series of the same blotches, less deep in color and not so well defined, extending the length of body and sometimes forming, with the upper series, more or less broken vertical bars; between the blotches a lighter shade of brown is interwoven with pale streaks of the ground-color; head nearly pale below, save some dark on chin and isthmus; two wide streaks from eye across cheek; opercle dark

brown; top of head with color of body; lips with brown and pale stripes; posterior half of maxillary pale; dorsal rather dark; caudal uniform gray or faintly barred; anal similar to dorsal in color, rays with pale tips forming a white edge; pectoral like caudal; ventrals pale.

A rather plainly marked species of different aspect from other Porto Rican species of *Malacoctenus*, but not differing widely in any important character, and as yet known only from Porto Rico. It seems most closely related to *M. lugubris*. Three specimens of about the same size; the type, No. 49369, U. S. N. M., 1.38 inches in length, from reef outside Culebra Harbor, February 9, 1899.

Malacoctenus culebræ Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 357, Culebra Island.

261. *Malacoctenus moorei* Evermann & Marsh.

Head 3.6; depth 3.7; eye 3.5; snout 3.4; maxillary 3.5; mandible 4.5; interorbital 4; scales 3-45-7, 11 in transverse series; D. XXI, 11; A. II, 20; pectoral 1 in head; ventral 1.2; caudal 1.2; longest dorsal spine 1.5, ray 1.2; longest anal ray 1.5. Body short, rather stout, compressed; head short, snout short, but pointed; mouth rather small, little oblique, gape scarcely reaching orbit; teeth in each jaw in a single series; gill-membranes broadly united across isthmus; eye small, interorbital space wide; dorsal outline rising abruptly to above eye, thence gently curved to origin of dorsal fin, and from there nearly straight to base of caudal fin; ventral outline regularly convex.




FIG. 97.—*Malacoctenus moorei*.

Color in alcohol: Light olivaceous, body crossed by about 9 or 10 dark broad vertical bars, which extend upon dorsal fin, these usually broadest above, pale interspaces therefore broadest on lower half of body; the fourth from last is a narrow dark line, the one following it is a double spot, the next narrow and indistinct, the last, at base of caudal, more distinct, followed by 3 small irregular white spots; top of head brown; side of head with fine punctulations; a dark line running forward from eye, a dark spot below eye, 2 or 3 dark blotches on anterior edge of opercle; under surface of head crossed by 3 or 4 irregular, indistinct dark lines; caudal and anal with fine dusky punctulations; pectoral and ventrals pale.

This species is close to *M. gilli*, from which it may be distinguished by the larger dorsal and anal fins, greater depth, wider interorbital, and the coloration. Known only from one specimen, 1.4 inches long, type No. 49370, U. S. N. M., collected at Culebra Island, February 11, 1899.

Malacoctenus moorei Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 358, Culebra Island.

262. *Malacoctenus puertoricensis* Evermann & Marsh.

Head 3.4; depth 3.4; eye 4; snout 3.5; maxillary 3.4; mandible 2.6; interorbital 7; preorbital 8; scales 4-44-8; D. XX, 10; A. II, 19; P. 14; V. 2; C. 13. Body short, stout, compressed; head rather long, snout long and pointed; mouth small, little oblique, maxillary scarcely reaching the front of orbit; teeth in a single row in each jaw; gill-membranes broadly united, free from isthmus; eye high up, interorbital narrow; caudal peduncle short, compressed, its least depth about 3 in head. Fins rather large; origin of dorsal over upper end of gill-opening, first spine slightly shorter than second, which is somewhat longer than third, whose length is about 2.2 in head; no notch behind third and fourth spines, all spines from third to fifteenth being about equal in length, the sixteenth and seventeenth

being somewhat shorter, the remaining three progressively longer; soft dorsal higher, its longest ray about 1.7 in head; longest anal ray 1.7; pectoral broad, 1.25 in head, reaching anal; ventral barely reaching origin of anal; a pair of slender ocular cirri, a small supraocular one, a short, slender, nasal cirrus and a few very slender ones at nape; scales large, not crowded anteriorly; lateral line well arched above pectoral.

Color in alcohol: Brown, much spotted and vermiculated with darker; top of head brown, side and under parts pale; side of body with about 6 or 7 broad, dark crossbars, broader than the paler interspaces, broadest and darkest above, and extending upon dorsal fin; under parts of body paler, more speckled; spinous dorsal with numerous small brown specks, a large black ocellus on base of 3 anterior spines, and a larger one on base of last 4 dorsal spines, being chiefly on body; soft dorsal, caudal, and anal each crossed by several series of small brown spots; pectoral and ventrals pale, pectoral with a few brown spots at base.

The above description from the type, a female, 2.5 inches long, No. 49371, U. S. N. M., obtained at Hucares, February 14. Three female cotypes, gotten at Fajardo, February 17, and one at Culebra, February 9, agree closely with the type; 2 of these, however, show faint traces of narrow horizontal lines along lower part of side.




FIG. 98.—*Malacoctenus puertoricensis*.

A male, 2.5 inches long, from Culebra, February 11, taken as one of the cotypes, may be described as follows: Head 3.5; depth 3.7; eye 3.8; snout 3.2; maxillary 3.1; mandible 2.4; interorbital 7; preorbital 6.2; scales 3-45-9; D. xx, 10; A. ii, 19; P. 14; V. 2; C. 13; longest dorsal spine 2 in head, longest ray 1.4; longest anal ray 1.5; pectoral 1; ventral 1.1; caudal 1.1. Color in alcohol: Tolerably uniform brown; crossbars on side very faint; longitudinal lines more evident than in female; throat and under parts of head mottled with white and light-brown; fins less speckled than in female, soft dorsal and anal pale, almost without spots.

Another male, 2.25 inches long, from Culebra, February 11, agrees with the large specimen just described, except that the crossbars on body are more distinct.

This species most closely resembles *M. bimaculatus* Steindachner, from which it differs in the larger head, greater depth, smaller mouth, narrower interorbital, and in the color. The tips of anal rays are not white, soft dorsal is spotted like caudal and anal, and there are no white spots on base of pectoral, as is said to be the case in *M. bimaculatus*.

Malacoctenus puertoricensis Evermann & Marsh, Report U. S. F. C. 1899 (Dec. 19), 358, Hucares, Porto Rico.

263. *Malacoctenus delalandi* (Cuvier & Valenciennes).

Head 3.4; depth 4; eye 3.5; snout 3.3; maxillary 3.3; mandible 2.6; interorbital 7; preorbital 6.2; scales 4-53-10; D. xx, 10; A. ii, 19; longest dorsal spine 2.2; longest ray 1.6; longest anal ray 1.8; pectoral 1.1; ventral 1.2; caudal lobes 1.4. Body compressed, heavy forward; head short, snout short, decurved; mouth rather large, maxillary scarcely reaching eye; gill-membranes free and broadly united across isthmus; teeth in a single series in each jaw; a few teeth on vomer, none on palatines; eye large,

above axis of body; small cirri above eye and a small nuchal fringe of filaments somewhat longer than ocular ones; scales moderate, lateral line arched above pectoral, strongly decurved posteriorly; belly naked anteriorly; first dorsal spine a little longer than second, which in turn is much longer than third, which is shorter than fourth; from the fifth to about the fourteenth increasing slightly in length, then regularly decreasing to the last but two, which is scarcely half as long as longest spines; last but one a little longer, and last about as long as fourth; soft dorsal considerably higher than spinous part; anal long, lower than soft dorsal, membrane deeply notched except among last 4 rays, which are longest.

Color, olivaceous-gray, darkest above; side crossed by about 6 irregular, broad vertical bars, first at base of pectoral, last at base of caudal fin, each of these bars consisting really of 3 large brownish blotches, middle one largest and plainest, the lower least distinct; belly and breast pale; head olivaceous; cheek and opercles crossed by darker bands; under side of head crossed by about 4 distinct dark bands separated by white of somewhat greater width; fins all uniform pale, except spinous dorsal, which has a dark blotch upon first 3 spines.

This species ranges from the West Indies to Brazil, and is also found on the west coast of Mexico north to Mazatlan, where it is the most abundant species occurring in the rock pools, if *Clinus zonifer* Jordan & Gilbert be the same species. We have compared specimens from the west coast with ours from Porto Rico and can detect no tangible differences. From Porto Rico we have 3 specimens, each about 1.75 inches long, from Ponce and Hucares.

Clinus delalandi Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 378, 1836, Brazil.

Clinus zonifer Jordan & Gilbert, Proc. U. S. N. M. 1881, 361, Mazatlan, Mexico.

Malacoctenus delalandi, Jordan & Evermann, l. c., 2358, 1898.

Genus 149. LABRISOMUS Swainson.

Body oblong, robust; head naked, short, compressed above; mouth rather large, with a row of stout, bluntish teeth in front of each jaw, behind which is a band of smaller teeth, broadest in lower jaw; teeth on vomer, no teeth on palatines; a tentacle above eye; side of neck with a tuft or series of fine filaments; dorsal fin continuous, with numerous slender spines and many soft rays, spines not very unequal; pectoral long; lateral line continuous; scales moderate or small, cycloid; shoulder-girdle without upturned hook-like process on its inner edge. Intestinal canal short, shorter than body.

The limits of this genus are not well defined. It differs from *Clinus* chiefly in the absence of an upturned spine-like process on inner edge of shoulder-girdle. This process is found on *Clinus acuminatus*, the type of the genus *Clinus*.

This genus has five American species, only one of which is yet known from Porto Rico.

- a. Scales moderate, about 70 in lateral line (so far as known); soft dorsal with 11 to 13 rays.
 b. Dorsal spines 16; anal rays 29; tentacles on nape..... *herminier*
 bb. Dorsal spines 18; no teeth on palatines; first ray of dorsal not longest; orbital tentacle well developed; nape with a conspicuous comb of fringes.
 c. Vomer with a cluster of small teeth..... *nuchipinnis*, 264
 bbb. Dorsal spines 20; teeth on palatines (?); first dorsal spine longest..... *bucciferus*
 aa. Scales very small, about 110; a comb of fringes at nape; first dorsal spines low; head with yellow spots.
microlepidotus

264. *Labrisomus nuchipinnis* (Quoy & Gaimard).

(PLATE 46.)

Head 3.5; depth 3.5; D. XVIII, 12; A. II, 17; scales 70. Body oblong, rather robust; head naked, thick, short, not very obtuse anteriorly, compressed above; mouth rather large, the maxillaries not prolonged backward, extending to opposite posterior part of eye, 2.5 in head; front teeth of jaws conic, strong, behind them a band of villiform teeth, broadest in lower jaw; vomer with a patch of smallish teeth; eye large; interorbital space very narrow; each side of neck with a long series of hair-like filaments, nearly as long as eye; orbital tentacle short and broad, multifid; nostril with a tufted barbel; lower jaw slightly projecting, its posterior teeth sometimes recurved; pectoral a little shorter than head, reaching vent. Dorsal spines rather slender, the 3 anterior spines scarcely shorter than the others, all the spines lower than soft rays; dorsal fin commencing near nape, spinous portion long; soft rays higher than the spines; caudal small; pectoral rather large; ventrals moderate; gill-membranes broadly united, free from isthmus; lateral line complete, high anteriorly, then abruptly decurved; membranes of vertical fins scaly; scales not very small, cycloid.

Color in life: Body mottled yellow-olivaceous, crossed by about 7 broad black vertical bars, the first extending from origin of dorsal to upper edge of opercle, second from fifth dorsal spine to base of pectoral, next 3 extending entirely across side, sixth from base of soft dorsal to median line, and seventh consisting of 2 or 3 black blotches upon upper part of caudal peduncle; a round jet-black spot as large as pupil upon opercle, surrounded by a narrow line of orange or yellow; cheek and upper parts of head like side; throat and breast pale-rosy; branchiostegals white with about 4 series of black spots on membranes; belly pale; dorsal fin yellowish at base, pale bluish-white on outer half, both spinous and soft portions with numerous small reddish spots; a round black spot between first and third dorsal spines; caudal pale; anal yellowish at base, paler on outer half; pectoral pale with numerous small reddish spots; ventral pale; nuchal filaments reddish-brown; iris reddish-brown.

The color on this species seems to be subject to considerable variation; we have the following note on a specimen obtained at Mayaguez: Side greenish-gray, with about 6 dark crossbars extending upon dorsal fin, bifid above; cheek and throat vermilion; inside of the mouth and breast vermilion, becoming golden posteriorly; cheek with many pale-blue spots; similar spots on top of head and base of pectoral and paler ones on lower part of side; a large black spot bordered by white on opercle, bounded on front and above by rich metallic-rosy; dorsal lemon barred with brown; anal lemon-yellow with pale darker bars; caudal plain lemon, faintly barred; pectoral lemon, with faint bars; ventral rosy; cirri barred with pale-yellow and dark-brown; 6 dark bars on spinous dorsal and 2 on soft, second one faint; a large black spot between first and third dorsal spines; iris with purplish-red spots.

West Indies, north to southern Florida, south to Brazil; generally common in rocky pools; also recorded from the Canary Islands. Length 6 to 8 inches. Common in Porto Rico, specimens being in the collection from San Antonio Bridge, Palo Seco, Mayaguez, Puerto Real, Ponce, Arroyo, Fajardo, Isabel Segunda, and Culebra.

Clinus nuchipinnis Quoy & Gaimard, Voy. Uranie et Physicienne, Zool., 255, 1824, Brazil.

Clinus pectinifer Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 374, 1836, Bahia.

Lepidoma cirrhosum De Kay, N. Y. Fauna: Fishes, 41, 1842, Florida.

Clinus canariensis Valenciennes, in Webb & Berthelot, Poiss. Iles Canaries, 60, 17, fig. 3, Canary Islands.

Labrisomus nuchipinnis, Jordan & Evermann, l. c., 2362, 1898.

Genus 150. AUCHENOPTERUS Günther.

Body moderately elongate, compressed, covered with rather large, cycloid scales; head shortish, naked, snout rather pointed; cheek full; mouth moderate, with a band of conical teeth in jaws and about one series on vomer, none on palatines; lower jaw prominent; gill-membranes united, free from isthmus; upper surface of head with filaments. Dorsal fin composed of stiff spines, with but a single soft ray, which is lower than the spines; first 3 spines more or less separated from others, stiff and rather wider set, sometimes higher than the others; anal fin low, with 2 short spines; ventrals jugular, well developed; pectoral broad; lateral line complete, strongly curved anteriorly.

An inhabitant of the warm seas. This genus differs from *Cristiceps* in the large scales and in having but 1 soft ray in dorsal fin. Of the dozen recognized American species of *Auchenopterus*, 5 are known from Porto Rico, and some of the others may occur there.

CORALLICOLA:

a. First 3 spines of dorsal forming a separate fin, being much higher than any of the spines in the posterior part of fin; snout rather acute.

b. Scales 33; dorsal with 1 ocellus, anal with none; a black crossbar at base of caudal; a yellow spot behind eye; snout pointed..... *nigripinnis*

bb. Scales 37 or 38.

c. First dorsal spine shorter than second; snout slender, very acute; caudal pale; dorsal with 2 ocelli, anal with 1; D. III-XXII, 1..... *marmoratus*.

AUCHENOPTERUS:

aa. First 3 spines of dorsal scarcely forming a separate fin, none of them higher than posterior spines; snout not very acute; anal without ocellus.

d. Caudal fin pale, usually with a dark bar at its base; a notch between third and fourth dorsal spines.

e. Dorsal spines about 31.

f. Scales 31 to 36.

g. Membrane of third dorsal spine joining fourth spine near its base..... *affinis*

gg. Membrane of third dorsal spine joining fourth spine near its tip.

h. Caudal fin white, with a broad, black bar on its base; body dark, no crossbars; scales 2-34-3, 7 in transverse series; D. XXX, 1..... *albicaudus*, 265

hh. Caudal fin mottled, a narrow white bar near base, before which is a broader dark bar; body with about 7 dark crossbars; scales 2-34-3, 7 in transverse series; D. XXIX, 1..... *fajardo*, 266

- hhh.* Caudal fin pale, no dark bar at base; body pale rosy, no crossbars; scales 2-33-3, 7 in transverse series; D. xxx, 1 *rubescens*, 267
- cc.* Dorsal spines about 28; membrane of third dorsal spine joining fourth spine near its tip; body with distinct crossbars.
- i.* Dorsal xxx, 1, with a distinct ocellus; scales about 2-31-3, 7 in transverse series..... *fasciatus*, 269
- ii.* Dorsal xxviii, without ocellus; scales 2-30-3, 7 in transverse series..... *cingulatus*, 268
- dd.* Caudal fin black; body chiefly black; head mottled with whitish; membrane of third dorsal spine joining fourth near its summit, fin not notched; dorsal spines 30; dorsal with 2 ocelli *nox*

265. *Auchenopterus albicaudus* Evermann & Marsh.

Head 3.2; depth 4; eye 4; snout 4.1; maxillary 2.2; mandible 1.6; interorbital 5.3; D. xxx, 1; A. ii, 17; pectoral 1.4; ventral 1.5; caudal 1.6; branchiostegals 6; scales 2-34-3, 7 in transverse series.* Body rather short, compressed; dorsal outline not elevated; head moderate, not broad; snout short, pointed; mouth large, oblique, maxillary extending to below middle of eye; lips broad, prominent; a band of conical teeth on each jaw, those on side somewhat enlarged and recurved; a patch of teeth on vomer, none on palatines; gill-membranes broadly united, free from isthmus; eye large, high up; nasal, supraocular, and nuchal region with fringed tuft-like cirri; a considerable notch between third and fourth dorsal spines, but not reaching base of membrane; longest anterior spine scarcely as long as those of posterior portion; scales large, reduced anteriorly; lateral line anteriorly separated from dorsal fin by only one scale; head naked.




FIG. 99.—*Auchenopterus albicaudus*.

Color, uniform dark brown on head and body, no dark crossbars; dorsal brown, mottled with lighter, narrowly edged with white; a black spot upon anterior 3 or 4 spines and a large black ocellus upon posterior portion of fin between twenty-second and twenty-fourth spines; anal rather darker, with narrow white edge; caudal peduncle black, fin abruptly white at base, entire fin being clear white, without specks; pectoral black at base, then barred with white and dark; ventral black at base, outer two-thirds barred with black and white.

This species seems to be related to the Pacific coast *Auchenopterus integripinnis*, which it closely resembles, differing in the larger scales, deeper body, and coloration.

Known only from the type, seined at Arroyo, Porto Rico, February 4.

Auchenopterus albicaudus Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 360, Arroyo, Porto Rico.

266. *Auchenopterus fajardo* Evermann & Marsh.

(PLATE 47.)

Head 3.25; depth 4.8; eye 4.2; snout 4.8; maxillary 1.7; mandible 1.5; interorbital 5.5; scales 2-34-3, 7 in transverse series; D. xxix, 1, the longest spine 2.3 in head; A. ii, 17; pectoral 1.4; ventral 1.7; caudal 1.4. Body elongate, strongly compressed posteriorly; head moderate, little compressed; mouth large, the long and slender maxillary reaching beyond posterior border of orbit; jaws subequal;

* In our descriptions of species of this family the scales are counted from near front of dorsal to and including lateral line, then in lateral line, then from origin of anal upward and backward to lateral line, the transverse series being counted from origin of anal upward and backward to dorsal fin. The half scale at base of dorsal and anal is not counted.

teeth of upper jaw conical and sharp, in a patch in front, becoming one row posteriorly; teeth in lower jaw similar, but fewer and weaker; vomerine teeth in two series. Nasal, ocular, and nuchal filaments present, all but the nasal about 5-branched. Dorsal origin over edge of preopercle, first 4 spines graduated, fourth shortest, thus forming a notch; dorsal ending with an unbranched soft ray, the joints visible under a strong lens; membrane of dorsal joined low to caudal; anal origin under eleventh dorsal spine and decurved portion of lateral line; pectoral reaching past front of anal; ventral moderate, of 3 rays, innermost shorter and slenderer.

Color in spirits: Body and head light-reddish, becoming a little paler posteriorly; body with traces of 6 or 8 dark vertical bars extending on the fins, their margins ill-defined; breast pale, 2 dark-reddish bars downward and backward from eye across upper and lower edge of cheek to opercle; maxillary blotched with dark; upper lip and tips of both jaws dark; lower part of head spotted with dark; a row of about 5 small dark spots on edge of preopercle; iris pink; dorsal and anal fins gray, except for extensions of dark bars of body and a few white spots on dorsal; a distinct ocellus on twenty-second, twenty-third, and twenty-fourth dorsal spines and their membranes; base of caudal gray, like ground-color of dorsal and anal; posterior part of caudal with gray mottlings on rays only, this portion separated from the basal part by a space without pigment on rays or membrane, making a distinct vertical bar; pectoral and ventrals mottled.

A handsomely colored blenny, of which the collection contains but one specimen, the type, No. 49376, U. S. N. M., 1.63 inches long, taken at Fajardo, Porto Rico, February 17, 1899.

Auchenopterus fajardo Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 361, Fajardo, Porto Rico.




FIG. 100.—*Auchenopterus rubescens*.

267. *Auchenopterus rubescens* Evermann & Marsh.

Head 3.4; depth 5; eye 5; snout 3.8; maxillary 2.6; interorbital 5.8; scales 2-33-3, 7 in transverse series; D. xxx, 1; A. ii, 18; pectoral 1.5; ventral 2; caudal 1.4. Body slender and compressed; head moderate, somewhat compressed above; snout pointed; mouth moderate, jaws equal, maxillary about reaching front of pupil; lips, especially upper, prominent; teeth small, conical, and sharp, in both jaws, in a numerous patch on front of upper jaw, fewer on sides; in lower jaw less numerous in front, a long single row of somewhat stronger teeth on sides; eye not large; a small nasal flap, and a 3 or 4 branched filament over eye and one at nape; scales rather large and regularly arranged; dorsal fin with a notch behind third spine, and with one unbranched soft ray at its end, membrane joined to caudal; origin of anal under eleventh dorsal spine.


Color in spirits: Everywhere a nearly uniform faded pink, save breast and lower side of head, which are paler; a small, inconspicuous dark round spot on dorsal fin, at twenty-third and twenty-fourth spines, a little nearer the base than margin, and made up of very small black punctulations. Indications of a yellow tinge on front of dorsal and base of anal in life; fins otherwise all pale.

Known only from the type, 1.3 inches long, collected at Puerto Real, Porto Rico.

Auchenopterus rubescens Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 360, Puerto Real, Porto Rico.

268. *Auchenopterus cingulatus* Evermann & Marsh.

Head 3; depth 4.4; eye 5; snout 4.2; maxillary 2.2; interorbital 6; scales 2-30-3, 7 in transverse series. D. xxviii, the longest spines 2.75 in head; A. ii, 16, the longest ray 2.75 in head; pectoral 1.3; ventral 1.8; caudal 1.6. Body rather long and slender, strongly compressed; head large, little compressed; snout moderately sharp; mouth large, maxillary reaching posterior border of eye, lips heavy, jaws subequal or lower very slightly projecting; teeth conical and sharp, in more than one row in each jaw, most numerous in front; a patch on vomer; a nasal filament, a 3 or 4 branched supraocular filament, and a 4-branched nuchal filament, branches of latter each with a dark dot on its anterior surface; dorsal originating over edge of preopercle, of spines only, second slightly longer than first; second, third, and fourth graduated, fourth comparatively short, thus forming a notch partly separating first spines from rest of fin; dorsal membrane joined low with caudal; anal free from caudal, about as high as dorsal, its thirteenth and fourteenth rays longest; first anal spine under tenth or eleventh dorsal spine; caudal rounded, shorter than head, of about 13 rays; pectoral large, reaching anal, of 12 rays; ventrals moderate, of 2 rays, spine not evident; lateral line running high to eleventh dorsal spine, here abruptly decurved two rows of scales, thence median to base of caudal.

FIG. 101.—*Auchenopterus cingulatus*.

Color in spirits: Body and head pale-yellow; body with 5 heavy dark-brown vertical bars, each about 4 rows of scales wide, extending on vertical fins; membrane of anterior dorsal spines, opercle, occipital, and scapular region blotched with same color; a dark bar backward and downward from eye across cheek, rather more than one-half width of eye; top of head between and behind eyes darkened; preorbital, maxillary, lips, and under part of head thickly punctulate with dark; dorsal and anal barred with extensions of the wide, dark body-bars, and with alternating narrower pale interspaces; caudal mottled or irregularly barred with grayish, its base with the plain pale-yellow ground-color, which is sharply separated from rest of fin by a curved dark line; posterior half of pectoral with dark bars formed of dots on rays, first bar plainest; basal half of pectoral pale; ventral with basal portion dark, rest barred like pectoral.

A pretty and strongly marked blenny, known only from Porto Rico; four specimens obtained from the coral reefs at Ponce, and one at Puerto Real. The type, No. 49375, U. S. N. M., from Ponce, is 0.8 inch long, and none of the cotypes exceeds 1 inch.

Auchenopterus cingulatus Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 361, Ponce, Porto Rico.

269. *Auchenopterus fasciatus* (Steindachner).

Head 3.5; depth 4.6; eye 4.5; snout 4.2; maxillary 2.3; mandible 2; interorbital 5.4; scale 2-31-3, 7 in transverse series; D. xxx, 1; A. ii, 17; P. 1.3 in head; V. 1.4; C. 1.4. Body slender, compressed, head narrow, snout pointed; mouth rather large, lower jaw slightly projecting, maxillary reaching posterior edge of pupil; gill-membranes broadly united across isthmus; eye small, above axis of body; profile rising gently and regularly from tip of snout to origin of dorsal fin, from which it continues in a nearly straight line to caudal peduncle; a small ocular cirrus and a small one on side of nape; fourth and fifth dorsal spines shorter than those before or after, others of approximately equal length; anal rays about as long as dorsal spines, the tips projecting beyond membrane; scales large, cycloid; belly,

breast, nape, cheek, and opercles scaled; lateral line complete, arched above pectoral, running on third row of scales from back for about 13 scales, where it drops to fifth row.

Color in alcohol: Somewhat rosy-brown, with faint traces of darker vertical bars; anterior part of eye dark, head uniform rosy-brown without markings; dorsal fin with 8 broad, dark bars, the first on first 2 or 3 spines and forming a small black spot near their tips, seventh between twentieth and twenty-fourth spines and containing a distinct ocellus, probably blue in life; anal with 5 broad black bars, but no ocellus; a dark bar across base of caudal, rest of fin pale; pectoral faintly and finely barred with dark.

This blenny reaches a length of 2 inches and is known from southern Florida (Cards Sound, Key West) and Porto Rico. One specimen, 1.25 inches long, obtained at Hucars, February 14, agrees in the main with current descriptions of this species. The dorsal spines, however, are more numerous and the color seems to be somewhat different. Without more material we hesitate to regard these as being of specific value.

Cremnobates fasciatus Steindachner, Ichth. Beiträge, V, 176, 1876, Florida Straits.

Auchenopterus fasciatus, Jordan & Evermann, l. c., 2373, 1898.

Genus 151. *AUCHENISTIUS* Evermann & Marsh.

This genus has the form of *Auchenopterus* and suggests that genus strongly. It differs in the absence of a lateral line, in the much smaller scales, in the absence of a notch at front of dorsal fin, and in the union of the membrane of the anal fin with that of the caudal. Only one species known.

Auchenistius Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 359 (*stahli*).




FIG. 102.—*Auchenistius stahli*.

270. *Auchenistius stahli* Evermann & Marsh.

Head 5; depth 6.5; eye 4.8; snout 6; maxillary 2.8; scales about 58, about 13 in transverse series; D. XLII or XLIII; A. I or II, 23 or 24; pectoral 2.5; ventral 2.2; caudal 1.3. Body elongate, somewhat compressed, especially posteriorly, dorsal and ventral outlines alike; head small, upper profile straight and descending; snout moderate, pointed; mouth large, maxillary reaching to or beyond middle of eye; jaws equal, heavy and projecting; teeth in lower jaw conical, short and strong, slightly recurved, in one row; teeth in upper jaw similar to those in lower, but a small patch of smaller teeth in front of jaw behind main row; teeth on vomer; gill-membranes joined to isthmus; nostrils with short tubes, a single flap above each eye and one on each side of nape; dorsal fin long, of spines only; last four spines somewhat longer than the preceding, forming a shallow notch, a feature not present in all examples; anal origin about midway between tip of snout and tip of caudal, fin similar to dorsal in shape, somewhat higher in type, but a trifle lower in cotype; membrane of dorsal and anal joined to caudal; caudal small, pointed; pectoral small, of 8 rays; ventral small, of 2 rays.

Color in spirits: Body everywhere with a very slight yellowish tinge, in some specimens a faded gray; one specimen has traces of 10 or 12 dark crossbars; fins all pale, in one case the dorsal and anal dark-edged.

This species is known only from the type, 1.2 inches long, No. 49372, U. S. N. M., from Ponce, February 1, 1899, and 13 cotypes, 8 from the coral and algae on the reefs at mouth of Culebra Harbor, February 11, and 5 from Puerto Real.

Named for Dr. A. Stahl, of Bayamon, Porto Rico, who, under many difficulties, has made considerable collections of the natural-history objects of Porto Rico.

Auchenistius stahli Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 359, Ponce, Porto Rico.

Genus 152. *CORALLIOZETUS* Evermann & Marsh.

Body slender and strongly compressed, without scales; head large, subcylindrical, bluntly pointed; mouth large; teeth not hooked, about 8 enlarged conical ones in front of each jaw, smaller ones behind; vomer with teeth; dorsal fin with a notch between rays and spines; the membrane slightly connected with caudal; caudal fin rounded; pectoral large, ventrals small and inserted slightly in advance of pectorals. A strongly marked genus, conspicuous in appearance by its heavy head and thin body; probably related to *Ophioblennius*, from which it is technically separated by absence of hooked canine teeth, convex caudal, and entire absence of a lateral line. Only one species known.

Coralliozetus Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 362 (*cardonæ*).

271. *Coralliozetus cardonæ* Evermann & Marsh.

Head 4; depth 5.6; eye 4; snout very short; maxillary 2; D. xvii, 11; A. 21; pectoral 1.3; ventral 1.8; caudal 1.4. Body scaleless, slender, much compressed; head large and heavy, not compressed nor depressed; snout very short and blunt; mouth large, horizontal, low in position, maxillary reaching far beyond eye; eyes small, close together, placed high and well forward; teeth conical, in a patch on the front of each jaw, an outer row of about 8 teeth (4 on a side) in each jaw, much enlarged; a single row of smaller teeth on sides of each jaw; teeth on vomer; a small flap at nostril and two short filaments




FIG. 103.—*Coralliozetus cardonæ*.

above eye, one much the smaller; no appendages at nape. Dorsal fin long and high, of slender, flexible spines, and longer, soft rays, a notch between fourth and fifth spines, last ray slightly connected with caudal; anal longer and lower than soft dorsal, free from caudal; caudal rounded; pectoral large, nearly as wide as body, reaching anal or beyond; ventral small, inserted before pectoral, of 3 rays, innermost very slender.

Color in spirits: Body dark-red, much paler in one specimen; head everywhere bluish-black, this color dusted upon body, particularly on anterior portion; a pale-gray bar downward and backward across cheek; fins pale, except ventrals and front of dorsal, which have color of head; a row of small rosy spots along bases of anal rays, seemingly in the flesh; sometimes a similar fainter row along base of dorsal.

Known only from Porto Rico; three specimens, 0.87 to 1 inch in length, taken on the coral reefs at Ponce on three successive days; type, No. 49377, U. S. N. M., 1 inch long, collected February 1, 1899.

Coralliozetus cardonæ Evermann & Marsh, Rept. U. S. F. C. 1899 (Dec. 19), 362, reef at the Cardona Light-house, Playa de Ponce, Porto Rico.

Genus 153. *EMBLEMARIA* Jordan & Gilbert.

Body slender, not eel-shaped, compressed, scaleless. Ventrals present, jugular, each of 1 spine and 2 soft rays. A single high dorsal fin beginning on nape and extending to caudal, with which it is not confluent; no notch between spinous and soft rays. Head cuboid, compressed, narrowed anteriorly. Symphysis of lower jaw forming a very acute angle. A single series of strong, blunt, conical teeth on each jaw, and on vomer and palatines. Vomer and palatine teeth larger, their series continuous, parallel to series in upper jaw. No cirri at nape; sometimes a cirrus on upper part of eyeball. Gill-openings very wide, membranes broadly united below, free from isthmus. Lateral line obsolete.

This genus bears some resemblance to *Bleinnius*, but the dentition is entirely different, approaching that of *Chenopsis*. An inhabitant of tropical America, in rather deep water; 4 species known, 2 of which are Atlantic.

- a. Eye without cirrus.
 b. Depth 5 in length; dorsal rays 33; ventrals not pure white..... *atlantica*
 aa. Eye with a long cirrus on eyeball above pupil.
 c. Maxillary longer, reaching posterior edge of orbit; body without dark crossbar; ventrals pale; tip of caudal not black..... *pandionis*, 272

272. *Emblemaria pandionis* Evermann & Marsh, new species.

Head 3.7; depth 5.8; eye 3.6; snout 5; maxillary 2; mandible 1.8; D. xvii, 18; A. ii, 23; P. 13; V. 1, 2; C. 13. Body slender, tapering, greatly compressed; head comparatively heavy; snout short, decurved; mouth large, horizontal, maxillary reaching posterior border of eye; each jaw with a patch of teeth in front, outer enlarged, bluntly conic, slightly incurved, these continued laterally on upper jaw in a single series of similar but smaller teeth, and on lower jaw in a single series of similar but




FIG. 104.—*Emblemaria pandionis*.

smaller, blunter teeth; palatines each with a row of blunt teeth; vomer probably without teeth; no teeth on tongue. Fins moderate, rays not filamentous; first 17 rays of dorsal simple, not jointed nor branched, remaining rays jointed and elevated, longest a little greater than snout and eye, the two anterior rays of anal not jointed nor branched; pectoral broad, short, its length 1.3 in head; ventral longer, filamentous, 1.2 in head; caudal short, pointed, 1.8 in head. An unbranched ocular cirrus on upper part of eye, its length about one-third that of eye; nasal tube slender and simple.

Color in alcohol: Pale flesh-color; middle of side with a series of about 15 small brown blotches, largest anteriorly; above these and along base of dorsal are 2 or 3 irregular series of smaller dark spots, those along base of dorsal quite black; a similar series along base of anal; head spotted like body; dorsal fin with numerous dark spots and blotches, most distinct anteriorly; caudal and anal fins somewhat dusky; pectoral and ventral pale; a few small brownish specks on throat and on breast in front of ventrals; a very dark-brown, almost black, oblong spot, placed obliquely on lower part of side above tenth anal ray, a narrow bar of same color just in front of it; this conspicuous color-marking present on left side only.

Known only from the type (No. 49535, U. S. N. M.), 1.5 inches long, caught in the tangle at Fish Hawk station 6084, between Vieques and Culebra islands, in 11 fathoms, and one cotype of same size taken in the dredge at Fish Hawk station 6086, 8.5 miles northeast from Isabel Segunda, Vieques Island, in 14.75 fathoms, the bottom at each station being coral, sand, and shells.

This interesting species is related to *E. oculocirris* Jordan, from La Paz, Lower California, from which it seems to differ chiefly in the longer maxillary and in the coloration.

Named for the U. S. Fish Commission steamer *Fish Hawk*, *Pandion* being the generic name of the fish-hawk or osprey, *Pandion carolinensis*.

Family LXXII. FIERASFERIDÆ. The Pearl-fishes.

Body elongate, compressed, tapering into a long and slender tail; no scales; teeth cardiform, on jaws, vomer, and palatines; canine teeth often present; no barbels; lower jaw included; vent at throat; gill-membranes somewhat united, free from isthmus; no pseudobranchie; no pyloric cæca; vertical fins very low, confluent, without spines; no ventral fins; pectoral fins present or absent.

Small shore-fishes of tropical seas, often living in shells of mollusks, echinoderms, etc., being especially often commensal with the pearl oyster and with the larger *Holothuria*. Only one genus in American waters.

Genus 154. FIERASFER Cuvier.

Gill-membranes little connected, leaving isthmus bare. No distinct caudal fin; pectoral fins developed.

The species of this genus are not well known, and their characters and nomenclatures are uncertain. It is not unlikely that the American species are all reducible to one, *Fierasfer affinis* or *dubius*, but our scanty material will not justify us in taking this view. (Jordan & Evermann.) Only one species known from Porto Rico.

273. *Fierasfer bermudensis* (Jones). Pearl-fish.

Head 8.5 in length; eye 4, longer than snout; mouth large, the maxillary reaching beyond orbit; pectoral 2.5 in head. Teeth small, acute, uniserial, 3 in a line on vomer; palatine teeth small. Color pale-brownish, a bluish streak crossing nape between opercles, 4 pale points on back. Vertebrae 100.

A small fish (length 3 to 6 inches) found in shallow water on coral shores in the West Indies. Two specimens, each about 4 inches long, obtained by us, one at Mayaguez, the other at Puerto Real. They certainly belong to the species described from Bermuda, which may not be different from Günther's *F. affinis* or Putnam's *F. dubius*.

?? *Fierasfer affinis* Günther, Cat., IV, 381, 1862, no locality given.

? *Fierasfer dubius* Putnam, Proc. Bost. Soc. Nat. Hist. 1874, 344, Pearl Islands.

Lefroyia bermudensis Jones, Zoologist, IX, 1874, 3838, Bermuda.

Fierasfer bermudensis, Jordan & Evermann, l. c., 2497, 1898.

Family LXXIII. PLEURONECTIDÆ. The Flounders.

Body strongly compressed, oval or elliptical in outline; head unsymmetrical, cranium twisted, both eyes being on same side of body, which is horizontal in life, the eyed side being uppermost and colored, the blind side lowermost and usually plain. In the very young fish the bones of the head are symmetrical, 1 eye on each side, and the body is vertical in the water. In most species the cranium becomes twisted, bringing the upper eye over with it. Eyes large, well separated. Mouth small or large, the dentition various, teeth always present; premaxillaries protractile; no supplemental maxillary bone; pseudobranchie present. Gills 4, a slit behind the fourth; lower pharyngeals separate; no air-bladder; preopercle with its margin usually distinct, not wholly adnate or hidden by skin of head; vent not far behind head, viscera confined to anterior part of body. Scales various, rarely absent, usually small. Lateral line usually present, extending on caudal fin, sometimes duplicated or wanting. Dorsal fin long, continuous, of soft rays only, beginning on head; anal similar, shorter; caudal various, sometimes coalescent with dorsal and anal; pectorals inserted rather high, rarely wanting; ventrals under pectorals, usually of several soft rays, one of them sometimes wanting.

Fishes mostly carnivorous, inhabiting sandy bottoms in all seas, some species ascending rivers. Many of them are important food-fishes. Genera about 55; species nearly 500.

Subfamilies of *Pleuronectidæ*.

- A. Ventral fins symmetrical, similar in position and in form of base, that of colored side not extended along ridge of abdomen.
- a. Mouth nearly symmetrical, dentition nearly equally developed on both sides, gape usually but not always wide. (Halibut tribe.) HIPPOGLOSSINÆ
- aa. Mouth unsymmetrical, jaws on eyed side with nearly straight outline, bones on blind side strongly curved; teeth chiefly on blind side.
- b. Eyes and color on right side (with occasional exceptions). (Flounder tribe.) PLEURONECTINÆ
- AA. Ventral fins unsymmetrical, dissimilar in position and usually also in form, that of eyed side being extended along ridge of abdomen. Eyes and color on left side. (Turbot tribe.) PSETTINÆ

HIPPOGLOSSINÆ. Halibut Tribe.

Large-mouthed flounders with the ventral fins symmetrical.—Mouth symmetrical, the jaws and dentition nearly equally developed on both sides; gape usually wide, maxillary more than one-third length of head. Lower pharyngeals narrow, usually with but 1 or 2 rows of sharp teeth; teeth in jaws usually acute. Eyes large; edge of preopercle free. Pectoral and ventral fins well developed, the ventral fins similar in position and in form of base, that of eyed side not being attached along ridge of abdomen. Septum of gill-cavity without foramen.

The only genera of *Hippoglossinæ* having species ranging as far south on our Atlantic coast as Florida are *Paralichthys*, *Ancylopssetta*, *Notosema*, and *Gastropssetta*. Though no species of this group is known from Porto Rico, we give a key to these 4 genera, as some or all of them may be hereafter found in Porto Rican waters.

- a. Dorsal fin beginning in advance of eye; teeth sharp, uniserial or smooth.
- b. Scales weakly ciliated; caudal fin with a distinct peduncle; mouth large; teeth unequal, some of anterior canine-like; gillrakers rather long and slender; no dorsal lobe nor produced ventral rays; vertebræ 35 to 41. PARALICHTHYS
- bb. Scales very strongly etenoid on both sides of body; mouth smallish, with small, sharp teeth; anterior rays of dorsal more or less exerted, thus forming a more or less distinct lobe; gill-membranes considerably united; gillrakers short and broad; caudal peduncle short; left ventral produced; vertebræ (in *A. quadrocellata*) 9 + 26 = 35. Lateral line with its tubes simple, not branched.
- c. Body broad, ovate, depth more than half length; dorsal lobe and left ventral moderately produced. ANCYLOPSETTA
- cc. Body elliptical, depth not more than one-half length; dorsal lobe and left ventral greatly produced. NOTOSEMA
- bbb. Scales entirely smooth; caudal peduncle short; mouth small; gillrakers short and thick; dorsal with an anterior lobe; left ventral elongate. GASTROPSSETTA

PSETTINÆ. Turbot Tribe.

Large-mouthed flounders, with ventral fins unsymmetrical.—Mouth symmetrical, the dentition nearly equally developed on both sides; gape usually wide (narrow in *Platophrys*, *Etropus*, etc.), maxillary commonly more than one-third length of head; lower pharyngeals narrow, each with one or more rows or a narrow band of small, sharp teeth; teeth in jaws acute; eyes not minute; pectorals and ventrals usually well developed; edge of preopercle free; ventral fins dissimilar in form or in position, that of left or eyed side inserted on ridge of abdomen, its base extended along this ridge, its rays more or less wide apart; caudal fin rounded or subtruncate; no accessory lateral line; anal spine usually weak or obsolete; a pelvic spine sometimes developed; vertebræ in moderate or small number, 31 to 45. Body sinistral.

Species chiefly tropical or subtropical in distribution. Of the 10 recognized American genera of this subfamily, 4 have Porto Rican representatives. In fact, all the species of flounders thus far known from Porto Rico belong in this group.

- a. Pectoral fin present on each side; septum of gill-cavity below gill-arches without foramen; a deep emargination near isthmus; ventral fins free from anal.
- b. Vomer with teeth; lateral line with a strong arch in front; teeth subequal, in villiform bands; body broadly ovate; caudal fin subsessile; interorbital area broad; scales small, cycloid; gillrakers long and slender; anterior dorsal rays produced; vertebræ 36. LOPHOPSETTA
- bb. Vomer toothless; ventral fins free from anal; caudal fin subsessile.
- c. Lateral line with a distinct arch in front; teeth small, uniserial, or imperfectly biserial.
- d. Interorbital space more or less broad, deeply concave, at least in males; form broad-ovate; gillrakers short and thick.
- e. Scales small, ctenoid, adherent, 75 to 100 or more; anterior rays of dorsal not elevated; pectoral of left side usually filamentous in male; vertebræ (in *P. lunatus*) 9 + 30 = 39. PLATOPHRYS, 155
- dd. Interorbital space a narrow ridge; dorsal not elevated in front.
- f. Gillrakers slender; right ventral elongate; scales ctenoid. TRICHOSETTA
- cc. Lateral line without arch in front.
- g. Teeth in upper jaw biserial, in lower uniserial, front teeth of upper jaw enlarged; vertebræ 35 or 36; gillrakers short; interorbital space broad in male. SYACIUM, 156
- gg. Teeth in each jaw uniserial; interorbital space very narrow, ridges coalescing between eyes.
- h. Mouth not very small, maxillary more than one-third length of head.
- i. Gillrakers very short and thick, tubercle-like.
- j. Scales cycloid, small, and firm. CYCLOPSETTA
- ii. Gillrakers slender, of moderate length; scales thin, deciduous, ciliated; vertebræ 34 to 40. CRITHARICHTHYS, 157
- hh. Mouth very small, teeth subequal, maxillary less than one-third length of head; scales thin; teeth uniserial; vertebræ 9 + 25 = 34. ETROPUS, 158
- aa. Pectoral fin of blind side wanting; eyes very close together; caudal fin subsessile; teeth small, uniserial; mouth moderate; lateral line of eyed side arched, that of right side nearly straight; dorsal fin beginning on snout, its anterior rays not exerted, its rays all simple and very numerous; gillrakers few and feeble; scales small; body thin, very elongate; vertebræ (in *M. sessilicauda*) 43; (deep-sea flounders). MONOLENE

Genus 155. **PLATOPHRYS** Swainson.

Eyes and color on left side. Body ovate, strongly compressed; mouth of the large type, but comparatively small; maxillary one-third or less length of head; teeth small, subequal, in one or two series; no teeth on vomer or palatines. Interorbital space broad and concave, broadest in adult males. Gillrakers moderate. Dorsal fin beginning in front of eye, all its rays simple; ventral of colored side on ridge of abdomen; caudal convex behind; pectoral of left side with usually one or more filamentous rays, longest in male. Scales very small, ctenoid, adherent; lateral line with a strong arch in front. Coloration usually variegated. The sexual differences are greater than usual among flounders, and the different sexes have often been taken for different species. As a rule, in males, pectoral fin on left side is much prolonged, interorbital area is much widened and very concave, and there are some tubercles about snout and lower eye. The young fishes, as is usually the case, resemble adult females. Lately, Dr. Emery has shown that the larval flounder, known as *Peloria heckeli*, is in all probability the young of *Pleuronectes podas*. The generic name, *Coccolus*, based on forms slightly more mature than those called *Peloria*, probably belongs here also. We have seen no larval forms so young as those which have been described as *Peloria heckeli*. We have, however, examined small transparent flounders, one with eyes quite symmetrical, taken in the Gulf Stream, and another with eyes on left side, taken at Key West. Both these may be larvæ of *Platophrys ocellatus*. The figures published by Emery seem to make it almost certain that the corresponding European forms belong to *P. podas*, although some doubt as to this is expressed by Facciola.

The species of *Platophrys* are widely distributed through warm seas, no tropical waters being wholly without them; all are extremely closely related and are distinguished with difficulty. On the other hand, variations due to differences of age and sex are greater than in any other of our genera.

Jordan & Evermann recognize 7 American species belonging to this genus. One of these (*P. maculifer*) is shown by our specimens to represent one of the younger stages of *P. lunatus*, and another (*P. ellipticus*) is a stage of the same species. Besides *P. lunatus*, the only other species known to occur in Porto Rico is *P. ocellatus*.

- a. Anal rays, at least anteriorly, each with a spinule at base (these formed by a slight widening of tip of interhæmal spines, each being covered by a little rough scale); front of dorsal with similar projections.
- b. Color brown, with pale rounded spots; fins dotted with brown; a faint dark spot at first one-third of lateral line; snout with horny points; mouth small, maxillary reaching front of eye..... *spinus*
- aa. Anal rays without spinules at their base.
- c. Anterior profile of head convex before interorbital area, the very short snout scarcely forming a reentrant angle at its base; form elliptic-ovate, outlines more regular than in *P. lunatus*.
- d. Dorsal rays 85 to 95.
- e. Scales not very small, about 75 pores in lateral line; no blue markings, at least in young.
- f. Mouth smaller, maxillary 3.33 in head. Color light-grayish, tinged with reddish, with small round spots of darker gray, and with lighter rings inclosing spaces of ground-color..... *ocellatus*, 274
- cc. Anterior profile of head strongly concave before interorbital area, the projecting snout leaving a marked reentrant angle above it.
- g. Mouth not very small; maxillary 3 in head. Color dark-olive, with many rings, curved spots, and small round dots of sky-blue edged with darker on body, these largest near middle of sides, where some are as large as eye; 3 obscure dark blotches on straight part of lateral line..... *lunatus*, 275

274. **Platophrys ocellatus** (Agassiz).

Head 4; depth 1.5; eye (lower) 3.4; snout 5; maxillary 3.33; D. 80; A. 60; scales 75 (pores); vertebrae 37. Body ovate, deep anteriorly, profile descending steeply, snout conspicuously projecting. Mouth very small and oblique, maxillary reaching vertical from the front of lower eye, 3.33 in head; tip of lower jaw entering profile. Teeth fine, conical, in two series in upper jaw, one in lower, those of the outer row in upper jaw larger and more widely separated than those of inner series. Snout very short, equaling interorbital width. Interorbital space narrow, deeply concave, closely scaled. Eyes large, lower in advance of upper; gillrakers obsolete, 7 rudiments on horizontal branch of anterior arch. Scales moderate, not extending on fins, those of colored side ctenoid, on blind side smooth; arch of lateral line short and high. Dorsal fin beginning opposite anterior nostril, rays nearly uniform in length, longest about one-half head; pectoral of colored side 4.75 in length; ventral of colored side beginning under middle of lower eye, with 6 rays; right ventral with 5 rays.

Color in life: Light-grayish, with reddish tinge, covered with small round spots of darker gray and with lighter rings inclosing spaces of ground-color; vertical fins similarly colored, with a small black spot near base of each ninth or tenth ray; 3 black spots on median line of body—one on straightest portion of lateral line just behind arch, another about midway between it, and a third much smaller one at caudal end of body or base of caudal peduncle; some other small black spots scattered over colored side. Eight specimens from Porto Rico present some variation of color; spots on fins not always as in above description, but the general color-pattern is about the same, varying mainly in the darkness or lightness of the shades.




FIG. 105.—*Platophrys ocellatus*.

The following shows the variation in some essential measurements of 4 of the largest specimens:

Total length.	Length to base of caudal.	Head in body.	Depth in body.	Eye in head.	Snout in head.	Maxillary in head.	Number of dorsal rays.	Number of anal rays.	Number of pores.
<i>mm.</i>	<i>mm.</i>								
73	63	3.7	1.65	3.4	4.85	3.4	80	67	75
71	62	4.13	1.8	3	3.75	3.33	80	60	73
50	42	3.8	1.7	3.14	3.18	3.38	80	59	75
46	40	4	1.8				80	60	75

Found in the western Atlantic from Long Island to Rio Janeiro, on sandy shores. Recorded from several Florida localities and recorded by Poey under other names from Cuba. The present collection contains specimens from San Antonio Bridge, San Juan, Culebra, and Boqueron; Mr. Gray's collection has one from San Geronimo.

Rhombus ocellatus Agassiz, in Spix, Pisc. Brasil., 85, pl. 46, 1829, Brazil.

Rhombus bahianus Castlenau, Anim. Nouv. et Rares Amér. du Sud, 1855, Bahia.

Platophrys nebularis Jordan & Gilbert, Proc. U. S. N. M. 1884, 31, 143, Key West.

Platophrys ocellatus, Jordan & Evermann, l. c., 2663, 1898.

275. *Platophrys lunatus* (Linnaeus).

Head 3.8; depth about 2; eye (lower) 6.25; interorbital space 3.12; snout (from lower eye) 3.6; maxillary 3.12; pectoral 1.22; D. 92; A. 74; scales (pores) 92. Body elliptical-ovate, strongly compressed; anterior profile concave, snout projecting, leaving a reentrant angle above it; mouth moderate, maxillary reaching to middle of pupil of lower eye; jaws subequal, lower with a well-developed knob

at symphysis; teeth small, in an irregular double series in each jaw; anterior end of maxillary with large blunt spine pointing somewhat outward and forward; interorbital very wide and deeply concave; orbital rim, below on upper orbit, above on lower, broken up into blunt papillae; eye with moderate cirri near upper margin of iris on eye membrane or cover. Gillrakers short and thick, 9 developed on lower part of arch, none on upper. Anterior part of interorbital, snout, maxillary, and mandible naked; scales of eyed side ctenoid, those of blind side smooth; arch of lateral line short and high, its base contained 5.5 in the straight portion. Dorsal fin beginning above the anterior nostril; ventral of colored side beginning under middle of lower eye, with 6 rays; left ventral inserted farther back, also of 6 rays.

Color in spirits: Ground-color of eyed side gray, thickly dotted with light-brown, giving it a brownish tint; speckled with dark-brown, numerous complete, broken, and incomplete rings of light-blue, edged outside with dark-brown or olive, surrounding areas of ground-color; diameter of longest ring about 10 mm., some of the rings with irregular or scalloped outline indicating coalescing spots; base of tail and head with blue spots and crescents; vertical fins and caudal with light-blue spots (becoming white in time); vertical fins brownish with black spots at intervals along the base; pectoral barred with black. A black blotch on lateral line at junction of straight portion and arch, another about midway the straight portion, and another at base of caudal peduncle.

The specimen upon which this description is based (taken from a specimen 213 mm. total length, from Culebra Island) differs somewhat from current descriptions chiefly in lacking the smaller spine on upper edge of maxillary behind larger one on snout; in the filamentous ray of pectoral; in having 6 rays in right ventral; presence of ocular cirri; and in having ctenoid scales on colored side. Notwithstanding these apparent differences, there can be no doubt regarding the identification. The pectoral is slightly filamentous, indicating the possibility of individuals, perhaps the other sex, having the long filament as found in other flat-fishes. The so-called "spine" on the snout is really a tubercle in the above-described specimen, and doubtless is more or less an age or sex character, if not both, and perhaps a more or less individual variation. As shown by other specimens, the scales are not always so strongly ctenoid, and they are not uniformly so in this one. The coloration is plainly that of *P. lunatus*. A series of fishes of this genus shows all grades of coloration and other variations which enable us to include in *P. lunatus* at least one other species formerly considered distinct, and very probably two. *Platophrys maculifer* represents a younger individual of *P. lunatus* in which the supposed differential marks are not so well developed. Jordan & Evermann, in Bulletin 47, U. S. N. M., state that they have never seen any young examples certainly referable to *P. lunatus*, and until its development is traced some of the species known from small examples only must be doubtful. This also leads us to believe that the present conclusions are correct.

The following notes will serve to show something of the color variations:

1. A specimen, 215 mm. long, from Aguadilla has the black blotches along side more diffuse and larger, the first being under end of pectoral below lateral line; cirri are present on both eyes.

2. A specimen from San Antonio Bridge, 131 mm. long, agrees perfectly with descriptions of *P. maculifer* (Poey) in coloration, and very closely in other respects. Head 3.33; depth about 1.8; eye (lower) 5.8; maxillary 3.3; snout 4.12; D. 90; A. 70; scales 86. Mouth small, oblique, the maxillary 3.4 in head; eye with cirri; no filamentous pectoral ray; arch of lateral line short and high. Color, reddish-gray, body covered with circles of round sky-blue spots, which are not confluent and not edged with darker color; besides these very few detached spots or other blue markings; head with similar blue spots, but no rings; area inclosed in blue rings not different from ground-color; caudal with blue spots, other fins with none; dorsal and anal mottled with black blotches each covering the bases of about 3 rays, with about 7 rays intervening between the blotches; a large, diffuse, dusky spot at front of straight part of lateral line; one better defined with lighter area on it anteriorly on middle of lateral line; a faint one under last part of vertical fins not far in front of caudal peduncle; pectoral grayish, with narrow dark bars.

3. A specimen 97 mm. long from Playa de Ponce may be described as follows: Head 3.5; depth about 1.7; eye about 4.33; snout 4; maxillary 3; D. 85; A. 73; pores 90; ocular cirri present. Color, gray, clouded with smoky-brown and some still more dusky mottlings; numerous small blue spots arranged in more or less complete rings and singly, some of the rings with a small whiter spot in center; large black blotch just behind arch of lateral line, one midway of straight portion, and another small faint one over last part of anal fin; first blotch has 4 gray areas in it, second has one large crescentic gray area, which is spotted or dotted with black; a fainter, lighter margin around front of point of the

most intense part of blotch, giving it an ocellated effect. Small ocelli, similar in character, scattered over body; fins all spotted and marked with black.

4. A specimen, 109 mm. total length, presents similar color pattern to the preceding, but of lighter shade. Head 3.8; depth 1.8; eye 4.16; snout 3.8; maxillary 2.7; D. 91; A. 70; scales (pores) 90; cirri present on eye membranes.

5. A specimen, 42 mm. long, possesses about the same general color arrangement as larger specimens, but lacks the blue spots. Head 3.33; depth 1.8; D. 94; A. 72; pores 87. Ocular cirri large.

6. Another young individual from Ponce, 45 mm. in total length, is very brightly and distinctly marked in the same general way as the others mentioned, but lacks the blue spots. Ocular cirri large. Head 3.33; depth 1.33; D. 94; A. 69; pores 82.

7. A young example from San Antonio Bridge, 50 mm. long, presents the same general appearance as the other young ones, but has longer ocular cirri. Head 3.5; depth 1.75; eye 4.8; snout 4; maxillary 3; pectoral 1.5; D. 96; A. 73; pores 83.

Life color: Pale-sandy, with numerous small ocellated brown spots with pale centers; a large black blotch on lateral line at posterior third of body; fins with numerous small white spots and a few larger irregular brown ones.

An inhabitant of the West Indies, north to Florida; common; recorded from Havana, Cienfuegos, Jamaica, Bahamas, Sombrero, St. Thomas, and from Green Turtle Key. Specimens at hand from Culebra, Aguadilla, San Antonio Bridge, San Juan, and Ponce, Porto Rico.

Solea lunata et punctata (the Sole), Catesby, Nat. Hist. Carolina, tab. 27, 1725, Bahamas.

Pleuronectes lunatus Linnæus, Syst. Nat., ed. X, 269, 1758, Bahamas; based on Catesby.

Pleuronectes argus Bloch, Ichth., tab. 48, 1783, Martinique; after Plumier.

?*Pleuronectes surinamensis* Bloch & Schneider, Syst. Ichth., 156, 1801, Surinam.

Pleuronectes maculiferus Pocy, Memorias, II, 316, 1860, Cienfuegos, Cuba.

Pleuronectes ellipticus Pocy, Memorias, II, 315, 1860, Cuba.

Platophrys lunatus, Jordan & Evermann, l. c., 2665, 1898.

Platophrys maculifer, Jordan & Evermann, l. c., 2664, 1898.

Platophrys ellipticus, Jordan & Evermann, l. c., 2665, 1898.

Genus 156. SYACIUM Ranzani.

Body elliptic-ovate, much compressed; interorbital space broad in males and more or less concave, narrowed in female; mouth moderate, gape curved; teeth in upper jaw biserial, in lower uniserial; front teeth of upper jaw enlarged; vomer toothless; scales rather large, ciliate; lateral line without arch in front; pectoral fins present on both sides; septum of gill-cavity below gill-arches without foramen; a deep emargination near isthmus; gillrakers short and thick; dorsal low, its anterior rays not elevated; pectorals both present; caudal sessile; no anal spine; pectorals produced in males; ventral fins short, that of colored side on ridge of abdomen.

This genus contains a considerable number of species, mostly American and African, which form a transition from *Platophrys* to *Citharichthys*. They fall readily into 2 groups distinguished by width of interorbital space. As this width is dependent on age, and as it is subject to various intergradations, the group *Aramaca* founded on it can not be admitted as a distinct genus.

a. Snout and orbits without spines or spinous processes.

b. Scales rather large, 50 to 57 in lateral line; interorbital space broad. Color, nearly plain brown, with darker dots or mottlings, no ring-like spots or ocelli; fins mottled; left pectoral barred; blind side sometimes wholly or partly dusky, especially in northern specimens..... *pectulum*

bb. Scales rather small, 58 to 70 in lateral line *micrurum*, 276

276. *Syacium micrurum* Ranzani.

Head 3.8 in length; depth 2.4; D. 87 to 92; A. 54 to 68; scales 65 to 70 (pores); eye 4 in head; maxillary 2.5 to 3. Form regularly elliptical, profile evenly convex to end of snout; eyes large, nearly even in front, male with interorbital space deeply concave, its width two-thirds vertical depth of eye (or more in Brazilian specimens); female with interorbital area much narrower, with a more or less perfect median groove, its width about equal to depth of pupil; mouth small, maxillary reaching to below middle of eye; teeth small, slender, in 2 rows above, in 1 row below, outer series in upper jaw somewhat enlarged, but hardly canine-like; gillrakers very short and thick, about 1 + 7 in number; scales small, firm, moderately ctenoid; pectoral 1.33 in head in female, reaching nearly to base of caudal in male; vertebrae 9 + 24 = 33.

Color, dark-brown, with many rings and spots of light-gray and blackish, some of the dark rings with a black central spot; a diffuse dusky blotch on lateral line above pectoral and one near base of caudal peduncle; fins with numerous inky spots and dark markings; blind side pale.

Fifteen specimens from Mayaguez appear to differ somewhat in general appearance from other specimens which can be referred to this species, but measurements and comparison reveal no really tangible constant differences. The largest, 163 millimeters long, presents the following characters: Head 3.9; depth 1.33; eye 4.5; snout 4.8; maxillary 2.4; interorbital 8; pectoral 1.3; D. 88; A. 68; scales 59 (68). Color, plain-brown, with a few faint or obscure grayish ocelli and darker spots; a faint brown blotch under end of pectoral, 1 on middle of straight line and 1 near base of caudal peduncle. All fins with light-brown crossbars.

For comparison measurements of an undoubted *S. micrurum* of similar size are given: Head 3.75; depth 2.4; eye 5.33; snout 4.8; maxillary 2.6; interorbital 16; pectoral 1.4; D. 88; A. 70; scales 59 (65).

West Indian fauna, Key West to Rio Janeiro, Cuba, Haiti, Jamaica, and Bahia, rather common; the Porto Rican examples are from Mayaguez, Ponce, Hucares, Aguadilla, Culebra, Boqueron, Palo Seco, and stations 6059 and 6061.

Syacium micrurum Ranzani, Nov. Spec. Pisc. Diss., Sec., 20, pl. 5, 1840, Brazil; Jordan & Evermann, l. c., 2672, 1898.

Hippoglossus ocellatus Poey, Memorias, II, 314, 1860, Cuba.

Hemirhombus aramaca Günther, Cat., IV, 42, 1862, Cuba and Jamaica.

Citharichthys athalion Jordan, Proc. U. S. N. M. 1886, 52, Havana.

Genus 157. CITHARICHTHYS Bleeker. Whiffs.

Eyes and color on left side. Body oblong; mouth of the large type, but comparatively small, with 1 series of small, sharp teeth in each jaw; no teeth on vomer or palatines. Gillrakers moderate, slender. Dorsal fin beginning just in front of eye; all fin rays simple; ventrals of colored side on ridge of abdomen; no anal spine; caudal fin convex or double truncate behind, none of the fins produced. Scales thin, deciduous, slightly ctenoid. Lateral line nearly straight, simple. Lower pharyngeals separate, each with a single row of teeth. Vertebrae 30 to 40.

This genus includes small flounders of weak organization, especially characteristic of sandy shores of tropical America. The sub-genus *Orthopsetta* includes species of more northern range and somewhat different form, and especially noteworthy as having an increased number of vertebrae. The two groups intergrade so perfectly that no sharp line of division can be drawn between them.

The genus *Citharichthys* contains 13 American species, 7 of which are Atlantic. Only 3 of these are known to occur in Porto Rico.

- a. Vertebrae 33 to 36; interorbital ridge low and narrow, head closely compressed.
- b. Eye large, 3 to 4.5 in head.
- c. Head large, 3 to 3.33 in length.
- d. Interorbital space very narrow, 5 in eye; snout with a spine; pectoral of eyed side elongate, one-third longer than head; maxillary 2.25 in head. D. 91; A. 73; scales 48. *dinoceros*
- ce. Head smaller, about 4 in length.
- e. Body comparatively elongate, depth about 2.5 in length; mouth very small; maxillary 3.5 in head; dorsal rays 83; anal 67; scales 40; eye 4 in head. *arctifrons*
- cc. Body comparatively broad, depth about half the length; mouth larger.
- f. Snout with a strong, sharp spine on eyed side, above upper lip; eye large, 2.3 to 3 in head; greatest depth of body over pectorals; interorbital space with a wide ridge, about half diameter of eye. *unicornis*, 277
- ff. Snout without distinct spine; eye moderate, 3.5 to 4.5 in head; greatest depth of body under middle of dorsal; interorbital space a narrow, sealy ridge with a slight median groove; maxillary 2.33 in head; teeth small, those in front slightly enlarged; body not very thin; gillrakers moderate, 6+13.
- g. Dorsal rays 68; anal 52; scales smaller, lateral line with about 53 pores; side with whitish blotches. *uhleri*
- gg. Dorsal rays 80; anal 56; scales large, 41 in lateral line; side and fins with dark blotches. *macrops*
- bb. Eye quite small, 5 to 8 in head; snout short, forming an angle with profile; mouth moderate, oblique, maxillary 2.5 to 2.66 in head; teeth small, anterior somewhat enlarged; dorsal about 80; anal 60; body and fins speckled.
- h. Scales not very large, 45 to 51 in lateral line; gillrakers long and slender, longer than pupil.
- i. Eye small, about 8 in head. *spilopterus*, 278
- ii. Eye larger, about 6 in head. *arenaceus*, 279

277. *Citharichthys unicornis* Goode.

Head 4.5; depth 2.33; eye 2.33; snout 4; maxillary 2.33; pectoral 1.55; D. 77; A. 60; scales 42. Body ovoidal, not particularly deep, but strongly compressed; head broad across front; mouth moderate; no spines, but a knob at symphysis of lower jaw; eye large; interorbital a narrow ridge dividing

anteriorly, becoming front rim of orbit, concave in front of eyes; pectoral short; longest rays of vertical fins moderate; scales large, deciduous.

Color, somewhat mottled light-brown.

This description is from a single specimen taken by the *Fish Hawk* at station 6063, in Mayaguez Harbor. Total length 57 mm., the only specimen of this species taken. After comparison with a large number of specimens in the U. S. National Museum, we have decided that the present specimen is a female, the lack of spines on head being coincident with a narrow interorbital. Our example corresponds exactly with many females of similar size in the National Museum collection. The wide interorbital in other species has been found associated constantly with the male when any difference of this kind existed. For comparison, and because the horned fish may be met with, a description of an individual, presumably a male, taken in deep water in the Gulf Stream, is here given: Total length 77 mm.; head 3.5; depth 2.25; eye 3; interorbital 6; snout 4.5; maxillary 2.25; pectoral 2.15; D. 74; A. 60; scales 45. Head broad, wider across front of eyes than in female; muzzle less sharp; interorbital wide anteriorly, narrowing behind, concave, a ridge extending across from back part of upper orbit to side of anterior orbit in front (in the female this makes the ridge between the eyes); a sharp slender spine projecting forward from edge of upper part of the snout; 2 smaller ones from point of upper orbit; small one from side of snout near tip extending somewhat to one side; 2 from front of orbital rim or lower eye; knob at symphysis of lower jaw; head, as in the female, everywhere scaled. Fins evidently once had brown blotches.

Formerly collected in deep water of Gulf Stream off southeast coast of New England; off Sand Key, Florida, in 44 fathoms; and in the Gulf of Mexico southward of Cape San Blas, in 60, 111, and 142 fathoms.

Citharichthys unicornis Goode, Proc. U. S. N. M. 1880, 342, Gulf Stream southeast of New England; Jordan & Evermann. l. e., 2683, 1898.

278. *Citharichthys spilopterus* Günther.

Head about 3.75; depth about 2.25; eye 7.8; snout 5.8; maxillary 2.5; pectoral 2; D. 82; A. 61; scales (pores) 48. Body moderately elongate, much compressed; snout short, forming an angle with profile; jaws strongly curved, upper somewhat hooked over lower; lower jaw slightly included; maxillary reaching to posterior margin of lower orbit; teeth small, in a single row, anterior a little enlarged; interorbital area a low, narrow ridge, which is divided anteriorly; gillrakers short and rather slender; scales finely ctenoid on colored side; origin of dorsal above anterior edge of upper eye, slightly on blind side; origin of anal slightly behind base of pectoral.

Color, light-brown mottled somewhat with darker; indistinct blotch on lateral line under tip of pectoral, one about middle of straight portion, and one at base of caudal; vertical fins with narrow elongate spots on rays vertically arranged, sometimes in pairs at more or less regular intervals.

Description taken from specimen, 148 mm. total length, from Mayaguez. Everywhere abundant on sandy shores of the warmer parts of western Atlantic, in shallow water; specimens recorded from South Carolina, Florida, Cuba, and Brazil. The Porto Rican collection contains numerous individuals from Mayaguez, Aguadilla, San Antonio Bridge, San Juan, Playa de Ponce, Palo Seco, Vieques, Caballo Blanco Reef, Puerto Real, Boqueron, and Fajardo.

Citharichthys spilopterus Günther, Cat., IV, 421, 1862, New Orleans, Santo Domingo, and Jamaica; Jordan & Evermann. l. e., 2685, 1898.

Citharichthys cayannensis Bleeker, Comptes Rendus Acad. Sci. Amsterdam, XIII, 1862, 6, Cayenne; name only.

Citharichthys guatemalensis Bleeker, Neder. Tydschr. Dierk., 1864, 73, Guatemala.

279. *Citharichthys arenaceus* Evermann & Marsh, new species.

Head 3.8; depth nearly 2; eye 6; snout 5; maxillary 2.2; pectoral 1.8; D. 74; A. 54; pores 51. Body elliptical, rather deep; head deep; mouth large; snout not very prominent, making but a slight notch in front of upper eye; interorbital narrow, concave; eyes small, not close together; lateral line not arched, but from upper end of gill-opening directed slightly downward to about in a line with tip of pectoral, whence it extends directly backward to caudal fin; scales of colored side (left) finely ctenoid, with accessory scales along region of lateral line; rays of vertical fins and caudal with small scales, as is usual in flat-fishes; ventrals with 6 rays each.

Color light-gray, thickly spotted with olive, giving it a granitic appearance, the rays of all the fins with narrow alternating olive and gray crossbars; some rays of vertical fins with short oblong spots extending in the direction of the ray; blind side pale.

A small-eyed *Citharichthys*, in this respect related to *C. spilopterus*, but, except in small examples, bearing scarcely any other resemblance to that species. It differs from *C. spilopterus* mainly in the general heavier appearance, being everywhere broader and thicker, and in its color. There are two unnamed specimens in the National Museum from Santo Domingo referable to this species, otherwise known only from Porto Rico. This description is from a specimen, 162 mm. in total length (type No. 49526, U. S. N. M.), 1 of 10 specimens from Mayaguez.

749536




FIG. 106.—*Citharichthys arenaceus*.

The following table of relative measurements of these specimens is given for comparison:

Comparative measurements of 10 cotypes of Citharichthys arenaceus.

No.	Total length.	Length to base of caudal.	Head.	Depth.	Eye.	Snout.	Maxillary.	Pectoral.	Dorsal.	Anal.	Pores.
1	162*	141	3.8	2	6.17	5.3	2.33	1.5	74	51	50
2	130	110	3.75	2	5.8	4.8	2.41	2.07	68	50	51
3	114	100	3.84	2.8	5.2	4.33	2.6	1.86	69	51	45
4	94	81	4.05	2.14	5	5	2.5	2	70	50	48
5	95	82	3.86	2.1	5.25	4.66	2.33	2.1	70	53	48
6	95	81	3.86	2.011	5.25	5.25	2.62	2.1	75	48	50
7	88	76	4	2.17	4.75	4.75	2.7	1.9	74	54	50
8	84	73	3.94	2.14	5.3	5.3	2.5	2.05	75	53	49
9	77	65	3.8	2.17	4.86	4.86	2.43	2.12	72	51	48
10	59	49	3.8	2.33	4.33	4.33	2.6	1.86	72	51	48

*The type.

One other specimen was collected January 19 at Mayaguez, 2 at Aguadilla January 18, and 1 at San Juan January 14. The two from Aguadilla may be described as follows:

Measurements.	No. 1.	No. 2.
Total length.....	125	123
Length to base of caudal.....	115	116
Head (about).....	4	4.14
Depth (about).....	2.16	2.33
Eye (about).....	5.8	6.22
Snout (about).....	5.25	5.9
Maxillary.....	2.63	2.54
Pectoral.....	2.25	2.17
Dorsal.....	73	72
Anal.....	52	51
Pores.....	46	50

Color of same general pattern, but with large brownish blotches over body and one on caudal peduncle.




FIG. 107.—*Etropus crossotus*.

Genus 158. **ETROPUS** Jordan & Gilbert.

Eyes and color on left side. Body regularly oval, deep, and compressed. Head small; mouth very small, teeth close-set, slender, and pointed, somewhat incurved, mostly on blind side; no teeth on vomer. Eyes small, separated by a narrow, scaleless ridge; margin of preopercle free. Ventrals free from anal, that of colored side inserted on ridge of abdomen, its base rather long. Dorsal fin beginning above eye; caudal double-truncate; anal without spine. Scales thin, deciduous, ctenoid on left side, cycloid on blind side. Lateral line simple, nearly straight.

This genus is very close to *Citharichthys*, from which it differs only in the very small size of mouth and in the correspondingly weak dentition. The 3 or 4 known species are similar in appearance to the species of *Citharichthys* and inhabit the same waters. Size small. *Thysanopssetta*, a South American genus, is also very close to *Etropus* and *Citharichthys*, but has teeth arranged in a band. The larval forms are translucent and symmetrical, as in *Platophrys*, *Monolene*, *Arnoglossus*, etc.

- a. Snout not acute; dorsal rays 75 to 85.
- b. Body comparatively elongate, depth rather less than one-half length.
- c. Dorsal rays 81; anal 58; head 4.25 in length; eye 3.5 in head; maxillary 4..... *microstomus*
- cc. Dorsal rays 77; anal 61; head 4 in length; eye 3.5 in head; maxillary 4.33..... *rimosus*
- bb. Body very deep, depth more than one-half length; eye 3.75 in head; maxillary 4; head 4.8; depth 1.8 to 2; D. 76 to 85; A. 56 to 67; scales 42 to 48; cirri on subopercle of blind side very numerous, white; olive ground, with darker blotches; fins sanded..... *crossotus*, 280

280. *Etropus crossotus* Jordan & Gilbert.

Head 4.8 in length; depth 1.75 to 2; D. 76 to 85; A. 56 to 67; V. 6; scales 42 to 48; vertebrae $9 + 25 = 34$. Body oval and strongly compressed, with the dorsal and ventral curves nearly equal; both outlines strongly arched anteriorly, body much deeper in the adult; head very small; snout short; mouth very small, its cleft not so long as diameter of orbit; teeth conical, pointed, close-set, strongly incurved, in a single series, those in upper jaw on the blind side only, those on lower jaw on both sides; eyes large, lower in advance of upper, separated by a very narrow scaleless ridge, which extends backward above the preopercle; edge of opercle on blind side with a row of conspicuous white cilia; upper nostril turned somewhat to blind side; anterior nostril on left side, with a very slender cirrus; dorsal fin commencing over front of upper eye, its middle rays highest, anterior not elevated; anal fin not preceded by a spine, its middle rays highest; caudal fin very sharply double-truncate, as long as head; pectorals short, that of left side the longer, about three-fourths length of head; ventral of colored side on ridge of abdomen, the membrane of its last rays nearly reaching base of first ray of anal; ventral of blind side longer than the other, one-half length of head, inserted farther forward than that of colored side; vent lateral, with well-developed anal papilla. Scales thin, large, etenoid on colored side, smooth on blind side, those on middle part of body larger; head entirely scaly, except the snout and interorbital ridge; rays of vertical fins with scales on basal half, on colored side; lateral line developed equally on both sides, nearly straight.

Color, olive-brown, with some darker blotches, most distinct in larger specimens; vertical fins finely mottled and streaked with black and gray; pectoral and ventral on left side spotted.

Tropical America on both coasts, north to Cerros Island and North Carolina, south to Panama and Rio Janeiro; recorded from Charleston, Cedar Keys, New Orleans, Galveston, Beaufort, N. C., Mazatlan, Panama, from localities along both sides of coast of Lower California, and from several places in Florida. Specimens before us are from Mayaguez, Palo Seco, and Arroyo, Porto Rico.

Etropus crossotus Jordan & Gilbert, Proc. U. S. N. M. 1881, 364, Mazatlan; Jordan & Evermann, l. c., 2689, 1898.

Family LXXIV. SOLEIDÆ. The Soles.

Body oblong or elongate, usually scaly; mouth very small, much twisted toward eyed side; teeth in villiform bands, very small or obsolete; eyes small, close together, with or without a bony ridge between them; edge of preopercle adnate, concealed by skin and scales; gill-openings narrow, gill-membranes adnate to shoulder-girdle above; pectoral fins small or wanting; ventral fins small, one or both sometimes wanting.

Small fishes living on sandy bottoms, similar to the *Pleuronectidæ* in structure, but much degraded, the fins and teeth having lost many distinctive qualities; the vertebrae usually in increased numbers.

The soles are numerous in the warm seas, and those of sufficient size are valued as food. They comprise about 12 genera and 150 species. They are naturally divisible into 3 subfamilies, each quite distinct from the others, and possibly independently descended or degraded from normal *Pleuronectidæ*. The North American species belong to 2 subfamilies, very different one from the other. The *Achirinae*, or American soles, are apparently allied to the *Psettinae*, the ventral fin of eyed side extending along ridge of abdomen. The *Soleinae*, or European soles, show in the insertion of ventral and in other respects a strong resemblance to the *Pleuronectinae*. The more aberrant *Cynoglossinae*, or tongue-fishes, are perhaps degraded *Soleinae*, but the eyes are sinistral, as in the *Psettinae*. In the *Soleinae* and *Achirinae* the eyes are dextral, as in the *Pleuronectinae*.

ACHIRINE:

- I. Soles with eyes on right side and separated by a distinct bony ridge; ventral with long base confluent with anal. Body oblong or ovate, with color on right side; eyes moderate or small, upper eye usually more or less in advance of lower; mouth small, more or less twisted toward blind side; teeth little developed, in villiform bands; edge of opercle adnate, usually concealed by scales; gill-openings more or less narrowed, gill-membranes adnate to shoulder-girdle above; blind side of head usually with fringes; pectoral fins small, sometimes wanting; ventral fins developed, one or both of them sometimes obsolete; scales usually etenoid, rarely wanting; lateral line straight, usually single; right ventral with extended base, confluent with anal fin.
 - a. Gill-openings of moderate extent, confluent below; vertical fins well separated; body ovate in outline, depth nearly one-half length; pectoral fins rudimentary or wanting; lateral line straight; scales well developed, etenoid more or less enlarged on head, those of blind side of head with fringes; vertebrae about 28..... ACHIRUS, 159
 - aa. Gill-openings very small, separate, each reduced to a small slit below angle of opercle; right ventral beginning at chin; pectoral fins minute or wanting; lateral line straight; snout dilated, dorsal beginning upon it.

- b. Scales present, etenoid; caudal somewhat confluent with dorsal.
 c. Left ventral rudimentary, with 2 rays..... APIONICHTHYS
 bb. Scales none; caudal free from dorsal and anal GYMNACHIRUS
 CYNAGLOSSINÆ:
 II. Soles with eyes on left side, not separated by a bony ridge. Body elongate, more or less lanceolate in outline, with color on left side.
 d. Ventral fin of eyed side only present, free from anal; no pectoral fins; no lateral line; head without fringes.

SYMPHURUS, 160

Genus 159. **ACHIRUS** Lacépède. American Soles.

Eyes and color on the right side. Body oblong, bluntly rounded anteriorly. Head small; eyes small, close together, upper eye in advance of lower, the two separated by a bony ridge; mouth small, somewhat turned toward colored side; nasal flaps present, nostril of blind side fringed; lip of colored side fringed; teeth very small, on blind side only; gill-openings rather narrow, but confluent below, not reduced to a slit; branchiostegal region scaled. Head closely scaled everywhere, scales on colored side similar to those on body, those of nape and chin much enlarged; scales on blind side anteriorly with their pectinations more or less produced, forming eirri; scales of both sides extremely rough, extending on fins. Lateral line straight, simple; edge of preopercle covered by scales. Dorsal beginning on snout, low in front and thickly scaled, its rays divided; anal fin similar, without spine; caudal fin free, convex; caudal peduncle very short and deep; pectoral fin of left side wanting, that of right side small or obsolete; ventral rays 3 or 4, ventral fin of colored side long, connected with anal by a membrane.

This strongly marked genus contains numerous species, all very closely related, and nearly all American. It has been united by Dr. Günther with *Solea*, but for no good reason, as the number of vertebrae is very much fewer than in the European soles, and the right ventral fin is decurrent along the abdomen and united with the anal in the American soles, while it is short and wholly free in all the European forms. The two groups belong in fact to distinct subfamilies. It is also worth noticing that the name *Achirus* is prior in date to that of *Solea*. The species with rudimentary pectoral fins have been set apart by Dr. Bean to form the genus *Baiostoma*, but the very slight development of these organs in some of the species and the evidently very close relationship of them all lead us to regard *Baiostoma* as a subgenus only. If we follow Kaup in restricting the name *Achirus* to the Asiatic group called *Pardachirus*, the present genus would receive the name of *Trinectes*. It seems to us, however, that both Lacépède and Cuvier regarded the species called by us *A. fasciatus* as the type of their genus *Achirus*.

BAIOSTOMA:

- a. Pectoral fins small, present at least on right side.
 b. Pectoral fins present on both sides, that of left side rudimentary, of a single ray; of eyed side with about 3 rays.
 c. Dorsal rays 60 to 67; anal rays about 48. Color brownish, irregularly spotted with darker, and with about 10 black vertical lines crossing lateral line..... *achirus*
 cc. Dorsal rays 53 to 57. Color olivaceous..... *inscriptus*, 281
 bb. Pectoral of right side only present.
 d. Dorsal rays 50 to 58; anal rays 38 to 48.
 c. Pectoral fin of 4 to 6 rays, considerably longer than eye; body with 8 to 10 narrow, vertical dark bars, these sometimes obsolete with age *lineatus*, 282

ACHIRUS:

- aa. Pectoral fins wholly wanting; dorsal rays 50 to 55; anal rays 37 to 46; right lower lip fringed; left nostril with some fringes; depth 1.8 in length; head 4; none of scales of eyed side with hair-like appendages. Color dusky-olive, more or less mottled with about 8 dark vertical stripes, these varying very much in width and number; caudal spotted..... *fasciatus*

281. **Achirus inscriptus** Gosse.

Head 3.75 in body; depth 1.75; D. 53 to 57; A. 40; scales 75 to 80; interorbital width less than eye; upper eye in advance of lower. Pectoral present on each side, that of left side rudimentary, usually of a single ray; that of eyed side with about 3; left ventral with 1 or 2 small rays, in some specimens entirely absent; right ventral joined to anal. Scales smaller and less rough than usual in this genus, those of nape scarcely enlarged on eyed side, those of blind side much fringed; scales of colored side with scattered hair-like appendages, some black, others pale. Vertebrae $8 + 20 = 28$.

Color, olivaceous; head, body, dorsal, and anal fins covered with a network of dark lines; traces of about 8 dark cross-streaks sometimes present; caudal base dusky. One specimen, 46 mm. long, from Palo Seco, has the color rather darker than usual, dark network so distinct that lighter ground-color shows as distinct light-gray spots; large round black spots scattered over side and vertical fins; no cross-streaks; caudal very pale with but a trace of dusky markings. Another individual, from San Antonio

Bridge, is very darkly colored, due to intensity of black network and darker gray ground-color; blind side somewhat dusted over with brown specks, which grade into solid blue-black on caudal peduncle; caudal fin abruptly pale with few faint dusky dots; vertical fins very dark on blind side, rays paler.

A specimen, 77 mm. long, from Aguadilla possesses 3 rays in each pectoral fin and is the only one of the collection having cross-streaks, of which there are about 8.

An inhabitant of the West Indies, north to southern Florida. It has been recorded from Haiti, Key West, and Cape Florida. This collection contains specimens from station 6063 (in Mayaguez Harbor in 75 fathoms), Fajardo, Palo Seco, Aguadilla, San Antonio Bridge, and San Juan.

Achirus inscriptus Gosse, Nat. Sojourn Jamaica, 52, pl. 1, fig. 4, 1851, Jamaica; Jordan & Evermann, l. c., 2696, 1898.
Monochir reticulatus Poey, Memorias, II, 317, 1861, Cuba.




FIG. 108.—*Achirus lineatus*.

282. *Achirus lineatus* (Linnæus).

Head 3.5; depth about 1.5; D. 49 to 58; A. 38 to 44; scales 75 to 85. Pectoral fin of right side only developed, of 4 to 6 rays, considerably longer than eye. Body with 8 to 10 narrow, vertical dark bars, these sometimes obsolete with age; vertical fins all with round dark spots, these usually especially distinct on caudal fin; some of scales of eyed side with black, hair-like appendages; pectoral fin with 5 or 6 rays, about 3 in head, its length equal to distance from outer edge of one eye to outer edge of other.

Color, brown, the young spotted with whitish, the adults sometimes with darker; body with about 8 narrow, vertical cross-streaks of blackish.

West Indies and Brazil, Florida Keys to Uruguay; common and variable. Recorded from Jamaica and Cienfuegos in the West Indies and from both coasts of Florida. Specimens at hand are from San Juan market, Palo Seco, Boqueron, and Vieques, Porto Rico.

Passer lineis transversis notatus Sloane, Jamaica, 2, 77, pl. 246, fig. 2, 1725, Jamaica.

Pleuronectes fuscus subrotundus glaber Browne, Jamaica, 445, 1756, Jamaica.

Pleuronectes lineatus Linnæus, Syst. Nat., ed. X, 268, 1758, Jamaica; based on Browne and Sloane; not of ed. XII, which is *Achirus fasciatus*.

Monochir maculipinnis Agassiz, Spix, Pisc. Brasil., 88, pl. 49, 1829, Brazil; Poey, Fauna Puerto-Riqueña, 341, 1881; Stahl, l. c., 166, 1883.

Balostoma brachiatris Bean, Proc. U. S. N. M., 1882, 413, Apalachicola Bay and south Florida.

Achirus comifer Jordan & Gilbert, Proc. U. S. N. M., 1884, 31, Key West.

Achirus lineatus, Jordan & Evermann, l. c., 2698, 1898.

Genus 160. SYMPHURUS Rafinesque. Tongue-fishes.

Body elongate, more or less lanceolate in outline, with eyes and color on left side; eyes small, very close together, with no distinct interorbital ridge between them; mouth small, twisted toward blind side; teeth little developed, in villiform bands; edge of preopercle covered by scales; gill-openings narrow, gill-membranes adnate to shoulder-girdle above, joined together and free from isthmus below; pectoral fins wanting (in adult); vertical fins more or less confluent; scales etenoid; lateral line wanting. Ventral fin of eyed side only present, free from anal; head without fringes.

SYMPHURUS:

- a. Scales not minute, etenoid, 65 to 105 in number; dorsal rays 86 to 100; anal rays 70 to 87.
 b. Scales rather large, about 65; head 4.33; depth 4.33; color, clouded-brown *piger*
 bb. Scales small, moderately etenoid, 75 to 105 in a longitudinal series.
 c. Dorsal and anal pale anteriorly, becoming more or less abruptly black posteriorly.
 d. Caudal fin abruptly pale, at least at tip.
 e. Body elongate, depth 4.5 in length; head 5.5. D. 96 to 100; A. 86 or 87; scales 88 to 90. Color, grayish, speckled with brown; dorsal and anal fins black on last tenth; caudal abruptly pale; tips of fin rays vermilion.... *marginatus*
 dd. Caudal fin black, as is a large part of dorsal and anal, black either continuous or in form of large spots. Color, brownish, often mottled, usually with more or less distinct darker crossbands and with longitudinal streaks along rows of scales, sometimes nearly plain brown.
 f. Scales rather larger, 75 to 90. Body less elongate, depth 3.1 to 3.66 in length; head 5.25 to 5.75. Dorsal 88 to 95; anal 75 to 80 *plagusia*, 283
 cc. Dorsal and anal pale throughout, or more or less mottled or spotted with darker, caudal similarly colored, not distinctly black. Body not very elongate, depth 3 to 3.33 in length. (Probably all varieties of *S. plagusia*.)
 g. Body with dark crossbands more or less distinct; fins mottled or speckled; upper eye slightly in advance of lower.
 h. Dorsal rays 86 to 95; anal rays 75 to 80; head 5 in length; depth 3.25; scales 85 to 93; crossbands more distinct than in related species..... *plagusia*
 hh. Dorsal rays 78 to 85; anal rays 70 to 72; head 5 in length; depth 3.5; scales 80 to 90. Color, light-brown, with darker crossbars, which become obsolete with age..... *pusillus*
 gg. Body uniform grayish, without crossbands; last part of dorsal and anal with 3 or 4 oblong black blotches; each somewhat larger than eye; upper eye directly above lower; head 5.66 in length. Scales 85; D. 92; A. 75. . . *diomedeanus*

ACEDIA:

- aa. Scales very small, etenoid, each with a medium dark streak, which simulates a keel, but is not a ridge; snout and jaws naked; fin rays in increased number.
 i. Head 5.66; depth 4.66; D. 119; A. 107; scales 120. Grayish, everywhere mottled with brown *nebulosus*

283. *Symphurus plagusia* Bloch & Schneider. Tongue-fish.

Head 5.25; depth about 3.5; D. 88; A. 76; scales 90. Body rather elongate. Color, brown, somewhat clouded; narrow longitudinal lines along rows of scales, posterior part of dorsal and anal and caudal black; dark bars on dorsal and anal in direction of rays. The black caudal has been considered the mark distinguishing this species from the closely-related form *S. plagiusa*, from which otherwise, according to descriptions, it differs but little.

Specimens at hand from Porto Rico show nearly all grades of coloration of the two forms which with measurements cover pretty well the two species. But all of the Porto Rican specimens are undoubtedly of the same species. They have been compared with specimens of *S. plagiusa*. The few specimens of the latter examined revealed none with black caudal, but they differed in no other respect, unless possibly in the slightly larger head and eyes. There can be little doubt but that the two forms are as nearly identical as are the two names.

Some individuals are plain brown, with longitudinal streaks of darker along rows of scales, or with distinct broad crossbars; or with irregular cross-mottling not arranged in definite bands. All of these colors may vary from faint and light to very dark and distinct. The fins are sometimes nearly plain, but usually with more or less numerous narrow dark bars running in direction of fin rays. Color of caudal rays ranging from pale, through longitudinally barred and dusky to very black.

West Indies to Brazil; common. Specimens have been taken at Havana and in Jamaica; in Porto Rico it is known from Palo Seco, Ponce, Mayaguez, Boqueron, and Hucars. The only record from Florida is "Off Key West, in about 20 fathoms" (Garman).

Measurements of 15 specimens of *Symphurus plagusia*, showing variations.

Locality.	Total length.	Length to base of caudal.	Head.	Depth.	D.	A.	Scales.	Color of tail.
Mayaguez	146	137	5.25	3.5	88	76	90	Black.
Do.	92	85	5.33	3.25	90	74	79	Pale.
Palo Seco.	120	113	5.33	3.05	88	77	82	Very black.
Do.	103	96	5.05	3.43	95	80	86	Black.
Do.	94	87	5.12	3.5	96	75	87	Do.
Do.	108	100	5.6	3.6	96	83	88	Do.
Do.	109	103	5.44	3.5	96	83	88	Do.
Boqueron	117	109	5.5	3.5	96	79	83	Very dusky.
Ponce	109	103	5.15	3.21	96	76	82	Dusky.
Do.	93.5	86	5.25	3.5	93	75	82	Slightly dusky.
Do.	88	82	5.12	3.25	91	75	78	Barred.
Do.	85	78	5.2	3.37	93	77	80	Very slightly dusky.
Do.	68	64	5	3.66	93	78	78	Plain pale.
Huacares	90	85	5.33	3.5	92	75	87	Pale.
Do.	91	84	5.25	3.5	91	79	82	Do.

Plagusia, Browne, Jamaica, 445, No. 1, 1756, Jamaica.

Pleuronectes plagusia Bloch & Schneider, Syst. Ichth., 162, 1801, Jamaica; after Browne.

Achirus ornatus Lacépède, Hist. Nat. Poiss., IV, 659, 1803, on a specimen "presented by Holland to France."

Plagusia tessellata Quoy & Gaimard, Voyage Uranie, Zoologie, 240, 1824, Rio Janeiro.

Plagusia brasiliensis Agassiz, Spix, Pisc. Brasil., 89, tab. 50, 1829, Brazil.

Aphoristia ornata, Poey, Fauna Puerto-Riqueña, 341, 1881; Stahl, l. c., 80 and 166, 1883.

Symphurus plagusia, Jordan & Evermann, 2709, 1898.

Family LXXV. ANTENNARIIDÆ. The Frog-fishes.

Head and body more or less compressed. Mouth vertical or very oblique, opening upward; lower jaw projecting; jaws with cardiform teeth; premaxillaries protractile. Gill-openings small, pore-like, in or behind lower axils of pectorals. No pseudobranchiæ. Gills 2.5 or 3; skin naked, smooth, or prickly. Pectoral members forming an elbow-like angle. Pseudobranchia long, with 3 actinosts. Ventral fins present, jugular, near together. Spinous dorsal of 1 to 3 separated, tentacle-like spines; soft dorsal long, larger than anal. Pyloric caeca none.

The *Antennariidae* comprise about 5 genera and 50 species, inhabitants of tropical seas, "living on floating seaweed, and enabled, by filling the capacious stomach with air, to sustain themselves on the surface of the water"; therefore widely dispersed by currents in the sea. The different species display the most grotesque forms and appearance; and, as Dr. Günther well says, there is none, perhaps, in which the singular organization of the fish is more distinctly seen to be in consonance with its habits; and there is none in which mimicry and protective coloration and organization are more strongly or more beautifully exemplified. "The habits of all are equally sluggish and inactive. They are found in all warm seas and are very bad swimmers. Those found near the coasts lie on the bottom of the sea, holding on with their arm-like pectoral fins to seaweeds or stones," among which they conceal themselves. Their colors and dermal filaments resemble the seaweeds very closely and render the fish extremely difficult of detection. The species of pelagic habits attach themselves to floating *Sargassum* or other floating seaweed and are carried about by the winds and currents.

- a. Head compressed; a rostral spine or tentacle, followed by 2 larger spines; palatine teeth developed; dorsal spines disconnected.
- b. Skin naked and smooth; ventral fins elongate PTEROPHRYNE, 161
- bb. Skin covered with prickles; ventral fins short..... ANTENNARIUS, 162
- aa. Head euboid; a single rostral spine or tentacle, received in a groove; soft dorsal low..... CHAUNAX, 163

Genus 161. PTEROPHRYNE Gill. Mouse-fish.

Body smooth or scarcely granular, short, somewhat compressed, with tumid abdomen; mouth small, oblique; palate with teeth; wrist and pectoral fin slender; ventrals elongated; soft dorsal and anal vertically expanded.

Small fishes of fantastic shape in the West Indies and the Gulf Stream.

- a. "Bait" on first dorsal spine bifurcate at tip..... *histro*
- aa. "Bait" on first dorsal spine bulbous, covered with fleshy filaments..... *gibba*, 284

284. *Pterophryne gibba* (Mitchill). *Sapo; Pescador.*

Garman refers to this species certain specimens obtained in Gulf weed about Key West and the Tortugas. These resemble *P. histrio*, but "differ markedly in certain respects. The bait on first dorsal spine, for instance, is bulbous and covered with slender fleshy filaments in our individuals, but in *P. histrio* it is bifurcate. *P. gibba* is fairly represented by Cuvier, 1817, in his *Chironectes lavigatus*. The formula for the individuals in hand is D. III, 12; A. 7; V. 5; P. 10; C. 9." (Garman.) West Indies north to Key West and the Tortugas; not seen by us in Porto Rico but recorded from there by Professor Poey and Dr. Stahl; probably not uncommon but often confounded with *P. histrio*.

Lophius gibbus Mitchill, Trans. Lit. and Phil. Soc. N. Y. 1815, I, pl. 4, fig. 9.

Chironectes lavigatus Cuvier, Mém. du Mus., III, 423, pl. 16, fig. 1, 1817, South Carolina.

?*Chironectes sonntagii* Baron J. W. von Müller, Reisen in den Vereinigten Staaten, Canada, und Mexico, Band I, 180, 1861, in floating seaweed; no exact locality stated.

Pterophryne lavigata, Poey, Fauna Puerto-Riqueña, 340, 1881; Stahl, l. c., 79 and 165, 1883.

Pterophryne gibba, Jordan & Evermann, l. c., 2717, 1898.

Genus 162. **ANTENNARIUS** Lacépède.

Body oblong, compressed, very deep through occipital region, tapering behind; breast tumid; mouth rather large, more or less oblique, or even vertical; cardiform teeth on jaws, vomer, and palatines; eye small; skin with small granules or spinules, these usually forked, and numerous fleshy slips. First dorsal spine developed as a small rostral tentacle; second and third dorsal spines strong, covered with skin, with numerous fleshy filaments; soft dorsal high and long; anal short and deep; caudal fin rounded, peduncle free; pectoral fins wide, with a rather wide wrist, at lower posterior angle of which are the very small gill-openings; ventral fins short.

Fantastic-looking fishes, very numerous in warm seas, especially in parts of the sea with floating vegetation; not rarely individuals are found far from their native latitudes, carried by currents to the coasts of Norway and New Zealand. Their power of swimming is very imperfect. When near the coast they conceal themselves among corals, stones, or fucus, holding on to the ground by means of their arm-like pectoral fins. The extraordinary range of some of the species which inhabit the Atlantic as well as the Indo-Pacific Ocean, is the consequence of their habit of attaching themselves to floating objects. Their coloration is so similar to their surroundings that it is hardly possible to distinguish the fish from a stone or coral covered with vegetation. Their way of attracting and seizing their prey is evidently the same as in the other fishes of this family. Almost all the species are highly colored, but the pattern of colors varies exceedingly. These fishes do not attain any considerable size, and probably never exceed a length of 10 inches. A great number of species have been distinguished by ichthyologists, but probably not more than twenty are known at present. (Günther.)

A dozen American species of this interesting genus have been recognized, 9 of which occur in the West Indies. The other 3 are found in the Gulf of California and southward on our Pacific coast. Of the 9 West Indian species only 3 have as yet been taken in Porto Rico.

- a. Bulbous tip or "bait" of first dorsal spine simple, undivided at tip.
- b. Skin smoothish except about eyes; first dorsal spine short, second rough. Body brown, with whitish spots; no ocelli *inops*, 285
- bb. Skin with prickles, velvety or shagreen-like.
- c. Prickles simple, none of them bifid.
- d. Color black; tips of pectorals and ventrals and one or two spots on side white (prickles undescribed).... *principis*
- dd. Color dusky; dorsal with 3 ocelli; caudal with many spots; first dorsal longer than second; no dermal flaps. *tenebrosus*
- cc. Prickles or spinules on body mostly bifid.
- e. Body with 3 large ocelli, 1 on dorsal, 1 on caudal, and 1 on middle of side, besides many black spots and streaks; tip of first dorsal spine fringed; mouth largely black within..... *ocellatus*
- aa. Bulbous tip or "bait" on first dorsal spine bifid at tip; skin shagreen-like.
- f. Color reddish, with brown spots, those about eye radiating.
- g. Dermal flaps numerous on body; spinules on skin short and stiff, rendering surface shagreen-like..... *scaber*, 286
- gg. Dermal flaps few; spinules on skin longer and slender, rendering surface velvety..... *tigris*
- ff. Color uniform black; surface of body rough, shagreen-like; inside of mouth white; first dorsal spine short, little longer than second..... *nuttingii*, 287
- aaa. Bulbous tip or "bait" of first dorsal spine trifid.
- h. First dorsal ray twice as long as second and as long as caudal; sides with numerous black ocelli, besides other streaks and dark spots; skin smoothish *multiocellatus*
- hh. First dorsal spine barely one-half longer than second; shorter than caudal; sides with dark streaks and reticulations; a large ocellus under middle of soft dorsal; body rough, with shagreen *radiosus*

285. *Antennarius inops* Poey.

Depth 2.75 with caudal. Skin lustrous, smooth, except for some points behind and below eye; third of first 3 dorsal rays largest, its membrane not reaching to vent; second ray also large, but shorter, placed between eyes; first spine developed as a fishing-rod, filiform, ending in a small, membranaceous lobe, its base close to that of second, and therefore distant from end of snout, its spine short, tip not reaching middle of second spine; short tentacles, like horns, on anterior part of third spine, over nostrils and under mouth; caudal rounded; pectoral so joined that it can not be turned forward as usual in this group, but rising obliquely backward and upward. Eye slightly longer than snout; mouth brown within.

Color brown, with white spots on body and median fins, 6 of the largest of these each with the center yellowish, largest from once to twice diameter of eye; spots on dorsal fins small; eye golden.

This species reaches a length of about 3 inches. It is known only from Porto Rico, whence it was originally described; not obtained by us.

Antennarius inops Poey, Anal. Soc. Esp. Hist. Nat., X, 1881, 340, Porto Rico; Jordan & Evermann, l. c., 2718, 1898.

† *Antennarius portoricensis*, Stahl, l. c., 246, 1882, Porto Rico.

286. *Antennarius scaber* (Cuvier).

(PLATE 48.)

D. 11-12; A. 7; P. 9 or 10. Anterior dorsal spine as long as second, and provided with 2 long and thick cutaneous flaps at its tip; third dorsal spine not continuous with the soft dorsal; soft dorsal fin terminating at some distance from the caudal, its last ray not extending to root of caudal, if laid backward; dorsal spines, head, back, and sides of body with more or less numerous cutaneous fringes, those of dorsal spines sometimes forming a dense cluster; skin very rough, covered with small spines.

Ground-color yellowish or reddish brown, brightest on back and underneath; body covered with large, irregularly-rounded or oblong brownish spots; head with similar oblong spots; a series of 9 or 10 brown lines radiating from pupil; fins all light-brown or rosy at base, paler on distal portion, each with a number of large roundish black or brownish spots, darkest on anal and pectoral; a series of pale-greenish blotches on interradial membranes near outer margin of each fin; bait pale-yellowish, stem paler; second and third dorsal spines rich brown.

A fish of small size from the Caribbean Sea; known from St. Lucia, Martinique, and Porto Rico; only one specimen, 3 inches long, seined on the coral reef at Mayaguez.

Chironectes scaber Cuvier, Mém. Mus., III, 425, pl. 16, fig. 2, 1817, Martinique.

Antennarius scaber, Jordan & Evermann, l. c., 2722, 1898.

287. *Antennarius nuttingii* Garman. "*Murciélago*."

(PLATE 49.)

D. 11-12; A. 7; V. 5; P. 11; C. 9. In form this species is shorter, more massive anteriorly, and less compressed than either *A. ocellatus* or *A. radiosus*. A transverse section across middle of body is a nearly equilateral triangle. Caudal region short. Head nearly as wide as high; cheeks swollen; forehead rather broad, converging forward on edges. Occipital concavity wide and deep, free from scales in a wide space below ends of first and second dorsal rays, this bare space being apparently for the reception of the fleshy bait-bulb, which latter has 2 elongate lobes. Snout as long as orbit, broad, truncate; chin vertical; symphyseal knob prominent. Mouth wide, subvertical. Eye small; orbit twice as long, hardly more than one-half interorbital space. First and second dorsal rays equal in length, not inclusive of the 2 elongate fleshy-fringed lobes surmounting first; base of first ray standing forward prominently over mouth, being free for some distance; greater portion of second ray free, while the third is connected with the dorsum by the skin, from base nearly to tip. This last ray is larger than either of its fellows. Soft dorsal large, middle rays longest, as long as distance from maxillary to hind edge of operculum, or as long as rays of caudal fin; fin not reaching back to base of caudal rays, fringed; hind margin of caudal convex, fringed; anal moderate, rays prominent in the margin, fin with a blunt angle on outer edge, subtending, when laid up against the tail, one-fourth or more of the length of the caudal rays; rays on pectoral fins extending out beyond margins more noticeably than those of other fins; ventrals small, in most instances with 6 points on outer margin, in one case having but 5; greatest length of caudal nearly one-fourth of total length; length of each maxillary two-thirds of caudal; scales short, small, close-set, harsh to the touch, having none of the velvety appearance.

Color in life: Entire body and head and bases of all fins black, with washing of rusty-brownish; middle of dorsal fin and outer parts of other fins more rusty; margin of dorsal black; each of the other fins with a distinct but narrow creamy-white border; bait white; inside of mouth black.

Until now this fish was known only from the Bahama Banks, where the type was collected by the Nutting expedition. Two specimens, each about 1.75 inches long, were obtained near the coral reef at Mayaguez, in company with the specimen of *A. scaber*.

Antennarius nuttingii Garman, Bull. Iowa Lab. Nat. Hist. 1896, 83, pl. II, Great Bahama Banks; Jordan & Evermann, l. c., 2723, 1898.

Genus 163. **CHAUNAX** Lowe.

Head very large, depressed, cuboid. Mouth large, subvertical; jaws and palate with bands of small teeth. Skin with small, sharp spines. Spinous dorsal reduced to a small tentacle above snout, retractile into a groove; soft dorsal moderate, low; anal short; ventrals small. Gills 2.5; no pseudobranchiæ. Muciferous channels very conspicuous, lateral line prominent, undulate; another series of mucous tubes extending from lower jaws to axil; still another extending backward from snout and maxillary to a point behind eye, when it ceases, uniting with a vertical line which extends from lateral line to lower line, these lines thus inclosing a quadrate area on cheek. Gill-opening small, well behind pectoral under front of soft dorsal.

Small, fantastic, deep-sea fishes.

a. Dorsal rays 11; anal 5; depth 2.5 in length *pictus*, 288
 aa. Dorsal rays 13; anal 7; depth 2.4 in length *nuttingii*




FIG. 109.—*Chaunax pictus*.

288. **Chaunax pictus** Lowe.

Head 1.6; depth 2.5; D. 1, 11; A. 5; P. 11; V. 4; C. 7. Rostral tentacle short, pedicellate; muciferous channels appearing as chain-like rows of pits.

Color in spirits: Pale, with faint rosy shades persisting on back, caudal peduncle, and sides of head.

Deep waters of the Atlantic; known from Madeira, Soudan, Cape Verde, Barbados, off Rhode Island, and elsewhere in the Gulf Stream, in 130 to 428 fathoms. Our single specimen, 2 inches in length, was taken in the beam trawl at Fish Hawk station 6070, 9 miles from Mayaguez, in 220 fathoms.

Chaunax pictus Lowe, Trans. Zool. Soc. Lond. 1846, 339, Madeira; Jordan & Evermann, l. c., 2726, 1898.

?*Chaunax fimbriatus* Hilgendorf, Sitzber. Ges. Naturf. Freunde 1879, 80, Sea of Japan.

?*Chaunax nuttingii* Garman, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 85, near Sand Key light, Florida, in 120 fathoms.

Family LXXVI. **OGCOEPHALIDÆ**. The Bat-fishes.

Head very broad and depressed, snout more or less elevated, trunk short and slender. Mouth not large, subterminal or inferior, lower jaw included; teeth villiform or eardiform. Gill-openings very small, above and behind axils of pectoral fins. Body and head covered with bony tubercles or spines. Spinous dorsal reduced to a small rostral tentacle, which is retractile into a cavity under a prominent process on forehead; in one genus the rostral tentacle is obsolete; soft dorsal and anal fins small and short; ventrals well developed; pectoral fin well developed, its base strongly angled, with long pseudobranchia and 3 aetionosts. Branchiostegals 5; no pseudobranchiæ. There are 8 genera and about 30 species of these fishes, chiefly American, some of them in the deep sea.

OGCOEPHALINÆ:

- a. Disk with frontal region elevated and the snout more or less produced forward; the tail stout; orbits lateral; teeth on vomer and palatines; rostral tentacle present.
- b. Gills 2.5; disk longer than broad..... OGCOEPHALUS, 164

HALIEUTINÆ:

- aa. Disk with frontal region depressed, not elevated above the rest; eyes partly superior; snout rounded, obtuse in front; tail slender.
- c. Dorsal fin present.
- d. Vomer and palatines with teeth..... HALIEUTICHTHYS, 165
- dd. Vomer and palatines toothless.
- e. Disk subcircular; gills 2.5.
- f. Mouth rather small, terminal; prickles feeble..... HALIEUTELLA
- ee. Disk subtriangular; gills 2; prickles very strong..... DIBRANCHUS




FIG. 110.—*Ogcocephalus vespertilio*, dorsal view.




FIG. 111.—*Ogcocephalus vespertilio*, front view.

Genus 164. OGCOEPHALUS Fischer. Sea Bats.

Body stoutish, tapering backward; head very broad and depressed, triangular in form, forehead elevated and produced. Eyes large, lateral. Mouth rather small, subinferior under snout; villiform teeth in bands, on jaws, vomer, and palatines. Skin covered with rough, bony tubercles. Dorsal and anal fins very small; rostral tentacle present, retractile into a cavity under a bony prominence on forehead; ventrals present, 1, 5, well separated; pectorals large, placed horizontally. Gills 2.5. No air-bladder; no pyloric caeca.

Tropical America, in shallow water. Small fishes of singular form, often regarded by the ignorant as venomous.

- a. Snout produced, rostral process pointed, 6 to 10 in length of body..... *vespertilio*, 289
- aa. Snout short, rostral process 12 to 15 times in length of body..... *nasutus*
- aaa. Snout short, rostral tubercle reduced to a button-like tubercle, which is about 25 times in length of body... *radiatus*

289. *Ogcocephalus vespertilio* (Linnaeus). *Bat-fish; Diablo; Murciélago de Mar.*

Head, from tip of upper jaw to gill-opening, nearly one-half the length; depth 5 in length from upper jaw to base of caudal; width 1.8; D. 4; A. 4; rostral process from 6 to 10 (9 in our specimens from Havana); pectoral 4.5; ventral 6; caudal 4.25. Body stoutish, much depressed, rostral process longer than in other species, variable in length; mouth small, maxillary reaching nearly to posterior margin of eye; villiform teeth in bands, on jaws, vomer, and palatines; interorbital flattish, its width less at anterior end of eyes than at posterior; rostral groove longer than broad; body covered with bony protuberances, variable in size, and not very definite in position, lower parts with a shagreen-like covering; posterior edge of pectorals much behind middle of body; ventrals long, reaching outward to edge of the disk-like anterior part of body; origin of dorsal over posterior edge of pectoral; anal under vertical of tips of dorsal rays, and reaching nearly to base of caudal. Pale grayish-brown above, reddish below; back with round black spots, conspicuous in life, but growing fainter and sometimes disappearing in spirits; belly in life a coppery-red; pectoral nearly plain dusky.

West Indies, north to the Florida Keys; common in shallow water. Length 12 inches.

Recorded from Porto Rico by Professor Poey and Dr. Stahl. Here described from a specimen from Havana, Cuba, about 10 inches in length. The length of snout is subject to great variation, but it is never short and button-like, as in *O. radiatus*.

Lophius vespertilio Linnaeus, Syst. Nat., ed. X, 1, 236, 1758, American Seas; after *Lophius fronti unicorni* of Artedi.

Mallucea longirostris Cuvier & Valenciennes, Hist. Nat. Poiss., XII, 452, 1837, Bahia.

Mallhe vespertilio, Poey, Fauna Puerto-Riqueña, 311, 1881; Stahl, l. c., 79 and 165, 1883.

Ogcocephalus vespertilio, Jordan & Evermann, l. c., 2737, 1898.

Genus 165. *HALIEUTICHTHYS* Poey.

Disk subcircular, anteriorly cordiform, head merging into body, very large and much depressed; cranial portion not elevated; interorbital space low and narrow; eyes partly superior; mouth terminal, horizontal, jaws subequal, the lower jaw nearly semicircular; teeth fine, on jaws and palate. Gills 2.5; no gillrakers; gill-openings anterior to pectoral; rostral tentacle very small, retractile; dorsal and anal few-rayed; pectoral large; carpus slender; caudal rounded; skin above sparsely armed with stellate tubercles; lower surface smooth.

a. Surface of body covered with brownish reticulations *aculeatus*, 290

aa. Surface of body blackish, not reticulate; pectoral with a broad black bar.

b. Eye 4; interorbital 2 in eye; nostrils comparatively remote *smithii*, 291

bb. Eye smaller; interorbital narrower; nostrils nearer together *caribbeus*

290. *Halieutichthys aculeatus* (Mitchill).

Head 1.8; depth 6; eye 5; interorbital 1.8 in eye, 8 or 9 in body; D. 5; A. 4; pectoral of 17 rays, 1.5 in head; caudal of 9 rays, 1.6 in head. Disk about as broad as long, its breadth equal to distance from tip of snout to base of last dorsal ray; body above with many conical spines with stellular bases, those at edge of disk usually 3-pointed, divergent; body smooth below; snout very short, obtuse, bridge over rostral cavity with a 3-pointed spine in front, a simple smaller one on each side; simple spines upon each supraorbital margin, and a row of about 5 on lower margin of eyeball, attached by a fleshy flexible membrane; many slender filaments on edge of disk and sides of trunk; supraoral cavity with a club-shaped tentacle.

Color in spirits: Body white below, darker above with obscure brownish markings; pectoral and caudal with 3 dark bars, outer widest and darkest, the 2 inner of the pectoral obscure in our specimen.

West Indies, Gulf of Mexico, and Gulf Stream, in water of moderate depth. One specimen, 2.75 inches long, taken by beam trawl in 75 fathoms at Fish Hawk station 6063, off Mayaguez, January 20.

Lophius aculeatus Mitchill, Amer. Month. Mag., II, 1818, 325, Straits of Bahama.

Halieutichthys aculeatus, Jordan & Evermann, l. c., 2739, 1898.

291. *Halieutichthys smithii* Evermann & Marsh, new species.

Head 2; depth 6; eye 4; interorbital 2 in eye; D. 1-5; A. 4; P. 17; V. 5; C. 9. Body not so wide as in *H. aculeatus*, the anterior edge being more evenly rounded, the lateral edges less divergent, and the posterior angles less salient; spines on body small and obscure, those on margin more prominent; tail smooth, without spines; snout broad; rostral spine small; eye very large; interorbital space wide; margin of body with a prominent fringe of filaments.

Color much as in *H. aculeatus* but lighter and without any reticulations; proximal half of pectoral pale, then a broad jet-black band on about nine of the interradial membranes only (the rays being pale), the outer fourth pale; caudal pale, somewhat dark at tip.




FIG. 112.—*Halieutichthys smithii*.

This species is close to *H. caribbeus* Garman, from which it differs in the much larger eye, wider interorbital, more remote nostrils and the coloration.

One specimen, the type (No. 49537, U. S. N. M.) 3.25 inches long, collected by the *Fish Hawk* at Station 6063, in Mayaguez Harbor, in 75 fathoms, taken by the beam trawl on sand and coral bottom.

We name this interesting species for Dr. Hugh M. Smith, chief of the Division of Scientific Inquiry of the United States Fish Commission.

INDEX.

	Page.		Page.
abildgaardii, Sparisoma	289	arenaceus, Citharichthys	326
Abudefduf	17, 22	arenatus, Priacanthus	166
saxatilis	20, 227	Arrayado	187
Acara Aya	174	Arribo	37
acerosa, Pterogorgia	19, 20	Arroyo	8, 21, 38, 45
Achirus inscriptus	330	Artibeus perspicillatus	16
lineatus	22, 331	ascensionis, Holocentrus	118
Acknowledgments	6	Asymmetron lucayanum	60
aculeatus, Halieutichthys	338	Atherina area	111
acuminatus, Eques	222	laticeps	111
Addison, Matthew G.	6	stipes	110
adscensionis, Epinephelus	152	atlantiens, Tarpon	80
adustus, Ophiocion	219	Atya seabra	15
Aëtobatus narinari	67	Auchenistius stabli	316
Agonostomus monticola	18, 114	Auchenopterus	53
Aguadilla	19, 34, 45	albicaudus	313
Aguaji	157	cingulatus	21, 315
Aguas Buenas	13	fajardo	313
Aguavina	159	faseiatus	315
Agujon	99	rubescens	314
Albacora	122	Augustini, Señor	6
albicaudus, Auchenopterus	313	Aulostomus maculatus	165
albifimbria, Scorpæna	275	aurorenatum, Sparisoma	238
albimentis, Lycodontis	78	aurorubens, Rhomboplites	181
Albula vulpes	82	Anxis thazard	19, 122
album, Haemulon	185	Awaous taïasia	18, 297
Algarrobo	36	aya, Neomanis	174
Algarrobo, Punta de	8	Bairdiella ronchus	218
Alphestes chloropterus	155	Bahama Lancelet	60
Alutera scripta	261	bahianus, Teuthis	254
alutus, Apogonichthys	144	bahiensis, Cypsilurus	104
analís, Neomanis	20, 176	Bajonado	202
Añasco	6	bajonado, Calamus	202
Anchoa Pelada	86	Baker, A. B.	5
Anchovy, Striped	88	Balaju	33, 101, 102
Angel-fish	246	Balao	102
Angel-fish, Blue	252	Baldwin, A. H.	5, 51, 161
Angelichthys ciliaris	252	baldwini, Prionodes	160
Anguilla chrysope	68	Balistes vetula	256
Anisotremus surinamensis	194	Bamboo Fish Pot	31
virginicus	195	Banana-fish	82
Antennarius	4, 19	Barbero	253, 254
inops	335	Barbu	117
nuttingii	335	Barbudo	33, 117
scaber	335	barraeuda, Sphyrana	115
antennatus, Chilomycterus	272	Barreto	300
Aphthalmichthys caribbeus	52, 71	Barriga Blanca	217
apodus, Neomanis	172	bartholomæi, Caranx	131
Apogonichthys alutus	144	batabana, Corvula	217
Apogon sellieauda	143	Bat-fish	285, 338
Apparatus and Methods	30	Bathystoma rimator	192
area, Atherina	111	striatum	193
Archosargus muimaculatus	204	Bayamon	34
arctifrons, Calamus	203	bayamonensis, Gobius	296
arenatus, Pomacanthus	251	Bean, Barton A	280
Ardea cerulea	16	Bean, Tarleton H	100, 280
herodias	16	beanorum, Pontinus	279
ardeola, Tylosurus	99	Beau Gregory	226
Areibo	18, 34, 45	Bello y Espinosa, Domingo	3, 4

	Page.		Page.
Benedict, James E	V	Butterfly	249, 250
Berdugo	222	Cabalito de Mar	109
Berg, Carlos	277	Cabellerote	170
bergii, Scorpaena	276	Cabeza de San Juan	8
Bergudo	222	Cabezón, Corbino	216
Bermuda Chub	211	Cabezote	111
bermudensis, Ficrasfer	319	Cabra Mora	152
Berrios, José V	6	Cabrilla	32, 149, 153
Bianchi, Don Antonio	6, 7	Cachicata	189, 190
bicaudalis, Laetophrys	264	Cachucho	183
Bien & Co., Julius	52	Caçonetta	62
Bigelow, Robert P.	V	Cadena, Punta de la	7
Big-eye	167	cærulea, Ardea	16
Big-eyed Herring	81	Cæsar	192
Scad	129	Cagon de lo Alto	181
bistrispinus, Ryphticus	163	Caguas	18
Bithynis jamaicensis	13	caguite, Sicydium	18, 291
olfersii	13, 18	Cailieu-Tassart	86
bivittatus, Iridio	232	Caji	172
Black Angel	251	Calamus aretifrons	203
Grouper	157	bajonado	202
Grunt	187	calamus	201
Guativere	150	kendalli	201
Moray	77	calidris, Vireo	16
Pilot	226	Caña Tiburones Lagoon	15
Soap-fish	163	Candil	118
Blanquillo	303	Cantherines pullus	258
bleekeriana, Ilisna	86	Canthigaster rostratus	269
Blennies	20	capistratus, Chaetodon	249
Bleniidæ	53	Capitan	230
Blow-fish, Spiny-back	268	Carangidæ	53
Blue Angel-fish	252	Caranx bartholomæi	131
Parrot-fish	239, 2 4	crysos	132
Tang	253	hippos	17, 131
Boea Colorado	189, 190	latus	132
Bocanegra, Susano	7	ruber	130
Bodianus punctatus	150	carbonarium, Hæmulon	188
ruber	150	Carbonero	130
Bollmannia boqueronensis	298	Carcharhinus falciformis	62
Bonaci Acara	157	limbatus	62
bonaci, Mycteroperca	157	Cardona Island	21
bonariense, Hæmulon	187	cardonæ, Coralliozetus	21, 317
Bone-fish	82	caretta, Thalassochelys	25
Bony-fish	81	caribæum, Branchiostoma	59
Boqueron, Ensenada del	21	caribbeus, Aphthalmichthys	52, 71
boqueronensis, Bollmannia	298	Carita	123
Borinquen, Cape	7	carolinus, Trachinotus	139
Bowers, George M	VI, 158	Casabe	136
bowersi, Mycteroperca	158	Catalineta	195, 251
brachiale, Sparisoma	242	Catalufa	166, 167
Brachydeuterus corvinæformis	197	Cataño	17
Brain Coral	19	catenata, Echidna	79
Branchiostoma caribæum	59	Caulolatilus cyanops	303
brasilianus, Gerres	209	Cayo Obispo	23
brasiliensis, Hemirhamphus	1 2	Cayo Sanquido	23
Mugil	112	Cayo Santiago	8, 23
Scorpaena	274	cayorum, Corythoichthys	23, 108
breviceps, Larimus	4, 216	Cavalla	124
bricei, Chaetodon	250	cavalla, Scomberomorus	124
Broad-head, Grubber	90	Cazon de Playa	62
broussonnetii, Gobioides	30	Ceiba	40
Brown Pelican	16, 22	Centropomus parallelus	146
Brownson Deep	8, 24	undecimalis	22, 146
Buchanan, James A	6	Cephalacanthus volitans	285
Bullon	244	Cero	124
Bumper	136	Ceryle alcyon	16
Bureteado	196	Chaetodipterus faber	21, 246
Busck, August	5	Chaetodon	19

	Page.		Page.
Chaetodon bricei	250	Congrio	70
capistratus	249	Conodon nobilis	19, 196
striatus	249	Cony, Horny	259
chalybeius, Chlorophthalmus	93	Cook, O. F.	V
Chapin	262, 263, 264	Coots	25
Characteristics of Aquatic Fauna	24	Coralliozetus cardonæ	21, 317
Chaunax pietus	336	Corbino Cabezon	216
Chelonia mydas	25	Corcobado	33, 133
Cherna	152, 154, 155	Cordillera Central	8, 11
Cherna Americana	154	Corega	37
Criolla	152	coriaceus, Rypiticus	163
de Vivera	154	Corjinua	33
Chicharro	129	Corocoro	186
Chilomycterus	21	coroides, Umbrina	220
antennatus	272	Corvina	33, 215, 216
Chilorhinus suensonii	72	corvinæformis, Brachydeuterus	197
Chinchorro	33	Corvula batabana	217
Chirivita	227, 251	sanctæ-luceæ	217
Chiro	81	Corythoichthys cayorum	23, 108
Chlorophthalmus chalybeius	93	Cotoro	240
chloropterus, Alphestes	155	Verde	239
Chloroscombrus cnrysurus	136	Cow-fish	264
chærostomus, Stolephorus	88	Crawfish	39
Chopa	33	Crevalle	132
Amarilla	204, 211	croco, Pomadasis	198
Blanca	211	croicensis, Searus	20, 244
chrysopteron, Sparisoma	239	cruentatus, Gonfopsis	16
chrysurus, Chloroscombrus	136	Petrometopon	149
Ocyurus	180	Priacanthus	167
chrysyra, Anguilla	68	crumenophthalmus, Trachurops	129
Chub	211	crynos, Caranx	132
Bermuda	211	cubanus, Stolephorus	88
Chucho	67	Cubera	169
Cibi Amarillo	131	Culebra Island	22, 40
Mancho	130	culebra, Malacoctenus	308
Cigar-fish	129	eurema, Mugil	22, 113
ciliaris, Angelichthys	252	Cutlas-fish	125
ciliatus, Monacanthus	258	cyanops, Caulolatilus	303
cinereum, Xystæma	207	cyanopterus, Neomænis	169
cingulatus, Auchenopterus	21, 315	Cynoscion jamaicensis	215
cirratum, Ginglymostoma	60	Cypsilurus bahiensis	104
Citharichthys arenaceus	326	Cytheræa dione	24, 25
spilopterus	326	Daetyloseopus tridigitatus	304
unicornis	325	Dajao	18, 111
Clark, Hubert L.	V	Dall, W. H.	V
Climate	9	Danish West Indies	40
Clupanodon pseudohispanicus	84	Dark-green Parrot-fish	240
Coasts	8	Dasyatis hastata	65
Cochino	256	say	65
Cockeye Pilot	20, 226, 227	Davenport, Richard G.	5
Coe, Wesley R.	V	Decapterus punctatus	129
ceruleus, Searus	244	decoris, Doratonotus	21, 231
Tenthis	253	Deep-water Gurnard	284
Cojinuda	132	delalandi, Malacoctenus	310
Colirubia	180	Demoiselle	227
Colorado	33	Dendroica petechia ruficapilla	16
Comico	166	dentex, Odontoscion	216
Commercial Fisheries	27	Diablo	338
Common Eel	68	Diodon holacanthus	271
Grunt	190	hystrix	271
Pompano	139	dione, Cytheræa	24, 25
Saw-fish	64	Diplectrum radiale	159
Spotted Moray	77	dispilurus, Dules	162
Trunk-fish	263	Diver, Sand	92
Conejo	266	Doctor-fish	254
Coney	149	Doderlein, Pietro	211
Conger Eel	70	Dog Snapper	171
conger, Leptocephalus	70	dominicensis, Tyrannus	16

	Page.		Page.
Donax	20, 25	Fish hawks	23
Doncella	232	Fish-peddlers	37
Dorado	3	Fistularia tabacaria	106
Doratonotus decoris	21, 234	flabellum, Rhipidogorgia	19, 20
megalepis	233	Flamencos Lagoon	15
Dormeur	164	Flasher	164
Dormitator maculatus	289	flavescens, Sparisoma	20, 240
dormitor, Philypnus	288	flavolineatum, Hæmulon	191
Dredging Stations	25	Flint, James M.	V
Drummer, Ground	218	floridae, Siphostoma	107
Jewsharp	221	Fly-catcher	16
Mongolar	215	Flying-fish	103
White-mouth	220	Flying Robin	285
Ducks	23	foetens, Synedus	92
Duerden, J. E.	V	Food-fishes of San Juan market	34
Dules dispilurus	162	Fool-fish	259
Ecaille, Grande	80	fonticola, Fundulus	96
Echeneis naucrates	301	Foster, Lieutenant	68
Echidna catenata	79	French Grunt	191
Eel, Common	68	Frigate Mackerel	122
Conger	70	Fry, Hog-mouth	88
Worm	72	Fundulus fonticola	96
Eleotris pisonis	290	funebri, Lycodontis	77
elongata, Xiphocaris	13, 15	furneri, Micropogon	19, 220
Elops saurus	19, 81	fuscus, Pelecanus	16
elucens, Siphostoma	108	Eupomacentrus	224
El Yunque	7, 8	gabonensis, Vomer	19, 133
Emblemaria pandionis	318	Gaff Topsail	137
Enjambre	149	Galliwasp	92
Ensenada del Boqueron	21	garmani, Stolephorus	89
Honda	24	Gata	60
Epilobocera sinuatifrons	13	Geology	9
Epinephelus adscensionis	152	Gerres brasilians	209
guttatus	32, 153	lineatus	17
morio	154	olisthostomus	209
striatus	152	plumieri	210
Eques acuminatus	222	rhombus	208
Eretmochelys imbricata	25	gibba, Pterophryne	334
Erizo	271	gilberti, Stolephorus	90
Escambron Point	17	Gillias jordanii	21, 307
Eseamuda, Sardina	85	Gill, Theodore	52, 75, 145, 182, 246
Escribano	101, 102	Gill nets	31
Esmeralda	296	Ginglymostoma cirratum	60
Etelis oculatus	183	gladius, Trachinotus	137
Eucinostomus gula	206	Globe-fish	269
harengulus	206	Goat-fish, Red	120
psudogula	205	Yellow	121
Eupomacentrus	17, 19, 22	Gobiesox	17
analis	224	tudes	305
fuscus	224	Gobiidae	53
leucostictus	226	Gobioides broussonnetii	300
exustus, Mytilus	16	Gobius bayamonensis	296
faber, Chaetodipterus	21, 246	lyricus	295
Fajardo	23, 40	oceanicus	296
fajardo, Auchenopterus	313	soporator	294
falcata, Seriola	128	Goggle-eye Jack	129
falcatus, Trachinotus	138	Goggler	129
faleiformis, Carcharhinus	62	gomesii, Ophichthus	75
Fan Coral	19	Goniopsis cruentatus	16
Fan-tail Mullet	113	Gonzalez, Don Luis	6
Fariña, Ricardo Amado	7	gracile, Peristedion	284
fasciatus, Auchenopterus	315	Grass Porgy	203
Fierasfer bermudensis	319	Gray, G. M.	17, 52
File-fish	259	Gray Grunt	186
Fino	150	Snapper	170
Fisheries and Fish Trade	29	Great Blue Heron	16
Fishery Products imported	41	Green Parrot-fish	245
Fishery Statistics	45	Turtle	25, 41

	Page.		Page.
griseus, Neomænis	170	Hind, Red	149
Ground Drummer	218	Hippocampus punctulatus	109
Sparring	91	hippos, Caranx	17
Grouper, Black	157	hispidus, Monacanthus	259
Nassau	152	Hog-choker	22
Red	154	Hog-fish	230
Grubber Broadhead	90	Hog-mouth Fry	88
Grunt, Black	187	holacanthus, Diodon	271
Common	190	Holacanthus tricolor	251
French	191	Holocentrus ascensionis	118
Gray	186	vexillarius	119
Margaret	185	Holothuria surinamensis	20
Open-mouthed	191	hoplomystax, Sparisoma	20, 237
Red-mouthed	192	Horny Cony	259
Striped	186	Horse-eye Jack	132
White	193	Hound-fish	99
Yellow	189	Hueares	8, 23, 40
Guacamaia	245	Hudson, C. B.	51
guacamaia, Pseudocarus	245	Humacao	8
Guachanche, Pélon	116	humeralis, Sardinella	85
guachancho, Sphyræna	116	Hydrography	11
Guanabano	271	Hyporhamphus unifasciatus	17, 101
Guanajibo, Punta de	8	hystrix, Diodon	271
Guanica	21	Ilfish : bleekeriana	86
Lagoon	15	imbricata, Eretnochelys	25
Guaseta	155	incisor, Kyphosus	211
Guativere, Black	150	inops, Antennarius	4, 335
Red	150	inscriptus, Achirus	330
Guavina	288, 297	intermedius, Synodus	92
guavina	22, 289	Iridio bivittatus	232
Guayama	8, 9, 12	kirschii	232
gula, Eucinostomus	206	Irish Pompano	209
Gundlach, Juan	3, 4	Isabelita	252
Gurnard	283	Isopora muricata	19
Deep-water	284	Jallao	185
guttatus, Epinephelus	32, 153	Jaboncillo	154
Hacienda Catalina	6	Jack, Goggle-eye	129
La Perla	6	Horse-eye	132
Romana	6	Yellow	131
Hæmulidae	53	jamaicensis, Bithynis	13
Hæmulon	53	Cynoscion	215
album	185	Jarea	33
bonariense	187	Jenkinsia lamprotenia	84
carbonarium	188	stolifera	84
flavolineatum	191	Jewsharp Drummer	221
macrostomum	186	Jobos Harbor	8
parra	187	jocu, Neomænis	171
plumieri	190	John Mariggle	81
scirrus	189	Jolt-head Porgy	202
Halicutichthys aculeatus	338	jonesi, Siphostoma	108
smithii	339	jordani, Gillias	21, 307
Hamlet	77, 152	Lycodontis	78
Hansard, Arthur C.	6	Jorobado	135
Hard-tail	132	Josea	113
harengulus, Eucinostomus	206	Jurel	33, 132
Hargitt, Chas. W.	V	Just, Emile	7
Harvest-fish	141	kendalli, Calamus	201
hastata, Dasyatis	65	Kendall, W. C.	7
Hawksbill turtle	25, 41	Key to Families of Fishes	54
Hechudo	90	King-fish	124
Hemirhamphus brasiliensis	102	Kingfishers	16, 23
hepatus, Teuthis	254	kirschii, Iridio	232
herodias, Ardea	16	Krug, Leopold	3
Heron, Little Blue	16	Kyphosus incisor	211
Great Blue	16	seetratrix	211
Herring, Big-eyed	81	Labrisomus nuchipinnis	311
Thread	86	Lachnolaimus maximus	230
Hill, Robert T.	8, 9	Lactophrys bicaudalis	264


	Page.		Page.
Lactophrys tricornis	264	mahogoni, Neomænis	179
trigonus	263	Mahogany Snapper	179
triqueter	262	Malacoctenus culebræ	308
Lady-fish	82	delalandi	310
levigatus, Lagocephalus	266	moorei	309
Lagarto	91, 92	puertoricensis	309
Lagocephalus levigatus	266	Manatee	25
lamprotenia, Jenkinsia	84	Mancheño	178
Lancelet, Bahama	60	Mancheva	178
West Indian	59	Mangrove Snapper	170
Lancelet-fish	254	Manjua	86, 88
Lane Snapper	178	Mansfield, Captain	7
Larimus	4	Mapiro	289
breviceps	216	Maps	294
laticeps, Atherina	111	Margaret Grunt	185
latirostris, Trichechus	25	Margate-fish	185
latus, Caranx	132	Mariposa	17, 20, 249, 250, 251, 252
Leather-fish	258, 259	marmoratus, Spheroides	268
Leather-jack	127	Marshall, R. A.	6
Lebranco	112	martiniensis, Menticirrhus	221
Le Fevre, Geo.	V	martiniensis, Upeneus	121
lefroyi, Ulæma	207	Masaguan	289
Leptocephalus conger	70	Matajuelo Real	81
lepturus, Trichiurus	125	maximus, Lachnolaimus	230
leucostictus, Eupomacentrus	226	Mayaguez	13, 35, 44
Lija Colorado	258	McAllister, Don Pablo	6
Lija Trompa	261	Medico	273, 254
limbatus, Carcharhinus	62	meeki, Microgobius	300
lineatus, Achirus	22, 331	megalepis, Doratunotus	233
Gerres	17	megalophthalmus, Neomænis	177
Lion-fish	277	Menticirrhus martiniensis	221
Lisa Francesca	81	Mero	154
Little Blue Heron	16	Cabrilla	158
Liza	17, 33, 112, 113, 258, 259	Guajiro	153
Lizard-fish	92	mesogaster, Parxocœtus	103
Lobotes surinamensis	17, 164	Microgobius meeki	300
Local Fisheries	33	Micropogon furnieri	19, 220
Investigations	15	micurum, Syacium	324
Loggerhead Turtle	25, 41	Millepora alaicornis	20
lorito, Sparisoma	210	Miller, Gerrit S., jr.	5
Loro	236, 238, 240, 241, 244	Milligan, J. D.	7
Colorado	239	Millspangh, Charles Frederick	4
Verde	239, 240	Mojarra	17, 295, 206, 207, 208, 209, 210
Incayanum, Asymmetron	60	Molinari, Nicholas	7
lunatus, Platophrys	322	Mona Passage	24
Luquillo	40, 76	Monacanthus	21
Lutjanide	53	ciliatus	258
Lycodontis albimentis	78	hispidus	259
funcbris	77	Mongolar Drummer	215
jordani	78	Monosira stahli	4
moringa	77	monticola, Agonostomus	18, 114
lyolepis, Stolephorus	89	Moon-fish	135
lyricus, Gobius	295	Moore, H. F.	V, 5
Macabi	82	Moore, J. Percy	V
Machete	125	moorei, Malacoctenus	309
Machuelo	86	Moray, Black	77
mackayi, Siphostoma	107	Moray, Common Spotted	77
Mackerel, Frigate	122	Morena	79
Spanish	123	Verde	77
Yellow	132	moringa, Lycodontis	77
macrolepis, Pontinus	280	morio, Epinephelus	154
macrophthalma, Sardinella	85	Moron	289, 290
macrostomum, Hæmulon	186	Mud-fish	243
maculatus, Aulostomus	105	Mud Parrot-fish	240
Dormitator	289	Mugil brasiliensis	112
Scomberomorus	33, 123	curema	22, 113
Upeneus	120	trichodon	113
Madregal	128	Mullet	17, 112

	Page.		Page.
Mullet, Fan-tail.....	113	parallelus, Centropomus.....	146
White.....	113	Parché.....	219
Muniama.....	207	Parexocetus mesogaster.....	103
Muraenesox savanna.....	71	Pargo.....	33, 176
Mureielago.....	285, 335	Amarilla.....	172
de Mar.....	338	de lo Alto.....	175
muricata palmata, Isopora.....	19	Colorado.....	171, 174
Mutton-fish.....	176, 209	Criollo.....	176
Mycteroperca bonaci.....	157	Guachinango.....	174
bowersi.....	158	Prieto.....	170
mydas, Chelonia.....	25	parra, Hæmulon.....	187
myops, Trachinocephalus.....	91	Parrot-fish.....	29, 236, 238, 240, 241
Myrichthys ocellatus.....	74	Blue.....	239, 244
Mytilus exustus.....	16	Dark-green.....	240
Naguabo.....	8	Green.....	245
narinari, Aëtobatus.....	67	Mud.....	240
Nassau Grouper.....	152	Red.....	239
naucrates, Echeneis.....	301	paru, Peprilus.....	141
Nègre.....	154	parvus, Upeneus.....	121
Neomantis analis.....	20, 176	Patillas River.....	38
apodus.....	172	Pearl-fish.....	319
aya.....	174	pectinatus, Pristis.....	64
cyanopterus.....	169	Pega.....	331
griseus.....	170	Pegador.....	301
jocu.....	171	Peje Puerco.....	256, 258
mahogoni.....	179	Pelada, Anchoa.....	86
megalophthalmus.....	177	Pelceanus fuscus.....	16
synagris.....	20, 178	Peprilus paru.....	141
vivanus.....	33, 175	perfasciatus, Stolephorus.....	88
Nerita peleronta.....	19	Peristedion gracile.....	284
New Species of Fishes.....	52	Permit.....	138
Nigger-fish.....	150	Perro Perro.....	230
niphobles, Sparisoma.....	20, 238	perspicillatus, Artibeus.....	16
nobilis, Conodon.....	19, 196	Peseador.....	334
nuchipinnis, Labrisomus.....	311	petechia ruficapilla, Dendroica.....	16
Nurse Shark.....	60	Peters, W.....	3
nuttingii, Antennarius.....	335	Petrometopon cruentatus.....	149
Obispo.....	67	Pez de Puerco.....	258
oceanicus, Gobius.....	296	Pez Sierra.....	64
Ocean Tang.....	254	Philypnus dormitor.....	288
ocellatus, Platyphrys.....	321	pietus, Chaunax.....	336
ocellatus, Etelis.....	183	Picuda.....	115
Myrichthys.....	74	picudilla, Sphyræna.....	116
Ocyurus chrysurus.....	180	Pilot, Black.....	226
Odontoscion dentex.....	216	Cockeye.....	226, 227
Ogcocephalus vespertilio.....	338	Pintado.....	124, 227
oglinum, Opisthonema.....	86	Piojo.....	81, 82
Ojanco.....	179	pisonis, Eleotris.....	290
Ojon.....	167	Pitirre.....	16
Ojudo.....	167	plagusia, Symphurus.....	332
Old Wench.....	256	Platophrys lunatus.....	322
Old Wife.....	243, 256	ocellatus.....	321
offersii, Bithynis.....	13, 18	Platygyra viridis.....	19
Oligoplites saurus.....	17, 127	Playa de Humacao.....	23
olisthostomus, Gerres.....	209	Pluma.....	201, 202
Open-mouthed Grunt.....	191	plumieri, Gerres.....	210
Ophichthus gomesii.....	75	Hæmulon.....	190
ophioneus, Sphagebranchus.....	73	Scorpæna.....	277
Ophioscion adustus.....	219	Sicydium.....	292
Opisthonema oglinum.....	17, 86	Pocilia vivipara.....	97
Ostrea virginica.....	16	Pocillidae.....	53
Ouatilibi.....	150	Poey, Felipe.....	4
Paguala.....	246	Polydactylus virginicus.....	17, 117
palmata, Isopora muricata.....	19	Pomacanthus arcuatus.....	251
Palmoneta.....	251	Pomadasis crocro.....	198
Palometa.....	137, 138, 141	ramosus.....	198
Palo Seco.....	17	Pompano.....	137
pandionis, Emblemaria.....	318	Common.....	139

	Page.		Page.
Pompano, Irish	209	Red Parrot-fish	239
Round	138	Red Snapper	174
Pompon	194	Redstart, American	16
Ponce	21, 37, 43	regalis, Scomberomorus	124
Pontinus beanorum	279	Rhipidogorgia flabellum	19, 20
macrolepis	280	rhombus, Gerres	208
Porcupine-fish	271	Rhomboplites aurorubens	181
Porgy, Grass	203	Riddle, Oscar	7, 17, 52, 176
Jolt-head	202	rimator, Bathystoma	192
Saucer-eye	201	Rincon	7
Shad	203	Río Añasco	14
Pork-fish	195	Arecibo	13
Porto Rico, Geological Features of	7	Bayamon	13
Harbors of	7	Blanco	14
Position and Size of	7	Caguinas	13
Portugais	251	Coamo	14
Priacanthus arenatus	166	Culebrinas	13
cruentatus	167	Descalabrado	14
Prices of Food-fishes	34	Don Alonzo	13
Prionodes baldwini	160	Fajardo	14
Prionotus punctatus	283	Grande de Arecibo	18
Pristis pectinatus	64	Guamani	14
productus, Stolephorus	90	Jacaguas	11
pseudohispanicus, Clupanodon	84	Loíza	12
pseudogula, Eucinostomus	205	Luquillo	15
Pseudoscarus guacamúia	245	Naguabo	14
Pterogorgia acerosa	19, 20	Portugues	14
Pterophryne gibba	334	Turabo	13
Puereo Espino	271	Yahueca	14
Puerto Arenas	40	Yauco	14
de Fajardo	8	Robalo	22, 33, 146
Isabel Segunda	40	Robin, Flying	285
Mosquito	40	Round	129
Negro	40	Rock Beauty	251
Real	37, 40	Rock-fish	158
Real de Cabo Rojo	8, 20	Rock-hind	152, 158
puertoricensis, Malacoctenus	309	Rock Salmon	128
Puertorriqueño	7	ronchus, Bairdiella	218
Puffer	269	Roneo	187, 218
Sharp-nosed	269	Amarillo	189
Smooth	266	Blanco	198
Southern	267	Carbonero	188
Spiny	272	Condenado	191
pullus, Cantherines	258	Prieto	187
punctatus, Bodianus	150	Ronco	190
Decapterus	129	rostratus, Canthigaster	269
Prionotus	283	Round Pompano	138
punctulatus, Hippocampus	109	Robin	129
Punta de Mata Redonda	8	Rubber-fish	211
de Santiago	8	ruber, Bodianus	150
Lima	8, 23	Caranx	130
Santiago	39	rubescens, Auchenopterus	314
Puppy-fish	141	rubripinne, Sparisoma	241
Quia-quia	129	Runner	127, 132
Quiebra	127	Rypticus bistris-pinus	163
Rabrubia	180	coriaceus	163
radiale, Diplectrum	159	saponaceus	163
Raiado	178	Sabalo	80
Rail	16	Sabanita	37
Ramírez, Miguel	7	Sable	33, 125
ramosus, Pomadasis	198	Saga	297
raphidoma, Tylosurus	99	Sailor's Choice	187
Rascacio	277	Salmonete	120
Rathbun, M. J.	V	Salmonete Amarilla	121
Raya	65	Salmon, Rock	128
Ray, Sting	65	Sama	164, 176
Red Goat-fish	120	San Antonio Bridge	17
Red Grouper	154	sanctæ-luciæ, Corvula	217
Red Guativero	150	Sand-diver	92
Red-hind	32, 149, 153	Sandpiper	16
Red-mouthed Grunt	192	San Francisco, Cape	7

	Page.		Page.
San Geronimo	17	Siphostoma jonesi	108
San Juan	33, 43	mackayi	107
San Juan Market	17	Sirajo	292
Santa Cruz	40	Sisi	195
Santa Cruz Fishermen	22	Slippery Dick	232
Sapo	334	Smith, H. M.	52, 339
saponaceus, Rypiticus	163	smithii, Halieutichthys	339
Sardina	17, 84, 85	Smooth Puffer	266
de España	84	Snapper, Dog	171
Escanuda	85	Gray	170
Sardine	33	Lane	178
Sardinella humeralis	85	Mahogany	179
macrophthalmia	85	Mangrove	170
Saucer-eye Porgy	201	Red	174
saurus, Elops	19, 81	Silk	175
Oligoplites	17, 127	Snook	146
Savalle	80	Soap-fish	92, 163
Savanilla	80	soporator, Gobius	294
savanna, Murænesox	71	Southern Puffer	267
Savola	125	Sting ray	65
Saw-fish, Common	64	Spade-fish	21, 246
saxatilis, Abudeiduf	20, 227	Spanish Mackerel	123
say, Dasyatis	65	Sparisoma	53
Scabbard-fish	125	abildgaardi	239
scaber, Antennarius	335	aurofrenatum	238
scabra, Atya	15	brachiale	242
Scad	129	chrysopteron	239
Scad, Big-eyed	129	flavescens	20, 240
Scamaroni, Guillermo R.	7	hoplomystax	20, 237
Scaridæ	53	lorito	240
Scarus œruleus	244	niphobles	20, 238
croicensis	20, 244	rubripinne	241
vetula	243	viride	240
Schoolmaster	172	xystrodon	20, 236
scurus, Hæmulon	189	Spearing, Ground	91
Scomberomorus cavalla	124	spengleri, Spheroides	267
maiculatus	33, 123	Sphagebranchus ophioneus	73
regalis	124	Spheroides marmoratus	268
Scorpaena albifimbria	275	spengleri	267
bergii	276	testudineus	269
brasiliensis	274	Sphyræna barracuda	115
grandicornis	277	guachancho	116
plumieri	277	pieudilla	116
scripta, Alutera	261	spilopterus, Citharichthys	326
Sea-cucumber	20	Spiny-back Blow-fish	268
Sea-horse	109	Spiny Puffer	272
Sea Serpent	75	Spotted Sting Ray	67
sectatrix, Kyphosus	211	Trunk-fish	264
Selene vomer	17, 135	Sprat	86
sellicauda, Apogon	143	Squirrel-fish	118
Seriola falcata	128	St. Thomas	40
Serranidæ	53	Fishermen	22
Seti	296	Stag-horn Coral	19
Setophaga ruticilla	16	Stahl, A	3, 4, 8, 316
Shad Porgy	203	stahli, Auchenistius	316
Shark, Nurse	60	Monosira	4
Shark Sucker	301	Stearns, Silas	154, 247
Sharp-nosed Puffer	269	Stevenson, Charles G	6
Shrimp	18	Sting Ray	65
Sicydium caguite	18, 291	Southern	65
plumieri	292	Spotted	67
Sierra	33	stipes, Atherina	110
Sierra de Cayey	8	Stolephorus	53
Sierra Luquillo	8	chærostomus	88
Silk Snapper	175	cnbanus	88
Silver-fish	125	garmani	89
Silver King	80	gilberti	90
Simpson, Chas. T.	V	lyolepis	89
sinuatifrons, Epiloboera	13	perfasciatus	88
Siphostoma elucens	108	productus	90
floridæ	107	stolifera, Jenkinsia	84

	Page.		Page.
striatum, Bathystoma	193	Tumble-nose	214
striatus, Chaetodon	249	Tylosurus ardeola	99
Epinephelus	152	raphidoma	99
Striped Anchovy	88	timucu	99
Grunt	186	Tyrannus dominicensis	16
Sucking-fish	301	Ulæma lefroyi	207
suensonii, Chilorhinus	72	Umbrina coroides	220
Summer Yellowbird	16	undecimalis, Centropomus	22, 146
Sundevall, Carl	4	unicornis, Citharichthys	325
surinamensis, Anisotremus	194	Unicorn-fish	261
Holothuria	20	unifasciatus, Hyporhamphus	17, 101
Lobotes	17, 164	unimaculatus, Archosargus	204
Swell-toad	267	Upeneus maculatus	120, 121
Syacium micurum	324	parvus	121
Symphurus plagusia	332	Urban, Ignatius	4
synagris, Neomænis	20, 178	Vaqueta de dos Colores	251
Synodus fetens	92	variegatus, Toxopneustes	20
intermedius	92	Vaughan, T. Wayland	V
tabacaria, Fistularia	106	Vega Baja	3
taiasica, Awaous	18, 297	Verrugato	220
Tambor	267, 269	vesperillo, Ogocephalus	338
Tamboril	266, 267, 269	vetula, Balistes	256
Tang, Blue	253	Scarus	243
Ocean	254	vexillarius, Holocentrus	119
Tarpon atlanticus	80	Vidal, Don Ignacio G	6, 20
Tarpum	80	Vieja, Colorado	240
Ten-pounder	81	Viejo	185, 198, 239, 241, 243
testudineus, Spheroides	269	Vieques Island	22, 40
Teuthis bahianus	254	Vireo calidris	16
ceruleus	253	virginica, Anisotremus	195
hepatus	254	Ostrea	16
Thalassochelys caretta	25	Polydaetylus	17, 117
thazard, Auxis	19, 122	viride, Sparisoma	240
Thread Herring	86	viridis, Platygyra	19
timucu, Tylosurus	99	vivanus, Neomænis	33, 175
Tom-tate	192	vivipara, Pecten	97
Tongue-fish	332	Volador	103, 104, 285
Topography	8	volitans, Cephalacanthus	285
Toro	166, 264	Vomer gabonensis	19, 133
Tortola	40	vomer, Selene	17, 135
Fishermen	22	vulpes, Albula	82
Tortuguero Lagoon	15	Ward, Henry B	V
Toxopneustes variegatus	20	Weirs	32
Trachinocephalus myops	91	West Indian Lancelet	59
Trachinotus	17	Wheeler, W. M	V
carolinus	139	White-bill	85
falcatius	138	White, S. Reynolds	6
glaucus	137	White Grunt	193
Trachurops erumcnophthalmus	129	White-mouth Drummer	220
Treadwell, A. L.	V	White Mullet	113
Tremba	303	Wilson, H. M	11, 12
Trichechus latirostris	25	Wilson, H. V	V
Trichurus lepturus	125	Wilson, J. B	5
trichodon, Mugil	113	Worm Eel	72
tricolor, Holacanthus	251	Wright, Lieutenant	6
tricornis, Lactophrys	264	Xiphocaris elongata	13, 15
tridigitatus, Dactyloscopus	304	Xurel	132
trigonus, Lactophrys	263	Xystema cinereum	207
Trinidad, Francisco	7	xystrodon, Sparisoma	20, 236
Triple-tail	164	Yellowbird, Summer	16
triqueter, Lactophrys	262	Yellow Goat-fish	121
Trompetero	105, 106	Grunt	189
Trumpet-fish	105, 106	Jack	131
Trunk-fish	262	Mackerel	132
Common	263	Tail	20, 180
Spotted	264	Zaga	297
tudes, Gobiesox	305	Zapatero	127




C. B. Hudson ad nat. del.

ANGUILLA CHRYSYPA RAFINESQUE. COMMON EEL; ANGUILLA.
ABOUT ONE-THIRD NATURAL SIZE.




A. H. Baldwin ad nat. del.

LYCODONTIS JORDANI EVERMANN & MARSH. TYPE.
ABOUT HALF NATURAL SIZE.




A. H. Baldwin ad nat. del.

HOLOCENTRUS ASCENSIONIS (OSBECK). SQUIRREL-FISH; CANDIL.
NATURAL SIZE.



A. H. Baldwin ad nat. del.

UPENEUS MACULATUS (BLOCH). RED GOAT-FISH; SALMONETE.
NATURAL SIZE.



A. H. Baldwin ad nat. del.

4 UPENEUS MARTINICUS CUVIER & VALENCIENNES. YELLOW GOAT-FISH; SALMONETE.
ABOUT 1/2 TIMES NATURAL SIZE.

JULIUS BIEN & CO. LITH. N. Y.






C. B. Hudson ad nat. del.

SCOMBEROMORUS MACULATUS (MITCHILL). SPANISH MACKEREL; SIERRA.
ABOUT HALF NATURAL SIZE.

JULIUS BIEN & CO. LITH. N. Y.




A. H. Baldwin ad nat. del.

OLIGOPLITES SAURUS (BLOCH & SCHNEIDER). LEATHER-JACK, ZAPATERO.
ABOUT NINE-TENTHS NATURAL SIZE.



C.B. Hudson ad nat del.

DECAPTERUS PUNCTATUS (AGASSIZ). SCAD, CABALLO AZUL.
ABOUT NATURAL SIZE.



C. B. Hudson ad. nat. del.


CARANX CRYSOS (MITCHILL). RUNNER.
ABOUT THREE-FIFTHS NATURAL SIZE.



C.B. Hudson ad nat. del.

TRACHINOTUS CAROLINUS (LINNÆUS). COMMON POMPANO, PÁMPANO.
ABOUT HALF NATURAL SIZE.

JULIUS BIENÉ LITH. N.º 11



C. B. Hudson ad nat. del.

EPINEPHELUS ADSCENSIONIS (OSBECK). ROCK HIND, CABRA MORA.
ABOUT THREE-FOURTHS NATURAL SIZE.

JULIUS BIEN & CO. LITH. N. Y.





C. B. Hudson ad nat. del.

EPINEPHELUS STRIATUS (BLOCH). NASSAU GROUPEE, CHERNA.
ABOUT THREE-FOURTHS NATURAL SIZE.



C. B. Hudson ad nat. del

EPINEPHELUS MACULOSUS (CUVIER & VALENCIENNES). RED HIND; CABRILLA.
ABOUT THREE-FIFTHS NATURAL SIZE.

JULIUS BIEN & CO. LITH. N. Y.





C. B. Hudson ad nat. del.

EPINEPHELUS MORIO (CUVIER & VALENCIENNES). RED GROUPER, MERO
NEARLY HALF NATURAL SIZE.

JULIUS BIEN & CO LITH N Y






A. H. Baldwin ad nat. del.

PRIONODES BALDWINI EVERMANN & MARSH. TYPE.

ABOUT 2½ TIMES NATURAL SIZE.



A. H. Baldwin ad nat. del.

PRIACANTHUS ARENATUS CUVIER & VALENCIENNES. CATALUFA, TORO.
ABOUT NINE-ELEVENTHS NATURAL SIZE.



C. B. Hudson ad nat. del.

NEOMÆNIS GRISEUS (LINNÆUS). GRAY SNAPPER; PARGO PRIETO.
ABOUT THREE-FOURTHS NATURAL SIZE.




C. B. Hudson ad nat. del.

NEOMÆNIS JOCU (BLOCH & SCHNEIDER). DOG SNAPPER; JOCÚ.
ABOUT THREE-FOURTHS NATURAL SIZE



C. B. Hudson ad nat. del.

NEOMÆNIS APODUS (WALBAUM). SCHOOLMASTER; CAJI
ABOUT FOUR-FIFTHS NATURAL SIZE.



A. H. Baldwin ad nat. del.


NEOMÆNIS AYA (BLOCH). RED SNAPPER, PARGO COLORADO.
ABOUT THREE-SEVENTHS NATURAL SIZE.






C. B. Hudson ad nat. del.

NEOMÆNIS ANALIS (CUVIER & VALENCIENNES). MUTTON-FISH; PARGO.
ABOUT TWO-FIFTHS NATURAL SIZE.




A. H. Balgwin ad nat. del.

NEOMÆNIS SYNAGRIS (LINNÆUS). LANE SNAPPER.
ABOUT THREE-FOURTHS NATURAL SIZE.



C.B. Hudson ad nat. del.

OCYURUS CHRYSURUS (BLOCH). YELLOW-TAIL; FABIRUBIA; COLORUBIA.
ABOUT THREE-FOURTHS NATURAL SIZE.




C. B. Hudson ad nat. del.

HÆMULON ALBUM CUVIER & VALENCIENNES. MARGATE-FISH.
ABOUT THREE-SEVENTHS NATURAL SIZE.




C. B. Hudson ad nat. del.

CALAMUS BAJONADO (BLOCH & SCHNEIDER). JOLT-HEAD PORGY; PLUMA.
ABOUT THREE-FOURTHS NATURAL SIZE.



A. H. Baldwin ad nat. del.


ARCHOSARGUS UNIMACULATUS (BLOCH).
NATURAL SIZE.



A. H. Baldwin ad nat. del.

EUPOMACENTRUS FUSCUS (CUVIER & VALENCIENNES).

ABOUT 1½ TIMES NATURAL SIZE.




A. H. Baldwin ad nat. del.

EUPOMACENTRUS LEUCOSTICTUS (MÜLLER & TROSCHEL).
NEARLY TWICE NATURAL SIZE.



A. H. Baldwin ad nat. del.

DORATONOTUS DECORIS EVERMANN & MARSH. TYPE.
ABOUT 2½ TIMES NATURAL SIZE.



A. H. Baldwin ad nat. del.

SPARISOMA ABILDGAARDI (BLOCH). RED PARROT-FISH, LORO COLORADO.
ABOUT THREE-FOURTHS NATURAL SIZE.


JULIUS BIEN & CO. LITH. N. Y.



A. H. Baldwin ad nat. del.


SCARUS VETULA BLOCH & SCHNEIDER
ABOUT THREE-FOURTHS NATURAL SIZE

JULIUS BIEN & CO. LITH. N.Y.



A. H. Baldwin ad nat. del.

SCARUS COERULEUS (BLOCH). BLUE PARROT-FISH; LORO.
ABOUT THREE-FIFTHS NATURAL SIZE




A. H. Baldwin ad nat. del.

CHÆTODIPTERUS FABER (BROUSSONNET). SPADE-FISH.
ABOUT THREE-FIFTHS NATURAL SIZE.



A. H. Baldwin ad nat. del.

CHAETODON STRIATUS LINNÆUS. BUTTERFLY-FISH; MARIPOSA.
NATURAL SIZE.



A. H. Baldwin ad natid.

CHÆTODON CAPISTRATUS LINNÆUS. BUTTERFLY-FISH, MARIPOSA.
NATURAL SIZE.



A. H. Baldwin ad. nat. del.


HOLACANTHUS TRICOLOR (BLOCH). ROCK BEAUTY; PALMONETA.

ABOUT 1 1/2 TIMES NATURAL SIZE.




A. H. Baldwin ad nat. del.

ANGELICHTHYS CILIARIS (LINNÆUS). BLUE ANGEL-FISH, MARIPOSA.
ABOUT THREE-FOURTHS NATURAL SIZE.



A. H. Baldwin ad nat. del.

TEUTHIS CŒRULEUS (BLOCH & SCHNEIDER). BLUE TANG; MEDICO.
ABOUT TWICE NATURAL SIZE.



A. H. Baldwin ad nat. del.

BALISTES VETULA LINNÆUS.
ABOUT THREE-FOURTHS NATURAL SIZE.




A. H. Baldwin ad nat. del.

LACTOPHRYS BICAUDALIS (LINNÆUS). SPOTTED TRUNK-FISH; CHAPIN.
NATURAL SIZE

JULIUS BIEN & CO LITH. N. Y.






A. H. Baldwin ad nat. del.


SPHEROIDES TESTUDINEUS (LINNÆUS). PUFFER, TAMBORIL.
ABOUT 1½ TIMES NATURAL SIZE.

JULIUS BIEN & CO. LITH. N. Y.



A. H. Baldwin ad nat. del.

CHILOMYCTERUS ANTENNATUS (CUVIER).
ABOUT 1½ TIMES NATURAL SIZE.



A. H. Baldwin ad nat. del.


PONTINUS MACROLEPIS GOODE & BEAN.
NATURAL SIZE.



A. H. Baldwin ad nat. del.

PERISTEDION GRACILE GOODE & BEAN
ABOUT 1 4/5 TIMES NATURAL SIZE.

JULIUS BIEN & CO. LITH. N. Y.




A. H. Baldwin ad nat. del.

DORMITATOR MACULATUS (BLOCH).
NATURAL SIZE.




A. H. Baldwin ad nat. del.

LABRISOMUS NUCHIPINNIS (QUOY & GAIMARD).
NATURAL SIZE.



A. H. Baldwin ad nat. del.

AUCHENOPTERUS FAJARDO EVERMANN & MARSH, TYPE.
NEARLY TWICE NATURAL SIZE.



A. H. Baldwin ad nat. det.

ANTENNARIUS SCABER (CUVIER).
ABOUT $1 \frac{1}{3}$ TIMES NATURAL SIZE.





A. H. Baldwin ad nat. del.

ANTENNARIUS NUTTINGII GARMAN. MURCIELAGO.
ABOUT 1½ TIMES NATURAL SIZE.



ABBREVIATIONS

Material	Color or Shade	Other qualities
M. - Mud	bl. - black	hd. - hard
S. - Sand	gr. - gray	sd. - soft
G. - Gravel	dk. - dark	sl. - fine
Sh. - Shells	gn. - green	cr. - coarse
Z. - Zircon	br. - brown	brk. - broken
Sp. - Spores	bl. - blue	ry. - rocky
C. - Clay	lt. - light	sk. - sticky

TIDES

	Mean
Plane of reference is Mean Low Water	Mean
Average time of High Water after Mean Meridian passage	0h 22m
Average time of Low Water after Mean Meridian passage	0 39
Average height of High Water above the plane of reference	1.1h.



The predicted time and height of the tide can be obtained from the **TIDE TABLES** published annually by the United States Coast and Geodetic Survey.

Note: Underlined figures refer to number of dredging Sta. U.S. Fish Commission Steamer Fish Hawk 1899.


ELEVATIONS
All elevations are referred to high water and are expressed in meters and by contour lines showing successive differences of 5 meters.


BUOYS
Red Buoy to be left to starboard hand
Black Buoy to be left on entering Port land
Black and Red horizontal stripes - Danger Buoy
Black and White perpendicular stripes - Channel Buoy
C.N. or S. signal flag, day or night

SOUNDINGS
The Soundings are expressed in fathoms and show the depth at mean low water.
The Short curve is shown thus:
- 12 -
- 18 -
- 24 -
.....



SAN JUAN HARBOR
PORTO RICO






Note: ...ington, D.C.

MAYAGUEZ BAY

From a survey in 1836 by the Department of Public Works Porto Rico. Obs. Spot @ Lat 18° 11' 56" N Long 67° 09' 00" W


The coast line and soundings shown in hair-line are from earlier surveys by the French Government in 1846 and by Capt. O. Kuhl in 1851



6065



MONA ISLAND.

From a Survey by Captain O. Kuhl in 1832. H.W.F. & C.V. 15' Rise about 2 feet. Playa Pajaro chimney Lat 18° 2' 15" N Long 67° 21' 20" W



AGUADILLA BAY.

From the Spanish Portulano 1618. Aguadilla: Lat 18° 25' 20" N Long 67° 09' 24" W. Good Anchorage off the centre of the town in 10 to 14 fathoms. Much swell and surf with the wind at N.W.




PUERTO DE GUANICA

South Side of Porto Rico. From a U.S. Government Survey in 1838, with additions from a Spanish Government Survey in 1875. Pta de la Meseta: Lat 17° 57' 35" N Long 66° 03' 42" W



View of the Entrance, distant 3 miles.




SOUNDINGS IN FATHOMS. HEIGHTS IN FEET.

Note: Underlined Figures refer to number of dredging Sta. U.S. Fish Commission Steamer Fish Hawk 1899.

Adapted from chart N° 372 Hydrographic Office Washington, D.C. Additions by U.S. Fish Commission, 1899.










GEOGRAPHICAL POSITION

Name	Latitude	Longitude West from Greenwich
Port San Juan Light	18° 28' 58"	66° 07' 28" 4 24" 23.9"

PORTO RICO



Note: Underlined figures refer to number of dredging stations U.S. Fish Commission Steamer Fish Hawk in 1899



TIDES

	San Juan	Port Ponce	Moná I.
The plane of reference is Mean Low Water			
Average time of High Water after Moon's Meridian passage	8 ^h 21 ^m	12 ^h 42 ^m	6 ^h 05 ^m
Average time of Low Water after Moon's Meridian passage	2 20	1 25	12 18
Average height of High Water above the plane of reference	1.1 ft.	0.8 ft.	2.0 ft.

