

The Old World Stenomidae:  
A Preliminary Survey of the Fauna,  
Notes on Relationships,  
and Revision of the Genus *Eriogenes*  
(Lepidoptera: Gelechioidea)

W. DONALD DUCKWORTH

## SERIAL PUBLICATIONS OF THE SMITHSONIAN INSTITUTION

The emphasis upon publications as a means of diffusing knowledge was expressed by the first Secretary of the Smithsonian Institution. In his formal plan for the Institution, Joseph Henry articulated a program that included the following statement: "It is proposed to publish a series of reports, giving an account of the new discoveries in science, and of the changes made from year to year in all branches of knowledge." This keynote of basic research has been adhered to over the years in the issuance of thousands of titles in serial publications under the Smithsonian imprint, commencing with *Smithsonian Contributions to Knowledge* in 1848 and continuing with the following active series:

*Smithsonian Annals of Flight*  
*Smithsonian Contributions to Anthropology*  
*Smithsonian Contributions to Astrophysics*  
*Smithsonian Contributions to Botany*  
*Smithsonian Contributions to the Earth Sciences*  
*Smithsonian Contributions to Paleobiology*  
*Smithsonian Contributions to Zoology*  
*Smithsonian Studies in History and Technology*

In these series, the Institution publishes original articles and monographs dealing with the research and collections of its several museums and offices and of professional colleagues at other institutions of learning. These papers report newly acquired facts, synoptic interpretations of data, or original theory in specialized fields. These publications are distributed by mailing lists to libraries, laboratories, and other interested institutions and specialists throughout the world. Individual copies may be obtained from the Smithsonian Institution Press as long as stocks are available.

S. DILLON RIPLEY  
*Secretary*  
Smithsonian Institution

The Old World Stenomidae:  
A Preliminary Survey of the Fauna,  
Notes on Relationships,  
and Revision of the Genus *Eriogenes*  
(Lepidoptera: Gelechioidea)

*W. Donald Duckworth*



SMITHSONIAN INSTITUTION PRESS

City of Washington

1973

## ABSTRACT

Duckworth, W. Donald. The Old World Stenomidae: A Preliminary Survey of the Fauna, Notes on Relationships, and Revision of the Genus *Eriogenes* (Lepidoptera: Gelechioidea). *Smithsonian Contributions to Zoology*, number 147, 21 pages, 9 figures, 7 maps, 1973.—A preliminary review of the presently known composition of the microlepidopterous family Stenomidae in the Old World is provided, including observations on distribution of the family, relationships with other families in the superfamily Gelechioidea, and a catalog of the Old World genera and species. The genus *Eriogenes* is revised and transferred from the Xyloryctidae to the Stenomidae, and one new species is described.

OFFICIAL PUBLICATION DATE is handstamped in a limited number of initial copies and is recorded in the Institution's annual report, *Smithsonian Year*. SI PRESS NUMBER 4777. SERIES COVER DESIGN: The coral *Montastrea cavernosa* (Linnaeus).

---

### Library of Congress Cataloging in Publication Data

Duckworth, W. Donald, 1935-

The Old World Stenomidae.

(Smithsonian contributions to zoology, no. 147)

Bibliography: p.

1. Stenomidae . I. Title. II. Series: Smithsonian Institution. Smithsonian contributions to zoology, no. 147.

QL1.S54 no. 147. [QL561.S8] 591'.08s [595.7'8] 72-13089

---

For sale by the Superintendent of Documents,  
U.S. Government Printing Office, Washington, D.C. 20402  
Price 40 cents domestic postpaid or 30 cents GPO Bookstore

# The Old World Stenomidae: A Preliminary Survey of the Fauna, Notes on Relationships, and Revision of the Genus *Eriogenes* (Lepidoptera: Gelechioidea)

*W. Donald Duckworth*

## Introduction

For many years the microlepidopterous family Stenomidae has been viewed as a uniquely New World family with little, if any, representation in the Old World fauna. The development of this concept is difficult to trace precisely; however, to a large extent it stems from the confusion among earlier workers over the relationship between the Stenomidae and the closely related family Xyloryctidae coupled with a lack of detailed studies dealing with either family.

Although both families were originally proposed by Edward Meyrick, his concepts concerning them fluctuated through the years primarily due to his steadfast refusal to utilize characters other than wing venation for his higher classification. This dependance on a single character source led to numerous misconceptions, particularly with the higher categories. As the venational characters he originally used to separate the xyloryctids and stenomids began to break down with the continuing discovery and description of new species, Meyrick (1915) merged the two families under the older name Xyloryctidae. From this point he pro-

ceeded during the remainder of his lifetime to describe hundreds of species from both the Old and New World tropics under the family Xyloryctidae.

Other workers, principally August Busck, continued to utilize the original two-family concept placing the New World species and genera in Stenomidae. This geographical distinction was further emphasized by Busck (1934) when in his Stenomidae portion of the *Lepidopterorum Catalogus* series he only included genera and species from the New World. Thus, through omission, the genera and species from areas other than North and South America remained largely unknown and, until the present, uncataloged. In addition, the Xyloryctidae (*sensu stricto*) have never been cataloged and remain essentially unstudied.

In recent years a number of papers have reflected the distinctiveness of the two families; however, due to the nature of the studies (faunistic, type catalog) they do not provide a sufficiently clear picture of the two families to serve as a basis for future studies. Clarke (1955a) provides a thorough account of Meyrick's treatment of xyloryctids and stenomids in his introductory material to the catalog of Meyrick types; however, his detailed treatment (1955b) of the type specimens does not include species from Australia where the majority of xyloryctids and Old World stenomids occur.

---

*W. Donald Duckworth, Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560.*

Diakonoff (1954) provided a complete and accurate definition of the two families based on his study of the Microlepidoptera of New Guinea. His descriptions coupled with those of Common (1970) based on the Australian fauna provide the most comprehensive characterization of the two families yet published. Both studies, however, are restricted in scope and the latter provides only fragmentary information below family level.

During the course of revisionary studies of the Western Hemisphere Stenomidae conducted by the present author in recent years, the lack of information concerning the Old World elements of the family has presented increasing difficulties, particularly in relation to higher category relationships and evolutionary trends. The present paper is intended to provide a partial remedy to this situation and, more importantly, to serve as a starting point for more comprehensive studies. An attempt has been made to provide as complete a catalog as possible of the previously described taxa that have been assigned to the family or that have been determined to be more appropriately placed in the family. No finality is suggested for this catalog, rather it is anticipated that numerous modifications will become necessary as additional information is obtained.

A detailed revision of the genus *Eriogenes* Meyrick is included and represents the first step toward a more thorough understanding of the Old World stenomid fauna and its relation to the remainder of the family. Previously assigned to the Xyloryctidae, the genus is herein transferred to the Stenomidae and a new species is described.

The comments which follow concerning the relationships between the Xyloryctidae and Stenomidae, as well as the statements concerning the distribution of the Old World fauna, are preliminary. A great deal more work remains to be accomplished, especially at the generic and specific levels, before more precise evaluations can be made. In the meantime, it seems appropriate to bring together as much of the existing information as possible in order to facilitate future studies.

The author wishes to acknowledge with gratitude the aid and cooperation of the following individuals and institutions who, through their support and encouragement, have materially aided the present study: Mr. Allan Watson, Mr. P. E. S. Whalley,

Dr. Klaus Sattler, British Museum (Natural History); Dr. J. F. Franclemont, Cornell University; Dr. I. F. B. Common, Commonwealth Scientific and Industrial Research Organization, Canberra, Australia, for lending types and other specimens in their charge for study. Special thanks are extended to Dr. Ian F. B. Common, C. S. I. R. O., Canberra, Australia and Mr. John S. Dugdale, Department of Scientific & Industrial Research, Nelson, New Zealand, for their aid in documenting the stenomid fauna of their respective areas.

The author also wishes to acknowledge the assistance of Mr. George Venable, departmental illustrator, for the artwork; Mrs. Vera Milbank, technician, for bibliographic assistance; and the National Museum of Natural History Photographic Laboratory for the photographs.

### Classification and Relationships

The greatest difficulty encountered in attempting to define the family Stenomidae is the lack of knowledge concerning the two most closely related families, Xyloryctidae and Oecophoridae. Fortunately, at least for purposes of the present study, the stenomids are relatively distinct morphologically, and while the oecophorids and xyloryctids are yet to be clearly separated from each other, it is possible to distinguish the stenomids from them to a satisfactory degree and to indicate certain relationships.

The Stenomidae are distinguished by the following combination of characters: labial palpus up-curved, generally robust, apical segment usually shorter than second segment; antenna long-ciliated in the male, scape without pecten; ocelli absent; maxillary palpus 4-segmented, usually partially encircling base of proboscis; forewing with veins 7 and 8 ( $R_4$  and  $R_5$ ) usually separate, vein 2 ( $CuA_2$ ) usually arising at or near lower angle of discal cell; hind wing with vein 8 ( $Sc + R_1$ ) approaching 7 ( $R_3$ ), or 7 ( $R_3$ ) curved upward near upper angle of discal cell, veins 6 ( $M_1$ ) and 7 ( $R_4$ ) usually stalked; abdomen without spines on terga; male genitalia with harpe simple to complex, bearing specialized setae which are frequently bifurcate apically.


As pointed out by Hodges (1966) many of the genera currently assigned to the Xyloryctidae ap-

pear to be improperly placed, and until the family is carefully studied it is impossible to provide an accurate analysis of family limits; however, in my opinion, the following characters provide the most satisfactory definition possible at this time: labial palpus upcurved; antenna simple, ciliated or pectinated in male, scape without pecten; maxillary palpus 3- or 4-segmented, usually partially encircling base of proboscis; forewing with veins 7 and 8 ( $R_4$  and  $R_5$ ) usually stalked, vein 2 ( $CuA_2$ ) arising well before lower angle of discal cell; hind wing with vein 8 ( $Sc + R_1$ ) separate and diverging from 7 ( $R_4$ ) well before upper angle of discal cell, veins 6 ( $M_1$ ) and 7 ( $R_4$ ) approximated, connate or stalked; abdomen with spines on posterior half of terga or along posterior margins; male genitalia with harpe usually simple, without specialized setae.


Diakonoff (1954) presented virtually the only discussion of the relationships between the Stenomidae and Xyloryctidae to be found in the modern literature. His conclusions were based primarily on the New Guinea fauna and thus do not necessarily define either family in relation to their total

range and composition. In fact, there is evidence which indicates that he did not always adhere to his family definitions for placement of the New Guinea genera. For example, the genus *Eriogenes*, revised and transferred to the Stenomidae in the present study, was included in the Xyloryctidae by Diakonoff (1954:127) with the following comment: "Meyrick regarded this genus to be related with *Agriophara* (Stenomidae) which is certainly erroneous. The insect is a true Xyloryctid, is in our opinion, allied to *Cryptophasa*, and does not show any connections with the Stenomidae." In fact, thorough study of the genus reveals that *Eriogenes* agrees in every respect with the characters Diakonoff lists as distinctive for the Stenomidae and, as discussed elsewhere in the present paper, demonstrates a closer relationship with the Neotropical stenomids than to any portion of the Old World fauna recognized to date.

As discussed in the introduction, the confusion concerning the families Xyloryctidae and Stenomidae has a long history and involves personalities as well as evaluation of characters. Personalities no longer intrude into the problem; however, the rela-



MAP 1.—Generalized distribution of Old World Stenomidae.




MAP 2.—Distribution of *Agriophara* species (type-localities only): 1, *A. asaphes* Diak.; 2, *A. atratella* (Wlk.); 3, *A. axesta* Meyr.; 4, *A. biornata* Diak.; 5, *A. bradleyi* Diak.; 6, *A. capnodes* Meyr.; 7, *A. cinderella* (Newman); 8, *A. cinerosa* Rosenstock; 9, *A. confertella* (Wlk.); 10, *A. cremnopsis* Lower; 11, *A. curia* Lucas; 12, *A. diminuta* Rosenstock; 13, *A. discoloba* Turner; 14, *A. dyscapna* Turner; 15, *A. fascifera* Meyr.; 16, *A. gravis* Meyr.; 17, *A. heterchroma* Diak.; 18, *A. horridula* Meyr.; 19, *A. hylinoia* Lower; 20, *A. leptosemela* Lower; 21, *A. leucanthes* Turner; 22, *A. leucosta* Turner; 23, *A. levis* Meyr.; 24, *A. lysimacha* Meyr.; 25, *A. musicolor* Meyr.; 26, *A. neoxantha* Meyr.; 27, *A. nephelopa* Diak.; 28, *A. nodigera* Turner; 29, *A. parallela* Diak.; 30, *A. parilis* Meyr.; 31, *A. phasmatopea* (Meyr.); 32, *A. plagiosema* Turner; 33, *A. platyscia* Lower; 34, *A. poliopepla* Turner; 35, *A. polistis* (Lower); 36, *A. salinaria* Meyr.; 37, *A. tephroptera* Lower; 38, *A. virescens* Diak.; 39, *A. coricopea* (Meyr.).

tive uncertainty over supergeneric classification, not only in these two families but throughout the Gelechioidea, continues to be the major obstacle to a better understanding of their interrelationships and distribution in the world fauna. Hodges (1966) and Powell (1973) have discussed this problem in relation to their studies of the gelechiids and ethmiids, respectively, and both emphasize the need for increased elucidation of suprageneric taxa within the Gelechioidea on a world basis.

### Geographical Distribution

A generalized diagram of the distribution of Old World stenomids is provided in Map 1. The patterns shown are based only on type-localities to avoid errors of misidentification in those groups not studied in detail. Maps 2–6 indicate the precise species localities for the various genera again based on type-localities only. Map 7 provides the total known distribution of the genus *Eriogenes* and is discussed in detail in the treatment of that genus.




MAP 3.—Distribution of *Phylomictis*, *Aproopta*, *Proscedes*, and *Nothochalara* species (type-localities only): 1, *Ph. arctans* Lucas; 2, *Ph. decretoria* Lucas; 3, *Ph. eclecta* Turner; 4, *Ph. idiotricha* Meyr.; 5, *Ph. leucopelta* Lower; 6, *Ph. linearia* Meyr.; 7, *Ph. maligna* Meyr.; 8, *Ph. monochroma* Lower; 9, *Ph. obliquata* Lucas; 10, *Ph. palaeomorpha* Turner; 11, *Ph. sarcinopa* Meyr.; 12, *A. melanchlaena* Turner; 13, *Pr. torquigera* Diak.; 14, *N. sordida* Diak.

Perhaps the most significant aspect of the distribution shown in Map 1 is the obvious absence of records from most of the Oriental Region. This reflects, of course, the serious lack of collecting of Microlepidoptera accomplished to date in this important area. The impact of our ignorance of the Oriental Region, as it relates to Microlepidoptera, is brought into focus when it is realized that there is increasing evidence, convincingly summarized and elaborated by Smith (1970), that the original center of distribution of flowering plants (angiosperms) was, in all probability, southeastern Asia and adjacent Malesia. Viewed in conjunction with the probability that the Order Lepidoptera owes its diversity and success to the development of the flowering plants, information concerning the Microlepidoptera from the Oriental Region be-

comes basic to an understanding of the origin and dispersal of the group. Thus, it seems reasonable to assume that until additional field sampling is achieved, any comments concerning relationships in the Old World Microlepidoptera are exceedingly tenuous.

With the previous statements in mind, it seems to me appropriate to examine the currently known distribution of the Old World stenomid fauna, albeit imperfect, and to consider possible explanations for the patterns observed.

When the various localities are examined in detail a definite tropical character to the distribution of Old World stenomids emerges. Also, most of the localities are in areas of tropical rain forest or tropical montane forest as defined by Eyre (1968) with the exception of southeastern Australia. This




MAP 4.—Distribution of *Synchalara* and *Neospastis* species (type-localities only).

- |                               |                                |
|-------------------------------|--------------------------------|
| ● <i>S. argoplaca</i> (Meyr.) | ● <i>S. rhombota</i> (Meyr.)   |
| ■ <i>S. byrsina</i> (Meyr.)   | ○ <i>N. calipidias</i> Meyr.   |
| ◆ <i>S. malacobryas</i> Meyr. | □ <i>N. encryphias</i> (Meyr.) |
| ★ <i>S. minax</i> (Meyr.)     | ☆ <i>N. ichnaea</i> (Meyr.)    |

pattern coincides with observations on the Neotropical stenomid fauna (Duckworth, 1969, 1971) and, in my opinion, it is in this type of habitat that stenomids have flourished and achieved their initial diversification in both the Old and New World. Secondary penetration of dryer habitats has occurred but appears to be less important to the overall family dispersal. The best example of this pattern can be seen in the genera from Madagascar (Maps 5 and 6) where only one species is described from the western savanna area, the remainder occurring in the tropical forests which extend along the eastern portion of the island.

A great deal more study at the generic level must be accomplished before additional interpretation of Old World stenomid distributions can be made. The one genus which has been examined in detail during the present study, *Eriogenes*, does indicate a distinct relationship between the Old and New


World elements of the family and, furthermore, appears to possess characteristics reflective of an ancestral condition. If this is found to be the case for other Old World genera, especially in the Southeast Asia-Malesia area, it would seem likely that the generalized model of angiosperm distribution proposed by Smith (1970) would hold true for stenomids as well. In the case of stenomid entry into the New World, data at this point would appear to favor Smith's southern route along eastern Australia and the New Caledonian-New Zealand insular chain into the West Antarctic Archipelago and, utilizing island stepping stones, into South America. There is no evidence, at this point, which suggests a route crossing Beringia to the north; the Nearctic stenomid fauna appears to have originated in the Neotropical Region and is only marginally successful in temperate habitats (Duckworth, 1971).



MAP 5.—Distribution of *Herbulotiana* species (type-localities only): 1, *H. abceda* Viette; 2, *H. benoistella* Viette; 3, *H. bernardiella* Viette; 4, *H. bicolorata* Viette; 5, *H. catalaella* Viette; 6, *H. collectella* Viette; 7, *H. halarcta* (Meyr.); 8, *H. longifascia* Viette; 9, *H. paulianella* Viette; 10, *H. rungSELLA* Viette; 11, *H. vadonella* Viette; 12, *H. violacea* Viette; 13, *H. atypicella* Viette; 14, *H. robustella* Viette; 15, *H. altitudinella* Viette; 16, *H. septella* Viette.

#### Revision of the Genus *Eriogenes* Meyrick

The genus *Eriogenes* was established by Meyrick (1925) with a single included species, *E. mesogypsa*, with the indication that it was apparently closely related to *Agriophara*. Diakonoff (1954) included *Eriogenes* in the family Xyloryctidae in his study of the New Guinea Microlepidoptera, noting that Meyrick's placement of the genus as a



MAP 6.—Distribution of *Amontes*, *Mocquerysiella*, and *Mnarolitia* species (type-localities only).

- *A. princeps* Viette
- *Mo. albicosta* Viette
- *Mo. bourginella* Viette
- *Mn. paulianellum* Viette
- ★ *Mn. sylvestrella* Viette

relative of *Agriophara* (Stenomidae) was in error. Contrary to that opinion, in the present study it has been determined that *Eriogenes* is, in fact, more appropriately placed in the Stenomidae and shows no relationship to the xyloryctids in any character other than gross superficial appearance. In addition, detailed study of the morphological characters within the genus, particularly the genitalia, suggest a relationship to certain genera in the Neotropical portion of the family.

## MORPHOLOGY

Superficially, the members of this genus are large, robust, densely scaled moths which resemble certain noctuids, or some of the larger xyloryctids such as *Cryptophasa*, in general appearance. Although comprised of only two species, the genus exhibits a number of morphological characters worthy of special emphasis.

**HEAD.**—In both species the eye index is 1.0 or greater, ranging from 1.0 to 1.3, suggesting that these moths are nocturnal in adult activity patterns. For additional discussion of the eye index and its significance refer to Duckworth (1971) and Powell (1973). The maxillary palpi are 4-segmented. The labial palpi are 3-segmented; the apical segment is reduced and variable in size and shape between the two species. The proboscis is reduced and hidden between the basal segments of the labial palpi when in the normal coiled position.

**THORAX.**—The thorax is very robust and clothed laterally and ventrally with long hair scales. The wing venation is homogeneous and no characters of specific significance were observed. The occurrence of a cross vein between the upper edge of the cell and vein 8 ( $Sc + R_1$ ) in the hind wing, mentioned by Diakonoff (1954), is variable and not a dependable character at either the generic or specific level.

**ABDOMEN.**—No structures of generic significance were observed on abdominal segments 1–8. Perhaps the most notable aspect of this area of the body is the absence of secondary sexual structures, with the exception of a small eversible ventral pouch covered with a small mass of erectile hair scales in the intersegmental membrane between segments 7 and 8 in *E. meyricki*.

**MALE GENITALIA.**—By far, the most significant and intriguing structural characters in this genus are found in the male genitalia. The harpe consists of a simple lobe rather uniformly clothed over the apical half with specialized setae. These specialized setae are broadened and multilobed apically, the number of lobes varying but apparently always more than two. The uncus is large, broadly articulated with the tegumen basally, and somewhat recurved near the midpoint. The gnathos is pendulous and produced into a heavily sclerotized

plate apically. The anellus consists of a simple plate with two lateral lobes. Both the shape of the gnathal plate and the size and shape of the lateral lobes of the anellus vary between the two species and serve to distinguish them specifically. The aedeagus is produced apically, and the vesica is armed with cornuti in *E. meyricki* and unarmed in *E. eriogenes*.


**FEMALE GENITALIA.**—No female specimens are known for the species in this genus.

## GEOGRAPHICAL DISTRIBUTION

This genus is presently known to occur only in New Guinea and adjacent islands, including Ceram in the Moluccas (Map 7). The lack of records for the two species of *Eriogenes* seriously restricts any effort to analyze distribution patterns or to project ranges in any definitive fashion. The absence of biological data, especially host plant information, further complicates matters and renders all observations made at this point tentative.

The most interesting aspect of the distribution patterns for the two species of *Eriogenes* is the divergence from mainland New Guinea into different adjacent island groups. *E. meyricki* is recorded from Ceram in the southern Moluccas, the Central Range on mainland New Guinea, and throughout the Bismark Archipelago. On the other hand, *E. mesogypsa* extends from the Idenburg River in West Irian, southeastward into the D'Entrecasteau and Louisiade island groups. Similar patterns have been noted for butterflies (Zeuner, 1943; Carpenter, 1953) and they appear to correlate with the geological history of New Guinea (Toxopeus, 1950; Holloway and Jardine, 1968).

In the Miocene the northern ranges of New Guinea were a part of a "Melanesian arc" of islands which included the North Moluccas and Talaud Islands to the west and the Bismarck and Solomon Islands to the east. Farther south, land which now constitutes the central range of New Guinea appeared in late Miocene, well isolated from the Melanesian arc. During the Pliocene-Pleistocene further uplifting occurred in the central range and gradually the central and northern ranges were joined. Other portions of New Guinea (e.g., Vogelkop) undoubtedly existed as separate islands during the Pleistocene. Thus, the butterfly fauna, as

MAP 7.—Distribution of *Eriogenes* species.■ *E. mesogypsa* Meyr● *E. meyricki* Duckworth

analyzed by Holloway and Jardine (1968) reflects three faunal elements; one-third endemic, one-third with Sunda-based affinities, and one-third Melanesian. They further conclude that this heterogeneity results from the Melanesian and endemic elements having risen separately through dispersal from Asia into the Melanesian arc and into the formerly separate southern part of New Guinea since the Miocene. Although insufficient data are available to confirm or reject a similar pattern in *Eriogenes*, it is interesting to note that *E. meyricki* displays a "Melanesian" pattern and *E. mesogypsa* appears to be more "endemic" in distribution.

The currently known distribution of the genus also suggests the possibility that species of *Eriogenes* may occur farther west into Indonesia. This area is virtually unknown insofar as Microlepidoptera are concerned (Diakonoff, 1954) and undoubtedly served as a major point of dispersal to the New Guinea region.

Available data suggest the possibility of a significant altitudinal difference for the two species

of *Eriogenes*. *E. mesogypsa* is much the better documented species in this regard and appears to occur at lower elevations up to approximately 2500 feet. One record, from Goodenough Island, indicates a range in elevation from 2500 feet to 4000 feet; however, the consistency of other records would suggest that this specimen was probably captured at the lower limits of this range. Also, there is a specimen from the Jimmi River, in the Western Highlands, with an altitude record of 4700 feet. This appears to be an error in that the Jimmi River is situated in a low valley at an elevation generally less than 2500 feet. At one point, according to Gressitt (1956), the river elevation is 460 meters (1508 ft). *E. meyricki*, on the other hand, is recorded from Ceram at an elevation of 4600 feet, Telefomin in central New Guinea at 4500 to 5500 feet, and the other specimens from Rook, St. Matthias and New Britain Islands are without elevation data. Rook Island has elevations above 2500 feet and thus the potential for correlation; however, St. Matthias is a much lower

island and would not appear to fit the concept. Yet Hartert (1924), in a paper on the birds of St. Matthias, indicates the island consists of a relatively high plateau (2130 ft) partly covered with thick forest which might be suitable habitat for species that occur at greater altitudes on the higher islands.

#### CLASSIFICATION AND RELATIONSHIPS

Although a great deal of the Old World stenomid fauna remains to be studied in detail, and undoubtedly is yet to be discovered, the genus *Eriogenes* is of considerable significance in that it appears to be more closely related morphologically to the Neotropical genus *Chlamydastis* and its allies than to any Old World genus. As pointed out in a previous paper (Duckworth, 1971), *Chlamydastis* and several related genera seem to constitute a major division within the family, characterized, in part, by their possession of multilobed, palmate-type specialized setae on the harpe in the male genitalia. *Eriogenes* has a less specialized type of multilobed setae on the harpe, which suggests the possibility that it represents a more primitive stage in the development of this character. In addition, virtually every structure of the male genitalia demonstrates both a relationship with *Chlamydastis* and a more generalized character state than is found in that genus. For example, in *Chlamydastis* the specialized setae are restricted to the outer costal margin of the harpe, there is a tendency for reduction in the length of the uncus, the harpe shows a tendency to divide into a costal lobe and saccular lobe, and the gnathos may be long and pendulous or entirely absent. In *Eriogenes* the specialized setae are uniformly distributed over the apical half of the harpe, the uncus is very long,

the harpe is a large simple lobe, and the gnathos is long and pendulous. Thus, in each of these structures, a clear relationship exists between the genera in terms of similarities of structure, with *Eriogenes* possessing less reduction and modification than *Chlamydastis*. After consideration of these and other characters (e.g., maculation and venation), *Eriogenes*, in my opinion, represents a distinct link between the Old and New World stenomid faunas and possibly displays characteristics reflective of an ancestral condition.

#### Genus *Eriogenes* Meyrick

*Eriogenes* Meyrick, 1925:159. [Type-species: *Eriogenes mesogyrsa* Meyrick by monotypy.]

Head densely scaled, lateral tufts raised. Male antenna heavily ciliated ventrally, slightly fasciculate. Labial palpus recurved, not reaching vertex, thickened with heavy scaling. Proboscis reduced, concealed between labial palpi; maxillary palpus 4-segmented. Thorax and abdomen densely covered with long hair scales. Legs with all tibia and tarsi expanded with hair scales. Forewing with costa slightly arched, apex rounded, termen oblique, tornus rounded; with 12 veins, all separate, discal cell extending beyond midpoint, 2 ( $CuA_2$ ) from end of cell, 7 ( $R_5$ ) to termen. Hind wing broader than forewing; with 8 veins, 3 and 4 ( $CuA_1$  and  $M_3$ ) connate, 6 and 7 ( $M_1$  and  $R_3$ ) connate, upper edge of cell curved upward toward vein 8 ( $Sc + R_1$ ) at apical four-fifths.

MALE GENITALIA.—Uncus simple, curved slightly ventrad from middle; gnathos pendulous, fused apically forming gnathal plate; harpe simple, apical half clothed with apically multilobed specialized setae; anellus with lateral lobes; vinculum fused;


#### Key to the Species of *Eriogenes*

##### BASED ON EXTERNAL FEATURES

- Apical segment of labial palpus one-third to one-half the length of second segment; forewing with discal spot on outer margin of median white fascia.....*E. mesogyrsa* Meyrick  
 Apical segment of labial palpus less than one-third the length of second segment; discal spot within median white fascia.....*E. meyricki*, new species

##### BASED ON THE MALE GENITALIA

- Anellus with lateral lobes long, falciform; vesica of aedeagus without cornuti.....  
 .....*E. mesogyrsa* Meyrick  
 Anellus with lateral lobes short, truncate; vesica of aedeagus with cornuti .....  
 .....*E. meyricki*, new species



FIGURES 1-4.—Head, legs, genitalia morphology: 1, caudolateral view of head of *E. mesogypsa*; 2, legs of *E. mesogypsa*; 3, male genitalia of *E. mesogypsa* (left harpe removed); 4, male genitalia of *E. meyricki* (left harpe removed).

aedeagus produced apically, vesica of aedeagus with or without cornuti.

FEMALE GENITALIA.—Unknown.

DISCUSSION.—This genus is readily separated from other Old World stenomid genera by the reduced proboscis, distinctive maculation of the forewings, robust thorax and abdomen covered with long hair scales, and distinctive male genitalia. In the latter, the long uncus, pendulous gnathos, and simple harpe with multilobed specialized setae are characteristic.

No female specimens of species in this genus are known and, consequently, it is impossible to provide distinguishing characteristics based on structures of the female genitalia.

### *Eriogenes mesogypsa* Meyrick

FIGURES 1-3, 5-8; MAP 7

*Eriogenes mesogypsa* Meyrick, 1925:159.

Alar expanse 30-35 mm.

Antenna with scape light brown suffused with white, shaft light brown dorsally. Head dark brown medially, whitish tufts posterior to antennae, face brown suffused with white; labial palpus dark brown, apical segment and apex of second segment sprinkled with white, apical segment one-third to one-half the length of second segment. Thorax and tegulae brown suffused with white dorsally. Legs white shaded with light brown; foreleg with coxa overcast with pale brown; femur, tibia, tarsi overcast with dark brown; midleg with tarsi brown; hind leg with tarsi white. Abdomen light brown to white. Forewing brown, darker basally, a median, triangulate, white fascia from costa to dorsum, discal spot dark brown on outer margin of white fascia, apical third brown along costa suffused with white toward tornus, a faint, outwardly curved brown line from costal three-fourths to tornus, cilia with interrupted brown line basally, white beyond. Hind wing light brown to white, cilia light brown basally, white beyond.

MALE GENITALIA (WDD 4199, lectotype).—Uncus rounded apically; gnathal plate sharply narrowing before apex, apex narrowly recurved; harpe broadly rounded apically; anellar plate with ventral margin straight, two upright, falciform lateral lobes; aedeagus long, narrow, more than one-half

the length of harpe, apex produced into a broad, knifelike flange; vesica without cornuti.

FEMALE GENITALIA.—Unknown.

TYPE.—Lectotype male, British Museum (Natural History).

TYPE-LOCALITY.—Kumusi River, Northeast British New Guinea at low elevation.



HOST PLANT.—Unknown.

DISTRIBUTION. — Presently known from New Guinea and adjacent islands at low elevations.

ADULT RECORDS. — NEW GUINEA: Kumusi River, low elevation; Bernhard Camp, Idenburg River, 50 m (Jan.); Hydrographer Mountains, 2500 ft (Jan., Feb.); Western Highlands, Jimi River, 4700 ft (July, Sept.). D'ENTRECASTEAU ISLANDS: Goodenough Island, 2500-4000 ft (April, May, June). MURUA (Woodlark Island); Kulamadau (Jan., May). LOUISIADE ARCHIPELAGO: Sudest Island (Tagula), Mount Riu, 2000 ft (Jan., Feb., March, April); Rossel Island, Mount Rossel, 2100 ft (Nov., Dec.); Rossel Island, Abaleti (Nov., Dec.); Misima Island, Umana Camp, 500 ft (June, July).

DISCUSSION.—Meyrick originally described this species from four specimens and noted that one of the four, from Ceram, was somewhat atypical. During the course of this study it was determined that the specimen from Ceram represented a separate species which, although very similar superficially, may be readily distinguished by both genitalia characters and maculation. In addition to the original three specimens, in excess of twenty additional specimens of *E. mesogypsa* from various localities have been studied.

For the most part, the material studied was collected many years ago and, consequently, it is impossible to determine the effect of time on the scale colors. Meyrick (1925) indicates a copper sheen in his original description which is not discernible in the specimens now, and it should be noted that the original series was collected at least ten years before Meyrick's description. In addition, most of the specimens, including the original series, are mouldy and somewhat greased. Thus, the colors are undoubtedly much duller than would be found in fresh specimens. I suspect the brown wing colors are more red-brown than they appear in the specimens now; however, additional information on this subject must await procurement of fresh material.



FIGURES 5-9.—Head morphology, wing venation, left wings: 5, head of *E. mesogyrsa*, frontal view, scales removed (labial palpus of *E. Meyricki* below); 6, maxilla of *E. mesogyrsa*; 7, wing venation of *E. mesogyrsa*; 8, left wings of *E. mesogyrsa*; 9, left wings of *E. meyricki*.

*Eriogenes meyricki*, new species

FIGURES 4, 9; MAP 7

Alar expanse 27–35mm.

Antenna with scape and dorsum of shaft light brown mixed with white. Head light brown medially, whitish tufts posterior to antennae, face light brown; labial palpus dark brown basally to middle of second segment, lighter brown mixed with white to apex, apical segment less than one-third length of second segment. Thorax and tegulae brown, heavily suffused with white. Legs as in *E. mesogypsa*. Abdomen pale brown to white ventrally, brown dorsally. Forewing brown, a median, irregularly shaped, white fascia from costa to dorsum, discal spot dark brown within the outer margin of white fascia, apical third with two relatively distinct transverse brown lines, one bordering the outer margin of white fascia, the other extending from apical three-fourths of costa to dorsum before tornus, cilia white with interrupted brown line basally. Hind wing brown, cilia brown.

MALE GENITALIA (WDD 4200, holotype).—Uncus truncate apically; gnathal plate gradually narrowing to broadly recurved apex; harpe acute apically; anellar plate with ventral margin indented at midpoint, two short, truncate lateral lobes; aedeagus short, broad, less than one-half the length of harpe, apex produced into a narrow, apically acute projection; vesica with two large, heavily sclerotized cornuti.

FEMALE GENITALIA.—Unknown.

TYPE.—Holotype male, British Museum (Natural History).

TYPE-LOCALITY.—Central Ceram, 4600 ft.

HOST PLANT.—Unknown.

DISTRIBUTION.—Presently known from the Moluccas, New Guinea, and the Bismarck Archipelago.

ADULT RECORDS.—MOLUCCAS: Ceram, 4600 ft (Jan.). NEW GUINEA: Telefomin (Eliptamin), 4500–5500 ft (June, Sept.). BISMARCK ARCHIPELAGO: Rook Island (July); New Britain, Keravat (no date). St. MATTHIAS GROUP: St. Matthias Island (July).

DISCUSSION.—Described from the male holotype: "Central Ceram, 4600 ft., Jan. '20, C. F. & J. Pratt"; two male paratypes: "St. Matthias Id., July, 1923, A. F. Eichhorn"; three male paratypes: "Rook Id., July, 1913, A. S. Meek, Joicey Bequest, Brit. Mus., 1934–120"; and one male paratype: "Bismarck Archipel., Rook Island, Meek, 6–7–13, M 768, Paravicini Coll., B. M. 1937–383." Two of the above paratypes deposited in the National Museum of Natural History, Smithsonian Institution, the remainder in the British Museum (Natural History). Two male paratypes: "New Guinea, Telefomin (Eliptamin), 4500–5500 ft., 19 June–14 Sept. 1959, W. W. Brandt, Genitalia Slide G617"; one male paratype: "NEW BRITAIN, Keravat, collected by Wm. Brandt, Sir Edward Hallstrom, Genitalia Slide G1616"; all deposited in the Australian National Insect Collection, C. S. I. R. O., Canberra, Australia.

This species seems very closely related to *E. mesogypsa* and may be separated by the characters indicated under the discussion pertaining to that species. In general, it seems *E. meyricki* is on the average smaller than *E. mesogypsa*, occurs at higher elevations, and is darker in maculation.

## Catalog of the Old World Stenomidae

Included in the following catalog are all those genera and species which have been assigned to the Stenomidae by previous workers or determined more appropriately placed with the family during the present study. The information provided concerning location of types has been gathered from various sources and where questionable the data is preceded by a question mark. The genera are arranged alphabetically and the included species are also listed alphabetically under the generic headings. Type-locality data are essentially as found in the original description, with an occasional addition (state, district, province) for clarity.

As mentioned in the introduction, no finality is implied for this catalog. Many of the species included have not been studied and the genera are, in many cases, poorly defined. Thus, the catalog is intended as a point of departure for future studies and a summation of the family's current composition in the Old World.

Genus *Agriophara* Rosenstock

MAP 2

*Agriophara* Rosenstock, 1885:39. [Type-species: *Agriophara cinerosa* Rosenstock by subsequent designation, Fletcher, 1929:9.]

*Hypeuryntis* Meyrick, 1897:389. [Type-species: *Hypeuryntis coricopa* Meyrick by monotypy.]

1. *A. asaphes* Diakonoff [*Agriophara*], 1948:193.  
Type-locality: Paniai Lake, New Guinea.  
Type: holotype female, Rijksmuseum van Natuurlijke Historie, Leiden.
2. *A. atratella* (Walker) [*Acrobasis*], 1866:1712.  
Type-locality: Moreton Bay, Queensland, Australia.  
Type: holotype male, British Museum (Natural History).
3. *A. axesta* Meyrick [*Agriophara*], 1890:79.  
Type-locality: Wirrabara, South Australia, Australia.  
Type: syntypes of both sexes, British Museum (Natural History).
4. *A. biornata* Diakonoff [*Agriophara*], 1954:147.  
Type-locality: Iebele Camp, 2250 m, West Irian, New Guinea.  
Type: holotype male, Rijksmuseum van Natuurlijke Historie, Leiden.
5. *A. bradleyi* Diakonoff [*Agriophara*], 1954:146.  
Type-locality: Mist Camp, 1800 m, West Irian, New Guinea.  
Type: holotype male, Rijksmuseum van Natuurlijke Historie, Leiden.
6. *A. capnodes* Meyrick [*Agriophara*], 1890:77.  
Type-locality: Mount Lofty, South Australia, Australia.  
Type: syntypes of both sexes, British Museum (Natural History).
7. *A. cinderella* (Newman) [*Chimabacche*], 1855:288.  
Type-locality: Mt. Alexander Range, Victoria, Australia.  
Type: holotype ?British Museum (Natural History).
8. *A. cinerosa* Rosenstock [*Agriophara*], 1885:439.  
Type-locality: South Australia.  
Type: holotype male, British Museum (Natural History).
9. *A. confertella* (Walker) [*Cryptolechia*], 1864:758.  
Type-locality: Moreton Bay, Queensland; Sydney, New South Wales, Australia.  
Type: syntypes of both sexes, British Museum (Natural History).
10. *A. coricopa* (Meyrick) [*Hypeuryntis*], 1897:389.  
Type-locality: Wellington, New Zealand.  
Type: holotype male, British Museum (Natural History).
11. *A. cremnopsis* Lower [*Agriophara*], 1894:93.  
Type-locality: Duaringa, Queensland, Australia.  
Type: holotype male, ?South Australia Museum, Adelaide.
12. *A. curta* Lucas [*Agriophara*], 1900:161.  
Type-locality: Brisbane, Queensland, Australia.  
Type: holotype male, South Australia Museum, Adelaide.
13. *A. diminuta* Rosenstock [*Agriophara*], 1885:440.  
Type-locality: South Australia.  
Type: holotype male, British Museum (Natural History).
14. *A. discobola* Turner [*Agriophara*], 1897:32.  
Type-locality: Gisborne, Victoria, Australia.  
Type: syntypes of both sexes, ?Australian National Insect Collection, Canberra.
15. *A. dyscapna* Turner [*Agriophara*], 1917:99.  
Type-locality: Gisborne, Victoria, Australia.  
Type: holotype male, ?Australian National Insect Collection, Canberra.
16. *A. fascifera* Meyrick [*Agriophara*], 1890:80.  
Type-locality: Sydney and Bathurst, 2500 ft, New South Wales, Australia.  
Type: syntypes of both sexes, British Museum (Natural History).
17. *A. gravis* Meyrick [*Agriophara*], 1890:77.  
Type-locality: Sydney, New South Wales, Australia; Deloraine, Tasmania.  
Type: syntypes of both sexes, British Museum (Natural History).
18. *A. heterchroma* Diakonoff [*Agriophara*], 1954:144.  
Type-locality: Iebele Camp, 2250 m, West Irian, New Guinea  
Type: holotype female, Rijksmuseum van Natuurlijke Historie, Leiden.
19. *A. horridula* Meyrick [*Agriophara*], 1890:77.  
Type-locality: Sydney, New South Wales, Australia.  
Type: holotype male, British Museum (Natural History).
20. *A. hyalinota* Lower [*Agriophara*], 1899:104.  
Type-locality: Parkside, South Australia; Duaringa, Queensland, Australia.

- Type: syntypes of both sexes, South Australia Museum, Adelaide.
21. *A. leptosemela* Lower [*Agriophara*], 1893:174.  
Type-locality: Parkside, South Australia, Australia.  
Type: holotype male (without abdomen), South Australia Museum, Adelaide.
  22. *A. leucanthes* Turner [*Agriophara*], 1897:31.  
Type-locality: Brisbane, Queensland, Australia.  
Type: syntype males, Australian National Insect Collection, Canberra.
  23. *A. leucosta* Lower [*Agriophara*], 1893:173.  
Type-locality: Parkside, South Australia, Australia.  
Type: syntypes of both sexes, ?South Australia Museum, Adelaide.
  24. *A. levis* Meyrick [*Agriophara*], 1921:446.  
Type-locality: Brisbane, Queensland, Australia.  
Type: holotype, British Museum (Natural History).
  25. *A. lysimacha* Meyrick [*Agriophara*], 1915:410.  
Type-locality: Woodlark Island, New Guinea.  
Type: holotype male, British Museum (Natural History).
  26. *A. murinella* (Walker) [*Cryptolechia*], 1860:757.  
Type-locality: Australia.  
Type: holotype female (without abdomen), British Museum (Natural History).
  27. *A. musicolor* Meyrick [*Agriophara*], 1930:560.  
Type-locality: Biagi, Mambare River, 5000 ft, New Guinea.  
Type: holotype male, ?Deutsches Entomologisches Institut, Berlin.
  28. *A. neoxanta* Meyrick [*Agriophara*], 1930:410.  
Type-locality: Woodlark Island, New Guinea; Cooktown, Queensland, Australia.  
Type: syntypes of both sexes, British Museum (Natural History).
  29. *A. nephelopa* Diakonoff [*Agriophara*], 1954:148.  
Type-locality: Moss Forest Camp, 2600–2800 m, West Irian, New Guinea.  
Type: male holotype, Rijksmuseum van Natuurlijke Historie, Leiden.
  30. *A. nodigera* Turner [*Agriophara*], 1900:11.  
Type-locality: Warwick, Queensland, Australia.
  - Type: holotype female, Australian National Insect Collection, Canberra.
  31. *A. parallela* Diakonoff [*Agriophara*], 1954:142.  
Type-locality: Lower Mist Camp, 1400–1600 m, West Irian, New Guinea.  
Type: holotype female, Rijksmuseum van Natuurlijke Historie, Leiden.
  32. *A. parilis* Meyrick [*Agriophara*], 1918:197.  
Type-locality: Setekwa River, 2000–3000 ft., New Guinea.  
Type: holotype male, British Museum (Natural History).
  33. *A. phasmatopa* (Meyrick) [*Stenoma*], 1910:460.  
Type-locality: Gizo, New Georgia, Solomon Islands.  
Type: syntype males, British Museum (Natural History).
  34. *A. plagiosema* Turner [*Agriophara*], 1897:32.  
Type-locality: Brisbane, Queensland, Australia.  
Type: holotype male, Australian National Insect Collection, Canberra.
  35. *A. platyscia* Lower [*Agriophara*], 1908:117.  
Type-locality: Tasmania.  
Type: holotype female (without abdomen), South Australia Museum, Adelaide.
  36. *A. poliopepla* Turner [*Agriophara*], 1897:31.  
Type-locality: Brisbane, Queensland, Australia.  
Type: syntypes of both sexes, Australian National Insect Collection, Canberra.
  37. *A. polistis* (Lower) [*Agriophara*], 1923:55.  
Type-locality: Dorrigo, New South Wales, Australia.  
Type: syntype males, South Australia Museum, Adelaide.
  38. *A. salinaria* Meyrick [*Agriophara*], 1931:43.  
Type-locality: Florida, Solomon Islands.  
Type: holotype male, ?British Museum (Natural History).
  39. *A. tephroptera* Lower [*Agriophara*], 1903:230.  
Type-locality: Broken Hill, New South Wales, Australia.  
Type: holotype male, South Australia Museum, Adelaide.
  40. *A. virescens* Diakonoff [*Agriophara*], 1954:143.  
Type-locality: Cyclops Mountains, 400–900 m, West Irian, New Guinea.  
Type: holotype male, Rijksmuseum van Natuurlijke Historie, Leiden.

**Genus *Amontes* Viette**

## MAP 6

*Amontes* Viette, 1958:116. [Type-species: *Amontes princeps* Viette by original designation.]

1. *A. princeps* Viette [*Amontes*], 1958:116.  
Type-locality: env. de Perinet, forêt d'Analamaoatra, 910 m, Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.

**Genus *Aproopta* Turner**

## MAP 3

*Aproopta* Turner, 1919:171. [Type-species: *Aproopta melanclaena* Turner by monotypy.]

1. *A. melanclaena* Turner [*Aproopta*], 1919:172.  
Type-locality: Katoomba, New South Wales, Australia.  
Type: holotype male, Australian Museum, Sydney.

**Genus *Eriogenes* Meyrick**

## MAP 7

*Eriogenes* Meyrick, 1925:159. [Type-species: *Eriogenes mesogypsa* Meyrick by monotypy.]

1. *E. mesogypsa* Meyrick [*Eriogenes*], 1925:159.  
Type-locality: Kumusi River, Northeast British New Guinea, at low elevation.  
Type: lectotype male, British Museum (Natural History).
2. *E. meyricki* Duckworth [*Eriogenes*], 1973:14.  
Type-locality: Central Ceram, 4600 ft, Moluccas.  
Type: holotype male, British Museum (Natural History).

**Genus *Herbulotiana* Viette**

## MAP 5

*Herbulotiana* Viette, 1954:17. [Type-species: *Herbulotiana abceda* Viette by original designation.]

1. *H. abceda* Viette [*Herbulotiana*], 1954:23.  
Type-locality: baie d'Antongil, Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
2. *H. altitudinella* Viette [*Herbulotiana*], 1962:24.

Type-locality: Plateau Soaindrana, 2070 m, Massif de l'Andringitra, Central Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.

3. *H. atypicella* Viette [*Herbulotiana*], 1956a:115.  
Type-locality: route d'Anosibe, km 57, Madagascar Est.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
4. *H. benoistella* Viette [*Herbulotiana*], 1954:18.  
Type-locality: baie d'Antongil, Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
5. *H. bernardiella* Viette [*Herbulotiana*], 1954:26.  
Type-locality: env. de Maroantsetra, forêt d'Ambodivoangy, Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
6. *H. bicolorata* Viette [*Herbulotiana*], 1954:20.  
Type-locality: baie d'Antongil, Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
7. *H. catalaella* Viette [*Herbulotiana*], 1954:22.  
Type-locality: env. de Maroantsetra, forêt d'Ambodivoangy, Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
8. *H. collectella* Viette [*Herbulotiana*], 1956a:111.  
Type-locality: env. de Maroantsetra, forêt d'Ambodivoangy, Nord-Est Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
9. *H. halarcta* (Meyrick) [*Agriophara*], 1917:59.  
Type-locality: Antananarivo, Madagascar.  
Type: holotype male, British Museum (Natural History).
10. *H. longifascia* Viette [*Herbulotiana*], 1954:26.  
Type-locality: baie d'Antongil, Madagascar.  
Type: holotype female, Museum National D'Histoire Naturelle, Paris.
11. *H. paulianella* Viette [*Herbulotiana*], 1954:24.  
Type-locality: env. de Maroantsetra, forêt d'Ambodivoangy, Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
12. *H. robustella* Viette [*Herbulotiana*], 1956a:113.  
Type-locality: env. de Perinet, forêt d'Analamaoatra, 910 m, Madagascar Est.

- Type: holotype male, Museum National D'Histoire Naturelle, Paris.
13. *H. rungsella* Viette [*Herbulotiana*], 1954:19.  
Type-locality: env. de Maroantsetra, forêt d'Ambodivoangy, Madagascar.  
Type: holotype female, Museum National D'Histoire Naturelle, Paris.
  14. *H. septella* Viette [*Herbulotiana*], 1956a:112.  
Type-locality: env. de Ranomafana, district d'Ifanadiana, 700 m, Madagascar Est.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
  15. *H. vadonella* Viette [*Herbulotiana*], 1956b:466.  
Type-locality: env. de Maroantsetra, forêt d'Ambodivoangy, Nord-Est Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
  16. *H. violacea* Viette [*Herbulotiana*], 1954:21.  
Type-locality: baie d'Antongil, Madagascar  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.

#### Genus *Mnarolitia* Viette

##### MAP 6

*Mnarolitia* Viette, 1954:16. [Type-species: *Mnarolitia paulianellum* Viette by original designation.]

1. *M. paulianellum* Viette [*Mnarolitia*], 1954:16.  
Type-locality: env. de Morafenobe, forêt de Mahajeby, Ouest Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.
2. *M. sylvestrella* Viette [*Mnarolitia*], 1968:84.  
Type locality: env. de Perinet, forêt d'Analamaoatra, 910 m, Madagascar Est.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.

#### Genus *Mocquerysiella* Viette

##### MAP 6

*Mocquerysiella* Viette, 1954:13. [Type-species: *Mocquerysiella alibcosta* Viette by original designation.]

1. *M. alibcosta* Viette [*Mocquerysiella*], 1954:14.  
Type-locality: baie d'Antongil, Nord-Est Madagascar.

Type: holotype male, Museum National D'Histoire Naturelle, Paris.

2. *M. bourginella* Viette [*Mocquerysiella*] 1954:15.  
Type-locality: env. de Maroantsetra, forêt d'Ambodivoangy, Nord-Est Madagascar.  
Type: holotype male, Museum National D'Histoire Naturelle, Paris.

#### Genus *Neospastis* Meyrick

##### MAP 4

*Neospastis* Meyrick, 1917:59. [Type-species: *Agriophara encryphias* Meyrick by original designation.]

1. *N. calpidias* Meyrick [*Neospastis*], 1917:59.  
Type-locality: Nilgiris, Pykara, 7000 ft, India.  
Type: lectotype male, British Museum (Natural History).
2. *N. encryphias* (Meyrick) [*Agriophara*], 1907a:743.  
Type-locality: Khasi Hills, Assam, India.  
Type: lectotype male, British Museum (Natural History).
3. *N. ichnaea* (Meyrick) [*Stenoma*], 1914:118.  
Type-locality: Anshi, Kanara, India.  
Type: lectotype male, British Museum (Natural History).

#### Genus *Nothochalara* Diakonoff

##### MAP 3

*Nothochalara* Diakonoff, 1954:135. [Type-species: *Nothochalara sordida* Diakonoff by original designation.]

1. *N. sordida* Diakonoff [*Nothochalara*], 1954:136.  
Type-locality: Araucaria Camp, 800 m, West Irian, New Guinea.  
Type: holotype female, Rijksmuseum van Natuurlijke Historie, Leiden.

#### Genus *Phylomictis* Meyrick

##### MAP 3

*Phylomictis* Meyrick, 1890:74. [Type-species: *Phylomictis maligna* Meyrick by monotypy.]

*Comoscotopa* Lower, 1902:239. [Type-species: *Comoscotopa leucopelta* Lower by monotypy.]

1. *P. arctans* Lucas [*Phylomictis*], 1900:159.

- Type-locality: Brisbane, Queensland, Australia.
- Type: syntypes of both sexes, ?South Australia Museum, Adelaide.
2. *P. decretoria* Lucas [*Phylomictis*], 1900:160.  
Type-locality: Brisbane, Queensland, Australia.  
Type: holotype female, ?South Australia Museum, Adelaide.
  3. *P. eclecta* Turner [*Phylomictis*], 1906:142.  
Type-locality: Burpengary, near Brisbane, Queensland, Australia.  
Type: holotype male, Australian National Insect Collection, Canberra.
  4. *P. idiotricha* Meyrick [*Phylomictis*], 1921:446.  
Type-locality: Brisbane, Queensland, Australia.  
Type: holotype male, British Museum (Natural History).
  5. *P. leucopelta* (Lower) [*Comoscotopa*], 1902:240.  
Type-locality: Mount Gambier, South Australia, Australia.  
Type: holotype male, ?South Australia Museum, Adelaide.
  6. *P. lintearia* Meyrick [*Phylomictis*], 1921:445.  
Type-locality: Brisbane, Queensland, Australia.  
Type: holotype male, British Museum (Natural History).
  7. *P. maligna* Meyrick [*Phylomictis*], 1890:75.  
Type-locality: Melbourne, Victoria, Australia.  
Type: holotype male, ?British Museum (Natural History).
  8. *P. monochroma* Lower [*Phylomictis*], 1892:17.  
Type-locality: Parkside, South Australia, Australia.  
Type: holotype female (without abdomen), South Australia Museum, Adelaide.
  9. *P. obliquata* Lucas [*Phylomictis*], 1900:160.  
Type-locality: May Orchard, Brisbane, Queensland, Australia.  
Type: syntypes of both sexes, ?South Australia Museum, Adelaide.
  10. *P. palaeomorpha* Turner [*Phylomictis*], 1897:30.  
Type-locality: Brisbane, Queensland, Australia.

Type: syntype male, Australian National Insect Collection, Canberra.

11. *P. sarcinopa* Meyrick [*Phylomictis*], 1920:322.  
Type-locality: Brisbane, Queensland, Australia.  
Type: holotype female, British Museum (Natural History).

### Genus *Proscedes* Diakonoff

#### MAP 3

*Proscedes* Diakonoff, 1954:137. [Type-species: *Proscedes torquigera* Diakonoff by original designation.]

1. *P. torquigera* Diakonoff [*Proscedes*], 1954:138.  
Type-locality: Rattan Camp, 1200 m, West Irian, New Guinea.  
Type: holotype male, Rijksmuseum van Natuurlijke Historie, Leiden.

### Genus *Synchalara* Meyrick

#### MAP 4

*Synchalara* Meyrick, 1917:60. [Type-species: *Agriophara rhombota* Meyrick by original designation.]

1. *S. argoplaca* (Meyrick) [*Agriophara*], 1907a:743.  
Type-locality: Maskeliya, Ceylon.  
Type: lectotype male, British Museum (Natural History).
2. *S. byrsina* (Meyrick) [*Agriophara*], 1907b:151.  
Type-locality: Khasi Hills, Assam, India.  
Type: lectotype male, British Museum (Natural History).
3. *S. malacobryas* Meyrick [*Synchalara*], 1938:518.  
Type-locality: Mt. Tafa, 8500 ft, New Guinea.  
Type: male holotype, British Museum (Natural History).
4. *S. minax* (Meyrick) [*Agriophara*], 1907b:152.  
Type-locality: Khasi Hills, Assam, India.  
Type: lectotype male, British Museum (Natural History).
5. *S. rhombota* (Meyrick) [*Agriophara*], 1907a:981.  
Type-locality: Khasi Hills, Assam, India.  
Type: lectotype male, British Museum (Natural History).

## Literature Cited

- Busk, A.  
1934. Stenomidae. In *Lepidopterorum Catalogus*, 67:1-73.
- Carpenter, G. D. H.  
1953. The Genus *Euploea* (Lep. Danaidae) in Micronesia, Melanesia, Polynesia and Australia. A Zoogeographic Study. *Transactions of the Zoological Society of London*, 28:1-165.
- Clarke, J. F. Gates  
1955a. *Catalogue of the Type Specimens of Microlepidoptera in the British Museum (Natural History) Described by Edward Meyrick*. Volume 1. London, England:British Museum.  
1955b. *Catalogue of the Type Specimens of Microlepidoptera in the British Museum (Natural History) Described by Edward Meyrick*. Volume 2. London, England:British Museum.
- Common, I. F. B.  
1970. Lepidoptera (Moths and Butterflies). In *The Insects of Australia*. Commonwealth Scientific and Industrial Research Organization, Canberra, Australia:Melbourne University Press.
- Diakonoff, A.  
1948. Microlepidoptera of the Wissel Lakes, West New Guinea, II. *Treubia*, 19 (2):183-195.  
1954. Microlepidoptera of New Guinea, Results of the Third Archbold Expedition (American-Netherlands Indian Expedition 1938-1939), part 4. *Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen, Afdeling Natuurkunde*, series 2, 50 (1):1-99.
- Duckworth, W. Donald  
1969. Bredin-Archbold-Smithsonian Biological Survey of Dominica: West Indian Stenomidae (Lepidoptera-Gelechioidea). *Smithsonian Contributions to Zoology*, 4:1-21.  
1971. Neotropical Microlepidoptera XX: Revision of the Genus *Setiostoma* (Lepidoptera:Stenomidae). *Smithsonian Contributions to Zoology*, 106:1-44.
- Eyre, S. R.  
1968. *Vegetation and Soils: A World Picture*. 2nd edition. London, England: Edward Arnold Publishers, Ltd.
- Fletcher, T. B.  
1929. A List of the Generic Names Used for Microlepidoptera. *Memoirs of the Department of Agriculture in India*, 11:1-244.
- Gressitt, J. L.  
1956. Entomological Investigations in New Guinea Mountains. *Proceedings, Hawaiian Entomological Society*, 16 (1):47-69.
- Hartert, E.  
1924. The Birds of St. Mathias Island. *Novitates Zoologicae*, 31:261-275.
- Hodges, R. W.  
1966. Revision of Nearctic Gelechiidae, I, The *Lita* Group (Lepidoptera:Gelechioidea). *Proceedings of the United States National Museum*, 119 (3547):1-66.
- Holloway, J. D., and N. Jardine  
1968. Two Approaches to Zoogeography: a Study Based on the Distributions of Butterflies, Birds and Bats in the Indo-Australian Area. *Proceedings of the Linnean Society of London*, 179 (2):153-188.
- Lower, O. B.  
1892. Descriptions of New South Australian Lepidoptera. *Transactions of the Royal Society of South Australia*, 15:5-17.  
1893. New Australian Lepidoptera. *Transactions of the Royal Society of South Australia*, 17:146-185.  
1894. New Australian Heterocera. *Transactions of the Royal Society of South Australia*, 18:77-113.  
1899. Descriptions of New Australian Lepidoptera. *Proceedings of the Linnean Society of New South Wales*, 24:83-116.  
1902. Descriptions of New Genera and Species of Australian Lepidoptera. *Transactions of the Royal Society of South Australia*, 26:212-247.  
1903. Descriptions of New Species of *Xysmatodoma* & c. *Transactions of the Royal Society of South Australia*, 27:216-239.  
1908. New Australian Lepidoptera. No. XXV. *Transactions of the Royal Society of South Australia*, 32:110-120.  
1923. Descriptions of New Australian Lepidoptera. *Transactions of the Royal Society of South Australia*, 47:54-57.
- Lucas, T. P.  
1900. New Species of Queensland Lepidoptera. *Proceedings of the Royal Society of Queensland*, 15:137-161.
- Meyrick, E.  
1890. Descriptions of Australian Lepidoptera. *Transactions of the Royal Society of South Australia*, 13:23-81.  
1897. XVI. Descriptions of New Lepidoptera from Australia and New Zealand. *Transactions of the Entomological Society of London*, 1897:367-390.  
1907a. Descriptions of Indian Microlepidoptera. *Journal of the Bombay Natural History Society*, 17:730-993.  
1907b. Descriptions of Indian Microlepidoptera. *Journal of the Bombay Natural History Society*, 18:137-159.  
1910. Descriptions of Malayan Microlepidoptera. *Transactions of the Entomological Society of London*, 1910:430-478.  
1914. Descriptions of Indian Microlepidoptera. *Journal of the Bombay Natural History Society*, 18:118-129.  
1915. *Exotic Microlepidoptera*, 1:368, 410. Marlborough, England: Published by the author.  
1917. *Exotic Microlepidoptera*, 2:59-60. Marlborough, England: Published by the author.  
1918. *Exotic Microlepidoptera*, 2:197. Marlborough, England: Published by the author.  
1920. *Exotic Microlepidoptera*, 2:322. Marlborough, England: Published by the author.  
1921. *Exotic Microlepidoptera*, 2:445-446. Marlborough, England: Published by the author.

1925. *Exotic Microlepidoptera*, 3:159. Marlborough, England: Published by the author.
1930. *Exotic Microlepidoptera*, 3:560. Marlborough, England: Published by the author.
1931. *Exotic Microlepidoptera*, 4:43. Marlborough, England: Published by the author.
1938. Papuan Microlepidoptera. *Transactions of the Royal Entomological Society of London*, 87:503-528.
- Newman, E.  
1855. Characters of a Few Australian Lepidoptera, Collected by Mr. Thomas R. Oxley. *Transactions of the Entomological Society of London*, new series, 3:281-300.
- Powell, J. A.  
1973. Systematic Monograph of New World Ethmiid Moths (Lepidoptera:Gelechioidea). *Smithsonian Contributions to Zoology*, 120:302 pages, 138 figures.
- Rosenstock, R.  
1885. Notes on Australian Lepidoptera, with Descriptions of New Species. *The Annals and Magazine of Natural History*, 16:421-443.
- Smith, A. C.  
1970. *The Pacific as a Key to Flowering Plant History*. University of Hawaii, Harold L. Lyon Arboretum Lecture Number One, 1-26.
- Toxopeus, L. J.  
1950. The Geological Principles of Species Evolution in New Guinea. *Eighth International Congress of Entomology, Stockholm*, 1948:508-522.
- Turner, A. J.  
1897. The Xyloryctidae of Queensland. *Annals of the Queensland Museum*, 4:3-33.  
1900. New Micro-Lepidoptera — Mostly from Queensland. *Transactions of the Royal Society of South Australia*, 24:6-23.  
1906. New Australian Lepidoptera, with Synonymic and Other Notes. *Transactions of the Royal Society of South Australia*, 30:118-142.
1917. Lepidopterological Gleanings. *Proceedings of the Royal Society of Queensland*, 24 (7) :70-106.
1919. The Australian Gelechiidae (Lepidoptera). *Proceedings of the Royal Society of Queensland*, 31 (10) :108-172.
- Viette, P.  
1954. Description de Nouveaux Tineides Malgaches (Lepidopteres). *Mémoires de L'Institut Scientifique de Madagascar, Series E*, 5:1-38.  
1956a. Microlépidoptères de Madagascar Nouveaux ou peu Connus. *Le Naturaliste Malgache*, 8 (1):107-127.  
1965b. Tineoidea (s. l.) Nouveaux ou peu Connus de Madagascar (Lepidoptera). *Bollettino del Laboratorio di Zoologia Generale e Agraria "Filippo Silvestri", Portici*, 23:460-472.  
1958. Microlépidoptères Nouveaux de Madagascar. *Revue Française d'Entomologie*, 25 (2) :111-124.  
1962. Descriptions de Quelques Nouveaux Microlépidoptères de Madagascar et des Comores. *Lambillionea*, 62 (3-6) :23-30.  
1968. Descriptions de Nouvelles Espèces de Microlépidoptères de Madagascar et de L'île Marion. *Bulletin Mensuel de la Société Linnéenne de Lyon*, 37 (2) :83-91.
- Walker, F.  
1864. *List of the Specimens of Lepidopterous Insects in the Collection of the British Museum*. Volume XXIX. London. England: British Museum.  
1866. *List of the Specimens of Lepidopterous Insects in the Collection of the British Museum*. Volume XXXV, Supplement, Part 5. London, England: British Museum.
- Zeuner, F. E.  
1943. Studies in the Systematics of *Troides* Hon. and its Allies. *Transactions of the Zoological Society of London*, 25:107-184.



## Publication in Smithsonian Contributions to Zoology

*Manuscripts* for serial publications are accepted by the Smithsonian Institution Press, subject to substantive review, only through departments of the various Smithsonian museums. Non-Smithsonian authors should address inquiries to the appropriate department. If submission is invited, the following format requirements of the Press will govern the preparation of copy.

*Copy* must be typewritten, double-spaced, on one side of standard white bond paper, with 1½" top and left margins, submitted in ribbon copy with a carbon or duplicate, and accompanied by the original artwork. Duplicate copies of all material, including illustrations, should be retained by the author. There may be several paragraphs to a page, but each page should begin with a new paragraph. Number consecutively all pages, including title page, abstract, text, literature cited, legends, and tables. The minimum length is 30 pages, including typescript and illustrations.

The *title* should be complete and clear for easy indexing by abstracting services. Taxonomic titles will carry a final line indicating the higher categories to which the taxon is referable: "(Hymenoptera: Sphecidae)." Include an *abstract* as an introductory part of the text. Identify the *author* on the first page of text with an unnumbered footnote that includes his professional mailing address. A *table of contents* is optional. An *index*, if required, may be supplied by the author when he returns page proof.

Two *headings* are used: (1) text heads (boldface in print) for major sections and chapters and (2) paragraph sideheads (caps and small caps in print) for subdivisions. Further headings may be worked out with the editor.

In *taxonomic keys*, number only the first item of each couplet; if there is only one couplet, omit the number. For easy reference, number also the taxa and their corresponding headings throughout the text; do not incorporate page references in the key.

In *synonymy*, use the short form (taxon, author, date:page) with a full reference at the end of the paper under "Literature Cited." Begin each taxon at the left margin with subsequent lines indented about three spaces. Within an entry, use a period-dash (.—) to separate each reference. Enclose with square brackets any annotation in, or at the end of, the entry. For *references within the text*, use the author-date system: "(Jones, 1910)" and "Jones (1910)." If the reference is expanded, abbreviate the data: "Jones (1910:122, pl. 20: fig. 1)."

Simple *tabulations* in the text (e.g., columns of data) may carry headings or not, but they should not contain rules. Formal *tables* must be submitted as pages separate from the text, and each table, no matter how large, should be pasted up as a single sheet of copy.

Use the *metric system* instead of, or in addition to, the English system.

*Illustrations* (line drawings, maps, photographs, shaded drawings) can be intermixed throughout the printed text. They will be termed *Figures* and should be numbered consecutively; however, if a group of figures is treated as a single figure, the components should be indicated by lowercase italic letters on the illustration, in the legend, and in text references: "Figure 9b." If illustrations (usually tone photographs) are printed separately from the text as full pages on a different stock of paper, they will be termed *Plates*, and individual components should be lettered (Plate 9b) but may be numbered (Plate 9: figure 2). Never combine the numbering system of text illustrations with that of plate illustrations. Submit all legends on pages separate from the text and not attached to the artwork. An instruction booklet for the preparation of illustrations is available from the Press on request.

In the *bibliography* (usually called "Literature Cited"), spell out book, journal, and article titles, using initial caps with all words except minor terms such as "and, of, the." For capitalization of titles in foreign languages, follow the national practice of each language. Underscore (for italics) book and journal titles. Use the colon-parentheses system for volume, number, and page citations: "10(2):5-9." Spell out such words as "figures," "plates," "pages."

For *free copies* of his own paper, a Smithsonian author should indicate his requirements on "Form 36" (submitted to the Press with the manuscript). A non-Smithsonian author will receive 50 free copies; order forms for quantities above this amount with instructions for payment will be supplied when page proof is forwarded.

