



1. Darwin, C. *The Descent of Man, And Selection in Relation to Sex* (Murray, London, 1871).
2. Elgar, M. A., Ghaffar, N. & Read, A. F. *J. Zool.* **222**, 455–470 (1990).
3. Vollrath, F. *Z. Tierpsychol.* **53**, 61–78 (1980).
4. Head, G. P. *Evolution* **49**, 776–781 (1995).
5. Gerhardt, U. *Z. Morphol. Okol. Tiere* **1**, 567–618 (1924).
6. Vollrath, F. & Parker, G. A. *Nature* **360**, 156–159 (1992).
7. Hormiga, G., Eberhard, W. G. & Coddington, J. A. *Antst. J. Zool.* **43**, 313–364 (1995).
8. Scharff, N. & Coddington, J. A. *Zool. J. Linn. Soc.* (in the press).
9. Griswold, C. E., Coddington, J. A., Hormiga, G. & Scharff, N. *Zool. J. Linn. Soc.* (in the press).
10. Bonnet, P. *Gnathosia* **29**, 183–200 (1975).
11. Robinson, M. H. & Robinson, B. *Smithson. Contr. Zool.* **218**, 1–22 (1976).
12. Felsenstein, J. *Am. Nat.* **125**, 1–15 (1985).
13. Farris, J. S. *Syst. Zool.* **19**, 83–92 (1970).
14. Swofford, D. & Maddison, W. P. *Math. Biosci.* **87**, 199–229 (1987).
15. Maddison, W. P. & Maddison, D. R. *MacClade: Analysis of Phylogeny and Character Evolution*, version 3.0 (Sinauer, Sunderland, MA, 1992).

Vollrath and Parker reply — Spider males are smaller than females, with very few exceptions<sup>1,2</sup>. This sexual dimorphism is an ancient trait: males are marginally smaller in liphistiid spiders ('living fossils' with a segmented abdomen)<sup>3</sup> and trapdoor spiders<sup>4</sup>. Compared with 'modern' labidognath spiders<sup>5–7</sup>, these 'primitive' orthognath spiders typically have more instars (moulting stages) and larger body size. Coddington *et al.* raise two important questions. First, does modern *Nephila* have larger females and smaller males than its ancestors? Second, did ancestral females have fewer instars, and males more, than the average for this taxon?

Answering these questions with morphological data alone is difficult, if not impossible, especially if the cladistic hypothesis is contested (see refs 8–10). It may be more profitable to seek answers in life-




Figure 3 Number of moults in 'tetragnathid' spiders. M, males; F, females. *N ma*, *Nephila maculata*; *N ms*, *N. madagascariensis*; *N c*, *N. clavipes*; *T m*, *Tetragnatha montana*; *T n*, *T. nigrata*; *P c*, *Pachygnathat clercki*; *Z x*, *Zygiella x-notata*; *M s*, *Meta segmentata*. The original data (refs given) have been normalized to count from the instar (stadium) that leaves the eggsac (all have 1 extra larval instar inside the eggsac). Note that in the ancestral *Atypus karschi*<sup>4</sup> males mature in 8 or 9 and females in 9–11 instars; males of the 'living fossil' *Liphistius*<sup>3</sup> have between 10 and 31 instars. Generally, the more instars, the larger the spider.

history data. Puzzling life-history traits (measured on the species level) are often best explained by models<sup>11</sup>. Our<sup>1</sup> puzzle was the very real observation that males of the golden orb spider *N. clavipes* are much smaller than their females, although all measured reproductive advantages suggested selection for a larger male size<sup>12</sup>. In females, large size always benefits fecundity<sup>13–15</sup>.

Our model<sup>1</sup> is designed to predict the extent of sexual dimorphism in a species (see also ref. 16). It trades off the mortality rate of each sex (which favours early reproduction, and hence smaller adult body size) against the reproductive advantages of later maturation (being bigger can mean laying more eggs or gaining greater mating prospects). The ratio of the evolutionarily stable body sizes for the two sexes generates the 'evolutionarily stable strategy' of sexual dimorphism. Thus we did not set out to prove absolute selection to reduce male size, but instead predicted quite generally the relative size of the two sexes from life-history traits that can be measured in the field and the laboratory.

When related to data about *N. clavipes* and similar species, the model predicted a general trend for dwarf males (dwarf relative to females) to occur in species with 'sit-and-wait' habits as opposed to searching and hunting habits: males must rove (and consequently suffer high mortality) to find females that are sedentary. A simple cladistic analysis (using accepted traits on a family level) seemed to confirm this trend and validate our model (Fig. 3 of ref. 1). The best confirmation is the discovery<sup>17</sup> of true dwarf males in a mygalomorph (ancestral) spider living in a marginal, high-mortality habitat, where the burrowing females are less at risk than the roving males.

When proposing female giantism (as opposed to male dwarfing) for *N. clavipes*, Coddington *et al.* make various assumptions. They claim that the males are typical for this taxon in size and instar number whereas the females are atypical in both. But in fact the data are ambiguous (Fig. 3). First, it seems that *Nephila* is unusual for tetragnathids (and most other spiders) in having non-overlapping male/female instar numbers. Second, *Nephila* can have rather few male instars. Third, the number of female instars is indeed large for tetragnathids, and so is body size. But this is an ancestral trait and could be fitted to phylogenetic hypotheses placing *Nephila* near the beginning of the ecribellate branch of orb weavers<sup>8,10</sup>. This would imply that other tetragnathids have a reduced number of instars. Finally, *Nephila* might not even be a true tetragnathid<sup>8,10</sup>.

It is never easy to decide how to read phylogenetic information. But the pertinent point of the controversy of mini males versus giant females is that we desperately need

more life-history data on unusual spiders, that is, species with extreme or absent sexual size dimorphism. This is the only way to solve a puzzle that intrigued Darwin and many researchers since: a puzzle that concerns the difference between male and female sizes, not their absolute values.

Fritz Vollrath

Department of Zoology, Universtetet B135, DK-8000 Aarhus C, Denmark

Geoff A. Parker

Department of Evolutionary Biology, PO Box 147, University of Liverpool, Liverpool L69 3BX, UK

1. Vollrath, F. & Parker, G. A. *Nature* **360**, 156–159 (1992).
2. Head, G. P. *Evolution* **49**, 776–781 (1995).
3. Schwendinger, P. *J. Zool. Scripta* **19**, 331–351 (1990).
4. Miyashita, T. *Acta Arachnol.* **41**, 177–186 (1992).
5. Schaefer, M. *Zool. Jb. Syst. Jena* **103**, 127–289 (1976).
6. Juberthie, C. *Bull. Soc. Hist. Nat. Toulouse* **89**, 299–318 (1954).
7. Bonnet, P. *Bull. Soc. Entomol. France* **1926**, 67–69 (1926).
8. Eberhard, W. G. *Evolution* **36**, 1067–1095 (1982).
9. Shear, W. A. in *Spiders: Webs, Behavior and Evolution* (ed. Shear, W. A.) 364–400 (Stanford Univ. Press, 1986).
10. Coddington, J. A. in *Spiders: Webs, Behavior and Evolution* (ed. Shear, W. A.) 319–363 (Stanford Univ. Press, 1986).
11. McNamara, J. M. & Houston, A. I. *Nature*, **380**, 215–221 (1996).
12. Vollrath, F. *Z. Tierpsychol.* **53**, 61–78 (1980).
13. Robinson, M. H. & Robinson, B. *Smithson. Contr. Zool.* **218**, 1–22 (1976).
14. Bonnet, P. *Bull. Soc. Zool. France* **54**, 501–523 (1929).
15. Vollrath, F. *Verh. Naturwiss. Verein. Hamburg* **26**, 277–289 (1983).
16. Parker, G. A. *J. Fish Biol.* **41**, (suppl. B) 1–20 (1992).
17. Main, B. Y. *Acta Zool. Fenn.* **190**, 273–278 (1990).