

Redescription of
Echinoderes dujardini (Kinorhyncha)
with Descriptions of
Closely Related Species

ROBERT P. HIGGINS

SERIES PUBLICATIONS OF THE SMITHSONIAN INSTITUTION

Emphasis upon publication as a means of “diffusing knowledge” was expressed by the first Secretary of the Smithsonian. In his formal plan for the Institution, Joseph Henry outlined a program that included the following statement: “It is proposed to publish a series of reports, giving an account of the new discoveries in science, and of the changes made from year to year in all branches of knowledge.” This theme of basic research has been adhered to through the years by thousands of titles issued in series publications under the Smithsonian imprint, commencing with *Smithsonian Contributions to Knowledge* in 1848 and continuing with the following active series:

Smithsonian Contributions to Anthropology
Smithsonian Contributions to Astrophysics
Smithsonian Contributions to Botany
Smithsonian Contributions to the Earth Sciences
Smithsonian Contributions to Paleobiology
Smithsonian Contributions to Zoology
Smithsonian Studies in Air and Space
Smithsonian Studies in History and Technology

In these series, the Institution publishes small papers and full-scale monographs that report the research and collections of its various museums and bureaux or of professional colleagues in the world of science and scholarship. The publications are distributed by mailing lists to libraries, universities, and similar institutions throughout the world.

Papers or monographs submitted for series publication are received by the Smithsonian Institution Press, subject to its own review for format and style, only through departments of the various Smithsonian museums or bureaux, where the manuscripts are given substantive review. Press requirements for manuscript and art preparation are outlined on the inside back cover.

S. Dillon Ripley
Secretary
Smithsonian Institution

SMITHSONIAN CONTRIBUTIONS TO ZOOLOGY • NUMBER 248

Redescription of
Echinoderes dujardini (Kinorhyncha)
with Descriptions of
Closely Related Species

Robert P. Higgins



SMITHSONIAN INSTITUTION PRESS

City of Washington

1977

ABSTRACT

Higgins, Robert P. Redescription of *Echinoderes dujardinii* (Kinorhyncha) with Descriptions of Closely Related Species. *Smithsonian Contributions to Zoology*, number 248, 26 pages, 31 figures, 2 tables, 1977.—A key to the adults of the genus *Echinoderes* is followed by a redescription of *E. dujardinii* from both adult and juvenile specimens. North American records for *E. dujardinii* are corrected by the description of *E. kozloffii*, new species. *Echinoderes pennaki*, a species sympatric with *E. kozloffii*, is redescribed, and the latter's sibling species, *E. pacificus*, is discussed. Synonomies and distribution records are given for species mentioned.

OFFICIAL PUBLICATION DATE is handstamped in a limited number of initial copies and is recorded in the Institution's annual report, *Smithsonian Year*. SERIES COVER DESIGN: The coral *Montastrea cavernosa* (Linnaeus).

Library of Congress Cataloging in Publication Data

Higgins, Robert P.

Redescription of *Echinoderes dujardinii* (Kinorhyncha) with descriptions of closely related species.

(Smithsonian contributions to zoology ; no. 248)

Bibliography: p.

Supt. of Docs. no.: SI 1.27:248

1. *Echinoderes dujardinii*. 2. *Echinoderes*. 3. Kinorhyncha—Classification. I. Title: Redescription of *Echinoderes dujardinii* (Kinorhyncha) II. Series: Smithsonian Institution. Smithsonian contributions to zoology ; no. 248.

QL1.S54 no. 248 [QL391.A2] 591'.08s [595'.183] 76-608300

Contents

	Page
Introduction	1
Methods	2
Acknowledgments	3
Key to Adults of the Genus <i>Echinoderes</i>	3
<i>Echinoderes dujardinii</i> Claparède	4
<i>Echinoderes kozloffii</i> , new species	13
<i>Echinoderes pennaki</i> Higgins	18
<i>Echinoderes pacificus</i> Schmidt	21
<i>Echinoderes brevicaudatus</i> , substitute name for <i>E. brevispinosus</i> Higgins ..	21
Literature Cited	24

Redescription of *Echinoderes dujardinii* (Kinorhyncha) with Descriptions of Closely Related Species

Robert P. Higgins

Introduction

The first published observation of a kinorhynch is that of the French naturalist Félix Dujardin (1851), who based his account on specimens found while examining material washed from algae collected near St. Malo on the coast of Normandy in 1841. In a manner not uncommon throughout the history of the study of meiofauna, Dujardin was unable to place the new invertebrate in an existing higher taxonomic category although he recognized its morphological affinities with acanthocephalans, rotifers, sipunculids, nematodes, tardigrades, and various crustaceans ("copepods without legs"). Ten years after his discovery, Dujardin (1851) introduced the first kinorhynch as "a tiny marine animal, *l'Echinodère*, constituting an intermediate form between the crustaceans and worms." A few years later Leuckart (1854) noted that he had seen "*l'Echinodère*" at Helgoland in 1846 but had assumed that it was a dipteran larva.

"*L'Echinodère*" became *Echinoderes dujardinii* Claparède, 1863 when the latter author described specimens from St. Vaast la Hougue, not far from St. Malo where Dujardin made his discovery. The following year, Gosse (1864), apparently unaware

of Claparède's publication, also described this species from the same locality, naming it *Echinodera dujardini*.

These descriptions suffered from the authors' lack of experience with the taxon, as have many subsequent ones, including my own. Despite Zelinka's (1928) more comprehensive description of *E. dujardinii* in his *Monographie der Echinodera*, certain taxonomic ambiguities have persisted. The lack of preserved specimens, especially types, has compounded the problem; consequently, the reported distribution of *E. dujardinii*, as well as other species, must be considered with due caution.

Since it was described, *E. dujardinii* has been reported from 26 localities including the northern and southern coasts of Europe, the Black Sea, Canary Islands, Japan, and the northwestern coast of the United States.

No specimens are available to confirm Chitwood's (1964a) report of *E. dujardinii* from Tomales Bay, California, one of the two northwestern U.S. localities. Neither an unpublished photograph by Chitwood (pers. comm.) of one of the specimens, nor sketches of specimens collected at the same locality and time (1960) by Dr. Tor G. Karling (pers. comm.), indicate that the species from Tomales Bay is *E. dujardinii*. Specimens sent to me from San Juan Island, Washington, by Dr. Eugene N. Kozloff, resembled *E. dujardinii* sufficiently to

Robert P. Higgins, Smithsonian Oceanographic Sorting Center, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560.

justify the tentative use of this name (Kozloff, 1972; Merriman and Corwin, 1973); however, both Kozloff and I expressed doubts that could be answered only by comparing the San Juan Island specimens with specimens from either the type-locality or localities (Naples and Trieste) where Zelinka (1928) obtained *E. dujardinii* from his redescription. My collection from the latter two localities failed to provide specimens of *E. dujardinii* but Kozloff, in 1973, was successful in collecting this species at Roscoff, not far from St. Vaast la Hougue.

The purpose of this paper is to redescribe *E. dujardinii*, clarifying both the original description by Claparède (1863) and the redescription by Zelinka (1928). In addition, this paper will correct the distribution record inasmuch as the San Juan Island specimens, reported as *E. dujardinii*, constitute a new species described below. Because of the co-occurrence of the new species with *E. pennaki* Higgins, 1960, the description of which reflects my own inexperience at the time, I shall redescribe it as well. The description of *E. pacificus* Schmidt, 1974 is included because it closely resembles the new species from San Juan Island. Finally, while studying the taxonomic history of *E. dujardinii*, I discovered that a name I proposed for a species from the Red Sea was preoccupied; therefore, a substitute name is offered for *E. brevispinosus* Higgins, 1966a.

METHODS.—Specimens were preserved in 70 percent ethyl alcohol, 5 percent formalin, or Duboscq and Brasil's fluid, and then transferred to a 70 percent ethyl alcohol-5 percent glycerin solution that was allowed to evaporate to glycerin. Most specimens were removed from the glycerin and individually placed in Hoyer's mounting medium, between two coverslips, and positioned on Cobb aluminum slide frames. This mounting procedure allows the slide to be placed on either of its surfaces so that both dorsal and ventral aspects of the specimen can be observed.

Hoyer's medium is necessary to soften the specimen so that, by judicious manipulation of the coverslip, the specimen will assume a dorsoventral position; this medium also clears the specimen, thus revealing the detailed structure of the exoskeleton.

A disadvantage of the Hoyer's medium is its tendency to clear the specimen too much, especially over a period of several years. This may be

partially overcome by reducing, by about 30 percent, the amount of chloral hydrate used in the medium and by sealing the preparation with Murrayite soon after the fluid has solidified. In this series of preparations, some specimens were remounted in glycerin once they had been oriented dorsoventrally. This is a procedure that should be used only when an adequate series of specimens are available, since in transferring them many dorsal spines are often broken. A few specimens were mounted in glycerin without first mounting them in Hoyer's medium. In all instances, coverslips were sealed with Murrayite.

Each specimen was studied with the use of Zeiss differential interference contrast optics and meristically analyzed. The resulting data are expressed in a standard format of abbreviations and terminology (Higgins, 1967, 1969a). Measurements are given in microns (μm); ratios (i.e., SW/TL) are expressed in percent of the total length (TL) measured on the midline, from the anterior margin of segment 3 (first trunk segment) to the posterior margin of segment 13, exclusive of spines. Maximum sternal width (MSW) is measured at the anteroventral margin of the widest pair of sternal plates as first encountered in measuring each segment from anterior to posterior. Sternal width at segment 12 (SW), or standard width, is measured at the anteroventral margin of the 12th sternal plates.

Middorsal spines (D), lateral spines (L), and lateral accessory appendages (LA) are numbered by segment and their cumulative mean length expressed by Dm, Lm, and LAm, respectively. Measurements are given for the lateral terminal spines (LTS), lateral terminal accessory spines (LTAS), midterminal spine (MTS), and penis spines (P) in males. The first penis spine (P-1) is usually the anteriormost of three such spines and is dorsally displaced; the second (P-2) is usually the shortest and often more truncate (probably the functional penis); the third (P-3) is usually adjacent to the second or slightly posterior to it. Both P-2 and P-3 are best observed in ventral aspect.

Several lateral spines appear to function as adhesive tubes. This study will introduce the adhesive tubes of the fourth segment (L-4) as homologues of other lateral spines. Appendages that function as adhesive tubes will be noted in the appropriate section of the text.

Specimens mentioned in this paper are deposited in the National Museum of Natural History, Smithsonian Institution, under the catalog numbers of the old United States National Museum (USNM).

ACKNOWLEDGMENTS.—I am grateful to Mr. John C. Boykin, University of Washington, the first to collect the new species from San Juan Island and recognize its taxonomic problems; and to our mutual friend and colleague from this same institution, Dr. Eugene N. Kozloff, who accepted the challenge of culturing the new species, and provided me with specimens both of the new species and of *E. dujardinii* that have made this study possible. I also acknowledge the cooperation of Dr. Jean Merriman, California State College at Sonoma, whose recent doctoral dissertation and publication have contributed to the knowledge of the new species from San Juan Island. The personal communications of Dr. Tor G. Karling, University of

Stockholm, and the late Dr. B. G. Chitwood were helpful in this study. Dr. Peter Schmidt, Institute für Zoologie der RWTH, Aachen, generously provided type specimens of *E. pacificus* for my use. Thanks also go to Prof. Peter Dohrn, Stazione Zoologica, Naples and to Dr. Jose Stirn, Marine Laboratory, Portoroz, Yugoslavia, for making their facilities available to me. Drs. Raymond B. Manning and Horton H. Hobbs, Jr., were generous in their advice and consultation during the preparation of this manuscript and, along with Dr. Richard S. Houbrick, in the reading of the final draft. I also wish to express my appreciation to Mrs. Eleanor Goldsmith, Mrs. Martha Brewster, and Mrs. Marie Wallace for their help in the preparation of this manuscript. Finally, I am pleased to acknowledge the Sumner Gerard Foundation and the Smithsonian Foreign Currency Program for their financial support.

Key to Adults of the Genus *Echinoderes*

- | | |
|---|--|
| 1. Middorsal spines absent | 2 |
| Middorsal spines present | 5 |
| 2. Lateral spines (except for terminal segment) absent | |
| <i>Echinoderes maxwelli</i> (Omer-Cooper, 1957) | 3 |
| Lateral spines present | 3 |
| 3. Lateral spines on segments 7, 10, and 12 only | <i>Echinoderes bengalensis</i> (Timm, 1958) |
| Lateral spines absent on segment 12 | 4 |
| 4. Lateral spines on segments 7, 10, and 11 only | <i>Echinoderes caribiensis</i> Kirsteuer, 1964 |
| Lateral spines on segments 7 and 10 only, segments 3-4 expanded laterally | <i>Echinoderes capitata</i> (Zelinka, 1928) |
| 5. Middorsal spines on segments 6-10 only | 6 |
| Middorsal spines with other arrangement | 13 |
| 6. Lateral spines on segments 7-12 only | 7 |
| Lateral spines with other arrangement | 21 |
| 7. Lateral accessory spine on segment 10 | <i>Echinoderes dujardinii</i> Claparède, 1863 |
| Lateral accessory spines absent on segments 3-11 | 8 |
| 8. Trunk length greater than 300 μ m | 9 |
| Trunk length less than 300 μ m | 11 |
| 9. Lateral spine on segment 12 short (12-17 μ m), less than half the length of lateral spine on segment 11 | <i>Echinoderes pacificus</i> Schmidt, 1974 |
| Lateral spines on segment 12 nearly equal to length of lateral spines on segment 11 | 10 |
| 10. Lateral terminal accessory spines 12%-15% of trunk length, midventral placid wider than adjacent placids | <i>Echinoderes kozloffii</i> , new species |
| Lateral terminal accessory spines about 8% of trunk length, midventral placid not wider than adjacent placids | <i>Echinoderes pilosus</i> Lang, 1949 |
| 11. Middorsal spine on segment 10 slightly longer than middorsal spine on segment 9 | <i>Echinoderes ehlersi</i> Zelinka, 1913 |
| Middorsal spine on segment 10 twice as long as middorsal spine on segment 9 | 12 |
| 12. Lateral terminal accessory spines about 41% of trunk length, no lateral accessory spine on segment 12 | <i>Echinoderes worthingi</i> Zelinka, 1928 |
| Lateral terminal accessory spines about 23% of trunk length, lateral accessory spine adjacent to lateral spine 12 | <i>Echinoderes ferrugineus</i> Zelinka, 1928 |
| 13. Middorsal spine on segment 11 (juvenile?) | <i>Echinoderes sonadiae</i> Timm, 1958 |
| Middorsal spine absent on segment 11 | 14 |

14. Middorsal spine on segment 7 15
 Middorsal spine absent on segment 7 18
15. Middorsal spines on segments 6, 7, and 9 only *Echinoderes setigera* Greeff, 1869
 Middorsal spines with other arrangement 16
16. Middorsal spines on segments 6-9 only *Echinoderes druxi* d'Hondt, 1973
 Middorsal spine on segment 5 17
17. Middorsal spines on segments 5-8 only *Echinoderes tchefouensis* Lou, 1934
 Middorsal spines on segments 5-10 only *Echinoderes borealis* Greeff, 1869
18. Middorsal spines on segments 6 and 9 only *Echinoderes citrinus* Zelinka, 1928
 Middorsal spines on segments 6, 8, and 10 only 19
19. Lateral accessory spines on segments 3, 8-10 with subdorsal spine on segment 3, terminal tergal extensions evenly tapering to point *Echinoderes newcaledoniensis* Higgins, 1967
 Lateral accessory spine on segment 10 or missing on all segments except terminal segment of female, terminal tergal extensions with uneven margins 20
20. Lateral accessory spine on segment 10 only, middorsal spines short, (15-18 μm). Terminal tergal extensions mesially recurved and somewhat rectangular with small protuberances at corners *Echinoderes riedli* Higgins, 1966a
 Lateral accessory spines absent on all segments except terminal segment of female, middorsal spines long (73-125 μm). Terminal tergal extensions rounded with elongate central spine *Echinoderes arlis* Higgins, 1966b
21. Posterior border of 12th tergal plate interrupted medially to form broad spine-shaped process. Small lateral spine or prominent hair (about 8 μm) on segment 12, similar spine or hair on segment 13 anterior to lateral terminal accessory spine of female *Echinoderes bookhouti* Higgins, 1964b
 Posterior border of 12th tergal plate even. Lateral spine or prominent hair absent on segment 12, segment 13 with lateral terminal accessory spine of female and lateral terminal spine only 22
22. Lateral spine on segments 7-11 only 23
 Lateral spines with other arrangement 25
23. Lateral terminal spines short, about 17% of trunk length. Spinous fringe most prominent at posteroventral margin of segment 4 and at lateral margins of sternal-tergal junctions of segments 9 and 10 *Echinoderes brevicaudatus*, substitute name for *Echinoderes brevispinosus* Higgins, 1966a
 Lateral terminal spines 24%-40% of trunk length, spinous fringe uniform on each segment where it occurs 24
24. Terminal sternal plates pointed, tergal plate extensions displaced medially from lateral terminal spines, midventral placid expanded at anterior margin, prominent spinous fringe along entire posteroventral margin of segment 3 *Echinoderes remanei* (Blake, 1930)
 Terminal sternal plates rounded, tergal plate extensions displaced laterally alongside lateral terminal spines, midventral placid evenly rounded or slightly truncate at anterior margin, spinous fringe more prominent near midline of posteroventral margin of segment 3 *Echinoderes pennaki* Higgins, 1960
25. Lateral spines on segments 6-9 only *Echinoderes canariensis* Greeff, 1869
 Lateral spines with other arrangement 26
26. Lateral spines on all segments (probably error in description) *Echinoderes steineri* (Chitwood, 1951)
 Lateral spines on segments 8-11 only 27
27. Lateral terminal spines about 50% of trunk length, lateral spine on segment 11 same length as or slightly longer than that of segment 10 *Echinoderes elongata* (Nyholm, 1947b)
 Lateral terminal spines equal to or longer than trunk length, lateral spine of segment 11 very long *Echinoderes levanderi* Karling, 1954

***Echinoderes dujardinii* Claparède**

FIGURES 1-12

"*Echinodère*" Dujardin, 1851:158, pl. 3: figs. 1-4 [St. Malo].—Schultze, 1853:253.

"Echinoceras" Leuckart, 1854:355 [erroneous spelling; Helgoland].

Echinoderes Dujardinii Claparède, 1863:90-92, pl. 16: figs. 7-13 [type-locality: St. Vaast la Hougue].—Leuckart, 1869:300.—Greeff, 1869:88-89, pl. 4: figs. 1-5 [by inference: Ostende, Nieuwpoort, Dieppe, Canary Islands; incorrectly


- cites Dujardin as author].—Panceri, 1876:4, 5, figs. 6, 7 [Ischia, incorrectly cites Greeff (sic) as author].—Reinhard, 1881:589 [Odessa, incorrectly cites Dujardin as author].—Moebius, 1884:7, 1887:117.—Reinhard, 1885:5.—Carus, 1885:184.—Ludwig, 1886:882.—Reinhard, 1887:438 [incorrectly cites Dujardin as author].—Perrier, 1893:453.—Zelinka, 1896:199 [incorrectly cites Greeff as author].—Grobben, 1905:336.—Schepotieff, 1907b:297, 298, pl. 17: figs. 1-6, 16a, pl. 18: figs. 22, 23, pl. 20: fig. 13 [Bergen, Naples, Brindisi, Rovigno].—Southern, 1914:69, 71 [Clew Bay, Blacksod Bay].—Zelinka, 1928:221 [*lapsus calami?*].—Abe, 1930:39-43.
- Echinodera Dujardinii* Gosse, 1864:403-404, pl. 2: fig. 16 [description taken from Dujardin, 1851 only].—Zelinka, 1928:228 [synonymy].
- Not *Echinoderes Dujardinii*.—Metschnikoff, 1865:459-461 [Helgoland, =*Echinoderes subfuscus* Zelinka, 1928]; 1869:190-193 [Selerno, =*Echinoderes subfuscus* Zelinka, 1928].
- Echinoderes brevispinosa* Metschnikoff, 1869:190 [Salerno].—Leuckart, 1869:300-301.—Reinhard, 1887:438.
- Echinoderes Sieboldii* Pagenstecher, 1875:117-123, pl. 7: figs. 1-6 [Palma]; 1877:88-89, fig. 69.—Carus, 1885:185.—Reinhard, 1887:438 [Odessa].—Schepotieff, 1907b:298, 299, pl. 17: fig. 7.—Abe, 1930:39-42.
- Echinoderes brevispinosus*.—Panceri, 1876:4 [correction of *E. brevispinosa* Metschnikoff, 1869].—Zelinka, 1928:228 [synonymy].
- Echinoderes meridionales* Panceri, 1876:5, 6, fig. 8 [Ischia].—Carus, 1885:185.—Zelinka, 1928:254, pl. 2: figs. 11-13 [synonymy, =*Habroderes meridionales*].
- Echinoderes dujardinii*(?).—Hartog, 1896:236, fig. 120 [Worthing].—McIntyre, 1962:503.
- Echinoderes dujardini*.—Hartog, 1896:237.—Remane, 1928:75, 76, figs. 13, 15 [incorrectly cites Greeff as author on p. 76 and in figure legends].—Johnston, 1938:5.—Higgins, 1960:88-90.—McIntyre, 1962:503.—Higgins, 1964a: 244, 246; 1964b:489-491; 1966a:120; 1967:75, 79; 1969b:113; 1971:25.—Kozloff, 1972:121.—Moore, 1973:349.—Higgins, 1974:512.—Boykin, 1974:40.—Schmidt, 1974:12.
- Echinoderes dujardinii*.—Levander, 1900:20.—Schepotieff, 1907a:147, 148, figs. 10-12 [Bergen, Naples].—Zelinka, 1907:132; 1908:630-641, figs. 1, 5, 6, 9 [incorrectly cites Greeff as author]; 1912:520-527 [incorrectly cites Greeff as author]; 1913:419, 420 [incorrectly cites Greeff as author]; 1928:221 [incorrectly cites Greeff as author], 228 [cites Greeff as author in synonymy], 228-235, figs. 17-19; pl. 3: figs. 1, 2, pl. 10: figs. 1-5, 7-23 [Naples, Trieste, Kiel].—Remane, 1928:60, 62, figs. 7, 9 [incorrectly cites Greeff as author].—Lou, 1934:3.—Remane, 1936: 333, figs. 215, 252, 266.—Zaneveld, 1938:261 [Scheveningen].—Nyholm, 1947a:424 [Gullmar Fjord]; 1947b:5, fig. 1.—Lang, 1949:3, 17.—Tokioka, 1949:67.—Hyman, 1951:180.—Karling, 1954:189-192.—Kirsteuer, 1964:389.—Marinov, 1964:62, 63, fig. 2 [Varna, doubtful record].
- Echinoderes sieboldii*.—Zelinka, 1928:228 [synonymy].—Remane, 1936:344 [synonymy].
- Habroderes dujardinii* Zelinka, 1928:248-250, pl. 4: figs. 1, 2 [described as "n. 1. (zu *Echinoderes dujardinii* Clap.)"].—Remane, 1936:332, 333, fig. 273 [noted as larval stage of *Echinoderes dujardinii*].
- Habroderes meridionales* Zelinka, 1928:254, pl. 2: figs. 11-13 [synonymy].—Remane, 1936:333 [noted as larval stage of *Echinoderes dujardinii*].
- Centropsis parallela* Zelinka, 1928:269, pl. 1: figs. 4, 5, 11, 12 [Naples, Trieste].—Remane, 1936:332, 333, fig. 273 [noted as larval stage of *Echinoderes dujardinii*].
- Echinoderes Masudai* Abe, 1930:42, 43, pl. 1: figs. 1, 2 [Hiroshima].—Remane, 1936:345, fig. 277.—Tokioka, 1949:67.
- Echinoderes sieboldii*.—Lou, 1934:3 [erroneous spelling].
- Echinoderes masudai*.—Lang, 1949:17.—Higgins, 1960:89 [nomen dubium]; 1966a:123; 1967:75; 1971:26.
- Echinoderes* aff. *dujardini*.—Băcescu et al., 1963:137, 138, figs. 1a-c [Agigea, probably not *Echinoderes dujardinii*].
- Not *Echinoderes brevispinosus* Higgins, 1966a:118-121, figs. 1, 2; 1966b:519.—Schmidt, 1974:14. [= *Echinoderes brevicaudatus*, substitute name].
- Not *Echinoderes dujardinii*.—Chitwood, 1964a:2 [Tomales Bay, Calif; identity uncertain].
- Not *Echinoderes dujardini*.—Merriman and Corwin, 1973:227-243, figs. 1-13 [San Juan Island, Washington; =*Echinoderes kozloffii*, new species].

REDESCRIPTION.—Adults (Figures 1-6), trunk length, 328-405 μm ; MSW-9, 78-85 μm , 19.9-24.9 percent of trunk length; SW, 70-83 μm , 18.9-24.9 percent of trunk length.


Second segment with 16 anteriorly rounded placids, midventral placid truncate, expanded slightly at anterior margin, distinctly larger than adjacent placids; trichoscalid plates on sides of midventral placid with medial indentation on anterior margin, posterior margin expanded laterally.

Segments 3-12 with short hairs, pattern distinctive (Figures 1, 2); posterior border of segments 5-11 with fine pectinate fringe ventrally, segments 6-11 with pectinate fringe dorsally; terminal segment with few hairlike processes along posterior margin, posterior margin of tergal plate deeply incised forming pointed extensions mesial to base of each lateral terminal spine, sternal plates broadly rounded with spinous extensions, 10-12 μm in length (Figures 3, 5).

Middorsal spines on segments 6-10, increasing uniformly in length, 13-29 μm ; lateral spines on segments 4 and 7-12, 14-29 μm in length; L-4, 7, 10 and 12 each with adhesive gland at base; L-10 accompanied by dorsally adjacent lateral accessory spine of nearly equal length, 21-26 μm ; L-4 usually longer (mean 24.2 μm) than remaining lateral spines; lateral terminal spines long, 160-192 μm , 40.7-54.8 percent of trunk length; lateral terminal accessory spines of female 52-62 μm in length; males without lateral terminal accessory spines but with penis spines in same position, anteriormost penis



FIGURES 1-2.—*Echinoderes dujardinii*, adult female (USNM 53342, RH125.14), neck and trunk segments: 1, ventral view; 2, dorsal view.



FIGURES 3-6.—*Echinoderes dujardinii*, segments 12-13: 3, ventral view, lateral half, adult female (USNM 53342, RH125.14); 4, dorsal view, lateral half of same adult female; 5, ventral view, lateral half, adult male (USNM 53342, RH125.35); 6, dorsal view, lateral half of same adult male.

spine (P-1) 23-33 μm in length, mesially adjacent penis spine (P-2) 24-36 μm , posteriorly adjacent penis spine (P-3) 26-39 μm in length.

Pachycycli (thickened anterior margins of trunk segments) well developed, forming a distinctive pattern at ventral midline and at attachments of lateral terminal spine muscles on segment 13; distinctive muscle scars on ventrolateral portion of first trunk segment, similar (but reversed orientation) lateral scars on eighth sternal plates; sensory spots, 2-3 μm in diameter, situated middorsally on segments 3-5, with two such spots on segment 12 (possibly one on each of segments 12 and 13), dorsolaterally on segments 6-11, and ventromesially on segments 5-12 (Figures 1, 2).

Morphometric data for adult specimens are shown in Table 1.

JUVENILE STAGES.—Preadult stage (J-6, "*Habroderes*-stage," Figures 7, 8) trunk length, 320-328 μm ; estimated MSW-9, 78-80 μm , 24.3-25.6 percent of trunk length; estimated SW, 65-72 μm , 19.8-23.5 percent of trunk length (estimated since tergal-sternal junctions are not defined in juvenile stages).

Second segment similar to that of adult, both placids and trichoscalid plates less well developed.

Trunk segments with fewer hairs than adult, pattern less distinctive and more variable; posterior borders of segments without pectinate fringe, with hairs (striations?) along border, group of prominent hairs (striations?) at ventral midline, terminal segment slightly incised dorsally and ventrally, sternal area without spinous extensions, tergal area with small extensions 2-4 μm in length, not evenly tapered.

Middorsal spines on segments 6-11, increasing uniformly in length, 17-39 μm ; lateral spines on segments 4 and 7-12, 17-26 μm in length; L-4, 7, 10 and 12 with poorly developed adhesive glands at base; L-10 and L-12 accompanied by a dorsally adjacent lateral accessory spine of nearly equal length, 21-25 μm , but thinner; L-4 more prominent than remaining lateral spines; lateral terminal spines long, 148-152 μm , 45.1-48.7 percent of trunk length; lateral terminal accessory spines 42-47 μm in length, sexes often indistinguishable in juvenile stages unless developing oocytes visible.

TABLE 1.—Measurements (μm) and indices (%) for *Echinoderes dujardini* adults

Character	Number	Range	Mean	Standard Deviation	Standard Error	Coefficient of Variability	Character	Number	Range	Mean	Standard Deviation	Standard Error	Coefficient of Variability		
TL	♂	26	328-380	350.5	15.4	3.0	L-4 (AT)	♂	16	17-26	23.0	2.3	0.6		
	♀	20	336-405	367.9	21.9	4.9		♀	17	22-28	25.2	1.6	0.4		
	♂♀	46	328-405	358.0	18.3	2.7		♂♀	33	17-28	24.2	1.9	0.3		
SW	♂	26	70-82	76.6	2.7	0.5	L-7	♂	19	13-25	17.8	3.6	0.8	20.2	
	♀	19	75-83	78.9	2.6	0.6		♀	17	14-26	18.9	3.5	0.8	18.3	
	♂♀	45	70-83	77.0	2.6	2.6		♂♀	36	13-26	18.3	3.5	0.6	19.0	
SW/TL	♂	26	19.3-24.9	21.6	1.3	0.2	L-8	♂	26	14-22	18.0	2.3	0.4	12.6	
	♀	19	18.9-24.4	21.5	1.5	0.4		♀	19	16-26	19.3	2.7	0.6	13.9	
	♂♀	45	18.9-24.9	21.5	1.4	0.2		♂♀	45	14-26	18.6	2.4	0.4	13.1	
MSW-9	♂	26	75-85	79.4	2.6	0.5	L-9	♂	26	17-23	20.4	2.1	0.4	10.5	
	♀	19	78-85	81.5	1.8	0.4		♀	19	16-28	20.6	2.8	0.6	13.6	
	♂♀	45	75-85	80.2	2.3	0.3		♂♀	45	16-28	20.5	2.4	0.4	11.8	
MSW/TL	♂	26	20.7-24.9	22.7	1.3	0.2	L-10	♂	26	20-25	22.2	1.2	0.2	5.6	
	♀	19	19.9-24.7	22.1	1.5	0.3		♀	20	18-26	22.2	1.6	0.4	7.4	
	♂♀	45	19.9-24.9	22.4	1.4	0.2		♂♀	46	18-26	22.2	1.4	0.2	6.3	
Dm	♂	10	15.8-18.8	17.3	1.1	0.4	LA-10	♂	22	22-26	24.6	1.5	0.3	6.2	
	♀	17	15.0-19.8	17.6	1.7	0.4		♀	18	21-26	24.1	1.6	0.4	6.5	
	♂♀	27	15.0-19.8	17.5	1.4	0.3		♂♀	40	21-26	24.4	1.5	0.2	6.3	
Dm/TL	♂	10	4.6-5.5	5.0	3.2	0.1	L-11	♂	25	17-28	22.5	2.1	0.4	9.5	
	♀	17	4.0-5.6	4.8	0.4	0.1		♀	20	20-27	22.9	1.8	0.4	7.8	
	♂♀	27	4.0-5.6	4.8	0.1	0.02		♂♀	45	17-28	22.7	2.0	0.3	8.6	
D-6	♂	11	13-16	14.0	1.0	0.3	L-12	♂	26	20-29	23.0	1.9	0.4	8.2	
	♀	19	13-16	14.2	1.1	0.3		♀	19	20-28	24.4	2.1	0.5	8.5	
	♂♀	30	13-16	14.1	1.1	0.2		♂♀	45	20-29	23.6	1.9	0.3	8.2	
D-7	♂	18	13-19	15.3	2.0	0.5	LTS	♂	26	160-184	175.5	6.1	1.2	3.5	
	♀	17	13-18	15.5	1.4	0.3		♀	20	160-192	176.6	8.6	1.9	4.9	
	♂♀	35	13-19	15.4	1.7	0.3		♂♀	46	160-192	175.6	7.2	1.1	4.1	
D-8	♂	18	11-23	16.3	1.1	0.3	LTS/TL	♂	26	44.3-54.8	50.2	2.9	0.6	5.7	
	♀	19	14-21	17.1	1.7	0.4		♀	20	40.7-54.0	47.9	3.8	0.8	7.8	
	♂♀	37	11-23	16.7	1.4	0.2		♂♀	46	40.7-54.8	49.2	3.0	0.5	6.6	
D-9	♂	20	13-23	18.6	2.6	0.6	LTAS	♀	19	52-62	56.6	3.1	0.7	5.4	
	♀	19	16-23	19.5	2.7	0.6		LTAS/TL	♂	19	13.7-17.5	15.3	1.2	0.3	7.8
	♂♀	39	13-23	19.0	2.6	0.4			♂	25	23-33	27.4	2.9	0.6	10.5
D-10	♂	23	20-26	22.6	2.4	0.5	P-1		♂	26	24-36	30.6	3.1	0.6	10.0
	♀	20	16-27	21.9	3.1	0.7		P-2	♂	26	24-36	30.6	3.1	0.6	10.0
	♂♀	43	16-27	22.3	2.7	0.4			P-3	♂	19	26-39	33.5	3.9	0.9
Lm	♂	18	17.7-22.8	20.6	1.4	0.3	Lm/TL			♂	18	4.6-6.7	5.8	0.5	0.1
	♀	17	19.2-25.0	21.2	1.5	0.4		♀		17	4.9-6.9	5.8	0.1	0.6	9.9
	♂♀	35	17.7-25.0	20.9	1.4	0.2		♂♀	35	4.6-6.9	5.8	0.5	0.1	9.0	

Pachycycli not well defined, muscle scars on ventrolateral areas of first trunk segment reduced, no scars on segment 8; sensory spots, 2-3 μm in diameter, situated middorsally on segments 4-5 and 12-13, laterodorsally on segments 6-13, mesially on sternal plates 4-11.

Fourth-stage juvenile (J-4, "*Habroderes*-stage," Figures 9, 10) trunk length, 224 μm ; estimated MSW-9, 72 μm , 32 percent of trunk length; estimated SW, 65 μm , 29 percent of trunk length.

Second segment as in preadult, less well developed.

Trunk segments similar to those of preadult but not well defined posteriorly so that only 10 trunk segments apparent; series of medial hairs (striations?) both dorsally and ventrally indicating area of presumptive segment 12; corresponding area of segment 13 less distinct; hairs less obvious than in


preadult, more scattered; terminal segment slightly incised ventrally with suggestions of pointed tergal extensions.

Middorsal spines on segments 6-11, D-6-10, 14-39 μm , increasing uniformly in length to segment 11, D-11, 79 μm , twice the length of D-10, extending slightly beyond terminal segment; lateral spines similar to preadult stage, 16-26 μm in length; lateral terminal spines 160 μm in length, 71.4 percent of trunk length; lateral terminal accessory spines 47 μm .


Third-stage juvenile (J-3, "*Hapaloderes*-stage," Figures 11, 12) trunk length, 200-208 μm ; estimated MSW-9, 72 μm , 34.3 percent of trunk length; estimated SW, 51 μm , 24.5 percent of trunk length.

Second segment as in J-4 stage, less well developed.


Trunk segments similar to those of J-4 stage, less



FIGURES 7-8.—*Echinoderes dujardini*, preadult (J-6) stage (USNM 53345, RH125.49), neck and trunk segments: 7, ventral view; 8, dorsal view.



FIGURES 9-10.—*Echinoderes dujardinii*, juvenile (J-4) stage (USNM 53344, RH125.53), neck and trunk segments: 9, ventral view; 10, dorsal view.



FIGURES 11-12.—*Echinoderes dujardinii*, juvenile (J-3) stage (USNM 53343, RH125.54), neck and trunk segments: 11, ventral view; 12, dorsal view.

well defined posteriorly, striations along posterior margins of segments 3-9 defining segments, stria-

tions incomplete on presumptive segment 10, restricted to midventral area, striations restricted to

mid-dorsal and midventral areas of presumptive segment 11, striations apparent on midventral area of presumptive segment 12; terminal segment blunt, without extensions.

Middorsal spines on segments 6–12 (D–12 essentially midterminal), D–6–9, 17–29 μm , increasing uniformly in length to segment 10, D–10, 39–43 μm in length, D–11, 98–109 μm in length, D–12 (midterminal) 136–170 μm in length, 60.7–85.0 percent of trunk length; lateral spines similar to J–4 stage except LA–12 absent, 18–22 μm in length; lateral terminal spines 60–64 μm , 28.8–30.0 percent of trunk length; lateral terminal accessory spines 46 μm .

MATERIAL EXAMINED.—46 adults (USNM 53342) consisting of 26 males and 20 females, 4 preadults (J–6, "*Hapaloderes*-stage") (USNM 53345), one fourth-stage juvenile (J–4, "*Hapaloderes*-stage") (USNM 53344), and 2 third-stage juveniles (J–3, "*Habroderes*-stage") (USNM 53343); col. Dr. Eugene N. Kozloff, 19 October 1973, Roscoff, Nord-Finistère, France (from the surface of sandy mud in the port).

REMARKS.—Dujardin (1851) and many subsequent authors prior to Zelinka (1928) introduced certain ambiguities into the description of *Echinoderes dujardinii*. The most common problems include the interpretation of the animal's length and the position of the spines on the trunk segments. Dujardin, for example, probably included the head and neck and may have included some juvenile stages in his total length measurements (300–550 μm). Prior to Zelinka (1928), most authors were not consistent in numbering segments; consequently, one must carefully compare the illustrations with the text when assessing the correct position of mid-dorsal and lateral spines.

The most accurate illustration of *E. dujardinii* published prior to Zelinka (1928) was that of Greeff (1869). One of the most important characteristics of this species, the presence of two lateral spines on segment 10, was noted by Dujardin (1851), but ignored by many subsequent authors.

Including *E. dujardinii*, the adults of seven species of *Echinoderes* have middorsal spines on segments 6–10 and lateral spines on segments 4, 7–12 (*E. pilosus* Lang, 1949 may have lateral spines or adhesive tubes on segment 4 although they are not included in the author's description). *Echinoderes ehlersi* Zelinka, 1913 (from Zanzibar) is 228 μm long, much smaller than *E. dujardinii*, and has significantly longer lateral terminal spines in relation

to the trunk length, 77 percent as contrasted with 41–55 percent in *E. dujardinii*. *Echinoderes ferrugineus* Zelinka, 1928 has been found with *E. dujardinii* at both Naples and Trieste (Zelinka, 1928). *Echinoderes ferrugineus* also is smaller than *E. dujardinii* and has longer lateral terminal spines relative to its trunk length, about 67–77 percent. *Echinoderes ferrugineus* differs from *E. ehlersi* in that D–10 in the former species is 55 μm , nearly twice the length of D–9; in *E. ehlersi*, the mid-dorsal spines increase uniformly from 8–14 μm in length.

Echinoderes worthingi Zelinka, 1928 also occurs with *E. dujardinii*. One specimen was collected along with *E. dujardinii* in this study and Southern (1914) found both species at Blacksod Bay, Ireland. *Echinoderes worthingi* is similar in size and other characteristics to both *E. ehlersi* and *E. ferrugineus* as noted by Zelinka (1928). *Echinoderes worthingi* most closely resembles *E. ferrugineus* in that the length of D–10, 45–50 μm , is twice that of D–9, 19–23 μm . The two species appear to be distinguishable by the shape of the terminal segment, a slightly smaller lateral terminal spine, and the presence of a prominent sensory hair (spine?) posteriorly adjacent to the lateral spine on segment 12 in *E. ferrugineus*; a similar hair exists in *E. pacificus* (Figures 28, 30).

Three additional species with the D–6–10, L–4, 7–12 spine combination are also similar to *E. dujardinii* in size. *Echinoderes pilosus* (from South Georgia Island) is poorly described, and as mentioned previously, I am including it because there is a chance that Lang (1949) overlooked the lateral spines (= adhesive tubes) on segment 4. *Echinoderes pilosus* differs from *E. pacificus* Schmidt, 1974 (from the Galapagos) in that the former has relatively longer lateral terminal spines, about 52 percent of the trunk length compared with 27–36 percent in the latter. If Lang is correct, a distinguishing feature of *E. pilosus* is the uniform size of the placids. In the other species, however, the midventral placid is much wider than the adjacent ones. Both *E. pacificus* and the remaining species, *E. kozloffii*, new species, are discussed more extensively below.

All six species mentioned in the foregoing discussion differ from *E. dujardinii* by their lack of a lateral accessory spine on segment 10.

Two additional species of *Echinoderes* share with

E. dujardinii the presence of a lateral accessory spine on segment 10. These include *E. riedli* Higgins, 1966a (from the Red Sea) and *E. newcaledoniensis* Higgins, 1967. *Echinoderes riedli* is smaller, 238 μm in length. *Echinoderes newcaledoniensis* is the only member of this genus having lateral accessory spines on segments 8–11 and is unique in possessing dorsolateral spines on segment 4. Both *E. riedli* and *E. newcaledoniensis* differ from *E. dujardinii* by having middorsal spines on segments 6, 8, and 10 only.

DISTRIBUTION.—*Echinoderes dujardinii* has been reported from 26 localities (Figure 13, also see annotations in synonymy). Most of the reports are European and include both the northern and southern coastal areas of the continent. These distribution records are probably reliable except for those from the Black Sea. Based on the evidence presented by Reinhard (1881), Băcescu et al. (1963), and Marinov (1964), I believe these records are questionable. Similarly, the reports of *E. dujardinii* from Japan (Tokiooka, 1949) and the Pacific coast of the United States (Chitwood, 1964a; Merriman and Corwin, 1972) are based on misidentifications of *E. kozloffii*, new species.

Echinoderes kozloffii, new species

FIGURES 14–21

Echinoderes.—Kozloff, 1972:121, figs. 1–18.—Higgins, 1974:514, figs. 11–16.

Echinoderes dujardinii.—Merriman and Corwin, 1973:227–243, figs. 1–13.

Echinoderes sp.—Boykin, 1974:40.

DIAGNOSIS.—Trunk length, 328–376 μm , mid-dorsal spines on segments 6–10, increasing uniformly in length; lateral spines on segments 4, 7–12, with adhesive glands at base of L-4, 7, 10 and 12, lateral spine of segment 12, 20–30 μm in length; lateral extensions of terminal tergal plate tapering evenly to point protruding about 5 μm beyond extensions of tergal plate (exclusive of prominent spines on latter).

DESCRIPTION.—Adults (Figures 14–19), trunk length, 328–376 μm ; MSW-9, 68–74 μm , 18.0–22.3 percent of trunk length; SW, 62–69 μm , 17.5–20.0 percent of trunk length.

Second segment with 16 anteriorly rounded placids; midventral placid truncate, not expanded laterally at anterior margin, distinctly larger than adjacent placids; trichoscalid plates on sides of





FIGURE 13.—*Echinoderes dujardinii*, distribution records, open circles indicate doubtful records.



FIGURES 14-15.—*Echinoderes kozloffi*, new species, holotypic female (USNM 53937), neck and trunk segments: 14, ventral view; 15, dorsal view.


midventral placid indented at anteromedial margin, posterior margin expanded laterally, four trichoscalid plates on dorsal placids much smaller than those on ventral placids.

Segments 3–12 with short hairs, pattern distinctive (Figures 14, 15); posterior border of segments 4–11 with fine pectinate fringe dorsally and ventrally; terminal segment with few hairlike processes along posterior margin; posterior margin of tergal plate deeply incised, laterally forming pointed, evenly tapered extensions mesial to base of each lateral terminal spine; sternal plates broadly rounded with spinous extensions, 15–22 μm in length (Figures 16, 18).

Middorsal spines on segments 6–10, increasing uniformly in length posteriorly, 20–45 μm ; lateral spines on segments 4 and 7–12, 24–32 μm in length; L-4, 7, 10 and 12 with adhesive gland at base; L-8 usually shorter (mean 22.9 μm) than L-7 (mean 27.5 μm) or L-9 (mean 26.1 μm); L-9–12 all similar in length, 22–32 μm ; lateral terminal spines long, 144–180 μm , 42.0–52.4 percent of trunk length; lateral

terminal accessory spines of female 46–59 μm in length; males without lateral terminal accessory spines but with penis spines in same position; anteriormost penis spine (P-1) 25–33 μm in length, mesially adjacent penis spine (P-2) 21–27 μm in length and distinctly broader, posteriorly adjacent penis spine (P-3) 35–42 μm in length, generally tending to curve mesially, crossing lateral terminal spine.

Pachycycli well developed, forming distinctive pattern at ventral midline and superficial to lateral terminal spine muscle attachments on segment 13; muscle scars on ventrolateral portion of first trunk segment almost indistinguishable, similar V-shaped scars on either side of ventral midline of segments 4–9; prominent scars on lateral margins of sternal plates 8–9, more centrally located on sternal plates of segment 10 (Figure 14); sensory spots, 2–3 μm in diameter, situated middorsally on segments 3–5 with two such spots on segment 12 (possibly one on each of segments 12 and 13), dorsolaterally on seg-



FIGURES 16–19.—*Echinoderes kozloffii*, new species, segments 12–13: 16, ventral view, lateral half, holotypic female (USNM 53337); 17, dorsal view, lateral half, holotypic female; 18, ventral view, lateral half, allotypic male (USNM 53338); 19, dorsal view, lateral half, allotypic male.

ments 3–11, ventromesially on segments 4–12, anterolaterally on ventral surface of segment 3, and possibly near lateral margin of sternal plates 5–7 (Figures 14, 15).

Morphometric data for adult specimens are shown in Table 2.

JUVENILE STAGES.—Preadult stage (J-6, "*Habroderes*-stage," Figures 20, 21) trunk length, 276–300 μm ; estimated MSW-9, 64–68 μm , 23–24 percent of trunk length; estimated SW, 62–64 μm , 22–23 percent of trunk length (estimated since tergal-sternal junctions not defined in juvenile stages).

Second segment similar to that of adult, both placids and trichoscalid plates less well developed.

Trunk segments with fewer hairs than adult although hairs as long as in adult (5–9 μm), pattern less distinctive and more variable; posterior borders of segments without pectinate fringe, with hairs (striations?) along border, group of prominent hairs (striations?) at ventral midline and second group

lateral to midline; terminal segment slightly incised dorsally and ventrally, sternal area without spinous extensions, tergal area with evenly tapered extensions, 2–4 μm in length.


Middorsal spines on segments 6–11, increasing uniformly in length, 17–62 μm (mean 27.9–34.0 μm , 10.7–12.0 percent of trunk length), D-11 extending beyond terminal tergal borders; lateral spines on segments 4, 7–12, 17–26 μm in length; L-4, 7, 10, and 12 with poorly developed adhesive glands at base; L-12 accompanied by a lateral accessory spine situated dorsally to it, slightly longer (22–26 μm) and thinner; L-4 also thinner than other lateral spines; lateral terminal spines long, 152–160 μm , 50.7–56.5 percent of trunk length; lateral terminal accessory spines 35–40 μm in length.

Pachycycli not well defined, muscle scars similar to those of preadult stage of *E. dujardinii*.

Fifth-stage juvenile (J-5, "*Habroderes*-stage") trunk length, 224–256 μm ; other measurements similar to preadult stage described above.

TABLE 2.—Measurements (μm) and indices (%) for *Echinoderes kozloffi*, new species, adults

Character	Number	Range	Mean	Standard Deviation	Standard Error	Coefficient of Variability	Character	Number	Range	Mean	Standard Deviation	Standard Error	Coefficient of Variability		
TL	♂ ♀ ♂♀.....	20 20 40	328-364 332-376 328-376	334.4 356.0 350.2	8.2 12.9 12.2	1.8 2.9 1.9	2.4 3.6 3.5	Lm/TL	♂ ♀ ♂♀.....	20 20 40	7.1-7.9 6.7-8.1 6.7-8.1	7.6 7.4 7.5	0.2 0.4 0.3	0.1 0.1 0.1	3.5 5.6 4.5
SW	♂ ♀ ♂♀.....	19 20 39	62-66 62-69 62-69	62.3 66.4 65.4	1.2 1.9 1.9	0.3 0.4 0.3	1.8 2.9 2.9	L-4 (AT)	♂ ♀ ♂♀.....	20 20 40	26-29 25-29 25-29	27.0 27.1 27.0	1.1 1.3 1.2	0.2 0.3 0.2	3.9 4.9 4.4
SW/TL	♂ ♀ ♂♀.....	19 20 39	17.7-19.8 17.5-20.0 17.5-20.0	18.5 18.6 18.6	0.5 0.8 0.7	0.1 0.2 0.1	2.5 4.4 3.6	L-7	♂ ♀ ♂♀.....	20 20 40	24-29 24-31 24-31	27.2 27.8 27.5	1.5 1.6 1.6	0.3 0.4 0.3	5.5 5.8 5.7
MSW-9	♂ ♀ ♂♀.....	19 20 39	68-72 69-74 68-74	69.5 71.8 70.7	1.3 1.8 1.9	0.3 0.4 0.3	1.9 2.5 2.7	L-8	♂ ♀ ♂♀.....	20 20 40	21-25 21-26 21-26	22.9 22.8 22.9	1.4 1.3 1.3	0.3 0.3 0.2	6.2 5.8 5.8
MSW/TL	♂ ♀ ♂♀.....	19 20 39	18.8-21.4 18.0-22.3 18.0-22.3	19.9 20.1 20.1	0.8 1.0 0.8	0.2 0.2 0.1	3.8 5.1 4.2	L-9	♂ ♀ ♂♀.....	20 20 40	24-28 23-29 23-29	26.2 26.0 26.1	0.9 1.3 1.1	0.2 0.3 0.2	3.4 5.0 4.2
Dm	♂ ♀ ♂♀.....	9 11 20	26.6-29.8 25.4-29.2 25.4-29.8	28.7 27.2 27.9	1.1 1.1 1.3	0.4 0.3 0.3	3.7 4.1 4.6	L-10	♂ ♀ ♂♀.....	20 20 40	26-30 26-29 26-30	27.7 27.5 27.6	1.4 1.2 1.3	0.3 0.3 0.2	4.9 4.3 4.6
Dm/TL	♂ ♀ ♂♀.....	9 11 20	7.6-9.1 7.1-8.1 7.1-9.1	8.3 7.6 7.9	0.5 0.3 0.5	0.2 0.1 0.1	5.4 4.0 6.5	L-11	♂ ♀ ♂♀.....	20 20 40	23-30 25-32 23-32	26.6 27.8 27.2	1.7 1.7 1.8	0.4 0.4 0.3	6.4 6.3 6.7
D-6	♂ ♀ ♂♀.....	10 14 24	20-26 20-26 20-26	23.3 22.0 22.5	2.0 1.9 2.0	0.6 0.5 0.4	8.6 8.6 8.9	L-12	♂ ♀ ♂♀.....	20 20 40	22-29 23-32 22-32	25.6 26.5 26.0	1.9 2.4 2.2	0.4 0.6 0.4	7.5 9.2 8.5
D-7	♂ ♀ ♂♀.....	13 12 25	21-27 22-26 21-27	24.4 23.4 23.9	1.8 1.2 1.6	0.5 0.3 0.3	7.2 5.0 6.5	LTS	♂ ♀ ♂♀.....	20 20 40	144-172 160-180 144-180	157.8 170.1 164.0	7.1 6.7 9.2	1.6 1.5 1.5	4.5 4.0 5.6
D-8	♂ ♀ ♂♀.....	13 15 28	23-27 22-26 22-27	25.7 24.4 25.0	1.0 1.7 1.5	0.3 0.4 0.3	4.0 6.8 6.0	LTS/TL	♂ ♀ ♂♀.....	20 20 40	40.0-50.0 43.5-52.4 42.0-52.4	45.9 47.9 46.9	2.5 2.5 2.7	0.6 0.6 0.4	5.5 5.2 5.7
D-9	♂ ♀ ♂♀.....	14 16 30	26-30 24-33 24-33	27.6 26.9 27.2	1.3 2.2 1.8	0.4 0.5 0.3	4.9 8.1 6.7	LTAS	♀ ♂♀.....	20 20	46-59 46-59	52.6 52.6	3.5 3.5	0.8 0.8	6.7 6.7
D-10	♂ ♀ ♂♀.....	20 20 40	38-45 36-43 36-45	41.6 39.6 40.6	2.5 1.8 2.4	0.6 0.4 0.4	6.1 4.6 5.9	LTAS/TL	♀ ♂♀.....	20 20	12.6-15.8 12.6-15.8	14.8 14.8	1.0 1.0	0.2 0.2	6.7 6.7
Lm	♂ ♀ ♂♀.....	20 20 40	24.7-27.7 24.0-27.7 24.0-27.7	26.0 26.4 26.2	0.8 0.9 0.9	0.2 0.2 0.1	2.9 3.5 3.3	P-1	♂ ♂♀.....	19 20	25-33 21-27	28.2 22.8	2.2 1.8	0.5 0.4	8.0 7.9
								P-2	♂ ♂♀.....	18 20	35-42 35-42	38.3 38.3	1.9 1.9	0.4 0.4	4.9 4.9



FIGURES 20-21.—*Echinoderes kozloffi*, new species, preadult (J-6) stage (USNM 53341, RH145.43), neck and trunk segments: 20, ventral view; 21, dorsal view.

Middorsal spines on segments 6–11, increasing uniformly in length, 15–72 μm (mean 33.0–39.8 μm), 14.7–15.8 percent of trunk length. Note: This latter statistic distinguishes J–5 from J–6.

HOLOTYPE.—Adult female, TL 344 μm ; North Bay, San Juan Island, Washington, USA (48°31.0'N, 123°01.0'W); 9 August 1975; col. E. N. Kozloff; USNM 53337.

ALLOTYPE.—Adult male, TL 348 μm ; as holotype; USNM 53338.

PARATYPES.—19 females and 19 males, TL 328–376 μm ; data as for holotype; USNM 53339; 5 preadult juveniles, TL 224–300 μm , 5 J–5? juveniles, 224–256 μm ; as holotype; USNM 53340; 4 females and 4 males, TL 328–364 μm ; Reid Harbor, Stuart Island, Washington, USA (48°40.0'N, 123°11.0'W); 31 July 1963; col. J. C. Boykin; USNM 53347; 3 preadult juveniles, TL 280–292 μm ; data as for above; USNM 53348.

REMARKS.—*Echinoderes kozloffii*, new species, most closely resembles *E. pacificus* from the Galapagos Islands. I consider the two sibling species. A similar sibling relationship involving the two geographic areas has been noted for two species of otoplanid turbellarians: *Philosyrtris sanjuanensis* Ax and Ax, 1967 (from San Juan Island, Washington) is a sibling species of *P. santacruzensis* Ax and Ax, 1974 (from the Galapagos Islands) according to these authors. *Echinoderes kozloffii* is similar to *E. pacificus* in total length, standard width, maximum sternal width, and general appearance. The range of the mean lengths of middorsal spines in *E. kozloffii* (25.4–29.8 μm) and those of *E. pacificus* (46.5–56.5 μm) is one of the significant differences between the two species. Other differences include the longer (25–29 μm) lateral spine on segment 4 in *E. kozloffii*, which contrasts with the shorter (12–17 μm) lateral spine on segment 4 in *E. pacificus*; in the former species the lateral terminal spines, 144–180 μm , 42.0–52.4 percent of the trunk length, are longer than those of the latter species, 90–118 μm , 27.0–36.0 percent of the trunk length. The lateral terminal accessory spines of *E. kozloffii* are slightly longer (46–59 μm) than those of *E. pacificus* (34–41 μm).

The relatively long (22–30 μm) lateral spine on segment 12 of *E. kozloffii* is a particularly noticeable feature that contrasts with the short (12–17 μm), blunt lateral spine on segment 12 of *E. pacificus*. The posterior margins of both tergal and

sternal plates of the terminal segment distinguish the two species. The mesial margins of the terminal sternal extensions of *E. pacificus* are interrupted and beset with a series of prominent hairs (Figure 28); in *E. kozloffii* these margins are even and have a less obvious series of hairs. Both species have hairs along the mesial margins of the terminal sternal extensions but, again, those of *E. pacificus* are more distinct, each hair associated with a steplike interruption of the margin. Also, the sternal extensions of both species have a spinous process. The processes in *E. pacificus*, however, are half the length (8–12 μm) of those in *E. kozloffii* (17–25 μm). *Echinoderes pacificus* is discussed later in this paper.

Echinoderes kozloffii is similar to *E. pilosus*. As noted above, *E. pilosus* is not well described but has the same general trunk length and spination formula as both *E. kozloffii* and *E. pacificus*. *Echinoderes pilosus* appears to differ from *E. kozloffii* in only two principal characters: *E. pilosus* has shorter lateral terminal accessory spines, about 35 μm , about 8 percent of the trunk length, and the midventral placid is the same width as adjacent placids. This latter character is unique within the genus.

Echinoderes kozloffii is sympatric with *E. pennaki* Higgins, 1960 but differs in that the latter species lacks a lateral spine on segment 12, has conspicuously longer lateral spines (mean 58.1 μm , 14 percent of the trunk length), a prominent ventral spinous fringe on the posterior border of the first trunk segment, lacks spinous processes on the terminal sternal plates, and has slightly longer, more pointed terminal tergal extensions. Other differences are mentioned in the discussion of *E. pennaki* that follows.


ETYMOLOGY.—This species is named in honor of Eugene N. Kozloff, a fellow student of the Kinorhyncha.

Echinoderes pennaki Higgins

FIGURES 22–25

Echinoderes pennaki Higgins, 1960:86; 1961:81.—Chitwood, 1964b:3.—Higgins, 1964a:246; 1964b:479.—Kirsteuer, 1964:389.—Higgins, 1966a:120; 1966b:519; 1967:75.—Schmidt, 1974:13.—Boykin, 1974:40.

REDESCRIPTION.—Holotypic adult female (Figures 22–25), trunk length, 404 μm ; MSW–8, 62 μm , 15




FIGURES 22-23.—*Echinoderes pennaki*, holotypic female (USNM 29746), neck and trunk segments; 22, ventral view; 23, dorsal view.

percent of trunk length; SW, 56 μm , 14 percent of trunk length.

Second segment with 16 anteriorly rounded placids; midventral placid slightly truncate, distinctly larger than adjacent placids; trichoscalid plates on sides of midventral placid indented slightly on anterior margin, short and broad, trichoscalid plates on dorsal surface similar but equally as broad as long.

Segments 3–12 with prominent hairs (10–15 μm), pattern distinctive (Figures 22–23); posterior border of segments 3–11 with pectinate fringe both dorsally and ventrally, fringe on ventral surface of segment 3 almost spinose, more finely pectinate on remaining segments, slightly evident near posterior border of 12th sternal plates; terminal segment with minute hairs on border of tergal and sternal plates, posterior margin of tergal plate deeply incised forming pointed extensions mesial to base of each lateral terminal spine, sternal plates broadly rounded without spinous extensions (Figures 24, 25).



FIGURES 24–25.—*Echinoderes pennaki*, segments 12–13 of holotypic female (USNM 29746): 24, ventral view, lateral half; 25, dorsal view, lateral half.

Middorsal spines on segments 6–10, increasing uniformly in length, 48–70 μm , mean length 59.6 μm , 15 percent of trunk length; lateral spines on segments 7–11, L-4 (=adhesive tube) absent although a scar, or perhaps a sensory spot, located slightly mesial to usual position of L-4; L-7 shorter (14 μm) than remaining lateral spines (32–36 μm) subequal in length, mean length of lateral spines 29.8 μm , 7 percent of trunk length; lateral terminal spines long, 156 μm , 39 percent of trunk length; lateral terminal accessory spines 52 μm in length.

Pachycycli well developed, forming distinctive pattern at ventral midline and superficial to attachment of lateral terminal spine muscles on segment 13; small muscle scars visible near midventral line of segments 3–4, 7, 11, and 12, more prominent scars dorsolaterally on segments 3, 5–12; sensory spots situated ventrolaterally on segment 3 and possibly on segment 4 (possibly adhesive tube openings?).

MATERIAL EXAMINED.—4 adult females including holotype (USNM 29746) and 3 paratypes (USNM 29747); col. P. L. Illg, 16 July 1958, East Sound, Orcas Island, Washington, USA, from a depth of 32 meters.

REMARKS.—Eighteen years of research on the Kinorhyncha has prompted me to reexamine my first species descriptions with better experience to guide me. The combination of this experience and better optical instruments has allowed me to illustrate and describe *E. pennaki* more accurately. For example, the range of "total length" given in my original description (Higgins, 1960) is "390–430 μm (taken along dorsal surface between anterior edge of second zonite and posterior edge of zonite 13)." More standard measurements are now taken from the anterior edge of segment 3 to the posterior edge of segment 13. Similarly, the maximum width of 80–90 μm originally given for this species differs from the more precise and standardized measurement of maximum sternal width as noted in previous publications (Higgins, 1967).

Echinoderes pennaki most closely resembles *E. remanei* (Blake, 1930) redescribed by me (Higgins, 1964a). The two species differ in size: *E. pennaki* is larger (380–404 μm) than *E. remanei* 282–358 μm) yet both have the same spination formula (D-6–10, L-4? 7–11). Both have a prominent spinous fringe on the posteroventral border of segment 3; this feature is more prominent in *E. remanei* and also occurs on the fourth segment. The midventral

placid of *E. remanei* is expanded laterally along its anterior margin, and the border of the terminal sternal plates is pointed, not rounded as in *E. penaki* (Figure 24). Our understanding of both species would benefit from an expanded study of their taxonomic characters based on larger numbers of individuals in a given sample.

Echinoderes pacificus Schmidt

FIGURES 26–31

Echinoderes pacificus Schmidt, 1974:1.

This species has been described with considerable accuracy by my colleague, Dr. Peter Schmidt (1974). In studying the holotype and paratype material sent to me by him, however, certain additional information was revealed by mounting specimens in Hoyer's and observing them with differential interference contrast optics. I have reillustrated *E. pacificus* (Figures 26–31) in order to facilitate the comparison of this species with others that might be confused with it.

Segments 3–12 are covered with short hairs in a distinctive pattern (Figures 26, 27). A fine pectinate fringe is evident ventrally on the posterior border of segments 4–13 although it is much less distinct on segment 12; segments 6–13 exhibit this fringe dorsally.

As noted in the discussion of *E. kozloffi*, the short (12–17 μm) lateral spine on segment 12 of *E. pacificus* is diagnostic. This spine tends to curve away from the trunk. Slightly posterior to it, at the junction of segments 12 and 13, there is a sensory hair (Figures 28–31) not reported in the original description.

Small muscle scars (2–3 μm in diameter), similar to sensory spots, are present anterolaterally on the ventral surface of the first trunk segment. Narrow, slitlike scars occur posterior to the sensory spots on sternal plates 7–9. More distinctive muscle scars (Figure 26) occur near the lateral margin of the sternal plates of segment 9; similar scars are

situated near the sensory spots of the 10th sternal plates. Sensory spots occur on either side of the ventral midline of segments 4–12. Three spots are present on the dorsal surface of the first trunk segment; only the median spot persists on the following two trunk segments (segments 4–5). Sensory spots on the segments bearing middorsal spines are on both sides of the midline. Two middorsal sensory spots appear to be on segment 12; one of these may be associated with segment 13.


MATERIAL EXAMINED.—Holotypic male, TL 372 μm (USNM 53335); 8 paratypic females and 7 paratypic males (USNM 53336); col. P. Schmidt, July–September 1972, Academy Bay, Station IX, 5c, upper subtidal sediments.

REMARKS.—As noted in the discussion of *E. kozloffi*, this species closely resembles *E. pacificus*. Both species share the same spination formula with *E. pilosus*, *E. ehlersi*, *E. worthingi*, and *E. ferrungineus* (*E. dujardinii* is similar but has a lateral accessory spine on segment 10).


Echinoderes ehlersi, *E. worthingi*, and *E. ferrungineus* are smaller, 210–260 μm trunk length, although the latter species shares with *E. pacificus* the presence of a sensory hair posterior to the lateral spine on segment 12. *Echinoderes pacificus* is most easily distinguished from all species having the same spination formula by its short lateral spine on segment 12 and the prominent border of hairs on the terminal sternal plates.

Echinoderes brevicaudatus, substitute name for *E. brevispinosus* Higgins

During the course of the present investigation, I discovered that the name *E. brevispinosa* was first used by Metschnikoff (1869:190), corrected to *E. brevispinosus* by Panceri (1876:4), and synonymized with *E. dujardinii* by Zelinka (1928:228). Since I inadvertently applied this preoccupied name to a species from the Red Sea (Higgins, 1966a), I now propose that it be replaced by the substitute name, *Echinoderes brevicaudatus*.



FIGURES 26-27.—*Echinoderes pacificus*, paratype female (USNM 53336 RH162.1), neck and trunk segments: 26, ventral view; 27, dorsal view.



FIGURES 28-31.—*Echinoderes pacificus*, segments 12-13: 28, ventral view, lateral half, paratypic female (USNM 53336, RH162.1); 29, dorsal view, lateral half of same paratypic female; 30, ventral view, lateral half, paratypic male (USNM 53336, RH162.9); 31, dorsal view, lateral half of same paratypic male.

Literature Cited

- Abe, Y.
1930. Das Vorkommen von *Echinoderes* in den japanischen Gewässern. *Journal of Science of the Hiroshima University*, series B, 1(1):39-44, 2 figures, 1 plate.
- Ax, P., and R. Ax
1967. Turbellaria Proseriata von der Pazifikküste der USA (Washington), I: Otoplanidae. *Zeitschrift für Morphologie und Ökologie der Tiere*, 61:215-254, 18 figures.
1974. Interstitielle Fauna von Galapagos, V: Otoplanidae (Turbellaria, Proseriata). *Mikrofauna des Meeresbodens*, 27:1-28, 11 figures.
- Băcescu, M., E. Dumitrescu, A. Marcus, G. Paladian, and R. Mayer
1963. Données quantitatives sur la faune pétricole de la mer noire à Agigea (Secteur Roumain), dans les conditions spéciales de l'année 1961. *Travaux du Muséum d'Histoire Naturelle "Grigore Antipa,"* 4:131-155, 3 figures, 1 plate.
- Blake, C. H.
1930. Three New Species of Worms Belonging to the Order Echinodera. *Biological Survey of the Mount Desert Region*, 4:3-10, 8 figures.
- Boykin, J. C.
1974. Phylum Kinorhyncha. Pages 39-41 in E. N. Kozloff, *Keys to the Marine Invertebrates of Puget Sound, the San Juan Archipelago, and Adjacent Regions*. vii + 266 pages, 1 unnumbered figure. Seattle: University of Washington Press.
- Carus, J. V.
1885. *Prodromus faunae Mediterraneae*. Volume 1, xi + 525 pages. Stuttgart: E. Schweizerbart'sche Verlagshandlung.
- Chitwood, B. G.
1951. *Echinoderella steineri*, new species (Scolecida, Echinodera). *Texas Journal of Science*, 3(1):113-114, 1 unnumbered figure.
1964a. European Kinorhynchs from Tomales Bay, California. Abstract 6, pages 2-3 in *Abstracts of Contributed Papers, Forty-fifth Annual Meeting of the Western Society of Naturalists*.
1964b. The Intertidal Occurrence of *Echinoderes pennaki* (Kinorhyncha) in the Straits of Juan de Fuca, Washington. Abstract 7, page 3 in *Abstracts of Contributed Papers, Forty-fifth Annual Meeting of the Western Society of Naturalists*.
- Claparède, E.
1863. *Beobachtungen über Anatomie und Entwicklungsgeschichte wirbelloser Tiere an der Küste der Normandie angestellt*. 120 pages, 18 plates. Leipzig: Wilhelm Engelmann.
- Dujardin, F.
1851. Sur un petit animal marin, *l'Echinodère*, formant un type intermédiaire entre les Crustacés et les Vers. *Annales des Sciences Naturelles, Zoologie*, series 3, 15:158-160, 1 plate.
- Gosse, P. H.
1864. The Natural History of the Hairy-backed Animalcules. *The Intellectual Observer: Review of Natural History, Microscopic Research, and Recreative Science*, 5:387-406, 2 plates.
- Greeff, R.
1869. Untersuchungen über einige merkwürdige Formen des Anthropoden- und Wurm-Typus. *Archiv für Naturgeschichte*, 35(1):71-100, 4 plates.
- Grobben, K.
1905. Kinorhyncha. Pages 335-336 in C. Claus and K. Grobben, *Lehrbuch der Zoologie*. x + 955 pages, 996 figures. Marburg: N. G. Elwert'sche Verlagshandlung.
- Hartog, M.
1896. Rotifera, Gastrotricha, and Kinorhyncha. Pages 197-238 (15 figures) in volume 2 in S. F. Harmer and A. E. Shipley, editors, *The Cambridge Natural History*. xii + 560 pages, 257 figures. London: MacMillan and Co., Ltd.
- Higgins, R. P.
1960. A New Species of *Echinoderes* (Kinorhyncha) from Puget Sound. *Transactions of the American Microscopical Society*, 79(1):85-91, 1 figure.
1961. Three New Homalorhage Kinorhynchs from the San Juan Archipelago, Washington. *Journal of the Elisha Mitchell Scientific Society*, 77(1):81-88, 14 figures.
1964a. Redescription of the Kinorhynch *Echinoderes remanei* (Blake, 1930) Karling, 1954. *Transactions of the American Microscopical Society*, 83(2):243-247, 3 figures.
1964b. Three New Kinorhynchs from the North Carolina Coast. *Bulletin of Marine Science of the Gulf and Caribbean*, 14(3):479-493, 18 figures.
1966a. Faunistic Studies in the Red Sea (in Winter, 1961-1962), Part II: Kinorhynchs from the Area of Al-Ghardaqa. *Zoologisches Jahrbücher, Systematik, Oekologie und Geographie der Tiere*, 93:118-126, 9 figures.
1966b. *Echinoderes arlis*, a New Kinorhynch from the Arctic Ocean. *Pacific Science*, 20(4):518-520, 2 figures.
1967. The Kinorhyncha of New Caledonia. In *Expedition Française sur Recifs Coralliens de la Nouvelle Calédonie*, 2:75-90, 12 figures.
1968. Taxonomy and Postembryonic Development of the Cryptorhagae, a New Suborder for the Mesopsam-

- mic Kinorhynch Genus *Cateria*. *Transactions of the American Microscopical Society*, 87(1):21-39, 25 figures.
- 1969a. Indian Ocean Kinorhyncha, 1: *Condyloderes* and *Sphenoderes*, New Cyclorhagid Genera. *Smithsonian Contributions to Zoology*, 14:1-13, 23 figures.
- 1969b. Indian Ocean Kinorhyncha, 2: Neocentrophyidae, a New Homalorhagid Family. *Proceedings of the Biological Society of Washington*, 87(7):113-128, 5 figures.
1971. A Historical Overview of Kinorhynch Research. Pages 25-31 in N. C. Hulings, editor, *Proceedings of the First International Conference on Meiofauna. Smithsonian Contributions to Zoology*, 76:1-205, 1 figure.
1974. Kinorhyncha. Chapter 11, pages 507-518 in volume 1 in A. C. Giese and J. S. Pearse, editors, *Reproduction of Marine Invertebrates*. xi + 546 pages, 18 figures. New York: Academic Press.
- d'Hondt, J. L.
1973. Contribution a l'Étude de la Microfaune Interstitielle des Plages de l'Ouest Algérien. *Vie et Milieu*, 23(2):227-241, 2 figures.
- Hyman, L. H.
1951. *The Invertebrates*. Volume 3, vi + 572 pages, 223 figures. New York: McGraw Hill.
- Johnston, T. H.
1938. Report on the Echinoderida. In *Scientific Reports of the Australasian Antarctic Expedition, 1911-1914*, series C, 10(7):1-13, 7 figures.
- Karling, T. G.
1954. *Echinoderes levanderi* n. sp. (Kinorhyncha) aus der Ostsee. *Arkiv för Zoologi*, series 2, 7(10):189-192, 6 figures.
- Kirsteuer, E.
1964. Zur Kenntnis der Kinorhynchen Venezuelas. *Zoologischer Anzeiger*, 173(6):388-393, 2 figures.
- Kozloff, E. N.
1972. Some Aspects of Development in *Echinoderes* (Kinorhyncha). *Transactions of the American Microscopical Society*, 91(2):119-130, 18 figures.
- Lang, K.
1949. Echinoderida. In N. H. Odhner, editor, *Further Zoological Results of the Swedish Antarctic Expedition, 1901-1903*, 4(2):1-22, 8 figures.
- Leuckart, R.
1854. Bericht über die Leistungen in der Naturgeschichte der niederen Thiere während der Jahre 1848-1853. *Archiv für Naturgeschichte*, 20(2):289-473.
1869. Bericht über die wissenschaftlichen Leistungen in der Naturgeschichte der niederen Thiere während der Jahre 1868-1869. *Archiv für Naturgeschichte*, 35(2):207-344.
- Levander, K. M.
1900. Über das Herbst- und Winterplankton im finnischen Meerbusen und in der Alands-See, 1898. *Acta Societatis pro Fauna et Flora Fennica*, 18(5):19-20.
- Lou, T. H.
1934. Sur la présence d'un nouveau Kinorhynque à Tchefou: *Echinoderes tchefouensis* sp. nov. *Contributions du Laboratoire de Zoologie, Académie Nationale de Peiping*, 1(4):1-9, 1 plate.
- Ludwig, H.
1886. *Dr. Johannes Leunis Synopsis der Tierkunde*. Volume 2, 1231 pages, 1160 figures. Hannover: Hahn'sche Buchhandlung.
- Marinov, T.
1964. On the Microzoobenthos Fauna of the Black Sea (Kinorhyncha and Halacaridae) [in Bulgarian with English summary]. *Bulletin of the Institute of Fish Culture and Fisheries, Varna KH*, 4:61-71, 11 figures.
- McIntyre, A. D.
1962. The Class Kinorhyncha (Echinoderida) in British Waters. *Journal of the Marine Biological Association of the United Kingdom*, 42:503-509, 2 figures.
- Merriman, J. A., and H. O. Corwin
1973. An Electron Microscopical Examination of *Echinoderes dujardini* Claparède (Kinorhyncha) [sic]. *Zeitschrift für Morphologie der Tiere*, 76:227-242, 13 figures.
- Metschnikoff, E.
1865. Ueber einige wenig bekannte niedere Thierformen. *Zeitschrift für wissenschaftliche Zoologie*, 15:450-461, 1 plate.
1869. Bemerkungen über Echinoderes: Melanges biologiques 7. *Bulletin de l'Académie des Sciences de Saint Pétersbourg*, 4:190-194.
- Möbius, K.
1884. Nachtrag zu dem im Jahre 1873 erschienenen Verzeichniss der wirbellosen Thiere der Ostsee. Pages 1-10 in volume 4 in *Bericht der Kommission zur wissenschaftlichen Untersuchung der deutschen Meere, Berlin*.
1887. Systematische Darstellung der Thiere des Plankton gewonnen in der westlichen Ostsee Fahrt von Kiel in den atlantischen ocean bis jenseit der Hebriden. Pages 111-124 in volume 5 in *Bericht der Kommission zur Untersuchung der deutschen Meer, Berlin*.
- Moore, P. G.
1973. *Campyloderes macquariae* Johnston, 1938 (Kinorhyncha: Cyclorhagida) from the Northern Hemisphere. *Journal of Natural History*, 7:341-354, 4 figures, 1 plate.
- Nyholm, K.-G.
- 1947a. Contributions to the Knowledge of the Postembryonic Development in Echinoderida Cyclorhagae. *Zoologiska bidrag från Uppsala*, 25:423-428, 4 figures.
- 1947b. Studies in the Echinoderida. *Arkiv für Zoologi*, 39A(14):1-36, 22 figures, 2 plates.
- Omer-Cooper, J.
1957. Deux nouvelles especes de Kinorhyncha en provenance de l'Afrique du Sud. *Bulletin Mensuel de la Société Linnéenne de Lyon*, 26(8):213-216, 3 figures.
- Pagenstecher, H. A.
1875. *Echinoderes Sieboldii*. *Zeitschrift für wissenschaftliche Zoologie*, supplement, 25(S):117-123, 1 plate.
1877. *Allgemeine Zoologie*. Part 2. vi + 528 pages, 206 figures. Berlin: Wiegandt, Hempel & Parey.
- Panceri, P.
1876. Osservazioni intorno a nuove forme di Vermi Nema-

- todi marini. *Atti della Reale Accademia delle Scienze Fisiche e Matematiche*, 7(10):1-9, 1 plate.
- Perrier, R.
1893. *Eléments d'Anatomie comparée*. 1208 pages. Paris: Librairie J.-B. Baillière et Fils.
- Reinhard, W.
1881. Über *Echinoderes* und *Desmoscolex* der Umgegend von Odessa. *Zoologischer Anzeiger* 4(97):588-592.
1885. *Kinorhyncha: Their Anatomical Structure and Position in the System* [in Russian]. 101 pages, 5 plates. Kharkov: University Printing Office.
1887. Kinorhyncha (*Echinoderes*), ihr anatomischer Bau und ihre Stellung im System. *Zeitschrift für wissenschaftliche Zoologie*, 45:401-467, 3 plates.
- Remane, A.
1928. Kinorhyncha. In G. Grimpe and E. Wagler, editors, *Die Tierwelt der Nord- und Ostsee*, 7(d₂):57-84, 20 figures.
1936. Gastrotricha und Kinorhyncha. Number 2 in part 1 in section 2 of volume 4 in H. G. Bronn, editor, *Klassen und Ordnungen des Tierreichs*. vi + 243-385, 297 figures.
- Schepotieff, A.
1907a. Zur Systematik der Nematoideen. *Zoologischer Anzeiger*, 31(5-6):132-161, 25 figures.
1907b. Die Echinoderiden. *Zeitschrift für wissenschaftliche Zoologie*, 88(2):291-326, 4 plates.
- Schmidt, P.
1974. Interstitielle Fauna von Galapagos, X: Kinorhyncha. *Mikrofauna des Meeresbodens*, 43:1-15, 2 figures.
- Schultze, M.
1853. Ueber *Chaetonotus* und *Ichtydium* (Ehrb.) und eine neue verwandte Gattung *Turbanella*. *Archiv für Anatomie, Physiologie und wissenschaftliche Medicin*. 1853:241-254, 1 plate.
- Southern, R.
1914. Nemathelminths, Kinorhyncha, and Chaetognatha. Part 54 in Clare Island Survey. *Proceedings of the Royal Irish Academy*, 31:1-80, 12 plates.
- Timm, R. W.
1958. Two New Species of *Echinoderella* (Phylum Kinorhyncha) from the Bay of Bengal. *Journal of the Bombay Natural History Society*, 55(1):107-109, 2 figures.
- Tokioka, T.
1949. Notes on *Echinoderes* Found in Japan. *Publications of the Seto Marine Biological Laboratory*, 1(2):67-69, 1 figure.
- Zaneveld, J. S.
1938. Marine Gastrotricha and Kinorhyncha from Scheveningen. *Zoologische Mededeelingen*, 20:257-262.
- Zelinka, C.
1896. Demonstration von Tafeln der Echinoderes-Monographie. *Verhandlungen der Deutschen Zoologischen Gesellschaft*, 6:197-199.
1907. Zur Kenntnis der Echinoderen. *Zoologischer Anzeiger*, 32(5):130-136.
1908. Zur Anatomie der Echinoderen. *Zoologischer Anzeiger*, 33(19-20):629-647, 11 figures.
1912. Die Spermatozoen der Echinoderen und ihre Genese. Pages 520-527 in *Verhandlungen des VIII Internationalen Zoologen-Kongress zu Graz vom 15.-20. August 1910*. 10 figures.
1913. Die Echinoderen der Deutschen Südpolar-Expedition 1901-1903. In *Deutschen Südpolar-Expedition 1901-1903*, 14(Zoology 6):419-436, 1 plate.
1928. *Monographie der Echinodera*. iv + 396 pages, 73 figures, 27 plates. Leipzig: Wilhelm Engelmann.

REQUIREMENTS FOR SMITHSONIAN SERIES PUBLICATION

Manuscripts intended for series publication receive substantive review within their originating Smithsonian museums or offices and are submitted to the Smithsonian Institution Press with approval of the appropriate museum authority on Form SI-36. Requests for special treatment—use of color, foldouts, casebound covers, etc.—require, on the same form, the added approval of designated committees or museum directors.

Review of manuscripts and art by the Press for requirements of series format and style, completeness and clarity of copy, and arrangement of all material, as outlined below, will govern, within the judgment of the Press, acceptance or rejection of the manuscripts and art.

Copy must be typewritten, double-spaced, on one side of standard white bond paper, with 1 $\frac{1}{4}$ " margins, submitted as ribbon copy (not carbon or xerox), in loose sheets (not stapled or bound), and accompanied by original art. Minimum acceptable length is 30 pages.

Front matter (preceding the text) should include: **title page** with only title and author and no other information, **abstract page** with author/title/series/etc., following the established format, **table of contents** with indents reflecting the heads and structure of the paper.

First page of text should carry the title and author at the top of the page and an unnumbered footnote at the bottom consisting of author's name and professional mailing address.

Center heads of whatever level should be typed with initial caps of major words, with extra space above and below the head, but with no other preparation (such as all caps or underline). Run-in paragraph heads should use period/dashes or colons as necessary.

Tabulations within text (lists of data, often in parallel columns) can be typed on the text page where they occur, but they should not contain rules or formal, numbered table heads.

Formal tables (numbered, with table heads, boxheads, stubs, rules) should be submitted as camera copy, but the author must contact the series section of the Press for editorial attention and preparation assistance before final typing of this matter.

Taxonomic keys in natural history papers should use the alined-couplet form in the zoology and paleobiology series and the multi-level indent form in the botany series. If cross-referencing is required between key and text, do not include page references within the key, but number the keyed-out taxa with their corresponding heads in the text.

Synonymy in the zoology and paleobiology series must use the short form (taxon, author, year:page), with a full reference at the end of the paper under "Literature Cited." For the botany series, the long form (taxon, author, abbreviated journal or book title, volume, page, year, with no reference in the "Literature Cited") is optional.

Footnotes, when few in number, whether annotative or bibliographic, should be typed at the bottom of the text page on which the reference occurs. Extensive notes must appear at the end of the text in a notes section. If bibliographic footnotes are required, use the short form (author/brief title/page) with the full reference in the bibliography.

Text-reference system (author/year/page within the text, with the full reference in a "Literature Cited" at the end of the text) must be used in place of bibliographic footnotes in all scientific series and is strongly recommended in the history and technology series: "(Jones, 1910:122)" or ". . . Jones (1910:122)."

Bibliography, depending upon use, is termed "References," "Selected References," or "Literature Cited." Spell out book, journal, and article titles, using initial caps in all major words. For capitalization of titles in foreign languages, follow the national practice of each language. Underline (for italics) book and journal titles. Use the colon-parentheses system for volume/number/page citations: "10(2):5-9." For alinement and arrangement of elements, follow the format of the series for which the manuscript is intended.

Legends for illustrations must not be attached to the art nor included within the text but must be submitted at the end of the manuscript—with as many legends typed, double-spaced, to a page as convenient.

Illustrations must not be included within the manuscript but must be submitted separately as original art (not copies). All illustrations (photographs, line drawings, maps, etc.) can be intermixed throughout the printed text. They should be termed **Figures** and should be numbered consecutively. If several "figures" are treated as components of a single larger figure, they should be designated by lowercase italic letters (underlined in copy) on the illustration, in the legend, and in text references: "Figure 9b." If illustrations are intended to be printed separately on coated stock following the text, they should be termed **Plates** and any components should be lettered as in figures: "Plate 9b." Keys to any symbols within an illustration should appear on the art and not in the legend.

A few points of style: (1) Do not use periods after such abbreviations as "mm, ft, yds, USNM, NNE, AM, BC." (2) Use hyphens in spelled-out fractions: "two-thirds." (3) Spell out numbers "one" through "nine" in expository text, but use numerals in all other cases if possible. (4) Use the metric system of measurement, where possible, instead of the English system. (5) Use the decimal system, where possible, in place of fractions. (6) Use day/month/year sequence for dates: "9 April 1976." (7) For months in tabular listings or data sections, use three-letter abbreviations with no periods: "Jan, Mar, Jun," etc.

Arrange and paginate sequentially EVERY sheet of manuscript—including ALL front matter and ALL legends, etc., at the back of the text—in the following order: (1) title page, (2) abstract, (3) table of contents, (4) foreword and/or preface, (5) text, (6) appendixes, (7) notes, (8) glossary, (9) bibliography, (10) index, (11) legends.

