

acute or even short pedicellate at base. In the hind wings the apical part is faintly fumose, and the pterostigma a little darker.

Expanse 38 mm.

Two specimens from Wellington, British Columbia (Theo. Bryant).

---

## TWO NEW FORMS OF *ÆNEIS* Huebner.

By HARRISON G. DYAR.

Mr. Merritt Cary collected a number of specimens of *Æneis* (*Chionobas*) in British America in 1903, and among them are two forms that do not agree with any known to me.

### *Æneis caryi*, n. var.

Gray brown above, washed with dark ferruginous red submarginally on fore wings, over most of surface of hind wings; two large black ocelli on fore wings above veins 2 and 5, respectively, one on the hind wings near anal angle, the three minutely white pupilled and repeated below. Fore wings below reddish on the disk and inner margin; hind wings contrastingly marbled in black and white, the median band strongly white edged.

One ♂, Smith Landing, Athabasca, June 13, 1903.

*Type*.—No. 8046, U. S. National Museum.

Near *norna* Thunberg, but the red color of the wings is much darker and more rust colored than in this form or in *katahdin* Newcomb.

### *Æneis nahanni*, n. sp.

Blackish above, washed with ferruginous brown, the veins darker, the markings of underside showing. A small ocellus or none above vein 5 on fore wings, two to five on hind wings, the one above vein 5 largest, the rest small or absent. Hind wings below black and white, coarsely strigose, somewhat as in *uhleri* Reakirt and *varuna* Edwards, but much more densely, the white of the wing being largely obscured. Median band weakly indicated; ocelli black with white pupils; fore wings shaded with red over the disk.

One ♂, one ♀, Nahanni Mts., Mackenzie, 2,500 feet, July 16, 1903.

*Type*.—No. 8047, U. S. National Museum.

Differs entirely in color from *uhleri*, and in the coarseness of the markings below from the forms of *norna*.

---