

it has a shorter wing and shorter tarsus, though its entire length is greater; it has also a square tail instead of a forked one. It measures seven inches and a half entire length; from the carpus to the end of the wing, five inches and three quarters; tarsus, three-quarters of an inch. I have called it *Thalassidroma castro*, as I am not aware that it has ever been described before." (A Sketch of Madeira, 1851, 123.) The specific name is derived from that of "*Roque de Castro*," given by the natives. The type locality is the Desertas Islands, near Madeira. — CHAS. W. RICHMOND, *Washington, D. C.*

Pelecanus occidentalis vs. *P. fuscus*. — Although "*Pelecanus fuscus*," credited to Linnæus, Syst. Nat., ed. 12, 1766, 215, has stood in our 'Check-List' since 1886, a glance at the first-mentioned work will convince any one that Linnæus used no such name. He divides *Pelecanus onocrotalus* into two varieties, α . *orientalis*, and β . *occidentalis*; under the latter are cited the "*Onocrotalus s. Pelecanus fuscus*" of Sloane's 'Jamaica,' the "*Onocrotalus americanus*" of Edwards, the "*Pelecanus subfuscus, gula distensili*" of Brown's 'Jamaica,' etc. The habitat of β is given as "America," and the references belong mainly to the Brown Pelican of Eastern North America. Varietal names, as used by Linnæus, were italicized and designated by a Greek letter instead of a separate number, but all such names were binomial marginal ones, and ought to be recognized. If ornithologists accept this view our Brown Pelican should stand as *Pelecanus occidentalis*; otherwise the *P. fuscus* of the 'Check-List' must be credited to Gmelin. — CHAS. W. RICHMOND, *Washington D. C.*

Old Squaw (*Clangula hyemalis*) in Indiana. — A few records have been given of individuals taken in the State, and in all instances they were probably blown inland by severe storms off Lake Michigan, where they are usually abundant in the winter season.

On Feb. 12, 1899, during intensely cold weather, a flock of thirteen was killed at English Lake, Ind., some thirty-five miles directly south of Lake Michigan. There was no open water, except a small space, some thirty yards square, where the ice had been cut and taken out for storing, and here the flock suddenly alighted. They were evidently in an exhausted condition, hunting for open water, as they paid no attention to twenty or thirty men working around the hole and floating on the ice, and only dove when struck at with pike poles. A gun was soon procured, and the whole flock dispatched, and a male specimen was sent to me. The following morning, February 13, three more Ducks of this species attempted to alight in the same hole, which had been kept open by the ice cutters, but a hungry Bald Eagle, who has a nest a half mile distant, stooped to them, without success however, and they continued a hurried flight over the frozen marshes.