

occur in this plateau region. Perhaps the Sparrow Hawk of the plateau region of Texas, in which Tom Green County is situated, will prove to be distinct from the eastern bird when more carefully studied.—G. H. RAGSDALE, *Gainsville, Texas.*

Great Gray Owl in Worcester County, Mass.—A Great Gray Owl (*Scotiaptex cinerea*) was killed in Princeton, Feb. 28, by E. T. Whitaker, a member of the Worcester Sportsmen's Club. The day was severely cold, the thermometer registering 4° below zero. This is the first record for this species in this County.—GEORGE B. CHURCHILL, *Worcester, Mass.*

Acadian Owl (*Nyctala acadica*) at Washington, D. C.—The Acadian Owl has always been considered of extremely rare occurrence at the Capital; in fact until the present winter but five specimens were known from this locality. The first of these was taken by Mr. Drexler some years ago, which was followed by an occasional capture from time to time. On December 12, 1890, an adult male was taken by Mr. Walter B. Barrows three miles east of the city at Brookland. This served as a stimulus for close search, and on January 4, 1891, reward came in the shape of three females taken by Mr. J. D. Figgins and myself. These were mostly in a dense thicket of pines, less than half an acre in extent, and the peculiarity especially noticeable was the fact that not a single male was to be found, although we searched the woods again and again, as well as neighboring patches of pine. On Feb. 4 or 5, 1891, one flew into the Smithsonian building where it was captured alive, and now serves as one of the attractions of the 'Zoo', making a total of five taken the past winter, or as many as all previous records taken together.

It would appear from this that the species is much more common than ordinarily supposed, but if such be the case it is strange that no more have been taken, as scarcely a week has passed that Mr. Figgins and myself have not spent at least one day in the woods, and always with an eye open for *Nyctala*, but since the record of January 4, not a feather has rewarded our efforts, and it must, I think, still be considered as one of the rarer birds of the district.—EDWIN M. HASBROUCK, *Washington, D. C.*

Occurrence of the Groove-billed Ani at Jupiter Inlet, Florida.—While at Palm Beach, Lake Worth, Fla., my friend, Mr. Franz Kinzel, a resident there, informed me that an example of *Crotophaga sulcirostris* Swains. had been shot during the first week in January last at Jupiter Inlet. Mr. Kinzel examined the bird, and identified it himself with the aid of Ridgway's 'Manual,' in which it is stated that this species has only occurred in the United States in the Valley of the lower Rio Grande in Texas, thence extending southwards to Peru.—A. S. PACKARD, *Brown University, Providence, R. I.*

Groove-billed Ani (*Crotophaga sulcirostris*) in Arizona.—A specimen of the Groove-billed Ani is now in my possession which was shot about