

Coralline algal rhodoliths form extensive benthic communities in the Gulf of Chiriqui, Pacific Panama

Received: 3 March 2008 / Accepted: 5 March 2008 / Published online: 18 March 2008
© Springer-Verlag 2008



Fig. 1 The research team collecting from an extensive rhodolith bed at 30 m



Fig. 2 An array of diverse coralline algal species typical of 10–30 m sedimentary habitats throughout the Gulf of Chiriqui



Fig. 3 Fleshy algal epiphytes attached to coralline algal nodules (rhodoliths)

Large areas of rhodolith beds (free-living, crustose, calcareous, Rhodophyta nodules) have been documented worldwide (Foster 2001), but are not well-known for Pacific Panama. We observed vast maërl expanses of calcareous red algal rhodoliths (Fig. 1) covering sedimentary bottoms between island groups throughout the Gulf of Chiriqui, during our September 2007 expedition. These predominant coralline algal communities (Fig. 2), mostly between 10–30 m deep, consisted of numerous crustose coralline algal species (e.g., *Lithophyllum divaricatum*, *Lithophyllum alternans*, *Lithothamnion indicum*, *Lithothamnion australe*, *Fosliella fertilis*) and serve as microhabitats and primary substrates for a high diversity of relatively small animals and fleshy red algae (Fig. 3; e.g., *Hypnea* spp., *Gelidium* spp., *Halymenia* spp., *Sebdenia* spp., *Peyssonnelia* spp.). Hypothetically, the instability/mobility of the fist-size rhodolith spheres precludes the development of highly structured macroalgal or macrofaunal communities. On the other hand, we posit that rhodoliths provide a two-dimensional hard-substrate refuge for smaller algae from the intense herbivory (limpets, chitons, sea urchins, and fishes) characteristic of Pacific Panama (reviewed by Birkeland 1988) and present throughout the heterogeneously structured coral reefs and broad (7 m tidal amplitude) rocky intertidal zones of the islands and outcrops we studied.

Acknowledgments Observations were made during a cruise aboard the R/V Urraca supported by the Smithsonian Tropical Research Institute and the Smithsonian Institution Marine Science Network. Barrett Brooks, Donald Hurlbert, Jennifer Dorton, Irving Bethancourt and the ship's crew contributed immeasurably to the success of the expedition. Photographs are by Donald Hurlbert, S.I. Photographer. Harbor Branch Oceanographic Institute Contribution No. 1697 and Smithsonian Marine Station at Fort Pierce Contribution No. 732.

References

- Birkeland C (1988) Geographic comparisons of coral-reef community processes. Proc 6th Int Coral Reef Symp 1:211–220
Foster MS (2001) Rhodoliths: between rocks and soft places. J Phycol 37:659–667

M. M. Littler (✉) · D. S. Littler
Department of Botany, National Museum of Natural History,
Smithsonian Institution, P.O. Box 37012, Washington,
DC 20013, USA
e-mail: littlerm@si.edu

Reef sites

Coral Reefs (2008) 27:553
DOI 10.1007/s00338-008-0368-5