

terminal punctiform. Hind wings: the postmedial shade broader, outwardly edged by a lunular line.

Expanse, 25 mm.

*Habitat*: Rockstone, Essequibo River.

*Freilla abluta*, sp. n.

*Male*.—Palpi outwardly brown, inwardly, and tips of second and third joints buff white. Head and thorax pale buff, the collar narrowly velvety black in front and then brown, but broadly pale buff behind. Abdomen slightly darker with broad transverse dull gray shades, and some black and brown irrorations. Wings pale buff, with some scattered dark irrorations; terminal dark-brown points. Fore wings: some dark striæ on costa; traces of an antemedial outangled brownish line; a fuscous spot irrorated with gray at end of cell; a faint postmedial line, outangled beyond cell, and inbent to middle of inner margin, followed by three similar lines only visible from below vein 3; a faint outer, indistinct line, outwardly paler edged, lunular, marked by small dark brown spots on interspaces; termen shaded with brownish gray at tornus, and between veins 4 and 8. Hind wings: antemedial dark points on inner margin; three fine medial dark-brown lines, followed by five pale lines, the fourth of these with some small black-brown spots; a subterminal brownish shade from below apex to anal angle, darker spotted, beyond which the termen is faintly shaded with gray. Wings below whitish buff, the fore wing shaded with pale reddish brown; an irregular outer fine fuscous line; fore wings with darker striæ on costal and inner margins; traces of an antemedial shade, a black point in cell, and a small spot at end of cell; hind wings with discal point, and geminate medial line.

Expanse, 28 mm.

*Habitat*: Cayenne, French Guiana.

#### A NEW ULOPHORA FROM FLORIDA.

(Lepidoptera, Pyralidae)

BY HARRISON G. DYAR.

*Ulophora caricæ*, new species.

Fore wings gray, whitish over the disk; a broad line of raised black scales near the base; a broad pale ochereous band beyond, oblique, followed by a faint reddish line; median space gray, whitish beyond of cell, discal dot double, very small and faint; outer line slightly excurved mesially and denticulate, double, reddish, followed by a dark purplish shade, which fills all the terminal space and is intensified at apex. Hind wing sordid whitish, gray at tip. Expanse, 17 mm.

*Cotypes*: Male and female, No. 15428, U. S. National Museum, Miami, Florida. Bred from larvæ on papaya. December 2, 1912 (W. W. Yothers).