

Iron-Coated Quartz as a Provenance and Paleoclimatic Marker in the Rhône Delta, France

Jean-Daniel Stanley and Thomas F. Jorstad

Deltas-Global Change Program
Paleobiology, E-206 NMNH
Smithsonian Institution
Washington, D.C. 20560 U.S.A.

ABSTRACT

STANLEY, J.-D. and JORSTAD, T.F., 2002. Iron-coated quartz as a provenance and paleoclimatic marker in the Rhône Delta, France. *Journal of Coastal Research*, 18(4), 712-729. West Palm Beach (Florida), ISSN 0749-0208.

Analyses of 430 surficial and core sediment samples in the Rhône delta and adjacent region in southern France record similar average proportions of iron-stained quartz (ISQ) on the delta plain and in the Rhône and Durance rivers (~30%). These two rivers, largely responsible for the formation of the Holocene delta, carried most of their sediment load from distal source terrains.

The much higher ISQ values in sand on the delta's modern coast (to >40%) and on older sand strand deposits (56%) subaerially exposed in the central and southern delta plain are primarily the result of post-depositional *in situ* formation of iron coating on quartz particles. However, some subsurface delta plain core samples with high stained-grain content (>40%) have a different origin. These layers were derived from Pleistocene and older sediment in proximal terrains, characterized by very high ISQ values (67%), that were transported to the delta plain. Evidence indicating displacement of highly-stained material eroded from these more local source areas is quartz with high ISQ values (50%) in the Vidourle and Gardon rivers with headwaters in adjacent uplands.

Strata with large ISQ values (>40%) in Holocene core sections are attributed primarily to periodically increased hydrological fluxes that resulted from marked paleoclimatic changes and, possibly, effects of human activity as seen in the archaeological record in southern France. Wetter climatic conditions increased the erosion and reworking of proximal ISQ-enriched terrigenous sediment to the delta. This study indicates that the stained-grain method can be used to help identify changes in sediment provenance and dispersal patterns related to Holocene paleoclimatic fluctuations and/or human activity in the Rhône delta region.

ADDITIONAL INDEX WORDS: *Archaeology, climatic cycles, coastal processes, Costières, Crau, distal sediment, floods, Gulf of Lion, Holocene, iron oxides, Pleistocene, proximal sediment, quartz coating, sediment dispersal, size-sorting, source terrains, weathering.*

INTRODUCTION

Deltas on marine coasts form primarily by the accretion of river sediment transported to the sea, and the erosion and removal of some of this material by wave-driven currents and other nearshore processes. The resulting deposits, accumulating by interaction of fluvial, coastal and nearshore processes, are composed of terrigenous sediment derived from both distal and proximal terrains in the river drainage basin. Most petrologic studies of deltas have emphasized the importance of fluvial transport from distal sources, a process of seaward displacement of material over time by episodic 'stop-and-go' events. These, triggered by repeated floods, typically involve episodes of erosion → downvalley and overbank transport → sediment deposition and temporary storage along the fluvial dispersal path → renewed sediment erosion and transport by subsequent flooding (MEADE, 1988; STANLEY and HAIT, 2000). In contrast, there is much less documentation on the more direct sediment dispersal to a delta plain from proximal source terrains, such as those located in the lower river valley close to a delta's apex, on adjacent delta margins, and seaward of a delta's coast.

The present study evaluates the use of iron-coated quartz grains of sand size as potential petrologic markers to identify Rhône delta sediment derived from proximal terrains in southern France and seaward of the coast in the Mediterranean (Figure 1). As in most deltas, quartz sand particles are present in all samples of the study area. The terms iron-stained and iron-coated quartz grains are used here interchangeably, and abbreviated ISQ. This approach is based on findings in previous investigations made in different world coastal settings. Holocene delta sediments generally comprise lower ISQ proportions than older (pre-Holocene) alluvial deposits buried beneath Holocene deltaic sections (COLEMAN and ROBERTS, 1988; STANLEY *et al.*, 2000). Also containing relatively high proportions of stained grains are Quaternary and older formations that are subaerially exposed near delta margins (MORGAN and MCINTYRE, 1959; STANLEY *et al.*, 2001), and seafloor sediment on continental shelves seaward of deltas that were once subaerially exposed (SHEPARD *et al.*, 1960; BERRYHILL, 1976; STANLEY and CHEN, 1991). Although commonly present in delta sequences, the ISQ grain component has only been used in a few instances to identify sediment provenance and dispersal from proximal terrains (STANLEY *et al.*, 2001).






Figure 1. Map showing the locations of surficial sediment samples collected in the Rhône delta study area in 1999 and 2000 (stained-grain data listed in Table 1). Massif Central and Rhône, Durance and Gardon rivers shown in inset; Vidourle River shown in map.

Focus herein is on ISQ distribution patterns in 13 specific surficial environmental settings in and near the Rhône delta so as to better interpret the sedimentary origin of stained grains in the subsurface Holocene record. This petrologic parameter is examined in delta cores to determine whether ISQ values remained generally constant in subsurface Holocene sections, or fluctuated markedly through time. From such an analysis of stained-grains in delta cores, it may be possible to test the following hypotheses: (1) older geologic formations in the lower Rhône River valley and other proximal regions served as sources of sediment dispersed to the Holocene delta; and (2) enriched stained-grain zones in subsurface sections, especially those that are laterally extensive, provide information on paleoclimatic fluctuations that affected Holocene sedimentation patterns in the Rhône depocenter.

BACKGROUND FOR A STAINED-GRAIN STUDY IN THE RHÔNE SYSTEM

The Rhône's fluvial, deltaic and coastal margin systems have been extensively studied. On the basis of what is already known of its Holocene history, the Rhône delta is ideally suited for a petrologic investigation to distinguish proximal from more distal terrigenous quartz-rich sediment supply, and to identify paleoclimatic fluctuations that likely affected depositional patterns. Compared to other world deltas, the Rhône is of intermediate size. The distance between the

delta apex just north of Arles and the mouth of its major distributary, the Grand Rhône, is ~42 km, and the delta shoreline is ~90 km in length (Figure 1). Margins on both the eastern and western delta plain comprise well-defined upland terrains, some with elevations exceeding 100 m.

Salient geographic characteristics of the delta are described by RUSSELL (1942), DEPARTEMENT DES BOUCHES DU RHÔNE (1970), CORRE (1992), L'HOMER (1992), and depicted in the detailed map (scale 1:50,000) compiled by the INSTITUT GEOGRAPHIQUE NATIONAL (1992). Noted among recent hydrologic investigations are those of VIVIAN (1989), FRUGET (1992), RODITIS and PONT (1993) and ARNAUD-FASSETTA (2000). For the geological attributes of the lower Rhône and contiguous delta regions, the reader is directed to BERTRAND and L'HOMER (1975), COLOMB and ROUX (1978) and L'HOMER *et al.* (1981). Results of geological analyses of the delta's offshore sector are presented by ALOISI *et al.* (1975), BELLAÏCHE and MART (1995), GENSOUS and TESSON (1996) and VELLA *et al.* (1998).

In addition to natural phenomena, this region has been increasingly modified by human activity from the Neolithic to the present. Among recent studies discussing specific anthropogenic modifications of the delta are those of JORDA and PROVANSAL (1990, 1996), CORRE (1992), STOUFF (1993), PASQUALINI and LANDURE (1995), MORHANGE *et al.* (1998), HENSEL *et al.* (1999), and STANLEY (2000).

Table 1. Proportions of partially + fully coated iron-stained grains (ISQ values) in 184 surficial samples (of modern, Holocene, and older age) collected in 13 depositional environments of the Rhône delta and adjacent region. Sample localities are shown in Figure 1; distribution of ISQ values in the different environments is depicted in Figure 4.

Rhône Delta Plain Setting													
Grand Rhône Margin (n = 13)		Petit Rhône Margin (n = 6)		Old Rhône Branch (n = 13)		Delta Plain (n = 29)		Lagoon and Wetland (n = 32)		Marsh (n = 17)		Beach (n = 17)	
202	28.07	248	29.14	207	33.52	216	20.47	201	40.50	204	17.52	200	32.75
203	39.59	252	15.64	228	7.98	218	16.15	230	24.52	224	27.64	222	42.77
208	38.30	254	12.38	234	6.25	221	46.11	235	15.13	229	45.97	232	20.05
213	69.88	272	19.37	237	19.87	227	50.30	236	25.87	255	33.71	250	20.25
214	34.00	280	24.26	238	12.39	233	30.69	239	10.19	257	28.80	267	34.41
215	12.69	303	21.29	274	7.35	241	13.65	240	17.01	261	14.85	287	20.67
217	17.71			278	27.80	243	16.43	251	12.08	269	31.00	288	20.31
220	9.03			294	24.31	244	33.54	262	26.95	270	35.57	289	15.54
225	23.05			368	7.34	245	28.78	265	9.03	293	35.80	292	34.59
226	30.66			369	49.24	253	16.57	266	27.49	296	46.92	348	37.21
242	29.97			373	48.42	258	9.07	273	16.62	301	22.49	351	36.15
247	15.41			374	54.68	259	7.59	276	23.08	307	28.21	356	23.18
364	50.48			379	15.89	260	24.00	285	11.65	322	2.41	359	37.91
						263	36.19	299	21.99	347	15.29	362	34.71
						271	33.76	312	10.69	378	30.30	366	45.59
						275	31.05	323	4.49	380	49.85	371	39.16
						277	44.65	336	8.11	383	1.31	376	29.03
						279	32.25	337	19.87				
						281	30.00	338	32.33				
						282	23.10	340	11.71				
						302	19.56	341	36.51				
						304	26.69	342	27.12				
						305	13.89	343	12.19				
						306	22.25	344	5.01				
						308	16.37	345	25.64				
						309	25.49	352	31.58				
						313	3.82	353	37.25				
						315	34.65	354	42.14				
						321	18.37	357	45.89				
								360	52.24				
								363	53.87				
								367	37.83				

Mineralogical analyses indicate that most sediments deposited in this delta were released by the Rhône river with headwaters in the high Alps of Switzerland. The river flows to France, first to the southwest, and then south, to the Mediterranean (Figure 1). Much of the material displaced toward the coast is derived from its major tributaries in France, including the Saône, Ain, Isère, Drôme and Durance, that drain geologically diverse terrains north and east of the delta. In addition to major distal sediment source areas in the Swiss Alps are the intermediate terrains in the Massif Central west of the Rhône river (drained by the Ardèche and other tributaries), the Jura and external Alpine chain in southeastern France (VAN ANDEL, 1955; VIVIAN, 1989; ARNAUD-FASSETTA, 2000). Much of the material transported from these regions is deposited in the delta plain as gray sand and olive gray to dark gray and dark olive gray silty mud (KRUIT, 1955; L'HOMER, 1975, 1987). The near-horizontal deposits forming this low-lying, fan-shaped, wave-dominated delta (DUBOUL-RAZAVET, 1956; GALLOWAY, 1975; L'HOMER, 1992) are positioned between the apex just north of Arles and the Gulf of Lion shelf south of the coast.

Distinguishing potential sources of ISQ grains transported from proximal areas to the Rhône drainage basin is of special

interest in this study. These could include highlands that comprise lithologically diverse formations near, and bordering, the delta proper (Figure 1). Proximal terrains include Pleistocene and older terrigenous strata (Costières and Crau terraces) and carbonate (Alpilles) units (COLOMB and ROUX, 1978; L'HOMER, 1987; JORDA *et al.*, 1990). Also recognized are pre-Holocene terrigenous units in the drainage basin of the southern Rhône that include reddish brown, orange and yellowish brown strata. Such deposits are identified in lithostratigraphic studies by general terms such as 'terre rouge à cailloux siliceux', 'sol rouge', 'marnes jaunes', 'vases jaunes', 'sable gris-jaunâtre', and 'cailloutis à matrice sableuse jaune' (COLLOT, 1904; BONIFAY, 1957; ALOISI and DUBOUL-RAZAVET, 1974; L'HOMER, 1975; COLOMB and ROUX, 1978).

Petrologic analyses of these strata identify oxidized lithologies that include coated quartz and limestone particles of sand to pebble size and sandy mud matrices. Colors of individual particles include orange-brown, yellow-brown and tan. Iron oxides in paleosols, intergranular matrices and particle coatings (ALOISI and DUBOUL-RAZAVET, 1974; JORDA *et al.*, 1991b) in these older formations previously have been attributed to several phenomena, including weathering of limestone carbonate terrains and pebbles, soil development, and

Table 1. *Continued.*

Rhône Delta Plain Setting				Rivers				Others			
Dune (n = 13)		Relict Sand Strand (n = 8)		Rhône and Durance Rivers Above Delta (n = 5)		Massif Central Tributaries (n = 7)		Delta Margin (n = 16)		Offshore Samples (n = 8)	
223	60.44	256	53.82	328	18.36	316	53.92	209	19.28	1	9.10
231	30.41	264	52.86	329	20.47	317	20.25	210	94.28	2	2.30
268	34.67	295	48.77	330	45.14	318	43.96	211	64.43	3	29.50
290	34.74	297	66.98	331	39.16	319	57.85	212	47.35	4	40.40
291	47.93	298	51.77	332	31.21	320	67.97	219	19.05	5	23.90
349	22.98	300	57.28			326	63.23	249	75.94	6	5.50
350	42.42	370	59.11			327	40.65	283	52.52	7	11.00
355	33.87	375	55.56					284	64.51	8	40.00
358	38.87							310	86.23		
361	34.07							311	74.76		
365	43.45							314	88.91		
372	38.87							324	67.62		
377	42.44							325	72.48		
								333	81.49		
								381	91.04		
								382	71.56		

staining by ground water flow in this region. In addition, it has been suggested that, after formation, some of these oxidized older materials in Provence and adjacent regions of southern France were extensively eroded, reworked and laterally displaced (BONIFAY, 1957). Other potentially important proximal sediment sources of ISQ grains transported to the delta include Quaternary and older formations eroded from exposures along the Mediterranean coast and from those now submerged in the Gulf of Lion shelf south of the delta. Such Quaternary and older deposits in coastal and offshore settings are commonly ochre and tan (ALOISI and DUBOUI-RAZAVET, 1974; LABEYRIE *et al.*, 1976).

METHODS

Sample Sites

Numerous surficial samples (Figure 1) and subsurface sediment in cores were collected in the Rhône delta proper and in adjacent regions in order to (1) distinguish material of proximal origin from more distal derivation, and (2) to identify evidence of changes in provenance and dispersal through time. Stained quartz grains were examined in a total of 430 sediment samples from surface sites and core sections. Of

these, 184 samples were collected in 1999 (numbers 200–324) and in 2000 (numbers 325 to 383) from the modern (most recent to past several centuries), Holocene (past 10,000 years) and older geographic settings numbered 1 to 5 below. ISQ data from surficial samples are listed in Table 1. Averaged ISQ values and range of proportions of the surficial samples are given in Table 2. Another 246 samples, recovered in sediment cores, are of Holocene to late Pleistocene age (settings 6 and 7 below); ISQ data for these subsurface samples are listed in Table 3.

(1) *Modern delta plain*, a total of 9 environments (n = 148), include the Grand Rhône distributary and its channel margin (n = 13 samples), Petit Rhône distributary and its channel margin (n = 6), old (earlier relict) Holocene Rhône channels still preserved on the delta plain (n = 13), delta plain proper between distributary channels (n = 29), delta lagoons (n = 32), marshes (n = 17), beaches (n = 17), coastal dunes (n = 13), and relict Holocene sand strands exposed on the southern to central delta plain (n = 8);

(2) *Modern Rhône and Durance rivers*, the two dominant fluvial systems that transported the bulk of sediment to the delta are the Rhône, flowing from the north, and the Durance

Table 3. ISQ values for 212 subsurface samples in 18 borings collected on the Rhône delta plain (core localities shown in Figure 9).

Depth (m)	Partial Plus Fully Stained	Depth (m)	Partial Plus Fully Stained	Depth (m)	Partial Plus Fully Stained	Depth (m)	Partial Plus Fully Stained	Depth (m)	Partial Plus Fully Stained
CORE #105 (n = 20)		30.30	3.3%	27.68	16.28%	25.16	22.4%	18.04	33.2%
0.96	39.0%	34.36	2.6%	30.13	19.42%	30.03	13.7%	24.31	21.2%
3.52	45.6%	40.04	1.6%	34.41	23.49%	34.62	4.7%	27.65	25.6%
6.52	42.4%	CORE # 108 (n = 10)		37.66	37.30%	38.02	17.9%	31.36	19.7%
9.22	32.2%	0.15	17.3%	40.05	9.82%	42.04	9.1%	34.36	26.2%
12.04	26.8%	3.99	19.3%	43.05	5.21%	44.00	19.2%	37.64	17.1%
15.61	35.1%	7.09	5.9%	45.85	6.98%	45.35	41.0%	38.65	12.9%
18.39	48.6%	9.19	24.9%	47.94	3.32%	CORE # 119 (n = 14)		CORE #126 (n = 11)	
21.84	28.4%	12.46	40.6%	51.76	7.57%	0.32	18.2%	0.32	23.2%
23.28	26.1%	15.56	21.4%	56.00	11.73%	3.21	19.6%	3.42	24.7%
27.82	38.0%	18.31	2.6%	58.69	7.26%	6.31	18.8%	5.71	10.5%
30.58	46.7%	20.22	48.9%	61.43	0.65%	9.61	22.9%	9.66	14.0%
33.83	32.8%	24.04	3.8%	67.53	0.67%	12.12	31.2%	12.52	22.6%
36.03	18.1%	27.06	24.9%	70.00	5.23%	15.83	28.3%	15.71	32.7%
40.00	7.6%	CORE #109 (n = 7)		70.26	3.79%	18.18	20.7%	18.35	27.4%
42.99	19.7%	0.11	31.0%	70.44	19.94%	21.41	4.4%	21.17	22.5%
46.31	3.6%	3.61	16.8%	CORE #116 (n = 17)		24.41	3.3%	24.67	14.9%
50.17	7.8%	6.34	2.9%	0.11	24.6%	27.83	3.2%	27.4	5.6%
53.72	10.6%	9.36	18.2%	3.54	22.7%	33.13	2.5%	29.49	9.1%
55.83	13.9%	12.15	28.5%	6.56	14.4%	36.05	19.7%	CORE #128 (n = 16)	
56.59	4.3%	15.04	18.5%	9.41	18.5%	39.31	37.6%	0.85	16.4%
CORE #106 (n = 12)		18.12	25.1%	11.46	20.4%	43.31	18.9%	1.16	6.1%
1.63	27.7%	CORE #110 (n = 11)		15.44	11.8%	CORE #122 (n = 4)		1.77	19.4%
9.24	29.5%	0.17	27.2%	18.06	9.5%	0.92	9.0%	2.64	6.9%
12.00	28.3%	3.13	5.5%	21.82	13.1%	2.71	10.1%	3.42	17.5%
14.15	42.9%	6.08	21.3%	24.79	20.4%	6.145	2.0%	4.07	4.6%
18.03	35.3%	9.46	21.3%	27.45	2.9%	9.035	10.2%	5.52	11.2%
24.56	3.9%	12.17	43.7%	31.73	0.0	CORE #124 (n = 13)		9.35	38.8%
30.17	4.7%	15.43	9.7%	33.84	0.0%	0.16	35.2%	15.07	32.1%
35.17	1.7%	17.58	43.2%	37.60	11.7%	3.01	12.3%	18.57	4.0%
40.65	32.5%	21.19	50.0%	41.52	4.0%	5.91	15.0%	19.77	19.3%
44.86	25.6%	24.34	9.3%	44.60	1.5%	9.12	14.7%	24.15	14.8%
48.54	27.0%	27.20	1.9%	47.36	23.7%	11.89	11.6%	27.52	4.8%
49.31	28.6%	30.37	24.5%	53.09	6.6%	16.02	27.0%	30.16	4.3%
CORE #107 (n = 14)		CORE # 112 (n = 6)		CORE #117 (n = 4)		18.93	27.0%	33.91	28.0%
0.43	28.1%	0.11	20.7%	23.89	9.0%	21.42	10.3%	36.75	29.2%
1.10	33.2%	3.04	4.56%	11.70	20.8%	24.57	9.8%	CORE #129 (n = 9)	
3.51	22.9%	5.73	1.00%	18.39	20.1%	27.71	8.8%	0.45	27.69%
6.01	31.8%	9.52	3.92%	28.09	1.3%	30.12	18.5%	1.38	38.82%
9.42	22.8%	11.26	30.65%	CORE #118 (n = 13)		33.81	21.3%	2.72	36.61%
12.90	18.0%	14.32	27.87%	1.90	30.0%	36.19	17.9%	4.27	50.45%
14.86	27.2%	CORE #114 (n = 20)		3.22	31.9%	CORE #125 (n = 11)		5.01	43.16%
17.68	25.2%	14.90	33.0%	5.72	20.6%	0.34	27.3%	6.01	28.93%
21.32	20.4%	18.50	32.11%	10.26	21.6%	3.03	93.5%	6.86	39.29%
24.11	24.1%	21.02	19.24%	15.10	31.7%	6.48	23.4%	7.41	34.60%
27.44	17.0%	24.70	6.15%	20.36	17.8%	9.31	32.9%	9.49	23.96%

(see Figure 1, inset), its major tributary flowing from the east and joining the Rhône at Avignon (n = 5; collected in 2000); (3) *Modern tributaries*, include those that transport substantial amounts of pre-Holocene material from proximal highlands: the Gardon River (or Gard), a tributary with headlands in the SE Massif Central (Figure 1, inset) that joins the southern stretch of the Rhône north of Arles; and the Vidourle River that disperses sediment from the Costières southward to and across the western part of the modern delta (n = 7; collected in 2000; see Figure 1); (4) *Older formations close to the delta margin*, comprise largely Pleistocene, Pliocene and older units that are subaerially exposed in upland terrains close to the delta plain (n = 16; collected in 1999 and 2000); (5) *Gulf of Lion continental shelf south of the delta*, includes

Holocene and Pleistocene formations and palimpsest-reworked relict sediment mixes thereof (n = 8); samples collected by the Cefrem Program, Université de Perpignan and provided courtesy of Dr. J.-C. Aloisi (ALOISI, 1973; ALOISI and CHARLET, 1975; ALOISI *et al.*, 1975; GENSOUS and TESSON, 1996); (6) *Holocene delta subsurface strata*, recovered in 19 sediment cores (n = 238; Table 3), includes 18 borings (n = 212) that range in length from 18 to 71 m (described by OOMKENS, 1970; TER KEURS, 1971), and 1 boring (PI, Piton-1; 18.3 m in length; n = 34) collected in the city of Arles just west of the Rhône (courtesy of G. Arnaud-Fassetta, 2001, personal communication); and (7) *Lower Holocene (H) to late Pleistocene (P) alluvial sections*, recovered at the base of 7 of the 19 borings listed in setting

(6) above ($n = 8$; cores 105, 106, 110, 114, 118 and 125 described by OOMKENS, 1970, and TER KEURS, 1971) and core PI ($n = 2$); these 10 subsurface samples are part of the total 246 core sample set.

Analyses

The quartz-grain method used here is described by STANLEY and others (2000). Proportions of quartz particles of sand size that are *clear* (transparent and translucent grains), *partially coated* (grains whose surfaces are stained to 75% by pigment), and *fully stained* (grains that are >75% coated) were determined for all of the 430 surficial and core samples (Tables 1, 3). Percentages of the three quartz grain categories were determined by making stereoscopic microscope counts of at least 300 quartz grains in each sample, selected randomly from the sand-size fraction (63–2000 μm). The more than 139,000 particles in the samples were examined by one operator (the second author) so as to make consistent recordings and obtain reproducible data. Quartz grain counts of sand size were examined using a Leitz Wetzlar binocular microscope using a Dyonics fiber-optic light source. Grain color was determined by using dry grains under fluorescent lighting and using standards in color charts (MUNSELL COLOR, 1975).

The averaged proportions of clear, partially stained, fully stained, and partially + fully stained grains (ISQ), and range of values in the 184 surficial samples from 13 environments in the modern, Holocene and older geographic settings 1 to 5 cited above are summarized in Table 2. The fully stained particle values in most samples is low (<3% in the different settings) and thus, in this study, ISQ value refers specifically to the percentage of partially plus fully stained grains.

Two replicate grain-size analyses were performed on all 184 surficial samples collected in the 13 environments with a Coulter Counter laser particle analyzer (LS200). From these, the following data were averaged: relative percent of sand (63–2000 μm), silt (0.5–63 μm), and clay (<0.5 μm); and mean grain size (in μm). The textural information for the surficial samples in delta and adjacent settings are listed in Table 2.

The elemental composition of grain coatings was determined for 4 representative samples in the study area: modern delta beach (surficial sample 366; Figure 2A); subsurface delta sample from an ISQ-enriched layer in the Piton core (PI) at Arles (depth of 10.65 m from the core top; Figure 2B); highland exposure (Costières) adjacent to the delta (surficial sample 381; Figure 3A); and margin of the Vidourle River in the western delta (surficial sample 298; Figure 3B). The selected grains were prepared by preferentially mounting the specimens in epoxy on standard glass microprobe discs. Each disc was then polished to expose a transverse view of the samples, thus providing maximum exposure of the stained area coating the grains. Approximately 12 grains from each of the four localities were analyzed using backscattered (BSI) and secondary (SEI) imaging with a JEOL JMA-840A high resolution scanning microscope. Analyses and elemental distribution of micro-areas for the four samples were performed

by Energy Dispersive Spectroscopy (EDS), with a working distance of 25 mm at 15 KeV and a beam current of 6 nA.

Analyses and interpretations of the mineralogical, faunal and floral components comprised in the sand-size fraction in each sample will be presented in a separate study (G. Randazzo and others, in preparation). A detailed petrologic study of core Piton collected at Arles is also presently in preparation by G. Arnaud-Fassetta, C. Beaudoin, H. Bruneton and others.

STAINED-GRAIN BASELINE DATA FOR THE STUDY AREA

The colors of the coated grains typically include dark red (2.5YR 3/6), dark reddish brown (2.5YR 3/4), dark yellowish brown (10YR 3/6), and orange (5YR 6/14). Interpretation of elemental component from the EDS analyses have identified the coatings on quartz particles as iron oxides (Dr. A. Logan, 2002, Personal Communication; Figures 2, 3). This study does not address any relationship or suggestion regarding the variability in iron-stained color.

The proportion of clear grains in the 9 delta environments averages ~70% and ranges from 44 to 80%. Four ISQ grain categories are identified on the basis of natural stained-grain groupings determined from surficial and subsurface sample sets in the study area (Figure 4 and Tables 1, 3): <30%; 30–39.9%; 40–49.9%; and >50%. Salient observations pertaining to ISQ values in the 13 environments of the 7 geographic sectors discussed in the previous sections are summarized below.

(1) *Modern Rhône delta plain*.—Most samples from the modern delta plain proper, from the Rhône valley just north of the apex to ~5 km landward (north) of the coast, but excluding relict sand strand deposits exposed on the delta surface, are characterized by relatively low and, in some cases, moderate, ISQ values (~30%). Calculated average values for Rhône distributary channel settings are as follows: Grand Rhône channel margin = 31%; Petit Rhône channel margin = 20%; and relict Holocene Rhône branches = 24%. Low average ISQ values are noted in three delta plain environments as follows: delta plain proper = 25%; lagoons on the delta plain = 23%; and marshes on the delta plain = 27.5%. Average values on beaches are 31% (example in Figure 2A), and in coastal dunes are 39%. In contrast, most samples collected on the southern and western delta plain surface within ~5 km of the coast record higher values, ranging from 30 to >50%. Sediment samples on the western delta plain in the vicinity of the Vidourle river also contain large ISQ values, commonly 40 to >50%. Samples from relict sand strands exposed on the southern and central plain are characterized by the highest average ISQ proportion (56%) on the delta proper.

(2) *Modern Rhône and Durance rivers*.—The average ISQ value is 31% in channel samples at the juncture of these 2 major rivers, and in the 35 km-long Rhône valley stretch between Avignon and the delta apex just north of Arles. This value is the same as that of samples along the Grand Rhône margin on the delta plain.

(3) *Modern rivers flowing from adjacent highlands*.—The Gardon and Vidourle, transporting sediment to the delta from proximal source terrains in the southern Massif Cen-




Figure 2. (A) SEM secondary electron image (SEI) along with corresponding dispersive energy spectrum for sample 366 grain collected on Rhône delta beach. The EDS spectrum corresponds to the positions of the coating (depicted by arrows) that surrounds most of this selected quartz grain. Bar scale = 1 mm. (B) SEM secondary electron image (SEI) for a Piton core sample collected at 10.65 m from core top. The EDS spectrum corresponds to the positions of the coating (depicted by arrows) that surround most of the quartz grain. Scale in μm .

tral, comprise an average ISQ value of $\sim 50\%$ (example in Figure 3B). This is significantly higher than that recorded by samples in the Rhône and Durance (31%, see setting 2 above), the two much larger rivers responsible for the deposition of most Holocene delta strata.

(4) *Older sequences close to the delta.*—By far the largest average ISQ value (67%; range 19 to 94%) in the study area is calculated for the 16 samples collected in Pleistocene and Pliocene formations exposed in the Costières and Crau highlands adjacent to the Rhône delta (example in Figure 3A).

(5) *Gulf of Lion shelf south of the delta.*—Seafloor samples from the shelf south of the Rhône delta record an average ISQ value of 20%, with a range of sample values from 2 to 40%. Proportions of coated grains in the offshore are higher in sand than in mud-rich deposits.

(6) *Holocene core sections.*—Most of the 246 samples from the 19 borings recovered in the modern delta plain are of Holocene age and record a remarkably large range of ISQ values

(from 0 to 94%). The following percentages of samples in the 4 coated grain categories are recorded: $<30\%$ ISQ occurs in 77% of all core samples; 30 to 39.9% ISQ in 15%; 40 to 49.9% ISQ in 5%; and $>50\%$ ISQ in 3% (Table 2). The distribution of subsurface strata that comprise high amounts ($>40\%$) of stained grains (example in Figure 2B) is shown in Figures 9 and 10.

(7) *Lower Holocene-Late Pleistocene core samples.*—The ISQ values (range from 4 to 41%) are determined for strata recovered at the base of 6 delta cores which were previously attributed (*cf.* OOMKENS, 1970; TER KEURS, 1971) to early Holocene to late Pleistocene to age. The stained-grain values of samples collected at the base of 5 of these cores, identified here as H/P (Figure 9), are comparable to those of sediment carried in the modern Rhône and Durance rivers and to most Holocene deposits on the present delta plain; these proportions are much lower than those recorded for Pleistocene and older sequences exposed in the adjacent delta highlands (see




Figure 3. (A) SEM backscattered image (BSI) along with corresponding dispersive energy spectrum for sample 381 material collected in uplands close to the Rhône delta. The EDS spectrum corresponds to the positions of the coating (depicted by arrows) that surrounds most of the selected quartz grain. Scale in μm . (B) SEM secondary electron image (SEI) along with dispersive energy spectrum for sample 298 collected in the Vidourle River, with headland in the delta margin uplands. Scale in μm .

4, above). The ISQ values range from 10 to 25% at the base of cores 110 and 114; based on overall lithology, these sections are likely late Pleistocene (P) strata. In marked contrast, analyses of basal samples in two cores, 118 (Figure 9) and PI (Figure 10), record ISQ values $>40\%$, more firmly indicating that the base of these borings actually recovered late Pleistocene material beneath the Holocene section.

EFFECTS OF SIZE-SORTING AND *IN SITU* FORMATION

The average ISQ values in the Rhône delta region ($\sim 30\%$) are high when compared to average stained-grain proportions

recorded in some other deltas, such as the Rio Grande (to $\sim 20\%$; STANLEY *et al.*, 2001) and the Ganges-Brahmaputra (8%) and Nile ($\sim 6\%$) (STANLEY *et al.*, 2000). Several phenomena could account for the unusually high ISQ values of Rhône delta samples: grain size and associated size-sorting factors; *in situ* post-depositional development of staining on the delta plain, including biologically-generated staining; and reworking and dispersal of older iron-stained sediment from source terrains that include large proportions of particles that were iron-stained prior to their deposition in the delta.

To evaluate potential influences of grain-size and size-sorting effects on ISQ distributions, the percentages of sand, silt,




Figure 4. Distribution of ISQ (partially + fully stained grains) values for 184 surficial samples collected in 13 depositional environments of the Rhône delta study area. Triangle and associated percentage depict averaged ISQ value for each environment.

clay, and mean size were calculated for each sediment sample collected on the Rhône delta plain and adjacent surficial environments. With this information, determination was made of the relation between each of these four textural parameters and relative percentages of each of the four stained-grain parameters (clear, partially stained, fully stained, fully + partially stained quartz). A set of 192 graphs was generated with the available ISQ and textural data for each surficial sample in 12 environments ($n = 176$; this survey excludes the 8 offshore samples). Representative stained grain versus size graphs are shown in Figures 5 to 8.

As a first step, 16 graphs were compiled using the above-

listed 4 stained-grain types versus the 4 grain-size attributes for all 176 surficial samples. The Coefficient of Determination (r^2 values) indicates a 'goodness of fit' correlation between variables, with r^2 values near 1.0 indicating a strong relationship between variables. The r^2 values determined for each of the 16 graphs range from 0.00987 to 0.220429. These low values and the random distribution patterns of data points on the graphs indicate an overall absent to weak statistical correlation between proportion of stained quartz and their grain-size attributes. Two representative graphs (Figures 5A,B) show that, when all sample data are evaluated, stained grains are independent of spe-




Figure 5. Graphs of surficial samples ($n = 176$) examined in this study (these data exclude offshore samples), showing plots of the proportion of partially + fully stained quartz grains versus (A) percentage of sand ($r^2 = 0.193$) and (B) mean grain size ($r^2 = 0.169$). r^2 = Coefficient of Determination.




Figure 6. Graphs showing ISQ values versus percentage of sand in surficial samples collected in (A) modern beach, (B) modern dunes and (C) older (relict Holocene) sand strand deposits on the Rhône delta plain. Note increasing ISQ values from A to C. r^2 = Coefficient of Determination.




Figure 7. Graphs showing ISQ values versus percentage of sand in surficial samples collected on the delta plain, including the (A) Grand Rhône distributary channel, (B) old (relict) Rhône distributary branches, and (C) delta plain proper. r^2 = Coefficient of Determination.

cific grain size is preferentially stained and that size-sorting effects are minimal.

To more critically detect possible iron stain-grain size relationships, an additional set of 16 graphs was generated using the same 8 above-listed parameters for each of 12 specific modern environments (those shown in Figure 4, except the offshore samples). This resulted in a total of 176 graphs, in addition to the 16 cited above.

- All graphs compiled for 7 of these 12 environments (Rhône and Durance rivers sampled north of the delta apex, Grand Rhône channel margin on the delta plain, lagoon, marsh, dune, relict Holocene sand strand, delta margin settings) show no statistical correlation between ISQ values and sand-size (Figures 6A–C, 7A, and 8).
- Stain versus size graphs for the other 5 of the 12 environments (relict Rhône branches, interdistributary delta plain, some coastal facies) record a weak correlation (Figures 7B, C). Of these, only several show a moderate correlation: those depicting stained quartz versus relative

percentage of clay in some near-coast environments and along the Petit Rhône channel; and those depicting stained grains versus relative percentages of clay and mean size of samples. These latter suggest size-sorting effects, probably the result of transport processes that preferentially affect the distribution of finer-size fractions (clay and silt) in the 2 delta environments.

The survey indicates that, for the most part, proportions of ISQ values are independent of, or only poorly related to, specific grain-size parameters of sediment transported to the Rhône delta. This implies that factors other than sediment transport, depositional mechanism, and size-sorting more directly influenced the ISQ values recorded in the study area.

We suggest two other more probable causes for the observed stained-grain variations and distributions: (1) post-depositional and *in situ* formation of iron-coated particles locally in the delta; and (2) deposition of older reworked quartz grains that were already iron-stained in their source terrains




Figure 8. Graph showing ISQ values (average = 67%) versus percentage of sand in samples collected in Pleistocene and older upland terrains located near the Rhône delta plain. r^2 = Coefficient of Determination.

prior to their transport to the delta. With respect to (1), conditions in some delta environments appear to have favored the formation of iron coatings on quartz particles after their deposition on the delta plain. Conditions leading to staining commonly involve the availability of iron in various forms, and chemical and/or biological alteration on grain surfaces that favor addition and/or removal of an iron film on particles (cf. VAN HOUTEN, 1968, 1973; GADEL and PAUC, 1973; DORN, 1998; NEWMAN, 2001; YAPP, 2001).

The processes that resulted in iron coating on quartz are not specified here. It appears that some post-depositional phenomenon may have resulted from the somewhat lower-than-average ISQ values of partially + fully stained grains recorded along the Petit Rhône margin (20%), in delta lagoons and wetlands (23%), and seafloor settings south of the delta margin (20%). This finding is in marked contrast to the much higher average values (39–56%) recorded in the sand-rich coastal (dune, relict sand strand) environments. The progressive increase of ISQ values (Figures 4, 6A–C), from modern beach (31%); to dune (39%) to older (relict) strand sand units on the southern to central delta plain (56%), indicates relatively recent Holocene to possibly ongoing formation of iron oxide film on particles in these predominantly coastal sand exposures. The absence of any apparent relation between grain size and staining as shown in Figure 6 is an additional, albeit indirect, indication for *in situ* formation of the higher than average ISQ values in these 3 environments, rather than the result of transport process.

ISQ VARIATIONS AS A FUNCTION OF PROVENANCE AND DISPERSAL

The ISQ distribution data provide evidence that a large proportion of particles transported by rivers to the Rhône delta was already partially or fully stained prior to reaching the modern delta plain environments. Sediment provenance of

older stained material, rather than transport process *per se*, appears to be a major factor controlling the marked variation of ISQ values recorded in the study area. The large proportions of clear, non-stained grains and moderate ISQ values (average of ~70%) in most delta samples correlates well with other petrologic attributes of Rhône delta deposits, such as the composition of heavy and light mineral assemblages (VAN ANDEL, 1955; ALOISI and CHARLET, 1975). Considered together, these findings support the contention by most workers that the bulk of sediment forming the modern delta plain surface was derived from distal (Alpine, Jura) and intermediate (Massif Central) sources. Significant in this respect is the similarity of average ISQ proportions (~24–31%) and stained-grain colors in samples of the modern lower Rhône and Durance channels, lower Rhône valley between Avignon and the delta apex at Arles, Grand Rhône channel, relict Rhône distributaries on the delta plain and other environments on the delta plain proper (Table 2 and Figure 4).

In marked contrast with the above findings are the ISQ values (67%) in proximal subaerial exposures of older formations in upland terrains of the lower Rhône region (average 67%) and in the rivers that drain them (average 50%). These are much higher average values than on much of the modern delta plain and in Rhône-Durance rivers (<31%) (Figures 1, 4). This finding indicates that, at least in recent time, there has been a much smaller amount of sediment transported to the delta from proximal sources, relative to material derived from intermediate and distal terrains. To determine whether the present ISQ delta plain pattern remained constant, or fluctuated, during the Holocene requires a review of stained-grain data obtained from subsurface samples (Table 3). The core data show that ISQ content in the study area is highly variable in time and space (Figures 9, 10).

A likely explanation for this is a fluctuating sediment input




Figure 9. Logs of 18 Rhône delta cores examined in this study, showing sampled strata (depth and ISQ data in Table 3) that record primarily moderate proportions of stained quartz grains (30% or less; in white), and a few layers with high ISQ values (>40%; in black). Samples from the base of cores 105, 106 and 125, collected at subsurface depths are attributed to the Pleistocene age by OOMKENS (1970) and TER KEURS (1971); range of ISQ values (4–29%), more likely indicate strata of Holocene (H/P) age. The base of cores 110 and 114, of possible Pleistocene age (P), has values that range from 10 to 25%. In contrast, the base of core 118, recording a much higher ISQ value (41%), is interpreted confidently as late Pleistocene (P).

to the delta from proximal sources (southern France, Provence, offshore). This possibility has been suggested recently by others (ARNAUD-FASSETTA and PROVANSAL, 1999; ARNAUD-FASSETTA, 2000). Delta sediments with enriched ISQ content (>40%) are attributed here to two distinct proximal sources, one terrestrial, the other seaward of the Rhône coast. The large ISQ values of samples at the delta's modern coastal margin (Figures 1, 2A) result, at least in part, to sediment derived from presently submerged sources on the continental shelf south of the delta. Yellowish to ochre sands and silts have been recovered between the coast and outer shelf in the Gulf of Lion (*cf.* ALOISI and DUBOUL-RAZAVET, 1974). However, our analyses here indicate that ISQ values in surficial




Figure 10. Layers with high proportions of stained-quartz strata (shown in black) in radiocarbon-dated Piton (PI) core collected in Arles (samples and dates provided courtesy of G. Arnaud-Fassetta, 2001, University of Paris-7) are correlated with Holocene periods of markedly altered deposition (shown in grey), paleoclimatic chronology, and cultural ages in southern France. The latter information, shown in the 3 columns in the right half of the figure, is modified from JORDA and PROVANSAL (1996).

offshore samples are low (average 20%), and that samples in 5 cores (112, 114, 116, 117, 118) recovered along the coast also record only low to modest ISQ values in subsurface sections (Figure 9). We propose two origins for delta coastal margin sediment that comprise high ISQ values: post-depositional iron-stain formation in Holocene to modern dune settings as indicated earlier; and dispersal of sediment from eroded older coastal exposures in the southern France region, as yet undetermined, that already comprise a high ISQ content and are moved along the coast. Coast-parallel transport is a response to the high-energy wave regime in this region (*cf.* CORRE, 1992; L'HOMER, 1992; BRUZZI and PROVANSAL, 1996; DURRIEU DE MADRON and PANOUS, 1996; SUANEZ and PROVANSAL, 1996; ESTOUREL *et al.*, 2001). An important portion of this reworked iron-stained material is driven landward onto the delta's nearshore margin primarily by strong winter storms (BRUZZI and PROVANSAL, 1996).

An origin other than the above, however, is needed to ex-

plain the large ISQ variations recorded in subsurface sections of Holocene non-marine sediment, especially those in borings collected in the northern and central sectors of the Rhône delta. Most ISQ sample values in subsurface sections are <30%. Available core data indicate that subsurface strata with large (>40%) ISQ values in this part of the delta constitute only a relatively small part of the total Holocene section, and it appears that these layers formed only periodically on the delta plain (Figures 9, 10). Of note in this respect are the relatively thin (<30 cm), enriched ISQ layers (values >40%) identified in 7 of the 19 cores examined; these high ISQ samples account for a total of 22 (or 9%) of the 246 samples examined. Although few in number, the stain-rich strata are widely distributed in the delta's subsurface sections (Figure 9): they occur in delta plain cores located close to adjacent uplands (core 129), in the central delta plain (for example, cores 108 and 110) and, locally, along the coast (cores 105 and 106).

STAINED-GRAIN VARIATIONS AND CLIMATIC FLUCTUATIONS

Distinct stained-grain changes occurred in the Rhône delta through Holocene time, as recorded in cores of the north and central plain. We attribute these variations to two factors: paleoclimatic fluctuations, especially those involving changes in precipitation; and human activity, including increased deforestation and intensified, or modified, agriculture practices that affected Provence and adjacent regions of southern France (JORDA and PROVANSAL, 1990, 1996; JORDA *et al.*, 1991a,b; PROVANSAL, 1995). Since they are often closely associated, it is difficult to distinguish between the relative effects of climate change and human activity (*cf.* BUTZER, 1982; PROVANSAL, 1995). Nevertheless, previous studies have shown that climatic conditions in this region fluctuated markedly between the early Holocene and the present, and that such changes altered the flow and sediment discharge patterns of the Rhône and Durance, smaller rivers and the tributaries flowing from the southern Massif Central (VIVIAN, 1989; TRICART and BRAVARD, 1991; BRAVARD *et al.*, 1992; FRUGET, 1992; RODITIS and PONT, 1993; ARNAUD-FASSETTA and PROVANSAL, 1999; PROVANSAL *et al.*, 1999; ARNAUD-FASSETTA, 2000).

To help interpret the delta's Holocene depositional evolution, it is useful to determine whether the large fluctuations of ISQ values recorded in Holocene delta subsurface sections are related in some manner to climatic change. Presently, transport of Rhône and Durance sediment from intermediate and distal source terrains prevails, with deposition in the delta region of relatively low (~30%) ISQ average proportions (Table 1). At times in the past when climate was similar to the present one, available stained-grain data for core sections show comparable ISQ values (<30%). It appears that major sediment input of intermediate and distal provenance masked the supply of material dispersed from more proximal sources, with its enriched ISQ values (>40%).

We envision a considerably different depositional regime in the past, however, when increased precipitation, or changed anthropogenic activity, or both, affected the southern France-

Mediterranean region. During wetter periods, proportions of sediment input derived from sheet-flow scour and erosion of source terrains closer to the delta increased substantially (*cf.* PICHARD, 1995). At such times, a larger amount of sediment characterized by high ISQ values were dispersed from adjacent uplands (Costières, Crau and others), southern Rhône valley sectors, southern Massif Central and from intermediate (including the Massif Central, western Alps) source terrains. Thus, this periodically increased influx of more proximal material (ISQ values >40%) would have masked the fluvial sediment of more distal derivation with lower ISQ values (~30%) that was transported more frequently by the Rhône and Durance systems to the delta.

If the above scenario is correct, it should be possible to correlate strata with enriched ISQ values in the 18 delta borings numbered 105 to 129 (shown in black in Figure 9) with phases of periodically marked hydrological fluxes and increased deposition. However, some enriched ISQ subsurface intervals were probably not detected in this investigation because: (1) some of the original high ISQ layers may have been removed by natural factors (erosion by lateral migration of distributary channels) during development of the delta; and (2) our sampling intervals may have missed such layers that can be quite thin (core depths listed in Table 3). Thus, strata that include high ISQ values cannot be correlated regionally in all cases. Moreover, the relatively small number of radiocarbon-dates presently available for these and other borings in the region (*cf.* OOMKENS, 1970; TER KEURS, 1971; LABEYRIE *et al.*, 1976; JORDA *et al.*, 1991b; PROVANSAL, 1995; GENSOUS and TESSON, 1996; MORHANGE *et al.*, 1998; ARNAUD-FASSETTA, 2000) precludes establishing a firm, well-defined stratigraphy based only on enriched ISQ core layers.

For these reasons, it is useful to investigate results of the ISQ study of the subsurface section in core PI collected in Arles: there are several radiocarbon dates for this boring, and numerous samples were collected at close interval. The core includes five zones characterized by high (>40%) ISQ values (consisting of 8 of the 34 examined samples). From core top (elevation of 5 m above m.s.l.) to base, these horizons (<20 cm thick) occur at depths of ~0.4–0.7 m, 2.96 m, ~3.5 m, 10.30 m and ~18.10–18.3 m. These strata can be used for an ISQ-paleoclimate correlation (Figure 10). Climate-related depositional phases have been defined independently in earlier studies by using diverse criteria, including lithology (JORDA *et al.*, 1991b), fossils (BAZILE *et al.*, 1986; CAMBON *et al.*, 1997), geomorphology (PROVANSAL, 1995), and archaeology (JORDA and PROVANSAL, 1996). Five periods, some involving markedly altered sedimentation patterns, occurred during the Neolithic, Bronze Age, Iron Age, post-Roman to end of Middle Age, and Little Ice Age (Figure 10).

Preliminary findings that associate some high stained-grain signals for the Rhône delta with dated paleoclimatic phases are encouraging. For example, three of the four upper enriched ISQ samples of Holocene age in core PI (Figure 10, layers a,c,d) appear to match periods of altered terrigenous sedimentation identified between ~5000 years ago and the present in Provence and the southern Alps (JORDA and PROVANSAL, 1996). Layer (b) remains less clearly defined stratigraphically. Additional radiocarbon-dating information for

the lower part of the boring is needed to help confirm that the two enriched ISQ samples (>40%) at the core base (layer e) are of late Pleistocene age.

This ISQ-based stratigraphy in the Rhône delta warrants further testing with more tightly sampled and well-dated cores.

CONCLUSIONS

Most surficial samples in Holocene core sections of the modern Rhône delta plain are characterized by high proportions of clear quartz grains (~70%) and moderate proportions (~30%) of partially plus fully iron-coated quartz particles. The ISQ values recorded on the modern delta plain are similar to the material presently carried by the Rhône and Durance, the major rivers that transported most material to the delta. These findings support the conclusion of earlier workers that most surficial and subsurface sediments forming the delta plain during the Holocene were derived from intermediate and distal source terrains in the Massif Central, Alpine chain in SE France, Jura and high Swiss Alps. Of special interest in this investigation are the higher ISQ values (>40%) recorded in some modern fluvial systems and deltaic environments, and in upland settings located close to the delta proper.

Large proportions of stained quartz (to >40%), for example, are recorded on the delta's modern coast, especially in sand-rich dunes. Some of these enriched ISQ sands were derived from the landward reworking of older iron-coated quartz-rich sediment eroded from coastal exposures and perhaps also from continental shelf terrains in the Gulf of Lion south of the delta. However, even higher average ISQ values (56%) are obtained for the relict sand strand deposits of mid to late Holocene age exposed on the central to southern delta plain. This marked increase of ISQ value in these progressively older coastal sand-rich sediment on the delta plain surface suggests post-depositional *in situ* formation of additional iron coatings of quartz during the Holocene.

A different origin is proposed for the high stained-grain content (>40%) recorded in some subsurface sections of the modern delta proper: periodic reworking and preferential transfer from proximal sources of sediment that was already characterized by high ISQ values. This displacement includes material eroded from Pleistocene and older geologic terrains exposed in the lower Rhône uplands close to the delta. Such reworked sediment is generally characterized by much higher average ISQ values (67%), and is readily distinguished from most modern and Holocene deposits forming the Rhône delta plain (~30%). Evidence favoring the displacement of eroded, highly stained material from proximal highlands to the delta is the high (50%) average ISQ value that characterizes the present sediment load of the Vidourle and Gardon rivers. Flowing from the southern Massif Central, these shorter rivers carry large proportions of reworked sediments with high-ISQ values, releasing these as overbank deposits in their lower valleys and on the Rhône delta plain.

We postulate that the Gardon, Vidourle and other tributaries that flow from the proximal upland terrains transported increased proportions of ISQ-enriched sediment to the

delta primarily during periods of higher precipitation. It appears that some subsurface strata with large ISQ values (>40%) in Holocene core sections of the central and northern delta can be correlated with increased hydrological fluxes that were periodically induced by paleoclimatic phases affecting southern France. Such wetter climatic conditions and, in some instances, associated human activity, resulted in increased erosion and displacement of terrigenous sediment with high ISQ values to the Rhône delta. It was during these episodes that material from local (including Costières, Crau) and intermediate (Massif Central) terrains masked the fluvial input from more distal Rhône and Durance sediment sources.

The stained-grain method holds promise as a practical petrologic and stratigraphic tool to help identify effects of paleoclimatic fluctuations and periodically altered sediment provenance and depositional patterns. This new information will enhance our understanding of Rhône delta development during the Holocene.

ACKNOWLEDGEMENTS

We thank Dr. V. Bostan, University of Geneva, Dr. G. Ranzazzo, University of Messina, and Mr. F. Lescayer, Reserve de Camargue, for assistance in sampling surficial sediment on the Rhône delta plain and in its wetlands and adjacent regions in 1999 and 2000. Professor P.L. de Boer kindly allowed us to sample borings in the University of Utrecht core collection, with assistance by P. Anten. Dr. W.J.E. van de Graaff generously donated sediment peels of Rhône cores from Shell International Exploration and Production in Rijswijk, The Netherlands. Dr. G. Arnaud-Fassetta, University of Paris 7, sent us samples and radiocarbon dates from core PI collected in the city of Arles, and Dr. J.-C. Aloïsi, University of Perpignan, provided a set of Cefrem seafloor samples from the Gulf of Lion shelf. Dr. A. Logan, Department of Mineral Sciences, National Museum of Natural History, Smithsonian Institution, provided the analytical expertise with the microprobe and data interpretation. We also acknowledge Ms. M. Parrish, Ms. K. McGoldrick and Ms. J. Siebrase at the Museum's Department of Paleobiology, for preparation of illustrations and technical assistance with the manuscript text and tables. Reviews of this study by Drs. A. Logan, E. Reinhardt and A.G. Warne were most helpful. Funding for this project (Deltas-Global Change Program) was provided by research awards from the National Geographic Society and the Walcott Fund of the National Museum of Natural History, Smithsonian Institution.

LITERATURE CITED

- ALOISI, J.-C., 1973. Les facteurs de la sédimentation récente sur le plateau continental languedocien. *Comptes Rendus, Académie des Sciences*, 277, 145-148.
- ALOISI, J.-C. and CHARLET, J.M., 1975. Intérêt de la thermoluminescence dans l'étude de l'alimentation en quartz du Golfe du Lion. *Revue de Géographie physique et de Géologie dynamique*, 17(2), 23-38.
- ALOISI, J.-C. and DUBOUL-RAZAVET, C.A., 1974. Deux exemples de sédimentation deltaïque actuelle en Méditerranée: les deltas du Rhône et de l'Ebre. *Bulletin du Centre de Recherches de Pau*, 8(1), 227-240.

- ALOISI, J.-C.; MONACO, A. N.; THOMMERET, J., and THOMMERET, Y., 1975. Evolution paléogéographique du plateau continental Languedocien dans le cadre du Golfe du Lion. *Revue de Géographie physique et de Géologie dynamique*, 17(2), 13–92.
- ARNAUD-FASSETTA, G., 2000. Quatre mille ans d'histoire hydrologique dans le delta du Rhône. *Grafiéo*, 11, 1–229.
- ARNAUD-FASSETTA, G. and PROVANSAL, M., 1999. High frequency variations of water flux and sediment discharge during the Little Ice Age (1586–1725 A.D.) in the Rhône delta (Mediterranean France). Relationship of the catchment basin. *Hydrobiologia*, 410, 241–250.
- BAZILE, F.; BAZILE-ROBERT, E.; DEBARD, E., and GUILLERAULT, P., 1986. Le Pléistocène terminal et l'Holocène en Languedoc rhodanien; domaines continental, littoral et marin. *Revue de Géologie dynamique et Géographie Physique*, 27, 95–103.
- BELLAÏCHE, G. and MART, Y., 1995. Morphostructure, growth patterns, and tectonic control of the Rhone and Nile deep-sea fans: a comparison. *American Association of Petroleum Geologists Bulletin*, 79(2), 259–284.
- BERRYHILL, H.L., JR. (Editor), 1976. Environmental studies, south Texas outer continental shelf. *U.S. Geological Survey Report to the Bureau of Land Management* PB-251 341, 1–270.
- BERTRAND, J.P. and L'HOMER, A., 1975. Le delta du Rhône. *IXème Congrès International de Sédimentologie*, Nice, France, 1–65.
- BONIFAY, E., 1957. Age et signification des sols rouges méditerranéens en Provence. *Comptes Rendus, Académie des Sciences*, 244, 3075–3077.
- BRAVARD, J.-P.; VEROT-BOURRELY, A., and SALVADOR, P.-G., 1992. Le climat d'après les informations fournies par les enregistrements sédimentaires fluviatiles étudiés sur des sites archéologiques. *Les Nouvelles de l'Archéologie*, 50, 7–13.
- BRUZZI, C. and PROVANSAL, M., 1996. Impacts morpho-sédimentaires des tempêtes sur les côtes de Provence. *Quaternaire*, 7, 129–137.
- BUTZER, K.W., 1982. *Archaeology as Human Ecology: Method and Theory for a Contextual Approach*. Cambridge University Press, New York, 364p.
- CAMBON, G.; SUC, J.-P.; ALOISI, J.-C.; GIRESSSE, P.; MONACO, A.; TOUZANI, A.; DUZER, D., and FERRIER, J., 1997. Modern pollen deposition in the Rhône delta area (lagoonal and marine sediments), France. *Grana*, 36, 105–113.
- COLLOT, L., 1904. Pliocène et Quaternaire de la région du bas-Rhône. *Bulletin de la Société Géologique de France*, Tome IV, 400–415.
- COLEMAN, J.M. and ROBERTS, H.H., 1988. Sedimentary development of the Louisiana continental shelf related to sea level cycles: Part 1—sedimentary sequences. *Geo-Marine Letters*, 8, 63–108.
- COLOMB, E. and ROUX, R.-M., 1978. La Crau. Données nouvelles et interprétations. *Géologie Méditerranéenne*, V, 303–324.
- CORRE, J.-J., 1992. Implications of climatic changes on the Golfe du Lion. In: JEFTIC, L.; MILLIMAN, J.D., and SESTINI, G. (eds.), *Climatic Change and the Mediterranean*. London: Edward Arnold, 328–427.
- DEPARTEMENT DES BOUCHES DU RHONE, 1970. *Camargue, Etude Hydrogéologique Pédologique et de Salinité*. Chapitre 3, pp. 3.1–3.85, Arles, France.
- DORN, R.I., 1998. Iron films. In: *Rock Coatings*. Amsterdam, Netherlands: Elsevier Science B.V., 144–184.
- DUBOUL-RAZAVET, C., 1956. Contribution à l'étude géologique et sédimentologique du delta du Rhône. *Mémoires de la Société Géologique de France*, 76, 1–234.
- DURRIEU DE MADRON, X. and PANOUSE, M., 1996. Transport de matière particulaire en suspension sur le plateau continental du Golfe du Lion. Situation estivale et hivernale. *Comptes Rendus, Académie des Sciences, Paris*, 322, 1061–1070.
- ESTOUREL, C.; BROCHE, P.; MARSALEIX, P.; DEVENON, J.-L.; AUCLAIR, F., and VEHL, R., 2001. The Rhône River plume in unsteady conditions: Numerical and experimental results. *Estuarine, Coastal, and Shelf Science*, 53, 25–38.
- FRUJGET, J.F., 1992. Ecology of the lower Rhône after 200 years of human influence: a review. *Regulated Rivers: Research & Management*, 7, 233–246.
- GADEL, F. and PAUC, H., 1973. Sédimentation récente à l'embouchure du Grand Rhône: données sédimentologiques et géochimiques. *Bulletin de l'Institut Géologique du Bassin d'Aquitaine*, 14, 127–141.
- GALLOWAY, W.E., 1975. Process framework for describing the morphologic and stratigraphic evolution of deltaic depositional systems. In: BROUSSARD, M.L. (ed.), *Deltas, Models for Exploration*. Houston, Texas: Houston Geological Society, 87–98.
- GENSOUS, B. and TESSON, M., 1996. Sequence stratigraphy, seismic profiles, and cores of Pleistocene deposits on the Rhône continental shelf. *Sedimentary Geology*, 105, 183–190.
- HENSEL, P.F.; DAY, JR., J.W., and PONT, D., 1999. Wetland vertical accretion and soil elevation change in the Rhône river delta, France: The importance of riverine flooding. *Journal of Coastal Research*, 15(3), 668–681.
- INSTITUT GEOGRAPHIQUE NATIONAL, 1992. *Parc Naturel Régional de Camargue* (map 3615 IGN, scale 1:50,000, Paris, France).
- JORDA, M.; MAGNIN, F., and PROVANSAL, M., 1991a. Un témoin de l'anthropisation progressive du milieu en Basse Provence: La nappe alluviale holocène du Vallat Neuf (Étang de Berre-Bouches-du-Rhône). *Comptes Rendus, Académie des Sciences, Paris*, 312, 921–927.
- JORDA, M.; PARRON, C.; PROVANSAL, M., and ROUX, M., 1991b. Erosion et détritisme holocènes en Basse Provence calcaire. L'impact de l'anthropisation. *Physio-Géographie*, 22/23, 37–47.
- JORDA, M. and PROVANSAL, M., 1990. Terrasses de culture et bilan érosif en région méditerranéenne. *Méditerranée*, 3(4), 55–62.
- JORDA, M. and PROVANSAL, M., 1996. Impact de l'anthropisation et du climat sur le détritisme en France du sud-est (Alpes du sud et Provence). *Bulletin de la Société Géologique de France*, no. 1, 159–168.
- JORDA, M.; PROVANSAL, M., and ROYET, R., 1990. L'histoire "naturelle" d'un site de l'âge du fer sur le piémont méridional des Alpes. Le domaine de Servanne (Bouches-du-Rhône). *Gallia*, 47, 57–66.
- KRUIT, C., 1955. *Sediments of the Rhone Delta: Grain Size and Microfauna*. Mouton & Co.: 'S-Gravenhage. 106p.
- LABEYRIE, J.; LALOU, C.; MONACO, A., and THOMMERET, J., 1976. Chronologie des niveaux eustatiques sur la côte du Roussillon de 33,000 ans BP à nos jours. *Comptes Rendus, Académie des Sciences, Paris*, 282, 349–352.
- L'HOMER, A., 1975. Notice explicative de la carte géologique des Saintes-Maries-de-la-Mer au 1/50000^e, *Bureau de Recherches Géologiques et Minières*, 34p.
- L'HOMER, A., 1987. Notice explicative de la carte géologique d'Arles-Camargue au 1/50000^e, *Bureau de Recherches Géologiques et Minières*, 72p.
- L'HOMER, A., 1992. Sea-level changes and impacts on the Rhône delta coastal lowlands. In: TOOLEY, M.J. and JELGERSMA, S. (eds.), *Impacts of Sea-level Rise on European Coastal Lowlands*. T.J. Press Ltd., Cornwall, 136–152.
- L'HOMER, A.; BAZILE, F.; THOMMERET, J., and THOMMERET, Y., 1981. Principales étapes de l'édification du delta du Rhône de 7000 BP à nos jours; variations du niveau marin. *Océanis*, 7(4), 389–408.
- MEADE, R.H., 1988. Movement and storage of sediment in river systems. In: LERMAN, A. and MEYBECK, M. (eds.), *Physical and Chemical Weathering in Geochemical Cycles*. Dordrecht, The Netherlands: Kluwer Academic, 165–179.
- MORGAN, J.P. and MCINTYRE, W.G., 1959. Quaternary geology of the Bengal basin, East Pakistan and India. *Bulletin of the Geological Society of America*, 70, 319–342.
- MORHANGE, C.; PROVANSAL, M.; VELLA, C.; ARNAUD, P.; BOURCIER, M., and LABOREL, J., 1998. Montée relative du niveau de la mer et mouvements du sol à l'Holocène en basse Provence (France, Méditerranée). *Annales de Géographie*, no. 600, 139–159.
- MUNSELL COLOR, 1975. *Munsell Soil Color Charts*. Baltimore, Maryland: Kollmorgen Corp. (not paginated).
- NEWMAN, D. K., 2001. How bacteria respire minerals. *Science*, 292, 1312–1313.
- OOMKENS, E., 1970. Depositional sequences and sand distribution in the postglacial Rhône delta complex. In: MORGAN, J.P. (ed.),

- Deltaic Sedimentation Modern and Ancient*. SEPM Special Publication, 15, 198–212.
- PASQUALINI, M. and LANDURE, C., 1995. *Delta du Rhône Programme de Recherche*. Service Régional de l'Archéologie de Provence-Alpes-Côte d'Azur, Direction Régionale des Affaires Culturelles, 246p.
- PICHARD, G., 1995. Les crues sur le bas Rhône de 1500 à nos jours. Pour une histoire hydro-climatique. *Méditerranée*, no. 3.4, 105–116.
- PROVANSAL, M., 1995. The role of climate in landscape morphogenesis since the Bronze Age in Provence, southeastern France. *The Holocene*, 5, 348–353.
- PROVANSAL, M.; BRAVARD, J.-P.; BERGER, J.-F.; SALVADOR, P.-G.; ARNAUD-FASSETTA, G.; BRUNETON, H., and VEROT-BOURELLY, A., 1999. Fluctuations hydromorphologiques du Rhône et sociétés fluviales, du Léman à la mer. *Gallia*, 56, 13–52.
- PROVANSAL, M. and MORHANGE, C., 1996. Seuils climatiques et réponses morphogénétiques en basse Provence depuis 5000 ans. *Quaternaire*, 5, 113–118.
- RODITIS, J.-C. and PONT, D., 1993. Dynamique fluviale et milieux de sédimentation du Rhône à l'amont immédiat de son delta. *Méditerranée*, 78, 5–18.
- RUSSELL, R.J., 1942. Geomorphology of the Rhone delta. *Association of American Geographers Annals*, 32, 149–254.
- SHEPARD, F.P.; PHLEGER, F.B., and VAN ANDEL, T.H., 1960. *Recent Sediments, Northwest Gulf of Mexico*. American Association of Petroleum Geologists: Tulsa, OK., 349 p.
- STANLEY, D.J., 2000. Radiocarbon dating the artificially contained surfaces of the Rhône deltaic plain, southern France. *Journal of Coastal Research*, 16, 1157–1161.
- STANLEY, D.J. and CHEN, Z., 1991. Distinguishing sand facies in the Nile delta, Egypt, by stained grain and compositional component analyses. *Journal of Coastal Research*, 7, 863–877.
- STANLEY, D. and HAIT, A., 2000. Deltas, radiocarbon dating, and measurements of sediment storage and subsidence. *Geology*, 28, 295–298.
- STANLEY, D.J.; HAIT, A.K., and JORSTAD, T.F., 2000. Iron-stained quartz to distinguish Holocene deltaic from Pleistocene alluvial deposits in small core samples. *Journal of Coastal Research*, 16, 357–367.
- STANLEY, J.-D.; RANDAZZO, G., and JORSTAD, T.F., 2001. Iron-stained quartz as record of recent reworking of older sediment by natural and anthropogenic processes, Rio Grande delta, Texas. *Journal of Coastal Research*, 17, 584–598.
- STOUFF, L., 1993. La lutte contre les eaux dans les pays du bas Rhône XII^e-XV^e siècles. L'exemple du pays d'Arles. *Méditerranée*, 3(4), 57–68.
- SUANEZ, S. and PROVANSAL, M., 1996. Morphosedimentary behaviour of the deltaic fringe in comparison to the relative sea-level rise on the Rhone delta. *Quaternary Science Reviews*, 15, 811–818.
- TER KEURS, D., 1971. Development of ostracoda and foraminifera assemblages in transgressive/regressive sequences. *Bulletin du Centre de Recherche, Pau-SNPA*, 5: 257–283.
- TRICART, J. and BRAVARD, J.-P., 1991. L'aménagement des trois plus grands fleuves européens: Rhin, Rhône et Danube. Problèmes et méfaits. *Annales de Géographie*, 561–562 and 668–713.
- VAN ANDEL, T.H., 1955. Sediments of the Rhône delta; sources and deposition of heavy minerals. *Verhandelingen van het Koninklijk Mijnbouwkundig Genootschap*, Geological series, no. 15, 502–555.
- VAN HOUTEN, F.B., 1968. Iron oxides in red beds. *Geological Society of America Bulletin*, 79, 399–416.
- VAN HOUTEN, F.B., 1973. Origins of red beds, a review 1961–1972. In: *Annual Review of Earth & Planetary Sciences*, 1, 39–61.
- VELLA, C.; BOURCIER, M., and PROVANSAL, M., 1998. Montée du niveau marin et sédimentation holocène sur la marge orientale du delta du Rhône, Provence, France. *Bulletin de la Société Géologique de France*, 169, 403–414.
- VIVIAN, H., 1989. Hydrological changes of the Rhône river. In: PETERS, G.E. (ed.). *Historical Change of Large Alluvial Rivers: Western Europe*. Chichester: John Wiley & Sons, 57–77.
- YAPP, C., 2001. Rusty relics of earth History: Iron (III) oxides, isotopes, and surficial environments. In: JEANLOZ, R.; ALBEE, A.L., and BURKE, K.C. (eds.), *Annual Review of Earth and Planetary Sciences*, Palo Alto, California, 29, pp. 165–199.

