

## How many species are there in Hawaii?<sup>1</sup>

LUCIUS G. ELDRIDGE<sup>2</sup> and SCOTT E. MILLER (Department of Natural Sciences, Bishop Museum, P.O. Box 19000A, Honolulu, Hawaii 96817, USA)

### Introduction

The Hawaiian Islands are the most isolated island group in the world. The 8 main southeastern islands, with their sequentially younger geological ages, great physiographic and climactic variation, are ideal natural laboratories for evolutionary and ecological research (see Howarth 1990, Kay 1994 for further background). Although the Hawaiian biota has been subjected to systematic study since shortly after the founding of Linnaean nomenclature, these results are scattered through publications and collections around the world. Moreover, many gaps remain in the knowledge of Hawaii's biota, especially for the smaller organisms such as viruses, bacteria, freshwater algae, protists, and soil invertebrates other than insects and mollusks. This paper is a first approximation by Bishop Museum's Hawaii Biological Survey to answer the frequently asked question: "How many species are there in Hawaii?"

In compiling this list, we have collected and assembled data and wisdom of the staff of Bishop Museum, libraries of the Bishop Museum and the University of Hawaii, and many colleagues in Hawaii and around the world. The marine portions draw heavily from the ongoing project *Reef and shore fauna of Hawaii*. The terrestrial invertebrate data derive primarily from an invertebrate survey pilot project supported by the John D. and Catherine T. MacArthur Foundation in the late 1980s. An early version of the terrestrial and freshwater portions of the list was circulated at the Hawaii Conservation Biology Initiative annual meeting in 1992. The interest of that group prompted us to continue the effort. We know that the present list is incomplete in several ways: 1) we expect that some literature has been missed; 2) we know that additional information, including unpublished new species, exist in museum collections; and 3) further fieldwork will produce additional records. The Hawaii Biological Survey is creating comprehensive databases for basic taxonomic, distributional, and biological information on all the organisms occurring in Hawaii, so we welcome additions and corrections to this list.

Classification of animals generally follows Parker (1982), with the use of kingdoms in the sense of Margulis & Schwartz (1988). Some artificial categories of convenience are used (for examples, marine mollusks). Freshwater categories generally include brackish water dwellers. Viruses and bacteria are not included because too few data are available, although the plant parasitic species were reviewed by Raabe *et al.* (1981). Human parasites are also not included. Unless noted, all numbers represent described species reported in the literature and often under-represent the eventual total which can be expected based on further exploration. On the other hand, there may be many more names in the literature for some groups because the taxa have not been reviewed or revised to eliminate the synonyms. The numbers from the *Reef and shore fauna* manuscripts include previously unpublished records.

The term endemic is problematic. In the terrestrial environment it is easy to determine the boundaries of the Hawaiian Islands, but the fact that a species is known only from Hawaii may mean only that it has not yet been reported elsewhere. This is especial-

---

1. Contribution No. 1995-002 to the Hawaii Biological Survey.

2. Executive Secretary, Pacific Science Association, P.O. Box 17801, Honolulu, Hawaii 96817, USA

## 4 BISHOP MUSEUM OCCASIONAL PAPERS: NO. 41, 1995

ly a problem in groups that require specialized sampling techniques and that may have been better studied in Hawaii than in many other places (e.g., planktonic crustaceans, soil mites). In the marine environment this problem is worse because the ocean provides a medium for movement throughout the Indo-west Pacific. Moreover, it is difficult to determine boundaries. Thus, we expect that most of the taxa currently listed as endemic to Hawaii's marine environment will ultimately prove to be more widely distributed.

**Definitions**

For the marine environment "Hawaii" includes the coastal areas to the deepest waters, including all the waters surrounding all of the islands of the archipelago.

*Nonindigenous species* (NIS) are those species that do not naturally occur in the Hawaiian Islands and have arrived either accidentally or intentionally through biological control efforts, aquaculture imports, etc. These species have also been referred to as alien, exotic, adventive, or introduced species (Frank & McCoy 1990).

*Indigenous species* are those species which naturally occur in the islands but are not endemic to the islands (i.e., they also occur naturally elsewhere).

For birds, *visitors* are nonresident species; *migratory* birds follow a repeatable pattern of occurrence in the islands.

*Subfossil/fossil* species in Hawaii are used in the context of birds and bats that became extinct prior to the arrival of Captain Cook. Invertebrate fossils discovered through core sampling, archaeological digs, and other techniques have not been included in the data tabulated below.

*Extinct* species are those forms that no longer exist in their natural habitat. It is difficult at best to attempt a tabulation of how many of the species originally described from the Hawaiian Islands have become extinct through natural processes or with the advent of humans and the concomitant destruction of native habitats and associated species through time. Data on known extinct vertebrates are included in the notes, but not in Table 1.

**Table 1. Estimates of Numbers of Species of the Hawaiian Biota**

Taxon	Total	Endemic	NIS	Reference(s)
<b>PROTOCTISTA</b>				
Algae				
freshwater	120+	2	?	Vis <i>et al.</i> 1994; I.A. Abbott, pers. comm.
marine	470	?	5	I.A. Abbott, pers. comm.
Myxomycota	101	0	?	Eliasson, 1991
Protozoa				
Foraminifera	1000+	?	0	Phillips, 1977; Chave, 1987
Other protozoa	42	2?	0	Ball, 1963; Phillips, 1977; Larsen & Patterson, 1990; Boyko, 1994
<b>FUNGI</b>				
Fungi	1300	?	?	Baker & Goos, 1972; Goos, 1977; Kohlmeyer, 1969
Lichens	723	240	?	Smith, 1991, 1993
<b>PLANTAE</b>				
Bryophytes				
mosses	244	112	11	Hoe, 1979
liverworts	220	?	?	Miller & Whittier, 1990

## RECORDS OF THE HAWAII BIOLOGICAL SURVEY FOR 1994

5

Taxon	Total	Endemic	NIS	Reference(s)
Fern Allies	24	7	2	W.H. & F. Wagner, pers. comm.
Ferns	205	124	27	W.H. & F. Wagner, pers. comm.
Gymnosperms	10?	0	10?	W. Appleby, pers. comm.
Angiosperms	1894	850	861	Wagner <i>et al.</i> , 1990; HBS Records for 1994*
<b>ANIMALIA</b>				
<b>Phagocytellozoa</b>				
Placozoa	1	0	0	Pearse, 1989
<b>Parazoa</b>				
Porifera				
marine	84	24	4+	Bergquist, 1977; Chave & Jones, 1991
freshwater	1	0	1	Svihla, 1941
<b>Eumetazoa</b>				
Cnidaria				
Hydroida				
marine	23	4	1	Cooke, 1977
deepwater	51	31	0	Nutting, 1905
freshwater	2	0	2	Matthews, 1966
Siphonophora	1	0	0	Eldredge & Devaney, 1977
Chondrophora	2	0	0	Eldredge & Devaney, 1977
Scyphozoa	13	0	5	Devaney & Eldredge, 1977; Cooke, 1984; Burch & Burch, 1995
Octocorallia	105	17	1	Devaney, 1977b; Grigg & Bayer, 1976; Muzik, 1978; Versevelt & Bayer, 1988; Bayer, 1990; Chave & Jones, 1991
Zoantharia	23	5	1	Cutress, 1977
Zoanthiniara	7	4	0	Walsh & Bowers, 1977
Scleractinia	99	12	0	Maragos, 1977, 1995; Grigg <i>et al.</i> , 1981; Cairns, 1984
Antipatharia	15	1	0	Grigg & Opresko, 1977; Chave & Jones, 1991
Ctenophora	10	0	1	Devaney, 1977a; R. Galt, pers. comm
Platyhelminthes				
marine	41	20	0	Poulter, 1987
freshwater/terrestrial	8	4	4	Hyman, 1939; Kawakatsu & Mitchell, 1984; Kawakatsu <i>et al.</i> , 1984
parasitic	519	371+	0	Yamaguti, 1968a, 1968b, 1970; Kaneko <i>et al.</i> , 1988; Dailey <i>et al.</i> , 1992
Nemertinea				
marine	26	5?	0	Coe, 1947; Devaney & Eldredge, 1987; J. Norenburg, pers. comm.
freshwater/terrestrial	2	0	2	Howarth & Moore, 1983; J. Norenburg, pers. comm.
Gnathostomulida	8	4	0	Sterrer, 1991
Rotifera				
marine	3	0	0	Hope, 1987
freshwater	27	?	0	P.N. Turner, pers. comm
Nematoda				
marine	54	23	0	Hope, 1987
plant/soil	127	?	?	Olveira, 1940; E.M. Noffsinger & E.P. Caswell (unpubl. 1989)
zooparasitic	33+	?	?	Chapin, 1925; Deardorff <i>et al.</i> , 1982; Deardorff, 1987; McKenzie & Davidson, 1989

## 6 BISHOP MUSEUM OCCASIONAL PAPERS: NO. 41, 1995

Taxon	Total	Endemic	NIS	Reference(s)
Priapulida	1	0	0	J. Bailey-Brock, pers. comm..
Mollusca				
Gastropoda				
marine	572	119	4	Kay, 1979; Kay & Palumbi, 1987
terrestrial	831±	759	50±	R.H. Cowie <i>et al.</i> , in press
freshwater	37±	7	30±	R.H. Cowie, pers. comm.
Bivalvia				
marine	129	66	1	Kay, 1979; Kay & Palumbi, 1987
freshwater	1	0	1	Burch, 1978; HBS Records for 1994
Polyplacophora	4	3	0	Kay, 1979
Scaphopoda	3	0	0	Kay, 1979
Cephalopoda	73	2	0	Kay, 1979; Berry, 1914; Roper & Young, 1975; Roper <i>et al.</i> , 1984; Young, 1991
Aplacophora	6	0	0	Kay, 1979
Annelida				
Oligochaeta				
marine	26	11	3?	Bailey-Brock, 1987; Erséus & Davis, 1989; Erséus, 1990
terrestrial	20	0	20	Nakamura, 1992; S.W. James, unpubl.
Archannelida	2	0	0	Bailey-Brock, 1987
Hirudinea				
marine	3	0	0	Bailey-Brock, 1987; Choy <i>et al.</i> , 1989
terrestrial	3	0	3	Moore, 1946
Polychaeta				
marine	250+	69+	4	Pettibone, 1986; Paxton & Bailey-Brock, 1986; Bailey-Brock & Hartman, 1987; Bailey-Brock, 1990, 1991; Fauchald, 1992
freshwater	1	0	0	Van Zwaluwenburg, 1948
Pogonophora	2	2	0	Southward, 1980
Echiura	6	0	0	Edmonds, 1987
Sipuncula	14	0	0	Edmonds, 1987
Arthropoda				
Insecta (nonmarine)				
Blattaria	19	0	19	Nishida, 1994
Coleoptera	1987	1358	597	Nishida, 1994; HBS Records for 1994
Collembola	168	95	71	Nishida, 1994
Dermaptera	24	9	14	Nishida, 1994
Diptera	1426	1048	631	Nishida, 1994
Heteroptera	316	214	96	Nishida, 1994; HBS Records for 1994
Homoptera	689	385	299	Nishida, 1994; HBS Records for 1994
Hymenoptera	1283	654	605	Nishida, 1994; HBS Records for 1994
Lepidoptera	1153	950	192	Nishida, 1994; HBS Records for 1994
Mallophaga	102	5	44	Nishida, 1994
Neuroptera	60	51	8	Nishida, 1994
Odonata	41	31	8	Nishida, 1994; HBS Records for 1994
Orthoptera	290	263	26	Nishida, 1994; Otte, 1994
Psocoptera	135	89	42	Nishida, 1994
Thysanoptera	147	29	116	Nishida, 1994
Remaining orders	62	6	45	Nishida, 1994
Insecta (marine)	4	1	0	Herring, 1961, 1965; N.L. Evenhuis, pers. comm.
Araneae	198	119	71	Nishida, 1994; HBS Records for 1994
Chilopoda	24	12	12	Nishida, 1994
Diplopoda	26	16	9	Nishida, 1994
Palpigrada	1	0	1	Nishida, 1994

## RECORDS OF THE HAWAII BIOLOGICAL SURVEY FOR 1994

7

Taxon	Total	Endemic	NIS	Reference(s)
Paupoda	2	0	1	Nishida, 1994
Pseudoscorpionida	12	12	0	Nishida, 1994
Shizomida	1	0	1	Nishida, 1994
Scorpionida	1	0	1	Nishida, 1994
Crustacea				
Branchiopoda	4	0	0	Ueno, 1936
Ostracoda				
marine	31	?	0	J.C. Holden, MS; Kornicker, 1976; Danielopol & Hartmann, 1986; Hartmann, 1991
freshwater	1	?	?	Vávra, 1906
Copepoda	100	?	?	L.G. Eldredge, MS; Motoda, 1963; Lewis, 1966, 1967; Humes, 1976; Wells, 1986; Kunz, 1993; HBS Records for 1994
Cirripedia	58	?	0	W.A. Newman, MS; Pilsbry, 1907; Boschma, 1953
Stomatopoda	17	11	1	L.G. Eldredge, MS
Mysidacea	14	?	0	L.G. Eldredge, MS; Ortmann, 1906
Tanaidacea	8	7	0	M. Miller, MS
Isopoda				
marine	27	?	?	M. Miller, MS; Richardson, 1906
terrestrial	51	15	26	Nishida, 1994
Amphipoda				
marine	172	?	8	Barnard, 1970; Brusca, 1973, 1978
terrestrial	9	6	3	Nishida, 1994
Euphausiacea	27	?	0	Ortmann, 1906; Brinton, 1962
Decapoda				
Penaeidae	20	0	0	L.G. Eldredge, MS
Sergestidae	10	0	0	L.G. Eldredge, MS
Stenopodidea	5	4	1	L.G. Eldredge, MS
Caridea	149	?	1	L.G. Eldredge, MS
Astacidea	1	0	1	Eldredge, 1994
Thalassinidea	8	?	1	L.G. Eldredge, MS
Palinuridea	13	2	0	L.G. Eldredge, MS
Anomura	43	?	0	L.G. Eldredge, MS
Brachyura	189	?	7	L.G. Eldredge, MS; HBS Records for 1994
Acari				
marine	19	17	2	Nishida, 1994
terrestrial	517	97	337	Nishida, 1994; HBS Records for 1994
Pycnogonida	12	?	0	J.H. Stock, MS; Child, 1972
Tardigrada	22	1	19	Nishida, 1994; McInnes, 1994
Phoronida	3	0	0	Emig & Bailey-Brock, 1987
Bryozoa	150+	?	?	Soule <i>et al.</i> , 1987
Entoprocta	?	?	?	Soule <i>et al.</i> , 1987
Brachiopoda	3	0	0	Emig, 1987; B.L. Burch, pers. comm.
Chaetognatha	8	1	0	Alvarino, 1978; Pierrot-Bults & Nair, 1991
Echinodermata				
Crinoida	16	15	0	L.G. Eldredge, MS; Clark, 1949
Asteroidea	82	53	0	L.G. Eldredge, MS; Clark, 1949
Echinoidea	75	35	0	L.G. Eldredge, MS; Clark, 1949
Holothuroidea	48+	19	0	L.G. Eldredge, MS; Clark, 1949
Ophiuroidea	57	28	0	D.M. Devaney & A.N. Baker, MS; Clark, 1949
Hemichordata	4	1	0	L.G. Eldredge, MS; Hadfield & Young, 1983; 1 undescribed sp.
Chordata				
Cephalochordata	1	0	0	Eldredge, 1967

Taxon	Total	Endemic	NIS	Reference(s)
Urochordata (Tunicata)				
Thaliacea	24	0	0	Metcalf & Hopkins, 1919; Yount, 1954; Soest, 1974a, 1974b
Larvacea	2	0	0	Taguchi, 1982; Kitalong, 1986
Ascidiacea	45	7?	2?	D.P. Abbott, MS; Tokioka, 1967
Pisces				
marine	1150+	134+	33	Eldredge, 1994; B. Mundy, pers. comm.
freshwater	45	5	40	Eldredge, 1994; Maciolek, 1984, Devick <i>et al.</i> , 1992
Amphibia	4	0	4	Vitousek <i>et al.</i> , 1987; C. Kishinami, pers. comm.
Reptilia	18	0	18	Vitousek <i>et al.</i> , 1987; Eldredge, 1994; C. Kishinami, pers. comm.
Aves	274	60	46	R.L. Pyle, pers. comm.
Mammalia				
terrestrial	20	1	19	Tomich, 1986
marine	24	0	0	Tomich, 1986
<b>totals</b>	<b>21,383</b>	<b>8,759</b>	<b>4,532</b>	

\* HBS Records for 1994 = articles in *Bishop Museum Occasional Papers* 41 and 42 of the Records of the Hawaii Biological Survey for 1994.

### Notes

**Algae:** Marine algae numbers are from an identification manual in preparation by I.A. Abbott & W. Magruder; numbers for freshwater and terrestrial species are only estimates, except for Vis *et al.* (1994) who reported 34 species; all other existing literature is outdated and unreliable (e.g., MacCaughey 1918).

**Protozoa:** Chave (1987) noted more than 1000 species in Hawaiian waters; Philipps (1977) reported on more than 400 species from depths less than 100 ft; Larsen & Patterson (1990) reported 31 benthic marine flagellate species, including 2 new species.

**Fungi:** Baker & Goos (1972), following Parris (1940) recorded 683 species, including Myxomycota (slime molds) but not lichens; Goos (1977) recorded 900 species of Fungi Imperfecti; Baker (1977) noted 1591 species of fungi from Hawaii and the central Pacific islands. Undoubtedly, the true number is much higher because the fungi are poorly studied in Hawaii (see also Goos & Gowing 1992). Hawksworth (1994) suggested that in a given geographical area there are about 6 times as many fungi in all habitats as there are native and naturalized plant species, which suggests that there may be 12,000 species of fungi. Endemism is very low (Baker & Goos 1972, Baker 1977, Goos 1977). Kohlmeier (1969) recorded 27 species of marine fungi including 2 nonindigenous species.

**Lichens:** Smith (1991) recorded 723 species, suggesting that there may be more than 800 species (his 1991 numbers include taxonomic changes published in Smith 1993). He recorded 240 species as supposedly endemic and suggested that the true number may be lower; 1 genus (*Ramalinopsis*) is endemic. See also Stenroos & Smith (1993).

**Ferns** (Polypodiophyta) and **fern allies** (Psilophyta and Lycophyta): Only native and naturalized (e.g., growing in the wild) species are listed; more than 200 additional species are cultivated (G. Staples, pers. comm.)

**Angiosperms:** Only native and naturalized species are listed; on the order of 8000 additional species are cultivated in agricultural settings and home gardens (G. Staples, pers. comm.).

**Porifera:** Of 80 species listed, 5 are known only from depths greater than 100 m, 5 recorded only from the Waikiki Aquarium, and 8 that might be considered doubtful species (Bergquist 1977).

**Cnidaria: Octocorallia:** Of 105 species listed, 4 are shallow water (Devaney 1977b),

## RECORDS OF THE HAWAII BIOLOGICAL SURVEY FOR 1994 9

remaining are 1 species of bamboo coral (Muzik 1978) and 101 species of deep-water gorgonians (Grigg & Bayer 1976, Versevelt & Bayer 1988, Bayer 1990).

**Cnidaria: Scleractinia:** Of 98 species listed, 42 found in shallow water (Maragos 1977), 49 in deeper water (Cairns 1984), 3 from northwest islands (Grigg *et al.* 1981), 5 new records in Maragos (1995); of 91 published species, 50 are exclusively ahermatypic, 37 are exclusively hermatypic, 4 are facultative species (Cairns 1984).

**Platyhelminthes (Parasitic):** Of 511 species listed, 147 monogenetic species including 132 species described from 121 species of Hawaiian fishes (Yamaguti 1968b). Of 357 digenetic species, 314 species including 227 species described from Hawaiian fishes. Also 30 species not cited (Yamaguti 1970), 13 from Yamaguti (1965), 7 species of cestodes (Yamaguti 1968a), and non-indigenous, marine, cage-cultured, freshwater tilapia with 1 species of marine monogenean (Kaneko *et al.* 1988).

**Nemertinea:** Of the 28 species in the table, 6 species from Coe (1947), 7 benthic and 12 pelagic undescribed/unreported species, 2 introduced terrestrial species (J. Norenburg, pers. comm.) plus 1 parasitic (Humes 1942).

**Annelida: Oligochaeta:** Of 26 species listed, 23 Tubificidae (Erséus & Davis 1989); 1 new species by Erséus (1990); the other 2 are unidentified species (Bailey-Brock 1987).

**Annelida: Polychaeta:** Approximately 100 additional species are known, but not yet reported from Hawaiian waters (J. Bailey-Brock, pers. comm.).

**Nematoda: zooparasitic:** The figure given is very incomplete. Of 33 species, 22 are from intermingling herds of mammals on Molokai (McKenzie & Davidson 1989); 2 from monk seals (Chapin 1935), 8 larval forms from fishes (Deardorff *et al.* 1982), 1 parasitic record from an elasmobranch (Deardorff 1987).

**Mollusca: Gastropoda:** There are 763 nomenclaturally correct native land species; all endemic except 2–4 indigenous but not endemic species (Cowie *et al.*, in press), approximately 50 nonindigenous species (R.H. Cowie, pers. comm.); 7 endemic freshwater snails, approximately 30 nonindigenous species (R.H. Cowie, pers. comm.).

**Mollusca: Cephalopoda:** Of 73 species in the table, only 4 considered benthic (Kay 1979; Roper *et al.* 1984); for other pelagic species see Roper & Young (1975) and Young (1991).

**Crustacea: Branchiopoda:** Four species from Mauna Kea at elevations from 6500–13,700 ft [1980–4175 m] (Ueno 1936).

**Crustacea: Ostracoda:** Numerous species; J.C. Holden (MS) made identifications to only genera, he noted more than 100 fossil species; 30 benthic species (Kornicker 1976, Danielopol & Hartmann 1986, Hartmann, 1991); 1 freshwater species (Vávra 1906).

**Crustacea: Copepoda:** The 92 species listed include planktonic, benthic, and ectoparasitic forms; 5 interstitial species (Wells 1986).

**Crustacea: Cirripedia:** Of 58 species in the table, 11 are from deep water including 8 new species (Pilsbry 1907); 46 shallow-water species includes 3 acrothoracican species (W.A. Newman, MS key); 1 rhizocephalan parasite (Boschma 1953).

**Crustacea: Isopoda:** Of 27 marine species, 3 were collected in deep water (Richardson 1906); 6 species are parasitic epicarideans (Danforth 1970); of 51 terrestrial species 15 are endemic and 25 nonindigenous species.

**Crustacea: Amphipoda:** Of 172 species listed, 121 gammarideans (Barnard 1970), 4 gammarideans (Brusca 1973), 7 caprellids, 40 hyperiids (Brusca 1973, 1978).

**Entoprocta:** Soule *et al.* (1987) provided only unidentified generic descriptions; 55 species reported from Kaneohe Bay (Dade & Honkalehto 1986).

**Chaetognatha:** Seven pelagic species from distribution maps (Pierrot-Bults & Nair 1991); 1 benthic species described from Kure Atoll, Hawaii (Alvarino 1978).

**Echinodermata: Crinoidea:** Of 16 species listed in the table, 11 described as new species by Clark (1908); no shallow-water forms.

**Echinodermata: Asteroidea:** Of 82 species listed, 18 found in shallow water and 64 in deeper water; 1 species and 2 subspecies of shallow-water forms considered endemic (Devaney & Baker MS); 50 species endemic to deeper waters described as new (Fisher 1906).

**Echinodermata: Echinoidea:** Of 75 species, 21 found in shallow water (Eldredge MS), 53 in deeper water; *Cylpeaster eurypetalus* known only from Pearl and Hermes Reef, and Midway and Kure atolls; 35 new species described from *Albatross* material.

**Echinodermata: Holothuroidea:** Of 48+ species in the table, 24 found in shallow water (Eldredge, MS), 24 in deeper water; 3 shallow-water and 1 deep-water for considered endemic (Clark & Rowe 1971); Fisher (1907) described 19 new species out of 46 reported; reports (unpublished) of additional species are known.

**Echinodermata: Ophiuroidea:** Of 57 species, 19 found in shallow water (Devaney & Baker MS), 38 in deeper water; 6 shallow-water and 22-deep-water forms endemic.

**Chordata: Urochordata: Thaliacea:** 19 species of Salpidae—cosmopolitan oceanic planktonic organisms in the circumglobal warm water zone (Yount 1958); 1 species of *Pyrosoma* (Metcalf & Hopkins 1919).

**Chordata: Pisces:** The 1150+ marine species listed include all species from all depths to 200 miles from shore—no freshwater, no undescribed, and no doubtful species are included; 1285 species when all conditions provided (B. Mundy, pers. comm.); Randall (1992) listed 536 species of shore fishes, with 25% endemism.

**Chordata: Aves:** The 274 species listed is an all-inclusive contemporary list: 60 resident native species, 46 resident nonindigenous species, 13 breeding visitor species, 155 nonbreeding visitor species. There are also 16 species extinct since Captain Cook, 35+ extinct before Captain Cook (subfossil), and 150± nonindigenous species not established bringing the total number of species known to occur or be sighted in Hawaii to 475± (R.L. Pyle, pers. comm). [Pyle (1992) listed 131 bird species; 53 nonindigenous species.] Thirty-two fossil species have been described by Olson & James (1991) and James & Olson (1991).

**Chordata: Mammalia:** Of 20 species of terrestrial mammals 19 are free-ranging species (the horse is no longer feral) and 1 species of bat (an additional undescribed subfossil species of bat is known; F.G. Howarth, pers. comm.); of 24 marine species, 2 are littoral marine (seals) and 22 are pelagic (whales and dolphins) (Tomich 1986).

### Summary

From literature and unpublished sources, approximately 21,383 species have been recorded from the Hawaiian Islands and surrounding waters. Of these, 8,759 are endemic to the Hawaiian Islands, and 4,532 are nonindigenous species of protists, fungi, plants, and animals. Of these approximately 15,000 species are terrestrial, 300 are found in freshwater, and 5,500 are marine-inhabiting. Endemism varies from very low (e.g., lower plants and many marine organisms) to very high (e.g., insects and snails). However, many gaps in the knowledge of Hawaii's biota (especially the smaller organisms—viruses, bacteria, protists, freshwater algae, etc.), still remain.

### Acknowledgements

The authors thank the following people and institutions for their assistance and support in compiling the data in this article: A. Allison, W. Appleby, B.L. Burch, T.A. Burch, R.H. Cowie, N.L. Evenhuis, D.R. Herbst, W.J. Hoe, F.G. Howarth, C. Imada, C. Kishinami, G.M. Nishida, D.A. Polhemus, R.L. Pyle, G. Staples, and A. Suzumoto (all from Bishop Museum, Honolulu, USA); I.A. Abbott, J. Bailey-Brock, M. Hadfield, C.W. Smith, and G. Wong (all from University of Hawai at Manoa, Honolulu, USA); E.P.



## RECORDS OF THE HAWAII BIOLOGICAL SURVEY FOR 1994 11

Caswell, E.M. Noffsinger (University of California, Davis, California, USA), R. Galt (California State University, Long Beach, California, USA), S.W. James (Maharishi International University, Fairfield, Iowa, USA), D.L. Hawksworth, P.M. Kirk (International Mycological Institute, Egham, Surrey, UK), G. Mason (University of Otago, Dunedin, New Zealand), D. Farr (U.S. Department of Agriculture, Beltsville, Maryland, USA), B. Mundy (National Marine Fisheries Service, Honolulu, Hawaii, USA), J. Norenberg and W.L. Wagner (Smithsonian Institution, Washington, D.C., USA), P.N. Turner (Christopher Newport University, Newport News, Virginia, USA); the staff of the libraries at the Bishop Museum and the University of Hawaii at Manoa; the National Science Foundation; and the John D. and Catherine T. MacArthur Foundation.

**References**

- Alvarino, A.** 1978. *Spadella gaetanoi*, a new benthic chaetognath from Hawaii. *Proc. Biol. Soc. Wash.* **91**(3): 650–57.
- Bailey-Brock, J.H.** 1987. Phylum Annelida, p. 213–16. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- . 1990. *Polydora nuchalis* (Polychaeta: Spionidae), a new Hawaiian record from aquaculture ponds. *Pac. Sci.* **44**(1): 81–87.
- . 1991. Tubeworms (Serpulidae, Polychaeta) collected from sewage outfalls, coral reefs and deep water off the Hawaiian Islands. *Bull. Mar. Sci.* **48**(2): 198–207.
- . & **O. Hartman.** 1987. Class Polychaeta, p. 216–454. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- Baker, G.E.** 1977. The prospect for mycology in the Central Pacific. *Harold L. Lyon Arboretum Lecture* **8**, 51 p.
- . & **R.D. Goos.** 1972. Endemism and evolution in the Hawaiian biota: Fungi, p. 409–31. In E. A. Kay, ed., *A natural history of the Hawaiian Islands: selected readings*. University of Hawaii Press, Honolulu.
- Ball, C.H.** 1963. *Cephaloidophora carpilodei* n. sp. and *C. pinquis* n. sp., gregarines of a xanthid crab in Hawaii. *J. Protozool.* **10**(3): 321–27.
- Barnard, J.L.** 1970. Sublittoral Gammaridea (Amphipoda) of the Hawaiian Islands. *Smithson. Contrib. Zool.* **3**, 286 p.
- Bayer, F.M.** 1990. A new isidid octocoral (Anthozoa: Gorgonacea) from New Caledonia, with descriptions of other new species from elsewhere in the Pacific Ocean. *Proc. Biol. Soc. Wash.* **103**(1): 205–28.
- Bergquist, P.R.** 1977. Porifera, p. 53–69. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- Berry, S.** 1914. The Cephalopoda of the Hawaiian Islands. *Bull. Bur. Fish.* **23**: 255–362.
- Boschma, H.** 1953. The Rhizocephala of the Pacific. *Zool. Meded.* **32**(17): 185–210.
- Boyko, C.B.** 1994. Catalog of recent type specimens in the Department of Invertebrates, American Museum of Natural History. I. Micro-invertebrates (phyla Sacromastigophora, Gnathostomula, Gastrotricha, Rotifera, and Tardigrada). *Am. Mus. Novit.* **3106**, 44 p.

## 12 BISHOP MUSEUM OCCASIONAL PAPERS: NO. 41, 1995

- Brinton, E.** 1962. The distribution of Pacific euphausiids. *Bull. Scripps Inst. Oceanogr.* **8**(2): 51–279.
- Brusca, G.J.** 1973. Pelagic Amphipoda from the waters near Oahu, Hawaii, excluding the family Scinidae. *Pac. Sci.* **27**(1): 8–27.
- . 1978. Contribution to the knowledge of hyperiid amphipods of the family Scinidae from near Hawaii, with a description of a new species, *Scina hawaiiensis*. *Pac. Sci.* **32**(3): 281–92.
- Burch, B.L.** 1978. Asian clam, *Corbicula*, threatens Hawaii. *The Nautilus* **92**(1): 54–55.
- Burch, B.L. & T.A. Burch.** 1995. New Hawaiian records for *Stephanoscyphus simplex* Kirkpatrick (Cnidaria: Scyphozoa). *Bishop Mus. Occas. Pap.* **42**: 53.
- Cairns, S.D.** 1984. New records of ahermatypic corals (Scleractinia) from the Hawaiian and Line Islands. *Occas. Pap. B. P. Bishop Mus.* **25**(10): 1–30.
- Chapin, E.A.** 1925. Descriptions of new internal parasites. *Proc. U.S. Natl. Mus.* **68**(2603): 1–4.
- Chave, E.H.** 1987. Common living benthic Foraminifera in Mamala Bay, Hawaii, with descriptions of two new species. *Bishop Mus. Occas. Pap.* **27**: 25–72.
- . & **A.T. Jones.** 1991. Deep-water megafauna of the Kohala and Haleakala slopes, Alenuihaha Channel, Hawaii. *Deep-Sea Res.* **38**(7): 781–803.
- Child, C.A.** 1972. A new species of *Anoropallene* (Pycnogonida) from the Hawaiian Islands. *Proc. Biol. Soc. Wash.* **85**(11): 147–50.
- Choy, B.K., G.H. Balazs & M. Dailey.** 1989. A new therapy for marine turtles parasitized by the piscicolid leech, *Ozobranchus brachiatus*. *Herp. Rev.* **20**(4): 89–90.
- Clark, A.H.** 1949. Ophiuroidea of the Hawaiian Islands. *B.P. Bishop Mus. Bull.* **195**, 133 p.
- Clark, A.M. & F.W.E. Rowe.** 1971. *Monograph of shallow-water Indo-west Pacific echinoderms*. British Museum (Natural History), London. 234 p.
- Coe, W.R.** 1947. Nemerteans of the Hawaiian and Marshall Islands. *Occas. Pap. B. P. Bishop Mus.* **19**(3): 101–06.
- Cooke, W.J.** 1977. Order Hydroida, p.71–104. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- . 1984. New scyphozoan records for Hawaii: *Anomalorhiza shawi* Light, 1921, and *Thysanostoma loriferum* (Ehrenberg, 1835); with notes on several other rhizostomes. *Proc. Biol. Soc. Wash.* **97**(3): 583–88.
- Cowie, R.H., N.L. Evenhuis & C.C. Christensen.** 1995. Catalog of the native land and freshwater molluscs of the Hawaiian Islands. *Bishop Mus. Bull. Zool.* **3**: in press.
- Cutress, C.E.** 1977. Orders Corallimorpharia, Actiniaria, Ceriantharia, p. 130–147. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 1. Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- Dade, W.B. & T. Honkalehto.** 1986. Common ectoproct bryozoans of Kaneohe Bay, p. 52–65. In: Jokieli, P.L., R.H. Richmond & R.A. Rogers, eds., *Coral reef population biology*. Hawaii Institute of Marine Biology Technical Report 37.
- Dailey, M.D., M.L. Fast & G.H. Balazs.** 1992. A survey of the Trematoda (Platyhelminthes: Digenea) parasitic in green turtles, *Chelonia mydas* (L.) from Hawaii. *Bull. South. Calif. Acad. Sci.* **91**(2): 84–91.

## RECORDS OF THE HAWAII BIOLOGICAL SURVEY FOR 1994 13

- Danforth, C.G.** 1970. Epicaridea (Isopoda) of Hawaii. *Bull. South. Calif. Acad. Sci.* **69**(1): 27–31.
- Danielopol, D.L. & G. Hartmann.** 1986. Ostracoda. Part 1: stygobiont Ostracoda from inland subterranean waters, p. 265–94. *In: Botosaneanu, L., ed., Stygo fauna mundi.* E.J. Brill, Leiden.
- Deardorff, T.L.** 1987. Redescription of *Pulchrascaris chiloscylli* (Johnston and Mawson, 1951) (Nematoda: Anisakidae), with comments on species in *Pulchrascaris* and *Terranova*. *Proc. Helminthol. Soc. Wash.* **54**(1): 28–39.
- . **M.M. Kliks, M.E. Rosenfeld, R.A. Rychlinski & R.S. Desowitz.** 1982. Larval ascaridoid nematodes from fishes near the Hawaiian Islands, with comments on pathogenicity experiments. *Pac. Sci.* **36**(2): 187–201.
- Devaney, D.M.** 1977a. Ctenophora, p. 262–69. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- . 1977b. Subclass Octocorallia, p. 119–29. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- Devaney, D.M. & L.G. Eldredge.** 1977. Class Scyphozoa, p. 108–18. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- . 1987. Phylum Nemertea (Rhynchocoela), p. 59–69. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- Devick, W.S., J.M. Fitzsimons & R.T. Nishimoto.** 1992. *Conservation of Hawaiian freshwater fishes.* State of Hawaii Division of Aquatic Resources. 26 p.
- Edmonds, S.J.** 1987. Phyla Sipuncula and Echiura, p. 185–212. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- Eliasson, U.H.** 1991. The myxomycete biota of the Hawaiian Islands. *Mycol. Res.* **95**: 257–67.
- Eldredge, L.G.** 1967. Record of a lancelet from Hawaii. *Pac. Sci.* **21**(4): 564.
- . 1994. *Introductions of commercially significant aquatic organisms to the Pacific islands.* South Pacific Commission, Noumea. 127 p.
- Eldredge, L.G. & D.M. Devaney.** 1977. Other hydrozoans, p. 105–07. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- Emig, C.C.** 1987. Phylum Brachiopoda, p. 167–70. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- . **& J.H. Bailey-Brock.** 1987. Phylum Phoronida, p. 171–81. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- Erséus, C.** 1990. Marine Oligochaeta of Hong Kong, p. 259–335. *In: Morton, B., ed., The*

## 14 BISHOP MUSEUM OCCASIONAL PAPERS: NO. 41, 1995

- marine flora and fauna of Hong Kong and southern China* 2. Vol. 1. Introduction and taxonomy. Hong Kong University Press, Hong Kong.
- . & **D. Davis**. 1989. The marine Tubificidae (Oligochaeta) of Hawaii. *Asian Mar. Biol.* **6**: 73–100.
- Fauchald, K.** 1992. A review of the genus *Eunice* (Polychaeta: Eunicidae) based upon type material. *Smithson. Contrib. Zool.* **523**, 422 p.
- Fisher, W.K.** 1906. The starfishes of the Hawaiian Islands. *Bull. U.S. Fish Commn.* **23**(3): 987–1130.
- . 1907. The holothurians of the Hawaiian Islands. *Proc. U.S. Natl. Mus.* **32**(1555): 637–744.
- Frank, J.H. & E.D. McCoy.** 1990. Endemics and epidemics of shibboleths and other things causing chaos. *Fla. Entomol.* **73**(1): 1–9.
- Goos, R.D.** 1977. Comparative observations of the Hyphomycetes of Hawaii and Central America. *Kavaka* **6**: 31–35.
- . & **D.P. Gowing.** 1992. Type specimens of fungi maintained at Herbarium Pacificum, Bernice P. Bishop Museum, Honolulu. *Mycotaxon* **43**: 177–98.
- Grigg, R.W. & F.M. Bayer.** 1976. Present knowledge of the systematic and zoogeography of the Order Gorgonacea in Hawaii. *Pac. Sci.* **30**(2): 167–75.
- . & **D. Opresko.** 1977. Order Antipatharia, p. 242–61. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- . **J.W. Wells & C. Wallace.** 1981. *Acropora* in Hawaii. Part 1. History of the scientific record, systematics, and ecology. *Pac. Sci.* **35**(1): 1–13.
- Hadfield, M.G. & R.E. Young.** 1983. *Planctosphaera* (Hemichordata: Enteropneusta) in the Pacific Ocean. *Mar. Biol.* **73**(2): 151–53.
- Hartmann, G.** 1991. Ostracoden von Hawaii, insbesondere aus dem marinen Interstitial. *Helgoländ. Meeresuntersuch.* **45**: 165–98.
- Hawksworth, D.L.** 1994. The tropical fungal biota: census, pertinence, prophylaxis, and prognosis, p. 265–93. In: Issac, S., J.C. Frankland, A.J. Whalley & R. Watling, eds., *Tropical mycology*. Cambridge University Press, Cambridge.
- Hering, J.L.** 1961. The genus *Halobates* (Hemiptera: Gerridae). *Pac. Insects* **3**(2–3): 223–305.
- . 1965. *Hermatobates*, a new generic record for the Atlantic Ocean, with descriptions of new species (Hemiptera: Gerridae). *Proc. U.S. Natl. Mus.* **117**(3510): 123–30.
- Hoe, W.J.** 1974. Annotated checklist of Hawaiian mosses. *Lyonia* **1**: 1–45.
- . 1979. The phytogeographical relationships of Hawaiian mosses. Unpubl. Ph.D. Dissertation, University of Hawaii, Honolulu. 357 p.
- Hope, W.D.** 1987. The Aschelminth complex, p. 71–81. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- Howarth, F.G.** 1990. Hawaiian terrestrial arthropods: an overview. *Bishop Mus. Occas. Pap.* **30**: 4–26.
- . & **J. Moore.** 1983. The land nemertine *Argonemertes dendyi* (Dakin) in Hawaii (Nemertinea: Hoplonemertinea: Prosorhochmidae). *Pac. Sci.* **37**(2): 141–44.

## RECORDS OF THE HAWAII BIOLOGICAL SURVEY FOR 1994 15

- Humes, A.G.** 1942. The morphology, taxonomy, and bionomics of the nemertean genus *Carcinonemertes*. *Ill. Biol. Monogr.* **18**(4): 3–105.
- . 1976. Distribution and hosts of *Stellicola* (Copepoda, Cyclopoida) associated with *Linckia* (Asteroidea) in the Indo-west Pacific. *Beaufortia* **25**(321): 49–61.
- Hyman, L.H.** 1939. Land planarians from the Hawaiian Islands. *Arch. Zool. Exper. Gen.* **80**(3): 116–24.
- James, H.F. & S.L. Olson.** 1991. Descriptions of thirty-two species of birds from the Hawaiian Islands: Part II. Passeriformes. *Ornithol. Monogr.* **46**: 1–88.
- Kaneko, J.J., II, R. Yamada, J.A. Block & R. Nakamura.** 1988. Infection of tilapia, *Oreochromis mossambicus* (Trewavas), by a marine monogenean, *Neobenedenia melleni* (MacCallum, 1927) Yamaguti, 1963 in Kaneohe Bay, Hawaii, USA, and its treatment. *J. Fish Diseases* **11**: 295–300.
- Kawakatsu, M. & R.W. Mitchell.** 1984. *Oahuhawaiiiana kazukolinda* gen. et sp. nov. (Turbellaria, Tricladida, Maricola), a new freshwater planarian from Honolulu, Oahu Island, Hawaii, U.S.A. *Zool. Sci.* **1**(3): 487–500.
- . **R.W. Mitchell, Y. Hirao & I. Tanaka.** 1984. Occurrence of *Dugesia dorotocephala* (Woodworth, 1897) (Turbellaria, Tricladida, Paludicola) in Honolulu, Hawaii. *Biol. Mag. (Okinawa)* **22**: 1–9.
- Kay, E.A.** 1979. *Hawaiian marine shells. Reef and shore fauna of Hawaii.* Section 4: Mollusca. Bishop Museum Press, Honolulu. 652 p.
- . 1994. *A natural history of the Hawaiian islands. Selected readings II.* University of Hawaii Press, Honolulu. xi + 520 p.
- . **& S.R. Palumbi.** 1987. Endemism and evolution in Hawaiian marine invertebrates. *Trends Ecol. Evol.* **2**(7): 183–86.
- Kitalong, A.E.** 1986. A preliminary study on the emergence patterns of microfauna in Kaneohe Bay, Hawaii, p. 414–23. *In: Jokiel, P.L., R.H. Richmond & R.A. Rogers, eds., Coral reef population biology.* Hawaii Institute of Marine Biology Technical Report 37.
- Kohlmeyer, J.** 1969. Marine fungi of Hawaii including the new genus *Helicascus*. *Canad. J. Bot.* **47**: 1469–1487.
- Kornicker, L.S.** 1976. Benthic marine Cypridinacea from Hawaii (Ostracoda). *Smithson. Contrib. Zool.* **231**, 24 p.
- Kunz, H.** 1993. Beitrag zur Kenntnis von Zwei *Psammopsyllus*-Arten (Copepoda, Harpacticoida, Cylindropsyllidae). *Crustaceana* **64**(2): 143–54.
- Larsen, J. & D.J. Patterson.** 1990. Some flagellates (Protista) from tropical marine sediments. *J. Nat. Hist.* **24**: 801–937.
- Lewis, A.G.** 1966. Copepod crustaceans parasitic on elasmobranch fishes of the Hawaiian Islands. *Proc. U.S. Natl. Mus.* **118**(3524): 57–154.
- . 1967. Copepod crustaceans parasitic on teleost fishes of the Hawaiian Islands. *Proc. U.S. Natl. Mus.* **121**(3574): 1–204.
- MacCaughy, V.** 1918. Algae of the Hawaiian Archipelago. *Bot. Gaz.* **65**: 42–57, 121–49.
- Maciolek, J.A.** 1984. Exotic fishes in Hawaii and other islands of Oceania, p. 131–61. *In: Courtenay, W.R., Jr. & J.R. Stauffer, eds., Distribution, biology, and management of exotic fishes.* Johns Hopkins University Press, Baltimore.
- Maragos, J.E.** 1977. Order Scleractinia, p. 158–241. *In Devaney, D.M. & L.G. Eldredge, eds., Reef and shore fauna of Hawaii.* Section 1: Protozoa through Ctenophora.

- Bishop Museum Press, Honolulu.
- . 1995. Revised checklist of extant shallow water stony coral species from Hawaii (Cnidaria: Anthozoa: Scleractinia). *Bishop Mus. Occas. Pap.* **42**: 53–54.
- Margulis, L. & K.V. Schwartz.** 1988. *Five kingdoms: an illustrated guide to the phyla of life on earth*. Second edition. W.H. Freeman & Co., New York. 376 p.
- Matthews, D.C.** 1966. A comparative study of *Craspedacusta sowerbyi* and *Calpasoma dactyloptera* life cycles. *Pac. Sci.* **20**(2): 246–59.
- McInnes, S.J.** 1994. Zoogeographic distribution of terrestrial/freshwater tardigrades from current literature. *J. Nat. Hist.* **28**(2): 257–352.
- McKenzie, M.E. & W.R. Davidson.** 1989. Helminth parasites of intermingling axis deer, wild swine and domestic cattle from the island of Molokai, Hawaii. *J. Wildl. Diseases* **25**(2): 252–57.
- Metcalf, M.M. & H. S. Hopkins.** 1919. *Pyrosoma*—taxonomic study based upon the collections of the United States Bureau of Fisheries and the United States National Museum. *Bull. U.S. Natl. Mus.* **100** 2(3): 195–272.
- Miller, H.A. & H. O. Whittier.** 1990. Bryophyte floras of tropical Pacific islands. *Trop. Bryol.* **2**: 167–75.
- Moore, J.P.** 1946. Leeches (Hirudinea) from the Hawaiian Islands, and two new species from the Pacific region in the Bishop Museum collection. *Occas. Pap. B. P. Bishop Mus.* **18**(11): 171–91.
- Motoda, S.** 1963. *Coryceus* and *Farranula* copepods (Cyclopoida) in Hawaiian waters. *Publ. Seto Mar. Biol. Lab.* **11**(2): 209–62.
- Muzik, K.** 1978. A bioluminescent gorgonian, *Lepidisis olapa*, new species (Coelenterata: Octocorallia), from Hawaii. *Bull. Mar. Sci.* **28**(4): 735–41.
- Nakamura, M.** 1992. An ecological study of earthworms in Papua New Guinea. *J. Chuo Univ.* **13**: 19–33.
- Nishida, G.M., ed.** 1994. Hawaiian terrestrial arthropod checklist. Second edition. *Bishop Mus. Tech. Rep.* **4**, 287 p.
- Nutting, C.C.** 1905. Hydroids of the Hawaiian Islands collected by the steamer *Albatross* in 1902. *Bull. U.S. Fish Commn.* **23**(3): 931–59.
- Oliveira, J.M.** 1940. Plant-parasitic and free-living nematodes in Hawaii. *B. P. Bishop Mus. Occas. Pap.* **15**(29): 361–73.
- Olson, S.L. & H.F. James.** 1991. Descriptions of thirty-two new species of birds from the Hawaiian Islands: Part I. Non-Passeriformes. *Ornithol. Monogr.* **45**: 1–88.
- Ortmann, A.E.** 1906. Schizopods of the Hawaiian Islands collected by the steamer *Albatross* in 1902. *Bull. U.S. Fish Commn.* **23**(3): 961–73.
- Otte, D.** 1994. *The crickets of Hawaii. Origin, systematics and evolution*. Orthopterist's Society, Philadelphia. vi + 396 p.
- Parker, S.P., ed.** 1982. *Synopsis and classification of living organisms*. 2 vols. McGraw-Hill Book Co., New York.
- Parris, G.K.** 1940. A check list of fungi, bacteria, nematodes, and viruses occurring in Hawaii, and their hosts. *Plant Disease Reporter; Supplement* **121**, 91 p.
- Paxton, H. & J.H. Bailey-Brock.** 1986. *Diopatra dexiognatha*, a new species of Onuphidae (Polychaeta) from Oahu, Hawaiian Islands. *Pac. Sci.* **40**(1-4): 16.
- Pearse, V.B.** 1989. Growth and behavior of *Trichoplax adhaerens*: first record of the phylum Placozoa in Hawaii. *Pac. Sci.* **43**(2): 117–21.

## RECORDS OF THE HAWAII BIOLOGICAL SURVEY FOR 1994 17

- Pettibone, M.H.** 1986. Review of the Iphioninae (Polychaeta: Polynoidae) and revision of *Iphione cimex* Quatrefages, *Gattyana deludens* Fauvel, and *Harmothoe iphionelloides* Johnson (Harmothoinae). *Smithson. Contrib. Zool.* **428**, 43 p.
- Phillips, F.J.** 1977. Protozoa, p. 12–52. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- Pierrot-Bults, A. C. & V. R. Nair.** 1991. Distribution patterns in Chaetognatha, p. 86–116. In: Bone, Q., H. Kapp & A.C. Pierrot-Bults, eds., *The biology of chaetognaths*. Oxford University Press, New York.
- Pilsbry, H. A.** 1907. Hawaiian Cirripedia. *Bull. Bur. Fish.* **26**: 179–190.0
- Poulter, J.** 1987. Phylum Platyhelminthes, p. 13–58. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- Pyle, R.L.** 1992. Checklist of the birds of Hawaii—1992. *'Elepaio* **42**: 43–62.
- Raabe, R.D.** 1981. *Checklist of plant diseases in Hawaii: including records of microorganisms, principally fungi, found in the state*. Enlarged and revised by Ibra L. Conners and Albert R. Martinez. Hawaii Institute of Tropical Agriculture and Human Resources, College of Tropical Agriculture and Human Resources, University of Hawaii at Manoa. 313 p.
- Randall, J.E.** 1992. Endemism of fishes in Oceania. *UNEP Regional Seas Rep. Stud.* **147**: 55–67.
- Richardson, H.** 1906. Isopods collected at the Hawaiian Islands by the U.S. Fish Commission steamer *Albatross*. *Bull. U.S. Fish Commn.* **23**(3): 819–26.
- Roper, C.F.E. & R.E. Young.** 1975. Vertical distribution of pelagic cephalopods. *Smithson. Contrib. Zool.* **209**, 51 p.
- . **M.J. Sweeney & C.E. Nauen.** 1984. FAO species catalogue. Vol. 3. Cephalopods of the world. An annotated and illustrated catalogue of species of interest to fisheries. *FAO Fish. Synop.* **125**, **3**: 1–277.
- Smith, C.W.** 1991. Lichen conservation in Hawaii, p. 35–45. In D.J. Galloway, ed., *Tropical lichens: their systematics, conservation, and ecology*. Systematics Association Special Volume 43. Clarendon Press, Oxford.
- . 1993. Notes on Hawaiian parmelioid lichens. *Bryologist* **96**: 326–32.
- Soest, R.W.M. van.** 1974a. Taxonomy of the subfamily Cyclosalpidae Yount, 1954 (Tunicata, Thaliacea), with descriptions of two new species. *Beaufortia* **22**(288): 17–55.
- . 1974b. A revision of the genera *Salpa* Forskål, 1775, *Pegea* Savigny, 1816, and *Ritteriella* Metcalf, 1919 (Tunicata, Thaliacea). *Beaufortia* **22**(293): 153–91.
- Soule, J.D., D.F. Soule & H.W. Chaney.** 1987. Phyla Entoprocta and Bryozoa (Ectoprocta), p. 83–166. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 2: Platyhelminthes through Phoronida and Section 3: Sipuncula through Annelida. Bishop Museum Press, Honolulu.
- Southward, E.C.** 1980. Two new species of Pogonophora from Hawaii. *Pac. Sci.* **34**(4): 371–78.
- Stenroos, S. & C.W. Smith.** 1993. Notes on the amphipacific relations of Hawaiian Cladoniaceae. *Trop. Biol.* **8**: 275–80.
- Sterrer, W.** 1991. Gnathostomulida from Hawaii. *Zool. Scripta* **20**(2): 129–36.
- Svihla, A.** 1941. The occurrence of freshwater sponges in the Hawaiian Islands. *Science* **94**(2438): 278–79.

## 18 BISHOP MUSEUM OCCASIONAL PAPERS: NO. 41, 1995

- Taguchi, S.** 1982. Seasonal study of fecal pellets and discarded houses of Appendicularia in a subtropical inlet, Kaneohe Bay, Hawaii. *Estuar. Coast. Shelf Sci.* **14**: 545–55.
- Tokioka, R.** 1967. Pacific Tunicata of the United States National Museum. *U.S. Natl. Mus. Bull.* **251**, 247 p.
- Tomich, P.Q.** 1986. *Mammals in Hawaii. A synopsis and notational bibliography*. Second edition. Bishop Museum Press, Honolulu. 375 p.
- Ueno, M.** 1936. Cladocera of Mauna Kea, Hawaii. *B. P. Bishop Mus. Occas. Pap.* **12**(11): 1–9.
- Van Zwaluwenburg, R.H.** 1948. *Lacustris hawaiiensis* Johnson. *Proc. Hawaii. Entomol. Soc.* **13**(2): 210.
- Vávra, W.** 1906. Ostracoden von Sumatra, Java, Siam, den Sandwich-Inseln und Japan. *Zool. Jahrb. (Syst.)* **23**: 413–38.
- Versevelt, J. & F.M. Bayer.** 1988. Revision of the genera *Bellonella*, *Eleutherobia*, *Nidallia* and *Nidaliopsis* (Octocorallia: Alcyoniidae and Nidaliidae), with descriptions of two new genera. *Zool. Verh.* **245**: 1–131.
- Vis, M.L., R.G. Sheath, J.A. Hambrook & K.M. Cole.** 1994. Stream macroalgae of the Hawaiian Islands: a preliminary study. *Pac. Sci.* **48**: 175–87.
- Vitousek, P.M., L.L. Loope & C.P. Stone.** 1987. Introduced species in Hawaii: biological effects and opportunities for ecological research. *Trends Ecol. Evol.* **2**(7): 224–27.
- Wagner, W.L., D.R. Herbst & S.H. Sohmer.** 1990. *Manual of the flowering plants of Hawai'i*. University of Hawaii Press & Bishop Museum Press, Honolulu. 1853 p.
- Walsh, G.E. & R.L. Bowers.** 1977. Order Zoanthiaria, p. 148–57. In Devaney, D.M. & L.G. Eldredge, eds., *Reef and shore fauna of Hawaii*. Section 1: Protozoa through Ctenophora. Bishop Museum Press, Honolulu.
- Wells, J.B.J.** 1986. Copepoda: marine-interstitial Harpacticoida, p. 356–81. In: Botosaneanu, L., ed., *Stygofauna mundi*. E.J. Brill, Leiden.
- Yamaguti, S.** 1965. New digenetic trematodes from Hawaiian fishes, I. *Pac. Sci.* **19**(4): 458–81.
- . 1968a. Cestode parasites of Hawaiian fishes. *Pac. Sci.* **22**(1): 21–36.
- . 1968b. *Monogenetic trematodes of Hawaiian fishes*. University of Hawaii Press, Honolulu. 287 p.
- . 1970. *Digenetic trematodes of Hawaiian fishes*. Keigaku Publishing Co., Tokyo. 436 p.
- Young, R.E.** 1991. Chiroteuthid and related paralarvae from Hawaiian waters. *Bull. Mar. Sci.* **49**(1–2): 162–85.
- Yount, J.L.** 1954. The taxonomy of the Salpidae (Tunicata) of the central Pacific Ocean. *Pac. Sci.* **8**(3): 276–330.