

Beyond Reserves: A Research Agenda for Conserving Biodiversity in Human-modified Tropical Landscapes

Robin L. Chazdon^{1,10}, Celia A. Harvey², Oliver Komar³, Daniel M. Griffith⁴, Bruce G. Ferguson⁵, Miguel Martínez-Ramos⁶, Helda Morales⁵, Ronald Nigh⁷, Lorena Soto-Pinto⁵, Michiel van Breugel⁸, and Stacy M. Philpott⁹

¹Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, Connecticut, U.S.A.

²Conservation International, 2011 Crystal Drive Suite 500, Arlington, Virginia 22202, U.S.A.

³Programa de Ciencias para la Conservación, SalvaNATURA, Colonia Flor Blanca, 33 Avenida Sur #640, San Salvador, El Salvador

⁴Biodiversity of BOSAWAS Biosphere Reserve, Saint Louis Zoo, Managua, Nicaragua

⁵Departamento de Agroecología, El Colegio de la Frontera Sur, Carretera Panamericana y Periférico Sur s-n, San Cristóbal de Las Casas, Chiapas, México

⁶Centro de Investigaciones en Ecosistemas, UNAM, AP 27-3 Santa María de Guido, CP 58089, Morelia, Michoacán, México

⁷Centro de Investigaciones y Estudios Superiores en Antropología Social, San Cristóbal de Las Casas, Chiapas, México

⁸Center for Tropical Forest Science, Smithsonian Tropical Forest Research Institute, Unit 0948, APO AA 34002, U.S.A.

⁹Department of Environmental Sciences, University of Toledo, 2801 W. Bancroft St., Toledo, Ohio 43606, U.S.A.

ABSTRACT

To truly understand the current status of tropical diversity and to forecast future trends, we need to increase emphasis on the study of biodiversity in rural landscapes that are actively managed or modified by people. We present an integrated landscape approach to promote research in human-modified landscapes that includes the effects of landscape structure and dynamics on conservation of biodiversity, provision of ecosystem services, and sustainability of rural livelihoods. We propose research priorities encompassing three major areas: biodiversity, human–environment interactions, and restoration ecology. We highlight key areas where we lack knowledge and where additional understanding is most urgent for promoting conservation and sustaining rural livelihoods. Finally, we recommend participatory and multidisciplinary approaches in research and management. Lasting conservation efforts demand new alliances among conservation biologists, agroecologists, agronomists, farmers, indigenous peoples, rural social movements, foresters, social scientists, and land managers to collaborate in research, co-design conservation programs and policies, and manage human-modified landscapes in ways that enhance biodiversity conservation and promote sustainable livelihoods.

Key words: agricultural matrix; agroecology; conservation value; ecosystem services; remnant vegetation; restoration; traditional knowledge.

THE ACTIVE DEBATE ON THE FUTURE OF TROPICAL BIODIVERSITY is largely driven by a deficit of information regarding the status of biodiversity in human-modified rural landscapes (Wright & Muller-Landau 2006, Gardner *et al.* 2007a). Biodiversity surveys and ecological studies have understandably focused on areas with a high concentration of plant and animal diversity—intact biological reserves and protected areas with low current levels of human intervention (Fazey *et al.* 2005). But these areas are not typical of most of the world's tropics, where more than 90 percent of tropical forest area lies beyond the borders of reserves and parks (WWF 2002). To truly understand the current status and forecast the future state of tropical diversity, we must also understand levels and patterns of biodiversity in landscapes actively managed and modified by humans for a wide variety of traditional and commercial purposes, including hunting and gathering, agriculture, extractive forestry, and plantations of native or exotic species. Further, we must investigate how these patterns are affected by different human practices,

land-use dynamics, spatial contexts, and socioeconomic contexts along a gradient of landscape modification, from smallholder agriculture to large-scale forestry and industrial commodity production (Bawa *et al.* 2004). The information obtained from such investigations is essential to identify and promote appropriate management strategies for conserving biodiversity in tropical regions (Zuidema & Sayer 2003, Lindenmayer *et al.* 2008).

The fates of biodiversity in protected areas and surrounding landscapes are inextricably linked (Schelhas & Greenberg 1996, McNeely & Scherr 2003, Zuidema & Sayer 2003, Vandermeer & Perfecto 2007, Harvey *et al.* 2008). Most protected areas in tropical regions are embedded within a matrix of heterogeneous land uses and are often directly or indirectly affected by forest fragmentation, road construction, agrochemicals, hunting, cattle grazing, agricultural incursions, fire, invasive species, over-harvest of non-timber forest products, logging, and mining (Janzen 1983, Schelhas & Greenberg 1996, DeFries *et al.* 2005, Hansen & DeFries 2007). These human activities often threaten species in protected areas (Laurance *et al.* 2006, Giraõ *et al.* 2007, Michalski *et al.* 2007). On the other hand, certain types of agriculture, agroforestry,

Received 16 January 2008; revision accepted 28 July 2008.

¹⁰Corresponding author; e-mail: Chazdon@uconn.edu

fallow vegetation, and forest patches surrounding protected areas can support significant levels of biodiversity (Daily *et al.* 2001, 2003; Mayfield *et al.* 2005; Peh *et al.* 2006; Barlow *et al.* 2007a), while also providing valuable ecosystem services, such as carbon sequestration and hydrological protection (Montagnini & Nair 2004, Potvin *et al.* 2007, Tschakert *et al.* 2007). Incorporation of 'biodiversity friendly' land uses into actively managed buffer zones or biological corridors can contribute to the long-term conservation value of protected areas (DeFries *et al.* 2007, Harvey *et al.* 2008). In landscapes lacking protected areas or intact forests, agriculture, agroforestry, remnant vegetation, plantations, and managed forest patches provide critical habitats and refugia for biodiversity (Harvey *et al.* 2006, Harvey & González 2007, Bhagwat *et al.* 2008).

Despite a growing recognition of the importance of assessing and conserving biodiversity in human-modified landscapes in the tropics (Schroth *et al.* 2004, Harvey & Sáenz 2007, Bhagwat *et al.* 2008), many key questions remain to be answered in order to provide clear guidelines for long-lasting conservation efforts (Fischer *et al.* 2006, Norris 2008). An integrated landscape approach is needed to understand the effects of landscape structure and dynamics on conservation of biodiversity, provision of ecosystem services, and sustainability of rural livelihoods (Tschardt *et al.* 2005). This integrated landscape approach was the basis for a companion paper by Harvey *et al.* (2008) that focused on policy recommendations within Mesoamerican countries. Here, we propose 12 priorities for investigation within human-modified landscapes in rural areas of the tropics, encompassing three major areas: biodiversity, human-environment interactions, and restoration ecology (Table 1). Our message is directed toward researchers and organizations that support research in tropical biology and conservation, rather than policy makers. We highlight key areas where we lack knowledge and where additional understanding is most urgent for promoting conservation and rural livelihoods (Table 1). Finally, we recommend that re-

TABLE 1. *A research agenda for conserving biodiversity in tropical human-modified landscapes. Each of the 12 areas of research focus are described in more detail in the text.*

Major area	Research focus
Bio-diversity status and landscape ecology	1. Population biology and long-term monitoring
	2. Animal dispersal and habitat use
	3. Effectiveness of buffer zones and corridors
	4. Effects of specific land-use practices
	5. Modeling impacts of climate change
Interactions between people and their environment	6. Ecosystem services and land use
	7. Relationships between biodiversity and ecosystem functions
	8. Social and economic impact of conservation activities
	9. Relationships between human communities, local resources, and sustainable management
Restoration ecology	10. Landscape-level restoration
	11. Costs and benefits of restoration objectives
	12. Effects of livestock on restoration

search and management be participatory and multidisciplinary, and should feed back into planning and managing landscapes within an adaptive framework. We advocate a broadening of focus beyond conservation *biology* toward a broader discipline of conservation *science*. Lasting conservation efforts demand new alliances among conservation biologists, agroecologists, agronomists, farmers, indigenous peoples, rural social movements, foresters, social scientists, land managers, and government agencies to collaborate in research, create conservation programs and policies, and to manage human-modified landscapes in ways that enhance biodiversity conservation (Pretty 1995, Cullen *et al.* 2005, Kaimowitz & Sheil 2007, Harvey *et al.* 2008). The authors' major areas of expertise are Mesoamerican agroecosystems and forests, and our research priorities emerge from intimate associations with Latin American landscapes and cultures. Nevertheless, our intention is to provide a broad framework that can be applied to other tropical regions and that will stimulate the development of research programs best suited to the unique human and biological history and landscape context of particular geographic regions.

AN INTEGRATED APPROACH TO RESEARCH IN HUMAN-MODIFIED LANDSCAPES

The burgeoning number of ecological studies in human-modified landscapes reflects an urgent need to examine biotic interactions between matrix habitats and embedded forest patches (Gascon *et al.* 1999, Jules & Shahani 2003, Klein *et al.* 2008, Lindenmayer *et al.* 2008). Ecological studies generally view the agricultural matrix as homogeneous and as a source of contamination of embedded forest patches (Janzen 1983, Nascimento *et al.* 2006), rather than viewing forest remnants as heterogeneous biodiversity sources and sinks within the broader landscape. By zooming out on the landscape matrix itself, we can investigate population dynamics and species interactions among component habitat types (agriculture, secondary vegetation, forest fragments) in a metapopulation or metacommunity context (Daily *et al.* 2001, Vandermeer & Carvajal 2001, Perfecto & Vandermeer 2002, Bennett *et al.* 2006, Kupfer *et al.* 2006, Pulido *et al.* 2007, Greenberg *et al.* 2008). Forest fragments and isolated remnant trees provide sources of propagules for re-populating surrounding areas and serve as resources, stepping stones, and refugia for wildlife that use multiple habitats (Bengtsson *et al.* 2003, Guevara *et al.* 2005). Like forest fragments, agroforestry systems can also function as biological corridors and stepping stones for many animal species (Estrada *et al.* 1997, Laurance 2004, Schroth *et al.* 2004). In some tropical regions, coffee, cacao, rubber, or other agroforestry systems are the dominant form of tree cover and therefore play a key role in biodiversity conservation at the landscape level (Alcorn 1990, Young 1994, Perfecto *et al.* 1996, Moguel & Toledo 1999, Peters 2000, Abarca 2006, Monro *et al.* 2006, Bhagwat *et al.* 2008).

Agricultural production systems vary widely in their impact on biodiversity, ecosystem services, land-use dynamics, and potential for regeneration when abandoned (Donald 2004, Chazdon *et al.*

2008, Philpott *et al.* 2008a). The negative impacts of large-scale industrial agriculture (cotton, soybeans, sugarcane, bananas, rubber, African oil palm) on biodiversity are widely acknowledged to be significantly greater than those of traditional, small-scale agroforestry systems (McNeely & Scherr 2003, Donald 2004, Schroth *et al.* 2004, Scherr & McNeely 2007). In heterogeneous human-modified landscapes, forest fragments provide ecosystem services that benefit crop production (Swift *et al.* 2004, Maass *et al.* 2005). Biodiversity in forest fragments and landscape heterogeneity can enhance pollinator activity for crops (Kremen *et al.* 2002, Ricketts *et al.* 2004, Balvanera *et al.* 2005, Klein *et al.* 2008), promote pest control (Pickett & Bugg 1998, Klein *et al.* 2006, Romero *et al.* 2006), and reduce fungal infection and weed growth (Soto-Pinto *et al.* 2002). Forest fragments also provide products for local use, protect watersheds, store carbon, and meet other economic and cultural needs (Khumbongmayum *et al.* 2005, Bongers *et al.* 2006). Empirical and theoretical studies show that conservation of crop diversity can enhance agricultural productivity and ecosystem services (Tscharrntke *et al.* 2005, Perrings *et al.* 2006, Jackson *et al.* 2007, Omer *et al.* 2007).

The objectives of an integrated approach to conservation within human-modified landscapes are not only to maximize protection of a wide range of taxa and ecosystem services, but also to improve agricultural productivity, food security, collective resource rights, and human welfare. These objectives are consistent with the findings of the International Assessment of Agricultural Knowledge, Science and Technology for Development (IAASTD 2008), which advocate a multifunctional perspective on agriculture, incorporating the need to protect biodiversity and ecosystem services. Trade-offs are unavoidable, but can be reduced in many cases through implementation of participatory, adaptive management (also referred to as adaptive co-management or community-based resource management). Adaptive management represents an important strategy for ongoing, systematic learning and adjustment to changing circumstances (Salafsky *et al.* 2008). This approach emphasizes *how* people learn and with *whom* rather than *what* they learn, and the ultimate goal is mobilization of local stakeholders and institutions leading to sustained action (Pretty 1995). Participation in research and decision making by multiple stakeholders is more likely to generate information and actions that meet both social and ecological needs than research by one or a few 'experts'. Although it is not a panacea, adaptive management can serve as an important tool to integrate conservation with sustainability of rural livelihoods. Further research is needed to improve learning methods and outcomes (Armitage *et al.* 2008), to promote equity of participation among stakeholders (Sanginga *et al.* 2006), and to determine the optimal level of experimentation to maximize ecological and social returns over different timescales (Hauser & Possingham 2008).

RESEARCH AGENDA

The research priorities discussed below address the urgent need for basic information required for adaptive management of human-modified ecosystems and landscapes, as well as for baseline data

for long-term studies. We highlight 12 research priorities within three major areas for investigation in human-modified landscapes: (1) biodiversity status and landscape ecology; (2) interactions between people and their environment; and (3) restoration ecology (Table 1).

BIODIVERSITY STATUS AND LANDSCAPE ECOLOGY.—In one of the first books to focus on the mosaic of landscape fragments in the tropics, Schelhas and Greenberg (1996) stated, "We must learn more about which species can survive and thrive in different types and arrangements of forest patches" (p. xvi; Introduction). We expand this question to include the full range of habitat types within existing human-modified landscapes, including agricultural systems, remnant trees in pastures or in nearby areas, riparian strips, living fences, shade trees in cropping systems, home gardens, secondary vegetation, and remnant forest patches. Further, we advocate approaches that look beyond individual habitat types to the linkages and dynamics across habitat types and landscapes. Few studies have quantified patterns of biodiversity within one or more taxa across entire gradients of landscape modification (Nichols *et al.* 2007, Philpott *et al.* 2008a). We highlight five research priorities addressing this broad realm.

1. Population biology and long-term monitoring within human modified landscapes

Monitoring studies provide baseline information to assess which component habitats in the landscape matrix can support particular taxa, and to evaluate short- and long-term effects of land-use change, landscape structure and heterogeneity, and successional dynamics on biodiversity and life-support systems. Which taxa can persist in human-modified landscapes and which cannot? These data will also provide a comparative context for monitoring studies conducted within intact forest areas within the same geographic region. Inventory data for particular taxa suggest that a subset of forest species can be found within human-modified landscapes (Dunn 2004, García-Estrada *et al.* 2006, Harvey *et al.* 2006, Sekercioglu *et al.* 2007, Barlow *et al.* 2007a, Gardner *et al.* 2007b, Kabir & Webb 2008), but we lack monitoring data over time to evaluate persistence of species in a range of habitats within and across landscapes (*e.g.*, Harvey *et al.* 2008, Suazo *et al.* 2008). Moreover, few studies have examined the effects of short-term or seasonal population fluctuations on observed landscape patterns of biodiversity (Barlow *et al.* 2007c).

We know virtually nothing about the long-term dynamics of populations and their genetic structure and variation within human-modified landscapes (but see Boshier *et al.* 2004), as most studies of plant and animal diversity within agricultural landscapes have focused solely on describing static patterns of abundance and species richness. How are population processes affected by different land-use practices, landscape configurations, and levels of landscape modification and degradation (Cascante *et al.* 2002, Boshier *et al.* 2004, Komar 2006, Pulido *et al.* 2007). The minimum amount of habitat needed to sustain species' population

dynamics in a predictable time frame is defined as landscape threshold (Reunanen *et al.* 2004). Landscape thresholds can be identified for certain taxa and study sites (Lindenmayer & Luck 2005, Radford *et al.* 2005), but for conservation planning we need much more information to identify these thresholds for keystone species, migratory species, commercial species, or species of conservation concern in different types of agricultural landscapes (Harvey 2007). These thresholds may also change over time, especially with the added stress of climate change.

Second-growth forests provide the only forested habitats within some human-modified landscapes. Long-term studies of natural regeneration are urgently needed to understand rates of vegetation change (Chazdon *et al.* 2007), suitability for management/harvesting, and conservation status of flora and fauna (Barlow *et al.* 2007b, Gardner *et al.* 2007b). We are far from understanding the role of plant–plant, microbial–plant, and animal–plant interactions on biodiversity change in successional environments. For example, we have scant information about effects of changes in seed dispersal, pollination, seed predation, and herbivory on vegetation composition in secondary forests in human-modified landscapes (Cramer *et al.* 2007).

2. Animal dispersal and habitat use within human-modified landscapes

Habitat utilization studies are needed to identify key habitats, foraging and nesting sites, and dispersal routes of animals in planning conservation units, buffer zones, and biological corridors and agricultural habitat components. How do habitat patches influence breeding behavior, dispersal, and species interactions within the agricultural mosaic? To what extent do remnant habitats serve as refugia for agricultural pests and their natural enemies (Schmidt *et al.* 2004)? Some taxa move readily across a complex landscape matrix, despite a high degree of habitat modification and fragmentation (*e.g.*, bats in pastoral and fragmented landscapes; Estrada *et al.* 2004, Bianconi *et al.* 2006, Medina *et al.* 2007). Other species are highly sensitive to even small changes in fragmentation and forest loss (*e.g.*, dung beetles; Klein 1989), insectivorous forest understory birds (Antongiovanni & Metzger 2005), small amphibians, and some turtle species (Suazo *et al.* 2008) or to structural changes in agricultural habitats (Cruz-Angón *et al.* 2008). Understanding animal movement is critical to reducing the negative effects of climate change on wildlife populations (Donald & Evans 2006) and will enable effective planning of biological corridors and stepping stones in human-modified landscapes.

3. Effectiveness of buffer zones and corridors for the conservation of target species, sources of forest regeneration, and production of ecosystem services

Despite some research in temperate regions (Tewksbury *et al.* 2002), we lack experimental studies to determine the most effective design and management of buffer zones and biological corridors (Fischer *et al.* 2006). What habitat types

and spatial arrangements provide effective buffer zones and corridors for different taxa? How can we incorporate existing agricultural lands, riparian strips, and remaining forest patches into buffer zones and biological corridors to increase the capacity of landscapes to protect biodiversity? How can we increase the value of buffer zones and biological corridors to local people through provision of ecosystem services that enhance agricultural productivity and protect against catastrophic landslides and floods (Schelhas & Greenberg 1996)? Presently, few landscape-level investigations address these urgent questions (Williams-Guillén *et al.* 2006, Anzures-Dadda & Manson 2007).

4. Effects of specific land-use practices on plant and animal communities

Comparative studies of biodiversity in alternative land-use scenarios are particularly important to inform policy makers about the potential impacts of changes in land use on both conservation and livelihood goals (Gillison *et al.* 2004). Further research is needed to evaluate the utility of indicator groups for monitoring changes in biodiversity under changing land-use practices (Schulze *et al.* 2004, Barlow *et al.* 2007a, Gardner *et al.* 2008). Few studies have examined how agrochemicals, fire, machinery, introduced plant species, rotational grazing, tree pruning, polycropping, harvesting of natural products, or combinations of these practices affect biodiversity in the agricultural matrix or in forest remnants (including biological reserves). Studies should also include effects of ‘cryptic’ habitat degradation in intact forests, resulting from fires, logging, hunting, or climate change (Barlow & Peres 2006).

Abrupt and large-scale changes in land use often occur as market prices for certain commodities rise or fall, as international trade agreements take effect, or as domestic or international policies or subsidies change. How do these land-use changes affect biodiversity? In the Talamanca region of Costa Rica, diverse cacao and banana agroforestry systems are being converted to plantain monocultures due to increased market demand, leading to the loss of mammal, bird, and dung beetle diversity (Harvey *et al.* 2006, Harvey & González 2007). Bats were negatively affected by the use of chemical products in intensive agriculture in Chiapas, Mexico, but not by organic agricultural production (García-Estrada *et al.* 2006). We lack information to assess the impacts on biodiversity of rapid, large-scale changes in agricultural land-use due to expanding global markets and increasing rate of biofuel production (Righelato & Spracklen 2007).

Determining species complementarity among different landscape patches and modeling landscape patch dynamics can provide insights for planning landscape configurations critical for the conservation of species assemblages (*e.g.*, Cuarón 2000). Few studies have examined to what degree current species assemblages are relicts of previously forested areas or reflect community assembly processes arising *de novo* within the human-modified landscape (Davis *et al.* 2001). Further, we have a poor understanding of the influence of habitat

connectivity on alpha, beta, and gamma diversity within these landscapes.

5. Modeling potential impacts of climate change on biodiversity and species migration across human-modified landscapes

Climate change will have major impacts on agricultural production as well as on biodiversity within both human-modified landscapes and protected areas throughout the tropics (Williams & Hilbert 2006). Climate change threatens biodiversity by changing the availability and distribution of suitable habitat and microclimates, thereby placing additional stress on species already threatened by deforestation, habitat degradation, hunting, and other human activities (Malhi *et al.* 2008). As temperatures increase and precipitation regimes change, many species will need to move to higher elevations or toward the poles to find suitable habitat, as occurred during early Holocene warming (Bush 2002). Migration to cooler and moister conditions will be impeded if human modifications create barriers for species movement. In regions such as the Amazonian-Andean ecotone, continuous habitat corridors across rainfall, elevational and latitudinal gradients will be needed to avoid catastrophic species loss due to climate change (Bush 2002). We urge the development of models to investigate landscape-level effects of climate change on biodiversity and to provide guidance in landscape planning to mitigate the effects of climate change (Hannah *et al.* 2002). On the ground, adaptive management will be crucial to enhance ecosystem resistance, resilience, and the ability to adapt to changing climates at regional and local scales (Millar *et al.* 2007). Research is needed to identify how agricultural landscapes can be carefully designed and managed to maximize carbon sequestration and reduce emissions from deforestation and degradation (REDD), thereby ensuring these landscapes contribute to reducing the rate of climate change (Verchot *et al.* 2007). Feasibility studies are needed to evaluate REDD policies on biodiversity conservation, displacement of land-use change within and between countries, and other ecosystem values (Miles & Kapos 2008).

INTERACTIONS BETWEEN PEOPLE AND THEIR ENVIRONMENT.—Tropical landscapes have been shaped by the people who have lived in them and used them in both sustainable and unsustainable ways over past centuries (Denevan 2001, Whitmore & Turner 2001, Heckenberger *et al.* 2003, Toledo *et al.* 2003). For example, local indigenous knowledge and innovation in Chiapas, Mexico have been instrumental in designing coffee agroforests for multiple productive and subsistence uses (Soto-Pinto *et al.* 2007). Traditional as well as modern forms of sustainable land-use emphasize the values of ecosystem services derived from productive landscapes. According to the Millennium Ecosystem Assessment framework (MEA 2003), ecosystem services affecting human well being (basic material for a good life, health, good social relations, security and freedom of choice and action) include four broad categories: (1) provisioning (*e.g.*, food, fresh water, genetic resources, fiber, fuelwood, biochem-

icals); (2) regulating (*e.g.*, climate regulation, water regulation and purification, erosion regulation, disease regulation, pollination); (3) cultural (*e.g.*, recreation, spiritual values, social relations, aesthetic values); and (4) supporting (*e.g.*, primary production, soil formation, nutrient cycling). Further research is needed to link production, ecosystem services, and biological conservation (Maass *et al.* 2005, Bennett & Balvanera 2007). Environmental service payments are increasingly applied as incentives for some regulating ecosystem services such as carbon capture and hydrological control through forest conservation, silvopastoral systems, agroforestry, and reforestation within tropical regions (Montagnini & Nair 2004, Pagiola *et al.* 2004, Tschakert *et al.* 2007). These payments could potentially apply to a broader range of land uses and ecosystem services. Understanding how human-modified landscapes provide these services is particularly crucial against the backdrop of rapid climatic change and the emergence of international carbon markets that can fund reforestation and forest conservation activities (Boyd *et al.* 2006, Wara 2007, Miles & Kapos 2008). We propose four research priorities in this broad area.

6. Assessing ecosystem services across a range of habitat types in human-modified landscapes

Further research is needed to quantify the life support value of hydrologic, nutrient storage, and carbon storage services in a wide range of habitat types within human-modified landscapes, including silvopastoral systems, swidden agriculture, and agroforestry systems (Soto-Pinto *et al.* 2005, Tschakert *et al.* 2007). Quantifying these costs and benefits will ensure a more rigorous scientific basis for targeting environmental payment schemes and other incentives for conservation (Tschakert *et al.* 2005, Wunder 2005, Steffan-Dewenter *et al.* 2007). How do different landscape configurations affect ecosystem services that farmers and local residents depend on and benefit from? How do ecosystem services vary across a gradient of human modification? Are levels of ecosystem services correlated with biodiversity across habitats and landscapes (Chan *et al.* 2006)?

7. Examining relationships between biodiversity and ecosystem functions

Over the last decade, relationships between biodiversity and ecosystem function have been heavily studied in grassland systems, with a primary focus on relationships between biomass accumulation and nutrient retention in relation to plant diversity (*e.g.*, Loreau *et al.* 2001). Yet, understanding relationships between all types of biodiversity and all ecosystem functions and services in human-modified landscapes is extremely important. Didham *et al.* (1996) called for ecologists to delve into understanding the consequences of insect biodiversity loss in forest fragments for ecological function. Some advances have been made in assessing ecosystem services in human-modified landscapes. Several studies have shown the importance of bee diversity and off-farm plant diversity for pollination of coffee and other crops (Klein *et al.* 2003, Ricketts *et al.* 2004), the importance of birds and bird diversity for predatory services (*e.g.*, Perfecto *et al.* 2004, VanBael *et al.* 2008), and how

changes in agricultural management affect predatory effects of ants (Philpott *et al.* 2008b), but more such studies are required. A major challenge in determining direct links between biodiversity and ecosystem function stems from the difficulty of assessing the mechanisms driving positive biodiversity and ecosystem function relationships. Although it is clear that species complementarity (*e.g.*, differences in species resource use), facilitation, and dominance may all be important (Loreau *et al.* 2001), the tools available for distinguishing between mechanisms are not applicable to the majority of ecosystem services. For pollination, evidence that biodiversity is important is somewhat substantial, but our understanding of the relationships between vital ecosystem services and animal, microbial, and fungal diversity is still in its infancy and should receive attention. To advance our understanding, we need empirical studies of biodiversity and ecosystem function relationships in agricultural and forestry systems in the tropics, and we need to develop analytical tools for detecting patterns (Balvanera *et al.* 2005) and understanding underlying mechanisms. For which services is biodiversity important? How does functional richness relate to species richness? What are the mechanisms that drive biodiversity–ecosystem function relationships in agroecosystems? What agroecosystems and landscape configurations provide the highest levels of ecosystem functions?

8. Assessing the social and economic impact of conservation activities within human-modified landscapes

Planning sustainable production landscapes requires evaluation of trade-offs and synergies (Brown 2005), so that appropriate schemes can be created to ensure adequate financial benefits, equity, rights, and choices for rural people whose livelihoods and well-being depend upon sustainable agricultural production or resource extraction. There is a critical need to estimate and model opportunity costs of conservation within and across landscapes, as exemplified by studies of Naidoo and Ricketts (2006) in the Atlantic Forests of Paraguay. In these studies, models were used to plan locations of proposed biological corridors that maximized biodiversity conservation as well as ecosystem services benefits. Further, assessments of the social or economic impact of existing conservation efforts, such as environmental services payments and agri-environment schemes, can help to refine and improve them within a framework of adaptive management (Pagiola *et al.* 2005, Donald & Evans 2006).

9. Understanding relationships between human communities, local resources, and sustainable management

Investigators have amassed a considerable body of research on traditional ecological knowledge and its relevance to conservation and environmental management issues (Berkes 1999, Inglis 1993). These topics have been a focus of collaboration between natural and social scientists for many decades (Posey & Balee 1989, Redford & Padoch 1992). In Africa, researchers in national programs and nongovernmental agencies examine traditional methods for natural resource management (Abate *et al.* 2000). In Venezuela, indigenous groups are building an online

database to encourage more widespread, equitable exchange and use of traditional knowledge in solving environmental problems (http://www.slais.ubc.ca/COURSES/libr500/05-06-wr2/www/D_Ionson/index.htm). Indigenous groups are demanding to be incorporated in the whole research process, to have access to data and published information, and to participate as active stakeholders in the design of conservation research agendas (Mauro & Hardison 2000). But conservation research is still failing to address both conservation and social needs throughout the tropics (Meijaard & Sheil 2007). We need to define strategies for greater participation of rural resource users and other stakeholders in conservation research, including exploration of the ethical and human rights aspects of conservation policies (West 2006). Scientific understanding of sustainable practices should be integrated with the knowledge and innovations of rural resource users, including indigenous and non-indigenous peoples, landowners, and landless peasants (Diemont *et al.* 2005, Sayer *et al.* 2007). How can we use local and scientific knowledge to satisfy both conservation and rural peoples' needs, including the needs of impoverished people (Kaimowitz & Sheil 2007)?

We need to investigate the social, political, economic, and institutional organization of local resource use, availability, access, and tenure and how these dynamics interact at local, regional, national, and transnational scales (Dietz *et al.* 2003). How do institutional, political, and legal frameworks constrain or support conservation in human-modified landscapes? Further, we need to develop and document the new relationships among agricultural, biological and social scientists, farmers, consumers, and local and regional governments that can arise from a landscape approach. Designing successful conservation strategies requires an understanding of how and why local residents manage their landscapes and adapt to environmental and demographic changes (Shanker *et al.* 2005, Harvey *et al.* 2007).

RESTORATION ECOLOGY.—Extensive areas of the tropics have been heavily degraded by inappropriate land use, especially extensive cattle grazing (Lamb *et al.* 2005). An estimated 350 million ha in the tropics are classified as degraded due to inappropriate use of fire, land clearing, poor grazing management, and destructive harvesting of ecosystem resources (Maginnis & Jackson 2005). In contrast with the global north, where most restoration research has taken place and where most people are urban, in the global south large, often impoverished rural populations live and work in direct contact with tropical landscapes. This situation creates a distinct set of challenges and opportunities for restoration (Armesto *et al.* 2007): (1) restoration budgets are minimal, requiring low-cost restoration approaches or techniques that pay for themselves within a production context; (2) local residents can often bring deep ecological knowledge and traditional management techniques to bear on restoration challenges; and (3) criteria for restoration success in tropical landscapes should include the well-being of local people, the strengthening of cultural ties with the landscape, and synergies between traditional and scientific knowledge. In such a context, restoration goals must

build from and respect cultural landscape history (Higgs 2003), while avoiding criteria rigidly based upon some previous historic moment or notion of pristine wilderness. Creative approaches to meeting these challenges will require collaboration among multidisciplinary teams, decision makers and local people (Chazdon 2008). During the past decade, many reforestation and restoration activities and agroforestry projects have been developed to meet specific conservation goals, but few studies have evaluated the impact of these programs on biodiversity or ecosystem services at the landscape scale (Chazdon 2008). Here, we present three research priorities in restoration ecology.

10. **Landscape-level restoration research**

Restoration research in degraded tropical lands has generally been conducted at small spatial scales, yet we need to begin to adopt a landscape-level approach to restoration of habitats as well as agricultural productivity (Holl *et al.* 2003, Dudley *et al.* 2005, Lindenmayer *et al.* 2008). This perspective requires restoration efforts linking existing forest remnants within the landscape to form buffer zones, regeneration nuclei, and biological corridors. How can degraded areas be rehabilitated to enhance agricultural productivity, biodiversity, and human welfare at the landscape level? What is the role of spatial configuration and matrix composition in site restoration (Shono *et al.* 2006)? What is the effect of local site restoration on neighboring forest patches, protected areas, or regenerating forests? Munro *et al.* (2007) emphasize the need to better understand the balance between quantity and quality of revegetation for assessing responses of different animal groups. Although planting of local native plant species is expected to benefit local fauna, few studies have directly addressed this assumption (Munro *et al.* 2007).

11. **Evaluating costs and benefits of different restoration objectives**

We lack a framework for assessing the costs and benefits of different objectives of habitat-based restoration. These costs and benefits need to be assessed on both economic and biological bases, and should include benefits of ecosystem services. Further, we lack experimental studies that evaluate the potential to combine a range of objectives through reforestation, such as commercial timber harvest, restoration of soil fertility, carbon sequestration, and wildlife habitat. Which of these goals are spatially and temporally compatible within a single restoration project? Can restoration projects have different short-term versus long-term goals?

12. **Evaluating effects of livestock on restoration**

Interactions among livestock, grazing management practices, fire regime, invasive species, and seed dispersal during early stages of succession are poorly investigated with regard to restoration processes (Miceli-Méndez *et al.* 2008). Although it is often assumed that active cattle pastures have little regenerative potential, studies indicate that extensively managed pastures often retain significant regenerative ability. For example, in Muy Muy, Nicaragua, 37 of 85 tree species were

able to regenerate successfully under extensive grazing systems (Esquivel *et al.* 2008). Although fencing off pasture areas around remnant trees can enhance regeneration (Laborde *et al.* 2008), in some cases livestock can potentially assist early stages of forest regeneration by dispersing seed and reducing grass cover and fuel loads, improving site conditions for seedling establishment (Janzen & Martin 1982, Posada *et al.* 2000). Cattle browsing on forest edges, for example, might be managed so as to spread seeds of useful trees in open pasture, improving habitat for birds and insects while diversifying fodder (Miceli-Méndez *et al.* 2008). In a landscape restoration context, these cattle-dispersed trees might then serve as regeneration nuclei, as do remnant trees in pastures (Laborde *et al.* 2008). Similarly, the inclusion of live fences and windbreaks within agricultural landscapes can help facilitate natural regeneration processes, by attracting native seed-dispersing animals, such as bats and birds, and by ameliorating microclimatic conditions for seedling establishment (Harvey 2000, Harvey *et al.* 2005).

CONCLUSIONS

The conservation challenges that we face today in the tropics appear more intractable than they were only a few decades ago. We have acquired a new vision of the complexity and interrelatedness of tropical rural landscapes that calls for a new approach to research and management. Conserving biodiversity requires taking bold steps beyond the protection of areas minimally impacted by past or present human activities. A new conservation paradigm must incorporate human-modified landscapes in assessment of biodiversity and ecosystem services, planning of corridors and buffer zones, and restoration of degraded lands. This paradigm requires strong scientific foundations across a range of land uses and landscapes, including smallholder agroforestry (Schroth *et al.* 2004, Bhagwat *et al.* 2008), swidden agriculture (Tschakert *et al.* 2007), rangelands (Harvey *et al.* 2007), monoculture plantations (Cyranski 2007, Koh & Wilcove 2007, Turner *et al.* 2008), and logged forests (Meijaard & Sheil 2007), to name a few. The big picture must also incorporate incentives and opportunity costs for multiple stakeholders.

The research agenda we propose is best accomplished within an interdisciplinary framework, involving teams of researchers from the biological and social sciences with backgrounds in ecology, taxonomy, systematics, agronomy, agroecology, economics, geography, forestry, communication, sociology, anthropology, law and other social sciences. Successful outcomes will advance by building new partnerships in research, management, assessment, and policy. Viewing human-modified landscapes as a research arena creates direct linkages between conservationists, social and natural scientists, and local communities, so that farmers and other peoples can enjoy sustainable rural livelihoods (Harvey *et al.* 2008). These steps will help to guide adaptive management responses to sustain biodiversity and ecosystem services in a rapidly changing world.

ACKNOWLEDGMENTS

This paper was written by a subgroup of the Working Group on Biodiversity and Conservation Value of Agricultural Landscapes of Mesoamerica supported by the National Center for Ecological Analysis and Synthesis (NCEAS). Funding for the working group was provided by National Science Foundation (grant DEB-0072909), the University of California, and the University of California's Santa Barbara campus. The working group was organized by R.C. and D.G. We thank E. Bruna, T. Gardner, S. Paladino, and three anonymous reviewers for constructive comments on earlier versions of this manuscript.

LITERATURE CITED

- ABARCA, E. L. 2006. Birds, traditional coffee plantations and spatial complexity: The diversity puzzle. PhD thesis, Wageningen University, The Netherlands.
- ABATE, T., A. VAN HUIS, AND J. K. O. AMPOFO. 2000. Pest management strategies in traditional agriculture: An African perspective. *Ann. Rev. Entomol.* 45: 631–659.
- ALCORN, J. 1990. Indigenous agroforestry systems in the Latin American tropics. *In* M. A. Altieri, and S. B. Hecht (Eds.). *Agroecology and small farm development*, pp. 203–213. CRC Press, Boca Raton, Florida.
- ANTONGIOVANNI, M., AND J. P. METZGER. 2005. Influence of matrix habitats on the occurrence of insectivorous bird species in Amazonian forest fragments. *Biol. Conserv.* 122: 441–451.
- ANZURES-DADDA, A., AND R. H. MANSON. 2007. Patch- and landscape-scale effects on howler monkey distribution and abundance in rainforest fragments. *Anim. Conserv.* 10: 69–76.
- ARMESTO, J. J., S. BAUTISTA, E. DEL VAL, B. FERGUSON, X. GARCÍA, A. GAXIOLA, H. GODINEZ-ALVAREZ, G. GUNN, FABIOLA LÓPEZ-BARRERA, R. MANSON, M. NÚÑEZ-AVILA, C. ORTIZ-ARRONA, P. TOGNETTI, AND G. WILLIAMS-LINERA. 2007. Towards an ecological restoration network of the Americas: Challenges and opportunities for reverting land degradation. *Front. Ecol. Environ.* 5: w1–w4.
- ARMITAGE, D., M. MARSCHKE, AND R. PLUMMER. 2008. Adaptive co-management and the paradox of learning. *Global Environ. Change* 18: 86–98.
- BALVANERA, P., C. K. KREMEN, AND M. MARTÍNEZ-RAMOS. 2005. Applying community structure analysis to ecosystem function: Examples from pollination and carbon storage. *Ecol. Appl.* 15: 360–375.
- BARLOW, J., AND C. PERES. 2006. Consequences of cryptic and recurring fire disturbances for ecosystem structure and biodiversity in Amazonian forests. *In* W. F. Laurance, and C. A. Peres (Eds.). *Emerging threats to tropical forests*. Chicago University Press, Chicago, Illinois.
- BARLOW, J., T. A. GARDNER, I. S. ARAUJO, T. C. ÁVILA-PIRES, A. B. BONALDO, J. E. COSTA, M. C. ESPOSITO, L. V. FERREIRA, J. HAWES, M. I. M. HERNANDEZ, M. S. HOOGMOED, R. N. LEITE, N. F. L-MAN-HUNG, J. R. MALCOLM, M. B. MARTINS, L. A. M. MESTRE, R. MIRANDA-SANTOS, A. L. NUNES-GUTJAHN, W. L. OVERAL, L. PARRY, S. L. PETERS, M. A. RIBERO-JUNIOR, M. N. F. DA SILVA, C. DA SILVA MOTTA, AND C. A. PERES. 2007a. Quantifying the biodiversity value of tropical primary, secondary, and plantation forests. *PNAS* 104: 18555–18560.
- BARLOW, J., L. A. M. MESTRE, T. A. GARDNER, AND C. A. PERES. 2007b. The value of primary, secondary and plantation forests for Amazonian birds. *Biol. Conserv.* 136: 212–231.
- BARLOW, J., W. L. OVERAL, I. S. ARAUJO, T. A. GARDNER, AND C. A. PERES. 2007c. The value of primary, secondary and plantation forests for fruit-feeding butterflies in the Brazilian Amazon. *J. Appl. Ecol.* 44: 1001–1012.
- BAWA, K. S., W. J. KRESS, N. M. NADKARNI, AND S. LELE. 2004. Beyond paradise—Meeting the challenges in tropical biology in the 21st century. *Biotropica* 36: 437–446.
- BENGTSSON, J., A. BERGMAN, M. OLSSON, AND J. ÖRBERG. 2003. Reserves, resilience and dynamic landscapes. *Ambio* 32: 389–396.
- BENNETT, A. E., J. Q. RADFORD, AND A. HASLEM. 2006. Properties of land mosaics: Implications for nature conservation in agricultural environments. *Biol. Conserv.* 133: 250–264.
- BENNETT, E. M., AND P. BALVANERA. 2007. The future of production systems in a globalized world. *Front. Ecol. Environ.* 5: 191–198.
- BERKES, F. 1999. *Sacred ecology: Traditional ecological knowledge and resource management*. Taylor and Francis, Philadelphia, Pennsylvania.
- BHAGWAT, S. A., K. J. WILLIS, H. J. B. BIRKS, AND R. J. WHITTAKER. 2008. Agroforestry: A refuge for tropical biodiversity? *Trends Ecol. Evol.* 23: 261–267.
- BIANCONI, G. V., S. B. MIKICH, AND W. A. PEDRO. 2006. Movements of bats (Mammalia, Chiroptera) in Atlantic Forest remnants in southern Brazil. *Rev. Bras. Zool.* 23: 1199–1206.
- BONGERS, F., A. WASSIE, F. STERCK, T. BEKELE, AND D. TEETAY. 2006. Ecological restoration and church forests in northern Ethiopia. *J. Drylands* 1: 35–45.
- BOSHIER, D. H., J. E. GORDON, AND A. J. BARRANCE. 2004. Prospects for *circa situm* tree conservation in Mesoamerican dry-forest agro-ecosystems. *In* G. W. Frankie, A. Mata, and S. B. Vinson (Eds.). *Biodiversity conservation in Costa Rica*, pp. 210–226. University of California Press, Berkeley, California.
- BOYD, E., M. GURIERREZ, AND M. CHANG. 2006. Small-scale forest carbon projects: Adapting CDM to low-income communities. *Global Environ. Change* 17: 250–259.
- BROWN, K. 2005. Addressing trade-offs in forest landscape restoration. *In* S. Mansourian, D. Vallauri, and N. Dudley (Eds.). *Forest restoration in landscapes: Beyond planting trees*, pp. 59–62. Springer, New York, New York.
- BUSH, M. B. 2002. Distributional change and conservation on the Andean flank: A paleoecological perspective. *Global Ecol. Biogeogr.* 11: 463–473.
- CASCANTE, A., M. QUESADA, J. J. LOBO, AND E. A. FUCHS. 2002. Effects of dry tropical forest fragmentation on the reproductive success and genetic structure of the tree *Samanea saman*. *Conserv. Biol.* 16: 137–147.
- CHAN, K. M. A., M. R. SHAW, D. R. CAMERON, E. C. UNDERWOOD, AND G. C. DAILY. 2006. Conservation planning for ecosystem services. *PLoS Biology* 4: e379.
- CHAZDON, R. L. 2008. Beyond deforestation: Restoring forests and ecosystem services on degraded lands. *Science* 320: 1458–1460.
- CHAZDON, R. L., C. A. HARVEY, M. MARTÍNEZ-RAMOS, P. BALVANERA, K. STONER, J. SCHONDUBE, L. D. AVILA CABADILLA, AND M. FLORES-HIDALGO. 2008. Tropical dry forest biodiversity and conservation value in agricultural landscapes of Mesoamerica. *In* R. Dirzo, H. A. Mooney, G. Ceballos, and H. Young (Eds.). *Ecology and conservation of Neotropical dry forests*. Island Press, Washington, DC.
- CHAZDON, R. L., S. G. LETCHER, M. VAN BREUGEL, M. MARTÍNEZ-RAMOS, F. BONGERS, AND B. FINEGAN. 2007. Rates of change in tree communities of secondary tropical forests following major disturbances. *Phil. Trans. R Soc. B* 362: 273–289.
- CRAMER, J. M. R., C. G. MESQUITA, T. VIZCARRA BENTOS, B. MOSER, AND G. B. WILLIAMSON. 2007. Forest fragmentation reduces seed dispersal of *Duckeodendron cestroides*, a Central Amazon endemic. *Biotropica* 39: 709–718.
- CRUZ-ANGÓN, A., T. S. SILLETT, AND R. GREENBERG. 2008. An experimental study of habitat selection by birds in a coffee plantation. *Ecology* 89: 921–927.
- CUARÓN, A. 2000. Effects of land-cover changes on mammals in a Neotropical region: A modeling approach. *Conserv. Biol.* 14: 1676–1692.
- CULLEN, L. JR., K. ALGER, AND D. M. RAMBALDI. 2005. Land reform and biodiversity conservation in Brazil in the 1990s: Conflict and the articulation of mutual interests. *Conserv. Biol.* 19: 747–755.

- CYRANOSKI, D. 2007. Logging: The new conservation. *Nature* 446: 608–610.
- DAILY, G. C., G. CEBALLOS, J. PACHECO, G. SUZAN, AND A. SANCHEZ-AZOFEIFA. 2003. Countryside biogeography of Neotropical mammals: Conservation opportunities in agricultural landscapes of Costa Rica. *Conserv. Biol.* 17: 1814–1826.
- DAILY, G. C., P. R. EHRlich, AND G. A. SANCHEZ-AZOFEIFA. 2001. Countryside biogeography: Use of human-dominated habitats by the avifauna of southern Costa Rica. *Ecol. Appl.* 11: 1–13.
- DAVIS, A. J., J. D. HOLLOWAY, H. HUIJBREGTS, J. KRIKKEEN, A. H. KIRK-SPRIGGS, AND S. L. SUTTON. 2001. Dung beetles as indicators of change in the forests of northern Borneo. *J. Appl. Ecol.* 38: 593–616.
- DEFRIES, R., A. HANSEN, A. C. NEWTON, AND M. C. HANSEN. 2005. Increasing isolation of protected areas in tropical forests over the past twenty years. *Ecol. Appl.* 15: 19–26.
- DEFRIES, R., A. HANSEN, B. L. TURNER, R. REID, AND J. G. LIU. 2007. Land use change around protected areas: Management to balance human needs and ecological function. *Ecol. Appl.* 17: 1031–1038.
- DENEVAN, W. M. 2001. *Cultivated landscapes of Native Amazonia and the Andes*. Oxford University Press, Oxford, UK.
- DIDHAM, R. K., J. GHAZOUL, N. E. STORK, AND A. J. DAVIS. 1996. Insects in fragmented forests: A functional approach. *Trends Ecol. Evol. Biol.* 11: 255–260.
- DIEMONT, S. A. W., J. E. MARTIN, S. I. LEVY-TACHER, R. B. NIGH, P. RAMIREZ LOPEZ, AND J. D. GOLICHER. 2005. Lacandon Maya forest management: Restoration of soil fertility using native tree species. *Ecol. Eng.* 28: 205–212.
- DIETZ, T., E. OSTROM, AND P. C. STERN. 2003. The struggle to govern the commons. *Science* 302: 1907–1912.
- DONALD, P. F. 2004. Biodiversity impacts of some agricultural commodity production systems. *Conserv. Biol.* 18: 17–38.
- DONALD, P. F., AND A. D. EVANS. 2006. Habitat connectivity and matrix restoration: The wider implications of agri-environment schemes. *J. Appl. Ecol.* 43: 209–218.
- DUDLEY, N., J. MORRISON, J. ARONSON, AND S. MANSOURIAN. 2005. Why do we need to consider restoration in a landscape context? *In* S. Mansourian, D. Vallauri, and N. Dudley (Eds.). *Forest restoration in landscapes: Beyond planting trees*, pp. 51–58. Springer, New York, New York.
- DUNN, R. R. 2004. Recovery of faunal communities during tropical forest regeneration. *Conserv. Biol.* 10: 302–309.
- ESQUIVEL, M. J., C. A. HARVEY, B. FINEGAN, F. CASANOVES, AND C. SKARPE. 2008. Effects of pasture management on the natural regeneration of Neotropical trees. *Appl. Ecol.* 45: 371–380.
- ESTRADA, A., R. COATES-ESTRADA, AND D. A. MERITT. 1997. Anthropogenic landscape changes and avian diversity at Los Tuxtlas, Mexico. *Biodiversity Conserv.* 6: 19–43.
- ESTRADA, A., C. JIMÉNEZ, A. RIVERA, AND E. FUENTES. 2004. General bat activity measured with an ultrasound detector in a fragmented tropical landscape in Los Tuxtlas, Mexico. *Anim. Biodivers. Conserv.* 27: 1–9.
- FAZEY, I., J. FISCHER, AND D. B. LINDENMAYER. 2005. What do conservation biologists publish? *Biol. Conserv.* 124: 63–73.
- FISCHER, J., D. B. LINDENMAYER, AND A. D. MANNING. 2006. Biodiversity, ecosystem function, and resilience: Ten guiding principles for commodity production landscapes. *Front. Ecol. Environ.* 4: 80–86.
- GARCÍA-ESTRADA, C., A. DAMON, C. SANCHEZ, L. SOTO-PINTO, G. IBARRA-NUÑEZ. 2006. Bat diversity in montane rainforest and shaded coffee under different management regimes in southeastern Chiapas, Mexico. *Biol. Conserv.* 132: 351–361.
- GARDNER, T. A., J. BARLOW, L. W. PARRY, AND C. A. PERES. 2007a. Predicting the uncertain future of tropical forest species in a data vacuum. *Biotropica* 39: 25–30.
- GARDNER, T. A., J. BARLOW, I. S. ARAUJO, T. C. AVILA-PIRES, A. B. BONALDO, J. E. COSTA, M. C. ESPOSITO, L. V. FERREIRA, J. HAWES, M. I. M. HERNANDEZ, M. S. HOOGMOED, R. N. LEITE, N. F. LO-MAN-HUNG, J. R. MALCOLM, M. B. MARTINS, L. A. M. MESTRE, R. MIRANDA-SANTOS, W. L. OVERAL, L. PARRY, S. L. PETERS, M. A. RIBEIRO-JUNIOR, M. N. F. DA SILVA, C. DA SILVA MOTTA, AND C. A. PERES. 2008. The cost-effectiveness of biodiversity surveys in tropical forests. *Ecol. Lett.* 11: 139–150.
- GARDNER, T. A., M. A. RIBEIRO-JUNIOR, J. BARLOW, T. C. S. ÁVILA-PIRES, M. S. HOOGMOED, AND C. A. PERES. 2007b. The value of primary, secondary, and plantation forests for a Neotropical herpetofauna. *Conserv. Biol.* 21: 775–787.
- GASCON, C., T. E. LOVEJOY, R. O. BIERREGAARD, J. R. MALCOLM, P. C. STOUFFER, H. L. VASCONCELOS, W. F. LAURANCE, B. ZIMMERMAN, M. TOCHER, AND S. BORGES. 1999. Matrix habitat and species richness in tropical forest remnants. *Biol. Conserv.* 91: 223–229.
- GILLISON, A. N., N. LISWANTI, S. BUDIDARSONO, M. VAN NOORDWIJ, AND T. P. TOMICH. 2004. Impact of cropping methods on biodiversity in coffee agroecosystems in Sumatra, Indonesia. *Ecol. Soc.* 9: 7. Available at: <http://www.ecologyandsociety.org/vol9/iss2/art7/>.
- GIRAÓ, L. C., A. V. LOPES, M. TABARELLI, AND E. M. BRUNA. 2007. Changes in tree reproductive traits reduce functional diversity in a fragmented Atlantic forest landscape. *PLoS ONE* 2: e908. doi:10.1371/journal.pone.0000908.
- GREENBERG, R., I. PERFECTO, AND S. M. PHILPOTT. 2008. Agroforests as model systems for tropical ecology. *Ecology* 89: 913–914.
- GUEVARA, S., J. LABORDE, AND G. SÁNCHEZ-RIOS. 2005. Los árboles que la selva dejó atrás. *Interciencia* 30: 595–601.
- HANNAH, L., G. F. MIDGLEY, AND D. MILLAR. 2002. Climate change-integrated conservation strategies. *Global Ecol. Biogeogr.* 11: 485–495.
- HANSEN, A. J., AND R. DEFRIES. 2007. Ecological mechanisms linking protected areas to surrounding lands. *Ecol. Appl.* 17: 974–988.
- HARVEY, C. A. 2000. Windbreaks enhance seed dispersal into agricultural landscapes in Monteverde, Costa Rica. *Ecol. Appl.* 10: 155–173.
- HARVEY, C. A. 2007. Designing agricultural landscapes for biodiversity conservation. *In* S. J. Scherr, and J. A. McNeely (Eds.). *Farming with nature: The science and practice of coagriculture*. Island Press, Washington, D.C.
- HARVEY, C. A., AND J. GONZÁLEZ. 2007. Agroforestry systems conserve species-rich but modified assemblages of tropical birds and bats. *Biodiv. Conserv.* 16: 2257–2292.
- HARVEY, C. A., AND J. C. SÁENZ (Eds.). 2007. *Evaluación y Conservación de la Biodiversidad en Agropaisajes de Mesoamérica*. INBio, Heredia, Costa Rica.
- HARVEY, C. A., J. GONZÁLEZ, AND E. SOMARRIBA. 2006. Dung beetle and mammal diversity in forests, indigenous agroforestry systems and plantation monocultures in Talamanca, Costa Rica. *Biodivers. Conserv.* 15: 555–585.
- HARVEY, C. A., O. KOMAR, R. CHAZDON, B. G. FERGUSON, B. FINEGAN, D. M. GRIFFITH, M. MARTÍNEZ-RAMOS, H. MORALES, R. NIGH, L. SOTO-PINTO, M. VAN BREUGEL, AND M. WISHNIE. 2008. Integrating agricultural landscapes with biodiversity conservation in the Mesoamerican hotspot: Opportunities and an action agenda. *Conserv. Biol.* 22: 8–15.
- HARVEY, C. A., C. VILLANUEVA, M. IBRAHIM, R. GÓMEZ, M. LÓPEZ, S. KUNTH, AND F. L. SINCLAIR. 2007. Finqueros, árboles y producción ganadera en paisajes Centroamericanos: Implicaciones para la conservación de biodiversidad. *In* C. A. Harvey, and J. C. Sáenz (Eds.). *Evaluación y Conservación de la Biodiversidad en Agropaisajes de Mesoamérica*. INBio, Heredia, Costa Rica.
- HARVEY, C. A., C. VILLANUEVA, J. VILLACÍS, M. CHACÓN, D. MUÑOZ, M. LÓPEZ, M. IBRAHIM, R. TAYLOR, J. L. MARTÍNEZ, A. NAVAS, J. SÁENZ, D. SÁNCHEZ, A. MEDINA, S. VÍLCHEZ, B. HERNÁNDEZ, A. PÉREZ, F. RUIZ, F. LÓPEZ, I. LANG, S. KUNTH, AND F. L. SINCLAIR. 2005. Contribution of live fences to the ecological integrity of agricultural landscapes. *Agric., Ecosyst. Environ.* 111: 200–230.
- HECKENBERGER, M. J., A. KUIKURO, U. T. KUIKURO, J. C. RUSSELL, M. SCHMIDT, C. FAUSTO, AND B. FRANCHETTO. 2003. Amazonia 1492: Pristine forest or cultural parkland? *Science* 302: 1710–1715.

- HAUSER, C. E., AND H. POSSINGHAM. 2008. Experimental or precautionary? Adaptive management over a range of time horizons. *J. Appl. Ecol.* 45: 72–81.
- HIGGS, E. 2003. *Nature by design: People, natural processes and ecological restoration*. Massachusetts Institute of Technology, Cambridge, Massachusetts.
- HOLL, K. D., E. E. CRONE, AND C. B. SCHULTZ. 2003. Landscape restoration: Moving from generalities to methodologies. *BioScience* 53: 491–502.
- INGLIS, J. T. (Ed). 1993. *Traditional ecological knowledge: Concepts and cases*. International Development Research Center, Ottawa, Canada.
- INTERNATIONAL ASSESSMENT OF AGRICULTURAL KNOWLEDGE, SCIENCE AND TECHNOLOGY FOR DEVELOPMENT (IAASTD). 2008. Available at: http://www.agassessment.org/index.cfm?Page=About_IAASTDandItemID=2. Accessed April 15, 2008.
- JACKSON, L. E., U. PASCUAL, AND T. HODGKIN. 2007. Utilizing and conserving agrobiodiversity in agricultural landscapes. *Agric., Ecosyst. Environ.* 121: 196–210.
- JANZEN, D. H. 1983. No park is an island: Increase in interference from outside as park size decreases. *Oikos* 41: 402–410.
- JANZEN, D. H., AND P. S. MARTIN. 1982. Neotropical anachronisms: The fruits the gomphotheres ate. *Science* 215: 19–27.
- JULES, E. S., AND P. SHAHANI. 2003. A broader ecological context to habitat fragmentation: Why matrix habitat is more important than we thought. *J. Veg. Sci.* 14: 459–464.
- KABIR, M. D. E., AND E. L. WEBB. 2008. Can homegardens conserve biodiversity in Bangladesh? *Biotropica* 40: 95–103.
- KAIMOWITZ, D., AND D. SHEIL. 2007. Conserving what and for whom? Why conservation should help meet basic human needs in the tropics. *Biotropica* 39: 567–574.
- KHUMBONGMAYUM, A. D., M. L. KHAN, AND R. S. TRIPATHI. 2005. Sacred groves of Manipur, northeast India: Biodiversity value, status and strategies for their conservation. *Biodivers. Conserv.* 14: 1541–1582.
- KLEIN, A. M., A. CUNNINGHAM, M. BOS, AND I. STEFFAN-DEWENTER. 2008. Advances in pollination ecology from tropical plantation crops. *Ecology* 89: 935–943.
- KLEIN, A. M., I. STEFFAN-DEWENTER, AND T. TSCHARNTKE. 2003. Fruit set of highland coffee increases with the diversity of pollinating bees. *Proc. R. Soc. Lond. B-Biol. Sci.* 270: 955–961.
- KLEIN, A. M., I. STEFFAN-DEWENTER, AND T. TSCHARNTKE. 2006. Rain forest promotes trophic interactions and diversity of trap-nesting Hymenoptera in adjacent agroforestry. *J. Anim. Ecol.* 75: 315–323.
- KLEIN, B. C. 1989. Effects of forest fragmentation on dung and carrion beetle communities in Central Amazonia. *Ecology* 70: 1715–1725.
- KOH, L. P., AND D. S. WILCOVE. 2007. Cashing in palm oil for conservation. *Nature* 448: 993–994.
- KOMAR, O. 2006. Ecology and conservation of birds in coffee plantations: A critical review. *Bird Conserv. Int.* 16: 1–23.
- KREMEN, C., WILLIAMS, N. M., AND R. W. THORP. 2002. Crop pollination from native bees at risk from agricultural intensification. *Proc. Natl. Acad. Sci. U.S.A.* 99: 16812–16816.
- KUPFER, J. A., G. P. MALANSON, AND S. B. FRANKLIN. 2006. Not seeing the ocean for the islands: The mediating influence of matrix-based processes on forest fragmentation effects. *Global Ecol. Biogeogr.* 15: 8–20.
- LABORDE, J., S. GUEVARA, AND G. SÁNCHEZ-RÍOS. 2008. Trees and shrub seed dispersal in pastures: The importance of rainforest trees outside forest fragments. *Ecoscience* 15: 6–16.
- LAMB, D., ERSKINE, P. D., AND PARROTTA, J. 2005. Restoration of degraded tropical forest landscapes. *Science* 310: 1628–1632.
- LAURANCE, S. G. W. 2004. Landscape connectivity and biological corridors. *In* G. A. Schroth, B. Fonseca, C. A. Harvey, C. Gascon, H. L. Vasconcelos, and A. M. N. Izac (Eds.). *Agroforestry and biodiversity conservation in tropical landscapes*, pp. 50–63. Island Press, Washington, DC.
- LAURANCE, W. F., H. E. M. NASCIMENTO, S. G. LAURANCE, A. C. ANDRADE, P. M. FLARNISIDE, J. E. L. RIBEIRO, AND R. L. CAPRETZ. 2006. Rain forest fragmentation and the proliferation of successional trees. *Ecology* 87: 469–482.
- LINDENMAYER, D. B., AND G. LUCK. 2005. Synthesis: Thresholds in conservation and management. *Biol. Conserv.* 124: 351–354.
- LINDENMAYER, D. B., R. J. HOBBS, R. MONTAGUE-DRAKE, J. ALEXANDRA, A. BENNETT, M. BURGMAN, P. CALE, A. CALHOUN, V. CRAMER, P. CULLEN, D. DRISCOLL, L. FAHRIG, J. FISCHER, J. FRANKLIN, Y. HALLA, M. HUNTER, P. GIBBONS, S. LAKE, G. LUCK, C. MACGREGOR, S. MCINTYRE, R. MAC NALLY, A. MANNING, J. MILLER, H. MOONEY, R. NOSS, H. POSSINGHAM, D. SAUNDERS, F. SCHMIEGELOW, M. SCOTT, D. SIMBERLOFF, T. SISK, G. TABOR, B. WALKER, J. WIENS, J. WOJNARSKI, E. ZAVALA. 2008. A checklist for ecological management of landscapes for conservation. *Ecol. Lett.* 11: 78–91.
- LOREAU, M., S. NAEEM, P. INCHAUSTI, J. BENGTSOON, J. P. GRIME, A. HECTOR, D. U. HOOPER, M. A. HUSTON, D. RAFFAELLI, B. SCHMID, D. TILMAN, AND D. A. WARDLE. 2001. Biodiversity and ecosystem functioning: Current knowledge and future challenges. *Science* 294: 804–808.
- MEA (MILLENNIUM ECOSYSTEM ASSESSMENT). 2003. *Ecosystems and human well being*. Island Press, Washington, DC.
- MAASS, J. M., P. BALVANERA, A. CASTILLO, G. C. DAILY, H. A. MOONEY, P. EHRLICH, M. QUESADA, A. MIRANDA, V. J. JARAMILLO, F. GARCÍA-OLIVA, A. MARTÍNEZ-YRIZAR, H. COTLER, J. LÓPEZ-BLANCO, J. A. PÉREZ-JIMÉNEZ, A. BÚRQUEZ, C. TINOCO, G. CEBALLOS, L. BARRAZA, R. AYALA, AND J. SARUKHÁN. 2005. Ecosystem services of tropical dry forests: Insights from long-term ecological and social research on the Pacific Coast of Mexico. *Ecol. Soc.* 10: 17 (online).
- MAGINNIS, S., AND W. JACKSON. 2005. Balancing restoration and development. *ITTO Tropical Forest Update* 15/2.
- MALHI, Y., J. T. ROBERTS, R. A. BETTS, T. J. KILEEN, W. LI, AND C. A. NOBRE. 2008. Climate change, deforestation, and the fate of the Amazon. *Science* 319: 169–172.
- MAURO, F., AND P. D. HARDISON. 2000. Traditional knowledge of indigenous and local communities: International debate and policy initiatives. *Ecol. Appl.* 10: 1263–1269.
- MAYFIELD, M. M. BONI M. F., DAILY, G. C., AND ACKERLY, D. 2005. Species and functional diversity of native and human-dominated plant communities. *Ecology* 86: 2365–2372.
- MCNEELY, J. A., AND S. J. SCHERR. 2003. *Ecoagriculture: Strategies to feed the world and save wild biodiversity*. Island Press, Washington, DC.
- MEDINA, A., C. A. HARVEY, D. SÁNCHEZ, S. VÍLCHEZ, AND B. HERNÁNDEZ. 2007. Bat diversity and movement in a Neotropical agricultural landscape. *Biotropica* 39: 120–128.
- MEIJAARD, E., AND D. SHEIL. 2007. Is wildlife research useful for wildlife conservation in the tropics? A review for Borneo with global implications. *Biodiv. Conserv.* 16: 3053–3065.
- MICELI-MÉNDEZ, C. L., B. G. FERGUSON, AND N. RAMÍREZ-MARCIAL. 2008. Seed dispersal by cattle: Natural history and applications to Neotropical forest restoration and agroforestry. *In* R. Myer (Ed.) *Post-Agricultural succession in the Neotropics*, pp. 165–191. Springer, New York, New York.
- MICHALSKI, F., I. NISHI, AND C. A. PERES. 2007. Disturbance-mediated drift in tree functional groups in Amazonian forest fragments. *Biotropica* 39: 691–701.
- MILES, L., AND V. KAPOS. 2008. Reducing greenhouse gas emissions from deforestation and forest degradation: Global land-use implications. *Science* 320: 1454–1455.
- MILLAR, C. I., N. L. STEPHENSON, AND S. L. STEPHENS. 2007. Climate change and forests of the future: Managing in the face of uncertainty. *Ecol. Appl.* 17: 2145–2151.
- MOGUEL, P., AND V. M. TOLEDO. 1999. Biodiversity conservation in traditional coffee systems of Mexico. *Conserv. Biol.* 13: 11–21.
- MONRO, A. K., D. T. JONES, AND M. E. ARAUJO. 2006. Taxonomic capacity can improve environmental and economic sustainability in biodiversity-rich shade coffee farms in El Salvador. *Syst. Biodivers.* 4: 1–8.
- MONTAGNINI, F., AND P. K. R. NAIR. 2004. Carbon sequestration: An underexploited environmental benefit of agroforestry systems. *Agroforestry Syst.* 61–62: 281–295.

- MUNRO, N. T., D. B. LINDENMAYER, AND J. FISCHER. 2007. Faunal response to revegetation in agricultural areas of Australia: A review. *Ecol. Manage. Rest.* 8: 199–207.
- NAIDOO, R., AND T. H. RICKETTS. 2006. Assessing the economic costs and benefits of conservation. *PLoS Biol.* 4: 2153–2164.
- NASCIMENTO, H. E. M., A. C. S. ANDRADE, J. L. C. CAMARGO, W. F. LAURANCE, S. G. LAURANCE, AND J. E. I. RIBIERO. 2006. Effects of the surrounding matrix on tree recruitment in Amazonian forest fragments. *Conserv. Biol.* 20: 853–860.
- NICHOLS, E., T. B. LARSEN, S. SPECTOR, A. L. V. DAVIS, F. ESCOBAR, M. FAVILA, K. VULINEC, AND T. S. R. NETWORK. 2007. Global dung beetle response to tropical forest modification and fragmentation: A quantitative literature review and meta-analysis. *Biol. Conserv.* 137: 1–19.
- NORRIS, K. 2008. Agriculture and biodiversity conservation: Opportunity knocks. *Conserv. Lett.* 1: 2–11.
- OMER, A., U. PASCUAL, AND N. P. RUSSELL. 2007. Biodiversity conservation and productivity in intensive agricultural systems. *J. Agric. Econ.* 58: 308–329.
- PAGIOLA, S., A. AGOSTINI, J. GOBBI, C. DE HAAN, M. IBRAHIM, E. RAMÍREZ, M. ROSALES, AND J. P. RUÍZ. 2004. Paying for biodiversity conservation services in agricultural landscapes. The World Bank Environment Department, Paper No. 96. The World Bank, Washington, DC.
- PAGIOLA, S., A. ARCENAS, AND G. PLATAIS. 2005. Can payments for environmental services help reduce poverty? An exploration of the issues and the evidence to date from Latin America. *World Dev.* 33: 237–253.
- PEH, K. S.-H., N. S. SODHI, J. DE JONG, C. H. SEKERCIOGLU, C. A.-M. YAP, AND S. L.-H. LIM. 2006. Conservation value of degraded habitats for forest birds in southern Peninsular Malaysia. *Divers. Distrib.* 12: 572–581.
- PERFECTO, I., AND J. VANDERMEER. 2002. Quality of agroecological matrix in a tropical montane landscape: Ants in coffee plantations in southern Mexico. *Conserv. Biol.* 16: 174–182.
- PERFECTO, I., R. A. RICE, R. GREENBERG, AND M. E. VANDERVOORT. 1996. Shade coffee: A disappearing refuge for biodiversity. *Bioscience* 46: 598–608.
- PERFECTO, I., J. VANDERMEER, G. L. BAUTISTA, G. I. NUÑEZ, R. GREENBERG, P. BICHIER, AND S. LANGRIDGE. 2004. Greater predation in shaded coffee farms: The role of resident Neotropical birds. *Ecology* 85: 2677–2681.
- PERRINGS, C., L. JACKSON, K. BAWA, L. BRUSSAARD, S. BRUSH, T. GAVIN, R. PAPA, U. PASCUAL, AND P. DE RUITER. 2006. Biodiversity in agricultural landscapes: Natural capital without losing interest. *Conserv. Biol.* 20: 263–264.
- PETERS, C. M. 2000. Precolumbian silviculture and indigenous management of Neotropical forests. In D. L. Lentz (Ed.): *Imperfect balance*, pp. 203–223. Columbia University Press, New York, New York.
- PHILPOTT, S. M., W. ARENDT, I. ARMBRECHT, P. BICHIER, T. DIETSCH, C. GORDON, R. GREENBERG, I. PERFECTO, L. SOTO-PINTO, C. TEJEDACRUZ, G. WILLIAMS, AND J. VALENZUELA. 2008a. Biodiversity loss in Latin American coffee landscapes: Reviewing evidence on ants, birds, and trees. *Conserv. Biol.* 22: 1093–1105.
- PHILPOTT, S. M., I. PERFECTO, AND J. VANDERMEER. 2008b. Effects of predatory ants on lower trophic levels across a gradient of coffee management complexity. *J. An. Ecol.* 77: 505–511.
- PICKETT, C. H., AND R. L. BUGG. 1998. *Enhancing biological control: Habitat management to promote natural enemies of agricultural pests*. University of California Press, Berkeley, California.
- POSADA, J. M., T. M. AIDE, AND J. CAVELIER. 2000. Cattle and weedy shrubs as restoration tools of tropical montane rainforest. *Restoration Ecol.* 8: 370–379.
- POSEY, D. A., AND W. L. BALEE (Eds.). 1989. *Resource management in Amazonia: Indigenous and folk strategies*. New York Botanical Garden, Bronx, New York.
- POTVIN, C., P. TSCHAKERT, F. LEBEL, K. KIBRY, H. BARRIOS, J. BOCARIZA, J. CAISAMO, L. CAISAMO, C. CANSARI, J. CASAMÁ, M. CASAMÁ, L. CHAMORRA, N. DUMASA, S. GOLDENBERG, V. GUAINORA, P. HAYES, T. MOORE, J. RUÍZ. 2007. A participatory approach to the establishment of a baseline scenario for a reforestation clean development mechanism project. *Mitig. Adapt. Strat. Glob. Change* 12: 1341–1362.
- PRETTY, J. N. 1995. Participatory learning for sustainable agriculture. *World Dev.* 23: 1247–1263.
- PULIDO, M. T., T. VALVERDE, AND J. CABALLERO. 2007. Variation in the population dynamics of the palm *Sabal yapa* in a landscape shaped by shifting cultivation in the Yucatan Peninsula, Mexico. *J. Trop. Ecol.* 23: 139–149.
- RADFORD, J. W., A. F. BENNETT, AND G. J. CHEERS. 2005. Landscape-level thresholds of habitat cover for woodland-dependent birds. *Biol. Conserv.* 124: 317–337.
- REDFORD, K., AND C. PADOCH (Eds.). 1992. *Conservation of Neotropical forests: Working from traditional resource use*. Columbia University Press, New York, New York.
- REUNANEN, P., M. MÖNKKÖNEN, A. NIKULA, E. HURME, AND V. NIVALA. 2004. Assessing landscape thresholds for the Siberian flying squirrel. *Ecol. Bull.* 51: 277–286.
- RICKETTS, T. H., G. C. DAILY, P. R. EHREICH, AND C. D. MICHENER. 2004. Economic value of tropical forest to coffee production. *Proc. Natl. Acad. Sci. U.S.A.* 101: 12579–12582.
- RIGHELATO, R., AND D. V. SPRACKLEN. 2007. Carbon mitigation by biofuels or by saving and restoring forests? *Science* 317: 902.
- ROMERO, K., NARANJO, E., MORALES, H., AND R. NIGH. 2006. Plagas de vertebrados en milpas lacandonas. *Interciencia* 31: 276–283.
- SALAFSKY, N., R. MARGOLUIS, AND K. REDFORD. 2008. Adaptive management: A tool for conservation practitioners. Biodiversity Support Program, Washington, DC. Available at: http://fosonline.org/resources/Publications/AdapManHTML/adman_1.html (accessed May 3, 2008).
- SANGINGA, P. C., J. TUMWINE, AND N. K. LILJA. 2006. Patterns of participation in farmers' research groups: Lessons from the highlands of southwestern Uganda. *Agric. Hum. Values* 23: 501–512.
- SAYER, J., B. CAMPBELL, L. PETHERAM, M. ALDRICH, M. R. PEREZ, D. ENDAMANA, Z.-L. N. DONGMO, L. DEFO, S. MARIKI, N. DOGGART, AND N. BURGESS. 2007. Assessing environment and development outcomes in conservation landscapes. *Biodivers. Conserv.* 16: 2677–2694.
- SCHELHAS, J., AND R. GREENBERG. 1996. Introduction: The value of forest patches. In J. Schelhas, and R. Greenberg (Eds.). *Forest patches in tropical landscapes*, pp. xv–xxxvi. Island Press, Washington, DC.
- SCHERR, S. J., AND J. A. MCNEELY (Eds.). 2007. *Farming with nature: The science and practice of ecoagriculture*. Island Press, Washington, DC.
- SCHMIDT, M. H., C. THIES, C. AND T. TSCHARNTKE. 2004. Landscape context of arthropod biological control. In G. M. Gurr, S. D. Wratten, and M. A. Altieri (Eds.). *Ecological engineering for pest management: Advances in habitat manipulation for arthropods*, pp. 55–63. CSIRO, Collingwood, VIC.
- SCHROTH, G. A., G. A. B. DA FONSECA, C. A. HARVEY, H. L. VASCONCELOS, C. GASCON, AND A.-M. N. IZAC. 2004. Introduction: The role of agroforestry in biodiversity conservation in tropical landscapes. In G. A. Schroth, B. Fonseca, C. A. Harvey, C. Gascon, H. L. Vasconcelos, and A. M. N. Izac (Eds.). *Agroforestry and biodiversity conservation in tropical landscapes*. Island Press, Washington, DC.
- SCHULZE, C. H., M. WALTERT, P. J. A. KESSLER, R. PITOPANG, SHAHABUDDIN, D. VEDDELER, M. MÜHLENBERG, S. R. GRADSTEIN, C. LEUSCHNER, I. STEFFAN-DEWENTER, AND T. TSCHARNTKE. 2004. Biodiversity indicator groups of tropical land-use systems: Comparing plants, birds, and insects. *Ecol. Appl.* 14: 1321–1333.
- SEKERCIOGLU, C. H., S. R. LOARIE, F. OVIEDO BRENES, P. R. EHREICH, AND G. C. DAILY. 2007. Persistence of forest birds in the Costa Rican agricultural countryside. *Conserv. Biol.* 21: 482–494.
- SHANKER, K., A. HIREMATH, AND K. BAWA. 2005. Linking biodiversity conservation and livelihoods in India. *PLoS Biol.* 3: e394. doi:10.1371/journal.pbio.0030394.
- SHONO, K., S. J. DAVIES, AND C. YEN KHENG. 2006. Regeneration of native plant species in restored forests on degraded lands in Singapore. *For. Ecol. Manage.* 237: 574–582.

- SOTO-PINTO, L., G. JIMÉNEZ-FERRER, A. VARGAS GUILLÉN, B. DE JONG BERGSMAN, E. ESQUIVEL-BAZÁN. 2005. Experiencia agroforestal para la captura de carbono en comunidades indígenas de México. *Revista Forestal Iberoamerica*. Available at: <http://www.e-ambiente.com.ar/Articulos/imprimiss.asp?IDArticulo=1121>.
- SOTO-PINTO, L., I. PERFECTO, AND J. CABALLERO-NIETO. 2002. Shade over coffee: Its effects on berry borer, leaf rust and spontaneous herbs in Chiapas, Mexico. *Agroforestry Syst.* 55: 37–45.
- SOTO-PINTO, L., V. VILLALVAZO, G. JIMENEZ-FERRER, N. RAMÍREZ-MARCIAL, G. MONTOYA, AND F. SINCLAIR. 2007. The role of coffee knowledge in determining shade composition of multistrata coffee systems in Chiapas, Mexico. *Biodivers. Conserv.* 16: 419–436.
- STEFFAN-DEWENTER, I., M. KESSLER, J. BARKMANN, M. M. BOS, D. BUCHORI, S. ERASMI, H. FAUST, G. GEROLD, K. GLENK, S. R. GRADSTEIN, E. GUHARDJA, M. HARTEVELD, D. HERTELD, P. HOHN, M. KAPPAS, S. KOHLER, C. LEUSCHNER, M. MAERTENS, R. MARGGRAF, S. MIGGELKLEIAN, J. MOGEA, R. PITOPANG, M. SCHAEFER, S. SCHWARZE, S. G. SPORN, A. STEINGREBE, S. S. TJITROSOEDIRDO, S. TJITROSOEMITO, A. TWELE, R. WEBER, L. WOLTMANN, M. ZELLER, AND T. TSCHARNTKE. 2007. Tradeoffs between income, biodiversity, and ecosystem functioning during tropical rainforest conversion and agroforestry intensification. *Proc. Natl. Acad. Sci. U.S.A.* 104: 4973–4978.
- SUAZO, I., J. ALVARADO, AND M. MARTÍNEZ-RAMOS. 2008. Effects of conversion of dry tropical forest to agricultural mosaic on herpetofaunal assemblages. *Conserv. Biol.* 22: 362–374.
- SWIFT, M. J., A.-M. N. IZAC, AND M. VAN NOORDWIJK. 2004. Biodiversity and ecosystem services in agricultural landscapes—Are we asking the right questions? *Agric., Ecosyst. Environ.* 104: 113–134.
- TEWKSBURY, J. J., D. J. LEVEY, N. M. HADDAD, S. SARGENT, J. L. ORROCK, A. WELDON, B. J. DANIELSON, J. BRINKERHOFF, E. I. DAMSCHEN, AND P. TOWNSEND. 2002. Corridors affect plants, animals, and their interactions in fragmented landscapes. *PNAS* 99: 12923–12926.
- TOLEDO, V. M., B. ORTIZ-ESPEJEL, L. CORTÉS, P. MOGUEL, AND M. D. J. ORDOÑEZ. 2003. The multiple use of tropical forests by indigenous peoples in Mexico: A case of adaptive management. *Conserv. Ecol.* 7: 9. Available at: <http://www.consecol.org/vol7/iss3/art9/>.
- TSCHAKERT, P., O. T. COOMES, AND C. POTVIN. 2007. Indigenous livelihoods, slash-and-burn agriculture, and carbon stocks in Eastern Panama. *Ecol. Econ.* 60: 807–820.
- TSCHARNTKE, T., A. M. KLEIN, A. KRUESS, I. STEFFAN-DEWENTER, AND C. THIES. 2005. Landscape perspectives on agricultural intensification and biodiversity—Ecosystem service management. *Ecol. Lett.* 8: 857–874.
- TURNER, E. G., J. L. SNADDON, T. M. FAYLE, AND W. A. FOSTER. 2008. Oil palm research in context: Identifying the need for biodiversity assessment. *PLoSOne* 3: e1572.
- VANBAEL, S. A., S. M. PHILPOTT, R. GREENBERG, P. BICHIER, N. BARBER, K. A. MOONEY, AND D. GRUNER. 2008. Birds as top predators across natural and managed systems. *Ecology* 89: 928–934.
- VANDERMEER, J., AND R. CARVAJAL. 2001. Metapopulation dynamics and the quality of the matrix. *Am. Nat.* 158: 211–220.
- VANDERMEER, J., AND I. PERFECTO. 2007. The agricultural matrix and a future paradigm for conservation. *Conserv. Biol.* 21: 274–277.
- VERCHOT, L. V., M. VAN NOORDWIJK, S. KANDJI, T. TOMICH, C. ONG, A. ALBRECHT, J. MACKENSEN, C. BANTILAN, K. V. ANUPAMA, AND C. PALM. 2007. Climate change: Linking adaptation and mitigation through agroforestry. *Mitigat. Adapt. Strateg. Global Change* 12: 901–918.
- WARA, M. 2007. Is the global carbon market working? *Nature* 445: 595–596.
- WEST, P. 2006. Conservation is our government now: The politics of ecology in Papua New Guinea (New ecologies for the twenty-first century). Duke University Press, Durham, North Carolina.
- WHITMORE, T. J., AND B. L. TURNER II. 2001. Cultivated landscapes of Middle America on the eve of the conquest. Oxford University Press, Oxford, UK.
- WILLIAMS-GUILLÉN, K. C. MCCANN, J. C. MARTÍNEZ SÁNCHEZ, AND E. KOONTZ. 2006. Resource availability and habitat use by mantled howling monkeys in a Nicaraguan coffee plantation: Can agroforests serve as core habitat for a forest mammal? *Anim. Conserv.* 9: 331–338.
- WILLIAMS, S. E., AND HILBERT, D. 2006. Climate change threats to the biodiversity of tropical rainforests in Australia. *In* W. F. Laurance, and C. Peres (Eds.). *Emerging threats to tropical forests*, pp. 33–52. University of Chicago Press, Chicago, Illinois.
- WRIGHT, S. J., AND H. C. MULLER-LANDAU. 2006. The future of tropical forest species. *Biotropica* 38: 207–301.
- WUNDER, S. 2005. Payments for environmental services: Some nuts and bolts. Center for International Forestry Research.
- WWF. 2002. Forest management outside protected areas. Position paper, WWF, Gland.
- YOUNG, A. M. 1994. The chocolate tree: A natural history of cacao. Smithsonian Institution, Washington, DC.
- ZUIDEMA, P. A., AND J. A. SAYER. 2003. Tropical forests in multi-functional landscapes: The need for new approaches to conservation and research. *In* P. A. Zuidema (Ed.): *Tropical forests in multi-functional landscapes*. Seminar proceedings. Prince Bernhard Centre, Utrecht University, Utrecht, The Netherlands.