Special Report

Weekly analysis of current issues

Amazonian forests falling faster 25 July 2003

by William F. Laurance



[Caption]

One of the world's most alarming environmental controversies appears to be worsening. Rates of forest destruction in Brazilian Amazonia exceeded the previous year's rate by about 40%. In total, 6.4 million acres were felled and burned, an area nearly the size of Belgium.

The latest deforestation numbers, based on analyses of satellite imagery by Brazil's National Space Agency, clearly signal a worsening trend for Amazonian forests. From 1990 to 1994, forest loss in the Brazilian Amazon slowed to around 3.4 million acres per year, but since then rates have climbed sharply, averaging 5 million acres annually. In 2002, an area the size of ten football fields was destroyed every minute.

"The latest numbers are just plain scary," said Carlos Peres, a Brazilian ecologist at the University of East Anglia, UK. "Forest clearing was appallingly high in 1995, but a lot of people attributed that to unusually robust economic conditions that year. What we're seeing now is that forest destruction can be really bad even when the Brazilian economy is slow."

It is particularly disappointing that the areas where deforestation rose most sharply in 2002 - including the densely populated states of Mato Grosso and Pará - had shown signs of improvement following intervention by the Brazilian government to slow rampant illegal forest clearing and wildfires (see image above). Illegal burning, in particular, had been substantially reduced in 2000 and 2001 by satellite monitoring and improved enforcement.

So why is forest destruction still increasing? Amazon-watchers agree on the main culprits - rapidly expanding road networks and the dramatic growth of Printer ready version E-mail article to a friend

See also:

Surface wildfires in central
Amazonia: short-term impact
on forest structure and carbon
loss

Torbjørn Haugaasen, Jos Barlow and Carlos A. Peres Changes in forest structure were examined 10-15 months after an unprecedented understorey wildfire burnt previously undisturbed primary forest in central Brazilian Amazonia, following the severe 1997-1998 El Niño dry season.

Forest Ecology and Management, 2003, 179:1-3:321-331

Slow burn: the insidious effects of surface fires on tropical forests [Research Focus]

William F. Laurance Accidental surface fires are emerging as one of the most pervasive threats to tropical forests. Although unimpressive in appearance, these fires can have industrial soybean farming. Since 2000, the Brazilian government has been engaged in an unprecedented effort to criss-cross the Amazon Basin with new highways, waterways, railroads, and other infrastructure projects. The goal is to greatly accelerate economic development - for timber, mining, and, especially, soybean farming.

Soybeans are big business in Brazil, which looks set to overtake the US as the world's largest soybean producer. The area of land under soybeans has grown sharply, increasing from 25 million acres in 1990 to over 45 million acres today. Most soybean production is concentrated along the southern margin of the Amazon, in drier woodlands and savannas, but each year the soybean farmers are encroaching further into the rainforest.

In addition to clearing immense expanses of woodland and forest, the soybean farmers have lobbied the government for major new highways and waterways, so they can transport their crops quickly and cheaply to the Amazon River, where they are then exported internationally. "These grandiose transportation projects are opening up the Amazonian frontier and promoting even greater forest loss at the hands of soybean farmers, cattle ranchers, and slash-and-burn farmers," said Philip Fearnside from Brazil's National Institute for Amazonian Research.

Political factors have also played a role in accelerating deforestation, according to Fearnside. "Last year, Blairo Magei - the biggest soybean entrepreneur in Brazil - was elected governor of Mato Grosso State. That sent a powerful signal to soybean growers that they could go full-steam ahead," said Fearnside. "Now we're seeing severe environmental consequences."

Much of Brazil's burgeoning soybean crop is exported to Europe, where it is used as feed for cattle and pigs. These exports, totaling millions of tons annually, are drawing increasing fire from European environmentalists, who argue that they promote industrial factory farming and help to subsidize rainforest destruction.

In light of the latest deforestation figures, Brazil's environment minister, Marina Silva, said that "emergency measures" would be taken to deal with the crisis. Such measures may include real-time monitoring of deforestation, and efforts to force all government

surprisingly potent impacts on rainforest plant and... Trends in Ecology and Evolution, 2003, 18:5:209-212

Forest negotiations at the United Nations: explaining cooperation and discord
David Humphreys
The paper examines the three rounds of intergovernmental negotiations that took place in the period 1990-2000. These are the UNCED forest negotiations (1990-1992), the Intergovernmental Panel on Forests (1995-1997) and...
Forest Policy and Economics, 2001, 3:3-4:125-135

ministries - including those planning the road and infrastructure developments - to consider the environment when enacting their policies.

But many doubt whether the government's efforts to slow deforestation will succeed. "You can't have it both ways," said Brazilian ecologist Peres. "What we're seeing is government-subsidized deforestation, because the Brazilian government is paying for the transportation infrastructure. Because of soybeans it's suddenly become hugely profitable to attack the frontier. And you can't rely on restraints from the private sector to limit deforestation-you need very proactive government programs to slow exploitation. The incentives for deforestation are still huge."

This pessimistic view is echoed by Thomas Lovejoy, president of the Heinz Center for Science, Economics and the Environment, in Washington, DC. "What makes the increase especially disturbing is that there had been some concerted efforts to contain Amazonian deforestation," said Lovejoy. "It appears those were overwhelmed by expansion of soybeans and increased road access, exacerbated by a very dry year. Clearly all efforts combined are losing ground."

William F. Laurance is a staff scientist at the Smithsonian Tropical Research Institute in Balboa, Panama

Caption: Because of greater road access, destructive wildfires are becoming increasingly prevalent in the Amazonian frontier (photo by William Laurance).

+++

Send us your comments for publication.

Sign up for BioMedNet News weekly email alerts.

Printer ready version E-mail article to a friend

Previous Special Reports

What a moo says about mood

11 July 2003



A cow's moo can reveal its mood and pigs could soon be weighed using a camera. These are just two inventions predicted by agricultural engineers to improve the lot of the livestock farmer.

Getting a Randle on metabolism

23 June 2003



Forty years on, the biochemical explanation of insulin resistance proposed by Philip Randle needs to be refined, says a leading biochemist.

Opium or IPM? Rehabilitating Afghan agriculture 10 June 2003



As this year's crop of opium poppy is harvested in Afghanistan, there are suspicions that production of this lucrative crop is on the increase once more. Could integrated pest management (IPM) help out?

Body, mind and brain: 100 years of somatosensory

neuroscience

2 June 2003

This year marks the 100th anniversary of a series of experiments that had enormous influence on scientific views of consciousness and bodily sensation. Much has changed since those pioneering studies, but their legacy

lives on.

more Special Reports