

Kensley, B. 1968

*With very best wishes
Brian Kensley*

B. F. KENSLEY

DEEP SEA DECAPOD CRUSTACEA FROM
WEST OF CAPE POINT, SOUTH AFRICA

June 1968 Junie
Volume 50 Band
Part 12 Deel

INVERTEBRATE
ZOOLOGY
Crustacea



ANNALS OF THE SOUTH AFRICAN MUSEUM
ANNALE VAN DIE SUID-AFRIKAANSE MUSEUM

Cape Town Kaapstad

LIBRARY
DIVISION OF CRUSTACEA

CARDED

The ANNALS OF THE SOUTH AFRICAN MUSEUM

are issued in parts at irregular intervals as material
becomes available

Obtainable from the South African Museum, P.O. Box 61, Cape Town
(Cash with order, post free)

Die ANNALE VAN DIE SUID-AFRIKAANSE MUSEUM

word uitgegee in dele op ongereelde tye na beskikbaarheid
van stof

Verkrygbaar van die Suid-Afrikaanse Museum, Posbus 61, Kaapstad
(Kontant met bestelling, posvry)

OUT OF PRINT/UIT DRUK

1, 2(1, 3, 5, 7-8), 3(1-2, 5, t.-p.i.), 5(2, 5, 7-9),
6(1, t.-p.i.), 7(1, 3), 8, 9(1-2), 10(1-3),
11(1-2, 7, t.-p.i.), 21, 24(2), 27, 31(1-3), 38,
44(4).

Price of this part/Prys van hierdie deel

R1.75

Trustees of the South African Museum © Trustees van die Suid-Afrikaanse Museum
1968

Printed in South Africa by
The Rustica Press, Pty., Ltd.
Court Road, Wynberg, Cape

In Suid-Afrika gedruk deur
Die Rustica-pers, Edms., Bpk.
Courtweg, Wynberg, Kaap

DEEP SEA DECAPOD CRUSTACEA FROM WEST OF CAPE POINT, SOUTH AFRICA

By

B. F. KENSLEY

South African Museum, Cape Town

(With 19 figures in the text)

CONTENTS

	PAGE
Introduction	283
List of species and stations	284
Description and notes ..	286
Summary	321
Acknowledgements	321
References	322

INTRODUCTION

In 1959 the research ship *Africana II* of the Division of Sea Fisheries carried out trawls at twelve stations off the west coast of the Cape Peninsula and off Cape Point, under the supervision of Dr. F. H. Talbot, then of the South African Museum. The trawls were done with a 15' beam trawl, between depths of 1098 and 3440 metres. The exact positions of the stations may be obtained from the Annual Report for the Division of Sea Fisheries for the period 1st April, 1959, to 31st March, 1960, published by the Government Printer, Pretoria. The fish, Crustacea, Mollusca, and other invertebrates obtained are of particular interest, as very little fishing has been done at these depths in South African waters. Also included in this paper are references to material obtained with an Isaacs-Kidd mid-water trawl in 1960-1.

Approximately 480 decapod Crustacea belonging to 35 species were obtained (excluding fragments and specimens damaged beyond identification, and the hermit crabs, which were sent to Mme Dechancé of the Paris Museum). These include anomurans, palinurans, penaeideans, and carideans. Of these, twelve are new records for the South African region and three are previously undescribed species. The two species of *Nematocarcinus* appear to be the most plentiful decapods at these depths, followed by *Pontophilus occidentalis* var. *indica*. Many more carideans than penaeideans were collected.

From such a small collection it is obviously impossible to draw any conclusions regarding the distribution of the species involved, but in very general terms they can be classified into those which appear to be endemic to the region, those which are cosmopolitan in distribution, and those which have Indo-

Pacific affinities. Eight species appear to be endemic to the South African region, viz. *Neolithodes capensis*, *N. asperrimus*, *Munidopsis chacei*, *M. barnardi*, *Willemoesia bona-spei*, *Gennadas gilchristi*, *G. kempi* and *Sclerocrangon bellmarleyi*. The following species are distributed throughout the Atlantic, Indian and Pacific Oceans: *Galacantha rostrata*, *Stereomastis sculpta*, *Gennadas bouvieri*, *Sergestes atlanticus*, *S. armatus*, *Systellaspis debilis*, *Acantheephyra haekelii*, *A. brevirostris*, while the following have affinities with the Indo-Pacific fauna: *Polycheles demani*, *Plesiopenaeus nitidus*, *Haliporus villosus*, *Sergestes regalis*, *S. prehensilis*, *Acantheephyra quadrispinosa*, *A. coralina*, *Nematocarcinus longirostris*, *N. parvidentatus*, *Pontophilus occidentalis* var. *indica*.

The whole collection is now in the South African Museum and catalogued with South African Museum catalogue numbers.

LIST OF SPECIES AND STATIONS

SAM = South African Museum catalogue numbers.

* = new record

IK = Isaacs-Kidd trawl specimens

Species	SAM	Station	Specimens	Depth (metres)
ANOMURA				
<i>Neolithodes capensis</i> Stebbing	A10542	A193	1	2745
	A10447	A190	1	2269
	A10469	A193	1	2745
	A10541	A319	1	2690-2727
<i>Neolithodes asperrimus</i> Barnard	A10445	A189	2	1098
	* <i>Munida</i> sp.	A10525	A318	1
* <i>Munidopsis chacei</i> n.sp.	A10470	A193	3	2745
* <i>Munidopsis barnardi</i> n.sp.	A10465	A193	2	2745
<i>Galacantha rostrata</i> A. Milne-Edwards	A10497	A317	1	2708-3038
	A10508	A317	3	2708-3038
	A10485	A315	3	2891-2965
	A10546	A322	2	2745
	A10449	A190	1	2260
	A10518	A318	1	2525-2782
	A10547	A193	14	2745
A10519	A318	3	2525-2782	
PALINURIDA				
<i>Polycheles demani</i> Stebbing	A10520	A318	1	2525-2782
	A10501	A317	2	2708-3038
	A10460	A192	3	2708
	A10568	A322	1	2745
	A10522	A318	3	2525-2782
	A10569	A193	2	2745
	A10453	A191	2	2745
	A10570	A317	2	2708-3038
	A10487	A315	3	2891-2965
	A10533	A319	4	2690-2727
	A10475	A315	5	2891-2965
<i>Stereomastis sculpta</i> (Smith)	A10443	A189	3	1098
<i>Stereomastis nana</i> (Smith)	A10559	A189	2	1098
* <i>Willemoesia bona-spei</i> n.sp.	A10509	A317	3	2708-3038
<i>Eryoneicus spinocolatus</i> Bouvier	A10473	A193	2	2745
	A10543	A322	3	2745
	A10448	A190	1	2269

Species	SAM	Station	Specimens	Depth (metres)
PENAEIDEA				
<i>Plesiopenaeus nitidus</i> Barnard A10441	A189	1	1098
	A10442	A189	5	1098
* <i>Haliporus villosus</i> Alcock & Anderson A10461	A192	1	2708
<i>Gennadas gilchristi</i> Calman A10577	A321	2	3239-3440
<i>Gennadas kemp</i> Stebbing A10575	A321	3	3239-3440
	A10576	A190	2	2269
* <i>Gennadas bouvieri</i> Kemp A10578	A321	1	3239-3440
* <i>Sergestes atlanticus</i> A. Milne-Edwards A12529	IK6	1	183 (west of C.T.)
<i>Sergestes armatus</i> Kröyer A10528	A318	1	2525-2782
	A10532	A319	1	2690-2727
* <i>Sergestes sargassi</i> Ortman A12528	IK6	1	183 (west of C.T.)
* <i>Sergestes corniculum</i> Kröyer A10502	A317	1	2708-3038
	A10556	A321	1	3239-3440
	A10571	A321	1	3239-3440
* <i>Sergestes regalis</i> Gordon A10574	A192	2	2708
<i>Sergestes prehensilis</i> Bate A10572	A321	1	3239-3440
	A10573	A317	1	2708-3038
	A10516	A318	1	2525-2782
	A10483	A315	1	2891-2965
CARIDEA				
<i>Systellaspis debilis</i> (A. Milne-Edwards) A10444	A189	1	1098
	A10494	A316	1	3148-3257
	A10544	A322	1	2745
	A10455	A191	1	2745
	A10495	A316	1	3148-3257
<i>Hymenodora glacialis</i> (Buchholz) A10566	A192	1	2708
	A10563	A190	1	2269
	A10562	A192	1	2708
	A10513	A317	2	2708-3038
<i>Notostomus westergreni</i> Faxon A10517	A318	1	2525-2782
<i>AcanthePHYra haeckelii</i> (Von Martens) A10536	A319	1	2690-2727
	A10439	A189	1	1098
	A10553	A321	1	3239-3440
	A10452	A190	1	2269
<i>AcanthePHYra quadrispinosa</i> Kemp A10561	A321	2	3239-3440
	A10458	A192	2	2708
	A10560	A319	2	2690-2727
* <i>AcanthePHYra gracilipes</i> Chace A10565	A190	1	2269
* <i>AcanthePHYra brevis</i> Smith A10564	A192	1	2269
* <i>AcanthePHYra corallina</i> (A. Milne-Edwards) A12531	A319	1	2708
	A12532	A319	1	2690-2727
	A12533	A190	1	2690-2727
	A10523	A318	4	2269
<i>Nematocarcinus longirostris</i> Bate A12541	A316	2	2525-2782
	A12542	A317	1	3148-3257
	A12543	A193	2	2708-3038
	A12544	A317	3	2745
	A12545	A315	3	2708-3038
	A12546	A192	2	2891-2965
	A12547	A189	1	2708
	A12548	A189	10	1098
	A12549	A190	8	2269
	A10471	A193	16	2745

Species	SAM	Station	Specimens	Depth (metres)
<i>Nematocarcinus longirostris</i> Bate A12551	A191	4	2745
 A12552	A317	14	2708-3038
<i>Nematocarcinus parvidentatus</i> Bate A12553	A190	9	2269
 A12554	A316	6	3148-3257
 A12555	A192	1	2708
 A10474	A193	7	2745
 A10551	A322	4	2745
 A12556	A318	3	2525-2782
 A12557	A317	3	2708-3038
 A12558	A318	14	2525-2782
 A12559	A317	12	2708-3038
 A12560	A319	7	2690-2727
 A12561	A316	20	3148-3257
 A12562	A193	10	2745
 A12563	A315	10	2891-2965
<i>Glyphocrangon sculptus</i> (Smith) A12534	A193	1	2745
 A10459	A192	2	2708
 A10547	A322	5	2745
 A10462	A193	2	2745
 A10464	A193	1	2745
<i>Glyphocrangon sculptus</i> (Smith) A10549	A322	1	2745
 A10539	A319	4	2690-2727
 A10535	A319	4	2690-2782
 A10521	A318	7	2525-2782
 A10451	A190	Sev.	2269
 A10446	A189	1	1098
<i>Sclerocrangon bellmarleyi</i> Stebbing A12535	A318	2	2525-2782
* <i>Pontophilus occidentalis</i> var. <i>indica</i> de Man A10512	A317	2	2708-3038
 A12536	A318	7	2525-2782
 A12537	A191	2	2745
 A12538	A318	5	2525-2782
 A10457	A192	10	2708
 A10476	A315	2	2891-2965
 A10503	A317	5	2908-3038
 A12539	A322	2	2745
 A12540	A316	7	3148-3257
 A10480	A315	Many	2891-2965
 A10482	A315	6	2891-2965
 A10524	A318	1	2525-2782
 A10489	A316	1	3148-3257
 A10500	A317	1	2708-3038
 A10468	A193	Many	2745

ANOMURA

Family Lithodidae

Neolithodes capensis Stebbing, 1905

Neolithodes capensis Stebbing, 1905: 70. Barnard, 1950: 410, fig. 77 a-c.

Previous records and distribution

Off Cape Point, 800 fathoms, (1570 m).

Material

- SAM.A10542, carapace length 13 mm. St. A193, 2745 m.
 SAM.A10447, carapace length 10 mm. St. A190, 2269 m.
 SAM.A10469, carapace length 15 mm. St. A193, 2745 m.
 SAM.A10541, carapace length 14 mm. St. A319, 2690-2727 m.

Remarks

All four specimens are juveniles, yet all have the finger of the right chela equal to the upper margin of the hand. In all, the dactyls are smooth and terete; these specimens can therefore tentatively be assigned to *capensis*.

Neolithodes asperrimus Barnard, 1947

Neolithodes asperrimus Barnard, 1947: 374. 1950: 411, fig. 77 d-f.

Previous records and distribution

Off Saldanha Bay, 500 fms (980 m); off Cape Point, 550 fms (1080 m).

Material

- SAM.A10445, ♀, carapace length 55 mm. St. A189, 1098 m.
 ♀, carapace length 45 mm. St. A189, 1098 m.

Remarks

The specimens agree with the original description in having the finger of the right chela one and a half times the length of the upper margin of the hand, distally flattened denticulate dactyls and more spines than *capensis*.

Family **Galatheidae***Munida* sp.*Description*

Eyes not wider than the eyestalks. Rostrum tridentate, with a row of six small spines at its base. Well-marked cervical and branchial grooves present. Whole carapace covered with transverse setiferous ridges. Two or three spines on lateral border of carapace anterior to the cervical groove, five spines posterior to it. Posterior margin of carapace a raised crenulated ridge. Second and third abdominal terga with a smooth transverse groove.

Distribution

M. subcaeca Bouvier: (Bouvier, 1922: 46. Chace, 1942: 43). Cuba, West Indies, Madeira, Canary Islands.

M. microphthalmal M. Edw.: (Benedict, 1903: 310. Bouvier, 1922: 45. Doflein & Balss, 1926: 172. Chace, 1942: 40). Atlantic coast of Europe, West Indies, Ceylon, Polynesia, tropical west coast of America, Cuba.

Material

- SAM.A10525, ♀ ovigerous, carapace length 13 mm. St. A318, 2525-2782.

Remarks


This specimen is unfortunately rather damaged, having the rostrum and accompanying spines broken and all the walking legs missing. The antennules and the third maxillipeds are present and agree exactly with Chace's figures of *M. microphthalma* (1942). The merus of the third maxilliped has two spines and not three as in *subcaeca*. The characteristic spinal armature of the second abdominal segment of *microphthalma* is lacking in this specimen, the second abdominal segment being smooth as in *subcaeca*. This, together with the poor state of the material, makes it difficult to give the specimen specific status.

Munidopsis chacei n.sp.

Figs 1, 3 a, b

Description

Carapace one and a quarter times longer than wide. Lateral margins slightly convex, front slightly narrower than posterior part of carapace. Rostrum half length of carapace, moderately slender, laterally armed with three or four spines. Distal portion of rostrum dorsally carinate, upcurved, ventrally flattened. No antennal spines. Antero-lateral angle formed by outwardly

FIG. 1. *Munidopsis chacei* n.sp.

directed conical spine. Three other conical spines along lateral branchial margin. Gastric region fairly well defined by smooth cervical groove. At base of rostrum a pair of prominent conical spines, followed by single median spine in mid-gastric region, and pair of spines in post-gastric region. Cardiac region armed with two pairs of spines. Posterior carapace margin thickened, armed with four short spines. Whole carapace covered with scattered short hairs, surface rough, lateral branchial regions with elongate transverse rugae. Eyestalk short, fairly mobile, produced on inner side into slender spine, slightly longer than diameter of eye. Latter unpigmented, spherical. Rest of eyestalk unarmed. Basal joint of antennular peduncle armed with four prominent spines, ventral one having denticulate margin, externo-lateral and dorsal spines longest, interno-lateral spine shortest. Second and third joints of equal length, unarmed. Basal joint of antenna with prominent outwardly directed spines, second joint with four spines on distal margin, lateral ones largest. Ischium of third maxilliped triangular in cross-section, innermost angle finely denticulate. Inner margin of merus with three or four small denticles. Chelipeds shorter than ambulatory pereopods. Ischia, meri, and carpi armed with prominent spines. Dactyl of chela equal to palm in length. Finger and thumb denticulate, teeth becoming obsolete proximally. Ischia, meri and carpi of ambulatory pereopods also armed with rows of prominent spines. Dactyls slender, two-thirds length of propodus, ventrally armed with short spines. Abdomen equal in width to carapace, dorsally rounded, unarmed. Second and third segments with smooth dorsal groove formed by two raised transverse ridges. Pleurae all ventrally rounded, that of second segment widest. Sixth segment with two prominent lateral lobes. Exo- and endopods of uropods equal in length, fringed with setae. Endopods with median ridge just off-centre. Telson slightly wider than long, distally broadly bilobed. Eggs 2 mm in diameter.

Material

SAM.A10470, ♀ ovigerous, carapace length 36 mm, overall length (including rostrum) 100 mm. (Holotype).

♀, carapace length 25 mm, overall length 62.5 mm.

♂, carapace length 25.9 mm, overall length 72 mm. St. A193, 2745 m.

Remarks


The present species most closely resembles *M. bairdii* (Smith, 1884) but differs in the following respects: there are three pairs of spines in the gastric region of *bairdii*, whereas the present species has an anterior and a posterior pair and a single median spine. In the cardiac region, *bairdii* has a pair of median spines followed by a single median spine, while in the present species there are two pairs of spines. The posterior carapace margin in *bairdii* is armed with ten spines, while there are only four in the present species. In *bairdii*, the dactyls are about three-quarters the propodus length (taken from Benedict's 1903 paper) while in this the dactyls are just over half the propodus length.

Munidopsis barnardi n.sp.

Figs 2, 3 c, d

Description

Carapace slightly longer than broad. Lateral margins very slightly convex, more or less parallel. Frontal margin between hepatic spines not narrower than rest of carapace. Rostrum triangular, about half length of carapace, curving distally upward, dorsally carinate, with minute tubercles. Base broad. Small broadly triangular antennal spine, followed by outwardly directed conical spine in hepatic region. Antero-lateral angle formed by prominent conical spine, followed by three or four smaller spines to weakly demarcated mid-branchial groove. Latter followed by one large spine, and several flattened rugae. Gastric region well defined by cervical groove. Two well-developed spines at base of rostrum. Posterior to these, transverse row of three spines. Posterior gastric region with pair of spines, and several transverse flattened setiferous rugae. Rest of carapace especially posterior portion covered with rugae. Posterior margin separated by narrow smooth groove. Posterior margin a double raised ridge, minutely crenulated. Eystalks short, only slightly movable, eyes unpigmented, almost embedded in stalk. Inner angle of latter produced into slender spine, longer than diameter of eye. Slight tubercle just below eye, exterior to eystalk spine. First antennular peduncle joint armed with two long spines on

FIG. 2. *Munidopsis barnardi* n.sp.

outer side, inner side with short spooned denticulate process. Merus of second maxilliped smooth. Chelipeds slightly longer than carapace (including rostrum). Merus equal to chela in length, with four prominent distal spines. Three pairs of ambulatory pereiopods slightly longer than chelipeds; propodi, carpi, and meri with tubercles along angles. Dactyls not very stout, longer than carpi, spinous on lower edge, spines becoming obsolete proximally. Abdomen equal in width to carapace, dorsally rounded and unarmed. Second and third segments with two




FIG. 3.

a. *Munidopsis chacei* n.sp. Lateral view of carapace. b. *Munidopsis chacei* n.sp. Ventral view of left antennule. c. *Munidopsis barnardi* n.sp. Lateral view of carapace. d. *Munidopsis barnardi* n.sp. Ventral view of left antennule.

raised dorsal ridges, separated by smooth transverse groove. Fourth and fifth segments merely with smooth transverse groove. Sixth segment posteriorly trilobed, median lobe wider than lateral lobes, latter longer than former. Pleurae all ventrally rounded, second broadest. Anterior three pleurae slightly tuberculate, with median curved ridge. Exo- and endopods of uropods sub-circular, fringed with setae, each having slight raised ridge. Telson equal in length to exopod of uropod.

Material

		♂ Carapace length	♀ Carapace length	♂ Overall length	♀ Overall length	Station	Depth (m)
SAM.A10497	..	15.5 mm		35 mm (including rostrum)		A317	2708-3038
SAM.A10465	..	14.0 mm		34.0 mm		A193	2745
SAM.A10508	..	14.5 mm		36.4 mm		A317	2708-3038
		17.0 mm		41.2 mm			
SAM.A10485	..	17.0 mm		42.0 mm		A315	2891-2965
		17.5 mm		43.0 mm			
SAM.A 12636 (Holotype)			15.5 mm		25.5 mm		
			15.0 mm		41.0 mm		

Remarks

This species is very similar to *M. crassa* Smith, 1884, but differs in the following respects: The rostrum, which in *crassa* is very nearly horizontal, is in the present species sharply upcurved; the frontal margin appears to be proportionally wider than in *crassa*, the telson differs in shape (see Chace in correspondence); the infero-mesial edge of the merus of the second maxilliped in *crassa* is armed with three conical spines, which are lacking in the present species; *crassa* also lacks the two extra rows of spines in the gastric region, found in the present species. The holotype female of *crassa* has an overall length of 125 mm, whereas the largest female in the present series is 41 mm long, several of the males also being of comparable size. There is thus a considerable difference in size between the two species.

Galacantha rostrata Milne-Edwards, 1880

Galacantha rostrata M. Edw. 1880: 52. Benedict, 1903: 304. Barnard, 1950: 494, fig. 92 e, f.

Previous records and distribution

Off Cape Point, Atlantic and Pacific coasts of North America, East Indies, Bay of Bengal, Arabian Sea.

Material

Twenty-one specimens of this species were taken from stations A190, A193, A318, A322, in depths varying from 2269-2782 m.

PALINURAFamily **Eryonidae***Polycheles demani* Stebbing, 1917

(?) *Polycheles beaumontii* (Alcock), Stebbing, 1908: 25. Stebbing, 1910: 377.
Polycheles demani Stebbing, 1917: 28. Barnard, 1950: 570, fig. 105 a-c.

Previous records and distribution

Off Cape Point, 500-1400 fms (980-2760 m).

<i>Material</i>					♀ Overall length	♂ Overall length	Station	Depth (m)
SAM.A10453	50 mm		A191	2745
SAM.A10460	41 mm		A192	2708
SAM.A10569	78 mm		A193	2745
SAM.A10487	51 mm		A315	2891-2965
SAM.A10475	41 mm	82 mm	A315	2745
SAM.A10570	43 mm		A317	2708-3038
SAM.A10501	36 mm		A317	2708-3038
SAM.A10522	64 mm		A318	2525-2782
SAM.A10520	62 mm		A318	2525-2782
SAM.A10533	53 mm		A319	2690-2727
SAM.A10568	65 mm		A322	2745
					54 mm			
					41 mm			
					39 mm			
					71 mm			
					67 mm			
					52 mm			
					41 mm			
					100 mm			
					38 mm			
					94 mm			
					101 mm			
					99 mm			
					85 mm			
					75 mm			

Stereomastis sculpta (Smith, 1882)*Pentacheles sculpta* Smith, 1882: 23.*Polycheles sculpta*: Stebbing, 1910: 377.*Stereomastis sculpta*: Barnard, 1950: 572, fig. 105 d.*Previous records and distribution*

Off Cape Point, 600 fms (1180 m), off Durban, 440 fms (865 m), Gibraltar, Canary Islands, East African coast, Mediterranean, East Indies, west coast of North America.

Material

SAM.A10443, ♀♀, overall length 92 mm, 87 mm, ♂, 69 mm. St. A189, 1098 fms.

Stereomastis nana (Smith, 1884)*Pentacheles nanus* Smith, 1884: 359.*Polycheles nana*: Stebbing, 1910: 377.*Stereomastis nana*: Barnard, 1950: 573, fig. 105 e.*Previous records*

Off Cape Point, 800 fms (1570 m), 1200 fms (2360 m), east and west coasts of North America, Gulf of Panama.

Material

SAM.A10559, ♀♀, overall length 55 mm, 47 mm. St.A189, 1098 fms.

Remarks

This species is very similar to the preceding one, the main differences being in the median carapace spines (2, 1, 2, 1-2, 2, 2 in *sculpta*, 2, 1, 1, 2, 1-2, 2, 2, in *nana*), the spinous structure of the sixth abdominal keel in *nana* and the slightly more spinous condition of the posterior portion of the carapace in *nana*. With reference to this, it is interesting to note that in two of the three specimens of *sculpta* in this collection there are traces of spines on the raised keel of abdominal segment six. The possibility exists that *nana* is a juvenile form of *sculpta* and that some of the spines are lost with development.

Willemoesia bona-spei n.sp.

Figs 4, 5

Description

Carapace one and a half times longer than broad, lateral margins of posterior carapace parallel, anterior margins converging. Antero-lateral angle formed by large spine (largest on carapace). Frontal margin emarginate. Two prominent spines above antennules, between which a prominent median spine (latter not marginal but just posterior to margin) projecting almost vertically from carapace. Medio-dorsal carina spine formula variable, three to five single spines, one pair and final single spine anterior to cervical groove. (1.1.1.1.2.1.C.2, . . .). Prominent posterior median carina in some specimens with pair of small spines just posterior to cervical groove. No regular arrangement posterior to this. Lateral branchial spine formula variable (6-9, 5-8, 18-30). Whole carapace covered with close-set tiny spinules. Latter scattered over orbito-gastric and post-median ridges. Supra-branchial ridge with about eight tiny spines. Anterior carapace margin and antero-lateral margins fringed with short hairs. Basal joints of antennules produced mesially into two wing-like processes, furnished with eight to ten small spines. Second and third antennular joints together equal in length to 1st joint. Antennal peduncle slightly longer than antennular peduncle. Mandible with thirteen to fourteen heavily chitinized teeth. Chelipeds almost three times longer than carapace. Finger of chela with spine at right angles to it, situated in distal half of finger. Latter and thumb equal in length to or slightly longer than palm. Latter with spines along both edges, and along outer edge of finger. Carpus two-thirds length of chela with spines along outer edge. Merus equal in length to chela, with inner margin spined.

Abdomen (excluding telson) three quarters carapace length. First five segments dorsally carinate, first four each with single small forwardly-directed tooth. Sixth segment dorsally smooth. Postero-lateral angle of sixth segment with small lobe. Telson triangular, apically acute, equal in length to posterior three and a half segments. Exopod of uropod subcircular, endopod elongate-




FIG. 4.
Willemoesia bona-spei n.sp.

oval, both fringed with setae. Pleuron of second segment almost circular, twice as broad as that of third. Pleurae of fifth and sixth segments ventrally pointed, anterior three ventrally rounded.

Material

SAM.A.10473	♀ Overall length	♂ Overall length	Station	Depth (m)	Spine formulae	
					Median	Lateral
..		108 mm	A193	2745	11121/2..	9/7/27
		74 mm			111121/...	9/6/27




FIG. 5. *Willemoesia bona-spei* n.sp.
 a. First pleopod of male. b. Mandible.
 c. Apex of maxilliped 2. d. Maxilla.

		♀ Overall length	♂ Overall length	Station	Depth (m)	Spine Median	Spine Lateral
SAM.A10509	..	87 mm		A317	2708-3038	111121/2..	8/7/30
			113 mm			11121/2...	7/5/27
			108 mm			11121/...	9/7/21
SAM.A10543 (Holotype)	..	109 mm (ovig.)		A322	2745	11121/....	6/8/30
SAM.A12637	..		107 mm			11121/2...	9/7/18
			94 mm			111121/2.	9/5/25

Remarks

From an assessment of the spine formulae of the present species, it would seem that it is closely related to several described species, such as *leptodactyla* (Willemoes-Suhm), *pacifica* Sund, or *challengeri* Sund. Specific delimitation in this genus is very difficult, particularly as the number of specimens available is very low. Both *leptodactyla* and *challengeri* have a thick 'fur', unlike the present species, the carapace of which is covered with tiny spines. The present species resembles Bate's 1888 plate 19 C (a specimen captured off the coast of Chile and named *pacificus* by Sund in 1920) both in the shape of the frontal margin, and in the presence of teeth on the supra-branchial ridge. This species differs from *pacificus* and indeed from all the described species in that the posterior portion of the median dorsal carina does not have any regular spine formula, but simply has a scattering of the tiny spines as found on the rest of the carapace.

Sund's *pacificus* has the orbital sinus more angular than the present species and the most anterior median spine is not marginal. In lateral view, the present species is only feebly arched, while in *pacificus* 'the carapace, when seen in profile, is strongly arched' (Sund, 1920). The present specimens are thus described as a new species, but every likelihood exists that with more material becoming available, it will be found to be synonymous with an already described species.

Eryoneicus spinoculatus Bouvier, 1905

Fig. 6

Eryoneicus spinoculatus Bouvier, 1905: 480. Bernard, 1953: 34. Belloc & Lorillou, 1961: 10.

Description

Carapace longer than wide, unarmed except for carinal spines. Rostrum a pair of small spines. Median dorsal carina spine formula—1, 1, 2, '1, 1, C, 2, 2, '1, 2, ('1 indicates a blunt spine). Lateral carina spine formula 6, 3, 7. Posterior carapace ridge, between median and lateral carinae, with ten or eleven spines. Orbito-cervical line with three small spines. Frontal margin rounded. First abdominal segment with two medio-dorsal spines, segments two to five with three medio-dorsal spines. Sixth with single posterior spine, anterior portion smooth. Single lateral spine on each abdominal tergum. Single spine on pleurae two to five. Pleuron of second segment twice as wide as that of third. Ventral margins of pleurae denticulate. Telson with two median spines, seven or eight lateral spines. First pleopods relatively undeveloped.

Material

SAM.A10448, one specimen, carapace length 24 mm, width 20 mm, overall length 44 mm. St. A190, 2269 m.




FIG. 6.
Eryoneicus spinoculatus Bouvier.

Remarks

From the median dorsal spine formula, it would seem that this specimen belongs to the *spinoculatus* group (Bernard, 1953). Using his key to the group, one arrives at *E. spinoculatus* var. *hibernicus* (Selbie), distinguished from *spinoculatus* s.s. by the lack of anterior spines on the sixth abdominal tergum. This variety has been recorded from 2100 metres off the coast of Ireland, from the north Atlantic and from the waters of Greenland.

PENAEIDEA

Family **Penaeidae**

Plesiopenaeus nitidus Barnard, 1947

Plesiopenaeus nitidus Barnard, 1947: 383. 1950: 622, fig. 116. Grindley & Penrith, 1965: 280.

Previous records

Off Cape Point, 475-630 fms (930-1240 m), south-west Indian Ocean.

Material

				♀	♂	Station	Depth (m)
				Carapace length	Carapace length		
SAM.A10441	17.2 mm		A189	1098
SAM.A10442	27.2 mm		A189	1098
				31.0 mm			
				31.6 mm			
					27.5 mm		
					22.7 mm		

Haliporus villosus Alcock & Anderson, 1874

Figs 7, 8

Haliporus villosus Alcock & Anderson, 1894: 146. Alcock, 1900: pl. 26, fig. 1.

Hymenopenaeus villosus: Burkenroad, 1936: 105.

Description

Integument soft and glabrous; carapace torn in places, rostrum missing. Carapace dorsally carinate, strongly arched, with distinct notch one third of carapace length from base of rostrum, formed by cervical groove. Mid-dorsal carina flattened for three millimetres just posterior to cervical notch. Eight dorsal teeth between cervical notch and rostral base. Strong post-antennal carinate spine present. Post-antennal carina meets obliquely descending cervical




FIG. 7. *Haliporus villosus* Alcock & Anderson.




FIG. 8. *Haliporus villosus* Alcock & Anderson.
 a. Telson and left uropod. b. Antennal scale. c. Mandible. d. Maxilla 1. e. Maxilla 2.
 f. Maxilliped 1. g. Eyestalk.

groove at hepatic spine, then runs posteriorly upward to posterior midline of carapace. A carina, anteriorly with two spines, stretches from the lateral midpoint of cervical groove, to lateral midpoint of post-antennal carina. No antennal spine but a blunt projection on carapace margin. Branchiostegal spine minute. Pterygostomial spine much larger. Pterygostomial carina stretches to below cervical groove. Another keel, ventral to pterygostomial carina, extends length of carapace to posterior margin. Several smaller branching keels in posterior region of carapace, meeting in posterior midline. Eyes slightly wider than eyestalks, reaching to end of first antennular peduncle joint. Antennular peduncle two and a half times length of eyestalk, basal joint with small spine on outer distal angle. Tiny tubercle on inner margin of eyestalk. Second joint of antennal peduncle with strong outwardly flared spine, just anterior to pterygostomial spine. Antennal scale broad, flexible, almost foliaceous, fringed with setae, tiny spine on outer margin. All mouthparts fringed with setae. Exopod of maxilliped 2 twice length of exopod of maxilliped 3. Maxillipeds 2 and 3, and pereiopods 1-4 with leaf-like epipods. Epipod of maxilliped 2 also has podobranch, maxilliped 3 with rudimentary podobranch. Epipods of pereiopods without podobranchs. Maxilliped 3 and pereiopod 4 reaching equally far forward, former with slender dactyl, propodus and carpus of almost equal length. Pereiopods long, fairly stout, 1 to 3 chelate. Fifth pereiopod longest, ending in very slender dactyl. All abdominal segments dorsally carinate, fourth, fifth, and sixth ending in slight raised spines. Each abdominal segment with two lateral ridges, latter meeting on posterior margin of each segment. Pleopods large, with long setiferous endo- and exopods. Telson apically acute, armed with four small lateral spines. Uropods almost foliaceous, setiferous, endopod slightly shorter than exopod. Latter with marginal tooth some distance from apex.

Previous records

Presumably from the Indian region (the paper containing the original description is not available in South Africa).

Material

SAM.A10461, ♀, carapace length 58.5 mm, overall length 168 mm. St. A192, 2798 m.

Remarks

This is a new record for the South African region. The species has apparently not been collected since it was recorded by the *Investigator* in 1894. The only difference between the present specimen and the holotype is that in the latter the integument is covered with short fine hairs, while the former has a glabrous integument.

Gennadas gilchristi Calman, 1925

Gennadas gilchristi Calman, 1925: 6. Barnard, 1950: 633, fig. 118 g, h.
Amalopenaeus elegans (non Smith) Stebbing, 1917: 31.

Previous record

Off Cape Point, 360–1014 fms (700–1990 m).

Material

SAM.A10577, ♂, carapace length 8·3 mm; ♀, carapace length 11·2 mm.
St. A.321, 3239–3440 m.

Gennadas kempi Stebbing, 1914

Gennadas kempi Stebbing, 1914: 283. Calman, 1925: 4. Balss, 1927: 260. Barnard, 1950: 630, fig. 118 a–d.

Previous records

Off Cape Point, 1000 fms (1970 m), south Atlantic.

Material

SAM.A10575, ♂, carapace length 8·5 mm. St. A321, 3239–3440 m.

SAM.A10576, ♂♂, carapace length 9 mm, 9 mm. St. A190, 1240 fms.

Gennadas bouvieri Kemp, 1909

Gennadas bouvieri Kemp, 1909: 727. Burkenroad, 1936: 80. Tirmizi, 1960: 360.

Amalopenaeus alcocki (non Kemp) Balss, 1935: 266.

Amalopenaeus bouvieri: Balss, 1925: 267.

Description

Rostrum of typical *Gennadas* type. Carapace carinate throughout its length. Thelycum of female with eighth thoracic sternite bearing a pair of slightly elongate antero-lateral projections, with setose tips.

Previous records and distribution

Arabian Sea, Zanzibar, eastern Pacific, Bahamas, Bermuda, Caribbean, south Atlantic (Ascension).

Material

SAM.A10578, ♀, carapace length 8·3 mm. St. A321, 3239–3440 m.

Remarks

On lifting the thelycal plate between the bases of the third and fourth pereopods, a pair of brown spermatophores could be seen. Each spermatophore has a rounded swollen base and a curved neck ending in a hook which is inserted into the spermatheca and makes removal difficult.

Apart from Balss's record of a male from the Ascension Island region, this species has not been found in the south Atlantic. This is thus a new record for the South African region.

Family **Sergestidae**Genus **SERGESTES** Milne Edwards

Up to and including 1950, the following species of *Sergestes* had been recorded from South African waters: *articus* Kröyer, *potens* Burkenroad (= *phorcus*

Faxon), *prehensilis* Bate (= *gloriosus* Stebbing), *armatus* Kröyer, *splendens* Sund.

The 1960 midwater collection of Decapoda yielded the following species: *potens* Burkenroad, *prehensilis* Bate, *regalis* Gordon, *armatus* Kröyer, *atlanticus* Milne Edwards, *sargassi* Ortmann, *corniculum* Kröyer, *splendens* Sund.

The present deep-water collection has yielded *regalis* Gordon, *armatus* Kröyer, *prehensilis* Bate, *corniculum* Kröyer.

Burkenroad (1937), in describing the Sergestidae of the Templeton Crocker Expedition to California, noted that only some species of *Sergestes* had specialized organs in the gastrohepatic region, first noted by Pesta in 1918. It is thought that these 'organs of Pesta' have a luminescent function. Burkenroad also noted that those species of *Sergestes* which lacked organs of Pesta invariably possessed dermal photophores, and went further to suggest that possibly the genus could be split into two natural groups. This has been done by Yaldwyn (1957). Those species possessing organs of Pesta fall into the subgenus *Sergestes*, while those possessing photophores fall into the subgenus *Sergia*. This classification has been followed in this work. In life, the organs of Pesta have a conspicuous colouring of red and blue, but in preserved material these organs, which are internal and difficult to see without damage to the specimen, become opaque-white.

Further division of the subgenus *Sergia* can be based on the type of photophore, whether it posses a translucent lens-like structure or whether merely an opaque-white dermal spot.

Sergestes (Sergestes) atlanticus Milne Edwards, 1830

Figs 9 d, 10 c, 11 d

Sergestes atlanticus Milne Edwards, 1830: 346. Bate, 1888: 389. Hansen, 1896: 951. 1903: 58. 1922: 41.

Sergestes (Sergestes) atlanticus: Yaldwyn, 1957: 8.

Description

Rostrum an apically acute spine. Prominent supra-orbital spine; well-developed hepatic spine. Feeble cervical groove midway along carapace. Slight gastrohepatic groove and suprabranchial ridge. Eyes wider than the eyestalk. Eyestalk half length of first antennular segment. Maxilliped 3 equals pereopod 3 in length. Two distal segments of pereopod 5 setose on only one margin. No dermal photophores.

Petasma of male short and stumpy. Processus uncifer not apically hooked. Processus ventralis apically acute, broadly triangular in shape. The lobus terminalis a short lobe below the oblong-oval lobus inermis. Lobus connectens and lobus armatus short; apically blunt lobes.

Previous records and distribution

North Atlantic, off South American coast, off Japan, Fiji, south of Australia.

Material

SAM.A12529, ♂ carapace length 27.9 mm. IK. St. 6, west of Cape Town; caught 200 metres in a depth of water 1600 metres.

Remarks

This appears to be the first record from South African waters of this species.

Sergestes (Sergestes) armatus Kröyer, 1855

Sergestes armatus: Hansen, 1922: 174. Calman, 1925: 26. Barnard, 1950: 643, fig. 120 m-p.
Sergestes (Sergestes) armatus: Yaldwyn, 1957: 8.

Material

SAM.A10528, ♀ carapace length \pm 17 mm. St. A318, 2525-2782 m.

SAM.A10532, ♀ carapace length 11.6 mm. St. A.319, 2690-2727 m.

Remarks

The most useful diagnostic feature of this species is the very obvious and well-developed third maxilliped, which is longer and stouter than the longest pereopod. This feature is also found in *Sergestes sargassi* to some extent, but that species can easily be distinguished by the two distal segments of the fifth pereopod which have setae on only one margin in *armatus*.

Previous records and distribution

Table Bay, 300 fms (590 m), Cape Point, 310 fms (600 m), north and south Atlantic, off Agulhas, off Natal coast, south-west Indian Ocean, South Australia, Mediterranean.

Sergestes (Sergestes) sargassi Ortmann, 1893

Figs 9 c, 10 d, 11 c

Sergestes sargassi Ortmann, 1893: 34. Hansen, 1922: 148.
Sergestes (Sergestes) sargassi: Yaldwyn, 1957: 8.

Description

Rostrum short, anterior margin almost vertical, topped by a short, sharp, horizontal spine. Carapace with well-defined cervical groove, prominent supra-branchial ridge, distinct gastrohepatic groove, minute supra-orbital and hepatic spines. Eyestalk twice as long as eye, latter slightly wider than the stalk. Eyestalk reaching half-way along first joint of antennular peduncle. Maxilliped 3 stouter and longer than the pereopods, two distal segments with internal margins having comb-like rows of bristles. Pereopod 3 longest, reaching slightly beyond the antennular peduncle. Pereopod 4 flattened with long setae on distal segments. Pereopod 5 about half length of fourth, two distal segments setose on both margins. No dermal photophores. Petasma lobes generally elongate, possessing lobus inermis (a small lobe on the lobus terminalis), stout processus ventralis, and smaller, thinner lobus armatus. Inner surface of processus ventralis armed with five stellate spines, followed by a marginal row of hooks, ending in two slightly larger apical hooks.




FIG. 9.

a. *Sergestes corniculum* Kröyer. Carapace. b. *Sergestes regalis* Gordon. Carapace.
 c. *Sergestes sargassi* Ortmann. Carapace. d. *Sergestes atlanticus* Milne Edwards. Carapace.




FIG. 10.

a. *Sergestes regalis* Gordon. Second ramus of antennule of male. b. *Sergestes corniculum* Kröyer. Second ramus of antennule of male. c. *Sergestes atlanticus* Milne Edwards. Second ramus of antennule of male. d. *Sergestes sargassi* Ortmann. Second ramus of antennule of male.




FIG. 11.

a. *Sergestes regalis* Gordon. Petasma. b. *Sergestes corniculum* Kröyer. Petasma. c. *Sergestes sargassi* Ortman. Petasma. d. *Sergestes atlanticus* Milne Edwards. Petasma.

Distribution

Off Madeira, Azores, Sargassum Sea.

Material

SAM.A12528, ♂, carapace length 8 mm, overall length 27 mm, IK St. 6, west of Cape Town caught at 200 metres in a depth of water of 1600 metres.

Remarks

It is remarkable that, like *Sergestes atlanticus*, only one specimen of this species was taken from all the hauls containing sergestids. This is the first record of this species from the South African region.

Sergestes (Sergestes) corniculum Kröyer, 1855

Figs 9 a, 10 b, 11 b

Sergestes corniculum Kröyer, 1855: 22. Pesta, 1918. Burkenroad, 1937: 316. Hansen, 1922: 126.
Sergestes (Sergestes) corniculum: Yaldwyn, 1957: 7.

Description

Rostrum with single acute apical spine, flanked by well-developed supra-orbital ridges, very seldom having a minute spine. Well-defined cervical groove about half-way along carapace. At base of cervical groove, a prominent ridge runs anteriorly to eye. Two well-defined ridges in the branchial region. A minute hepatic spine is sometimes present, more usually a blunt knob-like protuberance. Eye prominent, black, wider than the eye-stalk. Tiny tubercle on inner side of stalk, just posterior to eye. Eyestalk reaching half-way along first antennular peduncle segment. All pereopods laterally compressed. Second and third pereopods equally long and slender. Fifth pair about half the length of fourth. Two distal segments of pereopod 5 setose on both margins. No dermal photophores. Petasma of male with processus ventralis distally expanded, having eight to ten papilla-like protuberances. Lobus armatus stout, curved. Lobus connectens small, lobus terminalis apically blunt; lobus inermis longest, reaching furthest distally, apically acute, ending in two or three spines.

Previous records and distribution

Mediterranean, north Atlantic, off Durban, Agulhas, south-west Indian Ocean, off Cape Point.

Material

SAM.A10502, ♂, carapace length 15 mm. St. A317, 2708-3038 m.

SAM.A10556, ♀, carapace length 13 mm. St. A321, 3239-3440 m.

SAM.A10571, ♀, carapace length 14 mm. St. A321, 3239-3440 m.

Remarks

This appears to be the first record of this species from South African waters. It appears to be plentiful, occurring at most of the Isaacs-Kidd midwater stations in addition to the 1959 collection, in depths ranging from 200 metres to about 3700 metres.

Sergestes (Sergia) regalis Gordon, 1939

Figs 9 b, 10 a, 11 a

Sergestes regalis Gordon, 1939: 498.*Sergestes (Sergia) regalis*: Yaldwyn, 1957: 9.*Description*

Rostrum apically bifid. Carapace with well-defined supra-branchial ridge, with a less well-defined ridge ventral to the former. Cervical groove in posterior third of the carapace. Slight groove in antero-lateral third of carapace. Eye prominent, eyestalk two-thirds the length of first antennular peduncle segment. Pereiopods 1-3 slender, third longest, second and third chelate, with stiff red bristles. Pereiopods 4 and 5 shorter, flattened, fringed with long setae. Sixth abdominal segment ends in a spinule, telson medially grooved. Photophores of the 'opaque spot' type, difficult to detect. A variable number of photophores on the merus of the third pereiopod, usually about ten. Petasma of male with lobus connectens longer than lobus terminalis; lobus armatus and lobus connectens ending in retracted hooks, lobus armatus longer than processus ventralis.

Previous records and distribution

Mid south Atlantic, off Durban, south-west Indian Ocean.

Material

SAM.A10574, ♀, carapace length 26 mm, abdominal length 44 mm. St. A192, 2708 m.

Remarks

Only one large female was taken in this collection, but several more specimens were obtained from the Isaacs-Kidd midwater collection. This is the first record of the species from the South African region.

Sergestes (Sergia) prehensilis Bate, 1888*Sergestes prehensilis* Bate, 1888: 193. Gordon, 1935: 314.*Sergestes (Sergia) prehensilis*: Yaldwyn, 1957: 9.*Sergestes gloriosus* Stebbing, 1905: 84. 1910: 38. Barnard, 1950: 642, fig. 120 h-j.*Description*

Rostrum apically acute, sometimes a small denticle on the upper margin; indistinct supra-orbital ridge present, no supra-orbital spine. No hepatic spine but a blunt knob-like protuberance. Dermal photophores of the lens-like type.

Previous records and distribution

Off Sandy Point, 800 fms (1570 m), off Durban, 260 fms (510 m), East London region, Agulhas, south-west Indian Ocean, Japan.

Material

	♂		♀		Station	Depth (m)
	Carapace length		Carapace length			
SAM.A10483	A315	2891-2965
SAM.A10513	A317	2708-3038
SAM.A10516	A318	2525-2782
SAM.A10553	A321	3239-3440

Remarks

This species appears to be the most plentiful sergestid in South African waters. It was obtained at all the Isaacs-Kidd midwater stations in large quantities. It does not appear to be very plentiful in the hauls from the greater depths of the 1959 collection.

CARIDEA

Family **Oplophoridae***Systellaspis debilis* (Milne Edwards, 1881)

Systellaspis debilis: Chace, 1940: 181. Barnard, 1950: 663, fig. 124 a. Grindley & Penrith, 1965: 281.

Previous records and distribution

Off Cape Point, 1500 fms (2950 m), off Natal, north and west Atlantic, Indo-Pacific.

Material

				♀	♂	Station	Depth (m)
				Carapace length	Carapace length		
SAM.A10444	10.2 mm	A189	1098
SAM.A10455	12 mm		A191	2745
SAM.A10494	11.0 mm	A316	3148-3257
SAM.A10495	13 mm		A316	3148-3257
				(ovig.)			
SAM.A10544	12 mm		A322	2745

Remarks

The carapace length of ovigerous females varies from 12 mm to 14 mm in the Isaacs-Kidd material as well as the present collection. Egg size is 3-3.5 mm.

Hymenodora glacialis (Buchholz, 1874)

Hymenodora glacialis: Kemp, 1910: 72. Calman, 1925: 15. Barnard, 1950: 665, fig. 124 b.

Description

Number of rostral spines varying from three to six. Eyes very feebly pigmented, narrower than eyestalk. Telson broken in all the specimens.

Previous records and distribution

Off Cape Point, 1500 fms (2950 m), north Atlantic, west coast of Ireland, 1150 fms (2260 m), north and east Pacific.

Material

				♀	♂	Station	Depth (m)
				Carapace length	Carapace length		
SAM.A10563	11.0 mm		A190	2269
SAM.A10562		13.5 mm	A190	2708
SAM.A10566		15.0 mm	A192	2708
SAM.A10513	9.6 mm		A317	2708-3038
				12.0 mm			

Notostomus westergreni Faxon, 1893

Notostomus westergreni Faxon, 1893: 208. 1895: 171. Stebbing, 1905: 110. 1910: 395. Chace, 1940: 171.

Notostomus auriculatus Kemp (in MS) Barnard, 1950: 670, fig. 124 b, i.

Previous records and distribution

Off Cape Point, 800 fms (1570 m), off coast of Ecuador, off Bermuda, 900 fms (1770 m) off Keeling Islands, Indian Ocean.

Material

SAM.A10517, ♀, carapace length 38.6 mm, overall length (excluding rostrum), 97.5 mm. St. A318, 2525-2782 m (rostrum missing).

SAM.A12563, ♂, carapace length 42 mm, overall length (excluding rostrum), 90 mm, IK. St. 14, caught at 500 metres in a depth of 2000 metres.

Remarks

Stebbing (1905) noted that the specimen from Cape Point was perhaps a species other than *westergreni*, as it had a strong posterior tooth on the sixth abdominal segment. According to Barnard (1950), Kemp saw this specimen and named it *auriculatus* in MS.; this MS. could not be traced. Comparison of the present specimens and Stebbing's specimen with the original description of *westergreni* make it seem probable that all three specimens belong to this species. Although Faxon does not mention a spine on the sixth abdominal segment, the colour plate illustration (pl. F) shows one. Chace, 1940, also notes that the specimen of *westergreni* in the U.S. National Museum possesses a tooth on the sixth segment. There are minor variations in all the specimens. The lateral carina of the rostrum curves downward in Stebbing's specimen, while the above specimens are as in Faxon's plate, ending horizontally in the gastric region. The asymmetry of the antennal scales in Stebbing's specimen is almost certainly abnormal. The present specimens agree with Faxon's sketch; the outer apical spine extends some way past the apex of the scale and there is no asymmetry. The length of the rostral spines also seems to vary, but as these are very brittle, this variation may have no specific importance. Until further specimens are available for comparison the extent of variation of the species will remain unknown, and the name *westergreni* should be retained.

Acanthephyra haeckelii (Von Martens, 1868)

Ephyra haeckelii Von Martens, 1868: 54.

Acanthephyra haeckelii: Kemp, 1939: 575. Chace, 1940: 140. Barnard, 1950: 668. Grindley & Penrith, 1965: 280.

Previous records and distribution

Off Cape Point, 900 fms (1770 m), south-west Indian Ocean, north Atlantic, south Atlantic, Mediterranean, south Pacific.

Material				♀	♂	Station	Depth (m)	Telson spines
				Carapace length	Carapace length			
SAM.A10536				20.0 mm		A319	2690-2727	10
SAM.A10553					15.0 mm	A321	3239-3440	10
SAM.A10452				16.5 mm		A190	2269	9
SAM.A10439				11.5 mm		A189	1098	9

AcanthePHYra quadrispinosa Kemp, 1939

AcanthePHYra quadrispinosa Kemp, 1939: 576. Barnard, 1950: 668, fig. 124 g. Grindley & Penrith, 1965: 280.

AcanthePHYra batei (non Faxon) Stebbing, 1905: 107.

Previous records and distribution

Off Cape Point, 700-1800 fms (1380-3540 m), off Natal, 820 fms (1610 m), south-east of Agulhas, south-west Indian Ocean, Indo-Pacific.

Material				♀	♂	Station	Depth (m)
				Carapace length	Carapace length		
SAM.A10561				15.5 mm		A321	3239-3440
				9.5 mm			
SAM.A10560				14.0 mm	17.0 mm	A319	2690-2727
SAM.A10458				15.0 mm	11.0 mm	A192	2708

(ovig.)
Carapace lengths of ovigerous ♀♀ (from IK material): 14.0 mm, 14.3 mm, 15.0 mm, 15.5 mm, 17.6 mm, 18.4 mm, 19.3 mm.

AcanthePHYra gracilipes Chace, 1940

Figs 12, 14 a

AcanthePHYra gracilipes Chace, 1940: 149. Sivertsen & Holthuis, 1956: 6.

Description

Integument soft, thin; the specimen somewhat damaged. Eyestalk shorter than rostrum, widest at distal end. Small blunt tubercle on inner angle of stalk, just behind eye. Rostrum acutely triangular, with six dorsal teeth and no ventral teeth, reaching end of second joint of antennular peduncle. Dorsal carina ends before reaching the cervical groove. Tiny antennal spine present, also a slightly larger branchiostegal spine, unsupported by a keel. Distinct ridge-like keel above branchial region, stretching from hepatic region almost to the posterior margin. First two abdominal segments dorsally smooth, last four segments dorsally carinate, ending in short spines (spines of fifth segment broken). Distal portion of telson missing. Endopod of uropod equal in length to the sixth abdominal segment. Pereiopods long and slender.

Previous record

Off Bermuda.

Material

SAM.A10565, ♂, carapace length 16 mm, overall length \pm 50 mm. St. A190, 2269 m.




FIG. 12. *AcanthePHYRA gracilipes* Chace.

Remarks

This specimen agrees almost exactly with the original description of Chace (1940). The mandible is toothed over its entire length, confirming the genus. This is the first record of this species from South African waters and possibly from the southern hemisphere.

AcanthePHYRA brevirostris Smith, 1885

Figs 13, 14 b

AcanthePHYRA brevirostris Smith, 1885: 504. Balss, 1925: 252. Chace, 1940: 148. Sivertsen & Holthuis, 1956: 5.

Hymenodora duplex Bate, 1888: 843.

Description

Integument thin. Eye wider than eyestalk. Rostrum acutely triangular, reaching to end of second joint of antennular peduncle, with eight dorsal teeth; no ventral teeth. Posterior portion of carapace not carinate. Tiny antennal spine present, also a slightly larger branchiostegal spine, the anterior portion of which is supported by a slight keel. A slightly keeled ridge in the branchial region, reaching almost to posterior margin of the carapace. First two abdominal segments dorsally smooth, third to sixth segments dorsally carinate, each ending in a tooth; that of the third segment a large fleshy leaf-like structure, giving the species a distinctive appearance. Teeth of segments four to six small and sharp. Tip of telson missing.



FIG. 13. *AcanthePHYRA brevirostris* Smith.

FIG. 14.

a. *AcanthePHYRA gracilipes* Chace. Mandible. b. *AcanthePHYRA brevirostris* Smith. Mandible.

Previous records and distribution

North-east coast of U.S.A., Bermuda, Bahamas, off Portugal, West Africa, south-west Indian Ocean, off Pacific coast of Ecuador, off Marion Island.

Material

SAM.A10564, carapace length 19 mm, overall length 54 mm. St. A192, 2708 m.

Remarks

The mandible is almost identical to that of *A. gracilipes* and is dentate throughout its length. This is the first record of the specimens from South African waters. The closest record to South Africa was that of a specimen taken by the *Challenger*, off Marion Island.


Acanthephyra corallina (Milne Edwards, 1883)

Figs 15, 16, 17

Notostomus corallina Milne Edwards, 1883.*Acanthephyra valdiviae* Balss, 1914: 595. 1925: 260.*Acanthephyra corallina*: Chace, 1936: 27.*Description*

Integument firm. Carapace carinate throughout its length. A notch present in the dorsal carina, about two-thirds down the carapace. Rostrum stout, extending a little way past the antennal scales, curving slightly upward, supported by lateral keels which end some distance posterior to the orbits.

Rostral teeth: 18/3, 18/4, 17/3, 20/3, 22/3, 19/4, seven or eight teeth posterior to the orbit. The distance between the orbital groove and the dorsal carina equal to the distance between the orbital groove and the branchiostegal spine. Latter stout, outwardly flared, supported by a prominent keel. Antennal spine small. Strong hepatic spine at base of cervical groove, which is not well defined. Cervical groove joined by orbital groove. Branchiostegal keel joins with a well-developed keel in mid-branchial region. In the posterior portion of the carapace, this keel curves upwards and joins with a ridge which marks the upper border of the branchial region. Ventral branchial keel stretches from below branchiostegal keel to posterior margin of carapace, curving upward in this region. All abdominal segments dorsally keeled; segments 3-6 each ending

FIG. 15. *Acanthephyra corallina* (Milne Edwards).




FIG. 16. *Acanthephyra corallina* (Milne Edwards).
a. Mandible. b. Telson and left uropod.

in a strong tooth. Dorso-ventral length of second abdominal segment equals the greatest dorso-ventral length of the carapace. Eyes slightly wider than eyestalks. Antennal scales narrow. Mandibular palp short, three-jointed; seven teeth on cutting edge. Maxilliped 3 as stout as the pereiopods, reaching almost to the end of the antennal scale. Pereiopod 4 the longest, but only slightly longer than the other pereiopods. Meri of pereiopods 3, 4, 5 armed on posterior border with a single row of spinules. Dactyls of pereiopods 3 and 4 short, slender; that of pereiopod 5 reduced. Propodus of latter has a series of short stiff bristles at its distal end. Slender exopods on maxillipeds 2 and 3 and on all the pereiopods. Pleopods stout with well-developed endo- and exopods. Telson with four pairs of dorso-lateral spinules in distal half and pair of terminal spines flanking acute apex. The apex appears to be worn down with age, as some of the larger specimens do not have the sub-apical pair of spines. Exopod of uropod almost equal in length to telson, with two spines next to one another on the outer margin, some distance from the apex. Endopod slightly shorter than exopod.

Material

		♀	♂			
		Carapace length	Carapace length	Overall length	Station	Depth (m)
SAM.A12531		39.5 mm	149 mm	A319	2690-2727
SAM.A12532 (ovig.)	34.0 mm		138 mm	A319	2690-2727
SAM.A12533	27.0 mm		111 mm	A190	2269
SAM.A10523 (ovig.)	37.0 mm		134 mm	A318	2525-2782
		33.5 mm		132 mm	A318	2525-2782
			33.5 mm	129 mm	A318	2525-2782
			26.0 mm	99 mm	A318	2525-2782




FIG. 17. *Acantheephyra corallina* (Milne Edwards).
 a. Tip of pereopod 5. b. Tip of maxilliped 3. c. Maxilliped 2. d. Maxilliped 1.
 e. Maxilla 2. f. Pleopod 1 of male. g. Pleopod 2 of male.

Remarks

This is a new record for the South African region. The species was previously recorded from the mid-Indian Ocean, where a single male was caught by the *Valdivia*. The presence of a hepatic spine on the carapace together with the deep rostral base serves to distinguish this uncommon species.

Family **Nematocarcinidae***Nematocarcinus longirostris* Bate, 1888

Nematocarcinus longirostris Bate, 1888: 806. Stebbing, 1914: 44. Calman, 1925: 15. Barnard, 1950: 671, fig. 125 a-k.

Previous records and distribution

Off Cape Point, 1200 fms (2360 m), Marion Island, Japan, East Indies, west coast of South America.

Material

A total of 78 specimens of this species was obtained from the following stations: A189, A190, A191, A192, A193, A315, A316, A317, A318, A319. Of the 78, 10 were ovigerous females, with a carapace length varying from 28–34 mm, while 15 were mature males, with a carapace length varying from 23 to 29 mm.

Nematocarcinus parvidentatus Bate, 1888

Nematocarcinus parvidentatus Bate, 1888: 814. Stebbing, 1915: 99. Barnard, 1950: 674, fig. 125 l-o.

Previous records and distribution

Off Durban, 440 fms (865 m), off East London, 400 fms (780 m), off Cape Point, 900 fms (1770 m), Japan.

Material

A total of 108 specimens of this species was obtained from the following stations: A190, A192, A193, A315, A316, A317, A318, A319, A322. Of the 108, 15 were ovigerous females with a carapace length varying from 19–26 mm, while 20 were mature males with a carapace length varying from 17 to 22 mm.

Remarks

Several specimens of either *parvidentatus* or *longirostris* from most of the stations were so damaged that specific identification was impossible. As can be expected from two species so closely related, the mouthparts are almost identical and are of no use in distinguishing the species. The most useful characters are the lengths of the carapace at which the males and females become mature, as well as the rostral shape. In *parvidentatus* the minimum carapace length of mature males and females is usually less than that of *longirostris*. The rostrum is variable, relative length being unsatisfactory for specific separation. In general, *parvidentatus* has a rostrum broader in the vertical plane compared with its length than *longirostris*. The ventral sinuosity at the base of the rostrum in

parvidentatus noted by Barnard (1950) is also a useful character. A bopyrid isopod was found on a specimen of *longirostris* but is neither of the two bopyrids previously recorded on *Nematocarcinus* from South Africa.

Family **Glyphocrangonidae**

Glyphocrangon sculptus (S. I. Smith, 1883)

Rhachocaris sculpta Smith, 1883: 49.

Glyphocrangon sculptus: Smith, 1884: 365. Stebbing, 1908: 37. Barnard, 1950: 719, fig. 134 a-d.

Previous records and distribution

Off Cape Point, 1000 fms (1970 m), east coast of North America.

Material				♀	♂	Juveniles	Station	Depth (m)
				Carapace length	Carapace length			
SAM.A10521		20.5 mm 21.0 mm 22.8 mm	4	A318	2525-2782
SAM.A10515		22.8 mm	2	A318	2525-2782
SAM.A10535	21.0 mm	22.1 mm 22.5 mm		A319	2690-2727
SAM.A10451	26.0 mm (ovig.) 25.5 mm (ovig.) 25.0 mm (ovig.) 25.0 mm (ovig.) 24.0 mm (ovig.) 21.5 mm 22.0 mm 19.3 mm 22.8 mm 21.0 mm	21.1 mm 19.5 mm 19.0 mm 21.3 mm 21.0 mm 17.5 mm 18.9 mm 17.5 mm 18.9 mm 15.6 mm 15.6 mm		A190	2269
SAM.A12534			1	A193	2745
SAM.A10459			2	A192	2708
SAM.A10547			5	A322	2745
SAM.A10462			2	A193	2745
SAM.A10464			1	A193	2745
SAM.A10549			1	A322	2745
SAM.A10539			4	A319	2745

Family **Crangonidae**

Sclerocrangon bellmarleyi Stebbing, 1914

Sclerocrangon bellmarleyi Stebbing, 1914: 29. Barnard, 1950: 804, fig. 152.

Previous records and distribution

Off Durban, 440 fms (865 m), Natal coast, 400 fms (780 m).

Material

SAM.A10446, ♀ ovigerous, carapace length 13 mm, overall length 52 mm.
St. A189, 1098 m.

Remarks

This ovigerous female is the largest specimen of this species yet obtained. The eggs, of which only four remain, measure 2.5 mm in length and are in an advanced state of development, the shape of the larvae being clearly visible. This is the most southerly record of this apparently endemic species.

Pontophilus occidentalis Faxon var. *indica* de Man, 1918

Figs 18, 19

Pontophilus occidentalis var. *indica* de Man, 1918: 161. 1920: 264.

Description

Carapace with median dorsal keel bearing one cardiac and two gastric spines, the more anterior of which always the smaller. Two lateral carapace spines, one hepatic, one epibranchial, lying in an oblique plane. Well-developed antennal and branchiostegal spines, latter supported by a blunt keel. Minute post-orbital spinule above the post-orbital fissure. Antero-lateral angle of carapace with a minute spine. Rostrum varies in length in relation to eyes and in general shape; usually with one or two pairs of minute denticles at its base. Eyes large and rounded, tending to obscure the stalks. Pereiopod 1 stout, armed with the characteristic crangonid chela. Pereiopod 2 about half the




FIG. 18. *Pontophilus occidentalis* var. *indica* de Man.




FIG. 19. *Pontophilus occidentalis* var. *indica* de Man.

a. Rostrum and eyes, showing variations. b. Chela of pereopod 1. c. Tip of telson. d. Pleopod 2 of male.

length of pereopod 1; chelate, very slender. A rounded knob ending in a tiny spine present between bases of second pair of pereiopods. Pereiopod 3 longer than pereopod 1, very slender. Pereiopods 4 and 5 almost equal in length to pereopod 3, but slightly stouter. Abdomen dorsally smooth, the sixth segment at least twice the length of the fifth. Appendix masculina of pleopod 2 of the male shorter and stouter than the appendix interna. Telson with two pairs of minute lateral spines ending in three pairs of spines, the submedian pair being the longest.

Previous records and distribution

East Indies, in region of Makassar Straits.

Material

A total of about 90 specimens from the following stations was obtained: A191, A192, A193, A315, A316, A317, A318, A319, A321, A322.

The carapace lengths of ovigerous females varied between 11 and 12.5 mm.

Remarks

De Man (1920) distinguishes *Pontophilus occidentalis* Faxon from its variety *indica* by its smaller size (48 mm as against 73 mm), the almost microscopical size of the anterior gastric spine and the length of the abdomen. In *occidentalis* the abdomen is two and a half times the length of the carapace, while in *occidentalis* var. *indica* it is three times the carapace length. De Man (1920) noted that the closely related species *gracilis* Smith, *abyssi* Smith, *challengeri* Ortmann, *junceus* Bate, *profundus* Bate, *occidentalis* Faxon, and *occidentalis* var. *indica* de Man, might prove to be geographical races of a widely distributed species. It certainly is difficult to distinguish between the species and its variety. In all the present specimens, the first gastric spine is always smaller than the second (characteristic of the variety). The ratio between carapace length and abdominal length is very variable, being anything from 2.2 to 3. This criterion is thus not reliable in distinguishing the variety from the species. The largest specimen had an overall length of 53.1 mm, only slightly larger than de Man's limit of 48 mm. This is the first record of the species from the South African region. *Pontophilus gracilis* Smith, known from off the Cape Peninsula, has been recorded from depths of 190, 250, 470 fms (370, 490, 925 m), while the present species has been taken from depths of 2525–3440 m.

SUMMARY

A collection of deep-sea decapod Crustacea from west of Cape Point, South Africa, in depths between 1098 and 3440 metres, is described. The collection includes approximately 480 specimens of 35 species, of which 3 are new species and 12 are new records.

ACKNOWLEDGEMENTS

The trawling was done by courtesy of the Director, Division of Sea Fisheries, Cape Town, to whom we are very grateful.

I am indebted to Dr. M.-L. Penrith of the South African Museum for reading the manuscript and for making constructive criticisms and suggestions throughout the preparation of this work.

The Trustees of the South African Museum are grateful to the Council for Scientific and Industrial Research for the award of a grant to publish this paper.

REFERENCES

- BALSS, H. 1925. Macrura der Deutschen Tiefsee-Expedition. 2. Natantia, Teil A. *Wiss. Ergebn. dt. Tiefsee-Exped. 'Valdivia'* **20**: 217-315.
- BALSS, H. 1927. Macrura der Deutschen Tiefsee-Expedition. 3. Natantia, Teil B. *Wiss. Ergebn. dt. Tiefsee-Exped. 'Valdivia'* **23**: 247-275.
- BARNARD, K. H. 1947. Descriptions of new species of South African decapod Crustacea, with notes on synonymy and new records. *Ann. Mag. nat. Hist.* (11) **13**: 361-392.
- BARNARD, K. H. 1950. Descriptive catalogue of South African decapod Crustacea. *Ann. S. Afr. Mus.* **38**: 1-837.
- BATE, C. S. 1881. On the Penaeidae. *Ann. Mag. nat. Hist.* (5) **8**: 169-196.
- BATE, C. S. 1888. Report on the Crustacea Macrura collected by H.M.S. *Challenger* during the years 1873-1876. *Rep. Voy. Challenger, 1873-1876* **24**: 1-942.
- BELLOC, G. & LORILLOU, C. 1961. Catalogue des types de décapodes du Musée océanographique de Monaco. *Bull. Inst. oceanogr. Monaco* no. 1212: 1-19.
- BENEDICT, J. E. 1903. Descriptions of a new genus and forty-six new species of crustaceans of the family Galatheididae with a list of the known marine species. *Proc. U.S. natn. Mus.* **26**: 243-334.
- BERNARD, F. 1953. Decapoda Eryonidae (*Eryoneicus* et *Willemoesia*). *Dana Rep.* **6** (37): 1-93.
- BOUVIER, E. L. 1905. Sur les palinurides et la eryonides recueillies dans l'Atlantique oriental par les expéditions françaises et monagasques. *C.r. hebdom. Séanc. Acad. Sci., Paris* **140**: 479-482.
- BOUVIER, E. L. 1917. Crustacés décapodes (macroures marcheurs). *Résult. Camp. scient. Prince Albert* **1** **50**: 1-140.
- BOUVIER, E. L. 1922. Observations complémentaires sur les crustacés décapodes (abstraction faite des carides), provenant des campagnes de S.A.S. le Prince de Monaco. *Résult. Camp. scient. Prince Albert* **1** **62**: 1-103.
- BURKENROAD, M. D. 1936. The Aristaeinae, Solenocerinae and pelagic Penaeinae of the Bingham Oceanographic Collection. *Bull. Bingham oceanogr. Coll.* **5** (2): 1-51.
- BURKENROAD, M. D. 1937. The Templeton Crocker Expedition. 12. Sergestidae (Crustacea decapoda) from the lower Californian region, with descriptions of the two new species and some remarks on the organs of Pesta in *Sergestes*. *Zoologica, N.Y.* **22**: 315-329.
- CALMAN, W. T. 1925. On the macrurous decapod Crustacea collected in South African waters by the s.s. 'Pickle'. *Rep. Fish. mar. biol. Surv. Un. S. Afr.* **4** Spec. Report 3: 1-26.
- CHACE, F. A. 1936. Revision of the bathypelagic prawns of the family Acanthephyridae with notes on a new family Gomphonotidae. *J. Wash. Acad. Sci.* **26**: 24-31.
- CHACE, F. A. 1940. The plankton of the Bermuda Oceanographic Expedition. 9. The bathypelagic caridean Crustacea. *Zoologica, N.Y.* **25**: 117-209.
- CHACE, F. A. 1942. Report on the scientific results of the Atlantis Expedition to the West Indies, under the joint auspices of the University of Havana and Harvard University. *Torrea* **11**: 1-106.
- DOFLEIN, F. & BALSS, H. 1926. Die Galatheiden der Deutschen Tiefsee-Expedition. *Wiss. Ergebn. dt. Tiefsee-Exped. 'Valdivia'* **20**: 1-184.
- FAXON, W. 1893. Reports on the dredging operations off the west coast of central America to the Galapagos, to the west coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz, carried on by the U.S. Fish Commission steamer 'Albatross' during 1891, Lieut. Commander Z. L. Tanner, U.S.N. commanding. *Bull. Mus. comp. Zool. Harv.* **24**: 149-220.
- FAXON, W. 1895. Reports on an exploration of the west coasts of Mexico, central and South America, and off the Galapagos Islands, in charge of Alexander Agassiz, by the U.S. Fish Commission steamer 'Albatross' during 1891, Lieut. Commander Z. L. Tanner, U.S.N. commanding. *Mem. Mus. comp. Zool. Harv.* **18**: 1-292.
- GORDON, I. 1935. New or imperfectly known species of Crustacea Macrura. *J. Linn. Soc. Zool.* **39**: 307-351.
- GORDON, I. 1939. A new species of *Sergestes* (Crustacea, Decapoda) from the South Atlantic. *Ann. Mag. nat. Hist.* (11) **4**: 498-509.
- GRINDLEY, J. R. & PENRITH, M. J. 1965. Notes on the bathypelagic fauna of the seas around South Africa. *Zool. afr.* **1**: 275-295.
- HANSEN, H. J. 1896. On the development and the species of the crustaceans of the genus *Sergestes*. *Proc. zool. Soc. Lond.* **1896**: 936-970.

- HANSEN, H. J. 1903. On the Crustacea of the genera *Petalidium* and *Sergestes* from the 'Challenger' with an account of luminous organs in *Sergestes challengerii* n.sp. *Proc. zool. Soc. Lond.* **1903**: 52-79.
- HANSEN, H. J. 1922. Crustacés décapodes (Sergestides) provenant des campagnes des yacht *Hirondelle* et *Princesse-Alice* (1885-1915). *Résult. Camp. scient. Prince Albert I* **64**: 1-232.
- HOLTHUIS, L. B. 1955. The recent genera of the caridean and stenopodidean shrimps (class Crustacea, order Decapoda, supersection Natantia) with keys for their determination. *Zool. Verh., Leiden* **26**: 1-157.
- KEMP, S. 1909. The decapods of the genus *Gennadas* collected by H.M.S. 'Challenger'. *Proc. zool. Soc. Lond.* **11**: 718-730.
- KEMP, S. 1910. The decapod Natantia of the coasts of Ireland. *Scient. Invest. Fish. Brch Ire.* **1908** (1): 1-190.
- KEMP, S. 1939. On *Acanthephyra purpurea* and its allies (Crustacea, Decapoda, Hoplophoridae). *Ann. Mag. nat. Hist.* (11) **4**: 568-579.
- KRÖYER, H. 1855. Bidrag til kundskab om brebsdyrslægten *Sergestes*. *Over. K. danske Vidensk. Selsk. Forh.* **1855**: 22-23.
- MAN, J. G. de. 1918. Diagnoses of new species of macrurous decapod Crustacea from the Siboga Expedition. *Zool. Meded. Leiden* **4**: 159-166.
- MAN, J. G. de. 1920. The Decapoda of the Siboga Expedition. *Siboga Exped. monogr.* 39 a³: 1-320.
- MARTENS, E. von. 1868. Ueber einige ostasiatischen Süßwasserthiere. 11. Crustacea. *Arch. Naturgesch.* **34**: 1-64.
- MILNE-EDWARDS, A. 1880. Études préliminaires sur les crustacés. 1. *Bull. Mus. comp. Zool. Harv.* **8**: 1-67.
- MILNE-EDWARDS, A. 1930 after 1883. *Recueil de figures de Crustacés nouveau ou peu connus*. Paris.
- MILNE-EDWARDS, A. 1930. Description des genres *Glaucothoe*, *Sicyonie*, *Sergeste* et *Acète* de l'ordre des crustacés décapodes. *Annl. Sci. nat. Zool.* **19**: 333-352.
- MILNE-EDWARDS, A. & BOUVIER, E. L. 1894. Considerations générales sur la famille des Galatheides. *Annl. Sci. nat. Zool.* (7) **16**: 191-327.
- ORTMANN, A. 1893. Decapoden und Schizopoden der Plankton-Expedition. *Ergeb. Atlant. Ozean Planktonexped. Humboldt-Stift.* **2**: 1-210.
- PESTA, O. 1918. *Die Decapodenfauna der Adria. Versuch einer Monographie*. Leipzig & Wien. (not seen).
- PESTA, O. 1918. *Die Decapodenfauna der Adria. Versuch einer Monographie*. Leipzig & Wien.
- SIVERTSEN, E. & HOLTHUIS, L. G. 1956. Crustacea Decapoda (the Penaeidae and Stenopodidae excepted). *Rep. scient. Results Michael Sars. N. Atlant. deep Sea Exped.* **5** (12): 1-54.
- SMITH, S. I. 1882. Reports on the results of dredgings, under the supervision of Alexander Agassiz, on the east coast of the United States, during the summer of 1880, by the U.S. Coast Survey Steamer 'Blake', Commander J. R. Bartlett, U.S.N. commanding. Report on the Crustacea. Part I. Decapoda. *Bull. Mus. comp. Zool. Harv.* **10**: 1-108.
- SMITH, S. I. 1884. Crustacea of the 'Albatross' dredgings in 1883. *Rep. U.S. Commn. Fish.* **1882**: 356.
- SMITH, S. I. 1885. On some new or little known Decapoda Crustacea from recent fish commission dredgings off the east coast of the United States. *Proc. U.S. natn. Mus.* **7**: 493-511.
- STEBBING, T. R. R. 1905. South African Crustacea, part 3. *Mar. Invest. S. Afr.* **3**: 21-120.
- STEBBING, T. R. R. 1908. South African Crustacea, part 4. *Ann. S. Afr. Mus.* **6**: 1-96.
- STEBBING, T. R. R. 1910. General catalogue of the South African Crustacea (part 5). *Ann. S. Afr. Mus.* **6**: 281-293.
- STEBBING, T. R. R. 1914. South African Crustacea (part 7). *Ann. S. Afr. Mus.* **15**: 1-55.
- STEBBING, T. R. R. 1915. South African Crustacea (part 8). *Ann. S. Afr. Mus.* **15**: 57-104.
- STEBBING, T. R. R. 1917. South African Crustacea (part 9). *Ann. S. Afr. Mus.* **17**: 23-46.
- SUND, O. 1920. The 'Challenger' Eryonidae (Crustacea). *Ann. Mag. nat. Hist.* (9) **6**: 220.
- TIRMIZI, N. M. 1960. Crustacea, Penaeidae, part 2. Series Benthescyymae. *Scient. Rep. John Murray Exped.* **10**: 319-383.
- WILLEMOES-SUHM, R. von. 1873. In Thomson, C. W. Notes from the 'Challenger'. *Nature, Lond.* **8**: 51.
- YALDWYN, J. C. 1957. Deep water Crustacea of the genus *Sergestes* (Decapoda Natantia) from Cook Strait, New Zealand. *Zoology Publ. Vict. Univ. Wellington* **22**: 1-27.

INSTRUCTIONS TO AUTHORS

MANUSCRIPTS

In duplicate (one set of illustrations), type-written, double spaced with good margins, including TABLE OF CONTENTS and SUMMARY. Position of text-figures and tables must be indicated.

ILLUSTRATIONS

So proportioned that when reduced they will occupy not more than $4\frac{1}{2}$ in. = 7 in. ($7\frac{1}{2}$ in. including the caption). A scale (metric) must appear with all photographs.

REFERENCES

Authors' names and dates of publication given in text; full references at end of paper in alphabetical order of authors' names (Harvard system). References at end of paper must be given in this order:

Name of author, in capitals, followed by initials; names of joint authors connected by &, not 'and'. Year of publication; several papers by the same author in one year designated by suffixes a, b, etc. Full title of paper; initial capital letters only for first word and for proper names (except in German). Title of journal, abbreviated according to *World list of scientific periodicals* and underlined (italics). Series number, if any, in parenthesis, e.g. (3), (n.s.), (B.). Volume number in arabic numerals (without prefix 'vol.'). with wavy underlining (bold type). Part number, only if separate parts of one volume are independently numbered. Page numbers, first and last, preceded by a colon (without prefix 'p'). Thus:

SMITH, A. B. 1956. New *Plonia* species from South Africa. *Ann. Mag. nat. Hist.* (12) 9: 937-945.

When reference is made to a separate book, give in this order: Author's name; his initials; date of publication; title, underlined; edition, if any; volume number, if any, in arabic numerals, with wavy underlining; place of publication; name of publisher. Thus:

BROWN, X. Y. 1953. *Marine faunas*. 2nd ed. 2. London: Green.

When reference is made to a paper forming a distinct part of another book, give: Name of author of paper, his initials; date of publication; title of paper; 'In', underlined; name of author of book; his initials; title of book, underlined; edition, if any; volume number, if any, in arabic numerals, with wavy underlining; pagination of paper; place of publication; name of publisher. Thus:

SMITH, C. D. 1954. South African *Plonias*. In Brown, X. Y. *Marine faunas*. 2nd ed. 3: 63-95. London: Green.

SYNONYMY

Arranged according to chronology of names. Published scientific names by which a species has been previously designated (subsequent to 1758) are listed in chronological order, with abbreviated bibliographic references to descriptions or citations following in chronological order after each name. Full references must be given at the end of the paper. Articles and recommendations of the *International code of zoological nomenclature adopted by the XV International congress of zoology, London, July 1958*, are to be observed (particularly articles 22 and 51).

Examples: *Plonia capensis* Smith, 1954: 86, pl. 27, fig. 3. Green, 1955: 23, fig. 2.

When transferred to another genus:

Euplonia capensis (Smith) Brown, 1955: 259.

When misidentified as another species:

Plonia natalensis (non West), Jones, 1956: 18.

When another species has been called by the same name:

[non] *Plonia capensis*: Jones, 1957: 27 (= *natalensis* West).